

# *Algebra & Number Theory*

Volume 11  
2017  
No. 4

On Hilbert's 17th problem in low degree

Olivier Benoist





# On Hilbert's 17th problem in low degree

Olivier Benoist

Artin solved Hilbert's 17th problem, proving that a real polynomial in  $n$  variables that is positive semidefinite is a sum of squares of rational functions, and Pfister showed that only  $2^n$  squares are needed.

In this paper, we investigate situations where Pfister's theorem may be improved. We show that a real polynomial of degree  $d$  in  $n$  variables that is positive semidefinite is a sum of  $2^n - 1$  squares of rational functions if  $d \leq 2n - 2$ . If  $n$  is even or equal to 3 or 5, this result also holds for  $d = 2n$ .

## Introduction

**Hilbert's 17th problem.** Let  $\mathbf{R}$  be a real closed field, for instance the field  $\mathbb{R}$  of real numbers, and let  $n \geq 1$ . A polynomial  $f \in \mathbf{R}[X_1, \dots, X_n]$  is said to be positive semidefinite if  $f(x_1, \dots, x_n) \geq 0$  for all  $x_1, \dots, x_n \in \mathbf{R}$ . As an odd degree polynomial changes sign, such a polynomial has even degree.

Artin [1927] answered Hilbert's 17th problem by proving that a positive semidefinite polynomial  $f \in \mathbf{R}[X_1, \dots, X_n]$  is a sum of squares of rational functions.<sup>1</sup> This theorem was later improved by Pfister [1967, Theorem 1], who showed that it is actually the sum of  $2^n$  squares of rational functions. We refer to [Pfister 1995, Chapter 6] for a nice account of these classical results.

In two variables, the situation is very well understood. Hilbert [1888] showed that a positive semidefinite polynomial  $f \in \mathbf{R}[X_1, X_2]$  of degree  $\leq 4$  is a sum of 3 squares of rational functions,<sup>2</sup> and Cassels, Ellison and Pfister [Cassels et al. 1971] gave an example of a positive semidefinite polynomial  $f \in \mathbf{R}[X_1, X_2]$  of degree 6 that is not a sum of 3 squares of rational functions.

Our goal is to prove an analogue of Hilbert's result — that in low degree, less squares are needed — in more than two variables:

---

MSC2010: primary 11E25; secondary 14F20, 14P99.

Keywords: Hilbert's 17th problem, sums of squares, real algebraic geometry, Bloch–Ogus theory.

<sup>1</sup>Hilbert himself [1888] had given examples of positive semidefinite polynomials that are not sums of squares of polynomials.

<sup>2</sup>In fact, in this exceptional case, Hilbert actually showed that squares of polynomials suffice. We will not consider this question in what follows, and refer the interested reader to [Pfister and Scheiderer 2012].

**Theorem 0.1.** *Let  $f \in \mathbf{R}[X_1, \dots, X_n]$  be a positive semidefinite polynomial of degree  $d$ . Suppose that one of the following holds:*

- (i)  $d \leq 2n - 2$ .
- (ii)  $d = 2n$ , and either  $n$  is even,  $n = 3$ , or  $n = 5$ .

*Then  $f$  is a sum of  $2^n - 1$  squares in  $\mathbf{R}(X_1, \dots, X_n)$ .*

Of course, when  $d = 2$ , the classification of quadratic forms over  $\mathbf{R}$  shows the much stronger result that  $n + 1$  squares are enough. However, to the best of our knowledge, our theorem is already new for  $d = 4$  and  $n \geq 3$ .

**Dependence on the degree.** The question whether the bound  $2^n$  in Pfister's aforementioned theorem is optimal is natural and well known [Pfister 1971, §4, Problem 1]. It is often formulated in the following equivalent way, where the Pythagoras number  $p(K)$  of a field  $K$  is the smallest number  $p$  such that every sum of squares in  $K$  is a sum of  $p$  squares:

**Question 0.2.** Do we have  $p(\mathbf{R}(X_1, \dots, X_n)) = 2^n$ ?

When  $n \geq 2$ , the best known result is that  $n + 2 \leq p(\mathbf{R}(X_1, \dots, X_n)) \leq 2^n$  [Pfister 1995, p. 97], where the upper bound is Pfister's theorem and the lower bound is an easy consequence of the Cassels–Ellison–Pfister theorem.

Our main theorem does not address this question directly; it explores the opposite direction, that is, the values of the degree for which Pfister's bound may be improved. However, Theorem 0.1 gives insights into Question 0.2. The bound  $d \leq 2n$  has a natural geometric origin (it reflects the rational connectedness of an associated algebraic variety), and it would be natural to expect that Theorem 0.1 cannot be extended to degrees  $d \geq 2n + 2$ .

In view of Theorem 0.1, it is natural to ask whether the bound  $d \leq 2n - 2$  may be improved to  $d \leq 2n$  for every odd value of  $n$ . When  $n = 1$ , this is not the case because  $X_1^2 + 1$  is not a square. On the other hand, when  $n \geq 3$  is odd, we reduce this question to a geometric coniveau estimate (Proposition 6.3). When  $n = 3$ , it is very easy to check. We also verify it when  $n = 5$ , following an argument of Voisin. This explains the hypotheses on the degree in Theorem 0.1.

**Strategy of the proof.** In two variables, the theorems of Hilbert and Cassels–Ellison–Pfister quoted above have received geometric proofs by Colliot-Thélène [1992, Remark 2; 1993]. His idea is to consider the homogenization  $F$  of  $f$  and to introduce the algebraic surface  $Y := \{Z^2 + F = 0\}$ . Then, whether or not  $f$  may be written as a sum of three squares in  $\mathbf{R}(X_1, X_2)$  depends on the injectivity of the map  $\mathrm{Br}(\mathbf{R}) \rightarrow \mathrm{Br}(\mathbf{R}(Y))$ , which may be studied by geometrical methods.

We follow the same strategy in more variables. Proposition 3.2 and Proposition 3.3 translate the property that  $f$  is a sum of  $2^n - 1$  squares in  $\mathbf{R}(X_1, \dots, X_n)$  into a

cohomological property of (a resolution of singularities of) the variety  $Y$ . The group that plays a role analogous to that of the Brauer group in two variables is a degree  $n$  unramified cohomology group.

It remains to show that when the degree of  $f$  is small, some class in a degree  $n$  unramified cohomology group vanishes. This is more difficult than the corresponding result in two variables, as these groups are harder to control than Brauer groups. Our main tool to achieve this is Bloch–Ogus theory.

**Structure of the paper.** The first two sections gather general cohomological results for varieties over  $\mathbf{R}$ , which are used throughout the text. It will be very important for us to use cohomology with integral coefficients (as opposed to 2-torsion coefficients). For this reason, Section 1 is devoted to general properties of the 2-adic cohomology<sup>3</sup> of varieties over  $\mathbf{R}$ .

In Section 2, we recall the basics of Bloch–Ogus theory, then focus on the specific properties of it over real closed fields. In particular, we adapt to our needs a strategy of Colliot-Thélène and Scheiderer [1996] to compare the Bloch–Ogus theory of a variety over  $\mathbf{R}$  and over the algebraic closure  $\mathbf{C}$  of  $\mathbf{R}$ , and explain in our context consequences of the Bloch–Kato conjectures discovered by Bloch and Srinivas [1983] and extended by Colliot-Thélène and Voisin [2012].

We study when a positive semidefinite polynomial  $f \in \mathbf{R}[X_1, \dots, X_n]$  is a sum of  $2^n - 1$  squares of rational functions in Section 3. We successively relate this property to the level of the function field  $\mathbf{R}(Y)$  of the variety  $Y := \{Z^2 + F = 0\}$  in Proposition 3.2 (this is due to Pfister), to degree  $n$  unramified cohomology of  $Y$  in Proposition 3.3 (an important tool is Voevodsky's solution [2003] to the Milnor conjecture) and to degree  $n + 1$  cohomology of  $Y$  in Proposition 3.5 (this is the crucial step, which uses Bloch–Ogus theory, and where the rational connectedness of  $Y$  plays a role).

Section 4 contains the cohomological computations on the variety  $Y$  that are relevant to apply the results of Section 3. Section 4D will only be useful when  $n$  is odd and  $d = 2n$ , and is complemented by a geometric coniveau estimate in Section 5. The reader who is not interested in our partial and conditional results when  $n \geq 3$  is odd and  $d = 2n$  may skip them.

Section 6 completes the proof of Theorem 0.1. For a generic choice of  $f$  (that is, when the degree of  $f$  is maximal among the values allowed in the statement of Theorem 0.1, and  $Y$  is a smooth variety), this is an immediate consequence of the results obtained so far. In general, we do not know how to apply this argument directly, because we do not have a good control on the geometry of (a resolution of singularities of)  $Y$ . Instead, we rely on a specialization argument. This argument

---

<sup>3</sup>It would also have been possible to work with equivariant Betti cohomology over the field  $\mathbb{R}$  of real numbers [Krasnov 1994], and with its semialgebraic counterpart over a general real closed field.

reduces Theorem 0.1 to the generic case, but over a bigger real closed field. In particular, even if one is only interested in proving Theorem 0.1 over  $\mathbb{R}$ , one has to work over real closed fields that are not necessarily archimedean.

### 1. Cohomology of real varieties

Let  $\mathbf{R}$  be a real closed field and  $\mathbf{C}$  be an algebraic closure of  $\mathbf{R}$ . We denote the Galois group by  $G := \text{Gal}(\mathbf{C}/\mathbf{R}) \simeq \mathbb{Z}/2\mathbb{Z}$ . A variety over  $\mathbf{R}$  is a separated scheme of finite type over  $\mathbf{R}$ .

**1A. 2-adic cohomology.** If  $X$  is a variety over  $\mathbf{R}$ , we denote by  $H^k(X, \mathbb{Z}/2^r\mathbb{Z}(j))$  its étale cohomology groups. These cohomology groups are finite; this follows from the Hochschild–Serre spectral sequence

$$E_2^{p,q} = H^p(G, H^q(X_{\mathbf{C}}, \mathbb{Z}/2^r\mathbb{Z}(j))) \Rightarrow H^{p+q}(X, \mathbb{Z}/2^r\mathbb{Z}(j))$$

using the facts that  $X_{\mathbf{C}}$  has finite cohomological dimension [SGA 4<sub>3</sub> 1973, X Corollaire 4.3], that the groups  $H^q(X_{\mathbf{C}}, \mathbb{Z}/2^r\mathbb{Z}(j))$  are finite [SGA 4<sub>3</sub> 1973, XVI Théorème 5.1] and that a finite  $G$ -module has finite cohomology.

Let us define  $H^k(X, \mathbb{Z}_2(j)) := \varprojlim_r H^k(X, \mathbb{Z}/2^r\mathbb{Z}(j))$ . Since the Galois cohomology of finite  $G$ -modules is finite, [Jannsen 1988, Remark 3.5(c)] shows that these groups coincide with the continuous étale cohomology groups defined by Jannsen. In particular, we have a Hochschild–Serre spectral sequence [Jannsen 1988, Remark 3.5(b)]:

$$E_2^{p,q} = H^p(G, H^q(X_{\mathbf{C}}, \mathbb{Z}_2(j))) \Rightarrow H^{p+q}(X, \mathbb{Z}_2(j)). \quad (1-1)$$

We also freely use the cup-products, cohomology groups with support, cycle class maps and Gysin morphisms defined by Jannsen [1988].

Note that since  $G = \mathbb{Z}/2\mathbb{Z}$ , the sheaves  $\mathbb{Z}/2^r\mathbb{Z}(j)$  only depend on the parity of  $j$ , hence so do all the cohomology groups considered above.

Let  $\omega$  be the generator of  $H^1(\mathbf{R}, \mathbb{Z}_2(1)) \simeq \mathbb{Z}/2\mathbb{Z}$ . We denote also by  $\omega$  its reduction modulo 2: the generator of  $H^1(\mathbf{R}, \mathbb{Z}/2\mathbb{Z}) \simeq \mathbb{Z}/2\mathbb{Z}$ . If  $k \geq 1$ , their powers  $\omega^k$  generate  $H^k(\mathbf{R}, \mathbb{Z}_2(k)) \simeq \mathbb{Z}/2\mathbb{Z}$  and  $H^k(\mathbf{R}, \mathbb{Z}/2\mathbb{Z}) \simeq \mathbb{Z}/2\mathbb{Z}$ , and we still denote by  $\omega^k$  their pull-backs to any variety  $X$  over  $\mathbf{R}$ .

**1B. Comparison with geometric cohomology.** Let  $\pi : \text{Spec}(\mathbf{C}) \rightarrow \text{Spec}(\mathbf{R})$  be the base-change morphism, and fix  $j \in \mathbb{Z}$ . There is a natural short exact sequence of étale sheaves on  $\text{Spec}(\mathbf{R})$ :  $0 \rightarrow \mathbb{Z}/2^r\mathbb{Z}(j) \rightarrow \pi_*\mathbb{Z}/2^r\mathbb{Z} \rightarrow \mathbb{Z}/2^r\mathbb{Z}(j+1) \rightarrow 0$ , as one checks at the level of  $G$ -modules. They fit together to form a short exact sequence of 2-adic sheaves on  $\text{Spec}(\mathbf{R})$ :  $0 \rightarrow \mathbb{Z}_2(j) \rightarrow \pi_*\mathbb{Z}_2 \rightarrow \mathbb{Z}_2(j+1) \rightarrow 0$ .

Let  $X$  be a variety over  $\mathbf{R}$ , and let us still denote by  $\pi : X_{\mathbf{C}} \rightarrow X$  the base-change morphism. Notice that by the Leray spectral sequence, we have  $H^k(X, \pi_*\mathbb{Z}/2^r\mathbb{Z}) =$

$H^k(X_C, \mathbb{Z}/2^r\mathbb{Z})$ . Now pull-back the exact sequence of 2-adic sheaves  $0 \rightarrow \mathbb{Z}_2(j) \rightarrow \pi_*\mathbb{Z}_2 \rightarrow \mathbb{Z}_2(j+1) \rightarrow 0$  on  $X$  and take continuous étale cohomology. We obtain a long exact sequence

$$\begin{aligned} \cdots \rightarrow H^k(X, \mathbb{Z}_2(j)) &\xrightarrow{\pi^*} H^k(X_C, \mathbb{Z}_2) \\ &\xrightarrow{\pi_*} H^k(X, \mathbb{Z}_2(j+1)) \xrightarrow{\omega} H^{k+1}(X, \mathbb{Z}_2(j)) \rightarrow \cdots \end{aligned} \quad (1-2)$$

in which the boundary map  $H^k(X, \mathbb{Z}_2(j+1)) \rightarrow H^{k+1}(X, \mathbb{Z}_2(j))$  is the cup-product by the class of the extension  $0 \rightarrow \mathbb{Z}_2(j) \rightarrow \pi_*\mathbb{Z}_2 \rightarrow \mathbb{Z}_2(j+1) \rightarrow 0$ , which is the nonzero class  $\omega \in H^1(G, \mathbb{Z}_2(1)) \simeq \mathbb{Z}/2\mathbb{Z}$ .

**1C. Cohomological dimension.** Recall first the following well-known statement, which goes back to Artin.

**Proposition 1.1.** *Let  $X$  be an integral variety over  $\mathbf{R}$ . The following are equivalent:*

- (i)  $\mathbf{R}(X)$  is formally real, that is,  $-1$  is not a sum of squares in  $\mathbf{R}(X)$ .
- (ii)  $X$  has a smooth  $\mathbf{R}$ -point.
- (iii)  $X(\mathbf{R})$  is Zariski-dense in  $X$ .

*Proof.* By the Artin–Lang homomorphism theorem [Bochnak et al. 1998, Theorem 4.1.2], if (i) holds, every open affine subset of  $X$  contains an  $\mathbf{R}$ -point, proving (iii). Conversely, if  $X(\mathbf{R})$  were Zariski-dense in  $X$ ,  $-1$  could not be a sum of squares in  $\mathbf{R}(X)$ , because we would get a contradiction by evaluating this identity at an  $\mathbf{R}$ -point outside of the poles of the rational functions that appear. That (ii) implies (iii) is a consequence of the implicit function theorem [Bochnak et al. 1998, Corollary 2.9.8], and the converse is trivial.  $\square$

From this proposition, it is possible to deduce estimates on the cohomological dimension of varieties  $X$  over  $\mathbf{R}$  without  $\mathbf{R}$ -points. For the cohomological dimension of  $\mathbf{R}(X)$ , this follows from a theorem of Serre [1965] and Artin–Schreier theory. The cohomological dimension of an arbitrary variety  $X$  may then be controlled using [SGA 4<sub>3</sub> 1973, X Corollaire 4.2].

Here, we point out places in the literature where the statements we need are explicitly formulated.

**Proposition 1.2.** *Let  $X$  be an integral variety of dimension  $n$  over  $\mathbf{R}$  such that  $X(\mathbf{R}) = \emptyset$ .*

- (i)  $\mathbf{R}(X)$  has cohomological dimension  $n$ .
- (ii)  $X$  has étale cohomological dimension  $\leq 2n$ .
- (iii) If  $X$  is affine,  $X$  has étale cohomological dimension  $\leq n$ .

*Proof.* The first statement is [Colliot-Thélène and Parimala 1990, Proposition 1.2.1], where it is attributed to Ax.

The second (resp. third) statement follows from [Scheiderer 1994, Corollary 7.21], noticing that the real spectrum of  $X$  is empty by Proposition 1.1 and using that  $X_{\mathbb{C}}$  has étale cohomological dimension  $\leq 2d$  (resp.  $\leq d$ ) by [SGA 4<sub>3</sub> 1973, X Corollaire 4.3] (resp. [SGA 4<sub>3</sub> 1973, XIV Corollaire 3.2]).  $\square$

## 2. Bloch–Ogus theory

**2A. Gersten’s conjecture.** In this subsection, let  $X$  be a smooth variety over  $\mathbf{R}$ .

We want to apply Bloch–Ogus theory to the cohomology groups  $H^k(X, \mathbb{Z}_2(j))$ . For this purpose, one needs to check the validity of Gersten’s conjecture for this cohomology theory. There are two ways to do so.

First, the formal properties of continuous étale cohomology proven by Jannsen [1988] allow one to prove that associating to a variety  $X$  over  $\mathbf{R}$  its continuous étale cohomology groups  $H^k(X, \mathbb{Z}_2(j))$  is part of a Poincaré duality theory with supports in the sense of Bloch and Ogus [1974, Definition 1.3], in the same way as it is proven for étale cohomology with finite coefficients [Bloch and Ogus 1974, §2]. Then it is possible to apply [Bloch and Ogus 1974, Theorem 4.2].

Another possibility is to use the axioms of [Colliot-Thélène et al. 1997], which are easier to check. That these axioms hold for continuous étale cohomology is explained for instance in [Kahn 2012, §3C], allowing us to apply [Colliot-Thélène et al. 1997, Corollary 5.1.11].

Let us now explain the meaning of Gersten’s conjecture in our context. We define  $\mathcal{H}_X^k(j)$  to be the Zariski sheaf on  $X$  that is the sheafification of  $U \mapsto H^k(U, \mathbb{Z}_2(j))$ . Moreover, if  $z \in X$  is a point with closure  $Z \subset X$ , we define<sup>4</sup>

$$H_{\rightarrow}^k(z, \mathbb{Z}_2(j)) := \varinjlim_{U \subset Z} H^k(U, \mathbb{Z}_2(j)), \quad (2-1)$$

where  $U$  runs over all nonempty open subsets of  $Z$ . We define  $\iota_z : z \rightarrow X$  to be the inclusion, and we consider the skyscraper sheaves  $\iota_{z*} H_{\rightarrow}^k(z, \mathbb{Z}_2(j))$  on  $X$ . Finally, we set  $X^{(c)}$  to be the set of codimension  $c$  points in  $X$ . Then the sheaves  $\mathcal{H}_X^k(j)$  admit Cousin resolutions (see either [Bloch and Ogus 1974, (4.2.2)] or [Colliot-Thélène et al. 1997, Corollary 5.1.11] taking into account purity [Jannsen 1988, (3.21)] to obtain the precise form below):

$$\begin{aligned} 0 \rightarrow \mathcal{H}_X^k(j) \rightarrow \bigoplus_{z \in X^{(0)}} \iota_{z*} H_{\rightarrow}^k(z, \mathbb{Z}_2(j)) \rightarrow \bigoplus_{z \in X^{(1)}} \iota_{z*} H_{\rightarrow}^{k-1}(z, \mathbb{Z}_2(j-1)) \\ \rightarrow \dots \rightarrow \bigoplus_{z \in X^{(k)}} \iota_{z*} H_{\rightarrow}^0(z, \mathbb{Z}_2(j-k)) \rightarrow 0. \end{aligned} \quad (2-2)$$

<sup>4</sup>Beware that since continuous étale cohomology does not commute with inverse limit of schemes, this group does not coincide in general with the continuous Galois cohomology of the residue field of  $z$ .



The way this Cousin resolution is constructed, from a coniveau spectral sequence, shows that the arrows in (2-2) are given by maps in long exact sequences of cohomology with support, also called residue maps.

Since the sheaves in this resolution are flasque, the Cousin complex obtained by taking its global sections computes the Zariski cohomology of  $\mathcal{H}_X^k(j)$ . For instance, this implies that  $H^0(X, \mathcal{H}_X^k(j))$  coincides with the unramified cohomology group  $H_{\text{nr}}^k(X, \mathbb{Z}_2(j))$ , that is, the subgroup of  $H_{\rightarrow}^k(\eta, \mathbb{Z}_2(j))$  on which all residues at codimension 1 points of  $X$  vanish.

The exactness of (2-2) allows us to compute the second page of the coniveau spectral sequence for  $X$  mentioned above. As shown in [Bloch and Ogus 1974, Corollary 6.3] or [Colliot-Thélène et al. 1997, Corollary 5.1.11], it reads

$$E_2^{p,q} = H^p(X, \mathcal{H}_X^q(j)) \Rightarrow H^{p+q}(X, \mathbb{Z}_2(j)). \quad (2-3)$$

Recall that the filtration induced by this spectral sequence on  $H^k(X, \mathbb{Z}_2(j))$  is the coniveau filtration, where a class  $\alpha \in H^k(X, \mathbb{Z}_2(j))$  has coniveau  $\geq c$  if it vanishes in the complement of a closed subset of codimension  $c$  of  $X$ .

**2B. Bloch–Ogus theory over  $\mathbf{R}$ .** If  $X$  is a variety over  $\mathbf{R}$ , we still denote by  $\pi : X_{\mathbf{C}} \rightarrow X$  the natural morphism, and we naturally view  $X_{\mathbf{C}}$  as a variety over  $\mathbf{R}$ . The following proposition was proved in [Colliot-Thélène and Scheiderer 1996, Lemma 2.2.1] over  $\mathbb{R}$  and with 2-torsion coefficients, but the proof goes through, and we include it for completeness.

**Proposition 2.1.** *Let  $X$  be a smooth variety over  $\mathbf{R}$  and fix  $j \in \mathbb{Z}$ . Then there exists a long exact sequence of Zariski sheaves on  $X$ :*

$$\cdots \rightarrow \mathcal{H}_X^k(j) \rightarrow \pi_* \mathcal{H}_{X_{\mathbf{C}}}^k \rightarrow \mathcal{H}_X^k(j+1) \rightarrow \mathcal{H}_X^{k+1}(j) \rightarrow \cdots. \quad (2-4)$$

Moreover, the sheaf  $\pi_* \mathcal{H}_{X_{\mathbf{C}}}^k$  coincides with the sheafification of  $U \mapsto H^k(U_{\mathbf{C}}, \mathbb{Z}_2)$  and its cohomology groups are  $H^q(X, \pi_* \mathcal{H}_{X_{\mathbf{C}}}^k) = H^q(X_{\mathbf{C}}, \mathcal{H}_{X_{\mathbf{C}}}^k)$  for any  $k, q \geq 0$ .

*Proof.* Let  $x \in X$ . If  $V$  is a neighborhood of  $\pi^{-1}(x)$  in  $X_{\mathbf{C}}$ , the sheaf  $\mathcal{H}_V^k$  has a flasque Cousin resolution (2-2). Taking global sections and taking the limit over all such neighborhoods  $V$  gives a complex that is exact in positive degree (the argument for étale cohomology with finite coefficients is [Colliot-Thélène et al. 1997, Proposition 2.1.2], and the corresponding effaceability condition for continuous étale cohomology follows from [Colliot-Thélène et al. 1997, Theorem 5.1.10]). As a consequence,

$$\varinjlim_V H^p(V, \mathcal{H}_V^k) = 0 \quad \text{for } p > 0. \quad (2-5)$$

Considering the coniveau spectral sequences (2-3) for every  $V$  and taking (2-5) into account shows that

$$\varinjlim_V H^k(V, \mathbb{Z}_2) = \varinjlim_V H^0(V, \mathcal{H}_V^k). \quad (2-6)$$

Note that in both (2-5) and (2-6), it is possible to restrict to neighborhoods of the form  $U_C$  for  $U \subset X$  because they form a cofinal family.

Now, the exact sequences obtained by applying (1-2) to all open subsets of  $X$  fit together to induce a long exact sequence of Zariski presheaves on  $X$ . By exactness of sheafification, one obtains a long exact sequence of Zariski sheaves on  $X$ :

$$\cdots \rightarrow \mathcal{H}_X^k(j) \rightarrow \mathcal{F}^k \rightarrow \mathcal{H}_X^k(j+1) \rightarrow \mathcal{H}_X^{k+1}(j) \rightarrow \cdots,$$

where  $\mathcal{F}^k$  is the sheafification of  $U \mapsto H^k(U_C, \mathbb{Z}_2)$ . The universal property of sheafification gives a morphism  $\mathcal{F}^k \rightarrow \pi_* \mathcal{H}_{X_C}^k$ . The map induced on stalks at  $x \in X$  is precisely (2-6), hence an isomorphism. It follows that  $\mathcal{F}^k \simeq \pi_* \mathcal{H}_{X_C}^k$ , completing the construction of (2-4).

If  $k \geq 0$  and  $p > 0$ , the stalk of  $R^p \pi_* \mathcal{H}_{X_C}^k = 0$  at  $x \in X$  is given by (2-5), hence trivial. It follows that  $R^p \pi_* \mathcal{H}_{X_C}^k$  vanishes, and the Leray spectral sequence for  $\pi$  implies the last statement of the proposition.  $\square$

**2C. Consequences of the Bloch–Kato conjecture.** The following proposition is due to Bloch and Srinivas [1983, proof of Theorem 1] for  $k \leq 2$  and to Colliot-Thélène and Voisin [2012, Théorème 3.1] in general. Since both references work over an algebraically closed field, and since the latter uses Betti cohomology, we repeat the proof to emphasize that it works in our setting.

**Proposition 2.2.** *Let  $X$  be a smooth variety over  $\mathbf{R}$ . Then for every  $k \geq 0$ , the sheaf  $\mathcal{H}_X^{k+1}(k)$  is torsion free.*

*Proof.* Since it is a sheaf of  $\mathbb{Z}_2$ -modules, it suffices to prove that it has no 2-torsion. Consider the exact sequence of 2-adic sheaves on  $X$ :  $0 \rightarrow \mathbb{Z}_2(k) \xrightarrow{2} \mathbb{Z}_2(k) \rightarrow \mu_2^{\otimes k} \rightarrow 0$ . Taking long exact sequences of continuous cohomology over every open subset  $U \subset X$  to get a long exact sequence of presheaves on  $X$  and sheafifying it gives a long exact sequence of sheaves on  $X$ , part of which is

$$\mathcal{H}_X^k(k) \rightarrow \mathcal{H}_X^k(\mu_2^{\otimes k}) \rightarrow \mathcal{H}_X^{k+1}(k) \xrightarrow{2} \mathcal{H}_X^{k+1}(k),$$

where  $\mathcal{H}_X^k(\mu_2^{\otimes k})$  is the sheafification of  $U \mapsto H^k(U, \mu_2^{\otimes k})$ . Consequently, it suffices to prove the surjectivity of  $\mathcal{H}_X^k(k) \rightarrow \mathcal{H}_X^k(\mu_2^{\otimes k})$ .

On an open set  $U \subset X$ , the Kummer exact sequence  $0 \rightarrow \mu_2 \rightarrow \mathbb{G}_m \xrightarrow{2} \mathbb{G}_m \rightarrow 0$  induces a boundary map  $H^0(U, \mathcal{O}_U^*) \rightarrow H^1(U, \mu_2)$ . These maps sheafify to  $\mathcal{O}_X^* \rightarrow \mathcal{H}_X^1(\mu_2)$ , inducing via cup-products a morphism of sheaves  $(\mathcal{O}_X^*)^{\otimes k} \rightarrow \mathcal{H}_X^k(\mu_2^{\otimes k})$ . It is explained in [Colliot-Thélène and Voisin 2012, end of section 2.2] how Gersten’s conjecture for Milnor K-theory proven by Kerz [2009] and the Bloch–Kato conjecture proven by Rost and Voevodsky (since we only need this conjecture at the prime 2, Voevodsky’s work [2003, Corollary 7.4] on Milnor’s conjecture is sufficient here) imply the surjectivity of this morphism.

Over an open set  $U \subset X$ , the boundary maps  $H^0(U, \mathcal{O}_U^*) \rightarrow H^1(U, \mu_{2^r})$  for the Kummer exact sequences

$$0 \rightarrow \mu_{2^r} \rightarrow \mathbb{G}_m \xrightarrow{2^r} \mathbb{G}_m \rightarrow 0$$

fit together to induce a map  $H^0(U, \mathcal{O}_U^*) \rightarrow \varprojlim_r H^1(U, \mu_{2^r}) = H^1(U, \mathbb{Z}_2(1))$ . Again, this sheafifies to a morphism  $\mathcal{O}_X^* \rightarrow \mathcal{H}_X^1(1)$ , inducing via cup-products a morphism of sheaves  $(\mathcal{O}_X^*)^{\otimes k} \rightarrow \mathcal{H}_X^k(k)$  lifting  $(\mathcal{O}_X^*)^{\otimes k} \rightarrow \mathcal{H}_X^k(\mu_2^{\otimes k})$ . The surjectivity of  $\mathcal{H}_X^k(k) \rightarrow \mathcal{H}_X^k(\mu_2^{\otimes k})$  now follows from surjectivity of  $(\mathcal{O}_X^*)^{\otimes k} \rightarrow \mathcal{H}_X^k(\mu_2^{\otimes k})$ .  $\square$

In [Bloch and Srinivas 1983; Colliot-Thélène and Voisin 2012], the authors worked over an algebraically closed field, and the Tate twist was not essential for the result to hold. Here, it is very important; it is not true in general that the sheaf  $\mathcal{H}_X^k(k)$  has no torsion.

As in these references, the following are straightforward corollaries.

**Corollary 2.3.** *Let  $X$  be a smooth variety over  $\mathbf{R}$  and  $k \geq 0$ . Then*

$$H_{\text{nr}}^{k+1}(X, \mathbb{Z}_2(k)) = H^0(X, \mathcal{H}_X^{k+1}(k))$$

*is torsion free.*

**Corollary 2.4.** *Let  $X$  be an integral variety over  $\mathbf{R}$  with generic point  $\eta$  and  $k \geq 0$ . Then  $H_{\rightarrow}^{k+1}(\eta, \mathbb{Z}_2(k))$  is torsion free.*

*Proof.* If  $\alpha \in H_{\text{nr}}^{k+1}(U, \mathbb{Z}_2(k))$  is a torsion class on a smooth open subset  $U \subset X$ , it vanishes in  $H_{\text{nr}}^{k+1}(U, \mathbb{Z}_2(k))$  by Corollary 2.3, hence on an open subset  $V \subset U$ .  $\square$

Another application of Proposition 2.2 is as follows.

**Proposition 2.5.** *Let  $X$  be a smooth variety over  $\mathbf{R}$ . Then for every  $k \geq 0$ , there is an exact sequence*

$$0 \rightarrow \mathcal{H}_X^{k-1}(k) \rightarrow \pi_* \mathcal{H}_{X_C}^{k-1} \rightarrow \mathcal{H}_X^{k-1}(k+1) \rightarrow \mathcal{H}_X^k(k) \rightarrow \pi_* \mathcal{H}_{X_C}^k \rightarrow \mathcal{H}_X^k(k+1) \rightarrow 0.$$

*Proof.* Let us prove that the long exact sequence (2-4) splits into these shorter exact sequences. It suffices to prove that, for  $k \geq 0$ , the morphism  $\mathcal{H}_X^{k-1}(k) \rightarrow \pi_* \mathcal{H}_{X_C}^{k-1}$  is injective. The composition

$$\mathcal{H}_X^{k-1}(k) \rightarrow \pi_* \mathcal{H}_{X_C}^{k-1} \rightarrow \mathcal{H}_X^{k-1}(k)$$

is multiplication by 2. Consequently, the kernel of  $\mathcal{H}_X^{k-1}(k) \rightarrow \pi_* \mathcal{H}_{X_C}^{k-1}$  is of 2-torsion. Since  $\mathcal{H}_X^{k-1}(k)$  is torsion free by Proposition 2.2, this kernel is trivial, as required.  $\square$

**Proposition 2.6.** *Let  $X$  be an integral variety over  $\mathbf{R}$  with generic point  $\eta$ . Then for every  $k \geq 0$ , there is an exact sequence*

$$\begin{aligned} 0 \rightarrow H_{\rightarrow}^{k-1}(\eta, \mathbb{Z}_2(k)) &\rightarrow H_{\rightarrow}^{k-1}(\eta, \pi_* \mathbb{Z}_2) \rightarrow H_{\rightarrow}^{k-1}(\eta, \mathbb{Z}_2(k+1)) \\ &\rightarrow H_{\rightarrow}^k(\eta, \mathbb{Z}_2(k)) \rightarrow H_{\rightarrow}^k(\eta, \pi_* \mathbb{Z}_2) \rightarrow H_{\rightarrow}^k(\eta, \mathbb{Z}_2(k+1)) \rightarrow 0. \end{aligned}$$

*Proof.* Take the direct limit of the long exact sequence (1-2) applied to all open subsets of  $X$ ; it splits into exact sequences of length six by the same argument as in the proof of Proposition 2.5, using Corollary 2.4 instead of Corollary 2.3.  $\square$

### 3. Sums of squares and unramified cohomology

**3A. Sums of squares and level.** Let  $n \geq 1$ , consider a nonzero positive semidefinite polynomial  $f \in \mathbf{R}[X_1, \dots, X_n]$  and its homogenization  $F \in \mathbf{R}[X_0, \dots, X_n]$ . Notice that since an odd degree polynomial over  $\mathbf{R}$  changes sign,  $f$  and  $F$  must have even degree. This allows us to consider the double cover  $Y$  of  $\mathbb{P}_{\mathbf{R}}^n$  ramified over  $\{F = 0\}$  defined by the equation  $Y := \{Z^2 + F = 0\}$  in the weighted projective space  $\mathbb{P}(1, \dots, 1, \deg(F)/2)$ .

**Lemma 3.1.** *The variety  $Y$  is integral,  $\mathbf{R}(Y)$  is not formally real, and if  $\tilde{Y} \rightarrow Y$  is a resolution of singularities, then  $\tilde{Y}(\mathbf{R}) = \emptyset$ .*

*Proof.* To prove that  $Y$  is integral, one has to check that  $-f$  is not a square in  $\mathbf{R}(X_1, \dots, X_n)$ , or equivalently, that it is not a square in  $\mathbf{R}[X_1, \dots, X_n]$ . But if it were,  $f$  would be negative on  $\mathbf{R}^n$ , hence zero on  $\mathbf{R}^n$  by positivity, hence zero by Zariski-density of  $\mathbf{R}^n$  in  $\mathbf{C}^n$ . This is a contradiction.

The  $\mathbf{R}$ -points of  $\tilde{Y}$  necessarily lie above  $\mathbf{R}$ -points of  $Y$ , hence, by positivity of  $F$ , above zeroes of  $F$ . Consequently,  $\tilde{Y}(\mathbf{R})$  is not Zariski-dense in  $\tilde{Y}$ . Applying Proposition 1.1 using the smoothness of  $\tilde{Y}$  shows that  $\tilde{Y}(\mathbf{R}) = \emptyset$ , and that  $\mathbf{R}(Y)$  is not formally real.  $\square$

Recall that the level  $s(K) \in \mathbb{N}^* \cup \{\infty\}$  of a field  $K$  is  $\infty$  if  $-1$  is not a sum of squares in  $K$ , and the smallest  $s$  such that  $-1$  is a sum of  $s$  squares otherwise. In the latter case, it has been shown by Pfister [1965, Satz 4] to be a power of 2.

**Proposition 3.2.** *The polynomial  $f$  is a sum of  $2^n - 1$  squares in  $\mathbf{R}(X_1, \dots, X_n)$  if and only if  $\mathbf{R}(Y)$  has level  $< 2^n$ . Conversely, the polynomial  $f$  is not a sum of  $2^n - 1$  squares in  $\mathbf{R}(X_1, \dots, X_n)$  if and only if  $\mathbf{R}(Y)$  has level  $2^n$ .*

*Proof.* Proposition 1.1 shows that  $\mathbf{R}(X_1, \dots, X_n)$  is formally real and Artin's solution [1927] to Hilbert's 17th problem shows that  $f$  is a sum of squares in  $\mathbf{R}(X_1, \dots, X_n)$ .

Then [Lam 1980, Chapter 11, Theorem 2.7] applies and shows that  $f$  is a sum of  $2^n - 1$  squares in  $\mathbf{R}(X_1, \dots, X_n)$  if and only if  $\mathbf{R}(Y)$  has level  $< 2^n$  (this is essentially due to Pfister; the statement we have used is very close and its proof is identical to [Pfister 1965, Satz 5]).

Since  $\mathbf{R}(Y)$  is not formally real by Lemma 3.1, Pfister [1967, Theorem 2] has shown that its level is  $\leq 2^n$ . This concludes the proof.  $\square$

**3B. Level and unramified cohomology.** To apply Proposition 3.2, we need to control the level of the function field of a variety over  $\mathbf{R}$ . The following proposition relates it to one of its unramified cohomology groups. The equivalence (i)  $\Leftrightarrow$  (ii) is hinted at in [Colliot-Thélène 1993, bottom of p. 236], at least for  $n = 3$ . I am grateful to Olivier Wittenberg for explaining to me that the implication (ii)  $\Rightarrow$  (iii) holds.

**Proposition 3.3.** *Let  $X$  be a smooth integral variety over  $\mathbf{R}$ , and fix  $n \geq 1$ . The following assertions are equivalent:*

- (i) *The function field  $\mathbf{R}(X)$  has level  $< 2^n$ .*
- (ii) *The map  $H^n(\mathbf{R}, \mathbb{Z}/2\mathbb{Z}) \rightarrow H^n(\mathbf{R}(X), \mathbb{Z}/2\mathbb{Z})$  vanishes.*
- (iii) *The map  $H^n(\mathbf{R}, \mathbb{Z}_2(n)) \rightarrow H_{\text{nr}}^n(X, \mathbb{Z}_2(n))$  vanishes.*

*Proof.* Consider the property that the level of  $\mathbf{R}(X)$  is  $< 2^n$ . It is equivalent to the fact that  $-1$  is a sum of  $2^n - 1$  squares in  $\mathbf{R}(X)$ , hence to the fact that the Pfister quadratic form  $q := \langle 1, 1 \rangle^{\otimes n}$  is isotropic over  $\mathbf{R}(X)$ . By a theorem of Elman and Lam [1972, Corollary 3.3], this is equivalent to the vanishing of the symbol  $\{-1\}^n$  in the Milnor K-theory group  $K_n^M(\mathbf{R}(X))/2$ . By Voevodsky's proof [2003, Corollary 7.4] of the Milnor conjecture, the natural map

$$K_n^M(\mathbf{R}(X))/2 \rightarrow H^n(\mathbf{R}(X), \mathbb{Z}/2\mathbb{Z})$$

is an isomorphism, so our property is equivalent to the vanishing of  $H^n(\mathbf{R}, \mathbb{Z}/2\mathbb{Z}) \rightarrow H^n(\mathbf{R}(X), \mathbb{Z}/2\mathbb{Z})$ . We have proven that (i) and (ii) are equivalent.

Suppose that (iii) holds and let  $\eta$  be the generic point of  $X$ . The definition of  $H_{\text{nr}}^n(X, \mathbb{Z}_2(n))$  as a subgroup of  $H_{\rightarrow}^n(\eta, \mathbb{Z}_2(n))$  shows that  $H^n(\mathbf{R}, \mathbb{Z}_2(n)) \rightarrow H_{\rightarrow}^n(\eta, \mathbb{Z}_2(n))$  vanishes. Then we have a commutative diagram

$$\begin{array}{ccc} H^n(\mathbf{R}, \mathbb{Z}_2(n)) & \longrightarrow & H_{\rightarrow}^n(\eta, \mathbb{Z}_2(n)) \\ \downarrow \cong & & \downarrow \\ H^n(\mathbf{R}, \mathbb{Z}/2\mathbb{Z}) & \longrightarrow & H_{\rightarrow}^n(\eta, \mathbb{Z}/2\mathbb{Z}) \end{array}$$

where the groups on the right are defined as inductive limits on the open subsets of  $X$  as in (2-1), showing that  $H^n(\mathbf{R}, \mathbb{Z}/2\mathbb{Z}) \rightarrow H_{\rightarrow}^n(\eta, \mathbb{Z}/2\mathbb{Z})$  vanishes. Since étale cohomology commutes with such limits [SGA 4<sub>2</sub> 1972, VII Corollaire 5.8],  $H_{\rightarrow}^n(\eta, \mathbb{Z}/2\mathbb{Z})$  is nothing but the Galois cohomology group  $H^n(\mathbf{R}(X), \mathbb{Z}/2\mathbb{Z})$ , proving (ii).

Suppose conversely that (ii) holds, and let  $U \subset X$  be an open subset such that  $\omega^n$  vanishes in  $H^n(U, \mathbb{Z}/2\mathbb{Z})$ . Consider the following commutative exact diagram, where the lines are (1-2):

$$\begin{array}{ccccc}
H^{n-1}(U, \mathbb{Z}_2(n-1)) & \xrightarrow{\omega} & H^n(U, \mathbb{Z}_2(n)) & \longrightarrow & H^n(U_C, \mathbb{Z}_2) \\
\downarrow 2 & & \downarrow 2 & & \downarrow 2 \\
H^{n-1}(U, \mathbb{Z}_2(n-1)) & \xrightarrow{\omega} & H^n(U, \mathbb{Z}_2(n)) & \longrightarrow & H^n(U_C, \mathbb{Z}_2) \\
& & \downarrow & & \\
& & H^n(U, \mathbb{Z}/2\mathbb{Z}) & & 
\end{array}$$

Look at  $\omega^n \in H^n(U, \mathbb{Z}_2(n))$ . By hypothesis, it vanishes in  $H^n(U, \mathbb{Z}/2\mathbb{Z})$ , hence may be written  $2\alpha$  for some  $\alpha \in H^n(U, \mathbb{Z}_2(n))$ . Since  $\omega^n \in H^n(U, \mathbb{Z}_2(n))$  is the image of  $\omega^{n-1} \in H^{n-1}(U, \mathbb{Z}_2(n-1))$ ,  $\alpha_C \in H^n(U_C, \mathbb{Z}_2)$  is a 2-torsion class. By Corollary 2.4, any torsion class in  $H^n(U_C, \mathbb{Z}_2)$  vanishes on an open subset: up to shrinking  $U$ , we may assume that  $\alpha_C = 0$ , hence that there is  $\beta \in H^{n-1}(U, \mathbb{Z}_2(n-1))$  such that  $\beta \cdot \omega = \alpha$ . Then  $\omega^n = \beta \cdot 2\omega = 0 \in H^n(U, \mathbb{Z}_2(n))$ , proving (iii).  $\square$

**3C. From degree  $n$  to degree  $n+1$  cohomology.** Condition (iii) in Proposition 3.3 means that  $\omega^n$  has coniveau  $\geq 1$ . Proposition 3.5 uses Bloch–Ogus theory to relate this property to the coniveau of  $\omega^{n+1}$ .

Fix an integer  $n \geq 1$  and let  $X$  be a smooth variety over  $\mathbf{R}$ . The coniveau spectral sequence (2-3) induces two maps  $H^n(X, \mathbb{Z}_2(n)) \xrightarrow{\phi} H_{\text{nr}}^n(X, \mathbb{Z}_2(n))$  and  $K := \text{Ker}[H^{n+1}(X, \mathbb{Z}_2(n+1)) \rightarrow H_{\text{nr}}^{n+1}(X, \mathbb{Z}_2(n+1))] \xrightarrow{\psi} H^1(X, \mathcal{H}_X^n(n+1))$ .

Cup-product with  $\omega$  gives morphisms  $H^n(X, \mathbb{Z}_2(n)) \xrightarrow{\omega} H^{n+1}(X, \mathbb{Z}_2(n+1))$  and  $H_{\text{nr}}^n(X, \mathbb{Z}_2(n)) \xrightarrow{\omega} H_{\text{nr}}^{n+1}(X, \mathbb{Z}_2(n+1))$ . Let  $I := \{\alpha \in H^n(X, \mathbb{Z}_2(n)) \mid \alpha \cdot \omega \in K\}$  and  $I_{\text{nr}} := \{\alpha \in H_{\text{nr}}^n(X, \mathbb{Z}_2(n)) \mid \alpha \cdot \omega = 0\}$ .

Finally, Proposition 2.5 gives an exact sequence of sheaves on  $X$ :

$$0 \rightarrow \mathcal{H}_X^n(n+1) \rightarrow \pi_* \mathcal{H}_{X_C}^n \rightarrow \mathcal{H}_X^n(n) \xrightarrow{\omega} \mathcal{H}_X^{n+1}(n+1) \rightarrow \cdots \quad (3-1)$$

Taking cohomology, we obtain an exact sequence:

$$0 \rightarrow H_{\text{nr}}^n(X, \mathbb{Z}_2(n+1)) \rightarrow H_{\text{nr}}^n(X_C, \mathbb{Z}_2) \rightarrow I_{\text{nr}} \xrightarrow{\delta} H^1(X, \mathcal{H}_X^n(n+1)). \quad (3-2)$$

**Lemma 3.4.** *Let  $X$  be a smooth variety over  $\mathbf{R}$ . The diagram*

$$\begin{array}{ccc}
I & \xrightarrow{\omega} & K \\
\phi \downarrow & & \downarrow \psi \\
I_{\text{nr}} & \xrightarrow{\delta} & H^1(X, \mathcal{H}_X^n(n+1))
\end{array}$$

*constructed above commutes.*

*Proof.* Let  $\alpha \in I$ . By hypothesis, the class  $\alpha \cdot \omega \in H^{n+1}(X, \mathbb{Z}_2(n+1))$  vanishes on an open subset  $U \subset X$ . Let  $D := X \setminus U$  be endowed with its reduced structure.

The description of  $H^1(X, \mathcal{H}_X^n(n+1))$  as a cohomology group of the Cousin complex (2-2) shows that if  $X^\circ \subset X$  is an open subset whose complement has

codimension  $\geq 2$ , then the restriction  $H^1(X, \mathcal{H}_X^n(n+1)) \rightarrow H^1(X^\circ, \mathcal{H}_{X^\circ}^n(n+1))$  is injective. Consequently, to prove that  $\psi(\alpha \cdot \omega) = \delta \circ \phi(\alpha)$ , it is possible to remove from  $X$  a closed subset of codimension  $\geq 2$ . This allows us to suppose that  $D$  is smooth of pure codimension 1.

Our next task is to concretely identify  $\delta \circ \phi(\alpha)$ . The cohomology theory with supports in the sense of [Colliot-Thélène et al. 1997, Definition 5.1.1], which to a variety  $X$  over  $\mathbf{R}$  and a closed subset  $Z \subset X$  associates the groups  $H_Z^k(X, \pi_* \mathbb{Z}_2) = H_{Z_C}^k(X_C, \mathbb{Z}_2)$ , satisfies axioms COH1 and COH3 by [Colliot-Thélène et al. 1997, 5.5(1)], hence COH2 by [Colliot-Thélène et al. 1997, Proposition 5.3.2]. It follows from [Colliot-Thélène et al. 1997, Corollary 5.1.11] that the sheafification of  $U \mapsto H^n(U_C, \mathbb{Z}_2)$  (that is,  $\pi_* \mathcal{H}_{X_C}^n$  by Proposition 2.1) admits a Cousin resolution by flasque sheaves, and the same goes for  $\pi_* \mathcal{H}_{X_C}^{n+1}$ . These resolutions fit together with the Cousin resolutions (2-2) of  $\mathcal{H}_X^n(n+1)$ ,  $\mathcal{H}_X^n(n)$ ,  $\mathcal{H}_X^{n+1}(n+1)$  and  $\mathcal{H}_X^{n+1}(n)$ , giving rise to a diagram that is an exact sequence of flasque resolutions for the exact sequence of sheaves (3-1) by Proposition 2.6. Let us only draw the relevant part of the diagram containing the Cousin resolutions for  $\mathcal{H}_X^n(n+1)$ ,  $\pi_* \mathcal{H}_{X_C}^{n+1}$  and  $\mathcal{H}_X^n(n)$ :

$$\begin{array}{ccccc}
 \bigoplus_{z \in X^{(0)}} \iota_{z*} H_{\rightarrow}^n(z, \mathbb{Z}_2(n+1)) & \longrightarrow & \bigoplus_{z \in X^{(1)}} \iota_{z*} H_{\rightarrow}^{n-1}(z, \mathbb{Z}_2(n)) & \longrightarrow & \cdots \\
 \downarrow & & \downarrow & & \\
 \bigoplus_{z \in X^{(0)}} \iota_{z*} H_{\rightarrow}^n(z, \pi_* \mathbb{Z}_2) & \longrightarrow & \bigoplus_{z \in X^{(1)}} \iota_{z*} H_{\rightarrow}^{n-1}(z, \pi_* \mathbb{Z}_2) & \longrightarrow & \cdots \\
 \downarrow & & \downarrow & & \\
 \bigoplus_{z \in X^{(0)}} \iota_{z*} H_{\rightarrow}^n(z, \mathbb{Z}_2(n)) & \longrightarrow & \bigoplus_{z \in X^{(1)}} \iota_{z*} H_{\rightarrow}^{n-1}(z, \mathbb{Z}_2(n-1)) & \longrightarrow & \cdots
 \end{array} \quad (3-3)$$

It is now possible to give a description of  $\delta \circ \phi(\alpha)$  by a diagram chase in the diagram obtained by taking the global sections of (3-3). More precisely,  $\alpha$  induces a class

$$\phi(\alpha) \in \text{Ker} \left[ \bigoplus_{z \in X^{(0)}} H_{\rightarrow}^n(z, \mathbb{Z}_2(n)) \rightarrow \bigoplus_{z \in X^{(1)}} H_{\rightarrow}^{n-1}(z, \mathbb{Z}_2(n-1)) \right].$$

Lifting it in  $\bigoplus_{z \in X^{(0)}} H_{\rightarrow}^n(z, \pi_* \mathbb{Z}_2)$  by the hypothesis that  $\alpha \in I$ , pushing it to  $\bigoplus_{z \in X^{(1)}} H_{\rightarrow}^{n-1}(z, \pi_* \mathbb{Z}_2)$  and lifting it again to  $\bigoplus_{z \in X^{(1)}} H_{\rightarrow}^{n-1}(z, \mathbb{Z}_2(n))$  gives a cohomology class of degree one of the complex of global sections of the Cousin resolution of  $\mathcal{H}_X^n(n+1)$  representing  $\delta \circ \phi(\alpha) \in H^1(X, \mathcal{H}_X^n(n+1))$ .

At this point, consider the following commutative diagram, whose rows are exact sequences of cohomology with support, whose columns are instances of (1-2), and

where the coefficient ring  $\mathbb{Z}_2$  has been omitted:

$$\begin{array}{ccccccc}
 & & & & H^n(U, n+1) & \longrightarrow & H^{n-1}(D, n) \\
 & & & & \downarrow & & \downarrow \pi^* \\
 & & & & H^n(U_C) & \xrightarrow{\partial} & H^{n-1}(D_C) \\
 & & & & \downarrow \pi_* & & \downarrow \\
 H^{n-2}(D, n-1) & \longrightarrow & H^n(X, n) & \xrightarrow{j^*} & H^n(U, n) & \longrightarrow & H^{n-1}(D, n-1) \\
 \downarrow & & \downarrow \omega & & \downarrow & & \\
 H^{n-1}(D, n) & \xrightarrow{i_*} & H^{n+1}(X, n+1) & \twoheadrightarrow & H^{n+1}(U, n+1) & & \\
 \downarrow & & & & & & \\
 H^{n-1}(D_C) & & & & & & 
 \end{array}$$

Here we have denoted by  $i : D \rightarrow X$  and  $j : U \rightarrow X$  the inclusions, and by  $\partial$  the residue map. By our choice of  $U$ ,  $\alpha \in H^n(X, \mathbb{Z}_2(n))$  vanishes in  $H^{n+1}(U, \mathbb{Z}_2(n+1))$ . Chasing the diagram, there are two ways to construct a (not well-defined) class in  $H^{n-1}(D, \mathbb{Z}_2(n))$ . First, we may consider a class  $\beta \in H^{n-1}(D, \mathbb{Z}_2(n))$  such that  $i_*\beta = \alpha \cdot \omega$ . Second, we may lift  $j^*\alpha$  along  $\pi_*$ , apply the residue map  $\partial$ , and lift the resulting class along  $\pi^*$  to obtain  $\gamma \in H^{n-1}(D, \mathbb{Z}_2(n))$ .

Our diagram has been constructed from the diagram of distinguished triangles in the derived category of 2-adic sheaves on  $X$ :

$$\begin{array}{ccccccc}
 i_* Ri^! \mathbb{Z}_2(n+1) & \longrightarrow & \mathbb{Z}_2(n+1) & \longrightarrow & Rj_* j^* \mathbb{Z}_2(n+1) & \longrightarrow & \\
 \downarrow & & \downarrow & & \downarrow & & \\
 i_* Ri^! \pi_* \mathbb{Z}_2 & \longrightarrow & \pi_* \mathbb{Z}_2 & \longrightarrow & Rj_* j^* \pi_* \mathbb{Z}_2 & \longrightarrow & \\
 \downarrow & & \downarrow & & \downarrow & & \\
 i_* Ri^! \mathbb{Z}_2(n) & \longrightarrow & \mathbb{Z}_2(n) & \longrightarrow & Rj_* j^* \mathbb{Z}_2(n) & \longrightarrow & \\
 \downarrow & & \downarrow & & \downarrow & & 
 \end{array}$$

A homological algebra lemma due to Jannsen [2000, lemma, p. 268], applied exactly as in [Jannsen 2000, proof of Theorem 2], shows that the images of  $\beta$  and  $\gamma$  in  $H^{n-1}(D_C, \mathbb{Z}_2)$  that are well-defined up to the image of  $H^n(U, \mathbb{Z}_2(n+1))$  coincide up to a sign. It follows that  $\beta$  and  $\gamma$ , well-defined up to the images of  $H^n(U, \mathbb{Z}_2(n+1))$  and  $H^{n-2}(D, \mathbb{Z}_2(n-1))$  in  $H^{n-1}(D, \mathbb{Z}_2(n))$ , coincide up to a sign.



Now notice that  $\beta$  and  $\gamma$  induce classes in  $\bigoplus_{z \in X^{(1)}} H_{\rightarrow}^{n-1}(z, \mathbb{Z}_2(n))$ . Our explicit description of  $\delta \circ \phi(\alpha)$  shows that  $\beta$  is a representative of it as a cohomology class of degree one of the Cousin complex. On the other hand,  $\gamma$  has been constructed by lifting  $\alpha \cdot \omega$  along the Gysin morphism  $H^{n-1}(D, \mathbb{Z}_2(n)) \rightarrow H^{n+1}(X, \mathbb{Z}_2(n+1))$ . By construction of the coniveau spectral sequence [Bloch and Ogus 1974, §3; Colliot-Thélène et al. 1997, §1],  $\gamma$  is a representative of  $\psi(\alpha \cdot \omega)$  as a cohomology class of degree one of the Cousin complex.

At this point, we have proven that  $\psi(\alpha \cdot \omega) = [\gamma] = -[\beta] = -\delta \circ \phi(\alpha)$ . Since this element is 2-torsion because  $\omega$  is, one has in fact  $\psi(\alpha \cdot \omega) = \delta \circ \phi(\alpha)$ , as wanted.  $\square$

**Proposition 3.5.** *Let  $X$  be a smooth projective variety over  $\mathbf{R}$ , and fix  $n \geq 1$ . Consider the following assertions:*

- (i) *The class  $\omega^n \in H^n(X, \mathbb{Z}_2(n))$  has coniveau  $\geq 1$ .*
- (ii) *The class  $\omega^{n+1} \in H^{n+1}(X, \mathbb{Z}_2(n+1))$  has coniveau  $\geq 2$ .*

*Then (i) implies (ii). Moreover, if  $\mathrm{CH}_0(X_C)$  is supported on a closed subvariety of  $X_C$  of dimension  $n-1$ , then the converse holds.*

*Proof.* Either (i) or (ii) implies that  $\omega^{n+1}$  has coniveau  $\geq 1$ , or equivalently that it vanishes in  $H_{\mathrm{nr}}^{n+1}(X, \mathbb{Z}_2(n+1))$ . Let us suppose this is the case; in particular,  $\omega^n \in I$ .

By the coniveau spectral sequence (2-3),  $\omega^n$  has coniveau  $\geq 1$  in  $X$  if and only if its class in  $H_{\mathrm{nr}}^n(X, \mathbb{Z}_2(n))$  vanishes, and  $\omega^{n+1}$  has coniveau  $\geq 2$  if and only if its class in  $H^1(X, \mathcal{H}_X^n(n+1))$  vanishes. Then consider the diagram

$$\begin{array}{ccc} H^n(\mathbf{R}, \mathbb{Z}_2(n)) & \xrightarrow[\cong]{\omega} & H^{n+1}(\mathbf{R}, \mathbb{Z}_2(n+1)) \\ \downarrow & & \downarrow \\ I_{\mathrm{nr}} & \xrightarrow{\delta} & H^1(X, \mathcal{H}_X^n(n+1)) \end{array}$$

which is commutative by Lemma 3.4. Contemplating it shows that (i) implies (ii).

Conversely, if  $\mathrm{CH}_0(X_C)$  is supported on a closed subvariety of  $X_C$  of dimension  $n-1$ , we have  $H_{\mathrm{nr}}^n(X_C, \mathbb{Z}_2) = 0$  by [Colliot-Thélène and Voisin 2012, Proposition 3.3(ii)]. Indeed, the argument given there for Betti cohomology over  $\mathbb{C}$ , which relies on decomposition of the diagonal, works as well for 2-adic cohomology over  $\mathbf{C}$ . It then follows from the exact sequence (3-2) that  $\delta$  is injective, proving that (ii) implies (i).  $\square$

#### 4. Cohomology of smooth double covers

Recall the notation of Section 3A. The polynomial  $F \in \mathbf{R}[X_0, \dots, X_n]$  is the homogenization of a nonzero positive semidefinite polynomial  $f \in \mathbf{R}[X_1, \dots, X_n]$ .

Its degree  $d$  is even. We introduced the double cover  $Y$  of  $\mathbb{P}_{\mathbf{R}}^n$  ramified over  $\{F = 0\}$  defined by the equation  $Y := \{Z^2 + F = 0\}$ .

Throughout this section, we make the additional hypothesis that  $\{F = 0\}$  is smooth so that  $Y$  is smooth. By Lemma 3.1,  $Y(\mathbf{R}) = \emptyset$ . The main goal of this section is to prove Propositions 4.8 and 4.12.

**4A. Geometric cohomology.** We first collect some needed results on the cohomology of  $Y_C$ . They follow from general theorems on the cohomology of weighted complete intersections due to Dimca [1985]. When  $n = 3$ , we could also have applied [Clemens 1983, Corollary 1.19 and Lemma 1.23].

**Proposition 4.1.** *Let  $H_C \in H^2(Y_C, \mathbb{Z}_2(1))$  be the class of  $\mathcal{O}_{\mathbb{P}_C^n}(1)$ .*

- (i) *The cohomology groups  $H^k(Y_C, \mathbb{Z}_2)$  have no torsion.*
- (ii) *If  $k \neq n$  is odd,  $H^k(Y_C, \mathbb{Z}_2) = 0$ .*
- (iii) *If  $0 \leq l < \frac{n}{2}$ ,  $H^{2l}(Y_C, \mathbb{Z}_2) \simeq \mathbb{Z}_2(l)$  as a  $G$ -module, and is generated by  $H_C^l$ .*
- (iv) *If  $\frac{n}{2} < l \leq n$ ,  $H^{2l}(Y_C, \mathbb{Z}_2) \simeq \mathbb{Z}_2(l)$  as a  $G$ -module, and has a generator  $\alpha_l$  such that  $2\alpha_l = H_C^l$ .*

*Proof.* It suffices to prove the equalities as  $\mathbb{Z}_2$ -modules (this means that it is possible to forget the twist indicating the action of  $G$ ), because one recovers the correct twist by noticing that the relevant cohomology groups are rationally generated by algebraic cycles.

Using the fact that  $Y_C$  is defined over an algebraically closed subfield that may be embedded in  $\mathbb{C}$  together with the invariance of étale cohomology under an extension of algebraically closed fields, it suffices to prove the lemma when  $C = \mathbb{C}$ . Moreover, by comparison with Betti cohomology, it suffices to prove it for Betti cohomology.

Since  $Y_C$  is a strongly smooth weighted complete intersection in the sense of [Dimca 1985], its cohomology groups have no torsion by Proposition 6(ii) of that paper. Moreover, its Betti numbers in degree  $k \neq n$  are computed in Dimca's Proposition 6(i).

If  $l < \frac{n}{2}$ , the class  $H_C^l \in H^{2l}(Y_C, \mathbb{Z}_2)$  cannot be divisible by 2 because the intersection product  $\frac{1}{2}H_C^l \cdot \frac{1}{2}H_C^l \cdot H_C^{n-2l} = \frac{1}{2}$  would not be an integer. It follows that  $H_C^l$  generates  $H^{2l}(Y_C, \mathbb{Z}_2)$ .

If  $l > \frac{n}{2}$ , since  $H_C^l \cdot H_C^{n-l} = 2$ , it follows by Poincaré duality that  $H^{2l}(Y_C, \mathbb{Z}_2)$  is generated by a class  $\alpha_l$  such that  $2\alpha_l = H_C^l$ .  $\square$

**4B. Preparation for a deformation argument.** In the next subsections, we perform some computations on the cohomology of  $Y$ . One of the arguments, in the proofs of Lemma 4.7 and Proposition 4.12, is a reduction to the Fermat double cover  $Y^\dagger := \{Z^2 + F^\dagger = 0\}$ , where  $F^\dagger := X_0^d + \cdots + X_n^d$  is the Fermat equation. This

deformation argument relies on a little bit of semialgebraic geometry, so it is convenient to collect the relevant lemmas here.

Let  $V := \mathbf{R}[X_0, \dots, X_n]_d$  be the space of degree  $d$  homogeneous polynomials viewed as an algebraic variety over  $\mathbf{R}$ . The discriminant  $\Delta \subset V$  is the closed algebraic subvariety parametrizing equations that do not define smooth hypersurfaces in  $\mathbb{P}_{\mathbf{R}}^n$ . It is irreducible, and a general point of  $\Delta$  defines a hypersurface with only one ordinary double point as singularities. Let  $\Delta' \subset \Delta$  be the closed algebraic subvariety parametrizing singular hypersurfaces that do not have only one ordinary double point as singularities; it has codimension  $\geq 2$  in  $V$ . We view the sets of  $\mathbf{R}$ -points  $V(\mathbf{R})$ ,  $\Delta(\mathbf{R})$  and  $\Delta'(\mathbf{R})$  as semialgebraic sets. Define

$$\Pi := \{H \in V(\mathbf{R}) \mid H(x_0, \dots, x_n) > 0 \text{ for every } (x_0, \dots, x_n) \in \mathbf{R}^{n+1} \setminus (0, \dots, 0)\}.$$

**Lemma 4.2.** *The set  $\Pi \subset V(\mathbf{R})$  is convex, open and semialgebraic. Moreover, the polynomials  $F$  and  $F^\dagger$  belong to  $\Pi$ .*

*Proof.* It is immediate that  $\Pi$  is convex. We prove that the complement of  $\Pi$  is a closed semialgebraic set. By homogeneity of  $H$ , it coincides with the projection to  $V(\mathbf{R})$  of

$$Q := \{(H, x_0, \dots, x_n) \in V(\mathbf{R}) \times \mathbb{S}^n \mid H(x_0, \dots, x_n) \leq 0\},$$

where  $\mathbb{S}^n := \{(x_0, \dots, x_n) \in \mathbf{R}^{n+1} \mid x_0^2 + \dots + x_n^2 = 1\}$  is the unit sphere.

That it is semialgebraic follows from the Tarski–Seidenberg theorem [Bochnak et al. 1998, Theorem 2.2.1]. To check that it is closed, it suffices to check that its intersection with every closed hypercube in  $V(\mathbf{R})$  is closed, which follows from [Bochnak et al. 1998, Theorem 2.5.8].

That  $F^\dagger \in \Pi$  is clear. We know that  $F \geq 0$  because it is positive semidefinite. Moreover, it cannot vanish on  $\mathbf{R}^{n+1} \setminus (0, \dots, 0)$  because  $Y(\mathbf{R}) \neq \emptyset$  by Lemma 3.1. This shows that  $F \in \Pi$ .  $\square$

Now choose a general affine subspace  $W \subset V$  of dimension 2 that contains  $F$  and  $F^\dagger$ .

**Lemma 4.3.** *The set  $\Pi \cap W(\mathbf{R}) \cap \Delta(\mathbf{R})$  is finite.*

*Proof.* Let  $H \in \Pi \cap W(\mathbf{R}) \cap \Delta(\mathbf{R})$ . Since  $H \in \Delta(\mathbf{R})$ ,  $\{H = 0\} \subset \mathbb{P}_{\mathbf{R}}^n$  is a singular hypersurface. Since  $H \in \Pi$ ,  $\{H = 0\}$  has no real point. Consequently,  $\{H = 0\}$  has (geometrically) at least two singular points: any singular point and its distinct complex conjugate. This shows that  $\Pi \cap W(\mathbf{R}) \cap \Delta(\mathbf{R}) \subset W(\mathbf{R}) \cap \Delta'(\mathbf{R})$ . But if  $W$  has been chosen to properly intersect  $\Delta'$ , the variety  $W \cap \Delta'$  is already finite.  $\square$

**Lemma 4.4.** *There exists a variety  $S$  over  $\mathbf{R}$ , two points  $s, s^\dagger \in S(\mathbf{R})$  and a morphism  $\rho : S \rightarrow W \setminus \Delta$  such that  $S(\mathbf{R})$  is semialgebraically connected,  $\rho(s) = F$ ,  $\rho(s^\dagger) = F^\dagger$  and  $\rho(S(\mathbf{R})) \subset \Pi$ .*

*Proof.* Choose a coordinate system on  $W$  for which  $F$  has coordinate  $(-1, 0)$  and  $F^\dagger$  has coordinate  $(1, 0)$ . By Lemma 4.2, the segment  $[F, F^\dagger]$  is included in  $\Pi$  and  $W(\mathbf{R}) \setminus \Pi$  is closed and semialgebraic. Consequently, combining Proposition 2.2.8(ii) and Theorem 2.5.8 of [Bochnak et al. 1998], we see that the distance between  $[F, F^\dagger]$  and  $W(\mathbf{R}) \setminus \Pi$  is positive. It follows that if  $\varepsilon \in \mathbf{R}$  is small enough, the ellipse  $\{x^2 + y^2/\varepsilon \leq 1\} \subset W(\mathbf{R})$ , which contains  $F$  and  $F^\dagger$ , is included in  $\Pi$ .

Now, consider the double cover  $\rho : W' := \{x^2 + y^2/\varepsilon + z^2 = 1\} \rightarrow W$  and define  $S := \rho^{-1}(W \setminus \Delta) \subset W'$ . That  $\rho(S(\mathbf{R}))$  is included in  $\Pi$  and contains  $F$  and  $F^\dagger$  follows from our choice of the ellipse. The semialgebraic set  $W'(\mathbf{R})$  is a sphere  $\mathbb{S}^2$ , and  $S(\mathbf{R})$  is the complement of a finite number of points in it by Lemma 4.3. This allows us to show by hand that it is semialgebraically path-connected, hence semialgebraically connected by [Bochnak et al. 1998, Proposition 2.5.13].  $\square$

Over the base  $S$ , there is a smooth projective family  $\mathcal{Y} \xrightarrow{\rho} S$  obtained by pulling back by  $\rho$  the universal family of smooth double covers over  $W \setminus \Delta$ . In particular,  $\mathcal{Y}_s \simeq Y$  and  $\mathcal{Y}_{s^\dagger} \simeq Y^\dagger$ . Since  $\rho(S(\mathbf{R})) \subset \Pi$ , we see that  $\mathcal{Y}(\mathbf{R}) = \emptyset$ .

**4C. Cohomology over  $\mathbf{R}$  when  $d \equiv 0[4]$ .** We start with a general lemma.

**Lemma 4.5.** *Let  $X$  be a smooth projective geometrically integral variety of dimension  $n$  over  $\mathbf{R}$  such that  $X(\mathbf{R}) = \emptyset$ . Then:*

- (i)  $H^{2n}(X, \mathbb{Z}_2(n)) \simeq \mathbb{Z}_2$ .
- (ii)  $H^{2n}(X, \mathbb{Z}_2(n+1)) \simeq \mathbb{Z}/2\mathbb{Z}$ .

*Proof.* We use the exact sequence (1-2), as well as Proposition 1.2(ii).

Consider  $H^{2n}(X, \mathbb{Z}_2(n)) \rightarrow H^{2n}(X_C, \mathbb{Z}_2) \xrightarrow{\pi_*} H^{2n}(X, \mathbb{Z}_2(n+1)) \rightarrow 0$ . The cohomology class of a closed point in  $H^{2n}(X, \mathbb{Z}_2(n))$  pulls back to twice the cohomology class of a closed point in  $H^{2n}(X_C, \mathbb{Z}_2)$ . This shows that  $H^{2n}(X, \mathbb{Z}_2(n+1))$  is torsion. From  $H^{2n}(X, \mathbb{Z}_2(n+1)) \rightarrow H^{2n}(X_C, \mathbb{Z}_2) \rightarrow H^{2n}(X, \mathbb{Z}_2(n)) \rightarrow 0$ , we deduce that  $\mathbb{Z}_2 \simeq H^{2n}(X_C, \mathbb{Z}_2) \rightarrow H^{2n}(X, \mathbb{Z}_2(n))$  is an isomorphism. The composition  $H^{2n}(X, \mathbb{Z}_2(n)) \xrightarrow{\pi^*} H^{2n}(X_C, \mathbb{Z}_2) \xrightarrow{\pi_*} H^{2n}(X, \mathbb{Z}_2(n))$  being multiplication by 2, we see that the image of  $H^{2n}(X, \mathbb{Z}_2(n)) \xrightarrow{\pi^*} H^{2n}(X_C, \mathbb{Z}_2)$  has index 2, so that  $H^{2n}(X, \mathbb{Z}_2(n+1)) = \mathbb{Z}/2\mathbb{Z}$ .  $\square$

We need information about  $\omega^{2n} \in H^{2n}(Y, \mathbb{Z}_2(2n))$  provided by Lemma 4.7 below. As a first step towards this result, we deal with the Fermat double cover  $Y^\dagger := \{Z^2 + F^\dagger = 0\}$ , where  $F^\dagger := X_0^d + \cdots + X_n^d$ .

**Lemma 4.6.** *Suppose  $n$  is odd and  $d \equiv 0[4]$ . Then  $\omega^{2n} \in H^{2n}(Y^\dagger, \mathbb{Z}_2(2n))$  is zero.*

*Proof.* The morphism  $\mu : Y^\dagger \rightarrow Q^n$  to  $Q^n := \{Z^2 + T_0^2 + \cdots + T_n^2 = 0\} \subset \mathbb{P}_{\mathbf{R}}^{n+1}$  defined by  $T_i = X_i^{d/2}$  has even degree because  $d \equiv 0[4]$ . By Lemma 4.5 applied to

$Y^\dagger$  and  $Q^n$ , there is a commutative diagram with surjective vertical arrows:

$$\begin{array}{ccc} \mathbb{Z}_2 = H^{2n}(Q_{\mathbf{C}}^n, \mathbb{Z}_2) & \xrightarrow{\mu_{\mathbf{C}}^*} & H^{2n}(Y_{\mathbf{C}}^\dagger, \mathbb{Z}_2) = \mathbb{Z}_2 \\ \downarrow & & \downarrow \\ \mathbb{Z}/2\mathbb{Z} = H^{2n}(Q^n, \mathbb{Z}_2(2n)) & \xrightarrow{\mu^*} & H^{2n}(Y^\dagger, \mathbb{Z}_2(2n)) = \mathbb{Z}/2\mathbb{Z} \end{array}$$

Since  $\mu_{\mathbf{C}}^*$  is the multiplication by the even number  $\deg(\mu)$ ,  $\mu_{\mathbf{C}}^*$  vanishes. Hence so does the composite  $H^{2n}(\mathbf{R}, \mathbb{Z}_2(2n)) \rightarrow H^{2n}(Q^n, \mathbb{Z}_2(2n)) \xrightarrow{\mu^*} H^{2n}(Y^\dagger, \mathbb{Z}_2(2n))$ .  $\square$

We deduce the same result for  $Y$  using a deformation argument:

**Lemma 4.7.** *Suppose  $n$  is odd and  $d \equiv 0[4]$ . Then  $\omega^{2n} \in H^{2n}(Y, \mathbb{Z}_2(2n))$  is zero.*

*Proof.* Lemma 4.6 and the diagram

$$\begin{array}{ccc} H^{2n}(\mathbf{R}, \mathbb{Z}_2(2n)) & \xrightarrow{\cong} & H^{2n}(\mathbf{R}, \mathbb{Z}/2\mathbb{Z}) \\ \downarrow & & \downarrow \\ H^{2n}(Y^\dagger, \mathbb{Z}_2(2n)) & \longrightarrow & H^{2n}(Y^\dagger, \mathbb{Z}/2\mathbb{Z}) \end{array}$$

show that  $H^{2n}(\mathbf{R}, \mathbb{Z}/2\mathbb{Z}) \rightarrow H^{2n}(Y^\dagger, \mathbb{Z}/2\mathbb{Z})$  vanishes.

Now consider the family  $\mathcal{Y} \xrightarrow{p} S$  constructed at the end of Section 4B. The varieties  $Y$  and  $Y^\dagger$  are members of this family and  $S(\mathbf{R})$  is semialgebraically connected. If we were working over the field  $\mathbb{R}$  of real numbers, we would use topological arguments (namely a  $G$ -equivariant version of Ehresmann's theorem applied to the fibration  $p_{\mathbf{C}}^{-1}(S(\mathbb{R})) \rightarrow S(\mathbb{R})$ ) to show that  $H^{2n}(\mathbb{R}, \mathbb{Z}/2\mathbb{Z}) \rightarrow H^{2n}(Y, \mathbb{Z}/2\mathbb{Z})$  vanishes as well. Over an arbitrary real closed field  $\mathbf{R}$ , the corresponding tools have been developed by Scheiderer [1994] and the topological arguments may be replaced by [Scheiderer 1994, Corollary 17.21].

Let us explain more precisely how to apply this result. In doing so, we freely use the notations of [Scheiderer 1994]. Consider the composition

$$\begin{aligned} H^{2n}(\mathbf{R}, \mathbb{Z}/2\mathbb{Z}) &\rightarrow H^{2n}(\mathcal{Y}, \mathbb{Z}/2\mathbb{Z}) \xleftarrow{\sim} H^{2n}(\mathcal{Y}_b, \mathbb{Z}/2\mathbb{Z}) \\ &\rightarrow H^0(S_b, R^{2n}p_{b*}\mathbb{Z}/2\mathbb{Z}) \rightarrow H^0(S_r, i^*R^{2n}p_{b*}\mathbb{Z}/2\mathbb{Z}), \end{aligned}$$

where the isomorphism  $H^{2n}(\mathcal{Y}_b, \mathbb{Z}/2\mathbb{Z}) \xrightarrow{\sim} H^{2n}(\mathcal{Y}, \mathbb{Z}/2\mathbb{Z})$  follows from [Scheiderer 1994, Example 2.14], taking into account Proposition 1.1 and the fact that  $\mathcal{Y}(\mathbf{R}) = \emptyset$ . By proper base change [Scheiderer 1994, Theorem 16.2(b)] and comparing étale and  $b$ -cohomology using [Scheiderer 1994, Example 2.14] once again, we see that the stalk of  $i^*R^{2n}p_{b*}\mathbb{Z}/2\mathbb{Z}$  is  $H^{2n}(Y, \mathbb{Z}/2\mathbb{Z})$  at  $s$  and  $H^{2n}(Y^\dagger, \mathbb{Z}/2\mathbb{Z})$  at  $s^\dagger$ . We have proven above that  $H^{2n}(\mathbf{R}, \mathbb{Z}/2\mathbb{Z})$  vanishes in the stalk at  $s^\dagger$ . But we know

that the sheaf  $i^* R^{2n} p_{b*} \mathbb{Z}/2\mathbb{Z}$  is locally constant on  $S_r$  by [Scheiderer 1994, Corollary 17.20(b)], and that  $S_r$  is connected by [Bochnak et al. 1998, Proposition 7.5.1(i)] and because  $S(\mathbf{R})$  is semialgebraically connected. Consequently,  $H^{2n}(\mathbf{R}, \mathbb{Z}/2\mathbb{Z})$  also vanishes in the stalk at  $s$ , so that  $H^{2n}(\mathbf{R}, \mathbb{Z}/2\mathbb{Z}) \rightarrow H^{2n}(Y, \mathbb{Z}/2\mathbb{Z})$  is zero.

To conclude that  $\omega^{2n} \in H^{2n}(Y, \mathbb{Z}_2(2n))$  vanishes, consider the exact diagram

$$\begin{array}{ccccc} H^{2n}(\mathbf{R}, \mathbb{Z}_2(2n)) & \xrightarrow{\cong} & H^{2n}(\mathbf{R}, \mathbb{Z}/2\mathbb{Z}) & & \\ \downarrow & & \downarrow & & \\ H^{2n}(Y, \mathbb{Z}_2(2n)) & \xrightarrow{2} & H^{2n}(Y, \mathbb{Z}_2(2n)) & \longrightarrow & H^{2n}(Y, \mathbb{Z}/2\mathbb{Z}) \end{array}$$

and notice that the multiplication by 2 map is zero by Lemma 4.5.  $\square$

**Proposition 4.8.** *Suppose that  $d \equiv 0[4]$ . Then  $\omega^{n+1} \in H^{n+1}(Y, \mathbb{Z}_2(n+1))$  is zero.*

*Proof.* Suppose not, and let  $k \geq n+1$  be such that  $\omega^k \in H^k(Y, \mathbb{Z}_2(k))$  is nonzero and  $\omega^{k+1} \in H^{k+1}(Y, \mathbb{Z}_2(k+1))$  vanishes. By Proposition 1.2(ii),  $k$  exists and  $k \leq 2n$ .

Consider the short exact sequence (1-2) applied to  $Y$ :

$$H^k(Y, \mathbb{Z}_2(k+1)) \xrightarrow{\pi^*} H^k(Y_C, \mathbb{Z}_2) \xrightarrow{\pi_*} H^k(Y, \mathbb{Z}_2(k)) \xrightarrow{\omega} H^{k+1}(Y, \mathbb{Z}_2(k+1)).$$

By hypothesis,  $\omega^k \in \text{Im}(\pi_*)$ . By Proposition 4.1(ii), since  $\omega^k \in H^k(Y, \mathbb{Z}_2(k))$  is nonzero,  $k$  has to be even, say  $k = 2l$ .

If  $l$  were even, we would have  $H^k(Y_C, \mathbb{Z}_2(k+1))^G = 0$  by Proposition 4.1(iv), and the Hochschild–Serre spectral sequence (1-1) would show that

$$\pi^* : H^k(Y, \mathbb{Z}_2(k+1)) \rightarrow H^k(Y_C, \mathbb{Z}_2)$$

is zero. Consequently,  $\text{Im}(\pi_*)$  has no torsion by Proposition 4.1(iv). This is a contradiction and shows that  $l$  is odd.

Let  $H \in H^2(Y, \mathbb{Z}_2(1))$  be the class of  $\mathcal{O}_{\mathbb{P}_{\mathbf{R}}^n}(1)$ . Since  $\pi^* H^l = H_C^l$ , and taking into account Proposition 4.1(iv), the only class in  $\text{Im}(\pi_*)$  that may be nonzero is  $\pi_* \alpha_l$ . Consequently, we have  $\omega^k = \pi_* \alpha_l$ .

Choose by the Bertini theorem an  $l$ -dimensional linear subspace  $\mathbb{P}_{\mathbf{R}}^l \subset \mathbb{P}_{\mathbf{R}}^n$  that is transverse to the smooth hypersurface  $\{F=0\}$ , and define  $i : Z \hookrightarrow Y$  to be the inverse image of  $\mathbb{P}_{\mathbf{R}}^l$  in  $Y$ ; it is a smooth double cover of  $\mathbb{P}_{\mathbf{R}}^l$  ramified over  $\{F=0\} \cap \mathbb{P}_{\mathbf{R}}^l$ . Consider the following commutative diagram, where the left horizontal arrows are restrictions, the right horizontal arrows are Gysin morphisms and the vertical ones are those appearing in the exact sequence (1-2):

$$\begin{array}{ccccc} H^k(Y_C, \mathbb{Z}_2) & \xrightarrow{i_C^*} & H^k(Z_C, \mathbb{Z}_2) & \xrightarrow{ic_*} & H^{2n}(Y_C, \mathbb{Z}_2) \\ \downarrow \pi_* & & \downarrow \pi_* & & \downarrow \pi_* \\ H^k(Y, \mathbb{Z}_2(k)) & \xrightarrow{i^*} & H^k(Z, \mathbb{Z}_2(k)) & \xrightarrow{i_*} & H^{2n}(Y, \mathbb{Z}_2(n+l)) \end{array}$$

Look at  $\alpha_l \in H^k(Y_C, \mathbb{Z}_2)$ . We have  $i_* i^* \pi_* \alpha_l = i_* i^* \omega^k = i_* \omega^k = 0$  by Lemma 4.7 applied to  $Z$ . On the other hand,  $\pi_* i_{C*} i_C^* \alpha_l = \pi_*(\alpha_l \cdot [Z_C]) = \pi_*(\alpha_l \cdot H_C^{n-l}) = \pi_* \alpha_n \neq 0$  because it is the generator of  $H^{2n}(Y, \mathbb{Z}_2(n+l)) \simeq \mathbb{Z}/2\mathbb{Z}$  as seen in Lemma 4.5. This is a contradiction.  $\square$

**4D. Cohomology over  $R$  when  $n$  is odd.** As in the previous paragraph, we reduce the computations we need to the case of a quadric. To do this, we collect a few results about the cohomology of quadrics. Analogues with 2-torsion coefficients of some of these computations appear in [Kahn and Sujatha 2000, §4].

We denote by  $Q^n := \{Z^2 + T_0^2 + \cdots + T_n^2 = 0\} \subset \mathbb{P}_R^{n+1}$  the  $n$ -dimensional projective anisotropic quadric, by  $U^n := Q^n \setminus Q^{n-1}$  its affine counterpart and by  $H \in H^2(Q^n, \mathbb{Z}_2(1))$  the class of  $\mathcal{O}_{\mathbb{P}_R^{n+1}}(1)$ .

**Lemma 4.9.** *The cohomology groups of  $U^n$  are as follows:*

$$H^k(U^n, \mathbb{Z}_2(j)) = \begin{cases} \mathbb{Z}_2 & \text{if } k = 0 \text{ and } j \equiv 0[2], \\ \mathbb{Z}/2\mathbb{Z} \cdot \omega^k & \text{if } 1 \leq k \leq n \text{ and } j \equiv k[2], \\ \mathbb{Z}_2 & \text{if } k = n \text{ and } j \equiv n+1[2], \\ 0 & \text{otherwise.} \end{cases}$$

*Proof.* The geometric cohomology groups of  $U^n$  are easily computed as  $G$ -modules from the description of the geometric cohomology groups of  $Q^n$  and  $Q^{n-1}$  as  $G$ -modules given in [SGA 7<sub>II</sub> 1973, XII Théorème 3.3], and from the long exact sequence of cohomology with support associated with  $Q^{n-1} \subset Q^n$ :

$$\cdots \rightarrow H^{k-2}(Q^{n-1}, \mathbb{Z}_2(-1)) \rightarrow H^k(Q^n, \mathbb{Z}_2) \rightarrow H^k(U^n, \mathbb{Z}_2) \rightarrow \cdots$$

One gets

$$H^k(U^n, \mathbb{Z}_2) = \begin{cases} \mathbb{Z}_2 & \text{if } k = 0, \\ \mathbb{Z}_2(n+1) & \text{if } k = n, \\ 0 & \text{otherwise.} \end{cases}$$

Consider the Hochschild–Serre spectral sequence (1-1) for  $U^n$ . The only possibly nonzero arrows in this spectral sequence are the  $d_n : E_{n+1}^{p,n} \rightarrow E_{n+1}^{p+n+1,0}$ . These are necessarily surjective as  $H^k(U^n, \mathbb{Z}_2(j)) = 0$  for  $k > n$  by Proposition 1.2(iii). This allows us to compute the spectral sequence entirely, and to deduce the lemma.  $\square$

**Lemma 4.10.** *Suppose that  $n$  is odd. Then  $H^n(Q^n, \mathbb{Z}_2(n)) \simeq (\mathbb{Z}/2\mathbb{Z})^{\lceil n/4 \rceil}$  generated by  $\omega^n, \omega^{n-4}H^2, \dots, \omega^{n-4\lfloor n/4 \rfloor}H^{2\lfloor n/4 \rfloor}$ . Moreover, the 2-torsion subgroup of  $H^{n+1}(Q^n, \mathbb{Z}_2(n+1))$  is  $H^{n+1}(Q^n, \mathbb{Z}_2(n+1))[2] \simeq (\mathbb{Z}/2\mathbb{Z})^{\lceil n/4 \rceil}$ , generated by  $\omega^{n+1}, \omega^{n-3}H^2, \dots, \omega^{n+1-4\lfloor n/4 \rfloor}H^{2\lfloor n/4 \rfloor}$ .*

*Proof.* Fix  $r \geq 0$ . The Gysin morphism  $H^{n-2r-2}(Q^{n-r-1}, \mathbb{Z}_2(n-r-1)) \rightarrow H^{n-2r}(Q^{n-r}, \mathbb{Z}_2(n-r))$  is part of a long exact sequence of cohomology with

supports. In this exact sequence, the morphisms

$$\begin{aligned} H^{n-2r-1}(Q^{n-r}, \mathbb{Z}_2(n-r)) &\rightarrow H^{n-2r-1}(U^{n-r}, \mathbb{Z}_2(n-r)), \\ H^{n-2r}(Q^{n-r}, \mathbb{Z}_2(n-r)) &\rightarrow H^{n-2r}(U^{n-r}, \mathbb{Z}_2(n-r)) \end{aligned}$$

are surjective. Indeed, in the degrees that come up, all the cohomology of  $U^{n-r}$  comes from the base field by Lemma 4.9, hence a fortiori from  $Q^{n-r}$ .

It follows that this Gysin morphism is injective, with a cokernel naturally isomorphic to  $H^{n-2r}(U^{n-r}, \mathbb{Z}_2(n-r)) = H^{n-2r}(\mathbf{R}, \mathbb{Z}_2(n-r))$ . This allows us to compute  $H^{n-2r}(Q^{n-r}, \mathbb{Z}_2(n-r))$  by decreasing induction on  $r$  and shows, since  $n$  is odd, that  $H^n(Q^n, \mathbb{Z}_2(n)) \simeq (\mathbb{Z}/2\mathbb{Z})^{\lceil n/4 \rceil}$  generated by  $\omega^n, \omega^{n-4}H^2, \dots, \omega^{n-4\lfloor n/4 \rfloor}H^{2\lfloor n/4 \rfloor}$ .

Considering the long exact sequence (1-2), and using our knowledge of the geometric cohomology of  $Q^n$  to get

$$0 \rightarrow H^n(Q^n, \mathbb{Z}_2(n)) \xrightarrow{\omega} H^{n+1}(Q^n, \mathbb{Z}_2(n+1)) \rightarrow H^{n+1}(Q_C^n, \mathbb{Z}_2) \simeq \mathbb{Z}_2, \quad (4-1)$$

the lemma follows.  $\square$

In the remainder of this section, we continue to suppose that  $n$  is odd. We consider the generator  $\gamma$  of  $H^{n+1}(Q_C^n, \mathbb{Z}_2) \simeq \mathbb{Z}_2$  such that  $2\gamma = H_C^{(n+1)/2}$  [SGA 7<sub>II</sub> 1973, XII Théorème 3.3].

If  $n \equiv 1[4]$ , define  $\delta := \pi_*\gamma \in H^{n+1}(Q^n, \mathbb{Z}_2(n+1))$ . If  $n \equiv 3[4]$ , define  $\delta := \pi_*\gamma - H^{(n+1)/2} \in H^{n+1}(Q^n, \mathbb{Z}_2(n+1))$ .

**Lemma 4.11.** *Suppose that  $n$  is odd. Then  $\delta \in H^{n+1}(Q^n, \mathbb{Z}_2(n+1))[2]$ . Moreover,  $\delta$  does not belong to the subgroup of  $H^{n+1}(Q^n, \mathbb{Z}_2(n+1))[2]$  generated by  $\omega^{n-3}H^2, \dots, \omega^{n+1-4\lfloor n/4 \rfloor}H^{2\lfloor n/4 \rfloor}$ .*

*Proof.* We suppose that  $n \equiv 1[4]$ . The arguments when  $n \equiv 3[4]$  are analogous.

Since  $G$  acts on  $H^{n+1}(Q_C^n, \mathbb{Z}_2) \simeq \mathbb{Z}_2$  by multiplication by  $(-1)^{(n+1)/2}$  and since  $n+1 \not\equiv \frac{1}{2}(n+1)[2]$ , the natural morphism  $H^{n+1}(Q^n, \mathbb{Z}_2(n+1)) \rightarrow H^{n+1}(Q_C^n, \mathbb{Z}_2)$  is zero. From the exact sequence (4-1), this implies that  $H^{n+1}(Q^n, \mathbb{Z}_2(n+1))$  is a 2-torsion group, so that  $\delta$  is 2-torsion.

Let us check that  $\delta$  is nonzero. From the exact sequence

$$H^{n+1}(Q^n, \mathbb{Z}_2(n)) \xrightarrow{\pi^*} H^{n+1}(Q_C^n, \mathbb{Z}_2) \xrightarrow{\pi_*} H^{n+1}(Q^n, \mathbb{Z}_2(n+1)),$$

we see that it suffices to prove that there does not exist a cohomology class  $\zeta \in H^{n+1}(Q^n, \mathbb{Z}_2(n))$  such that  $\pi^*\zeta = \gamma$ . If such a class existed,  $\pi^*(\zeta \cdot H^{(n-1)/2})$  would be a generator of  $H^{2n}(Q_C^n, \mathbb{Z}_2)$ . This would contradict the fact, proven in Lemma 4.5, that the cokernel of

$$H^{2n}(Q^n, \mathbb{Z}_2(n)) \rightarrow H^{2n}(Q_C^n, \mathbb{Z}_2)$$

is  $H^{2n}(Q^n, \mathbb{Z}_2(n+1)) \simeq \mathbb{Z}/2\mathbb{Z}$ .



Consider now the commutative diagram below, whose horizontal arrows are Gysin morphisms:

$$\begin{array}{ccccc}
 H^{n-3}(Q^{n-2}, \mathbb{Z}_2(n-1)) & \rightarrow & H^{n-1}(Q^{n-1}, \mathbb{Z}_2(n)) & \rightarrow & H^{n+1}(Q^n, \mathbb{Z}_2(n+1)) \\
 \downarrow \omega & & \downarrow \omega & & \downarrow \omega \\
 H^{n-2}(Q^{n-2}, \mathbb{Z}_2(n)) & \rightarrow & H^n(Q^{n-1}, \mathbb{Z}_2(n+1)) & \rightarrow & H^{n+2}(Q^n, \mathbb{Z}_2(n+2))
 \end{array} \quad (4-2)$$

Let us show that the lower horizontal arrows of (4-2) are injective. Considering  $\omega^{n-1} \in H^{n-1}(Q^{n-1}, \mathbb{Z}_2(n+1))$  and using Lemma 4.9 shows that

$$H^{n-1}(Q^{n-1}, \mathbb{Z}_2(n+1)) \rightarrow H^{n-1}(U^{n-1}, \mathbb{Z}_2(n+1))$$

is surjective, so that the map

$$H^{n-2}(Q^{n-2}, \mathbb{Z}_2(n)) \rightarrow H^n(Q^{n-1}, \mathbb{Z}_2(n+1))$$

is injective. Since  $H^{n+1}(U^n, \mathbb{Z}_2(n+2)) = 0$  by Lemma 4.9,

$$H^n(Q^{n-1}, \mathbb{Z}_2(n+1)) \rightarrow H^{n+2}(Q^n, \mathbb{Z}_2(n+2))$$

is also injective.

Suppose for contradiction that  $\delta$  may be written as a linear combination of the classes  $\omega^{n-3}H^2, \dots, \omega^{n+1-4\lfloor n/4 \rfloor}H^{2\lfloor n/4 \rfloor}$ . Then it is the image by the Gysin morphism  $H^{n-3}(Q^{n-2}, \mathbb{Z}_2(n-1)) \rightarrow H^{n+1}(Q^n, \mathbb{Z}_2(n+1))$  of a class  $\epsilon$  that is a linear combination of  $\omega^{n-3}, \dots, \omega^{n+1-4\lfloor n/4 \rfloor}H^{2\lfloor n/4 \rfloor-2}$ . The image of  $\epsilon$  in  $H^{n+2}(Q^n, \mathbb{Z}_2(n+2))$  is  $\delta \cdot \omega = \pi_* \gamma \cdot \omega = 0$ . By the injectivity result just proved,  $\epsilon \cdot \omega = 0$ . It follows that  $\omega^{n-2}, \dots, \omega^{n+2-4\lfloor n/4 \rfloor}H^{2\lfloor n/4 \rfloor-2}$  are not independent in  $H^{n-2}(Q^{n-2}, \mathbb{Z}_2(n))$ , contradicting Lemma 4.10.  $\square$

We return to our double cover  $Y \rightarrow \mathbb{P}_R^n$ . We recall from Proposition 4.1 that  $H^{n+1}(Y_C, \mathbb{Z}_2) = \mathbb{Z}_2(n+1)$  with a generator  $\alpha := \alpha_{(n+1)/2}$  such that  $2\alpha = H_C^{(n+1)/2}$ .

If  $n \equiv 1[4]$ , define  $\beta := \pi_* \alpha \in H^{n+1}(Y, \mathbb{Z}_2(n+1))$ . If  $n \equiv 3[4]$ , define instead  $\beta := \pi_* \alpha - H^{(n+1)/2} \in H^{n+1}(Y, \mathbb{Z}_2(n+1))$ , where  $H \in H^2(Y, \mathbb{Z}_2(1))$  is the class of  $\mathcal{O}_{\mathbb{P}_R^n}(1)$ .

**Proposition 4.12.** *Suppose that  $n$  is odd. Then  $\omega^{n+1}$  is a linear combination of  $\omega^{n-3}H^2, \dots, \omega^{n+1-4\lfloor n/4 \rfloor}H^{2\lfloor n/4 \rfloor}$  and  $\beta$  in  $H^{n+1}(Y, \mathbb{Z}_2(n+1))$ .*

*Proof.* Using a deformation argument as in the proof of Lemma 4.7, one reduces to the case of the Fermat double cover  $Y^\dagger := \{Z^2 + F^\dagger = 0\}$ , where  $F^\dagger := X_0^d + \dots + X_n^d$ .

As in Lemma 4.6, we consider the morphism  $\mu : Y^\dagger \rightarrow Q^n$  to the quadric  $Q^n := \{Z^2 + T_0^2 + \dots + T_n^2 = 0\} \subset \mathbb{P}_R^{n+1}$ , defined by  $T_i = X_i^{d/2}$ . Note that  $\mu^*H = \left(\frac{d}{2}\right)H$ , so that  $\mu_C^* \gamma = \left(\frac{d}{2}\right)^{(n+1)/2} \alpha$  and  $\mu^* \delta = \left(\frac{d}{2}\right)^{(n+1)/2} \beta$ .

By Lemma 4.10, the group  $H^{n+1}(Q^n, \mathbb{Z}_2(n+1))[2]$  is freely generated by  $\omega^{n+1}, \omega^{n-3}H^2, \dots, \omega^{n+1-4\lfloor n/4 \rfloor}H^{2\lfloor n/4 \rfloor}$ . By Lemma 4.11,  $\delta$  does not belong to the

subgroup generated by  $\omega^{n-3}H^2, \dots, \omega^{n+1-4\lfloor n/4 \rfloor}H^{2\lfloor n/4 \rfloor}$ . Thus,  $\omega^{n+1}$  is a linear combination of  $\delta, \omega^{n-3}H^2, \dots, \omega^{n+1-4\lfloor n/4 \rfloor}H^{2\lfloor n/4 \rfloor}$  in  $H^{n+1}(Q^n, \mathbb{Z}_2(n+1))[2]$ .

Pulling back this relation by  $\mu$ , it follows that  $\omega^{n+1}$  is a linear combination of  $\beta, \omega^{n-3}H^2, \dots, \omega^{n+1-4\lfloor n/4 \rfloor}H^{2\lfloor n/4 \rfloor}$  in  $H^{n+1}(Y^\dagger, \mathbb{Z}_2(n+1))$ , as wanted.  $\square$

**Corollary 4.13.** *Suppose that  $n$  is odd. If  $\alpha \in H^{n+1}(Y_C, \mathbb{Z}_2)$  has coniveau  $\geq 2$ , then  $\omega^{n+1} \in H^{n+1}(Y, \mathbb{Z}_2(n+1))$  has coniveau  $\geq 2$ .*

*Proof.* The statement follows from Proposition 4.12, because the cohomology classes  $\omega^{n-3}H^2, \dots, \omega^{n+1-4\lfloor n/4 \rfloor}H^{2\lfloor n/4 \rfloor}$ , as well as  $H^{(n+1)/2}$  when  $n \neq 1$ , obviously have coniveau  $\geq 2$  as multiples of  $H^2$ .  $\square$

## 5. A geometric coniveau computation

In this section, we fix an odd integer  $n \geq 3$  and take  $d := 2n$ .

We work over an algebraically closed field  $C$  of characteristic 0. We consider  $F \in C[X_0, \dots, X_n]_d$  a homogeneous degree  $d$  polynomial such that  $\{F = 0\}$  is smooth. Let  $Y$  be the double cover of  $\mathbb{P}_C^n$  ramified over  $\{F = 0\}$  defined by the equation  $Y := \{Z^2 + F = 0\}$ , and  $H := \mathcal{O}_{\mathbb{P}_C^n}(1)$ .

By Proposition 4.1,  $H^{n+1}(Y, \mathbb{Z}_2)$  has a generator  $\alpha$  such that  $2\alpha = H^{(n+1)/2}$ . In order to apply Corollary 4.13, we need to answer positively the following:

**Question 5.1.** Does  $\alpha$  have coniveau  $\geq 2$ ?

When  $n = 3$ ,  $Y$  is a sextic double solid, and this is very easy:

**Lemma 5.2.** *If  $n = 3$ ,  $\alpha$  has coniveau  $\geq 2$ .*

*Proof.* A dimension count (see for instance Lemma 5.6 below) shows that  $Y$  contains a line, that is, a curve of degree 1 against  $H$ . The cohomology class of such a curve is  $\alpha$ , so that  $\alpha$  is algebraic, hence of coniveau 2.  $\square$

In what follows, we answer Question 5.1 positively when  $n = 5$ , following an argument of Voisin. We comment on the  $n \geq 7$  case in Section 5D.

**Proposition 5.3.** *If  $n = 5$ ,  $\alpha$  has coniveau  $\geq 2$ .*

**5A. Reductions.** The following reductions are standard. We include them because we do not know a convenient reference.

**Lemma 5.4.** *Let  $C \subset C'$  be an extension of algebraically closed fields. Then the answer to Question 5.1 is positive for  $Y$  over  $C$  if and only if it is for  $Y_{C'}$  over  $C'$ .*

*Proof.* It is clear that if  $\alpha$  has coniveau  $\geq 2$  over  $C$ , it has coniveau  $\geq 2$  over  $C'$ .

Suppose conversely that there is a closed subset  $Z \subset Y_{C'}$  of codimension  $\geq 2$  such that  $\alpha_{C'}$  vanishes in  $Y_{C'} \setminus Z$ . Taking an extension of finite type of  $C$  over which  $Z$  is defined, spreading out and shrinking the base gives a smooth integral variety  $B$  over  $C$ , and a subvariety  $Z_B \subset Y \times B$  of codimension  $\geq 2$  in the fibers

of  $p_2 : Y \times B \rightarrow B$  such that  $p_1^* \alpha$  vanishes in the generic geometric fiber of  $p_2 : (Y \times B \setminus Z_B) \rightarrow B$ . The existence of cospecialization maps for smooth morphisms [SGA 4<sub>1/2</sub> 1977, Arcata V (1.6)] implies that  $\alpha$  vanishes in every geometric fiber of  $p_2 : (Y \times B \setminus Z_B) \rightarrow B$ . Taking the fiber over a  $C$ -point of  $B$  shows that  $\alpha$  has coniveau  $\geq 2$ .  $\square$

**Lemma 5.5.** *In order to answer Question 5.1 positively in general, it suffices to answer it over the field  $\mathbb{C}$  of complex numbers, for a general choice of  $F$ .*

*Proof.* Let  $U \subset \mathbb{C}[X_0, \dots, X_n]_d$  be a Zariski-open subset of degree  $d$  polynomials  $F$  as in the hypothesis: for  $F \in U(\mathbb{C})$ ,  $\{F = 0\}$  is smooth and  $\alpha$  is of coniveau  $\geq 2$ .

Let  $K$  be an algebraic closure of the function field  $\mathbb{C}(U)$ , let  $F_K$  be the generic polynomial and let  $Y_K$  be the associated universal double cover. Choosing an isomorphism  $K \simeq \mathbb{C}$  such that the induced polynomial  $F_{\mathbb{C}}$  belongs to  $U$  shows that the answer to Question 5.1 is positive for  $Y_K$ .

Let us now deal with the case  $C = \mathbb{C}$ . It is possible to find the spectrum  $T$  of a strictly henselian discrete valuation ring and a morphism  $T \rightarrow \mathbb{C}[X_0, \dots, X_n]_d$  sending the closed point of  $T$  to the polynomial associated to  $Y$  and its generic point to the generic point of  $U$ . Let  $Y_T$  be the induced family of double covers. Up to replacing  $T$  by a finite extension, there exists a codimension 2 subset  $Z \subset Y_T$  flat over  $T$  such that  $\alpha$  vanishes in the complement of  $Z$  in the generic geometric fiber. Using cospecialization maps again shows that  $\alpha$  vanishes in the complement of  $Z$  in the special fiber, so that the answer to Question 5.1 is positive for  $Y$ .

In general, choose an algebraically closed subfield of finite transcendence degree of  $C$  over which  $Y$  is defined, embed it in  $\mathbb{C}$ , and apply Lemma 5.4.  $\square$

**5B. The variety of lines.** We define  $F(Y)$  to be the Fano variety of lines of  $Y$ , that is, the Hilbert scheme of  $Y$  parametrizing degree 1 curves in  $Y$ . We also introduce the universal family  $I \subset Y \times F(Y)$  and denote by  $q : I \rightarrow Y$  and  $p : I \rightarrow F(Y)$  the natural projections.

The following lemma is well known for hypersurfaces [Barth and Van de Ven 1978/79, §3] or complete intersections [Debarre and Manivel 1998, Théorème 2.1], and the proof in our situation is similar.

**Lemma 5.6.** *If  $Y$  is general,  $F(Y)$  is smooth, nonempty and of dimension  $n - 2$ .*

*Proof.* First, an easy dimension count shows that if  $F$  is general,  $\{F = 0\}$  contains no line [Barth and Van de Ven 1978/79, §3]. It follows that for such a general  $Y$ ,  $F(Y)$  is a double étale cover of the variety  $G(Y)$  of lines in  $\mathbb{P}_{\mathbb{C}}^n$  on which the restriction of  $F$  is a square.

One introduces the space  $V \subset \mathbb{C}[X_0, \dots, X_n]_d$  of degree  $d$  polynomials  $F$  such that  $\{F = 0\}$  is smooth. We consider the universal double cover  $Y_V \rightarrow V$ , and the universal Fano variety of lines  $F(Y)_V \rightarrow V$ , viewed as a double cover of  $G(Y)_V$

that is étale generically over  $V$ . Looking at the natural projection from  $G(Y)_V$  to the grassmannian of lines in  $\mathbb{P}_{\mathbb{C}}^n$ , one sees that  $G(Y)_V$  is smooth of dimension  $\dim(V) + n - 2$ . Since we are in characteristic 0, the general fiber of  $G(Y)_V \rightarrow V$  is smooth, and it remains to show that this morphism is dominant, or that  $F(Y)_V \rightarrow V$  is dominant. We do it by finding one point at which it is smooth.

To do so, we fix a line  $L$  with equations  $X_2 = \cdots = X_n = 0$ , and a degree  $d$  polynomial  $F$  such that the restriction of  $F$  to  $L$  is the square of  $H \in \mathbb{C}[X_0, X_1]_n$ . The double cover  $Y$  may be viewed naturally as the zero-locus of a section of  $\mathcal{O}(d)$  in the total space  $E \rightarrow \mathbb{P}_{\mathbb{C}}^n$  of the line bundle  $\mathcal{O}_{\mathbb{P}_{\mathbb{C}}^n}(n)$ . The inverse image of  $L$  in  $Y$  splits into the union of two lines. Let  $\Lambda$  be one of them. The normal exact sequence  $0 \rightarrow N_{\Lambda/Y} \rightarrow N_{\Lambda/E} \rightarrow N_{Y/E}|_{\Lambda} \rightarrow 0$  reads

$$0 \rightarrow N_{\Lambda/Y} \rightarrow \mathcal{O}(1)^{\oplus n-1} \oplus \mathcal{O}(n) \rightarrow \mathcal{O}(2n) \rightarrow 0.$$

The same computation as the one carried out in [Barth and Van de Ven 1978/79, §2] for hypersurfaces shows that the last arrow is given by  $(\partial F/\partial X_2, \dots, \partial F/\partial X_n, H)$ . Consequently, if  $H^0(\Lambda, \mathcal{O}(2n))$  is generated by multiples of  $\partial F/\partial X_2, \dots, \partial F/\partial X_n$  and  $H$ , then  $H^1(\Lambda, N_{\Lambda/Y}) = 0$  and  $\Lambda$  corresponds to a smooth point of the relative Hilbert scheme  $F(Y)_V \rightarrow V$ , as wanted. It is easy to find a polynomial  $F$  satisfying this condition.  $\square$

**Lemma 5.7.** *If there exists a smooth  $Y$  over  $\mathbb{C}$  such that  $F(Y)$  is smooth of dimension  $n - 2$ , and a cohomology class  $\zeta \in H^{n-1}(F(Y), \mathbb{Z}_2)$  such that  $q_* p^* \zeta$  is an odd multiple of  $\alpha$ , then Question 5.1 has a positive answer.*

*Proof.* By Lemma 5.5 and Lemma 5.6, it suffices to consider a double cover over the complex numbers whose variety of lines is smooth of dimension  $n - 2$ . By Ehresmann's theorem, the existence of a cohomology class  $\zeta$  as in our hypothesis does not depend on  $Y$  (as long as  $Y$  and  $F(Y)$  are smooth). Consequently, it suffices to answer Question 5.1 for a double cover  $Y$  for which such a  $\zeta$  exists.

On the one hand  $2\alpha = H^{(n+1)/2}$  is algebraic, hence of coniveau  $\geq 2$ . On the other hand  $\zeta$  has coniveau  $\geq 1$  because it vanishes on any affine open subset of  $F(Y)$ . It follows that  $q_* p^* \zeta$  has coniveau  $\geq 2$  because  $q : I \rightarrow Y$  is not dominant by dimension. Combining these two assertions, we see that  $\alpha$  has coniveau  $\geq 2$ .  $\square$

**5C. A degeneration argument.** In this subsection, we set  $n = 5$  and  $d = 10$ , and we prove Proposition 5.3 by checking the hypothesis of Lemma 5.7.

To do so, we choose four homogeneous polynomials  $P \in \mathbb{C}[X_0, \dots, X_5]_5$ ,  $G \in \mathbb{C}[X_1, X_2, X_3]_5$ , and  $Q_1, Q_2 \in \mathbb{C}[X_0, \dots, X_5]_9$ , and we set

$$F_0 := X_0^{10} + PG + Q_1 X_4 + Q_2 X_5 \in \mathbb{C}[X_0, \dots, X_5]_{10}.$$

The reason for this choice is that  $F_0$  restricts to a square on the cone  $\Gamma := \{G = X_4 = X_5 = 0\}$ , so that the inverse image of  $\Gamma$  in  $Y_0 := \{Z^2 + F_0 = 0\}$  has two irreducible

components, giving rise to two 1-dimensional families of lines in  $Y_0$ . We denote by  $\Phi_0 \subset F(Y_0)$  the curve corresponding to one of these families; it is naturally isomorphic to  $\{G = 0\} \subset \mathbb{P}_{\mathbb{C}}^2$ .

**Lemma 5.8.** *If  $P, G, Q_1, Q_2$  have been chosen general, then  $Y_0$  is smooth,  $\Phi_0$  is smooth and  $F(Y_0)$  is smooth of dimension 3 along  $\Phi_0$ .*

*Proof.* To check that the general zero-locus of such an  $F_0$  is smooth, it suffices to deal with equations of the form  $\lambda X_0^{10} + \mu X_1^{10} + Q_1 X_4 + Q_2 X_5 \in \mathbb{C}[X_0, \dots, X_5]_{10}$ . These form a linear system, so a general one among these is smooth outside of the base locus  $\{X_0 = X_1 = X_4 = X_5 = 0\}$  by the Bertini theorem. But there exists a particular one that is smooth on the base locus: take  $Q_1 = X_2^9$  and  $Q_2 = X_3^9$ . It follows that the general one is smooth everywhere.

To conclude, fix  $G$  such that  $\Phi_0 \simeq \{G = 0\} \subset \mathbb{P}_{\mathbb{C}}^2$  is smooth. It suffices to prove that for every  $\Lambda \in \Phi_0$ ,  $F(Y_0)$  is smooth of dimension 3 at  $\Lambda$ , with possible exceptions on a codimension 2 subset of the parameter space for  $P, Q_1$  and  $Q_2$ . By the computations in the proof of Lemma 5.6, we need to show that, outside such a subset,  $H^0(\Lambda, \mathcal{O}(10))$  is generated by multiples of  $\partial F_0 / \partial X_2, \dots, \partial F_0 / \partial X_5$  and  $X_0^5$ .

This amounts to showing that, outside of a codimension 2 subset of the parameter space for  $P \in \mathbb{C}[X_0, X_1]_5$  and  $Q_1, Q_2 \in \mathbb{C}[X_0, X_1]_9$ ,  $H^0(\mathbb{P}^1, \mathcal{O}(10))$  is generated by multiples of  $X_0^5, PX_1^4, Q_1$  and  $Q_2$ . This is easy to see, by exhibiting a complete curve in the projectivized parameter space avoiding the bad locus.  $\square$

Now, let  $\Delta$  be a small enough disk in  $\mathbb{C}[X_0, \dots, X_5]_{10}$  centered around the polynomial  $F_0$  given by Lemma 5.8. Let  $Y_{\Delta}$  be the family of double covers over it, and  $F(Y)_{\Delta}$  the corresponding family of varieties of lines.

Recall that  $\Phi_0$  is a smooth proper subvariety of the smooth locus of the special fiber  $F(Y_0)$ . Using the flow of a vector field as in the proof of Ehresmann's theorem, one sees that  $\Phi_0$  deforms (as a differentiable submanifold) to nearby fibers for which  $F(Y_t)$  is smooth, giving rise to a cohomology class  $\zeta_t = [\Phi_t] \in H^{n-1}(F(Y_t), \mathbb{Z}_2)$ . We compute  $q_* p^* \zeta_t = q_* [p^{-1}(\Phi_t)] = q_* [p^{-1}(\Phi_0)]$ ; it is the cycle class of the cone  $\Gamma$  that is equal to  $5\alpha$ .

**5D. Remarks.** When  $n \geq 7$ , an argument analogous to that of Section 5C fails, because one gets a double cover  $Y_0$  whose variety of lines is singular along a codimension 2 subset of the subvariety  $\Phi_0$  that we would like to deform to nearby fibers  $F(Y_t)$ .

It might still be possible to show, by another argument, the existence of a cohomology class  $\zeta$  allowing application of Lemma 5.7. To do so, one would need to compute part of the integral cohomology of  $F(Y)$ . The rational cohomology of  $F(Y)$  in the required degree is well understood thanks to [Debarre and Manivel 1998, Théorème 3.4] (where the computations are carried out in the analogous

setting of hypersurfaces or complete intersections). However, Debarre and Manivel's approach, relying on the Hodge decomposition, does not allow control of the integral cohomology groups of  $F(Y)$ .

## 6. Proof of the main theorem

We now come back to our main goal: the proof of Theorem 0.1.

**6A. The generic case.** Let us first put together what we have obtained so far. Fix  $n \geq 2$ . Define  $d(n)$  by setting  $d(n) := 2n$  if  $n$  is even or equal to 3 or 5 and  $d(n) := 2n - 2$  if  $n \geq 7$  is odd.

**Proposition 6.1.** *Let  $f \in \mathbf{R}[X_1, \dots, X_n]$  be a positive semidefinite polynomial of degree  $d(n)$  whose homogenization  $F$  defines a smooth hypersurface in  $\mathbb{P}_{\mathbf{R}}^n$ . Then  $f$  is a sum of  $2^n - 1$  squares in  $\mathbf{R}(X_1, \dots, X_n)$ .*

*Proof.* We consider the double cover  $Y$  of  $\mathbb{P}_{\mathbf{R}}^n$  ramified over  $\{F = 0\}$  defined by the equation  $Y := \{Z^2 + F = 0\}$ . The variety  $Y$  is smooth,  $Y(\mathbf{R}) = \emptyset$  by Lemma 3.1, and computing that the anticanonical bundle  $-K_Y = \mathcal{O}_{\mathbb{P}_{\mathbf{R}}^n}(n + 1 - d(n)/2)$  of  $Y$  is ample, one sees that  $Y_{\mathbf{C}}$  is Fano, hence rationally connected.

By Proposition 3.2, we need to show that the level of  $\mathbf{R}(Y)$  is  $< 2^n$ . Applying Proposition 3.3 (iii) $\Rightarrow$ (i), we have to prove that  $\omega^n \in H^n(Y, \mathbb{Z}_2(n))$  has coniveau  $\geq 1$ . Finally, since  $Y_{\mathbf{C}}$  is rationally connected, the converse Proposition 3.5 (ii) $\Rightarrow$ (i) holds: we only have to check that  $\omega^{n+1} \in H^{n+1}(Y, \mathbb{Z}_2(n+1))$  has coniveau  $\geq 2$ .

When  $n \neq 3$  or 5,  $d(n) \equiv 0[4]$  so that  $\omega^{n+1} \in H^{n+1}(Y, \mathbb{Z}_2(n+1))$  vanishes by Proposition 4.8.

When  $n = 3$  or 5,  $\omega^{n+1} \in H^{n+1}(Y, \mathbb{Z}_2(n+1))$  is seen to be of coniveau  $\geq 2$  by combining Corollary 4.13 and either Lemma 5.2 when  $n = 3$  or Proposition 5.3 when  $n = 5$ .  $\square$

**6B. A specialization argument.** We do not know how to deal with singular equations using the same arguments because one has too little control on the geometry of (a resolution of singularities of) the variety  $Y$ . Instead, we rely on a specialization argument, that will also take care of the lower values of the degree.

**Theorem 6.2** (Theorem 0.1). *Let  $f \in \mathbf{R}[X_1, \dots, X_n]$  be a positive semidefinite polynomial of degree  $\leq d(n)$ . Then  $f$  is a sum of  $2^n - 1$  squares in  $\mathbf{R}(X_1, \dots, X_n)$ .*

*Proof.* Consider  $g := f + t(1 + \sum_{i=1}^n X_i^{d(n)}) \in \mathbf{R}(t)[X_1, \dots, X_n]$ . It is a degree  $d(n)$  polynomial whose homogenization defines a smooth hypersurface in  $\mathbb{P}_{\mathbf{R}}^n$ , because so does its specialization  $1 + \sum_{i=1}^n X_i^{d(n)}$ . Let  $\mathbf{S} := \bigcup_r \mathbf{R}((t^{1/r}))$  be a real closed extension of  $\mathbf{R}(t)$ . By Artin's solution [1927] to Hilbert's 17th problem,  $f$  is a sum of squares in  $\mathbf{R}(X_1, \dots, X_n)$ , hence still a positive semidefinite polynomial viewed in  $\mathbf{S}[X_1, \dots, X_n]$ . Consequently, since  $t = (t^{1/2})^2$  is a square in  $\mathbf{S}$ ,

$g \in \mathbf{S}[X_1, \dots, X_n]$  is a positive semidefinite polynomial. Applying Proposition 6.1 over the real closed field  $\mathbf{S}$ , we see that  $g$  is a sum of  $2^n - 1$  squares in  $\mathbf{S}(X_1, \dots, X_n)$ : one has  $g = \sum_{i=1}^{2^n-1} h_i^2$ .

Consider the  $t$ -adic valuation on  $\mathbf{S}$ . Applying  $n$  times successively [Bourbaki 1985, Chapitre VI §10, Proposition 2], we can extend it to a valuation  $v$  on  $\mathbf{S}(X_1, \dots, X_n)$  that is trivial on  $\mathbf{R}(X_1, \dots, X_n)$ , and whose residue field is isomorphic to  $\mathbf{R}(X_1, \dots, X_n)$ . Note that these choices imply that  $v(g) = 0$  and that the reduction of  $g$  modulo  $v$  is  $f \in \mathbf{R}(X_1, \dots, X_n)$ .

Define  $m := \inf_i v(h_i)$  and notice that  $m \leq 0$  because  $v(g) = 0$ . Suppose for contradiction that  $m < 0$  and let  $j$  be such that  $v(h_j) = m$ . Then it is possible to reduce the equality  $gh_j^{-2} = \sum_{i=1}^{2^n-1} (h_i h_j^{-1})^2$  modulo  $v$ . This is absurd because we get a nontrivial sum of squares that is zero in  $\mathbf{R}(X_1, \dots, X_n)$ . This shows that  $m = 0$ . Consequently, it is possible to reduce the equality  $g = \sum_{i=1}^{2^n-1} h_i^2$  modulo  $v$ , showing that  $f$  is a sum of  $2^n - 1$  squares in  $\mathbf{R}(X_1, \dots, X_n)$  as wanted.  $\square$

We conclude by stating explicitly the following consequence of our proof.

**Proposition 6.3.** *If  $n \geq 3$  is odd and if Question 5.1 has a positive answer, then Theorem 0.1 also holds in  $n$  variables and degree  $d = 2n$ .*

### Acknowledgements

I have benefited from numerous discussions with Olivier Wittenberg, which have shaped my understanding of the cohomology of real algebraic varieties and have been very important for the completion of this work.

I am grateful to Claire Voisin for explaining to me the coniveau computation contained in Section 5 that was used to deal with the  $n = 5$  and  $d = 10$  case of Theorem 0.1.

### References

- [Artin 1927] E. Artin, “Über die Zerlegung definiter Funktionen in Quadrate”, *Abh. Math. Sem. Univ. Hamburg* **5**:1 (1927), 100–115. MR Zbl
- [Barth and Van de Ven 1978/79] W. Barth and A. Van de Ven, “Fano varieties of lines on hypersurfaces”, *Arch. Math. (Basel)* **31**:1 (1978/79), 96–104. MR Zbl
- [Bloch and Ogus 1974] S. Bloch and A. Ogus, “Gersten’s conjecture and the homology of schemes”, *Ann. Sci. École Norm. Sup. (4)* **7** (1974), 181–201. MR Zbl
- [Bloch and Srinivas 1983] S. Bloch and V. Srinivas, “Remarks on correspondences and algebraic cycles”, *Amer. J. Math.* **105**:5 (1983), 1235–1253. MR Zbl
- [Bochnak et al. 1998] J. Bochnak, M. Coste, and M.-F. Roy, *Real algebraic geometry*, *Ergebnisse der Mathematik* (3) **36**, Springer, Berlin, 1998. MR Zbl
- [Bourbaki 1985] N. Bourbaki, *Éléments de mathématique: Algèbre commutative, Chapitres 5 à 7*, Masson, Paris, 1985. MR Zbl

- [Cassels et al. 1971] J. W. S. Cassels, W. J. Ellison, and A. Pfister, “On sums of squares and on elliptic curves over function fields”, *J. Number Theory* **3** (1971), 125–149. MR Zbl
- [Clemens 1983] C. H. Clemens, “Double solids”, *Adv. in Math.* **47**:2 (1983), 107–230. MR Zbl
- [Colliot-Thélène 1992] J.-L. Colliot-Thélène, “Real rational surfaces without a real point”, *Arch. Math. (Basel)* **58**:4 (1992), 392–396. MR Zbl
- [Colliot-Thélène 1993] J.-L. Colliot-Thélène, “The Noether–Lefschetz theorem and sums of 4-squares in the rational function field  $\mathbf{R}(x, y)$ ”, *Compositio Math.* **86**:2 (1993), 235–243. MR Zbl
- [Colliot-Thélène and Parimala 1990] J.-L. Colliot-Thélène and R. Parimala, “Real components of algebraic varieties and étale cohomology”, *Invent. Math.* **101**:1 (1990), 81–99. MR Zbl
- [Colliot-Thélène and Scheiderer 1996] J.-L. Colliot-Thélène and C. Scheiderer, “Zero-cycles and cohomology on real algebraic varieties”, *Topology* **35**:2 (1996), 533–559. MR Zbl
- [Colliot-Thélène and Voisin 2012] J.-L. Colliot-Thélène and C. Voisin, “Cohomologie non ramifiée et conjecture de Hodge entière”, *Duke Math. J.* **161**:5 (2012), 735–801. MR Zbl
- [Colliot-Thélène et al. 1997] J.-L. Colliot-Thélène, R. T. Hoobler, and B. Kahn, “The Bloch–Ogus–Gabber theorem”, pp. 31–94 in *Algebraic K-theory* (Toronto, 1996), edited by V. P. Snaith, Fields Institute Communications **16**, American Mathematical Society, Providence, RI, 1997. MR Zbl
- [Debarre and Manivel 1998] O. Debarre and L. Manivel, “Sur la variété des espaces linéaires contenus dans une intersection complète”, *Math. Ann.* **312**:3 (1998), 549–574. MR Zbl
- [Dimca 1985] A. Dimca, “Monodromy and Betti numbers of weighted complete intersections”, *Topology* **24**:3 (1985), 369–374. MR Zbl
- [Elman and Lam 1972] R. Elman and T. Y. Lam, “Pfister forms and  $K$ -theory of fields”, *J. Algebra* **23** (1972), 181–213. MR Zbl
- [Hilbert 1888] D. Hilbert, “Ueber die Darstellung definiter Formen als Summe von Formenquadraten”, *Math. Ann.* **32**:3 (1888), 342–350. MR Zbl
- [Jannsen 1988] U. Jannsen, “Continuous étale cohomology”, *Math. Ann.* **280**:2 (1988), 207–245. MR Zbl
- [Jannsen 2000] U. Jannsen, “Letter from Jannsen to Gross on higher Abel–Jacobi maps”, pp. 261–275 in *The arithmetic and geometry of algebraic cycles* (Banff, AB, 1998), edited by B. B. Gordon et al., NATO Sci. Ser. C Math. Phys. Sci. **548**, Kluwer Academic Publishers, Dordrecht, Netherlands, 2000. MR Zbl
- [Kahn 2012] B. Kahn, “Classes de cycles motiviques étales”, *Algebra Number Theory* **6**:7 (2012), 1369–1407. MR Zbl
- [Kahn and Sujatha 2000] B. Kahn and R. Sujatha, “Motivic cohomology and unramified cohomology of quadrics”, *J. Eur. Math. Soc. (JEMS)* **2**:2 (2000), 145–177. MR Zbl
- [Kerz 2009] M. Kerz, “The Gersten conjecture for Milnor  $K$ -theory”, *Invent. Math.* **175**:1 (2009), 1–33. MR Zbl
- [Krasnov 1994] V. A. Krasnov, “On the equivariant Grothendieck cohomology of a real algebraic variety and its application”, *Izv. Ross. Akad. Nauk Ser. Mat.* **58**:3 (1994), 36–52. In Russian; translated in *Russian Acad. Sci. Izv. Math.* **44**:3 (1995), 461–477. MR Zbl
- [Lam 1980] T. Y. Lam, *The algebraic theory of quadratic forms*, Benjamin/Cummings Publishing Co., Reading, MA, 1980. MR Zbl
- [Pfister 1965] A. Pfister, “Zur Darstellung von  $-1$  als Summe von Quadraten in einem Körper”, *J. London Math. Soc.* **40** (1965), 159–165. MR Zbl
- [Pfister 1967] A. Pfister, “Zur Darstellung definiter Funktionen als Summe von Quadraten”, *Invent. Math.* **4** (1967), 229–237. MR Zbl



- [Pfister 1971] A. Pfister, "Sums of squares in real function fields", pp. 297–300 in *Actes du Congrès International des Mathématiciens, Tome 1* (Nice, 1970), Gauthier-Villars, Paris, 1971. MR Zbl
- [Pfister 1995] A. Pfister, *Quadratic forms with applications to algebraic geometry and topology*, London Mathematical Society Lecture Note Series **217**, Cambridge University Press, 1995. MR Zbl
- [Pfister and Scheiderer 2012] A. Pfister and C. Scheiderer, "An elementary proof of Hilbert's theorem on ternary quartics", *J. Algebra* **371** (2012), 1–25. MR Zbl
- [Scheiderer 1994] C. Scheiderer, *Real and étale cohomology*, Lecture Notes in Mathematics **1588**, Springer, Berlin, 1994. MR Zbl
- [Serre 1965] J.-P. Serre, "Sur la dimension cohomologique des groupes profinis", *Topology* **3** (1965), 413–420. MR Zbl
- [SGA 4<sub>1/2</sub> 1977] P. Deligne, *Cohomologie étale* (Séminaire de Géométrie Algébrique du Bois Marie), Lecture Notes in Mathematics **569**, Springer, Berlin, 1977. MR Zbl
- [SGA 4<sub>2</sub> 1972] M. Artin, A. Grothendieck, and J. L. Verdier, *Théorie des topos et cohomologie étale des schémas, Tome 2: Exposés V–VIII* (Séminaire de Géométrie Algébrique du Bois Marie 1963–1964), Lecture Notes in Mathematics **270**, Springer, Berlin, 1972. MR Zbl
- [SGA 4<sub>3</sub> 1973] M. Artin, A. Grothendieck, and J. L. Verdier, *Théorie des topos et cohomologie étale des schémas, Tome 3: Exposés IX–XIX* (Séminaire de Géométrie Algébrique du Bois Marie 1963–1964), Lecture Notes in Mathematics **305**, Springer, Berlin, 1973. MR Zbl
- [SGA 7<sub>II</sub> 1973] P. Deligne and N. Katz, *Groupes de monodromie en géométrie algébrique, II: Exposés X–XXII* (Séminaire de Géométrie Algébrique du Bois Marie 1967–1969), Lecture Notes in Mathematics **340**, Springer, Berlin, 1973. MR Zbl
- [Voevodsky 2003] V. Voevodsky, "Motivic cohomology with  $\mathbb{Z}/2$ -coefficients", *Publ. Math. Inst. Hautes Études Sci.* **98** (2003), 59–104. MR Zbl

Communicated by Jean-Louis Colliot-Thélène

Received 2016-07-11

Revised 2017-01-05

Accepted 2017-02-03

olivier.benoist@unistra.fr

*Institut de Recherche Mathématique Avancée,  
UMR 7501, Université de Strasbourg et CNRS,  
7 rue René Descartes, 67000 Strasbourg, France*



# Algebra & Number Theory

msp.org/ant

## EDITORS

### MANAGING EDITOR

Bjorn Poonen  
Massachusetts Institute of Technology  
Cambridge, USA

### EDITORIAL BOARD CHAIR

David Eisenbud  
University of California  
Berkeley, USA

## BOARD OF EDITORS

Dave Benson	University of Aberdeen, Scotland	Shigefumi Mori	RIMS, Kyoto University, Japan
Richard E. Borcherds	University of California, Berkeley, USA	Martin Olsson	University of California, Berkeley, USA
J-L. Colliot-Thélène	CNRS, Université Paris-Sud, France	Raman Parimala	Emory University, USA
Brian D. Conrad	Stanford University, USA	Jonathan Pila	University of Oxford, UK
Samit Dasgupta	University of California, Santa Cruz, USA	Anand Pillay	University of Notre Dame, USA
Hélène Esnault	Freie Universität Berlin, Germany	Michael Rapoport	Universität Bonn, Germany
Gavril Farkas	Humboldt Universität zu Berlin, Germany	Victor Reiner	University of Minnesota, USA
Hubert Flenner	Ruhr-Universität, Germany	Peter Sarnak	Princeton University, USA
Sergey Fomin	University of Michigan, USA	Joseph H. Silverman	Brown University, USA
Edward Frenkel	University of California, Berkeley, USA	Michael Singer	North Carolina State University, USA
Andrew Granville	Université de Montréal, Canada	Christopher Skinner	Princeton University, USA
Joseph Gubeladze	San Francisco State University, USA	Vasudevan Srinivas	Tata Inst. of Fund. Research, India
Roger Heath-Brown	Oxford University, UK	J. Toby Stafford	University of Michigan, USA
Craig Huneke	University of Virginia, USA	Pham Huu Tiep	University of Arizona, USA
Kiran S. Kedlaya	Univ. of California, San Diego, USA	Ravi Vakil	Stanford University, USA
János Kollár	Princeton University, USA	Michel van den Bergh	Hasselt University, Belgium
Yuri Manin	Northwestern University, USA	Marie-France Vignéras	Université Paris VII, France
Philippe Michel	École Polytechnique Fédérale de Lausanne	Kei-Ichi Watanabe	Nihon University, Japan
Susan Montgomery	University of Southern California, USA	Shou-Wu Zhang	Princeton University, USA

## PRODUCTION

production@msp.org

Silvio Levy, Scientific Editor

---

See inside back cover or [msp.org/ant](http://msp.org/ant) for submission instructions.

---

The subscription price for 2017 is US \$325/year for the electronic version, and \$520/year (+\$55, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to MSP.


---

Algebra & Number Theory (ISSN 1944-7833 electronic, 1937-0652 printed) at Mathematical Sciences Publishers, 798 Evans Hall #3840, c/o University of California, Berkeley, CA 94720-3840 is published continuously online. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices.

---

ANT peer review and production are managed by EditFLOW<sup>®</sup> from MSP.

PUBLISHED BY

 **mathematical sciences publishers**  
nonprofit scientific publishing

<http://msp.org/>

© 2017 Mathematical Sciences Publishers

# Algebra & Number Theory

Volume 11      No. 4      2017

---

Cup products of line bundles on homogeneous varieties and generalized PRV components of multiplicity one IVAN DIMITROV and MIKE ROTH	767
Mass formulas for local Galois representations and quotient singularities II: Dualities and resolution of singularities MELANIE MATCHETT WOOD and TAKEHIKO YASUDA	817
A new equivariant in nonarchimedean dynamics ROBERT RUMELY	841
On pairs of $p$ -adic $L$ -functions for weight-two modular forms FLORIAN SPRUNG	885
On Hilbert's 17th problem in low degree OLIVIER BENOIST	929
Gowers norms of multiplicative functions in progressions on average XUANCHENG SHAO	961
The degree of the Gauss map of the theta divisor GIULIO CODOGNI, SAMUEL GRUSHEVSKY and EDOARDO SERNESI	983
Correction to the article <i>Frobenius and valuation rings</i> ( <b>10:5</b> (2016), 1057–1090) RANKEYA DATTA and KAREN E. SMITH	1003



1937-0652(2017)11:4;1-H