

Algebra & Number Theory

Volume 11

2017

No. 4

**Gowers norms of multiplicative functions
in progressions on average**

Xuancheng Shao



Gowers norms of multiplicative functions in progressions on average

Xuancheng Shao

Let μ be the Möbius function and let $k \geq 1$. We prove that the Gowers U^k -norm of μ restricted to progressions $\{n \leq X : n \equiv a_q \pmod{q}\}$ is $o(1)$ on average over $q \leq X^{1/2-\sigma}$ for any $\sigma > 0$, where $a_q \pmod{q}$ is an arbitrary residue class with $(a_q, q) = 1$. This generalizes the Bombieri–Vinogradov inequality for μ , which corresponds to the special case $k = 1$.

1. Introduction

A basic problem in analytic number theory is to understand the distribution of primes, or other related arithmetic functions such as the Möbius function μ and the Liouville function λ , in arithmetic progressions when the modulus is relatively large. In this direction, the Bombieri–Vinogradov inequality leads us almost half way to the ultimate goal, if we average over the moduli.

Theorem (Bombieri–Vinogradov). *Let $X, Q \geq 2$, and let $A \geq 2$. Assume that $Q \leq X^{1/2}(\log X)^{-B}$ for some sufficiently large $B = B(A)$. Then for all but at most $Q(\log X)^{-A}$ moduli $q \leq Q$, we have*

$$\sup_{(a,q)=1} \left| \sum_{\substack{n \leq X \\ n \equiv a \pmod{q}}} \Lambda(n) - \frac{1}{\varphi(q)} \sum_{n \leq X} \Lambda(n) \right| \ll_A \frac{X}{Q(\log X)^A}.$$

The same statement holds for the Möbius function μ and the Liouville function λ .

See [Iwaniec and Kowalski 2004, Chapter 17] for its proof and applications. In this paper, we investigate a higher order generalization of the Bombieri–Vinogradov inequality, which measures more refined distributional properties. This higher order version involves Gowers norms, a central tool in additive combinatorics. We refer the readers to [Tao and Vu 2006, Chapter 11] for the basic definitions and applications. In particular, $\|f\|_{U^k(Y)}$ stands for the U^k -norm of the function f on the interval $[0, Y] \cap \mathbb{Z}$.

Shao was supported by a Glasstone Research Fellowship.

MSC2010: primary 11P32; secondary 11B30, 11N13.

Keywords: multiplicative functions, Bombieri–Vinogradov theorem, Gowers norms.

For any arithmetic function $f : \mathbb{Z} \rightarrow \mathbb{C}$ and any residue class $a \pmod{q}$, denote by $f(q \cdot + a)$ the function $m \mapsto f(qm + a)$. Precisely we study the Gowers U^k -norm of f restricted to progressions $\{n \leq X : n \equiv a \pmod{q}\}$, i.e., the U^k -norm of the functions $f(q \cdot + a)$ on $[0, X/q] \cap \mathbb{Z}$.

Corollary 1.1. *Let $X, Q \geq 2$, let k be a positive integer, let $A \geq 2$, and let $\varepsilon > 0$. Assume that $Q \leq X^{1/2}(\log X)^{-B}$ for some sufficiently large $B = B(k, A, \varepsilon)$. Then for all but at most $Q(\log X)^{-A}$ moduli $q \leq Q$, we have*

$$\sup_{\substack{0 \leq a < q \\ (a, q) = 1}} \|\mu(q \cdot + a)\|_{U^k(X/q)} \leq \varepsilon.$$

The same statement holds for the Liouville function λ .

The Bombieri–Vinogradov inequality is the $k = 1$ case of [Corollary 1.1](#) (qualitatively), since the U^1 -norm of a function is the same as the absolute value of its average. By the inverse theorem for Gowers norms [[Green et al. 2012](#)], [Corollary 1.1](#) is a straightforward consequence of the following result.

Theorem 1.2. *Let $X, Q \geq 2$ be parameters with $10Q^2 \leq X$. Associated to each $Q \leq q < 2Q$ we have*

- (1) a residue class $a_q \pmod{q}$ with $0 \leq a_q < q, (a_q, q) = 1$;
- (2) a nilmanifold G_q/Γ_q of dimension at most some $d \geq 1$, equipped with a filtration $(G_q)_\bullet$ of degree at most some $s \geq 1$ and a $(\log X)$ -rational Malcev basis \mathcal{X}_q ;
- (3) a polynomial sequence $g_q : \mathbb{Z} \rightarrow G_q$ adapted to $(G_q)_\bullet$;
- (4) a Lipschitz function $\varphi_q : G_q/\Gamma_q \rightarrow \mathbb{C}$ with $\|\varphi_q\|_{\text{Lip}(\mathcal{X}_q)} \leq 1$.

Let $\psi_q : \mathbb{Z} \rightarrow \mathbb{C}$ be the function defined by $\psi_q(n) = \varphi_q(g_q(n)\Gamma_q)$. Then for any $A \geq 2$, the bound

$$\left| \sum_{\substack{n \leq X \\ n \equiv a_q \pmod{q}}} \mu(n) \psi_q\left(\frac{n - a_q}{q}\right) \right| \ll_{A, d, s} \frac{X}{Q} \cdot \frac{\log \log X}{\log(X/Q^2)} \tag{1-1}$$

holds for all but at most $Q(\log X)^{-A}$ moduli $Q \leq q < 2Q$. The same statement holds for the Liouville function λ .

See [[Green and Tao 2012a](#)] for the precise definitions of nilmanifolds and the associated data appearing in the statement. To avoid confusions later on, we point out that the Lipschitz norm is defined by

$$\|\varphi_q\|_{\text{Lip}(\mathcal{X}_q)} = \|\varphi_q\|_\infty + \sup_{x \neq y} \frac{|\varphi_q(x) - \varphi_q(y)|}{d(x, y)},$$

where $d(\cdot, \cdot)$ is the metric induced by \mathcal{X}_q . In particular $\|\varphi_q\|_\infty \leq \|\varphi_q\|_{\text{Lip}(\mathcal{X}_q)}$.

To understand this paper, however, it is not essential to know these definitions, as long as one is willing to accept certain results about nilsequences as black boxes, many of which can be found in [Green and Tao 2012a]. The readers are thus encouraged to consider the following special case when the nilmanifolds are the torus \mathbb{R}/\mathbb{Z} , the polynomial sequences are genuine polynomials of degree at most s , and the Lipschitz functions are $\varphi(x) = e(x) = e^{2\pi i x}$.

Theorem (main theorem, special case). *Let $X, Q \geq 2$ be parameters with $10Q^2 \leq X$, and let $s \geq 1$. Then for any $A \geq 2$, the bound*

$$\sup_{\substack{0 \leq a < q \\ (a, q) = 1}} \sup_{\alpha_1, \dots, \alpha_s \in \mathbb{R}} \left| \sum_{\substack{n \leq X \\ n \equiv a \pmod{q}}} \mu(n) e(\alpha_s n^s + \dots + \alpha_1 n) \right| \ll_{A, s} \frac{X}{Q} \cdot \frac{\log \log X}{\log(X/Q^2)}$$

holds for all but at most $Q(\log X)^{-A}$ moduli $Q \leq q < 2Q$. The same statement holds for the Liouville function λ .

Without restricting to arithmetic progressions (i.e., when $Q = O(1)$), the disconnection between the Möbius function and nilsequences was studied by Green and Tao [2012b], as part of their program to count the number of solutions to linear equations in prime variables.

The rest of the paper is organized as follows. In Section 2 we reduce Theorem 1.2 to the minor arc case (Proposition 2.1). This reduction process is summarized in Lemma 2.4, using a factorization theorem for nilsequences [Green and Tao 2012a, Theorem 1.19]. In fact, one can obtain analogues of Theorem 1.2 for all 1-bounded multiplicative functions satisfying the Bombieri–Vinogradov estimate, such as indicator functions of smooth numbers (see [Fouvry and Tenenbaum 1996; Harper 2012] and the references therein). See [Frantzikinakis and Host 2017] for a previous work on Gowers norms of multiplicative functions, and also [Matthiesen 2016] for a generalization to some not necessarily bounded multiplicative functions. However, we will not seek for such generality here since any such result can be easily deduced from Lemma 2.4 and Proposition 2.1 as needed.

The rest of the argument applies to all bounded multiplicative functions. In Section 3 we consider the minor arc case using an orthogonality criterion. The idea, going back to Montgomery and Vaughan [1977] and Kátai [1986] (see also [Bourgain et al. 2013; Harper 2011]), is that one can make do with type-II estimates (or bilinear estimates) in a very restricted range when dealing with bounded multiplicative functions. This is the reason that we are unable to prove Theorem 1.2 for the primes, which would require type-II estimates in an inaccessible range, and also the reason that one saves no more than $\log X$ in the bound (1-1). In fact, to get this saving we use a quantitatively superior argument of Ramaré [2009], which received a lot of attention recently [Matomäki et al. 2015; Green 2016] following its use in

Matomäki and Radziwiłł’s recent breakthrough [2016]. Finally the required type-II estimates will be proved in Section 4.

2. Technical reductions

In this section, we reduce Theorem 1.2 to the following minor arc, or equidistributed, case. See [Green and Tao 2012a, Definition 1.2] for the precise definition about equidistribution of nilsequences.

Proposition 2.1. *Let $X, Q \geq 2$ be parameters with $10Q^2 \leq X$. Let $\eta \in (0, \frac{1}{2})$. Let $\mathcal{Q} \subset [Q, 2Q)$ be an arbitrary subset. Associated to each $q \in \mathcal{Q}$ we have*

- (1) *a residue class $a_q \pmod{q}$ with $0 \leq a_q < q$, $(a_q, q) = 1$, and an arbitrary interval $I_q \subset [0, X]$;*
- (2) *a nilmanifold G_q/Γ_q of dimension at most some $d \geq 1$, equipped with a filtration $(G_q)_\bullet$ of degree at most some $s \geq 1$ and an η^{-c} -rational Malcev basis \mathcal{X}_q for some sufficiently small $c = c(d, s) > 0$;*
- (3) *a polynomial sequence $g_q: \mathbb{Z} \rightarrow G_q$ adapted to $(G_q)_\bullet$ such that $\{g_q(m)\}_{1 \leq m \leq X/q}$ is totally η -equidistributed;*
- (4) *a Lipschitz function $\varphi_q: G_q/\Gamma_q \rightarrow \mathbb{C}$ with $\|\varphi_q\|_\infty \leq 1$ and $\int \varphi_q = 0$.*

Let $\psi_q: \mathbb{Z} \rightarrow \mathbb{C}$ be the function defined by $\psi_q(n) = \varphi_q(g_q(n)\Gamma_q)$. Let $f: \mathbb{Z} \rightarrow \mathbb{C}$ be a multiplicative function with $|f(n)| \leq 1$. Then

$$\sum_{q \in \mathcal{Q}} \left| \sum_{\substack{n \in I_q \\ n \equiv a_q \pmod{q}}} f(n) \psi_q\left(\frac{n - a_q}{q}\right) \right| \ll \frac{\log \eta^{-1}}{\log(X/Q^2)} \sum_{q \in \mathcal{Q}} \left(\frac{|I_q|}{q} + 1\right) + \eta^c X \log X \max_{q \in \mathcal{Q}} \|\varphi_q\|_{\text{Lip}(\mathcal{X}_q)},$$

for some constant $c = c(d, s) > 0$.

Thus one obtains a saving of (at most) $\log \eta^{-1} / \log(X/Q^2)$ compared to the trivial bound. The attentive reader may notice an extra factor $\log X$ in the second term of the bound, which prevents one from taking any $\eta = o(1)$ and still getting a nontrivial estimate. This extra factor mainly comes from the type-II estimate (Lemma 3.3); see the comments after its statement. It won’t be a concern for us since we will take η to be a large negative power of $\log X$.

To deduce Theorem 1.2 from Proposition 2.1, we may assume that A is sufficiently large depending on d and s , that X is sufficiently large depending on A, d and s , and that $Q \leq X^{1/2}(\log X)^{-B}$ for some sufficiently large $B = B(A)$, since otherwise the bound (1-1) is trivial. In particular, it suffices to establish the bound $O_A(Q(\log X)^{-2A})$ for the number of exceptional moduli.

Reducing to completely multiplicative functions. The first technical step of the reduction is to pass from the Möbius function μ to its completely multiplicative cousin λ . In this subsection we deduce [Theorem 1.2](#) for μ , assuming that it has already been proved for the Liouville function λ . This step is summarized in the following lemma.

Lemma 2.2. *Let $X \geq 2$ be large, and let $a \pmod{q}$ be a residue class with $(a, q) = 1$. Let $\varepsilon \in (0, 1)$, and assume that $q \leq \varepsilon X^{1/2}(\log X)^{-3}$. Let $f : \mathbb{Z} \rightarrow \mathbb{C}$ be a multiplicative function with $|f(n)| \leq 1$, and let $f' : \mathbb{Z} \rightarrow \mathbb{C}$ be the completely multiplicative function defined by $f'(p) = f(p)$ for each prime p . Let $c : \mathbb{Z} \rightarrow \mathbb{C}$ be an arbitrary function with $|c(n)| \leq 1$. If*

$$\left| \sum_{\substack{n \leq X \\ n \equiv a \pmod{q}}} f(n)c(n) \right| \geq \varepsilon \frac{X}{q},$$

then there is a positive integer $\ell \ll \varepsilon^{-3}$ with $(\ell, q) = 1$, such that

$$\left| \sum_{\substack{n \leq X/\ell \\ n \equiv a\ell^{-1} \pmod{q}}} f'(n)c(\ell n) \right| \gg \varepsilon \frac{X}{\ell q}.$$

To deduce [Theorem 1.2](#) for μ , apply [Lemma 2.2](#) with $f = \mu$ (so that $f' = \lambda$) and $\varepsilon = C \log \log X / \log(X/Q^2)$ for some large constant C depending on A . For each q satisfying

$$\left| \sum_{\substack{n \leq X \\ n \equiv a_q \pmod{q}}} \mu(n) \psi_q \left(\frac{n - a_q}{q} \right) \right| \geq \varepsilon \frac{X}{q}, \quad (2-1)$$

[Lemma 2.2](#) produces a positive integer $\ell = \ell_q \ll \varepsilon^{-3}$ with $(\ell_q, q) = 1$, such that

$$\left| \sum_{\substack{n \leq X/\ell_q \\ n \equiv a_q \ell_q^{-1} \pmod{q}}} \lambda(n) \psi_q \left(\frac{\ell_q n - a_q}{q} \right) \right| \gg \varepsilon \frac{X}{\ell_q q}.$$

For each $\ell \ll \varepsilon^{-3}$, apply [Theorem 1.2](#) for λ , with X replaced by X/ℓ , to conclude that there are at most $Q(\log X)^{-3A}$ moduli q satisfying (2-1) with $\ell_q = \ell$. It follows that the total number of moduli q satisfying (2-1) is $O(\varepsilon^{-3} Q(\log X)^{-3A}) = O(Q(\log X)^{-2A})$, as desired.

In the remainder of this subsection, we give a standard proof of [Lemma 2.2](#), starting with a basic lemma.

Lemma 2.3. *Let $f : \mathbb{Z} \rightarrow \mathbb{C}$ be a multiplicative function with $|f(n)| \leq 1$, and let $f' : \mathbb{Z} \rightarrow \mathbb{C}$ be the completely multiplicative function defined by $f'(p) = f(p)$ for*

each prime p . Let g be the multiplicative function with $f = f' * g$. Then for any $N \geq 2$ we have

$$\sum_{n \geq N} \frac{|g(n)|}{n} \ll N^{-1/2}(\log N)^2 \quad \text{and} \quad \sum_{n \leq N} |g(n)| \ll N^{1/2}(\log N)^2.$$

Proof. It is easy to see that $g(p) = 0$ and $|g(p^k)| \leq 2$ for every prime p . Set $\sigma = \frac{1}{2} + 1/(10 \log N)$ so that $\sigma \in (\frac{1}{2}, 1)$ and $N^\sigma \asymp N^{1/2}$. By Rankin's trick we have

$$\sum_{n \geq N} \frac{|g(n)|}{n} \leq \sum_n \frac{|g(n)|}{n} \left(\frac{n}{N}\right)^{1-\sigma} \ll N^{-1/2} \sum_n |g(n)|n^{-\sigma},$$

and similarly

$$\sum_{n \leq N} |g(n)| \leq \sum_n |g(n)| \left(\frac{N}{n}\right)^\sigma \ll N^{1/2} \sum_n |g(n)|n^{-\sigma}.$$

Thus it suffices to establish the bound

$$\sum_n |g(n)|n^{-\sigma} \ll (\log N)^2.$$

We may write the Dirichlet series associated to $|g|$ in terms of its Euler product:

$$\sum_n |g(n)|n^{-\sigma} = \prod_p (1 + |g(p^2)|p^{-2\sigma} + |g(p^3)|p^{-3\sigma} + \dots).$$

Since $|g(p^k)| \leq 2$, we may bound it by

$$\prod_p (1 + p^{-2\sigma} + p^{-4\sigma} + \dots)^2 (1 + p^{-3\sigma} + p^{-6\sigma} + \dots)^2 = \zeta(2\sigma)^2 \zeta(3\sigma)^2.$$

Since $\zeta(3\sigma) \ll 1$ and $\zeta(2\sigma) \ll (2\sigma - 1)^{-1}$, the desired bound follows immediately. \square

Proof of Lemma 2.2. Write $f = f' * g$ for some multiplicative function g . We have

$$\sum_{\substack{n \leq X \\ n \equiv a \pmod{q}}} f(n)c(n) = \sum_{\substack{\ell n \leq X \\ \ell n \equiv a \pmod{q}}} f'(\ell)g(\ell)c(\ell n) = \sum_{\substack{\ell \leq X \\ (\ell, q)=1}} g(\ell) \left(\sum_{\substack{n \leq X/\ell \\ n \equiv a\ell^{-1} \pmod{q}}} f'(n)c(\ell n) \right).$$

Let $L \geq 2$ be a parameter. Using the trivial bound $O(X/\ell q + 1)$ for the inner sum, we may apply Lemma 2.3 to bound the total contributions from those terms with $\ell \geq L$ by

$$\frac{X}{q} \sum_{\ell \geq L} \frac{|g(\ell)|}{\ell} + \sum_{\ell \leq X} |g(\ell)| \ll \frac{X}{qL^{1/2}} (\log L)^2 + X^{1/2} (\log X)^2.$$

We may choose $L \ll \varepsilon^{-3}$ such that the first term above is negligible compared to the lower bound $\varepsilon X/q$, and the second term is already negligible compared to $\varepsilon X/q$ by the assumption on q . It follows that

$$\sum_{\substack{\ell \leq L \\ (\ell, q) = 1}} |g(\ell)| \left| \sum_{\substack{n \leq X/\ell \\ n \equiv a\ell^{-1} \pmod{q}}} f'(n)c(\ell n) \right| \gg \varepsilon \frac{X}{q}.$$

Since $\sum |g(\ell)|\ell^{-1} \ll 1$, there is some $\ell \leq L$ with $(\ell, q) = 1$ such that

$$\left| \sum_{\substack{n \leq X/\ell \\ n \equiv a\ell^{-1} \pmod{q}}} f'(n)c(\ell n) \right| \gg \varepsilon \frac{X}{\ell q}.$$

This completes the proof of the lemma. □

Reducing to equidistributed nilsequences. We now use the factorization theorem [Green and Tao 2012a, Theorem 1.19] to reduce arbitrary nilsequences to equidistributed ones. This step is summarized in the following lemma, the proof of which is similar to arguments in [Green and Tao 2012b, Section 2].

Lemma 2.4. *Let $X \geq 2$ be large, and let $a \pmod{q}$ be a residue class with $0 \leq a < q$, $(a, q) = 1$. Let $\varepsilon \in (0, \frac{1}{2})$. Let $f : \mathbb{Z} \rightarrow \mathbb{C}$ be a completely multiplicative function with $|f(n)| \leq 1$. Given*

- a nilmanifold G/Γ of dimension at most some $d \geq 1$, equipped with a filtration G_\bullet of degree at most some $s \geq 1$ and a M_0 -rational Malcev basis \mathcal{X} for some $M_0 \geq \varepsilon^{-1}$;
- a polynomial sequence $g : \mathbb{Z} \rightarrow G$ adapted to G_\bullet ; and
- a Lipschitz function $\varphi : G/\Gamma \rightarrow \mathbb{C}$ with $\|\varphi\|_{\text{Lip}(\mathcal{X})} \leq 1$,

let $\psi : \mathbb{Z} \rightarrow \mathbb{C}$ be the function defined by $\psi(n) = \varphi(g(n)\Gamma)$. Assume that

$$\left| \sum_{\substack{n \leq X \\ n \equiv a \pmod{q}}} f(n)\psi\left(\frac{n-a}{q}\right) \right| \geq \varepsilon \frac{X}{q}.$$

For any $A \geq 2$ large enough depending on d and s , we may find $M_0 \leq M \leq M_0^{O_A(1)}$, an interval $I \subset [0, X]$ with $|I| \gg X/M^3$, a positive integer q' with $q \mid q'$ and $q' \leq qM$, a residue class $a' \pmod{q'}$ with $0 \leq a' < q'$, $(a', q') = 1$, and moreover

- a nilmanifold G'/Γ' of dimension at most d , equipped with a filtration G'_\bullet of degree at most s and a $M^{O_{d,s}(1)}$ -rational Malcev basis \mathcal{X}' ;
- a polynomial sequence $g' : \mathbb{Z} \rightarrow G'$ adapted to G'_\bullet such that $\{g'(m)\}_{1 \leq m \leq X/q}$ is totally M^{-A} -equidistributed; and

• a Lipschitz function $\varphi' : G' / \Gamma' \rightarrow \mathbb{C}$ with $\|\varphi'\|_\infty \leq 1$ and $\|\varphi'\|_{\text{Lip}(\mathcal{X}')} \leq M^{O_{d,s}(1)}$, such that

$$\left| \sum_{\substack{n \in I \\ n \equiv a' \pmod{q'}}} f(n) \psi' \left(\frac{n - a'}{q'} \right) \right| \gg \varepsilon \frac{|I|}{q'}$$

where $\psi' : \mathbb{Z} \rightarrow \mathbb{C}$ is the function defined by $\psi'(n) = \varphi'(g'(n)\Gamma')$.

To deduce [Theorem 1.2](#) for λ from [Proposition 2.1](#), apply [Lemma 2.4](#) with $f = \lambda$, $\varepsilon = C \log \log X / \log(X/Q^2)$ for some large constant C depending on A , and $M_0 = (\log X)^C$ for some large constant C depending on d and s . For each q satisfying

$$\left| \sum_{\substack{n \leq X \\ n \equiv a_q \pmod{q}}} \lambda(n) \psi_q \left(\frac{n - a_q}{q} \right) \right| \geq \varepsilon \frac{X}{q}, \tag{2-2}$$

[Lemma 2.4](#) produces $M_q, I_q, a' \pmod{q'}$, G' / Γ' , and $\psi' = \varphi' \circ g'$, all of which depend on q (some of these dependencies are suppressed for notational convenience), such that

$$\left| \sum_{\substack{n \in I_q \\ n \equiv a' \pmod{q'}}} \lambda(n) \psi' \left(\frac{n - a'}{q'} \right) \right| \gg \varepsilon \frac{|I_q|}{q'}$$

Divide the possible values of M_q into $O_A(1)$ subintervals of the form $[M^{1/2}, M]$ with $M_0 \leq M \leq M_0^{O_A(1)}$. Given M , let $\mathcal{Q} = \mathcal{Q}_M$ be the set of moduli $q \in \mathcal{Q}$ such that $M^{1/2} \leq M_q \leq M$, and let $\mathcal{Q}' = \mathcal{Q}'_M$ be the set of q' arising from $q \in \mathcal{Q}$. It suffices to show that

$$|\mathcal{Q}| \ll Q(\log X)^{-2A}.$$

Since q'/q is a positive integer at most M , each q' occurs with multiplicity at most M . Thus $|\mathcal{Q}| \leq M|\mathcal{Q}'|$. Before applying [Proposition 2.1](#) we need to ensure that each φ' has average 0. For $q' \in \mathcal{Q}'$ either

$$\left| \sum_{\substack{n \in I_q \\ n \equiv a' \pmod{q'}}} \lambda(n) \right| \gg \varepsilon \frac{|I_q|}{q'} \gg \frac{X}{q'(\log X)^{O_A(1)}} \tag{2-3}$$

or

$$\left| \sum_{\substack{n \in I_q \\ n \equiv a' \pmod{q'}}} \lambda(n) \left(\psi' \left(\frac{n - a'}{q'} \right) - \int \varphi' \right) \right| \gg \varepsilon \frac{|I_q|}{q'}. \tag{2-4}$$

To bound the number of q' satisfying (2-3), note that the Bombieri–Vinogradov inequality (for λ) is applicable since $q' \leq 2QM \leq X^{1/2}(\log X)^{-B/2}$ (recall the assumption that $Q \leq X^{1/2}(\log X)^{-B}$ for some large B). By choosing b large enough, we may ensure that the number of q' satisfying (2-3) is at most QM^{-A} .

Now let $\mathcal{Q}'_2 \subset \mathcal{Q}'$ be the set of $q' \in \mathcal{Q}'$ satisfying (2-4). To bound the size of \mathcal{Q}'_2 , we apply Proposition 2.1 after replacing each φ' by $\varphi' - \int \varphi'$ and dyadically dividing the possible values of q' . This leads to

$$\varepsilon \sum_{q' \in \mathcal{Q}'_2} \frac{|I_{q'}|}{q'} \ll \frac{\log M^A}{\log(X/Q^2 M^2)} \sum_{q' \in \mathcal{Q}'_2} \left(\frac{|I_{q'}|}{q'} + 1 \right) + M^{-cA} X (\log X)^2 M^{O_{d,s}(1)},$$

for some $c = c(d, s) > 0$. The first term on the right can be made negligible compared to the left-hand side, if the constant C in the choice of ε is taken large enough in terms of A . Hence

$$\varepsilon \frac{X}{M^3} \cdot \frac{|\mathcal{Q}'_2|}{QM} \ll \varepsilon \sum_{q' \in \mathcal{Q}'} \frac{|I_{q'}|}{q'} \ll M^{-cA + O_{d,s}(1)} X.$$

It follows that $|\mathcal{Q}'_2| \ll QM^{-cA + O_{d,s}(1)}$. Combining the estimates for the two types of q' together, we obtain

$$|\mathcal{Q}| \leq M|\mathcal{Q}'| \ll QM^{-cA + O_{d,s}(1)} \ll Q(\log X)^{-2A},$$

if the constant C in the choice of M is large enough depending on d and s . This completes the deduction of Theorem 1.2.

Proof of Lemma 2.4. Let C be a large constant (depending on d and s), and apply the factorization theorem [Green and Tao 2012a, Theorem 1.19] to find $CM_0 \leq M \leq M_0^{O_A(1)}$, a rational subgroup $\tilde{G} \subset G$, a Malcev basis $\tilde{\mathcal{X}}$ for $\tilde{G}/\tilde{\Gamma}$ (where $\tilde{\Gamma} = \Gamma \cap \tilde{G}$) in which each element is an M -rational combination of the elements of \mathcal{X} , and a decomposition $g = s\tilde{g}\gamma$ into polynomial sequences $s, \tilde{g}, \gamma : \mathbb{Z} \rightarrow G$ with the following properties:

- (1) s is $(M, X/q)$ -smooth in the sense that $d(s(n), \text{id}) \leq M$ and $d(s(n), s(n-1)) \leq qM/X$ for each $1 \leq n \leq X/q$.
- (2) \tilde{g} takes values in \tilde{G} ; moreover $\{\tilde{g}(n)\}_{1 \leq n \leq X/q}$ is totally M^{-CA} -equidistributed in $\tilde{G}/\tilde{\Gamma}$ (using the metric induced by the Malcev basis $\tilde{\mathcal{X}}$).
- (3) γ is M -rational in the sense that for each $n \in \mathbb{Z}$, $\gamma(n)^r \in \Gamma$ for some $1 \leq r \leq M$. Moreover, γ is periodic with period $t \leq M$.

We may assume that $X \geq qM^3$, since otherwise the conclusion holds trivially. After a change of variables $n = qm + a$, we may rewrite the assumption as

$$\left| \sum_{m \leq X/q} f(qm + a)\psi(m) \right| \gg \varepsilon \frac{X}{q}.$$

Dividing $[0, X/q]$ into $O(M^2)$ intervals of equal length and then further dividing them into residue classes modulo t , we may find an interval $J \subset [0, X]$ with

$|J| \asymp X/M^2$ and some residue class $b \pmod t$, such that

$$\left| \sum_{\substack{m \equiv b \pmod t \\ qm+a \in J}} f(qm+a)\varphi(s(m)\tilde{g}(m)\gamma(m)) \right| \gg \varepsilon \frac{|J|}{qt}. \tag{2-5}$$

Pick any m_0 counted in the sum (i.e., $m_0 \equiv b \pmod t$ and $qm_0+a \in J$), and note that we may replace $s(m)$ in (2-5) by $s(m_0)$ with a negligible error, since φ has Lipschitz norm at most 1 and

$$d(s(m)\tilde{g}(m)\gamma(m), s(m_0)\tilde{g}(m)\gamma(m)) = d(s(m), s(m_0)) \ll M^{-1},$$

for all m with $qm+a \in J$ by the right invariance of d and the smoothness property of s . Moreover, by the periodicity of γ , we may replace $\gamma(m)$ in (2-5) by $\gamma(m_0)$. Now let g' be the polynomial sequence defined by

$$g'(m) = \gamma(m_0)^{-1}\tilde{g}(tm+b)\gamma(m_0),$$

taking values in $G' = \gamma(m_0)^{-1}\tilde{G}\gamma(m_0)$, and let φ' be the automorphic function on G' defined by

$$\varphi'(x) = \varphi(s(m_0)\gamma(m_0)x).$$

The desired properties about G'/Γ' , g' , and φ' can be established via standard “quantitative nilinear algebra” (see the claim at the end of [Green and Tao 2012b, Section 2]). After a change of variables replacing m by $tm+b$, the inequality (2-5) can be rewritten as

$$\left| \sum_{m: qtm+c \in J} f(qtm+c)\varphi'(g'(m)) \right| \gg \varepsilon \frac{|J|}{qt},$$

where $c = qb+a$. This is almost what we need, but there is the slight issue that c may not be coprime with t . Let $d = (c, t)$ so that $d \leq M$. Let $q' = qt/d$ and $a' = c/d$ so that $(a', q') = 1$. Let $I = d^{-1}J$ so that $|I| \gg X/M^3$. Since f is completely multiplicative, we have

$$\left| \sum_{m: q'm+a' \in I} f(q'm+a')\varphi'(g'(m)) \right| \gg \varepsilon \frac{|I|}{q'}.$$

This completes the proof of the lemma. □

3. The minor arc case: Proof of Proposition 2.1

In this section we prove Proposition 2.1, which is the minor arc case of our main theorem and applies to all 1-bounded multiplicative functions. For convenience write

$$T = \sum_{q \in \mathcal{Q}} \left(\frac{|I_q|}{q} + 1 \right).$$

We may assume that

$$(X/Q^2)^{-1/20} < \eta < (\log X)^{-C},$$

for some sufficiently large $C = C(d, s) > 0$, since otherwise the bound is trivial. We may further assume that $|I_q| \geq X^{0.9}$ for each $q \in \mathcal{Q}$, since the contributions from those q with $|I_q| \leq X^{0.9}$ are trivially acceptable. After multiplying each φ_q by an appropriate scalar, it suffices to prove the desired inequality with the absolute value sign removed. Set

$$Y = \eta^{-1} \quad \text{and} \quad Z = \left(\frac{X}{Q^2}\right)^{1/20},$$

so that $2 \leq Y < Z \leq X^{1/20}$. Let F be the function defined by

$$F(n) = \sum_{\substack{q \in \mathcal{Q}, n \in I_q \\ n \equiv a_q \pmod{q}}} \psi_q \left(\frac{n - a_q}{q} \right).$$

Clearly F is supported on $[0, X]$. The desired bound can be rewritten as

$$\sum_{n \leq X} f(n)F(n) \ll \frac{\log Y}{\log Z} \cdot T + \eta^c X \log X \max_{q \in \mathcal{Q}} \|\varphi_q\|_{\text{Lip}(\mathcal{X}_q)}. \quad (3-1)$$

As alluded to in the introduction, this will be proved using an orthogonality criterion for multiplicative functions. A general principle of this type is given in [Green 2016, Proposition 2.2]. In the notations there, the terms giving rise to E_{triv} and E_{sieve} will be dealt with by Lemmas 3.1 and 3.2, respectively. In particular, the E_{sieve} term leads to the first bound in (3-1). The bilinear (type-II) sum E_{bilinear} will be dealt with in Lemma 3.3, leading to the second bound in (3-1).

Unfortunately we cannot directly apply [Green 2016, Proposition 2.2], since for example our function F is not necessarily bounded. In the remainder of this section we reproduce the argument from [Green 2016] with suitable modifications to prove (3-1). Recall the definition of Ramaré's weight function:

$$w(n) = \frac{1}{\#\{Y \leq p < Z : p | n\} + 1}.$$

Introduce also the function $\mu_{[Y, Z]}^2$, the indicator function of the set of integers n that are not divisible by the square of any prime $p \in [Y, Z]$. To prove (3-1), we first dispose of those terms with $\mu_{[Y, Z]}^2(n) = 0$:

$$\sum_{\substack{n \leq X \\ \mu_{[Y, Z]}^2(n) = 0}} |f(n)F(n)| \leq \sum_{Y \leq p < Z} \sum_{\substack{n \leq X \\ p^2 | n}} |F(n)|.$$

The following lemma will be used repeatedly.

Lemma 3.1. *For any positive integer $D \leq X^{0.4}$ we have*

$$\sum_{\substack{n \leq X \\ D|n}} |F(n)| \ll \frac{T}{D}.$$

Proof. Using the trivial bound

$$|F(n)| \leq \sum_{\substack{q \in \mathcal{Q}, n \in I_q \\ n \equiv a_q \pmod{q}}} 1, \tag{3-2}$$

we obtain

$$\sum_{\substack{n \leq X \\ D|n}} |F(n)| \leq \sum_{q \in \mathcal{Q}} \sum_{\substack{n \in I_q, D|n \\ n \equiv a_q \pmod{q}}} 1.$$

Since $(a_q, q) = 1$, the inner sum over n is nonempty unless $(q, D) = 1$, in which case it is $O(|I_q|/qD)$. The conclusion follows immediately. \square

Since $Z^2 \leq X^{0.4}$, **Lemma 3.1** implies that

$$\sum_{\substack{n \leq X \\ \mu_{[Y,Z]}^2(n)=0}} |f(n)F(n)| \ll T \sum_{Y \leq p < Z} \frac{1}{p^2} \ll \frac{T}{Y}.$$

Hence the contributions from those $n \leq X$ with $\mu_{[Y,Z]}^2(n) = 0$ are acceptable. If $\mu_{[Y,Z]}^2(n) = 1$, then we have the Ramaré identity

$$\sum_{\substack{Y \leq p < Z \\ p|n}} w\left(\frac{n}{p}\right) = \begin{cases} 1 & \text{if } p|n \text{ for some } Y \leq p < Z, \\ 0 & \text{otherwise.} \end{cases}$$

The following lemma disposes of those n not divisible by any $p \in [Y, Z)$:

Lemma 3.2. *We have*

$$\sum_{n \leq X} |F(n)| \cdot \mathbf{1}_{(n, \prod_{Y \leq p < Z} p)=1} \ll \frac{\log Y}{\log Z} \cdot T.$$

Proof. Using (3-2), we can bound the left-hand side by

$$\sum_{q \in \mathcal{Q}} \sum_{\substack{n \in I_q \\ n \equiv a_q \pmod{q}}} \mathbf{1}_{(n, \prod_{Y \leq p < Z} p)=1}.$$

Consider the inner sum for a fixed q . Writing $d = (q, \prod_{Y \leq p < Z} p)$, we may

bound the inner sum using a standard upper bound sieve (since $Z^2 \leq |I_q|/q$) to obtain

$$\sum_{\substack{n \in I_q \\ n \equiv a_q \pmod{q}}} \mathbf{1}_{(n, \prod_{Y \leq p < Z} p) = 1} \ll \frac{|I_q|}{q} \prod_{\substack{Y \leq p < Z \\ p|q}} \left(1 - \frac{1}{p}\right) \ll \frac{|I_q|}{q} \cdot \frac{\log Y}{\log Z} \cdot \frac{d}{\varphi(d)}.$$

On the other hand, since $d \mid \prod_{Y \leq p < Z} p$ and $d \leq q$ we have

$$\frac{d}{\varphi(d)} \leq \prod_{Y \leq p < W} \left(1 - \frac{1}{p}\right)^{-1},$$

where $W \sim Y + \log q$. Thus $d/\varphi(d) = O(1)$ since $Y \geq \log q$, and the conclusion of the lemma follows. \square

Thus we can restrict to those n with $\mu_{[Y, Z]}^2(n) = 1$ and having at least one prime divisor $p \in [Y, Z)$. By the Ramaré identity, we need to estimate

$$\Sigma := \sum_{\substack{n \leq X \\ \mu_{[Y, Z]}^2(n) = 1}} f(n) F(n) \sum_{\substack{Y \leq p < Z \\ p|n}} w\left(\frac{n}{p}\right).$$

Writing $m = n/p$ and using the multiplicativity of f , we obtain

$$\Sigma = \sum_{\substack{m \leq X/Y \\ \mu_{[Y, Z]}^2(m) = 1}} w(m) f(m) \sum_{\substack{Y \leq p < Z \\ p \leq X/m \\ (m, p) = 1}} f(p) F(pm). \quad (3-3)$$

The condition $\mu_{[Y, Z]}^2(m) = 1$ can be dropped since the contribution from those m divisible by \tilde{p}^2 for some $\tilde{p} \in [Y, Z)$ is at most

$$\sum_{Y \leq \tilde{p} < Z} \sum_{Y \leq p < Z} \sum_{\substack{m \leq X/p \\ \tilde{p}^2 | m}} |F(pm)| \ll T \sum_{Y \leq \tilde{p} < Z} \sum_{Y \leq p < Z} \frac{1}{p\tilde{p}^2} \ll \frac{T}{Y} \log \log X,$$

by an application of [Lemma 3.1](#). Similarly, the condition $(m, p) = 1$ in (3-3) can also be dropped since the contribution from the terms with $p \mid m$ is at most

$$\sum_{Y \leq p < Z} \sum_{\substack{m \leq X/p \\ p \mid m}} |F(pm)| \ll T \sum_{Y \leq p < Z} \frac{1}{p^2} \ll \frac{T}{Y},$$

by [Lemma 3.1](#). Both these bounds are acceptable. Thus it remains to bound

$$\Sigma' := \sum_{m \leq X/Y} w(m) f(m) \sum_{\substack{Y \leq p < Z \\ p \leq X/m}} f(p) F(pm).$$

Dyadically dividing the range $[Y, Z)$ for p , we consider

$$\Sigma'(P) := \sum_{m \leq X/P} w(m)f(m) \sum_{\substack{P \leq p < 2P \\ p \leq X/m}} f(p)F(pm)$$

for $P \in [Y, Z)$. Use the trivial bound $|w(m)f(m)| \leq 1$ and apply the Cauchy–Schwarz inequality to obtain

$$|\Sigma'(P)|^2 \ll \frac{X}{P} \sum_{m \leq X/P} \left| \sum_{\substack{P \leq p < 2P \\ p \leq X/m}} f(p)F(pm) \right|^2.$$

After expanding the square and changing the order of summation, we obtain

$$\begin{aligned} |\Sigma'(P)|^2 &\ll \frac{X}{P} \sum_{P \leq p, p' < 2P} f(p)\overline{f(p')} \sum_{m \leq \min(X/p, X/p')} F(pm)\overline{F(p'm)} \\ &\ll \frac{X}{P} \sum_{P \leq p, p' < 2P} \left| \sum_{m \leq \min(X/p, X/p')} F(pm)\overline{F(p'm)} \right|. \end{aligned}$$

Set $K = P$, $L = X/P$, and $\delta = \eta^c$ for some $c > 0$ small enough depending on d and s . The following lemma, whose proof will be given in Section 4, gives the necessary estimates for the type-II (bilinear) sums appearing above.

Lemma 3.3. *Let $K, L, Q \geq 2$ be parameters with $10Q^2 \leq L$. Let $\delta \in (0, \frac{1}{2})$. Associated to each $Q \leq q < 2Q$ we have*

- (1) *a residue class $a_q \pmod q$ with $0 \leq a_q < q$, $(a_q, q) = 1$, and an arbitrary interval I_q ;*
- (2) *a nilmanifold G_q/Γ_q of dimension at most some $d \geq 1$, equipped with a filtration $(G_q)_\bullet$ of degree at most some $s \geq 1$ and a δ^{-1} -rational Malcev basis \mathcal{X}_q ;*
- (3) *a polynomial sequence $g_q : \mathbb{Z} \rightarrow G_q$ adapted to $(G_q)_\bullet$;*
- (4) *a Lipschitz function $\varphi_q : G_q/\Gamma_q \rightarrow \mathbb{C}$ with $\|\varphi_q\|_{\text{Lip}(\mathcal{X}_q)} \leq 1$ and $\int \varphi_q = 0$.*

Let $\psi_q : \mathbb{Z} \rightarrow \mathbb{C}$ be the function defined by $\psi_q(n) = \varphi_q(g_q(n)\Gamma_q)$, and let F be the function defined by

$$F(n) = \sum_{\substack{Q \leq q < 2Q \\ n \in I_q \\ n \equiv a_q \pmod q}} \psi_q\left(\frac{n - a_q}{q}\right).$$

For each $k, k' \in [K, 2K)$, let $I(k, k') \subset [0, L]$ be an arbitrary interval. Suppose that

$$\sum_{K \leq k, k' < 2K} \left| \sum_{\ell \in I(k, k')} F(k\ell)\overline{F(k'\ell)} \right| \geq \delta K^2 L, \tag{3-4}$$

and that $K^{-c} < \delta < (\log Q)^{-1}$ for some sufficiently small $c = c(d, s) > 0$. Then the polynomial sequence $\{g_q(m)\}_{1 \leq m \leq KL/Q}$ fails to be totally $\delta^{O_{d,s}(1)}$ -equidistributed for some $Q \leq q < 2Q$.

To complete the proof of [Proposition 2.1](#), note that the hypotheses $10Q^2 \leq L$ and $K^{-c} < \delta < (\log Q)^{-1}$ in [Lemma 3.3](#) are satisfied by our choices of Y and Z . By setting $\varphi_q = 0$ for $q \notin \mathcal{Q}$ and renormalizing (replacing φ_q by $\varphi_q / \|\varphi_q\|_{\text{Lip}(\mathcal{X}_q)}$), we may apply [Lemma 3.3](#) to conclude that

$$|\Sigma'(P)|^2 \ll \frac{X}{P} \cdot \eta^c P X \max_{q \in \mathcal{Q}} \|\varphi_q\|_{\text{Lip}(\mathcal{X}_q)}^2.$$

The desired bound [\(3-1\)](#) follows after summing over P dyadically.

Remark 3.4. Instead of using simply the trivial bound $|w(n)| \leq 1$, one may appeal to [\[Green 2016, Lemma 2.1\]](#) to dispose of the extra $\log X$ factor that appeared when summing over P dyadically. We will, however, not bother with this since the type-II estimates we use already have an extra logarithmic factor anyways.

4. Type-II estimates

In this section we prove [Lemma 3.3](#). We start with the following lemma, needed to treat composite moduli.

Lemma 4.1. *Let $Q \geq 2$ and $Q \leq R \leq 4Q^2$. Let E be the set of pairs (q, q') with $Q \leq q, q' < 2Q$ and $R \leq [q, q'] < 2R$. For each $Q \leq q < 2Q$ and $R \leq r < 2R$, let*

$$m_q(r) = \#\{Q \leq q' < 2Q : (q, q') \in E, [q, q'] = r\}.$$

Then for any $m_0 \geq 1$ we have

$$\#\{(q, q') \in E : m_q([q, q']) \geq m_0\} \ll m_0^{-1} R \log Q.$$

Proof. By a dyadic division, it suffices to show that

$$\#\{(q, q') \in E : m_0 \leq m_q([q, q']) < 2m_0\} \ll m_0^{-1} R \log Q,$$

for any $m_0 \geq 1$. Call the left-hand side above $N(m_0)$. For any $D \geq 1$, let $\sigma_D(q)$ be the number of divisors of q in the range $[D, 8D]$. A moment's thought reveals that $m_q(r) \leq \sigma_D(q)$, where $D = Q^2/2R$. Indeed, each q' with $[q, q'] = r$ gives rise to a divisor (q, q') of q in the range $[D, 8D]$, and moreover (q, q') is uniquely determined by q' via $(q, q') = qq'/r$. It follows that

$$N(m_0) = \sum_{\substack{Q \leq q < 2Q \\ \sigma_D(q) \geq m_0}} \sum_{\substack{R \leq r < 2R \\ m_0 \leq m_q(r) < 2m_0}} m_q(r).$$

Since $m_q(r) = 0$ unless $q \mid r$, the inner sum over r is $O(m_0R/Q)$. It thus suffices to show that

$$\#\{Q \leq q < 2Q : \sigma_D(q) \geq m_0\} \ll m_0^{-2} Q \log Q,$$

for any $D \geq 1$. We may assume that $D \leq Q^{1/2}$ since otherwise we may replace D by $Q/8D$. By the second moment method, we have

$$\#\{Q \leq q < 2Q : \sigma_D(q) \geq m_0\} \leq \frac{1}{m_0^2} \sum_{Q \leq q < 2Q} \sigma_D(q)^2.$$

After expanding the square and changing the order of summation, the right-hand side above is

$$\frac{1}{m_0^2} \sum_{D \leq d_1, d_2 \leq 8D} \sum_{\substack{Q \leq q < 2Q \\ [d_1, d_2] \mid q}} 1 \ll \frac{Q}{m_0^2} \sum_{D \leq d_1, d_2 \leq 8D} \frac{1}{[d_1, d_2]} \ll \frac{Q}{m_0^2} \log D.$$

This completes the proof of the lemma. □

There are two places in the proof of [Lemma 3.3](#) where we lose a factor of $\log Q$ (and hence the assumption that $\delta < (\log Q)^{-1}$). One place is from dyadically decomposing the possible values of $[q, q']$, and the other from the conclusion of [Lemma 4.1](#). If one is only interested in prime moduli, then this extra loss can certainly be saved.

Proof of Lemma 3.3. In this proof, all implied constants are allowed to depend on d and s . For $k, k' \in [K, 2K)$, we may write

$$\sum_{\ell \in I(k, k')} F(k\ell) \overline{F(k'\ell)} = \sum_{Q \leq q, q' < 2Q} \sum_{\substack{\ell \in I(k, k', q, q') \\ k\ell \equiv a_q \pmod{q} \\ k'\ell \equiv a_{q'} \pmod{q'}}} \psi_q\left(\frac{k\ell - a_q}{q}\right) \overline{\psi_{q'}\left(\frac{k'\ell - a_{q'}}{q'}\right)}, \quad (4-1)$$

for some interval $I(k, k', q, q') \subset I(k, k')$. The solution to the simultaneous congruence conditions

$$k\ell \equiv a_q \pmod{q} \quad \text{and} \quad k'\ell \equiv a_{q'} \pmod{q'}$$

takes the form

$$\ell \equiv a(k, k', q, q') \pmod{[q, q']},$$

for some $0 \leq a(k, k', q, q') < [q, q']$. It is possible that no solutions exist, in which case we may simply set $I(k, k', q, q')$ to be empty and assign an arbitrary value to $a(k, k', q, q')$. After a change of variables $\ell = [q, q']m + a(k, k', q, q')$, the inner

sum over ℓ in (4-1) can be rewritten as

$$\sum_{m \in J(k, k', q, q')} \psi_q \left(\frac{k[q, q']}{q} m + b \right) \overline{\psi_{q'} \left(\frac{k'[q, q']}{q'} m + b' \right)},$$

for some interval $J(k, k', q, q') \subset [0, L/[q, q']]$, where

$$b = \frac{1}{q} (ka(k, k', q, q') - a_q) \quad \text{and} \quad b' = \frac{1}{q'} (k'a(k, k', q, q') - a_{q'}).$$

In principle b and b' depend on k, k', q and q' , but to simplify notations we drop this dependence, as the precise nature of b and b' is unimportant, apart from the obvious facts that $0 \leq b \leq k[q, q']/q$ and $0 \leq b' \leq k'[q, q']/q'$. Consider the polynomial sequence $g_{k, k', q, q'} : \mathbb{Z} \rightarrow G_q \times G_{q'}$ defined by

$$g_{k, k', q, q'}(m) = \left(g_q \left(\frac{k[q, q']}{q} m + b \right), g_{q'} \left(\frac{k'[q, q']}{q'} m + b' \right) \right),$$

and the Lipschitz function $\varphi_{q, q'} : G_q/\Gamma_q \times G_{q'}/\Gamma_{q'} \rightarrow \mathbb{C}$ defined by

$$\varphi_{q, q'}(x, x') = \varphi_q(x) \overline{\varphi_{q'}(x')}.$$

Then the type-II sum from (4-1) can be written as

$$\sum_{\ell \in I(k, k')} F(k\ell) \overline{F(k'\ell)} = \sum_{Q \leq q, q' < 2Q} \sum_{m \in J(k, k', q, q')} \varphi_{q, q'}(g_{k, k', q, q'}(m)).$$

After dyadically dividing the possible values of $[q, q']$, we deduce from the hypothesis (3-4) that

$$\sum_{K \leq k, k' < 2K} \sum_{\substack{Q \leq q, q' < 2Q \\ R \leq [q, q'] < 2R}} \left| \sum_{m \in J(k, k', q, q')} \varphi_{q, q'}(g_{k, k', q, q'}(m)) \right| \gg \delta^2 K^2 L, \quad (4-2)$$

for some $Q \leq R \leq 4Q^2$, where we used the assumption that $\delta < (\log Q)^{-1}$. For the rest of the proof fix such an R . Hence there is a subset T consisting of quadruples (k, k', q, q') with $R \leq [q, q'] < 2R$, such that

$$|T| \gg \delta^{O(1)} K^2 R, \quad (4-3)$$

and for $(k, k', q, q') \in T$ we have

$$\left| \sum_{m \in J(k, k', q, q')} \varphi_{q, q'}(g_{k, k', q, q'}(m)) \right| \gg \frac{\delta^2 L}{R}. \quad (4-4)$$

Since $\int \varphi_{q, q'} = 0$, (4-4) implies that the sequence $\{g_{k, k', q, q'}(m)\}_{0 \leq m \leq L/R}$ fails to be $\delta^{O(1)}$ -equidistributed. Hence by [Green and Tao 2012a, Theorem 2.9], there is

a nontrivial horizontal character $\chi_{q,q'} = \chi_{k,k',q,q'} : G_q \times G_{q'} \rightarrow \mathbb{C}$ with $\|\chi_{q,q'}\| \ll \delta^{-O(1)}$, such that

$$\|\chi_{q,q'} \circ g_{k,k',q,q'}\|_{C^\infty(L/R)} \ll \delta^{-O(1)}. \tag{4-5}$$

We have tacitly assumed that $\chi_{q,q'}$ is independent of k and k' , since this can be achieved after pigeonholing in the $\delta^{-O(1)}$ possible choices of $\chi_{q,q'}$ and enlarging the constant $O(1)$ in (4-3) appropriately. More explicitly, if we write

$$\chi_{q,q'} \circ g_{k,k',q,q'}(m) = \sum_{i=0}^s \beta_i(k, k', q, q') m^i, \tag{4-6}$$

for some coefficients $\beta_i(k, k', q, q') \in \mathbb{R}$, then (4-5) combined with [Green and Tao 2012b, Lemma 3.2] implies that there is a positive integer $r = O(1)$ such that

$$\|r\beta_i(k, k', q, q')\| \ll \delta^{-O(1)} \left(\frac{L}{R}\right)^{-i}, \tag{4-7}$$

for each $1 \leq i \leq s$ and $(k, k', q, q') \in T$. Write

$$\chi_{q,q'} = (\chi_{q,q'}^{(1)}, \chi_{q,q'}^{(2)}),$$

where $\chi_{q,q'}^{(1)}$ and $\chi_{q,q'}^{(2)}$ are horizontal characters on G_q and $G_{q'}$, respectively, with

$$\|\chi_{q,q'}^{(1)}\| \ll \delta^{-O(1)} \quad \text{and} \quad \|\chi_{q,q'}^{(2)}\| \ll \delta^{-O(1)}.$$

Write also

$$\chi_{q,q'}^{(1)} \circ g_q(n) = \sum_{i=0}^s \alpha_i(q, q') n^i \quad \text{and} \quad \chi_{q,q'}^{(2)} \circ g_{q'}(n) = \sum_{i=0}^s \alpha'_i(q, q') n^i,$$

for some coefficients $\alpha_i(q, q'), \alpha'_i(q, q') \in \mathbb{R}$.

Claim. *There exists a sequence of subsets $E_1 \subset \dots \subset E_s$ of pairs (q, q') with $R \leq [q, q'] < 2R$ and a sequence of positive integers $r_1 \geq \dots \geq r_s$ with $r_{i+1} \mid r_i$ for each i , such that $|E_1| \gg \delta^{O(1)} R$, $r_1 \ll \delta^{-O(1)}$, and moreover*

$$\|r_i \alpha_i(q, q')\| \ll \delta^{-O(1)} \left(\frac{KL}{Q}\right)^{-i} \quad \text{and} \quad \|r_i \alpha'_i(q, q')\| \ll \delta^{-O(1)} \left(\frac{KL}{Q}\right)^{-i},$$

for each $1 \leq i \leq s$ and $(q, q') \in E_i$.

Note that the claim actually implies the bounds

$$\|r_i \alpha_j(q, q')\| \ll \delta^{-O(1)} \left(\frac{KL}{Q}\right)^{-j} \quad \text{and} \quad \|r_i \alpha'_j(q, q')\| \ll \delta^{-O(1)} \left(\frac{KL}{Q}\right)^{-j},$$

for each $1 \leq i \leq j \leq s$ and $(q, q') \in E_j$.

Assuming the claim, we may conclude the proof of the lemma as follows: Pick an arbitrary pair $(q, q') \in E_1$. Since $\chi_{q,q'}$ is nontrivial, either $\chi_{q,q'}^{(1)}$ or $\chi_{q,q'}^{(2)}$

is nontrivial. Without loss of generality, assume that $\chi_{q,q'}^{(1)}$ is nontrivial. The diophantine information about $\alpha_i(q, q')$ from the claim implies that

$$\|r_1 \chi_{q,q'}^{(1)} \circ g_q\|_{C^\infty(KL/Q)} \ll \delta^{-O(1)}.$$

Thus by [Frantzikinakis and Host 2017, Lemma 5.3], the polynomial sequence g_q fails to be totally $\delta^{O(1)}$ -equidistributed.

It remains to establish the claim. Start by finding a subset E of pairs (q, q') with $R \leq [q, q'] < 2R$, such that the following properties hold:

- (1) $|E| \gg \delta^{O(1)} R$.
- (2) For each pair $(q, q') \in E$, there are at least $\delta^{O(1)} K^2$ pairs (k, k') such that $(k, k', q, q') \in T$.
- (3) For each pair $(q, q') \in E$, there are at most $\delta^{-O(1)}$ pairs $(q, \tilde{q}') \in E$ with $[q, q'] = [q, \tilde{q}']$, and similarly there are at most $\delta^{-O(1)}$ pairs $(\tilde{q}, q') \in E$ with $[q, q'] = [\tilde{q}, q']$.

Indeed, from the bound (4-3) we may first find E satisfying (1) and (2), and then apply Lemma 4.1 with $m_0 = \delta^{-C}$ for some sufficiently large C to remove a small number of pairs from E , so that property (3) is satisfied.

Construct $\{E_i\}$ and $\{r_i\}$ in the claim by downward induction on i as follows. Take E_{s+1} equal to the E just constructed and $r_{s+1} = r$ from (4-7). Now let $1 \leq i \leq s$, and suppose that E_j and r_j have already been constructed for $j > i$ satisfying the desired properties. First we show that for each pair $(q, q') \in E_{i+1}$, there is a positive integer $\tilde{r}(q, q') \ll \delta^{-O(1)}$ such that

$$\left\| \tilde{r}(q, q') r_{i+1} \alpha_i(q, q') \left(\frac{[q, q']}{q} \right)^i \right\| \ll \delta^{-O(1)} \left(\frac{KL}{R} \right)^{-i}, \quad (4-8)$$

and similarly with $\alpha_i(q, q')$ replaced by $\alpha'_i(q, q')$. To prove this, fix $(q, q') \in E_{i+1}$, and for the purpose of simplifying notations we drop the dependence on q and q' so that

$$\alpha_i = \alpha_i(q, q'), \quad \alpha'_i = \alpha'_i(q, q'), \quad \text{and} \quad \beta_i(k, k') = \beta_i(k, k', q, q').$$

From the definition of $g_{k,k',q,q'}$ we see the following relationship between the coefficients α_i, α'_i , and $\beta_i(k, k')$:

$$\beta_i(k, k') = \left(\frac{k[q, q']}{q} \right)^i \sum_{i \leq j \leq s} \alpha_j b^{j-i} + \left(\frac{k'[q, q']}{q'} \right)^i \sum_{i \leq j \leq s} \alpha'_j b^{j-i}. \quad (4-9)$$

Write $\tilde{\beta}_i(k, k')$ for the contribution from the term with $j = i$:

$$\tilde{\beta}_i(k, k') = \left(\frac{k[q, q']}{q} \right)^i \alpha_i + \left(\frac{k'[q, q']}{q'} \right)^i \alpha'_i.$$

By the induction hypothesis, $\|r_{i+1}\alpha_j\|$ and $\|r_{i+1}\alpha'_j\|$ are small for $j > i$. Combined with the bound $0 \leq b, b' \ll KR/Q$, this implies that the terms with $j > i$ are negligible:

$$\|r_{i+1}(\beta_i(k, k') - \tilde{\beta}_i(k, k'))\| \ll \delta^{-O(1)} \left(\frac{L}{R}\right)^{-i}.$$

It follows from (4-7) that

$$\|r_{i+1}\tilde{\beta}_i(k, k')\| \ll \delta^{-O(1)} \left(\frac{L}{R}\right)^{-i}, \quad (4-10)$$

whenever $(k, k', q, q') \in T$. Since $(q, q') \in E$, this holds for at least $\delta^{O(1)}K^2$ pairs (k, k') . Choose k' such that (4-10) holds whenever $k \in \mathcal{K}$, for some subset \mathcal{K} with $|\mathcal{K}| \gg \delta^{O(1)}K$. Since

$$\tilde{\beta}_i(k, k') - \tilde{\beta}_i(\tilde{k}, k') = \alpha_i \left(\frac{[q, q']}{q}\right)^i (k^i - \tilde{k}^i),$$

it follows that for $k, \tilde{k} \in \mathcal{K}$ we have

$$\left\| r_{i+1}\alpha_i \left(\frac{[q, q']}{q}\right)^i (k^i - \tilde{k}^i) \right\| \ll \delta^{-O(1)} \left(\frac{L}{R}\right)^{-i}.$$

Since $\delta > K^{-c}$ for some sufficiently small $c > 0$, the desired inequality (4-8) follows from a standard recurrence result such as [Green and Tao 2012a, Lemma 4.5]. The analogous bound for α'_i can be proved in a similar way.

Now that we have established (4-8), define $\tilde{E}_i \subset E_{i+1}$ to be a subset with $|\tilde{E}_i| \gg \delta^{O(1)}|E_{i+1}| \gg \delta^{O(1)}R$, such that $\tilde{r}(q, q')$ take a common value \tilde{r} for $(q, q') \in \tilde{E}_i$. We say that a pair $(q, q') \in \tilde{E}_i$ is *typical* if there are at least $\delta^{O(1)}R/Q$ pairs $(q, \tilde{q}') \in \tilde{E}_i$ with $\chi_{q, q'}^{(1)} = \chi_{q, \tilde{q}'}^{(1)}$, and similarly there are at least $\delta^{O(1)}R/Q$ pairs $(\tilde{q}, q') \in \tilde{E}_i$ with $\chi_{q, q'}^{(2)} = \chi_{\tilde{q}, q'}^{(2)}$. Define $E_i \subset \tilde{E}_i$ to be the set of typical pairs in \tilde{E}_i . By choosing the constant $O(1)$ in the definition of typical pairs sufficiently large, we may ensure that $|E_i| \gg \delta^{O(1)}R$.

Now let $(q, q') \in E_i$. Since (q, q') is typical, there exists a subset $\mathcal{Q}(q, q')$ with $|\mathcal{Q}(q, q')| \gg \delta^{O(1)}R/Q$, such that $(q, \tilde{q}') \in \tilde{E}_i$ and $\chi_{q, q'}^{(1)} = \chi_{q, \tilde{q}'}^{(1)}$ for all $\tilde{q}' \in \mathcal{Q}(q, q')$. Thus $\alpha(q, q') = \alpha(q, \tilde{q}')$ for all $\tilde{q}' \in \mathcal{Q}(q, q')$, and by applying (4-8) to (q, \tilde{q}') we obtain

$$\left\| \tilde{r}r_{i+1}\alpha_i(q, q') \left(\frac{[q, \tilde{q}']}{q}\right)^i \right\| \ll \delta^{-O(1)} \left(\frac{KL}{R}\right)^{-i},$$

for each $\tilde{q}' \in \mathcal{Q}(q, q')$. Since $\tilde{E}_i \subset E$, property (3) of the set E implies that

$$\#\left\{ \frac{[q, \tilde{q}']}{q} : \tilde{q}' \in \mathcal{Q}(q, q') \right\} \gg \delta^{O(1)}|\mathcal{Q}(q, q')| \gg \delta^{O(1)}\frac{R}{Q}.$$

By a standard recurrence result such as [Green and Tao 2012a, Lemma 4.5], there exists $r_i \ll \delta^{-O(1)} \tilde{r} r_{i+1} \ll \delta^{-O(1)}$ such that

$$\|r_i \alpha_i(q, q')\| \ll \delta^{-O(1)} \left(\frac{KL}{Q}\right)^{-i},$$

as desired. The analogous bound for $\alpha'_i(q, q')$ can be proved in a similar way. This finishes the proof of the claim, and also the proof of Lemma 3.3. \square

References

- [Bourgain et al. 2013] J. Bourgain, P. Sarnak, and T. Ziegler, “Disjointness of Moebius from horocycle flows”, pp. 67–83 in *From Fourier analysis and number theory to Radon transforms and geometry*, edited by H. M. Farkas et al., Dev. Math. **28**, Springer, New York, 2013. MR Zbl
- [Fouvry and Tenenbaum 1996] E. Fouvry and G. Tenenbaum, “Répartition statistique des entiers sans grand facteur premier dans les progressions arithmétiques”, *Proc. London Math. Soc.* (3) **72**:3 (1996), 481–514. MR Zbl
- [Frantzikinakis and Host 2017] N. Frantzikinakis and B. Host, “Higher order Fourier analysis of multiplicative functions and applications”, *J. Amer. Math. Soc.* **30**:1 (2017), 67–157. MR Zbl
- [Green 2016] B. Green, “A note on multiplicative functions on progressions to large moduli”, 2016. To appear in *Proc. Royal Soc. Edinburgh, Ser. A*. arXiv
- [Green and Tao 2012a] B. Green and T. Tao, “The quantitative behaviour of polynomial orbits on nilmanifolds”, *Ann. of Math.* (2) **175**:2 (2012), 465–540. MR Zbl
- [Green and Tao 2012b] B. Green and T. Tao, “The Möbius function is strongly orthogonal to nilsequences”, *Ann. of Math.* (2) **175**:2 (2012), 541–566. MR Zbl
- [Green et al. 2012] B. Green, T. Tao, and T. Ziegler, “An inverse theorem for the Gowers $U^{s+1}[N]$ -norm”, *Ann. of Math.* (2) **176**:2 (2012), 1231–1372. MR Zbl
- [Harper 2011] A. J. Harper, “A different proof of a finite version of Vinogradov’s bilinear sum inequality”, unpublished note, 2011, Available at <http://tinyurl.com/adamharper>.
- [Harper 2012] A. J. Harper, “Bombieri–Vinogradov and Barban–Davenport–Halberstam type theorems for smooth numbers”, preprint, 2012. arXiv
- [Iwaniec and Kowalski 2004] H. Iwaniec and E. Kowalski, *Analytic number theory*, American Mathematical Society Colloquium Publications **53**, American Mathematical Society, Providence, RI, 2004. MR Zbl
- [Kátai 1986] I. Kátai, “A remark on a theorem of H. Daboussi”, *Acta Math. Hungar.* **47**:1-2 (1986), 223–225. MR Zbl
- [Matomäki and Radziwiłł 2016] K. Matomäki and M. Radziwiłł, “Multiplicative functions in short intervals”, *Ann. of Math.* (2) **183**:3 (2016), 1015–1056. MR Zbl
- [Matomäki et al. 2015] K. Matomäki, M. Radziwiłł, and T. Tao, “An averaged form of Chowla’s conjecture”, *Algebra Number Theory* **9**:9 (2015), 2167–2196. MR Zbl
- [Matthiesen 2016] L. Matthiesen, “Generalized Fourier coefficients of multiplicative functions”, preprint, 2016. arXiv
- [Montgomery and Vaughan 1977] H. L. Montgomery and R. C. Vaughan, “Exponential sums with multiplicative coefficients”, *Invent. Math.* **43**:1 (1977), 69–82. MR Zbl
- [Ramaré 2009] O. Ramaré, *Arithmetical aspects of the large sieve inequality*, Harish-Chandra Research Institute Lecture Notes **1**, Hindustan Book Agency, New Delhi, 2009. MR Zbl
- [Tao and Vu 2006] T. Tao and V. Vu, *Additive combinatorics*, Cambridge Studies in Advanced Mathematics **105**, Cambridge Univ. Press, 2006. MR Zbl

Communicated by Peter Sarnak

Received 2016-07-15 Revised 2016-12-02 Accepted 2017-01-29

xuancheng.shao@maths.ox.ac.uk *Mathematical Institute, University of Oxford,
Radcliffe Observatory Quarter, Woodstock Road,
Oxford OX2 6GG, United Kingdom*

Algebra & Number Theory

msp.org/ant

EDITORS

MANAGING EDITOR

Bjorn Poonen
Massachusetts Institute of Technology
Cambridge, USA

EDITORIAL BOARD CHAIR

David Eisenbud
University of California
Berkeley, USA

BOARD OF EDITORS

Dave Benson	University of Aberdeen, Scotland	Shigefumi Mori	RIMS, Kyoto University, Japan
Richard E. Borcherds	University of California, Berkeley, USA	Martin Olsson	University of California, Berkeley, USA
J-L. Colliot-Thélène	CNRS, Université Paris-Sud, France	Raman Parimala	Emory University, USA
Brian D. Conrad	Stanford University, USA	Jonathan Pila	University of Oxford, UK
Samit Dasgupta	University of California, Santa Cruz, USA	Anand Pillay	University of Notre Dame, USA
Hélène Esnault	Freie Universität Berlin, Germany	Michael Rapoport	Universität Bonn, Germany
Gavril Farkas	Humboldt Universität zu Berlin, Germany	Victor Reiner	University of Minnesota, USA
Hubert Flenner	Ruhr-Universität, Germany	Peter Sarnak	Princeton University, USA
Sergey Fomin	University of Michigan, USA	Joseph H. Silverman	Brown University, USA
Edward Frenkel	University of California, Berkeley, USA	Michael Singer	North Carolina State University, USA
Andrew Granville	Université de Montréal, Canada	Christopher Skinner	Princeton University, USA
Joseph Gubeladze	San Francisco State University, USA	Vasudevan Srinivas	Tata Inst. of Fund. Research, India
Roger Heath-Brown	Oxford University, UK	J. Toby Stafford	University of Michigan, USA
Craig Huneke	University of Virginia, USA	Pham Huu Tiep	University of Arizona, USA
Kiran S. Kedlaya	Univ. of California, San Diego, USA	Ravi Vakil	Stanford University, USA
János Kollár	Princeton University, USA	Michel van den Bergh	Hasselt University, Belgium
Yuri Manin	Northwestern University, USA	Marie-France Vignéras	Université Paris VII, France
Philippe Michel	École Polytechnique Fédérale de Lausanne	Kei-Ichi Watanabe	Nihon University, Japan
Susan Montgomery	University of Southern California, USA	Shou-Wu Zhang	Princeton University, USA

PRODUCTION

production@msp.org

Silvio Levy, Scientific Editor


See inside back cover or msp.org/ant for submission instructions.

The subscription price for 2017 is US \$325/year for the electronic version, and \$520/year (+\$55, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to MSP.

Algebra & Number Theory (ISSN 1944-7833 electronic, 1937-0652 printed) at Mathematical Sciences Publishers, 798 Evans Hall #3840, c/o University of California, Berkeley, CA 94720-3840 is published continuously online. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices.

ANT peer review and production are managed by EditFLOW[®] from MSP.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2017 Mathematical Sciences Publishers

Algebra & Number Theory

Volume 11 No. 4 2017

Cup products of line bundles on homogeneous varieties and generalized PRV components of multiplicity one	767
IVAN DIMITROV and MIKE ROTH	
Mass formulas for local Galois representations and quotient singularities II: Dualities and resolution of singularities	817
MELANIE MATCHETT WOOD and TAKEHIKO YASUDA	
A new equivariant in nonarchimedean dynamics	841
ROBERT RUMELY	
On pairs of p -adic L -functions for weight-two modular forms	885
FLORIAN SPRUNG	
On Hilbert's 17th problem in low degree	929
OLIVIER BENOIST	
Gowers norms of multiplicative functions in progressions on average	961
XUANCHENG SHAO	
The degree of the Gauss map of the theta divisor	983
GIULIO CODOGNI, SAMUEL GRUSHEVSKY and EDOARDO SERNESI	
Correction to the article <i>Frobenius and valuation rings</i> (10:5 (2016), 1057–1090)	1003
RANKEYA DATTA and KAREN E. SMITH	