

Algebra & Number Theory

Volume 11

2017

No. 9

Adams operations on matrix factorizations

Michael K. Brown, Claudia Miller, Peder Thompson and Mark E. Walker



Adams operations on matrix factorizations

Michael K. Brown, Claudia Miller, Peder Thompson and Mark E. Walker

We define Adams operations on matrix factorizations, and we show these operations enjoy analogues of several key properties of the Adams operations on perfect complexes with support developed by Gillet and Soulé. As an application, we give a proof of a conjecture of Dao and Kurano concerning the vanishing of Hochster's θ pairing.

1. Introduction

We establish a theory of Adams operations on the Grothendieck group of matrix factorizations and use these operations to prove a conjecture of Dao and Kurano [2014, Conjecture 3.1(2)] concerning the vanishing of Hochster's θ pairing for a pair of modules defined on an isolated hypersurface singularity.

Let Q be a commutative Noetherian ring and let $f \in Q$. A *matrix factorization* of f in Q is a $\mathbb{Z}/2$ -graded, finitely generated projective Q -module $P = P_0 \oplus P_1$, equipped with an odd degree Q -linear endomorphism d satisfying $d^2 = f \operatorname{id}_P$. In other words, a matrix factorization is a pair of maps of finitely generated projective Q -modules, $(\alpha : P_1 \rightarrow P_0, \beta : P_0 \rightarrow P_1)$, satisfying $\alpha\beta = f \operatorname{id}_{P_0}$ and $\beta\alpha = f \operatorname{id}_{P_1}$.

When $f = 0$, a matrix factorization of f is the same thing as a $\mathbb{Z}/2$ -graded complex of finitely generated projective Q -modules. In this case, we have the evident $\mathbb{Z}/2$ -graded analogues of chain maps and homotopies of such. These, in fact, generalize to an arbitrary f . The matrix factorizations of $f \in Q$ form the objects of a category $\operatorname{mf}(Q, f)$, in which a morphism between objects P and P' of $\operatorname{mf}(Q, f)$ is a degree zero Q -linear map $g : P \rightarrow P'$ such that $d_{P'} \circ g = g \circ d_P$. In other words, a morphism is a pair of maps $g_0 : P_0 \rightarrow P'_0$ and $g_1 : P_1 \rightarrow P'_1$ causing the evident pair of squares to commute. A *homotopy* joining morphisms $g_1, g_2 : P \rightarrow P'$ in $\operatorname{mf}(Q, f)$ is a Q -linear map $h : P \rightarrow P'$ of odd degree such that $d_{P'}h + hd_P = g_1 - g_2$. The *homotopy category* of $\operatorname{mf}(Q, f)$ is the category $[\operatorname{mf}(Q, f)]$ obtained from $\operatorname{mf}(Q, f)$ by identifying homotopic morphisms. It is well-known that, when Q is

This work was partially supported by a grant from the Simons Foundation (#318705 for Mark Walker) and grants from the National Science Foundation (NSF Award DMS-0838463 for Michael Brown and Peder Thompson and NSF Award DMS-1003384 for Claudia Miller).

MSC2010: primary 13D15; secondary 13D02, 13D09, 13D22.

Keywords: Adams operations, matrix factorizations, Hochster's theta pairing.

regular and f is not a zero divisor, $[\mathrm{mf}(Q, f)]$ may be equipped with a canonical triangulated structure (see, for instance, [Orlov 2004] Section 3.1).

Much of the interest in matrix factorizations arises from the following result. For a Noetherian ring R , let $D^b(R)$ denote the bounded derived category of R . Objects of $D^b(R)$ are bounded complexes of finitely generated R -modules, and morphisms are obtained from chain maps by inverting the collection of quasiisomorphisms. Let $\mathrm{Perf}(R)$ denote the full triangulated subcategory of $D^b(R)$ consisting of bounded complexes of finitely generated and projective R -modules, and let $D_{\mathrm{sing}}(R)$ denote the Verdier quotient $D^b(R)/\mathrm{Perf}(R)$, called the *singularity category* of R . The following theorem is essentially due to work of Buchweitz [1986] and Eisenbud [1980]; this particular formulation of the result is proven by Orlov.

Theorem 1 [Orlov 2004, Theorem 3.9]. *If Q is regular and f is not a zero divisor, there is an equivalence of triangulated categories*

$$[\mathrm{mf}(Q, f)] \xrightarrow{\sim} D_{\mathrm{sing}}(Q/(f))$$

determined by sending a matrix factorization $(\alpha : P_1 \rightarrow P_0, \beta : P_0 \rightarrow P_1)$ to $\mathrm{coker}(\alpha)$.

Remark 1.1. In [Orlov 2004], Orlov assumes Q contains a field and has finite Krull dimension, but these assumptions are in fact not needed for this theorem to hold.

Let $R := Q/(f)$. Under the assumptions of Theorem 1, the Grothendieck group $K_0(\mathrm{mf}(Q, f))$ of the triangulated category $[\mathrm{mf}(Q, f)]$ is isomorphic to the quotient $G_0(R)/(\mathrm{im}(K_0(R) \rightarrow G_0(R)))$. So, defining a notion of Adams operations on $K_0(\mathrm{mf}(Q, f))$, in this setting, amounts to defining such operations on this quotient.

For a closed subset Z of $\mathrm{Spec}(Q)$, define $\mathcal{P}^Z(Q)$ to be the category of bounded complexes of finitely generated and projective Q -modules whose homology is supported on Z . Gillet–Soulé define lambda and Adams operations on the Grothendieck group $K_0^Z(Q) := K_0(\mathcal{P}^Z(Q))$ [Gillet and Soulé 1987, Sections 3 and 4]. It is tempting to mimic their approach to define Adams operations on $K_0(\mathrm{mf}(Q, f))$, since $\mathrm{mf}(Q, f)$ is somewhat analogous to $\mathcal{P}^{V(f)}(Q)$. But their construction relies on the Dold–Kan correspondence relating \mathbb{N} -graded complexes to simplicial modules; since matrix factorizations are $\mathbb{Z}/2$ -graded, such an approach is not available for $K_0(\mathrm{mf}(Q, f))$.

Instead, we model our approach after the construction of the *cyclic Adams operations* ψ_{cyc}^p on $K_0^Z(Q)$ developed by the authors in [BMTW 2017] (see also [Atiyah 1966; Haulion 2009; Köck 1997]). Let us give a brief summary of the construction of the operations ψ_{cyc}^p and some of their properties.

Fix a prime p . We assume that p is invertible in Q and that Q contains all p -th roots of unity (when Q is local, the case of primary interest to us, we can find such a prime p , at least after passing to a faithfully flat extension of Q). For a perfect

complex of \mathcal{Q} -modules X , let $T^p(X)$ denote the p -th tensor power of X , which comes equipped with a canonical left action by the symmetric group Σ_p . For a p -th root of unity $w \in \mathcal{Q}$, set $T^p(X)^{(w)}$ to be the eigenspace of eigenvalue w for the action of the p -cycle $(1\ 2\ \dots\ p)$ on $T^p(X)$. We define

$$\psi_{\text{cyc}}^p(X) = [T^p(X)^{(1)}] - [T^p(X)^{(\zeta)}]$$

where ζ is a primitive p -th root of unity.

In Sections 2 and 3 of [BMTW 2017], it is established that this formula induces a well-defined operation on $K_0^Z(\mathcal{Q})$ (see also [Hauton 2009]). In fact, by Corollary 6.14 of [BMTW 2017], if $p!$ is invertible in \mathcal{Q} , then ψ_{cyc}^p agrees with the p -th Adams operation on $K_0^Z(\mathcal{Q})$ defined by Gillet–Soulé. More generally, we have:

Theorem 2 [BMTW 2017, Theorem 3.7]. *If p is a prime, and \mathcal{Q} contains $1/p$ and all the p -th roots of unity, then the action of ψ_{cyc}^p on $K_0^Z(\mathcal{Q})$ satisfies the four Gillet–Soulé axioms defining a degree p Adams operation.*

We refer the reader to Theorem 3.7 of [BMTW 2017] for a precise statement of the four Gillet–Soulé axioms. A consequence of Theorem 2 is that the action of ψ_{cyc}^p on $K_0^Z(\mathcal{Q})_{\mathbb{Q}} := K_0^Z(\mathcal{Q}) \otimes \mathbb{Q}$ is diagonalizable: there is a “weight decomposition”

$$K_0^Z(\mathcal{Q})_{\mathbb{Q}} = \bigoplus_{i=c}^d K_0^Z(\mathcal{Q})_{\mathbb{Q}}^{(i)},$$

where $K_0^Z(\mathcal{Q})_{\mathbb{Q}}^{(i)}$ is the eigenspace of ψ_{cyc}^p of eigenvalue p^i , and c is the codimension of Z [loc. cit., Corollary 3.12].

In Section 2, we use the operations ψ_{cyc}^p as a model to construct cyclic Adams operations ψ_{cyc}^p on the Grothendieck group $K_0(\text{mf}(\mathcal{Q}, f))$, as well as more general versions for matrix factorizations with a support condition. In Theorem 2.10 and Proposition 2.13, we prove:

Theorem 3. *If p is prime, and \mathcal{Q} contains $1/p$ and all the p -th roots of unity, the operator ψ_{cyc}^p on $K_0(\text{mf}(\mathcal{Q}, f))$ satisfies the evident analogues of the four Gillet–Soulé axioms for a p -th Adams operation.*

Moreover, if \mathcal{Q} is regular and $f \in \mathcal{Q}$ is not a zero divisor, the canonical surjection

$$K_0^{V(f)}(\mathcal{Q}) \twoheadrightarrow K_0(\text{mf}(\mathcal{Q}, f))$$

is compatible with the action of ψ_{cyc}^p .

For \mathcal{Q} regular, f not a zero divisor, and $R = \mathcal{Q}/(f)$, given a finitely generated R -module M , let $[M]_{\text{stable}} \in K_0(\text{mf}(\mathcal{Q}, f))$ denote the image of $[M] \in G_0(R)$ under the canonical surjection $G_0(R) \twoheadrightarrow K_0(\text{mf}(\mathcal{Q}, f))$ given by Theorem 1.

Corollary 4. *Assume Q is a regular ring containing $1/p$ and all the p -th roots of unity for some prime p , and suppose $f \in Q$ is not a zero divisor. The action of ψ_{cyc}^p induces an eigenspace decomposition*

$$K_0(\text{mf}(Q, f))_{\mathbb{Q}} = \bigoplus_{i=1}^d K_0(\text{mf}(Q, f))_{\mathbb{Q}}^{(i)}.$$

Moreover, if M is a finitely generated R -module, then

$$[M]_{\text{stable}} \in \bigoplus_{i=\text{codim}_R M+1}^d K_0(\text{mf}(Q, f))_{\mathbb{Q}}^{(i)}.$$

In Section 3, we give an application of the above results. For the rest of this introduction, assume Q is a regular local ring with maximal ideal \mathfrak{m} , and assume f is a nonzero element of \mathfrak{m} . Assume also that $R = Q/(f)$ is an isolated singularity; that is, $R_{\mathfrak{p}}$ is regular for all $\mathfrak{p} \in \text{Spec}(R) \setminus \{\mathfrak{m}\}$. Then for any pair of finitely generated R -modules (M, N) , we have

$$\text{Tor}_i^R(M, N) \cong \text{Tor}_{i+2}^R(M, N) \quad \text{and} \quad \text{length Tor}_i^R(M, N) < \infty$$

for $i \gg 0$. This motivates the following definition.

Definition 1.2. With Q, f, R as above, for a pair of finitely generated R -modules (M, N) , set

$$\theta_R(M, N) = \text{length}(\text{Tor}_{2i}^R(M, N)) - \text{length}(\text{Tor}_{2i+1}^R(M, N))$$

for $i \gg 0$.

The pairing $\theta_R(-, -)$ is called *Hochster’s theta pairing*, since it first appeared in work of Hochster [1981]. The theta pairing should be regarded as the analogue, for the singularity category $D_{\text{sing}}(R)$, of the intersection multiplicity pairing that occurs, for example, in Serre’s multiplicity conjectures. There has been much recent work on better understanding the theta pairing, including when it vanishes and how it relates to more classical invariants. Buchweitz and van Straten [2012] show that, for complex isolated hypersurface singularities, the theta pairing can be recovered from the linking form on the link of an isolated singularity. In the same setting, Polishchuk and Vaintrob [2012] relate it to the classical residue pairing using the boundary bulk map. It was conjectured by Dao that θ vanishes for all isolated hypersurface singularities R such that $\dim(R)$ is even, and this has now been proven in almost all cases; see [Moore et al. 2011; Buchweitz and Van Straten 2012; Polishchuk and Vaintrob 2012; Walker 2017]. We refer the reader to Section 3 of [Dao and Kurano 2014] for additional history of the theta pairing and a list of several other conjectures.

One such conjecture, [Dao and Kurano 2014, Conjecture 3.1(2)], is an analogue of Serre’s vanishing conjecture (see the remark on page 111 of [Serre 2000]). This conjecture was proven by Dao in the case where R is excellent and contains a field, using a geometric approach [Dao 2013, Theorem 3.5]. As an application of the properties of Adams operations on matrix factorizations that we establish in Section 2, we prove this conjecture in full generality:

Theorem 5 (see Theorem 3.19). *Let (Q, \mathfrak{m}) be a regular local ring and $f \in \mathfrak{m}$ with $f \neq 0$. Suppose that $R = Q/(f)$ is an isolated singularity. If M and N are finitely generated R -modules such that*

$$\dim M + \dim N \leq \dim R$$

then $\theta_R(M, N) = 0$.

We close this introduction with a sketch of our proof of Theorem 5. We easily reduce to the case where there is a prime p such that Q contains $1/p$ and all p -th roots of unity. Given a matrix factorization $P = (\alpha : P_1 \rightarrow P_0, \beta : P_0 \rightarrow P_1)$ of f , one may obtain a matrix factorization P° of $-f$ by negating β . In Proposition 3.18, we show

$$\theta_R(M, N) = \chi([M]_{\text{stable}} \cup [N]_{\text{stable}}^\circ),$$

where $-\cup-$ is the pairing induced by tensor product of matrix factorizations, and χ denotes the Euler characteristic. The assumptions ensure that $[M]_{\text{stable}} \cup [N]_{\text{stable}}^\circ$ is a class in $K_0(\text{mf}^m(Q, 0))$, the Grothendieck group of $\mathbb{Z}/2$ -graded complexes of finitely generated projective Q -modules with finite length homology, so that χ is well-defined. By Corollary 4 and the linearity of χ , we may assume that the classes $[M]_{\text{stable}}$ and $[N]_{\text{stable}}^\circ$ lie in eigenspaces $K_0(\text{mf}(Q, 0))_{\mathbb{Q}}^{(i)}$ and $K_0(\text{mf}(Q, 0))_{\mathbb{Q}}^{(j)}$, respectively, where $i + j > d = \dim Q$. By properties of the operations ψ_{cyc}^p established in Theorem 3, $[M]_{\text{stable}} \cup [N]_{\text{stable}}^\circ \in K_0(\text{mf}^m(Q, 0))_{\mathbb{Q}}^{(i+j)}$.

At this point, one would like to argue that $K_0(\text{mf}^m(Q, 0))_{\mathbb{Q}} = K_0(\text{mf}^m(Q, 0))_{\mathbb{Q}}^{(d)}$, which would force $[M]_{\text{stable}} \cup [N]_{\text{stable}}^\circ = 0$. Indeed, one might expect $K_0(\text{mf}^m(Q, 0))$ to be generated by the $\mathbb{Z}/2$ -folding of the class of the Koszul complex on a regular sequence of generators of \mathfrak{m} , which lies in $K_0(\text{mf}^m(Q, 0))^{(d)}$ by the axioms in Theorem 3; this would be parallel to what occurs for bounded \mathbb{Z} -graded complexes. The proof of Theorem 5 sketched here would then be almost exactly the same as Gillet and Soulé’s proof of Serre’s vanishing conjecture.

We are not able to prove $K_0(\text{mf}^m(Q, 0))$ is generated by the Koszul complex, and indeed we have come to suspect this might be false (see Example 3.6). Fortunately, for the proof of Dao and Kurano’s conjecture, one needs only the weaker property that there is an equality of maps $\chi \circ \psi_{\text{cyc}}^p = p^d \chi$ from $K_0(\text{mf}^m(Q, 0))$ to \mathbb{Z} ; we prove this in Theorem 3.8.

2. Adams operations on matrix factorizations

In this section, we define cyclic Adams operations on matrix factorizations, closely following the construction of cyclic Adams operations on perfect complexes with support found in Sections 2 and 3 of [BMTW 2017]. We prove these operations enjoy analogues of many of the key properties of the operations on perfect complexes with support constructed in [loc. cit.].

2A. Construction. Let Q be a Noetherian commutative ring, $f \in Q$ any element (including possibly $f = 0$), and G a finite group. Let $\text{mf}(Q, f; G)$ be the category of G -equivariant matrix factorizations. When G is the trivial group, this is the category described in the introduction. More generally, an object of $\text{mf}(Q, f; G)$ is an object P of $\text{mf}(Q, f)$ equipped with a G -action (i.e., a group homomorphism $G \rightarrow \text{Aut}_{\text{mf}(Q, f)}(P)$), and a morphism is a G -equivariant morphism of matrix factorizations.

The category $\text{mf}(Q, f; G)$ is an exact category, with the notion of exactness given degree-wise in the evident manner.

Remark 2.1. We could equivalently define an object of $\text{mf}(Q, f; G)$ to consist of a pair of $Q[G]$ -modules P_0 and P_1 that are finitely generated and projective as Q -modules, together with a pair of morphisms of $Q[G]$ -modules, $(\alpha : P_1 \rightarrow P_0, \beta : P_0 \rightarrow P_1)$, such that $\alpha\beta$ and $\beta\alpha$ are each multiplication by f (which is central in $Q[G]$). Moreover, if $|G|$ is invertible in Q , we have $\text{mf}(Q, f; G) = \text{mf}(Q[G], f)$.

Example 2.2. If $f = 0$ (and G is trivial), $\text{mf}(Q, 0)$ is the category of $\mathbb{Z}/2$ -graded complexes of finitely generated projective Q -modules, with morphisms being chain maps.

A homotopy joining morphisms $g_1, g_2 : P \rightarrow P'$ in $\text{mf}(Q, f; G)$ is defined just as in the introduction, with the added condition that it be G -equivariant. In detail, it is a Q -linear, G -equivariant map $h : P \rightarrow P'$ of degree 1 such that $d_{P'}h + hd_P = g_1 - g_2$. The homotopy category of $\text{mf}(Q, f; G)$ is the category $[\text{mf}(Q, f; G)]$ obtained from $\text{mf}(Q, f; G)$ by identifying homotopic morphisms.

Given a ring homomorphism $Q \rightarrow Q'$ sending f to f' , there is an evident functor $\text{mf}(Q, f; G) \rightarrow \text{mf}(Q', f'; G)$ given by extension of scalars along $Q \rightarrow Q'$. When $Q' = Q_{\mathfrak{p}}$ for $\mathfrak{p} \in \text{Spec}(Q)$, we write this functor as $P \mapsto P_{\mathfrak{p}}$.

For an object $P \in \text{mf}(Q, f; G)$, define the support of P to be

$$\text{supp}(P) = \{\mathfrak{p} \in \text{Spec}(Q) \mid P_{\mathfrak{p}} \text{ is not homotopy equivalent to } 0 \text{ in } \text{mf}(Q_{\mathfrak{p}}, f; G)\}.$$

Given a closed subset Z of $\text{Spec}(Q)$, define $\text{mf}^Z(Q, f; G)$ to be the full subcategory of $\text{mf}(Q, f)$ consisting of objects P satisfying $\text{supp}(P) \subseteq Z$. Note that $\text{mf}^Z(Q, f; G)$ is a full, exact subcategory of $\text{mf}(Q, f; G)$, and $[\text{mf}^Z(Q, f; G)]$ is a full subcategory of $[\text{mf}(Q, f; G)]$.

We will mainly use the notion of supports for matrix factorizations when $f = 0$ and G is trivial, in which case objects of $\text{mf}(Q, 0)$ are $(\mathbb{Z}/2\text{-graded})$ complexes. One must be careful in this situation not to conflate the notion of being homotopy equivalent to 0 with being acyclic. The former implies the latter, but the latter does not imply the former in general. These conditions are equivalent, however, in the following case:

Lemma 2.3. *If Q is a regular ring, an object $P \in \text{mf}(Q, 0)$ is contractible if and only if $H_0(P) = H_1(P) = 0$.*

Proof. Suppose $P = (\alpha_0 : P_0 \rightarrow P_1, \alpha_1 : P_1 \rightarrow P_0)$ is acyclic, and set $M = \ker(\alpha_1) = \text{im}(\alpha_0)$ and $N = \ker(\alpha_0) = \text{im}(\alpha_1)$. We claim that M and N are projective. It suffices to prove $M_{\mathfrak{p}}$ and $N_{\mathfrak{p}}$ are free for all primes \mathfrak{p} . Since

$$0 \rightarrow M_{\mathfrak{p}} \rightarrow (P_1)_{\mathfrak{p}} \rightarrow (P_0)_{\mathfrak{p}} \rightarrow (P_1)_{\mathfrak{p}} \rightarrow \dots$$

is exact, we see that, for any d , $M_{\mathfrak{p}}$ is a d -th syzygy of some other $Q_{\mathfrak{p}}$ -module. Taking $d > \dim(Q_{\mathfrak{p}})$ gives that $M_{\mathfrak{p}}$ is free. Similarly, N is projective.

Choose splittings $\pi_0 : P_0 \rightarrow N$ and $\pi_1 : P_1 \rightarrow M$ of the inclusions $N \hookrightarrow P_0$ and $M \hookrightarrow P_1$. Define $A : P_0 \rightarrow N \oplus M$ and $B : P_1 \rightarrow N \oplus M$ to be given by $\begin{pmatrix} \pi_0 \\ \alpha_0 \end{pmatrix}$ and $\begin{pmatrix} \alpha_1 \\ \pi_1 \end{pmatrix}$, respectively. Set $E := \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix}$ and $F := \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}$.

We have the following isomorphism of matrix factorizations

$$\begin{array}{ccccc} P_0 & \xrightarrow{\alpha_0} & P_1 & \xrightarrow{\alpha_1} & P_0 \\ A \downarrow & & B \downarrow & & A \downarrow \\ N \oplus M & \xrightarrow{E} & N \oplus M & \xrightarrow{F} & N \oplus M \end{array}$$

and the bottom matrix factorization is clearly contractible. □

Remark 2.4. When Q is regular, f is not a zero divisor, and G is trivial, the support of any object of $\text{mf}(Q, f)$ is a subset of

$$\text{Sing}(R) := \{\mathfrak{p} \in \text{Spec}(R) \mid R_{\mathfrak{p}} \text{ is not regular}\}$$

where $R = Q/(f)$, and where we identify $\text{Spec } R$ with its image in $\text{Spec } Q$. Thus, in this case, we have

$$\text{mf}(Q, f) = \text{mf}^{\text{Sing}(R)}(Q, f).$$

Eventually, we will be making the additional assumption that R is an isolated singularity, meaning Q , and hence R , is local, and $\text{Sing}(R) = \{\mathfrak{m}\}$.

Define the Grothendieck group $K_0(\text{mf}^Z(Q, f; G))$ to be the abelian monoid given by isomorphism classes of objects of $\text{mf}^Z(Q, f; G)$ under the operation of direct sum, modulo the relations $[P] = [P'] + [P'']$ if there exists a short exact sequence $0 \rightarrow P' \rightarrow P \rightarrow P'' \rightarrow 0$ and $[P] = [P']$ if P and P' are homotopy

equivalent. As with the K -theory of complexes, $K_0(\text{mf}^Z(Q, f; G))$ is an abelian group, since $[P] + [\Sigma(P)] = 0$, where $\Sigma(P)$ denotes the suspension of P .

For $P \in \text{mf}(Q, f; G)$ and $P' \in \text{mf}(Q, f'; G')$, the tensor product $P \otimes_Q P'$ is the usual tensor product of Q -modules, with grading determined by $|p \otimes p'| = |p| + |p'|$ and differential $\partial(p \otimes p') = d_P(p) \otimes p' + (-1)^{|p|} p \otimes d_{P'}(p')$. The group $G \times G'$ acts in the evident manner, and the resulting object belongs to $\text{mf}(Q, f + f'; G \times G')$, since ∂^2 is multiplication by $f + f'$. Note, in particular, that the n -th tensor power of an object of $\text{mf}(Q, f)$ belongs to $\text{mf}(Q, nf)$.

We proceed to define cyclic Adams operations on $K_0(\text{mf}^Z(Q, f))$. The construction is closely parallel to that for $K_0^Z(Q)$ given in [BMTW 2017], with one minor exception: the need to “divide by p ”.

For an integer $n \geq 1$, we define a functor

$$T^n : \text{mf}^Z(Q, f) \rightarrow \text{mf}^Z(Q, nf; \Sigma_n)$$

given, on objects, by sending $P \in \text{mf}^Z(Q, f)$ to the matrix factorization

$$T^n(P) = \overbrace{P \otimes_Q \cdots \otimes_Q P}^{n \text{ times}}$$

equipped with the left action of Σ_n given by

$$\sigma(p_1 \otimes \cdots \otimes p_n) = \pm p_{\sigma^{-1}(1)} \otimes \cdots \otimes p_{\sigma^{-1}(n)}.$$

The sign is uniquely determined by the following rule: if σ is the transposition $(i \ i + 1)$ for some $1 \leq i \leq n - 1$ and p_1, \dots, p_n are homogenous elements of P , then

$$\sigma(p_1 \otimes \cdots \otimes p_n) = (-1)^{|p_i||p_{i+1}|} p_1 \otimes \cdots \otimes p_{i-1} \otimes p_{i+1} \otimes p_i \otimes p_{i+2} \otimes \cdots \otimes p_n.$$

The rule for morphisms is the evident one.

Following Section 2 of [BMTW 2017], for any i and j , let $\Sigma_{i,j}$ be the image of the canonical homomorphism $\Sigma_i \times \Sigma_j \hookrightarrow \Sigma_{i+j}$, and define a pairing

$$\star_{i,j} : K_0(\text{mf}^Z(Q, if); \Sigma_i) \times K_0(\text{mf}^Z(Q, jf); \Sigma_j) \rightarrow K_0(\text{mf}^Z(Q, (i+j)f); \Sigma_{i+j})$$

induced by the bifunctor $(P, P') \mapsto Q[\Sigma_{i+j}] \otimes_{Q[\Sigma_{i,j}]} P \otimes_Q P'$. This pairing is well-defined, commutative, and associative, by an argument identical to the proof of Lemma 2.4 in [loc. cit.].

The proof of Theorem 2.2 in [loc. cit.] also holds nearly verbatim for matrix factorizations and leads to a proof of:

Theorem 2.5. *For a commutative Noetherian ring Q , closed subset Z of $\text{Spec}(Q)$, element $f \in Q$, and integer $n \geq 1$, there is a function*

$$t_\Sigma^n : K_0(\text{mf}^Z(Q, f)) \rightarrow K_0(\text{mf}^Z(Q, nf; \Sigma_n))$$

such that, for an object $P \in \text{mf}^Z(Q, f)$, we have

$$t_{\Sigma}^n([P]) = [T^n(P)].$$

Remark 2.6. As in [BMTW 2017, §5], if k is a positive integer such that $k!$ is invertible in Q , then one can use Theorem 2.5 to establish an operation λ^k on $K_0(\text{mf}^Z(Q, f))$ that is induced from the k -th exterior power functor. Since we won't use such operations in this paper, we omit the details.

We now assume p is a prime that is invertible in Q , and we define C_p to be the subgroup of Σ_p generated by the p -cycle $(1\ 2\ \dots\ p)$. For any p -th root of unity ζ belonging to Q (including the case $\zeta = 1$), let Q_{ζ} denote the $Q[C_p]$ -module Q equipped with the C_p -action $\sigma q = \zeta q$. For $P \in \text{mf}^Z(Q, pf; C_p)$, we define

$$P^{(\zeta)} := \text{Hom}_{Q[C_p]}(Q_{\zeta}, P) = \ker(\sigma - \zeta : P \rightarrow P).$$

Since p is invertible and ζ belongs to Q , the module Q_{ζ} is a direct summand of $Q[C_p]$, and so $P \mapsto P^{(\zeta)}$ is an exact functor. It therefore induces a map

$$\phi_{\zeta}^p : K_0(\text{mf}^Z(Q, pf; C_p)) \xrightarrow{[P] \mapsto [P^{(\zeta)}]} K_0(\text{mf}^Z(Q, pf)),$$

and so we may form the composition

$$K_0(\text{mf}^Z(Q, f)) \xrightarrow{t_{\Sigma}^p} K_0(\text{mf}^Z(Q, pf; \Sigma_p)) \xrightarrow{\text{res}} K_0(\text{mf}^Z(Q, pf; C_p)) \xrightarrow{\phi_{\zeta}^p} K_0(\text{mf}^Z(Q, pf)).$$

We come upon the need to “divide by p ”. In general, if $u \in Q$ is a unit, we define an autoequivalence

$$\text{mult}_u : \text{mf}^Z(Q, f) \rightarrow \text{mf}^Z(Q, uf)$$

by sending a matrix factorization (α, β) to $(\alpha, u\beta)$. (Its inverse is given by $\text{mult}_{u^{-1}}$.) For example, in Section 3C, we will employ the functor mult_{-1} , which we will write as $\text{mult}_{-1}(P) = P^{\circ}$. Here, we use $\text{mult}_{1/p}$, and we define t_{ζ}^p to be the composition

$$K_0(\text{mf}^Z(Q, f)) \xrightarrow{\phi_{\zeta}^p \circ \text{res} \circ t_{\Sigma}^p} K_0(\text{mf}^Z(Q, pf)) \xrightarrow{\text{mult}_{1/p}} K_0(\text{mf}^Z(Q, f)).$$

Let A_p denote the subring of \mathbb{C} given by $\mathbb{Z}[1/p, e^{2\pi i/p}]$.

Definition 2.7. Assume p is a prime, Q is a (commutative, Noetherian) A_p -algebra, f is any element of Q , and Z is a closed subset of $\text{Spec}(Q)$. Define

$$\psi_{\text{cyc}}^p = \sum_{\zeta} \zeta t_{\zeta}^p : K_0(\text{mf}^Z(Q, f)) \rightarrow K_0(\text{mf}^Z(Q, f)),$$

where the sum ranges over all p -th roots of unity. (In this formula, the ζ occurring as a coefficient is interpreted as belonging to $\mathbb{Z}[e^{2\pi i/p}]$ whereas the ζ occurring as a subscript denotes its image in Q under the map $A_p \rightarrow Q$.)

Remark 2.8. The image of ψ_{cyc}^p is contained in the group $K_0(\text{mf}^Z(Q, f)) \otimes_{\mathbb{Z}} \mathbb{Z}[e^{2\pi i/p}]$. But, by an argument identical to the proof of Corollary 3.5 in [BMTW 2017], we have

$$\sum_{\zeta} \zeta t_{\zeta}^p = t_1^p - t_{\zeta'}^p$$

for any fixed primitive p -th root of unity ζ' , and thus the image of ψ_{cyc}^p can be taken to be $K_0(\text{mf}^Z(Q, f))$.

Remark 2.9. Setting $\phi^p = \sum_{\zeta} \zeta \phi_{\zeta}^p$, one gets another formulation

$$\psi_{\text{cyc}}^p = \text{mult}_{1/p} \circ \phi^p \circ \text{res} \circ \iota_{\Sigma}^p.$$

2B. Axioms for Adams operations on matrix factorizations à la Gillet–Soulé. In this subsection, we show the operations ψ_{cyc}^p satisfy the following analogues of the axioms of Gillet and Soulé (see Theorem 3.7 in [BMTW 2017]).

Theorem 2.10. *Assume p is a prime, Q is a (commutative, Noetherian) A_p -algebra, f, f_1, f_2 are any elements of Q , and Z is a closed subset of $\text{Spec}(Q)$:*

- (1) ψ_{cyc}^p is a group endomorphism of $K_0(\text{mf}^Z(Q, f))$.
- (2) For $\alpha \in K_0(\text{mf}^Z(Q, f_1))$ and $\beta \in K_0(\text{mf}^W(Q, f_2))$,

$$\psi_{\text{cyc}}^p(\alpha \cup \beta) = \psi_{\text{cyc}}^p(\alpha) \cup \psi_{\text{cyc}}^p(\beta) \in K_0(\text{mf}^{Z \cap W}(Q, f_1 + f_2)),$$

where \cup is the multiplication rule on Grothendieck groups induced by tensor product. The three operators ψ_{cyc}^p in the equation are, from left to right, acting on $K_0(\text{mf}^{Z \cap W}(Q, f_1 + f_2))$, $K_0(\text{mf}^Z(Q, f_1))$, and $K_0(\text{mf}^W(Q, f_2))$.

- (3) ψ_{cyc}^p is functorial in the following sense: Suppose $\rho : Q \rightarrow Q'$ is map of A_p -algebras, $f' = \rho(f)$, and $\tilde{\rho}^{-1}(Z) \subseteq Z'$ where $\tilde{\rho} : \text{Spec } Q' \rightarrow \text{Spec } Q$ is the induced map on spectra. Then extension of scalars along ρ induces a map $K_0(\text{mf}^Z(Q, f)) \rightarrow K_0(\text{mf}^{Z'}(Q', f'))$ that commutes with the actions of ψ_{cyc}^p .
- (4) If $f = gh$, so that $(g, h) := (Q \xrightarrow{g} Q, Q \xrightarrow{h} Q)$ is an object of $\text{mf}^{V(g,h)}(Q, f)$, we have

$$\psi_{\text{cyc}}^p[(g, h)] = p[(g, h)].$$

Proof. The proofs of (1)–(3) are essentially identical to the proofs of parts (1)–(3) of Theorem 3.7 in [BMTW 2017]. As for (4), let $(0, 0)$ denote the matrix factorization $(Q \xrightarrow{0} Q, Q \xrightarrow{0} Q)$ of 0, and let X denote the tensor product

$$(g, ph) \otimes_Q (0, 0) \otimes_Q \cdots \otimes_Q (0, 0).$$

Set $\zeta := e^{2\pi i/p}$ and $\sigma := (1\ 2\ \cdots\ p) \in C_p$. We equip X with a C_p action by letting σ act on the i -th factor of X in the following way: If x has odd degree, $\sigma \cdot x = \zeta^{i-1}x$. If x has even degree, $\sigma \cdot x = x$.

We claim that there is an isomorphism

$$T^p([g, h]) \cong (g, ph) \otimes_Q (0, 0) \otimes_Q \cdots \otimes_Q (0, 0)$$

in $\text{mf}^{V(g,h)}(Q, pf; C_p)$. To prove the claim, let V be a free Q -module of rank p with a fixed basis $\{e_0, \dots, e_{p-1}\}$. We identify the underlying Q -modules of $T^p((g, h))$ and X with the exterior algebra $\bigwedge V$ of V ; under this identification, the action of C_p on $T^p((g, h))$ is given by

$$\sigma(e_{i_1} \wedge \cdots \wedge e_{i_n}) = e_{\sigma^{-1}(i_1)} \wedge \cdots \wedge e_{\sigma^{-1}(i_n)},$$

and the action of C_p on X is given by

$$\sigma(e_{i_1} \wedge \cdots \wedge e_{i_n}) = \zeta^{i_1 + \cdots + i_n} e_{i_1} \wedge \cdots \wedge e_{i_n}.$$

For $0 \leq i \leq p-1$, define $v_i := 1/p \sum_j \zeta^{ij} e_j$. Then v_0, \dots, v_{p-1} form a basis of V . Let $\alpha : \bigwedge V \rightarrow \bigwedge V$ denote the Q -algebra automorphism given by $e_i \mapsto v_i$. Then α yields an isomorphism $T^p((g, h)) \xrightarrow{\sim} X$ of C_p -equivariant matrix factorizations; this proves the claim.

(In checking the details here, it is useful to note the following: The ‘‘differential’’ on $T^p((g, h))$ is given by $s_0 + s_1$, where s_0 is left-multiplication by $h(e_0 + \cdots + e_{p-1})$, and s_1 is given by the Koszul differential on the sequence (g, g, \dots, g) . Similarly, the ‘‘differential’’ on X is given by $t_0 + t_1$, where t_0 is left-multiplication by $ph e_0$ and t_1 is given by the Koszul differential on the sequence $(g, 0, \dots, 0)$.)

By [Remark 2.9](#), and the result analogous to Lemma 3.11 of [\[BMTW 2017\]](#) for matrix factorizations (with essentially the same proof), we have

$$\psi_{\text{cyc}}^p([g, h]) = \text{mult}_{1/p}(\phi^p([g, ph]) \cup \phi^p([(0, 0)]) \cup \cdots \cup \phi^p([(0, 0)])).$$

Here, ϕ^p acts as the identity on the first factor, which is equipped with the trivial action of C_p . Furthermore, direct calculation on the $(i+1)$ -st factor yields

$$\phi^p([(0, 0)]) = [I] + \zeta^i[\Sigma I] = (1 - \zeta^i)[I]$$

where I denotes the unit matrix factorization $(0 \xrightarrow{0} Q, Q \xrightarrow{0} 0)$. Thus, one obtains

$$\psi_{\text{cyc}}^p([g, h]) = \text{mult}_{1/p}([g, ph] \cup [I] \cup \cdots \cup [I]) \prod_{i=1}^{p-1} (1 - \zeta^i) = p[g, h],$$

since $\prod_{i=1}^{p-1} (1 - \zeta^i) = p$. □

Corollary 2.11. *If $a = (a_1, \dots, a_n)$ is a sequence of elements in an A_p -algebra Q and $K(a)$ is the associated $\mathbb{Z}/2$ -folded Koszul complex, regarded as an object of $\text{mf}^{V(a_1, \dots, a_n)}(Q, 0)$, then*

$$\psi_{\text{cyc}}^p([K(a)]) = p^n [K(a)] \in K_0(\text{mf}^{V(a_1, \dots, a_n)}(Q, 0)).$$

Proof. This follows from parts (2) and (4) of the theorem, because $K(a)$ is the tensor product of the matrix factorizations $(a_i, 0)$ and $\mathbb{Z}/2$ -folding commutes with tensor product. \square

2C. Diagonalizability. Suppose Q is a regular ring and $f \in Q$ is not a zero divisor. Recall, from the introduction, that $\mathcal{P}^{V(f)}(Q)$ denotes the category of bounded complexes of finitely generated and projective Q -modules whose homology is supported on $V(f)$, and $K_0^{V(f)}(Q)$ denotes its Grothendieck group. In this subsection, we construct a surjection

$$\rho_f : K_0^{V(f)}(Q) \twoheadrightarrow K_0(\text{mf}(Q, f))$$

that commutes with the actions of ψ_{cyc}^P . Using this, and Corollary 3.12 of [BMTW 2017] (the proof of which is really due to Gillet–Soulé), we deduce that the action of ψ_{cyc}^P on $K_0(\text{mf}(Q, f))_{\mathbb{Q}}$ decomposes the latter into eigenspaces of the expected weights.

Let K_f denote the Koszul dga associated to f , so that, as a Q -algebra, $K_f = Q[\epsilon]/(\epsilon^2)$ with $|\epsilon| = 1$, and it is equipped with the Q -linear differential d satisfying $d(\epsilon) = f$. Let $P(K_f/Q)$ denote the full subcategory of the category of dg- K_f -modules consisting of those that are finitely generated and projective as Q -modules. An object of $P(K_f/Q)$ is thus a bounded complex P of finitely generated projective Q -modules equipped with a degree one Q -linear map $s : P \rightarrow P_{+1}$ satisfying $d_P s + s d_P = f$ and $s^2 = 0$. (The map s is given by multiplication by ϵ .) A morphism from (P, d_P, s) to $(P', d_{P'}, s')$ is a chain map g such that $g s = s' g$. A homotopy from g_1 to g_2 is a degree one map h such that $d_{P'} h + h d_P = g_1 - g_2$ and $h s = s' h$.

There are functors

$$\mathcal{P}^{V(f)}(Q) \xleftarrow{F} P(K_f/Q) \xrightarrow{\text{Fold}} \text{mf}(Q, f),$$

where F is the forgetful functor that sends (P, d_P, s) to (P, d_P) , and Fold sends (P, d, s) to the following matrix factorization: the even degree part is $\bigoplus_i P_{2i}$, the odd degree part is $\bigoplus_i P_{2i+1}$ and the degree one endomorphism is $\partial := d + s$.

Define $K_0(P(K_f/Q))$ to be the Grothendieck group of objects modulo relations coming from short exact sequences and homotopy equivalences as usual.

Lemma 2.12. *If f is not a zero divisor in a regular ring Q , the functor F induces an isomorphism*

$$K_0(P(K_f/Q)) \xrightarrow{\sim} K_0^{V(f)}(Q).$$

Proof. Let $R = Q/(f)$. One has an evident quasiisomorphism $K_f \xrightarrow{\sim} R$ of dga's, and hence an equivalence of triangulated categories $D^b(R) \xrightarrow{\sim} D^b(K_f)$ induced by restriction of scalars. Thus, one has an isomorphism

$$G_0(R) = K_0(D^b(R)) \xrightarrow{\sim} K_0(D^b(K_f)).$$

We may model $D^b(K_f)$ by semiprojective K_f -modules with finitely generated homology. Since Q is regular, the good truncation of such a complex in sufficiently high degree is a complex of projective Q -modules. It thus follows from Quillen’s resolution theorem that the inclusion map determines an isomorphism

$$K_0(P(K_f/Q)) \xrightarrow{\sim} K_0(D^b(K_f)).$$

We thus obtain an isomorphism $G_0(R) \xrightarrow{\sim} K_0(P(K_f/Q))$, which we can describe explicitly as follows: If M is a finitely generated R -module, form a (possibly infinite) K_f -semiprojective resolution $P \xrightarrow{\sim} M$ of M . Then the map sends $[M]$ to $[P']$ where P' is a good truncation of P in sufficiently high degree.

We also have the more classical isomorphism $G_0(R) \xrightarrow{\sim} K_0^{V(f)}(Q)$, sending $[M]$ to the class of a Q -projective resolution of M . Since the complex P' constructed above is an example of such a resolution, it is clear that the triangle

$$\begin{array}{ccc}
 K_0(P(K_f/Q)) & \xrightarrow{F} & K_0^{V(f)}(Q) \\
 \swarrow \cong & & \searrow \cong \\
 & G_0(R) &
 \end{array}$$

commutes. □

The functor Fold induces a map from $K_0(P(K_f/Q))$ to $K_0(\text{mf}(Q, f))$, and thus, using the lemma, we obtain the desired map $\rho_f : K_0^{V(f)}(Q) \rightarrow K_0(\text{mf}(Q, f))$. Explicitly, the construction shows that if an object $P \in \mathcal{P}^{V(f)}(Q)$ admits a degree one map s satisfying $ds + sd = f$ and $s^2 = 0$, then $\rho_f([P]) = [\text{Fold}(P, d, s)]$. In particular, the map ρ_f is surjective, since for a matrix factorization $(\alpha : P_1 \rightarrow P_0, \beta : P_0 \rightarrow P_1) \in \text{mf}(Q, f)$, we have $(\alpha, \beta) = \text{Fold}(P, \alpha, \beta)$.

Since there exists an isomorphism $G_0(Q/(f)) \xrightarrow{\sim} K_0^{V(f)}(Q)$ which sends the class of a finitely generated $Q/(f)$ -module to the class of a chosen Q -projective resolution of it, we obtain a surjective map

$$G_0(Q/(f)) \twoheadrightarrow K_0(\text{mf}(Q, f)).$$

Note that this surjection agrees with the one induced by the inverse of the equivalence $[\text{mf}(Q, f)] \xrightarrow{\sim} D_{\text{sing}}(Q/(f))$ from [Theorem 1](#) of the introduction.

Given a finitely generated $Q/(f)$ -module M , let $[M]_{\text{stable}} \in K_0(\text{mf}(Q, f))$ denote the image of $[M]$ under the above surjection $G_0(Q/(f)) \twoheadrightarrow K_0(\text{mf}(Q, f))$. Explicitly, for such an M , one may find a Q -projective resolution (P, d) of it for which there exists a degree one endomorphism s of P satisfying $ds + sd = f$ and $s^2 = 0$ (by taking, for instance as above, a good truncation in sufficiently high degree of a K_f -semiprojective resolution $P \xrightarrow{\sim} M$). Then $[M]_{\text{stable}} = [\text{Fold}(P, d, s)]$.

We will use the following result to deduce the diagonalizability of ψ_{cyc}^p on the Grothendieck group of matrix factorizations from the corresponding result for complexes.

Proposition 2.13. *Assume Q is a regular A_p -algebra and $f \in Q$ is a not a zero divisor. The map ρ_f commutes with the Adams operations ψ_{cyc}^p .*

Proof. We need to show the diagram

$$\begin{array}{ccc}
 K_0^{V(f)}(Q) & \xrightarrow{\rho_f} & K_0(\text{mf}(Q, f)) \\
 \downarrow \psi_{\text{cyc}}^p & & \downarrow [Y] \mapsto [T^p(Y)^{(1)}] - [T^p(Y)^{(s)}] \\
 K_0^{V(f)}(Q) & \xrightarrow{\rho_{pf}} & K_0(\text{mf}(Q, pf)) \\
 \downarrow = & & \downarrow \text{mult}_{1/p} \\
 K_0^{V(f)}(Q) & \xrightarrow{\rho_f} & K_0(\text{mf}(Q, f))
 \end{array}$$

commutes.

It suffices to check the commutativity of the top square on classes $[P]$ for which there exists an s with $ds + sd = f$ and $s^2 = 0$. Recall that the induced differential $T^p(d)$ on $T^p(P)$ is given by

$$T^p(d)(x_1 \otimes \cdots \otimes x_p) = \sum_{i=1}^p (-1)^{|x_1| + \cdots + |x_{i-1}|} x_1 \otimes \cdots \otimes d(x_i) \otimes \cdots \otimes x_p,$$

and we define $T^p(s)$ to be the degree one map given by the same formula with s in place of d . Then $T^p(d)T^p(s) + T^p(s)T^p(d) = pf$ and $T^p(s)^2 = 0$. Moreover, it follows from the definitions that there is a canonical isomorphism

$$T^p(\text{Fold}(P, d, s)) \cong \text{Fold}(T^p(P), T^p(d), T^p(s)) \in \text{mf}(Q, pf),$$

and this isomorphism is equivariant for the action of Σ_p . The commutativity of the top square in the diagram follows.

The bottom square commutes by the more general lemma below. □

Lemma 2.14. *If Q is a regular, $f \in Q$ is not a zero divisor, and $u \in Q$ is a unit, the triangle*

$$\begin{array}{ccc}
 & & K_0(\text{mf}(Q, f)) \\
 & \nearrow \rho_f & \downarrow \text{mult}_u \\
 K_0^{V(f)}(Q) & & K_0(\text{mf}(Q, uf)) \\
 & \searrow \rho_{uf} &
 \end{array}$$

commutes.

Proof. Again, it suffices to check the commutativity of the diagram on classes $[P]$ such that P is a complex with differential d for which there exists an s with $ds + sd = f$ and $s^2 = 0$. If $[P]$ is such a class, $\rho_f([P]) = [\text{Fold}(P, d, s)]$.

Before applying ρ_{uf} , first replace (P, d) by the isomorphic complex (P', d') with $P'_i = P_i$ for all i and with $d'_i = d_i$ for i odd and $d'_i = ud_i$ for i even. Defining s' as $s'_i = s_i$ for i odd and $s'_i = us_i$ for i even, one has $d's' + s'd' = uf$. Then $\rho_{uf}([P]) = [\text{Fold}(P', d', s')] = \text{mult}_u([\text{Fold}(P, d, s)]) = (\text{mult}_u \circ \rho_f)([P])$. \square

Theorem 2.15. *Assume Q is a regular A_p -algebra of dimension d and $f \in Q$ is not a zero divisor. There is a decomposition*

$$K_0(\text{mf}(Q, f))_{\mathbb{Q}} = \bigoplus_{i=1}^d K_0(\text{mf}(Q, f))_{\mathbb{Q}}^{(i)},$$

which is independent of p , such that ψ_{cyc}^p acts on $K_0(\text{mf}(Q, f))_{\mathbb{Q}}^{(i)}$ as multiplication by p^i . Moreover, for a finitely generated $Q/(f)$ -module M , we have

$$[M]_{\text{stable}} \in \bigoplus_{i=\text{codim}_{Q/(f)} M+1}^d K_0(\text{mf}(Q, f))_{\mathbb{Q}}^{(i)}.$$

Proof. This follows from Corollary 3.12 of [BMTW 2017] and Proposition 2.13 by defining $K_0(\text{mf}(Q, f))_{\mathbb{Q}}^{(i)}$ to be the image of $K_0^{V(f)}(Q)_{\mathbb{Q}}^{(i)}$ under $\rho_f \otimes \mathbb{Q}$. \square

We close this subsection with a technical result needed below.

Corollary 2.16. *If Q is a regular A_p -algebra for a prime p , $f \in Q$ is not a zero divisor, and $u \in Q$ is a unit, we have an equality of maps $\psi_{\text{cyc}}^p \circ \text{mult}_u = \text{mult}_u \circ \psi_{\text{cyc}}^p$ from $K_0(\text{mf}(Q, f))$ to $K_0(\text{mf}(Q, uf))$.*

Proof. By Proposition 2.13, the diagonal maps in the commutative diagram of Lemma 2.14 commute with the action of ψ_{cyc}^p , and these maps are surjective. \square

3. Dao and Kurano’s Conjecture

In this section, we apply the results of Section 2 to give a proof of Theorem 5 from the introduction.

3A. Some properties of $\mathbb{Z}/2$ -graded complexes. We will need some general results about $\mathbb{Z}/2$ -graded complexes. Much of what we need holds in great generality, and so we start by working over a Noetherian commutative ring B .

Let $\text{LF}(B, 0)$ denote the abelian category of all $\mathbb{Z}/2$ -graded complexes of B -modules (“LF” stands for “linear factorization”), and let $\text{lf}(B, 0)$ denote the full subcategory of $\text{LF}(B, 0)$ consisting of complexes whose components are finitely generated B -modules. An object of $\text{LF}(B, 0)$ consists of a pair of B -modules, M^0 and M^1 , together with maps $d^0 : M^0 \rightarrow M^1$ and $d^1 : M^1 \rightarrow M^0$ such that

$d^1 \circ d^0 = 0 = d^0 \circ d^1$. Morphisms are given by the evident $\mathbb{Z}/2$ -graded analogues of chain maps. We also have the evident $\mathbb{Z}/2$ -versions of quasiisomorphisms and homotopies of chain maps. For objects $X, Y \in \text{LF}(B, 0)$, let $\text{Hom}_{\text{LF}}(X, Y)$ denote the $\mathbb{Z}/2$ -analogue of the mapping complex construction. So $\text{Hom}_{\text{LF}}(X, Y) \in \text{LF}(B, 0)$ with $\text{Hom}_{\text{LF}}(X, Y)^\epsilon = \bigoplus_{\epsilon'+\epsilon''=\epsilon} \text{Hom}_B(X^{\epsilon'}, Y^{\epsilon''})$. Note that the zero cycles in $\text{Hom}_{\text{LF}}(X, Y)$ are, by definition, the set of morphisms from X to Y in $\text{LF}(B, 0)$, and $H^0 \text{Hom}_{\text{LF}}(X, Y)$ is the set of morphisms modulo homotopy.

We write $X \otimes_{\text{LF}} Y \in \text{LF}(B, 0)$ for the evident $\mathbb{Z}/2$ -graded analogue of the tensor product of complexes, so that

$$(X \otimes_{\text{LF}} Y)^\epsilon = \bigoplus_{\epsilon=\epsilon'+\epsilon''} X^{\epsilon'} \otimes_B Y^{\epsilon''}.$$

We will also need the notion of the totalization $\text{Tot}(X.)$ of a bounded complex

$$X. := (0 \rightarrow X_m \rightarrow \dots \rightarrow X_0 \rightarrow 0)$$

of objects of $\text{LF}(B, 0)$, defined in a manner similar to the \mathbb{Z} -graded setting. In more detail, we have

$$\text{Tot}(X.)^\epsilon = \bigoplus_{i=0}^m X_i^{i+\epsilon},$$

with superscripts taken modulo 2. Moreover, if

$$0 \rightarrow X_m \rightarrow \dots \rightarrow X_0 \rightarrow M \rightarrow 0$$

is an exact sequence in $\text{LF}(B, 0)$, then there is a natural quasiisomorphism

$$\text{Tot}(X.) \xrightarrow{\sim} M$$

in $\text{LF}(B, 0)$.

For $M \in \text{LF}(B, 0)$, define $Z(M)$ to be the $\mathbb{Z}/2$ -graded module consisting of the kernels of the two maps comprising the complex M , and define $B(M)$ to be the $\mathbb{Z}/2$ -graded module given by the images of the two maps comprising M . Let $H(M)$ denote the $\mathbb{Z}/2$ -graded module consisting of the homology modules of M . Each of B , Z , and H can be interpreted as a functor from $\text{LF}(B, 0)$ to itself, and they restrict to functors from $\text{lf}(B, 0)$ to itself. Note that $B(M) \subseteq Z(M)$ and $H(M) = Z(M)/B(M)$.

Recall that $\text{mf}(B, 0)$ is the full subcategory of $\text{lf}(B, 0)$ consisting of complexes whose components are projective B -modules.

Definition 3.1. An object $X \in \text{mf}(B, 0)$ is called *proper* if $Z(X)$, $B(X)$ and $H(X)$ are all projective R -modules.

For $M \in \text{lf}(B, 0)$, an exact sequence of the form

$$\dots \rightarrow X_m \rightarrow \dots \rightarrow X_1 \rightarrow X_0 \rightarrow M \rightarrow 0$$

such that $X_i \in \text{mf}(B, 0)$ is proper for all i and each of the induced sequences

$$\begin{aligned} \cdots &\rightarrow B(X_m) \rightarrow \cdots \rightarrow B(X_1) \rightarrow B(X_0) \rightarrow B(M) \rightarrow 0, \\ \cdots &\rightarrow Z(X_m) \rightarrow \cdots \rightarrow Z(X_1) \rightarrow Z(X_0) \rightarrow Z(M) \rightarrow 0, \\ \cdots &\rightarrow H(X_m) \rightarrow \cdots \rightarrow H(X_1) \rightarrow H(X_0) \rightarrow H(M) \rightarrow 0 \end{aligned}$$

is also exact is called a *Cartan–Eilenberg resolution* of M . Such a resolution is *bounded* if $X_j = 0$ for all $j \gg 0$.

Lemma 3.2. *If B is a Noetherian commutative ring, and at least one of $X, Y \in \text{mf}(B, 0)$ is proper, then there is a natural isomorphism*

$$H(X) \otimes_{\text{LF}} H(Y) \xrightarrow{\sim} H(X \otimes_{\text{LF}} Y).$$

Proof. The proof is the same as for the classical Künneth Theorem. □

Lemma 3.3. *If B is a Noetherian commutative ring, then every $M \in \text{lf}(B, 0)$ admits a Cartan–Eilenberg resolution. If B is regular, every $M \in \text{lf}(B, 0)$ admits a bounded Cartan–Eilenberg resolution.*

Proof. Choose projective resolutions of $B^0(M)$, $B^1(M)$, $H^0(M)$ and $H^1(M)$, and make repeated use of the horseshoe lemma, just as in the proof of the classical version of this result. If B is regular, all of the chosen projective resolutions in the proof may be chosen to be bounded. □

Recall that $[\text{mf}(B, 0)]$ denotes the category with the same objects as $\text{mf}(B, 0)$ and with morphism sets given by $\text{Hom}_{[\text{mf}(B, 0)]}(X, Y) := H^0(\text{Hom}_{\text{LF}}(X, Y))$. We write $\mathcal{D}(\text{lf}(B, 0))$ for the category obtained from $\text{lf}(B, 0)$ by inverting all quasiisomorphisms.

Proposition 3.4. *If B is regular, the canonical functor*

$$[\text{mf}(B, 0)] \xrightarrow{\sim} \mathcal{D}(\text{lf}(B, 0))$$

is an equivalence.

Proof. Let M be an object in $\mathcal{D}(\text{lf}(B, 0))$. Applying [Lemma 3.3](#), choose a bounded Cartan–Eilenberg resolution X of M . Then the canonical map $\text{Tot}(X) \rightarrow M$ is a quasiisomorphism, and $\text{Tot}(X)$ is an object of $\text{mf}(B, 0)$; thus, the functor is essentially surjective. It is fully faithful by [Lemma 2.3](#). □

We are especially interested in complexes with finite length homology. Let $\text{lf}^{\text{fl}}(B, 0)$ and $\text{mf}^{\text{fl}}(B, 0)$ denote the full subcategories of $\text{lf}(B, 0)$ and $\text{mf}(B, 0)$ consisting of those complexes M such that $H^0(M)$ and $H^1(M)$ are finite length B -modules. Since this condition is preserved by quasiisomorphism, we may form $[\text{mf}^{\text{fl}}(B, 0)]$ and $\mathcal{D}(\text{lf}^{\text{fl}}(B, 0))$, and they may be identified as full subcategories of

$[\mathrm{mf}(B, 0)]$ and $\mathcal{D}(\mathrm{lf}(B, 0))$. Moreover, it follows from [Proposition 3.4](#) that the canonical functor induces an equivalence

$$[\mathrm{mf}^{\mathrm{fl}}(B, 0)] \xrightarrow{\sim} \mathcal{D}(\mathrm{lf}^{\mathrm{fl}}(B, 0)),$$

provided B is regular.

It will be convenient to give an alternative description of the category $\mathrm{LF}(B, 0)$ and of the constructions just described. Fix a degree two indeterminate t and form the \mathbb{Z} -graded algebra $\tilde{B} := B[t, t^{-1}]$, which we regard as a dg-ring with trivial differential. Recall that a dg- \tilde{B} -module is a graded \tilde{B} -module M equipped with a degree one \tilde{B} -linear map $d : M \rightarrow M$ such that $d^2 = 0$. Since t is a degree two invertible element, a dg- \tilde{B} -module is the same things as a \mathbb{Z} -graded complex of B -modules M together with a specified isomorphism $t : M \xrightarrow{\sim} M[2]$ of complexes. A morphism between two such pairs, say from (M, t) to (M', t') , is a chain map from M to M' that commutes with t and t' . There is an evident equivalence of abelian categories

$$\mathrm{dg}\text{-}\tilde{B}\text{-Mod} \xrightarrow{\sim} \mathrm{LF}(B, 0)$$

that sends a dg- \tilde{B} -module M to the object

$$(M^0 \xrightarrow{d} M^1 \xrightarrow{t^{-1}d} M^0)$$

of $\mathrm{LF}(B, 0)$. Moreover, the notions of mapping complex, tensor product, quasi-isomorphism, homotopy equivalence and totalization defined above for $\mathrm{LF}(B, 0)$ correspond to the standard notions for dg-modules. This equivalence thus allows us to employ standard results from differential graded algebra.

3B. Adams operations on $\mathbb{Z}/2$ -graded complexes with finite length homology.

Let Q be a regular local ring with maximal ideal \mathfrak{m} . Recall that $\mathrm{mf}^{\mathrm{m}}(Q, 0)$ is the category of $\mathbb{Z}/2$ -graded complexes of finite rank free Q -modules whose homology has support in $\{\mathfrak{m}\}$; notice that $\mathrm{mf}^{\mathrm{m}}(Q, 0) = \mathrm{mf}^{\mathrm{fl}}(Q, 0)$, where the right-hand side is as defined in [Section 3A](#).

Recall that $K_0^{\mathrm{m}}(Q)$ is the Grothendieck group of the category of bounded \mathbb{Z} -graded complexes of projective Q -modules whose homology has support in $\{\mathfrak{m}\}$. It is easy to prove that $K_0^{\mathrm{m}}(Q)$ is a free abelian group of rank one, generated by the class of the Koszul complex on a regular system of generators of \mathfrak{m} . One might thus expect the answer to the following question to be positive:

Question 3.5. *For a regular local ring (Q, \mathfrak{m}) , is $K_0(\mathrm{mf}^{\mathrm{m}}(Q, 0))$ a free abelian group of rank one, generated by the $\mathbb{Z}/2$ -folded Koszul complex?*

We know the answer to be “yes” if $\dim(Q) \leq 2$, but the general situation remains unknown. The following example illustrates the difficulty:

Example 3.6. Let (Q, \mathfrak{m}) be a regular local ring of dimension three, and suppose x, y, z form a regular sequence of generators for the maximal ideal \mathfrak{m} . Let

$$0 \rightarrow Q \xrightarrow{i} Q^3 \xrightarrow{A} Q^3 \xrightarrow{p} Q \rightarrow 0$$

be the usual Koszul complex on x, y, z (so that, for example, p is given by the row matrix (x, y, z)). The $\mathbb{Z}/2$ -folding of this Koszul complex,

$$K := \left(Q^3 \oplus Q \xrightarrow{\begin{bmatrix} A & 0 \\ 0 & 0 \end{bmatrix}} Q^3 \oplus Q \xrightarrow{\begin{bmatrix} 0 & i \\ p & 0 \end{bmatrix}} Q^3 \oplus Q \right),$$

determines a class $[K]$ in $K_0(\text{mf}^{\text{m}}(Q, 0))$.

Now define $B : Q^3 \rightarrow Q^3$ to be the map $i \circ p$. Then $AB = 0 = BA$, so that $X = (Q^3 \xrightarrow{A} Q^3 \xrightarrow{B} Q^3)$ is a $\mathbb{Z}/2$ -graded complex. Moreover, $\ker(B) = \text{im}(A)$ and $\ker(A)/\text{im}(B) \cong Q/\mathfrak{m}$, so that $X \in \text{mf}^{\text{m}}(Q, 0)$. We do not know whether $[X]$ is a multiple of $[K]$ in $K_0(\text{mf}^{\text{m}}(Q, 0))$.

To explain the relevance of [Question 3.5](#), let us define the *Euler characteristic* of an object $X \in \text{mf}^{\text{m}}(Q, 0)$ to be

$$\chi(X) = \text{length } H^0(X) - \text{length } H^1(X).$$

Then χ determines a group homomorphism

$$\chi : K_0(\text{mf}^{\text{m}}(Q, 0)) \rightarrow \mathbb{Z}.$$

For example, if K is the $\mathbb{Z}/2$ -folded Koszul complex on a regular system of generators for \mathfrak{m} , then $\chi(K) = 1$. Assume now that Q is a regular local A_p -algebra for a prime p (that is, assume p is invertible in Q and that Q contains a primitive p -th root of unity), so that the cyclic Adams operation ψ_{cyc}^p acts on $K_0(\text{mf}^{\text{m}}(Q, 0))$. We have $\psi_{\text{cyc}}^p([K]) = p^d [K]$, where $d = \dim(Q)$, by [Corollary 2.11](#). If the answer to [Question 3.5](#) were affirmative, we would obtain as an immediate consequence the identity

$$\chi \circ \psi_{\text{cyc}}^p = p^d \chi \tag{3.7}$$

of maps from $K_0(\text{mf}^{\text{m}}(Q, 0))$ to \mathbb{Z} . Moreover, this equation plays a key role in the proof of [Theorem 5](#).

Although we are unable to answer [Question 3.5](#), we are nevertheless able to prove an analogue to [\[Gillet and Soulé 1987, Proposition 7.1\]](#).

Theorem 3.8. *For a regular local ring Q of dimension d that is an A_p -algebra for some prime p , (3.7) holds.*

The proof of this theorem occupies the remainder of this subsection.

Fix a prime p , and let B be a commutative Noetherian A_p -algebra. Recall the functor t_ζ^p defined on $\text{mf}(B, 0)$ that sends X to $T^p(X)^{(\zeta)}$, where ζ is a p -th root of unity. It will be useful to interpret this functor as a composition

$$\text{mf}(B, 0) \xrightarrow{T^p} \text{mf}(B', 0) \xrightarrow{Y \mapsto Y^{(\zeta)}} \text{mf}(B, 0)$$

where we set $B' = B[C_p] = B[\sigma]/(\sigma^p - 1)$. Since B is an A_p -algebra, B' is isomorphic to a product of p copies of B equipped with an action of C_p . So, an object of $\text{mf}(B', 0)$ is the same thing as an object of $\text{mf}(B, 0)$ equipped with an action of C_p , and if B is regular, then so is B' .

The functors above preserve the condition that homology has finite length, and they send homotopic maps to homotopic maps, so that we have an induced functor

$$t_\zeta^p : [\text{mf}^{\text{fl}}(B, 0)] \rightarrow [\text{mf}^{\text{fl}}(B, 0)]$$

given as the composition of functors

$$[\text{mf}^{\text{fl}}(B, 0)] \xrightarrow{T^p} [\text{mf}^{\text{fl}}(B', 0)] \xrightarrow{Y \mapsto Y^{(\zeta)}} [\text{mf}^{\text{fl}}(B, 0)].$$

We will need a “derived” version of the functor t_ζ^p . When B is regular, then we may use the equivalence of [Proposition 3.4](#) to obtain a functor

$$\mathbf{t}_\zeta^p : \mathcal{D}(\text{lf}^{\text{fl}}(B, 0)) \rightarrow [\text{mf}^{\text{fl}}(B, 0)].$$

Explicitly, for $M \in \text{lf}^{\text{fl}}(B, 0)$, $\mathbf{t}_\zeta^p(M) = t_\zeta^p(P)$ where P is any object of $\text{mf}^{\text{fl}}(B, 0)$ for which there exists a quasiisomorphism $P \xrightarrow{\sim} M$.

Given $M \in \text{lf}(B, 0)$, recall that $H(M)$ denotes the object of $\text{lf}(B, 0)$ given by the $\mathbb{Z}/2$ -graded B -module with components $H^0(M)$ and $H^1(M)$, regarded as a complex with trivial differential. In terms of the dg-ring \tilde{B} , $H(M)$ corresponds to the homology of a dg- \tilde{B} -module, which is naturally a dg- \tilde{B} -module with trivial differential (since \tilde{B} has trivial differential). If $M \in \text{lf}^{\text{fl}}(B, 0)$, we define its Euler characteristic by

$$\chi(M) := \text{length } H^0(M) - \text{length } H^1(M),$$

as above.

Lemma 3.9. *If B is a regular A_p -algebra, then for any $M \in \text{lf}^{\text{fl}}(B, 0)$ and any p -th root of unity ζ , we have*

$$\chi(\mathbf{t}_\zeta^p(M)) = \chi(\mathbf{t}_\zeta^p(H(M))).$$

[Theorem 3.8](#) is a relatively easy consequence of [Lemma 3.9](#). Before proving [Lemma 3.9](#), we must introduce the following notation and establish one more preliminary result. For a bounded complex

$$X = (0 \rightarrow X_m \rightarrow X_{m-1} \rightarrow \cdots \rightarrow X_1 \rightarrow X_0 \rightarrow 0)$$

of objects of $\text{LF}(B, 0)$, we write $\mathcal{H}_q(X.) \in \text{LF}(B, 0)$ for its homology taken in the abelian category $\text{LF}(B, 0)$; that is,

$$\mathcal{H}_q(X.) = \ker(X_q \rightarrow X_{q-1}) / \text{im}(X_{q+1} \rightarrow X_q).$$

We write $H(X.)$ for the complex of objects of $\text{LF}(B, 0)$ obtained by applying H term-wise

$$H(X.) := (0 \rightarrow H(X_d) \rightarrow \cdots \rightarrow H(X_0) \rightarrow 0).$$

Note that $H(X.)$ is a complex of $\mathbb{Z}/2$ -graded modules, and we regard it as another complex of objects in $\text{LF}(B, 0)$.

Lemma 3.10. *For a Noetherian commutative ring B , assume*

$$Y. := (0 \rightarrow Y_m \rightarrow \cdots \rightarrow Y_0 \rightarrow 0)$$

is a complex in $\text{lf}(B, 0)$ such that both $\mathcal{H}_q H(Y.)$ and $H \mathcal{H}_q(Y.)$ have finite length for all q . Then $\text{Tot}(Y.)$ belongs to $\text{lf}^{\text{fl}}(B, 0)$, and we have

$$\begin{aligned} \chi(\text{Tot}(Y.)) &= \sum_{q \in \mathbb{Z}, \epsilon \in \mathbb{Z}/2} (-1)^{q+\epsilon} \text{length } \mathcal{H}_q(H^\epsilon(Y.)) \\ &= \sum_{q \in \mathbb{Z}, \epsilon \in \mathbb{Z}/2} (-1)^{q+\epsilon} \text{length } H^\epsilon(\mathcal{H}_q(Y.)). \end{aligned}$$

Proof. Our proof uses spectral sequences and is similar to the proof of the analogous fact concerning \mathbb{Z} -graded bicomplexes, but some care is needed to deal with the $\mathbb{Z}/2$ -grading.

We find it most convenient to work in the setting of $\text{dg-}\tilde{B}$ -modules. Recall that a $\text{dg-}\tilde{B}$ -module is the same thing as pair consisting of a \mathbb{Z} -graded complex of B -modules and a degree 2 automorphism. A graded \tilde{B} -module is a $\text{dg-}\tilde{B}$ -module with trivial differential.

Let us say that a graded \tilde{B} -module H has *finite length* if H^i has finite length as a B -module for each $i \in \mathbb{Z}$ (or, equivalently, for $i = 0, 1$). In this case, we define

$$\tilde{\chi}(H) = \text{length}_B(H^0) - \text{length}_B(H^1).$$

(Note that $\tilde{\chi}(H) = \text{length}_B(H^{2m}) - \text{length}_B(H^{2n+1})$ for any $m, n \in \mathbb{Z}$.) It is clear that if $Y \in \text{lf}^{\text{fl}}(B, 0)$, then

$$\chi(Y) = \tilde{\chi}(\tilde{H}(Y))$$

where χ is as defined before, and $\tilde{H}(Y)$ denotes the homology of Y regarded in the canonical way as a graded \tilde{B} -module.

We will need the following fact. If (M, d) is a $\text{dg-}\tilde{B}$ -module such that the underlying graded \tilde{B} -module M has finite length, then $H(M, d)$ also has finite length, and $\tilde{\chi}(H(M, d)) = \tilde{\chi}(M)$. This is seen to hold by a straightforward calculation.

We view Y as a bicomplex Y_{\cdot} with $m + 1$ rows, whose m -th row, for $0 \leq j \leq m$, is

$$\dots \rightarrow Y_j^{-1} \rightarrow Y_j^0 \rightarrow Y_j^1 \rightarrow \dots,$$

along with a degree $(2, 0)$ isomorphism of bicomplexes $t : Y_{\cdot} \xrightarrow{\sim} Y_{\cdot}^{+2}$. Since this bicomplex is uniformly bounded in the vertical direction, we have two strongly convergent spectral sequences of the form

$$\begin{aligned} 'E_2^{p,-q} = H_q(H^p(Y_{\cdot})) &\implies H^{p-q}(\text{Tot}(Y_{\cdot})) \quad \text{and} \\ ''E_2^{p,-q} = H^p(H_q(Y_{\cdot})) &\implies H^{p-q}(\text{Tot}(Y_{\cdot})). \end{aligned}$$

Let $E_r^{*,*}$, for $r \geq 2$, refer to either of these two spectral sequences. The isomorphism $t : Y_{\cdot} \xrightarrow{\sim} Y_{\cdot}^{+2}$ induces isomorphisms

$$t : E_r^{p,-q} \xrightarrow{\sim} E_r^{p+2,-q}$$

for each $r \geq 2$, and similarly on the underlying D_r -terms, and these isomorphisms commute with all the maps of the exact couple.

For any r , define a \mathbb{Z} -graded B -module $\text{Tot}(E_r)$ by

$$\text{Tot}(E_r)^n := \bigoplus_{p+q=n} E_r^{p,q}.$$

The isomorphism t induces an isomorphism of degree 2 on $\text{Tot}(E_r)$ making it into a graded \tilde{B} -module. For each r , the differential d_r on the E_r 's induces a degree one map (which we will also write as d_r) on $\text{Tot}(E_r)$, and since this map commutes with t , we have that $(\text{Tot}(E_r), d_r)$ is a dg- \tilde{B} -module. Finally, we have an identity

$$\text{Tot}(E_{r+1}) = H(\text{Tot}(E_r), d_r)$$

of graded \tilde{B} -modules.

Returning to the two specific instances of this spectral sequence, the assumptions give that each of $\text{Tot}'(E_2)$ and $\text{Tot}''(E_2)$ has finite length, and that we have

$$\begin{aligned} \tilde{\chi}(\text{Tot}'(E_2)) &= \sum_{q \in \mathbb{Z}, \epsilon \in \mathbb{Z}/2} (-1)^{q+\epsilon} \text{length } \mathcal{H}_q(H^\epsilon(Y)) \\ \tilde{\chi}(\text{Tot}''(E_2)) &= \sum_{q \in \mathbb{Z}, \epsilon \in \mathbb{Z}/2} (-1)^{q+\epsilon} \text{length } H^\epsilon(\mathcal{H}_q(Y)). \end{aligned} \tag{3.11}$$

By the general fact mentioned above, we get that each of $\text{Tot}(E_3), \text{Tot}(E_4), \dots$ also has finite length, and, moreover,

$$\tilde{\chi}(\text{Tot}(E_2)) = \tilde{\chi}(\text{Tot}(E_3)) = \dots = \tilde{\chi}(\text{Tot}(E_\infty)).$$

(Note that the spectral sequence degenerates after at most $m + 2$ steps, so that $E_{m+2} = E_{m+3} = \dots = E_\infty$.)

Now, for $\epsilon = 0, 1$, the B -module $H^\epsilon \text{Tot}(Y)$ admits a filtration by B -submodules whose subquotients are $E_\infty^{\epsilon,0}, E_\infty^{\epsilon-1,1}, \dots, E_\infty^{\epsilon-m,m}$, and hence

$$\begin{aligned} \chi(\text{Tot}(Y)) &= \tilde{\chi}(H(\text{Tot}(Y))) \\ &= \sum_q \text{length } E_\infty^{-q,q} - \sum_q \text{length } E_\infty^{1-q,q} \\ &= \tilde{\chi}(\text{Tot}(E_\infty)) = \tilde{\chi}(\text{Tot}(E_2)). \end{aligned}$$

By (3.11), the proof is complete. □

Proof of Lemma 3.9. We may assume, without loss of generality, that $M = P$ belongs to $\text{mf}^{\text{fl}}(B, 0)$. Let

$$\dots \rightarrow 0 \rightarrow X_m \rightarrow X_{m-1} \rightarrow \dots \rightarrow X_1 \rightarrow X_0 \rightarrow P \rightarrow 0$$

be a bounded Cartan–Eilenberg resolution of P . Since P is an object of $\text{mf}(B, 0)$, the induced quasiisomorphism $\text{Tot}(X.) \xrightarrow{\sim} P$ is a homotopy equivalence, a fact that will be used below.

Recall that X_i is proper. In particular, $H(X_i)$ is projective for all i , and the induced complex

$$\dots \rightarrow 0 \rightarrow H(X_m) \rightarrow H(X_{m-1}) \rightarrow \dots \rightarrow H(X_1) \rightarrow H(X_0) \rightarrow H(P) \rightarrow 0$$

is also exact. The latter gives, by definition,

$$\mathbf{t}_\zeta^p(H(P)) = \mathbf{t}_\zeta^p(\text{Tot}(H(X.))) = T^p(\text{Tot}(H(X.)))^{(\zeta)}. \tag{3.12}$$

For any bounded complex $Y.$ of objects of $\text{mf}(B, 0)$, write $T^p(Y.)$ for the complex of objects in $\text{mf}(B, 0)$ that, in degree j , is

$$T^p(Y.)_j = \bigoplus_{i_1+\dots+i_p=j} Y_{i_1} \otimes_{\text{LF}} \dots \otimes_{\text{LF}} Y_{i_p}.$$

For example, if $p = 2$, then $T^2(Y.)$ is the complex

$$\dots \rightarrow (Y_2 \otimes Y_0 \oplus Y_1 \otimes Y_1 \oplus Y_0 \otimes Y_2) \rightarrow (Y_1 \otimes Y_0 \oplus Y_0 \otimes Y_1) \rightarrow Y_0 \otimes Y_0 \rightarrow 0.$$

Each term of the complex $T^p(Y.)$ admits an evident signed action by C_p , and the maps of this complex respect these actions, so that we may regard $T^p(Y.)$ as a complex in $\text{mf}(B', 0)$, where $B' := B[C_p]$. We have an identity

$$T^p(\text{Tot}(Y.)) = \text{Tot}(T^p(Y.)) \tag{3.13}$$

of objects of $\text{mf}(B', 0)$.

Since B is an A_p -algebra, $(-)^{(\zeta)}$ is an exact functor from $\text{lf}(B', 0)$ to $\text{lf}(B, 0)$. In fact, B' is a product of copies of B , and this functor is given by extension of

scalars along one of the canonical projections $B' \rightarrow B$. In particular, we have

$$\text{Tot}(Y)_*^{(\zeta)} = \text{Tot}(Y_*^{(\zeta)}) \tag{3.14}$$

for any bounded complex Y of objects of $\text{lf}(B', 0)$, and

$$\text{H}(Y)_*^{(\zeta)} = \text{H}(Y_*^{(\zeta)}) \tag{3.15}$$

for any object $Y \in \text{lf}(B', 0)$.

Since each X_i is proper, [Lemma 3.2](#) implies that we have canonical isomorphisms

$$\text{H}(X_{i_1}) \otimes_{\text{LF}} \cdots \otimes_{\text{LF}} \text{H}(X_{i_p}) \xrightarrow{\cong} \text{H}(X_{i_1} \otimes_{\text{LF}} \cdots \otimes_{\text{LF}} X_{i_p})$$

which combine to give an isomorphism

$$T^p(\text{H}(X_*)) \xrightarrow{\cong} \text{H}(T^p(X_*)) \tag{3.16}$$

of complexes of objects of $\text{mf}(B', 0)$.

Combining these facts gives

$$\begin{aligned} \mathbf{t}_\zeta^p(\text{H}(P)) &= T^p(\text{Tot}(\text{H}(X_*)))^{(\zeta)}, && \text{by (3.12),} \\ &= (\text{Tot}(T^p(\text{H}(X_*))))^{(\zeta)}, && \text{by (3.13),} \\ &= \text{Tot}(T^p(\text{H}(X_*)))^{(\zeta)}, && \text{by (3.14),} \\ &= \text{Tot}(\text{H}(T^p(X_*)))^{(\zeta)}, && \text{by (3.16),} \\ &= \text{Tot}(\text{H}(T^p(X_*))^{(\zeta)}), && \text{by (3.15).} \end{aligned}$$

We now apply [Lemma 3.10](#) to the complex $Y := T^p(X_*)^{(\zeta)}$ of objects in $\text{mf}(B, 0)$, which gives

$$\sum_{q, \epsilon} (-1)^{q+\epsilon} \text{length } \mathcal{H}_q(\text{H}^\epsilon(Y_*)) = \sum_{q, \epsilon} (-1)^{q+\epsilon} \text{length } \text{H}^\epsilon(\mathcal{H}_q(Y_*)). \tag{3.17}$$

Since we have shown that $\text{Tot}(\text{H}(Y_*)) \cong \mathbf{t}_\zeta^p(\text{H}(P))$, the left-hand side of (3.17) is $\chi(\mathbf{t}_\zeta^p(\text{H}(P)))$.

Recall that, since P belongs to $\text{mf}(B, 0)$, the quasiisomorphism $\text{Tot}(X_*) \xrightarrow{\cong} P$ is a homotopy equivalence. It follows that the map

$$\text{Tot}(Y_*) \cong T^p(\text{Tot}(X_*))^{(\zeta)} \rightarrow T^p(P)^{(\zeta)}$$

is also a homotopy equivalence. We get

$$\text{H}^\epsilon(\mathcal{H}_q(Y_*)) \cong \begin{cases} \text{H}^\epsilon(\mathbf{t}_\zeta^p(P)) & \text{if } q = 0, \\ 0 & \text{otherwise,} \end{cases}$$

which shows that the right-hand side of (3.17) is $\chi(\mathbf{t}_\zeta^p(P))$. □

Proof of Theorem 3.8. Let $P \in \text{mf}^m(Q, 0) = \text{mf}^{\text{fl}}(Q, 0)$. By definition,

$$\chi(\psi_{\text{cyc}}^p([P])) = \sum_{\zeta} \zeta \chi(t_{\zeta}^p(P)).$$

By Lemma 3.9, the value of the right-hand side of this equation coincides with $\sum_{\zeta} \zeta \chi(t_{\zeta}^p(H(P)))$. Since $H(P)$ has trivial differential, the class

$$[H(P)] \in K_0(\mathcal{D}(\text{lf}^{\text{fl}}(Q, 0))) \cong K_0(\text{mf}^m(Q, 0))$$

is an integer multiple of the class of the residue field $k = Q/\mathfrak{m}$, which in turn coincides with the class of the folded Koszul complex $K \in \text{mf}^m(Q, 0)$. This proves that the equation of Theorem 3.8 holds in general provided it holds for the class $[K]$, and that special case is known to hold by Corollary 2.11. \square

3C. Proof of the conjecture. Throughout this section, we assume (Q, \mathfrak{m}) is a regular local ring and f is a nonzero element of \mathfrak{m} , and we set $R = Q/(f)$. We also assume R is an isolated singularity; that is, we assume $R_{\mathfrak{p}}$ is regular for all $\mathfrak{p} \in \text{Spec}(R) \setminus \{\mathfrak{m}\}$. Recall from the introduction that these conditions lead to a well-defined invariant for a pair (M, N) of finitely generated R -modules:

$$\theta_R(M, N) = \text{length}(\text{Tor}_{2n}^R(M, N)) - \text{length}(\text{Tor}_{2n+1}^R(M, N))$$

for $n \gg 0$.

For a finitely generated R -module M , $[M]_{\text{stable}}$ denotes its associated class in $K_0(\text{mf}(Q, f))$, given by the surjection $G_0(R) \twoheadrightarrow K_0(\text{mf}(Q, f))$ described in Section 2C. Recall that $[M]_{\text{stable}} = [\text{Fold}(P, d, s)]$, where P is a Q -projective resolution of M admitting a degree one endomorphism s that satisfies $ds + sd = f$ and $s^2 = 0$, that is, a Koszul resolution.

For a matrix factorization $X \in \text{mf}(Q, f)$, write X° for $\text{mult}_{-1} X \in \text{mf}(Q, -f)$. That is, if $X = (\alpha : P_1 \rightarrow P_0, \beta : P_0 \rightarrow P_1)$, then $X^\circ = (\alpha, -\beta)$. We also use the notation $(-)^\circ$ to denote the induced isomorphism $K_0(\text{mf}(Q, f)) \xrightarrow{\sim} K_0(\text{mf}(Q, -f))$. For a finitely generated R -module N , the class $[N]_{\text{stable}}^\circ$ is the image of $[N]$ under $G_0(R) \twoheadrightarrow K_0(\text{mf}(Q, -f))$, using that $Q/(f) = Q/(-f)$.

Proposition 3.18. For Q, \mathfrak{m}, f, R, M and N as in Definition 1.2,

$$\theta_R(M, N) = \chi([M]_{\text{stable}} \cup [N]_{\text{stable}}^\circ).$$

Proof. First note that, since f is an isolated singularity, one has

$$K_0(\text{mf}(Q, \pm f)) = K_0(\text{mf}^m(Q, \pm f))$$

and hence

$$[M]_{\text{stable}} \cup [N]_{\text{stable}}^\circ \in K_0(\text{mf}^m(Q, f + (-f))) = K_0(\text{mf}^m(Q, 0)).$$

Choose matrix factorizations $X = (d_1 : X_1 \rightarrow X_0, d_0 : X_0 \rightarrow X_1)$ and $Y = (d'_1 : Y_1 \rightarrow Y_0, d'_0 : Y_0 \rightarrow Y_1)$ such that $[X] = [M]_{\text{stable}}$ and $[Y] = [N]_{\text{stable}}^\circ$. Assume, without loss of generality, that N is maximal Cohen–Macaulay, and $N = \text{coker}(d'_1)$.

Let Z denote the object $(0 \rightarrow N, N \rightarrow 0)$ of $\text{lf}(Q, -f)$; here, 0 is in odd degree and N is in even degree. Let $\alpha : Y \rightarrow Z$ be the morphism in $\text{lf}(Q, -f)$ given by the canonical surjection in even degree and, of course, the zero map in odd degree. Since $\theta(M, N)$ clearly coincides with the Euler characteristic of $X \otimes Z$, it suffices to show that the morphism

$$\text{id} \otimes \alpha : X \otimes Y \rightarrow X \otimes Z$$

in $\text{lf}(Q, 0)$ is a quasiisomorphism. The map $\text{id} \otimes \alpha$ is clearly surjective, so it suffices to show that its kernel is acyclic. An easy calculation shows that $\ker(\text{id} \otimes \alpha) \cong X \otimes T$, where T is the object $(Y_1 \xrightarrow{\text{id}} Y_1, Y_1 \xrightarrow{-f} Y_1) \in \text{lf}(Q, -f)$. Since T is contractible, $X \otimes T$ is contractible; thus, $\text{id} \otimes \alpha$ is a quasiisomorphism. \square

We now prove the conjecture of Dao and Kurano:

Theorem 3.19. *Let (Q, \mathfrak{m}) be a regular local ring and $f \in \mathfrak{m}$ a nonzero element, and assume $R := Q/(f)$ is an isolated singularity. If M and N are finitely generated R -modules such that*

$$\dim M + \dim N \leq \dim R$$

then $\theta_R(M, N) = 0$.

Proof. Let p be any prime that is invertible in Q . We start by reducing to the case where Q contains a primitive p -th root of unity. If not, we form the faithfully flat extension $Q \subseteq Q'$ where Q' is the localization of $Q[x]/(x^p - 1)$ at any one of the maximal ideals lying over \mathfrak{m} , and set $R' = Q'/f \cong R \otimes_Q Q'$. Note that $R \subseteq R'$ is also faithfully flat, and thus

$$\text{Tor}_i^R(M, N) \otimes_R R' \cong \text{Tor}_i^{R'}(M \otimes_R R', N \otimes_R R').$$

It follows that

$$\theta_{R'}(M \otimes_R R', N \otimes_R R') = [R'/\mathfrak{m}' : R/\mathfrak{m}] \cdot \theta_R(M, N),$$

and so we may replace Q with Q' .

Set $d = \dim Q$, $c_M = \text{codim}_Q M$ and $c_N = \text{codim}_Q N$. The hypothesis that $\dim M + \dim N \leq \dim R = d - 1$ yields $c_M + c_N \geq d + 1$. By [Theorem 2.15](#), the classes $[M]_{\text{stable}}, [N]_{\text{stable}} \in K_0(\text{mf}(Q, f)) \otimes \mathbb{Q}$ decompose uniquely as

$$[M]_{\text{stable}} = \sum_{i=c_M}^d X_i \quad \text{and} \quad [N]_{\text{stable}} = \sum_{j=c_N}^d Y_j,$$

where X_i and Y_j are such that $\psi_{\text{cyc}}^p(X_i) = p^i X_i$ and $\psi_{\text{cyc}}^p(Y_j) = p^j Y_j$. Then

$$[N]_{\text{stable}}^{\circ} = \sum_{j=c_N}^d Y_j^{\circ}$$

and, by [Corollary 2.16](#), $\psi_{\text{cyc}}^p(Y_j^{\circ}) = p^j Y_j^{\circ}$ for all j .

By [Proposition 3.18](#), we have

$$\theta_R(M, N) = \chi([M]_{\text{stable}} \cup [N]_{\text{stable}}^{\circ}) = \sum_{i,j} \chi(X_i \cup Y_j^{\circ}),$$

and so it suffices to prove $\chi(X_i \cup Y_j^{\circ}) = 0$ for all i and j . For any i and j ,

$$\begin{aligned} p^d \chi(X_i \cup Y_j^{\circ}) &= \chi(\psi_{\text{cyc}}^p(X_i \cup Y_j^{\circ})) \\ &= \chi(\psi_{\text{cyc}}^p(X_i) \cup \psi_{\text{cyc}}^p(Y_j^{\circ})) \\ &= \chi(p^i X_i \cup p^j Y_j^{\circ}) \\ &= p^{i+j} \chi(X_i \cup Y_j^{\circ}), \end{aligned}$$

where the first equality is by [Theorem 3.8](#), the second is by [Theorem 2.10](#), and the third is by definition of X_i and Y_j . Since [Theorem 2.15](#) yields that $i + j \geq c_M + c_N > d$, we conclude that $\chi(X_i \cup Y_j^{\circ}) = 0$. \square

Acknowledgements

We thank Luchezar Avramov for helpful conversations in preparing this paper. We thank Dave Benson, Oliver Hauton, and Bernhard Köck for leading us to relevant literature [[Benson 1984](#); [Hauton 2009](#); [Köck 1997](#)] and Paul Roberts for making available to us his unpublished 1996 lecture notes on Adams operations.

References

- [Atiyah 1966] M. F. Atiyah, “Power operations in K -theory”, *Quart. J. Math. Oxford Ser. (2)* **17** (1966), 165–193. [MR Zbl](#)
- [Benson 1984] D. J. Benson, “Lambda and psi operations on Green rings”, *J. Algebra* **87**:2 (1984), 360–367. [MR Zbl](#)
- [BMTW 2017] M. K. Brown, C. Miller, P. Thompson, and M. E. Walker, “Cyclic Adams operations”, *J. Pure Appl. Algebra* **221**:7 (2017), 1589–1613. [MR Zbl](#)
- [Buchweitz 1986] R.-O. Buchweitz, “Maximal Cohen–Macaulay modules and Tate-cohomology over Gorenstein rings”, preprint, Univ. Hannover, 1986, <http://tinyurl.com/ragnarolaf>.
- [Buchweitz and Van Straten 2012] R.-O. Buchweitz and D. Van Straten, “An index theorem for modules on a hypersurface singularity”, *Mosc. Math. J.* **12**:2 (2012), 237–259. [MR Zbl](#)
- [Dao 2013] H. Dao, “Decent intersection and Tor-rigidity for modules over local hypersurfaces”, *Trans. Amer. Math. Soc.* **365**:6 (2013), 2803–2821. [MR Zbl](#)

- [Dao and Kurano 2014] H. Dao and K. Kurano, “Hochster’s theta pairing and numerical equivalence”, *J. K-Theory* **14**:3 (2014), 495–525. MR Zbl
- [Eisenbud 1980] D. Eisenbud, “Homological algebra on a complete intersection, with an application to group representations”, *Trans. Amer. Math. Soc.* **260**:1 (1980), 35–64. MR Zbl
- [Gillet and Soulé 1987] H. Gillet and C. Soulé, “Intersection theory using Adams operations”, *Invent. Math.* **90**:2 (1987), 243–277. MR Zbl
- [Hauton 2009] O. Hauton, *Steenrod operations and quadratic forms*, Ph.D. thesis, Université Paris VI, 2009, <http://www.mathematik.uni-muenchen.de/~hauton/these.pdf>.
- [Hochster 1981] M. Hochster, “The dimension of an intersection in an ambient hypersurface”, pp. 93–106 in *Algebraic geometry* (Chicago, 1980), edited by A. Libgober and P. Wagreich, Lecture Notes in Math. **862**, Springer, Berlin, 1981. MR Zbl
- [Köck 1997] B. Köck, “Adams operations for projective modules over group rings”, *Math. Proc. Cambridge Philos. Soc.* **122**:1 (1997), 55–71. MR Zbl
- [Moore et al. 2011] W. F. Moore, G. Piepmeyer, S. Spiroff, and M. E. Walker, “Hochster’s theta invariant and the Hodge–Riemann bilinear relations”, *Adv. Math.* **226**:2 (2011), 1692–1714. MR Zbl
- [Orlov 2004] D. O. Orlov, “Triangulated categories of singularities and D-branes in Landau–Ginzburg models”, *Tr. Mat. Inst. Steklova* **246** (2004), 240–262. In Russian; translated in *Proc. Steklov Inst. Math.* **246** (2004), 227–248. MR Zbl
- [Polishchuk and Vaintrob 2012] A. Polishchuk and A. Vaintrob, “Chern characters and Hirzebruch–Riemann–Roch formula for matrix factorizations”, *Duke Math. J.* **161**:10 (2012), 1863–1926. MR Zbl
- [Serre 2000] J.-P. Serre, *Local algebra*, Springer, Berlin, 2000. MR Zbl
- [Walker 2017] M. E. Walker, “On the vanishing of Hochster’s θ invariant”, *Ann. K-Theory* **2**:2 (2017), 131–174. MR Zbl

Communicated by Craig Huneke

Received 2016-12-31 Accepted 2017-08-09

- mkbrown5@math.wisc.edu Department of Mathematics,
University of Wisconsin-Madison, Madison, WI, United States
- clamille@syr.edu Mathematics Department, Syracuse University, Syracuse, NY,
United States
- peder.thompson@ttu.edu Department of Mathematics and Statistics,
Texas Tech University, Lubbock, TX, United States
- mark.walker@unl.edu Department of Mathematics, University of Nebraska-Lincoln,
Lincoln, NE, United States

Algebra & Number Theory

msp.org/ant

EDITORS

MANAGING EDITOR

Bjorn Poonen
Massachusetts Institute of Technology
Cambridge, USA

EDITORIAL BOARD CHAIR

David Eisenbud
University of California
Berkeley, USA

BOARD OF EDITORS

Richard E. Borcherds	University of California, Berkeley, USA	Martin Olsson	University of California, Berkeley, USA
J-L. Colliot-Thélène	CNRS, Université Paris-Sud, France	Raman Parimala	Emory University, USA
Brian D. Conrad	Stanford University, USA	Jonathan Pila	University of Oxford, UK
Samit Dasgupta	University of California, Santa Cruz, USA	Anand Pillay	University of Notre Dame, USA
Hélène Esnault	Freie Universität Berlin, Germany	Michael Rapoport	Universität Bonn, Germany
Gavril Farkas	Humboldt Universität zu Berlin, Germany	Victor Reiner	University of Minnesota, USA
Hubert Flenner	Ruhr-Universität, Germany	Peter Sarnak	Princeton University, USA
Sergey Fomin	University of Michigan, USA	Joseph H. Silverman	Brown University, USA
Edward Frenkel	University of California, Berkeley, USA	Michael Singer	North Carolina State University, USA
Andrew Granville	Université de Montréal, Canada	Christopher Skinner	Princeton University, USA
Joseph Gubeladze	San Francisco State University, USA	Vasudevan Srinivas	Tata Inst. of Fund. Research, India
Roger Heath-Brown	Oxford University, UK	J. Toby Stafford	University of Michigan, USA
Craig Huneke	University of Virginia, USA	Pham Huu Tiep	University of Arizona, USA
Kiran S. Kedlaya	Univ. of California, San Diego, USA	Ravi Vakil	Stanford University, USA
János Kollár	Princeton University, USA	Michel van den Bergh	Hasselt University, Belgium
Yuri Manin	Northwestern University, USA	Marie-France Vignéras	Université Paris VII, France
Philippe Michel	École Polytechnique Fédérale de Lausanne	Kei-Ichi Watanabe	Nihon University, Japan
Susan Montgomery	University of Southern California, USA	Shou-Wu Zhang	Princeton University, USA
Shigefumi Mori	RIMS, Kyoto University, Japan		

PRODUCTION

production@msp.org

Silvio Levy, Scientific Editor

See inside back cover or msp.org/ant for submission instructions.

The subscription price for 2017 is US \$325/year for the electronic version, and \$520/year (+\$55, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to MSP.

Algebra & Number Theory (ISSN 1944-7833 electronic, 1937-0652 printed) at Mathematical Sciences Publishers, 798 Evans Hall #3840, c/o University of California, Berkeley, CA 94720-3840 is published continuously online. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices.

ANT peer review and production are managed by EditFLOW[®] from MSP.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2017 Mathematical Sciences Publishers

Algebra & Number Theory

Volume 11 No. 9 2017

A nonarchimedean Ax–Lindemann theorem	1967
ANTOINE CHAMBERT-LOIR and FRANÇOIS LOESER	
A modular description of $\mathcal{X}_0(n)$	2001
KĘSTUTIS ČESNAVIČIUS	
Elementary equivalence versus isomorphism, II	2091
FLORIAN POP	
On the algebraic structure of iterated integrals of quasimodular forms	2113
NILS MATTHES	
On the density of zeros of linear combinations of Euler products for $\sigma > 1$	2131
MATTIA RIGHETTI	
Adams operations on matrix factorizations	2165
MICHAEL K. BROWN, CLAUDIA MILLER, PEDER THOMPSON and MARK E. WALKER	
Rationality does not specialize among terminal fourfolds	2193
ALEXANDER PERRY	
Topological noetherianity for cubic polynomials	2197
HARM DERKSEN, ROB H. EGGERMONT and ANDREW SNOWDEN	