

Algebra & Number Theory

Volume 13

2019

No. 1

**Extended eigenvarieties for
overconvergent cohomology**

Christian Johansson and James Newton



Extended eigenvarieties for overconvergent cohomology

Christian Johansson and James Newton

Recently, Andreatta, Iovita and Pilloni constructed spaces of overconvergent modular forms in characteristic p , together with a natural extension of the Coleman–Mazur eigencurve over a compactified (adic) weight space. Similar ideas have also been used by Liu, Wan and Xiao to study the boundary of the eigencurve. This all goes back to an idea of Coleman.

In this article, we construct natural extensions of eigenvarieties for arbitrary reductive groups G over a number field which are split at all places above p . If G is GL_2/\mathbb{Q} , then we obtain a new construction of the extended eigencurve of Andreatta–Iovita–Pilloni. If G is an inner form of GL_2 associated to a definite quaternion algebra, our work gives a new perspective on some of the results of Liu–Wan–Xiao.

We build our extended eigenvarieties following Hansen’s construction using overconvergent cohomology. One key ingredient is a definition of locally analytic distribution modules which permits coefficients of characteristic p (and mixed characteristic). When G is GL_n over a totally real or CM number field, we also construct a family of Galois representations over the reduced extended eigenvariety.

A correction was submitted on 27 October 2020 and posted online on 27 February 2021.

1. Introduction

1.1. The halo conjecture. The *eigencurve*, introduced by Coleman and Mazur [1998], is a rigid analytic curve $\mathcal{E}^{\mathrm{rig}}$ over \mathbb{Q}_p which parametrizes systems of Hecke eigenvalues of finite-slope overconvergent modular forms. It comes equipped with a morphism $\mathcal{E}^{\mathrm{rig}} \rightarrow \mathcal{W}_0^{\mathrm{rig}}$, called the weight map, whose target is known as *weight space*. $\mathcal{W}_0^{\mathrm{rig}}$ parametrizes continuous characters $\kappa : \mathbb{Z}_p^\times \rightarrow \overline{\mathbb{Q}}_p^\times$ and is a disjoint union of a finite number of open unit discs. There is also a morphism $\mathcal{E}^{\mathrm{rig}} \rightarrow \mathbb{G}_m^{\mathrm{rig}}$ which sends a system of Hecke eigenvalues to the U_p -eigenvalue; the p -adic valuation of the U_p -eigenvalue is known as the *slope*. The geometry of $\mathcal{E}^{\mathrm{rig}}$ encodes a wealth of information about congruences between finite-slope overconvergent modular forms, and it is therefore not surprising that its study remains a difficult topic. In particular, we know very little about the global geometry of $\mathcal{E}^{\mathrm{rig}}$ (for example, it is not known whether the number of irreducible components of $\mathcal{E}^{\mathrm{rig}}$ is finite or not).

Let $q = p$ if $p \neq 2$ and $q = 4$ if $p = 2$. The components of $\mathcal{W}_0^{\mathrm{rig}}$ are parametrized by the characters $(\mathbb{Z}/q\mathbb{Z})^\times \rightarrow (\mathbb{Z}/q\mathbb{Z})^\times$, and if we define $X := X_\kappa = \kappa(\exp(q)) - 1$, then X defines a parameter on each component. Very little is known about the global geometry of $\mathcal{E}^{\mathrm{rig}}$ over the centre $|X| \leq q^{-1}$ and it seems likely to be rather complicated. Near the boundary, however, the situation turns out to be rather

MSC2010: primary 11F33; secondary 11F80.

Keywords: p -adic automorphic forms, p -adic modular forms, eigenvarieties, Galois representations.

simple. Coleman and Mazur raised the question of whether the slope tends to zero as one moves along a component of \mathcal{E}^{rig} towards the boundary of $\mathcal{W}_0^{\text{rig}}$. Buzzard and Kilford [2005] investigated this question for $p = 2$ (and tame level 1) and proved a striking structure theorem: $\mathcal{E}^{\text{rig}}|_{\{|X| > 1/8\}}$ is a disjoint union of connected components $(E_i)_{i=0}^\infty$, and the weight map $E_i \rightarrow \{|X| > \frac{1}{8}\}$ is an isomorphism for every i . Moreover, the slope of a point on E_i with parameter X is $i \cdot v_2(X)$, where v_p is the p -adic valuation (normalized so that $v_p(p) = 1$). Out of this came a folklore conjecture; the following version is essentially [Liu et al. 2017, Conjecture 1.2]:

Conjecture. *For $r \in (0, 1)$ sufficiently close to 1, $\mathcal{E}^{\text{rig}}|_{\{|X| > r\}}$ is a disjoint union of connected components $(E_i)_{i=0}^\infty$ such that each E_i is finite over $\{|X| > r\}$. Moreover, there exist constants $\lambda_i \in \mathbb{R}_{\geq 0}$ for $i = 0, 1, \dots$, strictly increasing and tending to infinity, such that if x is a point on E_i with weight parameter X , then the slope of x is $\lambda_i v_p(X)$. The sequence $(\lambda_i)_{i=0}^\infty$ is a finite union of arithmetic progressions, after perhaps removing a finite number of terms.*

We will loosely refer to this as the “halo conjecture” (the “halo” in question is the (disjoint union of) annuli $\{|X| > r\}$). Let us assume $p \neq 2$ for simplicity. If κ is a point of $\mathcal{W}_0^{\text{rig}}$ then U_p acts compactly on the space of overconvergent modular forms M_κ^\dagger . The Fredholm determinants $\det(1 - T \cdot U_p | M_\kappa^\dagger) \in \overline{\mathbb{Q}_p}[[T]]$ interpolate to an entire series $F = \sum_{n=0}^\infty a_n T^n$ with coefficients in $\mathbb{Z}_p[[\mathbb{Z}_p^\times]] = \mathcal{O}(\mathcal{W}_0^{\text{rig}})^\circ$. Fix a character $\eta : \mathbb{Z}_p^\times \rightarrow \mathbb{F}_p^\times$ and consider the ideal $I_\eta = (p, [n] - \eta(n) \mid n = 1, \dots, p-1)$. The quotient ring $\mathbb{Z}_p[[\mathbb{Z}_p^\times]]/I_\eta$ is isomorphic to $\mathbb{F}_p[[X]]$ via the map sending $[\exp(p)]$ to $1 + X$. We may consider the reduction \bar{F}_η of F modulo I_η and the character

$$\bar{\kappa}_\eta : \mathbb{Z}_p^\times \rightarrow (\mathbb{Z}_p[[\mathbb{Z}_p^\times]]/I_\eta)^\times \cong \mathbb{F}_p[[X]]^\times.$$

In an unpublished note, Coleman conjectured that there should exist an $\mathbb{F}_p((X))$ -Banach space $\bar{M}_{\bar{\kappa}_\eta}^\dagger$ of “overconvergent modular forms of weight $\bar{\kappa}_\eta$ ” with a compact U_p -action such that $\det(1 - T \cdot U_p | \bar{M}_{\bar{\kappa}_\eta}^\dagger) = \bar{F}_\eta$, and promoted the idea that one should study the halo conjecture via integral models of the eigencurve near the boundary of weight space.

In [Andreatta et al. 2018], Andreatta, Iovita and Pilloni proved Coleman’s conjecture on the existence of $\bar{M}_{\bar{\kappa}_\eta}^\dagger$ and constructed an integral model \mathcal{E}' of \mathcal{E}^{rig} , which lives in the category of analytic adic spaces [Huber 1994]. $\mathcal{W}_0^{\text{rig}}$ has a natural formal scheme model $\text{Spf } \mathbb{Z}_p[[\mathbb{Z}_p^\times]]$, which may be viewed as an adic space \mathfrak{W}_0 over the affinoid ring $(\mathbb{Z}_p, \mathbb{Z}_p)$. Apart from the points corresponding to the adic incarnation of $\mathcal{W}_0^{\text{rig}}$, \mathfrak{W}_0 contains an additional $2(p-1)$ points in characteristic p , corresponding to the characters η and $\bar{\kappa}_\eta$. The latter points are analytic in Huber’s sense (the former are not) and one may consider the analytic locus $\mathcal{W}_0 = \mathcal{W}_0^{\text{rig}} \cup \{\bar{\kappa}_\eta \mid \eta\}$ of \mathfrak{W}_0 (viewing $\mathcal{W}_0^{\text{rig}}$ as an adic space). \mathcal{W}_0 may be viewed as a compactification of $\mathcal{W}_0^{\text{rig}}$, and Coleman’s idea may be interpreted as saying that one should study the behaviour of \mathcal{E}^{rig} near the boundary of \mathcal{W}_0 by extending the eigencurve to an adic space living over \mathcal{W}_0 , turning global behaviour into local behaviour “at infinity”. At a point $\bar{\kappa}_\eta$, one can no longer measure slopes using p . Instead, one has to use X . Noting that one can use X not only at $\bar{\kappa}_\eta$ but also “near” it, the halo conjecture says that the X -adic slope is constant as one approaches $\bar{\kappa}_\eta$. This supports

the idea that an extension of \mathcal{E}^{rig} exists, and that the E_i in the halo conjecture are X -adic Coleman families (i.e., subspaces which are finite over their images in weight space, and of constant slope). In this framework, the slopes of U_p should be the λ_i , with multiplicity the degree of X_i over $\{|X| > r\}$. The halo conjecture asserts, remarkably, that a Coleman family centred at a point over $\bar{\kappa}_\eta$ extends to some locus $\{|X| > r\}$ in the component corresponding to η , where r is independent of the family (and in particular its slope).

In [Liu et al. 2017], Liu, Wan and Xiao prove the halo conjecture for eigencurves for definite quaternion algebras. Here the construction of (overconvergent) automorphic forms is of a combinatorial nature. Those authors succeeded in proving the halo conjecture by calculations on some relatively explicit ad hoc integral models of spaces of overconvergent automorphic forms. They construct one space over the whole of \mathfrak{W}_0 , with a possibly noncompact U_p -action, and another model over $\{|X| > p^{-1}\}$ with a compact U_p -action. By the p -adic Jacquet–Langlands correspondence of [Chenevier 2005], this proves the halo conjecture for the components of the Coleman–Mazur eigencurve of (generically) Steinberg or supercuspidal type at some prime $q \neq p$.

1.2. Extended eigenvarieties for overconvergent cohomology. The main goal of this paper is to construct extensions of eigenvarieties for a very general class of connected reductive groups \mathbf{G} over \mathbb{Q} . In particular, we give a new construction of the extended eigencurve \mathcal{E}' appearing in [Andreatta et al. 2018]. Our construction also gives a conceptual framework for many of the results in [Liu et al. 2017] (and establishes a generalization of some of their results, which was described as an “optimistic expectation” in Remark 3.26(2) of that paper). See Theorem 6.3.4 for an interpretation of some of their results using the extended eigencurve.

Our construction of these eigenvarieties appears in Section 4.1. For the purposes of the introduction, we have the following vague statement:

Theorem A. *Let F be a number field and let \mathbf{H} be a connected reductive group over F which is split at all places above p . Set $\mathbf{G} = \text{Res}_{\mathbb{Q}}^F \mathbf{H}$. Then the eigenvarieties for \mathbf{G} constructed in [Hansen 2017] naturally extend to adic spaces $\mathcal{X}_{\mathbf{G}}$ over the extended weight space*

$$\mathcal{W} = \text{Spa}(\mathbb{Z}_p[[T'_0]], \mathbb{Z}_p[[T'_0]])^{\text{an}},$$

where T'_0 is a certain quotient of the \mathbb{Z}_p -points T_0 of a maximal torus in a suitable model of \mathbf{G} over \mathbb{Z}_p .

The assumption that \mathbf{H} is split at all places above p is made for convenience only; it makes it easy to define a “canonical” Iwahori subgroup. We believe that it should be relatively straightforward to generalize our constructions to general quasisplit \mathbf{G} over \mathbb{Q} (or to the setting of [Loeffler 2011]). The resulting theory would, however, be even more notationally cumbersome, so we have decided to stick to the simpler (but still very general) situation in this paper.

As a secondary goal, we show (Theorem 5.4.5) that when $\mathbf{G} = \text{Res}_{\mathbb{Q}}^F \text{GL}_{n/F}$, where F is a CM or totally real number field, the reduced eigenvariety that we construct carries a Galois determinant (in the language of [Chenevier 2014]) satisfying the expected compatibilities between Frobenii at unramified

places and the eigenvalues of Hecke operators. This shows that, in these cases, the new systems of Hecke eigenvalues that we construct in characteristic p carry arithmetic information.

Theorem B. *There exists an n -dimensional continuous determinant D of G_F with values in $\mathcal{O}^+(\mathcal{X}_G^{\text{red}})$ such that*

$$D(1 - X \text{Frob}_v) = P_v(X)$$

for unramified places v , where $P_v(X)$ is the usual Hecke polynomial (5.3.1).

Our proof of Theorem B is an adaptation of an argument due to the first author and David Hansen in the rigid setting, which will appear in a slightly refined form in [Hansen and Johansson \geq 2019]. It crucially uses Scholze’s results [2015] on Galois determinants attached to torsion classes, as well as filtrations on distribution modules constructed in [Hansen 2015].

To end our brief discussion of the results established in this paper, we explain one interpretation of the phrase “naturally extend” in Theorem A. Suppose for simplicity that $F = \mathbb{Q}$, $G(\mathbb{R})$ is compact modulo centre, and $G(\mathbb{Q}_p)$ may be identified with $\text{GL}_n(\mathbb{Q}_p)$. We let $T_0 \subseteq \text{GL}_n(\mathbb{Q}_p)$ denote the diagonal matrices with entries in \mathbb{Z}_p (in this case $T_0 = T'_0$). Modules of overconvergent automorphic forms for G (and some fixed tame level, which we suppress) were constructed in [Chenevier 2004] (see also [Loeffler 2011]). If we denote by U the Hecke operator corresponding to

$$\begin{pmatrix} 1 & & & \\ & p & & \\ & & \ddots & \\ & & & p^{n-1} \end{pmatrix}$$

then this acts compactly on the spaces of overconvergent automorphic forms, and so for each continuous character $\kappa : T_0 \rightarrow \overline{\mathbb{Q}}_p^\times$ there is a characteristic power series $F_\kappa \in \overline{\mathbb{Q}}_p[[T]]$ given by the determinant of $1 - TU$ on the space of overconvergent automorphic forms of weight κ . The following theorem is a consequence of our eigenvariety construction, together with Corollary 4.1.5.

Theorem C. *The characteristic power series F_κ glue together to $F_{\mathcal{W}} \in \mathcal{O}(\mathcal{W})\{\{T\}\}$, an entire function on affine 1-space over \mathcal{W} .*

Suppose $\bar{\kappa} : T_0 \rightarrow \mathbb{F}_q((X))^\times$ is a continuous character, with q a power of p . Then we give an interpretation of the specialization $F_{\mathcal{W}, \bar{\kappa}}$ of $F_{\mathcal{W}}$ at $\bar{\kappa}$ as the characteristic power series of U acting on an $\mathbb{F}_q((X))$ -Banach space of overconvergent automorphic forms of weight $\bar{\kappa}$.

The Fredholm hypersurface \mathcal{Z} cut out by $F_{\mathcal{W}}$ is locally quasifinite, flat and partially proper over \mathcal{W} and the eigenvariety \mathcal{X} comes equipped with a finite map to \mathcal{Z} .

1.3. Outline of the construction. The eigenvarieties that we extend are those constructed using overconvergent cohomology (sometimes also referred to as overconvergent modular symbols). Overconvergent cohomology was developed in [Stevens 1994; Ash and Stevens 2008], and the eigenvarieties were constructed in [Hansen 2017]. Let us recall their construction in the special case of the Coleman–Mazur eigencurve and $p \neq 2$. Let R be an affinoid \mathbb{Q}_p -algebra in the sense of rigid analytic geometry and let $\kappa : \mathbb{Z}_p^\times \rightarrow R^\times$ be a continuous homomorphism. It is well known that κ is locally analytic, and in

particular analytic on the cosets of $1 + p^s \mathbb{Z}_p$ for all sufficiently large s . For such s , we consider the Banach R -module $\mathcal{A}_\kappa[s]$ of functions $f : p\mathbb{Z}_p \rightarrow R$ which are analytic on the cosets of $p^s \mathbb{Z}_p$. The monoid

$$\Delta = \left\{ \gamma = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in M_2(\mathbb{Z}_p) \mid p \mid c, a \in \mathbb{Z}_p^\times, ad - bc \neq 0 \right\}$$

acts on $\mathcal{A}_\kappa[s]$ from the right by

$$(f, \gamma)(x) = \kappa(a + bx) f\left(\frac{c + dx}{a + bx}\right).$$

We consider the dual space $\mathcal{D}_\kappa[s] = \text{Hom}_{R, \text{cts}}(\mathcal{A}_\kappa[s], R)$, with the dual left action of Δ . A key point is that the matrix $t = \begin{pmatrix} 1 & 0 \\ 0 & p \end{pmatrix}$ acts compactly on $\mathcal{D}_\kappa[s]$; it factors through the compact injection $\mathcal{D}_\kappa[s] \hookrightarrow \mathcal{D}_\kappa[s+1]$. Fix an integer $N \geq 5$ (for simplicity) which is coprime to p , and consider the congruence subgroup $\Gamma = \Gamma_1(N) \cap \Gamma_0(p) \subseteq \Delta$. We may view $\mathcal{D}_\kappa[s]$ as a local system $\tilde{\mathcal{D}}_\kappa[s]$ on the complex modular curve $Y(\Gamma) = \Gamma \backslash \mathcal{H}$ and consider the singular cohomology group

$$H^1(Y(\Gamma), \tilde{\mathcal{D}}_\kappa[s]) = H^1(\Gamma, \mathcal{D}_\kappa[s]),$$

where the right-hand side is group cohomology (in general we would consider cohomology in all degrees, but it turns out that $H^i(\Gamma, \mathcal{D}_\kappa[s]) = 0$ if $i \neq 1$; see Section 6.1). It carries an action of the Hecke operator U_p . Considering these spaces for varying s and $R = \mathcal{O}(\mathcal{U})$, where $\mathcal{U} \subseteq \mathcal{W}_0^{\text{rig}}$ is an affinoid open subset, Hansen shows how to construct an eigenvariety from the Ash–Stevens cohomology groups by a clever adaptation of the eigenvariety construction of [Coleman 1997] (in the one-dimensional case) and [Buzzard 2007] (in the general case).¹ This eigenvariety turns out to equal the Coleman–Mazur eigencurve. To extend this construction to \mathcal{W}_0 , the key point is to define generalizations of the modules $\mathcal{D}_\kappa[s]$ for all open affinoid subsets $\mathcal{U} \subseteq \mathcal{W}_0$. Let $R = \mathcal{O}(\mathcal{U})$ and let $\kappa : \mathbb{Z}_p^\times \rightarrow R^\times$ be the induced character. The first thing to note is that κ is continuous but need not be locally analytic anymore, so one cannot directly copy the definition of $\mathcal{D}_\kappa[s]$. One could try to instead use the space \mathcal{A}_κ of all continuous functions $p\mathbb{Z}_p \rightarrow R$. This carries an action of Δ by the same formula, and we may consider its dual \mathcal{D}_κ . However, the action of t is no longer compact, so one has to do something different.

Let $f : p\mathbb{Z}_p \rightarrow R$ be a continuous function and let $f(x) = \sum_{n \geq 0} c_n \binom{x/p}{n}$ be its Mahler expansion. Recall that, when $\mathcal{U} \subseteq \mathcal{W}_0^{\text{rig}}$ (i.e., when R is a \mathbb{Q}_p -algebra), a theorem of Amice (see [Colmez 2010, Théorème I.4.7]) says that f is analytic on the cosets of $p^{s+1}\mathbb{Z}_p$ if and only if $|c_n|p^{n/p^s(p-1)} \rightarrow 0$ as $n \rightarrow \infty$. Here $|\cdot|$ is any \mathbb{Q}_p -Banach algebra norm such that $|p| = p^{-1}$. Dually, we may identify \mathcal{D}_κ with the ring of formal power series

$$\sum_{n \geq 0} d_n T^n,$$

where T^n is the distribution $f \mapsto c_n(f)$ and d_n is bounded as $n \rightarrow \infty$. The analytic distribution module $\mathcal{D}_\kappa[s]$ is defined by the weaker condition that $|d_n|p^{-n/p^s(p-1)}$ is bounded as $n \rightarrow \infty$. We may

¹Note that it is not clear how to topologize the R -modules $H^1(\Gamma, \mathcal{D}_\kappa[s])$, nor that they can be made into potentially ON-able Banach R -modules, so even in this special case we are combining Hansen’s eigenvariety construction with the Fredholm theory of Coleman and Buzzard, rather than using the Coleman–Mazur–Buzzard eigenvariety construction.

define norms $\| - \|_r$, for $r \in [1/p, 1)$, on \mathcal{D}_κ by $\| \sum_{n \geq 0} d_n T^n \|_r = \sup_n |d_n| r^n$. Let \mathcal{D}_κ^r denote the completion of \mathcal{D}_κ with respect to $\| - \|_r$; it may be explicitly described as the ring of power series $\sum_{n \geq 0} d_n T^n$, where $|d_n| r^n \rightarrow 0$. While the $\mathcal{D}_\kappa[s]$ are not among the \mathcal{D}_κ^r , one sees that $\varprojlim_{r \rightarrow 1} \mathcal{D}_\kappa^r = \varprojlim_{s \rightarrow \infty} \mathcal{D}_\kappa[s]$, so the norms allow one to recover the space of locally analytic distributions. As an aside, we remark it is possible to recover the $\mathcal{D}_\kappa[s]$ on the nose from the $\| - \|_r$, but we will not need them for the construction of eigenvarieties.

The upshot of considering the norms $\| - \|_r$ is that they may be constructed on \mathcal{D}_κ for any open affinoid $\mathcal{U} \subseteq \mathcal{W}_0$, by the formula given above. It is, however, not clear a priori that the norms interact well with the action of Δ . As a monoid Δ is generated by the Iwahori subgroup $I = \Delta \cap \mathrm{GL}_2(\mathbb{Z}_p)$ and the element t . The element t acts via multiplication by p on $p\mathbb{Z}_p$ and it is not too hard to see that it induces a norm-decreasing map $(\mathcal{D}_\kappa, \| - \|_r) \rightarrow (\mathcal{D}_\kappa, \| - \|_{r^{1/p}})$ and that the inclusions $\mathcal{D}_\kappa^s \subseteq \mathcal{D}_\kappa^r$ for $r < s$ are compact. Thus t induces a compact operator on \mathcal{D}_κ^r as desired. The action of I is more complicated to analyze, but it turns out that I acts by isometries for sufficiently large r (depending only on κ). To see this, it is useful to find a different description of $\| - \|_r$. This description, which we will outline below, is one of the key technical innovations of this paper. It is the analogue, in our setting of norms, of the observation in the rigid case that if κ is s -analytic then the I -action on \mathcal{A}_κ preserves $\mathcal{A}_\kappa[s]$.

Schneider and Teitelbaum [2003] generalized the norms defined above to the spaces $\mathcal{D}(G, L)$ of continuous distributions on a *uniform* pro- p group G [Dixon et al. 1999, Definition 4.1] valued in a finite extension L of \mathbb{Q}_p . To recall this construction briefly, a choice of a minimal set of topological generators of G induces an isomorphism $G \cong \mathbb{Z}_p^{\dim G}$ of p -adic manifolds and using multivariable Mahler expansions one may identify $\mathcal{D}(G, L)$ (as an L -Banach space) with $\mathcal{O}_L[[T_1, \dots, T_{\dim G}]] [1/p]$, and we put ($m = \dim G$)

$$\left\| \sum_{n_i \geq 0} d_{n_1, \dots, n_m} T_1^{n_1} \cdots T_m^{n_m} \right\|_r = \sup |d_{n_1, \dots, n_m}| r^{n_1 + \cdots + n_m}.$$

Schneider and Teitelbaum showed that these norms are submultiplicative and independent of the choice of minimal generating set. We generalize the construction of these norms to the module $\mathcal{D}(G, R)$ of distributions on a uniform group G valued in a certain class of normed \mathbb{Z}_p -algebras R that we call *Banach–Tate \mathbb{Z}_p -algebras*. These include the rings $R = \mathcal{O}(\mathcal{U})$ for $\mathcal{U} \subseteq \mathcal{W}_0$ open affinoid (for a suitable choice of norm) and generalize the constructions in the previous paragraph, which was the special case $G = p\mathbb{Z}_p$. Moreover, the action of $g \in G$ on $\mathcal{D}(G, R)$ via left or right translation is an isometry for $\| - \|_r$ (for any r).

Let $B_0 = \{\gamma \in I \mid c = 0\}$ be the upper triangular Borel and let $\bar{N}_1 = \left\{ \begin{pmatrix} 1 & 0 \\ x & 1 \end{pmatrix} \mid x \in p\mathbb{Z}_p \right\} \cong p\mathbb{Z}_p$; I has an Iwahori decomposition $I \cong \bar{N}_1 \times B_0$. Extend κ to a character of B_0 by setting $\kappa(\gamma) = \kappa(a)$. We have an I -equivariant injection $f \mapsto F$ of \mathcal{A}_κ into the space $\mathcal{C}(I, R)$ of continuous functions $F : G \rightarrow R$ given by $F(\bar{n}b) = f(\bar{n})\kappa(b)$, with $\bar{n} \in \bar{N}_1$ and $b \in B_0$. Here I acts on $\mathcal{C}(I, R)$ via left translation. The image is the set of functions F such that $F(gb) = \kappa(b)F(g)$ for all $g \in I$ and $b \in B_0$. Dually, we obtain an I -equivariant surjection $\mathcal{D}(I, R) \rightarrow \mathcal{D}_\kappa$. If we pretend, momentarily, that I is uniform, then we may

consider the quotient norm on \mathcal{D}_κ induced from $\| - \|_r$ on $\mathcal{D}(I, R)$ and one can show that for sufficiently large r , this quotient norm agrees with the previously defined $\| - \|_r$ on \mathcal{D}_κ . This shows that I acts by isometries on $(\mathcal{D}_\kappa, \| - \|_r)$ for sufficiently large r . In reality I is not uniform, but one can adapt the argument by working with a suitable open uniform normal subgroup of I .

This summarizes our construction of the modules \mathcal{D}_κ^r which we use to construct the eigenvariety. From the \mathcal{D}_κ^r , we construct variants $\mathcal{D}_\kappa^{<r}$ and function modules $\mathcal{A}_\kappa^r \subseteq \mathcal{A}_\kappa$ as well. When R is a Banach \mathbb{Q}_p -algebra, the $\mathcal{A}_\kappa[s]$ and $\mathcal{D}_\kappa[s]$ appearing in [Hansen 2017] are equal to \mathcal{A}_κ^r and $\mathcal{D}_\kappa^{<r}$, respectively, for $r = p^{-1/p^s(p-1)}$. It is easiest, however, to use the modules \mathcal{D}_κ^r to construct the eigenvariety since they are potentially orthonormalizable. Using the \mathcal{D}_κ^r , the construction of the eigenvariety follows [loc. cit.], and amounts largely to generalizing various well-known results from rigid geometry and nonarchimedean functional analysis. Our arguments also generalize from the case of $G = \mathrm{GL}_2/\mathbb{Q}$ to the general case considered in [loc. cit.] (as stated in Theorem A). In particular, our methods work for groups that do not have Shimura varieties (such as $G = \mathrm{Res}_{\mathbb{Q}}^F \mathrm{GL}_{n/F}$ for $n \geq 3$), which are intractable by the methods of [Andreatta et al. 2018] (see also [Andreatta et al. 2016]).

Remark 1.3.1. In independent work, Daniel Gulotta [2018] used a similar definition of distribution modules to extend Urban’s construction [2011] of equidimensional eigenvarieties for reductive groups possessing discrete series.

1.4. Questions and future work.

Generalizations of the halo conjecture. It is interesting to consider how the halo conjecture might generalize beyond the case of GL_2/\mathbb{Q} . For general G we raise the following questions:

Question 1.4.1. Does every irreducible component of the extended eigenvariety \mathcal{X}_G contain a point in the locus $p = 0$?

Question 1.4.2. Are there irreducible components of \mathcal{X}_G contained in the locus $p = 0$?

In the case of $G = \mathrm{GL}_2/\mathbb{Q}$ it is a consequence of the halo conjecture that every irreducible component contains a characteristic- p point. Similarly, when G is an inner form of GL_2/\mathbb{Q} associated to a definite quaternion algebra over \mathbb{Q} , it is a consequence of the results of [Liu et al. 2017] that every irreducible component contains a characteristic- p point (see Theorem 6.3.4). In general, we regard an affirmative answer to the first question as a very weak version of the halo conjecture.

If a component has a characteristic- p point, it becomes possible to study characteristic-0 points in the component (if they exist) by passing to the characteristic- p point, or to points approximating the characteristic- p point. In the case of GL_2/\mathbb{Q} (or its inner forms), components which have a characteristic- p point have a Zariski dense set of points corresponding to (twists of) classical modular forms of weight 2. One argument in this spirit appears in [Pottharst and Xiao 2014], which has been used by the authors in combination with the methods of [Liu et al. 2017] to establish new cases of the parity conjecture for the Bloch–Kato Selmer groups associated to Hilbert modular forms. We essentially do this by showing that there is a classical parallel weight-2 point on every irreducible component of an

eigenvariety for a definite quaternion algebra over a totally real field in which p splits completely. See [Johansson and Newton 2018] for more details.

Dimensions of irreducible components and functoriality. We note here that the theory of global irreducible components for the adic spaces we work with requires some explanation (see [Conrad 1999] for the rigid case). We have done this in a sequel to this paper [Johansson and Newton 2017], where we also generalize some of the results of [Hansen 2017]. In particular, we show that the lower bound for the dimension of irreducible components [loc. cit., Theorem 1.1.6] and (a variant of) the interpolation of Langlands functoriality [loc. cit., Theorem 5.1.6] generalize to our extended eigenvarieties.

One application of the interpolation of Langlands functoriality is that in the case of GL_2/\mathbb{Q} (or its inner forms) [Bergdall and Pollack 2016; Liu et al. 2017] show that the extended eigenvarieties contain the usual rigid eigenvarieties as a proper subspace. Applying functoriality (cyclic base change, for example) then shows that this is true for a larger class of groups. See [Johansson and Newton 2017] for more details.

Galois representations. In [Andreata et al. 2018] the natural question is raised as to whether the Galois representations attached to characteristic- p points of the extended eigencurve are trianguline (in an appropriate sense). One can similarly ask this question for the characteristic- p Galois representations constructed in this paper. Note that in our level of generality, it is still only conjectural that the characteristic-0 Galois representations carried by the eigenvariety are trianguline, but this is known, for example, in the case where G is a definite unitary group defined with respect to a CM field. It would also be interesting to construct a “patched extended eigenvariety” in this setting, extending the construction of [Breuil et al. 2017], and we hope to study this in the near future.

1.5. An outline of the paper. Let us describe the contents of the paper. Section 2 collects what we need about the eigenvariety machine and the notion of slope decompositions, and introduces some functional-analytic terminology that we will need throughout the paper. Since the key point of the paper is the construction of certain norms, we adopt terminology that puts emphasis on the norm, as opposed to merely the underlying topology. We give a definition of a slope decomposition (a concept introduced in [Ash and Stevens 2008]) that differs slightly from the definitions that appear in the literature. This is necessary since the definition given in [loc. cit.] neither localizes nor glues well, and so is not suitable for the construction of eigenvarieties. Our definition is a formalization of an informal definition that the first author learnt from conversations with David Hansen.

In Section 3, we carry out the construction of the norms on the \mathcal{D}_κ , following the outline above. We first discuss the generalization of the Schneider–Teitelbaum norms to distributions on a uniform group G valued in a certain class of normed \mathbb{Z}_p -algebras that we call Banach–Tate \mathbb{Z}_p -algebras. These include, for example, all Banach \mathbb{Q}_p -algebras in the usual sense, as well as Tate rings $R = \mathcal{O}(\mathcal{U})$ with \mathcal{U} an affinoid open subset of weight space (equipped with a suitable norm). We show that, in a precise sense, the completion of $\mathcal{D}(G, R)$ with respect to the family of norms $(\|\cdot\|_r)_{r \in [1/p, 1)}$ only depends on the underlying topology of R . Imposing some additional conditions on the norm (which is always possible in practice), we then construct the modules \mathcal{D}_κ^r , $\mathcal{D}_\kappa^{<r}$ and \mathcal{A}_κ^r as outlined above.

Section 4 then uses the modules \mathcal{D}_k^r to construct the eigenvariety, following the strategy in [Hansen 2017]. Since the \mathcal{D}_k^r are potentially orthonormalizable, the construction simplifies somewhat. We end the section by generalizing the “Tor-spectral sequence” [loc. cit., Theorem 3.3.1] to our setting, which is a key tool for analyzing the geometry of eigenvarieties, and use it give a description of the “points” of the eigenvariety valued in a local field. We use this description in Section 5 when we construct Galois determinants.

In Section 6 we discuss the relationship of our work with that of [Andreatta et al. 2018; Liu et al. 2017]. We show that when $G = \mathrm{GL}_2/\mathbb{Q}$, our construction, over the normalization of the weight space \mathcal{W}_0 discussed above, produces the same eigencurve as in [Andreatta et al. 2018] (this normalization is only different from \mathcal{W}_0 if $p = 2$). When G is the algebraic group over \mathbb{Q} associated with the units of a definite quaternion algebra over \mathbb{Q} , we show that our framework gives a conceptual proof of [Liu et al. 2017, Theorem 3.16], which is a key ingredient in their proof of the halo conjecture. In essence, the numerical estimate of [loc. cit., Theorem 3.16] falls out directly from our proof of compactness of the U_p -operator. Thus, it is possible to view our proof of compactness of suitable “ U_p -like” operators (known as controlling operators) as a generalization of [loc. cit., Theorem 3.16], as asked for in [loc. cit., Remark 3.26(2)]. Since this numerical estimate doesn’t appear strong enough to establish the halo conjecture in more general situations, we have restricted ourselves to proving the statement in the setting of [loc. cit.] as an illustration of our method.

Finally, the Appendix proves various results that we need on the class of Tate rings whose associated affinoid adic spaces appear as the local pieces of our eigenvarieties; some of these results might be of independent interest.

2. Preliminaries

The goal of this section is to set up some functional analytic terminology and theory. Specifically, we require the results of [Buzzard 2007, §2–3] on Fredholm determinants, Riesz theory and the construction of spectral varieties in a level of generality that is intermediate between the settings of [Buzzard 2007; Coleman 1997] (see also [Andreatta et al. 2018, Appendice B]). For example, we need to work over coefficient rings arising from affinoid opens in the adic space \mathcal{W}_0 discussed in our Introduction. These rings are complete topological rings which are *Tate* in the language of Huber [1993, §1]. The topology on these rings is induced by a norm, and to discuss the spectral theory of compact operators it is convenient to fix such a norm. This gives rise to a class of normed rings which we call Banach–Tate rings (see Definition 2.1.2).

The proofs in [Buzzard 2007] go through with little to no change when working over Banach–Tate rings, so we will be rather brief. All norms etc. will be nonarchimedean so we will ignore this adjective. All rings will be commutative unless otherwise specified.

2.1. Fredholm determinants over Banach–Tate rings.

Definition 2.1.1. Let R be a ring. A function $|\cdot| : R \rightarrow \mathbb{R}_{\geq 0}$ is called a *seminorm* if (for all $r, s \in R$)

- (1) $|0| = 0$ and $|1| = 1$;

$$(2) |r + s| \leq \max(|r|, |s|);$$

$$(3) |rs| \leq |r||s|.$$

If in addition $|r| = 0$ only if $r = 0$, then we say that $|\cdot|$ is a *norm*. A ring R together with a (semi-)norm will be called a (semi-)normed ring. A normed ring R is called a Banach ring if the metric induced by the norm is complete.

If $f : R \rightarrow S$ is a morphism of normed rings, we say that f is *bounded* if there is a constant $C > 0$ such that $|f(r)| \leq C|r|$ for all $r \in R$.

We say that two norms $|\cdot|, |\cdot|'$ on a ring R are *equivalent* if they induce the same topology. We say that they are *bounded-equivalent* if there are constants $C_1, C_2 > 0$ such that $C_1|r| \leq |r|' \leq C_2|r|$ for all $r \in R$ (note that this is stronger than equivalence, see Lemma 2.1.6). Let R be a normed ring. We say that $r \in R$ is *multiplicative* if $|rs| = |r||s|$ for all $s \in R$.

Definition 2.1.2. Let R be a normed ring. We say that R is *Tate* if R contains a multiplicative² unit ϖ such that $|\varpi| < 1$. We call such a ϖ a *multiplicative pseudouniformizer*. If R is also complete, we say that R is a *Banach–Tate ring*. If R is a Tate normed ring and ϖ is a multiplicative pseudouniformizer, then we define the corresponding valuation v_ϖ on R by $v_\varpi(r) = -\log_a |r|$, where $a = |\varpi^{-1}|$.

We remark that it is easy to see that a unit ϖ in a normed ring R is multiplicative if and only if $|\varpi^{-1}| = |\varpi|^{-1}$. A multiplicative pseudouniformizer ϖ is a uniform unit in the sense of [Kedlaya and Liu 2015, Remark 2.3.9(b)].

Remark 2.1.3. Let R be a Tate normed ring, with ϖ a multiplicative pseudouniformizer:

(1) The underlying topological ring is a Tate ring in the language of Huber; the unit ball R_0 is a ring of definition and ϖ is a topologically nilpotent unit. Conversely, assume R is a Tate ring and $\varpi \in R$ is a topologically nilpotent unit, contained in some ring of definition R_0 . If $a \in \mathbb{R}_{>1}$, then we may define a norm on R by $|r| = \inf\{a^{-n} \mid r \in \varpi^n R_0, n \in \mathbb{Z}\}$. Equipped with this norm, R is a Tate normed ring with unit ball R_0 and ϖ is a multiplicative pseudouniformizer.

(2) A Banach–Tate ring A is the same thing as a Banach algebra A satisfying $|A^m| \neq 1$ in the language of [Coleman 1997, §1] (and what we call a Banach ring is what Coleman calls a Banach algebra). Here A^m denotes the set of multiplicative units of A . Additionally, when R is a Banach–Tate ring and R^+ is a ring of integral elements, a choice of a multiplicative pseudouniformizer ϖ may be used to identify the Gelfand spectrum $\mathcal{M}(R)$ of bounded multiplicative seminorms on R [Berkovich 1990, §1.2] with the maximal compact Hausdorff quotient of the adic spectrum $\mathrm{Spa}(R, R^+)$ [Huber 1993]; see [Kedlaya and Liu 2015, Definition 2.4.6]. Concretely, ϖ gives us a natural way of viewing a rank-1 point in $\mathrm{Spa}(R, R^+)$ as a bounded multiplicative seminorm.

Definition 2.1.4. Let R be a normed ring. A *normed R -module* is an R -module M equipped with a function $\|\cdot\| : M \rightarrow \mathbb{R}_{\geq 0}$ such that (for all $m, n \in M$ and $r \in R$)

²That is, a unit which is multiplicative in the sense we just defined, as well as being a unit for multiplication!

- (1) $\|m\| = 0$ if and only if $m = 0$;
- (2) $\|m + n\| \leq \max(\|m\|, \|n\|)$;
- (3) $\|rm\| \leq |r| \|m\|$.

We remark that if $r \in R$ is a multiplicative unit, then one sees easily that $\|rm\| = |r| \|m\|$ for all $m \in M$. If R is a Banach ring and M is complete, we say that M is a *Banach R -module*.

Let R be a Tate normed ring. If M and N are normed R -modules, then a homomorphism $\phi : M \rightarrow N$ is a continuous R -linear map. In this case, continuity of an R -linear map ϕ is equivalent to boundedness; i.e., there exists $C \in \mathbb{R}_{>0}$ such that $\|\phi(m)\| \leq C \|m\|$ for all $m \in M$. In this case we set $|\phi| = \sup_{m \neq 0} |\phi(m)| \|m\|^{-1}$ as usual; $\text{Hom}_{R, \text{cts}}(M, N)$ becomes a normed R -module with respect to this norm. The open mapping theorem holds in this context; see, e.g., [Huber 1994, Lemma 2.4(i)].

Let R be a Noetherian Banach–Tate ring. The results of [Bosch et al. 1984, §3.7.2] hold in the context of Banach–Tate rings with the same proofs (thanks to the open mapping theorem), so R being Noetherian is equivalent to all ideals being closed. Moreover, the results of [loc. cit., §3.7.3] hold for *Noetherian* Banach–Tate rings with the same proofs. In particular, any finitely generated R -module carries a canonical complete topology, and any abstract R -linear map between two finitely generated R -modules is continuous and strict with respect to the canonical topology.

Definition 2.1.5. Let R be a Banach–Tate ring and let I be a set. We define $c_R(I)$ to be the set of sequences $(r_i)_{i \in I}$ in R tending to 0 (with respect to the filter of subsets of I with finite complement). It is a Banach R -module when equipped with the norm $\|(r_i)\| = \sup_{i \in I} |r_i|$.

We say that a Banach R -module M is (*potentially*) *orthonormalizable* (or (*potentially*) *ON-able* for short) if there exists a set I such that M is R -linearly isometric (resp. merely R -linearly homeomorphic) to $c_R(I)$. A set in M corresponding to the set $\{e_i = (\delta_{ij})_j \mid i \in I\} \subseteq c_R(I)$ under such a map is called an (*potential*) *ON-basis*.

Finally, we say that a Banach R -module M has *property (Pr)* if it is a direct summand of a potentially ON-able Banach R -module.

If $M \rightarrow N$ is a continuous morphism of ON-able Banach R -modules, then we may define its matrix for a fixed ON-basis on M and one on N as on [Buzzard 2007, p. 65], and the properties stated there hold in this situation as well. A morphism $\phi : M \rightarrow N$ between general Banach R -modules is said to be of *finite rank* if the image of ϕ is contained in a finitely generated submodule of N . More generally, ϕ is said to be *compact* (or *completely continuous*) if it is a limit of finite-rank operators in $\text{Hom}_{R, \text{cts}}(M, N)$. If R is Noetherian, [loc. cit., Lemma 2.3, Proposition 2.4] go through with the same proofs (using a multiplicative pseudouniformizer ϖ for what Buzzard calls ρ in the proof of Lemma 2.3) and we see that if $\phi : M \rightarrow N$ is a continuous R -linear map between ON-able Banach R -modules with matrix (a_{ij}) with respect to some bases $(e_i)_{i \in I}$ of M and $(f_j)_{j \in J}$ of N , then ϕ is compact if and only if $\lim_{j \rightarrow \infty} \sup_{i \in I} |a_{ij}| = 0$. When $M = N$ and $(e_i)_{i \in I} = (f_j)_{j \in J}$ this allows us to define the *characteristic power series*, or *Fredholm determinant*, $\det(1 - T\phi)$ of a compact ϕ using the recipe on [loc. cit., p. 67]

and one sees that $\det(1 - Tu) \in R\{\{T\}\}$, where

$$R\{\{T\}\} = \left\{ \sum_n a_n T^n \in R[[T]] \mid |a_m| M^m \rightarrow 0 \text{ for all } M \in \mathbb{R}_{\geq 0} \right\}$$

is the ring of entire power series in R .

Moving on, we remark that [loc. cit., Lemma 2.5, Corollary 2.6] are true in our setting with the same proofs. In particular, the notion of the Fredholm determinant extends to compact operators on potentially ON-able M , and may be computed using a potential ON-basis. It will be useful (at least psychologically) for us to know that these notions remain unchanged if we replace the norm on R by an equivalent one. First, we remark that changing the norm on R to an equivalent one doesn't change the topology on $c_R(I)$ (for I arbitrary). This can be seen directly, but it is also a consequence of the following lemma, which we will need later.

Lemma 2.1.6. *Let R be a complete Tate ring, and let $\varpi, \pi \in R$ be topologically nilpotent units. Assume that we have two equivalent norms $|\cdot|_\varpi$ and $|\cdot|_\pi$ on R (inducing the intrinsic topology) such that ϖ is multiplicative for $|\cdot|_\varpi$ and π is multiplicative for $|\cdot|_\pi$. Then we may find constants $C_1, C_2, s_1, s_2 > 0$ such that*

$$C_1 |a|_\pi^{s_1} \leq |a|_\varpi \leq C_2 |a|_\pi^{s_2}$$

for all $a \in R$.

Proof. We thank a referee for suggesting a more efficient argument for this proof. First, note that it suffices to find constants such that the inequalities hold for all nonzero $a \in R$, since it trivially holds for $a = 0$ and all choices of constants. We will first prove the second inequality. To start, pick $C' > 0$ such that $|a|_\pi \leq 1$ implies $|a|_\varpi \leq C'$ for all $a \in R$ (possible since the norms are equivalent). Since ϖ is topologically nilpotent we may find $m \in \mathbb{Z}_{\geq 1}$ such that $|\varpi^m|_\pi < 1$. It follows that $|\varpi^{-m}|_\pi \geq |\varpi^m|_\pi^{-1} > 1$.

For any nonzero $a \in R$, set

$$n = \left\lceil \frac{\log |a|_\pi}{\log |\varpi^m|_\pi^{-1}} \right\rceil.$$

We have $|\varpi^m|_\pi^n |a|_\pi \leq 1$. If $n \geq 0$, we deduce that $|\varpi^{mn} a|_\pi \leq |\varpi^m|_\pi^n |a|_\pi \leq 1$. If $n < 0$ we similarly deduce that

$$|\varpi^{mn} a|_\pi \leq |\varpi^{-m}|_\pi^{-n} |a|_\pi \leq |\varpi^m|_\pi^n |a|_\pi \leq 1.$$

Therefore we have $|\varpi^{mn} a|_\varpi \leq C'$. By multiplicativity of ϖ for $|\cdot|_\varpi$ we get $|a|_\varpi \leq C' |\varpi|_\varpi^{-mn}$. Setting $q = |\varpi|_\varpi^{-m} > 1$, we then have

$$|a|_\varpi \leq C' q^n \leq C' q^{(\log |a|_\pi / \log |\varpi^m|_\pi^{-1}) + 1} = C' q |a|_\pi^{s_2},$$

where we have put $s_2 = (\log_q |\varpi^m|_\pi^{-1})^{-1}$; note that $s_2 > 0$. Set $C_2 = C' q$; we get

$$|a|_\varpi \leq C_2 |a|_\pi^{s_2},$$

with $C_2, s_2 > 0$ as desired.

To get the first inequality, note that by symmetry we may find $D, t > 0$ such that $|a|_\pi \leq D|a|_\varpi^t$. Rearranging we obtain

$$C_1|a|_\pi^{s_1} \leq |a|_\varpi \leq C_2|a|_\pi^{s_2}$$

as desired, where $C_1 = D^{-1/t}$ and $s_1 = 1/t$. \square

Remark 2.1.7. If R has two equivalent norms $|\cdot|$ and $|\cdot|'$ with a common multiplicative pseudouniformizer ϖ such that $|\varpi| = |\varpi|'$, then the proof shows that $|\cdot|$ and $|\cdot|'$ are *bounded-equivalent*. This is also easy to see directly, and will be used freely throughout the paper.

Suppose then that $(M, |\cdot|)$ is a Banach $(R, |\cdot|)$ -module, and $(M, |\cdot|')$ is a Banach $(R, |\cdot|')$ -module, where $|\cdot|$ and $|\cdot|'$ are equivalent on both R and M . A Banach $(R, |\cdot|)$ -module isomorphism $(M, |\cdot|) \cong (c_R(I), |\cdot|)$ is then the same thing as a Banach $(R, |\cdot|')$ -module isomorphism $(M, |\cdot|) \cong (c_R(I), |\cdot|')$, since $(c_R(I), |\cdot|) \cong (c_R(I), |\cdot|')$ as *topological* R -modules via the identity map. Thus $(M, |\cdot|)$ is potentially ON-able if and only if $(M, |\cdot|')$ is potentially ON-able, and $(e_i)_{i \in I}$ is a potential ON-basis for $(M, |\cdot|)$ if and only if it is a potential ON-basis for $(M, |\cdot|')$. It follows, at least when R is Noetherian (which is all we need), that an operator ϕ is compact on a potentially ON-able $(M, |\cdot|)$ if and only if it is compact on a potentially ON-able $(M, |\cdot|')$, and the Fredholm determinant is the same.

We remark that the results [Buzzard 2007, Lemma 2.7–Corollary 2.10] hold over Noetherian Banach–Tate rings R , again with the same proofs. We can extend the notion of Fredholm determinants of compact operators on Banach R -modules with property (Pr) as on [loc. cit., pp. 72–73], and the results there hold over Noetherian Banach–Tate rings. One also sees that having property (Pr) is stable when changing the norms on (R, M) to equivalent ones, as is compactness of operators and the Fredholm determinants for compact operators are unchanged. We summarize the results of this section with the following proposition:

Proposition 2.1.8. *Let R be a Noetherian Banach–Tate ring. If M is a Banach R -module with property (Pr) and $\phi : M \rightarrow M$ is compact then there is a well-defined Fredholm determinant*

$$\det(1 - T\phi|M) \in R\{\{T\}\}.$$

If we change the norms on (R, M) to equivalent ones, then M still has property (Pr), ϕ is still compact, and the Fredholm determinant is unchanged.

2.2. Riesz theory, slope factorizations and slope decompositions. We continue to let R denote a Banach–Tate ring. If $Q \in R[T]$, we write $Q^*(T) := T^{\deg Q} Q(1/T)$. We recall the following definitions:

Definition 2.2.1. A *Fredholm series* is a formal power series $F = 1 + \sum_{n \geq 1} a_n T^n \in R\{\{T\}\}$. A polynomial $Q \in R[T]$ is called *multiplicative* if the leading coefficient of Q is a unit (in other words, if $Q^*(0) \in R^\times$).

Two entire series $P, Q \in R\{\{T\}\}$ are said to be *relatively prime* if the ideal (P, Q) is equal to $R\{\{T\}\}$.

The proof of [Buzzard 2007, Theorem 3.3] goes through without changes; we state it for completeness (see also [Andreatta et al. 2018, Théorème B.2]). Implicit in this is that [Buzzard 2007, Lemma 3.1] holds with the same proof; we will make use of this later.

Theorem 2.2.2. *Assume that R is Noetherian. Let M be a Banach R -module with property (Pr) and let $u : M \rightarrow M$ be a compact operator with $F = \det(1 - Tu)$. Assume that we have a factorization $F = QS$, where S is a Fredholm series, $Q \in R[T]$ is a multiplicative polynomial, and Q and S are relatively prime in $R\{\{T\}\}$. Then $\text{Ker } Q^*(u) \subseteq M$ is finitely generated and projective and has a unique u -stable closed complement N such that $Q^*(u)$ is invertible on N . The idempotent projectors $M \rightarrow \text{Ker } Q^*(u)$ and $M \rightarrow N$ lie in the closure of $R[u] \subseteq \text{End}_{R,\text{cts}}(M)$. The rank of $\text{Ker } Q^*(u)$ is $\deg Q$, and $\det(1 - Tu | \text{Ker } Q^*(u)) = Q$. Moreover, u is invertible on $\text{Ker } Q^*(u)$, and $\det(1 - Tu | N) = S$.*

Proof. Apart from the last sentence, this is (a minor reformulation of) [Buzzard 2007, Theorem 3.3]. To see that u is invertible on $\text{Ker } Q^*(u)$, note that

$$\det(u | \text{Ker } Q^*(u)) = Q^*(0) \in R^\times.$$

To see that $\det(1 - Tu | N) = S$, write $S' = \det(1 - Tu | N)$ and note that

$$F = \det(1 - Tu | \text{Ker } Q^*(u)) \det(1 - Tu | N) = QS'.$$

Hence $Q(S - S') = 0$, and Q is not a zero divisor since $Q(0) = 1$, so $S = S'$. \square

The following lemma may be extracted from the proof of [loc. cit., Lemma 5.6]; we give the short proof for completeness.

Lemma 2.2.3. *Assume that R is Noetherian. Let M and M' be two Banach R -modules with property (Pr) and assume that we have a continuous R -linear map $v : M \rightarrow M'$ and a compact R -linear map $i : M' \rightarrow M$. Set $u = iv$ and $u' = vi$. Then u and u' are both compact and $\det(1 - Tu) = \det(1 - Tu')$; call this entire power series F . If $F = QS$ is a factorization as in Theorem 2.2.2, then i restricts to an isomorphism between $\text{Ker } Q^*(u')$ and $\text{Ker } Q^*(u)$.*

Proof. Compactness of u and u' and the equality of their Fredholm determinants follows from [loc. cit., Proposition 2.7]. Now assume we have a factorization $F = QS$. If $x' \in \text{Ker } Q^*(u')$, then $Q^*(u)(i(x')) = i(Q^*(u')(x')) = 0$ so $i(\text{Ker } Q^*(u')) \subseteq \text{Ker } Q^*(u)$. Furthermore, if $i(x') = 0$ then $u'(x') = 0$, so $Q^*(u')(x') = Q^*(0).x' = 0$ and hence $x' = 0$, so i is injective on $\text{Ker } Q^*(u')$. For surjectivity onto $\text{Ker } Q^*(u)$, let $x \in \text{Ker } Q^*(u)$ and choose $y \in \text{Ker } Q^*(u)$ with $u(y) = x$ (possible by Theorem 2.2.2). Then one checks, similarly to the computation above, that $v(y) \in \text{Ker } Q^*(u')$, and hence $i(v(y)) = u(y) = x$, which gives us surjectivity and finishes the proof. \square

Next, we let K be a field, complete with respect to a nontrivial nonarchimedean absolute value. We briefly define the Newton polygon of a power series $F \in K[[T]]$, following [Ash and Stevens 2008, §4.2] (in this special case). A subset $\mathcal{N} \subseteq \mathbb{R}^2$ is said to be *sup-convex* if it is convex and, if a point (a, b) is in \mathcal{N} , then \mathcal{N} contains the whole half-line $\{(a, b + t) \mid t \geq 0\}$ above it. Given an arbitrary subset $S \subseteq \mathbb{R}^2$, there is a unique smallest sup-convex set containing S , which we will denote by $\mathcal{H}_+(S)$. If $I \subseteq \mathbb{Z}_{\geq 0}$ and $\omega : I \rightarrow \mathbb{R}$ is a function, then any set of the form $\mathcal{H}_+(\{(n, \omega(n)) \mid n \in I\})$ is called a *Newton polygon*. We refer to [loc. cit., §4.2] for the notions of vertices, edges and slopes of a Newton polygon.

Definition 2.2.4. Let $F = \sum_{n \geq 0} a_n T^n \in K[[T]]$. Fix a pseudouniformizer $\varpi \in K$ and consider the corresponding valuation v_ϖ . The *Newton polygon of F* is the Newton polygon $\mathcal{H}_+(S(F))$, where

$$\mathcal{S}(F) = \{(n, v_\varpi(a_n)) \mid n \in I_F\} \subseteq \mathbb{R}^2,$$

with $I_F = \{n \in \mathbb{Z}_{\geq 0} \mid a_n \neq 0\}$.

Let $h \in \mathbb{R}$. We say that a power series $F \in K[[T]]$ has slope $\leq h$ (or $> h$) if all slopes of its Newton polygon are $\leq h$ (or $> h$). Now consider a Banach–Tate ring R with a multiplicative pseudouniformizer ϖ . We say that $F \in R[[T]]$ has slope $\leq h$ (or $> h$) if, for any x in the Gelfand spectrum $\mathcal{M}(R)$ with residue field K_x , the specialization $F_x \in K_x[[T]]$ has slope $\leq h$ (or $> h$).

Definition 2.2.5. Let R be a Banach–Tate ring with a fixed multiplicative pseudouniformizer ϖ . Let $F \in R\{\{T\}\}$ be a Fredholm series and let $h \in \mathbb{R}$. A *slope $\leq h$ -factorization* of F is a factorization $F = QS$ in $R\{\{T\}\}$, where Q is a multiplicative polynomial of slope $\leq h$ and S is a Fredholm series of slope $> h$.

Remark 2.2.6. If R is a complete Tate ring with a fixed topologically nilpotent unit ϖ , then the notions of slope factorizations and slope $\leq h$ or $> h$ are independent of the choice of a norm on R with ϖ multiplicative. Moreover, one can define all these notions directly without choosing a norm on R .

Recall that an element $a \in R$ is called *quasinilpotent* if its spectral seminorm³ $|a|_{\text{sp}}$ is 0. This is equivalent to $|a|_x = 0$ for all $x \in \mathcal{M}(R)$ by [Berkovich 1990, Corollary 1.3.2]. The set of quasinilpotent elements forms an ideal of R , which is the kernel of the Gelfand transform $R \rightarrow \prod_{x \in \mathcal{M}(R)} K_x$. We note that it is easy to see that a quasinilpotent element is topologically nilpotent. For the kinds of rings R which appear in practice in this paper, the quasinilpotent elements are just the nilpotent elements (this follows from Theorem A.7), and the proof of the following lemma is simpler. However, we will avoid imposing additional technical assumptions at this stage.

Lemma 2.2.7. *Let R be a Banach–Tate ring with a fixed multiplicative pseudouniformizer ϖ and let $h \in \mathbb{Q}_{\geq 0}$. Let S be a Fredholm series of slope $> h$ and Q a multiplicative polynomial of slope $\leq h$. Then Q and S are relatively prime.*

Proof. We will use Coleman’s resultant Res , for which we refer to [Coleman 1997, §A3] (the reader may also benefit from the discussion on [Buzzard 2007, p. 74]). By [Coleman 1997, Lemma A3.7] it suffices to prove that $\text{Res}(Q, S)$ is a unit in $R\{\{T\}\}$. Pick $x \in \mathcal{M}(R)$ and specialize to K_x . Then $\text{Res}(Q, S)_x = \text{Res}(Q_x, S_x)$ and since Q_x has slope $\leq h$ and S_x has slope $> h$ we see that $\text{Res}(Q, S)_x \in K_x\{\{T\}\}^\times = K_x^\times$. By [Berkovich 1990, Corollary 1.2.4] we see that $\text{Res}(Q, S) = a_0 + T.F(T)$, where $a_0 \in R^\times$ and $F(T) \in R\{\{T\}\}$ has quasinilpotent coefficients. Multiplying by a_0^{-1} we see that it suffices to prove that if $F \in R\{\{T\}\}$ has quasinilpotent coefficients, then $1 - T.F(T) \in R\{\{T\}\}^\times$.

To prove this, we use an argument suggested to us by a referee, which is more efficient than our original argument. First note that the formal inverse of $1 - T.F(T)$ is $G(T) = \sum_{n \geq 0} T^n.F(T)^n$, so we need to show that this is entire. Setting $H(T) = T.F(T)$, it suffices to show that if $H(T)$ is any entire power

³The definition of the spectral seminorm is recalled in the Appendix.

series with $H(0) = 0$ and with quasinilpotent coefficients, then $\sum_{n \geq 0} H(T)^n$ converges and is entire. In fact, it suffices to prove that $\sum_{n \geq 0} H(T)^n \in R\langle T \rangle$, since if we have proved this we can apply this to $H(\varpi^{-N}T)$ for all N (which still has quasinilpotent coefficients) to deduce that $\sum_{n \geq 0} H(T)^n$ is entire.

So, to show this, it suffices to show that $H(T)$ is topologically nilpotent in $R\langle T \rangle$, where we equip $R\langle T \rangle$ with the Gauss norm coming from the norm on R . Since the coefficients of H tend to 0, we may write $H(T) = \sum_{n=1}^N h_n T^n + \sum_{n > N} h_n T^n$, with $|h_n| < 1$ for $n > N$. Then the tail $\sum_{n > N} h_n T^n$ is topologically nilpotent (it has Gauss norm < 1) and the terms $h_n T^n$ are topologically nilpotent for all n since h_n is quasinilpotent (and hence topologically nilpotent). So $H(T)$ is a finite sum of topologically nilpotent elements, and hence topologically nilpotent. \square

Continue to let R be a Banach–Tate ring with a fixed multiplicative pseudouniformizer ϖ . The following is a minor variation of [Ash and Stevens 2008, Definition 4.6.1].

Definition 2.2.8. Let M be an (abstract) R -module, let $u : M \rightarrow M$ be an R -linear map and let $h \in \mathbb{Q}$. An element $m \in M$ is said to have slope $\leq h$ with respect to u if there is a multiplicative polynomial $Q \in R[T]$ such that

- (1) $Q^*(u).m = 0$;
- (2) the slope of Q is $\leq h$.

We let $M_{\leq h} \subseteq M$ denote the subset of elements of slope $\leq h$.

Lemma 2.2.9 [Ash and Stevens 2008, Proposition 4.6.2]. $M_{\leq h}$ is an R -submodule of M , which is stable under u .

Proof. It is clear from the definition that $M_{\leq h}$ is closed under multiplication, and stable under u . It therefore suffices to prove that it is closed under addition, for which it suffices to prove that if Q_1 and Q_2 are two multiplicative polynomials of slope $\leq h$, then so is $Q_1 Q_2$. To see this it suffices to specialize to the case when R is a field and the norm is an absolute value. The assertion is then well known (for example, the argument in the proof of [loc. cit., Proposition 4.6.2] carries over without change). \square

Definition 2.2.10 [Ash and Stevens 2008, Definition 4.6.3]. Let M be an R -module with an R -linear map $u : M \rightarrow M$ and let $h \in \mathbb{Q}$. A slope $\leq h$ -decomposition of M is an $R[u]$ -module decomposition $M = M_h \oplus M^h$ such that

- (1) M_h is a finitely generated R -submodule of $M_{\leq h}$;
- (2) for every multiplicative polynomial $Q \in R[T]$ of slope $\leq h$, the map $Q^*(u) : M^h \rightarrow M^h$ is an isomorphism of R -modules.

Proposition 2.2.11. *We keep the above notation. If M has a slope $\leq h$ -decomposition $M_h \oplus M^h$, then it is unique, and $M_h = M_{\leq h}$ (in particular the latter is finitely generated over R). We will from now on write $M_{>h}$ for the unique complement. Moreover, slope decompositions satisfy the following functorial properties:*

- (1) Let $f : M \rightarrow N$ be a morphism of $R[u]$ -modules with slope $\leq h$ -decompositions. Then $f(M_{\leq h}) \subseteq N_{\leq h}$ and $f(M_{>h}) \subseteq N_{>h}$. Moreover, both $\text{Ker}(f)$ and $\text{Im}(f)$ have slope $\leq h$ -decompositions.
- (2) Let C^\bullet be a complex of $R[u]$ -modules and suppose that each C^i has a slope $\leq h$ -decomposition. Then every $H^i(C^\bullet)$ has a slope $\leq h$ -decomposition, explicitly given by $H^i(C^\bullet) = H^i(C_{\leq h}^\bullet) \oplus H^i(C_{>h}^\bullet)$.

Proof. The proof is identical to that of [Ash and Stevens 2008, Lemma 4.6.4]: one equates slope $\leq h$ -decompositions with \mathcal{S} -decompositions (as defined and studied in [loc. cit., §4.1]) for the set $\mathcal{S} \subseteq R[u]$ of all $Q^*(u)$, where Q is a multiplicative polynomial of slope $\leq h$. The properties then stated follow from general facts about \mathcal{S} -decompositions, recorded in [loc. cit., Proposition 4.1.2]. \square

Definition 2.2.12. Let R be a Banach–Tate ring with a fixed multiplicative pseudouniformizer ϖ and let M be a Banach R -module. Assume that M has a slope $\leq h$ -decomposition $M = M_{\leq h} \oplus M_{>h}$. If $f : R \rightarrow S$ is a bounded morphism of Banach–Tate rings such that $f(\varpi)$ is a multiplicative pseudouniformizer in S , we say that the slope $\leq h$ -decomposition is *functorial* for $R \rightarrow S$ if $M \widehat{\otimes}_R S = (M_{\leq h} \otimes_R S) \oplus (M_{>h} \widehat{\otimes}_R S)$ is a slope $\leq h$ -decomposition of $M \widehat{\otimes}_R S$ (using $f(\varpi)$ to define slopes for S). We say that the slope $\leq h$ -decomposition is *functorial* if it is functorial for all such bounded homomorphisms of Banach–Tate rings out of R .

Theorem 2.2.13. Let R be a Noetherian Banach–Tate ring with a fixed multiplicative pseudouniformizer ϖ , and let M be a Banach R -module with property (Pr). Let u be a compact R -linear operator on M , with Fredholm determinant $F(T) = \det(1 - Tu)$. If M has a slope $\leq h$ -decomposition which is functorial with respect to $R \rightarrow K_x$ for all $x \in \mathcal{M}(R)$, then F has a slope $\leq h$ -factorization. Conversely, if F has a slope $\leq h$ -factorization, then M has a functorial slope $\leq h$ -decomposition.

Proof. Assume that M has a slope $\leq h$ -decomposition $M = M_{\leq h} \oplus M_{>h}$ which is functorial with respect to $R \rightarrow K_x$ for all $x \in \mathcal{M}(R)$. Then both of these spaces satisfy property (Pr) and are u -stable, and hence we have

$$F = \det(1 - Tu \mid M_{\leq h}) \det(1 - Tu \mid M_{>h}).$$

We claim that this is a slope $\leq h$ -factorization. Put $Q = \det(1 - Tu \mid M_{\leq h})$, $S = \det(1 - Tu \mid M_{>h})$. Pick $x \in \mathcal{M}(R)$ with residue field K_x and specialize. We have $Q_x = \det(1 - Tu \mid M_{\leq h} \otimes_R K_x)$ and $S_x = \det(1 - Tu \mid M_{>h} \widehat{\otimes}_R K_x)$. By assumption $M \widehat{\otimes}_R K_x = (M_{\leq h} \otimes_R K_x) \oplus (M_{>h} \widehat{\otimes}_R K_x)$ is a slope $\leq h$ -decomposition, so Q_x has slopes $\leq h$ and S_x has slopes $> h$ and so $F = QS$ is a slope $\leq h$ -factorization.

Conversely, assume that F has a slope $\leq h$ -factorization $F = QS$. By Lemma 2.2.7 Q and S are relatively prime, so we may apply Theorem 2.2.2 to get a u -stable decomposition $M = \text{Ker } Q^*(u) \oplus N$. It is easy to see that this decomposition is functorial, as is a slope $\leq h$ -factorization, so it suffices to prove that this decomposition is a slope $\leq h$ -decomposition. First, since Q has slope $\leq h$ we see that $\text{Ker } Q^*(u) \subseteq M_{\leq h}$ (and we know it's finitely generated). It remains to show that for every multiplicative polynomial P of slope $\leq h$, $P^*(u)$ is invertible on N . By Lemma 2.2.7 P and S are relatively prime. Since $S = \det(1 - Tu \mid N)$ (by Theorem 2.2.2), it follows from [Buzzard 2007, Lemma 3.1] that P^* is invertible on N , as desired. \square

Corollary 2.2.14. *With notation and assumptions as in the theorem, a slope $\leq h$ -decomposition of M is functorial if and only if it is functorial for the natural map $R \rightarrow K_x$ for all $x \in \mathcal{M}(R)$.*

2.3. Fredholm hypersurfaces. In this section we discuss the notion of Fredholm hypersurfaces and relate this to slope factorizations and decompositions. We will use Huber’s adic spaces as our framework for nonarchimedean geometry, and we will use standard notions and notation from this theory freely, referring to the basic references [Huber 1994; 1996].

Any Tate ring R with a Noetherian ring of definition has an associated affinoid adic space $\mathrm{Spa}(R, R^+)$, for any ring of integral elements R^+ , by [Huber 1994, Theorem 2.5]. Fix an R^+ and consider $X = \mathrm{Spa}(R, R^+)$. We will frequently be interested in affine 1-space over X . As an adic space over (\mathbb{Z}, \mathbb{Z}) , we have $\mathbb{A}^1 = \mathrm{Spa}(\mathbb{Z}[T], \mathbb{Z})$; it represents the functor $X \mapsto \mathcal{O}(X)$ on the category of adic spaces (we note that the functor $X \mapsto \mathcal{O}^+(X)$ is represented by the “closed unit disc” $\mathrm{Spa}(\mathbb{Z}[T], \mathbb{Z}[T])$). The fibre product $\mathbb{A}_X^1 := X \times_{\mathrm{Spa}(\mathbb{Z}, \mathbb{Z})} \mathbb{A}^1$ exists, but it is no longer affinoid. Indeed, if we pick a topologically nilpotent unit $\varpi \in R$, it can be checked that the fibre product is given by

$$\mathbb{A}_X^1 = \bigcup_{m \geq 0} \mathrm{Spa}(R\langle \varpi^m T \rangle, R^+\langle \varpi^m T \rangle)$$

with respect to the transition maps coming from the natural inclusions. The ring of global functions on \mathbb{A}_X^1 is the ring of entire power series $R\{\{T\}\}$. Pick a topologically nilpotent unit $\varpi \in R$. If $h \in \mathbb{Q}$ then, writing $h = m/n$ with $m \in \mathbb{Z}$ and $n \in \mathbb{Z}_{\geq 1}$, we define an affinoid subset $\mathbb{B}_{X,h} \subseteq \mathbb{A}_X^1$ by

$$\mathbb{B}_{X,h} = \{|T^n| \leq |\varpi^{-m}|\} \subseteq \mathbb{A}_X^1.$$

We have $\mathbb{A}_X^1 = \bigcup_{h \in \mathbb{Q}} \mathbb{B}_{X,h}$.

Let R be a complete Tate ring with a Noetherian ring of definition and let $F \in R\{\{T\}\}$ be a Fredholm series. Put $X = \mathrm{Spa}(R, R^\circ)$. The closed subvariety $Z(F) := \{F = 0\} \subseteq \mathbb{A}_X^1$ is called the *Fredholm hypersurface* of F , or sometimes the *spectral variety* of F . It carries a projection map $Z(F) \rightarrow X$, which is flat, locally quasifinite and partially proper by [Andreatta et al. 2018, Théorème B.1].

Definition 2.3.1. Let R be a complete Tate ring with a Noetherian ring of definition, and pick a topologically nilpotent unit $\varpi \in R$. Let F be a Fredholm series with Fredholm hypersurface $Z = Z(F) \subseteq \mathbb{A}_X^1$, where $X = \mathrm{Spa}(R, R^\circ)$. Let $h \in \mathbb{Q}_{\geq 0}$ and let $U \subseteq X$ be an open affinoid in X ; put $Z_{U,h} = Z \cap \mathbb{B}_{U,h} \subseteq \mathbb{A}_X^1$ (this is an open affinoid subset of Z). We say that the pair (U, h) is a *slope datum* for (X, F) if $Z_{U,h} \rightarrow U$ is finite of constant degree (if the pair (X, F) is clear from the context, we occasionally just say that (U, h) is a slope datum).

Theorem 2.3.2. *Let R be a complete Tate ring with a Noetherian ring of definition, and pick a topologically nilpotent unit $\varpi \in R$. Let F be a Fredholm series over R with spectral variety $Z = Z(F) \subseteq \mathbb{A}_X^1$, where $X = \mathrm{Spa}(R, R^\circ)$. Let $U \subseteq X$ be an open affinoid and let $h \in \mathbb{Q}_{\geq 0}$. Then:*

- (1) *(U, h) is a slope datum for (X, F) if and only if F has a slope $\leq h$ -factorization in $\mathcal{O}_X(U)\{\{T\}\}$.*
- (2) *The collection of all $Z_{U,h}$ for all slope data (U, h) is an open cover of Z .*

Proof. This follows almost directly from [Andreatta et al. 2018, Théorème B.1, Corollaire B.1]. The second assertion is shown by tracing through the proof of [loc. cit., Lemme B.1, Théorème B.1]; adapting the proof of [loc. cit., Théorème B.1] slightly one sees that one may take the sets to be of the form $Z_{U,h}$ (the degree is locally constant on U , so constancy of the degree can be arranged). For the first assertion, the statement that if (U, h) is a slope datum then F has a slope $\leq h$ -factorization in $\mathcal{O}_X(U)\{\{T\}\}$ is [loc. cit., Corollaire B.1]. Conversely, if $F = QS$ is a slope $\leq h$ -factorization in $\mathcal{O}_X(U)\{\{T\}\}$, then S is a unit in $\mathcal{O}(\mathbb{B}_{U,h})$. Therefore $\mathcal{O}(Z_{U,h}) = \mathcal{O}(\mathbb{B}_{U,h})/(F) = \mathcal{O}(\mathbb{B}_{U,h})/(Q)$ is finite of constant degree equal to $\deg Q$ over U , and hence $Z_{U,h} \rightarrow U$ is finite of constant degree. \square

More generally, let X be an analytic adic space locally of the form $\mathrm{Spa}(R, R^\circ)$ for R a complete Tate ring with a Noetherian ring of definition, and let F be a Fredholm series over X with Fredholm hypersurface Z . If $U \subseteq X$ is an open affinoid and $h \in \mathbb{Q}_{\geq 0}$, we say that (U, h) is a slope datum for (X, F) if $\mathcal{O}(U)$ is Tate and there is a topologically nilpotent unit $\varpi \in \mathcal{O}(U)$ such that $Z_{U,h}$, defined using this choice of ϖ , is finite flat of constant degree over U .

When constructing eigenvarieties, it will be useful to consider a slightly more general notion. Let X be an analytic adic space as above and let F be a Fredholm series over X , with associated hypersurface Z . Write $\pi : Z \rightarrow X$ for the projection. We let $\mathcal{Cov}(Z)$ denote the set of all open affinoid $V \subseteq Z$ such that $\pi(V) \subseteq X$ is open affinoid, $\mathcal{O}(\pi(V))$ is Tate, and the map $\pi|_V : V \rightarrow \pi(V)$ is finite of constant degree. Then we have the following theorem.

Theorem 2.3.3. *Keep the notation and assumptions of the paragraph above. Then $\mathcal{Cov}(Z)$ is an open cover of Z . If $V \in \mathcal{Cov}(Z)$, then there exists a factorization $F = QS$ in $\mathcal{O}(\pi(V))\{\{T\}\}$, where Q is a multiplicative polynomial of degree $\deg \pi|_V$, S is a Fredholm series, Q and S are relatively prime, and we have*

$$\mathcal{O}(V) = \mathcal{O}(\pi(V))[T]/(Q) \quad \text{and} \quad \mathcal{O}^+(V) = (\mathcal{O}(\pi(V))[T]/(Q))^\circ.$$

Conversely, if such a factorization of F exists in $\mathcal{O}(U)\{\{T\}\}$, where $U \subseteq X$ is open affinoid and $\mathcal{O}(U)$ is Tate, then $V = \mathrm{Spa}(\mathcal{O}(U)[T]/(Q), (\mathcal{O}(U)[T]/(Q))^\circ)$ is naturally an element of $\mathcal{Cov}(Z)$.

Proof. The assertions about rings of integral elements follow immediately from the rest by Lemma A.3. The first two parts are [Andreatta et al. 2018, Théorème B.1, Corollaire B.1]. Note that if $\pi : V \rightarrow U$ is a finite flat morphism, with $U \subset X$ and $V \subset Z$ open affinoid, then π is open by [Huber 1996, Lemma 1.7.9]. For the last part, it is clear that $V \rightarrow U$ is finite and surjective of constant degree $\deg Q$, so it remains to see that V is naturally an open subset of Z . For this we may work locally over U . Set $B = \mathcal{O}(U)$. For each n we have compatible morphisms

$$B[T]/(Q) \rightarrow B\langle \varpi^n T \rangle / (Q) \leftarrow B\langle \varpi^n T \rangle / (F).$$

The second map is the projection onto the first factor in the decomposition

$$B\langle \varpi^n T \rangle / (F) \cong B\langle \varpi^n T \rangle / (Q) \times B\langle \varpi^n T \rangle / (S)$$

which results from the fact that Q and S are relatively prime. Thus $\{Q = 0\} \subseteq Z$ is open and closed in Z . Moreover, when n is sufficiently large, we claim that the first map is an isomorphism. To see this, consider the quotient map $p : B[T] \rightarrow B[T]/(Q)$ and equip the target with a submultiplicative norm that induces the canonical topology. For large enough n we will have $|p(\varpi^n T)| \leq 1$ and hence $p(B_0[\varpi^n T]) \subseteq (B[T]/(Q))_0$ (here we are using $-_0$ to denote unit balls), so p is continuous for the topology on $B[T]$ coming from the inclusion $B[T] \subseteq B\langle \varpi^n T \rangle$, and we may complete to obtain a morphism $B\langle \varpi^n T \rangle \rightarrow B[T]/(Q)$ with kernel $QB\langle \varpi^n T \rangle$. This gives an inverse to the map $B[T]/(Q) \rightarrow B\langle \varpi^n T \rangle/(Q)$, proving the claim. Thus we may identify V with $\{Q = 0\} \subseteq Z$, which shows that V is naturally an open subset of Z . \square

3. Relative distribution algebras

3.1. Relative distribution algebras and norms. A p -adic analytic group will in this paper always mean a \mathbb{Q}_p -analytic group. Let R be a Banach–Tate ring. We denote the unit ball of R by R_0 . If there exists a norm-decreasing homomorphism $\mathbb{Z}_p \rightarrow R$, where we equip \mathbb{Z}_p with the usual norm $|x|_p = p^{-\text{ord}_p(x)}$, we call such an R (together with the map $\mathbb{Z}_p \rightarrow R$) a *Banach–Tate \mathbb{Z}_p -algebra*. The goal of this section is to extend some of the constructions of [Schneider and Teitelbaum 2003, §4] to the case of continuous functions and distributions valued in such R . In particular, we construct R -valued analogues of (locally) analytic distribution algebras for compact p -adic analytic groups. We begin with a lemma on the existence of Banach–Tate \mathbb{Z}_p -algebra norms.

Lemma 3.1.1. *Let R be a Noetherian Banach–Tate ring with norm $|\cdot|$ and a multiplicative pseudouniformizer ϖ . Assume that there exists a continuous homomorphism $\mathbb{Z}_p \rightarrow R$ (necessarily unique). Then there exists a Banach–Tate \mathbb{Z}_p -algebra norm $|\cdot|'$ on R which is bounded-equivalent to $|\cdot|^s$ for some $s > 0$, and such that ϖ is a multiplicative pseudouniformizer for $|\cdot|'$.*

Proof. Note first that a norm $|\cdot|'$ on R is a Banach–Tate \mathbb{Z}_p -algebra norm if and only if $|p|' \leq p^{-1}$, so we need to check this. By continuity of $\mathbb{Z}_p \rightarrow R$ we have $p \in R^{\circ\circ}$. Choose $m \in \mathbb{Z}_{\geq 1}$ such that $|p|_{\text{sp}} < |\varpi|^{2/m}$ and consider the finite free R -algebra

$$S = R[\varpi^{1/m}] = R[X]/(X^m - \varpi).$$

We equip S with its canonical topology as a finite R -module; then the induced subspace topology on $R \subseteq S$ agrees with the original topology on S . Thus we have $p, \varpi^{1/m} \in S^{\circ\circ}$. Now equip S with a submultiplicative R -Banach module norm $|\cdot|_S$ that induces the canonical topology. Note that ϖ is a multiplicative pseudouniformizer for $|\cdot|_S$ with $|\varpi|_S = |\varpi|$, and that $(R, |\cdot|) \rightarrow (S, |\cdot|_S)$ is norm-decreasing. We then have $|p\varpi^{-1/m}|_{S, \text{sp}} < |\varpi|^{1/m} < 1$ by construction, so $p\varpi^{-1/m}$ is topologically nilpotent in S . We can then choose a ring of definition S_2 of S containing $\varpi^{1/m}$ and $p\varpi^{-1/m}$ and consider the norm

$$|s|_2 = \inf\{|\varpi|^{k/m} \mid s \in \varpi^{k/m} S_2\}.$$

Since $p \in \varpi^{1/m} S_2$ we have $|p|_2 < 1$, and we may hence find $s > 0$ such that $|p|_2^s \leq p^{-1}$. Restricting the norm $|\cdot|' := |\cdot|_2^s$ to $R \subseteq S$ then gives the desired norm. \square

Definition 3.1.2. Let X be a compact topological space and let A be a topological \mathbb{Z}_p -algebra:

- (1) We let $\mathcal{C}(X, A)$ denote the A -module of all continuous A -valued functions on X , and let $\mathcal{C}_{\text{sm}}(X, A)$ denote the subspace of all locally constant functions.
- (2) We put $\mathcal{D}(X, A) = \text{Hom}_{A, \text{cts}}(\mathcal{C}(X, A), A)$.

When A is a normed ring, we may topologize $\mathcal{C}(X, A)$ and $\mathcal{C}_{\text{sm}}(X, A)$ using the supremum norm, and we may give $\mathcal{D}(X, A)$ the corresponding dual/operator norm. If A is complete, this makes $\mathcal{C}(X, A)$ into a complete A -module. When the topology on X is profinite, $\mathcal{C}_{\text{sm}}(X, A)$ is dense in $\mathcal{C}(X, A)$ and, if R is a Banach–Tate \mathbb{Z}_p -algebra, the natural map $\mathcal{C}(X, \mathbb{Z}_p) \widehat{\otimes}_{\mathbb{Z}_p} R \rightarrow \mathcal{C}(X, R)$ is a topological isomorphism. Similarly $\mathcal{C}(X, \mathbb{Z}_p) \widehat{\otimes}_{\mathbb{Z}_p} R_0 \cong \mathcal{C}(X, R_0)$, where R_0 is the unit ball of R .

Continue to let X be a profinite set and R a Banach–Tate \mathbb{Z}_p -algebra with unit ball R_0 and a multiplicative pseudouniformizer ϖ . Note that $\mathcal{D}(X, R_0) = \text{Hom}_{R_0}(\mathcal{C}(X, R_0), R_0)$ (i.e., continuity with respect to the ϖ -adic topology is automatic) and that this is the unit ball in $\mathcal{D}(X, R)$. We may equip $\mathcal{D}(X, R_0)$ with the weak topology coming from the family of maps $\mathcal{D}(X, R_0) \rightarrow R_0$ given by $\mu \mapsto \mu(f)$ for $f \in \mathcal{C}(X, R_0)$ and the ϖ -adic topology on R_0 . We will refer to this topology as the *weak-star topology* on $\mathcal{D}(X, R_0)$. When $X = G$ is a profinite group, $\mathcal{D}(G, R)$ carries a convolution product

$$(\mu * \nu)(f) = \mu(g \mapsto \nu(h \mapsto f(gh))).$$

One checks directly that $\delta_g * \delta_h = \delta_{gh}$ for all $g, h \in G$, where δ_g denotes the Dirac distribution at g . This product preserves $\mathcal{D}(G, R_0)$. We sum up some basic properties of the weak-star topology.

Lemma 3.1.3. *If X is finite (hence discrete) the weak-star topology on $\mathcal{D}(X, R_0)$ coincides with the ϖ -adic topology. In general, if $X = \varprojlim_n X_n$ is an inverse limit of finite sets X_n , we have a natural isomorphism $\mathcal{D}(X, R_0) \cong \varprojlim_n \mathcal{D}(X_n, R_0)$ which identifies the weak-star topology on the source with the inverse limit topology, where $\mathcal{D}(X_n, R_0)$ is equipped with the ϖ -adic topology. When $X = G$ is a profinite group this is a ring homomorphism, and multiplication on $\mathcal{D}(G, R_0)$ is jointly continuous with respect to the weak-star topology.*

Proof. The first assertion is straightforward. For the second, note that the formation of $\mathcal{D}(X, R_0)$ is covariantly functorial in X , so the maps $X \rightarrow X_n$ induce a natural map $\mathcal{D}(X, R_0) \rightarrow \varprojlim_n \mathcal{D}(X_n, R_0)$ which is continuous by the first assertion when we equip the source and the target with the topologies in the statement of the lemma. Moreover it is easily checked to be a ring homomorphism when $X = G$ is a profinite group. Unraveling, we see that this morphism is the natural morphism

$$\text{Hom}_{R_0}(\mathcal{C}(X, R_0), R_0) \rightarrow \text{Hom}_{R_0}(\mathcal{C}_{\text{sm}}(X, R_0), R_0)$$

induced by the inclusion $\mathcal{C}_{\text{sm}}(X, R_0) \subseteq \mathcal{C}(X, R_0)$. Since this subspace is dense for the ϖ -adic topology, we see that the map is an isomorphism. To check that it also a homeomorphism, note first that by the same density one may define the weak-star topology using only locally constant functions. It is then straightforward to check that all basic opens from locally constant functions come by pullback from basic

opens on the $\mathcal{D}(X_n, R_0)$, which implies that the map is a homeomorphism. Finally, multiplication is jointly continuous for the ϖ -adic topology on $\mathcal{D}(G_n, R_0)$ for all n , and hence jointly continuous on the inverse limit. This implies the last assertion. \square

Let $X = \varprojlim_n X_n$ be a countable inverse limit of finite sets, viewed as a profinite set. We define $R_0\llbracket X \rrbracket := \varprojlim_n R_0[X_n]$ and $R\llbracket X \rrbracket := (\varprojlim_n R_0[X_n])[1/\varpi]$; these are R_0 - and R -modules, respectively, and independent of the choice of the X_n 's (here, if B is a ring and S is a finite set, $B[S]$ denotes the free B -module generated by the set S). If the X_n are groups (so X is a profinite group) then they carry natural algebra structures. We may topologize $R_0\llbracket X \rrbracket$ in two ways; either giving it the natural inverse limit topology or the ϖ -adic topology. We give $R\llbracket X \rrbracket$ the topology induced from the ϖ -adic topology on $R_0\llbracket X \rrbracket$, which is compatible with viewing $R\llbracket X \rrbracket$ as an R -Banach module with unit ball $R_0\llbracket X \rrbracket$.

Proposition 3.1.4. *Let $X = \varprojlim_n X_n$ be a profinite set. There is a natural R -Banach module isomorphism $R\llbracket X \rrbracket \rightarrow \mathcal{D}(X, R)$ sending $[x]$ to δ_x . It restricts to an R_0 -module isomorphism $R_0\llbracket X \rrbracket \rightarrow \mathcal{D}(X, R_0)$ which identifies the inverse limit topology on the source with the weak-star topology on the target. If $X = G$ is a profinite group, then these maps are ring homomorphisms.*

Proof. Define compatible maps $R_0[X_n] \rightarrow \text{Hom}_{R_0}(\mathcal{C}(X_n, R_0), R_0)$ by $[x] \mapsto \delta_x$; one checks directly that this is an isomorphism of topological R_0 -modules, and that it is a ring homomorphism when X is a profinite group. Taking inverse limits we get

$$R_0\llbracket X \rrbracket \xrightarrow{\sim} \text{Hom}_{R_0}(\mathcal{C}_{\text{sm}}(X, R_0), R_0) = \mathcal{D}(X, R_0).$$

Lemma 3.1.3 shows that this identifies the inverse limit topology on the source with the weak-star topology on the target. Inverting ϖ we get the desired isomorphism $R\llbracket X \rrbracket \rightarrow \mathcal{D}(X, R)$, which is clearly a Banach module isomorphism since it identifies the respective unit balls $R_0\llbracket X \rrbracket$ and $\mathcal{D}(X, R_0)$. \square

Recall the notion of a uniform pro- p group from [Dixon et al. 1999, Definition 4.1]. When G is a uniform pro- p group, $\mathbb{Z}_p\llbracket G \rrbracket$ may be identified with a ring of noncommutative formal power series

$$\left\{ \sum_{\alpha} d_{\alpha} \mathbf{b}^{\alpha} \mid d_{\alpha} \in \mathbb{Z}_p \right\},$$

where d is the dimension of G , $\alpha = (\alpha_1, \dots, \alpha_d) \in \mathbb{Z}_{\geq 0}^d$ is a multi-index, g_1, \dots, g_d is a minimal set of topological generators of G , $b_i = [g_i] - 1$ and $\mathbf{b}^{\alpha} := b_1^{\alpha_1} \cdots b_d^{\alpha_d}$. Our next goal is to show that the analogous description holds for $R_0\llbracket G \rrbracket$, with the same commutation relations between the \mathbf{b}^{α} .

Proposition 3.1.5. *Let $G = \mathbb{Z}_p^d$. For $\alpha \in \mathbb{Z}_{\geq 0}^d$, let $E_{\alpha} : \mathbb{Z}_p^d \rightarrow \mathbb{Z}_p$ denote the function*

$$E_{\alpha}(x_1, \dots, x_d) = \binom{x_1}{\alpha_1} \cdots \binom{x_d}{\alpha_d}.$$

Then the Amice transform

$$\mu \mapsto \sum_{\alpha} \mu(E_{\alpha}) T_1^{\alpha_1} \cdots T_d^{\alpha_d}$$

defines an algebra isomorphism $\mathcal{D}(\mathbb{Z}_p^d, R_0) \xrightarrow{\sim} R_0[[T_1, \dots, T_d]]$ which identifies the weak-star topology on the source with the product topology $R_0[[T_1, \dots, T_d]] = \prod_{\alpha} R_0 \cdot T_1^{\alpha_1} \cdots T_d^{\alpha_d}$ on the target.

Proof. Once again this is a simple extension of a well-known result, so we content ourselves with a sketch. The key observation is that $\mathcal{C}(\mathbb{Z}_p^d, R_0) \cong \mathcal{C}(\mathbb{Z}_p^d, \mathbb{Z}_p) \widehat{\otimes}_{\mathbb{Z}_p} R_0$. Then it is clear that

$$\mathcal{D}(\mathbb{Z}_p^d, R_0) \xrightarrow{\sim} \prod_{\alpha} R_0$$

via $\mu \mapsto (\mu(E_{\alpha}))_{\alpha}$ and it is straightforward to check that this identifies the weak topology on the source with the product topology on the target (it is the statement that the E_{α} suffice to define the weak-star topology). To finish, we remark that the computation that the algebra structures match up is identical to the well-known one in the case $R_0 = \mathbb{Z}_p$. \square

Note that the topology on $R_0[[T_1, \dots, T_d]]$ described in the proposition is equal to the $(\varpi, T_1, \dots, T_d)$ -adic topology. Let us return to the case of a general uniform pro- p group G . The ring $\mathbb{Z}_p[[G]]$ is described by formal power series as above. Following [Schneider and Teitelbaum 2003], let us define elements $c_{\beta\gamma, \alpha} \in \mathbb{Z}_p$ by

$$\mathbf{b}^{\beta} \mathbf{b}^{\gamma} = \sum_{\alpha} c_{\beta\gamma, \alpha} \mathbf{b}^{\alpha}. \quad (3.1.1)$$

We remark that, for fixed α , $c_{\beta\gamma, \alpha} \rightarrow 0$ as $|\beta| + |\gamma| \rightarrow +\infty$ (here and elsewhere, for a multi-index α we define $|\alpha| = \alpha_1 + \dots + \alpha_d$). This follows from [loc. cit., Lemma 4.1(ii)].

Proposition 3.1.6. *Let G be a uniform pro- p group and use the notation above. Then $R_0[[G]]$ may be identified with the ring of formal power series*

$$\left\{ \sum_{\alpha} d_{\alpha} \mathbf{b}^{\alpha} \mid d_{\alpha} \in R_0 \right\}$$

with multiplication given by

$$\left(\sum_{\beta} d_{\beta} \mathbf{b}^{\beta} \right) \left(\sum_{\gamma} e_{\gamma} \mathbf{b}^{\gamma} \right) = \sum_{\alpha} \left(\sum_{\beta, \gamma} d_{\beta} e_{\gamma} c_{\beta\gamma, \alpha} \right) \mathbf{b}^{\alpha}.$$

Proof. The choice g_1, \dots, g_d of a minimal (ordered) set of topological generators identifies G , as p -adic analytic manifold, with \mathbb{Z}_p^d . Thus we get a topological isomorphism $R_0[[G]] \cong R_0[[\mathbb{Z}_p^d]]$ of R_0 -modules (for both the weak-star and the ϖ -adic topology). Proposition 3.1.5 then implies the description in terms of power series. To see that the multiplication works out as described, note that the natural map $\mathbb{Z}_p[[G]] \rightarrow R_0[[G]]$ is an algebra homomorphism; hence the above formula is true for products of monomials. We can then deduce the formula in general by noting that R_0 is central in $R_0[[G]]$ and that the subring generated by R_0 and the image of $\mathbb{Z}_p[[G]]$ (for which the formula holds) is dense in $R_0[[G]]$ with respect to the weak-star topology, and that multiplication is jointly continuous for the weak-star topology. \square

Inverting ϖ we get an explicit description of $R[[G]]$ when G is uniform. Using this we may now define a family of norms on $R[[G]]$ following [Schneider and Teitelbaum 2003, §4]. We continue to fix a minimal ordered set of topological generators g_1, \dots, g_d .

Definition 3.1.7. Let $r \in [1/p, 1)$. We define the r -norm $\| - \|_r$ on $R[[G]]$ by the formula

$$\left\| \sum_{\alpha} d_{\alpha} \mathbf{b}^{\alpha} \right\|_r = \sup_{\alpha} |d_{\alpha}| r^{|\alpha|}.$$

Recall that we have fixed a choice of norm $| - |$ on R such that $|z| \leq |z|_p$ for all $z \in \mathbb{Z}_p$. This will be convenient for some calculations. Note that the definition of $\| - \|_r$ a priori depends on the choice of generators. We remark that for all $r \in [1/p, 1)$, $\| - \|_r$ induces the weak-star topology on $R_0[[G]]$ by Proposition 3.1.5 and a straightforward calculation. It follows that any homomorphism $G \rightarrow H$ of uniform groups induces a continuous homomorphism $R_0[[G]] \rightarrow R_0[[H]]$ when the source and target are equipped with (any) r -norms, since this is true for the weak-star topology using the characterization in Lemma 3.1.3.

Proposition 3.1.8. *The norm $\| - \|_r$ is independent of the choice of minimal ordered set of topological generators for G , and is submultiplicative. Finally, if we replace the norm $| - |$ on R by a bounded-equivalent \mathbb{Z}_p -algebra norm $| - |'$, then the resulting norm $\| - \|'_r$ is bounded-equivalent to $\| - \|_r$.*

Proof. For the proof of independence, we follow the discussion after the proof of [Schneider and Teitelbaum 2003, Theorem 4.10]. Let g'_1, \dots, g'_d be a different choice and set $b'_i = [g'_i] - 1 \in \mathbb{Z}_p[[G]]$ etc., and let $\| - \|'_r$ denote the r -norm with respect to this choice. We may write

$$\mathbf{b}'^{\beta} = \sum_{\alpha} c_{\beta, \alpha} \mathbf{b}^{\alpha}$$

in $\mathbb{Z}_p[[G]]$, and one has $|c_{\beta, \alpha}|_p r^{|\alpha|} \leq r^{|\beta|}$ (see [loc. cit.]). Transporting this to $R[[G]]$ we have the same identity, and the inequality $|c_{\beta, \alpha}| r^{|\alpha|} \leq r^{|\beta|}$ (since $|c_{\beta, \alpha}| \leq |c_{\beta, \alpha}|_p$). Expanding out a general element $\mu \in R[[G]]$ we then have

$$\mu = \sum_{\beta} d'_{\beta} \mathbf{b}'^{\beta} = \sum_{\alpha} \left(\sum_{\beta} d'_{\beta} c_{\beta, \alpha} \right) \mathbf{b}^{\alpha}.$$

We then have

$$\|\mu\|_r \leq \sup_{\beta, \alpha} |d'_{\beta}| |c_{\beta, \alpha}| r^{|\alpha|} \leq \sup_{\beta} |d'_{\beta}| r^{|\beta|} = \|\mu\|'_r.$$

By symmetry, we must have equality.

To prove submultiplicativity, we follow the proof of [Schneider and Teitelbaum 2003, Proposition 4.2]. Recall the $c_{\beta\gamma, \alpha}$ from (3.1.1). By [loc. cit., Lemma 4.1(ii)] we have $|c_{\beta\gamma, \alpha}| r^{|\alpha|} \leq |c_{\beta\gamma, \alpha}|_p r^{|\alpha|} \leq r^{|\beta|+|\gamma|}$ for all α, β, γ . Let $\mu = \sum_{\beta} d_{\beta} \mathbf{b}^{\beta}$ and $\nu = \sum_{\gamma} e_{\gamma} \mathbf{b}^{\gamma}$ be elements of $R[[G]]$; then we have

$$\mu * \nu = \sum_{\alpha} \left(\sum_{\beta, \gamma} d_{\beta} e_{\gamma} c_{\beta\gamma, \alpha} \right) \mathbf{b}^{\alpha}$$

and we can calculate

$$\|\mu\nu\|_r = \sup_{\alpha} \left| \sum_{\beta, \gamma} d_{\beta} e_{\gamma} c_{\beta\gamma, \alpha} \right| r^{|\alpha|} \leq \sup_{\beta, \gamma} |d_{\beta}| |e_{\gamma}| r^{|\beta|+|\gamma|} = \|\mu\|_r \|\nu\|_r,$$

where we use submultiplicativity and $|c_{\beta\gamma,\alpha}|r^{|\alpha|} \leq r^{|\beta|+|\gamma|}$ to obtain the middle inequality. This finishes the proof of submultiplicativity.

Finally, suppose we have $C_1|x| \leq |x'| \leq C_2|x|$ for all $x \in R$. Then $C_1|d_\alpha|r^{|\alpha|} \leq |d_\alpha|'r^{|\alpha|} \leq C_2|d_\alpha|r^{|\alpha|}$ for all α which implies

$$C_1 \left\| \sum_{\alpha} d_{\alpha} \mathbf{b}^{\alpha} \right\|_r \leq \left\| \sum_{\alpha} d_{\alpha} \mathbf{b}^{\alpha} \right\|_r' \leq C_2 \left\| \sum_{\alpha} d_{\alpha} \mathbf{b}^{\alpha} \right\|_r$$

as desired. \square

Before moving on to general compact p -adic analytic G , we record a few properties of the r -norms.

Lemma 3.1.9. *Let $r \in [1/p, 1)$. If $g \in G$, then $\|[g]\|_r = 1$ and $\|[g]\mu\|_r = \|\mu[g]\|_r = \|\mu\|_r$ for all $\mu \in R[[G]]$. Moreover, if ϕ is an automorphism of G (of p -adic analytic groups), then ϕ induces an automorphism of $R[[G]]$ satisfying $\|\phi(\mu)\|_r = \|\mu\|_r$ for all $\mu \in R[[G]]$.*

Proof. The first statement follows from the fact that the expansion of $[g]$ has coefficients in \mathbb{Z}_p and the constant term is 1. The second is an easy consequence of the first and submultiplicativity (since $[g]^{-1} = [g^{-1}]$ also has norm 1).

For the final statement, observe that if g_1, \dots, g_d is a set of topological generators then so are $\phi(g_1), \dots, \phi(g_d)$. Since the r -norms are independent of the choice of generators, we conclude that if $\mu = \sum_{\alpha} d_{\alpha} \mathbf{b}^{\alpha}$, then

$$\|\phi(\mu)\|_r = \left\| \sum_{\alpha} d_{\alpha} (\phi(\mathbf{b}))^{\alpha} \right\|_r = \sup_{\alpha} |d_{\alpha}| r^{|\alpha|} = \|\mu\|_r. \quad \square$$

We will use the last property mostly in the case when H is a compact p -adic analytic group, $N \subseteq H$ is a uniform open normal subgroup, and ϕ is the automorphism of N given by conjugation by some $h \in H$.

Now let G be an arbitrary compact p -analytic group. Pick a uniform open normal subgroup N and a set h_1, \dots, h_t of coset representatives of G/N . Any $\mu \in R[[G]]$ may be written uniquely as $\mu = \sum_i [h_i] \mu_i$ with $\mu_i \in R[[N]]$, and we define a norm $\| - \|_{N,r}$ on $R[[G]]$ by

$$\|\mu\|_{N,r} = \sup_i \|\mu_i\|_r.$$

We could alternatively take a right coset decomposition $\mu = \sum_i v_i [h_i]$ with $v_i \in R[[N]]$, and define $\| - \|_{N,r}^{\text{right}}$ on $R[[G]]$ by

$$\|\mu\|_{N,r}^{\text{right}} = \sup_i \|v_i\|_r.$$

Proposition 3.1.10. *We have $\| - \|_{N,r} = \| - \|_{N,r}^{\text{right}}$. The definition is also independent of the choice of coset representatives. Moreover, $\| - \|_{N,r}$ is submultiplicative and satisfies $\|[g]\|_{N,r} = 1$ and $\|[g]\mu\|_{N,r} = \|\mu[g]\|_{N,r} = \|\mu\|_{N,r}$ for all $g \in G$ and $\mu \in R[[G]]$.*

Proof. For left/right independence, note that $\mu = \sum_i ([h_i] \mu_i [h_i^{-1}]) [h_i]$ and hence

$$\|\mu\|_{N,r}^{\text{right}} = \sup_i \|[h_i] \mu_i [h_i^{-1}]\|_r = \sup_i \|\mu_i\|_r = \|\mu\|_{N,r}$$

by Lemma 3.1.9. For independence of the coset representatives, suppose we have h'_i a different set with $h_i = h'_i n_i$; then $\mu = \sum_i [h'_i]([n_i]\mu_i)$ and hence

$$\|\mu\|'_{N,r} = \sup_i \|[n_i]\mu_i\|_r = \sup_i \|\mu_i\|_r = \|\mu\|_{N,r}$$

by Lemma 3.1.9 again. Next we prove submultiplicativity. Define $k(i, j)$ by $h_i h_j = h_{k(i,j)} n_{ij}$. For $\mu = \sum_i [h_i]\mu_i$ and $\nu = \sum_j [h_j]\nu_j$ in $R[[G]]$, we have

$$\mu\nu = \sum_k [h_k] \left(\sum_{i,j:k(i,j)=k} [n_{ij}]([h_j^{-1}]\mu_i[h_j])\nu_j \right).$$

Using Lemma 3.1.9 and submultiplicativity for $\|\cdot\|_r$ on N one then sees easily that $\|\mu\nu\|_{N,r} \leq \|\mu\|_{N,r} \|\nu\|_{N,r}$.

Next, let $g \in G$. Writing $g = h_i n$ for some i and $n \in N$, we see that $\|[g]\|_{N,r} = \|[n]\|_r = 1$ by Lemma 3.1.9. Finally, the last property then follows by the same argument as in the proof of Lemma 3.1.9. \square

As the notation suggests, $\|\cdot\|_{N,r}$ *does* depend on the choice of N . For a study of how the completions change when one changes the subgroup in certain situations, see [Ardakov and Wadsley 2013, §10.6–10.8].

3.2. Completions. In this section we study the case when G is a uniform pro- p group in more detail. Let R be a Banach–Tate \mathbb{Z}_p -algebra with multiplicative pseudouniformizer ϖ as usual.

Definition 3.2.1. For $r \in [1/p, 1)$, define $\mathcal{D}^r(G, R)$ to be the completion of $\mathcal{D}(G, R)$ with respect to the norm $\|\cdot\|_r$, and we let $\mathcal{D}(G, R)$ denote the completion of $\mathcal{D}(G, R)$ with respect to the entire family of norms $(\|\cdot\|_r)_{r \in [1/p, 1)}$.

Remark 3.2.2. Note that if we change the norm on R to a bounded-equivalent one, then the completion $\mathcal{D}^r(G, R)$ is unchanged, by Proposition 3.1.8.

The motivation for this definition is that if R is a Banach \mathbb{Q}_p -algebra, $\mathcal{D}(G, R)$ is naturally isomorphic to the space of locally analytic R -valued distributions on G , by Proposition 3.2.9.

Note that there are natural norm-decreasing injective maps $\mathcal{D}^s(G, R) \rightarrow \mathcal{D}^r(G, R)$ whenever $r \leq s$ (which we will think of as inclusions), and that they fit together into an inverse system with limit $\mathcal{D}(G, R)$. The explicit description of $\mathcal{D}(G, R)$ from Proposition 3.1.6 gives us an explicit formal power series description

$$\mathcal{D}^r(G, R) = \left\{ \sum_{\alpha} d_{\alpha} \mathbf{b}^{\alpha} \mid d_{\alpha} \in R, |d_{\alpha}| r^{|\alpha|} \rightarrow 0 \right\}$$

and the norm is still given by $\|\sum d_{\alpha} \mathbf{b}^{\alpha}\|_r = \sup |d_{\alpha}| r^{|\alpha|}$. We see that, unlike $\mathcal{D}(G, R)$, the $\mathcal{D}^r(G, R)$ are naturally potentially ON-able, with a potential ON-basis given by the elements $\varpi^{-n(r, \varpi, \alpha)} \mathbf{b}^{\alpha}$, where

$$n(r, \varpi, \alpha) = \left\lfloor \frac{|\alpha| \log_p r}{\log_p |\varpi|} \right\rfloor. \quad (3.2.1)$$

Lemma 3.2.3. *If $r < s$ then the inclusion $\iota : \mathcal{D}^s(G, R) \hookrightarrow \mathcal{D}^r(G, R)$ is a compact map of R -Banach modules.*

Proof. For simplicity set $\mathcal{D}^r := \mathcal{D}^r(G, R)$ etc. for the duration of this proof. Choose a minimal set of topological generators g_1, \dots, g_d of G and set $b_i = [g_i] - 1$ as usual. For $n \geq 1$ define $T_n : \mathcal{D}^s \rightarrow \mathcal{D}^r$ by

$$T_n \left(\sum_{\alpha} d_{\alpha} \mathbf{b}^{\alpha} \right) = \sum_{|\alpha| < n} d_{\alpha} \mathbf{b}^{\alpha}.$$

By definition we see that T_n is of finite rank. Then if $\sum_{\alpha} d_{\alpha} \mathbf{b}^{\alpha}$ is in the unit ball of \mathcal{D}^s , i.e., $|d_{\alpha}| s^{|\alpha|} \leq 1$ for all α , we have

$$\left\| (\iota - T_n) \left(\sum_{\alpha} d_{\alpha} \mathbf{b}^{\alpha} \right) \right\|_r = \left\| \sum_{|\alpha| \geq n} d_{\alpha} \mathbf{b}^{\alpha} \right\|_r \leq (r/s)^n.$$

Thus $\|\iota - T_n\| \leq (r/s)^n$, where $\|\cdot\|$ is the operator norm on $\text{Hom}_{R, \text{cts}}(\mathcal{D}^s, \mathcal{D}^r)$, and hence $T_n \rightarrow \iota$ as $n \rightarrow \infty$, so ι is compact. \square

Lemma 3.2.4. *Let G and H be two uniform pro- p groups and assume that $f : G \rightarrow H$ is a homomorphism of p -adic analytic groups such that $f(G) \subseteq H^{p^n}$ for some $n \geq 0$. Then the induced map $f_* : \mathcal{D}(G, R) \rightarrow \mathcal{D}(H, R)$ is norm-decreasing when we equip $\mathcal{D}(G, R)$ with $\|\cdot\|_r$ and $\mathcal{D}(H, R)$ with $\|\cdot\|_{r^{1/p^n}}$. As a consequence, we get an induced map*

$$f_* : \mathcal{D}^r(G, R) \rightarrow \mathcal{D}^r(H, R)$$

which factors through the natural map $\mathcal{D}^{r^{1/p^n}}(H, R) \rightarrow \mathcal{D}^r(H, R)$. In particular, when $n \geq 1$, f_ is compact.*

Proof. We start with the first assertion. Note that the general case follows from two special cases: $n = 0$, f arbitrary, and $n = 1$, $G = H^p$ with f the inclusion. Indeed the general case can be written as a composition of these cases.

So, suppose first that $n = 0$. Scaling by powers of ϖ , it suffices to prove this for $\mathcal{D}(G, R_0)$. The map f_* is continuous with respect to the norm $\|\cdot\|_r$ (see the discussion after Definition 3.1.7). Let g_1, \dots, g_d be a minimal set of topological generators for G and let $b_i = [g_i] - 1$ as usual. Using that $\|[h] - 1\|_r \leq r$ for all $h \in H$, we see that

$$\left\| f_* \left(\sum d_{\alpha} \mathbf{b}^{\alpha} \right) \right\|_r = \left\| \sum d_{\alpha} f_*(\mathbf{b})^{\alpha} \right\|_r \leq \sup |d_{\alpha}| r^{|\alpha|} = \left\| \sum d_{\alpha} \mathbf{b}^{\alpha} \right\|_r$$

as desired, using continuity of f_* . This completes the proof of the first special case.

Next, we consider the second case $G = H^p \subseteq H$. Let $s = r^{1/p}$; since $r \geq p^{-1}$ we have $s \geq p^{-1/p} > p^{-1/(p-1)}$. Let h_1, \dots, h_d be a minimal set of topological generators for H . Then h_1^p, \dots, h_d^p form a minimal set of topological generators for H^p . Set $b_i = [h_i^p] - 1$ and $b'_i = [h_i] - 1$ (apologies for the mild abuse of notation). Then, inside $\mathcal{D}(H, R)$, we have

$$\|b_i\|_s = \|(1 + b'_i)^p - 1\|_s = \left\| \sum_{k=1}^p \binom{p}{k} (b'_i)^k \right\|_s = s^p = r$$

using $s > p^{-1/(p-1)}$ and $|p| \leq |p|_p = p^{-1}$. Thus, if $\sum d_\alpha \mathbf{b}^\alpha \in \mathcal{D}(H^p, R)$, then inside $\mathcal{D}(H, R)$ we have

$$\left\| \sum d_\alpha \mathbf{b}^\alpha \right\|_s \leq \sup |d_\alpha| r^\alpha,$$

which is equal to $\left\| \sum d_\alpha \mathbf{b}^\alpha \right\|_r$ computed inside $\mathcal{D}(H^p, R)$. This finishes the proof of first assertion. The remaining assertions are then easily verified (using Lemma 3.2.3 for last one). \square

Before discussing what happens when one changes the norm on R , we record the following important base change lemma.

Lemma 3.2.5. *Let R and S be Banach–Tate \mathbb{Z}_p -algebras, and let $f : R \rightarrow S$ be a bounded ring homomorphism. Suppose that there is a multiplicative pseudouniformizer $\varpi \in R$ such that $f(\varpi)$ is also multiplicative. Let $r \in [1/p, 1)$. Then the natural map $\mathcal{D}^r(G, R) \widehat{\otimes}_R S \rightarrow \mathcal{D}^r(G, S)$ is an isomorphism of Banach S -modules.*

Proof. Note that (since f is bounded) the fact that ϖ and $f(\varpi)$ are both multiplicative implies that $|\varpi| = |f(\varpi)|$. We recall the potential ON-bases $(\varpi^{-n(r, \varpi, \alpha)} \mathbf{b}^\alpha)_\alpha$ and $(f(\varpi)^{-n(r, f(\varpi), \alpha)} \mathbf{b}^\alpha)_\alpha$ of $\mathcal{D}^r(G, R)$ and $\mathcal{D}^r(G, S)$, respectively. It is straightforward to check that the tensor product norm on

$$\left(\bigoplus_\alpha R(\varpi^{-n(r, \varpi, \alpha)} \mathbf{b}^\alpha) \right) \otimes_R S = \bigoplus_\alpha S(f(\varpi)^{-n(r, f(\varpi), \alpha)} \mathbf{b}^\alpha)$$

induced by $\| - \|_r$ on $\mathcal{D}^r(G, R)$ and the norm on S is bounded-equivalent to the norm induced by $\| - \|_r$ on $\mathcal{D}^r(G, S)$, and so we obtain isomorphic completions, which gives the desired statement. \square

Our next goal is to prove that $\mathcal{D}(G, R)$ is independent, as a topological R -module, of the choice of norm on R . Recall that we only consider norms for which there exists a multiplicative pseudouniformizer, and such that the natural map $\mathbb{Z}_p \rightarrow R$ is norm-decreasing.

Proposition 3.2.6. *$\mathcal{D}(G, R)$ is independent of the choice of norm on R .*

Proof. Let $| - |$ and $| - |'$ be two equivalent such norms on R ; for $r \in [1/p, 1)$ we get the corresponding r -norms $\| - \|_r$ and $\| - \|'_r$ on $\mathcal{D}(G, R)$. Let $\mu = \sum_\alpha d_\alpha \mathbf{b}^\alpha \in \mathcal{D}(G, R)$ be an arbitrary element and use Lemma 2.1.6 to find constants $C_1, C_2, s_1, s_2 > 0$ such that

$$C_1 |a|^{s_1} \leq |a|' \leq C_2 |a|^{s_2}$$

for all $a \in R$. Then

$$C_1 |d_\alpha|^{s_1} r^{|\alpha|} \leq |d_\alpha|'_r r^{|\alpha|} \leq C_2 |d_\alpha|^{s_2} r^{|\alpha|}$$

for all α and hence

$$C_1 \|\mu\|_{r^{1/s_1}}^{s_1} \leq \|\mu\|'_r \leq C_2 \|\mu\|_{r^{1/s_2}}^{s_2}$$

for all r such that $r, r^{1/s_1}, r^{1/s_2} \in [1/p, 1)$. It follows that the families $(\| - \|_r)$ and $(\| - \|'_r)$ define the same topology on $\mathcal{D}(G, R)$, and hence the same completion, as required. \square

We now introduce a variant of the $\mathcal{D}^r(G, R)$, which, when R is a Banach \mathbb{Q}_p -algebra, recovers the analytic distribution algebra (with fixed radius of analyticity). Although we do not use this variant in our construction of eigenvarieties, it is used in Section 5 to construct Galois representations.

Let $r > s \geq 1/p$. Let $\mathcal{D}^{r,\circ}(G, R) = \{\sum d_\alpha \mathbf{b}^\alpha \mid |d_\alpha| r^{|\alpha|} \leq 1\}$ denote the unit ball of $\mathcal{D}^r(G, R)$. We define $\mathcal{D}^{<r,\circ}(G, R)$ to be the closure of $\mathcal{D}^{r,\circ}(G, R)$ in $\mathcal{D}^s(G, R)$. Note that $\mathcal{D}^{<r,\circ}(G, R)$ is an R_0 -module, a priori depending on s , which carries two natural topologies (the ϖ -adic topology and the subspace topology coming from $\mathcal{D}^s(G, R)$).

Proposition 3.2.7. *We have*

$$\mathcal{D}^{<r,\circ}(G, R) = \left\{ \sum_{\alpha} d_{\alpha} \mathbf{b}^{\alpha} \in \mathcal{D}^s(G, R) \mid |d_{\alpha}| r^{|\alpha|} \leq 1 \right\}$$

as a subset of $\mathcal{D}^s(G, R)$. Thus $\mathcal{D}^{<r,\circ}(G, R)$ is independent of s . The subspace topology corresponds to the weak topology with respect to the family of maps $(\mu \mapsto d_{\alpha}(\mu))_{\alpha}$ on the right-hand side (and is therefore also independent of s). The ϖ -adic topology is induced by the norm

$$\left\| \sum d_{\alpha} \mathbf{b}^{\alpha} \right\|_r = \sup_{\alpha} |d_{\alpha}|' r^{|\alpha|}$$

and is separated and complete.

Proof. Let $W = \{\sum d_{\alpha} \mathbf{b}^{\alpha} \in \mathcal{D}^s(G, R) \mid |d_{\alpha}| r^{|\alpha|} \leq 1\}$. Since the maps $\mu \mapsto d_{\alpha}(\mu)$ are continuous we see that W is closed. On other hand, any finite truncation of an element in W is in $\mathcal{D}^{r,\circ}(G, R)$, so $\mathcal{D}^{r,\circ}(G, R)$ is dense in W . It follows that $W = \mathcal{D}^{<r,\circ}(G, R)$. The subspace topology is given by the norm $\| - \|_s$, and one checks easily that this agrees with the weak topology in the statement of the proposition. The final statement is similarly easy to check; we leave it to the reader. \square

We then set $\mathcal{D}^{<r}(G, R) = \mathcal{D}^{<r,\circ}(G, R)[1/\varpi]$; this is naturally a Banach R -module which embeds into $\mathcal{D}^s(G, R)$ for all $s \in [1/p, r)$. The $(\mathcal{D}^{<r}(G, R))_{r > 1/p}$ form an inverse system and the natural map $\mathcal{D}^r(G, R) \rightarrow \mathcal{D}^s(G, R)$ factors over $\mathcal{D}^{<r}(G, R)$, so we therefore have

$$\mathcal{D}(G, R) = \varprojlim_r \mathcal{D}^{<r}(G, R)$$

as well. Recall the potential ON-basis $(\varpi^{-n(r,\varpi,\alpha)} \mathbf{b}^{\alpha})_{\alpha}$ of $\mathcal{D}^r(G, R)$; we have

$$\mathcal{D}^{<r}(G, R) = \left\{ \sum_{\alpha} d_{\alpha} \mathbf{b}^{\alpha} \mid |d_{\alpha}| r^{|\alpha|} \text{ bounded} \right\} = \left(\prod_{\alpha} R_0 \cdot \varpi^{-n(r,\varpi,\alpha)} \mathbf{b}^{\alpha} \right) \left[\frac{1}{\varpi} \right].$$

We remark that if $(\varpi^{-n(r,\varpi,\alpha)} \mathbf{b}^{\alpha})_{\alpha}$ is an ON-basis, we have $\mathcal{D}^{<r,\circ}(G, R) = \prod_{\alpha} R_0 \cdot \varpi^{-n(r,\varpi,\alpha)} \mathbf{b}^{\alpha}$ and the weak topology on the left-hand side is equal to the product topology on the right-hand side. Next, we define some function modules in a similar fashion. We put

$$\mathcal{C}^{<r,\circ}(G, R) = \{f \in \mathcal{C}(G, R) \mid |\mu(f)| \leq 1 \text{ for all } \mu \in \mathcal{D}(G, R) \cap \mathcal{D}^{r,\circ}(G, R)\}$$

and set $\mathcal{C}^{<r}(G, R) = \mathcal{C}^{<r,0}(G, R)[1/\varpi] \subseteq \mathcal{C}(G, R)$. We note that $f = \sum_{\alpha} c_{\alpha} E_{\alpha} \in \mathcal{C}^{<r}(G, R)$ if and only if $\varpi^{-n(r,\varpi,\alpha)} \mathbf{b}^{\alpha}(f) = c_{\alpha} \varpi^{-n(r,\varpi,\alpha)}$ is bounded as $\alpha \rightarrow \infty$; from this one sees that

$$\mathcal{C}^{<r}(G, R) = \left(\prod_{\alpha} R_0 \cdot \varpi^{n(r,\varpi,\alpha)} E_{\alpha} \right) \left[\frac{1}{\varpi} \right]$$

and that it is the dual space of $\mathcal{D}^r(G, R)$, and that the dual norm is given by $\|\sum_{\alpha} c_{\alpha} E_{\alpha}\|_r = \sup_{\alpha} |c_{\alpha}| r^{-|\alpha|}$. When $r > s \geq 1/p$, we let $\mathcal{C}^r(G, R)$ denote the closure of $\mathcal{C}^{<s}(G, R)$ inside $\mathcal{C}^{<r}(G, R)$. Tracing through the definitions we get an explicit description

$$\mathcal{C}^r(G, R) = \left\{ \sum_{\alpha} c_{\alpha} E_{\alpha} \mid |c_{\alpha}| r^{-|\alpha|} \rightarrow 0 \right\} = \widehat{\bigoplus_{\alpha} R \cdot \varpi^{n(r,\varpi,\alpha)} E_{\alpha}}$$

and see that the dual space of $\mathcal{C}^r(G, R)$ is $\mathcal{D}^{<r}(G, R)$.

Remark 3.2.8. The reader should compare our description of $\mathcal{C}^r(G, R)$ with the construction of [Liu et al. 2017, Section 5.4]. Here, the authors give a definition of a “modified” space of continuous functions on \mathbb{Z}_p with values in $\mathbb{Z}_p[[T, pT^{-1}]]$ which is morally (apart from the slight difference in coefficients and the fact that we have only defined $\mathcal{C}^r(G, R)$ for $r > 1/p$) the space $\mathcal{C}^{1/p}(\mathbb{Z}_p, R)$, where R is the Tate ring obtained as the rational localization

$$\mathbb{Z}_p[[T]] \left\langle \frac{p}{T} \right\rangle \left[\frac{1}{T} \right]$$

of $\mathbb{Z}_p[[T]]$. We give R the norm with unit ball $\mathbb{Z}_p[[T]] \langle p/T \rangle$ and $|T| = 1/p$, defined as in Remark 2.1.3. The definition which appears in [loc. cit.] is therefore a useful motivation for the general constructions of this article, although we will not use the modules $\mathcal{C}^r(G, R)$ in this article. We will see in Theorem 6.3.4 that one may use the module $\mathcal{D}^{1/p}(\mathbb{Z}_p, R)$ (which is defined) to prove the main results of [loc. cit.].

To finish this section we discuss the relationship between our constructions and the spaces of locally analytic functions or distributions when R is a Banach \mathbb{Q}_p -algebra. We may and do assume that \mathbb{Q}_p is isometrically embedded into R and that $|p| = 1/p$. Recall that the atlas on G induced from our choice of a topological basis identifies G^{p^n} with $(p^n \mathbb{Z}_p)^{\dim G}$. Amice’s theorem [Colmez 2010, Théorème I.4.7] tells us that the space of n -analytic functions on G with respect to this atlas is explicitly given as

$$\mathcal{C}^{n\text{-an}}(G, \mathbb{Q}_p) = \widehat{\bigoplus_{\alpha} \mathbb{Q}_p \cdot k_{\alpha} E_{\alpha}},$$

where $k_{\alpha} := \lfloor p^{-n} \alpha_1 \rfloor! \cdots \lfloor p^{-n} \alpha_{\dim G} \rfloor!$. The space of n -analytic R -valued functions on G is then

$$\mathcal{C}^{n\text{-an}}(G, R) = \widehat{\bigoplus_{\alpha} R \cdot k_{\alpha} E_{\alpha}} \subseteq \mathcal{C}(G, R).$$

It is well known that $|k_{\alpha}| \sim r_n^{|\alpha|}$ with $r_n = p^{-1/p^n(p-1)}$, and it follows that $\mathcal{C}^{n\text{-an}}(G, R) = \mathcal{C}^{r_n}(G, R)$ as \mathbb{Q}_p -Banach spaces. Since $r_n \rightarrow 1$ from below as $n \rightarrow \infty$ it follows that $\mathcal{C}(G, R)$ is the space of locally analytic R -valued functions on G , with its usual locally convex topology. Dually, $\mathcal{D}(G, R)$ is then the

space of locally analytic R -valued distributions with its usual locally convex topology. We sum up this discussion in a proposition.

Proposition 3.2.9. *When R is Banach \mathbb{Q}_p -algebra, $\mathcal{C}(G, R)$ is canonically the space of locally analytic R -valued functions on G and $\mathcal{D}(G, R)$ is dually the space of locally analytic R -valued distributions on G .*

3.3. Ash–Stevens distribution modules for Banach–Tate \mathbb{Z}_p -algebras. While the definitions in this subsection will be of a local nature, let us nevertheless start by introducing the global setup that we will need to define eigenvarieties. Let F be a number field. We put $\mathbf{G} = \text{Res}_{\mathbb{Q}}^F \mathbf{H}$, where \mathbf{H} is a connected reductive group over F split at all places $v \mid p$. \mathbf{G} is then a connected reductive group over \mathbb{Q} . When $v \mid p$ we write $\mathbf{H}_{\mathcal{O}_{F_v}}$ for a split model of \mathbf{H}_{F_v} over \mathcal{O}_{F_v} and choose a Borel subgroup \mathbf{B}_v (with unipotent radical \mathbf{N}_v) and a maximal torus $\mathbf{T}_v \subseteq \mathbf{B}_v$ of $\mathbf{H}_{\mathcal{O}_{F_v}}$. Set

$$\mathbf{G}_{\mathbb{Z}_p} = \prod_{v \mid p} \text{Res}_{\mathbb{Z}_p}^{\mathcal{O}_{F_v}} \mathbf{H}_{\mathcal{O}_{F_v}}, \quad \mathbf{B} = \prod_{v \mid p} \text{Res}_{\mathbb{Z}_p}^{\mathcal{O}_{F_v}} \mathbf{B}_v, \quad \mathbf{T} = \prod_{v \mid p} \text{Res}_{\mathbb{Z}_p}^{\mathcal{O}_{F_v}} \mathbf{T}_v$$

and also put $\mathbf{N} = \prod_{v \mid p} \text{Res}_{\mathbb{Z}_p}^{\mathcal{O}_{F_v}} \mathbf{N}_v$. We use overlines to denote opposite groups; e.g., $\bar{\mathbf{B}} = \prod_{v \mid p} \text{Res}_{\mathbb{Z}_p}^{\mathcal{O}_{F_v}} \bar{\mathbf{B}}_v$, where the $\bar{\mathbf{B}}_v$ are the opposite Borels of \mathbf{B}_v .

We will also need notation for various subgroups of $G := \mathbf{G}(\mathbb{Q}_p)$. Set $G_0 = \mathbf{G}_{\mathbb{Z}_p}(\mathbb{Z}_p)$, $B_0 = \mathbf{B}(\mathbb{Z}_p)$, $T_0 = \mathbf{T}(\mathbb{Z}_p)$, and $N_0 = \mathbf{N}(\mathbb{Z}_p)$. We let I denote the Iwahori subgroup of G defined as the preimage of $\mathbf{B}(\mathbb{F}_p)$ under the reduction map $G_0 \rightarrow \mathbf{G}_{\mathbb{Z}_p}(\mathbb{F}_p)$, and we let $K_s = \text{Ker}(G_0 \rightarrow \mathbf{G}_{\mathbb{Z}_p}(\mathbb{Z}/p^s))$ be the s -th principal congruence subgroup of G_0 for $s \geq 1$. Again for $s \geq 1$, set

$$\begin{aligned} T_s &= \text{Ker}(T_0 \rightarrow \mathbf{T}(\mathbb{Z}/p^s\mathbb{Z})), \\ N_s &= \text{Ker}(N_0 \rightarrow \mathbf{N}(\mathbb{Z}/p^s\mathbb{Z})), \\ \bar{N}_s &= \text{Ker}(\bar{\mathbf{N}}(\mathbb{Z}_p) \rightarrow \bar{\mathbf{N}}(\mathbb{Z}/p^s\mathbb{Z})). \end{aligned}$$

Set $B_s = T_s N_s$ for $s \geq 0$. We have Iwahori decompositions $I = \bar{N}_1 T_0 N_0$ and $K_s = \bar{N}_s T_s N_s$. Next, choose a splitting $T := \mathbf{T}(\mathbb{Q}_p) \rightarrow T_0$ of the inclusion $T_0 \subseteq T$ and put $\Sigma = \text{Ker}(T \rightarrow T_0)$. We set

$$\begin{aligned} \Sigma^+ &= \{t \in \Sigma \mid t \bar{N}_1 t^{-1} \subseteq \bar{N}_1\}, \\ \Sigma^{\text{cpt}} &= \{t \in \Sigma \mid t \bar{N}_1 t^{-1} \subseteq \bar{N}_2\}. \end{aligned}$$

We may then define $\Delta_p = I \Sigma^+ I$; this is a monoid and (Δ_p, I) is a Hecke pair (which means that I and $\delta I \delta^{-1}$ are commensurable for all $\delta \in \Delta_p$). The corresponding Hecke algebra (defined over \mathbb{Z}_p) will be denoted by $\mathbb{T}(\Delta_p, I)$.

With these preparations let us now move on to the definition of analytic and locally analytic distribution modules for general Banach–Tate \mathbb{Z}_p -algebras R . When R is a Banach \mathbb{Q}_p -algebra, these were defined in [Ash and Stevens 2008] using (locally) analytic functions on I . Recasting this in terms of the norms of the previous section, we are able to extend the definition to all Banach–Tate \mathbb{Z}_p -algebras.

Let $\kappa : T_0 \rightarrow R^\times$ be a continuous character. We will put some restrictions on the choice of norm on R , according to the following lemma. We set $\epsilon = 1$ if $p \neq 2$ and $\epsilon = 2$ if $p = 2$, and put $q = p^\epsilon$.

Lemma 3.3.1. *Keep the above notation and assume that R is Noetherian. Write $|\cdot|$ for the given norm on R and let ϖ be a multiplicative pseudouniformizer. Then there exists a Banach–Tate \mathbb{Z}_p -algebra norm $|\cdot|'$ on R , bounded-equivalent to $|\cdot|^s$ for some $s > 0$, such that $|\kappa(t)|' \leq 1$ for all $t \in T_0$, $|\kappa(t) - 1|' < 1$ for all $t \in T_\epsilon$, and ϖ is a multiplicative pseudouniformizer for $|\cdot|'$.*

Proof. The proof is similar to that of Lemma 3.1.1. Let t_1, \dots, t_a be a set of topological generators of T_0 and let t_{a+1}, \dots, t_b be a set of topological generators for T_ϵ ; it suffices to find $|\cdot|'$ such that $|\kappa(t_i)|' \leq 1$ for $i = 1, \dots, a$, $|\kappa(t_i) - 1|' < 1$ for $i = a + 1, \dots, b$, and $|p|' \leq p^{-1}$, which is bounded-equivalent to $|\cdot|^s$ for some $s > 0$.

Since T_0 is compact, $\kappa(T_0)$ is bounded, and hence t_i is powerbounded for all i . Moreover, T_ϵ is pro- p and $R^\circ/R^{\circ\circ}$ is a reduced discrete ring of characteristic p , so the continuous homomorphism $T_\epsilon \rightarrow (R^\circ/R^{\circ\circ})^\times$ induced from κ is trivial. We conclude that $\kappa(T_\epsilon) \subseteq 1 + R^{\circ\circ}$ and hence $\kappa(t_i) - 1 \in R^{\circ\circ}$ for $i = a + 1, \dots, b$. We may now choose m such that $|\kappa(t_i) - 1|_{\text{sp}} < |\varpi|^{2/m}$ for $i = a + 1, \dots, b$ and $|p|_{\text{sp}} < |\varpi|^{2/m}$. Arguing with $S = R[\varpi^{1/m}]$ as in Lemma 3.1.1 we may construct a norm $|\cdot|_2$ on R for which ϖ is a multiplicative pseudouniformizer with $|\varpi|_2 = |\varpi|$, $|\kappa(t_i)|_2 \leq 1$ for $i = 1, \dots, a$, $|\kappa(t_i) - 1|_2 < 1$ for $i = a + 1, \dots, b$, and $|p|_2 < 1$. Setting $|\cdot|' := |\cdot|_2^s$ for s sufficiently large then gives the desired norm. \square

Definition 3.3.2. Let R be a Banach–Tate \mathbb{Z}_p -algebra and $\kappa : T_0 \rightarrow R^\times$ a continuous character. We will say that the norm of R is *adapted to κ* if $\kappa(T_0) \subseteq R_0$ and $|\kappa(t) - 1| < 1$ for all $t \in T_\epsilon$. Note that then $|\kappa(t)| = 1$ for all $t \in T_0$ and there exists an $r < 1$ such that $|\kappa(t) - 1| \leq r$ for all $t \in T_\epsilon$.

For the rest of the subsection we consider a Banach–Tate \mathbb{Z}_p -algebra R and character $\kappa : T_0 \rightarrow R$ such that the norm on R is adapted to κ . We extend κ to a character of B_0 by making it trivial on N_0 . We define $\mathcal{A}_\kappa \subseteq \mathcal{C}(I, R)$ to be the subset of functions such that $f(gb) = \kappa(b)f(g)$ for all $g \in I$ and $b \in B_0$. \mathcal{A}_κ is naturally a Banach R -module and carries a continuous right action of I by left translation. Restricting a function from I to \bar{N}_1 gives a topological isomorphism $\mathcal{A}_\kappa \cong \mathcal{C}(\bar{N}_1, R)$. By definition, Σ^+ acts on the left on \bar{N}_1 by conjugation, and via the previous isomorphism this induces a right action of Σ^+ on \mathcal{A}_κ . These actions fit together into a right action of Δ_p on \mathcal{A}_κ . We let \mathcal{D}_κ denote the dual $\text{Hom}_{R, \text{cts}}(\mathcal{A}_\kappa, R)$, equipped with the dual left Δ_p -action. Since \mathcal{A}_κ is the set of B_0 -invariants of $\mathcal{C}(I, R)$ with respect to the action $(f.b)(g) = \kappa(b)^{-1}f(gb)$ ($b \in B_0$, $g \in I$), \mathcal{D}_κ is the Hausdorff B_0 -coinvariants of $\mathcal{D}(I, R)$ with respect to the dual (right) action. We record a more precise statement for future use:

Proposition 3.3.3. *The natural surjection $\mathcal{D}(I, R) \rightarrow \mathcal{D}_\kappa$ is equivariant for the natural left I -actions on both sides. Identifying \mathcal{D}_κ with $\mathcal{D}(\bar{N}_1, R)$, the map is given by $\delta_{\bar{n}b} \mapsto \kappa(b)\delta_{\bar{n}}$.*

Proof. The inverse to the restriction map $\mathcal{A}_\kappa \rightarrow \mathcal{C}(\bar{N}_1, R)$ is given by $f \mapsto (\bar{n}b \mapsto \kappa(b)f(\bar{n}))$ (here and above $\bar{n} \in \bar{N}_1$ and $b \in B_0$). From this one sees directly that the dual map sends $\delta_{\bar{n}b}$ to $\kappa(b)\delta_{\bar{n}}$. That this characterizes the maps follows from R -linearity and continuity for the weak-star topology. \square

To apply the results from the previous subsection we will need to know that some groups are uniform. The following result is presumably well known to experts but we have been unable to find a suitable

reference. The proof we give is due to Konstantin Ardakov, and we thank him for allowing us to include it here (any errors are due to the authors).

Proposition 3.3.4. *K_s and \bar{N}_s are uniform for $s \geq \epsilon$.*

Proof. By construction the groups are products $K_s = \prod_{v|p} K_{s,v}$, and similarly for $\bar{N}_{s,v}$, in a natural way, so it suffices to prove that each $K_{s,v}$ and each $\bar{N}_{s,v}$ is uniform. Fix $v|p$ and let $s \geq \epsilon$. First assume that $\mathbf{H}_v = \mathrm{GL}_n/\mathcal{O}_{F_v}$. Then we have the usual matrix logarithm $\log : K_{s,v} \rightarrow p^s M_n(\mathcal{O}_{F_v})$ and exponential $\exp : p^s M_n(\mathcal{O}_{F_v}) \rightarrow K_{s,v}$. They converge and are inverse to each other (by our assumption on s) and the Lie algebra $p^s M_n(\mathcal{O}_{F_v})$ is easily seen to be powerful by assumption, so by the definition of the correspondence between powerful Lie algebras and uniform pro- p groups [Dixon et al. 1999, Theorem 9.10] via the Campbell–Hausdorff series we see that $K_{s,v}$ is the uniform group corresponding to $p^s M_n(\mathcal{O}_{F_v})$. To get the result for $\bar{N}_{s,v}$ we may by conjugation assume that \bar{N}_v is the group of lower triangular unipotent matrices; the corresponding Lie algebra is that of lower triangular nilpotent matrices and we then argue similarly.

Now let \mathbf{H}_v be arbitrary and choose a closed immersion $\mathbf{H}_v \hookrightarrow \mathrm{GL}_n/\mathcal{O}_{F_v}$ for some n . We thereby identify \mathbf{H}_v with a closed subgroup of $\mathrm{GL}_n/\mathcal{O}_{F_v}$. Writing \mathbf{B}'_v for the upper triangular Borel of $\mathrm{GL}_n/\mathcal{O}_{F_v}$, \mathbf{N}'_v for its unipotent radical, $\bar{\mathbf{N}}'_v$ for the opposite of \mathbf{N}'_v and \mathbf{T}'_v for the diagonal torus, we may assume, after conjugating if necessary, that $\mathbf{T}_v \subseteq \mathbf{T}'_v$ etc. We write $K'_{s,v}$ etc. for the corresponding principal congruence subgroups. With this setup, we now give the rest of the proof for the $K_{s,v}$ only; the proof for $\bar{N}_{s,v}$ proceeds in the same way. Note that $K_{s,v} = \mathbf{H}_v(\mathcal{O}_{F_v}) \cap K'_{s,v}$. By [Dixon et al. 1999, Theorem 4.5] we see that $K'_{s,v}$ is torsion-free and that it suffices to prove that $K_{s,v}$ is powerful, i.e., that $[K_{s,v}, K_{s,v}] \subseteq K_{s,v}^q$, where $[K_{s,v}, K_{s,v}]$ is the derived subgroup of $K_{s,v}$ and $K_{s,v}^q$ is the subgroup generated by the q -th powers of elements in $K_{s,v}$ (any compact p -adic analytic group is topologically finitely generated). We remark that it is easy to see that $[K'_{s,v}, K'_{s,v}] \subseteq K'_{s+\epsilon,v} = (K'_{s,v})^q$. Using this and $K_{s,v} = \mathbf{H}_v(\mathcal{O}_{F_v}) \cap K'_{s,v}$, we see that $[K_{s,v}, K_{s,v}] \subseteq K'_{s+\epsilon,v} \cap \mathbf{H}_v(\mathcal{O}_{F_v}) = K_{s+\epsilon,v}$ and $K_{s,v}^q \subseteq K_{s+\epsilon,v}$.

It remains to prove that $K_{s+\epsilon,v} \subseteq K_{s,v}^q$. We have $T_{s+\epsilon,v} = T_{s,v}^q$ using the logarithm and exponential (\mathbf{T}_v is a split torus). We have an isomorphism of \mathcal{O}_{F_v} -schemes $\prod_{\alpha} x_{\alpha} : \prod_{\alpha} \mathbb{G}_a \xrightarrow{\sim} N_v$, where α ranges through the roots of \mathbf{H}_v whose root subgroups are contained in \mathbf{N}_v , and x_{α} is a corresponding root homomorphism. Under this isomorphism $N_{s,v}$ corresponds to $\prod_{\alpha} p^s \mathcal{O}_{F_v}$, and by standard properties of x_{α} one has $x_{\alpha}(q \cdot (p^s a)) = x_{\alpha}(p^s a)^q$ for any $a \in \mathcal{O}_{F_v}$. It follows that $N_{s+\epsilon,v} = N_{s,v}^q$. Similarly $\bar{N}_{s+\epsilon,v} = \bar{N}_{s,v}^q$. By the Iwahori decomposition we then see that $K_{s+\epsilon,v} \subseteq K_{s,v}^q$ as desired. \square

Remark 3.3.5. Note that the argument in the final paragraph of the above proof also implies that, for arbitrary p and $s \geq 1$, we have $\bar{N}_{s+1,v} = \bar{N}_{s,v}^p$.

Definition 3.3.6. Let $r \in [1/p, 1)$:

- (1) We define a norm $\|\cdot\|_r^{\mathrm{sub}}$ on \mathcal{D}_k by transporting the norm $\|\cdot\|_{\bar{N}_{\epsilon,r}}$ (defined before Proposition 3.1.10) on $\mathcal{D}(\bar{N}_1, R)$ to \mathcal{D}_k via the isomorphism $\mathcal{D}(\bar{N}_1, R) \cong \mathcal{D}_k$ obtained by restriction of functions from I to \bar{N}_1 .

- (2) We define a norm $\| - \|_r^{\text{quot}}$ on \mathcal{D}_κ as the quotient norm induced from $\| - \|_{K_\epsilon, r}$ (defined before Proposition 3.1.10) via the natural surjection $\mathcal{D}(I, R) \rightarrow \mathcal{D}_\kappa$.

Note that $\| - \|_r^{\text{quot}}$ is I -invariant; this follows from I -equivariance of the surjection and from Proposition 3.1.10. The following is the key result of this subsection:

Proposition 3.3.7. *Suppose that $|\kappa(t) - 1| \leq r$ for all $t \in T_\epsilon$. Then $\| - \|_r^{\text{quot}} = \| - \|_r^{\text{sub}}$ on \mathcal{D}_κ .*

Proof. Let us first assume that $p \neq 2$. First we claim that $\| - \|_r^{\text{quot}}$ is equal to the quotient norm on \mathcal{D}_κ coming from the surjection $\mathcal{D}(K_1, R) \rightarrow \mathcal{D}_\kappa$ and the norm $\| - \|_r$ on the source. To see this, pick a set $(b_i)_i$ of coset representatives of I/K_1 lying in B_0 and define a map $\pi_\kappa : \mathcal{D}(I, R) = \bigoplus_i \delta_{b_i} \mathcal{D}(K_1, R) \rightarrow \mathcal{D}(K_1, R)$ by

$$(\delta_{b_i} \mu_i)_i \mapsto \sum_i \kappa(b_i) \mu_i.$$

We remark that composing this map with the natural surjection $\mathcal{D}(K_1, R) \rightarrow \mathcal{D}_\kappa$ gives the natural surjection $\mathcal{D}(I, R) \rightarrow \mathcal{D}_\kappa$ (this follows from the explicit formula in Proposition 3.3.3). To prove the claim, it then suffices to prove that $\| - \|_r$ is the quotient norm of $\| - \|_{K_1, r}$ via π_κ . Write $\| - \|'_r$ for this quotient norm. For simplicity assume that 1 is one of the coset representatives b_i . Then our map is a section of the inclusion $\mathcal{D}(K_1, R) \subseteq \mathcal{D}(I, R)$. It is then clear that $\| - \|'_r \leq (\| - \|_{K_1, r})|_{\mathcal{D}(K_1, R)} = \| - \|_r$. Conversely, if $\mu = \sum_i \kappa(b_i) \mu_i \in \mathcal{D}(K_1, R)$ is the image of $\sum_i \delta_{b_i} \mu_i \in \mathcal{D}(I, R)$, then

$$\|\mu\|_r \leq \sup_i \|\kappa(b_i) \mu_i\|_r \leq \sup_i \|\mu_i\|_r = \left\| \sum_i \delta_{b_i} \mu_i \right\|_{K_1, r}.$$

Taking the infimum over such presentations we obtain $\| - \|_r \leq \| - \|'_r$, and hence equality.

Next, let $\bar{n}_1, \dots, \bar{n}_k$ (resp. n_1, \dots, n_k) be a minimal set of topological generators of \bar{N}_1 (resp. N_1), and let t_1, \dots, t_l be a set of topological generators of T_1 . Put $\mathbf{n}^\alpha = \prod_i (\delta_{n_i} - 1)^{\alpha_i}$ and similarly for T_1 and \bar{N}_1 . By Proposition 3.3.3, the map $\mathcal{D}(K_1, R) \rightarrow \mathcal{D}_\kappa$ is then given by

$$\sum_{\alpha, \beta, \gamma} d_{\alpha, \beta, \gamma} \bar{\mathbf{n}}^\alpha \mathbf{t}^\beta \mathbf{n}^\gamma \mapsto \sum_{\alpha} \left(\sum_{\beta} d_{\alpha, \beta, \underline{0}} \prod_i (\kappa(t_i) - 1)^{\beta_i} \right) \bar{\mathbf{n}}^\alpha.$$

We then make a computation similar to the one in the proof of the claim above. First, it's clear that $\| - \|_r^{\text{quot}} \leq \| - \|_r^{\text{sub}}$ since restricting $\| - \|_r$ on $\mathcal{D}(K_1, R)$ to $\mathcal{D}(\bar{N}_1, R)$ gives (the intrinsically defined) $\| - \|_r$, and the composition $\mathcal{D}(\bar{N}_1, R) \rightarrow \mathcal{D}(K_1, R) \rightarrow \mathcal{D}_\kappa \cong \mathcal{D}(\bar{N}_1, R)$ is the identity. Second, if

$$\sum_{\alpha} e_{\alpha} \bar{\mathbf{n}}^\alpha = \sum_{\alpha} \left(\sum_{\beta} d_{\alpha, \beta, \underline{0}} \prod_i (\kappa(t_i) - 1)^{\beta_i} \right) \bar{\mathbf{n}}^\alpha,$$

then

$$\left\| \sum_{\alpha} e_{\alpha} \bar{\mathbf{n}}^\alpha \right\|_r \leq \sup_{\alpha, \beta} \left(|d_{\alpha, \beta, \underline{0}}| \prod_i |\kappa(t_i) - 1|^{\beta_i} r^{|\alpha|} \right) \leq \sup_{\alpha, \beta} |d_{\alpha, \beta, \underline{0}}| r^{|\alpha| + |\beta|} \leq \left\| \sum_{\alpha, \beta, \gamma} d_{\alpha, \beta, \gamma} \bar{\mathbf{n}}^\alpha \mathbf{t}^\beta \mathbf{n}^\gamma \right\|_r,$$

where we have used the assumption $|\kappa(t_i) - 1| \leq r$ for all i to obtain the second inequality. Hence, taking the infimum over such presentations, we see that $\| - \|_r^{\text{sub}} \leq \| - \|_r^{\text{quot}}$ and equality follows.

The case $p = 2$ is similar. We identify \mathcal{D}_κ with $\mathcal{D}(\bar{N}_1, R)$ and consider the subspace $\mathcal{D}(\bar{N}_2, R)$. It carries the norm $\| - \|_r$ and also receives a quotient norm from the norm $\| - \|_r$ on $\mathcal{D}(K_2, R)$ via the surjection $\varphi : \mathcal{D}(K_2, R) \rightarrow \mathcal{D}(\bar{N}_2, R)$. These two norms are equal by the same type of argument as in the second part above. We then equip $\mathcal{D}(I, R)$ with the norm $\| - \|_{K_2, r}$ and $\mathcal{D}(\bar{N}_1, R)$ with the norm $\| - \|_r^{\text{sub}}$. Pick coset representatives $(\bar{n}_i)_i$ of \bar{N}_1/\bar{N}_2 and $(b_j)_j$ of B_0/B_2 , both containing 1. We may then write the map $\mathcal{D}(I, R) \rightarrow \mathcal{D}(\bar{N}_1, R)$ as

$$\bigoplus_{i,j} \delta_{\bar{n}_i b_j} \mathcal{D}(K_2, R) \rightarrow \bigoplus_i \delta_{\bar{n}_i} \mathcal{D}(\bar{N}_2, R), \quad (\delta_{\bar{n}_i b_j} \mu_{ij})_{i,j} \mapsto \left(\delta_{\bar{n}_i} \sum_j \kappa(b_j) \varphi(\mu_{ij}) \right)_i.$$

By a computation similar to that in the first part of the proof in the case $p \neq 2$ (using additionally the equality of the two norms on $\mathcal{D}(\bar{N}_2, R)$ asserted above) the norm $\| - \|_r^{\text{quot}}$ agrees with the norm $\| - \|_r^{\text{sub}}$, as desired. \square

Remark 3.3.8. It might happen that \bar{N}_1 is uniform when $p = 2$ (e.g., when $G = \text{Res}_{\mathbb{Q}}^F \text{GL}_{2/F}$). In this case, the norm $\| - \|_r^{\text{sub}}$ is bounded-equivalent to $\| - \|_{r^{1/2}}$ on $\mathcal{D}(\bar{N}_1, R)$.

Definition 3.3.9. Write r_κ for the minimal $r \in [1/p, 1)$ such that $|\kappa(t) - 1| \leq r$ for all $t \in T_\epsilon$. When $r \geq r_\kappa$, we write $\| - \|_r$ for the norm $\| - \|_r^{\text{sub}} = \| - \|_r^{\text{quot}}$ on \mathcal{D}_κ (we will never consider these norms when $r < r_\kappa$).

Let $r \geq r_\kappa$. We define \mathcal{D}_κ^r to be the completion of \mathcal{D}_κ with respect to $\| - \|_r$, and let $\mathcal{D}_\kappa = \varprojlim_r \mathcal{D}_\kappa^r$.

\mathcal{D}_κ^r is a Banach R -module with respect to its induced norm, and carries a left I -action (since I acts on \mathcal{D}_κ by $\| - \|_r$ -isometries, the action extends to the completion). When R is a Banach \mathbb{Q}_p -algebra, it follows from Proposition 3.2.9 that \mathcal{D}_κ is the locally analytic distribution module used in [Ash and Stevens 2008; Hansen 2017]. We will also need to extend the action of Σ^+ to these modules, and prove that elements of Σ^{cpt} give compact operators. For this, it is convenient to use the definition of $\| - \|_r$ as $\| - \|_r^{\text{sub}}$. We have a natural identification $\mathcal{D}_\kappa^r \cong \mathcal{D}^r(\bar{N}_1, R)$ when $p \neq 2$; when $p = 2$ we have $\mathcal{D}_\kappa^r \cong \bigoplus_{\bar{n}_i \in \bar{N}_1/\bar{N}_2} \delta_{\bar{n}_i} \mathcal{D}^r(\bar{N}_2, R)$.

Corollary 3.3.10. *If $t \in \Sigma^+$, then the action of t on \mathcal{D}_κ is norm-decreasing with respect to $\| - \|_r$ for any $r \geq r_\kappa$, and hence the action of t extends to \mathcal{D}_κ^r . If $t \in \Sigma^{\text{cpt}}$, then t acts compactly on \mathcal{D}_κ^r . Moreover, in this case the action of t is given by the composition of a norm-decreasing map*

$$\mathcal{D}_\kappa^r \rightarrow \mathcal{D}_\kappa^{r^{1/p}}$$

with the compact (norm-decreasing) inclusion

$$\mathcal{D}_\kappa^{r^{1/p}} \hookrightarrow \mathcal{D}_\kappa^r.$$

Proof. We use the identification $\mathcal{D}_\kappa \cong \mathcal{D}(\bar{N}_1, R)$, with regards to which t acts by the map induced from the homomorphism $\bar{N}_1 \rightarrow \bar{N}_1$ given by $\bar{n} \mapsto t\bar{n}t^{-1}$. First assume $p \neq 2$. The first assertion follows directly from Lemma 3.2.4. The second assertion follows from the third, so it remains to prove the third assertion. If $t \in \Sigma^{\text{cpt}}$ we have $t\bar{N}_1t^{-1} \subseteq \bar{N}_2 = (\bar{N}_1)^p$ by the definition of Σ^{cpt} and Remark 3.3.5, so the third assertion follows from Lemmas 3.2.3 and 3.2.4.

Now assume $p = 2$ and let $(\bar{n}_i)_i$ be a set of coset representatives of \bar{N}_1/\bar{N}_2 . We have $t\bar{N}_2t^{-1} \subseteq \bar{N}_2$ and $\bar{n} \mapsto t\bar{n}t^{-1}$ induces a map $\bar{N}_1/\bar{N}_2 \rightarrow \bar{N}_1/\bar{N}_2$; moreover if $t \in \Sigma^{\text{cpt}}$ then $t\bar{N}_2t^{-1} \subseteq \bar{N}_3$, by Remark 3.3.5. Writing $\mathcal{D}(\bar{N}_1, R) = \bigoplus_i \delta_{\bar{n}_i} \mathcal{D}(\bar{N}_2, R)$ we see that t acts by a sum of maps of the form $\mathcal{D}(\bar{N}_2, R) \rightarrow \mathcal{D}(\bar{N}_2, R)$ induced by $\bar{n} \mapsto t\bar{n}t^{-1}$, and the proof now proceeds as in the case when $p \neq 2$. \square

Continue to assume $r \geq r_\kappa$. In the rest of this subsection, for simplicity of notation we will assume $p \neq 2$ in the discussion, but everything we do works for $p = 2$ as well after minor adjustments, writing $\mathcal{D}_\kappa^r \cong \bigoplus_i \delta_{\bar{n}_i} \mathcal{D}^r(\bar{N}_2, R)$. Let $\bar{n}_1, \dots, \bar{n}_t$ be a topological basis of \bar{N}_1 and set $\mathbf{n}_i = [\bar{n}_i] - 1 \in \mathcal{D}_\kappa$ as usual. Then the description

$$\mathcal{D}^r(\bar{N}_1, R) = \widehat{\bigoplus_{\alpha} R \cdot \varpi^{-n(r, \varpi, \alpha)} \mathbf{n}^\alpha}$$

gives us an explicit description of \mathcal{D}_κ^r via the identification $\mathcal{D}_\kappa^r \cong \mathcal{D}^r(\bar{N}_1, R)$. In particular we remark that the \mathcal{D}_κ^r are potentially ON-able with a countable potential ON-basis that we can actually write down. This is in contrast with the compact distribution modules considered in [Ash and Stevens 2008], which are potentially ON-able but one cannot write down an explicit basis (and the dimension is uncountable), as well as the distribution modules considered in [Hansen 2017], which are not known to be potentially ON-able in general. Our next goal is to introduce variants of the modules \mathcal{D}_κ^r , as well as modules $\mathcal{A}_\kappa^r \subseteq \mathcal{A}_\kappa$, which are analogous to those considered in [loc. cit.].

Let $r > r_\kappa$ and pick $s \in [r_\kappa, r)$. The unit ball $\mathcal{D}_\kappa^{r, \circ}$ of \mathcal{D}_κ^r is Δ -stable since Δ acts by norm-decreasing operators. The natural map $\mathcal{D}_\kappa^r \rightarrow \mathcal{D}_\kappa^s$ is injective, and may naturally be thought of as an inclusion. Doing so, we define $\mathcal{D}_\kappa^{<r, \circ}$ to be the closure of $\mathcal{D}_\kappa^{r, \circ}$ inside \mathcal{D}_κ^s . It is a ϖ -torsion free R_0 -module and we set $\mathcal{D}_\kappa^{<r} = \mathcal{D}_\kappa^{<r, \circ}[1/\varpi]$; this is an R -module which naturally embeds into \mathcal{D}_κ^s . Since $\mathcal{D}_\kappa^{r, \circ} \subseteq \mathcal{D}_\kappa^s$ is Δ -stable we see that $\mathcal{D}_\kappa^{<r, \circ}$ and $\mathcal{D}_\kappa^{<r}$ are as well. We may then define

$$\mathcal{A}_\kappa^{<r, \circ} = \{f \in \mathcal{A}_\kappa \mid |\mu(f)| \leq 1 \text{ for all } \mu \in \mathcal{D}_\kappa \cap \mathcal{D}_\kappa^{r, \circ}\}$$

and $\mathcal{A}_\kappa^{<r} = \mathcal{A}_\kappa^{<r, \circ}[1/\varpi] \subseteq \mathcal{A}_\kappa$. Then $\mathcal{A}_\kappa^{<r}$ is the dual space of \mathcal{D}_κ^r . We equip it with the norm dual to $\|\cdot\|_r$, and define $\mathcal{A}_\kappa^r \subseteq \mathcal{A}_\kappa^{<r}$ to be the closure of $\mathcal{A}_\kappa^{<s} \subseteq \mathcal{A}_\kappa^{<r}$ with respect to this norm. These spaces are Δ -stable since $\mathcal{D}_\kappa \cap \mathcal{D}_\kappa^{r, \circ}$ is. Note that we have natural identifications

$$\mathcal{D}_\kappa^{<r} \cong \mathcal{D}^{<r}(\bar{N}_1, R), \quad \mathcal{A}_\kappa^{<r} \cong \mathcal{C}^{<r}(\bar{N}_1, R), \quad \mathcal{A}_\kappa^r \cong \mathcal{C}^r(\bar{N}_1, R),$$

so the discussion in Section 3.2 applies to give explicit descriptions of these spaces, and show that they are independent of the choice of s .

4. Overconvergent cohomology and eigenvarieties

In this section we establish the basic results on overconvergent cohomology needed to construct and analyze eigenvarieties. We retain the notation from Section 3.3, but we will change our point of view slightly, from a functional-analytic point of view to a geometric one. Instead of working with Banach–Tate \mathbb{Z}_p -algebras, we will work with complete Tate \mathbb{Z}_p -algebras, which we will always assume to have a

Noetherian ring of definition. A *weight* will therefore be a continuous homomorphism $\kappa : T_0 \rightarrow R^\times$, where R is a complete Tate \mathbb{Z}_p -algebra with a Noetherian ring of definition. We follow the strategy of [Hansen 2017, §3–4] to construct our eigenvarieties. A similar construction was also carried out in [Xiang 2012].

4.1. Eigenvarieties. We retain the global setup from the beginning of Section 3.3. To construct our eigenvarieties, we will need some more notation as well as some concepts from [Ash and Stevens 2008; Hansen 2017]. First, let us fix a compact open subgroup $K_\ell = K_\ell \subseteq \mathbf{G}(\mathbb{Q}_\ell)$, for each prime $\ell \neq p$, which is hyperspecial for all but finitely many ℓ , and set $K^p = \prod_{\ell \neq p} K_\ell$ (the tame level) and $K = K^p I$. We assume that K is neat (which is the case when K^p is sufficiently small).⁴ Let \mathbf{Z} denote the centre of \mathbf{G} and put $Z(K) = \mathbf{Z}(\mathbb{Q}) \cap K$. All weights in this section will be assumed to be trivial on $\overline{Z(K)} \subseteq T_0$.

We also fix a monoid $\Delta_\ell \subseteq \mathbf{G}(\mathbb{Q}_\ell)$ containing K_ℓ , which is equal to $\mathbf{G}(\mathbb{Q}_\ell)$ when K_ℓ is hyperspecial, such that (Δ_ℓ, K_ℓ) is a Hecke pair and the ℓ -Hecke algebra $\mathbb{T}(\Delta_\ell, K_\ell)$ (defined over \mathbb{Z}_p) is commutative. Set $\Delta^p = \prod' \Delta_\ell$ (restricted product with respect to the K_ℓ) and $\Delta = \Delta^p \Delta_p$ (recall that $\Delta_p = I \Sigma^+ I$). Next, as in [Hansen 2017, §2.1], we fix a choice $C_\bullet(K, -)$ of an *augmented Borel–Serre complex* and for any left Δ -module M we define $C^\bullet(K, M)$. Note that $C^\bullet(K, M)$ carries an action of the Hecke algebra $\mathbb{T}(\Delta, K)$. In general, if C^\bullet is a cochain complex we let $C^* = \bigoplus_{i \in \mathbb{Z}} C^i$ and, similarly, we use H^* to denote the direct sum of all cohomology groups when cohomology makes sense.

Fix once and for all an element $t \in \Sigma^{\text{cpt}}$. Let $\kappa : T_0 \rightarrow R^\times$ be a weight, and choose a Banach–Tate \mathbb{Z}_p -algebra norm on R which is adapted to κ . We let $\tilde{U}_{\kappa, r} = \tilde{U}_{t, \kappa, r}$ denote the corresponding Hecke operator on $C^\bullet(K, \mathcal{D}_\kappa^r)$ (here $r \geq r_\kappa$). This operator is compact and we let

$$F_\kappa^r(T) = \det(1 - T \tilde{U}_{\kappa, r} \mid C^*(K, \mathcal{D}_\kappa^r))$$

denote its Fredholm determinant, which exists since $C^*(K, \mathcal{D}_\kappa^r)$ is potentially ON-able (by basic properties of Borel–Serre complexes). Before proceeding, let us recall the definition of weight space.

Definition 4.1.1. Suppose (A, A^+) is a complete sheafy affinoid $(\mathbb{Z}_p, \mathbb{Z}_p)$ -algebra. The functor

$$(A, A^+) \mapsto \text{Hom}_{\text{cts}}(T_0/\overline{Z(K)}, A^\times)$$

from complete sheafy affinoid $(\mathbb{Z}_p, \mathbb{Z}_p)$ -algebras (A, A^+) to sets is representable by the affinoid ring $(\mathbb{Z}_p \llbracket T_0/\overline{Z(K)} \rrbracket, \mathbb{Z}_p \llbracket T_0/\overline{Z(K)} \rrbracket)$, and we let \mathfrak{W} denote the corresponding adic space. We remark that any continuous homomorphism $T_0/\overline{Z(K)} \rightarrow A^\times$ automatically lands in $(A^+)^\times$. To see this, note that T_0 is noncanonically isomorphic to $F \times \mathbb{Z}_p^r$ as a p -adic Lie group, where F is a finite group and $r \in \mathbb{Z}_{\geq 0}$. The image of F lands in the roots of unity $\mu_\infty(A)$ in A , and the image of \mathbb{Z}_p^r lands in $1 + A^{\circ\circ}$. Since A^+ is open and integrally closed in A (which is complete), $\mu_\infty(A)$ and $1 + A^{\circ\circ}$ are both subsets of $(A^+)^\times$.

We let \mathcal{W} denote the *analytic locus* of \mathfrak{W} ; this is an open subset. For any weight $\kappa : T_0 \rightarrow R^\times$ and ring of integral elements $R^+ \subseteq R^\circ$, we obtain a map $\mathcal{U} = \text{Spa}(R, R^+) \rightarrow \mathcal{W}$. If this map is an open immersion (so in particular we have $R^+ = R^\circ$, by Corollary A.6), we will conflate the weight κ and the open subset $\mathcal{U} \subseteq \mathcal{W}$ and refer to \mathcal{U} as an *open weight*. In this case, we will also replace κ by \mathcal{U} in our notation, writing $\mathcal{D}_\mathcal{U}$ etc.

⁴In fact it suffices to assume that K contains a neat open normal subgroup with index prime to p .

Proposition 4.1.2. *Let κ be a weight and choose an adapted Banach–Tate \mathbb{Z}_p -algebra norm on R . Then F_κ^r is independent of $r \geq r_\kappa$.*

Proof. Let $r_\kappa \leq r < s$; we wish to prove that $F_\kappa^r = F_\kappa^s$. Note that the general case follows from the case $s \leq r^{1/p}$, so we may assume this. Then, by Corollary 3.3.10, $\tilde{U}_{\kappa,r}$ factors as

$$C^*(K, \mathcal{D}_\kappa^r) \xrightarrow{\Phi} C^*(K, \mathcal{D}_\kappa^s) \xrightarrow{\iota} C^*(K, \mathcal{D}_\kappa^r),$$

where ι is induced by the natural compact inclusion $\mathcal{D}_\kappa^s \hookrightarrow \mathcal{D}_\kappa^r$. We have $\tilde{U}_{\kappa,s} = \Phi \circ \iota$, so the result follows from [Buzzard 2007, Lemma 2.7]. \square

In light of this we will from now on drop r from the notation and simply write F_κ . We remark that it currently depends on a choice of norm on R .

Proposition 4.1.3. *Let $\kappa : T_0 \rightarrow R^\times$ be a weight and choose an adapted Banach–Tate \mathbb{Z}_p -algebra norm on R . Let ϖ be a multiplicative pseudouniformizer in R :*

(1) *Assume that F_κ has a factorization $F_\kappa = QS$, where Q is a multiplicative polynomial, S is a Fredholm series, and Q and S are relatively prime. Let $s > r \geq r_\kappa$. The inclusion $C^\bullet(K, \mathcal{D}_\kappa^s) \subseteq C^\bullet(K, \mathcal{D}_\kappa^r)$ induces an equality $\text{Ker}^\bullet Q^*(\tilde{U}_{\kappa,s}) = \text{Ker}^\bullet Q^*(\tilde{U}_{\kappa,r})$ (here and elsewhere we write $\text{Ker}^\bullet Q^*(\tilde{U}_{\kappa,r})$ for the complex with i -th term being the kernel of $Q^*(\tilde{U}_{\kappa,r})$ acting on $C^i(K, \mathcal{D}_\kappa^r)$).*

(2) *Let R' be a complete Tate ring with a Noetherian ring of definition, which we assume to be equipped with a Banach–Tate \mathbb{Z}_p -algebra norm $|\cdot|'$ which induces the topology. Assume that we have a bounded homomorphism $\phi : R \rightarrow R'$ such that $|\cdot|'$ is adapted to $\kappa' = \kappa \circ \phi$ and $\phi(\varpi)$ is multiplicative for $|\cdot|'$. Then $F_{\kappa'} = \phi(F_\kappa)$.*

If we assume moreover that F_κ has a factorization as in the previous part, we have a canonical isomorphism $(\text{Ker}^\bullet Q^(\tilde{U}_{\kappa,r})) \otimes_R R' \cong \text{Ker}^\bullet Q^*(\tilde{U}_{\kappa',r})$.*

Proof. For assertion (1), we note that the general case follows from the case $s \leq r^{1/p}$, and then writing $\tilde{U}_{\kappa,r}$ and $\tilde{U}_{\kappa,s}$ as in the proof of Proposition 4.1.2 the result follows from Lemma 2.2.3. For part (2), the first assertion follows from Lemma 3.2.5 and [Buzzard 2007, Lemma 2.13], and the second assertion follows from Lemma 3.2.5 upon writing $C^\bullet(K, \mathcal{D}_\kappa^r) = \text{Ker}^\bullet Q^*(\tilde{U}_{\kappa,r}) \oplus N^\bullet$ as in Theorem 2.2.2 since $Q^*(\tilde{U}_{\kappa,r})$ is invertible on N^\bullet . \square

We can now prove that F_κ is independent of the choice of norm on R .

Proposition 4.1.4. *F_κ is independent of the choice of adapted Banach–Tate \mathbb{Z}_p -algebra norm on R .*

Proof. Let $|\cdot|$ and $|\cdot|'$ be two different such norms on R and denote the constructions coming from $|\cdot|$ as usual and the constructions coming from $|\cdot|'$ by adding a prime. By Lemma 2.1.6, we may find constants $C_1, C_2, s_1, s_2 > 0$ such that $C_1|a|^{s_1} \leq |a|' \leq C_2|a|^{s_2}$ for all $a \in R$, which, as in the proof of Proposition 3.2.6, implies that $\mathcal{D}_\kappa^{r^{1/s_2}} \subseteq \mathcal{D}_\kappa^{r','} \subseteq \mathcal{D}_\kappa^{r^{1/s_1}}$ for r suitably close to 1. Assume first that $s_2/s_1 < p$, or in other words that $r^{p/s_2} < r^{1/s_1}$. Then $\tilde{U}_{\kappa,r^{1/s_1}}$ factors through $\mathcal{D}_\kappa^{r^{1/s_2}}$, and hence through $\mathcal{D}_\kappa^{r','}$. Decreasing s_1 if necessary (still making sure that $s_2/s_1 \leq p$), the inclusion $\mathcal{D}_\kappa^{r','} \subseteq \mathcal{D}_\kappa^{r^{1/s_1}}$ is compact, and we may now argue as in the proof of Proposition 4.1.2.

We now do the general case. Choose multiplicative pseudouniformizers ϖ and ϖ' for $|-|$ and $|-|'$ respectively. Put $X = \mathrm{Spa}(R, R^\circ)$ and let $U \subseteq X$ be a rational subdomain. Write $R_U := \mathcal{O}_X(U)$ and ϕ_U for the natural map $R \rightarrow R_U$. Then it is easy to see, by (the proof of) Lemma 3.3.1, that we may find a Banach–Tate norm $|-|_U$ on R_U such that $\phi_U : (R, |-|) \rightarrow (R_U, |-|_U)$ is bounded, ϖ is a multiplicative pseudouniformizer for $|-|_U$, and for all t_U sufficiently large, $|-|_U^{t_U}$ is a Banach–Tate \mathbb{Z}_p -algebra norm on R_U which is adapted for $\kappa_U := \phi_U \circ \kappa$. We may find a norm $|-|'_U$ with the same properties in relation to $|-|'$ etc. By Proposition 4.1.3(2) we then have $F_{\kappa_U} = \phi_U(F_\kappa)$ and $F'_\kappa = \phi_U(F'_\kappa)$, using that changing a norm by raising it to a real positive power only reindexes the \mathcal{D}^r , and hence does not change the Fredholm determinant by Proposition 4.1.2. We also remark that raising $|-|$ and $|-|'$ to the same power $t > 0$ does not change the quantity s_2/s_1 . Thus, we aim to find an open cover $(U_i)_i$ of $\mathrm{Spa}(R, R^\circ)$ consisting of rational subdomains, such that, writing $R_i = \mathcal{O}(U_i)$ and equipping with norms of the form described above, the quantity s_1/s_2 is $< p$ for each i . This would then finish the proof.

It remains to construct the cover $(U_i)_i$. We recall from the proof of Lemma 2.1.6 that we may take

$$s_1 = \frac{\log |\varpi^m|'}{\log |\varpi^m|}, \quad s_2 = \frac{\log |(\varpi')^m|'}{\log |(\varpi')^m|}$$

for any m sufficiently large that $|\varpi^m|', |(\varpi')^m| < 1$. Let $\delta > 0$ be small, and choose m sufficiently large that $(|\varpi^m|')^{1/m} - |\varpi|'_{\mathrm{sp}} \leq \delta$ and $|(\varpi')^m|^{1/m} - |\varpi'|_{\mathrm{sp}} \leq \delta$. There is a continuous real-valued function Φ on $\mathcal{M}(R)$ given by

$$x \mapsto \Phi(x) = \frac{\log |\varpi'|' \log |\varpi|}{\log |\varpi'|_x \log |\varpi|'_x} = \frac{\log |\varpi'|'_x \log |\varpi|_x}{\log |\varpi'|_x \log |\varpi|'_x}.$$

Note the similarity between s_2/s_1 and Φ . We recall here that $\mathcal{M}(R)$ is the maximal Hausdorff quotient of $\mathrm{Spa}(R, R^\circ)$, and hence it does not depend on whether we used $|-|$ or $|-|'$ to construct it (though of course the functions $x \mapsto |-|_x$ and $x \mapsto |-|'_x$ are in general different). We compose this function with the projection $\mathrm{Spa}(R, R^\circ) \rightarrow \mathcal{M}(R)$ to get a function on $\mathrm{Spa}(R, R^\circ)$. We claim that it is constant and equal to 1. To see this, fix x and note that there is an $s > 0$ such that $|-|'_x = |-|_x^s$; one then checks easily that $\Phi(x) = 1$. Fix $x \in \mathrm{Spa}(R, R^\circ)$. Let U be a rational subdomain containing x and give $\mathcal{O}(U)$ two norms $|-|$ and $|-|'$ constructed as for R_i above. If U is small, $|\varpi|'_{\mathrm{sp}} = \sup_{y \in \mathcal{M}(\mathcal{O}(U)) \subseteq \mathcal{M}(R)} |\varpi|'_y$ is close to $|\varpi|'_x$, and similarly for $|\varpi'|_{\mathrm{sp}}$. Choosing δ small, we may then ensure that the quantity s_2/s_1 is close to $\Phi(x) = 1$ for U small; in particular it is $< p$ as desired. Picking such a U for every x gives the desired cover. \square

From the preceding two propositions, we can immediately deduce the following corollary.

Corollary 4.1.5. *Let $\mathcal{U}_1 \subseteq \mathcal{U}_2$ be open weights and let $\phi : \mathcal{O}_{\mathcal{W}}(\mathcal{U}_2) \rightarrow \mathcal{O}_{\mathcal{W}}(\mathcal{U}_1)$ be the induced map. Then $\phi(F_{\mathcal{U}_2}) = F_{\mathcal{U}_1}$. Therefore, the Fredholm determinants $(F_{\mathcal{U}})_{\mathcal{U}}$, where \mathcal{U} ranges over all open weights, glue together to a Fredholm series $F_{\mathcal{W}} \in \mathcal{O}(\mathcal{W})\{\{T\}\} = \mathcal{O}(\mathbb{A}_{\mathcal{W}}^1)$.*

We write \tilde{U}_κ for the Hecke operator on $C^\bullet(K, \mathcal{D}_\kappa)$ coming from our fixed $t \in \Sigma^{\mathrm{cpt}}$.

Proposition 4.1.6. *Let $\kappa : T_0 \rightarrow R^\times$ be a weight. Assume that F_κ has a factorization $F_\kappa = QS$, where Q is a multiplicative polynomial, S is a Fredholm series, and Q and S are relatively prime. Then $\text{Ker}^\bullet Q^*(\tilde{U}_\kappa)$ is a complex of projective R -modules. If $\phi : R \rightarrow R'$ is a continuous homomorphism, where R' is a complete Tate ring with a Noetherian ring of definition, then we have a canonical isomorphism $(\text{Ker}^\bullet Q^*(\tilde{U}_\kappa)) \otimes_R R' \cong \text{Ker}^\bullet Q^*(\tilde{U}_{\kappa'})$, where $\kappa' = \kappa \circ \phi$.*

Proof. We have $C^\bullet(K, \mathcal{D}_\kappa) = \varprojlim_{r \geq r_\kappa} C^\bullet(K, \mathcal{D}_\kappa^r)$. Since $\tilde{U}_\kappa = \varprojlim_{r \geq r_\kappa} \tilde{U}_{\kappa,r}$ the result follows from Proposition 4.1.3 upon noting that we may choose a topologically nilpotent unit $\varpi \in R$ and norms on R and R' such that ϕ and ϖ satisfies the assumptions of that proposition. \square

Next we study what happens when the factorization of F_κ changes. Keep the notation of Proposition 4.1.6. We make $\text{Ker}^\bullet Q^*(\tilde{U}_\kappa)$ into a complex of $R[T]/(Q(T))$ -modules by letting T act as \tilde{U}_κ^{-1} .

Proposition 4.1.7. *Let $\kappa : T_0 \rightarrow R^\times$ be a weight. Assume that F_κ has two factorizations $F_\kappa = Q_1 S_1 = Q_2 S_2$, where the Q_i are multiplicative polynomials, the S_i are Fredholm series, and for each i the Q_i and S_i are relatively prime. Assume further that $Q_1 \mid Q_2$. Then we have a canonical isomorphism*

$$\text{Ker}^\bullet Q_2^*(\tilde{U}_\kappa) \otimes_{R[T]/(Q_2)} R[T]/(Q_1) \cong \text{Ker}^\bullet (Q_1^*(\tilde{U}_\kappa)).$$

Proof. Let P be such that $Q_2 = PQ_1$; P is then a multiplicative polynomial and one checks that it is relatively prime to Q_1 , so we may find polynomials $A, B \in R[T]$ such that $PA + Q_1 B = 1$. We then have $R[T]/(Q_2) \cong R[T]/(Q_1) \times R[T]/(P)$ and $PA \in R[T]/(Q_2)$ corresponds to $(1, 0) \in R[T]/(Q_1) \times R[T]/(P)$. The proposition now amounts to showing that $PA \cdot \text{Ker}^\bullet Q_2^*(\tilde{U}_\kappa) = \text{Ker}^\bullet Q_1^*(\tilde{U}_\kappa)$. If $x \in \text{Ker}^\bullet Q_2^*(\tilde{U}_\kappa)$, then

$$Q_1^*(\tilde{U}_\kappa) \cdot PAx = \tilde{U}_\kappa^{\deg Q_1} A Q_2 x = 0,$$

which gives us one inclusion. For the other, assume that $y \in \text{Ker}^\bullet Q_1^*(\tilde{U}_\kappa)$. Note that $\deg PA = \deg Q_1 B$. Then

$$y = \tilde{U}_\kappa^{-\deg PA} (P^*(\tilde{U}_\kappa) A^*(\tilde{U}_\kappa) + B^*(\tilde{U}_\kappa) Q_1^*(\tilde{U}_\kappa)) y = P^*(\tilde{U}_\kappa) A^*(\tilde{U}_\kappa) \tilde{U}_\kappa^{-\deg PA} y,$$

which gives us the other inclusion. \square

We now return to the Fredholm series $F_\mathcal{W}$ from Corollary 4.1.5. Let $\mathcal{Z} \subseteq \mathbb{A}_\mathcal{W}^1$ denote its Fredholm hypersurface. Consider $\mathcal{Cov}(\mathcal{Z})$, the set of all open affinoid $V \subseteq Z$ such that $\pi(V) \subseteq X$ is open affinoid, $\mathcal{O}(\pi(V))$ is Tate, and the map $\pi|_V : V \rightarrow \pi(V)$ is finite of constant degree, where $\pi : \mathcal{Z} \rightarrow \mathcal{W}$ is the projection map. For $V \in \mathcal{Cov}(\mathcal{Z})$, let us write $F_\mathcal{W} = Q_V S_V$ for the associated factorization of $F_\mathcal{W}$ from Theorem 2.3.3.

Corollary 4.1.8. *The assignment $V \mapsto \text{Ker}^\bullet Q_V^*(\tilde{U}_{\pi(V)})$, with $V \in \mathcal{Cov}(\mathcal{Z})$, defines a bounded complex of coherent sheaves \mathcal{K}^\bullet on \mathcal{Z} .*

Proof. $\mathcal{Cov}(\mathcal{Z})$ is an open cover of \mathcal{Z} so we need to prove that whenever $V_1 \subseteq V_2$ are elements of $\mathcal{Cov}(\mathcal{Z})$, we have $(\text{Ker}^\bullet Q_{V_2}^*(\tilde{U}_{\pi(V_2)})) \otimes_{\mathcal{O}(V_2)} \mathcal{O}(V_1) \cong \text{Ker}^\bullet Q_{V_1}^*(\tilde{U}_{\pi(V_1)})$ canonically. Define $V_3 = \pi|_{V_2}^{-1}(\pi(V_1))$. Then we have $V_1 \subseteq V_3 \subseteq V_2$, so it suffices to treat the inclusions $V_1 \subseteq V_3$ and $V_3 \subseteq V_2$. In the first case

we have $\pi(V_1) = \pi(V_3)$ and the result follows from Proposition 4.1.7, since $V_1 \subseteq V_3$ forces $\mathcal{Q}_{V_1} \mid \mathcal{Q}_{V_3}$. For the second we have $V_3 = V_2 \times_{\pi(V_2)} \pi(V_1)$ and the result follows from Proposition 4.1.6. \square

This allows us to finish the construction of the eigenvariety. We define $\mathcal{H}^* = H^*(\mathcal{H}^\bullet)$; this is a coherent sheaf on \mathcal{Z} . Since the projectors $C^\bullet(K, \mathcal{D}_{\pi(V)}) \rightarrow \text{Ker}^\bullet Q_V^*(\tilde{U}_{\pi(V)})$ commute with the action of $\mathbb{T}(\Delta^p, K^p)$ (by construction, using the assertion about the projectors in Theorem 2.2.2), we get an induced action $\mathbb{T}(\Delta^p, K^p) \rightarrow \text{End}_{\mathcal{Z}}(\mathcal{H}^*)$. Let $\mathcal{T} \subseteq \text{End}_{\mathcal{Z}}(\mathcal{H}^*)$ denote the sub-presheaf generated over $\mathcal{O}_{\mathcal{Z}}$ by the image of $\mathbb{T}(\Delta^p, K^p)$. It is a sheaf by flatness of rational localization, hence a coherent sheaf of $\mathcal{O}_{\mathcal{Z}}$ -algebras, and we define the eigenvariety $\mathcal{X} = \mathcal{X}_{G, K^p}$ to be the relative $\text{Spa}(\mathcal{T}, \mathcal{T}^\circ) \rightarrow \mathcal{Z}$ (note that the sheaf of integral elements is determined by Lemma A.3). The morphism $q : \mathcal{X} \rightarrow \mathcal{Z}$ is finite by construction, and we have

$$\mathcal{O}(q^{-1}(V)) = \text{Im}(\mathbb{T}(\Delta^p, K^p) \otimes_{\mathbb{Z}_p} \mathcal{O}(V) \rightarrow \text{End}_{\mathcal{O}(V)}(H^*(\text{Ker}^\bullet Q_V^*)))$$

for all $V \in \text{Cov}(\mathcal{Z})$. In particular, if (\mathcal{U}, h) is a slope datum for $(\mathcal{W}, F_{\mathcal{W}})$, we write $\mathcal{T}_{\mathcal{U}, h} = \mathcal{O}(q^{-1}(\mathcal{Z}_{\mathcal{U}, h}))$ and have

$$\mathcal{T}_{\mathcal{U}, h} = \text{Im}(\mathbb{T}(\Delta^p, K^p) \otimes_{\mathbb{Z}_p} \mathcal{O}(\mathcal{Z}_{\mathcal{U}, h}) \rightarrow \text{End}_{\mathcal{O}(\mathcal{Z}_{\mathcal{U}, h})}(H^*(K, \mathcal{D}_{\mathcal{U}})_{\leq h})).$$

Remark 4.1.9. Our eigenvariety \mathcal{X} contains the eigenvariety constructed in [Hansen 2017, §4] as the open subset $\{p \neq 0\}$. Indeed, our construction specializes to his over Banach \mathbb{Q}_p -algebras, with the minor difference that we use the complexes $C^\bullet(K, \mathcal{D}_{\mathcal{U}}^r)$ to construct the auxiliary Fredholm hypersurface \mathcal{Z} , whereas the complexes $C_\bullet(K, \mathcal{A}_{\mathcal{U}}^r)$ (in our notation) are used in [loc. cit.], giving a different auxiliary Fredholm hypersurface. However, working over the union of the two Fredholm hypersurfaces, one sees that the coherent sheaf \mathcal{H}^* on $\mathbb{A}_{\mathcal{W}^{\text{rig}}}^1$, with its $\mathbb{T}(\Delta^p, K^p)$ -action, is equal to the sheaf \mathcal{M}^* on $\mathbb{A}_{\mathcal{W}^{\text{rig}}}^1$ (in the notation of [loc. cit., §4.3]), with its $\mathbb{T}(\Delta^p, K^p)$ -action (here we have used \mathcal{W}^{rig} to denote the locus $\{p \neq 0\} \subseteq \mathcal{W}$).

Remark 4.1.10. Like the other constructions, our construction of overconvergent cohomology and eigenvarieties has numerous variations, which are sometimes useful to keep in mind. For example, one may use compactly supported cohomology, homology or Borel–Moore homology instead (see [loc. cit., §3.3]), and/or one could use the modules $\mathcal{A}_{\mathcal{K}}$ instead of the $\mathcal{D}_{\mathcal{K}}$. One can also add (or remove) Hecke operators, or work over some restricted family of weights, rather than the universal one.

4.2. The Tor-spectral sequence. We now give the analogue of the Tor spectral sequence in [Hansen 2017, Theorem 3.3.1], which is a key tool in analyzing the eigenvarieties. We phrase it in terms of slope decompositions and Banach–Tate rings, though we could have formulated it more generally for elements in $\text{Cov}(\mathcal{Z})$.

Theorem 4.2.1. *Let $h \in \mathbb{Q}_{\geq 0}$ and let $\kappa : T_0 \rightarrow R^\times$ be a weight. We fix an adapted Banach–Tate \mathbb{Z}_p -algebra norm on R , and suppose that $C^\bullet(K, \mathcal{D}_{\mathcal{K}}^r)$ has a slope $\leq h$ -decomposition for some $r \geq r_{\mathcal{K}}$. Let $R \rightarrow S$ be a bounded homomorphism of Banach–Tate \mathbb{Z}_p -algebras with adapted norms and write κ_S for the induced*

weight $T_0 \rightarrow S^\times$. Then there is a convergent Hecke-equivariant (cohomological) second quadrant spectral sequence

$$E_2^{ij} = \mathrm{Tor}_{-i}^R(H^j(K, \mathcal{D}_\kappa)_{\leq h}, S) \Rightarrow H^{i+j}(K, \mathcal{D}_{\kappa_S})_{\leq h}.$$

Proof. We follow the proof of [Hansen 2017, Theorem 3.3.1]. Define a chain complex \mathcal{C}_\bullet by $\mathcal{C}_i = C^{-i}(K, \mathcal{D}_\kappa)_{\leq h}$ and the obvious differentials (i.e., we are just reindexing and viewing $C^\bullet(K, \mathcal{D}_\kappa)_{\leq h}$ as a chain complex). This is a bounded chain complex of finite projective R -modules. Thus

$$\mathbf{Tor}_{i+j}^R(\mathcal{C}_\bullet, S) = H_{i+j}(\mathcal{C}_\bullet \otimes_R S),$$

where \mathbf{Tor} denotes hyper-Tor. The hyper-Tor spectral sequence then gives us a homological spectral sequence

$$E_{ij}^2 = \mathrm{Tor}_i^R(H_j(\mathcal{C}_\bullet), S) \Rightarrow H_{i+j}(\mathcal{C}_\bullet \otimes_R S)$$

which is concentrated in the fourth quadrant. Reindexing we may turn this into a cohomological spectral sequence (see [Weibel 1994, Dual definition 5.2.3])

$$E_2^{ij} = \mathrm{Tor}_{-i}^R(H_{-j}(\mathcal{C}_\bullet), S) \Rightarrow H_{-i-j}(\mathcal{C}_\bullet \otimes_R S)$$

which is concentrated in the second quadrant. Since $H_{-j}(\mathcal{C}_\bullet) = H^j(K, \mathcal{D}_\kappa)_{\leq h}$ (by definition) and $H_{-i-j}(\mathcal{C}_\bullet \otimes_R S) \cong H^{i+j}(K, \mathcal{D}_{\kappa_S})_{\leq h}$ (canonically, by our previous results) this gives the desired spectral sequence. Finally, Hecke-equivariance follows from the functoriality of the hyper-Tor spectral sequence. \square

As an application, we prove the following analogue of [Hansen 2017, Theorem 4.3.3]. We will use it in the next section when we construct Galois representations.

Proposition 4.2.2. *Let (\mathcal{U}, h) be a slope datum and let $\mathfrak{m} \subseteq \mathcal{O}_{\mathcal{W}}(\mathcal{U})$ be a maximal ideal corresponding to a weight $\kappa : T_0 \rightarrow L^\times$, where $L = \mathcal{O}_{\mathcal{W}}(\mathcal{U})/\mathfrak{m}$; this is a local field by Lemma A.12. Fix an absolute value on L with $|p| \leq p^{-1}$ (i.e., an adapted Banach–Tate \mathbb{Z}_p -algebra norm). Let $\mathbb{T} \subseteq \mathbb{T}(\Delta, K)$ be a \mathbb{Z}_p -subalgebra and put*

$$\begin{aligned} \mathbb{T}_{\kappa, h} &= \mathrm{Im}(\mathbb{T} \otimes_{\mathbb{Z}_p} L \rightarrow \mathrm{End}_L(H^*(K, \mathcal{D}_\kappa)_{\leq h})), \\ \mathbb{T}_{\mathcal{U}, h} &= \mathrm{Im}(\mathbb{T} \otimes_{\mathbb{Z}_p} \mathcal{O}_{\mathcal{W}}(\mathcal{U}) \rightarrow \mathrm{End}_{\mathcal{O}_{\mathcal{W}}(\mathcal{U})}(H^*(K, \mathcal{D}_{\mathcal{U}})_{\leq h})). \end{aligned}$$

Then there is a natural isomorphism $(\mathbb{T}_{\mathcal{U}, h} \otimes_{\mathcal{O}_{\mathcal{W}}(\mathcal{U})} L)^{\mathrm{red}} \cong \mathbb{T}_{\kappa, h}^{\mathrm{red}}$.

Proof. By the Tor-spectral sequence we see that, if $T \in \mathbb{T}$ acts as 0 on $H^*(K, \mathcal{D}_{\mathcal{U}})_{\leq h}$, then it acts nilpotently on $H^*(K, \mathcal{D}_\kappa)_{\leq h}$. It follows that we have a surjection $\mathbb{T}_{\mathcal{U}, h} \otimes_{\mathcal{O}_{\mathcal{W}}(\mathcal{U})} L \twoheadrightarrow \mathbb{T}_{\kappa, h}^{\mathrm{red}}$ of finite-dimensional commutative L -algebras. To finish the proof it suffices to show that if \mathfrak{q} is a maximal ideal of $\mathbb{T}_{\mathcal{U}, h} \otimes_{\mathcal{O}_{\mathcal{W}}(\mathcal{U})} L$ then the localization $(\mathbb{T}_{\kappa, h})_{\mathfrak{q}}$ is nonzero. Let j be maximal such that $(H^j(K, \mathcal{D}_{\mathcal{U}})_{\leq h})_{\mathfrak{q}} \neq 0$ and localize the entire Tor-spectral sequence with respect to \mathfrak{q} . Then the entry $(E_2^{0, j})_{\mathfrak{q}}$ is stable (i.e., $(E_2^{0, j})_{\mathfrak{q}} = (E_\infty^{0, j})_{\mathfrak{q}}$) and it follows that $(H^j(K, \mathcal{D}_\kappa)_{\leq h})_{\mathfrak{q}} \neq 0$. Thus we must have $(\mathbb{T}_{\kappa, h})_{\mathfrak{q}} \neq 0$ as desired. \square

Corollary 4.2.3. *We retain the notation of Proposition 4.2.2. If we let U_t be the double coset operator $[KtK] \in \mathbb{T}(\Delta, K)$ for our fixed $t \in \Sigma^{\mathrm{cpt}}$, consider the commutative subalgebra $\mathbb{T}(\Delta^p, K^p)[U_t] \subseteq \mathbb{T}(\Delta, K)$. Then we have a natural isomorphism $(\mathbb{T}_{\mathcal{U}, h} \otimes_{\mathcal{O}_{\mathcal{W}}(\mathcal{U})} L)^{\mathrm{red}} \cong \mathbb{T}(\Delta^p, K^p)[U_t]_{\kappa, h}^{\mathrm{red}}$.*

Proof. By Proposition 4.2.2 we have a natural isomorphism

$$(\mathbb{T}(\Delta^p, K^p)[U_t]_{\mathcal{U},h} \otimes_{\mathcal{O}(\mathcal{U})} L)^{\text{red}} \cong \mathbb{T}(\Delta^p, K^p)[U_t]_{\kappa,h}^{\text{red}}$$

so it suffices to show that $\mathcal{T}_{\mathcal{U},h} \cong \mathbb{T}(\Delta^p, K^p)[U_t]_{\mathcal{U},h}$, which is clear from the definitions (note that $\text{End}_{\mathcal{O}(\mathcal{T}_{\mathcal{U},h})}(H^*(K, \mathcal{T}_{\mathcal{U}})_{\leq h}) \subseteq \text{End}_{\mathcal{O}(\mathcal{U})}(H^*(K, \mathcal{T}_{\mathcal{U}})_{\leq h})$). \square

5. Galois representations

We continue to assume that all weights are trivial on $\overline{Z(K)}$. Let $G = \text{Res}_{\mathbb{Q}}^F \text{GL}_n/F$, with F a totally real or CM field. In this section we construct a Galois determinant (in the language of [Chenevier 2014]) valued in the global sections of the *reduced* extended eigenvariety, satisfying the expected compatibility between Hecke eigenvalues and the characteristic polynomial of Frobenius at all unramified primes. The construction is an adaptation of a construction due to the first author and David Hansen, to appear in [Hansen and Johansson \geq 2019] (in a slightly refined form), which produces a Galois determinant over the reduced rigid eigenvariety as constructed in [Hansen 2017]. The key step is to produce the desired Galois determinant for all “points” of the extended eigenvariety. In the rigid analytic setting one can then glue these individual determinants together by an argument due to Bellaïche and Chenevier; we prove a version of this gluing technique in our setting. We will not assume that $G = \text{Res}_{\mathbb{Q}}^F \text{GL}_n/F$ until Section 5.3.

5.1. Filtrations. Let $\kappa : T_0 \rightarrow L^\times$ be a weight, where L is a local field equipped with an adapted absolute value; we let ϖ be a uniformizer of L . Let $r > r_\kappa$ and choose an auxiliary $s \in (r_\kappa, r)$. In this subsection, we construct a filtration on the unit ball $\mathcal{D}_\kappa^{<r,\circ}$ of $\mathcal{D}_\kappa^{<r}$ with finite graded pieces, generalizing the filtrations constructed in [Hansen 2015]. We define

$$\text{Fil}^j \mathcal{D}_\kappa^{<r,\circ} := \mathcal{D}_\kappa^{<r,\circ} \cap \varpi^j \mathcal{D}_\kappa^{s,\circ}.$$

When κ and r are clear from the context we will simply write Fil^j for $\text{Fil}^j \mathcal{D}_\kappa^{<r,\circ}$ (we will always omit the choice of s). By the definitions the Fil^j are open and closed in the subspace topology on $\mathcal{D}_\kappa^{<r,\circ}$ coming from \mathcal{D}_κ^s (we recall that $\mathcal{D}_\kappa^{<r,\circ}$ is compact with respect to this subspace topology since \mathcal{O}_L is compact). Therefore the $\mathcal{D}_\kappa^{<r,\circ} / \text{Fil}^j$ are finite discrete \mathcal{O}_L -torsion modules and we have $\mathcal{D}_\kappa^{<r,\circ} = \varprojlim_j \mathcal{D}_\kappa^{<r,\circ} / \text{Fil}^j$ topologically. Note that Fil^j is Δ -stable (being the intersection of two Δ -stable subsets in \mathcal{D}_κ^s), so the \mathcal{O}_L -torsion modules $\mathcal{D}_\kappa^{<r,\circ} / \text{Fil}^j$ inherit a Δ -action and the equality in the previous sentence is Δ -equivariant. We also record the following lemma.

Lemma 5.1.1. *We have*

$$H^i(K, \mathcal{D}_\kappa^{<r,\circ}) = \varprojlim_j H^i(K, \mathcal{D}_\kappa^{<r,\circ} / \text{Fil}^j) \quad \text{for all } i.$$

Proof. On Borel–Serre complexes we have $C^\bullet(K, \mathcal{D}_\kappa^{<r,\circ}) = \varprojlim_j C^\bullet(K, \mathcal{D}_\kappa^{<r,\circ} / \text{Fil}^j)$ and these are bounded complexes of compact abelian Hausdorff groups. This category is abelian and has exact inverse limits (e.g., by [Neukirch 1999, Proposition IV.2.7]), which gives us the result. \square

We remark that $C^\bullet(K, \mathcal{D}_\kappa^{<r})$ has a slope $\leq h$ -decomposition for any $h \in \mathbb{Q}_{\geq 0}$ (since we are working over the field L) and $C^\bullet(K, \mathcal{D}_\kappa^{<r})_{\leq h} = C^\bullet(K, \mathcal{D}_\kappa^r)_{\leq h}$ (since it is sandwiched between $C^\bullet(K, \mathcal{D}_\kappa^r)_{\leq h}$ and $C^\bullet(K, \mathcal{D}_\kappa^s)_{\leq h}$, and these are equal). Thus, we may use the $\mathcal{D}_\kappa^{<r}$ instead of the \mathcal{D}_κ^r to define overconvergent cohomology. We will do this to define Galois representations because of the fact that $\mathcal{D}_\kappa^{<r, \circ}$ is profinite with respect to topology coming from the Fil^j .

5.2. A morphism of Hecke algebras. We continue with the notation of the previous subsection. In this subsection we will fix a \mathbb{Z}_p -subalgebra $\mathbb{T} \subseteq \mathbb{T}(\Delta, K)$ and, if A is any \mathbb{Z}_p -algebra, we will write \mathbb{T}_A for $\mathbb{T} \otimes_{\mathbb{Z}_p} A$. Then $\mathbb{T}_{\mathcal{O}_L}$ acts on $H^*(K, \mathcal{D}_\kappa^{<r, \circ} / \text{Fil}^j)$ for all r and j and we set

$$\begin{aligned} \mathbb{T}_{\kappa, j}^r &= \text{Im}(\mathbb{T}_{\mathcal{O}_L} \rightarrow \text{End}_{\mathcal{O}_L}(H^*(K, \mathcal{D}_\kappa^{<r, \circ} / \text{Fil}^j))), \\ \mathbb{T}_{\kappa, \text{Fil}}^r &= \text{Im}\left(\mathbb{T}_{\mathcal{O}_L} \rightarrow \prod_j \mathbb{T}_{\kappa, j}^r\right). \end{aligned}$$

We equip each $\mathbb{T}_{\kappa, j}^r$ with the discrete topology (they are finite rings) and give $\prod_j \mathbb{T}_{\kappa, j}^r$ the product topology; we then give $\mathbb{T}_{\kappa, \text{Fil}}^r$ the subspace topology. We let $\widehat{\mathbb{T}}_{\kappa, \text{Fil}}^r$ denote the completion of $\mathbb{T}_{\kappa, \text{Fil}}^r$ in $\prod_j \mathbb{T}_{\kappa, j}^r$; this is a compact Hausdorff ring.

Fix $h \in \mathbb{Q}_{\geq 0}$. We have natural maps $H^*(K, \mathcal{D}_\kappa^{<r, \circ}) \rightarrow H^*(K, \mathcal{D}_\kappa^{<r}) \rightarrow H^*(K, \mathcal{D}_\kappa^{<r})_{\leq h}$ and we define $H^*(K, \mathcal{D}_\kappa^{<r, \circ})_{\leq h}$ to be the image of the composition.

Lemma 5.2.1. $H^*(K, \mathcal{D}_\kappa^{<r, \circ})_{\leq h}$ is an open and bounded \mathcal{O}_L -submodule of $H^*(K, \mathcal{D}_\kappa^{<r})_{\leq h}$ (and hence a finite free \mathcal{O}_L -module).

Proof. It is an \mathcal{O}_L -submodule which spans $H^*(K, \mathcal{D}_\kappa^{<r})_{\leq h}$ essentially by construction, so it suffices to show that it is finitely generated. The morphisms $H^*(K, \mathcal{D}_\kappa^{<r, \circ}) \rightarrow H^*(K, \mathcal{D}_\kappa^{<r}) \rightarrow H^*(K, \mathcal{D}_\kappa^{<r})_{\leq h}$ are induced by the morphisms

$$C^\bullet(K, \mathcal{D}_\kappa^{<r, \circ}) \rightarrow C^\bullet(K, \mathcal{D}_\kappa^{<r}) \rightarrow C^\bullet(K, \mathcal{D}_\kappa^{<r})_{\leq h}$$

of complexes. Let $C^\bullet(K, \mathcal{D}_\kappa^{<r, \circ})_{\leq h}$ denote the image of the composition. Note that each $C^i(K, \mathcal{D}_\kappa^{<r, \circ})$ is bounded in $C^i(K, \mathcal{D}_\kappa^{<r})$, so $C^i(K, \mathcal{D}_\kappa^{<r, \circ})_{\leq h}$ is bounded in $C^i(K, \mathcal{D}_\kappa^{<r})_{\leq h}$ by continuity of the projection. Thus $C^\bullet(K, \mathcal{D}_\kappa^{<r, \circ})_{\leq h}$ is a bounded complex of finite free \mathcal{O}_L -modules. Since $H^*(K, \mathcal{D}_\kappa^{<r, \circ})_{\leq h} \subseteq \text{Im}(H^*(C^\bullet(K, \mathcal{D}_\kappa^{<r, \circ})_{\leq h}) \rightarrow H^*(K, \mathcal{D}_\kappa^{<r})_{\leq h})$, finite generation of the former follows. \square

We define

$$\mathbb{T}_{\kappa, \leq h}^{r, \circ} = \text{Im}(\mathbb{T}_{\mathcal{O}_L} \rightarrow \text{End}_{\mathcal{O}_L}(H^*(K, \mathcal{D}_\kappa^{<r, \circ})_{\leq h}));$$

by the above lemma this is a finite \mathcal{O}_L -algebra and hence naturally a compact Hausdorff ring. Note that if $T \in \mathbb{T}_{\mathcal{O}_L}$ is 0 in $\mathbb{T}_{\kappa, \text{Fil}}^r$, i.e., acts as 0 on all $H^*(K, \mathcal{D}_\kappa^{<r, \circ} / \text{Fil}^j)$, then it acts as 0 on $H^*(K, \mathcal{D}_\kappa^{<r, \circ})_{\leq h}$ and so is 0 in $\mathbb{T}_{\kappa, \leq h}^{r, \circ}$. In other words we have a natural (surjective) map $\mathbb{T}_{\kappa, \text{Fil}}^r \rightarrow \mathbb{T}_{\kappa, \leq h}^{r, \circ}$. The goal of this section is to show that this map is continuous and so extends to the completion $\widehat{\mathbb{T}}_{\kappa, \text{Fil}}^r$.

To do this we introduce some special open sets. Let $\text{pr}_j : \mathbb{T}_{\kappa, \text{Fil}}^r \rightarrow \mathbb{T}_{\kappa, j}^r$ denote the projection. We put $U_j = \text{Ker}(\text{pr}_j)$; this is an open ideal of $\mathbb{T}_{\kappa, \text{Fil}}^r$. On $\mathbb{T}_{\kappa, \leq h}^{r, \circ}$ the opens that we will use are more delicate to

construct. We have a commutative diagram

$$\begin{array}{ccc} H^*(K, \mathcal{D}_\kappa^{<r,\circ}) & \longrightarrow & H^*(K, \mathcal{D}_\kappa^{<r})_{\leq h} \\ \downarrow & & \downarrow \\ H^*(K, \mathcal{D}_\kappa^{s,\circ}) & \longrightarrow & H^*(K, \mathcal{D}_\kappa^s)_{\leq h} \end{array}$$

where the right vertical map is an isomorphism, which we think of as an equality (the reader may trace through the definitions and see that this is a natural thing to do). Defining $H^*(K, \mathcal{D}_\kappa^{s,\circ})_{\leq h}$ to be the image of $H^*(K, \mathcal{D}_\kappa^{s,\circ}) \rightarrow H^*(K, \mathcal{D}_\kappa^s)_{\leq h}$ we see that $H^*(K, \mathcal{D}_\kappa^{<r,\circ})_{\leq h} \subseteq H^*(K, \mathcal{D}_\kappa^{s,\circ})_{\leq h}$ are both open lattices in $H^*(K, \mathcal{D}_\kappa^{<r})_{\leq h} = H^*(K, \mathcal{D}_\kappa^s)_{\leq h}$ (Lemma 5.2.1 holds for $H^*(K, \mathcal{D}_\kappa^{s,\circ})_{\leq h}$ as well, with the same proof).

Lemma 5.2.2. *The ideals*

$$V_j = \{T \in \mathbb{T}_{\kappa, \leq h}^{r,\circ} \mid T(H^*(K, \mathcal{D}_\kappa^{<r,\circ})_{\leq h}) \subseteq \varpi^j H^*(K, \mathcal{D}_\kappa^{s,\circ})_{\leq h}\}$$

form a basis of open neighbourhoods of 0 in $\mathbb{T}_{\kappa, \leq h}^{r,\circ}$.

Proof. It's easy to check that the V_j are ideals. By the preceding remarks the subgroups $\varpi^j H^*(K, \mathcal{D}_\kappa^{s,\circ})_{\leq h}$ form a basis of neighborhoods of 0 in $H^*(K, \mathcal{D}_\kappa^{<r,\circ})_{\leq h}$ (for $j \gg 0$). Using this the lemma is elementary. \square

We may now prove continuity. Denote the map $\mathbb{T}_{\kappa, \text{Fil}}^r \rightarrow \mathbb{T}_{\kappa, \leq h}^{r,\circ}$ by ϕ .

Proposition 5.2.3. *We have $\phi(U_j) \subseteq V_j$. Thus ϕ is continuous and extends to a continuous map $\widehat{\mathbb{T}}_{\kappa, \text{Fil}}^r \rightarrow \mathbb{T}_{\kappa, \leq h}^{r,\circ}$.*

Proof. The second statement follows directly from the first (by general properties of linearly topologized groups and the fact that $\mathbb{T}_{\kappa, \leq h}^{r,\circ}$ is complete). The first statement amounts, by the definitions, to proving that if T acts as 0 on $H^*(K, \mathcal{D}_\kappa^{<r,\circ} / \text{Fil}^j)$, then it maps $H^*(K, \mathcal{D}_\kappa^{<r,\circ})_{\leq h}$ into $\varpi^j H^*(K, \mathcal{D}_\kappa^{s,\circ})_{\leq h}$, or in other words that T acts as 0 on

$$\text{Im}(H^*(K, \mathcal{D}_\kappa^{<r,\circ})_{\leq h} \rightarrow H^*(K, \mathcal{D}_\kappa^{s,\circ})_{\leq h} / \varpi^j).$$

By definition this image is equal to

$$\text{Im}(H^*(K, \mathcal{D}_\kappa^{<r,\circ}) \rightarrow H^*(K, \mathcal{D}_\kappa^{s,\circ})_{\leq h} / \varpi^j).$$

Assume now that T acts as 0 on $H^*(K, \mathcal{D}_\kappa^{<r,\circ} / \text{Fil}^j)$. We may factor $H^*(K, \mathcal{D}_\kappa^{<r,\circ}) \rightarrow H^*(K, \mathcal{D}_\kappa^{s,\circ})_{\leq h} / \varpi^j$ through $H^*(K, \mathcal{D}_\kappa^{s,\circ}) / \varpi^j$ so it suffices to prove that T acts as 0 on

$$\text{Im}(H^*(K, \mathcal{D}_\kappa^{<r,\circ}) \rightarrow H^*(K, \mathcal{D}_\kappa^{s,\circ}) / \varpi^j).$$

From the long exact sequence attached to the short exact sequence

$$0 \rightarrow \mathcal{D}_\kappa^{s,\circ} \xrightarrow{\varpi^j} \mathcal{D}_\kappa^{s,\circ} \rightarrow \mathcal{D}_\kappa^{s,\circ} / \varpi^j \rightarrow 0$$

we see that

$$H^*(K, \mathcal{D}_\kappa^{s,\circ}) / \varpi^j \hookrightarrow H^*(K, \mathcal{D}_\kappa^{s,\circ} / \varpi^j).$$

By definition, we have a natural map $\mathcal{D}_\kappa^{<r,\circ}/\text{Fil}^j \rightarrow \mathcal{D}_\kappa^{s,\circ}/\varpi^j$. Assembling the last few sentences, we have a commutative diagram

$$\begin{array}{ccc} H^*(K, \mathcal{D}_\kappa^{<r,\circ}) & \longrightarrow & H^*(K, \mathcal{D}_\kappa^{s,\circ}/\varpi^j) \\ \downarrow & & \downarrow \\ H^*(K, \mathcal{D}_\kappa^{<r,\circ}/\text{Fil}^j) & \longrightarrow & H^*(K, \mathcal{D}_\kappa^{s,\circ}/\varpi^j) \end{array}$$

where the right vertical map is injective. By assumption T acts as 0 on $H^*(K, \mathcal{D}_\kappa^{<r,\circ}/\text{Fil}^j)$; hence it acts as 0 on $\text{Im}(H^*(K, \mathcal{D}_\kappa^{<r,\circ}) \rightarrow H^*(K, \mathcal{D}_\kappa^{s,\circ}/\varpi^j))$. But, since the right vertical map is injective, this image is isomorphic to $\text{Im}(H^*(K, \mathcal{D}_\kappa^{<r,\circ}) \rightarrow H^*(K, \mathcal{D}_\kappa^{s,\circ}/\varpi^j))$. Thus T acts as 0 on this, which is what we wanted to prove. \square

5.3. Galois representations. We now specialize to the case $G = \text{Res}_{\mathbb{Q}}^F \text{GL}_n/F$, where F is a totally real or CM number field, and $n \geq 2$. When discussing Galois representations we will, for simplicity of referencing, use the same conventions as in [Scholze 2015, §5]. Let S' denote the set of places w of \mathbb{Q} such that either $w = \infty$, or if w is finite then K_w is *not* hyperspecial. This is a finite set containing p and the primes which ramify in F . We let S denote the set of places in F lying above those in S' . We set

$$\mathbb{T} = \bigotimes_{v \notin S} \mathbb{T}_v,$$

where $\mathbb{T}_v = \mathbb{Z}_p[\text{GL}_n(F_v)/\text{GL}_n(\mathcal{O}_{F_v})]$ is the usual spherical Hecke algebra (we assume that $K \subseteq \text{GL}_n(\mathcal{O}_F \otimes_{\mathbb{Z}} \widehat{\mathbb{Z}})$). We have $\mathbb{T} \subseteq \mathbb{T}(\Delta, K)$ and we will use the notation and results of the previous subsection for this choice of \mathbb{T} . Let q_v be the size of the residue field at v . We have the (unnormalized) Satake isomorphism

$$\mathbb{T}_v[q_v^{1/2}] \cong \mathbb{Z}_p[q_v^{1/2}][x_1^{\pm 1}, \dots, x_n^{\pm 1}]^{S_n},$$

where S_n is the symmetric group on $\{1, \dots, n\}$ permuting the variables x_1, \dots, x_n . If we let $T_{i,v}$ denote the i -th elementary symmetric polynomial in x_1, \dots, x_n , then $q_v^{i(n+1)/2} T_{i,v} \in \mathbb{T}_v$ and we define

$$P_v(X) = 1 - q_v^{(n+1)/2} T_{1,v} X + q_v^{n+1} T_{2,v} X^2 - \dots + (-1)^n q_v^{n(n+1)/2} T_{n,v} X^n \in \mathbb{T}_v[X]. \quad (5.3.1)$$

The following theorem is essentially a special case of [Scholze 2015, Theorem 5.4.1]. We let $G_{F,S}$ denote the Galois group of the maximal algebraic extension of F unramified outside S . For the notion of a determinant we refer to [Chenevier 2014].

Theorem 5.3.1. (Scholze) *There exists an integer M depending only on $[F : \mathbb{Q}]$ and n such that the following is true: for any j and r there exists an ideal $I_{\kappa,j}^r \subseteq \mathbb{T}_{\kappa,j}^r$ with $(I_{\kappa,j}^r)^M = 0$ and an n -dimensional continuous determinant D of $G_{F,S}$ with values in $\mathbb{T}_{\kappa,j}^r/I_{\kappa,j}^r$ such that*

$$D(1 - X \text{Frob}_v) = P_v(X)$$

for all $v \notin S$.

We remark that the local systems corresponding to the $\mathcal{D}_\kappa^{<r,\circ}/\text{Fil}^j$ are not necessarily included in the formulation of [Scholze 2015, Theorem 5.4.1], but the proof works the same: one first applies the Hochschild–Serre spectral sequence to reduce to the case of a trivial local system.

Corollary 5.3.2. *Keep the notation of Theorem 5.3.1:*

- (1) *There exists a closed ideal $I \subseteq \widehat{\mathbb{T}}_{\kappa, \text{Fil}}^r$ such that $I^M = 0$ and an n -dimensional continuous determinant D of $G_{F,S}$ with values in $\widehat{\mathbb{T}}_{\kappa, \text{Fil}}^r/I$ such that $D(1 - X \text{Frob}_v) = P_v(X)$ for all $v \notin S$.*
- (2) *Let $h \in \mathbb{Q}_{\geq 0}$. Then there exists a closed ideal $J \subseteq \mathbb{T}_{\kappa, \leq h}^{r,\circ}$ such that $J^M = 0$ and an n -dimensional continuous determinant D of $G_{F,S}$ with values in $\mathbb{T}_{\kappa, \leq h}^{r,\circ}/J$ such that $D(1 - X \text{Frob}_v) = P_v(X)$ for all $v \notin S$.*

Proof. Part (1) follows from the theorem and the definition and compactness of $\widehat{\mathbb{T}}_{\kappa, \text{Fil}}^r$ via [Chenevier 2014, Example 2.32]; one sets $I = \widehat{\mathbb{T}}_{\kappa, \text{Fil}}^r \cap \prod_j I_{\kappa, j}^r$. Assertion (2) then follows from (1) and Proposition 5.2.3. \square

5.4. Gluing over the reduced eigenvariety. We now finish our construction of a Galois determinant over the reduced eigenvariety $\mathcal{X}^{\text{red}} = \mathcal{X}_G^{\text{red}}$. See Definition A.10 for the definition of the reduced subspace. Keep the notation of the previous subsection. Let (\mathcal{U}, h) be a slope datum. We have a corresponding open affinoid $\mathcal{X}_{\mathcal{U}, h} \subseteq \mathcal{X}$. If \mathfrak{m}_κ is a maximal ideal of $\mathcal{O}(\mathcal{U})$ corresponding to the weight $\kappa : T_0 \rightarrow L^\times$ with $L = \mathcal{O}(\mathcal{U})/\mathfrak{m}_\kappa$ a local field, then Corollary 4.2.3 gives us a natural identification

$$(\mathcal{O}(\mathcal{X}_{\mathcal{U}, h})/\mathfrak{m}_\kappa)^{\text{red}} = (\mathcal{O}(\mathcal{X}_{\mathcal{U}, h}^{\text{red}})/\mathfrak{m}_\kappa)^{\text{red}} \cong \mathbb{T}(\Delta^p, K^p)[U_t]_{\kappa, h}^{\text{red}}.$$

Fix r . Note that $\mathbb{T}_{\kappa, \leq h}^r = \mathbb{T}_{\kappa, \leq h}^{r,\circ}[1/\varpi]$ is naturally a closed L -subalgebra of $\mathbb{T}(\Delta^p, K^p)[U_t]_{\kappa, h}$. From Corollary 5.3.2 it follows that we have a Galois determinant into $\mathbb{T}(\Delta^p, K^p)[U_t]_{\kappa, h}^{\text{red}}$ and therefore into $(\mathcal{O}(\mathcal{X}_{\mathcal{U}, h}^{\text{red}})/\mathfrak{m}_\kappa)^{\text{red}}$. We record this discussion in the following convenient form:

Lemma 5.4.1. *Let (\mathcal{U}, h) be a slope datum and let \mathfrak{m} be a maximal ideal of $\mathcal{O}(\mathcal{X}_{\mathcal{U}, h}^{\text{red}})$. Then there exists an n -dimensional continuous determinant D of $G_{F,S}$ with values in $\mathcal{O}(\mathcal{X}_{\mathcal{U}, h}^{\text{red}})/\mathfrak{m}$ such that $D(1 - X \text{Frob}_v) = P_v(X)$ for all $v \notin S$.*

Proof. This follows from the discussion above since the map $\text{Spec } \mathcal{O}(\mathcal{X}_{\mathcal{U}, h}^{\text{red}}) \rightarrow \text{Spec } \mathcal{O}(\mathcal{U})$ sends maximal ideals to maximal ideals (the map is finite). \square

Before we can glue we need some preparations. Let A be a reduced \mathbb{Z}_p -algebra which is finite and free as a \mathbb{Z}_p -module; we equip it with the p -adic topology. Consider the adic space $Z^{\text{an}} = \mathbb{A}_{S^{\text{an}}}^1$, where $S = \text{Spa}(A[[X_1, \dots, X_d]], A[[X_1, \dots, X_d]])$; here $A[[X_1, \dots, X_d]]$ carries the (p, X_1, \dots, X_d) -adic topology. Fix an index $i \in \{1, \dots, d\}$. Let T be a coordinate on $\mathbb{A}_{S^{\text{an}}}^1$. We are interested in the open affinoid subsets $V_m = \{|p^m|, |X_1^m|, \dots, |X_d^m| \leq |X_i| \neq 0, |X_i^m T| \leq 1\}$ of Z^{an} for $m \in \mathbb{Z}_{\geq 1}$. Note that the union of the V_m is the locus $V = \{|X_i| \neq 0\} \subseteq Z^{\text{an}}$. The ring $\mathcal{O}(V_m)$ has a ring of definition

$$R_m = A[[X_1, \dots, X_d]] \left\langle \frac{p^m}{X_i}, \frac{X_1^m}{X_i}, \dots, \frac{X_d^m}{X_i} \right\rangle \langle X_i^m T \rangle$$

(with the X_i -adic topology), and we have natural maps $R_{m+1} \rightarrow R_m$ for all m . Note that $\mathcal{O}(V_m) = R_m[1/X_i]$ and that $\mathcal{O}^+(V_m)$ is the integral closure of R_m in $\mathcal{O}(V_m)$. By Theorem A.5 (and its proof), in fact we have $\mathcal{O}^+(V_m) = \mathcal{O}(V_m)^\circ$ and $\mathcal{O}(V_m)^\circ$ is a finite R_m -algebra.

Lemma 5.4.2. *The image of R_{m+1} in R_m/X_i^n is finite for all m and n . More generally, if M is a finitely generated R_{m+1} -module, then the image of M in $M \otimes_{R_{m+1}} R_m/X_i^n$ is finite.*

Proof. Fix m and n . We start with the first assertion. It suffices to check that the kernel contains the ideal

$$I = \left(p^{mn}, X_1^{mn}, \dots, X_d^{mn}, \left(\frac{p^{m+1}}{X_i} \right)^{mn}, \left(\frac{X_1^{m+1}}{X_i} \right)^{mn}, \dots, \left(\frac{X_d^{m+1}}{X_i} \right)^{mn}, (X_i^{m+1}T)^n \right)$$

since it's straightforward to check that R_{m+1}/I is finite. It's also straightforward to check that the given generators of I are in the kernel. This finishes the (sketch of) proof of the first assertion. For the second, there is an integer $q \geq 0$ and a surjection $R_{m+1}^q \twoheadrightarrow M$ of R_{m+1} -modules, which gives us a commuting diagram

$$\begin{array}{ccc} R_{m+1}^q & \longrightarrow & R_m^q/X_i^n \\ \downarrow & & \downarrow \\ M & \longrightarrow & M \otimes_{R_{m+1}} R_m/X_i^n \end{array}$$

where the vertical maps are surjections, and the second assertion then follows from the first. \square

Proposition 5.4.3. *With notation as above, let $Y \rightarrow V = \{|X_i| \neq 0\} \subset Z^{\text{an}}$ be a finite morphism of adic spaces, and assume that Y is reduced. Then $\mathcal{O}^+(Y)$ is compact.*

Proof. Write Y_m for the pullback of V_m ; these form an increasing cover of Y consisting of open affinoids. Since $Y_m \rightarrow V_m$ is finite and $\mathcal{O}^+(V_m) = \mathcal{O}(V_m)^\circ$ we know that $\mathcal{O}^+(Y_m)$ is integral over $\mathcal{O}(V_m)^\circ$ (by definition of a finite morphism). Since $\mathcal{O}(Y_m)$ is reduced and $\mathcal{O}(V_m)^\circ$ is Nagata (it is finite over R_m , which is Nagata by the proof of Theorem A.5) it follows that $\mathcal{O}^+(Y_m)$ is finite over $\mathcal{O}(V_m)^\circ$. In particular, $\mathcal{O}^+(Y_m)$ is finite over R_m , and hence Noetherian. We may also deduce from Lemma A.2 that $\mathcal{O}^+(Y_m) = \mathcal{O}(Y_m)^\circ$.

For any n the map $\mathcal{O}^+(Y_{m+1}) \rightarrow \mathcal{O}^+(Y_m)/X_i^n$ factors through $\mathcal{O}^+(Y_{m+1}) \otimes_{R_{m+1}} R_m/X_i^n$ and hence the image is finite by Lemma 5.4.2. It follows that $\mathcal{O}^+(Y_{m+1})$, when equipped with the weak topology with respect to the map $\mathcal{O}^+(Y_{m+1}) \rightarrow \mathcal{O}^+(Y_m)$, is compact. We deduce that $\mathcal{O}^+(Y) = \varprojlim_m \mathcal{O}^+(Y_m)$ is compact, as desired. \square

Corollary 5.4.4. *Assume that $X \rightarrow Z^{\text{an}}$ is a finite morphism of adic spaces, with X reduced. Then $\mathcal{O}^+(X)$ is compact.*

Proof. For each $i \in \{1, \dots, d\}$ consider the locus $Z_i = \{|X_i| \neq 0\} \subseteq Z^{\text{an}}$ (i.e., what was previously denoted by V) and set $Z_0 = \{|p| \neq 0\}$. Let X_i , for $i \in \{0, \dots, d\}$, denote the corresponding pullbacks to X . Then we have a strict inclusion $\mathcal{O}^+(X) \subseteq \prod_{i=0}^d \mathcal{O}^+(X_i)$ with closed image and the $\mathcal{O}^+(X_i)$ are compact (for $i = 1, \dots, d$ this is Proposition 5.4.3 and for $i = 0$ this is [Bellaïche and Chenevier 2009, Lemma 7.2.11]; note that X_0 is nested in the terminology of that paper), so $\mathcal{O}^+(X)$ is compact as well. \square

We may then prove the main result of this section.

Theorem 5.4.5. *There is an n -dimensional continuous determinant D of $G_{F,S}$ with values in $\mathcal{O}^+(\mathcal{X}^{\text{red}})$ such that*

$$D(1 - X \text{Frob}_v) = P_v(X)$$

for all $v \notin S$.

Proof. Fix a collection $\{(\mathcal{U}, h)\}$ of slope data such that the $\mathcal{X}_{\mathcal{U},h}^{\text{red}}$ cover \mathcal{X}^{red} . We then have injections

$$\mathcal{O}^+(\mathcal{X}^{\text{red}}) \hookrightarrow \prod_{(\mathcal{U},h)} \mathcal{O}(\mathcal{X}_{\mathcal{U},h}^{\text{red}}) \hookrightarrow \prod_{(\mathcal{U},h)} \prod_{\mathfrak{m}} \mathcal{O}(\mathcal{X}_{\mathcal{U},h}^{\text{red}})/\mathfrak{m},$$

where the \mathfrak{m} range over all maximal ideals of $\mathcal{O}(\mathcal{X}_{\mathcal{U},h}^{\text{red}})$. Note that we have an injection $\mathcal{O}(\mathcal{X}_{\mathcal{U},h}^{\text{red}}) \hookrightarrow \prod_{\mathfrak{m}} \mathcal{O}(\mathcal{X}_{\mathcal{U},h}^{\text{red}})/\mathfrak{m}$ by Lemma A.1, so the second morphism really is an injection. Then note that $\mathcal{O}^+(\mathcal{X}^{\text{red}})$ is compact; since \mathcal{X}^{red} is finite over $\text{Spa}(\mathbb{Z}_p[[T_0]], \mathbb{Z}_p[[T_0]])^{\text{an}} \times \mathbb{A}^1$ this follows from Corollary 5.4.4. To see that $\mathbb{Z}_p[[T_0]]$ is of the form $A[[X_1, \dots, X_d]]$, we write $T_0 \cong T_0^{\text{tor}} \times T_0^{\text{free}}$, where T_0^{tor} is the torsion subgroup of T_0 (which is finitely generated) and $T_0^{\text{free}} \cong \mathbb{Z}_p^{\dim T_0}$ is a free complement (see, e.g., [Neukirch 1999, Proposition II.5.7]). Set $A = \mathbb{Z}_p[T_0^{\text{tor}}]$; then

$$\mathbb{Z}_p[[T_0]] \cong A \otimes_{\mathbb{Z}_p} \mathbb{Z}_p[[T_0^{\text{free}}]] \cong A[[X_1, \dots, X_d]].$$

Thus we may glue the determinants from Lemma 5.4.1 into the desired determinant using [Chenevier 2014, Example 2.32]. \square

6. The Coleman–Mazur eigencurve

In this section we give a short discussion of the special case of the Coleman–Mazur eigencurve and the relationship between our work and that of Andreatta, Iovita and Pilloni [Andreatta et al. 2018] and Liu, Wan and Xiao [Liu et al. 2017].

6.1. The case $G = \text{GL}_2/\mathbb{Q}$. Let us consider the special case $G = \text{GL}_2/\mathbb{Q}$. We begin by fixing choices of groups and Hecke algebras/operators. Let B be the upper triangular Borel, I the corresponding Iwahori and T the diagonal torus. We use the element $t = \begin{pmatrix} 1 & 0 \\ 0 & p \end{pmatrix} \in \Sigma^{\text{cpt}}$; the corresponding Hecke operator is the U_p -operator. We choose the tame level

$$K_1(N) = \left\{ g \in \text{GL}_2(\widehat{\mathbb{Z}}^p) \mid g \equiv \begin{pmatrix} * & * \\ 0 & 1 \end{pmatrix} \pmod{N} \right\},$$

with $N \in \mathbb{Z}_{\geq 5}$ prime to p . Put

$$\Delta_\ell = \begin{cases} \text{GL}_2(\mathbb{Q}_\ell) & \text{if } \ell \nmid N, \\ \text{GL}_2(\mathbb{Z}_\ell) & \text{if } \ell \mid N. \end{cases}$$

With these choices, everything else is determined and we use the notation of the main part of the paper.

In this case, overconvergent modular symbols were first constructed in [Stevens 1994], and the corresponding eigencurve was constructed and shown to agree with the Coleman–Mazur eigencurve

in [Bellaïche 2010]. Stevens and Bellaïche worked with the compactly supported cohomology H_c^1 , which admits a very explicit description in this case (it is given by the functor denoted by Symb in [Bellaïche 2010; Stevens 1994]). Ordinary cohomology was first considered in [Hansen 2017]. For ordinary cohomology, one has

$$H^*(K, \mathcal{D}_\kappa) = H^1(K, \mathcal{D}_\kappa)$$

for all weights κ , upon noting that the H^i vanish automatically for all $i \geq 2$ and that H^0 vanishes by (a simpler version of) the argument in the proof of [Chojcecki et al. 2017, Lemma 5.4]. A consequence is the following lemma:

Lemma 6.1.1. *Let $\kappa : T_0 \rightarrow R^\times$ be a weight, and assume that F_κ has a slope $\leq h$ -factorization for some $h \in \mathbb{Q}_{\geq 0}$ (slopes with respect to some multiplicative pseudouniformizer $\varpi \in R$). Then $H^1(K, \mathcal{D}_\kappa)_{\leq h}$ is a finite projective R -module and is compatible with arbitrary base change.*

Proof. Let $f : R \rightarrow S$ be a continuous homomorphism of complete Tate rings and equip S with a Banach–Tate \mathbb{Z}_p -algebra norm such that $f(\varpi)$ is a multiplicative pseudouniformizer. Put $\kappa_S = f \circ \kappa$. By the vanishing of the H^i for $i \neq 1$ the Tor-spectral sequence collapses and gives us that

$$\begin{aligned} H^1(K, \mathcal{D}_\kappa)_{\leq h} \otimes_R S &= H^1(K, \mathcal{D}_{\kappa_S})_{\leq h}, \\ \text{Tor}_1^R(H^1(K, \mathcal{D}_\kappa)_{\leq h}, S) &= 0. \end{aligned}$$

The first line is compatibility with base change. Putting $S = R/J$ for an arbitrary ideal $J \subseteq R$ (automatically closed) we see from the second line that $H^1(K, \mathcal{D}_\kappa)_{\leq h}$ is a finite flat R -module, and hence is finite projective. \square

If $\kappa : T_0 \rightarrow R^\times$ is a weight, we write $\kappa_i, i = 1, 2$, for the characters $\mathbb{Z}_p^\times \rightarrow R^\times$ defined by

$$\kappa \left(\begin{pmatrix} a & 0 \\ 0 & d \end{pmatrix} \right) = \kappa_1(a) \kappa_2(d)$$

and we identify \bar{N}_1 with $p\mathbb{Z}_p$ via $\begin{pmatrix} 1 & 0 \\ x & 1 \end{pmatrix} \mapsto x$. If we consider the eigenvariety $\mathcal{X}^{\text{rig}} = \mathcal{X}_G^{\text{rig}}$ constructed in [Hansen 2017], then it is equidimensional of dimension 2 by Proposition B.1 of that paper since $H^* = H^1$. This object is usually referred to as the “eigensurface”. If we instead do the eigenvariety construction over the part $\mathcal{W}_0^{\text{rig}}$ of weight space where $\kappa_2 = 1$, we obtain an eigenvariety that turns out to equal the Coleman–Mazur eigencurve; it is in particular reduced, equidimensional of dimension 1, and flat over $\mathcal{W}_0^{\text{rig}}$. Let us denote this eigenvariety by \mathcal{E}^{rig} ; the properties of \mathcal{E}^{rig} stated in the previous sentence are presumably well known to experts but we will give a brief sketch of the proofs below. To begin with, it is equidimensional of dimension 1 (by the same argument as above for \mathcal{X}^{rig}). For weights with $\kappa_2 = 1$ we conflate κ and κ_1 , and we may write the action on \mathcal{A}_κ explicitly as

$$(f.\gamma)(x) = \kappa(a + bx) f \left(\frac{c + dx}{a + bx} \right)$$

for $\gamma = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in M_2(\mathbb{Z}_p)$ such that $a \in \mathbb{Z}_p^\times$, $c \in p\mathbb{Z}_p$ (this defines the submonoid $\Delta_{p,0}$ of Δ_p generated by I and t), and $x \in p\mathbb{Z}_p$ [Hansen 2017, §2.2]. Using the anti-involution

$$\begin{pmatrix} a & b \\ c & d \end{pmatrix} \mapsto \begin{pmatrix} a & c/p \\ pb & d \end{pmatrix}$$

on $\Delta_{p,0}$ and rescaling f by p^{-1} in the argument to a function on \mathbb{Z}_p one sees this right action corresponds to the left action defined in [Stevens 1994]. To prove that \mathcal{E}^{rig} is reduced one uses a criterion of [Chenevier 2005, Proposition 3.9]; see [Bellaïche 2010, Theorem IV.2.1] for the analogous statement for the H_c^1 . Unfortunately, the notion of a “classical structure” in [Chenevier 2005] is based on the input data for the eigenvariety construction in [Buzzard 2007] and is therefore not directly applicable to the situation in [Hansen 2017]. Let us state a version of [Chenevier 2005, Proposition 3.9] applicable to the situation in [Hansen 2017, Definition 4.2.1]. We use the notation and terminology of [Hansen 2017, §4–5] freely; in particular we use the language of classical rigid geometry for this proposition.

Proposition 6.1.2. *Let $\mathfrak{D} = (\mathcal{W}, \mathcal{Z}, \mathcal{M}, T, \psi)$ be an eigenvariety datum. If (U, h) is a slope datum, assume that $\mathcal{M}(\mathcal{Z}_{U,h})$ is a projective $\mathcal{O}(U)$ -module. Assume moreover that there exists a very Zariski dense set $\mathcal{W}^{\text{cl}} \subseteq \mathcal{W}$ such that, if (U, h) is a slope datum, there is a Zariski open and dense subset $W_{U,h}$ of $\mathcal{W}^{\text{cl}} \cap U$ such that $\mathcal{M}(\mathcal{Z}_{U,h})_x$ is a semisimple $T[T^{-1}]$ -module. Here T is the parameter on $\mathbb{A}_{\mathcal{W}}^1$, which naturally acts invertibly on $\mathcal{M}(\mathcal{Z}_{U,h})$, and the set \mathcal{W}^{cl} is given the Zariski topology. Then the eigenvariety $\mathcal{X}_{\mathfrak{D}}$ is reduced.*

The proof is virtually identical to that of [Chenevier 2005, Proposition 3.9]; we omit it. Using this, one proves that \mathcal{E}^{rig} is reduced in the same way as in the proof of [Bellaïche 2010, Theorem IV.2.1(i)], using the control theorem of Stevens (see [Hansen 2017, Theorem 3.2.5]); recall that our Hecke algebra $\mathbb{T}(\Delta^p, K^p)$ contains no Hecke operators at primes dividing N . That \mathcal{E}^{rig} is equal to the Coleman–Mazur eigencurve is then proved using the control theorems of Coleman and Stevens and the Eichler–Shimura isomorphism together with [Hansen 2017, Theorem 5.1.2] (this type of argument is well known to experts; see for example the proof of [Bellaïche 2010, Theorem IV.2.1(i)]). The argument for flatness will be given below. This finishes our review of the basic properties of \mathcal{E}^{rig} .

Let us now return to the constructions of this paper. Our eigenvariety construction gives an extension \mathcal{E} of \mathcal{E}^{rig} defined over the locus $\mathcal{W}_0 \subseteq \mathcal{W}$ where $\kappa_2 = 1$. Another such extension \mathcal{E}' was constructed by Andreatta, Iovita and Pilloni [Andreatta et al. 2018]. Note that \mathcal{W}_0 is naturally the analytic locus of the formal weight space \mathfrak{W}_0 with $\kappa_2 = 1$. We have

$$\mathfrak{W}_0 \cong \text{Spa}(\mathbb{Z}_p[[\mathbb{Z}_p^\times]], \mathbb{Z}_p[[\mathbb{Z}_p^\times]]).$$

When $p \neq 2$, $\mathbb{Z}_p[[\mathbb{Z}_p^\times]]$ is a regular ring. When $p = 2$ this is no longer the case; we have $\mathbb{Z}_2[[\mathbb{Z}_2^\times]] \cong \mathbb{Z}_2[\mathbb{Z}/2][[X]]$ and $\mathbb{Z}_2[\mathbb{Z}/2]$ is not regular. We will instead work over the normalization A of $\mathbb{Z}_2[[\mathbb{Z}_2^\times]]$, which is isomorphic to $\mathbb{Z}_2[[X]] \times \mathbb{Z}_2[[X]]$. The normalization map

$$\mathbb{Z}_2[\mathbb{Z}/2][[X]] \rightarrow \mathbb{Z}_2[[X]] \times \mathbb{Z}_2[[X]]$$

is explicitly given by

$$\sum_{n \geq 0} (a_n + b_n g) X^n \mapsto \left(\sum_{n \geq 0} (a_n + b_n) X^n, \sum_{n \geq 0} (a_n - b_n) X^n \right),$$

where $a_n, b_n \in \mathbb{Z}_2$ and $g \in \mathbb{Z}/2$ is the nontrivial element. Using this map we get a weight into A , and we let $\tilde{\mathcal{W}}_0 = \text{Spa}(A, A)$ and put $\tilde{\mathcal{W}}_0 = \tilde{\mathcal{W}}_0^{\text{an}}$. We remark that $\tilde{\mathcal{W}}_0^{\text{rig}} \cong \mathcal{W}_0^{\text{rig}}$ canonically via the normalization map. To make our notation uniform, we set $\tilde{\mathcal{W}}_0 = \mathcal{W}_0$ when $p \neq 2$. We may perform our construction over $\tilde{\mathcal{W}}_0$, and one may pull back the Banach modules that are used to construct the eigencurve in [Andreatta et al. 2018] to $\tilde{\mathcal{W}}_0$, and construct the eigencurve over $\tilde{\mathcal{W}}_0$ instead. Let us denote the corresponding eigenvarieties by $\tilde{\mathcal{E}}$ and $\tilde{\mathcal{E}}'$, though we hasten to remark that these should *not* be thought of as normalizations of \mathcal{E} and \mathcal{E}' .

Lemma 6.1.3. *Let $\mathcal{U} \subset \tilde{\mathcal{W}}_0$ be a connected open affinoid subset such that $\mathcal{O}(\mathcal{U})$ is a Tate ring. Then $\mathcal{O}(\mathcal{U})$ is a Dedekind domain.*

Proof. We first show that $\mathcal{O}(\mathcal{U})$ is regular of Krull dimension 1. The connected components of $\tilde{\mathcal{W}}_0$ are isomorphic to $\text{Spa}(\mathbb{Z}_p[[X]], \mathbb{Z}_p[[X]])$ so we may take \mathcal{U} to be an open affinoid subset of $\text{Spa}(\mathbb{Z}_p[[X]], \mathbb{Z}_p[[X]])$. Let \mathfrak{q} be a maximal ideal of $\mathcal{O}(\mathcal{U})$ and let \mathfrak{p} be its preimage in $A = \mathbb{Z}_p[[X]]$. By Proposition A.15 the natural map $A_{\mathfrak{p}} \rightarrow \mathcal{O}(\mathcal{U})_{\mathfrak{q}}$ induces an isomorphism on completions. By Lemma A.13 \mathfrak{p} defines a closed point of $\text{Spec } A \setminus \{(p, X)\}$. Since A is a regular local ring of dimension 2, it follows that $A_{\mathfrak{p}}$ is a regular local ring of dimension 1, and hence the same is true for $\mathcal{O}(\mathcal{U})_{\mathfrak{q}}$ (since if R is a Noetherian local ring with completion \hat{R} , then R is regular if and only if \hat{R} is regular, and $\dim R = \dim \hat{R}$). It follows that $\mathcal{O}(\mathcal{U})$ is a product of regular integral domains of dimension 1. Since \mathcal{U} is connected $\mathcal{O}(\mathcal{U})$ does not contain any nontrivial idempotents, so $\mathcal{O}(\mathcal{U})$ is an integral domain. \square

Let (\mathcal{U}, h) be a connected slope datum for $\tilde{\mathcal{E}}$ (by which we mean a slope datum for the construction that produces $\tilde{\mathcal{E}}$ such that \mathcal{U} is connected; we will use the terminology “(connected) slope datum for $\tilde{\mathcal{E}}$ ” similarly). Then $\mathcal{O}(\tilde{\mathcal{E}}_{\mathcal{U}, h})$ is, by definition, an $\mathcal{O}(\mathcal{U})$ -submodule of $\text{End}_{\mathcal{O}(\mathcal{U})}(H^1(K, \mathcal{D}_{\mathcal{U}})_{\leq h})$. The latter is projective by Lemma 6.1.1, so the former is also projective since $\mathcal{O}(\mathcal{U})$ is Dedekind. Thus the natural map $\tilde{\mathcal{E}}_{\mathcal{U}, h} \rightarrow \mathcal{U}$ is finite flat, and hence $\tilde{\mathcal{E}} \rightarrow \tilde{\mathcal{W}}_0$ is flat. The same applies to $\tilde{\mathcal{E}}'$.

Now let (\mathcal{U}, h) be a slope datum for $\tilde{\mathcal{E}}$ and $\tilde{\mathcal{E}}'$. Let $(\mathcal{U}_i)_{i \in I}$ be an open affinoid cover of \mathcal{U}^{rig} . Then the natural map $\mathcal{O}(\mathcal{U}) \rightarrow \prod_{i \in I} \mathcal{O}(\mathcal{U}_i)$ is an injection (since $\mathcal{U} \setminus \mathcal{U}^{\text{rig}}$ does not contain any open subset of \mathcal{U}) so tensoring with the finite projective $\mathcal{O}(\mathcal{U})$ -module $\mathcal{O}(\tilde{\mathcal{E}}_{\mathcal{U}, h})$ we get an injection

$$\mathcal{O}(\tilde{\mathcal{E}}_{\mathcal{U}, h}) \hookrightarrow \left(\prod_{i \in I} \mathcal{O}(\mathcal{U}_i) \right) \otimes_{\mathcal{O}(\mathcal{U})} \mathcal{O}(\tilde{\mathcal{E}}_{\mathcal{U}, h}) \cong \prod_{i \in I} \mathcal{O}(\tilde{\mathcal{E}}_{\mathcal{U}_i, h}),$$

which in particular shows that $\mathcal{O}(\tilde{\mathcal{E}}_{\mathcal{U}, h})$ is reduced. The image of $\mathcal{O}(\tilde{\mathcal{E}}_{\mathcal{U}, h})$ inside $\prod_{i \in I} \mathcal{O}(\tilde{\mathcal{E}}_{\mathcal{U}_i, h})$ is equal to the $\mathcal{O}(\mathcal{U})$ -span of the image of $\mathbb{T}(\Delta^p, K^p)[U_p]$ in $\prod_{i \in I} \mathcal{O}(\tilde{\mathcal{E}}_{\mathcal{U}_i, h})$. The same holds replacing $\tilde{\mathcal{E}}$ by $\tilde{\mathcal{E}}'$, so since we have canonical isomorphisms

$$\mathcal{O}(\tilde{\mathcal{E}}_{\mathcal{U}_i, h}) \cong \mathcal{O}(\tilde{\mathcal{E}}'_{\mathcal{U}_i, h}),$$

we obtain a canonical isomorphism $\mathcal{O}(\tilde{\mathcal{E}}_{\mathcal{U},h}) \cong \mathcal{O}(\tilde{\mathcal{E}}'_{\mathcal{U},h})$, compatible with the way the eigencurves are built. As a result we have a canonical isomorphism $\tilde{\mathcal{E}} \cong \tilde{\mathcal{E}}'$ extending the canonical isomorphism $\tilde{\mathcal{E}}^{\text{rig}} \cong (\tilde{\mathcal{E}}')^{\text{rig}}$. We summarize the discussion above in the following theorem:

Theorem 6.1.4. *The eigenvariety $\tilde{\mathcal{E}}$ is reduced and flat over $\tilde{\mathcal{W}}_0$. Moreover, it is canonically isomorphic to the eigencurve $\tilde{\mathcal{E}}'$ constructed by Andreatta, Iovita and Pilloni [Andreatta et al. 2018].*

Remark 6.1.5. In fact, it is possible to show that \mathcal{E} and \mathcal{E}' are isomorphic for all p (i.e., including $p = 2$), using the interpolation theorem [Johansson and Newton 2017, Theorem 3.2.1], since both \mathcal{E} and \mathcal{E}' are reduced with Zariski dense sets of classical points which naturally match up.

Fix a character $\eta : (\mathbb{Z}/q)^\times \rightarrow \mathbb{F}_p^\times$ (recall that $q = 4$ if $p = 2$ and $q = p$ otherwise). We have a natural isomorphism $\mathbb{Z}_p^\times \cong (\mathbb{Z}/q)^\times \times (1 + q\mathbb{Z}_p)$ defined by $z \mapsto (\bar{z}, z/\omega(\bar{z}))$, where an overline denotes reduction modulo q and ω denotes the Teichmüller lift. Let us write $\langle z \rangle := z/\omega(\bar{z})$. Then we may define a character $\kappa_\eta : \mathbb{Z}_p^\times \rightarrow \mathbb{Z}_p[[X]]^\times$ by

$$\kappa_\eta(z) = \omega(\eta(\bar{z})) \sum_{n=0}^{\infty} \binom{p^{-1} \log \langle z \rangle}{n} X^n.$$

We let $\bar{\kappa}_\eta$ denote its reduction modulo p . This is a character $\mathbb{Z}_p^\times \rightarrow \mathbb{F}_p((X))^\times$ which we may think of as a character $T_0 \rightarrow \mathbb{F}_p((X))^\times$ with $\kappa_2 = 1$. We remark that if $p = 2$ then η , and hence $\bar{\kappa}_\eta$, is unique.

Corollary 6.1.6. *There are infinitely many (nonordinary) finite-slope U_p -eigenvectors in $H^1(K, \mathcal{D}_{\bar{\kappa}_\eta})$.*

Proof. By Corollary 4.2.3, its analogue for $\tilde{\mathcal{E}}'$ (which is simpler, and is essentially [Buzzard 2007, Lemma 5.9]) and Theorem 6.1.4 we see that $H^1(K, \mathcal{D}_{\bar{\kappa}_\eta})$ and the module $\bar{M}_{\bar{\kappa}_\eta}^\dagger(N)$ of overconvergent modular forms of weight $\bar{\kappa}_\eta$ and tame level N constructed in [Andreatta et al. 2018] contain the same finite-slope systems of Hecke eigenvalues. By [Bergdall and Pollack 2016, Corollary A.1], $\bar{M}_{\bar{\kappa}_\eta}^\dagger(N)$ has infinitely many finite-slope U_p -eigenvectors, so we are done. \square

Remark 6.1.7. It is possible to prove Corollary 6.1.6 directly from [Bergdall and Pollack 2016, Theorem A] (using the observation in the remark following [loc. cit., Corollary A.2]) without any reference to [Andreatta et al. 2018].

6.2. Estimates for the Newton polygon of U_p . In this and the following section we give a short proof of the estimates obtained in [Liu et al. 2017, Theorem 3.16] for the Newton polygon of U_p acting on spaces of overconvergent automorphic forms for a definite quaternion algebra over \mathbb{Q} .

We fix an odd prime p and assume that we are in the setting of Section 3.3. Suppose that $\bar{N}_1 \cong \mathbb{Z}_p$ and fix a topological generator \bar{n} . Let f be a norm-decreasing R -linear map

$$f : \bigoplus_{i=1}^t \mathcal{D}_\kappa^r \rightarrow \bigoplus_{i=1}^t \mathcal{D}_\kappa^{r^{1/p}}$$

and recall that we have a compact inclusion

$$\iota : \bigoplus_{i=1}^t \mathcal{D}_\kappa^{r^{1/p}} \hookrightarrow \bigoplus_{i=1}^t \mathcal{D}_\kappa^r.$$

We set $U = \iota \circ f$. Then U is a compact endomorphism of $M = \bigoplus_{i=1}^t \mathcal{D}_\kappa^r$. Recall that we have a potential ON-basis for \mathcal{D}_κ^r given by the elements $e_{r,\alpha} := \varpi^{-n(r,\varpi,\alpha)} \bar{\mathfrak{n}}^\alpha$ for $\alpha \in \mathbb{Z}_{\geq 0}$.

We consider the potential ON-basis for M given by

$$e_{r,0}^1 = (e_{r,0}, 0, \dots, 0), \quad \dots, \quad e_{r,0}^t = (0, \dots, 0, e_{r,0}), \quad e_{r,1}^1 = (e_{r,1}, 0, \dots, 0), \quad \dots$$

Lemma 6.2.1. *Assume that there is no $x \in R$ with $1 < |x| < |\varpi|^{-1}$. Then U maps $e_{r,\alpha}^i$ to a sum $\sum_{j,\beta} a_{\beta}^j e_{r,\beta}^j$ with*

$$|a_{\beta}^j| \leq |\varpi|^{n(r,\varpi,\beta) - n(r^{1/p},\varpi,\beta)}.$$

If we define

$$\lambda(0) = 0, \quad \lambda(i+1) = \lambda(i) + n(r, \varpi, \lfloor i/t \rfloor) - n(r^{1/p}, \varpi, \lfloor i/t \rfloor),$$

the Fredholm series

$$\det(1 - TU|M) = \sum_{n \geq 0} c_n T^n \in R\{\{T\}\}$$

satisfies $|c_n| \leq |\varpi|^{\lambda(n)}$.

Proof. We first prove the estimate on the matrix coefficients of U . Apply f to $e_{r,\alpha}^i$. We get a sum $\sum_{j,\beta} b_{\beta}^j e_{r^{1/p},\beta}^j$, and the fact that f is norm-decreasing is equivalent to

$$|b_{\beta}^j| \leq |\varpi|^{n(r^{1/p},\varpi,\beta) - n(r,\varpi,\alpha)} r^{|\alpha| - |\beta|/p}$$

for all j, β . Since $|\varpi| < |\varpi|^{-n(r,\varpi,\alpha)} r^{|\alpha|}$, $|\varpi|^{-n(r^{1/p},\varpi,\beta)} r^{|\beta|/p} \leq 1$ by construction, we deduce that $|b_{\beta}^j| < |\varpi|^{-1}$ for all j, β . By our assumption on R , it follows that $|b_{\beta}^j| \leq 1$ for all j, β . We then have

$$e_{r^{1/p},\beta}^j = \varpi^{-n(r^{1/p},\varpi,\beta)} \bar{\mathfrak{n}}^\beta = \varpi^{n(r,\varpi,\beta) - n(r^{1/p},\varpi,\beta)} e_{r,\beta}^j$$

so we conclude that

$$U e_{r,\alpha}^i = \sum_{j,\beta} a_{\beta}^j e_{r,\beta}^j,$$

where $a_{\beta}^j = \varpi^{n(r,\varpi,\beta) - n(r^{1/p},\varpi,\beta)} b_{\beta}^j$, and therefore

$$|a_{\beta}^j| \leq |\varpi|^{n(r,\varpi,\beta) - n(r^{1/p},\varpi,\beta)}.$$

In other words, the i -th row of the matrix for U (we begin indexing rows at $i = 0$) has entries with norm $\leq |\varpi|^{n(r,\varpi,\lfloor i/t \rfloor) - n(r^{1/p},\varpi,\lfloor i/t \rfloor)}$. We deduce immediately that $|c_n| \leq |\varpi|^{\lambda(n)}$, since c_n is an alternating sum of products of matrix entries coming from n distinct rows [Serre 1962, Proposition 7], and each of these products has norm $\leq |\varpi|^{\lambda(n)}$. \square

6.3. Definite quaternion algebras over \mathbb{Q} . As an application of Lemma 6.2.1, we give a new proof of [Liu et al. 2017, Theorem 3.16]. In this section we assume that p is odd. We need to set up things so that we can apply the machinery of Sections 3.3 and 4. Let D/\mathbb{Q} be a definite quaternion algebra, split at p , and let G be the reductive group over \mathbb{Q} defined by $G(R) = (D \otimes_{\mathbb{Q}} R)^{\times}$, for \mathbb{Q} -algebras R . We fix an isomorphism $D_p \cong M_2(\mathbb{Q}_p)$ and henceforth identify D_p with $M_2(\mathbb{Q}_p)$ via this isomorphism. Let $G_{\mathbb{Z}_p}$ be

the \mathbb{Z}_p -model for G given by $G_{\mathbb{Z}_p}(R) = (M_2(\mathbb{Z}_p) \otimes_{\mathbb{Z}_p} R)^\times$ for \mathbb{Z}_p -algebras R . We let B be the upper triangular Borel in $G_{\mathbb{Z}_p}$ and let T be the diagonal maximal torus.

Fix a tame level $K^p = \prod_{l \neq p} K_l$ with $K_l \subset G(\mathbb{Q}_l)$ compact open, let $K = K^p I$, and assume that K is neat.

Fix a character $\eta : \mathbb{Z}_p^\times \rightarrow \mathbb{F}_p^\times$ and let $\Lambda = \mathbb{Z}_p[[\mathbb{Z}_p^\times]]$. We have an induced map $\pi_\eta : \Lambda \rightarrow \mathbb{F}_p$ and we denote its kernel by \mathfrak{m}_η . We write Λ_η for the localization $\Lambda_{\mathfrak{m}_\eta}$. We have a universal character

$$[\cdot]_\eta : \mathbb{Z}_p^\times \rightarrow \Lambda_\eta^\times.$$

We give the complete local ring Λ_η the \mathfrak{m}_η -adic topology. Fixing a topological generator γ of $1 + p\mathbb{Z}_p$ gives an isomorphism

$$\mathbb{Z}_p[[X]] \rightarrow \Lambda_\eta, \quad X \mapsto [\gamma]_\eta - 1.$$

Let $\mathfrak{W}_\eta = \text{Spa}(\Lambda_\eta, \Lambda_\eta)$, denote its analytic locus by \mathcal{W}_η and let $\mathcal{U}_1 \subset \mathcal{W}_\eta$ be the rational subdomain of \mathfrak{W}_η

$$\mathcal{U}_1 = \{|p| \leq |X| \neq 0\}.$$

Pulling back \mathcal{U}_1 to the open unit disc $\mathcal{W}_\eta^{\text{rig}}$ gives the “boundary annulus” $|X|_p \geq p^{-1}$.

We let $R_\eta = \mathcal{O}(\mathcal{U}_1)$. More explicitly, $R_\eta = R_\eta^\circ[[1/X]]$, where R_η° is a ring of definition for R_η , given by the X -adic completion of $\mathbb{Z}_p[[X]][p/X]$, with the X -adic topology.

Even more explicitly, we can describe the elements of R_η° as formal power series

$$\left\{ \sum_{n \in \mathbb{Z}} a_n X^n \mid a_n \in \mathbb{Z}_p, |a_n|_p p^{-n} \leq 1, |a_n|_p p^{-n} \rightarrow 0 \text{ as } n \rightarrow -\infty \right\}$$

X is a topologically nilpotent unit in R_η and so equipping R_η with the norm

$$|r| = \inf\{p^{-n} \mid r \in X^n R_\eta^\circ, n \in \mathbb{Z}\}$$

makes R_η into a Banach–Tate \mathbb{Z}_p -algebra. This norm has the explicit description

$$\left| \sum_{n \in \mathbb{Z}} a_n X^n \right| = \sup\{|a_n|_p p^{-n}\}. \quad (6.3.1)$$

Note that if $r \in \Lambda_\eta$, we have $|r| = \inf\{p^{-n} \mid r \in \mathfrak{m}_\eta^n, n \in \mathbb{Z}\}$.

We now define a continuous character

$$\kappa_\eta : T_0 \rightarrow R_\eta^\times$$

by

$$\kappa_\eta \begin{pmatrix} a & 0 \\ 0 & d \end{pmatrix} = [a]_\eta.$$

Lemma 6.3.1. *The norm we have defined on R_η is adapted to κ_η . Moreover, for $t \in T_1$ we have $|\kappa_\eta(t) - 1| \leq 1/p$.*

Proof. If $t \in T_1$ we have $\kappa(t) - 1 = (1 + X)^\alpha - 1 = \sum_{n \geq 1} \binom{\alpha}{n} X^n$ for some $\alpha \in \mathbb{Z}_p$. So $|\kappa_\eta(t) - 1| \leq 1/p$. \square

We can now apply the theory of Section 4 to the space of overconvergent automorphic forms $H^0(K, \mathcal{D}_{\kappa_\eta}^{1/p})$. Note that we have the concrete description

$$H^0(K, \mathcal{D}_{\kappa_\eta}^{1/p}) = \{f : D^\times \backslash (D \otimes \mathbb{A}^f)^\times / K^p \rightarrow \mathcal{D}_{\kappa_\eta}^{1/p} \mid f(gk) = k^{-1}f(g) \text{ for } k \in I\}$$

and $H^0(K, \mathcal{D}_{\kappa_\eta}^{1/p})$ is a Banach R_η -module with norm $|f| = \sup_{g \in (D \otimes \mathbb{A}^f)^\times} \|f(g)\|_{1/p}$.

In particular, we consider the action on $H^0(K, \mathcal{D}_{\kappa_\eta}^{1/p})$ of the Hecke operator U_{κ_η} attached to the element $\begin{pmatrix} 1 & 0 \\ 0 & p \end{pmatrix} \in \Sigma^{\text{cpt}}$. As a simple consequence of our results, we obtain the following theorem, which is essentially due to Liu–Wan–Xiao — compare with [Liu et al. 2017, Theorem 3.16, §5.4].

Theorem 6.3.2. *The Hecke operator U_{κ_η} is compact. Consider the Fredholm series*

$$F_{\kappa_\eta}(T) = \sum_{n \geq 0} c_n T^n = \det(1 - T U_{\kappa_\eta} | H^0(K, \mathcal{D}_{\kappa_\eta}^{1/p})).$$

Let $t = |D^\times \backslash (D \otimes \mathbb{A}^f)^\times / K|$. We have $c_n \in \Lambda_\eta$ and moreover we have

$$c_n \in \mathfrak{m}_\eta^{\lambda(n)} \quad \text{for } n \in \mathbb{Z}_{\geq 0},$$

where $\lambda(0) = 0, \lambda(1), \dots$ is a sequence of integers determined by

$$\lambda(0) = 0, \quad \lambda(i+1) = \lambda(i) + \lfloor i/t \rfloor - \lfloor i/pt \rfloor.$$

Proof. Compactness of U_{κ_η} follows from Corollary 3.3.10. The fact that $c_n \in \Lambda_\eta$ follows from Corollary 4.1.5, since $F_{\kappa_\eta}(T)$ extends to a Fredholm series over \mathcal{W}_η , and $\mathcal{O}(\mathcal{W}_\eta) = \Lambda_\eta$.

The rest of the theorem follows from Lemma 6.2.1 (note that the norm on R_η satisfies the assumption of that lemma), using the fact that if we choose representatives g_1, \dots, g_t for the double cosets $D^\times \backslash (D \otimes \mathbb{A}^f)^\times / K$ and $r \in [p^{-1}, 1)$ we have an isomorphism of potentially ON-able R_η -modules:

$$H^0(K, \mathcal{D}_{\kappa_\eta}^r) \cong \bigoplus_{i=1}^t \mathcal{D}_{\kappa_\eta}^r, \quad f \mapsto (f(g_i))_{i=1}^t.$$

We take $\varpi = X$, and compute that

$$\begin{aligned} n(p^{-1}, X, \lfloor i/t \rfloor) &= \lfloor i/t \rfloor, \\ n(p^{-1/p}, X, \lfloor i/t \rfloor) &= \lfloor 1/p \lfloor i/t \rfloor \rfloor = \lfloor i/pt \rfloor. \end{aligned}$$

□

As in Section 6.1, our eigenvariety construction, applied to the modules $H^0(K, \mathcal{D}_\kappa^r)$ with $\kappa_2 = 1$, gives an eigenvariety \mathcal{E}_η which is flat over \mathcal{W}_η . The open subspace $\mathcal{E}_\eta^{\text{rig}}$ defined by $|p| \neq 0$ is the eigenvariety constructed in [Buzzard 2004].

We end this section with our interpretation of [Liu et al. 2017, Theorems 1.3 and 1.5]. First we need some extra notation. For $m \leq n$ positive integers we define

$$\mathcal{U}_{m/n} = \{|p^m| \leq |X^n| \neq 0\} \subseteq \mathcal{W}_\eta.$$

We set $\mathcal{U} = \mathcal{U}_1$.

For a real number $\alpha \in (0, 1]$ we denote by $\mathcal{W}_\eta^{>p^{-\alpha}}$ the open subspace of \mathcal{W}_η obtained as the union of open affinoids

$$\mathcal{W}_\eta^{>p^{-\alpha}} = \bigcup_{m/n < \alpha} \mathcal{U}_{m/n}.$$

Note that X is a topologically nilpotent unit in $\mathcal{O}(\mathcal{U}_{m/n})$ for all m, n . We denote the pullback of \mathcal{E}_η to $\mathcal{W}_\eta^{>p^{-\alpha}}$ by $\mathcal{E}_\eta^{>p^{-\alpha}}$. The eigenvariety \mathcal{E}_η comes equipped with a map to the spectral variety $Z(F_{\kappa_\eta})$, and therefore it comes equipped with a map to \mathbb{A}_L^1 . For $h = m/n \in \mathbb{Q}$ with $m \in \mathbb{Z}$ and $n \in \mathbb{Z}_{\geq 1}$ we define an affinoid subspace $\mathbb{B}_{\mathcal{U},=h} \subset \mathbb{A}_L^1$ by

$$\mathbb{B}_{\mathcal{U},=h} = \{|T^n| = |X^{-m}|\}.$$

Similarly, if $h = m/n \leq h' = m'/n$ are rational numbers, we define

$$\mathbb{B}_{\mathcal{U},[h,h']} = \{|X^{-m}| \leq |T^n| \leq |X^{-m'}|\}.$$

Lemma 6.3.3. *Let L/\mathbb{Q}_p be a finite extension and let $x \in L$ with $|x|_p = p^{-\alpha}$, where $0 < \alpha \leq 1$. Consider the closed immersion $\iota : \mathrm{Spa}(L, \mathcal{O}_L) \hookrightarrow \mathcal{U}$ induced by the continuous \mathbb{Z}_p -algebra map $R_\eta \rightarrow L$ sending X to x . Let $\mathbb{B}_{x,=h}$ be the pullback of $\mathbb{B}_{\mathcal{U},=h}$ along ι . Then $\mathbb{B}_{x,=h} \hookrightarrow \mathbb{A}_L^1$ is the affinoid open defined by*

$$\mathbb{B}_{x,=h} = \{|T|_p = p^{-\alpha h}\}.$$

Proof. The affinoid $\mathbb{B}_{x,=h}$ is given by $\{|T^n| = |x^{-m}| = |p^{\alpha m}|\} \subset \mathbb{A}_L^1$. □

Theorem 6.3.4 [Liu et al. 2017]. *The space $\mathcal{E}_\eta^{>p^{-1}}$ is a disjoint countable union of adic spaces finite and flat over $\mathcal{W}_\eta^{>p^{-1}}$.*

Moreover, there is an explicit α depending only on K^p , with $0 < \alpha < 1$, such that

$$\mathcal{E}_\eta^{>p^{-\alpha}} = \bigsqcup_{i \geq 0} \mathcal{X}_{\eta,i}$$

with $\mathcal{X}_{\eta,i}$ finite flat over $\mathcal{W}_\eta^{>p^{-\alpha}}$ and each piece $\mathcal{X}_{\eta,i}$ of the eigenvariety has constant slope, in the sense that each map $\mathcal{X}_{\eta,i} \rightarrow \mathbb{A}_L^1$ factors through the affinoid subspace $\mathbb{B}_{\mathcal{U},=h_i} \subset \mathbb{A}_L^1$ for some $h_i \in \mathbb{Q}_{\geq 0}$. In particular, if we measure slopes on $\mathcal{X}_{\eta,i}^{\mathrm{rig}}$ with the usual p -adic valuation, then the slope of a point in $\mathcal{X}_{\eta,i}^{\mathrm{rig}}$ is given by $h_i v_p(T)$.

Proof. This follows from Theorems 1.3, 1.5 and Remark 3.25 of [Liu et al. 2017]. Remark 3.25 shows that, after restricting to $\mathcal{W}_\eta^{>p^{-1}}$, the Fredholm series F_{κ_η} factorizes as a countable product of multiplicative polynomials $\prod_{i \geq 0} P_i$, with each finite product $\prod_{i \geq 0}^N P_i$ a factor in a slope factorization over every affinoid subspace of $\mathcal{W}_\eta^{>p^{-1}}$. This establishes the claim about $\mathcal{E}_\eta^{>p^{-1}}$. Theorem 1.5 shows moreover that (for some explicit α) the restriction of F_{κ_η} to $\mathcal{W}_\eta^{>p^{-\alpha}}$ factorizes as $\prod_{i \geq 0} Q_i$, such that the specialization of Q_i at every classical rigid analytic point of $\mathcal{W}_\eta^{>p^{-\alpha}}$ has constant slope equal to h_i for some $h_i \in \mathbb{Q}_{\geq 0}$ (independent of the specialization). We obtain a decomposition of $Z(F_{\kappa_\eta})$ as a disjoint union of spaces Z_i , finite flat over $\mathcal{W}_\eta^{>p^{-\alpha}}$, such that every classical rigid analytic point of Z_i is contained in $\mathbb{B}_{\mathcal{U},=h_i}$. The space $\mathcal{X}_{\eta,i}$ is defined to be the inverse image of Z_i in $\mathcal{E}_\eta^{>p^{-\alpha}}$. It now remains to show that every point

of Z_i is contained in $\mathbb{B}_{\mathcal{U},=h_i}$. First we check this for rank-1 points: a rank-1 point of Z_i which is not in $\mathbb{B}_{\mathcal{U},=h_i}$ is contained in $\mathbb{B}_{\mathcal{U},[h,h']}$ for some interval $[h, h']$ which does not contain h_i . But then $\mathbb{B}_{\mathcal{U},[h,h']} \cap Z_i$ is a nonempty open subset in Z_i which contains no classical rigid analytic point, which is impossible since Z_i is finite flat over $\mathcal{W}_\eta^{>p^{-\alpha}}$ (in particular the map $Z_i \rightarrow \mathcal{W}_\eta^{>p^{-\alpha}}$ is open). Let V be an affinoid open in $\mathcal{W}_\eta^{>p^{-\alpha}}$. Then $Z_i|_V \cap \mathbb{B}_{\mathcal{U},=h_i}$ is an affinoid open in $Z_i|_V$ such that the complement contains no rank-1 point. We have $Z_i|_V = \text{Spa}(A, A^\circ)$ for some Tate ring A with a Noetherian ring of definition. So [Huber 1993, Lemma 3.4] (see the proof of Corollary 4.2 of that paper) implies that rank-1 points are dense in the constructible topology of $\text{Spa}(A, A^\circ)$ and we deduce that $Z_i|_V \cap \mathbb{B}_{\mathcal{U},=h_i} = Z_i|_V$. Therefore Z_i is contained in $\mathbb{B}_{\mathcal{U},=h_i}$, as desired. The final sentence of the theorem follows from Lemma 6.3.3. \square

Note that [Liu et al. 2017] proves moreover that the slopes appearing in the above theorem (with multiplicities) are given by a finite union of arithmetic progressions.

Appendix: Some algebraic properties of Tate rings

In this section we prove some properties of the kinds of Tate rings and adic spaces that we need. We start with a ring-theoretic lemma.

Lemma A.1. *Let R be a complete Tate ring with a Noetherian ring of definition R_0 . Then R is Jacobson.*

Proof. Let $\varpi \in R_0$ be a topologically nilpotent unit in R . Because R_0 is a Zariski ring when equipped with its ϖ -adic topology (since it is complete), $\text{Spec } R_0 \setminus \{\varpi = 0\}$ is a Jacobson scheme by [EGA IV₃ 1966, (10.5.7)]. But $\text{Spec } R_0 \setminus \{\varpi = 0\} = \text{Spec } R$, so R is Jacobson as desired. \square

We record another simple lemma that will prove to be useful.

Lemma A.2. *Let R be a complete Tate ring with a Noetherian ring of definition R_0 . If $S \subseteq R$ is an open and bounded subring (i.e., a ring of definition) containing R_0 , then S is a finitely generated R_0 -module, hence Noetherian, and integral over R_0 . Moreover, R° is the integral closure of R_0 in R .*

Proof. Pick a topologically nilpotent unit $\varpi \in R$ contained in R_0 . Since S is bounded we have $S \subseteq \varpi^{-N} R_0$ for some N , and hence S is an R_0 -submodule of the cyclic R_0 -module $\varpi^{-N} R_0$. The lemma now follows since R_0 is Noetherian. For the last assertion, first note that the integral closure is contained in R° . Since R° is the union of all open and bounded subrings and any two open bounded subrings are contained in a third, the assertion follows from the first part. \square

One consequence of the above lemma is a version for Tate rings (with our Noetherian hypothesis) of [Bosch et al. 1984, 6.3.4/Proposition 1]:

Lemma A.3. *Let R be a complete Tate ring with a Noetherian ring of definition. Let S be a finite R -algebra, equipped with the natural R -module topology. Then S is a complete Tate ring with a Noetherian ring of definition.*

Moreover, the integral closure of R° in S is equal to S° . In particular, the morphism $(R, R^\circ) \rightarrow (S, S^\circ)$ is a finite morphism of affinoid rings (see [Huber 1996, 1.4.2]) and $\text{Spa}(S, S^\circ) \rightarrow \text{Spa}(R, R^\circ)$ is a finite morphism of adic spaces (see [loc. cit., 1.4.4]).

Proof. We let R_0 denote a Noetherian ring of definition for R , let ϖ denote a topologically nilpotent unit in R and let s_1, \dots, s_n denote R -module generators of S . Each s_i is integral over R , and multiplying by a large enough power of ϖ we may assume that each s_i is integral over R_0 . Let S_0 be the subring of S generated by R_0 and s_1, \dots, s_n . Now S_0 is a finite R_0 -module and is in particular a Noetherian ring. Moreover S_0 with the ϖ -adic topology is an open subring of S . In particular, S is an f -adic topological ring, and since ϖ is a topologically nilpotent unit we see that S is a Tate ring with a Noetherian ring of definition. Completeness of S follows from completeness of finitely generated modules over Noetherian adic rings (this is [Huber 1994, Lemma 2.3(ii)]).

Finally we show that the integral closure R° in S is equal to S° . It is clear from the definition of the topology on S that R° maps to S° so the integral closure of R° in S is contained in S° . Conversely, by Lemma A.2, S° is the integral closure of S_0 in S . Since S_0 is integral over R_0 , we see that S° is integral over R_0 , and therefore it is integral over R° . \square

Next we recall the notion of uniformity. If R is a normed ring, then the spectral seminorm $|\cdot|_{\text{sp}}$ on R is defined by $|r|_{\text{sp}} = \lim_{n \rightarrow \infty} |r^n|^{1/n}$. It is well known that this limit exists and defines a power-multiplicative seminorm. Whenever it is a norm, we will refer to it as the spectral norm on R . Conversely, if we mention “the spectral norm of R ”, we are implicitly stating (or assuming) that the spectral seminorm is a norm.

Definition A.4. Let R be a complete Tate ring. We say that R is *uniform* if the set of power-bounded elements R° is bounded. We say that R is *stably uniform* if any rational localization of R is also uniform. If R is a Banach–Tate ring, we say that R is *uniform* if the norm is power-multiplicative.

Note that, if R is Banach–Tate ring whose underlying complete Tate ring is uniform, then the given norm on R is equivalent to the corresponding spectral norm, which is power-multiplicative. In this case, [Berkovich 1990, Theorem 1.3] says that the spectral norm is equal to the Gelfand norm $\sup_{x \in \mathcal{M}(R)} |\cdot|_x$. If R is in addition stably uniform, then if $\varpi \in R$ is a multiplicative pseudouniformizer and $U \subseteq X = \text{Spa}(R, R^+)$ is a rational subdomain, we may equate $\mathcal{M}(\mathcal{O}_X(U))$ with the rank-1 points in U using ϖ and equip $\mathcal{O}_X(U)$ with the corresponding Gelfand norm.

We may extend the definition of stable uniformity to arbitrary *analytic* adic spaces, i.e., those that are locally the adic spectra of complete Tate rings. We say that such an X is stably uniform if there is a cover of open affinoid subsets $U_i \subseteq X$ such that $\mathcal{O}_X(U_i)$ is stably uniform. We remark that if R is a complete sheafy Tate ring such that $\text{Spa}(R, R^+)$ is stably uniform, then R is stably uniform (this is a short argument; see [Kedlaya and Liu 2015, Remark 2.8.12]). When R has a Noetherian ring of definition, many naturally occurring complete Tate rings are stably uniform. Below we will prove some results in this direction.

Theorem A.5. *Let A be a reduced quasiexcellent ring. Let I be an ideal of A and give A the I -adic topology. If U is a rational subdomain of $X = \text{Spa}(A, A)$ and $\mathcal{O}_X(U)$ is Tate, then $\mathcal{O}_X(U)$ is uniform. In other words, the analytic locus $X^{\text{an}} \subseteq X$ is stably uniform. We also have $\mathcal{O}_X^+(U) = \mathcal{O}_X(U)^\circ$.*

Moreover, if $U \subset X^{\text{an}}$ is an arbitrary open affinoid then $\mathcal{O}_{X^{\text{an}}}^+(U) = \mathcal{O}_{X^{\text{an}}}(U)^\circ$ and $\mathcal{O}_{X^{\text{an}}}(U)^\circ$ is bounded in $\mathcal{O}_{X^{\text{an}}}(U)$.

Proof. Let $f_1, \dots, f_n, g \in A$ such that f_1, \dots, f_n generate an open ideal and let

$$U = \{|f_1|, \dots, |f_n| \leq |g| \neq 0\}.$$

Put $R = \mathcal{O}_X(U)$; recall that R may be constructed as completion of the f -adic ring $T = A[1/g]$ with ring of definition $T_0 = A[f_1/g, \dots, f_n/g] \subseteq T$ and ideal of definition $J = I[f_1/g, \dots, f_n/g] \subseteq T_0$. Let R_0 be the J -adic completion of T_0 ; this is a ring of definition of R , with ideal of definition JR_0 . Since A is reduced, so is T and hence T_0 ; moreover T_0 is quasiexcellent since it is finitely generated over A . Recall that a Noetherian ring is reduced if and only if Serre's conditions R_0 and S_1 hold (we apologize for the unfortunate clash of notations). By [EGA IV₂ 1965, (7.8.3.1)], R_0 inherits these properties from T_0 and is therefore reduced. Moreover, T_0 , and hence $T_0/J = R_0/JR_0$, are Nagata (since they are finitely generated over A , which is quasiexcellent, and hence Nagata). By [Marot 1975, Proposition 2.3], R_0 is Nagata (note that there is a trivial misprint in the reference).

Pick a topologically nilpotent unit $\varpi \in R$ (recall that R is Tate by assumption); without loss of generality assume $\varpi \in R_0$. Then $R = R_0[1/\varpi]$, so R is contained in the total ring of fractions $\mathcal{Q}(R_0)$ of R_0 . Since R_0 is reduced and Nagata, it follows that the integral closure R' of R_0 in R is a finitely generated R_0 -module, and hence is bounded. Now $R' = R^\circ$ by Lemma A.2, so R° is bounded as desired.

For the assertion about $\mathcal{O}_X^+(U)$, let T^+ denote the integral closure of T_0 in T . By definition, the completion of T^+ is $R^+ := \mathcal{O}_X^+(U)$. In particular, R^+ contains R_0 and the assertion now follows from Lemma A.2.

To check the assertion about an open affinoid $U = \text{Spa}(\mathcal{O}_X(U), \mathcal{O}_X^+(U))$, note that U has a finite cover by Tate rational subdomains $(U_i)_{i \in I}$ of X . Since the maps $\mathcal{O}(U) \rightarrow \mathcal{O}(U_i)$ are bounded (the U_i are also rational subdomains of U), the strict embedding

$$\mathcal{O}(U) \hookrightarrow \prod_{i \in I} \mathcal{O}(U_i)$$

induces an embedding

$$\mathcal{O}(U)^\circ \hookrightarrow \mathcal{O}(U) \cap \prod_{i \in I} \mathcal{O}(U_i)^\circ$$

but the right-hand side equals

$$\mathcal{O}(U) \cap \prod_{i \in I} \mathcal{O}^+(U_i) = \mathcal{O}^+(U)$$

by the first part of the theorem, so we are done because by definition $\mathcal{O}^+(U) \subseteq \mathcal{O}(U)^\circ$, which implies that we have equality. Finally, the boundedness of $\mathcal{O}_{X^{\text{an}}}(U)^\circ$ follows from the boundedness of the $\mathcal{O}(U_i)^\circ$. \square

Corollary A.6. *Let \mathcal{O} be a complete discrete valuation ring and let A be a reduced complete Noetherian adic ring formally of finite type over \mathcal{O} , i.e., such that A/A° is a finitely generated \mathcal{O} -algebra. Then the analytic locus $X^{\text{an}} \subseteq X$ is stably uniform. Moreover, if $U \subset X^{\text{an}}$ is an open affinoid subspace then $\mathcal{O}_{X^{\text{an}}}^+(U) = \mathcal{O}_{X^{\text{an}}}(U)^\circ$ and $\mathcal{O}_{X^{\text{an}}}(U)^\circ$ is bounded in $\mathcal{O}_{X^{\text{an}}}(U)$.*

Proof. A is excellent by [Valabrega 1975, Proposition 7; 1976, Theorem 9] (see [Conrad 1999], near the end of the Introduction). Thus, Theorem A.5 applies. \square

We also note that the proof of Theorem A.5 applies essentially verbatim to prove the following similar result.

Theorem A.7. *Let R be a complete Tate ring and assume that R has a ring of definition R_0 which is quasiexcellent and reduced. Then R is stably uniform. Moreover, if $X = \mathrm{Spa}(R, R^\circ)$, and $U \subset X$ is an open affinoid subspace, then $\mathcal{O}_X^+(U) = \mathcal{O}_X(U)^\circ$ and $\mathcal{O}_X(U)^\circ$ is bounded in $\mathcal{O}_X(U)$. In particular, $\mathcal{O}_X(U)$ is reduced.*

This theorem also allows us to develop the theory of the nilreduction of an adic space. We only give a sketch here — one can check that everything in [Bosch et al. 1984, §9.5.1] works in our setting.

Definition A.8. Let X be an adic space. Define the *nilradical* $\mathrm{rad} \mathcal{O}_X$ to be the sheaf associated to the presheaf $U \mapsto \mathrm{rad}(\mathcal{O}_X(U))$, where $\mathrm{rad}(\mathcal{O}_X(U))$ is the nilradical of the ring $\mathcal{O}_X(U)$.

Proposition A.9. *Let R be a complete Tate ring and assume that R has a ring of definition R_0 which is quasiexcellent. Let $X = \mathrm{Spa}(R, R^\circ)$. Then $\mathrm{rad} \mathcal{O}_X \subset \mathcal{O}_X$ is a coherent \mathcal{O}_X -ideal, associated to the ideal $\mathrm{rad}(R)$ of R .*

More generally, if X is an adic space which is locally of the form $\mathrm{Spa}(R, R^\circ)$ where R is a complete Tate ring with a quasiexcellent ring of definition, then $\mathrm{rad} \mathcal{O}_X$ is a coherent \mathcal{O}_X -ideal.

Proof. The key point is that if $U \subset X = \mathrm{Spa}(R, R^\circ)$ is a rational subdomain, then

$$\mathrm{Spa}(\mathcal{O}_X(U)/\mathrm{rad}(R), (\mathcal{O}_X(U)/\mathrm{rad}(R))^\circ) \rightarrow \mathrm{Spa}(R^{\mathrm{red}}, (R^{\mathrm{red}})^\circ)$$

is a rational subdomain, so Theorem A.7 implies that $\mathcal{O}_X(U)/\mathrm{rad}(R)$ is reduced, which implies that $\mathrm{rad}(\mathcal{O}_X(U)) = \mathrm{rad}(R)\mathcal{O}_X(U)$. \square

Definition A.10. Let X be an adic space which is locally of the form $\mathrm{Spa}(R, R^\circ)$, where R is a complete Tate ring with a quasiexcellent ring of definition. Then we define X^{red} to be the closed subspace of X cut out by $\mathrm{rad} \mathcal{O}_X$ (see [Huber 1996, 1.4]).

In this paper the analytic adic spaces encountered will locally be of the form $\mathrm{Spa}(R, R^\circ)$, where R is a complete Tate ring with a ring of definition R_0 which is formally of finite type over \mathbb{Z}_p . We will need a few properties of these rings, all of which follow from the material in [Abbes 2010]. We recall the following definition from that paper, specialized to our Noetherian situation.

Definition A.11. A Noetherian adic ring B is called a *1-valuative order* if it is an integral domain which is local of Krull dimension 1, and has no J -torsion, where J is an ideal of definition (this is independent of the choice of ideal of definition).

This is [Abbes 2010, Definition 1.11.1], except that we demand that B is Noetherian. If B is a 1-valuative order, then the integral closure \bar{B} in $L = \mathrm{Frac}(B)$ is finite over B and is a complete discrete valuation ring, so L is a complete discrete valuation field [loc. cit., Proposition 1.11.4]. If A is any

Noetherian adic ring with an ideal of definition I and $\mathfrak{p} \in \operatorname{Spec} A$, then A/\mathfrak{p} is a 1-valuation order if and only if \mathfrak{p} is a closed point in $\operatorname{Spec} A \setminus V(I)$ [loc. cit., Proposition 1.11.8].

Lemma A.12. *Let R be a complete Tate ring with a ring of definition R_0 which is formally of finite type over \mathbb{Z}_p , and let $\mathfrak{m} \subseteq R$ be a maximal ideal. Then R/\mathfrak{m} is a local field.*

Proof. Let $\mathfrak{p} = R_0 \cap \mathfrak{m}$ and let $\varpi \in R_0$ be a topologically nilpotent unit. Then \mathfrak{p} is a closed point in $\operatorname{Spec} R_0 \setminus V((\varpi)) = \operatorname{Spec} R$, so R_0/\mathfrak{p} is a 1-valuation order and hence its fraction field R/\mathfrak{m} is a complete discrete valuation field. It remains to prove that the residue field is finite. For this, it suffices to show that the residue field of the local ring R_0/\mathfrak{p} is finite since the integral closure of R_0/\mathfrak{p} in R/\mathfrak{m} is finite over R_0/\mathfrak{p} . Pick an adic surjection $A = \mathbb{Z}_p[[T_1, \dots, T_m]]\langle X_1, \dots, X_n \rangle \twoheadrightarrow R_0$ for some $m, n \in \mathbb{Z}_{\geq 0}$. The maximal ideal of R_0/\mathfrak{p} is open and so corresponds to an open maximal ideal of A , and hence to a maximal ideal of $\mathbb{F}_p[X_1, \dots, X_n]$ in a way that preserves residue fields. It follows that R_0/\mathfrak{p} is finite as desired. \square

Lemma A.13. *Let $f : A \rightarrow B$ be a morphism of topologically finite type between Noetherian adic rings. Let I be an ideal of definition of A and assume that A/I is Jacobson. Let $J = IB$; this is an ideal of definition of B . If $\mathfrak{q} \in \operatorname{Spec} B \setminus V(J)$ is a closed point, then $\mathfrak{p} = f^{-1}(\mathfrak{q})$ is a closed point in $\operatorname{Spec} A \setminus V(I)$.*

Proof. The morphism $A \rightarrow B/\mathfrak{q}$ is topologically of finite type and B/\mathfrak{q} is a 1-valuation order, so by [Abbes 2010, Proposition 1.11.2] $A \rightarrow B/\mathfrak{q}$ is finite. It is then easy to check that this forces A/\mathfrak{p} to be a 1-valuation order as well, and hence \mathfrak{p} to be closed in $\operatorname{Spec} A \setminus V(I)$ by [loc. cit., Proposition 1.11.8]. \square

Corollary A.14. *Let $g : R \rightarrow S$ be a continuous morphism between two complete Tate rings with a ring of definition that is formally of finite type over \mathbb{Z}_p . Then g is topologically of finite type⁵ and pulls back maximal ideals to maximal ideals.*

Proof. Choose a ring of definition R_0 for R which is formally of finite type over \mathbb{Z}_p . By [Huber 1993, Proposition 1.10] g is adic, and therefore $g(R_0)$ is contained in a ring of definition for S . Since any two rings of definition are contained in another, we can find a ring of definition S_0 for S such that $g(R_0) \subseteq S_0$ and S_0 contains a ring of definition S_1 which is formally of finite type over \mathbb{Z}_p . It follows from Lemma A.2 that S_0 is finite over S_1 , and hence S_0 is also formally of finite type over \mathbb{Z}_p .

Let $\varpi \in R_0$ be a topologically nilpotent unit in R . Then $I = \varpi R_0$ and $J = g(\varpi)S_0$ are ideals of definition, and $R_0/I \rightarrow S_0/J$ is of finite type. Therefore $R_0 \rightarrow S_0$ is topologically of finite type; hence so is g . This proves the first assertion. The second then follows from Lemma A.13, since maximal ideals of R and S correspond to closed points in $\operatorname{Spec} R = \operatorname{Spec} R_0 \setminus V(I)$ and $\operatorname{Spec} S = \operatorname{Spec} S_0 \setminus V(J)$ respectively. \square

Proposition A.15. *Let S be a complete Tate ring with a Noetherian ring of definition S_0 and a topologically nilpotent unit $\pi \in S_0$ such that $S_0/\pi S_0$ is Jacobson:*

(1) *Let A be a Noetherian adic ring with an ideal of definition I such that A/I is Jacobson. Let $f : A \rightarrow S$ be a continuous morphism such that the induced map $\operatorname{Spa}(S, S^\circ) \rightarrow \operatorname{Spa}(A, A)$ is an open immersion,*

⁵That is, g factors through a surjective, continuous and open morphism $R\langle X_1, \dots, X_n \rangle \rightarrow S$; see [Huber 1994, Lemma 3.3].

and let \mathfrak{q} be a maximal ideal of S with preimage $\mathfrak{p} = f^{-1}(\mathfrak{q})$ in A . Then the natural map $A_{\mathfrak{p}} \rightarrow S_{\mathfrak{q}}$ induces an isomorphism on completions (with respect to the maximal ideals).

(2) Let R be a complete Tate ring with a Noetherian ring of definition R_0 and a topologically nilpotent unit $\varpi \in R_0$ such that R_0/ϖ is Jacobson. Let $h : R \rightarrow S$ be a continuous morphism such that the induced map $\mathrm{Spa}(S, S^\circ) \rightarrow \mathrm{Spa}(R, R^\circ)$ is an open immersion, and let \mathfrak{q}' be a maximal ideal of S with preimage $\mathfrak{p}' = h^{-1}(\mathfrak{q}')$ in R . Then the natural map $R_{\mathfrak{p}'} \rightarrow S_{\mathfrak{q}'}$ induces an isomorphism on completions (with respect to the maximal ideals).

Proof. We prove part (1); the proof of part (2) is virtually identical. Since S/\mathfrak{q} is a complete discretely valued field it defines a point v in $\mathrm{Spa}(S, S^\circ) \subseteq \mathrm{Spa}(A, A)$; let $U = \{|f_1|, \dots, |f_n| \leq |g| \neq 0\}$ be a rational subdomain of $\mathrm{Spa}(A, A)$ which contains this point and is contained in $\mathrm{Spa}(S, S^\circ)$. Let

$$T = A[f_1/g, \dots, f_n/g] \subseteq A[1/g]$$

and let \widehat{T} be the IT -adic completion of T . Since $\widehat{T}[1/g] = \mathcal{O}(U)$ we see that the valuation v extends to a valuation w on $\widehat{T}[1/g]$, and hence \mathfrak{q} extends to a maximal ideal $\mathfrak{r} = \mathrm{Ker} w$ of $\widehat{T}[1/g]$. We will abuse notation and let \mathfrak{r} denote its preimage in any of the rings T , $T[1/g]$ and \widehat{T} as well; then \mathfrak{r} is a closed point in $\mathrm{Spec} \widehat{T} \setminus V(IT)$.

By [Abbes 2010, Proposition 1.12.18], the natural map $T_{\mathfrak{r}} \rightarrow \widehat{T}_{\mathfrak{r}}$ induces an isomorphism of completions. We claim that the natural maps $A_{\mathfrak{p}} \rightarrow T_{\mathfrak{r}}$ and $\widehat{T}_{\mathfrak{r}} \rightarrow \widehat{T}[1/g]_{\mathfrak{r}}$ are isomorphisms. For the second map this is clear (by the general fact that if B is any ring, $f \in B$, and $P \in \mathrm{Spec} B[1/f] \subseteq \mathrm{Spec} B$, then the natural map $B_P \rightarrow B[1/f]_P$ is an isomorphism). For the first map, we have natural maps $A_{\mathfrak{p}} \rightarrow T_{\mathfrak{r}} \rightarrow T[1/g]_{\mathfrak{r}} = A[1/g]_{\mathfrak{r}}$ and it is clear that the second map and the composite are isomorphisms, so the first map is an isomorphism as well. Summing up, we see that the natural map $A_{\mathfrak{p}} \rightarrow \mathcal{O}(U)_{\mathfrak{r}}$ induces an isomorphism on completions. By an almost identical argument, the natural map $S_{\mathfrak{q}} \rightarrow \mathcal{O}(U)_{\mathfrak{r}}$ induces an isomorphism on completions. It then follows that the natural map $A_{\mathfrak{p}} \rightarrow S_{\mathfrak{q}}$ induces an isomorphism on completions, as desired. \square

Acknowledgments

This paper was inspired by [Andreatta et al. 2018], and Johansson wishes to thank Vincent Pilloni for some interesting discussions on that paper and David Hansen for innumerable and invaluable discussions on overconvergent cohomology and eigenvarieties. He would also like to thank Konstantin Ardakov for patiently answering several questions, and both authors would like to thank him for providing us with a proof of Proposition 3.3.4, and allowing us to include it here. We wish to thank John Bergdall and Kevin Buzzard for various conversations about the boundary of the eigencurve, as well as helpful comments on an earlier version of this paper. Finally, we wish to thank the anonymous referees for their careful reading and helpful comments. Johansson was supported by NSF grant DMS-1128155 during the main work on this paper, and by the Herchel Smith Foundation during the revision. Newton was supported by ERC Starting Grant 306326. Work on this project was also supported by the EPSRC Platform Grant EP/I019111.

References

- [Abbes 2010] A. Abbes, *Éléments de géométrie rigide, I: Construction et étude géométrique des espaces rigides*, Progr. Math. **286**, Springer, 2010. MR Zbl
- [Andreatta et al. 2016] F. Andreatta, A. Iovita, and V. Pilloni, “The adic, cuspidal, Hilbert eigenvarieties”, *Res. Math. Sci.* **3** (2016), art. id. 34. MR Zbl
- [Andreatta et al. 2018] F. Andreatta, A. Iovita, and V. Pilloni, “Le halo spectral”, *Ann. Sci. Éc. Norm. Supér. (4)* **51**:3 (2018), 603–655. MR Zbl
- [Ardakov and Wadsley 2013] K. Ardakov and S. Wadsley, “On irreducible representations of compact p -adic analytic groups”, *Ann. of Math. (2)* **178**:2 (2013), 453–557. MR Zbl
- [Ash and Stevens 2008] A. Ash and G. Stevens, “ p -adic deformations of arithmetic cohomology”, preprint, 2008, available at <http://math.bu.edu/people/ghs/preprints/Ash-Stevens-02-08.pdf>.
- [Bellaïche 2010] J. Bellaïche, “Eigenvarieties and p -adic L -functions”, preprint, 2010, available at <http://people.brandeis.edu/~jbellaic/preprint/coursebook.pdf>.
- [Bellaïche and Chenevier 2009] J. Bellaïche and G. Chenevier, *Families of Galois representations and Selmer groups*, Astérisque **324**, Société Mathématique de France, Paris, 2009. MR Zbl
- [Bergdall and Pollack 2016] J. Bergdall and R. Pollack, “Arithmetic properties of Fredholm series for p -adic modular forms”, *Proc. Lond. Math. Soc. (3)* **113**:4 (2016), 419–444. MR Zbl
- [Berkovich 1990] V. G. Berkovich, *Spectral theory and analytic geometry over non-Archimedean fields*, Math. Surveys and Monographs **33**, Amer. Math. Soc., Providence, RI, 1990. MR Zbl
- [Bosch et al. 1984] S. Bosch, U. Güntzer, and R. Remmert, *Non-Archimedean analysis: a systematic approach to rigid analytic geometry*, Grundlehren der Math. Wissenschaften **261**, Springer, 1984. MR Zbl
- [Breuil et al. 2017] C. Breuil, E. Hellmann, and B. Schraen, “Une interprétation modulaire de la variété trianguline”, *Math. Ann.* **367**:3–4 (2017), 1587–1645. MR Zbl
- [Buzzard 2004] K. Buzzard, “On p -adic families of automorphic forms”, pp. 23–44 in *Modular curves and abelian varieties* (Bellaterra, Spain, 2002), edited by J. Cremona et al., Progr. Math. **224**, Birkhäuser, Basel, 2004. MR Zbl
- [Buzzard 2007] K. Buzzard, “Eigenvarieties”, pp. 59–120 in *L -functions and Galois representations* (Durham, 2004), edited by D. Burns et al., London Math. Soc. Lecture Note Ser. **320**, Cambridge Univ. Press, 2007. MR Zbl
- [Buzzard and Kilford 2005] K. Buzzard and L. J. P. Kilford, “The 2-adic eigencurve at the boundary of weight space”, *Compos. Math.* **141**:3 (2005), 605–619. MR Zbl
- [Chenevier 2004] G. Chenevier, “Familles p -adiques de formes automorphes pour GL_n ”, *J. Reine Angew. Math.* **570** (2004), 143–217. MR Zbl
- [Chenevier 2005] G. Chenevier, “Une correspondance de Jacquet–Langlands p -adique”, *Duke Math. J.* **126**:1 (2005), 161–194. MR Zbl
- [Chenevier 2014] G. Chenevier, “The p -adic analytic space of pseudocharacters of a profinite group and pseudorepresentations over arbitrary rings”, pp. 221–285 in *Automorphic forms and Galois representations, I* (Durham, 2011), edited by F. Diamond et al., London Math. Soc. Lecture Note Ser. **414**, Cambridge Univ. Press, 2014. MR Zbl
- [Chojecki et al. 2017] P. Chojecki, D. Hansen, and C. Johansson, “Overconvergent modular forms and perfectoid Shimura curves”, *Doc. Math.* **22** (2017), 191–262. MR Zbl
- [Coleman 1997] R. F. Coleman, “ p -adic Banach spaces and families of modular forms”, *Invent. Math.* **127**:3 (1997), 417–479. MR Zbl
- [Coleman and Mazur 1998] R. Coleman and B. Mazur, “The eigencurve”, pp. 1–113 in *Galois representations in arithmetic algebraic geometry* (Durham, 1996), edited by A. J. Scholl and R. L. Taylor, London Math. Soc. Lecture Note Ser. **254**, Cambridge Univ. Press, 1998. MR Zbl
- [Colmez 2010] P. Colmez, “Fonctions d’une variable p -adique”, pp. 13–59 Astérisque **330**, Société Mathématique de France, Paris, 2010. MR Zbl

- [Conrad 1999] B. Conrad, “Irreducible components of rigid spaces”, *Ann. Inst. Fourier (Grenoble)* **49**:2 (1999), 473–541. MR Zbl
- [Dixon et al. 1999] J. D. Dixon, M. P. F. du Sautoy, A. Mann, and D. Segal, *Analytic pro- p -groups*, 2nd ed., Cambridge Studies in Adv. Math. **61**, Cambridge Univ. Press, 1999. MR Zbl
- [EGA IV₂ 1965] A. Grothendieck, “Eléments de géométrie algébrique, IV: Étude locale des schémas et des morphismes de schémas, II”, *Inst. Hautes Études Sci. Publ. Math.* **24** (1965), 5–231. MR Zbl
- [EGA IV₃ 1966] A. Grothendieck, “Eléments de géométrie algébrique, IV: Étude locale des schémas et des morphismes de schémas, III”, *Inst. Hautes Études Sci. Publ. Math.* **28** (1966), 5–255. MR Zbl
- [Gulotta 2018] D. R. Gulotta, *Equidimensional adic eigenvarieties for groups with discrete series*, Ph.D. thesis, Columbia University, 2018, available at <https://search.proquest.com/docview/2031119725>.
- [Hansen 2015] D. Hansen, “Iwasawa theory of overconvergent modular forms, I: Critical p -adic L -functions”, preprint, 2015. arXiv
- [Hansen 2017] D. Hansen, “Universal eigenvarieties, trianguline Galois representations, and p -adic Langlands functoriality”, *J. Reine Angew. Math.* **730** (2017), 1–64. MR Zbl
- [Hansen and Johansson \geq 2019] D. Hansen and C. Johansson, “Completed and overconvergent cohomology”, in preparation.
- [Huber 1993] R. Huber, “Continuous valuations”, *Math. Z.* **212**:3 (1993), 455–477. MR Zbl
- [Huber 1994] R. Huber, “A generalization of formal schemes and rigid analytic varieties”, *Math. Z.* **217**:4 (1994), 513–551. MR Zbl
- [Huber 1996] R. Huber, *Étale cohomology of rigid analytic varieties and adic spaces*, Aspects of Math. **E30**, Vieweg, Wiesbaden, 1996. MR Zbl
- [Johansson and Newton 2017] C. Johansson and J. Newton, “Irreducible components of extended eigenvarieties and interpolating Langlands functoriality”, 2017. To appear in *Math. Res. Lett.* arXiv
- [Johansson and Newton 2018] C. Johansson and J. Newton, “Parallel weight 2 points on Hilbert modular eigenvarieties and the parity conjecture”, preprint, 2018. arXiv
- [Kedlaya and Liu 2015] K. S. Kedlaya and R. Liu, *Relative p -adic Hodge theory: foundations*, Astérisque **371**, Société Mathématique de France, Paris, 2015. MR Zbl arXiv
- [Liu et al. 2017] R. Liu, D. Wan, and L. Xiao, “The eigencurve over the boundary of weight space”, *Duke Math. J.* **166**:9 (2017), 1739–1787. MR Zbl
- [Loeffler 2011] D. Loeffler, “Overconvergent algebraic automorphic forms”, *Proc. Lond. Math. Soc.* (3) **102**:2 (2011), 193–228. MR Zbl
- [Marot 1975] J. Marot, “Sur les anneaux universellement japonais”, *Bull. Soc. Math. France* **103**:1 (1975), 103–111. MR Zbl
- [Neukirch 1999] J. Neukirch, *Algebraic number theory*, Grundlehren der Math. Wissenschaften **322**, Springer, 1999. MR Zbl
- [Pottharst and Xiao 2014] J. Pottharst and L. Xiao, “On the parity conjecture in finite-slope families”, preprint, 2014. arXiv
- [Schneider and Teitelbaum 2003] P. Schneider and J. Teitelbaum, “Algebras of p -adic distributions and admissible representations”, *Invent. Math.* **153**:1 (2003), 145–196. MR Zbl
- [Scholze 2015] P. Scholze, “On torsion in the cohomology of locally symmetric varieties”, *Ann. of Math.* (2) **182**:3 (2015), 945–1066. MR Zbl
- [Serre 1962] J.-P. Serre, “Endomorphismes complètement continus des espaces de Banach p -adiques”, *Inst. Hautes Études Sci. Publ. Math.* **12** (1962), 69–85. MR Zbl
- [Stevens 1994] G. Stevens, “Rigid analytic modular symbols”, preprint, 1994, available at <http://math.bu.edu/people/ghs/research.d/RigidSymbs.pdf>.
- [Urban 2011] E. Urban, “Eigenvarieties for reductive groups”, *Ann. of Math.* (2) **174**:3 (2011), 1685–1784. MR Zbl
- [Valabrega 1975] P. Valabrega, “On the excellent property for power series rings over polynomial rings”, *J. Math. Kyoto Univ.* **15**:2 (1975), 387–395. MR Zbl
- [Valabrega 1976] P. Valabrega, “A few theorems on completion of excellent rings”, *Nagoya Math. J.* **61** (1976), 127–133. MR Zbl

[Weibel 1994] C. A. Weibel, *An introduction to homological algebra*, Cambridge Studies in Adv. Math. **38**, Cambridge Univ. Press, 1994. MR Zbl

[Xiang 2012] Z. Xiang, “A construction of the full eigenvariety of a reductive group”, *J. Number Theory* **132**:5 (2012), 938–952. MR Zbl

Communicated by Bjorn Poonen

Received 2017-06-15 Revised 2018-06-28 Accepted 2018-09-25

chrjohv@chalmers.se

*Department of Mathematical Sciences, Chalmers University of Technology
and the University of Gothenburg, Gothenburg, Sweden*

j.newton@kcl.ac.uk

Department of Mathematics, King’s College London, London, United Kingdom

Algebra & Number Theory

msp.org/ant

EDITORS

MANAGING EDITOR

Bjorn Poonen
Massachusetts Institute of Technology
Cambridge, USA

EDITORIAL BOARD CHAIR

David Eisenbud
University of California
Berkeley, USA

BOARD OF EDITORS

Richard E. Borcherds	University of California, Berkeley, USA	Martin Olsson	University of California, Berkeley, USA
Antoine Chambert-Loir	Université Paris-Diderot, France	Raman Parimala	Emory University, USA
J-L. Colliot-Thélène	CNRS, Université Paris-Sud, France	Jonathan Pila	University of Oxford, UK
Brian D. Conrad	Stanford University, USA	Anand Pillay	University of Notre Dame, USA
Samit Dasgupta	University of California, Santa Cruz, USA	Michael Rapoport	Universität Bonn, Germany
Hélène Esnault	Freie Universität Berlin, Germany	Victor Reiner	University of Minnesota, USA
Gavril Farkas	Humboldt Universität zu Berlin, Germany	Peter Sarnak	Princeton University, USA
Hubert Flenner	Ruhr-Universität, Germany	Joseph H. Silverman	Brown University, USA
Sergey Fomin	University of Michigan, USA	Michael Singer	North Carolina State University, USA
Edward Frenkel	University of California, Berkeley, USA	Christopher Skinner	Princeton University, USA
Andrew Granville	Université de Montréal, Canada	Vasudevan Srinivas	Tata Inst. of Fund. Research, India
Joseph Gubeladze	San Francisco State University, USA	J. Toby Stafford	University of Michigan, USA
Roger Heath-Brown	Oxford University, UK	Pham Huu Tiep	University of Arizona, USA
Craig Huneke	University of Virginia, USA	Ravi Vakil	Stanford University, USA
Kiran S. Kedlaya	Univ. of California, San Diego, USA	Michel van den Bergh	Hasselt University, Belgium
János Kollár	Princeton University, USA	Marie-France Vignéras	Université Paris VII, France
Philippe Michel	École Polytechnique Fédérale de Lausanne	Kei-Ichi Watanabe	Nihon University, Japan
Susan Montgomery	University of Southern California, USA	Shou-Wu Zhang	Princeton University, USA
Shigefumi Mori	RIMS, Kyoto University, Japan		

PRODUCTION

production@msp.org
Silvio Levy, Scientific Editor

See inside back cover or msp.org/ant for submission instructions.

The subscription price for 2019 is US \$385/year for the electronic version, and \$590/year (+\$60, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to MSP.

Algebra & Number Theory (ISSN 1944-7833 electronic, 1937-0652 printed) at Mathematical Sciences Publishers, 798 Evans Hall #3840, c/o University of California, Berkeley, CA 94720-3840 is published continuously online. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices.

ANT peer review and production are managed by EditFlow® from MSP.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2019 Mathematical Sciences Publishers

Algebra & Number Theory

Volume 13 No. 1 2019

Ordinary algebraic curves with many automorphisms in positive characteristic GÁBOR KORCHMÁROS and MARIA MONTANUCCI	1
Variance of arithmetic sums and L -functions in $\mathbb{F}_q[t]$ CHRIS HALL, JONATHAN P. KEATING and EDVA RODITTY-GERSHON	19
Extended eigenvarieties for overconvergent cohomology CHRISTIAN JOHANSSON and JAMES NEWTON	93
A tubular variant of Runge's method in all dimensions, with applications to integral points on Siegel modular varieties SAMUEL LE FOURN	159
Algebraic cycles on genus-2 modular fourfolds DONU ARAPURA	211
Average nonvanishing of Dirichlet L -functions at the central point KYLE PRATT	227



1937-0652(2019)13:1;1-C