



Moments of random multiplicative functions, II: High moments

Adam J Harper

We determine the order of magnitude of $\mathbb{E}|\sum_{n \le x} f(n)|^{2q}$ up to factors of size $e^{O(q^2)}$, where f(n) is a Steinhaus or Rademacher random multiplicative function, for all real $1 \le q \le c \log x / \log \log x$. In the Steinhaus case, we show that $\mathbb{E}|\sum_{n \le x} f(n)|^{2q} = e^{O(q^2)} x^q (\log x / (q \log(2q)))^{(q-1)^2}$ on this whole

In the Steinhaus case, we show that $\mathbb{E}\left|\sum_{n \le x} f(n)\right|^{2q} = e^{O(q^2)} x^q (\log x/(q \log(2q)))^{(q-1)^2}$ on this whole range. In the Rademacher case, we find a transition in the behavior of the moments when $q \approx (1 + \sqrt{5})/2$, where the size starts to be dominated by "orthogonal" rather than "unitary" behavior. We also deduce some consequences for the large deviations of $\sum_{n \le x} f(n)$.

The proofs use various tools, including hypercontractive inequalities, to connect $\mathbb{E}|\sum_{n\leq x} f(n)|^{2q}$ with the *q*-th moment of an Euler product integral. When *q* is large, it is then fairly easy to analyze this integral. When *q* is close to 1 the analysis seems to require subtler arguments, including Doob's L^p maximal inequality for martingales.

1. Introduction

In this sequence of papers, we are interested in the moments $\mathbb{E} \left| \sum_{n \le x} f(n) \right|^{2q}$ of random multiplicative functions f(n).

We consider two different models for f(n), a *Steinhaus random multiplicative function* and a *Rade-macher random multiplicative function*. We obtain a Steinhaus random multiplicative function by letting $(f(p))_{p \text{ prime}}$ be a sequence of independent Steinhaus random variables (i.e., distributed uniformly on the unit circle $\{|z| = 1\}$), and then setting $f(n) := \prod_{p^a \parallel n} f(p)^a$ for all natural numbers n, where $p^a \parallel n$ means that p^a is the highest power of the prime p that divides n. We obtain a Rademacher random multiplicative function by letting $(f(p))_{p \text{ prime}}$ be independent Rademacher random variables (i.e., taking values ± 1 with probability $\frac{1}{2}$ each), and then setting $f(n) := \prod_{p \mid n} f(p)$ for all squarefree n, and f(n) = 0 when n is not squarefree.

Random multiplicative functions have attracted quite a lot of attention as models for functions of number theoretic interest: for example, Rademacher random multiplicative functions were introduced by Wintner [1944] as a model for the Möbius function $\mu(n)$. There are also probabilistic and analytic

When this work was started, the author was supported by a research fellowship at Jesus College, Cambridge. The work continued while the author was in residence at the Mathematical Sciences Research Institute in Berkeley, California (supported by the National Science Foundation under Grant No. DMS-1440140), during the Spring 2017 semester. *MSC2010:* primary 11N56; secondary 11K65, 11L40.

Keywords: random multiplicative functions, random Euler products, moments, orthogonal behavior, unitary behavior, martingales.

motivations for studying them, see Saksman and Seip's open problems paper [2016], for example. The introduction to the previous paper [Harper 2017] in this sequence contains a more extensive discussion of some of these connections.

Harper [2017] showed that for Steinhaus or Rademacher random multiplicative f(n), for all large x we have

$$\mathbb{E} \left| \sum_{n \le x} f(n) \right|^{2q} \asymp \left(\frac{x}{1 + (1 - q)\sqrt{\log \log x}} \right)^q \quad \forall 0 \le q \le 1.$$

In particular, taking $q = \frac{1}{2}$ this implies that $\mathbb{E} \left| \sum_{n \le x} f(n) \right| \asymp \sqrt{x} / (\log \log x)^{1/4}$, which proved a conjecture of Helson [2010] that the first absolute moment should be $o(\sqrt{x})$.

Our goal here is to investigate the case where $q \ge 1$. When $q \in \mathbb{N}$ is fixed, one can expand the 2q-th power and reduce the calculation of $\mathbb{E} |\sum_{n \le x} f(n)|^{2q}$ to a number theoretic counting problem. For example, in the Steinhaus case one has

$$\mathbb{E}\bigg|\sum_{n\leq x} f(n)\bigg|^{2q} = \#\bigg\{n_1,\ldots,n_{2q}\leq x: \prod_{i=1}^q n_i = \prod_{i=q+1}^{2q} n_i\bigg\}.$$

Starting from this, one can obtain an asymptotic for the moment as $x \to \infty$, which was carried out by Harper, Nikeghbali and Radziwiłł [Harper et al. 2015], and also independently by Heap and Lindqvist [2016], and (in the Steinhaus case) in unpublished work of Granville and Soundararajan. The result is that, for fixed $q \in \mathbb{N}$ and Steinhaus random multiplicative f(n), one has

$$\mathbb{E}\left|\sum_{n\leq x} f(n)\right|^{2q} \sim C_{\mathrm{St}}(q) x^q \log^{(q-1)^2} x \quad \text{as } x \to \infty,$$
(1-1)

where the constant $C_{\text{St}}(q)$ satisfies $C_{\text{St}}(q) = e^{-q^2 \log q - q^2 \log \log q + O(q^2)}$ for large q. For Rademacher random multiplicative f(n), when q = 1 we have that $\mathbb{E} \left| \sum_{n \le x} f(n) \right|^2 = \sum_{n \le x, n \text{ squarefree}} 1 \sim (6/\pi^2)x$, and for fixed integer $q \ge 2$ we have

$$\mathbb{E}\left|\sum_{n\leq x} f(n)\right|^{2q} \sim C_{\mathrm{Rad}}(q) x^q \log^{q(2q-3)} x \quad \text{as } x \to \infty,$$

where the constant $C_{\text{Rad}}(q)$ satisfies $C_{\text{Rad}}(q) = e^{-2q^2 \log q - 2q^2 \log \log q + O(q^2)}$ for large q. As described in [Harper et al. 2015; Heap and Lindqvist 2016], we actually have much more precise information about the constants $C_{\text{St}}(q)$, $C_{\text{Rad}}(q)$ (for example they factor into explicit "arithmetic" and "geometric" parts), but this will not be important for our purposes here.

We would like to have information about $\mathbb{E} |\sum_{n \le x} f(n)|^{2q}$ when $q \ge 1$ is not necessarily integral, and that allows q to vary as a function of x rather than being fixed.

Regarding uniformity in q, Theorem 4.1 of [Granville and Soundararajan 2001] implies that for Steinhaus random multiplicative f(n), and uniformly for all large x and integers $q \ge 1$ such that $q^{eq} \le x$,

we have

$$e^{-q^2\log q - q^2\log\log(2q)} \left(\log\left(\frac{\log x}{q\log 2q}\right)\right)^{-O(q^2)} \le \frac{\mathbb{E}\left|\sum_{n\le x} f(n)\right|^{2q}}{x^q\log^{(q-1)^2} x} \le e^{-q^2\log q + O(q^2)}$$

This range of q is essentially the largest on which one could expect a result of a similar shape to (1-1). Indeed, if $q \ge A \log x / \log \log x$ for some $A \ge 1$ (say) then we have

$$e^{-q^2\log q - q^2\log\log(2q)}x^q\log^{(q-1)^2}x \le ((1+o(1))A)^{-q^2}x^q,$$

which becomes incompatible with the lower bound $\mathbb{E} |\sum_{n \le x} f(n)|^{2q} \ge (\mathbb{E} |\sum_{n \le x} f(n)|^2)^q = \lfloor x \rfloor^q$ coming from Hölder's inequality.¹ But the bounds of Granville and Soundararajan are imperfect, as the upper bound doesn't include the factor $e^{-q^2 \log \log(2q)}$ that we expect to appear, and the lower bound features the extraneous factor $(\log(\log x/(q \log 2q)))^{-O(q^2)}$. They also remain restricted to *integer q*. There are various other results in the literature that study the Steinhaus moments $\mathbb{E} |\sum_{n \le x} f(n)|^{2q}$, and variants of them, for integer q, especially for small integers where one can try to obtain lower order terms in the known asymptotics. See e.g., the preprint of Shi and Weber [2016]. However, the author is not aware of any work giving sharp moment bounds for noninteger q, nor improving the dependence on q in Granville and Soundararajan's bounds [2001] for the large integer case.

We shall prove the following uniform estimate for all real q.

Theorem 1.1. There exists a small absolute constant c > 0 such that the following is true. If f(n) is a Steinhaus random multiplicative function, then uniformly for all large x and real $1 \le q \le c \log x / (\log \log x)$ we have

$$\mathbb{E}\left|\sum_{n\leq x} f(n)\right|^{2q} = e^{-q^2\log q - q^2\log\log(2q) + O(q^2)} x^q \log^{(q-1)^2} x.$$

To avoid any confusion, we restate this first result more explicitly: on the stated range of q and x, we always have

$$e^{-q^2\log q - q^2\log\log(2q) - Cq^2} \le \frac{\mathbb{E} \left| \sum_{n \le x} f(n) \right|^{2q}}{x^q \log^{(q-1)^2} x} \le e^{-q^2\log q - q^2\log\log(2q) + Cq^2}.$$

for a certain absolute constant C. We do *not* know how to prove an asymptotic like (1-1) when q is not a fixed natural number.

¹In this paper we are not particularly concerned with the case where $q \ge \log x/\log \log x$, but for completeness we make a few indicative remarks. Section 6 of Granville and Soundararajan [2001] contains various results on this range of q. Setting $v = \log(2q(\log q)/\log x) \gg 1$, and redoing the calculations on page 2293 with the Rankin shift $1 + q/\log x$ replaced by $1 + v/\log(q \log x)$ and with q^2 -smooth numbers replaced by $q \log x$ -smooth numbers, one can show that $\mathbb{E}|\sum_{n\le x} f(n)|^{2q} \le x^{q(1+v/\log(q \log x)+o(1))}$ uniformly for $q \ge \log x/\log \log x$. In particular, if $q = \log^{1+a} x$ for any fixed $a \ge 0$ then we have $\mathbb{E}|\sum_{n\le x} f(n)|^{2q} \le x^{q(1+a/(a+2)+o(1))}$. By only considering the contribution to the expectation from the event that f(p) is very close to 1 for all primes $p \le q \log x/\log \log x = \log^{2+a} x/\log \log x$, one can obtain a comparable lower bound for $\mathbb{E}|\sum_{n\le x} f(n)|^{2q}$ (as in Corollary 6.3 of Granville and Soundararajan [2001]).

In the Rademacher case, even conjecturally the behavior of $\mathbb{E} \left| \sum_{n \le x} f(n) \right|^{2q}$ is perhaps not obvious. On a wide range of real $q \ge 2$, we might expect that

$$\mathbb{E}\left|\sum_{n\leq x} f(n)\right|^{2q} = e^{-2q^2\log q - 2q^2\log\log(2q) + O(q^2)} x^q \log^{q(2q-3)} x^{q}$$

as in the known asymptotics. But this certainly cannot be the answer for all $1 \le q \le 2$, since on some of that range the exponent q(2q-3) of the logarithm would be negative. (And, by Hölder's inequality, we must at least have $\mathbb{E} \left| \sum_{n \le x} f(n) \right|^{2q} \ge \left(\mathbb{E} \left| \sum_{n \le x} f(n) \right|^2 \right)^q \gg x^q$.)

Theorem 1.2. Let $q_0 = (1 + \sqrt{5})/2 \approx 1.618$. There exists a small absolute constant c > 0 such that the following is true. If f(n) is a Rademacher random multiplicative function, then uniformly for all large x and real $1 \le q \le c \log x / \log \log x$ we have

$$\mathbb{E}\left|\sum_{n\leq x} f(n)\right|^{2q} = e^{-2q^2\log q - 2q^2\log\log(2q) + O(q^2)} \left(1 + \min\{\log\log x, \frac{1}{|q-q_0|}\}\right) x^q \log^{\max\{(q-1)^2, q(2q-3)\}} x.$$

With hindsight, the exponent of $\log x$ we obtain in Theorem 1.2 is perhaps quite natural, since one doesn't expect slower growth in the Rademacher than the Steinhaus case (where there is "more room" for the complex valued random variables to cancel), and we expect q(2q-3) to be the correct exponent eventually. Notice that the golden ratio q_0 is the value at which q(2q-3) becomes larger than $(q-1)^2$. But the additional factor min{log log x, $1/|q - q_0|$ } that appears for q close to q_0 seems genuinely unexpected, and hard to understand except through an inspection of the proof of the theorem.

Next we shall discuss the proofs. Once q is moderately large, namely when $q \ge \log \log x$, we can prove the upper bounds in Theorems 1.1 and 1.2 by fairly simple arguments. See Section 3. This is because, for such q, terms like $\log^{O(q)} x$ can be absorbed into the factor $e^{O(q^2)}$ in our theorems, so we can afford to use simple techniques that are a bit wasteful (e.g., involving Hölder's inequality to reduce to the case of integer q) to reduce matters to a counting problem. Then Rankin's trick is almost sufficient to perform the relevant counts. To obtain the terms $e^{-q^2 \log \log(2q)}$ and $e^{-2q^2 \log \log(2q)}$ in the theorems, we use Rankin's trick along with a slightly more careful treatment of small prime factors.

Our main work is to prove Theorems 1.1 and 1.2 for $1 \le q \le \log \log x$, and also the lower bounds for larger q. Let $F(s) = \sum_{n=1,p|n \Rightarrow p \le x}^{\infty} f(n)/n^s$ denote the Dirichlet series corresponding to f(n), on x-smooth numbers (i.e., numbers with all their prime factors $\le x$). We can also write F(s) as an Euler product, namely $F(s) = \prod_{p \le x} (1 - f(p)/p^s)^{-1}$ in the Steinhaus case and $F(s) = \prod_{p \le x} (1 + f(p)/p^s)$ in the Rademacher case. In the author's treatment [Harper 2017] of low moments, the first step was to show (roughly) that $\mathbb{E} |\sum_{n \le x} f(n)|^{2q} \approx x^q \mathbb{E} (\frac{1}{\log x} \int_{-1/2}^{1/2} |F(\frac{1}{2} + it)|^2 dt)^q$ when $\frac{2}{3} \le q \le 1$. Similarly, our first step here is to show that

$$\mathbb{E} \left| \sum_{n \le x} f(n) \right|^{2q} \approx e^{O(q^2)} x^q \mathbb{E} \left(\frac{1}{\log x} \int_{-1/2}^{1/2} \left| F\left(\frac{1}{2} + \frac{q}{\log x} + it \right) \right|^2 dt \right)^q.$$
(1-2)

Note the shift by $q/\log x$ in the integral, which is analogous to the use of Rankin's trick in our elementary upper bound argument for $q \ge \log \log x$. The basic strategy for proving something like (1-2) is the same as in [Harper 2017], namely conditioning on the behavior of f(n) on smaller primes; using fairly standard moment inequalities, like Khintchine's inequality, to show the conditional expectation behaves like a power of a mean square average; and using Parseval's identity to relate the mean square to an integral average of the Euler product. In [Harper 2017] one could bound terms by using Hölder's inequality to pass to the second moment, whereas here we need suitable rough bounds for high moments. These are supplied by a pair of hypercontractive inequalities, see Probability Result 2.3 in Section 2. Applying the hypercontractive inequalities introduces various divisor functions $d_{\lceil q \rceil}(n)$, $d_{2\lceil q \rceil - 1}(n)$ into our calculations, requiring a bit more number theoretic work as compared with the low moments argument of [Harper 2017]. We refer to the beginning of Section 4 for a rigorous formulation of (1-2), and more technical comparison of this part of the argument with the low moments case [Harper 2017].

Next, we observe that the right-hand side of (1-2) is

$$\approx e^{O(q^2)} x^q \mathbb{E}\left(\frac{1}{\log^2 x} \sum_{|n| \le (\log x)/2} \left|F\left(\frac{1}{2} + \frac{q}{\log x} + i\frac{n}{\log x}\right)\right|^2\right)^q,$$
(1-3)

since heuristically the value of $\left|F\left(\frac{1}{2} + \frac{q}{\log x} + it\right)\right|$ doesn't change much on *t* intervals of length $1/\log x$. One can obtain rigorous statements of this kind using Hölder's inequality in the upper bound arguments, and Jensen's inequality in the lower bound arguments, see Sections 5 and 6. Now we can see heuristically why Theorems 1.1 and 1.2 might hold. In the Steinhaus case, the Euler product F(s) behaves on average like an *L*-function from a *unitary* family, and then since we have $q \ge 1$ (and very differently than in the low moments case [Harper 2017]) the sum over *n* essentially gives us $\log x$ independent tries at obtaining a large value of F(s). So the right-hand side of (1-3) is $\approx e^{O(q^2)}x^q \frac{1}{\log^{2q}x} \log x \mathbb{E} \left|F\left(\frac{1}{2} + \frac{q}{\log x}\right)\right|^{2q} \approx e^{O(q^2)}x^q \frac{1}{\log^{2q-1}x}\left(\frac{\log x}{q\log(2q)}\right)^{q^2}$, as in Theorem 1.1. In the Rademacher case, $F\left(\frac{1}{2} + \frac{q}{\log x} + it\right)$ behaves like an *L*-function from an *orthogonal* family when $t \approx 0$, and like an *L*-function from a unitary family² when $t \approx 1$. Thus, thanks to those $(\log x)/4 \le |n| \le (\log x)/2$ (say) we get a contribution $e^{O(q^2)}x^q \frac{1}{\log^{2q-1}x}\left(\frac{\log x}{q\log(2q)}\right)^{q^2}$ to the right-hand side of (1-3), and thanks to the n = 0 term we get a contribution $\approx e^{O(q^2)}x^q \frac{1}{\log^{2q}x}\mathbb{E} \left|F\left(\frac{1}{2} + \frac{1}{q\log(2q)}\right)^{q^2}$ to the right-hand side of (1-3), and thanks to the n = 0 term we get a contribution $\approx e^{O(q^2)}x^q \frac{1}{\log^{2q}x}\mathbb{E} \left|F\left(\frac{1}{2} + \frac{1}{\log^{2q}x}\right)\right|^{2q}$ are see according to the right form 1.2 arises because of the contribution from intermediate values of n.

To prove the lower bounds in Theorems 1.1 and 1.2 rigorously, as we do in Section 6, roughly speaking it suffices to note that (1-3) is $\geq e^{O(q^2)} x^q \frac{1}{\log^{2q} x} \mathbb{E} \sum_{|n| \leq (\log x)/2} \left| F\left(\frac{1}{2} + \frac{q}{\log x} + i \frac{n}{\log x}\right) \right|^{2q}$, and then compute $\mathbb{E} \left| F\left(\frac{1}{2} + \frac{q}{\log x} + i \frac{n}{\log x}\right) \right|^{2q}$. In practice the details are slightly more complicated because the precise version of (1-3) involves some other terms, including subtracted error terms that must be upper bounded. However,

²At first glance, one might expect $F(\frac{1}{2} + \frac{q}{\log x} + it)$ to behave like a *symplectic L*-function when $t \approx 0$, because averaging over Rademacher f(n) models averaging over quadratic Dirichlet characters. The reason we actually have orthogonal behavior is because we restrict our sums $\sum_{n \leq x} f(n)$ to squarefree terms. For some other contexts where a transition from orthogonal/symplectic to unitary behavior arises, as for large *t* here, see the papers of Florea [2017], Keating and Odgers [2008], and Soundararajan and Young [2010], for example.

we can obtain suitable upper bounds from our main Section 5 argument for proving the upper bounds in Theorems 1.1 and 1.2.

To prove those upper bounds rigorously, we need to capture the fact that typically there will only be a few large terms in the sum over *n* in (1-3). When $q \ge 2$, a careful application of Hölder's inequality lets us bound (1-3) by estimating terms of the form $\mathbb{E} \left| F\left(\frac{1}{2} + \frac{q}{\log x} + i \frac{n}{\log x}\right) \right|^2 \left| F\left(\frac{1}{2} + \frac{q}{\log x} + i \frac{m}{\log x}\right) \right|^{2(q-1)}$. These decrease in size quite rapidly as |m - n| becomes large (and, in the Rademacher case, as |m|, |n| become large), because the parts of the two Euler products over primes $> x^{1/|m-n|}$ become decorrelated rather than reinforcing one another. This indeed says that one doesn't expect large contributions from many different *m*, *n*. When 1 < q < 2, such a direct argument doesn't expect large contributions from many different *m*, *n*. When 1 < q < 2, such a direct argument doesn't over "small" and "large" primes differently, so after a (different) careful application of Hölder's inequality, one is led to expectations where different parts of the Euler product appear to different exponents, to maximize the decorrelation we capture. The most difficult situation is where *q* is very close to 1 (i.e., q = 1 + o(1) as $x \to \infty$). To handle this without picking up any terms that blow up as *q* approaches 1, we use a martingale maximal inequality (see Probability Result 2.5 in Section 2) that essentially lets us maximize over several different splittings of the Euler product simultaneously.

As just described, we go to quite a lot of trouble to prove Theorems 1.1 and 1.2 when q is just a little larger than 1. It is satisfying to have a uniform result (and a method capable of proving one), but in addition this range of q turns out to be relevant for deducing the following corollary.

Corollary 1.3. Let x be large, and let f(n) be a Steinhaus or Rademacher random multiplicative function. For all $2 \le \lambda \le \sqrt{\log x}$, say, we have

$$\mathbb{P}\left(\left|\sum_{n\leq x} f(n)\right| \geq \lambda\sqrt{x}\right) \ll \frac{1}{\lambda^2} e^{-(\log^2 \lambda)/\log\log x}.$$

Proof of Corollary 1.3. For any $1 \le q \le \frac{3}{2}$, say, Theorems 1.1 and 1.2 imply that

$$\mathbb{P}\left(\left|\sum_{n\leq x} f(n)\right| \geq \lambda\sqrt{x}\right) \leq \frac{\mathbb{E}\left|\sum_{n\leq x} f(n)\right|^{2q}}{(\lambda\sqrt{x})^{2q}} \ll \frac{\log^{(q-1)^2} x}{\lambda^{2q}} = \frac{1}{\lambda^2} e^{(q-1)^2 \log\log x - 2(q-1)\log\lambda}.$$

Calculus implies that the right-hand side is minimized if we choose $q - 1 = \frac{\log \lambda}{\log \log x}$, and inserting this choice proves Corollary 1.3.

In the paper [Harper 2017] on low moments, by considering $\mathbb{E} |\sum_{n \le x} f(n)|^{2q}$ with q a little smaller than 1 the author showed that $\mathbb{P}(|\sum_{n \le x} f(n)| \ge z\sqrt{x}/(\log \log x)^{1/4}) \ll \min\{\log z, \sqrt{\log \log x}\}/z^2$ for all $z \ge 2$. Corollary 1.3 is weaker than this when $\lambda \le e^{\sqrt{\log \log x}}$, but stronger for larger λ . In [loc. cit.] the author also showed (see Corollary 2 there, and the subsequent discussion) that $\mathbb{P}(|\sum_{n \le x} f(n)| \ge z\sqrt{x}/(\log \log x)^{1/4}) \gg e^{-(\log^2 z)/\log \log x}/z^2(\log \log x)^{O(1)}$ on a wide range of z. Together all these results give a fairly complete description of the tail behavior of $\sum_{n \le x} f(n)$, up to factors $(\log \log x)^{O(1)}$.

We end this introduction with a few remarks on other possible approaches to Theorems 1.1 and 1.2, and connections with the wider literature.

The quantity $\frac{1}{\log x} \int_{-1/2}^{1/2} |F(\frac{1}{2} + \frac{q}{\log x} + it)|^2 dt$ in (1-2) is closely related to (the total mass of a truncation of) a probabilistic object called *critical multiplicative chaos*. This connection is discussed extensively in the introduction to the low moments paper [Harper 2017], since in that case the techniques for analyzing $\mathbb{E}(\frac{1}{\log x} \int_{-1/2}^{1/2} |F(\frac{1}{2} + \frac{q}{\log x} + it)|^2 dt)^q$ are heavily motivated by ideas from the multiplicative chaos literature. When q > 1 the analogous problem does not seem to have been investigated for critical multiplicative chaos, since the *q*-th moment of the integral will diverge as $x \to \infty$ and this seems to be all the information that was wanted in that case (where the usual interest is in letting $x \to \infty$ and obtaining a limiting measure whose properties can be investigated). Theorems 1.1 and 1.2 show very different behavior in the Steinhaus and Rademacher cases when *q* is large, whereas in the usual problems of multiplicative chaos one finds rather universal behavior (and indeed the Steinhaus and Rademacher moments are of the same order when $q \le 1$).

Assuming the generalized Riemann hypothesis for Dirichlet *L*-functions, Munsch [2017] proved almost sharp upper bounds for the 2*k*-th moment of theta functions $\theta(1, \chi)$ as the character χ varies over nonprincipal Dirichlet characters mod q, for each fixed $k \in \mathbb{N}$. He did this by writing $\theta(1, \chi)$ as a Perron integral involving the *L*-function $L(s, \chi)$, and then expanding the 2*k*-th power and bounding the averages of products $\prod_{j=1}^{2k} |L(\frac{1}{2} + it_j, \chi)|$ that emerge. This is interesting here because for even characters χ , $\theta(1, \chi)$ behaves roughly like $\sum_{n \leq \sqrt{q}} \chi(n)$, which is modeled by the sum $\sum_{n \leq \sqrt{q}} f(n)$ of a Steinhaus random multiplicative function. In our case, using Perron's formula we have

$$\mathbb{E} \bigg| \sum_{n \le x} f(n) \bigg|^{2q} \approx \mathbb{E} \bigg| \frac{1}{2\pi} \int_{-\sqrt{x}}^{\sqrt{x}} F\big(\frac{1}{2} + \frac{q}{\log x} + it\big) \frac{x^{1/2 + q/\log x + it}}{1/2 + q/\log x + it} \, dt \bigg|^{2q} \\ \le x^q e^{O(q^2)} \mathbb{E} \bigg(\int_{-\sqrt{x}}^{\sqrt{x}} \big| F\big(\frac{1}{2} + \frac{q}{\log x} + it\big) \big| \frac{dt}{|1/2 + q/\log x + it|} \bigg)^{2q},$$

say. We already have asymptotics for $\mathbb{E} |\sum_{n \le x} f(n)|^{2q}$ for fixed $q \in \mathbb{N}$, but we might hope to get an alternative proof of sharp upper bounds for $q \notin \mathbb{N}$ by using Hölder's inequality in some way. A direct application, producing a term $|F(\frac{1}{2} + \frac{q}{\log x} + it)|^{2q}$, cannot give sharp bounds because it doesn't recognize that the size of the expectation will be dominated by the integral of $F(\frac{1}{2} + \frac{q}{\log x} + it)$ over a very short (random) *t* interval. To detect this, one could pull out a few (say *d*) copies of the bracket before applying Hölder's inequality to the remaining ones. This would produce a multiple integral of terms of the form $\mathbb{E}(\prod_{j=1}^{d} |F(\frac{1}{2} + \frac{q}{\log x} + it_j)|)|F(\frac{1}{2} + \frac{q}{\log x} + iu)|^{2q-d}$, and the biggest contribution comes when all of the t_j are approximately equal to *u*, so indeed we would capture the localization of the largest contributions. Based on a few rough calculations, it appears this alternative method can prove sharp upper bounds if we take d = 3 (we need to pull out enough terms to adequately detect the localization), and if $q \ge 5$, say. But for smaller *q* this kind of argument doesn't seem operable to prove sharp bounds, indeed one has already lost too much information in applying the triangle inequality to the Perron integral. Nevertheless, it might permit a relatively straightforward extension of Munsch's results [2017] to noninteger $k \ge 5$.

A standard strategy for proving lower bounds is to calculate

$$\mathbb{E}\left(\sum_{n\leq x}f(n)\right)R_{x,q}(f) \quad \text{and} \quad \mathbb{E}|R_{x,q}(f)|^{2q/(2q-1)},$$

where $R_{x,q}(f)$ is some function that is chosen as a proxy for $\left(\sum_{n \le x} f(n)\right)^{2q-1}$ that is easier to understand. Then Hölder's inequality gives

$$\mathbb{E}\left|\sum_{n\leq x} f(n)\right|^{2q} \geq \frac{\left|\mathbb{E}\left(\sum_{n\leq x} f(n)\right) R_{x,q}(f)\right|^{2q}}{(\mathbb{E}|R_{x,q}(f)|^{2q/(2q-1)})^{2q-1}}.$$

If we can estimate the expectations in the numerator and denominator, and $R_{x,q}(f)$ is well chosen so that both of them do behave like $\mathbb{E}|\sum_{n\leq x} f(n)|^{2q}$ (up to scaling factors that would cancel out), then one obtains a sharp lower bound for the 2q-th moment. Munsch and Shparlinski [2016] proved sharp lower bounds for the 2k-th moments of theta functions $\theta(1, \chi)$, for fixed $k \in \mathbb{N}$, by implementing this strategy with a power of a short character sum chosen as the "proxy" object. Our analysis shows that for Rademacher random multiplicative functions, we can imagine heuristically that $|\sum_{n\leq x} f(n)| \approx \frac{\sqrt{x}}{\log x} |F(\frac{1}{2} + \frac{q}{\log x})|$ (when studying 2q-th moments with $q > q_0$). Motivated by this, we could try taking $R_{x,q}(f) = |F(\frac{1}{2} + \frac{q}{\log x})|^{2q-1}$, or perhaps a small variant of this where primes smaller than $q^{O(1)}$ are excluded from the Euler product. In the Rademacher case, rough calculations suggest this will indeed yield the lower bound $\mathbb{E}|\sum_{n\leq x} f(n)|^{2q} \geq e^{O(q^2)}x^q (\log x/(q \log(2q)))^{q(2q-3)}$, which is sharp when $q > q_0$. For smaller q, and in the Steinhaus case, our analysis suggests taking $R_{x,q}(f) = \sum_{|m|\leq \log x} |F(\frac{1}{2} + \frac{q}{\log x} + i \frac{m}{\log x})|^{2q-1}$. This choice actually won't quite work, but rough calculations suggest that comparing $\mathbb{E}|\sum_{n\leq x} f(n)|^2 \sum_{|m|\leq \log x} |F(\frac{1}{2} + \frac{q}{\log x} + i \frac{m}{\log x})|^{2(q-1)}$ will yield sharp lower bounds for $\mathbb{E}|\sum_{n\leq x} f(n)|^{2q}$. This does not seem simpler than our original proofs of the lower bounds in Theorems 1.1 and 1.2, however.

Notation and references. We will say a number *n* is *y*-smooth if all prime factors of *n* are $\leq y$. We will generally use *p* to denote primes. Unless mentioned otherwise, the letters *c*, *C* will denote positive constants, *c* usually being a small constant and *C* a large one. We write f(x) = O(g(x)) and $f(x) \ll g(x)$, both of which mean that there exists *C* such that $|f(x)| \leq Cg(x)$, for all *x*. Sometimes this notation will be adorned with a subscript parameter (e.g., $O_{\epsilon}(\cdot)$ and \ll_{δ}), meaning that the implied constant *C* is allowed to depend on that parameter. We write $f(x) \approx g(x)$ to mean that $g(x) \ll f(x) \ll g(x)$, in other words that $cg(x) \leq |f(x)| \leq Cg(x)$ for some *c*, *C*, for all *x*.

The books of Gut [2013] and of Montgomery and Vaughan [2007] may be consulted as excellent general references for probabilistic and number theoretic background for this paper.

2. Preliminary results

Random Euler products. We begin with some "two point" estimates for the expectation of the 2α -th power of a random Euler product, multiplied by the 2β -th power of an imaginary shift of that product.

These estimates, and small variants of them, will be basic tools throughout our work. The calculations are closely related to computations of shifted moments of L-functions, as in the papers of Chandee [2011] and of Soundararajan and Young [2010], for example.

Euler Product Result 2.1. If f is a Steinhaus random multiplicative function, then for any real α , $\beta \ge 0$, any real $100(1 + \max\{\alpha^2, \beta^2\}) \le x \le y$, and any real $\sigma \ge -1/\log y$ and t, we have

$$\mathbb{E} \prod_{x$$

If we also have $\sigma \leq 1/\log y$, then the above is

$$= e^{O(\max\{\alpha,\beta,\alpha^2,\beta^2\}(1+|t|/\log^{100} x))} \left(\frac{\log y}{\log x}\right)^{\alpha^2+\beta^2} \left(1+\min\left\{\frac{\log y}{\log x},\frac{1}{|t|\log x}\right\}\right)^{2\alpha\beta}$$

In particular, for any real $\alpha \ge 0$, any real $100(1 + \alpha^2) \le x \le y$, and any real $\sigma \ge -1/\log y$, we have

$$\mathbb{E}\prod_{x$$

and if $\sigma \leq 1/\log y$ as well then this is $= e^{O(\max\{\alpha, \alpha^2\})} (\log y/\log x)^{\alpha^2}$.

Proof of Euler Product Result 2.1. For concision in writing the proof, let us temporarily set $M = M(\alpha, \beta) := \max\{\alpha, \beta, \alpha^3, \beta^3\}$.

Firstly, using the Taylor expansion of the logarithm we may rewrite

$$\begin{split} \left|1 - \frac{f(p)}{p^{1/2+\sigma}}\right|^{-2\alpha} \left|1 - \frac{f(p)}{p^{1/2+\sigma+it}}\right|^{-2\beta} \\ &= \exp\left\{-2\alpha \Re \log\left(1 - \frac{f(p)}{p^{1/2+\sigma}}\right) - 2\beta \Re \log\left(1 - \frac{f(p)}{p^{1/2+\sigma+it}}\right)\right\} \\ &= \exp\left\{\frac{2\alpha \Re f(p)}{p^{1/2+\sigma}} + \frac{\alpha \Re f(p)^2}{p^{1+2\sigma}} + \frac{2\beta \Re f(p)p^{-it}}{p^{1/2+\sigma}} + \frac{\beta \Re f(p)^2 p^{-2it}}{p^{1+2\sigma}} + O\left(\frac{\max\{\alpha,\beta\}}{p^{3/2+3\sigma}}\right)\right\}. \end{split}$$

Next, if $y \ge p > x \ge 100 \max\{\alpha^2, \beta^2\}$ then every term in the exponential here has size at most $2 \max\{\alpha, \beta\}/p^{1/2+\sigma} = 2 \max\{\alpha, \beta\}e^{-\sigma \log p}/p^{1/2} \le e/5$. Therefore we may apply the series expansion of the exponential function, finding the above is

$$= 1 + \frac{2(\alpha \Re f(p) + \beta \Re f(p)p^{-it})}{p^{1/2+\sigma}} + \frac{(\alpha \Re f(p)^2 + \beta \Re f(p)^2 p^{-2it})}{p^{1+2\sigma}} + \frac{2(\alpha \Re f(p) + \beta \Re f(p)p^{-it})^2}{p^{1+2\sigma}} + O\left(\frac{M}{p^{3/2+3\sigma}}\right).$$

Now taking expectations, by symmetry we have $\mathbb{E}\Re f(p) = \mathbb{E}\Re f(p)^2 = 0$, similarly for $\mathbb{E}\Re f(p)p^{-it}$ and $\mathbb{E}\Re f(p)^2 p^{-2it}$. A simple trigonometric calculation also shows that $\mathbb{E}(\Re f(p))^2 = \frac{1}{2}$, and similarly $\mathbb{E}\Re f(p)\Re f(p)p^{-it} = \cos(t \log p)/2$. So we get

$$\begin{split} \mathbb{E} \left| 1 - \frac{f(p)}{p^{1/2+\sigma}} \right|^{-2\alpha} \left| 1 - \frac{f(p)}{p^{1/2+\sigma+it}} \right|^{-2\beta} \\ &= 1 + \frac{2\mathbb{E}(\alpha \Re f(p) + \beta \Re f(p)p^{-it})^2}{p^{1+2\sigma}} + O\left(\frac{M}{p^{3/2+3\sigma}}\right) \\ &= 1 + \frac{2(\alpha^2 \mathbb{E}(\Re f(p))^2 + 2\alpha\beta \mathbb{E}\Re f(p)\Re f(p)p^{-it} + \beta^2 \mathbb{E}(\Re f(p)p^{-it})^2)}{p^{1+2\sigma}} + O\left(\frac{M}{p^{3/2+3\sigma}}\right) \\ &= 1 + \frac{\alpha^2 + \beta^2 + 2\alpha\beta \cos(t\log p)}{p^{1+2\sigma}} + O\left(\frac{M}{p^{3/2+3\sigma}}\right) \\ &= \exp\left\{\frac{\alpha^2 + \beta^2 + 2\alpha\beta \cos(t\log p)}{p^{1+2\sigma}} + O\left(\frac{M}{p^{3/2+3\sigma}}\right)\right\}. \end{split}$$

Combining the above calculation with the independence of f on distinct primes, and using that, for $p \le y$,

$$p^{3/2+\sigma} = e^{\sigma \log p} p^{3/2} \ge e^{-1} p^{3/2}$$
 and $\sum_{p>x} \frac{1}{p^{3/2}} \ll \frac{1}{\sqrt{x} \log x}$,

we deduce that the quantity

$$\mathbb{E}\prod_{x$$

in the statement of the result is

$$= \exp\bigg\{\sum_{x
$$= \exp\bigg\{\sum_{x$$$$

To deduce the second part of Euler Product Result 2.1, we can use standard estimates from prime number theory. Indeed, the Chebychev and Mertens estimates for sums over primes imply that

$$\sum_{x
$$= (\alpha^2 + \beta^2) \log\left(\frac{\log y}{\log x}\right) + O(\max\{\alpha^2, \beta^2\}),$$$$

using that $e^{-2\sigma \log p} - 1 \ll |\sigma| \log p \ll \log p/\log y$ for $|\sigma| \le 1/\log y$. We may remove the nuisance factor $p^{2\sigma}$ from the sum $\sum_{x with the same error term. Then using the prime$

number theorem in the form $\pi(z) := \#\{p \le z : p \text{ prime}\} = \int_2^z 1/\log u \, du + O(z/(\log^{100} z))$, we have

$$\sum_{x
$$= \int_x^y \frac{\cos(t \log z)}{z \log z} dz + O\left(\frac{1+|t|}{\log^{100} x}\right)$$
$$= \int_{\log x}^{\log y} \frac{\cos(tu)}{u} du + O\left(\frac{1+|t|}{\log^{100} x}\right).$$$$

Now if $|t| \log y \le 1$, then the estimate $\cos(tu) = 1 + O((tu)^2)$ shows the integral is $\log \log y - \log \log x + O((t \log y)^2) = \log(\log y/\log x) + O(1)$. If instead we have $|t| \log x \le 1$ but $|t| \log y > 1$, then we can evaluate the part of the integral with $u \le 1/|t|$ using the estimate $\cos(tu) = 1 + O((tu)^2)$, and estimate the rest using integration by parts, yielding an overall estimate $\log(1/(|t| \log x)) + O(1)$. If $|t| \log x > 1$ then integration by parts shows the whole integral is O(1). In any case, the second part of Euler Product Result 2.1 is proved.

The third part follows by setting $\beta = 0$ in the preceding statements.

We will need a version of the above result for Rademacher random multiplicative functions. Unlike in the Steinhaus case, the distribution of $f(n)n^{-it}$ is not the same for all real t in the Rademacher case, so our general statement must allow two different imaginary shifts in our two Euler product factors.

Euler Product Result 2.2. If f is a Rademacher random multiplicative function, then for any real $\alpha, \beta \ge 0$, any real $100(1 + \max\{\alpha^2, \beta^2\}) \le x \le y$, and any real $\sigma \ge -1/\log y$ and t_1, t_2 , we have

$$\mathbb{E} \prod_{x$$

If we also have $\sigma \leq 1/\log y$, then the above is

$$=e^{O(\max\{\alpha,\beta,\alpha^{2},\beta^{2}\}(1+\frac{|t_{1}|+|t_{2}|}{\log^{100}x}))}\left(1+\min\left\{\frac{\log y}{\log x},\frac{|t_{1}|^{-1}}{\log x}\right\}\right)^{\alpha^{2}-\alpha}\left(1+\min\left\{\frac{\log y}{\log x},\frac{|t_{2}|^{-1}}{\log x}\right\}\right)^{\beta^{2}-\beta}\cdot\left(\frac{\log y}{\log x}\right)^{\alpha^{2}+\beta^{2}}\left(\left(1+\min\left\{\frac{\log y}{\log x},\frac{|t_{1}+t_{2}|^{-1}}{\log x}\right\}\right)\left(1+\min\left\{\frac{\log y}{\log x},\frac{|t_{1}-t_{2}|^{-1}}{\log x}\right\}\right)\right)^{2\alpha\beta}.$$

As an upper bound, we may replace the error term $e^{O(\max\{\alpha,\beta,\alpha^2,\beta^2\}(1+(|t_1|+|t_2|)/(\log^{100}x)))}$ by

$$e^{O(\max\{\alpha,\beta,\alpha^2,\beta^2\})}\min\left\{\frac{\log y}{\log x},\,1+\frac{(|t_1|+|t_2|)^{1/100}}{\log x}\right\}^{|\alpha^2-\alpha|+|\beta^2-\beta|+4\alpha\beta},$$

and as a lower bound we may replace it by

$$e^{O(\max\{\alpha,\beta,\alpha^2,\beta^2\})}\min\left\{\frac{\log y}{\log x}, 1+\frac{(|t_1|+|t_2|)^{1/100}}{\log x}\right\}^{-(|\alpha^2-\alpha|+|\beta^2-\beta|+4\alpha\beta)}.$$

The estimation of the error terms here is rather crude, but will be sufficient as they only depend quite mildly on the t_i .

Proof of Euler Product Result 2.2. The proof is a fairly straightforward adaptation of the proof of Euler Product Result 2.1. We again temporarily set $M = M(\alpha, \beta) := \max\{\alpha, \beta, \alpha^3, \beta^3\}$. In the first place we have

$$\begin{split} \left| 1 + \frac{f(p)}{p^{1/2 + \sigma + it_1}} \right|^{2\alpha} \left| 1 + \frac{f(p)}{p^{1/2 + \sigma + it_2}} \right|^{2\beta} \\ &= \exp\left\{ 2\alpha \Re \log\left(1 + \frac{f(p)}{p^{1/2 + \sigma + it_1}}\right) + 2\beta \Re \log\left(1 + \frac{f(p)}{p^{1/2 + \sigma + it_2}}\right) \right\} \\ &= \exp\left\{ \frac{2\alpha \Re f(p) p^{-it_1}}{p^{1/2 + \sigma}} - \frac{\alpha \Re f(p)^2 p^{-2it_1}}{p^{1+2\sigma}} + \frac{2\beta \Re f(p) p^{-it_2}}{p^{1/2 + \sigma}} - \frac{\beta \Re f(p)^2 p^{-2it_2}}{p^{1+2\sigma}} + O\left(\frac{\max\{\alpha, \beta\}}{p^{3/2 + 3\sigma}}\right) \right\} \\ &= 1 + \frac{2(\alpha \Re f(p) p^{-it_1} + \beta \Re f(p) p^{-it_2})}{p^{1/2 + \sigma}} - \frac{(\alpha \Re f(p)^2 p^{-2it_1} + \beta \Re f(p)^2 p^{-2it_2})}{p^{1+2\sigma}} + \frac{2(\alpha \Re f(p) p^{-it_1} + \beta \Re f(p) p^{-it_2})^2}{p^{1+2\sigma}} + O\left(\frac{M}{p^{3/2 + 3\sigma}}\right). \end{split}$$

Furthermore, in the Rademacher case we have $f(p)^2 \equiv 1$, whilst still

$$\mathbb{E}\Re f(p)p^{-it} = \cos(t\log p)\mathbb{E}f(p) = 0.$$

So we get

$$\mathbb{E} \left| 1 + \frac{f(p)}{p^{1/2 + \sigma + it_1}} \right|^{2\alpha} \left| 1 + \frac{f(p)}{p^{1/2 + \sigma + it_2}} \right|^{2\beta} \\ = 1 - \frac{(\alpha \cos(2t_1 \log p) + \beta \cos(2t_2 \log p))}{p^{1 + 2\sigma}} + \frac{2(\alpha \cos(t_1 \log p) + \beta \cos(t_2 \log p))^2}{p^{1 + 2\sigma}} + O\left(\frac{M}{p^{3/2 + 3\sigma}}\right),$$

and using standard cosine identities this is all

$$= 1 + \frac{\alpha^{2} + \beta^{2} + (\alpha^{2} - \alpha)\cos(2t_{1}\log p) + (\beta^{2} - \beta)\cos(2t_{2}\log p)}{p^{1+2\sigma}} + \frac{2\alpha\beta(\cos((t_{1} + t_{2})\log p) + \cos((t_{1} - t_{2})\log p)))}{p^{1+2\sigma}} + O\left(\frac{M}{p^{3/2+3\sigma}}\right)$$
$$= \exp\left\{\frac{\alpha^{2} + \beta^{2} + (\alpha^{2} - \alpha)\cos(2t_{1}\log p) + (\beta^{2} - \beta)\cos(2t_{2}\log p)}{p^{1+2\sigma}} + \frac{2\alpha\beta(\cos((t_{1} + t_{2})\log p) + \cos((t_{1} - t_{2})\log p)))}{p^{1+2\sigma}} + O\left(\frac{M}{p^{3/2+3\sigma}}\right)\right\}.$$

The first two conclusions of Euler Product Result 2.2 now follow exactly as in the proof of Euler Product Result 2.1.

For the final claimed inequalities, we note that the source of the unwanted error term

$$O\left(\max\{\alpha, \beta, \alpha^2, \beta^2\} \frac{|t_1| + |t_2|}{\log^{100} x}\right)$$

in the exponent lies in our using the prime number theorem to estimate the various sums

$$\sum_{x
$$\sum_{x
$$\sum_{x
$$\sum_{x$$$$$$$$

Instead, if $|t_1| \ge \log^{100} x$ (which is the only case where it might produce a large error term) we can upper bound $\sum_{x by$

$$\sum_{\substack{x$$

As in the proof of Euler Product Result 2.1, the second sum here is $\ll \max\{\alpha, \alpha^2\}$ (we can use the prime number theorem to estimate it, since the lower end point is now sufficiently large that we don't pick up a big error term), and the first sum is

$$|\alpha^{2} - \alpha| \left(\sum_{x$$

We can handle the other sums similarly when t_2 , $t_1 + t_2$, $t_1 - t_2$ are large. In the worst case, as an upper bound this will produce an extra multiplicative factor

$$\exp\left\{(|\alpha^2 - \alpha| + |\beta^2 - \beta| + 4\alpha\beta) \min\left\{\log\left(\frac{\log y}{\log x}\right), \log\left(1 + \frac{(|t_1| + |t_2|)^{1/100}}{\log x}\right)\right\}\right\}.$$

An exactly similar argument gives a lower bound with $(|\alpha^2 - \alpha| + |\beta^2 - \beta| + 4\alpha\beta)$ replaced by $-(|\alpha^2 - \alpha| + |\beta^2 - \beta| + 4\alpha\beta)$.

Probabilistic preparations. Next we record some moment estimates, mostly fairly simple yet interesting, that will be input to our arguments in various places.

Probability Result 2.3 (Rough hypercontractive inequalities). For any real $q \ge 1$, the following is true.

If f(n) is a Steinhaus random multiplicative function, then for any sequence of complex numbers $(a_n)_{n \leq N}$ we have

$$\mathbb{E}\left|\sum_{n\leq N}a_nf(n)\right|^{2q}\leq \left(\sum_{n\leq N}|a_n|^2d_{\lceil q\rceil}(n)\right)^q,$$

where $d_k(\cdot)$ denotes the k-fold divisor function (i.e., the number of k-tuples of natural numbers whose product is \cdot , or equivalently the Dirichlet series coefficient of $\zeta(s)^k$), and $\lceil q \rceil$ denotes the ceiling of q.

If f(n) is a Rademacher random multiplicative function, then for any sequence of complex numbers $(a_n)_{n \leq N}$ we have

$$\mathbb{E}\left|\sum_{n\leq N}a_nf(n)\right|^{2q}\leq \left(\sum_{n\leq N}|a_n|^2d_{2\lceil q\rceil-1}(n)\right)^q.$$

Proof of Probability Result 2.3. By Hölder's inequality, for any real $q \ge 1$ we have

$$\mathbb{E}\left|\sum_{n\leq N}a_nf(n)\right|^{2q}\leq \left(\mathbb{E}\left|\sum_{n\leq N}a_nf(n)\right|^{2\lceil q\rceil}\right)^{q/\lceil q\rceil},$$

so (replacing q by $\lceil q \rceil$) it suffices to treat the case where q is a natural number.

For Steinhaus f(n), expanding the 2q-th power and taking expectations we get

$$\mathbb{E}\left|\sum_{n\leq N}a_nf(n)\right|^{2q}=\sum_{n_1,\ldots,n_q\leq N}a_{n_1}\cdots a_{n_q}\sum_{m_1,\ldots,m_q\leq N}\overline{a_{m_1}\cdots a_{m_q}}\mathbf{1}_{\prod_{i=1}^q n_i=\prod_{i=1}^q m_i},$$

where 1 denotes the indicator function. Using the upper bound

$$|a_{n_1}\cdots a_{n_q}\overline{a_{m_1}\cdots a_{m_q}}| \le (\frac{1}{2})(|a_{n_1}\cdots a_{n_q}|^2 + |a_{m_1}\cdots a_{m_q}|^2),$$

together with the symmetry of the n_i and the m_i , we deduce that

$$\mathbb{E}\left|\sum_{n\leq N}a_{n}f(n)\right|^{2q} \leq \sum_{n_{1},\dots,n_{q}\leq N}|a_{n_{1}}\cdots a_{n_{q}}|^{2}\sum_{m_{1},\dots,m_{q}\leq N}\mathbf{1}_{\prod_{i=1}^{q}n_{i}=\prod_{i=1}^{q}m_{i}} \leq \sum_{n_{1},\dots,n_{q}\leq N}|a_{n_{1}}\cdots a_{n_{q}}|^{2}d_{q}\left(\prod_{i=1}^{q}n_{i}\right).$$

Finally, since the function $d_q(\cdot)$ is submultiplicative we find the above is

$$\leq \sum_{n_1,...,n_q \leq N} |a_{n_1} \cdots a_{n_q}|^2 d_q(n_1) \cdots d_q(n_q) = \left(\sum_{n \leq N} |a_n|^2 d_q(n)\right)^q.$$

In the Rademacher case, one needs a bit more involved argument. We refer the reader to Lemma 2 of Halász [1983], where this result is proved by induction on the exponent 2q. We may remark that, since Rademacher f(n) is only supported on squarefree n, we may assume that a_n is only nonzero for squarefree n, and then $d_{2\lceil q\rceil-1}(n) = (2\lceil q\rceil-1)^{\Omega(n)}$ where $\Omega(n)$ is the number of prime factors of n. The ultimate source of the factors $d_{2\lceil q\rceil-1}(n)$ is that, when one expands the expectation in the inductive proof,

the only surviving terms are those where the product $n_1 \cdots n_{2q}$ is a perfect square, so all the prime factors of n_{2q} must be repeated somewhere amongst the other terms n_1, \ldots, n_{2q-1} .

We describe the inequalities in Probability Result 2.3 as "rough hypercontractive inequalities" because (if we take 2*q*-th roots of both sides) they upper bound an L^{2q} -norm by a weighted L^2 norm without any other terms, but the weights $d_{\lceil q \rceil}(n)$, $d_{2\lceil q \rceil - 1}(n)$ will not generally be the sharpest possible unless *q* is an integer. One can prove more precise results for noninteger *q* using more subtle interpolation techniques, see Section 2 of Bondarenko, Brevig, Saksman, Seip and Zhao [Bondarenko et al. 2018] for the Steinhaus case, and Chapitre III of Bonami [1970] for the Rademacher case (expressed in rather different notation). However, for our applications the extra precision in these inequalities will not be needed.

Probability Result 2.4. Let $(\epsilon_n)_{n \le N}$ be a sequence of independent random variables, each satisfying $\mathbb{E}\epsilon_n = 0$ and $\mathbb{E}|\epsilon_n|^2 = 1$, and let $(a_n)_{n \le N}$ be a sequence of complex numbers. Then for any real $q \ge 1$, we have

$$\mathbb{E}\left|\sum_{n\leq N}a_{n}\epsilon_{n}\right|^{2q}\geq\left(\sum_{n\leq N}|a_{n}|^{2}\right)^{q}$$

Proof of Probability Result 2.4. Since we assume that $q \ge 1$, simply applying Hölder's inequality we get

$$\left(\sum_{n\leq N}|a_n|^2\right)^q = \left(\mathbb{E}\left|\sum_{n\leq N}a_n\epsilon_n\right|^2\right)^q \leq \mathbb{E}\left|\sum_{n\leq N}a_n\epsilon_n\right|^{2q}.$$

If the ϵ_n are Rademacher or Steinhaus random variables,³ then Khintchine's inequality (see e.g., Lemma 3.8.1 of Gut [2013]) in fact implies that $\mathbb{E} |\sum_{n \le N} a_n \epsilon_n|^{2q} \asymp_q (\sum_{n \le N} |a_n|^2)^q$ for all real $q \ge 0$. For our purposes here we will only require the simple lower bound in Probability Result 2.4, but it is useful to keep Khintchine's inequality in mind since it means that when we apply the lower bound, we are doing something sharp.

The final result we shall record is more sophisticated, and requires some terminology before we can state it. Suppose that $(\Omega, \mathcal{F}, \mathbb{P})$ is a probability space, and $(\mathcal{F}_n)_{n\geq 0}$ is a *filtration* on \mathcal{F} , in other words a sequence of sub- σ -algebras satisfying $\mathcal{F}_0 \subseteq \mathcal{F}_1 \subseteq \cdots \subseteq \mathcal{F}$. We say a sequence of random variables $(X_n)_{n\geq 0}$ on $(\Omega, \mathcal{F}, \mathbb{P})$ is a *submartingale* (relative to $(\mathcal{F}_n)_{n\geq 0}$ and \mathbb{P}) if it satisfies:

- (i) (adapted) X_n is measurable with respect to \mathcal{F}_n , for all $n \ge 0$.
- (ii) (integrable) $\mathbb{E}|X_n|$ is finite, for all $n \ge 0$;
- (iii) (nondecreasing on average) For all $n \ge 1$, the conditional expectation $\mathbb{E}(X_n | \mathcal{F}_{n-1}) \ge X_{n-1}$ almost surely.

Condition (iii) says that a submartingale is nondecreasing on average, in quite a strong sense: for any given value of X_{n-1} (or, informally speaking, any other "information" from the sigma algebra \mathcal{F}_{n-1}), the conditional expectation of X_n will be at least as large. One can apply this property to partition the sample

³We emphasize that here we are referring to ordinary Rademacher or Steinhaus random variables, not random *multiplicative* functions.

space Ω in useful ways, and prove that the moments of the random variables comprising a submartingale satisfy the following useful bound. We will use this as an ingredient in proving our 2*q*-th moment upper bounds when *q* is close to 1.

Probability Result 2.5 (Doob's L^p maximal inequality, see Theorem 9.4 of Gut [2013]). Let $(X_n)_{n\geq 0}$ be a nonnegative submartingale (on some probability space and with respect to some filtration). Then for any p > 1, we have

$$\mathbb{E}(\max_{0 \le k \le n} X_k)^p \le \left(\frac{p}{p-1}\right)^p \mathbb{E}X_n^p$$

Some miscellaneous lemmas. As in the first paper [Harper 2017] in this sequence, we will need the following version of Parseval's identity for Dirichlet series to help with relating $\mathbb{E} |\sum_{n \le x} f(n)|^{2q}$ to an Euler product average.

Harmonic Analysis Result 2.6 (See (5.26) in Section 5.1 of [Montgomery and Vaughan 2007]). Let $(a_n)_{n=1}^{\infty}$ be any sequence of complex numbers, and let $A(s) := \sum_{n=1}^{\infty} a_n/n^s$ denote the corresponding Dirichlet series, and σ_c denote its abscissa of convergence. Then for any $\sigma > \max\{0, \sigma_c\}$, we have

$$\int_0^\infty \frac{\left|\sum_{n \le x} a_n\right|^2}{x^{1+2\sigma}} dx = \frac{1}{2\pi} \int_{-\infty}^\infty \left|\frac{A(\sigma+it)}{\sigma+it}\right|^2 dt$$

We will use the following estimate to handle sums of divisor-type functions that appear in our calculations.

Number Theory Result 2.7. Let $0 < \delta < 1$, let $m \ge 1$, and suppose that $\max\{3, 2m\} \le y \le z \le y^{10}$ and that $1 < u \le v(1 - y^{-\delta})$. As usual, let $\Omega(d)$ denote the total number of prime factors of d (counted with multiplicity). Then

$$\sum_{\substack{u \le d \le v, \\ p \mid d \Rightarrow y \le p \le z}} m^{\Omega(d)} \ll_{\delta} \frac{(v-u)m}{\log y} \prod_{y \le p \le z} \left(1 - \frac{m}{p}\right)^{-1}.$$

This is a slight generalization of a result of Lau, Tenenbaum and Wu [2013, Lemma 2.1] (see also [Halász 1983, Lemma 3]). See [Harper 2017, Section 2.1] for the full (short) proof.

3. Easier cases of the theorems

As remarked in the Introduction, since we allow a multiplicative error term $e^{O(q^2)}$ in our theorems, it turns out that proving our claimed upper bounds when $\log \log x \le q \le c \log x/\log \log x$ is somewhat straightforward. We present these arguments in this section. Some of the techniques involved, including the use of Rankin's trick with an exponent roughly like $1 + q/\log x$, and a special treatment of prime factors that are $\ll q^2$, will recur later when we develop our main arguments.

The upper bound in the Steinhaus case, for very large q. For $q \ge \log \log x$ we have $\log^{(q-1)^2} x = \log^{q^2+O(q)} x = e^{O(q^2)} \log^{q^2} x$. Thus to establish the upper bound part of Theorem 1.1 for $\log \log x \le q \le c \log x/\log \log x$, it will suffice to show that

$$\left\|\sum_{n \le x} f(n)\right\|_{2q} \le e^{-(q/2)\log q - (q/2)\log\log(2q) + O(q)}\sqrt{x}\log^{q/2} x,$$

where as usual we write $\|\cdot\|_r := (\mathbb{E}|\cdot|^r)^{1/r}$.

To prove this, we first apply Minkowski's inequality to obtain that

$$\left\|\sum_{n \le x} f(n)\right\|_{2q} = \left\|\sum_{\substack{m \le x, \\ m \text{ is } q^2 \text{ smooth }}} f(m) \sum_{\substack{n \le x/m, \\ p|n \Rightarrow p > q^2}} f(n)\right\|_{2q} \le \sum_{\substack{m \le x, \\ m \text{ is } q^2 \text{ smooth }}} \left\|\sum_{\substack{n \le x/m, \\ p|n \Rightarrow p > q^2}} f(n)\right\|_{2q}$$

Recall here that a number is said to be q^2 -smooth if all of its prime factors are $\leq q^2$. Using the first part of Probability Result 2.3, and then using Rankin's trick of upper bounding $\mathbf{1}_{n \leq x/m}$ by $(x/(nm))^{1+q/\log x}$ (and recalling that the divisor function $d_{\lceil q \rceil}(n)$ is the Dirichlet series coefficient of $\zeta(s)^{\lceil q \rceil} = \sum_{n=1}^{\infty} d_{\lceil q \rceil}(n)/n^s = \prod_p (1-1/p^s)^{-\lceil q \rceil}$), we get

$$\begin{split} \left\| \sum_{\substack{n \le x/m, \\ p \mid n \Rightarrow p > q^2}} f(n) \right\|_{2q} &\leq \left(\sum_{\substack{n \le x/m, \\ p \mid n \Rightarrow p > q^2}} d_{\lceil q \rceil}(n) \right)^{1/2} \\ &\leq \left(\left(\frac{x}{m} \right)^{1+q/\log x} \sum_{\substack{n \le x/m, \\ p \mid n \Rightarrow p > q^2}} \frac{d_{\lceil q \rceil}(n)}{n^{1+q/\log x}} \right)^{1/2} \\ &\leq \sqrt{\frac{x}{m}} e^{O(q)} \prod_{p > q^2} \left(1 - \frac{1}{p^{1+q/\log x}} \right)^{-\lceil q \rceil/2}. \end{split}$$

Finally, the product over primes here is $\zeta(1+q/\log x)^{\lceil q \rceil/2} \prod_{p \le q^2} (1-1/p^{1+q/\log x})^{\lceil q \rceil/2}$. Using the fact that the zeta function has a simple pole at 1, this equals $e^{O(q)}(\log x/q)^{\lceil q \rceil/2}e^{-\sum_{p \le q^2} (\lceil q \rceil/2)/(p^{1+q/\log x})} = e^{O(q)}(\log x/q \log q)^{q/2}$ on our range log log $x \le q \le c \log x/\log \log x$. And when we sum over *m* we have $\sum_{m \le x,m \text{ is } q^2 \text{ smooth }} 1/\sqrt{m} \le e^{\sum_{p \le q^2} O(1/\sqrt{p})} \le e^{O(q)}$, so putting everything together we get an acceptable upper bound for $\|\sum_{n \le x} f(n)\|_{2q}$.

The upper bound in the Rademacher case, for very large q. Similarly as in the Steinhaus case, to prove the upper bound part of Theorem 1.2 for $\log \log x \le q \le c \log x / \log \log x$ it will suffice to show that, for Rademacher random multiplicative f(n), we have

$$\left\|\sum_{n \le x} f(n)\right\|_{2q} \le e^{-q \log q - q \log \log(2q) + O(q)} \sqrt{x} \log^q x.$$

Using Minkowski's inequality, the second part of Probability Result 2.3, and then Rankin's trick, we get

$$\begin{split} \sum_{n \le x} f(n) \Big\|_{2q} &\le \sum_{\substack{m \le x, \\ m \text{ is } q^2 \text{ smooth }}} \Big\| \sum_{\substack{n \le x/m, \\ p \mid n \Rightarrow p > q^2}} f(n) \Big\|_{2q} \\ &\le \sum_{\substack{m \le x, \\ m \text{ is } q^2 \text{ smooth }}} \left(\sum_{\substack{n \le x/m, \\ p \mid n \Rightarrow p > q^2}} d_{2\lceil q \rceil - 1}(n) \right)^{1/2} \\ &\le \sum_{\substack{m \le x, \\ m \text{ is } q^2 \text{ smooth }}} \sqrt{\frac{x}{m}} e^{O(q)} \prod_{p > q^2} \left(1 - \frac{1}{p^{1+q/\log x}} \right)^{-(2\lceil q \rceil - 1)/2}. \end{split}$$

We can estimate the product over primes as in the Steinhaus case, finding it is

$$e^{O(q)} \left(\frac{\log x}{q}\right)^{(2\lceil q\rceil - 1)/2} e^{-\sum_{p \le q^2} ((2\lceil q\rceil - 1)/2)/p^{1+q/\log x}} = e^{O(q)} \left(\frac{\log x}{q\log q}\right)^q$$

on our range $\log \log x \le q \le c \log x / \log \log x$. And when we sum over *m* we again have

$$\sum_{\substack{m \le x, \\ \text{is } q^2 \text{ smooth}}} \frac{1}{\sqrt{m}} \le e^{\sum_{p \le q^2} O(1/\sqrt{p})} \le e^{O(q)},$$

so putting everything together we get an acceptable upper bound for $\left\|\sum_{n < x} f(n)\right\|_{2a}$.

т

4. The reduction to Euler products

In this section we shall prove four propositions that make precise the assertion in (1-2) that $\mathbb{E} \left| \sum_{n \le x} f(n) \right|^{2q}$ may be bounded by studying integrals of Euler products.

Upper bounds: statement of the propositions. We will need a little notation, which is exactly the same as in the author's previous paper [Harper 2017] dealing with low moments. Given a random multiplicative function f(n) (either Steinhaus or Rademacher, depending on the context), and an integer $0 \le k \le \log \log x$, let F_k denote the partial Euler product of f(n) over $x^{e^{-(k+1)}}$ -smooth numbers. Thus for all complex *s* with $\Re(s) > 0$, we have

$$F_k(s) = \prod_{p \le x^{e^{-(k+1)}}} \left(1 - \frac{f(p)}{p^s}\right)^{-1} = \sum_{\substack{n=1, \\ n \text{ is } x^{e^{-(k+1)}} \text{ smooth}}}^{\infty} \frac{f(n)}{n^s}$$

in the Steinhaus case, and

$$F_k(s) = \prod_{p \le x^{e^{-(k+1)}}} \left(1 + \frac{f(p)}{p^s} \right) = \sum_{\substack{n=1, \\ n \text{ is } x^{e^{-(k+1)}} \text{ smooth}}}^{\infty} \frac{f(n)}{n^s}$$

in the Rademacher case (the product taking a different form because f(n) is only supported on squarefree numbers in that case).

Proposition 4.1. Let f(n) be a Steinhaus random multiplicative function, let x be large, and set $\mathcal{L} := \lfloor (\log \log x)/10 \rfloor$. Uniformly for all $1 \le q \le \log^{0.05} x$, we have

$$\left\|\sum_{n \le x} f(n)\right\|_{2q} \le \sqrt{\frac{x}{\log x}} e^{O(q)} \sum_{0 \le k \le \mathcal{L}} \left\|\int_{-1/2}^{1/2} \left|F_k\left(\frac{1}{2} + \frac{q-k}{\log x} + it\right)\right|^2 dt\right\|_q^{1/2} + e^{O(q)} \sqrt{\frac{x}{\log x}} + e^{O(q)} \sqrt{$$

In the low moments case, Proposition 1 of [Harper 2017] gives an analogous upper bound for all $\frac{2}{3} \le q \le 1$, but with the quantity \mathcal{L} replaced by the smaller quantity $\mathcal{K} = \lfloor \log \log \log x \rfloor$, and the shift $(q-k)/\log x$ in the Euler product replaced by $-k/\log x$.

The additional shift by $q/\log x$ here corresponds to applying Rankin's trick with exponent $1 + q/\log x$ in our treatment of very large q in Section 3. We can introduce this at the acceptable cost of a prefactor $e^{O(q)}$ in the proposition, and it means that when we analyze the Euler product we can restrict attention to numbers that are $x^{1/q}$ -smooth, which is crucial to obtaining the desired factor $e^{-q^2 \log q}$ in Theorem 1.1. The significant contribution from very smooth numbers, when q becomes large, also explains why we must let k run over a wider range than in the low moments case to obtain acceptable bounds. Finally, we remark that the range $1 \le q \le \log^{0.05} x$ allowed in Proposition 4.1 is somewhat artificial, but more than sufficient since we already proved the Theorem 1.1 upper bound for all $\log \log x \le q \le c \log x/\log \log x$ in Section 3. It could be increased somewhat, but it seems hard to obtain an upper bound of a similar shape to Proposition 4.1 on the full range $1 \le q \le c \log x/\log \log x$, since for very large q the significant contribution from very smooth numbers changes the behavior in parts of the proof.

Proposition 4.2. Let f(n) be a Rademacher random multiplicative function, let x be large, and set $\mathcal{L} := \lfloor (\log \log x)/10 \rfloor$. Uniformly for all $1 \le q \le \log^{0.05} x$, we have

$$\left\| \sum_{n \le x} f(n) \right\|_{2q} \le \sqrt{\frac{x}{\log x}} e^{O(q)} \sum_{0 \le k \le \mathcal{L}} \max_{N \in \mathbb{Z}} \frac{1}{(|N|+1)^{1/8}} \left\| \int_{N-1/2}^{N+1/2} \left| F_k \left(\frac{1}{2} + \frac{q-k}{\log x} + it \right) \right|^2 dt \right\|_q^{1/2} + e^{O(q)} \sqrt{\frac{x}{\log x}}.$$

One has to deal with translates by N in the Rademacher case because, unlike in the Steinhaus case, the distribution of $(f(n)n^{it})$ is *not* the same (for $t \neq 0$) as the distribution of (f(n)) for Rademacher random multiplicative f(n). However, as in the low moments argument in [Harper 2017], the main contribution will come from small N.

Lower bounds: statement of the propositions. For our work on lower bounds, we again connect the size of $\|\sum_{n \le x} f(n)\|_{2q}$ with a certain integral average, and thence with random Euler products. Let *F* denote the partial Euler product of f(n), either Steinhaus or Rademacher, over *x*-smooth numbers. (Thus $F = F_{-1}$, if we slightly abuse our earlier notation).

Proposition 4.3. If f(n) is a Steinhaus random multiplicative function, and x is large, then uniformly for all $q \ge 1$ we have

$$\left\|\sum_{n \le x} f(n)\right\|_{2q} \gg \sqrt{\frac{x}{\log x}} \left\|\int_{1}^{x^{1/4}} \left|\sum_{m \le z} f(m)\right|^{2} \frac{dz}{z^{2}}\right\|_{q}^{1/2}.$$

In particular, for any large quantity $V \leq (\log x)/q$ we have that $\left\|\sum_{n \leq x} f(n)\right\|_{2q}$ is

$$\gg \sqrt{\frac{x}{\log x}} \left(\left\| \int_{-1/2}^{1/2} \left| F\left(\frac{1}{2} + \frac{4Vq}{\log x} + it\right) \right|^2 dt \right\|_q^{1/2} - \frac{C}{e^{Vq/2}} \left\| \int_{-1/2}^{1/2} \left| F\left(\frac{1}{2} + \frac{2Vq}{\log x} + it\right) \right|^2 dt \right\|_q^{1/2} \right),$$

where C > 0 is an absolute constant.

Notice that we don't need to impose any upper bound on q here (although, for the second statement, there is an implicit upper bound $q \ll \log x$ in order that we can choose large $V \le (\log x)/q$). This means we can use Proposition 4.3 to prove the lower bound in Theorem 1.1 on the full range of q there.

Proposition 4.4. If f(n) is a Rademacher random multiplicative function, then the first bound in *Proposition 4.3 continues to hold, and the second bound may be replaced by the statement that*

$$\begin{split} \left\|\sum_{n \le x} f(n)\right\|_{2q} \gg \sqrt{\frac{x}{\log x}} \left(\left\|\int_{-1/2}^{1/2} \left|F\left(\frac{1}{2} + \frac{4Vq}{\log x} + it\right)\right|^2 dt\right\|_q^{1/2} \\ &- \frac{C}{e^{Vq/2}} \max_{N \in \mathbb{Z}} \frac{1}{(|N|+1)^{1/8}} \left\|\int_{N-1/2}^{N+1/2} \left|F\left(\frac{1}{2} + \frac{2Vq}{\log x} + it\right)\right|^2 dt\right\|_q^{1/2} \right). \end{split}$$

These results are again of the same general shape as the corresponding Propositions 3 and 4 of [Harper 2017] from the low moments case. In fact, the propositions here are a little simpler as they don't involve an additional subtracted error term $-C\sqrt{x/\log x}$. This is accomplished by some reorganization of the proof, and shrinking the range of integration over z to $[1, x^{1/4}]$ rather than $[1, \sqrt{x}]$ from the low moments case, which makes no difference when applying the results. The other difference, similarly as in Section 3 and in our discussion of upper bounds, is that here we introduce shifts of the shape $4Vq/\log x$ in our Euler products, as opposed to $4V/\log x$ in the low moments analogues.

Proof of Propositions 4.1 and 4.2. We begin with Proposition 4.1. Let P(n) denote the largest prime factor of n, and recall that a number n is said to be y-smooth if $P(n) \le y$. Recall also that the divisor function $d_{\lceil q \rceil}(n)$ is the Dirichlet series coefficient of $\zeta(s)^{\lceil q \rceil} = \prod_p (1 - 1/p^s)^{-\lceil q \rceil}$. By Minkowski's inequality, we have

$$\left\|\sum_{n \le x} f(n)\right\|_{2q} \le \sum_{0 \le k \le \mathcal{L}} \left\|\sum_{\substack{n \le x, \\ x^{e^{-(k+1)}} < P(n) \le x^{e^{-k}}} f(n)\right\|_{2q} + \left\|\sum_{\substack{n \le x, \\ P(n) \le x^{e^{-(\mathcal{L}+1)}}} f(n)\right\|_{2q}.$$

Furthermore, the first part of Probability Result 2.3, followed by Rankin's trick with exponent $1-1/\log^{0.9} x$ (bounding $\mathbf{1}_{n \le x}$ by $\left(\frac{x}{n}\right)^{1-1/\log^{0.9} x} = xe^{-\log^{0.1} x} \frac{1}{n^{1-1/\log^{0.9} x}}$), implies that

$$\left\|\sum_{\substack{n \le x, \\ P(n) \le x^{e^{-(\mathcal{L}+1)}}} f(n)\right\|_{2q} \le \left[\sum_{\substack{n \le x, \\ P(n) \le x^{e^{-(\mathcal{L}+1)}}} d_{\lceil q \rceil}(n)\right]^{1/2} \le \left[xe^{-\log^{0.1}x} \sum_{\substack{n \le x, \\ P(n) \le x^{e^{-(\mathcal{L}+1)}}} \frac{d_{\lceil q \rceil}(n)}{n^{1-1/\log^{0.9}x}}\right]^{1/2}$$

Here the sum over n is $\leq \prod_{p \leq x^{e^{-(\mathcal{L}+1)}}} (1 - 1/p^{1-1/\log^{0.9} x})^{-\lceil q \rceil}$, and recalling that $\mathcal{L} := \lfloor (\log \log x)/10 \rfloor$ this is $\leq \prod_{p \leq e^{\log^{0.9} x}} (1 - 1/p^{1-1/\log^{0.9} x})^{-\lceil q \rceil}$, which is $= e^{O(q)} \prod_{p \leq e^{\log^{0.9} x}} (1 - 1/p)^{-\lceil q \rceil} = \log^{O(q)} x$ by standard Chebychev and Mertens estimates for sums over primes. Since we assume in Proposition 4.1 that $q \leq \log^{0.05} x$, this whole contribution is $\ll \sqrt{x}e^{-c\log^{0.1} x}$, which is more than acceptable.

Next, if we let $\mathbb{E}^{(k)}$ denote expectation conditional on $(f(p))_{p \le x^{e^{-(k+1)}}}$, then the first part of Probability Result 2.3 applied, after conditioning on $(f(p))_{p \le x^{e^{-(k+1)}}}$, with

$$a_m = \mathbf{1}_{p \mid m \Rightarrow x^{e^{-(k+1)}}$$

implies

$$\begin{split} \sum_{0 \le k \le \mathcal{L}} \left\| \sum_{\substack{n \le x, \\ x^{e^{-(k+1)}} < P(n) \le x^{e^{-k}}}} f(n) \right\|_{2q} &= \sum_{0 \le k \le \mathcal{L}} \left\| \sum_{\substack{1 < m \le x, \\ p \mid m \Rightarrow x^{e^{-(k+1)}} < p \le x^{e^{-k}}}} f(m) \sum_{\substack{n \le x/m, \\ n \text{ is } x^{e^{-(k+1)}} - \text{smooth}}} f(n) \right\|_{2q} \\ &= \sum_{0 \le k \le \mathcal{L}} \left(\mathbb{E}\mathbb{E}^{(k)} \right\| \sum_{\substack{1 < m \le x, \\ p \mid m \Rightarrow x^{e^{-(k+1)}} < p \le x^{e^{-k}}}} f(m) \sum_{\substack{n \le x/m, \\ n \text{ is } x^{e^{-(k+1)}} - \text{smooth}}} f(n) \right\|_{2q}^{2q} \Big)^{1/2q} \\ &\le \sum_{0 \le k \le \mathcal{L}} \left(\mathbb{E}\left(\sum_{\substack{1 < m \le x, \\ p \mid m \Rightarrow x^{e^{-(k+1)}} < p \le x^{e^{-k}}}} d_{\lceil q \rceil}(m) \right\| \sum_{\substack{n \le x/m, \\ n \text{ is } x^{e^{-(k+1)}} - \text{smooth}}} f(n) \Big|^{2} \Big)^{q} \Big)^{1/2q}. \end{split}$$

To proceed further, we want to replace $\left|\sum_{n \le x/m, n \text{ is } x^{e^{-(k+1)}}\text{-smooth } f(n)\right|^2$ in the above by a smoothed version. Set $X = e^{\sqrt{\log x}}$, say, and note that (uniformly for any $1 \le q \le \log^{0.05} x$) the above is

$$\sum_{0 \le k \le \mathcal{L}} \left\| \sum_{\substack{1 < m \le x, \\ p \mid m \Rightarrow x^{e^{-(k+1)}} < p \le x^{e^{-k}}}} d_{\lceil q \rceil}(m) \right\| \sum_{\substack{n \le x/m, \\ n \text{ is } x^{e^{-(k+1)}} - \text{smooth}}} f(n) \Big|^2 \Big\|_q^{1/2} \\ \ll \sum_{0 \le k \le \mathcal{L}} \left\| \sum_{\substack{1 < m \le x, \\ p \mid m \Rightarrow x^{e^{-(k+1)}} < p \le x^{e^{-k}}}} d_{\lceil q \rceil}(m) \frac{X}{m} \int_m^{m(1+1/X)} \Big\| \sum_{\substack{n \le x/t, \\ x^{e^{-(k+1)}} - \text{smooth}}} f(n) \Big|^2 dt \Big\|_q^{1/2} \\ + \sum_{0 \le k \le \mathcal{L}} \left\| \sum_{\substack{1 < m \le x, \\ p \mid m \Rightarrow x^{e^{-(k+1)}} < p \le x^{e^{-k}}}} d_{\lceil q \rceil}(m) \frac{X}{m} \int_m^{m(1+1/X)} \Big\| \sum_{\substack{x \le x/t, \\ x^{e^{-(k+1)}} - \text{smooth}}} f(n) \Big|^2 dt \Big\|_q^{1/2}.$$
(4-1)

We next want to show that the second term in (4-1) may be discarded as an error term. Using Minkowski's inequality again, followed by Hölder's inequality with exponent q applied to the normalized integral $\frac{X}{m} \int_{m}^{m(1+1/X)} dt$, this second term is

$$\leq \sum_{0 \leq k \leq \mathcal{L}} \left[\sum_{\substack{1 < m \leq x, \\ p \mid m \Rightarrow x^{e^{-(k+1)}} < p \leq x^{e^{-k}}}} d_{\lceil q \rceil}(m) \left\| \frac{X}{m} \int_{m}^{m(1+1/X)} \left| \sum_{\substack{x/t < n \leq x/m, \\ x^{e^{-(k+1)}} - \text{smooth}}} f(n) \right|^{2} dt \right\|_{q} \right]^{1/2}$$

$$\leq \sum_{0 \leq k \leq \mathcal{L}} \left[\sum_{\substack{1 < m \leq x, \\ p \mid m \Rightarrow x^{e^{-(k+1)}} < p \leq x^{e^{-k}}}} d_{\lceil q \rceil}(m) \left(\frac{X}{m} \int_{m}^{m(1+1/X)} \mathbb{E} \left| \sum_{\substack{x/t < n \leq x/m, \\ x^{e^{-(k+1)}} - \text{smooth}}} f(n) \right|^{2q} dt \right)^{1/q} \right]^{1/2}$$

The length of the sum over *n* here is $x(t-m)/(mt) \le x/(mX)$, so when $x/X \le m \le x$ there will be at most one term in the sum, and we simply have

$$\mathbb{E} \left| \sum_{\substack{x/t < n \le x/m, \\ x^{e^{-(k+1)}} \text{-smooth}}} f(n) \right|^{2q} \le 1.$$

When $x^{e^{-(k+1)}} < m < x/X$, we take a fairly crude approach and use the Cauchy–Schwarz inequality, obtaining that

$$\mathbb{E} \left| \sum_{\substack{x/t < n \le x/m, \\ x^{e^{-(k+1)}} \text{-smooth}}} f(n) \right|^{2q} \le \left[\mathbb{E} \left| \sum_{\substack{x/t < n \le x/m, \\ x^{e^{-(k+1)}} \text{-smooth}}} f(n) \right|^{2} \mathbb{E} \left| \sum_{\substack{x/t < n \le x/m, \\ x^{e^{-(k+1)}} \text{-smooth}}} f(n) \right|^{2(2q-1)} \right]^{1/2} \\ \ll \left[\frac{x}{mX} \mathbb{E} \left| \sum_{\substack{x/t < n \le x/m, \\ x^{e^{-(k+1)}} \text{-smooth}}} f(n) \right|^{2(2q-1)} \right]^{1/2} \\ \ll \left[\frac{x}{mX} \left(\frac{x}{m} \right)^{2q-1} \log^{O(q^2)} x \right]^{1/2} \\ = \left(\frac{x}{m} \right)^{q} \frac{\log^{O(q^2)} x}{X^{1/2}}.$$

Here the crude upper bound $(x/m)^{2q-1}\log^{O(q^2)} x$ for the 2(2q-1)-th moment may be proved as in Section 3.

Putting things together, we find that the second term in (4-1) is

$$\ll \sum_{0 \le k \le \mathcal{L}} \left[\frac{x \log^{O(q)} x}{X^{1/2q}} \sum_{\substack{1 < m < x/X, \\ p \mid m \Rightarrow x^{e^{-(k+1)}} < p \le x^{e^{-k}}}} \frac{d_{\lceil q \rceil}(m)}{m} + \sum_{\substack{x/X \le m \le x, \\ p \mid m \Rightarrow x^{e^{-(k+1)}} < p \le x^{e^{-k}}}} d_{\lceil q \rceil}(m) \right]^{1/2}.$$

To bound the first of these sums we use the simple estimate

$$\sum_{\substack{1 < m < x/X, \\ p \mid m \Rightarrow x^{e^{-(k+1)}} < p \le x^{e^{-k}}}} \frac{d_{\lceil q \rceil}(m)}{m} \le \prod_{x^{e^{-(k+1)}} < p \le x^{e^{-k}}} \left(1 - \frac{1}{p}\right)^{-\lceil q \rceil},$$

which is $= e^{O(q)}$ by the Mertens estimate for products over primes. To bound the second sum, by submultiplicativity of $d_{\lceil q \rceil}(\cdot)$ we always have $d_{\lceil q \rceil}(m) \le \lceil q \rceil^{\Omega(m)}$, where $\Omega(m)$ is the total number of prime factors of *m*. And we note (to obtain good dependence on *k*) that if $m \ge x/X$ only has prime factors from the interval $(x^{e^{-(k+1)}}, x^{e^{-k}}]$, then we must have $\Omega(m) \ge e^k/2$, say. So we get that

$$\sum_{\substack{x/X \le m \le x, \\ p \mid m \Rightarrow x^{e^{-(k+1)}}$$

is at most

$$5^{-e^{k}/2} \sum_{\substack{x/X \le m \le x, \\ p \mid m \Rightarrow x^{e^{-(k+1)}}$$

where the first inequality uses Number Theory Result 2.7. Recalling that we have

$$q \le \log^{0.05} x$$
, $\mathcal{L} = \lfloor (\log \log x) / 10 \rfloor$ and $X = e^{\sqrt{\log x}}$.

the second term in (4-1) is

$$\leq \sum_{0 \leq k \leq \mathcal{L}} \left[\frac{x \log^{O(q)} x}{X^{1/2q}} + e^{O(q)} 2^{-e^k} \frac{x}{\log x} \right]^{1/2} \leq e^{O(q)} \sqrt{\frac{x}{\log x}},$$

which is an acceptable contribution for Proposition 4.1.

Turning to the remaining first sum in (4-1), this is equal to

$$\sum_{0 \le k \le \mathcal{L}} \left\| \int_{x^{e^{-(k+1)}}}^{x} \left| \sum_{\substack{n \le x/t, \\ x^{e^{-(k+1)}} - \text{smooth}}} f(n) \right|^2 \sum_{\substack{t/(1+1/X) \le m \le t, \\ p \mid m \Rightarrow x^{e^{-(k+1)}}$$

Now we set $u = u(k, t) := e^k (\log t) / \log x$, and notice that (by submultiplicativity) $d_{\lceil q \rceil}(m) \le \lceil q \rceil^{\Omega(m)}$, and if $m \ge t/(1 + 1/X)$ only has prime factors from the interval $(x^{e^{-(k+1)}}, x^{e^{-k}}]$ then we must have $\Omega(m) \ge u - 1$. So using Number Theory Result 2.7 (whose conditions are satisfied since $X = e^{\sqrt{\log x}}$ isn't too large, and $k \le (\log \log x)/10$) we get

$$\begin{split} \sum_{\substack{t/(1+1/X) \le m \le t, \\ p \mid m \Rightarrow x^{e^{-(k+1)}}$$

provided x is sufficiently large. Consequently, the first sum in (4-1) is

$$\leq e^{O(q)} \sum_{0 \leq k \leq \mathcal{L}} \left\| \int_{x^{e^{-(k+1)}}}^{x} \left\| \sum_{\substack{n \leq x/t, \\ x^{e^{-(k+1)}} - \text{smooth}}} f(n) \right\|_{q}^{2} \frac{dt}{\log t} \right\|_{q}^{1/2}$$
$$= e^{O(q)} \sum_{0 \leq k \leq \mathcal{L}} \sqrt{x} \left\| \int_{1}^{x^{1-e^{-(k+1)}}} \left\| \sum_{\substack{n \leq z, \\ x^{e^{-(k+1)}} - \text{smooth}}} f(n) \right\|_{2}^{2} \frac{dz}{z^{2} \log(x/z)} \right\|_{q}^{1/2},$$

where the second line follows from making the substitution z = x/t.

To obtain a satisfactory dependence on k in our final estimations, we now note that if $z \le \sqrt{x}$ we have $\log(x/z) \gg \log x$, whereas if $\sqrt{x} < z \le x^{1-e^{-(k+1)}}$ we have $\log(x/z) \gg e^{-k} \log x$. Thus in any case we have $\log(x/z) \gg z^{-2k/\log x} \log x$. As discussed earlier, we also want to introduce a Rankin style shift, which we will achieve by adding a factor $(x/z)^{2q/\log x} = e^{O(q)} z^{-2q/\log x}$ into the integral. Inserting these estimates, we find the first sum in (4-1) is

$$\leq \frac{\sqrt{x}e^{O(q)}}{\sqrt{\log x}} \sum_{0 \leq k \leq \mathcal{L}} \left\| \int_{1}^{x^{1-e^{-(k+1)}}} \left\| \sum_{\substack{n \leq z, \\ x^{e^{-(k+1)}} \text{-smooth}}} f(n) \right\|^{2} \frac{dz}{z^{2+2q/\log x - 2k/\log x}} \right\|_{q}^{1/2}.$$

Finally, using Harmonic Analysis Result 2.6 and then Minkowski's inequality, all of the above is

$$\leq \sqrt{\frac{x}{\log x}} e^{O(q)} \sum_{0 \leq k \leq \mathcal{L}} \left\| \int_{-\infty}^{\infty} \frac{|F_k(1/2 + q/\log x - k/\log x + it)|^2}{|1/2 + q/\log x - k/\log x + it|^2} dt \right\|_q^{1/2} \\ \leq \sqrt{\frac{x}{\log x}} e^{O(q)} \sum_{0 \leq k \leq \mathcal{L}} \left[\sum_{n \in \mathbb{Z}} \frac{1}{n^2 + 1} \left\| \int_{n-1/2}^{n+1/2} \left| F_k\left(\frac{1}{2} + \frac{q - k}{\log x} + it\right) \right|^2 dt \right\|_q^2 \right]^{1/2}$$

where F_k denotes the partial Euler product of f(n) over $x^{e^{-(k+1)}}$ -smooth numbers. In the Steinhaus case, since the law of the random function f(n) is the same as the law of $f(n)n^{it}$ for any fixed $t \in \mathbb{R}$ we have

$$\left\|\int_{n-1/2}^{n+1/2} \left|F_k\left(\frac{1}{2} + \frac{q-k}{\log x} + it\right)\right|^2 dt\right\|_q = \left\|\int_{-1/2}^{1/2} \left|F_k\left(\frac{1}{2} + \frac{q-k}{\log x} + it\right)\right|^2 dt\right\|_q \quad \forall n.$$

Proposition 4.1 now follows on putting everything together.

The proof of Proposition 4.2, covering the Rademacher case, is very similar to the Steinhaus case. We use the Rademacher part of Probability Result 2.3, producing various terms $d_{2\lceil q\rceil-1}(n)$ in place of $d_{\lceil q\rceil}(n)$, but this doesn't alter the analysis. The only nontrivial change comes at the very end of the proof, where (since it is no longer the case that the law of the random function f(n) is the same as the law of $f(n)n^{it}$) we apply the bound

$$\begin{split} \left[\sum_{n\in\mathbb{Z}}\frac{1}{n^2+1}\left\|\int_{n-1/2}^{n+1/2}\left|F_k\left(\frac{1}{2}+\frac{q-k}{\log x}+it\right)\right|^2dt\right\|_q\right]^{1/2} \\ \ll \left[\max_{N\in\mathbb{Z}}\frac{1}{(|N|+1)^{1/4}}\left\|\int_{N-1/2}^{N+1/2}\left|F_k\left(\frac{1}{2}+\frac{q-k}{\log x}+it\right)\right|^2dt\right\|_q\right]^{1/2}. \quad \Box \end{split}$$

Proof of Propositions 4.3 and 4.4. We proceed somewhat similarly to Section 2.5 of [Harper 2017] or Section 2.2 of [Harper et al. 2015].

Again we let P(n) denote the largest prime factor of n, and we introduce an auxiliary Rademacher random variable ϵ that is independent of everything else. Then we find that

$$\begin{split} \left\| \sum_{\substack{n \le x, \\ P(n) > x^{3/4}}} f(n) \right\|_{2q} &= \frac{1}{2} \left\| \sum_{\substack{n \le x, \\ P(n) > x^{3/4}}} f(n) + \sum_{\substack{n \le x, \\ P(n) > x^{3/4}}} f(n) + \sum_{\substack{n \le x, \\ P(n) > x^{3/4}}} f(n) + \sum_{\substack{n \le x, \\ P(n) > x^{3/4}}} f(n) \right\|_{2q} + \left\| \sum_{\substack{n \le x, \\ P(n) > x^{3/4}}} f(n) - \sum_{\substack{n \le x, \\ P(n) > x^{3/4}}} f(n) \right\|_{2q} \right) \\ &\leq \left\| \epsilon \sum_{\substack{n \le x, \\ P(n) > x^{3/4}}} f(n) + \sum_{\substack{n \le x, \\ P(n) \le x^{3/4}}} f(n) \right\|_{2q} \\ &\leq \left\| \epsilon \sum_{\substack{n \le x, \\ P(n) > x^{3/4}}} f(n) + \sum_{\substack{n \le x, \\ P(n) \le x^{3/4}}} f(n) \right\|_{2q} \\ &= \left\| \sum_{n \le x} f(n) \right\|_{2q}. \end{split}$$

Here the first inequality is Minkowski's inequality; the second is Hölder's inequality (with exponent 2q) applied only to the averaging over ϵ ; and the final equality follows since the law of

$$\epsilon \sum_{\substack{n \le x, \\ P(n) > x^{3/4}}} f(n) = \epsilon \sum_{x^{3/4}$$

conditional on the values $(f(p))_{p \le x^{3/4}}$ is the same as the law of $\sum_{n \le x, P(n) > x^{3/4}} f(n)$.

Now in the decomposition

$$\sum_{n \le x, P(n) > x^{3/4}} f(n) = \sum_{x^{3/4}$$

the inner sums are determined by the values $(f(p))_{p \le x^{3/4}}$ (and in fact by the values $(f(p))_{p \le x^{1/4}}$), which are independent of the outer random variables $(f(p))_{x^{3/4} . So conditioning on the values <math>(f(p))_{p \le x^{3/4}}$

determining the inner sums and applying Probability Result 2.4 with $a_p = \sum_{m < x/p} f(m)$, it follows that

$$\left\|\sum_{\substack{n \le x, \\ P(n) > x^{3/4}}} f(n)\right\|_{2q} \ge \left\|\sum_{x^{3/4}$$

Next we want to replace the sum over p by an integral average. We can rewrite

$$\sum_{x^{3/4}$$

and noting that $x/r - x/(r+1) = x/(r(r+1)) \gg (x/r)^{2/3}$ on our range of *r*, a Hoheisel-type prime number theorem in short intervals (see, e.g., Theorem 12.8 of [Ivić 2003]) implies that

$$\sum_{x^{3/4}$$

Making a substitution z = x/t, we see this integral is the same as $x \int_{1}^{x^{1/4}} \left| \sum_{m \le z} f(m) \right|^2 \frac{dz}{z^2}$. Checking back, this completes the proof of the first part of Proposition 4.3.

To deduce the second part of Proposition 4.3, we note that for any large V and any $q \ge 1$ we have

$$\begin{split} \left\| \int_{1}^{x^{1/4}} \left| \sum_{m \le z} f(m) \right|^{2} \frac{dz}{z^{2}} \right\|_{q} \\ & \ge \left\| \int_{1}^{x^{1/4}} \left| \sum_{\substack{m \le z, \\ x - \text{smooth}}} f(m) \right|^{2} \frac{dz}{z^{2+8Vq/\log x}} \right\|_{q} \\ & \ge \left\| \int_{1}^{\infty} \left| \sum_{\substack{m \le z, \\ x - \text{smooth}}} f(m) \right|^{2} \frac{dz}{z^{2+8Vq/\log x}} \right\|_{q} - \left\| \int_{x^{1/4}}^{\infty} \left| \sum_{\substack{m \le z, \\ x - \text{smooth}}} f(m) \right|^{2} \frac{dz}{z^{2+8Vq/\log x}} \right\|_{q} \\ & \ge \left\| \int_{1}^{\infty} \left| \sum_{\substack{m \le z, \\ x - \text{smooth}}} f(m) \right|^{2} \frac{dz}{z^{2+8Vq/\log x}} \right\|_{q} - \frac{1}{e^{Vq}} \left\| \int_{1}^{\infty} \left| \sum_{\substack{m \le z, \\ x - \text{smooth}}} f(m) \right|^{2} \frac{dz}{z^{2+4Vq/\log x}} \right\|_{q}. \end{split}$$

By Harmonic Analysis Result 2.6, provided that $V \leq (\log x)/q$ (so that $Vq/\log x$ is uniformly bounded) the first term here is $\gg \left\| \int_{-1/2}^{1/2} \left| F\left(\frac{1}{2} + \frac{4Vq}{\log x} + it\right) \right|^2 dt \right\|_q$ and the subtracted second term is $\ll e^{-Vq} \left\| \int_{-\infty}^{\infty} \left| F\left(\frac{1}{2} + \frac{2Vq}{\log x} + it\right) \right|^2 / \left| \frac{1}{2} + \frac{2Vq}{\log x} + it \right|^2 dt \right\|_q$, which in the Steinhaus case is $\ll e^{-Vq} \left\| \int_{-1/2}^{1/2} \left| F\left(\frac{1}{2} + \frac{2Vq}{\log x} + it\right) \right|^2 dt \right\|_q$ by "translation invariance in law". Putting everything together, this finishes the proof of Proposition 4.3.

The arguments in the Rademacher case are exactly the same until the final line, where we don't have "translation invariance" so we must upper bound $\|\int_{-\infty}^{\infty} |F(\frac{1}{2} + \frac{2Vq}{\log x} + it)|^2 / |\frac{1}{2} + \frac{2Vq}{\log x} + it|^2 dt \|_q$ by $\max_{N \in \mathbb{Z}} \frac{1}{(|N|+1)^{1/4}} \|\int_{N-1/2}^{N+1/2} |F(\frac{1}{2} + \frac{2Vq}{\log x} + it)|^2 dt \|_q$, say.

5. Proofs of the upper bounds in Theorems 1.1 and 1.2

In view of Proposition 4.1, the key to obtaining the upper bound in Theorem 1.1 will lie in proving the following. Recall here that $F_k(s)$ denotes the partial Euler product of f(n) over $x^{e^{-(k+1)}}$ -smooth numbers, and in the special case where k = -1 we usually write F(s) (rather than $F_{-1}(s)$) for the partial Euler product over *x*-smooth numbers.

Key Proposition 5.1. Let f(n) be a Steinhaus random multiplicative function. For all large x, and uniformly for $1 \le q \le \log^{100} x$ (say) and $-1 \le k \le \mathcal{L} = \lfloor (\log \log x)/10 \rfloor$ and $-e^k/\log x \le \sigma \le 1/(100 \log(2q))$ (say), we have

$$\mathbb{E}\left(\int_{-1/2}^{1/2} \left| F_k\left(\frac{1}{2} + \sigma + it\right) \right|^2 dt \right)^q \ll \frac{e^{O(q^2)}}{\log^{q-1} x} \left(\frac{\log x}{\log 2q}\right)^{q^2} \min\left\{\frac{1}{e^{k+1}}, \frac{1}{|\sigma|\log x}\right\}^{q^2-q+1}.$$

Key Proposition 5.1 is actually much more general, in terms of the allowed range of q and σ , than we immediately need (and the proof would let us extend the range of q quite a lot further if we wished, all it really requires is something like $e^k/\log x \le 1/(100 \log(2q))$). The increased generality will be useful in Section 6, where Key Proposition 5.1 will play an auxiliary role, and also in clarifying the essential features of the proof.

Proof of the upper bound in Theorem 1.1, assuming Key Proposition 5.1. In view of the discussion in Section 3, it will suffice to prove the Theorem 1.1 upper bound for $1 \le q \le \log \log x$. And to do that, in view of Proposition 4.1 it will suffice to show that

$$\sum_{0 \le k \le \mathcal{L}} \left\| \int_{-1/2}^{1/2} \left| F_k \left(\frac{1}{2} + \frac{q-k}{\log x} + it \right) \right|^2 dt \right\|_q^{1/2} \le e^{-(q/2)\log q - (q/2)\log\log(2q) + O(q)} \log^{q/2 - 1/2 + 1/2q} x.$$

Applying Key Proposition 5.1 with $\sigma = (q - k)/\log x$ (which is indeed $\leq 1/(100\log(2q))$ on our range of q), we find the left-hand side is

$$\leq \sum_{0 \leq k \leq \mathcal{L}} \left(\frac{e^{O(q^2)}}{\log^{q-1} x} \left(\frac{\log x}{\log 2q} \right)^{q^2} \min\left\{ \frac{1}{e^{k+1}}, \frac{1}{|q-k|} \right\}^{q^2-q+1} \right)^{1/2q}$$
$$= \sum_{0 \leq k \leq \mathcal{L}} \left(e^{O(q^2)} \left(\frac{\log x}{\log 2q} \min\left\{ \frac{1}{e^{k+1}}, \frac{1}{q} \right\} \right)^{q^2-q+1} \right)^{1/2q}.$$

It is easy to see that this satisfies our desired bound.

For Theorem 1.2, we need a Rademacher analogue of the above.

Key Proposition 5.2. Let f(n) be a Rademacher random multiplicative function. For all large x, and uniformly for $1 \le q \le \log^{100} x$ (say) and $-1 \le k \le \mathcal{L} = \lfloor (\log \log x)/10 \rfloor$ and $-e^k/\log x \le \sigma \le 1/(100 \log(2q))$ (say), we have

$$\mathbb{E}\left(\int_{-1/2}^{1/2} |F_k(\frac{1}{2} + \sigma + it)|^2 dt\right)^q \\ \ll \frac{e^{O(q^2)}}{\log^q x} \left(1 + \min\left\{\log\log x, \frac{1}{|q - q_0|}\right\}\right) \left(\frac{\log x}{\log 2q}\right)^{\max\{2q^2 - q, q^2 + 1\}} \min\left\{\frac{1}{e^{k+1}}, \frac{1}{|\sigma|\log x}\right\}^{\max\{2q^2 - 2q, q^2 - q + 1\}},$$

where $q_0 = (1 + \sqrt{5})/2$.

Furthermore, for any $|N| \ge 1$ *we have*

$$\mathbb{E}\left(\int_{N-\frac{1}{2}}^{N+\frac{1}{2}} \left|F_{k}\left(\frac{1}{2}+\sigma+it\right)\right|^{2} dt\right)^{q} \\ \ll \min\left\{|N|^{1/100}, \frac{\log x}{e^{k+1}\log 2q}, \frac{1}{|\sigma|\log 2q}\right\}^{q(q+1)} \frac{e^{O(q^{2})}}{\log^{q-1}x} \left(\frac{\log x}{\log 2q}\right)^{q^{2}} \min\left\{\frac{1}{e^{k+1}}, \frac{1}{|\sigma|\log x}\right\}^{q^{2}-q+1}.$$

Proof of the upper bound in Theorem 1.2, assuming Key Proposition 5.2. Similarly as in the Steinhaus case, in view of Proposition 4.2 and the discussion in Section 3 it will suffice to show that for all $1 \le q \le \log \log x$, we have

$$\sum_{0 \le k \le \mathcal{L}} \max_{N \in \mathbb{Z}} \frac{1}{(|N|+1)^{1/8}} \left\| \int_{N-1/2}^{N+1/2} \left| F_k \left(\frac{1}{2} + \frac{q-k}{\log x} + it \right) \right|^2 dt \right\|_q^{1/2} \\ \le e^{-q \log q - q \log \log(2q) + O(q)} \left(1 + \min \left\{ \log \log x, \frac{1}{|q-q_0|} \right\} \right)^{1/2q} (\log x)^{\max\{q-1, q/2 - 1/2 + 1/2q\}}.$$

We apply Key Proposition 5.2 with $\sigma = (q - k)/\log x$ (which is indeed $\leq 1/(100 \log(2q))$) on our range of q). When $q \leq 15$, say, we have $\min\{|N|^{1/100}, \log x/(e^{k+1}\log 2q), 1/(|\sigma|\log 2q)\}^{q(q+1)} \leq |N|^{q/5} \leq |N|^{2q/8}$, so (on taking 2q-th roots in Key Proposition 5.2 and then multiplying by the prefactor $1/(|N|+1)^{1/8}$) we see the contribution from $|N| \geq 1$ to $\max_{N \in \mathbb{Z}}$ will never exceed the contribution from the N = 0 term. So overall, when $1 \leq q \leq 15$ the left-hand side will be

$$\leq \sum_{0 \leq k \leq \mathcal{L}} \left(\frac{e^{O(q^2)}}{\log^q x} \min\left\{ \log \log x, \frac{1}{|q-q_0|} \right\} \left(\frac{\log x}{\log 2q} \right)^{\max\{2q^2 - q, q^2 + 1\}} \left(\frac{1}{e^{k+1}} \right)^{\max\{2q^2 - 2q, q^2 - q + 1\}} \right)^{1/2q} \\ \ll \sum_{0 \leq k \leq \mathcal{L}} \left(\min\left\{ \log \log x, \frac{1}{|q-q_0|} \right\} \left(\frac{\log x}{e^{k+1}} \right)^{\max\{2q(q-1), q^2 - q + 1\}} \right)^{1/2q}.$$

This certainly gives our desired bound for $1 \le q \le 15$.

When $15 \le q \le \log \log x$, we note first that $\max\{2q^2 - 2q, q^2 - q + 1\} = 2q^2 - 2q$. (In fact this is true as soon as $q \ge q_0$.) So using the bound

$$\min\left\{|N|^{1/100}, \frac{\log x}{e^{k+1}\log 2q}, \frac{1}{|\sigma|\log 2q}\right\}^{q(q+1)} \le |N|^{(2q+1)/100} \min\left\{\frac{\log x}{e^{k+1}\log 2q}, \frac{1}{|\sigma|\log 2q}\right\}^{q^2-q-1},$$

we again find that the contribution from $|N| \ge 1$ to $\max_{N \in \mathbb{Z}}$ will never exceed the contribution from the N = 0 term. Overall, in this case we get a bound

$$\ll \sum_{0 \le k \le \mathcal{L}} \left(e^{O(q^2)} \left(\frac{\log x}{\log 2q} \min\left\{ \frac{1}{e^{k+1}}, \frac{1}{q} \right\} \right)^{2q^2 - 2q} \right)^{1/2q} \ll \left(e^{O(q^2)} \left(\frac{\log x}{q \log 2q} \right)^{2q^2 - 2q} \right)^{1/2q},$$
ared.

as desired.

We shall prove Key Propositions 5.1 and 5.2 in several steps over the course of this section. For convenience in the writing we set $\mathcal{X} := \min\{\log x/e^{k+1}, 1/|\sigma|\}$, and note that under our hypotheses this is always $\geq 100 \log(2q)$. The point of this definition is that the contribution from primes $p > e^{\mathcal{X}}$ in our Euler products will ultimately contribute only to the $e^{O(q^2)}$ term.

Preliminary maneuvres. We begin with a few manipulations to discretize and set up the problem, in both the Steinhaus and Rademacher cases. For any $q \ge 1$, we have

$$\left\| \int_{-1/2}^{1/2} \left| F_k \left(\frac{1}{2} + \sigma + it \right) \right|^2 dt \right\|_q \leq \left\| \int_{-1/(2\mathcal{X})}^{1/(2\mathcal{X})} \sum_{|n| \leq \mathcal{X}/2+1} \left| F_k \left(\frac{1}{2} + \sigma + i\left(\frac{n}{\mathcal{X}} + t \right) \right) \right|^2 dt \right\|_q$$
$$= \frac{1}{\mathcal{X}} \left\| \int_{-1/2\mathcal{X}}^{1/2\mathcal{X}} \mathcal{X} \sum_{|n| \leq \mathcal{X}/2+1} \left| F_k \left(\frac{1}{2} + \sigma + i\left(\frac{n}{\mathcal{X}} + t \right) \right) \right|^2 dt \right\|_q$$

Applying Hölder's inequality with exponent q to the normalized integral $\int_{-1/(2\mathcal{X})}^{1/(2\mathcal{X})} \mathcal{X} dt$, we see the right-hand side is

$$\leq \frac{1}{\mathcal{X}} \bigg(\int_{-1/(2\mathcal{X})}^{1/(2\mathcal{X})} \mathcal{X} \mathbb{E} \bigg(\sum_{|n| \leq \mathcal{X}/2+1} \big| F_k \big(\frac{1}{2} + \sigma + i \big(\frac{n}{\mathcal{X}} + t \big) \big) \big|^2 \bigg)^q dt \bigg)^{1/q}.$$

In the Steinhaus case, where $|F_k(\frac{1}{2} + \sigma + i(\frac{n}{\lambda} + t))|^2$ has the same distribution for any given shift *t*, we can simplify the above to give the bound

$$\left\|\int_{-1/2}^{1/2} \left|F_k(\frac{1}{2} + \sigma + it)\right|^2 dt\right\|_q \le \frac{1}{\mathcal{X}} \left\|\sum_{|n| \le \mathcal{X}/2 + 1} \left|F_k(\frac{1}{2} + \sigma + \frac{in}{\mathcal{X}})\right|^2\right\|_q,$$

so we have indeed passed to studying a discrete sum rather than an integral. Finally, we rewrite the right-hand side as

$$\frac{1}{\mathcal{X}} \left(\mathbb{E} \sum_{|n| \leq \mathcal{X}/2+1} \left| F_k \left(\frac{1}{2} + \sigma + \frac{in}{\mathcal{X}} \right) \right|^2 \left(\sum_{|m| \leq \mathcal{X}/2+1} \left| F_k \left(\frac{1}{2} + \sigma + \frac{im}{\mathcal{X}} \right) \right|^2 \right)^{q-1} \right)^{1/q} \\
\ll \frac{1}{\mathcal{X}} \left(\mathcal{X} \mathbb{E} \left| F_k \left(\frac{1}{2} + \sigma \right) \right|^2 \left(\sum_{|m| \leq \mathcal{X}} \left| F_k \left(\frac{1}{2} + \sigma + \frac{im}{\mathcal{X}} \right) \right|^2 \right)^{q-1} \right)^{1/q}, \quad (5-1)$$

where the inequality again uses the distributional "translation invariance" (shifting *n* to zero in the outer sum, and replacing *m* by m - n in the second sum).

In the case of Rademacher f(n), if we mimic the above calculations we obtain that

$$\left\| \int_{N-1/2}^{N+1/2} \left| F_k \left(\frac{1}{2} + \sigma + it \right) \right|^2 dt \right\|_q$$

$$\leq \frac{1}{\mathcal{X}} \left(\int_{-1/2\mathcal{X}}^{1/2\mathcal{X}} \mathcal{X} \mathbb{E} \sum_{|n| \leq \mathcal{X}/2+1} \left| F_k \left(\frac{1}{2} + \sigma + i \left(\frac{n}{\mathcal{X}} + N + t \right) \right) \right|^2 \left(\sum_{|m| \leq \mathcal{X}/2+1} \left| F_k \left(\frac{1}{2} + \sigma + i \left(\frac{m}{\mathcal{X}} + N + t \right) \right) \right|^2 \right)^{q-1} dt \right)^{1/q}.$$
(5-2)

Proof of Key Proposition 5.1, for $q \ge 2$. When $q \ge 2$, we are helped by the fact that we can use Hölder's inequality again to analyze $\left(\sum_{|m|\le\chi} \left|F_k\left(\frac{1}{2}+\sigma+\frac{im}{\chi}\right)\right|^2\right)^{q-1}$ in (5-1). If we let $\mu := \sum_{|m|\le\chi} 1/(|m|+1)^2$, so that $\mu \asymp 1$, then first we have

$$\left(\sum_{|m|\leq\mathcal{X}} \left|F_k\left(\frac{1}{2} + \sigma + \frac{im}{\mathcal{X}}\right)\right|^2\right)^{q-1} = \mu^{q-1} \left(\frac{1}{\mu} \sum_{|m|\leq\mathcal{X}} \frac{1}{(|m|+1)^2} (|m|+1)^2 \left|F_k\left(\frac{1}{2} + \sigma + \frac{im}{\mathcal{X}}\right)\right|^2\right)^{q-1}$$

Using Hölder's inequality with exponent q - 1, we deduce

$$\left(\sum_{|m|\leq\mathcal{X}} \left|F_k\left(\frac{1}{2} + \sigma + \frac{im}{\mathcal{X}}\right)\right|^2\right)^{q-1} \leq \mu^{q-1} \cdot \frac{1}{\mu} \sum_{|m|\leq\mathcal{X}} \frac{1}{(|m|+1)^2} ((|m|+1)^2 \left|F_k\left(\frac{1}{2} + \sigma + \frac{im}{\mathcal{X}}\right)\right|^2)^{q-1} \\ = e^{O(q)} \sum_{|m|\leq\mathcal{X}} \frac{1}{(|m|+1)^2} (|m|+1)^{2(q-1)} \left|F_k\left(\frac{1}{2} + \sigma + \frac{im}{\mathcal{X}}\right)\right|^{2(q-1)}.$$
(5-3)

We remark that the choice of weights $1/(|m|+1)^2$ that we introduced is fairly arbitrary. The key point is that we expect, in (5-1), that the only significant contribution should come from small *m* (for which $|F_k(\frac{1}{2} + \sigma + \frac{im}{\chi})|^2$ will be highly correlated with the outer term $|F_k(\frac{1}{2} + \sigma)|^2$), so we don't want to pick up a factor like χ (inefficiently reflecting the total length of the sum) in our application of Hölder's inequality.

In view of the above computation, to bound the right-hand side of (5-1) when $q \ge 2$ we need to bound terms of the form

$$(|m|+1)^{2(q-1)}\mathbb{E}\left|F_k\left(\frac{1}{2}+\sigma\right)\right|^2\left|F_k\left(\frac{1}{2}+\sigma+\frac{im}{\mathcal{X}}\right)\right|^{2(q-1)}.$$

Recall that $-e^k/\log x \le \sigma \le 1/100\log(2q)$ here. Inserting the definition of $F_k(s)$, and using a trivial bound $e^{O\left(\sum_{p\le 100q^2} q/p^{1/2+\sigma}\right)} = e^{O\left(\sum_{p\le 100q^2} q/\sqrt{p}\right)} = e^{O(q^2/\log q)}$ for the parts of the Euler products over primes $\le 100q^2$, this is

$$e^{O(q^2/\log q)}(|m|+1)^{2(q-1)}\mathbb{E}\prod_{100q^2$$

Now if $e^{k+1}/\log x \le \sigma \le 1/100 \log(2q)$, (and so $\mathcal{X} := \min\{\log x/e^{k+1}, 1/|\sigma|\} = 1/\sigma$), then the first part of Euler Product Result 2.1 implies that the expectation of the part of the Euler product over primes $e^{1/\sigma} is equal to <math>\exp\{O(\sum_{e^{1/\sigma} , which is all <math>e^{O(q^2)}$. Using

this fact, as well as the independence of f(p) for different primes p, we find the above is always equal to

$$e^{O(q^2)}(|m|+1)^{2(q-1)}\mathbb{E}\prod_{100q^2$$

Notice that our size assumptions on σ , k, q guarantee that e^{χ} is larger than $100q^2$. Finally, the second part of Euler Product Result 2.1 implies this is all equal to

$$e^{O(q^2)}(|m|+1)^{2(q-1)} \left(\frac{\mathcal{X}}{\log q}\right)^{1+(q-1)^2} \left(1+\frac{\mathcal{X}}{(|m|+1)\log q}\right)^{2(q-1)} = e^{O(q^2)} \left(\frac{\mathcal{X}}{\log q}\right)^{q^2}.$$

Putting this together with (5-3) and (5-1), we find

$$\begin{split} \left\| \int_{-1/2}^{1/2} \left| F_k \left(\frac{1}{2} + \sigma + it \right) \right|^2 dt \right\|_q &\ll \frac{1}{\mathcal{X}} \left(\mathcal{X} \cdot e^{O(q^2)} \sum_{|m| \le \mathcal{X}} \frac{1}{(|m| + 1)^2} \left(\frac{\mathcal{X}}{\log q} \right)^{q^2} \right)^{1/q} \\ &= \frac{1}{\mathcal{X}} \left(\mathcal{X} \cdot e^{O(q^2)} \left(\frac{\mathcal{X}}{\log q} \right)^{q^2} \right)^{1/q}. \end{split}$$

Raising everything to the power *q* and inserting the fact that $\mathcal{X} = \min\{\log x/e^{k+1}, 1/|\sigma|\}$, this gives the statement of Key Proposition 5.1.

Proof of Key Proposition 5.2, for $q \ge 2$ **.** We begin with the second part of Key Proposition 5.2, where $|N| \ge 1$. Then similarly as in the deduction of (5-3) in the Steinhaus case, for any $|n| \le \chi/2 + 1$ and any $|t| \le 1/(2\chi)$ we can use Hölder's inequality to show

$$\left(\sum_{|m| \le \frac{\chi}{2} + 1} \left| F_k \left(\frac{1}{2} + \sigma + i \left(\frac{m}{\chi} + N + t \right) \right) \right|^2 \right)^{q-1} \le e^{O(q)} \sum_{|m| \le \frac{\chi}{2} + 1} \frac{1}{(|m-n|+1)^2} (|m-n|+1)^{2(q-1)} \left| F_k \left(\frac{1}{2} + \sigma + i \left(\frac{m}{\chi} + N + t \right) \right) \right|^{2(q-1)}.$$

So to bound the right-hand side of (5-2), we need to bound terms of the form

$$\left(|m-n|+1\right)^{2(q-1)}\mathbb{E}\left|F_k\left(\frac{1}{2}+\sigma+i\left(\frac{n}{\mathcal{X}}+N+t\right)\right)\right|^2\left|F_k\left(\frac{1}{2}+\sigma+i\left(\frac{m}{\mathcal{X}}+N+t\right)\right)\right|^{2(q-1)}$$

As in the Steinhaus case, the contribution from primes $p \le 100q^2$ to this expectation is trivially $e^{O(q^2/\log q)}$. Using the first part of Euler Product Result 2.2, the contribution from primes $e^{\mathcal{X}} is <math>e^{O(q^2)}$, and overall (noting that the imaginary shifts $n/\mathcal{X} + N + t$, $m/\mathcal{X} + N + t$ are $\gg |N| \gg 1$ and also $\ll |N|$) the above expression is at most

$$e^{O(q^2)} \min\left\{\frac{\mathcal{X}}{\log q}, |N|^{\frac{1}{100}}\right\}^{(q-1)^2 + 3(q-1)} (|m-n|+1)^{2(q-1)} \left(\frac{\mathcal{X}}{\log q}\right)^{1 + (q-1)^2} \left(\frac{\mathcal{X}}{(|m-n|+1)\log q}\right)^{2(q-1)}.$$

Apart from the factor $\min\{\mathcal{X}/\log q, |N|^{1/100}\}^{(q-1)(q+2)}$, this is precisely analogous to the estimate we had in the Steinhaus case, so when $|N| \ge 1$ we indeed get the same bound as in the Steinhaus case apart

from a multiplier

$$\min\left\{\frac{\mathcal{X}}{\log q}, |N|^{1/100}\right\}^{(q-1)(q+2)} \leq \min\left\{\frac{\mathcal{X}}{\log q}, |N|^{1/100}\right\}^{q(q+1)} = \min\left\{\frac{\log x}{e^{k+1}\log q}, \frac{1}{|\sigma|\log q}, |N|^{1/100}\right\}^{q(q+1)}.$$

It remains to address the first part of Key Proposition 5.2, where N = 0. In this case, when $q \ge 2$ we expect the main contribution to (5-2) to come from terms with $n, m \approx 0$, so rather than splitting up the sum over *m* according to the size of |m - n| we shall just split it up according to the size of *m*. Proceeding in this way, using Hölder's inequality as in the Steinhaus case we find for any $|t| \le 1/(2\mathcal{X})$ that

$$\sum_{|n| \le \frac{\chi}{2} + 1} \left| F_k \left(\frac{1}{2} + \sigma + i \left(\frac{n}{\chi} + t \right) \right) \right|^2 \left(\sum_{|m| \le \chi/2 + 1} \left| F_k \left(\frac{1}{2} + \sigma + i \left(\frac{m}{\chi} + t \right) \right) \right|^2 \right)^{q-1} \\ \le e^{O(q)} \sum_{|n| \le \chi/2 + 1} \left| F_k \left(\frac{1}{2} + \sigma + i \left(\frac{n}{\chi} + t \right) \right) \right|^2 \sum_{|m| \le \chi/2 + 1} \frac{1}{(|m| + 1)^2} (|m| + 1)^{2(q-1)} \left| F_k \left(\frac{1}{2} + \sigma + i \left(\frac{m}{\chi} + t \right) \right) \right|^{2(q-1)}.$$
(5-4)

So to bound the right-hand side of (5-2), we again need to bound terms of the form

$$(|m|+1)^{2(q-1)}\mathbb{E}\left|F_k\left(\frac{1}{2}+\sigma+i\left(\frac{n}{\chi}+t\right)\right)\right|^2\left|F_k\left(\frac{1}{2}+\sigma+i\left(\frac{m}{\chi}+t\right)\right)\right|^{2(q-1)}$$

Using the second part of Euler Product Result 2.2, this is

$$e^{O(q^2)}(|m|+1)^{2(q-1)} \left(1 + \frac{\mathcal{X}}{(|m|+1)\log q}\right)^{(q-1)^2 - (q-1)} \left(\frac{\mathcal{X}}{\log q}\right)^{1 + (q-1)^2} \cdot \left(\left(1 + \frac{\mathcal{X}}{(|m-n|+1)\log q}\right) \left(1 + \frac{\mathcal{X}}{(|m+n|+1)\log q}\right)\right)^{2(q-1)}.$$

Now depending on the signs of *m*, *n*, one of the terms |m - n|, |m + n| will be equal to ||m| - |n|| and the other will equal $|m| + |n| \ge |m|$. So the above is always

$$\leq e^{O(q^2)} (|m|+1)^{2(q-1)} \left(1 + \frac{\mathcal{X}}{(|m|+1)\log q}\right)^{q(q-1)} \left(\frac{\mathcal{X}}{\log q}\right)^{1 + (q-1)^2} \left(\frac{\mathcal{X}}{\left(\left||m| - |n|\right| + 1\right)\log q}\right)^{2(q-1)}.$$

Putting this together with (5-4) and (5-2), if we first perform the sum over $|n| \leq \mathcal{X}/2 + 1$ we get that $\left\| \int_{-1/2}^{1/2} \left| F_k \left(\frac{1}{2} + \sigma + it \right) \right|^2 dt \right\|_q$ is at most

$$\frac{e^{O(q)}}{\mathcal{X}} \bigg(\sum_{|m| \le \mathcal{X}/2+1} \frac{(|m|+1)^{2(q-1)}}{(|m|+1)^2} \bigg(1 + \frac{\mathcal{X}}{(|m|+1)\log q} \bigg)^{q(q-1)} \bigg(\frac{\mathcal{X}}{\log q} \bigg)^{1+(q-1)^2+2(q-1)} \bigg)^{1/q}.$$

Finally performing the sum over *m*, the dominant contribution comes from small terms (note that for terms with $|m| > \mathcal{X}/\log q$ we have $(1 + \mathcal{X}/((|m| + 1)\log q))^{q(q-1)} = e^{O(q^2)}$, so overall these contribute at most $e^{O(q^2)}\mathcal{X}^{2q-3}(\mathcal{X}/\log q)^{1+(q-1)^2+2(q-1)} = e^{O(q^2)}(\mathcal{X}/\log q)^{q^2+2q-3}$ inside the bracket), and gives us a bound $\ll [e^{O(q)}/\mathcal{X}](\mathcal{X}/\log q)^{(2q^2-q)/q}$. Raising everything to the power $q \ge 2$ and inserting the fact that $\mathcal{X} = \min\{\log x/e^{k+1}, 1/|\sigma|\}$, this gives the bound claimed in Key Proposition 5.2.

Proof of Key Proposition 5.1, for 1 < q < 2**.** When 1 < q < 2, it is not immediately obvious how to analyze the term $\mathbb{E} |F_k(\frac{1}{2} + \sigma)|^2 (\sum_{|m| \le \mathcal{X}} |F_k(\frac{1}{2} + \sigma + \frac{im}{\mathcal{X}})|^2)^{q-1}$ in (5-1). We may begin by letting $C = C(q) = e^{1/(q-1)}$, and noting that $\mathbb{E} |F_k(\frac{1}{2} + \sigma)|^2 (\sum_{|m| \le \mathcal{X}} |F_k(\frac{1}{2} + \sigma + \frac{im}{\mathcal{X}})|^2)^{q-1}$ is

$$\leq \mathbb{E} \left| F_k \left(\frac{1}{2} + \sigma \right) \right|^2 \left(\sum_{d \leq (q-1)\log \mathcal{X} + 1} \sum_{C^{d-1} \leq |m| \leq C^d} \left| F_k \left(\frac{1}{2} + \sigma + \frac{im}{\mathcal{X}} \right) \right|^2 \right)^{q-1}$$

Here we adopt the convention that the term m = 0 is included in $\sum_{C^{d-1} \le |m| \le C^d}$ when d = 1, and that any terms with $|m| > \mathcal{X}$ are omitted from all sums (so the imaginary shift in the second copy of F_k always has size $|m/\mathcal{X}| \le 1$). The motivation for splitting things up like this is that we expect our estimates for all terms with $\sum_{C^{d-1} \le |m| \le C^d}$ to be roughly the same, up to a factor $C^{O(1)}$. And, when everything is raised to the power q - 1, this factor simply becomes a constant multiplier. Next, if we let $D = D(q) \in \mathbb{N}$ be a parameter, to be fixed later, we can split things up further and find

$$\mathbb{E} |F_{k}(\frac{1}{2}+\sigma)|^{2} \left(\sum_{|m|\leq\mathcal{X}} |F_{k}(\frac{1}{2}+\sigma+\frac{im}{\mathcal{X}})|^{2}\right)^{q-1} \\ \leq \sum_{r\leq(q-1)\log\mathcal{X}/D+1} \mathbb{E} |F_{k}(\frac{1}{2}+\sigma)|^{2} \left(\sum_{(r-1)D(5-5)$$

Notice that we may further assume that all terms *d* for which $C^{d-1} > \mathcal{X}$ are omitted here, since for those the sum over *m* is empty (by our earlier convention).

Now in the sum over *m*, we expect (thinking about Euler Product Result 2.1) that the part of the Euler product $F_k(\frac{1}{2} + \sigma + \frac{im}{\chi})$ on primes $\leq e^{\chi/C^d}$ will be roughly the same size for all $C^{d-1} \leq |m| \leq C^d$, and indeed roughly the same size as the corresponding part of $F_k(\frac{1}{2} + \sigma)$. To simplify our writing about this, for each $d \geq 1$ and $|m| \leq \chi$ let us set

$$G_d(m) := \prod_{p \le e^{\mathcal{X}/C^d}} \left| 1 - \frac{f(p)}{p^{1/2 + \sigma + im/\mathcal{X}}} \right|^{-2}, \quad \text{and} \quad H_d(m) := \frac{\left| F_k \left(\frac{1}{2} + \sigma + \frac{im}{\mathcal{X}} \right) \right|^2}{G_d(m)}.$$

(These quantities of course depend on x, k, σ as well, but we suppress that in our notation.) We will also set $G_d := G_d(0)$ and $H_d := H_d(0)$. Then the expectation in (5-5) may be written as

$$\mathbb{E} \max_{(r-1)D < d \le rD} G_d H_d \left(\sum_{C^{d-1} \le |m| \le C^d} G_d(m) H_d(m) \right)^{q-1}$$

We want to apply Hölder's inequality to this expectation, in such a way that the bracketed sum is raised to the power 1/(q-1), and so we can connect up the expectation with the terms inside. Prior to doing

this, we rewrite the expectation again as

$$\mathbb{E} \max_{(r-1)D < d \le rD} G_d^{1-(q-1)^2} H_d^{1-(q-1)} C^{2(q-1)(2-q)d} \left(\frac{G_d^{q-1} H_d}{C^{2(2-q)d}} \sum_{C^{d-1} \le |m| \le C^d} G_d(m) H_d(m) \right)^{q-1}.$$

Simplifying the various exponents, this is all

$$\leq \mathbb{E}\left(\max_{(r-1)D < d \leq rD} G_d^{q(2-q)} H_d^{2-q} C^{2(q-1)(2-q)d}\right) \left(\max_{(r-1)D < d \leq rD} \frac{G_d^{q-1} H_d}{C^{2(2-q)d}} \sum_{C^{d-1} \leq |m| \leq C^d} G_d(m) H_d(m)\right)^{q-1},$$

and now using Hölder's inequality with exponents 1/(2-q) and 1/(q-1), we get a bound

$$\leq \left(\mathbb{E} \max_{(r-1)D < d \le rD} G_d^q H_d C^{2(q-1)d} \right)^{2-q} \left(\mathbb{E} \max_{(r-1)D < d \le rD} \frac{G_d^{q-1} H_d}{C^{2(2-q)d}} \sum_{C^{d-1} \le |m| \le C^d} G_d(m) H_d(m) \right)^{q-1} \\ \leq \left(\mathbb{E} \max_{(r-1)D < d \le rD} G_d^q H_d C^{2(q-1)d} \right)^{2-q} \left(\sum_{(r-1)D < d \le rD} \mathbb{E} \frac{G_d^{q-1} H_d}{C^{2(2-q)d}} \sum_{C^{d-1} \le |m| \le C^d} G_d(m) H_d(m) \right)^{q-1}.$$

We remark that the motivation for the uneven splitting of the Euler products here (moving $G_d^{(q-1)^2}$ and H_d^{q-1} into the second bracket) is that, as noted above, $\mathbb{E}G_d^{q-1}G_d(m)$ will behave in approximately the same way as $\mathbb{E}G_d^q$, but on the large primes the shifts im/\mathcal{X} provide extra cancellation in $\mathbb{E}H_dH_d(m)$. So the best way to split up the G_d terms is "evenly", i.e., such that the total exponent of G_d terms in both brackets after Hölder's inequality remains q, whereas for the H_d terms it is better to move a larger piece inside the second bracket (with the sum over m) to maximize the cancellation we pick up. As we shall see, the powers of C that we have introduced will serve to balance the final sizes of all the terms.

Using the independence of f(p) for different primes, together with Euler Product Result 2.1, the sums inside the second bracket are

$$\begin{split} &\sum_{(r-1)D < d \le rD} \frac{1}{C^{2(2-q)d}} \sum_{C^{d-1} \le |m| \le C^d} \mathbb{E}G_d^{q-1}G_d(m)\mathbb{E}H_dH_d(m) \\ &\ll \sum_{(r-1)D < d \le rD} \frac{1}{C^{2(2-q)d}} \sum_{C^{d-1} \le |m| \le C^d} \left(1 + \frac{\mathcal{X}}{C^d}\right)^{1 + (q-1)^2} \left(\frac{\mathcal{X}}{1 + |m|}\right)^{2(q-1)} (\min\{\mathcal{X}, C^d\})^2 \left(\frac{\min\{\mathcal{X}, C^d\}}{1 + |m|}\right)^2 \\ &\ll \sum_{(r-1)D < d \le rD} \frac{C^{O(1)}}{C^{2(2-q)d}} \mathcal{X}^{q^2} C^{(3-q^2)d} \\ &= C^{O(1)} \mathcal{X}^{q^2} \sum_{(r-1)D < d \le rD} C^{-(q-1)^2d} = C^{O(1)} \mathcal{X}^{q^2} C^{-(q-1)^2(r-1)D}. \end{split}$$

When performing this calculation, we noted that the contribution to $\mathbb{E}H_dH_d(m)$ from primes $p > e^{\mathcal{X}}$ is uniformly bounded (by the first part of Euler Product Result 2.1), similarly as in our analysis of the case $q \ge 2$. Some of our estimates here were a bit crude, but there seems to be no way to avoid losing some

factors $C^{O(1)}$, which further explains why our choice of $C = e^{1/(q-1)}$ is essentially the largest we can make without incurring unacceptable losses.

To bound $\mathbb{E} \max_{(r-1)D < d \le rD} G_d^q H_d C^{2(q-1)d}$, where we need to handle the maximum in a nontrivial way rather than replacing it by a sum (because this term will not be raised to the small power q-1), we will use Probability Result 2.5. To do this, we first rewrite $\mathbb{E} \max_{(r-1)D < d \le rD} G_d^q H_d C^{2(q-1)d}$ as

$$\mathbb{E} |F_k(\frac{1}{2} + \sigma)|^2 \max_{(r-1)D < d \le rD} (G_d C^{2d})^{q-1} \\ = \mathbb{E} |F_k(\frac{1}{2} + \sigma)|^2 \tilde{\mathbb{E}} \max_{(r-1)D < d \le rD} \prod_{\substack{p < e^{\mathcal{X}/C^d}}} \left|1 - \frac{f(p)}{p^{1/2+\sigma}}\right|^{-2(q-1)d} C^{2(q-1)d},$$

where $\tilde{\mathbb{E}}$ is expectation under the "tilted" measure defined by $\tilde{\mathbb{P}}(A) = \mathbb{E} |F_k(\frac{1}{2} + \sigma)|^2 \mathbf{1}_A / \mathbb{E} |F_k(\frac{1}{2} + \sigma)|^2$ for each event *A* (and **1** denotes the indicator function). We note, for use in a little while, that if *A* is an event not involving certain primes then those terms factor out from the expectation and cancel between numerator and denominator in the definition of $\tilde{\mathbb{P}}(A)$. Furthermore, the random variables f(p) are still independent under the measure $\tilde{\mathbb{P}}$, since if *A*, *B* are events involving disjoint sets of primes then we can split up the Euler product $|F_k(\frac{1}{2} + \sigma)|^2$ into subproducts over the corresponding sets, and then the expectation \mathbb{E} will split up correspondingly.

Now Euler Product Result 2.1 implies that $\mathbb{E}|F_k(\frac{1}{2}+\sigma)|^2 \simeq \mathcal{X}$. Furthermore, if we write

$$L_d := \prod_{p \le e^{\mathcal{X}/C^d}} \left| 1 - \frac{f(p)}{p^{1/2+\sigma}} \right|^{-2(q-1)/1.01}$$

(say), and $\lambda_d := \tilde{\mathbb{E}}L_d$, then Euler Product Result 2.1 and the independence of the f(p) imply that

$$\lambda_{d} = \frac{\mathbb{E} \prod_{p \le e^{\mathcal{X}/C^{d}}} |1 - f(p)/p^{1/2 + \sigma}|^{-2(1 + (q-1)/1.01)}}{\mathbb{E} \prod_{p \le e^{\mathcal{X}/C^{d}}} |1 - f(p)/p^{1/2 + \sigma}|^{-2}} \asymp \left(1 + \frac{\mathcal{X}}{C^{d}}\right)^{2(q-1)/1.01 + (q-1)^{2}/1.01}$$

We similarly get that $\tilde{\mathbb{E}}L_d^{1,01} \approx (1 + \mathcal{X}/C^d)^{2(q-1)+(q-1)^2}$. So we have shown that

$$\mathbb{E} \max_{(r-1)D < d \le rD} G_d^q H_d C^{2(q-1)d} \asymp \mathcal{X} \cdot \tilde{\mathbb{E}} \max_{(r-1)D < d \le rD} \left(\frac{L_d}{\lambda_d}\right)^{1.01} (C^d + \mathcal{X})^{2(q-1)} \left(1 + \frac{\mathcal{X}}{C^d}\right)^{(q-1)^2/1.01} \\ \ll \mathcal{X}^{1+2(q-1)} \left(1 + \frac{\mathcal{X}}{C^{(r-1)D+1}}\right)^{(q-1)^2/1.01} \tilde{\mathbb{E}} \max_{(r-1)D < d \le rD} \left(\frac{L_d}{\lambda_d}\right)^{1.01},$$

where we used the fact that $C^d \leq C\mathcal{X}$ (given our convention that those *d* for which $C^{d-1} > \mathcal{X}$ are omitted) and $C^{2(q-1)} \ll 1$. Finally, since the f(p) are independent under the measure $\tilde{\mathbb{P}}$ (and so the "increments" of different primes in the Euler product are independent), the sequence of random variables $(L_{rD}/\lambda_{rD}), (L_{rD-1}/\lambda_{rD-1}), \ldots, (L_{(r-1)D+1}/\lambda_{(r-1)D+1})$ (taken in that order) form a nonnegative submartingale relative to $\tilde{\mathbb{P}}$ and to the sigma algebras generated by $(f(p))_{p \leq e^{\mathcal{X}/C^{rD}}}, (f(p))_{p \leq e^{\mathcal{X}/C^{rD-1}}}, \ldots, (f(p))_{p < e^{\mathcal{X}/C^{r-1}/D+1}}$. For example, we may calculate the conditional expectation

$$\begin{split} \tilde{\mathbb{E}}\left(\!\left(\frac{L_{rD-1}}{\lambda_{rD-1}}\right) \mid (f(p))_{p \le e^{\mathcal{X}/C^{rD}}}\right) &= \frac{L_{rD}}{\lambda_{rD-1}} \tilde{\mathbb{E}}\left(\prod_{e^{\mathcal{X}/C^{rD}}$$

which gives the "nondecreasing on average" property from the definition of a submartingale. Thus Probability Result 2.5 is applicable, and gives that $\tilde{\mathbb{E}} \max_{(r-1)D < d \le rD} (L_d/\lambda_d)^{1.01}$ is

$$\ll \tilde{\mathbb{E}} \left(\frac{L_{(r-1)D+1}}{\lambda_{(r-1)D+1}} \right)^{1.01} \asymp \frac{\left(1 + \frac{\mathcal{X}}{C^{(r-1)D+1}} \right)^{2(q-1)+(q-1)^2}}{\lambda_{(r-1)D+1}^{1.01}} \asymp \left(1 + \frac{\mathcal{X}}{C^{(r-1)D+1}} \right)^{(q-1)^2 - \frac{(q-1)^2}{1.01}}$$

Putting together (5-1), (5-5), and the above calculations, we get that

$$\left\| \int_{-1/2}^{1/2} \left| F_k \left(\frac{1}{2} + \sigma + it \right) \right|^2 dt \right\|_q \\ \ll \frac{1}{\mathcal{X}} \left(\mathcal{X} D^{q-1} \sum_{r \le ((q-1)\log \mathcal{X})/D+1} \left(\frac{\mathcal{X}^{q^2}}{C^{(r-1)D(q-1)^2}} \right)^{2-q} (C^{O(1)} \mathcal{X}^{q^2} C^{-(q-1)^2(r-1)D})^{q-1} \right)^{1/q}.$$

Recalling that $C = e^{1/(q-1)}$ and collecting terms together, we find this is all

$$\ll \frac{1}{\mathcal{X}} \left(\mathcal{X}^{1+q^2} D^{q-1} \sum_{r \le (q-1)\log \mathcal{X}/D+1} C^{-(r-1)D(q-1)^2} \right)^{1/q} = \frac{1}{\mathcal{X}} \left(\mathcal{X}^{1+q^2} D^{q-1} \sum_{r \le (q-1)\log \mathcal{X}/D+1} e^{-(r-1)D(q-1)} \right)^{1/q}.$$

So if we finally choose $D := \lfloor 1/(q-1) \rfloor$, then both the sum over *r* and the term D^{q-1} will be $\ll 1$. Recalling that $\mathcal{X} = \min\{(\log x)/e^{k+1}, 1/|\sigma|\}$, we see this bound is as claimed in Key Proposition 5.1.

Proof of Key Proposition 5.2, for 1 < q < 2**.** We again begin with the second part of the proposition, where $|N| \ge 1$. In this case we can analyze the terms $\mathbb{E} |F_k(\frac{1}{2} + \sigma + i(\frac{n}{\chi} + N + t))|^2 (\sum_{|m| \le \frac{\chi}{2} + 1} |F_k(\frac{1}{2} + \sigma + i(\frac{m}{\chi} + N + t))|^2)^{q-1}$ in (5-2) by splitting the sum over *m* into subsums where $C^{d-1} \le |m-n| \le C^d$, and otherwise following the argument from the Steinhaus case. We obtain the same estimates as there, except the error term in Euler Product Result 2.2 produces an additional factor

$$\min\left\{1+\frac{\mathcal{X}}{C^{d}}, |N|^{1/100}\right\}^{|(q-1)^{2}-(q-1)|+4(q-1)} \min\left\{C^{d}, \mathcal{X}, 1+\frac{|N|^{1/100}}{1+\mathcal{X}/C^{d}}\right\}^{4} \ll \min\{\mathcal{X}, |N|^{1/100}\}^{4}$$

when estimating (the analogue of) the terms $\mathbb{E}G_d^{q-1}G_d(m)\mathbb{E}H_dH_d(m)$, and an additional factor min $\{1 + \mathcal{X}/C^{(r-1)D+1}, |N|^{1/100}\}^{q(q-1)} \ll \min\{\mathcal{X}, |N|^{1/100}\}^{q(q-1)}$ when estimating (the analogue of) the term $\mathbb{E}\max_{(r-1)D < d \le rD} G_d^q H_d C^{2(q-1)d}$. So overall we get the same bound as in the Steinhaus case, apart from

a factor

$$(\min\{\mathcal{X}, |N|^{1/100}\}^{q(q-1)})^{2-q} (\min\{\mathcal{X}, |N|^{1/100}\}^4)^{(q-1)} \ll \min\{\mathcal{X}, |N|^{1/100}\}^{(q-1)(4+2q-q^2)}$$

A small calculation shows that for $1 \le q \le 2$, we have $(q-1)(4+2q-q^2) \le 5(q-1) \le q(q+1)$, giving the factor $\min\{\mathcal{X}, |N|^{1/100}\}^{q(q+1)} = \min\{(\log x)/e^{k+1}, 1/|\sigma|, |N|^{1/100}\}^{q(q+1)}$ claimed in the second part of Key Proposition 5.2.

When N = 0, to prove Key Proposition 5.2 we need to bound

$$\sum_{|n|\leq \mathcal{X}/2+1} \mathbb{E} \left| F_k \left(\frac{1}{2} + \sigma + i \left(\frac{n}{\mathcal{X}} + t \right) \right) \right|^2 \left(\sum_{|m|\leq \mathcal{X}/2+1} \left| F_k \left(\frac{1}{2} + \sigma + i \left(\frac{m}{\mathcal{X}} + t \right) \right) \right|^2 \right)^{q-1}$$

in (5-2). Following the same argument that led to the bound (5-5) in the Steinhaus case, but now splitting the sum over *m* according to the size of |m| - |n| rather than the size of |m|, one obtains that

$$\sum_{|n| \le \mathcal{X}/2+1} \mathbb{E} |F_{k}(\frac{1}{2} + \sigma + i(\frac{n}{\mathcal{X}} + t))|^{2} \left(\sum_{|m| \le \mathcal{X}/2+1} |F_{k}(\frac{1}{2} + \sigma + i(\frac{m}{\mathcal{X}} + t))|^{2} \right)^{q-1} \\ \ll D^{q-1} \sum_{|n| \le \mathcal{X}/2+1} \sum_{r \le (q-1) \log \mathcal{X}/D+1} \mathbb{E}_{(r-1)D < d \le rD} \left|F_{k}(\frac{1}{2} + \sigma + i(\frac{n}{\mathcal{X}} + t))\right|^{2} \\ \cdot \left(\sum_{\substack{|m| \le \mathcal{X}/2+1, \\ C^{d-1} \le ||m| - |n|| \le C^{d}}} |F_{k}(\frac{1}{2} + \sigma + i(\frac{m}{\mathcal{X}} + t))|^{2} \right)^{q-1}.$$
(5-6)

Here we again have $C = e^{1/(q-1)}$, and $D = D(q) \in \mathbb{N}$ is a parameter, and we adopt our usual conventions (analogously to the Steinhaus case) about including the |m| = |n| term when d = 1 and omitting overly large terms from all sums.

Now for each $d \ge 1$ and $|m| \le \mathcal{X}/2 + 1$, and treating t as fixed, we shall set

$$G_d(m) := \prod_{p \le e^{\mathcal{X}/C^d}} \left| 1 + \frac{f(p)}{p^{1/2 + \sigma + i(m/\mathcal{X} + i)}} \right|^2, \quad \text{and} \quad H_d(m) := \frac{\left| F_k \left(\frac{1}{2} + \sigma + i\left(\frac{m}{\mathcal{X}} + i\right) \right) \right|^2}{G_d(m)}$$

This is the same notation that we used in the Steinhaus case, but with the Euler products now replaced by their Rademacher versions (supported on squarefree numbers only). Splitting the expectation and applying Hölder's inequality as in the Steinhaus case, it follows that (5-6) is

$$\ll D^{q-1} \sum_{|n| \le \mathcal{X}/2+1} \sum_{r \le (q-1)\log \mathcal{X}/D+1} \left(\mathbb{E} \max_{(r-1)D < d \le rD} G_d(n)^q H_d(n) C^{2(q-1)d} \right)^{2-q} \cdot \left(\sum_{(r-1)D < d \le rD} \mathbb{E} \frac{G_d(n)^{q-1} H_d(n)}{C^{2(2-q)d}} \sum_{\substack{|m| \le \mathcal{X}/2+1, \\ C^{d-1} \le ||m| - |n|| \le C^d}} G_d(m) H_d(m) \right)^{q-1}.$$

Continuing to follow the argument from the Steinhaus case, but using Euler Product Result 2.2 in place of Euler Product Result 2.1, we can bound these terms further. Proceeding to do this, and noting that one

of the terms |m - n|, |m + n| that arise in Euler Product Result 2.2 will always equal ||m| - |n|| and the other will equal $|m| + |n| \ge |n|$, we find the sums in the second bracket are

$$\sum_{(r-1)D < d \le rD} \frac{1}{C^{2(2-q)d}} \sum_{\substack{|m| \le \mathcal{X}/2+1, \\ C^{d-1} \le ||m| - |n|| \le C^d}} \mathbb{E}G_d(n)^{q-1}G_d(m)\mathbb{E}H_d(n)H_d(m)$$

$$\ll \sum_{(r-1)D < d \le rD} \frac{1}{C^{2(2-q)d}} \sum_{\substack{C^{d-1} \le ||m| - |n|| \le C^d}} \min\left\{\frac{\mathcal{X}}{C^d}, \frac{\mathcal{X}}{|n|}\right\}^{(q-1)^2 - (q-1)} \left(1 + \frac{\mathcal{X}}{C^d}\right)^{1 + (q-1)^2} \cdot \left(\frac{\mathcal{X}}{1 + ||m| - |n||} \min\left\{\frac{\mathcal{X}}{C^d}, \frac{\mathcal{X}}{|n|}\right\}\right)^{2(q-1)} (\min\{\mathcal{X}, C^d\})^2 \left(\frac{\min\{\mathcal{X}, C^d\}}{1 + ||m| - |n||} \left(1 + \frac{\min\{\mathcal{X}, C^d\}}{1 + |m| + |n|}\right)\right)^2.$$

Collecting terms together, and then upper bounding $\min\{\mathcal{X}/C^d, \mathcal{X}/|n|\}^{q(q-1)}$ by $(\mathcal{X}/(1+|n|))^{q(q-1)}$ and upper bounding $\min\{\mathcal{X}, C^d\}$ everywhere else by C^d , the above is

$$\ll \left(\frac{\mathcal{X}}{1+|n|}\right)^{q(q-1)} \sum_{(r-1)D < d \le rD} \frac{C^{O(1)}}{C^{2(2-q)d}} \mathcal{X}^{q^2} C^{(3-q^2)d} = \left(\frac{\mathcal{X}}{1+|n|}\right)^{q(q-1)} C^{O(1)} \mathcal{X}^{q^2} C^{-(q-1)^2(r-1)D}.$$

We can also adapt the Steinhaus argument to bound $\mathbb{E} \max_{(r-1)D < d \le rD} G_d(n)^q H_d(n) C^{2(q-1)d}$. In this case we again have $\mathbb{E} |F_k(\frac{1}{2} + \sigma + i(\frac{n}{\mathcal{X}} + t))|^2 \asymp \mathcal{X}$, and we may define the "tilted" measure $\tilde{\mathbb{P}}$ and set $L_d := \prod_{p \le e^{\mathcal{X}/C^d}} |1 + f(p)/p^{1/2 + \sigma + i(n/\mathcal{X}+t)}|^{2(q-1)/1.01}$ analogously to the Steinhaus case. Then Euler Product Result 2.2 implies that $\tilde{\mathbb{E}}L_d \asymp \min{\{\mathcal{X}/C^d, \mathcal{X}/|n|\}^{(1+(q-1)/1.01)(q-1)/1.01}(1 + \mathcal{X}/C^d)^{2(q-1)/1.01+(q-1)^2/1.01^2}}$ and $\tilde{\mathbb{E}}L_d^{1.01} \asymp \min{\{\mathcal{X}/C^d, \mathcal{X}/|n|\}^{q(q-1)}(1 + \mathcal{X}/C^d)^{2(q-1)+(q-1)^2}}$. So the same submartingale argument as in the Steinhaus case shows that

$$\mathbb{E} \max_{(r-1)D < d \le rD} G_d(n)^q H_d(n) C^{2(q-1)d} \ll \mathcal{X}^{1+2(q-1)} \left(1 + \frac{\mathcal{X}}{C^{(r-1)D+1}} \right)^{(q-1)^2} \min\left\{ \frac{\mathcal{X}}{C^{(r-1)D+1}}, \frac{\mathcal{X}}{|n|} \right\}^{q(q-1)} \\ \ll \frac{\mathcal{X}^{q^2}}{C^{(r-1)D(q-1)^2}} \left(\frac{\mathcal{X}}{1+|n|} \right)^{q(q-1)}.$$

Putting everything together, recalling that $C = e^{1/(q-1)}$ and choosing $D := \lfloor 1/(q-1) \rfloor$, we deduce that (5-6) is

$$\ll D^{q-1} \sum_{|n| \le \frac{\mathcal{X}}{2} + 1} \sum_{r \le \frac{(q-1)\log \mathcal{X}}{D} + 1} \frac{\mathcal{X}^{q^2}}{C^{(r-1)D(q-1)^2}} \left(\frac{\mathcal{X}}{1+|n|}\right)^{q(q-1)} \ll \mathcal{X}^{q^2} \sum_{|n| \le \frac{\mathcal{X}}{2} + 1} \left(\frac{\mathcal{X}}{1+|n|}\right)^{q(q-1)}$$

Since $q_0 = (1 + \sqrt{5})/2 \approx 1.618$ satisfies $q_0(q_0 - 1) = 1$, and we have 1 < q < 2, the sum over n here is $\ll \mathcal{X}^{\max\{1,q(q-1)\}} \min\{\log \mathcal{X}, 1/|q - q_0|\}$. Substituting into (5-2), and recalling that $\mathcal{X} := \min\{\log x/e^{k+1}, 1/|\sigma|\}$, this gives the first (N = 0) bound claimed in Key Proposition 5.2 when 1 < q < 2.

6. Proofs of the lower bounds in Theorems 1.1 and 1.2

Recall that F(s) denotes the Euler product of f(n) over x-smooth numbers.

The lower bound in the Steinhaus case. To prove the lower bound part of Theorem 1.1, in view of Proposition 4.3 our main work will be to prove a suitable lower bound for $\|\int_{-1/2}^{1/2} |F(\frac{1}{2} + \frac{4Vq}{\log x} + it)|^2 dt\|_q$, where *V* is a large fixed constant. (We also need an upper bound for the subtracted quantity $\|\int_{-1/2}^{1/2} |F(\frac{1}{2} + \frac{2Vq}{\log x} + it)|^2 dt\|_q$, but this will follow directly from Key Proposition 5.1.)

To obtain our lower bound, we note first that for any $q \ge 1$ we have

$$\left(\int_{-1/2}^{1/2} \left| F\left(\frac{1}{2} + \frac{4Vq}{\log x} + it\right) \right|^2 dt \right)^q \ge \left(\sum_{|k| \le (\log x - 1)/2} \int_{(k-1/2)/\log x}^{(k+1/2)/\log x} \left| F\left(\frac{1}{2} + \frac{4Vq}{\log x} + it\right) \right|^2 dt \right)^q \\ \ge \sum_{|k| \le (\log x - 1)/2} \left(\int_{(k-1/2)/\log x}^{(k+1/2)/\log x} \left| F\left(\frac{1}{2} + \frac{4Vq}{\log x} + it\right) \right|^2 dt \right)^q.$$

This step could be wasteful if many of the pieces $\int_{(k-1/2)/\log x}^{(k+1/2)/\log x} |F(\frac{1}{2} + \frac{4Vq}{\log x} + it)|^2 dt$ made substantial contributions to the full integral. But for large q we expect instead that the dominant contribution should come from just a few large (and therefore rare) contributions, so we will not lose too much. In the Steinhaus case, since the distribution of $\int_{(k-1/2)/\log x}^{(k+1/2)/\log x} |F(\frac{1}{2} + \frac{4Vq}{\log x} + it)|^2 dt$ is independent of k we get the simpler lower bound

$$\mathbb{E}\bigg(\int_{-1/2}^{1/2} \left| F\bigg(\frac{1}{2} + \frac{4Vq}{\log x} + it\bigg) \right|^2 dt \bigg)^q \gg \log x \cdot \mathbb{E}\bigg(\int_{-1/(2\log x)}^{1/(2\log x)} \left| F\bigg(\frac{1}{2} + \frac{4Vq}{\log x} + it\bigg) \right|^2 dt \bigg)^q.$$

Now we want to remove the remaining short integral over t, which is a technical obstacle to connecting up the expectation with the random product F. Heuristically, since the Euler product shouldn't vary much on intervals of length $1/\log x$ we should simply obtain something like $\left(\frac{1}{\log x}\left|F\left(\frac{1}{2}+\frac{4Vq}{\log x}\right)\right|^2\right)^q$ in the bracket. It turns out that a neat way to handle this issue is using Jensen's inequality (applied to the normalized integral $\int_{-1/2\log x}^{1/2\log x} \log x \, dt$), which implies that

$$\begin{split} \mathbb{E}\bigg(\int_{-1/(2\log x)}^{1/(2\log x)} \left| F\bigg(\frac{1}{2} + \frac{4Vq}{\log x} + it\bigg) \right|^2 dt \bigg)^q \\ &= \frac{1}{\log^q x} \mathbb{E}\bigg(\int_{-1/(2\log x)}^{1/(2\log x)} \log x \cdot e^{2\log|F(1/2 + 4Vq/\log x + it)|} dt \bigg)^q \\ &\geq \frac{1}{\log^q x} \mathbb{E}\bigg(\exp\bigg\{\int_{-1/(2\log x)}^{1/(2\log x)} \log x \cdot 2\log \bigg| F\bigg(\frac{1}{2} + \frac{4Vq}{\log x} + it\bigg) \bigg| dt\bigg\}\bigg)^q \\ &= \frac{1}{\log^q x} \mathbb{E}\exp\bigg\{2q \int_{-1/(2\log x)}^{1/(2\log x)} \log x \cdot \log \bigg| F\bigg(\frac{1}{2} + \frac{4Vq}{\log x} + it\bigg) \bigg| dt\bigg\}. \end{split}$$

Here the exponential, inside the expectation, may be rewritten as

$$\begin{split} \prod_{p \le x} \exp\left\{-2q \int_{-1/(2\log x)}^{1/(2\log x)} \log x \cdot \Re \log\left(1 - \frac{f(p)}{p^{1/2 + 4Vq/\log x + it}}\right) dt\right\} \\ &= \prod_{p \le x} \exp\left\{2q \log x \Re\left(\frac{f(p)}{p^{1/2 + 4Vq/\log x}} \int_{-1/(2\log x)}^{1/(2\log x)} e^{-it\log p} dt + \frac{f(p)^2}{2p^{1 + 8Vq/\log x}} \int_{-1/(2\log x)}^{1/(2\log x)} e^{-2it\log p} dt\right) + O\left(\frac{q}{p^{3/2}}\right)\right\} \\ &= e^{O(q)} \prod_{p \le x} \exp\left\{2q \Re\left(\frac{f(p)}{p^{1/2 + 4Vq/\log x}} \log x \int_{-1/(2\log x)}^{1/(2\log x)} e^{-it\log p} dt + \frac{f(p)^2}{2p^{1 + 8Vq/\log x}}\right)\right\}. \end{split}$$

The first equality here uses the Taylor expansion of the logarithm, and the second equality uses the estimate

$$\int_{-1/(2\log x)}^{1/(2\log x)} e^{-2it\log p} dt = \int_{-1/(2\log x)}^{1/(2\log x)} (1+O(|t|\log p)) dt = \frac{1}{\log x} + O\left(\frac{\log p}{\log^2 x}\right)$$

and also the fact that

$$\sum_{p \le x} \frac{\log p}{p^{1+8Vq/\log x}} \ll \log x.$$

Putting things together, using the independence of f(p) for different primes p to move the expectation inside the product, we have shown that our original object

$$\mathbb{E}\left(\int_{-1/2}^{1/2} \left| F\left(\frac{1}{2} + \frac{4Vq}{\log x} + it\right) \right|^2 dt \right)^q \\ \ge \frac{e^{O(q)}}{\log^{q-1} x} \prod_{p \le x} \mathbb{E} \exp\left\{2q \Re\left(\frac{f(p)}{p^{1/2 + 4Vq/\log x}} \log x \int_{-1/(2\log x)}^{1/(2\log x)} e^{-it\log p} dt + \frac{f(p)^2}{2p^{1+8Vq/\log x}}\right)\right\}.$$
(6-1)

It will be convenient to note some simple bounds for the quantity inside the exponential, which we will use shortly. Firstly, this quantity is always trivially bounded by $O(q/\sqrt{p})$. Secondly, using our previous calculation that

$$\int_{-1/(2\log x)}^{1/(2\log x)} e^{-it\log p} dt = \frac{1}{\log x} + O\left(\frac{\log p}{\log^2 x}\right),$$

we can obtain that

$$2q\Re\left(\frac{f(p)}{p^{1/2+4Vq/\log x}}\log x\int_{-1/(2\log x)}^{1/(2\log x)}e^{-it\log p}\,dt + \frac{f(p)^2}{2p^{1+8Vq/\log x}}\right)$$
$$= \frac{2q\Re f(p)}{p^{1/2+4Vq/\log x}} + O\left(\frac{q\log p}{p^{1/2+4Vq/\log x}\log x} + \frac{q}{p}\right).$$

To conclude, we note that certainly when $100q^2 \le p \le x$ we have, in view of the Taylor expansion of the exponential and the simple bounds noted above and the fact that $\mathbb{E}(\Re f(p))^2 = \frac{1}{2}$, that

$$\begin{split} \mathbb{E} \exp \bigg\{ 2q \Re \bigg(\frac{f(p)}{p^{1/2 + 4Vq/\log x}} \log x \int_{-1/(2\log x)}^{1/(2\log x)} e^{-it\log p} dt + \frac{f(p)^2}{2p^{1+8Vq/\log x}} \bigg) \bigg\} \\ &= \mathbb{E} \bigg(1 + 2q \Re \bigg(\frac{f(p)}{p^{1/2 + 4Vq/\log x}} \log x \int_{-1/(2\log x)}^{1/(2\log x)} e^{-it\log p} dt + \frac{f(p)^2}{2p^{1+8Vq/\log x}} \bigg) + \frac{2q^2 (\Re f(p))^2}{p^{1+8Vq/\log x}} \\ &+ O\bigg(\frac{q^2\log p}{p^{1+8Vq/\log x}\log x} + \frac{q^3}{p^{3/2}} \bigg) \bigg) \\ &= 1 + \frac{q^2}{p^{1+8Vq/\log x}} + O\bigg(\frac{q^2\log p}{p^{1+8Vq/\log x}\log x} + \frac{q^3}{p^{3/2}} \bigg). \end{split}$$

When $p < 100q^2$, we shall instead use the trivial bound $\exp\{O(q/\sqrt{p})\}$. Inserting these into (6-1), we get an overall lower bound

$$\begin{split} & \mathbb{E}\bigg(\int_{-1/2}^{1/2} \left|F\big(\frac{1}{2} + \frac{4Vq}{\log x} + it\big)\right|^2 dt\bigg)^q \\ & \geq \frac{e^{O(q)}}{\log^{q-1} x} \prod_{p<100q^2} \exp\bigg\{O\bigg(\frac{q}{\sqrt{p}}\bigg)\bigg\} \prod_{100q^2 \le p \le x} \exp\bigg\{\frac{q^2}{p^{1+8Vq/\log x}} + O\bigg(\frac{q^2\log p}{p^{1+8Vq/\log x}\log x} + \frac{q^3}{p^{3/2}}\bigg)\bigg\} \\ & = \frac{e^{O(q^2/\log(2q))}}{\log^{q-1} x} \prod_{100q^2 \le p \le x} \exp\bigg\{\frac{q^2}{p^{1+8Vq/\log x}}\bigg\} \\ & = \frac{e^{O(q^2)}}{\log^{q-1} x} \bigg(\frac{\log x}{Vq\log(2q)}\bigg)^{q^2}. \end{split}$$

Here the final equality follows because, similarly as in the calculations in Section 3, we have

$$\prod_{100q^2 \le p \le x} \exp\left\{\frac{q^2}{p^{1+8Vq/\log x}}\right\} = e^{O(q^2)} \zeta \left(1 + \frac{8Vq}{\log x}\right)^{q^2} e^{-\sum_{p < 100q^2} \frac{q^2}{p^{1+8Vq/\log x}}}.$$

Then

$$\zeta \left(1 + \frac{8Vq}{\log x}\right)^{q^2} = e^{O(q^2)} \left(\frac{\log x}{Vq}\right)^{q^2} \quad \text{and} \quad e^{-\sum_{p < 100q^2} \frac{q^2}{p^{1+8Vq/\log x}}} = \frac{e^{O(q^2)}}{\log^{q^2}(2q)}$$

by Mertens' estimates for sums over primes.

Inserting this into Proposition 4.3 we find that

$$\left\|\sum_{n \le x} f(n)\right\|_{2q} \gg \sqrt{\frac{x}{\log x}} \left(\frac{e^{O(q)}}{\log^{(q-1)/2q} x} \left(\frac{\log x}{Vq \log(2q)}\right)^{q/2} - \frac{C}{e^{Vq/2}} \left\|\int_{-1/2}^{1/2} \left|F\left(\frac{1}{2} + \frac{2Vq}{\log x} + it\right)\right|^2 dt\right\|_q^{1/2}\right)\right\|_q^{1/2} + \frac{C}{\log^{q-1}(q-1)/2q} \left\|\int_{-1/2}^{1/2} \left|F\left(\frac{1}{2} + \frac{2Vq}{\log x} + it\right)\right|^2 dt\right\|_q^{1/2}\right)\right\|_q^{1/2} + \frac{C}{\log^{q-1}(q-1)/2q} \left\|\int_{-1/2}^{1/2} \left|F\left(\frac{1}{2} + \frac{2Vq}{\log x} + it\right)\right|^2 dt\right\|_q^{1/2}\right)\right\|_q^{1/2} + \frac{C}{\log^{q-1}(q-1)/2q} \left\|\int_{-1/2}^{1/2} \left|F\left(\frac{1}{2} + \frac{2Vq}{\log x} + it\right)\right|^2 dt\right\|_q^{1/2}\right\|_q^{1/2} + \frac{C}{\log^{q-1}(q-1)/2q} \left\|\int_{-1/2}^{1/2} \left|F\left(\frac{1}{2} + \frac{2Vq}{\log x} + it\right)\right|^2 dt\right\|_q^{1/2}\right\|_q^{1/2} + \frac{C}{\log^{q-1}(q-1)/2q} \left\|\int_{-1/2}^{1/2} \left|F\left(\frac{1}{2} + \frac{2Vq}{\log x} + it\right)\right|^2 dt\right\|_q^{1/2}\right\|_q^{1/2} + \frac{C}{\log^{q-1}(q-1)/2q} \left\|\int_{-1/2}^{1/2} \left|F\left(\frac{1}{2} + \frac{2Vq}{\log x} + it\right)\right|^2 dt\right\|_q^{1/2}\right\|_q^{1/2} + \frac{C}{\log^{q-1}(q-1)/2q} \left\|\int_{-1/2}^{1/2} \left|F\left(\frac{1}{2} + \frac{2Vq}{\log x} + it\right)\right|^2 dt\right\|_q^{1/2}\right\|_q^{1/2} + \frac{C}{\log^{q-1}(q-1)/2q} \left\|\int_{-1/2}^{1/2} \left|F\left(\frac{1}{2} + \frac{2Vq}{\log x} + it\right)\right\|_q^{1/2} dt\right\|_q^{1/2} + \frac{C}{\log^{q-1}(q-1)/2q} \left\|\int_{-1/2}^{1/2} \left|F\left(\frac{1}{2} + \frac{2Vq}{\log x} + it\right)\right\|_q^{1/2} dt\right\|_q^{1/2} dt\right\|_q^{1/2} dt$$

And using Key Proposition 5.1 with k = -1 to control the subtracted term, provided that $2Vq/\log x \le 1/(100\log(2q))$ we can bound everything below by

$$\sqrt{\frac{x}{\log x}} \left(\frac{e^{O(q)}}{\log^{(q-1)/2q} x} \left(\frac{\log x}{Vq \log(2q)} \right)^{q/2} - \frac{Ce^{O(q)}}{e^{Vq/2}} \frac{(Vq)^{(q-1)/2q}}{\log^{(q-1)/2q} x} \left(\frac{\log x}{Vq \log(2q)} \right)^{q/2} \right)$$

If we set V to be a sufficiently large fixed constant, the subtracted term will be negligible compared with the first term, and our Theorem 1.1 lower bound will be proved. It only remains to note that the condition $2Vq/\log x \le 1/(100\log(2q))$ is then satisfied provided $q \le (c \log x)/\log \log x$, for a sufficiently small fixed constant c > 0.

The lower bound in the Rademacher case. To prove the lower bound part of Theorem 1.2, we shall invoke Proposition 4.4 and adapt the argument from the previous subsection to lower bound $\|\int_{-1/2}^{1/2} |F(\frac{1}{2} + \frac{4Vq}{\log x} + it)|^2 dt\|_q$, where V is a large fixed constant and now F(s) denotes the Rademacher random Euler product. Indeed, exactly the same argument as on page 2315 gives, for any $q \ge 1$, that

$$\mathbb{E}\left(\int_{-1/2}^{1/2} \left|F\left(\frac{1}{2} + \frac{4Vq}{\log x} + it\right)\right|^2 dt\right)^q \ge \sum_{|k| \le (\log x - 1)/2} \mathbb{E}\left(\int_{(k-1/2)/\log x}^{(k+1/2)/\log x} \left|F\left(\frac{1}{2} + \frac{4Vq}{\log x} + it\right)\right|^2 dt\right)^q.$$

Using Jensen's inequality, also as on page 2315, shows this is all

$$\geq \frac{1}{\log^{q} x} \sum_{|k| \leq (\log x - 1)/2} \mathbb{E} \exp\left\{ 2q \int_{(k - 1/2)/\log x}^{(k + 1/2)/\log x} \log x \cdot \log \left| F\left(\frac{1}{2} + \frac{4Vq}{\log x} + it\right) \right| dt \right\}.$$

And recalling that in the Rademacher case we have $F(s) = \prod_{p \le x} (1 + f(p)/p^s)$, with $f(p) \in \{\pm 1\}$, we find this is all

$$\begin{split} &= \frac{1}{\log^{q} x} \sum_{|k| \le (\log x - 1)/2} \prod_{p \le x} \mathbb{E} \exp\left\{ 2q \int_{(k-1/2)/\log x}^{(k+1/2)/\log x} \log x \cdot \Re \log\left(1 + \frac{f(p)}{p^{1/2 + 4Vq/\log x + it}}\right) dt \right\} \\ &= \frac{1}{\log^{q} x} \sum_{|k| \le (\log x - 1)/2} \prod_{p \le x} \mathbb{E} \exp\left\{ 2q \int_{(k-1/2)/\log x}^{(k+1/2)/\log x} \log x \cdot \left(\frac{f(p)\cos(t\log p)}{p^{1/2 + 4Vq/\log x}} - \frac{\cos(2t\log p)}{2p^{1+8Vq/\log x}} + O\left(\frac{1}{p^{3/2}}\right)\right) dt \right\}. \end{split}$$

At this stage we cannot efficiently remove the integral of $\cos(t \log p)$ in the first term, but for the second term we can write $\cos(2t \log p) = \cos(2k \log p/\log x) + O(\log p/\log x)$. The total contribution from these "big Oh" terms for all p, as well as from the $O(1/p^{3/2})$ term, is a multiplicative factor $e^{O(q)}$. So we obtain that

$$\mathbb{E}\left(\int_{-1/2}^{1/2} \left|F\left(\frac{1}{2} + \frac{4Vq}{\log x} + it\right)\right|^2 dt\right)^q \\ \ge \frac{e^{O(q)}}{\log^q x} \sum_{|k| \le (\log x - 1)/2} \prod_{p \le x} \mathbb{E} \exp\left\{2q\left(\frac{f(p)}{p^{1/2 + 4Vq/\log x}}\log x\right)\right) \\ \int_{(k-1/2)/\log x}^{(k+1/2)/\log x} \cos(t\log p) dt - \frac{\cos(2k\log p/\log x)}{2p^{1+8Vq/\log x}}\right)\right\}, \quad (6-2)$$

which is the Rademacher analogue of (6-1) from the Steinhaus case.

Next, when $100q^2 \le p \le x$ we have, in view of the Taylor expansion of the exponential (and the fact that $f(p)^2 \equiv 1$), that

$$\mathbb{E} \exp\left\{ 2q \left(\frac{f(p)}{p^{1/2+4Vq/\log x}} \log x \int_{(k-1/2)/\log x}^{(k+1/2)/\log x} \cos(t\log p) dt - \frac{\cos(2k\log p/\log x)}{2p^{1+8Vq/\log x}} \right) \right\}$$

$$= \mathbb{E} \left(1 + 2q \left(\frac{f(p)}{p^{1/2+4Vq/\log x}} \log x \int_{(k-1/2)/\log x}^{(k+1/2)/\log x} \cos(t\log p) dt - \frac{\cos(2k\log p/\log x)}{2p^{1+8Vq/\log x}} \right)$$

$$+ \frac{2q^2 \cos^2(k\log p/\log x)}{p^{1+8Vq/\log x}} + O\left(\frac{q^2\log p}{p^{1+8Vq/\log x}\log x} + \frac{q^3}{p^{3/2}} \right) \right).$$

Using the cosine identity $\cos^2(k \log p / \log x) = (\frac{1}{2})(1 + \cos(2k \log p / \log x))$, and the fact that $\mathbb{E}f(p) = 0$, we find the above is

$$= 1 + \frac{q^2 + (q^2 - q)\cos(2k\log p/\log x)}{p^{1 + 8Vq/\log x}} + O\left(\frac{q^2\log p}{p^{1 + 8Vq/\log x}\log x} + \frac{q^3}{p^{3/2}}\right)$$

When $p < 100q^2$, we shall instead use the trivial bound $\exp\{O(q/\sqrt{p})\}$. Inserting these into (6-2), we get

$$\mathbb{E}\left(\int_{-1/2}^{1/2} \left|F\left(\frac{1}{2} + \frac{4Vq}{\log x} + it\right)\right|^2 dt\right)^q \\ \ge \frac{e^{O(q^2/\log(2q))}}{\log^q x} \sum_{|k| \le (\log x - 1)/2} \prod_{100q^2 \le p \le x} \exp\left\{\frac{q^2 + (q^2 - q)\cos(2k\log p/\log x)}{p^{1 + 8Vq/\log x}}\right\}$$

Now when $2 \le q \le c \log x / \log \log x$, say, we can afford to discard all the terms in this lower bound except the k = 0 term, which gives us that

$$\mathbb{E}\left(\int_{-1/2}^{1/2} \left|F\left(\frac{1}{2} + \frac{4Vq}{\log x} + it\right)\right|^2 dt\right)^q \ge \frac{e^{O(q^2/\log(2q))}}{\log^q x} \prod_{100q^2 \le p \le x} \exp\left\{\frac{2q^2 - q}{p^{1+8Vq/\log x}}\right\}$$
$$= \frac{e^{O(q^2)}}{\log^q x} \left(\frac{\log x}{Vq\log(2q)}\right)^{2q^2 - q}.$$

Inserting this into Proposition 4.4, and applying Key Proposition 5.2 with k = -1 and $\sigma = 2Vq/\log x$ to control the subtracted term there, we find that $\left\|\sum_{n \le x} f(n)\right\|_{2q}$ is

$$\gg \sqrt{\frac{x}{\log x}} \left(\frac{e^{O(q)}}{\log^{1/2} x} \left(\frac{\log x}{Vq \log(2q)}\right)^{q-1/2} - \frac{Ce^{O(q)}}{e^{Vq/2}} \frac{(Vq)^{1/2}}{\log^{1/2} x} \left(\frac{\log x}{Vq \log(2q)}\right)^{q-1/2}\right).$$

If V is a sufficiently large constant, the subtracted term is negligible compared with the first term and we obtain the lower bound claimed in Theorem 1.2.

When $1 \le q < 2$ (or really when $1 \le q \le q_0 = (1 + \sqrt{5})/2$), we cannot afford to take quite such a crude approach. Using Chebychev's estimates and the prime number theorem as in the proof of Euler Product

Result 2.1, we have

$$\sum_{100q^2 \le p \le x} \frac{\cos(2k \log p/\log x)}{p^{1+8Vq/\log x}} = \sum_{2 \le p \le x^{1/V}} \frac{\cos(2k \log p/\log x)}{p} + O(1)$$
$$= \int_{\log 2}^{\log(x^{1/V})} \frac{\cos((2k/\log x)u)}{u} \, du + O(1)$$
$$= \log \min\{\log(x^{1/V}), \log(x^{1/(1+|k|)})\} + O(1)$$

Using this estimate, we get a lower bound for $\mathbb{E}\left(\int_{-1/2}^{1/2} \left|F\left(\frac{1}{2} + \frac{4Vq}{\log x} + it\right)\right|^2 dt\right)^q$ that is

$$\gg \frac{1}{\log^{q} x} \left(\frac{\log x}{V}\right)^{q^{2}} \sum_{|k| \le (\log x - 1)/2} \min\left\{\frac{\log x}{V}, \frac{\log x}{1 + |k|}\right\}^{q^{2} - q},$$

and (remembering that q_0 satisfies $q_0^2 - q_0 = 1$) this is

$$\gg \frac{1}{\log^{q} x} \left(\frac{\log x}{V}\right)^{q^{2} + \max\{1, q^{2} - q\}} \min\left\{\log\log x, \frac{1}{|q - q_{0}|}\right\}.$$

Again, inserting this in Proposition 4.4 produces the lower bound claimed in Theorem 1.2.

Acknowledgements

The author would like to thank the anonymous referee, for helpful comments and suggestions.

References

- [Bonami 1970] A. Bonami, "Étude des coefficients de Fourier des fonctions de $L^p(G)$ ", Ann. Inst. Fourier (Grenoble) 20:2 (1970), 335–402. MR Zbl
- [Bondarenko et al. 2018] A. Bondarenko, O. F. Brevig, E. Saksman, K. Seip, and J. Zhao, "Pseudomoments of the Riemann zeta function", *Bull. Lond. Math. Soc.* **50**:4 (2018), 709–724. MR Zbl
- [Chandee 2011] V. Chandee, "On the correlation of shifted values of the Riemann zeta function", *Q. J. Math.* **62**:3 (2011), 545–572. MR Zbl
- [Florea 2017] A. Florea, "The fourth moment of quadratic Dirichlet *L*-functions over function fields", *Geom. Funct. Anal.* **27**:3 (2017), 541–595. MR Zbl
- [Granville and Soundararajan 2001] A. Granville and K. Soundararajan, "Large character sums", J. Amer. Math. Soc. 14:2 (2001), 365–397. MR Zbl
- [Gut 2013] A. Gut, Probability: a graduate course, 2nd ed., Springer, 2013. MR Zbl
- [Halász 1983] G. Halász, "On random multiplicative functions", pp. 74–96 in *Hubert Delange colloquium* (Orsay, France, 1982), Publ. Math. Orsay **83**, Univ. Paris XI, 1983. MR Zbl
- [Harper 2017] A. J. Harper, "Moments of random multiplicative functions, I: Low moments, better than squareroot cancellation, and critical multiplicative chaos", preprint, 2017. To appear in *Forum of Mathematics, Pi*. arXiv
- [Harper et al. 2015] A. J. Harper, A. Nikeghbali, and M. Radziwiłł, "A note on Helson's conjecture on moments of random multiplicative functions", pp. 145–169 in *Analytic number theory*, edited by C. Pomerance and M. T. Rassias, Springer, 2015. MR Zbl
- [Heap and Lindqvist 2016] W. P. Heap and S. Lindqvist, "Moments of random multiplicative functions and truncated characteristic polynomials", *Q. J. Math.* **67**:4 (2016), 683–714. MR Zbl

[Helson 2010] H. Helson, "Hankel forms", Studia Math. 198:1 (2010), 79-84. MR Zbl

[Ivić 2003] A. Ivić, The Riemann zeta-function: Theory and applications, Dover, Mineola, NY, 2003. Zbl

- [Keating and Odgers 2008] J. P. Keating and B. E. Odgers, "Symmetry transitions in random matrix theory and *L*-functions", *Comm. Math. Phys.* **281**:2 (2008), 499–528. MR Zbl
- [Lau et al. 2013] Y.-K. Lau, G. Tenenbaum, and J. Wu, "On mean values of random multiplicative functions", *Proc. Amer. Math. Soc.* 141:2 (2013), 409–420. MR Zbl
- [Montgomery and Vaughan 2007] H. L. Montgomery and R. C. Vaughan, *Multiplicative number theory, I: Classical theory*, Cambridge Stud. Adv. Math. **97**, Cambridge Univ. Press, 2007. MR Zbl
- [Munsch 2017] M. Munsch, "Shifted moments of *L*-functions and moments of theta functions", *Mathematika* 63:1 (2017), 196–212. MR Zbl
- [Munsch and Shparlinski 2016] M. Munsch and I. E. Shparlinski, "Upper and lower bounds for higher moments of theta functions", *Q. J. Math.* **67**:1 (2016), 53–73. MR Zbl
- [Saksman and Seip 2016] E. Saksman and K. Seip, "Some open questions in analysis for Dirichlet series", pp. 179–191 in *Recent progress on operator theory and approximation in spaces of analytic functions*, edited by C. Bénéteau et al., Contemp. Math. **679**, Amer. Math. Soc., Providence, RI, 2016. MR Zbl
- [Shi and Weber 2016] S. Y. Shi and M. Weber, "On the equation $n_1 \cdots n_{k+1} = n_{k+2} \cdots n_{2(k+1)}$ and some variant with restricted unknowns", preprint, 2016. arXiv
- [Soundararajan and Young 2010] K. Soundararajan and M. P. Young, "The second moment of quadratic twists of modular *L*-functions", *J. Eur. Math. Soc.* **12**:5 (2010), 1097–1116. MR Zbl

[Wintner 1944] A. Wintner, "Random factorizations and Riemann's hypothesis", Duke Math. J. 11 (1944), 267–275. MR Zbl

Communicated by Roger Heath-Brown

Received 2018-04-27 Revised 2019-04-24 Accepted 2019-07-05

a.harper@warwick.ac.uk

University of Warwick, Coventry, United Kingdom

Algebra & Number Theory

msp.org/ant

EDITORS

MANAGING EDITOR

Bjorn Poonen Massachusetts Institute of Technology Cambridge, USA EDITORIAL BOARD CHAIR David Eisenbud University of California Berkeley, USA

BOARD OF EDITORS

Bhargav Bhatt	University of Michigan, USA	Raman Parimala	Emory University, USA
Richard E. Borcherds	University of California, Berkeley, USA	Jonathan Pila	University of Oxford, UK
Antoine Chambert-Loir	Université Paris-Diderot, France	Irena Peeva	Cornell University, USA
J-L. Colliot-Thélène	CNRS, Université Paris-Sud, France	Anand Pillay	University of Notre Dame, USA
Brian D. Conrad	Stanford University, USA	Michael Rapoport	Universität Bonn, Germany
Samit Dasgupta	Duke University, USA	Victor Reiner	University of Minnesota, USA
Hélène Esnault	Freie Universität Berlin, Germany	Peter Sarnak	Princeton University, USA
Gavril Farkas	Humboldt Universität zu Berlin, Germany	Joseph H. Silverman	Brown University, USA
Hubert Flenner	Ruhr-Universität, Germany	Michael Singer	North Carolina State University, USA
Sergey Fomin	University of Michigan, USA	Christopher Skinner	Princeton University, USA
Edward Frenkel	University of California, Berkeley, USA	Vasudevan Srinivas	Tata Inst. of Fund. Research, India
Wee Teck Gan	National University of Singapore	J. Toby Stafford	University of Michigan, USA
Andrew Granville	Université de Montréal, Canada	Shunsuke Takagi	University of Tokyo, Japan
Ben J. Green	University of Oxford, UK	Pham Huu Tiep	University of Arizona, USA
Joseph Gubeladze	San Francisco State University, USA	Ravi Vakil	Stanford University, USA
Christopher Hacon	University of Utah, USA	Michel van den Bergh	Hasselt University, Belgium
Roger Heath-Brown	Oxford University, UK	Akshay Venkatesh	Institute for Advanced Study, USA
János Kollár	Princeton University, USA	Marie-France Vignéras	Université Paris VII, France
Philippe Michel	École Polytechnique Fédérale de Lausanne	Kei-Ichi Watanabe	Nihon University, Japan
Susan Montgomery	University of Southern California, USA	Melanie Matchett Wood	University of California, Berkeley, USA
Shigefumi Mori	RIMS, Kyoto University, Japan	Shou-Wu Zhang	Princeton University, USA
Martin Olsson	University of California, Berkeley, USA		

PRODUCTION

production@msp.org

Silvio Levy, Scientific Editor

See inside back cover or msp.org/ant for submission instructions.

The subscription price for 2019 is US \$385/year for the electronic version, and \$590/year (+\$60, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to MSP.

Algebra & Number Theory (ISSN 1944-7833 electronic, 1937-0652 printed) at Mathematical Sciences Publishers, 798 Evans Hall #3840, c/o University of California, Berkeley, CA 94720-3840 is published continuously online. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices.

ANT peer review and production are managed by EditFLOW® from MSP.

PUBLISHED BY

mathematical sciences publishers

nonprofit scientific publishing

http://msp.org/ © 2019 Mathematical Sciences Publishers

Algebra & Number Theory

Volume 13 No. 10 2019

The elliptic KZB connection and algebraic de Rham theory for unipotent fundamental groups of elliptic curves MA LUO	2243
Moments of random multiplicative functions, II: High moments ADAM J HARPER	2277
Artin–Mazur–Milne duality for fppf cohomology CYRIL DEMARCHE and DAVID HARARI	2323
Betti numbers of Shimura curves and arithmetic three-orbifolds MIKOŁAJ FRĄCZYK and JEAN RAIMBAULT	2359
Combinatorial identities and Titchmarsh's divisor problem for multiplicative functions SARY DRAPPEAU and BERKE TOPACOGULLARI	2383
The construction problem for Hodge numbers modulo an integer MATTHIAS PAULSEN and STEFAN SCHREIEDER	2427

