

# ***Algebra & Number Theory***

Volume 13

2019

No. 10

**The construction problem for Hodge numbers  
modulo an integer**

Matthias Paulsen and Stefan Schreieder



# The construction problem for Hodge numbers modulo an integer

Matthias Paulsen and Stefan Schreieder

For any integer  $m \geq 2$  and any dimension  $n \geq 1$ , we show that any  $n$ -dimensional Hodge diamond with values in  $\mathbb{Z}/m\mathbb{Z}$  is attained by the Hodge numbers of an  $n$ -dimensional smooth complex projective variety. As a corollary, there are no polynomial relations among the Hodge numbers of  $n$ -dimensional smooth complex projective varieties besides the ones induced by the Hodge symmetries, which answers a question raised by Kollár in 2012.

## 1. Introduction

Hodge theory allows one to decompose the  $k$ -th Betti cohomology of an  $n$ -dimensional compact Kähler manifold  $X$  into its  $(p, q)$ -pieces for all  $0 \leq k \leq 2n$ :

$$H^k(X, \mathbb{C}) = \bigoplus_{\substack{p+q=k \\ 0 \leq p, q \leq n}} H^{p,q}(X), \quad \overline{H^{p,q}(X)} = H^{q,p}(X).$$

The  $\mathbb{C}$ -linear subspaces  $H^{p,q}(X)$  are naturally isomorphic to the Dolbeault cohomology groups  $H^q(X, \Omega_X^p)$ .

The integers  $h^{p,q}(X) = \dim_{\mathbb{C}} H^{p,q}(X)$  for  $0 \leq p, q \leq n$  are called Hodge numbers. One usually arranges them in the so called Hodge diamond:

$$\begin{array}{ccccccc}
 & & & h^{n,n} & & & \\
 & & & \vdots & & & \\
 & & h^{n,n-1} & & h^{n-1,n} & & \\
 & \ddots & & \vdots & & \ddots & \\
 & h^{n,1} & & \vdots & & h^{1,n} & \\
 h^{n,0} & & h^{n-1,1} & \dots & h^{1,n-1} & & h^{0,n} \\
 & h^{n-1,0} & & & & h^{0,n-1} & \\
 & \ddots & & \vdots & & \ddots & \\
 & & h^{1,0} & & h^{0,1} & & \\
 & & h^{0,0} & & & & 
 \end{array}$$

The sum of the  $k$ -th row of the Hodge diamond equals the  $k$ -th Betti number. We always assume that a Kähler manifold is compact and connected, so we have  $h^{0,0} = h^{n,n} = 1$ .

*MSC2010:* primary 32Q15; secondary 14C30, 14E99, 51M15.

*Keywords:* Hodge numbers, Kähler manifolds, construction problem.

Complex conjugation and Serre duality induce the symmetries

$$h^{p,q} = h^{q,p} = h^{n-p,n-q} \quad \text{for all } 0 \leq p, q \leq n. \quad (1)$$

Additionally, we have the Lefschetz inequalities

$$h^{p,q} \leq h^{p+1,q+1} \quad \text{for } p+q < n. \quad (2)$$

While Hodge theory places severe restrictions on the geometry and topology of Kähler manifolds, Simpson [2004] points out that very little is known to which extent the theoretically possible phenomena actually occur. This leads to the following construction problem for Hodge numbers:

**Question 1.** Let  $(h^{p,q})_{0 \leq p,q \leq n}$  be a collection of nonnegative integers with  $h^{0,0} = 1$  obeying the Hodge symmetries (1) and the Lefschetz inequalities (2). Does there exist a Kähler manifold  $X$  such that  $h^{p,q}(X) = h^{p,q}$  for all  $0 \leq p, q \leq n$ ?

After results in dimensions two and three (see e.g., [Hunt 1989]), significant progress has been made by Schreieder [2015]. For instance, it is shown in [loc. cit., Theorem 3] that the above construction problem is fully solvable for large parts of the Hodge diamond in arbitrary dimensions. In particular, the Hodge numbers in a given weight  $k$  may be arbitrary (up to a quadratic lower bound on  $h^{p,p}$  if  $k = 2p$  is even) and so the outer Hodge numbers can be far larger than the inner Hodge numbers (see [loc. cit., Theorem 1]), contradicting earlier expectations formulated in [Simpson 2004]. Weaker results with simpler proofs, concerning the possible Hodge numbers in a given weight, have later been obtained by Arapura [2016].

In [Schreieder 2015], it was also observed that one cannot expect a positive answer to Question 1 in its entirety. For example, any 3-dimensional Kähler manifold  $X$  with  $h^{1,1}(X) = 1$  and  $h^{2,0}(X) \geq 1$  satisfies  $h^{2,1}(X) < 12^6 \cdot h^{3,0}(X)$ , see [loc. cit., Proposition 28]. Therefore, a complete classification of all possible Hodge diamonds of Kähler manifolds or smooth complex projective varieties seems hopelessly complicated.

While these inequalities aggravate the construction problem for Hodge numbers, one might ask whether there also exist number theoretic obstructions for possible Hodge diamonds. For example, the Chern numbers of Kähler manifolds satisfy certain congruences due to integrality conditions implied by the Hirzebruch–Riemann–Roch theorem.

For an arbitrary integer  $m \geq 2$ , let us consider the Hodge numbers of a Kähler manifold in  $\mathbb{Z}/m\mathbb{Z}$ , which forces all inequalities to disappear. The purpose of this paper is to show that Question 1 is modulo  $m$  completely solvable even for smooth complex projective varieties.

**Theorem 2.** *Let  $m \geq 2$  be an integer. For any integer  $n \geq 1$  and any collection of integers  $(h^{p,q})_{0 \leq p,q \leq n}$  such that  $h^{0,0} = 1$  and  $h^{p,q} = h^{q,p} = h^{n-p,n-q}$  for  $0 \leq p, q \leq n$ , there exists a smooth complex projective variety  $X$  of dimension  $n$  such that*

$$h^{p,q}(X) \equiv h^{p,q} \pmod{m}$$

for all  $0 \leq p, q \leq n$ .

Therefore, the Hodge numbers of Kähler manifolds do not follow any number theoretic rules, and the behavior of smooth complex projective varieties is the same in this aspect.

As a consequence of [Theorem 2](#), we show:

**Corollary 3.** *Up to the Hodge symmetries (1), there are no polynomial relations among the Hodge numbers of smooth complex projective varieties of the same dimension.*

In particular, there are no polynomial relations in the strictly larger class of Kähler manifolds, which was a question raised by Kollár after a colloquium talk of Kotschick at the University of Utah in fall 2012. For linear relations among Hodge numbers, this question was settled in work of Kotschick and Schreieder [\[2013\]](#).

We call the Hodge numbers  $h^{p,q}(X)$  with  $p \in \{0, n\}$  or  $q \in \{0, n\}$  (i.e., the ones placed on the border of the Hodge diamond) the *outer Hodge numbers* of  $X$  and the remaining ones the *inner Hodge numbers*. Note that the outer Hodge numbers are birational invariants and are thus determined by the birational equivalence class of  $X$ .

Our proof shows (see [Theorem 5](#) below) that any smooth complex projective variety is birational to a smooth complex projective variety with prescribed inner Hodge numbers in  $\mathbb{Z}/m\mathbb{Z}$ . As a corollary, there are no polynomial relations among the inner Hodge numbers within a given birational equivalence class. This is again a generalization of the corresponding result for linear relations obtained in [\[Kotschick and Schreieder 2013, Theorem 2\]](#).

The proof of [Theorem 2](#) can thus be divided into two steps: First we solve the construction problem modulo  $m$  for the outer Hodge numbers. This is done in [Section 2](#). Then we show the aforementioned result that the inner Hodge numbers can be adjusted arbitrarily in  $\mathbb{Z}/m\mathbb{Z}$  via birational equivalences (in fact, via repeated blow-ups). This is done in [Section 3](#). Finally, in [Section 4](#) we deduce that no nontrivial polynomial relations between Hodge numbers exist, thus answering Kollár's question.

## 2. Outer Hodge numbers

We prove the following statement via induction on the dimension  $n \geq 1$ .

**Proposition 4.** *For any collection of integers  $(h^{p,0})_{1 \leq p \leq n}$ , there exists a smooth complex projective variety  $X_n$  of dimension  $n$  together with a very ample line bundle  $L_n$  on  $X_n$  such that*

$$h^{p,0}(X_n) \equiv h^{p,0} \pmod{m}$$

*for all  $1 \leq p \leq n$  and*

$$\chi(L_n^{-1}) \equiv 1 \pmod{m}.$$

*Proof.* We take  $X_1$  to be a curve of genus  $g$  where  $g \equiv h^{1,0} \pmod{m}$ . Further, we take  $L_1$  to be a line bundle of degree  $d$  on  $X_1$  where  $d > 2g$  and  $d \equiv -g \pmod{m}$ . Then  $L_1$  is very ample and by the Riemann–Roch theorem we have  $\chi(L_1^{-1}) \equiv 1 \pmod{m}$ .

Now let  $n > 1$ . We define a collection of integers  $(k^{p,0})_{-1 \leq p \leq n-1}$  recursively via

$$k^{-1,0} = 0, \quad k^{0,0} = 1, \quad k^{p,0} = h^{p,0} - 2k^{p-1,0} - k^{p-2,0} \text{ for } 1 \leq p \leq n-1.$$

We choose  $X_{n-1}$  and  $L_{n-1}$  by induction hypothesis such that  $h^{p,0}(X_{n-1}) \equiv k^{p,0} \pmod{m}$  for all  $1 \leq p \leq n-1$ .

Let  $E$  be a smooth elliptic curve and let  $L$  be a very ample line bundle of degree  $d$  on  $E$  such that  $d \equiv 1 \pmod{m}$ . Let  $e$  be a positive integer such that

$$e \equiv 1 + \sum_{p=1}^n (-1)^p h^{p,0} \pmod{m}.$$

Let  $X_n \subset X_{n-1} \times E \times E$  be a hypersurface defined by a general section of the very ample line bundle

$$P_n = \mathrm{pr}_1^* L_{n-1} \otimes \mathrm{pr}_2^* L^{m-1} \otimes \mathrm{pr}_3^* L^e$$

on  $X_{n-1} \times E \times E$ . By Bertini's theorem, we may assume  $X_n$  to be smooth and irreducible. Let  $L_n$  be the restriction to  $X_n$  of the very ample line bundle

$$Q_n = \mathrm{pr}_1^* L_{n-1} \otimes \mathrm{pr}_2^* L \otimes \mathrm{pr}_3^* L$$

on  $X_{n-1} \times E \times E$ . Then  $L_n$  is again very ample.

By the Lefschetz hyperplane theorem, we have

$$h^{p,0}(X_n) = h^{p,0}(X_{n-1} \times E \times E)$$

for all  $1 \leq p \leq n-1$ . Since the Hodge diamond of  $E \times E$  is

$$\begin{array}{ccccc} & & 1 & & \\ & 2 & & 2 & \\ 1 & & 4 & & 1, \\ & 2 & & 2 & \\ & & 1 & & \end{array}$$

Künneth's formula yields

$$h^{p,0}(X_n) = h^{p,0}(X_{n-1}) + 2h^{p-1,0}(X_{n-1}) + h^{p-2,0}(X_{n-1}) \equiv k^{p,0} + 2k^{p-1,0} + k^{p-2,0} = h^{p,0} \pmod{m}$$

for all  $1 \leq p \leq n-1$ . Therefore, it only remains to show that  $h^{n,0}(X_n) \equiv h^{n,0} \pmod{m}$  and  $\chi(L_n^{-1}) \equiv 1 \pmod{m}$ . Since

$$\chi(\mathcal{O}_{X_n}) = 1 + \sum_{p=1}^n (-1)^p h^{p,0}(X_n),$$

the congruence  $h^{n,0}(X_n) \equiv h^{n,0} \pmod{m}$  is equivalent to  $\chi(\mathcal{O}_{X_n}) \equiv e \pmod{m}$ .

By definition of  $X_n$ , the ideal sheaf on  $X_{n-1} \times E \times E$  of regular functions vanishing on  $X_n$  is isomorphic to the sheaf of sections of the dual line bundle  $P_n^{-1}$ . Hence, there is a short exact sequence

$$0 \rightarrow P_n^{-1} \rightarrow \mathcal{O}_{X_{n-1} \times E \times E} \rightarrow i_* \mathcal{O}_{X_n} \rightarrow 0 \quad (3)$$

of sheaves on  $X_{n-1} \times E \times E$  where  $i: X_n \rightarrow X_{n-1} \times E \times E$  denotes the inclusion. Together with Künneth's formula and the Riemann–Roch theorem, we obtain

$$\chi(\mathcal{O}_{X_n}) = \chi(\mathcal{O}_{X_{n-1} \times E \times E}) - \chi(P_n^{-1}) = \chi(\mathcal{O}_{X_{n-1}}) \underbrace{\chi(\mathcal{O}_E)^2}_{=0} - \underbrace{\chi(L_{n-1}^{-1})}_{\equiv 1} \underbrace{\chi(L^{1-m})}_{\equiv 1} \underbrace{\chi(L^{-e})}_{\equiv -e} \equiv e \pmod{m}.$$

Tensoring (3) with  $Q_n^{-1}$  yields the short exact sequence

$$0 \rightarrow P_n^{-1} \otimes Q_n^{-1} \rightarrow Q_n^{-1} \rightarrow i_* i^* Q_n^{-1} \rightarrow 0$$

and thus

$$\chi(L_n^{-1}) = \chi(Q_n^{-1}) - \chi(P_n^{-1} \otimes Q_n^{-1}) = \underbrace{\chi(L_{n-1}^{-1})}_{\equiv 1} \underbrace{\chi(L^{-1})^2}_{\equiv 1} - \chi(L_{n-1}^{-2}) \underbrace{\chi(L^{-m})}_{\equiv 0} \chi(L^{-e-1}) \equiv 1 \pmod{m}.$$

This finishes the induction step. □

### 3. Inner Hodge numbers

We now show the following result, which significantly improves [Kotschick and Schreieder 2013, Theorem 2].

**Theorem 5.** *Let  $X$  be a smooth complex projective variety of dimension  $n$  and let  $(h^{p,q})_{1 \leq p,q \leq n-1}$  be any collection of integers such that  $h^{p,q} = h^{q,p} = h^{n-p,n-q}$  for  $1 \leq p, q \leq n-1$ . Then  $X$  is birational to a smooth complex projective variety  $X'$  such that*

$$h^{p,q}(X') \equiv h^{p,q} \pmod{m}$$

for all  $1 \leq p, q \leq n-1$ .

Together with Proposition 4, this will complete the proof of Theorem 2.

Let us recall the following result on blow-ups, see e.g., [Voisin 2002, Theorem 7.31]: If  $\tilde{X}$  denotes the blow-up of a Kähler manifold  $X$  along a closed submanifold  $Z \subset X$  of codimension  $c$ , we have

$$H^{p,q}(\tilde{X}) \cong H^{p,q}(X) \oplus \bigoplus_{i=1}^{c-1} H^{p-i,q-i}(Z).$$

Therefore,

$$h^{p,q}(\tilde{X}) = h^{p,q}(X) + \sum_{i=1}^{c-1} h^{p-i,q-i}(Z). \quad (4)$$

In order to prove Theorem 5, we first show that we may assume that  $X$  contains certain subvarieties, without modifying its Hodge numbers modulo  $m$ .

**Lemma 6.** *Let  $X$  be a smooth complex projective variety of dimension  $n$ . Let  $r, s \geq 0$  be integers such that  $r + s \leq n - 1$ . Then  $X$  is birational to a smooth complex projective variety  $X'$  of dimension  $n$  such that  $h^{p,q}(X') \equiv h^{p,q}(X) \pmod{m}$  for all  $0 \leq p, q \leq n$  and such that  $X'$  contains at least  $m$  disjoint smooth closed subvarieties that are all isomorphic to a projective bundle of rank  $r$  over  $\mathbb{P}^s$ .*

*Proof.* We first blow up  $X$  in a point and denote the result by  $\tilde{X}$ . The exceptional divisor is a subvariety in  $\tilde{X}$  isomorphic to  $\mathbb{P}^{n-1}$ . In particular,  $\tilde{X}$  contains a copy of  $\mathbb{P}^s \subset \mathbb{P}^{n-1}$ . Now we blow up  $\tilde{X}$  along  $\mathbb{P}^s$  to obtain  $\hat{X}$ . The exceptional divisor in  $\hat{X}$  is the projectivization of the normal bundle of  $\mathbb{P}^s$  in  $\tilde{X}$ . Since  $\mathbb{P}^s$  is contained in a smooth closed subvariety of dimension  $r + s + 1$  in  $\tilde{X}$  (choose either  $\mathbb{P}^{r+s+1} \subset \mathbb{P}^{n-1}$  if  $r + s < n - 1$  or  $\tilde{X}$  if  $r + s = n - 1$ ), the normal bundle of  $\mathbb{P}^s$  in  $\tilde{X}$  contains a vector subbundle of rank  $r + 1$ , and hence its projectivization contains a projective subbundle of rank  $r$ . Therefore,  $\hat{X}$  admits a subvariety isomorphic to the total space of a projective bundle of rank  $r$  over  $\mathbb{P}^s$ .

By (4), the above construction only has an additive effect on the Hodge diamond, i.e., the differences between respective Hodge numbers of  $\hat{X}$  and  $X$  are constants independent of  $X$ . Hence, we may apply the above construction  $m - 1$  more times to obtain a smooth complex projective variety  $X'$  containing  $m$  disjoint copies of the desired projective bundle and satisfying  $h^{p,q}(X') \equiv h^{p,q}(X) \pmod{m}$ .  $\square$

In the following, we consider the primitive Hodge numbers

$$l^{p,q}(X) = h^{p,q}(X) - h^{p-1,q-1}(X)$$

for  $p + q \leq n$ . Clearly, it suffices to show [Theorem 5](#) for a given collection  $(l^{p,q})_{(p,q) \in I}$  of primitive Hodge numbers instead, where

$$I = \{(p, q) \mid 1 \leq p \leq q \leq n - 1 \text{ and } p + q \leq n\}.$$

This is because one can get back the original Hodge numbers from the primitive Hodge numbers via the relation

$$h^{p,q}(X) = h^{0,q-p}(X) + \sum_{i=1}^p l^{i,q-p+i}(X)$$

for  $p \leq q$  and  $p + q \leq n$ , and  $h^{0,q-p}(X)$  is a birational invariant.

We define a total order  $<$  on  $I$  via

$$(r, s) < (p, q) \iff r + s < p + q \text{ or } (r + s = p + q \text{ and } s < q).$$

**Proposition 7.** *Let  $X$  be a smooth complex projective variety of dimension  $n$ . Let  $(r, s) \in I$ . Then  $X$  is birational to a smooth complex projective variety  $X'$  of dimension  $n$  such that*

$$l^{r,s}(X') \equiv l^{r,s}(X) + 1 \pmod{m} \quad \text{and} \quad l^{p,q}(X') \equiv l^{p,q}(X) \pmod{m}$$

for all  $(p, q) \in I$  with  $(r, s) < (p, q)$ .

*Proof.* By [Lemma 6](#), we may assume that  $X$  contains  $m$  disjoint copies of a projective bundle of rank  $r - 1$  over  $\mathbb{P}^{s-r+1}$ . Therefore, it is possible to blow up  $X$  along a projective bundle  $B_d$  of rank  $r - 1$  over



smooth hypersurfaces  $Y_d \subset \mathbb{P}^{s-r+1}$  of degree  $d$  (in case of  $r = s$ ,  $Y_d$  just consists of  $d$  distinct points in  $\mathbb{P}^1$ ) and we may repeat this procedure  $m$  times and with different values for  $d$ . The Hodge numbers of  $B_d$  are the same as for the trivial bundle  $Y_d \times \mathbb{P}^{r-1}$ , see e.g., [Voisin 2002, Lemma 7.32].

By the Lefschetz hyperplane theorem, the Hodge diamond of  $Y_d$  is the sum of the Hodge diamond of  $Y_1 \cong \mathbb{P}^{s-r}$ , having nonzero entries only in the middle column, and of a Hodge diamond depending on  $d$ , having nonzero entries only in the middle row. It is well known (e.g., by computing Euler characteristics as in Section 2) that the two outer entries of this middle row are precisely  $\binom{d-1}{s-r+1}$ .

Now we blow up  $X$  once along  $B_{s-r+2}$  and  $m-1$  times along  $B_1$  and denote the resulting smooth complex projective variety by  $X'$ . Due to (4) and Künneth's formula, this construction affects the Hodge numbers modulo  $m$  in the same way as if we would blow up a single subvariety  $Z \times \mathbb{P}^{r-1} \subset X$ , where  $Z$  is a (formal)  $(s-r)$ -dimensional Kähler manifold whose Hodge diamond is concentrated in the middle row and has outer entries equal to  $\binom{s-r+2-1}{s-r+1} = 1$ . In particular, we have  $h^{p,q}(Z \times \mathbb{P}^{r-1}) = 0$  unless  $s-r \leq p+q \leq s+r-2$  (and  $p+q$  has the same parity as  $s-r$ ) and  $|p-q| \leq s-r$ . On the other hand,  $h^{p,q}(Z \times \mathbb{P}^{r-1}) = 1$  if  $s-r \leq p+q \leq s+r-2$  and  $|p-q| = s-r$ .

Taking differences in (4), it follows that

$$l^{p,q}(X') \equiv l^{p,q}(X) + h^{p-1,q-1}(Z \times \mathbb{P}^{r-1}) - h^{p-n+s-1,q-n+s-1}(Z \times \mathbb{P}^{r-1}) \pmod{m}$$

for all  $p+q \leq n$ . But we have

$$(p-n+s-1) + (q-n+s-1) = p+q-2n+2s-2 \leq 2s-n-2 \leq s-r-2$$

and hence  $h^{p-n+s-1,q-n+s-1}(Z \times \mathbb{P}^{r-1}) = 0$  for all  $(p,q) \in I$  by the above remark.

Further,

$$l^{r,s}(X') \equiv l^{r,s}(X) + h^{r-1,s-1}(Z \times \mathbb{P}^{r-1}) = l^{r,s}(X) + 1 \pmod{m}$$

since  $s-r \leq (r-1) + (s-1) \leq s+r-2$  and  $|r-s| = s-r$ .

Finally,  $r+s < p+q$  implies  $(p-1) + (q-1) > s+r-2$ , while  $r+s = p+q$  and  $s < q$  imply  $|p-q| > s-r$ , so we have  $h^{p-1,q-1}(Z \times \mathbb{P}^{r-1}) = 0$  in both cases and thus

$$l^{p,q}(X') \equiv l^{p,q}(X) + h^{p-1,q-1}(Z \times \mathbb{P}^{r-1}) = l^{p,q}(X) \pmod{m}$$

for all  $(p,q) \in I$  with  $(r,s) < (p,q)$ . □

*Proof of Theorem 5.* The statement is an immediate consequence of applying Proposition 7 inductively  $t_{p,q}$  times to each  $(p,q) \in I$  in the descending order induced by  $\prec$ , where  $t_{p,q} \equiv l^{p,q} - l^{p,q}(X_{p,q}) \pmod{m}$  and  $X_{p,q}$  is the variety obtained in the previous step. □

#### 4. Polynomial relations

The following principle seems to be classical.

**Lemma 8.** *Let  $N \geq 1$  and  $S \subset \mathbb{Z}^N$  be a subset such that its reduction modulo  $m$  is the whole of  $(\mathbb{Z}/m\mathbb{Z})^N$  for infinitely many integers  $m \geq 2$ . If  $f \in \mathbb{C}[x_1, \dots, x_N]$  is a polynomial vanishing on  $S$ , then  $f = 0$ .*



*Proof.* Let  $f \in \mathbb{C}[x_1, \dots, x_N]$  be a nonzero polynomial vanishing on  $S$ . By choosing a  $\mathbb{Q}$ -basis of  $\mathbb{C}$  and a  $\mathbb{Q}$ -linear projection  $\mathbb{C} \rightarrow \mathbb{Q}$  which sends a nonzero coefficient of  $f$  to 1, we see that we may assume that the coefficients of  $f$  are rational, hence even integral. Since  $f \neq 0$ , there exists a point  $z \in \mathbb{Z}^N$  such that  $f(z) \neq 0$ . Choose an integer  $m \geq 2$  from the assumption which does not divide  $f(z)$ . Then  $f(z) \not\equiv 0 \pmod{m}$ . However, we have  $z \equiv s \pmod{m}$  for some  $s \in S$  and thus  $f(z) \equiv f(s) = 0 \pmod{m}$ , because  $f \in \mathbb{Z}[x_1, \dots, x_N]$ . This is a contradiction.  $\square$

*Proof of Corollary 3.* This follows by applying Lemma 8 to the set  $S$  of possible Hodge diamonds, where we consider only a nonredundant quarter of the diamond to take the Hodge symmetries into account. Theorem 2 guarantees that the reductions of  $S$  modulo  $m$  are surjective even for all integers  $m \geq 2$ .  $\square$

In the same way Theorem 2 implies Corollary 3, Theorem 5 yields the following.

**Corollary 9.** *There are no nontrivial polynomial relations among the inner Hodge numbers of all smooth complex projective varieties in any given birational equivalence class.*

### Acknowledgements

The second named author thanks J. Kollár and D. Kotschick for independently making him aware of Kollár’s question (answered in Corollary 3 above), which was the starting point of this paper. The authors are grateful to the referees for useful suggestions. This work is supported by the DFG project “Topologische Eigenschaften von algebraischen Varietäten” (project no. 416054549).

### References

- [Arapura 2016] D. Arapura, “Geometric Hodge structures with prescribed Hodge numbers”, pp. 414–421 in *Recent advances in Hodge theory*, edited by M. Kerr and G. Pearlstein, Lond. Math. Soc. Lecture Note Ser. **427**, Cambridge Univ. Press, 2016. [MR](#) [Zbl](#)
- [Hunt 1989] B. Hunt, “Complex manifold geography in dimension 2 and 3”, *J. Differential Geom.* **30**:1 (1989), 51–153. [MR](#) [Zbl](#)
- [Kotschick and Schreieder 2013] D. Kotschick and S. Schreieder, “The Hodge ring of Kähler manifolds”, *Compos. Math.* **149**:4 (2013), 637–657. [MR](#) [Zbl](#)
- [Schreieder 2015] S. Schreieder, “On the construction problem for Hodge numbers”, *Geom. Topol.* **19**:1 (2015), 295–342. [MR](#) [Zbl](#)
- [Simpson 2004] C. Simpson, “The construction problem in Kähler geometry”, pp. 365–402 in *Different faces of geometry*, edited by S. Donaldson et al., Int. Math. Ser. **3**, Springer, 2004. [MR](#) [Zbl](#)
- [Voisin 2002] C. Voisin, *Hodge theory and complex algebraic geometry, I*, Cambridge Stud. Adv. Math. **76**, Cambridge Univ. Press, 2002. [MR](#) [Zbl](#)

Communicated by János Kollár

Received 2019-03-13      Revised 2019-06-13      Accepted 2019-07-29

[paulsen@math.lmu.de](mailto:paulsen@math.lmu.de)

Mathematisches Institut, LMU München, Germany

[schreieder@math.lmu.de](mailto:schreieder@math.lmu.de)

Mathematisches Institut, LMU München, Germany

# Algebra & Number Theory

[msp.org/ant](http://msp.org/ant)

## EDITORS

### MANAGING EDITOR

Bjorn Poonen  
Massachusetts Institute of Technology  
Cambridge, USA

### EDITORIAL BOARD CHAIR

David Eisenbud  
University of California  
Berkeley, USA

## BOARD OF EDITORS

Bhargav Bhatt	University of Michigan, USA	Raman Parimala	Emory University, USA
Richard E. Borcherds	University of California, Berkeley, USA	Jonathan Pila	University of Oxford, UK
Antoine Chambert-Loir	Université Paris-Diderot, France	Irena Peeva	Cornell University, USA
J-L. Colliot-Thélène	CNRS, Université Paris-Sud, France	Anand Pillay	University of Notre Dame, USA
Brian D. Conrad	Stanford University, USA	Michael Rapoport	Universität Bonn, Germany
Samit Dasgupta	Duke University, USA	Victor Reiner	University of Minnesota, USA
Hélène Esnault	Freie Universität Berlin, Germany	Peter Sarnak	Princeton University, USA
Gavril Farkas	Humboldt Universität zu Berlin, Germany	Joseph H. Silverman	Brown University, USA
Hubert Flenner	Ruhr-Universität, Germany	Michael Singer	North Carolina State University, USA
Sergey Fomin	University of Michigan, USA	Christopher Skinner	Princeton University, USA
Edward Frenkel	University of California, Berkeley, USA	Vasudevan Srinivas	Tata Inst. of Fund. Research, India
Wee Teck Gan	National University of Singapore	J. Toby Stafford	University of Michigan, USA
Andrew Granville	Université de Montréal, Canada	Shunsuke Takagi	University of Tokyo, Japan
Ben J. Green	University of Oxford, UK	Pham Huu Tiep	University of Arizona, USA
Joseph Gubeladze	San Francisco State University, USA	Ravi Vakil	Stanford University, USA
Christopher Hacon	University of Utah, USA	Michel van den Bergh	Hasselt University, Belgium
Roger Heath-Brown	Oxford University, UK	Akshay Venkatesh	Institute for Advanced Study, USA
János Kollár	Princeton University, USA	Marie-France Vignéras	Université Paris VII, France
Philippe Michel	École Polytechnique Fédérale de Lausanne	Kei-Ichi Watanabe	Nihon University, Japan
Susan Montgomery	University of Southern California, USA	Melanie Matchett Wood	University of California, Berkeley, USA
Shigefumi Mori	RIMS, Kyoto University, Japan	Shou-Wu Zhang	Princeton University, USA
Martin Olsson	University of California, Berkeley, USA		

## PRODUCTION

[production@msp.org](mailto:production@msp.org)

Silvio Levy, Scientific Editor

---

See inside back cover or [msp.org/ant](http://msp.org/ant) for submission instructions.

---

The subscription price for 2019 is US \$385/year for the electronic version, and \$590/year (+\$60, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to MSP.


---

Algebra & Number Theory (ISSN 1944-7833 electronic, 1937-0652 printed) at Mathematical Sciences Publishers, 798 Evans Hall #3840, c/o University of California, Berkeley, CA 94720-3840 is published continuously online. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices.

---

ANT peer review and production are managed by EditFlow® from MSP.

PUBLISHED BY

 **mathematical sciences publishers**  
nonprofit scientific publishing

<http://msp.org/>

© 2019 Mathematical Sciences Publishers

# Algebra & Number Theory

Volume 13

No. 10

2019

---

<a href="#">The elliptic KZB connection and algebraic de Rham theory for unipotent fundamental groups of elliptic curves</a>	2243
MA LUO	
<a href="#">Moments of random multiplicative functions, II: High moments</a>	2277
ADAM J HARPER	
<a href="#">Artin–Mazur–Milne duality for fppf cohomology</a>	2323
CYRIL DEMARCHE and DAVID HARARI	
<a href="#">Betti numbers of Shimura curves and arithmetic three-orbifolds</a>	2359
MIKOŁAJ FRĄCZYK and JEAN RAIMBAULT	
<a href="#">Combinatorial identities and Titchmarsh’s divisor problem for multiplicative functions</a>	2383
SARY DRAPPEAU and BERKE TOPACOGULLARI	
<a href="#">The construction problem for Hodge numbers modulo an integer</a>	2427
MATTHIAS PAULSEN and STEFAN SCHREIEDER	