

Algebra & Number Theory

Volume 13

2019

No. 5

**Contragredient representations over local fields
of positive characteristic**

Wen-Wei Li



Contragredient representations over local fields of positive characteristic

Wen-Wei Li

It was conjectured by Adams, Vogan and Prasad that under the local Langlands correspondence, the L -parameter of the contragredient representation equals that of the original representation composed with the Chevalley involution of the L -group. We verify a variant of their prediction for all connected reductive groups over local fields of positive characteristic, in terms of the local Langlands parametrization of A. Genestier and V. Lafforgue. We deduce this from a global result for cuspidal automorphic representations over function fields, which is in turn based on a description of the transposes of Lafforgue's excursion operators.

1. Introduction	1197
2. Review of representation theory	1203
3. Statement of a variant of the conjecture	1210
4. Overview of the global Langlands parametrization	1219
5. The transposes of excursion operators	1231
Acknowledgements	1240
References	1240

1. Introduction

Let F be a local field. Choose a separable closure $\bar{F}|F$ and let W_F be the Weil group of F . For a connected reductive F -group G , the *local Langlands conjecture* asserts the existence of a map

$$\text{LLC} : \Pi(G) \rightarrow \Phi(G)$$

where $\Pi(G)$ is the set of isomorphism classes of irreducible smooth representations π of $G(F)$ (or Harish-Chandra modules when F is archimedean), and $\Phi(G)$ is the set of $\hat{G}$ -conjugacy classes of L -parameters $W_F \xrightarrow{\phi} {}^L G$. Here the representations and the L -groups are taken over $\mathbb{C}$, but we will soon switch to the setting of nonarchimedean F and ℓ -adic coefficients.

It is expected that the L -packets $\Pi_\phi := \text{LLC}^{-1}(\phi)$ are finite sets; if $\pi \in \Pi_\phi$, we say ϕ is the L -parameter of π . The local Langlands correspondence also predicates on the internal structure of Π_ϕ when ϕ is a *tempered* parameter; this requires additional structures as follows:

MSC2010: primary 11F70; secondary 11R58, 22E55.

Keywords: contragredient representation, function field, local Langlands conjecture.

- When G is quasisplit, choose a Whittaker datum $\mathfrak{w} = (U, \chi)$ of G , taken up to $G(F)$ -conjugacy, where $U \subset G$ is a maximal unipotent subgroup and χ is a generic smooth character of $U(F)$. The individual members of Π_ϕ are described in terms of

$$S_\phi := Z_{\hat{G}}(\text{im}(\phi)), \quad \mathcal{S}_\phi := \pi_0(S_\phi).$$

Specifically, to each $\pi \in \Pi_\phi$ one should be able to attach an irreducible representation ρ of the finite group $\mathcal{S}_\phi$ (up to isomorphism), such that a $\mathfrak{w}$ -generic $\pi \in \Pi_\phi$ maps to $\rho = \mathbf{1}$.

- For nonsplit G , one needs to connect G to a quasisplit group by means of a pure inner twist, or more generally a rigid inner twist [Kaletha 2016b]; in parallel, the L -packets will extend across various inner forms of G . We refer to [loc. cit., §5.4] for a discussion in this generality.

One natural question is to describe various operations on $\Pi(G)$ in terms of L -parameters. Among them, we consider the *contragredient* $\check{\pi}$ of π . The question is thus:

$$\text{How is } \pi \mapsto \check{\pi} \text{ in } \Pi(G) \text{ reflected on } \Phi(G)?$$

Despite its immediate appearance, this question has not been considered in this generality until the independent work of Adams and Vogan [2016, Conjecture 1.1] and D. Prasad [2018, §4]. The answer hinges on the *Chevalley involution* ${}^L\theta$ on ${}^L G$ to be reviewed in Section 3.1.

Conjecture 1.1 (Adams and Vogan; Prasad). Let π be an irreducible smooth representation of $G(F)$.

- (1) If π has L -parameter ϕ , then $\check{\pi}$ has L -parameter ${}^L\theta \circ \phi$.
- (2) Assume for simplicity that G is quasisplit and fix a Whittaker datum $\mathfrak{w}$. If a tempered representation $\pi \in \Pi_\phi$ corresponds to an irreducible representation ρ of $\mathcal{S}_\phi$, then $\check{\pi}$ corresponds to $(\rho \circ {}^L\theta)^\vee$ tensored with a character ξ of $\mathcal{S}_\phi$.

To define ξ , we use the general recipe [Kaletha 2013, Lemma 4.1]:

$$\begin{array}{ccc} \pi_0(S_\phi / Z_{\hat{G}}^{\text{Gal}(\bar{F}|F)}) & \longrightarrow & \ker[H^1(W_F, Z_{\hat{G}^{\text{sc}}}) \rightarrow H^1(W_F, Z_{\hat{G}})] \\ \uparrow & & \downarrow \\ \mathcal{S}_\phi & & \left(\frac{G^{\text{ad}}(F)}{\text{im}[G(F) \rightarrow G^{\text{ad}}(F)]} \right)^{\text{Pontryagin dual}} \end{array}$$

Let B be the Borel subgroup of G included in the Whittaker datum, and choose a maximal torus $T \subset B$. Take the $\kappa \in T^{\text{ad}}(F)$ acting as -1 on each $\mathfrak{g}_\alpha$ where α is any B -simple root. This furnishes the character ξ of $\mathcal{S}_\phi$. When G is not quasisplit, we have to endow it with a pure or rigid inner twist alluded to above.

Conjecture 1.1 comprises two layers: the second one is due to Prasad [2018]. In this article, we will focus exclusively on the first layer.

A precondition of the Adams–Vogan–Prasad conjecture is the existence of a map $\Pi(G) \rightarrow \Phi(G)$, baptized the *Langlands parametrization*, which has been constructed for many groups in various ways:

- When F is archimedean, the local Langlands correspondence is Langlands’ paraphrase of Harish-Chandra’s works. The “first layer” of the Adams–Vogan–Prasad conjecture is established by Adams and Vogan [2016], and Kaletha [2013, Theorem 5.9] obtained the necessary refinement for the “second layer”.
- When F is nonarchimedean of characteristic zero and G is a symplectic or quasisplit orthogonal group, Kaletha [2013, Theorem 5.9, Corollary 5.10] verified the Adams–Vogan–Prasad conjecture in terms of Arthur’s *endoscopic classification* of representations, which offers the local Langlands correspondence for these groups.
- For nonarchimedean F and general G , Kaletha [2013, §6] also verified the conjecture for the depth-zero and epipelagic supercuspidal L -packets, constructed by DeBacker, Reeder and Kaletha using induction from open compact subgroups.

The aim of this article is to address the first layer of Conjecture 1.1 when F is a nonarchimedean local field of characteristic $p > 0$ and G is arbitrary, in terms of the Langlands parametrization $\Pi(G) \rightarrow \Phi(G)$ of A. Genestier and V. Lafforgue [2017]. Their method is based on the geometry of the moduli stack of *restricted chtoucas*, intimately related to the global Langlands parametrization of cuspidal automorphic representations by Lafforgue [2018]. Accordingly, our representations π will be realized on $\overline{\mathbb{Q}}_\ell$ -vector spaces, where ℓ is a prime number not equal to p , and the L -group ${}^L G$ is viewed as a $\overline{\mathbb{Q}}_\ell$ -group. As $\mathbb{C} \simeq \overline{\mathbb{Q}}_\ell$ as abstract fields, passing to $\overline{\mathbb{Q}}_\ell$ does not alter the smooth representation theory of $G(F)$. On the other hand, there are subtle issues such as the independence of ℓ in the Langlands parametrization, which we refer to [Lafforgue 2018, §12.2.4] for further discussions.

Our main local result is.

Theorem 1.2 (Theorem 3.2.2). *Let F be a nonarchimedean local field of characteristic $p > 0$ and G be a connected reductive F -group. Fix $\ell \neq p$ as above. If an irreducible smooth representation π of $G(F)$ has parameter $\phi \in \Phi(G)$ under the Langlands parametrization of Genestier and Lafforgue, then $\check{\pi}$ has parameter ${}^L \theta \circ \phi$.*

Remark 1.3. The prefix L for local parameters and local packets is dropped for the following reason. The parameters of Genestier and Lafforgue are always *semisimple* or completely reducible in the sense of Serre [2005]; in other words, the monodromy part of the Weil–Deligne parameter is trivial; see Lemma 2.4.4. As mentioned in [Genestier and Lafforgue 2017], one expects that their parameter is the semisimplification of the “true” L -parameter of π . Hence the packets Π_ϕ in question are larger than expected, and the Langlands parametrization we adopt is coarser, unless when ϕ does not factorize through any Levi ${}^L M \hookrightarrow {}^L G$, i.e., ϕ is *semisimple and elliptic*.

Our strategy is to reduce it into a global statement. Let $\mathring{F}$ be a global field of characteristic $p > 0$, say $\mathring{F} = \mathbb{F}_q(X)$ for a geometrically irreducible smooth proper $\mathbb{F}_q$ -curve X , and set $\mathbb{A} = \mathbb{A}_{\mathring{F}}$. Let G be a connected reductive $\mathring{F}$ -group. Fix a level $N \subset X$, whence the corresponding congruence subgroup $K_N \subset G(\mathbb{A})$ and the Hecke algebra $C_c(K_N \backslash G(\mathbb{A})/K_N; \overline{\mathbb{Q}}_\ell)$. Also fix a cocompact lattice Ξ in $A_G(\mathring{F}) \backslash A_G(\mathbb{A})$

where $A_G \subset G$ is the maximal central split torus. *Grosso modo*, the global Langlands parametrization in [Lafforgue 2018] is deduced from a commutative $\overline{\mathbb{Q}}_\ell$ -algebra $\mathcal{B}$ acting on the Hecke module

$$H_{\{0\},1} := C_c^{\text{cusp}}(\text{Bun}_{G,N}(\mathbb{F}_q)/\Xi; \overline{\mathbb{Q}}_\ell) = \bigoplus_{\alpha \in \ker^1(\mathring{F}, G)} C_c^{\text{cusp}}(G_\alpha(\mathring{F}) \backslash G_\alpha(\mathbb{A})/K_N \Xi; \overline{\mathbb{Q}}_\ell)$$

of $\overline{\mathbb{Q}}_\ell$ -valued cusp forms, extended across pure inner forms indexed by $\ker^1(\mathring{F}, G)$ (finite in number). The algebra $\mathcal{B}$ is generated by the *excursion operators* $S_{I,f,\vec{\gamma}}$. For any character $\nu : \mathcal{B} \rightarrow \overline{\mathbb{Q}}_\ell$ of algebras, denote by $\mathfrak{H}_\nu$ the generalized ν -eigenspace of $H_{\{0\},1}$. Then $H_{\{0\},1} = \bigoplus_\nu \mathfrak{H}_\nu$ as Hecke modules. Moreover, Lafforgue's machinery of ${}^L G$ -pseudocharacters associates a semisimple L -parameter $\sigma : \text{Gal}(\overline{\mathbb{F}}|\mathring{F}) \rightarrow {}^L G(\overline{\mathbb{Q}}_\ell)$ to ν . In fact ν is determined by σ , so that we may write $\mathfrak{H}_\sigma = \mathfrak{H}_\nu$.

There is an evident Hecke-invariant bilinear form on $H_{\{0\},1}$, namely the *integration pairing*

$$\langle h, h' \rangle := \sum_{\alpha \in \ker^1(\mathring{F}, G)} \int_{G_\alpha(\mathring{F}) \backslash G_\alpha(\mathbb{A})/\Xi} hh', \quad h, h' \in H_{\{0\},1},$$

with respect to some Haar measure on $G(\mathbb{A}) = G_\alpha(\mathbb{A})$ which is $\mathbb{Q}$ -valued on compact open subgroups. It is nondegenerate as easily seen by passing to $\overline{\mathbb{Q}}_\ell \simeq \mathbb{C}$. Now comes our global theorem.

Theorem 1.4 (Theorem 3.3.2). *If σ, σ' are two semisimple L -parameters for G such that $\langle \cdot, \cdot \rangle$ is nontrivial on $\mathfrak{H}_\sigma \otimes \mathfrak{H}_{\sigma'}$, then $\sigma' = {}^L\theta \circ \sigma$ up to $\hat{G}(\overline{\mathbb{Q}}_\ell)$ -conjugacy.*

Our local-global argument runs by first reducing Theorem 1.2 to the case that π is integral supercuspidal such that ω_π has finite order when restricted to A_G ; this step makes use of the compatibility of Langlands parametrization with parabolic induction, as established in [Genestier and Lafforgue 2017]. The second step is to globalize π into a cuspidal automorphic representation $\hat{\pi}$ with a suitable global model of G and Ξ , satisfying $\hat{\pi}^{K_N} \neq \{0\}$. The subspaces $\mathfrak{H}_\sigma$ of $H_{\{0\},1}$ might have isomorphic irreducible constituents in common, but upon modifying the automorphic realization, one can always assume that $\hat{\pi}^{K_N}$ lands in some $\mathfrak{H}_\sigma$. An application of Theorem 1.4 and the local-global compatibility of Langlands parametrization [Genestier and Lafforgue 2017] will conclude the proof.

The proof of Theorem 1.4 relies upon the determination of the transpose $S \mapsto S^*$ of excursion operators with respect to $\langle \cdot, \cdot \rangle$, namely the Lemma 5.3.3:

$$S_{I,f,\vec{\gamma}}^* = S_{I,f^\dagger,\vec{\gamma}^{-1}}$$

where $f \in \mathcal{O}(\hat{G} \backslash ({}^L G)^I // \hat{G})$, the finite set I and $\vec{\gamma} \in \text{Gal}(\overline{\mathbb{F}}|\mathring{F})^I$ are the data defining excursion operators, and $f^\dagger(\vec{g}) = f({}^L\theta(\vec{g})^{-1})$ for $\vec{g} \in ({}^L G)^I$. This property entails that if $\nu : \mathcal{B} \rightarrow \overline{\mathbb{Q}}_\ell$ corresponds to σ , then $\nu^* : S \mapsto \nu(S^*)$ corresponds to ${}^L\theta \circ \sigma$ (Proposition 5.3.4).

The starting point of the computation of the transpose is the fact that $\langle \cdot, \cdot \rangle$ is of geometric origin: it stems from the Verdier duality on the moduli stack $\text{Cht}_{N,I}^{(I_1, \dots, I_k)} / \Xi$ of chtoucas. The Chevalley involution intervenes ultimately in describing the effect of Verdier duality in geometric Satake equivalence, which is in turn connected to $\text{Cht}_{N,I}^{(I_1, \dots, I_k)} / \Xi$ via certain canonical smooth morphisms.

These geometric ingredients are already implicit in [Lafforgue 2018]. We just recast the relevant parts into our needs and supply some more details. In fact, the pairing $\langle \cdot, \cdot \rangle$ and its geometrization were used in a crucial way in older versions of [Lafforgue 2018]; that usage is now deprecated, and this article finds another application thereof.

Our third main result concerns the *duality involution* proposed by Prasad [2018, §3]. Assume that G is quasisplit. Fix an additive character ψ of G , an F -pinning $\mathcal{P}$ of G and the corresponding Whittaker datum $\mathfrak{w}$; replacing ψ by ψ^{-1} yields another Whittaker datum $\mathfrak{w}'$. Prasad defined an involution $\iota_{G,\mathcal{P}}$ as the commuting product of the Chevalley involution $\theta = \theta_{\mathcal{P}}$ of G and some inner automorphism ι_- which calibrates the Whittaker datum. Up to $G(F)$ -conjugation, this recovers the MVW involutions on classical groups [Mœglin et al. 1987, Chapitre 4] as well as the transpose-inverse on $\mathrm{GL}(n)$, whose relation with contragredient is well known.

Theorem 1.5 (Theorem 3.5.4). *Let $\phi \in \Phi(G)$ be a semisimple parameter such that Π_{ϕ} contains a unique $\mathfrak{w}$ -generic member π . Then $\Pi_{\iota_{\theta\phi}}$ satisfies the same property with respect to $\mathfrak{w}'$, and $\tilde{\pi} \simeq \pi \circ \iota_{G,\mathcal{P}} \in \Pi_{\iota_{\theta\phi}}$.*

Besides the crucial assumption which is expected to hold for tempered parameters if one works over $\mathbb{C}$ with true L -packets (called Shahidi’s tempered L -packet conjecture [1990]), the main inputs are Theorem 1.2 and the local “trivial functoriality” applied to $\iota_{G,\mathcal{P}}$ (see [Genestier and Lafforgue 2017, Théorèmes 0.1 and 8.1]). Due to these assumptions and the coarseness of our LLC, one should regard this result merely as some heuristic for Prasad’s conjectures [2018].

To conclude this introduction, let us mention two important issues that are left unanswered in this article:

- As in [Lafforgue 2018; Genestier and Lafforgue 2017], these techniques can be generalized to some *metaplectic coverings*, i.e., central extensions of locally compact groups

$$1 \rightarrow \mu_m(F) \rightarrow \tilde{G} \rightarrow G(F) \rightarrow 1$$

where $\mu_m(R) = \{z \in R^{\times} : z^m = 1\}$ as usual; it is customary to assume $\mu_m(F) = \mu_m(\bar{F})$ here. Fix a character $\zeta : \mu_m \hookrightarrow \overline{\mathbb{Q}_{\ell}^{\times}}$. One studies the irreducible smooth representations π of $\tilde{G}$ that are ζ -genuine, i.e., $\pi(\varepsilon) = \zeta(\varepsilon) \cdot \mathrm{id}$ for all $\varepsilon \in \mu_m(F)$. The most satisfactory setting for metaplectic coverings is due to Brylinski and Deligne [2001] that classifies the central extensions of G by $\mathbf{K}_2$ as sheaves over $(\mathrm{Spec} F)_{\mathrm{Zar}}$. Taking F -points and pushing-out from $\mathbf{K}_2(F)$ by norm-residue symbols yields a central extension above.

The L -group ${}^L\tilde{G}_{\zeta}$ associated to a Brylinski–Deligne $\mathbf{K}_2$ -central extension, m and ζ has been constructed in many situations; see the references in [Lafforgue 2018, §14]. Now consider the metaplectic variant of Conjecture 1.1. If π is ζ -genuine, $\tilde{\pi}$ will be ζ^{-1} -genuine so one needs a canonical L -isomorphism ${}^L\theta : {}^L\tilde{G}_{\zeta} \rightarrow {}^L\tilde{G}_{\zeta^{-1}}$; this is further complicated by the fact that ${}^L\tilde{G}_{\zeta}$ is not necessarily a split extension of groups. Although some results seem within reach when G is split, it seems more reasonable to work in the broader $\mathbf{K}_2$ -setting and incorporate the framework of Gaitsgory and Lysenko [2018] for the geometric part. Nonetheless, this goes beyond the scope of the present article.

- With powerful tools from p -adic Hodge theory, Fargues and Scholze proposed a program to obtain a local Langlands parametrization in characteristic zero, akin to that of Genestier and Lafforgue; see

[Fargues 2016] for an overview. It would certainly be interesting to try to adopt our techniques to characteristic zero. However, our key tools are global adélic in nature, whilst the setting of Fargues and Scholze is global in a different sense (over the Fargues–Fontaine curve). This hinders a direct translation into the characteristic zero setting.

Organization of this article. In Section 2, we collect the basic backgrounds on cusp forms, the integration pairing $\langle \cdot, \cdot \rangle$, contragredient representations and L -parameters, all in the ℓ -adic setting.

In Section 3, we begin by defining the Chevalley involution with respect to a chosen pinning and its extension to the L -group. Then we state the main Theorems 3.2.2 and 3.3.2 in the local and global cases, respectively. The local-global argument and the heuristic on duality involutions (Theorem 3.5.4) are also given there.

We give a brief overview of some basic vocabulary of [Lafforgue 2018] in Section 4. The only purpose of this section is to fix notation and serve as a preparation of the next section. As in [Lafforgue 2018; Genestier and Lafforgue 2017], we allow nonsplit groups as well.

The transposes of excursion operators are described in Section 5. It boils down to explicating the interplay between Verdier duality and partial Frobenius morphisms on the moduli stack of chtoucas. As mentioned before, a substantial part of this section can be viewed as annotations to [Lafforgue 2018], together with a few new computations. The original approach in Section 5 in an earlier manuscript has been substantially simplified following suggestions of Lafforgue.

In Sections 4 and 5, we will work exclusively in the global setting.

Conventions. Throughout this article, we fix a prime number ℓ distinct from the characteristic $p > 0$ of the fields under consideration. We also fix an algebraic closure $\overline{\mathbb{Q}_\ell}$ of the field $\mathbb{Q}_\ell$ of ℓ -adic numbers.

The six operations on ℓ -adic complexes are those defined in [Laszlo and Olsson 2008a; 2008b], for algebraic stacks locally of finite type over a reasonable base scheme, for example over $\mathrm{Spec} \mathbb{F}_q$ where q is some power of p . Given a morphism f of finite type between such stacks, the symbols $f_!$, f_* , etc. will always stand for the functors between derived categories $D_c^b(\dots, E)$ unless otherwise specified, where the field of coefficients E is some algebraic extension of $\mathbb{Q}_\ell$. The perverse t -structure on such stacks is defined in [Laszlo and Olsson 2009]; further normalizations will be recalled in Section 4.1. The constant sheaf associated to E on such a stack $\mathcal{X}$ is denoted by $E_{\mathcal{X}}$.

We use the notation $C_c(X; E)$ to indicate the space of compactly supported smooth E -valued functions on a topological space X , where E is any ring. Since we work exclusively over totally disconnected locally compact spaces, smoothness here means locally constant.

For a local or global field F , we denote by W_F the Weil group F with respect to a choice of separable closure $\overline{F}|F$, and by $I_F \subset \mathrm{Gal}(\overline{F}|F)$ the inertia subgroup. The arithmetic Frobenius automorphism is denoted by Frob . If F is local nonarchimedean, $\mathfrak{o}_F$ will stand for its ring of integers.

If $\mathring{F}$ is a global field, we write $\mathbb{A} = \mathbb{A}_{\mathring{F}} := \prod'_v \mathring{F}_v$ for its ring of adèles, where v ranges over the places of $\mathring{F}$. We also write $\mathfrak{o}_v = \mathfrak{o}_{\mathring{F}_v}$ in this setting.

For a scheme T , we write:

- $\Delta : T \hookrightarrow T^I$ for the diagonal morphism, where I is any set.
- $\pi_1(T, \bar{t})$ for the étale fundamental group with respect to a geometric point $\bar{t} \rightarrow T$, when T is connected, normal and locally Noetherian.
- $\mathcal{O}(T)$ for the ring of regular functions on T .
- $T_B := T \times_{\text{Spec } A} \text{Spec } B$ if T is a scheme over $\text{Spec } A$, and B is a commutative A -algebra.
- $E(T) := \text{Frac } \mathcal{O}(T)$ for the function field, when T is an irreducible variety over a field E .

Suppose that T is a variety over a field. The geometric invariant-theoretic quotient of T under the right action of some group variety Q , if it exists, is written as $T // Q$. Similar notation pertains to left or bilateral actions.

Let G be a connected reductive group over a field F . For any F -algebra A , denote the group of A -points of G by $G(A)$, endowed with a topology whenever A is. Denote by Z_G , G^{der} , G^{ad} for the center, derived subgroup and the adjoint group of G , respectively. Normalizers and centralizers in G are written as $N_G(\cdot)$ and $Z_G(\cdot)$. If $T \subset G$ is a maximal torus, we write T^{ad} , etc. for the corresponding subgroups in G^{ad} , etc. The character and cocharacter groups of a torus T are denoted by $X^*(T)$ and $X_*(T)$ as $\mathbb{Z}$ -modules, respectively.

The L -group and Langlands dual group of G are denoted by ${}^L G$ and $\hat{G}$, respectively. We use the Galois form of L -groups: details will be given in Section 2.4.

For an affine algebraic group H over some field E , the additive category of finite-dimensional algebraic representations of H will be denoted as $\text{Rep}_E(H)$. The trivial representation is denoted by $\mathbf{1}$. For any object $W \in \text{Rep}_E(H)$, we write $\check{W}$ or $W^\vee$ for its contragredient representation on $\text{Hom}_E(W, E)$. For any automorphism θ of H , write W^θ for the representation on W such that every $h \in H$ acts by $w \mapsto \theta(h) \cdot w$.

The same notation $\check{\pi}$ applies to the contragredient of a smooth representation π of a locally compact totally disconnected group. This will be the topic of Section 2.3. We denote the central character of an irreducible smooth representation π as ω_π .

2. Review of representation theory

2.1. Cusp forms. Let $\mathring{F}$ be a global field of characteristic $p > 0$. We may write $\mathring{F} = \mathbb{F}_q(X)$ where q is some power of p , and X is a smooth, geometrically irreducible proper curve over $\mathbb{F}_q$. Denote $\mathbb{A} = \mathbb{A}_{\mathring{F}}$. Fix a closed subscheme $N \subset X$ which is finite over $\mathbb{F}_q$, known as the level.

Let G be a connected reductive group over $\mathring{F}$. We associate to N a compact open subgroup

$$K_N := \ker \left[G \left(\prod_{v \in |X|} \mathfrak{o}_v \right) \rightarrow G(\mathcal{O}(N)) \right] \subset G(\mathbb{A}).$$

Denote the maximal split central torus in G by A_G . It is also known that there is a cocompact lattice

$$\Xi \subset A_G(\mathring{F}) \backslash A_G(\mathbb{A}),$$

which we fix once and for all. The space $G(\mathring{F}) \backslash G(\mathbb{A}) / \Xi$ is known to have finite volume with respect to any Haar measure on $G(\mathbb{A})$.

In what follows, we use a Haar measure on $G(\mathbb{A})$ such that $\text{mes}(K) \in \mathbb{Q}$ for any compact open subgroup K . The existence of such measures is established in [Vignéras 1996, Théorème 2.4]. The same convention pertains to the subgroups of G .

For all subextension $E | \mathbb{Q}_\ell$ of $\overline{\mathbb{Q}_\ell} | \mathbb{Q}_\ell$, we have the space

$$C_c(G(\mathring{F}) \backslash G(\mathbb{A}) / \Xi; E) = \bigcup_{N:\text{levels}} C_c(G(\mathring{F}) \backslash G(\mathbb{A}) / K_N \Xi; E)$$

of smooth E -valued functions of compact support on $G(\mathring{F}) \backslash G(\mathbb{A}) / \Xi$. Then $G(\mathbb{A})$ acts on the left of $C_c(G(\mathring{F}) \backslash G(\mathbb{A}) / \Xi; E)$ by $(gf)(x) = f(xg)$. Accordingly, $C_c(G(\mathring{F}) \backslash G(\mathbb{A}) / K_N \Xi; E)$, the space of K_N -invariants, is a left module under the unital E -algebra $C_c(K_N \backslash G(\mathbb{A}) / K_N; E)$, the Hecke algebra under convolution $\star$.

Our convention on Haar measures means that we can integrate E -valued smooth functions on $G(\mathring{F}) \backslash G(\mathbb{A}) / \Xi$, etc.

The subspace of $C_c(G(\mathring{F}) \backslash G(\mathbb{A}) / \Xi; E)$ of cuspidal functions

$$C_c^{\text{cusp}}(G(\mathring{F}) \backslash G(\mathbb{A}) / \Xi; E) = \bigcup_{N:\text{levels}} C_c^{\text{cusp}}(G(\mathring{F}) \backslash G(\mathbb{A}) / K_N \Xi; E)$$

is defined by either

- requiring that the constant terms $f_P(x) = \int_{U(\mathring{F}) \backslash U(\mathbb{A})} f(ux) du$ are zero whenever $P = MU \subsetneq G$ is a parabolic subgroup, or
- using the criterion in terms of Hecke-finiteness in [Lafforgue 2018, Proposition 8.23].

We record two more basic facts:

- The E -vector space $C_c^{\text{cusp}}(G(\mathring{F}) \backslash G(\mathbb{A}) / K_N \Xi; E)$ is finite-dimensional. This result is originally due to Harder, and can be deduced from the uniform bound on supports of such functions in [Mœglin and Waldspurger 1994, I.2.9].
- As a smooth $G(\mathbb{A})$ -representation, $C_c^{\text{cusp}}(G(\mathring{F}) \backslash G(\mathbb{A}) / \Xi; E)$ is absolutely semisimple, i.e., it is semisimple after $- \otimes_E \overline{\mathbb{Q}_\ell}$; see [Bourbaki 2012, VIII.226]. Indeed, the semisimplicity in the case $E = \overline{\mathbb{Q}_\ell} \simeq \mathbb{C}$ is well known.

In parallel, $C_c^{\text{cusp}}(G(\mathring{F}) \backslash G(\mathbb{A}) / K_N \Xi; E)$ is also absolutely semisimple as a $C_c(K_N \backslash G(\mathbb{A}) / K_N; E)$ -module. Recall the module structure: $f \in C_c(K_N \backslash G(\mathbb{A}) / K_N; E)$ acts on h as

$$(f \cdot h)(x) := \int_{K_N \backslash G(\mathbb{A}) / K_N} h(xg) f(g) dg = (h \star \check{f})(x), \quad x \in G(\mathbb{A}) \quad (2-1)$$

where $\check{f}(g) = f(g^{-1})$ and the convolution $\star$ is defined in the usual manner.

We record the following standard result for later use.

Proposition 2.1.1. *For every $G(\mathbb{A})$ -representation $\dot{\pi}$, assumed to be smooth, let $\dot{\pi}^{K_N}$ be the space of K_N -invariant vectors. It is a left module under $C_c(K_N \backslash G(\mathbb{A})/K_N; \overline{\mathbb{Q}_\ell})$.*

- (i) *For all irreducible $G(\mathbb{A})$ -representations $\dot{\pi}_1, \dot{\pi}_2$ generated by K_N -invariants, we have $\dot{\pi}_1 \simeq \dot{\pi}_2 \iff \dot{\pi}_1^{K_N} \simeq \dot{\pi}_2^{K_N}$ as simple $C_c(K_N \backslash G(\mathbb{A})/K_N; \overline{\mathbb{Q}_\ell})$ -modules.*
- (ii) *Given any decomposition $C_c^{\text{cusp}}(G(\dot{F}) \backslash G(\mathbb{A})/\Xi; \overline{\mathbb{Q}_\ell}) = \bigoplus_{\dot{\pi} \in \Pi} \dot{\pi}$ into irreducibles, where Π is a set (with multiplicities) of irreducible subrepresentations, we have*

$$C_c^{\text{cusp}}(G(\dot{F}) \backslash G(\mathbb{A})/K_N \Xi; \overline{\mathbb{Q}_\ell}) = \bigoplus_{\dot{\pi} \in \Pi, \dot{\pi}^{K_N} \neq 0} \dot{\pi}^{K_N}$$

in which each $\dot{\pi}^{K_N}$ is simple.

- (iii) *For every irreducible $G(\mathbb{A})$ -representation $\dot{\pi}$ generated by K_N -invariants, we have a natural isomorphism of multiplicity spaces*

$$\begin{aligned} \text{Hom}_{G(\mathbb{A})\text{-Rep}}(\dot{\pi}, C_c^{\text{cusp}}(G(\dot{F}) \backslash G(\mathbb{A})/\Xi; \overline{\mathbb{Q}_\ell})) \\ \xrightarrow{\sim} \text{Hom}_{C_c(K_N \backslash G(\mathbb{A})/K_N; \overline{\mathbb{Q}_\ell})\text{-Mod}}(\dot{\pi}^{K_N}, C_c^{\text{cusp}}(G(\dot{F}) \backslash G(\mathbb{A})/K_N \Xi; \overline{\mathbb{Q}_\ell})). \end{aligned}$$

Property (i) actually holds for representations of $G(\dot{F}_v)$ and of its Hecke algebras, for any place v of $\dot{F}$.

The C_c^{cusp} in (ii) and (iii) can be replaced by $\bigoplus_{\alpha \in \ker^1(\dot{F}, G)} C_c^{\text{cusp}}(G_\alpha(\dot{F}) \backslash G(\mathbb{A})/\Xi; \overline{\mathbb{Q}_\ell})$; see (2-2).

Proof. By semisimplicity, $C_c^{\text{cusp}}(G(\dot{F}) \backslash G(\mathbb{A})/\Xi; \overline{\mathbb{Q}_\ell})$ (or the $\bigoplus_\alpha$ version) decomposes uniquely into $W \oplus W'$ such that

- W is a subrepresentation isomorphic to a direct sum of irreducibles, each summand is generated by K_N -invariants;
- W' is a subrepresentation satisfying $(W')^{K_N} = \{0\}$.

For (ii)–(iii), it suffices to look at the $G(\mathbb{A})$ -representation W and the $C_c(K_N \backslash G(\mathbb{A})/K_N; \overline{\mathbb{Q}_\ell})$ -module W^{K_N} ; both are semisimple. The required assertions follow from the standard equivalences between categories in [Renard 2010, I.3 and III.1.5] and Schur’s lemma [Renard 2010, III.1.8 and B.II]. $\square$

Next, we introduce the moduli stack $\text{Bun}_{G,N}$ over $\mathbb{F}_q$ of G -torsors on X with level N structures: it maps any $\mathbb{F}_q$ -scheme S to the groupoid

$$\text{Bun}_{G,N}(S) = \left\{ (\mathcal{G}, \psi) \left| \begin{array}{l} \mathcal{G} \text{ a } G\text{-torsor over } X \times S \text{ and} \\ \psi: \mathcal{G}|_{N \times S} \xrightarrow{\sim} G|_{N \times S} \text{ a trivialization over } N \end{array} \right. \right\}, \quad \text{Bun}_G := \text{Bun}_{G,\emptyset}.$$

For this purpose, we need suitable models of G over X . Let $U \subset X$ be the maximal open subscheme such that G extends to a connected reductive U -group scheme. We follow [Lafforgue 2018, §12.1] to take parahoric models at the formal neighborhoods of all points of $X \setminus U$. Glue these parahoric models with the smooth model over U , à la Beauville–Laszlo, to yield a smooth affine X -group scheme with geometrically connected fibers, known as a *Bruhat–Tits group scheme* over X ; see also [Heinloth 2010, §1].

Regard $\text{Bun}_{G,N}(\mathbb{F}_q)$ as a set, on which Ξ acts naturally. As explained in [Lafforgue 2018], we have

$$\text{Bun}_{G,N}(\mathbb{F}_q) = \bigsqcup_{\alpha \in \ker^1(\mathring{F}, G)} G_\alpha(\mathring{F}) \backslash G_\alpha(\mathbb{A}) / K_N \quad (2-2)$$

where

- $\ker^1(\mathring{F}, G)$ is the kernel of $H^1(\mathring{F}, G) \rightarrow \prod_{v \in |X|} H^1(\mathring{F}_v, G)$;
- to each $\alpha \in \ker^1(\mathring{F}, G)$ is attached a locally trivial pure inner twist G_α of G , and we fix an identification $G_\alpha(\mathbb{A}) \simeq G(\mathbb{A})$.

The decomposition is compatible with Ξ -actions. The pointed set $\ker^1(\mathring{F}, G)$ is finite; it is actually trivial when G is split. As before, we have the spaces

$$C_c^{\text{cusp}}(\text{Bun}_{G,N}(\mathbb{F}_q) / \Xi; E) = \bigoplus_{\alpha \in \ker^1(\mathring{F}, G)} C_c^{\text{cusp}}(G_\alpha(\mathring{F}) \backslash G_\alpha(\mathbb{A}) / K_N \Xi; E).$$

The cuspidality on the left-hand side can be defined in terms of Hecke-finiteness as before. We shall also use compatible Haar measures on various $G_\alpha(\mathbb{A})$.

From the viewpoint of harmonic analysis, the mere effect of working with $\text{Bun}_{G,N}(\mathbb{F}_q)$ is to consider all the inner twists from $\ker^1(\mathring{F}, G)$ at once. See also [Lafforgue 2018, §12.2.5].

2.2. Integration pairing. Let E be a subextension of $\overline{\mathbb{Q}_\ell} | \mathbb{Q}_\ell$.

Definition 2.2.1. With the Haar measures as in Section 2.1, we define the *integration pairing*

$$\langle \cdot, \cdot \rangle : C_c^{\text{cusp}}(G(\mathring{F}) \backslash G(\mathbb{A}) / \Xi; E) \otimes_E C_c^{\text{cusp}}(G(\mathring{F}) \backslash G(\mathbb{A}) / \Xi; E) \rightarrow E$$

$$h \otimes h' \mapsto \langle h, h' \rangle := \int_{G(\mathring{F}) \backslash G(\mathbb{A}) / \Xi} hh'.$$

The pairing is clearly E -bilinear, symmetric and $G(\mathbb{A})$ -invariant. There is an obvious variant for not necessarily cuspidal functions.

Lemma 2.2.2. *The pairing $\langle \cdot, \cdot \rangle$ above is absolutely nondegenerate, i.e., its radical equals $\{0\}$ after $-\otimes_E \overline{\mathbb{Q}_\ell}$.*

Proof. It is legitimate to assume $E = \overline{\mathbb{Q}_\ell}$, and there exists an isomorphism of fields $\overline{\mathbb{Q}_\ell} \simeq \mathbb{C}$. The nondegeneracy over $\mathbb{C}$ is well known: we have $\int h \bar{h} \geq 0$, and equality holds if and only if $h = 0$. $\square$

Remark 2.2.3. For a chosen level $N \subset X$, we have an analogous pairing

$$\langle \cdot, \cdot \rangle : C_c^{\text{cusp}}(G(\mathring{F}) \backslash G(\mathbb{A}) / K_N \Xi; E) \otimes_E C_c^{\text{cusp}}(G(\mathring{F}) \backslash G(\mathbb{A}) / K_N \Xi; E) \rightarrow E$$

$$h \otimes h' \mapsto \langle h, h' \rangle := \int_{G(\mathring{F}) \backslash G(\mathbb{A}) / K_N \Xi} hh'.$$

The integration here is actually a “stacky” sum over $G(\mathring{F}) \backslash G(\mathbb{A}) / K_N \Xi$, i.e., $\langle h, h' \rangle$ equals that of Definition 2.2.1 if one starts with a Haar measure on $G(\mathbb{A})$ with $\text{mes}(K_N) = 1$. It is also E -bilinear, symmetric, absolutely nondegenerate and invariant in the sense that

$$\langle f \cdot h, h' \rangle = \langle h, \check{f} \cdot h' \rangle, \quad f \in C_c(K_N \backslash G(\mathbb{A}) / K_N; E);$$

see (2-1). There is an obvious variant for not necessarily cuspidal functions.

The spaces in question being finite-dimensional, it makes sense to talk about the *transpose* of a linear operator. For example, the transpose of the left multiplication by f is given by that of $\check{f}$.

As in Section 2.1, the integration pairing extends to

$$\langle \cdot, \cdot \rangle : C_c^{\text{cusp}}(\text{Bun}_{G,N}(\mathbb{F}_q)/\Xi; E) \otimes_E C_c^{\text{cusp}}(\text{Bun}_{G,N}(\mathbb{F}_q)/\Xi; E) \rightarrow E$$

$$h \otimes h' \mapsto \int_{\text{Bun}_{G,N}(\mathbb{F}_q)/\Xi} hh'.$$

This is the orthogonal sum of the integrations pairings on various $G_\alpha(\mathbb{A})$.

2.3. Representations. In this subsection, we let F be a local field of characteristic $p > 0$. Denote the cardinality of the residue field of F as q . Let G be a connected reductive F -group. The smooth representations of $G(F)$ will always be realized on $\overline{\mathbb{Q}_\ell}$ -vector spaces. Irreducible smooth representation of $G(F)$ are admissible; see [Renard 2010, VI.2.2].

The smooth characters of $G(F)$ are homomorphisms $G(F) \rightarrow \overline{\mathbb{Q}_\ell}^\times$ with open kernel. We will need to look into a class of particularly simple characters, namely those trivial on the open subgroup

$$G(F)^1 := \bigcap_{\chi \in X^*(G)} \ker |\chi|_F \quad (2-3)$$

of $G(F)$, where $X^*(G) := \text{Hom}_{\text{alg. grp}}(G, \mathbb{G}_m)$ and

$$|\cdot|_F : F^\times \rightarrow q^{\mathbb{Z}} \subset \mathbb{Q}^\times$$

is the normalized absolute value on F . Note that $G(F)/G(F)^1 \simeq \mathbb{Z}^r$ with $r := \text{rk}_{\mathbb{Z}} X^*(G)$. Moreover, $G(F)^1 \supset G^{\text{der}}(F) = G^{\text{der}}(F)^1$.

For any smooth character ω of $Z_G(F)$, denote by $C_c(G(F), \omega)$ the space of functions $f : G(F) \rightarrow \overline{\mathbb{Q}_\ell}$ such that $f(zg) = \omega(z)f(g)$ for all $z \in Z_G(F)$ and $\text{Supp}(f)$ is compact modulo $Z_G(F)$.

Let $\text{Ind}_P^G(\cdot)$ denote the unnormalized parabolic induction from the Levi quotient of $P \subset G$. Let δ_P denote the modulus character of $P(F)$ taking values in $q^{\mathbb{Z}}$. Upon choosing $q^{1/2} \in \overline{\mathbb{Q}_\ell}$, we can also form the *normalized parabolic induction* $I_P^G(\cdot) := \text{Ind}_P^G(\cdot \otimes \delta_P^{1/2})$.

We need the notion [Renard 2010, VI.7.1] of the *cuspidal support* (M, τ) of an irreducible smooth representation π . Here $M \subset G$ is a Levi subgroup and τ is a supercuspidal irreducible representation of $M(F)$, such that π is a subquotient of $I_P^G(\tau)$ for any parabolic subgroup $P \subset G$ with Levi component M . The cuspidal support is unique up to $G(F)$ -conjugacy. It is known that one can choose P with Levi component M such that $\pi \hookrightarrow I_P^G(\tau)$. See [Renard 2010, VI.5.4].

We collect below a few properties of an irreducible smooth representation π of $G(F)$:

(1) Suppose that π is supercuspidal. There exists a finite extension E of $\mathbb{Q}_\ell$ such that π is defined over E . Indeed, since the central character ω_π can be defined over some finite extension of $\mathbb{Q}_\ell$, so is $\pi \hookrightarrow C_c(G(F), \omega_\pi)$.

From this and the discussion on cuspidal supports, it follows that every π can be defined over some finite extension E of $\mathbb{Q}_\ell$.

(2) We say π is *integral* if it admits an $\mathfrak{o}_E$ -model of finite type, where E is a finite extension of $\mathbb{Q}_\ell$. See [Vignéras 2001, §1.4] for details. Then an irreducible supercuspidal π is integral if and only if ω_π has ℓ -adically bounded image in $\overline{\mathbb{Q}_\ell}^\times$.

Again, this is a consequence of $\pi \hookrightarrow C_c(G(F), \omega_\pi)$. It also implies the notion of integrality stated in the beginning of [Genestier and Lafforgue 2017].

(3) Let V be the underlying vector space of π . The *contragredient representation* $\check{\pi}$ of a smooth representation π is realized on the space $V^\vee$ of the smooth vectors in $\text{Hom}_{\overline{\mathbb{Q}_\ell}}(V, \overline{\mathbb{Q}_\ell})$. It satisfies $\langle \check{\rho}(g)\check{v}, v \rangle = \langle \check{v}, \rho(g^{-1})v \rangle$. If π is defined over E , so is $\check{\pi}$. Taking contragredient preserves irreducibility and supercuspidality. It is clear that $(\pi \otimes \chi)^\vee = \check{\pi} \otimes \chi^{-1}$ for any smooth character $\chi : G(F) \rightarrow \overline{\mathbb{Q}_\ell}^\times$.

(4) Moreover, $(\pi)^{\vee\vee} \simeq \pi$ for all smooth irreducible π ; see [Renard 2010, III.1.7]. Also, $\omega_{\check{\pi}} = \omega_\pi^{-1}$.

Proposition 2.3.1. *If π is an irreducible smooth representation of $G(F)$ with cuspidal support (M, τ) , then $\check{\pi}$ has cuspidal support $(M, \check{\tau})$.*

Proof. Choose a parabolic subgroup $P \subset G$ with M as Levi component such that $\pi \hookrightarrow I_P^G(\tau)$. Once the Haar measures are chosen, we have $I_P^G(\tau)^\vee \simeq I_P^G(\check{\tau})$ canonically; see [Bushnell and Henniart 2006, §3.5]. Dualizing, we deduce $I_P^G(\check{\tau}) \twoheadrightarrow \check{\pi}$. Thus $\check{\pi}$ is a subquotient of $I_P^G(\check{\tau})$. $\square$

2.4. L-parameters. Let F be a local or global field of characteristic $p > 0$. For a connected reductive F -group G , we denote by $\tilde{F}|F$ the splitting field of G , which is a finite Galois extension inside a chosen separable closure $\bar{F}$.

Denote by W_F the absolute Weil group of F . It comes with canonical continuous homomorphisms (i) $W_F \rightarrow \text{Gal}(\bar{F}|F)$, and (ii) $W_{F_v} \rightarrow W_F$ if F is global and v is a place of F . For (ii) we choose an embedding $\bar{F} \hookrightarrow \bar{F}_v$ of separable closures.

Definition 2.4.1. The *Langlands dual group* $\hat{G}$ of G is a pinned connected reductive $\mathbb{Q}_\ell$ -group (in fact, definable over $\mathbb{Z}$), on which $\text{Gal}(\tilde{F}|F)$ operates by pinned automorphisms. Throughout this article, we use the *finite Galois forms* of the L -group of G , namely

$${}^L G := \hat{G} \rtimes \text{Gal}(\tilde{F}|F)$$

viewed as an affine algebraic group.

If $M \hookrightarrow G$ is a Levi subgroup, we obtain a the corresponding embedding ${}^L M \rightarrow {}^L G$ of standard Levi subgroup.

Definition 2.4.2. An L -parameter for G is a homomorphism $\sigma : W_F \rightarrow {}^L G(\overline{\mathbb{Q}_\ell})$ such that:

- The following diagram commutes:

$$\begin{array}{ccc} W_F & \xrightarrow{\sigma} & {}^L G \\ & \searrow & \swarrow \\ & \text{Gal}(\tilde{F}|F) & \end{array}$$

- σ is continuous with respect to the ℓ -adic topology on ${}^L G(\overline{\mathbb{Q}}_\ell)$.
- σ is *relevant* in the sense of [Borel 1979, §8.2], which matters only when G is not quasisplit.
- (The local case) σ is Frobenius semisimple: $\rho(\sigma(\text{Frob}))$ is semisimple for every algebraic representation $\rho : {}^L G(\overline{\mathbb{Q}}_\ell) \rightarrow \text{GL}(N, \overline{\mathbb{Q}}_\ell)$, where Frob stands for any Frobenius element in W_F (see [Bushnell and Henniart 2006, 32.7 Proposition] for more discussions on Frobenius-semisimplicity).
- (The global case) σ is semisimple in the sense of [Serre 2005], to be described below. We do not require Frobenius-semisimplicity here because for ℓ -adic representations of geometric origin, that property is a long-standing conjecture in étale cohomology.

The set of $\hat{G}(\overline{\mathbb{Q}}_\ell)$ -conjugacy classes of L -parameters is denoted as $\Phi(G)$. By [Borel 1979, §3.4], there is a natural map $\Phi(M) \rightarrow \Phi(G)$ for any Levi subgroup M .

Remark 2.4.3. Since ${}^L G(\overline{\mathbb{Q}}_\ell)$ carries the ℓ -adic topology and σ is required to be continuous, when F is local we get rid of the Weil–Deligne group in the usual formulation in terms of ${}^L G(\mathbb{C})$. Besides, we do not consider Arthur parameters in this article.

As recalled earlier, the structure of Weil groups allows us to

- localize a global L -parameter at a place v ;
- talk about L -parameters of the form $\text{Gal}(\overline{F}|F) \rightarrow {}^L G(\overline{\mathbb{Q}}_\ell)$ and their localizations when F is global.

Next, we recall the *semisimplicity* of L -parameters following [Lafforgue 2018; Serre 2005]: a continuous homomorphism $\sigma : W_F \rightarrow {}^L G(\overline{\mathbb{Q}}_\ell)$ is called semisimple if the Zariski closure of $\text{im}(\sigma)$ is reductive in ${}^L G(\overline{\mathbb{Q}}_\ell)$, in the sense that its identity connected component is reductive. When G is split, this is exactly the definition of complete reducibility in [Serre 2005, 3.2.1], say by applying [loc. cit., Proposition 4.2].

Lemma 2.4.4. *Assume F is local. The following are equivalent for any L -parameter σ for G :*

- σ is semisimple.
- The Weil–Deligne parameter associated to σ has trivial nilpotent part.

Proof. By composing σ with any faithful algebraic representation $\rho : {}^L G(\overline{\mathbb{Q}}_\ell) \hookrightarrow \text{GL}(N, \overline{\mathbb{Q}}_\ell)$, we may assume that σ is an ℓ -adic representation $W_F \rightarrow \text{GL}(N, \overline{\mathbb{Q}}_\ell)$. To σ is associated the Weil–Deligne representation $\text{WD}(\sigma)$: it comes with a nilpotent operator n . For details, see [Bushnell and Henniart 2006, 32.5].

(i) $\Rightarrow$ (ii): The line $\overline{\mathbb{Q}}_\ell n$ is preserved by $\text{im}(\sigma)$ -conjugation. Since $\exp(tn) \in \text{im}(\sigma)$ for $t \in \mathbb{Z}_\ell$ with $|t| \ll 1$, the semisimplicity of σ forces $n = 0$.

(ii) $\Rightarrow$ (i): As $n = 0$, the smooth representation underlying $\text{WD}(\sigma)$ is just σ , hence σ is semisimple as a smooth representation of W_F by [Bushnell and Henniart 2006, 32.7 Theorem]. The reductivity (or complete reducibility) of the Zariski closure of $\text{im}(\sigma)$ then follows from the theory in [Serre 2005]. $\square$

Finally, we define parabolic subgroups of ${}^L G$ as in [Borel 1979, 3.2]. They are subgroups of the form $N_{L_G}(\hat{P})$ where $\hat{P} \subset \hat{G}$ is a parabolic subgroups, and whose projection to $\text{Gal}(\tilde{F}|F)$ has full image. Define the unipotent radical of such a parabolic subgroup to be that of $\hat{P}$. We still have the notion of Levi decomposition in this setting; see [Borel 1979, 3.4].

Following [Lafforgue 2018, §13], the *semisimplification* σ^{ss} of an L -parameter σ is defined as follows:

- First, take the smallest parabolic subgroup ${}^L P \subset {}^L G$ containing $\text{im}(\sigma)$.
- Project to the Levi quotient.
- Then embed back into ${}^L G$ using some Levi decomposition.

The resulting parameter is well-defined up to $\hat{G}(\overline{\mathbb{Q}_\ell})$ -conjugacy.

By definition, an L -homomorphism ${}^L H \rightarrow {}^L G$ between L -groups is an algebraic homomorphism respecting the projections to $\text{Gal}(\tilde{F}|F)$.

Lemma 2.4.5. *Up to $\hat{G}(\overline{\mathbb{Q}_\ell})$ -conjugacy, semisimplification commutes with L -automorphisms of ${}^L G$.*

Proof. Indeed, an L -automorphism permutes the parabolic subgroups of ${}^L G$ together with their Levi decompositions. $\square$

3. Statement of a variant of the conjecture

3.1. Chevalley involutions. To begin with, we consider a split connected reductive group H over a field, equipped with a pinning $\mathcal{P} = (B, T, (X_\alpha)_{\alpha \in \Delta_0})$, where

- (B, T) is a Borel pair of H , and
- X_α is a nonzero vector in the root subspace $\mathfrak{h}_\alpha$, where α ranges over the set Δ_0 of B -simple roots.

Definition 3.1.1. The *Chevalley involution* $\theta = \theta_{\mathcal{P}}$ is the unique pinned automorphism of H acting as $t \mapsto w_0(t^{-1})$ on T , where w_0 stands for the longest element in the Weyl group associated to T .

This is the definition in [Prasad 2018, §4], and it is clear that $\theta^2 = \text{id}_H$.

The Chevalley involution will be considered in the following settings. Let F be a field with separable closure $\bar{F}$.

- (1) Let $H = \hat{G}$ be the dual group of G , which is connected reductive over F . The dual group is endowed with a pinning and we obtain $\theta : \hat{G} \rightarrow \hat{G}$. Since $\text{Gal}(\tilde{F}|F)$ operates by pinned automorphisms on $\hat{G}$, the Chevalley involution extends to

$${}^L \theta : {}^L G \rightarrow {}^L G, \quad g \rtimes \sigma \mapsto \theta(g) \rtimes \sigma,$$

which is still an involution.

- (2) Let G be a quasisplit connected reductive group over F . Then G admits an F -pinning $\mathcal{P}$, i.e., a Galois-invariant pinning of $H := G_{\bar{F}}$. Therefore the Chevalley involution $\theta = \theta_{\mathcal{P}}$ for $G_{\bar{F}}$ descends to G .

Furthermore, observe that if $H = \prod_{i=1}^r H_i$ and $\mathcal{P}$ decomposes into $(\mathcal{P}_1, \dots, \mathcal{P}_r)$ accordingly, the corresponding Chevalley involution $\theta_{\mathcal{P}}$ equals $\prod_{i=1}^r \theta_{\mathcal{P}_i}$.

3.2. The local statement. Let F be a local field of characteristic $p > 0$. Let G be a connected reductive F -group. The set of isomorphism classes of irreducible smooth representations over $\overline{\mathbb{Q}}_\ell$ of $G(F)$ will be denoted by $\Pi(G)$. The local statement to follow presumes a given *Langlands parametrization* of representations, namely an arrow

$$\Pi(G) \rightarrow \Phi(G)$$

$$\pi \mapsto \phi.$$

This is the “automorphic to Galois” direction of the local Langlands correspondence for G . We say that ϕ is the parameter of π , and denote by $\Pi_\phi \subset \Pi(G)$ the fiber over ϕ , called the *packet* associated to ϕ .

For the local statement, we employ the Langlands parametrization furnished by Genestier and Lafforgue [2017]. It is actually an arrow

$$\Pi(G) \rightarrow \{\text{semisimple } L\text{-parameters}\} / \hat{G}(\overline{\mathbb{Q}}_\ell)\text{-conj.} \subset \Phi(G).$$

Remark 3.2.1. The Genestier–Lafforgue parameters are expected to be the semisimplifications of authentic (yet hypothetical) Langlands parameters. As a consequence, the packets Π_ϕ for general Genestier–Lafforgue parameters are expected to be a disjoint union of authentic L -packets, unless when ϕ is an elliptic parameter (see Lemma 2.4.4), i.e., $\text{im}(\phi)$ is ${}^L G$ -ir in the sense of [Serre 2005, 3.2.1].

Further descriptions and properties of the Genestier–Lafforgue parametrization will be reviewed in due course. Let us move directly to the main local statement.

Theorem 3.2.2. *Let $\phi \in \Phi(G)$ be a semisimple L -parameter. In terms of the Langlands parametrization of Genestier–Lafforgue, we have*

$$\{\check{\pi} : \pi \in \Pi_\phi\} = \Pi_{\iota_\theta \circ \phi},$$

where $\iota_\theta : {}^L G \rightarrow {}^L G$ is the Chevalley involution in Section 3.1.

If the Genestier–Lafforgue parametrization is replaced by an authentic Langlands parametrization, the statement above becomes [Adams and Vogan 2016, Conjecture 1.1]; it is also a part of [Prasad 2018, §4, Conjecture 2], but Prasad’s conjecture also predicates on the internal structure of L -packets. The conjecture of Adams, Vogan, and Prasad applies to any local field F ; known cases in this generality include:

- The case $F = \mathbb{R}$ in [Adams and Vogan 2016, Theorem 7.1(a)], with admissible representations of $G(\mathbb{R})$ over $\mathbb{C}$.
- The tempered L -packets for symplectic groups $\text{Sp}(2n)$ and quasisplit SO groups over nonarchimedean local fields F of characteristic zero in terms of Arthur’s endoscopic classification, see [Kaletha 2013, Corollary 5.10].
- The depth-zero and epipelagic L -packets for many p -adic groups [Kaletha 2013, §6].

Each case above requires a different construction of L -packets, applicable to different groups or parameters, whereas the Theorem 3.2.2 furnishes a uniform statement. On the other hand, Theorem 3.2.2 is weaker since the Langlands parametrization here is coarser, in view of the Remark 3.2.1.

The proof of Theorem 3.2.2 will occupy Section 3.4.

3.3. The global statement. Theorem 3.2.2 will be connected to the global result below.

Let $\mathring{F} = \mathbb{F}_q(X)$ and fix the level $N \subset X$ as in Section 2.1. Let G , K_N and Ξ be as in Section 2.1, so that $\text{Bun}_{G,N}$ is defined. Note that we need to choose a model of G over X which is a Bruhat–Tits group scheme, still denoted as G . Let $U \subset X$ denote the (open) locus of good reduction of G , and set

$$\hat{N} := N \cup (X \setminus U). \quad (3-1)$$

This is a finite closed $\mathbb{F}_q$ -subscheme of X , the “unramified locus”. Let $\eta \rightarrow X$ be the generic point of X ; fix a geometric generic point $\bar{\eta} \rightarrow \eta$ of X .

The main global result of Lafforgue [2018, Théorème 12.3] gives a canonical decomposition of $C_c(K_N \backslash G(\mathbb{A})/K_N; \overline{\mathbb{Q}_\ell})$ -modules

$$C_c^{\text{cusp}}(\text{Bun}_{G,N}(\mathbb{F}_q)/\Xi; \overline{\mathbb{Q}_\ell}) = \bigoplus_{\sigma} \mathfrak{H}_{\sigma} \quad (3-2)$$

indexed by L -parameters $\sigma : \text{Gal}(\bar{F}|\mathring{F}) \rightarrow {}^L G(\overline{\mathbb{Q}_\ell})$ up to $\hat{G}(\overline{\mathbb{Q}_\ell})$ -conjugacy that

- are semisimple, and
- factor continuously through $\text{Gal}(\bar{F}|\mathring{F}) \rightarrow \pi_1(X \setminus \hat{N}, \bar{\eta})$.

Remark 3.3.1. Since the left-hand side of (3-2) is a semisimple module, of finite dimension over $\overline{\mathbb{Q}_\ell}$, so are its submodules $\mathfrak{H}_{\sigma}$. To each σ we may associate a set (with multiplicities) of simple submodules C_{σ} , such that

$$\mathfrak{H}_{\sigma} = \bigoplus_{\mathcal{L} \in C_{\sigma}} \mathcal{L}, \quad \text{hence} \quad C_c^{\text{cusp}}(\text{Bun}_{G,N}(\mathbb{F}_q)/\Xi; \overline{\mathbb{Q}_\ell}) = \bigoplus_{\sigma} \bigoplus_{\mathcal{L} \in C_{\sigma}} \mathcal{L}$$

as $C_c(K_N \backslash G(\mathbb{A})/K_N; \overline{\mathbb{Q}_\ell})$ -modules.

The decomposition (3-2) is built on two pillars: the theories of excursion operators and pseudocharacters for ${}^L G$. As in the local case, we defer the necessary details of [Lafforgue 2018] to Section 4.

Theorem 3.3.2. Suppose that $\mathfrak{H}_{\sigma}, \mathfrak{H}_{\sigma'}$ are two nonzero summands in (3-2) such that the restriction

$$\langle \cdot, \cdot \rangle_{\sigma, \sigma'} : \mathfrak{H}_{\sigma} \otimes_{\overline{\mathbb{Q}_\ell}} \mathfrak{H}_{\sigma'} \rightarrow \overline{\mathbb{Q}_\ell}$$

of the integration pairing $\langle \cdot, \cdot \rangle$ of Remark 2.2.3 (extended to $\text{Bun}_{G,N}(\mathbb{F}_q)/\Xi$) is not identically zero. Then we have

$$\sigma' = {}^L \theta \circ \sigma \quad \text{in } \Phi(G);$$

here ${}^L \theta$ is the Chevalley involution of ${}^L G$.

The proof of Theorem 3.3.2 will be accomplished at the end of Section 5.3.

3.4. Local-global argument. Consider a connected reductive group G over a local field F of characteristic p as in the local setting Section 3.2. As usual, A_G stands for the maximal central split torus in G , and $\tilde{F}|F$ stands for the splitting field of G . Take a maximal torus $T \subset G$ with splitting field equal to $\tilde{F}$. Let $H^1(W_F, Z_{\hat{G}})$ denote the continuous cohomology with values in $Z_{\hat{G}}(\overline{\mathbb{Q}_\ell})$ with discrete topology.

The first lemma concerns the Langlands parametrization of smooth characters of $G(F)$. The general case turns out to be delicate: by the discussion in [Lapid and Mao 2015, Appendix A], the usual cohomological construction actually yields an arrow in the opposite direction:

$$\begin{array}{ccc} H^1(W_F, Z_{\hat{G}}) & \longrightarrow & \{\eta : G(F) \rightarrow \overline{\mathbb{Q}_\ell}^\times, \text{ smooth character} \} \\ \downarrow & & \\ \Phi(G) & & \end{array}$$

It is injective but not necessarily surjective. However, we only need the invert it when $\eta|_{G(F)^1}$ is trivial. This is well known to experts, and below is a sketch.

Lemma 3.4.1. *For G as above, there is a canonical homomorphism of groups*

$$\{\eta : G(F)/G(F)^1 \rightarrow \overline{\mathbb{Q}_\ell}^\times, \text{ a smooth character} \} \rightarrow H^1(W_F, Z_{\hat{G}}).$$

Here we do not assume $\text{char}(F) > 0$.

Proof. Fix η . First, one can take a z -extension of G as in [Lapid and Mao 2015, Proof of Lemma A.1], i.e., a central extension

$$1 \rightarrow C \rightarrow G_1 \xrightarrow{p} G \rightarrow 1, \quad C \text{ is an induced torus, } G_1^{\text{der}} \text{ simply connected.}$$

Then $\eta_1 := \eta \circ p$ is trivial on $G_1(F)^1$. We know that $H^1(F, G_1^{\text{der}})$ is trivial. Put $S := G_1/G_1^{\text{der}}$ so that $G_1(F)/G_1^{\text{der}}(F) \xrightarrow{\sim} S(F)$ and $\hat{S} \simeq Z_{\hat{G}_1} = Z_{\hat{G}_1}^\circ$. Then $G_1^{\text{der}}(F) \subset G_1(F)^1$ implies that η_1 factors through $S(F)$. The local classfield theory affords an element $a \in H^1(W_F, Z_{\hat{G}_1})$. Since $\eta_1|_C = 1$, we infer that a has trivial image in $H^1(W_F, \hat{C})$.

Furthermore, using the fact that C is induced, in [loc. cit.] the following natural isomorphism is constructed:

$$H^1(W_F, Z_{\hat{G}}) \simeq \ker[H^1(W_F, Z_{\hat{G}_1}) \rightarrow H^1(W_F, \hat{C})].$$

All in all, we obtain $a \in H^1(W_F, Z_{\hat{G}})$. It is routine to check that $\eta \mapsto a$ is independent of the choice of z -extensions, see [loc. cit.]. $\square$

In fact, η corresponds to some class in $H^1(W_F/I_F, Z_{\hat{G}}^{I_F})$. To see this, one readily reduces to the case of a torus S as above. Since $S(F)^1$ contains the parahoric subgroup, one can infer, for example by the Satake isomorphism [Haines and Rostami 2010, Proposition 1.0.2] for S , that we obtain a parameter in $H^1(W_F/I_F, \hat{S}^{I_F})$.

The second lemma concerns the globalization of groups.

Lemma 3.4.2. *Given G and F as above, one can choose*

- $\mathring{F}$: a global field of characteristic p ;
- $\mathring{G}$: a connected reductive $\mathring{F}$ -group with maximal $\mathring{F}$ -torus $\mathring{T}$, sharing the same splitting field $\mathring{F}|\mathring{F}$;
- v : a place of $\mathring{F}$, and w is the unique place of $\tilde{F}$ lying over v , in particular $\text{Gal}(\tilde{F}|\mathring{F})$ equals the decomposition group $\Gamma_w := \text{Gal}(\tilde{F}_w|\mathring{F}_v)$;

such that

- there exist isomorphisms $\mathring{F}_v \simeq F$, $\tilde{F}_w \simeq \tilde{F}$, which identify $\Gamma := \text{Gal}(\tilde{F}|F)$ with Γ_w ;
- under the identifications above, there is an isomorphism

$$\begin{array}{ccc} \mathring{G}_{\mathring{F}_v} & \xrightarrow{\sim} & G \\ \cup & & \cup \\ \mathring{T}_{\mathring{F}_v} & \xrightarrow{\sim} & T \end{array},$$

i.e., $\mathring{G} \supset \mathring{T}$ is an $\mathring{F}$ -model of $G \supset T$;

- $\mathring{G}$ and G share the same root datum endowed with actions of $\Gamma \simeq \Gamma_w$, relative to $\mathring{T}$ and T respectively.

Proof. Standard. See for instance [Arthur 1988, p.526] or [Vignéras 2001, 3.12]. $\square$

Remark 3.4.3. The matching of root data in Lemma 3.4.2 also implies that $A_{\mathring{G}}$ is “the same” as A_G . Hereafter, we shall drop the clumsy notation $\mathring{G}$, $\mathring{T}$ or $A_{\mathring{G}}$, and denote them abusively as G , T or A_G instead.

For any closed discrete subgroup $\Xi \subset A_G(F)$ isomorphic to $\mathbb{Z}^{\dim A_G}$, its isomorphic image in $A_G(\mathring{F}) \backslash A_G(\mathbb{A})$ will also be denoted by Ξ . Another consequence of Lemma 3.4.2 is that Ξ is a cocompact lattice in $A_G(\mathring{F}) \backslash A_G(\mathbb{A})$ satisfying the requirements in Section 2.1.

Proof of Theorem 3.2.2 from Theorem 3.3.2. In what follows, we write $\pi \rightsquigarrow \phi$ if $\pi \in \Pi(G)$ has Genestier–Lafforgue parameter $\phi \in \Phi(G)$. It suffices to show that for every $\pi \in \Pi(G)$,

$$(\pi \rightsquigarrow \phi) \implies (\check{\pi} \rightsquigarrow {}^L\theta \circ \phi). \quad (3-3)$$

Indeed, this assertion amounts to $\{\check{\pi} : \pi \in \Pi_\phi\} \subset \Pi_{{}^L\theta \circ \phi}$. The reverse inclusion will follow by applying (3-3) to any $\pi_1 \in \Pi(G)$ with $\pi_1 \rightsquigarrow {}^L\theta \circ \phi$, which in turn yields $\pi := \check{\pi}_1 \rightsquigarrow {}^L\theta \circ {}^L\theta \circ \phi = \phi$ whilst $\pi_1 = \check{\pi}$.

The assertion (3-3) will be established in steps:

Step 1. We reduce (3-3) to the case π supercuspidal. Indeed, let (M, τ) be the cuspidal support of π reviewed in Section 2.3. By Proposition 2.3.1, $\check{\pi}$ has cuspidal support $(M, \check{\tau})$.

On the dual side, choose an L -embedding $\iota : {}^L M \hookrightarrow {}^L G$ as reviewed in Section 2.4. Suppose that $\tau \rightsquigarrow \phi_\tau$ in M . By [Genestier and Lafforgue 2017, Théorème 0.1], ϕ equals to the composite of $W_F \xrightarrow{\phi_\tau} {}^L M \hookrightarrow {}^L G$ up to $\hat{G}(\overline{\mathbb{Q}_\ell})$ -conjugacy. The same relation holds for the parameters for $\check{\pi}$ and $\check{\tau}$. Denoting ${}^L\theta_M$ the

Chevalley involution on ${}^L M$, the diagram

$$\begin{array}{ccc} {}^L M & \xhookrightarrow{\iota} & {}^L G \\ {}^L \theta_M \downarrow & & \downarrow {}^L \theta \\ {}^L M & \xhookrightarrow{\iota} & {}^L G \end{array}$$

is commutative up to an explicit $\hat{G}(\overline{\mathbb{Q}_\ell})$ -conjugacy, by [Prasad 2018, §5, Lemma 4]. Upon replacing (G, π) by (M, τ) , we have reduced (3-3) to the supercuspidal case.

Step 2. Consider the smooth character $\omega := \omega_\pi|_{A_G(F)}$. We reduce (3-3) to the case that ω is of finite order as follows (see also Remark 3.4.4). First, recalling (2-3), there exists a character

$$\eta_0 : A_G(F)/A_G(F)^1 \rightarrow \overline{\mathbb{Q}_\ell}^\times$$

such that $\eta_0 \otimes \omega$ is of finite order. Indeed, this is easily reduced to the case $A_G \simeq \mathbb{G}_m$, and it suffices to take $\eta_0(\varpi) = \omega(\varpi)^{-1}$ where $\varpi \in F^\times$ is some uniformizer.

Secondly, the inclusion of discrete free commutative groups of finite type

$$A_G(F)/A_G(F)^1 = A_G(F)/A_G(F) \cap G(F)^1 \hookrightarrow G(F)/G(F)^1$$

has finite cokernel, whereas $\overline{\mathbb{Q}_\ell}^\times$ is divisible. Therefore η_0 extends to a smooth character

$$\eta : G(F)/G(F)^1 \rightarrow \overline{\mathbb{Q}_\ell}^\times.$$

The central character of $\pi \otimes \eta$ has finite order when restricted to $A_G(F)$.

Attach $a \in H^1(W_F, Z_{\hat{G}})$ to η by Lemma 3.4.1; it can be used to twist elements of $\Phi(G)$ by the homomorphism

$$W_F \ltimes (Z_{\hat{G}} \times \hat{G}) \rightarrow W_F \ltimes \hat{G}, \quad w \ltimes (z, g) \mapsto w \ltimes (zg)$$

by choosing any cocycle representative of a ; see [Genestier and Lafforgue 2017, Remarque 0.2].

In the construction above, $-\otimes \eta^{-1}$ corresponds to twisting a parameter by a^{-1} . We have $(\pi \otimes \eta)^\vee \simeq \tilde{\pi} \otimes \eta^{-1}$. Concurrently, ${}^L \theta \circ (\phi \cdot a) = ({}^L \theta \circ \phi) \cdot a^{-1}$ since Chevalley involution acts as $z \mapsto z^{-1}$ on the center. Therefore, by replacing π by $\pi \otimes \eta$, it suffices to prove (3-3) when ω has finite order.

Step 3. Now we can assume π to be integral supercuspidal (see Section 2.3) with $\omega := \omega_\pi|_{A_G(F)}$ of finite order. By [Genestier and Lafforgue 2017], we know that the parameter ϕ of π factors through $\text{Gal}(\bar{F}|F)$. Take a global $\mathring{F}$ -model of $G \supset A_G$ as in Lemma 3.4.2 with $\mathring{F}_v \simeq F$. As A_G is split over $\mathring{F}$, by reducing to $\mathbb{G}_m$ and applying [Artin and Tate 1968, Chapter X, §2, Theorem 5], there exists an automorphic character

$$\dot{\omega} = \bigotimes_u \dot{\omega}_u : A_G(\mathring{F}) \backslash A_G(\mathbb{A}) \rightarrow \overline{\mathbb{Q}_\ell}^\times$$

of finite order, such that $\dot{\omega}_v = \omega$.

Since ω is smooth, there exists a closed discrete subgroup $\Xi \subset A_G(F)$ such that $\omega|_\Xi = 1$ and $\Xi \simeq \mathbb{Z}^{\dim A_G}$. In view of Remark 3.4.3, Ξ also affords the cocompact lattice in $A_G(\mathring{F}) \backslash A_G(\mathbb{A})$ required in Section 2.1.

Claim: there exists a cuspidal automorphic representation $\hat{\pi} = \bigotimes_u \hat{\pi}_u$ of $G(\mathbb{A})$ (in the extended sense that we consider all G_α simultaneously, $\alpha \in \ker^1(\hat{F}, G)$) such that:

- The central character of $\hat{\pi}$ equals $\hat{\omega}$ on $A_G(\mathbb{A})$.
- We have $\hat{\pi}_v \simeq \pi$.
- Relative to the chosen lattice Ξ and a sufficiently deep level N , the $C_c(K_N \backslash G(\mathbb{A})/K_N; \overline{\mathbb{Q}_\ell})$ -module $\hat{\pi}^{K_N}$ can be embedded in some summand $\mathfrak{H}_\sigma$ in (3-2).

This can be achieved by the following variant of the argument in [Henniart 1983, Appendice 1] (which works over $\mathbb{C}$) via Poincaré series; see also the proof of [Genestier and Lafforgue 2017, Lemme 1.4]. For each place u of $\hat{F}$, choose a smooth function $f_u \in C_c(G(\hat{F}_u), \hat{\omega}_u)$ such that:

- There exists a finite set S of places of $\hat{F}$ containing v and the ramification locus of G , such that when $u \notin S$, the function f_u is right $G(\mathfrak{o}_u)$ -invariant, supported on $A_G(\hat{F}_u)G(\mathfrak{o}_u)$ and $f_u(1) = 1$, where $G(\mathfrak{o}_u)$ is the hyperspecial subgroup arising from some reductive model of G over the ring of S -integers in $\hat{F}$.
- We require f_v to a matrix coefficient of π and assume $f_v(1) \neq 0$.
- For every $u \in S \setminus \{v\}$, we require that $C_u := \text{Supp}|f_u|$ is a sufficiently small neighborhood of 1 modulo $A_G(F_u)$, so that the image of

$$\text{Supp}(f_v) \times \prod_{u \in S \setminus \{v\}} C_u \times \prod_{u \notin S} G(\mathfrak{o}_u)A_G(\hat{F}_u)$$

in $A_G(\mathbb{A}) \backslash G(\mathbb{A}) = (A_G \backslash G)(\mathbb{A})$ intersects $A_G(\hat{F}) \backslash G(\hat{F}) = (A_G \backslash G)(\hat{F})$ only at 1. To see why this can be achieved, embed $A_G \backslash G$ into some affine space over F .

Take $f := \prod_u f_u : G(\mathbb{A}) \rightarrow \overline{\mathbb{Q}_\ell}$ and form

$$P_f(g) = \sum_{\gamma \in (A_G \backslash G)(\hat{F})} f(\gamma g), \quad g \in G(\mathbb{A}).$$

The sum is finite when g is constrained in any compact subset modulo $A_G(\mathbb{A})$. By choosing N sufficiently deep, it furnishes an element of $C_c(G(\hat{F}) \backslash G(\mathbb{A})/K_N \Xi; \overline{\mathbb{Q}_\ell})$. Moreover, $P_f(1) = f(1) \neq 0$ by the condition on supports. By looking at f_v , we see that P_f is a cusp form.

Decompose $C_c^{\text{cusp}}(\text{Bun}_{G,N}(\mathbb{F}_q)/\Xi; \overline{\mathbb{Q}_\ell})$ into simple submodules as in Remark 3.3.1. There exists a summand $\mathcal{L}$ contained in some $\mathfrak{H}_\sigma$ such that P_f has nonzero component in $\mathcal{L}$. Let $\hat{\pi}$ be the cuspidal automorphic representation corresponding to $\mathcal{L}$ via Proposition 2.1.1 (realized in $\bigoplus_\alpha C_c^{\text{cusp}}(G_\alpha(\hat{F}) \backslash G(\mathbb{A}) \cdots)$) where $\alpha \in \ker^1(\hat{F}, G)$ so that $\hat{\pi}^{K_N} = \mathcal{L} \hookrightarrow \mathfrak{H}_\sigma$. Then $\hat{\pi}$ has central character $\hat{\omega}$ on $A_G(\mathbb{A})$ and $\hat{\pi}_v \simeq \pi$, since P_f and $\mathcal{L}$ have similar properties under $C_c(K_N \backslash G(\mathbb{A})/K_N; \overline{\mathbb{Q}_\ell})$.

Step 4. Since the integration pairings $\langle \cdot, \cdot \rangle$ of Remark 2.2.3 are nondegenerate, $\hat{\pi}^{K_N} \subset \mathfrak{H}_\sigma$ must pair nontrivially with some simple $C_c(K_N \backslash G(\mathbb{A})/K_N; \overline{\mathbb{Q}_\ell})$ -submodule of some $\mathfrak{H}_{\sigma'}$. Proposition 2.1.1 implies that the simple submodule takes the form $(\hat{\pi}')^{K_N} \subset \mathfrak{H}_{\sigma'}$ for some cuspidal automorphic representation $\hat{\pi}'$.

Theorem 3.3.2 then asserts $\sigma' = {}^L\theta \circ \sigma$ in $\Phi(G)$ (global version). On the other hand, $\check{\pi}'$ pairs nontrivially with $\check{\pi}$ under the integration pairing $\langle \cdot, \cdot \rangle$ of Definition 2.2.1. The invariance of $\langle \cdot, \cdot \rangle$ therefore implies that, as $G(\mathbb{A})$ -representations,

$$\bigotimes_u (\check{\pi}_u)^\vee = \check{\pi}^\vee \simeq \check{\pi}'.$$

The local-global compatibility in [Genestier and Lafforgue 2017, Théorème 0.1(b)] says that

$$\begin{aligned} \pi &\simeq \check{\pi}_v \rightsquigarrow (\sigma|_{\text{Gal}(\bar{F}|F)})^{\text{ss}}, \\ \check{\pi} &\simeq (\check{\pi}_v)^\vee \simeq \check{\pi}'_v \rightsquigarrow (\sigma'|_{\text{Gal}(\bar{F}|F)})^{\text{ss}} = ({}^L\theta \circ \sigma|_{\text{Gal}(\bar{F}|F)})^{\text{ss}}. \end{aligned}$$

Here we choose an embedding of the separable closure of $\hat{F}$ into $\bar{F}$, and the semisimplification is defined as in Section 2.4. In particular, $\phi = (\sigma|_{\text{Gal}(\bar{F}|F)})^{\text{ss}}$ in $\Phi(G)$ (local version).

By Lemma 2.4.5 we have $({}^L\theta \circ \sigma|_{\text{Gal}(\bar{F}|F)})^{\text{ss}} = {}^L\theta \circ (\sigma|_{\text{Gal}(\bar{F}|F)})^{\text{ss}}$. Summarizing,

$$\check{\pi} \rightsquigarrow {}^L\theta \circ (\sigma|_{\text{Gal}(\bar{F}|F)})^{\text{ss}} = {}^L\theta \circ \phi$$

holds in $\Phi(G)$ (local version). This establishes (3-3) and the Theorem 3.2.2 follows. $\square$

Remark 3.4.4. As pointed out by a referee, Lemma 3.4.1 can be avoided in Step 2 by the following arguments. Restrict $\text{quot} : G \rightarrow T := G/G_{\text{der}}$ to an isogeny $A_G \rightarrow T$. The same arguments show that some smooth character $\eta : T(F) \rightarrow \overline{\mathbb{Q}_\ell}^\times$ pulls back to our given $\eta_0 : A_G(F) \rightarrow \overline{\mathbb{Q}_\ell}^\times$. To complete Step 2, it remains to compare (a) the parameters of η and η^{-1} and (b) the parameters of π and $\pi \otimes \eta$. For (a), apply local trivial functoriality [Genestier and Lafforgue 2017, Théorème 8.1] to the automorphism $t \mapsto t^{-1}$ of T . For (b), apply it to the homomorphism $G \xrightarrow{(\text{id}, \text{quot})} G \times T$ with normal image, as performed in [Genestier and Lafforgue 2017, Remarque 0.3].

From Section 4 onwards, we will focus exclusively on Theorem 3.3.2 and the underlying geometric considerations.

3.5. Remarks on the duality involution. Conserve the assumptions for the local statement in Section 3.2 and assume G is quasisplit. Fix an F -pinning $\mathcal{P} = (B, T, (X_\alpha)_\alpha)$ of G . Choose the unique $\kappa \in T^{\text{ad}}(F)$ such that $\kappa X_\alpha \kappa^{-1} = -X_\alpha$, for all simple root α with respect to (B, T) . Observe that $\kappa^2 = 1$ in G^{ad} .

Let $\theta = \theta_{\mathcal{P}}$ be the Chevalley involution of G , and ι_- be the inner involution $g \mapsto \kappa g \kappa^{-1}$. Observe that $\iota_- \theta = \theta \iota_-$. Indeed, $\iota_- \theta \iota_-$ is seen to preserve $\mathcal{P}$ and coincides with θ on T , hence $\iota_- \theta \iota_- = \theta$ by the characterization of the Chevalley involution.

Definition 3.5.1 [Prasad 2018, §3]. Relative to the F -pinning $\mathcal{P}$, set $\iota_{G, \mathcal{P}} := \iota_- \theta = \theta \iota_-$. It is called the *duality involution* of G .

Recall that $\iota_{G, \mathcal{P}}$ induces a pinned automorphism of $\hat{G}$, called the *dual* automorphism of $\iota_{G, \mathcal{P}}$, which depends only on $\iota_{G, \mathcal{P}}$ modulo $G^{\text{ad}}(F)$; see [Borel 1979, §2.5] for the general set-up. This recipe applies to any base field F .

Lemma 3.5.2. *The Chevalley involution on $\hat{G}$ is the dual of $\iota_{G, \mathcal{P}}$ in the sense above. This result holds over any field F .*

Proof. Since ι_- comes from $G^{\text{ad}}(F)$ -action, $\iota_{G, \mathcal{P}}$ and θ have the same dual. It suffices to show that the Chevalley involution of $\hat{G}$ is dual to that of G . Since both automorphisms are pinned, it suffices to show that the induced automorphisms on $X_*(T_{\bar{F}})$ and $X^*(T_{\bar{F}})$ are mutually dual. Recall that the Chevalley involution of G and $\hat{G}$ act on $X_*(T_{\bar{F}})$ and $X^*(T_{\bar{F}})$, respectively, as $x \mapsto -w_0(x)$, where w_0 is the longest element in the Weyl group. Since $w_0^2 = 1$, these two automorphisms are indeed mutually dual. $\square$

Fix a nontrivial smooth character $\psi : F \rightarrow \overline{\mathbb{Q}_\ell}^\times$. From the F -pinning $\mathcal{P} = (B, T, (X_\alpha)_\alpha)$ we produce a Whittaker datum $\mathfrak{w} := (U, \chi)$ for G taken up to $G(F)$ -conjugacy, that is,

- U is the unipotent radical of B ,
- $\chi : U(F) \rightarrow \overline{\mathbb{Q}_\ell}^\times$ is the composition of ψ with the algebraic character $U \rightarrow \mathbb{G}_a$ mapping each X_α to 1.

The automorphisms of G act on F -pinnings, thereby act on Whittaker data. Put

$$\mathfrak{w}' := (U, \chi^{-1}) = \iota_- \mathfrak{w}.$$

Fix ψ , $\mathcal{P}$ and the associated Whittaker datum $\mathfrak{w}$ for G . Let $\phi \in \Phi(G)$ be a semisimple parameter. Define the Genestier–Lafforgue packet Π_ϕ as in Section 3.2. We say that *Shahidi’s property* holds for Π_ϕ and $\mathfrak{w}$, if

$$\exists! \pi \in \Pi_\phi \quad \text{such that } \pi \text{ is } \mathfrak{w}\text{-generic.} \quad (3-4)$$

Further discussions about this property will be given in Remark 3.5.5.

Lemma 3.5.3. *The following are equivalent for an irreducible smooth representation π of $G(F)$:*

- (i) π is $\mathfrak{w}$ -generic.
- (ii) $\pi \circ \theta$ is $\mathfrak{w}$ -generic.
- (iii) $\check{\pi}$ is $\mathfrak{w}'$ -generic.
- (iv) $\pi \circ \iota_-$ is $\mathfrak{w}'$ -generic.

Proof. (i) $\iff$ (ii) since θ preserves $\mathcal{P}$. (i) $\iff$ (iii) is [Prasad 2018, §4, Lemma 2]. (i) $\iff$ (iv) follows from transport of structure by the involution ι_- . $\square$

The following result serves as a partial heuristic for [Prasad 2018, §3, Conjecture 1].

Theorem 3.5.4. *Define the Whittaker data $\mathfrak{w}$ and $\mathfrak{w}'$ as above. Let $\phi \in \Phi(G)$ be a semisimple parameter such that Π_ϕ satisfies Shahidi’s property (3-4) with respect to $\mathfrak{w}$. Then the following hold:*

- (i) *The packet $\Pi_{L_{\theta \circ \phi}}$ satisfies Shahidi’s property (3-4) with respect to $\mathfrak{w}'$.*
- (ii) *Let π be the unique $\mathfrak{w}$ -generic member of Π_ϕ , then $\check{\pi}$ is the unique $\mathfrak{w}'$ -generic member of $\Pi_{L_{\theta \circ \phi}}$.*
- (iii) *If $\pi \in \Pi_\phi$ is $\mathfrak{w}$ -generic, then $\check{\pi} \simeq \pi \circ \iota_{G, \mathcal{P}}$.*

Proof. Parts (i), (ii) follow immediately from Lemma 3.5.3 and Theorem 3.2.2, which says that $\Pi_{\mathbf{L}_{\theta \circ \phi}} = \{\tilde{\pi} : \pi \in \Pi_{\phi}\}$.

Now consider (iii). We claim that $\pi \circ \iota_{G, \mathcal{P}} \in \Pi_{\mathbf{L}_{\theta \circ \phi}}$. In view of Lemma 3.5.2, the corresponding statement for global Langlands parametrization of cuspidal automorphic representations follows from the “trivial functoriality” (under the dual of $\iota_{G, \mathcal{P}}$) in [Genestier and Lafforgue 2017, Théorème 0.1 and 8.1].

Using Lemma 3.5.3, we see that $\pi \circ \iota_{G, \mathcal{P}} = (\pi \circ \theta) \circ \iota_{-}$ is also a $\mathfrak{w}'$ -generic member of $\Pi_{\mathbf{L}_{\theta \circ \phi}}$. It follows from (ii) that $\tilde{\pi} \simeq \pi \circ \iota_{G, \mathcal{P}}$. $\square$

Remark 3.5.5. Choose an isomorphism $\overline{\mathbb{Q}_{\ell}} \xrightarrow{\sim} \mathbb{C}$ and let $\phi \in \Phi(G)$ be semisimple. By a conjecture of Shahidi [1990, Conjecture 9.4], one expects that when ϕ is a tempered L -parameter, (3-4) will hold for the *authentic L -packet* associated to Π_{ϕ} and for any $\mathfrak{w}$.

On the other hand, [Gross and Prasad 1992, Conjecture 2.6] proposes a characterization of L -parameters satisfying (3-4). It is stated in terms of adjoint L -factors, thus applies directly to the ℓ -adic case. The author is grateful to Yeansu Kim for this comment.

Because of the semisimplified nature of our packet Π_{ϕ} , see Remark 3.2.1, we expect (3-4) to hold only when ϕ is not the semisimplification of any other L -parameter. This occurs when ϕ is elliptic, in which case every $\pi \in \Pi_{\phi}$ is supercuspidal: otherwise the compatibility of the parametrization $\pi \rightsquigarrow \phi$ with cuspidal supports will force ϕ to factor through some proper Levi. It is believed that the authentic L -packets for elliptic ϕ have the same property. Many constructions of such L -packets have been proposed, such as in [Kaletha 2016a]. Nonetheless, the precise relation of these packets to the Langlands parametrization of Genestier and Lafforgue [2017] remains to be settled.

Remark 3.5.6. As shown in [Prasad 2018], up to $G(F)$ -conjugacy, $\iota_{G, \mathcal{P}}$ reduces to the well-known MVW involution when G is classical; it reduces to $g \mapsto {}^t g^{-1}$ when $G = \mathrm{GL}(n)$. According to [Prasad 2018, §3, Corollary 1], when Z_G is an elementary 2-group, $\iota_{G, \mathcal{P}}$ is independent of $\mathcal{P}$ up to $G(F)$ -conjugacy.

4. Overview of the global Langlands parametrization

4.1. Geometric setup. Fix some power q of a prime number p . Take $E \subset \overline{\mathbb{Q}_{\ell}}$ to be a finite extension of $\mathbb{Q}_{\ell}$ containing a square root $q^{1/2}$ of q , which we fix once and for all. The sheaves and complexes under consideration will be E -linear.

Suppose that S is a smooth $\mathbb{F}_q$ -scheme of finite type and of pure dimension d . For any reasonable algebraic stack $\mathcal{X}$ equipped with a morphism $\mathfrak{p} : \mathcal{X} \rightarrow S$, define the *normalized perverse sheaves* on $\mathcal{X}$ with respect to S to be of the form

$$\mathcal{F}[-d]\left(-\frac{d}{2}\right), \quad \mathcal{F} \text{ a nonnormalized perverse sheaf.}$$

The usual operations on constructible complexes continue to hold in the normalized setting, with the proviso that the dualizing complex in [Laszlo and Olsson 2008b, §7.3] becomes

$$\Omega_{\mathcal{X}} := (\text{the nonnormalized one})[-2d](-d) \simeq \mathfrak{p}^!(E_S)$$

and the duality operator becomes $\mathbb{D} = R\mathcal{H}om(-, \Omega_{\mathcal{X}})$ accordingly. This formalism extends to ind-stacks, etc. with a morphism to S . When $S = \operatorname{Spec} \mathbb{F}_q$, we revert to the usual definitions.

Next, assume $\mathring{F} = \mathbb{F}_q(X)$ is a global field and G is a connected reductive $\mathring{F}$ -group with a chosen Bruhat–Tits model over X , as in Section 2.1. Fix a maximal $\mathring{F}$ -torus $T \subset G$. Also recall that $\hat{G}$ carries a Galois-stable pinning $(\hat{B}, \hat{T}, (X_\alpha)_\alpha)$. Enlarging E if necessary, we can assume that:

All irreducible $\overline{\mathbb{Q}_\ell}$ -representations of ${}^L G$ are realized over E .

Fix a partition of a finite set

$$I = I_1 \sqcup \cdots \sqcup I_k$$

used to label points on X and a level $N \subset X$. Set $\hat{N} = |N| \cup (X \setminus U)$ as in (3-1). Define the *Hecke stack* $\operatorname{Hecke}_{N,I}^{(I_1, \dots, I_k)}$ that maps each $\mathbb{F}_q$ -scheme S to the groupoid

$$\operatorname{Hecke}_{N,I}^{(I_1, \dots, I_k)}(S) = \left\{ \begin{array}{l} (x_i)_i \in (X \setminus \hat{N})(S)^I, \\ ((\mathcal{G}_j, \psi_j) \in \operatorname{Bun}_{G,N}(S))_{j=0}^k, \\ \phi_j : \mathcal{G}_{j-1} \dashrightarrow \mathcal{G}_j \end{array} \middle| \begin{array}{l} \phi_j \text{ defined off } \bigcup_{i \in I_j} \Gamma_{x_i}, \\ \psi_j \phi_j|_{N \times S} = \psi_{j-1} \\ \forall j = 1, \dots, k \end{array} \right\} \quad (4-1)$$

where Γ_{x_i} stands for the graph of $x_i : S \rightarrow X$. The points $(x_i)_{i \in I}$ are known as the “paws”.

The reason for partitioning I into $I_1, \dots, I_k$ is to define *partial Frobenius morphisms*, see Section 4.3.

The ind-scheme $\operatorname{Gr}_I^{(I_1, \dots, I_k)}$, the factorization version of affine Grassmannian of Beilinson–Drinfeld, is the space classifying the same data (4-1) as $\operatorname{Hecke}_{I, \emptyset}^{(I_1, \dots, I_k)}$ together with a trivialization θ of $\mathcal{G}_k$. It also admits a morphism of “paws” to X^I . In fact $\operatorname{Gr}_I^{(I_1, \dots, I_k)}$ is ind-projective; we refer to [Lafforgue 2018, §1] for further details. When I is a singleton and $k = 1$, the usual Beilinson–Drinfeld Grassmannian over X is recovered.

The *factorization structure* here means that given a surjection $\zeta : I \rightarrow J$, we have, for $U_\zeta := \{(x_i)_{i \in I} : \zeta(a) \neq \zeta(b) \implies x_a \neq x_b\} \subset X^I$ and $I'_a := I_a \cap \zeta^{-1}(j)$, $\forall j$, the canonical isomorphism

$$\operatorname{Gr}_I^{(I_1, \dots, I_k)} \times_{X^I} U_\zeta \xrightarrow{\sim} \prod_{j \in J} \operatorname{Gr}_{\zeta^{-1}(j)}^{(I'_1, \dots, I'_k)}$$

over U_ζ see [Lafforgue 2018, Remarque 1.9]. The factorization structure is mainly to be employed together with the complexes that are *universally locally acyclic*, hereafter abbreviated as *ULA*, with respect to the base (say X^I). This property (see [Richarz 2014, §3.2] or [Braverman and Gaiitsgory 2002, §5.1]) is immensely useful for “spreading out” certain properties of complexes from some open subset in the base, see e.g., [Richarz 2014, Theorem 3.16].

Given $(n_i)_i \in \mathbb{Z}_{\geq 0}^I$. Define $\Gamma_{\sum_i n_i x_i} \subset X \times X^I$ as the closed subscheme Zariski-locally defined by $\prod_{i \in I} t_i^{n_i}$, with t_i being a local equation for x_i in X , where $(x_i)_{i \in I}$ are the aforementioned “paws”. Then define

$$G_{\sum_{i \in I} n_i x_i} := \text{the Weil restriction of } G \text{ with respect to } \Gamma_{\sum_i n_i x_i} \rightarrow X^I.$$

One interprets $G_{\sum_i \infty x_i}$ in the same manner by considering formal neighborhoods, but we won't go into the details.

As in the discussion preceding [Lafforgue 2018, Proposition 1.10], there is a notion of $G_{\sum_i \infty x_i}$ -action on $\mathrm{Gr}_I^{(I_1, \dots, I_k)}$, namely by altering the trivialization θ of $\mathcal{G}_k$ at $\Gamma_{\sum_i \infty x_i}$.

Let $\mathrm{Perv}_{G_{\sum_i \infty x_i}}(\mathrm{Gr}_I^{(I_1, \dots, I_k)})$ denote the category of $G_{\sum_i \infty x_i}$ -equivariant normalized perverse sheaves on the ind-scheme $\mathrm{Gr}_I^{(I_1, \dots, I_k)}$ relative to X^I ; for nonsplit G , we confine ourselves to $(X \setminus \hat{N})^I$ as in [Lafforgue 2018, §12.3.1]. The factorization version of *geometric Satake equivalence* [Lafforgue 2018, Théorèmes 1.17 and 12.16] gives an additive functor

$$\begin{aligned} \mathrm{Rep}_E({}^L G^I) &\rightarrow \mathrm{Perv}_{G_{\sum_i \infty x_i}}(\mathrm{Gr}_I^{(I_1, \dots, I_k)}) \\ W &\mapsto \mathcal{S}_{I, W, E}^{(I_1, \dots, I_k)}. \end{aligned}$$

For later reference, we record some of the basic properties of this functor, all of which can be found in [loc. cit.]:

- (1) The normalized perverse sheaves $\mathcal{S}_{I, W, E}^{(I_1, \dots, I_k)}$ are ULA relative to the morphism to X^I (or $(X \setminus \hat{N})^I$).
- (2) When $|I| = 1$, the geometric Satake equivalence [Richarz 2014; Zhu 2015] yields $W \mapsto \mathcal{S}_{I, W, E}^{(I)}$. This extends to general I and “factorizable” W using the factorization structure on affine Grassmannians, see [Lafforgue 2018]. Namely, for any family $(W_i)_{i \in I}$ of objects in $\mathrm{Rep}_E({}^L G)$, one can associate $\mathcal{S}_{I, \boxtimes_i W_i, E}^{(I_1, \dots, I_k)}$ in $\mathrm{Perv}_{G_{\sum_i \infty x_i}}(\mathrm{Gr}_I^{(I_1, \dots, I_k)})$.
- (3) Write ${}^L G^I$ for $({}^L G)^I$. In order to obtain a functorial construction in all $W \in \mathrm{Rep}_E({}^L G^I)$, we take the ${}^L G^I \times {}^L G^I$ -representation $\mathcal{R} := \boxtimes_{i \in I} \mathcal{O}({}^L G)$ over E . This becomes an ind-object of $\mathrm{Rep}_E({}^L G^I)$ using the ${}^L G^I$ -action on the first slot, and this ind-object carries a ${}^L G^I$ -action from the second slot. Take a system of representatives of irreducible objects $V \in \mathrm{Rep}_E({}^L G)$. As ${}^L G^I \times {}^L G^I$ -representations, we have

$$\bigoplus_{V: \text{irred}} V \otimes_E V^\vee \xrightarrow{\sim} \mathcal{R} \quad \text{by taking matrix coefficients.}$$

The decomposition above and the available $\mathcal{S}_{I, V, E}^{(I_1, \dots, I_k)}$ define a normalized ind-perverse sheaf $\mathcal{S}_{I, \mathcal{R}, E}^{(I_1, \dots, I_k)}$, with the $\hat{G}^I$ -action inherited from the second slot of $\mathcal{R}$. Now we can define, for each $W \in \mathrm{Rep}_E({}^L G^I)$,

$$\mathcal{S}_{I, W, E}^{(I_1, \dots, I_k)} := (\mathcal{S}_{I, \mathcal{R}, E}^{(I_1, \dots, I_k)} \otimes_E W)^{{}^L G^I} \simeq \bigoplus_{V: \text{irred}} \mathcal{S}_{I, V, E}^{(I_1, \dots, I_k)} \otimes_E (V^\vee \otimes_E W)^{{}^L G^I} \simeq \bigoplus_{V: \text{irred}} \mathcal{S}_{I, V, E}^{(I_1, \dots, I_k)} \otimes_E \mathfrak{W}_V, \quad (4-2)$$

where: (a) W is viewed as a constant sheaf on $\mathrm{Gr}_I^{(I_1, \dots, I_k)}$. (b) ${}^L G^I$ acts diagonally. (c) $\mathfrak{W}_V$ stands for the multiplicity space of V in W . Functoriality in W is clear, and it is readily seen to agree with the previous step if $W = V$, up to isomorphism.

Given $W \in \mathrm{Rep}_E({}^L G^I)$, we define the reduced closed subscheme

$$\mathrm{Gr}_{I, W}^{(I_1, \dots, I_k)} := \mathrm{Supp} \mathcal{S}_{I, W, E}^{(I_1, \dots, I_k)} \subset \mathrm{Gr}_I^{(I_1, \dots, I_k)}.$$

In this manner, the objects of $\mathrm{Rep}_E({}^L G^I)$ will serve as truncation parameters for $\mathrm{Gr}_{I, W}^{(I_1, \dots, I_k)}$. For the traditional definition in terms of weights and relative positions, see [Lafforgue 2018, Définition 1.12].

When $|I| = 1$ and W is irreducible, $\mathcal{S}_{I,W,E}^{(I)}$ is well known to be isomorphic to the normalized IC-complex of the stratum $\mathrm{Gr}_{I,W}^{(I)}$.

We move to the *moduli stack of chtoucas* with level structures, whose the details can be found in [Lafforgue 2018, §2, §12.3.2]. For $I = I_1 \sqcup \cdots \sqcup I_k$ and N as before, $\mathrm{Cht}_{N,I}^{(I_1, \dots, I_k)}$ is defined by a pull-back diagram

$$\begin{array}{ccc} \mathrm{Cht}_{N,I}^{(I_1, \dots, I_k)} & \longrightarrow & \mathrm{Hecke}_{N,I}^{(I_1, \dots, I_k)} \\ \downarrow & \square & \downarrow (\mathcal{G}_0, \mathcal{G}_k) \\ \mathrm{Bun}_{G,N} & \xrightarrow{\mathrm{id} \times \mathrm{Frob}} & \mathrm{Bun}_{G,N} \times \mathrm{Bun}_{G,N} \end{array}$$

of ind-stacks over $\mathbb{F}_q$. It classifies the chains

$$(\mathcal{G}_0, \psi_0) \xrightarrow{\phi_1} \cdots \xrightarrow{\phi_{k-1}} (\mathcal{G}_k, \psi_k) \xrightarrow{\phi_k} ({}^\tau \mathcal{G}_0, {}^\tau \psi_0)$$

of G -torsors with N -level structures (see (4-1)). Here, for every $\mathbb{F}_q$ -scheme S and $(\mathcal{G}, \psi) \in \mathrm{Bun}_{G,N}(S)$ we set

$$({}^\tau \mathcal{G}, {}^\tau \psi) := (\mathrm{id}_X \times \mathrm{Frob}_S)^*(\mathcal{G}, \psi)$$

and similarly for the morphisms in $\mathrm{Bun}_{G,N}(S)$. Note that $\mathrm{Cht}_{N,I}^{(I_1, \dots, I_k)}$ is an ind-stack of ind-finite type over $\mathbb{F}_q$ endowed with a morphism of “paws”

$$\mathfrak{p}_{N,I}^{(I_1, \dots, I_k)} : \mathrm{Cht}_{N,I}^{(I_1, \dots, I_k)} \rightarrow (X \setminus \hat{N})^I$$

coming from that of $\mathrm{Hecke}_{N,I}^{(I_1, \dots, I_k)}$. Stability conditions of Harder–Narasimhan type attached to dominant coweights $\mu \in X_*(T^{\mathrm{ad}})$ of G^{ad} on the datum $\mathcal{G}_0$ gives rise to the truncated piece $\mathrm{Cht}_{N,I}^{(I_1, \dots, I_k), \leq \mu}$. Choose any Borel subgroup (over the separable closure) of G containing T . For coweights μ, μ' , write

$$\mu' \geq \mu \iff \mu' - \mu \in \sum_{\check{\alpha}: \text{simple coroot}} \mathbb{Q}_{\geq 0} \cdot \check{\alpha}.$$

As μ grows with respect to $\geq$, we have the filtered limit

$$\mathrm{Cht}_{N,I}^{(I_1, \dots, I_k)} = \varinjlim_{\mu} \mathrm{Cht}_{N,I}^{(I_1, \dots, I_k), \leq \mu}.$$

Exactly as in the case of affine Grassmannians, there is another truncation indexed by $W \in \mathrm{Rep}_E(({}^L G)^I)$; see [Lafforgue 2018, §2] for details. They give rise to

$$\mathrm{Cht}_{N,I,W}^{(I_1, \dots, I_k)} \supset^{\mathrm{open}} \mathrm{Cht}_{N,I,W}^{(I_1, \dots, I_k), \leq \mu}.$$

By [Lafforgue 2018, Proposition 2.6], $\mathrm{Cht}_{N,I,W}^{(I_1, \dots, I_k)}$ is a reduced Deligne–Mumford stack locally of finite type over $(X \setminus \hat{N})^I$, for any W . The connected components of an open substack of the form $\mathrm{Cht}_{N,I,W}^{(I_1, \dots, I_k), \leq \mu}$ are quotients of quasiprojective $(X \setminus \hat{N})^I$ -schemes by finite groups; when N is large relative to μ and to the highest weights of W , those connected components are even quasiprojective $(X \setminus \hat{N})^I$ -schemes. The

last property can serve to justify some geometric reasoning over such stacks, by reducing them to the usual scheme-theoretic setting.

We have $Z_G(\mathring{F}) \backslash Z_G(\mathbb{A}) \hookrightarrow \text{Bun}_{Z_G, N}(\mathbb{F}_q)$, and the latter acts on $\text{Cht}_{N, I}^{(I_1, \dots, I_k)}$ by twisting G -torsors by Z_G -torsors. This action leaves each truncated piece invariant. In particular, for a lattice $\Xi \subset Z_G(\mathring{F}) \backslash Z_G(\mathbb{A})$ chosen as in Section 2.1, we have Ξ -action on $\text{Cht}_{N, I, W}^{(I_1, \dots, I_k), \leq \mu}$, etc. One can shrink Ξ to make it act freely, and consider the quotients $\text{Cht}_{N, I, W}^{(I_1, \dots, I_k), \leq \mu} / \Xi$, etc.

By the discussions before [Lafforgue 2018, Définition 2.14], $\text{Cht}_{N, I, W}^{(I_1, \dots, I_k), \leq \mu} / \Xi$ is a Deligne–Mumford stack of finite type.

4.2. Cohomologies. We keep the notation from Section 4.1. In what follows, normalization of perverse sheaves will always be with respect to the base $(X \setminus \hat{N})^I$.

The first ingredient [Lafforgue 2018, Proposition 2.8] is a canonical smooth morphism

$$\epsilon_{(I), W, \underline{n}}^{(I_1, \dots, I_k)} : \text{Cht}_{N, I, W}^{(I_1, \dots, I_k)} \rightarrow \text{Gr}_{I, W}^{(I_1, \dots, I_k)} / G_{\sum_{i \in I} n_i x_i}$$

where $\underline{n} = (n_i)_{i \in I} \in \mathbb{Z}_{\geq 0}^I$ is sufficiently positive with respect to $W \in \text{Rep}_E(({}^L G)^I)$, so that the $G_{\sum_i \infty x_i}$ -action factors through $G_{\sum_i n_i x_i}$. Assume furthermore that $W = \boxtimes_{j=1}^k W_j$ where each $W_j \in \text{Rep}_E(({}^L G)^{I_j})$ is irreducible. In [Lafforgue 2018, (2.5)] the canonical smooth morphism

$$\epsilon_{(I_1, \dots, I_k), W, \underline{n}}^{(I_1, \dots, I_k)} : \text{Cht}_{N, I, W}^{(I_1, \dots, I_k)} \rightarrow \prod_{j=1}^k \text{Gr}_{I_j, W_j}^{(I_j)} / G_{\sum_{i \in I_j} n_i x_i}$$

is constructed. These two are related by the canonical smooth morphism [Lafforgue 2018, (1.12)]

$$\kappa_{I, W}^{(I_1, \dots, I_k)} : \text{Gr}_{I, W}^{(I_1, \dots, I_k)} \rightarrow \prod_{j=1}^k \text{Gr}_{I_j, W_j}^{(I_j)} / G_{\sum_{i \in I_j} n_i x_i}$$

that chops a chain $\mathcal{G}_0 \dashrightarrow \mathcal{G}_1 \dashrightarrow \dots \dashrightarrow \mathcal{G}_k$ (the trivialization forgotten) classified by $\text{Gr}_I^{(I_1, \dots, I_k)}$ into segments indexed by I_j . By [Lafforgue 2018, (1.13)], when $m_i \gg n_i$ it factorizes through a smooth

$$\tilde{\kappa}_{I, W}^{(I_1, \dots, I_k)} : \text{Gr}_{I, W}^{(I_1, \dots, I_k)} / G_{\sum_{i \in I} m_i x_i} \rightarrow \prod_{j=1}^k \text{Gr}_{I_j, W_j}^{(I_j)} / G_{\sum_{i \in I_j} n_i x_i}.$$

For an interesting result on *local models* of $\text{Cht}_{N, I, W}^{(I_1, \dots, I_k)}$ based on these morphisms, see [Lafforgue 2018, Proposition 2.11]. However, we do not need that result in this article.

As an application, for each $W \in \text{Rep}_E(({}^L G)^I)$ we take the normalized perverse sheaf $\mathcal{S}_{I, W, E}^{(I_1, \dots, I_k)}$ on $\text{Gr}_{I, W}^{(I_1, \dots, I_k)}$. Descend this complex to $\text{Gr}_{I, W}^{(I_1, \dots, I_k)} / G_{\sum_{i \in I} n_i x_i}$ by its equivariance given by geometric Satake. Hence on can form the complex $(\epsilon_{I, W, \underline{n}}^{(I_1, \dots, I_k)})^* \mathcal{S}_{I, W, E}^{(I_1, \dots, I_k)}$.

Since $\mathcal{S}_{I, W, E}^{(I_1, \dots, I_k)}$ is ULA with respect to $(X \setminus \hat{N})^I$, so is its inverse image via the smooth morphism $\epsilon_{I, W, \underline{n}}^{(I_1, \dots, I_k)}$; see [Braverman and Gaitsgory 2002, 5.1.2, item 2]. We claim that the complex $(\epsilon_{I, W, \underline{n}}^{(I_1, \dots, I_k)})^* \mathcal{S}_{I, W, E}^{(I_1, \dots, I_k)}$ is moreover normalized perverse on $\text{Cht}_{N, I, W}^{(I_1, \dots, I_k)}$ for irreducible $W = \boxtimes_{j=1}^k W_j$.

Indeed, the claim is a routine consequence of the factorization structure on $\mathrm{Gr}_{I,W}^{(I_1, \dots, I_k)}$ and the ULA property, smoothness, etc.

This completes our construction when G is semisimple. In general, one has to consider a lattice Ξ as in Section 2.1. According to [Lafforgue 2018, Remarque 1.20], $\mathcal{S}_{I,W,E}^{(I_1, \dots, I_k)}$ descends to $\mathrm{Gr}_{I,W}^{(I_1, \dots, I_k)} / G_{\sum_i n_i x_i}^{\mathrm{ad}}$. By the discussions after [loc. cit., Définition 2.14], $\epsilon_{I,W,\underline{n}}^{(I_1, \dots, I_k)}$ induces

$$\epsilon_{N,I,W,\underline{n}}^{(I_1, \dots, I_k), \Xi} : \mathrm{Cht}_{N,I,W}^{(I_1, \dots, I_k)} / \Xi \rightarrow \mathrm{Gr}_{I,W}^{(I_1, \dots, I_k)} / G_{\sum_i n_i x_i}^{\mathrm{ad}}$$

which is smooth of relative dimension equal to $\dim G_{\sum_i n_i x_i}^{\mathrm{ad}}$. We define accordingly

$$\mathcal{F}_{N,I,W,\Xi,E}^{(I_1, \dots, I_k)} := (\epsilon_{N,I,W,\underline{n}}^{(I_1, \dots, I_k), \Xi})^* \mathcal{S}_{I,W,E}^{(I_1, \dots, I_k)}. \quad (4-3)$$

This is still a normalized perverse sheaf on $\mathrm{Cht}_{N,I,W}^{(I_1, \dots, I_k)} / \Xi$. In [loc. cit.], one actually deduces that $\mathcal{F}_{N,I,W,\Xi,E}^{(I_1, \dots, I_k)}$ is isomorphic to the normalized IC-sheaf on $\mathrm{Cht}_{N,I,W}^{(I_1, \dots, I_k)} / \Xi$.

Thus far we have assumed $W = \boxtimes_{j=1}^k W_j$. A general definition, functorial in arbitrary $W \in \mathrm{Rep}_E(({}^L G)^I)$, can be crafted by repeating the construction for $W \mapsto \mathcal{S}_{I,W,E}^{(I_1, \dots, I_k)}$ reviewed in Section 4.1. The result still takes the form (4-3), except that the right-hand side is now constructed functorially in $W \in \mathrm{Rep}_E(({}^L G)^I)$; see [loc. cit., §4.5].

Next, introduce the other truncation parameter μ from Section 4.1. The morphism of paws induces

$$\mathfrak{p}_{N,I}^{(I_1, \dots, I_k), \leq \mu} : \mathrm{Cht}_{N,I,W}^{(I_1, \dots, I_k), \leq \mu} / \Xi \rightarrow (X \setminus \hat{N})^I.$$

Recall that $\mathrm{Cht}_{N,I,W}^{(I_1, \dots, I_k), \leq \mu}$ is open and Ξ -invariant in $\mathrm{Cht}_{N,I,W}^{(I_1, \dots, I_k)}$. Define

$$\begin{aligned} \mathcal{H}_{N,I,W}^{\leq \mu, E} &:= (\mathfrak{p}_{N,I}^{(I_1, \dots, I_k), \leq \mu})^* \mathcal{F}_{N,I,W,\Xi,E}^{(I_1, \dots, I_k)} \big|_{\mathrm{Cht}_{N,I,W}^{(I_1, \dots, I_k), \leq \mu} / \Xi}, \\ \mathcal{H}_{N,I,W}^{i, \leq \mu, E} &:= \mathrm{H}^i \mathcal{H}_{N,I,W}^{\leq \mu, E}, \end{aligned} \quad (4-4)$$

$i \in \mathbb{Z}$, here H^i is taken with respect to the ordinary t -structure on $\mathrm{D}_c((X \setminus \hat{N})^I, E)$.

- By using the forgetful morphisms as in [Lafforgue 2018, Construction 2.7 and Corollaire 2.18], these complexes are seen to be independent of the partition $(I_1, \dots, I_k)$. The notation in (4-4) is thus justified.
- For $\mu \leq \mu'$, the open immersion $j : \mathrm{Cht}_{N,I,W}^{(I_1, \dots, I_k), \leq \mu} / \Xi \rightarrow \mathrm{Cht}_{N,I,W}^{(I_1, \dots, I_k), \leq \mu'} / \Xi$ induces a canonical arrow $\mathcal{H}_{N,I,W}^{\leq \mu, E} \rightarrow \mathcal{H}_{N,I,W}^{\leq \mu', E}$. This is a standard consequence of the formalism of six operations as $j^* = j^!$.
- They also respect the *coalescence* of paws with respect to any map $\zeta : I \rightarrow J$. We refer to [Lafforgue 2018, Proposition 4.12] for further explanations.

Let I be a finite set and $W \in \mathrm{Rep}_E(({}^L G)^I)$ arbitrary. Denote the generic point of X and X^I by η and η^I , respectively, and choose geometric points over them

$$\bar{\eta} \rightarrow \eta, \quad \bar{\eta}^I \rightarrow \eta^I.$$

Let $\Delta : X \rightarrow X^I$ be the diagonal embedding. Following [Lafforgue 2018, §8] or [Varshavsky 2007, §1.3], we choose an *arrow of specialization*

$$\mathfrak{sp} : \overline{\eta^I} \rightarrow \Delta(\overline{\eta}),$$

i.e., a morphism $(X^I)_{(\overline{\eta^I})} \rightarrow (X^I)_{(\Delta(\overline{\eta}))}$ or equivalently $\overline{\eta^I} \rightarrow (X^I)_{(\Delta(\overline{\eta}))}$, where the subscripts indicate strict Henselizations at the corresponding geometric points. By [Lafforgue 2018, Proposition 8.24], the induced pull-back morphism

$$\mathfrak{sp}^* : \varinjlim_{\mu} \mathcal{H}_{N,I,W}^{0,\leq\mu,E} \big|_{\Delta(\overline{\eta})} \rightarrow \varinjlim_{\mu} \mathcal{H}_{N,I,W}^{0,\leq\mu,E} \big|_{\overline{\eta^I}}$$

between E -vector spaces is injective.

Now comes the Hecke action. Let $f \in C_c(K_N \backslash G(\mathbb{A})/K_N; E)$. According to [Lafforgue 2018, Corollaire 6.5], taking a coweight $\kappa \gg 0$ with respect to f , there is an induced morphism

$$T(f) : \mathcal{H}_{N,I,W}^{\leq\mu,E} \rightarrow \mathcal{H}_{N,I,W}^{\leq\mu+\kappa,E} \quad (4-5)$$

in $D_c^b((X \setminus \hat{N})^I, E)$, with various compatibilities. It is E -linear in f and satisfies $T(ff') = T(f)T(f')$. After passing to $\varinjlim_{\mu}$, we are led to the left $C_c(K_N \backslash G(\mathbb{A})/K_N; E)$ -module

$$H_{I,W} := \left(\varinjlim_{\mu} \mathcal{H}_{N,I,W}^{0,\leq\mu,E} \big|_{\Delta(\overline{\eta})} \right)^{\text{Hf}} \quad (4-6)$$

where “Hf” means Hecke-finite with respect to the action (4-5). This definition is clearly functorial in W . The following properties are established in [Lafforgue 2018, §§8–9]:

- Compatibility with coalescence of paws. Namely, every map $\zeta : J \rightarrow I$ induces a canonical isomorphism $\chi_{\zeta} : H_{I,W} \xrightarrow{\sim} H_{J,W^{\zeta}}$, where $W^{\zeta} \in \text{Rep}_E({}^L G^J)$ denotes the pull-back of W via ζ .
- The arrow $\mathfrak{sp}^*$ commutes with Hecke action since the latter is defined on the level of $D_c^b((X \setminus \hat{N})^I, E)$. Moreover, it induces an isomorphism

$$\mathfrak{sp}^* : H_{I,W} \xrightarrow{\sim} \left(\varinjlim_{\mu} \mathcal{H}_{N,I,W}^{0,\leq\mu,E} \big|_{\overline{\eta^I}} \right)^{\text{Hf}}.$$

- We have ${}^L G^{\varnothing} = \{1\}$, $\eta^{\varnothing} = \text{Spec } \mathbb{F}_q$ when $I = \varnothing$. There are natural isomorphisms

$$H_{\{0\},1} \xleftarrow{\chi} H_{\varnothing,1} \xrightarrow{\sim} C_c^{\text{cusp}}(\text{Bun}_{G,N}(\mathbb{F}_q)/\Xi; E). \quad (4-7)$$

The arrow χ is induced by coalescence via the unique map $\varnothing \rightarrow \{0\}$. The rightward arrow stems from the fact [Varshavsky 2004, Proposition 2.16(c)] that $\text{Cht}_{N,\varnothing,1}/\Xi$ is the constant stack $\text{Bun}_{G,N}(\mathbb{F}_q)/\Xi$ over $\text{Spec } \mathbb{F}_q$, which implies a canonical isomorphism

$$\varinjlim_{\mu} \mathcal{H}_{N,\varnothing,1}^{0,\leq\mu,E} \big|_{\Delta(\overline{\eta})} \xrightarrow{\sim} C_c(\text{Bun}_{G,N}(\mathbb{F}_q)/\Xi; E) \quad (4-8)$$

of $C_c(K_N \backslash G(\mathbb{A})/K_N; E)$ -modules.

- For $I = \{1\}$, $W = \mathbf{1}$, coalescence induces $\mathrm{Cht}_{N,\{1\},\mathbf{1}}^{(\{0\})} / \Xi \xrightarrow{\sim} (\mathrm{Cht}_{N,\emptyset,\mathbf{1}} / \Xi) \times_{\mathrm{Spec} \mathbb{F}_q} (X \setminus \hat{N})$ by [Lafforgue 2018, (8.4)]. In this case, $\mathcal{H}_{N,\{0\},\mathbf{1}}^{0,\leq\mu,E}$ is a constant sheaf and the $\varinjlim_{\mu}$ of its stalk at $\bar{\eta}$ is still $C_c(\mathrm{Bun}_{G,N}(\mathbb{F}_q) / \Xi; E)$.
- Via these isomorphisms, the $C_c(K_N \backslash G(\mathbb{A}) / K_N; E)$ -module structures on $H_{\emptyset,\mathbf{1}}$ and $H_{\{0\},\mathbf{1}}$ match the one on $C_c^{\mathrm{cusp}}(\mathrm{Bun}_{G,N}(\mathbb{F}_q) / \Xi; E)$ recorded in Section 2.1. See [Lafforgue 2018, §8].

The last item above is how harmonic analysis enters the geometric picture.

4.3. Partial Frobenius morphisms and Galois actions. We conserve the previous conventions and review the partial Frobenius morphisms. Let $J \subset I$ be finite sets. Choose a partition $I = I_1 \sqcup \cdots \sqcup I_k$ with $I_1 = J$, together with a specialization arrow $\mathrm{sp} : \bar{\eta}^I \rightarrow \Delta(\bar{\eta})$. The choice of partition intervenes in the constructions, but will disappear in the final results.

Let $\mathrm{Frob}_J = \mathrm{Frob}_{I_1} : (X \setminus \hat{N})^I \rightarrow (X \setminus \hat{N})^I$ be the morphism that equals Frob on the coordinates indexed by I_1 , and id elsewhere.

Take $W \in \mathrm{Rep}_E({}^L G)^I$ as well the lattice Ξ as in Section 4.2. In [Lafforgue 2018, §3] is defined the partial Frobenius morphism

$$\mathrm{Frob}_{I_1,N}^{(I_1,\dots,I_k)} : \mathrm{Cht}_{N,I,W}^{(I_1,\dots,I_k)} \rightarrow \mathrm{Cht}_{N,I,W}^{(I_2,\dots,I_k,I_1)} \quad (4-9)$$

covering Frob_{I_1} , that respects Ξ -actions. In terms of the notations in Section 4.1, it sends the chain

$$(\mathcal{G}_0, \psi_0) \xrightarrow{\phi_1} \cdots \rightarrow (\mathcal{G}_k, \psi_k) \rightarrow (\tau \mathcal{G}_0, \tau \psi_0)$$

into

$$(\mathcal{G}_1, \psi_1) \xrightarrow{\phi_2} \cdots \rightarrow (\mathcal{G}_k, \psi_k) \rightarrow (\tau \mathcal{G}_0, \tau \psi_0) \xrightarrow{\tau \phi_1} (\tau \mathcal{G}_1, \tau \psi_1)$$

whereas the paws are transformed accordingly by Frob_{I_1} . The cyclic composition of k partial Frobenius morphism equals the total Frobenius endomorphism of $\mathrm{Cht}_{N,I,W}^{(I_1,\dots,I_k)}$. An easy consequence is that Frob_{I_1} is a *universal homeomorphism*; see [Stacks 2005–, Tag 04DC].

The induced morphism between the quotients by Ξ is also named $\mathrm{Frob}_{I_1,N}^{(I_1,\dots,I_k)}$. Now introduce the dominant coweight μ of G^{ad} in Section 4.2 as truncation parameter. A basic fact is that whenever $\mu' \gg \mu$ with respect to W ,

$$(\mathrm{Frob}_{I_1,N}^{(I_1,\dots,I_k)})^{-1} \mathrm{Cht}_{N,I,W}^{(I_2,\dots,I_k,I_1),\leq\mu} \subset \mathrm{Cht}_{N,I,W}^{(I_1,\dots,I_k),\leq\mu'}. \quad (4-10)$$

When $k = 1$, we have the usual Frobenius correspondence $\Phi : \mathrm{Frob}^* \mathcal{S}_{I,W,E}^{(I)} \xrightarrow{\sim} \mathcal{S}_{I,W,E}^{(I)}$ between normalized perverse sheaves on $\mathrm{Gr}_{I,W}^{(I)} / G_{\sum_i n_i x_i}^{\mathrm{ad}}$. In general, by writing

$$\mathrm{Frob}_{I_1}(x_i)_{i \in I} = (x'_i)_{i \in I}, \quad (x_i)_{i \in I} \in (X \setminus \hat{N})^I(S) \quad \forall S : \mathbb{F}_q\text{-scheme}$$

and supposing $W = \boxtimes_{j=1}^k W_j$ is irreducible, there is a commutative diagram:

$$\begin{array}{ccc}
 \mathrm{Cht}_{N,I,W}^{(I_1,\dots,I_k)} / \Xi & \xrightarrow{\mathrm{Frob}_{I_1,N}^{(I_1,\dots,I_k)}} & \mathrm{Cht}_{N,I,W}^{(I_2,\dots,I_k,I_1)} / \Xi \\
 \downarrow \epsilon_{N,(I_1,\dots,I_k),\Xi}^{(I_1,\dots,I_k)} & & \downarrow \epsilon_{N,(I_2,\dots,I_k,I_1),\Xi}^{(I_2,\dots,I_k,I_1)} \\
 \prod_{j=1}^k \mathrm{Gr}_{I_j,W_j}^{(I_j)} / G_{\sum_{i \in I_j} n_i x_i}^{\mathrm{ad}} & \xrightarrow{\mathrm{Frob} \times \mathrm{id} \times \dots \times \mathrm{id}} & \prod_{j=1}^k \mathrm{Gr}_{I_j,W_j}^{(I_j)} / G_{\sum_{i \in I_j} n_i x_i'}^{\mathrm{ad}}
 \end{array}$$

In view of the constructions in Section 4.2 using the smooth morphisms $\epsilon_{\dots}$, the ULA property, etc., we obtain a canonical isomorphism in $D_c^b(\mathrm{Cht}_{N,I,W}^{(I_1,\dots,I_k)} / \Xi, E)$

$$F_{I_1,N,W}^{(I_1,\dots,I_k)} : (\mathrm{Frob}_{I_1,N}^{(I_1,\dots,I_k)})^* \mathcal{F}_{N,I,W,\Xi,E}^{(I_2,\dots,I_k,I_1)} \xrightarrow{\sim} \mathcal{F}_{N,I,W,\Xi,E}^{(I_1,\dots,I_k)} \quad (4-11)$$

extending the previous case $k = 1$. See [Lafforgue 2018, Proposition 3.4]. This isomorphism can be extended functorially to arbitrary $W \in \mathrm{Rep}_E(({}^L G)^I)$ by repeating the construction for $W \mapsto \mathcal{S}_{I,W,E}^{(I_1,\dots,I_k)}$.

Abbreviate the $\mathrm{Frob}_{I_1,N}^{(I_1,\dots,I_k)}$ on $\mathrm{Cht}_{N,I,W}^{(I_1,\dots,I_k)} / \Xi$ as a_1 . It fits into the commutative diagram

$$\begin{array}{ccccc}
 \mathrm{Cht}_{N,I,W}^{(I_2,\dots,I_k), \leq \mu} / \Xi & \xleftarrow{a_1} & a_1^{-1}(\mathrm{Cht}_{N,I,W}^{(I_2,\dots,I_k), \leq \mu} / \Xi) & \xleftarrow{a_2} & \mathrm{Cht}_{N,I,W}^{(I_1,\dots,I_k), \leq \mu'} / \Xi \\
 \downarrow \mathfrak{p} & & \downarrow \mathfrak{p} & & \downarrow \mathfrak{p} \\
 (X \setminus \hat{N})^I & \xleftarrow{\mathrm{Frob}_{I_1}} & (X \setminus \hat{N})^I & \xlongequal{\quad} & (X \setminus \hat{N})^I
 \end{array} \quad (4-12)$$

where $\mathfrak{p}$ denotes the self-evident morphisms of paws, a_1 is a universal homeomorphism and a_2 is an open immersion. Hence (4-11) affords a *cohomological correspondence* between bounded constructible complexes in the sense of [Varshavsky 2007, §1]: for a_1, a_2 in (4-12),

$$\mathrm{Frob}_{I_1,N,W}^{(I_1,\dots,I_k)} : a_1^* \underbrace{\mathcal{F}_{N,I,W,\Xi,E}^{(I_2,\dots,I_k,I_1)}}_{\text{on the } \leq \mu \text{ part}} \rightarrow a_2^! \underbrace{\mathcal{F}_{N,I,W,\Xi,E}^{(I_1,\dots,I_k)}}_{\text{on the } \leq \mu' \text{ part}}, \quad a_1^* = a_2^! \quad (4-13)$$

The left square of (4-12) is not Cartesian; however, in the commutative diagram defined with Cartesian square

$$\begin{array}{ccccc}
 & & a_1 & & a_1^{-1}(\mathrm{Cht}_{N,I,W}^{(I_2,\dots,I_k), \leq \mu} / \Xi) \\
 & & \swarrow & & \swarrow \exists! \varphi \\
 \mathrm{Cht}_{N,I,W}^{(I_2,\dots,I_k), \leq \mu} / \Xi & \xleftarrow{\tilde{a}_1} & \mathrm{Frob}_{I_1}^*(\mathrm{Cht}_{N,I,W}^{(I_2,\dots,I_k), \leq \mu} / \Xi) & & \\
 \downarrow \mathfrak{p} & \square & \downarrow \tilde{\mathfrak{p}} & & \downarrow \mathfrak{p} \\
 (X \setminus \hat{N})^I & \xleftarrow{\mathrm{Frob}_{I_1}} & (X \setminus \hat{N})^I & \xleftarrow{\quad} & (X \setminus \hat{N})^I
 \end{array}$$

the arrow φ is a universal homeomorphism since both Frob_{I_1} and a_1 are. Therefore we obtain

$$\mathrm{BC} : \mathrm{Frob}_{I_1}^* \mathfrak{p}_! \xrightarrow{\sim} \tilde{\mathfrak{p}}_! \tilde{a}_1^* \xleftarrow{\sim} \tilde{\mathfrak{p}}_! \varphi_! \varphi^* \tilde{a}_1^* \simeq \mathfrak{p}_! a_1^*. \quad (4-14)$$

Indeed, bc is the isomorphism of base change by the universal homeomorphism Frob_{I_1} [Laszlo and Olsson 2008b, 12.2]; the second isomorphism is induced by $\varphi_! \varphi^* \xrightarrow{\sim} \mathrm{id}$, which is in turn due to the topological invariance of the étale topos (see [SGA 4₁ 1972; SGA 4₂ 1972; SGA 4₃ 1973, Exp VIII, Théorème 1.1] or [Stacks 2005–, Tag 04DY]) under the universal homeomorphism φ .

In view of (4-4), we can now define

$$F_J = F_{I_1} : \mathrm{Frob}_{I_1}^* \mathcal{H}_{N,I,W}^{\leq \mu, E} \rightarrow \mathcal{H}_{N,I,W}^{\leq \mu', E} \quad (4-15)$$

as the composite in $\mathrm{D}_c^b((X \setminus \hat{N})^I, E)$

$$\mathrm{Frob}_{I_1}^* \mathfrak{p}_! \underbrace{\mathcal{F}_{N,I,W,\Xi,E}^{(I_2, \dots, I_k)}}_{\text{on } \leq \mu} \xrightarrow{\sim} \mathfrak{p}_! a_1^* \underbrace{\mathcal{F}_{N,I,W,\Xi,E}^{(I_2, \dots, I_k)}}_{\text{on } \leq \mu} \xrightarrow{\mathfrak{p}_! \mathrm{Frob}_{I_1, N, W}^{(I_1, \dots, I_k)}} \mathfrak{p}_! a_2^! \underbrace{\mathcal{F}_{N,I,W,\Xi,E}^{(I_1, \dots, I_k)}}_{\text{on } \leq \mu'} \rightarrow \underbrace{\mathcal{F}_{N,I,W,\Xi,E}^{(I_1, \dots, I_k)}}_{\text{on } \leq \mu'}$$

the last arrow arising from $\mathfrak{p}_! a_2^! = \mathfrak{p}_! (a_2)_! a_2^! \xrightarrow{(a_2)_! a_2^! \rightarrow \mathrm{id}} \mathfrak{p}_!$. It is functorial in $W \in \mathrm{Rep}_E(({}^L G)^I)$ and is shown to be compatible with the coalescence of paws in [Lafforgue 2018, §§3–4]. Hence the dependence is only on $J \subset I$.

Consequently, the morphism F_J also acts E -linearly on $\varinjlim_{\mu} \mathcal{H}_{N,I,W}^{0, \leq \mu, E} |_{\overline{\eta}^I}$. Given any partition $I = I_1 \sqcup \dots \sqcup I_k$, the actions of $F_{I_1}, \dots, F_{I_k}$ form a commuting family whose cyclic composition equals the total Frobenius action on $\varinjlim_{\mu} \mathcal{H}_{N,I,W}^{0, \leq \mu, E} |_{\overline{\eta}^I}$.

On the other hand, the standard theory [Stacks 2005–, Tag 03QW] yields a continuous representation of $\pi_1(\eta^I, \overline{\eta}^I)$ on $\mathcal{H}_{N,I,W}^{0, \leq \mu, E} |_{\overline{\eta}^I}$ which passes to $\varinjlim_{\mu}$.

To conclude this subsection, we recall briefly the following extension of groups

$$1 \rightarrow \ker[\pi_1(\eta^I, \overline{\eta}^I) \rightarrow \hat{\mathbb{Z}}] \rightarrow \mathrm{FWeil}(\eta^I, \overline{\eta}^I) \rightarrow \mathbb{Z}^I \rightarrow 0.$$

We refer to [Lafforgue 2018, Remarque 8.18] and the subsequent discussions for all further details. When $|I| = 1$, it becomes the Weil group $W_{\tilde{F}}$ of $\tilde{F} = \mathbb{F}_q(X)$; in general there is a surjection $\mathrm{FWeil}(\eta^I, \overline{\eta}^I) \twoheadrightarrow W_{\tilde{F}}^I$ depending on the choice of $\mathfrak{sp}$. The surjection induces an isomorphism from the profinite completion $\mathrm{FWeil}(\eta^I, \overline{\eta}^I)$ to that of $W_{\tilde{F}}^I$, i.e., $\pi_1(\eta, \overline{\eta})^I$. As mentioned in [loc. cit.], the action on $\varinjlim_{\mu} \mathcal{H}_{N,I,W}^{0, \leq \mu, E} |_{\overline{\eta}^I}$ of

- the partial Frobenius morphisms F_J for various $J \subset I$, and
- that of $\pi_1(\eta^I, \overline{\eta}^I)$

meld into an action of $\mathrm{FWeil}(\eta^I, \overline{\eta}^I)$. The upshot of [loc. cit., §8] is to produce a continuous E -linear $\pi_1(\eta, \overline{\eta})^I$ -action on $(\varinjlim_{\mu} \mathcal{H}_{N,I,W}^{0, \leq \mu, E} |_{\overline{\eta}^I})^{\mathrm{Hf}}$ therefrom. In other words, one wants to factorize the $\mathrm{FWeil}(\eta^I, \overline{\eta}^I)$ -action through its profinite completion continuously.

The key for the passage to $\pi_1(\eta, \overline{\eta})^I$ -action is *Drinfeld's Lemma*. This method requires some finiteness conditions which in turn involve the Eichler–Shimura relations. These important issues are addressed at length in [loc. cit., §8], but they are not needed in this article.

The aforementioned continuous representation transports to $H_{I,W}$, namely

$$\vec{\gamma} \cdot f := (\mathfrak{sp}^*)^{-1}(\vec{\gamma} \cdot (\mathfrak{sp}^* f)), \quad \vec{\gamma} \in \pi_1(\eta, \bar{\eta})^I, f \in H_{I,W}.$$

This action turns out to be independent of the choice of $\bar{\eta}^I$ and $\mathfrak{sp}$, by [loc. cit., Lemme 9.4].

4.4. Excursion operators and pseudocharacters. Consider finite sets I, J and $W \in \text{Rep}_E(({}^L G)^I)$ and $U \in \text{Rep}_E(({}^L G)^J)$. Let ζ_I, ζ_J be the unique maps from I, J into the singleton $\{0\}$. The diagonal action on W gives $W^{\zeta_I} \in \text{Rep}_E({}^L G)$; the space of $\hat{G}$ -invariants $(W^{\zeta_I})^{\hat{G}}$ is therefore a representation of $\text{Gal}(\tilde{F}|F)$. Denote by $(W^{\zeta_I})^{\hat{G}}|_{X \setminus \hat{N}}$ the E -lisse sheaf on $X \setminus \hat{N}$ obtained by descent. Likewise, we have $(W^{\zeta_I})^{\hat{G}}|_{X \setminus \hat{N}}$ by taking the maximal quotient of W^{ζ_I} on which $\hat{G}$ acts trivially. A pair of morphisms

$$\begin{aligned} \mathcal{H}_{N,J,U}^{\leq \mu, E} \boxtimes (W^{\zeta_I})^{\hat{G}}|_{X \setminus \hat{N}} &\rightarrow \mathcal{H}_{N,J \sqcup I, U \boxtimes W}^{\leq \mu, E}|_{(X \setminus \hat{N})^J \times \Delta(X \setminus \hat{N})} \\ \mathcal{H}_{N,J,U}^{\leq \mu, E} \boxtimes (W^{\zeta_I})^{\hat{G}}|_{X \setminus \hat{N}} &\leftarrow \mathcal{H}_{N,J \sqcup I, U \boxtimes W}^{\leq \mu, E}|_{(X \setminus \hat{N})^J \times \Delta(X \setminus \hat{N})} \end{aligned}$$

in $D_c^b((X \setminus \hat{N})^{J \sqcup \{0\}}, E)$ are constructed in [loc. cit., (12.18), (12.19)]. Roughly speaking, they are defined via coalescence and the functoriality of $\mathcal{H}$ with respect to $(W^{\zeta_I})^{\hat{G}} \hookrightarrow W^{\zeta_I} \twoheadrightarrow (W^{\zeta_I})^{\hat{G}}$.

Now take $J = \emptyset$ and $U = \mathbf{1}$. Let $x \in W$ and $\xi \in W^\vee$ be $\hat{G}$ -invariant under the diagonal action, viewed as maps $E \rightarrow (W^{\zeta_I})^{\hat{G}}$ and $(W^{\zeta_I})^{\hat{G}} \rightarrow E$, respectively. Taking $\varinjlim_\mu H^0(\cdots|_{\bar{\eta}})$ yields the creation and annihilation operators (see [loc. cit., Définitions 5.1, 5.2 and 12.3.4])

$$\varinjlim_\mu \mathcal{H}_{N,\emptyset,\mathbf{1}}^{0,\leq \mu, E} \xrightleftharpoons[C_\xi^b]{C_x^\sharp} \varinjlim_\mu \mathcal{H}_{N,I,W}^{0,\leq \mu, E}|_{\Delta(\bar{\eta})}$$

between E -vector spaces. Restriction to Hecke-finite parts yields arrows

$$H_{\emptyset,\mathbf{1}} \simeq H_{\{0\},\mathbf{1}} \xrightleftharpoons[C_\xi^b]{C_x^\sharp} H_{I,W}, \quad \text{see (4-6).}$$

Given I, W, x, ξ as above and $\vec{\gamma} = (\gamma_i)_{i \in I} \in \pi_1(\eta, \bar{\eta})^I$, the *excursion operator* $S_{I,W,x,\xi,\vec{\gamma}}$ is the composite

$$\begin{array}{ccccc} H_{\{0\},\mathbf{1}} & & & & H_{\{0\},\mathbf{1}} \\ \downarrow C_x^\sharp & & & & \uparrow C_\xi^b \\ H_{I,W} & \xrightarrow[\sim]{\mathfrak{sp}^*} & (\varinjlim_\mu \mathcal{H}_{N,I,W}^{0,\leq \mu, E}|_{\bar{\eta}^I})^{\text{Hf}} & \xrightarrow[\sim]{\vec{\gamma}} & (\varinjlim_\mu \mathcal{H}_{N,I,W}^{0,\leq \mu, E}|_{\bar{\eta}^I})^{\text{Hf}} \xrightarrow[\sim]{(\mathfrak{sp}^*)^{-1}} H_{I,W} \end{array}$$

Here $\vec{\gamma}$ acts in the manner reviewed in Section 4.3. Upon recalling (4-7), we obtain

$$S_{I,W,x,\xi,\vec{\gamma}} \in \text{End}_E(H_{\{0\},\mathbf{1}}) \simeq \text{End}_E(C_c^{\text{cusp}}(\text{Bun}_{G,N}(\mathbb{F}_q)/\Xi; E)).$$

Moreover, by [loc. cit., Définition-Proposition 9.1]

- we have $S_{I,W,x,\xi,\vec{\gamma}} \in \text{End}_{C_c(K_N \backslash G(\mathbb{A})/K_N; E)}(H_{\{0\}, \mathbf{1}})$;
- the formation of $S_{I,W,x,\xi,\vec{\gamma}}$ is E -bilinear in x, ξ and continuous in $\vec{\gamma}$ for the topology on the finite-dimensional space $\text{End}_E(H_{\{0\}, \mathbf{1}})$ induced by E ;
- let $\mathcal{B}_E$ be the E -subalgebra of $\text{End}_{C_c(K_N \backslash G(\mathbb{A})/K_N; E)}(H_{\{0\}, \mathbf{1}})$ generated by $S_{I,W,x,\xi,\vec{\gamma}}$ for all quintuples $(I, W, x, \xi, \vec{\gamma})$. Then $\mathcal{B}_E$ is a finite-dimensional commutative E -algebra by [loc. cit., (10.2)].

The foregoing constructions behave well under finite extensions of the field E of coefficients. Define the $\overline{\mathbb{Q}_\ell}$ -algebra

$$\mathcal{B} := \mathcal{B}_E \otimes_E \overline{\mathbb{Q}_\ell} \subset \text{End}_{\overline{\mathbb{Q}_\ell}}(H_{\{0\}, \mathbf{1}} \otimes_E \overline{\mathbb{Q}_\ell}).$$

Upon enlarging E , we may assume that all homomorphisms $\nu : \mathcal{B} \rightarrow \overline{\mathbb{Q}_\ell}$ (finitely many) of $\overline{\mathbb{Q}_\ell}$ -algebras are defined over E . There is a decomposition of $C_c(K_N \backslash G(\mathbb{A})/K_N; E)$ -modules into generalized eigenspaces

$$H_{\{0\}, \mathbf{1}} = \bigoplus_{\nu} \mathfrak{H}_{\nu}, \quad \mathfrak{H}_{\nu} := \{f \in H_{\{0\}, \mathbf{1}} : \forall T \in \mathcal{B}_E, \exists d \geq 1 \mid (T - \nu(T))^d f = 0\}. \quad (4-16)$$

Here ν ranges over the characters of $\mathcal{B}$, and one may take $d = \dim_E H_{\{0\}, \mathbf{1}}$. The same holds after passing to $\overline{\mathbb{Q}_\ell}$. All in all,

$$C_c^{\text{cusp}}(\text{Bun}_{G,N}(\mathbb{F}_q)/\Xi; \overline{\mathbb{Q}_\ell}) = \bigoplus_{\nu: \mathcal{B} \rightarrow \overline{\mathbb{Q}_\ell}} \mathfrak{H}_{\nu} \quad \text{in } C_c(K_N \backslash G(\mathbb{A})/K_N; \overline{\mathbb{Q}_\ell})\text{-Mod}.$$

It is conjectured that $\mathcal{B}$ is reduced, which will imply that $d = 1$ suffices.

The next step is to reencode the excursion operators $S_{I,W,x,\xi,\vec{\gamma}}$. Let $f(\vec{g}) = \langle \xi, \vec{g} \cdot x \rangle_{W^\vee \otimes W}$ where $\vec{g} \in ({}^L G)^I$ and $\langle \cdot, \cdot \rangle_{W^\vee \otimes W}$ is the duality pairing $W^\vee \otimes_E W \rightarrow E$. Then $f \in \mathcal{O}(\hat{G} \backslash ({}^L G)^I // \hat{G})$, where $\hat{G}$ acts by bilateral translations through diagonal embedding. By [loc. cit., Lemme 10.6], $S_{I,W,x,\xi,\vec{\gamma}}$ depends only on $(I, f, \vec{\gamma})$. Using some algebraic version of the Peter–Weyl theorem, one can uniquely define the operators

$$S_{I,f,\vec{\gamma}} \in \mathcal{B}_E, \quad f \in \mathcal{O}(\hat{G} \backslash ({}^L G)^I // \hat{G}),$$

in a manner compatible with the original $S_{I,W,x,\xi,\vec{\gamma}}$, such that if f comes from a function $\text{Gal}(\tilde{F}|\mathring{F})^I \rightarrow E$, then $S_{I,f,\vec{\gamma}} = f(\vec{\gamma}) \cdot \text{id}$. See [loc. cit., Remarque 12.20] for further explanations.

Take $n \in \mathbb{Z}_{\geq 1}$ and $I := \{0, \dots, n\}$. Then $\hat{G}$ acts on $({}^L G)^n$ by simultaneous conjugation. There is a natural map

$$\begin{aligned} \mathcal{O}({}^L G)^n // \hat{G} &\rightarrow \mathcal{O}({}^L G \backslash ({}^L G)^{\{0, \dots, n\}} // \hat{G}) \subset \mathcal{O}(\hat{G} \backslash ({}^L G)^{\{0, \dots, n\}} // \hat{G}) \\ f &\mapsto [\tilde{f} : (g_0, \dots, g_n) \mapsto f(g_0^{-1} g_1, \dots, g_0^{-1} g_n)]. \end{aligned} \quad (4-17)$$

When n is fixed, the operators

$$\Theta_n(f)(\vec{\gamma}) := S_{\{0, \dots, n\}, \tilde{f}, (1, \vec{\gamma})}, \quad n \in \mathbb{Z}_{\geq 1}, \vec{\gamma} \in \pi_1(\eta, \bar{\eta})^n, f \in \mathcal{O}({}^L G)^n // \hat{G} \quad (4-18)$$

in $\mathcal{B}_E$ afford a homomorphism $\mathcal{O}(({}^L G)^n // \hat{G}) \rightarrow C(\pi_1(X \setminus \hat{N}, \bar{\eta})^n, \mathcal{B}_E)$ between E -algebras, where $C(\dots)$ denotes the algebra of continuous functions under pointwise operations. See [loc. cit., Proposition 10.10] for the passage to $\pi_1(X \setminus \hat{N}, \bar{\eta})$.

Since $\mathcal{O}({}^L G // ({}^L G)^{\{0, \dots, n\}} // \hat{G}) \subsetneq \mathcal{O}(\hat{G} // ({}^L G)^{\{0, \dots, n\}} // \hat{G})$ in general, the map (4-17) is not always surjective. Nonetheless, the operators $S_{\{0, \dots, n\}, \tilde{f}, (1, \vec{\gamma})}$ still generate $\mathcal{B}_E$ as $n, f, \vec{\gamma}$ vary; see [loc. cit., Remarque 12.20].

Finally, the machinery of ${}^L G$ -pseudocharacters associates a semisimple L -parameter $\sigma \in \Phi(G)$ to any character $\nu : \mathcal{B} \rightarrow \overline{\mathbb{Q}_\ell}$, characterized as follows:

- Version 1: for all $n \in \mathbb{Z}_{\geq 1}$, $\vec{\gamma} = (\gamma_1, \dots, \gamma_n)$ and $f \in \mathcal{O}(({}^L G)^n // \hat{G})$, we have (see [loc. cit., Proposition 11.7])

$$f(\sigma(\gamma_1), \dots, \sigma(\gamma_n)) = \nu \circ \Theta_n(f)(\vec{\gamma}).$$

- Version 2: for all $n \in \mathbb{Z}_{\geq 1}$, $\vec{\gamma} = (\gamma_1, \dots, \gamma_n)$ and $\tilde{f} \in \mathcal{O}(\hat{G} // ({}^L G)^{\{0, \dots, n\}} // \hat{G})$, we have

$$\tilde{f}(\sigma(1), \sigma(\gamma_1), \dots, \sigma(\gamma_n)) = \nu(S_{\{0, \dots, n\}, \tilde{f}, (1, \vec{\gamma})}). \quad (4-19)$$

The version 2 above is *a priori* stronger, but they are actually equivalent by the preceding remarks on generators.

By the discussions preceding [loc. cit., Remarque 12.21], the map $\text{Hom}_{\overline{\mathbb{Q}_\ell}\text{-Alg}}(\mathcal{B}, \overline{\mathbb{Q}_\ell}) \rightarrow \Phi(G)$ above is injective. Hence we may write $\mathfrak{H}_\sigma = \mathfrak{H}_\nu$ if $\nu \mapsto \sigma \in \Phi(G)$, and set $\mathfrak{H}_\sigma = \{0\}$ if σ does not match any ν . This leads to the desired decomposition (3-2).

5. The transposes of excursion operators

5.1. On Verdier duality. Retain the notation of Section 4.2. Among them, we recall only two points:

- The duality operator $\mathbb{D}$ is normalized with respect to $(X \setminus \hat{N})^I$.
- $W^{\vee, \theta}$ denotes the contragredient of $W \in \text{Rep}_E({}^L G)^I$ twisted by the Chevalley involution of $({}^L G)^I$.

The following results are recorded in [loc. cit., Remarque 5.4]. For the benefit of the readers, we will give some more details below.

Proposition 5.1.1. *There is a canonical isomorphism*

$$\mathbb{D} \mathcal{S}_{I, W, E}^{(I_1, \dots, I_k)} \xrightarrow{\sim} \mathcal{S}_{I, W^{\vee, \theta}, E}^{(I_1, \dots, I_k)}$$

between functors from $W \in \text{Rep}_E({}^L G)^I$ to $\text{Perv}_{\bar{G}_{\sum_i \infty x_i}}(\text{Gr}_I^{(I_1, \dots, I_k)})$.

Proof. As noted in [Braverman and Gaitsgory 2002, §B.6], $\mathbb{D}$ preserves the ULA property with respect to $\text{Gr}_I^{(I_1, \dots, I_k)} \rightarrow (X \setminus \hat{N})^I$. Since $\mathcal{S}_{I, W, E}^{(I_1, \dots, I_k)}$ is ULA, the factorization structure on $\text{Gr}_I^{(I_1, \dots, I_k)}$ reduces the affairs to the case $|I| = 1$, i.e., the Beilinson–Drinfeld affine Grassmannian used in [Mirković and Vilonen 2007; Richarz 2014; Zhu 2015]. Consider its fiber Gr_x over some point $x \in |X \setminus \hat{N}|$. In the notation from Section 4.1, there is a left $G_{\infty x}$ -action on Gr_x .

In the local setting above, denote the usual duality operator $\mathrm{Perv}_{G_\infty}(\mathrm{Gr}_x)$ by $\mathbb{D}$; normalization is not an issue here. The main ingredients are

- the Satake functor $\mathrm{Rep}_E({}^L G)^I \rightarrow \mathrm{Perv}_{G_\infty}(\mathrm{Gr}_x)$, written as $W \mapsto \mathcal{S}_{W,E}$;
- a canonical isomorphism between functors in W :

$$\mathbb{D}\mathcal{S}_{W,E} \xrightarrow{\sim} \mathcal{S}_{W^{\vee,\theta},E}.$$

Granting these ingredients, for general $|I|$ we obtain canonical isomorphisms $\mathbb{D}\mathcal{S}_{I,W,E}^{(I_1,\dots,I_k)} \xrightarrow{\sim} \mathcal{S}_{I,W^{\vee,\theta},E}^{(I_1,\dots,I_k)}$

Let us explain the two ingredients in the local setting. The functor $W \mapsto \mathcal{S}_{W,E}$ is obtained in [Richarz 2014; Zhu 2015, Theorem A.12], which are based on the case over separably closed fields in [Mirković and Vilonen 2007]. In order to explain the effect of ${}^L\theta$, we shall review the case over the separable closure $\mathbb{k}$ of $\mathbb{F}_x$ first. The canonical isomorphism $\mathbb{D}\mathcal{S}_{W,E} \xrightarrow{\sim} \mathcal{S}_{W^{\vee,\theta},E}$ over $\mathbb{k}$ can be found in [Bezrukavnikov and Finkelberg 2008, Lemma 14], for example, where a stronger equivariant version is established; they work over $\mathbb{C}$, but the argument is largely formal.

Next, apply Galois descent as explicated in [Richarz 2014, §6; Zhu 2015, Appendix]. Let $\mathcal{C} := \mathrm{Perv}_{G_\infty}(\mathrm{Gr}_x)$ and set $\mathcal{C}'$ to be its avatar over $\mathbb{k}$. The absolute Galois group Υ of $\mathbb{F}_x$ operates on $\mathcal{C}'$ via $\otimes$ -equivalences, in a manner compatible with the fiber functor (total cohomology), thus Υ acts on the Tannakian group $\hat{G}$ as well. By [Richarz 2014, p.237], $\mathcal{C}$ is equivalent as an abelian category to $(\mathcal{C}' + \text{continuous descent data under } \Upsilon)$. The Satake equivalence over $\mathbb{k}$ and the machinery from [loc. cit.] furnish an equivalence of $\otimes$ -categories

$$\mathrm{Perv}_{G_\infty}(\mathrm{Gr}_x) \rightarrow \mathrm{Rep}_E(\hat{G} \rtimes_{\mathrm{geom}} \Upsilon),$$

where c means continuity, and “geom” means the Tannakian or “geometric” Υ -action on $\hat{G}$. See [Lafforgue 2018, Remarque 1.19] for the choice of commutativity constraints.

By [Zhu 2015, Proposition A.6; Richarz 2014, Corollary 6.8], the geometric Υ -action on $\hat{G}$ differs from the familiar “algebraic” one by the adjoint action via $\rho_{\hat{B}} \circ \chi_{\mathrm{cycl}} : \Upsilon \rightarrow \mathbb{Z}_\ell^\times \rightarrow \hat{G}^{\mathrm{ad}}(\mathbb{Q}_\ell)$, where χ_{cycl} is the ℓ -adic cyclotomic character and $\rho_{\hat{B}}$ is the half-sum of positive roots in $\hat{B}$; in particular, $\hat{G} \rtimes_{\mathrm{geom}} \Upsilon \simeq \hat{G} \rtimes_{\mathrm{alg}} \Upsilon$ (= absolute Galois form of the L -group) continuously. Since $\theta \in \mathrm{Aut}(\hat{G})$ stabilizes $\rho_{\hat{B}}$, the isomorphism matches $\theta \rtimes_{\mathrm{geom}} \mathrm{id}$ with the Chevalley involution $\theta \rtimes_{\mathrm{alg}} \mathrm{id} =: {}^L\theta$.

All in all, we obtain the Satake functor $W \mapsto \mathcal{S}_{W,E}$ as well as the canonical isomorphisms $\mathbb{D}\mathcal{S}_{W,E} \xrightarrow{\sim} \mathcal{S}_{W^{\vee,\theta},E}$. This completes the proof. $\square$

Note that the equivariance can be upgraded to $G_{\sum_i \infty x_i}^{\mathrm{ad}}$ or $G_{\sum_i n_i x_i}^{\mathrm{ad}}$ where $n_i \gg 0$ relative to W , see [Lafforgue 2018, Remarque 1.20].

Proposition 5.1.2. *There is a canonical isomorphism*

$$\mathbb{D}\mathcal{F}_{N,I,W,\Xi,E}^{(I_1,\dots,I_k)} \xrightarrow{\sim} \mathcal{F}_{N,I,W^{\vee,\theta},\Xi,E}^{(I_1,\dots,I_k)}$$

between functors from $W \in \mathrm{Rep}_E({}^L G)^I{}^{\mathrm{op}}$ to $\mathrm{Perv}(\mathrm{Cht}_{N,I,W}^{(I_1,\dots,I_k)} / \Xi)$ that is compatible with coalescence of paws.

Proof. First, by combining [Lafforgue 2018, Proposition 2.8] and the explanations before Corollaire 2.15 of [loc. cit.], the smooth morphisms

$$\mathrm{Cht}_{N,I,W}^{(I_1,\dots,I_k)} / \Xi \xrightarrow{\epsilon_{N,I,W,n}^{(I_1,\dots,I_k),\Xi}} \mathrm{Gr}_{I,W}^{(I_1,\dots,I_k)} / G_{\sum_i n_i x_i}^{\mathrm{ad}}, \quad \mathrm{Gr}_{I,W}^{(I_1,\dots,I_k)} \rightarrow \mathrm{Gr}_{I,W}^{(I_1,\dots,I_k)} / G_{\sum_i n_i x_i}^{\mathrm{ad}}$$

have the same relative dimension; denote it by d .

Proposition 5.1.1 gives a functorial isomorphism between descent data of shifted perverse sheaves from $\mathrm{Gr}_{I,W}^{(I_1,\dots,I_k)}$ to $\mathrm{Gr}_{I,W}^{(I_1,\dots,I_k)} / G_{\sum_i n_i x_i}^{\mathrm{ad}}$, abbreviated as $\mathbb{D}\mathcal{S}_1 \xrightarrow{\sim} \mathcal{S}_2$. Denote the corresponding shifted perverse sheaves on $\mathrm{Gr}_{I,W}^{(I_1,\dots,I_k)} / G_{\sum_i n_i x_i}^{\mathrm{ad}}$ as $\mathcal{S}_1^b$ and $\mathcal{S}_2^b$. By standard results, see [Laszlo and Olsson 2008b, 9.1.2], the isomorphism above descends to

$$(\mathbb{D}\mathcal{S}_1^b)[2d](d) \xrightarrow{\sim} \mathcal{S}_2^b.$$

Since $\mathcal{F}_{N,I,W,\Xi,E}^{(I_1,\dots,I_k)}$ and $\mathcal{F}_{N,I,W^{\vee,\theta},\Xi,E}^{(I_1,\dots,I_k)}$ are defined in (4-3) as $(\epsilon_{N,I,W,n}^{(I_1,\dots,I_k),\Xi})^* \mathcal{S}_1^b$ and $(\epsilon_{N,I,W^{\vee,\theta},n}^{(I_1,\dots,I_k),\Xi})^* \mathcal{S}_2^b$, respectively, the assertion follows immediately by the same standard result. $\square$

Take any partition $I = I_1 \sqcup \dots \sqcup I_k$, truncation parameter μ and $W \in \mathrm{Rep}_E(({}^L G)^I)$. As a consequence of Propositions 5.1.1 and 5.1.2, we deduce that $\mathrm{Gr}_{I,W}^{(I_1,\dots,I_k)} = \mathrm{Gr}_{I,W^{\vee,\theta}}^{(I_1,\dots,I_k)}$ and $\mathrm{Cht}_{N,I,W}^{(I_1,\dots,I_k),\leq\mu} = \mathrm{Cht}_{N,I,W^{\vee,\theta}}^{(I_1,\dots,I_k),\leq\mu}$.

Remark 5.1.3. Below is a review of the cup product of $!$ -pushforward. Let S be a regular scheme and let $p : \mathcal{X} \rightarrow S$ be an algebraic stack of finite type over S . Let $\mathcal{L}, \mathcal{L}'$ be in $D_c^-(\mathcal{X}, E)$. Our goal is to define a canonical arrow

$$p_! \mathcal{L} \otimes^L p_! \mathcal{L}' \rightarrow p_! (\mathcal{L} \otimes^L \mathcal{L}').$$

Denote by Δ and $p \times p$ the diagonal morphisms $\mathcal{X} \rightarrow \mathcal{X} \times_S \mathcal{X}$ and $\mathcal{X} \times_S \mathcal{X} \rightarrow S$, respectively. The Künneth formula [Laszlo and Olsson 2008b, 11.0.14 Theorem] yields a canonical isomorphism in $D_c^-(S)$

$$p_! \mathcal{L} \otimes^L p_! \mathcal{L}' \simeq (p \times p)_! (\mathcal{L} \boxtimes \mathcal{L}'),$$

where $\boxtimes$ denotes the external tensor product. Since $\mathcal{L} \otimes^L \mathcal{L}' = \Delta^* (\mathcal{L} \boxtimes \mathcal{L}')$, to obtain the desired arrow, it remains to use the

$$(p \times p)_! \rightarrow (p \times p)_! \Delta_! \Delta^* = p_! \Delta^*$$

arising from $\mathrm{id} \rightarrow \Delta_* \Delta^* = \Delta_! \Delta^*$, as Δ is a closed immersion.

Consider the normalized dualizing complex Ω on $\mathrm{Cht}_{N,I,W}^{(I_1,\dots,I_k),\leq\mu} / \Xi$. The *trace map*

$$\mathrm{Tr} : (p_{N,I}^{(I_1,\dots,I_k),\leq\mu})_! \Omega \xrightarrow{\mathrm{Tr}} E_{(X \setminus \hat{N})^I}$$

in Verdier duality is obtained by adjunction from $\Omega \xrightarrow{\sim} (p_{N,I}^{(I_1,\dots,I_k),\leq\mu})^! E_{(X \setminus \hat{N})^I}$.

On the other hand, Proposition 5.1.2 affords a canonical arrow $\mathcal{F}_{N,I,W^{\vee,\theta},\Xi,E}^{(I_1,\dots,I_k)} \overset{\mathbb{L}}{\otimes} \mathcal{F}_{N,I,W,\Xi,E}^{(I_1,\dots,I_k)} \rightarrow \Omega$. Apply the cup-product construction to the stack $\text{Cht}_{N,I,W}^{(I_1,\dots,I_k),\leq\mu} / \Xi$ over $(X \setminus \hat{N})^I$ to obtain canonical arrows in $D_c^b((X \setminus \hat{N})^I, E)$:

$$\begin{array}{ccc}
 \mathcal{H}_{N,I,W^{\vee,\theta}}^{\leq\mu,E} \overset{\mathbb{L}}{\otimes} \mathcal{H}_{N,I,W}^{\leq\mu,E} & & E_{(X \setminus \hat{N})^I} \\
 \parallel & & \uparrow \\
 (\mathfrak{p}_{N,I}^{(I_1,\dots,I_k),\leq\mu})_!(\mathcal{F}_{N,I,W^{\vee,\theta},\Xi,E}^{(I_1,\dots,I_k)}) \overset{\mathbb{L}}{\otimes} (\mathfrak{p}_{N,I}^{(I_1,\dots,I_k),\leq\mu})_!(\mathcal{F}_{N,I,W,\Xi,E}^{(I_1,\dots,I_k)}) & & \text{Tr} \\
 \downarrow & & \uparrow \\
 (\mathfrak{p}_{N,I}^{(I_1,\dots,I_k),\leq\mu})_!(\mathcal{F}_{N,I,W^{\vee,\theta},\Xi,E}^{(I_1,\dots,I_k)}) \overset{\mathbb{L}}{\otimes} \mathcal{F}_{N,I,W,\Xi,E}^{(I_1,\dots,I_k)} & \longrightarrow & (\mathfrak{p}_{N,I}^{(I_1,\dots,I_k),\leq\mu})_!\Omega
 \end{array} \quad (5-1)$$

By homological common sense (see [Kashiwara and Schapira 1990, Example I.24(ii)] for example), taking $H^\bullet$ in (5-1) with respect to the ordinary t -structure on $(X \setminus \hat{N})^I$ yield natural arrows between E -sheaves over $(X \setminus \hat{N})^I$

$$\mathfrak{B}_{N,I,W}^{\Xi,E} : \mathcal{H}_{N,I,W^{\vee,\theta}}^{i,\leq\mu,E} \otimes_E \mathcal{H}_{N,I,W}^{-i,\leq\mu,E} \rightarrow E_{(X \setminus \hat{N})^I}, \quad i \in \mathbb{Z};$$

we will only use the case $i = 0$ in this article.

Following [Lafforgue 2018, Remarque 9.2], we may even pass to $\varinjlim_\mu$ and look at the stalk at $\xi \in \{\eta^I, \Delta(\bar{\eta})\}$, thereby obtain from $\mathfrak{B}_{N,I,W}^{\Xi,E}$ the E -bilinear pairings

$$\langle \cdot, \cdot \rangle_\xi : \varinjlim_\mu \mathcal{H}_{N,I,W^{\vee,\theta}}^{0,\leq\mu,E} \big|_\xi \otimes_E \varinjlim_\mu \mathcal{H}_{N,I,W}^{0,\leq\mu,E} \big|_\xi \longrightarrow E.$$

Their relation with the arrow $\mathfrak{sp}$ of specialization is given by [loc. cit., (9.6)]

$$\langle \mathfrak{sp}^* h, \mathfrak{sp}^* h' \rangle_{\eta^I} = \langle h, h' \rangle_{\Delta(\bar{\eta})}. \quad (5-2)$$

In the discussions surrounding (4-8), we have seen that $\text{Cht}_{N,\emptyset,1} / \Xi$ is the constant stack $\text{Bun}_{G,N}(\mathbb{F}_q) / \Xi$ over $\text{Spec } \mathbb{F}_q$. The upshot is that, as in [loc. cit., Remarque 9.2], the pairing $\langle \cdot, \cdot \rangle_{\Delta(\bar{\eta})}$ for $I = \emptyset$ reduces to the integration pairing on $C_c(G(\check{F}) \backslash G(\mathbb{A}) / K_N \Xi; E)$, assuming $\text{mes}(K_N) = 1$. Upon restriction to Hecke-finite part, we get the pairing $\langle \cdot, \cdot \rangle$ for $H_{\emptyset,1}$ in Remark 2.2.3. The same holds for $H_{\{0\},1}$ by coalescence (4-7).

5.2. Frobenius invariance. For every morphism f between reasonable schemes or stacks, we will denote by “can” the canonical isomorphisms exchanging $f^* \leftrightarrow f^!$ and $f_* \leftrightarrow f_!$ under $\mathbb{D}$. When f is a universal homeomorphism or open immersion, we have $f_* = f_!$ and $f^* = f^!$ or only $f^* = f^!$, respectively.

Merge the conventions from Sections 5.1 and 4.3. Let $J \subset I$ be finite sets, $I = I_1 \sqcup \dots \sqcup I_k$ with $J = I_1$. We are going to explicate the compatibility between $\mathfrak{B}_{N,I,W}^{\Xi,E}$ and $\text{Frob}_J^* \mathcal{H}_{N,I,W}^{\leq\mu,E} \xrightarrow{F_J} \mathcal{H}_{N,I,W}^{\leq\mu',E}$, i.e., (4-15).

Lemma 5.2.1. *In $D_c^b(\text{Cht}_{N,I,W}^{(I_1,\dots,I_k)} / \Xi, E)$, there is a commutative diagram whose arrows are all invertible:*

$$\begin{array}{ccc}
 (\text{Frob}_{I_1,N,W^{\vee,\theta}}^{(I_1,\dots,I_k)})^* \mathcal{F}_{N,I,W^{\vee,\theta},E}^{(I_2,\dots,I_1)} & \longrightarrow & (\text{Frob}_{I_1,N,W}^{(I_1,\dots,I_k)})^* \mathbb{D} \mathcal{F}_{N,I,W,\Xi,E}^{(I_2,\dots,I_1)} \xrightarrow[\sim]{\text{can}} \mathbb{D}(\text{Frob}_{I_1,N,I}^{(I_1,\dots,I_k)})^* \mathcal{F}_{N,I,W,\Xi,E}^{(I_2,\dots,I_1)} \\
 \downarrow F_{I,N,W^{\vee,\theta}}^{(I_1,\dots,I_k)} & & \nearrow \mathbb{D} F_{I,N,W}^{(I_1,\dots,I_k)} \\
 \mathcal{F}_{N,I,W^{\vee,\theta},\Xi,E}^{(I_1,\dots,I_k)} & \longrightarrow & \mathbb{D} \mathcal{F}_{N,I,W,\Xi,E}^{(I_1,\dots,I_k)}
 \end{array}$$

where $F_{I,N,\dots}^{(I_1,\dots,I_k)}$ is from (4-11), and the horizontal arrows except can are induced by Proposition 5.1.2.

Proof. We may assume $W = \boxtimes_{j=1}^k W_j$ is irreducible. Using the definition (4-3), the smoothness of $\epsilon_{N,I,W,\underline{n}}^{(I_1,\dots,I_k),\Xi}$ as well as the ULA properties of $\mathcal{F}$ and $\mathcal{S}$, the desired commutativity eventually reduces to that of

$$\begin{array}{ccc}
 \text{Frob}^* \mathcal{S}_{I_j,W_j^{\vee,\theta},E}^{(I_j)} & \longrightarrow & \text{Frob}^* \mathbb{D} \mathcal{S}_{I_j,W_j,E}^{(I_j)} \xrightarrow[\sim]{\text{can}} \mathbb{D} \text{Frob}^* \mathcal{S}_{I_j,W_j,E}^{(I_j)} \\
 \downarrow \Phi & & \nearrow \mathbb{D} \Phi \\
 \mathcal{S}_{I_j,W_j^{\vee,\theta},E}^{(I_j)} & \longrightarrow & \mathbb{D} \mathcal{S}_{I_j,W_j,E}^{(I_j)}
 \end{array}$$

in $D_c^b(\text{Gr}_{I_j,W_j}^{(I_j)} / G_{\sum_{i \in I_j} n_i x_i}^{\text{ad}}, E)$, for each $1 \leq j \leq k$. Here Φ stands for the usual Frobenius correspondences, and the horizontal arrows except can are from Proposition 5.1.1. The square commutes by the functoriality of Φ .

for the triangular part, [Laszlo and Olsson 2008a, 4.8.2 Corollary] says that $\text{can} : \text{Frob}^* \mathbb{D} \xrightarrow{\sim} \mathbb{D} \text{Frob}^*$ equals

$$\text{Frob}^* \text{R}\mathcal{H}om(-, \Omega) \xrightarrow{\text{natural}} \text{R}\mathcal{H}om(\text{Frob}^*(-), \text{Frob}^* \Omega) \xrightarrow{f_*} \text{R}\mathcal{H}om(\text{Frob}^*(-), \Omega),$$

where Ω is the dualizing complex and $f : \text{Frob}^* \Omega \xrightarrow{\sim} \Omega$ is the canonical isomorphism furnished by [loc. cit.]. Both f and “can” reflect the fact that universal homeomorphisms conserve duality. In our case, that fact is also realized by transport of structure via Frobenius, i.e., we have $f = \Phi_\Omega$, the Frobenius correspondence for Ω . The desired commutativity thus reduces to that of

$$\begin{array}{ccc}
 \text{Frob}^* \text{R}\mathcal{H}om(\mathcal{S}, \Omega) & \xrightarrow{\text{natural}} & \text{R}\mathcal{H}om(\text{Frob}^* \mathcal{S}, \text{Frob}^* \Omega) \\
 \downarrow \Phi_{\text{R}\mathcal{H}om(\mathcal{S}, \Omega)} & \nwarrow (\Phi_\Omega^{-1})^* \circ (\Phi_\Omega)_* & \\
 \text{R}\mathcal{H}om(\mathcal{S}, \Omega) & &
 \end{array}$$

for all $\mathcal{S} \in D_c^b(\text{Gr}_{I_j,W_j}^{(I_j)} / G_{\sum_{i \in I_j} n_i x_i}^{\text{ad}}, E)$. This is by now standard. $\square$

Reintroduce the truncation parameters $\mu' \gg \mu$ so that (4-10) holds with respect to both W and $W^{\vee,\theta}$. Let a_1 (universal homeomorphism) and a_2 (open immersion) be as in (4-12). As μ increases, $\text{Cht}_{N,I,W}^{\dots, \leq \mu}$ and $\text{Cht}_{N,I,W}^{\dots, \leq \mu'}$ form open coverings of $\text{Cht}_{N,I,W}^{\dots}$.

To state the next result, we write $\Omega^{\leq \mu}$ and Ω for the normalized dualizing complex on $\text{Cht}_{N,I,W}^{\cdots, \leq \mu} / \Xi$ and $a_1^{-1} \text{Cht}_{N,I,W}^{(I_2, \dots, I_1), \leq \mu} / \Xi$, respectively. Recall that dualizing complexes are unique up to unique isomorphisms [Laszlo and Olsson 2008a, 3.4.5]. There are canonical isomorphisms $a_1^* \Omega^{\leq \mu} \xrightarrow{\sim} \Omega \xleftarrow{\sim} a_2^! \Omega^{\leq \mu'}$, since $a_1^* = a_1^!$.

Lemma 5.2.2. *In $D_c^b(a_1^{-1} \text{Cht}_{N,I,W}^{(I_2, \dots, I_1), \leq \mu} / \Xi, E)$, there is a commutative diagram*

$$\begin{array}{ccc}
 a_1^* \underbrace{\mathcal{F}_{N,I,W^{\vee,\theta},\Xi,E}^{(I_2, \dots, I_1)}}_{\text{on } \leq \mu} \overset{\mathbb{L}}{\otimes} a_1^* \underbrace{\mathcal{F}_{N,I,W,\Xi,E}^{(I_2, \dots, I_1)}}_{\text{on } \leq \mu} & \longrightarrow & a_1^* \Omega^{\leq \mu} \\
 \downarrow \text{Frob}_{I_1,N,W^{\vee,\theta}}^{(I_1, \dots, I_k)} \overset{\mathbb{L}}{\otimes} \text{Frob}_{I_1,N,W}^{(I_1, \dots, I_k)} & & \downarrow \cong \\
 a_2^! \underbrace{\mathcal{F}_{N,I,W^{\vee,\theta},\Xi,E}^{(I_1, \dots, I_k)}}_{\text{on } \leq \mu'} \overset{\mathbb{L}}{\otimes} a_2^! \underbrace{\mathcal{F}_{N,I,W,\Xi,E}^{(I_1, \dots, I_k)}}_{\text{on } \leq \mu'} & \longrightarrow & a_2^! \Omega^{\leq \mu'}
 \end{array}$$

$\begin{array}{ccc} & \searrow \cong & \\ & \Omega & \\ & \nearrow \cong & \end{array}$

where the arrows from $\cdots \overset{\mathbb{L}}{\otimes} \cdots$ to Ω are induced from Lemma 5.2.1.

Proof. It suffices to show the commutativity of the outer pentagon, since the triangle is defined to be commutative. Recall the passage from (4-11) to (4-13): $\text{Frob}_{I_1,N,\dots}^{(I_1, \dots, I_k)}$ is obtained by restricting $F_{I_1,N,\dots}^{(I_1, \dots, I_k)}$ to the open substacks cut out by the conditions $\leq \mu$ and $\leq \mu'$. It remains to apply Lemma 5.2.1; note that the effect of arrows $a_1^* \Omega^{\leq \mu} \xrightarrow{\sim} \Omega \xleftarrow{\sim} a_2^! \Omega^{\leq \mu'}$ match the morphism “can” in Lemma 5.2.1. $\square$

Proposition 5.2.3. *Write $J := I_1$. There is a commutative diagram in $D_c^b((X \setminus \hat{N}), E)$*

$$\begin{array}{ccc}
 \text{Frob}_J^* \mathcal{H}_{N,I,W^{\vee,\theta}}^{\leq \mu, E} \overset{\mathbb{L}}{\otimes} \text{Frob}_J^* \mathcal{H}_{N,I,W}^{\leq \mu, E} & \xrightarrow{\text{Frob}_J^* (5-1)} & \text{Frob}_J^* E_{(X \setminus \hat{N})^I} \\
 \downarrow F_J \overset{\mathbb{L}}{\otimes} F_J & & \downarrow F_J \\
 \mathcal{H}_{N,I,W^{\vee,\theta}}^{\leq \mu', E} \overset{\mathbb{L}}{\otimes} \mathcal{H}_{N,I,W}^{\leq \mu', E} & \xrightarrow{(5-1)} & E_{(X \setminus \hat{N})^I}
 \end{array}$$

where

- $F_J \overset{\mathbb{L}}{\otimes} F_J$ is induced from the F_J in (4-15),
- the F_J on the right is the evident partial Frobenius morphism for $E_{(X \setminus \hat{N})^I}$.

Proof. Retain the notation for Lemma 5.2.2 and let $\mathfrak{p} := \mathfrak{p}_{N,I}^{(I_1, \dots, I_k)}$. Upon recalling the formalism of $\text{K}\tilde{\mathcal{A}}_4$ -nneth formula, cup products (Remark 5.1.3) and the trace maps Tr (see (5-1)), Lemma 5.2.2 produce a diagram in $D_c^b((X \setminus \hat{N})^I, E)$:

$$\begin{array}{ccccc}
 \text{Frob}_J^* \mathfrak{p}_! \underbrace{\mathcal{F}_{N,I,W^{\vee,\theta},\Xi,E}^{(I_2,\dots,I_1)}}_{\text{on } \leq \mu} \overset{\mathbb{L}}{\otimes} \text{Frob}_J^* \mathfrak{p}_! \underbrace{\mathcal{F}_{N,I,W,\Xi,E}^{(I_2,\dots,I_1)}}_{\text{on } \leq \mu} & \longrightarrow & \text{Frob}_J^* \mathfrak{p}_! \Omega^{\leq \mu} & \xrightarrow{\text{Frob}_J^* \text{Tr}} & \text{Frob}_J^* E_{(X \setminus \hat{N})^I} \\
 \downarrow \text{BC} \overset{\mathbb{L}}{\otimes} \text{BC} \simeq & & \downarrow \simeq \text{BC} & & \downarrow \simeq F_J \\
 \mathfrak{p}_! a_1^* \mathcal{F}_{N,I,W^{\vee,\theta},\Xi,E}^{(I_2,\dots,I_1)} \overset{\mathbb{L}}{\otimes} \mathfrak{p}_! a_1^* \mathcal{F}_{N,I,W,\Xi,E}^{(I_2,\dots,I_1)} & \longrightarrow & \mathfrak{p}_! a_1^* \Omega^{\leq \mu} \simeq \mathfrak{p}_! \Omega & \xrightarrow{\text{Tr}} & E_{(X \setminus \hat{N})^I} \\
 \downarrow \mathfrak{p}_! \text{Frob}_{I_1,N,W^{\vee,\theta}}^{(I_1,\dots,I_k)} \overset{\mathbb{L}}{\otimes} \mathfrak{p}_! \text{Frob}_{I_1,N,W}^{(I_1,\dots,I_k)} & & \parallel & & \uparrow \text{Tr} \\
 \mathfrak{p}_! a_2^! \mathcal{F}_{N,I,W^{\vee,\theta},\Xi,E}^{(I_1,\dots,I_k)} \overset{\mathbb{L}}{\otimes} \mathfrak{p}_! a_2^! \mathcal{F}_{N,I,W,\Xi,E}^{(I_1,\dots,I_k)} & \longrightarrow & \mathfrak{p}_! a_2^! \Omega^{\leq \mu'} \simeq \mathfrak{p}_! \Omega & & \\
 \downarrow & & \downarrow & & \\
 \mathfrak{p}_! \underbrace{\mathcal{F}_{N,I,W^{\vee,\theta},\Xi,E}^{(I_1,\dots,I_k)}}_{\text{on } \leq \mu'} \overset{\mathbb{L}}{\otimes} \mathfrak{p}_! \underbrace{\mathcal{F}_{N,I,W,\Xi,E}^{(I_1,\dots,I_k)}}_{\text{on } \leq \mu'} & \longrightarrow & \mathfrak{p}_! \Omega^{\leq \mu'} & &
 \end{array}$$

where BC and $\mathfrak{p}_! a_2^! \rightarrow \mathfrak{p}_!$ are the arrows in (4-14) and explained after (4-15), respectively. The diagram commutes, indeed:

- The first two rows form a commutative diagram by the naturality of BC , which is ultimately based on the topological invariance of the étale topos together with the fact that universal homeomorphisms respect duality [Laszlo and Olsson 2008b, 9.1.5 Proposition and 12.2].
- The commutativity of the middle square comes from Lemma 5.2.2, by applying $\mathfrak{p}_!$.
- The remaining pieces commute by the naturality of $\mathfrak{p}_! a_2^! \rightarrow \mathfrak{p}_!$ and of Tr .

The composite of the last row is (5-1), and that of the first row is its Frob_J^* -image (now for the $\leq \mu$ part). The composite of the leftmost column yields $F_J \overset{\mathbb{L}}{\otimes} F_J : \text{Frob}_J^* \mathcal{H}_{N,I,W^{\vee,\theta}}^{\leq \mu,E} \overset{\mathbb{L}}{\otimes} \text{Frob}_J^* \mathcal{H}_{N,I,W}^{\leq \mu,E} \rightarrow \mathcal{H}_{N,I,W^{\vee,\theta}}^{\leq \mu',E} \overset{\mathbb{L}}{\otimes} \mathcal{H}_{N,I,W}^{\leq \mu',E}$ by the very definition of F_J . This completes the proof. $\square$

The case $k = 1$, i.e., when F_J is the total Frobenius morphism, is relatively straightforward; see the proof of Lemma 5.2.1.

Recall from Section 4.3 that F_J furnishes an E -linear endomorphism of $\varinjlim_{\mu} \mathcal{H}_{N,I,W}^{0,\leq \mu,E} |_{\eta^I}$, still denoted as F_J .

Corollary 5.2.4. *The pairing $\langle \cdot, \cdot \rangle_{\eta^I}$ in (5-2) is invariant under F_J for all $J \subset I$.*

Proof. After taking H^0 and $\varinjlim_{\mu}$, Proposition 5.2.3 implies that

$$\langle h_1, h_2 \rangle_{\eta^I} = \langle F_J(h_1), F_J(h_2) \rangle_{\eta^I}$$

for all h_1, h_2 in $\varinjlim_{\mu} \mathcal{H}_{N,I,W^{\vee,\theta}}^{0,\leq \mu,E} |_{\eta^I}$ and $\varinjlim_{\mu} \mathcal{H}_{N,I,W}^{0,\leq \mu,E} |_{\eta^I}$, respectively. $\square$

The cautious reader might worry about a missing power of p in Corollary 5.2.4 due to Tate twists. It does not occur here by our normalizations of $\mathcal{S}$, $\mathcal{F}$ and $\mathbb{D}$.

5.3. Computation of the transpose. We adopt the notation of Section 4.4. The integration pairing $\langle \cdot, \cdot \rangle$ of Remark 2.2.3 is nondegenerate symmetric on the finite-dimensional E -vector space $H_{\{0\},1} \simeq H_{\emptyset,1}$. The transpose S^* of any $S \in \text{End}_E(H_{\{0\},1})$, characterized by $\langle h', Sh \rangle = \langle S^*h', h \rangle$ for all $h, h' \in H_{\{0\},1}$. Identify $\langle \cdot, \cdot \rangle$ with the pairing $\langle \cdot, \cdot \rangle_{\Delta(\bar{\eta})}$ in (5-2).

Lemma 5.3.1. *For all data I, W, ξ, x and $\vec{\gamma} = (\gamma_i)_i \in \pi_1(\eta, \bar{\eta})^I$ for excursion operators, we have*

$$S_{I,W,\xi,x,\vec{\gamma}}^* = S_{I,W^{\vee,\theta},x,\xi,\vec{\gamma}^{-1}}.$$

In particular, the E -algebra $\mathcal{B}_E$ is closed under transpose $S \mapsto S^$.*

Note that the roles of x, ξ are switched when one passes from W to $W^{\vee,\theta}$. The transpose-invariance of $\mathcal{B}_E$ has already been sketched in [Lafforgue 2018, Remarque 12.15].

Proof. Recall from Section 4.3 that by choosing $\bar{\eta}^I$ and $\mathfrak{sp}$, there is a homomorphism $\text{FWeil}(\eta^I, \bar{\eta}^I) \rightarrow \mathbb{W}_{\bar{F}}^I$ inducing an isomorphism between profinite completions. As $S_{I,W,\xi,x,\vec{\gamma}}$ and $S_{I,W^{\vee,\theta},x,\xi,\vec{\gamma}^{-1}}$ are both continuous in $\vec{\gamma}$, it suffices to consider that case when $\vec{\gamma}$ comes from $\text{FWeil}(\eta^I, \bar{\eta}^I)$.

By [Lafforgue 2018, Remarque 5.4, (9.8)], $\mathcal{C}_{\xi}^b$ and $\mathcal{C}_{\xi}^{\sharp}$ are already transposes of each other on the sheaf level with respect to $\mathfrak{B}_{N,I,W}^{\Xi,E}$; in particular they commute with $\mathfrak{sp}^*$. Ditto for $\mathcal{C}_x^{\sharp}$ and $\mathcal{C}_x^b$. Therefore, for all $h, h' \in H_{\{0\},1}$, we infer by using (5-2) that

$$\begin{aligned} \langle h', S_{I,W,x,\xi,\vec{\gamma}}(h) \rangle_{\Delta(\bar{\eta})} &= \langle h', \mathcal{C}_{\xi}^b(\mathfrak{sp}^*)^{-1}(\vec{\gamma} \cdot \mathfrak{sp}^* \mathcal{C}_x^{\sharp} h) \rangle_{\Delta(\bar{\eta})} \\ &= \langle \mathfrak{sp}^*(h'), \mathfrak{sp}^*(\mathcal{C}_{\xi}^b(\mathfrak{sp}^*)^{-1}(\vec{\gamma} \cdot \mathfrak{sp}^* \mathcal{C}_x^{\sharp} h)) \rangle_{\bar{\eta}^I} \\ &= \langle \mathfrak{sp}^* \mathcal{C}_{\xi}^{\sharp}(h'), \vec{\gamma} \cdot \mathfrak{sp}^* \mathcal{C}_x^{\sharp}(h) \rangle_{\bar{\eta}}, \\ \langle S_{I,W^{\vee,\theta},x,\xi,\vec{\gamma}^{-1}}(h'), h \rangle_{\Delta(\bar{\eta})} &= \langle \mathcal{C}_x^b(\mathfrak{sp}^*)^{-1}(\vec{\gamma}^{-1} \cdot \mathfrak{sp}^* \mathcal{C}_{\xi}^{\sharp} h'), h \rangle_{\Delta(\bar{\eta})} \\ &= \langle \mathfrak{sp}^*(\mathcal{C}_x^b(\mathfrak{sp}^*)^{-1}(\vec{\gamma}^{-1} \cdot \mathfrak{sp}^* \mathcal{C}_{\xi}^{\sharp} h')), \mathfrak{sp}^*(h) \rangle_{\bar{\eta}^I} \\ &= \langle \vec{\gamma}^{-1} \cdot \mathfrak{sp}^* \mathcal{C}_{\xi}^{\sharp}(h'), \mathfrak{sp}^* \mathcal{C}_x^{\sharp}(h) \rangle_{\bar{\eta}^I}. \end{aligned}$$

It remains to show that $\langle \cdot, \cdot \rangle_{\bar{\eta}^I}$ is $\text{FWeil}(\eta^I, \bar{\eta}^I)$ -invariant. Recall that the $\text{FWeil}(\eta^I, \bar{\eta}^I)$ -action unites those from $\pi_1(\eta^I, \bar{\eta}^I)$ and partial Frobenius morphisms F_J . The $\pi_1(\eta^I, \bar{\eta}^I)$ -action leaves $\langle \cdot, \cdot \rangle_{\bar{\eta}^I}$ invariant since the latter comes from the sheaf-level pairing $\mathfrak{B}_{N,I,W}^{\Xi,E}$ over $(X \setminus \hat{N})^I$. The F_J -invariance of $\langle \cdot, \cdot \rangle$ for all $J \subset I$ is assured by Corollary 5.2.4. $\square$

We are now able to describe the transpose of excursion operators.

Definition 5.3.2. For every $f \in \mathcal{O}(\hat{G} \llbracket {}^L G \rrbracket^I \parallel \hat{G})$, set $f^{\dagger}(\vec{g}) := f({}^L \theta(\vec{g}^{-1}))$ where $\vec{g} \in ({}^L G)^I$ and ${}^L \theta$ stands for the Chevalley involution of $({}^L G)^I$. Then $f \mapsto f^{\dagger}$ defines an involution of $\mathcal{O}(\hat{G} \llbracket {}^L G \rrbracket^I \parallel \hat{G})$.

Lemma 5.3.3. *For all $I, f \in \mathcal{O}(\hat{G} \llbracket {}^L G \rrbracket^I \parallel \hat{G})$ and $\vec{\gamma} = (\gamma_i)_i \in \pi_1(\eta, \bar{\eta})^I$, we have $S_{I,f,\vec{\gamma}}^* = S_{I,f^{\dagger},\vec{\gamma}^{-1}}$.*

Proof. It suffices to consider the case $f(\vec{g}) = \langle \xi, \vec{g} \cdot x \rangle_{W^{\vee} \otimes W}$, where $\xi \in W^{\vee}, x \in W$ are as in Lemma 5.3.1, and $\langle \cdot, \cdot \rangle_{W^{\vee} \otimes W}$ is the evident duality pairing. As before, denote by $W^{\theta}, W^{\vee,\theta}$ be the ${}^L \theta$ -twists of the

representations W , $W^\vee$ etc., and the preceding convention on pairing still applies. Note that $(W^{\vee,\theta})^\vee \simeq W^\theta$ canonically in $\text{Rep}_E({}^L G)^I$.

For every $\vec{g} \in ({}^L G)^I$, we have

$$\begin{aligned} f^\dagger(\vec{g}) &= \langle \xi, \underbrace{{}^L\theta(g)^{-1} \cdot x}_{\text{original action}} \rangle_{W^\vee \otimes W} \\ &= \langle \xi, \underbrace{\vec{g}^{-1} \cdot x}_{{}^L\theta\text{-twisted}} \rangle_{W^{\theta,\vee} \otimes W^\theta} \\ &= \underbrace{\langle \vec{g} \cdot \xi \rangle}_{{}^L\theta\text{-twisted}}, x_{W^{\theta,\vee} \otimes W^\theta} \\ &= \langle x, \vec{g} \cdot \xi \rangle_{(W^{\vee,\theta})^\vee \otimes W^{\vee,\theta}}. \end{aligned}$$

In view of Lemma 5.3.1, we deduce that $S_{I,f,\vec{\gamma}}^* = S_{I,W,\xi,x,\vec{\gamma}}^*$ equals $S_{I,W^{\vee,\theta},x,\xi,\vec{\gamma}^{-1}} = S_{I,f^\dagger,\vec{\gamma}^{-1}}$, as asserted. $\square$

Consider any homomorphism $\nu : \mathcal{B} := \mathcal{B}_E \otimes_E \overline{\mathbb{Q}_\ell} \rightarrow \overline{\mathbb{Q}_\ell}$ of $\overline{\mathbb{Q}_\ell}$ -algebras. As $\mathcal{B}$ is commutative and closed under transpose, $\nu^* : S \mapsto \nu(S^*)$ is also a homomorphism of $\overline{\mathbb{Q}_\ell}$ -algebras.

Proposition 5.3.4. *If $\sigma \in \Phi(G)$ is attached to $\nu : \mathcal{B} \rightarrow \overline{\mathbb{Q}_\ell}$, then ${}^L\theta \circ \sigma$ is attached to ν^* .*

Proof. Fix $n \in \mathbb{Z}_{\geq 0}$ and let $I := \{0, \dots, n\}$. Given the characterization (4-19) of the L -parameters attached to ν , ν^* , it boils down to the observation that for all $\vec{\gamma} = (\gamma_0, \dots, \gamma_n) \in \pi_1(\eta, \bar{\eta})^I$ and $f \in \mathcal{O}(\hat{G} \parallel {}^L G^I \parallel \hat{G})$,

$$\nu^*(S_{I,f,\vec{\gamma}}) = \nu(S_{I,f,\vec{\gamma}}^*) = \nu(S_{I,f^\dagger,\vec{\gamma}^{-1}}) = f^\dagger(\sigma(\gamma_0)^{-1}, \dots, \sigma(\gamma_n)^{-1}) = f({}^L\theta\sigma(\gamma_0), \dots, {}^L\theta\sigma(\gamma_n)),$$

in which the second equality stems from Lemma 5.3.3. $\square$

Write $\mathfrak{H}_\sigma := \mathfrak{H}_\nu$ if $\sigma \in \Phi(G)$ is attached to ν , and write $\langle \cdot, \cdot \rangle_{\sigma,\sigma'} := \langle \cdot, \cdot \rangle|_{\mathfrak{H}_\sigma \otimes \mathfrak{H}_{\sigma'}}$, for all $\sigma, \sigma' \in \Phi(G)$.

Proof of Theorem 3.3.2. Enlarge E so that all homomorphisms $\nu : \mathcal{B} \rightarrow \overline{\mathbb{Q}_\ell}$ are defined over E . Fix a ν such that $\mathfrak{H}_\nu \neq \{0\}$. Since $\mathcal{B}_E$ is closed under transpose, the subspace $\mathfrak{H}_\nu^\perp \subset H_{\{0\},1}$ defined relative to $\langle \cdot, \cdot \rangle$ is $\mathcal{B}_E$ -stable as well. Since $\langle \cdot, \cdot \rangle$ is nondegenerate, $\mathfrak{H}_\nu^\perp \neq H_{\{0\},1}$. Use the $\mathcal{B}_E$ -invariance to decompose $\mathcal{B}_E$ -modules as follows

$$\mathfrak{H}_\nu^\perp = \bigoplus_\mu \mathfrak{H}_\nu^\perp \cap \mathfrak{H}_\mu, \quad \frac{H_{\{0\},1}}{\mathfrak{H}_\nu^\perp} = \bigoplus_\mu \frac{\mathfrak{H}_\mu}{\mathfrak{H}_\nu^\perp \cap \mathfrak{H}_\mu} \neq \{0\}.$$

We contend that $\mathfrak{H}_\mu \not\subset \mathfrak{H}_\nu^\perp$ only if $\mu = \nu^*$, or equivalently $\mu^* = \nu$.

Indeed, $\langle \cdot, \cdot \rangle$ induces a nondegenerate pairing

$$\langle \cdot, \cdot \rangle_\nu : \mathfrak{H}_\nu \otimes_E \bigoplus_\mu \frac{\mathfrak{H}_\mu}{\mathfrak{H}_\nu^\perp \cap \mathfrak{H}_\mu} \rightarrow E.$$

Let $d := \dim H_{\{0\},1}$. For every $S \in \mathcal{B}_E$, write $S_v := S|_{\mathfrak{H}_v}$. Then $(S_v - v(S))^d = 0$. Taking transpose with respect to $\langle \cdot, \cdot \rangle_v$ yields $(S_v^* - v(S))^d = 0$ in $\text{End}_E(\mathfrak{H}_\mu/(\mathfrak{H}_\mu \cap \mathfrak{H}_v^\perp))$, for each μ .

On the other hand, the transpose $S^* \in \mathcal{B}_E$ with respect to $\langle \cdot, \cdot \rangle$ satisfies $(S^* - \mu(S^*))^d = 0$ on $\mathfrak{H}_\mu$, and $S^*|_{\mathfrak{H}_\mu}$ induces $S^{*,\mu} \in \text{End}_E(\mathfrak{H}_\mu/(\mathfrak{H}_\mu \cap \mathfrak{H}_v^\perp))$ satisfying $(S^{*,\mu} - \mu(S^*))^d = 0$. Clearly $S^{*,\mu} = S_v^*$. All in all, we deduce that $\mu^*(S) := \mu(S^*) = v(S)$ whenever $\mathfrak{H}_\mu \neq \mathfrak{H}_v^\perp \cap \mathfrak{H}_\mu$.

It follows from the claim that if $\langle \cdot, \cdot \rangle_{\sigma, \sigma'}$ is not identically zero, then the corresponding $v, v' : \mathcal{B} \rightarrow \overline{\mathbb{Q}_\ell}$ satisfy $v^* = v'$. Now Proposition 5.3.4 implies ${}^L\theta \circ \sigma = \sigma'$. $\square$

Acknowledgements

The author is deeply grateful to Alain Genestier, Vincent Lafforgue, Dipendra Prasad and Changjian Su for their helpful comments, answers and corrections. Thanks also goes to the referees for pertinent suggestions.

References

- [Adams and Vogan 2016] J. Adams and D. A. Vogan, Jr., “Contragredient representations and characterizing the local Langlands correspondence”, *Amer. J. Math.* **138**:3 (2016), 657–682. MR Zbl
- [Arthur 1988] J. Arthur, “The invariant trace formula, II: Global theory”, *J. Amer. Math. Soc.* **1**:3 (1988), 501–554. MR Zbl
- [Artin and Tate 1968] E. Artin and J. Tate, *Class field theory*, W. A. Benjamin, New York, 1968. MR Zbl
- [Bezrukavnikov and Finkelberg 2008] R. Bezrukavnikov and M. Finkelberg, “Equivariant Satake category and Kostant–Whittaker reduction”, *Mosc. Math. J.* **8**:1 (2008), 39–72. MR Zbl
- [Borel 1979] A. Borel, “Automorphic L -functions”, pp. 27–61 in *Automorphic forms, representations and L -functions, II*, edited by A. Borel and W. Casselman, Proc. Sympos. Pure Math. **33**, Amer. Math. Soc., Providence, RI, 1979. MR Zbl
- [Bourbaki 2012] N. Bourbaki, *Algèbre, Chapitre VIII*, 2nd revised ed., Springer, 2012. MR Zbl
- [Braverman and Gaitsgory 2002] A. Braverman and D. Gaitsgory, “Geometric Eisenstein series”, *Invent. Math.* **150**:2 (2002), 287–384. MR Zbl
- [Brylinski and Deligne 2001] J.-L. Brylinski and P. Deligne, “Central extensions of reductive groups by $\mathbb{K}_2$ ”, *Publ. Math. Inst. Hautes Études Sci.* **94** (2001), 5–85. MR Zbl
- [Bushnell and Henniart 2006] C. J. Bushnell and G. Henniart, *The local Langlands conjecture for $\text{GL}(2)$* , Grundlehren der Math. Wissenschaften **335**, Springer, 2006. MR Zbl
- [Fargues 2016] L. Fargues, “Geometrization of the local Langlands correspondence: an overview”, preprint, 2016. arXiv
- [Gaitsgory and Lysenko 2018] D. Gaitsgory and S. Lysenko, “Parameters and duality for the metaplectic geometric Langlands theory”, *Selecta Math. (N.S.)* **24**:1 (2018), 227–301. MR Zbl
- [Genestier and Lafforgue 2017] A. Genestier and V. Lafforgue, “Chtoucas restreints pour les groupes réductifs et paramétrisation de Langlands locale”, preprint, 2017. arXiv
- [Gross and Prasad 1992] B. H. Gross and D. Prasad, “On the decomposition of a representation of SO_n when restricted to SO_{n-1} ”, *Canad. J. Math.* **44**:5 (1992), 974–1002. MR Zbl
- [Haines and Rostami 2010] T. J. Haines and S. Rostami, “The Satake isomorphism for special maximal parahoric Hecke algebras”, *Represent. Theory* **14** (2010), 264–284. MR Zbl
- [Heinloth 2010] J. Heinloth, “Uniformization of $\mathcal{G}$ -bundles”, *Math. Ann.* **347**:3 (2010), 499–528. MR Zbl
- [Henniart 1983] G. Henniart, *La conjecture de Langlands locale pour $\text{GL}(3)$* , Mém. Soc. Math. France **11-12**, Soc. Math. France, Paris, 1983. MR Zbl
- [Kaletha 2013] T. Kaletha, “Genericity and contragredience in the local Langlands correspondence”, *Algebra Number Theory* **7**:10 (2013), 2447–2474. MR Zbl

- [Kaletha 2016a] T. Kaletha, “Regular supercuspidal representations”, preprint, 2016. [arXiv](#)
- [Kaletha 2016b] T. Kaletha, “Rigid inner forms of real and p -adic groups”, *Ann. of Math. (2)* **184**:2 (2016), 559–632. [MR](#) [Zbl](#)
- [Kashiwara and Schapira 1990] M. Kashiwara and P. Schapira, *Sheaves on manifolds*, Grundlehren der Math. Wissenschaften **292**, Springer, 1990. [MR](#) [Zbl](#)
- [Lafforgue 2018] V. Lafforgue, “Chtoucas pour les groupes réductifs et paramétrisation de Langlands globale”, *J. Amer. Math. Soc.* **31**:3 (2018), 719–891. [MR](#) [Zbl](#)
- [Lapid and Mao 2015] E. Lapid and Z. Mao, “A conjecture on Whittaker–Fourier coefficients of cusp forms”, *J. Number Theory* **146** (2015), 448–505. [MR](#) [Zbl](#)
- [Laszlo and Olsson 2008a] Y. Laszlo and M. Olsson, “The six operations for sheaves on Artin stacks, I: Finite coefficients”, *Publ. Math. Inst. Hautes Études Sci.* **107** (2008), 109–168. [MR](#) [Zbl](#)
- [Laszlo and Olsson 2008b] Y. Laszlo and M. Olsson, “The six operations for sheaves on Artin stacks, II: Adic coefficients”, *Publ. Math. Inst. Hautes Études Sci.* **107** (2008), 169–210. [MR](#) [Zbl](#)
- [Laszlo and Olsson 2009] Y. Laszlo and M. Olsson, “Perverse t -structure on Artin stacks”, *Math. Z.* **261**:4 (2009), 737–748. [MR](#) [Zbl](#)
- [Mirković and Vilonen 2007] I. Mirković and K. Vilonen, “Geometric Langlands duality and representations of algebraic groups over commutative rings”, *Ann. of Math. (2)* **166**:1 (2007), 95–143. [MR](#) [Zbl](#)
- [Mœglin and Waldspurger 1994] C. Mœglin and J.-L. Waldspurger, *Décomposition spectrale et séries d’Eisenstein*, Progress in Math. **113**, Birkhäuser, Basel, 1994. [MR](#) [Zbl](#)
- [Mœglin et al. 1987] C. Mœglin, M.-F. Vignéras, and J.-L. Waldspurger, *Correspondances de Howe sur un corps p -adique*, Lecture Notes in Math. **1291**, Springer, 1987. [MR](#) [Zbl](#)
- [Prasad 2018] D. Prasad, “Generalizing the MVW involution, and the contragredient”, *Trans. Amer. Math. Soc.* (online publication November 2018).
- [Renard 2010] D. Renard, *Représentations des groupes réductifs p -adiques*, Cours Spécialisés **17**, Soc. Math. France, Paris, 2010. [MR](#) [Zbl](#)
- [Richarz 2014] T. Richarz, “A new approach to the geometric Satake equivalence”, *Doc. Math.* **19** (2014), 209–246. [MR](#) [Zbl](#)
- [Serre 2005] J.-P. Serre, “Complète réductibilité”, exposé 932, pp. 195–217 in *Séminaire Bourbaki*, 2003/2004, Astérisque **299**, 2005. [MR](#) [Zbl](#)
- [SGA 4₁ 1972] M. Artin, A. Grothendieck, and J. L. Verdier, *Théorie des topos et cohomologie étale des schémas, Tome 1: Théorie des topos, Exposés I–IV* (Séminaire de Géométrie Algébrique du Bois Marie 1963–1964), Lecture Notes in Math. **269**, Springer, 1972. [MR](#) [Zbl](#)
- [SGA 4₂ 1972] M. Artin, A. Grothendieck, and J. L. Verdier, *Théorie des topos et cohomologie étale des schémas, Tome 2: Exposés V–VIII* (Séminaire de Géométrie Algébrique du Bois Marie 1963–1964), Lecture Notes in Math. **270**, Springer, 1972. [MR](#) [Zbl](#)
- [SGA 4₃ 1973] M. Artin, A. Grothendieck, and J. L. Verdier, *Théorie des topos et cohomologie étale des schémas, Tome 3: Exposés IX–XIX* (Séminaire de Géométrie Algébrique du Bois Marie 1963–1964), Lecture Notes in Math. **305**, Springer, 1973. [MR](#) [Zbl](#)
- [Shahidi 1990] F. Shahidi, “A proof of Langlands’ conjecture on Plancherel measures; complementary series for p -adic groups”, *Ann. of Math. (2)* **132**:2 (1990), 273–330. [MR](#) [Zbl](#)
- [Stacks 2005–] P. Belmans, A. J. de Jong, et al., “The Stacks project”, electronic reference, 2005–, Available at <http://stacks.math.columbia.edu>.
- [Varshavsky 2004] Y. Varshavsky, “Moduli spaces of principal F -bundles”, *Selecta Math. (N.S.)* **10**:1 (2004), 131–166. [MR](#) [Zbl](#)
- [Varshavsky 2007] Y. Varshavsky, “Lefschetz–Verdier trace formula and a generalization of a theorem of Fujiwara”, *Geom. Funct. Anal.* **17**:1 (2007), 271–319. [MR](#) [Zbl](#)
- [Vignéras 1996] M.-F. Vignéras, *Représentations l -modulaires d’un groupe réductif p -adique avec $l \neq p$* , Progress in Math. **137**, Birkhäuser, Boston, 1996. [MR](#) [Zbl](#)

[Vignéras 2001] M.-F. Vignéras, “Correspondance de Langlands semi-simple pour $GL(n, F)$ modulo $l \neq p$ ”, *Invent. Math.* **144**:1 (2001), 177–223. MR Zbl

[Zhu 2015] X. Zhu, “The geometric Satake correspondence for ramified groups”, *Ann. Sci. Éc. Norm. Supér. (4)* **48**:2 (2015), 409–451. MR Zbl

Communicated by Marie-France Vignéras

Received 2018-10-16 Revised 2019-01-09 Accepted 2019-03-10

wwli@bicmr.pku.edu.cn

*Beijing International Center for Mathematical Research, Peking University,
Beijing, China*

Algebra & Number Theory

msp.org/ant

EDITORS

MANAGING EDITOR

Bjorn Poonen
Massachusetts Institute of Technology
Cambridge, USA

EDITORIAL BOARD CHAIR

David Eisenbud
University of California
Berkeley, USA

BOARD OF EDITORS

Richard E. Borcherds	University of California, Berkeley, USA	Raman Parimala	Emory University, USA
Antoine Chambert-Loir	Université Paris-Diderot, France	Jonathan Pila	University of Oxford, UK
J-L. Colliot-Thélène	CNRS, Université Paris-Sud, France	Anand Pillay	University of Notre Dame, USA
Brian D. Conrad	Stanford University, USA	Michael Rapoport	Universität Bonn, Germany
Samit Dasgupta	University of California, Santa Cruz, USA	Victor Reiner	University of Minnesota, USA
Hélène Esnault	Freie Universität Berlin, Germany	Peter Sarnak	Princeton University, USA
Gavril Farkas	Humboldt Universität zu Berlin, Germany	Joseph H. Silverman	Brown University, USA
Hubert Flenner	Ruhr-Universität, Germany	Michael Singer	North Carolina State University, USA
Sergey Fomin	University of Michigan, USA	Christopher Skinner	Princeton University, USA
Edward Frenkel	University of California, Berkeley, USA	Vasudevan Srinivas	Tata Inst. of Fund. Research, India
Andrew Granville	Université de Montréal, Canada	J. Toby Stafford	University of Michigan, USA
Joseph Gubeladze	San Francisco State University, USA	Pham Huu Tiep	University of Arizona, USA
Roger Heath-Brown	Oxford University, UK	Ravi Vakil	Stanford University, USA
Craig Huneke	University of Virginia, USA	Michel van den Bergh	Hasselt University, Belgium
Kiran S. Kedlaya	Univ. of California, San Diego, USA	Akshay Venkatesh	Institute for Advanced Study, USA
János Kollár	Princeton University, USA	Marie-France Vignéras	Université Paris VII, France
Philippe Michel	École Polytechnique Fédérale de Lausanne	Kei-Ichi Watanabe	Nihon University, Japan
Susan Montgomery	University of Southern California, USA	Melanie Matchett Wood	University of Wisconsin, Madison, USA
Shigefumi Mori	RIMS, Kyoto University, Japan	Shou-Wu Zhang	Princeton University, USA
Martin Olsson	University of California, Berkeley, USA		

PRODUCTION

production@msp.org
Silvio Levy, Scientific Editor

See inside back cover or msp.org/ant for submission instructions.

The subscription price for 2019 is US \$/year for the electronic version, and \$/year (+\$, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to MSP.

Algebra & Number Theory (ISSN 1944-7833 electronic, 1937-0652 printed) at Mathematical Sciences Publishers, 798 Evans Hall #3840, c/o University of California, Berkeley, CA 94720-3840 is published continuously online. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices.

ANT peer review and production are managed by EditFlow® from MSP.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2019 Mathematical Sciences Publishers

Algebra & Number Theory

Volume 13 No. 5 2019

Surjectivity of Galois representations in rational families of abelian varieties	995
AARON LANDESMAN, ASHVIN A. SWAMINATHAN, JAMES TAO and YUJIE XU	
A unified and improved Chebotarev density theorem	1039
JESSE THORNER and ASIF ZAMAN	
On the Brauer–Siegel ratio for abelian varieties over function fields	1069
DOUGLAS ULMER	
A five-term exact sequence for Kac cohomology	1121
CÉSAR GALINDO and YIBY MORALES	
On the paramodularity of typical abelian surfaces	1145
ARMAND BRUMER, ARIEL PACETTI, CRIS POOR, GONZALO TORNARÍA, JOHN VOIGHT and DAVID S. YUEN	
Contragredient representations over local fields of positive characteristic	1197
WEN-WEI LI	



1937-0652(2019)13:5;1-8