

Algebra & Number Theory

Volume 13
2019
No. 6

Examples of hypergeometric twistor \mathcal{D} -modules

Alberto Castaño Domínguez, Thomas Reichelt and Christian Sevenheck



Examples of hypergeometric twistor \mathcal{D} -modules

Alberto Castaño Domínguez, Thomas Reichelt and Christian Sevenheck

We show that certain one-dimensional hypergeometric differential systems underlie objects of the category of irregular mixed Hodge modules, which was recently introduced by Sabbah, and compute the irregular Hodge filtration for them. We also provide a comparison theorem between two different types of Fourier–Laplace transformation for algebraic integrable twistor \mathcal{D} -modules.

1. Introduction

In a series of papers Sabbah and Yu (partly joint with Esnault) [Yu 2014; Sabbah and Yu 2015; Esnault et al. 2017; Sabbah 2018] considered a so-called irregular Hodge filtration on certain cohomology groups and on certain irregular \mathcal{D} -modules. It can be seen as a generalization of the Hodge filtration on a mixed Hodge module in the sense of M. Saito. Geometrically, such a filtration arises by considering a version of the twisted de Rham cohomology of certain proper maps, and it plays (conjecturally) a role in Hodge theoretic mirror symmetry (see [Katzarkov et al. 2017]). Sabbah [2018] has defined a category of irregular mixed Hodge modules, which is (up to a technical equivalence) a certain subcategory of T. Mochizuki’s category of (integrable) mixed twistor \mathcal{D} -modules. He has proved that a rigid irreducible \mathcal{D} -module on the projective line can be uniquely upgraded to an irregular Hodge module if and only if its formal local monodromies are unitary. Consequently, these objects come equipped with an irregular Hodge filtration and one can define irregular Hodge numbers for them. They should be seen as interesting numerical invariants attached to these differential systems, contrary to the case of arbitrary mixed twistor \mathcal{D} -modules, where there is no obvious way to define such numbers. In [Castaño Domínguez and Sevenheck 2019], the first and the third named author have computed that filtration and its corresponding numbers for the purely irregular hypergeometric modules, that is for systems of the form $\mathcal{D}_{\mathbb{G}_m}/\mathcal{D}_{\mathbb{G}_m}P$, where P is the operator

$$P = \prod_{i=1}^n (t\partial_t - \alpha_i) - t$$

The authors are partially supported by the project SISYPH: ANR-13-IS01-0001-01/02 and DFG grant SE 1114/5-1. Castaño Domínguez is also partially supported by MTM2014-59456-P, the ERDF and GI-2136. Reichelt is supported by a DFG Emmy Noether Fellowship (RE 3567/1-1).

MSC2010: primary 14F10; secondary 32C38.

Keywords: \mathcal{D} -modules, irregular Hodge filtration, twistor \mathcal{D} -modules, Fourier–Laplace transformation, hypergeometric \mathcal{D} -modules.

for real numbers $\alpha_1, \dots, \alpha_n$. Let us consider the noncommutative ring $R_{\mathbb{G}_m}^{\text{int}} := \mathbb{C}[z, t^{\pm}] \langle z^2 \partial_z, tz \partial_t \rangle$. A crucial point was to show that a certain quotient of the corresponding sheaf $\mathcal{R}_{\mathbb{G}_m}^{\text{int}}$ on \mathbb{G}_m which restricts to the $\mathcal{D}_{\mathbb{G}_m, t}$ -module $\mathcal{D}_{\mathbb{G}_m} / \mathcal{D}_{\mathbb{G}_m} P$ on $z = 1$ actually underlies an object in the category $\text{IrrMHM}(\mathbb{G}_m)$ and the latter can be uniquely extended to an object in $\text{IrrMHM}(\mathbb{P}^1)$.

In this paper we discuss the case of more general hypergeometric \mathcal{D} -module, that is, for quotients $\mathcal{D}_{\mathbb{G}_m} / \mathcal{D}_{\mathbb{G}_m} P$, where now P is of the form

$$P = \prod_{i=1}^n (t \partial_t - \alpha_i) - t \prod_{j=1}^m (t \partial_t - \beta_j)$$

for positive integers m, n and real numbers $\alpha_1, \dots, \alpha_n, \beta_1, \dots, \beta_m$ such that there is no integer difference between any α_i and β_j (this is the irreducibility assumption). It is worth noticing that the presence of the factor $\prod_{j=1}^m (t \partial_t - \beta_j)$ rules out the usage of the geometric arguments of [Castaño Domínguez and Sevenheck 2019]. We obtain (see Theorem 5.7) that for certain such systems, the corresponding quotient of $\mathcal{R}_{\mathbb{G}_m}^{\text{int}}$ still underlies an object of $\text{IrrMHM}(\mathbb{G}_m)$. As an application, we can completely determine the irregular Hodge filtration for all systems $\mathcal{D} / \mathcal{D} P$ as above, where n is arbitrary and where $m = 1$.

The strategy of the proof (which is rather different from that of [Castaño Domínguez and Sevenheck 2019]) of the main theorem is to reduce these differential systems from (Fourier–Laplace transformed) A-hypergeometric \mathcal{D} -modules (the so-called GKZ-systems of Gelfand, Graev, Zelevinski and Kapranov, see [Gelfand et al. 1987; Gelfand et al. 1989]), but at the level of (algebraic, integrable, mixed) twistor \mathcal{D} -modules. Notice that the paper [Mochizuki 2015b] also studies twistor structures on GKZ-systems, by considering twistor \mathcal{D} -modules associated to meromorphic functions. We use instead a central result of [Reichelt and Sevenheck 2015], where the Hodge filtration on certain GKZ-systems has been computed explicitly. Technically, the main point in our proof consists in showing that for an \mathcal{R} -module underlying an integrable mixed twistor \mathcal{D} -module on the affine space, the algebraic Fourier–Laplace transformation (which is defined very much the same as in the case of algebraic \mathcal{D} -modules) coincides with the Fourier–Laplace transformation that can be defined inside the category MTM, or even IrrMHM. Along the way, we also obtain (see Theorem 4.7) that an \mathcal{R} -module version of the GKZ- \mathcal{D} -module underlies an irregular Hodge module provided that the parameter $\beta \in \mathbb{C}^d$ of this system satisfies a natural combinatorial condition. Notice that for the special case $\beta = 0$, this theorem can also be deduced from [Mochizuki 2015b, Proposition 1.4].

Our results give concrete representations for objects in the categories MTM and IrrMHM, which usually are difficult to describe explicitly. We hope that a similar approach can be used to understand the irregular Hodge filtration for some higher dimensional analogues of the classical hypergeometric systems, also called Horn systems, which occur in the mirror symmetry picture for toric varieties.

2. Some results on \mathcal{R} - and mixed twistor \mathcal{D} -modules

Let X be a complex manifold of dimension d . We denote by \mathcal{O}_X the sheaf of holomorphic functions and \mathcal{D}_X the sheaf of differential operators with holomorphic coefficients. Recall that \mathcal{D}_X is generated by the

tangent sheaf Θ_X . We put $\mathcal{X} := \mathbb{A}_z^1 \times X$, where the subscript means that z is the canonical coordinate on \mathbb{A}^1 . Denote by $p_z : \mathcal{X} \rightarrow X$ the projection. We denote by $\mathcal{R}_{\mathcal{X}}$ the sheaf of subalgebras of $\mathcal{D}_{\mathcal{X}}$ generated by $zp_z^* \Theta_X$ over $\mathbb{C}_{\mathcal{X}}$ and by $\mathcal{R}_{\mathcal{X}}^{\text{int}}$ the sheaf of subalgebras of $\mathcal{D}_{\mathcal{X}}$ generated by $zp_z^* \Theta_X$ and $z^2 \partial_z$ over $\mathbb{C}_{\mathcal{X}}$. In local coordinates x_1, \dots, x_d , they are given by $\mathbb{C}_{\mathcal{X}} \langle z \partial_{x_1}, \dots, z \partial_{x_d} \rangle$ and $\mathbb{C}_{\mathcal{X}} \langle z^2 \partial_z, z \partial_{x_1}, \dots, z \partial_{x_d} \rangle$, respectively. We set $\Omega_{\mathcal{X}}^1 := z^{-1} p_z^* \Omega_X^1$ as a subsheaf of $p_z^* \Omega_X^1 \otimes \mathbb{C}_{\mathcal{X}}(*(\{0\} \times X))$, $\Omega_{\mathcal{X}}^p := \bigwedge^p \Omega_{\mathcal{X}}^1$ and $\omega_{\mathcal{X}} := \Omega_{\mathcal{X}}^d$.

Let $f : X \rightarrow Y$ be a morphism of complex manifolds. We consider the transfer \mathcal{R} -modules, given by $\mathcal{R}_{\mathcal{X} \rightarrow \mathcal{Y}} := \mathbb{C}_{\mathcal{X}} \otimes_{f^{-1} \mathbb{C}_{\mathcal{Y}}} f^{-1} \mathcal{R}_{\mathcal{Y}}$ and $\mathcal{R}_{\mathcal{Y} \leftarrow \mathcal{X}} := \omega_{\mathcal{X}} \otimes \mathcal{R}_{\mathcal{X} \rightarrow \mathcal{Y}} \otimes f^{-1} \omega_{\mathcal{Y}}$, being respectively a $(\mathcal{R}_{\mathcal{X}}, f^{-1} \mathcal{R}_{\mathcal{Y}})$ -bimodule and a $(f^{-1} \mathcal{R}_{\mathcal{Y}}, \mathcal{R}_{\mathcal{X}})$ -bimodule. We have the inverse image and direct image functors

$$f^+(\mathcal{N}) := \mathcal{R}_{\mathcal{X} \rightarrow \mathcal{Y}} \overset{L}{\otimes}_{f^{-1} \mathcal{R}_{\mathcal{Y}}} f^{-1} \mathcal{N}, \quad f_+(\mathcal{M}) := \mathbf{R}f_*(\mathcal{R}_{\mathcal{Y} \leftarrow \mathcal{X}} \overset{L}{\otimes}_{\mathcal{R}_{\mathcal{X}}} \mathcal{M}), \quad (1)$$

between the bounded derived categories $D^b(\mathcal{R}_{\mathcal{X}})$ and $D^b(\mathcal{R}_{\mathcal{Y}})$.

If $f : X \times Y \rightarrow Y$ is a projection and $\dim X = d$, then $f_+(\mathcal{M})$ is given by

$$f_+(\mathcal{M}) = \mathbf{R}f_* \text{DR}_{\mathcal{X} \times \mathcal{Y}/\mathcal{Y}}(\mathcal{M})[d],$$

where $\text{DR}_{\mathcal{X} \times \mathcal{Y}/\mathcal{Y}}(\mathcal{M})$ is the relative de Rham complex with differential

$$d(\eta \otimes m) = d\eta \otimes m + \sum_{i=1}^d \left(\frac{dx_i}{z} \wedge \eta \right) \otimes z \partial_{x_i} m,$$

the $(x_i)_{1 \leq i \leq d}$ being local coordinates on X .

Let $\sigma : \mathbb{G}_{m,z} \rightarrow \mathbb{G}_{m,z}$ be the automorphism $z \mapsto -\bar{z}^{-1}$. Set $\mathbf{S} := \{z \in \mathbb{A}_z^1 \mid |z| = 1\}$. If $\lambda \in \mathbf{S}$ then $\sigma(\lambda) = -\lambda$. Let $\mathcal{E}_{\mathbf{S} \times X/\mathbf{S},c}^{(d,d)}(V)$ be the space of C^∞ -sections of $\Omega_{\mathbf{S} \times X/\mathbf{S}}^{d,d}$ over any open subset V of $\mathbf{S} \times X$ with compact support and $C_c^0(\mathbf{S})$ the space of continuous functions on \mathbf{S} with compact support. The space of $C^\infty(\mathbf{S})$ -linear maps $\mathcal{E}_{\mathbf{S} \times X/\mathbf{S},c}^{(n,n)}(V) \rightarrow C_c^0(\mathbf{S})$ is denoted by $\mathfrak{D}\mathbf{b}_{\mathbf{S} \times X/\mathbf{S}}(V)$. This gives rise to the sheaf $\mathfrak{D}\mathbf{b}_{\mathbf{S} \times X/\mathbf{S}}$. The abelian category $\mathcal{R}\text{-Tri}(X)$ consists of triples $\mathcal{T} = (\mathcal{M}_1, \mathcal{M}_2, C)$ where $\mathcal{M}_1, \mathcal{M}_2$ are $\mathcal{R}_{\mathcal{X}}$ -modules and $C : \mathcal{M}_1|_{\mathbf{S} \times X} \otimes \sigma^* \mathcal{M}_2|_{\mathbf{S} \times X} \rightarrow \mathfrak{D}\mathbf{b}_{\mathbf{S} \times X/\mathbf{S}}$ is a $\mathcal{R}_{\mathcal{X}}|_{\mathbf{S} \times X} \otimes \sigma^* \mathcal{R}_{\mathcal{X}}|_{\mathbf{S} \times X}$ -linear morphism. If $D \subset X$ is a hypersurface, one similarly defines a category $\mathcal{R}\text{-Tri}(X, D)$ using $\mathcal{R}_{\mathcal{X}}(*D) := \mathcal{R}_{\mathcal{X}} \otimes_{\mathbb{C}_{\mathcal{X}}} \mathbb{C}_{\mathcal{X}}(*(\mathbb{A}_z^1 \times D))$ -modules (see [Mochizuki 2015a, §2.1] for details).

Now let $X := X_0 \times \mathbb{A}_t^1$ and let $\Theta_X(\log X_0)$ be the sheaf of vector fields on X which are logarithmic along X_0 . Let $V_0 \mathcal{R}_{\mathcal{X}}$ be the sheaf of subalgebras in $\mathcal{R}_{\mathcal{X}}$ which is generated by $zp_z^* \Theta_X(\log X_0)$. For $z_0 \in \mathbb{A}_z^1$ we denote by $\mathcal{X}^{(z_0)}$ a small neighborhood of $\{z_0\} \times X$. A coherent $\mathcal{R}_{\mathcal{X}}$ -module is called strictly specializable along t at z_0 if $\mathcal{M}|_{\mathcal{X}^{(z_0)}}$ is equipped with an increasing and exhaustive filtration $V_a^{(z_0)}(\mathcal{M}|_{\mathcal{X}^{(z_0)}})_{a \in \mathbb{R}}$ by coherent $(V_0 \mathcal{R}_{\mathcal{X}})|_{\mathcal{X}^{(z_0)}}$ -modules satisfying certain conditions (see [Mochizuki 2015a, §§2.1.2.1, 2.1.2.2]). This filtration is unique if it exists. \mathcal{M} is called strictly specializable along t if it is strictly specializable along t for any z_0 .

Remark 2.1. If \mathcal{M} is itself a coherent $V_0 \mathcal{R}_{\mathcal{X}}$ -module, then \mathcal{M} is automatically specializable along t and the corresponding filtration $V_a(\mathcal{M})$ exists globally and is trivial, i.e., $V_a(\mathcal{M}) = V_b(\mathcal{M})$ for all $a, b \in \mathbb{R}$.

If \mathcal{M} is a coherent $\mathcal{R}_{\mathcal{X}}(*t)$ -module, we define similarly a filtration $V_a^{(z_0)}(\mathcal{M}_{|\mathcal{X}^{(z_0)}})$ and the notion of strict specializability along t (see [Mochizuki 2015a, §3.1.1]). In this case we define the $\mathcal{R}_{\mathcal{X}}$ -submodules $\mathcal{M}[*t]$ and $\mathcal{M}![t]$ of \mathcal{M} , which are locally generated by $V_0^{(z_0)}\mathcal{M}$ and $V_{<0}^{(z_0)}\mathcal{M}$, respectively.

Remark 2.2. If the coherent $\mathcal{R}_{\mathcal{X}}(*t)$ -module \mathcal{M} is itself $V_0\mathcal{R}_{\mathcal{X}}$ coherent, then

$$\mathcal{M}![t] = \mathcal{M}[*t] = \mathcal{M}(*t) = \mathcal{M}.$$

Given an $\mathcal{R}_{\mathcal{X}}(*t)$ -triple $\mathcal{T} = (\mathcal{M}_1, \mathcal{M}_2, C)$ which is strictly specializable along t we can define

$$\mathcal{T}![t] := (\mathcal{M}_1[*t], \mathcal{M}_2![t], C![t]), \quad \mathcal{T}[*t] := (\mathcal{M}_1![t], \mathcal{M}_2[*t], C[*t])$$

(see [Mochizuki 2015a, Proposition 3.2.1] for details).

The category of filtered $\mathcal{R}_{\mathcal{X}}$ -triples (i.e., $\mathcal{R}_{\mathcal{X}}$ -triples equipped with a finite increasing filtration W) underlies the category $\text{MTM}(X)$ of mixed twistor \mathcal{D} -modules (see [Mochizuki 2015a, Definition 7.2.1]). The full subcategory of objects $\mathcal{T} \in \text{MTM}(X)$ satisfying $\mathcal{T} = \mathcal{T}[*D]$ for some hypersurface $D \subset X$ is denoted by $\text{MTM}(X, [*D])$.

If X is a smooth, algebraic variety, we denote by X^{an} the corresponding complex manifold. Let \bar{X} be a smooth, complete, algebraic variety such that $X \hookrightarrow \bar{X}$ is an open immersion and $D := \bar{X} \setminus X$ is a hypersurface. We can define the category of (integrable) algebraic, mixed twistor \mathcal{D} -modules as

$$\text{MTM}_{\text{alg}}^{(\text{int})}(X) := \text{MTM}^{(\text{int})}(\bar{X}^{\text{an}}, [*D]). \quad (2)$$

We remark that this definition is independent of the completion up to an equivalence of categories [Mochizuki 2015a, Lemma 14.1.3].

Let $f : X \rightarrow Y$ be a quasiprojective morphism of smooth, algebraic varieties. We take completions $X \subset \bar{X}$ and $Y \subset \bar{Y}$ as above, such that $D_X := \bar{X} \setminus X$ and $D_Y := \bar{Y} \setminus Y$ and we have a projective morphism $\bar{f} : \bar{X} \rightarrow \bar{Y}$ which restricts to f . For $\mathcal{T} \in \text{MTM}_{\text{alg}}(X)$, corresponding to $\bar{\mathcal{T}} \in \text{MTM}(\bar{X}, [*D_X])$, we define

$$f_*^i \mathcal{T} := \mathcal{H}^i \bar{f}_* \bar{\mathcal{T}},$$

where \bar{f}_* is the direct image functor for mixed twistor \mathcal{D} -modules arising from the one for \mathcal{R} -modules depicted in (1).

If X is an algebraic variety, we denote by \mathcal{D}_X the sheaf of algebraic differential operators and by $\mathcal{R}_{\mathcal{X}}$ the sheaf of z -differential operators, where here $\mathcal{X} := \mathbb{A}_{\mathbb{C}}^1 \times X$. We define the inverse and direct image functor in the category of algebraic $\mathcal{R}_{\mathcal{X}}$ -modules as in (1). Analogously to the construction of $\mathcal{R}_{\mathcal{X}}$, we can consider the projection $p : \mathbb{P}^1 \times X \rightarrow X$, and construct the sheaf of subalgebras of $\mathcal{D}_{\mathbb{P}^1 \times X}(*(\{\infty\} \times X))$ generated by $z^2 \partial_z$ and $zp^* \Theta_X$ over $\mathcal{O}_{\mathbb{P}^1 \times X}$ (see [Mochizuki 2015a, §14.4.1.1]), which will be denoted by $\mathcal{R}_{\mathbb{P}^1 \times X}^{\text{int}}(*\infty)$. In that sense, an algebraic integrable $\mathcal{R}_{\mathcal{X}}$ -module gives rise to a unique $\mathcal{R}_{\mathbb{P}^1 \times X}^{\text{int}}(*\infty)$ -module (see [ibid., Theorem 14.4.8]).

The following lemma, which will be needed later, is due to T. Mochizuki.

Lemma 2.3. *Given two good $\mathcal{R}_{\mathbb{P}^1 \times X}^{\text{int}}(*\infty)$ -modules $\mathcal{P}_1, \mathcal{P}_2$ and an analytic isomorphism $f : \mathcal{P}_1^{\text{an}} \rightarrow \mathcal{P}_2^{\text{an}}$, then f is induced by a unique algebraic isomorphism between \mathcal{P}_1 and \mathcal{P}_2 .*

Proof. Take a coherent $\mathcal{O}_{\mathbb{P}^1 \times X}$ -submodule $\mathcal{N}_1 \subset \mathcal{P}_1$ such that $\mathcal{R}_{\mathbb{P}^1 \times X}^{\text{int}}(*\infty) \otimes \mathcal{N}_1 \rightarrow \mathcal{P}_1$ is surjective and a coherent $\mathcal{O}_{\mathbb{P}^1 \times X}$ -module $\mathcal{N}_2 \subset \mathcal{P}_2$ such that both $\mathcal{R}_{\mathbb{P}^1 \times X}^{\text{int}}(*\infty) \otimes \mathcal{N}_2 \rightarrow \mathcal{P}_2$ is surjective and $f(\mathcal{N}_1^{\text{an}}) \subset \mathcal{N}_2^{\text{an}}$. According to GAGA we have a morphism $g : \mathcal{N}_1 \rightarrow \mathcal{N}_2$ which after analytification is equal to the morphism $\mathcal{N}_1^{\text{an}} \rightarrow \mathcal{N}_2^{\text{an}}$ induced by f . Denote by \mathcal{K}_1 the kernel of $\mathcal{R}_{\mathbb{P}^1 \times X}^{\text{int}}(*\infty) \otimes \mathcal{N}_1 \rightarrow \mathcal{P}_1$. This gives a morphism $\mathcal{K}_1 \rightarrow \mathcal{P}_2$ which one obtains as the composition $\mathcal{K}_1 \rightarrow \mathcal{R}_{\mathbb{P}^1 \times X}^{\text{int}}(*\infty) \otimes \mathcal{N}_1 \xrightarrow{\varphi} \mathcal{R}_{\mathbb{P}^1 \times X}^{\text{int}}(*\infty) \otimes \mathcal{N}_2 \rightarrow \mathcal{P}_2$, where φ is induced by g . Because the induced morphism $(\mathcal{R}_{\mathbb{P}^1 \times X}^{\text{int}}(*\infty) \otimes \mathcal{N}_1)^{\text{an}} \rightarrow \mathcal{P}_2^{\text{an}}$ factors through $\mathcal{P}_1^{\text{an}}$, the induced morphism $\mathcal{K}_1^{\text{an}} \rightarrow \mathcal{P}_2^{\text{an}}$ is 0. Hence, we obtain that $\mathcal{K}_1 \rightarrow \mathcal{P}_2$ is 0, which means that $\mathcal{R}_{\mathbb{P}^1 \times X}^{\text{int}}(*\infty) \otimes \mathcal{N}_1 \rightarrow \mathcal{P}_2$ factors through \mathcal{P}_1 . This shows the existence. The uniqueness follows from [Serre 1955–1956, Proposition 10]. \square

Since an algebraic, integrable, mixed twistor \mathcal{D} -module on X gives rise to an analytic $\mathcal{R}_{\mathbb{P}^1 \times X}^{\text{int}}(*\infty)$ -module which underlies an algebraic $\mathcal{R}_{\mathbb{P}^1 \times X}^{\text{int}}(*\infty)$ -module by [Mochizuki 2015a, Theorem 14.4.8], the lemma above shows that we can define functors (up to canonical isomorphism)

$$\begin{aligned} \text{For}_i : \text{MTM}_{\text{alg}}^{\text{int}}(X) &\rightarrow \text{Mod}(\mathcal{R}_{\mathcal{X}}^{\text{int}}) \\ (\mathcal{M}_1, \mathcal{M}_2, C) &\mapsto \mathcal{M}_i \quad \text{for } i = 1, 2, \end{aligned}$$

which become faithful if we impose goodness.

3. Fourier transformation of twistor \mathcal{D} -modules

In this section we define the Fourier–Laplace transformation in the categories of integrable \mathcal{R} -modules and integrable, algebraic, mixed twistor \mathcal{D} -modules, and we prove that these two transformations are compatible.

Consider the diagram

$$\begin{array}{ccc} \mathbb{A}^N \times \hat{\mathbb{A}}^N & \xrightarrow{j} & \mathbb{P}^N \times \hat{\mathbb{P}}^N, \\ p \swarrow & & \downarrow \bar{q} \\ \mathbb{A}^N & \xrightarrow{q} \hat{\mathbb{A}}^N & \xrightarrow{\hat{j}} \hat{\mathbb{P}}^N \end{array}$$

where p and q are the projections to the first and second factor respectively. Consider the function $\varphi = \sum_{i=1}^N w_i \cdot \lambda_i$ on $\mathbb{A}^N \times \hat{\mathbb{A}}^N$.

Let $\mathcal{A}_{\text{aff}}^{\varphi/z}$ be the $\mathcal{R}_{\mathbb{A}^1 \times \mathbb{A}^N \times \hat{\mathbb{A}}^N}$ -module $\mathcal{O}_{\mathbb{A}^1 \times \mathbb{A}^N \times \hat{\mathbb{A}}^N}$ equipped with the z -connection $zd + d\varphi$, and consider the reduced divisor $D := (\mathbb{P}^N \times \hat{\mathbb{P}}^N) \setminus (\mathbb{A}^N \times \hat{\mathbb{A}}^N)$. Then $\mathcal{A}_*^{\varphi/z} := j_* \mathcal{A}_{\text{aff}}^{\varphi/z}$ carries a natural structure of an $\mathcal{R}_{\mathbb{A}^1 \times \mathbb{P}^N \times \hat{\mathbb{P}}^N}(*D)$ -module.

We denote by $\mathcal{E}_*^{\varphi/z}$ the analytification of $\mathcal{A}_*^{\varphi/z}$, which is an $\mathcal{R}_{\mathbb{A}^1 \times \mathbb{P}^N \times \hat{\mathbb{P}}^N}(*D)$ -module.

Lemma 3.1. $\mathcal{E}_*^{\varphi/z}$ is strictly specializable along D and

$$\mathcal{E}^{\varphi/z} := \mathcal{E}_*^{\varphi/z}[*D] = \mathcal{E}_*^{\varphi/z}.$$

Proof. We denote the coordinates on $\mathbb{P}^N \times \hat{\mathbb{P}}^N$ by $((w_0 : w_1 : \dots : w_N), (\lambda_0 : \lambda_1 : \dots : \lambda_N))$, where the chart $\mathbb{A}^N \times \hat{\mathbb{A}}^N$ is embedded via the map $j : (w_1, \dots, w_N, \lambda_1, \dots, \lambda_N) \mapsto ((1 : w_1, \dots, w_N), (1 : \lambda_1 : \dots : \lambda_N))$. By symmetry it is enough to prove the claim in the charts $\{w_1 \neq 0, \lambda_0 \neq 0\}$, $\{w_1 \neq 0, \lambda_1 \neq 0\}$ and $\{w_1 \neq 0, \lambda_2 \neq 0\}$. We will assume $N \geq 2$ and consider the chart $X := \{w_1 \neq 0, \lambda_2 \neq 0\}$; the arguments with the other charts and when $N = 1$ go similarly. The chart X is embedded as $(x_1, \dots, x_N, \mu_1, \dots, \mu_N) \mapsto ((x_1 : 1 : x_2 : \dots : x_N), (\mu_1 : \mu_2 : 1 : \mu_3 : \dots : \mu_N))$, so that the map φ is given on X by

$$\frac{1}{x_1 \mu_1} \left(\mu_2 + x_2 + \sum_{i \geq 3} \mu_i x_i \right).$$

Set $\mathcal{D}_X := \mathbb{A}^1 \times (D \cap X) = \mathbb{A}^1 \times \{x_1 \cdot \mu_1 = 0\}$. The module $(\mathcal{E}_*^{\varphi/z})|_X$ is a cyclic $\mathcal{R}_{\mathbb{A}^1 \times X}(*\mathcal{D}_X)$ -module $\mathcal{R}_{\mathbb{A}^1 \times X}(*\mathcal{D}_X)/\mathcal{I}$, where the left ideal \mathcal{I} is generated by

$$\begin{aligned} z\partial_{x_1} + \frac{1}{x_1^2 \mu_1} \left(\mu_2 + x_2 + \sum_{i \geq 3} \mu_i x_i \right), \quad z\partial_{x_2} - \frac{1}{x_1 \mu_1}, \quad z\partial_{x_j} - \frac{\mu_j}{x_1 \mu_1}, \\ z\partial_{\mu_1} + \frac{1}{x_1 \mu_1^2} \left(\mu_2 + x_2 + \sum_{i \geq 3} \mu_i x_i \right), \quad z\partial_{\mu_2} - \frac{1}{x_1 \mu_1}, \quad z\partial_{\mu_j} - \frac{x_j}{x_1 \mu_1}, \end{aligned}$$

where $j \geq 3$. Consider the map $i_g : X \rightarrow \mathbb{A}_t^1 \times X$ given by

$$(x_1, \dots, x_N, \mu_1, \dots, \mu_N) \mapsto (x_1 \cdot \mu_1, x_1, \dots, x_N, \mu_1, \dots, \mu_N).$$

The direct image $i_{g,+}(\mathcal{R}_{\mathcal{X}}(*\mathcal{D}_X)/\mathcal{I})$ is a cyclic $\mathcal{R}_{\mathbb{A}_t^1 \times \mathcal{X}}(*(\mathbb{A}_t^1 \times \mathcal{D}_X))$ -module $\mathcal{R}_{\mathbb{A}_t^1 \times \mathcal{X}}(*(\mathbb{A}_t^1 \times \mathcal{D}_X))/\mathcal{I}'$ where \mathcal{I}' is generated by

$$\begin{aligned} z\partial_{x_1} + \mu_1 z\partial_t + \frac{1}{x_1^2 \mu_1} \left(\mu_2 + x_2 + \sum_{i \geq 3} \mu_i x_i \right), \quad z\partial_{x_2} - \frac{1}{x_1 \mu_1}, \quad z\partial_{x_j} - \frac{\mu_j}{x_1 \mu_1}, \\ z\partial_{\mu_1} + x_1 z\partial_t + \frac{1}{x_1 \mu_1^2} \left(\mu_2 + x_2 + \sum_{i \geq 3} \mu_i x_i \right), \quad z\partial_{\mu_2} - \frac{1}{x_1 \mu_1}, \quad z\partial_{\mu_j} - \frac{x_j}{x_1 \mu_1}, \quad t - x_1 \mu_1, \end{aligned}$$

where $j \geq 3$. Define the cyclic $\mathcal{R}_{\mathbb{A}_t^1 \times \mathcal{X}}(*t)$ -module $\mathcal{R}_{\mathbb{A}_t^1 \times \mathcal{X}}(*t)/\mathcal{I}$ where \mathcal{I} is generated by

$$\begin{aligned} z\partial_{x_1} + \mu_1 z\partial_t + \frac{\mu_1}{t^2} \left(\mu_2 + x_2 + \sum_{i \geq 3} \mu_i x_i \right), \quad z\partial_{x_2} - \frac{1}{t}, \quad z\partial_{x_j} - \frac{\mu_j}{t}, \\ z\partial_{\mu_1} + x_1 z\partial_t + \frac{x_1}{t^2} \left(\mu_2 + x_2 + \sum_{i \geq 3} \mu_i x_i \right), \quad z\partial_{\mu_2} - \frac{1}{t}, \quad z\partial_{\mu_j} - \frac{x_j}{t}, \quad t - x_1 \mu_1, \end{aligned}$$

where $j \geq 3$. Then we have the following $\mathcal{R}_{\mathbb{A}_t^1 \times \mathcal{X}}$ -linear isomorphism

$$\begin{aligned} \mathcal{R}_{\mathbb{A}_t^1 \times \mathcal{X}}(*(\mathbb{A}_t^1 \times \mathcal{D}_X))/\mathcal{I}' &\rightarrow \mathcal{R}_{\mathbb{A}_t^1 \times \mathcal{X}}(*t)/\mathcal{I} \\ P \cdot \frac{1}{(x_1 \mu_1)^k} &\mapsto P \cdot \frac{1}{t^k}. \end{aligned}$$

Consider the V -filtration along $t = 0$. The relations $1/t^k = (z\partial_{\mu_2})^k$,

$$z\partial_t = -\frac{1}{t} \left(z\partial_{x_1} x_1 + \frac{1}{t} \left(\mu_2 + x_2 + \sum_{i \geq 3} \mu_i x_i \right) \right) = -z\partial_{x_1} x_1 z\partial_{\mu_2} - \left(\mu_2 + x_2 + \sum_{i \geq 3} \mu_i x_i \right) (z\partial_{\mu_2})^2$$

and a straightforward induction over k for $(z\partial_t)^k$ show that $i_{g,+}(\mathcal{R}_{\mathcal{X}}(*\mathcal{D}_X)/\mathcal{F})$ is a cyclic, hence also coherent, $V_0\mathcal{R}_{\mathbb{A}^1_{\mathcal{X}} \times \mathcal{X}}$ -module. It follows from Remark 2.1 that $i_{g,+}(\mathcal{R}_{\mathcal{X}}(*\mathcal{D}_X)/\mathcal{F}) = i_{g,+}(\mathcal{R}_{\mathcal{X}}(*\mathcal{D}_X)/\mathcal{F})[*t]$, and as a consequence, we are done by applying [Mochizuki 2015a, §3.3.1.1] and Remark 2.2. \square

It follows from [Sabbah and Yu 2015, Proposition 3.3] that $\mathcal{E}^{\varphi/z}$ underlies an object $\mathcal{T}^{\varphi/z} \in \text{MTM}_{\text{alg}}^{\text{int}}(\mathbb{A}^N \times \hat{\mathbb{A}}^N)$. Let us notice that the preceding lemma, as well as the similar Lemma 3.6 below, are related to a more general statement in [Mochizuki 2015b, Corollary 3.12] on mixed twistor \mathcal{D} -modules associated to nondegenerate functions. However, in order to keep the paper self-contained, we prefer to give direct proofs here.

We will now define a Fourier–Laplace transformation for algebraic $\mathcal{R}_{\mathbb{A}^1 \times \mathbb{A}^N}^{\text{int}}$ -modules.

Definition 3.2. The Fourier–Laplace transformation functor from the category of algebraic $\mathcal{R}_{\mathbb{A}^1 \times \mathbb{A}^N}^{\text{int}}$ -modules to the category of algebraic $\mathcal{R}_{\mathbb{A}^1 \times \hat{\mathbb{A}}^N}^{\text{int}}$ -modules is defined as

$$\widehat{\mathcal{M}} := \text{FL}(\mathcal{M}) := \mathcal{H}^0 q_+((p^+ \mathcal{M}) \otimes \mathcal{A}_{\text{aff}}^{\varphi/z}),$$

for any \mathcal{M} in $\text{Mod}(\mathcal{R}_{\mathbb{A}^1 \times \mathbb{A}^N}^{\text{int}})$.

Remark 3.3. Let $M := \Gamma(\mathbb{A}^1 \times \mathbb{A}^N, \mathcal{M})$ be the $R_{\mathbb{A}^1 \times \mathbb{A}^N}^{\text{int}}$ -module of global sections of \mathcal{M} . The $R_{\mathbb{A}^1 \times \hat{\mathbb{A}}^N}^{\text{int}}$ -module $\widehat{M} := \Gamma(\mathbb{A}^1 \times \hat{\mathbb{A}}^N, \widehat{\mathcal{M}})$ is isomorphic to M as a $\mathbb{C}[z]$ -module and the full $R_{\mathbb{A}^1 \times \hat{\mathbb{A}}^N}^{\text{int}}$ -structure is given by

$$\lambda_i \cdot m := -z\partial_{w_i} \cdot m, \quad z\partial_{\lambda_i} \cdot m := w_i \cdot m \quad \text{and} \quad z^2\partial_z \cdot m := \left(z^2\partial_z - \sum_{i=1}^N z\partial_{w_i} w_i \right) \cdot m.$$

On the other hand, there is a similar definition of a Fourier–Laplace transformation in the category of algebraic $\mathcal{D}_{\mathbb{A}^N}$ -modules (see e.g., [Reichelt 2014, Definition 1.2]) which we also denote by FL.

The Fourier–Laplace transformation for algebraic, integrable, mixed twistor \mathcal{D} -modules is defined in the following way.

Definition 3.4. The Fourier–Laplace transformation in the category of algebraic, integrable mixed twistor \mathcal{D} -modules on \mathbb{A}^N is defined by

$$\text{FL}_{\text{MTM}}(\mathcal{M}) := \mathcal{H}^0 q_*((p^* \mathcal{M}) \otimes \mathcal{T}^{\varphi/z}),$$

where $\mathcal{M} \in \text{MTM}_{\text{alg}}^{\text{int}}(\mathbb{A}^N)$.

Recall that for $\mathcal{M} = (\mathcal{M}_1, \mathcal{M}_2, C) \in \text{MTM}_{\text{alg}}^{\text{int}}(X)$ we denote by For_i the forgetful functors $\text{For}_i(\mathcal{M}) = \mathcal{M}_i$ for $i = 1, 2$.

Proposition 3.5. *Let $\mathcal{M} \in \text{MTM}_{\text{alg}}^{\text{int}}(\mathbb{A}^N)$. Then*

$$\text{For}_1(\text{FL}_{\text{MTM}}(\mathcal{M})) = \text{FL}(\text{For}_1(\mathcal{M})) \quad \text{and} \quad \text{For}_2(\text{FL}_{\text{MTM}}(\mathcal{M})) = z^{-N} \text{FL}(\text{For}_2(\mathcal{M})).$$

Proof. By [Mochizuki 2015a, §14.3.3.3] it is clear that For_i almost commutes with p^* , more precisely we have

$$\text{For}_1(p^*(\mathcal{M})) = z^N p^+(\text{For}_1(\mathcal{M})) \quad \text{and} \quad \text{For}_2(p^*(\mathcal{M})) = p^+(\text{For}_2(\mathcal{M})).$$

Then it is enough to prove for $\mathcal{N} \in \text{MTM}_{\text{alg}}^{\text{int}}(\mathbb{A}^N \times \hat{\mathbb{A}}^N)$ that $q_+(\text{For}_i(\mathcal{N}) \otimes \mathcal{A}_{\text{aff}}^{\varphi/z}) \cong \text{For}_i(q_*(\mathcal{N} \otimes \mathcal{T}^{\varphi/z}))$. We have

$$\begin{aligned} \hat{j}_+ q_+(\text{For}_i(\mathcal{N}) \otimes \mathcal{A}_{\text{aff}}^{\varphi/z}) &\cong \bar{q}_+ j_+(\text{For}_i(\mathcal{N}) \otimes \mathcal{A}_{\text{aff}}^{\varphi/z}) \\ &\cong \bar{q}_+ j_*(\text{For}_i(\mathcal{N}) \otimes \mathcal{A}_{\text{aff}}^{\varphi/z}) \\ &\cong R\bar{q}_* \text{DR}_{\mathbb{P}^N \times \hat{\mathbb{P}}^N} j_*(\text{For}_i(\mathcal{N}) \otimes \mathcal{A}_{\text{aff}}^{\varphi/z}). \end{aligned}$$

Since $\mathcal{N}, \mathcal{T}^{\varphi/z} \in \text{MTM}_{\text{alg}}^{\text{int}}(\mathbb{A}^N \times \hat{\mathbb{A}}^N)$, there exist mixed twistor \mathcal{D} -modules $\bar{\mathcal{N}}, \bar{\mathcal{T}}^{\varphi/z} \in \text{MTM}^{\text{int}}(\mathbb{P}^N \times \hat{\mathbb{P}}^N, [*D])$ whose underlying \mathcal{R} -modules are (after stupid localization along D) analytifications of the $j_* \text{For}_i(\mathcal{N})$ and $j_* \mathcal{A}_{\text{aff}}^{\varphi/z}$. Hence

$$(j_*(\text{For}_i(\mathcal{N}) \otimes \mathcal{A}_{\text{aff}}^{\varphi/z}))^{\text{an}} \cong \text{For}_i(\bar{\mathcal{N}} \otimes \bar{\mathcal{T}}^{\varphi/z})(*D) \cong \text{For}_i(\bar{\mathcal{N}} \otimes \bar{\mathcal{T}}^{\varphi/z}),$$

where the last equation follows from Lemma 3.1. We therefore get

$$\begin{aligned} (\hat{j}_+ p_+(\text{For}_i(\mathcal{N}) \otimes \mathcal{A}_{\text{aff}}^{\varphi/z}))^{\text{an}} &\cong R\bar{q}_* \text{DR}_{\mathbb{P}^N \times \hat{\mathbb{P}}^N}^{\text{an}} (j_*(\text{For}_i(\mathcal{N}) \otimes \mathcal{A}_{\text{aff}}^{\varphi/z}))^{\text{an}} \\ &\cong R\bar{q}_* \text{DR}_{\mathbb{P}^N \times \hat{\mathbb{P}}^N}^{\text{an}} \text{For}_i(\bar{\mathcal{N}} \otimes \bar{\mathcal{T}}^{\varphi/z}) \\ &\cong \text{For}_i(\bar{q}_*(\bar{\mathcal{N}} \otimes \bar{\mathcal{T}}^{\varphi/z})). \end{aligned}$$

The claim follows now from Lemma 2.3, noting that the goodness is a consequence of Lemma 3.1 and [Mochizuki 2015a, Lemma 14.4.15]. \square

We have the following variant, which will be used in the next section. Consider the diagram

$$\begin{array}{ccccc} \mathbb{A}^N \times \mathbb{G}_m & \xrightarrow{j} & \mathbb{P}^N \times \mathbb{P}^1 \\ p \swarrow & & \downarrow \bar{q} \\ \mathbb{A}^N & \xrightarrow{q} & \mathbb{G}_m \xrightarrow{\hat{j}} \mathbb{P}^1 \end{array}$$

and let $\psi := w_1 \cdot t + w_2 + \cdots + w_N$.

As above we define the $\mathcal{R}_{\mathbb{A}^1 \times \mathbb{A}^N \times \mathbb{G}_m}$ -module $\mathcal{A}_{\text{aff}}^{\psi/z}$, being $\mathcal{O}_{\mathbb{A}^1 \times \mathbb{A}^N \times \mathbb{G}_m}$ endowed with the z -connection $z d + d\psi$. As in the other case, we can consider the divisor $H := (\mathbb{P}^N \times \mathbb{P}^1) \setminus (\mathbb{A}^N \times \mathbb{G}_m)$ and obtain the $\mathcal{R}_{\mathbb{A}^1 \times \mathbb{P}^N \times \mathbb{P}^1}(*H)$ -module $\mathcal{A}_*^{\psi/z} := j_* \mathcal{A}_{\text{aff}}^{\psi/z}$. In the same vein as before, we will denote by $\mathcal{E}_*^{\psi/z}$ the $\mathcal{R}_{\mathbb{A}^1 \times \mathbb{P}^N \times \mathbb{P}^1}(*H)$ -module being the analytification of $\mathcal{A}_*^{\psi/z}$. The following lemma is similar to Lemma 3.1.

Lemma 3.6. $\mathcal{E}_*^{\psi/z}$ is strictly specializable along H and

$$\mathcal{E}^{\psi/z} := \mathcal{E}_*^{\psi/z}[*H] = \mathcal{E}_*^{\psi/z}.$$

Proof. We denote the coordinates on $\mathbb{P}^N \times \mathbb{P}^1$ by $((w_0 : w_1 : \dots : w_N), (u : t))$, where the chart $\mathbb{A}^N \times \mathbb{G}_m$ is embedded via the map $j : (w_1, \dots, w_N, t) \mapsto ((1 : w_1 : \dots : w_N), (1 : t))$. We will assume $N \geq 3$ and consider the chart $X := \{w_2 \neq 0, u \neq 0\}$; the other charts behave similarly, as do the cases $N = 1, 2$. The chart X is embedded as $(x_1, \dots, x_N, u) \mapsto ((x_1 : x_2 : 1 : x_3 : \dots : x_N), (u : 1))$. On this chart the map ψ is given by $\frac{1}{x_1} \left(\frac{x_2}{u} + 1 + x_3 + \dots + x_N \right)$. Set $\mathcal{H}_X := \mathbb{A}^1 \times (H \cap X) = \mathbb{A}^1 \times \{x_1 \cdot u = 0\}$. The module $(\mathcal{E}_*^{\psi/z})|_X$ is a cyclic $\mathcal{R}_{\mathcal{X}}(*\mathcal{H}_X)$ -module $\mathcal{R}_{\mathcal{X}}(*\mathcal{H}_X)/\mathcal{I}$, where the left ideal \mathcal{I} is generated by

$$z\partial_{x_1} + \frac{1}{x_1^2} \left(\frac{x_2}{u} + 1 + x_3 + \dots + x_N \right), \quad z\partial_{x_2} - \frac{1}{x_1 u}, \quad z\partial_{x_j} - \frac{1}{x_1}, \quad z\partial_u + \frac{x_2}{x_1 u^2},$$

with $j \geq 3$. Consider the map $i_g : X \rightarrow \mathbb{A}_s^1 \times X$ given by

$$(x_1, \dots, x_N, u) \mapsto (x_1 \cdot u, x_1, \dots, x_N, u).$$

Analogous to Lemma 3.1, the direct image $i_{g,+}(\mathcal{R}_{\mathcal{X}}(*\mathcal{H}_X)/\mathcal{I})$ is a cyclic $\mathcal{R}_{\mathbb{A}_s^1 \times \mathcal{X}}(*(\mathbb{A}_s^1 \times \mathcal{H}_X))$ -module $\mathcal{R}_{\mathbb{A}_s^1 \times \mathcal{X}}(*(\mathbb{A}_s^1 \times \mathcal{H}_X))/\mathcal{I}'$ where \mathcal{I}' is the left ideal generated by

$$z\partial_{x_1} + uz\partial_s + \frac{1}{x_1^2} \left(\frac{x_2}{u} + 1 + x_3 + \dots + x_N \right), \quad z\partial_{x_2} - \frac{1}{x_1 u}, \quad z\partial_{x_j} - \frac{1}{x_1}, \quad z\partial_u + x_1 z\partial_s + \frac{x_2}{x_1 u^2}, \quad s - x_1 u,$$

and $j \geq 3$. Define the cyclic $\mathcal{R}_{\mathbb{A}_s^1 \times \mathcal{X}}(*s)$ -module $\mathcal{R}_{\mathbb{A}_s^1 \times \mathcal{X}}(*s)/\mathcal{I}$ where \mathcal{I} is generated by

$$z\partial_{x_1} + uz\partial_s + \frac{1}{s^2} (x_2 u + u^2 + x_3 u^2 + \dots + x_N u^2), \quad z\partial_{x_2} - \frac{1}{s}, \quad z\partial_{x_j} - \frac{u}{s}, \quad z\partial_u + x_1 z\partial_s + \frac{x_1 x_2}{s^2}, \quad s - x_1 u,$$

and where $j \geq 3$. We have the following $\mathcal{R}_{\mathbb{A}_s^1 \times \mathcal{X}}$ -linear isomorphism

$$\begin{aligned} \mathcal{R}_{\mathbb{A}_s^1 \times \mathcal{X}}(*(\mathbb{A}_s^1 \times \mathcal{H}_X))/\mathcal{I}' &\rightarrow \mathcal{R}_{\mathbb{A}_s^1 \times \mathcal{X}}(*s)/\mathcal{I} \\ P \frac{1}{(x_1 u)^k} &\mapsto P \frac{1}{s^k}. \end{aligned}$$

Consider the V -filtration along $s = 0$. The relations $1/s^k = (z\partial_{x_2})^k$,

$$z\partial_s = -\frac{1}{s} \left(z + uz\partial_u + \frac{x_2}{s} \right) = -z \cdot z\partial_{x_2} - uz\partial_u z\partial_{x_2} - x_2 (z\partial_{x_2})^2$$

and a straightforward induction over k for $(z\partial_s)^k$ show that $i_{g,+}(\mathcal{R}_{\mathcal{X}}(*D_X)/\mathcal{I})$ is a coherent $V_0 \mathcal{R}_{\mathbb{A}_s^1 \times \mathcal{X}}$ -module. As in the previous lemma, this shows the claim. \square

It follows again from [Sabbah and Yu 2015, Proposition 3.3] that $\mathcal{E}^{\psi/z}$ underlies an object $\mathcal{T}^{\psi/z} \in \text{MTM}_{\text{alg}}^{\text{int}}(\mathbb{A}^n \times \mathbb{G}_m)$.

Definition 3.7. (1) The Fourier–Laplace transformation with respect to the kernel ψ in the category of algebraic $\mathcal{R}_{\mathbb{A}^1 \times \mathbb{A}^N}$ -modules is defined as

$$\text{FL}^{\psi}(\mathcal{M}) := \mathcal{H}^0 q_+((p^+ \mathcal{M}) \otimes \mathcal{A}_{\text{aff}}^{\psi/z}),$$

for any $\mathcal{M} \in \text{Mod}(\mathcal{R}_{\mathbb{A}^N})$.

- (2) Analogously, the Fourier–Laplace transformation with respect to the kernel ψ in the category of algebraic, integrable twistor \mathcal{D} -modules on \mathbb{A}^N is defined by

$$\mathrm{FL}_{\mathrm{MTM}}^{\psi}(\mathcal{M}) := \mathcal{H}^0 q_*((p^* \mathcal{M}) \otimes \mathcal{T}^{\psi/z}),$$

for any $\mathcal{M} \in \mathrm{MTM}_{\mathrm{alg}}^{\mathrm{int}}(\mathbb{A}^N)$.

We get the following result for the kernel ψ .

Proposition 3.8. *Let $\mathcal{M} \in \mathrm{MTM}_{\mathrm{alg}}^{\mathrm{int}}(\mathbb{A}^N)$. Then*

$$\mathrm{For}_1(\mathrm{FL}_{\mathrm{MTM}}^{\psi}(\mathcal{M})) = z^{1-N} \mathrm{FL}^{\psi}(\mathrm{For}_1(\mathcal{M})) \quad \text{and} \quad \mathrm{For}_2(\mathrm{FL}_{\mathrm{MTM}}^{\psi}(\mathcal{M})) = z^{-N} \mathrm{FL}^{\psi}(\mathrm{For}_2(\mathcal{M})).$$

Proof. We have, by [Mochizuki 2015a, §14.3.3.3],

$$\mathrm{For}_1(p^*(\mathcal{M})) = zp^+(\mathrm{For}_1(\mathcal{M})) \quad \text{and} \quad \mathrm{For}_2(p^*(\mathcal{M})) = p^+(\mathrm{For}_2(\mathcal{M})).$$

The rest of the proof carries over almost word for word from Proposition 3.5, using Lemma 3.6. \square

4. GKZ systems and irregular Hodge modules

Let $A = (a_{ki})$ be a $d \times N$ integer matrix with columns $(\underline{a}_1, \dots, \underline{a}_N)$. We define

$$\mathbb{N}A := \sum_{i=1}^N \mathbb{N}\underline{a}_i \subset \mathbb{Z}^d$$

and similarly for $\mathbb{Z}A$ and $\mathbb{R}_{\geq 0}A$. Throughout this section we assume

$$\mathbb{Z}A = \mathbb{Z}^d \quad \text{and} \quad \mathbb{N}A = \mathbb{Z}^d \cap \mathbb{R}_{\geq 0}A.$$

Set $\mathbb{A}^N := \mathrm{Spec}(\mathbb{C}[w_1, \dots, w_N])$ and $\hat{\mathbb{A}}^N := \mathrm{Spec}(\mathbb{C}[\lambda_1, \dots, \lambda_N])$ and

$$\mathbb{L}_A := \left\{ \ell = (\ell_1, \dots, \ell_N) \in \mathbb{Z}^N : \sum_{i=1}^N \ell_i \underline{a}_i \right\}.$$

Definition 4.1. The GKZ-hypergeometric system \mathcal{M}_A^{β} is the cyclic $\mathcal{D}_{\hat{\mathbb{A}}^N}$ -module $\mathcal{D}_{\hat{\mathbb{A}}^N}/\mathcal{I}$, where \mathcal{I} is the left ideal generated by

$$E_k := \sum_{i=1}^N a_{ki} \lambda_i \partial_{\lambda_i} - \beta_k, \quad \text{for } k = 1, \dots, d,$$

and

$$\square_{\ell} := \prod_{\ell_i > 0} \partial_{\lambda_i}^{\ell_i} - \prod_{\ell_i < 0} \partial_{\lambda_i}^{-\ell_i}, \quad \text{for } \ell \in \mathbb{L}_A.$$

The GKZ-hypergeometric system \mathcal{M}_A^β is the Fourier–Laplace transform of the cyclic $\mathcal{D}_{\mathbb{A}^N}$ -module $\check{\mathcal{M}}_A^\beta := \mathcal{D}_{\mathbb{A}^N} / \mathcal{I}$, where \mathcal{I} is the left ideal generated by

$$\check{E}_k := \sum_{i=1}^N a_{ki} \partial_{w_i} w_i + \beta_k, \quad \text{for } k = 1, \dots, d,$$

and

$$\check{\square}_\ell := \prod_{\ell_i > 0} w_i^{\ell_i} - \prod_{\ell_i < 0} w_i^{-\ell_i}, \quad \text{for } \ell \in \mathbb{L}_A.$$

The semigroup ring $\mathbb{C}[\mathbb{N}A] \subset \mathbb{C}[t_1^\pm, \dots, t_d^\pm]$ is naturally a $\mathbb{C}[w_1, \dots, w_N]$ -module under the isomorphism

$$\begin{aligned} \mathbb{C}[w_1, \dots, w_N] / ((\check{\square}_\ell)_{\ell \in \mathbb{L}_A}) &\rightarrow \mathbb{C}[\mathbb{N}A] \\ w_i &\mapsto t^{a_i}, \end{aligned}$$

where we are using the multiindex notation $t^{a_i} := \prod_{k=1}^d t_k^{a_{ki}}$. We set $S_A := \mathbb{C}[\mathbb{N}A]$. Notice that the rings $\mathbb{C}[w_1, \dots, w_N]$ and S_A carry a natural \mathbb{Z}^d -grading given by $\deg(w_i) = \underline{a}_i$. This is compatible with the grading on the Weyl algebra $\mathcal{D}_{\mathbb{A}^N} := \Gamma(\mathbb{A}^N, \mathcal{D}_{\mathbb{A}^N})$ given by $\deg(w_i) = \underline{a}_i$ and $\deg(\partial_{w_i}) = -\underline{a}_i$.

Definition 4.2 [Matusevich et al. 2005, Definition 5.2]. Let P be a finitely generated \mathbb{Z}^d -graded $\mathbb{C}[w_1, \dots, w_N]$ -module. An element $\alpha \in \mathbb{Z}^d$ is called a true degree of P if the graded part P_α is nonzero. A vector $\alpha \in \mathbb{C}^d$ is called a quasidegree of P if α lies in the complex Zariski closure $\text{qdeg}(P)$ of the true degrees of P via the natural embedding $\mathbb{Z}^d \hookrightarrow \mathbb{C}^d$.

Consider the set of strongly resonant parameters of A :

$$\text{sRes}(A) := \bigcup_{j=1}^N \text{sRes}_j(A),$$

where

$$\text{sRes}_j(A) := \{\beta \in \mathbb{C}^d \mid \beta \in -(\mathbb{N} + 1)\underline{a}_j + \text{qdeg}(S_A / (t^{a_j}))\}.$$

Consider as well the torus $\mathbb{G}_m^d := \text{Spec}(\mathbb{C}[t_1^\pm, \dots, t_d^\pm])$, together with the torus embedding

$$\begin{aligned} h : \mathbb{G}_m^d &\rightarrow \mathbb{A}^N \\ (t_1, \dots, t_d) &\mapsto (t^{a_1}, \dots, t^{a_N}). \end{aligned}$$

The following proposition is a slight generalization of the results of Schulze and Walther [2009, Theorem 3.6, Corollary 3.8].

Proposition 4.3 [Reichelt and Sevenheck 2015, Proposition 3.11]. *Let A be a $d \times N$ integer matrix satisfying $\mathbb{Z}A = \mathbb{Z}^d$ and $\mathbb{N}A = \mathbb{Z}^d \cap \mathbb{R}_{\geq 0}A$. Assume that $\beta \notin \text{sRes}(A)$. Then*

$$\mathcal{H}^0(h_+ \mathcal{O}_{\mathbb{G}_m^d}^\beta) \cong \check{\mathcal{M}}_A^\beta,$$

where $\mathcal{O}_{\mathbb{G}_m^d}^\beta \cong \mathcal{D}_{\mathbb{G}_m^d} / \mathcal{D}_{\mathbb{G}_m^d} \cdot (\partial_{t_1} t_1 + \beta_1, \dots, \partial_{t_d} t_d + \beta_d)$.

For $\beta \in \mathbb{R}^d$, the \mathcal{D} -module $\mathcal{O}_{\mathbb{G}_m^d}^\beta$ underlies the complex mixed Hodge module ${}^p\mathbb{C}_{\mathbb{G}_m^d}^{H,\beta}$. Hence for $\beta \in \mathbb{R}^d \setminus \text{sRes}(A)$ the \mathcal{D} -module \mathcal{M}_A^β underlies the complex mixed Hodge module $\mathcal{H}^0 h_* {}^p\mathbb{C}_{\mathbb{G}_m^d}^{H,\beta}$. The Hodge filtration on $\check{\mathcal{M}}_A^\beta$ can be explicitly computed, provided that β belongs to a certain set \mathfrak{A}_A of so-called admissible parameters β . We recall its definition from [Reichelt and Sevenheck 2015, Formula (14) right before Lemma 4.4]: let $\underline{c} := \underline{a}_1 + \cdots + \underline{a}_N$ and define for all facets F of $\mathbb{R}_{\geq 0} A$ the uniquely determined primitive, inward-pointing, normal vector \underline{n}_F of F , such that $\langle \underline{n}_F, F \rangle = 0$ and $\langle \underline{n}_F, \mathbb{N}A \rangle \subset \mathbb{Z}_{\geq 0}$. Set $e_F := \langle \underline{n}_F, \underline{c} \rangle \in \mathbb{Z}_{>0}$. The set of admissible parameters of A is then defined by

$$\mathfrak{A}_A := \bigcap_{F \text{ facet}} \left\{ \mathbb{R} \cdot F - \left[0, \frac{1}{e_F} \right) \cdot \underline{c} \right\}.$$

Theorem 4.4 [Reichelt and Sevenheck 2015, Theorem 4.17]. *For $\beta \in \mathfrak{A}_A$ the Hodge filtration on $\check{\mathcal{M}}_A^\beta$ is equal to the order filtration shifted by $N - d$, i.e.,*

$$F_{p+N-d}^H \check{\mathcal{M}}_A^\beta = F_p^{\text{ord}} \check{\mathcal{M}}_A^\beta.$$

Let us define the cyclic $\mathcal{R}_{\mathbb{A}^1 \times \mathbb{A}^N}$ -module $\check{\mathcal{N}}_A^\beta := \mathcal{R}_{\mathbb{A}^1 \times \mathbb{A}^N} / \mathcal{I}_z$, where \mathcal{I}_z is the left ideal generated by

$$\check{E}_k^z = \sum_{i=1}^N a_{ki} z \partial_{w_i} w_i + z \beta_k, \quad \text{for } k = 1, \dots, d,$$

and

$$\check{\square}_\ell = \prod_{\ell_i > 0} w_i^{\ell_i} - \prod_{\ell_i < 0} w_i^{-\ell_i}, \quad \text{for } \ell \in \mathbb{L}_A.$$

We will denote by $\check{M}_A^\beta := \Gamma(\mathbb{A}^N, \check{\mathcal{M}}_A^\beta)$ and $\check{N}_A^\beta := \Gamma(\mathbb{A}^1 \times \mathbb{A}^N, \check{\mathcal{N}}_A^\beta)$ the modules of global sections of $\check{\mathcal{M}}_A^\beta$ and $\check{\mathcal{N}}_A^\beta$, respectively.

We will also consider the Rees module of \check{M}_A^β with respect to the order filtration F_\bullet^{ord} , which is given by $R^{F^{\text{ord}}} \check{M}_A^\beta := \sum_{k \geq 0} z^k F_k^{\text{ord}} \check{M}_A^\beta$. An easy computation shows $R^{F^{\text{ord}}} \check{M}_A^\beta = \check{N}_A^\beta$, hence

$$R^{F^H} \check{M}_A^\beta = z^{N-d} \check{N}_A^\beta. \quad (3)$$

Definition 4.5. The \mathcal{R} -GKZ-hypergeometric system \mathcal{N}_A^β is the cyclic $\mathcal{R}_{\mathbb{A}^1 \times \hat{\mathbb{A}}^N}^{\text{int}}$ -module $\mathcal{R}_{\mathbb{A}^1 \times \hat{\mathbb{A}}^N}^{\text{int}} / \mathcal{I}$, where the left ideal \mathcal{I} is generated by

$$\begin{aligned} E_0^z &:= z^2 \partial_z + \sum_{i=1}^N \lambda_i z \partial_{\lambda_i}, \\ E_k^z &:= \sum_{i=1}^N a_{ki} \lambda_i z \partial_{\lambda_i} - z \beta_k, \quad \text{for } k = 1, \dots, d, \end{aligned}$$

and

$$\square_\ell^z := \prod_{\ell_i > 0} (z \partial_{\lambda_i})^{\ell_i} - \prod_{\ell_i < 0} (z \partial_{\lambda_i})^{-\ell_i}, \quad \text{for } \ell \in \mathbb{L}_A.$$

Remark 4.6. Note that, considering $\check{\mathcal{N}}_A^\beta$ as an $\mathcal{R}_{\mathbb{A}^1 \times \hat{\mathbb{A}}^N}^{\text{int}}$ -module with the trivial action of $z^2 \partial_z$, \mathcal{N}_A^β is its Fourier–Laplace transform as $\mathcal{R}_{\mathbb{A}^1 \times \hat{\mathbb{A}}^N}^{\text{int}}$ -modules, according to Remark 3.3.

Theorem 4.7. *Let A be a $d \times N$ -matrix and $\beta \in \mathfrak{A}_A$ an admissible parameter. The \mathcal{R} -GKZ-hypergeometric system $z^{-d} \mathcal{N}_A^\beta$ underlies an algebraic, integrable, mixed twistor \mathcal{D} -module $\mathcal{T}\mathcal{M}_A^\beta$.*

Proof. By the remark above, we know that $\mathcal{N}_A^\beta = \text{FL}(\check{\mathcal{N}}_A^\beta)$, which in turn, thanks to the choice of β , Theorem 4.4 and formula (3), is equal to $\text{FL}(z^{d-N} \mathcal{R}^{F^H} \check{\mathcal{M}}_A^\beta)$. Since $\mathcal{R}^{F^H} \check{\mathcal{M}}_A^\beta$ is the Rees module of a mixed Hodge module on \mathbb{A}^N , it gives rise to an algebraic, integrable mixed twistor \mathcal{D} -module on \mathbb{A}^N , say $\mathcal{T}\check{\mathcal{M}}_A^\beta$. Then we can apply Proposition 3.5 and get

$$\mathcal{N}_A^\beta = z^{d-N} \text{FL}(\text{For}_2(\mathcal{T}\check{\mathcal{M}}_A^\beta)) = z^d \text{For}_2(\text{FL}_{\text{MTM}}(\mathcal{T}\check{\mathcal{M}}_A^\beta)).$$

The result follows from writing $\mathcal{T}\mathcal{M}_A^\beta := \text{FL}_{\text{MTM}}(\mathcal{T}\check{\mathcal{M}}_A^\beta)$. \square

Corollary 4.8. *The analytification of $\mathcal{T}\mathcal{M}_A^\beta$ gives rise to an irregular mixed Hodge module on \mathbb{A}^N which has a natural extension to an $\mathcal{R}_{\mathbb{A}^1 \times \mathbb{P}^N}^{\text{int}}$ -module underlying an object of $\text{IrrMHM}(\mathbb{P}^N)$.*

Proof. This follows from applying [Sabbah 2018, Corollary 0.5] to the operations performed to get $\mathcal{T}\mathcal{M}_A^\beta$. \square

5. Application to confluent hypergeometric systems

In this section we are going to use the results achieved so far for the special case of the matrix

$$A = \left(\begin{array}{c|c|c} \mathbf{1}_m & \mathbf{0}_{m \times (n-1)} & \text{Id}_m \\ \hline \mathbf{1}_{n-1} & -\text{Id}_{n-1} & \mathbf{0}_{(n-1) \times m} \end{array} \right).$$

For the sake of simplicity, we will write $N = n + m$ in what follows. Before going on, let us introduce the main object of study of this section and state some of its basic properties, extending what we mentioned in the introduction.

Definition 5.1. Let $(n, m) \neq (0, 0)$ be a pair of nonnegative integers, and let $\alpha_1, \dots, \alpha_n$ and β_1, \dots, β_m be elements of \mathbb{C} . The hypergeometric \mathcal{D} -module of type (n, m) associated with the α_i and the β_j is defined as the quotient of $\mathcal{D}_{\mathbb{G}_m}$ by the left ideal generated by the so-called hypergeometric operator

$$\prod_{i=1}^n (t \partial_t - \alpha_i) - t \prod_{j=1}^m (t \partial_t - \beta_j).$$

We will denote it by $\mathcal{H}(\alpha_i; \beta_j)$.

Proposition 5.2. *Let $\mathcal{H} := \mathcal{H}(\alpha_i; \beta_j)$ be a hypergeometric \mathcal{D} -module of type (n, m) , and let η be any complex number. Then we have the following:*

- (1) *If we denote the Kummer \mathcal{D} -module $\mathcal{D}_{\mathbb{G}_m}/(t \partial_t - \eta)$ by \mathcal{K}_η , then $\mathcal{H} \otimes_{\mathcal{O}_{\mathbb{G}_m}} \mathcal{K}_\eta \cong \mathcal{H}(\alpha_i + \eta; \beta_j + \eta)$. In particular, an overall integer shift of the parameters gives us an isomorphic \mathcal{D} -module.*
- (2) *\mathcal{H} is irreducible if and only if for any pair (i, j) of indices, $\alpha_i - \beta_j$ is not an integer.*

- (3) If \mathcal{H} is irreducible, its isomorphism class depends only on the classes modulo \mathbb{Z} of the α_i and the β_j , so we can choose such parameters on a fundamental domain of \mathbb{C}/\mathbb{Z} .

Proof. A simple calculation shows (1). (2) follows from [Katz 1990, Propositions 2.11.9 and 3.2], whereas (3) is part of [ibid., Proposition 3.2]. \square

As we mentioned in the introduction, we can express any one-dimensional hypergeometric \mathcal{D} -module as the inverse image of a GKZ hypergeometric \mathcal{D} -module (see [Castaño Domínguez and Sevenheck 2019, Corollary 2.9]). Notice that there is a similar statement at the level of \mathcal{R} -modules (see [ibid., Lemma 2.12]), yielding a description of the $\mathcal{R}_{\mathbb{A}_z^1 \times \mathbb{G}_{m,t}}^{\text{int}}$ -module $\widehat{\mathcal{H}}$ from Theorem 5.7 below as an inverse image of a GKZ-hypergeometric \mathcal{R} -module (as defined in [ibid., Definition 2.10]).

Proposition 5.3. *Let $\mathcal{H}(\alpha_i; \beta_j)$ be a hypergeometric $\mathcal{D}_{\mathbb{G}_m}$ -module of type (n, m) with $\alpha_1 = 0$, let $A \in M((N-1) \times N, \mathbb{Z})$ as above, and let $\gamma = (\beta_1, \dots, \beta_m, \alpha_2, \dots, \alpha_n)^t$. Let $\iota : \mathbb{G}_m \rightarrow \mathbb{A}^N$ be given by $t \mapsto (t, 1, \dots, 1)$. Then*

$$\mathcal{H}(\alpha_i; \beta_j) \cong \iota^+ \mathcal{M}_A^\gamma.$$

Since the restriction map ι is not smooth we do not know a priori whether taking inverse image by it preserves irregular mixed Hodge modules. In order to show that $\mathcal{H}(\alpha_i; \beta_j)$ can be upgraded to an element of $\text{IrrMHM}(\mathbb{G}_m)$ we use Proposition 3.8, where the reduction procedure is built in by the use of the Fourier kernel $\psi = w_1 \cdot t + w_2 + \dots + w_N$.

Let $A \in M((N-1) \times N, \mathbb{Z})$ as above and $\gamma = (\gamma_1, \dots, \gamma_{N-1})^t \in \mathfrak{A}_A$. The $\mathcal{D}_{\mathbb{A}^N}$ -module $\check{\mathcal{M}}_A^\gamma$ underlies a mixed Hodge module on \mathbb{A}^N , so that the Rees module $\mathcal{R}^{F^H}(\check{\mathcal{M}}_A^\gamma)$ then gives rise to an algebraic, integrable mixed twistor \mathcal{D} -module on \mathbb{A}^N that we denote by $\mathcal{T}\check{\mathcal{M}}_A^\gamma$. Then we have the following concrete description of its Fourier–Laplace transform $\text{FL}_{\text{MTM}}^\psi(\mathcal{T}\check{\mathcal{M}}_A^\gamma) = q_*(p^*(\mathcal{T}\check{\mathcal{M}}_A^\gamma) \otimes \mathcal{T}^{\psi/z})$.

Proposition 5.4. *Let A and γ be as before. Then the $\mathcal{R}_{\mathbb{A}^1 \times \mathbb{G}_m}^{\text{int}}$ -module $\text{For}_2(\text{FL}_{\text{MTM}}^\psi(\mathcal{T}\check{\mathcal{M}}_A^\gamma))$ can be expressed as $\mathcal{R}_{\mathbb{A}^1 \times \mathbb{G}_m}^{\text{int}}/(P, H)$, where*

$$P = z^2 \partial_z + (n-m)t z \partial_t + \varepsilon z \quad \text{and} \quad H = z t \partial_t \prod_{i=1}^{n-1} z(t \partial_t - \gamma_{m+i}) - t \prod_{j=1}^m z(t \partial_t - \gamma_j),$$

with $\varepsilon = \sum_{j=1}^m \gamma_j - \sum_{i=m+1}^{N-1} \gamma_i + N - 1$.

Proof. As said after Theorem 4.4, for any γ inside the domain \mathfrak{A}_A of admissible parameters, the Hodge filtration of $\check{\mathcal{M}}_A^\gamma$ is the order filtration shifted by $N - (N-1) = 1$. Therefore, for such values of γ we can give an explicit expression of the Rees module of the filtered module $(\check{\mathcal{M}}_A^\gamma, F_\bullet^H)$. Namely, we have the isomorphism of $\mathcal{R}_{\mathbb{A}^1 \times \mathbb{A}^N}^{\text{int}}$ -modules

$$\mathcal{R}^{F^H}(\check{\mathcal{M}}_A^\gamma) \cong {}_z\check{\mathcal{N}}_A^\gamma := \mathcal{R}_{\mathbb{A}^1 \times \mathbb{A}^N}^{\text{int}}/(\check{E}_i^z, \check{E}_j^z, \check{\square}, z^2 \partial_z - z),$$

where

$$\begin{aligned}\check{E}_i^z &= z\partial_{w_1}w_1 - z\partial_{w_i}w_i + \gamma_{m+i-1}z, \quad \text{for } i = 2, \dots, n, \\ \check{E}_j^z &= z\partial_{w_1}w_1 + z\partial_{w_{n+j}}w_{n+j} + \gamma_jz, \quad \text{for } j = 1, \dots, m, \\ \check{\square} &= \prod_{i=1}^n w_i - \prod_{j=1}^m w_{n+j}.\end{aligned}$$

First we compute $\mathrm{FL}^\psi(z\check{\mathcal{N}}_A^\gamma)$, which involves performing three operations with $z\check{\mathcal{N}}_A^\gamma$: inverse image by $p : \mathbb{G}_m \times \mathbb{A}^N \rightarrow \mathbb{A}^N$, tensor product with the $\mathcal{R}_{\mathbb{A}^1 \times \mathbb{G}_m \times \mathbb{A}^N}^{\mathrm{int}}$ -module $\mathcal{A}_{\mathrm{aff}}^{\psi/z}$ and direct image by $q : \mathbb{G}_m \times \mathbb{A}^N \rightarrow \mathbb{G}_m$. The first one is pretty easy. Namely

$$p^+z\check{\mathcal{N}}_A^\gamma \cong \mathcal{R}_{\mathbb{A}^1 \times \mathbb{G}_m \times \mathbb{A}^N}^{\mathrm{int}} / (\check{E}_i^z, \check{E}_j^z, \check{\square}, z^2\partial_z - z, z\partial_t).$$

Let us tensor now $p^+z\check{\mathcal{N}}_A^\gamma$ with $\mathcal{A}_{\mathrm{aff}}^{\psi/z}$. This $\mathcal{R}^{\mathrm{int}}$ -module can be presented as $\mathcal{R}_{\mathbb{A}^1 \times \mathbb{G}_m \times \mathbb{A}^N}^{\mathrm{int}} \cdot e^{\psi/z} = \mathcal{R}_{\mathbb{A}^1 \times \mathbb{G}_m \times \mathbb{A}^N}^{\mathrm{int}} / \mathcal{I}^\psi$, where \mathcal{I}^ψ is the left ideal generated by

$$z^2\partial_z + w_1t + w_2 + \dots + w_N, \quad z\partial_t - w_1, \quad z\partial_{w_1} - t, \quad z\partial_{w_i} - 1, i = 2, \dots, N.$$

For $n \in p^+z\check{\mathcal{N}}_A^\gamma$, we will call n^ψ the tensor $n \otimes e^{\psi/z}$. Then we can obtain the formulas

$$\begin{aligned}(z\partial_{w_1}w_1n \otimes e^{\psi/z}) &= z\partial_{w_1}(w_1n \otimes e^{\psi/z}) - t(n \otimes w_1e^{\psi/z}) = (z\partial_{w_1}w_1 - tz\partial_t) \cdot n^\psi, \\ (z\partial_{w_k}w_kn \otimes e^{\psi/z}) &= z\partial_{w_k}(w_kn \otimes e^{\psi/z}) - (n \otimes w_ke^{\psi/z}) = (z\partial_{w_k}w_k - w_k) \cdot n^\psi, \quad \text{for } k = 2, \dots, N, \\ (z^2\partial_zn \otimes e^{\psi/z}) &= z^2\partial_z \cdot n^\psi - (n \otimes (-\psi)e^{\psi/z}) = (z^2\partial_z + w_1t + w_2 + \dots + w_N) \cdot n^\psi, \\ (z\partial_tn \otimes e^{\psi/z}) &= z\partial_t \cdot n^\psi - (n \otimes w_1e^{\psi/z}) = (z\partial_t - w_1) \cdot n^\psi.\end{aligned}$$

Hence $p^+z\check{\mathcal{N}}_A^\gamma \otimes \mathcal{A}_{\mathrm{aff}}^{\psi/z}$ is the cyclic $\mathcal{R}_{\mathbb{A}^1 \times \mathbb{G}_m \times \mathbb{A}^N}^{\mathrm{int}}$ -module $\mathcal{R}_{\mathbb{A}^1 \times \mathbb{G}_m \times \mathbb{A}^N}^{\mathrm{int}} / \mathcal{J}^\psi$, with \mathcal{J}^ψ being the left ideal generated by

$$\begin{aligned}\prod_{i=1}^n w_i - \prod_{j=1}^m w_{n+j}, \quad z^2\partial_z - z + w_1t + w_2 + \dots + w_N, \quad z\partial_t - w_1, \\ z\partial_{w_1}w_1 - tz\partial_t - z\partial_{w_i}w_i + w_i + \gamma_{m+i-1}z, \quad \text{for } i = 2, \dots, n, \\ z\partial_{w_1}w_1 - tz\partial_t + z\partial_{w_{n+j}}w_{n+j} - w_{n+j} + \gamma_jz, \quad \text{for } j = 1, \dots, m.\end{aligned}$$

We now consider the zeroth cohomology $\mathcal{H}^0q_+(p^+z\check{\mathcal{N}}_A^\gamma \otimes \mathcal{A}_{\mathrm{aff}}^{\psi/z})$, which is in turn the N -th cohomology of the de Rham complex $q_*\mathrm{DR}_{\mathbb{A}^1 \times \mathbb{G}_m \times \mathbb{A}^N / \mathbb{A}^1 \times \mathbb{G}_m}(p^+z\check{\mathcal{N}}_A^\gamma \otimes \mathcal{A}_{\mathrm{aff}}^{\psi/z})$. This is given by the cyclic $\mathcal{R}_{\mathbb{A}^1 \times \mathbb{G}_m}^{\mathrm{int}}$ -module $\mathcal{R}_{\mathbb{A}^1 \times \mathbb{G}_m}^{\mathrm{int}} / (P', H')$, where the operators P' and H' are given by

$$P' := z^2\partial_z + (n-m)tz\partial_t + \varepsilon'z, \quad H' := zt\partial_t \prod_{i=1}^{n-1} (zt\partial_t - \gamma_{m+i}z) - (-1)^mt \prod_{j=1}^m (zt\partial_t - \gamma_jz)$$

and $\varepsilon' := \sum_{j=1}^m \gamma_j - \sum_{i=m+1}^{N-1} \gamma_i - 1$. Replacing t by $(-1)^m t$ we obtain that $\mathrm{FL}^\psi(z\check{\mathcal{M}}_A^\gamma) \cong \mathcal{R}_{\mathbb{A}^1 \times \mathbb{G}_m}^{\mathrm{int}}/(P', H)$, with

$$H := zt\partial_t \prod_{i=1}^{n-1} (zt\partial_t - \gamma_{m+i}z) - t \prod_{j=1}^m (zt\partial_t - \gamma_j z).$$

Now it follows from Proposition 3.8 that

$$\mathrm{For}_2(\mathrm{FL}_{\mathrm{MTM}}^\psi(\check{\mathcal{M}}_A^\gamma)) \cong z^{-N} \mathrm{FL}^\psi(z\check{\mathcal{M}}_A^\gamma) \cong \mathcal{R}_{\mathbb{A}^1 \times \mathbb{G}_m}^{\mathrm{int}}/(P, H)$$

with

$$P = z^2\partial_z + (n-m)tz\partial_t + \varepsilon z \quad \text{and} \quad H = zt\partial_t \prod_{i=1}^{n-1} z(t\partial_t - \gamma_{m+i}) - t \prod_{j=1}^m z(t\partial_t - \gamma_j),$$

and $\varepsilon = \sum_{j=1}^m \gamma_j - \sum_{i=m+1}^{N-1} \gamma_i + N - 1$. □

Remark 5.5. As a matter of fact, we do not have to restrict ourselves to the region \mathfrak{A}_A to find our admissible parameters. If we have $\gamma \in \mathfrak{A}_A$ and add to it an integer vector $\underline{k} \in \mathbb{Z}^{N-1}$ with no negative entries, then $\gamma + \underline{k} \notin \mathrm{sRes}(A)$ by definition (see the proof of [Reichelt and Sevenheck 2015, Lemma 4.5]). Therefore, since $\mathcal{O}_{\mathbb{G}_m^d}^\gamma \cong \mathcal{O}_{\mathbb{G}_m^d}^{\gamma+\underline{k}}$ for any integer vector \underline{k} , we have $\check{\mathcal{M}}_A^\gamma \cong \check{\mathcal{M}}_A^{\gamma+\underline{k}}$ by Proposition 4.3 and the statement of the proposition holds true after changing \mathfrak{A}_A by $\mathfrak{A}_A + \mathbb{N}^{N-1}$.

We will also make use of the following result, which calculates the admissible domain \mathfrak{A}_A for the matrix A in our particular context.

Lemma 5.6. *Let $A \in M((N-1) \times N, \mathbb{Z})$ be the matrix defined at the beginning of the section. Consider a point $p = (p_1, \dots, p_m, q_1, \dots, q_{n-1}) \in [0, 1)^{N-1}$. Let us define*

$$p_- := \min((\{p_1, \dots, p_m\} \setminus \{0\}) \cup \{1\}) \quad \text{and} \quad p_+ := \max\{p_1, \dots, p_m\},$$

that is, the minimum of the p_i that do not vanish (taking $p_- = 1$ if all of them are zero) and the maximum of them all.

Then, p belongs to $(\mathfrak{A}_A + \mathbb{N}^{N-1}) \subset \mathbb{R}^{N-1}$ if and only if, for all $i = 1, \dots, n-1$

- $q_i \in [0, p_-)$ if some p_i vanishes, or
- $q_i \in [0, p_-) \cup [p_+, 1)$, otherwise.

Proof. We will first find the expression for the admissible region \mathfrak{A}_A . For this purpose, we must find a set of hyperplanes containing the facets of the cone $C := \mathbb{R}_{\geq 0} A \subset \mathbb{R}^{N-1}$. Denote by $\{\underline{u}_1, \dots, \underline{u}_{N-1}\}$ the canonical basis of \mathbb{R}^{N-1} and write x_1, \dots, x_{N-1} for the corresponding coordinates.

Since any face of a cone is generated by a subset of its generators, and for our given matrix A , any $(N-1) \times (N-1)$ -minor is nonzero (so that any subset of $N-1$ columns generates a full-dimensional cone), we see that any facet can contain at most $N-2$ columns. On the other hand, such facet must be $(N-2)$ -dimensional, so it cannot be generated by fewer columns. Therefore, we can conclude that it contains exactly $N-2$ columns.

Any linear functional h defining a facet of C must satisfy that $h(C) \geq 0$. Denote by $H_{k,l}$ the hyperplane not containing \underline{a}_k and \underline{a}_l . There are five classes of these hyperplanes: $H_{1,i}$, $H_{1,n+j}$, H_{i_1,i_2} , $H_{i,n+j}$, $H_{n+j_1,n+j_2}$ with $i, i_1, i_2 \in \{2, \dots, n\}$ and $j, j_1, j_2 \in \{1, \dots, m\}$. The linear functionals defining them are, respectively,

$$\begin{aligned} h_{1,i} &:= x_{m+i-1}, \\ h_{1,n+j} &:= x_j, \\ h_{i_1,i_2} &:= x_{m+i_1-1} - x_{m+i_2-1}, \\ h_{i,n+j} &:= x_j - x_{m+i-1}, \\ h_{n+j_1,n+j_2} &:= x_{j_1} - x_{j_2}. \end{aligned}$$

All of the linear forms $h_{1,i}$, h_{i_1,i_2} and $h_{n+j_1,n+j_2}$ (for the corresponding values of i, i_1, i_2, j_1, j_2) take both negative and positive values on some columns of A , so the associated hyperplanes do not contain any facet.

We conclude that each facet of C is contained in one of the following hyperplanes:

$$\begin{aligned} H_{1,n+j} : x_j &= 0 \quad \text{for } j = 1, \dots, m, \\ H_{i,n+j} : x_j - x_{m+i-1} &= 0 \quad \text{for } i = 2, \dots, n, j = 1, \dots, m. \end{aligned} \tag{4}$$

These hyperplanes are different from each other and the respective functionals satisfy $h_{1,n+j}(C) \geq 0$ and $h_{i,n+j}(C) \geq 0$. Hence each of them contains a different facet of the cone C .

The primitive, inward-pointing normal vectors of the hyperplanes $H_{1,n+j}$ and $H_{i,n+j}$ are $\underline{n}_{1,n+j} := \underline{u}_j$ and $\underline{n}_{i,n+j} := \underline{u}_j - \underline{u}_{m+i-1}$, respectively. Denote by \underline{c} the sum of all columns of A . We have $\underline{c} = 2(\underline{u}_1 + \dots + \underline{u}_m)$ and $e_{k,l} := \langle \underline{n}_{k,l}, \underline{c} \rangle = 2$, where k and l take the admissible values corresponding to the hyperplanes we consider in (4) (i.e., we have either $(k, l) = (1, n+j)$ or $(k, l) = (i, n+j)$ for $i = 2, \dots, n$ and $j = 1, \dots, m$). Define

$$\begin{aligned} \mathfrak{A}_{k,l} &:= H_{k,l} - \left[0, \frac{1}{e_{k,l}}\right] \cdot \underline{c} \\ &= H_{k,l} - [0, 1] \cdot (\underline{u}_1 + \dots + \underline{u}_m) \\ &= \begin{cases} H_{1,n+j} - [0, 1] \cdot \underline{u}_j & \text{for } j = 1, \dots, m, \\ H_{i,n+j} - [0, 1] \cdot \underline{u}_j & \text{for } i = 2, \dots, n, j = 1, \dots, m, \end{cases} \end{aligned}$$

since for $(k, l) = (1, n+j)$ and $(k, l) = (i, n+j)$, the vectors $\underline{u}_1, \dots, \underline{u}_{j-1}, \underline{u}_{j+1}, \dots, \underline{u}_m$ are contained in $H_{1,n+j}$ and $H_{i,n+j}$, respectively. Then we have

$$\mathfrak{A}_{1,n+j} = H_{1,n+j} - [0, 1] \cdot \underline{u}_j = \{(x_1, \dots, x_{N-1}) \in \mathbb{R}^{N-1} \mid -1 < x_j \leq 0\}$$

for all $j = 1, \dots, m$ and

$$\mathfrak{A}_{i,n+j} = H_{i,n+j} - [0, 1] \cdot \underline{u}_j = \{(x_1, \dots, x_{N-1}) \in \mathbb{R}^{N-1} \mid -1 < x_j - x_{m+i-1} \leq 0\}$$

for all $i = 2, \dots, n, j = 1, \dots, m$. According to the construction given before Theorem 4.4, we can conclude that

$$\mathfrak{A}_A = \bigcap_{F \text{ facet}} \{\mathbb{R} \cdot F - [0, \frac{1}{e_F}) \cdot \underline{c}\} = \bigcap_{k, l \text{ from (4)}} \mathfrak{A}_{k, l},$$

so we can describe the admissible region \mathfrak{A}_A as

$$\mathfrak{A}_A : \begin{cases} -1 < x_j \leq 0 & \text{for } j = 1, \dots, m, \\ -1 < x_j - x_{m+i-1} \leq 0 & \text{for } i = 2, \dots, n, j = 1, \dots, m \end{cases} \subset \mathbb{R}^{N-1}.$$

Now let us pick a point $p \in [0, 1)^{N-1} \cap (\mathfrak{A}_A + \mathbb{N}^{N-1})$, and take $\underline{k} = (k_1, \dots, k_{N-1}) \in \mathbb{N}^{N-1}$ such that $p \in [0, 1)^{N-1} \cap (\mathfrak{A}_A + \underline{k})$. The shifted domain is given by

$$\mathfrak{A}_A + \underline{k} : \begin{cases} -1 + k_j < x_j \leq k_j & \text{for } j = 1, \dots, m, \\ -1 + k_j - k_{m+i-1} < x_j - x_{m+i-1} \leq k_j - k_{m+i-1} & \text{for } i = 2, \dots, n, j = 1, \dots, m \end{cases} \subset \mathbb{R}^{N-1}.$$

Assume first there is a vanishing coordinate p_{j_0} . Then we must have $k_{j_0} = 0$. For such an index and any $i = 1, \dots, n-1$, we can consider the $n-1$ inequalities

$$-1 - k_{m+i} < -q_i \leq -k_{m+i},$$

from where we deduce that every q_i belongs to $[k_{m+i}, k_{m+i} + 1) \cap [0, 1)$, for $i = 1, \dots, n-1$. In order for those intersections to be nonempty, we must have $k_{m+i} + 1 > 0$ and $k_{m+i} < 1$, so necessarily $k_{m+i} = 0$ for all i (and hence q_i must lie within $[0, 1)$, which is no new information).

Now, for any nonvanishing p_j , it is clear that $k_j = 1$. Then, if we look at the remaining inequalities, we see that

$$0 < p_j - q_i \leq 1,$$

for every $i = 1, \dots, n-1$, and any $j \in \{1, \dots, m\}$ such that $p_j \neq 0$. Therefore, every q_i belongs to $[0, 1) \cap \bigcap_{p_j \neq 0} [p_j - 1, p_j) = [0, p_-)$. Obviously, if $p_j = 0$ for all $j = 1, \dots, m$, we obtain that the q_i belong all to $[0, 1) = [0, p_-)$.

Assume now that no p_j vanishes. Then $k_1 = \dots = k_m = 1$. It follows that we can express the shifted region $\mathfrak{A}_A + \underline{k}$ as

$$\mathfrak{A}_A + \underline{k} : \begin{cases} 0 < x_j \leq 1 & \text{for } j = 1, \dots, m, \\ -k_{m+i-1} < x_j - x_{m+i-1} \leq 1 - k_{m+i-1} & \text{for } i = 2, \dots, n, j = 1, \dots, m \end{cases} \subset \mathbb{R}^{N-1}.$$

Then, for any $j = 1, \dots, m$, we have $q_i \in [0, 1) \cap [p_j + k_{m+i} - 1, p_j + k_{m+i})$, for $i = 1, \dots, n-1$. As before, this implies that $p_j + k_{m+i} > 0$ and $p_j + k_{m+i} - 1 < 1$, for each $j = 1, \dots, m$. Since each p_j lives in $(0, 1)$, the k_{m+i-1} can only be either 0 or 1.

Pick an $i \in \{1, \dots, n-1\}$ such that $k_{m+i} = 0$. Then, as before,

$$q_i \in \bigcap_{j=1}^m [p_j - 1, p_j) \cap [0, 1) = [0, p_-).$$

If our index i is such that $k_{m+i} = 1$, then

$$q_i \in \bigcap_{j=1}^m [p_j, p_j + 1) \cap [0, 1) = [p_+, 1),$$

and one direction of the statement is done.

To show the other implication of the lemma, suppose now that every q_i lies within $[0, p_-) \cup [p_+, 1)$ for $i = 1, \dots, n-1$, and no p_j vanishes. We can rewrite this as a disjunction: either $q_i \in \bigcap_{j=1}^m [0, p_j) = [0, 1) \cap \bigcap_{j=1}^m [p_j - 1, p_j)$ or $q_i \in \bigcap_{j=1}^m [p_j, 1) = [0, 1) \cap \bigcap_{j=1}^m [p_j, p_j + 1)$. If $q_i \in [0, p_-)$, define $k_{m+i} := 0$. Otherwise, we take $k_{m+i} := 1$. Summing up, it is clear that

$$p \in (\mathfrak{A}_A + (1, \overset{(m)}{\cdot}, 1, k_{m+1}, \dots, k_{N-1})) \cap [0, 1)^{N-1}.$$

If some p_j vanishes, and every q_i belongs to $[0, p_-)$, we can do the same as above to see that

$$p \in (\mathfrak{A}_A + (k_1, \dots, k_m, 0, \dots, 0)) \cap [0, 1)^{N-1},$$

where k_j vanishes if so does p_j and is equal to 1 if $p_j \neq 0$. □

As a consequence of the above calculation of the set of admissible parameters, let us prove a result extending [Castaño Domínguez and Sevenheck 2019, Theorem 2.13].

Theorem 5.7. *Let $\alpha_1, \dots, \alpha_n$ and β_1, \dots, β_m be real numbers, lying on the interval $[0, 1)$ and increasingly ordered. Assume moreover that:*

- *No difference $\alpha_i - \beta_j$ is zero, for any $i = 1, \dots, n$ and $j = 1, \dots, m$.*
- *After applying the bijection $[0, 1) \rightarrow S^1$ given by $x \mapsto e^{2\pi i x}$, all the images of the α_i are at one arc of the unit circle, while those of the β_j find themselves at the complementary arc. (In other words and going back to the interval $[0, 1)$, either no α_i belongs to any interval (β_j, β_{j+1}) or vice versa.)*

Consider the operators P and H given by

$$P = z^2 \partial_z + (n - m) t z \partial_t + \varepsilon z \quad \text{and} \quad H = \prod_{i=1}^n z(t \partial_t - \alpha_i) - t \prod_{j=1}^m z(t \partial_t - \beta_j),$$

with $\varepsilon = \sum_{j=1}^m \beta_j - \sum_{i=1}^n \alpha_i + N - 1$. Let $\widehat{\mathcal{H}}(\alpha_i; \beta_j)$ be the $\mathcal{R}_{\mathbb{A}_z^1 \times \mathbb{G}_m}^{\text{int}}$ -module

$$\widehat{\mathcal{H}}(\alpha_i; \beta_j) := \mathcal{O}_{\mathbb{A}_z^1 \times \mathbb{G}_m} \langle z^2 \partial_z, z t \partial_t \rangle / (P, H).$$

Then, $\widehat{\mathcal{H}}(\alpha_i; \beta_j)$ underlies a unique object of $\text{IrrMHM}(\mathbb{G}_m)$ with associated $\mathcal{D}_{\mathbb{G}_m}$ -module $\mathcal{H}(\alpha_i; \beta_j)$. It can be uniquely extended to an irreducible $\mathcal{R}_{\mathbb{A}_z^1 \times \mathbb{P}^1}^{\text{int}}$ -module underlying an object of $\text{IrrMHM}(\mathbb{P}^1)$.

Proof. Let us assume first that $\alpha_1 = 0$. Then, by the first assumption on the α_i and the β_j , we have $\beta_j \neq 0$ for every j . By the second assumption we can deduce that no α_i is between any two β_j , but all of the β_j must be between two certain α_i . Thanks to Lemma 5.6, this means that $\gamma := (\beta_1, \dots, \beta_m, \alpha_2, \dots, \alpha_n)$

belongs to $\mathfrak{A}_A + \mathbb{N}^{N-1}$, where A is the matrix of the beginning of the section. As a consequence, by Proposition 5.4 and Remark 5.5 we have that

$$\mathrm{For}_2(\mathrm{FL}_{\mathrm{MTM}}^\psi(\check{\mathcal{M}}_A^\vee)) \cong \widehat{\mathcal{H}}(\alpha_i; \beta_j)$$

(recall that $\check{\mathcal{M}}_A^\vee$ is the algebraic integrable mixed twistor \mathcal{D} -module with underlying $\mathcal{R}_{\mathbb{A}^N}^{\mathrm{int}}$ -module $\mathcal{R}^{F^H} \check{\mathcal{M}}_A^\vee$, i.e., such that $\mathrm{For}_2(\check{\mathcal{M}}_A^\vee) = \mathcal{R}^{F^H} \check{\mathcal{M}}_A^\vee$). We have moreover that $\check{\mathcal{M}}_A^\vee \in \mathrm{IrrMHM}(\mathbb{A}^N)$ and thanks to [Sabbah 2018, Corollary 0.5], we know that the functors entering in the definition of $\mathrm{FL}_{\mathrm{MTM}}^\psi$ preserve the category of irregular mixed Hodge modules, so we conclude that $\widehat{\mathcal{H}}(\alpha_i; \beta_j)$ underlies an element of $\mathrm{IrrMHM}(\mathbb{G}_m)$.

Assume now that $\alpha_1 > 0$. For any real number η , denote by $\widehat{\mathcal{K}}_\eta$ the Kummer $\mathcal{R}_{\mathbb{A}^1 \times \mathbb{G}_m}$ -module $\mathcal{R}_{\mathbb{A}^1 \times \mathbb{G}_m}^{\mathrm{int}} / (z^2 \partial_z, tz \partial_t - z\eta)$.

The tensor product of $\mathcal{R}_{\mathbb{A}^1 \times \mathbb{G}_m}^{\mathrm{int}}$ -modules $\widehat{\mathcal{H}}(\alpha_i; \beta_j) \otimes_{\mathcal{O}_{\mathbb{A}^1 \times \mathbb{G}_m}} \widehat{\mathcal{K}}_{-\alpha_1}$ gives rise to the corresponding tensor product of twistor \mathcal{D} -modules on \mathbb{G}_m . This product can be presented as $\widehat{\mathcal{H}}(\alpha'_i; \beta'_j)$, where $\alpha'_i = \alpha_i - \alpha_1$ for every i and $\beta'_j = \beta_j - \alpha_1$ for every j . The assumptions on the parameters imply that $\alpha'_1 = 0$ and the vector $(\beta'_1, \dots, \beta'_m, \alpha'_2, \dots, \alpha'_n)$ lives in $\mathfrak{A}_A + \mathbb{N}^{N-1}$. Then, arguing as before, such tensor product is an irregular mixed Hodge module of exponential-Hodge origin. Since $\widehat{\mathcal{K}}_{\alpha_1}$ is the faithful image of a mixed Hodge module on \mathbb{G}_m , the tensor product with it preserves the condition of being in $\mathrm{IrrMHM}(\mathbb{G}_m)$ due to [Sabbah 2018, Corollary 0.5], and so is the case of our original $\mathcal{R}_{\mathbb{A}^1 \times \mathbb{G}_m}^{\mathrm{int}}$ -module

$$\widehat{\mathcal{H}}(\alpha_i; \beta_j) \cong \widehat{\mathcal{H}}(\alpha'_i; \beta'_j) \otimes_{\mathcal{O}_{\mathbb{A}^1 \times \mathbb{G}_m}} \widehat{\mathcal{K}}_{\alpha_1}.$$

This ends the statement on the existence. Let us prove now the claims on the unicity, as in [Castaño Domínguez and Sevenheck 2019, Theorem 2.13], noting that the condition on the differences $\alpha_i - \beta_j$ is equivalent to \mathcal{H} being irreducible, and thus rigid (see [ibid., Proposition 2.5], noting that all the parameters belong to $[0, 1)$).

Consider now any twistor \mathcal{D} -module $\widehat{\mathcal{H}}'$ on $\mathbb{G}_{m,t}$ whose underlying $\mathcal{D}_{\mathbb{G}_{m,t}}$ -module is \mathcal{H} . Since the functor Ξ_{DR} is faithful by [Mochizuki 2015a, Remark 7.2.9], we have an injection of Hom groups

$$\mathrm{Hom}_{\mathrm{MTM}(\mathbb{G}_{m,t})}(\widehat{\mathcal{H}}, \widehat{\mathcal{H}}') \hookrightarrow \mathrm{Hom}_{\mathcal{D}_{\mathbb{G}_{m,t}}}(\mathcal{H}, \mathcal{H}).$$

But \mathcal{H} is irreducible, so its only endomorphism is the identity and then the twistor \mathcal{D} -module underlying \mathcal{H} is unique.

On the other hand, let $j : \mathbb{G}_{m,t} \hookrightarrow \mathbb{P}^1$ be the canonical inclusion and consider the $\mathcal{D}_{\mathbb{P}^1}$ -module $\mathcal{H}_{pr} := j_+ \mathcal{H}$. It is an irreducible holonomic $\mathcal{D}_{\mathbb{P}^1}$ -module, because so is \mathcal{H} by the assumption on the α_i and the β_j . Then it gives rise to a unique pure integrable twistor \mathcal{D} -module $\widehat{\mathcal{H}}_{pr}$ on \mathbb{P}^1 by [Mochizuki 2011, Theorem 1.4.4; Sabbah 2018, Remark 1.39]. In addition, its underlying $\mathcal{D}_{\mathbb{P}^1}$ -module \mathcal{H}_{pr} is rigid, as \mathcal{H} was. As a consequence, we can invoke [ibid., Theorem 0.7] and claim that such twistor \mathcal{D} -module on \mathbb{P}^1 is in fact an object of $\mathrm{IrrMHM}(\mathbb{P}^1)$. Take now $\widehat{\mathcal{H}}' := j^+ \widehat{\mathcal{H}}_{pr}$, which is an irregular mixed Hodge module whose underlying $\mathcal{D}_{\mathbb{G}_{m,t}}$ -module is \mathcal{H} , by [Mochizuki 2015a, Proposition 14.1.24]. Then we must have, as was just shown, $\widehat{\mathcal{H}}' \cong \widehat{\mathcal{H}}$, so that the extension $\widehat{\mathcal{H}}_{pr}$ of $\widehat{\mathcal{H}}$ is unique, and we are done. \square

Remark 5.8. Let us consider the last theorem for the case $m = n$, that is, the case of regular hypergeometric systems. Consider $\widehat{\mathcal{H}}$ as a $\mathcal{R}_{\mathbb{A}_z^1 \times \mathbb{G}_m}$ -module only, as such it is isomorphic to $\mathcal{R}_{\mathbb{A}_z^1 \times \mathbb{G}_m}/(H)$, where now

$$H = \prod_{i=1}^m z(t\partial_t - \alpha_i) - t \prod_{j=1}^m z(t\partial_t - \beta_j).$$

$\mathcal{R}_{\mathbb{A}_z^1 \times \mathbb{G}_m}$ is graded by degree in z (where z has degree 1), and since H is homogenous (which is not the case if $n \neq m$), we see that $\widehat{\mathcal{H}}$ is a *graded* $\mathcal{R}_{\mathbb{A}_z^1 \times \mathbb{G}_m}$ -module. It is obviously *strict*, i.e., it has no z -torsion, and then by [Sabbah and Schnell 2018, A.7(5)], we see that $\widehat{\mathcal{H}}$ is the Rees module of a filtered $\mathcal{D}_{\mathbb{G}_m}$ -module, namely, the (regular) hypergeometric module $\mathcal{H}(\alpha_i; \beta_j)$ together with the filtration by order of differential operators. Notice also that if $n = m$, we have $P = z^2\partial_z + \varepsilon z$, which implies that $\widehat{\mathcal{H}}$ has an action by $z\partial_z$ and that if we write $\widehat{\mathcal{H}} = \bigoplus_k \widehat{\mathcal{H}}_k$ (grading with respect to z), then for any $m \in \widehat{\mathcal{H}}_k$, we have $(z\partial_z)(m) = (k - \epsilon)m$.

Now suppose that we have $n = m$ and that additionally the hypotheses of the last theorem are satisfied, then since $\widehat{\mathcal{H}}(\alpha_i; \beta_j)$ is the unique object in $\text{IrrMHM}(\mathbb{G}_m)$ (lying actually in the essential image of $\text{MHM}(\mathbb{G}_m)$) with underlying $\mathcal{D}_{\mathbb{G}_m}$ -module $\mathcal{H}(\alpha_i; \beta_j)$, it is the Rees module of the filtered module $(\mathcal{H}(\alpha_i; \beta_j), F_\bullet^H)$, where F_\bullet^H denotes the Hodge filtration of the complex variation of Hodge structures on $\mathcal{H}(\alpha_i; \beta_j)$. Hence $F_\bullet^H \mathcal{H}(\alpha_i; \beta_j) = F_\bullet^{\text{ord}} \mathcal{H}(\alpha_i; \beta_j)$ in this case. Moreover, if we put

$$R_k := \prod_{i=1}^k (t\partial_t - \alpha_i)$$

for $k = 0, \dots, n-1$ (where $R_0 := 1$), then $(R_k)_{k=0, \dots, n-1}$ is an $\mathcal{O}_{\mathbb{G}_m}$ -basis of $\mathcal{H}(\alpha_i, \beta_j)$ and yields a splitting of the Hodge filtration F_\bullet^H . In particular, we obtain that the Hodge numbers $h^p(\mathcal{H}(\alpha_i; \beta_j)) = \dim(F_k^H / F_{k-1}^H)$ are all equal to one. This is consistent with [Fedorov 2018, Theorem 1] (up to an overall shift, as noticed in that theorem) in the version of [Castaño Domínguez and Sevenheck 2019, Proposition 2.6], since under the assumption of Theorem 5.7, the function $\#\{j : \beta_j < \alpha_k\}$ is constant.

We will finish this section with a calculation of an irregular Hodge filtration, similar to the last section of [Castaño Domínguez and Sevenheck 2019]. In that reference, the authors computed such a filtration in the case where the hypergeometric \mathcal{D} -module had a purely irregular singularity at infinity, that is, it was of type $(n, 0)$. It is immediate to see that for modules of type $(n, 1)$, the second assumption of Theorem 5.7 holds true, so that we obtain an explicit description of the $\mathcal{R}_{\mathbb{A}_z^1 \times \mathbb{G}_m}^{\text{int}}$ -module underlying the irregular Hodge module with associated $\mathcal{D}_{\mathbb{G}_m}$ -module $\mathcal{H}(\alpha_1, \dots, \alpha_n; \beta)$. In the sequel, we are going to compute the irregular Hodge filtration of such modules of type $(n, 1)$.

Let us recall the conventions and notations used in [Castaño Domínguez and Sevenheck 2019, §4] (compare [Sabbah 2018, Notation 2.1]). We will deal with the classical hypergeometric \mathcal{D} -module $\mathcal{H} = \mathcal{H}(\alpha_i; \beta)$, where the α_i and β are $n+1$ real numbers belonging to the interval $[0, 1)$. We will denote by $\widehat{\mathcal{H}}$ both its associated algebraic, integrable twistor \mathcal{D} -module on \mathbb{G}_m and its underlying $\mathcal{R}_{\mathbb{A}_z^1 \times \mathbb{G}_m}^{\text{int}}$ -module (as in the statement of Theorem 5.7). From now on, we will write \mathcal{X} , ${}^\theta\mathcal{X}$ and ${}^\tau\mathcal{X}$ meaning

the products $\mathbb{A}_z^1 \times \mathbb{G}_{m,t}$, $\mathcal{X} \times \mathbb{G}_{m,\theta}$, and $\mathcal{X} \times \mathbb{A}_t^1$, respectively, where $\theta = 1/\tau$. Finally, we will write ${}^\tau\mathcal{X}_0 = \mathcal{X} \times \{\tau = 0\} \subset {}^\tau\mathcal{X}$.

Theorem 5.9. *Let real numbers $\alpha_1, \dots, \alpha_n, \beta \in [0, 1]$ be given. Suppose that $\alpha_1 \leq \dots \leq \alpha_n$ and that moreover $\alpha_i - \beta \notin \mathbb{Z}$ for all $i = 1, \dots, n$. For each $k = 1, \dots, n$, set $\rho(k) = -(n-1)\alpha_k + k$. Then the jumping numbers of the irregular Hodge filtration of $\mathcal{H} = \mathcal{H}(\alpha_i; \beta)$ are, up to an overall real shift, the numbers $\rho(k)$. The irregular Hodge numbers are the multiplicities of those jumping numbers, or equivalently, the nonzero values of $|\rho^{-1}(x)|$, for x real.*

Moreover, for $r = 0, \dots, n-1$, let $v_\alpha(r) = \lceil -\alpha + r - \varepsilon - (n-1)\alpha_{r+1} \rceil$ (recall from Theorem 5.7 that $\varepsilon = \beta - \sum_{i=1}^n \alpha_i + n$). Let us consider the operators

$$\bar{Q}_r = (-(n-1))^r \prod_{i=1}^r (t\partial_t - \alpha_i)$$

for $r = 0, \dots, n-2$ (where the empty product equals one) and

$$\bar{Q}_{n-1} = (-(n-1))^{n-1} \prod_{i=1}^{n-1} (t\partial_t - \alpha_i) + \frac{(-(n-1))^{n-1} t(\beta - \alpha_1)}{1 + \alpha_1 - \alpha_n} \bar{Q}_0.$$

Then, the irregular Hodge filtration $F_\bullet^{\text{irr}}\mathcal{H}$ is given by

$$F_{\alpha+j}^{\text{irr}}\mathcal{H} = \bigoplus_{k: j \geq v_\alpha(k)} \mathcal{O}_X \bar{Q}_k.$$

Remark 5.10. In general, the procedure given below can be of use to find an explicit expression for the irregular Hodge filtration, not only the numbers, of any hypergeometric of type (n, m) , provided both assumptions from Theorem 5.7 are fulfilled. However, the calculations become soon too cumbersome to be included here.

Proof. We will mimic the arguments of [Castaño Domínguez and Sevenheck 2019, §4], providing almost no proof of the claims which are similar to some therein.

We must first consider the rescaling of $\widehat{\mathcal{H}}$: this is the inverse image ${}^\theta\widehat{\mathcal{H}} := \mu^*\mathcal{H}$ (as $\mathcal{O}_{\theta\mathcal{X}}$ -module), endowed with a natural action of $\mathcal{R}_{\theta\mathcal{X}}^{\text{int}}$ as depicted in [Sabbah 2018, (2.4)] (note that $\theta = \tau^{-1}$), where μ is the morphism given in [ibid., Notation 2.1] by

$$\begin{aligned} \mu : {}^\theta\mathcal{X} &\rightarrow \mathcal{X} \\ (z, t, \theta) &\mapsto (z\theta, t). \end{aligned}$$

In this sense, we can apply the same argument of [Castaño Domínguez and Sevenheck 2019, Proposition 4.1] to get that the $\mathcal{R}_{\theta\mathcal{X}}^{\text{int}}$ -module ${}^\theta\widehat{\mathcal{H}}$ associated with $\widehat{\mathcal{H}}$ can be presented as $\mathcal{R}_{\theta\mathcal{X}}^{\text{int}}/(P, {}^\theta R, {}^\theta H)$, where $P = z^2\partial_z + (n-m)t z\partial_t + \varepsilon z$ as in Theorem 5.7, ${}^\theta R = z^2\partial_z - z\theta\partial_\theta$ and

$${}^\theta H = \prod_{i=1}^n z\theta(t\partial_t - \alpha_i) - t z\theta(t\partial_t - \beta).$$

Now we have to invert θ to obtain an $\mathcal{R}_{\tau\mathcal{X}}^{\text{int}}(*^\tau\mathcal{X}_0)$ -module ${}^\tau\widehat{\mathcal{H}}$, to work in the setting given by [Sabbah 2018, §2.3]. In this sense, we will denote by ${}^\tau\widehat{\mathcal{H}}$ the $\mathcal{R}_{\tau\mathcal{X}}^{\text{int}}(*^\tau\mathcal{X}_0)$ -module $(\text{id}_{\mathcal{X}} \times (j \circ \text{inv}))_* {}^\theta\widehat{\mathcal{H}}$, where $\text{inv} : \mathbb{G}_{m,\theta} \rightarrow \mathbb{G}_{m,\tau}$ is the inversion operator $\theta \mapsto \tau^{-1}$ and $j : \mathbb{G}_{m,\tau} \hookrightarrow \mathbb{A}_\tau^1$ is the canonical inclusion. Then it is easy to see that ${}^\tau\widehat{\mathcal{H}} = \mathcal{R}_{\tau\mathcal{X}}^{\text{int}}(*^\tau\mathcal{X}_0)/(P, {}^\tau R, {}^\tau H)$, with P as always, ${}^\tau R = z^2\partial_z + z\tau\partial_\tau$ and

$${}^\tau H = \prod_{i=1}^n \frac{z}{\tau} (t\partial_t - \alpha_i) - t \frac{z}{\tau} (t\partial_t - \beta).$$

The next step is forming the basis of ${}^\tau\widehat{\mathcal{H}}$ as a $\mathcal{O}_{\tau\mathcal{X}}(*^\tau\mathcal{X}_0)$ -module. Let it be given by

$$Q_k = (-(n-1))^k \prod_{i=1}^k \frac{z}{\tau} (t\partial_t - \alpha_i)$$

for $k = 0, \dots, n-2$ and

$$Q_{n-1} = (-(n-1))^{n-1} \prod_{i=1}^{n-1} \frac{z}{\tau} (t\partial_t - \alpha_i) + \frac{(-(n-1))^{n-1} t(\beta - \alpha_1)}{1 + \alpha_1 - \alpha_n} Q_0.$$

It is indeed a basis: we can use the expressions of ${}^\tau R$ and P to replace the classes of $z\tau\partial_\tau$ and $z^2\partial_z$, respectively, in terms of $zt\partial_t$. Now ${}^\tau\widehat{\mathcal{H}}$ is generated as a $\mathcal{O}_{\tau\mathcal{X}}(*^\tau\mathcal{X}_0)$ -module by the powers of $zt\partial_t$, and we can get rid of those of exponent greater than $n-1$ using ${}^\tau H$. The remaining n powers can be expressed as a linear combination of the Q_i , forming a triangular matrix (almost diagonal in fact), so the latter conform a basis as well.

One could wonder about the odd expression of the Q_i . In the case with no betas of [Castaño Domínguez and Sevenheck 2019], the basis considered there was formed just by the successive products $\prod_{i=1}^k \frac{z}{\tau} (t\partial_t - \alpha_i)$, up to some constant. In this case, such a basis does not provide a connection matrix solving the Birkhoff problem with a diagonal matrix as a coefficient of the pole at infinity in z , which would give us a way to read the spectrum from that matrix (see [de Gregorio et al. 2009, Proposition 4.8]). As a consequence, we have to adapt such initial basis, and that is how we get the Q_i . Let us write the connection matrix explicitly.

Let $c = (\beta - \alpha_1)/(1 + \alpha_1 + \alpha_n)$, in such a way that

$$Q_{n-1} = (-(n-1))^{n-1} \prod_{i=1}^{n-1} \frac{z}{\tau} (t\partial_t - \alpha_i) + (-(n-1))^{n-1} ct Q_0.$$

A similar (but longer) calculation to the proof of [Castaño Domínguez and Sevenheck 2019, Lemma 4.3] shows that the integrable connection arising from the $\mathcal{R}_{\tau\mathcal{X}}^{\text{int}}(*^\tau\mathcal{X}_0)$ -module structure associated with ${}^\tau\widehat{\mathcal{H}}$ has the following matrix form:

$$\nabla \underline{Q} = \underline{Q} \left((\tau A_0 + z A_\infty) \frac{dz}{z^2} + (-\tau A_0 + z A'_\infty) \frac{dt}{(n-1)zt} - (\tau A_0 + z A_\infty) \frac{d\tau}{z\tau} \right).$$

There, if $n > 2$, A_0 , A'_∞ and A_∞ are the matrices

$$A_0 = \begin{pmatrix} 0 & \cdots & -(-(n-1))^{n-1}ct & 0 \\ 1 & \ddots & & (-(-(n-1))^{n-1}(c+1)t \\ & \ddots & 0 & \vdots \\ & & 1 & 0 \end{pmatrix}, \quad (5)$$

$$A'_\infty = \text{diag}((n-1)\alpha_1, \dots, (n-1)\alpha_n), \text{ and}$$

$$A_\infty = \text{diag}(0, 1, \dots, n-1) - \varepsilon I_n - A'_\infty.$$

If $n = 2$, we have

$$A_0 = \begin{pmatrix} ct & c(c+1)t^2 \\ 1 & (c+1)t \end{pmatrix}, \quad A'_\infty = \begin{pmatrix} \alpha_1 & 0 \\ 0 & \alpha_2 \end{pmatrix} \quad \text{and} \quad A_\infty = \text{diag}(0, 1) - \varepsilon I_2 - A'_\infty. \quad (6)$$

Finally, the irregular Hodge filtration is obtained from a suitable V -filtration along the divisor $\tau = 0$ defined on ${}^\tau\widehat{\mathcal{H}}$, which is called ${}^\tau V$ -filtration (the new symbol ${}^\tau V$ is to make clear the variety over which we are working; note the same convention from Remarks 2.20 on in [Sabbah 2018]). We are actually defining a filtration on ${}^\tau\widehat{\mathcal{H}}$, and then prove that it equals the ${}^\tau V$ -filtration, following [Mochizuki 2015a, §2.1.2].

Let us consider

$${}^\tau U_\alpha {}^\tau\widehat{\mathcal{H}} := \left\{ \sum_{k=0}^{n-1} f_k \tau^{v_k} Q_k : f_k \in \mathcal{O}_{\tau\mathcal{X}}, \max(k - (n-1)\alpha_{k+1} - \varepsilon - v_k) \leq \alpha \right\}, \quad (7)$$

$${}^\tau U_{<\alpha} {}^\tau\widehat{\mathcal{H}} := \left\{ \sum_{k=0}^{n-1} f_k \tau^{v_k} Q_k : f_k \in \mathcal{O}_{\tau\mathcal{X}}, \max(k - (n-1)\alpha_{k+1} - \varepsilon - v_k) < \alpha \right\},$$

for any $\alpha \in \mathbb{R}$.

The ${}^\tau U_\alpha {}^\tau\widehat{\mathcal{H}}$ form an increasing filtration, indexed by the real numbers but with a discrete set of jumping numbers, such that $\tau {}^\tau U_\alpha {}^\tau\widehat{\mathcal{H}} = {}^\tau U_{\alpha-1} {}^\tau\widehat{\mathcal{H}}$ for any α (those are conditions i and ii' in [Mochizuki 2015a, §2.1.2]). As usual, the graded piece associated with α is $\text{Gr}_\alpha^{{}^\tau U} {}^\tau\widehat{\mathcal{H}} = {}^\tau U_\alpha {}^\tau\widehat{\mathcal{H}} / {}^\tau U_{<\alpha} {}^\tau\widehat{\mathcal{H}}$.

In (7), all the exponents v_k of the powers of τ accompanying the $f_k Q_k$ satisfy that $v_k \geq -\alpha + k - (n-1)\alpha_{k+1} - \varepsilon$. Then we can define the steps of the filtration in the same alternative way as in [Castaño Domínguez and Sevenheck 2019, Remark 4.5] as the free $\mathcal{O}_{\tau\mathcal{X}}$ -modules of finite rank

$${}^\tau U_\alpha {}^\tau\widehat{\mathcal{H}} = \bigoplus_{k=0}^{n-1} \mathcal{O}_{\tau\mathcal{X}} \cdot \tau^{v_\alpha(k)} Q_k, \quad (8)$$

where $v_\alpha(k) = \lceil -\alpha + k - \varepsilon - (n-1)\alpha_{k+1} \rceil$. With that expression, it is clear that the graded pieces $\text{Gr}_\alpha^{{}^\tau U} {}^\tau\widehat{\mathcal{H}}$ are

$$\text{Gr}_\alpha^{{}^\tau U} {}^\tau\widehat{\mathcal{H}} = \bigoplus_{k=0}^{n-1} \mathcal{O}_{\tau\mathcal{X}} \cdot \tau^{v_\alpha(k)} Q_k,$$

which are strict $\mathcal{R}_{\mathcal{X}}$ -modules (condition iv in [Mochizuki 2015a, §2.1.2]).

The next step in the proof is proving that $\widehat{\tau\mathcal{H}}$ is strictly \mathbb{R} -specializable along ${}^{\tau}\mathcal{X}_0$ and its ${}^{\tau}V$ -filtration is actually given by the ${}^{\tau}U_{\alpha}\widehat{\tau\mathcal{H}}$. Although the proof is similar to that of [Castaño Domínguez and Sevenheck 2019, Proposition 4.6], we have to adapt it a bit to our case here.

After what we already showed, it remains to show conditions iii' and v of [Mochizuki 2015a, §2.1.2] and prove that the ${}^{\tau}U_{\alpha}\widehat{\tau\mathcal{H}}$ are coherent $V_0\mathcal{R}_{\mathcal{X}}$ -modules. Let us start by the second condition. Consider then the mappings $\mathfrak{p}, \mathfrak{e}$ given by

$$\begin{aligned} (\mathfrak{p}, \mathfrak{e}) : \mathbb{R} \times \mathbb{C} &\rightarrow \mathbb{R} \times \mathbb{C} \\ (\beta, \omega) &\mapsto (\beta + 2\Re(z\bar{\omega}), -\beta z + \omega - \bar{\omega}z^2). \end{aligned}$$

We must check that the operator $z\tau\partial_{\tau} - \mathfrak{e}(\beta, \omega)$ is nilpotent on the graded pieces $\text{Gr}_{\alpha}^U \widehat{\tau\mathcal{H}}$ only for a finite amount of $(\beta, \omega) \in \mathcal{K} := \{\beta + 2\Re(z_0\bar{\omega}) = \alpha\}$, for any value z_0 of z . Moreover, those (β, ω) should belong in fact to $\mathbb{R} \times \{0\}$ (see [Sabbah 2018, §1.3.a]), if we want to obtain the \mathbb{R} -specializability.

Take then $(\beta, \omega) \in \mathcal{K}$ and $f\tau^{\nu}Q_k \in {}^{\tau}U_{\alpha}\widehat{\tau\mathcal{H}}$, with $f \in \mathcal{O}_{\tau\mathcal{X}}$. We must have that $k - (n-1)\alpha_{k+1} - \varepsilon - \nu \leq \alpha$. Assume that $n > 2$ and $k < n-2$. Thanks to the matrix form (5) we know that

$$(z\tau\partial_{\tau} - \mathfrak{e}(\beta, \omega))f\tau^{\nu}Q_k = (z\tau\partial_{\tau} + (\nu + (n-1)\alpha_{k+1} + \varepsilon - k + \beta)z - \omega + \bar{\omega}z^2)(f)\tau^{\nu}Q_k - f\tau^{\nu+1}Q_{k+1}.$$

Recall that the α_i are increasingly ordered, lying within the interval $[0, 1)$. Thus $f\tau^{\nu+1}Q_{k+1}$ lives in ${}^{\tau}U_{\alpha}\widehat{\tau\mathcal{H}}$, for

$$k+1 - (n-1)\alpha_{k+2} - \varepsilon - \nu - 1 \leq ((k+1) - (n-1)\alpha_{k+2} - \varepsilon) - (k - (n-1)\alpha_{k+1} - \varepsilon) - 1 + \alpha \leq \alpha.$$

Now we should look at what happens to the class of $f\tau^{\nu+1}Q_{k+1}$ in the α -graded piece of $\widehat{\tau\mathcal{H}}$.

Note that $[f\tau^{\nu}Q_k] \neq 0$ if and only if $\nu + (n-1)\alpha_{k+1} + \varepsilon - k + \alpha = 0$, so

$$\begin{aligned} (z\tau\partial_{\tau} - \mathfrak{e}(\beta, \omega))f\tau^{\nu}Q_k &= (z\tau\partial_{\tau} + (\beta - \alpha)z - \omega + \bar{\omega}z^2)(f)\tau^{\nu}Q_k - f\tau^{\nu+1}Q_{k+1} \\ &= (z\tau\partial_{\tau} - 2\Re(z_0\bar{\omega})z - \omega + \bar{\omega}z^2)(f)\tau^{\nu}Q_k - f\tau^{\nu+1}Q_{k+1}. \end{aligned}$$

Now notice that τ divides $\tau\partial_{\tau}(f)$, so in fact $z\tau\partial_{\tau}(f)\tau^{\nu}Q_k \in {}^{\tau}U_{\alpha-1}\widehat{\tau\mathcal{H}}$ and then we can further reduce our expression to

$$(z\tau\partial_{\tau} - \mathfrak{e}(\beta, \omega))f\tau^{\nu}Q_k = (-\omega - 2\Re(z_0\bar{\omega})z + \bar{\omega}z^2)f\tau^{\nu}Q_k - f\tau^{\nu+1}Q_{k+1}.$$

On the other hand, $\tau^{\nu+1}Q_{k+1}$ does not vanish either in $\text{Gr}_{\alpha}^U \widehat{\tau\mathcal{H}}$ if and only if $\alpha_{k+2} = \alpha_{k+1}$. Indeed, we know that $\nu + (n-1)\alpha_{k+1} + \varepsilon - k + \alpha = 0$, so doing the same as before, $k+1 - (n-1)\alpha_{k+2} - \varepsilon - \nu - 1 = \alpha + (n-1)(\alpha_{k+2} - \alpha_{k+1})$ and the claim follows. Furthermore, in order to $(z\tau\partial_{\tau} - \mathfrak{e}(\beta, \omega))$ to vanish, we should impose that $\omega = 0$, just by looking at the coefficients of the powers of z in the expression for f .

If $k = n-2$, we obtain from (5) that

$$\begin{aligned} (z\tau\partial_{\tau} - \mathfrak{e}(\beta, \omega))f\tau^{\nu}Q_{n-2} &= (z\tau\partial_{\tau} + (\nu + (n-1)\alpha_{n-1} + \varepsilon - (n-2) + \beta)z - \omega + \bar{\omega}z^2)(f)\tau^{\nu}Q_{n-2} \\ &\quad - f\tau^{\nu+1}Q_{n-1} + f\tau^{\nu+1}(-(n-1))^{n-1}ctQ_0. \end{aligned}$$

Since $-(n-1)\alpha_1 - \varepsilon - \nu - 1 \leq -(n-1)(\alpha_1 - \alpha_{n-1} + 1) + \alpha < \alpha$ because $\alpha_{n-1} < \alpha_1 + 1$, the last summand above belongs to ${}^\tau U_{<\alpha} {}^\tau \widehat{\mathcal{H}}$, and then the argument can follow as with $k < n-2$.

Now if $k = n-1$, then everything would be the same again as before except we get the additional summand $-f\tau^{\nu+1}Q_{k+1}$, which becomes $-f\tau^{\nu+1}(-(n-1))^{n-1}(c+1)tQ_1$, whose class vanishes in the graded piece under consideration, too. Indeed,

$$1 - (n-1)\alpha_2 - \varepsilon - \nu - 1 \leq -(n-1)(\alpha_2 - \alpha_n + 1) + \alpha < \alpha,$$

for $\alpha_n < \alpha_2 + 1$.

In conclusion, $(z\tau\partial_\tau - \mathfrak{e}(\beta, \omega))^l f\tau^\nu Q_k$ can only vanish in $\mathrm{Gr}_\alpha^U {}^\tau \widehat{\mathcal{H}}$ if $\alpha = \beta$ (and then $\omega = 0$), and does not do so until we get to an index $k+l$ such that α_{k+l} is strictly bigger than α_k . Since there is a finite set of indexes, $(z\tau\partial_\tau - \mathfrak{e}(\beta, \omega))$ is nilpotent, of nilpotency index n at most.

When $n = 2$, we notice from (6) that we have two possibilities. If $k = 0$, everything is the same as with $k = n-2$ for $n > 2$, and if $k = 1$,

$$\begin{aligned} (z\tau\partial_\tau - \mathfrak{e}(\beta, \omega))f\tau^\nu Q_1 \\ = (z\tau\partial_\tau + (\nu + \alpha_2 + \varepsilon - 1 + \beta)z - \omega + \bar{\omega}z^2)(f)\tau^\nu Q_1 + f\tau^{\nu+1}(c+1)tQ_1 + f\tau^{\nu+1}c(c+1)t^2Q_0. \end{aligned}$$

Here the argument runs similarly as in the general case.

Condition iii' can be rephrased as $z\tau\partial_\tau {}^\tau U_\alpha {}^\tau \widehat{\mathcal{H}} \subseteq {}^\tau U_\alpha {}^\tau \widehat{\mathcal{H}}$, using that ${}^\tau U_\alpha {}^\tau \widehat{\mathcal{H}} = \tau {}^\tau U_{\alpha+1} {}^\tau \widehat{\mathcal{H}}$, and that follows essentially from the same argument used to prove condition v above. Last, since $V_0\mathcal{R}_\mathcal{X} = \mathcal{O}_{\tau\mathcal{X}}\langle z\partial_t, z\tau\partial_\tau \rangle$, it is clear from the computations above and the alternative expression (8) for the filtration steps that they are cyclic $V_0\mathcal{R}_\mathcal{X}$ -modules, and then coherent. Summing up and noting that all the calculations performed were in fact independent of z_0 , ${}^\tau \widehat{\mathcal{H}}$ is strictly \mathbb{R} -specializable along ${}^\tau \mathcal{X}_0$ and the ${}^\tau U_\bullet {}^\tau \widehat{\mathcal{H}}$ form its ${}^\tau V$ -filtration.

We can finally show the expression for the irregular Hodge filtration and then the irregular Hodge numbers like in [Castaño Domínguez and Sevenheck 2019, Theorem 4.7]. Since we know that $\widehat{\mathcal{H}}$ underlies an object in $\mathrm{IrrMHM}(\mathbb{G}_{m,t})$ by Theorem 5.7, we deduce by [Sabbah 2018, Definition 2.52] that $\widehat{\mathcal{H}}$ is well-rescalable (see [ibid., Definition 2.19]) and so we can apply [ibid., Definition 2.22]. After formula (8), we clearly have

$$i_{\tau=z}^* {}^\tau V_\alpha {}^\tau \widehat{\mathcal{H}} = {}^\tau V_\alpha {}^\tau \widehat{\mathcal{H}} / (\tau - z) {}^\tau V_\alpha {}^\tau \widehat{\mathcal{H}} = \bigoplus_k \mathcal{O}_\mathcal{X} z^{v_\alpha(k)} \bar{Q}_k,$$

which is free z -graded of finite rank. Denote by π the projection $\mathcal{X} \rightarrow \mathbb{G}_{m,t}$. Then, the z -adic filtration on $\pi^* \mathcal{H}[z^{-1}]$ induces a filtration on $i_{\tau=z}^* {}^\tau V_\alpha {}^\tau \widehat{\mathcal{H}}$, given by

$$F_r i_{\tau=z}^* {}^\tau V_\alpha {}^\tau \widehat{\mathcal{H}} := \bigoplus_{s \leq r} \left(\bigoplus_{k: v_\alpha(k) \leq s} \mathcal{O}_{\mathbb{G}_{m,t}} \bar{Q}_k \right) z^s.$$

Then, $\mathrm{Gr}^F(i_{\tau=z}^* {}^\tau U_\alpha {}^\tau \widehat{\mathcal{H}})$ is the Rees module associated to a new good filtration $F_{\alpha+\bullet}^{\mathrm{irr}} \mathcal{H}$ on \mathcal{H} , which is the irregular Hodge filtration. More concretely, $F_\bullet^{\mathrm{irr}} \mathcal{H}$ is given by

$$F_{\alpha+j}^{\mathrm{irr}} \mathcal{H} = \bigoplus_{k: v_\alpha(k) \leq j} \mathcal{O}_{\mathbb{G}_{m,t}} \bar{Q}_k.$$

Therefore, its jumping numbers are $-\varepsilon + i - 1 - (n - 1)\alpha_i$ for $i = 1, \dots, n$. Since the irregular Hodge filtration is defined up to an overall real shift, we can normalize the jumping numbers to $i - (n - 1)\alpha_i$ and the irregular Hodge numbers will be their multiplicities. \square

Acknowledgements

We would like to thank Takuro Mochizuki for communicating the proof of Lemma 2.3 to us. We would further like to thank Takuro Mochizuki and Claude Sabbah for their interest in our work and for many stimulating discussions. We are grateful to the participants of the workshop *Mixed Twistor \mathcal{D} -modules* in Heidelberg in 2017 for their time and effort. We also thank the Max Planck Institute for Mathematics in the Sciences, where some part of the work presented here has been carried out.

References

- [Castaño Domínguez and Sevenheck 2019] A. Castaño Domínguez and C. Sevenheck, “Irregular Hodge filtration of some confluent hypergeometric systems”, *J. Inst. Math. Jussieu* (online publication May 2019).
- [Esnault et al. 2017] H. Esnault, C. Sabbah, and J.-D. Yu, “ E_1 -degeneration of the irregular Hodge filtration”, *J. Reine Angew. Math.* **729** (2017), 171–227. MR Zbl
- [Fedorov 2018] R. Fedorov, “Variations of Hodge structures for hypergeometric differential operators and parabolic Higgs bundles”, *Int. Math. Res. Not.* **2018**:18 (2018), 5583–5608. MR Zbl
- [Gelfand et al. 1987] I. M. Gelfand, M. I. Graev, and A. V. Zelevinski, “Holonomic systems of equations and series of hypergeometric type”, *Dokl. Akad. Nauk SSSR* **295**:1 (1987), 14–19. In Russian; translated in *Soviet Math. Dokl.* **36**:1 (1988), 5–10. MR Zbl
- [Gelfand et al. 1989] I. M. Gelfand, A. V. Zelevinski, and M. M. Kapranov, “Гипергеометрические функции и торические многообразия” “(Hypergeometric functions and toric varieties)”, *Funktsional. Anal. i Prilozhen.* **23**:2 (1989), 12–26. Translated as “Hypergeometric functions and toral manifolds”, *Funct. Anal. Appl.* **23**:2 (1989), 94–106. MR Zbl
- [de Gregorio et al. 2009] I. de Gregorio, D. Mond, and C. Sevenheck, “Linear free divisors and Frobenius manifolds”, *Compos. Math.* **145**:5 (2009), 1305–1350. MR Zbl
- [Katz 1990] N. M. Katz, *Exponential sums and differential equations*, Annals of Mathematics Studies **124**, Princeton University Press, 1990. MR Zbl
- [Katzarkov et al. 2017] L. Katzarkov, M. Kontsevich, and T. Pantev, “Bogomolov–Tian–Todorov theorems for Landau–Ginzburg models”, *J. Differential Geom.* **105**:1 (2017), 55–117. MR Zbl
- [Matusevich et al. 2005] L. F. Matusevich, E. Miller, and U. Walther, “Homological methods for hypergeometric families”, *J. Amer. Math. Soc.* **18**:4 (2005), 919–941. MR Zbl
- [Mochizuki 2011] T. Mochizuki, *Wild harmonic bundles and wild pure twistor \mathcal{D} -modules*, Astérisque **340**, Société Mathématique de France, Paris, 2011. MR Zbl
- [Mochizuki 2015a] T. Mochizuki, *Mixed twistor \mathcal{D} -modules*, Lecture Notes in Mathematics **2125**, Springer, 2015. MR Zbl
- [Mochizuki 2015b] T. Mochizuki, “Twistor property of GKZ-hypergeometric systems”, preprint, 2015. arXiv
- [Reichelt 2014] T. Reichelt, “Laurent polynomials, GKZ-hypergeometric systems and mixed Hodge modules”, *Compos. Math.* **150**:6 (2014), 911–941. MR Zbl
- [Reichelt and Sevenheck 2015] T. Reichelt and C. Sevenheck, “Hypergeometric Hodge modules”, preprint, 2015. To appear in *Algebraic Geometry*. arXiv
- [Sabbah 2018] C. Sabbah, *Irregular Hodge theory*, Mém. Soc. Math. Fr. (N.S.) **156**, Société Mathématique de France, Paris, 2018. MR Zbl
- [Sabbah and Schnell 2018] C. Sabbah and C. Schnell, “The MHM project”, electronic reference, 2018, Available at <http://www.cmls.polytechnique.fr/perso/sabbah/MHMPProject/mhm.html>.

- [Sabbah and Yu 2015] C. Sabbah and J.-D. Yu, “On the irregular Hodge filtration of exponentially twisted mixed Hodge modules”, *Forum Math. Sigma* **3** (2015), art. id. e9, 71 pp. MR Zbl
- [Schulze and Walther 2009] M. Schulze and U. Walther, “Hypergeometric D-modules and twisted Gauß-Manin systems”, *J. Algebra* **322**:9 (2009), 3392–3409. MR Zbl
- [Serre 1955–1956] J.-P. Serre, “Géométrie algébrique et géométrie analytique”, *Ann. Inst. Fourier, Grenoble* **6** (1955–1956), 1–42. MR Zbl
- [Yu 2014] J.-D. Yu, “Irregular Hodge filtration on twisted de Rham cohomology”, *Manuscripta Math.* **144**:1-2 (2014), 99–133. MR Zbl

Communicated by Hélène Esnault

Received 2018-07-17 Revised 2019-01-28 Accepted 2019-03-08

alberto.castano@usc.es

Instituto de Matemáticas, Universidade de Santiago de Compostela, Spain

Fakultät für Mathematik, Technische Universität Chemnitz, Chemnitz, Germany

treichelt@mathi.uni-heidelberg.de

Mathematisches Institut, Universität Heidelberg, Heidelberg, Germany

christian.sevenheck@mathematik.tu-chemnitz.de

Fakultät für Mathematik, Technische Universität Chemnitz, Chemnitz, Germany

Algebra & Number Theory

msp.org/ant

EDITORS

MANAGING EDITOR

Bjorn Poonen
Massachusetts Institute of Technology
Cambridge, USA

EDITORIAL BOARD CHAIR

David Eisenbud
University of California
Berkeley, USA

BOARD OF EDITORS

Richard E. Borcherds	University of California, Berkeley, USA	Martin Olsson	University of California, Berkeley, USA
Antoine Chambert-Loir	Université Paris-Diderot, France	Raman Parimala	Emory University, USA
J-L. Colliot-Thélène	CNRS, Université Paris-Sud, France	Jonathan Pila	University of Oxford, UK
Brian D. Conrad	Stanford University, USA	Anand Pillay	University of Notre Dame, USA
Samit Dasgupta	University of California, Santa Cruz, USA	Michael Rapoport	Universität Bonn, Germany
Hélène Esnault	Freie Universität Berlin, Germany	Victor Reiner	University of Minnesota, USA
Gavril Farkas	Humboldt Universität zu Berlin, Germany	Peter Sarnak	Princeton University, USA
Hubert Flenner	Ruhr-Universität, Germany	Joseph H. Silverman	Brown University, USA
Sergey Fomin	University of Michigan, USA	Michael Singer	North Carolina State University, USA
Edward Frenkel	University of California, Berkeley, USA	Christopher Skinner	Princeton University, USA
Wee Teck Gan	National University of Singapore	Vasudevan Srinivas	Tata Inst. of Fund. Research, India
Andrew Granville	Université de Montréal, Canada	J. Toby Stafford	University of Michigan, USA
Ben J. Green	University of Oxford, UK	Pham Huu Tiep	University of Arizona, USA
Joseph Gubeladze	San Francisco State University, USA	Ravi Vakil	Stanford University, USA
Roger Heath-Brown	Oxford University, UK	Michel van den Bergh	Hasselt University, Belgium
Craig Huneke	University of Virginia, USA	Akshay Venkatesh	Institute for Advanced Study, USA
Kiran S. Kedlaya	Univ. of California, San Diego, USA	Marie-France Vignéras	Université Paris VII, France
János Kollár	Princeton University, USA	Kei-Ichi Watanabe	Nihon University, Japan
Philippe Michel	École Polytechnique Fédérale de Lausanne	Melanie Matchett Wood	University of Wisconsin, Madison, USA
Susan Montgomery	University of Southern California, USA	Shou-Wu Zhang	Princeton University, USA
Shigefumi Mori	RIMS, Kyoto University, Japan		

PRODUCTION

production@msp.org
Silvio Levy, Scientific Editor


See inside back cover or msp.org/ant for submission instructions.

The subscription price for 2019 is US \$385/year for the electronic version, and \$590/year (+\$60, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to MSP.

Algebra & Number Theory (ISSN 1944-7833 electronic, 1937-0652 printed) at Mathematical Sciences Publishers, 798 Evans Hall #3840, c/o University of California, Berkeley, CA 94720-3840 is published continuously online. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices.

ANT peer review and production are managed by EditFlow® from MSP.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2019 Mathematical Sciences Publishers

Algebra & Number Theory

Volume 13 No. 6 2019

Positivity functions for curves on algebraic varieties	1243
BRIAN LEHMANN and JIAN XIAO	
The congruence topology, Grothendieck duality and thin groups	1281
ALEXANDER LUBOTZKY and TYAKAL NANJUNDIAH VENKATARAMANA	
On the ramified class field theory of relative curves	1299
QUENTIN GUIGNARD	
Blow-ups and class field theory for curves	1327
DAICHI TAKEUCHI	
Algebraic monodromy groups of l -adic representations of $\mathrm{Gal}(\overline{\mathbb{Q}}/\mathbb{Q})$	1353
SHIANG TANG	
Weyl bound for p -power twist of $\mathrm{GL}(2)$ L -functions	1395
RITABRATA MUNSHI and SAURABH KUMAR SINGH	
Examples of hypergeometric twistor \mathcal{D} -modules	1415
ALBERTO CASTAÑO DOMÍNGUEZ, THOMAS REICHELT and CHRISTIAN SEVENHECK	
Ulrich bundles on K3 surfaces	1443
DANIELE FAENZI	
Unlikely intersections in semiabelian surfaces	1455
DANIEL BERTRAND and HARRY SCHMIDT	
Congruences of parahoric group schemes	1475
RADHIKA GANAPATHY	
An improved bound for the lengths of matrix algebras	1501
YAROSLAV SHITOV	



1937-0652(2019)13:6;1-7