

Algebra & Number Theory

Volume 13

2019

No. 9

Positivity of anticanonical divisors
and F -purity of fibers

Sho Ejiri



Positivity of anticanonical divisors and F -purity of fibers

Sho Ejiri

We prove that given a flat generically smooth morphism between smooth projective varieties with F -pure closed fibers, if the source space is Fano, weak Fano or a variety with nef anticanonical divisor, respectively, then so is the target space. We also show that, in arbitrary characteristic, a generically smooth surjective morphism between smooth projective varieties cannot have nef and big relative anticanonical divisor, if the target space has positive dimension.

1. Introduction

Let X be a smooth projective variety over an algebraically closed field. The positivity of the anticanonical divisor $-K_X$ on X is an important notion that helps us know certain geometric properties of X . Let $f : X \rightarrow Y$ be a surjective morphism from X to another smooth projective variety Y . Kollár, Miyaoka and Mori [Kollár et al. 1992, Corollary 2.9] proved that, under the assumption that f is smooth, if X is a Fano variety, that is $-K_X$ is ample, then so is Y . It follows from an analogous argument that, under the same assumption, if $-K_X$ is nef, then so is $-K_Y$ [Miyaoka 1993; Fujino and Gongyo 2014, Theorem 1.1; Debarre 2001, Corollary 3.15(a)]. Based on these results, Yasutake asked “what positivity condition is passed from $-K_X$ to $-K_Y$?” Some answers to this question are known in characteristic 0. Fujino and Gongyo [2012, Theorem 1.1] proved that, under the assumption that f is smooth, if X is a weak Fano variety, that is $-K_X$ is nef and big, then so is Y . Birkar and Chen [2016, Theorem 1.1] showed that, under the same assumption, if $-K_X$ is semiample, then so is $-K_Y$. Furthermore, similar but weaker results hold even if f is not smooth (but the characteristic of k is still 0). For example, a result of Prokhorov and Shokurov [2009, Lemma 2.8] (see also [Fujino and Gongyo 2012, Corollary 3.3]) implies that if $-K_X$ is nef and big, then $-K_Y$ is big. Chen and Zhang [2013, Main Theorem] also proved that if $-K_X$ is nef, then $-K_Y$ is pseudoeffective.

In contrast, little was known about the positive characteristic case. In this paper, assuming that the geometric generic fiber has only F -pure or strongly F -regular singularities, we prove that (generalizations of) the statements above hold in positive characteristic, except the one concerning semiampleness. F -purity and strong F -regularity are mild singularities defined in terms of Frobenius splitting properties

MSC2010: primary 14D06; secondary 14J45.

Keywords: Fano variety, weak Fano variety, anticanonical divisor, restricted base locus, augmented base locus.

(Definition 2.4), which have a close connection to log canonical and Kawamata log terminal singularities, respectively.

Suppose that the base field k is an algebraically closed field of characteristic $p > 0$. Let $f : X \rightarrow Y$ be a surjective morphism between smooth projective varieties, Δ an effective \mathbb{Q} -divisor on X with index $\text{ind } \Delta$, and D a \mathbb{Q} -divisor on Y . Let $X_{\bar{\eta}}$ denote the geometric generic fiber of f . Then our main theorem is stated as follows:

Theorem 1.1 (Theorem 4.1). *Let y be a scheme-theoretic point in Y such that the following conditions hold:*

- (i) $\dim f^{-1}(y) = \dim X - \dim Y$.
- (ii) *The support of Δ does not contain any irreducible component of $f^{-1}(y)$.*
- (iii) $(X_{\bar{y}}, \Delta|_{X_{\bar{y}}})$ *is F -pure, where $X_{\bar{y}}$ is the geometric fiber over y .*

*Suppose that $p \nmid \text{ind}(\Delta)$ and $-(K_X + \Delta + f^*D)$ is nef. Then y is not in the Zariski closure of $\mathbf{B}_-(-(K_Y + D))$.*

Here, \mathbf{B}_- denotes the restricted base locus (Definition 2.8). This locus is empty (resp. has nonempty complement) if and only if the divisor is nef (resp. pseudoeffective). Theorem 1.1 implies, in the case of $\Delta = 0$, that every closed fiber over $\mathbf{B}_-(-(K_Y + D))$ is “bad”, where “bad” means the fiber is not F -pure or has dimension larger than that of the general fiber.

The following two theorems are corollaries of Theorem 1.1.

Theorem 1.2 (Corollary 4.5). *Suppose that conditions (i)–(iii) in Theorem 1.1 hold for every closed point in Y :*

- (1) *Assume $p \nmid \text{ind}(\Delta)$. If $-(K_X + \Delta + f^*D)$ is nef, then so is $-(K_Y + D)$.*
- (2) *If $-(K_X + \Delta + f^*D)$ is ample, then so is $-(K_Y + D)$.*

Theorem 1.3 (Corollary 4.6). *Suppose that $(X_{\bar{\eta}}, \Delta|_{X_{\bar{\eta}}})$ is F -pure:*

- (1) *If $p \nmid \text{ind}(\Delta)$ and $-(K_X + \Delta + f^*D)$ is nef, then $-(K_Y + D)$ is pseudoeffective.*
- (2) *If $-(K_X + \Delta + f^*D)$ is ample, then $-(K_Y + D)$ is big.*
- (3) *If $(X_{\bar{\eta}}, \Delta|_{X_{\bar{\eta}}})$ is strongly F -regular and $-(K_X + \Delta + f^*D)$ is nef and big, then $-(K_Y + D)$ is big.*

Theorem 1.2 is a generalization of [Kollár et al. 1992, Corollary 2.9] and [Debarre 2001, Corollary 3.15] in positive characteristic. One can also recover [Kollár et al. 1992, Corollary 2.9] in characteristic zero from Theorem 1.2, using reduction to characteristic p . Our proof relies on a study of the positivity of direct image sheaves in terms of the Grothendieck trace of the relative Frobenius morphism. This is completely different from the proof [Kollár et al. 1992, Corollary 2.9] that is an application of their great study regarding rational curves on varieties. Theorem 1.3 should be compared with [Prokhorov and Shokurov 2009, Lemma 2.8] and [Chen and Zhang 2013, Main Theorem].

The following two theorems are direct consequences of Theorems 1.2 and 1.3.

Theorem 1.4 (Corollary 4.7). *Suppose that $(X_{\bar{\eta}}, \Delta|_{X_{\bar{\eta}}})$ is F -pure. If $p \nmid \text{ind}(\Delta)$ and $K_X + \Delta$ is numerically equivalent to $f^*(K_Y + L)$ for some \mathbb{Q} -divisor L on Y , then L is pseudoeffective.*

Theorem 1.5 (Corollary 4.9). *Suppose that f is a flat morphism such that every closed fiber is F -pure and the geometric generic fiber is strongly F -regular. If X is a weak Fano variety, that is, $-K_X$ is nef and big, then so is Y .*

Theorem 1.5 is a positive characteristic counterpart of [Fujino and Gongyo 2012, Theorem 1.1].

For another application of Theorem 1.1, we return to the situation where k is of arbitrary characteristic. Suppose that $f : X \rightarrow Y$ is a generically smooth surjective morphism between smooth projective varieties of positive dimension.

Theorem 1.6 (Corollary 4.10 and Theorem 5.4). *The relative anticanonical divisor $-K_{X/Y}$ cannot be both nef and big.*

Theorem 1.7 (Corollary 4.11 and Theorem 5.5). *Suppose that $\omega_{X/\bar{\eta}}^{-m}$ is globally generated for an integer $m > 0$. Then $f_*\omega_{X/Y}^{-m}$ is not big in the sense of Definition 2.6.*


In both the theorems, the characteristic zero case is proved by reduction to positive characteristic. Theorem 1.6 includes [Kollár et al. 1992, Corollary 2.8] which states that $-K_{X/Y}$ is not ample. Theorem 1.7 generalizes [Miyaoka 1993, Corollary 2'] which says that if $\omega_{X/Y}^{-1}$ is f -ample and $\omega_{X/Y}^{-m}$ is f -free for some $m \in \mathbb{Z}_{>0}$, then $f_*\omega_{X/Y}^{-m}$ is not an ample vector bundle.

Notation. Let k be a field. A k -scheme is a separated scheme of finite type over k . A variety is an integral k -scheme. Let $\varphi : S \rightarrow T$ be a morphism of k -schemes and T' a T -scheme. Then, $S_{T'}$ and $\varphi_{T'} : S_{T'} \rightarrow T'$ denote the fiber product $S \times_T T'$ and its second projection, respectively. For a Cartier or \mathbb{Q} -Cartier divisor D on S (resp. an \mathcal{O}_S -module \mathcal{G}), the pullback of D (resp. \mathcal{G}) to $S_{T'}$ is written as $D_{T'}$ (resp. $\mathcal{G}_{T'}$) if it is well defined. Similarly, for a homomorphism $\alpha : \mathcal{F} \rightarrow \mathcal{G}$ of \mathcal{O}_S -modules, $\alpha_{T'} : \mathcal{F}_{T'} \rightarrow \mathcal{G}_{T'}$ is the pullback of α to $S_{T'}$. Assume that k is of characteristic $p > 0$. We say that k is F -finite if the field extension k/k^p is finite. Let X be a k -scheme. Then, $F_X : X \rightarrow X$ denotes the absolute Frobenius morphism of X . We often write the source of F_X^e as X^e . Let $f : X \rightarrow Y$ be a morphism between k -schemes. The same morphism is denoted by $f^{(e)} : X^e \rightarrow Y^e$ when we regard X (resp. Y) as X^e (resp. Y^e). We define the e -th relative Frobenius morphism of f to be the morphism $F_{X/Y}^{(e)} := (F_X^e, f^{(e)}) : X^e \rightarrow X \times_Y Y^e =: X_{Y^e}$. We write the localization of \mathbb{Z} at $(p) = p\mathbb{Z}$ as $\mathbb{Z}_{(p)}$.

2. Preliminaries

2A. Relative Frobenius morphisms and trace maps. In this subsection, given a morphism between varieties, we consider the relative Frobenius morphism and its trace map. Let k be an F -finite field and $f : X \rightarrow Y$ a morphism from a pure dimensional Gorenstein k -scheme X to a regular variety Y . For each

$d, e \in \mathbb{Z}_{>0}$ we use the notation defined by the following commutative diagram:



Since F_Y is flat, every horizontal morphism in the diagram is a Gorenstein morphism, so every object is a pure dimensional Gorenstein k -scheme. Let ω_X be the dualizing sheaf on X . Let $\text{Tr}_{F_{X/Y}^{(1)}} : F_{X/Y}^{(1)} \omega_{X^1} \rightarrow \omega_{X_{Y^1}}$ denote the morphism obtained by applying the functor $\mathcal{H}om_{\mathcal{O}_{X_{Y^1}}}((\cdot), \omega_{X_{Y^1}})$ to the natural morphism $F_{X/Y}^{(1)} : \mathcal{O}_{X_{Y^1}} \rightarrow F_{X/Y}^{(1)} \mathcal{O}_{X^1}$. Take a Cartier divisor K_X satisfying $\mathcal{O}_X(K_X) \cong \omega_X$. Set $K_{X/Y} := K_X - f^* K_Y$. For each $e \in \mathbb{Z}_{>0}$ we define

$$\begin{aligned} \phi_{X/Y}^{(1)} &:= \text{Tr}_{F_{X/Y}^{(1)}} \otimes \mathcal{O}_{X_{Y^1}}(-K_{X_{Y^1}}) : F_{X/Y}^{(1)} \mathcal{O}_{X^1}((1-p)K_{X^1/Y^1}) \rightarrow \mathcal{O}_{X_{Y^1}}, \quad \text{and} \\ \phi_{X/Y}^{(e+1)} &:= (\phi_{X/Y}^{(e)})_{Y^{e+1}} \circ F_{X_{Y^1}/Y^1}^{(e)} (\phi_{X/Y}^{(1)} \otimes \mathcal{O}_{X_{Y^{e+1}}}((1-p^e)K_{X_{Y^{e+1}}/Y^{e+1}})) \\ &\quad : F_{X/Y}^{(e+1)} \mathcal{O}_X((1-p^{e+1})K_{X^{e+1}/Y^{e+1}}) \rightarrow \mathcal{O}_{X_{Y^{e+1}}}. \end{aligned}$$

Let E be an effective Cartier divisor on X , let $a > 0$ be an integer not divisible by p , and let $d > 0$ be the minimum integer satisfying $a|(p^d - 1)$. Note that an integer $e > 0$ satisfies $a|(p^e - 1)$ if and only if $d|e$. Set $\Delta := E \otimes \frac{1}{a} \in \text{Car}(X) \otimes_{\mathbb{Z}} \mathbb{Z}_{(p)}$. For each $e \in d\mathbb{Z}_{>0}$ we define

$$\begin{aligned} \mathcal{L}_{(X/Y, \Delta)}^{(e)} &:= \mathcal{O}_{X^e}((1-p^e)(K_{X^e/Y^e} + \Delta^e)) \subseteq \mathcal{O}_{X^e}((1-p^e)K_{X^e/Y^e}), \\ \phi_{(X/Y, \Delta)}^{(d)} &: F_{X/Y}^{(d)} \mathcal{L}_{(X/Y, \Delta)}^{(d)} \rightarrow F_{X/Y}^{(d)} \mathcal{O}_{X^d}((1-p^d)K_{X^d/Y^d}) \xrightarrow{\phi_{X/Y}^{(d)}} \mathcal{O}_{X_{Y^d}}, \quad \text{and} \\ \phi_{(X/Y, \Delta)}^{(e+d)} &:= (\phi_{(X/Y, \Delta)}^{(e)})_{Y^e} \circ F_{X_{Y^d}/Y^d}^{(e)} (\phi_{(X/Y, \Delta)}^{(d)} \otimes (\mathcal{L}_{(X/Y, \Delta)}^{(e)})_{Y^{e+d}}) : F_{X/Y}^{(e+d)} \mathcal{L}_{(X/Y, \Delta)}^{(e+d)} \rightarrow \mathcal{O}_{X_{Y^{e+d}}}. \end{aligned}$$

In order to generalize the definitions above, we recall the notion of generalized divisors on a k -scheme. Let X be a k -scheme of pure dimension satisfying S_2 and G_1 . An *AC divisor* (or *almost Cartier divisor*) on

X is a reflexive coherent subsheaf of the sheaf of total quotient rings on X that is invertible in codimension one (see [Hartshorne 1994; Miller and Schwede 2012]). For an AC divisor D , the coherent sheaf defining D is denoted by $\mathcal{O}_X(D)$. The set of AC divisors $\text{WSh}(X)$ has a structure of additive group [Hartshorne 1994, Corollary 2.6]. A $\mathbb{Z}_{(p)}$ -AC divisor (resp. \mathbb{Q} -AC divisor) is an element of $\text{WSh}(X) \otimes_{\mathbb{Z}} \mathbb{Z}_{(p)}$ (resp. $\text{WSh}(X) \otimes_{\mathbb{Z}} \mathbb{Q}$). An AC divisor D is said to be *effective* if $\mathcal{O}_X \subseteq \mathcal{O}_X(D)$, and a $\mathbb{Z}_{(p)}$ -AC (resp. \mathbb{Q} -AC) divisor Δ is said to be *effective* if $\Delta = D \otimes r$ for an effective AC divisor D and an $r \geq 0$. For two AC divisors D and E , the notation $D \leq E$ means that $E - D$ is effective. We use the same notation for $\mathbb{Z}_{(p)}$ -AC (resp. \mathbb{Q} -AC) divisors.

Remark 2.1. Each of the natural morphisms

$$\text{WSh}(X) \xrightarrow{(\cdot) \otimes 1} \text{WSh}(X) \otimes_{\mathbb{Z}} \mathbb{Z}_{(p)} \rightarrow \text{WSh}(X) \otimes_{\mathbb{Z}} \mathbb{Q}$$

is not necessarily injective (see Example 2.2). Let D and E be AC divisors. Then, D and E are equal as $\mathbb{Z}_{(p)}$ -AC (resp. \mathbb{Q} -AC) divisors if and only if $mD = mE$ for some nonzero $m \in \mathbb{Z} \setminus p\mathbb{Z}$ (resp. $m \in \mathbb{Z}$). Furthermore, D and E can be equal as $\mathbb{Z}_{(p)}$ -AC (resp. \mathbb{Q} -AC) divisors even when D is effective but E is not (see Example 2.2).

Example 2.2 [Corti 1992, (16.1.2)]. Set $X := \text{Spec } k[x, y, z, z^{-1}]/(x^n - zy^n)$ for an $n \geq 2$. Note that X is integral and Gorenstein but not normal. Let D and E be AC divisors on X defined by $x^{-1}\mathcal{O}_X$ and $y^{-1}\mathcal{O}_X$, respectively. For an $m \geq 1$, one has

$$mD = mE \iff x^{-m}\mathcal{O}_X = y^{-m}\mathcal{O}_X \iff n \mid m.$$

Hence, we see that

- $D \neq E$ as AC divisors,
- $D \otimes 1 = E \otimes 1$ as $\mathbb{Z}_{(p)}$ -AC divisors if and only if $p \nmid n$, and
- $D \otimes 1 = E \otimes 1$ as \mathbb{Q} -AC divisors.

Furthermore, $D - E$ is not effective but $n(D - E) = 0$ is effective.

Remark 2.3. Let E and K be two AC divisors, take an $\varepsilon \in \mathbb{Z}_{(p)}$ (resp. $\varepsilon \in \mathbb{Q}$) and set $\Delta := E \otimes \varepsilon$. When we consider the $\mathbb{Z}_{(p)}$ -AC (resp. \mathbb{Q} -AC) divisor $K + \Delta$, for each $m \in \mathbb{Z}$ with $\varepsilon m \in \mathbb{Z}$, we let $m(K + \Delta)$ denote the AC divisor $mK + (\varepsilon m)E$.

Let $f : X \rightarrow Y$ be a morphism from a pure dimensional k -scheme X to a regular variety Y and assume that X satisfies S_2 and G_1 . Let E be an effective AC divisor on X , and fix a Gorenstein open subset $U \subseteq X$ such that $\text{codim}_X(X \setminus U) \geq 2$ and $E|_U$ is Cartier. Let $U \xhookrightarrow{\iota} X$ denote the open immersion. Then,

for each $e \in \mathbb{Z}_{>0}$, we have the following commutative diagram:

$$\begin{array}{ccccc}
 U^e & \xrightarrow{F_{U/Y}^{(e)}} & U_{Y^e} & \longrightarrow & U \\
 \downarrow \iota^{(e)} & & \downarrow \iota_{Y^e} & & \downarrow \iota \\
 X^e & \xrightarrow{F_{X/Y}^{(e)}} & X_{Y^e} & \longrightarrow & X \\
 & \searrow f^{(e)} & \downarrow f_{Y^e} & & \downarrow f \\
 & & Y^e & \xrightarrow{F_Y^e} & Y
 \end{array}$$

Take an integer $a > 0$ not divisible by p , set $\Delta := E \otimes \frac{1}{a}$, and let d be the minimum positive integer satisfying $a \mid (p^d - 1)$. For each $e \in d\mathbb{Z}_{>0}$, we define

$$\mathcal{L}_{(X/Y, \Delta)}^{(e)} := \iota^{(e)*} \mathcal{L}_{(U/Y, \Delta|_U)}^{(e)} \quad \text{and} \quad \phi_{(X/Y, \Delta)}^{(e)} := \iota_{Y^e*}(\phi_{(U/Y, \Delta|_U)}^{(e)}) : F_{X/Y}^{(e)} \mathcal{L}_{(X/Y, \Delta)}^{(e)} \rightarrow \mathcal{O}_{X_{Y^e}}.$$

Fix $e \in d\mathbb{Z}_{>0}$. Set $\mathcal{L}_{(X, \Delta)}^{(e)} := \mathcal{O}_{X^e}((1 - p^e)(K_{X^e} + \Delta^e))$. We can define the morphism $\phi_{(X, \Delta)}^{(e)} : F_{X*}^e \mathcal{L}_{(X, \Delta)}^{(e)} \rightarrow \mathcal{O}_X$ by a procedure similar to the one above, replacing $F_{X/Y}^{(e)}$ by F_X^e . In the case where k is perfect and $Y = \text{Spec } k$, one may identify, respectively, F_X^e , $\mathcal{L}_{(X, \Delta)}^{(e)}$ and $\phi_{(X, \Delta)}^{(e)}$ with $F_{X/Y}^{(e)}$, $\mathcal{L}_{(X/Y, \Delta)}^{(e)}$ and $\phi_{(X/Y, \Delta)}^{(e)}$.

We next introduce singularities of pairs defined by the Grothendieck trace map of the Frobenius morphism.

Definition 2.4. Let X be a k -scheme of pure dimension satisfying S_2 and G_1 , and let Δ be an effective \mathbb{Q} -AC divisor on X :

- (i) We say that (X, Δ) is *F-pure* if for each $e \in \mathbb{Z}_{>0}$ and every effective AC divisor E' with $\Delta' := E' \otimes 1/(p^e - 1) \leq \Delta$, the morphism

$$\phi_{(X, \Delta')}^{(e)} : F_{X*}^e \mathcal{O}_X((1 - p^e)(K_X + \Delta')) \rightarrow \mathcal{O}_X$$

is surjective.

- (ii) [Schwede 2008, Definition 3.1] Assume that X is a normal variety. We say that (X, Δ) is *strongly F-regular* if every effective Cartier divisor D on X , the morphism

$$F_{X*}^e \mathcal{O}_X(\lfloor (1 - p^e)(K_X + \Delta) - D \rfloor) = F_{X*}^e \mathcal{O}_X((1 - p^e)(K_X + \Delta')) \xrightarrow{\phi_{(X, \Delta')}^{(e)}} \mathcal{O}_X$$

is surjective for some $e \in \mathbb{Z}_{>0}$, where $\Delta' := \frac{1}{p^e - 1} \lfloor (p^e - 1)\Delta + D \rfloor$. Here, $\lfloor \cdot \rfloor$ denotes the round down.

We simply say that X is *F-pure* (resp. *strongly F-regular*) if $(X, 0)$ is *F-pure* (resp. *strongly F-regular*).

Remarks 2.5. (1) With the notation as above, assume that X is normal and affine. Then the above definition of *F-purity* is equivalent to that in [Hara and Watanabe 2002]. This follows from the fact that $\lfloor (p^e - 1)\Delta \rfloor$ is the greatest element of the set S of all divisors E' such that $E' \leq (p^e - 1)\Delta$.

- (2) When X is not normal, S in (1) does not necessarily have a greatest element. Let X , D and E be as in Example 2.2 with $p \nmid n$ and $n \nmid l$, where $l := p^e - 1$. Then, $\Delta := D \otimes 1 = E \otimes 1$ as $\mathbb{Z}_{(p)}$ -divisors and $lD \neq lE$. If S has a greatest element G , then $G \geq lD$ and $G \otimes 1 = l\Delta$, from which one can get $G = lD$. In the same way, we get $G = lE$, so $lD = lE$, a contradiction.
- (3) Let (X, Δ) be a strongly F -regular pair, and Δ' an effective \mathbb{Q} -divisor on X . Then there is $\varepsilon \in \mathbb{Q}_{>0}$ such that $(X, \Delta + \varepsilon\Delta')$ is again strongly F -regular.

2B. Weak positivity. In this subsection, we recall the notion of weak positivity introduced by Viehweg [1983]. The definition employed in this paper is slightly different from Viehweg's original one. We work over a field k in this subsection.

Definition 2.6. Let Y be a quasiprojective normal variety, let \mathcal{G} and \mathcal{G}' be coherent sheaves on Y , and let \mathcal{H} be an ample line bundle on Y . Take a subset S of the underlying topological space of Y such that the stalk of \mathcal{G} at any point in S is free, i.e., there is an open subset $Y_0 \subset Y$ such that $S \subseteq Y_0$ and $\mathcal{G}|_{Y_0}$ is locally free:

- (i) We say that a morphism $\mathcal{G} \rightarrow \mathcal{G}'$ is *surjective over S* if S and the support of the cokernel do not intersect.
- (ii) We say that \mathcal{G} is *globally generated over S* if the natural morphism $H^0(Y, \mathcal{G}) \otimes_k \mathcal{O}_Y \rightarrow \mathcal{G}$ is surjective over S .
- (iii) We say that \mathcal{G} is *weakly positive over S* if for every $\alpha \in \mathbb{Z}_{>0}$, there is $\beta \in \mathbb{Z}_{>0}$ such that $(S^{\alpha\beta} \mathcal{G})^{**} \otimes \mathcal{H}^\beta$ is globally generated over S . Here, $S^{\alpha\beta}(\)$ and $(\)^{**}$ denote the $\alpha\beta$ -th symmetric product and the double dual, respectively.
- (iv) We say that \mathcal{G} is *big over S* if there is $\gamma \in \mathbb{Z}_{>0}$ such that $(S^\gamma \mathcal{G}) \otimes \mathcal{H}^{-1}$ is weakly positive over S .

We simply say that \mathcal{G} is *generically globally generated* if \mathcal{G} is globally generated over $\{\eta\}$, where η is the generic point of Y . Furthermore, we simply say that \mathcal{G} is *weakly positive* (resp. *big*) if it is weakly positive (resp. big) over $\{\eta\}$.

Remark 2.7. Let Y , \mathcal{G} , S and \mathcal{H} be as above:

- (1) The above definitions are independent of the choice of \mathcal{H} [Viehweg 1995, Lemma 2.14].
- (2) Suppose that \mathcal{G} is a vector bundle over a smooth projective curve Y . Then \mathcal{G} is weakly positive (resp. big) over Y if and only if \mathcal{G} is nef (resp. ample).

2C. Augmented and restricted base loci. In this subsection, we recall the definition of the augmented and restricted base locus of a \mathbb{Q} -Cartier divisor. In this subsection, we work over a field k .

Definition 2.8 [Ein et al. 2006; Mustařa 2013]. Let Y be a quasiprojective variety and D a \mathbb{Q} -Cartier divisor on Y :

- (i) The *stable base locus* $\mathbf{B}(D)$ of D is defined as the reduced base locus of mD for sufficiently large and divisible integer m .

(ii) The *augmented base locus* is given by

$$\mathbf{B}_+(D) := \bigcap_A \mathbf{B}(D - A),$$

where A runs over all the ample \mathbb{Q} -Cartier divisors on Y .

(iii) The *restricted base locus* (also called the *nonnef locus* or the *diminished base locus*) is defined by

$$\mathbf{B}_-(D) := \bigcup_A \mathbf{B}(D + A),$$

where A runs over all the ample \mathbb{Q} -Cartier divisors on Y .

Remarks 2.9. (1) In [Ein et al. 2006], the variety Y is assumed to be projective.

(2) Assume that Y is projective. Then the following hold:

- $\mathbf{B}_+(D) = \emptyset$ if and only if D is ample [loc. cit., Example 1.7].
- $\mathbf{B}_+(D) \neq Y$ if and only if D is big [loc. cit., Example 1.7].
- $\mathbf{B}_-(D) = \emptyset$ if and only if D is nef [loc. cit., Example 1.18].
- When k is uncountable, $\mathbf{B}_-(D) \neq Y$ if and only if D is pseudoeffective [Mustață 2013, Section 2].

(3) Assume that Y is a normal projective variety and D is Cartier. Let S be a subset of the underlying topological space of Y . Then, the weak positivity (resp. bigness) of $\mathcal{O}_Y(D)$ is equivalent to saying that $S \cap \mathbf{B}_-(D) = \emptyset$ (resp. $S \cap \mathbf{B}_+(D) = \emptyset$).

The next lemma can be proved in the same way as in the proof of [Ein et al. 2006, Proposition 1.19].

Lemma 2.10 [Ein et al. 2006, Propositions 1.19]. *Let the notation be as in Definition 2.8. Let H be an ample \mathbb{Q} -Cartier divisor on Y and $\{a_m\}$ a sequence of positive rational numbers that converges to zero. Then $\mathbf{B}_-(D) = \bigcup_m \mathbf{B}(D + a_m H)$.*

3. Auxiliary lemmas

In this section, we prove several lemmas used in the proofs of the main theorems. Throughout this section, the base field k is assumed to be an F -finite field of characteristic $p > 0$.

Lemma 3.1. *Let W be a normal quasiprojective variety and $W_0 \subseteq W$ an open subset. Let \mathcal{H} be an ample line bundle on W and \mathcal{G} a coherent sheaf on W such that $\mathcal{G}|_{W_0}$ is locally free. Then there exists $m_0 \in \mathbb{Z}_{>0}$ such that, for every $m \geq m_0$, there exists a homomorphism $\theta : \bigoplus^n \mathcal{G} \rightarrow \mathcal{H}^m$ of \mathcal{O}_W -modules which is surjective over W_0 .*

Proof. Take $m \in \mathbb{Z}_{>0}$ such that $\mathcal{G}^* \otimes \mathcal{H}^m$ is generated by its global sections. Since W_0 is Noetherian, we get a surjective morphism $\theta' : \bigoplus^n \mathcal{O}_W \twoheadrightarrow \mathcal{G}^* \otimes \mathcal{H}^m$ for some $n \in \mathbb{Z}_{>0}$. We then obtain

$$\bigoplus^n \mathcal{G} \cong \left(\bigoplus^n \mathcal{O}_W \right) \otimes \mathcal{G} \xrightarrow{\theta' \otimes \mathcal{G}} (\mathcal{G}^* \otimes \mathcal{H}^m) \otimes \mathcal{G} \cong \mathcal{H}om(\mathcal{G}, \mathcal{H}^m) \otimes \mathcal{G} \rightarrow \mathcal{H}^m.$$

Here, the last morphism is the evaluation morphism, which is surjective over W_0 , since $\mathcal{G}|_{W_0}$ is locally free. Hence, the composite of the above morphisms is the desired morphism. \square

Lemma 3.2. *Let W be a normal quasiprojective variety and D a Cartier divisor on W . Let $W_0 \subseteq W$ be an open subset, \mathcal{E} a coherent sheaf that is globally generated over W_0 , and \mathcal{G} a coherent sheaf on W such that $\mathcal{G}|_{W_0}$ is free. Suppose that there exists a morphism*

$$\varphi : \mathcal{E} \otimes \left(\bigotimes^{p^e} \mathcal{G} \right) \rightarrow \mathcal{O}_W(D) \otimes (F_W^e)^* \mathcal{G}$$

that is surjective over W_0 . Then $W_0 \cap \mathbf{B}_-(D) = \emptyset$.

Proof. Obviously, we may assume that $\mathcal{E} = \bigoplus^d \mathcal{O}_W$ for some $d \in \mathbb{Z}_{>0}$. Then $\mathcal{E} \otimes (\bigotimes^{p^e} \mathcal{G}) \cong \bigoplus^d (\bigotimes^{p^e} \mathcal{G})$. Take an ample line bundle \mathcal{H} on W such that $\mathcal{G} \otimes \mathcal{H}$ is globally generated. Since we have $(\bigotimes^{p^e} \mathcal{G}) \otimes \mathcal{H}^{p^e} \cong \bigotimes^{p^e} (\mathcal{G} \otimes \mathcal{H})$ and $((F_Y^e)^* \mathcal{G}) \otimes \mathcal{H}^{p^e} \cong (F_Y^e)^* (\mathcal{G} \otimes \mathcal{H})$, replacing \mathcal{G} (resp. φ) by $\mathcal{G} \otimes \mathcal{H}$ (resp. $\varphi \otimes \mathcal{H}^{p^e}$), we may assume that \mathcal{G} is globally generated. Let $S(\mathcal{G}) \subseteq \mathbb{Q}$ be the set of rational numbers r satisfying the following condition: there is $h \in \mathbb{Z}_{>0}$ such that $p^h r \in \mathbb{Z}$ and that the sheaf

$$\mathcal{O}_W(p^h r D) \otimes (F_W^h)^* \mathcal{G}$$

is globally generated over W_0 . We then have $0 \in S(\mathcal{G})$. We prove that $S(\mathcal{G})$ is *not* bounded from above. Choose $r \in S$ and $h \in \mathbb{Z}_{>0}$ so that the above conditions hold. We then have the following sequence of morphisms:

$$\begin{aligned} \bigoplus^d \left(\bigotimes^{p^e} (\mathcal{O}_W(p^h r D) \otimes (F_W^h)^* \mathcal{G}) \right) &\cong \mathcal{O}_W(p^{e+h} r D) \otimes (F_W^h)^* \left(\bigoplus^d \left(\bigotimes^{p^e} \mathcal{G} \right) \right) \\ &\xrightarrow{\psi} \mathcal{O}_W(p^{e+h} r D) \otimes (F_W^h)^* (\mathcal{O}_W(D) \otimes (F_W^e)^* \mathcal{G}) \\ &\cong \mathcal{O}_W((p^h + p^{e+h} r) D) \otimes (F_W^{e+h})^* \mathcal{G} \end{aligned}$$

Here, $\psi := ((F_W^h)^* \varphi) \otimes \mathcal{O}_W(p^h r D)$, so it is surjective over W_0 , which implies that the last sheaf is globally generated over W_0 . We then see that $1/p^e + r = (p^h + p^{e+h} r)/p^{e+h} \in S(\mathcal{G})$, and hence $S(\mathcal{G})$ can not be bounded from above. Next, we show the assertion. Lemma 3.1 shows that we have an ample Cartier divisor H on W and a morphism $\theta : \bigoplus^n \mathcal{G} \rightarrow \mathcal{H} := \mathcal{O}_W(H)$ that is surjective over W_0 . One can easily check that $S(\mathcal{G}) \subseteq S(\mathcal{H})$, so $S(\mathcal{H})$ is also not bounded from above. Take $0 < r \in S(\mathcal{H})$. Then for some $h \gg 0$, the sheaf $\mathcal{O}_W(p^h r D) \otimes (F_W^h)^* \mathcal{H} \cong \mathcal{O}_W(p^h r (D + \frac{1}{r} H))$ is globally generated over W_0 , and so $\mathbf{B}(D + \frac{1}{r} H) \subseteq W \setminus W_0$. Hence, we conclude from Lemma 2.10 that $\mathbf{B}_-(D) \subseteq W \setminus W_0$. \square

Before stating the next lemma, we recall Keeler's vanishing theorem, which is a relative version of Fujita's vanishing theorem.

Theorem 3.3 [Keeler 2003, Theorem 1.5]. *Let $f : X \rightarrow Y$ be a projective morphism between Noetherian schemes, \mathcal{F} a coherent sheaf on X , and \mathcal{L} an f -ample line bundle on X . Then there exists $m_0 \in \mathbb{Z}_{>0}$ such that*

$$R^i f_*(\mathcal{F} \otimes \mathcal{L}^m \otimes \mathcal{N}) = 0$$

for each $m \geq m_0$ and every f -nef line bundle \mathcal{N} on X .

Note that, in the situation of the theorem, a line bundle on X is said to be f -nef if the restriction to each fiber is nef.

Lemma 3.4. *Let $f : X \rightarrow Y$ be a surjective morphism between projective varieties, and \mathcal{A} an ample line bundle on X . Then there exists $m_0 \in \mathbb{Z}_{>0}$ such that $f_*(\mathcal{F} \otimes \mathcal{A}^m \otimes \mathcal{N})$ is generated by its global sections for each $m \geq m_0$ and every nef line bundle \mathcal{N} on X .*

Proof. Let \mathcal{H} be an ample line bundle on Y that is generated by its global sections. Set $n := \dim Y$. Take $m_1 > 0$ so that $\mathcal{A}^{m_1} \otimes f^*\mathcal{H}^{-n}$ is nef. Applying Theorem 3.3, we get $m_2 > 0$ such that $H^i(X, \mathcal{F} \otimes \mathcal{A}^m \otimes \mathcal{N}) = 0$ and $R^i f_*(\mathcal{F} \otimes \mathcal{A}^m \otimes \mathcal{N}) = 0$ for each $m \geq m_2$, $i > 0$ and every nef line bundle \mathcal{N} on X . The Leray spectral sequence then implies that $H^i(Y, f_*(\mathcal{F} \otimes \mathcal{A}^m \otimes \mathcal{N})) = 0$ for each $i > 0$. Fix $m \geq m_0 := m_1 + m_2$ and a nef line bundle \mathcal{N} on X , and set $\mathcal{M} := \mathcal{A}^m \otimes \mathcal{N}$. We then have $\mathcal{F} \otimes \mathcal{M} \otimes f^*\mathcal{H}^{-i} \cong \mathcal{F} \otimes \mathcal{A}^{m-m_1} \otimes (\mathcal{A}^{m_1} \otimes f^*\mathcal{H}^{-n}) \otimes f^*\mathcal{H}^{n-i}$, so the above argument tells us that the i -th cohomology of $f_*(\mathcal{F} \otimes \mathcal{M} \otimes f^*\mathcal{H}^{-i}) \cong (f_*(\mathcal{F} \otimes \mathcal{M})) \otimes \mathcal{H}^{-i}$ vanishes for $0 < i \leq n$. This means that $f_*(\mathcal{F} \otimes \mathcal{M})$ is 0-regular with respect to \mathcal{H} , so this sheaf is generated by its global sections thanks to the Castelnuovo–Mumford regularity (see [Lazarsfeld 2004, Theorem 1.8.5]). \square

Lemma 3.5. *Let $g : V \rightarrow W$ be a surjective projective morphism from a k -scheme V to a variety W , let \mathcal{A} be a g -ample line bundle on V , and let \mathcal{F} be a coherent sheaf on V that is flat over W . Then, there exists $m_0 \in \mathbb{Z}_{>0}$ such that $g_*(\mathcal{F} \otimes \mathcal{A}^m \otimes \mathcal{N})$ is locally free for each $m \geq m_0$ and every g -nef line bundle \mathcal{N} on V .*

Proof. By Theorem 3.3, there is $m_0 \in \mathbb{Z}_{>0}$ such that $R^i g_*(\mathcal{F} \otimes \mathcal{A}^m \otimes \mathcal{N}) = 0$ for each $m \geq m_0$, $i > 0$ and every g -nef line bundle \mathcal{N} on V . Fix $m \geq m_0$ and a g -nef line bundle \mathcal{N} on V , and set $\mathcal{M} := \mathcal{A}^m \otimes \mathcal{N}$. For each $i \geq 0$, define the function $h^i : W \rightarrow \mathbb{Z}$ by $h^i(w) := \dim_{k(w)} H^i(V_w, (\mathcal{F} \otimes \mathcal{M})|_{V_w})$. By the choice of m_0 and cohomology and base change (see [Hartshorne 1977, Theorem III 12.11]), we obtain that $h^i = 0$ for each $i > 0$, so $\chi((\mathcal{F} \otimes \mathcal{M})|_{V_w}) = h^0(w)$ for every $w \in W$. Then [Hartshorne 1977, Theorem III 9.9 and its proof] implies that h^0 is constant. Hence, our claim follows from Grauert’s theorem (see [Hartshorne 1977, Corollary III 12.9]). \square

Lemma 3.6. *Let the notation be as in Lemma 3.5. Let \mathcal{L} be a line bundle on V :*

- (1) *If \mathcal{L} is g -free, then there exists $n_0 \in \mathbb{Z}_{>0}$ such that the natural morphism*

$$g_*\mathcal{L}^m \otimes g_*(\mathcal{F} \otimes \mathcal{L}^n \otimes g^*\mathcal{P}) \rightarrow g_*(\mathcal{F} \otimes \mathcal{L}^{m+n} \otimes g^*\mathcal{P})$$

is surjective for each $n \geq n_0$, $m > 0$ and every line bundle \mathcal{P} on W .

- (2) *If \mathcal{L} is g -ample and g -free, then there exists $n_0 \in \mathbb{Z}_{>0}$ such that the natural morphism*

$$g_*\mathcal{L}^m \otimes g_*(\mathcal{F} \otimes \mathcal{L}^n \otimes \mathcal{N}) \rightarrow g_*(\mathcal{F} \otimes \mathcal{L}^{m+n} \otimes \mathcal{N})$$

is surjective for each $n \geq n_0$, $m > 0$ and every g -nef line bundle \mathcal{N} on V .

Proof. We first show that (2) implies (1). Since \mathcal{L} is g -free, g can be decomposed as

$$g : V \xrightarrow{\sigma} Z \xrightarrow{\tau} W$$

and $\mathcal{L} \cong \sigma^* \mathcal{M}$ for a τ -ample and τ -free line bundle \mathcal{M} on Z . Then we have

$$\begin{array}{ccc}
 (\tau_* \mathcal{M}^m) \otimes \tau_*((\sigma_* \mathcal{F}) \otimes \mathcal{M}^n \otimes \tau^* \mathcal{P}) & \longrightarrow & \tau_*((\sigma_* \mathcal{F}) \otimes \mathcal{M}^{m+n} \otimes \tau^* \mathcal{P}) \\
 \downarrow & & \downarrow \cong \\
 (\tau_* \sigma_* \sigma^* \mathcal{M}^m) \otimes \tau_*((\sigma_* \mathcal{F}) \otimes \mathcal{M}^n \otimes \tau^* \mathcal{P}) & & \\
 \downarrow \cong & & \\
 \tau_* \sigma_* \mathcal{L}^m \otimes \tau_* \sigma_* (\mathcal{F} \otimes \mathcal{L}^n \otimes \sigma^* \tau^* \mathcal{P}) & \longrightarrow & \tau_* \sigma_* (\mathcal{F} \otimes \mathcal{L}^{m+n} \otimes \sigma^* \tau^* \mathcal{P})
 \end{array}$$

Here, the isomorphisms follow from the projection formula. If (2) holds, then the top horizontal arrow is surjective, and hence so is the bottom horizontal arrow, so (1) holds. We show (2). Theorem 3.3 tells us that we have $n_0 \in \mathbb{Z}_{>0}$ such that for each $n \geq n_0$ and every g -nef line bundle \mathcal{N} on V , the sheaf $\mathcal{F} \otimes \mathcal{L}^n \otimes \mathcal{N}$ is 0-regular with respect to \mathcal{L} and g . Hence, in the case when $m = 1$, our claim follows from the relative Castelnuovo–Mumford regularity (see [Lazarsfeld 2004, Example 1.8.24]). Using this repeatedly, we see that the natural morphism

$$\left(\bigotimes_{i=1}^m g_* \mathcal{L} \right) \otimes g_*(\mathcal{F} \otimes \mathcal{L}^n \otimes \mathcal{N}) \rightarrow g_*(\mathcal{F} \otimes \mathcal{L}^{m+n} \otimes \mathcal{N})$$

is surjective for each $m \in \mathbb{Z}_{>0}$. This morphism factors through

$$(g_* \mathcal{L}^m) \otimes g_*(\mathcal{F} \otimes \mathcal{L}^n \otimes \mathcal{N}) \rightarrow g_*(\mathcal{F} \otimes \mathcal{L}^{m+n} \otimes \mathcal{N}),$$

which completes the proof. \square

Lemma 3.7. *Let V be a k -scheme of pure dimension satisfying S_2 and G_1 , let W be a regular variety, and let $g : V \rightarrow W$ be a flat projective morphism. Let $E \geq 0$ be an AC divisor on V such that $aK_V + E$ is Cartier for some $a \in \mathbb{Z}_{>0}$ with $p \nmid a$. Set $\Delta := E \otimes \frac{1}{a}$. Let $U \subseteq V$ be a Gorenstein open subset. Suppose that the codimension of $(V \setminus U)|_{V_w}$ (resp. $E|_{V_w}$) in V_w is at least 2 (resp. 1) for every $w \in W$:*

- (1) [Patakfalvi et al. 2018, Corollary 3.31] *The set $W_0 := \{w \in W \mid (V_w, \overline{\Delta|_{U_w}}) \text{ is } F\text{-pure}\}$ is an open subset of W . Here, V_w is the geometric fiber over w and $\overline{\Delta|_{U_w}}$ is the $\mathbb{Z}_{(p)}$ -AC divisor on V_w obtained as the unique extension of $\Delta|_{U_w}$.*
- (2) *Assume that W_0 is nonempty. Set $V_0 := g^{-1}(W_0)$. Let \mathcal{A} be a line bundle on V such that $\mathcal{A}|_{V_0}$ is $g|_{V_0}$ -ample. Then there exists $m_0 \in \mathbb{Z}_{>0}$ such that*

$$g_{W^e *} (\phi_{(V/W, \Delta)}^{(e)} \otimes \mathcal{A}_{W^e}^m \otimes \mathcal{N}_{W^e}) : g^{(e)}_* (\mathcal{L}_{(V/W, \Delta)}^{(e)} \otimes \mathcal{A}^{p^e m} \otimes \mathcal{N}^{p^e}) \rightarrow g_{W^e *} (\mathcal{A}_{W^e}^m \otimes \mathcal{N}_{W^e})$$

is surjective over W_0 for each $e \in \mathbb{Z}_{>0}$ with $a \mid (p^e - 1)$, $m \geq m_0$ and every line bundle \mathcal{N} on V whose restriction $\mathcal{N}|_{V_0}$ to V_0 is $g|_{V_0}$ -nef

Proof. One can prove (1) by the same argument as that in [Patakfalvi et al. 2018]. We prove (2). Let $d > 0$ be the minimum integer such that $a \mid (p^d - 1)$. For simplicity, let $\phi^{(e)}$ (resp. $\mathcal{L}^{(e)}$) denote $\phi_{(V/W, \Delta)}^{(e)}$ (resp. $\mathcal{L}_{(V/W, \Delta)}^{(e)}$) for each $e \in d\mathbb{Z}_{>0}$. Replacing W by W_0 , we may assume that $W_0 = W$. The morphism $\phi^{(e)}|_{V_w} \cong \phi_{(V_w/w, \Delta_w)}^{(e)}$ is then surjective for every $w \in W$ and $e \in d\mathbb{Z}_{>0}$, so $\phi^{(e)}$ is surjective. Applying

Theorem 3.3 to the kernel of $\phi^{(d)}$, we obtain $m'_0 \in \mathbb{Z}_{>0}$ such that $g_{W^d*}(\phi^{(d)} \otimes \mathcal{A}_{W^d}^m \otimes \mathcal{N}_{W^d})$ is surjective for every $m \geq m'_0$ and g -nef line bundle \mathcal{N} on V . Take $m_0 \geq m'_0$ so that $m_0 A - (K_{V/W} + \Delta)$ is g -nef, where A is a Cartier divisor on V satisfying $\mathcal{O}_V(A) \cong \mathcal{A}$. We fix $m \geq m_0$ and a g -nef line bundle \mathcal{N} on V . Set $\mathcal{M} := \mathcal{A}^m \otimes \mathcal{N}$. We show that $\psi^{(e)} := g_{W^e*}(\phi^{(e)} \otimes \mathcal{M}_{W^e})$ is surjective for each $e \in d\mathbb{Z}_{>0}$. We have already seen that $\psi^{(d)}$ is surjective. Take $e \in d\mathbb{Z}_{>0}$. Assuming the surjectivity of $\psi^{(e)}$, we show that $\psi^{(e+d)}$ is surjective. By the definition of $\phi^{(e+d)}$, we have

$$\begin{aligned} \psi^{(e+d)} &= g_{W^{e+d}*}(\phi^{(e+d)} \otimes \mathcal{M}_{W^{e+d}}) \\ &\cong (F_W^d)^*(g_{W^e*}(\phi^{(e)} \otimes \mathcal{M}_{W^e})) \circ g_{W^{e+d}*}^{(e)}(\phi_{(V^e/W^e, \Delta^e)}^{(d)} \otimes (\mathcal{L}^{(e)} \otimes \mathcal{M}^{p^e})_{W^{e+d}}) \\ &= (F_W^d)^*(\psi^{(e)}) \circ g_{W^{e+d}*}^{(e)}(\phi_{(V^e/W^e, \Delta^e)}^{(d)} \otimes (\mathcal{L}^{(e)} \otimes \mathcal{M}^{p^e})_{W^{e+d}}). \end{aligned}$$

Note that the surjectivity of $\psi^{(e)}$ induces that of $(F_W^d)^*(\psi^{(e)})$, so we only need to show that

$$g_{W^{e+d}*}^{(e)}(\phi_{(V^e/W^e, \Delta^e)}^{(d)} \otimes (\mathcal{L}^{(e)} \otimes \mathcal{M}^{p^e})_{W^{e+d}})$$

is surjective. We can rewrite this morphism as

$$g_{W^d*}(\phi^{(d)} \otimes (\mathcal{O}_V((1-p^e)(K_{V/W} + \Delta)) \otimes \mathcal{M}^{p^e})_{W^d})$$

identifying $g^{(e)} : V^e \rightarrow W^e$ with $g : V \rightarrow W$. This morphism is surjective if

$$\mathcal{P} := \mathcal{O}_V((1-p^e)(K_{V/W} + \Delta)) \otimes \mathcal{M}^{p^e} \otimes \mathcal{A}^{-m_0}$$

is g -nef, by the choice of m'_0 . This g -nefness follows from the isomorphisms

$$\mathcal{P} \cong \mathcal{O}_V((1-p^e)(K_{V/W} + \Delta)) \otimes \mathcal{A}^{mp^e - m_0} \otimes \mathcal{N}^{p^e} \cong \mathcal{O}_V((p^e - 1)(m_0 A - (K_{V/W} + \Delta))) \otimes \mathcal{A}^{(m-m_0)p^e} \otimes \mathcal{N}^{p^e}$$

and the choice of m_0 . □

4. Main theorems and corollaries

In this section, we prove the main theorems. After this, we give several corollaries.

4A. Main theorems. In this subsection, we prove Theorems 4.1 and 4.2. *Throughout this subsection, we use the following notation:*

Fix an F -finite field k . Let $f : X \rightarrow Y$ be a surjective projective morphism from a pure dimensional quasiprojective k -scheme X satisfying S_2 and G_1 to a normal quasiprojective variety Y . Let E be an effective AC-divisor on X and $a > 0$ an integer such that $aK_X + E$ is Cartier. Set $\Delta := E \otimes a^{-1}$. Let $U \subset X$ be the Gorenstein locus and $W \subseteq Y$ the maximal regular open subset such that $g := f|_V : V \rightarrow W$ is flat, where $V := f^{-1}(W)$. Suppose that there exists a scheme-theoretic point $w \in W$ with the following properties:

- (i) $\text{codim}_{X_w}(X_w \setminus U_w) \geq 2$.
- (ii) $\text{Supp } E$ does not contain any irreducible component of X_w .
- (iii) $(X_{\bar{w}}, \overline{\Delta|_{U_{\bar{w}}}})$ is F -pure.

Here, $X_{\bar{w}}$ (resp. $U_{\bar{w}}$) is the geometric fiber of f (resp. $f|_U$) over w , and $\overline{\Delta|_{U_{\bar{w}}}}$ is the $\mathbb{Z}_{(p)}$ -AC divisor on $X_{\bar{w}}$ obtained as the unique extension of $\Delta|_{U_{\bar{w}}}$. Let D be a \mathbb{Q} -Cartier divisor on Y .

We now have the following commutative diagram whose squares are cartesian:

$$\begin{array}{ccccc}
 U_{\bar{w}} & \xrightarrow{\quad\quad\quad} & & \xrightarrow{\quad\quad\quad} & U & \text{(Gorenstein)} \\
 \downarrow \scriptstyle (\text{codim} \geq 2) & & & & \downarrow \scriptstyle (\text{codim} \geq 2) & \\
 X_{\bar{w}} & \xrightarrow{\quad\quad\quad} & V & \hookrightarrow & X & \\
 \downarrow \scriptstyle f_{\bar{w}} & & \downarrow \scriptstyle \text{(flat)} \quad g & & \downarrow \scriptstyle f & \\
 \bar{w} := \operatorname{Spec} \bar{k}(w) & \xrightarrow{\quad\quad\quad} & W & \hookrightarrow & Y & \\
 & & \text{(regular)} & & &
 \end{array}$$

Here, “(codim ≥ 2)” means the morphism is an open immersion whose complement is of codimension at least 2.

In this situation, we prove the following two theorems:

Theorem 4.1. *Let the notation be as above. Assume that X is projective and $K_X + \Delta$ is \mathbb{Q} -Cartier:*

- (1) *If $p \nmid a$ and $-(K_X + \Delta + f^*D)$ is nef, then the closure of $\mathbf{B}_-(-(K_W + D|_W))$ in Y does not contain w .*
- (2) *If K_X is \mathbb{Q} -Cartier and $-(K_X + \Delta + f^*D)$ is ample, then $\mathbf{B}_+(-(K_W + D|_W))$ does not contain w .*

Theorem 4.2. *Let the notation be as above. Take $b \in \mathbb{Z}_{>0}$ so that bD is Cartier and set $\mathcal{M} := \mathcal{O}_X(-ab(K_X + \Delta + f^*D))$. Assume that $\mathcal{M}|_U$ is $f|_U$ -free for some open subset $U \subseteq X$ containing $f^{-1}(w)$. If $p \nmid a$ and if $f_*\mathcal{M}$ is weakly positive over $\{w\}$, then $w \notin \mathbf{B}_-(-(K_W + D|_W))$.*

Remarks 4.3. (1) When X is normal, we may choose a as the Cartier index of $K_X + \Delta$.

- (2) When $w \in W$ is the generic point, assumptions (i) and (ii) above hold. However, assumption (iii) does not necessarily hold even if X is smooth and $\Delta = 0$.
- (3) If X is a variety, then $\operatorname{codim}_Y(Y \setminus W) \geq 2$. Furthermore, if $\operatorname{codim}_Y(Y \setminus W) \geq 2$ and K_Y is \mathbb{Q} -Cartier, then $\mathbf{B}_-(-(K_W + D|_W)) = \mathbf{B}_-(-(K_Y + D))|_W$.

Remark 4.4. Take $m \in \mathbb{Z}$. In the following proof, $m\Delta$ (resp. mf^*D) denotes $\frac{m}{a}E$ (resp. $f^*(mD)$) if $a \mid m$ (resp. mD is Cartier). Note that there may be two distinct AC divisors on X that are equal as $\mathbb{Z}_{(p)}$ -AC divisors.

Proof of Theorem 4.1. We first show that (1) implies (2). By the assumption in (2), Δ is \mathbb{Q} -Cartier. Write $a = mp^c$, where $m, c \in \mathbb{Z}_{\geq 0}$ with $p \nmid m$. Take $e \gg 0$. Put $a' := m(p^e + 1)$. Then $p \nmid a'$. Set $\Delta' := (p^{e-c}E) \otimes \frac{1}{a'}$. We then have $\Delta' = p^{e-c}a/a'\Delta = p^e/(p^e + 1)\Delta \leq \Delta$, so (X, Δ') satisfies assumption (iii). Since $e \gg 0$, we may assume that $-(K_X + \Delta' + f^*D) = -(K_X + \Delta + f^*D) - 1/(p^e + 1)\Delta$ is ample. Let H be an ample \mathbb{Q} -Cartier divisor on Y such that $-(K_X + \Delta' + f^*(D + H))$ is nef. Then, (1) implies that

$w \notin \mathbf{B}_-(-(K_W + D|_W + H|_W))$. Putting $\Gamma := -(K_W + D|_W)$, we obtain that $\mathbf{B}_+(\Gamma) \subseteq \mathbf{B}(\Gamma - \frac{1}{2}H|_W) \subseteq \mathbf{B}_-(\Gamma - H|_W) \subseteq W \setminus \{w\}$.

We begin the proof of (1). Define W_0 to be the subset of points in W satisfying conditions (i)–(iii). We first claim that $W_0 \subseteq W$ is open. Lemma 3.7 (1) tells us that (iii) is an open condition on W . Set $r := \dim X - \dim Y$ and $Z := (X \setminus U)_{\text{red}}$. Then, condition (i) (resp. (ii)) is equivalent to saying that $\dim Z_w \leq r - 2$ (resp. $\dim E_w \leq r - 1$). Hence, our claim follows from Chevalley's theorem [EGA IV₃ 1966, Corollaire 13.1.5], which says that the function $\delta(w) := \dim Z_w$ (resp. $\delta(w) := \dim E_w$) on W is upper semicontinuous. Next, let us introduce some notation:

(n1) Take $d \in \mathbb{Z}_{>0}$ with $a \mid d$ such that dD and $d(K_X + \Delta)$ are Cartier.

(n2) Let \mathcal{A}' be an ample line bundle on X and put $\mathcal{A}'|_V := \mathcal{A}$.

(n3) Denote $g_*\mathcal{A}$ by \mathcal{G} .

Lemmas 3.4–3.7 tell us that, by replacing \mathcal{A}' , we may assume that the following conditions hold:

(a1) For every nef line bundle \mathcal{N}' on X and each $0 \leq i < d$ with $a \mid i$, the sheaf

$$g_*(\mathcal{O}_V(-i(K_V + \Delta|_V)) \otimes \mathcal{A} \otimes \mathcal{N}) \cong (f_*(\mathcal{O}_X(-i(K_X + \Delta)) \otimes \mathcal{A}' \otimes \mathcal{N}'))|_W$$

is a locally free sheaf generated by its global sections, where $\mathcal{N} := \mathcal{N}'|_V$. In particular, $\mathcal{G} = g_*\mathcal{A}$ is locally free (Lemmas 3.4 and 3.5).

(a2) For a g -nef line bundle \mathcal{N} on V , the natural morphism

$$\mathcal{G} \otimes g_*(\mathcal{A}^n \otimes \mathcal{N}) = (g_*\mathcal{A}) \otimes g_*(\mathcal{A}^n \otimes \mathcal{N}) \rightarrow g_*(\mathcal{A}^{n+1} \otimes \mathcal{N})$$

is surjective for each $n \in \mathbb{Z}_{>0}$ (Lemma 3.6).

(a3) For each $e \in \mathbb{Z}_{>0}$ with $a \mid (p^e - 1)$, there exists a morphism

$$g_*(\mathcal{L}_{(V/W, \Delta|_V)}^{(e)} \otimes \mathcal{A}^{p^e}) \rightarrow g_{W^e*}\mathcal{A}_{W^e} \cong (F_W^e)^*g_*\mathcal{A} = (F_W^e)^*\mathcal{G}$$

that is surjective over W_0 (Lemma 3.7).

We continue to introduce some notation:

(n4) Take an ample Cartier divisor H on W such that for each $0 \leq i < d$, there is a surjective morphism $\bigoplus^t \omega_Y^{-i} \otimes \mathcal{G} \rightarrow \mathcal{H} := \mathcal{O}_W(H)$. Such an H exists as shown in Lemma 3.1.

(n5) Fix $e \in \mathbb{Z}_{>0}$ with $a \mid (p^e - 1)$ and write $p^e - 1 = dq + r$ for $q, r \in \mathbb{Z}$ with $0 \leq r < d$. Note that $a \mid r$.

(n6) Set \mathcal{N}' to be the nef line bundle $\mathcal{O}_X(-dq(K_X + \Delta + f^*D))$ and

$$\mathcal{N} := \mathcal{N}'|_V \cong \mathcal{O}_V(-dq(K_V + \Delta|_V + g^*(D|_W))).$$

Also, set $\mathcal{P} := \mathcal{O}_W(-dq(K_W + D|_W))$.

(n7) Recall that $\mathcal{L}_{(V/W, \Delta|_V)}^{(e)} := \mathcal{O}_V((1 - p^e)(K_{V/W} + \Delta|_V))$. Here, we identify V^e (resp. W^e) with V (resp. W). Note that $\mathcal{L}_{(V/W, \Delta|_V)}$ is g -nef in this situation.

We prove the assertion. We have

$$\mathcal{N} \otimes g^* \mathcal{P}^{-1} \underset{(n6)}{\cong} \mathcal{O}_V(-dq(K_{V/W} + \Delta|_V)) \underset{(n7)}{\cong} \mathcal{L}_{(V/W, \Delta|_V)}^{(e)} \otimes \mathcal{O}_V(r(K_{V/W} + \Delta|_V)).$$

We then obtain

$$\mathcal{O}_V(-r(K_V + \Delta|_V)) \otimes \mathcal{A} \otimes \mathcal{N} \cong \mathcal{L}_{(V/W, \Delta|_V)}^{(e)} \otimes \mathcal{A} \otimes g^*(\mathcal{P} \otimes \omega_W^{-r}),$$

so the projection formula implies that

$$g_*(\mathcal{O}_V(-r(K_V + \Delta|_V)) \otimes \mathcal{A} \otimes \mathcal{N}) \cong g_*(\mathcal{L}_{(V/W, \Delta|_V)}^{(e)} \otimes \mathcal{A}) \otimes \mathcal{P} \otimes \omega_W^{-r}. \quad (*)$$

It then follows from (a1) that the right-hand side is globally generated. Hence, we may apply Lemma 3.2 to the composition of the following morphisms which are surjective over W_0 :

$$\begin{aligned} \left(\bigotimes^{p^e} \mathcal{G} \right) \otimes \bigoplus^t ((g_*(\mathcal{L}_{(V/W, \Delta|_V)}^{(e)} \otimes \mathcal{A})) \otimes \mathcal{P} \otimes \omega_Y^{-r}) &\cong \left(\bigotimes^{p^e-1} \mathcal{G} \right) \otimes (g_*(\mathcal{L}_{(V/W, \Delta|_V)}^{(e)} \otimes \mathcal{A})) \otimes \mathcal{P} \otimes \left(\bigoplus^t \mathcal{G} \otimes \omega_Y^{-r} \right) \\ &\xrightarrow{(n4)} \left(\bigotimes^{p^e-1} \mathcal{G} \right) \otimes (g_*(\mathcal{L}_{(V/W, \Delta|_V)}^{(e)} \otimes \mathcal{A})) \otimes \mathcal{P} \otimes \mathcal{H} \\ &\xrightarrow{(a2)} (g_*(\mathcal{L}_{(V/W, \Delta|_V)}^{(e)} \otimes \mathcal{A}^{p^e})) \otimes \mathcal{P} \otimes \mathcal{H} \\ &\xrightarrow{(a3)} ((F_Y^e)^* \mathcal{G}) \otimes \mathcal{P} \otimes \mathcal{H}. \end{aligned}$$

Note that $\mathcal{P} \otimes \mathcal{H} \cong \mathcal{O}_W(dq(-(K_W + D|_W) + \frac{1}{dq}H))$. Thus, we obtain

$$\mathbf{B}(-(K_Y + D) + \frac{2}{dq}H) \underset{\text{by def}}{\subseteq} \mathbf{B}_-(-(K_Y + D) + \frac{1}{dq}H) \underset{\text{Lemma 3.2}}{\subseteq} W \setminus W_0.$$

Since $\frac{2}{dq}$ goes to zero as $e \rightarrow \infty$, we conclude from Lemma 2.10 that $\mathbf{B}_-(-(K_Y + D)) \subseteq W \setminus W_0$. \square

Proof of Theorem 4.2. Replacing $f : X \rightarrow Y$ by $g : V \rightarrow W$, we may assume that $X = V$ and $Y = W$. Then f is flat. Set $Y_0 := Y \setminus (f(X \setminus U))$. Note that $Y_0 \subseteq Y$ is an open subset containing w . We may assume $U = f^{-1}(Y_0)$ by shrinking U . Put $f_0 := f|_U : U \rightarrow Y_0$. Since $\mathcal{M}|_U$ is f_0 -free by assumption, f_0 can be decomposed as

$$f_0 : U \xrightarrow{\sigma} T \xrightarrow{\tau} Y_0$$

and $\mathcal{M}|_U \cong \sigma^* \mathcal{R}$ for a τ -ample line bundle \mathcal{R} on T . For each $c \in \mathbb{Z}_{>0}$, the projection formula says that $\sigma_*(\mathcal{M}^c|_U) \cong (\sigma_* \mathcal{O}_U) \otimes \mathcal{R}^c$, so we get

$$(f_* \mathcal{M}^c)|_{Y_0} \cong (f_0)_*(\mathcal{M}^c|_U) \cong \tau_* \sigma_*(\mathcal{M}^c|_U) \cong \tau_*((\sigma_* \mathcal{O}_U) \otimes \mathcal{R}^c).$$

The last sheaf is locally free if $c \gg 0$, as shown in [Hartshorne 1977, Theorem III 9.9 and its proof]. Fix such a c . Replacing b by bc , we may assume that $(f_* \mathcal{M})|_{Y_0}$ is locally free. We then have a closed subset $Z \subset Y$ of codimension at least 2 such that $Y_0 \subseteq Y \setminus Z$ and $(f_* \mathcal{M})|_{Y \setminus Z}$ is locally free. Shrinking Y to $Y \setminus Z$, we may assume that $f_* \mathcal{M}$ is locally free. Take $\alpha \in \mathbb{Z}_{>0}$ and an ample Cartier divisor H on Y . Set

$\mathcal{H} := \mathcal{O}_Y(H)$. Then, there is $\beta \in \mathbb{Z}_{>0}$ such that $(S^{\alpha\beta} f_* \mathcal{M}) \otimes \mathcal{H}^\beta$ is globally generated over $\{w\}$. We may assume that this sheaf is globally generated over Y_0 , shrinking Y_0 to a neighborhood of w . We set

$$(n8) \quad d := ab\alpha\beta$$

and use notation (n2) and (n3) in the proof of Theorem 4.1. Assume that \mathcal{A} satisfies conditions (a2) and (a3). Furthermore, we add the following notation:

(n9) Take $n_0 \in \mathbb{Z}_{>0}$ with $a \mid n_0$ such that for each $n \geq n_0$, $0 \leq i < d$ and every line bundle \mathcal{Q} on Y , the natural morphism

$$(f_* \mathcal{M}) \otimes f_*(\mathcal{M}^n \otimes \mathcal{O}_X(-i(K_X + \Delta))) \otimes \mathcal{A} \otimes f^* \mathcal{Q} \rightarrow f_*(\mathcal{M}^{n+1} \otimes \mathcal{O}_X(-i(K_X + \Delta))) \otimes \mathcal{A} \otimes f^* \mathcal{Q}$$

is surjective over Y_0 . We can find such an n_0 by Lemma 3.6.

(n10) Choose $v \in \mathbb{Z}_{>0}$ so that

- $\mathcal{H}^v \otimes f_* \mathcal{O}_X(-i(K_X + \Delta))$ is generated by its global sections for each $i \in a\mathbb{Z}$ with $abn_0 \leq i < abn_0 + d$, and
- there is a morphism $\bigoplus^t \mathcal{G} \rightarrow \mathcal{H}^v$ that is surjective over Y_0 .

The existence of such a v is ensured by Lemma 3.1.

(n11) Fix $e \in \mathbb{Z}_{>0}$ with $a \mid (p^e - 1)$ and write $p^e - 1 = dq + r$ for $q, r \in \mathbb{Z}$ with $abn_0 \leq r < abn_0 + d$. Note that $a \mid r$.

We also use notation (n6) and (n7) in the proof of Theorem 4.1. Then,

$$\begin{aligned} \mathcal{N} &= \mathcal{O}_X(-dq(K_X + \Delta + f^* D)) \cong \mathcal{M}^{\alpha\beta q} \quad \text{and} \\ \mathcal{O}_X(-r(K_X + \Delta)) &\cong \mathcal{M}^{n_0} \otimes \mathcal{O}_X(-(abn_0 - r)(K_X + \Delta)) \otimes g^* \mathcal{O}_Y(abn_0 D). \end{aligned}$$

Note that $0 \leq abn_0 - r < d$. Therefore, the morphisms

$$\begin{aligned} (S^q(\mathcal{H}^\beta \otimes S^{\alpha\beta} f_* \mathcal{M})) \otimes \mathcal{H}^v \otimes f_*(\mathcal{O}_X(-r(K_X + \Delta))) \otimes \mathcal{A} \\ \cong \mathcal{H}^{\beta q + v} \otimes (S^q(S^{\alpha\beta} f_* \mathcal{M})) \otimes f_*(\mathcal{O}_X(-r(K_X + \Delta))) \otimes \mathcal{A} \\ \xrightarrow{(n9)} \mathcal{H}^{\beta q + v} \otimes f_*(\mathcal{O}_X(-r(K_X + \Delta))) \otimes \mathcal{A} \otimes \mathcal{M}^{\alpha\beta q} \\ \cong \mathcal{H}^{\beta q + v} \otimes f_*(\mathcal{O}_X(-r(K_X + \Delta))) \otimes \mathcal{A} \otimes \mathcal{N} \\ \cong \mathcal{H}^{\beta q + v} \otimes f_*(\mathcal{L}_{(X/Y, \Delta)}^{(e)} \otimes \mathcal{A}) \otimes \mathcal{P} \otimes \omega_Y^{-r} \\ (*) \end{aligned}$$

are surjective over Y_0 . Here, the last isomorphism is $(*)$ in the proof of Theorem 4.1. By the choice of β and v , we see that the first sheaf is globally generated over Y_0 , and hence so is the last sheaf. Now, we

have the following sequence of morphisms that are surjective over Y_0 :

$$\begin{aligned}
 (\otimes^{p^e} \mathcal{G}) \otimes \bigoplus^t (\mathcal{H}^{\beta q + v} \otimes f_*(\mathcal{L}_{(X/Y, \Delta)}^{(e)} \otimes \mathcal{A}) \otimes \mathcal{P} \otimes \omega_Y^{-r}) \\
 \cong (\otimes^{p^e - 1} \mathcal{G}) \otimes (f_*(\mathcal{L}_{(X/Y, \Delta)}^{(e)} \otimes \mathcal{A})) \otimes \mathcal{P} \otimes (\bigoplus^t \mathcal{G} \otimes \omega_Y^{-r}) \otimes \mathcal{H}^{\beta q + v} \\
 \xrightarrow{(n10)} (\otimes^{p^e - 1} \mathcal{G}) \otimes (f_*(\mathcal{L}_{(X/Y, \Delta)}^{(e)} \otimes \mathcal{A})) \otimes \mathcal{P} \otimes \mathcal{H}^{\beta q + 2v} \\
 \xrightarrow{(a2)} (f_*(\mathcal{L}_{(X/Y, \Delta)}^{(e)} \otimes \mathcal{A}^{p^e})) \otimes \mathcal{P} \otimes \mathcal{H}^{\beta q + 2v} \\
 \xrightarrow{(a3)} ((F_Y^e)^* \mathcal{G}) \otimes \mathcal{P} \otimes \mathcal{H}^{\beta q + 2v}.
 \end{aligned}$$

Note that $\mathcal{P} \otimes \mathcal{H}^{\beta q + 2v} \cong \mathcal{O}_X(dq(-(K_Y + D) + (\beta q + 2v)/(dq)H))$. Replacing e by some larger one if necessary, we may assume that $\beta q > 2v$. Then,

$$\mathbf{B}\left(-(K_Y + D) + \frac{2\beta q}{dq}H\right) \subseteq_{\text{by def}} \mathbf{B}_-\left(-(K_Y + D) + \frac{\beta q + 2v}{dq}H\right) \subseteq_{\text{Lemma 3.2}} Y \setminus Y_0.$$

Since $\frac{2\beta q}{dq} = \frac{2}{ab\alpha}$ goes to zero as $\alpha \rightarrow \infty$, we conclude from Lemma 2.10 that $\mathbf{B}_-(-(K_Y + D)) \subseteq Y \setminus Y_0$. \square

4B. Corollaries. In this subsection, we give several corollaries of the main theorems. *Throughout this subsection, we use the following notation:*

Let $f : X \rightarrow Y$ be a surjective morphism between regular projective varieties over an F -finite field, Δ an effective \mathbb{Q} -divisor on X , and a the Cartier index of Δ . Let D be a \mathbb{Q} -divisor on Y . Let $\bar{\eta}$ denote the geometric generic point of Y .

Corollary 4.5. *Assume that f is flat. Suppose that $\text{Supp } \Delta$ does not contain any component of any fiber, and $(X_{\bar{y}}, \Delta_{\bar{y}})$ is F -pure for every point $y \in Y$:*

- (1) *If $p \nmid a$ and if $-(K_X + \Delta + f^*D)$ is nef, then so is $-(K_Y + D)$.*
- (2) *If $-(K_X + \Delta + f^*D)$ is ample, then so is $-(K_Y + D)$.*

Proof. This follows from Theorem 4.1 and Remarks 2.9 immediately. \square

The author learned the proof of Corollary 4.6(3) below from professor Yoshinori Gongyo.

Corollary 4.6. *Assume that $(X_{\bar{\eta}}, \Delta_{\bar{\eta}})$ is F -pure:*

- (1) *If $p \nmid a$ and if $-(K_X + \Delta + f^*D)$ is nef, then $-(K_Y + D)$ is pseudoeffective.*
- (2) *If $-(K_X + \Delta + f^*D)$ is ample, then $-(K_Y + D)$ is big.*
- (3) *If $(X_{\bar{\eta}}, \Delta_{\bar{\eta}})$ is strongly F -regular and if $-(K_X + \Delta + f^*D)$ is nef and big, then $-(K_Y + D)$ is big.*

Proof. By Remarks 2.9, we see that (1) and (2) follow from (1) and (2) of Theorem 4.1, respectively. We prove (3). By Kodaira's lemma, there is a \mathbb{Q} -divisor $\Delta' \geq \Delta$ on X such that $-(K_X + \Delta' + f^*D)$ is ample and $(X_{\bar{\eta}}, \Delta'_{\bar{\eta}})$ is again strongly F -regular. Hence (3) follows from (2). \square

Corollary 4.7. *Assume that $(X_{\bar{\eta}}, \Delta_{\bar{\eta}})$ is F -pure. If $p \nmid a$ and if $K_X + \Delta$ is numerically equivalent to $f^*(K_Y + L)$ for some \mathbb{Q} -divisor L on Y , then L is pseudoeffective.*

Proof. Set $D := -(K_Y + L)$. Then, $K_X + \Delta + f^*D$ is numerically trivial, so it is nef. Hence, by Corollary 4.6(1), we obtain the assertion. \square

Remarks 4.8. (1) In the situation of Corollary 4.7, it is known that if $(X_{\bar{\eta}}, \Delta_{\bar{\eta}})$ is globally F -split, then $\kappa(L) \geq 0$ (see [Das and Schwede 2017, Theorem B] or [Ejiri 2017, Theorem 3.18]). Of course, $(X_{\bar{\eta}}, \Delta_{\bar{\eta}})$ is not necessary globally F -split even if $X_{\bar{\eta}}$ is a smooth curve and $\Delta = 0$. At the same time, Chen and Zhang proved that the relative canonical divisor of an elliptic fibration has nonnegative Kodaira dimension [Chen and Zhang 2015, 3.2].

(2) In the case when $\dim Y = 1$, Corollary 4.7 follows from a result of Patakfalvi [2014, Theorem 1.6].

Corollary 4.9. *Assume that f is flat and every geometric fiber is F -pure:*

- (1) *If X is a Fano variety, that is, $-K_X$ is ample, then so is Y .*
- (2) *Suppose that the geometric generic fiber of f is strongly F -regular. If X is a weak Fano variety, that is, $-K_X$ is nef and big, then so is Y .*

Proof. Putting $\Delta = 0$ and $D = 0$, we see that the assertions follow from Corollaries 4.5(2) and 4.6(3). \square

Corollary 4.10. *Assume that Y has positive dimension:*

- (1) *If $(X_{\bar{\eta}}, \Delta_{\bar{\eta}})$ is F -pure, then $-(K_{X/Y} + \Delta)$ is not ample.*
- (2) *If $(X_{\bar{\eta}}, \Delta_{\bar{\eta}})$ is strongly F -regular, then $-(K_{X/Y} + \Delta)$ cannot be both nef and big.*

Proof. Set $D := -K_Y$. Then $-(K_X + \Delta + f^*D) = -(K_{X/Y} + \Delta)$. Since $-(K_Y + D) = 0$ is not big, the assertions follows from Corollary 4.6(2) and (3). \square

Corollary 4.11. *Assume that Y has positive dimension. Suppose that $\mathcal{O}_X(-ab(K_X + \Delta))|_{X_{\bar{\eta}}}$ is globally generated for some $b \in \mathbb{Z}_{>0}$. If $p \nmid a$ and if $(X_{\bar{\eta}}, \Delta_{\bar{\eta}})$ is F -pure, then $f_*\mathcal{O}_X(-ab(K_{X/Y} + \Delta))$ is not big.*

Proof. Set $\mathcal{G}_{(-l)} := f_*\mathcal{O}_X(al(K_X + \Delta))$ for each $l \in \mathbb{Z}$. Suppose that $\mathcal{G}_{(b)}$ is big. Then, $\mathcal{H}^{-1} \otimes S^\gamma \mathcal{G}_{(b)}$ is weakly positive for some $\gamma \in \mathbb{Z}_{>0}$ and an ample line bundle \mathcal{H} on Y . Take $n_0 \in \mathbb{Z}_{>0}$ so that the natural morphism $\mathcal{G}_{(b)} \otimes \mathcal{G}_{(n)} \rightarrow \mathcal{G}_{(b+n)}$ is generically surjective for each $n \geq n_0$. We can find such an n_0 by Lemma 3.6. Choose $\nu \in \mathbb{Z}_{>0}$ so that $\mathcal{H}^\nu \otimes \mathcal{G}_{(n_0)}$ is globally generated. Fix $l \in \mathbb{Z}$ with $l > \nu$. Using the natural morphism $S^l(S^\gamma \mathcal{G}_{(b)}) \otimes \mathcal{G}_{(n_0)} \rightarrow \mathcal{G}_{(bl\gamma + n_0)}$ and [Viehweg 1995, Lemma 2.16], we see that $\mathcal{H}^{\nu-l} \otimes \mathcal{G}_{(bl\gamma + n_0)}$ is weakly positive. Let H be a Cartier divisor on Y satisfying $\mathcal{O}_Y(H) \cong \mathcal{H}$ and set $H' := (l - \nu)/(a(bl\gamma + n_0))H$. The projection formula then shows that

$$\mathcal{H}^{\nu-l} \otimes \mathcal{G}_{(bl\gamma + n_0)} \cong f_*\mathcal{O}_X(-a(bl\gamma + n_0)(K_{X/Y} + \Delta + f^*H')).$$

It then follows from Theorem 4.2 that $\mathbf{B}_-(-H') \neq \emptyset$, i.e., $-H'$ is pseudoeffective, a contradiction. \square

5. Results in arbitrary characteristic

In this section we generalize several results in Section 4 to arbitrary characteristic. In particular, we prove the characteristic zero counterparts of Corollaries 4.10 and 4.11 (Theorems 5.4 and 5.5). We also deal with a morphism that is special but not necessarily smooth, and show that the image of a Fano variety is again a Fano variety.

To begin with, let us recall the following definition:

Definition 5.1. Let X be a normal variety over a field k of characteristic zero, and Δ an effective \mathbb{Q} -Weil divisor on X . Let (X_R, Δ_R) be a model of (X, Δ) over a finitely generated \mathbb{Z} -subalgebra R of k . We say that (X, Δ) is of *dense F -pure type* (resp. *strongly F -regular type*) if there exists a dense (resp. dense open) subset $S \subseteq \operatorname{Spec} R$ such that (X_μ, Δ_μ) is F -pure (resp. strongly F -regular) for all closed points $\mu \in S$.

Remark 5.2. The above definition can be generalized in an obvious way to the case where X is a finite disjoint union of varieties over k .

Theorem 5.3 [Takagi 2004, Corollary 3.4]. *Let X be a normal variety over a field of characteristic zero, and Δ an effective \mathbb{Q} -Weil divisor on X such that $K_X + \Delta$ is \mathbb{Q} -Cartier. Then (X, Δ) is klt if and only if it is of strongly F -regular type.*

Theorem 5.4. *Let k be an algebraically closed field of characteristic zero. Let $f : X \rightarrow Y$ be a surjective morphism between smooth projective varieties of positive dimension, and Δ an effective \mathbb{Q} -divisor on X . If (X_y, Δ_y) is of dense F -pure type (resp. klt) for every general closed point $y \in Y$, then $-(K_{X/Y} + \Delta)$ cannot be ample (resp. both nef and big).*

Proof. Let X_R, Δ_R, Y_R, y_R and f_R be models of X, Δ, Y, y and f over a finitely generated \mathbb{Z} -algebra R , respectively. We may assume that $(X_R)_{y_R}$ is a model of X_y over R . We first suppose that (X_y, Δ_y) is of dense F -pure type for a general closed point $y \in Y$. Then, there is a dense subset $S \subseteq \operatorname{Spec} R$ such that $((X_y)_\mu, \Delta_\mu)$ is F -pure for every $\mu \in S$. Note that $(X_y)_\mu \cong (X_\mu)_{y_\mu}$ and $(\Delta_y)_\mu = (\Delta_\mu)_{y_\mu}$. Corollary 4.10 then implies that $-(K_{X_\mu/Y_\mu} + \Delta_\mu)$ is not ample, which means that $-(K_{X/Y} + \Delta)$ is not ample. We next suppose that (X_y, Δ_y) is klt for every general closed point $y \in Y$. If $-(K_{X/Y} + \Delta)$ is nef and big, then by Kodaira's lemma, there is $\Delta' \geq \Delta$ such that $-(K_{X/Y} + \Delta')$ is ample and (X_y, Δ'_y) is klt for a general closed point $y \in Y$. However, Theorem 5.3 tells us that (X_y, Δ'_y) is of dense F -pure type, which contradicts the above arguments. \square

Theorem 5.5. *Let k be an algebraically closed field of characteristic zero. Let $f : X \rightarrow Y$ be a surjective morphism between smooth projective varieties of positive dimension, and Δ an effective \mathbb{Q} -divisor on X . Assume that (X_y, Δ_y) is of dense F -pure type for a general closed point $y \in Y$. Let $\bar{\eta}$ be a geometric generic point of Y . If $\mathcal{O}_X(-m(K_{X/Y} + \Delta))|_{X_{\bar{\eta}}}$ is globally generated for some $m > 0$ such that $m\Delta$ is integral, then $f_*\mathcal{O}_X(-m(K_{X/Y} + \Delta))$ is not big.*

Proof. Set $\mathcal{G} := f_*\mathcal{O}_X(-m(K_{X/Y} + \Delta))$ and $r := \operatorname{rank} \mathcal{G}$. Since $y \in Y$ is general, f is flat at every point in $f^{-1}(y)$ and $\dim H^0(X_y, -m(K_{X_y} + \Delta_y)) = r$. Let X_R, Δ_R, Y_R, y_R and f_R be models of X, Δ, Y, y and f ,

respectively. By replacing R if necessary, we may assume that $f_{R*}\mathcal{O}_{X_R}(-m(K_{X_R/Y_R} + \Delta_R))$ and $(X_R)_{y_R}$ are respectively models of \mathcal{G} and X_y . We may further assume that $\dim H^0((X_\mu)_{y_\mu}, -m(K_{X_\mu/Y_\mu} + \Delta_\mu)) = r$ for every $\mu \in \operatorname{Spec} R$. Then, [Hartshorne 1977, Corollary 12.9] implies that the natural morphism

$$\mathcal{G}_\mu = f_{R*}\mathcal{O}_{X_R}(-m(K_{X_R/Y_R} + \Delta_R))|_{Y_\mu} \rightarrow f_{\mu*}\mathcal{O}_{X_\mu}(-m(K_{X_\mu/Y_\mu} + \Delta_\mu))$$

is surjective over y_μ . Since $f_{\mu*}\mathcal{O}_{X_\mu}(-m(K_{X_\mu/Y_\mu} + \Delta_\mu))$ is not big as shown in Corollary 4.11, \mathcal{G}_μ is also not big. Hence, the lemma below completes the proof. \square

Lemma 5.6. *Let \mathcal{G} be a torsion-free coherent sheaf on a smooth quasiprojective variety Y over an algebraically closed field of characteristic zero. Let Y_R and \mathcal{G}_R be models of Y and \mathcal{G} respectively over a finitely generated \mathbb{Z} -algebra R . If \mathcal{G} is big, then there exists a dense open subset $S \subseteq \operatorname{Spec} R$ such that \mathcal{G}_μ is big for every $\mu \in S$.*

Proof. Let $Z \subset Y$ be a closed subset of codimension at least 2 such that $\mathcal{G}|_{Y \setminus Z}$ is locally free. Replacing Y by $Y \setminus Z$, we may assume that \mathcal{G} is locally free. By the definition, we have $\gamma \in \mathbb{Z}_{>0}$ such that $\mathcal{H}^{-1} \otimes S^\gamma \mathcal{G}$ is weakly positive for some ample line bundle \mathcal{H} on Y . Then, there is $\beta \in \mathbb{Z}_{>0}$ such that $\mathcal{H}^\beta \otimes S^{2\beta}(\mathcal{H}^{-1} \otimes S^\gamma \mathcal{G}) \cong \mathcal{H}^{-\beta} \otimes S^{2\beta}(S^\gamma \mathcal{G})$ is generically globally generated. Using the natural morphism $S^{2\beta}(S^\gamma \mathcal{G}) \rightarrow S^{2\beta+\gamma} \mathcal{G}$, we see that $\mathcal{F} := \mathcal{H}^{-\beta} \otimes S^{2\beta+\gamma} \mathcal{G}$ is generically globally generated, i.e., there is a morphism $\theta : \bigoplus^t \mathcal{O}_Y \rightarrow \mathcal{F}$ that is surjective over a dense open subset $V \subseteq Y$, where $t \in \mathbb{Z}_{>0}$. Let θ_R , H_R and V_R be models of θ , H and V over R , respectively. Replacing R if necessary, we may assume that θ_R is surjective over V_R . Thus for every closed point $\mu \in \operatorname{Spec} R$, the morphism $\theta_\mu : \bigoplus^t \mathcal{O}_{X_\mu} \rightarrow \mathcal{H}_\mu^{-\beta} \otimes S^{2\beta+\gamma} \mathcal{G}_\mu$ is surjective over V_μ , which means that \mathcal{G}_μ is big. \square

Kollár, Miyaoka and Mori [1992, Corollary 2.9] (compare [Miyaoka 1993, Theorem 3]) proved that images of Fano varieties under smooth morphisms are again Fano varieties. The rest of this paper is devoted to extending this result to toroidal morphisms.

Definition 5.7 [Abramovich and Karu 2000; Kawamata 2002]. Let k be an algebraically closed field of arbitrary characteristic:

- (i) Let X be a normal variety and U an open subset of X . We say that the embedding $U \subseteq X$ is *toroidal* if for every closed point $x \in X$, there exists
 - a toric variety V with torus T ,
 - a closed point $v \in V$ and
 - an isomorphism $\hat{\mathcal{O}}_{X,x} \cong \hat{\mathcal{O}}_{V,v}$ of complete local k -algebras such that the ideal of $B := X \setminus U$ maps isomorphically to that of $V \setminus T$.

Such a pair (V, v) is called a *local model* at $x \in X$. The pair (X, B) is often called a *toroidal variety*.

- (ii) Let (X, B) and (Y, C) be toroidal varieties. A *toroidal morphism* $f : (X, B) \rightarrow (Y, C)$ is a dominant morphism $f : X \rightarrow Y$ with $f(X \setminus B) \subseteq Y \setminus C$ such that for every closed point $x \in X$, there exist
 - local models (V, v) and (W, w) at x and $y := f(x)$, respectively, and

- a toric morphism $g : V \rightarrow W$ such that the following diagram commutes:

$$\begin{array}{ccc} \hat{\mathcal{O}}_{X,x} & \xrightarrow{\cong} & \hat{\mathcal{O}}_{V,v} \\ \hat{f}^\# \uparrow & & \uparrow \hat{g}^\# \\ \hat{\mathcal{O}}_{Y,y} & \xrightarrow{\cong} & \hat{\mathcal{O}}_{W,w} \end{array}$$

The next theorem is a generalization of [Kollár et al. 1992, Corollary 2.9].

Theorem 5.8. *Let k be an algebraically closed field of any characteristic $p \geq 0$. Let $f : X \rightarrow Y$ be a surjective morphism between smooth projective varieties and B a reduced divisor on X . Let Δ be a \mathbb{Q} -divisor on X such that $0 \leq \Delta \leq B$ and that $a\Delta$ is integral for some $0 < a \in \mathbb{Z} \setminus p\mathbb{Z}$. Assume that the following conditions hold:*

- (i) *f induces a toroidal morphism $f : (X, B) \rightarrow (Y, \emptyset)$.*
- (ii) *f is equidimensional.*
- (iii) *Every closed fiber of f is reduced.*
- (iv) *$\text{Supp } \Delta$ does not contain any irreducible component of any fiber.*

*In this situation, if $-(K_X + \Delta + f^*D)$ is ample for some \mathbb{Q} -divisor D on Y , then so is $-(K_Y + D)$.*

Proof. Let $x_\lambda \in X$ be a closed point and set $y_\lambda := f(x_\lambda)$. By assumption (i), there is a local model (V_λ, v_λ) (resp. (W_λ, w_λ)) at x_λ (resp. y_λ) and a toric morphism $g_\lambda : V_\lambda \rightarrow W_\lambda$. Using Artin's approximation theorem [1969, Corollary 2.6], we obtain a commutative diagram

$$\begin{array}{ccccc} X & \xleftarrow{\rho_\lambda} & T_\lambda & \xrightarrow{\mu_\lambda} & V_\lambda \\ f \downarrow & & h_\lambda \downarrow & & \downarrow g_\lambda \\ Y & \xleftarrow{\sigma_\lambda} & U_\lambda & \xrightarrow{\nu_\lambda} & W_\lambda \end{array} \quad (*)$$

such that

- T_λ and U_λ are varieties,
- all the horizontal morphisms are étale, and
- there is a closed point $t_\lambda \in T_\lambda$ such that $\rho_\lambda(t_\lambda) = x_\lambda$ and $\mu_\lambda(t_\lambda) = v_\lambda$.

Let $k[v_1, v_1^{-1}, \dots, v_m, v_m^{-1}]$ (resp. $k[w_1, w_1^{-1}, \dots, w_n, w_n^{-1}]$) be the coordinate ring of the torus of V_λ (resp. W_λ). Set $t_i := \mu_\lambda^* v_i$ and $u_i := \nu_\lambda^* w_i$ for each i . We then see from assumptions (ii) and (iii) that

$$h_\lambda^* u_j = h_\lambda^* \nu_\lambda^* w_j = \mu_\lambda^* g_\lambda^* w_j = \mu_\lambda^* \prod_{l_{j-1} < i \leq l_j} v_i = \prod_{l_{j-1} < i \leq l_j} t_i$$

for $j = 1, \dots, n$, where $0 = l_0 < l_1 < \dots < l_n \leq m$.

Shrinking T_λ if necessary, we may assume that for any closed point $t \in T_\lambda$, there are $a_1, \dots, a_m \in k$ such that $\mathfrak{m}_t = (t_1 - a_1, \dots, t_m - a_m)$. We may also assume a similar condition for U_λ and u_1, \dots, u_n .

Let Λ be a *finite* set of λ such that $X = \bigcup_{\lambda \in \Lambda} \rho_\lambda(T_\lambda)$. When $p = 0$, one can check that diagram (*) can be reduced to characteristic $p \gg 0$ for all $\lambda \in \Lambda$ simultaneously. For this reason, we consider the case of $p > 0$.

We see from [Matsumura 1986, Corollary of Theorem 23.1] that f is flat. Therefore, in order to apply Corollary 4.5, we only need to show that $(Z, \Delta|_Z)$ is F -pure for every closed fiber Z of f . This holds if $(S, (\rho_\lambda^* \Delta)|_S)$ is F -pure for every closed fiber S of h_λ , since ρ_λ is étale. Fix a closed fiber S over $u \in U_\lambda$ and a closed point $t \in T_\lambda$ contained in S . Then, there are $a_1, \dots, a_m, b_1, \dots, b_n \in k$ such that $\mathfrak{m}_t = (t_1 - a_1, \dots, t_m - a_m)$ and $\mathfrak{m}_u = (u_1 - b_1, \dots, u_n - b_n)$. Put $t'_i := t_i - a_i$ and $u'_i := u_i - b_i$ for each i . We then have

$$h_\lambda^* u'_j = \prod_{l_{j-1} < i \leq l_j} (t'_i + a_i) - \prod_{l_{j-1} < i \leq l_j} a_i$$

for $j = 1, \dots, n$. Set $\delta := \prod_{l_n < i \leq m} t_i$. Now, one can easily check that

- the sequence $(h_\lambda^* u'_1, \dots, h_\lambda^* u'_n, \delta)$ is $\mathcal{O}_{T_\lambda, t}$ -regular, and
- $(h_\lambda^* u'_1 \cdots h_\lambda^* u'_n)^{q-1} \cdot \delta^{q-1} \notin \mathfrak{m}_t^{[q]}$ for every $q = p^e$.

Then, [Hara and Watanabe 2002, Corollary 2.7] tells us that $(S, \text{div}(\delta)|_S)$ is F -pure around t . Since $\rho_\lambda^* \Delta \leq \text{div}(\delta)$, we conclude that $(S, (\rho_\lambda^* \Delta)|_S)$ is F -pure around t . \square

Example 5.9. Let $\{e_1, e_2, e_3\}$ be the canonical basis of \mathbb{R}^3 . For integers $m, n \geq 0$, we define $\mathbf{v}_{m,n} := (1, m, n) \in \mathbb{R}^3$. Let $\Sigma_{m,n}$ be the fan consisting of all the faces of the following cones:

$$\begin{aligned} &\langle \mathbf{v}_{m,n}, e_2, e_2 + e_3 \rangle, \langle \mathbf{v}_{m,n}, e_2 + e_3, e_3 \rangle, \langle \mathbf{v}_{m,n}, -e_2, e_3 \rangle, \langle \mathbf{v}_{m,n}, e_2, -e_3 \rangle, \langle \mathbf{v}_{m,n}, -e_2, -e_3 \rangle, \\ &\langle -e_1, e_2, e_2 + e_3 \rangle, \langle -e_1, e_2 + e_3, e_3 \rangle, \langle -e_1, -e_2, e_3 \rangle, \langle -e_1, e_2, -e_3 \rangle, \langle -e_1, -e_2, -e_3 \rangle. \end{aligned}$$

Let $X_{m,n}$ be the smooth toric 3-fold corresponding to the fan $\Sigma_{m,n}$ with respect to the lattice $\mathbb{Z}^3 \subset \mathbb{R}^3$. Then $X_{m,n}$ is a Fano variety if and only if $m, n \in \{0, 1\}$. The projection $\mathbb{R}^3 \rightarrow \mathbb{R}^2 : (x, y, z) \mapsto (x, y)$ induces a toric morphism $f : X_{m,n} \rightarrow Y_m$ from $X_{m,n}$ to the Hirzebruch surface $Y_m := \mathbb{P}_{\mathbb{P}^1}(\mathcal{O}_{\mathbb{P}^1} \oplus \mathcal{O}_{\mathbb{P}^1}(m))$. Set $\Delta = 0$. Then one can check that f satisfies the assumptions of Theorem 5.8, but it is not smooth. Hence by Theorem 5.8, we see that Y_m is a Del Pezzo surface if $m = 0, 1$. In fact, it is well known that Y_m is a Del Pezzo surface if and only if $m = 0, 1$.

Acknowledgements

The author is greatly indebted to his supervisor Professor Shunsuke Takagi for suggesting several problems studied in this paper and for much valuable advice. He wishes to express his gratitude to Professor Yoshinori Gongyo for showing him the proof of Corollary 4.9(2) and for fruitful conversations. He is grateful to Professors Zsolt Patakfalvi and Hiromu Tanaka for suggesting improvements of several results in a preprint version of this paper. He would like to thank Professors Caucher Birkar, Yifei Chen and Osamu Fujino for stimulating discussions and helpful comments. He also would like to thank Professor Akiyoshi Sannai, and Doctors Takeru Fukuoka, Akihiro Kanemitsu, Eleonora Anna Romano, Kenta Sato,

Antoine Song and Fumiaki Suzuki for useful comments. Finally, he is grateful to the referees for a careful reading and for suggesting an improvement of Theorem 5.8. He was supported by JSPS KAKENHI Grant Number 15J09117 and the Program for Leading Graduate Schools, MEXT, Japan.

References

- [Abramovich and Karu 2000] D. Abramovich and K. Karu, “Weak semistable reduction in characteristic 0”, *Invent. Math.* **139**:2 (2000), 241–273. MR Zbl
- [Artin 1969] M. Artin, “Algebraic approximation of structures over complete local rings”, *Inst. Hautes Études Sci. Publ. Math.* **36** (1969), 23–58. MR Zbl
- [Birkar and Chen 2016] C. Birkar and Y. Chen, “Images of manifolds with semi-ample anti-canonical divisor”, *J. Algebraic Geom.* **25**:2 (2016), 273–287. MR Zbl
- [Chen and Zhang 2013] M. Chen and Q. Zhang, “On a question of Demailly–Petersen–Schneider”, *J. Eur. Math. Soc.* **15**:5 (2013), 1853–1858. MR Zbl
- [Chen and Zhang 2015] Y. Chen and L. Zhang, “The subadditivity of the Kodaira dimension for fibrations of relative dimension one in positive characteristics”, *Math. Res. Lett.* **22**:3 (2015), 675–696. MR Zbl
- [Corti 1992] A. Corti, “Adjunction of log divisors”, pp. 171–182 in *Flips and abundance for algebraic threefolds* (Salt Lake City, UT, 1991), edited by J. Kollár, Astérisque **211**, Soc. Math. France, Paris, 1992. Zbl
- [Das and Schwede 2017] O. Das and K. Schwede, “The F -different and a canonical bundle formula”, *Ann. Sc. Norm. Super. Pisa Cl. Sci. (5)* **17**:3 (2017), 1173–1205. MR Zbl
- [Debarre 2001] O. Debarre, *Higher-dimensional algebraic geometry*, Springer, 2001. MR Zbl
- [EGA IV₃ 1966] A. Grothendieck, “Éléments de géométrie algébrique, IV: Étude locale des schémas et des morphismes de schémas, III”, *Inst. Hautes Études Sci. Publ. Math.* **28** (1966), 5–255. MR Zbl
- [Ein et al. 2006] L. Ein, R. Lazarsfeld, M. Mustață, M. Nakamaye, and M. Popa, “Asymptotic invariants of base loci”, *Ann. Inst. Fourier (Grenoble)* **56**:6 (2006), 1701–1734. MR Zbl
- [Ejiri 2017] S. Ejiri, “Weak positivity theorem and Frobenius stable canonical rings of geometric generic fibers”, *J. Algebraic Geom.* **26**:4 (2017), 691–734. MR Zbl
- [Fujino and Gongyo 2012] O. Fujino and Y. Gongyo, “On images of weak Fano manifolds”, *Math. Z.* **270**:1–2 (2012), 531–544. MR Zbl
- [Fujino and Gongyo 2014] O. Fujino and Y. Gongyo, “On images of weak Fano manifolds, II”, pp. 201–207 in *Algebraic and complex geometry* (Hannover, Germany, 2012), edited by A. Frühbis-Krüger et al., Springer Proc. Math. Stat. **71**, Springer, 2014. MR Zbl
- [Hara and Watanabe 2002] N. Hara and K.-I. Watanabe, “ F -regular and F -pure rings vs. log terminal and log canonical singularities”, *J. Algebraic Geom.* **11**:2 (2002), 363–392. MR Zbl
- [Hartshorne 1977] R. Hartshorne, *Algebraic geometry*, Graduate Texts in Math. **52**, Springer, 1977. MR Zbl
- [Hartshorne 1994] R. Hartshorne, “Generalized divisors on Gorenstein schemes”, *K-Theory* **8**:3 (1994), 287–339. MR Zbl
- [Kawamata 2002] Y. Kawamata, “On algebraic fiber spaces”, pp. 135–154 in *Contemporary trends in algebraic geometry and algebraic topology* (Tianjin, 2000), edited by S.-S. Chern et al., Nankai Tracts Math. **5**, World Sci., River Edge, NJ, 2002. MR Zbl
- [Keeler 2003] D. S. Keeler, “Ample filters of invertible sheaves”, *J. Algebra* **259**:1 (2003), 243–283. MR Zbl
- [Kollár et al. 1992] J. Kollár, Y. Miyaoka, and S. Mori, “Rational connectedness and boundedness of Fano manifolds”, *J. Differential Geom.* **36**:3 (1992), 765–779. MR Zbl
- [Lazarsfeld 2004] R. Lazarsfeld, *Positivity in algebraic geometry, II: Positivity for vector bundles, and multiplier ideals*, *Ergebnisse der Mathematik (3)* **49**, Springer, 2004. MR Zbl
- [Matsumura 1986] H. Matsumura, *Commutative ring theory*, Cambridge Studies in Adv. Math. **8**, Cambridge Univ. Press, 1986. MR Zbl

- [Miller and Schwede 2012] L. E. Miller and K. Schwede, “Semi-log canonical vs F -pure singularities”, *J. Algebra* **349**:1 (2012), 150–164. MR Zbl
- [Miyaoaka 1993] Y. Miyaoka, “Relative deformations of morphisms and applications to fibre spaces”, *Comment. Math. Univ. St. Paul.* **42**:1 (1993), 1–7. MR Zbl
- [Mustață 2013] M. Mustață, “The non-nef locus in positive characteristic”, pp. 535–551 in *A celebration of algebraic geometry* (Cambridge, MA, 2011), edited by B. Hassett et al., Clay Math. Proc. **18**, Amer. Math. Soc., Providence, RI, 2013. MR Zbl
- [Patakfalvi 2014] Z. Patakfalvi, “Semi-positivity in positive characteristics”, *Ann. Sci. École Norm. Sup. (4)* **47**:5 (2014), 991–1025. MR Zbl
- [Patakfalvi et al. 2018] Z. Patakfalvi, K. Schwede, and W. Zhang, “ F -singularities in families”, *Algebr. Geom.* **5**:3 (2018), 264–327. MR Zbl
- [Prokhorov and Shokurov 2009] Y. G. Prokhorov and V. V. Shokurov, “Towards the second main theorem on complements”, *J. Algebraic Geom.* **18**:1 (2009), 151–199. MR Zbl
- [Schwede 2008] K. Schwede, “Generalized test ideals, sharp F -purity, and sharp test elements”, *Math. Res. Lett.* **15**:6 (2008), 1251–1261. MR Zbl
- [Takagi 2004] S. Takagi, “An interpretation of multiplier ideals via tight closure”, *J. Algebraic Geom.* **13**:2 (2004), 393–415. MR Zbl
- [Viehweg 1983] E. Viehweg, “Weak positivity and the additivity of the Kodaira dimension for certain fibre spaces”, pp. 329–353 in *Algebraic varieties and analytic varieties* (Tokyo, 1981), edited by S. Iitaka, Adv. Stud. Pure Math. **1**, North-Holland, Amsterdam, 1983. MR Zbl
- [Viehweg 1995] E. Viehweg, *Quasi-projective moduli for polarized manifolds*, Ergebnisse der Mathematik (3) **30**, Springer, 1995. MR Zbl

Communicated by Shigefumi Mori

Received 2018-05-22 Revised 2019-05-09 Accepted 2019-06-13

s-ejiri@cr.math.sci.osaka-u.ac.jp

*Department of Mathematics, Graduate School of Science, Osaka University,
Japan*

Algebra & Number Theory

msp.org/ant

EDITORS

MANAGING EDITOR

Bjorn Poonen
Massachusetts Institute of Technology
Cambridge, USA

EDITORIAL BOARD CHAIR

David Eisenbud
University of California
Berkeley, USA

BOARD OF EDITORS

Richard E. Borcherds	University of California, Berkeley, USA	Martin Olsson	University of California, Berkeley, USA
Antoine Chambert-Loir	Université Paris-Diderot, France	Raman Parimala	Emory University, USA
J-L. Colliot-Thélène	CNRS, Université Paris-Sud, France	Jonathan Pila	University of Oxford, UK
Brian D. Conrad	Stanford University, USA	Anand Pillay	University of Notre Dame, USA
Samit Dasgupta	University of California, Santa Cruz, USA	Michael Rapoport	Universität Bonn, Germany
Hélène Esnault	Freie Universität Berlin, Germany	Victor Reiner	University of Minnesota, USA
Gavril Farkas	Humboldt Universität zu Berlin, Germany	Peter Sarnak	Princeton University, USA
Hubert Flenner	Ruhr-Universität, Germany	Joseph H. Silverman	Brown University, USA
Sergey Fomin	University of Michigan, USA	Michael Singer	North Carolina State University, USA
Edward Frenkel	University of California, Berkeley, USA	Christopher Skinner	Princeton University, USA
Wee Teck Gan	National University of Singapore	Vasudevan Srinivas	Tata Inst. of Fund. Research, India
Andrew Granville	Université de Montréal, Canada	J. Toby Stafford	University of Michigan, USA
Ben J. Green	University of Oxford, UK	Pham Huu Tiep	University of Arizona, USA
Joseph Gubeladze	San Francisco State University, USA	Ravi Vakil	Stanford University, USA
Roger Heath-Brown	Oxford University, UK	Michel van den Bergh	Hasselt University, Belgium
Craig Huneke	University of Virginia, USA	Akshay Venkatesh	Institute for Advanced Study, USA
Kiran S. Kedlaya	Univ. of California, San Diego, USA	Marie-France Vignéras	Université Paris VII, France
János Kollár	Princeton University, USA	Kei-Ichi Watanabe	Nihon University, Japan
Philippe Michel	École Polytechnique Fédérale de Lausanne	Melanie Matchett Wood	University of Wisconsin, Madison, USA
Susan Montgomery	University of Southern California, USA	Shou-Wu Zhang	Princeton University, USA
Shigefumi Mori	RIMS, Kyoto University, Japan		

PRODUCTION

production@msp.org
Silvio Levy, Scientific Editor


See inside back cover or msp.org/ant for submission instructions.

The subscription price for 2019 is US \$385/year for the electronic version, and \$590/year (+\$60, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to MSP.

Algebra & Number Theory (ISSN 1944-7833 electronic, 1937-0652 printed) at Mathematical Sciences Publishers, 798 Evans Hall #3840, c/o University of California, Berkeley, CA 94720-3840 is published continuously online. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices.

ANT peer review and production are managed by EditFlow® from MSP.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2019 Mathematical Sciences Publishers

Algebra & Number Theory

Volume 13 No. 9 2019

Proof of a conjecture of Colliot-Thélène and a diophantine excision theorem JAN DENEUF	1983
Irreducible characters with bounded root Artin conductor AMALIA PIZARRO-MADARIAGA	1997
Frobenius–Perron theory of endofunctors JIANMIN CHEN, ZHIBIN GAO, ELIZABETH WICKS, JAMES J. ZHANG, XIAOHONG ZHANG and HONG ZHU	2005
Positivity of anticanonical divisors and F -purity of fibers SHO EJIRI	2057
A probabilistic approach to systems of parameters and Noether normalization JULIETTE BRUCE and DANIEL ERMAN	2081
The structure of correlations of multiplicative functions at almost all scales, with applications to the Chowla and Elliott conjectures TERENCE TAO and JONI TERÄVÄINEN	2103
VI-modules in nondescribing characteristic, part I ROHIT NAGPAL	2151
Degree of irrationality of very general abelian surfaces NATHAN CHEN	2191
Lower bounds for the least prime in Chebotarev ANDREW FIORI	2199
Brody hyperbolicity of base spaces of certain families of varieties MIHNEA POPA, BEHROUZ TAJI and LEI WU	2205



1937-0652(2019)13:9;1-4