

Algebra & Number Theory

Volume 14
2020
No. 2

On the definition of quantum Heisenberg category

Jonathan Brundan, Alistair Savage and Ben Webster

On the definition of quantum Heisenberg category

Jonathan Brundan, Alistair Savage and Ben Webster

We introduce a diagrammatic monoidal category $\mathcal{Heis}_k(z, t)$ which we call the *quantum Heisenberg category*; here, $k \in \mathbb{Z}$ is “central charge” and z and t are invertible parameters. Special cases were known before: for central charge $k = -1$ and parameters $z = q - q^{-1}$ and $t = -z^{-1}$ our quantum Heisenberg category may be obtained from the deformed version of Khovanov’s Heisenberg category introduced by Licata and Savage by inverting its polynomial generator, while $\mathcal{Heis}_0(z, t)$ is the affinization of the HOMFLY-PT skein category. We also prove a basis theorem for the morphism spaces in $\mathcal{Heis}_k(z, t)$.

1. Introduction

Fix a commutative ground ring \mathbb{k} and parameters $z, t \in \mathbb{k}^\times$. This paper introduces a family of pivotal monoidal categories $\mathcal{Heis}_k(z, t)$, one for each *central charge* $k \in \mathbb{Z}$. We refer to these categories as *quantum Heisenberg categories*. The terminology is due to a connection to Khovanov’s Heisenberg category [2014]: our category for central charge $k = -1$ is a two parameter deformation of the category from [loc. cit.], and is closely related to the one parameter deformation introduced already by Licata and Savage [2013]. The category $\mathcal{Heis}_0(z, t)$ has also already appeared in the literature: it is the *affine HOMFLY-PT skein category* from [Brundan 2017, Section 4]. For more general central charges, our categories are new. They were discovered by mimicking the approach of Brundan [2018], where the definition of the degenerate Heisenberg categories introduced in [Mackaay and Savage 2018] was reformulated.

In fact, we will give three different monoidal presentations of $\mathcal{Heis}_k(z, t)$. They all start from the affine Hecke algebra AH_n associated to the symmetric group \mathfrak{S}_n . It is convenient to assemble these algebras for all $n \geq 0$ into a single monoidal category $\mathcal{AH}(z)$. By definition, this is the strict \mathbb{k} -linear monoidal category generated by one object \uparrow and two morphisms $x : \uparrow \rightarrow \uparrow$ and $\tau : \uparrow \otimes \uparrow \rightarrow \uparrow \otimes \uparrow$, subject to the

This article is based upon work done while the first author was in residence at the Mathematical Sciences Research Institute in Berkeley, California during the Spring 2018 semester, which was supported by the National Science Foundation grant DMS-1440140. The first author was also supported by DMS-1700905. The second and third authors were supported by Discovery Grants from the Natural Sciences and Engineering Research Council of Canada. The research of the third author was also supported in part by Perimeter Institute for Theoretical Physics. Research at Perimeter Institute is supported by the Government of Canada through the Department of Innovation, Science and Economic Development Canada and by the Province of Ontario through the Ministry of Research, Innovation and Science.

MSC2010: primary 17B10; secondary 18D10.

Keywords: Heisenberg category, categorification, affine Hecke algebra.

relations

$$\tau \circ (1_{\uparrow} \otimes x) \circ \tau = x \otimes 1_{\uparrow}, \quad (1.1)$$

$$\tau \circ \tau = z\tau + 1_{\uparrow \otimes \uparrow}, \quad (1.2)$$

$$(\tau \otimes 1_{\uparrow}) \circ (1_{\uparrow} \otimes \tau) \circ (\tau \otimes 1_{\uparrow}) = (1_{\uparrow} \otimes \tau) \circ (\tau \otimes 1_{\uparrow}) \circ (1_{\uparrow} \otimes \tau). \quad (1.3)$$

The second relation here implies that τ is invertible. We also require that x is invertible, i.e., there is another generator x^{-1} such that

$$x \circ x^{-1} = x^{-1} \circ x = 1_{\uparrow}. \quad (1.4)$$

Adopting the usual string calculus for strict monoidal categories, we represent τ , τ^{-1} , x , and more generally $x^{\circ a}$ for any $a \in \mathbb{Z}$, by the diagrams

$$\tau = \text{Diagram of } \tau, \quad \tau^{-1} = \text{Diagram of } \tau^{-1}, \quad x = \text{Diagram of } x, \quad x^{\circ a} = \text{Diagram of } x^{\circ a}. \quad (1.5)$$

Then the relations (1.1)–(1.3) are equivalent to the following diagrammatic relations:

$$\text{Diagram of } \tau = \text{Diagram of } \tau^{-1}, \quad \text{Diagram of } x = \text{Diagram of } x^{\circ 1}, \quad (1.6)$$

$$\text{Diagram of } \tau - \text{Diagram of } \tau = z \uparrow \uparrow, \quad (1.7)$$

$$\text{Diagram of } x = \text{Diagram of } x^{\circ a}, \quad \text{Diagram of } x^{\circ a} = \text{Diagram of } x^{\circ 1}. \quad (1.8)$$

The affine Hecke algebra AH_n itself may be identified with $\text{End}_{\mathcal{AH}(z)}(\uparrow^{\otimes n})$, with its standard generators x_i and τ_j coming from a dot on the i -th string and the positive crossing of the j -th and $(j+1)$ -th strings, respectively; our convention for this numbers strings $1, \dots, n$ from right to left. It is often convenient to assume (passing to a quadratic extension if necessary) that \mathbb{k} contains a root q of the quadratic equation $x^2 - zx - 1 = 0$, so that $z = q - q^{-1}$. The quadratic relation in AH_n may then be written as $(\tau_j - q)(\tau_j + q^{-1}) = 0$. Such a choice of parameter q is not needed in sections 2–4, but is essential for the applications in sections 5–10.

To obtain the quantum Heisenberg category $\mathcal{Heis}_k(z, t)$ from $\mathcal{AH}(z)$, we adjoin a right dual \downarrow to the object \uparrow , i.e., we add an additional generating object \downarrow and additional generating morphisms

$$c = \text{Diagram of } c: \mathbb{1} \rightarrow \downarrow \otimes \uparrow \quad \text{and} \quad d = \text{Diagram of } d: \uparrow \otimes \downarrow \rightarrow \mathbb{1}$$

subject to the relations

$$\text{Diagram of } c = \text{Diagram of } \downarrow, \quad \text{Diagram of } d = \text{Diagram of } \downarrow. \quad (1.9)$$

Then we add several more generating morphisms subject to relations which ensure that the resulting monoidal category is strictly pivotal, and moreover that there is a distinguished isomorphism

$$\uparrow \otimes \downarrow \cong \downarrow \otimes \uparrow \oplus \mathbb{1}^{\oplus k} \quad \text{if } k \geq 0$$

or

$$\uparrow \otimes \downarrow \oplus \mathbb{1}^{\oplus(-k)} \cong \downarrow \otimes \uparrow \text{ if } k \leq 0.$$

There are various equivalent ways to accomplish this in practice; see Sections 2–4. In these sections, we establish the equivalence of the three approaches and record many other useful relations which follow from the defining ones, including the property already mentioned that $\mathcal{H}eis_k(z, t)$ admits a strictly pivotal structure.

In this paragraph, we explain the approach from Section 4 in the special case $k = -1$. According to Definition 4.1 and (4.14), $\mathcal{H}eis_{-1}(z, t)$ is the strict \mathbb{k} -linear monoidal category generated by objects \uparrow , \downarrow and morphisms

$$\begin{array}{c} \nearrow \\ \times \end{array}, \quad \begin{array}{c} \nearrow \\ \times \end{array}, \quad \curvearrowleft, \quad \curvearrowright, \quad \curvearrowleft \quad \text{and} \quad \curvearrowright$$

subject to (1.7)–(1.9), the relations

$$\begin{array}{c} \text{Diagram} \\ \text{with } x \text{ and } t \\ \text{and } z \end{array} = \begin{array}{c} \text{Diagram} \\ \text{with } \uparrow \text{ and } \downarrow \end{array}, \quad \begin{array}{c} \text{Diagram} \\ \text{with } x \text{ and } t \\ \text{and } z \end{array} = \begin{array}{c} \text{Diagram} \\ \text{with } \uparrow \text{ and } \downarrow \end{array} + t z \begin{array}{c} \text{Diagram} \\ \text{with } \uparrow \text{ and } \downarrow \end{array}, \quad \begin{array}{c} \text{Diagram} \\ \text{with } x \text{ and } t \\ \text{and } z \end{array} = 0, \quad \begin{array}{c} \text{Diagram} \\ \text{with } x \text{ and } t \\ \text{and } z \end{array} = -t^{-1}z^{-1}1_{\mathbb{1}},$$

and one more relation, which is equivalent to (1.4). We have *not* included the generating morphism x since, due to a special feature of the $k = -1$ case, it can be recovered from the other generators via the formula

$$x = \begin{array}{c} \text{Diagram} \\ \text{with } x \text{ and } t \\ \text{and } z \end{array} := t \begin{array}{c} \text{Diagram} \\ \text{with } \uparrow \text{ and } \downarrow \end{array} - t^2 \begin{array}{c} \text{Diagram} \\ \text{with } \uparrow \text{ and } \downarrow \end{array}.$$

The relations in Definition 4.1 which involve x such as (1.6) are consequences of the other relations with one exception: we must still impose that x is invertible, that is, relation (1.4).

The deformed Heisenberg category $\mathcal{H}(q^2)$ introduced in [Licata and Savage 2013] is (the additive envelope of) the strict \mathbb{k} -linear monoidal category defined by the same presentation as in the previous paragraph, with the parameters satisfying $tz = -1$, but *without* the relation (1.4). This follows easily on comparing our presentation with the one in [loc. cit.], using also the fact that our category is strictly pivotal. The generator x denoted by a dot here is not the same as the morphism denoted by a dot in [loc. cit.] (that is simply equal to the right curl); instead, our dot is the “star dot” of [Cautis et al. 2018] (up to renormalization). The Hecke algebra generator $T = \begin{array}{c} \text{Diagram} \\ \text{with } x \text{ and } t \\ \text{and } z \end{array}$ from [Licata and Savage 2013, Definition 2.1] is related to our τ by $T = q\tau$ (so that the quadratic relation becomes $(T_j - q^2)(T_j + 1) = 0$). Also the generator X appearing just before [loc. cit., Lemma 3.8] is our $-x$. In fact, the category $\mathcal{H}(q^2)$ may be identified with the monoidal subcategory of our category $\mathcal{H}eis_{-1}(z, -z^{-1})$ consisting of all objects and all morphisms which do not involve negative powers of x .

For any \mathbb{k} -linear category \mathcal{C} , there is an associated strict \mathbb{k} -linear monoidal category $\mathcal{E}nd_{\mathbb{k}}(\mathcal{C})$ consisting of \mathbb{k} -linear endofunctors and natural transformations. Then one can consider “representations” of $\mathcal{H}eis_k(z, t)$ by considering \mathbb{k} -linear monoidal functors into $\mathcal{E}nd_{\mathbb{k}}(\mathcal{C})$ for different choices of \mathcal{C} . The motivation for the definition of $\mathcal{H}eis_k(z, t)$ comes from the fact that it acts in this way on other well-known categories

appearing in representation theory. If $k = 0$ and $t = q^n$ then $\mathcal{Heis}_k(z, t)$ acts on representations of $U_q(\mathfrak{gl}_n)$, with the generating objects \uparrow and \downarrow acting by tensoring with the natural $U_q(\mathfrak{gl}_n)$ -module and its dual, respectively; see [Section 5](#). This action is an extension of the monoidal functor from the HOMFLY-PT skein category to the category of finite-dimensional $U_q(\mathfrak{gl}_n)$ -modules constructed originally by Turaev [\[1989\]](#). If $k \neq 0$ then $\mathcal{Heis}_k(z, t)$ acts on representations of the cyclotomic Hecke algebras of level $|k|$ from [\[Ariki and Koike 1994\]](#), with \uparrow and \downarrow acting by induction and restriction functors if $k < 0$, or vice versa if $k > 0$; see [Section 6](#). When $k = -1$, this specializes to the action of the deformed Heisenberg category on modules over the usual (finite) Hecke algebras associated to the symmetric groups constructed already in [\[Licata and Savage 2013\]](#). The action of $\mathcal{Heis}_{-l}(z, t)$ on representations of cyclotomic Hecke algebras extends to an action on category \mathcal{O} over the rational Cherednik algebras of type $\mathfrak{S}_n \wr \mathbb{Z}/l$ for all $n \geq 0$, with \uparrow and \downarrow acting by certain Bezrukavnikov–Etingof induction and restriction functors from; [\[Bezrukavnikov and Etingof 2009\]](#); see [Section 7](#).

We also prove a basis theorem for the morphism spaces in $\mathcal{Heis}_k(z, t)$; see [Section 10](#) for the precise statement. In particular, our basis theorem implies that the *center* $\text{End}_{\mathcal{Heis}_k(z, t)}(\mathbb{1})$ of the quantum Heisenberg category is the tensor product $\text{Sym} \otimes \text{Sym}$ of *two* copies of the algebra of symmetric functions. In the degenerate case studied in [\[Brundan 2018\]](#), the basis theorem was proved by treating the cases $k = 0$ and $k \neq 0$ separately, appealing to results from [\[Brundan et al. 2017\]](#) and [\[Mackaay and Savage 2018\]](#); the proofs in [\[loc. cit.\]](#) ultimately exploited analogs of the categorical actions mentioned above, on representations of degenerate cyclotomic Hecke algebras and representations of $\mathfrak{gl}_n(\mathbb{C})$, respectively. In the quantum case, it is still possible to prove the basis theorem when $k = 0$ by such an argument, but for nonzero k the approach from [\[loc. cit.\]](#) seems to be unmanageable due to the larger center. Instead, we prove the basis theorem here by following the technique developed in the degenerate case in [\[Brundan et al. 2018, Theorem 6.4\]](#) (and earlier, in the context of Kac–Moody 2-categories, in [\[Webster 2018\]](#)). It depends crucially on the existence of an action of $\mathcal{Heis}_k(z, t)$ on a “sufficiently large” module category, which is obtained by choosing $l \gg 0$ then taking the tensor product of actions of $\mathcal{Heis}_{-l}(z, t)$ and $\mathcal{Heis}_{k+l}(z, 1)$ on representations of suitably generic cyclotomic Hecke algebras of levels l and $k+l$, respectively.

The construction of this categorical tensor product involves a remarkable monoidal functor from $\mathcal{Heis}_k(z, t)$ to a certain localization of the symmetric product

$$\mathcal{Heis}_l(z, u) \odot \mathcal{Heis}_m(z, v)$$

for $k = l+m$ and $t = uv$. This functor is defined in [Section 8](#) and is the quantum analog of the categorical comultiplication from [\[Brundan et al. 2018, Theorem 5.4\]](#). The particular tensor products exploited to prove the basis theorem are generic examples of *generalized cyclotomic quotients* of $\mathcal{Heis}_k(z, t)$; see [Section 9](#) for the general definition. In fact, these \mathbb{k} -linear categories first appeared in [\[Webster 2015, Proposition 5.6\]](#), but in a rather different form; the precise relationship between the categories of [\[loc. cit.\]](#) and the ones here will be explained in [\[Brundan et al. 2019\]](#).

We have stopped short of proving any results about the *decategoryification* of $\mathcal{Heis}_k(z, t)$ here, but let us make some remarks about this. There are two complementary points of view:

- One can consider the *Grothendieck ring* $K_0(\text{Kar}(\mathcal{H}eis_k(z, t)))$ of the additive Karoubi envelope of $\mathcal{H}eis_k(z, t)$. For generic z (i.e., when q is not a root of unity), we expect that this is isomorphic to a \mathbb{Z} -form for a central reduction of the universal enveloping algebra of the infinite-dimensional Heisenberg Lie algebra, just as was established in the degenerate case in [Brundan et al. 2018, Theorem 1.1]. However, there is a significant obstruction to proving this result in the quantum case: we do not know how to show that the split Grothendieck group $K_0(AH_n)$ of the affine Hecke algebra is isomorphic to that of the finite Hecke algebra.
- Alternatively, one can pass to the *trace* (or zeroth Hochschild homology). In [Cautis et al. 2018], this was computed already for the category $\mathcal{H}(q^2)$ of [Licata and Savage 2013], revealing an interesting connection to the elliptic Hall algebra. Using the basis theorem proved here, we expect it should be possible to extend the calculations made in [loc. cit.] to give a description of the trace of the full category $\mathcal{H}eis_k(z, t)$ for all $k \in \mathbb{Z}$.

In the main body of the article, proofs of all lemmas involving purely diagrammatic manipulations have been omitted. However, we have attempted to give enough details for the reader familiar with the analogous calculations in the degenerate case from [Brundan 2018, Section 2] and [Brundan et al. 2018, Section 5] to be able to reconstruct the proofs. Brundan, Savage and Webster [≥ 2020] are currently preparing a sequel in which we incorporate a (symmetric) Frobenius algebra into the definition of $\mathcal{H}eis_k(z, t)$, in a similar way to the Frobenius Heisenberg categories defined in the degenerate case in [Savage 2019]. We will include full proofs of all of the diagrammatic lemmas in the more general Frobenius setting in this sequel.

2. First approach

Before formulating our first definition of $\mathcal{H}eis_k(z, t)$, let us make some general remarks. We refer to the relation (1.7) as the *upward skein relation*. Rotating it through $\pm 90^\circ$ or 180° , one obtains three more skein relations; for example, here is the *leftward skein relation*

$$\begin{array}{c} \diagup \diagdown \\ \diagdown \diagup \end{array} - \begin{array}{c} \diagup \diagdown \\ \diagdown \diagup \end{array} = z \begin{array}{c} \curvearrowleft \\ \curvearrowright \end{array}. \quad (2.1)$$

At present, this has no meaning since we have not defined the leftward cups, caps or crossings which it involves! However, already in the monoidal category obtained from $\mathcal{AH}(z)$ by adjoining a right dual \downarrow to \uparrow as explained in the introduction, we can introduce the rightward crossings

$$\begin{array}{c} \diagup \diagdown \\ \diagdown \diagup \end{array} := \begin{array}{c} \curvearrowright \\ \curvearrowleft \end{array}, \quad \begin{array}{c} \diagup \diagdown \\ \diagdown \diagup \end{array} := \begin{array}{c} \curvearrowleft \\ \curvearrowright \end{array}, \quad (2.2)$$

and then we see that the rightward skein relation holds from (1.7). Rotating the two rightward crossings once more by a similar procedure, we obtain positive and negative downward crossings satisfying the downward skein relation. We also define the downward dot

$$y = \begin{array}{c} \downarrow \\ \uparrow \end{array} := \begin{array}{c} \curvearrowright \\ \curvearrowleft \end{array}. \quad (2.3)$$

It is immediate from these definitions and (1.9) that dots and crossings slide past rightward cups and caps

$$\begin{array}{c} \text{cup} \\ \text{dot} \end{array} = \begin{array}{c} \text{dot} \\ \text{cup} \end{array}, \quad \begin{array}{c} \text{cup} \\ \text{crossing} \end{array} = \begin{array}{c} \text{crossing} \\ \text{cup} \end{array}, \quad (2.4)$$

$$\begin{array}{c} \text{dot} \\ \text{cup} \end{array} = \begin{array}{c} \text{cup} \\ \text{dot} \end{array}, \quad \begin{array}{c} \text{dot} \\ \text{crossing} \end{array} = \begin{array}{c} \text{crossing} \\ \text{dot} \end{array}, \quad \begin{array}{c} \text{cup} \\ \text{crossing} \end{array} = \begin{array}{c} \text{crossing} \\ \text{cup} \end{array}, \quad (2.5)$$

$$\begin{array}{c} \text{cup} \\ \text{crossing} \end{array} = \begin{array}{c} \text{crossing} \\ \text{cup} \end{array}, \quad \begin{array}{c} \text{dot} \\ \text{crossing} \end{array} = \begin{array}{c} \text{crossing} \\ \text{dot} \end{array}, \quad \begin{array}{c} \text{cup} \\ \text{crossing} \end{array} = \begin{array}{c} \text{crossing} \\ \text{cup} \end{array}, \quad (2.6)$$

Also, the following relations are easily deduced by attaching rightward cups and caps to the relations in (1.8), then rotating the pictures using the definitions of the rightward/downward crossings:

$$\begin{array}{c} \text{cup} \\ \text{dot} \end{array} = \begin{array}{c} \text{dot} \\ \text{cup} \end{array}, \quad \begin{array}{c} \text{dot} \\ \text{crossing} \end{array} = \begin{array}{c} \text{crossing} \\ \text{dot} \end{array}, \quad \begin{array}{c} \text{cup} \\ \text{crossing} \end{array} = \begin{array}{c} \text{crossing} \\ \text{cup} \end{array}, \quad \begin{array}{c} \text{dot} \\ \text{crossing} \end{array} = \begin{array}{c} \text{crossing} \\ \text{dot} \end{array}. \quad (2.7)$$

The following lemma will be used repeatedly (often without reference). There are analogous dot slide relations for the rightward and downward crossings (obtained by rotation).

Lemma 2.1. *The following relations hold for $a \in \mathbb{Z}$:*

$$\begin{array}{c} \text{dot} \\ \text{crossing} \end{array}^a = \begin{cases} \text{dot}^a - z \sum_{\substack{b+c=a \\ b,c>0}} b \begin{array}{c} \text{dot} \\ \text{cup} \end{array} \begin{array}{c} \text{dot} \\ \text{cup} \end{array} & \text{if } a > 0 \\ \text{dot}^a + z \sum_{\substack{b+c=a \\ b,c\leq 0}} b \begin{array}{c} \text{dot} \\ \text{cup} \end{array} \begin{array}{c} \text{dot} \\ \text{cup} \end{array} & \text{if } a \leq 0; \end{cases} \quad \begin{array}{c} \text{dot} \\ \text{crossing} \end{array}^a = \begin{cases} \text{dot}^a - z \sum_{\substack{b+c=a \\ b,c\geq 0}} b \begin{array}{c} \text{dot} \\ \text{cup} \end{array} \begin{array}{c} \text{dot} \\ \text{cup} \end{array} & \text{if } a \geq 0, \\ \text{dot}^a + z \sum_{\substack{b+c=a \\ b,c< 0}} b \begin{array}{c} \text{dot} \\ \text{cup} \end{array} \begin{array}{c} \text{dot} \\ \text{cup} \end{array} & \text{if } a < 0; \end{cases} \quad (2.8)$$

$$\begin{array}{c} \text{dot} \\ \text{crossing} \end{array}^a = \begin{cases} \text{dot}^a + z \sum_{\substack{b+c=a \\ b,c>0}} b \begin{array}{c} \text{dot} \\ \text{cup} \end{array} \begin{array}{c} \text{dot} \\ \text{cup} \end{array} & \text{if } a > 0, \\ \text{dot}^a - z \sum_{\substack{b+c=a \\ b,c\leq 0}} b \begin{array}{c} \text{dot} \\ \text{cup} \end{array} \begin{array}{c} \text{dot} \\ \text{cup} \end{array} & \text{if } a \leq 0; \end{cases} \quad \begin{array}{c} \text{dot} \\ \text{crossing} \end{array}^a = \begin{cases} \text{dot}^a + z \sum_{\substack{b+c=a \\ b,c\geq 0}} b \begin{array}{c} \text{dot} \\ \text{cup} \end{array} \begin{array}{c} \text{dot} \\ \text{cup} \end{array} & \text{if } a \geq 0, \\ \text{dot}^a - z \sum_{\substack{b+c=a \\ b,c< 0}} b \begin{array}{c} \text{dot} \\ \text{cup} \end{array} \begin{array}{c} \text{dot} \\ \text{cup} \end{array} & \text{if } a < 0. \end{cases} \quad (2.9)$$

Now we can explain the first way to complete the definition of the quantum Heisenberg category following the scheme outlined in the introduction. The idea is to invert the morphism

$$\left\{ \begin{array}{c} \text{dot} \\ \text{crossing} \\ \text{cup} \\ \text{dot} \\ \vdots \\ k-1 \text{dot} \end{array} \right\} : \uparrow \otimes \downarrow \rightarrow \downarrow \otimes \uparrow \oplus \mathbb{1}^{\oplus k} \quad \text{if } k \geq 0, \quad (2.10)$$

$$\left[\begin{array}{c} \text{dot} \\ \text{crossing} \\ \text{cup} \\ \text{dot} \\ \vdots \\ k-1 \text{dot} \end{array} \right] : \uparrow \otimes \downarrow \oplus \mathbb{1}^{\oplus (-k)} \rightarrow \downarrow \otimes \uparrow \quad \text{if } k < 0,$$

in $\text{Add}(\mathcal{H}eis_k(z, t))$ (where Add denotes the additive envelope).

Definition 2.2. The *quantum Heisenberg category* $\mathcal{H}\text{eis}_k(z, t)$ is the strict \mathbb{k} -linear monoidal category obtained from $\mathcal{AH}(z)$ by adjoining a right dual \downarrow to \uparrow as explained in the introduction, together with the matrix entries of the following morphism which we declare to be a two-sided inverse to the morphism (2.10):

$$\left\{ \begin{array}{l} \left[\begin{array}{c|c|c|c} \text{Diagram 1} & \cdots & \text{Diagram } k-1 \\ \hline \text{Diagram } 0 & & & \end{array} \right] : \downarrow \otimes \uparrow \oplus \mathbb{1}^{\oplus k} \rightarrow \uparrow \otimes \downarrow \quad \text{if } k \geq 0, \\ \left[\begin{array}{c|c|c|c} \text{Diagram 1} & \cdots & \text{Diagram } -k-1 \\ \hline \text{Diagram } 0 & & & \end{array} \right] : \downarrow \otimes \uparrow \rightarrow \uparrow \otimes \downarrow \oplus \mathbb{1}^{\oplus (-k)} \quad \text{if } k < 0. \end{array} \right. \quad (2.11)$$

We impose one more essential relation:

$$\bigcirc = tz^{-1}1_{\mathbb{1}} \text{ if } k > 0, \quad \bigcirc = (tz^{-1} - t^{-1}z^{-1})1_{\mathbb{1}} \text{ if } k = 0, \quad \bigcirc_{-k} = tz^{-1}1_{\mathbb{1}} \text{ if } k < 0, \quad (2.12)$$

where the leftward cups and caps are defined by the formulas

$$\begin{array}{c} \text{Diagram 1} \\ \text{Diagram 0} \\ \text{Diagram } -k \end{array} := \begin{cases} -t^{-1}z^{-1} \text{ Diagram } k-1 & \text{if } k > 0, \\ t \text{ Diagram } 0 & \text{if } k = 0, \\ t^{-1} \text{ Diagram } -k & \text{if } k < 0; \end{cases} \quad \begin{array}{c} \text{Diagram 1} \\ \text{Diagram } 0 \end{array} := \begin{cases} t^k \text{ Diagram } 0 & \text{if } k \geq 0, \\ -t^{-1}z^{-1} \text{ Diagram } 0 & \text{if } k < 0. \end{cases} \quad (2.13)$$

To complete the definition, we introduce a few more shorthands for morphisms. We have already introduced one of the two leftward crossings; define the other one so that the leftward skein relation (2.1) holds. Also set

$$\text{Diagram } 0 := \text{Diagram } 0 + z \text{ Diagram } 1 \quad \text{if } k > 0, \quad \text{Diagram } a := \text{Diagram } a \quad \text{if } 0 < a < k, \quad (2.14)$$

$$\text{Diagram } 0 := \text{Diagram } 0 + z \text{ Diagram } 1 \quad \text{if } k < 0, \quad \text{Diagram } a := \text{Diagram } a \quad \text{if } 0 < a < -k. \quad (2.15)$$

Next, introduce the following *(+)-bubbles* assuming $a \leq 0$:

$$\text{Diagram } a := \begin{cases} -tz^{-1} \text{ Diagram } -a & \text{if } a > -k, \\ tz^{-1}1_{\mathbb{1}} & \text{if } a = -k, \\ 0 & \text{if } a < -k; \end{cases} \quad a\text{Diagram } a := \begin{cases} t^{-1}z^{-1} \text{ Diagram } -a & \text{if } a > k, \\ -t^{-1}z^{-1}1_{\mathbb{1}} & \text{if } a = k, \\ 0 & \text{if } a < k. \end{cases} \quad (2.16)$$

Finally, define the *(+)-bubbles* with label $a > 0$ to be the usual bubbles with a dots:

$$\text{Diagram } a := \text{Diagram } a, \quad a\text{Diagram } a := a\text{Diagram } a. \quad (2.17)$$

Then define $(-)$ -bubbles for all $a \in \mathbb{Z}$ by setting

$$\ominus^a := \bigcirc \circ^a - \oplus^a, \quad a\ominus := a \circ \circ - a \oplus. \quad (2.18)$$

In the case $k = 0$, the assertion that (2.10) and (2.11) are two-sided inverses means that

$$\begin{array}{c} \text{Diagram} \\ \text{if } k = 0, \end{array} \quad \begin{array}{c} \text{Diagram} \\ \text{if } k = 0. \end{array} \quad (2.19)$$

In fact, the defining relations for $\mathcal{Heis}_0(z, t)$ from [Definition 2.2](#) are exactly the same as the ones for the affine HOMFLY-PT skein category $\mathcal{AOS}(z, t)$ from [\[Brundan 2017, Theorem 1.1 and Section 4\]](#). Thus,

$$\mathcal{Heis}_0(z, t) = \mathcal{AOS}(z, t).$$

In this case, most of the other relations that we need have already been proved in [\[loc. cit.\]](#). However, the arguments there exploit a theorem of Turaev [\[1989, Lemma I.3.3\]](#) to establish all of the relations that do not involve dots; the approach described below reproves all of these relations in a way that is independent of Turaev's work.

When $k > 0$, the assertion that the morphisms (2.10) and (2.11) are two-sided inverses implies the following relations:

$$\begin{array}{c} \text{Diagram} \\ \text{if } k > 0, \end{array} \quad \begin{array}{c} \text{Diagram} \\ \text{if } k > 0, \end{array} \quad (2.20)$$

$$\begin{array}{c} \text{Diagram} \\ = 0 \text{ if } k > 0, \end{array} \quad \begin{array}{c} \text{Diagram} \\ = 0 \text{ if } 0 \leq a < k, \end{array} \quad \begin{array}{c} \text{Diagram} \\ = -\delta_{a,k} t^{-1} z^{-1} 1_{\mathbb{1}} \text{ if } 0 < a \leq k. \end{array} \quad (2.21)$$

To derive these relations, we multiplied the matrices (2.10) and (2.11) in both orders, then equated the result with the appropriate identity matrix. The following useful relation is an easy exercise at this point; one needs to use (2.8), (2.12), (2.13) and (2.21):

$$a \circ \begin{array}{c} \text{Diagram} \\ = \delta_{a,0} t \text{ if } 0 \leq a \leq k. \end{array} \quad (2.22)$$

Finally, when $k < 0$, we will need the following relations which are deduced from (2.10) and (2.11) by the same argument as explained in the previous paragraph

$$\begin{array}{c} \text{Diagram} \\ = \begin{array}{c} \text{Diagram} \\ - \sum_{a=0}^{k-1} \begin{array}{c} \text{Diagram} \\ \text{if } k < 0, \end{array} \end{array} \end{array} \quad (2.23)$$

$$\begin{array}{c} \text{Diagram} \\ = 0 \text{ if } k < 0, \end{array} \quad \begin{array}{c} \text{Diagram} \\ = 0 \text{ if } 0 \leq a < -k, \end{array} \quad \begin{array}{c} \text{Diagram} \\ = -\delta_{a,0} t^{-1} z^{-1} 1_{\mathbb{1}} \text{ if } 0 \leq a < -k. \end{array} \quad (2.24)$$

Now we are going to consider the counterpart of the morphism (2.10) defined using the negative instead of positive rightward crossing:

$$\left\{ \begin{array}{c} \left[\begin{array}{c} \times \\ \curvearrowleft \\ \circlearrowleft \\ \vdots \\ k-1 \circlearrowleft \end{array} \right] : \uparrow \otimes \downarrow \rightarrow \downarrow \otimes \uparrow \oplus \mathbb{1}^{\oplus k} \\ \left[\begin{array}{c} \times \\ \curvearrowleft \\ \circlearrowleft \\ \vdots \\ k-1 \circlearrowleft \\ \curvearrowleft \circlearrowleft \cdots \circlearrowleft \end{array} \right] : \uparrow \otimes \downarrow \oplus \mathbb{1}^{\oplus(-k)} \rightarrow \downarrow \otimes \uparrow \end{array} \right. \begin{array}{l} \text{if } k > 0, \\ \text{if } k \leq 0. \end{array} \quad (2.25)$$

Lemma 2.3. *The morphism (2.25) is invertible with two-sided inverse*

$$\left\{ \begin{array}{c} \left[\begin{array}{c} \times \\ \curvearrowleft \\ \circlearrowleft \\ \vdots \\ -k-1 \end{array} \right] : \downarrow \otimes \uparrow \oplus \mathbb{1}^{\oplus k} \rightarrow \uparrow \otimes \downarrow \quad \text{if } k > 0, \\ \left[\begin{array}{c} \times \\ \curvearrowleft \\ \circlearrowleft \\ \vdots \\ -k-1 \\ \curvearrowleft \end{array} \right] : \downarrow \otimes \uparrow \rightarrow \uparrow \otimes \downarrow \oplus \mathbb{1}^{\oplus(-k)} \quad \text{if } k \leq 0. \end{array} \right. \quad (2.26)$$

Moreover, we have that

$$k \circlearrowleft = -t^{-1}z^{-1}1_{\mathbb{1}} \text{ if } k > 0, \quad \circlearrowleft = (tz^{-1} - t^{-1}z^{-1})1_{\mathbb{1}} \text{ if } k = 0, \quad \circlearrowleft = -t^{-1}z^{-1}1_{\mathbb{1}} \text{ if } k < 0, \quad (2.27)$$

$$\curvearrowleft = \begin{cases} tz^{-1} \circlearrowleft & \text{if } k > 0, \\ t^{-1} \circlearrowleft & \text{if } k \leq 0, \end{cases} \quad \curvearrowright = \begin{cases} t^k \circlearrowright & \text{if } k > 0, \\ t^{-1} \circlearrowright & \text{if } k = 0, \\ tz^{-1} \circlearrowright & \text{if } k < 0. \end{cases} \quad (2.28)$$

3. Second approach

Our second presentation for $\mathcal{H}eis_k(z, t)$ is very similar to the first presentation, but we invert the morphism (2.25) instead of (2.10).

Definition 3.1. The *quantum Heisenberg category* $\mathcal{H}eis_k(z, t)$ is the strict \mathbb{k} -linear monoidal category obtained from $\mathcal{AH}(z)$ by adjoining a right dual \downarrow to \uparrow as explained in the introduction, together with the matrix entries of the morphism (2.26), which we declare to be a two-sided inverse to (2.25). In addition, we impose the relation (2.27) for the leftward cups and caps which are defined in this approach from

(2.28). Define the other leftward crossing, i.e., the one which does not appear in (2.26), so the leftward skein relation (2.1) holds. Also set

$$\begin{array}{ll} \text{Diagram 1} := \text{Diagram 2} - z \text{Diagram 3} & \text{if } k > 0, \\ \text{Diagram 1} := \text{Diagram 2} & \text{if } 0 < a < k, \end{array} \quad (3.1)$$

$$\begin{array}{ll} \text{Diagram 1} := \text{Diagram 2} - z \text{Diagram 3} & \text{if } k < 0, \\ \text{Diagram 1} := \text{Diagram 2} & \text{if } 0 < a < -k. \end{array} \quad (3.2)$$

Finally define the (+)- and (-)-bubbles from (2.16)–(2.18) as before.

Theorem 3.2. *Definitions 2.2 and 3.1 give two different presentations for the same monoidal category, with all of the named morphisms introduced in the two definitions being the same. Moreover, there is a unique isomorphism of \mathbb{k} -linear monoidal categories*

$$\Omega_k : \mathcal{H}eis_k(z, t) \rightarrow \mathcal{H}eis_{-k}(z, t^{-1})^{\text{op}} \quad (3.3)$$

sending

$$\begin{array}{ll} \text{Diagram 1} \mapsto \text{Diagram 1}, & \text{Diagram 2} \mapsto -\text{Diagram 2}, \\ \text{Diagram 3} \mapsto \text{Diagram 3}, & \text{Diagram 4} \mapsto \text{Diagram 4}. \end{array}$$

The effect of Ω_k on the other morphisms is as follows:

$$\begin{array}{llll} \text{Diagram 1} \mapsto \text{Diagram 1}, & \text{Diagram 2} \mapsto -\text{Diagram 2}, & \text{Diagram 3} \mapsto -\text{Diagram 3}, & \text{Diagram 4} \mapsto -\text{Diagram 4}, \\ \text{Diagram 5} \mapsto -\text{Diagram 5}, & \text{Diagram 6} \mapsto -\text{Diagram 6}, & \text{Diagram 7} \mapsto -\text{Diagram 7}, & \text{Diagram 8} \mapsto -\text{Diagram 8}, \\ \text{Diagram 9} \mapsto \text{Diagram 9}, & \text{Diagram 10} \mapsto \text{Diagram 10}, & \text{Diagram 11} \mapsto \text{Diagram 11}, & \text{Diagram 12} \mapsto \text{Diagram 12}, \\ \text{Diagram 13} \mapsto -\text{Diagram 13}, & \text{Diagram 14} \mapsto -\text{Diagram 14}, & \text{Diagram 15} \mapsto -\text{Diagram 15}, & \text{Diagram 16} \mapsto -\text{Diagram 16}. \end{array}$$

Proof. To avoid confusion, denote the category $\mathcal{H}eis_k(z, t)$ from Definition 2.2 by $\mathcal{H}eis_k^{\text{old}}(z, t)$ and the one from Definition 3.1 by $\mathcal{H}eis_k^{\text{new}}(z, t)$. The relations and other definitions for the category $\mathcal{H}eis_k^{\text{new}}(z, t)$ in Definition 3.1 and the ones for $\mathcal{H}eis_{-k}^{\text{old}}(z, t^{-1})$ from Definition 2.2 are related by reflecting all diagrams in a horizontal plane and multiplying by $(-1)^{x+y}$, where x is the number of crossings and y is the number of leftward cups and caps (including leftward cups and caps in (+)- and (-)-bubbles but not ones labeled by \diamond or \heartsuit). It follows that there are mutually inverse isomorphisms

$$\mathcal{H}eis_{-k}^{\text{old}}(z, t^{-1}) \xrightleftharpoons[\Omega_+]{\Omega_-} \mathcal{H}eis_k^{\text{new}}(z, t)^{\text{op}}$$

both defined in the same way as the functor Ω_k in the statement of the theorem. Now we apply Lemma 2.3 and Definition 3.1 to construct a strict \mathbb{k} -linear monoidal functor

$$\Theta_k : \mathcal{H}eis_k^{\text{new}}(z, t) \rightarrow \mathcal{H}eis_k^{\text{old}}(z, t)$$

which is the identity on diagrams. This functor is an isomorphism because it has a two-sided inverse, namely, $\Omega_+ \circ \Theta_{-k} \circ \Omega_-$. Thus, using Θ_k , we may identify $\mathcal{H}eis_k^{\text{new}}(z, t)$ and $\mathcal{H}eis_k^{\text{old}}(z, t)$. Finally, $\Omega_k := \Omega_+$ gives the required symmetry. \square

In the remainder of the section, we record some further consequences of the defining relations, thereby showing that $\mathcal{H}eis_k(z, t)$ is strictly pivotal. The first lemma explains how dots slide past leftward cups, caps and crossings. Its generalization to dots with arbitrary multiplicities $n \in \mathbb{Z}$ may also be deduced using induction and the leftward skein relation like in [Lemma 2.1](#).

Lemma 3.3. *The following relations hold:*

$$\text{Diagram 1} = \text{Diagram 2}, \quad (3.4)$$

$$\text{Diagram 3} = \text{Diagram 4}, \quad (3.5)$$

Let Sym be the algebra of symmetric functions over \mathbb{k} . This is an infinite rank polynomial algebra with two sets of algebraically independent generators, namely, the *elementary symmetric functions* e_1, e_2, \dots and the *complete symmetric functions* h_1, h_2, \dots . Adopting the convention that $e_n = h_n := \delta_{n,0}$ for $n \leq 0$, the elementary and complete symmetric functions are related by the following well-known identity [[Macdonald 1995](#), (I.2.6)]:

$$\sum_{r+s=n} (-1)^s e_r h_s = \delta_{n,0}. \quad (3.6)$$

The following lemma, which we may refer to as the *infinite Grassmannian relation* (following Lauda), shows that there is a well-defined homomorphism

$$\beta : \text{Sym} \otimes \text{Sym} \rightarrow \text{End}_{\mathcal{H}eis_k(z, t)}(\mathbb{1}) \quad (3.7)$$

such that

$$h_n \otimes 1 \mapsto (-1)^{n-1} t z^{n+k} \text{Diagram 1}, \quad 1 \otimes h_n \mapsto (-1)^n t^{-1} z^{-n} \text{Diagram 2}, \quad (3.8)$$

$$e_n \otimes 1 \mapsto t^{-1} z \text{Diagram 3}, \quad 1 \otimes e_n \mapsto -t z \text{Diagram 4}. \quad (3.9)$$

We will prove in [Corollary 10.2](#) that β is actually an *isomorphism*.

Lemma 3.4. *For any $a \in \mathbb{Z}$, we have that*

$$\sum_{\substack{b,c \in \mathbb{Z} \\ b+c=a}} \text{Diagram 1} = \sum_{\substack{b,c \in \mathbb{Z} \\ b+c=a}} \text{Diagram 2} = -\delta_{a,0} z^{-2} \mathbb{1}_{\mathbb{1}}. \quad (3.10)$$

Moreover

$$\text{Diagram 1}^a = \delta_{a,-k} t z^{-1} \mathbb{1}_{\mathbb{1}} \quad \text{if } a \leq -k, \quad a \text{Diagram 1} = -\delta_{a,k} t^{-1} z^{-1} \mathbb{1}_{\mathbb{1}} \quad \text{if } a \leq k, \quad (3.11)$$

$$a \text{Diagram 2} = \delta_{a,0} t z^{-1} \mathbb{1}_{\mathbb{1}} \quad \text{if } a \geq 0, \quad \text{Diagram 2}^a = -\delta_{a,0} t^{-1} z^{-1} \mathbb{1}_{\mathbb{1}} \quad \text{if } a \geq 0. \quad (3.12)$$

Corollary 3.5. *For an indeterminate w , we have that*

$$\oplus(w)\oplus(w) = \ominus(w)\ominus(w) = 1_{\mathbb{1}}, \quad (3.13)$$

where

$$\oplus(w) := t^{-1}z \sum_{n \in \mathbb{Z}} \oplus_n w^{-n} \in w^k 1_{\mathbb{1}} + w^{k-1} \text{End}_{\mathcal{H}eisk(z,t)}(\mathbb{1})[[w^{-1}]], \quad (3.14)$$

$$\oplus(w) := -tz \sum_{n \in \mathbb{Z}} n \oplus_n w^{-n} \in w^{-k} 1_{\mathbb{1}} + w^{-k-1} \text{End}_{\mathcal{H}eisk(z,t)}(\mathbb{1})[[w^{-1}]], \quad (3.15)$$

$$\ominus(w) := -tz \sum_{n \in \mathbb{Z}} \ominus_n w^{-n} \in 1_{\mathbb{1}} + w \text{End}_{\mathcal{H}eisk(z,t)}(\mathbb{1})[[w]], \quad (3.16)$$

$$\ominus(w) := t^{-1}z \sum_{n \in \mathbb{Z}} n \ominus_n w^{-n} \in 1_{\mathbb{1}} + w \text{End}_{\mathcal{H}eisk(z,t)}(\mathbb{1})[[w]]. \quad (3.17)$$

Using the next relations plus (2.14) and (3.2), the leftward cups and caps decorated by \diamond or \heartsuit can be eliminated from any diagram.

Lemma 3.6. *The following relations hold:*

$$\begin{array}{c} \uparrow \\ \curvearrowleft \\ a \end{array} = -z^2 \sum_{b \geq 1} b \begin{array}{c} \uparrow \\ \curvearrowleft \\ \end{array} \oplus_{-a-b} \quad \text{if } 0 \leq a < k, \quad (3.18)$$

$$\begin{array}{c} a \\ \curvearrowleft \\ \uparrow \end{array} = -z^2 \sum_{b \geq 1} \begin{array}{c} \curvearrowleft \\ \uparrow \\ b \end{array} \ominus_{-a-b} \oplus \quad \text{if } 0 \leq a < -k. \quad (3.19)$$

The next lemma shows that \downarrow is left dual to \uparrow (as well as being right dual by the original construction). Thus, the monoidal category $\mathcal{H}eisk(z, t)$ is rigid.

Lemma 3.7. *The following relations hold:*

$$\begin{array}{c} \uparrow \\ \curvearrowleft \end{array} = \uparrow, \quad \begin{array}{c} \curvearrowleft \\ \uparrow \end{array} = \downarrow. \quad (3.20)$$

The final lemma together with (3.4) implies that $\mathcal{H}eisk(z, t)$ is strictly pivotal, with duality functor

$$*: \mathcal{H}eisk(z, t) \xrightarrow{\sim} (\mathcal{H}eisk(z, t)^{\text{op}})^{\text{rev}} \quad (3.21)$$

defined on morphisms by rotating diagrams through 180° .

Lemma 3.8. *The following relations hold:*

$$\begin{array}{c} \curvearrowleft \\ \curvearrowleft \end{array} = \begin{array}{c} \curvearrowleft \\ \uparrow \end{array}, \quad \begin{array}{c} \curvearrowleft \\ \uparrow \end{array} = \begin{array}{c} \curvearrowleft \\ \curvearrowleft \end{array}, \quad \begin{array}{c} \curvearrowleft \\ \uparrow \end{array} = \begin{array}{c} \uparrow \\ \curvearrowleft \end{array}, \quad \begin{array}{c} \curvearrowleft \\ \uparrow \end{array} = \begin{array}{c} \curvearrowleft \\ \curvearrowleft \end{array}, \quad (3.22)$$

$$\begin{array}{c} \curvearrowleft \\ \curvearrowleft \end{array} = \begin{array}{c} \curvearrowleft \\ \downarrow \end{array}, \quad \begin{array}{c} \curvearrowleft \\ \downarrow \end{array} = \begin{array}{c} \curvearrowleft \\ \curvearrowleft \end{array}, \quad \begin{array}{c} \curvearrowleft \\ \downarrow \end{array} = \begin{array}{c} \downarrow \\ \curvearrowleft \end{array}, \quad \begin{array}{c} \curvearrowleft \\ \downarrow \end{array} = \begin{array}{c} \curvearrowleft \\ \curvearrowleft \end{array}. \quad (3.23)$$

4. Third approach

Now we have enough relations in hand to formulate our third presentation for $\mathcal{H}eis_k(z, t)$. This presentation does not involve any leftward cups or caps decorated by \diamond or \heartsuit ; [Lemma 3.6](#) showed already that these are redundant as generators.

Definition 4.1. The *quantum Heisenberg category* $\mathcal{H}eis_k(z, t)$ is the strict \mathbb{k} -linear monoidal category obtained from $\mathcal{AH}(z)$ by adjoining a right dual \downarrow to \uparrow as explained in the introduction, plus two more generating morphisms \curvearrowleft and \curvearrowright subject to the following additional relations:

$$\curvearrowleft = \left| \begin{array}{c} \uparrow \curvearrowright \\ \downarrow \end{array} \right| - t^{-1}z \curvearrowright \curvearrowleft + z^2 \sum_{a,b>0} \begin{array}{c} a \curvearrowleft \\ b \curvearrowright \end{array} \oplus_{-a-b}, \quad (4.1)$$

$$\curvearrowright = \left| \begin{array}{c} \uparrow \curvearrowleft \\ \downarrow \end{array} \right| + tz \curvearrowleft \curvearrowright + z^2 \sum_{a,b>0} \begin{array}{c} -a-b \oplus \\ a \curvearrowright \\ b \curvearrowleft \end{array}, \quad (4.2)$$

$$\begin{array}{c} \uparrow \\ \curvearrowleft \end{array} = \delta_{k,0} t^{-1} \quad \text{if } k \geq 0, \quad \begin{array}{c} a+k \curvearrowleft \\ \oplus \end{array} = (\delta_{a,-k} t z^{-1} - \delta_{a,0} t^{-1} z^{-1}) 1_{\mathbb{1}} \quad \text{if } -k \leq a \leq 0, \quad (4.3)$$

$$\begin{array}{c} \uparrow \\ \curvearrowright \end{array} = \delta_{k,0} t \quad \text{if } k \leq 0, \quad \begin{array}{c} \curvearrowright \\ \oplus_{a-k} \end{array} = (\delta_{a,0} t z^{-1} - \delta_{a,k} t^{-1} z^{-1}) 1_{\mathbb{1}} \quad \text{if } k \leq a \leq 0. \quad (4.4)$$

Here, we have used the leftward crossings which are defined in this approach by

$$\begin{array}{c} \uparrow \\ \curvearrowleft \end{array} := \begin{array}{c} \uparrow \curvearrowright \\ \downarrow \end{array}, \quad \begin{array}{c} \uparrow \\ \curvearrowright \end{array} := \begin{array}{c} \uparrow \curvearrowleft \\ \downarrow \end{array}, \quad (4.5)$$

and the (+)-bubbles which are defined for $a \leq k$ or $a \leq -k$, respectively, by

$$\begin{array}{c} \oplus_{a-k} \\ a+k \end{array} := t^{a+1} z^{a-1} \det(\begin{array}{c} \curvearrowleft \\ k+i-j+1 \end{array} \curvearrowright)_{i,j=1,\dots,a}, \quad (4.6)$$

$$\begin{array}{c} \oplus \\ a+k \end{array} := -t^{-a-1} z^{a-1} \det(-\begin{array}{c} \curvearrowright \\ a-k+i-j+1 \end{array})_{i,j=1,\dots,a}, \quad (4.7)$$

interpreting the determinants as $\delta_{a,0}$ in case $a \leq 0$. Finally, define the (+)-bubbles with label $a > 0$ to be the usual bubbles with a dots as in [\(2.17\)](#), then define the (-)-bubbles for all $a \in \mathbb{Z}$ so that [\(2.18\)](#) holds.

Before proving the equivalence of this definition with the earlier ones, we make some remarks about the relations [\(4.1\)–\(4.7\)](#). If $k \leq 1$, the relation [\(4.1\)](#) is equivalent to

$$\curvearrowleft = \left| \begin{array}{c} \uparrow \curvearrowright \\ \downarrow \end{array} \right| - t^{-1}z \curvearrowright \curvearrowleft. \quad (4.8)$$

This follows immediately from the definition of the (+)-bubbles from (4.6). Similarly, when $k \geq -1$, the relation (4.2) is equivalent to

$$\text{Diagram} = \left| \begin{array}{c} \uparrow \\ +tz \end{array} \right| \text{.} \quad (4.9)$$

Here are some other useful consequences of these relations:

$$\text{Diagram} = \delta_{k,0} t^{-1} \text{Diagram} \quad \text{if } k \geq 0, \quad \text{Diagram} = t \text{Diagram} \quad \text{if } k \geq 0, \quad (4.10)$$

$$\text{Diagram} = \delta_{k,0} t \text{Diagram} \quad \text{if } k \leq 0, \quad \text{Diagram} = t^{-1} \text{Diagram} \quad \text{if } k \leq 0, \quad (4.11)$$

$$\text{Diagram} = \delta_{k,0} t^{-1} \text{Diagram} \quad \text{if } k \geq 0, \quad \text{Diagram} = t \text{Diagram} \quad \text{if } k \geq 0, \quad (4.12)$$

$$\text{Diagram} = \delta_{k,0} t \text{Diagram} \quad \text{if } k \leq 0, \quad \text{Diagram} = t^{-1} \text{Diagram} \quad \text{if } k \leq 0. \quad (4.13)$$

These follow from (4.3)–(4.4) on expanding the definitions of the sideways crossings. Then, using (4.13) and the leftward skein relation to convert the negative crossings in (4.8) to positive ones, relation (4.8) can be further simplified in case that $k < 0$: it is equivalent to

$$\text{Diagram} = \left| \begin{array}{c} \uparrow \\ \downarrow \end{array} \right| \text{.} \quad (4.14)$$

Similarly, (4.9) is equivalent to the following when $k > 0$:

$$\text{Diagram} = \left| \begin{array}{c} \uparrow \\ \downarrow \end{array} \right| \text{.} \quad (4.15)$$

Finally, when $k = 0$, the relations (4.8)–(4.9) together are equivalent to the single assertion

$$\text{Diagram} = \left(\text{Diagram} \right)^{-1}, \quad (4.16)$$

i.e., both of the relations from (2.19).

Theorem 4.2. *The category $\mathcal{H}eis_k(z, t)$ defined by Definition 4.1 is the same as the one from Definitions 2.2 and 3.1, with all morphisms introduced in the third definition being the same as the ones from before.*

Proof. To avoid confusion in the proof, we denote the category from the equivalent Definitions 2.2 and 3.1 by $\mathcal{H}eis_k^{\text{old}}(z, t)$, and the one from Definition 4.1 by $\mathcal{H}eis_k^{\text{new}}(z, t)$. From the evident symmetry in the relations (4.1)–(4.7), it follows that there is an isomorphism

$$\Omega_k : \mathcal{H}eis_k^{\text{new}}(z, t) \rightarrow \mathcal{H}eis_{-k}^{\text{new}}(z, t^{-1})^{\text{op}}$$

which reflects diagrams in a horizontal plane and multiplies by $(-1)^{x+y}$ where x is the number of crossings and y is the number of leftward cups and caps. Combining this with (3.3), we are reduced to proving the theorem under the assumption that $k \leq 0$.

We first check that all of the defining relations (4.1)–(4.7) of $\mathcal{H}eis_k^{\text{new}}(z, t)$ are satisfied in $\mathcal{H}eis_k^{\text{old}}(z, t)$, so that there is a strict \mathbb{k} -linear monoidal functor

$$\Theta : \mathcal{H}eis_k^{\text{new}}(z, t) \rightarrow \mathcal{H}eis_k^{\text{old}}(z, t)$$

which is the identity on diagrams. For this, note to start with that (4.5) holds in $\mathcal{H}eis_k^{\text{old}}(z, t)$ as we have shown that the latter category is strictly pivotal. The relation (4.6) is almost trivial when $k \leq 0$ and holds thanks to (3.11). For (4.7), the identity holds if $a - k \leq 0$ due again to (3.11), so assume that $a - k > 0$. Then the desired identity is the image under the homomorphism β from (3.7) of the identity

$$(-1)^{a-k-1} t^{-1} z^{-1} h_{a-k} \otimes 1 = -z^{a-k-1} t^{-a+k-1} \det(-tz^{-1} e_{i-j+1} \otimes 1)_{i,j=1,\dots,a-k}$$

in $\text{Sym} \otimes \text{Sym}$. This follows from the well-known identity $h_n = \det(e_{i-j+1})_{i,j=1,\dots,n}$; see [Macdonald 1995, Exercise I.2.8]. It remains to check the relations (4.1)–(4.4). For (4.1)–(4.2) when $k = 0$, we just need to check the equivalent form (4.16), which follows by (2.19). For (4.1) when $k < 0$, we check the equivalent form (4.14), which holds due to the second relation from (2.23). For (4.2) when $k < 0$, we use the first relation from (2.23), expanding the leftward caps decorated by \heartsuit using (2.13) when $a = 0$ or (2.15) and (3.19) when $a > 0$. Finally, the relations (4.3)–(4.4) follow easily from (2.24), (2.12)–(2.13) and (2.27)–(2.28).

Now we want to show that Θ is an isomorphism. We do this by using the presentation from Definition 2.2 to construct a two-sided inverse

$$\Phi : \mathcal{H}eis_k^{\text{old}}(z, t) \rightarrow \mathcal{H}eis_k^{\text{new}}(z, t),$$

still assuming that $k \leq 0$. We define Φ on morphisms by declaring that it takes the rightward cup, the rightward cap, and all dots and crossings (with any orientation) to the corresponding morphisms in $\mathcal{H}eis_k^{\text{new}}(z, t)$, and also

$$\Phi \left(\begin{array}{c} 0 \\ \downarrow \heartsuit \end{array} \right) := -tz \curvearrowleft \quad \text{if } k < 0, \quad \Phi \left(\begin{array}{c} a \\ \downarrow \heartsuit \end{array} \right) := -z^2 \sum_{b \geq 1} \curvearrowleft \circ b \quad -a-b \bigoplus \quad \text{if } 0 < a < -k.$$

To see that Φ is well defined, we must verify the relations from Definition 2.2. For (2.12), we must check the following in $\mathcal{H}eis_k^{\text{new}}(z, t)$:

$$t \circlearrowleft = (tz^{-1} - t^{-1}z^{-1})1_{\mathbb{1}} \quad \text{if } k = 0, \quad \circlearrowleft^{-k} = tz^{-1}1_{\mathbb{1}} \quad \text{if } k < 0.$$

These follow from (4.4) and (4.12). Then the main work is to show that the images under Φ of the morphisms (2.10) and (2.11) are two-sided inverses in $\mathcal{H}eis_k^{\text{new}}(z, t)$. When $k = 0$, this is immediate from

(4.16), so suppose that $k < 0$. The images under Φ of the two equations in (2.23) are precisely the known relations (4.2) and (4.14). We are left with checking that the images under Φ of the relations

$$\text{Diagram: } \text{A crossing with a dot on the top-left arc and a dot on the bottom-right arc, labeled } a. \quad \text{Diagram: } \text{A crossing with a dot on the top arc and a dot on the bottom arc, labeled } b. \quad \text{Diagram: } \text{A crossing with a dot on the top-left arc and a dot on the bottom-left arc, labeled } a, \text{ with a dot on the top arc labeled } b. \quad \text{Equation: } \text{Diagram} = 0, \quad \text{Diagram} = 0, \quad \text{Diagram} = \delta_{a,b} 1_{\mathbb{1}}$$

hold in $\mathcal{H}eis_k^{\text{new}}(z, t)$ for $0 \leq a, b < -k$. The first of these when $a = 0$ follows by (4.13). To see it for $0 < a < -k$, we first apply the leftward skein relation, then slide the dots past the crossing using the leftward analog of (2.9) which may be deduced from the definition (4.5), and finally appeal to (4.4). The second and third relations follow from (4.11) and (4.4) in the case that $b = 0$. To prove them when $0 < b < -k$, we must show that

$$\sum_{c \geq 1} \text{Diagram} = 0, \quad \sum_{c \geq 1} \text{Diagram} = -\delta_{a,b} z^{-2} 1_{\mathbb{1}}$$

in $\mathcal{H}eis_k^{\text{new}}(z, t)$. For the first identity, it is zero if $b \geq -k$ as the (+)-bubble vanishes by (1.3). To see it for $0 < b < -k$, use the skein relation, commute the dots past the crossing, then appeal to (4.4) and (4.11). For the second identity, define a homomorphism $\gamma : \text{Sym} \rightarrow \text{End}_{\mathcal{H}eis_k^{\text{new}}(z, t)}(\mathbb{1})$ by sending $e_n \mapsto t^{-1} z \text{Diagram}$ for $n \geq 0$. Using $h_n = \det(e_{i-j+1})_{i,j=1,\dots,n}$ and (4.7), it follows that γ sends $h_n \mapsto (-1)^{n-1} t z \text{Diagram}$ for $n \leq -k$. Then the identity we are trying to prove follows by applying γ to the identity $\sum_{c \geq 1} (-1)^{-k-b-c} e_{k+a+c} h_{-k-b-c} = \delta_{a,b}$, which is (3.6).

To complete the proof, we must show that Θ and Φ are indeed two-sided inverses. To check that $\Theta \circ \Phi = \text{Id}$, the only difficulty is to see that

$$\Theta(\Phi(\text{Diagram})) = \text{Diagram}.$$

When $a = 0$, this is immediate from (2.13), while if $0 < a < -k$ it follows from (2.15) and (3.19). To check that $\Phi \circ \Theta = \text{Id}$, the only difficulty is to see that

$$\Phi(\text{Diagram}) = \text{Diagram}, \quad \Phi(\text{Diagram}) = \text{Diagram}.$$

These follow from (2.13) and (4.12)–(4.13). \square

Lemma 4.3. *Suppose that \mathcal{C} is a strict \mathbb{k} -linear monoidal category containing objects \uparrow and \downarrow and morphisms $\hat{\phi}$, $\hat{\times}$, $\hat{\times}$, $\hat{\cup}$ and $\hat{\cap}$ satisfying (1.6)–(1.9). Then \mathcal{C} contains at most one pair of morphisms $\hat{\cup}$ and $\hat{\cap}$ which satisfy (4.1)–(4.4) (for the sideways crossings and the (+)-bubbles defined via (2.2) and (4.5)–(4.7)).*

Proof. If $k \leq 0$, Theorem 4.2 implies that the morphism (2.10) is invertible in \mathcal{C} , and $\hat{\times}$ is the $(1, 1)$ -entry of the inverse matrix. This property characterizes $\hat{\times}$ uniquely as a morphism in \mathcal{C} when $k \leq 0$, independent of the choices of $\hat{\cup}$ or $\hat{\cap}$. Similarly, when $k \geq 0$, the morphism (2.25) is invertible in \mathcal{C} , and $\hat{\times}$ is the $(1, 1)$ -entry of the inverse matrix. Thus $\hat{\times}$ is characterized uniquely when $k \leq 0$. To complete the proof when $k = 0$, it remains to use (4.12)–(4.13), since these show how to express $\hat{\cup}$ and

\curvearrowleft in terms of \curvearrowleft and \curvearrowright and the two leftward crossings. To complete the proof when $k < 0$, we note instead that the $(2, 1)$ -entry of the inverse of (2.10) is $-tz\curvearrowleft$, so \curvearrowleft is uniquely determined in \mathcal{C} . Then \curvearrowright may be recovered uniquely using the relation (2.13) and our knowledge of \times . Finally when $k > 0$, the $(1, 2)$ -entry of the inverse of (2.25) gives $t^{-1}z\curvearrowright$ and then \curvearrowright may be recovered using (2.28) and our knowledge of \times . \square

To conclude the section, we formulate three more important sets of relations. The first of these explains how to expand *curls*. It is quite surprising that we have never needed to simplify left curls when $k > 0$ (or right curls when $k < 0$) before this point.

Lemma 4.4. *The following relations hold for any $a \in \mathbb{Z}$:*

$$a \circlearrowleft = z \sum_{b \geq 0} \oplus_{a-b} \circlearrowleft_b - z \sum_{b < 0} \ominus_{a-b} \circlearrowleft_b, \quad (4.17)$$

$$a \circlearrowright = z \sum_{b > 0} \oplus_{a-b} \circlearrowright_b - z \sum_{b \leq 0} \ominus_{a-b} \circlearrowright_b, \quad (4.18)$$

$$\circlearrowleft \circlearrowleft_a = z \sum_{b \leq 0} b \circlearrowleft_{a-b} \ominus - z \sum_{b > 0} b \circlearrowleft_{a-b} \oplus, \quad (4.19)$$

$$\circlearrowright \circlearrowright_a = z \sum_{b < 0} b \circlearrowright_{a-b} \ominus - z \sum_{b \geq 0} b \circlearrowright_{a-b} \oplus. \quad (4.20)$$

The following lemma gives a braid relation for *alternating crossings*. All other variations on the braid relation can be deduced from this plus the original braid relation from (1.8), by arguments similar to the proof of the braid relations in (2.7).

Lemma 4.5. *The following relation holds:*

$$\times \times - \times \times = z^3 \sum_{\substack{a, b \geq 0 \\ c > 0}} \oplus_{-a-b-c} \circlearrowleft_b \circlearrowright_c \quad \text{if } k \geq 0, \quad (4.21)$$

$$\times \times - \times \times = z^3 \sum_{\substack{a, b \geq 0 \\ c > 0}} \circlearrowleft_c \circlearrowright_{-a-b-c} \oplus \quad \text{if } k \leq 0. \quad (4.22)$$

Finally we have the *bubble slides*.

Lemma 4.6. *The following relations hold for any $a \in \mathbb{Z}$:*

$$\circlearrowleft \oplus_a = \circlearrowleft_a \oplus - z^2 \sum_{\substack{b \geq 0 \\ c > 0}} b+c \circlearrowleft_{a-b-c} \oplus, \quad (4.23)$$

$$\circlearrowright \oplus_a = \oplus_a \circlearrowright - z^2 \sum_{\substack{b \geq 0 \\ c > 0}} \oplus_{a-b-c} \circlearrowright_{b+c}, \quad (4.24)$$

$$a \overset{\circ}{\ominus} = \overset{\circ}{\ominus} a - z^2 \sum_{\substack{b \leq 0 \\ c < 0}} b+c \overset{\circ}{\ominus} a-b-c \overset{\circ}{\ominus}, \quad (4.25)$$

$$\overset{\circ}{\ominus} a = \overset{\circ}{\ominus} a - z^2 \sum_{\substack{b \leq 0 \\ c < 0}} \overset{\circ}{\ominus} a-b-c \overset{\circ}{\ominus} b+c. \quad (4.26)$$

5. Action on representations of quantum \mathbf{GL}_n

In this section, we construct an action of $\mathcal{Heis}_0(z, t)$ on the category of modules over $U_q(\mathfrak{gl}_n)$ and use this action to produce a family of generators for the center of $U_q(\mathfrak{gl}_n)$. These central elements were introduced originally by Bracken, Gould and Zhang [Gould et al. 1991]. We also determine their images under the Harish-Chandra homomorphism, giving a new approach to some results of Li [2010]. Throughout the section, we work in the generic case, setting

$$\mathbb{k} := \mathbb{Q}(q), \quad z := q - q^{-1}, \quad t := q^n$$

for an indeterminate q . In fact, the formulae which we derive are defined over $\mathbb{Z}[q, q^{-1}]$, hence, they make sense over any ground ring for any invertible q (including roots of unity).

For the precise definition of $U_q(\mathfrak{gl}_n)$, we follow the conventions of [Brundan 2017, Section 3], denoting its standard generators by $\{e_i, f_i, d_j^{\pm 1} \mid i = 1, \dots, n-1, j = 1, \dots, n\}$. The usual diagonal generator k_i of the subalgebra $U_q(\mathfrak{sl}_n)$ is $d_i d_{i+1}^{-1}$. The subalgebras of $U_q(\mathfrak{gl}_n)$ generated by the e_i, f_i and d_j^{\pm} are $U_q(\mathfrak{gl}_n)^+$, $U_q(\mathfrak{gl}_n)^-$ and $U_q(\mathfrak{gl}_n)^0$, respectively. We also have the Borel subalgebras $U_q(\mathfrak{gl}_n)^{\pm} := U_q(\mathfrak{gl}_n)^0 U_q(\mathfrak{gl}_n)^{\pm}$ and $U_q(\mathfrak{gl}_n)^{\mp} := U_q(\mathfrak{gl}_n)^0 U_q(\mathfrak{gl}_n)^{\pm}$. We will often cite Lusztig's book [1993], noting that our q and k_i are Lusztig's v^{-1} and K_i^{-1} .

The natural module V^+ and dual natural module V^- are the left $U_q(\mathfrak{gl}_n)$ -modules with bases

$$\{v_i^+ \mid 1 \leq i \leq n\} \quad \text{and} \quad \{v_i^- \mid 1 \leq i \leq n\},$$

respectively, on which the generators act by

$$f_i v_j^+ = \delta_{i,j} v_{i+1}^+, \quad e_i v_j^+ = \delta_{i+1,j} v_i^+, \quad d_i v_j^+ = q^{\delta_{i,j}} v_j^+, \quad (5.1)$$

$$f_i v_j^- = \delta_{i+1,j} v_i^-, \quad e_i v_j^- = \delta_{i,j} v_{i+1}^-, \quad d_i v_j^- = q^{-\delta_{i,j}} v_j^-. \quad (5.2)$$

We denote the weight of v_i^+ by ε_i ; then v_i^- is of weight $-\varepsilon_i$. Let $\Lambda := \bigoplus_{i=1}^n \mathbb{Z} \varepsilon_i$ be the *weight lattice* with inner product (\cdot, \cdot) defined so that $\varepsilon_1, \dots, \varepsilon_n$ are orthonormal. The *positive roots* are $\{\varepsilon_i - \varepsilon_j \mid 1 \leq i < j \leq n\}$. By a *weight module* we mean a $U_q(\mathfrak{gl}_n)$ -module V that is the sum of its weight spaces $V_{\lambda} := \{v \in V \mid d_i v = q^{(\lambda, \varepsilon_i)} v\}$ for all $\lambda \in \Lambda$. The *Weyl group* is the symmetric group \mathfrak{S}_n . It acts in obvious ways on Λ and on $U_q(\mathfrak{gl}_n)^0 = \mathbb{k}[d_1^{\pm 1}, \dots, d_n^{\pm 1}]$, permuting the generators. Denote the longest element of \mathfrak{S}_n by w_0 .

We work with the Hopf algebra structure on $U_q(\mathfrak{gl}_n)$ whose comultiplication Δ satisfies

$$\Delta(e_i) = d_i^{-1} d_{i+1} \otimes e_i + e_i \otimes 1, \quad \Delta(f_i) = 1 \otimes f_i + f_i \otimes d_i d_{i+1}^{-1}, \quad \Delta(d_j) = d_j \otimes d_j. \quad (5.3)$$

We also need various (anti)automorphisms. First, we have the *bar involution*, which is the antilinear automorphism $- : U_q \rightarrow U_q$ defined from $\bar{e}_i := e_i$, $\bar{f}_i := f_i$ and $\bar{d}_i := d_i^{-1}$. Then there are linear antiautomorphisms T and G defined from

$$T(e_i) := f_i, \quad T(f_i) := e_i, \quad T(d_i) := d_i, \quad (5.4)$$

$$G(e_i) := e_{n-i}, \quad G(f_i) := f_{n-i}, \quad G(d_i) := d_{n+1-i}. \quad (5.5)$$

The maps $-$, T and G commute with each other. Finally, we have Lusztig's braid group action, under which the i -th generator of the braid group acts by the automorphism $T_i : U_q(\mathfrak{gl}_n) \rightarrow U_q(\mathfrak{gl}_n)$ (which is $T_{i,-}''$ from [Lusztig 1993, Section 37.1.3]) defined for $|j - i| > 1$ and $k \neq i, i + 1$ by

$$\begin{aligned} T_i(e_i) &= -f_i d_i d_{i+1}^{-1}, & T_i(e_{i \pm 1}) &= e_i e_{i \pm 1} - q^{-1} e_{i \pm 1} e_i, & T_i(e_j) &= e_j, \\ T_i(f_i) &= -d_i^{-1} d_{i+1} e_i, & T_i(f_{i \pm 1}) &= f_{i \pm 1} f_i - q f_i f_{i \pm 1}, & T_i(f_j) &= f_j, \\ T_i(d_i) &= d_{i+1}, & T_i(d_{i+1}) &= d_i, & T_i(d_k) &= d_k. \end{aligned}$$

A key role is played by the *R-matrix*. We recall its definition following the approach from [Lusztig 1993, Section 32.1]. Let Θ be the *quasi-R-matrix* from [loc. cit., Section 4.1]. This is an infinite sum of components $\Theta_\alpha \in U_q(\mathfrak{gl}_n)_{-\alpha}^- \otimes U_q(\mathfrak{gl}_n)_\alpha^+$ as α runs over the positive root lattice $\bigoplus_{i=1}^{n-1} \mathbb{N}(\varepsilon_i - \varepsilon_{i+1})$. Let $P : V \otimes W \rightarrow W \otimes V$ be the tensor flip. Assuming in addition that V and W are weight modules, let $\Pi : V \otimes W \rightarrow V \otimes W$ be the diagonal map defined from

$$\Pi(v \otimes w) := q^{(\lambda, \mu)} v \otimes w$$

for v of weight λ and w of weight μ . Then, for finite-dimensional weight modules V and W , the *R-matrix*

$$R_{V,W} : V \otimes W \xrightarrow{\sim} W \otimes V \quad (5.6)$$

is the $U_q(\mathfrak{gl}_n)$ -module isomorphism defined by the composition $\Theta \circ P \circ \Pi$, which makes sense since all but finitely many of the components Θ_α act as zero. The inverse $R_{V,W}^{-1} : W \otimes V \rightarrow V \otimes W$ is the map $\Pi^{-1} \circ P^{-1} \circ \bar{\Theta}$, where $\bar{\Theta}$ is obtained from Θ by applying the bar involution to each tensor factor. For finite-dimensional weight modules U , V and W , we have the *hexagon property*

$$R_{U,W} \otimes \text{id}_V \circ \text{id}_U \otimes R_{V,W} = R_{U \otimes V, W}, \quad \text{id}_V \otimes R_{U,W} \circ R_{U,V} \otimes \text{id}_W = R_{U,V \otimes W}. \quad (5.7)$$

This is proved in [Lusztig 1993, Proposition 32.2.2] (our $R_{V,W}$ is Lusztig's $f\mathcal{R}_{W,V}$ taking the function f from [loc. cit., Section 31.1.3] to be $f(\lambda, \mu) := -(\lambda, \mu)$).

In fact, to define the isomorphism $R_{V,W}$, one only needs *one* of the modules V or W to be a finite-dimensional weight module; the other can be an arbitrary $U_q(\mathfrak{gl}_n)$ -module. To see this, one just needs to observe that Π extends to a linear map $V \otimes W \rightarrow V \otimes W$ when just one of V or W is a weight module on setting

$$\Pi(v \otimes w) := \begin{cases} (d_\lambda \otimes 1)(v \otimes w) & \text{if } w \text{ is a weight vector of weight } \lambda, \\ (1 \otimes d_\lambda)(v \otimes w) & \text{if } v \text{ is a weight vector of weight } \lambda, \end{cases}$$

where $d_\lambda := d_1^{(\lambda, \varepsilon_1)} \cdots d_n^{(\lambda, \varepsilon_n)}$. Then the same formula $R_{V,W} := \Theta \circ P \circ \Pi$ makes sense when only one of V or W is a finite-dimensional weight module, and it still gives an isomorphism of $U_q(\mathfrak{gl}_n)$ -modules. Moreover, the hexagon property (5.7) remains true if only two of U , V and W are finite-dimensional weight modules. These assertions follow from the known results in the previous paragraph. For example, to prove that $R_{V,W}$ is an isomorphism assuming that W is a finite-dimensional weight module, let $\rho_W: U_q(\mathfrak{gl}_n) \rightarrow \text{End}_{\mathbb{K}}(W)$ be the corresponding representation. Then

$$(\rho_W \otimes 1)(\Theta) \in \text{End}_{\mathbb{K}}(W) \otimes U_q(\mathfrak{gl}_n) \quad \text{and} \quad (\rho_W \otimes 1)(\bar{\Theta}) \in \text{End}_{\mathbb{K}}(W) \otimes U_q(\mathfrak{gl}_n).$$

It suffices to show that these are inverse to each other, since then $R_{V,W} = (\rho_W \otimes 1)(\Theta) \circ P \circ \Pi$ has inverse $\Pi^{-1} \circ P^{-1} \circ (\rho_W \otimes 1)(\bar{\Theta})$ for any module V . We have that

$$(\rho_W \otimes 1)(\Theta) \circ (\rho_W \otimes 1)(\bar{\Theta}) \in \text{End}_{\mathbb{K}}(W) \otimes U_q(\mathfrak{gl}_n)$$

and, for any finite-dimensional weight module V with corresponding representation ρ_V , we have

$$(1 \otimes \rho_V)((\rho_W \otimes 1)(\Theta) \circ (\rho_W \otimes 1)(\bar{\Theta})) = 1$$

by the known result. Since the intersection of the annihilators of all finite-dimensional weight modules is zero, this implies that $(\rho_W \otimes 1)(\Theta) \circ (\rho_W \otimes 1)(\bar{\Theta}) = 1$. The proof that $(\rho_W \otimes 1)(\bar{\Theta}) \circ (\rho_W \otimes 1)(\Theta) = 1$ is analogous, as is the proof of the hexagon property when just two of the modules are finite-dimensional weight modules.

The goal now is to derive explicit formulae for $R_{V^\pm, M}$ and R_{M, V^\pm} for any module M . Similar formulae were established already in [Gould et al. 1991, Section III] following the older conventions of Drinfeld and Jimbo. They involve the *higher root elements* defined as follows. Let

$$e_{i,i} = f_{i,i} := z^{-1}, \quad e_{i,i+1} := e_i, \quad f_{i,i+1} := f_i. \quad (5.8)$$

Then when $j - i > 1$ we recursively define

$$e_{i,j} := e_{i,r} e_{r,j} - q^{-1} e_{r,j} e_{i,r}, \quad f_{i,j} := f_{r,j} f_{i,r} - q^{-1} f_{i,r} f_{r,j}, \quad (5.9)$$

where r is any index chosen so that $i < r < j$. It is an induction exercise to see that these elements are well defined independent of the choice of r ; see the proof of the following lemma for a more conceptual explanation of this. Alternatively, $e_{i,j}$ and $f_{i,j}$ can be defined using the braid group action: we have that

$$e_{i,j} = T_{j-1} \cdots T_{i+1}(e_i), \quad f_{i,j} = \overline{T_{j-1} \cdots T_{i+1}(f_i)}.$$

Note that

$$T(e_{i,j}) = f_{i,j}, \quad T(f_{i,j}) = e_{i,j}, \quad (5.10)$$

$$G(e_{i,j}) = e_{n+1-j, n+1-i}, \quad G(f_{i,j}) = f_{n+1-j, n+1-i}. \quad (5.11)$$

However, the bar involution does not fix $e_{i,j}$ or $f_{i,j}$ (except when $j = i + 1$).

Lemma 5.1. *For any $i < j$, the $(\varepsilon_i - \varepsilon_j)$ -component $\Theta_{i,j}$ of the quasi- R -matrix Θ satisfies*

$$\Theta_{i,j} = \sum_{\substack{r \geq 1 \\ i=i_0 < i_1 < \dots < i_r = j}} z^r f_{i_{r-1}, i_r} \cdots f_{i_0, i_1} \otimes \overline{e_{i_{r-1}, i_r} \cdots e_{i_0, i_1}} = \sum_{\substack{r \geq 1 \\ i=i_0 < i_1 < \dots < i_r = j}} z^r \overline{f_{i_0, i_1} \cdots f_{i_{r-1}, i_r}} \otimes e_{i_0, i_1} \cdots e_{i_{r-1}, i_r}.$$

Proof. It suffices to derive the first expression. Then the second follows using (5.10) and the identity $(T \otimes T)(\Theta_\alpha) = P(\Theta_\alpha)$, which may easily be deduced from the characterization in [Lusztig 1993, Theorem 4.1.2(a)]. To prove the first expression, we appeal to further results of Lusztig from [loc. cit.]. Let f be Lusztig's "half" quantum group with its standard generators $\theta_1, \dots, \theta_{n-1}$; see also [Brundan et al. 2014, Section 2.1] which follows the same conventions as here. There are two isomorphisms

$$(-)^+ : f \xrightarrow{\sim} U_q(\mathfrak{gl}_n)^+, \quad \theta_i^+ := e_i, \quad (-)^- : f \xrightarrow{\sim} U_q(\mathfrak{gl}_n)^-, \quad \theta_i^- := f_i.$$

Consider the convex ordering on the positive roots defined so that $\varepsilon_i - \varepsilon_j < \varepsilon_p - \varepsilon_q$ if either $i < p$ or ($i = p$ and $j < q$); this is the "standard order" as in [loc. cit., Example A.1]. Let $\theta_{i,j}$ be Lusztig's higher root element associated to this ordering, which was denoted $r_{\varepsilon_i - \varepsilon_j}$ in [loc. cit., Section 2.4]. Noting that $(\varepsilon_m - \varepsilon_j, \varepsilon_i - \varepsilon_m)$ is a minimal pair for $\varepsilon_i - \varepsilon_j$, [loc. cit., Theorem 4.2] implies that these satisfy the following recursion: $\theta_{i,i+1} = \theta_i$ and $\theta_{i,j} = \theta_{i,r} \theta_{r,j} - q \theta_{r,j} \theta_{i,r}$ for any $i < r < j$. Comparing with (5.9), it follows that $\theta_{i,j}^+ = \overline{e_{i,j}}$ and $\theta_{i,j}^- = (-q)^{j-i-1} f_{i,j}$; in particular, these equalities justify the independence of r in (5.9). Then we appeal to [loc. cit., Theorem 2.7] (which was extracted from [Lusztig 1993]) to see that $\{\theta_{i_{r-1}, i_r} \cdots \theta_{i_0, i_1} \mid r \geq 1, i = i_0 < \dots < i_r = j\}$ and $\{(1-q^2)^r \theta_{i_{r-1}, i_r} \cdots \theta_{i_0, i_1} \mid r \geq 1, i = i_0 < \dots < i_r = j\}$ are a pair of dual bases for $f_{\varepsilon_i - \varepsilon_j}$ with respect to Lusztig's form. Finally the formula from [Lusztig 1993, Theorem 4.1.2(b)] gives that

$$\Theta_{i,j} = \sum_{\substack{r \geq 1 \\ i=i_0 < \dots < i_r = j}} (-q)^{i-j} (1-q^2)^r \theta_{i_{r-1}, i_r}^- \cdots \theta_{i_0, i_1}^- \otimes \theta_{i_{r-1}, i_r}^+ \cdots \theta_{i_0, i_1}^+.$$

This simplifies to the desired formula. \square

For $1 \leq i, j \leq n$ let $e_{i,j}^+ \in \text{End}_\mathbb{k}(V^+)$ (resp. $e_{i,j}^- \in \text{End}_\mathbb{k}(V^-)$) be the ij -matrix unit with respect to the basis v_1^+, \dots, v_n^+ (resp. v_1^-, \dots, v_n^-). Then for $i < j$ and $v^\pm \in V^\pm$ we have that

$$e_{i,j} v^+ = e_{i,j}^+ v^+, \quad f_{i,j} v^+ = e_{j,i}^+ v^+, \quad e_{i,j} v^- = (-q)^{i-j+1} e_{j,i}^- v^-, \quad f_{i,j} v^- = (-q)^{i-j+1} e_{i,j}^- v^-, \quad (5.12)$$

$$\overline{e_{i,j}} v^+ = e_{i,j}^+ v^+, \quad \overline{f_{i,j}} v^+ = e_{j,i}^+ v^+, \quad \overline{e_{i,j}} v^- = (-q)^{j-i-1} e_{j,i}^- v^-, \quad \overline{f_{i,j}} v^- = (-q)^{j-i-1} e_{i,j}^- v^-. \quad (5.13)$$

These follow easily by induction on $j - i$ using (5.1)–(5.2) and (5.9). Also let

$$x_{i,j} := z^2 \sum_{r=1}^{\min(i,j)} e_{r,i} d_r f_{r,j} d_j, \quad y_{i,j} := z^2 \sum_{r=\max(i,j)}^n d_i f_{i,r} d_r e_{j,r} \quad (5.14)$$

for any $1 \leq i, j \leq n$. Then for $m \geq 0$ we set

$$x_{i,j}^{(m)} := \sum_{i=i_0, i_1, \dots, i_{m-1}, i_m=j} x_{i_0, i_1} \cdots x_{i_{m-1}, i_m}, \quad y_{i,j}^{(m)} := \sum_{i=i_0, i_1, \dots, i_{m-1}, i_m=j} y_{i_0, i_1} \cdots y_{i_{m-1}, i_m}. \quad (5.15)$$

In particular, $x_{i,j}^{(0)} = y_{i,j}^{(0)} = \delta_{i,j}$. From (5.11), we get that

$$G(x_{i,j}^{(m)}) = y_{n+1-j,n+1-i}^{(m)}, \quad G(y_{i,j}^{(m)}) = x_{n+1-j,n+1-i}^{(m)}. \quad (5.16)$$

Lemma 5.2. *For any $U_q(\mathfrak{gl}_n)$ -module M , the endomorphisms $R_{V^\pm, M}$ and R_{M, V^\pm} and their inverses are given explicitly by the following operators:*

$$\begin{aligned} R_{V^+, M} &= zP \circ \sum_{i \leq j} e_{i,j}^+ \otimes f_{i,j} d_j, & R_{V^+, M}^{-1} &= -zP \otimes \sum_{i \leq j} \overline{d_i f_{i,j}} \otimes e_{i,j}^+, \\ R_{M, V^+} &= zP \circ \sum_{i \leq j} e_{i,j} d_i \otimes e_{j,i}^+, & R_{M, V^+}^{-1} &= -zP \circ \sum_{i \leq j} e_{j,i}^+ \otimes \overline{d_j e_{i,j}}, \\ R_{V^-, M} &= -zP \circ \sum_{i \leq j} (-q)^{i-j} e_{j,i}^- \otimes \overline{d_i f_{i,j}}, & R_{V^-, M}^{-1} &= zP \circ \sum_{i \leq j} (-q)^{i-j} f_{i,j} d_j \otimes e_{j,i}^-, \\ R_{M, V^-} &= -zP \circ \sum_{i \leq j} (-q)^{i-j} \overline{d_j e_{i,j}} \otimes e_{i,j}^-, & R_{M, V^-}^{-1} &= zP \circ \sum_{i \leq j} (-q)^{i-j} e_{i,j}^- \otimes e_{i,j} d_i. \end{aligned}$$

Proof. These are all proved by similar calculations, so we just go through the argument for R_{M, V^-} . Take $v \otimes v_j^- \in M \otimes V^-$. By definition, $R_{M, V^-}(v \otimes v_j^-) = \Theta(v_j^- \otimes d_j^{-1}v)$. To compute the action of Θ , we observe by weight considerations that only its weight components $\Theta_{\varepsilon_i - \varepsilon_j}$ for $i \leq j$ are nonzero on $v_j^- \otimes d_j^{-1}v$. Moreover, in the first expression for $\Theta_{i,j}$ from Lemma 5.1, all of the monomials with $r > 1$ act on v_j^- as zero. We deduce that

$$R_{M, V^-}(v \otimes v_j^-) = v_j^- \otimes d_j^{-1}v + z \sum_{i < j} f_{i,j} v_j^- \otimes \overline{e_{i,j} d_j} v.$$

Then we use (5.12) to replace $f_{i,j}$ with $(-q)^{i-j+1} e_{i,j}^-$, the relation $e_{i,j} d_j = q d_j e_{i,j}$, and the definition $\overline{e_{j,j}} = -z^{-1}$ to get

$$R_{M, V^-}(v \otimes v_j^-) = -z e_{j,j}^- v_j^- \otimes \overline{d_j e_{j,j}} v - z \sum_{i < j} (-q)^{i-j} e_{i,j}^- v_j^- \otimes \overline{e_{i,j} d_j} v.$$

Now observe that the expression on the right-hand side of the formula we are trying to prove acts on $v \otimes v_j^-$ in the same way. \square

Corollary 5.3. *For any $U_q(\mathfrak{gl}_n)$ -module M and $m \in \mathbb{Z}$, we have that*

$$(R_{M, V^+} \circ R_{V^+, M})^m = \begin{cases} \sum_{i,j=1}^n e_{i,j}^+ \otimes x_{i,j}^{(m)} & \text{if } m \geq 0, \\ \sum_{i,j=1}^n e_{i,j}^+ \otimes \overline{y_{i,j}^{(-m)}} & \text{if } m \leq 0. \end{cases}$$

Proof. This follows from Lemma 5.2 and Definitions (5.14)–(5.15). \square

Now we return to the Heisenberg category $\mathcal{H}eis_0(z, t)$ taking $t := q^n$. Let $\mathcal{OS}(z, t)$ be the *HOMFLY-PT skein category* as defined in the introduction of [Brundan 2017], which is Turaev's Hecke category [1989].

By [Brundan 2017, Theorem 1.1], $\mathcal{OS}(z, t)$ has a presentation by generators and relations which is very similar to the presentation of $\mathcal{Heis}_0(z, t)$ from [Definition 2.2](#) but *without* the morphism x . Consequently, there is a strict \mathbb{k} -linear monoidal functor $\mathcal{OS}(z, t) \rightarrow \mathcal{Heis}_0(z, t)$. By [[loc. cit.](#), Lemma 4.2], this functor is faithful, so we may use it to *identify* $\mathcal{OS}(z, t)$ with a subcategory of $\mathcal{Heis}_0(z, t)$. Thus, $\mathcal{OS}(z, t)$ is the monoidal subcategory of $\mathcal{Heis}_0(z, t)$ consisting of all objects and all morphisms which do not involve dots (i.e., x or y). In fact, as noted already after [Definition 2.2](#), $\mathcal{Heis}_0(z, t)$ is the *affine* HOMFLY-PT skein category from [[loc. cit.](#), Section 4].

Let $U_q(\mathfrak{gl}_n)\text{-mod}$ be the category of all left $U_q(\mathfrak{gl}_n)$ -modules. By [[loc. cit.](#), Lemma 3.1] (although the result is much older, e.g., it was exploited already in [[Turaev 1989](#)]), there is a monoidal functor

$$\Psi : \mathcal{OS}(z, t) \rightarrow U_q(\mathfrak{gl}_n)\text{-mod} \quad (5.17)$$

to the category of left $U_q(\mathfrak{gl}_n)$ -modules. The functor Ψ sends the generating objects \uparrow and \downarrow to V^+ and V^- , respectively. It maps the various generating morphisms to the following $U_q(\mathfrak{gl}_n)$ -module homomorphisms:

$$\begin{array}{l} \nearrow : v_i^+ \otimes v_j^+ \mapsto \begin{cases} v_j^+ \otimes v_i^+ & \text{if } i < j, \\ q v_j^+ \otimes v_i^+ & \text{if } i = j, \\ v_j^+ \otimes v_i^+ + z v_i^+ \otimes v_j^+ & \text{if } i > j; \end{cases} \\ \searrow : v_i^+ \otimes v_j^- \mapsto \begin{cases} v_j^- \otimes v_i^+ & \text{if } i \neq j, \\ q^{-1} v_j^- \otimes v_i^+ - z \sum_{r=1}^{i-1} (-q)^{-r} v_{j-r}^- \otimes v_{i-r}^+ & \text{if } i = j; \end{cases} \end{array} \quad (5.18)$$

$$\begin{array}{l} \nearrow : v_i^- \otimes v_j^- \mapsto \begin{cases} v_j^- \otimes v_i^- & \text{if } i > j, \\ q v_j^- \otimes v_i^- & \text{if } i = j, \\ v_j^- \otimes v_i^- + z v_i^- \otimes v_j^- & \text{if } i < j; \end{cases} \\ \searrow : v_i^- \otimes v_j^+ \mapsto \begin{cases} v_j^+ \otimes v_i^- & \text{if } i \neq j, \\ q^{-1} v_j^+ \otimes v_i^- - z \sum_{r=1}^{n-i} (-q)^{-r} v_{j+r}^+ \otimes v_{i+r}^- & \text{if } i = j; \end{cases} \end{array} \quad (5.19)$$

$$\begin{array}{l} \nearrow : v_i^+ \otimes v_j^+ \mapsto \begin{cases} v_j^+ \otimes v_i^+ & \text{if } i > j, \\ q^{-1} v_j^+ \otimes v_i^+ & \text{if } i = j, \\ v_j^+ \otimes v_i^+ - z v_i^+ \otimes v_j^+ & \text{if } i < j; \end{cases} \\ \searrow : v_i^+ \otimes v_j^- \mapsto \begin{cases} v_j^- \otimes v_i^+ & \text{if } i \neq j, \\ q v_j^- \otimes v_i^+ + z \sum_{r=1}^{n-i} (-q)^r v_{j+r}^- \otimes v_{i+r}^+ & \text{if } i = j; \end{cases} \end{array} \quad (5.20)$$

$$\begin{array}{l} \nearrow : v_i^- \otimes v_j^- \mapsto \begin{cases} v_j^- \otimes v_i^- & \text{if } i < j, \\ q^{-1} v_j^- \otimes v_i^- & \text{if } i = j, \\ v_j^- \otimes v_i^- - z v_i^- \otimes v_j^- & \text{if } i > j; \end{cases} \\ \searrow : v_i^- \otimes v_j^+ \mapsto \begin{cases} v_j^+ \otimes v_i^- & \text{if } i \neq j, \\ q v_j^+ \otimes v_i^- + z \sum_{r=1}^{i-1} (-q)^r v_{j-r}^+ \otimes v_{i-r}^- & \text{if } i = j; \end{cases} \end{array} \quad (5.21)$$

$$\begin{array}{l} \nearrow : v_i^+ \otimes v_j^+ \mapsto \begin{cases} v_j^+ \otimes v_i^+ & \text{if } i > j, \\ q^{-1} v_j^+ \otimes v_i^+ & \text{if } i = j, \\ v_j^+ \otimes v_i^+ - z v_i^+ \otimes v_j^+ & \text{if } i < j; \end{cases} \\ \searrow : v_i^+ \otimes v_j^- \mapsto \begin{cases} v_j^- \otimes v_i^+ & \text{if } i < j, \\ q^{-1} v_j^- \otimes v_i^+ & \text{if } i = j, \\ v_j^- \otimes v_i^+ - z v_i^- \otimes v_j^+ & \text{if } i > j; \end{cases} \end{array} \quad (5.22)$$

$$\begin{array}{l} \nearrow : v_i^- \otimes v_j^- \mapsto \begin{cases} v_j^- \otimes v_i^- & \text{if } i < j, \\ q^{-1} v_j^- \otimes v_i^- & \text{if } i = j, \\ v_j^- \otimes v_i^- - z v_i^- \otimes v_j^- & \text{if } i > j; \end{cases} \\ \searrow : v_i^- \otimes v_j^+ \mapsto \begin{cases} v_j^+ \otimes v_i^- & \text{if } i \neq j, \\ q v_j^+ \otimes v_i^- + z \sum_{r=1}^{i-1} (-q)^r v_{j-r}^+ \otimes v_{i-r}^- & \text{if } i = j; \end{cases} \end{array} \quad (5.23)$$

$$\begin{array}{l} \nearrow : v_i^- \otimes v_j^+ \mapsto \begin{cases} v_j^+ \otimes v_i^- & \text{if } i \neq j, \\ q^{-1} v_j^+ \otimes v_i^- & \text{if } i = j, \\ v_j^+ \otimes v_i^- - z v_i^- \otimes v_j^+ & \text{if } i > j; \end{cases} \\ \searrow : v_i^- \otimes v_j^- \mapsto \begin{cases} v_j^- \otimes v_i^+ & \text{if } i < j, \\ q^{-1} v_j^- \otimes v_i^+ & \text{if } i = j, \\ v_j^- \otimes v_i^+ - z v_i^+ \otimes v_j^- & \text{if } i > j; \end{cases} \end{array} \quad (5.24)$$

$$\begin{array}{l} \nearrow : v_i^- \otimes v_j^+ \mapsto \begin{cases} v_j^+ \otimes v_i^- & \text{if } i \neq j, \\ q v_j^+ \otimes v_i^- + z \sum_{r=1}^{i-1} (-q)^r v_{j-r}^+ \otimes v_{i-r}^- & \text{if } i = j; \end{cases} \\ \searrow : v_i^- \otimes v_j^- \mapsto \begin{cases} v_j^- \otimes v_i^+ & \text{if } i < j, \\ q^{-1} v_j^- \otimes v_i^+ & \text{if } i = j, \\ v_j^- \otimes v_i^+ - z v_i^+ \otimes v_j^- & \text{if } i > j; \end{cases} \end{array} \quad (5.25)$$

$$\begin{array}{l} \cup : 1 \mapsto \sum_{j=1}^n (-1)^j q^j v_j^- \otimes v_j^+, \\ \cap : v_i^+ \otimes v_j^- \mapsto (-1)^i q^{-i} \delta_{i,j}, \end{array} \quad \begin{array}{l} \cup : 1 \mapsto \sum_{j=1}^n (-1)^j q^{n+1-j} v_j^+ \otimes v_j^-, \\ \cap : v_i^- \otimes v_j^+ \mapsto (-1)^i q^{i-n-1} \delta_{i,j}. \end{array} \quad (5.26) \quad (5.27)$$

These formulae are recorded in many places in the literature going back to the original work [Turaev 1989], but one finds many different choices of normalization. For our choices, (5.18)–(5.21) and (5.22)–(5.25) follow from the formulae for the R -matrix and its inverse from [Lemma 5.2](#), while the formulae (5.26)–(5.27) are derived in [\[Brundan 2017, Section 3\]](#).

Theorem 5.4. *Assuming $t = q^n$ and $z = q - q^{-1}$, there is a strict \mathbb{k} -linear monoidal functor*

$$\widehat{\Psi} : \mathcal{H}eis_0(z, t) \rightarrow \mathcal{E}nd_{\mathbb{k}}(U_q(\mathfrak{gl}_n)\text{-mod})$$

such that $\Psi = \text{Ev} \circ \widehat{\Psi}|_{\mathcal{O}S(z, t)}$, where Ev denotes evaluation on the trivial module. On objects, $\widehat{\Psi}$ takes X to the endofunctor $\Psi(X) \otimes -$, e.g., $\widehat{\Psi}(\uparrow) = V^+ \otimes -$ and $\widehat{\Psi}(\downarrow) = V^- \otimes -$. On morphisms, $\widehat{\Psi}$ sends $f \in \text{Hom}_{\mathcal{O}S(z, t)}(X, Y)$ to the natural transformation $\Psi(f) \otimes 1 : \Psi(X) \otimes - \rightarrow \Psi(Y) \otimes -$. Finally, on the additional generating morphism x , it is defined by

$$\widehat{\Psi}(x)_M := R_{M, V^+} \circ R_{V^+, M} : V^+ \otimes M \rightarrow V^+ \otimes M, \quad v_j^+ \otimes m \mapsto \sum_{i=1}^n v_i^+ \otimes x_{i,j} m.$$

Proof. We just need to verify that the relations from [Definition 2.2](#) are satisfied. All of the ones that do not involve x follow immediately since they are already satisfied by the morphisms in the image of the monoidal functor Ψ . Also $R_{V^+, M} \circ R_{M, V^+}$ is invertible since each of these R -matrices is invertible. It just remains to check the relation (1.6). In fact, this is a formal consequence of the hexagon property; see e.g., [\[Virk 2011, Proposition 3.1.1\]](#). The argument goes as follows. By (5.7), we have for any $U_q(\mathfrak{gl}_n)$ -module M that

$$R_{V^+ \otimes M, V^+} \circ R_{V^+, V^+ \otimes M} = R_{V^+, V^+} \otimes \text{id}_M \circ \text{id}_{V^+} \otimes R_{M, V^+} \circ \text{id}_{V^+} \otimes R_{V^+, M} \circ R_{V^+, V^+} \otimes \text{id}_M.$$

This establishes that the image under $\widehat{\Psi}$ of the relation

$$\begin{array}{c} \uparrow \quad \uparrow \\ \circ \quad | = \quad \circ \\ \uparrow \quad \uparrow \end{array}$$

is satisfied, from which (1.6) easily follows. \square

Let $Z_q(\mathfrak{gl}_n)$ be the center of $U_q(\mathfrak{gl}_n)$. It is identified with the endomorphism algebra of the identity functor $\text{Id}_{U_q(\mathfrak{gl}_n)\text{-mod}}$; indeed, evaluation on the identity element of the regular representation defines a canonical algebra isomorphism $\text{End}(\text{Id}_{U_q(\mathfrak{gl}_n)\text{-mod}}) \xrightarrow{\sim} Z_q(\mathfrak{gl}_n)$. Dotted bubbles are endomorphisms of the unit object of $\mathcal{H}eis_0(z, t)$. Applying the monoidal functor $\widehat{\Psi}$ from [Theorem 5.4](#), we obtain natural transformations

$$\widehat{\Psi}(\bigcirc \circ^m) : \text{Id}_{U_q(\mathfrak{gl}_n)\text{-mod}} \rightarrow \text{Id}_{U_q(\mathfrak{gl}_n)\text{-mod}}, \tag{5.28}$$

hence, central elements $z_m \in Z(U_q(\mathfrak{gl}_n))$ for each $m \in \mathbb{Z}$. A calculation using (5.26)–(5.27) and [Corollary 5.3](#) shows that

$$z_m = \begin{cases} \sum_{i=1}^n q^{2i-n-1} x_{i,i}^{(m)} & \text{if } m \geq 0, \\ \sum_{i=1}^n q^{2i-n-1} \overline{y_{i,i}^{(-m)}} & \text{if } m \leq 0. \end{cases} \tag{5.29}$$

We have trivially that $z_0 = [n]_q$. The goal in the remainder of the section is to compute explicit formulae for the images of all the others under the Harish-Chandra homomorphism.

Our argument uses the Harish-Chandra homomorphism in two different forms adapted to the positive and negative Borel subalgebras, respectively. To review the definitions, let $\rho_+ := -\varepsilon_2 - 2\varepsilon_3 - \dots - (n-1)\varepsilon_n$ and $\rho_- := -(n-1)\varepsilon_1 - \dots - 2\varepsilon_{n-2} - \varepsilon_{n-1}$, i.e., $\rho_- = w_0(\rho_+)$. For any $\lambda \in \Lambda$, we have the *shift automorphism*

$$S_\lambda : U_q(\mathfrak{gl}_n)^0 \rightarrow U_q(\mathfrak{gl}_n)^0, \quad d_i \mapsto q^{(\lambda, \varepsilon_i)} d_i. \quad (5.30)$$

For example, $S_{-\rho_+}(d_i) = q^{i-1} d_i$ and $S_{-\rho_-}(d_i) = q^{n-i} d_i$. Let $U_q(\mathfrak{gl}_n)_0$ be the zero weight space of $U_q(\mathfrak{gl}_n)$, which is a subalgebra containing $U_q(\mathfrak{gl}_n)^0$. Let I_+ (resp. I_-) be the intersection of $U_q(\mathfrak{gl}_n)_0$ with the left ideal of $U_q(\mathfrak{gl}_n)$ generated by e_1, \dots, e_{n-1} (resp. f_1, \dots, f_{n-1}). Equivalently, I_+ (resp. I_-) is the intersection of $U_q(\mathfrak{gl}_n)_0$ with the right ideal generated by f_1, \dots, f_{n-1} (resp. e_1, \dots, e_{n-1}). It follows that I_\pm is a two-sided ideal of $U_q(\mathfrak{gl}_n)_0$. Let $\text{pr}_\pm : U_q(\mathfrak{gl}_n)_0 \rightarrow U_q(\mathfrak{gl}_n)^0$ be the algebra homomorphism defined by projection along the direct sum decomposition $U_q(\mathfrak{gl}_n)_0 = U_q(\mathfrak{gl}_n)^0 \oplus I_\pm$. The two versions of the *Harish-Chandra homomorphism* are

$$HC_\pm := S_{-\rho_\pm} \circ \text{pr}_\pm : U_q(\mathfrak{gl}_n)_0 \rightarrow U_q(\mathfrak{gl}_n)^0. \quad (5.31)$$

The following is an extension of the well-known description of $Z_q(\mathfrak{sl}_n)$ from e.g., [Jantzen 1996, 6.25].

Lemma 5.5 [Li 2010, Lemma 2.1]. *The restriction $HC := HC_+|_{Z_q(\mathfrak{gl}_n)}$ defines an algebra isomorphism between $Z_q(\mathfrak{gl}_n)$ and the algebra $\mathbb{k}[(d_1 \cdots d_n)^{-1}, d_1^2, \dots, d_n^2]^{\mathfrak{S}_n}$.*

The following facts are also well known, but we could not find a suitable reference.

Lemma 5.6. *Each braid group generator $T_i : U_q(\mathfrak{gl}_n) \rightarrow U_q(\mathfrak{gl}_n)$ fixes $Z_q(\mathfrak{gl}_n)$ pointwise.*

Proof. Take $c \in Z_q(\mathfrak{gl}_n)$. Let V be an integrable highest weight module. Since V is irreducible, both c and $T_i(c)$ act on V as scalars. These scalars are equal because there is an automorphism $T_i : V \rightarrow V$ such that $T_i(cv) = T_i(c)T_i(v)$; see [Lusztig 1993, Section 37.1.1]. This shows that $c - T_i(c)$ acts as zero on every integrable highest weight module. The intersection of the annihilators of all integrable highest weight modules is zero, so this proves that $c = T_i(c)$. \square

Lemma 5.7. *The restriction $HC = HC_+|_{Z_q(\mathfrak{gl}_n)}$ is equal also to the restriction $HC_-|_{Z_q(\mathfrak{gl}_n)}$.*

Proof. Let T_{w_0} be the product of simple braid group generators T_i taken in some order corresponding to a reduced expression of w_0 . This is an automorphism of $U_q(\mathfrak{gl}_n)$ which switches $U_q(\mathfrak{gl}_n)^\sharp$ and $U_q(\mathfrak{gl}_n)^\flat$, and it sends $d_i \mapsto d_{n+1-i}$. It follows that

$$HC_\mp \circ T_{w_0} = T_{w_0} \circ HC_\pm. \quad (5.32)$$

Clearly, T_{w_0} fixes $\mathbb{k}[(d_1 \cdots d_n)^{-1}, d_1^2, \dots, d_n^2]^{\mathfrak{S}_n}$ pointwise. It also fixes $Z_q(\mathfrak{gl}_n)$ pointwise by Lemma 5.6. Hence, $HC_-|_{Z_q(\mathfrak{gl}_n)} = HC_- \circ T_{w_0}|_{Z_q(\mathfrak{gl}_n)} = T_{w_0} \circ HC_+|_{Z_q(\mathfrak{gl}_n)} = HC_+|_{Z_q(\mathfrak{gl}_n)}$. \square

Lemma 5.8. *The antiautomorphism G fixes $Z_q(\mathfrak{gl}_n)$ pointwise.*

Proof. We have that

$$HC_{\mp} \circ G = G \circ HC_{\pm}. \quad (5.33)$$

Combined with [Lemma 5.7](#), it follows that $HC_+ \circ G|_{Z_q(\mathfrak{gl}_n)} = G \circ HC_+|_{Z_q(\mathfrak{gl}_n)}$. Also G clearly fixes $\mathbb{K}[(d_1 \cdots d_n)^{-1}, d_1^2, \dots, d_n^2]^{\mathfrak{S}_n}$ pointwise. Hence, $HC_+ \circ G|_{Z_q(\mathfrak{gl}_n)} = HC_+|_{Z_q(\mathfrak{gl}_n)}$, which implies the result since HC_+ is injective on $Z_q(\mathfrak{gl}_n)$. \square

In particular, this shows that $G(z_m) = z_m$, hence, on applying G to the right-hand side of [\(5.29\)](#) using [\(5.16\)](#), we obtain another formula for z_m :

$$z_m = \begin{cases} \sum_{i=1}^n q^{n+1-2i} y_{i,i}^{(m)} & \text{if } m \geq 0, \\ \sum_{i=1}^n q^{n+1-2i} \overline{x_{i,i}^{(-m)}} & \text{if } m \leq 0. \end{cases} \quad (5.34)$$

Comparing with [\(5.29\)](#), it follows that

$$z_{-m} = \overline{z_m} \quad (5.35)$$

for every $m \in \mathbb{Z}$. From now on, we only consider z_m for $m \geq 1$.

Finally, consider the *modified complete symmetric polynomials*

$$\tilde{h}_m(x_1, \dots, x_n) := \sum_{1 \leq i_1 \leq \dots \leq i_m \leq n} (q^{-1}z)^{\#\{i_1, \dots, i_m\}-1} x_{i_1} \cdots x_{i_m}. \quad (5.36)$$

We will use these for all values of $n \geq 0$ (not just the n fixed above for \mathfrak{gl}_n). We have that

$$\tilde{h}_m(x_1, \dots, x_n) = qz^{-1} \text{ if } m = 0 \quad \text{and} \quad \tilde{h}_m(x_1, \dots, x_n) = 0 \text{ if } m > 0 \text{ but } n = 0.$$

These elements obviously satisfy the recurrence relation

$$\tilde{h}_m(x_1, \dots, x_n) = \tilde{h}_m(x_1, \dots, x_{n-1}) + q^{-1}z \sum_{r=1}^m \tilde{h}_{m-r}(x_1, \dots, x_{n-1}) x_n^r \quad (5.37)$$

for $n > 0$.

Lemma 5.9. $\tilde{h}_m(x_1, \dots, x_n) = \tilde{h}_m(x_1, \dots, x_{n-1}) + \tilde{h}_{m-1}(x_1, \dots, x_n) x_n - q^{-2} \tilde{h}_{m-1}(x_1, \dots, x_{n-1}) x_n$.

Proof. By [\(5.37\)](#) with m replaced by $m-1$, we have that

$$\begin{aligned} \tilde{h}_{m-1}(x_1, \dots, x_n) x_n &= \tilde{h}_{m-1}(x_1, \dots, x_{n-1}) x_n + q^{-1}z \sum_{r=1}^{m-1} \tilde{h}_{m-r-1}(x_1, \dots, x_{n-1}) x_n^{r+1} \\ &= \tilde{h}_{m-1}(x_1, \dots, x_{n-1}) x_n + q^{-1}z \sum_{r=2}^m \tilde{h}_{m-r}(x_1, \dots, x_{n-1}) x_n^r \\ &= q^{-2} \tilde{h}_{m-1}(x_1, \dots, x_{n-1}) x_n + q^{-1}z \sum_{r=1}^m \tilde{h}_{m-r}(x_1, \dots, x_{n-1}) x_n^r. \end{aligned}$$

Given this, it is easy to see that the right-hand side of the identity we are trying to prove is equal to the right-hand side of [\(5.37\)](#). \square

Theorem 5.10. *For any $m \geq 1$ we have that $HC(z_m) = q^{n-1}\tilde{h}_m(d_1^2, \dots, d_n^2)$.*

Proof. Noting that $q^{1-n}z_m = \sum_{i=1}^n q^{2i-2n}x_{i,i}^{(m)}$ according to (5.29), this follows from the following claim: *for any $m \geq 1$ and $i = 1, \dots, n$, we have that*

$$HC_+(x_{i,i}^{(m)}) = \tilde{h}_m(d_1^2, \dots, d_i^2) - q^{-2}\tilde{h}_m(d_1^2, \dots, d_{i-1}^2). \quad (5.38)$$

To prove (5.38), we proceed by induction on $m+n$. The result is easy to check when $n=1$. Now assume that $n > 1$. The Harish-Chandra homomorphism HC_+ is compatible with the usual “top left corner” embedding of $U_q(\mathfrak{gl}_{n-1})$ into $U_q(\mathfrak{gl}_n)$. This follows because the restriction of ρ_+ for \mathfrak{gl}_n is the weight ρ_+ for \mathfrak{gl}_{n-1} . Also the elements $x_{1,1}^{(m)}, \dots, x_{n-1,n-1}^{(m)}$ of $U_q(\mathfrak{gl}_{n-1})$ are the same as these elements in $U_q(\mathfrak{gl}_n)$. Thus we get (5.38) for each $i < n$ from the induction hypothesis. It remains to prove (5.38) when $i=n$. We have that

$$q^{1-n}HC_-(z_m) = \sum_{i=1}^n q^{2i-2n} \sum_{j_1, \dots, j_m} HC_-(z^{2m}e_{j_1,i}d_{j_1}f_{j_1,j_2}d_{j_2} \cdots e_{j_m,j_{m-1}}d_{j_m}f_{j_m,i}d_i).$$

By the definition of HC_- , the terms in this expansion are zero if either $j_1 < i$ or $j_m < i$. Thus, the sum simplifies to give

$$q^{1-n}HC_-(z_m) = \sum_{i=1}^n q^{2i-2n} HC_-(y_{i,i}^{(m-1)} d_i^2) = \sum_{i=1}^n HC_-(y_{i,i}^{(m-1)}) d_i^2.$$

Now we apply G , using Lemma 5.8, (5.33) and (5.11), to see that

$$q^{1-n}HC_+(z_m) = \sum_{i=1}^n HC_+(x_{i,i}^{(m-1)}) d_i^2.$$

Remembering (5.29), we have now proved that

$$\sum_{i=1}^n q^{2i-2n} HC_+(x_{i,i}^{(m)}) = \sum_{i=1}^n HC_+(x_{i,i}^{(m-1)}) d_i^2. \quad (5.39)$$

The same identity with n replaced by $(n-1)$ gives

$$\sum_{i=1}^{n-1} q^{2i-2(n-1)} HC_+(x_{i,i}^{(m)}) = \sum_{i=1}^{n-1} HC_+(x_{i,i}^{(m-1)}) d_i^2. \quad (5.40)$$

By the induction hypothesis, the left-hand side of (5.40) is equal to $\tilde{h}_m(d_1^2, \dots, d_{n-1}^2)$. Hence, (5.39) can be rewritten to obtain

$$\begin{aligned} HC_+(x_{n,n}^{(m)}) + q^{-2}\tilde{h}_m(d_1^2, \dots, d_{n-1}^2) \\ = HC_+(x_{n,n}^{(m-1)}) d_n^2 + \tilde{h}_m(d_1^2, \dots, d_{n-1}^2) \\ = \tilde{h}_m(d_1^2, \dots, d_{n-1}^2) + \tilde{h}_{m-1}(d_1^2, \dots, d_n^2) d_n^2 - q^{-2}\tilde{h}_{m-1}(d_1^2, \dots, d_{n-1}^2) d_n^2, \end{aligned}$$

where we have used the induction hypothesis again to establish the second equality. This is equal to $\tilde{h}_m(d_1^2, \dots, d_n^2)$ thanks to [Lemma 5.9](#). The conclusion follows. \square

Corollary 5.11 [Li 2010, Theorem 4.1]. $Z_q(\mathfrak{gl}_n)$ is generated by z_1, \dots, z_n and $(d_1 \cdots d_n)^{-1}$.

Proof. This follows from [Lemma 5.5](#) and [Theorem 5.10](#) since $\mathbb{k}[x_1, \dots, x_n]^{\mathfrak{S}_n}$ is generated by the modified complete symmetric functions $\tilde{h}_1(x_1, \dots, x_n), \dots, \tilde{h}_n(x_1, \dots, x_n)$. \square

6. Action on modules over cyclotomic Hecke algebras

Throughout the section, we assume that we are given a polynomial

$$f(w) = f_0 w^l + f_1 w^{l-1} + \cdots + f_l \in \mathbb{k}[w] \quad (6.1)$$

of degree $l \geq 0$ such that $f_0 = 1$ and $f_l = t^2$. Recall from the introduction that the affine Hecke algebra AH_n with its standard generators $x_1, \dots, x_n, \tau_1, \dots, \tau_{n-1}$ is identified with the endomorphism algebra $\text{End}_{A\mathcal{H}(z)}(\uparrow^{\otimes n})$ so that x_i is the dot on the i -th string and τ_j is the positive crossing of the j -th and $(j+1)$ -th strings (numbering strings $1, \dots, n$ from right to left). The *cyclotomic Hecke algebra* H_n^f of level l associated to the polynomial $f(w)$ is the quotient of AH_n by the two-sided ideal generated by $f(x_1)$. We also include the possibility $n = 0$ with the convention that $H_0^f = \mathbb{k}$.

The basis theorem proved in [\[Ariki and Koike 1994, Theorem 3.10\]](#) shows that the following gives a basis for H_n^f as a free \mathbb{k} -module:

$$\{x_1^{r_1} \cdots x_n^{r_n} \tau_g \mid 0 \leq r_1, \dots, r_n < l, g \in \mathfrak{S}_n\}, \quad (6.2)$$

where τ_g denotes the element of the finite Hecke algebra defined from a reduced expression for the permutation g . By the basis theorem, the obvious homomorphism $H_n^f \rightarrow H_{n+1}^f$ sending the generators x_i and τ_j to the elements of H_{n+1}^f with the same names is *injective*. So we may identify H_n^f with a subalgebra of H_{n+1}^f . We denote the induction and restriction functors by

$$\text{ind}_n^{n+1} := H_{n+1}^f \otimes_{H_n^f} - : H_n^f\text{-mod} \rightarrow H_{n+1}^f\text{-mod}, \quad (6.3)$$

$$\text{res}_n^{n+1} : H_{n+1}^f\text{-mod} \rightarrow H_n^f\text{-mod}. \quad (6.4)$$

We are going to make the Abelian category $\bigoplus_{n \geq 0} H_n^f\text{-mod}$ into a left $\mathcal{H}\text{eis}_{-l}(z, f_0^{-1})$ -module category, with \uparrow and \downarrow acting as induction and restriction, respectively. In order to do this, we need the *Mackey theorem* for H_n^f : there is an isomorphism of functors

$$\text{ind}_{n-1}^n \circ \text{res}_{n-1}^n \oplus \text{Id}^{\oplus l} \xrightarrow{\sim} \text{res}_n^{n+1} \circ \text{ind}_n^{n+1}. \quad (6.5)$$

The standard proof shows that the map

$$H_n^f \otimes_{H_{n-1}^f} H_n^f \oplus \bigoplus_{r=0}^{l-1} H_n^f \rightarrow H_{n+1}^f, \quad (u \otimes v, w_0, \dots, w_{l-1}) \mapsto u \tau_n v + \sum_{r=0}^{l-1} w_r x_{n+1}^r \quad (6.6)$$

is an isomorphism of (H_n^f, H_n^f) -bimodules. This implies that there is a unique (H_n^f, H_n^f) -bimodule homomorphism

$$\text{tr}_n^f : H_{n+1}^f \rightarrow H_n^f \quad (6.7)$$

such that $\text{tr}_n^f(\tau_n) = 0$ and $\text{tr}_n^f(x_{n+1}^r) = \delta_{r,0}$ for $0 \leq r < l$.

Lemma 6.1. *For any $n \geq 0$, we have that $\text{tr}_n^f(f(x_{n+1})) = 0$.*

Proof. For $u, v \in H_{n+1}^f$, write $u \equiv_n v$ as shorthand for $u = v$ in case $n = 0$, or $u - v \in H_n^f \tau_n H_n^f$ in case $n > 0$. We first show by induction on $n = 0, 1, \dots$ that

$$\tau_n \cdots \tau_1 x_1^a \tau_1 \cdots \tau_n \equiv_n \begin{cases} \sum_{\substack{b+c_1+\cdots+c_n=a \\ b>0, c_1, \dots, c_n \geq 0}} (\prod_{i \text{ with } c_i \neq 0} (-z^2 c_i)) x_{n+1}^b x_n^{c_n} \cdots x_1^{c_1} & \text{if } a > 0, \\ \sum_{\substack{b+c_1+\cdots+c_n=a \\ b \leq 0, c_1, \dots, c_n \leq 0}} (\prod_{i \text{ with } c_i \neq 0} (z^2 c_i)) x_{n+1}^b x_n^{c_n} \cdots x_1^{c_1} & \text{if } a \leq 0. \end{cases} \quad (6.8)$$

We explain this in detail in the case $a > 0$, since the case $a \leq 0$ is similar. The base case is trivial. For the induction step, using the relations depicted in (2.8)–(2.9), we have that

$$\begin{aligned} \tau_n x_n^a \tau_n &= \tau_n \tau_n^{-1} x_n^a \tau_n - z \sum_{\substack{b+c=a \\ b, c > 0}} \tau_n x_{n+1}^b x_n^c \\ &= x_{n+1}^a - z \sum_{\substack{b+c=a \\ b, c > 0}} \tau_n^{-1} x_{n+1}^b x_n^c - z^2 \sum_{\substack{b+c=a \\ b, c > 0}} x_{n+1}^b x_n^c \\ &\equiv_n x_{n+1}^a - z^2 \sum_{\substack{b+c+d=a \\ b, c, d > 0}} x_{n+1}^b x_n^{c+d} - z^2 \sum_{\substack{b+c=a \\ b, c > 0}} x_{n+1}^b x_n^c = x_{n+1}^a - z^2 \sum_{\substack{b+c=a \\ b, c > 0}} c x_{n+1}^b x_n^c. \end{aligned}$$

Now take the expression for $\tau_{n-1} \cdots \tau_1 x_1^a \tau_1 \cdots \tau_{n-1}$ given by the induction hypothesis, multiply on left and right by τ_n , and use the above identity plus the observation

$$\tau_n (H_{n-1}^f \tau_{n-1} H_{n-1}^f) \tau_n = H_{n-1}^f \tau_n \tau_{n-1} \tau_n H_{n-1}^f = H_{n-1}^f \tau_{n-1} \tau_n \tau_{n-1} H_{n-1}^f \subseteq H_n^f \tau_n H_n^f.$$

Finally, to deduce the lemma, we multiply (6.8) by f_{l-a} and sum over $a = 0, 1, \dots, l$ to show

$$\tau_n \cdots \tau_1 f(x_1) \tau_1 \cdots \tau_n \equiv_n f_l + \sum_{a=1}^l f_{l-a} \sum_{\substack{b+c_1+\cdots+c_n=a \\ b>0, c_1, \dots, c_n \geq 0}} \left(\prod_{i \text{ with } c_i \neq 0} (-z^2 c_i) \right) x_{n+1}^b x_n^{c_n} \cdots x_1^{c_1}.$$

The left-hand side is zero by the cyclotomic relation in H_{n+1}^f . The right-hand side is equal to $f(x_{n+1})$ plus terms in the kernel of tr_n^f . \square

Theorem 6.2. *There is a unique strict \mathbb{k} -linear monoidal functor*

$$\Psi_f : \mathcal{H}eis_{-l}(z, t) \rightarrow \mathcal{E}nd_{\mathbb{k}} \left(\bigoplus_{n \geq 0} H_n^f \text{-mod} \right)$$

sending the generating object \uparrow (resp. \downarrow) to the additive endofunctor that takes an H_n^f -module M to $\text{ind}_n^{n+1} M$ (resp. $\text{res}_{n-1}^n M$), and the generating morphisms x, τ, c and d to the natural transformations defined on the H_n^f -module M as follows:

- $\Psi_f(x)_M : H_{n+1}^f \otimes_{H_n^f} M \rightarrow H_{n+1}^f \otimes_{H_n^f} M, u \otimes v \mapsto ux_{n+1} \otimes v.$
- $\Psi_f(\tau)_M : H_{n+2}^f \otimes_{H_n^f} M \rightarrow H_{n+2}^f \otimes_{H_n^f} M, u \otimes v \mapsto u\tau_{n+1} \otimes v$ (where we have identified $\text{ind}_{n+1}^{n+2} \circ \text{ind}_n^{n+1}$ with ind_n^{n+2} in the obvious way).
- $\Psi_f(c)_M : M \rightarrow \text{res}_n^{n+1}(H_{n+1}^f \otimes_{H_n^f} M), v \mapsto 1 \otimes v$, i.e., it is the unit of the canonical adjunction making $(\text{ind}_n^{n+1}, \text{res}_n^{n+1})$ into an adjoint pair of functors.
- $\Psi_f(d)_M : H_n^f \otimes_{H_{n-1}^f} (\text{res}_{n-1}^n M) \rightarrow M, u \otimes v \mapsto uv$, i.e., it is the counit of the canonical adjunction making $(\text{ind}_{n-1}^n, \text{res}_{n-1}^n)$ into an adjoint pair of functors.

Proof. We use the presentation for $\mathcal{H}eis_{-l}(z, t)$ from [Definition 2.2](#). Let us first treat the case $l = 0$. In this case, the polynomial $f(w)$ from [\(6.1\)](#) is 1 and $t^2 = 1$. The category $\bigoplus_{n \geq 0} H_n^f\text{-mod}$ is simply the category of left \mathbb{k} -modules, and all of the induction and restriction functors are zero. Consequently, almost of the relations are trivially true. The only one that requires any thought is the relation $\bigcirc = (tz^{-1} - t^{-1}z^{-1})1_{\mathbb{1}}$ from [\(2.12\)](#). This holds because the scalar on the right-hand side is zero as $t^2 = 1$.

Henceforth, we assume that $l > 0$. Then $\mathcal{H}eis_{-l}(z, t)$ is generated by the objects \uparrow and \downarrow and morphisms x, τ, c and d subject to the relations [\(1.6\)–\(1.9\)](#), plus two more relations:

- (1) $\left[\begin{array}{c} \nearrow \\ \searrow \end{array} \cup \cup \hat{\cup} \dots \hat{\cup} \hat{\cup}^{l-1} \right]$ is invertible where $\sigma := \begin{array}{c} \nearrow \\ \searrow \end{array}$ is defined by [\(2.2\)](#).
- (2) $\bigcirc^l = tz^{-1}1_{\mathbb{1}}$ where $\gamma := \bigcirc$ is defined by [\(2.13\)](#), i.e., it is $-t^{-1}z^{-1}$ times the $(2, 1)$ -entry of the inverse of the matrix in (1).

The relations [\(1.6\)–\(1.9\)](#) are straightforward to check. On H_n^f -modules, $\Psi_f(\sigma)$ comes from the (H_n^f, H_n^f) -bimodule homomorphism $H_n^f \otimes_{H_{n-1}^f} H_n^f \rightarrow H_{n+1}^f, u \otimes v \mapsto u\tau_n v$. So we get the relation (1) since [\(6.6\)](#) is invertible by the proof of the Mackey theorem. Moreover, we see from [\(6.6\)](#) and the definition that $\Psi_f(\gamma)$ comes from the (H_n^f, H_n^f) -bimodule homomorphisms $-t^{-1}z^{-1} \text{tr}_n^f : H_{n+1}^f \rightarrow H_n^f$ for all $n \geq 0$. So for (2) we must show that $-t^{-1}z^{-1} \text{tr}_n^f(x_{n+1}^l) = tz^{-1}$. This follows from [Lemma 6.1](#) and the definition of tr_n^f , remembering that $t^2 = f_l$. \square

If we switch the roles of induction and restriction, we can reformulate [Theorem 6.2](#) in terms of Heisenberg categories of positive central charge. We prefer for this to replace the induction functor ind_n^{n+1} from before (which is the canonical left adjoint to restriction) with the *coinduction functor*

$$\text{coind}_n^{n+1} := \text{Hom}_{H_n^f}(H_{n+1}^f, -) : H_n^f\text{-mod} \rightarrow H_{n+1}^f\text{-mod} \quad (6.9)$$

which is its canonical right adjoint.

Theorem 6.3. *There is a unique strict \mathbb{k} -linear monoidal functor*

$$\Psi_f^{\vee} : \mathcal{H}eis_l(z, t^{-1}) \rightarrow \mathcal{E}nd_{\mathbb{k}}\left(\bigoplus_{n \geq 0} H_n^f\text{-mod}\right)$$

sending the generating object \uparrow (resp. \downarrow) to the additive endofunctor that takes an H_n^f -module M to $\text{res}_{n-1}^n M$ (resp. $\text{coind}_n^{n+1} M$), and the generating morphisms x, τ, c and d to the natural transformations defined on the H_n^f -module M as follows:

- $\Psi_f^\vee(x)_M : \text{res}_{n-1}^n M \rightarrow \text{res}_{n-1}^n M, v \mapsto x_n v$.
- $\Psi_f^\vee(\tau)_M : \text{res}_{n-2}^n M \rightarrow \text{res}_{n-2}^n M, v \mapsto -\tau_{n-1}^{-1} v$.
- $\Psi_f^\vee(c)_M : M \rightarrow \text{Hom}_{H_{n-1}^f}(H_n^f, \text{res}_{n-1}^n M), v \mapsto (u \mapsto uv)$, i.e., it is the unit of the canonical adjunction making $(\text{res}_{n-1}^n, \text{coind}_{n-1}^n)$ into an adjoint pair of functors.
- $\Psi_f^\vee(d)_M : \text{res}_n^{n+1}(\text{Hom}_{H_n^f}(H_{n+1}^f, M)) \rightarrow M, \theta \mapsto \theta(1)$, i.e., it is the counit of the canonical adjunction making $(\text{res}_n^{n+1}, \text{coind}_n^{n+1})$ into an adjoint pair of functors.

Proof. This may be proved directly in a similar way to the proof of [Theorem 6.2](#). One uses the presentation for $\mathcal{Heis}_l(z, t^{-1})$ from [Definition 3.1](#) instead of the one from [Definition 2.2](#), plus the Mackey isomorphism [\(6.6\)](#) and [Lemma 6.1](#) as before. We leave the details to the reader. \square

In fact, we have that $\text{ind}_n^{n+1} \cong \text{coind}_n^{n+1}$. This follows by the uniqueness of adjoints, since [Lemma 3.7](#) and [Theorem 6.2](#) (resp. [Theorem 6.3](#)) implies that ind_n^{n+1} is right adjoint to restriction as well as being left adjoint (resp. coind_n^{n+1} is left adjoint to restriction as well as being right adjoint). It follows that all three functors (induction, coinduction and restriction) send finitely generated projective modules to finitely generated projective modules. Hence:

Lemma 6.4. *The restrictions of the functors Ψ_f and Ψ_f^\vee constructed in Theorems 6.2–6.3 give strict \mathbb{k} -linear monoidal functors*

$$\Psi_f : \mathcal{Heis}_{-l}(z, t) \rightarrow \mathcal{End}_{\mathbb{k}}\left(\bigoplus_{n \geq 0} H_n^f\text{-pmod}\right), \quad \Psi_f^\vee : \mathcal{Heis}_l(z, t^{-1}) \rightarrow \mathcal{End}\left(\bigoplus_{n \geq 0} H_n^f\text{-pmod}\right),$$

where H_n^f -pmod denotes the category of finitely generated projective left H_n^f -modules.

7. Action on category \mathcal{O} for rational Cherednik algebras

The Heisenberg action on $\bigoplus_{n \geq 0} H_n^f$ -mod from [Theorem 6.2](#) can also be extended to an action on the category \mathcal{O} for rational Cherednik algebras, following an argument of Shan. To explain this in more detail, assume that $\mathbb{k} = \mathbb{C}$, and consider the complex reflection group $G(l, 1, n) \cong \mathfrak{S}_n \wr \mathbb{Z}/l\mathbb{Z}$ for $l \geq 1$, with reflection representation \mathbb{k}^n defined as in [\[Shan 2011, Section 3.1\]](#). Defining a rational Cherednik algebra requires a choice of parameters, for which there are a bewildering number of different parametrizations. We have:

- A single parameter $\kappa \in \mathbb{k}$, which is the parameter $k_{H,1}$ in [\[Ginzburg et al. 2003, Remark 3.2\]](#) for a reflecting hyperplane H on which the difference of two coordinates vanish.
- An l -tuple $(\kappa_1, \dots, \kappa_l) \in \mathbb{k}^l$ of parameters, which corresponds to the family $\{k_{H,i}\}_{0 \leq i \leq l}$ of parameters in [\[loc. cit., Remark 3.2\]](#) associated to a reflecting hyperplane H on which a single coordinate

vanishes so that $\kappa_i = k_{H,i}$. In [loc. cit.], it is assumed that $k_{H,0} = k_{H,l} = 0$, but adding a constant to all $k_{H,i}$ leaves the algebra unchanged. It is useful for us to incorporate an additional degree of freedom, so we drop the vanishing condition here: our parameter κ_l may be nonzero.

Let H_n be the rational Cherednik algebra attached to these parameters as in [loc. cit., Section 3].

Let $q := \exp(\sqrt{-1}\pi\kappa)$ and $q_i := \exp(\sqrt{-1}\pi(\kappa_i - i/\ell))$ for $i = 1, \dots, l$. One can relate these to the parameters in [Shan 2011] by choosing integers $e \geq 2$ and (s_1, \dots, s_l) then letting $\kappa := 1/e$ and $\kappa_i := \kappa s_i + i/\ell$, so $q_i = q^{s_i}$, for $i = 1, \dots, l$; note that the parameter q in [loc. cit.] is our q^2 . Let $\mathcal{O} = \mathcal{O}_{\kappa; \kappa_1, \dots, \kappa_l} := \bigoplus_{n \geq 0} \mathcal{O}_n$ where \mathcal{O}_n is the category of H_n -modules introduced in [Ginzburg et al. 2003, Section 3]. Also define

$$f(w) := \prod_{i=1}^l (w + q_i^2), \quad t := q_1 \cdots q_l.$$

By [Ginzburg et al. 2003, Theorem 5.16], there is an exact functor

$$KZ : \mathcal{O} \rightarrow \bigoplus_{n \geq 0} H_n^f\text{-mod}. \quad (7.1)$$

Note that this functor depends on a choice for each n of a basepoint in the subset of \mathbb{C}^n where all entries are distinct and nonzero. Different basepoints give isomorphic functors, but the isomorphism depends on the homotopy class of a path between the basepoints. For simplicity, we assume these basepoints are chosen to lie in the set $\{(b_1, \dots, b_n) \in \mathbb{R}^n \mid 0 < b_1 < \dots < b_n\}$. Since this is a contractible space, the resulting KZ functors are all canonically isomorphic, and there is no need for us to be more specific.

Matching with the formulae in [Ginzburg et al. 2003; Shan 2011] requires using the isomorphism from the cyclotomic Hecke algebra in [Shan 2011, Section 3.1] to ours that sends the generators T_0, T_1, \dots, T_{n-1} to $-x_1, q\tau_1, \dots, q\tau_{n-1}$. The Hecke algebra generators T_i ($i = 1, \dots, n-1$) in [Shan 2011] are of the form $-T$ for Hecke algebra generators T from [Ginzburg et al. 2003, Section 5.2.5] associated to reflections in the first type of hyperplane above. Also, T_0 is a scalar multiple (depending on the choice of κ_l) of the Hecke algebra generator T in [loc. cit., Section 5.2.5] associated to a reflection of the second type. The key point in all of this is that the minimal polynomials for x_1 and τ_i ($i = 1, \dots, n-1$) arising from the key formula in [loc. cit., Section 5.2.5] are $f(w)$ and $(w - q)(w + q^{-1})$ (up to scalars), i.e., we do indeed get defining relations of H_n^f .

The functor KZ is fully faithful on projectives [loc. cit., Theorem 5.16]. Moreover, it intertwines the Bezrukavnikov–Etingof induction and restriction functors denoted $\text{ind}_{b_{n+1}}$ and $\text{res}_{b_{n+1}}$ in [Shan 2011, Section 3.2] with the functors ind_n^{n+1} and res_n^{n+1} thanks to [loc. cit., Theorem 2.1]. These induction and restriction functors also depend on a choice of basepoint with a particular stabilizer, which following Shan we fix to be $(0, 0, \dots, 0, 1)$. (It would be more philosophically consistent with our previous conventions to say that whenever we choose a basepoint for restriction, we choose one of the form $(b_1, \dots, b_n) \in \mathbb{R}^n$ such that $0 \leq b_1 \leq b_2 \leq \dots \leq b_n$; whether we have equality or strict inequality depends on which stabilizer

we wish to have under the action of $G(l, 1, n)$. As before, all such choices give canonically isomorphic functors.)

Theorem 7.1. *There is a strict \mathbb{k} -linear monoidal functor*

$$\widehat{\Psi}_f : \mathcal{H}eis_{-l}(z, t) \rightarrow \mathcal{E}nd_{\mathbb{k}}(\mathcal{O}). \quad (7.2)$$

that makes \mathcal{O} into a module category over $\mathcal{H}eis_{-l}(z, t)$, with \uparrow and \downarrow acting as Bezrukavnikov–Etingof induction and restriction functors, respectively. This can be done in such a way that KZ is a morphism of $\mathcal{H}eis_{-l}(z, t)$ -module categories, viewing $\bigoplus_{n \geq 0} H_n^f$ -mod as a module category via the functor Ψ_f from Theorem 6.2.

Proof. Our argument is exactly as in the proof of [Shan 2011, Theorem 5.1] using [loc. cit., Lemma 2.4]. We need to show that there are certain natural transformations of functors satisfying specific relations. Theorem 6.2 allows us to define these on the image of the functor KZ via the action of $\mathcal{H}eis_{-l}(z, t)$. The full-faithfulness of KZ allows us to transfer this to an action on the full subcategory of projectives in \mathcal{O} . Since \mathcal{O} has enough projectives by [Ginzburg et al. 2003, Corollary 2.8], this action can be extended to an arbitrary object X by presenting X as the cokernel of a map between projectives. The resulting action is well-defined due to the fact that endomorphisms of an object lift to any projective resolution uniquely up to homotopy. \square

Remark 7.2. This quantum Heisenberg action is in many ways more convenient for working with category \mathcal{O} over Cherednik algebras than a Kac–Moody 2-category action, since the Heisenberg action requires no special assumptions on parameters. In fact, this action is still well defined if \mathbb{k} is replaced by a complete local ring, so one can extend the Heisenberg action to deformed category \mathcal{O} .

8. Categorical comultiplication

In this section, we construct the quantum analog of the categorical comultiplication from [Brundan et al. 2018, Theorem 5.4]. As discussed in [loc. cit., Theorem 1.3], the name “categorical comultiplication” derives from the relationship of this map to the usual comultiplication on the universal enveloping algebra of the Heisenberg Lie algebra. Since in the quantum case an explicit description of $K_0(\text{Kar}(\mathcal{H}eis_k(z, t)))$ analogous to that of [loc. cit., Theorem 1.1] is not available, we will not make a precise statement along these lines here, but we fully expect an analog of [loc. cit., Theorem 1.3] to hold in all situations where the Grothendieck ring has the expected form. As well as the quantum Heisenberg category $\mathcal{H}eis_k(z, t)$, we will work with $\mathcal{H}eis_l(z, u)$ and $\mathcal{H}eis_m(z, v)$ for $l, m \in \mathbb{Z}$ and $u, v \in \mathbb{k}^\times$ chosen so that

$$k = l + m, \quad t = uv. \quad (8.1)$$


To avoid confusion between these different categories, the reader will want to view the material in this section in color.

Let $\mathcal{H}eis_l(z, u) \odot \mathcal{H}eis_m(z, v)$ be the symmetric product of $\mathcal{H}eis_l(z, u)$ and $\mathcal{H}eis_m(z, v)$ as defined [Brundan et al. 2018, Section 3]. This is the strict \mathbb{k} -linear monoidal category defined by first taking


the free product of $\mathcal{Heis}_l(z, u)$ and $\mathcal{Heis}_m(z, v)$, i.e., the strict \mathbb{k} -linear monoidal category defined by the disjoint union of the given generators and relations of $\mathcal{Heis}_l(z, u)$ and of $\mathcal{Heis}_m(z, v)$, then adjoining isomorphisms $\sigma_{X,Y} : X \otimes Y \xrightarrow{\sim} Y \otimes X$ for each pair of objects $X \in \mathcal{Heis}_l(z, u)$ and $Y \in \mathcal{Heis}_m(z, v)$ subject to the relations

$$\begin{aligned}\sigma_{X_1 \otimes X_2, Y} &= (\sigma_{X_1, Y} \otimes 1_{X_2}) \circ (1_{X_1} \otimes \sigma_{X_2, Y}), & \sigma_{X_2, Y} \circ (f \otimes 1_Y) &= (1_Y \otimes f) \circ \sigma_{X_1, Y}, \\ \sigma_{X, Y_1 \otimes Y_2} &= (1_{Y_1} \otimes \sigma_{X, Y_2}) \circ (\sigma_{X, Y_1} \otimes 1_{Y_2}), & \sigma_{X, Y_2} \circ (1_X \otimes g) &= (g \otimes 1_X) \circ \sigma_{X, Y_1}\end{aligned}$$

for all $X, X_1, X_2 \in \mathcal{Heis}_l(z, u)$, $Y, Y_1, Y_2 \in \mathcal{Heis}_m(z, v)$ and $f : X_1 \rightarrow X_2$, $g : Y_1 \rightarrow Y_2$. Morphisms in $\mathcal{Heis}_l(z, u) \odot \mathcal{Heis}_m(z, v)$ are linear combinations of diagrams colored both blue and red. In these diagrams, as well as the generating morphisms of $\mathcal{Heis}_l(z, u)$ and $\mathcal{Heis}_m(z, v)$, we have the additional two-color crossings



which represent the isomorphisms $\sigma_{X,Y}$ for $X \in \{\uparrow, \downarrow\}$ and $Y \in \{\uparrow, \downarrow\}$, and their inverses



Definition 8.1. Given a diagram D representing a morphism in $\mathcal{Heis}_l(z, u) \odot \mathcal{Heis}_m(z, v)$ and two generic points in this diagram, one on a red string and the other on a blue string, we will denote the morphism represented by

$$(D \text{ with an extra dot at the red point}) - (D \text{ with an extra dot at the blue point})$$

by labeling the points with dots joined by a dotted line. For example

$$\begin{array}{c} \uparrow \quad \uparrow \\ \circ \cdots \circ \end{array} := \begin{array}{c} \uparrow \quad \uparrow \\ \circ \quad - \quad \circ \end{array} \quad (8.2)$$

Let $\mathcal{Heis}_l(z, u) \overline{\odot} \mathcal{Heis}_m(z, v)$ be the strict \mathbb{k} -linear monoidal category obtained by localizing at $\begin{array}{c} \uparrow \quad \uparrow \\ \circ \cdots \circ \end{array}$. This means that we adjoin a two-sided inverse to this morphism, which we denote as a dumbbell

$$\begin{array}{c} \uparrow \quad \uparrow \\ \circ - \circ \end{array} := \left(\begin{array}{c} \uparrow \quad \uparrow \\ \circ \cdots \circ \end{array} \right)^{-1}. \quad (8.3)$$

Just as explained in the degenerate case in [Brundan et al. 2018, Sections 4–5], all morphisms whose string diagram is that of an identity morphism with a horizontal dotted line joining two points of different colors are also automatically invertible in the localized category. We also denote the inverses of such morphisms by using a solid dumbbell in place of the dotted one. For instance

$$\begin{array}{c} \uparrow \quad \uparrow \quad | \quad \downarrow \quad \downarrow \\ \circ \quad - \quad \circ \end{array} = \begin{array}{c} \text{twisted crossing} \\ \text{diagram} \end{array} = \left(\begin{array}{c} \text{twisted crossing} \\ \text{diagram} \end{array} \right)^{-1} = \left(\begin{array}{c} \uparrow \quad \uparrow \quad | \quad \downarrow \quad \downarrow \\ \circ \cdots \circ \end{array} \right)^{-1}.$$

We also need the following morphisms, which we refer to as *internal bubbles*:

$$\text{Diagram 1} := z \sum_{a \geq 0} \text{Diagram 2} + z \text{Diagram 3}, \quad \text{Diagram 4} := z \sum_{a \geq 0} \text{Diagram 5} - a \text{Diagram 6} + z \text{Diagram 7}, \quad (8.4)$$

$$\text{Diagram 8} := z \sum_{a \geq 0} \text{Diagram 9} - z \text{Diagram 10}, \quad \text{Diagram 11} := z \sum_{a \geq 0} \text{Diagram 12} - a \text{Diagram 13} - z \text{Diagram 14}. \quad (8.5)$$

The category $\mathcal{Heis}_l(z, u) \overline{\odot} \mathcal{Heis}_m(z, v)$ possesses various symmetries which are often useful. Derived from (3.3), we have the strict \mathbb{k} -linear monoidal isomorphism

$$\Omega_{l|m} : \mathcal{Heis}_l(z, u) \overline{\odot} \mathcal{Heis}_m(z, v) \xrightarrow{\sim} (\mathcal{Heis}_{-l}(z, u^{-1}) \overline{\odot} \mathcal{Heis}_{-m}(z, v^{-1}))^{\text{op}}, \quad (8.6)$$

which takes a diagram to its mirror image in a horizontal plane multiplied by $(-1)^{x+y}$ where x is the number of one-colored crossings and y is the number of leftward cups and caps (including ones in (+)-, (-)- and internal bubbles). Also, we have

$$\text{flip} : \mathcal{Heis}_l(z, u) \overline{\odot} \mathcal{Heis}_m(z, v) \xrightarrow{\sim} \mathcal{Heis}_m(z, v) \overline{\odot} \mathcal{Heis}_l(z, u) \quad (8.7)$$

defined on diagrams by switching the colors blue and red then multiplying by $(-1)^z$ where z is the total number of dumbbells (both solid and dotted) in the picture. Finally, the category $\mathcal{Heis}_l(z, u) \overline{\odot} \mathcal{Heis}_m(z, v)$ is strictly pivotal, with duality functor

$$* : \mathcal{Heis}_l(z, u) \overline{\odot} \mathcal{Heis}_m(z, v) \xrightarrow{\sim} ((\mathcal{Heis}_l(z, u) \overline{\odot} \mathcal{Heis}_m(z, v))^{\text{op}})^{\text{rev}} \quad (8.8)$$

defined by rotating diagrams through 180° just like in (3.21).

We denote the duals of the internal bubbles (8.4)–(8.5) by

$$\text{Diagram 1}^*, \quad \text{Diagram 2}^*, \quad \text{Diagram 3}^*, \quad \text{Diagram 4}^*.$$

This definition ensures that internal bubbles commute past cups and caps in all possible configurations. For example

$$\text{Diagram 8}^* = \text{Diagram 12}, \quad \text{Diagram 11}^* = \text{Diagram 14}.$$

Again as in [Brundan et al. 2018, Sections 4–5], there are many other obvious commuting relations, such as

$$\text{Diagram 15} = \text{Diagram 16},$$

$$\text{Diagram 17} = \text{Diagram 18}, \quad \text{Diagram 19} = \text{Diagram 20},$$

$$\text{Diagram 21} = \text{Diagram 22},$$

$$\text{Diagram 23} = \text{Diagram 24},$$

$$\text{Diagram 25} = \text{Diagram 26},$$

$$\text{Diagram 27} = \text{Diagram 28}, \quad \text{Diagram 29} = \text{Diagram 30}.$$

$$\begin{array}{c} \text{Diagram 1} \\ \text{Diagram 2} \end{array} = \begin{array}{c} \text{Diagram 1} \\ \text{Diagram 2} \end{array}, \quad \begin{array}{c} \text{Diagram 3} \\ \text{Diagram 4} \end{array} = \begin{array}{c} \text{Diagram 4} \\ \text{Diagram 3} \end{array}, \quad \begin{array}{c} \text{Diagram 5} \\ \text{Diagram 6} \end{array} = \begin{array}{c} \text{Diagram 6} \\ \text{Diagram 5} \end{array},$$

as well as the mirror images of these under the symmetries $\Omega_{l|m}$, flip and $*$. We will appeal to all such relations below without further mention.

Here are some more interesting relations. The first shows how to “teleport” dots across dumbbells (plus a correction term):

$$\begin{array}{c} \text{Diagram 7} \end{array} = \begin{array}{c} \text{Diagram 8} \end{array} + \sum_{\substack{b+c=a-1 \\ b,c \geq 0}} \begin{array}{c} \text{Diagram 9} \end{array} - \sum_{\substack{b+c=a-1 \\ b,c < 0}} \begin{array}{c} \text{Diagram 10} \end{array} \quad (8.9)$$

for any $a \in \mathbb{Z}$. We also have the following relations to commute dumbbells past one-color crossings:

$$\begin{array}{c} \text{Diagram 11} \end{array} = \begin{array}{c} \text{Diagram 12} \end{array} + z \begin{array}{c} \text{Diagram 13} \end{array}, \quad (8.10)$$

$$\begin{array}{c} \text{Diagram 14} \end{array} = \begin{array}{c} \text{Diagram 15} \end{array} + z \begin{array}{c} \text{Diagram 16} \end{array}, \quad (8.11)$$

$$\begin{array}{c} \text{Diagram 17} \end{array} = \begin{array}{c} \text{Diagram 18} \end{array} + z \begin{array}{c} \text{Diagram 19} \end{array}, \quad (8.12)$$

$$\begin{array}{c} \text{Diagram 20} \end{array} = \begin{array}{c} \text{Diagram 21} \end{array} + z \begin{array}{c} \text{Diagram 22} \end{array}, \quad (8.13)$$

These are all straightforward to prove: one first cancels the solid dumbbells by composing on the top and bottom with their inverses then uses the affine Hecke algebra relations (1.6)–(1.7) to commute dots past crossings in the result. For example, to prove the first relation in (8.10), we have

$$\begin{array}{c} \text{Diagram 23} \end{array} = \begin{array}{c} \text{Diagram 24} \end{array} - \begin{array}{c} \text{Diagram 25} \end{array} \stackrel{(1.6)}{=} \begin{array}{c} \text{Diagram 26} \end{array} - \begin{array}{c} \text{Diagram 27} \end{array} + z \begin{array}{c} \text{Diagram 28} \end{array} = \begin{array}{c} \text{Diagram 29} \end{array} + z \begin{array}{c} \text{Diagram 30} \end{array}.$$

We then compose on the top with a solid dumbbell connecting the red strand and the leftmost blue strand, and compose on the bottom with a solid dumbbell connecting the red strand and the rightmost blue strand.

The following seven lemmas are the quantum analogs of [Brundan et al. 2018, Lemmas 5.6–5.12]. Their proofs are quite similar to the degenerate case.

Lemma 8.2. *We have that*

$$\begin{array}{c} \text{Diagram 31} \end{array} = - \left(\begin{array}{c} \text{Diagram 32} \end{array} \right)^{-1}.$$

Lemma 8.3. *For any $a \in \mathbb{Z}$, we have that*

$$\begin{array}{c} \text{Diagram 33} \end{array} + \begin{array}{c} \text{Diagram 34} \end{array} = z \sum_{\substack{b \in \mathbb{Z} \\ b < a \text{ or } b > 0}} \begin{array}{c} \text{Diagram 35} \end{array} - z \sum_{0 \leq b \leq a} \begin{array}{c} \text{Diagram 36} \end{array}.$$

Lemma 8.4. *The following relations hold:*

$$\begin{aligned} \text{Diagram 1} &= \text{Diagram 2} + z^2 \text{Diagram 3} - z^2 \sum_{\substack{a \geq 0 \\ b \geq 0}} \text{Diagram 4}_a \text{Diagram 5}_b, \\ \text{Diagram 1} &= \text{Diagram 2} + z^2 \text{Diagram 3} - z^2 \sum_{\substack{a \geq 0 \\ b > 0}} \text{Diagram 4}_a \text{Diagram 5}_b. \end{aligned}$$

Lemma 8.5. *We have that*

$$\text{Diagram 6} = z \text{Diagram 7} - z^2 \sum_{\substack{a \geq 0 \\ b \in \mathbb{Z}}} \text{Diagram 8}_a \text{Diagram 9}_b.$$

Lemma 8.6. *We have that*

$$\text{Diagram 10} = \text{Diagram 11} - tz \text{Diagram 12} - z^2 \text{Diagram 13} + z^3 \sum_{\substack{a, b > 0 \\ c \in \mathbb{Z}}} \text{Diagram 14}_{-b-c} \text{Diagram 15}_{-a+c} \text{Diagram 16}_a.$$

Lemma 8.7. *We have that*

$$\text{Diagram 17} = z^2 \sum_{\substack{a, b > 0 \\ c \in \mathbb{Z}}} \text{Diagram 18}_a \text{Diagram 19}_b - t \text{Diagram 20}.$$

Lemma 8.8. *We have that*

$$\text{Diagram 21} = \text{Diagram 22} + z^2 \text{Diagram 23}.$$

Using these, we can prove the main theorem of the section:

Theorem 8.9. *For $k = l + m$ and $t = uv$, there is a unique strict \mathbb{k} -linear monoidal functor*

$$\Delta_{l|m} : \mathcal{H}eis_k(z, t) \rightarrow \text{Add}(\mathcal{H}eis_l(z, u) \overline{\odot} \mathcal{H}eis_m(z, v))$$

such that $\uparrow \mapsto \uparrow \oplus \uparrow$, $\downarrow \mapsto \downarrow \oplus \downarrow$, and on morphisms

$$\text{Diagram 24} \mapsto \text{Diagram 25} + \text{Diagram 26}, \tag{8.14}$$

$$\text{Diagram 27} \mapsto \text{Diagram 28} + \text{Diagram 29} + q \text{Diagram 30} + q \text{Diagram 31} + z \text{Diagram 32} - z \text{Diagram 33} + z \text{Diagram 34} + z \text{Diagram 35}, \tag{8.15}$$

$$\text{Diagram 36} \mapsto \text{Diagram 37} + \text{Diagram 38} + q^{-1} \text{Diagram 39} + q^{-1} \text{Diagram 40} + z \text{Diagram 41} - z \text{Diagram 42} + z \text{Diagram 43}, \tag{8.16}$$

$$\text{Diagram 44} \mapsto \text{Diagram 45} + \text{Diagram 46}, \quad \text{Diagram 47} \mapsto \text{Diagram 48} + \text{Diagram 49}. \tag{8.17}$$

Moreover, we have that

$$\Delta_{l|m}(\text{---}) = \text{---} + \text{---}, \quad \Delta_{l|m}(\text{---}) = -\text{---} - \text{---}. \quad (8.18)$$

Also, the following hold for all $a \in \mathbb{Z}$:

$$\Delta_{\textcolor{blue}{I}|\textcolor{red}{m}}(\textcolor{blue}{\oplus} a) = z \sum_{b \in \mathbb{Z}} \frac{\textcolor{blue}{\oplus} b}{\textcolor{blue}{\oplus} a-b}, \quad \Delta_{\textcolor{blue}{I}|\textcolor{red}{m}}(a \textcolor{blue}{\oplus}) = -z \sum_{b \in \mathbb{Z}} \frac{b \textcolor{blue}{\oplus}}{a-b \textcolor{blue}{\oplus}}, \quad (8.19)$$

$$\Delta_{\textcolor{blue}{I}|\textcolor{red}{m}}(\bigodot a) = -z \sum_{b \in \mathbb{Z}} \frac{\bigodot b}{\bigodot a-b}, \quad \Delta_{\textcolor{blue}{I}|\textcolor{red}{m}}(a \bigodot) = z \sum_{b \in \mathbb{Z}} \frac{b \bigodot}{a-b \bigodot}. \quad (8.20)$$

Equivalently, in terms of the generating functions (3.14)–(3.17) and their analogs in $\mathcal{H}eis_l(z, u)$ and $\mathcal{H}eis_m(z, v)$

$$\Delta_{l|m}(\oplus(w)) = \textcolor{blue}{\oplus}(w) \textcolor{red}{\oplus}(w), \quad \Delta_{l|m}(\textcolor{red}{\oplus}(w)) = \textcolor{blue}{\oplus}(w) \textcolor{red}{\oplus}(w), \quad (8.21)$$

$$\Delta_{l|m}(\bigodot(w)) = \textcolor{blue}{\bigodot}(w) \textcolor{red}{\bigodot}(w), \quad \Delta_{l|m}(\bigotimes(w)) = \textcolor{blue}{\bigotimes}(w) \textcolor{red}{\bigotimes}(w). \quad (8.22)$$

Remark 8.10. For the proof, it is helpful to notice that $\text{flip} \circ \Delta_{l|m} = \Delta_{m|l}$ (on extending flip to the additive envelopes in the obvious way). However, $\Delta_{l|m}$ does not commute with either of the other symmetries Ω or $*$. In fact, the map $\Omega_{-l|-m} \circ \Delta_{-l|-m} \circ \Omega_k$ would be an equally good alternative choice for the categorical comultiplication map. The only change to the above formulae if one uses this alternative is that one needs to replace q with $-q^{-1}$ in (8.15)–(8.16); this is the “Galois symmetry” in the choice of the root q of the equation $x^2 - zx - 1 = 0$.

Proof. In view of the uniqueness from Lemma 4.3, we may take (8.14)–(8.18) as the definition of $\Delta_{l|m}$ on generating morphisms, and must check that the images of the relations (1.6)–(1.9) and (4.1)–(4.4) from Definition 4.1 are all satisfied in $\text{Add}(\mathcal{H}eis_l(z, u) \bar{\odot} \mathcal{H}eis_m(z, v))$; we must also check (8.19)–(8.20). The details are sufficiently similar to the degenerate case from the proof of [Brundan et al. 2018, Theorem 5.4] that we only sketch the steps needed below.

First one checks (1.6)–(1.8). For example, to check the skein relation, the image under $\Delta_{l|m}$ of  is $A + \text{flip}(A)$ where

$$A := \left(\begin{array}{c} \nearrow \searrow \\ \diagup \diagdown \end{array} - \begin{array}{c} \nearrow \searrow \\ \diagdown \diagup \end{array} \right) + z \left(\begin{array}{c} \uparrow \uparrow \\ \circ \circ \circ \end{array} - \begin{array}{c} \uparrow \uparrow \\ \circ \circ \circ \end{array} \right) + z \left(\begin{array}{c} \nearrow \searrow \\ \diagup \diagdown \end{array} + \begin{array}{c} \nearrow \searrow \\ \diagdown \diagup \end{array} - \begin{array}{c} \nearrow \searrow \\ \diagup \diagdown \end{array} \right).$$

Using the skein relation in $\mathcal{Heis}_l(z, u)$ plus (8.9), A simplifies to $B := z \uparrow \uparrow + z \uparrow \uparrow$. This is what is required since the image under $\Delta_{l|m}$ of $z \uparrow \uparrow$ is $B + \text{flip}(B)$. The other relations here are checked by similarly explicit calculations. The one for the braid relation is rather long.

The relation (1.9) is easy.

To check (8.19)–(8.20), we assume to start with that $k \geq 0$. Consider the clockwise (+)-bubble $a \oplus$. When $a \leq 0$, this is just a scalar (usually zero) due to (3.11) and the assumption $k \geq 0$, and the relation

to be checked is trivial. So assume that $a > 0$. Then $a \oplus = a \circlearrowleft$, hence, its image under $\Delta_{l|m}$ is $-a \circlearrowleft - a \circlearrowright$, which is indeed equal to $-z \sum_{b \in \mathbb{Z}} b \oplus a-b \oplus$ by Lemma 8.3. This establishes the left-hand identity in (8.19), hence, the left-hand identity in (8.21). The right-hand identity in (8.21) then follows using (3.13), thereby establishing the right-hand identity in (8.19) as well. Next, consider the clockwise $(-)$ -bubble $a \ominus$. This time the relation to be checked is trivial when $a \geq 0$, so assume that $a < 0$. Then, using the assumption $k \geq 0$ again, we have that $a \ominus = a \circlearrowright$, hence, its image under $\Delta_{l|m}$ is $-a \circlearrowleft - a \circlearrowright$, which is equal to $z \sum_{b \in \mathbb{Z}} b \ominus a-b \ominus$ by Lemma 8.3 (noting when $a < 0 \leq k$ that the term involving $(+)$ -bubbles is zero). Then we complete the proof of (8.20) using the equivalent form (8.22) and (3.13) once again. It remains to treat $k \leq 0$. This follows by similar arguments; one starts by considering the counterclockwise $(+)$ - and $(-)$ -bubbles using the identities obtained by applying $\Omega_{l|m}$ to Lemma 8.3, then gets the clockwise ones using (3.13).

Consider (4.3)–(4.4). The relations involving bubbles follow easily from (8.19)–(8.20). Next consider the right curl relation in (4.3), so $k \geq 0$. Applying $\Delta_{l|m}$ to the relation reveals that we must show that $A + \text{flip}(A) = B + \text{flip}(B)$ where

$$A := z \circlearrowleft \circlearrowleft \circlearrowleft - \circlearrowright \circlearrowright \circlearrowright, \quad B := \delta_{k,0} t^{-1}.$$

This follows from Lemma 8.5, noting that the only nonzero term in the summation on the right-hand side of that identity is the one with $a = b = 0$ due to the assumption that $k \geq 0$. The argument for the left curl in (4.4) is entirely similar; it uses the identity obtained by applying $* \circ \Omega_{l|m}$ to Lemma 8.5.

Finally, one must check (4.1)–(4.2). This is a calculation just like in the final paragraph of the proof of [Brundan et al. 2018, Theorem 5.4]; ultimately one uses Lemmas 8.6–8.8. \square

9. Generalized cyclotomic quotients

In this section, we define some \mathbb{k} -linear categories, namely, the generalized cyclotomic quotients of $\mathcal{H}eis_k(z, t)$. Recall that $x = \uparrow$ and $y = \downarrow$.

Definition 9.1. Suppose we are given monic polynomials

$$f(w) = f_0 w^l + f_1 w^{l-1} + \cdots + f_l \in \mathbb{k}[w], \quad (9.1)$$

$$g(w) = g_0 w^m + g_1 w^{m-1} + \cdots + g_m \in \mathbb{k}[w] \quad (9.2)$$

such that $k = m - l$ and $t^2 = f_l/g_m$. Define

$$\mathbb{O}^+(w) = t^{-1} z \sum_{n \in \mathbb{Z}} \mathbb{O}_n^+ w^{-n} := g(w)/f(w) \in w^k + w^{k-1} \mathbb{k}[[w^{-1}]], \quad (9.3)$$

$$\widetilde{\mathbb{O}}^+(w) = -tz \sum_{n \in \mathbb{Z}} \widetilde{\mathbb{O}}_n^+ w^{-n} := f(w)/g(w) \in w^{-k} + w^{-k-1} \mathbb{k}[[w^{-1}]], \quad (9.4)$$

$$\mathbb{O}^-(w) = -tz \sum_{n \in \mathbb{Z}} \mathbb{O}_n^- w^{-n} := t^2 g(w)/f(w) \in 1 + w \mathbb{k}[[w]], \quad (9.5)$$

$$\tilde{\mathbb{O}}^-(w) = t^{-1}z \sum_{n \in \mathbb{Z}} \tilde{\mathbb{O}}_n^- w^{-n} := t^{-2} f(w)/g(w) \in 1 + w\mathbb{k}[[w]]; \quad (9.6)$$

cf. (3.14)–(3.17). Let $\mathcal{I}(f|g)$ be the left tensor ideal generated by the morphisms

$$\{f(x), \bigoplus_{n=-k}^l \mathbb{O}_n^+ \mathbb{1} \mid -k < n < l\}. \quad (9.7)$$

The *generalized cyclotomic quotient* associated to the polynomials $f(w)$ and $g(w)$ is the quotient category

$$\mathcal{H}(f|g) := \mathcal{H}eis_k(z, t)/\mathcal{I}(f|g). \quad (9.8)$$

It is a module category over $\mathcal{H}eis_k(z, t)$.

The following is the quantum analog of [Brundan 2018, Lemma 1.8]; see also [Brundan and Davidson 2017, Lemma 4.14] for the analog in the setting of Kac–Moody 2-categories.

Lemma 9.2. *In the setup of Definition 9.1, $\mathcal{I}(f|g)$ may be defined equivalently as the left tensor ideal generated by*

$$\{g(y), \bigoplus_{n=k}^m \tilde{\mathbb{O}}_n^+ \mathbb{1} \mid k < n < m\}. \quad (9.9)$$

Moreover, it contains $\bigoplus_{n=-k}^l \mathbb{O}_n^+ \mathbb{1}$, $\bigoplus_{n=-k}^l \mathbb{O}_n^- \mathbb{1}$, $\bigoplus_{n=k}^m \tilde{\mathbb{O}}_n^+ \mathbb{1}$ and $\bigoplus_{n=k}^m \tilde{\mathbb{O}}_n^- \mathbb{1}$ for all $n \in \mathbb{Z}$.

Proof. For morphisms $\theta, \phi : X \rightarrow Y$, we will write $\theta \equiv \phi$ as shorthand for $\theta - \phi \in \mathcal{I}(f|g)$. By (3.11)–(3.12), we have automatically that $\bigoplus_{n=-k}^l \mathbb{O}_n^+ \mathbb{1} \equiv \mathbb{O}_n^+ \mathbb{1}$ when $n \leq -k$, $\bigoplus_{n=k}^m \tilde{\mathbb{O}}_n^+ \mathbb{1} \equiv \tilde{\mathbb{O}}_n^+ \mathbb{1}$ when $n \leq k$, $\bigoplus_{n=0}^l \mathbb{O}_n^- \mathbb{1} \equiv \mathbb{O}_n^- \mathbb{1}$ when $n \geq 0$, and $\bigoplus_{n=0}^m \tilde{\mathbb{O}}_n^- \mathbb{1} \equiv \tilde{\mathbb{O}}_n^- \mathbb{1}$ when $n \geq 0$.

In this paragraph, we use ascending induction on n to show that $\bigoplus_{n=-k}^l \mathbb{O}_n^+ \mathbb{1} \equiv \mathbb{O}_n^+ \mathbb{1}$ for all $n \in \mathbb{Z}$. This is immediate from (9.7) if $n < l$, so assume that $n \geq l$. The fact that $f(x) \equiv 0$ implies that

$$\sum_{a=0}^l f_a \bigoplus_{n=a}^l \mathbb{O}_n^+ \mathbb{1} + \sum_{a=0}^l f_a \bigoplus_{n=a}^l \mathbb{O}_n^- \mathbb{1} = \sum_{a=0}^l f_a \bigcirc_{n=a}^l \mathbb{O}_n^+ \mathbb{1} \equiv 0.$$

On the left-hand side of this, the only nonzero $(-)$ -bubble arises when $n = a = l$, so it shows that $\sum_{a=0}^l f_a \bigoplus_{n=a}^l \mathbb{O}_n^+ \mathbb{1} \equiv \delta_{l,n} f_l t^{-1} z^{-1} \mathbb{1}$. Using the induction hypothesis and $f_l = g_m t^2$, we deduce that $\bigoplus_{n=-k}^l \mathbb{O}_n^+ \mathbb{1} \equiv \delta_{l,n} g_m t z^{-1} \mathbb{1}$. Equating w^{l-n} -coefficients in $f(w) \mathbb{O}^+(w) = g(w)$, we get that $\sum_{a=0}^l f_a \mathbb{O}_{n-a}^+ \mathbb{1} \equiv \delta_{l,n} g_m t z^{-1} \mathbb{1}$. Hence, $\bigoplus_{n=-k}^l \mathbb{O}_n^+ \mathbb{1} \equiv \mathbb{O}_n^+ \mathbb{1}$ as claimed.

Next, we show by descending induction on n that $\bigoplus_{n=-k}^l \mathbb{O}_n^- \mathbb{1} \equiv \mathbb{O}_n^- \mathbb{1}$ for all $n \in \mathbb{Z}$. We may assume that $n < 0$. Equating w^{-n} -coefficients in $f(w) \mathbb{O}^+(w) = t^{-2} f(w) \mathbb{O}^-(w)$ gives that

$$\sum_{a=0}^l f_{l-a} \mathbb{O}_{a+n}^+ \mathbb{1} = - \sum_{a=0}^l f_{l-a} \mathbb{O}_{a+n}^- \mathbb{1}.$$

Using the induction hypothesis plus the previous paragraph, we deduce that

$$\sum_{a=0}^l f_{l-a} \bigoplus_{n=a}^l \mathbb{O}_n^+ \mathbb{1} + f_l \mathbb{O}_n^- \mathbb{1} + \sum_{a=1}^l f_{l-a} \bigoplus_{n=a}^l \mathbb{O}_n^- \mathbb{1} \equiv 0.$$

But also from $f(x) \equiv 0$ we get that

$$\sum_{a=0}^l f_{l-a} \oplus_{a+n} + \sum_{a=0}^l f_{l-a} \ominus_{a+n} = \sum_{a=0}^l f_{l-a} \odot_{a+n} \equiv 0.$$

Taking the difference of these two identities establishes the induction step.

Using the notation of (3.14)–(3.17), we have now shown that $\oplus(w) \equiv \mathbb{O}^\pm(w)\mathbb{1}$. Taking inverses using (3.13), we deduce that $\oplus(w) \equiv \widetilde{\mathbb{O}}^\pm(w)\mathbb{1}$. Hence, $n \oplus \equiv \widetilde{\mathbb{O}}_n^\pm \mathbb{1}$ for all $n \in \mathbb{Z}$. So we have established the last assertion from the lemma.

Equating w^b -coefficients in $g(w) = f(w)\mathbb{O}^+(w)$ shows that $g_{m-b} = t^{-1}z \sum_{a=0}^l f_{l-a} \mathbb{O}_{a-b}^+$. Hence

$$g(y) = \sum_{a=0}^l t^{-1} f_{l-a} \left(z \sum_{b \geq 0} \begin{array}{c} b \\ \oplus \\ a-b \end{array} \right) \stackrel{(4.17)}{=} \sum_{a=0}^l t^{-1} f_{l-a} \begin{array}{c} \oplus \\ a \end{array} \equiv 0.$$

We have now shown that $\mathcal{I}(f|g)$, the left tensor ideal generated by (9.7), contains (9.9). Similarly, the left tensor ideal generated by (9.9) contains (9.7). This completes the proof. \square

We assume for the rest of the section that \mathbb{k} is a field, and that we are given a factorization $t = uv^{-1}$ for $u, v \in \mathbb{k}^\times$ such that $u^2 = f_l$ and $v^2 = g_m$. Let $\mathcal{V}(f)$ and $\mathcal{V}(g)^\vee$ denote $\bigoplus_{n \geq 0} H_n^f$ -pmod and $\bigoplus_{n \geq 0} H_n^g$ -pmod viewed as a module categories over $\mathcal{Heis}_{-l}(z, u)$ and $\mathcal{Heis}_m(z, v^{-1})$ via the monoidal functors Ψ_f and Ψ_g^\vee from Lemma 6.4. Let

$$\mathcal{V}(f|g) := \mathcal{V}(f) \boxtimes \mathcal{V}(g)^\vee \tag{9.10}$$

be their linearized Cartesian product, i.e., the \mathbb{k} -linear category with objects that are pairs (X, Y) for $X \in \mathcal{V}(f)$, $Y \in \mathcal{V}(g)^\vee$, and morphisms

$$\text{Hom}_{\mathcal{V}(f|g)}((X, Y), (U, V)) := \text{Hom}_{\mathcal{V}(f)}(X, U) \otimes \text{Hom}_{\mathcal{V}(g)^\vee}(Y, V)$$

with the obvious composition law. There is an equivalence of categories

$$\mathcal{V}(f|g) \rightarrow \bigoplus_{r,s \geq 0} (H_r^f \otimes H_s^g)\text{-pmod},$$

hence, $\mathcal{V}(f|g)$ is additive Karoubian. Moreover, $\mathcal{V}(f|g)$ is a module category over the symmetric product $\mathcal{Heis}_{-l}(z, u) \odot \mathcal{Heis}_m(z, v^{-1})$.

Lemma 9.3. *Let V be a finite-dimensional AH_2 -module. All eigenvalues of x_2 on V are of the form $\lambda, q^2\lambda$ or $q^{-2}\lambda$ for eigenvalues λ of x_1 on V .*

Proof. We may assume for the proof that \mathbb{k} is algebraically closed. Suppose that $v \in V$ is a simultaneous eigenvector for the commuting operators x_1 and x_2 of eigenvalues λ_1 and λ_2 , respectively. If $\tau_1 v = q v$ (resp. $\tau_1 v = -q^{-1}v$) then $\lambda_2 = q^2\lambda_1$ (resp. $\lambda_2 = q^{-2}\lambda_1$), as follows easily from the relation $x_2(\tau_1 - z)v = \tau_1 x_1 v$. Otherwise, v and $\tau_1 v$ are linearly independent, in which case the matrix describing the action of x_1 on the subspace with basis $\{v, \tau_1 v\}$ is $\begin{pmatrix} \lambda_1 & -z\lambda_2 \\ 0 & \lambda_2 \end{pmatrix}$. So λ_2 is another eigenvalue of x_1 on V . \square

Lemma 9.4. *Assume that $f(w)$ and $g(w)$ split as products of linear factors in $\mathbb{k}[w]$, and moreover assume that $\lambda\mu^{-1} \notin \{q^{2i} \mid i \in \mathbb{Z}\}$ for all roots λ of $f(w)$ and μ of $g(w)$. Then the categorical action of $\mathcal{Heis}_{-l}(z, u) \odot \mathcal{Heis}_m(z, v^{-1})$ on $\mathcal{V}(f|g)$ defined above extends to an action of the localization $\mathcal{Heis}_{-l}(z, u) \overline{\odot} \mathcal{Heis}_m(z, v^{-1})$ from Definition 8.1.*

Proof. Lemma 9.3 implies that the eigenvalues of x_1, \dots, x_n on any finite-dimensional H_n^f -module are of the form $q^{2i}\lambda$ for $i \in \mathbb{Z}$ and a root λ of $f(w)$. Consequently, the commuting endomorphisms defined by evaluating $\uparrow \downarrow \phi$ and $\phi \uparrow \downarrow$ on an object of $\mathcal{V}(f|g)$ have eigenvalues contained in the sets $\{q^{2i}\lambda \mid i \in \mathbb{Z}, \lambda \text{ a root of } f(w)\}$ and $\{q^{2j}\mu \mid j \in \mathbb{Z}, \mu \text{ a root of } g(w)\}$, respectively. By the genericity assumption, these sets are disjoint, hence, all eigenvalues of the endomorphism defined by $\uparrow \cdots \downarrow \phi = \uparrow \downarrow \phi - \uparrow \phi \uparrow$ lie in \mathbb{k}^\times . Consequently, this endomorphism is invertible. \square

Lemma 9.4 shows for suitably generic $f(w), g(w)$ that there is a strict \mathbb{k} -linear monoidal functor $\Psi_f \overline{\odot} \Psi_g^\vee : \mathcal{Heis}_{-l}(z, u) \overline{\odot} \mathcal{Heis}_m(z, v^{-1}) \rightarrow \mathcal{End}_\mathbb{k}(\mathcal{V}(f|g))$. Composing this functor with the functor $\Delta_{-l|m}$ from Theorem 8.9, we obtain a strict \mathbb{k} -linear monoidal functor

$$\Psi_{f|g} := \Psi_f \overline{\odot} \Psi_g^\vee \circ \Delta_{-l|m} : \mathcal{Heis}_k(z, t) \rightarrow \mathcal{End}_\mathbb{k}(\mathcal{V}(f|g)). \quad (9.11)$$

Thus, we have made $\mathcal{V}(f|g)$ into a module category over $\mathcal{Heis}_k(z, t)$.

Theorem 9.5. *Assume that $f(w), g(w)$ satisfy the genericity assumption from Lemma 9.4 so that (9.11) is defined. Let $\text{Ev} : \mathcal{End}_\mathbb{k}(\mathcal{V}(f|g)) \rightarrow \mathcal{V}(f|g)$ be the \mathbb{k} -linear functor defined by evaluation on $S := (H_0^f, H_0^g) \in \mathcal{V}(f|g)$. The composition $\text{Ev} \circ \Psi_{f|g}$ factors through the generalized cyclotomic quotient $\mathcal{H}(f|g)$ to induce an equivalence of $\mathcal{Heis}_k(z, t)$ -module categories*

$$\psi_{f|g} : \text{Kar}(\mathcal{H}(f|g)) \rightarrow \mathcal{V}(f|g).$$

Proof. We first show that $\Psi_{f|g}(\oplus(w))_S \in w^k \text{End}(S)[[w^{-1}]]$ equals $\oplus^+(w)1_S$. Recalling that $\oplus^+(w)$ is the expansion at $w = \infty$ of the rational function $g(w)/f(w)$, this follows because

$$\Psi_{f|g}(\oplus(w))_S = \Psi_f(\oplus(w))_{H_0^f} \otimes \Psi_g^\vee(\oplus(w))_{H_0^g}$$

thanks to (8.21), and also $\Psi_f(\oplus(w))_{H_0^f} = 1/f(w)$ and $\Psi_g^\vee(\oplus(w))_{H_0^g} = g(w)$. To see the last two assertions, we first apply Lemma 9.2 to see that $\mathcal{I}(f|1)$, the left tensor ideal of $\mathcal{Heis}_{-l}(z, u)$ generated by $f(x)$, contains all coefficients of the series $\oplus(w) - 1/f(w)\mathbb{1}$; all elements of this ideal act as zero on H_0^f since its generator $f(x)$ acts as zero. Then we apply Lemma 9.2 again to see that $\mathcal{I}(1|g)$, the left tensor ideal of $\mathcal{Heis}_m(z, v^{-1})$ generated by $g(y)$, contains all coefficients of $\oplus(w) - g(w)\mathbb{1}$; all elements of this act as zero on H_0^g .

The previous paragraph shows that $\oplus_n - \oplus_n^+ \mathbb{1}$ acts as zero on S for all $n \in \mathbb{Z}$. Also it is obvious that $f(x)$ acts as zero on S . So the left tensor ideal $\mathcal{I}(f|g)$ acts as zero on S , which proves that $\text{Ev} \circ \Psi_{f|g}$ factors through the quotient $\mathcal{H}(f|g) = \mathcal{Heis}_k(z, t)/\mathcal{I}(f|g)$ to induce a \mathbb{k} -linear functor $\mathcal{H}(f|g) \rightarrow \mathcal{V}(f) \boxtimes \mathcal{V}(g)^\vee$. Since $\mathcal{V}(f|g)$ is additive Karoubian, this extends to the Karoubi envelope to induce the functor $\psi_{f|g}$ from the statement of the theorem. Moreover, it is automatic from the definition that $\psi_{f|g}$ is a morphism

of $\mathcal{H}eis_k(z, t)$ -module categories. It just remains to show that $\psi_{f|g}$ is an equivalence, which we do by showing that it is full, faithful and dense.

First we show that $\psi_{f|g}$ is full and faithful. It suffices to check this on objects $X = X_r \otimes \cdots \otimes X_1$ and $Y = Y_s \otimes \cdots \otimes Y_1$ that are words in \uparrow and \downarrow . We assume moreover that $k \geq 0$; a similar argument with the roles of \uparrow and \downarrow interchanged does the job when $k \leq 0$ too. Let $X^* = X_1^* \otimes \cdots \otimes X_r^*$ be the dual object (here, $\uparrow^* = \downarrow$, $\downarrow^* = \uparrow$). By rigidity, we have a canonical isomorphism $\text{Hom}_{\mathcal{H}(f|g)}(X, Y) \cong \text{Hom}_{\mathcal{H}(f|g)}(\mathbb{1}, X^* \otimes Y)$, from which we get a commuting diagram

$$\begin{array}{ccc} \text{Hom}_{\mathcal{H}(f|g)}(X, Y) & \xrightarrow{\sim} & \text{Hom}_{\mathcal{H}(f|g)}(\mathbb{1}, X^* \otimes Y) \\ \psi_{f|g} \downarrow & & \downarrow \psi_{f|g} \\ \text{Hom}_{\mathcal{V}(f|g)}(X \otimes S, Y \otimes S) & \xrightarrow{\sim} & \text{Hom}_{\mathcal{V}(f|g)}(S, X^* \otimes Y \otimes S). \end{array}$$

The left-hand vertical map in this diagram is an isomorphism if and only if the right-hand vertical map is one. We claim that the left-hand vertical map is an isomorphism when $X = Y = \uparrow^{\otimes n}$. To prove this, the usual straightening algorithm (see the beginning of the proof of [Theorem 10.1](#) for details) shows that $\text{End}_{\mathcal{H}eis_k(z, t)}(\uparrow^{\otimes n})$ is spanned by diagrams in the image of the canonical homomorphism $AH_n \rightarrow \text{End}_{\mathcal{H}eis_k(z, t)}(\uparrow^{\otimes n})$, with some number of bubbles added to the right-hand edge. Thus we have an induced homomorphism $H_n^f \rightarrow \text{End}_{\mathcal{H}(f|g)}(\uparrow^{\otimes n})$ which is surjective since bubbles on the right-hand edge are scalars in the generalized cyclotomic quotient. On the other hand, $\text{End}_{\mathcal{V}(f|g)}(\uparrow^{\otimes n} \otimes S) = \text{End}_{H_n^f}(H_n^f) = H_n^f$. The claim follows. Hence, the right-hand vertical map is an isomorphism when $X^* \otimes Y = \downarrow^{\otimes n} \otimes \uparrow^{\otimes n}$. Using this, we can show that the right hand vertical map is an isomorphism in general. All of the morphism spaces are zero unless $X^* \otimes Y$ has the same number of \uparrow 's as \downarrow 's. If all \downarrow 's are to the left of all \uparrow 's, we are done already, so we may assume that $X^* \otimes Y$ involves $\uparrow \otimes \downarrow$ as a subword. Let U be $X^* \otimes Y$ with the two letters in this subword interchanged and V be $X^* \otimes Y$ with these two letters deleted. Using the isomorphism $\uparrow \otimes \downarrow \cong \downarrow \otimes \uparrow \oplus \mathbb{1}^{\oplus k}$ from [\(2.10\)](#), we get a commuting diagram:

$$\begin{array}{ccc} \text{Hom}_{\mathcal{H}(f|g)}(\mathbb{1}, X^* \otimes Y) & \xrightarrow{\sim} & \text{Hom}_{\mathcal{H}(f|g)}(\mathbb{1}, U \oplus V^{\oplus k}) \\ \psi_{f|g} \downarrow & & \downarrow \psi_{f|g} \\ \text{Hom}_{\mathcal{V}(f|g)}(S, X^* \otimes Y \otimes S) & \xrightarrow{\sim} & \text{Hom}_{\mathcal{V}(f|g)}(S, U \otimes S \oplus V \otimes S^{\oplus k}) \end{array}$$

By induction, the right-hand vertical map is an isomorphism, hence, so too is the left-hand one.

Finally, we explain why $\psi_{f|g}$ is dense. Let Q be an indecomposable object in $\mathcal{V}(f|g)$. We have that $\downarrow^{\otimes m} \otimes \uparrow^{\otimes n} \otimes S = \downarrow^{\otimes m} \otimes (H_n^f, H_m^g) = (H_n^f, H_m^g) \oplus M$ where M is a direct sum of summands of $(H_{n'}^f, H_{m'}^g)$ with $n' < n$ and $m' < m$. It follows that Q is isomorphic to the image of some idempotent in $\text{End}_{\mathcal{V}(f|g)}(\downarrow^{\otimes m} \otimes \uparrow^{\otimes n} \otimes S)$ for some $m, n \geq 0$. Since we have shown already that $\psi_{f|g}$ is full and faithful, there is a corresponding idempotent in $\text{End}_{\mathcal{H}(f|g)}(\downarrow^{\otimes m} \otimes \uparrow^{\otimes n})$. The latter idempotent defines an object P of $\text{Kar}(\mathcal{H}(f|g))$ such that $\psi_{f|g}(P) \cong Q$. \square

Remark 9.6. If $g(w) = 1$ the genericity assumption is vacuous, so [Theorem 9.5](#) gives us an equivalence of categories $\psi_{f|1} : \text{Kar}(\mathcal{H}(f|1)) \rightarrow \mathcal{V}(f)$. In other words, the generalized cyclotomic quotient $\mathcal{H}(f|1)$

is Morita equivalent to the “usual” cyclotomic quotient defined by the cyclotomic Hecke algebras H_n^f for all $n \geq 0$. This statement is the quantum analog of [Brundan 2018, Theorem 1.7]; see also [Rouquier 2012, Theorem 4.25] for the analogous result in the setting of Kac–Moody 2-categories.

Remark 9.7. More generally, suppose that there are factorizations $f(w) = f_1(w)f_2(w)$ and $g(w) = g_1(w)g_2(w)$ such that the genericity assumption $\lambda\mu^{-1} \notin \{q^{2i} \mid i \in \mathbb{Z}\}$ holds for λ a root of $f_1(w)$ or $g_1(w)$, and μ a root of $f_2(w)$ or $g_2(w)$. Then a similar argument to the proof of [Theorem 9.5](#) can be used to show that the categories $\text{Kar}(\mathcal{H}(f|g))$ and $\text{Kar}(\mathcal{H}(f_1|g_1) \boxtimes \mathcal{H}(f_2|g_2))$ are equivalent. In particular, applying this to $\text{Kar}(\mathcal{H}(f|1))$ and using the previous remark, it follows that the cyclotomic Hecke algebra H_n^f is Morita equivalent to $\bigoplus_{n_1+n_2=n} H_{n_1}^{f_1} \otimes H_{n_2}^{f_2}$, thereby recovering a result of Dipper and Mathas [2002].

10. Basis theorem

Finally, we prove a basis theorem for the morphism spaces in $\mathcal{H}eis_k(z, t)$. Our proof of this is very similar to the argument in the degenerate case from [Brundan et al. 2018, Theorem 6.4]. Let $X = X_r \otimes \cdots \otimes X_1$ and $Y = Y_s \otimes \cdots \otimes Y_1$ be objects of $\mathcal{H}eis_k(z, t)$ for $X_i, Y_j \in \{\uparrow, \downarrow\}$. An (X, Y) -*matching* is a bijection between $\{i \mid X_i = \uparrow\} \sqcup \{j \mid Y_j = \downarrow\}$ and $\{i \mid X_i = \downarrow\} \sqcup \{j \mid Y_j = \uparrow\}$. A *reduced lift* of an (X, Y) -matching means a diagram representing a morphism $X \rightarrow Y$ such that

- the endpoints of each string are points which correspond under the given matching;
- there are no floating bubbles and no dots on any string;
- there are no self-intersections of strings and no two strings cross each other more than once.

Fix a set $B(X, Y)$ consisting of a choice of reduced lift for each of the (X, Y) -matchings. Let $B_o(X, Y)$ be the set of all morphisms that can be obtained from the elements of $B(X, Y)$ by adding dots labeled with integer multiplicities near to the terminus of each string. Also recall the homomorphism $\beta : \text{Sym} \otimes \text{Sym} \rightarrow \text{End}_{\mathcal{H}eis_k(z, t)}(\mathbb{1})$ from (3.7). Using it, we can make the morphism space $\text{Hom}_{\mathcal{H}eis_k(z, t)}(X, Y)$ into a right $\text{Sym} \otimes \text{Sym}$ -module: $\phi\theta := \phi \otimes \beta(\theta)$.

Theorem 10.1. *For any ground ring \mathbb{k} , parameters $z, t \in \mathbb{k}^\times$, and objects $X, Y \in \mathcal{H}eis_k(z, t)$, the morphism space $\text{Hom}_{\mathcal{H}eis_k(z, t)}(X, Y)$ is a free right $\text{Sym} \otimes \text{Sym}$ -module with basis $B_o(X, Y)$.*

Proof. We just prove this when $k \leq 0$; the result for $k \geq 0$ then follows by applying Ω_k . Let $X = X_r \otimes \cdots \otimes X_1$ and $Y = Y_s \otimes \cdots \otimes Y_1$ be two objects.

We first observe that $B_o(X, Y)$ spans $\text{Hom}_{\mathcal{H}eis_k(z, t)}(X, Y)$ as a right $\text{Sym} \otimes \text{Sym}$ -module. The defining relations and the additional relations derived in [Sections 2, 3](#) and [4](#) give Reidemeister-type relations modulo terms with fewer crossings, plus a skein relation and bubble and dot sliding relations. These relations allow diagrams for morphisms in $\mathcal{H}eis_k(z, t)$ to be transformed in a similar way to the way oriented tangles are simplified in skein categories, modulo diagrams with fewer crossings. Hence, there is a straightening algorithm to rewrite any diagram representing a morphism $X \rightarrow Y$ as a linear combination of the ones in $B_o(X, Y)$.

It remains to prove the linear independence. We say $\phi \in B_o(X, Y)$ is *positive* if it only involves nonnegative powers of dots. It suffices to show just that the positive morphisms in $B_o(X, Y)$ are linearly independent. Indeed, given any linear relation of the form $\sum_{i=1}^N \phi_i \otimes \beta(\theta_i) = 0$ for morphisms $\phi_i \in B_o(X, Y)$ and coefficients $\theta_i \in \text{Sym} \otimes \text{Sym}$, we can “clear denominators” by multiplying the termini of the strings by sufficiently large positive powers of dots to reduce to the positive case.

The main step now is to prove the linear independence in the special case that $X = Y = \uparrow^{\otimes n}$. To do this, we need to allow the ground ring \mathbb{k} to change, so we will add a subscript to our notation, denoting $\mathcal{H}eis_k(z, t)$, $\mathcal{V}(f|g)$, $\text{Sym} \otimes \text{Sym}$, … by ${}_{\mathbb{k}}\mathcal{H}eis_k(z, t)$, ${}_{\mathbb{k}}\mathcal{V}(f|g)$, ${}_{\mathbb{k}}\text{Sym} \otimes {}_{\mathbb{k}}\text{Sym}$, … to avoid any confusion. It suffices to prove the linear independence of positive elements of $B_o(X, Y)$ in the special case that $\mathbb{k} = \mathbb{Z}[z^{\pm 1}, t^{\pm 1}]$; one can then use the canonical \mathbb{k} -linear monoidal functor ${}_{\mathbb{k}}\mathcal{H}eis_k(z, t) \rightarrow {}_{\mathbb{k}} \otimes_{\mathbb{Z}[z^{\pm 1}, t^{\pm 1}]} \mathbb{Z}[z^{\pm 1}, t^{\pm 1}] {}_{\mathbb{k}}\mathcal{H}eis_k(z, t)$ to deduce the linear independence over an arbitrary ground ring \mathbb{k} and for arbitrary parameters.

So assume now that $\mathbb{k} = \mathbb{Z}[z^{\pm 1}, t^{\pm 1}]$ and take a linear relation $\sum_{i=1}^N \phi_i \otimes \beta(\theta_i) = 0$ for positive $\phi_i \in B_o(X, Y)$. Choose a so that the multiplicities of dots in all ϕ_i arising in this linear relation are $\leq a$. Also choose $b, c \geq 0$ so that all of the symmetric functions $\theta_i \in {}_{\mathbb{k}}\text{Sym} \otimes {}_{\mathbb{k}}\text{Sym}$ are polynomials in the elementary symmetric functions $e_1 \otimes 1, \dots, e_b \otimes 1$ and $1 \otimes e_1, \dots, 1 \otimes e_c$. Then choose l, m so that $a < l$, $b + c < m$ and $k = m - l$. Note that $l \geq m$ due to our standing assumption that $k \leq 0$. Let u_1, \dots, u_b and v_1, \dots, v_c be indeterminates and \mathbb{K} be the algebraic closure of the field $\mathbb{Q}(z, t, u_1, \dots, u_b, v_1, \dots, v_c)$. Pick $q \in \mathbb{K}^\times$ so that $z = q - q^{-1}$ and consider the cyclotomic Hecke algebras ${}_{\mathbb{K}}H_n^f$ and ${}_{\mathbb{K}}H_n^g$ over \mathbb{K} associated to the polynomials

$$f(w) := w^l + t^2, \quad g(w) = w^m + u_1 w^{m-1} + \dots + u_b w^{m-b} + v_c w^c + \dots + v_1 w + 1.$$

Note the formula for $g(w)$ makes sense because $b + c < m$. Consider the ${}_{\mathbb{k}}\mathcal{H}eis_k(z, t)$ -module category ${}_{\mathbb{k}}\mathcal{V}(f|g)$ from (9.11) (taking $u := t$ and $v := 1$). Since $\mathbb{k} \hookrightarrow \mathbb{K}$, there is a canonical \mathbb{k} -linear monoidal functor ${}_{\mathbb{k}}\mathcal{H}eis_k(z, t) \rightarrow {}_{\mathbb{k}}\mathcal{H}eis_k(z, t)$, allowing us to view ${}_{\mathbb{k}}\mathcal{V}(f|g)$ also as a module category over ${}_{\mathbb{k}}\mathcal{H}eis_k(z, t)$. Then we evaluate the relation $\sum \phi_i \otimes \beta(\theta_i) = 0$ on ${}_{\mathbb{k}}S := ({}_{\mathbb{k}}H_0^f, {}_{\mathbb{k}}H_0^g)$ to obtain a relation in ${}_{\mathbb{k}}H_n^f$. By the basis theorem for ${}_{\mathbb{k}}H_n^f$ from (6.2) and the assumption that $a < l$, the images of ϕ_1, \dots, ϕ_N in ${}_{\mathbb{k}}H_n^f$ are linearly independent over \mathbb{K} , so we deduce that the image of $\beta(\theta_i)$ in \mathbb{K} is zero for each i . To deduce from this that $\theta_i = 0$, recall that θ_i is a polynomial in $e_1 \otimes 1, \dots, e_b \otimes 1, 1 \otimes e_1, \dots, 1 \otimes e_c$. So we need to show that the images of $\beta(e_1 \otimes 1), \dots, \beta(e_b \otimes 1), \beta(1 \otimes e_1), \dots, \beta(1 \otimes e_c)$ in \mathbb{K} are algebraically independent. In fact, we claim that these images are the indeterminates $u_1, \dots, u_b, v_1, \dots, v_c$, respectively. To prove this, note that the low degree terms of $\mathbb{O}^\pm(w)$ are

$$\begin{aligned} \mathbb{O}^+(w) &= g(w)/f(w) = w^k + u_1 w^{k-1} + \dots + u_b w^{k-b} + \dots \in w^k \mathbb{K}[[w^{-1}]], \\ \mathbb{O}^-(w) &= t^2 g(w)/f(w) = 1 + v_1 w + \dots + v_c w^c + \dots \in \mathbb{K}[[w]]. \end{aligned}$$

By (3.9), (9.3)–(9.5) and Lemma 9.2, the images of $\beta(e_n \otimes 1)$ and $\beta(1 \otimes e_n)$ are the w^{k-n} - and w^n -coefficients of $\mathbb{O}^+(u)$ and $\mathbb{O}^-(u)$, respectively, and the claim follows.

We have now proved the linear independence when $X = Y = \uparrow^{\otimes n}$. Returning to the general case, we can use the canonical isomorphism $\text{Hom}_{\mathcal{H}\text{eisen}(z,t)}(X, Y) \cong \text{Hom}_{\mathcal{H}\text{eisen}(z,t)}(\mathbb{1}, X^* \otimes Y)$ arising from the rigidity to see that the $\text{Sym} \otimes \text{Sym}$ -linear independence of the positive morphisms in $B_o(X, Y)$ is equivalent to the $\text{Sym} \otimes \text{Sym}$ -linear independence of the positive morphisms in $B_o(\mathbb{1}, X^* \otimes Y)$. Thus, we are reduced to the case that $X = \mathbb{1}$. Assume this from now on. The set $B_o(\mathbb{1}, Y)$ is empty unless Y has the same number n of \uparrow 's as \downarrow 's. Also we have already proved the linear independence in the case $Y = \downarrow^{\otimes n} \otimes \uparrow^{\otimes n}$. So we may assume that Y has a subword $\uparrow \otimes \downarrow$. Let Z be Y with the two letters in the subword interchanged. By induction, we may assume the linear independence has already been established for $B_o(\mathbb{1}, Z)$. Consider a linear relation $\sum_{i=1}^N \phi_i \otimes \beta(\theta_i)$ for positive $\phi_i \in B_o(\mathbb{1}, Y)$. Recalling the isomorphism $\uparrow \otimes \downarrow \oplus \mathbb{1}^{\oplus(-k)} \xrightarrow{\sim} \downarrow \otimes \uparrow$ from (2.25), multiplying the subword $\uparrow \otimes \downarrow$ on top by the sideways crossing  defines a $\text{Sym} \otimes \text{Sym}$ -linear map $s : \text{Hom}_{\mathcal{H}\text{eisen}(z,t)}(\mathbb{1}, Y) \hookrightarrow \text{Hom}_{\mathcal{H}\text{eisen}(z,t)}(\mathbb{1}, Z)$. Unfortunately, s does not send $B_o(\mathbb{1}, Y)$ into $B_o(\mathbb{1}, Z)$. However, the image of $B_o(\mathbb{1}, Y)$ is related to $B_o(\mathbb{1}, Z)$ in a triangular way, which is good enough to complete the argument. The full explanation of this is almost exactly the same as in the degenerate case, so we refer the reader to the last paragraph of the proof of [Brundan et al. 2018, Theorem 6.4] for the details. \square

Corollary 10.2.

$$\text{End}_{\mathcal{H}\text{eisen}(z,t)}(\mathbb{1}) \cong \text{Sym} \otimes \text{Sym}.$$

References

- [Ariki and Koike 1994] S. Ariki and K. Koike, “A Hecke algebra of $(\mathbb{Z}/r\mathbb{Z}) \wr \mathfrak{S}_n$ and construction of its irreducible representations”, *Adv. Math.* **106**:2 (1994), 216–243. [MR](#) [Zbl](#)
- [Bezrukavnikov and Etingof 2009] R. Bezrukavnikov and P. Etingof, “Parabolic induction and restriction functors for rational Cherednik algebras”, *Selecta Math. (N.S.)* **14**:3-4 (2009), 397–425. [MR](#) [Zbl](#)
- [Brundan 2017] J. Brundan, “Representations of the oriented skein category”, preprint, 2017. [arXiv](#)
- [Brundan 2018] J. Brundan, “On the definition of Heisenberg category”, *Algebr. Comb.* **1**:4 (2018), 523–544. [MR](#) [Zbl](#)
- [Brundan and Davidson 2017] J. Brundan and N. Davidson, “Categorical actions and crystals”, pp. 105–147 in *Categorification and higher representation theory*, edited by A. Beliakova and A. D. Lauda, *Contemp. Math.* **683**, Amer. Math. Soc., Providence, RI, 2017. [MR](#) [Zbl](#)
- [Brundan et al. 2014] J. Brundan, A. Kleshchev, and P. J. McNamara, “Homological properties of finite-type Khovanov–Lauda–Rouquier algebras”, *Duke Math. J.* **163**:7 (2014), 1353–1404. [MR](#) [Zbl](#)
- [Brundan et al. 2017] J. Brundan, J. Comes, D. Nash, and A. Reynolds, “A basis theorem for the affine oriented Brauer category and its cyclotomic quotients”, *Quantum Topol.* **8**:1 (2017), 75–112. [MR](#) [Zbl](#)
- [Brundan et al. 2018] J. Brundan, A. Savage, and B. Webster, “The degenerate Heisenberg category and its Grothendieck ring”, preprint, 2018. [arXiv](#)
- [Brundan et al. 2019] J. Brundan, A. Savage, and B. Webster, “Heisenberg and Kac–Moody categorification”, preprint, 2019. [arXiv](#)
- [Brundan et al. ≥ 2020] J. Brundan, A. Savage, and B. Webster, “Quantum Frobenius Heisenberg categorification”, in preparation.
- [Cautis et al. 2018] S. Cautis, A. D. Lauda, A. M. Licata, P. Samuelson, and J. Sussan, “The elliptic Hall algebra and the deformed Khovanov Heisenberg category”, *Selecta Math. (N.S.)* **24**:5 (2018), 4041–4103. [MR](#) [Zbl](#)
- [Dipper and Mathas 2002] R. Dipper and A. Mathas, “Morita equivalences of Ariki–Koike algebras”, *Math. Z.* **240**:3 (2002), 579–610. [MR](#) [Zbl](#)
- [Ginzburg et al. 2003] V. Ginzburg, N. Guay, E. Opdam, and R. Rouquier, “On the category \mathcal{O} for rational Cherednik algebras”, *Invent. Math.* **154**:3 (2003), 617–651. [MR](#) [Zbl](#)

[Gould et al. 1991] M. D. Gould, R. B. Zhang, and A. J. Bracken, “Generalized Gelfand invariants and characteristic identities for quantum groups”, *J. Math. Phys.* **32**:9 (1991), 2298–2303. [MR](#) [Zbl](#)

[Jantzen 1996] J. C. Jantzen, *Lectures on quantum groups*, Grad. Stud. Math. **6**, Amer. Math. Soc., Providence, RI, 1996. [MR](#) [Zbl](#)

[Khovanov 2014] M. Khovanov, “Heisenberg algebra and a graphical calculus”, *Fund. Math.* **225**:1 (2014), 169–210. [MR](#) [Zbl](#)

[Li 2010] J. Li, “The quantum Casimir operators of $U_q(\mathfrak{gl}_n)$ and their eigenvalues”, *J. Phys. A* **43**:34 (2010), art. id. 345202. [MR](#) [Zbl](#)

[Licata and Savage 2013] A. Licata and A. Savage, “Hecke algebras, finite general linear groups, and Heisenberg categorification”, *Quantum Topol.* **4**:2 (2013), 125–185. [MR](#) [Zbl](#)

[Lusztig 1993] G. Lusztig, *Introduction to quantum groups*, Progr. Math. **110**, Birkhäuser, Boston, 1993. [MR](#) [Zbl](#)

[Macdonald 1995] I. G. Macdonald, *Symmetric functions and Hall polynomials*, 2nd ed., Oxford Math. Monogr., Oxford Univ. Press, 1995. [MR](#) [Zbl](#)

[Mackaay and Savage 2018] M. Mackaay and A. Savage, “Degenerate cyclotomic Hecke algebras and higher level Heisenberg categorification”, *J. Algebra* **505** (2018), 150–193. [MR](#) [Zbl](#)

[Rouquier 2012] R. Rouquier, “Quiver Hecke algebras and 2-Lie algebras”, *Algebra Colloq.* **19**:2 (2012), 359–410. [MR](#) [Zbl](#)

[Savage 2019] A. Savage, “Frobenius Heisenberg categorification”, *Algebr. Comb.* **2**:5 (2019), 937–967. [Zbl](#)

[Shan 2011] P. Shan, “Crystals of Fock spaces and cyclotomic rational double affine Hecke algebras”, *Ann. Sci. Éc. Norm. Sup. (4)* **44**:1 (2011), 147–182. [MR](#) [Zbl](#)

[Turaev 1989] V. G. Turaev, “Operator invariants of tangles, and R -matrices”, *Izv. Akad. Nauk SSSR Ser. Mat.* **53**:5 (1989), 1073–1107. In Russian; translated in *Mat. USSR-Uzv.* **35**:2 (1990), 411–444. [Zbl](#)

[Virk 2011] R. Virk, “Derived equivalences and \mathfrak{sl}_2 -categorifications for $U_q(\mathfrak{gl}_n)$ ”, *J. Algebra* **346** (2011), 82–100. [MR](#) [Zbl](#)

[Webster 2015] B. Webster, “Canonical bases and higher representation theory”, *Compos. Math.* **151**:1 (2015), 121–166. [MR](#) [Zbl](#)

[Webster 2018] B. Webster, “Unfurling Khovanov–Lauda–Rouquier algebras”, preprint, 2018. [arXiv](#)

Communicated by J. Toby Stafford

Received 2019-01-27

Revised 2019-07-22

Accepted 2019-09-02

brundan@uoregon.edu

Department of Mathematics, University of Oregon, Eugene, OR, United States

alistair.savage@uottawa.ca

Department of Mathematics and Statistics, University of Ottawa, ON, Canada

ben.webster@uwaterloo.ca

Department of Pure Mathematics, University of Waterloo, Waterloo, ON, Canada

Perimeter Institute for Theoretical Physics, Waterloo, ON, Canada

Algebra & Number Theory

msp.org/ant

EDITORS

MANAGING EDITOR

Bjorn Poonen
Massachusetts Institute of Technology
Cambridge, USA

EDITORIAL BOARD CHAIR

David Eisenbud
University of California
Berkeley, USA

BOARD OF EDITORS

Bhargav Bhatt	University of Michigan, USA	Martin Olsson	University of California, Berkeley, USA
Richard E. Borcherds	University of California, Berkeley, USA	Raman Parimala	Emory University, USA
Antoine Chambert-Loir	Université Paris-Diderot, France	Jonathan Pila	University of Oxford, UK
J.-L. Colliot-Thélène	CNRS, Université Paris-Sud, France	Irena Peeva	Cornell University, USA
Brian D. Conrad	Stanford University, USA	Anand Pillay	University of Notre Dame, USA
Samit Dasgupta	Duke University, USA	Michael Rapoport	Universität Bonn, Germany
Hélène Esnault	Freie Universität Berlin, Germany	Victor Reiner	University of Minnesota, USA
Gavril Farkas	Humboldt Universität zu Berlin, Germany	Peter Sarnak	Princeton University, USA
Hubert Flenner	Ruhr-Universität, Germany	Joseph H. Silverman	Brown University, USA
Sergey Fomin	University of Michigan, USA	Michael Singer	North Carolina State University, USA
Edward Frenkel	University of California, Berkeley, USA	Christopher Skinner	Princeton University, USA
Wee Teck Gan	National University of Singapore	Vasudevan Srinivas	Tata Inst. of Fund. Research, India
Andrew Granville	Université de Montréal, Canada	J. Toby Stafford	University of Michigan, USA
Ben J. Green	University of Oxford, UK	Shunsuke Takagi	University of Tokyo, Japan
Joseph Gubeladze	San Francisco State University, USA	Pham Huu Tiep	University of Arizona, USA
Christopher Hacon	University of Utah, USA	Ravi Vakil	Stanford University, USA
Roger Heath-Brown	Oxford University, UK	Michel van den Berg	Hasselt University, Belgium
János Kollár	Princeton University, USA	Akshay Venkatesh	Institute for Advanced Study, USA
Philippe Michel	École Polytechnique Fédérale de Lausanne	Marie-France Vignéras	Université Paris VII, France
Susan Montgomery	University of Southern California, USA	Melanie Matchett Wood	University of California, Berkeley, USA
Shigefumi Mori	RIMS, Kyoto University, Japan	Shou-Wu Zhang	Princeton University, USA

PRODUCTION

production@msp.org

Silvio Levy, Scientific Editor

See inside back cover or msp.org/ant for submission instructions.

The subscription price for 2020 is US \$415/year for the electronic version, and \$620/year (+\$60, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to MSP.

Algebra & Number Theory (ISSN 1944-7833 electronic, 1937-0652 printed) at Mathematical Sciences Publishers, 798 Evans Hall #3840, c/o University of California, Berkeley, CA 94720-3840 is published continuously online. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices.

ANT peer review and production are managed by EditFLOW® from MSP.

PUBLISHED BY

 **mathematical sciences publishers**

nonprofit scientific publishing

<http://msp.org/>

© 2020 Mathematical Sciences Publishers

Algebra & Number Theory

Volume 14 No. 2 2020

On the definition of quantum Heisenberg category	275
JONATHAN BRUNDAN, ALISTAIR SAVAGE and BEN WEBSTER	
Characteristic cycles and Gevrey series solutions of A -hypergeometric systems	323
CHRISTINE BERKESCH and MARÍA-CRUZ FERNÁNDEZ-FERNÁNDEZ	
Singularity categories of deformations of Kleinian singularities	349
SIMON CRAWFORD	
Iwasawa main conjecture for Rankin–Selberg p -adic L -functions	383
XIN WAN	
Positivity results for spaces of rational curves	485
ROYA BEHESHTI and ERIC RIEDL	
Generalized Schur algebras	501
ALEXANDER KLESHCHEV and ROBERT MUTH	