

Algebra & Number Theory

Volume 14
2020
No. 2

**Singularity categories of deformations
of Kleinian singularities**

Simon Crawford



Singularity categories of deformations of Kleinian singularities

Simon Crawford

Let G be a finite subgroup of $\mathrm{SL}(2, \mathbb{k})$ and let $R = \mathbb{k}[x, y]^G$ be the coordinate ring of the corresponding Kleinian singularity. In 1998, Crawley-Boevey and Holland defined deformations \mathcal{O}^λ of R parametrised by weights λ . In this paper, we determine the singularity categories $\mathcal{D}_{\mathrm{sg}}(\mathcal{O}^\lambda)$ of these deformations, and show that they correspond to subgraphs of the Dynkin graph associated to R . This generalises known results on the structure of $\mathcal{D}_{\mathrm{sg}}(R)$. We also provide a generalisation of the intersection theory appearing in the geometric McKay correspondence to a noncommutative setting.

1. Introduction

1A. Background. Throughout let \mathbb{k} be an algebraically closed field of characteristic 0. The Kleinian singularities \mathbb{k}^2/G , where G is a finite subgroup of $\mathrm{SL}(2, \mathbb{k})$, are ubiquitous in algebraic geometry, representation theory, and singularity theory. In this paper, we shall study the latter of these for a family of (generically noncommutative) algebras.

The notion of the singularity category of a ring R was introduced by Buchweitz [1986] as a particular Verdier quotient of $\mathcal{D}^b(\mathrm{mod}\text{-}R)$. More specifically, writing $\mathrm{Perf}(R)$ for the full subcategory of perfect complexes in $\mathcal{D}^b(\mathrm{mod}\text{-}R)$, Buchweitz defined the singularity category as the Verdier quotient category

$$\mathcal{D}_{\mathrm{sg}}(R) := \frac{\mathcal{D}^b(\mathrm{mod}\text{-}R)}{\mathrm{Perf}(R)}.$$

By construction, this category possesses the structure of a triangulated category. Buchweitz also showed that, when R is Gorenstein, the singularity category is triangle equivalent to $\underline{\mathrm{MCM}}\text{-}R$, the stable category of maximal Cohen–Macaulay R -modules (that this latter category is triangulated also follows from a general result of Happel [1988]). The singularity category of a commutative ring R is also closely related to the category of reduced matrix factorisations of R by [Eisenbud 1980, Corollary 6.3], and under mild hypotheses these categories are in fact equivalent.

From the above definition, it is not difficult to see that $\mathcal{D}_{\mathrm{sg}}(R)$ is trivial precisely when R has finite global dimension. However, in general it is difficult to give an adequate description of the singularity category of an arbitrary Gorenstein ring of infinite global dimension. Recent work includes [Chen 2011; 2018] which describes the singularity category when R has radical square zero or when it is a quadratic

MSC2010: primary 14J17; secondary 16G20, 16G50, 18E30.

Keywords: singularity categories, Kleinian singularities, preprojective algebras.

monomial algebra, and [Kalek 2015] which provides a description when R is a so-called gentle algebra. Moreover, in [Amiot et al. 2015] the authors determine the singularity categories of some commutative Gorenstein isolated singularities.

The standard examples of commutative surface singularities are the Kleinian singularities, which are very well understood. The main aim of this paper is to provide a concrete description of the singularity category of certain noncommutative deformations of the coordinate ring of a Kleinian singularity.

Very little is known about the singularities of noncommutative rings, particularly those which are not finite over their centre. For example, given a singular noncommutative ring S , it is not known whether and under what circumstances one can find commutative rings R_1, \dots, R_k such that we have an equivalence of triangulated categories $\mathcal{D}_{\text{sg}}(S) \simeq \bigoplus_{i=1}^k \mathcal{D}_{\text{sg}}(R_i)$. If this is the case, one can think of S as having the same singularities as those of the varieties $\text{Spec } R_i$. Our main result, Theorem 1.1, shows that we have such a decomposition of singularity categories for the algebras of interest in this paper, and can be seen as a first step towards better understanding singularities of noncommutative surfaces.

Crawley-Boevey and Holland [1998] defined a family of algebras $\mathcal{O}^\lambda(\tilde{Q})$ depending on the data of an extended Dynkin quiver \tilde{Q} and a so-called weight for \tilde{Q} . Write Q for the Dynkin quiver obtained from \tilde{Q} by removing an extending vertex, and write R_Q for the coordinate ring of the corresponding Kleinian singularity. Then the algebras $\mathcal{O}^\lambda(\tilde{Q})$ may be thought of as deformations of R_Q in the sense that there exists a filtration \mathcal{F} of $\mathcal{O}^\lambda(\tilde{Q})$ satisfying $\text{gr}_{\mathcal{F}} \mathcal{O}^\lambda(\tilde{Q}) \simeq R_Q$. These deformations are generically noncommutative, a property which depends on the weight λ , and it is easy to determine when this is the case. When $\tilde{Q} = \tilde{A}_n$, if $\mathcal{O}^\lambda(\tilde{Q})$ is noncommutative then it is an example of a generalised Weyl algebra, as studied in [Bavula 1992; Hodges 1993]. If $\mathcal{O}^\lambda(\tilde{Q})$ is commutative, a description of its singularity category follows from [Iyama and Wemyss 2014, Theorem 3.2], where quite geometric techniques are employed. Through a completely ring-theoretic approach, we determine $\mathcal{D}_{\text{sg}}(\mathcal{O}^\lambda(\tilde{Q}))$ irrespective of whether the deformation is commutative or noncommutative.

Our main result can be stated as follows, where undefined terms will be defined in Section 2.

Theorem 1.1 (Theorem 3.6, Theorem 4.13). *Let \tilde{Q} be an extended Dynkin quiver with vertex set $\{0, 1, \dots, n\}$, where 0 is an extending vertex, and write Q for the full subquiver obtained by deleting vertex 0. Let λ be a weight for \tilde{Q} . Then there exists a subset $J = J(\lambda)$ of $\{1, \dots, n\}$ such that, if $Q^{(1)} \sqcup \dots \sqcup Q^{(r)}$ is the full subquiver of Q obtained by deleting the vertices in J , so that the $Q^{(i)}$ are connected and therefore necessarily Dynkin, there is a triangle equivalence*

$$\mathcal{D}_{\text{sg}}(\mathcal{O}^\lambda(\tilde{Q})) \simeq \bigoplus_{i=1}^r \mathcal{D}_{\text{sg}}(R_{Q^{(i)}}).$$

One can show that we may restrict our attention to the case where the weight λ is *quasidominant* (see Definition 2.16). For example, when $\mathbb{k} = \mathbb{C}$, a weight is quasidominant if, for $1 \leq i \leq n$,

$$\lambda_i \in \{z \in \mathbb{C} \mid \text{Re } z > 0, \text{ or } \text{Re } z = 0 \text{ and } \text{Im } z \geq 0\},$$

and where λ_0 can be arbitrary. In this case, the subset J in the above theorem is $J = \{i \in \{1, \dots, n\} \mid \lambda_i = 0\}$.

The result in Theorem 1.1 coincides with the intuition coming from commutative singularity theory which says that deforming a singularity should make it no worse; in our context, deforming a singularity corresponds to making weights at certain vertices of \tilde{Q} nonzero, and the above theorem says that this makes the singularity category simpler, in a precise sense. In the Appendix, we also provide a simple proof which illustrates how the translation functor acts on the triangulated category $\mathcal{D}_{\text{sg}}(\mathcal{O}^\lambda(\tilde{Q}))$.

Since the first version of this paper appeared online, an alternative proof of Theorem 1.1 has been given in [Kalck and Yang 2018, Theorem 9.4] using relative singularity categories.

Now suppose that the weight $\lambda \in \mathbb{k}^{n+1}$ is given by $\lambda_0 = 1$ and $\lambda_i = 0$ for $1 \leq i \leq n$, and in this case write $\lambda = \varepsilon_0$. We then consider $\mathcal{O}^\lambda(\tilde{Q})$ to be a noncommutative analogue of R_Q . This viewpoint is partially justified by the following immediate corollary:

Corollary 1.2. *Retain the notation of Theorem 1.1, and suppose that $\lambda = \varepsilon_0$, as above. Then there is a triangle equivalence*

$$\mathcal{D}_{\text{sg}}(\mathcal{O}^\lambda(\tilde{Q})) \simeq \mathcal{D}_{\text{sg}}(R_Q).$$

Another family of results concerning Kleinian singularities is what is often called the geometric McKay correspondence, which concerns the intersection theory of the minimal resolution of $\text{Spec } R_Q$. In the last section, we prove a result which may be seen as a generalisation of this to a noncommutative setting. We give an imprecise statement of this result below, and a more precise statement in Section 5.

Theorem 1.3 (Theorem 5.8). *Let \tilde{Q} be an extended Dynkin quiver with corresponding Dynkin quiver Q and let $\lambda = \varepsilon_0$. Then $\mathcal{O}^\lambda(\tilde{Q})$ has a noncommutative resolution, and the intersection theory of the exceptional objects in this resolution is the same as that of the exceptional curves in the minimal resolution of a Kleinian singularity of type corresponding to Q .*

In particular, this result further supports the viewpoint that $\mathcal{O}^\lambda(\tilde{Q})$ can be viewed as a noncommutative analogue of R_Q when $\lambda = \varepsilon_0$.

We now take a moment to provide an overview of the proof of Theorem 1.1. For all of our calculations, we work in $\text{MCM-}\mathcal{O}^\lambda(\tilde{Q})$ rather than $\mathcal{D}_{\text{sg}}(\mathcal{O}^\lambda(\tilde{Q}))$; as mentioned previously, these two categories are triangle equivalent, see Theorem 2.7. The first important observation to make is that we can restrict our attention to weights λ which are *quasidominant*. In Section 3, this restriction allows us to give a concrete description of $\mathcal{D}_{\text{sg}}(\mathcal{O}^\lambda(\tilde{Q}))$ as a \mathbb{k} -linear category in terms of an auxiliary Krull–Schmidt category. We also find that the isoclasses V_i of indecomposable objects in $\mathcal{D}_{\text{sg}}(\mathcal{O}^\lambda(\tilde{Q}))$ are indexed by those vertices $i \geq 1$ with $\lambda_i = 0$. This auxiliary category allows us to establish the equivalence of Theorem 1.1, but only as a \mathbb{k} -linear equivalence.

In a previous version of this paper, a lengthy case-by-case analysis was then used to show that we had the desired triangle equivalence. A result of Keller which subsequently appeared significantly reduces the amount of work that needs to be done. In fact, it now suffices to show that the direct sum appearing on the right-hand side of the (a priori \mathbb{k} -linear) equivalence of Theorem 1.1 is a decomposition into algebraic triangulated subcategories. This is shown in Section 4.

1B. Organisation of the paper. This paper is organised as follows. In Section 2, we recall some basic definitions and facts, and introduce the notation used throughout the paper. In Section 3, the singularity categories of $O^\lambda(\tilde{Q})$ are determined as \mathbb{k} -linear categories. We then complete the proof of Theorem 1.1 in Section 4. In Section 5, we provide a noncommutative version of the geometric McKay correspondence, and in the Appendix we detail how the translation functor behaves on objects of $\mathcal{D}_{\text{sg}}(R_Q)$.

2. Preliminaries

We now recall some of the definitions and results that we will make use of throughout this paper. In this section, R will denote an arbitrary ring.

2A. Conventions. As stated in the introduction, throughout \mathbb{k} will denote an algebraically closed field of characteristic 0. We write $\text{mod-}R$ (respectively, $R\text{-mod}$) for the category of finitely generated right (respectively, left) R -modules; in this paper, we shall use right modules unless otherwise stated. We also write $\text{proj-}R$ for the full subcategory of $\text{mod-}R$ consisting of finitely generated projective modules. We write $M^* := \text{Hom}_R(M, R)$ for the dual of an R -module M , which is an $(R, \text{End}_R(M))$ -bimodule. We write $\text{p.dim } M$ and $\text{i.dim } M$ for the projective and injective dimensions of $M \in \text{mod-}R$, respectively, and $\text{gl.dim } R$ for the global dimension of R .

2B. Definitions and basic results.

Definition 2.1. A *quiver* Q is a directed multigraph, and we write Q_0 for the set of vertices and Q_1 for the set of arrows. We equip Q with head and tail maps $h, t : Q_1 \rightarrow Q_0$ which take an arrow to the vertices that are its head and tail respectively. A *nontrivial path* in the quiver is a sequence of arrows $p = \alpha_1 \alpha_2 \dots \alpha_\ell$ with $h(\alpha_i) = t(\alpha_{i+1})$ for $1 \leq i \leq \ell - 1$ (that is, we compose arrows from left to right), and such a path is said to have *length* ℓ . Moreover, for each vertex $i \in Q_0$ there is a *trivial path* e_i of length 0, with head and tail vertex both equal to i .

Definition 2.2. Given a field \mathbb{k} and a quiver Q , we define the path algebra $\mathbb{k}Q$ of Q as follows: as a \mathbb{k} -vector space, $\mathbb{k}Q$ has a basis given by paths in the quiver, and we define multiplication by concatenation of paths:

$$p \cdot q = \begin{cases} pq & \text{if } h(p) = t(q), \\ 0 & \text{otherwise.} \end{cases}$$

If R is a commutative ring, then $\text{Spec } R$ is nonsingular if and only if R has finite global dimension. It is therefore sensible to say that a (possibly noncommutative) ring is *nonsingular* if it has finite global dimension, and *singular* otherwise. Before we are able to define the singularity category of a ring, we must make a few more definitions.

Definition 2.3. Given R -modules M and N , write $\underline{\text{Hom}}_R(M, N) = \text{Hom}_R(M, N) / \sim$, where $f \sim f'$ if and only if $f - f'$ factors through a finitely generated projective module. The *stable module category* of R , denoted $\underline{\text{mod-}}R$, is then the category whose objects are the same as those of $\text{mod-}R$, and for modules

M, N , has morphisms $\underline{\text{Hom}}_R(M, N)$. Given a full subcategory $\text{abc-}R$ of $\text{mod-}R$, we write $\underline{\text{abc-}}R$ for the full subcategory of $\underline{\text{mod-}}R$ whose objects are the same as those of $\text{abc-}R$.

Noting that an element of $\sum_{i=1}^k n_i \otimes f_i$ of $N \otimes_R M^*$ gives rise to a homomorphism $M \rightarrow N$ via $m \mapsto \sum_{i=1}^k n_i f_i(m)$, it is not hard to show that a module homomorphism $f : M \rightarrow N$ factors through a projective module if and only if f is the image of some element of $N \otimes_R M^*$. Abusing notation, this allows us to identify $\underline{\text{Hom}}_R(M, N)$ with $\text{Hom}_R(M, N)/(N \otimes_R M^*)$, which will be useful in later calculations. In this paper, we are often in the situation where $R = e\Lambda e$, $M = e_1\Lambda e$, and $N = e_2\Lambda e$, where Λ is some ring and e, e_1 , and e_2 are pairwise orthogonal idempotents, and we are able to make identifications

$$M^* \cong e\Lambda e_1, \quad \text{Hom}_R(M, N) \cong e_2\Lambda e_1, \quad \underline{\text{Hom}}_R(M, N) \cong \frac{e_2\Lambda e_1}{e_2\Lambda e\Lambda e_1}.$$

In the stable module category, we have a weaker notion of an isomorphism than in the usual module category. Indeed, [Auslander and Bridger 1969, Proposition 1.44] shows that two R -modules M, N are isomorphic in $\underline{\text{mod-}}R$ if and only if there exist projective modules P and Q such that $M \oplus P \cong N \oplus Q$ in $\text{mod-}R$.

The *first syzygy* ΩM of $M \in \text{mod-}R$ is defined to be the kernel of any surjection $R^n \twoheadrightarrow M$. The observation in the previous paragraph combined with [Rotman 1979, Proposition 8.5] implies that ΩM is uniquely determined in $\underline{\text{mod-}}R$.

Definition 2.4. A ring R is said to be *Gorenstein* if it is noetherian (i.e., left and right noetherian) and both $\text{i.dim } R_R$ and $\text{i.dim } {}_R R$ are finite. By [Zaks 1969, Lemma A], under these hypotheses the values $\text{i.dim } R_R$ and $\text{i.dim } {}_R R$ coincide, and we call this common value the (*injective*) *dimension* of R .

Definition 2.5. Suppose that R is Gorenstein. A finitely generated R -module M is said to be *maximal Cohen–Macaulay* (MCM) if it satisfies $\text{Ext}_R^i(M, R) = 0$ for all $i \geq 1$. We write $\text{MCM-}R$ for the full subcategory of $\text{mod-}R$ consisting of maximal Cohen–Macaulay R -modules.

For commutative local rings, the above definition coincides with the usual (commutative) definition of maximal Cohen–Macaulay modules in terms of depth [Buchweitz 1986, Section 4.2]. Maximal Cohen–Macaulay modules have the following elementary properties, proofs of which can be found in [loc. cit.]:

Lemma 2.6. (1) *Any finitely generated projective module is MCM.*

(2) *MCM modules are reflexive.*

(3) *Finite direct sums and direct summands of MCM modules are MCM.*

(4) *An MCM module is either projective or has infinite projective dimension.*

With these definitions in hand, we now recall a theorem which identifies a category that is triangle equivalent to the singularity category in the case of a Gorenstein ring R :

Theorem 2.7 [Buchweitz 1986, Theorem 4.4.1]. *Suppose that R is Gorenstein. Then the full subcategory $\underline{\text{MCM-}}R$ of $\underline{\text{mod-}}R$ whose objects are MCM R -modules is a triangulated category, with translation functor Σ given by $\Sigma M = \Omega^{-1}M$. Moreover, there is a triangle equivalence $\mathcal{D}_{\text{sg}}(R) \simeq \underline{\text{MCM-}}R$.*

While the term “singularity category” is more suggestive (which was our main reason for using this terminology in the introduction), since every example that we consider in this paper satisfies the hypotheses of this theorem, we instead focus our attention on determining $\underline{\text{MCM}}\text{-}R$.

Theorem 2.7 is a specific example of a more general result due to Happel, which we now briefly recall. An *exact category* \mathcal{C} is an additive category possessing a class of *conflations* (sometimes called exact sequences) which are triples of objects connected by arrows $X \rightarrow Y \rightarrow Z$, and which satisfy a number of axioms; see [Chen 2012, Section 2] for more details. An object $P \in \mathcal{C}$ is *projective* if the functor $\text{Hom}_{\mathcal{C}}(P, -)$ sends conflations to exact sequences, and we say that \mathcal{C} has *enough projectives* if every object $Z \in \mathcal{C}$ fits into a conflation $X \rightarrow P \rightarrow Z$ with P projective. Dually, one has a notion of an *injective object* and of having *enough injectives*. An exact category \mathcal{C} is said to be *Frobenius* provided that it has enough projectives and enough injectives, and the class of projective objects coincides with the class of injective objects. Given a Frobenius category \mathcal{C} , we may form its stable category $\underline{\mathcal{C}}$ in the same way we formed the stable category $\underline{\text{MCM}}\text{-}R$. Then [Happel 1988] shows that this category is triangulated, and if $X \rightarrow Y \rightarrow Z$ is a conflation in \mathcal{C} then there exists a triangle of the form $X \rightarrow Y \rightarrow Z \rightarrow \Sigma X$ in $\underline{\mathcal{C}}$. If \mathcal{T} is a triangulated category which is triangle equivalent to the stable category of a Frobenius category, then we say that \mathcal{T} is *algebraic*.

If R is a Gorenstein ring, then $\text{MCM}\text{-}R$ is Frobenius and so Happel’s result implies that $\underline{\text{MCM}}\text{-}R$ is triangulated; this triangulated structure is precisely the one given in Theorem 2.7. In $\text{MCM}\text{-}R$, every conflation $X \rightarrow Y \rightarrow Z$ arises from a short exact sequence $0 \rightarrow X \rightarrow Y \rightarrow Z \rightarrow 0$ of $\text{MCM } R$ -modules.

Finally, we recall two useful results that will be helpful when identifying the maximal Cohen–Macaulay modules of a ring. Given an additive category \mathcal{C} and an object $C \in \mathcal{C}$, we write $\text{add}(C)$ for the full subcategory of \mathcal{C} consisting of direct summands of finite direct sums of C . This is the smallest additive subcategory of \mathcal{C} which contains C and is closed under taking direct summands. The following result is due to Auslander, but we provide a proof.

Proposition 2.8 (Auslander). *Suppose that R is Gorenstein and that $M \in \text{MCM}\text{-}R$ is a generator (for example, this occurs if M has R as a direct summand or if R is simple). If $\text{gl.dim End}_R(M) \leq 2$, then $\text{add } M = \text{MCM}\text{-}R$.*

Proof. Write $\Lambda = \text{End}_R(M)$. Since $\text{mod-}R$ has split idempotents (a fact which holds for any ring R), [Krause 2015, Proposition 2.3] implies that the functor $\text{Hom}_R(M, -) : \text{mod-}R \rightarrow \text{mod-}\Lambda$ restricts to an equivalence

$$\text{add } M \xrightarrow{\cong} \text{proj-}\Lambda. \quad (2.9)$$

We also note that since M is a generator, $R^n \in \text{add } M$ for any $n \geq 1$.

That $\text{add } M \subseteq \text{MCM}\text{-}R$ is clear, so suppose that $N \in \text{MCM}\text{-}R$. Since R is noetherian, N^* is finitely presented, so we have an exact sequence of left R -modules of the form

$$R^m \rightarrow R^n \rightarrow N^* \rightarrow 0.$$

Applying $\text{Hom}_R(-, R)$ and noting that N is MCM and therefore reflexive, we obtain an exact sequence

$$0 \rightarrow N \rightarrow R^n \rightarrow R^m.$$

Applying $\text{Hom}_R(M, -)$ then gives an exact sequence

$$0 \rightarrow \text{Hom}_R(M, N) \rightarrow \text{Hom}_R(M, R^n) \xrightarrow{\theta} \text{Hom}_R(M, R^m) \rightarrow \text{coker } \theta \rightarrow 0,$$

where, since M is a generator, $\text{Hom}_R(M, R^n)$ and $\text{Hom}_R(M, R^m)$ are both projective Λ -modules by (2.9). Since $\text{gl.dim } \Lambda \leq 2$ we have $\text{p.dim coker } \theta \leq 2$, and therefore $\text{Hom}_R(M, N)$ is also a projective Λ -module. By (2.9), it follows that $N \in \text{add } M$. \square

Remark 2.10. If R has injective dimension at most 2 then the converse of Proposition 2.8 also holds; see [Simon 2018, Proposition 2.2.11].

When R has injective dimension at most 2, we also have the following:

Lemma 2.11. *Let R be a Gorenstein ring of injective dimension at most 2. Then $M \in \text{mod-}R$ is reflexive if and only if it is maximal Cohen–Macaulay.*

Proof. (\Leftarrow) This is Lemma 2.6 (2), and doesn't require the hypothesis on injective dimension.

(\Rightarrow) Suppose now that M is reflexive. Since R is noetherian, M^* is finitely presented, so we have an exact sequence of the form

$$R^m \rightarrow R^n \rightarrow M^* \rightarrow 0.$$

Applying $\text{Hom}_R(-, R)$ and noting that M is reflexive yields an exact sequence

$$0 \rightarrow M \rightarrow R^n \xrightarrow{\theta} R^m \rightarrow \text{coker } \theta \rightarrow 0.$$

But then, by [Rotman 1979, Corollary 6.55], $\text{Ext}_R^i(M, R) \cong \text{Ext}_R^{i+2}(\text{coker } \theta, R) = 0$ for all $i \geq 1$, where the last equality follows since $\text{i.dim } R \leq 2$. That is, M is maximal Cohen–Macaulay. \square

2C. The deformations of Crawley-Boevey and Holland. In Crawley-Boevey and Holland [1998] introduced the notion of the *deformed preprojective algebra* of a quiver Q , and, if \tilde{Q} is extended Dynkin, a family of \mathbb{k} -algebras $\mathcal{O}^\lambda(\tilde{Q})$ which may be thought of as deformations of the coordinate ring of a Kleinian singularity. We now recall these definitions, noting that our definition of $\mathcal{O}^\lambda(\tilde{Q})$ differs slightly from that of Crawley-Boevey and Holland, but is consistent with their definition by [loc. cit., Theorem 0.1].

Definition 2.12. Let Q be a quiver without loops. The *double* of Q is the quiver \bar{Q} obtained from Q by adding a *reverse arrow* $\bar{\alpha} : j \rightarrow i$ for each arrow $\alpha : i \rightarrow j$ in Q . We call the arrows in \bar{Q} which are not reverse arrows *ordinary arrows*. Given a weight $\lambda \in \mathbb{k}^{Q_0}$ for Q , the corresponding *deformed preprojective algebra* is the \mathbb{k} -algebra

$$\Pi^\lambda(Q) := \mathbb{k}\bar{Q}/I$$

where I is the two-sided ideal of $\mathbb{k}\bar{Q}$ with generators

$$\sum_{\substack{\alpha \in Q_1 \\ t(\alpha)=i}} \alpha \bar{\alpha} - \sum_{\substack{\alpha \in Q_1 \\ h(\alpha)=i}} \bar{\alpha} \alpha - \lambda_i e_i$$

for each vertex $i \in Q_0$. It is easy to see that we can equivalently define I as being the two-sided ideal with the single generator

$$\sum_{\alpha \in Q_1} (\alpha \bar{\alpha} - \bar{\alpha} \alpha) - \sum_{i \in Q_0} \lambda_i e_i.$$

It is helpful to think of a weight as a label from \mathbb{k} at each vertex of Q , and we will often refer to λ_i as the weight at vertex i .

Now suppose that \tilde{Q} is extended Dynkin, with vertices and arrows (of its double) labelled as in Figure 1. Throughout this paper, it will be our convention that \tilde{Q} denotes an extended Dynkin quiver, while Q will denote the corresponding Dynkin quiver obtained by removing the extending vertex 0, where the orientation of the arrows comes from Figure 1.

We are now able to define the algebras of interest to us. We write $\mathcal{O}^\lambda(\tilde{Q})$ for the algebra

$$\mathcal{O}^\lambda(\tilde{Q}) := e_0 \Pi^\lambda(\tilde{Q}) e_0.$$

The elements of $\mathcal{O}^\lambda(\tilde{Q})$ may be thought of as linear combinations of (equivalence classes of) paths in the double of \tilde{Q} which start and end at the extending vertex 0.

If $\lambda = \mathbf{0}$, then $\Pi(Q) := \Pi^\lambda(Q)$ is the (undeformed) preprojective algebra of Gelfand and Ponomarev [1979], and in this case we also write $\mathcal{O}(\tilde{Q}) := \mathcal{O}^\lambda(\tilde{Q})$. We will often write Π^λ and \mathcal{O}^λ (or Π and \mathcal{O} if $\lambda = \mathbf{0}$) when the corresponding quiver is either unimportant or understood.

For our purposes, it is important to know precisely when the rings \mathcal{O}^λ are noncommutative. This depends on the weight λ and also on a vector $\delta \in \mathbb{N}^{\tilde{Q}_0}$, which we now define. Let G be the finite subgroup of $\mathrm{SL}(2, \mathbb{k})$ corresponding to \tilde{Q} by the McKay correspondence. Then each vertex of \tilde{Q} corresponds to an irreducible representation W_i of G , and we set $\delta_i := \dim_{\mathbb{k}} W_i$. If we number the vertices of \tilde{Q} as in Figure 1, then

$$\begin{aligned} \tilde{A}_n : \quad \delta &= (\underbrace{1, 1, \dots, 1}_{n+1 \text{ times}}, 1) \\ \tilde{D}_n : \quad \delta &= (1, 1, \underbrace{2, 2, \dots, 2}_{n-3 \text{ times}}, 2, 1, 1) \\ \tilde{E}_6 : \quad \delta &= (1, 2, 1, 2, 3, 2, 1) \\ \tilde{E}_7 : \quad \delta &= (1, 2, 3, 4, 3, 2, 1, 2) \\ \tilde{E}_8 : \quad \delta &= (1, 2, 3, 4, 5, 6, 4, 2, 3). \end{aligned}$$

Theorem 2.13 [Crawley-Boevey and Holland 1998, Theorem 0.1, Theorem 0.4(1)]. *\mathcal{O}^λ is commutative if and only if $\lambda \cdot \delta = \sum_{i \in \tilde{Q}_0} \lambda_i \delta_i = 0$. In the case when $\lambda = \mathbf{0}$, \mathcal{O} is isomorphic to the coordinate ring of the Kleinian singularity corresponding to \tilde{Q} .*




Figure 1. Doubles of extended Dynkin graphs with the labelling of vertices and arrows that will be used throughout this paper. We have labelled the arrows only for those quivers in which we will need to refer to specific paths.

In fact, by [Crawley-Boevey and Holland 1998, Lemma 2.2], one may always assume that $\lambda \cdot \delta$ is either 0 or 1. When λ is a weight with $\lambda \cdot \delta = 0$ (so that \mathcal{O}^λ is commutative), if we define $\lambda' = (\lambda_0 + 1, \lambda_1, \dots, \lambda_n)$ then we consider $\mathcal{O}^{\lambda'}$ to be a noncommutative analogue of \mathcal{O}^λ . It is natural to ask if there is any relationship between the singularity categories of \mathcal{O}^λ and $\mathcal{O}^{\lambda'}$. In fact, it follows from Theorem 1.1 that the singularity categories of these \mathbb{k} -algebras are triangle equivalent.

When $\tilde{Q} = \tilde{A}_n$, it can be checked that \mathcal{O}^λ has a presentation of the form

$$\overline{\mathbb{k}\langle x, y, z \rangle}, \quad (2.14)$$

$$\left\langle \begin{array}{l} xz = (z + \lambda \cdot \delta)x, \quad xy = \prod_{i=0}^n (z + \sum_{j=1}^i \lambda_j) \\ yz = (z - \lambda \cdot \delta)y, \quad yx = \prod_{i=0}^n (z - \lambda \cdot \delta + \sum_{j=1}^i \lambda_j) \end{array} \right\rangle$$

When \mathcal{O}^λ is noncommutative, since there is no loss in generality in assuming $\lambda \cdot \delta = 1$, these are precisely the algebras considered by Hodges [1993] and Bavula [1992], where in the latter they were called generalised Weyl algebras.

In [Crawley-Boevey and Holland 1998, Section 7], the authors prove a number of results in the case where the weight λ is *dominant*, a term which we now define. Fix a total ordering $<$ on \mathbb{k} which also satisfies the following:

- (1) If $a < b$, then $a + c < b + c$ for all $c \in \mathbb{k}$.
- (2) On the integers, $<$ coincides with the usual order.
- (3) For any $a \in \mathbb{k}$, there exists $m \in \mathbb{Z}$ with $a < m$.

For example, when $\mathbb{k} = \mathbb{C}$ we may define $<$ by $z < z'$ if and only if $\operatorname{Re} z < \operatorname{Re} z'$, or $\operatorname{Re} z = \operatorname{Re} z'$ and $\operatorname{Im} z < \operatorname{Im} z'$. We then say that a weight $\lambda \in \mathbb{k}^{Q_0}$ is *dominant* if $\lambda_i \geq 0$ for all $i \in Q_0$.

When λ is dominant and Q is (extended) Dynkin, and Crawley-Boevey and Holland showed that it is easy to determine certain representation-theoretic properties of $\Pi^\lambda(Q)$. For example, we have the following useful result which we will use frequently in later sections:

Lemma 2.15 [Crawley-Boevey and Holland 1998, Lemma 7.1(1)]. *Suppose that Q is Dynkin, and let λ be a dominant weight for Q . Write Q_λ for the full subquiver supported on those vertices i with $\lambda_i = 0$. Then $\Pi^\lambda(Q) \cong \Pi(Q_\lambda)$. In particular, the projective $\Pi^\lambda(Q)$ -modules are the modules $e_i \Pi^\lambda(Q)$, where i is a vertex with $\lambda_i = 0$.*

2D. Restriction to quasidominant weights. A weaker version of dominance will play an important role in this paper, which we now define.

Definition 2.16. If \tilde{Q} is extended Dynkin, we say that a weight λ is *quasidominant* if $\lambda_i \geq 0$ for all $i \neq 0$, where $<$ is a total ordering on \mathbb{k} as above.

It turns out that we are able to restrict attention to quasidominant weights for the remainder of this paper. We now state this as an assumption, before explaining why this is the case.

Assumption 2.17. If λ is a weight for an extended Dynkin quiver \tilde{Q} , then we always assume that the weight λ is quasidominant unless explicitly stated otherwise.

To explain why this restriction is possible, we first recall a definition. Let Q be a quiver, and let $C = 2I - A$ be the generalised Cartan matrix of Q , where A is the adjacency matrix of the underlying graph of Q . For each loop-free vertex $i \in Q_0$, define the *dual reflection* $r_i : \mathbb{k}^{Q_0} \rightarrow \mathbb{k}^{Q_0}$ by

$$(r_i \lambda)_j = \lambda_j - C_{ij} \lambda_i.$$

It is easy to see that if \tilde{Q} is extended Dynkin then $\lambda \cdot \delta = (r_i \lambda) \cdot \delta$. We then have the following result, which appears in unpublished work of Boddington and Levy [2007].

Lemma 2.18. *Suppose that λ is a weight for an extended Dynkin quiver \tilde{Q} , and let ρ be a sequence of dual reflections at vertices other than the extending vertex 0. Then $\mathcal{O}^\lambda \cong \mathcal{O}^{\rho(\lambda)}$.*

This is a strengthening of [Crawley-Boevey and Holland 1998, Lemma 7.9], in which the authors established only a Morita equivalence between these two rings, rather than an isomorphism. Combining Lemma 2.18 with [loc. cit, Lemma 7.8], we have the following result, which justifies the restriction given in Assumption 2.17.

Lemma 2.19. *Suppose that λ is a weight for an extended Dynkin quiver \tilde{Q} . Then there exists a quasidominant weight λ' with $\mathcal{O}^\lambda \cong \mathcal{O}^{\lambda'}$.*

We will see later that this assumption allows one to easily read off a number of useful facts about the module category of \mathcal{O}^λ , and ultimately its singularity category as well. As a first example, if we restrict our attention to quasidominant weights then it is easy to detect whether \mathcal{O}^λ is singular.

Lemma 2.20. *If λ is a quasidominant weight for an extended Dynkin quiver \tilde{Q} , then \mathcal{O}^λ is singular if and only if $\lambda_i = 0$ for some $i \neq 0$.*

Proof. By [Crawley-Boevey and Holland 1998, Theorem 0.4(4)], \mathcal{O}^λ is singular if and only if $\lambda \cdot \alpha = 0$ for some Dynkin root α . The possible values of these Dynkin roots are not important to us; it suffices to know that they have the form $(0, \alpha') \in \mathbb{Z}^{n+1}$ where, in particular, α' has entirely nonnegative or nonpositive entries, and has at least one nonzero entry. In addition, $\varepsilon_i \in \mathbb{Z}^{n+1}$ for $1 \leq i \leq n$ is always a Dynkin root, where ε_i is the i -th coordinate vector (here the entries are indexed from 0 to n). Therefore, if $\lambda_i = 0$ for some $i \neq 0$ then $\lambda \cdot \alpha = 0$ for the Dynkin root $\alpha = \varepsilon_i$, while if $\lambda_i \neq 0$ for all $i \neq 0$, then necessarily $\lambda \cdot \alpha \neq 0$ for all Dynkin roots α . The result then follows. \square

3. The singularity category of $\mathcal{O}^\lambda(\tilde{Q})$ as a \mathbb{k} -linear category

Our first step in determining $\text{MCM-}\mathcal{O}^\lambda$ is to determine its structure as an additive category, or indeed as a \mathbb{k} -linear category. We first identify an important module.

Lemma 3.1. *$\Pi^\lambda e_0$ is a finitely generated \mathcal{O}^λ -module, and it satisfies $\text{End}_{\mathcal{O}^\lambda}(\Pi^\lambda e_0) = \Pi^\lambda$. Moreover, $\Pi^\lambda e_0$ is maximal Cohen–Macaulay.*

Proof. The first part of the statement follows from [Montgomery and Small 1981, Lemma 1]. To determine the endomorphism ring, first note that, by [Crawley-Boevey and Holland 1998, Lemma 1.4, Corollary 3.5], Π^λ is Morita equivalent to a ring which is a maximal order and hence is itself a maximal order. The claim then follows from the results in [Crawley-Boevey 1999, Section 5.4].

For the final claim, note that $\Pi^\lambda e_0$ is a reflexive \mathcal{O}^λ -module by [Crawley-Boevey 1999, Section 5.4]. Therefore, since $\Pi^\lambda e_0$ is finitely generated, and since $\text{i. dim } \mathcal{O}^\lambda \leq 2$ by [Crawley-Boevey and Holland 1998, Theorem 1.6], Lemma 2.11 implies that $\Pi^\lambda e_0$ is maximal Cohen–Macaulay. \square

Write $V_i = e_i \Pi^\lambda e_0$; we shall refer to these \mathcal{O}^λ -modules as *vertex modules*, and they will play an important role in determining $\underline{\text{MCM}}\text{-}\mathcal{O}^\lambda$. Using Lemma 3.1, we are able to calculate the Hom spaces between the vertex modules.

Corollary 3.2. *We have $\text{Hom}_{\mathcal{O}^\lambda}(V_i, V_j) = e_j \Pi^\lambda e_i$.*

Proof. By Lemma 3.1, $\Pi^\lambda = \text{End}_{\mathcal{O}^\lambda}(\Pi^\lambda e_0) = \bigoplus_{k,\ell} \text{Hom}_{\mathcal{O}^\lambda}(e_k \Pi^\lambda e_0, e_\ell \Pi^\lambda e_0)$. Multiplying on the left by e_j kills each Hom space with $\ell \neq j$, while multiplying on the right by e_i kills each Hom space with $k \neq i$. It follows that

$$e_j \Pi^\lambda e_i = e_j \left(\bigoplus_{k,\ell} \text{Hom}_{\mathcal{O}^\lambda}(V_k, V_\ell) \right) e_i = e_j \text{Hom}_{\mathcal{O}^\lambda}(V_i, V_j) e_i = \text{Hom}_{\mathcal{O}^\lambda}(V_i, V_j),$$

as claimed. \square

This allows us to determine the stable endomorphism ring of $\Pi^\lambda e_0$. We fix some notation which will be used throughout the rest of this paper: write Q_λ for the full subquiver of \tilde{Q} with vertex set $I_\lambda := \{i \in \{1, \dots, n\} \mid \lambda_i = 0\}$.

Lemma 3.3. *We have $\underline{\text{End}}_{\mathcal{O}^\lambda}(\Pi^\lambda e_0) \cong \Pi(Q_\lambda)$.*

Proof. Write $\mu = (\lambda_1, \dots, \lambda_n)$. By Corollary 3.2, we have that

$$(\Pi^\lambda e_0)^* = \bigoplus_i \text{Hom}_{\mathcal{O}^\lambda}(e_i \Pi^\lambda e_0, e_0 \Pi^\lambda e_0) = \bigoplus_i e_0 \Pi^\lambda e_i = e_0 \Pi^\lambda.$$

Then, noting that $\Pi^\lambda e_0 (\Pi^\lambda e_0)^* = \Pi^\lambda e_0 \Pi^\lambda$, we have $\underline{\text{End}}_{\mathcal{O}^\lambda}(\Pi^\lambda e_0) \cong \Pi^\lambda / (\Pi^\lambda e_0 \Pi^\lambda) \cong \Pi^\mu(Q)$. Since the entries of μ are all ≥ 0 by Assumption 2.17 and Q is Dynkin, Lemma 2.15 tells us that $\Pi^\mu(Q)$ is isomorphic to the preprojective algebra supported on the vertices i of Q with $\mu_i = 0$; that is, $\Pi^\mu(Q) \cong \Pi(Q_\lambda)$. Therefore $\underline{\text{End}}_{\mathcal{O}^\lambda}(\Pi^\lambda e_0) \cong \Pi(Q_\lambda)$. \square

We are also able to determine when a vertex module is projective. It turns out that this is the case precisely when the corresponding vertex is deleted when passing from \tilde{Q} to Q_λ .

Lemma 3.4. *If $i = 0$ or $\lambda_i \neq 0$, then V_i is a projective \mathcal{O}^λ -module.*

Proof. When $i = 0$ this is clear. So suppose that $i \neq 0$ and $\lambda_i \neq 0$. Then, as in the proof of Lemma 3.3, $e_i = 0$ in $\Pi^\lambda / \Pi^\lambda e_0 \Pi^\lambda$ and so $e_i \in \Pi^\lambda e_0 \Pi^\lambda$. But then, using Corollary 3.2, $V_i V_i^* = e_i \Pi^\lambda e_0 \Pi^\lambda e_i \ni e_i^3 = e_i$, where e_i is the identity element of $\text{End}_{\Pi^\lambda}(V_i) = e_i \Pi^\lambda e_i$, and so V_i is projective by the dual basis lemma (see [Lam 1999, (2.9)]). \square

It follows that the vertex modules V_i satisfying $\lambda_i \neq 0$ are equal to the zero object in the singularity category, so that $\Pi^\lambda e_0$ and $\bigoplus_{i \in I_\lambda} V_i$ are isomorphic in $\underline{\text{MCM}}\text{-}\mathcal{O}^\lambda$. When working in the stable module category, we will sometimes refer to those vertex modules whose corresponding weight is zero as nonprojective vertex modules.

Proposition 3.5. (1) $\text{MCM}\text{-}\mathcal{O}^\lambda = \text{add } \Pi^\lambda e_0$.

(2) $\underline{\text{MCM}}\text{-}\mathcal{O}^\lambda = \text{add } \Pi^\lambda e_0 = \text{add}(\bigoplus_{i \in I_\lambda} V_i)$.

Proof. (1) First note that \mathcal{O}^λ is Gorenstein and that, using [Crawley-Boevey and Holland 1998, Theorem 1.5],

$$\text{gl.dim End}_{\mathcal{O}^\lambda}(\Pi^\lambda e_0) = \text{gl.dim } \Pi^\lambda \leq 2.$$

Since $\Pi^\lambda e_0$ has \mathcal{O}^λ as a direct summand, the first claim then follows from Proposition 2.8.

(2) Part (1) immediately implies that $\underline{\text{MCM}}\text{-}\mathcal{O}^\lambda = \text{add } \Pi^\lambda e_0 = \text{add}(\bigoplus_i V_i)$. But projective modules become the zero object when passing to the stable module category, so the result follows by Lemma 3.4. \square

We recall that an additive category is said to be *Krull–Schmidt* if every object decomposes into a finite direct sum of objects, each of which has a local endomorphism ring. By [Krause 2015, Theorem 4.2], this decomposition is unique up to reordering.

Theorem 3.6. *The functor $\underline{\text{Hom}}_{\mathcal{O}^\lambda}(\Pi^\lambda e_0, -)$ induces a \mathbb{k} -linear equivalence*

$$\underline{\text{MCM}}\text{-}\mathcal{O}^\lambda \simeq \text{proj-}\Pi(Q_\lambda).$$

Proof. By [Krause 2015, Proposition 2.3], the functor

$$\underline{\text{Hom}}_{\mathcal{O}^\lambda}(\Pi^\lambda e_0, -) : \underline{\text{mod}}\text{-}\mathcal{O}^\lambda \rightarrow \text{mod-}\underline{\text{End}}_{\mathcal{O}^\lambda}(\Pi^\lambda e_0) = \text{mod-}\Pi(Q_\lambda)$$

induces a fully faithful \mathbb{k} -linear functor $\text{add } \Pi^\lambda e_0 \rightarrow \text{proj-}\Pi(Q_\lambda)$, where $\text{add } \Pi^\lambda e_0 = \underline{\text{MCM}}\text{-}\mathcal{O}^\lambda$ by Proposition 3.5. Since $\Pi(Q_\lambda)$ is finite-dimensional [Białkowski et al. 2007, Proposition 2.1], $\text{mod-}\Pi(Q_\lambda)$ is Krull–Schmidt and hence so too is $\text{proj-}\Pi(Q_\lambda)$. Therefore, to establish essential surjectivity of the functor $\underline{\text{Hom}}_{\mathcal{O}^\lambda}(\Pi^\lambda e_0, -)$, it suffices to show that we can hit each indecomposable projective $e_i \Pi(Q_\lambda)$, where $i \in I_\lambda$. Indeed, we have

$$\underline{\text{Hom}}_{\mathcal{O}^\lambda}(\Pi^\lambda e_0, V_i) = \frac{e_i \Pi^\lambda}{e_i \Pi^\lambda e_0 \Pi^\lambda} = \frac{e_i \Pi^\lambda}{e_i \Pi^\lambda \cap \Pi^\lambda e_0 \Pi^\lambda} = e_i \frac{\Pi^\lambda}{\Pi^\lambda e_0 \Pi^\lambda} = e_i \Pi(Q_\lambda),$$

and so the functor is also essentially surjective. We therefore have the claimed equivalence. \square

It follows that $\underline{\text{MCM}}\text{-}\mathcal{O}^\lambda$ is nontrivial if and only if $\lambda_i = 0$ for some $i \neq 0$ which, by Lemma 2.20, happens precisely when \mathcal{O}^λ is singular; this is consistent with the more general fact that $\mathcal{D}_{\text{sg}}(R)$ is nontrivial if and only if R is singular. Moreover, the vertex modules V_i with $i = 0$, or with $i \neq 0$ and $\lambda_i \neq 0$, are those which are projective and hence vanish in $\underline{\text{MCM}}\text{-}\mathcal{O}^\lambda$. This is reflected by the fact that these are the vertices which are deleted to obtain Q_λ .

As an immediate consequence of (the proof of) Theorem 3.6, we have the following result:

Corollary 3.7. *$\underline{\text{MCM}}\text{-}\mathcal{O}^\lambda$ is a Krull–Schmidt category.*

Remark 3.8. By Proposition 3.5, the objects of $\underline{\text{MCM}}\text{-}\mathcal{O}^\lambda$ are direct summands of finite direct sums of the nonprojective vertex modules. Since these vertex modules are indecomposable and $\underline{\text{MCM}}\text{-}\mathcal{O}^\lambda$ is Krull–Schmidt, in fact every object of $\underline{\text{MCM}}\text{-}\mathcal{O}^\lambda$ is isomorphic to a finite direct sum of vertex modules.

The following two corollaries are then immediate from Theorem 3.6:

Corollary 3.9. Suppose that \tilde{Q} is an extended Dynkin quiver and $Q_\lambda = Q^{(1)} \sqcup \cdots \sqcup Q^{(r)}$ is a disjoint union of connected quivers $Q^{(i)}$, which are therefore necessarily Dynkin. Then there is a \mathbb{k} -linear equivalence

$$\underline{\text{MCM}}\text{-}\mathcal{O}^\lambda \simeq \bigoplus_{i=1}^r \text{proj-}\Pi(Q^{(i)}).$$

Corollary 3.10. Let \tilde{Q} and \tilde{Q}' be extended Dynkin quivers (not necessarily of the same type) and let λ and λ' be quasidominant weights for \tilde{Q} and \tilde{Q}' , respectively. If $Q_\lambda \cong Q_{\lambda'}$, then there is a \mathbb{k} -linear equivalence

$$\underline{\text{MCM}}\text{-}\mathcal{O}^\lambda(\tilde{Q}) \simeq \underline{\text{MCM}}\text{-}\mathcal{O}^{\lambda'}(\tilde{Q}').$$

It is illustrative to apply Theorem 3.6 (and its corollaries) to an example.

Example 3.11. Suppose that $\tilde{Q} = \tilde{\mathbb{A}}_5$, and consider the deformation \mathcal{O}^λ where the weight λ is indicated in red on the left-hand quiver in Figure 2.

By Corollary 3.9, there is a \mathbb{k} -linear equivalence $\underline{\text{MCM}}\text{-}\mathcal{O}^\lambda \simeq \text{proj-}\Pi(\mathbb{A}_3) \oplus \text{proj-}\Pi(\mathbb{A}_1)$. In more suggestive notation, we can write this equivalence as

$$\mathcal{D}_{\text{sg}}(\mathcal{O}^\lambda) \simeq \mathcal{D}_{\text{sg}}(R_{\mathbb{A}_3}) \oplus \mathcal{D}_{\text{sg}}(R_{\mathbb{A}_1}),$$

and so it is sensible to consider \mathcal{O}^λ as having an \mathbb{A}_3 singularity and an \mathbb{A}_1 singularity.

If we more concretely set $\lambda = (-1, 0, 0, 0, 1, 0)$ (respectively, $\lambda = (0, 0, 0, 0, 1, 0)$) then \mathcal{O}^λ is commutative (respectively, noncommutative). In this case we can use (2.14) to write down a presentation for \mathcal{O}^λ . In particular, we have a \mathbb{k} -linear equivalence

$$\mathcal{D}_{\text{sg}} \frac{\mathbb{k}[x, y, z]}{\langle xy - z^4(z+1)^2 \rangle} \simeq \mathcal{D}_{\text{sg}} \frac{\mathbb{k}\langle x, y, z \rangle}{\left\langle \begin{array}{l} xz = (z+1)x, \quad xy = z^4(z+1)^2 \\ yz = (z-1)y, \quad yx = (z-1)^4 z^2 \end{array} \right\rangle}.$$




Figure 2. A weight λ for an $\tilde{\mathbb{A}}_5$ quiver, and the corresponding category $\underline{\text{MCM}}\text{-}\mathcal{O}^\lambda$.

4. The singularity category of $\mathcal{O}^\lambda(\tilde{Q})$ as a triangulated category

We are now able to complete the proof of Theorem 1.1 which gives a complete description of the singularity category of $\mathcal{O}^\lambda(\tilde{Q})$. To do this, we need to show that the induced triangulated structures on the right-hand sides of the \mathbb{k} -linear equivalences

$$\mathcal{D}_{\text{sg}}(\mathcal{O}^\lambda(\tilde{Q})) \simeq \underline{\text{MCM}}\text{-}\mathcal{O}^\lambda(\tilde{Q}) \simeq \bigoplus_{i=1}^r \text{proj-}\Pi(Q^{(i)}), \quad (4.1)$$

$$\bigoplus_{i=1}^r \mathcal{D}_{\text{sg}}(R_{Q^{(i)}}) \simeq \bigoplus_{i=1}^r \text{proj-}\Pi(Q^{(i)}) \quad (4.2)$$

are the same. We achieve this in by showing that the summands $\text{proj-}\Pi(Q^{(i)})$ in (4.1) are in fact triangulated subcategories of $\bigoplus_{i=1}^r \text{proj-}\Pi(Q^{(i)})$, and that the triangulated structure on each $\text{proj-}\Pi(Q^{(i)})$ is essentially unique. To establish the latter of these, we use the following result:

Theorem 4.3 [Keller 2018, Corollary 2]. *Let \mathcal{T} and \mathcal{T}' be Krull–Schmidt \mathbb{k} -linear triangulated categories which are finite, connected, algebraic and standard. If \mathcal{T} and \mathcal{T}' are equivalent as \mathbb{k} -linear categories, then they are in fact equivalent as triangulated categories.*

We note that, if Q is Dynkin, $\text{proj-}\Pi(Q)$ (and the \mathbb{k} -linearly equivalent category $\mathcal{D}_{\text{sg}}(R_Q)$) are finite, connected, and standard since they are \mathbb{k} -linearly equivalent to certain orbit categories which are known to have these properties (see [Amiot et al. 2015, Remark 5.9]). Therefore, if we can show that each $\text{proj-}\Pi(Q^{(i)})$ is an algebraic triangulated subcategory under the \mathbb{k} -linear equivalence (4.1), then each \mathbb{k} -linear equivalence $\text{proj-}\Pi(Q^{(i)}) \simeq \mathcal{D}_{\text{sg}}(R_{Q^{(i)}})$ is in fact a triangle equivalence, which will prove Theorem 1.1 from the introduction.

We must first show that the translation functor Σ induced on the right-hand side of (4.1) *preserves connected components*, in the sense that it restricts to an autoequivalence of each of the subcategories $\text{proj-}\Pi(Q^{(i)})$. Writing P_i for the indecomposable projective $\Pi(Q_\lambda)$ -module corresponding to vertex i in Q_λ , it follows from the Krull–Schmidt property of $\text{proj-}\Pi(Q_\lambda)$ that Σ permutes the P_i . We write σ for the induced permutation of the vertices. This allows us to make the following observation:

Lemma 4.4. *With the above setup, σ is a graph automorphism of Q_λ .*

Proof. First note that the spaces $\text{Hom}_{\Pi(Q_\lambda)}(P_i, P_j) = e_j \Pi(Q_\lambda) e_i$ can be graded by path length, and that vertex i and vertex j are adjacent in Q_λ if and only if there is a degree 1 morphism in $\text{Hom}_{\Pi(Q_\lambda)}(P_i, P_j)$. Applying Σ , this is equivalent to $\text{Hom}_{\Pi(Q_\lambda)}(P_{\sigma(i)}, P_{\sigma(j)})$ containing a degree 1 morphism, which happens if and only if $\sigma(i)$ and $\sigma(j)$ are adjacent in Q_λ . That is, σ is a graph automorphism of Q_λ . \square

If the $Q^{(i)}$ are pairwise nonisomorphic, then the fact that the induced translation functor has to be a graph automorphism forces it to preserve connected components, as required. This leaves only the cases where some of the $Q^{(i)}$ are isomorphic, and one might hope to abstractly prove that the translation functor preserves connected components. Unfortunately, the following example shows that one should not expect this to be the case.

Example 4.5. Let T be a Krull–Schmidt \mathbb{k} -linear category with only two indecomposable objects U and V , and suppose these objects satisfy $\text{Hom}_T(U, V) = 0 = \text{Hom}_T(V, U)$ and $\text{End}_T(U) = \mathbb{k} = \text{End}_T(V)$. For example, this is the case for $\text{MCM-}\mathcal{O}^\lambda(\tilde{\mathbb{A}}_3)$ when $(\lambda_0, \lambda_1, \lambda_2, \lambda_3) = (0, 0, 1, 0)$, since in this case it is \mathbb{k} -linearly equivalent to $\text{proj-}\Pi(\mathbb{A}_1) \oplus \text{proj-}\Pi(\mathbb{A}_1)$. This category has two possible triangulated structures: the first has $\Sigma = \text{id}$, and the distinguished triangles are isomorphic to direct sums and rotations of

$$U \xrightarrow{\text{id}} U \rightarrow 0 \rightarrow U \quad \text{and} \quad V \xrightarrow{\text{id}} V \rightarrow 0 \rightarrow V,$$

and the second option has $\Sigma U = V$ and $\Sigma V = U$, and the distinguished triangles are isomorphic to direct sums and rotations of

$$U \xrightarrow{\text{id}} U \rightarrow 0 \rightarrow V.$$

The first example decomposes into a direct sum of two triangulated subcategories, while the second example does not.

While the above example shows that one should not expect to be able to abstractly prove that the translation functor preserves connected components, this is essentially the only counterexample. The following proof is due to Jeremy Rickard, and we thank him for allowing us to reproduce it:

Lemma 4.6. *Suppose that \mathcal{T} is a Krull–Schmidt \mathbb{k} -linear triangulated category with finitely many indecomposables which decomposes as a \mathbb{k} -linear category as*

$$\mathcal{T} = \bigoplus_{i=1}^n \mathcal{T}_i.$$

Suppose that the translation functor Σ satisfies $\Sigma \mathcal{T}_i = \mathcal{T}_j$ for some $i \neq j$. Then \mathcal{T}_i and \mathcal{T}_j each have only one isoclass of indecomposable objects.

Proof. Let $\alpha : X \rightarrow Y$ be a nonzero morphism between two indecomposable objects of \mathcal{T}_i , and complete to a triangle

$$X \xrightarrow{\alpha} Y \xrightarrow{\beta} Z \xrightarrow{\gamma} \Sigma X,$$

where $\Sigma X \in \mathcal{T}_j$ by assumption. We claim that every indecomposable summand of Z lies in \mathcal{T}_j . To this end, suppose that $Z = Z' \oplus Z''$ where $Z' \in \mathcal{T}_j$ and $Z'' \in \bigoplus_{k \neq j} \mathcal{T}_k$, and write $\gamma = (\gamma', 0)$. The map γ' gives rise to a triangle $Z' \xrightarrow{\gamma'} \Sigma X \rightarrow Y' \rightarrow \Sigma Z'$ and rotating yields the triangle $X \rightarrow \Sigma^{-1} Y' \rightarrow Z' \xrightarrow{\gamma'} \Sigma X$. The direct sum of this triangle with the triangle $0 \rightarrow Z'' \rightarrow Z'' \rightarrow 0$ is a triangle isomorphic to the triangle $X \rightarrow Y \rightarrow Z \rightarrow \Sigma X$, and so $Y \cong \Sigma^{-1} Y' \oplus Z''$. By indecomposability of Y , we therefore have $\Sigma^{-1} Y' = 0$ or $Z'' = 0$. If $\Sigma^{-1} Y' = 0$ then $Y \cong Z''$ and $\Sigma X \cong Z'$. Our original triangle becomes

$$X \xrightarrow{\alpha} Z'' \rightarrow \Sigma X \oplus Z'' \rightarrow \Sigma X$$

which is isomorphic to the direct sum of the triangles $X \rightarrow 0 \rightarrow \Sigma X \rightarrow \Sigma X$ and $0 \rightarrow Z'' \rightarrow Z'' \rightarrow 0$. This means that α is the zero map, contrary to our assumption, and so we must have $Z'' = 0$, establishing

the claim. Now, since every indecomposable summand of Z lies in \mathcal{T}_j , β is the zero map. Applying $\text{Hom}(Y, -)$, we get an exact sequence

$$\text{Hom}(Y, X) \xrightarrow{\alpha \circ -} \text{Hom}(Y, Y) \rightarrow \text{Hom}(Y, Z)$$

where the last term is 0. By exactness, there exists $\alpha' : Y \rightarrow X$ with $\alpha\alpha' = \text{id}_Y$. Since \mathcal{T} is Krull–Schmidt the endomorphism ring of X is local, which implies that the idempotent map $\alpha'\alpha$ is a unit and therefore equal to id_X . Therefore $\alpha : X \rightarrow Y$ is an isomorphism, and so \mathcal{T}_i (and hence \mathcal{T}_j) has only one indecomposable object, up to isomorphism. \square

Therefore, to show that the induced translation functor on $\bigoplus_{i=1}^r \text{proj-}\Pi(Q^{(i)})$ from the second \mathbb{k} -linear equivalence in (4.1) preserves connected components, we only need to consider the case when there exist $Q^{(i)}$ and $Q^{(j)}$, $i \neq j$, with $Q^{(i)} = \mathbb{A}_1 = Q^{(j)}$. It suffices to show that, for the corresponding objects $V_i, V_j \in \underline{\text{MCM}}\text{-}\mathcal{O}^\lambda(\tilde{Q})$, we have $\Sigma V_i = V_i$ and $\Sigma V_j = V_j$. To this end, we first have the following result:

Proposition 4.7. *Let Q be a non-Dynkin quiver with no oriented cycles, and with vertices labelled $\{0, 1, \dots, n\}$. Write $\Pi(Q)$ for the preprojective algebra of Q , and write $V_i = e_i \Pi(Q) e_0$, which is a right $e_0 \Pi(Q) e_0$ -module. Then, for any $i \neq 0$, there exists a short exact sequence of $e_0 \Pi(Q) e_0$ -modules*

$$0 \rightarrow V_i \rightarrow \bigoplus_{j \in \partial i} V_j \rightarrow V_i \rightarrow 0,$$

where ∂i is the set of vertices adjacent to i in Q .

Proof. By [Brenner et al. 2002, Proposition 4.2], there is an exact sequence of $\Pi(Q)$ -modules

$$0 \rightarrow e_i \Pi(Q) \rightarrow \bigoplus_{j \in \partial i} e_j \Pi(Q) \rightarrow e_i \Pi(Q) \rightarrow S_i \rightarrow 0, \quad (4.8)$$

where S_i is the simple module at vertex i . Noting that $e_0 \Pi(Q)$ is a direct summand of $\Pi(Q)$ and hence projective, applying $\text{Hom}_{\Pi(Q)}(e_0 \Pi(Q), -)$ yields an exact sequence

$$\begin{aligned} 0 \rightarrow \text{Hom}_{\Pi(Q)}(e_0 \Pi(Q), e_i \Pi(Q)) &\rightarrow \bigoplus_{j \in \partial i} \text{Hom}_{\Pi(Q)}(e_0 \Pi(Q), e_j \Pi(Q)) \\ &\rightarrow \text{Hom}_{\Pi(Q)}(e_0 \Pi(Q), e_i \Pi(Q)) \rightarrow \text{Hom}_{\Pi(Q)}(e_0 \Pi(Q), S_i) \rightarrow 0. \end{aligned}$$

We also have $\text{Hom}_{\Pi(Q)}(e_0 \Pi(Q), e_k \Pi(Q)) = V_k$ and, since $i \neq 0$, $\text{Hom}_{\Pi(Q)}(e_0 \Pi(Q), S_i) = 0$. We therefore have exactness of

$$0 \rightarrow V_i \rightarrow \bigoplus_{j \in \partial i} V_j \rightarrow V_i \rightarrow 0,$$

as claimed. \square

In particular this result holds for extended Dynkin quivers, where we remark that if we wish to apply it to an $\tilde{\mathbb{A}}_n$ quiver then we must orient the arrows so that there are no oriented cycles; this does not change the isomorphism class of $\Pi(\tilde{\mathbb{A}}_n)$ [Crawley-Boevey and Holland 1998, Lemma 2.2].

Remark 4.9. The above result may or may not fail for Dynkin quivers, depending on how the vertices are labelled. For example, when $Q = \mathbb{A}_3$ where the vertices are labelled as follows,

$$0 \text{ --- } 1 \text{ --- } 2$$

then the complexes of interest to us are

$$0 \rightarrow V_1 \rightarrow V_0 \oplus V_2 \rightarrow V_1 \rightarrow 0, \quad \text{and} \quad 0 \rightarrow V_2 \rightarrow V_1 \rightarrow V_2 \rightarrow 0.$$

Since $\dim_{\mathbb{k}} V_0 = 1$, $\dim_{\mathbb{k}} V_1 = 1$, and $\dim_{\mathbb{k}} V_2 = 1$, the first of these is exact while the second is not. If instead we label the vertices of Q as follows,

$$1 \text{ --- } 0 \text{ --- } 2$$

then the complexes of interest to us are

$$0 \rightarrow V_1 \rightarrow V_0 \rightarrow V_1 \rightarrow 0, \quad \text{and} \quad 0 \rightarrow V_2 \rightarrow V_0 \rightarrow V_2 \rightarrow 0,$$

and both of these are exact since $\dim_{\mathbb{k}} V_0 = 2$, $\dim_{\mathbb{k}} V_1 = 1$, and $\dim_{\mathbb{k}} V_2 = 1$.

We now use Proposition 4.7 to show that the induced translation functor on $\bigoplus_{i=1}^r \text{proj-}\Pi(Q^{(i)})$ preserves connected components.

Proposition 4.10. *Let \tilde{Q} be an extended Dynkin quiver and λ be a quasidominant weight for \tilde{Q} . Write $Q_\lambda = Q^{(1)} \sqcup \cdots \sqcup Q^{(r)}$ as a disjoint union of connected quivers $Q^{(i)}$, which are therefore necessarily Dynkin. Consider the triangulated structure on $\bigoplus_{i=1}^r \text{proj-}\Pi(Q^{(i)})$ induced by the \mathbb{k} -linear equivalence*

$$\underline{\text{MCM}}\text{-}\mathcal{O}^\lambda \simeq \bigoplus_{i=1}^r \text{proj-}\Pi(Q^{(i)})$$

of Corollary 3.9, and let Σ be the translation functor. Then each $\text{proj-}\Pi(Q^{(i)})$ is invariant under Σ .

Proof. By Lemma 4.6 and the discussion following it, the only situation in which there exist $\text{proj-}\Pi(Q^{(i)})$ which are not necessarily invariant under Σ is when we have multiple $Q^{(i)}$ equal to \mathbb{A}_1 . Working in $\underline{\text{MCM}}\text{-}\mathcal{O}^\lambda(\tilde{Q})$, this happens if and only if there is some vertex i with $\lambda_i = 0$, and if j is adjacent to i then either $j = 0$ or $\lambda_j \neq 0$; in particular, the modules V_j corresponding to these vertices are projective as $\mathcal{O}^\lambda(\tilde{Q})$ -modules by Lemma 3.4. By Proposition 4.7, we have an exact sequence of $e_0\Pi(\tilde{Q})e_0$ -modules

$$0 \rightarrow V_i \xrightarrow{\phi} \bigoplus_{j \in \partial i} V_j \xrightarrow{\psi} V_i \rightarrow 0. \quad (4.11)$$

Now consider (4.11) as a sequence of modules over $\mathcal{O}^\lambda(\tilde{Q})$. It is a complex since the composition $\psi\phi$ is equal to the (undeformed) preprojective relation at vertex i , which is equal to $\lambda_i e_i = 0$. Filtering $\Pi(\tilde{Q})$ and $\mathcal{O}^\lambda(\tilde{Q})$ by path length we obtain a sequence of associated graded modules, which is in fact the exact sequence (4.11). It is standard (see [McConnell and Robson 2001, Proposition 7.6.14]) that this implies

that (4.11) is exact as a sequence of modules over $\mathcal{O}^\lambda(\tilde{Q})$. To summarise, we have an exact sequence of $\mathcal{O}^\lambda(\tilde{Q})$ -modules

$$0 \rightarrow V_i \rightarrow \bigoplus_{j \in \partial i} V_j \rightarrow V_i \rightarrow 0$$

whose middle term is projective. It follows from the definition of the translation functor that $\Sigma V_i = V_i$ in $\underline{\text{MCM}}\text{-}\mathcal{O}^\lambda(\tilde{Q})$. Thus each $\text{proj-}\Pi(Q^{(i)})$ is invariant under the induced translation functor on $\bigoplus_{i=1}^r \text{proj-}\Pi(Q^{(i)})$. \square

We now seek to prove Theorem 1.1. Retaining all of the above notation, for each $1 \leq i \leq r$, define

$$\mathcal{W}_i := \{V_j \mid j \in Q_0^{(i)}\}, \quad \mathcal{C}_i := \text{add} \left(V_0 \oplus \bigoplus_{j \in Q_0^{(i)}} V_j \right) \quad \text{and} \quad \mathcal{T}_i := \text{add} \left(\bigoplus_{j \in Q_0^{(i)}} V_j \right),$$

where the latter two are viewed as subcategories of $\text{MCM-}\mathcal{O}^\lambda$ and $\underline{\text{MCM}}\text{-}\mathcal{O}^\lambda$, respectively. It will also be convenient to write

$$M_i = \bigoplus_{j \in (Q_\lambda)_0 \setminus Q_0^{(i)}} V_j,$$

and to set

$$\mathcal{W}_i^c := \{V_j \mid j \in (Q_\lambda)_0 \setminus Q_0^{(i)}\}, \quad \mathcal{C}_i^c := \text{add}(V_0 \oplus M_i) \quad \text{and} \quad \mathcal{T}_i^c := \text{add } M_i.$$

Observe that we can decompose $\underline{\text{MCM}}\text{-}\mathcal{O}^\lambda$ as

$$\underline{\text{MCM}}\text{-}\mathcal{O}^\lambda = \bigoplus_{i=1}^r \mathcal{T}_i$$

as \mathbb{k} -linear categories. We wish to show that this is also a decomposition into triangulated subcategories. To do this, we first prove a result which shows that the \mathcal{C}_i are Frobenius subcategories of the Frobenius category $\text{MCM-}\mathcal{O}^\lambda$. We call a subcategory \mathcal{B} of an exact category \mathcal{A} *extension-closed* if whenever we have a conflation $X \rightarrow Y \rightarrow Z$ with $X, Z \in \mathcal{B}$ then necessarily $Y \in \mathcal{B}$. Furthermore, an extension-closed subcategory \mathcal{B} is called *admissible* provided that every $B \in \mathcal{B}$ fits into conflations $B \rightarrow P \rightarrow B'$ and $B'' \rightarrow Q \rightarrow B$ with $B', B'' \in \mathcal{B}$ and where P, Q are projective in \mathcal{A} . We remark that an admissible subcategory of a Frobenius category is itself Frobenius; see [Chen 2012, Section 2].

Lemma 4.12. *For each i , the subcategory \mathcal{C}_i satisfies the following property: if $X \rightarrow Y \rightarrow Z$ is a conflation in $\text{MCM-}\mathcal{O}^\lambda$ such that two of the three objects are in \mathcal{C}_i , then the third object is also in \mathcal{C}_i . Consequently, \mathcal{C}_i is a Frobenius subcategory of $\text{MCM-}\mathcal{O}^\lambda$.*

Proof. We only show that if $X \rightarrow Y \rightarrow Z$ is a conflation with $X, Y \in \mathcal{C}_i$ then $Z \in \mathcal{C}_i$, with the other cases being similar. So suppose that we have such a conflation. Since $\underline{\text{MCM}}\text{-}\mathcal{O}^\lambda \simeq \text{proj-}\Pi(Q_\lambda)$ and this category is Krull–Schmidt, we have $Z \oplus P \cong U \oplus U' \oplus Q$ in $\text{MCM-}\mathcal{O}^\lambda$, where $U \in \mathcal{W}_i$, $U' \in \mathcal{W}_i^c$,

and P, Q are projective. This conflation gives rise to a triangle $X \rightarrow Y \rightarrow Z \rightarrow \Sigma X$ in $\underline{\text{MCM}}\text{-}\mathcal{O}^\lambda$, and applying the functor $\underline{\text{Hom}}_{\mathcal{O}^\lambda}(M_i, -)$ yields an exact sequence

$$\underline{\text{Hom}}_{\mathcal{O}^\lambda}(M_i, Y) \rightarrow \underline{\text{Hom}}_{\mathcal{O}^\lambda}(M_i, Z) \rightarrow \underline{\text{Hom}}_{\mathcal{O}^\lambda}(M_i, \Sigma X).$$

Now $\Sigma X \in \mathcal{C}_i$ by Proposition 4.10 and $Y \in \mathcal{C}_i$ by definition, while $M_i \in \mathcal{C}_i^c$, so both of the flanking terms are 0. This implies that the middle term, which is equal to $\underline{\text{Hom}}_{\mathcal{O}^\lambda}(M_i, U')$, is also 0. But this means that $U' = 0$, and hence $Z \oplus P \in \mathcal{C}_i$. Since, by definition, \mathcal{C}_i is closed under direct summands, it follows that $Z \in \mathcal{C}_i$ as required.

For the final claim, first notice that the above paragraph tells us that \mathcal{C}_i is extension-closed. Moreover, given an object $C \in \mathcal{C}_i$, since $\text{MCM-}\mathcal{O}^\lambda$ is Frobenius we can always find conflations $C \rightarrow P \rightarrow Z$ and $X \rightarrow Q \rightarrow C$ with $X, Z \in \text{MCM-}\mathcal{O}^\lambda$ and P, Q projective. Since projective \mathcal{O}^λ -modules are direct summands of sums of copies of \mathcal{O}^λ , we have $P, Q \in \mathcal{C}_i$ by definition, and then the previous paragraph tells us that $X, Z \in \mathcal{C}_i$. Therefore \mathcal{C}_i is admissible and hence Frobenius. \square

This allows us to prove our main theorem.

Theorem 4.13. *Let \tilde{Q} and Q_λ be as in Corollary 3.9. Then the \mathbb{k} -linear equivalence*

$$\underline{\text{MCM}}\text{-}\mathcal{O}^\lambda \simeq \bigoplus_{i=1}^r \text{proj-}\Pi(Q^{(i)}),$$

of Corollary 3.9 is a triangle equivalence, where the right-hand side is a decomposition into triangulated subcategories satisfying $\text{proj-}\Pi(Q^{(i)}) \simeq \mathcal{D}_{\text{sg}}(R_{Q^{(i)}})$.

Proof. By Lemma 4.12, we know that \mathcal{C}_i is a Frobenius subcategory of $\text{MCM-}\mathcal{O}^\lambda$. Using [Arentz-Hansen 2019, Theorem 3.15(2)], it follows that \mathcal{T}_i is equal to the stable category of the Frobenius category \mathcal{C}_i for $1 \leq i \leq r$, and so the decomposition

$$\underline{\text{MCM}}\text{-}\mathcal{O}^\lambda = \bigoplus_{i=1}^r \mathcal{T}_i$$

is in fact a decomposition into triangulated subcategories. If we set $e^{(i)} = \sum_{j \in Q_0^{(i)}} e_j$, [Krause 2015, Proposition 2.3] implies that the functor

$$\underline{\text{Hom}}_{\mathcal{O}^\lambda}(e^{(i)} \Pi^\lambda e_0, -) : \underline{\text{mod}}\text{-}\mathcal{O}^\lambda \rightarrow \text{mod-}\underline{\text{End}}_{\mathcal{O}^\lambda}(e^{(i)} \Pi^\lambda e_0)$$

restricts to a \mathbb{k} -linear equivalence $\mathcal{T}_i \simeq \text{proj-}\Pi(Q^{(i)})$. This equivalence also induces an algebraic triangulated structure on $\text{proj-}\Pi(Q^{(i)})$. Since this category is \mathbb{k} -linearly equivalent to $\mathcal{D}_{\text{sg}}(R_{Q^{(i)}})$, Theorem 4.3 implies that they are triangle equivalent, completing the proof. \square

Remark 4.14. In the first version of this paper, [Crawford 2016], a much longer argument was used to establish Theorem 4.13. The original proof made use of the so-called knitting algorithm from [Iyama and Wemyss 2010] to construct short exact sequences of \mathcal{O}^λ -modules, and explicitly listed many such sequences. The techniques used and the exact sequences given may be of independent interest.

5. A noncommutative geometric McKay correspondence

We now look at a generalisation of the intersection theory of the minimal resolution of a Kleinian singularity $\text{Spec } R_Q$ to a noncommutative setting. We begin by recalling the result in the commutative setting, whereby it is often referred to as the geometric McKay correspondence.

Let R_Q be a Kleinian singularity with corresponding extended Dynkin quiver \tilde{Q} . Then the affine variety $\text{Spec } R_Q$ is an isolated surface singularity which has a unique minimal resolution. It is well-known (see, for example, [Leuschke and Wiegand 2012, Section 6.4]) that the exceptional fibre of this minimal resolution is a union of n irreducible curves γ_i , where n is the number of vertices of Q . Moreover, each γ_i is isomorphic to \mathbb{P}^1 and has self-intersection -2 , and $\gamma_i \cap \gamma_j$ is either empty or a point. In fact, the dual graph of the exceptional fibre is given by the underlying graph of Q . Let Γ be the $n \times n$ matrix with entries

$$\Gamma_{ij} = \begin{cases} -2 & \text{if } i = j, \\ 1 & \text{if } \gamma_i \text{ and } \gamma_j \text{ intersect,} \\ 0 & \text{otherwise,} \end{cases}$$

so that Γ_{ij} is equal to the intersection multiplicity $\gamma_i \bullet \gamma_j$. With an appropriate labelling of the curves γ_i , we have $\Gamma = -C$ where C is the Cartan matrix corresponding to the Dynkin type of Q ; explicitly, $C = 2I - A$, where A is the adjacency matrix of the underlying graph of Q .

Now let \tilde{Q} be an extended Dynkin quiver with $n + 1$ vertices, let Q be the quiver obtained by removing the extending vertex, and let $\lambda = \varepsilon_0 = (1, 0, \dots, 0)$; that is, the weight at the extending vertex is 1, and 0 for all of the other vertices. We may then consider $\mathcal{O}^\lambda(\tilde{Q})$ to be a noncommutative analogue of R_Q , the coordinate ring of the corresponding Kleinian singularity; indeed, these rings have equivalent singularity categories by Theorem 1.1. We now seek to generalise the geometric McKay correspondence to $\mathcal{O}^\lambda(\tilde{Q})$ for this particular choice of λ . To do so, we need an appropriate analogue of a resolution of the singular ring $\mathcal{O}^\lambda(\tilde{Q})$; we use the definition given in [Qin et al. 2019], where such a resolution is a noncommutative ring satisfying certain properties which we will recall below. The role of the exceptional curves in the resolution will be played by finite-dimensional simple modules, of which our resolution of $\mathcal{O}^\lambda(\tilde{Q})$ has finitely many, and the intersection multiplicity of any two such modules is provided by [Mori and Smith 2001]. In this section, we will prove the following:

Theorem 5.1 (Theorem 5.14). *Let \tilde{Q} be an extended Dynkin quiver with corresponding Dynkin quiver Q and let $\lambda = \varepsilon_0$. Then $\mathcal{O}^\lambda(\tilde{Q})$ has a noncommutative resolution (which in fact can take the form $\mathcal{O}^\mu(\tilde{Q})$ for some weight μ and so is a deformation), and the exceptional objects (the finite-dimensional simple modules) in this resolution may be indexed so that the corresponding intersection matrix is $-C$, where C is the Cartan matrix corresponding to Q .*

5A. Noncommutative quasirepant resolutions. We now give a precise definition of the noncommutative resolutions which appear in the above theorem, which is taken from [Qin et al. 2019], and we prove a useful general result. We first recall a definition:

Definition 5.2. Let A be a \mathbb{k} -algebra and $X, Y \in \text{Mod-}A$. We say that X and Y are n -isomorphic, written $X \cong_n Y$, if there exists a third module Z and homomorphisms

$$\phi : Z \rightarrow X \quad \text{and} \quad \psi : Z \rightarrow Y$$

such that the kernels and cokernels of ϕ and ψ each have GK dimension at most n .

We can now give the precise definition of a *noncommutative quasicrepant resolution* from [Qin et al. 2019, Definition 3.15].

Definition 5.3. Let A be a noetherian Auslander–Gorenstein \mathbb{k} -algebra with $\text{GKdim}(A) = d$. Then a *noncommutative quasicrepant resolution* (NQCR) of A is a triple (B, M, N) where B is a noetherian Auslander regular, Cohen–Macaulay \mathbb{k} -algebra with $\text{GKdim}(B) = d$ and where ${}_B M_A$ and ${}_A N_B$ are finitely generated bimodules which are reflexive on both sides and which satisfy

$$M \otimes_A N \cong_{d-2} B \quad \text{and} \quad N \otimes_B M \cong_{d-2} A.$$

NQCRs have the following useful property, which is not proven in [Qin et al. 2019] but provides a useful complement to [loc. cit., Theorem 0.6]. We remark that the following statement remains true if one replaces “noncommutative quasicrepant resolution” by “noncommutative quasiresolution” (see [loc. cit., Definition 3.2]), the latter notion being slightly weaker.

Lemma 5.4. *If B is a noncommutative quasicrepant resolution of A and C is Morita equivalent to B , then C is also a noncommutative quasicrepant resolution of A .*

Proof. Let $d = \text{GKdim } A$. Since B is a NQCR of A , it is Auslander regular and Cohen–Macaulay, has GK dimension d , and there exist bimodules ${}_B M_A$ and ${}_A N_B$ which are finitely generated and reflexive on both sides and which satisfy

$$M \otimes_A N \cong_{d-2} B \quad \text{and} \quad N \otimes_B M \cong_{d-2} A.$$

Moreover, since C is Morita equivalent to B , there exists a progenerator $P \in \text{mod-}B$ with $C \cong \text{End}(P_B)$, and we may view P as a (C, B) -bimodule.

First note that C has GK dimension d by standard Morita theory, and by applying [Yekutieli and Zhang 2002, Proposition 4.3], we deduce that C is also Auslander regular and Cohen–Macaulay. Writing ${}_B Q_C = \text{Hom}({}_C P, C) \cong \text{Hom}({}_B P, B)$, define two bimodules ${}_C \tilde{M}_A$ and ${}_A \tilde{N}_C$ as follows:

$$\tilde{M} = P \otimes_B M \quad \text{and} \quad \tilde{N} = N \otimes_B Q.$$

Since we have pairs of mutually inverse equivalences

$$\begin{aligned} - \otimes_C Q : \text{mod-}B &\rightarrow \text{mod-}C, & - \otimes_B P : \text{mod-}C &\rightarrow \text{mod-}B, \\ P \otimes_B - : B\text{-mod} &\rightarrow C\text{-mod} & Q \otimes_C - : C\text{-mod} &\rightarrow B\text{-mod} \end{aligned}$$

and since M and N are reflexive on both sides, it follows that \tilde{M} and \tilde{N} are reflexive on both sides.

It remains to show that $\tilde{M} \otimes_A \tilde{N} \cong_{d-2} C$ and $\tilde{N} \otimes_C \tilde{M} \cong_{d-2} A$. The latter of these is immediate, since

$$\tilde{N} \otimes_C \tilde{M} = N \otimes_B Q \otimes_C P \otimes_B M \cong N \otimes_B M \cong_{d-2} A,$$

where the first isomorphism follows from [Lam 1999, 18.17 Proposition] and the $(d-2)$ -isomorphism follows since (B, M, N) is a NQCR of A . We now wish to show that $\tilde{M} \otimes_A \tilde{N} \cong_{d-2} C$. Since $M \otimes_A N \cong_{d-2} B$, there exists a bimodule ${}_B Z_B$ and morphisms

$$\phi : Z \rightarrow M \otimes_A N \quad \text{and} \quad \psi : Z \rightarrow B$$

such that the kernels and cokernels of ϕ and ψ have GK dimension at most $d-2$. Define $\tilde{Z} = P \otimes_B Z \otimes_B Q$ and morphisms

$$\begin{aligned} \tilde{\phi} : \tilde{Z} &\rightarrow \tilde{M} \otimes_A \tilde{N}, & \tilde{\phi} &= \text{id}_P \otimes \phi \otimes \text{id}_Q, \\ \tilde{\psi} : \tilde{Z} &\rightarrow C, & \tilde{\psi} &(p \otimes x \otimes q) = q(p\psi(x)). \end{aligned}$$

We claim that the kernels and cokernels of these maps have GK dimension at most $d-2$, which will complete the proof.

We first consider the kernel and cokernel of $\tilde{\phi}$. Since P and Q are projective (hence flat) on the right and left respectively, we can make an identification $\ker \tilde{\phi} \cong P \otimes_B \ker \phi \otimes_B Q$. Since P and Q are finitely generated modules, [McConnell and Robson 2001, Proposition 8.3.14] implies that $\text{GKdim} \ker \tilde{\phi} \leq \text{GKdim} \ker \phi \leq d-2$, as required. Flatness of P and Q also allows us to make identifications

$$\text{im } \tilde{\phi} \cong P \otimes_B \text{im } \phi \otimes_B Q \quad \text{and} \quad \text{coker } \tilde{\phi} \cong P \otimes_B \text{coker } \phi \otimes_B Q,$$

and so $\text{GKdim} \text{coker } \tilde{\phi} \leq \text{GKdim} \text{coker } \phi \leq d-2$.

We now turn our attention to $\tilde{\psi}$. Observe that we can view $\tilde{\psi}$ as the composition

$$\tilde{\psi} : P \otimes_B Z \otimes_B Q \xrightarrow{\text{id}_P \otimes \psi \otimes \text{id}_Q} P \otimes_B B \otimes_B Q \xrightarrow{m \otimes \text{id}_Q} P \otimes_B Q \xrightarrow{\mu} C$$

where $m : P \otimes_B B \rightarrow P$ is the multiplication map and $\mu : P \otimes_B Q \rightarrow C$ is the evaluation map. Flatness of Q implies that $m \otimes \text{id}_Q$ is an isomorphism, while μ is an isomorphism because P is a progenerator. Thus $\ker \tilde{\psi} = \ker(\text{id}_P \otimes \psi \otimes \text{id}_Q)$, and again we can identify this with $P \otimes_B \ker \psi \otimes_B Q$, which has GK dimension at most $d-2$ using the same argument as in the previous paragraph. Similarly, we have an identification $\text{im } \tilde{\psi} \cong P \otimes_B \text{im } \psi \otimes_B Q$, and arguing again as above, we find that $\text{im } \tilde{\psi}$ has GK dimension at most $d-2$, completing the proof. \square

5B. Intersection theory for a family of noncommutative resolutions. We return now to the \mathbb{k} -algebra of interest, namely \mathcal{O}^λ where $\lambda = \varepsilon_0$. Our first aim is to identify an appropriate NQCR, which we have in fact already done.

Lemma 5.5. Π^λ is a NQCR of \mathcal{O}^λ .

Proof. Since \mathcal{O}^λ is Auslander–Gorenstein, Π^λ is Auslander regular and Cohen–Macaulay, and they both have GK dimension 2 [Crawley-Boevey and Holland 1998, Theorem 1.5, Theorem 1.6], it suffices to

show that $\Pi^\lambda e_0 \in \text{bimod}-(\Pi^\lambda, \mathcal{O}^\lambda)$ and $e_0 \Pi^\lambda \in \text{bimod}-(\mathcal{O}^\lambda, \Pi^\lambda)$ are reflexive on both sides, and that

$$\Pi^\lambda e_0 \otimes_{\mathcal{O}^\lambda} e_0 \Pi^\lambda \cong_0 \Pi^\lambda \quad \text{and} \quad e_0 \Pi^\lambda \otimes_{\Pi^\lambda} \Pi^\lambda e_0 \cong_0 \mathcal{O}^\lambda.$$

By Lemma 3.1, $\Pi^\lambda e_0$ is reflexive as a right \mathcal{O}^λ -module, and since it is a generator for $\text{mod-}\mathcal{O}^\lambda$, [Lam 1999, Proposition 18.17] implies it is also reflexive as left Π^λ -module. Similarly, $e_0 \Pi^\lambda$ is a reflexive module on both sides.

The 0-isomorphism $e_0 \Pi^\lambda \otimes_{\Pi^\lambda} \Pi^\lambda e_0 \cong_0 \mathcal{O}^\lambda$ follows from the fact that these two modules are actually isomorphic. To see that $\Pi^\lambda e_0 \otimes_{\mathcal{O}^\lambda} e_0 \Pi^\lambda \cong_0 \Pi^\lambda$, it suffices to show that the multiplication map

$$m : \Pi^\lambda e_0 \otimes_{\mathcal{O}^\lambda} e_0 \Pi^\lambda \rightarrow \Pi^\lambda$$

has finite-dimensional kernel and cokernel. The cokernel of m is $\Pi^\lambda / \Pi^\lambda e_0 \Pi^\lambda \cong \Pi(Q_\lambda) = \Pi(Q)$, which is finite-dimensional. To see that $K = \ker m$ is finite-dimensional, factor m as

$$m : \Pi^\lambda e_0 \otimes_{\mathcal{O}^\lambda} e_0 \Pi^\lambda \xrightarrow{\pi} \Pi^\lambda e_0 \Pi^\lambda \rightarrow \Pi^\lambda$$

where π is surjective and hence $\ker \pi = K$. We then have a short exact sequence

$$0 \rightarrow K \rightarrow \Pi^\lambda e_0 \otimes_{\mathcal{O}^\lambda} e_0 \Pi^\lambda \xrightarrow{\pi} \Pi^\lambda e_0 \Pi^\lambda \rightarrow 0.$$

Since $\Pi^\lambda e_0$ is a finitely generated \mathcal{O}^λ -module, $\Pi^\lambda e_0 \otimes_{\mathcal{O}^\lambda} e_0 \Pi^\lambda$ is a finitely generated Π^λ -module, and so K is also a finitely generated Π^λ -module. Applying $-\otimes_{\Pi^\lambda} \Pi^\lambda e_0$ to the above sequence, which is an exact functor since $\Pi^\lambda e_0$ is a projective left Π^λ -module, we obtain the short exact sequence

$$0 \rightarrow K \otimes_{\Pi^\lambda} \Pi^\lambda e_0 \rightarrow \Pi^\lambda e_0 \otimes_{\mathcal{O}^\lambda} e_0 \Pi^\lambda \otimes_{\Pi^\lambda} \Pi^\lambda e_0 \rightarrow \Pi^\lambda e_0 \Pi^\lambda \otimes_{\Pi^\lambda} \Pi^\lambda e_0 \rightarrow 0.$$

It is easy to see that the above sequence is in fact

$$0 \rightarrow K e_0 \rightarrow \Pi^\lambda e_0 \xrightarrow{\sim} \Pi^\lambda e_0 \rightarrow 0,$$

and so $K e_0 = 0$. It follows that K has the structure of a finitely generated right $\Pi^\lambda / \Pi^\lambda e_0 \Pi^\lambda$ -module, and is therefore finite-dimensional. By definition, we find that Π^λ is a NQCR of \mathcal{O}^λ . \square

We can actually obtain infinitely many noncommutative resolutions of \mathcal{O}^λ using the dual reflections r_i of [Crawley-Boevey and Holland 1998], the definition of which was given prior to Lemma 2.18. It is clear that the r_i preserve the \mathbb{Z}^{n+1} lattice inside \mathbb{k}^{n+1} . It was also noted earlier that $\lambda \cdot \delta = r_i \lambda \cdot \delta$ for all $\lambda \in \mathbb{k}^{\tilde{Q}_0}$ and $i \in \tilde{Q}_0$, so that the r_i preserve the affine hyperplanes $\{\lambda \in \mathbb{k}^{n+1} \mid \lambda \cdot \delta = c\}$ for each $c \in \mathbb{k}$; since $\varepsilon_0 \cdot \delta = 1$, we are primarily interested in the case $c = 1$. Crawley-Boevey and Holland [1998, Corollary 5.2] proved the following useful result:

Lemma 5.6. *Let ρ be a composition of dual reflections. Then Π^λ is Morita equivalent to $\Pi^{\rho(\lambda)}$.*

By combining Lemmas 5.4, 5.5 and 5.6, we obtain the following:

Corollary 5.7. *$\Pi^{\rho(\lambda)}$ is a NQCR of \mathcal{O}^λ for any composition of dual reflections ρ .*

As stated previously, we need to identify an analogue of the exceptional curves appearing in the minimal resolution of a Kleinian singularity. When $\lambda = \varepsilon_0$, by Lemma 2.15, Π^λ has precisely n isoclasses of finite-dimensional simple modules, and hence by Morita equivalence so does $\Pi^{\rho(\lambda)}$ for any composition of dual reflections ρ . These will play the role of the exceptional objects in our noncommutative resolution.

We also require a notion of intersection multiplicity for the exceptional objects, which is provided by [Mori and Smith 2001]. Given a nonsingular noetherian ring S and $M, N \in \text{mod-}S$ which satisfy $\dim_{\mathbb{k}} \text{Ext}_S^\ell(M, N) < \infty$ for all $\ell \geq 0$, we define the *intersection multiplicity* of M and N to be

$$M \bullet N := \sum_{\ell \geq 0} (-1)^{\ell+1} \dim_{\mathbb{k}} \text{Ext}_S^\ell(M, N)$$

(note that this sum has finitely many terms since S is nonsingular).

We are now in a position to prove a preliminary version of Theorem 5.1:

Theorem 5.8. *Let \tilde{Q} be an extended Dynkin quiver with $n+1$ vertices, and let $\lambda = \varepsilon_0$. Let $\mu = \rho(\lambda)$, where ρ is any composition of dual reflections, so that Π^μ is a NQCR of \mathcal{O}^λ . Then Π^μ has precisely n finite-dimensional simple modules S_i up to isomorphism, and with a suitable indexing of them, the intersection matrix Γ with entries $\Gamma_{ij} = S_i \bullet S_j$ is $-C$, where C is the Cartan matrix corresponding to Q .*

Proof. The discussion after Corollary 5.7 shows that Π^μ has n finite-dimensional simple modules S_i up to isomorphism, so it remains to prove the result on the intersection multiplicities.

Since Morita equivalence preserves dimensions of Hom and Ext groups, we are able to calculate the intersection numbers of the finite-dimensional Π^μ -modules by performing the calculations over Π^λ instead. Identifying Π^λ -modules with representations of \tilde{Q} which satisfy the relations coming from Π^λ , [Crawley-Boevey and Holland 1998, Lemma 7.2(6), Theorem 7.4] tells us that the dimension vector of S_i is $\varepsilon_i \in \mathbb{N}^{n+1}$. It follows that

$$S_i \cong \frac{e_i \Pi^\lambda}{\bigoplus_{\substack{\alpha \in \tilde{Q}_1 \\ t(\alpha)=i}} \alpha \Pi^\lambda}. \quad (5.9)$$

Also observe that

$$\text{Hom}_{\Pi^\lambda}(e_i \Pi^\lambda, S_j) = \begin{cases} \mathbb{k} e_i & \text{if } i = j, \\ 0 & \text{if } i \neq j. \end{cases} \quad (5.10)$$

The proof of [Crawley-Boevey and Holland 1998, Lemma 10.1] shows that, for each $i \neq 0$, there is an exact sequence of Π^λ -modules

$$0 \rightarrow e_i \Pi^\lambda \xrightarrow{\phi} \bigoplus_{k \in \partial i} e_k \Pi^\lambda \xrightarrow{\psi} e_i \Pi^\lambda \rightarrow S_i \rightarrow 0.$$

Since the modules $e_k \Pi^\lambda$ are direct summands of Π^λ and hence projective, this is in fact a projective resolution of S_i . Now let $1 \leq j \leq n$. Seeking to calculate the extension groups between S_i and S_j , we apply $\text{Hom}_{\Pi^\lambda}(-, S_j)$ to the corresponding deleted resolution to obtain the complex

$$0 \rightarrow \text{Hom}_{\Pi^\lambda}(e_i \Pi^\lambda, S_j) \rightarrow \bigoplus_{k \in \partial i} \text{Hom}_{\Pi^\lambda}(e_k \Pi^\lambda, S_j) \rightarrow \text{Hom}_{\Pi^\lambda}(e_i \Pi^\lambda, S_j) \rightarrow 0. \quad (5.11)$$

We now consider three distinct cases when computing the homology of this complex. If $j = i$ then, using (5.10), as a complex of vector spaces (5.11) becomes

$$0 \rightarrow \mathbb{k} \rightarrow 0 \rightarrow \mathbb{k} \rightarrow 0$$

and so we can immediately read off that

$$\dim_{\mathbb{k}} \operatorname{Hom}_{\Pi^\lambda}(S_i, S_i) = 1 = \dim_{\mathbb{k}} \operatorname{Ext}_{\Pi^\lambda}^2(S_i, S_i), \quad \dim_{\mathbb{k}} \operatorname{Ext}_{\Pi^\lambda}^\ell(S_i, S_i) = 0 \quad \text{for } \ell = 1 \text{ or } \ell \geq 3,$$

and so $S_i \bullet S_i = -1 + 0 - 1 = -2$. If $j \in \partial i$, then (5.11) becomes

$$0 \rightarrow 0 \rightarrow \mathbb{k} \rightarrow 0 \rightarrow 0$$

and so

$$\dim_{\mathbb{k}} \operatorname{Ext}_{\Pi^\lambda}^1(S_i, S_i) = 1, \quad \dim_{\mathbb{k}} \operatorname{Ext}_{\Pi^\lambda}^\ell(S_i, S_i) = 0 \quad \text{for } \ell = 0 \text{ or } \ell \geq 2.$$

That is, if i and j are adjacent in \tilde{Q} , then

$$S_i \bullet S_j = 0 + 1 + 0 = 1.$$

Finally, if $j \neq i$ and $j \notin \partial i$ then (5.11) becomes

$$0 \rightarrow 0 \rightarrow 0 \rightarrow 0 \rightarrow 0$$

and clearly

$$\dim_{\mathbb{k}} \operatorname{Ext}_{\Pi^\lambda}^\ell(S_i, S_i) = 0 \quad \text{for } \ell \geq 0,$$

and so $S_i \bullet S_j = 0$ in this case. It follows that the intersection matrix Γ satisfies $\Gamma = -C$. \square

The above result should be seen as a noncommutative analogue of the geometric McKay correspondence. However, we can strengthen this result by showing that \mathcal{O}^λ possesses a NQCR which is actually a “deformation”: that is, a NQCR of the form \mathcal{O}^μ for some weight μ . Since we are restricting our attention to quasidominant weights, the fact that \mathcal{O}^μ is nonsingular forces $\mu_i > 0$ for all $i \geq 1$ (see Lemma 2.20). It is not immediately clear that such a deformation exists; we prove its existence in the next subsection.

5C. \mathcal{O}^λ has a NQCR which is a deformation. The dual reflections defined earlier also appear in the so-called *numbers game* of [Mozes 1990]. The relationship between this game and our setting is that the moves considered by Mozes can equivalently be described as an application of a dual reflection to a weight λ . This allows us to make use of some of the results from this paper; in particular we are able to prove that, for $\lambda = \varepsilon_0$, NQCRs of \mathcal{O}^λ which are also deformations exist.

Lemma 5.12. *Let \tilde{Q} be an extended Dynkin quiver with $n + 1$ vertices. Then there exists a sequence of dual reflections ρ such that $\rho(\varepsilon_0)_i > 0$ for all $i \neq 0$; in particular, $\rho(\varepsilon_0)$ is quasidominant.*

Proof. It suffices to show that we can find such a sequence of dual reflections when we work over the field \mathbb{R} , since any such sequence will also have the desired effect on ε_0 when we work over our algebraically closed field \mathbb{k} of characteristic 0. Accordingly, we write $<$ instead of \prec for the total order. Write G for the group

generated by the dual reflections, which is simply the Weyl group of type corresponding to Q . Lemma 5.5 of [Mozes 1990], when translated into our notation, says that $\{\lambda \in \mathbb{R}^{n+1} \mid \lambda_i \geq 0 \text{ for all } 0 \leq i \leq n\}$ is a fundamental domain for the action of G on $\{\lambda \in \mathbb{R}^{n+1} \mid \lambda \cdot \delta > 0\}$. Recalling that G preserves the affine hyperplane $V := \{\lambda \in \mathbb{R}^{n+1} \mid \lambda \cdot \delta = 1\}$, it follows that $V = \bigcup_{\rho \in G} \rho U$, where U is the n -simplex $\{\lambda \in \mathbb{R}^{n+1} \mid \lambda_i \geq 0 \text{ for all } 0 \leq i \leq n \text{ and } \lambda \cdot \delta = 1\}$. Let $H = \{\lambda \in V \mid \lambda_i > 0 \text{ for all } i \neq 0\}$, which is a convex subset of V containing open balls of arbitrarily large diameter. Since each ρU has the same finite diameter, there exists some $\rho \in G$ with $\rho U \subseteq H$. In particular, $\rho(\varepsilon_0) \in H$; that is, $\rho(\varepsilon_0)_i > 0$ for all $i \neq 0$. \square

Remark 5.13. By playing Mozes' numbers game, one can often determine an explicit sequence of dual reflections ρ satisfying the hypotheses of Lemma 5.12. For example, if $\tilde{Q} = \tilde{A}_4$, then the numbers game starting with the initial configuration $(-3, 1, 1, 1, 1)$ terminates at ε_0 , and so by applying the corresponding dual reflections in reverse we obtain the desired ρ . More generally, [Gashi et al. 2012, Proposition 5.1] tells us that when \tilde{Q} is of type \tilde{A}_{2m} , \tilde{D}_{4m} , \tilde{D}_{4m+1} , \tilde{E}_6 or \tilde{E}_8 , where m is a positive integer, then the numbers game starting with the initial configuration $(1 - \sum_{i=1}^n \delta_i, 1, 1, \dots, 1)$ terminates at ε_0 , and so this determines a sequence of dual reflections ρ such that $\rho(\lambda)_i > 0$ for all $i \neq 0$.

We are now in a position to prove Theorem 5.1.

Theorem 5.14. *Let \tilde{Q} be an extended Dynkin quiver with $n + 1$ vertices, and let $\lambda = \varepsilon_0$. Then \mathcal{O}^λ has a NQCR of the form \mathcal{O}^μ , where \mathcal{O}^μ has precisely n finite-dimensional simple modules S_i up to isomorphism. With a suitable indexing of the S_i , the intersection matrix Γ with entries $\Gamma_{ij} = S_i \bullet S_j$ is $-C$, where C is the Cartan matrix corresponding to Q .*

Proof. Lemma 5.12 tells us that there exists a sequence of dual reflections ρ such that \mathcal{O}^μ is nonsingular, where $\mu = \rho(\lambda)$. Since Π^λ is a resolution of \mathcal{O}^λ and there are Morita equivalences between Π^λ , Π^μ , and \mathcal{O}^μ (by [Crawley-Boevey and Holland 1998, Corollary 5.2, Corollary 9.6]), it follows that \mathcal{O}^μ is a NQCR of \mathcal{O}^λ . Finally, these Morita equivalences combined with Theorem 5.8 tells us that \mathcal{O}^μ has precisely n finite-dimensional simple modules S_i up to isomorphism, and since Morita equivalences preserve dimensions of Hom and Ext groups, the claimed intersection multiplicities follow from Theorem 5.8 as well. \square

Appendix: Uniqueness of the translation functor on objects of $\text{proj-}\Pi(Q)$ when Q is Dynkin

In this appendix, we show that if Q is Dynkin and $\text{proj-}\Pi(Q)$ has the structure of a (not necessarily algebraic) triangulated category, then the translation functor Σ is uniquely determined on objects of $\text{proj-}\Pi(Q)$. In particular, this tells us how the translation functor acts on objects in Theorem 4.13. Although this is known from that result, we believe that an elementary and relatively short proof of this fact may be of independent interest.

For the remainder of this section, write P_1, \dots, P_n for the n indecomposable projective right $\Pi(Q)$ -modules corresponding to the vertices of Q . Write W_0, \dots, W_n for the $n + 1$ irreducible representations of the finite group G corresponding to Q . Since $\text{proj-}\Pi(Q)$ is Krull–Schmidt, it is easy to see that $\Sigma P_i = P_j$ for some j , so write σ for the permutation of the vertices of Q satisfying $\Sigma P_i = P_{\sigma(i)}$. The map $W_i \rightarrow W_i^*$

sending a representation to its dual is an involution of $\{W_1, \dots, W_n\}$ (where we intentionally omit W_0), and we can view this map as an automorphism ν of Q . Throughout this section, all Hom spaces are over $\Pi(Q)$, and we omit this subscript. The aim of this section is to prove the following result, which we achieve by analysing cases.

Theorem A.1. *Consider the category $\text{proj-}\Pi(Q)$ with some triangulated structure with translation functor Σ . Then $\sigma = \nu$ as automorphisms of Q .*

Remark A.2. Explicitly, ν is the identity automorphism of Q when Q is \mathbb{A}_1 , \mathbb{D}_n (n even), \mathbb{E}_7 , or \mathbb{E}_8 , and it is the unique graph automorphism of order 2 when Q is \mathbb{A}_n ($n \geq 2$), \mathbb{D}_n (n odd), or \mathbb{E}_6 .

Adapting the proof of Lemma 4.4, σ is necessarily a graph automorphism of Q . Therefore, since the automorphism group of an \mathbb{A}_1 , \mathbb{E}_7 , or \mathbb{E}_8 graph is trivial, it immediately follows that σ is the identity in these cases. For the remaining cases, we argue that Σ is uniquely determined using Lemma 4.4 and by considering the dimensions of the Hom spaces between the P_i . We record the dimensions of these Hom spaces in the following lemma; the \mathbb{E}_7 and \mathbb{E}_8 cases are unnecessary and hence omitted, but they can be established in the same way.

Lemma A.3. *Let Q be a Dynkin quiver with n vertices and let P_1, \dots, P_n be the n indecomposable projective right $\Pi(Q)$ -modules corresponding to the vertices of Q . Let $H(Q)$ be the matrix with*

$$H(Q)_{ij} = \dim_{\mathbb{k}} \text{Hom}(P_j, P_i) = \dim_{\mathbb{k}} e_i \Pi(Q) e_j.$$

(1) *If $Q = \mathbb{A}_n$ then*

$$H(\mathbb{A}_n) = \begin{pmatrix} 1 & 1 & 1 & \cdots & 1 & 1 & 1 \\ 1 & 2 & 2 & \cdots & 2 & 2 & 1 \\ 1 & 2 & 3 & \cdots & 3 & 2 & 1 \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots & \vdots \\ 1 & 2 & 3 & \cdots & 3 & 2 & 1 \\ 1 & 2 & 2 & \cdots & 2 & 2 & 1 \\ 1 & 1 & 1 & \cdots & 1 & 1 & 1 \end{pmatrix}.$$

(2) *If $Q = \mathbb{D}_n$ then*

$$H(\mathbb{D}_n) = \begin{pmatrix} 2 & 2 & 2 & \cdots & 2 & 1 & 1 \\ 2 & 4 & 4 & \cdots & 4 & 2 & 2 \\ 2 & 4 & 6 & \cdots & 6 & 3 & 3 \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots & \vdots \\ 2 & 4 & 6 & \cdots & 2(n-2) & n-2 & n-2 \\ 1 & 2 & 3 & \cdots & n-2 & \lfloor \frac{n-1}{2} \rfloor & \lfloor \frac{n-1}{2} \rfloor \\ 1 & 2 & 3 & \cdots & n-2 & \lfloor \frac{n-1}{2} \rfloor & \lfloor \frac{n-1}{2} \rfloor \end{pmatrix}.$$

(3) If $Q = \mathbb{E}_6$ then

$$H(\mathbb{E}_6) = \begin{pmatrix} 4 & 2 & 4 & 6 & 4 & 2 \\ 2 & 2 & 3 & 4 & 3 & 2 \\ 4 & 3 & 6 & 8 & 6 & 3 \\ 6 & 4 & 8 & 12 & 8 & 4 \\ 4 & 3 & 6 & 8 & 6 & 3 \\ 2 & 2 & 3 & 4 & 3 & 2 \end{pmatrix}.$$

Proof. These can be calculated using [Erdmann and Snashall 1998a, Section 4; 1998b, 3.4; Malkin et al. 2006, Theorem 2.3.b]. \square

We now begin our case-by-case argument. In each case, the technique is the same: seeking a contradiction, we show that if σ is a graph automorphism of Q different from the one given in Theorem A.1 then we arrive at a contradiction. We begin with the type \mathbb{A} case.

Proposition A.4. *Let σ be the graph automorphism of \mathbb{A}_n induced by the translation functor Σ on $\text{proj-}\Pi(\mathbb{A}_n)$. Then σ is the identity when $n = 1$, and it is the unique order 2 graph automorphism when $n \geq 2$.*

Proof. We have already established the $n = 1$ case, so suppose $n \geq 2$. By Lemma 4.4, σ is either the identity or has order 2 so, seeking a contradiction, suppose that σ is the identity; that is $\Sigma P_i = P_i$ for all i . Consider the nonzero morphism $P_1 \rightarrow P_n$ given by left multiplication by $\bar{\alpha}_{n-1}\bar{\alpha}_{n-2} \cdots \bar{\alpha}_1$, which gives rise to a distinguished triangle

$$P_1 \rightarrow P_n \rightarrow M \rightarrow P_1$$

for some $M \in \text{proj-}\Pi(\mathbb{A}_n)$. Applying $\text{Hom}(-, P_n)$, this gives rise to an exact sequence

$$\begin{array}{ccccccc} \text{Hom}(P_n, P_n) & \xrightarrow{\bar{\alpha}_{n-1}\bar{\alpha}_{n-2}\cdots\bar{\alpha}_1} & \text{Hom}(P_1, P_n) & \xrightarrow{\beta} & \text{Hom}(M, P_n) & \xrightarrow{\gamma} & \text{Hom}(P_n, P_n) \xrightarrow{\bar{\alpha}_{n-1}\bar{\alpha}_{n-2}\cdots\bar{\alpha}_1} \text{Hom}(P_1, P_n) \\ \parallel & & \parallel & & \parallel & & \parallel \\ \mathbb{k}e_n & & \mathbb{k}\bar{\alpha}_{n-1}\bar{\alpha}_{n-2}\cdots\bar{\alpha}_1 & & \mathbb{k}e_n & & \mathbb{k}\bar{\alpha}_{n-1}\bar{\alpha}_{n-2}\cdots\bar{\alpha}_1 \end{array}$$

where we use Lemma A.3 to write down bases for each of the Hom spaces. Now the left-hand map is surjective, so exactness implies that β is the zero map, which forces γ to be injective. Moreover, the right-hand map is injective, so that γ is the zero map. In particular, Lemma A.3 implies that we have $\text{Hom}(M, P_n) = 0$ and so $M = 0$, but this tells us that $P_1 \cong P_n$ which is absurd. Therefore σ must be the unique order 2 graph automorphism of \mathbb{A}_n . \square

We now turn our attention to the type \mathbb{E} cases.

Proposition A.5. *Let σ be the graph automorphism of \mathbb{E}_n induced by the translation functor Σ on $\text{proj-}\Pi(\mathbb{E}_n)$, where $n \in \{6, 7, 8\}$. Then σ is the identity when $n \neq 6$, and it is the unique order 2 graph automorphism when $n = 6$.*

Proof. Again, the \mathbb{E}_7 and \mathbb{E}_8 cases are immediate from Lemma 4.4, so consider \mathbb{E}_6 . By Lemma 4.4, σ is either the identity or has order 2 so, seeking a contradiction, suppose that σ is the identity. Consider the nonzero morphism $P_2 \rightarrow P_6$ given by left multiplication by $\alpha_5 \bar{\alpha}_4 \alpha_3 \bar{\alpha}_2$, which gives rise to a distinguished triangle

$$P_2 \rightarrow P_6 \rightarrow M \rightarrow P_2$$

for some $M \in \text{proj-}\Pi(\mathbb{E}_6)$. Applying $\text{Hom}(-, P_6)$, this gives rise to an exact sequence

$$\begin{array}{ccccccc} \text{Hom}(P_6, P_6) & \xrightarrow{\cdot \alpha_5 \bar{\alpha}_4 \alpha_3 \bar{\alpha}_2} & \text{Hom}(P_2, P_6) & \xrightarrow{\beta} & \text{Hom}(M, P_6) & \xrightarrow{\gamma} & \text{Hom}(P_6, P_6) \xrightarrow{\cdot \alpha_5 \bar{\alpha}_4 \alpha_3 \bar{\alpha}_2} \text{Hom}(P_2, P_6) \\ \parallel & & \parallel & & \parallel & & \parallel \\ \mathbb{k}e_6 & & \mathbb{k}\alpha_5 \bar{\alpha}_4 \alpha_3 \bar{\alpha}_2 & & \mathbb{k}e_6 & & \mathbb{k}\alpha_5 \bar{\alpha}_4 \alpha_3 \bar{\alpha}_2 \\ \oplus & & \oplus & & \oplus & & \oplus \\ \mathbb{k}\alpha_5 \bar{\alpha}_4 \alpha_1 \bar{\alpha}_1 \alpha_4 \bar{\alpha}_5 & & \mathbb{k}\alpha_5 \bar{\alpha}_4 \alpha_1 \bar{\alpha}_1 \alpha_4 \bar{\alpha}_5 \alpha_5 \bar{\alpha}_4 \alpha_3 \bar{\alpha}_2 & & \mathbb{k}\alpha_5 \bar{\alpha}_4 \alpha_1 \bar{\alpha}_1 \alpha_4 \bar{\alpha}_5 & & \mathbb{k}\alpha_5 \bar{\alpha}_4 \alpha_1 \bar{\alpha}_1 \alpha_4 \bar{\alpha}_5 \alpha_5 \bar{\alpha}_4 \alpha_3 \bar{\alpha}_2 \end{array}$$

where again we use Lemma A.3 to write down bases for each of the Hom spaces. We see that the left-hand map is surjective and so β is the zero map, and exactness implies that γ is injective. Since the right-hand map is injective it follows that γ is the zero map. Therefore by Lemma A.3 $\text{Hom}(M, P_6) = 0$ and so $M = 0$, but this tells us that $P_2 \cong P_6$ which is absurd. Therefore σ must be the unique order 2 graph automorphism of \mathbb{E}_6 . \square

Finally we consider the type \mathbb{D} cases. Since we claim that Σ behaves differently depending on whether n is odd or even, we have to consider these two cases separately; additionally, we consider the $n = 4$ case separately since $\text{Aut}(\mathbb{D}_4) \cong S_3$ instead of it having order 2.

Proposition A.6. *Let σ be the graph automorphism of \mathbb{D}_4 induced by the translation functor Σ on $\text{proj-}\Pi(\mathbb{D}_4)$. Then σ is the identity.*

Proof. By Lemma 4.4, σ is either the identity, a two-cycle which swaps a pair of vertices $i \neq 2 \neq j$, or it cycles the vertices 1, 3, 4. We rule out the latter two possibilities.

First suppose that σ is a two-cycle: without loss of generality, $\sigma = (3\ 4)$. Consider the nonzero morphism $P_3 \rightarrow P_4$ given by left multiplication by $\bar{\alpha}_4 \alpha_3$, which gives rise to a distinguished triangle

$$P_3 \rightarrow P_4 \rightarrow M \rightarrow P_4$$

for some $M \in \text{proj-}\Pi(\mathbb{D}_4)$. Applying $\text{Hom}(-, P_3)$, this gives rise to an exact sequence

$$\begin{array}{ccccccc} \text{Hom}(P_3, P_3) & \xrightarrow{\cdot \bar{\alpha}_3 \alpha_4} & \text{Hom}(P_4, P_3) & \xrightarrow{\beta} & \text{Hom}(M, P_3) & \xrightarrow{\gamma} & \text{Hom}(P_4, P_3) \xrightarrow{\cdot \bar{\alpha}_4 \alpha_3} \text{Hom}(P_3, P_3) \\ \parallel & & \parallel & & \parallel & & \parallel \\ \mathbb{k}e_3 \oplus \mathbb{k}\bar{\alpha}_3 \alpha_4 \bar{\alpha}_4 \alpha_3 & & \mathbb{k}\bar{\alpha}_3 \alpha_4 & & \mathbb{k}\bar{\alpha}_3 \alpha_4 & & \mathbb{k}e_3 \oplus \mathbb{k}\bar{\alpha}_3 \alpha_4 \bar{\alpha}_4 \alpha_3 \end{array}$$

Clearly the left-hand map surjects, so exactness forces β to be the zero map, which in turn implies that γ is injective. The right-hand map is injective, and exactness forces γ to be the zero map. In particular we have $\text{Hom}(M, P_3) = 0$ and so $M = 0$, but this tells us that $P_3 \cong P_4$ which is absurd. Therefore σ is not a two-cycle.

Now suppose that σ is a three-cycle: without loss of generality, $\sigma = (1\ 3\ 4)$. We now consider the triangle obtained from the morphism $\alpha_1\alpha_3 \cdot : P_3 \rightarrow P_1$,

$$P_3 \rightarrow P_1 \rightarrow M \rightarrow P_4$$

and seek to obtain contradiction. Applying the functor $\text{Hom}(-, P_3)$, we get exactness of the following sequence:

$$\begin{array}{ccccccc} \text{Hom}(P_3, P_3) & \xrightarrow{\cdot \bar{\alpha}_3 \alpha_4} & \text{Hom}(P_4, P_3) & \xrightarrow{\beta} & \text{Hom}(M, P_3) & \xrightarrow{\gamma} & \text{Hom}(P_1, P_3) \xrightarrow{\cdot \alpha_1 \alpha_3} \text{Hom}(P_3, P_3) \\ \parallel & & \parallel & & \parallel & & \parallel \\ \mathbb{k}e_3 \oplus \mathbb{k}\bar{\alpha}_3 \bar{\alpha}_1 \alpha_1 \alpha_3 & & \mathbb{k}\bar{\alpha}_3 \alpha_4 & & \mathbb{k}\bar{\alpha}_3 \bar{\alpha}_1 & & \mathbb{k}e_3 \oplus \mathbb{k}\bar{\alpha}_3 \bar{\alpha}_1 \alpha_1 \alpha_3 \end{array}$$

Again the left-hand map is surjective, forcing β to be the zero map and hence γ to be injective. Moreover, the right-hand map is injective, and so γ must be the zero map. In particular we have $\text{Hom}(M, P_3) = 0$ and so $M = 0$, but this tells us that $P_1 \cong P_3$ which is absurd. Therefore σ is not a three-cycle, and hence must be the identity. \square

Proposition A.7. *Let $n \geq 5$ be odd and let σ be the graph automorphism of \mathbb{D}_n induced by the translation functor Σ on $\text{proj-}\Pi(\mathbb{D}_n)$. Then σ is the unique graph automorphism of order 2.*

Proof. By Lemma 4.4, σ is either the identity or $(n-1\ n)$ so, seeking a contradiction, assume it is the former; that is, $\Sigma P_i = P_i$ for all i . Consider the morphism $P_n \rightarrow P_1$ given by left multiplication by $\alpha_1\alpha_2\alpha_3 \cdots \alpha_{n-3}\alpha_n$. This gives rise to a distinguished triangle

$$P_n \rightarrow P_1 \rightarrow M \rightarrow P_n$$

for some $M \in \text{proj-}\Pi(\mathbb{D}_n)$. Applying $\text{Hom}(-, P_1)$ gives rise to the following exact sequence,

$$\begin{array}{ccccccc} \text{Hom}(P_1, P_1) & \xrightarrow{\cdot \alpha_1 \alpha_2 \cdots \alpha_{n-3} \alpha_n} & \text{Hom}(P_n, P_1) & \xrightarrow{\beta} & \text{Hom}(M, P_1) & \xrightarrow{\gamma} & \text{Hom}(P_1, P_1) \xrightarrow{\cdot \alpha_1 \alpha_2 \cdots \alpha_{n-3} \alpha_n} \text{Hom}(P_n, P_1) \\ \parallel & & \parallel & & \parallel & & \parallel \\ \mathbb{k}e_1 \oplus \mathbb{k}p & & \mathbb{k}\alpha_1 \alpha_2 \cdots \alpha_{n-3} \alpha_n & & \mathbb{k}e_1 \oplus \mathbb{k}p & & \mathbb{k}\alpha_1 \alpha_2 \cdots \alpha_{n-3} \alpha_n \end{array}$$

where here p is some path. The left-hand map is surjective, so β is the zero map and therefore γ is injective. The kernel of the right-hand map is one-dimensional, and exactness tells us that γ has rank 1. In particular we have $\dim \text{Hom}(M, P_1) = 1$ and so M is either P_{n-1} or P_n by Lemma A.3. If we instead apply $\text{Hom}(-, P_n)$, the resulting exact sequence is

$$\begin{array}{ccccccc} \text{Hom}(P_1, P_n) & \xrightarrow{\cdot \alpha_1 \alpha_2 \cdots \alpha_{n-3} \alpha_n} & \text{Hom}(P_n, P_n) & \xrightarrow{\beta} & \text{Hom}(M, P_n) & \xrightarrow{\gamma} & \text{Hom}(P_1, P_n) \xrightarrow{\cdot \alpha_1 \alpha_2 \cdots \alpha_{n-3} \alpha_n} \text{Hom}(P_n, P_n) \\ \parallel & & \parallel & & \parallel & & \parallel \\ \mathbb{k}\bar{\alpha}_n \bar{\alpha}_{n-3} \cdots \bar{\alpha}_2 \bar{\alpha}_1 & & \mathbb{k}^{(n-1)/2} & & \mathbb{k}\bar{\alpha}_n \bar{\alpha}_{n-3} \cdots \bar{\alpha}_2 \bar{\alpha}_1 & & \mathbb{k}^{(n-1)/2} \end{array}$$

Since n is odd, the shortest path from vertex n to vertex 1 and back to vertex n is zero in $\Pi(\mathbb{D}_n)$, so the first and the last maps both have rank zero. Therefore β has full rank, forcing the kernel of γ to have dimension $(n-1)/2$. Exactness also forces γ to have rank 1, and therefore $\dim \operatorname{Hom}(M, P_n) = (n-1)/2 + 1$. Now we have already seen that M is either P_{n-1} or P_n , but $\dim \operatorname{Hom}(P_{n-1}, P_n) = (n-1)/2 = \dim \operatorname{Hom}(P_n, P_n)$, so we have a contradiction. Therefore σ is the unique graph automorphism of order 2. \square

Proposition A.8. *Let $n \geq 6$ be even and let σ be the graph automorphism of \mathbb{D}_n induced by the translation functor Σ on $\operatorname{proj}\text{-}\Pi(\mathbb{D}_n)$. Then σ is the identity.*

Proof. By Lemma 4.4, σ is either the identity or $(n-1) n$ so, seeking a contradiction, assume it is the latter. Consider the morphism $P_n \rightarrow P_1$ given by left multiplication by $\alpha_1 \alpha_2 \cdots \alpha_{n-3} \alpha_n$, and extend this to a distinguished triangle

$$P_n \rightarrow P_1 \rightarrow M \rightarrow P_{n-1}$$

for some $M \in \operatorname{proj}\text{-}\Pi(\mathbb{D}_n)$. If we apply $\operatorname{Hom}(-, P_1)$ we get the following exact sequence

$$\begin{array}{ccccccc} \operatorname{Hom}(P_1, P_1) & \xrightarrow{\cdot \alpha_1 \alpha_2 \cdots \alpha_{n-3} \alpha_n} & \operatorname{Hom}(P_{n-1}, P_1) & \xrightarrow{\beta} & \operatorname{Hom}(M, P_1) & \xrightarrow{\gamma} & \operatorname{Hom}(P_1, P_1) \xrightarrow{\cdot \alpha_1 \alpha_2 \cdots \alpha_{n-3} \alpha_n} \operatorname{Hom}(P_n, P_1) \\ \parallel & & \parallel & & & & \parallel & \parallel \\ \mathbb{k}e_1 \oplus \mathbb{k}p & & \mathbb{k}\alpha_1 \alpha_2 \cdots \alpha_{n-3} \alpha_{n-1} & & & & \mathbb{k}e_1 \oplus \mathbb{k}p & \mathbb{k}\alpha_1 \alpha_2 \cdots \alpha_{n-3} \alpha_n \end{array}$$

where p is some path. The left-hand map surjects, so $\beta = 0$ and therefore γ injects. The right-hand map has a one-dimensional kernel, so γ has rank 1. It follows that $\dim \operatorname{Hom}(M, P_1) = 1$, which implies that M is either P_{n-1} or P_n . If we instead apply $\operatorname{Hom}(-, P_n)$ we get

$$\begin{array}{ccccccc} \operatorname{Hom}(P_1, P_n) & \xrightarrow{\cdot \alpha_1 \alpha_2 \cdots \alpha_{n-3} \alpha_n} & \operatorname{Hom}(P_{n-1}, P_n) & \xrightarrow{\theta} & \operatorname{Hom}(M, P_n) & \xrightarrow{\eta} & \operatorname{Hom}(P_1, P_n) \xrightarrow{\cdot \alpha_1 \alpha_2 \cdots \alpha_{n-3} \alpha_n} \operatorname{Hom}(P_n, P_n) \\ \parallel & & \parallel & & & & \parallel & \parallel \\ \mathbb{k}\bar{\alpha}_n \bar{\alpha}_{n-3} \cdots \bar{\alpha}_2 \bar{\alpha}_1 & & \mathbb{k}^{n/2-1} & & & & \mathbb{k}\bar{\alpha}_n \bar{\alpha}_{n-3} \cdots \bar{\alpha}_2 \bar{\alpha}_1 & \mathbb{k}^{n/2} \end{array}$$

Since n is even, the shortest path from vertex n to vertex 1 and then to vertex $n-1$ is zero in $\Pi(\mathbb{D}_n)$, while the shortest path from vertex n to vertex 1 and back to vertex n is nonzero. It follows that the left-hand map is the zero map, while the right-hand map has rank 1. Therefore θ has full rank which implies that the kernel of η has dimension $n/2 - 1$. Moreover, the right-hand map is injective, so that η has rank 0 and so $\dim \operatorname{Hom}(M, P_n) = n/2 - 1$. Combining this with our earlier restriction on M , this forces $M = P_{n-1}$, and our distinguished triangle is therefore

$$P_n \rightarrow P_1 \rightarrow P_{n-1} \rightarrow P_{n-1}.$$

Since $\operatorname{Hom}(P_1, P_{n-1})$ is spanned by $\bar{\alpha}_{n-1} \bar{\alpha}_{n-3} \cdots \bar{\alpha}_2 \bar{\alpha}_1$ and γ is not the zero map, we can assume that the map $P_1 \rightarrow P_{n-1}$ in this triangle is given by left multiplication by (a scalar multiple of) $\bar{\alpha}_{n-1} \bar{\alpha}_{n-3} \cdots \bar{\alpha}_2 \bar{\alpha}_1$. Moreover, $\operatorname{Hom}(P_{n-1}, P_{n-1}) = \operatorname{span}\{e_{n-1}, p_2, \dots, p_{n/2}\}$ where the p_i are paths of length ≥ 4 . Since θ is not the zero map and the composition $P_1 \rightarrow P_{n-1} \rightarrow P_{n-1}$ must be zero, the map $P_{n-1} \rightarrow P_{n-1}$ in this triangle lies in $\operatorname{span}\{p_2, \dots, p_{n/2}\}$. But then $\theta : \operatorname{Hom}(P_{n-1}, P_n) \rightarrow \operatorname{Hom}(P_{n-1}, P_n)$ maps the longest

path in $\text{Hom}(P_{n-1}, P_n)$ to zero, contradicting the fact that θ has trivial kernel. It follows that σ is not a two-cycle. \square

Acknowledgements

The work contained in this paper was completed while the author was an EPSRC-funded student at the University of Edinburgh, and the material contained in this paper appears in an adapted form in his PhD thesis, [Simon 2018]. The author would like to thank his supervisor Susan J. Sierra for suggesting this problem and providing guidance, Michael Wemyss for a number of useful discussions, and the anonymous referee whose suggestions ultimately led to a much shorter argument than that given in the first version of this paper. The author also thanks the EPSRC.

References

- [Amiot et al. 2015] C. Amiot, O. Iyama, and I. Reiten, “Stable categories of Cohen–Macaulay modules and cluster categories”, *Amer. J. Math.* **137**:3 (2015), 813–857. MR Zbl
- [Arentz-Hansen 2019] E. Arentz-Hansen, “Classifying subcategories in quotients of exact categories”, *Appl. Categ. Structures* **27**:5 (2019), 535–547. MR Zbl
- [Auslander and Bridger 1969] M. Auslander and M. Bridger, *Stable module theory*, Mem. Amer. Math. Soc. **94**, Amer. Math. Soc., Providence, RI, 1969. MR Zbl
- [Bavula 1992] V. V. Bavula, “Generalized Weyl algebras and their representations”, *Algebra i Analiz* **4**:1 (1992), 75–97. In Russian; translated in *St. Petersburg Math. J.* **4**:1 (1993), 71–92. MR Zbl
- [Białkowski et al. 2007] J. Białkowski, K. Erdmann, and A. Skowroński, “Deformed preprojective algebras of generalized Dynkin type”, *Trans. Amer. Math. Soc.* **359**:6 (2007), 2625–2650. MR Zbl
- [Boddington and Levy 2007] P. Boddington and P. Levy, “Deformations of Kleinian singularities of type E_6 , E_7 and E_8 ”, unpublished paper, 2007.
- [Brenner et al. 2002] S. Brenner, M. C. R. Butler, and A. D. King, “Periodic algebras which are almost Koszul”, *Algebr. Represent. Theory* **5**:4 (2002), 331–367. MR Zbl
- [Buchweitz 1986] R.-O. Buchweitz, *Maximal Cohen–Macaulay modules and Tate-cohomology over Gorenstein rings*, Univ. Hannover, 1986.
- [Chen 2011] X.-W. Chen, “The singularity category of an algebra with radical square zero”, *Doc. Math.* **16** (2011), 921–936. MR Zbl
- [Chen 2012] X.-W. Chen, “Three results on Frobenius categories”, *Math. Z.* **270**:1-2 (2012), 43–58. MR Zbl
- [Chen 2018] X.-W. Chen, “The singularity category of a quadratic monomial algebra”, *Q. J. Math.* **69**:3 (2018), 1015–1033. MR Zbl
- [Crawford 2016] S. Crawford, “Singularity categories of deformations of Kleinian singularities”, 2016. To appear in *Algebra Number Theory*. arXiv
- [Crawley-Boevey 1999] W. Crawley-Boevey, “DMV lectures on representations of quivers, preprojective algebras and deformations of quotient singularities”, lecture notes, 1999, Available at <https://www.math.uni-bielefeld.de/~wcrawley/dmvlects.pdf>.
- [Crawley-Boevey and Holland 1998] W. Crawley-Boevey and M. P. Holland, “Noncommutative deformations of Kleinian singularities”, *Duke Math. J.* **92**:3 (1998), 605–635. MR Zbl
- [Eisenbud 1980] D. Eisenbud, “Homological algebra on a complete intersection, with an application to group representations”, *Trans. Amer. Math. Soc.* **260**:1 (1980), 35–64. MR Zbl
- [Erdmann and Snashall 1998a] K. Erdmann and N. Snashall, “On Hochschild cohomology of preprojective algebras, I”, *J. Algebra* **205**:2 (1998), 391–412. MR Zbl

- [Erdmann and Snashall 1998b] K. Erdmann and N. Snashall, “Preprojective algebras of Dynkin type, periodicity and the second Hochschild cohomology”, pp. 183–193 in *Algebras and modules, II* (Geiranger, Norway, 1996), edited by I. Reiten et al., CMS Conf. Proc. **24**, Amer. Math. Soc., Providence, RI, 1998. MR Zbl
- [Gashi et al. 2012] Q. R. Gashi, T. Schedler, and D. E. Speyer, “Looping of the numbers game and the alcoved hypercube”, *J. Combin. Theory Ser. A* **119**:3 (2012), 713–730. MR Zbl
- [Gelfand and Ponomarev 1979] I. M. Gelfand and V. A. Ponomarev, “Model algebras and representations of graphs”, *Funkt. Anal. i Prilozhen.* **13**:3 (1979), 1–12. In Russian; translated in *Funct. Anal. Appl.* **13**:3 (1979) 157–166. MR
- [Happel 1988] D. Happel, *Triangulated categories in the representation theory of finite dimensional algebras*, Lond. Math. Soc. Lect. Note Ser. **119**, Cambridge Univ. Press, 1988. MR Zbl
- [Hodges 1993] T. J. Hodges, “Noncommutative deformations of type-A Kleinian singularities”, *J. Algebra* **161**:2 (1993), 271–290. MR Zbl
- [Iyama and Wemyss 2010] O. Iyama and M. Wemyss, “The classification of special Cohen–Macaulay modules”, *Math. Z.* **265**:1 (2010), 41–83. MR Zbl
- [Iyama and Wemyss 2014] O. Iyama and M. Wemyss, “Singular derived categories of \mathbb{Q} -factorial terminalizations and maximal modification algebras”, *Adv. Math.* **261** (2014), 85–121. MR Zbl
- [Kalck 2015] M. Kalck, “Singularity categories of gentle algebras”, *Bull. Lond. Math. Soc.* **47**:1 (2015), 65–74. MR Zbl
- [Kalck and Yang 2018] M. Kalck and D. Yang, “Relative singularity categories, II: DG models”, preprint, 2018. arXiv
- [Keller 2018] B. Keller, “A remark on a theorem by Claire Amiot”, *C. R. Math. Acad. Sci. Paris* **356**:10 (2018), 984–986. MR Zbl
- [Krause 2015] H. Krause, “Krull–Schmidt categories and projective covers”, *Expo. Math.* **33**:4 (2015), 535–549. MR Zbl
- [Lam 1999] T. Y. Lam, *Lectures on modules and rings*, Grad. Texts Math. **189**, Springer, 1999. MR Zbl
- [Leuschke and Wiegand 2012] G. J. Leuschke and R. Wiegand, *Cohen–Macaulay representations*, Math. Surv. Monogr. **181**, Amer. Math. Soc., Providence, RI, 2012. MR Zbl
- [Malkin et al. 2006] A. Malkin, V. Ostrik, and M. Vybornov, “Quiver varieties and Lusztig’s algebra”, *Adv. Math.* **203**:2 (2006), 514–536. MR Zbl
- [McConnell and Robson 2001] J. C. McConnell and J. C. Robson, *Noncommutative Noetherian rings*, revised ed., Grad. Stud. Math. **30**, Amer. Math. Soc., Providence, RI, 2001. MR Zbl
- [Montgomery and Small 1981] S. Montgomery and L. W. Small, “Fixed rings of Noetherian rings”, *Bull. Lond. Math. Soc.* **13**:1 (1981), 33–38. MR Zbl
- [Mori and Smith 2001] I. Mori and S. P. Smith, “Bézout’s theorem for non-commutative projective spaces”, *J. Pure Appl. Algebra* **157**:2-3 (2001), 279–299. MR Zbl
- [Mozes 1990] S. Mozes, “Reflection processes on graphs and Weyl groups”, *J. Combin. Theory Ser. A* **53**:1 (1990), 128–142. MR Zbl
- [Qin et al. 2019] X.-S. Qin, Y.-H. Wang, and J. J. Zhang, “Noncommutative quasi-resolutions”, *J. Algebra* **536** (2019), 102–148. MR Zbl
- [Rotman 1979] J. J. Rotman, *An introduction to homological algebra*, Pure Appl. Math. **85**, Academic, New York, 1979. MR Zbl
- [Simon 2018] C. Simon, *Singularities of noncommutative surfaces*, Ph.D. thesis, University of Edinburgh, 2018, Available at <http://hdl.handle.net/1842/31543>.
- [Yekutieli and Zhang 2002] A. Yekutieli and J. J. Zhang, “Dualizing complexes and tilting complexes over simple rings”, *J. Algebra* **256**:2 (2002), 556–567. MR Zbl
- [Zaks 1969] A. Zaks, “Injective dimension of semi-primary rings”, *J. Algebra* **13** (1969), 73–86. MR Zbl

Communicated by J. Toby Stafford

Received 2017-02-24 Revised 2019-07-26 Accepted 2019-09-14

simon.crawford@uwaterloo.ca

Department of Pure Mathematics, University of Waterloo, ON, Canada

Algebra & Number Theory

msp.org/ant

EDITORS

MANAGING EDITOR

Bjorn Poonen
Massachusetts Institute of Technology
Cambridge, USA

EDITORIAL BOARD CHAIR

David Eisenbud
University of California
Berkeley, USA

BOARD OF EDITORS

Bhargav Bhatt	University of Michigan, USA	Martin Olsson	University of California, Berkeley, USA
Richard E. Borcherds	University of California, Berkeley, USA	Raman Parimala	Emory University, USA
Antoine Chambert-Loir	Université Paris-Diderot, France	Jonathan Pila	University of Oxford, UK
J.-L. Colliot-Thélène	CNRS, Université Paris-Sud, France	Irena Peeva	Cornell University, USA
Brian D. Conrad	Stanford University, USA	Anand Pillay	University of Notre Dame, USA
Samit Dasgupta	Duke University, USA	Michael Rapoport	Universität Bonn, Germany
Hélène Esnault	Freie Universität Berlin, Germany	Victor Reiner	University of Minnesota, USA
Gavril Farkas	Humboldt Universität zu Berlin, Germany	Peter Sarnak	Princeton University, USA
Hubert Flenner	Ruhr-Universität, Germany	Joseph H. Silverman	Brown University, USA
Sergey Fomin	University of Michigan, USA	Michael Singer	North Carolina State University, USA
Edward Frenkel	University of California, Berkeley, USA	Christopher Skinner	Princeton University, USA
Wee Teck Gan	National University of Singapore	Vasudevan Srinivas	Tata Inst. of Fund. Research, India
Andrew Granville	Université de Montréal, Canada	J. Toby Stafford	University of Michigan, USA
Ben J. Green	University of Oxford, UK	Shunsuke Takagi	University of Tokyo, Japan
Joseph Gubeladze	San Francisco State University, USA	Pham Huu Tiep	University of Arizona, USA
Christopher Hacon	University of Utah, USA	Ravi Vakil	Stanford University, USA
Roger Heath-Brown	Oxford University, UK	Michel van den Bergh	Hasselt University, Belgium
János Kollár	Princeton University, USA	Akshay Venkatesh	Institute for Advanced Study, USA
Philippe Michel	École Polytechnique Fédérale de Lausanne	Marie-France Vignéras	Université Paris VII, France
Susan Montgomery	University of Southern California, USA	Melanie Matchett Wood	University of California, Berkeley, USA
Shigefumi Mori	RIMS, Kyoto University, Japan	Shou-Wu Zhang	Princeton University, USA

PRODUCTION

production@msp.org
Silvio Levy, Scientific Editor


See inside back cover or msp.org/ant for submission instructions.

The subscription price for 2020 is US \$415/year for the electronic version, and \$620/year (+\$60, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to MSP.

Algebra & Number Theory (ISSN 1944-7833 electronic, 1937-0652 printed) at Mathematical Sciences Publishers, 798 Evans Hall #3840, c/o University of California, Berkeley, CA 94720-3840 is published continuously online. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices.

ANT peer review and production are managed by EditFlow® from MSP.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2020 Mathematical Sciences Publishers

Algebra & Number Theory

Volume 14 No. 2 2020

On the definition of quantum Heisenberg category	275
JONATHAN BRUNDAN, ALISTAIR SAVAGE and BEN WEBSTER	
Characteristic cycles and Gevrey series solutions of A -hypergeometric systems	323
CHRISTINE BERKESCH and MARÍA-CRUZ FERNÁNDEZ-FERNÁNDEZ	
Singularity categories of deformations of Kleinian singularities	349
SIMON CRAWFORD	
Iwasawa main conjecture for Rankin–Selberg p -adic L -functions	383
XIN WAN	
Positivity results for spaces of rational curves	485
ROYA BEHESHTI and ERIC RIEDL	
Generalized Schur algebras	501
ALEXANDER KLESHCHEV and ROBERT MUTH	



1937-0652(2020)14:2;1-#