

Algebra & Number Theory

Volume 14
2020
No. 4

A moving lemma for relative 0-cycles

Amalendu Krishna and Jinhyun Park



A moving lemma for relative 0-cycles

Amalendu Krishna and Jinhyun Park

We prove a moving lemma for the additive and ordinary higher Chow groups of relative 0-cycles of regular semilocal k -schemes essentially of finite type over an infinite perfect field. From this, we show that the cycle classes can be represented by cycles that possess certain finiteness, surjectivity, and smoothness properties. It plays a key role in showing that the crystalline cohomology of smooth varieties can be expressed in terms of algebraic cycles.

1. Introduction	991
2. The fs and sfs-cycles	996
3. The sfs-moving lemma in affine spaces	1005
4. The fs-property of residual cycles	1014
5. Regularity of the original cycle over residual points	1025
6. Vertical separation of residual fibers	1033
7. Regularity of residual cycles over finite closed points	1040
8. The main results	1046
Acknowledgments	1053
References	1053

1. Introduction

Just as the classical Chow moving lemma played a fundamental role in studies of Chow groups of smooth algebraic varieties over a field, the moving lemma of Bloch [1986; 1994] played a significant role in studies of higher Chow groups of smooth algebraic varieties, i.e., the motivic cohomology. One limitation of those moving lemmas however is that they focus only on the proper intersection properties of the given cycles. Occasionally, the given circumstances require us to know more about the cycles beyond such proper intersection properties. For instance, we often need to know whether the given cycles are finite over the base scheme, and smooth, or, if not, whether they can be moved to such cycles. Such questions require more subtle treatments and may hold under special circumstances only.

The goal of this article is to prove a moving lemma of this sort for higher relative 0-cycles of a regular semilocal scheme essentially of finite type over an infinite perfect field k . Here, “essentially of finite type” means it is obtained by localizing a quasiprojective k -scheme at a finite set Σ of points. Achieving suitable finiteness and regularity of the cycles is the main characteristic of the moving lemma we seek.

MSC2010: primary 14C25; secondary 14F42, 19E15.

Keywords: algebraic cycles, moving lemma, higher Chow group, additive higher Chow group, linear projection, Grassmannian.

In the introduction, we state the main results, explain the motivation, and give an outline of the article.

1A. The sfs-moving lemma. Let k be an infinite perfect field. Let R be a regular semilocal k -algebra essentially of finite type. Let $V = \text{Spec}(R)$ and let Σ denote the set of closed points of V . Let $\text{Tz}^q(V, \bullet; m)$ be the nondegenerate additive cycle complex of V in codimension $q \geq 1$ and with modulus $m \geq 1$. Let $\text{TCH}^q(V, n; m)$ denote the associated homology groups, called the additive higher Chow groups of V (see Section 2A).

For $n \geq 1$, let $\text{Tz}_{\text{sfs}}^n(V, n; m)$ denote the subgroup of sfs-cycles in $\text{Tz}^n(V, n; m)$ (see Section 2E). Roughly speaking, an sfs-cycle is an element $\alpha \in \text{Tz}^n(V, n; m)$ such that every irreducible component of α intersects $\Sigma \times F$ properly for every face $F \subset \square_k^{n-1}$, is finite and surjective over an irreducible component of V , and the image under every projection $V \times \square_k^{n-1} \rightarrow V \times \square_k^j$ ($0 \leq j \leq n - 1$) is a regular scheme. Those cycles have the trivial boundaries (see Lemma 2.21). Let $\text{TCH}_{\text{sfs}}^n(V, n; m)$ denote the image of the canonical map $\text{Tz}_{\text{sfs}}^n(V, n; m) \rightarrow \text{TCH}^n(V, n; m)$ (see Section 2F). The goal of this article is to prove the following result.

Theorem 1.1 (the sfs-moving lemma). *Let k be an infinite perfect field. Let $m, n \geq 1$ be integers. Let V be a smooth semilocal k -scheme essentially of finite type. Then the canonical map $\text{TCH}_{\text{sfs}}^n(V, n; m) \rightarrow \text{TCH}^n(V, n; m)$ is an isomorphism.*

For the same V as above, let $z^q(V, \bullet)$ denote the cubical version of Bloch’s cycle complex (see [Krishna and Levine 2008, Section 1]) and let $\text{CH}^q(V, n)$ denote the associated higher Chow groups. We can define the subgroup $z_{\text{sfs}}^n(V, n)$ of sfs-cycles and the higher Chow group $\text{CH}_{\text{sfs}}^n(V, n)$ of sfs-cycles analogous to the additive higher Chow group of sfs-cycles. There is a canonical map $\text{CH}_{\text{sfs}}^n(V, n) \rightarrow \text{CH}^n(V, n)$. As a byproduct of the discussions toward the proof of Theorem 1.1, we can recover the following result, stated in [Elbaz-Vincent and Müller-Stach 2002].¹

Theorem 1.2. *Let k be an infinite perfect field. Let $V = \text{Spec}(R)$ be a smooth semilocal k -scheme essentially of finite type. Let $n \geq 1$ be an integer. Then the canonical map $\text{CH}_{\text{sfs}}^n(V, n) \rightarrow \text{CH}^n(V, n)$ is an isomorphism.*

Theorem 1.1 provides the main geometric ground for the proof of the following result and a few of its consequences in the paper [Krishna and Park 2015a], discussed separately due to the huge size and complexities of the proofs of the current article. In particular, it allows one to describe the crystalline cohomology of a smooth scheme in positive characteristic in terms of algebraic cycles.

Theorem 1.3 [Krishna and Park 2015a]. *Let k be any field and let R be a smooth semilocal k -algebra essentially of finite type. Let $m, n \geq 1$ be integers. Then there is a natural isomorphism*

$$\tau_R : \mathbb{W}_m \Omega_R^{n-1} \xrightarrow{\cong} \text{TCH}^n(R, n; m),$$

where $\mathbb{W}_m \Omega_R^\bullet$ is the big de Rham–Witt complex of Hesselholt and Madsen.

¹In [Elbaz-Vincent and Müller-Stach 2002, Lemma 3.11], Theorem 1.2 is claimed for arbitrary fields, but we do not know if this can be achieved using the techniques of linear projections.

When R is a field, this was first proven by Rülling [2007]; the above is a higher dimensional generalization, but it also relies on the theorem of Rülling.

1B. The presentation lemma. We deduce Theorem 1.1 from the following general presentation lemma for residual cycles of linear projections. This has the flavor (hence the name) of Gabber’s geometric presentation lemma (see [Colliot-Thélène et al. 1997]). Of course, our assertions are different and intricate.

Let k be an infinite perfect field. Given a finite map $h : Y' \rightarrow Y$ of k -schemes and a reduced closed subscheme $Z \subset Y'$, let $h^+(Z)$ be the closure of $h^{-1}(h(Z)) \setminus Z$ in Y' with the reduced induced closed subscheme structure. We call this the “residual scheme of Z ” with respect to h .

Let $n \geq 1$ and let $\hat{A}_0, \dots, \hat{A}_{n-1}$ be smooth projective and geometrically integral k -schemes of positive dimensions. For $0 \leq j \leq n - 1$, let $A_j \subset \hat{A}_j$ be a nonempty affine open subset. Set $C_0 := \text{Spec}(k)$ and $C_j := \prod_{i=0}^{j-1} A_i$ for $j \geq 1$. Let $\pi_j : C_n \rightarrow C_j$ be the obvious projection. For any map $f : Y' \rightarrow Y$, let $f_j : Y' \times C_j \rightarrow Y \times C_j$ be the map $f \times \text{id}_{C_j}$.

Let $\bar{X} \subset \mathbb{P}_k^m$ be a reduced closed subscheme of pure dimension $r \geq 1$ and let $X \subset \bar{X}$ be the complement of a hyperplane in \mathbb{P}_k^m such that X is regular and integral. Let $\Sigma \subset X$ be a finite set of closed points. Let $Z \subset X \times C_n$ be an integral closed subscheme of dimension r such that the projection $Z \rightarrow C_n$ is not constant, and the projection $Z \rightarrow X$ is finite and surjective.

The presentation lemma for the residual schemes that we prove is the following.

Theorem 1.4. *Let k be an infinite perfect field. There exist a closed embedding $\bar{X} \hookrightarrow \mathbb{P}_k^N$, a hyperplane $H \subset \mathbb{P}_k^N$ with $X = \bar{X} \setminus H$, and a dense open subset $\mathcal{U} \subset \text{Gr}(N - r - 1, H)$ of the Grassmannian variety such that for each $L \in \mathcal{U}(k)$, the linear projection $\phi_L : \mathbb{P}_k^N \setminus L \rightarrow \mathbb{P}_k^r$ away from L defines a finite surjective morphism $\phi : \bar{X} \rightarrow \mathbb{P}_k^r$ such that the following hold:*

(1) *There exists a Cartesian square:*

$$\begin{array}{ccc} X & \hookrightarrow & \bar{X} \\ \phi \downarrow & & \downarrow \phi \\ \mathbb{A}_k^r & \hookrightarrow & \mathbb{P}_k^r \end{array}$$

- (2) ϕ is étale over an affine open neighborhood of $\phi(\Sigma)$.
- (3) $\phi(x) \neq \phi(x')$ for each pair $x \neq x'$ of points in Σ .
- (4) The map $k(\phi(x)) \rightarrow k(x)$ is an isomorphism for each $x \in \Sigma$.
- (5) The induced map $Z \rightarrow \phi_n(Z)$ is birational.
- (6) The map $\phi_n^+(Z) \rightarrow X$ is finite and surjective.
- (7) $\pi_j(\phi_n^+(Z))$ is regular at all points lying over Σ for each $0 \leq j \leq n$.

1C. Outline of proofs and remarks. We first remark that although V may be in general obtained by localizing a quasiprojective k -scheme at a finite set Σ of not necessarily closed points, for the proof of

the sfs-moving lemma, we can easily reduce to the case of closed points. See Proposition 2.19. Then the proof the sfs-moving lemma can be broadly divided into two parts.

In the first part, we prove it when the underlying semilocal ring is the localization of an affine space \mathbb{A}_k^r at a finite set of closed points. To solve this case, we rely on two key ingredients: the lemma of Bloch [1986, Lemma 1.2] and the moving lemma for cycles with modulus on affine spaces by Kai [2019]. (N.B., Part of what we need in this article from [loc. cit.] is also available in [Krishna and Park 2016].) The moving lemma of Kai allows us to ensure that our cycles can be made to intersect the closed points of the semilocal scheme V properly. After this, we apply an “spread out and specialize” type of argument using [Bloch 1986, Lemma 1.2] to achieve our goal.

Roughly speaking, we argue that we can equip the sfs-property to cycles after moving them via a certain kind of twisted translations by a general set of k -rational points of \mathbb{A}_k^r . This requires us to use that the ground field k is infinite. The rest of the argument is to construct a homotopy between the new and the original cycle. The plain translations by the rational points do not work and the twisted translations make the argument more involved than the classical case. This is done in Section 3.

In the second part, we prove the general case of the sfs-moving lemma by combining the affine space case and the presentation lemma (Theorem 1.4). The proof of the presentation lemma is an intricate application of the method of linear projections and moduli in algebraic geometry.

The reason for this intricacy lies in the fact that it is not sufficient for us to find enough linear projections which give finite and flat morphisms from a projective variety X to projective spaces. We need to invoke a more delicate linear projection in such a way that if we project a subvariety in some smooth family over X to a similar family over the projective space, the resulting residual scheme has certain desired geometric properties, e.g., regularity along a given set of fibers in the family. Even more, we need to ensure that if we project this smooth family over X to a smaller dimensional family via proper maps, then the images of the residual scheme continue to enjoy the good properties.

Showing that one can find enough such linear projections that do the above jobs lies at the heart of the argument. We see that the moduli spaces of linear subspaces that we encounter in the process are all rational, and we find enough rational lines in them. We then reduce the argument to studies of a family of linear subspaces parametrized by a rational line (pencil of linear subspaces). This simplifies the problem.

Along the proofs, we need to separate the cases of algebraically closed and general infinite perfect fields. We first prove the results over algebraically closed fields. Over a general infinite perfect field k , we argue that we can find enough linear subspaces after going to an algebraic closure \bar{k} so that all desired properties are achieved (over \bar{k}) in such a generality that they remain to be satisfied for the original cycle over k after descent. One of these generalities we ensure over \bar{k} is that the whole residual *scheme* is regular, and not just its irreducible components (even if the latter case suffices for the sfs-moving lemma). We then show that there are enough such linear subspaces defined over k . This is achieved using a Galois descent.

Carrying out this program rigorously takes up from Section 4 to Section 7. We combine them to prove the main results in Section 8.

We now make some remarks on our assumption on the ground field. We need k to be infinite to ensure that our moduli spaces have enough k -rational points. We need it to be perfect to achieve the regularity of various residual subvarieties. Although we only need the regularity of cycles, our argument at some stage uses the condition that some regular schemes that we encounter in the middle are actually smooth over k (e.g., see the last part of the proof of Proposition 7.8). The perfectness requirement is evident even in the proof of the sfs-moving lemma in affine space, where we need to use a specialization argument. To make sure that we do not destroy the regularity during the specialization, we need our over-field to be separably generated over k (e.g., see the proof of Lemma 3.11). This requires k to be perfect.

Recall that the moving lemma of Bloch and Chow hold over all fields. One proves this for infinite perfect fields first. The case of finite field reduces to the case of infinite perfect fields using the techniques of pro- ℓ -extensions and the push-pull operators on the Chow groups. However, we cannot use this technique in our case because the smoothness property of the sfs-cycles are not well-behaved under the push-forward operators. However, based on Theorem 1.1, we prove Theorem 1.3 in [Krishna and Park 2015a] over all base fields with different methods.

Finally, the reader may notice that our sfs-moving lemma is stated and proven in this paper for

$$\mathrm{TCH}^n(V, n; m) \quad \text{for } m \geq 1.$$

However, we remark that one does not miss out on anything by this assumption because it is shown in [Krishna and Park 2017, Theorem 1.5] that

$$\mathrm{TCH}^n(V, n; 0) = 0.$$

In particular, $\mathrm{TCH}_{\mathrm{sfs}}^n(V, n; 0) = 0$.

The main result of this article plays essential roles in [Gupta and Krishna 2019b; 2019a; Krishna and Park 2015a]. Apart from these applications, we hope that our presentation lemma through linear projection techniques as well as various results and ideas of manipulating locally closed subsets of the Grassmannian will be useful in the future to anyone in the mathematics community (in particular, those working with algebraic cycles) who uses the linear projection machines in the tool box.

1D. Conventions. Unless we specify otherwise, k is a fixed field. We shall assume later that k is infinite and perfect for our main results. A k -scheme is a separated scheme of finite type over k . An affine k -scheme is a k -scheme which is affine. A k -variety is an equidimensional reduced k -scheme. The product $X \times Y$ means $X \times_k Y$, unless we specify otherwise. We let \mathbf{Sch}_k be the category of k -schemes and \mathbf{Sm}_k of smooth k -schemes. A scheme essentially of finite type is a scheme obtained by localizing at a finite subset of (not necessarily closed) points of a quasiprojective subscheme of a finite type k -scheme. We include the case of not localizing at all. For $\mathcal{C} = \mathbf{Sch}_k, \mathbf{Sm}_k$, we let $\mathcal{C}^{\mathrm{ess}}$ be the extension of the category \mathcal{C} , whose objects are either those in \mathcal{C} or those obtained by localizing an object of \mathcal{C} at a finite subset.

Given $X \in \mathcal{C}^{\mathrm{ess}}$ and a finite set of points $\Sigma \subset X$, we write X_Σ for the localization of X along Σ . If $Y \subset X$ is an inclusion of a reduced locally closed subscheme, then the closure of Y is considered a closed

subscheme of X with the reduced induced structure. The image of a reduced closed subset under a proper map is considered a closed subscheme of the target scheme with the reduced induced structure.

2. The fs and sfs-cycles

After recalling the definition of higher Chow groups and additive higher Chow groups, we define our main objects of study: the fs and sfs-cycles. We prove some preliminary results about these cycles.

2A. Higher Chow groups and additive higher Chow groups. Let k be a field. First recall (see [Bloch 1986]) the definition of higher Chow groups. Let $X \in \mathbf{Sch}_k^{\text{ess}}$ be equidimensional. Let $\mathbb{P}_k^1 = \text{Proj } k[Y_0, Y_1]$, and $\square^n = (\mathbb{P}_k^1 \setminus \{1\})^n$. Let $(y_1, \dots, y_n) \in \square^n$ be the coordinates. A *face* of \square^n is a closed subscheme defined by a set of equations $\{y_{i_1} = \epsilon_1, \dots, y_{i_s} = \epsilon_s\}$, where $\epsilon_j \in \{0, \infty\}$. For $1 \leq i \leq n$ and $\epsilon = 0, \infty$, let $t_i^\epsilon : \square^{n-1} \rightarrow \square^n$ be the inclusion given by $(y_1, \dots, y_{n-1}) \mapsto (y_1, \dots, y_{i-1}, \epsilon, y_i, \dots, y_{n-1})$. Its image gives a codimension 1 face.

Let $q, n \geq 0$. When X is obtained by localizing at a nonclosed point, for closed subschemes in $X \times \square^n$, the notion of dimensions could be ambiguous but the codimensions are well-defined. So, we use dimensions only when there is no ambiguity.

Let $z^q(X, n)$ be the free abelian group on the set of integral closed subschemes of $X \times \square^n$ of codimension q , that intersect properly with $X \times F$ for each face F of \square^n . We define the boundary map $\partial_i^\epsilon(Z) := [(\text{Id}_X \times t_i^\epsilon)^*(Z)]$. This collection of data gives a cubical abelian group $(n \mapsto z^q(X, n))$ in the sense of [Krishna and Levine 2008, Section 1.1], and the groups $z^q(X, n) := z^q(X, n)/z^q(X, n)_{\text{degn}}$ (in the notations of [loc. cit.]) give a complex of abelian groups, whose boundary map at level n is given by $\partial := \sum_{i=1}^n (-1)^i (\partial_i^\infty - \partial_i^0)$. The homology $\text{CH}^q(X, n) := H_n(z^q(X, \bullet), \partial)$ is called the higher Chow group of X .

We recall the definition of additive higher Chow groups from [Krishna and Park 2015b, Section 2] (see also [Park 2009]). Let $X \in \mathbf{Sch}_k^{\text{ess}}$ be equidimensional. Let $\mathbb{A}_k^1 = \text{Spec } k[t]$, $\mathbb{G}_m = \text{Spec } k[t, t^{-1}]$, and $\bar{\square} = \mathbb{P}_k^1$. For $n \geq 1$, let $B_n = \mathbb{A}_k^1 \times \square^{n-1}$, $\bar{B}_n = \mathbb{A}_k^1 \times \bar{\square}^{n-1}$ and $\hat{B}_n = \mathbb{P}_k^1 \times \bar{\square}^{n-1} \supset \bar{B}_n$. Let $(t, y_1, \dots, y_{n-1}) \in \bar{B}_n$ be the coordinates.

On \bar{B}_n , define the Cartier divisors $F_{n,i}^1 := \{y_i = 1\}$ for $1 \leq i \leq n-1$, $F_{n,0} := \{t = 0\}$, and let $F_n^1 := \sum_{i=1}^{n-1} F_{n,i}^1$. A *face* of B_n is a closed subscheme defined by a set of equations of the form $y_{i_1} = \epsilon_1, \dots, y_{i_s} = \epsilon_s$, where $\epsilon_j \in \{0, \infty\}$. For $1 \leq i \leq n-1$ and $\epsilon = 0, \infty$, let $t_{n,i}^\epsilon : B_{n-1} \rightarrow B_n$ be the inclusion $(t, y_1, \dots, y_{n-2}) \mapsto (t, y_1, \dots, y_{i-1}, \epsilon, y_i, \dots, y_{n-2})$. Its image is a codimension 1 face.

The additive higher Chow complex is defined similarly using the spaces B_n instead of \square^n , but together with proper intersections with all faces, we impose additional conditions called the *modulus conditions*, that control how the cycles should behave at “infinity”: (see [Krishna and Park 2015b, Definition 2.1]) let X be a k -scheme, and let V be an integral closed subscheme of $X \times B_n$. Let \bar{V} denote the Zariski closure of V in $X \times \bar{B}_n$ and let $v : \bar{V}^N \rightarrow \bar{V} \subset X \times \bar{B}_n$ be the normalization of \bar{V} . Let $m, n \geq 1$ be integers. We say that V satisfies the *modulus m condition* on $X \times B_n$, if as Weil divisors on \bar{V}^N we have $(m+1)[v^*(F_{n,0})] \leq [v^*(F_n^1)]$. When $n = 1$, we have $F_1^1 = \emptyset$, so it means $v^*(F_{1,0}) = 0$, or $\{t = 0\} \cap \bar{V} = \emptyset$.

If V is a cycle on $X \times B_n$, we say that V satisfies the modulus m condition if each of its irreducible components satisfies the modulus m condition. When m is understood, often we just say that V satisfies the modulus condition. Note that since $F_{n,0} = \{t = 0\} \subset \bar{B}_n$, replacing \bar{B}_n by \hat{B}_n in the definition does not change the nature of the modulus condition on V .

For an equidimensional $X \in \mathbf{Sch}_k^{\text{ess}}$, and integers $m, n, q \geq 1$, we first define $\underline{\text{Tz}}^q(X, 1; m)$ to be the free abelian group on integral closed subschemes Z of $X \times \mathbb{A}^1$ of codimension q , satisfying the modulus condition (see [Krishna and Park 2015b, Definition 2.5]). For $n > 1$, $\underline{\text{Tz}}^q(X, n; m)$ is the free abelian group on integral closed subschemes Z of $X \times B_n$ of codimension q such that for each face F of B_n , Z intersects $X \times F$ properly on $X \times B_n$, and Z satisfies the modulus m condition on $X \times B_n$. For each $1 \leq i \leq n - 1$ and $\epsilon = 0, \infty$, let $\partial_i^\epsilon(Z) := [(\text{Id}_X \times \iota_{n,i}^\epsilon)^*(Z)]$. The proper intersection with faces ensures that $\partial_i^\epsilon(Z)$ are well-defined. The cycles in $\underline{\text{Tz}}^q(X, n; m)$ are called the *admissible cycles* (or, often as *additive higher Chow cycles, or additive cycles*).

This gives the cubical abelian group $(\underline{n} \mapsto \underline{\text{Tz}}^q(X, n + 1; m))$ in the sense of [Krishna and Levine 2008, Section 1.1]. Using the containment lemma [Krishna and Park 2012, Proposition 2.4], that each face $\partial_i^\epsilon(Z)$ lies in $\underline{\text{Tz}}^q(X, n - 1; m)$ is implied from the defining conditions.

For a cycle $\sum_{i=1}^s n_i Z_i$, we let $|\alpha|$ be the closed subscheme $\bigcup_{i=1}^s Z_i$ with its reduced structure. This is called the support of α . If $f : Y \rightarrow X$ is flat and $\alpha \in \underline{\text{Tz}}^q(X, n; m)$, we write $f^*(\alpha)$ often as α_Y . This shorthand is more evident when f is a localization morphism.

Definition 2.1 [Krishna and Park 2015b, Definition 2.6]. Let $X \in \mathbf{Sch}_k^{\text{ess}}$ be equidimensional. The *additive higher Chow complex*, or just the *additive cycle complex*, $\text{Tz}^q(X, \bullet; m)$ of X in codimension q with modulus m is the nondegenerate complex associated to the cubical abelian group $(\underline{n} \mapsto \underline{\text{Tz}}^q(X, n + 1; m))$, i.e., $\text{Tz}^q(X, n; m)$ is the quotient $\underline{\text{Tz}}^q(X, n; m) / \underline{\text{Tz}}^q(X, n; m)_{\text{degn}}$.

The boundary map of this complex at level n is given by $\partial := \sum_{i=1}^{n-1} (-1)^i (\partial_i^\infty - \partial_i^0)$, and it satisfies $\partial^2 = 0$. The homology $\text{TCH}^q(X, n; m) := H_n(\text{Tz}^q(X, \bullet; m))$ for $n \geq 1$ is the *additive higher Chow group* of X with modulus m .

2B. Subcomplexes associated to some algebraic subsets. Let $X \in \mathbf{Sch}_k^{\text{ess}}$ be a variety. Here are some subgroups of $\text{Tz}^q(X, n; m)$ with a finer intersection property with a given finite set \mathcal{W} of locally closed algebraic subsets of X :

Definition 2.2 [Krishna and Park 2012, Definition 4.2]. Define $\underline{\text{Tz}}_{\mathcal{W}}^q(X, n; m)$ to be the subgroup of $\underline{\text{Tz}}^q(X, n; m)$ generated by integral closed subschemes $Z \subset X \times B_n$ that additionally satisfy

$$\text{codim}_{W \times F}(Z \cap (W \times F)) \geq q \text{ for all } W \in \mathcal{W} \text{ and all faces } F \subset B_n. \tag{2-1}$$

The groups $\underline{\text{Tz}}_{\mathcal{W}}^q(X, n + 1; m)$ for $n \geq 0$ form a cubical subgroup of $(\underline{n} \mapsto \underline{\text{Tz}}^q(X, n + 1; m))$ and they give the subcomplex $\text{Tz}_{\mathcal{W}}^q(X, \bullet; m) \subset \text{Tz}^q(X, \bullet; m)$ by modding out by the degenerate cycles. The homology groups are denoted by $\text{TCH}_{\mathcal{W}}^q(X, n; m)$.

2C. Schemes with finite closed points. Recall that (see [Gabber et al. 2013, Section 2.2]) we say a scheme X is an FA-scheme if for any finite subset $\Sigma \subset X$, there exists an *affine* open neighborhood $U \subset X$ of Σ . We have the following [loc. cit.]:

Lemma 2.3. *Any quasiprojective k -scheme is FA. Any open subset of an FA-scheme is FA. Given any finite subset Σ of a quasiprojective k -scheme, and an open subset $U \subset X$ containing Σ , there exists an affine open neighborhood $W \subset U$ of Σ .*

Recall (Section 1D) that a semilocal k -scheme V is essentially of finite type if there is a quasiprojective k -scheme whose localization at a finite subset Σ of points gives V . By Lemma 2.3, we may obtain it by localizing an *affine* k -scheme of finite type.

Definition 2.4. For any semilocal k -scheme V essentially of finite type, a pair (X, Σ) consisting of an affine k -scheme X of finite type and a finite set Σ of points such that $V = \text{Spec}(\mathcal{O}_{X, \Sigma})$, is called an *atlas* for V . A smooth (resp. regular) atlas (X, Σ) is an atlas such that X is smooth over k (resp. regular).

Lemma 2.5. *Let $V = \text{Spec}(R)$ be a semilocal k -scheme obtained by localizing at a finite set Σ of points of a quasiprojective k -variety X . For a cycle α on $V \times B_n$, let $\bar{\alpha}$ be its Zariski closure in $X \times B_n$.*

Then $\alpha \in \text{Tz}_{\Sigma}^q(V, n; m)$ if and only if there exists an affine open neighborhood $U \subset X$ of Σ such that $\bar{\alpha}_U \in \text{Tz}_{\Sigma}^q(U, n; m)$.

Here, if $\partial(\alpha) = 0$, then we can assume that $\partial(\bar{\alpha}_U) = 0$. If α is a boundary, then we can assume $\bar{\alpha}_U$ is also a boundary. If V is smooth over k , then we may take (U, Σ) to be a smooth atlas.

Proof. The first three assertions were proven in [Krishna and Park 2016, Lemmas 4.13 and 4.14]. For the last one, choose any X of finite type using the first assertion. Since V is smooth, we have $X_{\text{sing}} \cap V = \emptyset$ and $X_{\text{sm}} = X \setminus X_{\text{sing}} \supset \Sigma$. By Lemma 2.3, we can choose an affine open $U \subset X_{\text{sm}}$ containing Σ . \square

2D. The fs-cycles. Recall that for higher Chow groups of a semilocal k -scheme V in the Milnor range, [Elbaz-Vincent and Müller-Stach 2002, Lemma 3.11] used the notions called fs-cycles and sfs-cycles. An fs-cycle in [loc. cit.] is a cycle α on $V \times \square_k^n$ such that for each irreducible component Z , the morphism $Z \rightarrow V$ is finite and surjective. However, a moment's thought gives that it is not a good notion. For instance, if V is reducible, then one can almost never achieve the surjection part.

Even if we modify the definition a bit by requiring instead that the support $|\alpha| \rightarrow V$ is finite and surjective, still there is a problem when V is not irreducible: suppose $V = V_1 \cup V_2$ is a disjoint union of irreducible components. Suppose for $i = 1, 2$, we have an irreducible closed subscheme Z_i on $V \times \square_k^n$ such that $Z_i \rightarrow V_i$ is finite surjective. Then $W := Z_1 + Z_2$ and $W' := Z_1 + 2Z_2$ are both fs-cycles in this updated sense. But, then $W' - W = Z_2$ is still finite over V , while it is no longer surjective over V . As a result the set of fs-cycles in the above sense is not even closed under basic summation of cycles, thus they do not form a group.

The natural notion to work with is the following:

Definition 2.6. Let $X, Y \in \mathbf{Sch}_k^{\text{ess}}$. First suppose that Y is irreducible. In this case, we say that a morphism $Y \rightarrow X$ of k -schemes is *fs over X* (or an *fs-morphism*, or simply *fs* when X is understood) if it is finite and it is surjective to an irreducible component of X .

In case Y is not necessarily irreducible, we say $Y \rightarrow X$ is *fs over X* if for each irreducible component $Y_j \subset Y$, the induced map $Y_j \rightarrow X$ is fs over X .

We generalize it further: let $f : Y \rightarrow X$ be a morphism in $\mathbf{Sch}_k^{\text{ess}}$ and let $U \rightarrow X$ be a flat morphism. We say that $Y \rightarrow X$ is *fs over U* , if the fiber product $f' : Y \times_X U \rightarrow U$ is fs.

This notion coincides with the naïve notion mentioned above when X is irreducible. Unlike the naïve notion, this notion of fs-morphisms behaves well under base changes.

Lemma 2.7. *Let $f : Y \rightarrow X$ be an fs morphism in $\mathbf{Sch}_k^{\text{ess}}$. Let $U \rightarrow X$ be a flat morphism in $\mathbf{Sch}_k^{\text{ess}}$. Then the fiber product $f' : Y \times_X U \rightarrow U$ is fs.*

Proof. That the base change of a finite morphism is again finite is apparent. The remaining part on surjectivity over an irreducible component follows by [EGA IV₂ 1965, Proposition (2.3.7)(ii), page 16], where the dominance there is equivalent to surjectivity under finiteness. □

Lemma 2.8. *Let Z be a cycle on $Y \times B$ such that Z is fs over Y in the sense that each irreducible component of Z is fs over Y .*

Let $f : Y \rightarrow X$ be a finite surjective morphism in $\mathbf{Sch}_k^{\text{ess}}$ of irreducible schemes. Then the finite push-forward $f_(Z)$ on $X \times B$ is fs over X .*

Proof. We may assume Z is irreducible. Since $Z \rightarrow Y$ is finite surjective and $Y \rightarrow X$ is finite surjective, the composite $Z \rightarrow Y \rightarrow X$ is finite surjective. □

Here is one simple criterion on finiteness

Lemma 2.9 (finiteness criterion). *Let X be an equidimensional affine k -scheme essentially of finite type. Let \hat{B} be a smooth projective geometrically integral k -scheme of finite type of dimension $n > 0$ and let $B \subset \hat{B}$ be a nonempty affine open subset.*

Let $Z \in z^n(X \times B)$ be an irreducible cycle. Then $Z \rightarrow X$ is fs over X if and only if Z is closed in $X \times \hat{B}$.

Proof. Let $f : Z \hookrightarrow X \times \hat{B} \rightarrow X$ be the composite map. Suppose f is fs over X . Since the second map is projective, by [Hartshorne 1977, Corollary II-4.8(e), Theorem II-4.9, pages 102–103], the first map is a closed immersion. This proves (\Rightarrow).

Conversely, suppose that Z is closed in $X \times \hat{B}$, i.e., the first map is a closed immersion (thus projective). Since the second map is projective, the composite f is projective. Hence, f is a projective morphism of affine schemes, so that it must be finite by [Hartshorne 1977, Exercise II-4.6, page 106]. Moreover, $Z \rightarrow X_i$ being a finite map of irreducible affine schemes of the same dimension, where X_i is the irreducible component that receives Z , this morphism must also be surjective. This proves (\Leftarrow). □

Lemma 2.10. *Let $V = \text{Spec}(R)$ be a semilocal k -scheme essentially of finite type with the set of closed points Σ . Let $B \subset \hat{B}$ be as in Lemma 2.9. Let $F := \hat{B} \setminus B$. Let $Z \in z^n(V \times B)$ be an irreducible cycle and let \hat{Z} be the Zariski closure of Z in $V \times \hat{B}$.*

Suppose that $\hat{Z} \cap (\Sigma \times F) = \emptyset$. Then given any affine atlas (X, Σ) for V , there exists an affine open subatlas (U, Σ) for V such that for the Zariski closure \bar{Z} of Z in $X \times B$, the projection map $\bar{Z}_U \rightarrow U$ is fs over U .

If V is smooth over k from the first place, then we can choose (U, Σ) such that U is smooth over k as well.

Proof. Let (X, Σ) be a given atlas. Let $\hat{\bar{Z}}$ be the Zariski closure of \bar{Z} in $X \times \hat{B}$ and let $\hat{f}: \hat{\bar{Z}} \hookrightarrow X \times \hat{B} \rightarrow X$ be the composition with the projection. Let $Y := \hat{f}(\hat{\bar{Z}} \cap (X \times F))$. Since \hat{f} is projective and since $\hat{\bar{Z}} \cap (\Sigma \times F) = \hat{Z} \cap (\Sigma \times F) = \emptyset$, we see that $Y \subset X$ is a closed subset disjoint from Σ . Hence, $X \setminus Y$ is an open neighborhood of Σ such that $\hat{\bar{Z}} \cap ((X \setminus Y) \times F) = \emptyset$. By Lemma 2.3, we can find an affine open neighborhood U of Σ in $X \setminus Y$, so we have $\hat{\bar{Z}} \cap (U \times F) = \emptyset$. In particular, $\hat{\bar{Z}} \cap (U \times \hat{B}) = \bar{Z} \cap (U \times \hat{B})$. This means \bar{Z}_U is closed in $U \times \hat{B}$. Hence, by Lemma 2.9, the map $\bar{Z}_U \rightarrow U$ is fs over U .

In case V is smooth, then by excising the singular locus of X , which is disjoint from Σ , we may assume that X is smooth. Then the open subset $U \subset X$ is also smooth. \square

Let X be an equidimensional quasiprojective k -scheme and let $\Sigma \subset X$ be a finite set of points. By Lemma 2.3, we may replace X be an affine k -scheme. We have the following two notions of fs-cycles:

Definition 2.11. Let $V = X_\Sigma$. Let $m, n \geq 1$ be integers:

- (1) A cycle $\alpha \in \text{Tz}_\Sigma^n(X, n; m)$ is said to be an fs-cycle along Σ if there is an affine open neighborhood $U \subset X$ of Σ such that each irreducible component of α_U is fs over U . The group of fs-cycles along Σ is denoted by $\text{Tz}_{\Sigma, \text{fs}}^n(X, n; m)$.
- (2) A cycle $\alpha \in \text{Tz}_\Sigma^n(V, n; m)$ is said to be an fs-cycle if each irreducible component of α is fs over V . The group of fs-cycles is denoted by $\text{Tz}_{\text{fs}}^n(V, n; m)$.

These two notions are related as follows:

Corollary 2.12. *Let X be an equidimensional affine k -scheme and let $\Sigma \subset X$ be a finite set of points. Let $V = X_\Sigma$. Let $m, n \geq 1$ be integers. Then a cycle $\alpha \in \text{Tz}_\Sigma^n(X, n; m)$ is an fs-cycle along Σ if and only if $\alpha_V \in \text{Tz}_\Sigma^n(V, n; m)$ is an fs-cycle.*

Proof. (\Rightarrow) Since the localization map $V \rightarrow X$ is flat and it factors through any open neighborhood $U \subset X$ of Σ , one can pull-back by Lemma 2.7 to prove this direction.

(\Leftarrow) By Lemma 2.5, there exists an affine open subatlas (U_1, Σ) of (X, Σ) for V such that the closure $\bar{\alpha}$ of Z in $U_1 \times B_n$ is in $\text{Tz}_\Sigma^n(U_1, n; m)$.

For each irreducible component Z of α , let \hat{Z} be its Zariski closure in $V \times \hat{B}$. Since Z is fs over V , by Lemma 2.9 Z is already closed in $V \times \hat{B}_n$, thus $Z = \hat{Z}$. In particular, $\hat{Z} \cap (\Sigma \times F_n) = \emptyset$. Hence by Lemma 2.10 there exists an affine open subatlas (U_Z, Σ) for V of (U_1, Σ) such that for the Zariski closure

\bar{Z} of Z in $U_1 \times B_n$, the base change $\bar{Z}_{U_Z} \rightarrow U_Z$ is fs. By taking $U := \bigcap_Z U_Z$ where the intersection is taken over all (finitely many) irreducible components of α , we deduce that $\bar{Z}_U \rightarrow U$ is fs. This proves the corollary. \square

We have the following a bit different characterization of the cycles centered around $\text{Tz}_{\text{fs}}^n(V, n; m)$:

Proposition 2.13. *Let $V = \text{Spec}(R)$ be a semilocal k -scheme of geometric type with the set Σ of closed points. Let $m, n \geq 1$. Let $Z \in \text{Tz}_{\Sigma}^n(V, n; m)$ be an irreducible cycle. Then Z is an fs-cycle if and only if there is an atlas (X, Σ) for V such that for the closures \bar{Z} in $X \times B_n$ and \hat{Z} in $V \times \hat{B}_n$, we have $\bar{Z} \in \text{Tz}_{\Sigma}^n(X, n; m)$ and $\hat{Z} \cap (\Sigma \times F_n) = \emptyset$.*

Here, V is smooth over k if and only if we can choose (X, Σ) in the above such that X is smooth over k as well.

Proof. For the first assertion, suppose that Z is an fs-cycle. By Lemma 2.5, there is a affine atlas (X, Σ) for V such that $\bar{Z} \in \text{Tz}_{\Sigma}^n(X, n; m)$. Since $Z \rightarrow V$ is fs over V , by Lemma 2.9, $\hat{Z} \cap (\Sigma \times F_n) = \emptyset$.

Conversely, suppose that for an atlas (X, Σ) and the closure \bar{Z} in $X \times B_n$, we have $\bar{Z} \in \text{Tz}_{\Sigma}^n(X, n; m)$ and $\hat{Z} \cap (\Sigma \times F_n) = \emptyset$. Then, by Lemma 2.10, we may shrink (X, Σ) to an affine open atlas (U, Σ) such that $\bar{Z}_U \rightarrow U$ fs over U . Hence $\bar{Z}_U \in \text{Tz}_{\Sigma, \text{fs}}^n(U, n; m)$. Now by Corollary 2.12, we have $Z \in \text{Tz}_{\text{fs}}^n(V, n; m)$.

For the second assertion, in case V was smooth, then we could have take X to be smooth here by the last assertion of Lemma 2.5. Conversely, a localization of a smooth scheme is smooth again, so that V is smooth over k . \square

2E. The sfs-cycles. For $1 \leq j \leq n$, let $\pi_j : B_n \rightarrow B_j$ and $\hat{\pi}_j : \hat{B}_n \rightarrow \hat{B}_j$ be the projection maps. Let $X \in \mathbf{Sch}_k^{\text{ess}}$ equidimensional. We shall often denote the maps $\text{id}_X \times \pi_j : X \times B_n \rightarrow X \times B_j$ and $\text{id}_X \times \hat{\pi}_j : X \times \hat{B}_n \rightarrow X \times \hat{B}_j$ simply by π_j and $\hat{\pi}_j$, respectively, if the scheme X is fixed in a given context.

For any reduced closed subscheme $Z \subset X \times B_n$ and $1 \leq j \leq n$, let $Z^{(j)} = (\text{id}_X \times \pi_j)(Z)$ be the scheme-theoretic image of Z . Let $Z^{(0)}$ be the scheme-theoretic image of Z in X . Note that if the projection $Z \rightarrow X$ is proper, then $(\text{id}_X \times \pi_j)(Z)$ is closed in $X \times B_j$ and, with its reduced induced closed subscheme structure, coincides with $Z^{(j)}$. The same holds for $Z^{(0)}$. We shall use $Z^{(j)}$ when $Z \rightarrow X$ is in fact finite.

Definition 2.14. Let $X \in \mathbf{Sch}_k^{\text{ess}}$ be smooth over k and let $\Sigma \subset X$ be a finite set of points. Let $m, n \geq 1$ be integers. An integral cycle $[Z] \in \underline{\text{Tz}}^n(X, n; m)$ is called an *sfs-cycle along Σ* , if $[Z] \in \underline{\text{Tz}}_{\Sigma}^n(X, n; m)$, and there exists an affine neighborhood $U \subset X$ of Σ such that the following hold:

- (1) Z_U is finite and surjective over an irreducible component of U , i.e., $Z_U \rightarrow U$ is an fs-morphism.
- (2) The scheme $(Z^{(j)})_U$ is smooth over k for every $0 \leq j \leq n$.

A cycle $\alpha \in \underline{\text{Tz}}^n(X, n; m)$ is called an *sfs-cycle along Σ* if every irreducible component of α is an sfs-cycle along Σ .

Lemma 2.15. *Let X be an equidimensional smooth affine k -scheme and let $\Sigma \subset X$ be a finite set of points. Let $V = X_\Sigma$. Let $m, n \geq 1$ be integers. Then $\alpha \in \underline{\mathrm{Tz}}_\Sigma^n(X, n; m)$ is an sfs-cycle along Σ if and only if $\alpha_V \in \underline{\mathrm{Tz}}_\Sigma^n(V, n; m)$ is an fs-cycle such that $Z^{(j)}$ is smooth over k for each $0 \leq j \leq n$ and for each irreducible component Z of α_V .*

Proof. Under Corollary 2.12, the (\Rightarrow) direction is obvious. We prove (\Leftarrow) . By Corollary 2.12, together with Lemma 2.3, we can find an affine open neighborhood $U' \subset X$ of Σ such that the closure $\alpha_{U'} \in \underline{\mathrm{Tz}}_\Sigma^n(U', n; m)$ is an fs-cycle along Σ . Now let $Y \subset U'$ be the union of the images of the finite maps $(Z_{U'}^{(j)})_{\mathrm{sing}} \rightarrow U'$, where Z runs over all irreducible components of α and $0 \leq j \leq n$. Since $Z_{U'} \rightarrow U'$ is finite for each Z , this $Y \subset U'$ is a closed subset that does not meet Σ . By Lemma 2.3, we can choose an affine open neighborhood $U \subset U' \setminus Y$ of Σ . Then for each component Z of α and each $0 \leq j \leq n$, the scheme $Z_U^{(j)}$ is smooth over k . Note $(Z_U)^{(j)} = (Z^{(j)})_U$ naturally. This shows that α_U is an sfs-cycle along Σ . \square

Another property that sfs-cycles enjoy is the following:

Lemma 2.16. *Let $\phi : X \rightarrow Y$ be an étale morphism of smooth affine k -schemes. Let $\Sigma \subset Y$ be a finite set of points and let $\Sigma' = \phi^{-1}(\Sigma)$. Let $Z \in \underline{\mathrm{Tz}}^n(Y, n; m)$ be an integral sfs-cycle along Σ . Then the flat pull-back $\phi^*(Z) \in \underline{\mathrm{Tz}}^n(X, n; m)$ is an sfs-cycle along Σ' .*

Proof. It is easy to see that $\phi^*(Z) \in \underline{\mathrm{Tz}}_{\Sigma'}^n(X, n; m)$. We now prove the other properties. We can shrink Y and assume that $Z \rightarrow Y$ is finite and surjective, and $Z^{(j)}$ is smooth over k for $0 \leq j \leq n$. Let $W := \phi^*(Z)$. It follows from Lemma 2.7 that W is an fs-cycle along Σ' . To prove that each $W^{(j)}$ is smooth over k , let $W_j := \phi^*(Z^{(j)})$ and consider the commutative diagram

$$\begin{array}{ccccccc}
 W & \twoheadrightarrow & W^{(j)} & \longrightarrow & W_j & \longrightarrow & X \\
 \downarrow & & & & \downarrow & & \downarrow \phi \\
 Z & \twoheadrightarrow & & & Z^{(j)} & \longrightarrow & Y.
 \end{array} \tag{2-2}$$

Here, the map $W^{(j)} \rightarrow W_j$ exists uniquely since the right square is Cartesian. The outer big square is also Cartesian, and this implies that so is the left square. In particular, the vertical arrows are all étale, the horizontal arrows are all finite and surjective and all schemes in (2-2) are reduced. In particular, $W^{(j)} \twoheadrightarrow W_j$. On the other hand, as $W \rightarrow X$ is finite, $W^{(j)} = \pi_j(W)$ is a reduced closed subscheme of W_j . Thus $W^{(j)} = W_j$. Since Z and $Z^{(j)}$ are smooth over k and ϕ is étale, it follows that W and W_j are smooth over k . In particular, $W^{(j)} = W_j$ is smooth over k . This finishes the proof. \square

2F. Additive higher Chow groups of fs and sfs-cycles. The goal of this paper is to prove the “sfs-moving lemma” which will show that the cycle class groups of sfs-cycles coincide with the additive higher Chow groups in the Milnor range for a smooth semilocal k -scheme essentially of finite type when k is an infinite perfect field.

Let $m, n \geq 1$. Let X be a smooth affine k -scheme and let $\Sigma \subset X$ be a finite set of points. It follows from Definition 2.14 that $\mathrm{Tz}_{\Sigma, \mathrm{sfs}}^n(X, n; m)$ is a subgroup of $\mathrm{Tz}_{\Sigma, \mathrm{fs}}^n(X, n; m)$.

Definition 2.17. We let

$$\begin{aligned} \widetilde{\text{TCH}}_{\Sigma}^n(X, n; m) &= \frac{\ker(\partial : \text{Tz}_{\Sigma}^n(X, n; m) \rightarrow \text{Tz}^n(X, n - 1; m))}{\text{im}(\partial : \text{Tz}^n(X, n + 1; m) \rightarrow \text{Tz}^n(X, n; m)) \cap \text{Tz}_{\Sigma}^n(X, n; m)}, \\ \text{TCH}_{\Sigma, \text{fs}}^n(X, n; m) &= \frac{\ker(\partial : \text{Tz}_{\Sigma, \text{fs}}^n(X, n; m) \rightarrow \text{Tz}^n(X, n - 1; m))}{\text{im}(\partial : \text{Tz}^n(X, n + 1; m) \rightarrow \text{Tz}^n(X, n; m)) \cap \text{Tz}_{\Sigma, \text{fs}}^n(X, n; m)}, \\ \text{TCH}_{\Sigma, \text{sfs}}^n(X, n; m) &= \frac{\ker(\partial : \text{Tz}_{\Sigma, \text{sfs}}^n(X, n; m) \rightarrow \text{Tz}^n(X, n - 1; m))}{\text{im}(\partial : \text{Tz}^n(X, n + 1; m) \rightarrow \text{Tz}^n(X, n; m)) \cap \text{Tz}_{\Sigma, \text{sfs}}^n(X, n; m)}. \end{aligned}$$

We similarly define $\widetilde{\text{TCH}}_{\Sigma}^n(V, n; m)$, $\text{TCH}_{\text{fs}}^n(V, n; m)$, and $\text{TCH}_{\text{sfs}}^n(V, n; m)$.

If X is not necessarily connected, note that the groups for X are obtained simply by taking the direct sums of the corresponding groups over all connected components of X .

In the above, the definition of the group $\widetilde{\text{TCH}}_{\Sigma}^n(X, n; m)$ is slightly different from that of $\text{TCH}_{\Sigma}^n(X, n; m)$ in Definition 2.2. However, we have:

Lemma 2.18. *The natural surjection $\text{TCH}_{\Sigma}^n(X, n; m) \twoheadrightarrow \widetilde{\text{TCH}}_{\Sigma}^n(X, n; m)$ is an isomorphism. Similarly, $\text{TCH}_{\Sigma}^n(V, n; m) \twoheadrightarrow \widetilde{\text{TCH}}_{\Sigma}^n(V, n; m)$ is an isomorphism.*

Proof. By the moving lemma for additive higher Chow groups of smooth affine schemes of W. Kai [2019] (see [Krishna and Park 2016, Theorem 4.1] for a sketch of its proof), the composition $\text{TCH}_{\Sigma}^n(X, n; m) \twoheadrightarrow \widetilde{\text{TCH}}_{\Sigma}^n(X, n; m) \rightarrow \text{TCH}^n(X, n; m)$ is an isomorphism. Hence, the first arrow is injective. The proof for the second one is similar, except that we use [Krishna and Park 2016, Theorem 4.10]. \square

We thus have canonical maps

$$\text{TCH}_{\text{sfs}}^n(V, n; m) \rightarrow \text{TCH}_{\text{fs}}^n(V, n; m) \rightarrow \text{TCH}_{\Sigma}^n(V, n; m) \rightarrow \text{TCH}^n(V, n; m), \tag{2-3}$$

where the last map is an isomorphism by Lemma 2.18 and [Kai 2019]. Our goal is to show that all other maps are also isomorphisms.

2G. Reduction to localization at closed points. The semilocal k -schemes essentially of finite type we consider are obtained by localizing an affine k -scheme (see Lemma 2.3) at a finite set Σ of points which may not necessarily be closed. In Section 2G, we show that for the sfs-moving lemma, it is possible to reduce to the case when all points of Σ are actually closed. The following is the goal:

Proposition 2.19. *Suppose the natural map $\text{TCH}_{\text{sfs}}^n(V, n; m) \rightarrow \text{TCH}^n(V, n; m)$ is an isomorphism for every smooth semilocal k -scheme V essentially of finite type, obtained by localizing at a finite set of closed points. Then the natural map $\text{TCH}_{\text{sfs}}^n(V, n; m) \rightarrow \text{TCH}^n(V, n; m)$ is an isomorphism for every smooth semilocal k -scheme V essentially of finite type.*

We prove the following first:

Lemma 2.20. *Let V be a smooth semilocal k -scheme essentially of finite type, obtained by localizing an affine k -scheme X at a finite set Σ of, not necessarily closed, points. Let $\alpha \in \underline{\text{Tz}}^n(V, n; m)$.*

Then there exist (1) a smooth semilocal k -scheme V' essentially of finite type, obtained by localizing an affine k -scheme at a finite set Σ' of closed points with a flat localization map $V \rightarrow V'$ and (2) a cycle $\alpha' \in \underline{\mathrm{Tz}}^n(V', n; m)$ such that the flat pull-back map $\phi_V^{V'} : \underline{\mathrm{Tz}}^n(V', n; m) \rightarrow \underline{\mathrm{Tz}}^n(V, n; m)$ satisfies $\phi_V^{V'}(\alpha') = \alpha$. If $\partial\alpha = 0$, we can ensure $\partial\alpha' = 0$.

Proof. By Lemma 2.3, we may assume that $V = X_\Sigma$, where X is a smooth affine k -scheme of finite type. For the cycle $\alpha \in \underline{\mathrm{Tz}}^n(V, n; m)$, by Lemma 2.5, there exists a smooth affine open neighborhood $U \subset X$ containing Σ such that the Zariski closure α_U of α in $U \times B_n$ is in $\underline{\mathrm{Tz}}^n(U, n; m)$. If $\partial\alpha = 0$, we can shrink U further (if necessary) so that $\partial\alpha_U = 0$.

For each $p \in \Sigma$, there exists a closed point $\mathfrak{m}_p \in U$ that is a specialization of p . (It exists by the basic fact in commutative algebra that any proper ideal of a commutative ring with unit is contained in a maximal ideal.) We choose it so that a distinct pair of points of Σ gives a distinct pair of points. Let $\Sigma' := \{\mathfrak{m}_p \mid p \in \Sigma\}$, and take $V' := U_{\Sigma'}$. Here, $\alpha_U \in \underline{\mathrm{Tz}}^n(U, n; m)$, and let $\alpha' \in \underline{\mathrm{Tz}}^n(V', n; m)$ be its flat pull-back via the localization map $V' \rightarrow U$. This satisfies $\partial\alpha' = 0$ if $\partial\alpha = 0$. By the construction of V' , we also have the localization map $V \rightarrow V'$ and the flat pull-back map $\phi_V^{V'} : \underline{\mathrm{Tz}}^n(V', n; m) \rightarrow \underline{\mathrm{Tz}}^n(V, n; m)$. By the construction of α' , we have $\phi_V^{V'}(\alpha') = \alpha$. This proves the lemma. \square

We remark however that Lemma 2.20 does not say that the map $\phi_V^{V'}$ is surjective. It simply says that for each element α , there is some V' such that α can be an image of a cycle over V' .

Proof of Proposition 2.19. Since the map $\mathrm{TCH}_{\mathrm{sfs}}^n(V, n; m) \rightarrow \mathrm{TCH}^n(V, n; m)$ is automatically injective, it is enough to prove that this is surjective. Let $\alpha \in \mathrm{TCH}^n(V, n; m)$ be an arbitrary cycle class, and choose its cycle representative in $\underline{\mathrm{Tz}}^n(V, n; m)$, also denoted by α . Being a cycle representing a class in $\mathrm{TCH}^n(V, n; m)$, we have $\partial\alpha = 0$.

By Lemma 2.20, there exists now a smooth semilocal k -scheme (V', Σ') essentially of finite type, obtained by localizing at a finite set of closed points, a cycle class $\alpha' \in \mathrm{TCH}^n(V', n; m)$ and the localization map $\phi_V^{V'} : \mathrm{TCH}^n(V', n; m) \rightarrow \mathrm{TCH}^n(V, n; m)$ sends α' to α .

On the other hand, the localization map $\phi_V^{V'}$ sends the sfs-cycles over V' to the sfs-cycles over V . To see this, we first note that this map sends $\underline{\mathrm{Tz}}_{\Sigma'}^n(V', n; m)$ to $\underline{\mathrm{Tz}}_{\Sigma}^n(V, n; m)$ because the localization does not increase the dimensions of schemes, thus the proper intersection condition with Σ' implies the proper intersection condition with Σ . Now, the sfs-cycles are preserved under $\phi_V^{V'}$ because the localization (flat pull-back) of fs-morphisms are fs-morphisms by Lemma 2.7, while it is a basic fact in commutative algebra that a localization of a regular local ring is again a regular local ring. Hence, we have a commutative diagram:

$$\begin{array}{ccc}
 \mathrm{TCH}_{\mathrm{sfs}}^n(V', n; m) & \xrightarrow{\phi_{\mathrm{sfs}}} & \mathrm{TCH}_{\mathrm{sfs}}^n(V, n; m) \\
 \mathrm{sfs}_{V'} \downarrow & & \downarrow \mathrm{sfs}_V \\
 \mathrm{TCH}^n(V', n; m) & \xrightarrow{\phi} & \mathrm{TCH}^n(V, n; m),
 \end{array} \tag{2-4}$$

where $\phi = \phi_V^{V'}$ and ϕ_{sfs} is the restriction of ϕ . By construction, we have $\phi(\alpha') = \alpha$. By the given assumption, we have that $\text{sfs}_{V'}$ is surjective, so that there exists $\alpha'' \in \text{TCH}_{\text{sfs}}^n(V', n; m)$ such that $\text{sfs}_{V'}(\alpha'') = \alpha'$. Hence $\alpha = \phi(\alpha') = \phi \circ \text{sfs}_{V'}(\alpha'') = \dagger \text{sfs}_V \circ \phi_{\text{sfs}}(\alpha'')$, where \dagger holds by the commutativity of the diagram (2-4). In particular, $\alpha \in \text{im}(\text{sfs}_V)$. Since α was arbitrary in $\text{TCH}^n(V, n; m)$, this shows that sfs_V is surjective, hence an isomorphism. \square

We have one further result.

Lemma 2.21. *Let (V, Σ) be a smooth semilocal k -scheme essentially of finite type. Let $m, n \geq 1$ be integers. Let $\alpha \in \underline{\text{Tz}}_{\Sigma}^n(V, n; m)$ be such that $|\alpha|$ is finite over V . Then α does not intersect $V \times F$ for any proper face $F \subset B_n$ at all. In particular, $\partial(\alpha) = 0$.*

Proof. We may assume that $\alpha = [Z]$ is an irreducible cycle and V is integral. We prove that $Z \cap (V \times F)$ is empty.

The composite $Z \cap (V \times F) \hookrightarrow Z \rightarrow V$ is finite by the given assumption. Hence, its image in V is closed and therefore must intersect Σ nontrivially if nonempty. It suffices therefore to show that the fiber product $\Sigma \times_V Z \times_{B_n} F = Z \cap (\Sigma \times F)$ is empty.

However, by the given assumption that $Z \in \underline{\text{Tz}}_{\Sigma}^n(V, n; m)$, the proper intersection condition with Σ reads $\text{codim}_{\Sigma \times F} Z \cap (\Sigma \times F) \geq n$. Equivalently,

$$\dim Z \cap (\Sigma \times F) \leq \dim(\Sigma \times F) - n = \dim F - n < 0.$$

But this means $Z \cap (\Sigma \times F) = \emptyset$. This proves the lemma. \square

Convention. Using Proposition 2.19, from now on, when we say a semilocal k -scheme essentially of finite type, it will mean that it is obtained by localizing at a finite set of closed points, unless we say otherwise.

3. The sfs-moving lemma in affine spaces

In this section, we prove a special case of Theorem 1.1 when the underlying semilocal scheme is a localization an affine space over k . This will be a ground for the general case of the theorem.

3A. The Set-up for affine spaces. We fix some notations that we shall use throughout this section.

Let k be an infinite perfect field. Let $m, n, r \geq 1$ be integers. We let $\Sigma \subset \mathbb{A}_k^r = \text{Spec}(k[x_1, \dots, x_r])$ be a finite set of closed points. Let V be the localization of \mathbb{A}_k^r at Σ . Let $j : V \rightarrow \mathbb{A}_k^r$ be the inclusion map. Let $p_n : \mathbb{A}_k^r \times \mathbb{A}_k^1 \times \bar{\square}^{n-1} \rightarrow \mathbb{A}_k^r \times \mathbb{A}_k^1$ and $q : \mathbb{A}_k^r \times \mathbb{A}_k^1 \rightarrow \mathbb{A}_k^r$ denote the projection maps and let $q_n = q \circ p_n$. Using the automorphism $y \mapsto 1/(1-y)$ of \mathbb{P}_k^1 , we replace $(\square, \infty, 0)$ by $(\mathbb{A}_k^1, 0, 1)$, and write $\square = \mathbb{A}_k^1$.

For any closed subset $Y \subset \mathbb{A}_k^r \times \mathbb{A}_k^1 \times \square^{n-1}$, let \bar{Y} be its closure in $\mathbb{A}_k^r \times \mathbb{A}_k^1 \times \bar{\square}^{n-1}$. We let $Z \in \text{Tz}_{\Sigma}^n(\mathbb{A}_k^r, n; m)$ be an irreducible cycle. For an integer $s \geq 0$ and a point $g \in \mathbb{A}_k^r$, we consider the map (see [Kai 2019])

$$\begin{aligned} \phi_{g,s} : \mathbb{A}_k^r \times \mathbb{A}_k^1 \times \square \times \bar{\square}^{n-1} &\rightarrow \mathbb{A}_k^r \times \mathbb{A}_k^1 \times \bar{\square}^{n-1}; \\ (\underline{x}, t, y, y_1, \dots, y_{n-1}) &\mapsto (\underline{x} + yt^{s(m+1)}g, t, y_1, \dots, y_{n-1}). \end{aligned} \tag{3-1}$$

Note that $\phi_{g,s}$ is strictly speaking defined over the residue field of g , but to simplify notation we often won't make it explicit. If needed, one can take the scalar extension to the residue field of g to turn g into a rational point. For $a \in \square(k)$, we let $\phi_{g,s,a}$ be the composite map

$$\mathbb{A}_k^r \times \mathbb{A}_k^1 \times \bar{\square}^{n-1} \hookrightarrow \mathbb{A}_k^r \times \mathbb{A}_k^1 \times \square \times \bar{\square}^{n-1} \xrightarrow{\phi_{g,s}} \mathbb{A}_k^r \times \mathbb{A}_k^1 \times \bar{\square}^{n-1},$$

where the first arrow takes (\underline{x}, t, y) to (\underline{x}, t, a, y) .

The evaluation of $\phi_{g,s}$ at $y = 1$ defines an isomorphism $\mathbb{A}_{k(g)}^r \times \mathbb{A}_{k(g)}^1 \rightarrow \mathbb{A}_{k(g)}^r \times \mathbb{A}_{k(g)}^1$, given by $\phi_{g,s,1}(\underline{x}, t) = (\underline{x} + t^{s(m+1)}g, t)$. Let $\phi_{g,s,1}^\sharp : k(g)[\underline{x}, t] \rightarrow k(g)[\underline{x}, t]$ be the corresponding $k(g)$ -algebra isomorphism.

3B. Some properties of the twisted translations. Note that $\phi_{g,s}$ is a flat morphism. In particular, $\phi_{g,s}^*(Z)$ is an algebraic cycle on $\mathbb{A}_k^r \times \mathbb{A}_k^1 \times \square^n$. In the next few lemmas, we verify some algebraic and geometric properties of $\phi_{g,s}^*(Z)$.

Lemma 3.1. *Let $f(\underline{x}, t) \in k[\underline{x}, t]$ be a nonzero polynomial. Then there is a nonempty open subset $U \subset \mathbb{A}_k^r$ such that for each $g \in U$ and sufficiently large $s \gg 0$ (not depending on g), the polynomial $\phi_{g,s,1}^\sharp(f)$ is monic in t over $k(g)[\underline{x}]$, i.e., integral over $k(g)[\underline{x}]$.*

Proof. Let $M := \deg_t f$ and write $f(\underline{x}, t) = \sum_{i=0}^M f_i(\underline{x})t^{M-i}$ for some $f_i \in k[\underline{x}]$ and $M \geq 0$. Since $f \neq 0$, we have $f_0(\underline{x}) \neq 0$. Let $d_i = \deg_{\underline{x}}(f_i)$, which is the total degree in \underline{x} . We first consider the case $r = 1$ and take $U = \mathbb{A}_k^1 \setminus \{0\}$. Let $c_i \in k$ be the coefficient of the highest degree term of $f_i(\underline{x})$. Since $f_0(\underline{x}) \neq 0$, we have $c_0 \in k^\times$. Then,

$$f(x + t^{s(m+1)}g, t) = \sum_{i=0}^M f_i(x + t^{s(m+1)}g)t^{M-i} = \sum_{i=0}^M c_i(g^{d_i} t^{d_i s(m+1) + M - i} + (\text{lower degree terms in } t)).$$

Let i_0 be the smallest integer such that $d_{i_0} = \max\{d_0, d_1, \dots, d_M\}$. Here, $c_{i_0} \in k^\times$ by definition.

If $d_{i_0} = 0$, then each $f_i(x)$ is a constant, so $f(x + t^{s(m+1)}g, t)$ gives an integral dependence in t as desired. Suppose $d_{i_0} > 0$. If $i_0 = 0$, then for each $i > 0$ and each $s > 0$, we have

$$d_0 s(m+1) + M \geq d_i s(m+1) + M > d_i s(m+1) + M - i.$$

Hence, the leading coefficient of the highest degree term in t is $c_0 g^{d_0} \in k(g)^\times$, so, after dividing by this unit $c_0 g^{d_0}$, we get a monic polynomial in t . Hence it is integral.

If $i_0 > 0$, then for each $i > i_0$ and each $s > 0$, we have

$$d_{i_0} s(m+1) + M - i_0 \geq d_i s(m+1) + M - i_0 > d_i s(m+1) + M - i,$$

while for $0 \leq i < i_0$, we have $d_i < d_{i_0}$ so that for every sufficiently large $s > 0$, we have

$$d_i s(m+1) + M - i < d_{i_0} s(m+1) + M - i_0.$$

Note that this choice of s depends only on f and not on g . Hence, for every sufficiently large $s > 0$ (not depending on g), again the leading coefficient of highest degree in t is $c_{i_0}g^{d_{i_0}} \in k(g)^\times$. Hence after dividing by this unit, it gives the desired integral dependence relation.

In case $r \geq 2$, the backbone of the proof is the same, but one problem is a possible cancellation of the highest degree terms in t , namely, if d_i is the total degree of $f_i(x_1, \dots, x_r)$, then possibly a multiple number of monomials in $\phi_{g,s,1}^\sharp(f)$ could have the same total degree d_i . However, such g 's form a closed subscheme of \mathbb{A}_k^r (depends on $f(\underline{x}, t)$), so for a general $g \in U$ for some nonempty open subset $U \subset \mathbb{A}_k^r$, we can avoid it. \square

W. Kai [2019, Proposition 2.3] (or see [Krishna and Park 2016, Claim of proof of Theorem 4.1]) defines a positive integer $s(Z)$ associated to Z , which plays a crucial role in proving the modulus condition for $\phi_{g,s}^*(Z)$.

Lemma 3.2. *Let $s \geq s(Z)$ be any integer. Then $\phi_{g,s}^*(Z) \in \text{Tz}^n(\mathbb{A}_k^r, n + 1; m)$ for any $g \in \mathbb{A}_k^r$.*

Proof. The modulus condition for $\phi_{g,s}^*(Z)$ follows from [Kai 2019, Proposition 2.3] (see also [Krishna and Park 2016, Proof of Theorem 4.1]). We show that $\phi_{g,s}^*(Z)$ intersects all faces of \square^n properly. Let F be a face of \square^n . If $F = \{0\} \times F'$ for some face F' of \square^{n-1} , then the proper intersection follows directly from that of Z with F' since the map $\phi_{g,s,0}$ is identity. If $F = \{1\} \times F'$ for some face F' of \square^{n-1} , then the proper intersection also follows from that of Z with F' since the map $\phi_{g,s,1} : \mathbb{A}_k^r \times \mathbb{A}_k^1 \times F' \rightarrow \mathbb{A}_k^r \times \mathbb{A}_k^1 \times F'$ is an isomorphism. If $F = \square \times F'$ for some face F' of \square^{n-1} , then the map $\mathbb{A}_k^r \times \mathbb{A}_k^1 \times \square \times F' \rightarrow \mathbb{A}_k^r \times \mathbb{A}_k^1 \times F'$ is flat of relative dimension one and hence we get

$$\begin{aligned} \dim(\phi_{g,s}^*(Z) \cap (\mathbb{A}_k^r \times \mathbb{A}_k^1 \times \square \times F')) &= \dim(\phi_{g,s}^*(Z \cap F')) \\ &= \dim(Z \cap F') + 1 \leq \dim(Z) + 1 - \text{codim}_{\square^{n-1}}(F') \\ &= \dim(\phi_{g,s}^*(Z)) - \text{codim}_{\square^n}(\square \times F') \\ &= \dim(\phi_{g,s}^*(Z)) - \text{codim}_{\square^n}(F). \end{aligned}$$

This proves the desired proper intersection of $\phi_{g,s}^*(Z)$. \square

Lemma 3.3. *Assume that $n = 1$. For $g \in \mathbb{A}_k^r \setminus \{0\}$ and $s \gg 0$ as in Lemma 3.1, $\phi_{g,s,1}^*(Z)$ is finite and surjective over $\mathbb{A}_{k(g)}^r$.*

Proof. Since $\mathbb{A}_k^r \times \mathbb{A}_k^1$ is factorial, there exists an irreducible polynomial $f(\underline{x}, t) \in k[\underline{x}, t]$ such that $Z = \text{Spec}(k[\underline{x}, t]/(f(\underline{x}, t)))$. The modulus condition mandates that this cycle does not intersect the divisor $\{t = 0\}$ in $\mathbb{A}_k^r \times \mathbb{A}_k^1$, so that after scaling f by a constant in k^\times , we must have $f = th - 1$ for some $h(\underline{x}, t) \in k[\underline{x}, t]$. By Lemma 3.1, $\phi_{g,s,1}^\sharp(th - 1)$ is monic in t for $g \in \mathbb{A}_k^r \setminus \{0\}$ and $s \gg 0$ up to scaling by a unit in $k(g)^\times$. This is equivalent to saying that $\phi_{g,s,1}^*(Z_{k(g)}) \rightarrow \mathbb{A}_{k(g)}^r$ is finite. As both have the same dimension and $\mathbb{A}_{k(g)}^r$ is integral, this morphism is automatically surjective. \square

3C. The three types of cycles. In order to generalize Lemma 3.3 to $n \geq 2$ case, we need to consider three types of cycles.

Lemma 3.4. *Suppose that the projection to the first factor $Z \rightarrow \mathbb{A}_k^r$ is dominant. Then there is a dense open subset $U \subset \mathbb{A}_k^r$ such that each $g \in U$ and integer $s > 0$, the projection to the first factor $\phi_{g,s,1}^*(Z_{k(g)}) \rightarrow \mathbb{A}_{k(g)}^r$ is still dominant.*

Proof. This is immediate from the definition of $\phi_{g,s}$. \square

Lemma 3.5. *Assume that (a) the projection $q_n : Z \rightarrow \mathbb{A}_k^r$ is not dominant while (b) the projection $\text{pr}_2 : Z \rightarrow \mathbb{A}_k^1$ is dominant. Then there is a dense open subset $U \subset \mathbb{A}_k^r$ such that for each $g \in U$ and $s > 0$, we have*

(1) $\dim(q_n(\phi_{g,s,1}^*(Z_{k(g)}))) = \dim(q_n(Z_{k(g)})) + 1$ and

(2) the projection $\text{pr}_2 : \phi_{g,s,1}^*(Z_{k(g)}) \rightarrow \mathbb{A}_{k(g)}^1$ is dominant.

Proof. By (b), the map pr_2 is a dominant morphism to a regular curve, thus it is flat by [Hartshorne 1977, Proposition III-9.7, page 256]. In particular, $\text{pr}_2(Z) \subset \mathbb{A}_k^1$ is a dense open subset. For each $g \in \mathbb{A}_k^r$ and $s > 0$, we have a surjection $\Phi : q_n(Z_{k(g)}) \times \text{pr}_2(Z_{k(g)}) \rightarrow q_n(\phi_{g,s,1}^*(Z_{k(g)}))$, given by sending (x, t) to $x + t^{s(m+1)}g$. Thus, $\dim q_n(\phi_{g,s,1}^*(Z_{k(g)})) \leq \dim q_n(Z_{k(g)}) + 1$.

On the other hand, for each fixed closed point $t_0 \in \text{pr}_2(Z)$, the set $\Phi(q_n(Z_{k(g)}), t_0)$ has the same dimension as that of $q_n(Z_{k(g)})$, while it is an equidimensional proper closed subset of $q_n(\phi_{g,s,1}^*(Z_{k(g)}))$ when g is a general member, i.e., in an open subset of \mathbb{A}_k^r . Since $\text{pr}_2(Z)$ is dense open in \mathbb{A}_k^1 and hence of positive dimension, we must have $\dim(q_n(\phi_{g,s,1}^*(Z_{k(g)}))) > \dim(q_n(Z_{k(g)}))$. This proves (1). Property (2) is obvious because $\phi_{g,s}$ does not modify the \mathbb{A}_k^1 -coordinate. \square

Lemma 3.6. *Assume that neither of the projections $q_n : Z \rightarrow \mathbb{A}_k^r$ nor $\text{pr}_2 : Z \rightarrow \mathbb{A}_k^1$ is dominant. Let $s \geq 1$ be any integer. Then there is a dense open subset $U \subset \mathbb{A}_k^r$ such that for each $g \in U$, there is an open neighborhood $\mathcal{W}_g \subset \mathbb{A}_{k(g)}^r$ of Σ such that $\phi_{g,s,1}^*(Z_{k(g)})$ restricted over \mathcal{W}_g is empty.*

Proof. Since $\text{pr}_2 : Z \rightarrow \mathbb{A}_k^1$ is not dominant and Z is irreducible, $\text{pr}_2(Z)$ must be a singleton closed subset $\{t_0\}$. By the modulus condition that Z satisfies, we must have $t_0 \neq 0$ and $Z \subset \mathbb{A}_k^r \times \{t_0\} \times \square_k^{n-1}$. It is therefore sufficient to prove the lemma by replacing k by $k(t_0)$ and Σ by $\pi_{t_0}^{-1}(\Sigma)$, where $\pi_{t_0} : \text{Spec}(k(t_0)) \rightarrow \text{Spec}(k)$ is the base change. We can thus assume that $t_0 \in k^\times$. Consider the proper closed subset $\overline{q_n(Z)} \subset \mathbb{A}_k^r$ of dimension $< r$ and the dense open complement $\mathcal{U}_0 = \mathbb{A}_k^r \setminus \overline{q_n(Z)}$.

Because Z restricted over \mathcal{U}_0 is empty, we see that the translation $\phi_{g,s,1}^*(Z_{k(g)})$ restricted to the translation $\phi_{g,s,1}^*(\mathcal{U}_0)$ is empty for every $g \in \mathbb{A}_k^r$. Hence, it is enough to show that for an open subset $U \subset \mathbb{A}_k^r$, the set $\mathcal{W}_g := \phi_{g,s,1}^*(\mathcal{U}_0)$ contains Σ for each $g \in U$. However, this is evident because Σ is a finite set of closed point of \mathbb{A}_k^r while \mathcal{U}_0 is a dense open subset of \mathbb{A}_k^r , and $\phi_{g,s,1}^*$ is translation by a nonzero constant factor $(t_0^{s(m+1)})$ of g . This proves the lemma. \square

3D. Key lemmas. The key to our sfs-moving lemma for the localizations of \mathbb{A}_k^r are the following two lemmas.

Let $W \subset \mathbb{A}_k^r \times B_n$ be a reduced closed subscheme and let \overline{W} be its closure in $\mathbb{A}_k^r \times \overline{B}_n$ with reduced closed subscheme structure. We let $\overline{W}^o = \overline{W} \cap (\mathbb{A}_k^r \times \mathbb{G}_{m,k} \times \square^{n-1})$. We fix a closed point $x \in \Sigma$ and integers $m, s \geq 1$.

Define

$$P_1 : \mathbb{A}_k^r \times \mathbb{A}_k^r \times \mathbb{A}_k^1 \times \bar{\square}^{n-1} \rightarrow \mathbb{A}_k^r,$$

$$P_2 : \mathbb{A}_k^r \times \mathbb{A}_k^r \times \mathbb{A}_k^1 \times \bar{\square}^{n-1} \rightarrow \mathbb{A}_k^r \times \mathbb{A}_k^1 \times \bar{\square}^{n-1}$$

to be the projection to the first factor, and the projection to the remaining factors. For a fixed $x \in \mathbb{A}_k^r$, define $\iota_x : \mathbb{A}_k^r \times \mathbb{A}_k^1 \times \bar{\square}^{n-1} \rightarrow \mathbb{A}_k^r \times \mathbb{A}_k^r \times \mathbb{A}_k^1 \times \bar{\square}^{n-1}$ to be the map $(g, t, \underline{y}) \mapsto (g, x + t^{s(m+1)}g, t, \underline{y})$. Let $\theta_x := P_2 \circ \iota_x$ and $\omega_{\bar{W},x} := (P_1 \circ \iota_x)|_{\theta_x^{-1}(\bar{W})}$, where $\theta_x^{-1}(\bar{W})$ is given its reduced induced closed subscheme structure. We then have the commutative diagram

$$\begin{array}{ccccccc}
 \theta_x^{-1}(\bar{W}^o) & \hookrightarrow & \mathbb{A}_k^r \times \mathbb{G}_{m,k} \times \bar{\square}^{n-1} & \xrightarrow{\iota_x} & \mathbb{A}_k^r \times \mathbb{A}_k^r \times \mathbb{G}_{m,k} \times \bar{\square}^{n-1} & \xrightarrow{P_2} & \mathbb{A}_k^r \times \mathbb{G}_{m,k} \times \bar{\square}^{n-1} & \hookleftarrow & \bar{W}^o \\
 & \searrow & \downarrow & & \downarrow & & \downarrow & & \downarrow \\
 \theta_x^{-1}(W) & \hookrightarrow & \theta_x^{-1}(\bar{W}) & \xrightarrow{\iota_x} & \mathbb{A}_k^r \times \mathbb{A}_k^r \times \mathbb{A}_k^1 \times \bar{\square}^{n-1} & \xrightarrow{P_2} & \mathbb{A}_k^r \times \mathbb{A}_k^1 \times \bar{\square}^{n-1} & \hookleftarrow & \bar{W} \\
 & \searrow & \searrow & & \downarrow P_1 & & & & \\
 & & & & \mathbb{A}_k^r & & & & \\
 & & \omega_{\bar{W},x} & & & & & & \\
 & & \omega_{W,x} & & & & & &
 \end{array}$$

(3-2)

where the top row's ι_x, P_2 are the restrictions of the second row, and $\omega_{W,x}$ is the natural composition. The vertical arrows are canonical open immersions. It is easy to check that ι_x is a closed immersion and θ_x is an isomorphism on the top row. Using (3-1) and (3-2), one immediately verifies the following observation which we shall use often.

Lemma 3.7. *Let $x \in \mathbb{A}_k^r$ be fixed. Then for each $g \in \mathbb{A}_k^r$, the map*

$$\omega_{\bar{W},x}^{-1}(g) \rightarrow \phi_{g,s,1}^*(\bar{W}),$$

$(g, t, \underline{y}) \mapsto (x, t, \underline{y})$, is an isomorphism. The same holds for W and \bar{W}^o as well.

Another lemma we shall use is the following.

Lemma 3.8 [Bloch 1986, Lemma 1.2]. *Let X be an algebraic k -scheme and G a connected algebraic k -group acting on X . Let $A, B \subset X$ be closed subsets, and assume the fibers of the map $G \times A \rightarrow X$, $(g, a) \mapsto g \cdot a$ all have the same dimension, and that this map is dominant.*

Moreover, suppose that for an over-field $K \supset k$ and a K -morphism $\psi : X_K \rightarrow G_K$, there is a nonempty open subset $U \subset X$ such that for every $x \in U_K$, a scheme point, we have

$$\text{tr. deg}_k k(\varphi \circ \psi(x), \pi(x)) \geq \dim(G),$$

where $\pi : X_K \rightarrow X_k$ and $\varphi : G_K \rightarrow G_k$ are the projection maps. Define $\phi : X_K \rightarrow X_K$ by $\phi(x) = \psi(x) \cdot x$ and suppose ϕ is an isomorphism. Then the intersection $\phi(A_K \cap U_K) \cap B_K$ is proper.

3E. Applications of the key lemmas. We apply the above two lemmas to our cycle Z and various other closed subsets associated to it. Let $\eta \in \mathbb{A}_k^r$ denote the generic point and let $K := k(\eta)$. We can regard $\eta \in \mathbb{A}_k^r(K)$. Apply Lemma 3.8 with

$$X = \mathbb{A}_k^r \times \mathbb{A}_k^1 \times \overline{\square}_k^{n-1}, \quad G = \mathbb{A}_k^r, \quad \psi(\underline{x}, t, \underline{y}) = (\eta)t^{s(m+1)}, \quad A = \Sigma \times \mathbb{A}_k^1 \times \overline{\square}_k^{n-1}, \quad \text{and} \quad B = \overline{Z},$$

where G acts on $\mathbb{A}_k^r \times \mathbb{A}_k^1 \times \overline{\square}_k^{n-1}$ by $g \cdot (\underline{x}, t, \underline{y}) = (g + \underline{x}, t, \underline{y})$. We let $\phi : X_K \rightarrow X_K$ be given by $\phi(\underline{x}, t, \underline{y}) = ((\eta)t^{s(m+1)} + \underline{x}, t, \underline{y})$. One checks immediately that the conditions of Lemma 3.8 are satisfied and we conclude that $\phi(A_K) \cap \overline{Z}_K$ has dimension at most zero. Comparing this with (3-2) and using Lemma 3.7, this is equivalent to saying that the generic fiber of $\omega_{\overline{Z}, x}$ is finite for every $x \in \Sigma$.

It follows that if Z' is an irreducible component of $\theta_x^{-1}(\overline{Z})$, then either the map $\omega_{\overline{Z}, x} : Z' \rightarrow \mathbb{A}_k^r$ is not dominant or it is dominant and generically quasifinite. In the dominant case, Chevalley's theorem on fiber dimensions (e.g., see [Hartshorne 1977, Exercise II-3.22, page 95]) tells us that we must have $\dim(Z') = r$ and $Z' \rightarrow \mathbb{A}_k^r$ is generically finite. In any case, it follows that there is a dense open subset of \mathbb{A}_k^r over which $Z' \rightarrow \mathbb{A}_k^r$ is quasifinite (with possibly empty fibers).

By taking the finite intersection of such dense open subsets, running over all irreducible components of $\theta_x^{-1}(\overline{Z})$ and all $x \in \Sigma$, we conclude that there is a dense open subset $U \subset \mathbb{A}_k^r$ such that for each $x \in \Sigma$, the map $\omega_{\overline{Z}, x}^{-1}(U) \rightarrow U$ is quasifinite. Using Lemma 3.7, equivalently we get:

Lemma 3.9. *For any integer $s \geq 1$, there is a dense open subset $U \subset \mathbb{A}_k^r$ such that for every $g \in U$, the set $(\Sigma \times \overline{B}_n)_{k(g)} \cap \overline{\phi_{g,s,1}^*(\overline{Z})}_{k(g)} = (\Sigma \times \overline{B}_n)_{k(g)} \cap \overline{\phi_{g,s,1}^*(Z_{k(g)})}$ is finite.*

We can now show the following:

Lemma 3.10. *Let $s \gg 0$ be as in Lemma 3.1. Assume that Z is either dominant over \mathbb{A}_k^r or restricts to zero on V . Then we can find a dense open $U \subset \mathbb{A}_k^r$ such that for $g \in U$, the scheme $\overline{\phi_{g,s,1}^*(Z)}|_V$ is either empty or finite and surjective over V .*

Proof. We can assume $n \geq 2$ by Lemma 3.3. We let $U_1 \subset \mathbb{A}_k^r$ be the intersection of open subsets obtained in Lemmas 3.6 and 3.9. We can therefore assume that $\overline{\phi_{g,s,1}^*(Z_{k(g)})} \rightarrow \mathbb{A}_{k(g)}^r$ is dominant for all $g \in U_1$.

For $g \in U_1$, there is a commutative diagram

$$\begin{array}{ccc} \mathbb{A}_{k(g)}^r \times \mathbb{A}_{k(g)}^1 \times \overline{\square}_{k(g)}^{n-1} & \xrightarrow{p_n} & \mathbb{A}_{k(g)}^r \times \mathbb{A}_{k(g)}^1 \\ \phi_{g,s,1} \downarrow & & \downarrow \phi_{g,s,1} \\ \mathbb{A}_{k(g)}^r \times \mathbb{A}_{k(g)}^1 \times \overline{\square}_{k(g)}^{n-1} & \xrightarrow{p_n} & \mathbb{A}_{k(g)}^r \times \mathbb{A}_{k(g)}^1, \end{array} \quad (3-3)$$

where the horizontal arrows are the projections.

If we let $W = p_n(\overline{Z}_{k(g)})$, it follows from Lemma 3.9 that the composite map $\overline{\phi_{g,s,1}^*(Z_{k(g)})} \rightarrow \overline{\phi_{g,s,1}^*(W)} \rightarrow \mathbb{A}_{k(g)}^r$ is quasifinite over $\Sigma_{k(g)}$. Since $\overline{\phi_{g,s,1}^*(Z_{k(g)})} \rightarrow \mathbb{A}_{k(g)}^r$ is dominant by Lemma 3.4, it follows from Chevalley's theorem on fiber dimensions (see [Hartshorne 1977, Exercise II-3.22, page 95]) that there is an open neighborhood $U_g \subset \mathbb{A}_{k(g)}^r$ of $\Sigma_{k(g)}$ over which the map $\overline{\phi_{g,s,1}^*(Z_{k(g)})} \rightarrow \mathbb{A}_{k(g)}^r$ is

quasifinite with nonempty fibers. We then get maps

$$\overline{\phi_{g,s,1}^*(Z_{k(g)})} \cap q_n^{-1}(U_g) \xrightarrow{p_n} \phi_{g,s,1}^*(W) \cap q^{-1}(U_g) \xrightarrow{q} U_g,$$

where the first map is projective and the composite map is quasifinite with nonempty fibers. This implies that the first map is also quasifinite, and hence, it is finite. Since $\bar{Z} \rightarrow W$ is dominant, so is the map $\overline{\phi_{g,s,1}^*(Z_{k(g)})} \rightarrow \phi_{g,s,1}^*(W)$ by (3-3). It follows that $\overline{\phi_{g,s,1}^*(Z_{k(g)})} \rightarrow \phi_{g,s,1}^*(W)$ is finite and surjective over U_g .

On the other hand, we have shown in Lemma 3.3 that $\phi_{g,s,1}^*(W) \rightarrow \mathbb{A}_{k(g)}^r$ is finite and surjective over \mathbb{A}_k^r for our choice of $s \gg 0$ and $g \in \mathbb{A}_k^r \setminus \{0\}$. We conclude that there is an open neighborhood $U_g \subset \mathbb{A}_{k(g)}^r$ of $\Sigma_{k(g)}$ over which $\phi_{g,s,1}^*(\bar{Z}_{k(g)}) \rightarrow \mathbb{A}_{k(g)}^r$ is finite and surjective.

To show this property for $\phi_{g,s,1}^*(Z_{k(g)})$, we fix $x \in \Sigma$ and use the diagram (3-2) where we take $\bar{W} = Y := \bar{Z} \setminus Z$. To understand the generic fiber of $\omega_{Y,x}$, we apply Lemma 3.8 with

$$X = \mathbb{A}_k^r \times \mathbb{A}_k^1 \times \bar{\square}_k^{n-1}, G = \mathbb{A}_k^r, \psi(x, t, y) = (\eta)t^{s(m+1)}, A = \Sigma \times \mathbb{A}_k^1 \times \bar{\square}_k^{n-1}, B = Y, \quad (3-4)$$

where G acts on $\mathbb{A}_k^r \times \mathbb{A}_k^1 \times \bar{\square}_k^{n-1}$ by $g \cdot (x, t, y) = (g + x, t, y)$ as before. One checks immediately that the conditions of Lemma 3.8 are satisfied. It follows that the intersection $\phi_{\eta,s,1}(A_{k(\eta)}) \cap B_{k(\eta)}$ is proper. By a dimension counting, this means that $\phi_{\eta,s,1}(A_{k(\eta)}) \cap B_{k(\eta)} = \emptyset$. Equivalently, we have $A_{k(\eta)} \cap \phi_{\eta,s,1}^*(Y_{k(\eta)}) = \emptyset$. We conclude by Lemma 3.7 that for every $x \in \Sigma$, the map $\omega_{Y,x} : \theta_x^{-1}(Y) \rightarrow \mathbb{A}_k^r$ is not dominant. We can therefore find a dense open subset $U \subset U_1 \subset \mathbb{A}_k^r$ such that the fiber of $\omega_{Y,x} : \theta_x^{-1}(Y) \rightarrow \mathbb{A}_k^r$ is empty over U for every $x \in \Sigma$. In other words, for every $g \in U$, the intersection $\phi_{g,s,1}^*(Y_{k(g)}) \cap A_{k(g)} = (\overline{\phi_{g,s,1}^*(Z_{k(g)})} \setminus \phi_{g,s,1}^*(Z_{k(g)})) \cap A_{k(g)}$ is empty. But this means that the map $\phi_{g,s,1}^*(Z_{k(g)}) \rightarrow \mathbb{A}_{k(g)}^r$ is finite and surjective over an affine neighborhood of $\Sigma_{k(g)}$ (see Lemma 2.10). \square

Lemma 3.11. *Assume that $Z \in \text{Tz}_{\Sigma}^n(\mathbb{A}_k^r, n; m)$ is an irreducible cycle such that $Z \rightarrow \mathbb{A}_k^r$ is finite and surjective over an affine neighborhood of Σ . We can then find $s \gg 0$ and a dense open subset $U \subset \mathbb{A}_k^r$ such that for each $1 \leq j \leq n$ and for each $g \in U$, the scheme $(\phi_{g,s,1}^*(Z_{k(g)}))^{(j)}$ is regular over an affine neighborhood of $\Sigma_{k(g)}$.*

Proof. We take $W = Z_{\text{sing}}$, the singular locus of Z , in (3-2) and consider the map $\omega_{Z_{\text{sing}},x} : \theta_x^{-1}(Z_{\text{sing}}) \rightarrow \mathbb{A}_k^r$ for $x \in \Sigma$. We had seen previously that the map θ_x on the top row of (3-2) is an isomorphism. In particular, the map $\theta_x : \theta_x^{-1}(Z_{\text{sing}}) \rightarrow Z_{\text{sing}}$ is an isomorphism. But this implies that $\dim(\theta_x^{-1}(Z_{\text{sing}})) = \dim(Z_{\text{sing}}) \leq r - 1$. It follows that the map $\omega_{Z_{\text{sing}},x} : \theta_x^{-1}(Z_{\text{sing}}) \rightarrow \mathbb{A}_k^r$ is not dominant. We can therefore find a dense open subset $U \subset \mathbb{A}_k^r$ such that the fibers of ω_x over U are empty. By shrinking U further, we can assume that this holds for all $x \in \Sigma$.

It follows from Lemma 3.7 that for every $g \in U$, the closed subscheme $(\phi_{g,s,1}^*(Z_{k(g)}))_{\text{sing}} = \phi_{g,s,1}^*((Z_{k(g)})_{\text{sing}}) = \phi_{g,s,1}^*((Z_{\text{sing}})_{k(g)})$ does not meet $(\Sigma \times B_n)_{k(g)}$. Here, the last equality uses the perfectness of k . But this means that $\phi_{g,s,1}^*(Z_{k(g)})$ is regular at all points lying over $\Sigma_{k(g)}$. By choosing $s \gg 0$ as in Lemma 3.1, shrinking U further, and using Lemma 3.10, we can assume that $\phi_{g,s,1}^*(Z_{k(g)})$ is

finite and surjective over an affine neighborhood of $\Sigma_{k(g)}$. But then $\phi_{g,s,1}^*(Z_{k(g)})$ must be regular over an affine neighborhood of $\Sigma_{k(g)}$.

Let $Z^{(j)} \subset \mathbb{A}_k^r \times B_j$ be the projection of Z to B_j as in Section 2E for $1 \leq j \leq n$. Since $Z \rightarrow \mathbb{A}_k^r$ is finite and surjective over an affine neighborhood of Σ , each $Z^{(j)}$ is also finite and surjective over an affine neighborhood of Σ . We can therefore repeat the above process successively for each $Z^{(j)}$ by shrinking U further each time. In the end, we get a dense open subset $U \subset \mathbb{A}_k^r$ such that each $1 \leq j \leq n$ and for each $g \in U$, the scheme $\phi_{g,s,1}^*(Z_{k(g)}^{(j)})$ is regular over a common affine neighborhood of $\Sigma_{k(g)}$. Since the diagram

$$\begin{array}{ccc}
 \mathbb{A}_{k(g)}^r \times \mathbb{A}_{k(g)}^1 \times \square_{k(g)}^{n-1} & \xrightarrow{\pi_j} & \mathbb{A}_{k(g)}^r \times \mathbb{A}_{k(g)}^1 \times \square_{k(g)}^{j-1} \\
 \phi_{g,s,1} \downarrow & & \downarrow \phi_{g,s,1} \\
 \mathbb{A}_{k(g)}^r \times \mathbb{A}_{k(g)}^1 \times \bar{\square}_{k(g)}^{n-1} & \xrightarrow{\pi_j} & \mathbb{A}_{k(g)}^r \times \mathbb{A}_{k(g)}^1 \times \square_{k(g)}^{j-1}
 \end{array} \tag{3-5}$$

commutes and the vertical maps are isomorphisms, it follows that $\phi_{g,s,1}^*(Z_{k(g)}^{(j)}) = (\phi_{g,s,1}^*(Z_{k(g)}))^{(j)}$. We have therefore shown that there is a dense open subset $U \subset \mathbb{A}_k^r$ such that for every $g \in U$ and $1 \leq j \leq n$, the scheme $(\phi_{g,s,1}^*(Z_{k(g)}))^{(j)}$ is regular over a common affine neighborhood of $\Sigma_{k(g)}$. This finishes the proof. \square

Lemma 3.12. *For every integer $s \geq 1$, there is a dense open subset $U \subset \mathbb{A}_k^r$ such that for every $g \in U$, one has $\phi_{g,s,1}^*(Z_{k(g)}) \cap (\Sigma \times \mathbb{A}_k^1 \times F)_{k(g)} = \emptyset$ for every proper face F of \square^{n-1} .*

Proof. We let F be a proper face of \square^{n-1} and let $W = Z \cap (\mathbb{A}_k^r \times \mathbb{A}_k^1 \times F)$. We fix a point $x \in \Sigma$ and consider the diagram (see (3-2)):

$$\begin{array}{ccccccc}
 \theta_x^{-1}(W) & \hookrightarrow & \mathbb{A}_k^r \times \mathbb{G}_{m,k} \times F & \xrightarrow{\iota_x} & \mathbb{A}_k^r \times \mathbb{A}_k^r \times \mathbb{G}_{m,k} \times F & \xrightarrow{P_2} & \mathbb{A}_k^r \times \mathbb{G}_{m,k} \times F \longleftarrow W \\
 & \searrow \omega_{W,x} & & & \downarrow P_1 & & \\
 & & & & \mathbb{A}_k^r & &
 \end{array} \tag{3-6}$$

As in (3-2), the map $\theta_x = P_2 \circ \iota_x$ is an isomorphism. Note also that (see Lemma 3.7) for any $g \in \mathbb{A}_k^r$, the map $\omega_{W,x}^{-1}(g) \rightarrow \phi_{g,s,1}^*(Z) \cap (\{x\} \times \mathbb{A}_k^1 \times F)$, which sends (g, t, \underline{y}) to (x, t, \underline{y}) , is an isomorphism. It follows therefore that the map $\omega_{W,x}$ is not dominant. Equivalently, there exists a dense open $U \subset \mathbb{A}_k^r$ such that the fibers of $\omega_{W,x}$ over U are empty. Shrinking U further if necessary, we can assume that this happens for all $x \in \Sigma$. It is clear that for every $g \in U$, the set $\phi_{g,s,1}^*(Z_{k(g)}) \cap (\Sigma \times \mathbb{A}_k^1 \times F)_{k(g)}$ is empty. This proves the lemma. \square

3F. The proof of the moving lemma for affine spaces. We can now prove the main result of this section, the sfs-moving lemma for the localizations of \mathbb{A}_k^r . We begin with the following intermediate modification step.

Lemma 3.13. *Let k be an infinite field and let $\alpha \in \mathrm{Tz}^n(\mathbb{A}_k^r, n; m)$. Let $V = \mathrm{Spec}(\mathcal{O}_{\mathbb{A}_k^r, \Sigma})$ for a finite subset $\Sigma \subset \mathbb{A}_k^r$ of closed points, with the localization map $j : V \rightarrow \mathbb{A}_k^r$. Assume that $\partial(j^*(\alpha)) = 0$. Then there are cycles $\beta \in \mathrm{Tz}^n(\mathbb{A}_k^r, n; m)$ and $\gamma \in \mathrm{Tz}^n(\mathbb{A}_k^r, n + 1; m)$ with $\partial(j^*(\gamma)) = j^*(\alpha) - j^*(\beta)$ such that each component of β is either dominant over \mathbb{A}_k^r or restricts to zero on V .*

Proof. We choose an integer $s \gg 0$ which is at least as large as the integer $s(Z)$ and the one chosen in Lemmas 3.5 and 3.6 for every irreducible component Z of α . It follows from Lemma 3.2 that $\phi_{g,s}^*(\alpha)$ intersects all faces of \square^n properly. Taking the face $F = \{1\} \times \square^{n-1}$ (and using the containment lemma [Krishna and Park 2016, Proposition 2.2]), we see that $\phi_{g,s,1}^*(\alpha) \in \mathrm{Tz}^n(\mathbb{A}_{k(g)}^r, n; m)$ for all $g \in \mathbb{A}_k^r$. We can also assume that $s \gg 0$ is large enough so that Lemma 3.2 holds also for each boundary of each component of α .

We let $U \subset \mathbb{A}_k^r$ be any dense open which is contained in the intersection of the ones given by Lemmas 3.5 and 3.6 for all irreducible components of $|\alpha|$. We let $g \in U(k)$ be any element. It follows by our choice of g that if Z is a component of α , then $\phi_{g,s,1}^*(Z)$ is either dominant over \mathbb{A}_k^r , or it restricts to zero on V , or satisfies conditions (1) and (2) of Lemma 3.5.

We now compute

$$\begin{aligned} \phi_{g,s}^* \circ \partial(\alpha) &= \phi_{g,s}^* \left(\sum_{i=1}^{n-1} (-1)^i (\partial_i^1 - \partial_i^0)(\alpha) \right) \\ &= \dagger \sum_{i=1}^{n-1} (-1)^i (\partial_{i+1}^1 - \partial_{i+1}^0)(\phi_{g,s}^*(\alpha)) \\ &= - \sum_{i=2}^n (-1)^i (\partial_i^1 - \partial_i^0)(\phi_{g,s}^*(\alpha)), \end{aligned}$$

where $=\dagger$ follows from (3-1). On the other hand, we have

$$\begin{aligned} \partial \circ \phi_{g,s}^*(\alpha) &= \sum_{i=1}^n (-1)^i (\partial_i^1 - \partial_i^0)(\phi_{g,s}^*(\alpha)) \\ &= (-1)(\partial_1^1 - \partial_1^0)(\phi_{g,s}^*(\alpha)) + \sum_{i=2}^n (-1)^i (\partial_i^1 - \partial_i^0)(\phi_{g,s}^*(\alpha)). \end{aligned}$$

It follows that $\partial(\phi_{g,s}^*(\alpha)) + \phi_{g,s}^*(\partial(\alpha)) = (\partial_1^0 - \partial_1^1)(\phi_{g,s}^*(\alpha)) = \alpha - \phi_{g,s,1}^*(\alpha)$. Lemma 3.2 says that $\phi_{g,s}^*(\alpha) \in \mathrm{Tz}^n(\mathbb{A}_k^r, n + 1; m)$. If we let $\gamma = \phi_{g,s}^*(\alpha)$ and $\beta = \phi_{g,s,1}^*(\alpha)$, we see that $\partial(j^*(\gamma)) = j^*(\alpha) - j^*(\beta)$. It also follows that $\partial(j^*(\beta)) = 0$.

We now replace α by β in $\mathrm{Tz}^n(\mathbb{A}_k^r, n; m)$ and repeat the above process. It follows from Lemmas 3.4, 3.5 and 3.6 that after finite steps, we arrive at new cycles $\beta \in \mathrm{Tz}^n(\mathbb{A}_k^r, n; m)$ and $\gamma \in \mathrm{Tz}^n(\mathbb{A}_k^r, n + 1; m)$ such that $\partial(j^*(\gamma)) = j^*(\alpha) - j^*(\beta)$. Moreover, each component of β is either dominant over \mathbb{A}_k^r or restricts to zero on V . \square

Theorem 3.14. *Let k be an infinite perfect field and let $\alpha \in \text{Tz}^n(\mathbb{A}_k^r, n; m)$. Let $V = \text{Spec}(\mathcal{O}_{\mathbb{A}_k^r, \Sigma})$ for a finite subset $\Sigma \subset \mathbb{A}_k^r$ of closed points, with the localization map $j : V \rightarrow \mathbb{A}_k^r$. Assume that $\partial(j^*(\alpha)) = 0$. Then there are cycles $\beta \in \text{Tz}_{\text{sfs}}^n(V, n; m)$ and $\gamma \in \text{Tz}^n(V, n + 1; m)$ such that $\partial(\gamma) = j^*(\alpha) - \beta$.*

Proof. By applying Lemma 3.13 and removing those components of the resulting new cycle α which restrict to zero on V , we can assume that every component of α is dominant over \mathbb{A}_k^r . Note that this does not change $\partial(j^*(\alpha))$.

We now choose an integer $s \gg 0$ which is at least as large as the integer $s(Z)$ and the one chosen in Lemmas 3.10 and 3.11 for every irreducible component Z of α . It follows from Lemma 3.2 that $\phi_{g,s}^*(\alpha)$ intersects all faces of \square^n properly and $\phi_{g,s,1}^*(\alpha) \in \text{Tz}^n(\mathbb{A}_{k(g)}^r, n; m)$ for all $g \in \mathbb{A}_k^r$ (see the proof of Lemma 3.13). We can also assume that $s \gg 0$ is large enough so that Lemma 3.2 holds also for each boundary of each component of α .

We let $U \subset \mathbb{A}_k^r$ be any dense open which is contained in the intersection of the ones given by Lemmas 3.10, 3.11 and 3.12 for all irreducible components of α . Since U is rational and k is infinite, $U(k)$ is a dense subset of U . We let $g \in U(k)$ be any element. We claim that $j^*(\phi_{g,s,1}^*(\alpha)) \in \text{Tz}_{\text{sfs}}^n(V, n; m)$, where $\phi_{g,s}^*(-)$ is defined on $\text{Tz}^n(\mathbb{A}_k^r, n; m)$ by the usual linear extension. By Lemmas 3.10 and 3.11, we only need to show that $\phi_{g,s,1}^*(\alpha) \in \text{Tz}_{\Sigma}^n(\mathbb{A}_k^r, n; m)$. But this is equivalent to showing that $(\Sigma \times \mathbb{A}_k^1 \times F) \cap |\phi_{g,s,1}^*(\alpha)| = \emptyset$ for every proper face F of \square^{n-1} , which in turn follows from Lemma 3.12. The claim is thus proven.

A computation identical to the one in the proof of Lemma 3.13 shows that

$$\partial(\phi_{g,s}^*(\alpha)) + \phi_{g,s}^*(\partial(\alpha)) = (\partial_1^0 - \partial_1^1)(\phi_{g,s}^*(\alpha)) = \alpha - \phi_{g,s,1}^*(\alpha).$$

Lemma 3.2 says that $\phi_{g,s}^*(\alpha) \in \text{Tz}^n(\mathbb{A}_k^r, n + 1; m)$. If $\partial(j^*(\alpha)) = 0$, we can set $\gamma = j^* \circ \phi_{g,s}^*(\alpha)$ and $\beta = j^*(\phi_{g,s,1}^*(\alpha))$. We get $\partial(\gamma) = j^*(\alpha) - \beta$ and we have shown above that $\beta \in \text{Tz}_{\text{sfs}}^n(V, n; m)$. The theorem is now proven. □

Remark 3.15. The proof of Theorem 3.14 (where we take $n \geq 2$, replace B_n by \square^{n-1} and take $s = 0$ everywhere in the proof) also shows that if $n \geq 1$ and $\alpha \in z^n(\mathbb{A}_k^r, n)$ is a higher Chow cycle with $\partial(j^*(\alpha)) = 0$, then we can find $\gamma \in z^n(V, n + 1)$ and $\beta \in z_{\text{sfs}}^n(V, n)$ such that $\partial(\gamma) = j^*(\alpha) - \beta$. Note that $n = 0$ case of this result is trivial.

4. The fs-property of residual cycles

Let k be an infinite perfect field. In this section, we discuss some results on linear projections in projective spaces, and show how these projections can be used to equip the residual cycle of a given cycle with certain finiteness properties over the base scheme. The main result of Section 4 is Theorem 4.15. It will be used later in proving the fs-moving lemma (see Lemma 8.7), a precursor to the final sfs-moving lemma.

For $0 \leq n < N$ and a linear subspace $H \subset \mathbb{P}_k^N$ defined over k , let $\text{Gr}(n, H)$ be the Grassmannian scheme of n -dimensional linear subspaces of \mathbb{P}_k^N contained in H . This is a homogeneous space of dimension

$(\dim(H) - n)(n + 1)$. Unless we specify the field of definition, a linear subspace of \mathbb{P}_k^N will mean a k -linear subspace.

Given two closed subschemes $Y, Y' \subset \mathbb{P}_k^N$, let $\text{Sec}(Y, Y') \subset \mathbb{P}_k^N$ be the union of all lines $\ell_{yy'}$ joining distinct points $y \in Y, y' \in Y'$. In general, we have $\dim(\text{Sec}(Y, Y')) \leq \dim(Y) + \dim(Y') + 1$. If $Y = Y'$, the scheme $\text{Sec}(Y, Y') = \text{Sec}(Y)$ is the secant variety of Y . If $Y' = L$ is a linear subspace, then $\text{Sec}(Y, L) = C_L(Y)$ is the cone over Y with vertices in L .

4A. Containment and avoidance. Let $0 \leq m \leq n < N$ be integers and let $S, T \subset \mathbb{P}_k^N$ be two disjoint subsets.

Definition 4.1. We denote the set of n -dimensional linear subspaces of \mathbb{P}_k^N containing S by $\text{Gr}_S(n, \mathbb{P}_k^N)$. We write $\text{Gr}_S(n, \mathbb{P}_k^N)$ as $\text{Gr}_x(n, \mathbb{P}_k^N)$ if $S = \{x\}$ is a closed point. We denote the set of n -dimensional linear subspaces of \mathbb{P}_k^N which do not intersect S by $\text{Gr}(S, n, \mathbb{P}_k^N)$. If $S = \{x\}$, we write $\text{Gr}(S, n, \mathbb{P}_k^N)$ as $\text{Gr}(x, n, \mathbb{P}_k^N)$. We let $\text{Gr}_S(T, n, \mathbb{P}_k^N) := \text{Gr}_S(n, \mathbb{P}_k^N) \cap \text{Gr}(T, n, \mathbb{P}_k^N)$. For any linear subspace $L \subset \mathbb{P}_k^N$, we define $\text{Gr}_S(n, L)$ and $\text{Gr}(T, n, L)$ similarly.

One checks that, when $M \subset \mathbb{P}_k^N$ is a linear subspace of dimension m , then $\text{Gr}_M(n, \mathbb{P}_k^N)$ is a homogeneous space which is an irreducible closed subscheme of $\text{Gr}(n, \mathbb{P}_k^N)$ of dimension $(N - n)(n - m)$. The following result is elementary. We leave the proof as an exercise.

Lemma 4.2. *Let $N > n$. (1) If $S' \subset S$, then $\text{Gr}(S, n, \mathbb{P}_k^N) \subset \text{Gr}(S', n, \mathbb{P}_k^N)$. (2) For any finite closed set $S \subset \mathbb{P}_k^N$, $\text{Gr}(S, n, \mathbb{P}_k^N) \subset \text{Gr}(n, \mathbb{P}_k^N)$ is a dense open subset.*

Lemma 4.3. *Let $X \subset \mathbb{P}_k^N$ be a closed subscheme of dimension $r \geq 1$ with $N \gg r$ and let $H \subset \mathbb{P}_k^N$ be a hyperplane, not containing any irreducible component of X . Then $\text{Gr}(X, N - r - 1, H)$ is a dense open subset of $\text{Gr}(N - r - 1, H)$.*

Proof. Consider the incidence scheme $S = \{(x, L) \in X \times \text{Gr}(N - r - 1, H) \mid x \in L\}$. We have the obvious projection maps $X \xleftarrow{\pi_1} S \xrightarrow{\pi_2} \text{Gr}(N - r - 1, H)$.

Each fiber of π_1 over $X \setminus (X \cap H)$ is empty. It is a smooth morphism over $X \cap H$ with its fiber over $x \in X \cap H$ to be $\text{Gr}_x(N - r - 1, H)$, whose dimension is $((N - 1) - (N - r - 1))(N - r - 1 - 0) = r(N - r - 1)$. It follows that $\dim(S) \leq \dim(X \cap H) + \dim \text{Gr}_x(N - r - 1, H) = r - 1 + r(N - r - 1) = r(N - r) - 1$. Thus, $\pi_2(S)$ is a closed subscheme of $\text{Gr}(N - r - 1, H)$ of dimension $\leq r(N - r) - 1$ which is less than $\dim \text{Gr}(N - r - 1, H) = r(N - r)$. Hence, $\text{Gr}(X, N - r - 1, H) = \text{Gr}(N - r - 1, H) \setminus \pi_2(S)$ is a dense open subset. \square

4B. Transverse intersection. For a reduced scheme X , let $X_{\text{sing}} \subset X$ be the singular locus of X and let X_{sm} be its complement. For a closed subscheme $X \subset \mathbb{P}_k^N$, let $\text{Gr}^{\text{tr}}(X, n, \mathbb{P}_k^N)$ denote the set of n -dimensional linear subspaces which *do not* intersect X_{sing} , and whose intersection with X_{sm} is transverse (if not empty). We let

$$\text{Gr}^{\text{tr}}(X, S, n, \mathbb{P}_k^N) = \text{Gr}(S, n, \mathbb{P}_k^N) \cap \text{Gr}^{\text{tr}}(X, n, \mathbb{P}_k^N) \quad \text{and} \quad \text{Gr}_S^{\text{tr}}(X, n, \mathbb{P}_k^N) = \text{Gr}_S(n, \mathbb{P}_k^N) \cap \text{Gr}^{\text{tr}}(X, n, \mathbb{P}_k^N).$$

For a linear subspace $H \subset \mathbb{P}_k^N$, we define $\text{Gr}^{\text{tr}}(X, S, n; H)$ and $\text{Gr}_S^{\text{tr}}(X, n; H)$ similarly.

Lemma 4.4. *Let $r \geq 2$ be an integer and suppose $N \gg r$. Let $H \subset \mathbb{P}_k^N$ be a hyperplane. Let $L \subset \mathbb{P}_k^N$ be a linear subspace of dimension $N - r + 1$ intersecting H transversely and let $X \subset L$ be a curve (not necessarily connected) none of whose components is contained in H . Then the set of linear subspaces in $\text{Gr}^{\text{tr}}(L, X, N - 2, H)$ is a dense open subset of $\text{Gr}(N - 2, H)$.*

Proof. Observe that $\text{Gr}^{\text{tr}}(L, N - 2, H)$ is a dense open subset of $\text{Gr}(N - 2, H)$. Consider the map $v_L : \text{Gr}^{\text{tr}}(L, N - 2, H) \rightarrow \text{Gr}(N - r - 1, L \cap H)$ given by $v_L(M) = L \cap M$. This v_L is a smooth surjective morphism of relative dimension $2(r - 1)$. It follows from Lemma 4.3 that $\text{Gr}(X, N - r - 1, L \cap H)$ is a dense open subset of $\text{Gr}(N - r - 1, L \cap H)$, so $v_L^{-1}(\text{Gr}(X, N - r - 1, L \cap H))$ is a dense open subset of $\text{Gr}^{\text{tr}}(L, N - 2, H)$, and hence a dense open subset of $\text{Gr}(N - 2, H)$. \square

4C. Affine Veronese embedding and linear projection. Recall that for positive integers $m, d \geq 1$, the Veronese embedding $v_{m,d} : \mathbb{P}_k^m \hookrightarrow \mathbb{P}_k^N$ is a closed embedding given by $v_{m,d}([\underline{x}]) = [M_0(\underline{x}), \dots, M_N(\underline{x})] = [M(\underline{x})]$, where $N = \binom{m+d}{m} - 1$ and $\{M_0, \dots, M_N\}$ are all monomials in $\{x_0, \dots, x_m\}$ of degree d , arranged in the lexicographic order.

If $[y_0, \dots, y_N] \in \mathbb{P}_k^N$ denotes the projective coordinates, it is clear that $v_{m,d}^{-1}(\{y_0 = 0\}) = \{x_0^d = 0\}$. In particular, the Veronese embedding yields Cartesian squares

$$\begin{array}{ccccc}
 \mathbb{A}_k^m & \longrightarrow & \mathbb{P}_k^m & \longleftarrow & dH_{m,0} \\
 v_{m,d} \downarrow & & \downarrow v_{m,d} & & \downarrow v_{m,d} \\
 \mathbb{A}_k^N & \longrightarrow & \mathbb{P}_k^N & \longleftarrow & H_{N,0},
 \end{array} \tag{4-1}$$

where $H_{m,0} \subset \mathbb{P}_k^m$ is the hyperplane $\{x_0 = 0\}$ and the vertical arrows are all closed embeddings. The closed embedding $v_{m,d} : \mathbb{A}_k^m \hookrightarrow \mathbb{A}_k^N$ is given by $v_{m,d}(y_1, \dots, y_m) = (M'_1, \dots, M'_N)$, where $\{M'_1, \dots, M'_N\}$ is the induced ordered set of all monomials in $\{y_1, \dots, y_m\}$ of degree bounded by d .

Let $1 \leq r < N$ be two integers. Recall (e.g., see [Krishna and Park 2012, Lemma 6.1]) that when $L \subset \mathbb{P}_k^N$ is a linear subspace of dimension $N - r - 1$, there is an associated projection map $\phi_L : \mathbb{P}_k^N \setminus L \rightarrow \mathbb{P}_k^r$, where \mathbb{P}_k^r is a linear subspace of \mathbb{P}_k^N such that $L \cap \mathbb{P}_k^r = \emptyset$. This map ϕ_L defines a vector bundle over \mathbb{P}_k^r of rank $N - r$, whose fiber over a point $x \in \mathbb{P}_k^r$ is the affine space $C_x(L) \setminus L$, where $C_x(L) = \text{Sec}(\{x\}, L)$.

Remark 4.5. The referee asked whether the above vector bundle $\phi_L : \mathbb{P}_k^N \setminus L \rightarrow \mathbb{P}_k^r$ is isomorphic to $\mathcal{O}_{\mathbb{P}_k^r}(1)^{\oplus(N-r)}$. Indeed, ϕ_L is (up to an isomorphism) the projection map of quotient stacks $\phi_L : [((\mathbb{A}^{r+1} \setminus \{0\}) \times_k V)/\mathbb{G}_m] \rightarrow [(\mathbb{A}^{r+1} \setminus \{0\})/\mathbb{G}_m]$, where $V = k^{N-r}$ and the \mathbb{G}_m -action everywhere is by scalar multiplication. Since $[((\mathbb{A}^{r+1} \setminus \{0\}) \times_k V)/\mathbb{G}_m] \cong [(\mathbb{A}^{r+1} \setminus \{0\})/\mathbb{G}_m] \times_{B\mathbb{G}_m} [V/\mathbb{G}_m]$, one identifies ϕ_L with the map $\mathbb{P}_k^r \times_{B\mathbb{G}_m} \pi^*(V(1)^{\oplus(N-r)}) \rightarrow \mathbb{P}_k^r$, where $V(1)$ is the line bundle on $B\mathbb{G}_m := [\text{Spec}(k)/\mathbb{G}_m]$ associated to the 1-dimensional \mathbb{G}_m -representation given by the scalar multiplication on k , and $\pi : [(\mathbb{A}^{r+1} \setminus \{0\})/\mathbb{G}_m] \rightarrow B\mathbb{G}_m$ is the canonical projection.

Note that in general, if we let \mathbb{G}_m act on k by weight $n \in \mathbb{Z}$ (i.e., $\lambda \cdot x = \lambda^n x$) and let $V(n)$ denote the corresponding line bundle on $B\mathbb{G}_m$, then $\pi^*(V(n))$ is isomorphic to $\mathcal{O}_{\mathbb{P}_k^r}(n)$. Hence the above $\pi^*(V(1)^{\oplus(N-r)})$ is isomorphic to $\mathcal{O}_{\mathbb{P}_k^r}(1)^{\oplus(N-r)}$, as wished.

Definition 4.6. Recall that if $X \subset \mathbb{P}_k^N$ is a closed subscheme with $X \cap L = \emptyset$, then ϕ_L restricted to X defines a projection $\phi_L : \phi_L|_X : X \rightarrow \mathbb{P}_k^r$. We call it the linear projection of X away from L . Since this is a morphism of projective schemes with affine fibers, it must be a finite morphism. In particular, $\dim(X) \leq r$.

We shall use the following situation often: let $H \subset \mathbb{P}_k^N$ be a hyperplane containing L and $X \subset \mathbb{P}_k^N$ a closed subscheme with $X \cap L = \emptyset$ and $X \not\subset H$. Then ϕ_L defines the Cartesian squares of morphisms

$$\begin{array}{ccccc}
 X \setminus H & \longrightarrow & X & \longleftarrow & X \cap H \\
 \downarrow & & \downarrow & & \downarrow \\
 \mathbb{P}_k^r \setminus H & \longrightarrow & \mathbb{P}_k^r & \longleftarrow & \mathbb{P}_k^r \cap H.
 \end{array} \tag{4-2}$$

Together with (4-1), we deduce the following fact, which we use often:

Lemma 4.7. Let $X \hookrightarrow \mathbb{A}_k^m$ be an affine scheme of dimension $r \geq 1$ and let $\bar{X} \hookrightarrow \mathbb{P}_k^m$ be its projective closure. Then, for $d \geq 1$, the Veronese embedding $v_{m,d} : \mathbb{P}_k^m \hookrightarrow \mathbb{P}_k^N$ and the linear projection away from $L \in \text{Gr}(N - r - 1, \mathbb{P}_k^N)(k)$ yield a Cartesian diagram with finite vertical maps

$$\begin{array}{ccc}
 X & \longrightarrow & \bar{X} \\
 \phi_L \downarrow & & \downarrow \phi_L \\
 \mathbb{A}_k^r & \longrightarrow & \mathbb{P}_k^r
 \end{array} \tag{4-3}$$

if $L \in \text{Gr}(\bar{X}, N - r - 1, H_{N,0})(k)$, where $H_{N,0} = \{y_0 = 0\} \subset \mathbb{P}_k^N$ as in (4-1).

4D. The Set-up. Let k be an infinite perfect field. Here, we introduce the basic Set-up that will be used for most of the paper. This set of assumptions will be referred to as the Set-up of Section 4D.

(1) *The spaces:* Let X be an equidimensional reduced projective k -scheme of dimension $r \geq 1$ with a given closed embedding $\eta : X \hookrightarrow \mathbb{P}_k^N$ with $N \gg r$ and of degree $d + 1 \gg 0$. We let \hat{B} be a smooth projective geometrically integral k -scheme of positive dimension and let $B \subset \hat{B}$ be a nonempty affine open subset with $F := \hat{B} \setminus B$. Let $\Sigma \subset X_{\text{sm}}$ be a finite set of closed points.

(2) *The linear projections:* Suppose that $H \subset \mathbb{P}_k^N$ is a hyperplane not meeting Σ , and that $X \setminus (X \cap H) \subset X_{\text{sm}}$. For $L \in \text{Gr}(X, N - r - 1, H)(k)$, let $\phi_L : X \rightarrow \mathbb{P}_k^r$ be the linear projection away from L . If L is fixed in a given context, we often drop it from the notation of ϕ_L and write as $\phi : X \rightarrow \mathbb{P}_k^r$. We write $\hat{\phi} = \hat{\phi}_L = \phi_L \times \text{id}_{\hat{B}} : X \times \hat{B} \rightarrow \mathbb{P}_k^r \times \hat{B}$.

(3) *The cycles:* Let $Z \subset X \times \hat{B}$ be a reduced closed subscheme with irreducible components $\{Z_1, \dots, Z_s\}$, each of dimension r . We suppose that both $X \times F$ and $H \times \hat{B}$ intersect properly with each irreducible component of Z . We let $\hat{f} : Z \rightarrow X$ and $\hat{g} : Z \rightarrow \hat{B}$ denote the restrictions of the projection maps. Let $E \subset \hat{B}$ be a closed subset containing F such that no component of Z is contained in $\hat{g}^{-1}(E)$. We suppose that each projection $Z_i \rightarrow \hat{B}$ is nonconstant.

(4) *The residual schemes and residual sets:* Let $L^+(Z)$ be the closure of $\hat{\phi}^{-1}(\hat{\phi}(Z)) \setminus Z$ in $X \times \hat{B}$ with the reduced closed subscheme structure. For any closed point $x \in \bar{X}$, we write $L^+(\{x\})$ as $L^+(x)$. We let $L^+(\Sigma) = \bigcup_{x \in \Sigma} L^+(x)$.

4E. A Nisnevich property of linear projections. The first result on “moving” our cycle Z is the following:

Lemma 4.8. *We are under the Set-up of Section 4D. After replacing the embedding $\eta : X \hookrightarrow \mathbb{P}_k^N$ by a bigger one via a Veronese embedding if necessary, there exists a dense open subset $\mathcal{U} \subset \text{Gr}(X, N-r-1, H)$ such that each $L \in \mathcal{U}(k)$ satisfies the following:*

- (1) ϕ_L is étale at Σ .
- (2) $\phi_L(x) \neq \phi_L(x')$ for each pair of distinct points $x \neq x' \in \Sigma$.
- (3) $k(\phi_L(x)) \xrightarrow{\sim} k(x)$ for all $x \in \Sigma$.
- (4) $L^+(x) \neq \emptyset$ for all $x \in \Sigma$.
- (5) $L^+(x) \cap \hat{f}(\hat{g}^{-1}(E)) = \emptyset$ for all $x \in \Sigma$.
- (6) $L^+(x) \cap \hat{f}(Z_i) = \emptyset$ for all $x \in \Sigma$ if $\hat{f} : Z_i \rightarrow X$ is not dominant over any irreducible component of X .

Proof. Replacing the given embedding $X \hookrightarrow \mathbb{P}_k^N$ by its composition with a Veronese embedding, we may begin with a closed embedding $X \hookrightarrow \mathbb{P}_k^N$ such that $N \gg r$ and the degree of X in \mathbb{P}_k^N is bigger than one.

Step 1. *First suppose that k is algebraically closed.* It follows from our assumption that $\dim(\hat{g}^{-1}(E)) \leq r-1$. Since \hat{f} is projective, it follows that $\hat{f}(\hat{g}^{-1}(E))$ is a closed subset of X of dimension at most $r-1$. We let $W \subset X$ be the union of X_{sing} , $\hat{f}(\hat{g}^{-1}(E))$ and the images of all components of Z which are not dominant over X . This is a closed subset of X such that $\dim(W) \leq r-1$. In particular, $\dim(\text{Sec}(D_1, W \cup D_2)) \leq r$ for any finite closed subsets $D_1, D_2 \subset X$. It follows from Lemma 4.3 that $\mathcal{U}_1 := \bigcap_{x \in \Sigma} \text{Gr}(X \cup \text{Sec}(\{x\}, W \cup (\Sigma \setminus \{x\})), N-r-1, H)$ is dense open in $\text{Gr}(N-r-1, H)$. Furthermore, any $L \in \mathcal{U}_1(k)$ satisfies (5) and (6) by construction.

We continue the proof of the rest of the properties. Let $T_{\Sigma, X} \subset \mathbb{P}_k^N$ be the union of the tangent spaces to X at all points of Σ . Since $\Sigma \subset X_{\text{sm}}$, we have $T_{\Sigma, X} = T_{\Sigma, X_{\text{sm}}}$, which is a finite union of linear subspaces of dimension r . For each $x \in \Sigma$, the set $\mathcal{Z}_x = X \cup T_{\Sigma, X} \cup \text{Sec}(\{x\}, X_{\text{sing}} \cup (\Sigma \setminus \{x\}))$ is closed in \mathbb{P}_k^N of dimension r . Therefore, the set $\mathcal{U} = \bigcap_{x \in \Sigma} \text{Gr}(\mathcal{Z}_x, N-r-1, H) \cap \mathcal{U}_1$ is dense open in $\text{Gr}(N-r-1, H)$ by Lemma 4.3. By construction, each $L \in \mathcal{U}(k)$ defines the finite surjective map $\phi_L : X \rightarrow \mathbb{P}_k^r$, which is unramified at Σ and separates the points of Σ . In particular, (2) holds.

Since X_{sm} is regular and dense in X , it follows that $\phi_L|_{X_{\text{sm}}} : X_{\text{sm}} \rightarrow \mathbb{P}_k^r$ is a dominant and quasifinite morphism between regular k -schemes. In particular, the map $\mathcal{O}_{\mathbb{P}_k^r, \phi_L(x)} \rightarrow \mathcal{O}_{X, x}$ is a local homomorphism of regular local rings with the finite closed fiber for each $x \in \Sigma$. It follows from [EGA IV₂ 1965, Proposition (6.1.5), page 136] (or [Matsumura 1986, Theorem 23.1, page 179]) that ϕ_L is flat at each point of Σ . Hence ϕ_L is étale at Σ , being flat and unramified, proving (1).

Since $k = \bar{k}$, the isomorphisms of the residue fields, (3) is evident. Property (4) follows because $\deg(\phi_L) > 1$ by the assumptions on the chosen Veronese embedding of X . This proves the lemma in Step 1 when k is algebraically closed.

Step 2. *Now suppose that k is any infinite perfect field.* Let \bar{k} be an algebraic closure. For any k -scheme A , let $\pi_A : A_{\bar{k}} \rightarrow A$ be the base change to \bar{k} . We have that $\Sigma_{\bar{k}} = \pi_X^{-1}(\Sigma)$ is still a finite closed set of the regular scheme $X_{\text{sm},\bar{k}}$. By Step 1 applied to $X_{\bar{k}}$, $H_{\bar{k}}$ and $\Sigma_{\bar{k}}$, there exists a dense open $\mathcal{U}' \subset \text{Gr}(N - r - 1, H_{\bar{k}})$ where the mentioned properties (1)–(6) hold.

Since k is perfect, there exists a finite Galois extension $k \subset k'$ in \bar{k} such that \mathcal{U}' is defined over k' . Let $\mathcal{U} := \bigcap_{\sigma \in \text{Gal}(k'/k)} \sigma \cdot \mathcal{U}'$. This is a nonempty open subset defined over the radical closure of k in k' , but since k is perfect, this radical closure is equal to k . Hence $\mathcal{U} \subset \text{Gr}(N - r - 1, H)$ and it is defined over k (see [Colliot-Thélène et al. 1997, Lemma 3.4.3]). Here we have $\mathcal{U}_{\bar{k}} \subset \mathcal{U}'$. Now, for each $L \in \mathcal{U}(k)$, we have $X \cap L = \emptyset$ by our choice of the open set. So, we get a finite surjective map $\phi_L : X \rightarrow \mathbb{P}_k^r$ over k .

We prove that ϕ_L is étale at each point $x \in \Sigma$. Let $y := \phi_L(x)$. By the faithfully flat descent [EGA IV₄ 1967, Corollaire (17.7.3)(ii), page 72], the map $\phi_L : \text{Spec}(\mathcal{O}_{X,x}) \rightarrow \text{Spec}(\mathcal{O}_{\mathbb{P}_k^r,y})$ is étale if and only if its faithfully flat base change $\phi_{L,\bar{k}} : \text{Spec}(\mathcal{O}_{X_{\bar{k}},x_{\bar{k}}}) \rightarrow \text{Spec}(\mathcal{O}_{\mathbb{P}_{\bar{k}}^r,y_{\bar{k}}})$ of the semilocal schemes via $\text{Spec}(\bar{k}) \rightarrow \text{Spec}(k)$ is étale. Here, $x_{\bar{k}} := \pi_X^{-1}(x)$ and $y_{\bar{k}} := \pi_{\mathbb{P}_k^r}^{-1}(y)$. But Step 1 shows that the latter map $\phi_{L,\bar{k}}$ is étale at each point of the set $x_{\bar{k}} \subset \Sigma_{\bar{k}}$, thus so is the former ϕ_L at x . This proves (1).

Since $\phi_{L,\bar{k}}$ separates the points of $\Sigma_{\bar{k}}$ by construction, (2) is obvious. Furthermore, this shows that for each $x \in \Sigma$, the map $\phi_{L,\bar{k}} : \pi_X^{-1}(x) \rightarrow \pi_{\mathbb{P}_k^r}^{-1}(y)$ is injective, where $y = \phi_L(x)$. Hence by Lemma 4.9 below, we have $k(x) = k(y)$, which proves (3). Property (4) is evident because $\deg(\phi_L) > 1$ and $k(\phi(x)) \simeq k(x)$ for each $x \in \Sigma$ by (3).

Conditions (5) and (6) are apparent for any $L \in \mathcal{U}(k)$ because for every $x \in \Sigma$, we have that $(L_{\bar{k}})^+(x') \cap \hat{f}_{\bar{k}}(\hat{g}_{\bar{k}}^{-1}(E_{\bar{k}})) = \emptyset = (L_{\bar{k}})^+(x') \cap \hat{f}_{\bar{k}}(Z_{i,\bar{k}})$ for all x' lying in the finite set $\pi_X^{-1}(x) \subset \Sigma_{\bar{k}}$. Note here that if Z_i is not dominant over a component of X , then no component of $Z_{i,\bar{k}}$ can be dominant over any component of $X_{\bar{k}}$. This finishes the proof of the lemma. □

We used the following in the middle of the proof of the above Lemma 4.8.

Lemma 4.9. *Let k be an infinite perfect field and let $\phi : X \rightarrow Y$ be a finite morphism of k -schemes. Consider the base change Cartesian square:*

$$\begin{array}{ccc}
 X_{\bar{k}} & \xrightarrow{\phi_{\bar{k}}} & Y_{\bar{k}} \\
 \pi_X \downarrow & & \downarrow \pi_Y \\
 X & \xrightarrow{\phi} & Y
 \end{array}
 \tag{4-4}$$

Let $x \in X$ be a closed point and let $y := \phi(x)$. Then one has $|\pi_Y^{-1}(y)| \leq |\pi_X^{-1}(x)|$. The equality holds if and only if $[k(x) : k(y)] = 1$. Furthermore, this equality holds when the map $\phi_{\bar{k}} : \pi_X^{-1}(x) \rightarrow \pi_Y^{-1}(y)$ is injective.

Proof. Since k is perfect, we have $|\pi_X^{-1}(x)| = [k(x) : k]$ and $|\pi_Y^{-1}(y)| = [k(y) : k]$. So, the field extensions $k \hookrightarrow k(y) \hookrightarrow k(x)$ imply the first and the second assertions. If the map $\phi_{L_k} : \pi_X^{-1}(x) \rightarrow \pi_Y^{-1}(y)$ is injective, then $|\pi_Y^{-1}(y)| \geq |\pi_X^{-1}(x)|$. The last part of the lemma thus follows. \square

4F. Some algebraic results. We discuss some algebraic results that will be needed.

Lemma 4.10. *Let $f : A \rightarrow B$ be an injective finite unramified local homomorphism of noetherian local rings that induces an isomorphism of the residue fields. Then f is an isomorphism.*

Proof. Let \mathfrak{m}_A and \mathfrak{m}_B be the maximal ideals of A and B , respectively. Since f is finite, to show that f is surjective it suffices to show that $A/\mathfrak{m}_A \rightarrow B/(\mathfrak{m}_A B)$ is surjective by Nakayama’s lemma. But this follows because the map $A/\mathfrak{m}_A \rightarrow B/\mathfrak{m}_B$ is an isomorphism and so is the map $B/(\mathfrak{m}_A B) \rightarrow B/\mathfrak{m}_B$ as f is unramified. \square

Lemma 4.11. *Let $f : Y' \rightarrow Y$ be a finite surjective morphism of regular k -schemes. Let $W \subset Y$ be an irreducible closed subset and let $y \in W$ be a closed point. Let $S = f^{-1}(y)$ and $W' = f^{-1}(W)$. Let $x \in S$ and let $Z \subset W'$ be an irreducible component passing through x . Suppose that f is étale at x and $k(y) \xrightarrow{\sim} k(x)$. Then $Z \cap S = \{x\}$ if and only if Z is the only component of W' passing through x .*

Proof. We first observe that f must be a flat morphism (see [Hartshorne 1977, Exercise III-10.9, page 276]). We next note that any irreducible component of W' that passes through x will be in the connected component of Y' containing x . So, we may assume Y' is connected. On the other hand, $W \subset Y$ being irreducible, it must belong to a unique connected component of Y . Hence, we may also assume that Y is connected.

Now, first suppose $S = \{x\}$. We claim that f is an isomorphism locally around y , so that the lemma holds trivially. Indeed, it follows from Lemma 4.10 that the map $\mathcal{O}_{Y,y} \rightarrow \mathcal{O}_{Y',x}$ is an isomorphism. This implies that f is a finite and flat map with $[k(Y') : k(Y)] = 1$ (see [Liu 2002, Exercise 5.1.25, page 176]) and hence must be an isomorphism.

We now suppose $|S| > 1$. Consider the commutative diagram of semilocal rings

$$\begin{array}{ccccc}
 \mathcal{O}_{Y,y} & \xrightarrow{\alpha_1} & \mathcal{O}_{Y',S} & \xrightarrow{\alpha_2} & \mathcal{O}_{Y',x} \\
 \downarrow \beta_1 & & \downarrow \beta_2 & & \downarrow \beta_3 \\
 \mathcal{O}_{W,y} & \xrightarrow{\alpha_3} & \mathcal{O}_{W',S} & \xrightarrow{\alpha_4} & \mathcal{O}_{W',x} \\
 & \searrow \gamma & \downarrow \beta_4 & & \downarrow \beta_5 \\
 & & \mathcal{O}_{Z,S} & \xrightarrow{\alpha_5} & \mathcal{O}_{Z,x} \\
 & \nearrow \gamma' & & &
 \end{array} \tag{4-5}$$

where $\gamma := \beta_4 \circ \alpha_3$ and $\gamma' := \alpha_5 \circ \gamma$. Here, α_1 and α_3 are finite and flat, and $\alpha_2 \circ \alpha_1$ is étale. The lemma is equivalent to that α_5 is an isomorphism if and only if β_5 is.

Suppose α_5 is an isomorphism. Since β_4 is surjective and α_3 is finite, the map γ is finite. Thus, γ' is a finite map of local rings. Since $\alpha_2 \circ \alpha_1$ is étale, the map $\alpha_4 \circ \alpha_3$ is also étale. Since β_5 is surjective, we see

that γ' is unramified. Thus, γ' is a finite and unramified map of local rings. Since $Z \rightarrow W$ is surjective and $k(y) \simeq k(x)$, the map γ' is an isomorphism by Lemma 4.10. In particular, $\alpha_4 \circ \alpha_3$ is an étale map of local rings such that $\beta_5 \circ \alpha_4 \circ \alpha_3$ is an isomorphism, in particular, étale. It follows that β_5 is étale, by [EGA IV₄ 1967, Proposition (17.3.4), page 62]. Thus, β_5 is a surjective étale map of local rings. But it can happen only if β_5 is an isomorphism.

Conversely, suppose that β_5 is an isomorphism. Let \mathfrak{p} be the minimal prime of $\mathcal{O}_{W',S}$ such that $\mathcal{O}_{W',S}/\mathfrak{p} = \mathcal{O}_{Z,S}$ and let $\{\mathfrak{p}_1, \dots, \mathfrak{p}_m\}$ denote the set of distinct minimal primes of $\mathcal{O}_{W',S}$ different from \mathfrak{p} . To show that α_5 is an isomorphism, we need to show that $\mathfrak{p} + \mathfrak{p}_i = \mathcal{O}_{W',S}$ for all $1 \leq i \leq m$.

Claim 1. $\mathfrak{p}_i \mathcal{O}_{W',x} = \mathcal{O}_{W',x}$ for all $1 \leq i \leq m$.

Proof. Note that $\mathcal{O}_{W',x}$ is an integral domain because $\mathcal{O}_{Z,x}$ is an integral domain and β_5 is an isomorphism. Thus, we must have either $\mathfrak{p}_i \mathcal{O}_{W',x} = 0$ or $\mathfrak{p}_i \mathcal{O}_{W',x} = \mathcal{O}_{W',x}$. In the first case, we have $\mathfrak{p}_i \mathcal{O}_{Z,x} = 0$ as β_5 is an isomorphism. Equivalently, $\alpha_5 \circ \beta_4(\mathfrak{p}_i) = 0$. Since $\mathfrak{p}_i \neq \mathfrak{p}$, and $\mathfrak{p}_i, \mathfrak{p}$ are minimal, there is $a_i \in \mathfrak{p}_i \setminus \mathfrak{p}$ such that $\beta_4(a_i) \neq 0$. Hence, $\alpha_5 \circ \beta_4(a_i) \neq 0$, because α_5 is injective being a localization of an integral domain. This is a contradiction. Thus, we must have $\mathfrak{p}_i \mathcal{O}_{W',x} = \mathcal{O}_{W',x}$ for each i , proving Claim 1. \square

Let \mathfrak{m} be the maximal ideal of $\mathcal{O}_{W',S}$ defining the closed point x . By Claim 1, for any $1 \leq i \leq m$ there exists $a_i \in \mathfrak{p}_i \setminus \mathfrak{m}$ in $\mathcal{O}_{W',S}$ such that $\alpha_4(a_i)$ is invertible. Let $a = \prod_{i=1}^m a_i$. We see that there are nonzero elements $b, c \in \mathcal{O}_{Z',S}$ with $c \notin \mathfrak{m}$ such that $c(1 - ab) = 0$.

Claim 2. $1 - ab \in \mathfrak{p}$.

Proof. Let $v = 1 - ab$. Then, we have $cv = 0 \in \mathfrak{m}$ with $c \notin \mathfrak{m}$, so that $v \in \mathfrak{m}$ and $\alpha_4(v) = 0$. Toward contradiction, suppose $v \notin \mathfrak{p}$. Then $v \in \mathfrak{m} \setminus \mathfrak{p}$, so that $\beta_4(v) \neq 0$. Thus $\beta_5 \circ \alpha_4(v) = \alpha_5 \circ \beta_4(v) \neq 0$ because α_5 is injective. But this contradicts that $\alpha_4(v) = 0$. Hence, we have $v \in \mathfrak{p}$, proving Claim 2. \square

By Claim 2, we have $v \in \mathfrak{p}$, $ab \in \mathfrak{p}_i$ for all i , while $v - ab = 1$. This shows that $\mathfrak{p} + \mathfrak{p}_i = \mathcal{O}_{W',S}$ for all $1 \leq i \leq m$. Thus, α_5 is an isomorphism. \square

4G. Birationality under linear projections. Using Lemma 4.8, we shall show that the linear projections often give birational morphisms when restricted to a given integral closed subscheme. But first, we derive the following consequence of the results we proved in Section 4E and Section 4F. We continue to work with the Set-up of Section 4D.

We use a trick of “marking” irreducible components: for each $1 \leq i \leq s$, we fix a closed point $\alpha_i \in (Z_i)_{\text{sm}}$ such that (1) $\alpha_i \notin Z_j$ for $j \neq i$, (2) $x_i = \hat{f}(\alpha_i) \in X_{\text{sm}}$ but not in Σ , and (3) $b_i = \hat{g}(\alpha_i) \in B$. Note here that $\alpha_i \in (Z_i)_{\text{sm}}$ and $x_i \in X_{\text{sm}}$ can be achieved as follows: by the assumptions of the Set-up of Section 4D, each Z_i intersects $H \times \hat{B}$ properly and $X \setminus (X \cap H) \subset X_{\text{sm}}$. Then any choice of a point in $Z_i|_{(X \setminus (X \cap H)) \times B}$ maps to a point of X_{sm} . Moreover, perfectness of k implies that $(Z_i)_{\text{sm}} \cap Z_i|_{(X \setminus (X \cap H)) \times B} \neq \emptyset$. Let $\Xi = \{x_1, \dots, x_s\} \cup \Sigma$ and $E = \{b_1, \dots, b_s\} \cup F$. Since $Z_i \not\subset X \times F$ and $Z_i \rightarrow \hat{B}$ is nonconstant by the Set-up of Section 4D, no component of Z lies in $\hat{g}^{-1}(E)$.

Lemma 4.12. *After replacing the embedding $X \hookrightarrow \mathbb{P}_k^N$ by a bigger one via a Veronese embedding if necessary, there is a dense open subset $\mathcal{U} \subset \text{Gr}(X, N - r - 1, H)$ such that each $L \in \mathcal{U}(k)$ has the property that $Z_i \cap \hat{\phi}_L^{-1}(\hat{\phi}_L(\alpha_i)) = \{\alpha_i\}$ for all $1 \leq i \leq s$.*

Proof. We let $\pi : \text{Spec}(\bar{k}) \rightarrow \text{Spec}(k)$ denote the base change map. For any $A \in \mathbf{Sch}_k$, we shall write $\pi_A : A_{\bar{k}} \rightarrow A$ simply as π using a shorthand.

We fix i . Let $\beta_i := \hat{\phi}_L(\alpha_i)$. Let $\pi^{-1}(\alpha_i) = \{\alpha_{ij}\}_j$, which is a finite set of points, and let $x_{ij} := \hat{f}_{\bar{k}}(\alpha_{ij})$, $b_{ij} := \hat{g}_{\bar{k}}(\alpha_{ij})$. Note that all of α_{ij} and x_{ij} lie in the smooth loci of $(Z_i)_{\bar{k}}$ and $X_{\bar{k}}$, respectively.

We let $\Xi_i := \{x_{ij}\}_j \cup \Sigma_{\bar{k}}$ and $E_i := \{b_{ij}\}_j \cup F_{\bar{k}}$.

Applying Lemma 4.8 over \bar{k} for the above Ξ_i (for Σ there) and E_i (for E there), we obtain a dense open subset $\mathcal{U}'_i \subset \text{Gr}(X_{\bar{k}}, N - r - 1, H_{\bar{k}})$ such that every $L \in \mathcal{U}'_i(k)$ satisfies the properties (1)–(6) there. Repeating the argument of Lemma 4.8 in Step 2, we obtain a dense open subset $\mathcal{U}_i \subset \text{Gr}(X, N - r - 1, H)$ such that for every $L \in \mathcal{U}_i(k)$, we have $L_{\bar{k}} \in \mathcal{U}'_i(\bar{k})$.

We show that the following map is bijective:

$$\hat{\phi}_{L, \bar{k}} : \pi^{-1}(\alpha_i) \rightarrow \pi^{-1}(\beta_i). \tag{4-6}$$

Suppose this is not injective, i.e., for some $j < j'$, we have $\hat{\phi}_{L, \bar{k}}(\alpha_{ij}) = \hat{\phi}_{L, \bar{k}}(\alpha_{ij'})$. Then $b_{ij} = \hat{g}_{\bar{k}}(\alpha_{ij}) = \hat{g}_{\bar{k}}(\alpha_{ij'}) = b_{ij'}$. Since \bar{k} is algebraically closed, we can write $\alpha_{ij} = (x_{ij}, b_{ij})$ and $\alpha_{ij'} = (x_{ij'}, b_{ij'})$. Since $b_{ij} = b_{ij'}$ and $\alpha_{ij} \neq \alpha_{ij'}$, we must have $x_{ij} \neq x_{ij'}$.

But at the same time, we have

$$\hat{\phi}_{L, \bar{k}}(x_{ij}) = \hat{f}_{\bar{k}}(\hat{\phi}_{L, \bar{k}}(\alpha_{ij})) = \hat{f}_{\bar{k}}(\hat{\phi}_{L, \bar{k}}(\alpha_{ij'})) = \hat{\phi}_{L, \bar{k}}(x_{ij'}).$$

In particular, $x_{ij'} \in L_{\bar{k}}^+(x_{ij})$. Since $\hat{g}_{\bar{k}}(\alpha_{ij'}) = b_{ij'} \in E_i$, we thus have $\hat{f}_{\bar{k}}(\alpha_{ij'}) = x_{ij'} \in (L_{\bar{k}})^+(x_{ij}) \cap \hat{f}_{\bar{k}}(\hat{g}_{\bar{k}}^{-1}(E_i))$. But this contradicts property (5) of Lemma 4.8 satisfied by $L_{\bar{k}}$. Hence the map (4-6) is injective.

On the other hand, we have

$$\pi^{-1}(\beta_i) \times_{\text{Spec}(k(\beta_i))} \text{Spec}(k(\alpha_i)) = \text{Spec}((\bar{k} \otimes_k k(\beta_i)) \otimes_{k(\beta_i)} k(\alpha_i)) = \text{Spec}(\bar{k} \otimes_k k(\alpha_i)) = \pi^{-1}(\alpha_i)$$

so that it follows that the map (4-6) is surjective, as well, thus bijective.

Going back to the proof of the lemma, first note that we clearly have $Z_i \cap \hat{\phi}_L^{-1}(\beta_i) \supset \{\alpha_i\}$. For the inclusion in the other direction, toward contradiction suppose there is $\alpha' \in \hat{\phi}_L^{-1}(\beta_i) \setminus \{\alpha_i\}$ such that $\alpha' \in Z_i$. Clearly we have $\pi^{-1}(\alpha') \cap \pi^{-1}(\alpha_i) = \emptyset$. On the other hand, we have $\hat{\phi}_{L, \bar{k}}(\pi^{-1}(\alpha')) \subset \pi^{-1}(\beta_i) = \hat{\phi}_{L, \bar{k}}(\pi^{-1}(\alpha_i))$, where the second equality holds by the bijectivity of (4-6).

Hence there is some $\alpha'_j \in \pi^{-1}(\alpha')$ and $\alpha_{ij'} \in \pi^{-1}(\alpha_i)$ such that

$$(a) \ \alpha'_j \neq \alpha_{ij'}, \quad \text{while} \quad (b) \ \hat{\phi}_{L, \bar{k}}(\alpha'_j) = \hat{\phi}_{L, \bar{k}}(\alpha_{ij'}).$$

Property (b) implies that $\hat{g}_{\bar{k}}(\alpha'_j) = \hat{g}_{\bar{k}}(\alpha_{ij'}) = b_{ij'}$. Since \bar{k} is algebraically closed, for $x' := \hat{f}_{\bar{k}}(\alpha'_j)$, we can express $\alpha'_j = (x', b_{ij'})$ and $\alpha_{ij'} = (x_{ij'}, b_{ij'})$. Because $\alpha'_j \notin \alpha_{ij'}$ by (a), we must have $x' \neq x_{ij'} = \hat{f}_{\bar{k}}(\alpha_{ij'})$. In particular, $x' \in L_{\bar{k}}^+(x_{ij'})$. But $\hat{g}_{\bar{k}} = b_{ij'} \in E_i$ so that we obtain $x' \in L_{\bar{k}}^+(x_{ij'}) \cap \hat{f}_{\bar{k}}(\hat{g}_{\bar{k}}^{-1}(E_i))$. But, this

contradicts property (5) of Lemma 4.8 satisfied by $L_{\bar{k}}$. Hence no such α' exists. Our proof then is over by taking $\mathcal{U} := \bigcap_{i=1}^s \mathcal{U}_i$. \square

Combined with Lemma 4.11, we immediately have:

Corollary 4.13. *For each linear projection L as in Lemma 4.12 and each $1 \leq i \leq s$, one has that Z_i is the only irreducible component of $\hat{\phi}_L^{-1}(\hat{\phi}_L(Z_i))$ passing through a given marked point $\alpha_i \in Z_i \setminus \bigcup_{j \neq i} Z_j$.*

We can now prove the birationality of a given finite set of integral closed subschemes of $X \times \hat{B}_n$ under suitable linear projections.

Lemma 4.14. *For a suitable choice of the set E in the Set-up of Section 4D, after replacing the embedding $X \hookrightarrow \mathbb{P}_k^N$ by a bigger one via a Veronese embedding if necessary, there is a dense open subset $\mathcal{U} \subset \text{Gr}(X, N - r - 1, H)$ such that for each $L \in \mathcal{U}(k)$, the induced map $\hat{\phi}_L : Z_i \rightarrow \hat{\phi}_L(Z_i)$ is birational for all $1 \leq i \leq s$.*

Proof. We follow the choices of $\alpha_i \in Z_i$, Ξ and E that we made just before Lemma 4.12. We shall prove the lemma for this E . We let $\mathcal{U} \subset \text{Gr}(X, N - r - 1, H)$ be as given by Lemma 4.12 and fix $L \in \mathcal{U}(k)$. We let $T_i := \hat{\phi}_L(Z_i)$ and $\beta_i := \hat{\phi}_L(\alpha_i)$. To show that $\hat{\phi}_L : Z_i \rightarrow T_i$ is birational, we prove a stronger assertion that the map $\mathcal{O}_{T_i, \beta_i} \rightarrow \mathcal{O}_{Z_i, \beta_i}$ of semilocal rings is an isomorphism, where $\mathcal{O}_{Z_i, \beta_i} := \mathcal{O}_{Z_i, Z_i \cap \hat{\phi}_L^{-1}(\beta_i)}$. Consider the maps

$$\mathcal{O}_{T_i, \beta_i} \rightarrow \mathcal{O}_{Z_i, \beta_i} \rightarrow \mathcal{O}_{Z_i, \alpha_i}. \tag{4-7}$$

It follows from Lemma 4.12 that $Z_i \cap \hat{\phi}_L^{-1}(\beta_i) = \{\alpha_i\}$. In particular, the second map of (4-7) is an isomorphism, actually the identity map. By condition (1) of Lemma 4.8, the map ϕ_L is étale in an affine open neighborhood U' of Ξ , and thus ϕ_L is étale at α_i . In particular, the composite map in (4-7) is unramified. By condition (3) of Lemma 4.8, we have $k(\beta_i) \xrightarrow{\sim} k(\alpha_i)$. Hence, the first map of (4-7) is an injective finite unramified map of local rings, that induces an isomorphism of the residue fields. It is therefore an isomorphism by Lemma 4.10. This completes the proof. \square

4H. A presentation lemma for moving to fs-cycles. The final result of Section 4 is the following Theorem 4.15, that will be used in the proof of the fs-moving lemma, specifically, in the proof of Lemma 8.7.

Theorem 4.15. *Under the Set-up of Section 4D, let $Z_i^0 := Z_i|_{X \times B}$ and $Z^0 := Z|_{X \times B}$.*

Then for a suitable choice of the set E in the Set-up, after replacing the embedding $X \hookrightarrow \mathbb{P}_k^N$ by its composition with a suitable Veronese embedding, there is a dense open subset $\mathcal{U} \subset \text{Gr}(X, N - r - 1, H)$ such that each $L \in \mathcal{U}(k)$ satisfies the following:

- (1) ϕ_L is étale at Σ .
- (2) ϕ_L separates the points of Σ .
- (3) $k(\phi_L(x)) \xrightarrow{\sim} k(x)$ for all $x \in \Sigma$.
- (4) *There exists an affine open neighborhood $U \subset X$ of Σ such that:*

- (4a) If Z_i^0 is an irreducible component of Z^0 that is dominant over an irreducible component of X , then for each component Z' of $L^+(Z_i^0)$, the map $Z'_U \rightarrow U$ is fs over U .
- (4b) If Z_i^0 is an irreducible component of Z^0 that is not dominant over any irreducible component of X , then $L^+(Z_i^0)_U = 0$.

Proof. For $1 \leq i \leq s$, choose closed points $\alpha_i \in Z_i^0 \setminus (\bigcup_{j \neq i} Z_j^0)$ such that

$$x_i := \hat{f}(\alpha_i) \in X_{\text{sm}} \quad \text{and} \quad b_i := \hat{g}(\alpha_i) \in B,$$

as we did in Lemma 4.12. Let $\Xi := \{x_1, \dots, x_r\} \cup \Sigma \subset X_{\text{sm}}$ and $E := \{b_1, \dots, b_r\} \cup F \subset \hat{B}$. Since $Z_i \not\subset X \times F$ and $Z_i \rightarrow \hat{B}$ is nonconstant, it is not contained in $\hat{g}^{-1}(E)$. We choose $\mathcal{U} \subset \text{Gr}(X, N-r-1, H)$ as given by Lemma 4.12 and fix $L \in \mathcal{U}(k)$. In particular, all the properties of Lemma 4.8 holds, so that we have conditions (1)–(3) of the theorem.

To prove (4), first note that the irreducible components of $L^+(Z_i^0)$ are exactly the restrictions to $X \times B$ of the irreducible components of $L^+(Z_i)$. Let Z_i be an irreducible component of Z dominant over an irreducible component of X . Let Z' be an irreducible component of $L^+(Z_i)$. We prove that $Z' \cap (\Sigma \times F) = \emptyset$.

Suppose, on the contrary, that there is a closed point $\lambda \in Z' \cap (\Sigma \times F)$. This means that there is a closed point $\lambda' \in Z_i$ such that $\hat{\phi}_L(\lambda) = \hat{\phi}_L(\lambda')$. We claim in this case that

$$\{\lambda'\} = \hat{\phi}_L^{-1}(\hat{\phi}_L(\lambda)) \cap Z_i = \{\lambda\}. \tag{4-8}$$

Suppose we have shown that $\lambda' = \lambda$. Then we get $\lambda \in Z_i$ and (4-8) becomes equivalent to showing that $\hat{\phi}_L^{-1}(\hat{\phi}_L(\lambda)) \cap Z_i = \{\lambda\}$. But the proof of this equality is simply a repetition of the argument of Lemma 4.12. Hence, the claim is reduced to showing that $\lambda' = \lambda$.

Let's do it. First consider the case when k is algebraically closed. We can then uniquely write $\lambda = (x, b)$ for some closed points $x \in \Sigma$ and $b \in F$, and $\lambda' = (x', b)$, where $x' \in \hat{\phi}_L^{-1}(\hat{\phi}_L(x))$. If $x' \neq x$, then $x' \in L^+(x)$ and $x' \in \hat{f}(\hat{g}^{-1}(E))$, which contradicts condition (5) of Lemma 4.8. Hence, we must have $x' = x$ so that $\lambda' = \lambda$.

If k is not algebraically closed, we argue as in the proof of Lemma 4.12. Suppose again that $\lambda' \neq \lambda$. Then for the base change map $\pi : \text{Spec}(\bar{k}) \rightarrow \text{Spec}(k)$, we have $\pi^{-1}(\lambda') \cap \pi^{-1}(\lambda) = \emptyset$. Let $\beta' := \hat{\phi}_L(\lambda') = \hat{\phi}_L(\lambda)$. We show as in the argument of Lemma 4.12 that the map $\hat{\phi}_{L, \bar{k}} : \pi^{-1}(\lambda') \rightarrow \pi^{-1}(\beta')$ is bijective. Using this, we continue following the proof of Lemma 4.12, to get closed points $\tilde{\lambda} \in \pi^{-1}(\lambda)$ and $\tilde{\lambda}' \in \pi^{-1}(\lambda')$ such that $\hat{f}_{\bar{k}}(\tilde{\lambda}) \in (L_{\bar{k}})^+(\hat{f}_{\bar{k}}(\tilde{\lambda}')) \cap \hat{f}_{\bar{k}}(\hat{g}_{\bar{k}}^{-1}(E_i))$. But this contradicts property (5) of Lemma 4.8 for $L_{\bar{k}}$, which violates our choice of L . This proves (4-8).

Coming back to the proof of $Z' \cap (\Sigma \times F) = \emptyset$, we now note using Corollary 4.13 that $Z' \neq Z_i$. So, the two deductions $\lambda \in Z' \cap Z_i$ and $\hat{\phi}_L^{-1}(\hat{\phi}_L(\lambda)) \cap Z_i = \{\lambda\}$ from (4-8) together contradict Lemma 4.11. Hence, we must have $Z' \cap (\Sigma \times F) = \emptyset$, as desired.

Now, by Lemma 2.10, there is an affine open neighborhood $U_{i,Z'} \subset X_{\text{sm}}$ of Σ such that $Z'_{U_{i,Z'}} \rightarrow U_{i,Z'}$ is fs. We take $U_1 := \bigcap U_{i,Z'}$ where the intersection is taken over all i such that Z_i dominant over a component of X and the irreducible components Z' . This open set U_1 works for (4a).

About property (4b), let Z_i be an irreducible component of Z which is not dominant over X . Let Z' be a component of $L^+(Z_i)$. In this case, we repeat the proof of (4a) above, where we now apply condition (6) of Lemma 4.8, to conclude that $Z' \cap (\Sigma \times \hat{B}) = \emptyset$.

It follows that $\hat{f}(L^+(Z_i))$ is a closed subset of X disjoint from Σ . Hence, we can apply Lemma 2.3 to obtain an affine open neighborhood U'_i of Σ in X such that $L^+(Z_i)_{U'_i} = \emptyset$. We take $U_2 := \bigcap U'_i$, where the intersection is taken over all i such that Z_i is not dominant over any component of X . This open set U_2 works for (4b). Taking $U := U_1 \cap U_2$, we have (4), and this concludes the proof of the theorem. \square

5. Regularity of the original cycle over residual points

The focus of the remaining sections is to achieve the sfs-property of the residual cycle of Z along Σ via more refined linear projections. In order to achieve this, we first ensure that our original cycle Z is regular at all points lying over the residual set $L^+(\Sigma)$ of $\Sigma \subset X$. We later show that this regularity of Z at all points lying over $L^+(\Sigma)$ implies the regularity of the residual cycle of Z along Σ . The goal of this section is to achieve the first one when k is algebraically closed. The general case will be considered later.

5A. A basic algebraic result. We first discuss the following:

Lemma 5.1. *Let k be an algebraically closed field. Let $X \subset \mathbb{P}_k^N$ be a reduced closed subscheme of dimension 1. Suppose $N \gg 1$ and let $x \neq y$ be two closed points on X_{sm} . Let $\text{Gr}_{x+2y}(N-1, \mathbb{P}_k^N) \subset \text{Gr}(N-1, \mathbb{P}_k^N)$ be the set of hyperplanes containing $\{x, y\}$ that do not intersect X transversely at y . Then $\text{Gr}_{\{x,y\}}(N-1, \mathbb{P}_k^N) \simeq \mathbb{P}_k^{N-2}$ and $\text{Gr}_{x+2y}(N-1, \mathbb{P}_k^N) \simeq \mathbb{P}_k^{N-3}$.*

Proof. Recall that $\text{Gr}_{\{x,y\}}(N-1, \mathbb{P}_k^N) \subset \text{Gr}(N-1, \mathbb{P}_k^N)$ is the set of hyperplanes containing $\{x, y\}$. Since $x \neq y$, by elementary linear algebra on ranks of linear systems, we immediately have $\text{Gr}_{\{x,y\}}(N-1, \mathbb{P}_k^N) \simeq \mathbb{P}_k^{N-2}$. We prove the second assertion. Since $N \gg 1$, we can find a linear form $s_1 \in W = H^0(\mathbb{P}_k^N, \mathcal{O}(1))$ which does not vanish anywhere in $\{x, y\}$. This yields a k -linear map $\alpha : W \rightarrow \mathcal{O}_{X,\{x,y\}}/\mathfrak{m}_x \mathfrak{m}_y^2 =: \mathcal{O}_{\{x+2y\}}$ given by $\alpha(s) = s/s_1$. Since k is algebraically closed, the ideal \mathfrak{m}_y is generated by linear forms vanishing at y . Hence, the composite map $W \rightarrow \mathcal{O}_{X,\{x,y\}}/\mathfrak{m}_x \mathfrak{m}_y^2 \rightarrow \mathcal{O}_{X,y}/\mathfrak{m}_y^2 =: \mathcal{O}_{\{2y\}}$ is surjective and $\alpha^{-1}(\mathfrak{m}_y^2)$ is precisely the set of linear forms in W not transverse to X at y .

We first claim that α is surjective. Since x, y are two distinct regular closed points of X , the set $\text{Gr}_y(x, N-1, \mathbb{P}_k^N)$ is nonempty and hence, $\mathfrak{m}_y/\mathfrak{m}_x \mathfrak{m}_y \xrightarrow{\sim} \mathcal{O}_{\{x\}}$ and there is a commutative diagram of short exact sequences:

$$\begin{array}{ccccccc}
 0 & \longrightarrow & \alpha^{-1}(\mathfrak{m}_x \mathfrak{m}_y) & \longrightarrow & \alpha^{-1}(\mathfrak{m}_y) & \longrightarrow & \mathcal{O}_{\{x\}} \longrightarrow 0 \\
 & & \downarrow & & \downarrow & & \parallel \\
 0 & \longrightarrow & \mathfrak{m}_x \mathfrak{m}_y / \mathfrak{m}_x \mathfrak{m}_y^2 & \longrightarrow & \mathfrak{m}_y / \mathfrak{m}_y^2 & \longrightarrow & \mathcal{O}_{\{x\}} \longrightarrow 0
 \end{array} \tag{5-1}$$

In particular, the first vertical map is surjective. Since $\text{Gr}_x(y, N - 1, \mathbb{P}_k^N) \neq \emptyset$, we conclude that α is surjective.

To finish the proof, we look at the commutative diagram with exact rows:

$$\begin{array}{ccccccc}
 0 & \longrightarrow & \ker(\alpha) & \longrightarrow & W & \xrightarrow{\alpha} & \mathcal{O}_{\{x+2y\}} \longrightarrow 0 \\
 & & \downarrow & & \parallel & & \downarrow \\
 0 & \longrightarrow & \alpha^{-1}(\mathfrak{m}_y^2) & \longrightarrow & W & \longrightarrow & \mathcal{O}_{\{2y\}} \longrightarrow 0
 \end{array} \tag{5-2}$$

Since the last vertical arrow is surjective with one-dimensional kernel, by the snake lemma, the first vertical arrow is injective with one-dimensional cokernel. Since $\mathbb{P}(\alpha^{-1}(\mathfrak{m}_y^2)) \simeq \mathbb{P}_k^{N-2}$, we conclude that $\text{Gr}_{x+2y}(N - 1, \mathbb{P}_k^N) \simeq \mathbb{P}(\ker(\alpha)) \simeq \mathbb{P}_k^{N-3}$. \square

5B. The Set-up+(fs). We suppose k is an infinite perfect field. The Set-up we now use repeatedly is the following situation, that we call the Set-up+(fs):

- (1) *The Set-up:* We still suppose the Set-up of Section 4D, not necessarily specifying some closed subset $E \subset \hat{B}$.
- (2) *The fs-property:* There exists an affine open neighborhood $X_{\text{fs}} \subset X_{\text{sm}}$ of Σ , that is dense open in X , such that the projection $Z \rightarrow X$ is fs over X_{fs} .

5C. Regularity of the original cycle over residual points. We now discuss two central results: Lemmas 5.2 and 5.10. Recall that X is equidimensional under the above assumptions.

Lemma 5.2. *Let k be an algebraically closed field. Suppose $r = 1$. We are under the Set-up+(fs) of Section 5B. Let $x \in X_{\text{fs}}$ be a closed point and let $S \subset X \setminus \{x\}$ be another finite set of closed points.*

After replacing \mathbb{P}_k^N by a bigger projective space via a Veronese embedding if necessary, there exists a dense open subset $\mathcal{U}_S \subset \text{Gr}_x(N - 1, \mathbb{P}_k^N)$ such that each $L \in \mathcal{U}_S(k)$ satisfies the following:

- (1) $L \cap ((X \setminus X_{\text{fs}}) \cup S) = \emptyset$.
- (2) L intersects X_{fs} transversely.
- (3) $L \cap X$ consists of $(d + 1)$ -distinct closed points $c_0 = x, c_1, \dots, c_d$.
- (4) Z is regular at all points lying over $\{c_1, \dots, c_d\}$. In particular, each component Z_i does not meet other irreducible components at points lying over $\{c_1, \dots, c_d\}$.

Proof. Since $\dim(Z_{\text{sing}}) = 0$, we see that $\hat{f}(Z_{\text{sing}})$ is a finite closed subset of X . Since X_{fs} is dense in X , we have $|X \setminus X_{\text{fs}}| < \infty$. Hence, $T := (\hat{f}(Z_{\text{sing}}) \cup (X \setminus X_{\text{fs}}) \cup S) \setminus \{x\}$ is a finite closed subset of X . Thus the hyperplanes disjoint from T form a dense open subset $\text{Gr}(T, N - 1, \mathbb{P}_k^N)$ of $\text{Gr}(N - 1, \mathbb{P}_k^N)$ by Lemma 4.2. The set $\mathcal{U}_1 := \text{Gr}^{\text{tr}}(X, N - 1, \mathbb{P}_k^N) \cap \text{Gr}(T, N - 1, \mathbb{P}_k^N)$ is dense open in $\text{Gr}(N - 1, \mathbb{P}_k^N)$. If we show that $\mathcal{U}_S := \mathcal{U}_1 \cap \text{Gr}_x(N - 1, \mathbb{P}_k^N) \neq \emptyset$, then this set will be dense open in $\text{Gr}_x(N - 1, \mathbb{P}_k^N)$. It is moreover clear that any $L \in \mathcal{U}_S(k)$ satisfies (1)–(4). It remains to show that $\text{Gr}^{\text{tr}}(X, N - 1, \mathbb{P}_k^N) \cap \text{Gr}_x(N - 1, \mathbb{P}_k^N)$ and $\text{Gr}(T, N - 1, \mathbb{P}_k^N) \cap \text{Gr}_x(N - 1, \mathbb{P}_k^N)$ are both nonempty.

Let V be the set of linear forms in $H^0(\mathbb{P}_k^N, \mathcal{O}(1))$ that vanish at x . Note that $\dim|V| = N - 1$ and that the maximal ideal $\mathfrak{m}_x \subset \mathcal{O}_{X,x}$ is generated by the members of V . Let $\mathcal{B} \subset X \times |V|$ be the incidence scheme consisting of pairs (y, L) such that L passes through y , but not transverse to X at y . We study the fiber of $\pi_1 : \mathcal{B} \rightarrow X$ over each $y \in X_{\text{sm}} \setminus \{x\}$.

Choose $s_1 \in V$ such that $s_1(x) = 0$ but $s_1(y) \neq 0$. Consider the map $\beta : V \rightarrow \mathcal{O}_{X,y}/\mathfrak{m}_y^2$ given by $\beta(s) = s/s_1$. Since $\dim|V| = N - 1$, while $\text{Gr}_{\{x,y\}}(N - 1, \mathbb{P}^N) \simeq \mathbb{P}^{N-2}$ and $\text{Gr}_{\{x+2y\}}(N - 1, \mathbb{P}^N) \simeq \mathbb{P}^{N-3}$ by at most $N - 3$, because $\dim_k(\mathcal{O}_{X,y}/\mathfrak{m}_y^2) = 2$. Thus, $\dim(\mathcal{B}) \leq \dim X + \dim(\pi_1^{-1}(y)) \leq 1 + N - 3 = N - 2$. Hence its image in $|V|$ under the projection $\pi_2 : X \times |V| \rightarrow |V|$ is a proper closed subset (note that X is projective). Since $N \gg 0$, its complement $\text{Gr}_x^{\text{tr}}(X, N - 1, \mathbb{P}_k^N)$ in $\text{Gr}_x(N - 1, \mathbb{P}_k^N)$ is a dense open subset. Since $\dim(\text{Gr}_x(N - 1, \mathbb{P}_k^N)) = N - 1$ and $T \subset X$ is a finite set of closed points different from x , the assertion that $\text{Gr}(T, N - 1, \mathbb{P}_k^N) \cap \text{Gr}_x(N - 1, \mathbb{P}_k^N)$ is nonempty follows from Lemma 5.1. We have therefore finished the proof. \square

In Section 6A, we will obtain a slightly stronger version of Lemma 5.2. This is done in Lemma 5.9. The difference in the latter lemma from the former is that (following the notations of Lemma 5.2), after a possible reembedding, we may impose an additional property that for $L \cap X = \{c_0 = x, c_1, \dots, c_d\}$, no three points of them are collinear.

At one bad extreme case, suppose X is contained in a 2-dimensional projective space. Then for any hyperplane L , which is a line, the hyperplane section $L \cap X$ is entirely collinear. This is an important obstacle to avoid. We will show in Lemma 5.7 that, after taking a Veronese reembedding for a high enough degree $d \geq 3$, we can always avoid it. It will be improved for the higher dimensional case in Lemma 5.8. These two are some technical grounds needed in Section 6A.

Once we can avoid the above extreme case using a Veronese reembedding, then one can employ the following well-known general result (see [Arbarello et al. 1985, Chapter III, page 109]):

Theorem 5.3 (general position theorem). *Let $N \geq 2$. Let $C \subset \mathbb{P}^N$ be an irreducible nondegenerate, possibly singular, curve of degree d . Then a general hyperplane meets C in d points, any N of which are linearly independent.*

Recall that a closed embedding $X \subset \mathbb{P}_k^n$ of an integral projective scheme X is said to be nondegenerate if no hyperplane of \mathbb{P}_k^n contains X . We won't give the proof of Theorem 5.3 here. We mention that Theorem 5.3 for $N = 2$ is immediate, while, for $N \geq 3$ reduces to the following special case (see [loc. cit.]), that is more relevant to the paper:

Lemma 5.4. *Let $C \subset \mathbb{P}^N$ with $N \geq 3$ be an irreducible nondegenerate, possibly singular, curve of degree d . Then a general hyperplane meets C in d points, no three of which are collinear.*

Remark 5.5. To give a bit of the flavor of the proof of Lemma 5.4, we remark that with some efforts (see [Arbarello et al. 1985, pages 110–111] or imitate [Hartshorne 1977, Proposition IV-3.8, page 311]), one can argue that if Lemma 5.4 fails, then all tangent lines to C passes through a single fixed point $p \in C$. Then a linear projection from p would shrink the entire curve C to a point in \mathbb{P}^{N-1} . Since C is nondegenerate, we can argue this cannot happen.

Such a curve in \mathbb{P}^N all of whose tangent lines pass through a fixed point is called *strange* (see [Hartshorne 1977, page 311]). We remark that in case C is nonsingular, it is known that the only nonsingular strange curves in any \mathbb{P}^N are either a line or a conic in \mathbb{P}^2 in characteristic 2 (see [Samuel 1966, Theorem, Appendix to Chapter II, page 76] or [Hartshorne 1977, Theorem IV-3.9, page 312]).

We thank the referee for pointing to us that some technical part of our construction of the paper is relevant to noncollinearity of configurations of points and strange curves. \square

Combined with the Bertini theorem ([Kleiman and Altman 1979, Theorem 1] or [Jouanolou 1983]), we immediately extend Lemma 5.4 to the following higher dimensional version, which we use:

Proposition 5.6 (linear general position theorem). *Let $X \subset \mathbb{P}^N$ with $N \geq 3$ be a nondegenerate, possibly singular, variety of degree d . Let $r = \dim X \geq 1$. Then for a general sequence of hyperplanes H_1, \dots, H_r in \mathbb{P}^N , the intersection $X \cap H_1 \cap \dots \cap H_r$ has d points, no three of which are collinear.*

Note that the above Proposition 5.6 holds for schemes that are nondegenerate in the projective spaces of dimension at least 3. This is another view of why we had a pathology about noncollinearity when X was contained in a 2-dimensional projective space in the paragraph before Theorem 5.3.

As said before, to avoid this problem, we need to replace the embedding by a bigger Veronese embedding. This is discussed now in the following:

Lemma 5.7. *Let $C \subset \mathbb{P}_k^n$ be a reduced projective curve. Suppose that there exists a 2-dimensional linear subspace $L \subset \mathbb{P}_k^n$ such that $C \subset L$. Let $\vartheta : \mathbb{P}_k^n \hookrightarrow \mathbb{P}_k^N$ be the d -uple Veronese embedding with $d \geq 3$. Then the image of each irreducible component of C via ϑ does not lie inside a 2-dimensional linear subspace of \mathbb{P}_k^N .*

Proof. We can assume C is an irreducible curve in order to prove the lemma. After a linear change of coordinates in \mathbb{P}_k^n , we may assume that $\mathbb{P}_k^n = \mathbb{P}(V)$ and $L = \mathbb{P}(W)$, where V is an $(n+1)$ -dimensional k -vector space with a basis $\{x_0, \dots, x_n\}$ and $W = \text{Span}_k\{x_0, x_1, x_2\}$ is a subspace of V . For any closed embedding $f : C \hookrightarrow \mathbb{P}_k^m$, we let $d_f(C)$ denote the degree of C under f .

Let $\iota : C \hookrightarrow L$ be the closed embedding as given in the assumption of the lemma. Let $d_0 := d_\iota(C) \geq 1$. Since L is linear in \mathbb{P}_k^n , the degree of C under the composite of the embeddings $C \hookrightarrow L \hookrightarrow \mathbb{P}_k^n$ is also d_0 .

Toward contradiction, suppose that there is a 2-dimensional linear subspace $L' \subset \mathbb{P}_k^N$ such that $\vartheta(C) \subset L'$, where $\vartheta : \mathbb{P}_k^n \hookrightarrow \mathbb{P}_k^N$ is the d -uple Veronese embedding with $d \geq 3$. We denote the resulting embedding $C \hookrightarrow L'$ by $\vartheta|_C$.

By our choice of the embedding $L \hookrightarrow \mathbb{P}_k^n$, we have a commutative diagram

$$\begin{array}{ccccc}
 C & \xrightarrow{\iota} & L & \hookrightarrow & \mathbb{P}_k^n \\
 & \searrow & \downarrow \vartheta' & & \downarrow \vartheta \\
 & & M & \hookrightarrow & \mathbb{P}_k^N
 \end{array} \tag{5-3}$$

where $M \cong \mathbb{P}_k^r$ (with $r = (d+1)(d+2)/2 - 1$). The horizontal arrows in the right square are linear embeddings and the vertical arrows are the d -uple Veronese embeddings.

The linearity of the inclusion $L' \hookrightarrow \mathbb{P}_k^N$ implies that $d_{\vartheta|_C}(C)$ coincides with the degree of C under the composite closed embedding $C \hookrightarrow L' \hookrightarrow \mathbb{P}_k^N$. By the same argument, the degree of C for this composite embedding coincides with the degree of C for the composite embedding $C \hookrightarrow L \hookrightarrow M$. Since ϑ' is the d -uple Veronese embedding, it follows that the degree of C for the latter composite embedding is d_0d . We conclude that $d_{\vartheta|_C}(C) = d_0d$.

If we now apply the degree-genus adjunction formula for plane curves to the embedding ι , we get $g_a(C) = \frac{1}{2}(d_0 - 1)(d_0 - 2)$, where $g_a(C)$ is the arithmetic genus of C . The same formula for the embedding $\vartheta|_C$ yields $g_a(C) = \frac{1}{2}(d_0d - 1)(d_0d - 2)$.

Hence $(d_0 - 1)(d_0 - 2) = (d_0d - 1)(d_0d - 2)$, i.e., $d_0^2(d^2 - 1) - 3d_0(d - 1) = 0$. This factors into

$$d_0(d - 1)(d_0(d + 1) - 3) = 0. \tag{5-4}$$

Since $d_0 \geq 1$ and $d \geq 3$, the left hand side of (5-4) is $\geq 1 \cdot 2 \cdot (1 \cdot 4 - 3) > 0$, so that the equality of (5-4) cannot hold, thus a contradiction. This proves the lemma. \square

An analogue of Lemma 5.7 in higher dimensions is the following.

Lemma 5.8. *Let $\iota : X \hookrightarrow \mathbb{P}_k^n$ be a reduced projective scheme of pure dimension $r \geq 2$. Assume that the degree of each irreducible component of X in \mathbb{P}_k^n is at least two. Let $\Sigma \subset X$ be a finite set of closed points. For an integer $d \geq 1$, let $\vartheta : \mathbb{P}_k^n \hookrightarrow \mathbb{P}_k^N$ be the d -uple Veronese embedding.*

Then for all sufficiently large $d \geq 3$, (depending on X, Σ, n and the degrees of the irreducible components of X in \mathbb{P}_k^n), a general intersection $H_1 \cap \dots \cap H_{r-1} \cap \vartheta(X)$ of X with hyperplanes H_i in $\text{Gr}_\Sigma(N - 1, \mathbb{P}_k^N)(k)$ is a reduced curve, none of whose irreducible component is contained in a 2-dimensional linear subspace of \mathbb{P}_k^N .

Proof. By the Bertini theorems of Kleiman and Altman [1979, Theorem 1], an intersection of $\vartheta(X)$ with $(r - 1)$ general hyperplanes containing Σ in a large enough d -uple Veronese embedding ϑ is a curve C , whose intersection with every irreducible component of $\vartheta(X)$ is again irreducible. Since k is perfect and X is reduced, it is actually geometrically reduced. It follows therefore from the Bertini theorem of Jouanolou [1983, Théorème 6.3] that C can be chosen to be reduced.

Let X_1, \dots, X_t be the irreducible components of X and let C_1, \dots, C_t denote the irreducible components of C .

Let s_i be the degree of X_i in \mathbb{P}_k^n so that the degree of X in \mathbb{P}_k^n is $s = \sum_{i=1}^r s_i$ (see [Hartshorne 1977, Proposition I-7.6, page 52]). Let $C = H_1 \cap \dots \cap H_{r-1} \cap \vartheta(X)$ be as above. Let $d_i(C_i)$ denote the degree of C_i in \mathbb{P}_k^n via the inclusion $\iota : C \hookrightarrow X \hookrightarrow \mathbb{P}_k^n$ and let $d_\vartheta(C_i)$ denote the degree of C_i in \mathbb{P}_k^N . Each of the hyperplanes $H_1, \dots, H_{r-1} \subset \mathbb{P}_k^N$ restricts to a unique hypersurface of degree d in \mathbb{P}_k^n . Since these hyperplanes are sufficiently general, an elementary degree computation shows that $d_i(C_i) = d^{r-1}s_i$ and $d_\vartheta(C_i) = d^r s_i$ for each $1 \leq i \leq t$. We need to show that if d is sufficiently large, then each $C_i = H_1 \cap \dots \cap H_{r-1} \cap \vartheta(X_i)$ is not contained in a 2-dimensional linear subspace of \mathbb{P}_k^N . To show this, we can assume that X and C are irreducible. In particular, $d_i(C) = sd^{r-1}$ and $d_\vartheta(C) = sd^r$.

We shall prove our assertion as an application of Castelnuovo’s bound for the genus of curves. Let $3 \leq n' \leq n$ be the smallest integer such that $X \subset \mathbb{P}_k^{n'} \subset \mathbb{P}_k^n$, where the first embedding is nondegenerate and the second embedding is linear. Note that the lower bound on n' is forced by our assumption on the lower bounds of the dimension of X and its degree in \mathbb{P}_k^n .

Since H_1, \dots, H_{r-1} restrict to general hypersurfaces of degree d in \mathbb{P}_k^n , we see that they restrict to hypersurfaces of the same degree in $\mathbb{P}_k^{n'}$. Since a hypersurface (of degree at least two) section of a nondegenerate closed subvariety of a projective space is necessarily nondegenerate (looking at the homogeneous coordinate rings), we conclude that the composite embedding $C \hookrightarrow X \hookrightarrow \mathbb{P}_k^{n'}$ is also nondegenerate. Furthermore, the degrees of X and C inside $\mathbb{P}_k^{n'}$ are the same as their respective degrees inside \mathbb{P}_k^n .

Let $m \geq 1$ and $0 \leq \epsilon < n' - 1$ be two integers such that $sd^{r-1} - 1 = m(n' - 1) + \epsilon$. It follows from Castelnuovo’s bound on the arithmetic genus (see [Harris 1982, Chapter 3; Arbarello et al. 1985, Chapter III, page 116] and see [Ballico 1989, Remark following Lemma 2.1] for singular curves) of C that

$$g_a(C) \leq \frac{(n' - 1)m(m - 1)}{2} + m\epsilon. \tag{5-5}$$

Since $n' - 1 \geq 2$ and d is sufficiently large, we can assume $m < sd^{r-1} - 1$. We thus get

$$\begin{aligned} 2g_a(C) &\leq (n' - 1)m(m - 1) + 2m\epsilon \\ &< (n' - 1)m(m - 1) + 2m(n' - 1) \\ &= (n' - 1)m(m + 1) \\ &\leq (sd^{r-1} - 1)(sd^{r-1} - 1) \\ &= (sd^{r-1} - 1)^2. \end{aligned} \tag{5-6}$$

Now toward contradiction, suppose that inside \mathbb{P}_k^N , the curve C is contained in a 2-dimensional linear subspace $L \subset \mathbb{P}_k^N$. Since $d_\vartheta(C)$ is equal to the degree of C inside L , the degree-genus adjunction formula for the embedding $C \hookrightarrow L \cong \mathbb{P}_k^2$, yields $2g_a(C) = (sd^r - 1)(sd^r - 2)$. Note that if we let $e' := sd^{r-1} - 1$, then

$$\begin{aligned} 2g_a(C) &= (sd^r - 1)(sd^r - 2) \\ &= (d(sd^{r-1} - 1) + d - 1)(d(sd^{r-1} - 1) + d - 2) \\ &= d^2(e')^2 + (2d - 3)d(e') + (d - 1)(d - 2), \end{aligned} \tag{5-7}$$

and because $d \geq 3$ and $s > 0$, we have $2g_a(C) > (e')^2 + e' + 0 \geq (e')^2$.

On the other hand, from (5-6) we had $2g_a(C) \leq (e')^2$. This is a contradiction. □

We now present the aforementioned improvement of Lemma 5.2.

Lemma 5.9. *Let $X \hookrightarrow \mathbb{P}_k^N$ and $x \in X_{\text{fs}}$ be as in Lemma 5.2. After replacing \mathbb{P}_k^N by a bigger projective space via a Veronese embedding, there exists a dense open $\mathcal{U}_S \subset \text{Gr}_x(N - 1, \mathbb{P}_k^N)$ such that every $L \in \mathcal{U}_S(k)$ satisfies the following:*

- (1) *Conditions (1)–(4) of Lemma 5.2.*
- (2) *No three points of $L \cap X = \{x = c_0, c_1, \dots, c_d\}$ are collinear.*

Proof. Suppose first that X does not lie inside any 2-dimensional linear subspace of \mathbb{P}_k^N . In this case, we choose \mathcal{U}_S just as in Lemma 5.2 so that (1) holds. Condition (2) holds by Proposition 5.6. Hence the lemma is proven in this case.

Suppose now that X lies inside a 2-dimensional linear space of \mathbb{P}_k^N . In this case, we choose a suitable Veronese embedding $\mathbb{P}_k^N \hookrightarrow \mathbb{P}_k^{N'}$ such that the image of each irreducible component of X does not lie in any 2-dimensional linear subspace of $\mathbb{P}_k^{N'}$ applying Lemma 5.7. Then after reembedding if necessary, we have a nonempty open subset $\mathcal{U}_S \subset \text{Gr}_x(N' - 1, \mathbb{P}_k^{N'})$ such that conditions (1)–(4) of Lemma 5.2 hold.

In doing so, we can make sure that X is nondegenerate in a projective space of dimension at least 3. Then condition (1) holds by the choice of \mathcal{U}_S , while condition (2) holds by Proposition 5.6. This proves the lemma. \square

The following result generalizes Lemma 5.9 to higher dimensional $r \geq 1$.

Lemma 5.10. *Let k be an algebraically closed field. Suppose $r \geq 1$. We are under the Set-up+(fs) of Section 5B. Let $x \in X_{\text{fs}}$ be a closed point and let $S \subset X \setminus \{x\}$ be another finite set of closed points.*

After replacing \mathbb{P}_k^N by a bigger projective space via a Veronese embedding if necessary, we have the following property: given any hyperplane $H_0 \subset \mathbb{P}_k^N$ disjoint from $S \cup \{x\}$ and a general $L_0 \in \text{Gr}_{S \cup \{x\}}^{\text{tr}}(H_0, N - r + 1, \mathbb{P}_k^N)(k)$, there exists a dense open subset $\mathcal{U}_S \subset \text{Gr}_x^{\text{tr}}(L_0, N - 1, \mathbb{P}_k^N)$ such that each $L \in \mathcal{U}_S(k)$ satisfies the following:

- (1) $L \cap L_0 \cap ((X \setminus X_{\text{fs}}) \cup S) = \emptyset$.
- (2) $L \cap L_0$ intersects X_{fs} transversely.
- (3) $L \cap L_0 \cap X$ has $(d+1)$ -distinct closed points $c_0 = x, c_1, \dots, c_d$.
- (4) Z is regular at all points lying over $\{c_1, \dots, c_d\}$. In particular, each component Z_i does not meet other irreducible components at points lying over $\{c_1, \dots, c_d\}$.
- (5) $L_0 \cap X$ is an equidimensional reduced curve none of whose irreducible component lies inside a 2-dimensional linear subspace of \mathbb{P}_k^N .
- (6) No three points of $L \cap L_0 \cap X = \{x = c_0, c_1, \dots, c_d\}$ are collinear.

Proof. In case $r = 1$, we have $\text{Gr}(N - r + 1, \mathbb{P}_k^N) = \text{Gr}(N, \mathbb{P}_k^N) = \{\mathbb{P}_k^N\}$ so that $L_0 = \mathbb{P}_k^N$ and Lemma 5.10 follows from Lemmas 5.7 and 5.9. Hence we may assume $r \geq 2$. Let X_1, \dots, X_t be the irreducible components of X .

We saw in the proof of Lemma 5.8 that the Bertini theorems of Kleiman and Altman [1979, Theorem 1] and Jouanolou [1983] imply that an intersection of X with $(r-1)$ general hyperplanes containing $S \cup \{x\}$ in a large enough Veronese embedding of \mathbb{P}_k^N is a reduced curve C whose intersection with every irreducible component of X is irreducible. This curve C contains $S \cup \{x\}$. We can also ensure that no component of C is contained in $\hat{f}(Z_{\text{sing}}) \cup (X \setminus X_{\text{fs}})$, it is regular at points away from X_{sing} , and for each component of $Z|_{C \times \hat{B}}$, its projection to \hat{B} is nonconstant.

Hence, after replacing the embedding $\eta : X \hookrightarrow \mathbb{P}_k^N$ by its composition with a Veronese embedding of \mathbb{P}_k^N , we can find an $(r-1)$ -tuple of general hyperplanes (H_1, \dots, H_{r-1}) , each in $\text{Gr}_{S \cup \{x\}}(N-1, \mathbb{P}_k^N)$, such that the linear subspace $L_0 = H_1 \cap \dots \cap H_{r-1}$ has the following properties:

- (a) L_0 is transverse to H_0 .
- (b) $C = L_0 \cap X$ is a reduced curve none of whose components lies in $\hat{f}(Z_{\text{sing}}) \cup (X \setminus X_{\text{fs}})$.
- (c) $C \cap X_i$ is irreducible for each $1 \leq i \leq t$.
- (d) C is regular at points away from X_{sing} .
- (e) For each component of $Z|_{C \times \hat{B}}$, the projection to \hat{B} is nonconstant.

Let $S' := (C \setminus \{x\}) \cap (\hat{f}(Z_{\text{sing}}) \cup (X \setminus X_{\text{fs}}) \cup S)$, which is a finite closed subset of C .

Note from the definition of the degree of the embedding $\eta : X \hookrightarrow \mathbb{P}_k^N$ that a general hyperplane inside L_0 will intersect C at $(d+1)$ distinct closed points. Applying Lemma 5.9 to the curve C , the finite set S' , and $L_0 \simeq \mathbb{P}_k^{N-r+1}$ (which is regarded as the ambient projective space for C), there exists a dense open subset $\mathcal{U}_{C,S'} \subset \text{Gr}_x(N-r, L_0)$ that satisfies the assertions (1)–(2) of Lemma 5.9. Note that as $N \gg r$, the subset $\text{Gr}^{\text{tr}}(L_0, N-1, \mathbb{P}_k^N)$ is dense open in $\text{Gr}(N-1, \mathbb{P}_k^N)$.

Consider the regular map

$$\theta_{L_0} : \text{Gr}^{\text{tr}}(L_0, N-1, \mathbb{P}_k^N) \rightarrow \text{Gr}(N-r, L_0), \tag{5-8}$$

given by $\theta_{L_0}(L) = L \cap L_0$.

One checks that θ_{L_0} is a surjective smooth morphism of relative dimension $r-1$. Since θ_{L_0} is a smooth and surjective morphism such that $\theta_{L_0}^{-1}(\text{Gr}_x(N-r, L_0)) = \text{Gr}_x^{\text{tr}}(L_0, N-1, \mathbb{P}_k^N)$, we see that $\mathcal{U}_S := \theta_{L_0}^{-1}(\mathcal{U}_{C,S'})$ is a dense open subset of $\text{Gr}_x^{\text{tr}}(L_0, N-1, \mathbb{P}_k^N)$.

We want to show that each $L \in \mathcal{U}_S(k)$ satisfies the desired conditions (1)–(4). This is a tautology, but let us write it in detail: suppose $L \in \mathcal{U}_S(k)$, i.e., $\theta_{L_0}(L) \cap S' = \emptyset$ and $\theta_{L_0}(L) = L \cap L_0$ satisfies (1)–(4) with Z replaced by $Z|_{C \times \hat{B}}$. Since $\theta_{L_0}(L) \cap ((X \setminus X_{\text{fs}}) \cup S) = L \cap (L_0 \cap X) \cap ((X \setminus X_{\text{fs}}) \cup S) = \theta_{L_0}(L) \cap C \cap ((X \setminus X_{\text{fs}}) \cup S) \subset \theta_{L_0}(L) \cap S'$, and since $x \in X_{\text{fs}}$, we see that $\theta_{L_0}(L) \cap ((X \setminus X_{\text{fs}}) \cup S) = \emptyset$, proving (1).

Since L intersects L_0 transversely, which in turn intersects X transversely along X_{sm} by (b) and (d) above, we see that $\theta_{L_0}(L)$ intersects X transversely along X_{sm} , proving (2). Also, $\theta_{L_0}(L) \cap X = \theta_{L_0}(L) \cap C = \{x = c_0, c_1, \dots, c_d\}$ with $c_i \neq c_j$ for $i \neq j$, proving (3). Finally, since $(C \cap \hat{f}(Z_{\text{sing}})) \setminus \{x\} \subset S'$ and since $\theta_{L_0}(L) \cap S' = \emptyset$, we see that Z is regular at all points lying over c_i for $1 \leq i \leq d$, proving (4).

We now prove (5). First of all, if the degree of any irreducible component of X inside \mathbb{P}_k^N was less than or equal to two, before we do anything else, we first could have replaced \mathbb{P}_k^N by its suitable Veronese embedding so as to ensure that the degree of any irreducible component of X is bigger than 2. In doing so, we see using Lemma 5.8 that the intersection L_0 of general $(r-1)$ hyperplanes H_1, \dots, H_{r-1} lying in $\text{Gr}_{S \cup \{x\}}(N-1, \mathbb{P}_k^N)$ will have the property that L_0 will satisfy the above (a)–(e), and $L_0 \cap X$ will be a reduced curve none of whose irreducible component is contained in a 2-dimensional linear subspace

of \mathbb{P}_k^N . Note that since X is equidimensional and L_0 is general, the curve $L_0 \cap X$ will have this property too. This proves (5). The last property (6) is a direct consequence of (5), condition (2) of Lemma 5.9, which we already achieved from the beginning, and Proposition 5.6. \square

Later, the set $\{c_1, \dots, c_d\}$ that we obtained in Lemma 5.10 will be taken to be $L^+(x)$ for $x \in \Sigma$, where Σ is the given set of finitely many closed regular points of X . This means the regularity of Z at points lying over the residual points $L^+(\Sigma)$. We will come back to this discussion, and it will be finished in Proposition 7.2.

6. Vertical separation of residual fibers

In this section, we prove some results which we shall need in order to prove the regularity of the residual cycle of Z along Σ . The main goal is to show that the distinct fibers, of the projection $Z \rightarrow X$ to the “horizontal axis” over the residual points of Σ (for a suitable linear projection) are mapped to disjoint sets under the projection $\hat{g} : Z \rightarrow \hat{B}$ to the “vertical axis”. We call this property of linear projections, *the vertical separation of residual fibers*. We continue to use the Set-up+(fs) of Section 5B.

6A. Separating residual fibers of Z along \hat{B} : the local case. Let k be an algebraically closed field. In Lemma 5.2, under certain assumptions, we found a nonempty open subset of a Grassmannian such that each member L satisfies the properties (1)–(4) there. In Lemma 5.9, after choosing a Veronese reembedding into a bigger projective space, we achieved an additional noncollinearity of any three points of the hyperplane sections. It was generalized to Lemma 5.10 for $r \geq 1$.

In Section 6A, we want to further strengthen them, by constructing a nonempty open subset for which we have an additional separation property, which will be called property (I) . This is eventually done in Proposition 6.5.

Up to Lemma 6.4, we assume the following. We suppose $r = 1$. We let $x \in X_{fs}$ be a closed point and let $S \subset X \setminus \{x\}$ be another finite set of closed points. For any map $W \rightarrow X$ and a closed point $y \in X$, let W_y be the reduced fiber of W over y . We work under the Set-up of Lemma 5.9, which includes Lemma 5.2.

Since we want to prove a property called (I) by a kind of double induction argument on the pairs of numbers (m, n) with $0 \leq m \leq n \leq d - 1$, we find it convenient to temporarily introduce some intermediate notations.

Definition 6.1. For $1 \leq n \leq d - 1$ and $0 \leq m \leq n$, we say that a member $H = (H, c_1, \dots, c_d) \in \text{Gr}_x(N - 1, \mathbb{P}_k^N)(k) \times X^d$ is (Z, x, m, n) -admissible, if H satisfies the properties (1) and (2) of Lemma 5.9 with $H \cap X = \{x = c_0, c_1, \dots, c_d\}$, together with the additional property

$$(I)_{m,n} := \begin{cases} \hat{g}(Z_{c_i}) \cap \hat{g}(Z_{c_j}) = \emptyset & \text{for } 0 \leq i \neq j \leq n, \\ \hat{g}(Z_{c_i}) \cap \hat{g}(Z_{c_{n+1}}) = \emptyset & \text{for } 0 \leq i \leq m. \end{cases} \tag{6-1}$$

We remark that for $n = 0$ (thus we have just $(I)_{0,0}$), the first condition of (6-1) is empty.

Before anything else, we note the following elementary fact:

Lemma 6.2. *The projections $\hat{f} : Z \rightarrow X$ and $\hat{g} : Z \rightarrow \hat{B}$ are finite and the sets $\hat{g}(Z_x) \subset \hat{B}$ and $\hat{g}^{-1}(\hat{g}(Z_x)) \subset Z$ are finite subsets of closed points.*

Proof. Note that $\hat{f} : Z \rightarrow X$ is a projective morphism of reduced curves such that its restriction over the dense open subset X_{fs} of X is fs. Hence \hat{f} is a projective quasifinite morphism, hence a finite morphism. Since \hat{g} is a projective morphism from a curve which is nonconstant on each component of the source, it must also be finite. Since Z_x is a finite set, as Z is fs over X_{fs} and $x \in X_{fs}$, the lemma now follows. \square

Let $V_d \subset X^d$ be the nonempty open subset whose coordinates are all distinct from each other and distinct from x as well. More precisely, this is the complement of the union of all the small diagonals $\Delta_{i,j} \subset X^d$ defined by the equation $y_i = y_j$ for $1 \leq i < j \leq d$ as well as the subschemes given by $y_i = x$ for $1 \leq i \leq d$. Let $\pi : X^d \rightarrow \text{Sym}^d(X) = X^d/\mathfrak{S}_d$ be the quotient map for the action by the symmetric group \mathfrak{S}_d which permutes the coordinates. Since \mathfrak{S}_d acts freely on $V_d \subset X^d$, the restriction $\pi : V_d \rightarrow \pi(V_d)$ is finite étale of degree $d!$.

Inside V_d , we consider the following subsets of “bad points” that do not satisfy the analogue of condition $(I)_{m,n}$ for $(y_1, \dots, y_d) \in V_d$. That is, for $y_0 := x$, let $D_0 := \emptyset$, while for $n \geq 1$, let $D_n \subset V_d$ be the subset of points (y_1, \dots, y_d) such that $\hat{g}(Z_{y_i}) \cap \hat{g}(Z_{y_j}) \neq \emptyset$ for some $0 \leq i \neq j \leq n$ and $G_m^n \subset V_d$ be the subset of points such that $\hat{g}(Z_{y_i}) \cap \hat{g}(Z_{y_{n+1}}) \neq \emptyset$ for some $0 \leq i \leq m$.

Express $D_n = D_{n,1} \cup D_{n,2}$, where $D_{n,1}$ consists of the points $(y_1, \dots, y_d) \in V_d$ such that $\hat{g}(Z_{y_i}) \cap \hat{g}(Z_{y_j}) \neq \emptyset$ for some $1 \leq i \neq j \leq n$, while $D_{n,2}$ consists of the points $(y_1, \dots, y_d) \in V_d$ such that $\hat{g}(Z_{y_0}) \cap \hat{g}(Z_{y_i}) \neq \emptyset$ for some $1 \leq i \leq n$. We also write $G_m^n = \bigcup_{i=0}^m G_{m,i}^n$, where $G_{m,i}^n$ consists of the points $(y_1, \dots, y_d) \in V_d$ such that $\hat{g}(Z_{y_i}) \cap \hat{g}(Z_{y_{n+1}}) \neq \emptyset$ for $0 \leq i \leq m$. We check these “bad sets” are closed.

Lemma 6.3. *The subsets $D_{n,i}$ for $i = 1, 2$ and $G_{m,i}^n$ for $0 \leq i \leq m$ are closed subsets of V_d . In particular, D_n and G_m^n are closed subsets of V_d .*

Proof. Let $E_{n,1} \subset \hat{B}^d$ be the subset of points (b_1, \dots, b_d) such that $b_i = b_j$ for some $1 \leq i \neq j \leq n$. Let $E_{n,2} \subset \hat{B}^d$ be the subset of points (b_1, \dots, b_d) such that $b_i \in \hat{g}(Z_x)$ for some $1 \leq i \leq n$. The set $E_{n,1}$ is certainly closed in \hat{B}^d , while $E_{n,2}$ is closed in \hat{B}^d because $\hat{g}(Z_x)$ is finite by Lemma 6.2. One checks that $D_{n,i} = \hat{f}^{\times d}((\hat{g}^{\times d})^{-1}(E_{n,i})) \cap V_d$ for $i = 1, 2$, where $\hat{f}^{\times d} : Z^d \rightarrow X^d$ and $\hat{g}^{\times d} : Z^d \rightarrow \hat{B}^d$ are the direct products of \hat{f} and \hat{g} . Since $\hat{f}^{\times d}$ is finite by Lemma 6.2, this shows that $D_{n,i}$ is closed in V_d .

Similarly, let $J_{m,0}^n \subset \hat{B}^d$ be the subset of points (b_1, \dots, b_d) such that $b_{n+1} \in \hat{g}(Z_x)$. This is closed since $\hat{g}(Z_x)$ is finite by Lemma 6.2. For $1 \leq i \leq m$, let $J_{m,i}^n \subset \hat{B}^d$ be the subset of points (b_1, \dots, b_d) such that $b_i = b_{n+1}$. This is also closed. One checks that $G_{m,i}^n = \hat{f}^{\times d}((\hat{g}^{\times d})^{-1}(J_{m,i}^n)) \cap V_d$, and this shows that $G_{m,i}^n$ is closed in V_d for $0 \leq i \leq m$. \square

Coming back to the story, we let $\mathcal{U}_S \subset \text{Gr}_x(N-1, \mathbb{P}_k^N)$ be the nonempty open set of Lemma 5.9. Let $\mathcal{U}_S \rightarrow \text{Sym}^d(X)$ be the map given by $L \mapsto \sum_{i=1}^d [c_i]$, where $L \cap X = \{x = c_0, c_1, \dots, c_d\}$. By condition (3)

of Lemma 5.2, its image is in $\pi(V_d)$. Define \mathcal{V}_S by the Cartesian diagram

$$\begin{array}{ccc} \mathcal{V}_S & \xrightarrow{e} & V_d \\ \psi \downarrow & & \downarrow \pi \\ \mathcal{U}_S & \longrightarrow & \pi(V_d), \end{array} \tag{6-2}$$

so that ψ is a finite surjective étale map. The set $\mathcal{V}_S \setminus e^{-1}(D_n \cup G_m^n)$ is open in \mathcal{V}_S by Lemma 6.3. Via the open map ψ , we define the open subset $\mathcal{U}_{m,n}^S := \psi(\mathcal{V}_S \setminus e^{-1}(D_n \cup G_m^n)) \subset \mathcal{U}_S$. This is open in $\text{Gr}_x(N-1, \mathbb{P}_k^N)$.

Lemma 6.4. *For $0 \leq n \leq d-1$ and $0 \leq m \leq n$, the subset $\mathcal{U}_{m,n}^S \subset \text{Gr}_x(N-1, \mathbb{P}_k^N)$ is nonempty. In particular, it is a dense open subset of $\text{Gr}_x(N-1, \mathbb{P}_k^n)$.*

Proof. Step 1. $\mathcal{U}_{0,0}^S \neq \emptyset$. Note that condition $(I)_{0,0}$ is independent of the choice of an x -fixing order on $L \cap X$. Let $T = S \cup (\hat{f}(\hat{g}^{-1}(\hat{g}(Z_x))) \setminus \{x\})$. This is a finite closed subset of X by Lemma 6.2. Applying Lemma 5.2 to T (in the place of S there), we obtain a dense open subset \mathcal{U}_T of $\mathcal{U}_S \subset \text{Gr}_x(N-1, \mathbb{P}_k^N)$. On the other hand, condition (1) (in Lemma 5.2) for T implies that for each $L \in \mathcal{U}_T(k)$, we have $L \cap (\hat{f}(\hat{g}^{-1}(\hat{g}(Z_x))) \setminus \{x\}) = \emptyset$, which shows that $\hat{g}(Z_{c_0}) \cap \hat{g}(Z_{c_j}) = \emptyset$ for each $j \neq 0$ when $L \cap X = \{x = c_0, c_1, \dots, c_d\}$, for every x -fixing order on $L \cap X$. Thus $(I)_{0,0}$ holds, and $\mathcal{U}_T \subset \mathcal{U}_{0,0}^S$, in particular $\mathcal{U}_{0,0}^S \neq \emptyset$.

Step 2. For $0 \leq n \leq d-2$, if $\mathcal{U}_{n,n}^S \neq \emptyset$, then $\mathcal{U}_{0,n+1}^S \neq \emptyset$. If $\mathcal{U}_{n,n}^S \neq \emptyset$, then it is a dense open subset of $\text{Gr}_x(N-1, \mathbb{P}_k^N)$. In particular, for the dense open subset $\mathcal{U}_T \subset \text{Gr}_x(N-1, \mathbb{P}_k^N)$ of Step 1, the intersection $\mathcal{U}_{n,n}^S \cap \mathcal{U}_T$ is dense open in $\text{Gr}_x(N-1, \mathbb{P}_k^N)$. But, by definition, one notes that $\mathcal{U}_{n,n}^S \cap \mathcal{U}_T \subset \mathcal{U}_{0,n+1}^S$ so that $\mathcal{U}_{0,n+1}^S \neq \emptyset$.

Step 3. For $0 \leq n \leq d-1$ and $0 \leq m \leq n-1$, if $\mathcal{U}_{m,n}^S \neq \emptyset$, then $\mathcal{U}_{m+1,n}^S \neq \emptyset$. If $\mathcal{U}_{m,n}^S \neq \emptyset$, then it is dense open in $\text{Gr}_x(N-1, \mathbb{P}_k^N)$. For the dense open subset $\mathcal{U}_T \subset \text{Gr}_x(N-1, \mathbb{P}_k^N)$ of Step 1, the intersection $\mathcal{U}_{m,n}^S \cap \mathcal{U}_T$ is therefore nonempty dense open in $\text{Gr}_x(N-1, \mathbb{P}_k^N)$.

Fix an element $L'_0 \in (\mathcal{U}_{m,n}^S \cap \mathcal{U}_T)(k)$ and let $L'_0 \cap X = \{x = c_0, c_1, \dots, c_d\}$. Since every k -point of $\mathcal{U}_{m,n}^S$ satisfies condition (2) of Lemma 5.9, we know that no three points of $L'_0 \cap X$ are collinear. Thus $\{c_0, c_{m+1}, c_{n+1}\}$ are not collinear so that when $\ell = \text{Sec}(\{c_0\}, \{c_{m+1}\})$ is the line joining c_0 and c_{m+1} , it does not pass through c_{n+1} .

We let $P = \text{Sec}(\{c_{n+1}\}, \ell)$. The subspace $\text{Gr}_\ell(N-1, \mathbb{P}_k^N)$ is of dimension $N-2$ and $\text{Gr}_P(N-1, \mathbb{P}_k^N)$ is a closed subspace of $\text{Gr}_\ell(N-1, \mathbb{P}_k^N)$ of dimension $N-3$ (see Lemma 5.1). Because we may assume $N \geq 3$, there is a one-parameter family (actually isomorphic to \mathbb{P}_k^1) \mathcal{B} in $\text{Gr}_x(N-1, \mathbb{P}_k^N)$ such that (i) $\{L'_0\} \in \mathcal{B}$, (ii) every member of the family \mathcal{B} passes through both of $\{c_0, c_{m+1}\}$ and (iii) a general member does not pass through c_{n+1} . Since $\mathcal{U}_{m,n}^S \cap \mathcal{U}_T$ is dense open in $\text{Gr}_x(N-1, \mathbb{P}_k^N)$ and $L'_0 \in \mathcal{U}_{m,n}^S \cap \mathcal{U}_T \cap \text{Gr}_L(N-1, \mathbb{P}_k^N)$, the latter is dense open in $\text{Gr}_\ell(N-1, \mathbb{P}_k^N)$. Hence, a general member of \mathcal{B} is contained in $\mathcal{U}_{m,n}^S \cap \mathcal{U}_T$.

Let $W \subset \mathcal{B} \cap \mathcal{U}_{m,n}^S \cap \mathcal{U}_T$ be a smooth affine irreducible (rational) curve passing through $\{L'_0\}$. Consider again the quotient map $\pi : X^d \rightarrow \text{Sym}^d(X) = X^d/\mathcal{S}_d$, and the finite étale map $\pi : V_d \rightarrow \pi(V_d)$ for the open set V_d defined previously in (6-2). Consider the map $W \rightarrow \pi(V_d)$ given by $L \mapsto \sum_{i=1}^d [y_i]$, where

$L \cap X = \{x = y_0, y_1, \dots, y_d\}$. This yields the Cartesian product

$$\begin{array}{ccccc}
 & & e & & \\
 & & \curvearrowright & & \\
 W' & \longrightarrow & \tilde{W} & \longrightarrow & V_d \\
 & \searrow & \downarrow \psi & & \downarrow \pi \\
 & & W & \longrightarrow & \pi(V_d)
 \end{array} \tag{6-3}$$

so that ψ is finite and étale. Note also that the members of \tilde{W} can be represented by $\underline{L} = (L, y_1, \dots, y_d) \in W \times V_d$ such that $L \cap X = \{x = y_0, y_1, \dots, y_d\}$. We let $W' \subset \tilde{W}$ be the component containing the point (L'_0, c_1, \dots, c_d) . For the “bad” closed subsets $D_n, G_{m+1}^n \subset V_d$ of Lemma 6.3, we have:

Claim. $\mathcal{Y} := e^{-1}(D_n \cup G_{m+1}^n)$ is a proper closed subset of W' .

Proof. That this is a closed subset of W' follows by Lemma 6.3. We need to show that this is a proper subset. Note that $D_n = D_{n,1} \cup D_{n,2}$ and $G_{m+1}^n = \bigcup_{i=0}^{m+1} G_{m+1,i}^n$. We analyze each piece of them in what follows.

Case 1: We first show that $e^{-1}(D_{n,2}) = \emptyset$ and $e^{-1}(G_{m+1,0}^n) = \emptyset$.

Note that we had $W \subset \mathcal{B} \cap \mathcal{U}_{m,n}^S \cap \mathcal{U}_T$, where \mathcal{U}_T is as in Lemma 5.2. Here, condition (1) of Lemma 5.2 (and S replaced by T) reads as ‘ $L \cap ((X \setminus X_{\text{fs}}) \cup T) = \emptyset$ ’ for each $L \in \mathcal{U}_T(k)$. So, for every $L \in W(k)$, this is disjoint from $T = S \cup (\hat{f}(\hat{g}^{-1}(\hat{g}(Z_x))) \setminus \{x\})$. Hence, if $e^{-1}(D_{n,2}) \neq \emptyset$, then it gives an element $L \in W(k)$ such that $L \cap X = \{x = y_0, y_1, \dots, y_d\}$ satisfies $\hat{g}(Z_{y_0}) \cap \hat{g}(Z_{y_i}) \neq \emptyset$ for some $1 \leq i \leq n$, so that L intersects with a point of T , contradicting the above choice of W . Hence $e^{-1}(D_{n,2}) = \emptyset$. An identical argument shows that $e^{-1}(G_{m+1,0}^n) = \emptyset$.

Case 2: We now show that $e^{-1}(D_{n,1})$ and $e^{-1}(G_{m+1,i}^n)$ for $1 \leq i \leq m$ are finite.

To do so, it is enough to show that these closed subsets are proper in W' , as W' is an irreducible curve. Suppose $e^{-1}(D_{n,1}) = W'$. In particular $\underline{L}'_0 := (L'_0, c_1, \dots, c_d) \in e^{-1}(D_{n,1})$, so that $(c_1, \dots, c_d) \in D_{n,1}$, so $\hat{g}(Z_{c_i}) \cap \hat{g}(Z_{c_j}) \neq \emptyset$ for some $1 \leq i \neq j \leq n$. But, this contradicts that $L'_0 \in \mathcal{U}_{m,n}^S(k)$. Hence, $e^{-1}(D_{n,1})$ is proper closed in W' . By the same argument, we have $|e^{-1}(G_{m+1,i}^n)| < \infty$.

Case 3: It remains to show that $|e^{-1}(G_{m+1,m+1}^n)| < \infty$.

To do so, we will make use of our choice of W that $W \subset \mathcal{B}$. Recall that $\mathcal{B} \subset \text{Gr}_x(N-1, \mathbb{P}_k^N)$ is a one-parameter family containing $\{L'_0\}$ such that every member of \mathcal{B} passes through $\{c_0, c_{m+1}\}$, while a general member does not pass through c_{n+1} .

Consider the composite $q : W' \xrightarrow{e} V_d \rightarrow X^2$, where the last arrow takes (y_1, \dots, y_d) to $(y_{m+1}, y_{n+1}) \in X^2$. Since every $L \in W(k) \subset \mathcal{B}(k)$ contains c_{m+1} by construction, the composition of q with the first projection $X^2 \rightarrow X$, taking (y_{m+1}, y_{n+1}) to y_{m+1} , is the constant map that takes all of W' to $c_{m+1} \in X$. On the other hand, the general member $L \in W(k)$ does not contain c_{n+1} . This implies that the composite of q with the second projection $X^2 \rightarrow X$, taking (y_{m+1}, y_{n+1}) to y_{n+1} , is nonconstant. Hence, the map q is nonconstant and the image $q(W')$ in X^2 is an irreducible curve contained in $\{c_{m+1}\} \times X \cong X$ (recall that k is assumed to be algebraically closed).

Write it as $W' \xrightarrow{u} q(W') \xrightarrow{v} X$, where u is induced by q and v is the projection to the coordinate y_{n+1} . Since both u and v are nonconstant morphisms of irreducible curves, they are dominant and quasifinite. In particular, the composite $v \circ u$ is quasifinite. Note that by definition, $e^{-1}(G_{m+1, m+1}^n) \subset \{(L, y_1, \dots, y_d) \in W' \mid y_{n+1} \in S_1\} = (v \circ u)^{-1}(S_1)$, where $S_1 := f(\hat{g}^{-1}(\hat{g}(Z_{c_{m+1}})))$. Since \hat{f} and \hat{g} are finite by Lemma 6.2, the set S_1 is finite, thus $(v \circ u)^{-1}(S_1)$ is a finite set. Hence, we have $|e^{-1}(G_{m+1, m+1}^n)| < \infty$, being a subset of a finite set. This finishes the proof of the Claim. \square

Back to the proof of Step 3, since the set \mathcal{Y} of the Claim is finite, the subset $W' \setminus \mathcal{Y} \subset W'$ is nonempty open. Since ψ is an open map and $W' \subset \tilde{W}$ is open subset such that $W' \rightarrow W$ is surjective, it follows that $\psi(W' \setminus \mathcal{Y}) \subset W$ is a nonempty (thus dense) open subset. By construction, $\psi(W' \setminus \mathcal{Y}) \subset \mathcal{U}_{m+1, n}^S$. In particular, we get $\mathcal{U}_{m+1, n}^S \neq \emptyset$. This proves Step 3.

Back to the proof of the lemma, by inductively applying the above three steps, we deduce that each $\mathcal{U}_{m, n}^S$ is a dense open subset of $\text{Gr}_x(N - 1, \mathbb{P}_k^N)$. \square

Now we allow $r \geq 1$. We can strengthen Lemma 5.10 as follows:

Proposition 6.5. *We follow the notations and the assumptions of Lemma 5.10. Let $r \geq 1$. After replacing \mathbb{P}_k^N by a bigger projective space via Veronese if necessary, we have the following property: given any hyperplane $H_0 \subset \mathbb{P}_k^N$ disjoint from $S \cup \{x\}$ and a general $L_0 \in \text{Gr}_{S \cup \{x\}}^{\text{tr}}(H_0, N - r + 1, \mathbb{P}_k^N)(k)$, there exists a dense open subset $\mathcal{U}_x^S \subset \text{Gr}_x^{\text{tr}}(L_0, N - 1, \mathbb{P}_k^N)$ such that each $L \in \mathcal{U}_x^S(k)$ satisfies the properties (1)–(6) of Lemma 5.10 as well as the additional property (I) : $\hat{g}(Z_{c_i}) \cap \hat{g}(Z_{c_j}) = \emptyset$ for each pair $0 \leq i \neq j \leq d$.*

Proof. The $r = 1$ case of the proposition follows from Lemma 6.4 with $(m, n) = (d - 1, d - 1)$. So we assume $r \geq 2$.

We use an argument of reduction to the $r = 1$ case as we did in Lemma 5.10. Using the notations there, choose a reembedding $\eta : X \hookrightarrow \mathbb{P}_k^N$, a general $L_0 \in \text{Gr}(N - r + 1, \mathbb{P}_k^N)(k)$ and $C = L_0 \cap X$ as in Lemma 5.10. Let $S' := (C \setminus \{x\}) \cap (\hat{f}(Z_{\text{sing}}) \cup (X \setminus X_{\text{fs}}) \cup S)$ and $W = Z|_{C \times \hat{B}}$. Applying the “ $r = 1$ ” case of the proposition (proven in Lemma 6.4) to C, S' and W , with the identification $L_0 \simeq \mathbb{P}_k^{N-r+1}$, there is a dense open subset $\mathcal{U}' \subset \text{Gr}_x(N - r, L_0)$ that satisfies the properties of Proposition 6.5 for $r = 1$ case. (In terms of the notations of Lemma 6.4, we have $\mathcal{U}' = \mathcal{U}_{d-1, d-1}^{S'}$.) Note that Lemma 6.4 is applicable to C by property (6) of Lemma 5.10.

Recall now that we had a smooth surjective morphism of varieties

$$\theta_{L_0} : \text{Gr}_x^{\text{tr}}(L_0, N - 1, \mathbb{P}_k^N) \rightarrow \text{Gr}_x(N - r, L_0)$$

from (5-8). So, the inverse image $\mathcal{U}_x^S := \theta_{L_0}^{-1}(\mathcal{U}')$ is a dense open subset of $\text{Gr}_x^{\text{tr}}(L_0, N - 1, \mathbb{P}_k^N)$. We claim that this \mathcal{U}_x^S fulfills the requirements of the proposition for $r \geq 2$ case.

Indeed, since $W = Z|_{C \times \hat{B}}$, we see that $Z_y = W_y$ and hence $\hat{g}(Z_y) = \hat{g}(W_y)$ for any closed point $y \in C$. Hence, for $L \in \text{Gr}_x^{\text{tr}}(L_0, N - 1, \mathbb{P}_k^N)(k)$ with $\theta_{L_0}(L) \cap X = (L \cap L_0) \cap C = \{x = c_0, c_1, \dots, c_d\}$, condition (I) is satisfied if and only if condition (I) is satisfied for $\theta_{L_0}(L)$ with X replaced by the curve C . This means $L \in \mathcal{U}_x^S(k)$ satisfies the proposition, as desired. \square

6B. Separating residual fibers of Z along \hat{B} : the semilocal case. Note that in the statement of Proposition 6.5, the dense open subset that we found depends on the choice of a single regular closed point $x \in X$. We want to extend it to a finite subset Σ of regular points. This issue will be completely resolved in Proposition 7.2 by using the “cone admissibility” condition, which we develop as property (3) of the following Proposition 6.6. One further aspect on étaleness is studied in Section 6C.

Recall that when $M \subset \mathbb{P}_k^N$ is a linear subspace and $x \in \mathbb{P}_k^N$ is a closed point, after the base change $\text{Spec}(k(x)) \rightarrow \text{Spec}(k)$, the cone $C_x(M) = \text{Sec}(\{x\}, M)$ is the smallest linear subspace containing both x and M . When $x \notin M$, we have $\dim(C_x(M)) = \dim(M) + 1$. In this article, we need to use the cones only when k is algebraically closed, so that no confusion will arise.

Proposition 6.6. *Let k be an algebraically closed field. We are under the Set-up+(fs) of Section 5B. After replacing the embedding $X \hookrightarrow \mathbb{P}_k^N$ by a bigger one via a Veronese embedding if necessary, we have the following: for the given hyperplane $H \subset \mathbb{P}_k^N$ disjoint from Σ and a general $L_0 \in \text{Gr}^{\text{tr}}(H, N - r + 1, \mathbb{P}_k^N)(k)$, there exists a dense open subset $\mathcal{W} \subset \text{Gr}(N - 2, H)$ such that each $M \in \mathcal{W}(k)$ satisfies the following properties:*

- (1) M intersects L_0 transversely.
- (2) $M \cap L_0 \cap X = \emptyset$.
- (3) For each $x \in \Sigma$, the cone $C_x(M)$ lies in $\mathcal{U}_x^{\Sigma \setminus \{x\}}(k)$ for the open subset $\mathcal{U}_x^{\Sigma \setminus \{x\}} \subset \text{Gr}_x^{\text{tr}}(L_0, N - 1, \mathbb{P}_k^N)$ of Proposition 6.5.

Proof. Note that if $\Sigma = \{x_1, \dots, x_n\}$, then condition (3) consists of conditions $(3)_i : C_{x_i}(M) \in \mathcal{U}_{x_i}^{\Sigma \setminus \{x_i\}}(k)$ for $1 \leq i \leq n$. Suppose we proved the existence of a dense open subset $\mathcal{W}_i \subset \text{Gr}(N - 2, H)$ for which each member $M \in \mathcal{W}_i(k)$ satisfies conditions (1), (2), and $(3)_i$ for each $1 \leq i \leq n$. Then we can take $\mathcal{W} := \bigcap_{i=1}^n \mathcal{W}_i$, which is again a dense open subset of $\text{Gr}(N - 2, H)$. Hence, it is enough to prove the existence of those \mathcal{W}_i . Without loss of generality, we may assume $i = 1$. For notational simplicity, we let $x := x_1$ and $T := \Sigma \setminus \{x_1\}$. We note also that when $r = 1$, we have $\text{Gr}^{\text{tr}}(H, N - r + 1, \mathbb{P}_k^N) = \text{Gr}(N, \mathbb{P}_k^N) = \{\mathbb{P}_k^N\}$ so that the choice of L_0 plays no role. We prove the proposition for the cases of $r = 1$ and $r \geq 2$ separately.

Step 1. Suppose $r = 1$. Consider the affine morphism of schemes

$$\vartheta_x : \text{Gr}(x, N - 2, \mathbb{P}_k^N) \rightarrow \text{Gr}_x(N - 1, \mathbb{P}_k^N), L \mapsto C_x(L). \tag{6-4}$$

This is a smooth surjective morphism, and defines a vector bundle of rank $N - 1$. For the closed irreducible subscheme $\text{Gr}(N - 2, H) \hookrightarrow \text{Gr}(x, N - 2, \mathbb{P}_k^N)$, the restriction $\vartheta_{x,H} : \text{Gr}(N - 2, H) \rightarrow \text{Gr}_x(N - 1, \mathbb{P}_k^N)$ of ϑ_x , is an isomorphism.

Let $\mathcal{U}_x^T \subset \text{Gr}_x(N - 1, \mathbb{P}_k^N)$ be the dense open subset of Proposition 6.5, applied to x, T and $H_0 = H$ for $r = 1$. Then $\vartheta_{x,H}^{-1}(\mathcal{U}_x^T)$ is a dense open subset of $\text{Gr}(N - 2, H)$. Since $\text{Gr}(X, N - 2, H)$ is its dense open subset by Lemma 4.3, so is the intersection $\mathcal{W}_1 := \vartheta_{x,H}^{-1}(\mathcal{U}_x^T) \cap \text{Gr}(X, N - 2, H)$ in $\text{Gr}(N - 2, H)$. One checks that this satisfies the required conditions (1), (2), and $(3)_1$, proving the proposition for $r = 1$.

Step 2. Suppose now that $r \geq 2$. As we did previously in Lemma 5.10 with $H_0 = H$ via a Bertini argument of [Kleiman and Altman 1979], we choose a reembedding $\eta : X \hookrightarrow \mathbb{P}_k^N$, a general $L_0 \in \text{Gr}^{\text{tr}}(H, N - r + 1, \mathbb{P}_k^N)(k)$, a curve $C = L_0 \cap X$, and $Z|_{C \times \hat{B}}$. Consider again the map in (6-4). When L_0 contains x , this ϑ_x induces a smooth surjective map $\vartheta_x^{L_0} : \text{Gr}^{\text{tr}}(L_0, x, N - 2, \mathbb{P}_k^N) \rightarrow \text{Gr}_x^{\text{tr}}(L_0, N - 1, \mathbb{P}_k^N)$, where we recall that $\text{Gr}^{\text{tr}}(L_0, x, n, \mathbb{P}_k^N) := \text{Gr}^{\text{tr}}(L_0, n, \mathbb{P}_k^N) \cap \text{Gr}(x, n, \mathbb{P}_k^N)$. This restricts to give $\vartheta_{x,H} : \text{Gr}^{\text{tr}}(L_0, N - 2, H) \rightarrow \text{Gr}_x^{\text{tr}}(L_0, N - 1, \mathbb{P}_k^N)$. One checks that this map is an inclusion whose image is the dense open subset $\text{Gr}_x^{\text{tr}}(L_0 \cap H, N - 1, \mathbb{P}_k^N)$. As $H \cap \{x\} = \emptyset$, we see that $\text{Gr}_x^{\text{tr}}(L_0 \cap H, N - 1, \mathbb{P}_k^N)$ coincides with $\text{Gr}_x^{\text{tr}}(L_0, N - 1, \mathbb{P}_k^N)$. This implies that $\vartheta_{x,H}$ is an isomorphism.

Let $\mathcal{U}_x^T \subset \text{Gr}_x^{\text{tr}}(L_0, N - 1, \mathbb{P}_k^N)$ be the dense open subset of Proposition 6.5 applied to x, T , and $H_0 = H$ for $r \geq 2$. Since $\vartheta_{x,H}$ is an isomorphism, $\vartheta_{x,H}^{-1}(\mathcal{U}_x^T)$ is dense open in $\text{Gr}^{\text{tr}}(L_0, N - 2, H)$, thus dense open in $\text{Gr}(N - 2, H)$. Combining this with Lemma 4.3, we conclude that $\mathcal{W}_1 := \vartheta_{x,H}^{-1}(\mathcal{U}_x^T) \cap \text{Gr}(C, N - 2, H)$ is dense open in $\text{Gr}(N - 2, H)$. One checks that each $M \in \mathcal{W}(k)$ satisfies the required conditions (1), (2), and (3)₁. This finishes the proof. \square

6C. Étaleness of linear projections at $L^+(\Sigma)$. Recall that we had obtained a linear projection $\phi_L : X \rightarrow \mathbb{P}_k^r$ that is étale at each point of Σ in condition (1) of Lemma 4.8. Unfortunately, this is not quite enough for us. We need to have L such that ϕ_L is étale at each point of $L^+(\Sigma)$ as well. We show that we can achieve this as a geometric consequence of condition (3) of Proposition 6.6.

Part of the requirement of Proposition 6.6 that $C_x(M)$ lies in $\mathcal{U}_x^{\Sigma \setminus \{x\}}(k)$ for the open set $\mathcal{U}_x^{\Sigma \setminus \{x\}}$ is that $C_x(M)$ intersects $X_{\text{fs}} \subset X_{\text{sm}}$ transversely. This comes from condition (2) of Lemma 5.10. Here is its geometric meaning.

Lemma 6.7. *Let k be an algebraically closed field and let $L \in \text{Gr}(X, N - r - 1, \mathbb{P}_k^N)(k)$. Let \mathbb{P}_k^r be a linear subspace of \mathbb{P}_k^N such that $L \cap \mathbb{P}_k^r = \emptyset$. Let $y \in \mathbb{P}_k^r$ be a closed point such that $C_y(L) \cap X_{\text{sing}} = \emptyset$. Then $C_y(L)$ intersects X transversely if and only if the linear projection $\phi_L : X \rightarrow \mathbb{P}_k^r$ away from L is finite and étale over an affine neighborhood of y in \mathbb{P}_k^r .*

Proof. (\Rightarrow) Suppose that $C_y(L)$ intersects X transversely and let $E := C_y(L) \cap X$ be this scheme-theoretic intersection. Since k is perfect while $C_y(L)$ and X_{sm} have the complementary dimensions $N - r$ and r in \mathbb{P}_k^N , respectively, the transverse intersection is equivalent to saying that E is smooth, $|E| < \infty$, and each point of E is a simple regular point of X_{sm} . Because we are given that $X \cap L = \emptyset$ and $L \subset C_y(L)$, we see that $C_y(L) \cap X = (C_y(L) \setminus L) \cap X$, which is precisely the scheme-theoretic fiber $\phi_L^{-1}(y)$ over $y \in \mathbb{P}_k^r$.

Since $C_y(L) \cap X_{\text{sing}} = \emptyset$, we see that $\phi_L^{-1}(y) \cap X_{\text{sing}} = \emptyset$. Since ϕ_L is finite, $\phi_L(X_{\text{sing}})$ is a closed subscheme of \mathbb{P}_k^r not meeting y . Hence, there is an affine open $U \subset \mathbb{P}_k^r$ containing y such that $\phi_L^{-1}(U)$ is regular. We therefore get a Cartesian square

$$\begin{array}{ccc}
 E & \longrightarrow & \phi_L^{-1}(U) \\
 \phi_L^y \downarrow & & \downarrow \phi_L \\
 \text{Spec}(k(y)) & \longrightarrow & U
 \end{array} \tag{6-5}$$

such that ϕ_L^y is smooth. Since ϕ_L is a finite map of regular affine schemes over k , it is flat by [Hartshorne 1977, Exercise III-10.9, page 276] (or [EGA IV₂ 1965, Proposition (6.1.5), page 136]). It follows therefore by [Hartshorne 1977, Exercise III-10.2, page 275] (or [EGA IV₃ 1966, Théorème (12.2.4)(iii), page 183]) that there is an affine neighborhood of y in U over which the restriction of the map ϕ_L is smooth, thus finite and étale.

(\Leftarrow) If ϕ_L is étale over a neighborhood of y , then its base change to $\text{Spec}(k(y))$, i.e., the map $\phi_L^y : E = C_y(L) \cap X \rightarrow \text{Spec}(k(y))$ from the scheme-theoretic intersection is étale. Since $k = k(y)$, this means E is smooth over k so that the intersection is transverse. \square

Corollary 6.8. *Let k be an algebraically closed field. Let $L \in \text{Gr}(X, N - r - 1, \mathbb{P}_k^N)(k)$ and realize the linear projection $\phi_L : X \rightarrow \mathbb{P}_k^r$ for a linear subspace $\mathbb{P}_k^r \subset \mathbb{P}_k^N$ such that $L \cap \mathbb{P}_k^r = \emptyset$. Suppose that for each $x \in \Sigma$, we have $C_x(L) \cap X_{\text{sing}} = \emptyset$ and $C_x(L)$ intersects X_{sm} transversely. Then there is an affine open neighborhood $U \subset \mathbb{A}_k^r$ of $\phi_L(\Sigma)$ such that $\phi_L : \phi_L^{-1}(U) \rightarrow U$ is finite and étale. In particular, $\phi_L : X \rightarrow \mathbb{P}_k^r$ is étale at every point of $\phi_L^{-1}(\phi_L(\Sigma))$.*

Proof. Note that for each $x \in \Sigma$, we have $C_x(L) = C_{\phi_L(x)}(L)$ since ϕ_L is given with a chosen internal linear subspace $\mathbb{P}_k^r \subset \mathbb{P}_k^N$. Since $C_{\phi_L(x)}(L) \cap X_{\text{sing}} = \emptyset$ and $C_{\phi_L(x)}(L)$ intersects X transversely, Lemma 6.7 says that there is an affine open neighborhood $U_x \subset \mathbb{P}_k^r$ of $\phi_L(x)$ such that $\phi_L : \phi_L^{-1}(U_x) \rightarrow U_x$ is finite and étale. Hence, for $U := \bigcup_{x \in \Sigma} U_x$, the restriction $\phi_L : \phi_L^{-1}(U) \rightarrow U$ is finite and étale. By Lemma 2.3, we may shrink this U into an affine open neighborhood of $\phi_L(\Sigma)$. This implies the corollary. \square

7. Regularity of residual cycles over finite closed points

Our goal in Section 7 is to study the regularity of the residual cycles using the technique of vertical separation of residual fibers studied in Section 6. We continue to work with the Set-up+(fs) of Section 5B. In particular, for each irreducible component Z_i of Z , the projection $Z_i \rightarrow \hat{B}$ is nonconstant and the projection $Z_i \rightarrow X$ is fs over X_{fs} .

7A. Admissible sets. Property (I) in Proposition 6.5 encourages the following definition, that encodes a set of data needed to achieve the remaining properties of residual cycles.

Definition 7.1. Let k be an infinite perfect field. Let $x \in X_{\text{fs}}$ be a closed point. A finite subset $D \subset X_{\text{fs}}$ of distinct closed points is called (Z, x) -admissible if (1) $x \in D$, (2) Z is regular at all points lying over $D \setminus \{x\}$, and (3) $\hat{g}(Z_{x_1}) \cap \hat{g}(Z_{x_2}) = \emptyset$ for each distinct pair $x_1 \neq x_2$ in D .

The following application of Proposition 6.6 will be a basis for our proof of the regularity of the residual cycles along Σ . We study it for $k = \bar{k}$ case, but it will soon be generalized gradually.

Proposition 7.2. *Let k be an algebraically closed field. We are under the Set-up+(fs) of Section 5B. Let $Y \subset X$ be a closed subset of dimension at most $r-1$. After replacing the embedding $\eta : X \hookrightarrow \mathbb{P}_k^N$ by a bigger one via a Veronese embedding if necessary, we have the following: for the given hyperplane $H \subset \mathbb{P}_k^N$ disjoint from Σ , there is a dense open subset $\mathcal{U} \subset \text{Gr}(N - r - 1, H)$ such that for each $L \in \mathcal{U}(k)$,*

we have $L \cap X = \emptyset$ so that there is a finite and surjective linear projection $\phi_L : X \rightarrow \mathbb{P}_k^r$. Furthermore, it satisfies the following properties:

- (1) The map $\phi_L : \phi_L^{-1}(U) \rightarrow U$ is étale for some affine open $U \subset \mathbb{P}_k^r$ containing $\phi_L(\Sigma)$.
- (2) $\phi_L(x) \neq \phi_L(x')$ for each pair $x \neq x' \in \Sigma$.
- (3) $k(\phi_L(x)) \xrightarrow{\sim} k(x)$ for each $x \in \Sigma$.
- (4) $L^+(x) \cap Y = \emptyset$ for each $x \in \Sigma$.
- (5) $\phi_L^{-1}(\phi_L(x))$ is (Z, x) -admissible for each $x \in \Sigma$.

Proof. As in the proof of Proposition 6.5, we can choose a reembedding $\eta : X \hookrightarrow \mathbb{P}_k^N$ and a dense open subset $\mathcal{U}_1 \subset \text{Gr}_\Sigma(N - r + 1, \mathbb{P}_k^N)$ such that each $L' \in \mathcal{U}_1(k)$ satisfies the condition that $L' \cap X$ is a reduced curve none of whose components is contained in Y , is regular away from X_{sing} , and for each component of $Z|_{X \times \hat{B}}$, the projection to \hat{B} is nonconstant. Since $H \cap \Sigma = \emptyset$, we see that $\mathcal{U}_0 := \text{Gr}_\Sigma(N - r + 1, \mathbb{P}_k^N) \cap \text{Gr}^{\text{tr}}(H, N - r + 1, \mathbb{P}_k^N) \neq \emptyset$. It follows that this intersection is dense open in $\text{Gr}_\Sigma(N - r + 1, \mathbb{P}_k^N)$. Letting $\mathcal{U}'_1 := \mathcal{U}_0 \cap \mathcal{U}_1$, we see that \mathcal{U}'_1 is a dense open subset of $\text{Gr}_\Sigma(N - r + 1, \mathbb{P}_k^N)$ such that each $L' \in \mathcal{U}'_1(k)$ intersects H transversely and $L' \cap X$ is a curve of the above type.

Choose $L_0 \in \mathcal{U}'_1(k)$. (N.B., When $r = 1$, there is a unique choice $L_0 = \mathbb{P}_k^N$ automatically, and we have $C = X$.) We now apply Proposition 6.6. It follows that there exists a dense open subset $\mathcal{W} \subset \text{Gr}(N - 2, H)$ such that each $M \in \mathcal{W}(k)$ satisfies conditions (1)–(3) of Proposition 6.6.

On the other hand, the subset $\text{Gr}^{\text{tr}}(L_0, \text{Sec}(\Sigma, Y \cap C), N - 2, H) \subset \text{Gr}(N - 2, H)$ is a dense open subset by Lemmas 4.3 and 4.4. Hence $\mathcal{V}' := \mathcal{W} \cap \text{Gr}^{\text{tr}}(L_0, \text{Sec}(\Sigma, Y \cap C), N - 2, H) \subset \text{Gr}(N - 2, H)$ is a dense open subset. Since L_0 intersects H transversely, the map $\text{Gr}(N - 2, H) \rightarrow \text{Gr}(N - r - 1, H)$, given by $M \mapsto L_0 \cap M$, is smooth and surjective (note that $N \gg r$). In particular, the image $\mathcal{U}_2 := \{L_0 \cap M \in \text{Gr}(N - r - 1, H) \mid M \in \mathcal{V}'\}$ of \mathcal{V}' is a dense open subset of $\text{Gr}(N - r - 1, H)$. Let $\mathcal{U}'_2 \subset \text{Gr}(X, N - r - 1, H)$ be the dense open set of Lemma 4.8 so that $\mathcal{U} := \mathcal{U}_2 \cap \mathcal{U}'_2 \subset \text{Gr}(X, N - r - 1, H)$ is a dense open subset.

Claim. *Each $L \in \mathcal{U}(k)$ satisfies the properties (1)–(5) of the proposition.*

We ignore L from the notation of ϕ_L for simplicity. Before we prove the claim, we note that $\phi^{-1}(\phi(\Sigma)) \subset X_{\text{fs}}$, as follows from condition (3) of Proposition 6.6 which includes condition (1) of Lemma 5.10.

Now, condition (3) of Proposition 6.6 also implies that, by Corollary 6.8, there is an affine neighborhood U of $\phi(\Sigma)$ such that $\phi^{-1}(U) \rightarrow U$ is finite étale. This proves (1).

Since our open set \mathcal{U} is contained in the open set of Lemma 4.8, we can use the properties there, too. Condition (2) of Lemma 4.8 is that the map ϕ is injective on Σ , proving (2). Condition (3) is obvious because k is assumed to be algebraically closed. Condition (4) follows from our choice of M (thus of L) that it avoids the cone involving Y .

We now prove (5). We need to verify the three conditions of the (Z, x) -admissibility of Definition 7.1 for each $x \in \Sigma$. Condition (1) of Definition 7.1 that $x \in \phi^{-1}(\phi(x))$ is obvious.

We prove condition (2) of Definition 7.1. Condition (3) of Proposition 6.6 says that condition (4) of Lemma 5.10 applied to $C_x(L) \cap X$ holds. Note that the cone $C_x(L)$ plays the role of the linear space in the statement of Lemma 5.10. That is, each point of Z lying over a point of $(C_x(L) \cap X) \setminus \{x\}$ is a regular point of Z . This means that each point of Z lying over a point of $\phi^{-1}(\phi(x)) \setminus \{x\}$ is regular. This proves condition (2) of Definition 7.1 for $\phi^{-1}(\phi(x))$.

Condition (3) of Definition 7.1 for the (Z, x) -admissibility of $\phi^{-1}(\phi(x))$ for $x \in \Sigma$ follows from condition (I) of Proposition 6.5, which is part of condition (3) of Proposition 6.6. This proves (5). We have thus proven the claim, and hence, the proposition. \square

7B. Regularity of residual cycles: $k = \bar{k}$ case. We now prove regularity of residual cycles at points lying over Σ using Proposition 7.2 when k is algebraically closed. Recall (Section 4D) that for a linear projection $\phi_L : X \rightarrow \mathbb{P}_k^r$, the residual scheme $L^+(Z)$ is the closure of $\hat{\phi}_L^{-1}(\hat{\phi}_L(Z)) \setminus Z$ in $X \times \hat{B}$ with the reduced induced closed subscheme structure.

We let $T := \hat{\phi}_L(Z) = \hat{\phi}_L(L^+(Z)) \subset \mathbb{P}_k^r \times \hat{B}$ with the reduced subscheme structure and let $\tilde{Z} := T \times_{(\mathbb{P}_k^r \times \hat{B})} (X \times \hat{B}) = \hat{\phi}_L^{-1}(T) = \hat{\phi}_L^{-1}(\hat{\phi}_L(Z))$ as a scheme. We first have:

Lemma 7.3. *We are under the Set-up+(fs) of Section 5B. Let $x \in X_{\text{fs}}$ be a closed point. Suppose in addition that Z is irreducible.*

Let $\phi_L : X \rightarrow \mathbb{A}_k^r$ be a finite surjective morphism obtained by a linear projection as before such that $\phi_L^{-1}(\phi_L(x))$ satisfies condition (3) of Definition 7.1 of (Z, x) -admissibility. Let $\alpha \in Z$ be a point lying over a point of $\phi_L^{-1}(\phi_L(x))$. Let $S = \hat{\phi}_L^{-1}(\hat{\phi}_L(\alpha))$.

Then $Z \cap S = \{\alpha\}$ and the natural map $\mathcal{O}_{Z, Z \cap S} \rightarrow \mathcal{O}_{Z, \alpha}$ is an isomorphism of local rings.

Proof. Suppose $\alpha \in Z$ lies over $x_1 \in \phi_L^{-1}(\phi_L(x))$. Toward contradiction, suppose there is a point $\alpha' \in Z$ lying over some $x_2 \in \phi_L^{-1}(\phi_L(x)) \setminus \{x_1\}$. Since $\hat{\phi}_L(\alpha) = \hat{\phi}_L(\alpha')$, we have $\hat{g}(\alpha) = \hat{g}(\alpha')$ in B , where $\hat{g} : X \times \hat{B} \rightarrow \hat{B}$ is the projection. Let b_0 be this common closed point. This $Z \rightarrow B$ is nonconstant and we have $\alpha \in Z_{x_1}$ and $\alpha' \in Z_{x_2}$ so that $\hat{g}(Z_{x_1}) \cap \hat{g}(Z_{x_2}) \ni b_0$, contradicting condition (3) of Definition 7.1 for the set $\phi_L^{-1}(\phi_L(x))$. \square

Lemma 7.4. *Let k be algebraically closed. Let $L \in \mathcal{U}(k) \subset \text{Gr}(N - r - 1, H)(k)$ be as in Proposition 7.2. Suppose Z is irreducible and let $\alpha = (a, b) \in Z$ be a closed point such that $a \in \phi_L^{-1}(\phi_L(\Sigma))$. Assume that Z is irreducible and $\alpha \in Z$. Then $\mathcal{O}_{\tilde{Z}, \alpha} \rightarrow \mathcal{O}_{Z, \alpha}$ is an isomorphism. In particular, Z is the only irreducible component of \tilde{Z} which passes through α , with multiplicity 1, and the cycle $[\tilde{Z}] - [Z]$ has no component equal to Z .*

Proof. We shall write ϕ_L simply as ϕ . Let $y = \phi(a)$ and $\beta = \hat{\phi}(\alpha) = (\phi(a), b) = (y, b)$. We let $x \in \Sigma$ be such that $y = \phi(x)$ and let $S = \phi^{-1}(y) \times \{b\} = \hat{\phi}^{-1}(\beta) \subset X \times \hat{B}$.

Let $U \subset \mathbb{P}_k^r$ be as in condition (1) of Proposition 7.2. Since $\hat{\phi}$ is finite and étale over $U \times \hat{B}$, it follows that the map $\tilde{Z} \rightarrow T$ is finite and étale over $T \cap (U \times \hat{B})$. In particular, the map of rings $\mathcal{O}_{T, \beta} \rightarrow \mathcal{O}_{\tilde{Z}, S}$ is finite and étale. This in turn implies that the map $\mathcal{O}_{T, \beta} \rightarrow \mathcal{O}_{Z, Z \cap S}$ is finite and unramified.

On the other hand, by condition (5) of Proposition 7.2 that $\phi_L^{-1}(\phi_L(x))$ is (Z, x) -admissible, we deduce that for each $x \in \Sigma$, the map $\mathcal{O}_{Z, Z \cap S} \rightarrow \mathcal{O}_{Z, \alpha}$ is an isomorphism by Lemma 7.3. Hence, the map $\mathcal{O}_{T, \beta} \rightarrow \mathcal{O}_{Z, \alpha}$ is an injective (since $Z \twoheadrightarrow T$), finite and unramified map of local rings which induces isomorphism between the residue fields (as k is algebraically closed). Lemma 4.10 therefore says that the map $\mathcal{O}_{T, \beta} \rightarrow \mathcal{O}_{Z, \alpha}$ must be an isomorphism.

We next observe that as $\mathcal{O}_{T, \beta} \rightarrow \mathcal{O}_{\tilde{Z}, S}$ is finite and étale, the map $\mathcal{O}_{T, \beta} \rightarrow \mathcal{O}_{\tilde{Z}, \alpha}$ is étale. In particular, the map $\hat{\mathcal{O}}_{T, \beta} \rightarrow \hat{\mathcal{O}}_{\tilde{Z}, \alpha}$ of completions is finite and étale. Since it induces an isomorphism between the residue fields, it follows again from Lemma 4.10 that $\hat{\mathcal{O}}_{T, \beta} \rightarrow \hat{\mathcal{O}}_{\tilde{Z}, \alpha}$ is an isomorphism. Hence, there are local homomorphisms of complete local rings

$$\hat{\mathcal{O}}_{T, \beta} \rightarrow \hat{\mathcal{O}}_{\tilde{Z}, \alpha} \twoheadrightarrow \hat{\mathcal{O}}_{Z, \alpha}, \tag{7-1}$$

where both the first map and the composite map are isomorphisms. Thus, the second map is an isomorphism too. The second map in (7-1) being *a priori* a surjection, the Krull intersection theorem [Matsumura 1986, Theorem 8.10, page 60] says that this map is an isomorphism if and only if $\mathcal{O}_{\tilde{Z}, \alpha} \twoheadrightarrow \mathcal{O}_{Z, \alpha}$ (without completion) is an isomorphism. This in turn is equivalent to that Z is the only irreducible component of \tilde{Z} passing through α , and Z has multiplicity 1 in \tilde{Z} . We have thus proven the lemma. \square

Lemma 7.5. *Let k be algebraically closed and $L \in \mathcal{U}(k) \subset \text{Gr}(N - r - 1, H)(k)$ as in Proposition 7.2. Suppose that Z is irreducible. Then $L^+(Z)$ is regular at all points lying over Σ .*

Proof. We continue with the notations of the proof of Lemma 7.4. Let $\alpha = (x, b) \in X \times \hat{B}$ with $x \in \Sigma$ be such that $\alpha \in L^+(Z)$. Let $\beta = \hat{\phi}(\alpha) = (\phi(x), b) := (y, b)$. It follows from Lemma 7.4 that Z does not pass through α . This implies that the canonical map $\mathcal{O}_{\tilde{Z}, \alpha} \rightarrow \mathcal{O}_{L^+(Z), \alpha}$ is an isomorphism. Therefore, it suffices therefore to show that $\mathcal{O}_{\tilde{Z}, \alpha}$ is regular.

Since $\alpha \in L^+(Z)$, there must exist a closed point $\alpha' = (x', b) \in Z$ with $x' \in \phi^{-1}(y)$. As $\alpha \notin Z$, we must have $x' \neq x$. It follows again from Lemma 7.4 that $\mathcal{O}_{\tilde{Z}, \alpha'} \xrightarrow{\cong} \mathcal{O}_{Z, \alpha'}$. We have also shown in the middle of the proof of Lemma 7.4 that the map $\hat{\mathcal{O}}_{T, \beta} \rightarrow \hat{\mathcal{O}}_{\tilde{Z}, (a, b)}$ of completions in (7-1) is an isomorphism for every $a \in \phi^{-1}(y)$. We thus get the commutative diagram of local rings

$$\begin{array}{ccccc} \mathcal{O}_{\tilde{Z}, \alpha} & \longleftarrow & \mathcal{O}_{T, \beta} & \longrightarrow & \mathcal{O}_{\tilde{Z}, \alpha'} & \xrightarrow{\cong} & \mathcal{O}_{Z, \alpha'} \\ \downarrow & & \downarrow & & \downarrow & & \\ \hat{\mathcal{O}}_{\tilde{Z}, \alpha} & \xleftarrow{\cong} & \hat{\mathcal{O}}_{T, \beta} & \xrightarrow{\cong} & \hat{\mathcal{O}}_{\tilde{Z}, \alpha'}, & & \end{array} \tag{7-2}$$

where the vertical arrows are completion maps.

Since $x' \neq x$, it follows from condition (2) in Definition 7.1 and condition (5) in Proposition 7.2 that $\mathcal{O}_{Z, \alpha'}$ is regular. It follows from (7-2) that all rings of the bottom of the diagram are regular, using a basic fact in commutative algebra that (\star) a noetherian local ring is regular if and only if its completion is a regular local ring [Matsumura 1986, proof of Theorem 19.5, page 157]. Equivalently, all rings of the top of the diagram are regular by (\star) again. In particular, $\mathcal{O}_{\tilde{Z}, \alpha}$ is regular. This finishes the proof. \square

To extend Lemma 7.5 to reducible subschemes Z in Lemma 7.7, we first consider the following:

Lemma 7.6. *Let k be algebraically closed. We are under the Set-up+(fs) of Section 5B. Here, Z is not necessarily irreducible. Then after replacing the embedding $X \hookrightarrow \mathbb{P}_k^N$ into a bigger space via a Veronese embedding if necessary, there is a dense open subset $\mathcal{U} \subset \text{Gr}(X, N - r - 1, H)$ such that for each $L \in \mathcal{U}(k)$, the induced map $\hat{\phi}_L$ takes distinct components of Z to distinct components of $\hat{\phi}_L(Z)$.*

Proof. As we did previously in Lemma 4.12, for each $1 \leq i \leq s$, choose a closed point $\alpha_i = (x_i, b_i) \in Z_i \setminus (\cup_{j \neq i} Z_j)$, so that $x_i := \hat{f}(\alpha_i) \in X_{\text{sm}}$ and $b_i := \hat{g}(\alpha_i) \in B$. We observe that if $j \neq i$, then $Z_j \not\subset X \times \{b_i\}$, because $Z_j \rightarrow \hat{B}$ is nonconstant.

Let $A_i = \bigcup_{j \neq i} \hat{f}(Z_j \cap (X \times \{b_i\}))$. This is a closed subset of dimension $\leq r - 1$. In particular, $\dim(\text{Sec}(A_i, \{x_i\})) \leq r$. Note that $x_i \notin A_i$. Let

$$\mathcal{U} := \text{Gr}(X, N - r - 1, H) \cap \bigcap_{i=1}^s \text{Gr}(\text{Sec}(A_i, \{x_i\}), N - r - 1, H).$$

This is dense open in $\text{Gr}(N - r - 1, H)$ by Lemma 4.3.

Suppose now that $\phi_L : X \rightarrow \mathbb{P}_k^r$ is the projection obtained by any $L \in \mathcal{U}(k)$. We fix an integer $1 \leq i \leq s$ and let $\beta_i := \hat{\phi}_L(\alpha_i)$. It is clear that $\beta_i \in \hat{\phi}_L(Z_i)$. We claim that $\beta_i \notin \hat{\phi}_L(Z_j)$ for $j \neq i$. To see this, note that $\beta_i \in \hat{\phi}_L(Z_j)$ if and only if $Z_j \cap (L^+(x_i) \times \{b_i\}) \neq \emptyset$. Equivalently, there exists a closed point $x'_j \neq x_i$ such that $\phi_L(x'_j) = \phi_L(x_i)$ and $x'_j \in A_i$. But this implies that $L \cap \text{Sec}(A_i, \{x_i\}) \neq \emptyset$, which contradicts the choice of L . This proves the claim and hence the lemma. □

Lemma 7.7. *Let k be algebraically closed. We are under the Set-up+(fs) of Section 5B. Here, Z is not necessarily irreducible. Let $\mathcal{U} \subset \text{Gr}(N - r - 1, H)$ be the intersection of the dense open subsets of Proposition 7.2 and Lemma 7.6. Then for each $L \in \mathcal{U}(k)$, the residual scheme $L^+(Z)$ is regular at all points lying over Σ .*

Proof. For a choice of L , for simplicity write $\phi := \phi_L$. For $1 \leq i \leq s$, let $T_i = \hat{\phi}(Z_i)$ with the reduced closed subscheme structure and let $\tilde{Z}_i = T_i \times_{(\mathbb{P}_k^r \times \hat{B})} (X \times \hat{B}) = \hat{\phi}^{-1}(\hat{\phi}(Z_i))$ as a scheme.

The first claim is that \tilde{Z}_i and \tilde{Z}_j share no common component if $i \neq j$. Indeed, if they do share a common component, this would imply that $T_i = T_j$, which contradicts the choice of L as in Lemma 7.6.

Our second claim is that $L^+(Z_i)$ and $L^+(Z_j)$ do not meet at points lying over Σ if $i \neq j$. Suppose on the contrary that there is a closed point $\alpha = (x, b) \in L^+(Z_i) \cap L^+(Z_j)$ with $x \in \Sigma$. This implies that there are closed points $\alpha_i = (x_i, b) \in Z_i$ and $\alpha_j = (x_j, b) \in Z_j$ such that $x_i, x_j \in \phi^{-1}(y)$, where $y = \phi(x)$. It follows from Lemma 7.4 that $x_i, x_j \in \phi^{-1}(y) \setminus \{x\}$.

If $x_i = x_j$, then two components Z_i and Z_j of Z meet at $\alpha_i = \alpha_j$ that lies over $x_i = x_j$ in $\phi^{-1}(y) \setminus \{x\}$. In particular, Z is singular at a point lying over $x_i = x_j$ in $\phi^{-1}(y) \setminus \{x\}$, which contradicts condition (2) of Definition 7.1, which is part of condition (5) of Proposition 7.2. Hence we must have $x_i \neq x_j$. In this case, we get $b \in \hat{g}(Z_{x_i}) \cap \hat{g}(Z_{x_j}) \neq \emptyset$ for two distinct points $x_i, x_j \in \phi^{-1}(y) \setminus \{x\}$. This time, it contradicts condition (3) of Definition 7.1, which is part of condition (5) of Proposition 7.2. Hence, we proved the second claim.

It follows from the two claims that $L^+(Z)$ is regular at all points lying over Σ if and only if $L^+(Z_i)$ is so for every $1 \leq i \leq s$. Since we proved the latter holds in Lemma 7.5, we finished the proof of the lemma. \square

7C. Regularity of residual cycles: general case. We can now generalize Proposition 7.2 to all infinite perfect field as follows. This includes the regularity of the residual cycle along Σ .

Proposition 7.8. *Let k be any infinite perfect field. We are under the Set-up+(fs) of Section 5B. Let $Y \subset X$ be a closed subset of dimension at most $r - 1$. Then after replacing the embedding $\eta : X \hookrightarrow \mathbb{P}_k^N$ by a bigger one via a Veronese embedding if necessary, we have the following: for the given hyperplane $H \subset \mathbb{P}_k^N$ disjoint from Σ , there is a dense open subset $\mathcal{U} \subset \text{Gr}(N - r - 1, H)$ such that for each $L \in \mathcal{U}(k)$, we have $L \cap X = \emptyset$ so that there is a finite and surjective linear projection $\phi_L : X \rightarrow \mathbb{P}_k^r$. Moreover, it satisfies the following properties:*

- (1) $\hat{\phi}_L(Z_i) \neq \hat{\phi}_L(Z_j)$ if $i \neq j$.
- (2) The map $\phi_L : \phi_L^{-1}(U) \rightarrow U$ is étale for some affine open $U \subset \mathbb{P}_k^r$ containing $\phi_L(\Sigma)$.
- (3) $\phi_L(x) \neq \phi_L(x')$ for each pair of distinct points $x \neq x' \in \Sigma$.
- (4) $k(\phi_L(x)) \xrightarrow{\sim} k(x)$ for each $x \in \Sigma$.
- (5) $L^+(x) \cap Y = \emptyset$ for each $x \in \Sigma$.
- (6) $L^+(Z)$ is regular at all points lying over Σ .
- (7) The map $\hat{\phi}_L : Z \rightarrow \hat{\phi}_L(Z)$ is birational.

Proof. If k is algebraically closed, the proposition follows from Proposition 7.2 and Lemmas 7.6 and 7.7. In general, let \bar{k} be an algebraic closure of k and let $\pi_X : X_{\bar{k}} \rightarrow X$ be the projection map from the base change to \bar{k} . We have $\Sigma_{\bar{k}} = \bigcup_{x \in \Sigma} \pi_X^{-1}(x)$. Choose a sufficiently large closed embedding $\eta : X \hookrightarrow \mathbb{P}_k^N$ so that for the induced embedding $X_{\bar{k}} \hookrightarrow \mathbb{P}_{\bar{k}}^N$, there exists a dense open subset $\tilde{\mathcal{U}} \subset \text{Gr}(N - r - 1, H_{\bar{k}})$ for which all assertions of Proposition 7.2 as well as Lemmas 7.6 and 7.7 applied to $X_{\bar{k}}, Z_{\bar{k}}$ and the set $\Sigma_{\bar{k}} \subset X_{\bar{k}}$ hold. (N.B., Under the base change to \bar{k} , the irreducible components Z_i of Z may decompose further into irreducible components Z_{ij} of $Z_{i,\bar{k}}$. At least $Z_{\bar{k}}$ and $Z_{i,\bar{k}}$ for all i stay reduced because the extension \bar{k} over k is separable.)

Then we can argue via a Galois descent as in the Step 2 of the proof of Lemma 4.8 to find a dense open $\mathcal{U}_1 \subset \text{Gr}(N - r - 1, H)$ defined over k such that $(\mathcal{U}_1)_{\bar{k}} \subset \tilde{\mathcal{U}}$. We take $\mathcal{U} := \mathcal{U}_1 \cap \mathcal{U}_2$, where \mathcal{U}_2 is the open set in Lemma 4.8 so that we can also use the assertions of Lemma 4.8 as well.

Now, for each $L \in \mathcal{U}(k)$, we have $X \cap L = \emptyset$ by our choice of the open set. So, we get a finite linear projection map $\phi_L : X \rightarrow \mathbb{P}_k^r$ over k . We write this map as ϕ . Condition (1) is clear now by construction together with Lemma 7.6. Conditions (2), (3), (4) hold by conditions (2), (1), (3) of Lemma 4.8, respectively. Condition (5) follows immediately from condition (4) of Proposition 7.2.

To prove (6), as we did at the beginning of Section 7B, let $T := \hat{\phi}(Z) = \hat{\phi}(L^+(Z)) \subset \mathbb{P}_k^r \times \hat{B}$ with the reduced subscheme structure, and let $\tilde{Z} := T \times_{(\mathbb{P}_k^r \times \hat{B})} (X \times \hat{B}) = \hat{\phi}^{-1} \hat{\phi}(Z)$ as a scheme. Then we have

the commutative diagram

$$\begin{array}{ccccc}
 Z_{\bar{k}} & \hookrightarrow & \tilde{Z}_{\bar{k}} & \xrightarrow{\hat{\phi}_{\bar{k}}} & T_{\bar{k}} \\
 \downarrow & & \downarrow & & \downarrow \\
 Z & \hookrightarrow & \tilde{Z} & \xrightarrow{\hat{\phi}} & T,
 \end{array} \tag{7-3}$$

where the vertical arrows are the base changes to \bar{k} . Note that the map $\hat{\phi} : Z \rightarrow T$ is surjective by definition. As both squares are Cartesian and the vertical maps are smooth, it follows that $L^+(Z_{\bar{k}}) \xrightarrow{\cong} L^+(Z)_{\bar{k}}$. By the choice of our open set \mathcal{U} , Lemma 7.5 shows that $L^+(Z_{\bar{k}})$ is regular at all points lying over $\Sigma_{\bar{k}}$, i.e., $L^+(Z)_{\bar{k}}$ is regular at all points lying over $\Sigma_{\bar{k}}$.

We replace Z by $Z|_V$ and consider the induced Cartesian squares

$$\begin{array}{ccccc}
 L^+(Z)_{\bar{k}} & \longrightarrow & V_{\bar{k}} & \longrightarrow & \text{Spec}(\bar{k}) \\
 \downarrow & & \downarrow & & \downarrow \\
 L^+(Z) & \longrightarrow & V & \longrightarrow & \text{Spec}(k),
 \end{array} \tag{7-4}$$

where the vertical arrows are the base changes to \bar{k} . Since $L^+(Z)_{\bar{k}}$ is regular and \bar{k} is perfect, the top horizontal composite map is smooth. Hence, by the faithfully flat descent [EGA IV₂ 1965, Corollaire (17.7.3)(ii), page 72], the bottom horizontal composite map is smooth. In particular, $L^+(Z)$ is regular. This proves (6). Property (7) is a direct consequence of Lemma 4.14. \square

Remark 7.9. Condition (4) of Proposition 7.2 or condition (5) of Proposition 7.8 that $L^+(x) \cap Y = \emptyset$ for each $x \in \Sigma$ is no longer needed in this version of the article toward the proof of the main theorems. However, we decided to keep them in this article because the property that the residual points of a projection can be made to avoid the given proper closed subscheme Y is nontrivial, and may be useful in an analysis of algebraic cycles in the future.

8. The main results

In this final section, we use various results of the previous sections to prove our main theorems: the presentation lemma and the sfs-moving lemma. The Set-up for the main results is as in Section 8A. This differs a bit from the Set-up of Section 4D and the Set-up+(fs) of Section 5B.

8A. The Set-up(★). Let k be an infinite perfect field and $n \geq 1$ an integer. We work under the following setting:

(1) *The box coordinates:* For $0 \leq i \leq n - 1$, let \hat{A}_i be a smooth projective geometrically integral k -scheme of positive dimension and let $A_i \subset \hat{A}_i$ be a nonempty affine open subset. Let $C_0 = \text{Spec}(k) = \hat{C}_0$. For $1 \leq j \leq n$, we write $C_j = \prod_{i=0}^{j-1} A_i$ and $\hat{C}_j = \prod_{i=0}^{j-1} \hat{A}_i$. Let $\pi_j : \hat{C}_n \rightarrow \hat{C}_j$ be the projection map. We write $B = C_n$ and $\hat{B} = \hat{C}_n$. Let $F := \hat{B} \setminus B$.

(2) *The base scheme and the cycles:* Let $X \subset \mathbb{A}_k^m$ be an integral smooth affine closed subscheme of dimension $r \geq 1$ and let $\bar{X} \hookrightarrow \mathbb{P}_k^m$ be its closure with the reduced subscheme structure. Let $\Sigma \subset X$ be a finite set of closed points.

Let $Z \subset X \times B$ be a reduced closed subscheme of pure dimension r , and let $\{Z_1, \dots, Z_s\}$ be all of its irreducible components. Suppose $Z \rightarrow X$ is an fs-morphism, i.e., finite and surjective because X is integral. Let $E \subset \hat{B}$ be a closed subset containing F such that no irreducible component of Z is contained in $X \times E$.

Let $\hat{Z} \subset \bar{X} \times \hat{B}$ denote the closure of Z in $\bar{X} \times \hat{B}$ with the reduced structure. Similarly, \hat{Z}_i denotes the closure of Z_i in $\bar{X} \times \hat{B}$. We let $\hat{f} : \hat{Z} \rightarrow \bar{X}$ and $\hat{g} : \hat{Z} \rightarrow \hat{B}$ denote the projection maps.

For each $0 \leq j \leq n$, we define $Z^{(j)} = \pi_j(Z) := (\text{id}_X \times \pi_j)(Z)$. Because $Z \rightarrow X$ is fs, this definition makes sense. Similarly we define $\hat{Z}^{(j)}$ for $0 \leq j \leq n$.

(3) *The linear projections:* Suppose we are given a Veronese embedding $\mathbb{P}_k^m \hookrightarrow \mathbb{P}_k^N$ with $N \gg m$. For $L \in \text{Gr}(\bar{X}, N - r - 1, H)(k)$, where $H = \mathbb{P}_k^N \setminus \mathbb{A}_k^N$ as in Lemma 4.7, let $\phi_L : \bar{X} \rightarrow \mathbb{P}_k^r$ be the linear projection away from L which restricts to a finite map $\phi_L : X \rightarrow \mathbb{A}_k^r$. If L is fixed in a given context, we often drop it from ϕ_L and write ϕ .

For $0 \leq j \leq n$, let $\phi_j = \phi \times \text{id}_{C_j} : X \times C_j \rightarrow \mathbb{A}_k^r \times C_j$, $\tilde{\phi}_j = \phi \times \text{id}_{\hat{C}_j} : X \times \hat{C}_j \rightarrow \mathbb{A}_k^r \times \hat{C}_j$ and $\hat{\phi}_j = \phi \times \text{id}_{\hat{C}_j} : \bar{X} \times \hat{C}_j \rightarrow \mathbb{P}_k^r \times \hat{C}_j$ be the induced maps. We let $L^+(Z)$ denote the closure of $\phi_n^{-1}(\phi_n(Z)) \setminus Z$ in $X \times B$ with the reduced structure. We define $L^+(\hat{Z})$ similarly.

8B. The residual cycle. For $L \in \text{Gr}(\bar{X}, N - r - 1, H)(k)$ as in the Set-up (\star) of Section 8A, the morphism $\phi = \phi_L : X \rightarrow \mathbb{A}_k^r$ is a finite surjective morphism of affine k -schemes so that it is automatically flat by [Hartshorne 1977, Exercise III-10.9, page 276] (or [EGA IV₂ 1965, Proposition (6.1.5), page 136]). Hence, for algebraic cycles on $X \times C_j$, we have the proper push-forward ϕ_{j*} and the flat pull-back ϕ_j^* operations. (See [Fulton 1984, Sections 1.4 and 1.7].) For $X \times \hat{C}_j$, we have similar operations $\tilde{\phi}_{j*}$ and $\tilde{\phi}_j^*$.

Definition 8.1. If $Z \subset X \times C_j$ is an integral closed subscheme, the *residual cycle* by $\phi = \phi_L$ is defined to be

$$L^*([Z]) := \phi_j^* \phi_{j*}([Z]) - [Z].$$

We extend it \mathbb{Z} -linearly to all cycles on $X \times C_j$. Similarly, for cycles on $X \times \hat{C}_j$, we define the *residual cycle* by

$$L^*([Z]) := \tilde{\phi}_j^* \tilde{\phi}_{j*}([Z]) - [Z].$$

Note that by definition, $L^+([Z]) = |L^*([Z])|$.

Lemma 8.2. *We are under the Set-up (\star) of Section 8A. In particular, $Z \rightarrow X$ is an fs-morphism. Suppose that Z is integral. Then for each L in the Set-up (\star) , the morphism $L^+(Z) \rightarrow X$ is also fs.*

Proof. Let $T = \phi_n(Z) \subset \mathbb{A}_k^r \times C_n$. Let $\tilde{Z} := T \times_{(\mathbb{A}_k^r \times C_n)} (X \times C_n) = \phi_n^{-1}(T)$ as a scheme. It suffices to show that the map $\tilde{Z} \rightarrow X$ is fs. Consider the commutative diagram

$$\begin{array}{ccccc}
 Z & \xrightarrow{\iota} & \tilde{Z} & \xrightarrow{\phi_n} & T \\
 & \searrow & \downarrow \hat{f} & & \downarrow \hat{f}' \\
 & & X & \xrightarrow{\phi} & \mathbb{A}_k^r,
 \end{array} \tag{8-1}$$

where the vertical arrows are the projection maps and the right square is Cartesian, and ι is the closed immersion.

Since $Z \rightarrow X$ is an fs-morphism and ϕ is an fs-morphism, the composite $(\phi \circ \hat{f})|_Z = \phi \circ \hat{f} \circ \iota$ is an fs-morphism. By the commutativity, this means $\hat{f}' \circ \phi_n \circ \iota$ is an fs-morphism. But since $Z \rightarrow T$ is surjective (as T being the image of Z under ϕ_n by definition), it follows that \hat{f}' is finite (e.g., see [Liu 2002, Proposition 3.16(f), page 104]). Hence \hat{f}' is an fs-morphism. Now, ϕ is flat, so by Lemma 2.7, the morphism \hat{f} is an fs-morphism. □

Lemma 8.3. *We are under the Set-up(★) of Section 8A. In particular, $Z \rightarrow X$ is an fs-morphism. Suppose that Z is integral. Suppose that there is an integer $1 \leq j \leq n$ such that the projection map $Z^{(j)} \rightarrow C_j$ is nonconstant.*

Then for each $L \in \text{Gr}(N - r - 1, H)(k)$ satisfying Lemma 4.14 and Proposition 7.8 for all $Z^{(i)}$ over $j \leq i \leq n$, we have the equalities

$$[L^+(Z^{(j)})] = L^*([Z^{(j)}]) \quad \text{and} \quad \pi_{j*}(L^*([Z])) = m_j L^*([Z^{(j)}]),$$

where $m_j = [k(Z) : k(Z^{(j)})]$.

Proof. First of all, note that by Lemma 8.2, every component of $L^+(Z)$ is fs over X . In particular, by the finiteness criterion Lemma 2.9, each irreducible component of $L^+(Z)$ is closed in $X \times \hat{C}_n$. The push-forward $\pi_{j*}([L^*(Z)])$ is given by the projective map $\pi_j : X \times \hat{C}_n \rightarrow X \times \hat{C}_j$ is projective.

To prove the first equality, replacing Z by $Z^{(j)}$, we may assume $n = j$ and $Z^{(j)} = Z$. Let $T = \tilde{\phi}_n(Z) \subset \mathbb{A}_k^r \times \hat{C}_n$.

Note that the map $Z \rightarrow T$ is birational by Lemma 4.14. Hence, by the definition of the proper push-forward and flat pull-back of cycles, the first equality is equivalent to showing that $\tilde{Z} := T \times_{(\mathbb{A}_k^r \times \hat{C}_n)} (X \times \hat{C}_n) = \tilde{\phi}_n^{-1}(T)$ is a reduced scheme.

To show that \tilde{Z} is reduced, let $U \subset \mathbb{A}_k^r$ be an affine open neighborhood of Σ as in condition (3) of Proposition 7.8. Since $Z \rightarrow X$ is finite and surjective, the open subset $T \cap (U \times \hat{C}_n)$ is dense in T . The map $\tilde{\phi}_n$ is étale over this dense open subset of T . Hence, $\tilde{Z} = \tilde{\phi}_n^{-1}(T)$ is reduced over this dense open subset of T . However, $\tilde{\phi}_n^{-1}(T) \rightarrow T$ is finite and flat everywhere, it means $\tilde{Z} = \tilde{\phi}_n^{-1}(T)$ is reduced. This proves the first equality.

For the second equality, consider the commutative diagram

$$\begin{array}{ccc}
 X \times \hat{C}_n & \xrightarrow{\tilde{\phi}_n} & \mathbb{A}_k^r \times \hat{C}_n \\
 \pi_j \downarrow & & \downarrow \pi_j \\
 X \times \hat{C}_j & \xrightarrow{\tilde{\phi}_j} & \mathbb{A}_k^r \times \hat{C}_j.
 \end{array} \tag{8-2}$$

This is a Cartesian square in which the vertical arrows are projective and the horizontal arrows are finite and flat. Hence, by [Fulton 1984, Proposition 1.7], we have

$$\begin{aligned}
 \pi_{j*}(L^*([Z])) &= \pi_{j*}(\tilde{\phi}_n^* \circ \tilde{\phi}_{n*}([Z]) - [Z]) \\
 &= \pi_{j*} \circ \tilde{\phi}_n^* \circ \tilde{\phi}_{n*}([Z]) - \pi_{j*}([Z]) \\
 &= \tilde{\phi}_j^* \circ \pi_{j*} \circ \tilde{\phi}_{n*}([Z]) - \pi_{j*}([Z]) \\
 &= \tilde{\phi}_j^* \circ \tilde{\phi}_{j*} \circ \pi_{j*}([Z]) - \pi_{j*}([Z]) \\
 &= \tilde{\phi}_j^* \circ \tilde{\phi}_{j*}(m_j[Z^{(j)}]) - m_j[Z^{(j)}] \\
 &= m_j(\tilde{\phi}_j^* \circ \tilde{\phi}_{j*}([Z^{(j)}]) - [Z^{(j)}]) \\
 &= m_j L^*([Z^{(j)}]),
 \end{aligned}$$

which proves the second equality. □

The following complements Lemma 8.3:

Lemma 8.4. *We are under the Set-up(★) of Section 8A. In particular, $Z \rightarrow X$ is an fs-morphism. Suppose that Z is integral such that $Z \rightarrow C_n$ is nonconstant. Suppose that for an integer $0 \leq j \leq n - 1$, the projection $Z^{(j)} \rightarrow C_j$ is constant. Then for each $L \in \text{Gr}(N - r - 1, H)(k)$ satisfying the conditions of Proposition 7.8 for Z , we have the equality $\pi_j(Z') = \pi_j(Z)$ for each irreducible component Z' of $L^+(Z)$.*

Proof. Toward contradiction, suppose that there is an irreducible component Z' of $L^+(Z)$ such that $\pi_j(Z') \neq \pi_j(Z)$. In particular, this implies that $L^+(Z^{(j)}) \neq \emptyset$. On the other hand, we are given that $Z^{(j)} = \pi_j(Z) = X \times \{c_j\}$ for some closed point $c_j \in \hat{C}_j$. In this case, $\tilde{\phi}_j(Z^{(j)}) = \mathbb{A}_k^r \times \{c_j\}$ so that $\tilde{\phi}_j^{-1} \tilde{\phi}_j(Z^{(j)}) = X \times \{c_j\} = Z^{(j)}$. Hence, $L^+(Z^{(j)}) = \emptyset$. This is a contradiction. □

8C. The presentation lemma. We now prove the presentation lemma for residual cycles under linear projections. We are under the Set-up(★) in Section 8A.

Theorem 8.5. *Let k be an infinite perfect field. Let $Z \subset X \times C_n$ be an integral closed subscheme such that $Z \rightarrow X$ is finite surjective, and the projection $Z \rightarrow C_n$ is nonconstant.*

Then there exist an embedding $\eta : \bar{X} \hookrightarrow \mathbb{P}_k^N$ and a dense open subset $\mathcal{U} \subset \text{Gr}(N - r - 1, H)$, where $H := \mathbb{P}_k^N \setminus \mathbb{A}_k^N$, such that for each $L \in \mathcal{U}(k)$, the linear projection $\phi_L : \mathbb{P}_k^N \setminus L \rightarrow \mathbb{P}_k^r$ away from L defines a finite surjective morphism $\phi : \bar{X} \rightarrow \mathbb{P}_k^r$ satisfying the following properties:

(1) *There exists a Cartesian square*

$$\begin{array}{ccc}
 X & \hookrightarrow & \bar{X} \\
 \downarrow \phi & & \downarrow \phi \\
 \mathbb{A}_k^r & \hookrightarrow & \mathbb{P}_k^r.
 \end{array}$$

(2) *ϕ is étale over an affine open neighborhood of $\phi(\Sigma)$.*

(3) *$\phi(x) \neq \phi(x')$ for every pair $x \neq x'$ in Σ .*

(4) *The map $k(\phi(x)) \rightarrow k(x)$ is an isomorphism for each $x \in \Sigma$.*

(5) *The induced map $Z \rightarrow \phi_n(Z)$ is birational.*

(6) *The map $L^+(Z) \rightarrow X$ is finite surjective.*

(7) *For each $0 \leq j \leq n$, the scheme $\pi_j(L^+(Z))$ is regular at all points lying over Σ .*

Proof. Since $Z \rightarrow X$ is finite surjective, for each $0 \leq j \leq n$ the morphism $Z^{(j)} \rightarrow X$ is also finite surjective. Let $i_0 \in \{0, \dots, n\}$ be the largest integer i such that $Z^{(i)} \subset X \times \{b\}$ for some closed point $b \in C_i$. Note that $Z^{(0)} = X = X \times_k C_0$, so such i_0 exists. Note also that $i_0 \leq n - 1$ by our assumption.

Choose a large enough Veronese embedding $\mathbb{P}_k^m \hookrightarrow \mathbb{P}_k^N$ such that for the composite embedding $\eta : \bar{X} \hookrightarrow \mathbb{P}_k^N$, and the hyperplane $H = H_{N,0}$ as in Lemma 4.7, there are open dense subsets $\mathcal{U}_j \subset \text{Gr}(N - r - 1, H)$ such that each $L \in \mathcal{U}_j(k)$ satisfies Lemma 4.14 and conditions (1)–(6) of Proposition 7.8 for $Z^{(j)}$ over all $i_0 + 1 \leq j \leq n$. We let $\mathcal{U} = \bigcap_{j=i_0+1}^n \mathcal{U}_j$.

Condition (1) of the theorem automatically follows from our choice of H and Lemma 4.7. Conditions (2)–(4) follow directly from conditions (2)–(4) of Proposition 7.8. Condition (5) follows from Lemma 4.14. Condition (6) follows from Lemma 8.2.

We prove (7). We have to show that every irreducible component of $\pi_j(L^+(Z))$ is regular at all points lying over Σ and no two components of $\pi_j(L^+(Z))$ meet at points lying over Σ . We first assume that $i_0 + 1 \leq j \leq n$.

Let Z' be an irreducible component of $L^+(Z)$. Since $j > i_0$, Lemma 8.3 says that $\pi_j(Z')$ is a component of the effective cycle $\pi_{j*}(L^*([Z])) = m_j L^*([Z^{(j)}]) = m_j [L^+(Z^{(j)})]$ with $m_j \geq 1$. Since Z' was arbitrary, it follows that the irreducible components of $\pi_j(L^+(Z))$ are the same as those of $L^+(Z^{(j)})$. On the other hand, condition (6) of Proposition 7.8 (with our choice of L) says that $L^+(Z^{(j)})$ is regular at all points lying over Σ . It follows that each irreducible component of $\pi_j(L^+(Z))$ is regular at all points lying over Σ , and in particular no two components meet at points lying over Σ .

If $0 \leq j \leq i_0$, then Lemma 8.4 says that $\pi_j(L^+(Z))$ coincides with $\pi_j(Z)$, which in turn is of the form $X \times \{b\}$ for some closed point $b \in C_j$. In particular, $\pi_j(L^+(Z))$ is irreducible. As X is regular everywhere, in particular at all points lying over Σ , it follows that $\pi_j(L^+(Z))$ is regular at all points lying over Σ . This completes the proof of the theorem. □

8D. The sfs-moving lemma. We now prove the sfs-moving lemma for additive higher Chow groups of relative 0-cycles over semilocal k -schemes. A similar argument also proves the sfs-moving lemma for Bloch’s higher Chow groups of relative 0-cycles over semilocal k -schemes.

Let k be an infinite perfect field. We apply Theorem 8.5 with $\hat{A}_i := \mathbb{P}_k^1$ for $0 \leq i \leq n - 1$, while $A_0 := \mathbb{A}_k^1$ and $A_1 = \cdots = A_{n-1} = \square_k^1$ so that $C_j = B_j = \mathbb{A}_k^1 \times \square_k^{j-1}$ for $j \geq 1$. The sfs-moving lemma for additive higher Chow groups of relative 0-cycles is the following:

Theorem 8.6. *Let R be a regular semilocal k -scheme essentially of finite type of dimension $r \geq 0$ over an infinite perfect field k . Let $V = \text{Spec}(R)$ and let $m, n \geq 1$ be integers. Then the canonical map $\text{TCH}_{\text{sfs}}^n(V, n; m) \rightarrow \text{TCH}^n(V, n; m)$ is an isomorphism.*

This theorem is proven in steps. Since R is regular, it is a product of regular semilocal k -domains, and each k -domain corresponds to a connected component of $\text{Spec}(R)$. Thus we may reduce to the case when R is integral. We also remark that by Proposition 2.19, we may assume that R is obtained by localizing an integral smooth affine k -scheme at a finite set of closed points. Note that Theorem 8.6 is obvious for $r = 0$, so we may assume $r \geq 1$. We have injective maps (using Lemma 2.18),

$$\text{TCH}_{\text{sfs}}^n(V, n; m) \rightarrow \text{TCH}_{\text{fs}}^n(V, n; m) \rightarrow \text{TCH}_{\Sigma}^n(V, n; m) \rightarrow \text{TCH}^n(V, n; m).$$

The last arrow is an isomorphism by [Krishna and Park 2016, Theorem 4.10]. We show that the middle arrow is an isomorphism, which we call the *fs-moving lemma*:

Lemma 8.7. *The map $\text{TCH}_{\text{fs}}^n(V, n; m) \rightarrow \text{TCH}_{\Sigma}^n(V, n; m)$ is an isomorphism.*

Proof. By the above discussion, we assume V is integral. Since this map is injective, we only have to show that it is surjective. Let $\gamma \in \text{Tz}_{\Sigma}^n(V, n; m)$ be a cycle with $\partial(\gamma) = 0$.

First suppose that there is an atlas (\mathbb{A}_k^r, Σ) so that γ lifts to a cycle $\bar{\gamma} \in \text{Tz}_{\Sigma}^n(\mathbb{A}_k^r, n; m)$. In this case, we can apply Theorem 3.14 and write $\gamma = \gamma_1 + \partial(\gamma_2)$, where $\gamma_1 \in \text{Tz}_{\text{sfs}}^n(V, n; m) \subset \text{Tz}_{\text{fs}}^n(V, n; m)$ and $\gamma_2 \in \text{Tz}^n(V, n + 1; m)$. One immediately has $\partial(\gamma_1) = 0$, proving the desired surjectivity in this case.

In general, we write $\gamma = \alpha + \beta$, where no component of α is an fs-cycle and β is an fs-cycle. Lemma 2.5 says that there is a connected smooth affine atlas (X, Σ) for V , and cycles $\bar{\alpha}, \bar{\beta}, \bar{\gamma} \in \text{Tz}_{\Sigma}^n(X, n; m)$ such that $\bar{\alpha}_V = \alpha, \bar{\beta}_V = \beta, \bar{\gamma}_V = \gamma, \bar{\gamma} = \bar{\alpha} + \bar{\beta}$ and $\partial(\bar{\gamma}) = 0$.

Since no component of α is fs over V , it follows that the projection of every component of $\bar{\alpha}$ to B_n must be nonconstant. We can therefore apply Theorem 4.15 to obtain a finite flat map $\phi : X \rightarrow \mathbb{A}_k^r$ such that α satisfies all the properties there. Let $\Sigma' = \phi(\Sigma)$, which consists of finitely many closed points of \mathbb{A}_k^r . Let $V' = \text{Spec}(\mathcal{O}_{\mathbb{A}_k^r, \Sigma'})$ and $W := X \times_{\mathbb{A}_k^r} V'$. We have inclusions $\Sigma \subset V \subset W \subset X$, and a finite flat morphism $\phi : W \rightarrow V'$.

Write $\bar{\alpha} = \bar{\alpha}_1 + \bar{\alpha}_2$, where each component of $\bar{\alpha}_1$ is dominant over X and no component of $\bar{\alpha}_2$ is dominant over X . As β is an fs-cycle over V , after shrinking X if needed, $\bar{\beta}$ is an fs-cycle over X along Σ by Corollary 2.12.

We now have

$$\bar{\gamma} = \bar{\alpha}_1 + \bar{\alpha}_2 + \bar{\beta} = (\bar{\alpha}_1 - \phi_n^* \phi_{n*}(\bar{\alpha}_1)) + (\bar{\alpha}_2 - \phi_n^* \phi_{n*}(\bar{\alpha}_2)) + (\bar{\beta} - \phi_n^* \phi_{n*}(\bar{\beta})) + \phi_n^* \phi_{n*}(\bar{\gamma}).$$

Let $\bar{\alpha}'_i := \bar{\alpha}_i - \phi_n^* \phi_{n*}(\bar{\alpha}_i)$ for $i = 1, 2$, and $\bar{\beta}' := \bar{\beta} - \phi_n^* \phi_{n*}(\bar{\beta})$. Since $\bar{\beta}$ is an fs-cycle on X along Σ and ϕ is finite, $\phi_{n*}(\bar{\beta})$ is an fs-cycle over \mathbb{A}_k^r by Lemma 2.8. Since $X \rightarrow \mathbb{A}_k^r$ is flat, by Lemma 2.7 $\phi_n^* \phi_{n*}(\bar{\beta})$ is an fs-cycle over X along Σ . In particular, $\bar{\beta}' \in \text{Tz}_{\Sigma, \text{fs}}^n(X, n; m)$. On the other hand, by Theorem 4.15, we have $(\bar{\alpha}'_2)_V = 0$ and $(\bar{\alpha}'_1)_V \in \text{Tz}_{\Sigma, \text{fs}}^n(V, n; m)$.

Since $\bar{\gamma} \in \text{Tz}_{\Sigma}^n(X, n; m)$ with $\partial(\bar{\gamma}) = 0$, it follows that $\phi_*(\bar{\gamma}) \in \text{Tz}_{\Sigma'}^n(\mathbb{A}_k^r, n; m)$ with $\partial(\phi_{n*}(\bar{\gamma})) = 0$. By the previous case, there are cycles $\eta_1 \in \text{Tz}_{\text{fs}}^n(V', n; m)$, and $\eta_2 \in \text{Tz}^n(V', n + 1; m)$ such that $j^*(\phi_{n*}(\bar{\gamma})) = \eta_1 + \partial\eta_2$. Equivalently, $\phi_{n*}(\bar{\gamma}_W) = \eta_1 + \partial\eta_2$. Hence, $\phi_n^* \phi_{n*}(\bar{\gamma}_W) = \phi_n^*(\eta_1) + \phi_n^*(\partial\eta_2) = \phi_n^*(\eta_1) + \partial(\phi_n^*(\eta_2))$. Moreover, $\phi_n^*(\eta_1)$ is an fs-cycle by Lemma 2.7. Combining these, we have

$$\gamma = (\bar{\gamma})_V = (\bar{\alpha}'_1)_V + \bar{\beta}'_V + (\phi_n^*(\eta_1))_V + \partial((\phi_n^*(\eta_2))_V) = \gamma_1 + \partial((\phi_n^*(\eta_2))_V),$$

where $\gamma_1 := (\bar{\alpha}'_1)_V + \bar{\beta}'_V + (\phi_n^*(\eta_1))_V \in \text{Tz}_{\text{fs}}^n(V, n; m)$. Since $\partial\gamma = 0$, we also deduce that $\partial\gamma_1 = 0$. This completes the proof of the lemma. \square

Proof of Theorem 8.6. We may assume that V is integral. Using Lemma 8.7, it suffices to show that the map $\text{TCH}_{\text{sfs}}^n(V, n; m) \rightarrow \text{TCH}_{\text{fs}}^n(V, n; m)$ is surjective. Let $\alpha \in \text{Tz}_{\text{fs}}^n(V, n; m)$ be an fs-cycle, which always has $\partial(\alpha) = 0$ by Lemma 2.21. Write $\alpha = \alpha_1 + \alpha_2$, where $\alpha_2 \in \text{Tz}_{\text{sfs}}^n(V, n; m)$, while $\alpha_1 \in \text{Tz}_{\text{fs}}^n(V, n; m)$ but no component of α_1 lies in $\text{Tz}_{\text{sfs}}^n(V, n; m)$. Note that $\partial(\alpha_i) = 0$ for $i = 1, 2$ by Lemma 2.21 again. It is enough to prove that α_1 is equivalent to a cycle in $\text{Tz}_{\text{sfs}}^n(V, n; m)$. Replacing α by α_1 , we may therefore assume that no component of α lies in $\text{Tz}_{\text{sfs}}^n(V, n; m)$.

Apply Lemma 2.5 to choose a connected smooth affine atlas (X, Σ) for V and a cycle $\bar{\alpha} \in \text{Tz}_{\Sigma}^n(X, n; m)$ such that $\partial(\bar{\alpha}) = 0$. If $X \simeq \mathbb{A}_k^r$, we can apply Theorem 3.14 to write $\alpha = \beta + \partial(\gamma)$, where $\beta \in \text{Tz}_{\text{sfs}}^n(V, n; m) \subset \text{Tz}_{\text{fs}}^n(V, n; m)$ and $\gamma \in \text{Tz}^n(V, n + 1; m)$. This solves the problem in this case.

Suppose that X is not an affine space. If Z is a component of α whose projection to B_n is constant, then Z is already an sfs-cycle. But, we supposed no component of α is an sfs-cycle. Hence, $Z \rightarrow B_n$ is nonconstant for each irreducible component Z . It follows that Lemma 8.3 and Theorem 8.5 apply to every component of α . Let $\phi : X \rightarrow \mathbb{A}_k^r$ be the finite and flat map as in Theorem 8.5 and let $\Sigma' = \phi(\Sigma)$. By shrinking $\mathcal{U} \subset \text{Gr}(N - r - 1, H)$ if necessary, we can assume that conditions (1)–(7) of Theorem 8.5 hold for each $L \in \mathcal{U}(k)$ and for each component of α .

Let $V' = \text{Spec}(\mathcal{O}_{\mathbb{A}_k^r, \Sigma'})$ and let $W = X \times_{\mathbb{A}_k^r} V'$. We have inclusions $\Sigma \subset V \subset W \subset X$ and a finite and flat morphism $\phi_{\Sigma} : W \rightarrow V'$ of smooth semilocal k -schemes. Let $j : V \rightarrow W$ be the localization map.

We can write $\bar{\alpha}_W = (\bar{\alpha}_W - \phi_n^* \phi_{n*}(\bar{\alpha}_W)) + \phi_n^* \phi_{n*}(\bar{\alpha}_W)$. We have $\partial(\phi_{n*}(\bar{\alpha}_W)) = \phi_{n*}(\partial(\bar{\alpha}_W)) = 0$. By the previous case of affine space atlas, we can write $\phi_{n*}(\bar{\alpha}_W) = \eta_1 + \partial(\eta_2)$, where $\eta_1 \in \text{Tz}_{\text{sfs}}^n(V', n; m)$ and $\eta_2 \in \text{Tz}^n(V', n + 1; m)$. This yields $\phi_n^* \phi_{n*}(\bar{\alpha}_W) = \phi_n^*(\eta_1) + \partial(\phi_n^*(\eta_2))$. Since $\phi : W \rightarrow V'$ is finite and étale, it follows by Lemmas 2.7 and 2.16 that $\phi_n^*(\eta_1) \in \text{Tz}_{\text{sfs}}^n(W, n; m)$.

It follows from Lemma 8.3 and Theorem 8.5 that $j^*(\bar{\alpha}_W - \phi_n^* \phi_{n*}(\bar{\alpha}_W)) \in \text{Tz}_{\text{sfs}}^n(V, n; m)$. Let $\beta = j^*(\bar{\alpha}_W - \phi_n^* \phi_{n*}(\bar{\alpha}_W)) + j^*(\phi_n^*(\eta_1)) \in \text{Tz}_{\text{sfs}}^n(V, n; m)$ and $\gamma = j^*(\phi_n^*(\eta_2)) \in \text{Tz}^n(V, n + 1; m)$. Then, we

get

$$\begin{aligned}\alpha &= j^*(\bar{\alpha}_W) = j^*(\bar{\alpha}_W - \phi_n^* \phi_{n*}(\bar{\alpha}_W)) + j^* \phi_n^*(\eta_1) + j^*(\partial(\phi_n^*(\eta_2))) \\ &= j^*(\bar{\alpha}_W - \phi_n^* \phi_{n*}(\bar{\alpha}_W)) + j^* \phi_n^*(\eta_1) + \partial(j^* \phi_n^*(\eta_2)) \\ &= \beta + \partial(\gamma).\end{aligned}$$

Since $\partial(\alpha) = 0$, we must have $\partial(\beta) = 0$ as well. This proves the theorem. \square

Proof of Theorem 1.2. We take $n \geq 2$, $A_0 = \hat{A}_0 = \text{Spec}(k)$, $A_i = \square_k$ and $\hat{A}_i := \mathbb{P}_k^1$ for $1 \leq i \leq n-1$ in Theorem 8.5. We now repeat the proof of Theorem 8.6 verbatim using Remark 3.15. \square

Acknowledgments

We thank Spencer Bloch, H el ene Esnault, Najmuddin Fakhruddin, Bruno Kahn, Marc Levine, and Kay R ulling for some conversations related to the question studied here throughout the time we were working on it. We thank the handling editors and, the anonymous referee of Algebra & Number Theory, who read the article with thoroughness and patience, and supplied numerous suggestions, which helped the authors in improving the article.

We acknowledge that part of this work was done during Park’s visits to TIFR and Krishna’s visits to KAIST, and we thank both the institutions. Park would like to thank Juya and Damy for his peace of mind at home through the time this paper was written. Park was partially supported by the National Research Foundation of Korea (NRF) grant (2015R1A2A2A01004120 and 2018R1A2B6002287) funded by the Korean government (MSIP), and TJ Park Junior Faculty Fellowship funded by POSCO TJ Park Foundation.

References

- [Arbarello et al. 1985] E. Arbarello, M. Cornalba, P. A. Griffiths, and J. Harris, *Geometry of algebraic curves, I*, Grundlehren der Math. Wissenschaften **267**, Springer, 1985. MR Zbl
- [Ballico 1989] E. Ballico, “On singular curves in the case of positive characteristic”, *Math. Nachr.* **141** (1989), 267–273. MR Zbl
- [Bloch 1986] S. Bloch, “Algebraic cycles and higher K -theory”, *Adv. Math.* **61**:3 (1986), 267–304. MR Zbl
- [Bloch 1994] S. Bloch, “The moving lemma for higher Chow groups”, *J. Algebraic Geom.* **3**:3 (1994), 537–568. MR Zbl
- [Colliot-Th el ene et al. 1997] J.-L. Colliot-Th el ene, R. T. Hoobler, and B. Kahn, “The Bloch–Ogus–Gabber theorem”, pp. 31–94 in *Algebraic K-theory* (Toronto, 1996), edited by V. P. Snaith, Fields Inst. Commun. **16**, Amer. Math. Soc., Providence, RI, 1997. MR Zbl
- [EGA IV₂ 1965] A. Grothendieck, “El ements de g eom etrie alg ebrique, IV:  tude locale des sch emas et des morphismes de sch emas, II”, *Inst. Hautes  tudes Sci. Publ. Math.* **24** (1965), 5–231. MR Zbl
- [EGA IV₃ 1966] A. Grothendieck, “El ements de g eom etrie alg ebrique, IV:  tude locale des sch emas et des morphismes de sch emas, III”, *Inst. Hautes  tudes Sci. Publ. Math.* **28** (1966), 5–255. MR Zbl
- [EGA IV₄ 1967] A. Grothendieck, “El ements de g eom etrie alg ebrique, IV:  tude locale des sch emas et des morphismes de sch emas, IV”, *Inst. Hautes  tudes Sci. Publ. Math.* **32** (1967), 5–361. MR Zbl
- [Elbaz-Vincent and M uller-Stach 2002] P. Elbaz-Vincent and S. M uller-Stach, “Milnor K -theory of rings, higher Chow groups and applications”, *Invent. Math.* **148**:1 (2002), 177–206. MR Zbl
- [Fulton 1984] W. Fulton, *Intersection theory*, Ergebnisse der Mathematik (3) **2**, Springer, 1984. MR Zbl

- [Gabber et al. 2013] O. Gabber, Q. Liu, and D. Lorenzini, “The index of an algebraic variety”, *Invent. Math.* **192**:3 (2013), 567–626. MR Zbl
- [Gupta and Krishna 2019a] R. Gupta and A. Krishna, “Relative K -theory via 0-cycles in finite characteristic”, preprint, 2019. arXiv
- [Gupta and Krishna 2019b] R. Gupta and A. Krishna, “Zero-cycles with modulus and relative K -theory”, preprint, 2019. arXiv
- [Harris 1982] J. Harris, *Curves in projective space*, Sémin. Math. Sup. **85**, Univ. Montreal Press, 1982. MR Zbl
- [Hartshorne 1977] R. Hartshorne, *Algebraic geometry*, Graduate Texts in Math. **52**, Springer, 1977. MR Zbl
- [Jouanolou 1983] J.-P. Jouanolou, *Théorèmes de Bertini et applications*, Prog. Math. **42**, Birkhäuser, Boston, 1983. MR Zbl
- [Kai 2019] W. Kai, “A moving lemma for algebraic cycles with modulus and contravariance”, *Int. Math. Res. Not.* (online publication May 2019).
- [Kleiman and Altman 1979] S. L. Kleiman and A. B. Altman, “Bertini theorems for hypersurface sections containing a subscheme”, *Comm. Algebra* **7**:8 (1979), 775–790. MR Zbl
- [Krishna and Levine 2008] A. Krishna and M. Levine, “Additive higher Chow groups of schemes”, *J. Reine Angew. Math.* **619** (2008), 75–140. MR Zbl
- [Krishna and Park 2012] A. Krishna and J. Park, “Moving lemma for additive higher Chow groups”, *Algebra Number Theory* **6**:2 (2012), 293–326. MR Zbl
- [Krishna and Park 2015a] A. Krishna and J. Park, “de Rham-Witt sheaves via algebraic cycles”, preprint, 2015. arXiv
- [Krishna and Park 2015b] A. Krishna and J. Park, “DGA-structure on additive higher Chow groups”, *Int. Math. Res. Not.* **2015**:1 (2015), 1–54. MR Zbl
- [Krishna and Park 2016] A. Krishna and J. Park, “On additive higher Chow groups of affine schemes”, *Doc. Math.* **21** (2016), 49–89. MR Zbl
- [Krishna and Park 2017] A. Krishna and J. Park, “A moving lemma for cycles with very ample modulus”, *Ann. Sc. Norm. Super. Pisa Cl. Sci. (5)* **17**:4 (2017), 1521–1549. MR Zbl
- [Liu 2002] Q. Liu, *Algebraic geometry and arithmetic curves*, Oxford Grad. Texts Math. **6**, Oxford Univ. Press, 2002. MR Zbl
- [Matsumura 1986] H. Matsumura, *Commutative ring theory*, Cambridge Stud. Adv. Math. **8**, Cambridge Univ. Press, 1986. MR Zbl
- [Park 2009] J. Park, “Regulators on additive higher Chow groups”, *Amer. J. Math.* **131**:1 (2009), 257–276. MR Zbl
- [Rülling 2007] K. Rülling, “The generalized de Rham–Witt complex over a field is a complex of zero-cycles”, *J. Algebraic Geom.* **16**:1 (2007), 109–169. MR Zbl
- [Samuel 1966] P. Samuel, *Lectures on old and new results on algebraic curves*, Tata Inst. Fund. Res. Lect. Math. **36**, Tata Inst. Fund. Res., Bombay, 1966. MR Zbl

Communicated by Hélène Esnault

Received 2019-04-13 Revised 2019-11-18 Accepted 2019-12-16

amal@math.tifr.res.in

School of Mathematics, Tata Institute of Fundamental Research, Mumbai, India

jinhyun@kaist.edu

Department of Mathematical Sciences, KAIST, Daejeon, South Korea

Algebra & Number Theory

msp.org/ant

EDITORS

MANAGING EDITOR

Bjorn Poonen
Massachusetts Institute of Technology
Cambridge, USA

EDITORIAL BOARD CHAIR

David Eisenbud
University of California
Berkeley, USA

BOARD OF EDITORS

Bhargav Bhatt	University of Michigan, USA	Martin Olsson	University of California, Berkeley, USA
Richard E. Borcherds	University of California, Berkeley, USA	Raman Parimala	Emory University, USA
Antoine Chambert-Loir	Université Paris-Diderot, France	Jonathan Pila	University of Oxford, UK
J-L. Colliot-Thélène	CNRS, Université Paris-Sud, France	Irena Peeva	Cornell University, USA
Brian D. Conrad	Stanford University, USA	Anand Pillay	University of Notre Dame, USA
Samit Dasgupta	Duke University, USA	Michael Rapoport	Universität Bonn, Germany
Hélène Esnault	Freie Universität Berlin, Germany	Victor Reiner	University of Minnesota, USA
Gavril Farkas	Humboldt Universität zu Berlin, Germany	Peter Sarnak	Princeton University, USA
Hubert Flenner	Ruhr-Universität, Germany	Michael Singer	North Carolina State University, USA
Sergey Fomin	University of Michigan, USA	Christopher Skinner	Princeton University, USA
Edward Frenkel	University of California, Berkeley, USA	Vasudevan Srinivas	Tata Inst. of Fund. Research, India
Wee Teck Gan	National University of Singapore	J. Toby Stafford	University of Michigan, USA
Andrew Granville	Université de Montréal, Canada	Shunsuke Takagi	University of Tokyo, Japan
Ben J. Green	University of Oxford, UK	Pham Huu Tiep	University of Arizona, USA
Joseph Gubeladze	San Francisco State University, USA	Ravi Vakil	Stanford University, USA
Christopher Hacon	University of Utah, USA	Michel van den Bergh	Hasselt University, Belgium
Roger Heath-Brown	Oxford University, UK	Akshay Venkatesh	Institute for Advanced Study, USA
János Kollár	Princeton University, USA	Marie-France Vignéras	Université Paris VII, France
Philippe Michel	École Polytechnique Fédérale de Lausanne	Melanie Matchett Wood	University of California, Berkeley, USA
Susan Montgomery	University of Southern California, USA	Shou-Wu Zhang	Princeton University, USA
Shigefumi Mori	RIMS, Kyoto University, Japan		

PRODUCTION

production@msp.org
Silvio Levy, Scientific Editor

See inside back cover or msp.org/ant for submission instructions.

The subscription price for 2020 is US \$415/year for the electronic version, and \$620/year (+\$60, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to MSP.

Algebra & Number Theory (ISSN 1944-7833 electronic, 1937-0652 printed) at Mathematical Sciences Publishers, 798 Evans Hall #3840, c/o University of California, Berkeley, CA 94720-3840 is published continuously online. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices.

ANT peer review and production are managed by EditFLOW® from MSP.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2020 Mathematical Sciences Publishers

Algebra & Number Theory

Volume 14 No. 4 2020

The distribution of p -torsion in degree p cyclic fields	815
JACK KLYS	
On the motivic class of an algebraic group	855
FEDERICO SCAVIA	
A representation theory approach to integral moments of L -functions over function fields	867
WILL SAWIN	
Deformations of smooth complete toric varieties: obstructions and the cup product	907
NATHAN ILTEN and CHARLES TURO	
Mass equidistribution on the torus in the depth aspect	927
YUEKE HU	
The basepoint-freeness threshold and syzygies of abelian varieties	947
FEDERICO CAUCCI	
On the Ekedahl–Oort stratification of Shimura curves	961
BENJAMIN HOWARD	
A moving lemma for relative 0-cycles	991
AMALENDU KRISHNA and JINHYUN PARK	



1937-0652(2020)14:4;1-Y