

# *Algebra & Number Theory*

Volume 14

2020

No. 5

On the group of purely inseparable points of  
an abelian variety defined over  
a function field of positive characteristic, II

Damian Rössler



# On the group of purely inseparable points of an abelian variety defined over a function field of positive characteristic, II

Damian Rössler

Let  $A$  be an abelian variety over the function field  $K$  of a curve over a finite field. We describe several mild geometric conditions ensuring that the group  $A(K^{\text{perf}})$  is finitely generated and that the  $p$ -primary torsion subgroup of  $A(K^{\text{sep}})$  is finite. This gives partial answers to questions of Scanlon, Ghioca and Moosa, and Poonen and Voloch. We also describe a simple theory (used to prove our results) relating the Harder–Narasimhan filtration of vector bundles to the structure of finite flat group schemes of height one over projective curves over perfect fields. Finally, we use our results to give a complete proof of a conjecture of Esnault and Langer on Verschiebung divisibility of points in abelian varieties over function fields.

1. Introduction	1123
2. Intermediate results	1132
3. Semistable sheaves on curves	1133
4. Finite flat group schemes over curves	1135
5. Proofs of the claims made in Section 2A	1144
6. Proofs of the claims made in Section 2B	1146
7. Proof of Theorem 1.1	1149
8. Proof of Theorem 1.2	1150
9. Proof of Theorem 1.4	1152
Appendix A. Rational points in families	1164
Appendix B. Ampleness of the Hodge bundle and inseparable points	1166
Appendix C. Specialisation of the Mordell–Weil group	1168
Acknowledgments	1170
References	1171

## 1. Introduction

Let  $k$  be a finite field characteristic  $p > 0$  and let  $S$  be a smooth, projective and geometrically connected curve over  $k$ . Let  $K := \kappa(S)$  be its function field. Let  $A$  be an abelian variety of dimension  $g$  over  $K$ . Choose an algebraic closure  $\bar{K}$  of  $K$ . Let  $K^{\text{perf}} \subseteq \bar{K}$  be the maximal purely inseparable extension of  $K$ , let  $K^{\text{sep}} \subseteq \bar{K}$  be the maximal separable extension of  $K$  and let  $K^{\text{unr}} \subseteq K^{\text{sep}}$  be the maximal separable

*MSC2010:* primary 11J95; secondary 11G10, 14G25.

*Keywords:* abelian varieties, rational points, purely inseparable extensions, Frobenius, Verschiebung.

extension of  $K$ , which is unramified above every place of  $K$ . Finally, we let  $\mathcal{A}$  be a smooth commutative group scheme over  $S$  such that  $\mathcal{A}_K = A$ . We shall write  $\omega_{\mathcal{A}} := \epsilon_{\mathcal{A}/S}^*(\Omega_{\mathcal{A}/S})$  for the restriction of the cotangent sheaf of  $\mathcal{A}$  over  $S$  via the zero section  $\epsilon_{\mathcal{A}/S} : S \rightarrow \mathcal{A}$  of  $\mathcal{A}$ . We shall say that  $\omega_{\mathcal{A}}$  is the Hodge bundle of  $\mathcal{A}$ .

If  $G$  is an abelian group, we shall write

$$\mathrm{Tor}_p(G) := \{x \in G \mid \exists n \geq 0 : p^n \cdot x = 0\} \quad \text{and} \quad \mathrm{Tor}^p(G) := \{x \in G \mid \exists n \geq 0 : n \cdot x = 0 \wedge (n, p) = 1\}.$$

The aim of this text is to prove the following two theorems and to give a proof of a conjecture of Esnault and Langer (see further below).

**Theorem 1.1.** (a) *Suppose that  $A$  is geometrically simple. If  $A(K^{\mathrm{perf}})$  is finitely generated and of rank  $> 0$ , then  $\mathrm{Tor}_p(A(K^{\mathrm{sep}}))$  is a finite group.*

(b) *Suppose that  $A$  is an ordinary (not necessarily simple) abelian variety. If  $\mathrm{Tor}_p(A(K^{\mathrm{sep}}))$  is a finite group, then  $A(K^{\mathrm{perf}})$  is finitely generated.*

**Theorem 1.2.** *Suppose that  $\mathcal{A}$  is a semiabelian scheme and that  $A$  is a geometrically simple abelian variety over  $K$ . If  $\mathrm{Tor}_p(A(K^{\mathrm{sep}}))$  is infinite, then*

- (a)  $\mathcal{A}$  is an abelian scheme;
- (b) there is  $r_A \geq 0$  such that  $p^{r_A} \cdot \mathrm{Tor}_p(A(K^{\mathrm{sep}})) \subseteq \mathrm{Tor}_p(A(K^{\mathrm{unr}}))$ .

Furthermore, there is

- (c) an abelian scheme  $\mathcal{B}$  over  $S$ ;
- (d) an  $S$ -isogeny  $\mathcal{A} \rightarrow \mathcal{B}$ , whose degree is a power of  $p$  and such that the corresponding isogeny  $\mathcal{A}_K \rightarrow \mathcal{B}_K$  is étale;
- (e) an étale  $S$ -isogeny  $\mathcal{B} \rightarrow \mathcal{B}$  whose degree is  $> 1$  and is a power of  $p$ ,

and

- (f) (Voloch) if  $A$  is ordinary then the Kodaira–Spencer rank of  $A$  is not maximal;
- (g) if  $\dim(A) \leq 2$  then  $\mathrm{Tr}_{\bar{K}|\bar{k}}(A_{\bar{K}}) \neq 0$ ;
- (h) for all closed points  $s \in S$ , the  $p$ -rank of  $\mathcal{A}_s$  is  $> 0$ .

Here  $\mathrm{Tr}_{\bar{K}|\bar{k}}(A_{\bar{K}})$  is the  $\bar{K}|\bar{k}$ -trace of  $A_{\bar{K}}$ . This is an abelian variety over  $\bar{k}$ . See [Section 9A](#).

Theorems [1.1](#) and [1.2](#) (b) have applications in the context of the work of Poonen and Voloch on the Brauer–Manin obstruction over function fields. In particular Theorems [1.1](#) and [1.2](#) (b) show that the conclusion of [[Poonen and Voloch 2010](#), Theorem B] holds whenever the underlying abelian variety is geometrically simple, has semistable reduction and violates any of the conditions in [Theorem 1.2](#), in particular if it has a point of bad reduction. Theorems [1.1](#) and [1.2](#) (b) also feed into the “full” Mordell–Lang conjecture. See [[Scanlon 2005](#), after Claim 4.4; [Abramovich and Voloch 1992](#), Introduction] for this conjecture. In particular, in conjunction with the main result of [[Ghioca and Moosa 2006](#)], Theorems [1.1](#)

and 1.2 (b) show that the “full” Mordell–Lang conjecture holds if the underlying abelian variety is ordinary, geometrically simple, has semistable reduction and violates any of the conditions in [Theorem 1.2](#), in particular if it has a point of bad reduction.

Let now  $L$  be a field, which is finitely generated as a field over an algebraically closed field  $l_0$  of characteristic  $p$ . Let  $C$  be an abelian variety over  $L$ .

**Conjecture 1.3** (Esnault-Langer). *Suppose that for all  $\ell \geq 0$  we are given a point  $x_\ell \in C^{(p^\ell)}(L)$  and suppose that for all  $\ell \geq 1$ , we have  $V_{C^{(p^\ell)}/L}(x_\ell) = x_{\ell-1}$ . Then the image of  $x_0$  in  $C(L)/\mathrm{Tr}_{L|l_0}(C)(l_0)$  is a torsion point, which is of order prime to  $p$ .*

See [\[Esnault and Langer 2013, Remark 6.3 and after Lemma 6.5\]](#). This conjecture is important in the theory of stratified bundles in positive characteristic; see [\[Esnault and Langer 2013, Question 3 in the introduction\]](#) for details.

Here  $C^{(p^\ell)}$  is the base change of  $C$  by the  $\ell$ -th power of the absolute Frobenius morphism on  $\mathrm{Spec} L$  and  $V_{C^{(p^\ell)}/L} : C^{(p^\ell)} \rightarrow C^{(p^{\ell-1})}$  is the Verschiebung morphism. The abelian variety  $\mathrm{Tr}_{L|l_0}(C)$  is the  $L|l_0$ -trace of  $C$  (see [Section 9A](#)). It is an abelian variety over  $l_0$  and the variety  $\mathrm{Tr}_{L|l_0}(C)_L$  comes with an injective morphism to  $C$ . This gives in particular an injective map  $\mathrm{Tr}_{L|l_0}(C)(l_0) \rightarrow C(L)$ . The Lang–Néron theorem (see [\[Lang 1983, Chapter 6, Theorem 2\]](#)) asserts that  $C(L)/\mathrm{Tr}_{L|l_0}(C)(l_0)$  is a finitely generated group. Thus  $\mathrm{Tr}_{L|l_0}(C)(l_0) \subseteq C(L)$  is precisely the subgroup of  $C(L)$  consisting of divisible elements (i.e., elements divisible by any integer).

In the present text, we shall call a point  $x_0 \in C(L)$  with the property described in [Conjecture 1.3](#) an *indefinitely Verschiebung divisible point*. We shall write  $\mathrm{IVD}(C) = \mathrm{IVD}(C, L) \subseteq C(L)$  for the subgroup of indefinitely Verschiebung divisible points.

We prove:

**Theorem 1.4.** *Conjecture 1.3 holds.*

Note that [Theorem 1.4](#) has the following consequence, which is of independent interest: if  $C$  is as in [Conjecture 1.3](#),  $C$  is ordinary and  $\mathrm{Tr}_{L^{\mathrm{perf}}|l_0}(C_{L^{\mathrm{perf}}}) = 0$  then

$$\bigcap_{j \geq 0} p^j \cdot C(L^{\mathrm{perf}}) = \mathrm{Tor}^p(C(L^{\mathrm{perf}})).$$

To see this, let  $x \in C(L^{\mathrm{perf}})$ . Let  $L_1|L$  be a finite purely inseparable extension, which is a field of definition for  $x$ . Remember that the multiplication by  $p$  endomorphism of  $C$  is the composition of the Verschiebung morphism with the relative Frobenius morphism, which is purely inseparable. Also, recall that since  $C$  is ordinary, the Verschiebung morphism is (by definition) separable. Note finally that since  $\mathrm{Tr}_{L^{\mathrm{perf}}|l_0}(C_{L^{\mathrm{perf}}}) = 0$  we also have  $\mathrm{Tr}_{L_1|l_0}(C_{L_1}) = 0$ . In particular, if  $x \in \bigcap_{j \geq 0} p^j \cdot C(L^{\mathrm{perf}})$  then  $x$  is an indefinitely Verschiebung divisible element of  $C(L_1)$  and thus must lie in  $\mathrm{Tor}^p(C(L_1)) \subseteq \mathrm{Tor}^p(C(L^{\mathrm{perf}}))$  according to [Theorem 1.4](#). The inclusion  $\mathrm{Tor}^p(C(L^{\mathrm{perf}})) \subseteq \bigcap_{j \geq 0} p^j \cdot C(L^{\mathrm{perf}})$  is straightforward.

**Outline of the paper.** The basic strategy of the paper hinges on [Lemma 4.8](#) below. This Lemma associates a maximal multiplicative subgroup scheme with any finite flat group scheme of height one over  $S$ . The existence of this subgroup scheme is not straightforward and follows from an analysis of the Harder–Narasimhan filtration of (a Frobenius twist of) the coLie algebra of the group scheme. This analysis is carried out in [Section 4B](#).

One can apply [Lemma 4.8](#) to the kernel of the relative Frobenius morphism  $F_{\mathcal{A}/S} : \mathcal{A} \rightarrow \mathcal{A}^{(p)}$ , replace  $\mathcal{A}$  by the resulting quotient and repeat this construction ad infinitum, stopping only when the maximal multiplicative subgroup scheme is trivial.

It is then a basic (unresolved) question to determine minimal geometric conditions on  $\mathcal{A}$  ensuring that the resulting sequence of semiabelian schemes stops. This also makes sense (and seems important to us) if  $k$  is replaced by any perfect field of characteristic  $p > 0$  (not only when  $k$  is finite).

This question turns out to be intimately related to [Theorems 1.1, 1.2 and 1.4](#). To explain why, we shall first quote a result, which improves on (and elucidates) [Lemma B.2](#) in the Appendix. This result is proven in [\[Rössler 2019a\]](#), which builds on the present article. We shall only need [Lemma B.2](#) in the present text but for conceptual clarity, we shall present the improved result in this outline. Let  $E \subseteq S$  be the finite set of points  $s \in S$  where  $\mathcal{A}_s$  is not an abelian variety. Let  $U := S \setminus E$ . We first recall a classical result:

**Theorem 1.5** (Artin–Milne). *There is a canonical injective group homomorphism*

$$A^{(p)}(K)/F_{A/K}(A(K)) \hookrightarrow \mathrm{Hom}_K(F_K^*(\omega_K), \Omega_{K/k}).$$

Here  $F_K$  is the absolute Frobenius endomorphism of  $K$  (the  $p$ -th power map). See [\[Artin and Milne 1976, III.3.5.6\]](#) for the proof, which works in a more general setting. In [\[Rössler 2019a\]](#) this is refined as follows:

**Theorem 1.6** (R.). *The image of the Artin–Milne map lies inside the subgroup  $\mathrm{Hom}_C(F_S^*(\omega), \Omega_{S/k}(E))$  of  $\mathrm{Hom}_K(F_K^*(\omega_K), \Omega_{K/k})$ .*

Here  $F_S$  is the absolute Frobenius endomorphism of  $S$ . Here we write  $\Omega_{S/k}(E) := \Omega_{S/k}(E) \otimes \mathcal{O}_S(E)$  and  $E$  is understood as a divisor with no multiplicities. [Theorem 1.6](#) refines [Lemma B.2](#) below (for the knowledgeable reader, in [\[Rössler 2019a\]](#) it is even proven that the image of the Selmer group of the relative Frobenius morphism lies in  $\mathrm{Hom}_C(F_S^*(\omega), \Omega_{S/k}(E))$ ). The group  $\mathrm{Hom}_C(F_S^*(\omega), \Omega_{S/k}(E))$  can be understood as the target of an Abel–Jacobi map in logarithmic Higgs cohomology, although to give a precise meaning to this interpretation would require the development of a good theory of Higgs bundles in positive characteristic (which does not exist at the moment, to the author’s knowledge). This theorem is proven by providing a geometric interpretation for the Artin–Milne map and analysing its poles, making essential use of Faltings and Chai’s semistable compactification of the universal abelian scheme. The existence of this compactification allows us to show that the poles are at most logarithmic, which is in essence the content of [Theorem 1.6](#). Let us now explain why [Theorem 1.6](#) is relevant for [Theorem 1.1](#).

Consider, e.g., (b) in [Theorem 1.1](#). Suppose that  $A(K^{\text{perf}})$  is not finitely generated. We have

$$A(K^{\text{perf}}) = \bigcup_{i \geq 0} A(K^{p^{-i}})$$

and by the Lang–Néron theorem (see also [Section 9A](#))  $A(K^{p^{-i}})$  is finitely generated. Hence for infinitely many  $i \geq 0$ , we must have

$$A^{(p^{i+1})}(K)/F_{A^{(p^i)}/K}(A(K)) \simeq A(K^{p^{-i-1}})/A(K^{p^{-i}}) \neq 0.$$

In particular, for infinitely many  $i \geq 0$ , we must have

$$\text{Hom}_C(F_S^{\circ(i+1),*}(\omega), \Omega_{S/k}(E)) \neq 0$$

according to [Theorem 1.6](#). If now the vector bundle  $\omega$  were ample, this would lead to a contradiction, because if  $i$  is large enough and  $\omega$  is ample then there cannot be any morphism from  $F_S^{\circ(i+1),*}(\omega)$  to  $\Omega_{S/k}(E)$ . This was already noticed in the earlier article [[Rössler 2015](#)], where details are given. One can refine this line of reasoning as follows. If  $\omega$  is not ample and  $A$  is ordinary then one can show that  $\omega$  must have a certain nontrivial quotient, which is semistable of degree 0. This nontrivial quotient turns out to be induced by the maximal multiplicative subgroup scheme mentioned above. Calling it  $G_{\mathcal{A}}$ , we may then replace  $\mathcal{A}$  by  $\mathcal{A}/G_{\mathcal{A}}$ . The group  $(\mathcal{A}/G_{\mathcal{A}})_K(K^{\text{perf}})$  will again be infinitely generated, since the morphism  $A \rightarrow (\mathcal{A}/G_{\mathcal{A}})_K$  has finite kernel. Hence we can repeat the above reasoning for  $\mathcal{A}/G_{\mathcal{A}}$  and we obtain an infinite sequence of isogenous abelian varieties. The next step in the proof of [Theorem 1.1](#) (b) is to show that in this sequence, there are finitely many isomorphism classes. This follows from the fact that the degrees of  $\omega_{\mathcal{A}}$  and  $\mathcal{A}/G_{\mathcal{A}}$  are the same and more generally the degrees of the Hodge bundles of all the semiabelian schemes in the sequence are the same. This is a consequence of a computation involving the cotangent complex of the quotient morphism (see [Lemma 4.12](#)). It then follows from a classical reasoning involving moduli spaces of abelian varieties, familiar from Zarhin’s proof of the Tate conjecture over function fields, that the sequence contains only finitely many isomorphism classes. We can thus conclude that, up to isogeny,  $A$  contains a nontrivial finite endomorphism, whose kernel is multiplicative. The dual of this endomorphism is then separable and this shows that  $\text{Tor}_p(A^\vee)(K^{\text{sep}})$  is infinite (consider the kernels of its powers). Since  $A^\vee$  is isogenous to  $A$ , we see that  $\text{Tor}_p(A)(K^{\text{sep}})$  is also infinite. This concludes our outline of the proof of [Theorem 1.1](#) (b).

For [Theorem 1.1](#) (a), we consider the quotients of  $A$  by finite subgroups of  $\text{Tor}_p(A)(K^{\text{sep}})$  of increasing size. These quotients also run through finitely many isomorphism classes by a similar reasoning and we thus see that if  $\text{Tor}_p(A)(K^{\text{sep}})$  is infinite then, up to isogeny,  $A$  is endowed with a separable finite endomorphism. The dual of this endomorphism is then purely inseparable and of degree a positive power of  $p$ , and if  $A^\vee(K)$  is not finite, we may show that  $A^\vee(K^{\text{perf}})$  is infinitely generated by considering the inverse images of  $A(K)$  under the powers of this endomorphism. If now  $A^\vee(K^{\text{perf}})$  is not finitely generated, neither is  $A(K^{\text{perf}})$ , since  $A$  and  $A^\vee$  are isogenous. This concludes our outline of the proof of [Theorem 1.1](#) (a).

In [Theorem 1.2](#), we start out as in [Theorem 1.1](#) (a) and we again obtain, up to isogeny, a separable finite endomorphism of degree a positive power of  $p$ . The rest of the theorem investigates the geometric

consequences of the existence of this endomorphism. The most interesting consequence is the fact that it implies that  $\mathcal{A}$  must be an abelian scheme (if  $A$  is geometrically simple). This is (a) in [Theorem 1.2](#). The main point here is that the endomorphism extends to an étale endomorphism of  $\mathcal{A}$ . If  $\mathcal{A}$  had a fibre with a toric part then the endomorphism would induce an automorphism of the toric part, because tori only have infinitesimal  $p$ -primary subgroups in characteristic  $p$  and these are only étale if they are trivial. This fact forces the whole endomorphism to be an automorphism, which is impossible. The proof of (c), (d) and (e) are straightforward and not much more than a rewording of the fact that there are only finitely many isomorphism classes in the set of quotients described above. The proof of (b) follows essentially from a variant of the fact that, under (a), the above endomorphism extends to an everywhere étale and finite endomorphism of  $\mathcal{A}$ . This also easily gives a proof of (h). The proof of (g) is based on class field theory and the Serre–Tate theory of canonical liftings. First, up to a finite extension, the field extension generated by the points of  $\mathrm{Tor}_p(A)(K^{\mathrm{sep}})$  is everywhere unramified by (a) and (b). If  $\mathrm{Tor}_p(A)(K^{\mathrm{sep}}) = \mathrm{Tor}_p(A)(\bar{K})$  then a simple application of the Serre–Tate theory of canonical liftings shows that  $A_{\bar{K}}$  is the base change of an abelian variety defined over  $\bar{k}$ . Hence it must be contained in the Hilbert class field of  $K$ , which is but a constant field extension (i.e., comes from an extension of  $k$ ), up to a finite extension. So if  $\mathrm{Tor}_p(A)(K^{\mathrm{sep}})$  is infinite then it is an infinite torsion subset of  $A(K\bar{k})$ , which is finitely generated by the Lang–Néron theorem if the trace of  $A$  vanishes. This is a contradiction.

We now turn to [Theorem 1.4](#). Esnault and Langer [[2013](#), Theorem 6.2], using a height argument due to Raynaud, proved that the image of  $x_0$  in  $C(L)/\mathrm{Tr}_{L|l_0}(C)(l_0)$  is a torsion point under the assumption that  $C$  has everywhere potential good reduction in codimension one. Their argument works as follows. Choose a polarisation on  $C$ . This induces polarisations on all the  $C^{(p^\ell)}$  by base change. A simple computation shows that if a point  $x \in C(L)$  has a preimage  $y$  in  $C^{(p)}(K)$  under the Verschiebung map then the height of  $x$  with respect to the polarisation is  $p$  times the height of  $y$  with respect to the base changed polarisation. Now if  $C$  has everywhere good reduction in codimension one, there is an abelian scheme  $\mathcal{C}$  extending  $C$  on an open subset with complement of codimension  $\geq 2$  of a normal complete model  $V$  of  $L$  and the polarisations on  $C$  and  $C^{(p)}$  naturally extend to this open subset. This implies that the heights of  $x$  are  $y$  (with respect to the polarisations and a choice of ample line bundle on  $V$ ) are integers, because they can then be computed in a completely geometric fashion. In particular, the height of  $x$  is an integer divisible by  $p$ . Repeating this argument with  $y$ , one sees that the height of  $x$  is divisible by arbitrarily high powers of  $p$  and one concludes that it must vanish. Then the conclusion follows from a theorem of Lang (see [[Conrad 2006](#), Theorem 9.15]). The argument described above breaks down in the presence of bad reduction in codimension one because the orders of the component groups of the special fibres of the local Néron models of the varieties  $C^{(p^\ell)}$  increase with  $\ell$  if they are not trivial and this introduces denominators in the heights.

Our approach to [Theorem 1.4](#) is again via the infinite sequence of quotients described at the beginning of the outline. This sequence will effectively replace the sequence of the  $C^{(p^\ell)}$ . It has the advantage over the sequence of the  $C^{(p^\ell)}$  that it falls inside a bounded family of abelian varieties (see below), making it possible to control the order of the (analogues of the) images of the  $x_\ell$  in the component groups of the Néron models. This makes a similar height computation possible. The proof is in several steps.

Step (0). Reduction to the case where  $L$  is the function field of a smooth and projective curve  $B$  over  $l_0$ . This follows from a Bertini type argument—see [Appendix C](#).

Step (1). We consider the images of the  $x_\ell$  under the Artin–Milne map. A crucial point is that these images must be compatible under the Verschiebung morphisms (see diagram (8) below) and this constrains the image of  $x_1$  under the Artin–Milne map. Using [Lemma B.2](#) (or [Theorem 1.6](#)), the theory of semistable sheaves in positive characteristic and various global results on finite flat group schemes of height one in a global situation proven in [Section 4](#), we show that the image of  $x_1$  under the Artin–Milne map must factor through the coLie algebra of the maximal multiplicative subgroup  $(\ker F_{C/B})_\mu$  of  $\ker F_{C/B}$ . This implies that the image of  $x_1$  in  $(C^{(p)}/(\ker F_{C/B})_{\mu,L}^{(p)})(L) = (C^{(p)}/G_{C^{(p)},L})(L)$  maps to 0 under the Artin–Milne map. From the definitions, this means that the image of  $x_0$  in  $(C/G_{C,L})(L)$  is divisible by  $p$  in  $(C/G_{C,L})(L)$ . Suppose for simplicity that  $C$  has a semiabelian model  $\mathcal{C}$  over  $B$ . We can now repeat this process and we obtain a sequence of purely inseparable morphisms  $\psi_i : C \rightarrow C_i$  of increasing degree, such that  $\psi_{i,L}(x_0)$  in  $C_i$  is divisible by  $p^i$  in  $C_i(L)$ .

Step (2). We choose a polarisation  $\phi_{D_0} : C \rightarrow C^\vee$ . The image of  $x_0$  under  $\phi_{D_0}$  is of course also indefinitely Verschiebung divisible. We identify  $\phi_{D_0}(x_0)$  with a line bundle  $M$  on  $C$ . Since  $\phi_{D_0}(x_0)$  is indefinitely Verschiebung divisible, there are line bundles  $M_i$  on  $C^{(p^i)}$  such that  $M$  is the pull-back of  $M_i$  by the morphism  $C \rightarrow C^{(p^i)}$  arising by composing relative Frobenii. The morphism  $C \rightarrow C^{(p^i)}$  factors through  $\psi_{i,L}$  by construction. Hence there are line bundles  $J_i$  on the  $C_i$  such that  $\psi_{i,L}^*(J_i) = M$ .

Step (3). We now compute the height pairing between  $x_0$  and  $M$ . This can easily be seen to equal the height pairing between  $\psi_{i,L}(x_0)$  and  $J_i$ . Since  $\psi_{i,L}(x_0)$  is divisible by  $p^i$ , we see that the height pairing between  $x_0$  and  $M$  is divisible by  $p^i$ . If the  $C_i$  were all abelian schemes we could deduce (like Raynaud–Esnault–Langer above) that the height pairing between  $x_0$  and  $M$  must vanish, because then all the values of the various height pairing would be integral. However, we cannot assume this.

Step (4). All the  $C_i$  are essentially part of a bounded family of abelian varieties over  $L$  because the degrees of the Hodge bundles of the  $C_i$  are all equal (see above in the outline). Using this, one can prove that there is an infinite set  $I_0 \subseteq \mathbb{N}$  such that if  $i \in I_0$  the image of any element of  $C_i(L)$  in the component groups of the Néron model of  $C_i$  has an order, which is bounded independently of  $i$ . This follows from [Proposition A.2](#) (a) in the Appendix. The gist of the argument is that in a bounded family of semiabelian varieties over  $B$ , it is possible to smoothly compactify the generic fibre, up to normalisation in a finite extension of the function field of the parameter space. This would follow from resolution of singularities but in the present situation is a consequence of the work of Mumford, Chai–Faltings and Künnemann (see [\[Künnemann 1998, Theorem 4.2\]](#)). This means that the abelian varieties in the family almost all have regular compactifications with a bounded number of geometric fibres over  $B$ . This bound is also a bound for the order of the image of a rational point in the component groups of the Néron model.

Step (5). In view of Step (4), if we replace  $x_0$  by a certain multiple of  $x_0$ , all the height pairing in sight are integers. Hence the divisibility argument envisaged in Step (3) can be carried out and yields that the

height pairing of  $x_0$  and  $M$  vanishes. This pairing is by construction twice the Néron–Tate height of  $x_0$  with respect to the polarisation  $\phi_{D_0}$  and we conclude from a theorem of Lang [1983] that the image of  $x_0$  in  $C(L)/\mathrm{Tr}_{L|l_0}(C)(l_0)$  is a torsion point. It remains to show that its order is prime to  $p$ .

Step (6). We first show that we may suppose that  $\mathrm{Tr}_{\bar{L}|l_0}(C_{\bar{L}}) = 0$ . This is not completely straightforward, because when one passes to a finite extension in [Conjecture 1.3](#), one loses control of part of the torsion of  $C(L)/\mathrm{Tr}_{L|l_0}(C)(l_0)$ . However, although the parasitical torsion subgroup that might appear is not known, its exponent only depends on the degree of the extension. This degree can be taken to be the same for all the Frobenius twists of  $C$  and the information one gathers from this suffices to prove the conjecture, provided one can prove it for a finite extension. Thus we may suppose that  $\dim(\mathrm{Tr}_{\bar{L}|l_0}(C_{\bar{L}})) = \dim(\mathrm{Tr}_{L|l_0}(C))$  and then, after quotienting by  $\mathrm{Tr}_{L|l_0}(C)$ , that  $\mathrm{Tr}_{\bar{L}|l_0}(C_{\bar{L}}) = 0$ . Now recall that the  $C_i$  are essentially part of a bounded family of abelian varieties over  $L$  (see step (4)). Using this, and the fact that now  $\mathrm{Tr}_{\bar{L}|l_0}(C_{i,\bar{L}}) = 0$  for all  $i \geq 1$ , one can prove that there is an infinite set  $I_0 \subseteq \mathbb{N}$  such that if  $i \in I_0$ , the cardinality of the torsion subgroup of  $C_i(L)$  is uniformly bounded. This follows from [Proposition A.2](#) (b) in the Appendix. To finish the proof of [Theorem 1.4](#), suppose that  $x_0$  is a nonzero torsion point, which is indefinitely Verschiebung divisible. Since the image of  $x_0$  in  $C_i(L)$  is divisible by  $p^i$ , we see that the torsion group of  $C_i(L)$  has an element of order  $p^{i+1}$ . This contradicts the above uniformity statement and shows that the order of  $x_0$  must be prime to  $p$ .

The argument to prove the uniformity statement alluded to in Step (6) goes roughly as follows. One first notices that the torsion subgroup of a trace free abelian variety coincides with the set of elements of vanishing Néron–Tate height by the already quoted theorem of Lang. Thus they can be described as the points of a moduli space of sections, which is of finite type over  $l_0$ , at least for those torsion points, whose image in the component groups of the Néron model of the abelian variety is trivial. Since the abelian variety is trace free, the torsion subgroup is finite and thus this moduli space is finite. Using the uniformity statement in Step (4), we may assume that the torsion points of the  $C_i(L)$  have trivial images in the components of the corresponding Néron models, up to multiplication by a fixed integer (independent of  $i$  running through an infinite set). The number of irreducible components of the moduli space of each  $C_i$  is now uniformly bounded, since the  $C_i$  are part of a bounded family. This gives a uniform bound for the torsion subgroups of the  $C_i(L)$ .

The reader may enjoy the talk [\[Rössler 2019b\]](#) as an introduction to parts of the present article.

The structure of the article is as follows. In [Section 2](#), we state various intermediate results, from which we shall deduce [Theorems 1.1](#) and [1.2](#). [Theorem 2.1](#) in [Section 2A](#) is of independent interest and is (we feel) likely to be useful for the study of the geometry of (especially ordinary) abelian varieties in general. The results in [Section 2A](#) are deduced from some results in the theory of finite flat groups schemes of height one over  $S$ , most of which follow from the existence of a Harder–Narasimhan filtration on their Lie algebras. These results on finite flat group schemes are proven in [Section 4](#) and for the convenience of the reader, we included a section ([Section 3](#)) listing the results on semistable sheaves over curves in positive characteristic that we need. To the knowledge of the author, there are very few general results

on the structure of finite flat group schemes in a global situation (e.g., when the base is not affine) and it seems that it is the first time that the theory of semistability of vector bundles is being used in this context. In [Bost 2004] a similar idea is used in characteristic 0, where it is applied to the study of formal groups over curves (recall that all groups schemes are smooth in characteristic 0, so the Lie algebras of finite flat group schemes vanish in characteristic 0). Lemma 4.4 below (which concerns finite flat group schemes of height one) is inspired by [Bost 2004, Lemma 2.9]. A prototype of Lemma 4.4 can be found in [Shepherd-Barron 1992, Lemma 9.1.3.1] but it is not applied to the study of group schemes there. The key results here are the Lemmata 4.4 and 4.8, which will hopefully lead to further generalisations (e.g., in the situation when the base scheme is of dimension higher than one - in this direction, see [Langer 2015, Theorem 7.3]). The results in Section 2B do not require the theory of semistable sheaves and are based on geometric class field theory, the theory of Serre–Tate canonical liftings and on the existence of moduli schemes for abelian varieties. In Section 5, we prove the various claims made in Section 2A and in Section 6 we prove the claims made in Section 2B. In Section 7, we prove Theorem 1.1 and in Section 8 we prove Theorem 1.2. In Section 9B, we give a proof of Theorem 1.4. The proof of Theorem 1.4 is quite long and uses virtually all the other results proven in this text.

In his very interesting recent preprint, Xinyi Yuan [2018] uses some techniques which are also used in the present paper. They were discovered independently. His text focusses on the case where the base curve is the projective line. In particular, the “quotient process” used in step (2) of the proof of Theorem 1.4 and also in the proof of Theorem 1.2 also appears (over the projective line) in section 2.2 of [Yuan 2018]. Theorem 2.9 of [Yuan 2018] overlaps with the proof of Lemma 4.11.

The prerequisites for this article are algebraic geometry at the level of the EGA, familiarity with the basic theory of finite flat group schemes, as expounded in [Tate 1997] and a good knowledge of the theory of abelian schemes and varieties, as presented in [Milne 1986; Mumford 1970; Moret-Bailly 1985]. We also expect the reader to be familiar with the basic properties of Néron models (as in the chapter on basics of [Bosch et al. 1990]) and to have a working knowledge of Grothendieck topologies.

**Notation.** If  $X$  is an integral scheme, we write  $\kappa(X)$  for the local ring at the generic point of  $X$  (which is a field). If  $X$  is a scheme of characteristic  $p$ , we denote the absolute Frobenius endomorphism of  $X$  by  $F_X$ . If  $f : X \rightarrow Y$  is a morphism between two schemes of characteristic  $p$  and  $\ell > 0$ , abusing language, we denote by  $X^{(p^\ell)}$  the fibre product of  $f$  and  $F_Y^{\circ\ell}$ , where  $F_Y^{\circ\ell}$  is the  $\ell$ -th power of the Frobenius endomorphism  $F_Y$  of  $Y$ . If  $G \rightarrow X$  is a group scheme, we write  $\epsilon_{G/X} : X \rightarrow G$  for the zero section of  $G$  and

$$\omega_{G/X} = \omega_G := \epsilon_{G/X}^*(\Omega_{G/X}).$$

If  $X$  is of characteristic  $p$ , we shall write  $F_{G/X} : G \rightarrow G^{(p)}$  for the relative Frobenius morphism. If in addition  $G$  is flat and commutative, we shall write  $V_{G^{(p)}/X} : G^{(p)} \rightarrow G$  for the corresponding Verschiebung morphism; we shall write

$$F_{G/X}^{(n)} : G \rightarrow G^{(p^n)} \quad \text{and} \quad V_{G^{(p^n)}/X}^{(j)} : G^{(p^n)} \rightarrow G^{(p^{n-j})}$$

for the compositions of morphisms

$$F_{G^{(p^n-1)}/X} \circ \cdots \circ F_{G/X} \quad \text{and} \quad V_{G^{(p^n-j+1)}/X} \circ V_{G^{(p^n-j+2)}/X} \circ \cdots \circ V_{G^{(p^n)}/X},$$

respectively. See [SGA 3, 2011, Exposé VII<sub>A</sub>, §4, “Frobeniusseries”] for the definition of the relative Frobenius morphism and the Verschiebung. If  $G$  is finite flat and commutative, we shall write  $G^\vee$  for the Cartier dual of  $G$ .

## 2. Intermediate results

We keep the notations and terminology of the introduction.

**2A. Consequences of infinite generation of  $A(K^{\text{perf}})$ .** We shall write

$$\overline{\text{rk}}_{\min}(\omega_{\mathcal{A}}) := \lim_{\ell \rightarrow \infty} \text{rk}((F_S^{\circ\ell,*}(\omega_{\mathcal{A}}))_{\min}) \quad \text{and} \quad \overline{\mu}_{\min}(\omega_{\mathcal{A}}) := \lim_{\ell \rightarrow \infty} \frac{\text{deg}((F_S^{\circ\ell,*}(\omega_{\mathcal{A}}))_{\min})}{p^\ell \cdot \text{rk}((F_S^{\circ\ell,*}(\omega_{\mathcal{A}}))_{\min})}.$$

Here  $F_S^{\circ\ell}$  is the  $\ell$ -th power of the absolute Frobenius endomorphism of  $S$  and  $(F_S^{\circ\ell,*}(\omega_{\mathcal{A}}))_{\min}$  is the semistable quotient with minimal slope of the vector bundle  $F_S^{\circ\ell,*}(\omega_{\mathcal{A}})$ . See Section 3 for details. Our main tool will be the following theorem.

**Theorem 2.1.** *There exists a (necessarily unique) multiplicative subgroup scheme  $G_{\mathcal{A}} \hookrightarrow \ker F_{\mathcal{A}/S}$ , with the following property: if  $H$  is a finite, flat, multiplicative group scheme of height one over  $S$  and  $f : H \rightarrow \ker F_{\mathcal{A}/S}$  is a morphism of group schemes, then  $f$  factors through  $G_{\mathcal{A}}$ .*

*If  $\mathcal{A}$  is ordinary and  $\omega_{\mathcal{A}}$  is not ample then the order of  $G_{\mathcal{A}}$  is  $p^{\overline{\text{rk}}_{\min}(\omega_{\mathcal{A}})}$ .*

*If  $\phi : \mathcal{A} \rightarrow \mathcal{B}$  is a morphism of smooth commutative group schemes over  $S$ , then the restriction of  $\phi$  to  $G_{\mathcal{A}}$  factors through  $G_{\mathcal{B}}$ . Furthermore, we have  $\text{deg}(\omega_{\mathcal{A}}) = \text{deg}(\omega_{\mathcal{A}/G_{\mathcal{A}}})$ .*

Here  $\mathcal{A}/G_{\mathcal{A}}$  is the “fppf quotient” of  $\mathcal{A}$  by  $G_{\mathcal{A}}$ , which is also a smooth commutative group scheme over  $S$ . See Proposition 4.1 below for details.

**Remark 2.2.** Note that  $\overline{\mu}_{\min}(\omega_{\mathcal{A}}) > 0$  is equivalent to  $\omega_{\mathcal{A}}$  being ample (see [Barton 1971]).

**Remark 2.3.** Theorem 2.1 holds more generally if  $k$  is only supposed to be perfect (the proof does not use the fact that  $k$  is finite).

**Remark 2.4.** It would be interesting to provide an explicit example of an abelian variety  $A$  as in the introduction to this article, such that  $A$  is ordinary,  $\mathcal{A}$  is semiabelian,  $\text{Tr}_{\overline{k}|\overline{k}}(A_{\overline{k}}) = 0$  and  $G_{\mathcal{A}} \neq 0$ . It should be possible to construct such an example by considering mod  $p$  reductions of the abelian variety constructed in [Catanese and Dettweiler 2016, Theorem 1.3]. We hope to return to this question in a later article. The following question is also of interest: is there an ordinary abelian variety  $A$  as above, such that  $A$  has maximal Kodaira–Spencer rank,  $\mathcal{A}$  is semiabelian and  $G_{\mathcal{A}} \neq 0$ ?

**Proposition 2.5.** *Suppose that  $A$  is ordinary and that  $\mathcal{A}$  is semiabelian. Suppose that  $A(K^{\text{perf}})$  is not finitely generated. Then  $G_{\mathcal{A}}$  is of order  $> 1$  and  $\mathcal{A}/G_{\mathcal{A}}$  is also semiabelian.*

**Proposition 2.6.** *Suppose that  $A$  is ordinary and that  $\mathcal{A}$  is semiabelian over  $S$ . Suppose that  $A(K^{\text{perf}})$  is not finitely generated.*

*Then there is a finite flat morphism*

$$\phi : \mathcal{A} \rightarrow \mathcal{B},$$

*where  $\mathcal{B}$  is a semiabelian over  $S$  and a finite flat morphism*

$$\lambda : \mathcal{B} \rightarrow \mathcal{B}$$

*such that  $\ker(\phi)$  and  $\ker(\lambda)$  are multiplicative group schemes and such that the order of  $\ker(\lambda)$  is  $> 1$ .*

## 2B. Consequences of infiniteness of $\text{Tor}_p(A(K^{\text{sep}}))$ or $\text{Tor}_p(A(K^{\text{unr}}))$ .

**Theorem 2.7.** *Suppose that  $\text{Tr}_{\bar{K}|\bar{k}}(A_{\bar{K}}) = 0$ . Suppose that the action of  $\text{Gal}(K^{\text{sep}}|K)$  on  $\text{Tor}_p(A(K^{\text{unr}}))$  factors through  $\text{Gal}(K^{\text{sep}}|K)^{\text{ab}}$ . Then  $\text{Tor}_p(A(K^{\text{unr}}))$  is finite.*

Here  $\text{Gal}(K^{\text{sep}}|K)^{\text{ab}}$  is the maximal abelian quotient of  $\text{Gal}(K^{\text{sep}}|K)$ .

**Proposition 2.8.** *Suppose that  $\dim(A) \leq 2$  and that  $\text{Tr}_{\bar{K}|\bar{k}}(A_{\bar{K}}) = 0$ . Then  $\text{Tor}_p(A(K^{\text{unr}}))$  is finite.*

**Theorem 2.9.** *Suppose that  $\text{Tor}_p(A(K^{\text{sep}}))$  is infinite. Then there is an étale  $K$ -isogeny*

$$\phi : A \rightarrow B,$$

*where  $B$  is an abelian variety over  $K$  and there is an étale  $K$ -isogeny*

$$\lambda : B \rightarrow B$$

*such that the order of  $\ker(\lambda)$  is  $> 1$  and such that the orders of  $\ker(\lambda)$  and  $\ker(\phi)$  are powers of  $p$ .*

**Theorem 2.10.** *Suppose that there exists an étale  $K$ -isogeny  $\phi : A \rightarrow A$ , such that  $\deg(\phi) = p^r$  for some  $r > 0$ . Suppose also that  $A$  is a geometrically simple abelian variety and that  $\mathcal{A}$  is a semiabelian scheme.*

*Then  $\mathcal{A}$  is an abelian scheme and  $\phi$  extends to an étale (necessarily finite)  $S$ -morphism  $\mathcal{A} \rightarrow \mathcal{A}$  of group schemes.*

## 3. Semistable sheaves on curves

Let  $Y$  be a scheme, which is smooth, projective and geometrically connected of relative dimension one over a field  $t_0$ .

Suppose to begin with that  $t_0$  is algebraically closed.

If  $V$  is a nonzero coherent locally free sheaf on  $Y$ , we write (as is customary)

$$\mu(V) = \deg(V) / \text{rk}(V),$$

where

$$\deg(V) := \int_Y c_1(V)$$

and  $\text{rk}(V)$  is the rank of  $V$ . The quantity  $\mu(V)$  is called the *slope* of  $V$ . Recall that a nonzero locally free coherent sheaf  $V$  on  $Y$  is called semistable if for any nonzero coherent subsheaf  $W \subseteq V$ , we have

$\mu(W) \leq \mu(V)$ . Let  $V/Y$  be a nonzero locally free coherent sheaf on  $Y$ . There is a unique filtration by coherent subsheaves

$$0 = V_0 \subsetneq V_1 \subsetneq V_2 \subsetneq \cdots \subsetneq V_{\text{hn}(V)} = V$$

such that all the sheaves  $V_i/V_{i-1}$  ( $1 \leq i \leq \text{hn}(V)$ ) are (locally free and) semistable and such that the sequence  $\mu(V_i/V_{i-1})$  is strictly decreasing. This filtration is called the *Harder–Narasimhan filtration* of  $V$  (shorthand: HN filtration). One then defines

$$V_{\min} := V/V_{\text{hn}(V)-1}, \quad V_{\max}(V) := V_1 \quad \text{and} \quad \mu_{\max}(V) := \mu(V_1), \quad \mu_{\min}(V) := \mu(V_{\min}).$$

Let now  $r \in \mathbb{Q}$ . Suppose that  $r \in \{\mu(V_1), \dots, \mu(V/V_{\text{hn}(V)-1})\}$ . Let  $i(r) \in \mathbb{N}$  be the unique natural number such that  $\mu(V_{i(r)}/V_{i(r)-1}) = r$ . We shall write

$$V_{=r} := V_{i(r)}/V_{i(r)-1} \quad \text{and} \quad V_{\geq r} := V_{i(r)}.$$

We shall also write

$$V_{>r} := V_{j(r)},$$

where  $j(r) \in \mathbb{N}$  is the largest natural number such that  $\mu(V_{j(r)}/V_{j(r)-1}) > r$ .

One basic property of semistable sheaves that we shall use repeatedly is the following. If  $V$  and  $W$  are nonzero coherent locally free sheaves on  $Y$  and  $\mu_{\min}(V) > \mu_{\max}(W)$  then  $\text{Hom}_Y(V, W) = 0$ . This follows from the definitions.

See [Brenner et al. 2008, Chapter 5] (for instance) for all these notions.

If  $V$  is a nonzero coherent locally free sheaf on  $Y$  and  $t_0$  has positive characteristic, we say that  $V$  is *Frobenius semistable* if  $F_Y^{\text{or},*}(V)$  is semistable for all  $r \in \mathbb{N}$ . The terminology *strongly semistable* also appears in the literature.

**Theorem 3.1.** *Let  $V$  be a nonzero coherent locally free sheaf on  $Y$ . There is an  $\ell_0 = \ell_0(V) \in \mathbb{N}$  such that the quotients of the Harder–Narasimhan filtration of  $F_Y^{\text{ol},*}(V)$  are all Frobenius semistable.*

*Proof.* See, e.g., [Langer 2004, Theorem 2.7, p. 259]. □

Theorem 3.1 shows in particular that the definitions

$$\begin{aligned} \bar{\mu}_{\min}(V) &:= \lim_{\ell \rightarrow \infty} \mu_{\min}(F_Y^{\text{ol},*}(V))/p^\ell, & \bar{\mu}_{\max}(V) &:= \lim_{\ell \rightarrow \infty} \mu_{\max}(F_Y^{\text{ol},*}(V))/p^\ell, \\ \bar{\text{rk}}_{\min}(V) &:= \lim_{\ell \rightarrow \infty} \text{rk}((F_Y^{\text{ol},*}(V))_{\min}), & \bar{\text{rk}}_{\max}(V) &:= \lim_{\ell \rightarrow \infty} \text{rk}((F_Y^{\text{ol},*}(V))_{\max}), \end{aligned}$$

make sense if  $V$  is a nonzero locally free and coherent sheaf on  $Y$ .

Suppose now that  $t_0$  is only perfect (not necessarily algebraically closed). If  $V$  is a nonzero coherent sheaf on  $Y$ , then we shall write  $\mu(V) := \mu(V_{t_0})$  and we shall say that  $V$  is semistable if  $V_{t_0}$  is semistable. The HN filtration of  $V_{t_0}$  is invariant under  $\text{Gal}(\bar{t}_0|t_0)$  by unicity and by a simple descent argument, we see that there is a unique filtration by coherent subsheaves

$$V_0 \subsetneq V_1 \subsetneq V_2 \subsetneq \cdots \subsetneq V_{\text{hn}(V)}$$

such that

$$V_{0, \bar{t}_0} \subsetneq V_{1, \bar{t}_0} \subsetneq V_{2, \bar{t}_0} \subsetneq \cdots \subsetneq V_{\text{hn}(V), \bar{t}_0}$$

is the HN filtration of  $V_{\bar{t}_0}$ . We then define as before

$$\mu_{\max}(V) := \mu(V_1) \quad \text{and} \quad \mu_{\min}(V) := \mu(V/V_{\text{hn}(V)-1}).$$

Notice that we have  $\mu_{\max}(V) = \mu_{\max}(V_{\bar{t}_0})$  and  $\mu_{\min}(V) = \mu_{\min}(V_{\bar{t}_0})$ .

Notice that if  $V$  and  $W$  are nonzero coherent locally free coherent sheaves on  $Y$  and  $\mu_{\min}(V) > \mu_{\max}(W)$  then we still have  $\text{Hom}_Y(V, W) = 0$ , since there is a natural inclusion

$$\text{Hom}_Y(V, W) \subseteq \text{Hom}_{Y_{\bar{t}_0}}(V_{\bar{t}_0}, W_{\bar{t}_0}).$$

If  $t_0$  has positive characteristic, we shall say that  $V$  is Frobenius semistable if  $V_{\bar{t}_0}$  is Frobenius semistable. Since Frobenius morphisms commute with all morphisms, this is equivalent to requiring that  $F_Y^{r,*}(V)$  is semistable for all  $r \in \mathbb{N}$  (with our extended definition of semistability).

We can now extend the range of the terminology introduced above:

$$\begin{aligned} V_{\min} &:= V/V_{\text{hn}(V)-1}, & V_{\max} &:= V_1, \\ \bar{\mu}_{\min}(V) &:= \lim_{\ell \rightarrow \infty} \mu_{\min}(F_Y^{\circ\ell,*}(V))/p^\ell, & \bar{\mu}_{\max}(V) &:= \lim_{\ell \rightarrow \infty} \mu_{\max}(F_Y^{\circ\ell,*}(V))/p^\ell, \\ \bar{\text{rk}}_{\min}(V) &:= \lim_{\ell \rightarrow \infty} \text{rk}((F_Y^{\circ\ell,*}(V))_{\min}), & \bar{\text{rk}}_{\max}(V) &:= \lim_{\ell \rightarrow \infty} \text{rk}((F_Y^{\circ\ell,*}(V))_{\max}). \end{aligned}$$

Note that we have  $\bar{\mu}_{\min}(V) = \bar{\mu}_{\min}(V_{\bar{t}_0})$ ,  $\bar{\mu}_{\max}(V) = \bar{\mu}_{\max}(V_{\bar{t}_0})$ ,  $\bar{\text{rk}}_{\min}(V) = \bar{\text{rk}}_{\min}(V_{\bar{t}_0})$ ,  $\bar{\text{rk}}_{\max}(V) = \bar{\text{rk}}_{\max}(V_{\bar{t}_0})$  as expected.

If  $V$  is a nonzero coherent locally free coherent sheaf on  $Y$  such that all the quotients of the HN filtration of  $V$  are Frobenius semistable, we shall say that  $V$  has a Frobenius semistable HN filtration. Note that by [Theorem 3.1](#) above, for any nonzero coherent locally free coherent sheaf  $V$  on  $Y$ , the sheaf  $F^{or,*}(V)$  has a Frobenius semistable HN filtration for all but finitely many  $r \in \mathbb{N}$ .

The following simple lemma will also prove very useful. It was suggested by J.-B. Bost.

**Lemma 3.2.** *Let  $V$  and  $W$  be coherent locally free sheaves on  $Y$ . Suppose that  $\mu(V) = \mu(W)$  and that  $\text{rk}(V) = \text{rk}(W)$ . Let  $\phi : V \rightarrow W$  be a monomorphism of  $\mathcal{O}_Y$ -modules. Then  $\phi$  is an isomorphism.*

*Proof.* We may suppose that  $V$  and  $W$  are of positive rank, otherwise the lemma is tautologically true. Let  $M := \det(W) \otimes \det(V)^\vee$ . The assumptions imply that  $\deg(M) = 0$ . Let  $\det(\phi) \in H^0(Y, M)$  be the section induced by  $\phi$ . The zero scheme  $Z(\det(\phi))$  of  $\det(\phi)$  is a torsion sheaf since  $\det(\phi)$  is nonzero at the generic point of  $Y$  and the length of  $Z(\det(\phi))$  is equal to the degree of  $M$  so  $Z(\det(\phi))$  must be empty. In other words,  $M$  is the trivial sheaf and  $\det(\phi)$  is a constant nonzero section of  $M$ . In particular,  $\phi$  is an isomorphism. □

#### 4. Finite flat group schemes over curves

The terminology of this section is independent of the introduction.

**4A. Quotients by proper flat group schemes.** Let  $Y$  be a noetherian scheme. Let  $G$  be a commutative strongly quasiprojective flat group scheme over  $Y$ . See [Bosch et al. 1990, 8.2, p. 211] for the definition of strong quasiprojectivity. Note that if  $Y$  is regular then  $G$  is strongly quasiprojective over  $Y$  if it is quasiprojective over  $Y$ .

Suppose that  $H$  is a closed subgroup scheme of  $G$ , which is proper and flat over  $Y$ . The  $Y$ -scheme  $G$  (resp.  $H$ ) defines a functor  $\underline{G}$  (resp.  $\underline{H}$ ) from the category of  $Y$ -schemes to the category of abelian groups. Both functors are fppf sheaves by a classical result of Grothendieck. We may thus form the quotient  $\underline{G}/\underline{H}$  of  $\underline{G}$  and  $\underline{H}$  in the category of fppf sheaves.

The following proposition describes the quotient construction that we use in this text.

**Proposition 4.1.** *The fppf sheaf  $\underline{G}/\underline{H}$  is representable by a group scheme  $G/H$  over  $Y$ , which is also strongly quasiprojective. The natural morphism  $q : G \rightarrow G/H$  is proper and faithfully flat and makes  $G$  into an  $H_{G/H}$ -torsor over  $G/H$ .*

*Proof.* See [Bosch et al. 1990, Theorem 8.12, p. 220]. □

Note that if  $G$  is semiabelian and  $Y$  is normal then  $G$  is quasiprojective over  $Y$  (combine [Moret-Bailly 1985, VI.3.1] with [Raynaud 1970, XI.1.4]). In particular if  $Y$  is regular and  $G$  is semiabelian then  $G$  is strongly quasiprojective over  $Y$ .

**4B. The HN-filtration on the Lie algebra of a finite flat group scheme of height one.** Let  $S$  be a smooth, projective and geometrically connected curve over a perfect field  $k$ . Suppose that  $\text{char}(k) = p > 0$ .

The following preliminary lemma will be very useful.

**Lemma 4.2.** *Let  $G$  be a finite flat commutative group scheme over  $S$ . Let  $T \rightarrow S$  be a flat, radicial and finite morphism and let  $\phi : H \hookrightarrow G_T$  be a closed subgroup scheme, which is finite, flat and multiplicative. Then there is a finite flat closed subgroup scheme  $\phi_0 : H_0 \hookrightarrow G$ , such that  $\phi_{0,T} \simeq \phi$ .*

*Proof.* Taking Cartier duals, we get a morphism

$$\phi^\vee : G_T^\vee \rightarrow H^\vee.$$

Notice that  $H^\vee$  is étale over  $T$ , since  $H$  is multiplicative. By radicial invariance of étale morphisms, there is a finite flat group scheme  $J_0 \rightarrow S$ , such that  $J_{0,T} \simeq H^\vee$ . Notice also that the morphism  $\phi^\vee$  is given by a section of the first projection

$$G_T^\vee \times_T H^\vee \rightarrow G_T^\vee$$

and since  $H^\vee$  is étale over  $T$ , the image of this section is open and closed (see [Milne 1980, Corollary 3.12]). Since the projection morphism

$$G_T^\vee \times_T H^\vee \rightarrow G^\vee \times_S J_0$$

is also radicial, this open set comes from a unique open subset of  $G \times_S J_0$  and this open subset defines an open and closed subscheme of  $G^\vee \times_S J_0$ , which is isomorphic to  $G^\vee$  via the first projection. Hence the morphism  $\phi^\vee$  comes from a unique morphism  $G^\vee \rightarrow J_0$ . Taking the Cartier dual of this morphism gives the morphism  $\phi_0$ . □

Recall that a commutative finite flat group scheme  $\psi : G \rightarrow S$  over  $S$  is said to be *of height one* if  $F_{G/S} = \epsilon_{G/S} \circ \psi$ . Recall also that a (sheaf in) commutative  $p$ -Lie (resp.  $p$ -coLie) algebras  $V$  over  $S$  is a coherent locally free sheaf  $V$  on  $S$  together with a morphism of  $\mathcal{O}_S$ -modules  $F_S^*(V) \rightarrow V$  (resp.  $V \rightarrow F_S^*(V)$ ). A morphism of commutative  $p$ -Lie (resp.  $p$ -coLie) algebras  $V \rightarrow W$  is a morphism of  $\mathcal{O}_S$ -modules from  $V$  to  $W$  satisfying an evident compatibility condition. There is a covariant functor  $\text{Lie}(\cdot)$  (resp. contravariant functor  $\text{coLie}(\cdot)$ ) from the category of commutative finite flat group schemes of height one over  $S$  to the category of commutative  $p$ -Lie (resp.  $p$ -coLie) algebras, which sends a group scheme  $G$  over  $S$  to  $\text{Lie}(G) := \epsilon_{G/S}^*(\Omega_{G/S})^\vee$  (resp.  $\text{coLie}(G) := \epsilon_{G/S}^*(\Omega_{G/S})$ , together with the morphism

$$\begin{aligned} \text{Lie}(V_{G^{(p)}/S}) &:= (V_{G^{(p)}/S}^*)^\vee : F_S^*(\text{Lie}(G)) = \text{Lie}(G^{(p)}) \rightarrow \text{Lie}(G) \\ (\text{resp. } \text{coLie}(V_{G^{(p)}/S}) &:= V_{G^{(p)}/S}^* : \text{coLie}(G) \rightarrow F_S^*(\text{coLie}(G^{(p)})) = \text{coLie}(G^{(p)})). \end{aligned}$$

Here  $(V_{G^{(p)}/S}^*)^\vee$  (resp.  $V_{G^{(p)}/S}^*$ ) is the dual of the pull-back morphism  $V_{G^{(p)}/S}^*$  (resp. is the pull-back morphism) on differentials induced by the Verschiebung morphism  $V_{G^{(p)}/S}$ .

The category of sheaves in commutative  $p$ -Lie algebras is tautologically antiequivalent to the category of sheaves in commutative  $p$ -coLie algebras.

It can be shown that  $\text{Lie}$  is an equivalence of additive categories (see [SGA 3, 2011, Exposé VIIA, Remark 7.5]). In particular, a sequence of finite flat group schemes of height one

$$0 \rightarrow G' \rightarrow G \rightarrow G'' \rightarrow 0$$

is exact if and only if the sequence

$$0 \rightarrow \text{Lie}(G') \rightarrow \text{Lie}(G) \rightarrow \text{Lie}(G'') \rightarrow 0$$

is a sequence of commutative  $p$ -Lie algebras. Furthermore, we have

$$\text{order}(G) = p^{\text{rk}(\text{Lie}(G))}$$

(see [Mumford 1970, Proof of Theorem, p. 139, paragraph 14]).

**Lemma 4.3.** *Let  $\phi : V \rightarrow W$  be a morphism of commutative  $p$ -Lie algebras. Then the image  $\text{Im}(\phi)$  (resp. the kernel  $\ker(\phi)$ ) of  $\phi$  as a morphism of  $\mathcal{O}_S$ -modules is endowed with a unique structure of commutative  $p$ -Lie algebra, such that the morphism  $\text{Im}(\phi) \rightarrow W$  (resp.  $\ker(\phi) \rightarrow V$ ) is a morphism of commutative  $p$ -Lie algebras.*

*Proof.* The proof is left to the reader. □

If  $\phi : V \rightarrow W$  is an injective morphism of commutative  $p$ -Lie algebras, we shall say that  $\text{Im}(\phi)$  is a subsheaf in commutative  $p$ -Lie algebras. Beware that in this situation, the arrow  $\phi$  might have no cokernel in the category of commutative  $p$ -Lie algebras. So in particular,  $\text{Im}(\phi)$  might not correspond to a subgroup scheme. On the other hand, if the quotient of  $\mathcal{O}_S$ -modules  $W/\text{Im}(\phi)$  is locally free, then  $W/\text{Im}(\phi)$  can be endowed with an evident commutative  $p$ -Lie algebra structure, making it into a cokernel of  $W$  by  $\text{Im}(\phi)$  in the category of commutative  $p$ -Lie algebras. In that case,  $\text{Im}(\phi)$  corresponds to a subgroup scheme.

We shall say that a finite flat commutative group scheme  $G$  of height one (or its associated commutative  $p$ -Lie algebra) is *biinfinitesimal* if the associated morphism  $F_S^*(\text{Lie}(G)) \rightarrow \text{Lie}(G)$  is nilpotent. To say that  $F_S^*(\text{Lie}(G)) \rightarrow \text{Lie}(G)$  is nilpotent means that for some  $n \geq 1$ , the composition

$$F_S^{\circ n,*}(\text{Lie}(G)) \rightarrow F_S^{\circ(n-1),*}(\text{Lie}(G)) \rightarrow \dots \rightarrow F_S^*(\text{Lie}(G)) \rightarrow \text{Lie}(G) \rightarrow 0$$

vanishes. We notice without proof that if

$$0 \rightarrow G' \rightarrow G \rightarrow G'' \rightarrow 0$$

is an exact sequence of commutative finite flat group schemes, then  $G'$  and  $G''$  are biinfinitesimal if and only if  $G$  is biinfinitesimal. Note also that a finite flat commutative group scheme  $G$  of height one is multiplicative if and only if the associated morphism  $F_S^*(\text{Lie}(G)) \rightarrow \text{Lie}(G)$  is an isomorphism. This implies that if  $G_1$  and  $G_2$  are finite flat group schemes of height one over  $S$ , where  $G_1$  is biinfinitesimal and  $G_2$  is multiplicative then there are no nonzero morphisms of group schemes from  $G_1$  to  $G_2$  and also no nonzero morphisms of group schemes from  $G_2$  to  $G_1$ .

We inserted the following alternative proof of a special case of [Lemma 4.2](#) to show the mechanics of  $p$ -Lie algebras at work in a simple situation.

*Second proof of Lemma 4.2 when  $G$  is of height one and  $T$  is smooth.* We may assume that  $T \simeq S$  and that  $T \rightarrow S$  is a power  $F_S^{\circ n}$  of  $F_S$ . By induction on  $n$ , we are reduced to prove the statement for  $n = 1$ .

We are given a commutative diagram with exact rows and columns

$$\begin{array}{ccccccc} & & 0 & & & & \\ & & \downarrow & & & & \\ 0 & \longrightarrow & F_T^*(\text{Lie}(H)) & \xrightarrow{F_T^*(\text{Lie}(\phi))} & F_T^*(\text{Lie}(G)_T) & & \\ & & \downarrow \text{Lie}(V_{H/T}) & & \downarrow \text{Lie}(V_{G_T/T}) & & \\ 0 & \longrightarrow & \text{Lie}(H) & \xrightarrow{\text{Lie}(\phi)} & \text{Lie}(G)_T & & \\ & & \downarrow & & & & \\ & & 0 & & & & \end{array}$$

With the above reductions in place, this gives a commutative diagram with exact rows and columns

$$\begin{array}{ccccccc} & & 0 & & & & \\ & & \downarrow & & & & \\ 0 & \longrightarrow & F_S^*(\text{Lie}(H)) & \xrightarrow{F_S^*(\text{Lie}(\phi))} & F_S^{\circ 2,*}(\text{Lie}(G)) & & \\ & & \downarrow \text{Lie}(V_{H/S}) & & \downarrow F_S^*(\text{Lie}(V_{G/S})) & & \\ 0 & \longrightarrow & \text{Lie}(H) & \xrightarrow{\text{Lie}(\phi)} & F_S^*(\text{Lie}(G)) & & \\ & & \downarrow & & & & \\ & & 0 & & & & \end{array}$$

Now consider the commutative diagram

$$\begin{array}{ccc}
 F_S^*(\mathrm{Lie}(H)) & \xrightarrow{F_S^*(\mathrm{Lie}(\phi))} & F_S^{\circ 2,*}(\mathrm{Lie}(G)) \\
 \downarrow \mathrm{Lie}(V_{H/S}) & \searrow & \downarrow F_S^*(\mathrm{Lie}(V_{G/S})) \\
 \mathrm{Lie}(H) & \xrightarrow{\mathrm{Lie}(\phi)} & F_S^*(\mathrm{Lie}(G)) \\
 & \searrow & \downarrow \mathrm{Lie}(G) \\
 & & \mathrm{Lie}(G)
 \end{array}$$

where the diagonal arrows are defined so that the diagram becomes commutative. The labelling of the arrows shows that the upper triangle is the base change by  $F_S$  of the lower triangle. Hence the image of  $\mathrm{Lie}(\phi)$  is the base change by  $F_S$  of the image of  $\mathrm{Lie}(H)$  in  $\mathrm{Lie}(G)$ , since  $\mathrm{Lie}(V_{H/S})$  is an isomorphism. So  $H_0$  can be defined as the group scheme of height one associated with the image of  $\mathrm{Lie}(H)$  in  $\mathrm{Lie}(G)$ .  $\square$

**Lemma 4.4.** *Let  $V$  be a sheaf in commutative  $p$ -Lie algebras  $V$  over  $S$ . Suppose that the HN filtration*

$$0 = V_0 \subsetneq V_1 \subsetneq V_2 \subsetneq \cdots \subsetneq V_{\mathrm{hn}(V)} = V$$

*of  $V$  is Frobenius semistable. Then for any  $V_i$  such that  $\mu_{\min}(V_i) \geq 0$ ,  $V_i$  is a subsheaf in commutative  $p$ -Lie algebras  $V$  over  $S$ . If  $\mu_{\min}(V_i) > 0$  then  $V_i$  is biinfiniteesimal.*

*Proof.* For the first statement, consider the morphism  $\phi : F_S^*(V_i) \rightarrow V$  given by the composition of the inclusion  $F_S^*(V_i) \rightarrow F_S^*(V)$  with the morphism  $F_S^*(V) \rightarrow V$  given by the commutative  $p$ -Lie algebra structure. We have to check that the image of  $\phi$  lies in  $V_i$ . The composition of  $\phi$  with the quotient morphism  $V \rightarrow V/V_i$  gives a morphism  $F_S^*(V_i) \rightarrow V/V_i$  and it is equivalent to check that this morphism vanishes. Now we compute

$$\mu_{\min}(F_S^*(V_i)) = p \cdot \mu(V_i/V_{i-1}) \quad \text{and} \quad \mu_{\max}(V/V_i) = \mu(V_{i+1}/V_i) < \mu(V_i/V_{i-1}),$$

and thus  $\mu_{\min}(F_S^*(V_i)) > \mu_{\max}(V/V_i)$ . We conclude that  $\mathrm{Hom}_S(F_S^*(V_i), V/V_i) = 0$  (see the discussion after [Theorem 3.1](#)) which concludes the proof of the first statement. To prove the second statement, it is sufficient by the remarks preceding the lemma to show that  $V_i/V_{i-1}$  is biinfiniteesimal for all indices  $i$  such that  $\mu(V_i/V_{i-1}) > 0$ . By the above computation, we have

$$\mu_{\min}(F_S^*(V_i/V_{i-1})) = \mu(F_S^*(V_i/V_{i-1})) = p \cdot \mu(V_i/V_{i-1})$$

and thus  $\mu_{\min}(F_S^*(V_i/V_{i-1})) > \mu(V_i/V_{i-1})$ . Again, this implies that  $\mathrm{Hom}_S(F_S^*(V_i/V_{i-1}), V_i/V_{i-1}) = 0$ , showing that  $V_i/V_{i-1}$  is biinfiniteesimal.  $\square$

**Remark 4.5.** As explained in the introduction, a characteristic 0 analogue of [Lemma 4.4](#) can be found in [\[Bost 2004, Lemma 2.9\]](#). See also [\[Shepherd-Barron 1992, Lemma 9.1.3.1\]](#), where a variant of a special case of [Lemma 4.4](#) is proven under the assumption that  $p$  is sufficiently large.

**Lemma 4.6.** *Let  $G$  be a commutative finite flat group scheme of height one over  $S$  and suppose given an exact sequence*

$$0 \rightarrow G_{\text{binf}} \rightarrow G \rightarrow G_\mu \rightarrow 0$$

*of finite flat group schemes such that  $G_\mu$  is multiplicative and  $G_{\text{binf}}$  is biinfinitesimal. Then the sequence splits and this splitting is unique.*

*Proof.* Consider the commutative diagram with exact rows and columns

$$\begin{array}{ccccccc}
 0 & \longrightarrow & \ker(\text{Lie}(V_{G_{\text{binf}}^{(p^n)}/S}^{(n)})) & \longrightarrow & \ker(\text{Lie}(V_{G^{(p^n)}/S}^{(n)})) & \longrightarrow & 0 \\
 & & \downarrow \simeq & & \downarrow & & \downarrow \\
 0 & \longrightarrow & F_S^{\text{on},*}(\text{Lie}(G_{\text{binf}})) & \longrightarrow & F_S^{\text{on},*}(\text{Lie}(G)) & \longrightarrow & F_S^{\text{on},*}(\text{Lie}(G_\mu)) \longrightarrow 0 \\
 & & \downarrow =0 & & \downarrow & & \downarrow \simeq \\
 0 & \longrightarrow & \text{Lie}(G_{\text{binf}}) & \longrightarrow & \text{Lie}(G) & \longrightarrow & \text{Lie}(G_\mu) \longrightarrow 0
 \end{array}$$

where  $n \geq 0$  is chosen so that  $V_{G_{\text{binf}}^{(p^n)}/S}^{(n),*} = 0$ . Then the image of the arrow

$$F_S^{\text{on},*}(\text{Lie}(G)) \rightarrow \text{Lie}(G)$$

splits the bottom sequence. For the unicity of the splitting, note that for any two splittings  $\sigma_1, \sigma_2$  of the bottom sequence the morphism  $\sigma_1 - \sigma_2 : \text{Lie}(G_\mu) \rightarrow \text{Lie}(G)$  of vector bundles factors through the image of  $\text{Lie}(G_{\text{binf}})$ . It thus defines a morphism of vector bundles  $\text{Lie}(G_\mu) \rightarrow \text{Lie}(G_{\text{binf}})$ , which is by construction a morphism of  $p$ -Lie algebras. Such a morphism must vanish (see the discussion after Lemma 4.3). Thus  $\sigma_1 = \sigma_2$ . □

**Lemma 4.7.** *Let  $G$  be a commutative finite flat group scheme of height one over  $S$ . Suppose that  $\text{Lie}(G)$  is Frobenius semistable of slope 0. Let  $n \geq 0$  be such that  $\text{rk}(\ker(V_{G^{(p^n)}/S}^{(n),*}))$  is maximal. Then there is a canonical decomposition*

$$G^{(p^n)} \simeq H_{\text{binf}} \times_S H_\mu,$$

where  $H_{\text{binf}}$  (resp.  $H_\mu$ ) is a biinfinitesimal (resp. multiplicative) finite flat group scheme over  $S$ .

*Proof.* Consider the commutative diagram

$$\begin{array}{ccccccc}
 0 & \longrightarrow & F_S^{\text{on},*}(\ker(\text{Lie}(V_{G^{(p^n)}/S}^{(n)}))) & \longrightarrow & F_S^{\text{on},*}(\ker(\text{Lie}(V_{G^{(p^n)}/S}^{(n)}))) & \longrightarrow & 0 \\
 & & \downarrow \sim & & \downarrow & & \downarrow \\
 0 & \longrightarrow & F_S^{\text{on},*}(\ker(\text{Lie}(V_{G^{(p^n)}/S}^{(n)}))) & \longrightarrow & F_S^{\text{on}(2n),*}(\text{Lie}(G)) & \longrightarrow & F_S^{\text{on},*}(W) \longrightarrow 0 \\
 & & \downarrow =0 & & \downarrow & & \downarrow \\
 0 & \longrightarrow & \ker(\text{Lie}(V_{G^{(p^n)}/S}^{(n)})) & \longrightarrow & F_S^{\text{on},*}(\text{Lie}(G)) & \longrightarrow & W \longrightarrow 0
 \end{array}$$

where  $n \geq 0$  is such that  $\text{rk}(\ker(\text{Lie}(V_{G^{(p^n)}/S}^{(n)})))$  is maximal and  $W$  is the image of  $\text{Lie}(V_{G^{(p^n)}/S}^{(n)})$ . The two bottom rows and the two leftmost columns in this diagram are exact by construction. Furthermore the map  $F_S^{(n),*}W \rightarrow W$  is a monomorphism for otherwise  $\text{rk}(\ker(\text{Lie}(V_{G^{(p^n)}/S}^{(n)})))$  is not maximal. The diagram thus has exact rows and columns. Since the second row gives a surjection

$$F_S^{\circ(2n),*}(\text{Lie}(G)) \rightarrow F_S^{\circ n,*}(W)$$

we have  $\mu_{\min}(F_S^{\circ n,*}(W)) \geq 0$ . Also, since the second column gives an injection

$$F_S^{\circ n,*}(W) \hookrightarrow F_S^{(n),*}(\text{Lie}(G))$$

we have  $\mu_{\max}(F_S^{\circ n,*}(W)) \leq 0$ . Thus  $F_S^{\circ n,*}(W)$  is of slope 0. Thus  $W$  is also of slope 0. Hence by [Lemma 3.2](#), the monomorphism

$$F_S^{\circ n,*}(W) \rightarrow W$$

is an isomorphism. Now we see that the image of the morphism  $F_S^{\circ(2n),*}(\text{Lie}(G)) \rightarrow F_S^{\circ n,*}(\text{Lie}(G))$  splits the bottom sequence. □

**Lemma 4.8.** *Let  $G$  be a finite flat commutative group scheme of height one over  $S$ . There exists a (necessarily unique) multiplicative subgroup scheme  $G_\mu \hookrightarrow G$ , such that if  $H$  is a multiplicative subgroup scheme of height one over  $S$  and  $f : H \rightarrow G$  is a morphism of group schemes, then  $f$  factors through  $G_\mu$ . Furthermore, for any  $n \geq 0$ , we have  $(G_\mu)^{(p^n)} = (G^{(p^n)})_\mu$ . If  $G$  is multiplicative over a dense open subset of  $S$  and  $\text{Lie}(G)$  has Frobenius semistable HN filtration then  $\text{Lie}(G) = \text{Lie}(G)_{\leq 0}$  and  $G_\mu$  corresponds to the subgroup scheme associated with  $\text{Lie}(G)_{=0}$ .*

*Proof.* In view of [Lemma 4.2](#), we may replace  $G$  by  $G^{(p^n)}$  for any  $n \geq 0$  and in particular suppose that  $\text{Lie}(G)$  has a Frobenius semistable HN filtration. Let  $f : H \rightarrow G$  be a morphism of group schemes and consider the corresponding map

$$\text{Lie}(f) : \text{Lie}(H) \rightarrow \text{Lie}(G).$$

Since  $H$  is multiplicative,  $\text{Lie}(H)$  is Frobenius semistable of slope 0 (this is a consequence of [Theorem 3.1](#)). Thus the image of  $\text{Lie}(f)$  lies in  $\text{Lie}(G)_{\geq 0}$ . According to [Lemma 4.4](#) there is an exact sequence of  $p$ -Lie algebras

$$0 \rightarrow \text{Lie}(G)_{>0} \rightarrow \text{Lie}(G)_{\geq 0} \xrightarrow{\pi} \text{Lie}(G)_{=0} \rightarrow 0$$

and we may assume according to [Lemma 4.7](#) that there is a splitting

$$\text{Lie}(G)_{=0} \simeq \text{Lie}(G)_{=0,\text{binf}} \oplus \text{Lie}(G)_{=0,\mu}$$

of  $\text{Lie}(G)_{=0}$  into multiplicative and biinfinitesimal part (we might have to twist  $G$  some more for this). The inverse image of  $\text{Lie}(G)_{=0,\mu}$  by  $\pi$  gives a  $p$ -Lie subalgebra  $\pi^*(\text{Lie}(G)_{=0,\mu})$  of  $\text{Lie}(G)_{\geq 0}$ . This gives an exact sequence

$$0 \rightarrow \pi^*(\text{Lie}(G)_{=0,\mu}) \rightarrow \text{Lie}(G)_{\geq 0} \rightarrow \text{Lie}(G)_{=0,\text{binf}} \rightarrow 0$$

Since  $\text{Lie}(H)$  is multiplicative, the image of  $\text{Lie}(H)$  in  $\text{Lie}(G)_{=0, \text{binf}}$  vanishes and thus the image of  $\text{Lie}(H)$  lies in  $\pi^*(\text{Lie}(G)_{=0, \mu})$ . On the other hand by [Lemma 4.6](#) and [Lemma 4.4](#), we have again a canonical decomposition

$$\pi^*(\text{Lie}(G)_{=0, \mu})_{\mu} \oplus \pi^*(\text{Lie}(G)_{=0, \mu})_{\text{binf}}$$

into multiplicative and biinfinitesimal part and thus the image of  $\text{Lie}(f)$  lies in  $\pi^*(\text{Lie}(G)_{=0, \mu})_{\mu}$ . Now  $\pi^*(\text{Lie}(G)_{=0, \mu})_{\mu}$  is a multiplicative  $p$ -Lie subalgebra of  $\text{Lie}(G)$  and it defines the required subgroup scheme.

If  $G$  is multiplicative over an open subset of  $S$  then we have an injection

$$F_S^{n, *}(\text{Lie}(G)) \hookrightarrow \text{Lie}(G)$$

(obtained by composition) for any  $n \geq 0$  and thus if  $\text{Lie}(G)$  has Frobenius semistable HN filtration then we must have  $\text{Lie}(G) = \text{Lie}(G)_{\leq 0}$ . Secondly the morphism  $F_S^*(\text{Lie}(G)) \hookrightarrow \text{Lie}(G)$  then induces an injection

$$F_S^*(\text{Lie}(G)_{=0}) \hookrightarrow \text{Lie}(G)_{=0}$$

and since both source and target in this map have the same rank and the same slope, we deduce from [Lemma 3.2](#) that this map must be an isomorphism. Thus  $\text{Lie}(G)_{=0}$  is multiplicative and by the explicit construction above, it is associated with  $G_{\mu}$ . □

**Remark 4.9.** Note that the “connected étale” decomposition of  $G_K^{\vee}$  (see the beginning of [\[Tate 1997\]](#)) gives a canonical exact sequence of group schemes

$$0 \rightarrow (G_K^{\vee})_{\text{inf}} \rightarrow G_K^{\vee} \rightarrow (G_K^{\vee})_{\text{et}} \rightarrow 0$$

over  $K$ , where  $(G_K^{\vee})_{\text{inf}}$  is an infinitesimal group scheme and  $(G_K^{\vee})_{\text{et}}$  is an étale group scheme over  $K$ . The group scheme  $(G_K^{\vee})_{\text{et}}$  corresponds to a representation of  $\text{Gal}(K^{\text{sep}}|K)$  into a finite  $p$ -group  $E$  and one might be tempted to think that  $G_{\mu}$  is the Cartier dual of the group scheme corresponding to the largest unramified quotient of  $E$ , i.e., the largest quotient of  $E$ , such that the action of  $\text{Gal}(K^{\text{sep}}|K)$  factors through the fundamental group  $\pi_1(S)$ . This not so, however. Indeed, consider a finite flat commutative group scheme  $G$  of height one, which is such that  $\bar{\mu}_{\text{max}}(\text{Lie}(G)) < 0$ . Then  $G_{\mu} = 0$  and for any finite flat base change  $S' \rightarrow S$ , we also have  $(G_{S'})_{\mu} = 0$ . On the other hand  $(G_K^{\vee})_{\text{et}}$  will become constant (and hence entirely unramified) after a finite separable field extension  $K'|K$ .

**4C. Quotients of semiabelian schemes by finite flat multiplicative group schemes.** Let  $S$  be a smooth, projective and geometrically connected curve over a perfect field  $k$ .

**Lemma 4.10.** *Let  $\mathcal{A} \rightarrow S$  be a semiabelian scheme. Suppose that there is an open dense subset  $U \subseteq S$ , such that  $\mathcal{A}_U \rightarrow U$  is an abelian scheme. Suppose that  $G \hookrightarrow \mathcal{A}$  is a finite, flat, closed subgroup scheme. Then the quotient scheme  $\mathcal{A}/G$  is also a semiabelian scheme and  $(\mathcal{A}/G)_U \rightarrow U$  is an abelian scheme.*

*Proof.* Since the quotient morphism  $q : \mathcal{A} \rightarrow \mathcal{A}/G$  is faithfully flat, the group scheme  $\mathcal{A}/G$  also has geometrically regular fibres (and is flat). Hence  $\mathcal{A}/G$  is smooth over  $S$ . Over  $U$ , its fibres are proper since

the quotient morphism is also proper and they are thus abelian varieties. In other words,  $(\mathcal{A}/G)_U \rightarrow U$  is an abelian scheme. Now let  $s \in S$ . Since  $(\mathcal{A}/G)_s$  is smooth, we know by the Barsotti–Chevalley theorem (see [Milne 2017, Theorem 10.25, p. 157]) that  $(\mathcal{A}/G)_s$  sits in the middle of an exact sequence

$$0 \rightarrow E_1 \rightarrow (\mathcal{A}/G)_s \rightarrow A_1 \rightarrow 0, \tag{1}$$

where  $A_1$  is an abelian variety over  $s$  and  $E_1$  is a connected affine algebraic group variety over  $s$ . The subgroup variety  $E_1$  is maximal among connected affine subgroup varieties of  $(\mathcal{A}/G)_s$  (see [Milne 2017, Theorem 10.5, p. 153 and proof, and Theorem 10.24, p. 156]). Finally it has the form  $E_1 = T_1 \times_s U$ , where  $T_1$  is a torus and  $U$  is a connected unipotent group variety (see [Milne 2017, Chapter 10 (i), p. 161]). When we write that the sequence (1) is exact, we mean that the third morphism is faithfully flat and that its kernel is  $E_1$ .

By assumption, the corresponding presentation for  $\mathcal{A}_s$  is

$$0 \rightarrow T \rightarrow \mathcal{A}_s \rightarrow A_0 \rightarrow 0,$$

where  $T$  is a torus and  $A_0$  is an abelian variety, both over  $s$ .

Let  $D$  be the identity component of the closed subgroup scheme  $q_s^{-1}(U \times 0)$  of  $\mathcal{A}_s$  (see [Milne 2017, Proposition 1.14] for details). Since  $s$  is perfect the closed subscheme  $D_{\text{red}}$  of  $D$  is a closed subgroup scheme of  $D$  (see [Milne 2017, Corollary 1.25, p. 24]). Moreover  $D$  and hence  $D_{\text{red}}$  is affine, since  $q_s$  is finite. Since  $T$  is the maximal connected affine subgroup variety of  $\mathcal{A}_s$ , we see that  $D_{\text{red}}$  must be contained in  $T$ . However, every closed subgroup scheme of a multiplicative group over  $s$  is multiplicative (see [SGA 3<sub>II</sub> 1970, 8.1, Exposé IX]) and thus  $D_{\text{red}}$  is multiplicative. Thus  $D_{\text{red}}$  is contained in the kernel of the morphism  $q_s^{-1}(U \times 0) \rightarrow U \times 0$  (because there are no nontrivial morphisms between multiplicative and unipotent algebraic groups — see [Milne 2017, Corollary 15.18, p. 255]). Now notice that  $q_s^{-1}(U \times 0)(\bar{s})/D(\bar{s})$  is a finite set (see [Milne 2017, Proposition 1.14, p. 21]). On the other hand  $q_s(D(\bar{s})) = \{0\}$  by the above so  $U(\bar{s})$  must be finite. Since  $U$  is smooth, it must thus be trivial. This shows that  $(\mathcal{A}/G)_s$  is an extension of an abelian variety by a torus. Since  $s \in S$  was arbitrary, we see that  $\mathcal{A}/G$  is a semiabelian scheme. □

**Lemma 4.11.** *Let  $G \rightarrow S$  be a finite flat group scheme of multiplicative type. Then there is a finite étale morphism  $T \rightarrow S$  such that  $G_T$  is a diagonalisable group scheme.*

*Proof.* See [SGA 3<sub>II</sub> 1970, Exposé IX, Introduction]. □

**Lemma 4.12.** *Let  $\mathcal{A} \rightarrow S$  be a smooth commutative group scheme. Suppose that  $G \hookrightarrow \mathcal{A}$  is a finite, flat, closed subgroup scheme, which is multiplicative. Then*

$$\deg(\omega_{\mathcal{A}}) = \deg(\omega_{\mathcal{A}/G}).$$

*Proof.* By Lemma 4.11, we may assume that  $G$  is diagonalisable. In particular, we may assume that there is a finite group scheme  $G_0 \rightarrow \text{Spec}(k)$  such that  $G_{0,S} \simeq G$ . Let  $\mathcal{B} := \mathcal{A}/G$ . Let  $f : \mathcal{A} \rightarrow S$  and  $g : \mathcal{B} \rightarrow S$

be the structural morphisms and let  $\pi : \mathcal{A} \rightarrow \mathcal{B}$  be the quotient morphism. The triangle of cotangent complexes associated with the morphisms  $\pi$ ,  $g$  and  $f$  gives an exact sequence

$$0 \rightarrow \mathcal{H}_1(\text{CT}(\pi)) \rightarrow \pi^*(\Omega_g) \rightarrow \Omega_f \rightarrow \Omega_\pi \rightarrow 0, \tag{2}$$

where  $\text{CT}(\pi)$  is the cotangent complex of  $\pi$  and  $\mathcal{H}_1(\text{CT}(\pi))$  is its first homology sheaf. Now  $\pi$  makes  $\mathcal{A}$  into a torsor over  $\mathcal{B}$  and under  $G_B$ . Hence there is a faithfully flat morphism  $T \rightarrow \mathcal{B}$  (for instance, we may take  $T = \mathcal{A}$ ), such that  $\mathcal{A}_T \simeq (G_B) \times_{\mathcal{B}} T$ . In particular we have

$$\Omega_{\pi_T} \simeq \Omega_{G_0/k, T} \quad \text{and} \quad \mathcal{H}_1(\text{CT}(\pi_T)) \simeq \mathcal{H}_1(\text{CT}(G_0/k))_T$$

because the homology sheaves of the cotangent complex of  $G_0$  over  $k$  are flat (since they are  $k$ -vector spaces).

On the other hand, since  $T \rightarrow \mathcal{B}$  is flat, we have

$$\Omega_{\pi_T} \simeq \Omega_{\pi, T} \quad \text{and} \quad \mathcal{H}_1(\text{CT}(\pi_T)) \simeq \mathcal{H}_1(\text{CT}(\pi))_T.$$

Finally, notice that  $\Omega_{G_0/k, T}$  and  $\mathcal{H}_1(\text{CT}(G_0/k))_T$  are flat and thus by flat descent, the sheaves  $\mathcal{H}_1(\text{CT}(\pi))$  and  $\Omega_\pi$  are flat (in other words: locally free). Hence the sequence

$$0 \rightarrow \epsilon_{\mathcal{A}/S}^*(\mathcal{H}_1(\text{CT}(\pi))) \rightarrow \epsilon_{\mathcal{B}/S}^*(\Omega_g) \rightarrow \epsilon_{\mathcal{A}/S}^*(\Omega_f) \rightarrow \epsilon_{\mathcal{A}/S}^*(\Omega_\pi) \rightarrow 0 \tag{3}$$

is also exact. Furthermore, we then have

$$\epsilon_{\mathcal{A}/S}^*(\mathcal{H}_1(\text{CT}(\pi))) \simeq \mathcal{H}_1(\text{CT}(G_0/k))_S \quad \text{and} \quad \epsilon_{\mathcal{A}/S}^*(\Omega_\pi) \simeq \Omega_{G_0/k, S}$$

and thus the sheaves  $\epsilon_{\mathcal{A}/S}^*(\mathcal{H}_1(\text{CT}(\pi)))$  and  $\epsilon_{\mathcal{A}/S}^*(\Omega_\pi)$  are trivial sheaves. In particular, we have that  $\text{deg}(\epsilon_{\mathcal{A}/S}^*(\mathcal{H}_1(\text{CT}(\pi)))) = \text{deg}(\epsilon_{\mathcal{A}/S}^*(\Omega_\pi)) = 0$  and by the additivity of  $\text{deg}(\cdot)$ , we deduce from the existence of the sequence (3) that  $\text{deg}(\omega_{\mathcal{A}}) = \text{deg}(\omega_{\mathcal{A}/G})$ . □

**Remark 4.13.** The computation of the cotangent complex made in the proof of [Lemma 4.11](#) is in essence also contained in [\[Ekedahl 1988, Proposition 1.1\]](#) (but the assumptions made there are not quite the right ones for us).

### 5. Proofs of the claims made in [Section 2A](#)

We now use the terminology of the introduction. So let  $k$  be a finite field of characteristic  $p > 0$  and let  $S$  be a smooth, projective and geometrically connected curve over  $k$ . Let  $K := \kappa(S)$  be its function field. Let  $A$  be an abelian variety of dimension  $g$  over  $K$ . Fix an algebraic closure  $\bar{K}$  of  $K$ . Let  $K^{\text{perf}} \subseteq \bar{K}$  be the maximal purely inseparable extension of  $K$  and let  $K^{\text{unr}} \subseteq K^{\text{sep}}$  be the maximal separable extension of  $K$ , which is unramified above every place of  $K$ . Finally, we let  $\mathcal{A}$  be a smooth commutative group scheme over  $S$  such that  $\mathcal{A}_K = A$ .

*Proof of [Theorem 2.1](#).* Recall the statement: there exists a (necessarily unique) multiplicative subgroup scheme  $G_{\mathcal{A}} \hookrightarrow \ker F_{\mathcal{A}/S}$ , with the following property: if  $H$  is a multiplicative, finite and flat group

scheme of height one over  $S$  and  $f : H \rightarrow \ker F_{\mathcal{A}/S}$  is a morphism of group schemes, then  $f$  factors through  $G_{\mathcal{A}}$ . If  $A$  is ordinary and  $\omega_{\mathcal{A}}$  is not ample then the order of  $G_{\mathcal{A}}$  is  $p^{\overline{\text{rk}}_{\min}(\omega_{\mathcal{A}})}$ . If  $\phi : \mathcal{A} \rightarrow \mathcal{B}$  is a morphism of smooth commutative group schemes over  $S$ , then the restriction of  $\phi$  to  $G_{\mathcal{A}}$  factors through  $G_{\mathcal{B}}$ . Furthermore, we have  $\deg(\omega_{\mathcal{A}}) = \deg(\omega_{\mathcal{A}/G_{\mathcal{A}}})$ .

In spite of its lengthy statement, the proof [Theorem 2.1](#) readily follows from [Lemmata 4.8](#) and [4.12](#). More precisely, we simply have to define  $G_{\mathcal{A}} := (\ker F_{\mathcal{A}/S})_{\mu}$  in the notation of [Lemma 4.8](#). The equality  $\deg(\omega_{\mathcal{A}}) = \deg(\omega_{\mathcal{A}/G_{\mathcal{A}}})$  now follows from [Lemma 4.12](#).  $\square$

*Proof of [Proposition 2.5](#).* Recall the assumptions of [Proposition 2.5](#):  $A$  is ordinary,  $\mathcal{A}$  is semiabelian and  $A(K^{\text{perf}})$  is not finitely generated. We have to prove that  $G_{\mathcal{A}}$  is of order  $> 1$  and that  $\mathcal{A}/G_{\mathcal{A}}$  is also semiabelian.

We know that  $\bar{\mu}_{\min}(\omega_{\mathcal{A}/S}) \geq 0$  by [Lemma 4.8](#) and since  $A(K^{\text{perf}})$  is not finitely generated, we know by [Theorem B.1](#) that  $\bar{\mu}_{\min}(\omega_{\mathcal{A}/S}) = 0$ . [Proposition 2.5](#) now follows from [Theorem 2.1](#) and [Lemma 4.10](#).  $\square$

*Proof of [Proposition 2.6](#).* Recall the assumptions of [Proposition 2.6](#):  $A$  is ordinary,  $\mathcal{A}$  is semiabelian over  $S$  and  $A(K^{\text{perf}})$  is not finitely generated. We have to prove that there a finite flat morphism

$$\phi : \mathcal{A} \rightarrow \mathcal{B},$$

where  $\mathcal{B}$  is a semiabelian over  $S$  and a finite flat morphism

$$\lambda : \mathcal{B} \rightarrow \mathcal{B}$$

such that  $\ker(\phi)$  are  $\ker(\lambda)$  are multiplicative group schemes and such that the order of  $\ker(\lambda)$  is  $> 1$ .

Consider now  $\mathcal{A}_1 := \mathcal{A}/G_{\mathcal{A}}$ . By [Lemma 4.10](#), the group scheme  $\mathcal{A}_1$  is also semiabelian and of course  $A_1 := \mathcal{A}_{1,K}$  is also an ordinary abelian variety. We also have that  $A_1(K^{\text{perf}})$  is not finitely generated, since the natural map  $A(K^{\text{perf}}) \rightarrow A_1(K^{\text{perf}})$  has finite kernel. Finally, the quotient morphism is  $\mathcal{A} \rightarrow \mathcal{A}_1$  is finite, flat, with multiplicative kernel and  $G_{\mathcal{A}}$  is nontrivial by [Proposition 2.5](#).

Repeating the above procedure for  $\mathcal{A}_1$  in place of  $\mathcal{A}$  and continuing this way, we obtain an infinite sequence of semiabelian schemes over  $S$

$$\mathcal{A} \rightarrow \mathcal{A}_1 \rightarrow \mathcal{A}_2 \rightarrow \dots, \tag{4}$$

where all the connecting morphisms are finite, flat, of degree  $> 1$  and with multiplicative kernel. Applying [Lemma 4.12](#), we see that

$$\deg(\omega_{\mathcal{A}}) = \deg(\omega_{\mathcal{A}_1}) = \deg(\omega_{\mathcal{A}_2}) = \dots$$

Let now  $K'$  be a finite separable extension of  $K$  such that  $A(K)[n] \simeq (\mathbb{Z}/n\mathbb{Z})^{2 \dim(A)}$  for some  $n \geq 3$  such that  $(p, n) = 1$ . Let  $S'$  be the normalisation of  $S$  in  $K'$ . After base-change, we obtain an infinite sequence of semiabelian schemes over  $S'$

$$\mathcal{A}_{S'} \rightarrow \mathcal{A}_{1,S'} \rightarrow \mathcal{A}_{2,S'} \rightarrow \dots \tag{5}$$

and applying a theorem of Zarhin (see [\[Rössler 2013, Theorem 3.1\]](#) for a statement, explanations and further references), we conclude that in the sequence [\(5\)](#), there are only finitely many isomorphism

classes of semiabelian schemes over  $S'$ . On the other hand, applying a basic finiteness result in Galois cohomology proven by Borel and Serre (see [Zarkhin and Parshin 1989, paragraph 3, p. 69]), we can now conclude that in the sequence (4), there are also only finitely many isomorphism classes of semiabelian schemes over  $S$ .

Hence there are integers  $j > i \geq 0$  and an isomorphism

$$I : \mathcal{A}_i \simeq \mathcal{A}_j$$

over  $S$ . Letting  $\phi : \mathcal{A} \rightarrow \mathcal{A}_i$  be the constructed morphism and letting  $\lambda$  be the constructed morphism  $\mathcal{A}_i \rightarrow \mathcal{A}_j$  composed with  $I^{-1}$ , we can now conclude the proof of Proposition 2.6.  $\square$

### 6. Proofs of the claims made in Section 2B

**Theorem 2.7.** *Suppose that  $\text{Tr}_{\bar{k}|\bar{k}}(A_{\bar{k}}) = 0$ . Suppose that the action of  $\text{Gal}(K^{\text{sep}}|K)$  on  $\text{Tor}_p(A(K^{\text{unr}}))$  factors through  $\text{Gal}(K^{\text{sep}}|K)^{\text{ab}}$ . Then  $\text{Tor}_p(A(K^{\text{unr}}))$  is finite.*

*Proof.* Let  $L|K$  be the maximal subextension of  $K^{\text{unr}}|K$ , which is Galois with abelian Galois group. Since  $S$  is geometrically integral,  $K \otimes_k \bar{k}$  is a field and  $L$  contains a subfield isomorphic to  $K \otimes_k \bar{k}$  (note that  $\bar{k} = k^{\text{sep}}$  and that  $\text{Gal}(\bar{k}|k) \simeq \widehat{\mathbb{Z}}$ , which is an abelian group). Furthermore, geometric class field theory (see, e.g., [Szamuely 2010, Corollary 1.3]) tells us that  $\text{Gal}(L | K \otimes_k \bar{k})$  is a finite group. In particular, the field  $L$  is finitely generated (as a field) over  $\bar{k}$ , since  $K \otimes_k \bar{k}$  is finitely generated over  $\bar{k}$ . Now suppose to obtain a contradiction that  $\text{Tor}_p(A(K^{\text{unr}}))$  were infinite. By assumption, we have

$$\text{Tor}_p(A(K^{\text{unr}})) \subseteq \text{Tor}_p(A(L)).$$

Thus  $\text{Tor}_p(A(L))$  is infinite as well. By the Lang–Néron theorem, this implies that

$$\text{Tr}_{L|\bar{k}}(A_L) \neq 0,$$

contradicting the first assumption.  $\square$

**Proposition 2.8.** *Suppose that  $\dim(A) \leq 2$  and that  $\text{Tr}_{\bar{k}|\bar{k}}(A_{\bar{k}}) = 0$ . Then  $\text{Tor}_p(A(K^{\text{unr}}))$  is finite.*

*Proof.* Notice that if  $\text{Tor}_p(A(K^{\text{unr}}))$  is infinite then we have

$$\bigcap_{\ell \geq 0} p^\ell \cdot \text{Tor}_p(A(K^{\text{unr}})) \neq 0$$

This follows from the fact that for each  $n \geq 0$ , the set

$$\{x \in \text{Tor}_p(A(K^{\text{unr}})) \mid p^n \cdot x = 0\}$$

is finite (the details are left to the reader). Let  $G \subseteq \bigcap_{\ell \geq 0} p^\ell \cdot (\text{Tor}_p(A(K^{\text{unr}})))$  be the subgroup of elements annihilated by the multiplication by  $p$  map.

If  $G = 0$  then there the conclusion holds, because then  $\bigcap_{\ell \geq 0} p^\ell \cdot (\text{Tor}_p(A(K^{\text{unr}}))) = 0$  and thus  $\text{Tor}_p(A(K^{\text{unr}}))$  is finite by the above remark.

Suppose now that  $\#G = p$ . Then  $\bigcap_{\ell \geq 0} p^\ell \cdot \text{Tor}_p(A(K^{\text{unr}}))$  is infinite and it is a union of cyclic groups of  $p$ -power order (use the classification theorem for finite abelian groups). Thus the action of  $\text{Gal}(K^{\text{sep}}|K)$  on  $\bigcap_{\ell \geq 0} p^\ell \cdot (\text{Tor}_p(A(K^{\text{unr}})))$  factors through  $\text{Gal}(K^{\text{sep}}|K)^{\text{ab}}$ . But this contradicts [Theorem 2.7](#) and thus we must have  $\#G > p$ . If  $\#G > p$  then by the assumption that  $\dim(A) \leq 2$ , we see that we must have  $\#G = p^2$  and thus the inclusions

$$\text{Tor}_p(A(K^{\text{unr}})) \subseteq \text{Tor}_p(A(K^{\text{sep}})) \subseteq \text{Tor}_p(A(\overline{K}))$$

are both equalities. In particular,  $A$  is an ordinary abelian surface. Let now  $s \in S$  be a closed point such that  $\mathcal{A}_s$  is an ordinary abelian variety over  $s$ . Let  $W := \text{Spec}(\widehat{\mathcal{O}}_{S,s}^{\text{sh}})$  be the spectrum of the completion of the strict henselisation of the local ring at  $s$  and write  $\widehat{K}_s^{\text{sh}}$  for the fraction field of  $\widehat{\mathcal{O}}_{S,s}^{\text{sh}}$ . The abelian scheme  $\mathcal{A}_W \rightarrow W$  gives rise to an element  $e$  of

$$\text{Hom}_{\mathbb{Z}_p}(T_p(\mathcal{A}_{\overline{s}}(\overline{s})) \otimes T_p(\mathcal{A}_{\overline{s}}^\vee(\overline{s})), \widehat{\mathcal{O}}_{S,s}^{\text{sh}*}).$$

Here  $T_p(\mathcal{A}_{\overline{s}}(\overline{s}))$  and  $T_p(\mathcal{A}_{\overline{s}}^\vee(\overline{s}))$  are the  $p$ -adic Tate modules of  $\mathcal{A}_{\overline{s}}$  and  $\mathcal{A}_{\overline{s}}^\vee$  respectively and  $\widehat{\mathcal{O}}_{S,s}^{\text{sh}*}$  is the group of multiplicative units of  $\widehat{\mathcal{O}}_{S,s}^{\text{sh}}$ . The element  $e$  is called the Serre–Tate pairing associated with  $\mathcal{A}_W$ . See [\[Katz 1981\]](#) for the construction of this pairing. We have  $e = 0$  if and only if  $\mathcal{A}_W \simeq \mathcal{A}_{\overline{s}} \times_{\overline{s}} W$ . Furthermore, the fact that

$$\text{Tor}_p(\mathcal{A}(W)) = \text{Tor}_p(A(\widehat{K}_s^{\text{sh}})) = \text{Tor}_p(A(K^{\text{unr}})) = \text{Tor}_p(A(\overline{K}_s^{\text{sh}}))$$

in our situation shows that  $e = 0$ . This follows directly from the definition of the Serre–Tate pairing in the ordinary case (see the definition of the morphism “ $p^n$ ” in [\[Katz 1981, p. 151\]](#)). Thus we have  $\mathcal{A}_W \simeq \mathcal{A}_{\overline{s}} \times_{\overline{s}} W$  and in particular  $\text{Tr}_{\overline{K}|\overline{k}}(A_{\overline{K}}) \neq 0$  by [Proposition 9.1 \(c\)](#) below. This contradicts one of our assumptions. We conclude that  $G = 0$ , so the conclusion must hold.  $\square$

**Theorem 2.9.** *Suppose that  $\text{Tor}_p(A(K^{\text{sep}}))$  is infinite. Then there is an étale  $K$ -isogeny*

$$\phi : A \rightarrow B,$$

where  $B$  is an abelian variety over  $K$  and there is an étale  $K$ -isogeny

$$\lambda : B \rightarrow B$$

such that the order of  $\ker(\lambda)$  is  $> 1$  and such that the orders of  $\ker(\lambda)$  and  $\ker(\phi)$  are powers of  $p$ .

*Proof.* Note that in [\[Rössler 2013, Theorem 1.4\]](#), this statement is proven under the supplementary assumption that there exist  $n \in \mathbb{Z}$ , such that  $(n, p) = 1$  and  $n > 3$  and such that  $A[n](\overline{K}) \simeq (\mathbb{Z}/n\mathbb{Z})^{2 \dim(A)}$ . Using [\[Zarkhin and Parshin 1989, paragraph 3, “Finiteness Theorem for Forms”, p. 69\]](#) in the proof, it can be seen that this assumption is not necessary. A completely parallel argument is described in the proof of [Proposition 2.6](#). We leave the details to the reader.  $\square$

**Theorem 2.10.** *Suppose that there exists an étale  $K$ -isogeny  $\phi : A \rightarrow A$ , such that  $\deg(\phi)$  is strictly larger than 1 and that  $\deg(\phi) = p^r$  for some  $r > 0$ . Suppose also that  $A$  is a geometrically simple abelian variety and that  $\mathcal{A}$  is a semiabelian scheme.*

*Then  $\mathcal{A}$  is an abelian scheme and  $\phi$  extends to an étale  $S$ -morphism  $\mathcal{A} \rightarrow \mathcal{A}$  of group schemes.*

*Proof.* Notice first that by a result of Raynaud [1970, IX, Corollary 1.4, p. 130], the morphism  $\phi$  extends uniquely to an  $S$ -morphism  $\bar{\phi} : \mathcal{A} \rightarrow \mathcal{A}$  of group schemes. Since  $\bar{\phi}$  is étale over  $K$ , we have an exact sequence of coherent sheaves

$$0 \rightarrow \bar{\phi}^*(\Omega_{\mathcal{A}/S}) \rightarrow \Omega_{\mathcal{A}/S}$$

on  $\mathcal{A}$ . Let  $\sigma \in H^0(\mathcal{A}, \det(\bar{\phi}^*(\Omega_{\mathcal{A}/S}))^\vee \otimes \det(\Omega_{\mathcal{A}/S}))$  be the corresponding section. Since

$$\sigma_K \in H^0(A, \det(\phi^*(\Omega_{A/K}))^\vee \otimes \det(\Omega_{A/K}))$$

has an empty zero-scheme, the zero scheme  $Z(\sigma)$  is supported on a finite number of closed fibres of  $\mathcal{A}$ . Hence there exists a finite number  $P_1, \dots, P_n$  of closed points of  $S$ , such that  $Z(\sigma) = \coprod_{i=1}^n n_i \mathcal{A}_{P_i}$  (as Weil divisors) for some  $n_i \geq 0$ . On the other hand, the Weil divisor  $Z(\sigma)$  is rationally equivalent to 0, since  $\det(\phi^*(\Omega_{\mathcal{A}/S}))^\vee \otimes \det(\Omega_{\mathcal{A}/S}) \simeq \det(\Omega_{\mathcal{A}/S})^\vee \otimes \det(\Omega_{\mathcal{A}/S}) \simeq \mathcal{O}_{\mathcal{A}}$ . Now notice that the morphism  $p^* : \text{Pic}(S) \rightarrow \text{Pic}(\mathcal{A})$  of Picard groups is injective, because it is split by the map  $\epsilon_{\mathcal{A}/S}^* : \text{Pic}(\mathcal{A}) \rightarrow \text{Pic}(S)$ . Hence the Weil divisor  $\coprod_{i=1}^n n_i P_i$  is rationally equivalent to 0 on  $S$ , which implies that  $n_i = 0$  for all  $i = 1, \dots, n$ . In other words, we have  $Z(\sigma) = \emptyset$  and thus the morphism  $\bar{\phi}^*(\Omega_{\mathcal{A}/S}) \rightarrow \Omega_{\mathcal{A}/S}$  is an isomorphism. By [Hartshorne 1977, III, Proposition 10.4], this implies that  $\bar{\phi}$  is étale.

Let now  $s \in S$  be a closed point such that  $\mathcal{A}_s$  has a presentation

$$0 \rightarrow G \xrightarrow{\iota} \mathcal{A}_s \rightarrow A_0^0 \rightarrow 0,$$

where  $G$  is a torus over  $s$  of dimension  $d > 0$  and  $A_0^0$  is an abelian variety over  $s$ . The morphism  $\bar{\phi}_s|_G : G \rightarrow \mathcal{A}_s$  factors through  $G$ , since there is no nonconstant  $s$ -morphism  $G \rightarrow A_0^0$ . Call  $\gamma : G \rightarrow G$  the resulting morphism. The morphism  $\gamma$  is étale. Indeed, we have a commutative diagram

$$\begin{array}{ccccc} \gamma^*(\iota^*(\Omega_{\mathcal{A}_s/s})) & \longrightarrow & \gamma^*(\Omega_{G/s}) & \longrightarrow & \Omega_{G/s} \\ \downarrow \sim & & & & \downarrow = \\ \iota^*(\bar{\phi}_s^*(\Omega_{\mathcal{A}_s/s})) & \longrightarrow & \iota^*(\Omega_{\mathcal{A}_s/s}) & \longrightarrow & \Omega_{G/s} \end{array}$$

and in the lower row of this diagram all the arrows are surjective. Thus the arrow

$$\gamma^*(\Omega_{G/s}) \rightarrow \Omega_{G/s}$$

must also be surjective and hence an isomorphism. Since  $G$  is smooth over  $\kappa(s)$ , we conclude that  $\gamma$  is smooth by [Hartshorne 1977, III, Proposition 10.4]. In particular  $\gamma$  is faithfully flat, because it is a morphism of group schemes and  $G$  is connected (see, e.g., [SGA 3<sub>I</sub> 2011, Exposé IV-B, Corollary 1.3.2]). Now recall that there is a  $K$ -morphism  $\psi : A \rightarrow A$  such that  $\psi \circ \phi = [p^{\deg(\phi)}]_A$  (because finite commutative group schemes over  $K$  are annihilated by their order; see [Tate and Oort 1970, Theorem (Deligne), p. 4]). The morphism  $\psi$  extends uniquely to  $\bar{\psi} : \mathcal{A} \rightarrow \mathcal{A}$  and thus by unicity, we have  $\bar{\psi} \circ \bar{\phi} = [p^{\deg(\phi)}]_{\mathcal{A}}$ . In particular,  $\ker(\gamma)$  is a closed subscheme of  $\ker([p^{\deg(\phi)}]_G)$ . Since  $\ker([p^{\deg(\phi)}]_G)$  is an infinitesimal group scheme and  $\gamma$  is étale, we see that  $\ker(\gamma) = 0$  (since  $\ker(\gamma)$  is étale over  $s$ ). Thus  $\gamma$  is an isomorphism.

Now choose a  $\bar{s}$ -isomorphism  $G_{\bar{s}} \simeq \mathbb{G}_m^d$  (here  $\bar{s}$  is the spectrum of the algebraic closure of  $\kappa(s)$ ). The morphism  $\gamma_{\bar{s}}$  is described by a matrix  $M \in \mathrm{GL}_d(\mathbb{Z})$  (because the group scheme dual to  $G_{\bar{s}}$  is the diagonalisable group scheme over  $\bar{s}$  associated with  $\mathbb{Z}^d$ ). Hence there exists a monic polynomial  $P(x) \in \mathbb{Z}[x]$ , such that  $P(0) = \pm 1$  and such that  $P(\gamma_{\bar{s}}) = 0$ .

Finally, choose a prime  $l \neq p$ . Let  $\widehat{\mathcal{O}}_s^{\mathrm{sh}}$  be the completion of the strict henselisation of the local ring of  $S$  at  $s$ . Let  $\widehat{K}_s^{\mathrm{sh}}$  be the fraction field of  $\widehat{\mathcal{O}}_s^{\mathrm{sh}}$  and let  $j \in \mathbb{N}$ . The closed subgroup scheme  $G_{\bar{s}}[l^j]$  of  $G_{\bar{s}}$  extends uniquely to a finite and étale subgroup scheme  $\widetilde{G}_{lj}$  of  $\mathcal{A}_{\widehat{\mathcal{O}}_s^{\mathrm{sh}}}$  over  $\widehat{\mathcal{O}}_s^{\mathrm{sh}}$ . See [SGA 3<sub>II</sub> 1970, Theorem 3.6 and Theorem 3.6 bis]. Furthermore the natural map  $\widetilde{G}_{lj}(\widehat{\mathcal{O}}_s^{\mathrm{sh}}) \rightarrow G_{\bar{s}}[l^j](\bar{s})$  is a bijection, since  $\widehat{\mathcal{O}}_s^{\mathrm{sh}}$  is strictly henselian and  $\widetilde{G}_{lj}$  is étale (see [Milne 1980, Proposition I.4.4]). Hence  $P(\phi)(\widetilde{G}_{lj}(\widehat{\mathcal{O}}_s^{\mathrm{sh}})) = 0$ . On the other hand, the image of the group  $\bigcup_{j \in \mathbb{N}} \widetilde{G}_{lj}(\widehat{\mathcal{O}}_s^{\mathrm{sh}})$  in  $A_{\widehat{K}_s^{\mathrm{sh}}}$  is dense, because  $A$  is geometrically simple and the group  $\bigcup_{j \in \mathbb{N}} \widehat{G}_{lj, \bar{s}}(\mathcal{O}_s^{\mathrm{sh}})$  is infinite. Hence  $P(\phi) = 0$  and since  $P(0) = \pm 1$ , we see that  $\phi$  is an automorphism, which is a contradiction.  $\square$

### 7. Proof of Theorem 1.1

**Theorem 1.1.** (a) *Suppose that  $A$  is geometrically simple. If  $A(K^{\mathrm{perf}})$  is finitely generated and of rank  $> 0$  then  $\mathrm{Tor}_p(A(K^{\mathrm{sep}}))$  is a finite group.*

(b) *Suppose that  $A$  is an ordinary (not necessarily simple) abelian variety. If  $\mathrm{Tor}_p(A(K^{\mathrm{sep}}))$  is a finite group then  $A(K^{\mathrm{perf}})$  is finitely generated.*

We shall need the following:

**Lemma 7.1.** *Let  $B$  be an abelian variety over  $K$  and let  $\gamma : B \rightarrow B$  be a  $K$ -isogeny such that  $\deg(\phi) > 1$ . Suppose that  $B$  is geometrically simple. Let  $H \subseteq A(\bar{K})$  be a finitely generated subgroup. Then the set*

$$\bigcap_{r \geq 0} \gamma^{or}(H)$$

*is a finite group.*

*Proof of Lemma 7.1.* Let  $G := \bigcap_{r \geq 0} \gamma^{or}(H)$ . Let  $F := G/\mathrm{Tor}(G)$  be the quotient of  $G$  by its torsion subgroup. We may suppose without restriction of generality that  $\mathrm{rk}(G) > 0$  for otherwise the lemma is proven. Since  $\gamma$  is a group homomorphism, we have  $\gamma(\mathrm{Tor}(G)) \subseteq \mathrm{Tor}(G)$  and thus  $\gamma$  gives rise to a group homomorphism  $F \rightarrow F$  that we also denote by  $\gamma$ . By construction, we have  $\gamma(F) = F$  and thus  $\gamma : F \rightarrow F$  is a bijection, since  $F$  is a finitely generated free  $\mathbb{Z}$ -module. Let

$$P(t) := t^n + a_{n-1}t^{n-1} + \cdots + a_1t + a_0 \in \mathbb{Z}[t]$$

be the characteristic polynomial of  $\gamma : F \rightarrow F$ . We have  $P(\gamma) = 0$  by the Cayley–Hamilton theorem and since  $\gamma$  is an automorphism, we have

$$P(0) = a_0 = \pm 1 = \det(\gamma).$$

Hence

$$(-a_0)^{-1} \cdot (\gamma^{\circ, n-1} + a_{n-1} \cdot \gamma^{\circ, n-2} + \dots a_1 \cdot \text{Id}_F)$$

is the inverse of  $\gamma : F \rightarrow F$ . Now let  $\tilde{\gamma}$  be the  $K$ -group scheme homomorphism

$$\tilde{\gamma} := (-a_0)^{-1} \cdot (\gamma^{\circ, n-1} + a_{n-1} \cdot \gamma^{\circ, n-2} + \dots a_1 \cdot \text{Id}_B)$$

from  $B$  to  $B$ . Suppose first that the morphism of  $K$ -group schemes  $\tilde{\gamma} \circ \gamma - \text{Id}_B$  is not the zero morphism. Then it is surjective, because  $B$  is simple. Furthermore the group  $G$  is dense in  $B_{\bar{K}}$ , since  $B$  is geometrically simple. Thus the group  $(\tilde{\gamma} \circ \gamma - \text{Id}_B)(G)$  is dense in  $B_{\bar{K}}$ . On the other hand, by construction  $(\tilde{\gamma} \circ \gamma - \text{Id}_B)(G) \subseteq \text{Tor}(G)$ . Since  $\text{Tor}(G)$  is a finite group, it is not dense in  $B_{\bar{K}}$  and thus we deduce that  $\tilde{\gamma} \circ \gamma - \text{Id}_B$  must be the zero morphism. Hence  $\gamma$  is invertible (with inverse  $\tilde{\gamma}$ ), which contradicts the assumption that  $\text{deg}(\gamma) > 1$ . We conclude that we cannot have  $\text{rk}(G) > 0$  and thus  $G = \text{Tor}(G)$  is a finite group.  $\square$

*Proof of Theorem 1.1.* For statement (a), suppose first that  $\text{Tor}_p(A(K^{\text{sep}}))$  is not a finite group. Then by Theorem 2.9, there exists an abelian variety  $B$  over  $K$ , which is  $K$ -isogenous to  $A$  and which carries an étale  $K$ -endomorphism  $B \rightarrow B$ , whose degree is  $> 1$  and is a power of  $p$ . The dual of  $B$  hence carries an isogeny  $\phi$ , which is purely inseparable (because the dual of a finite étale group scheme over a field is an infinitesimal group scheme) and thus we have

$$B^\vee(K^{\text{perf}}) = \bigcap_{r \geq 0} \phi^{or}(B^\vee(K^{\text{perf}}))$$

By Lemma 7.1,  $B^\vee(K^{\text{perf}})$  is thus either finite or not finitely generated and the same holds for  $A$ , since  $A$  is isogenous to  $B^\vee$ . This proves (a).

We now turn to the proof of statement (b). Note that by Grothendieck’s semiabelian reduction theorem, there is a finite and separable extension  $K_1|K$  such that  $A_{K_1}$  extends to a semiabelian scheme over the normalisation  $S_1$  of  $S$  in  $K_1$ . The scheme  $S_1$  might not be geometrically connected over  $k$  but there a finite extension  $k_1$  of  $k$ , such that the connected components of  $S_{1, k_1}$  are geometrically connected. We choose one of these connected components, say  $S_2$ . The extension of function fields corresponding to the morphism  $S_2 \rightarrow S$  is separable by construction so we may (and do) assume that  $A$  is semiabelian to begin with. Suppose that  $A(K^{\text{perf}})$  is not finitely generated and that  $A$  is ordinary. Then by Proposition 2.6, there is an abelian variety  $B$  over  $K$ , which is  $K$ -isogenous to  $A$  and which carries a  $K$ -isogeny  $B \rightarrow B$ , whose kernel is a multiplicative group scheme of order  $> 1$ . The dual  $\phi$  of this isogeny is an étale isogeny of  $B^\vee$ , which has degree  $p^r$  for some  $r > 0$ . Thus  $\text{Tor}_p(B^\vee(K^{\text{sep}}))$  is an infinite group and the same holds for  $A$ , since  $A$  is isogenous to  $B^\vee$ . This proves (b).  $\square$

### 8. Proof of Theorem 1.2

**Theorem 1.2.** *Suppose that  $\mathcal{A}$  is a semiabelian scheme and that  $A$  is a geometrically simple abelian variety over  $K$ . If  $\text{Tor}_p(A(K^{\text{sep}}))$  is infinite, then*

- (a)  $A$  is an abelian scheme;
- (b) there is  $r_A \geq 0$  such that  $p^{r_A} \cdot \text{Tor}_p(A(K^{\text{sep}})) \subseteq \text{Tor}_p(A(K^{\text{unr}}))$ .

Furthermore, there is

- (c) an abelian scheme  $\mathcal{B}$  over  $S$ ;
- (d) a generically étale  $S$ -isogeny  $\mathcal{A} \rightarrow \mathcal{B}$ , whose degree is a power of  $p$ ;
- (e) an étale  $S$ -isogeny  $\mathcal{B} \rightarrow \mathcal{B}$  whose degree is  $> 1$  and is a power of  $p$ .

Finally

- (f) if  $A$  is ordinary then the Kodaira–Spencer rank of  $A$  is not maximal;
- (g) if  $\dim(A) \leq 2$  then  $\mathrm{Tr}_{\bar{K}|\bar{k}}(A_{\bar{K}}) \neq 0$ .
- (h) for all closed points  $s \in S$ , the  $p$ -rank of  $\mathcal{A}_s$  is  $> 0$ .

*Proof.* Proof of (a): Note that by [Theorem 2.9](#), the abelian variety  $A$  is isogenous to an abelian variety  $B$  over  $K$ , which is endowed with an étale endomorphism of degree a positive power of  $p$ . Since  $A$  extends to a semiabelian scheme over  $S$  so does  $B$ . This is a consequence of a theorem of Grothendieck (see [\[Abbes 2000, 5.\]](#) for a nice presentation). Thus, by [Theorem 2.10](#) we see that  $B$  extends to an abelian scheme  $\mathcal{B}$  over  $S$ . Using the criterion of Néron–Ogg–Shafarevich (see [\[Serre and Tate 1968\]](#)), we see that  $A$  also extends to an abelian scheme over  $S$ . By the uniqueness of semiabelian models (see [\[Raynaud 1970, IX, Corollary 1.4, p. 130\]](#)), this extension must be  $\mathcal{A}$  and thus  $\mathcal{A}$  is an abelian scheme.

Proof of (b): Let  $H := \mathrm{Gal}(K^{\mathrm{sep}}|K^{\mathrm{unr}})$ . For  $i \geq 0$ , let  $G_i := A(K^{\mathrm{sep}})[p^i]$ . The group  $G_i$  is the group of  $K$ -rational points of an étale finite group scheme  $\underline{G}_i$  over  $K$ , which is naturally a closed subgroup scheme of  $A$ . Let  $A_i := A/\underline{G}_i$  and for  $i \leq j$  let  $\phi_{i,j} : A_i \rightarrow A_j$  be the natural morphism. Let  $\mathcal{A}_i$  be the connected component of the zero section of the Néron model of  $A_i$  over  $S$ . By (a) and the criterion of Néron–Ogg–Shafarevich (see [\[Serre and Tate 1968\]](#)), this is an abelian scheme. Furthermore, by [\[Raynaud 1970, IX, Corollary 1.4, p. 130\]](#) the morphisms  $\phi_{i,j}$  extend to morphisms  $\bar{\phi}_{i,j} : \mathcal{A}_i \rightarrow \mathcal{A}_j$  and we have the classical exact sequence

$$\bar{\phi}_{i,j}^*(\Omega_{\mathcal{A}_j/S}) \rightarrow \Omega_{\mathcal{A}_i/S} \rightarrow \Omega_{\bar{\phi}_{i,j}} \rightarrow 0.$$

Now the morphism  $\bar{\phi}_{i,j}^*(\Omega_{\mathcal{A}_j/S}) \rightarrow \Omega_{\mathcal{A}_i/S}$  is injective over the generic point of  $\mathcal{A}_i$ , because  $\phi_{i,j} = \bar{\phi}_{i,j,K}$  is smooth by construction. On the other hand both  $\bar{\phi}_{i,j}^*(\Omega_{\mathcal{A}_j/S})$  and  $\Omega_{\mathcal{A}_i/S}$  are locally free and thus it follows that  $\bar{\phi}_{i,j}^*(\Omega_{\mathcal{A}_j/S}) \rightarrow \Omega_{\mathcal{A}_i/S}$  is also injective. Hence we have an exact sequence

$$0 \rightarrow \bar{\phi}_{i,j}^*(\Omega_{\mathcal{A}_j/S}) \rightarrow \Omega_{\mathcal{A}_i/S} \rightarrow \Omega_{\bar{\phi}_{i,j}} \rightarrow 0. \tag{6}$$

Let  $\pi_i : \mathcal{A}_i \rightarrow S$  be the structural morphism. We have a functorial isomorphism

$$\Omega_{\mathcal{A}_i} \simeq \pi_i^*(\pi_{i,*}(\Omega_{\mathcal{A}_i/S}))$$

and thus there is a coherent sheaf  $T_{i,j}$  on  $S$ , which is a torsion sheaf, such that  $\pi_i^*(T_{i,j}) \simeq \Omega_{\bar{\phi}_{i,j}}$  and the sequence (6) is the pull-back by  $\pi_i^*$  of a sequence

$$0 \rightarrow \pi_{j,*}(\Omega_{\mathcal{A}_j/S}) \rightarrow \pi_{i,*}(\Omega_{\mathcal{A}_i/S}) \rightarrow T_{i,j} \rightarrow 0$$

and in particular

$$\deg_S(\pi_{j,*}(\Omega_{\mathcal{A}_j/S})) + \deg_S(T_{i,j}) = \deg_S(\pi_{i,*}(\Omega_{\mathcal{A}_i/S})).$$

Now recall that  $\text{deg}_S(\pi_{i,*}(\Omega_{\mathcal{A}_i/S})) \geq 0$  for all  $i \geq 0$  (see [Faltings and Chai 1990, V, Proposition 2.2, p. 164]). Thus, for  $i = 0, 1, \dots$ , the sequence  $\text{deg}_S(\pi_{i,*}(\Omega_{\mathcal{A}_i/S}))$  is a nonincreasing sequence of natural numbers. Hence for large enough  $i$ , say  $i_0$ , it reaches its minimum. We conclude that  $T_{i_0,j} = 0$  for  $j > i_0$ , so that the morphism  $\bar{\phi}_{i_0,j}$  is étale and finite. Now  $\phi_{0,i_0}(G_j(K^{\text{sep}}))$  lies by construction in the kernel of  $\phi_{i_0,j}$ . Thus

$$\phi_{0,i_0}(G_j(K^{\text{sep}})) \subseteq A_{i_0}(K^{\text{unr}})$$

when  $j > i_0$ . In other words, for any  $x \in G_j(K^{\text{sep}})$  and any  $\gamma \in H$ , we have

$$\gamma(x) - x \in G_{i_0}(K^{\text{sep}}).$$

In particular, we have

$$\gamma(p^{i_0} \cdot x) = p^{i_0} \cdot \gamma(x) = p^{i_0} \cdot x$$

In particular, since  $j > i_0$  was arbitrary, we see that

$$\gamma(p^{i_0} \cdot x) = p^{i_0} \cdot x$$

for all  $x \in \text{Tor}_p(A(K^{\text{sep}}))$  and all  $\gamma \in H$ . Setting  $r_A = i_0$  concludes the proof of (b).

Proof of the existence statements (c), (d), (e): this is a consequence of (a) and Theorems 2.9 and 2.10.

Proof of (f): this is contained in a theorem of J.-F. Voloch; see [Voloch 1995, Proposition on p. 1093].

Proof of (g): this is a consequence of (b) and Proposition 2.8.

Proof of (h): This follows from (a) and (e). □

### 9. Proof of Theorem 1.4

**9A. The trace of an abelian variety over a function field: basic facts.** Let  $E$  be an abelian over a field  $F$ . Let  $F_0 \subseteq F$  be a subfield.

The  $F|F_0$  trace  $(\text{Tr}_{F|F_0}(E), \lambda)$  (if it exists) of  $E$  over  $F_0$  is an abelian variety  $\text{Tr}_{F|F_0}(E)$  over  $F_0$  together with a homomorphism  $\lambda : \text{Tr}_{F|F_0}(E)_F \rightarrow E$  of abelian varieties over  $F$ . They have the following universal property. For any abelian  $E_0$  over  $F_0$  and a homomorphism  $\phi : E_{0,F} \rightarrow E$  of abelian varieties, there is a unique morphism  $\tilde{\phi} : E_{0,F} \rightarrow \text{Tr}_{F|F_0}(E)_F$  such that  $\phi = \lambda \circ \tilde{\phi}$ . This means that  $\text{Tr}_{F|F_0}(E)$  and  $\lambda$  are uniquely determined if they exist.

Here are some known facts about  $\text{Tr}_{F|F_0}(E)$ . Before stating them, we record the fact for any finite morphism of abelian varieties  $f : E' \rightarrow E$  over  $F$ , the natural morphism  $E'/\ker(f) \rightarrow E$  is a closed immersion. Here  $E'/\ker(f)$  is the quotient described in Proposition 4.1. To see this, consider that the morphism  $E'/\ker(f) \rightarrow E$  is by definition a monomorphism of fppf sheaves over  $F_0$  and hence a monomorphism of schemes. On the other hand, it is proper and of finite type and thus a closed immersion (see [EGA IV<sub>4</sub> 1967, p. 182] for this). We shall call  $\text{Im}(f)$  the abelian variety  $E'/\ker(f)$  viewed as an abelian subvariety of  $E$ .

The field extension  $F|F_0$  is called primary (resp. regular) if the algebraic closure of  $F_0$  in  $F$  is purely inseparable over  $F_0$  (if  $F_0$  is algebraically closed in  $F$  and  $F$  is separable over  $F_0$ ). Note that if  $F$  is the function field of a smooth and geometrically integral variety over  $F_0$  then  $F|F_0$  is regular.

**Proposition 9.1** (see [Conrad 2006, Theorems 6.4 and 6.12]). (a) *If  $F|F_0$  is primary then the  $F|F_0$  trace  $(\mathrm{Tr}_{F|F_0}(E), \lambda)$  of  $E$  over  $F_0$  exists and the kernel of  $\lambda$  is finite over  $F$ .*

(b) *If  $F|F_0$  is regular then the kernel of the morphism  $\lambda$  is connected and so is its Cartier dual.*

(c) *If  $F_1|F$  and  $F|F_0$  are primary extensions then  $(\mathrm{Tr}_{F|F_0}(E)_{F_1}, \lambda_{F_1})$  is an  $F_1|F_0$ -trace of  $E_{F_1}$ .*

(d) *We have  $\mathrm{Tr}_{F|F_0}(A/\mathrm{Im}(\lambda)) = 0$ .*

We also recall the *Lang–Néron theorem* (see [Conrad 2006, Theorem 7.1; Lang 1983, Chapter 6, Theorem 2]): if  $F|F_0$  is a finitely generated regular extension then the quotient group  $E(F)/\mathrm{Tr}_{F|F_0}(E)(F_0)$  is finitely generated. Here  $\mathrm{Tr}_{F|F_0}(E)(F_0)$  is viewed as a subgroup of  $E(F)$  via  $\lambda$  and the natural base change map from  $F_0$  to  $F$ .

**9B. The proof.** We now use the notations of [Conjecture 1.3](#).

Let  $\lambda : \mathrm{Tr}_{L|l_0}(C) \rightarrow C$  be the canonical morphism. We write  $C/\mathrm{Im}(\lambda)$  for the quotient of  $C$  by  $\mathrm{Im}(\lambda)$  in the sense of [Proposition 4.1](#).

We begin with:

**Proposition 9.2.** *If  $\mathrm{IVD}(C/\mathrm{Im}(\lambda), L) \subseteq \mathrm{Tor}^P((C/\mathrm{Im}(\lambda))(L))$  then  $\mathrm{IVD}(C, L) \subseteq \mathrm{Tor}^P(C(L))$ .*

For the proof of [Proposition 9.2](#), we shall need the following:

**Lemma 9.3.** *Let  $N$  be a finite flat infinitesimal group scheme over a field  $J$  of characteristic  $p$ . There is a finite field extension  $J'|J$  such that for any  $n \geq 0$  and any element  $\alpha \in H^1(J, N^{(p^n)})$ , the image  $\alpha_{J'}$  of  $\alpha$  in  $H^1(J', N^{(p^n)}_{J'})$  vanishes.*

Here  $H^1(J, N^{(p^n)})$  is the first cohomology group of  $N^{(p^n)}$  viewed as a sheaf in the fppf topology. More concretely, it is the group of isomorphism classes of torsors of  $N^{(p^n)}$  over  $J$ . In the following proof, we shall write  $J^{p^{-m}} \subseteq \bar{J}$  for the subfield of  $\bar{J}$  consisting of elements of the form  $x^{p^{-m}}$ , where  $x \in J$ .

*Proof of Lemma 9.3.* First suppose that  $N$  has a filtration by finite closed subgroup schemes, whose quotients are isomorphic to either  $\alpha_{p,J}$  or  $\mu_{p,J}$ . Let  $m \geq 0$  be the number of nonvanishing quotients. We shall prove by induction on  $m$  that the image of  $\alpha$  in  $H^1(J^{p^{-m}}, N^{(p^n)})$  vanishes for all  $n \geq 0$  (under the supplementary assumption on  $N$ ), for any field  $J$  of characteristic  $p$ . If  $m = 0$  the statement holds tautologically, so we shall suppose that it holds for  $1, \dots, m - 1$ . Let

$$0 \rightarrow F_1 \rightarrow N_{J_1} \rightarrow F_2 \rightarrow 0$$

be a presentation of  $N$  where  $F_2$  is isomorphic to either  $\alpha_{p,J}$  or  $\mu_{p,J}$  and  $F_1$  has a filtration as above, whose number of nonvanishing quotients is  $\leq m - 1$ . This induces exact sequences

$$\begin{aligned} 0 \rightarrow H^1(J^{p^{-1}}, (F_{1,J^{p^{-1}}})^{(p^n)}) &\rightarrow H^1(J^{p^{-1}}, (N_{J^{p^{-1}}})^{(p^n)}) \rightarrow H^1(J^{p^{-1}}, (F_{2,J^{p^{-1}}})^{(p^n)}) \\ 0 \rightarrow H^1(J^{p^{-m}}, (F_{1,J^{p^{-m}}})^{(p^n)}) &\rightarrow H^1(J^{p^{-m}}, (N_{J^{p^{-m}}})^{(p^n)}) \rightarrow H^1(J^{p^{-m}}, (F_{2,J^{p^{-m}}})^{(p^n)}) \end{aligned}$$

(observe that  $H^0(J^{p^{-m}}, (F_{2,J^{p^{-m}}})^{(p^n)}) = 0$  since  $F_2$  is infinitesimal). Since  $F_2^{(p^n)}$  is of height one, the image of  $\alpha$  in  $H^1(J^{p^{-1}}, (F_{2,J^{p^{-1}}})^{(p^n)})$  vanishes by [Milne 2006, Lemma III.3.5.7]. The element  $\alpha$  is thus

the image of an element  $\beta \in H^1(J^{p^{-1}}, (F_{1,J^{p^{-1}}})^{(p^n)})$ . By the inductive hypothesis, the image of  $\beta$  in  $H^1(J^{p^{-m}}, (F_{1,J^{p^{-m}}})^{(p^n)})$  vanishes and thus the image of  $\alpha$  in  $H^1(J^{p^{-m}}, (N_{J^{p^{-m}}})^{(p^n)})$  vanishes, proving the claim.

Now according to [Grothendieck 1974, §2.4, p. 28] there is a finite extension  $J_1$  of  $J$  such that  $N_{J_1}$  has a filtration by finite closed subgroup schemes, whose quotients are isomorphic to either  $\alpha_{p,J_1}$  or  $\mu_{p,J_1}$ . This extension will by construction also work for all the group schemes  $N^{(p^n)}$  and the number of nonvanishing quotients of all the group schemes  $N^{(p^n)}_{J_1}$  is constant, say it is  $m$ . Hence the extension  $J' := J_1^{p^{-m}}$  has the required property. □

*Proof of Proposition 9.2.* . Now suppose that  $\text{IVD}(C/\text{Im}(\lambda), L) \subseteq \text{Tor}^p((C/\text{Im}(\lambda))(L))$ . We want to show that  $\text{IVD}(C, L) \subseteq \text{Tor}^p(C(L))$ .

Write

$$\lambda^{(p^n)} : \text{Tr}_{L|l_0}(C)^{(p^n)} \rightarrow C^{(p^n)}$$

for the base change of  $\lambda$  by  $F_L^{\circ n}$ . We have an exact sequence

$$0 \rightarrow \text{Im}(\lambda)(L) \rightarrow C(L) \rightarrow (C/\text{Im}(\lambda))(L)$$

and we have  $(C/\text{Im}(\lambda))^{(p^n)} \simeq C^{(p^n)}/\text{Im}(\lambda^{(p^n)})$ . Let now

$$x_0 \in C(L), \quad x_1 \in C^{(p)}(L), \quad x_2 \in C^{(p^2)}(L), \quad \dots,$$

be a sequence of points such  $V_{C^{(p)}/L}(x_1) = x_0$ ,  $V_{C^{(p^2)}/L}(x_2) = x_1$ , etc. Then we know from the above supposition that the image of  $x_n$  in  $(C^{(p^n)}/\text{Im}(\lambda^{(p^n)}))(L)$  is a prime to  $p$  torsion point for all  $n \geq 0$ . In particular, the order  $m$  of the image of  $x_n$  in  $(C^{(p^n)}/\text{Im}(\lambda^{(p^n)}))(L)$  is independent of  $n$ , because the degree of the Verschiebung is always a power of  $p$ . Let  $m$  be the order of  $x_0$  (and hence of all the  $x_n$ ). Then  $m \cdot x_n \in \text{Im}(\lambda^{(p^n)})(L)$  for all  $n$  and thus  $m \cdot x_0$  is indefinitely Verschiebung divisible in  $\text{Im}(\lambda)(L)$  (because the Verschiebung morphism commutes with morphisms of commutative group schemes). It now suffices to prove that  $m \cdot x_0$  is of finite and prime to  $p$  order in  $\text{Im}(\lambda)(L)$ . Hence, we may and do assume that the morphism  $\lambda : \text{Tr}_{L|l_0}(C) \rightarrow C$  is a surjection.

Now  $\lambda$  is also finite and purely inseparable by [Conrad 2006, Theorem 6.12] and it is thus a bijection. We are now given infinitely many  $L$ -morphisms

$$\dots (\lambda^{(p^n)})^*(x_n) \rightarrow \dots \rightarrow (\lambda^{(p)})^*(x_1) \rightarrow \lambda^*(x_0),$$

where  $(\lambda^{(p^n)})^*(x_n)$  is the base change by  $\lambda^{(p^n)}$  of  $x_n$  viewed as a closed subscheme of  $C^{(p^n)}$ . The  $L$ -scheme  $(\lambda^{(p^n)})^*(x_n)$  is a torsor under the group scheme  $(\ker \lambda)^{(p^n)} \simeq \ker \lambda^{(p^n)}$  and according to Lemma 9.3, there is a finite extension  $L'$ , which splits all the  $(\lambda^{(p^n)})^*(x_n)$ . We thus obtain an indefinitely Verschiebung divisible point  $x'_0$  in  $\text{Tr}_{L|l_0}(C)(L')$ , whose image in  $C(L')$  is  $x_0$ . Now  $\text{Tr}_{L|l_0}(C)_{L'}$  is by definition the base change to  $L'$  of an abelian variety over  $l_0$ ; so we are reduced to showing Theorem 1.4 for abelian varieties  $C$  that arise by base-change from  $l_0$ . Lemma 9.4 below thus concludes the proof. □

**Lemma 9.4.** *We have  $\text{IVD}(C, L) \subseteq \text{Tor}^p(C(L))$  if  $C \simeq C_0 \times_{l_0} L$ , where  $C_0$  is an abelian variety over  $l_0$ .*

*Proof of Lemma 9.4.* By [Esnault and Langer 2013, Theorem 6.2 and afterwards] there is an  $m \geq 1$  so that  $m \cdot x_0 \in C_0(l_0)$ . Since  $l_0$  is algebraically closed, this implies that  $x_0 \in C_0(l_0)$ , concluding the proof.  $\square$

*Proof of Theorem 1.4.* We begin with a couple of reductions.

(1) *We may assume in the statement of Theorem 1.4 that  $L$  is the function field of a smooth and proper curve  $B$  over  $l_0$ .*

Using Proposition 9.2 and Proposition 9.1 (d), we see that when carrying out reduction (1), we may assume that  $\text{Tr}_{L|l_0}(C) = 0$ . Reduction (1) now follows from a standard spreading out argument together with Proposition C.1 in the Appendix. Here one could probably appeal instead to Hilbert’s irreducibility theorem (as in [Lang 1983, Chapter 9, Corollary 6.3]) but for lack of an adequate reference in the case of function fields, we prefer to use Proposition C.1.

(2) *We may assume in the statement of Theorem 1.4 that  $\dim(\text{Tr}_{\bar{L}|l_0}(C_{\bar{L}})) = \dim(\text{Tr}_{L|l_0}(C))$ .*

To see this, suppose for the space of this paragraph that we know that Theorem 1.4 is true in general under restrictions (1) and (2). Let  $L'|L$  be a finite extension such that  $\dim(\text{Tr}_{L'|l_0}(C_{L'}))$  is maximal among all finite extensions of  $L$ . In particular we then have  $\dim(\text{Tr}_{L'|l_0}(C_{L'})) = \dim(\text{Tr}_{\bar{L}|l_0}(C_{\bar{L}}))$ . According to Proposition 9.1 (c), we may assume that  $L'|L$  is separable. Replacing  $L'$  by the Galois closure of  $L'$  over  $L$ , we may even suppose that  $L'|L$  is Galois. Let  $y_0 \in C(L)$  be an indefinitely Verschiebung divisible element. Suppose  $y_0 \neq 0$ . Applying our assumptions to  $C_{L'}$  and to the normalisation  $B'$  of  $B$  in  $L'$ , we see that the image of  $y_0$  in  $C_{L'}(L')$  is indefinitely Verschiebung divisible. Thus for some integer  $m_{y_0}$ , which is prime to  $p$ , the element  $m_{y_0} \cdot y_0$  is divisible in the group  $C_{L'}(L')$ . Now there is a natural group homomorphism  $u : C_{L'}(L') \rightarrow C(L)$  (the trace) given by the formula

$$u(z) = \sum_{\sigma \in \text{Gal}(L'|L)} \sigma(z).$$

Hence  $m_{y_0} \cdot u(y_0) = m_{y_0} \cdot [L' : L] \cdot y_0$  is divisible in the group  $C(L)$  and hence

$$m_{y_0} \cdot [L' : L] \cdot y_0 \in \text{Tr}_{L|l_0}(C)(l_0).$$

Now if the order of the image of  $y_0$  in  $C(L)/\text{Tr}_{L|l_0}(C(l_0))$  is prime to  $p$  then we are done. Otherwise, we may (and do) replace  $y_0$  by a multiple such that the image in  $C(L)/\text{Tr}_{L|l_0}(C(l_0))$  of  $y_0$  is a nonzero element of order  $p$ . In the rest of the argument, we shall derive a contradiction from the existence of this element. Let  $i \geq 1$ . Let  $y_i \in C^{(p^i)}(L)$  be such that  $V_{C^{(p^i)}/L}^{(i)}(y_1) = y_0$ . The variety

$$(C^{(p^i)})_{L'} = (C_{L'})^{(p^i)} \cong C_{L'}^{(p^i)}$$

also has the property that  $\dim(\text{Tr}_{L'|l_0}(C_{L'}^{(p^i)})) = \dim(\text{Tr}_{\bar{L}|l_0}(C_{\bar{L}}^{(p^i)}))$  since  $C^{(p^i)}$  is isogenous to  $C$  over  $L$ . Therefore, repeating the above reasoning, there is an integer  $m_{y_i}$ , which is prime to  $p$ , such that  $m_{y_i} \cdot [L' : L] \cdot y_i \in \text{Tr}_{L|l_0}(C^{(p^i)}(L))$ . Now according to Proposition 9.1 (c), the natural morphism

$\mathrm{Tr}_{L|l_0}(C)_L^{(p^i)} \rightarrow C^{(p^i)}$  obtained by base change under  $F_C^{(i)}$  from the morphism  $\mathrm{Tr}_{L|l_0}(C)_L \rightarrow C$  makes  $\mathrm{Tr}_{L|l_0}(C)^{(p^i)}$  into the trace of  $C^{(p^i)}$ . Thus the map  $V_{C^{(p^i)}/L}^{(i)}(\bar{L})$  induces a surjective map

$$C^{(p^i)}(\bar{L})/\mathrm{Tr}_{L|l_0}(C)^{(p^i)}(l_0) \rightarrow C(\bar{L})/\mathrm{Tr}_{L|l_0}(C)(l_0)$$

and the map  $F_{C/L}^{(i)}(\bar{L})$  induces a bijective map

$$C(\bar{L})/\mathrm{Tr}_{L|l_0}(C)(l_0) \rightarrow C^{(p^i)}(\bar{L})/\mathrm{Tr}_{L|l_0}(C)^{(p^i)}(l_0).$$

Since  $V_{C^{(p^i)}/L}^{(i)}(\bar{L}) \circ F_{C/L}^{(i)}(\bar{L}) = p^i$ , we see that the order of  $y_i$  in

$$C^{(p^i)}(L)/\mathrm{Tr}_{L|l_0}(C)^{(p^i)}(l_0) \subseteq C^{(p^i)}(\bar{L})/\mathrm{Tr}_{L|l_0}(C)^{(p^i)}(l_0)$$

is  $p^{i+1}$ . This is a contradiction if  $i$  is chosen large enough so that  $p^i$  is not a divisor of  $[L' : L]$ . We conclude that the order of the image of  $y_0$  in  $C(L)/\mathrm{Tr}_{L|l_0}(C)(l_0)$  is prime to  $p$  and this concludes reduction step (2).

We now assume that we are given an abelian variety  $C$  over  $L$  and that  $C$  satisfies the assumptions of [Theorem 1.4](#) as well as (1) and (2).

Let as before  $\lambda : \mathrm{Tr}_{L|l_0}(C)_L \rightarrow C$  be the canonical morphism. According to [Proposition 9.2](#), it will be sufficient to prove that  $\mathrm{IVD}(C/\mathrm{Im}(\lambda), L) \subseteq \mathrm{Tor}^p((C/\mathrm{Im}(\lambda))(L))$ . By [Proposition 9.1](#) (d), we have  $\mathrm{Tr}_{L|l_0}(C/\mathrm{Im}(\lambda)) = 0$  and since we work under supplementary assumption (2), we even have  $\mathrm{Tr}_{\bar{L}|l_0}(C/\mathrm{Im}(\lambda)) = 0$ . Thus we may replace  $C$  by  $C/\mathrm{Im}(\lambda)$  and assume from now on that  $\mathrm{Tr}_{\bar{L}|l_0}(C) = 0$ . Finally, since we have  $\mathrm{Tr}_{\bar{L}|l_0}(C) = 0$ , we may replace without restriction of generality  $L$  by a finite extension  $L'$  and  $B$  by its normalisation  $B'$  in  $L'$ . We may thus assume that there is an integer  $m \geq 3$ , with  $(m, p) = 1$  and such that  $C[m] \simeq (\mathbb{Z}/m\mathbb{Z})^{2\dim(C)}$  and  $C^\vee[m] \simeq (\mathbb{Z}/m\mathbb{Z})^{2\dim(C^\vee)}$ .

By a theorem of Raynaud (see [\[Abbes 2000, Proposition 5.10\]](#)), the connected component of the Néron model of  $C$  will then be a semiabelian scheme. We call it  $\mathcal{C}$ .

Now suppose as in the statement of [Conjecture 1.3](#) that we are given points  $x_\ell \in C^{(p^\ell)}(L)$  and suppose that for all  $\ell \geq 1$ , we have  $V_{C^{(p^\ell)}/L}(x_\ell) = x_{\ell-1}$ . We want to show that  $x_0 \in \mathrm{Tor}^p(C(L))$ .

By [Lemma B.2](#) and the discussion preceding it we have a canonical map

$$\alpha : C^{(p)}(L) \rightarrow \mathrm{Hom}_B(\omega_{\mathcal{C}^{(p)}}, \Omega_{B/l_0}(E)) \tag{7}$$

such that  $\alpha(x) = 0$  if and only if  $x \in F_{C/L}(C(L))$ . Here  $E = E(C)$  is the reduced divisor, which is the union of the closed point  $b \in B$  such that  $\mathcal{C}_b$  is not proper over  $\kappa(b)$ . Note that we have  $E(C) = E(C^{(p)}) = E(C^{(p^2)}) = \dots$ . The map  $\alpha$  is naturally compatible with isogenies (we skip the verification) and so there is an infinite commutative diagram

$$\begin{array}{ccc} C^{(p)}(L) & \longrightarrow & \mathrm{Hom}_B(\omega_{\mathcal{C}^{(p)}}, \Omega_{B/l_0}(E)) \\ \uparrow V_{C^{(p^2)}/L} & & \uparrow V_{C^{(p^2)}/B}^* \\ C^{(p^2)}(L) & \longrightarrow & \mathrm{Hom}_B(\omega_{\mathcal{C}^{(p^2)}}, \Omega_{B/l_0}(E)) \\ \uparrow & & \uparrow \\ \vdots & \longrightarrow & \vdots \end{array} \tag{8}$$

Remember that we have

$$\omega_{\mathcal{C}(p^n)} \simeq F_B^{\circ n, *}(\omega_{\mathcal{C}}).$$

Now choose  $n_1 \geq 1$  so that

- $\omega_{\mathcal{C}(p^{n_1})}$  has a Frobenius semistable HN filtration;
- $(\omega_{\mathcal{C}(p^{n_1})})_{=0} \simeq (\omega_{\mathcal{C}(p^{n_1})})_{=0, \text{binf}} \oplus (\omega_{\mathcal{C}(p^{n_1})})_{=0, \mu}$  splits into a biinfinitesimal and a multiplicative commutative coLie-algebra (see Lemmata 4.4 and 4.7).

Note that if some  $n_1 \geq 1$  has the two above properties, than any higher  $n_1$  will as well (by definition for the first property and tautologically for the second one).

Choose  $n_2 > n_1$  so that

(I) the image of the map

$$V_{\mathcal{C}(p^{n_2})/B}^{(n_2-n_1),*} : \omega_{\mathcal{C}(p^{n_1})} \rightarrow \omega_{\mathcal{C}(p^{n_2})}$$

lies in  $(\omega_{\mathcal{C}(p^{n_2})})_{\geq 0} \simeq F_B^{\circ(n_2-n_1),*}((\omega_{\mathcal{C}(p^{n_1})})_{\geq 0})$ ;

(II) the image of the map of coLie algebras

$$V_{\mathcal{C}(p^{n_2})/B}^{(n_2-n_1),*} : (\omega_{\mathcal{C}(p^{n_1})})_{=0} \rightarrow F_B^{\circ(n_2-n_1),*}((\omega_{\mathcal{C}(p^{n_1})})_{=0}) = (\omega_{\mathcal{C}(p^{n_2})})_{=0}$$

is  $F_B^{\circ(n_2-n_1),*}((\omega_{\mathcal{C}(p^{n_1})})_{=0, \mu})$ . Note that this is possible because the biinfinitesimal part of  $(\omega_{\mathcal{C}(p^{n_1})})_{=0}$  will be sent to 0 by sufficiently many composed Verschiebung morphisms (by definition).

Note that under (I) for any  $n_3 > n_2$  the image of the map

$$V_{\mathcal{C}(p^{n_3})/B}^{(n_3-n_2),*} : (\omega_{\mathcal{C}(p^{n_2})})_{\geq 0} \rightarrow \omega_{\mathcal{C}(p^{n_3})}$$

and hence of the map

$$V_{\mathcal{C}(p^{n_3})/B}^{(n_3-n_1),*} : \omega_{\mathcal{C}(p^{n_1})} \rightarrow \omega_{\mathcal{C}(p^{n_3})}$$

automatically lies in  $(\omega_{\mathcal{C}(p^{n_3})})_{\geq 0} \simeq F_B^{\circ(n_3-n_1),*}((\omega_{\mathcal{C}(p^{n_1})})_{\geq 0})$ .

Choose  $n_3 > n_2$  so that

(III) the map

$$\omega_{\mathcal{C}(p^{n_3})} \rightarrow \Omega_{B/l_0}(E)$$

given by  $x_{n_3}$  factors through its quotient  $(F_B^{\circ n_3, *}(\omega_{\mathcal{C}}))_{\leq 0} \simeq F_B^{\circ(n_3-n_1),*}((\omega_{\mathcal{C}(p^{n_1})})_{\leq 0})$ ;

(IV) the image of the map

$$V_{\mathcal{C}(p^{n_3})/B}^{(n_3-n_2),*} : F_B^{\circ(n_2-n_1),*}((\omega_{\mathcal{C}(p^{n_1})})_{=0}) \rightarrow F_B^{\circ(n_3-n_2),*}((\omega_{\mathcal{C}(p^{n_2})})_{=0})$$

is  $F_B^{\circ(n_3-n_2),*}((\omega_{\mathcal{C}(p^{n_2})})_{=0, \mu}) \simeq F_B^{\circ(n_3-n_1),*}((\omega_{\mathcal{C}(p^{n_1})})_{=0, \mu})$ .

Now we shall exploit the compatibility between the morphism

$$\omega_{\mathcal{C}(p^{n_1})} \xrightarrow{c(x_{n_1})} \Omega_{B/k}(E)$$

induced by  $x_{n_1}$  and the morphism

$$\omega_{\mathcal{C}(p^{n_3})} \xrightarrow{c(x_{n_3})} \Omega_{B/k}(E)$$

induced by  $x_{n_3}$ . According to the diagram (8), this compatibility gives the equality

$$c(x_{n_3}) \circ V_{\mathcal{C}(p^{n_3-n_1})/B}^* = c(x_{n_1}).$$

In other words the composition of morphisms

$$\omega_{\mathcal{C}(p^{n_1})} \xrightarrow{V_{\mathcal{C}(p^{n_3-n_1})/B}^*} \omega_{\mathcal{C}(p^{n_3})} \xrightarrow{c(x_{n_3})} \Omega_{B/k}(E)$$

is  $c(x_{n_1})$ . Furthermore, in view of (I) and (III) the map  $c(x_{n_1})$  factors as

$$\omega_{\mathcal{C}(p^{n_1})} \xrightarrow{V_{\mathcal{C}(p^{n_3})/B}^{(n_3-n_1),*}} F_B^{\circ(n_3-n_1),*}((\omega_{\mathcal{C}(p^{n_1})})_{\geq 0}) \rightarrow F_B^{\circ(n_3-n_1),*}((\omega_{\mathcal{C}(p^{n_1})})_{=0}) \rightarrow F_B^{\circ(n_3-n_1),*}((\omega_{\mathcal{C}(p^{n_1})})_{\leq 0}) \rightarrow \Omega_{B/k}(E)$$

and by (I) the map

$$\omega_{\mathcal{C}(p^{n_1})} \xrightarrow{V_{\mathcal{C}(p^{n_3})/B}^{(n_3-n_1),*}} F_B^{\circ(n_3-n_1),*}((\omega_{\mathcal{C}(p^{n_1})})_{=0})$$

factors as

$$\omega_{\mathcal{C}(p^{n_1})} \xrightarrow{V_{\mathcal{C}(p^{n_1})/B}^{(n_2-n_1),*}} F_B^{\circ(n_2-n_1),*}((\omega_{\mathcal{C}(p^{n_1})})_{\geq 0}) \rightarrow F_B^{\circ(n_2-n_1),*}((\omega_{\mathcal{C}(p^{n_1})})_{=0}) \xrightarrow{V_{\mathcal{C}(p^{n_1})/B}^{(n_3-n_2),*}} F_B^{\circ(n_3-n_1),*}((\omega_{\mathcal{C}(p^{n_1})})_{=0})$$

and thus by (IV) and (II) the image of this last map is precisely  $F_B^{\circ(n_3-n_1),*}((\omega_{\mathcal{C}(p^{n_1})})_{=0,\mu})$ .


We have thus constructed a multiplicative quotient of the  $p$ -coLie algebra  $\omega_{\mathcal{C}(p^{n_1})}$ . On the other hand the  $p$ -coLie algebra  $\omega_{\mathcal{C}(p^{n_1})}$  is the  $p$ -coLie algebra of the finite flat group scheme  $\ker F_{\mathcal{C}(p^{n_1})/B}$ . By the equivalence of categories recalled in Section 4B, this quotient corresponds to a multiplicative subgroup scheme of  $\ker F_{\mathcal{C}(p^{n_1})/B}$ . By Lemma 4.8, this subgroup scheme embeds in the canonical largest multiplicative subgroup scheme  $(\ker F_{\mathcal{C}(p^{n_1})/B})_{\mu}$  of  $\ker F_{\mathcal{C}(p^{n_1})/B}$  (in fact, it coincides with it, but we shall not need this). Finally note that

$$(\ker F_{\mathcal{C}(p^{n_1})/B})_{\mu} \simeq ((\ker F_{C/B})_{\mu})^{(p^{n_1})},$$

by the last part of Lemma 4.8.

Let  $G := (\ker F_{C/B})_{\mu}$ . Note that  $G = G_C$  in the notation of Theorem 2.1. Now consider the quotient  $\mathcal{C}_1 := \mathcal{C}/G$  (which is a semiabelian scheme by 4.10) and let  $\psi_1 : \mathcal{C} \rightarrow \mathcal{C}_1$  be the quotient morphism. The

point  $x_{n_1}$  and its image  $y_{n_1}$  in  $\mathcal{C}_1(L)$  give a commutative diagram



where the left column is an exact sequence and  $c(y_{n_1})$  is the morphism induced by  $y_{n_1}$ .

Thus  $c(y_{n_1})$  vanishes. In particular,  $y_{n_1}$  lies in the image of  $F_{\mathcal{C}_1^{(p^{n_1-1})}/B}(\mathcal{C}_1^{(p^{n_1-1})}(L))$ . Using the fact that

$$[p]_{\mathcal{C}_1^{(p^{n_1-1})}} = V_{\mathcal{C}_1^{(p^{n_1})}/B} \circ F_{\mathcal{C}_1^{(p^{n_1-1})}/B},$$

we conclude that  $y_{n_1-1}$  has a  $p$ -th root in  $\mathcal{C}_1^{(p^{n_1-1})}(L)$ . Hence  $y_0$  also has a  $p$ -th root in  $\mathcal{C}_1(L)$ . Now since  $G$  is independent of  $x_0$ , we conclude that the image of any indefinitely Verschiebung divisible point of  $\mathcal{C}(L)$  in  $\mathcal{C}_1(L)$  has a  $p$ -th root. Since  $G$  is compatible with twists, we also see that for any  $n \geq 0$  the image of any indefinitely Verschiebung divisible point of  $\mathcal{C}^{(p^n)}(L)$  in  $\mathcal{C}_1^{(p^n)}(L)$  has a  $p$ -th root. From this, by an elementary combinatorial consideration, we see that the image of any indefinitely Verschiebung divisible point of  $\mathcal{C}(L)$  in  $\mathcal{C}_1(L)$  has a  $p$ -th root, which is indefinitely Verschiebung divisible.

By the discussion above, the image of  $\text{IVD}(\mathcal{C})$  in  $\mathcal{C}_1(L)$  lies in  $p \cdot \text{IVD}(\mathcal{C}_{1,L})$ . This is the crucial fact that the rest of the proof will exploit.

Let  $\mathcal{C}_1 := \mathcal{C}/G_{\mathcal{C}}, \mathcal{C}_2/G_{\mathcal{C}_1}, \dots$  be the sequence of smooth commutative group schemes obtained by successively quotienting by the canonical subgroup schemes described in Theorem 2.1. Note that all the  $\mathcal{C}_i$  are semiabelian by Lemma 4.10. We shall denote by  $\psi_i$  the morphism  $\mathcal{C} \rightarrow \mathcal{C}_i$  obtained by composition. We write  $\mathcal{C}_i := \mathcal{C}_{i,L}$  for convenience.

Let  $m_{00}$  be an integer such that  $m_{00} \cdot x_0 =: v_0$  extends to an element  $\tilde{v}_0$  of  $\mathcal{C}(B)$ .

Now let  $D_0$  be a line bundle on  $\mathcal{C}$ . We suppose that  $[-1]_{\mathcal{C}}^*(D_0) \simeq D_0$  (i.e.,  $D_0$  is symmetric), where  $[-1]_{\mathcal{C}}$  is the inversion morphism given by the group scheme structure of  $\mathcal{C}$  over  $B$ . We also suppose that  $D_0$  is a relatively ample line bundle. If  $x \in \mathcal{C}(B)$ , write  $\tau_x : \mathcal{C} \rightarrow \mathcal{C}$  for the translation by  $x$  morphism. We use the same notation for  $x \in \mathcal{C}(L)$ .

Now consider the isogeny  $\phi_{D_0} : \mathcal{C} \rightarrow \mathcal{C}^{\vee}$  from  $\mathcal{C}$  to its dual abelian variety, which is induced by  $D_0$  (this is the polarisation induced by  $D_0$ ). Since  $v_0 \in \text{IVD}(\mathcal{C})$ , we also have  $\phi_{D_0}(v_0) \in \text{IVD}(\mathcal{C}^{\vee})$ , since

relative Frobenius morphisms are naturally compatible with morphisms of abelian varieties. The point  $\phi_{D_0}(v_0)$  corresponds to the line bundle

$$M = \tau_{v_0}^*(D_0) \otimes D_0^\vee$$

on  $C$  (see [Mumford 1970, III.13]). Since the morphism dual to the Verschiebung morphism is the relative Frobenius morphism (this is very often the definition of the Verschiebung), we see that the fact that  $\phi_{D_0}(v_0) \in \text{IVD}(C^\vee)$  translates to the fact that there exist line bundles  $M_i$  on  $C^{(p^i)}$  for all  $i \geq 1$ , such that

$$F_{C/L}^*(M_1) \simeq M, \quad F_{C^{(p)}/L}^*(M_2) \simeq M_1, \quad F_{C^{(p^2)}/L}^*(M_3) \simeq M_2, \quad \dots$$

Since  $\psi_i$  factors by construction through  $F_{C^{(p^{i-1})}/L} \circ F_{C^{(p^{i-2})}/L} \circ \dots \circ F_{C/L}$ , we see that for each  $i \geq 1$ , there is a line bundle  $J_i$  on  $C_i$  such that  $\psi_{i,L}^*(J_i) \simeq M$ .

Now recall that  $D_0$  extends uniquely (up to isomorphism) to a line bundle  $\mathcal{D}_0$  on  $\mathcal{C}$ , if we require  $D_0$  to be trivial along the unit section of  $\mathcal{C}$  (see [Moret-Bailly 1985, Proposition 2.6, p. 21]). Similarly the line bundle  $M$  extends uniquely (up to isomorphism) to a line bundle  $\mathcal{M}$  on  $\mathcal{C}$  with the same property. We shall write  $\mathcal{J}_i$  for the line bundle similarly associated with  $J_i$  on  $C_i$ . Notice that by unicity, we have  $\psi_i^*(\mathcal{J}_i) \simeq \mathcal{M}$ .

We shall now make a height computation. We shall need:

**Lemma 9.5.** *Let  $\mathcal{W}$  be a line bundle on  $\mathcal{C}$ , which is trivial when restricted to the unit section and such that  $\mathcal{W}_L$  is algebraically equivalent to 0. Let  $x \in C(B)$ . Then  $\text{deg}(x^*(\mathcal{W}))$  is the Néron–Tate height pairing of  $x_L \in C(L)$  and  $\mathcal{W}_L$ .*

*Proof.* This follows from [Moret-Bailly 1985, III.3.2 and 3.3] and the definition of polarisations. □

**Proposition 9.6.** (a) *There exists a constant  $m_0 \in \mathbb{N}^*$  and an infinite set  $I_0 \subseteq \mathbb{N}^*$  such that for any  $i \in I_0$  and any  $P \in C_i(L)$ , the element  $m_0 \cdot P$  extends to an element of  $C_i(B)$ .*

(b) *There is a constant  $c_0 \in \mathbb{N}^*$  and an infinite set  $I_0 \subseteq \mathbb{N}^*$  such that for any  $i \in I_0$  and any  $P \in \text{Tor}(C_i(L))$  we have  $c_0 \cdot P = 0$ .*

We shall prove this proposition later, using Proposition A.2 in the Appendix.

Let  $i \in I_0$ . For the next computation, recall that  $\psi_{i,L}(v_0)$  is divisible by  $p^i$  in  $C_i(L)$ . Let  $z_i$  be an element of  $C_i(L)$  such that  $p^i \cdot z_i = \psi_{i,L}(v_0)$ . According to Proposition 9.6 (a),  $m_0 \cdot z_i$  extends to an element  $u_i$  of  $C_i(B)$ . By construction, we have  $p^i \cdot u_i = m_0 \cdot \psi_i(\tilde{v}_0)$ . We compute

$$\begin{aligned} \text{deg}([m_0](\tilde{v}_0)^*(\mathcal{M})) &= \text{deg}([m_0](\tilde{v}_0)^*(\psi_i^*(J_i))) = \text{deg}([m_0](\psi_i(\tilde{v}_0))^*(J_i)) \\ &= \text{deg}([p^i](u_i)^*(J_i)) = \text{deg}(u_i^*([p^i]^*(J_i))) \\ &= \text{deg}(u_i^*(J_i^{\otimes p^i})) = p^i \cdot \text{deg}(u_i^*(J_i)). \end{aligned}$$

Here  $[m_0]$  refers to the multiplication by  $m_0$  morphism (in particular  $[m_0](\tilde{v}_0) = m_0 \cdot \tilde{v}_0$ ). Suppose for contradiction that  $\text{deg}([m_0](\tilde{v}_0)^*(\mathcal{M})) \neq 0$ . If we choose  $i$  large enough so that  $p^i$  is not a divisor of  $\text{deg}([m_0](\tilde{v}_0)^*(\mathcal{M}))$  then we get a contradiction. Thus  $\text{deg}([m_0](\tilde{v}_0)^*(\mathcal{M})) = 0$ . We may also compute

$$\text{deg}([m_0](\tilde{v}_0)^*(\mathcal{M})) = \text{deg}(\tilde{v}_0^*([m_0]^*(\mathcal{M}))) = \text{deg}(\tilde{v}_0^*(\mathcal{M}^{\otimes m_0})) = m_0 \cdot \text{deg}(\tilde{v}_0^*(\mathcal{M})).$$

In particular, by [Lemma 9.5](#), the Néron–Tate height pairing of  $v_0$  and  $M$  vanishes. Now notice that  $M$  is by definition the image of  $v_0$  under the polarisation induced by the symmetric ample line bundle  $D_0$ . Hence the Néron–Tate pairing of  $v_0$  and  $M$  is twice the Néron–Tate height of  $v_0$  with respect to the polarisation induced by  $D_0$ . In particular, the Néron–Tate height of  $v_0$  with respect to  $D_0$  vanishes. By a theorem of Lang (see [\[Conrad 2006, Theorem 9.15\]](#)) we conclude that the image of  $v_0$  in  $C(L)$  is an element of finite order. Thus the image of  $x_0$  in  $C(L)$  is also an element of finite order.

Now we show that  $x_0 \in \text{Tor}^p(C(L))$ . For contradiction, suppose that  $x_0 \notin \text{Tor}^p(C(L))$ . We thus may (and do) replace  $x_0$  by one of its multiples and suppose that  $p \cdot x_0 = 0$  and  $x_0 \neq 0$ . We know that  $\psi_{i,L}(x_0)$  is divisible by  $p^i$  in  $C_i(L)$  and since  $\psi_{i,L}$  is injective we conclude that there is an element of order  $p^{i+1}$  in  $C_i(L)$  for all  $i \geq 1$ . This contradicts [Proposition 9.6](#) (b) so we are done.  $\square$

*Proof of Proposition 9.6.* We need some preliminaries on moduli spaces of abelian varieties. Let  $n \geq 3$  with  $(n, p) = 1$  and  $g \geq 1$ . We shall choose particular values for  $g$  and  $n$  later.

Let  $A_{g,n}$  be the functor from the category of locally noetherian  $\mathbb{F}_p$ -schemes to the category of sets, such that

$$A_{g,n}(B) = \left\{ \begin{array}{l} \text{isomorphism classes of the following objects: principally polarised abelian schemes} \\ \text{over } B \text{ endowed with a symplectic isomorphism } (\mathbb{Z}/n\mathbb{Z})_B^{2g} \simeq \mathcal{A}[n] \end{array} \right\}$$

D. Mumford proved (see [\[Mumford et al. 1994\]](#)) that the functor  $A_{g,n}$  is representable by a scheme, which is separated and of finite type over  $\mathbb{F}_p$ . We shall also denote this scheme by  $A_{g,n}$ .

Furthermore, C. Chai and G. Faltings [\[1990, V, 2., Theorem 2.5\]](#) proved that there exists

- a scheme  $\bar{A}_{g,n}$  (resp.  $A_{g,n}^*$ ), which is proper over  $\mathbb{F}_p$ ;
- an open immersion  $A_{g,n} \hookrightarrow \bar{A}_{g,n}$  (resp. an open immersion  $A_{g,n} \hookrightarrow A_{g,n}^*$ );
- a semiabelian scheme  $\mathcal{U}$  over  $\bar{A}_{g,n}$ , such that  $\mathcal{U}_{A_{g,n}}$  is isomorphic to the universal abelian scheme over  $A_{g,n}$ ;
- a morphism  $\bar{\pi} : \bar{A}_{g,n} \rightarrow A_{g,n}^*$  compatible with the above open immersions of  $A_{g,n}$ ;
- a line bundle  $\omega^0$  on  $A_{g,n}^*$ , which is ample and such that  $\bar{\pi}^*(\omega^0) = \omega_{\mathcal{U}/\bar{A}_{g,n}}$ .

Write  $Z := B \times_{l_0} A_{g,n,l_0}^*$ . Recall that the Hilbert scheme  $\text{Hilb}(Z/l_0)$  is a scheme, representing the functor

$$T \mapsto \{ \text{closed subschemes of } Z_T, \text{ which are proper and flat over } T \}$$

from the category of locally noetherian scheme  $T$  over  $l_0$  to the category of sets. It is locally of finite type over  $l_0$  (see [\[Grothendieck 1966\]](#)).

Let  $\Phi \in \mathbb{Q}[\lambda]$  be a polynomial with rational coefficients and  $L_0/Z$  an ample line bundle. By definition, the  $l_0$ -scheme  $\text{Hilb}_\Phi(Z/l_0)$  represents the functor

$$T \mapsto \left\{ \begin{array}{l} \text{closed subschemes } W \text{ of } Z_T, \text{ which are proper and flat over } T \\ \text{and such that } \chi(W_t, L_{0,W_t}^{\otimes \lambda}) = \Phi(\lambda) \text{ for all } \lambda \in \mathbb{N} \text{ and all } t \in T \end{array} \right\}$$

from the category of locally noetherian scheme  $T$  over  $l_0$  to the category of sets.

Here  $W_t$  is the fibre at  $t \in T$  of the morphism  $W \rightarrow T$  and  $L_{0,W_t}$  is the pull-back of  $L$  to  $W_t$  by the natural morphism  $W_t \rightarrow Z$ . The symbol  $\chi(\cdot)$  refers to the Euler characteristic. By definition

$$\chi(W_t, L_{W_t}^{\otimes \lambda}) = \sum_{r \geq 0} (-1)^r \dim_{\kappa(t)} H^r(W_t, L_{W_t}^{\otimes \lambda}).$$

(this is called the Hilbert polynomial of  $W_t$  with respect to  $L_{W_t}$ ). It is shown in [Grothendieck 1966], that  $\text{Hilb}_\Phi(Z/l_0)$  is projective over  $l_0$  (as a consequence of the projectivity of  $Z$ ). Notice that by construction, we have a disjoint union

$$\text{Hilb}(Z/l_0) = \coprod_{\Phi \in \mathbb{Q}[\lambda]} \text{Hilb}_\Phi(Z/l_0)$$

Finally, it is shown in [Fantechi et al. 2005, part II, 5.23] that the functor  $\text{Mor}_{l_0}(B, A_{g,n}^*)$  from locally noetherian  $l_0$ -schemes  $T$  to the category of sets, such that

$$\text{Mor}_{l_0}(B, A_{g,n,l_0}^*)(T) = \{T\text{-morphisms from } B_T \text{ to } A_{g,n,T}^*\}$$

is representable by an open subscheme of  $\text{Hilb}(Z/l_0)$ . More precisely, the natural transformation of functors

$$T\text{-morphism } f \text{ from } B_T \text{ to } A_{g,n,T}^* \mapsto \text{graph of } f$$

is represented by an open immersion

$$\text{Mor}_{l_0}(B, A_{g,n}^*) \hookrightarrow \text{Hilb}(B \times_{l_0} A_{g,n,l_0}^*/l_0).$$

Let now  $D$  be an ample line bundle on  $B$ . We choose  $L_0$  to be the line bundle  $D \boxtimes \omega_{l_0}^0$  on  $Z = B \times_{l_0} A_{g,n,l_0}^*$ .

Recall that the Hodge bundles of the  $C_i$  all have the same degree by Lemma 4.12. Let  $d_0 := \deg(\omega_{C/B})$  be this common degree. Our aim is to use this to show that all the  $C_i$  embed in a bounded family of abelian varieties and apply Proposition A.2.

Notice that  $C_i[m] \simeq (\mathbb{Z}/m\mathbb{Z})^{2 \dim(C_i)}$  and  $C_i^\vee[m] \simeq (\mathbb{Z}/m\mathbb{Z})^{2 \dim(C_i^\vee)}$ . Indeed, since  $\psi_{i,L}$  is purely inseparable, it induces an isomorphism  $C[m] \rightarrow C_i[m]$  and thus  $C_i[m] \simeq (\mathbb{Z}/m\mathbb{Z})^{2 \dim(C_i)}$  by (IV) above. For the isomorphism  $C_i^\vee[m] \simeq (\mathbb{Z}/m\mathbb{Z})^{2 \dim(C_i^\vee)}$ , notice that the dual morphism  $\psi_{i,L}^\vee : C_i^\vee \rightarrow C^\vee$  is separable (because its kernel is the Cartier dual of a multiplicative group scheme) and of order a power of  $p$ . Hence, since  $(p, m) = 1$  it also induces an isomorphism  $C_i^\vee[m] \rightarrow C^\vee[m]$  (we leave the details to the reader).

Now let  $E_i := (C_i \times_L C_i^\vee)^4$ . By Zarhin’s trick (see [Moret-Bailly 1985, IX.1.1])  $E_i$  carries a principal polarisation. Furthermore, by the last paragraph, we also have  $E_i[m] \simeq (\mathbb{Z}/m\mathbb{Z})^{2 \dim(E_i)}$ . Notice also that the identity component of the Néron model of  $C_i$  is semiabelian, since  $C_i$  is semiabelian. Hence the identity component of the Néron model of  $C_i^\vee$  is also semiabelian, since  $C_i^\vee$  is isogenous to  $C_i$  (see [Abbes 2000, Proposition 5.8 (4)] for a neat presentation). Since the formation of the Néron model is compatible with products, we conclude that the identity component  $\mathcal{E}_i$  of the Néron model of  $E_i$  is also semiabelian. We also see  $\mathcal{E}_i|_{B \setminus E(C)}$  is an abelian scheme over  $B \setminus E(C)$  (where  $E(C)$  is as in (7)). Finally, we have  $\deg(\omega_{\mathcal{E}_i/B}) = 8 \cdot d_0$  by [Faltings and Chai 1990, V.3, Lemma 3.4, p. 166].

Let now  $g = \dim(E_i) = 8 \cdot \dim(C)$  and  $n = m$ . By definition,  $E_i$  is associated with an  $l_0$ -morphism  $\text{Spec } L \rightarrow \mathbf{A}_{g,n,l_0}^*$ . By the valuative criterion of properness, this morphism extends to a morphism  $\phi_i : B \rightarrow \mathbf{A}_{g,n,l_0}^*$  (resp. to a morphism  $\bar{\phi}_i : B \rightarrow \bar{\mathbf{A}}_{g,n,l_0}$ ). By unicity, we have  $\bar{\pi} \circ \bar{\phi}_i = \phi_i$ . Thus, since semiabelian extensions are unique (see [Raynaud 1970, IX, Corollary 1.4, p. 130]), we have  $\phi_i^*(\omega_{l_0}^0) \simeq \omega_{\mathcal{E}_i/B}$ . The morphism  $\phi_i$  is by definition associated with an element of  $\text{Mor}_{l_0}(B, \mathbf{A}_{g,n,l_0}^*)(l_0)$ . We can now compute the Hilbert polynomial of the graph  $\Gamma_{\phi_i}$  of  $\phi_i$  with respect to the line bundle  $L_0$ :

$$\begin{aligned} \chi(\Gamma_{\phi_i}, L_0^{\otimes \lambda}) &=: Q(\lambda) = \chi(B, (D \otimes \phi_i^*(\omega_{l_0}^0))^{\otimes \lambda}) \\ &= \deg_B((D \otimes \phi_i^*(\omega_{l_0}^0))^{\otimes \lambda}) + 1 - g(B) \\ &= \lambda \cdot \deg_B(D \otimes \phi_i^*(\omega_{l_0}^0)) + 1 - g(B) \\ &= \lambda \cdot \deg_B(D \otimes \omega_{\mathcal{E}_i/B}) + 1 - g(B) \\ &= \lambda \cdot \deg_B(D) + \lambda \cdot \deg_B(\omega_{\mathcal{E}_i/B}) + 1 - g(B) \\ &= \lambda \cdot \deg_B(D) + \lambda \cdot 8 \cdot d_0 + 1 - g(B). \end{aligned} \tag{9}$$

Here  $g(B)$  is the genus of  $B$ . The second equality is justified by the Riemann–Roch theorem on  $B$ . We thus see that the Hilbert polynomial  $Q(\lambda)$  of the graph of  $\phi_i$  with respect to  $L_0$  is  $Q(\lambda)$  is independent of  $i$ . Thus the element of  $\text{Mor}_{l_0}(B, \mathbf{A}_{g,n,l_0}^*)(l_0)$  corresponding to  $\mathcal{E}_i$  lies in the scheme

$$\text{Mor}_{l_0}(B, \mathbf{A}_{g,n,l_0}^*)(l_0) \cap \text{Hilb}_{Q(\lambda)}(B \times_{l_0} \mathbf{A}_{g,n,l_0}^*/l_0)$$

which is of finite type over  $l_0$  by the above discussion. We now let  $Y$  be the Zariski closure in

$$\text{Mor}_{l_0}(B, \mathbf{A}_{g,n,l_0}^*)(l_0) \cap \text{Hilb}_{Q(\lambda)}(B \times_{l_0} \mathbf{A}_{g,n,l_0}^*/l_0)$$

of the set all the elements of  $(\text{Mor}_{l_0}(B, \mathbf{A}_{g,n,l_0}^*)(l_0) \cap \text{Hilb}_{Q(\lambda)}(B \times_{l_0} \mathbf{A}_{g,n,l_0}^*/l_0))(l_0)$  which correspond to some  $\phi_i$  ( $i \geq 0$ ). Finally we let  $H_{00}$  be some irreducible component of  $Y$ , which meets infinitely many such points. Let  $\eta_{00} := \kappa(H_{00})$ . By construction, we have an  $H_{00}$ -morphism

$$B \times_{l_0} H_{00} \rightarrow \mathbf{A}_{g,n,H_{00}}^*$$

which sends  $(B \setminus E(C))_{\eta_{00}}$  into  $\mathbf{A}_{g,n,\eta_{00}} \subseteq \mathbf{A}_{g,n,\eta_{00}}^*$  (because by construction,  $(B \setminus E(C))_x$  is sent into  $\mathbf{A}_{g,n,x}$  for a dense sent of points  $x \in H_{00}$ ). Let

$$\gamma_0 : B_{\eta_{00}} \rightarrow \mathbf{A}_{g,n,\eta_{00}}^*$$

be the induced morphism over  $\eta_{00}$ . Now recall that there is a proper morphism  $\bar{\pi} : \bar{\mathbf{A}}_{g,n} \rightarrow \mathbf{A}_{g,n}^*$ . By the valuative criterion of properness, there is a unique  $\eta_{00}$ -morphism  $\gamma : B_{\eta_{00}} \rightarrow \bar{\mathbf{A}}_{g,n,\eta_{00}}$  such that  $\bar{\pi}_{\eta_{00}} \circ \gamma = \gamma_0$ . The morphism  $\gamma$  extends over an open subset  $H_0$  of  $H_{00}$ , yielding an  $H_0$ -morphism

$$\tilde{\gamma} : B \times_{l_0} H_0 \rightarrow \bar{\mathbf{A}}_{g,n,H_0}.$$

Replacing  $H_0$  by one of its open subsets, we may suppose that  $H_0$  is normal. Let now  $\mathcal{B}_0$  be the base change of  $\mathcal{U}$  by  $\tilde{\gamma}$ . A theorem of Moret-Bailly [1985, VI.3.1] together with a result of Raynaud [1970, XI.1.4] then shows that  $\mathcal{B}_0$  can be endowed with a relatively ample line bundle, which is symmetric and

trivial along the zero section. Let also  $t_0 := l_0$ ,  $C := B \times_{l_0} H_0$ . If we now apply [Proposition A.2](#) (a) with this choice of  $H_0$ ,  $t_0$ ,  $C$  and  $\mathcal{B}_0$ , we reach the conclusion that there is an infinite set  $I_0 \subseteq \mathbb{N}^*$  and a constant  $n_0$ , such that for  $i \in I_0$ , and any  $P \in E_i(L)$ , the element  $n_0 \cdot P$  extends to an element of  $\mathcal{E}_i(L)$ . Since  $\mathcal{C}_i$  is a direct factor of  $\mathcal{E}_i$ , we may replace  $E_i$  (resp.  $\mathcal{E}_i$ ) by  $\mathcal{C}_i$  (resp.  $\mathcal{C}_i$ ) in the last sentence. This proves (a), with  $m_0 = n_0$ . For (b), note that  $\text{Tr}_{L|l_0}(E_i) = 0$  (since  $E_i$  is a product of abelian varieties isogenous to  $C$ ) and apply [Proposition A.2](#) (b) to the same situation.  $\square$

### Appendix A: Rational points in families

The terminology of this appendix is independent of the terminology of the rest of the article and appendices.

Let  $t_0$  be an algebraically closed field. Let  $H_0$  be an integral scheme of finite type over  $t_0$ . Let  $\pi : C \rightarrow H_0$  be a smooth curve over  $H_0$ , with geometrically connected fibres. Let  $\mathcal{B}_0$  be a semiabelian scheme over  $C$ . Suppose that there exists a line bundle  $L$  on  $\mathcal{B}_0$ , which is ample relatively to  $C$ , symmetric and trivial along the zero section. Let  $\eta_0 := \kappa(H_0)$  and let  $\lambda_0 := \kappa(C)$ . Note that  $\lambda_0$  lies over  $\eta_0$  via  $\pi$  and that  $\lambda_0$  is also the generic point of  $C_{\eta_0}$  viewed as a subset of  $C$ . We suppose that  $\mathcal{B}_{0,\lambda_0}$  is an abelian variety over  $\lambda_0$ .

In the next proposition, we shall need the following lemma, which is well known from the theory of minimal models of curves.

**Lemma A.1.** *Let  $\phi : X \rightarrow Y$  be a morphism of smooth varieties over  $t_0$ . Suppose also that there is a dense open set  $Y_1 \subseteq Y$ , such that  $\phi|_{Y_1} : \phi^{-1}(Y_1) \rightarrow Y_1$  is smooth. Denote by  $X^{\text{sm}}$  the maximal open subscheme of  $X$ , such that  $\phi|_{X^{\text{sm}}} \rightarrow Y$  is smooth.*

*Let  $\sigma \in X(Y)$  be a section of  $\phi$ . Then  $\sigma \in X^{\text{sm}}(Y) \subseteq X(Y)$ .*

*Proof.* See [[Liu 2002](#), Example 4.3.25].  $\square$

**Proposition A.2.** (a) *There is a natural number  $n_0$  and a dense open set  $V \subseteq H_0$  with the following properties. For any  $x \in V(t_0)$ ,  $\mathcal{B}_{0,\kappa(C_x)}$  is an abelian variety and for any  $P_x \in \mathcal{B}_0(\kappa(C_x))$ , the point  $n_0 \cdot P_x \in \mathcal{B}_0(\kappa(C_x))$  extends to an element of  $\mathcal{N}(\mathcal{B}_{0,\kappa(C_x)})^0(C_x)$ .*

(b) *Suppose that  $C$  is proper over  $H_0$ . Suppose that there is a set  $T_0 \subseteq H_0(t_0)$ , which is dense in  $H_0$  and such that for any  $x \in T_0$  we have  $\text{Tr}_{\kappa(C_x)|t_0}(\mathcal{B}_{0,\kappa(C_x)}) = 0$ . Then there is a dense open set  $V \subseteq H_0$  and a natural number  $b_0$  such that for all  $x \in V(t_0)$ , we have  $\#\text{Tor}(\mathcal{B}_0(\kappa(C_x))) \leq b_0$ .*

Here  $\mathcal{N}(\mathcal{B}_{0,\kappa(C_x)})^0$  is the connected component of the identity of the Néron model  $\mathcal{N}(\mathcal{B}_{0,\kappa(C_x)})$  of  $\mathcal{B}_{0,\kappa(C_x)}$  over  $C_x$ .

*Proof.* We start with (a). We shall write  $\bar{\eta}_0$  for an algebraic closure of  $\eta_0$ . Consider the semiabelian scheme  $\mathcal{B}_{0,\bar{\eta}_0}$  over  $C_{\bar{\eta}_0}$ . According to [[Künnemann 1998](#), Theorem 4.2], there is an open immersion

$$\mathcal{B}_{0,\bar{\eta}_0} \hookrightarrow S_1 \tag{10}$$

of  $C_{\bar{\eta}_0}$ -schemes, with the following properties:  $S_1$  is a regular scheme, which is projective over  $C_{\bar{\eta}_0}$  and the open immersion  $\mathcal{B}_{0,\bar{\eta}_0} \hookrightarrow S_1$  is an isomorphism when restricted to the open subset of  $C_{\bar{\eta}_0}$  over

which  $\mathcal{B}_{0,\bar{\eta}_0}$  is an abelian scheme. In particular  $S_1$  is smooth over  $\bar{\eta}_0$ , since  $\bar{\eta}_0$  is perfect. There is a finite field extension  $\eta \rightarrow \eta_0$  and a morphism

$$\mathcal{B}_{0,\eta} \rightarrow S \tag{11}$$

of  $C_\eta$ -schemes, which is model of (10). By flat descent, the morphism  $\mathcal{B}_{0,\eta} \rightarrow S$  is also an open immersion and  $S$  is also smooth over  $\eta$  and projective over  $C_\eta$ . Again by flat descent  $\mathcal{B}_{0,\eta} \rightarrow S$  is an isomorphism when restricted to the open subset of  $C_\eta$  over which  $\mathcal{B}_{0,\eta}$  is an abelian scheme.

We now let  $g : H \rightarrow H_0$  be the normalisation of  $H_0$  in  $\eta$ . Slightly abusing notation, we also denote by  $\eta$  the generic point of  $H$ . Note that  $g$  is a finite morphism (see, e.g., [EGA IV<sub>2</sub> 1965, p. 214–218]). We let  $\mathcal{B}$  be the semiabelian scheme on  $C_H$  obtained by base change and we let  $\lambda$  be the generic point of  $C_H$ . Again  $\lambda$  lies over  $\eta$  via the second projection and is also the generic point of the  $C_\eta$ . By an elementary constructibility argument, there is a nonempty open set  $U \subseteq H$  and an open immersion

$$\mathcal{B}_{C_U} \hookrightarrow \tilde{\mathcal{S}}$$

of  $C_U$ -schemes, where  $\tilde{\mathcal{S}}$  is smooth over  $U$  and projective over  $C_U$ . Furthermore, we may assume that there is an open subset  $U' \subseteq C_U$ , which surjects onto  $U$ , with the property that  $\mathcal{B}_{U'}$  is an abelian scheme over  $U'$  and that the induced morphism  $\mathcal{B}_{U'} \hookrightarrow \tilde{\mathcal{S}}_{U'}$  is an isomorphism.

Let  $N_0$  be the supremum of the set of values of the function, which associates with any  $q \in C_U$  the number of geometric irreducible components of the fibre  $\tilde{\mathcal{S}}_q$  of  $\tilde{\mathcal{S}}$  over  $q$ . This function is constructible (see [EGA IV<sub>3</sub> 1966, p. 82]) and so  $N_0$  is finite.

Now let  $y \in U(t_0)$ . By construction  $\mathcal{B}_{C_y}$  is then a generically abelian semiabelian scheme over  $C_y$ . We have a canonical  $C_y$ -morphism  $f : (\tilde{\mathcal{S}}_{C_y})^{\text{sm}} \rightarrow \mathcal{N}(\mathcal{B}_{\kappa(C_y)})$  by the definition of the Néron model. Let  $P_y \in \mathcal{B}(\kappa(C_y))$ . The section  $P_y$  extends uniquely to a element of  $(\tilde{\mathcal{S}}_{C_y})^{\text{sm}}(C_y)$  by the valuative criterion of properness and Lemma A.1. It also extends uniquely to an element of  $\mathcal{N}(\mathcal{B}_{\kappa(C_y)})(C_y)$  by the definition of the Néron model. By unicity, these two extensions are compatible with the morphism  $f$ . Let  $s \in C_y(t_0)$ . Since the number of irreducible components of  $(\tilde{\mathcal{S}}_{C_y})_s^{\text{sm}}$  is  $\leq N_0$ , we see that the images of the multiples  $P_y, 2 \cdot P_y, \dots$  of  $P_y$  in  $\mathcal{N}(\mathcal{B}_{\kappa(C_y)})(s)$  are contained in at most  $N_0$  components of  $\mathcal{N}(\mathcal{B}_{\kappa(C_y)})_s$ . Hence the order of the image of  $P_y$  in the component group of  $\mathcal{N}(\mathcal{B}_{\kappa(C_y)})_s$  is  $\leq N_0$ . Since  $s$  was arbitrary, we see that  $N_0! \cdot P_y$  extends to an element of  $\mathcal{N}(\mathcal{B}_{\kappa(C_y)})^0(C_y)$ . Note also (for use in (b) below) that since  $\mathcal{B}_{C_y}$  is semiabelian,  $\mathcal{N}(\mathcal{B}_{\kappa(C_y)})^0(C_y)$  naturally identifies with  $\mathcal{B}_{C_y}$  by the unicity of semiabelian extensions.

Finally let  $V$  be the open set  $H_0 \setminus g(H \setminus U)$ . By construction, we have  $g^{-1}(V) \subseteq U$ . Thus every point of  $V(t_0)$  lifts to a point of  $U(t_0)$  (since  $g$  is finite) and we see that  $V$  has the required properties.

For the proof of (b) we first let  $U$  be as in the proof of (a). We let  $\text{Sec}_U^0(\mathcal{B}_{C_U}/C_U)$  the functor from locally noetherian  $U$ -schemes  $T$  to sets, such that

$$\text{Sec}_U^0(\mathcal{B}_{C_U}/C_U)(T) = \{\text{sections } \sigma \text{ of } \mathcal{B}_{C_T} \rightarrow C_T \text{ such that } \deg((\sigma^*(L))_{C_t}) = 0 \text{ for all } t \in T\}.$$

As  $\mathcal{B}_{C_U}$  is quasiprojective over  $U$ , this functor is representable by a scheme  $\text{Sec}_U^0(\mathcal{B}_{C_U}/C_U)$  of finite type over  $U$ . See, e.g., [Nitsure 2005, Example before 5.6.3]. See the proof of Proposition 9.6 for a similar

construction. We leave the details to the reader. Now let  $x \in g^{-1}(T_0) \cap U$ . We have an identification

$$\begin{aligned} \text{Sec}_U^0(\mathcal{B}_{C_U}/C_U)_x(t_0) &= \text{Sec}_x^0(\mathcal{B}_{C_x}/C_x)(t_0) \\ &= \{P \in \mathcal{B}_{C_x}(C_x) \mid \text{the Néron–Tate height of } P \text{ with respect to } L_{\mathcal{B}_{C_x}} \text{ vanishes}\}. \end{aligned}$$

See [Moret-Bailly 1985, III.3.2 and 3.3]. Since  $\text{Tr}_{k(C_x)|t_0}(\mathcal{B}_{0,\kappa(C_x)}) = 0$ , a theorem of Lang (see [Conrad 2006, Theorem 9.15]) implies that  $\text{Sec}_U^0(\mathcal{B}_{C_U}/C_U)_x(t_0)$  consists of torsion sections. Furthermore, by the Lang–Néron theorem,  $\text{Sec}_U^0(\mathcal{B}_{C_U}/C_U)_x(t_0)$  is finite. Hence  $\text{Sec}_U^0(\mathcal{B}_{C_U}/C_U)_x$  is quasifinite. Since quasifiniteness is a constructible property (see [EGA IV<sub>3</sub> 1966, p. 71]) and  $g^{-1}(T_0) \cap U$  is dense in  $U$  (because  $g$  is finite and  $T_0$  is dense in  $H_0$ ), this implies that the scheme  $\text{Sec}_U^0(\mathcal{B}_{C_U}/C_U)$  is quasifinite over an open subset of  $U$ . Now replace  $U$  by one of its open subschemes so that  $\text{Sec}_U^0(\mathcal{B}_{C_U}/C_U)$  becomes quasifinite over  $U$ . Let  $b_{00}$  be an upper bound for the cardinality of the fibres of  $\text{Sec}_U^0(\mathcal{B}_{C_U}/C_U) \rightarrow U$ . Using (a), we conclude that we have

$$\#(n_0 \cdot \text{Tor}(\mathcal{B}_0(\kappa(x)))) \leq b_{00}$$

for all  $x \in U(t_0)$ . In particular  $b_{00}! \cdot n_0 \cdot \text{Tor}(\mathcal{B}_0(\kappa(x)))$  is the trivial group. Thus by the structure of finite subgroups of abelian varieties, we have

$$\#\text{Tor}(\mathcal{B}_0(\kappa(x))) \leq (b_{00}! \cdot n_0)^{2 \dim(\mathcal{B}_{C_U}/C_U)},$$

and we choose  $b_0 := (b_{00}! \cdot n_0)^{2 \dim(\mathcal{B}_{C_U}/C_U)}$ . Finally we let as before  $V$  be the open set  $H_0 \setminus g(H \setminus U)$ . By construction, we have  $g^{-1}(V) \subseteq U$ . Thus every point of  $V(t_0)$  lifts to a point of  $U(t_0)$  (since  $g$  is finite) and we see that  $V$  has the required properties. □

### Appendix B: Ampleness of the Hodge bundle and inseparable points

The terminology of this appendix is independent of the terminology of the rest of the article and appendices. In this appendix, we shall prove a mild extension of the main result of [Rössler 2015].

Let  $k$  be a perfect field and let  $S$  be a geometrically connected, smooth and proper curve over  $k$ . Let  $K := \kappa(S)$  be its function field. Suppose from now on that  $k$  has characteristic  $p > 0$ .

Let  $\pi : \mathcal{A} \rightarrow S$  be a smooth commutative group scheme and let  $A := \mathcal{A}_K$  be the generic fibre of  $\mathcal{A}$ . Let  $\epsilon_{\mathcal{A}/S} : S \rightarrow \mathcal{A}$  be the zero-section and let  $\omega := \epsilon_{\mathcal{A}/S}^*(\Omega_{\mathcal{A}/S}^1)$  be the Hodge bundle of  $\mathcal{A}$  over  $S$ .

**Theorem B.1.** *Suppose that  $\mathcal{A}/S$  is semiabelian, that  $A$  is an abelian variety and that  $\bar{\mu}_{\min}(\omega) > 0$ . Then there exists  $\ell_0 \in \mathbb{N}$  such the natural injection  $A(K^{p^{-\ell_0}}) \hookrightarrow A(K^{\text{perf}})$  is surjective (and hence a bijection).*

N.B. In [Rössler 2015, Theorem 1.1], Theorem B.1 was proven under the assumption that  $A$  is principally polarised and that  $k$  is algebraically closed. In can be shown that the condition  $\bar{\mu}_{\min}(\omega) > 0$  is equivalent to the requirement that  $\omega$  is an ample bundle (see [Rössler 2015, Introduction] for detailed references).

*Proof.* Notice first that in our proof of Theorem B.1, we may replace  $K$  by a finite extension field  $K'$  without restriction of generality. We may thus suppose that  $A$  is endowed with an  $m$ -level structure for some  $m \geq 3$  with  $(m, p) = 1$ .

If  $Z \rightarrow W$  is a  $W$ -scheme and  $W$  is a scheme of characteristic  $p$ , then for any  $n \geq 0$  we shall write  $Z^{[n]} \rightarrow W$  for the  $W$ -scheme given by the composition of arrows

$$Z \rightarrow W \xrightarrow{F_W^n} W.$$

Now fix  $n \geq 1$  and suppose that  $A(K^{p^{-n}}) \setminus A(K^{p^{-n+1}}) \neq \emptyset$ .

Fix  $P \in A^{(p^n)}(K) \setminus A^{(p^{n-1})}(K) = A(K^{p^{-n}}) \setminus A(K^{p^{-n+1}})$ . The point  $P$  corresponds to a commutative diagram of  $k$ -schemes

$$\begin{array}{ccc} & & A \\ & \nearrow P & \downarrow \\ \text{Spec } K^{[n]} & \xrightarrow{F_K^n} & \text{Spec } K \end{array}$$

such that the residue field extension  $K|\kappa(P(\text{Spec } K^{[n]}))$  is of degree 1 (in other words  $P$  is birational onto its image). In particular, the map of  $K$ -vector spaces  $P^*(\Omega_{A/k}^1) \rightarrow \Omega_{K^{[n]}/k}^1$  arising from the diagram is nonzero.

Now recall that there is a canonical exact sequence

$$0 \rightarrow \pi_K^*(\Omega_{K/k}^1) \rightarrow \Omega_{A/k}^1 \rightarrow \Omega_{A/K}^1 \rightarrow 0.$$

Furthermore the map  $F_K^{n,*}(\Omega_{K/k}^1) \xrightarrow{F_K^{n,*}} \Omega_{K^{[n]}/k}^1$  vanishes. Also, we have a canonical identification  $\Omega_{A/K}^1 = \pi_K^*(\omega_K)$  (see [Bosch et al. 1990, Chapter 4, Proposition 2]). Thus the natural surjection  $P^*(\Omega_{A/k}^1) \rightarrow \Omega_{K^{[n]}/k}^1$  gives rise to a nonzero map

$$\phi_n = \phi_{n,P} : F_K^{n,*}(\omega_K) \rightarrow \Omega_{K^{[n]}/k}^1.$$

The next crucial lemma examines the poles of the morphism  $\phi_n$ .

We let  $E$  be the reduced closed subset, which is the union of the points  $s \in S$ , such that the fibre  $\mathcal{A}_s$  is not complete.

**Lemma B.2.** *The morphism  $\phi_n$  extends to a morphism of vector bundles*

$$F_S^{n,*}(\omega) \rightarrow \Omega_{S^{[n]}/k}^1(E).$$

*Proof of Lemma B.2.* First notice that there is a natural identification  $\Omega_{S^{[n]}/k}^1(\log E) = \Omega_{S^{[n]}/k}^1(E)$ , because there is a sequence of coherent sheaves

$$0 \rightarrow \Omega_{S^{[n]}/k} \rightarrow \Omega_{S^{[n]}/k}^1(\log E) \rightarrow \mathcal{O}_E \rightarrow 0,$$

where the morphism onto  $\mathcal{O}_E$  is the residue morphism. Here the sheaf  $\Omega_{S^{[n]}/k}^1(\log E)$  is the sheaf of differentials on  $S^{[n]} \setminus E$  with logarithmic singularities along  $E$ . See [Illusie 1990, Introduction] for this result and more details on these notions.

We may also suppose without restriction of generality that  $A$  is principally polarised. Indeed, consider the following reasoning. By Zarhin's trick, the abelian variety  $B := (A \times_K A^\vee)^4$  is principally polarised. Also,  $B$  can be endowed with an  $m$ -level structure compatible with the given  $m$ -level structure on  $A$ ,

since  $A^\vee$  is isogenous to  $A$ . Let  $\mathcal{B} := (\mathcal{A} \times_K \mathcal{A}^\vee)^4$ , where (abusing language) we have written  $\mathcal{A}^\vee$  for the connected component of the zero-section of the Néron model of  $A^\vee$ . The group scheme  $\mathcal{A}^\vee$  is also semiabelian, since  $A^\vee$  is isogenous to  $A$  over  $K$ . The morphism  $P \times 0 \times 0 \times \cdots \times 0$  (seven times) gives a point in  $B^{(p^n)}(K)$  and there is a commutative diagram

$$\begin{array}{ccc}
 F_K^{n,*}(\omega_{\mathcal{B},K}) & \xrightarrow{\phi_{n,P \times 0 \times \dots}} & \Omega_{K^{[n]}/k}^1 \\
 \downarrow & & \uparrow \\
 F_K^{n,*}(\omega_{\mathcal{A},K}) & \xrightarrow{\phi_{n,P}} & \Omega_{K^{[n]}/k}^1
 \end{array} \tag{12}$$

where the vertical arrow on the left is the pull-back map induced by the closed immersion  $\lambda \mapsto \lambda \times 0 \times 0 \times \cdots \times 0$  (seven times). Now since  $B$  is principally polarised, we know that if [Lemma B.2](#) holds for principally polarised abelian varieties, the upper row of the diagram (12) extends to a morphism  $F_S^{n,*}(\omega_{\mathcal{B}}) \rightarrow \Omega_{S^{[n]}/k}^1(E)$  (note that the set of points, where  $\mathcal{B}$  is not complete coincides with the set of points, where  $\mathcal{A}$  is not complete). Since  $F_S^{n,*}(\omega_{\mathcal{A}})$  is a direct summand of  $F_S^{n,*}(\omega_{\mathcal{B}})$ , we see that [Lemma B.2](#) holds for  $A$  if it holds for  $B$ , thus completing the reduction of [Lemma B.2](#) to the principally polarised case.  $\square$

The rest of the proof of [Theorem B.1](#) is identical word for word with the proof of [Theorem 1.1](#) in [\[Rössler 2015\]](#) (from the beginning of the proof of [Lemma 2.1](#)).  $\square$

### Appendix C: Specialisation of the Mordell–Weil group

The terminology of this appendix is independent of the terminology of the rest of the article and appendices.

In this appendix, we shall prove a geometric analogue of Néron’s result on the specialisation of the generic Mordell–Weil group to a fibre in a family of abelian varieties over number fields (see [\[Lang 1983, Chapter 9, Corollary 6.3\]](#)). The following results are reminiscent of some results proven by Hrushovski [\[1998\]](#) in a mixed characteristic context and they are probably already known to many people but we include complete proofs for lack of a reference.

Let  $l_0$  be an algebraically closed field. Let  $U$  be a smooth and connected quasiprojective variety over  $l_0$ . Let  $\mathcal{B}$  be an abelian scheme over  $U$ . Suppose given an immersion  $\iota : U \hookrightarrow \mathbb{P}^N$  for some  $N \geq 0$ . Let  $K$  be the function field of  $U$  and let  $B := \mathcal{B}_K$ .

**Proposition C.1.** *Suppose that  $\mathcal{B}(U)$  is finitely generated. For almost all linear subspaces  $L \subseteq \mathbb{P}^N$  of codimension  $\dim(U) - 1$ , the intersection  $C := L \cap U$  is smooth, connected, nonempty, the specialisation map*

$$\mathcal{B}(U) \rightarrow \mathcal{B}_C(C)$$

*is injective and  $\text{Tr}_{\kappa(C)|l_0}(\mathcal{B}_{\kappa(C)}) = 0$ .*

Recall that the linear subspaces  $L \subseteq \mathbb{P}^N$  of codimension  $\dim(U) - 1$  are classified by the Grassmannian  $\text{Gr}(\dim(U) - 1, N)$ , which is smooth and projective over  $l_0$ . The words “almost all” stand for “for all the  $l_0$ -rational points of some dense Zariski open subset of  $\text{Gr}(\dim(U) - 1, N)$ ”.

Recall that by a theorem of Weil, the restriction map  $\mathcal{B}(U) \rightarrow B(K)$  is a bijection. Thus, by the Lang–Néron theorem, the condition that  $\mathcal{B}(U) = B(K)$  is finitely generated is equivalent to the condition  $\mathrm{Tr}_{K|k_0}(B) = 0$ .

For the proof of [Proposition C.1](#), we shall need a few lemmata:

**Lemma C.2.** *Let  $N$  be a finite étale group scheme over  $U$ . Let  $t \in H_{\mathrm{et}}^1(U, N)$  and suppose that  $t \neq 0$ . Then for almost all linear subspaces  $L \subseteq \mathbb{P}^N$  of codimension  $\dim(U) - 1$ , the intersection  $C := L \cap U$  is smooth, connected, nonempty and the restriction  $t_C \in H_{\mathrm{et}}^1(C, N_C)$  of  $t$  to  $C$  does not vanish.*

*Proof.* Let  $T \rightarrow U$  be a torsor under  $N$ . Note that the torsor  $T$  is nontrivial if and only if for all the irreducible components  $T'$  of  $T$ , the (automatically flat and finite) morphism  $T' \rightarrow U$  has degree  $> 1$ . The same remark applies to the restriction of  $T$  to a smooth and connected closed subscheme of  $U$ .

Let  $(T_i)$  be the set of irreducible components of  $T$ .

By Bertini’s theorem in Jouanolou’s presentation [[1983](#), p. 89, Corollary 6.11], for almost all linear subspaces  $L \subseteq \mathbb{P}^N$  of codimension  $\dim(U) - 1$ ,

- the intersection  $C := L \cap U$  is smooth, connected and nonempty;
- all the  $T_{i,C}$  are irreducible.

Let  $C$  be in this class. Suppose that  $T \rightarrow U$  is not trivial. By construction, the irreducible components of  $T_C$  are the  $T_{i,C}$ . Since  $T_{i,C} \rightarrow C$  is flat and finite of the same degree as  $T_i \rightarrow U$ , we see that the irreducible components of  $T_C$  all have degree  $> 1$  over  $C$ . Hence the torsor  $T_C$  is not trivial. □

**Lemma C.3.** *Let  $N$  be a finite étale group scheme over  $U$ . Suppose that  $N(U) = 0$ . Then for almost all linear subspaces  $L \subseteq \mathbb{P}^N$  of codimension  $\dim(U) - 1$ , the intersection  $C := L \cap U$  is smooth, connected, nonempty and  $N_C(C) = 0$ .*

*Proof.* Let  $(N_i)$  be the set of irreducible components of  $N$ , excluding the component of the identity. The condition that  $N(U) = 0$  is equivalent to the condition that for all  $i$ , the morphism  $N_i \rightarrow U$  has degree  $> 1$ .

As before, by Bertini’s theorem, for almost all linear subspaces  $L \subseteq \mathbb{P}^N$  of codimension  $\dim(U) - 1$ ,

- the intersection  $C := L \cap U$  is smooth and connected;
- all the  $N_{i,C}$  are irreducible.

Let  $C$  be in this class. By construction, the irreducible components of  $N_C$  outside of the component of the identity are the  $N_{i,C}$ . Since  $N_{i,C} \rightarrow C$  is flat and finite of the same degree as  $N_i \rightarrow U$ , we see that the irreducible components of  $N_C$  outside of the component of the identity all have degree  $> 1$  over  $C$ . Hence  $N_C(C) = 0$ . □

**Lemma C.4.** *Let  $G \subseteq \mathcal{B}(U)$  be a finite group. For almost all linear subspaces  $L \subseteq \mathbb{P}^N$  of codimension  $\dim(U) - 1$ , the intersection  $C := L \cap U$  is smooth and connected, and the reduction map*

$$G \rightarrow \mathcal{B}_C(C)$$

*is injective.*

*Proof.* The proof is left to the reader. □

Finally, we need an elementary but very insightful lemma, due to in essence to Néron. The following version is due to Hrushovski [1998, Lemma 1]:

**Lemma C.5** (Néron–Hrushovski). *Let  $r : G \rightarrow H$  be a map of abelian groups. Let  $l$  be a prime number. Suppose that  $\mathrm{Tor}_l(H) = 0$  and that the induced map  $G/lG \rightarrow H/lH$  is injective. Then  $\ker r \subseteq \bigcap_{j \geq 0} l^j G$ .*

*Proof.* Let  $g \in \ker r$ . Suppose for contradiction that  $g \notin \bigcap_{j \geq 0} l^j G$ . Let  $m \geq 0$  be the smallest natural number such that  $g \notin l^m G$ . Then there is  $g' \in G$  such that  $l^{m-1}g' = g$  and thus  $r(g') \in \mathrm{Tor}_l(H)$  so that from the assumptions we have  $r(g') = 0$ . Since the map  $G/lG \rightarrow H/lH$  is injective, there is  $g'' \in G$  such that  $lg'' = g'$ . Hence  $g = l^m g''$ , a contradiction.  $\square$

*Proof of Proposition C.1.* Let  $l$  be a prime number such that  $\mathrm{Tor}_l(\mathcal{B}(U)) = 0$  and such that  $l$  is not the characteristic of  $l_0$ . Notice that for any closed subscheme  $C$  of  $U$ , we have an injection  $\delta_C : \mathcal{B}(C)/l\mathcal{B}(C) \hookrightarrow H_{\mathrm{et}}^1(C, \ker [l]_{\mathcal{B},C})$  and this injection is functorial for restrictions to smaller closed subschemes  $C_1 \hookrightarrow C$ . According to Lemmata C.2, C.3 and C.4, for almost all linear subspaces  $L \subseteq \mathbb{P}^N$  of codimension  $\dim(U) - 1$ ,

- the intersection  $C := L \cap U$  is smooth and connected;
- the restriction map  $H^1(U, \ker [l]_{\mathcal{B}}) \rightarrow H^1(C, \ker [l]_{\mathcal{B},C})$  is injective on the image of  $\delta_U$ ;
- $(\ker [l]_{\mathcal{B},C})(C) = 0$ ;
- the restriction map  $\mathrm{Tor}(\mathcal{B}(U)) \rightarrow \mathcal{B}(C)$  is injective.

Let  $C$  be in this class. By construction, the map  $\mathcal{B}(U)/l\mathcal{B}(U) \rightarrow \mathcal{B}(C)/l\mathcal{B}(C)$  is injective and  $\mathrm{Tor}_l(\mathcal{B}(C)) = 0$ . Let  $F$  be a free subgroup of  $\mathcal{B}(U)$ , which is a direct summand of  $\mathrm{Tor}(\mathcal{B}(U))$ . We have  $F \cap (\bigcap_{j \geq 0} l^j \mathcal{B}(U)) = 0$  since  $\mathcal{B}(U)$  is finitely generated and  $F$  is free. Applying Lemma C.5 to  $G = \mathcal{B}(U)$  and  $H = \mathcal{B}(C)$ , we see that the restriction map  $F \rightarrow \mathcal{B}(C)$  is injective. Since the restriction map  $\mathrm{Tor}(\mathcal{B}(U)) \rightarrow \mathcal{B}(C)$  is also injective, we thus see that the restriction map  $\mathcal{B}(U) \rightarrow \mathcal{B}(C)$  is injective. Finally, we have  $\mathrm{Tr}_{\kappa(C)/l_0}(\mathcal{B}_{\kappa(C)}) = 0$ , for otherwise, we would have  $\mathrm{Tor}_l(\mathcal{B}(C)) \neq 0$ .  $\square$

### Acknowledgments

My warm thanks to the referee for his/her careful reading and for many suggestions. The article would be much less clear without his/her help and encouragement. I would like to thank J.-B. Bost for his feedback, especially for pointing out the article [Catanese and Dettweiler 2016], for suggesting Remark 2.4 and for providing [Bost 2004, Lemma 2.9], whose positive characteristic analogue is technically at the root of the present text. Minhyong Kim’s article [1997] also played a fundamental role in the genesis of the present text; the construction described there pointed me in (what I hope is) the right direction when I started studying purely inseparable points on abelian varieties. I had many interesting discussions with him about his article. I am very grateful to J.-F. Voloch for many exchanges on the material of this article and for his remarks on the text and to P. Ziegler for many discussions on and around the “full” Mordell–Lang conjecture. Many thanks also to T. Scanlon for his interest and for interesting discussions

Purely inseparable points of an abelian variety defined over a function field of positive characteristic 1171

around the group  $A(K^{\text{perf}})$ . Last but not least, many thanks to H el ene Esnault and her student Marco d’Addezio for their interest and for many enlightening discussions around [Theorem 1.4](#). I also benefitted from A.-J. de Jong’s and F. Oort’s vast knowledge; they both very kindly took the time to answer some rather speculative messages.

## References

- [Abbes 2000] A. Abbes, “R eduction semi-stable des courbes d’apr es Artin, Deligne, Grothendieck, Mumford, Saito, Winters, . . .”, pp. 59–110 in *Courbes semi-stables et groupe fondamental en g eom etrie alg ebrique* (Luminy, 1998), edited by J.-B. Bost et al., Progr. Math. **187**, Birkh user, Basel, 2000. [MR](#) [Zbl](#)
- [Abramovich and Voloch 1992] D. Abramovich and J. F. Voloch, “Toward a proof of the Mordell–Lang conjecture in characteristic  $p$ ”, *Int. Math. Res. Not.* **5** (1992), 103–115. [MR](#) [Zbl](#)
- [Artin and Milne 1976] M. Artin and J. S. Milne, “Duality in the flat cohomology of curves”, *Invent. Math.* **35** (1976), 111–129. [MR](#) [Zbl](#)
- [Barton 1971] C. M. Barton, “Tensor products of ample vector bundles in characteristic  $p$ ”, *Amer. J. Math.* **93** (1971), 429–438. [MR](#) [Zbl](#)
- [Bosch et al. 1990] S. Bosch, W. L utkebohmert, and M. Raynaud, *N eron models*, Ergebnisse der Mathematik und ihrer Grenzgebiete (3) **21**, Springer, 1990. [MR](#) [Zbl](#)
- [Bost 2004] J.-B. Bost, “Germes of analytic varieties in algebraic varieties: canonical metrics and arithmetic algebraization theorems”, pp. 371–418 in *Geometric aspects of Dwork theory*, vol. I, edited by A. Adolphson et al., Walter de Gruyter, Berlin, 2004. [MR](#) [Zbl](#)
- [Brenner et al. 2008] H. Brenner, J. Herzog, and O. Villamayor, *Three lectures on commutative algebra*, edited by G. Colom -Nin et al., University Lecture Series **42**, Amer. Math. Soc., Providence, RI, 2008. [MR](#) [Zbl](#)
- [Catanese and Dettweiler 2016] F. Catanese and M. Dettweiler, “Vector bundles on curves coming from variation of Hodge structures”, *Internat. J. Math.* **27**:7 (2016), art. id. 1640001, 25 pp. [MR](#) [Zbl](#)
- [Conrad 2006] B. Conrad, “Chow’s  $K/k$ -image and  $K/k$ -trace, and the Lang–N eron theorem”, *Enseign. Math.* (2) **52**:1–2 (2006), 37–108. [MR](#) [Zbl](#)
- [EGA IV<sub>2</sub> 1965] A. Grothendieck, “El ements de g eom etrie alg ebrique, IV:  tude locale des sch emas et des morphismes de sch emas, II”, *Inst. Hautes  tudes Sci. Publ. Math.* **24** (1965), 5–231. [MR](#) [Zbl](#)
- [EGA IV<sub>3</sub> 1966] A. Grothendieck, “El ements de g eom etrie alg ebrique, IV:  tude locale des sch emas et des morphismes de sch emas, III”, *Inst. Hautes  tudes Sci. Publ. Math.* **28** (1966), 5–255. [MR](#) [Zbl](#)
- [EGA IV<sub>4</sub> 1967] A. Grothendieck, “El ements de g eom etrie alg ebrique, IV:  tude locale des sch emas et des morphismes de sch emas, IV”, *Inst. Hautes  tudes Sci. Publ. Math.* **32** (1967), 5–361. [MR](#) [Zbl](#)
- [Ekedahl 1988] T. Ekedahl, “Canonical models of surfaces of general type in positive characteristic”, *Inst. Hautes  tudes Sci. Publ. Math.* **67** (1988), 97–144. [MR](#) [Zbl](#)
- [Esnault and Langer 2013] H. Esnault and A. Langer, “On a positive equicharacteristic variant of the  $p$ -curvature conjecture”, *Doc. Math.* **18** (2013), 23–50. [MR](#) [Zbl](#)
- [Faltings and Chai 1990] G. Faltings and C.-L. Chai, *Degeneration of abelian varieties*, Ergebnisse der Mathematik und ihrer Grenzgebiete (3) **22**, Springer, 1990. [MR](#) [Zbl](#)
- [Fantechi et al. 2005] B. Fantechi, L. G ottsche, L. Illusie, S. L. Kleiman, N. Nitsure, and A. Vistoli, *Fundamental algebraic geometry: Grothendieck’s FGA explained*, Mathematical Surveys and Monographs **123**, Amer. Math. Soc., Providence, RI, 2005. [MR](#) [Zbl](#)
- [Ghioca and Moosa 2006] D. Ghioca and R. Moosa, “Division points on subvarieties of isotrivial semi-abelian varieties”, *Int. Math. Res. Not.* **2006** (2006), art. id. 65437, 23 pp. [MR](#) [Zbl](#)
- [Grothendieck 1966] A. Grothendieck, “Techniques de construction et th eor emes d’existence en g eom etrie alg ebrique IV: Les sch emas de Hilbert”, expos e 221 in *S minaire Bourbaki*, 1960/1961, W. A. Benjamin, Amsterdam, 1966. Reprinted as pp. 249–276 in *S minaire Bourbaki* **6**, Soc. Math. France, Paris, 1995. [MR](#)

- [Grothendieck 1974] A. Grothendieck, *Groupes de Barsotti–Tate et cristaux de Dieudonné* (Montréal, 1970), Les Presses de l'Université de Montréal, 1974. [MR](#) [Zbl](#)
- [Hartshorne 1977] R. Hartshorne, *Algebraic geometry*, Graduate Texts in Mathematics **52**, Springer, 1977. [MR](#) [Zbl](#)
- [Hrushovski 1998] E. Hrushovski, “Proof of Manin’s theorem by reduction to positive characteristic”, pp. 197–205 in *Model theory and algebraic geometry*, edited by E. Bouscaren, Lecture Notes in Math. **1696**, Springer, 1998. [MR](#) [Zbl](#)
- [Illusie 1990] L. Illusie, “Réduction semi-stable et décomposition de complexes de de Rham à coefficients”, *Duke Math. J.* **60**:1 (1990), 139–185. [MR](#) [Zbl](#)
- [Jouanolou 1983] J.-P. Jouanolou, *Théorèmes de Bertini et applications*, Progress in Mathematics **42**, Birkhäuser, Boston, 1983. [MR](#) [Zbl](#)
- [Katz 1981] N. Katz, “Serre–Tate local moduli”, pp. 138–202 in *Algebraic surfaces* (Orsay, 1976–78), edited by J. Giraud et al., Lecture Notes in Math. **868**, Springer, 1981. [MR](#) [Zbl](#)
- [Kim 1997] M. Kim, “Purely inseparable points on curves of higher genus”, *Math. Res. Lett.* **4**:5 (1997), 663–666. [MR](#) [Zbl](#)
- [Künnemann 1998] K. Künnemann, “Projective regular models for abelian varieties, semistable reduction, and the height pairing”, *Duke Math. J.* **95**:1 (1998), 161–212. [MR](#) [Zbl](#)
- [Lang 1983] S. Lang, *Fundamentals of Diophantine geometry*, Springer, 1983. [MR](#) [Zbl](#)
- [Langer 2004] A. Langer, “Semistable sheaves in positive characteristic”, *Ann. of Math. (2)* **159**:1 (2004), 251–276. [MR](#) [Zbl](#)
- [Langer 2015] A. Langer, “Generic positivity and foliations in positive characteristic”, *Adv. Math.* **277** (2015), 1–23. [MR](#) [Zbl](#)
- [Liu 2002] Q. Liu, *Algebraic geometry and arithmetic curves*, Oxford Graduate Texts in Mathematics **6**, Oxford University Press, 2002. [MR](#) [Zbl](#)
- [Milne 1980] J. S. Milne, *Étale cohomology*, Princeton Mathematical Series **33**, Princeton University Press, 1980. [MR](#) [Zbl](#)
- [Milne 1986] J. S. Milne, “Abelian varieties”, pp. 103–150 in *Arithmetic geometry* (Storrs, CT, 1984), edited by G. Cornell and J. H. Silverman, Springer, 1986. [MR](#) [Zbl](#)
- [Milne 2006] J. S. Milne, *Arithmetic duality theorems*, 2nd ed., BookSurge, Charleston, SC, 2006. [MR](#) [Zbl](#)
- [Milne 2017] J. S. Milne, *Algebraic groups: the theory of group schemes of finite type over a field*, Cambridge Studies in Advanced Mathematics **170**, Cambridge University Press, 2017. [MR](#) [Zbl](#)
- [Moret-Bailly 1985] L. Moret-Bailly, *Pinceaux de variétés abéliennes*, Astérisque **129**, Société Mathématique de France, Paris, 1985. [MR](#) [Zbl](#)
- [Mumford 1970] D. Mumford, *Abelian varieties*, Tata Institute of Fundamental Research Studies in Mathematics **5**, Oxford University Press, London, 1970. [MR](#) [Zbl](#)
- [Mumford et al. 1994] D. Mumford, J. Fogarty, and F. Kirwan, *Geometric invariant theory*, 3rd ed., Ergebnisse der Mathematik und ihrer Grenzgebiete (2) **34**, Springer, 1994. [MR](#) [Zbl](#)
- [Nitsure 2005] N. Nitsure, “Construction of Hilbert and Quot schemes”, pp. 105–137 in *Fundamental algebraic geometry: Grothendieck’s FGA explained*, Math. Surveys Monogr. **123**, Amer. Math. Soc., Providence, RI, 2005. [MR](#)
- [Poonen and Voloch 2010] B. Poonen and J. F. Voloch, “The Brauer–Manin obstruction for subvarieties of abelian varieties over function fields”, *Ann. of Math. (2)* **171**:1 (2010), 511–532. [MR](#) [Zbl](#)
- [Raynaud 1970] M. Raynaud, *Faisceaux amples sur les schémas en groupes et les espaces homogènes*, Lecture Notes in Mathematics **119**, Springer, 1970. [MR](#) [Zbl](#)
- [Rössler 2013] D. Rössler, “Infinitely  $p$ -divisible points on abelian varieties defined over function fields of characteristic  $p > 0$ ”, *Notre Dame J. Form. Log.* **54**:3-4 (2013), 579–589. [MR](#) [Zbl](#)
- [Rössler 2015] D. Rössler, “On the group of purely inseparable points of an abelian variety defined over a function field of positive characteristic”, *Comment. Math. Helv.* **90**:1 (2015), 23–32. [MR](#) [Zbl](#)
- [Rössler 2019a] D. Rössler, “Le groupe de Selmer des isogénies de hauteur un”, preprint, 2019. [arXiv](#)
- [Rössler 2019b] D. Rössler, “Perfect points on abelian varieties in positive characteristic”, video, 2019, available at <https://www.youtube.com/watch?v=TKLEIpNsYDI>.
- [Scanlon 2005] T. Scanlon, “A positive characteristic Manin–Mumford theorem”, *Compos. Math.* **141**:6 (2005), 1351–1364. [MR](#) [Zbl](#)

- [Serre and Tate 1968] J.-P. Serre and J. Tate, “Good reduction of abelian varieties”, *Ann. of Math. (2)* **88** (1968), 492–517. [MR](#) [Zbl](#)
- [SGA 3<sub>I</sub> 2011] M. Demazure and A. Grothendieck, *Schémas en groupes, Tome I: Propriétés générales des schémas en groupes, Exposés I–VII* (Séminaire de Géométrie Algébrique du Bois Marie 1962–1964), revised ed., edited by P. Gille and P. Polo, Documents Mathématiques (Paris) **7**, Société Mathématique de France, Paris, 2011. [MR](#) [Zbl](#)
- [SGA 3<sub>II</sub> 1970] M. Demazure and A. Grothendieck, *Schémas en groupes, Tome II: Groupes de type multiplicatif, et structure des schémas en groupes généraux, Exposés VIII–XVIII* (Séminaire de Géométrie Algébrique du Bois Marie 1962–1964), Lecture Notes in Math. **152**, Springer, 1970. [MR](#) [Zbl](#)
- [Shepherd-Barron 1992] N. I. Shepherd-Barron, “Miyaoka’s theorems on the generic seminegativity of  $T_X$  and on the Kodaira dimension of minimal regular threefolds”, pp. 103–114 in *Flips and abundance for algebraic threefolds*, Astérisque **211**, Société Mathématique de France, Paris, 1992. [Zbl](#)
- [Szamuely 2010] T. Szamuely, “Corps de classes des schémas arithmétiques”, exposé 1006, pp. 257–286 in *Séminaire Bourbaki, 2008/2009*, Astérisque **332**, Société Mathématique de France, Paris, 2010. [MR](#) [Zbl](#)
- [Tate 1997] J. Tate, “Finite flat group schemes”, pp. 121–154 in *Modular forms and Fermat’s last theorem* (Boston, MA, 1995), edited by G. Cornell et al., Springer, 1997. [MR](#) [Zbl](#)
- [Tate and Oort 1970] J. Tate and F. Oort, “Group schemes of prime order”, *Ann. Sci. École Norm. Sup. (4)* **3** (1970), 1–21. [MR](#) [Zbl](#)
- [Voloch 1995] J. F. Voloch, “Diophantine approximation on abelian varieties in characteristic  $p$ ”, *Amer. J. Math.* **117**:4 (1995), 1089–1095. [MR](#) [Zbl](#)
- [Yuan 2018] X. Yuan, “Positivity of Hodge bundles of abelian varieties over some function fields”, preprint, 2018. [arXiv](#)
- [Zarkhin and Parshin 1989] Y. G. Zarkhin and A. N. Parshin, “Finiteness problems in Diophantine geometry”, pp. 35–102 in *American Mathematical Society translations*, edited by B. Silver, American Mathematical Society Translations (2) **143**, Amer. Math. Soc., Providence, RI, 1989. [Zbl](#)

Communicated by Hélène Esnault

Received 2018-09-20    Revised 2019-11-19    Accepted 2019-12-17

[damian.rossler@maths.ox.ac.uk](mailto:damian.rossler@maths.ox.ac.uk)

*Mathematical Institute, University of Oxford, Oxford, United Kingdom*

# Algebra & Number Theory

[msp.org/ant](http://msp.org/ant)

## EDITORS

### MANAGING EDITOR

Bjorn Poonen  
Massachusetts Institute of Technology  
Cambridge, USA

### EDITORIAL BOARD CHAIR

David Eisenbud  
University of California  
Berkeley, USA

### BOARD OF EDITORS

Bhargav Bhatt	University of Michigan, USA	Martin Olsson	University of California, Berkeley, USA
Richard E. Borcherds	University of California, Berkeley, USA	Raman Parimala	Emory University, USA
Antoine Chambert-Loir	Université Paris-Diderot, France	Jonathan Pila	University of Oxford, UK
J-L. Colliot-Thélène	CNRS, Université Paris-Sud, France	Irena Peeva	Cornell University, USA
Brian D. Conrad	Stanford University, USA	Anand Pillay	University of Notre Dame, USA
Samit Dasgupta	Duke University, USA	Michael Rapoport	Universität Bonn, Germany
Hélène Esnault	Freie Universität Berlin, Germany	Victor Reiner	University of Minnesota, USA
Gavril Farkas	Humboldt Universität zu Berlin, Germany	Peter Sarnak	Princeton University, USA
Hubert Flenner	Ruhr-Universität, Germany	Michael Singer	North Carolina State University, USA
Sergey Fomin	University of Michigan, USA	Christopher Skinner	Princeton University, USA
Edward Frenkel	University of California, Berkeley, USA	Vasudevan Srinivas	Tata Inst. of Fund. Research, India
Wee Teck Gan	National University of Singapore	J. Toby Stafford	University of Michigan, USA
Andrew Granville	Université de Montréal, Canada	Shunsuke Takagi	University of Tokyo, Japan
Ben J. Green	University of Oxford, UK	Pham Huu Tiep	University of Arizona, USA
Joseph Gubeladze	San Francisco State University, USA	Ravi Vakil	Stanford University, USA
Christopher Hacon	University of Utah, USA	Michel van den Bergh	Hasselt University, Belgium
Roger Heath-Brown	Oxford University, UK	Akshay Venkatesh	Institute for Advanced Study, USA
János Kollár	Princeton University, USA	Marie-France Vignéras	Université Paris VII, France
Philippe Michel	École Polytechnique Fédérale de Lausanne	Melanie Matchett Wood	University of California, Berkeley, USA
Susan Montgomery	University of Southern California, USA	Shou-Wu Zhang	Princeton University, USA
Shigefumi Mori	RIMS, Kyoto University, Japan		

## PRODUCTION

[production@msp.org](mailto:production@msp.org)

Silvio Levy, Scientific Editor

---

See inside back cover or [msp.org/ant](http://msp.org/ant) for submission instructions.

The subscription price for 2020 is US \$415/year for the electronic version, and \$620/year (+\$60, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to MSP.

Algebra & Number Theory (ISSN 1944-7833 electronic, 1937-0652 printed) at Mathematical Sciences Publishers, 798 Evans Hall #3840, c/o University of California, Berkeley, CA 94720-3840 is published continuously online. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices.

---

ANT peer review and production are managed by EditFLOW<sup>®</sup> from MSP.

PUBLISHED BY

 **mathematical sciences publishers**  
nonprofit scientific publishing

<http://msp.org/>

© 2020 Mathematical Sciences Publishers

# Algebra & Number Theory

Volume 14    No. 5    2020

---

The universal family of semistable $p$ -adic Galois representations URS HARTL and EUGEN HELLMANN	1055
On the group of purely inseparable points of an abelian variety defined over a function field of positive characteristic, II DAMIAN RÖSSLER	1123
Mixed Tate motives and the unit equation II ISHAI DAN-COHEN	1175
$p$ -adic distribution of CM points and Hecke orbits I: Convergence towards the Gauss point SEBASTIÁN HERRERO, RICARDO MENARES and JUAN RIVERA-LETÉLIER	1239
Roots of $L$ -functions of characters over function fields, generic linear independence and biases CORENTIN PERRET-GENTIL	1291