

Algebra & Number Theory

Volume 14
2020
No. 8

Auslander correspondence for triangulated categories

Norihiro Hanihara



Auslander correspondence for triangulated categories

Norihiro Hanihara

We give analogues of the Auslander correspondence for two classes of triangulated categories satisfying certain finiteness conditions. The first class is triangulated categories with additive generators and we consider their endomorphism algebras as the Auslander algebras. For the second one, we introduce the notion of [1]-additive generators and consider their graded endomorphism algebras as the Auslander algebras. We give a homological characterization of the Auslander algebras for each class. Along the way, we also show that the algebraic triangle structures on the homotopy categories are unique up to equivalence.

1. Introduction

The main concern in representation theory of algebras is to understand the module categories. Among such categories, those with *finitely many* indecomposable objects, or equivalently the *representation-finite* algebras, are most fundamental. Let us recall the following famous theorem due to Auslander [1971]:

Theorem 1.1 (Auslander correspondence). *There exists a bijection between the set of Morita equivalence classes of finite dimensional algebras Λ of finite representation type and the set of Morita equivalence classes of finite dimensional algebras Γ such that $\text{gl. dim } \Gamma \leq 2$ and $\text{dom. dim } \Gamma \geq 2$.*

This theorem states that a categorical property (=representation-finiteness) of $\text{mod } \Lambda$ can be characterized by homological invariants (=gl. dim and dom. dim) of Γ , called the *Auslander algebra* of $\text{mod } \Lambda$. There are many results of this type giving the relationships between categorical properties of those appearing naturally in representation theory, and homological properties of their “Auslander algebras”, for example, [Iyama 2005; 2007; Enomoto 2018].

The aim of this paper is to find an analogue of these results for triangulated categories [Neeman 2001]. Let k be an arbitrary field and \mathcal{T} be a k -linear, Hom-finite, idempotent-complete triangulated category. We consider two kinds of finiteness conditions on triangulated categories.

The first one is a direct analogue of representation-finiteness: \mathcal{T} is *finite*, that is, \mathcal{T} has finitely many indecomposable objects up to isomorphism. In this case, \mathcal{T} has an additive generator M . We call $\text{End}_{\mathcal{T}}(M)$ the *Auslander algebra* of \mathcal{T} , which is uniquely determined by \mathcal{T} up to Morita equivalence. The first main result of this paper is the following homological characterization of the Auslander algebras of triangulated categories. We say that a finite dimensional algebra A is *twisted n -periodic* if it is self-injective and there

This work is supported by JSPS KAKENHI Grant Number JP19J21165.

MSC2010: primary 18E30; secondary 16E05, 16E65, 16G70.

Keywords: triangulated category, Auslander correspondence, periodic algebra, Cohen–Macaulay module.

exists an automorphism α of A such that $\Omega^n \simeq (-)_\alpha$ as functors on $\underline{\text{mod}} A$. We refer to Corollary 2.2 for equivalent characterizations.

Theorem 1.2. *Let k be a perfect field. The following are equivalent for a basic finite dimensional k -algebra A :*

- (1) *A is the Auslander algebra of a k -linear, Hom-finite, idempotent-complete triangulated category which is finite.*
- (2) *A is twisted 3-periodic.*

This result shows a close connection between periodic algebras [Erdmann and Skowroński 2008] and triangulated categories. Our proof depends on Amiot’s result (Proposition 3.2). This is a complement of Heller’s classical observation [1968, 16.4] which gives a parametrization of pretriangle structures on a pretriangulated category \mathcal{T} in terms of isomorphisms $\Omega^3 \simeq [-1]$ on $\underline{\text{mod}} \mathcal{T}$. Later practice of this property of the third syzygy in representation theory can be seen in [Auslander and Reiten 1996; Yoshino 2005; Amiot 2007; Iyama and Oppermann 2013].

Moreover, with some additional assumptions on \mathcal{T} , we give a bijection between finite triangulated categories and certain algebras, which is a more precise form of the above theorem; see Theorem 3.4. Furthermore, after submitting this article, a similar result by Muro [2020] appeared. His main result enables us to state Theorem 3.4 with less additional assumptions; see Remark 3.5.

The second finiteness condition is the following:

- (S1) There is an object $M \in \mathcal{T}$ such that $\mathcal{T} = \text{add}\{M[n] \mid n \in \mathbb{Z}\}$.
- (S2) For any $X, Y \in \mathcal{T}$, $\text{Hom}_{\mathcal{T}}(X, Y[n]) = 0$ holds for almost all n .

If these conditions are satisfied, we say \mathcal{T} is [1]-finite and call M as in (S1) a [1]-additive generator. For example, the bounded derived categories of representation-finite hereditary algebras are [1]-finite, and additive generators for module categories are [1]-additive generators for the derived categories. There are various studies on [1]-finite triangulated categories, for example [Rouquier 2008; Xiao and Zhu 2005; Amiot 2007]. Note that [1]-finite triangulated categories have infinitely many indecomposable objects unless $\mathcal{T} = 0$.

For a [1]-finite triangulated category \mathcal{T} with a [1]-additive generator M , we call

$$C = \bigoplus_{n \in \mathbb{Z}} \text{Hom}_{\mathcal{T}}(M, M[n])$$

the [1]-Auslander algebra of \mathcal{T} , which is naturally a \mathbb{Z} -graded algebra and is uniquely determined by \mathcal{T} up to graded Morita equivalence. Thanks to our condition (S2), C is finite dimensional. To study it, we prepare some results on “graded projectivization” in Section 4 (see Proposition 4.2). Such constructions of graded algebras appear naturally in various contexts [Artin and Zhang 1994; Asashiba 2017].

Our second main result is the Auslander correspondence for [1]-finite triangulated categories. To state it, we have to restrict to a nice class of triangulated categories called *algebraic*. Recall that they are

the stable categories of Frobenius categories [Happel 1988, I.2.6]. Algebraic triangulated categories are enhanced by differential graded categories [Keller 2006], and play a central role in tilting theory [Angeleri Hügel et al. 2007].

Now we can formulate the following second main result of this paper in terms of algebraic triangulated categories and graded algebras. We say that a finite dimensional \mathbb{Z} -graded algebra A is *(a)-twisted n-periodic* if it is self-injective and there exists a graded automorphism α of A such that $P_\alpha \simeq P$ for all $P \in \text{proj}^{\mathbb{Z}} A$ and $\Omega^n \simeq (-)_\alpha(a)$ as functors on $\underline{\text{mod}}^{\mathbb{Z}} A$. We refer to Corollary 2.4 for equivalent conditions.

Theorem 1.3. *Let k be an algebraically closed field. There exists a bijection between the following:*

- (1) *The set of triangle equivalence classes of k -linear, Hom-finite, idempotent-complete, algebraic triangulated categories \mathcal{T} which are [1]-finite.*
- (2) *The graded Morita equivalence classes (see Definition 4.3) of finite dimensional graded k -algebra C which are (-1) -twisted 3-periodic.*
- (3) *A disjoint union of Dynkin diagrams of type A, D, and E.*

The correspondences are given as follows:

- From (1) to (2): Taking the [1]-Auslander algebra of \mathcal{T} .
- From (1) to (3): Taking the tree type of the AR-quiver of \mathcal{T} .
- From (2) to (1): $C \mapsto \text{proj}^{\mathbb{Z}} C$.
- From (3) to (1): $Q \mapsto k(\mathbb{Z}Q)$, where $k(\mathbb{Z}Q)$ is the mesh category associated with $\mathbb{Z}Q$.

Moreover, we have the following explicit descriptions of (1) and (2) in the above theorem.

Theorem 1.4 (Theorem 5.3, Proposition 6.1). *The classes (1) and (2) in Theorem 1.3 are the same as (1') and (2'), respectively:*

- (1') *The set of triangle equivalence classes of the bounded derived categories $D^b(\text{mod } kQ)$ of the path algebra kQ for a disjoint union Q of Dynkin quivers of type A, D, and E.*
- (2') *The orbit algebras $k(\mathbb{Z}Q)/[1]$ for a disjoint union Q of Dynkin quivers of type A, D, and E.*

Compared to Theorem 1.2, Theorem 1.3 is more strict in the point that the Auslander algebras C correspond *bijectionally* to the triangulated categories. This can be done by the classification of [1]-finite triangulated categories as is stated in (1'). These results suggest that [1]-finite triangulated categories are easier than finite ones in controlling their triangle structures as well as their additive structures.

Our classification is deduced from the following uniqueness of the triangle structures on the homotopy categories, which is somehow surprising; compare [Keller 2018].

Theorem 1.5 (Theorem 5.1). *Let Λ be a ring such that $K^b(\text{proj } \Lambda)$ is a Krull–Schmidt category and Λ does not have a semisimple ring summand, and let \mathcal{C} be an algebraic triangulated category. If \mathcal{C} and $K^b(\text{proj } \Lambda)$ are equivalent as additive categories, then they are equivalent as triangulated categories.*

For example, $K^b(\text{proj } \Lambda)$ is Krull–Schmidt if Λ is a module-finite algebra over a complete Noetherian local ring. We actually see that the possible triangle structure on a given Krull–Schmidt additive category is unique in the sense that the suspensions and the mapping cones are uniquely determined as objects, see Proposition 5.5 for details.

As an application of our classification Theorem 1.4 of [1]-finite triangulated categories, we recover the main result of [Chen et al. 2008] stating that any finite dimensional algebra over an algebraically closed field with derived dimension 0 is piecewise hereditary of Dynkin type.

We also apply Theorem 1.4 to Cohen–Macaulay representation theory. A rich source of [1]-finite triangulated categories is given by CM-finite Iwanaga–Gorenstein algebras [Curtis and Reiner 1981; 1987; Leuschke and Wiegand 2012; Simson 1992; Yoshino 1990], for example, simple singularities and trivial extension algebras of representation-finite hereditary algebras. We consequently obtain the following result, which states that $\underline{\text{CM}}^{\mathbb{Z}} \Lambda$ is triangle equivalent to the derived category of a Dynkin quiver under some mild assumptions.

Corollary 1.6 (Theorem 7.3). *Let k be an algebraically closed field and $\Lambda = \bigoplus_{n \geq 0} \Lambda_n$ be a positively graded CM-finite Iwanaga–Gorenstein algebra such that each Λ_n is finite dimensional over k and Λ_0 has finite global dimension. Then, the stable category $\underline{\text{CM}}^{\mathbb{Z}} \Lambda$ is [1]-finite and therefore, it is triangle equivalent to $D^b(\text{mod } kQ)$ for a disjoint union Q of some Dynkin quivers of type A, D, and E.*

This partially recovers [Kajiura et al. 2007; Buchweitz et al. 2020, 2.2] in a quite different way. Note that our result is more general, but less explicit in the sense that Corollary 1.6 does not give the type of Q from given Λ .

As this application suggests, our classification shows that the “easiest” triangulated categories are very likely to be the derived category of Dynkin quivers, and provides a completely different method (from a direct construction of tilting objects) of giving a triangle equivalence for such categories.

Notations and conventions. We denote by k a field. For a category \mathcal{C} , we denote by $\text{Hom}_{\mathcal{C}}(-, -)$ or simply $\mathcal{C}(-, -)$ the Hom-spaces between the objects and by $J_{\mathcal{C}}(-, -)$ the Jacobson radical of \mathcal{C} . A \mathcal{C} -module is a contravariant functor from \mathcal{C} to the category of abelian groups. A \mathcal{C} -module M is finitely presented if there is an exact sequence

$$\mathcal{C}(-, X) \rightarrow \mathcal{C}(-, Y) \rightarrow M \rightarrow 0$$

for some $X, Y \in \mathcal{C}$. We denote by $\text{mod } \mathcal{C}$ the category of finitely presented \mathcal{C} -modules. If \mathcal{C} is graded by a group G , the category of finitely presented graded functor is denoted by $\text{mod}^G \mathcal{C}$, and its projectives by $\text{proj}^G \mathcal{C}$. The morphism space in $\text{mod}^G \mathcal{C}$ is denoted by $\text{Hom}_{\mathcal{C}}(-, -)_0$ or $\mathcal{C}(-, -)_0$. The category $\text{mod}^G \mathcal{C}$ is endowed with the grade shift functor (g) for each $g \in G$, defined by $M(g) = M$ as an ungraded module and $(M(g)(X))_h = (MX)_{gh}$ for each $X \in \mathcal{C}$.

Similarly, for a k -algebra A , the Jacobson radical of A is denoted by J_A . A module over A means a finitely generated right module. We denote by $\text{mod } A$ (resp. $\text{proj } A$) the category of (projective) A -modules. If A is graded, the category of graded (projective) A -modules is denoted by $\text{mod}^G A$ (resp. $\text{proj}^G A$).

2. Periodicity of syzygies

Let A be a k -algebra. We denote by A^e the enveloping algebra $A^{\text{op}} \otimes_k A$ and by Ω_A (resp. Ω_{A^e}) the syzygy, that is, the kernel of the projective cover in $\text{mod } A$ (resp. $\text{mod } A^e$). In this section, we generalize for our purpose the result of Green, Snashall and Solberg [Green et al. 2003] which relates the periodicity of syzygy of simple A -modules and that of A considered as a bimodule over itself. The following theorem and its proof is a graded and twisted version of [loc. cit., 1.4].

Theorem 2.1. *Let G be an abelian group and A be a finite dimensional, ring-indecomposable, non-semisimple G -graded k -algebra. Assume that $J_A = J_{A_0} \oplus (\bigoplus_{i \neq 0} A_i)$ and that A/J_A is separable over k . Then, the following are equivalent for $a \in G$ and $n > 0$:*

- (1) $\Omega_A^n(A/J_A) \simeq A/J_A(a)$ in $\text{mod}^G A$.
- (2) A is self-injective and there exists a graded algebra automorphism α of A such that $\Omega^n \simeq (-)_\alpha(a)$ as functors on $\text{mod}^G A$.
- (3) There exists a graded algebra automorphism α of A such that $\Omega_{A^e}^n(A) \simeq {}_1 A_\alpha(a)$ in $\text{mod}^G A^e$.

Proof. By the original case, we have that A is self-injective under the assumption (3). Then the implication (3) \Rightarrow (2) follows. Also, (2) \Rightarrow (1) is clear.

It remains to prove (1) implies (3). Note that by our assumption on J_A , it is graded and any simple object in $\text{mod}^G A$ is simple in $\text{mod } A$. Assume (1) holds and set $B = \Omega_{A^e}^n(A)$. This is a projective A -module on each side.

Step 1: $S \otimes_A B$ is simple for all graded simple (right) A -modules S .

Let S be a graded simple A -module. Then, applying $S \otimes_A -$ to the minimal projective resolution $P: \cdots \rightarrow P_i \xrightarrow{d_i} P_{i-1} \rightarrow \cdots \rightarrow P_0$ of A in $\text{mod}^G A^e$ yields the minimal projective resolution of S in $\text{mod}^G A$. Indeed, since A/J_A is separable over k , we have $J_{A^e} = J_A \otimes_k A + A \otimes_k J_A$. Then, $\text{Im } d_i \subset P_{i-1} J_{A^e} = J_A P_{i-1} + P_{i-1} J_A$ by the minimality of P and therefore, $\text{Im}(S \otimes d_i) \subset S \otimes_A P_{i-1} J_A$ by $S \otimes_A J_A P_{i-1} = 0$. This shows $S \otimes_A P$ is minimal. Therefore we have $S \otimes_A B \simeq \Omega_A^n(S)$, which is simple by assumption (1).

It follows by induction that the exact functor $- \otimes_A B$ preserves length.

Step 2: $B \simeq A(a)$ in $\text{mod}^G A$.

Consider the exact sequence $0 \rightarrow J_A \rightarrow A \rightarrow A/J_A \rightarrow 0$ in $\text{mod}^G A$. Applying $- \otimes_A B$ yields $B \rightarrow A/J_A(a) \rightarrow 0$. This shows that the module B contains $A/J_A(a)$ in its top. But since B is a projective (right) A -module having the same length as A by the remark following Step 1, we see that $B \simeq A(a)$ in $\text{mod}^G A$.

Step 3: There exists a graded algebra automorphism α of A such that $\Omega_{A^e}^n(A) \simeq {}_1 A_\alpha(a)$.

By Step 2, there exists a graded algebra endomorphism α of A such that $B \simeq {}_\alpha A_1(a)$ in $\text{mod}^G A^e$. Indeed, fix an isomorphism $\varphi: A(a) \rightarrow B$ in $\text{mod}^G A$, put $x = \varphi(1)$, and set $\alpha(u) = \varphi^{-1}(ux)$ for $u \in A$. Then, α is of degree 0, since x and φ are, and it is easily checked that α is an algebra endomorphism

and that $\varphi: {}_{\alpha}A_1(a) \rightarrow B$ is an isomorphism in $\text{mod}^G A^e$. Now we show that α is an isomorphism. Let I be the kernel of α . Since $B \simeq {}_{\alpha}A$ is a projective left A -module, the inclusion $I \subset A$ in $\text{mod}^G A$ stays injective by applying $-\otimes_A \alpha A$. But since the map $I \otimes_A \alpha A \rightarrow {}_{\alpha}A$ is zero, we have $I \otimes_A \alpha A = 0$, and we conclude that $I = 0$ by the remark following Step 1.

This finishes the proof of (1) \Rightarrow (3). \square

We need the following two particular cases. The first one, which we will use in Section 3 is the following result for $G = \{1\}$, where the special case “ $\Omega^n(S) \simeq S$ for all simples” is [Green et al. 2003, 1.4].

Corollary 2.2. *Let A be a ring-indecomposable, nonsemisimple finite dimensional k -algebra such that A/J_A is separable over k . Then, the following are equivalent for $n > 0$:*

- (1) $\Omega_A^n(A/J_A) \simeq A/J_A$.
- (2) A is self-injective and there exists an automorphism α of A such that $\Omega^n \simeq (-)_{\alpha}$ as functors on $\text{mod } A$.
- (3) There exists an automorphism α of A such that $\Omega_{A^e}^n(A) \simeq {}_1A_{\alpha}$ in $\text{mod } A^e$.

We name such algebras as follows:

Definition 2.3. A finite dimensional algebra is *twisted n -periodic* if it is a direct product of simple algebras or algebras satisfying the equivalent conditions in Corollary 2.2.

The second one is the following for $G = \mathbb{Z}$ and the permutation of simples is the identity, which will be used in Section 6.

Corollary 2.4. *Let A be a finite dimensional, ring-indecomposable, nonsemisimple \mathbb{Z} -graded k -algebra such that A/J_A is separable over k . Then, the following are equivalent for $a \in \mathbb{Z}$ and $n > 0$:*

- (1) $\Omega_A^n(S) \simeq S(a)$ in $\text{mod}^{\mathbb{Z}} A$ for any simple objects in $\text{mod}^{\mathbb{Z}} A$.
- (2) A is self-injective and there exists a graded algebra automorphism α of A such that $\Omega^n \simeq (-)_{\alpha}(a)$ as functors on $\text{mod}^{\mathbb{Z}} A$ and $P_{\alpha} \simeq P$ in $\text{mod}^{\mathbb{Z}} A$ for all $P \in \text{proj}^{\mathbb{Z}} A$.
- (3) There exists a graded algebra automorphism α of A such that $\Omega_{A^e}^n(A) \simeq {}_1A_{\alpha}(a)$ in $\text{mod}^{\mathbb{Z}} A^e$ and $P_{\alpha} \simeq P$ in $\text{mod}^{\mathbb{Z}} A$ for all $P \in \text{proj}^{\mathbb{Z}} A$.

Similarly, we name these algebras as follows:

Definition 2.5. A finite dimensional graded algebra is *(a)-twisted n -periodic* if it is a direct product of simple algebras or algebras satisfying the equivalent conditions in Corollary 2.4.

3. Auslander correspondence

We now prove the first main result Theorem 1.2 of this paper, which gives a homological characterization of the Auslander algebras of finite triangulated categories.

First, we give the properties of the endomorphism algebra of a basic additive generator for a finite triangulated category, proving Theorem 1.2(1) \Rightarrow (2).

Proposition 3.1. *Let \mathcal{T} be a k -linear, Hom-finite idempotent-complete triangulated category. Assume \mathcal{T} has an additive generator M . Take M to be basic and set $C = \text{End}_{\mathcal{T}}(M)$. Let α be the automorphism of C induced by $[1]$; precisely, fix an isomorphism $a : M \rightarrow M[1]$ and define α by $\alpha(f) = a^{-1} \circ f[1] \circ a$ for $f \in \text{End}_{\mathcal{T}}(M)$. Then, C is a finite dimensional algebra which is twisted 3-periodic.*

Proof. Since $\text{mod } \mathcal{T} \simeq \text{mod } C$ and $\text{mod } \mathcal{T}$ is a Frobenius category (see [Krause 2007, 4.2]), C is self-injective. Also, since the triangles in \mathcal{T} yield projective resolutions of C -modules, the third syzygy is induced by the automorphism α , that is, we have $\Omega^3 \simeq (-)_{\alpha}$ on $\text{mod } C$. Then C is twisted 3-periodic by Corollary 2.2. \square

For the converse implication, we need the following result due to Amiot, which allows one to introduce a triangle structure on the category of projectives in a Frobenius category.

Proposition 3.2 [Amiot 2007, 8.1]. *Let \mathcal{P} be an idempotent complete k -linear category such that the functor category $\text{mod } \mathcal{P}$ is naturally a Frobenius category. Let S be an autoequivalence of \mathcal{P} and extend this to $\text{mod } \mathcal{P} \rightarrow \text{mod } \mathcal{P}$. Assume there exists an exact sequence of exact functors from $\text{mod } \mathcal{P}$ to $\text{mod } \mathcal{P}$*

$$0 \rightarrow 1 \rightarrow X^0 \rightarrow X^1 \rightarrow X^2 \rightarrow S \rightarrow 0,$$

where X^i take values in $\mathcal{P} = \text{proj } \mathcal{P}$. Then, \mathcal{P} has a structure of a triangulated category with suspension S . The triangles are ones isomorphic to $X^0 M \rightarrow X^1 M \rightarrow X^2 M \rightarrow SX^0 M$ for $M \in \text{mod } \mathcal{P}$.

Combining this with Corollary 2.2, we can prove Theorem 1.2(2) \Rightarrow (1). Let us summarize the proof below.

Proof of Theorem 1.2. (1) \Rightarrow (2) is Proposition 3.1.

(2) \Rightarrow (1) Since A is self-injective, Ω^3 permutes the simples, so by Corollary 2.2, there exists an exact sequence

$$0 \rightarrow A \rightarrow P^0 \rightarrow P^1 \rightarrow P^2 \rightarrow {}_1A_{\alpha} \rightarrow 0$$

of (A, A) -bimodules, with P^i 's projective and α is an automorphism of A . Then, we can apply Proposition 3.2 for $\mathcal{P} = \text{proj } A$, $S = - \otimes_A A_{\alpha}$, and $X^i = - \otimes_A P^i$. \square

Applying a recent result of Keller [2018], we can formulate Theorem 1.2 in terms of bijection between triangulated categories and algebras under some assumptions on triangulated categories. Let us recall the relevant definitions. Let \mathcal{T} be a k -linear triangulated category with Auslander–Reiten triangles, and Γ its AR-quiver. Then Γ together with the AR-translation τ forms a translation quiver. For each pair of vertices $x, y \in \Gamma$, we denote by $\{x \rightarrow y\}$ the set of arrows from x to y . Fix a bijection $\sigma : \{y \rightarrow x\} \rightarrow \{\tau x \rightarrow y\}$, and define $m_x = \sum_{a \in \{y \rightarrow x\}} \sigma(a)a$ which is a morphism in the path category $k\Gamma$. Let I be the ideal of $k\Gamma$ generated by $\{m_x \mid x \in \Gamma\}$.

Definition 3.3 [Riedtmann 1980; Happel 1988]. In the above setting, we call the category $k\Gamma/I$ the *mesh category* of the translation quiver Γ . We say that \mathcal{T} is *standard* if it is k -linearly equivalent its mesh category of the AR-quiver.

We have the following version of Theorem 1.2 under the standardness of \mathcal{T} .

Theorem 3.4. *Let k be an algebraically closed field. Then, there exists a bijection between the following:*

- (1) *The set of triangle equivalence classes of k -linear, Hom-finite, idempotent-complete triangulated categories which are finite, algebraic, and standard.*
- (2) *The set of isomorphism classes of finite dimensional mesh algebras over k .*

The correspondence from (1) to (2) is given by taking the basic Auslander algebra, and from (2) to (1) by taking the category of projective modules.

Proof. We first check that each map is well-defined.

Let \mathcal{T} be a triangulated category as in (1). Then, the standardness of \mathcal{T} implies that its basic Auslander algebra is a mesh algebra.

Suppose next that A is a finite dimensional mesh algebra. We want to show that $\text{proj } A$ has the unique structure of an algebraic triangulated category up to equivalence. Since the third syzygy of simple A -modules are simple, $\mathcal{T} = \text{proj } A$ has a structure of a triangulated category by Theorem 1.2. Also, this is standard since A is a mesh algebra. We claim that $\text{proj } A$ admits a triangle structure which is algebraic. Since \mathcal{T} is a finite, standard triangulated category, there exists a Dynkin quiver Q , a k -linear automorphism F of $D^b(\text{mod } kQ)$, and a k -linear equivalence $D^b(\text{mod } kQ)/F \simeq \text{proj } A$ [Riedtmann 1980]. As in the proof of [Keller 2018], F is isomorphic to $-\otimes_k^L X$ for some (kQ, kQ) -bimodule complex X . Then by [Keller 2005], $D^b(\text{mod } kQ)/F$ admits an algebraic triangle structure as a triangulated orbit category, hence so does $\text{proj } A$. This finishes the proof of the claim. Now, this algebraic triangle structure is unique up to equivalence by the main result of [Keller 2018]. This shows the well-definedness.

It is clear that these maps are mutually inverse. □

Remark 3.5. One can show that using the main result of [Muro 2020], the assumption “standardness” can be dropped.

4. Graded projectivization

In this section, we formulate the method of realizing certain additive categories, which we call G -finite additive categories on which a group G acts with some finiteness conditions, as the category of graded projective modules over a G -graded algebra. This generalizes the classical “projectivization” [Auslander et al. 1995, II.2], which realizes a finite additive category as the category of projectives over an algebra.

Let \mathcal{A} be an additive category with an action of a group G . Precisely, an automorphism F_g of \mathcal{A} is given for each $g \in G$ so that $F_{gh} = F_h \circ F_g$ for all $g, h \in G$. Then the action of G extends to an automorphism of $\text{mod } \mathcal{A}$ by $F_g M = M \circ F_g^{-1}$. For example, the action on the representable functors is $F_g \mathcal{A}(-, X) = \mathcal{A}(-, F_g X)$.

Recall that the orbit category \mathcal{A}/G has the same objects as \mathcal{A} and the morphism space

$$(\mathcal{A}/G)(X, Y) = \bigoplus_{g \in G} \mathcal{A}(X, F_g Y)$$

and the composition $b \circ a$ of $a \in \mathcal{T}(X, F_g Y)$ and $b \in \mathcal{T}(Y, F_h Z)$ is given by $b \circ a = F_g(b)a$, where the right hand side is the composition in \mathcal{A} . Then, \mathcal{A}/G is naturally a G -graded category whose degree g part is $\mathcal{A}(X, F_g Y)$.

Proposition 4.1. *Let \mathcal{A} be an additive category with an action of a group G . Consider the orbit category $\mathcal{C} = \mathcal{A}/G$. Then, the following assertions hold:*

- (1) *The Yoneda embedding $\mathcal{A} \rightarrow \text{proj}^G \mathcal{C}$ is fully faithful. It is an equivalence if \mathcal{A} is idempotent-complete.*
- (2) *There exists an equivalence $\text{mod } \mathcal{A} \simeq \text{mod}^G \mathcal{C}$ such that the action of F_g on $\text{mod } \mathcal{A}$ corresponds to the grade shift (g) on $\text{mod}^G \mathcal{C}$, that is, we have the following commutative diagram of functors:*

$$\begin{array}{ccc}
 \text{mod } \mathcal{A} & \xrightarrow{\simeq} & \text{mod}^G \mathcal{C} \\
 F_g \downarrow & & \downarrow (g) \\
 \text{mod } \mathcal{A} & \xrightarrow{\simeq} & \text{mod}^G \mathcal{C}
 \end{array}$$

Proof. (1) We have the Yoneda lemma for graded functors: $\text{Hom}_{\text{mod}^G \mathcal{C}}(\mathcal{C}(-, X), M) = (MX)_0$. It follows that the Yoneda embedding $\mathcal{A} \rightarrow \text{proj}^G \mathcal{C}$ is fully faithful. Also, if \mathcal{A} is idempotent-complete, the projectives in $\text{mod}^G \mathcal{C}$ are representable, and therefore the Yoneda embedding is dense.

(2) It is clear that the functor in (1) induces an equivalence $\text{mod } \mathcal{A} \simeq \text{mod}^G \mathcal{C}$. Also, the degree h part of the functor $\mathcal{C}(-, F_g X)$ is $\mathcal{A}(-, F_h F_g X) = \mathcal{A}(-, F_{gh} X)$, which is equal to the same degree part of $\mathcal{C}(-, X)(g)$. Thus we have the commutative diagram. □

Now we impose the following finiteness conditions on the G -action:

- (G1) There is $M \in \mathcal{A}$ such that $\mathcal{A} = \text{add}\{F_g M \mid g \in G\}$.
- (G2) For any $X, Y \in \mathcal{A}$, $\text{Hom}_{\mathcal{A}}(X, F_g Y) = 0$ for almost all $g \in G$.

If these conditions are satisfied, we say that an additive category \mathcal{A} with an action of G is G -finite. If \mathcal{A} is a G -finite additive category, we say $M \in \mathcal{A}$ as in (G1) is a G -additive generator. If G is generated by a single element F , we use the term F -finite for G -finiteness, and F -additive generator for G -additive generator. Note that if G is the trivial group, G -finiteness is nothing but finiteness, and a G -additive generator is an additive generator.

Let us reformulate Proposition 4.1 in terms of the graded endomorphism algebra below. Note that this generalizes the classical “projectivization” for finite additive categories, which is the case G is trivial, to “graded projectivization” for G -finite categories. Although this is rather formal, it will be useful in the sequel.

Proposition 4.2. *Let \mathcal{A} be a k -linear, Hom-finite, idempotent-complete category with an action of G , which is G -finite. Let $M \in \mathcal{A}$ be a G -additive generator and set $C = \text{End}_{\mathcal{A}/G}(M)$. Then, the following assertions hold:*

- (1) C is a finite dimensional G -graded algebra.
- (2) The functor $\mathcal{A} \rightarrow \text{proj}^G C$, $X \mapsto \bigoplus_{g \in G} \text{Hom}_{\mathcal{A}}(M, F_g X)$ is an equivalence.
- (3) There exists an equivalence $\text{mod } \mathcal{A} \simeq \text{mod}^G C$ such that the action of g on $\text{mod } \mathcal{A}$ corresponds to the grade shift (g) on $\text{mod}^G C$.

Proof. (1) C is finite dimensional by (G2).

(2) Since we have an equivalence $\text{proj}^G \mathcal{A}/G \rightarrow \text{proj}^G C$ by substituting M , the assertion follows from Proposition 4.1(1).

(3) This is the same as Proposition 4.1(2). □

Definition 4.3. G -Graded rings A and B are *graded Morita equivalent* if there is an equivalence $\text{mod}^G A \simeq \text{mod}^G B$ which commutes with grade shift functors (g) for all $g \in G$.

Let us note the following remark.

Proposition 4.4. Assume (G1) is satisfied and set $C = \text{End}_{\mathcal{A}/G}(M)$.

- (1) The ungraded algebra C does not depend on the choice of M up to Morita equivalence.
- (2) The graded algebra C does not depend on the choice of M up to graded Morita equivalence.

Proof. (1) Since C is the endomorphism algebra of an additive generator of the category \mathcal{A}/G , the assertion follows.

(2) This follows from Proposition 4.2(3). □

As a direct application of this graded projectivization, we present as an example the following graded version of the Auslander correspondence. For simplicity, we consider \mathbb{Z} -graded algebras. A graded algebra Λ is *representation-finite* if $\text{mod}^{\mathbb{Z}} \Lambda$ has finitely many indecomposables up to grade shift. This is equivalent to the representation-finiteness of the ungraded algebra Λ [Gordon and Green 1982].

Proposition 4.5. There exists a bijection between the following:

- (1) The set of graded Morita equivalence classes of finite dimensional \mathbb{Z} -graded algebras Λ of finite representation type.
- (2) The set of graded Morita equivalence classes of finite dimensional \mathbb{Z} -graded algebras Γ with $\text{gl. dim } \Gamma \leq 2 \leq \text{dom. dim } \Gamma$.

The correspondence is given as follows:

- From (1) to (2): $\Gamma = \text{End}_{\Lambda}(M) = \bigoplus_{n \in \mathbb{Z}} \text{Hom}_{\Lambda}(M, M(n))_0$ for a (1)-additive generator M for $\text{mod}^{\mathbb{Z}} \Lambda$.
- From (2) to (1): $\Lambda = \text{End}_{\Gamma}(Q) = \bigoplus_{n \in \mathbb{Z}} \text{Hom}_{\Gamma}(Q, Q(n))_0$ for a (1)-additive generator Q for the category of graded projective-injective Γ -modules.

Proof. Note that Γ (resp. Λ) does not depend on the choice of M (resp. Q) by Proposition 4.4(2). The rest of the proof follows by the same argument as in Theorem 1.1; see [Auslander et al. 1995, VI.5]. □

Notice that this correspondence $\Lambda \leftrightarrow \Gamma$ is the same as the ungraded case, thus it is a refinement of Theorem 1.1 on how much grading Λ or Γ have up to graded Morita equivalence.

5. Uniqueness of triangle structures

The aim of this section is to prove some results which state the uniqueness of triangle structures on certain additive categories. We say that an additive category \mathcal{C} has a unique algebraic triangle structure up to equivalence if $\mathcal{C}_1 = (\mathcal{C}, [1], \Delta)$ and $\mathcal{C}_2 = (\mathcal{C}, [1]', \Delta')$ are algebraic triangle structures on \mathcal{C} , then there exists a triangle equivalence $F: \mathcal{C}_1 \xrightarrow{\sim} \mathcal{C}_2$ such that $F(X) \simeq X$ in \mathcal{C} for all $X \in \mathcal{C}$.

The following is the main result of this section.

Theorem 5.1. *Let Λ be a ring with no simple ring summands such that $\mathbb{K}^b(\text{proj } \Lambda)$ is Krull–Schmidt. Then, the additive category $\mathbb{K}^b(\text{proj } \Lambda)$ has a unique algebraic triangle structure up to equivalence.*

We give applications of Theorem 5.1. For a quiver Q , let $\mathbb{Z}Q$ be the associated infinite translation quiver [Assem et al. 2006; Happel 1988], and let $k(\mathbb{Z}Q)$ be its mesh category [Happel 1988].

Corollary 5.2. *Let Q be a disjoint union of Dynkin quivers which does not contain A_1 . Then, the mesh category $k(\mathbb{Z}Q)$ has a unique algebraic triangle structure up to equivalence.*

As a consequence, we have the classification of [1]-finite algebraic triangulated categories.

Theorem 5.3. *Let k be an algebraically closed field. Any [1]-finite algebraic triangulated category over k is triangle equivalent to the bounded derived category $\text{D}^b(\text{mod } kQ)$ of the path algebra kQ for a disjoint union Q of Dynkin quivers of type A , D , and E .*

Now we start the preparations for the proofs of the above results. Recall that an additive category is Krull–Schmidt if any object is a finite direct sum of objects whose endomorphism rings are local. This is the case if the category is idempotent-complete and Hom-finite over a complete Noetherian local ring. A Krull–Schmidt category \mathcal{C} is purely nonsemisimple if for each $X \in \mathcal{C}$, $J_{\mathcal{C}}(-, X) \neq 0$ or $J_{\mathcal{C}}(X, -) \neq 0$ holds. Note that these conditions are equivalent if \mathcal{C} is triangulated.

First we observe that the suspension and the terms appearing in triangles in a triangulated category are determined by its additive structure under some Krull–Schmidt assumptions. Recall from [Auslander et al. 1995, I.2] that a morphism $f: X \rightarrow Y$ in a Krull–Schmidt category is right minimal if for any direct summand X' of X , the restriction $f|_{X'}$ is nonzero. We dually define left minimality.

Lemma 5.4. *Let \mathcal{C} be a Krull–Schmidt additive category. Assume \mathcal{C} has a structure of a triangulated category. Let $f: X \rightarrow Y$ be a right minimal morphism in $J_{\mathcal{C}}$:*

- (1) *The mapping cone of f is the minimal weak cokernel of f .*
- (2) *$X[1]$ is the minimal weak cokernel of the minimal weak cokernel of f .*

Proof. Complete f to a triangle $X \xrightarrow{f} Y \xrightarrow{g} Z \xrightarrow{h} X[1]$.

(1) We have to show that g is the minimal weak cokernel of f . We only have to show the left minimality of g . If this is not the case, then h has a summand $W \xrightarrow{1_W} W$ for a common nonzero summand W of Z and $X[1]$. This contradicts the right minimality of f .

(2) We want to show that h is the minimal weak cokernel of g . Again, we only have to show the left minimality of h . If this is not the case, then $f[1]$ has a summand $V \xrightarrow{1_V} V$ for a common nonzero summand V of $X[1]$ and $Y[1]$. This contradicts $f \in J_{\mathcal{C}}(X, Y)$. \square

We deduce that the possible triangle structures on a given purely nonsemisimple Krull–Schmidt additive category is roughly unique in the following sense. We denote by $\text{cone}_{\Delta}(f)$ the mapping cone of f in a triangle structure Δ .

Proposition 5.5. *Let \mathcal{C} be a purely nonsemisimple Krull–Schmidt additive category. If $(\mathcal{C}, [1], \Delta)$ and $(\mathcal{C}, [1]', \Delta')$ are triangle structures on \mathcal{C} , then we have the following:*

- (1) $X[1] \simeq X[1]'$ for all objects $X \in \mathcal{C}$.
- (2) $\text{cone}_{\Delta}(f) \simeq \text{cone}_{\Delta'}(f)$ in \mathcal{C} for all morphisms f in \mathcal{C} .

Proof. (1) Let $X \in \mathcal{C}$ be an indecomposable object. Since \mathcal{C} is purely nonsemisimple, there exists a nonzero morphism $f: X \rightarrow Y$ in $J_{\mathcal{C}}$. Then, f is a right minimal radical map, and hence the assertion follows from Lemma 5.4(2).

(2) Let $f: X \rightarrow Y$ be an arbitrary morphism in \mathcal{C} . By removing the summands isomorphic to $W \xrightarrow{1} W$, which does not affect the mapping cone, we may assume $f \in J_{\mathcal{C}}$. Then, f has a decomposition $X_1 \oplus X_2 \xrightarrow{(f_1, 0)} Y$ with right minimal $f_1 \in J_{\mathcal{C}}$ and the mapping cone of f is the direct sum of that of f_1 and $X_2[1]$. Now the mapping cone of f_1 is determined by Lemma 5.4(1) and since \mathcal{C} is purely nonsemisimple, $[1]$ is determined by the additive structure by (1). This proves the assertion. \square

For Theorem 5.1, we need the following result of Keller on algebraic triangulated categories.

Proposition 5.6 [Keller 1994, 4.3]. *Let \mathcal{T} be an algebraic triangulated category and $T \in \mathcal{T}$ be a tilting object. Then, there exists a triangle equivalence $\mathcal{T} \simeq \mathcal{K}^b(\text{proj End}_{\mathcal{T}}(T))$.*

Note that we have the following observation, which will be crucial for the proof.

Lemma 5.7. *Let \mathcal{C} be a purely nonsemisimple Krull–Schmidt additive category. Assume $\mathcal{C}_1 = (\mathcal{C}, [1], \Delta)$ and $\mathcal{C}_2 = (\mathcal{C}, [1]', \Delta')$ are triangle structures on \mathcal{C} . Then, an object $T \in \mathcal{C}$ is a tilting object in \mathcal{C}_1 if and only if it is a tilting object in \mathcal{C}_2 .*

Proof. Indeed, we have $\mathcal{C}_1(T, T[n]) = \mathcal{C}_2(T, T[n]')$ by Proposition 5.5(1), which shows that the vanishing of extensions does not depend on the triangle structure. Also, by Proposition 5.5(2), T generates \mathcal{C}_1 if and only if T generates \mathcal{C}_2 . This shows the assertion. \square

Now we are ready to prove our results.

Proof of Theorem 5.1. Let \mathcal{C} be the underlying additive category of $\mathcal{K} = \mathbb{K}^b(\text{proj } \Lambda)$. Assume \mathcal{C} is triangulated. We show that \mathcal{C} is triangle equivalent to \mathcal{K} by finding a tilting object whose endomorphism ring is Λ . Note that $\mathcal{C} = \mathcal{K}$ is purely nonsemisimple and Krull–Schmidt by our assumption on Λ . Let $T \in \mathcal{C}$ be the object corresponding to $\Lambda \in \mathcal{K}$. Then, T is a tilting object by Lemma 5.7 and clearly $\text{End}_{\mathcal{C}}(T) = \Lambda$. By our assumption that \mathcal{C} is algebraic, we deduce that \mathcal{C} is triangle equivalent to \mathcal{K} by Proposition 5.6. \square

For the proof of Corollary 5.2, let us recall the following standardness theorem of Riedtmann.

Proposition 5.8 [Riedtmann 1980]. *Let k be a field and \mathcal{T} be a k -linear, Hom-finite idempotent-complete triangulated category whose AR-quiver is $\mathbb{Z}Q$ for some acyclic quiver Q . Assume the endomorphism algebra of an indecomposable object of \mathcal{T} is k . Then, \mathcal{T} is k -linearly equivalent to the mesh category $k(\mathbb{Z}Q)$.*

A well known application of this result is an equivalence $\mathbb{K}^b(\text{proj } kQ) \simeq k(\mathbb{Z}Q)$ for a Dynkin quiver Q [Happel 1988, I.5.6].

Proof of Corollary 5.2. Since $k(\mathbb{Z}Q) \simeq \mathbb{K}^b(\text{proj } kQ)$ as additive categories, Theorem 5.1 gives the result. \square

A k -linear triangulated category \mathcal{T} is *locally finite* [Xiao and Zhu 2005] if for each indecomposable $X \in \mathcal{T}$, we have $\sum_{Y:\text{indec.}} \dim_k \text{Hom}_{\mathcal{T}}(X, Y) < \infty$. This condition is equivalent to its dual [loc. cit.]. Clearly, our [1]-finite triangulated categories are locally finite. The classification of [1]-finite triangulated category depends on the following result.

Proposition 5.9 [Xiao and Zhu 2005, 2.3.5]. *Let k be an algebraically closed field and \mathcal{T} be a locally finite triangulated category which does not contain a nonzero finite triangulated subcategory. Then, the AR-quiver of \mathcal{T} is $\mathbb{Z}Q$ for a disjoint union Q of Dynkin quivers of type A , D , and E .*

Proof of Theorem 5.3. The AR-quiver of a [1]-finite triangulated category is $\mathbb{Z}Q$ for some Dynkin quiver Q by Proposition 5.9. Moreover, it is equivalent to $k(\mathbb{Z}Q)$ by Proposition 5.8. Thus Corollary 5.2 applies. \square

We end this section by noting the following lemma, which we use later. This lemma states in particular, that for mesh categories, the suspension is unique up to isomorphism of functors

Lemma 5.10. *Let Q be a Dynkin quiver and α be an automorphism of the mesh category $k(\mathbb{Z}Q)$ such that $\alpha X \simeq X$ for all $X \in k(\mathbb{Z}Q)$. Then, α is isomorphic as functors to the identity functor.*

Proof. Since Q is Dynkin, we can inductively construct a natural isomorphism between α and id . \square

6. [1]-Auslander correspondence

In this section, we prove the second main result, Theorem 1.3, of this paper. In the first subsection, we give the correspondence from triangulated categories to algebras, and the converse one in the second subsection. We will prove the main theorem in the final subsection.

6A. From triangulated categories to algebras. We apply the graded projectivization prepared in Section 4 to triangulated categories. Let \mathcal{T} be a k -linear, Hom-finite, idempotent-complete triangulated category. Consider the action on \mathcal{T} of $G = \mathbb{Z}$, generated by the suspension $[1]$. Then, the G -finiteness in this case are:

(S1) There is $M \in \mathcal{T}$ such that $\mathcal{T} = \text{add}\{M[n] \mid n \in \mathbb{Z}\}$.

(S2) For any $X, Y \in \mathcal{T}$, $\text{Hom}_{\mathcal{T}}(X, Y[n]) = 0$ for almost all n .

According to the terminology in Section 4, we say \mathcal{T} is $[1]$ -finite, and call M as in (S1) a $[1]$ -additive generator.

The following proposition gives the correspondence from triangulated categories to algebras.

Proposition 6.1. *Let \mathcal{T} be a k -linear, Hom-finite, idempotent-complete, triangulated category which is $[1]$ -finite. Let $M \in \mathcal{T}$ be a $[1]$ -additive generator and set $C = \text{End}_{\mathcal{T}/[1]}(M)$. Then, C is a finite-dimensional graded self-injective algebra such that $\Omega^3 L \simeq L(-1)$ for any graded C -module L .*

Proof. C is finite dimensional by (S2). Also, since $\text{mod } \mathcal{T} \simeq \text{mod}^{\mathbb{Z}} C$ by Proposition 4.2(2) and $\text{mod } \mathcal{T}$ is Frobenius, C is self-injective. It remains to show the statement on the third syzygy. Let L be a graded C -module and let $Q \rightarrow R \rightarrow L \rightarrow 0$ be a projective presentation of L in $\text{mod}^{\mathbb{Z}} C$. Take the map $X \rightarrow Y$ in \mathcal{T} corresponding to $Q \rightarrow R$ and complete it to a triangle $W \rightarrow X \rightarrow Y \rightarrow W[1]$. Put $P_Z = \bigoplus_{n \in \mathbb{Z}} \text{Hom}_{\mathcal{T}}(M, Z[n])$ for each $Z \in \mathcal{T}$. This is the graded projective C -module corresponding to Z . Note that $P_{Z[1]} = P_Z(1)$, where (1) is the grade shift functor on $\text{mod}^{\mathbb{Z}} C$. The triangle above yields an exact sequence $P_X(-1) \rightarrow P_Y(-1) \rightarrow P_W \rightarrow P_X \rightarrow P_Y \rightarrow P_W(1)$. Since $P_X = Q$ and $P_Y = R$, we see that $\Omega^3 L \simeq L(-1)$. □

Example 6.2. Let Q be a Dynkin quiver and $\mathcal{T} = \text{D}^b(\text{mod } kQ)$. Let M be an additive generator for $\text{mod } kQ$. Then, M is a $[1]$ -additive generator for \mathcal{T} and we have $C = \text{End}_{kQ}(M) \oplus \text{Ext}_{kQ}^1(M, M)$. The degree 0 part of C is the Auslander algebra of $\text{mod } kQ$.

Let Q' be another Dynkin quiver with the same underlying graph Δ as Q . Since kQ and kQ' are derived equivalent, we have $\text{D}^b(\text{mod } kQ') = \mathcal{T}$. Similarly as above, an additive generator M' for $\text{mod } kQ'$ is a $[1]$ -additive generator for \mathcal{T} . The corresponding graded algebra C' is $\text{End}_{kQ'}(M') \oplus \text{Ext}_{kQ'}^1(M', M')$, with the Auslander algebra of $\text{mod } kQ'$ in the degree 0 part.


By Proposition 4.4, C and C' are isomorphic as ungraded algebras (but not as graded algebras). In this way, $C \simeq C'$ contains the Auslander algebras of module categories over Δ for any orientation of Δ .

Let us give a more specific example.


Example 6.3. Let Q be the following Dynkin quiver of type A_3 , and \mathcal{T} be its derived category $\text{D}^b(\text{mod } kQ)$.

$$a \leftarrow b \leftarrow c.$$

Then, the AR-quiver of \mathcal{T} is as follows:



where $1, \dots, 6$ denotes the objects from $\text{mod } kQ$. Take $M = \bigoplus_{i=1}^6 M_i$, where M_i is the indecomposable kQ -module corresponding to the vertex i . Then, $C = \text{End}_{kQ}(M) \oplus \text{Ext}_{kQ}^1(M, M)$. It is easily verified that C is presented by the quiver $\mathbb{Z}A_3/[1]$ and the mesh relations. The quiver of C looks as follows:



where the vertices with the same number are identified, with mesh relations along the dotted lines. The arrows $1 \rightarrow 5$ and $2 \rightarrow 6$ have degree 1 and all the others have degree 0.

Now, let Q' be the quiver obtained by reflecting Q at vertex a :

$$a \rightarrow b \leftarrow c.$$

Fix an equivalence $\text{D}^b(\text{mod } kQ') \simeq \text{D}^b(\text{mod } kQ)$ so that $M' = M_2 \oplus \dots \oplus M_6 \oplus M_1[1]$ is an additive generator for $\text{mod } kQ'$. Then, $C' = \text{End}_{kQ'}(M') \oplus \text{Ext}_{kQ'}^1(M', M')$ is presented by the same quiver with relations as C , with arrows $2 \rightarrow 1$ and $2 \rightarrow 6$ having degree 1 and all the others degree 0. Thus $C \simeq C'$ as ungraded algebras but not as graded algebras.

Nevertheless, C and C' are graded Morita equivalent. Here we give a direct equivalence $\text{mod}^{\mathbb{Z}}C \rightarrow \text{mod}^{\mathbb{Z}}C'$. Let e_i be the idempotent of C corresponding to M_i ($1 \leq i \leq 6$) and set $P = e_2C \oplus \dots \oplus e_6C \oplus e_1C(1)$. Then, we have $\text{End}_C(P) \simeq C'$ as graded algebras and $\text{Hom}_C(P, -)$ gives a desired equivalence.

6B. From algebras to triangulated categories. We can give the converse correspondence as in Section 3. Setting $a = -1$ in the following proposition gives the result.

Proposition 6.4. *Let A be a finite dimensional graded algebra such that A/J_A is separable over k and $\Omega^3 S \simeq S(a)$ for any graded simple module S . Then, $\text{proj}^{\mathbb{Z}}A$ has a structure of a triangulated category. If k is algebraically closed and $a \neq 0$, then the suspension is isomorphic to $(-a)$ and the algebraic triangle structure on $\text{proj}^{\mathbb{Z}}A$ is unique up to equivalence.*

Proof. By Corollary 2.4, A is self-injective and there exists an exact sequence

$$0 \rightarrow A \rightarrow P^0 \rightarrow P^1 \rightarrow P^2 \rightarrow {}_1A_{\alpha}(-a) \rightarrow 0$$

in $\text{mod}^{\mathbb{Z}}A^e$, where P^i , $i = 0, 1, 2$ are projectives, and α is a graded algebra automorphism of A such that $P_{\alpha} \simeq P$ for all $P \in \text{proj}^{\mathbb{Z}}A$. Then, we can apply Proposition 3.2 for $\mathcal{P} = \text{proj}^{\mathbb{Z}}A$, $X^i = - \otimes_A P^i$ and $S = (-)_{\alpha}(-a)$ to see that $\text{proj}^{\mathbb{Z}}A$ is triangulated with suspension $(-)_{\alpha}(-a)$. Now assume k is

algebraically closed and $a \neq 0$. Since we have $\text{Hom}_{\text{proj}^{\mathbb{Z}} A}(X, Y(-na)) = 0$ for almost all $n \in \mathbb{Z}$ for each $X, Y \in \text{proj}^{\mathbb{Z}} A$, the triangulated category $\text{proj}^{\mathbb{Z}} A$ is [1]-finite, and therefore, it is equivalent to the mesh category $k(\mathbb{Z}Q)$ for some Dynkin diagram Q by Propositions 5.9 and 5.8. Then, by changing the triangle structure if necessary, $\text{proj}^{\mathbb{Z}} A$ has a structure of an algebraic triangulated category, which is unique up to equivalence by Corollary 5.2. Also, $(-)_\alpha(-a)$ and $(-a)$ are isomorphic as functors by Lemma 5.10. \square

6C. Proof of Theorem 1.3. Combining the previous results, we can now prove the second main result of this paper.

Proof of Theorem 1.3. For M as in (1), C is as stated in (2) by Proposition 6.1. Also, the graded Morita equivalence class of C does not depend on the choice of M by Proposition 4.4. This shows the well-definedness of (1) to (2).

For the map from (2) to (1), it is well-defined since $\text{proj}^{\mathbb{Z}} C$ has the unique structure of an algebraic triangulated category up to equivalence by Proposition 6.4.

It is easily checked that these maps are mutually inverse.

The bijection between (1) and (3) is Proposition 5.9 and Theorem 5.3. \square

Remark 6.5. The algebra C in Theorem 1.3 satisfies [3] \simeq (1) as functors on $\text{mod}^{\mathbb{Z}} C$ by Proposition 6.1.

7. Applications to Cohen–Macaulay modules

Applying our classification in Theorem 5.3 of [1]-finite triangulated categories, we show that the stable categories $\underline{\text{CM}}^{\mathbb{Z}} \Lambda$ of some CM-finite Iwanaga–Gorenstein algebras, in particular, of (commutative) graded simple singularities are triangle equivalent to the derived categories of Dynkin quivers.

A Noetherian algebra Λ is *Iwanaga–Gorenstein* if $\text{id}_{\Lambda} \Lambda = \text{id}_{\Lambda^{\text{op}}} \Lambda < \infty$. A typical example of Iwanaga–Gorenstein algebra is given by commutative Gorenstein rings of finite Krull dimension. For an Iwanaga–Gorenstein algebra Λ , we have the category

$$\text{CM} \Lambda = \{X \in \text{mod} \Lambda \mid \text{Ext}_{\Lambda}^i(X, \Lambda) = 0 \text{ for all } i > 0\}$$

of *Cohen–Macaulay* Λ -modules. It is naturally a Frobenius category and we have a triangulated category $\underline{\text{CM}} \Lambda$.

Now consider the case Λ is graded: let $\Lambda = \bigoplus_{n \geq 0} \Lambda_n$ is a positively graded Noetherian algebra such that each Λ_n is finite dimensional over a field k . If Λ is a graded Iwanaga–Gorenstein algebra, we similarly have the category

$$\text{CM}^{\mathbb{Z}} \Lambda = \{X \in \text{mod}^{\mathbb{Z}} \Lambda \mid \text{Ext}_{\Lambda}^i(X, \Lambda) = 0 \text{ for all } i > 0\}$$

of graded Cohen–Macaulay modules. It is again Frobenius and hence the stable category $\underline{\text{CM}}^{\mathbb{Z}} \Lambda$ is triangulated. A graded Iwanaga–Gorenstein algebra is *CM-finite* if $\text{CM}^{\mathbb{Z}} \Lambda$ has finitely many indecomposable objects up to grade shift.

We now show that CM-finite Iwanaga–Gorenstein algebras give a large class of examples of [1]-finite triangulated categories.

Proposition 7.1. *Let Λ be a positively graded CM-finite Iwanaga–Gorenstein algebra with $\text{gl. dim } \Lambda_0 < \infty$. Then, the triangulated category $\underline{\text{CM}}^{\mathbb{Z}}\Lambda$ is [1]-finite.*

To prove this, we need an observation for general Noetherian algebras, which is motivated by [Yamaura 2013, 3.5]. Let us fix some notations. We denote by $\text{Ext}_{\Lambda}^i(-, -)_0$ the Ext groups on $\text{mod}^{\mathbb{Z}}\Lambda$. Note that for $M, N \in \text{mod}^{\mathbb{Z}}\Lambda$, the Ext groups on $\text{mod } \Lambda$ are graded k -vector spaces: $\text{Ext}_{\Lambda}^i(M, N) = \bigoplus_{n \in \mathbb{Z}} \text{Ext}_{\Lambda}^i(M, N(n))_0$, ($i \geq 0$). For each $M \in \text{mod}^{\mathbb{Z}}\Lambda$ and $n \in \mathbb{Z}$, we denote by $M_{\geq n}$ the Λ -submodule of M consisting of components of degree $\geq n$.

Lemma 7.2. *Let Λ be a positively graded Noetherian algebra with $\text{gl. dim } \Lambda_0 < \infty$. Then, for any $X, Y \in \text{mod}^{\mathbb{Z}}\Lambda$, we have $\text{Hom}_{\Lambda}(X, \Omega^n Y)_0 = 0$ for sufficiently large n .*

Proof. Take a minimal graded projective resolution of Y : $\cdots \rightarrow P_2 \rightarrow P_1 \rightarrow P_0 \rightarrow Y \rightarrow 0$. We will show that for each $i \in \mathbb{Z}$, $P_n = (P_n)_{\geq i}$ holds for $n \gg 0$. For this, it suffices to show that $P_n = (P_n)_{\geq 1}$ for $Y = Y_{\geq 0}$. Note that the degree 0 part of the minimal projective resolution of Y yields a Λ_0 -projective resolution of Y_0 . By our assumption that $\text{gl. dim } \Lambda_0 < \infty$, we have $(P_n)_0 = 0$, hence $(P_n)_{\geq 1} = P_n$ for sufficiently large n . Now, we have $\text{Hom}_{\Lambda}(X, \Lambda(-n))_0 = \text{Hom}_{\Lambda}(X, \Lambda)_{-n} = 0$ for $n \gg 0$. Indeed, this is certainly true if X is projective. For general X , take a surjection $P \twoheadrightarrow X$ from a projective module P . Then we have an injection $\text{Hom}_{\Lambda}(X, \Lambda) \hookrightarrow \text{Hom}_{\Lambda}(P, \Lambda)$ and our assertion follows from the case X is projective. Therefore, we conclude that $\text{Hom}_{\Lambda}(X, P_n)_0 = 0$, thus $\text{Hom}_{\Lambda}(X, \Omega^{n+1}Y)_0 = 0$ for sufficiently large n . □

Proof of Proposition 7.1. We verify the conditions (S1) and (S2) found in Section 6A.

First we show (S2): $\underline{\text{Hom}}_{\Lambda}(X, \Omega^n Y)_0 = 0$ for almost all n for each $X, Y \in \underline{\text{CM}}^{\mathbb{Z}}(\Lambda)$. The case $n \gg 0$ is done in Lemma 7.2, so it remains to prove the case $n \ll 0$. Since Λ is CM-finite, $\underline{\text{CM}}^{\mathbb{Z}}\Lambda$ has the AR duality, and we have $D \underline{\text{Hom}}(X, \Omega^n Y)_0 \simeq \underline{\text{Hom}}(Y, \Omega^{-n-1} \tau X)_0$, hence the assertion follows from the case of $n \gg 0$.

Next we show (S1): $\underline{\text{CM}}^{\mathbb{Z}}\Lambda$ has only finitely many indecomposables up to suspension. Since Λ is of finite CM type, there exists $0 \neq n \in \mathbb{Z}$ such that $\Omega^n X \simeq X$ up to grade shift for any indecomposable $X \in \underline{\text{CM}}^{\mathbb{Z}}\Lambda$. By (S2), $\Omega^n X$ and X are not actually isomorphic in $\underline{\text{CM}}^{\mathbb{Z}}\Lambda$. Therefore, $\underline{\text{CM}}^{\mathbb{Z}}\Lambda$ has only finitely many indecomposables up to Ω^n , in particular up to Ω^{-1} .

These assertions show that $\underline{\text{CM}}^{\mathbb{Z}}\Lambda$ is [1]-finite. □

As an application of Theorem 5.3, we immediately obtain the following result.

Theorem 7.3. *Let k be algebraically closed and let $\Lambda = \bigoplus_{n \geq 0} \Lambda_n$ is a positively graded Iwanaga–Gorenstein algebra such that each Λ_n is finite dimensional over k . Suppose Λ is CM-finite and $\text{gl. dim } \Lambda_0 < \infty$. Then, the AR-quiver of $\underline{\text{CM}}^{\mathbb{Z}}\Lambda$ is $\mathbb{Z}\Delta$ for a disjoint union Δ of some Dynkin diagrams of type A, D and E. Moreover, $\underline{\text{CM}}^{\mathbb{Z}}\Lambda$ is triangle equivalent to $\text{D}^b(\text{mod } kQ)$ for any orientation Q of Δ .*

Proof. The statement for the AR-quiver follows from Propositions 7.1 and 5.9. The triangle equivalence follows from Proposition 7.1 and Theorem 5.3. □

A well-known class of commutative Gorenstein rings of finite representation type is given by simple singularities. Here we assume that k is algebraically closed of characteristic 0. Then, they are classified up to isomorphism by the Dynkin diagrams for each $d = \dim \Lambda$ and have the form $k[x, y, z_2, \dots, z_d]/(f)$ with

$$(A_n) \quad f = x^2 + y^{n+1} + z_2^2 + \dots + z_d^2, \quad (n \geq 1),$$

$$(D_n) \quad f = x^2y + y^{n-1} + z_2^2 + \dots + z_d^2, \quad (n \geq 4),$$

$$(E_6) \quad f = x^3 + y^4 + z_2^2 + \dots + z_d^2,$$

$$(E_7) \quad f = x^3 + xy^3 + z_2^2 + \dots + z_d^2,$$

$$(E_8) \quad f = x^3 + y^5 + z_2^2 + \dots + z_d^2;$$

see [Leuschke and Wiegand 2012, Chapter 9]. We admit any grading on Λ so that each variable and f are homogeneous of positive degrees. Then, Λ is CM-finite (in the graded sense) since its completion $\hat{\Lambda}$ at the maximal ideal $\Lambda_{>0}$ is CM-finite, that is, $\text{CM } \hat{\Lambda}$ has only finitely many indecomposable objects [Yoshino 1990, Chapter 15].


Corollary 7.4. *Let k be an algebraically closed field of characteristic zero and $\Lambda = k[x, y, z_2, \dots, z_d]/(f)$ with f one of the above. Give a grading on Λ so that each variable and f are homogeneous of positive degrees. Then, the stable category $\underline{\text{CM}}^{\mathbb{Z}} \Lambda$ is triangle equivalent to the derived category $D^b(\text{mod } kQ)$ of the path algebra kQ of a disjoint union Q of Dynkin quivers.*

We give several more examples. First we consider the case Λ is finite dimensional.

Example 7.5. Let

$$\Lambda = \Lambda_n = k[x]/(x^n)$$


with $\deg x = 1$. Then, Λ is a finite dimensional self-injective algebra. In this case we have $\text{CM}^{\mathbb{Z}} \Lambda = \text{mod}^{\mathbb{Z}} \Lambda$. It is of finite representation type with indecomposable Λ -modules Λ_i ($1 \leq i \leq n$), and $\Lambda_0 = k$ has finite global dimension. We can easily compute its AR-quiver (for $n = 4$) to be




where the top of Λ_i is in degree 0. We see that the AR-quiver of $\text{mod}^{\mathbb{Z}} \Lambda$ is $\mathbb{Z}A_{n-1}$. Consequently, we have a triangle equivalence $\text{mod}^{\mathbb{Z}} \Lambda \simeq D^b(\text{mod } kQ)$ for a quiver Q of type A_{n-1} .

The next one is a finite dimensional Iwanaga–Gorenstein algebra.


Example 7.6. Let Λ be the algebra presented by the following quiver with relations:



with $\deg x = \deg y = 1$ and all other arrows having degree 0. Then, it is an Iwanaga–Gorenstein algebra of dimension 1. (In fact, this is the 3-preprojective algebra [Iyama and Oppermann 2013] of its degree 0 part.) We can compute the AR-quiver of $\text{mod}^{\mathbb{Z}} \Lambda$ to be the following:



Here, each module is graded so that its top is concentrated in degree 0, or equivalently, its lowest degree is at 0. We then compute the category $\text{CM}^{\mathbb{Z}} \Lambda$ to be the circled modules and it is verified that the AR-quiver of $\text{CM}^{\mathbb{Z}} \Lambda$ is



We see that this is $\mathbb{Z}A_2$ and consequently $\text{CM}^{\mathbb{Z}} \Lambda \cong \text{D}^b(\text{mod } kQ)$ for a quiver Q of type A_2 .


We consider as a final example a *Gorenstein order*: let $R = k[x_1, \dots, x_d]$ be a polynomial ring. A Noetherian R -algebra Λ is an R -order if it is projective as an R -module. An R -order Λ is *Gorenstein* if $\text{Hom}_R(\Lambda, R)$ is projective as a Λ -module. In this case, Cohen–Macaulay Λ -modules are Λ -modules which are projective as R -modules.

Example 7.7. Let $R = k[x]$ be a graded polynomial ring with $\deg x = 1$ and let

$$\Lambda = \begin{pmatrix} R & R \\ (x^n) & R \end{pmatrix}.$$

This is a Gorenstein R -order of dimension 1. Its indecomposable CM modules up to grade shift are given by the row vectors $M_i = ((x^i) \ R)$ for $0 \leq i \leq n$, and M_0 and M_n are the projectives. We define the

gradings on the M_i so that their top (0 k) is in degree 0. Then, the AR-quiver of $\text{CM}^{\mathbb{Z}}\Lambda$ (for $n = 4$) is computed to be



where the upgoing arrows are natural inclusions, the downgoing arrows are the multiplications by x , and the dotted lines indicate the AR-translations. By deleting the projective vertices, we see that the AR-quiver of $\underline{\text{CM}}^{\mathbb{Z}}\Lambda$ is $\mathbb{Z}A_{n-1}$, and consequently $\underline{\text{CM}}^{\mathbb{Z}}\Lambda \simeq D^b(\text{mod } kQ)$ for a quiver Q of type A_{n-1} .

Acknowledgement

The author is deeply grateful to his supervisor Osamu Iyama for many helpful suggestions and careful instructions.

References

- [Amiot 2007] C. Amiot, “On the structure of triangulated categories with finitely many indecomposables”, *Bull. Soc. Math. France* **135**:3 (2007), 435–474. MR Zbl
- [Angeleri Hugel et al. 2007] L. Angeleri Hugel, D. Happel, and H. Krause (editors), *Handbook of tilting theory*, Lond. Math. Soc. Lect. Note Ser. **332**, Cambridge Univ. Press, 2007. MR Zbl
- [Artin and Zhang 1994] M. Artin and J. J. Zhang, “Noncommutative projective schemes”, *Adv. Math.* **109**:2 (1994), 228–287. MR Zbl
- [Asashiba 2017] H. Asashiba, “A generalization of Gabriel’s Galois covering functors, II: 2-categorical Cohen–Montgomery duality”, *Appl. Categ. Structures* **25**:2 (2017), 155–186. MR Zbl
- [Assem et al. 2006] I. Assem, D. Simson, and A. Skowronski, *Elements of the representation theory of associative algebras, I: Techniques of representation theory*, Lond. Math. Soc. Student Texts **65**, Cambridge Univ. Press, 2006. MR Zbl
- [Auslander 1971] M. Auslander, “Representation dimension of Artin algebras”, lecture notes, Queen Mary College, 1971.
- [Auslander and Reiten 1996] M. Auslander and I. Reiten, “DTr-periodic modules and functors”, pp. 39–50 in *Representation theory of algebras* (Cocoyoc, Mexico, 1994), edited by R. Bautista et al., CMS Conf. Proc. **18**, Amer. Math. Soc., Providence, RI, 1996. MR Zbl
- [Auslander et al. 1995] M. Auslander, I. Reiten, and S. O. Smal, *Representation theory of Artin algebras*, Cambridge Stud. Adv. Math. **36**, Cambridge Univ. Press, 1995. MR Zbl
- [Buchweitz et al. 2020] R.-O. Buchweitz, O. Iyama, and K. Yamaura, “Tilting theory for Gorenstein rings in dimension one”, *Forum Math. Sigma* **8** (2020), Paper No. e36, 37. MR
- [Chen et al. 2008] X.-W. Chen, Y. Ye, and P. Zhang, “Algebras of derived dimension zero”, *Comm. Algebra* **36**:1 (2008), 1–10. MR Zbl
- [Curtis and Reiner 1981] C. W. Curtis and I. Reiner, *Methods of representation theory with applications to finite groups and orders, I*, Wiley, New York, 1981. MR Zbl

- [Curtis and Reiner 1987] C. W. Curtis and I. Reiner, *Methods of representation theory with applications to finite groups and orders, II*, Wiley, New York, 1987. MR Zbl
- [Enomoto 2018] H. Enomoto, “Classifications of exact structures and Cohen–Macaulay-finite algebras”, *Adv. Math.* **335** (2018), 838–877. MR Zbl
- [Erdmann and Skowroński 2008] K. Erdmann and A. Skowroński, “Periodic algebras”, pp. 201–251 in *Trends in representation theory of algebras and related topics*, edited by A. Skowroński, Eur. Math. Soc., Zürich, 2008. MR Zbl
- [Gordon and Green 1982] R. Gordon and E. L. Green, “Representation theory of graded Artin algebras”, *J. Algebra* **76**:1 (1982), 138–152. MR Zbl
- [Green et al. 2003] E. L. Green, N. Snashall, and Ø. Solberg, “The Hochschild cohomology ring of a selfinjective algebra of finite representation type”, *Proc. Amer. Math. Soc.* **131**:11 (2003), 3387–3393. MR Zbl
- [Happel 1988] D. Happel, *Triangulated categories in the representation theory of finite-dimensional algebras*, Lond. Math. Soc. Lect. Note Ser. **119**, Cambridge Univ. Press, 1988. MR Zbl
- [Heller 1968] A. Heller, “Stable homotopy categories”, *Bull. Amer. Math. Soc.* **74** (1968), 28–63. MR Zbl
- [Iyama 2005] O. Iyama, “The relationship between homological properties and representation theoretic realization of Artin algebras”, *Trans. Amer. Math. Soc.* **357**:2 (2005), 709–734. MR Zbl
- [Iyama 2007] O. Iyama, “Auslander correspondence”, *Adv. Math.* **210**:1 (2007), 51–82. MR Zbl
- [Iyama and Oppermann 2013] O. Iyama and S. Oppermann, “Stable categories of higher preprojective algebras”, *Adv. Math.* **244** (2013), 23–68. MR Zbl
- [Kajiura et al. 2007] H. Kajiura, K. Saito, and A. Takahashi, “Matrix factorizations and representations of quivers, II: Type ADE case”, *Adv. Math.* **211**:1 (2007), 327–362. MR Zbl
- [Keller 1994] B. Keller, “Deriving DG categories”, *Ann. Sci. École Norm. Sup. (4)* **27**:1 (1994), 63–102. MR Zbl
- [Keller 2005] B. Keller, “On triangulated orbit categories”, *Doc. Math.* **10** (2005), 551–581. MR Zbl
- [Keller 2006] B. Keller, “On differential graded categories”, pp. 151–190 in *International Congress of Mathematicians, II* (Madrid, 2006), edited by M. Sanz-Solé et al., Eur. Math. Soc., Zürich, 2006. MR Zbl
- [Keller 2018] B. Keller, “A remark on a theorem by Claire Amiot”, *C. R. Math. Acad. Sci. Paris* **356**:10 (2018), 984–986. MR Zbl
- [Krause 2007] H. Krause, “Derived categories, resolutions, and Brown representability”, pp. 101–139 in *Interactions between homotopy theory and algebra* (Chicago, 2004), edited by L. L. Avramov et al., Contemp. Math. **436**, Amer. Math. Soc., Providence, RI, 2007. MR Zbl
- [Leuschke and Wiegand 2012] G. J. Leuschke and R. Wiegand, *Cohen–Macaulay representations*, Math. Surv. Monogr. **181**, Amer. Math. Soc., Providence, RI, 2012. MR Zbl
- [Muro 2020] F. Muro, “Enhanced finite triangulated categories”, *Journal of the Institute of Mathematics of Jussieu* (2020), 1–43.
- [Neeman 2001] A. Neeman, *Triangulated categories*, Ann. of Math. Stud. **148**, Princeton Univ. Press, 2001. MR Zbl
- [Riedtmann 1980] C. Riedtmann, “Algebren, Darstellungsköcher, Überlagerungen und zurück”, *Comment. Math. Helv.* **55**:2 (1980), 199–224. MR Zbl
- [Rouquier 2008] R. Rouquier, “Dimensions of triangulated categories”, *J. K-Theory* **1**:2 (2008), 193–256. MR Zbl
- [Simson 1992] D. Simson, *Linear representations of partially ordered sets and vector space categories*, Algebra Logic Appl. **4**, Gordon and Breach, Montreux, Switzerland, 1992. MR Zbl
- [Xiao and Zhu 2005] J. Xiao and B. Zhu, “Locally finite triangulated categories”, *J. Algebra* **290**:2 (2005), 473–490. MR Zbl
- [Yamaura 2013] K. Yamaura, “Realizing stable categories as derived categories”, *Adv. Math.* **248** (2013), 784–819. MR Zbl
- [Yoshino 1990] Y. Yoshino, *Cohen–Macaulay modules over Cohen–Macaulay rings*, Lond. Math. Soc. Lect. Note Ser. **146**, Cambridge Univ. Press, 1990. MR Zbl
- [Yoshino 2005] Y. Yoshino, “A functorial approach to modules of G-dimension zero”, *Illinois J. Math.* **49**:2 (2005), 345–367. MR Zbl

2058

Norihiro Hanihara

Communicated by Michel Van den Bergh

Received 2018-09-28 Revised 2019-12-01 Accepted 2020-04-23

m17034e@math.nagoya-u.ac.jp

*Graduate School of Mathematics, Nagoya University, Furocho, Chikusa-ku,
Nagoya, Japan*

Algebra & Number Theory

msp.org/ant

EDITORS

MANAGING EDITOR

Bjorn Poonen
Massachusetts Institute of Technology
Cambridge, USA

EDITORIAL BOARD CHAIR

David Eisenbud
University of California
Berkeley, USA

BOARD OF EDITORS

Jason P. Bell	University of Waterloo, Canada	Susan Montgomery	University of Southern California, USA
Bhargav Bhatt	University of Michigan, USA	Martin Olsson	University of California, Berkeley, USA
Richard E. Borcherds	University of California, Berkeley, USA	Raman Parimala	Emory University, USA
Frank Calegari	University of Chicago, USA	Jonathan Pila	University of Oxford, UK
Antoine Chambert-Loir	Université Paris-Diderot, France	Irena Peeva	Cornell University, USA
J-L. Colliot-Thélène	CNRS, Université Paris-Sud, France	Anand Pillay	University of Notre Dame, USA
Brian D. Conrad	Stanford University, USA	Michael Rapoport	Universität Bonn, Germany
Samit Dasgupta	Duke University, USA	Victor Reiner	University of Minnesota, USA
Hélène Esnault	Freie Universität Berlin, Germany	Peter Sarnak	Princeton University, USA
Gavril Farkas	Humboldt Universität zu Berlin, Germany	Michael Singer	North Carolina State University, USA
Sergey Fomin	University of Michigan, USA	Christopher Skinner	Princeton University, USA
Edward Frenkel	University of California, Berkeley, USA	Vasudevan Srinivas	Tata Inst. of Fund. Research, India
Wee Teck Gan	National University of Singapore	Shunsuke Takagi	University of Tokyo, Japan
Andrew Granville	Université de Montréal, Canada	Pham Huu Tiep	University of Arizona, USA
Ben J. Green	University of Oxford, UK	Ravi Vakil	Stanford University, USA
Joseph Gubeladze	San Francisco State University, USA	Michel van den Bergh	Hasselt University, Belgium
Christopher Hacon	University of Utah, USA	Akshay Venkatesh	Institute for Advanced Study, USA
Roger Heath-Brown	Oxford University, UK	Marie-France Vignéras	Université Paris VII, France
János Kollár	Princeton University, USA	Melanie Matchett Wood	University of California, Berkeley, USA
Michael J. Larsen	Indiana University Bloomington, USA	Shou-Wu Zhang	Princeton University, USA
Philippe Michel	École Polytechnique Fédérale de Lausanne		

PRODUCTION

production@msp.org
Silvio Levy, Scientific Editor

See inside back cover or msp.org/ant for submission instructions.

The subscription price for 2020 is US \$415/year for the electronic version, and \$620/year (+\$60, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to MSP.

Algebra & Number Theory (ISSN 1944-7833 electronic, 1937-0652 printed) at Mathematical Sciences Publishers, 798 Evans Hall #3840, c/o University of California, Berkeley, CA 94720-3840 is published continuously online. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices.

ANT peer review and production are managed by EditFLOW® from MSP.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2020 Mathematical Sciences Publishers

Algebra & Number Theory

Volume 14 No. 8 2020

Toroidal orbifolds, destackification, and Kummer blowings up DAN ABRAMOVICH, MICHAEL TEMKIN and JAROSŁAW WŁODARCZYK	2001
Auslander correspondence for triangulated categories NORIHIRO HANIHARA	2037
Supersingular locus of Hilbert modular varieties, arithmetic level raising and Selmer groups YIFENG LIU and YICHAO TIAN	2059
Burch ideals and Burch rings HAILONG DAO, TOSHINORI KOBAYASHI and RYO TAKAHASHI	2121
Sous-groupe de Brauer invariant et obstruction de descente itérée YANG CAO	2151
Most words are geometrically almost uniform MICHAEL JEFFREY LARSEN	2185
On a conjecture of Yui and Zagier YINGKUN LI and TONGHAI YANG	2197
On iterated product sets with shifts, II BRANDON HANSON, OLIVER ROCHE-NEWTON and DMITRII ZHELEZOV	2239
The dimension growth conjecture, polynomial in the degree and without logarithmic factors WOUTER CASTRYCK, RAF CLUCKERS, PHILIP DITTMANN and KIEN HUU NGUYEN	2261



1937-0652(2020)14:8;1-U