

# *Algebra & Number Theory*

Volume 17  
2023  
No. 2

**Some refinements of the Deligne–Illusie theorem**

Piotr Achinger and Junecue Suh





# Some refinements of the Deligne–Illusie theorem

Piotr Achinger and Junecue Suh

We extend the results of Deligne and Illusie on liftings modulo  $p^2$  and decompositions of the de Rham complex in several ways. We show that for a smooth scheme  $X$  over a perfect field  $k$  of characteristic  $p > 0$ , the truncations of the de Rham complex in  $\max(p-1, 2)$  consecutive degrees can be reconstructed as objects of the derived category in terms of its truncation in degrees at most one (or, equivalently, in terms the obstruction class to lifting modulo  $p^2$ ). Consequently, these truncations are decomposable if  $X$  admits a lifting to  $W_2(k)$ , in which case the first nonzero differential in the conjugate spectral sequence appears no earlier than on page  $\max(p, 3)$  (these corollaries have been recently strengthened by Drinfeld, by Bhatt and Lurie, and by Li and Mondal). Without assuming the existence of a lifting, we describe the gerbes of splittings of two-term truncations and the differentials on the second page of the conjugate spectral sequence, answering a question of Katz.

The main technical result used in the case  $p > 2$  belongs purely to homological algebra. It concerns certain commutative differential graded algebras whose cohomology algebra is the exterior algebra, dubbed by us *abstract Koszul complexes*, of which the de Rham complex in characteristic  $p$  is an example.

In the Appendix, we use the aforementioned stronger decomposition result to prove that Kodaira–Akizuki–Nakano vanishing and Hodge–de Rham degeneration both hold for  $F$ -split  $(p+1)$ -folds.

## 1. Introduction

**1A. Decompositions of the de Rham complex.** Deligne and Illusie [1987] showed that for a smooth scheme  $X$  over a perfect field  $k$  of characteristic  $p > 0$ , a flat lifting of the Frobenius twist  $X' = F_k^* X$  to  $W_2(k)$  induces a splitting of the truncation of the de Rham complex in degrees  $[0, 1]$ , i.e., an isomorphism in the derived category

$$\mathcal{O}_{X'} \oplus \Omega_{X'/k}^1[-1] \xrightarrow{\sim} \tau_{\leq 1}(F_{X/k,*}\Omega_{X/k}^\bullet).$$

Using the algebra structure of the de Rham complex, they further show that it induces an isomorphism

$$\bigoplus_{i < p} \Omega_{X'/k}^i[-i] \xrightarrow{\sim} \tau_{< p}(F_{X/k,*}\Omega_{X/k}^\bullet).$$

With their method, it is unclear if one could extend this further to an isomorphism between  $\bigoplus_{i \geq 0} \Omega_{X'/k}^i[-i]$  and  $F_{X/k,*}\Omega_{X/k}^\bullet$  if  $\dim X \geq p$ , i.e., whether the de Rham complex  $\Omega_{X/k}^\bullet$  is *decomposable*. As a step further, Deligne and Illusie prove using duality that this is the case if  $\dim X = p$ .

MSC2020: primary 14F40; secondary 14G17.

Keywords: de Rham cohomology, Koszul complex, Deligne–Illusie, lifting modulo  $p^2$ , conjugate spectral sequence,  $F$ -splitting.

© 2023 MSP (Mathematical Sciences Publishers). Distributed under the Creative Commons Attribution License 4.0 (CC BY). Open Access made possible by subscribing institutions via [Subscribe to Open](#).

It is as of today an open problem whether there exists a smooth  $X$  over  $k$  liftable to  $W_2(k)$ , necessarily of dimension  $\dim X > p$ , for which the de Rham complex is not decomposable.<sup>1</sup> In this paper, as a small contribution to this question, we investigate the ways in which the truncation  $\tau_{\leq 1}(F_{X/k,*}\Omega_{X/k}^\bullet)$  determines the truncations  $\tau_{[a,b]}(F_{X/k,*}\Omega_{X/k}^\bullet)$ . Our first result is the following:

**Theorem 1.1.** *Let  $X$  be a smooth scheme over a perfect field  $k$  of characteristic  $p > 0$  which is liftable to  $W_2(k)$ . Then the truncations*

$$\tau_{[a,b]}(F_{X/k,*}\Omega_{X/k}^\bullet)$$

*are decomposable for  $a \leq b < a + p - 1$  when  $p > 2$ , and for  $a \leq b \leq a + 1$  when  $p = 2$ .*

The above result immediately implies that in the conjugate spectral sequence

$$E_2^{ij} = H^i(X', \Omega_{X'/k}^j) \Rightarrow H^{i+j}(X, \Omega_{X/k}^\bullet) \tag{1-1}$$

the differentials  $d_r^{ij}$  are zero for  $2 \leq r < p$  when  $p > 2$ , and for  $r = 2$  when  $p = 2$ . As a sample corollary, we obtain the following criterion for degeneration of spectral sequences.

**Corollary 1.2.** *For  $X$  as in Theorem 1.1, suppose that*

$$H^i(X, \Omega_{X/k}^j) = 0 \text{ for } |i - j| \geq p.$$

*Then the conjugate spectral sequence (1-1) degenerates. If moreover  $X$  is proper over  $k$ , then the Hodge to de Rham spectral sequence*

$$E_1^{ij} = H^j(X, \Omega_{X/k}^i) \Rightarrow H^{i+j}(X, \Omega_{X/k}^\bullet)$$

*degenerates as well.*

**1B. Truncations of the de Rham complex.** Our methods give information about truncations of the de Rham complex without assuming liftability modulo  $p^2$ . Our results in this direction are the strongest and most explicit for truncations in two consecutive degrees. Namely, for a general smooth  $X$  over  $k$  (not necessarily liftable to  $W_2(k)$ ) and for  $q \geq 1$ , the truncated complex  $\tau_{[q-1,q]}(F_{X/k,*}\Omega_{X/k}^\bullet)$  can be described as the mapping fiber of  $\delta^q[-q]$  for a map

$$\delta^q : \Omega_{X'/k}^q \rightarrow \Omega_{X'/k}^{q-1}[2],$$

that is, a class

$$\delta^q \in \text{Ext}^2(\Omega_{X'/k}^q, \Omega_{X'/k}^{q-1}),$$

which is the ‘‘cup product’’ with the negative of the deformation obstruction class

$$\delta^1 = -\text{obs}(X'/k/W_2(k)) \in \text{Ext}^2(\Omega_{X'/k}^1, \mathcal{O}_{X'}) \simeq H^2(X', T_{X'/k}) \tag{1-2}$$

to the existence of a lifting of  $X'$  to  $W_2(k)$  (see Corollary 4.3). The result in particular implies that the two-term truncation  $\tau_{[q-1,q]}(F_{X/k,*}\Omega_{X/k}^\bullet)$  is decomposable if  $X'$  lifts to  $W_2(k)$ , and yields a description

---

<sup>1</sup>*Added in proof:* This problem has been recently resolved by Petrov [2023], who constructed such a variety.

of the differentials on the second page of the conjugate spectral sequence — answering a natural question of Katz.

**Theorem 1.3** (see Corollary 4.4). *In the above situation, the differential*

$$d_2^{ij} : H^i(X', \Omega_{X'/k}^j) \rightarrow H^{i+2}(X', \Omega_{X'/k}^{j-1})$$

*in the conjugate spectral sequence (1-1) is induced by the cup product with the negative of the obstruction class  $\text{obs}(X'/k/W_2(k))$ .*

Deligne and Illusie [1987, § 3] define the *gerbe of splittings*  $\text{sc } K$  of a two-term complex  $K$ , and relate the gerbe of splittings of  $\tau_{\leq 1}(F_{X/k,*}\Omega_{X/k}^\bullet)$  to the gerbe of liftings of  $X'$  to  $W_2(k)$ . This provides a “categorification” of the equality (1-2). In the same vein, for  $p > 2$ , our description of the class of  $\tau_{[q-1,q]}(F_{X/k,*}\Omega_{X/k}^\bullet)$  can be upgraded to a morphism of gerbes (see Theorem 3.9)

$$\wedge^q : \text{sc}(\tau_{\leq 1}(F_{X/k,*}\Omega_{X/k}^\bullet)) \rightarrow \text{sc}(\tau_{[q-1,q]}(F_{X/k,*}\Omega_{X/k}^\bullet)).$$

Let us now discuss longer truncations of the de Rham complex. The assertion of Theorem 1.1 is subsumed by a recent beautiful observation of Drinfeld [2020, § 5.12.1] (a proof appeared in Bhatt and Lurie [2022]): a lifting of  $X'$  to  $W_2(k)$  induces a  $\mu_p$ -action on the de Rham complex  $F_{X/k,*}\Omega_{X/k}^\bullet$ , which one can use to show that the truncations  $\tau_{[q-p+1,q]}(F_{X/k,*}\Omega_{X/k}^\bullet)$  are decomposable for all  $q$  (even more recently, Li and Mondal [2021] found an independent proof). However, the method of proof of Theorem 1.1 is completely different and provides interesting information even if  $X$  is not liftable to  $W_2(k)$ . It is deduced from the following result (when  $p > 2$ ) and Corollary 4.3 (when  $p = 2$ ) alluded to above.

**Theorem 1.4.** *Let  $X$  be a smooth scheme over a perfect field  $k$  of characteristic  $p > 0$ , let  $q$  be an integer, and let  $m < p - 1$ . One then has an isomorphism in the derived category of  $X'$ ,*

$$\tau_{[q-m,q]}(F_{X/k,*}\Omega_{X/k}^\bullet) \simeq \tau_{\geq q-m}(L\Gamma^q(\tau_{\leq 1}F_{X/k,*}\Omega_{X/k}^\bullet)),$$

where  $L\Gamma^q$  is the derived  $q$ -th divided power.

**1C. Abstract Koszul complexes.** The proof of Theorem 1.4 has very little to do with algebraic geometry. To state the main technical result behind it, we need the notion of an *abstract Koszul complex* (Definition 2.1), which is a certain commutative differential graded algebra (cdga)  $K$  in a ringed topos for which the multiplication induces isomorphisms

$$\wedge^q \mathcal{H}^1(K) \xrightarrow{\sim} \mathcal{H}^q(K) \quad \text{for all } q \geq 0.$$

Thanks to the Cartier isomorphism, the de Rham complexes  $F_{X/k,*}\Omega_{X/k}^\bullet$  in characteristic  $p > 0$  are examples of such, and hence the result below immediately implies Theorem 1.4.

**Theorem 1.5** (see Theorem 2.8). *Let  $K$  be an abstract Koszul complex in a ringed topos  $(X, \mathcal{O})$  satisfying the flatness condition (2-1), and let  $q \geq m \geq 1$  be integers such that  $m!$  is invertible in  $\mathcal{O}$ . Suppose that*

either  $q = m$  or that  $m + 1$  is a nonzero divisor in  $\mathcal{O}$ . Then there exists an isomorphism in the derived category

$$\tau_{\geq q-m}(L\Gamma^q(\tau_{\leq 1}(K))) \xrightarrow{\sim} \tau_{[q-m,q]}(K). \tag{1-3}$$

In (1-3),  $L\Gamma^q$  is again the derived  $q$ -th divided power, and the source of the map can be more concretely realized as  $\tau_{\geq q-m}$  of the Koszul complex

$$\dots \rightarrow \underbrace{\Gamma^{q-i}(K^0) \otimes \wedge^i(\mathcal{Z}^1 K)}_{\text{degree } i} \rightarrow \dots \rightarrow K^0 \otimes \wedge^{q-1}(\mathcal{Z}^1 K) \rightarrow \underbrace{\wedge^q(\mathcal{Z}^1 K)}_{\text{degree } q} \rightarrow 0 \rightarrow \dots$$

For  $m = 1$  (and assuming that 2 is a nonzerodivisor), we again obtain more refined information regarding  $\tau_{[q-1,q]}K$ , including the differentials on the second page of the spectral sequence

$$E_2^{ij} = H^i(X, \mathcal{H}^j(K)) \Rightarrow H^{i+j}(X, K) \quad (\text{see Corollary 4.3}).$$

As observed by Kato [1989], logarithmic de Rham complexes are abstract Koszul complexes, and hence Theorem 1.1 works also in the log case. The inspiration for Theorem 2.8 came from the result of Steenbrink [1995, § 2.8] describing the nearby cycle complex  $R\Psi\mathbb{Q}$  for a complex semistable degeneration in terms of the logarithmic structure; see also [Achinger and Ogus 2020, § 4]. It is an interesting question whether Steenbrink’s result can be extended to work with integral coefficients; the nearby cycles  $R\Psi\mathbb{Z}$  are coconnective  $E_\infty$ -algebra versions of abstract Koszul complexes, but we do not know whether they admit cdga models (see Remark 2.11 and Example 2.3). An affirmative answer would give an application unrelated to the Deligne–Illusie theorem, refining [Achinger and Ogus 2020, Theorem 4.2.2(1)], providing a description of the two-step truncations  $\tau_{[q-1,q]}$  of certain logarithmic nearby cycle complexes.

**1D. The case  $p = 2$  (Theorem 4.1).** The description of the truncations  $\tau_{[q-1,q]}(F_{X/k,*}\Omega_{X/k}^\bullet)$  and its corollary, Theorem 1.3, can be deduced from the “abstract Koszul complex” machinery and Theorem 1.4, but only for  $p > 2$ . In contrast, the assertion of Theorem 1.5 is vacuous if  $2 \cdot \mathcal{O} = 0$ . Accordingly, the computation of the class of  $\tau_{[q-1,q]}F_{X/k,*}\Omega_{X/k}^\bullet$ , occupying the entire Section 4 is much harder in the case  $p = 2$ , and uses more information about the de Rham complex than merely its abstract Koszul complex structure. For this technical point, we highlight the passage from (4-3) to (4-4).

It could be worthwhile to extend the methods used in the case of  $p = 2$  in order to “compute” the truncations  $\tau_{[q-p+1,p]}$  in  $p$  consecutive degrees, and it would be interesting to extract the exact abstract properties of the de Rham complex in positive characteristic needed for the proof. Its relationship with the aforementioned result of Drinfeld, Bhatt–Lurie, and Li–Mondal remains elusive.

The results concerning the truncations  $\tau_{[q-1,q]}(F_{X/k,*}\Omega_{X/k}^\bullet)$  and Theorem 1.3, including the case  $p = 2$ , presented here are due to the second author and appeared in his 2007 Ph.D. thesis. After the first author proved Theorem 1.4, the authors decided to publish their results together.

**1E. Application to  $F$ -split  $(p+1)$ -folds.** As an illustration of this circle of ideas, using the refinement of the Deligne–Illusie theorem due to Drinfeld, Bhatt–Lurie, and Li–Mondal, we prove in the Appendix that

the Kodaira–Akizuki–Nakano vanishing theorem and the degeneration of the Hodge to de Rham spectral sequence both hold for  $F$ -split  $(p+1)$ -folds in characteristic  $p$ .<sup>2</sup>

**Notation.** If  $K$  is a cochain complex in an abelian category, we write  $\mathcal{Z}^i K = \ker(d : K^i \rightarrow K^{i+1})$ ,  $\mathcal{B}^i K = \operatorname{im}(d : K^{i-1} \rightarrow K^i)$ , and  $\mathcal{H}^i(K) = \mathcal{Z}^i K / \mathcal{B}^i K$ , denote by  $\tau_{\leq b}(K)$  the subcomplex

$$\tau_{\leq b}(K) = [\dots \rightarrow K^{b-1} \rightarrow \mathcal{Z}^b K \rightarrow 0 \rightarrow \dots]$$

and by  $\tau_{\geq a}(K)$  the quotient  $K/\tau_{\leq a}(K)$ , and define  $\tau_{[a,b]}(K) = \tau_{\geq a}\tau_{\leq b}(K)$ . We call  $K$  *decomposable* if it is isomorphic in the derived category to the complex with zero differential  $\bigoplus \mathcal{H}^i(K)[-i]$ .

A *commutative differential graded algebra (cdga)* is an associative graded ring  $K = \bigoplus_{n \in \mathbb{Z}} K^n$  which is graded-commutative (i.e.,  $xy = (-1)^{mn}yx$  for  $x \in K^m, y \in K^n$ ), endowed with a differential  $d : K \rightarrow K$  mapping  $K^n$  to  $K^{n+1}$  and satisfying  $d(xy) = dx \cdot y + (-1)^n x \cdot dy$  for  $x \in K^n$ . We say that  $K$  is *coconnective* if  $K^n = 0$  for  $n < 0$ .

## 2. Abstract Koszul complexes

**2A. Definition and examples.** We work in a ringed topos  $(X, \mathcal{O})$ .

**Definition 2.1** (abstract Koszul complex). A coconnective commutative differential graded  $\mathcal{O}$ -algebra  $K$  is called an *abstract Koszul complex* if the following conditions are satisfied:

- (i)  $\mathcal{O} \rightarrow \mathcal{H}^0(K)$  is an isomorphism.
- (ii) For every  $q \geq 1$ , the multiplication map  $\mathcal{H}^1(K)^{\otimes q} \rightarrow \mathcal{H}^q(K)$  factors through an isomorphism

$$\mu^q : \bigwedge^q \mathcal{H}^1(K) \xrightarrow{\sim} \mathcal{H}^q(K).$$

**Example 2.2** (De Rham complex in characteristic  $p > 0$ ). Let  $X$  be a smooth scheme over a perfect field  $k$  of characteristic  $p > 0$ , and let  $F_{X/k} : X \rightarrow X'$  be its relative Frobenius. Let  $K = F_{X/k,*}\Omega_{X/k}^\bullet$  be the de Rham complex, treated as a cdga over  $\mathcal{O}_{X'}$ . Then the Cartier isomorphisms

$$C : \mathcal{H}^i(F_{X/k,*}\Omega_{X/k}^\bullet) \xrightarrow{\sim} \Omega_{X'/k}^i$$

are multiplicative, and hence  $K$  is an abstract Koszul complex over  $(X', \mathcal{O}_{X'})$ .

More generally, if  $f : (X, \mathcal{M}_X) \rightarrow (S, \mathcal{M}_S)$  is a morphism of fine log schemes over  $\mathbb{F}_p$  which is smooth and of Cartier type, then the log de Rham complex  $F_{X/S,*}\Omega_{(X,\mathcal{M}_X)/(S,\mathcal{M}_S)}^\bullet$  is an abstract Koszul complex [Kato 1989, Theorem 4.12].

**Example 2.3** (nearby cycle complexes; see, e.g., [Steenbrink 1995, § 2]). Let  $X$  be a complex manifold and let  $D = \bigcup D_\alpha$  be a divisor with simple normal crossings on  $X$ . Let  $j : U = X \setminus D \hookrightarrow X$  be the complementary open immersion, and let  $K = Rj_*\mathbb{Q}_U$ . Since we are working with rational coefficients,

<sup>2</sup>Added in proof: These results have recently been improved upon by Petrov (private communication, 2023), who showed that the assumption on dimension is not necessary.

we can find a cdga model for  $K$  (e.g., [Kříž and May 1995, Part II, Corollary 1.5]). The purity theorem implies that

$$R^i j_* \mathbb{Q}_U = \wedge^i \left( \bigoplus \mathbb{Q}_{D_\alpha} \right),$$

so any cdga model of  $K$  is an abstract Koszul complex over  $(X, \mathbb{Q}_X)$ . Moreover, one has an isomorphism in the derived category [Steenbrink 1995, Lemma 2.7] (see also [Achinger and Ogus 2020, § 4])

$$\tau_{\leq 1}(R^1 j_* \mathbb{Z}_U) \simeq [\mathcal{O}_X \xrightarrow{\text{exp}} \mathcal{M}^{\text{gp}}],$$

where  $\mathcal{M}^{\text{gp}}$  is the sheaf of meromorphic functions without zeros or poles on  $U$ . Variants of this construction exist for the nearby cycle complexes  $R\Psi\mathbb{Q}$  for a semistable degeneration over a disc, and there exist analogs in  $\ell$ -adic étale cohomology (with  $\mathbb{Q}_\ell$  coefficients).

Recall [SGA 4<sub>2</sub> 1972, exposé V, Definition 1.1] that an  $\mathcal{O}$ -module  $M$  is *flat* if the functor  $(-)\otimes_{\mathcal{O}} M$  is exact on the category of all  $\mathcal{O}$ -modules. By the Deligne–Lazard theorem [SGA 4<sub>2</sub> 1972, exposé V, théorème 8.2.12], an  $\mathcal{O}$ -module is flat if and only if it is a local inductive limit (see [SGA 4<sub>2</sub> 1972, exposé V, § 8.1]) of free  $\mathcal{O}$ -modules of finite rank.

In the following, we will work with abstract Koszul complexes satisfying the additional flatness condition:

$$\text{the } \mathcal{O}\text{-modules } K^0, \mathcal{B}^1 K, \mathcal{Z}^1 K, \text{ and } \mathcal{H}^1(K) \text{ are flat.} \quad (2-1)$$

In particular, this implies that the modules  $\mathcal{H}^q(K) \simeq \wedge^q \mathcal{H}^q(K)$  are flat for all  $q \geq 0$ . The above condition is satisfied in the situation of Examples 2.2 and 2.3.

**2B. Koszul complexes.** Our goal is to show that to a certain extent, the underlying complex of an abstract Koszul complex satisfying the flatness condition (2-1) is determined by its truncation in degrees  $\leq 1$  (Theorem 2.8). We achieve this using the notion of the Koszul complex of a map  $u : P \rightarrow Q$ ; see [Illusie 1971, chapitre I, § 4.3] and [Kato and Saito 2004, § 1.1–1.2].

Recall first that if  $M$  and  $N$  are  $\mathcal{O}$ -modules, then for every  $q \geq 0$  there is a natural decomposition of the divided (resp. exterior) power

$$\Gamma^q(M \oplus N) = \bigoplus_{a+b=q} \Gamma^a M \otimes \Gamma^b N \quad \left( \text{resp. } \wedge^q(M \oplus N) = \bigoplus_{a+b=q} \wedge^a M \otimes \wedge^b N \right).$$

In what follows, we will use the *comultiplication maps*

$$\eta^q : \Gamma^q M \rightarrow (\Gamma^{q-1} M) \otimes M \quad (\text{resp. } \eta^q : \wedge^q M \rightarrow M \otimes \wedge^{q-1} M)$$

obtained as the composition of  $\Gamma^q$  (resp.  $\wedge^q$ ) of the diagonal map  $M \rightarrow M \oplus M$  and the projection to the  $(a, b) = (q-1, 1)$ -part (resp.  $(a, b) = (1, q-1)$ -part) in the above decomposition of  $\Gamma^q(M \oplus M)$



(resp. of  $\wedge^q(M \oplus M)$ ). Explicitly, we have

$$\eta^q(x_1^{[e_1]} \cdots x_r^{[e_r]}) = \sum_{i=1}^r (x_1^{[e_1]} \cdots x_i^{[e_i-1]} \cdots x_r^{[e_r]}) \otimes x_i \quad (e_1 + \cdots + e_r = q),$$

$$\eta^q(x_1 \wedge \cdots \wedge x_q) = \sum_{i=1}^q (-1)^{i-1} x_i \otimes x_1 \wedge \cdots \wedge \widehat{x}_i \wedge \cdots \wedge x_q.$$

Sometimes we omit the superscript  $q$  when it is clear from the context.

**Definition 2.4** (Koszul complex  $\text{Kos}^q(u)$ ). Let  $u : P \rightarrow Q$  be a map of  $\mathcal{O}$ -modules, and let  $q \geq 0$  be an integer. Then the  $q$ -th Koszul complex  $\text{Kos}^q(u)$  is the cochain complex whose  $i$ -th term is

$$\text{Kos}^q(u)^i = \Gamma^{q-i}(P) \otimes \wedge^i(Q),$$

with differential  $d : \text{Kos}^q(u)^i \rightarrow \text{Kos}^q(u)^{i+1}$  defined as the composition

$$\begin{array}{ccc} \Gamma^{q-i}(P) \otimes \wedge^i(Q) & \xrightarrow{\eta \otimes \text{id}} & \Gamma^{q-i-1}(P) \otimes P \otimes \wedge^i(Q) \\ & & \downarrow \text{id} \otimes u \otimes \text{id} \\ & & \Gamma^{q-i-1}(P) \otimes Q \otimes \wedge^i(Q) \xrightarrow{\text{id} \otimes \wedge} \Gamma^{q-i-1}(P) \otimes \wedge^{i+1}(Q) \end{array}$$

Concretely, with  $e_1 + \cdots + e_r = q - i$ ,  $x_1, \dots, x_r \in P$ , and  $y \in \wedge^i Q$ ,

$$d(x_1^{[e_1]} \cdots x_r^{[e_r]} \otimes y) = \sum_{j=1}^r (x_1^{[e_1]} \cdots x_j^{[e_j-1]} \cdots x_r^{[e_r]}) \otimes u(x_j) \wedge y.$$

We note here that our convention differs slightly from that in [Kato and Saito 2004] and [Illusie 1971], who use  $\wedge^i(Q) \otimes \Gamma^{q-i}(P)$  as the  $i$ -th term. The two complexes, ours and theirs, are isomorphic via the map which is  $(-1)^i$  times the map switching the two tensor factors in degree  $i$ . The reason for this convention is that later in Proposition 3.2 we will obtain the left comultiplication maps on exterior powers, which in applications to the de Rham complex will be compatible with interior multiplication of differential forms.

**Proposition 2.5.** Let  $u : P \rightarrow Q$  be a map of flat  $\mathcal{O}$ -modules, and let  $F(u) = [P \xrightarrow{u} Q]$  be the two-term cochain complex with  $P$  in degree zero (the mapping fiber). There exist natural isomorphisms in the derived category

$$\text{Kos}^q(u) \simeq L\Lambda^q(F[1])[-q] \simeq L\Gamma^q(F),$$

where  $L\Lambda^q$  (resp.  $L\Gamma^q$ ) is the derived exterior (resp. divided) power.

*Proof.* Combine [Kato and Saito 2004, Corollary 1.2.7] with [Illusie 1971, chapitre I, 4.3.2.1]. See also [Illusie 1972, chapitre VIII, lemme 2.1.2.1]. □

**Corollary 2.6.** For a map  $u : P \rightarrow Q$  between flat  $\mathcal{O}$ -modules, the complex  $\text{Kos}^q(u)$ , treated as an object of the derived category, depends only on  $F(u) = [P \rightarrow Q]$  up to quasi-isomorphism. In particular, if  $F(u)$  is decomposable, then so is  $\text{Kos}^q(u)$ .

**Proposition 2.7** (compare [Steenbrink 1995, Lemma 1.4]). *Let  $u : P \rightarrow Q$  be a map of  $\mathcal{O}$ -modules. There exist unique arrows*

$$\Gamma^{q-i}(\ker u) \otimes \wedge^i(Q) \rightarrow \mathcal{Z}^i(\mathrm{Kos}^q(u)) \quad \text{and} \quad \alpha^i : \Gamma^{q-i}(\ker u) \otimes \wedge^i(\mathrm{cok} u) \rightarrow \mathcal{H}^i(\mathrm{Kos}^q(u))$$

*making the following diagram commute:*

$$\begin{array}{ccc} \Gamma^{q-i}(P) \otimes \wedge^i(Q) & \xlongequal{\quad} & \mathrm{Kos}^q(u)^i \\ \uparrow & & \uparrow \\ \Gamma^{q-i}(\ker u) \otimes \wedge^i(Q) & \longrightarrow & \mathcal{Z}^i(\mathrm{Kos}^q(u)) \\ \downarrow & & \downarrow \\ \Gamma^{q-i}(\ker u) \otimes \wedge^i(\mathrm{cok} u) & \xrightarrow{\alpha^i} & \mathcal{H}^i(\mathrm{Kos}^q(u)) \end{array}$$

*Moreover, the map  $\alpha^i$  is an isomorphism if  $P$ ,  $Q$ ,  $\ker u$ ,  $\mathrm{im} u$ , and  $\mathrm{cok} u$  are all flat.*

*Proof.* The first assertion is straightforward. The second is reduced as in [Illusie 1971, chapitre I, 4.3.1.6] using the Deligne–Lazard theorem to the case where  $P$ ,  $Q$ ,  $\ker u$ ,  $\mathrm{im} u$ , and  $\mathrm{cok} u$  are free  $\mathcal{O}$ -modules of finite rank. In this case, splitting the surjection  $Q \rightarrow \mathrm{cok} u$  one can write  $u = u' \oplus u''$ , where  $u' : P \rightarrow \mathrm{im} u$  and  $u'' : 0 \rightarrow \mathrm{cok} u$ . The assertion then holds for  $u'$  (by [Illusie 1971, chapitre I, 4.3.1.6]) and for  $u''$  (trivially), for all  $q$ , and then the assertion for  $u = u' \oplus u''$  follows from the isomorphism [Illusie 1971, chapitre I, 4.3.1.5]

$$\mathrm{Kos}^\bullet(u) = \mathrm{Kos}^\bullet(u') \otimes \mathrm{Kos}^\bullet(u''),$$

where  $\mathrm{Kos}^\bullet(u) = \bigoplus_{q \geq 0} \mathrm{Kos}^q(u)[q]$ . □

**2C. Truncations of abstract Koszul complexes.** The following theorem is the main result of this section.

**Theorem 2.8.** *Let  $m \geq 0$  be an integer such that  $m!$  is invertible in  $\mathcal{O}$ , and let  $q \geq m$ . Suppose that either  $q = m$ , or that  $m + 1$  is not a zero divisor in  $\mathcal{O}$ . Let  $K$  be an abstract Koszul complex on  $(X, \mathcal{O})$  satisfying the flatness condition (2-1), and write*

$$\tau_{\leq 1} K = [K^0 \xrightarrow{\partial} Z^1 K]$$

*for its truncation in degrees  $\leq 1$ . Then the multiplication maps*

$$\mathrm{Kos}^q(\partial)^i = \Gamma^{q-i}(K^0) \otimes \wedge^i(Z^1 K) = \mathrm{Sym}^{q-i}(K^0) \otimes \wedge^i(Z^1 K) \rightarrow K^i$$

*for  $q - m \leq i \leq q$  (where we can identify  $\Gamma^{q-i}$  with  $\mathrm{Sym}^{q-i}$  as  $q - i \leq m$ , so that  $(q - i)!$  is invertible in  $\mathcal{O}$ ) induce a quasi-isomorphism*

$$\tau_{\geq q-m}(L\Gamma^q(\tau_{\leq 1}(K))) = \tau_{\geq q-m} \mathrm{Kos}^q(\partial) \xrightarrow{\sim} \tau_{[q-m, q]}(K). \tag{2-2}$$

*Proof.* The multiplication maps define a morphism of “naive truncations”

$$\begin{array}{ccc} \mathrm{Kos}^q(\partial)^{\geq q-m} = [\mathrm{Sym}^m(K^0) \otimes \wedge^{q-m}(\mathcal{Z}^1 K) \longrightarrow \cdots \longrightarrow \wedge^q(\mathcal{Z}^1 K)] & & \\ \mu \downarrow & & \downarrow \\ \tau_{\leq q}(K)^{\geq q-m} = & & [K^{q-m} \longrightarrow \cdots \longrightarrow Z^q K] \end{array}$$

To obtain the desired morphism  $\mu : \tau_{\geq q-m} \mathrm{Kos}^q(\partial) \rightarrow \tau_{[q-m, q]}(K)$ , we need to check that the map

$$\mathrm{Kos}^q(\partial)^{q-m} \rightarrow K^{q-m}$$

takes the image of  $\mathrm{Kos}^q(\partial)^{q-m-1} \rightarrow \mathrm{Kos}^q(\partial)^{q-m}$  into  $\mathcal{B}^{q-m} K = dK^{q-m-1}$ . This is clear if  $q = m$ , so suppose that  $(m + 1)$  is not a zero divisor.

Let  $z \in \mathrm{Kos}^q(\partial)^{q-m}$  be the image of  $w \in \mathrm{Kos}^q(\partial)^{q-m-1}$ , and consider  $(m + 1)w$  as an element of the submodule

$$\mathrm{Sym}^{m+1}(K^0) \otimes \wedge^{q-m-1}(\mathcal{Z}^1 K) \subseteq \Gamma^{m+1}(K^0) \otimes \wedge^{q-m-1}(\mathcal{Z}^1 K).$$

Let  $u \in K^{q-m-1}$  be the image of  $(m + 1)w$  under the multiplication map

$$\mathrm{Sym}^{m+1}(K^0) \otimes \wedge^{q-m-1}(\mathcal{Z}^1 K) \rightarrow K^{q-m-1}.$$

Then  $du = (m + 1)\mu(z)$  in  $K^{q-m}$ , where  $\mu(z)$  is the image of  $z$  under the multiplication map, and hence  $\mu(z)$  gives an  $(m + 1)$ -torsion class in  $\mathcal{H}^{q-m}(K)$ . Since by assumption  $\mathcal{H}^{q-m}(K) \simeq \wedge^{q-m} \mathcal{H}^1(K)$  is flat and  $m + 1$  is not a zero divisor,  $\mu(z) \in dK^{m-q-1}$  as desired.

Finally, the maps induced by  $\mu : \tau_{\geq q-m} \mathrm{Kos}^q(\partial) \rightarrow \tau_{[q-m, q]}(K)$  on cohomology can, thanks to Proposition 2.7, be identified with the maps

$$\mu^i : \wedge^i \mathcal{H}^1(K) \rightarrow \mathcal{H}^i(K) \quad \text{for } q - m \leq i \leq q,$$

which are isomorphisms by assumption. □

**Remark 2.9.** Implicit in the above proof is the subcomplex  $\widetilde{\mathrm{Kos}}^q(u)$  of  $\mathrm{Kos}^q(u)$  whose  $i$ -th term equals  $\mathrm{Sym}^{q-i} K \otimes \wedge^i(\mathcal{Z}^1 K)$ . The two complexes agree in degrees  $\geq q - m$ , and more generally the quotient  $\mathrm{Kos}^q(u)^{q-i} / \widetilde{\mathrm{Kos}}^q(u)^{q-i}$  is annihilated by  $i!$ . This subcomplex probably does not have any “derived meaning”, (for example, it is not clear that it is decomposable if  $\tau_{\leq 1}(K)$  is), but its advantage is that there is a multiplication map  $\mu : \widetilde{\mathrm{Kos}}^q(u) \rightarrow K$ .

**Remark 2.10.** For an illustration of Theorem 2.8, let us see what happens in the “minimal” situation where it does not apply. To this end, let us consider the de Rham complex  $K = [\mathbb{F}_p[x] \rightarrow \mathbb{F}_p[x] dx]$  of the polynomial ring  $\mathbb{F}_p[x]$  over  $\mathbb{F}_p$ , treated as a complex of modules over  $\mathbb{F}_p[x^p]$ . Set  $m = p - 1$  and  $q = p$ , and let us check that the intermediate assertion in the proof of Theorem 2.8, that  $\mu : \mathrm{Kos}^q(\partial)^{q-m} \rightarrow K^{q-m}$  takes the image of  $d_{\mathrm{Kos}} : \mathrm{Kos}^q(\partial)^{q-m-1} \rightarrow \mathrm{Kos}^q(\partial)^{q-m}$  into  $\mathcal{B}^{q-m} K = dK^{q-m-1}$ , does not hold. Explicitly,

the groups and maps in question form the diagram

$$\begin{array}{ccc}
 \Gamma_{\mathbb{F}_p[x^p]}^p(\mathbb{F}_p[x]) & \xrightarrow{d_{\text{Kos}}} & \Gamma_{\mathbb{F}_p[x^p]}^{p-1}(\mathbb{F}_p[x]) \otimes \mathbb{F}_p[x] dx \quad \simeq \quad \text{Sym}_{\mathbb{F}_p[x^p]}^{p-1}(\mathbb{F}_p[x]) \otimes \mathbb{F}_p[x] dx \\
 \downarrow ? & & \downarrow \mu \\
 d(\mathbb{F}_p[x])^{\subset} & \xrightarrow{\hspace{10em}} & \mathbb{F}_p[x] dx
 \end{array}$$

Let us consider the element  $x^{[p]}$  in the top left corner. Its image under the Koszul differential is  $x^{[p-1]} \otimes dx = -x^{p-1} \otimes dx$ , whose image under  $\mu$  is  $-x^{p-1} dx$ , a nonexact form.

This calculation suggests a link with the Cartier operator in the situation where  $K$  is the de Rham complex of a smooth scheme in characteristic  $p$ . And indeed, we shall see it again in the proof of Theorem 4.1.

**Remark 2.11.** Our proof of Theorem 2.8 makes use of an explicit model of the cdga  $K$ . Thus, for example, if  $K$  and  $K'$  are equivalent cdgas to which the theorem applies, it is not obvious whether the isomorphisms (2-2) we obtain for  $K$  and  $K'$  are compatible. More importantly, it does not apply to the more general case of coconnective  $E_\infty$ -algebras or cosimplicial commutative rings whose cohomology algebras satisfy axioms (i)–(ii) of Definition 2.1.

**Corollary 2.12.** *Let  $K$  be an abstract Koszul complex, and let  $n$  be such that  $n!$  is invertible in  $\mathcal{O}$ . Suppose that  $\tau_{\leq 1}(K)$  is decomposable. Then for  $a \leq b < a + n$ , the complex  $\tau_{[a,b]}(K)$  is decomposable. Moreover, the complex  $\tau_{\leq n}(K)$  is decomposable as well.*

**2D. Application to de Rham cohomology.** We now establish some of the straightforward consequences for de Rham cohomology mentioned in the introduction. The remaining ones shall be established at the end of Section 4.

*Proof of Theorem 1.1, case  $p > 2$ .* Let  $K = F_{X/k,*} \Omega_{X/k}^\bullet$ . By Example 2.2, this is an abstract Koszul complex over the ringed space  $(X', \mathcal{O}_{X'})$ . By [Deligne and Illusie 1987, théorème 2.1], the liftability assumption implies that the complex  $\tau_{\leq 1}(K)$  is decomposable. Corollary 2.12 with  $m = p - 1$  implies that  $\tau_{[a,b]}(K)$  is decomposable for  $a \leq b < a + p - 1$ , as desired. □

*Proof of Corollary 1.2.* The differentials on the  $E_r$ -page of (1-1) depend only on the truncations  $\tau_{[a,b]}(\Omega_{X/k}^\bullet)$  with  $a \leq b < a + r$ , and hence all differentials on the pages  $E_r$  with  $r < p$  vanish. Suppose that  $d_r^{ij} \neq 0$ . Then in particular  $H^i(X, \Omega_{X/k}^j)$  and  $H^{i+r}(X, \Omega_{X/k}^{j-r+1})$  are both nonzero, and hence

$$|i - j| < p \quad \text{and} \quad |(i + r) - (j - r + 1)| = |(i - j) + 2r - 1| < p,$$

which implies  $r < p$ , but that forces  $d_r^{ij} = 0$ , hence a contradiction. Therefore (1-1) degenerates.

For  $X$  proper over  $k$ , one can deduce the degeneration of the Hodge to de Rham spectral sequence as in [Deligne and Illusie 1987, corollaire 2.4]. □

**Remark 2.13.** As in [Deligne and Illusie 1987, § 4] and [Kato 1989, Theorem 4.12(2)], analogous assertions hold for a smooth and separated morphism of  $\mathbb{F}_p$ -schemes  $X \rightarrow S$ , or more generally for a

smooth morphism of Cartier type  $f : (X, \mathcal{M}_X) \rightarrow (S, \mathcal{M}_S)$  between fine log schemes over  $\mathbb{F}_p$ , assuming that there exists a fine log scheme  $(\tilde{S}, \mathcal{M}_{\tilde{S}})$  over  $\mathbb{Z}/p^2\mathbb{Z}$  such that  $\tilde{S}$  is flat over  $\mathbb{Z}/p^2\mathbb{Z}$  and a smooth lifting

$$\tilde{f}' : (\tilde{X}', \mathcal{M}_{\tilde{X}'}) \rightarrow (\tilde{S}, \mathcal{M}_{\tilde{S}})$$

of  $f'$  (the base change  $f$  under the absolute Frobenius  $(S, \mathcal{M}_S) \rightarrow (S, \mathcal{M}_S)$ ). Here  $\mathbb{F}_p$  and  $\mathbb{Z}/p^2\mathbb{Z}$  are given the trivial log structure.

### 3. Truncations in two consecutive degrees and gerbes of splittings

In the following, we make a more detailed analysis of the truncations  $\tau_{[q-1, q]}K$  for an abstract Koszul complex  $K$ , as well as their associated gerbes of splittings. We keep working in the category of modules in a ringed topos  $(X, \mathcal{O})$ .

**3A. First-order attachment maps.** For a complex  $K$  and an integer  $q$ , the truncation

$$\tau_{[q-1, q]}K = [\cdots \rightarrow 0 \rightarrow K^{q-1}/\mathcal{B}^{q-1} \rightarrow \mathcal{Z}^q K \rightarrow 0 \rightarrow \cdots]$$

fits inside the functorial exact triangle

$$\mathcal{H}^{q-1}(K)[1-q] \rightarrow \tau_{[q-1, q]}K \rightarrow \mathcal{H}^q(K)[-q] \xrightarrow{\delta_K^q[-q]} \mathcal{H}^{q-1}(K)[2-q],$$

yielding a morphism

$$\delta_K^q : \mathcal{H}^q(K) \rightarrow \mathcal{H}^{q-1}(K)[2]$$

such that  $\delta_K^q[-q]$  is the unique morphism making the above triangle distinguished (see [Achinger and Ogus 2020, Proposition 2.1.1]). Thus the truncation  $\tau_{[q-1, q]}K$  is determined by the map  $\delta_K^q$ , as the mapping fiber of  $\delta_K^q[-q]$ ; it is decomposable if and only if  $\delta_K^q = 0$ . We note for future reference the effect of the shift functor on the maps  $\delta_K^q$ :

$$\delta_{K[p]}^q = (-1)^p \delta_K^{p+q}. \tag{3-1}$$

The maps  $\delta_K^q$  describe the differentials on the second page of the spectral sequence

$$E_2^{pq} = H^p(X, \mathcal{H}^q(K)) \Rightarrow H^{p+q}(X, K).$$

Namely, the differential

$$d_2^{pq} : H^p(X, \mathcal{H}^q(K)) \rightarrow H^{p+2}(X, \mathcal{H}^{q-1}(K)) = H^p(X, \mathcal{H}^{q-1}(K)[2])$$

is the map induced by  $\delta_K^q$  on  $H^p(X, -)$ .

**3B. Gerbe of splittings.** We recall the gerbe of splittings described in [Deligne and Illusie 1987]. Let

$$K = [K^0 \xrightarrow{d} K^1]$$

be a two-term complex (i.e.,  $K^i = 0$  for  $i \neq 0, 1$ ), and suppose that the two conditions below hold:

- (1)  $\mathcal{H}^1(K)$  is locally free of finite rank, and
- (2) the projection of  $K^0$  onto  $\mathcal{B}^1 = \text{im}(d)$  locally admits a section.<sup>3</sup>

One then constructs the gerbe  $\text{sc}(K)$  under  $\underline{\text{Hom}}(\mathcal{H}^1(K), \mathcal{H}^0(K))$  over  $X$  [Deligne and Illusie 1987, § 3.2] as the stackification of the prestack  $\text{sc}'(K)$  whose objects are local splittings

$$s : \mathcal{H}^1(K) \rightarrow K^1$$

of the projection  $K^1 = \mathcal{Z}^1 K \rightarrow \mathcal{H}^1(K)$ , and where morphisms  $s \rightarrow s'$  are maps

$$h : \mathcal{H}^1(K) \rightarrow K^0$$

such that  $dh = s' - s$ . The automorphisms of an object  $s$  are then identified with  $\underline{\text{Hom}}(\mathcal{H}^1(K), \mathcal{H}^0(K))$ , and this makes  $\text{sc}(K)$  into a gerbe under  $\underline{\text{Hom}}(\mathcal{H}^1(K), \mathcal{H}^0(K))$ . We denote by

$$\text{cl sc } K \in H^2(X, \underline{\text{Hom}}(\mathcal{H}^1(K), \mathcal{H}^0(K))) = \text{Ext}^2(\mathcal{H}^1(K), \mathcal{H}^0(K)) = \text{Hom}(\mathcal{H}^1(K), \mathcal{H}^0(K)[2])$$

the class of the gerbe  $\text{sc } K$ . The following result relates this class to the map  $\delta_K^1$  defined previously.

**Lemma 3.1** [Deligne and Illusie 1987, proposition 3.3]. *Let  $K = [K^0 \rightarrow K^1]$  be a two-term complex satisfying (1) and (2) above. Then, one has the following equality in  $\text{Hom}(\mathcal{H}^1(K), \mathcal{H}^0(K)[2])$ :*

$$\text{cl sc } K = -\delta_K^1.$$

A bit more generally, suppose that  $q$  is an integer and  $K$  a complex satisfying the following conditions:

- (1)  $K^i = 0$  for  $i \neq q - 1, q$ ,
- (2)  $\mathcal{H}^q(K)$  is locally free of finite rank, and
- (3) the projection of  $K^{q-1}$  onto  $\mathcal{B}^q = \text{im}(d)$  locally admits a section.

Then we denote by  $\text{sc}_{[q-1, q]}(K)$  the gerbe of splittings of the complex

$$\dots \rightarrow 0 \rightarrow K^{q-1} \rightarrow K^q \rightarrow 0 \rightarrow \dots$$

concentrated in degrees 0 and 1 and with  $d$  being equal to the original differential of  $K$ , rather than  $(-1)^{q-1}$  times that; this convention has the consequence that

$$\text{cl sc}_{[q-1, q]}(K) = (-1)^{q-1} \text{cl sc}(K[q - 1])$$

in  $H^2(X, \underline{\text{Hom}}(\mathcal{H}^q, \mathcal{H}^{q-1}))$ . Combined with Lemma 3.1 and (3-1), this implies the following generalization of Lemma 3.1:

$$\text{cl sc}_{[q-1, q]}(K) = -\delta_K^q.$$

When there is no confusion as to what  $q$  is, we simply write  $\text{sc}(K)$  for  $\text{sc}_{[q-1, q]}(K)$ .

**3C. Truncated Koszul complexes.** Let  $K = [K^0 \xrightarrow{d} K^1]$  be a two-term complex of modules over  $(\mathcal{X}, \mathcal{O})$ , and let  $q \geq 1$ . Using the Koszul complex, one can build another two-term complex, concentrated in

---

<sup>3</sup>As pointed out to us by the referee, condition (2) is in fact not needed for the construction. Indeed, if  $s' - s : \mathcal{H}^1(K) \rightarrow \mathcal{B}^1$ , then locally there exists an  $h : \mathcal{H}^1 \rightarrow K^0$  such that  $s' - s = dh$ .

degrees  $[q - 1, q]$ :

$$\tau_{\geq q-1}(\mathbf{Kos}^q(d)) = \left[ \cdots \rightarrow 0 \rightarrow \frac{K^0 \otimes \wedge^{q-1} K^1}{d(\Gamma^2 K^0 \otimes \wedge^{q-2} K^1)} \rightarrow \wedge^q K^1 \rightarrow 0 \rightarrow \cdots \right].$$

By Proposition 2.7 we have morphisms

$$\alpha^q : \wedge^q \mathcal{H}^1(K) \rightarrow \mathcal{H}^q(\mathbf{Kos}^q(d)) \quad \text{and} \quad \alpha^{q-1} : \mathcal{H}^0(K) \otimes \wedge^{q-1} \mathcal{H}^1(K) \rightarrow \mathcal{H}^{q-1}(\mathbf{Kos}^q(d)), \quad (3-2)$$

which are isomorphisms if  $K^0$ ,  $K^1$ ,  $\mathcal{H}^0(K)$ , and  $\mathcal{H}^1(K)$  are flat. The following result describes the maps  $\delta_{\mathbf{Kos}^q(d)}^q$  and hence the truncation  $\tau_{\geq q-1}(\mathbf{Kos}^q(d))$ .

**Proposition 3.2.** *Let  $K = [K^0 \xrightarrow{d} K^1]$  be a two-term complex and let  $q \geq 1$ . Suppose that  $K^0$ ,  $K^1$ ,  $\mathcal{H}^0(K)$ , and  $\mathcal{H}^1(K)$  are flat. Then the following diagram is commutative:*

$$\begin{array}{ccc} \wedge^q \mathcal{H}^1(K) & \xrightarrow{\eta^q} & \mathcal{H}^1(K) \otimes \wedge^{q-1} \mathcal{H}^1(K) & \xrightarrow{\delta_K^1 \otimes \text{id}} & \mathcal{H}^0(K) \otimes \wedge^{q-1} \mathcal{H}^1(K)[2] \\ \alpha^q \downarrow & & & & \downarrow \alpha^{q-1} \\ \mathcal{H}^q(\mathbf{Kos}^q(d)) & \xrightarrow{\delta_{\mathbf{Kos}^q(d)}^q} & & & \mathcal{H}^{q-1}(\mathbf{Kos}^q(d))[2] \end{array}$$

In other words, using the identifications (3-2), we have the equality

$$\delta_{\mathbf{Kos}^q(d)}^q = (\delta_K^1 \otimes \text{id}) \circ \eta^q$$

of maps  $\wedge^q \mathcal{H}^1(K) \rightarrow \mathcal{H}^0(K) \otimes \wedge^{q-1} \mathcal{H}^1(K)[2]$ .

*Proof.* Let us abbreviate  $\mathcal{H}^i(K)$  to  $\mathcal{H}^i$ . We first check that the two-term complexes  $\tau_{\geq q-1} \mathbf{Kos}^q(d)$  and the naive  $(q-1)$ -shift of  $K \otimes \wedge^{q-1} \mathcal{H}^1$  form the middle square inside a commutative diagram with exact rows

$$\begin{array}{ccccccccc} 0 & \longrightarrow & \mathcal{H}^0 \otimes \wedge^{q-1} \mathcal{H}^1 & \longrightarrow & \frac{K^0 \otimes \wedge^{q-1} K^1}{d(\Gamma^2(K^0) \otimes \wedge^{q-2} K^1)} & \longrightarrow & \wedge^q K^1 & \longrightarrow & \wedge^q \mathcal{H}^1 & \longrightarrow & 0 \\ & & \text{id} \downarrow & & \downarrow \beta & & \downarrow \alpha & & \downarrow \eta^q & & \\ 0 & \longrightarrow & \mathcal{H}^0 \otimes \wedge^{q-1} \mathcal{H}^1 & \longrightarrow & K^0 \otimes \wedge^{q-1} \mathcal{H}^1 & \xrightarrow{d \otimes \text{id}} & K^1 \otimes \wedge^{q-1} \mathcal{H}^1 & \longrightarrow & \mathcal{H}^1 \otimes \wedge^{q-1} \mathcal{H}^1 & \longrightarrow & 0 \end{array}$$

We define the maps  $\alpha$  and  $\beta$  as follows. The map  $\beta$  is uniquely determined by

$$\beta(w \otimes z_1 \wedge \cdots \wedge z_{q-1}) \pmod{d(\Gamma^2(K^0) \otimes \wedge^{q-2} K^1)} = w \otimes [z_1] \wedge \cdots \wedge [z_{q-1}].$$

It is well defined because elements of the form

$$d(w^{[2]} \otimes z_1 \wedge \cdots \wedge z_{q-1}) = w \otimes dw \wedge z_1 \wedge \cdots \wedge z_{q-1}$$

or

$$d(wv \otimes z_1 \wedge \cdots \wedge z_{q-1}) = v \otimes dw \wedge z_1 \wedge \cdots \wedge z_{q-1} + w \otimes dv \wedge z_1 \wedge \cdots \wedge z_{q-1}$$

are sent to zero, since  $[dw] = 0 = [dv]$ . The map  $\alpha$  is the composition

$$\wedge^q K^1 \xrightarrow{\eta^q} K^1 \otimes \wedge^{q-1} K^1 \xrightarrow{\text{id} \otimes \text{proj.}} K^1 \otimes \wedge^{q-1} \mathcal{H}^1.$$

The commutativity of the left- and rightmost squares is trivial to check. To see that the middle square commutes, we take (the class of)  $w \otimes z_1 \wedge \cdots \wedge z_{q-1} \in K^0 \otimes \wedge^{q-1} K^1$ , and compute

$$\begin{aligned} \alpha(d(w \otimes z_1 \wedge \cdots \wedge z_{q-1})) &= \alpha(dw \wedge z_1 \wedge \cdots \wedge z_{q-1}) \\ &= \sum_{i=1}^q (-1)^i z_i \otimes [dw] \wedge [z_1] \wedge \cdots \wedge \widehat{[z_i]} \wedge \cdots \wedge [z_{q-1}] + dw \otimes [z_1] \wedge \cdots \wedge [z_{q-1}] \\ &= dw \otimes [z_1] \wedge \cdots \wedge [z_{q-1}] = (d \otimes \text{id})(\beta(w \otimes z_1 \wedge \cdots \wedge z_{q-1})). \end{aligned}$$

Now, our commutative diagram of complexes translates into a commutative square in the derived category,

$$\begin{array}{ccc} \wedge^q \mathcal{H}^1 & \xrightarrow{\delta_{\text{Kos}^q(d)}^q} & \mathcal{H}^0 \otimes \wedge^{q-1} \mathcal{H}^1[2] \\ \eta^q \downarrow & & \downarrow \text{id} \\ \mathcal{H}^1 \otimes \wedge^{q-1} \mathcal{H}^1 & \xrightarrow{\delta_K^1 \otimes \text{id}} & \mathcal{H}^0 \otimes \wedge^{q-1} \mathcal{H}^1[2] \end{array}$$

This implies the required assertion. □

**3D. Two-term truncations of abstract Koszul complexes.** The following result relates the maps  $\delta_K^q$  and  $\delta_K^1$  for a cdga  $K$ .

**Proposition 3.3.** *Suppose 2 is a nonzerodivisor in  $\mathcal{O}$ . Let  $K$  be a coconnective commutative differential graded algebra such that  $K^0$ ,  $\mathcal{Z}^1 K$ ,  $\mathcal{H}^0(K)$ , and  $\mathcal{H}^1(K)$  are flat. Let  $q \geq 1$  be an integer such that  $\mathcal{H}^{q-1}(K)$  is flat. Then the following diagram commutes:*

$$\begin{array}{ccccc} \wedge^q \mathcal{H}^1(K) & \xrightarrow{\eta^q} & \mathcal{H}^1(K) \otimes \wedge^{q-1} \mathcal{H}^1(K) & \xrightarrow{\delta_K^1 \otimes \text{id}} & \mathcal{H}^0(K) \otimes \wedge^{q-1} \mathcal{H}^1(K)[2] \\ \text{mult.} \downarrow & & & & \downarrow \text{mult.} \\ \mathcal{H}^q(K) & \xrightarrow{\delta_K^q} & & & \mathcal{H}^{q-1}(K)[2] \end{array}$$

*Proof.* Write  $\tau_{\leq 1} K = [K^0 \xrightarrow{\partial} \mathcal{Z}^1 K]$ . The proof of Theorem 2.8 (with  $m = 1$ ) provides a morphism of complexes

$$\mu : \tau_{\geq q-1} \text{Kos}^q(\partial) \rightarrow \tau_{[q-1, q]} K.$$

By functoriality of the maps  $\delta^q$ , we have a commutative square

$$\begin{array}{ccc} \mathcal{H}^q(\tau_{\geq q-1} \text{Kos}^q(\partial)) & \longrightarrow & \mathcal{H}^{q-1}(\tau_{\geq q-1} \text{Kos}^q(\partial))[2] \\ \mu \downarrow & & \downarrow \\ \mathcal{H}^q(\tau_{[q-1, q]} K) & \longrightarrow & \mathcal{H}^{q-1}(\tau_{[q-1, q]} K)[2] \end{array}$$



That is, a commutative square

$$\begin{array}{ccc} \wedge^q \mathcal{H}^1(K) & \longrightarrow & \mathcal{H}^0(K) \otimes \wedge^{q-1} \mathcal{H}^1(K)[2] \\ \downarrow & & \downarrow \\ \mathcal{H}^q(K) & \longrightarrow & \mathcal{H}^{q-1}(K)[2] \end{array}$$

The assertion then follows from Proposition 3.2. □

**Remark 3.4.** The proof of [Achinger and Ogus 2020, Theorem 4.2.2(1)] implies the assertion of Proposition 3.3 under the stronger assumption that  $q!$  is invertible in  $\mathcal{O}$ . However, the argument does not use the cdga structure of  $K$ , only a weaker structure of a commutative monoid in the derived category  $K \otimes^{\mathbb{L}} K \rightarrow K$ . In particular, the assertion holds for some  $E_\infty$ -algebras which are not a priori equivalent to cdgas.

**Remark 3.5.** In [Achinger and Ogus 2020, Lemma 2.1.1], it is shown that the maps  $\delta_K^q$  are compatible with the derived tensor product in the following way. If  $K$  and  $L$  are complexes and  $i$  and  $j$  are integers such that  $\mathcal{H}^i(K)$  and  $\mathcal{H}^j(L)$  are flat  $\mathcal{O}$ -modules, then the following square commutes:

$$\begin{array}{ccc} \mathcal{H}^i(K) \otimes \mathcal{H}^j(L) & \xrightarrow{\delta_K^i \otimes 1 + (-1)^i \otimes \delta_L^j} & (\mathcal{H}^{i-1}(K)[2] \otimes \mathcal{H}^j(L)) \oplus (\mathcal{H}^i(K) \otimes \mathcal{H}^{j-1}(L)[2]) \\ \downarrow & & \downarrow \\ \mathcal{H}^{i+j}(K \otimes^{\mathbb{L}} L) & \xrightarrow{\delta_{K \otimes^{\mathbb{L}} L}^{i+j}} & \mathcal{H}^{i+j-1}(K \otimes^{\mathbb{L}} L)[2] \end{array}$$

If  $q!$  is invertible in  $\mathcal{O}$ , so that  $\wedge^q \mathcal{H}^1(K)$  is a direct summand of  $\mathcal{H}^1(K)^{\otimes q}$ , the assertion of Proposition 3.3 can be deduced from this result.

For illustration, let us see how to do this for  $q = 2$ . We set  $L = K$  and  $i = j = 1$  in the above diagram, obtaining the middle square of the diagram below:

$$\begin{array}{ccccc} \wedge^2 \mathcal{H}^1(K) & \xrightarrow{\eta^2} & \mathcal{H}^1(K) \otimes \mathcal{H}^1(K) & \xrightarrow{\text{id} \otimes \delta_K^1} & \mathcal{H}^1(K) \otimes \mathcal{H}^0(K)[2] \\ \downarrow x \wedge y \mapsto \frac{1}{2}(x \otimes y - y \otimes x) & & \downarrow & & \downarrow \frac{1}{2}(\text{shuffle}, \text{id}) \\ \mathcal{H}^1(K) \otimes \mathcal{H}^1(K) & \xrightarrow{\delta_K^1 \otimes 1 - 1 \otimes \delta_K^1} & (\mathcal{H}^0(K)[2] \otimes \mathcal{H}^1(K)) \oplus (\mathcal{H}^1(K) \otimes \mathcal{H}^0(K)[2]) & & \\ \downarrow & & \downarrow & & \\ \mathcal{H}^2(K \otimes^{\mathbb{L}} K) & \xrightarrow{\delta_{K \otimes^{\mathbb{L}} K}^2} & \mathcal{H}^1(K \otimes^{\mathbb{L}} K)[2] & & \\ \downarrow & & \downarrow & & \\ \mathcal{H}^2(K) & \xrightarrow{\delta_K^2} & \mathcal{H}^1(K)[2] & & \end{array}$$

Here, the bottom square certifies the functoriality of  $\delta^2$  with respect to the multiplication map  $K \otimes^{\mathbb{L}} K \rightarrow K$ . Commutativity of the top square is easy to check. Then, commutativity of the exterior square gives the required assertion.

**Corollary 3.6.** *Suppose 2 is a nonzerodivisor in  $\mathcal{O}$ . Let  $K$  be an abstract Koszul complex satisfying the flatness condition (2-1) and let  $q \geq 1$ . We have the following commutative diagram:*

$$\begin{array}{ccccc} \wedge^q \mathcal{H}^1(K) & \xrightarrow{\eta^q} & \mathcal{H}^1(K) \otimes \wedge^{q-1} \mathcal{H}^1(K) & \xrightarrow{\delta_K^1 \otimes \text{id}} & \wedge^{q-1} \mathcal{H}^1(K)[2] \\ \wr \downarrow & & & & \downarrow \wr \\ \mathcal{H}^q(K) & \xrightarrow{\delta_K^q} & & & \mathcal{H}^{q-1}(K)[2] \end{array}$$

In other words, using the vertical identifications, we have the equality

$$\delta_K^q = (\delta_K^1 \otimes \text{id}) \circ \eta^q$$

in  $\text{Hom}(\mathcal{H}^q(K), \mathcal{H}^{q-1}(K)[2])$ .

**Corollary 3.7.** *Suppose 2 is a nonzerodivisor in  $\mathcal{O}$ . Let  $K$  be an abstract Koszul complex satisfying the flatness condition (2-1) and let  $q \geq 1$ . Then*

$$\text{cl sc}_{[q-1,q]}(K) = \eta^q(\text{cl sc}(\tau_{\leq 1} K)).$$

**Corollary 3.8.** *Suppose that 2 is a nonzerodivisor in  $\mathcal{O}$ . Let  $K$  be an abstract Koszul complex satisfying the flatness condition (2-1). Then the differential*

$$d_2^{p,q} : H^p(X, \mathcal{H}^q(K)) \rightarrow H^{p+2}(X, \mathcal{H}^{q-1}(K))$$

equals the cup product with the class

$$\delta_K^1 = -\text{cl sc}(\tau_{\leq 1} K) \in H^2(X, \underline{\text{Hom}}(\mathcal{H}^1(K), \mathcal{H}^0(K))),$$

followed by evaluation.

**3E. Morphisms of gerbes of splittings.** Let  $K$  be an abstract Koszul complex satisfying the flatness condition (2-1). In Corollary 3.7, under the assumption that 2 is a nonzerodivisor in  $\mathcal{O}$ , we calculated the gerbe classes  $\text{cl sc}_{[q-1,q]} K$  in terms of the class  $\text{cl sc}(\tau_{\leq 1} K)$ . Below, under the stronger assumption that 2 is invertible, we promote this equality into a morphism of gerbes.

**Theorem 3.9.** *For each integer  $q \geq 1$ , there is a morphism*

$$\wedge^q : \text{sc}(\tau_{\leq 1} K) \rightarrow \text{sc}(\tau_{[q-1,q]} K)$$

of gerbes over  $X$ , under which the obstruction classes correspond by the relation

$$\text{cl sc}(\tau_{[q-1,q]} K) = \text{ctr}^q(\text{cl sc } \tau_{\leq 1}(K)),$$

where  $\text{ctr}^q : \underline{\text{Hom}}(\mathcal{H}^1, \mathcal{H}^0) \rightarrow \underline{\text{Hom}}(\mathcal{H}^q, \mathcal{H}^{q-1})$  denotes the morphism which maps a local section  $f$  of the source to the one of the target by the formula

$$\text{ctr}^q(f) : \omega_1 \wedge \cdots \wedge \omega_q \mapsto \sum_{j=1}^q (-1)^{j-1} f(\omega_j) \omega_1 \wedge \cdots \wedge \omega_{j-1} \wedge \omega_{j+1} \wedge \cdots \wedge \omega_q.$$

(Compare the formula for  $\text{ctr}^q$  with the explicit formula for  $\eta^q$  in Section 2B.)

**Notation.** Before proceeding to the proof, we gather some notation concerning Čech cohomology. We denote by  $\check{C}(U_\bullet, K^\bullet)$  the Čech resolution of a complex  $K^\bullet$  with respect to a hypercovering  $U_\bullet$ . The differential induced by that of  $K^\bullet$  will still be denoted by  $d$ , while the Čech differential on the component  $\check{C}(U_p, K^q)$ ,

$$(-1)^q \sum_{i=0}^{p+1} (-1)^i d_i^*,$$

will be denoted by  $\check{d}$ . Then the total differential

$$D = d + \check{d}$$

is the differential of the total complex  $\check{C}(U_\bullet, K^\bullet)$ .

When we compute the obstruction classes, we will use some notation which may not be standard. As usual, for each integer  $m \geq -1$ , we denote by  $[m]$  the set of integers  $i$  such that  $0 \leq i \leq m$  (empty set for  $[-1]$ ). And we denote by  $d_{ij} : [m-2] \rightarrow [m]$  the unique increasing injection omitting  $i$  and  $j$ , where  $0 \leq i < j \leq m$ . For example, for  $m = 2$ , we have

$$d_{02} = d_2 \circ d_0 = d_0 \circ d_1 : [0] \rightarrow [2]$$

(which maps 0 onto 1), where  $d_i : [m-1] \rightarrow [m]$  denotes the unique increasing injection omitting  $i$ .

On the other hand, we denote by  $\text{pr}_i : [0] \rightarrow [m]$  (resp.  $\text{pr}_{ij} : [1] \rightarrow [m]$ ) the unique map sending 0 to  $i$  (resp. 0 to  $i$  and 1 to  $j$ ) for  $0 \leq i \leq m$  (resp. for  $0 \leq i < j \leq m$ ).

*Proof of Theorem 3.9.* In order to prove Theorem 3.9, we first describe the morphism, show that it is well defined, and then calculate the obstruction class.

**Construction of the functor  $\wedge^q$ .** We construct  $\wedge^q : \text{sc } \tau_{\leq 1} K \rightarrow \text{sc } \tau_{[q-1, q]} K$  by stackifying a morphism between the corresponding prestacks:  $\text{sc}' \tau_{\leq 1} K \rightarrow \text{sc}' \tau_{[q-1, q]} K$ .

Given an object of  $\text{sc}'(\tau_{\leq 1} K)$  over  $U$ , that is, a section  $s : \mathcal{H}^1 \rightarrow \mathcal{Z}^1$  of the projection  $\mathcal{Z}^1 \rightarrow \mathcal{H}^1$  over  $U$ , we define  $\wedge^q(s)$  as the composite morphism

$$\mathcal{H}^q \simeq \wedge^q(\mathcal{H}^1) \xrightarrow{\wedge^q(s)} \wedge^q(\mathcal{Z}^1) \xrightarrow{\text{prod}} \mathcal{Z}^q,$$

where  $\text{prod}$  means product; it is clearly a section of  $\mathcal{Z}^q \rightarrow \mathcal{H}^q$  over  $U$ .

Let  $s_0$  and  $s_1$  be two objects of  $\text{sc}'(\tau_{\leq 1}K)$  over  $U$  and let  $h$  be a homotopy from  $s_0$  to  $s_1$ . Then we need to define a corresponding homotopy  $\wedge^q(h)$  from  $\wedge^q(s_0)$  to  $\wedge^q(s_1)$ . We first define a map

$$\wedge^q(h) : (\mathcal{H}^1)^{\otimes q} \rightarrow K^{q-1}/\mathcal{B}^{q-1}$$

by letting it send  $\omega_1 \otimes \cdots \otimes \omega_q$  (where  $\omega_j$  are local sections of  $\mathcal{H}^1$ ) to the class of

$$\sum_{j=1}^q (-1)^{j-1} h(\omega_j) s_0(\omega_1) \wedge \cdots \wedge s_0(\omega_{j-1}) \wedge s_1(\omega_{j+1}) \wedge \cdots \wedge s_1(\omega_q)$$

modulo  $\mathcal{B}^{q-1}$ .

It is easy to show that it factors through  $\wedge^q \mathcal{H}^1$ : if, say,  $\omega_1 = \omega_2 = \omega$ , then the alternating sum on the right reduces to the difference of the first two terms

$$h(\omega) s_1(\omega) \wedge s_1(\omega_3) \wedge \cdots \wedge s_1(\omega_q) - h(\omega) s_0(\omega) \wedge s_1(\omega_3) \wedge \cdots \wedge s_1(\omega_q),$$

which is equal to

$$h(\omega) dh(\omega) \wedge s_1(\omega_3) \wedge \cdots \wedge s_1(\omega_q),$$

which is a coboundary *since 2 is invertible*. Thus we have defined  $\wedge^q(h)$ :

$$\begin{array}{ccc} (\mathcal{H}^1)^{\otimes q} & & \\ \downarrow & \searrow \wedge^q(h) & \\ \wedge^q(\mathcal{H}^1) & \xrightarrow[\wedge^q(h)]{} & K^{q-1}/\mathcal{B}^{q-1} \end{array}$$

Then the following calculation shows that  $\wedge^q(h)$  is really a homotopy:

$$\begin{aligned} & (\wedge^q(s_1) - \wedge^q(s_0))(\omega_1 \wedge \cdots \wedge \omega_q) \\ &= \sum_{j=1}^q s_0(\omega_1) \wedge \cdots \wedge s_0(\omega_{j-1}) \wedge \{s_1(\omega_j) - s_0(\omega_j)\} \wedge s_1(\omega_{j+1}) \wedge \cdots \wedge s_1(\omega_q) \\ &= \sum_{j=1}^q s_0(\omega_1) \wedge \cdots \wedge s_0(\omega_{j-1}) \wedge \{dh(\omega_j)\} \wedge s_1(\omega_{j+1}) \wedge \cdots \wedge s_1(\omega_q) \\ &= \sum_{j=1}^q (-1)^{j-1} dh(\omega_j) \wedge s_0(\omega_1) \wedge \cdots \wedge s_0(\omega_{j-1}) \wedge s_1(\omega_{j+1}) \wedge \cdots \wedge s_1(\omega_q) \\ &= d[\wedge^q(h)(\omega_1 \wedge \cdots \wedge \omega_q)]. \end{aligned}$$

**Functoriality of  $\wedge^q$ .** Now in order to show that the morphism  $\wedge^q$  is a functor, we must show that it is compatible with the composition of homotopies; so let  $h : s_0 \Rightarrow s_1$  and  $h' : s_1 \Rightarrow s_2$  be two such in the

source. We first define a second homotopy operator

$$H_2^q(h, h') : (\mathcal{H}^1)^{\otimes q} \rightarrow K^{q-2}, \quad \omega_1 \otimes \cdots \otimes \omega_q \mapsto \sum_{1 \leq j < k \leq q} (-1)^{j+k+1} h(\omega_j) h'(\omega_k) s(j, k),$$

where  $s(j, k)$  is equal to

$$s_0(\omega_1) \wedge \cdots \wedge s_0(\omega_{j-1}) \wedge s_1(\omega_{j+1}) \wedge \cdots \wedge s_1(\omega_{k-1}) \wedge s_2(\omega_{k+1}) \wedge \cdots \wedge s_2(\omega_q).$$

To show that  $\wedge^q(h + h')$  and  $\wedge^q(h) + \wedge^q(h')$  are the same homotopies, it suffices to demonstrate the formula

$$[\wedge^q(h + h') - \{\wedge^q(h) + \wedge^q(h')\}](\omega_1 \wedge \cdots \wedge \omega_q) = dH_2^q(\omega_1 \otimes \cdots \otimes \omega_q).$$

One expands the left-hand side and groups the terms involving  $h$  and  $h'$  separately:

$$\begin{aligned} & \sum_{j=1}^q (-1)^{j-1} \{h(\omega_j) + h'(\omega_j)\} s_0(\omega_1) \wedge \cdots \wedge s_0(\omega_{j-1}) \wedge s_2(\omega_{j+1}) \wedge \cdots \wedge s_2(\omega_q) \\ & \quad - \sum_{j=1}^q (-1)^{j-1} h(\omega_j) s_0(\omega_1) \wedge \cdots \wedge s_0(\omega_{j-1}) \wedge s_1(\omega_{j+1}) \wedge \cdots \wedge s_1(\omega_q) \\ & \quad \quad - \sum_{j=1}^q (-1)^{j-1} h'(\omega_j) s_1(\omega_1) \wedge \cdots \wedge s_1(\omega_{j-1}) \wedge s_2(\omega_{j+1}) \wedge \cdots \wedge s_2(\omega_q) \\ & = \sum_{j=1}^q (-1)^{j-1} h(\omega_j) s_0(\omega_1) \wedge \cdots \wedge s_0(\omega_{j-1}) \wedge \{s_2(\omega_{j+1}) \wedge \cdots \wedge s_2(\omega_q) - s_1(\omega_{j+1}) \wedge \cdots \wedge s_1(\omega_q)\} \\ & \quad + \sum_{k=1}^q (-1)^{k-1} h'(\omega_k) \{s_0(\omega_1) \wedge \cdots \wedge s_0(\omega_{k-1}) - s_1(\omega_1) \wedge \cdots \wedge s_1(\omega_{k-1})\} \wedge s_2(\omega_{k+1}) \wedge \cdots \wedge s_2(\omega_q). \end{aligned}$$

The differences in the curly brackets are themselves alternating sums, so the last expression is equal to

$$\begin{aligned} & \sum_{j=1}^q (-1)^{j-1} h(\omega_j) s_0(\omega_1) \wedge \cdots \wedge s_0(\omega_{j-1}) \\ & \quad \wedge \left\{ \sum_{k>j} (-1)^{k-(j+1)} dh'(\omega_k) \wedge s_1(\omega_{j+1}) \wedge \cdots \wedge s_1(\omega_{k-1}) \wedge s_2(\omega_{k+1}) \wedge \cdots \wedge s_2(\omega_q) \right\} \\ & \quad + \sum_{k=1}^q (-1)^{k-1} h'(\omega_k) \left\{ \sum_{j<k} (-1)^{j-1} (-dh(\omega_j)) \wedge s_0(\omega_1) \wedge \cdots \wedge s_0(\omega_{j-1}) \wedge s_1(\omega_{j+1}) \wedge \cdots \wedge s_1(\omega_{k-1}) \right\} \\ & \quad \quad \quad \wedge s_2(\omega_{k+1}) \wedge \cdots \wedge s_2(\omega_q) \\ & = \sum_{1 \leq j < k \leq q} (-1)^{j+k+1} \{h(\omega_j) dh'(\omega_k) + h'(\omega_k) dh(\omega_j)\} \wedge s(j, k), \end{aligned}$$

and this is now equal to

$$dH_2^q(h, h')(\omega_1 \otimes \cdots \otimes \omega_q). \tag{3-3}$$

This completes the proof of the fact that  $\wedge^q$  is a functor.

**Calculation of obstruction classes.** Finally, we relate the obstruction elements. Let  $U_\bullet \rightarrow X$  be an open hypercovering such that one has

- (1) a section  $s : \mathcal{H}^1 \rightarrow \mathcal{Z}^1$  of the canonical projection  $\mathcal{Z}^1 \rightarrow \mathcal{H}^1$  over  $U_0$ , and
- (2) a homotopy  $h : \mathcal{H}^1 \rightarrow K^0$  over  $U_1$  satisfying

$$d_1^*s - d_0^*s = dh. \quad (3-4)$$

Then, by definition, the class of

$$\text{obs} = \text{obs}_1 = d_0^*h - d_1^*h + d_2^*h \in \Gamma(U_2, \underline{\text{Hom}}(\mathcal{H}^1, \mathcal{H}^0))$$

in  $H^2(X, \underline{\text{Hom}}(\mathcal{H}^1, \mathcal{H}^0))$  is cl sc  $\tau_{\leq 1}K$ . On the other hand, by applying  $\wedge^q$  to  $s$  and  $h$ , one sees that the class of

$$\text{obs}_q = d_0^*(\wedge^q h) - d_1^*(\wedge^q h) + d_2^*(\wedge^q h) \in \Gamma(U_2, \underline{\text{Hom}}(\mathcal{H}^q, \mathcal{H}^{q-1}))$$

in  $H^2(X, \underline{\text{Hom}}(\mathcal{H}^q, \mathcal{H}^{q-1}))$  is cl sc  $\tau_{[q-1, q]}K$ .

Now let  $\omega_1 \wedge \cdots \wedge \omega_q$  be a local section of  $\mathcal{H}^q \simeq \wedge^q \mathcal{H}^1$ . Then the evaluation of  $\text{obs}_q$  at  $\omega_1 \wedge \cdots \wedge \omega_q$  is equal to

$$\begin{aligned} & \sum_{j=1}^q (-1)^{j+1} d_0^*h(\omega_j) d_{01}^*s(\omega_1) \wedge \cdots \wedge d_{01}^*s(\omega_{j-1}) \wedge d_{02}^*s(\omega_{j+1}) \wedge \cdots \wedge d_{02}^*s(\omega_q) \\ & - \sum_{j=1}^q (-1)^{j+1} d_1^*h(\omega_j) d_{01}^*s(\omega_1) \wedge \cdots \wedge d_{01}^*s(\omega_{j-1}) \wedge d_{12}^*s(\omega_{j+1}) \wedge \cdots \wedge d_{12}^*s(\omega_q) \\ & + \sum_{j=1}^q (-1)^{j+1} d_2^*h(\omega_j) d_{02}^*s(\omega_1) \wedge \cdots \wedge d_{02}^*s(\omega_{j-1}) \wedge d_{12}^*s(\omega_{j+1}) \wedge \cdots \wedge d_{12}^*s(\omega_q). \end{aligned}$$

One groups the terms around the second sum and gets

$$\begin{aligned} & \sum_{j=1}^q (-1)^{j+1} (d_0^*h - d_1^*h + d_2^*h)(\omega_j) \cdot d_{01}^*s(\omega_1) \wedge \cdots \wedge d_{01}^*s(\omega_{j-1}) \wedge d_{12}^*s(\omega_{j+1}) \wedge \cdots \wedge d_{12}^*s(\omega_q) \\ & + \sum_{j=1}^q (-1)^{j+1} d_0^*h(\omega_j) d_{01}^*s(\omega_1) \wedge \cdots \wedge d_{01}^*s(\omega_{j-1}) \\ & \quad \wedge \{d_{02}^*s(\omega_{j+1}) \wedge \cdots \wedge d_{02}^*s(\omega_q) - d_{12}^*s(\omega_{j+1}) \wedge \cdots \wedge d_{12}^*s(\omega_q)\} \\ & + \sum_{j=1}^q (-1)^{j+1} d_2^*h(\omega_j) \{d_{02}^*s(\omega_1) \wedge \cdots \wedge d_{02}^*s(\omega_{j-1}) - d_{01}^*s(\omega_1) \wedge \cdots \wedge d_{01}^*s(\omega_{j-1})\} \\ & \quad \wedge d_{12}^*s(\omega_{j+1}) \wedge \cdots \wedge d_{12}^*s(\omega_q). \end{aligned}$$

The first alternating sum reduces to the “main” term we want, when taken modulo the coboundaries ( $\mathcal{B}^{q-1}$ ). In the last two sums, one first notes that, as  $h$  is a homotopy from  $d_0^*s$  to  $d_1^*s$ , it follows that  $d_0^*h$

is a homotopy from  $d_0^* d_0^* s$  to  $d_0^* d_1^* s$ , that is,

$$d_0^* h : d_{01}^* s \Rightarrow d_{02}^* s.$$

Similarly,  $d_2^* h$  is a homotopy from  $d_{02}^* s$  to  $d_{12}^* s$ . Essentially by repeating the last three equalities leading up to (3-3), this time with a minus sign, one sees that the last two sums add up to

$$-dH_2^q(d_0^* h, d_2^* h)(\omega_1 \otimes \cdots \otimes \omega_q),$$

which is a coboundary. Therefore, reducing modulo  $\mathcal{B}^{q-1}$ , one gets

$$\text{ev}(\text{obs}_q, \omega_1 \wedge \cdots \wedge \omega_q) = \text{ev}(\text{ctr}^q(\text{obs}_1), \omega_1 \wedge \cdots \wedge \omega_q).$$

This means

$$\text{cl sc } \tau_{[q-1, q]} K = \text{ctr}^q \text{ cl sc } \tau_{\leq 1} K,$$

which completes the proof. □

**Remark 3.10.** The construction of the map between gerbes can also be carried out using the language of higher topos theory [Lurie 2009]. Let us give a brief outline.

Let  $p : Y \rightarrow Z$  be a map of spaces, or more generally in any  $\infty$ -category  $\mathcal{C}$ . One can then build the space  $\text{sc}(p)$  of splittings of  $p$  as the homotopy fiber of

$$p : \text{Hom}_{\mathcal{C}}(Z, Y) \rightarrow \text{Hom}_{\mathcal{C}}(Z, Z)$$

over the identity  $\text{id}_Z$ . Similarly, if  $p : Y \rightarrow Z$  is a map in the derived  $\infty$ -category of a ringed topos  $(X, \mathcal{O})$ , one obtains a sheaf of spaces  $\underline{\text{sc}}(p)$  of splittings of  $p$ .

In the special case when  $Y$  is a two-term complex  $K = [K^0 \xrightarrow{d} K^1]$  satisfying the conditions in Section 3B and  $Y \rightarrow Z$  is the projection  $K \rightarrow \mathcal{H}^1(K)[-1]$ , the sheaf  $\underline{\text{sc}}(p)$  is a sheaf of groupoids (a stack) and can be identified with the gerbe of splittings  $\text{sc } K$ .

Applying the functor  $\tau_{\geq q-1} L\Gamma^q$  to the map  $p : K \rightarrow \mathcal{H}^1(K)[-1]$  one obtains (simply by functoriality) a morphism of sheaves of spaces

$$\underline{\text{sc}}(p) \rightarrow \underline{\text{sc}}(\tau_{\geq q-1} L\Gamma^q(p)).$$

By inspection, the map  $\tau_{\geq q-1} L\Gamma^q(p)$  is the projection

$$\tau_{[q-1, q]} \text{Kos}^q(d) \rightarrow \mathcal{H}^q(\text{Kos}^q(d))[-q] = \wedge^q \mathcal{H}^1(K)[-q].$$

This way one obtains by abstract nonsense a morphism of gerbes  $\text{sc } K \rightarrow \text{sc}(\tau_{[q-1, q]} \text{Kos}^q(d))$ .

In the case when  $K$  is an abstract Koszul complex satisfying the flatness condition (2-1), the morphism of gerbes  $\text{sc } \tau_{\leq 1} K \rightarrow \text{sc } \tau_{[q-1, q]} K$  obtained this way should agree with the one constructed in Theorem 3.9, though we did not check it.

#### 4. Gerbes of splittings of the de Rham complex

Our method of explicating the truncations  $\tau_{[q-1, q]}K$  for an abstract Koszul complex  $K$  in terms of the truncation  $\tau_{\leq 1}K$  requires that 2 be a nonzerodivisor. In this section, we describe these two-term truncations in the case of the de Rham complex in characteristic  $p > 0$  by calculating the class

$$\text{cl sc}(\tau_{[q-1, q]}F_*\Omega_{X/S}^\bullet).$$

The calculation uses more information about the de Rham complex than its being an abstract Koszul complex, namely the nature of the Cartier isomorphism (which we use only for  $p = 2$ ). As a corollary, we deduce that  $\tau_{[q-1, q]}(F_*\Omega_{X/S}^\bullet)$  is decomposable if  $\tau_{\leq 1}(F_*\Omega_{X/S}^\bullet)$  is, and obtain a description of the  $d_2$  differentials in the conjugate spectral sequence.

**Theorem 4.1.** *Let  $S$  be a scheme of characteristic  $p > 0$  and  $X/S$  a smooth separated scheme of finite type. Then for each integer  $q$ , the class*

$$\text{cl sc}(\tau_{[q-1, q]}F_*\Omega_{X/S}^\bullet)$$

is the image of the class

$$\text{cl sc}(\tau_{\leq 1}F_*\Omega_{X/S}^\bullet)$$

under the contraction map (described in Theorem 3.9).

*Proof.* We put  $K = F_*\Omega_{X/S}^\bullet$ , with  $F : X \rightarrow X'$  the relative Frobenius of  $X/S$ .

To calculate the class, we take an open hypercovering  $U_\bullet \rightarrow X'$  such that

(1) over  $U_0$ , one has a section  $s : \mathcal{H}^1 \rightarrow \mathcal{Z}^1$  of the projection  $\mathcal{Z}^1 \rightarrow \mathcal{H}^1$  and a section

$$\sigma^{(q)} : \mathcal{H}^q \rightarrow (\mathcal{H}^1)^{\otimes q}$$

of the canonical projection  $(\mathcal{H}^1)^{\otimes q} \rightarrow \mathcal{H}^q$ , and

(2) over  $U_1$ , one has a homotopy  $h : \mathcal{H}^1 \rightarrow K^0$  such that

$$dh = d_1^*s - d_0^*s : \mathcal{H}^1 \rightarrow \mathcal{Z}^1.$$

(Let us recall that  $\mathcal{H}^1$  is locally free over  $\mathcal{O}_{X'}$ , and hence so is  $\mathcal{H}^q = \wedge^q \mathcal{H}^1$  for all integers  $q$ .) The locally free kernel of the projection  $(\mathcal{H}^1)^{\otimes q} \rightarrow \mathcal{H}^q$  being denoted by  $\mathcal{J}^q$ , the 1-cocycle

$$d_0^*\sigma^{(q)} - d_1^*\sigma^{(q)} \in \Gamma(U_1, \underline{\text{Hom}}_{\mathcal{O}_{X'}}(\mathcal{H}^q, \mathcal{J}^q))$$

represents the obstruction, in  $H^1(X', \underline{\text{Hom}}(\mathcal{H}^q, \mathcal{J}^q)) = \text{Ext}_{\mathcal{O}_{X'}}^1(\mathcal{H}^q, \mathcal{J}^q)$ , to the global existence of a section.



Let us calculate the class

$$\text{cl sc } \tau_{[q-1,q]}K \in H^2(X', \underline{\text{Hom}}(\mathcal{H}^q, \mathcal{H}^{q-1}))$$

in characteristic  $p \geq 2$ . For ease of notation, we denote  $\sigma^{(q)}$  simply by  $\sigma$  when no confusion is likely.

To do so, we may choose the composite morphism

$$\mathcal{H}^q \xrightarrow{\sigma^{(q)}} (\mathcal{H}^1)^{\otimes q} \xrightarrow{s^{\otimes q}} (\mathcal{Z}^1)^{\otimes q} \xrightarrow{\wedge} \mathcal{Z}^q,$$

which we denote by  $(s^{\wedge q}) \circ \sigma$ , as the section of the projection  $\mathcal{Z}^q \rightarrow \mathcal{H}^q$  over  $U_0$ .

Then one forms (the negative of) the Čech difference

$$d_1^*(s^{\wedge q}) \circ d_1^* \sigma - d_0^*(s^{\wedge q}) \circ d_0^* \sigma = [(d_1^* s)^{\wedge q} - (d_0^* s)^{\wedge q}] \circ d_0^* \sigma - (d_1^* s)^{\wedge q} \circ (d_0^* \sigma - d_1^* \sigma).$$

One notes that the second term is zero, since the image of  $d_0^* \sigma - d_1^* \sigma$  is contained in  $\mathcal{J}^q$ , which in turn is annihilated by  $(d_1^* s)^{\wedge q}$ , for the wedge product is strictly graded commutative.

Then one expresses — over  $U_1$  — the remaining first term as the differential of something:

$$\begin{aligned} & ((d_1^* s)^{\wedge q} - (d_0^* s)^{\wedge q})(\omega_1 \otimes \cdots \otimes \omega_q) \\ &= \sum_{j=1}^q (d_0^* s) \omega_1 \wedge \cdots \wedge (d_0^* s) \omega_{j-1} \wedge (d_1^* s - d_0^* s) \omega_j \wedge (d_1^* s) \omega_{j+1} \wedge \cdots \wedge (d_1^* s) \omega_q \\ &= d \sum_{j=1}^q (-1)^{j+1} h(\omega_j) (d_0^* s) \omega_1 \wedge \cdots \wedge (d_0^* s) \omega_{j-1} \wedge (d_1^* s) \omega_{j+1} \wedge \cdots \wedge (d_1^* s) \omega_q. \end{aligned}$$

One defines  $\eta = \eta^{(q)} = \eta(\omega_1 \otimes \cdots \otimes \omega_q)$  to be

$$\sum_{j=1}^q (-1)^{j+1} h(\omega_j) (d_0^* s) \omega_1 \wedge \cdots \wedge (d_0^* s) \omega_{j-1} \wedge (d_1^* s) \omega_{j+1} \wedge \cdots \wedge (d_1^* s) \omega_q$$

in order to have a commutative diagram

$$\begin{array}{ccc} \mathcal{H}^q & \xrightarrow{d_1^*(s^{\wedge q}) \circ d_1^* \sigma - d_0^*(s^{\wedge q}) \circ d_0^* \sigma} & \mathcal{Z}^q \\ & \searrow d_0^* \sigma & \nearrow d \\ & (\mathcal{H}^1)^{\otimes q} \xrightarrow{\bar{\eta}} & K^{q-1}/\mathcal{B}^{q-1} \end{array}$$

in which  $\bar{\eta}$  means the composite of  $\eta$  followed by  $K^{q-1} \rightarrow K^{q-1}/\mathcal{B}^{q-1}$ .

With this, we calculate the class of the gerbe by forming the Čech difference over  $U_2$ :

$$\begin{aligned} (d_0^* - d_1^* + d_2^*)(\bar{\eta} \circ d_0^* \sigma) &= d_0^* \bar{\eta} \circ d_{01}^* \sigma - d_1^* \bar{\eta} \circ d_{01}^* \sigma + d_2^* \bar{\eta} \circ d_{02}^* \sigma \\ &= (d_0^* - d_1^* + d_2^*) \bar{\eta} \circ d_{01}^* \sigma - d_2^* \bar{\eta} \circ (d_{01}^* \sigma - d_{02}^* \sigma). \end{aligned} \tag{4-1}$$

Let us put  $\text{obs}_1 = (d_0^* - d_1^* + d_2^*)h$ , which represents the class of  $\text{sc } \tau_{\leq 1} K$ . Then the first summand in the second line of (4-1) can be expressed in terms of  $\text{obs}_1$ :

$$\begin{aligned}
& (d_0^* - d_1^* + d_2^*)\bar{\eta}(\omega_1 \otimes \cdots \otimes \omega_q) \\
&= \sum_{j=1}^q (-1)^{j+1} d_0^* h(\omega_j) d_{01}^* s(\omega_1) \wedge \cdots \wedge d_{01}^* s(\omega_{j-1}) \wedge d_{02}^* s(\omega_{j+1}) \wedge \cdots \wedge d_{02}^* s(\omega_q) \\
&\quad - \sum_{j=1}^q (-1)^{j+1} d_1^* h(\omega_j) d_{01}^* s(\omega_1) \wedge \cdots \wedge d_{01}^* s(\omega_{j-1}) \wedge d_{12}^* s(\omega_{j+1}) \wedge \cdots \wedge d_{12}^* s(\omega_q) \\
&\quad + \sum_{j=1}^q (-1)^{j+1} d_2^* h(\omega_j) d_{02}^* s(\omega_1) \wedge \cdots \wedge d_{02}^* s(\omega_{j-1}) \wedge d_{12}^* s(\omega_{j+1}) \wedge \cdots \wedge d_{12}^* s(\omega_q) \\
&= \sum_{j=1}^q (-1)^{j+1} \text{obs}_1(\omega_j) d_{01}^* s(\omega_1) \wedge \cdots \wedge d_{01}^* s(\omega_{j-1}) \wedge d_{12}^* s(\omega_{j+1}) \wedge \cdots \wedge d_{12}^* s(\omega_q) \\
&\quad + \sum_{j=1}^q (-1)^{j+1} d_0^* h(\omega_j) d_{01}^* s(\omega_1) \wedge \cdots \wedge d_{01}^* s(\omega_{j-1}) \wedge \{ (d_{02}^* s^{\wedge(q-j)} - d_{12}^* s^{\wedge(q-j)})(\omega_{j+1} \otimes \cdots \otimes \omega_q) \} \\
&\quad + \sum_{j=1}^q (-1)^{j+1} d_2^* h(\omega_j) \{ (d_{02}^* s^{\wedge(j-1)} - d_{01}^* s^{\wedge(j-1)})(\omega_1 \otimes \cdots \otimes \omega_{j-1}) \} \wedge d_{12}^* s(\omega_{j+1}) \wedge \cdots \wedge s(\omega_q).
\end{aligned}$$

Again, as in the three equalities leading up to (3-3), the differences in the curly brackets are themselves alternating sums, and one sees that the sum of the last two alternating sums is equal to

$$-dH_2^q(d_0^*h, d_2^*h)(\omega_1 \otimes \cdots \otimes \omega_q),$$

hence is zero modulo  $\mathcal{B}^{q-1}$ . On the other hand, the first alternating sum is equal to

$$\text{ev}(\text{ctr}(\text{obs}_1), \omega_1 \wedge \cdots \wedge \omega_q).$$

Now we analyze the second summand in the second line of (4-1). It is the cup product of two cohomology classes:

$$\bar{\eta}|_{\mathcal{J}^q} \in \Gamma(U_1, \underline{\text{Hom}}(\mathcal{J}^q, \mathcal{H}^{q-1})) \text{ representing } [\bar{\eta}|_{\mathcal{J}^q}] \in \text{Ext}^1(\mathcal{J}^q, \mathcal{H}^{q-1})$$

and

$$d_0^* \sigma - d_1^* \sigma \in \Gamma(U_1, \underline{\text{Hom}}(\mathcal{H}^q, \mathcal{J}^q)) \text{ representing } [\sigma] \in \text{Ext}^1(\mathcal{H}^q, \mathcal{J}^q).$$

When  $q$  is less than  $p = \text{char}(S)$ ,  $[\sigma]$  is zero, for in this case one disposes of a canonical section of

$$(\mathcal{H}^1)^{\otimes q} \rightarrow \mathcal{H}^q,$$

namely the antisymmetrization.

On the other hand, if  $p$  is odd, then  $[\bar{\eta}|_{\mathcal{J}^q}]$  is zero, because (even more strongly)  $\bar{\eta}$  itself kills  $\mathcal{J}^q$ : for example, it maps a local section

$$\omega \otimes \omega \otimes \omega_3 \otimes \cdots \otimes \omega_q$$

of  $\mathcal{J}^q$  to the element

$$[h(\omega) d_1^* s(\omega) - h(\omega) d_0^* s(\omega)] \wedge d_1^* s(\omega_3) \wedge \cdots \wedge d_1^* s(\omega_q) = h(\omega) dh(\omega) \wedge \omega_3 \wedge \cdots \wedge \omega_q \pmod{\mathcal{B}^{q-1}},$$

which is a coboundary when 2 is invertible ( $d(h(\omega)^2) = 2h(\omega) dh(\omega)$ ).

So let us restrict our attention to the case  $p = 2$  and show that the class  $[\bar{\eta}|_{\mathcal{J}^q}]$  is still zero. First, one can easily check that  $\bar{\eta} : (\mathcal{H}^1)^{\otimes q} \rightarrow K^{q-1}/\mathcal{B}^{q-1}$ , and a fortiori  $\bar{\eta}|_{\mathcal{J}^q} : \mathcal{J}^q \rightarrow \mathcal{Z}^{q-1}/\mathcal{B}^{q-1} = \mathcal{H}^{q-1}$ , is *symmetric* in the sense that any element of the form

$$\omega_1 \otimes \cdots \otimes \omega_{j-1} \otimes (\omega_j \otimes \omega_{j+1} + \omega_{j+1} \otimes \omega_j) \otimes \omega_{j+2} \otimes \cdots \otimes \omega_q$$

(when  $1 = -1$ , adding is subtracting) maps to zero under  $\bar{\eta}$ . Therefore, one has a commutative diagram

$$\begin{array}{ccccc} \mathcal{J}^q & \longrightarrow & \mathcal{J}^q/\mathcal{J}^q & \xrightarrow{\bar{\eta}} & \mathcal{H}^{q-1} \\ \downarrow & & \downarrow & & \downarrow \\ (\mathcal{H}^1)^{\otimes q} & \longrightarrow & \text{Sym}^q(\mathcal{H}^1) & \longrightarrow & K^{q-1}/\mathcal{B}^{q-1} \end{array}$$

where the composite of the two horizontal arrows in the first row (resp. in the second row) is equal to  $\bar{\eta}|_{\mathcal{J}^q}$  (resp.  $\bar{\eta}$ ), and  $\mathcal{J}^q$  denotes the (locally free) kernel of the projection  $(\mathcal{H}^1)^{\otimes q} \rightarrow \text{Sym}^q(\mathcal{H}^1)$ .

We get a notational advantage by taking the quotient by  $\mathcal{J}^q$ : now  $\mathcal{J}^q/\mathcal{J}^q$  is generated by the images of local sections of the form

$$\omega \otimes \omega \otimes \omega_3 \otimes \cdots \otimes \omega_q. \tag{4-2}$$

Such a local section is mapped under  $\bar{\eta}$  onto

$$h(\omega) dh(\omega) \wedge d_0^* s(\omega_3) \wedge \cdots \wedge d_0^* s(\omega_q) \pmod{\mathcal{B}^{q-1}} = [h(\omega) dh(\omega) \pmod{\mathcal{B}^1}] \wedge \omega_3 \wedge \cdots \wedge \omega_q. \tag{4-3}$$

We prove that  $[\bar{\eta}|_{\mathcal{J}^q}]$  is zero by finding a 0-cochain  $z$  with coefficients in  $\text{Hom}(\mathcal{J}^q, \mathcal{H}^{q-1})$ , that is, a section of this sheaf over  $U_0$ , such that  $\bar{\eta}|_{\mathcal{J}^q} = d_0^* z - d_1^* z$ . As we know that  $\bar{\eta}|_{\mathcal{J}^q}$  factors through  $\bar{\eta} : \mathcal{J}^q/\mathcal{J}^q \rightarrow \mathcal{H}^{q-1}$ , it suffices to find  $\tilde{z} : \mathcal{J}^q/\mathcal{J}^q \rightarrow \mathcal{H}^{q-1}$  such that

$$\bar{\eta} = d_1^* \tilde{z} - d_0^* \tilde{z}.$$

But from (4-3) and the fact that  $C^{-1}W^* dh(\omega) = [h(\omega) dh(\omega)]$ , one sees that

$$\bar{\eta}(\omega \otimes \omega \otimes \omega_3 \otimes \cdots \otimes \omega_q) = (C^{-1}W^* dh(\omega)) \wedge \omega_3 \wedge \cdots \wedge \omega_q. \tag{4-4}$$

We denote here by  $W$  the base change of the absolute Frobenius endomorphism of  $S$ , so that the diagram

$$\begin{array}{ccc} X' & \xrightarrow{W} & X \\ \downarrow & & \downarrow \\ S & \xrightarrow{\text{frob}_S} & S \end{array}$$

is cartesian, by  $W^*$  the pullback morphism of differential forms

$$\Omega_{X/S}^1 \rightarrow W_* \Omega_{X'/S}^1$$

and by  $C^{-1}$  the (inverse) Cartier operation

$$\Omega_{X'/S}^1 \rightarrow \mathcal{H}^1(F_* \Omega_{X/S}^\bullet)$$

(see [Katz 1970, § 7] and recall  $p = 2$ ). Thus the last expression is the same as

$$C^{-1} W^*(d_1^* s \omega) \wedge \omega_3 \wedge \cdots \wedge \omega_q - C^{-1} W^*(d_0^* s \omega) \wedge \omega_3 \wedge \cdots \wedge \omega_q,$$

and one is led to define over  $U_0$

$$\tilde{z} : \mathcal{J}^q / \mathcal{J}^q \rightarrow \mathcal{H}^{q-1}$$

so that it maps the local section (4-2) modulo  $\mathcal{J}^q$  to

$$C^{-1} W^*(s \omega) \wedge \omega_3 \wedge \cdots \wedge \omega_q.$$

As pointed out earlier, local sections of the form (4-2) generate  $\mathcal{J}^q / \mathcal{J}^q$ , so such  $\tilde{z}$  is unique if exists at all. Now its existence can be shown locally: if one has a basis  $e_1, \dots, e_d$  of  $\mathcal{H}^1$  over  $\mathcal{O}_{X'}$ , then the images of the sections

$$\{e_{j_1} \otimes \cdots \otimes e_{j_q} : 1 \leq j_1 \leq \cdots \leq j_q \leq d \text{ with at least one repetition}\}$$

under  $\mathcal{J}^q \rightarrow \mathcal{J}^q / \mathcal{J}^q$  form a local basis of  $\mathcal{J}^q / \mathcal{J}^q$ , and then one can let  $\tilde{z}$  map the class of  $e_{j_1} \otimes \cdots \otimes e_{j_q}$  to

$$C^{-1} W^*(s(\omega_{j_k})) \wedge (\text{the rest}),$$

where  $j_k$  is an index that repeats: if two or more indices repeat, whichever one is chosen, the result is zero, and if an index repeats itself three or more times, it doesn't matter which consecutive terms are chosen, for  $1 = -1$  and the sign doesn't matter.

Then one needs to show that any local section of the form (4-2) is mapped as desired under  $\tilde{z}$  thus defined. One expresses the sections  $\omega, \omega_3, \dots, \omega_q$  as linear combinations of the  $\{e_i\}$  and one sees that it boils down to showing the linearity in each variable  $\omega_3, \dots, \omega_q$ , which is evident, as well as the linearity “in the variable  $\omega \otimes \omega$ ”, which is less so.

Let  $\omega = \alpha\xi + \beta\theta$ , where  $\alpha, \beta$  are sections of  $\mathcal{O}_{X'}$  and  $\xi, \theta$  sections of  $\mathcal{H}^1$ . Then one calculates

$$\begin{aligned} C^{-1}W^*(s(\alpha\xi + \beta\theta)) &= C^{-1}W^*(F^*\alpha \cdot s(\xi) + F^*\beta \cdot s(\theta)) \\ &= C^{-1}(\alpha^2W^*s(\xi) + \beta^2W^*s(\theta)) \\ &= (F^*\alpha)^2C^{-1}W^*s(\xi) + (F^*\beta)^2C^{-1}W^*s(\theta), \end{aligned}$$

where  $F^* : \mathcal{O}_{X'} \rightarrow F_*\mathcal{O}_X$  is the canonical pullback morphism; here one uses the fact that  $W \circ F$  is equal to the absolute Frobenius of  $X$ .

On the other hand, if one expands  $\omega \otimes \omega$  as  $\alpha^2\xi \otimes \xi + \beta^2\theta \otimes \theta + \alpha\beta(\xi \otimes \theta + \theta \otimes \xi)$ , then the last term is symmetric (i.e., lies in  $\mathcal{J}^2$ ), and hence we get the same result this way.

This can also be explained with the diagram

$$\begin{array}{c} \mathcal{Z}^1 \subseteq F_*\Omega_{X/S}^1 \xrightarrow{F_*(W^*)} F_*W_*\Omega_{X'/S}^1 \xrightarrow{F_*W_*C^{-1}} F_*W_*\mathcal{H}^1(K) = (F_{X'})_*\mathcal{H}^1(K) \\ \uparrow s \\ \mathcal{H}^1(K) \end{array}$$

over  $U_0$ , where  $F_{X'}$  denotes the absolute Frobenius of  $X'$ ; it shows that the map

$$\omega \mapsto C^{-1}W^*s(\omega)$$

is 2-linear, while “extracting”  $\omega$  out of  $\omega \otimes \omega$  would be  $2^{-1}$ -linear; hence these nonlinearities cancel each other and the map  $\omega \otimes \omega \mapsto C^{-1}W^*s(\omega)$  is linear.

This shows that  $\tilde{z}$ , hence the 0-cochain  $z$  which is obtained by composing  $\tilde{z}$  with the projection  $\mathcal{J}^q \rightarrow \mathcal{J}^q/\mathcal{J}^q$ , is well defined and has the desired property. Therefore the class  $[\tilde{\eta}]_{\mathcal{J}^q}$  is zero, and the only thing that contributes to the class (4-1) is the  $q$ -th contraction of  $\text{obs}_1$ . This ends the proof.  $\square$

By the construction (Theorem 3.9) and the calculation (Theorem 4.1), we immediately get:

**Corollary 4.2.** *With the notation as in Theorem 4.1, suppose that  $X'/S$  is liftable to  $\tilde{S}$ . Then for each integer  $q$ , the truncation  $\tau_{[q-1,q]}F_*\Omega_{X/S}^\bullet$  of length 2 is decomposable in the derived category  $D(X', \mathcal{O}_{X'})$ .*

*Proof.* This follows from [Deligne and Illusie 1987, 3.5] (which identifies the obstruction to liftability to the decomposability of  $\tau_{\leq 1}F_{X/S,*}\Omega_{X/S}^\bullet$ ) and Theorems 3.9 and 4.1 (which relate the decomposability of  $\tau_{[0,1]}F_{X/S,*}\Omega_{X/S}^\bullet$  with that of  $\tau_{[q-1,q]}F_{X/S,*}\Omega_{X/S}^\bullet$ ).  $\square$

In particular, we extend the (special) case of Corollary 3.6 applied to the de Rham complex in characteristic  $p > 2$ , even to the case of  $p = 2$ .

**Corollary 4.3.** *Let  $X$  be a smooth variety over a perfect field  $k$ . Then we have the equality*

$$\delta_{F_{X/k,*}\Omega_{X/k}^\bullet}^q = (\text{id} \otimes \delta_K^1) \circ \eta^q$$

in  $\text{Hom}(\Omega_{X'/k}^q, \Omega_{X'/k}^{q-1}[2])$ .

Finally, we answer the question of Katz:

**Corollary 4.4.** *Let  $S$  be a scheme of characteristic  $p > 0$ ,  $f : X \rightarrow S$  a smooth separated morphism of finite type, and  $f' : X' \rightarrow S$  (resp.  $F : X \rightarrow X'$ ) the base change of  $f$  by the Frobenius endomorphism of  $S$  (resp. the relative Frobenius). Suppose  $\tilde{S}$  is a flat  $\mathbb{Z}/p^2$ -scheme whose reduction modulo  $p$  yields  $S$ . Then the morphism in the conjugate spectral sequence*

$$d_2^{ij} : R^i f'_* \Omega_{X'/S}^j \rightarrow R^{i+2} f'_* \Omega_{X'/S}^{j-1},$$

where one identifies  $\mathcal{H}^j(F_* \Omega_{X/S}^\bullet)$  with  $\Omega_{X'/S}^j$  via the Cartier isomorphism, can be canonically regarded as the cup product with the additive inverse of the obstruction class (in  $H^2(X', T_{X'/S})$ ) to lifting  $X'/S$  over  $\tilde{S}$ .

*Proof.* We first remark that by [Deligne and Illusie 1987, 3.9] it can be directly seen that the obstruction class to lifting does not depend on the choice of a flat  $\mathbb{Z}/p^2$ -lifting  $\tilde{S}$  of  $S$ . Then the corollary follows from [Deligne and Illusie 1987, 3.5] and Theorems 3.9 and 4.1. □

### Appendix: $F$ -split schemes of dimension $p + 1$

Let  $k$  be a perfect field of characteristic  $p > 0$ . As mentioned in the introduction, Drinfeld, Bhatt–Lurie, and Li–Mondal (see [Li and Mondal 2021, Remark 5.7] for context) have obtained the following result.

**Theorem A.1** [Drinfeld 2020, § 5.12.1; Bhatt and Lurie 2022, Remark 4.7.18; Li and Mondal 2021, Corollary 5.5]. *Let  $X$  be a smooth scheme over a perfect field  $k$  of characteristic  $p > 0$ . Suppose that  $X$  is liftable to  $W_2(k)$ . Then the truncations*

$$\tau_{[q-p+1, q]} F_{X/k, *} \Omega_{X/k}^\bullet$$

are decomposable for all  $q$ .

Below, we employ this in order to show Kodaira–Akizuki–Nakano vanishing and Hodge–de Rham degeneration for  $F$ -split smooth projective schemes of dimension at most  $p + 1$ .

Recall [Mehta and Ramanathan 1985] that a  $k$ -scheme  $X$  is  $F$ -split if the morphism  $F_X^* : \mathcal{O}_X \rightarrow F_{X, *} \mathcal{O}_X$  is a split injection. Since  $k$  is perfect, this is equivalent to the splitting of  $F_{X/k}^* : \mathcal{O}_{X'} \rightarrow F_{X/k, *} \mathcal{O}_X$ . It is well known that every  $F$ -split scheme over  $k$  admits a flat lifting to  $W_2(k)$  (see [Illusie 1996, § 8.5] for the smooth case or [Langer 2015, § 8, Proposition 4] for the general case).

If  $X$  is  $F$ -split and if  $L$  is a line bundle on  $X$ , then tensoring the split injection  $\mathcal{O}_X \rightarrow F_{X, *} \mathcal{O}_X$  with  $L$  and taking cohomology shows that for all  $i$ ,  $H^i(X, L)$  is a direct summand of  $H^i(X, L \otimes F_{X, *} \mathcal{O}_X)$ . By the projection formula and the fact that Frobenius is affine this latter summand equals  $H^i(X, F_X^* L) = H^i(X, L^p)$ , and hence the Frobenius pullback maps

$$F_X^* : H^i(X, L) \rightarrow H^i(X, L^p)$$

are injective. Thus, if  $H^i(X, L^m) = 0$  for  $m \gg 0$ , then already  $H^i(X, L) = 0$ . Consequently, if  $X$  is moreover smooth (or just Gorenstein) and projective, then  $H^i(X, L^{-1}) = 0$  for  $i < \dim X$  and  $L$  ample,

that is, Kodaira vanishing holds on  $X$ . Similar reasoning with  $L = \mathcal{O}_X$  shows that

$$F_X^* : H^i(X, \mathcal{O}_X) \rightarrow H^i(X, \mathcal{O}_X)$$

is bijective for all  $i \geq 0$ .

**Theorem A.2** (Kodaira–Akizuki–Nakano vanishing). *Let  $X$  be a smooth projective scheme over  $k$  of dimension  $d = p + 1$ . If  $X$  is  $F$ -split, then Kodaira–Akizuki–Nakano vanishing holds for  $X$ , i.e., for every ample line bundle  $L$ , we have*

$$H^i(X, L^{-1} \otimes \Omega_{X/k}^j) = 0 \quad \text{for } i + j < d = p + 1.$$

*Proof.* By Serre vanishing, the assertion holds for  $L^{p^m}$  for  $m \gg 0$ . Therefore we may assume that it holds for  $L^p$ . Following the proof of [Deligne and Illusie 1987, lemme 2.9], we form the complex  $K^\bullet = (L')^{-1} \otimes F_{X/k,*} \Omega_{X/k}^\bullet$ , where  $L'$  is the pullback of  $L$  to  $X'$ , and write the two spectral sequences

$${}_I E_1^{ij} = H^j(X', (L')^{-1} \otimes F_{X/k,*} \Omega_{X/k}^i) \Rightarrow H^{i+j}(X', K^\bullet) \tag{A-1}$$

and

$${}_{II} E_2^{ij} = H^i(X', (L')^{-1} \otimes \Omega_{X'/k}^j) \Rightarrow H^{i+j}(X', K^\bullet). \tag{A-2}$$

Now the projection formula gives  ${}_I E_1^{ij} = H^j(X, L^{-p} \otimes \Omega_{X/k}^i)$ , which vanishes for  $i + j \leq p$  by assumption. Consequently the abutment  $H^r(X', K^\bullet)$  vanishes for  $r \leq p$ .

We now investigate the second spectral sequence. Since  $X$  is  $F$ -split, it lifts to  $W_2(k)$ . Theorem A.1 implies that the differentials on  ${}_{II} E_r^{ij}$  are zero for  $r \leq p$ . For dimensional reasons, there are no nonzero differentials for  $r > p + 1$ , and the only two nonzero differentials on  ${}_{II} E_{p+1}^{ij}$  are

$$d_{p+1}^{0,p} : H^0(X', (L')^{-1} \otimes \Omega_{X'/k}^p) \rightarrow H^{p+1}(X', (L')^{-1})$$

and

$$d_{p+1}^{0,p+1} : H^0(X', (L')^{-1} \otimes \omega_{X'/k}) \rightarrow H^{p+1}(X', (L')^{-1} \otimes \Omega_{X'/k}^1).$$

We will show that  $d_{p+1}^{0,p} = 0$ , which will then imply that in (A-2) we have

$${}_{II} E_2^{ij} = {}_{II} E_{p+1}^{ij} = {}_{II} E_\infty^{ij} = 0 \quad \text{for } i + j \leq p.$$

Note that  $H^p(X', K^\bullet) = 0$  implies  $0 = {}_{II} E_{p+2}^{0,p} = \ker(d_{p+1}^{0,p})$ , i.e.,  $d_{p+1}^{0,p}$  is injective.

By Lemma A.3 below applied to  $E = L^{-1}$  and the map  $F_{X^-/k} : X^- \rightarrow X$ , where  $X^- = (F_k)^{-1}(X)$  is the Frobenius *untwist* of  $X$ , we have a commutative square:

$$\begin{array}{ccc} H^0(X', (L')^{-1} \otimes \Omega_{X'/k}^p) & \xrightarrow{d_{p+1}^{0,p}(L)} & H^{p+1}(X', (L')^{-1}) \\ \downarrow & & \downarrow F_{X'/k}^* \\ 0 = H^0(X, L^{-p} \otimes \Omega_{X/k}^p) & \xrightarrow{d_{p+1}^{0,p}(F_{X^-/k}^* L^{-1})} & H^{p+1}(X, L^{-p}) \end{array}$$

(In fact the left vertical map is zero.) Here we use the identification  $(F_{X^-/k}^*(L^{-1}))' = F_{X/k}^*(L') - 1 = L^{-p}$  and the bottom map is deduced similarly for (the pullback to  $X^-$  of)  $L^p$  in place of  $L$ . The vertical right map is injective because  $X$  is  $F$ -split and we have a commutative diagram

$$\begin{array}{ccccc}
 & & & & F_X^* \\
 & & & \curvearrowright & \\
 H^{p+1}(X, L^{-1}) & \xrightarrow{\sim} & H^{p+1}(X', (L')^{-1}) & \xrightarrow{F_{X/k}^*} & H^{p+1}(X, L^{-p}) \\
 & W^* & & & 
 \end{array}$$

( $W : X' \rightarrow X$  being the projection), and the horizontal maps are injective by the previous paragraph. We conclude that  $H^0(X', (L')^{-1}) \otimes \Omega_{X'/k}^p = 0$ . □

In the proof above, as well as in the proof of Hodge–de Rham degeneration below, we need the following functoriality result.

**Lemma A.3.** *For a vector bundle  $E$  on a smooth  $k$ -scheme  $X$ , write  $K^\bullet(E)$  to denote the complex  $E' \otimes F_{X/k,*} \Omega_{X/k}^\bullet$ . Let  $f : Y \rightarrow X$  be a map of smooth  $k$ -schemes. Then  $f$  induces a map of complexes  $f^* K^\bullet(E) \rightarrow K^\bullet(f^* E)$  and hence a map of spectral sequences*

$$\begin{array}{ccc}
 {}_{II} E_2^{ij}(E) = H^i(X', E' \otimes \Omega_{X'/k}^j) & \Rightarrow & H^{i+j}(X', K^\bullet(E)) \\
 \downarrow & & \downarrow \\
 {}_{II} E_2^{ij}(f^* E) = H^i(Y', (f^* E)' \otimes \Omega_{Y'/k}^j) & \Rightarrow & H^{i+j}(Y', K^\bullet(f^* E))
 \end{array}$$

where the maps  $H^i(X', E' \otimes \Omega_{X'/k}^j) \rightarrow H^i(Y', (f^* E)' \otimes \Omega_{Y'/k}^j)$  are induced by the composition

$$H^i(X', E' \otimes \Omega_{X'/k}^j) \rightarrow H^i(Y', (f')^*(E') \otimes (f')^* \Omega_{X'/k}^j) \rightarrow H^i(Y', (f')^*(E') \otimes \Omega_{Y'/k}^j).$$

*Proof.* This follows from the commutative diagram below:

$$\begin{array}{ccccc}
 Y & \longrightarrow & Y' & \longrightarrow & Y \\
 \downarrow f & & \downarrow f' & & \downarrow f \\
 X & \longrightarrow & X' & \longrightarrow & X \\
 & \searrow & \downarrow & & \downarrow \\
 & & \text{Spec}(k) & \xrightarrow{F_k} & \text{Spec}(k)
 \end{array}$$

Note that  $(f')^*(E') \simeq (f^* E)'$ . □

**Theorem A.4** (Hodge–de Rham degeneration). *Let  $X$  be a smooth projective scheme over  $k$  of dimension  $d = p + 1$ . If  $X$  is  $F$ -split, then the Hodge to de Rham spectral sequence*

$${}_I E_1^{ij} = H^j(X, \Omega_{X/k}^i) \Rightarrow H^{i+j}(X, \Omega_{X/k}^\bullet)$$

*degenerates.*



*Proof.* Since  $X$  is proper, it is enough to show that the conjugate spectral sequence

$${}_{II}E_2^{ij} = H^i(X', \Omega_{X'/k}^j) \Rightarrow H^{i+j}(X, \Omega_{X/k}^\bullet)$$

degenerates. Since  $X$  is  $F$ -split, it lifts to  $W_2(k)$ , and then Theorem A.1 implies that the differentials on the page  ${}_{II}E_r^{ij}$  of the conjugate spectral sequence are zero for  $r \leq p$ .

Since  $\dim X = p + 1$ , the only possibly nonzero differentials in this spectral sequence are therefore

$$d_{p+1}^{0,p} : H^0(X', \Omega_{X'/k}^p) \rightarrow H^{p+1}(X', \mathcal{O}_{X'})$$

and

$$d_{p+1}^{0,p+1} : H^0(X', \omega_{X'/k}) \rightarrow H^{p+1}(X', \Omega_{X'/k}^1).$$

We will show that  $d_{p+1}^{0,p} = 0$ . Indeed, functoriality of the above maps with respect to Frobenius (Lemma A.3 with  $E = \mathcal{O}_X$  and the relative Frobenius  $F_{X^-/k}$ ) gives a commutative square

$$\begin{array}{ccc} H^0(X', \Omega_{X'/k}^p) & \xrightarrow{d_{p+1}^{0,p}} & H^{p+1}(X', \mathcal{O}_{X'}) \\ F_{X'/k}^* \downarrow & & \downarrow F_{X'/k}^* \\ H^0(X, \Omega_{X/k}^p) & \xrightarrow{d_{p+1}^{0,p} \text{ for } X^-} & H^{p+1}(X, \mathcal{O}_X) \end{array}$$

where  $X^- = (F_k^{-1})^* X$  is again the Frobenius untwist of  $X$ . Since the Frobenius is zero on  $\Omega^i$  for  $i > 0$ , the left vertical map is zero. On the other hand, since  $X$  is  $F$ -split, the right vertical map is an isomorphism. Therefore the top map  $d_{p+1}^{0,p}$  is zero.

Finally, we obtain the vanishing of  $d_{p+1}^{0,p+1}$  by comparing dimensions and duality. Indeed, we have

$$\begin{aligned} \dim H^{p+2}(X, \Omega_{X/k}^\bullet) &= \dim H^p(X, \Omega_{X/k}^\bullet) && \text{(Poincaré duality)} \\ &= \sum_{i+j=p} \dim H^i(X', \Omega_{X'/k}^j) && \text{(since } d_{p+1}^{0,p} = 0) \\ &= \sum_{i+j=p+2} \dim H^i(X', \Omega_{X'/k}^j) && \text{(Serre duality),} \end{aligned}$$

so  $d_{p+1}^{0,p+1} = 0$ . □

**Remark A.5** (see [Li and Mondal 2021, Corollary 5.6] and [Bhatt and Lurie 2022]). In fact, the results of Drinfeld, Bhatt–Lurie, and Li–Mondal yield more than we have stated in Theorem A.1. Namely, for  $X$  smooth over  $k$  and liftable to  $W_2(k)$ , there exists a decomposition in the derived category

$$F_{X/k,*} \Omega_{X/k}^\bullet \simeq \bigoplus_{i=0}^{p-1} K_i$$

where  $\mathcal{H}^j(K_i) = 0$  unless  $i$  and  $j$  are congruent modulo  $p$ . This implies that in the conjugate spectral sequence, as well as in the second spectral sequence used in the proof of Theorem A.2, nonzero differentials may appear only on pages  $E_r$  where  $r$  is congruent to one modulo  $p$ . We only used this with  $r \leq p$ , and it would be interesting to obtain new vanishing and degeneration theorems using this stronger fact.

### Acknowledgements

Achinger is grateful to Luc Illusie, Shizhang Li, Arthur Ogus, and Vadim Vologodsky for useful comments and stimulating discussions. Suh thanks Nicholas Katz for raising the question that we answer in Theorem 1.3, and Pierre Deligne and Luc Illusie for helpful comments. Both authors thank the referees for helpful comments and suggestions.

Achinger was supported by NCN SONATA grant number UM0-2017/26/D/ST1/00913.

### References

- [Achinger and Ogus 2020] P. Achinger and A. Ogus, “Monodromy and log geometry”, *Tunis. J. Math.* **2**:3 (2020), 455–534. MR Zbl
- [Bhatt and Lurie 2022] B. Bhatt and J. Lurie, “Absolute prismatic cohomology”, preprint, 2022. arXiv 2201.06120
- [Deligne and Illusie 1987] P. Deligne and L. Illusie, “Relèvements modulo  $p^2$  et décomposition du complexe de de Rham”, *Invent. Math.* **89**:2 (1987), 247–270. MR Zbl
- [Drinfeld 2020] V. Drinfeld, “Prismatization”, preprint, 2020. arXiv 2005.04746
- [Illusie 1971] L. Illusie, *Complexe cotangent et déformations, I*, Lecture Notes in Mathematics **239**, Springer, 1971. MR Zbl
- [Illusie 1972] L. Illusie, *Complexe cotangent et déformations, II*, Lecture Notes in Mathematics **283**, Springer, 1972. MR Zbl
- [Illusie 1996] L. Illusie, “Frobenius et dégénérescence de Hodge”, pp. 113–168 in *Introduction à la théorie de Hodge*, Panor. Synthèses **3**, Soc. Math. France, Paris, 1996. MR Zbl
- [Kato 1989] K. Kato, “Logarithmic structures of Fontaine–Illusie”, pp. 191–224 in *Algebraic analysis, geometry, and number theory* (Baltimore, MD, 1988), edited by J.-I. Igusa, Johns Hopkins Univ. Press, Baltimore, MD, 1989. MR Zbl
- [Kato and Saito 2004] K. Kato and T. Saito, “On the conductor formula of Bloch”, *Publ. Math. Inst. Hautes Études Sci.* **100** (2004), 5–151. MR Zbl
- [Katz 1970] N. M. Katz, “Nilpotent connections and the monodromy theorem: applications of a result of Turrittin”, *Inst. Hautes Études Sci. Publ. Math.* **39** (1970), 175–232. MR Zbl
- [Kříž and May 1995] I. Kříž and J. P. May, *Operads, algebras, modules and motives*, Astérisque **233**, 1995. MR Zbl
- [Langer 2015] A. Langer, “Bogomolov’s inequality for Higgs sheaves in positive characteristic”, *Invent. Math.* **199**:3 (2015), 889–920. MR Zbl
- [Li and Mondal 2021] S. Li and S. Mondal, “On endomorphisms of the de Rham cohomology functor”, preprint, 2021. To appear in *Geom. Topol.* arXiv 2109.04303
- [Lurie 2009] J. Lurie, *Higher topos theory*, Annals of Mathematics Studies **170**, Princeton Univ. Press, Princeton, NJ, 2009. MR Zbl
- [Mehta and Ramanathan 1985] V. B. Mehta and A. Ramanathan, “Frobenius splitting and cohomology vanishing for Schubert varieties”, *Ann. of Math. (2)* **122**:1 (1985), 27–40. MR Zbl
- [Petrov 2023] A. Petrov, “Non-decomposability of the de Rham complex and non-semisimplicity of the Sen operator”, preprint, 2023. arXiv 2302.11389
- [SGA 4<sub>2</sub> 1972] M. Artin, A. Grothendieck, and J. L. Verdier, *Théorie des topos et cohomologie étale des schémas, Tome 2: Exposés V–VIII* (Séminaire de Géométrie Algébrique du Bois Marie 1963–1964), Lecture Notes in Math. **270**, Springer, 1972. MR Zbl
- [Steenbrink 1995] J. H. M. Steenbrink, “Logarithmic embeddings of varieties with normal crossings and mixed Hodge structures”, *Math. Ann.* **301**:1 (1995), 105–118. MR Zbl

Communicated by Bhargav Bhatt

Received 2021-09-13    Revised 2022-02-10    Accepted 2022-03-25

pachinger@impan.pl

*Institute of Mathematics of the Polish Academy of Sciences, Warsaw, Poland*

jusuh@ucsc.edu

*Mathematics Department, University of California, Santa Cruz, CA, United States*

# Algebra & Number Theory

msp.org/ant

## EDITORS

MANAGING EDITOR  
Antoine Chambert-Loir  
Université Paris-Diderot  
France

EDITORIAL BOARD CHAIR  
David Eisenbud  
University of California  
Berkeley, USA

## BOARD OF EDITORS

Jason P. Bell	University of Waterloo, Canada	Philippe Michel	École Polytechnique Fédérale de Lausanne
Bhargav Bhatt	University of Michigan, USA	Martin Olsson	University of California, Berkeley, USA
Frank Calegari	University of Chicago, USA	Irena Peeva	Cornell University, USA
J.-L. Colliot-Thélène	CNRS, Université Paris-Saclay, France	Jonathan Pila	University of Oxford, UK
Brian D. Conrad	Stanford University, USA	Anand Pillay	University of Notre Dame, USA
Samit Dasgupta	Duke University, USA	Bjorn Poonen	Massachusetts Institute of Technology, USA
Hélène Esnault	Freie Universität Berlin, Germany	Victor Reiner	University of Minnesota, USA
Gavril Farkas	Humboldt Universität zu Berlin, Germany	Peter Sarnak	Princeton University, USA
Sergey Fomin	University of Michigan, USA	Michael Singer	North Carolina State University, USA
Edward Frenkel	University of California, Berkeley, USA	Vasudevan Srinivas	Tata Inst. of Fund. Research, India
Wee Teck Gan	National University of Singapore	Shunsuke Takagi	University of Tokyo, Japan
Andrew Granville	Université de Montréal, Canada	Pham Huu Tiep	Rutgers University, USA
Ben J. Green	University of Oxford, UK	Ravi Vakil	Stanford University, USA
Christopher Hacon	University of Utah, USA	Akshay Venkatesh	Institute for Advanced Study, USA
Roger Heath-Brown	Oxford University, UK	Melanie Matchett Wood	Harvard University, USA
János Kollár	Princeton University, USA	Shou-Wu Zhang	Princeton University, USA
Michael J. Larsen	Indiana University Bloomington, USA		

## PRODUCTION

production@msp.org  
Silvio Levy, Scientific Editor

---

See inside back cover or [msp.org/ant](http://msp.org/ant) for submission instructions.

---

The subscription price for 2023 is US \$485/year for the electronic version, and \$705/year (+\$65, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to MSP.

---

Algebra & Number Theory (ISSN 1944-7833 electronic, 1937-0652 printed) at Mathematical Sciences Publishers, 798 Evans Hall #3840, c/o University of California, Berkeley, CA 94720-3840 is published continuously online.

---

ANT peer review and production are managed by EditFLOW<sup>®</sup> from MSP.

PUBLISHED BY

 **mathematical sciences publishers**  
nonprofit scientific publishing

<http://msp.org/>

© 2023 Mathematical Sciences Publishers

# Algebra & Number Theory

Volume 17 No. 2 2023

---

Torsion points on elliptic curves over number fields of small degree MAARTEN DERICKX, SHELDON KAMIENNY, WILLIAM STEIN and MICHAEL STOLL	267
Tame fundamental groups of pure pairs and Abhyankar's lemma JAVIER CARVAJAL-ROJAS and AXEL STÄBLER	309
Constructions of difference sets in nonabelian 2-groups T. APPLEBAUM, J. CLIKEMAN, J. A. DAVIS, J. F. DILLON, J. JEDWAB, T. RABBANI, K. SMITH and W. YOLLAND	359
The principal block of a $\mathbb{Z}_\ell$ -spets and Yokonuma type algebras RADHA KESSAR, GUNTER MALLE and JASON SEMERARO	397
Geometric properties of the Kazhdan–Lusztig Schubert basis CRISTIAN LENART, CHANGJIAN SU, KIRILL ZAINOULLINE and CHANGLONG ZHONG	435
Some refinements of the Deligne–Illusie theorem PIOTR ACHINGER and JUNECUE SUH	465
A transference principle for systems of linear equations, and applications to almost twin primes PIERRE-YVES BIENVENU, XUANCHENG SHAO and JONI TERÄVÄINEN	497