

# *Algebra & Number Theory*

Volume 18

2024

No. 11

**The distribution of large quadratic character sums  
and applications**

Youness Lamzouri





# The distribution of large quadratic character sums and applications

Youness Lamzouri

*Dedicated to Andrew Granville on the occasion of his 60th birthday*

We investigate the distribution of the maximum of character sums over the family of primitive quadratic characters attached to fundamental discriminants  $|d| \leq x$ . In particular, our work improves results of Montgomery and Vaughan, and gives strong evidence that the Omega result of Bateman and Chowla for quadratic character sums is optimal. We also obtain similar results for real characters with prime discriminants up to  $x$ , and deduce the interesting consequence that almost all primes with large Legendre symbol sums are congruent to 3 modulo 4. Our results are motivated by a recent work of Bober, Goldmakher, Granville and Koukoulopoulos, who proved similar results for the family of nonprincipal characters modulo a large prime. However, their method does not seem to generalize to other families of Dirichlet characters. Instead, we use a different and more streamlined approach, which relies mainly on the quadratic large sieve. As an application, we consider a question of Montgomery concerning the positivity of sums of Legendre symbols.

## 1. Introduction

Character sums play a central role in modern number theory through their numerous applications in the study of various arithmetic, analytic, algebraic and geometric objects. One important and basic example is that of quadratic Dirichlet characters, which include the Legendre symbol. The study of such characters and related sums has a long and rich history stretching back to the work of Gauss on binary quadratic forms. Let  $\chi$  be a Dirichlet character modulo  $q$ . One quantity that was extensively studied over the past century is

$$M(\chi) := \max_{t \leq q} \left| \sum_{n \leq t} \chi(n) \right|.$$

In 1918, Pólya and Vinogradov independently proved that

$$M(\chi) \ll \sqrt{q} \log q.$$

---

*MSC2020:* primary 11L40, 11N64; secondary 11K65.

*Keywords:* character sums, Dirichlet L-functions, Pólya–Vinogradov inequality, quadratic large sieve, Kronecker symbol, Legendre symbol, positivity of partial sums.

On the other hand, an easy argument (based on applying Parseval's theorem to the Pólya Fourier series (2-1) attached to  $\chi$ ) shows that  $M(\chi) \gg \sqrt{q}$  for all primitive characters modulo  $q$ . Though one can establish the Pólya–Vinogradov inequality using only basic Fourier analysis, improving on it has proved to be a very difficult problem, and resisted substantial progress outside of special cases. We should also note that any such improvement would have important consequences on several important quantities in analytic number theory, including class numbers of quadratic fields, short character sums, and the least quadratic nonresidue (see, for example, [Bober and Goldmakher 2016; Granville and Mangerel 2023; Mangerel 2020]). Granville and Soundararajan [2007] made an important breakthrough by showing that the Pólya–Vinogradov inequality can substantially be improved for characters of a fixed odd order. Further improvements to this case were obtained by Goldmakher [2012], and Lamzouri and Mangerel [2022].

Since  $\sqrt{q} \ll M(\chi) \ll \sqrt{q} \log q$  for all primitive characters modulo  $q$ , a natural question is to determine the maximal order of  $M(\chi)$ . Assuming the generalized Riemann hypothesis (GRH), Montgomery and Vaughan [1977] proved that  $M(\chi) \ll \sqrt{q} \log \log q$ . This turns out to be optimal (up to a constant factor) in view of an older result of Paley [1932] who proved the existence of an infinite family of quadratic characters for which  $M(\chi)$  is that large. Granville and Soundararajan [2007] refined Montgomery and Vaughan's conditional result and showed that

$$M(\chi) \leq (2C_\chi + o(1))\sqrt{q} \log \log q, \quad (1-1)$$

for all primitive characters  $\chi$  modulo  $q$ , where  $C_\chi = e^\gamma/\pi$  if  $\chi$  is odd, and  $C_\chi = e^\gamma/(\sqrt{3}\pi)$  if  $\chi$  is even. On the other hand, Paley's result was refined by Bateman and Chowla [1950], who proved the existence of an infinite sequence of moduli  $q$ , and primitive quadratic characters  $\chi \pmod{q}$ , such that

$$M(\chi) \geq \left(\frac{e^\gamma}{\pi} + o(1)\right)\sqrt{q} \log \log q. \quad (1-2)$$

This result was extended to the family of primitive characters modulo a large prime  $q$  by several authors (see, for example, Theorem 3 of [Granville and Soundararajan 2007]). Finally, we should note that Granville and Soundararajan [2007] conjectured that Bateman and Chowla's Omega result should correspond to the true extreme values of  $M(\chi)$ , namely that

$$M(\chi) \leq (C_\chi + o(1))\sqrt{q} \log \log q, \quad (1-3)$$

for all primitive characters  $\chi$ .

**1A. The distribution of character sums.** In view of (1-1), (1-2) and (1-3) it is natural to renormalize  $M(\chi)$  by defining

$$m(\chi) := \frac{e^{-\gamma}\pi}{\sqrt{q}} M(\chi).$$

Montgomery and Vaughan [1979] were the first to study the distribution of  $m(\chi)$  over families of Dirichlet characters. In particular, they showed that  $m(\chi)$  is bounded (and hence  $M(\chi) \ll \sqrt{q}$ ) for most characters.

Let  $q$  be a large prime and

$$\Phi_q(\tau) := \frac{1}{\varphi(q)} |\{\chi \neq \chi_0 \pmod{q} : m(\chi) > \tau\}|,$$

where  $\varphi(q)$  is Euler’s totient function. It follows from [Montgomery and Vaughan 1979] that

$$\Phi_q(\tau) \ll_A \tau^{-A},$$

for any constant  $A \geq 1$ . This estimate was improved by Bober and Goldmakher [2013] for fixed  $\tau$ , and subsequently by Bober, Goldmakher, Granville and Koukoulopoulos [Bober et al. 2018] who showed that uniformly for  $2 \leq \tau \leq \log \log q - M$  (where  $M \geq 4$  is a parameter) we have

$$\exp\left(-\frac{e^{\tau+A_0-\eta}}{\tau}(1 + O(E_1(\tau, M)))\right) \leq \Phi_q(\tau) \leq \exp\left(-\frac{e^{\tau-2-\eta}}{\tau}(1 + O(E_2(\tau)))\right), \tag{1-4}$$

where  $E_1(\tau, M) = (\log \tau)^2 / \sqrt{\tau} + e^{-M/2}$ ,  $E_2(\tau) = (\log \tau) / \tau$ ,  $\eta := e^{-\gamma} \log 2$ , and  $A_0 = 0.088546\dots$  is an explicit constant which can be expressed as a sum of integrals over the modified Bessel function of the first kind. In particular, this result gives strong evidence to the Granville–Soundararajan conjecture (1-3) for odd primitive characters modulo a large prime  $q$ .

Although the family of quadratic characters was the first for which large character sums were exhibited by Paley [1932], and then later by Bateman and Chowla [1950], no such distribution results are known in this case. In fact, the only known result for real characters is a result of Montgomery and Vaughan [1979] who showed that  $\max_t |\sum_{n \leq t} (\frac{n}{p})| \ll \sqrt{p}$  for most primes  $p \leq x$ . The main reason which explains why the results of [Bober and Goldmakher 2013; Bober et al. 2018] do not carry over to this setting is the fact that they rely heavily on the orthogonality relations for characters modulo  $q$ . Indeed, the key ingredient in the proof of (1-4) is to estimate the off-diagonal terms when bounding large moments of the tail of the sum in Pólya’s Fourier expansion (2-1) below, which the authors of [Bober et al. 2018] successfully achieved using intricate estimates involving divisor functions.

In this paper, we overcome this problem for the family of quadratic characters by using a different and more streamlined approach which relies on the quadratic large sieve. Before stating our results we need some notation. For a fundamental discriminant  $d$  we let  $\chi_d(\cdot) = (\frac{\cdot}{d})$  be the Kronecker symbol modulo  $|d|$ . It is useful here to consider the cases of positive and negative discriminants separately, since as Theorems 1.1 and 1.2 below show, the distribution of large values of  $m(\chi_d)$  behaves differently in each case. The difference between these cases lies in the fact that the character  $\chi_d$  is even if  $d$  is positive, and is odd if  $d$  is negative. Thus, in view of Conjecture (1-3) we expect the extreme values of  $m(\chi_d)$  for  $d > 0$  to be smaller by a factor of  $\sqrt{3}$  compared to the case  $d < 0$ . We denote by  $\mathcal{F}(x)$  the set of fundamental discriminants  $d$  such that  $|d| \leq x$ , and let  $\mathcal{F}^+(x)$  (respectively  $\mathcal{F}^-(x)$ ) be the subset of  $\mathcal{F}(x)$  consisting of positive (respectively negative) discriminants. Then we have the following standard estimates (see, for example, Lemma 4.1 of [Granville and Soundararajan 2006])

$$|\mathcal{F}^\pm(x)| = \frac{3}{\pi^2}x + O_\varepsilon(x^{1/2+\varepsilon}).$$

Our goal is to estimate the distribution functions

$$\Psi_x^\pm(\tau) := \frac{1}{|\mathcal{F}^\pm(x)|} |\{d \in \mathcal{F}^\pm(x) : m(\chi_d) > \tau\}|,$$

uniformly for  $\tau$  in the range  $2 \leq \tau \leq (1 + o(1)) \log \log x$  in the case of  $\Psi_x^-(\tau)$ , and the range  $2 \leq \tau \leq (1/\sqrt{3} + o(1)) \log \log x$  in the case of  $\Psi_x^+(\tau)$ . Conjecture (1-3) implies that these ranges are best possible up to the term  $o(\log \log x)$ . Here and throughout we shall denote by  $\log_k$  the  $k$ -th iteration of the natural logarithm. We prove the following results.

**Theorem 1.1.** *Let  $\eta = e^{-\gamma} \log 2$ , and  $x$  be a large real number. Uniformly for  $\tau$  in the range  $2 \leq \tau \leq \log_2 x + \log_5 x - \log_4 x - C$  (where  $C > 0$  is a suitably large constant) we have*

$$\exp\left(-\frac{e^{\tau-\eta-B_0}}{\tau} \left(1 + O\left(\frac{(\log \tau)^2}{\sqrt{\tau}}\right)\right)\right) \leq \Psi_x^-(\tau) \leq \exp\left(-\frac{e^{\tau-\eta-\log 2-2}}{\tau} \left(1 + O\left(\frac{\log \tau}{\tau}\right)\right)\right),$$

where

$$B_0 = \int_0^1 \frac{\tanh y}{y} dy + \int_1^\infty \frac{\tanh y - 1}{y} dy = 0.8187\dots \quad (1-5)$$

**Theorem 1.2.** *Let  $B_0$  be the constant in Theorem 1.1. There exists positive constants  $C_1$  and  $C_2$  such that uniformly for  $\tau$  in the range  $2 \leq \tau \leq (\log_2 x + \log_5 x - \log_4 x - C_1)/\sqrt{3}$  we have*

$$\exp\left(-\frac{e^{\sqrt{3}\tau-B_0}}{\sqrt{3}\tau} \left(1 + O\left(\frac{1}{\tau}\right)\right)\right) \leq \Psi_x^+(\tau) \ll \exp\left(-\frac{e^{\sqrt{3}\tau}}{\tau^{C_2}}\right).$$

**Remark 1.3.** Theorems 1.1 and 1.2 (the latter is the analogue of Theorem 1.3 of [Bober et al. 2018]) give strong evidence to the Granville–Soundararajan conjecture (1-3) for the family of quadratic characters. Moreover, a direct consequence of these results is the fact that almost all fundamental discriminants  $|d| \leq x$  for which  $M(\chi_d)$  is large are negative.

**Remark 1.4.** The lower bounds in (1-4) and Theorems 1.1 and 1.2 are consequences of the works of Granville and Soundararajan [2003; 2006] on the distribution of  $|L(1, \chi)|$  (over nonprincipal characters modulo  $q$ ) in the case of (1-4), and  $L(1, \chi_d)$  (over fundamental discriminants  $|d| \leq x$ ) in the case of Theorems 1.1 and 1.2 (see Section 2 for more details). Moreover, note that we obtain a slightly different constant in the upper bound of Theorem 1.1 compared to (1-4). We believe that this is caused by the different nature of the family of quadratic characters. Indeed in our case, the difference between the constants in the upper and lower bounds of Theorem 1.1 is  $2 + \log 2 - B_0 \approx 1.8744$ , which is slightly smaller than the analogous difference  $2 + A_0 \approx 2.0885$  for the family of nonprincipal characters modulo a large prime  $q$  in (1-4). Finally, we note that by using our approach, we can derive an easier proof of the upper bound of (1-4) in the range  $2 \leq \tau \leq \log_2 q + \log_5 q - \log_4 q - C$ . This is achieved by following the exact same argument of the proof of the upper bound of Theorem 1.1, and replacing the quadratic

large sieve inequalities of Heath-Brown and Elliott (see Lemma 4.1 below) by the following large sieve estimate of [Montgomery 1971, Theorem 6.2]

$$\sum_{\chi \bmod q} \left| \sum_{n \leq N} a_n \chi(n) \right|^2 \ll (q + N) \sum_{n \leq N} |a_n|^2,$$

which holds for an arbitrary complex sequence  $\{a_n\}_{n \geq 1}$ , and all integers  $q \geq 2$ . Although our approach gives a slightly smaller range of  $\tau$  in this case, it has the advantage of extending the upper bound of (1-4) to all moduli  $q$ .

**Remark 1.5.** Using our work we can show that the structure results for large character sums obtained by Bober, Goldmakher, Granville and Koukoulopoulos in Section 2 of [Bober et al. 2018] for the family of nonprincipal characters modulo a large prime  $q$ , hold verbatim for the family of quadratic characters  $\chi_d$  attached to fundamental discriminants  $|d| \leq x$ . Since these results are technical, and the statements are exactly the same, we prefer to not state them here and refer the reader to the exact statements in [Bober et al. 2018]. The proofs follow along the same lines of [Bober et al. 2018] by using the auxiliary lemmas therein which hold for all primitive characters (and are derived using the “pretentious” theory of character sums developed by Granville and Soundararajan [2007]), and replacing the ingredients of the proof of Theorem 1.1 in [Bober et al. 2018] by those of the proof of Theorem 1.1 of our paper.

**1B. Analogous results for prime discriminants.** Using our approach we establish similar results to Theorems 1.1 and 1.2 over prime discriminants. The analogous lower bounds are direct consequences of newly established results on the distribution of  $L(1, (\frac{\cdot}{p}))$ , which we shall describe in the next section. Furthermore, the analogous upper bounds will be obtained using the same methods of proofs of Theorems 1.1 and 1.2, together with the large sieve inequality of Montgomery and Vaughan [1979] for prime discriminants (see Lemma 5.1 below). Since the Legendre symbol modulo  $p$  is even if  $p \equiv 1 \pmod 4$ , and is odd if  $p \equiv 3 \pmod 4$ , we shall consider the cases of primes congruent to 1 and 3 modulo 4 separately. For  $a \in \{1, 3\}$ , let  $\Psi_{x,a}^{\text{prime}}(\tau)$  be the proportion of primes  $p \leq x$  such that  $p \equiv a \pmod 4$  and  $m((\frac{\cdot}{p})) > \tau$ .

**Theorem 1.6.** *Let  $\eta$  and  $B_0$  be the constants in Theorem 1.1. There exists positive constants  $C_1, C_2$  such that:*

1. *Uniformly in the range  $2 \leq \tau \leq \log_2 x + \log_5 x - \log_4 x - C_1$  we have*

$$\Psi_{x,3}^{\text{prime}}(\tau) \leq \exp\left(-\frac{e^{\tau-\eta-\log 2-2}}{\tau} \left(1 + O\left(\frac{\log \tau}{\tau}\right)\right)\right).$$

2. *Uniformly in the range  $2 \leq \tau \leq (\log_2 x)/2 - 2 \log_3 x$  we have*

$$\Psi_{x,3}^{\text{prime}}(\tau) \geq \exp\left(-\frac{e^{\tau-\eta-B_0}}{\tau} (1 + o(1))\right). \tag{1-6}$$

3. *Uniformly in the range  $(\log_2 x)/2 - 2 \log_3 x \leq \tau \leq \log_2 x - \log_3 x - C_1$  we have*

$$\Psi_{x,3}^{\text{prime}}(\tau) \geq \exp(-C_2 \tau e^\tau).$$

**Theorem 1.7.** *Let  $B_0$  be the constant in Theorem 1.1. There exists positive constants  $C_1, C_2$  and  $C_3$  such that:*

1. *Uniformly in the range  $2 \leq \tau \leq (\log_2 x - \log_3 x - C_1)/\sqrt{3}$  we have*

$$\exp(-C_3 \tau e^{\sqrt{3}\tau}) \leq \Psi_{x,1}^{\text{prime}}(\tau) \ll \exp\left(-\frac{e^{\sqrt{3}\tau}}{\tau C_2}\right).$$

2. *Moreover, uniformly in the smaller range  $2 \leq \tau \leq ((\log_2 x)/2 - 2 \log_3 x)/\sqrt{3}$  we have the improved lower bound*

$$\Psi_{x,1}^{\text{prime}}(\tau) \geq \exp\left(-\frac{e^{\sqrt{3}\tau - B_0}}{\sqrt{3}\tau}(1 + o(1))\right). \quad (1-7)$$

**Remark 1.8.** An interesting consequence of Theorems 1.6 and 1.7 is that almost all primes  $p \leq x$  for which  $M\left(\left(\frac{\cdot}{p}\right)\right)$  is large are congruent to 3 modulo 4. Moreover, we note that conditionally on the GRH, the lower bound (1-6) (respectively (1-7)) holds in the extended range  $2 \leq \tau \leq \log_2 x - 2 \log_3 x - C_4$  (respectively  $2 \leq \tau \leq (\log_2 x - 2 \log_3 x - C_4)/\sqrt{3}$ ), for some positive constant  $C_4$ . See Remark 6.1 below for a justification of this fact.

**Remark 1.9.** Theorem 1.4 of [Bober et al. 2018] establishes the existence of a limiting distribution for  $m(\chi)$ , as  $\chi$  varies over nonprincipal characters modulo a large prime  $q$ , when  $q \rightarrow \infty$ . In [Hussain and Lamzouri 2023], we established an analogous result for  $M\left(\left(\frac{\cdot}{p}\right)\right)$  as  $p$  varies over the primes in a large dyadic interval  $[Q, 2Q]$  and  $Q \rightarrow \infty$ .

**1C. The distribution of  $L\left(1, \left(\frac{\cdot}{p}\right)\right)$ .** In order to prove the lower bounds of Theorems 1.6 and 1.7 we need estimates on the distribution of  $L\left(1, \left(\frac{\cdot}{p}\right)\right)$ , as  $p$  varies over the primes. It turns out that this case is harder than the case of the bigger family of primitive quadratic characters attached to fundamental discriminants  $d$ , which was investigated by Granville and Soundararajan [2003] (see the precise statement of their result in Theorem 2.2 below). By the law of quadratic reciprocity, this is due to the fact that current bounds on character sums over primes are much weaker than analogous bounds for character sums over the integers.<sup>1</sup> In particular, such bounds are heavily affected by the possible existence of a Landau–Siegel exceptional discriminant.

Joshi [1970] extended Littlewood’s Omega result [1928], by establishing the existence of infinitely many primes  $p$  such that

$$L\left(1, \left(\frac{\cdot}{p}\right)\right) \geq (e^\gamma + o(1)) \log \log p.$$

<sup>1</sup>Bounds for character sums over primes are ultimately connected to the distribution of primes in arithmetic progressions. The weakness of such bounds explains for example why the strongest version of the prime number theorem for arithmetic progressions, namely the Siegel–Walfisz theorem, only holds for very small moduli.

We improve on this result by obtaining estimates for the proportion of primes  $p \leq x$  such that  $L(1, (\frac{\cdot}{p})) > e^\gamma \tau$  uniformly for  $\tau$  in the range  $2 \leq \tau \leq (1 - o(1)) \log \log p$ , which is believed to be best possible up to the factor  $o(\log \log p)$  (see for example the conjectures of Montgomery and Vaughan [1999] and Granville and Soundararajan [2003]). For  $\tau > 0$  and  $a \in \{1, 3\}$  we define

$$F_{x,a}(\tau) := \frac{2}{\pi(x)} \left| \left\{ p \leq x : p \equiv a \pmod{4}, L\left(1, \left(\frac{\cdot}{p}\right)\right) > e^\gamma \tau \right\} \right|.$$

**Theorem 1.10.** *Let  $a \in \{1, 3\}$  and  $x$  be large. In the range  $2 \leq \tau \leq (\log_2 x)/2 - 2 \log_3 x$  we have*

$$F_{x,a}(\tau) = \exp\left(-\frac{e^{\tau-B_0}}{\tau} (1 + o(1))\right). \tag{1-8}$$

Moreover, there exists positive constants  $C_1, C_2 > 0$  such that in the range  $(\log_2 x)/2 - 2 \log_3 x \leq \tau \leq \log_2 x - \log_3 x - C_2$  we have

$$\exp(-C_1 \tau e^\tau) \leq F_{x,a}(\tau) \leq \exp\left(-\frac{e^{\tau+\log 2-2}}{\tau} \left(1 + O\left(\frac{\log \tau}{\tau}\right)\right)\right). \tag{1-9}$$

Furthermore, the same estimates hold for the proportion of primes  $p \leq x$  such that  $p \equiv a \pmod{4}$  and  $L(1, (\frac{\cdot}{p})\chi_{-3}) > (2e^\gamma/3)\tau$ , where  $\chi_{-3}$  is the nonprincipal character modulo 3.

**Remark 1.11.** One can compare our results with those of [Granville and Soundararajan 2003] for the distribution of  $L(1, \chi_d)$  over fundamental discriminants  $|d| \leq x$  (see Theorem 2.2 below). In this case, the same estimate (1-8) holds for the proportion of fundamental discriminants  $|d| \leq x$  such that  $L(1, \chi_d) > e^\gamma \tau$ , uniformly for  $\tau$  in the larger range  $2 \leq \tau \leq \log_2 x + (1 - o(1)) \log_4 x$ . We should also note that conditionally on GRH, the estimate (1-8) holds in the extended range  $2 \leq \tau \leq \log_2 x - 2 \log_3 x - C_2$  (for some positive constant  $C_2$ ) by a result of Holmin, Jones, Kurlberg, McLeman and Petersen [Holmin et al. 2019] (see Remark 6.1 below). Finally, we note that in the case  $a = 3$ , our proof yields a better constant in the upper bound of (1-9) than if we just apply Theorem 1.6 directly (using the inequality (2-8) below).

To prove the precise estimate (1-8) in the smaller range  $2 \leq \tau \leq (1/2 - o(1)) \log_2 x$ , we use our previous work [Lanzouri 2017], where we established asymptotic formulas for the complex moments of  $L(1, (\frac{\cdot}{p}))$  involving a secondary term which is coming from a possible Landau–Siegel exceptional discriminant. Although we could not rule out that this term might be as large as the main term, we were able to show that it does not affect the leading term in the tail of the distribution of  $L(1, (\frac{\cdot}{p}))$ . However, we note that the error term  $o(1)$  inside the estimate (1-8) is not effective, due to the use of Siegel’s theorem. The upper bound of (1-9) will be a consequence of Theorem 2.7 below, which is the key ingredient in the proofs of the upper bounds of Theorems 1.6 and 1.7. Finally, to obtain the lower bound of (1-9), we combine Theorem 2.7 with a strong form of Linnik’s theorem established by Bombieri [1987] using his zero density estimates for Dirichlet  $L$ -functions.

**1D. An application to a question of Montgomery.** As an application of our results we consider the problem of the positivity of sums of the Legendre symbol. Let  $p \geq 3$  be a prime and

$$S_p(t) = \sum_{n \leq t} \left( \frac{n}{p} \right).$$

The question of determining when  $S_p(t)$  is positive was considered by several mathematicians including Fekete, Chowla, Montgomery and others. Since  $S_p(t)$  is periodic of period  $p$ , one can renormalize the variable  $t$  and define for  $\alpha \geq 0$  the function

$$f_p(\alpha) := \sum_{0 \leq n \leq \alpha p} \left( \frac{n}{p} \right),$$

which is periodic of period 1. One can extend  $f_p$  to all  $\alpha \in \mathbb{R}$  by periodicity. Montgomery [1976] studied the following natural question: How frequently is  $f_p(\alpha)$  positive for a prime  $p \equiv 3 \pmod{4}$ ? More precisely, he investigated the quantity

$$\lambda(p) := \mu(\{\alpha \in [0, 1) : f_p(\alpha) > 0\}),$$

for such primes<sup>2</sup>, where  $\mu(\mathcal{C})$  denotes the Lebesgue measure of a measurable set  $\mathcal{C}$ . He proved in [Montgomery 1976] that for all primes  $p \equiv 3 \pmod{4}$  we have  $\lambda(p) > 1/50$ . He also established the existence of infinitely many such primes such that  $\lambda(p) < 1/3 + \varepsilon$  for any fixed  $\varepsilon > 0$ . In her Master's thesis, Mehkari [2005] slightly improved the constant  $1/50$  in the first result of Montgomery. She also ran extensive numerical computations which suggest that the value  $1/3$  for his second result is optimal. Furthermore, she proved conditionally on GRH that  $\lambda(p) \leq 0.764$  for a positive proportion of the primes  $p \equiv 3 \pmod{4}$ , and that  $\lambda(p) \geq 0.285$  for a positive proportion of the primes  $p \equiv 3 \pmod{4}$ . Montgomery [1976] also wrote "the ideas found in our proof can also be used to show that there are infinitely many primes  $p \equiv 3 \pmod{4}$  such that  $\lambda(p) > 1 - \varepsilon$ ." Using a different method, based on the proof of Theorem 1.6, we improve on these results by showing that for any  $\varepsilon > 0$ , both inequalities  $\lambda(p) > 1 - \varepsilon$  and  $\lambda(p) < 1/3 + \varepsilon$  hold for a positive proportion of the primes  $p \equiv 3 \pmod{4}$ . We also quantify these proportions in terms of  $\varepsilon$  and consider the question of uniformity by letting  $\varepsilon \rightarrow 0$  slowly as a function of  $x$ , if we vary over the primes  $p \leq x$  such that  $p \equiv 3 \pmod{4}$ .

**Theorem 1.12.** *Let  $\nu > 0$  be a small fixed constant. Let  $x$  be large and*

$$1 < T \leq \exp\left((1 - \nu) \log_2 x \log_3 x / (\log_4 x)\right)$$

*be a real number. The number of primes  $p \leq x$  with  $p \equiv 3 \pmod{4}$  and such that  $\lambda(p) > 1 - 1/T$  is*

$$\gg \pi(x) \exp\left(-\exp\left(\frac{\log T \log_3 T}{\log_2 T} (1 + o(1))\right)\right). \quad (1-10)$$

<sup>2</sup>Note that  $f_p$  is even if  $p \equiv 3 \pmod{4}$  and is odd if  $p \equiv 1 \pmod{4}$ . The analogous question in the latter case becomes: how often is  $f_p(\alpha) > 0$  for  $0 < \alpha < 1/2$ ? We shall only consider the former case in this paper since our methods can be extended to handle the case of primes  $p \equiv 1 \pmod{4}$ .

In particular, this quantity is

$$\gg_v \pi(x) \exp(-T^v).$$

**Theorem 1.13.** *Let  $x$  be large and  $T > 1$ . There exists positive constants  $c_1, c_2, c_3$  such that the number of primes  $p \leq x$  with  $p \equiv 3 \pmod{4}$  and such that  $\lambda(p) < 1/3 + 1/T$  is at least*

$$\pi(x) \exp(-c_1 T^2 (\log T)^3), \tag{1-11}$$

if  $1 < T \leq c_2 (\log_2 x)^{1/2} / (\log_3 x)^2$ , and is at least

$$\pi(x) \exp(-c_1 T^2 (\log T)^5), \tag{1-12}$$

if  $c_2 (\log_2 x)^{1/2} / (\log_3 x)^2 \leq T \leq c_3 (\log x)^{1/2} / (\log_2 x)^{5/2}$ .

**1E. Further applications.** We should note that our method could be adapted to investigate the distribution of  $M(\chi)$  over the family of primitive cubic Dirichlet characters with conductor up to  $x$ . We are planning to pursue this direction in a future paper. Moreover, we mention that the ideas of the proof of Theorem 1.6 are used in [Hussain and Lamzouri 2023], concerning the limiting distribution of character paths attached to the family of Legendre symbols modulo primes. Hussain [2022b] previously established a similar result for character paths attached to nonprincipal characters modulo a large prime  $q$ . Furthermore, in her PhD thesis [Hussain 2022a], she proved conditionally on GRH that character paths attached to the family of Legendre symbols converge in law (in the space of continuous functions) to a random Fourier series constructed using Rademacher random multiplicative functions. Our forthcoming work will establish this result unconditionally.

## 2. Outline and key ingredients of the proofs of Theorems 1.1, 1.2 and 1.6

**2A. The overall strategy of the proof of Theorem 1.1.** We first start by describing the strategy and key ideas of the proof of Theorem 1.1, since the method for proving Theorems 1.2, 1.6 and 1.7 is similar. In particular, it will be useful to compare the argument with [Bober et al. 2018]. The character sum  $\sum_{n \leq t} \chi_d(n)$  has a simple Fourier expansion first obtained by Pólya in the following quantitative form [Montgomery and Vaughan 2007, equation (9.19), p. 311]

$$\sum_{n \leq t} \chi_d(n) = \frac{\mathcal{G}(\chi_d)}{2\pi i} \sum_{1 \leq |n| \leq Z} \frac{\chi_d(n)(1 - e(-nt/|d|))}{n} + O\left(1 + \frac{|d| \log |d|}{Z}\right), \tag{2-1}$$

where  $\mathcal{G}(\chi_d)$  is the Gauss sum attached to  $\chi_d$ . Let  $\delta := 1/100$ , and define  $\mathcal{F}^*(x)$  to be the set of fundamental discriminants  $x^{1-\delta} \leq |d| \leq x$ . We note that the proportion of those discriminants  $d$  with  $|d| \leq x^{1-\delta}$  is  $\ll x^{-\delta}$ , which is much smaller than the distribution function  $\Psi_x^-(\tau)$  in the range  $\tau \leq \log \log x$ . Hence, we will only focus on the fundamental discriminants  $d \in \mathcal{F}^*(x)$ . For these discriminants we have

$$m(\chi_d) = \frac{e^{-\gamma}}{2} \max_{\alpha \in [0,1)} \left| \sum_{1 \leq |n| \leq Z} \frac{\chi_d(n)(1 - e(\alpha n))}{n} \right| + O(x^{-\delta}), \tag{2-2}$$

where  $Z = x^{21/40}$ . We first describe a heuristic argument that will help us isolate the key ingredient in the proof of Theorem 1.1. This will also explain why we should expect the tail of the distribution of  $m(\chi_d)$  to behave like  $\exp(-e^\tau/\tau)$ . A standard idea, which goes back to the work of Montgomery and Vaughan [1977] on exponential sums with multiplicative coefficients, is to split the sum on the right-hand side of (2-2) into two parts

$$\sum_{1 \leq |n| \leq Z} \frac{\chi_d(n)(1 - e(\alpha n))}{n} = \sum_{\substack{1 \leq |n| \leq Z \\ P^+(n) \leq y}} \frac{\chi_d(n)(1 - e(\alpha n))}{n} + \sum_{\substack{1 \leq |n| \leq Z \\ P^+(n) > y}} \frac{\chi_d(n)(1 - e(\alpha n))}{n}, \tag{2-3}$$

and show that uniformly over  $\alpha$ , the bulk of the distribution comes from the first part, over  $y$ -friable integers, with a suitable choice of the parameter  $y$ . Here  $P^+(n)$  is the largest prime factor of  $n$ , and an integer  $n$  is called  $y$ -friable (or  $y$ -smooth) if  $P^+(n) \leq y$ . To understand what choice of the parameter  $y$  we should make we observe that

$$\max_{\alpha \in [0,1]} \left| \sum_{\substack{1 \leq |n| \leq Z \\ P^+(n) \leq y}} \frac{\chi_d(n)(1 - e(\alpha n))}{n} \right| \ll \sum_{\substack{n \geq 1 \\ P^+(n) \leq y}} \frac{1}{n} \ll \log y \tag{2-4}$$

by Mertens' theorem. Hence, if the main part of the contribution to  $m(\chi_d) > \tau$  is coming from  $y$ -friable integers, we should aim for a choice of  $y$  such that<sup>3</sup>  $y \approx e^\tau$ . Heuristically, for small  $y$  ( $y \leq \log x$  say) we can prescribe the values of  $\chi_d(p)$  for  $p \leq y$  with probability<sup>4</sup>  $2^{-\pi(y)} = \exp(-(\log 2 + o(1))y/\log y)$  by the prime number theorem. Therefore, if  $y \asymp e^\tau$  this probability looks like  $\exp(-c_1 e^\tau/\tau)$  for some positive constant  $c_1$ , which agrees with the statement of Theorem 1.1. Thus, in order for this heuristic to work, we need to efficiently control the second part in (2-3) uniformly over  $\alpha \in [0, 1)$ . We achieve this in the following theorem, which is the key ingredient in the proofs of Theorems 1.1 and 1.2.

**Theorem 2.1.** *Let  $h(n)$  be a completely multiplicative function such that  $|h(n)| \leq 1$  for all  $n$ . Let  $x$  be large and put  $Z = x^{21/40}$ . There exists a constant  $c > 0$  such that for all real numbers  $2 \leq y \leq c \log x \log_4 x / (\log_3 x)$  and  $1/\log y \leq A \leq 4$ , the number of fundamental discriminants  $|d| \leq x$  such that*

$$\max_{\alpha \in [0,1]} \left| \sum_{\substack{1 \leq n \leq Z \\ P^+(n) > y}} \frac{\chi_d(n)h(n)e(n\alpha)}{n} \right| > e^y A$$

is

$$\ll x \exp\left(-\frac{A^2 y}{2 \log y} \left(1 + O\left(\frac{\log_2 y}{\log y} + \frac{\log_4 x}{A \log_2 x \log_3 x}\right)\right)\right).$$

<sup>3</sup>This is not completely correct, since the above argument shows that we rather have  $y \approx e^{c\tau}$  for some constant  $c > 0$ . However, it turns out that the optimal choice of  $c$  is  $c = 1$ .

<sup>4</sup>This is correct in the range  $y \leq \log \log x$ , as proved in Lemma 7.1 below. It also follows from GRH in the larger range  $y \leq c \log x \log \log x$  for some small constant  $c > 0$ ; see Theorem 13.5 of [Montgomery 1971].

Bober, Goldmakher, Granville and Koukoulopoulos [Bober et al. 2018] proved a similar result in the case of nonprincipal characters modulo a large prime  $q$ . However our proof differs from theirs as we shall now explain. Let  $z = q^{21/40}$ . In order to prove the analogue of Theorem 2.1 for the family of nonprincipal characters modulo  $q$ , the authors of [Bober et al. 2018] established nontrivial upper bounds for the  $2k$ -th moment of

$$\max_{\alpha \in [0,1)} \left| \sum_{\substack{1 \leq n \leq z \\ P^+(n) > y}} \frac{\chi(n)e(n\alpha)}{n} \right| \tag{2-5}$$

for  $k$  roughly up to  $\log q$ . By decoupling the  $\alpha$  from the character  $\chi$  and expanding the  $2k$ -th moment of the inner sum in (2-5), one needs to control terms which have size as large as  $z^k > q^{k/2}$  with  $k$  up to  $\log q$ . This was possible in the case of the family of nonprincipal characters modulo  $q$  thanks to the orthogonality relations of characters, which imply that the off-diagonal terms in these moments are given by  $m \equiv n \pmod q$  with  $m \neq n$  and  $m, n \leq z^k$ . The authors of [Bober et al. 2018] proceed to bound these off-diagonal terms using intricate estimates involving the  $k$ -th divisor function.

This argument is no longer valid for families of quadratic characters, since in this case only “quasi-orthogonality relations” are known (see, for example, Lemma 4.1 of [Granville and Soundararajan 2003]), which allow one to control terms up to size  $x^c$  for some  $c < 1$ , if we run over fundamental discriminants  $|d| \leq x$ . To overcome this problem, we made the key observation that when  $N < n < 2N$  and  $N$  is very large, one only needs to compute a small moment over the corresponding sum over the interval  $[N, 2N]$ , in order to show that this sum is small for most characters. More specifically, our approach consists of first splitting the sum in (2-5) (in the case of quadratic characters associated to fundamental discriminants  $|d| \leq x$ ) into two parts: the first over  $n \leq Y$  and the second over  $Y < n \leq Z$ , where  $Y = (\log x)^{W(x)}$  is relatively “small” (our method allows one to choose  $W(x) = \log_3 x / (\log_4 x)$ ). We first bound the  $2k$ -th moments of the first sum over  $n \leq Y$  and show that only the diagonal terms contribute<sup>5</sup> to these moments if  $k \leq (\log x) / (3 \log Y)$  say. We then proceed to show that the second part over the large  $n$ 's is very small for most characters. To this end, we split it in dyadic intervals  $N \leq n \leq 2N$  and then bound the  $2\ell_N$  moment of

$$\max_{\alpha \in [0,1)} \left| \sum_{N < n < 2N} \frac{\chi(n)e(n\alpha)}{n} \right|$$

where  $\ell_N$  is chosen such that  $N^{\ell_N}$  is roughly of size  $x$ . This allows us to show that each of these parts is small for most characters using *tailored* moments according to the size of  $N$ . In the range  $Y < N < x^\epsilon$  we use (4-1) below, which is an easy version of the quadratic large sieve first due to Elliott, while in the remaining range  $x^\epsilon < N < Z$  we use the quadratic large sieve of Heath-Brown (see (4-2) below).

---

<sup>5</sup>We prefer to do this using an easy version of the quadratic large sieve (see (4-1) below), rather than employing the quasiorthogonality relations, in order to get a clean argument.

Deducing the upper bound of Theorem 1.1 from Theorem 2.1. Let  $Z = x^{21/40}$  and  $\delta = 1/100$ . By (2-2) we have

$$m(\chi_d) \leq \frac{e^{-\gamma}}{2} \max_{\alpha \in [0,1]} \left| \sum_{\substack{1 \leq |n| \leq Z \\ P^+(n) \leq y}} \frac{\chi_d(n)(1 - e(\alpha n))}{n} \right| + 2e^{-\gamma} \max_{\alpha \in [0,1]} \left| \sum_{\substack{1 \leq |n| \leq Z \\ P^+(n) > y}} \frac{\chi_d(n)e(\alpha n)}{n} \right| + O(x^{-\delta}), \tag{2-6}$$

for any fundamental discriminant  $-x \leq d \leq -x^{1-\delta}$ . Since we do not have control over the first part over  $y$ -friable integers, we shall bound it using Corollary 3.5 of [Bober et al. 2018] which gives

$$\max_{\alpha \in [0,1]} \left| \sum_{\substack{1 \leq |n| \leq Z \\ P^+(n) \leq y}} \frac{\chi_d(n)(1 - e(\alpha n))}{n} \right| \leq 2e^\gamma \log y + 2 \log 2 + O\left(\frac{\log \log y}{\log y}\right). \tag{2-7}$$

We now set  $y = e^{\tau - e^{-\gamma} \log 2 - 2B}$ , where  $B > 0$  is a parameter to be chosen. Combining this last estimate with (2-6) and (2-7) we deduce that

$$m(\chi_d) \leq \tau - 2B + 2e^{-\gamma} \max_{\alpha \in [0,1]} \left| \sum_{\substack{1 \leq |n| \leq Z \\ P^+(n) > y}} \frac{\chi_d(n)e(\alpha n)}{n} \right| + O\left(\frac{\log \tau}{\tau}\right).$$

Thus, the proportion of fundamental discriminants  $-x \leq d \leq -x^{1-\delta}$  such that  $m(\chi_d) > \tau$  is bounded by the proportion of fundamental discriminants  $|d| \leq x$  such that

$$\max_{\alpha \in [0,1]} \left| \sum_{\substack{1 \leq |n| \leq Z \\ P^+(n) > y}} \frac{\chi_d(n)e(\alpha n)}{n} \right| > e^\gamma \left( B - C_0 \frac{\log \tau}{\tau} \right),$$

for some suitably large constant  $C_0 > 0$ . Choosing  $B = 1$  and appealing to Theorem 2.1 completes the proof. □

We now turn our attention to the lower bound of Theorem 1.1 which is easier. For  $d < 0$  one has the identity (see Theorem 9.21 of [Montgomery and Vaughan 2007])

$$\sum_{n \leq |d|/2} \chi_d(n) = (2 - \chi(2)) \frac{\mathcal{G}(\chi_d)}{i\pi} L(1, \chi_d),$$

which implies

$$m(\chi_d) \geq e^{-\gamma} L(1, \chi_d). \tag{2-8}$$

Therefore, one can immediately deduce a corresponding lower bound for the distribution function  $\Psi_x^-(\tau)$  from the following result, which follows from a straightforward adaptation of the work of Granville and Soundararajan [2003] on the distribution of  $L(1, \chi_d)$ .

**Theorem 2.2** (Granville–Soundararajan). *Let  $B_0$  be the constant in Theorem 1.1,  $\psi$  be a character modulo some  $b \in \{1, 3\}$ , and  $1 \leq \tau \leq \log_2 x + \log_4 x - 20 - M$  for some  $M \geq 0$ . Then we have*

$$\frac{1}{|\mathcal{F}^\pm(x)|} \left| \left\{ d \in \mathcal{F}^\pm(x) : L(1, \chi_d \psi) > \frac{\phi(b)}{b} e^\gamma \tau \right\} \right| = \exp\left(-\frac{e^{\tau-B_0}}{\tau} \left(1 + O\left(\frac{1}{\tau} + e^{-e^M}\right)\right)\right).$$

One can do a little better by combining our Theorem 2.1 with the following result, which we extract from the proof of Theorem 1.2 of [Bober et al. 2018].

**Theorem 2.3** (Bober, Goldmakher, Granville and Koukoulopoulos). *Let  $C$  be a positive constant. Let  $q$  be a large positive integer, and  $y \leq (\log q)^2$  be a large real number. Let  $\chi$  be a primitive odd character modulo  $q$  such that*

$$|L(1, \chi)| > e^\gamma \log y - C \quad \text{and} \quad \max_{\alpha \in [0,1)} \left| \sum_{\substack{n \geq 1 \\ P^+(n) > y}} \frac{\chi(n)e(n\alpha)}{n} \right| \leq 1.$$

Then we have

$$m(\chi) > e^{-\gamma} (|L(1, \chi)| + \log 2) + O\left(\frac{(\log \log y)^2}{\sqrt{\log y}}\right).$$

**Remark 2.4.** We note that  $q$  is assumed to be prime in the statement of Theorem 1.2 of [Bober et al. 2018]. However, this is only used to show that many nonprincipal characters  $\chi \pmod q$  satisfying the assumptions of Theorem 2.3 exist. Indeed, the proof of Theorem 2.3 (see the end of Section 4 of [Bober et al. 2018]) only uses estimates on exponential sums over  $y$ -friable integers [Bober et al. 2018, Lemmas 3.2 and 3.4] together with ideas of [Bober 2014] on averages of character sums to arbitrary moduli.

*Deducing the lower bound of Theorem 1.1 from Theorems 2.1, 2.2 and 2.3.* Let  $Z = x^{21/40}$  and  $\delta = 1/100$ . By partial summation and the Pólya–Vinogradov inequality, it follows that

$$\max_{\alpha \in [0,1)} \left| \sum_{n > Z} \frac{\chi_d(n)e(n\alpha)}{n} \right| \ll \frac{\sqrt{|d|} \log |d|}{Z} \ll x^{-\delta}, \tag{2-9}$$

for any fundamental discriminant  $|d| \leq x$ . Furthermore, by Lemma 3.2 of [Bober et al. 2018] we have

$$\max_{\alpha \in [0,1)} \left| \sum_{\substack{n > Z \\ P^+(n) \leq y}} \frac{\chi_d(n)e(n\alpha)}{n} \right| \leq \sum_{\substack{n > Z \\ P^+(n) \leq y}} \frac{1}{n} \ll e^{-\sqrt{\log y}}, \tag{2-10}$$

for any real number  $2 \leq y \leq (\log x)^2$ . Combining these estimates implies that

$$\max_{\alpha \in [0,1)} \left| \sum_{\substack{n > Z \\ P^+(n) > y}} \frac{\chi_d(n)e(n\alpha)}{n} \right| \ll e^{-\sqrt{\log y}}, \tag{2-11}$$

for any fundamental discriminant  $|d| \leq x$  and any real number  $2 \leq y \leq (\log x)^2$ .

Let  $C_1, C_2 > 0$  be suitably large constants and put  $y = e^{\tau+C_1}$ . If  $C_1$  is large enough, then combining Theorems 2.1 and 2.2 we deduce that the number of fundamental discriminants  $-x < d < 0$  such that

$$L(1, \chi_d) > e^\gamma \tau - \log 2 + C_1 \frac{(\log \tau)^2}{\sqrt{\tau}} \quad \text{and} \quad \max_{\alpha \in [0,1)} \left| \sum_{\substack{1 \leq n \leq Z \\ P^+(n) > y}} \frac{\chi_d(n)e(n\alpha)}{n} \right| \leq 1 - \frac{C_2}{\tau}$$

is

$$\geq \exp\left(-\frac{e^{\tau-\eta-B_0}}{\tau} \left(1 + O\left(\frac{(\log \tau)^2}{\sqrt{\tau}}\right)\right)\right).$$

By (2-11) we deduce that for any such discriminant  $d$  we have

$$\max_{\alpha \in [0,1)} \left| \sum_{\substack{n \geq 1 \\ P^+(n) > y}} \frac{\chi_d(n)e(n\alpha)}{n} \right| \leq 1,$$

and hence it follows from Theorem 2.3 that  $m(\chi_d) > \tau$  if  $C_1$  is suitably large. This completes the proof.  $\square$

**2B. The case of positive discriminants: Proof of Theorem 1.2.** Recall that  $\chi_d$  is an even character if  $d$  is a positive fundamental discriminant. In this case, we shall appeal to the following structure result for even characters with large sums, which we extract from the proof of Theorem 2.3 of [Bober et al. 2018].

**Theorem 2.5** (Bober, Goldmakher, Granville and Koukoulopoulos). *Let  $C$  be a positive constant. Let  $q$  be a large positive integer and  $y \leq (\log q)^2$  be a large real number. Let  $\psi \pmod q$  be a primitive even character such that*

$$\max_{\alpha \in [0,1)} \left| \sum_{\substack{n \geq 1 \\ P^+(n) \leq y}} \frac{\psi(n)e(n\alpha)}{n} \right| > \frac{e^\gamma}{\sqrt{3}} \log y - C \log \log y, \quad \text{and} \quad \max_{\alpha \in [0,1)} \left| \sum_{\substack{n \geq 1 \\ P^+(n) > y}} \frac{\psi(n)e(n\alpha)}{n} \right| \leq 1.$$

Then we have

$$m(\psi) = \frac{e^{-\gamma} \sqrt{3}}{2} |L(1, \psi \chi_{-3})| + O(\log \log y).$$

**Remark 2.6.** Here again we note that  $q$  is assumed to be prime in Theorem 2.3 of [Bober et al. 2018], but this is only used to show that many nonprincipal characters  $\psi \pmod q$  satisfying the assumptions of Theorem 2.5 exist. Indeed, all of the ingredients of the proof of Theorem 2.5 hold for an arbitrary primitive even character  $\psi$  (see Section 8 of [Bober et al. 2018]), and are derived using the “pretentious” theory of character sums developed by Granville and Soundararajan [2007], along with estimates of Montgomery and Vaughan [1977] on exponential sums with multiplicative coefficients.

*Deducing Theorem 1.2 from Theorems 2.1, 2.2 and 2.5.* The lower bound follows immediately from Theorem 2.2 together with the standard inequality (see for example the beginning of Section 4 of [Bober et al. 2018])

$$M(\psi) \geq \frac{\sqrt{3q}}{2\pi} |L(1, \psi \chi_{-3})|, \tag{2-12}$$

which is valid for any primitive even character  $\psi$  modulo  $q$ .

We now prove the upper bound. First, it follows from (2-1) that for all fundamental discriminants  $0 < d < x$  we have

$$m(\chi_d) = e^{-\gamma} \max_{\alpha \in [0,1)} \left| \sum_{1 \leq n \leq Z} \frac{\chi_d(n) \sin(2\pi n\alpha)}{n} \right| + O(1), \tag{2-13}$$

since  $\chi_d$  is even. Let  $C_1, C_2 > 0$  be suitably large constants,  $y = e^{\sqrt{3}\tau + C_1}$ , and define  $\mathcal{D}^+(x, y)$  to be the set of fundamental discriminants  $0 < d \leq x$  such that

$$m(\chi_d) > \tau \quad \text{and} \quad \max_{\alpha \in [0,1)} \left| \sum_{\substack{1 \leq n \leq Z \\ P^+(n) > y}} \frac{\chi_d(n)e(n\alpha)}{n} \right| \leq 1 - \frac{C_2}{\tau}.$$

By combining the lower bound of Theorem 1.2 with Theorem 2.1 we deduce that if  $C_1$  is suitably large then

$$\frac{|\mathcal{D}^+(x, y)|}{|\mathcal{F}^+(x)|} = \Psi_x^+(\tau)(1 + o(1)). \tag{2-14}$$

Now, let  $d \in \mathcal{D}^+(x, y)$ . By (2-10) and (2-13) together with our assumption on  $d$  we have

$$m(\chi_d) = e^{-\gamma} \max_{\alpha \in [0,1)} \left| \sum_{\substack{1 \leq n \leq Z \\ P^+(n) \leq y}} \frac{\chi_d(n) \sin(2\pi n\alpha)}{n} \right| + O(1) = e^{-\gamma} \max_{\alpha \in [0,1)} \left| \sum_{\substack{n \geq 1 \\ P^+(n) \leq y}} \frac{\chi_d(n) \sin(2\pi n\alpha)}{n} \right| + O(1).$$

Since  $m(\chi_d) > \tau$  we deduce that

$$\max_{\alpha \in [0,1)} \left| \sum_{\substack{n \geq 1 \\ P^+(n) \leq y}} \frac{\chi_d(n)e(n\alpha)}{n} \right| \geq e^\gamma m(\chi_d) + O(1) > \frac{e^\gamma}{\sqrt{3}} \log y - C_3,$$

for some positive constant  $C_3$ . Moreover, by (2-11) and our assumption on  $d$  we get

$$\max_{\alpha \in [0,1)} \left| \sum_{\substack{n \geq 1 \\ P^+(n) > y}} \frac{\chi_d(n)e(n\alpha)}{n} \right| \leq 1 - \frac{C_2}{\tau} + O(e^{-\sqrt{\tau}}) \leq 1,$$

if  $C_2$  is suitably large. Thus  $\chi_d$  satisfies the conditions of Theorem 2.5, which implies that

$$m(\chi_d) = \frac{e^{-\gamma} \sqrt{3}}{2} |L(1, \chi_d \chi_{-3})| + O(\log \tau).$$

Therefore, for all  $d \in \mathcal{D}^+(x, y)$  we have

$$|L(1, \chi_d \chi_{-3})| > \frac{2e^\gamma}{\sqrt{3}} \tau - C_4 \log \tau,$$

for some constant  $C_4 > 0$ . Hence, it follows from Theorem 2.2 that

$$|\mathcal{D}^+(x, y)| \ll |\mathcal{F}^+(x)| \exp\left(-\frac{e^{\sqrt{3}\tau}}{\tau C_5}\right),$$

for some constant  $C_5 > 0$ . Combining this estimate with (2-14) completes the proof. □

**2C. The family of Legendre symbols: Outline of the proofs of Theorems 1.6 and 1.7.** In order to prove the upper and lower bounds of Theorems 1.6 and 1.7, we shall follow the same strategy of the proofs of Theorems 1.1 and 1.2. To this end we establish the following key result, which is the analogue of Theorem 2.1 for prime discriminants. To shorten our notation, we let  $\psi_p$  denote the Legendre symbol modulo  $p$  throughout the remaining part of the paper.

**Theorem 2.7.** *Let  $h(n)$  be a completely multiplicative function such that  $|h(n)| \leq 1$  for all  $n$ . Let  $x$  be large and put  $Z = x^{21/40}$ . There exists a constant  $c > 0$  such that for all real numbers  $2 \leq y \leq c \log x \log_4 x / (\log_3 x)$  and  $1/\log y \leq A \leq 4$ , the number of primes  $p \leq x$  such that*

$$\max_{\alpha \in [0,1)} \left| \sum_{\substack{1 \leq n \leq Z \\ P^+(n) > y}} \frac{\psi_p(n)h(n)e(n\alpha)}{n} \right| > e^\gamma A$$

is

$$\ll \pi(x) \exp\left(-\frac{A^2 y}{2 \log y} \left(1 + O\left(\frac{\log_2 y}{\log y} + \frac{\log_4 x}{A \log_2 x \log_3 x}\right)\right)\right).$$

We end this section by deducing Theorems 1.6 and 1.7 from this result along with Theorem 1.10 on the distribution of large values of  $L(1, \psi_p)$ .

*Proof of Theorem 1.6 assuming Theorems 1.10 and 2.7.* First, Part 1 of Theorem 1.6 can be derived along the same lines of the proof of the upper bound of Theorem 1.1 by replacing Theorem 2.1 by Theorem 2.7.

Now, the proof of Part 2 follows along the same lines of the proof of the lower bound of Theorem 1.1 by replacing Theorems 2.1 and 2.2 by Theorem 2.7 and the estimate (1-8) respectively. Finally, the proof of Part 3 follows from (1-9) together with the lower bound (2-8). □

*Proof of Theorem 1.7 assuming Theorems 1.10 and 2.7.* Part 2 and the lower bound of Part 1 of Theorem 1.7 follow immediately from Theorem 1.10 together with the lower bound (2-12). Finally, the upper bound of Part 1 can be derived along the same lines of the proof of the upper bound of Theorem 1.2 with the choice  $y = \tau^2 e^{\sqrt{3}\tau + C_1}$  for some suitably large constant  $C_1$ , by using Theorem 2.5 and replacing Theorems 2.1 and 2.2 by Theorems 2.7 and 1.10 respectively. □

**2D. The plan of the paper.** The plan of the paper is as follows. In Section 3 we gather several preliminary results on sums of divisor functions and moments of Random multiplicative functions, which will shall use throughout the paper. Section 4 will be devoted to the proof of Theorem 2.1. Theorem 2.7 will be established in Section 5. In Section 6 we investigate the distribution of large values of  $L(1, \psi_p)$  and prove Theorem 1.10. Finally, in Section 7, we investigate the positivity of sums of Legendre symbols and prove Theorems 1.12 and 1.13.

### 3. Sums of divisor functions and random multiplicative functions

In this section we gather together several preliminary results concerning sums of divisor functions, which are related to certain moments over random multiplicative functions. We let  $\{\mathbb{X}(n)\}_{n \geq 1}$  be Rademacher

random multiplicative functions, that is  $\{\mathbb{X}(p)\}_{p \text{ prime}}$  are I.I.D. random variables taking the values  $\pm 1$  with equal probability  $1/2$ , and we extend  $\mathbb{X}(n)$  multiplicatively to all positive integers by setting  $\mathbb{X}(1) = 1$  and  $\mathbb{X}(n) = \prod_{p^\ell || n} \mathbb{X}(p)^\ell$ . We start with the following lemma.

**Lemma 3.1.** *Let  $k$  be a large real number. Then for any  $0 \leq \alpha \leq \log_3 k / (2 \log k)$  we have*

$$\sum_{n=1}^{\infty} \frac{d_k(n)^2}{n^{2-\alpha}} = \exp(O(k \log \log k)), \tag{3-1}$$

and

$$\sum_{n=1}^{\infty} \frac{d_k(n^2)}{n^{2-\alpha}} = \exp(O(k \log \log k)). \tag{3-2}$$

Moreover, for all  $y > k$  we have

$$\sum_{P^-(n) > y} \frac{d_k(n^2)}{n^2} = \exp\left(O\left(\frac{k^2}{y \log y}\right)\right), \tag{3-3}$$

where here and throughout  $P^-(n)$  denotes the smallest prime factor of  $n$ .

*Proof.* The bound (3-1) follows from Lemma 3.3 of [Lamzouri 2011]. Furthermore, the estimate (3-2) follows from (3-1) upon noting that  $d_k(n^2) \leq d_k(n)^2$ . We now establish (3-3). Let  $p > y$  be a prime number. Then we have

$$\begin{aligned} \mathbb{E}\left(\left(1 - \frac{\mathbb{X}(p)}{p}\right)^{-k}\right) &= \frac{1}{2}\left(\left(1 - \frac{1}{p}\right)^{-k} + \left(1 + \frac{1}{p}\right)^{-k}\right) \\ &= \frac{1}{2}\left(\exp\left(\frac{k}{p} + O\left(\frac{k}{p^2}\right)\right) + \exp\left(-\frac{k}{p} + O\left(\frac{k}{p^2}\right)\right)\right) \\ &= 1 + O\left(\frac{k^2}{p^2}\right). \end{aligned} \tag{3-4}$$

Hence we derive

$$\begin{aligned} \sum_{P^-(n) > y} \frac{d_k(n^2)}{n^2} &= \mathbb{E}\left(\sum_{P^-(n) > y} \frac{d_k(n)\mathbb{X}(n)}{n}\right) = \prod_{p > y} \mathbb{E}\left(\left(1 - \frac{\mathbb{X}(p)}{p}\right)^{-k}\right) \\ &= \exp\left(O\left(k^2 \sum_{p > y} \frac{1}{p^2}\right)\right) = \exp\left(O\left(\frac{k^2}{y \log y}\right)\right), \end{aligned}$$

as desired. □

In order to prove Theorems 2.1 and 2.7 we need a uniform bound for the moments

$$\mathcal{M}_y(k) := \mathbb{E}\left(\left|\sum_{\substack{n > 1 \\ P^-(n) > y}} \frac{\mathbb{X}(n)}{n}\right|^{2k}\right). \tag{3-5}$$

To this end we establish the following result.

**Proposition 3.2.** *Let  $k \geq 2$  be a positive integer and  $(k \log k)/10 < y$  be real numbers. Then we have*

$$\mathcal{M}_y(k) \leq e^{O(k \log \log y / \log y)} \left( \frac{2k}{ey \log y} \right)^k.$$

**Remark 3.3.** A similar bound was obtained by Bober, Goldmakher, Granville and Koukoulopoulos in the case of Steinhaus Random multiplicative functions (see the end of Section 5 of [Bober et al. 2018]). However, our argument is easier, and can be adapted to recover this case as well.

*Proof of Proposition 3.2.* By expanding the moment and using that

$$\mathbb{E}(\mathbb{X}(n)) = \begin{cases} 1 & \text{if } n \text{ is a square,} \\ 0 & \text{otherwise,} \end{cases}$$

we get

$$\mathcal{M}_y(k) = \mathbb{E} \left( \sum_{P^-(n) > y} \frac{\tilde{d}_{2k}(n)}{n} \mathbb{X}(n) \right) = \sum_{P^-(n) > y} \frac{\tilde{d}_{2k}(n^2)}{n^2}, \tag{3-6}$$

where

$$\tilde{d}_k(n) := \left| \left\{ (n_1, \dots, n_k) \in \mathbb{N}^k, \text{ such that } n_j > 1 \text{ for all } j, \text{ and } n_1 \cdots n_k = n \right\} \right|.$$

Therefore, it follows from (3-3) that

$$\mathcal{M}_y(k) \leq \sum_{P^-(n) > y} \frac{d_{2k}(n^2)}{n^2} = \exp \left( O \left( \frac{k^2}{y \log y} \right) \right). \tag{3-7}$$

To obtain a better bound for  $\mathcal{M}_y(k)$  we consider the following “good” event

$$\mathcal{A} := \{ |\mathbb{Y}| \leq \varepsilon \}, \quad \text{where } \mathbb{Y} := \sum_{p > y} \frac{\mathbb{X}(p)}{p},$$

and  $0 < \varepsilon < 1$  is a small parameter to be chosen. Note that

$$\mathbb{P}(\mathcal{A}^c) \leq \varepsilon^{-2\ell} \mathbb{E}(|\mathbb{Y}|^{2\ell}) \ll \left( \frac{3\ell}{e\varepsilon^2 y \log y} \right)^\ell,$$

since

$$\mathbb{E}(|\mathbb{Y}|^{2\ell}) = \sum_{\substack{p_1, \dots, p_{2\ell} > y \\ p_1 \cdots p_{2\ell} = \square}} \frac{1}{p_1 \cdots p_{2\ell}} \leq \frac{(2\ell)!}{2^\ell \ell!} \left( \sum_{p > y} \frac{1}{p^2} \right)^\ell \ll e^{O(\ell / \log y)} \left( \frac{2\ell}{ey \log y} \right)^\ell, \tag{3-8}$$

which follows from the bounds  $(2\ell)! / (2^\ell \ell!) \ll (2\ell/e)^\ell$  by Stirling’s formula, and

$$\sum_{p > y} 1/p^2 = 1/(y \log y) + O(1/(y(\log y)^2))$$

by the prime number theorem. Choosing  $\ell = \lfloor (\varepsilon^2 y \log y) / 3 \rfloor$  gives

$$\mathbb{P}(\mathcal{A}^c) \ll \exp \left( -\frac{\varepsilon^2 y \log y}{3} \right). \tag{3-9}$$

We now split the moment  $\mathcal{M}_y(k)$  into two parts

$$\mathcal{M}_y(k) = \mathbb{E}\left(\left|\sum_{\substack{n>1 \\ P^-(n)>y}} \frac{\mathbb{X}(n)}{n}\right|^{2k} \cdot \mathbf{1}_{\mathcal{A}}\right) + \mathbb{E}\left(\left|\sum_{\substack{n>1 \\ P^-(n)>y}} \frac{\mathbb{X}(n)}{n}\right|^{2k} \cdot \mathbf{1}_{\mathcal{A}^c}\right), \tag{3-10}$$

where  $\mathbf{1}_{\mathcal{B}}$  is the indicator function of the event  $\mathcal{B}$ . By the Cauchy–Schwarz inequality and the estimates (3-7) and (3-9), the contribution of the second part is

$$\mathbb{E}\left(\left|\sum_{\substack{n>1 \\ P^-(n)>y}} \frac{\mathbb{X}(n)}{n}\right|^{2k} \cdot \mathbf{1}_{\mathcal{A}^c}\right) \leq \mathcal{M}_y(2k)^{1/2} \cdot \mathbb{P}(\mathcal{A}^c)^{1/2} \ll \exp\left(-\frac{\varepsilon^2 y \log y}{6} + O\left(\frac{k^2}{y \log y}\right)\right). \tag{3-11}$$

Next, we shall estimate the contribution of the first part in (3-10). Note that on the event  $\mathcal{A}$  we have  $|e^{\mathbb{Y}} - 1| \leq e^\varepsilon |\mathbb{Y}|$  and

$$\sum_{\substack{n>1 \\ P^-(n)>y}} \frac{\mathbb{X}(n)}{n} = -1 + \prod_{p>y} \left(1 - \frac{\mathbb{X}(p)}{p}\right)^{-1} = -1 + e^{\mathbb{Y} + O(1/y \log y)} = e^{\mathbb{Y}} - 1 + O\left(\frac{1}{y \log y}\right).$$

Therefore, using Minkowski’s inequality and (3-8) we derive

$$\mathbb{E}\left(\left|\sum_{\substack{n>1 \\ P^-(n)>y}} \frac{\mathbb{X}(n)}{n}\right|^{2k} \cdot \mathbf{1}_{\mathcal{A}}\right)^{1/2k} \leq e^\varepsilon \mathbb{E}(|\mathbb{Y}|^{2k})^{1/2k} + O\left(\frac{1}{y \log y}\right) \leq e^{\varepsilon + O(1/\log y)} \sqrt{\frac{2k}{ey \log y}}.$$

Choosing  $\varepsilon = (\log \log y)/\log y$  and combining this estimate with (3-11) completes the proof. □

#### 4. The distribution of the tail in Pólya’s Fourier expansion: Proof of Theorem 2.1

We start by recording two large sieve inequalities for quadratic characters, the most important of which is due to Heath-Brown [1995].

**Lemma 4.1.** *Let  $x, N \geq 2$ . Then for arbitrary complex numbers  $a_n$  we have*

$$\sum_{d \in \mathcal{F}(x)} \left| \sum_{n \leq N} a_n \chi_d(n) \right|^2 \ll (x + N^2 \log N) \sum_{\substack{m, n \leq N \\ mn = \square}} |a_m a_n|, \tag{4-1}$$

and for any  $\varepsilon > 0$  we have

$$\sum_{d \in \mathcal{F}(x)} \left| \sum_{n \leq N} a_n \chi_d(n) \right|^2 \ll_\varepsilon (xN)^\varepsilon (x + N) \sum_{\substack{m, n \leq N \\ mn = \square}} |a_m a_n|. \tag{4-2}$$

*Proof.* The first inequality is standard and is a straightforward application of the Pólya–Vinogradov inequality. It can be found for example in Lemma 1 of [Baker and Montgomery 1990], and can be traced back to [Elliott 1970]. The second inequality, which is deeper, was established by Heath-Brown [1995, Corollary 2]. □

Let  $Y := \exp(C \log_2 x \log_3 x / \log_4 x)$ , where  $C > 0$  is a suitably large constant, and  $Y \leq N \leq x^{21/40}$  be a real number. Using Lemma 4.1, we shall first prove that for all fundamental discriminants  $|d| \leq x$  except for a small exceptional set  $\mathcal{E}(x)$ , the quantity

$$\max_{\alpha \in [0,1)} \left| \sum_{N \leq n \leq 2N} \frac{\chi_d(n) a_n e(n\alpha)}{n} \right| \tag{4-3}$$

is small, where  $\{a_n\}_{n \geq 1}$  is an arbitrary sequence of complex numbers such that  $|a_n| \leq 1$  for all  $n$ . We shall use Heath-Brown’s large sieve (4-2) if  $N$  is in the range  $x^\varepsilon \leq N \leq x^{21/40}$ , and Elliott’s large sieve (4-1) in the remaining range  $Y \leq N \leq x^\varepsilon$ .

**Proposition 4.2.** *Let  $A, \varepsilon > 0$  be fixed and put  $\delta = 1/100$ . Let  $\{a_n\}_{n \geq 1}$  be an arbitrary sequence of complex numbers such that  $|a_n| \leq 1$  for all  $n$ . Let  $N_1, N_2$  be real numbers such that  $x^\varepsilon \leq N_1 < N_2 \leq 2N_1 \leq x^{21/40}$  be real numbers. Then there are at most  $O_{\varepsilon,A}(x^{1-\delta})$  fundamental discriminants  $|d| \leq x$  such that*

$$\max_{\alpha \in [0,1)} \left| \sum_{N_1 \leq n \leq N_2} \frac{\chi_d(n) a_n e(n\alpha)}{n} \right| \geq \frac{1}{(\log N_1)^A}.$$

**Proposition 4.3.** *Let  $A > 0$  be a fixed constant. Let  $\{a_n\}_{n \geq 1}$  be an arbitrary sequence of complex numbers such that  $|a_n| \leq 1$  for all  $n$ . There exist positive constants  $\varepsilon$  and  $C$  (which depend at most on  $A$ ) such that for all real numbers  $N_1, N_2$  verifying  $\exp(C \log_2 x \log_3 x / \log_4 x) \leq N_1 < N_2 \leq 2N_1 \leq x^\varepsilon$ , the number of fundamental discriminants  $|d| \leq x$  such that*

$$\max_{\alpha \in [0,1)} \left| \sum_{N_1 \leq n \leq N_2} \frac{\chi_d(n) a_n e(n\alpha)}{n} \right| \geq \frac{1}{(\log N_1)^A}$$

is

$$\ll x \exp\left(-\frac{\log x \log_4 x}{10 \log_2 x}\right).$$

To prove these results we shall bound suitable moments of (4-3). We start with the following easy lemma which reduces the problem of bounding these moments to a setting where we can apply the large sieve.

**Lemma 4.4.** *Let  $\{a_n\}_{n \geq 1}$  be an arbitrary sequence of complex numbers such that  $|a_n| \leq 1$  for all  $n$ . Let  $\mathcal{D}$  be a set of Dirichlet characters, and  $2 \leq N_1 < N_2 \leq R$  be real numbers. Define  $\mathcal{A} = \{b/R : 1 \leq b \leq R\}$ . Then for any positive integer  $k \geq 1$  we have*

$$\begin{aligned} & \left( \sum_{\chi \in \mathcal{D}} \max_{\alpha \in [0,1)} \left| \sum_{N_1 \leq n \leq N_2} \frac{\chi(n) a_n e(n\alpha)}{n} \right|^{2k} \right)^{1/2k} \\ & \leq \left( \sum_{\alpha \in \mathcal{A}} \sum_{\chi \in \mathcal{D}} \left| \sum_{N_1^k \leq n \leq N_2^k} \frac{\chi(n) g_{N_1, N_2, k}(n, \alpha)}{n} \right|^2 \right)^{1/2k} + O(|\mathcal{D}|^{1/2k} N_2 R^{-1}), \end{aligned}$$

where

$$g_{N_1, N_2, k}(n, \alpha) := \sum_{\substack{N_1 \leq n_1, \dots, n_k \leq N_2 \\ n_1 \cdots n_k = n}} \prod_{j=1}^k a_{n_j} e(\alpha n_j), \tag{4-4}$$

and the implicit constant in the error term is absolute.

*Proof.* First, we observe that for all  $\alpha \in [0, 1)$  there exists  $\beta_\alpha \in \mathcal{A}$  such that  $|\alpha - \beta_\alpha| \leq 1/R$ . In this case we have  $e(\alpha n) = e(\beta_\alpha n) + O(n/R)$ , and hence

$$\max_{\alpha \in [0,1)} \left| \sum_{N_1 \leq n \leq N_2} \frac{\chi(n)a_n e(\alpha n)}{n} \right| = \max_{\beta \in \mathcal{A}} \left| \sum_{N_1 \leq n \leq N_2} \frac{\chi(n)a_n e(\beta n)}{n} \right| + O\left(\frac{N_2}{R}\right).$$

Therefore, it follows from Minkowski's inequality that

$$\begin{aligned} & \left( \sum_{\chi \in \mathcal{D}} \max_{\alpha \in [0,1)} \left| \sum_{N_1 \leq n \leq N_2} \frac{\chi(n)a_n e(\alpha n)}{n} \right|^{2k} \right)^{1/2k} \\ & \leq \left( \sum_{\chi \in \mathcal{D}} \max_{\alpha \in \mathcal{A}} \left| \sum_{N_1 \leq n \leq N_2} \frac{\chi(n)a_n e(\alpha n)}{n} \right|^{2k} \right)^{1/2k} + O(|\mathcal{D}|^{1/2k} N_2 R^{-1}) \\ & \leq \left( \sum_{\alpha \in \mathcal{A}} \sum_{\chi \in \mathcal{D}} \left| \sum_{N_1 \leq n \leq N_2} \frac{\chi(n)a_n e(\alpha n)}{n} \right|^{2k} \right)^{1/2k} + O(|\mathcal{D}|^{1/2k} N_2 R^{-1}). \end{aligned} \quad (4-5)$$

The lemma follows upon noting that

$$\left| \sum_{N_1 \leq n \leq N_2} \frac{\chi(n)a_n e(\alpha n)}{n} \right|^{2k} = \left| \sum_{N_1^k \leq n \leq N_2^k} \frac{\chi(n)g_{N_1, N_2, k}(n, \alpha)}{n} \right|^2. \quad \square$$

*Proof of Proposition 4.2.* Let

$$k := \begin{cases} 2 & \text{if } \sqrt{x} \leq N_1 \leq x^{21/40}, \\ \lfloor \log x / \log N_1 \rfloor & \text{if } x^\varepsilon \leq N_1 < \sqrt{x}. \end{cases}$$

We observe that  $2 \leq k \leq 1/\varepsilon$  by our assumption on  $N_1$ , and that  $|g_{N_1, N_2, k}(n, \alpha)| \leq d_k(n)$  for all  $N_1, N_2$  and  $\alpha$ . Let  $\delta = 1/100$  and  $\nu > 0$  be a small parameter to be chosen. Using Lemma 4.4 with the choice  $R = N_1^{1+\delta}$  together with the large sieve inequality (4-2) and the easy inequality  $|a + b|^k \leq 2^k(|a|^k + |b|^k)$  we obtain

$$\begin{aligned} & \sum_{d \in \mathcal{F}(x)} \max_{\alpha \in [0,1)} \left| \sum_{N_1 \leq n \leq N_2} \frac{\chi_d(n)a_n e(\alpha n)}{n} \right|^{2k} \\ & \ll_\varepsilon \sum_{\alpha \in \mathcal{A}} \sum_{d \in \mathcal{F}(x)} \left| \sum_{N_1^k \leq n \leq N_2^k} \frac{\chi_d(n)g_{N_1, N_2, k}(n, \alpha)}{n} \right|^2 + \frac{x}{N_1^{2k\delta}} \\ & \ll_{\varepsilon, \nu} (N_2^k x)^\nu (x + N_2^k) \sum_{\alpha \in \mathcal{A}} \sum_{\substack{N_1^k \leq n_1, n_2 \leq N_2^k \\ n_1 n_2 = \square}} \frac{|g_{N_1, N_2, k}(n_1, \alpha)g_{N_1, N_2, k}(n_2, \alpha)|}{n_1 n_2} + \frac{x}{N_1^{2k\delta}} \\ & \ll_{\varepsilon, \nu} x^{21/20+3\nu} N_1^{1+\delta} \sum_{\substack{N_1^k \leq n_1, n_2 \leq N_2^k \\ n_1 n_2 = \square}} \frac{d_k(n_1)d_k(n_2)}{n_1 n_2} + \frac{x}{N_1^{2k\delta}}, \end{aligned} \quad (4-6)$$

since  $N_2^k \ll_\varepsilon x^{21/20}$  by our assumption on  $k$  and  $N_1$ . Now, writing  $n_1 n_2 = n^2$ , we get

$$\sum_{\substack{N_1^k \leq n_1, n_2 \leq N_2^k \\ n_1 n_2 = \square}} \frac{d_k(n_1) d_k(n_2)}{n_1 n_2} \leq \sum_{n \geq N_1^k} \frac{d_{2k}(n^2)}{n^2} \leq N_1^{-k(1-\nu)} \sum_{n=1}^{\infty} \frac{d_{2k}(n^2)}{n^{1+\nu}} \ll_{\varepsilon, \nu} N_1^{-k(1-\nu)}. \tag{4-7}$$

Now, since  $k \geq 2$ , then  $N_1^{3k/2} \geq N_1^{k+1} \geq x$ , by our definition of  $k$ , which implies that  $N_1^k \geq x^{2/3}$ . Thus, choosing  $\nu$  to be suitably small, and combining this estimate with (4-6) gives

$$\sum_{d \in \mathcal{F}(x)} \max_{\alpha \in [0,1]} \left| \sum_{N_1 \leq n \leq N_2} \frac{\chi_d(n) a_n e(\alpha n)}{n} \right|^{2k} \ll_{\varepsilon, \nu} x^{21/20+3\nu} N_1^{1+\delta} x^{-2/3(1-\nu)} + \frac{x}{N_1^{2k\delta}} \ll_\varepsilon x^{1-4\delta/3}.$$

Finally, the number of fundamental discriminants  $d \in \mathcal{F}(x)$  such that

$$\max_{\alpha \in [0,1]} \left| \sum_{N_1 \leq n \leq N_2} \frac{\chi_d(n) a_n e(\alpha n)}{n} \right| \geq \frac{1}{(\log N_1)^A}$$

is

$$\leq (\log N_1)^{2Ak} \sum_{d \in \mathcal{F}(x)} \max_{\alpha \in [0,1]} \left| \sum_{N_1 \leq n \leq N_2} \frac{\chi_d(n) a_n e(\alpha n)}{n} \right|^{2k} \ll_{\varepsilon, A} x^{1-\delta},$$

which completes the proof. □

*Proof of Proposition 4.3.* We proceed similarly to the proof of Proposition 4.2 but we choose  $k = \lfloor \log x / (3 \log N_1) \rfloor$  in this case. Then we have

$$1/(3\varepsilon) - 1 \leq k \leq (\log x \log_4 x) / (3C \log_2 x \log_3 x)$$

by our assumption on  $N_1$ . Using Lemma 4.4 with  $R = N_1^2$  together with the large sieve inequality (4-1) and the easy inequality  $|a + b|^k \leq 2^k(|a|^k + |b|^k)$  we obtain

$$\begin{aligned} \sum_{d \in \mathcal{F}(x)} \max_{\alpha \in [0,1]} \left| \sum_{N_1 \leq n \leq N_2} \frac{\chi_d(n) a_n e(\alpha n)}{n} \right|^{2k} &\ll e^{O(k)} \sum_{\alpha \in \mathcal{A}} \sum_{d \in \mathcal{F}(x)} \left| \sum_{N_1^k \leq n \leq N_2^k} \frac{\chi_d(n) g_{N_1, N_2, k}(n, \alpha)}{n} \right|^2 + \frac{x e^{O(k)}}{N_1^{2k}} \\ &\ll x e^{O(k)} \sum_{\alpha \in \mathcal{A}} \sum_{\substack{N_1^k \leq n_1, n_2 \leq N_2^k \\ n_1 n_2 = \square}} \frac{|g_{N_1, N_2, k}(n_1, \alpha) g_{N_1, N_2, k}(n_2, \alpha)|}{n_1 n_2} + \frac{x e^{O(k)}}{N_1^{2k}} \\ &\ll x N_1^2 e^{O(k)} \sum_{n \geq N_1^k} \frac{d_{2k}(n^2)}{n^2} + \frac{x}{N_1^{2k}} \end{aligned} \tag{4-8}$$

by (4-7), since  $|g_{N_1, N_2, k}(n, \alpha)| \leq d_k(n)$  for all  $N_1, N_2$  and  $\alpha$ . To bound the sum over  $n$ , we shall use Rankin's trick. Choosing  $\nu = \log_3 k / (2 \log k)$  and using (3-2) we obtain

$$\sum_{n \geq N_1^k} \frac{d_{2k}(n^2)}{n^2} \leq N_1^{-k\nu} \sum_{n=1}^{\infty} \frac{d_{2k}(n^2)}{n^{2-\nu}} \ll \exp\left(-\frac{k \log_3 k \log N_1}{2 \log k} + O(k \log_2 k)\right).$$

Inserting this estimate in (4-8) we deduce that

$$\sum_{d \in \mathcal{F}(x)} \max_{\alpha \in [0,1]} \left| \sum_{N_1 \leq n \leq N_2} \frac{\chi_d(n) a_n e(\alpha n)}{n} \right|^{2k} \ll x \exp\left(-\frac{k \log_3 k \log N_1}{2 \log k} + O(k \log_2 k + \log N_1)\right).$$

Therefore, the number of fundamental discriminants  $d \in \mathcal{F}(x)$  such that

$$\max_{\alpha \in [0,1]} \left| \sum_{N_1 \leq n \leq N_2} \frac{\chi_d(n) a_n e(\alpha n)}{n} \right| \geq \frac{1}{(\log N_1)^A}$$

is

$$\begin{aligned} &\leq (\log N_1)^{2Ak} \sum_{d \in \mathcal{F}(x)} \max_{\alpha \in [0,1]} \left| \sum_{N_1 \leq n \leq N_2} \frac{\chi_d(n) a_n e(\alpha n)}{n} \right|^{2k} \\ &\ll x \exp\left(-\frac{k \log_3 k \log N_1}{2 \log k} + O(k \log_2 k + Ak \log_2 N_1 + \log N_1)\right) \\ &\ll x \exp\left(-\frac{\log x \log_4 x}{10 \log_2 x}\right) \end{aligned}$$

by our assumption on  $N_1$  and  $k$ , if  $\varepsilon$  is suitably small and  $C$  is suitably large. This completes the proof.  $\square$

Using Propositions 4.2 and 4.3 reduces the proof of Theorem 2.1 to studying the distribution of the maximum over  $\alpha \in [0, 1)$  of the very short sum

$$\left| \sum_{\substack{1 \leq n \leq Y \\ P^+(n) > y}} \frac{\chi_d(n) h(n) e(\alpha n)}{n} \right|.$$

To this end we shall use the large sieve to bound its  $2k$ -th moments for every  $k \leq (\log x) / (3 \log Y)$ .

**Proposition 4.5.** *Let  $h(n)$  be a completely multiplicative function such that  $|h(n)| \leq 1$  for all  $n$ . Let  $C > 0$  be a suitably large constant and put  $Y = \exp(C \log_2 x \log_3 x / \log_4 x)$ . For any positive integer  $2 \leq k \leq \log x / (3 \log Y)$  and real number  $(k \log k) / 10 \leq y \leq k^2$  we have*

$$\sum_{d \in \mathcal{F}(x)} \max_{\alpha \in [0,1]} \left| \sum_{\substack{1 \leq n \leq Y \\ P^+(n) > y}} \frac{\chi_d(n) h(n) e(\alpha n)}{n} \right|^{2k} \ll x \left( e^{O\left(\frac{k \log_2 k}{\log k}\right)} \left( \frac{2e^{2\gamma} k \log y}{ey} \right)^k + O\left(\frac{1}{(\log k)^{10k}}\right) \right).$$

*Proof of Proposition 4.5.* We first define

$$S_d(y, Y) = \max_{\alpha \in [0,1]} \left| \sum_{\substack{2 \leq n \leq Y \\ P^-(n) > y}} \frac{\chi_d(n)h(n)e(\alpha n)}{n} \right|. \tag{4-9}$$

Then we have

$$\sum_{\substack{1 \leq n \leq Y \\ P^+(n) > y}} \frac{\chi_d(n)h(n)e(\alpha n)}{n} = \sum_{\substack{1 \leq a \leq Y \\ P^+(a) \leq y}} \frac{\chi_d(a)h(a)}{a} \sum_{\substack{1 < b \leq Y/a \\ P^-(b) > y}} \frac{\chi_d(b)h(b)e(\alpha ab)}{b},$$

and hence

$$\max_{\alpha \in [0,1]} \left| \sum_{\substack{1 \leq n \leq Y \\ P^+(n) > y}} \frac{\chi_d(n)h(n)e(\alpha n)}{n} \right| \leq \sum_{\substack{1 \leq a \leq Y \\ P^+(a) \leq y}} \frac{S_d(y, Y/a)}{a}.$$

Therefore, using Hölder’s inequality we obtain

$$\begin{aligned} \sum_{d \in \mathcal{F}(x)} \max_{\alpha \in [0,1]} \left| \sum_{\substack{1 \leq n \leq Y \\ P^+(n) > y}} \frac{\chi_d(n)h(n)e(\alpha n)}{n} \right|^{2k} &\leq \sum_{d \in \mathcal{F}(x)} \left( \sum_{P^+(a) \leq y} \frac{1}{a} \right)^{2k-1} \sum_{\substack{1 \leq a \leq Y \\ P^+(a) \leq y}} \frac{S_d(y, Y/a)^{2k}}{a} \\ &\leq (e^\gamma \log y + O(1))^{2k-1} \sum_{\substack{1 \leq a \leq Y \\ P^+(a) \leq y}} \frac{1}{a} \sum_{d \in \mathcal{F}(x)} S_d(y, Y/a)^{2k}. \end{aligned} \tag{4-10}$$

by Mertens’ theorem. We shall now bound the moments

$$\sum_{d \in \mathcal{F}(x)} S_d(y, z)^{2k}$$

uniformly in  $2 \leq y < z \leq Y$ . To this end we split the inner sum over  $n$  into two parts  $y < n \leq N$ , and  $N < n \leq z$ , where

$$N = \min(z, \exp(C \log k \log_2 k / \log_3 k)). \tag{4-11}$$

Note that the second part will be empty unless  $z > \exp(C \log k \log_2 k / \log_3 k)$ . Using Minkowski’s inequality we get

$$\begin{aligned} \left( \sum_{d \in \mathcal{F}(x)} S_d(y, z)^{2k} \right)^{1/2k} &\leq \left( \sum_{d \in \mathcal{F}(x)} \max_{\alpha \in [0,1]} \left| \sum_{\substack{2 \leq n \leq N \\ P^-(n) > y}} \frac{\chi_d(n)h(n)e(\alpha n)}{n} \right|^{2k} \right)^{1/2k} \\ &\quad + \left( \sum_{d \in \mathcal{F}(x)} \max_{\alpha \in [0,1]} \left| \sum_{\substack{N < n \leq z \\ P^-(n) > y}} \frac{\chi_d(n)h(n)e(\alpha n)}{n} \right|^{2k} \right)^{1/2k}. \end{aligned} \tag{4-12}$$

We start by bounding the first term. By Lemma 4.4 with  $R := \exp(2C \log k \log_2 k / \log_3 k)$  and

$$a_n = \begin{cases} h(n) & \text{if } P^-(n) > y \\ 0 & \text{otherwise,} \end{cases}$$

we obtain

$$\begin{aligned} & \left( \sum_{d \in \mathcal{F}(x)} \max_{\alpha \in [0,1)} \left| \sum_{\substack{2 \leq n \leq N \\ P^-(n) > y}} \frac{\chi_d(n)h(n)e(\alpha n)}{n} \right|^{2k} \right)^{1/2k} \\ & \leq \left( \sum_{\alpha \in \mathcal{A}} \sum_{d \in \mathcal{F}(x)} \left| \sum_{\substack{n \leq N^k \\ P^-(n) > y}} \frac{\chi_d(n)g_{2,N,k}(n, \alpha)}{n} \right|^2 \right)^{1/2k} + O(x^{1/2k}R^{-1/2}), \end{aligned} \quad (4-13)$$

where  $\mathcal{A} = \{b/R : 1 \leq b \leq R\}$ . We now proceed similarly to the proof of (4-8). Using the large sieve inequality (4-1), and noting that  $N^k \leq x^{1/3}$  and  $|g_{2,N,k}(n, \alpha)| \leq \tilde{d}_k(n)$  for all  $\alpha$  and  $N$  we derive

$$\begin{aligned} \sum_{d \in \mathcal{F}(x)} \left| \sum_{\substack{n \leq N^k \\ P^-(n) > y}} \frac{\chi_d(n)g_{2,N,k}(n, \alpha)}{n} \right|^2 & \ll x \sum_{\substack{n_1, n_2 \leq N^k \\ P^-(n_1 n_2) > y \\ n_1 n_2 = \square}} \frac{\tilde{d}_k(n_1)\tilde{d}_k(n_2)}{n_1 n_2} \leq x \sum_{P^-(n) > y} \frac{\tilde{d}_{2k}(n^2)}{n^2} \\ & \ll x e^{O(k \log \log y / \log y)} \left( \frac{2k}{ey \log y} \right)^k, \end{aligned} \quad (4-14)$$

where the last inequality follows from (3-6) and Proposition 3.2. Inserting this estimate in (4-13) implies that

$$\left( \sum_{d \in \mathcal{F}(x)} \max_{\alpha \in [0,1)} \left| \sum_{\substack{2 \leq n \leq N \\ P^-(n) > y}} \frac{\chi_d(n)h(n)e(\alpha n)}{n} \right|^{2k} \right)^{1/2k} \leq x^{1/(2k)} \left( e^{O(\log_2 k / \log k)} \left( \frac{2k}{ey \log y} \right)^{1/2} + O\left( \frac{1}{R^{1/2}} \right) \right). \quad (4-15)$$

We now assume that  $z > \exp(C \log k \log_2 k / \log_3 k)$  and bound the second term on the right-hand side of (4-12). We shall split the inner sum over  $n$  into dyadic intervals. Let  $J_1 = \lfloor \log N / \log 2 \rfloor$ ,  $J_2 = \lfloor \log z / \log 2 \rfloor$ , and define  $t_{J_1} = N$ ,  $t_{J_2+1} = z$ , and  $t_j := 2^j$  for  $J_1 + 1 \leq j \leq J_2$ . Using Hölder's inequality we obtain

$$\begin{aligned} \left| \sum_{\substack{N < n \leq z \\ P^-(n) > y}} \frac{\chi_d(n)h(n)e(\alpha n)}{n} \right|^{2k} & = \left| \sum_{J_1 \leq j \leq J_2} \frac{1}{j^2} \left( j^2 \sum_{\substack{t_j < n \leq t_{j+1} \\ P^-(n) > y}} \frac{\chi_d(n)h(n)e(\alpha n)}{n} \right) \right|^{2k} \\ & \leq \left( \sum_{J_1 \leq j \leq J_2} \frac{1}{j^{4k}} \right)^{2k-1} \sum_{J_1 \leq j \leq J_2} j^{4k} \left| \sum_{\substack{t_j < n \leq t_{j+1} \\ P^-(n) > y}} \frac{\chi_d(n)h(n)e(\alpha n)}{n} \right|^{2k} \\ & \leq \left( \frac{C_1}{\log N} \right)^{2k+1} \sum_{J_1 \leq j \leq J_2} j^{4k} \left| \sum_{\substack{t_j < n \leq t_{j+1} \\ P^-(n) > y}} \frac{\chi_d(n)h(n)e(\alpha n)}{n} \right|^{2k}, \end{aligned} \quad (4-16)$$

for some constant  $C_1 > 0$ . Therefore, this reduces the problem to bounding the following moments over dyadic intervals  $[t_j, t_{j+1}]$ :

$$\sum_{d \in \mathcal{F}(x)} \max_{\alpha \in [0,1)} \left| \sum_{\substack{t_j < n \leq t_{j+1} \\ P^-(n) > y}} \frac{\chi_d(n)h(n)e(\alpha n)}{n} \right|^{2k}.$$

Let  $\mathcal{B}_j = \{b/4^j : 1 \leq b \leq 4^j\}$ . By the easy inequality  $|a + b|^k \leq 2^k(|a|^k + |b|^k)$  together with Lemma 4.4 with  $R = 4^j$  and the same choice of  $a_n$  as before, we derive

$$\begin{aligned} \sum_{d \in \mathcal{F}(x)} \max_{\alpha \in [0,1)} \left| \sum_{\substack{t_j < n \leq t_{j+1} \\ P^-(n) > y}} \frac{\chi_d(n)h(n)e(\alpha n)}{n} \right|^{2k} \\ \ll e^{O(k)} \sum_{\alpha \in \mathcal{B}_j} \sum_{d \in \mathcal{F}(x)} \left| \sum_{\substack{t_j^k < n \leq t_{j+1}^k \\ P^-(n) > y}} \frac{\chi_d(n)g_{t_j, t_{j+1}, k}(n, \alpha)}{n} \right|^2 + \frac{x e^{O(k)}}{2^{2jk}}, \end{aligned} \quad (4-17)$$

since  $t_j \asymp t_{j+1} \asymp 2^j$ . Furthermore, similarly to the proof of (4-8), it follows from the large sieve inequality (4-1) that the main term on the right-hand side of (4-17) is

$$\ll x e^{O(k)} \sum_{\alpha \in \mathcal{B}_j} \sum_{\substack{t_j^k < n_1, n_2 \leq t_{j+1}^k \\ P^-(n_1 n_2) > y \\ n_1 n_2 = \square}} \frac{|g_{t_j, t_{j+1}, k}(n_1, \alpha)g_{t_j, t_{j+1}, k}(n_2, \alpha)|}{n_1 n_2} \ll e^{O(k)} 4^j x \sum_{n \geq t_j^k} \frac{d_{2k}(n^2)}{n^2}, \quad (4-18)$$

since  $t_{j+1}^k \leq z^k \leq Y^k \leq x^{1/3}$ . We now put  $\nu = \log_3 k / (2 \log k)$  and use (3-2) to get

$$\sum_{n \geq t_j^k} \frac{d_{2k}(n^2)}{n^2} \leq t_j^{-k\nu} \sum_{n=1}^{\infty} \frac{d_{2k}(n^2)}{n^{2-\nu}} \ll \exp\left(-\frac{jk \log_3 k}{4 \log k} + O(k \log_2 k)\right). \quad (4-19)$$

Inserting these estimates in (4-17) gives

$$\sum_{d \in \mathcal{F}(x)} \max_{\alpha \in [0,1)} \left| \sum_{\substack{t_j < n \leq t_{j+1} \\ P^-(n) > y}} \frac{\chi_d(n)h(n)e(\alpha n)}{n} \right|^{2k} \ll x \exp\left(-\frac{jk \log_3 k}{5 \log k} + O(k \log_2 k)\right).$$

Using this last estimate together with (4-16), and noting that  $j^4 \leq \exp(j \log_3 k / (20 \log k))$  for all  $j \geq J_1$  by our choice of  $N$  if  $C$  is suitably large, we derive

$$\begin{aligned} \sum_{d \in \mathcal{F}(x)} \max_{\alpha \in [0,1)} \left| \sum_{\substack{N < n \leq z \\ P^-(n) > y}} \frac{\chi_d(n)h(n)e(\alpha n)}{n} \right|^{2k} &\ll x e^{O(k \log_2 k)} \sum_{J_1 \leq j \leq J_2} \exp\left(-\frac{jk \log_3 k}{20 \log k}\right) \\ &\ll x \exp\left(-\frac{(\log N)k \log_3 k}{20 \log k} + O(k \log_2 k)\right) \\ &\ll x \exp\left(-\frac{C}{40} k \log_2 k\right). \end{aligned}$$

Combining this estimate with (4-12) and (4-15) we obtain

$$\sum_{d \in \mathcal{F}(x)} S_d(y, z)^{2k} \ll x \left( e^{O(\log_2 k / \log k)} \left( \frac{2k}{ey \log y} \right)^{1/2} + O\left( \frac{1}{(\log k)^{C/80}} \right) \right)^{2k},$$

uniformly for  $y < z \leq Y$ . Therefore, inserting this estimate in (4-10) we deduce

$$\sum_{d \in \mathcal{F}(x)} \max_{\alpha \in [0, 1)} \left| \sum_{\substack{n \leq Y \\ P^+(n) > y}} \frac{\chi_d(n) h(n) e(\alpha n)}{n} \right|^{2k} \ll x \left( e^{O(\frac{\log_2 k}{\log k})} \left( \frac{2e^{2\gamma} k \log y}{ey} \right)^{1/2} + O\left( \frac{1}{(\log k)^{C/80-1}} \right) \right)^{2k}.$$

The result follows upon choosing  $C$  to be suitably large, and using the basic inequality<sup>6</sup>  $(a + b)^{2m} \leq (a^{2m} + \sqrt{b}^{2m})(1 + \sqrt{b})^{2m} \leq (a^{2m} + b^m)e^{2m\sqrt{b}}$ , which is valid for all real numbers  $a, b > 0$  and positive integers  $m$ . □

We end this section by deducing Theorem 2.1 from Propositions 4.2, 4.3, and 4.5.

*Proof of Theorem 2.1.* Let  $Y := \exp(C \log_2 x \log_3 x / \log_4 x)$  for some suitably large constant  $C > 0$ . Let  $L_1 := \lfloor \log Y / \log 2 \rfloor$ ,  $L_2 := \lfloor \log Z / \log 2 \rfloor$ , and define  $s_{L_1} := Y$ ,  $s_{L_2+1} := Z$ , and  $s_\ell := 2^\ell$  for  $L_1 + 1 \leq \ell \leq L_2$ . Then we have

$$\begin{aligned} \max_{\alpha \in [0, 1)} \left| \sum_{\substack{1 \leq n \leq Z \\ P^+(n) > y}} \frac{\chi_d(n) h(n) e(\alpha n)}{n} \right| \\ \leq \max_{\alpha \in [0, 1)} \left| \sum_{\substack{1 \leq n \leq Y \\ P^+(n) > y}} \frac{\chi_d(n) h(n) e(\alpha n)}{n} \right| + \sum_{L_1 \leq \ell \leq L_2} \max_{\alpha \in [0, 1)} \left| \sum_{\substack{s_\ell \leq n \leq s_{\ell+1} \\ P^+(n) > y}} \frac{\chi_d(n) h(n) e(\alpha n)}{n} \right|. \end{aligned} \quad (4-20)$$

Using Propositions 4.2 and 4.3 with  $A = 2$  and

$$a_n = \begin{cases} h(n) & \text{if } P^+(n) > y, \\ 0 & \text{otherwise,} \end{cases}$$

we deduce that

$$\sum_{L_1 \leq \ell \leq L_2} \max_{\alpha \in [0, 1)} \left| \sum_{\substack{s_\ell \leq n \leq s_{\ell+1} \\ P^+(n) > y}} \frac{\chi_d(n) h(n) e(\alpha n)}{n} \right| \ll \sum_{L_1 \leq \ell \leq L_2} \frac{1}{\ell^2} \ll \frac{\log_4 x}{\log_2 x \log_3 x}, \quad (4-21)$$

for all fundamental discriminants  $|d| \leq x$  except for a set  $\mathcal{E}(x)$  of size

$$|\mathcal{E}(x)| \ll L_2 x \exp\left(-\frac{\log x \log_4 x}{10 \log_2 x}\right) \ll x \exp\left(-\frac{\log x \log_4 x}{20 \log_2 x}\right).$$

For  $B > 0$  we let  $\tilde{\Psi}_{x,y}(B)$  be the proportion of fundamental discriminants  $|d| \leq x$  such that

$$\max_{\alpha \in [0, 1)} \left| \sum_{\substack{1 \leq n \leq Y \\ P^+(n) > y}} \frac{\chi_d(n) h(n) e(\alpha n)}{n} \right| > e^\gamma B.$$

<sup>6</sup>This inequality simply follows by considering the two cases  $a \leq \sqrt{b}$  and  $a \geq \sqrt{b}$ .

Let  $k$  be a positive integer satisfying the assumptions of Proposition 4.5. Then, it follows from this result that

$$\begin{aligned} \tilde{\Psi}_{x,y}(B) &\leq (e^\gamma B)^{-2k} \frac{1}{|\mathcal{F}(x)|} \sum_{d \in \mathcal{F}(x)} \max_{\alpha \in [0,1]} \left| \sum_{\substack{1 \leq n \leq Y \\ P^+(n) > y}} \frac{\chi_d(n)h(n)e(\alpha n)}{n} \right|^{2k} \\ &\ll e^{O\left(\frac{k \log_2 k}{\log k}\right)} \left(\frac{2k \log y}{eB^2y}\right)^k + O\left(\frac{1}{(e^\gamma B(\log k)^5)^{2k}}\right). \end{aligned}$$

We now assume that  $1/(\log y)^2 \leq B \leq \sqrt{20}$  and choose  $k = \lfloor (B^2y)/(2 \log y) \rfloor$ . This gives

$$\tilde{\Psi}_{x,y}(B) \ll \exp\left(-\frac{B^2y}{2 \log y} \left(1 + O\left(\frac{\log_2 y}{\log y}\right)\right)\right). \tag{4-22}$$

Combining this estimate with (4-20) and (4-21) and choosing  $B = A - C_0 \log_4 x / (\log_2 x \log_3 x)$  for some suitably large constant  $C_0$  completes the proof. □

**5. The distribution of the tail of the Pólya Fourier series for prime discriminants:  
Proof of Theorem 2.7**

In order to prove Theorem 2.7 we follow the same lines of the proof of Theorem 2.1 but we replace the large sieve inequality (4-1) by the following large sieve inequality for prime discriminants, which is a special case of Lemma 9 of [Montgomery and Vaughan 1979] (see also Lemma 1 of [Baker and Montgomery 1990]).

**Lemma 5.1** [Montgomery and Vaughan 1979, Lemma 9]. *Let  $x, N$  be real numbers such that  $x \geq 2$  and  $2 \leq N \leq x^{1/3}$ . Then for arbitrary complex numbers  $a_1, \dots, a_N$  we have*

$$\sum_{p \leq x} \left| \sum_{n \leq N} a_n \psi_p(n) \right|^2 \ll \frac{x}{\log x} \sum_{\substack{m, n \leq N \\ mn = \square}} |a_m a_n|.$$

Since the number of primes up to  $x$  is smaller by a factor of size  $\log x$  compared to the number of fundamental discriminants up to  $x$ , it suffices to establish the analogue of Proposition 4.5 for prime discriminants. Indeed, the savings in Propositions 4.2 and 4.3 are much larger than  $\log x$ , and hence we can use these results in this setting as well by simply embedding the set of primes  $p \leq x$  in the set of fundamental discriminants  $d$  with  $|d| \leq x$ .

**Proposition 5.2.** *Let  $h(n)$  be a completely multiplicative function such that  $|h(n)| \leq 1$  for all  $n$ . Let  $Y = \exp(C \log_2 x \log_3 x / \log_4 x)$ , for some suitably large constant  $C$ . Then, for any positive integer  $2 \leq k \leq \log x / (3 \log Y)$  and real number  $(k \log k) / 10 \leq y \leq k^2$  we have*

$$\sum_{p \leq x} \max_{\alpha \in [0,1]} \left| \sum_{\substack{1 \leq n \leq Y \\ P^+(n) > y}} \frac{\psi_p(n)h(n)e(\alpha n)}{n} \right|^{2k} \ll \pi(x) \left( e^{O\left(\frac{k \log_2 k}{\log k}\right)} \left(\frac{2e^{2\gamma} k \log y}{ey}\right)^k + O\left(\frac{1}{(\log k)^{10k}}\right) \right).$$

*Proof.* We shall closely follow the proof of Proposition 4.5 and only indicate where the main changes occur. First we define

$$\tilde{S}_p(y, Y) = \max_{\alpha \in [0,1)} \left| \sum_{\substack{2 \leq n \leq Y \\ P^-(n) > y}} \frac{\psi_p(n)h(n)e(\alpha n)}{n} \right|. \quad (5-1)$$

Then, similarly to (4-10) we have

$$\sum_{p \leq x} \max_{\alpha \in [0,1)} \left| \sum_{\substack{n \leq Y \\ P^+(n) > y}} \frac{\psi_p(n)h(n)e(\alpha n)}{n} \right|^{2k} \leq (e^y \log y + O(1))^{2k-1} \sum_{\substack{a \leq Y \\ P^+(a) \leq y}} \frac{1}{a} \sum_{p \leq x} \tilde{S}_p(y, Y/a)^{2k}. \quad (5-2)$$

Let  $y < z \leq Y$  be a real number and  $N$  be defined by (4-11). Using Minkowski's inequality as in (4-12) we get

$$\begin{aligned} \left( \sum_{p \leq x} \tilde{S}_p(y, z)^{2k} \right)^{1/2k} &\leq \left( \sum_{p \leq x} \max_{\alpha \in [0,1)} \left| \sum_{\substack{2 \leq n \leq N \\ P^-(n) > y}} \frac{\psi_p(n)h(n)e(\alpha n)}{n} \right|^{2k} \right)^{1/2k} \\ &\quad + \left( \sum_{p \leq x} \max_{\alpha \in [0,1)} \left| \sum_{\substack{N < n \leq z \\ P^-(n) > y}} \frac{\psi_p(n)h(n)e(\alpha n)}{n} \right|^{2k} \right)^{1/2k}. \end{aligned} \quad (5-3)$$

We start by bounding the first term. Let  $R = \exp(2C \log k \log_2 k / \log_3 k)$ . Using the same argument leading to (4-15) and replacing the large sieve inequality (4-1) by Lemma 5.1 we obtain

$$\begin{aligned} \left( \sum_{p \leq x} \max_{\alpha \in [0,1)} \left| \sum_{\substack{2 \leq n \leq N \\ P^-(n) > y}} \frac{\psi_p(n)h(n)e(\alpha n)}{n} \right|^{2k} \right)^{1/2k} \\ \leq \pi(x)^{1/(2k)} \left( e^{O(\log_2 k / \log k)} \left( \frac{2k}{ey \log y} \right)^{1/2} + O\left( \frac{1}{R^{1/2}} \right) \right). \end{aligned} \quad (5-4)$$

We now assume that  $z > \exp(C \log k \log_2 k / \log_3 k)$  and bound the second term on the right-hand side of (5-3). By (4-16) we have

$$\left| \sum_{\substack{N < n \leq z \\ P^-(n) > y}} \frac{\psi_p(n)h(n)e(\alpha n)}{n} \right|^{2k} \leq \left( \frac{C_1}{\log N} \right)^{2k+1} \sum_{J_1 \leq j \leq J_2} j^{4k} \left| \sum_{\substack{t_j < n \leq t_{j+1} \\ P^-(n) > y}} \frac{\psi_p(n)h(n)e(\alpha n)}{n} \right|^{2k}, \quad (5-5)$$

for some constant  $C_1 > 0$ , where  $J_1 = \lfloor \log N / \log 2 \rfloor$ ,  $J_2 = \lfloor \log z / \log 2 \rfloor$ ,  $t_{J_1} = N$ ,  $t_{J_2+1} = z$ , and  $t_j = 2^j$  for  $J_1 + 1 \leq j \leq J_2$ . As before we let  $\mathcal{B}_j = \{b/4^j : 1 \leq b \leq 4^j\}$ . Combining Lemmas 4.4 and 5.1 with (4-18) and (4-19) we deduce that

$$\begin{aligned} \sum_{p \leq x} \max_{\alpha \in [0,1)} \left| \sum_{\substack{t_j < n \leq t_{j+1} \\ P^-(n) > y}} \frac{\psi_p(n)h(n)e(\alpha n)}{n} \right|^{2k} &\ll e^{O(k)} \sum_{\alpha \in \mathcal{B}_j} \sum_{p \leq x} \left| \sum_{\substack{t_j^k < n \leq t_{j+1}^k \\ P^-(n) > y}} \frac{\psi_p(n)g_{t_j, t_{j+1}, k}(n, \alpha)}{n} \right|^2 + \frac{\pi(x)e^{O(k)}}{2^{2jk}} \\ &\ll \pi(x) \exp\left( -\frac{jk \log_3 k}{5 \log k} + O(k \log_2 k) \right). \end{aligned}$$

Using this last estimate together with (5-5) we derive

$$\sum_{p \leq x} \max_{\alpha \in [0,1)} \left| \sum_{\substack{N < n \leq z \\ P^-(n) > y}} \frac{\psi_p(n)h(n)e(\alpha n)}{n} \right|^{2k} \ll \pi(x) \exp\left(-\frac{C_1}{40}k \log_2 k\right).$$

We have now established the key estimates over prime discriminants. Thus we continue the proof along the exact same lines as the proof of Proposition 4.5, by combining this last estimate with (5-2), (5-3) and (5-4). This yields the desired result.  $\square$

*Proof of Theorem 2.7.* As before we let  $Y = \exp(C \log_2 x \log_3 x / \log_4 x)$  for some suitably large constant  $C > 0$ , and put  $L_1 = \lfloor \log Y / \log 2 \rfloor$ ,  $L_2 = \lfloor \log Z / \log 2 \rfloor$ ,  $s_{L_1} = Y$ ,  $s_{L_2+1} = Z$ , and  $s_\ell = 2^\ell$  for  $L_1+1 \leq \ell \leq L_2$ . By (4-20) we have

$$\begin{aligned} \max_{\alpha \in [0,1)} \left| \sum_{\substack{1 \leq n \leq Z \\ P^+(n) > y}} \frac{\psi_p(n)h(n)e(\alpha n)}{n} \right| \\ \leq \max_{\alpha \in [0,1)} \left| \sum_{\substack{1 \leq n \leq Y \\ P^+(n) > y}} \frac{\psi_p(n)h(n)e(\alpha n)}{n} \right| + \sum_{L_1 \leq \ell \leq L_2} \max_{\alpha \in [0,1)} \left| \sum_{\substack{s_\ell \leq n \leq s_{\ell+1} \\ P^+(n) > y}} \frac{\psi_p(n)h(n)e(\alpha n)}{n} \right|. \end{aligned}$$

By embedding the set of primes  $3 \leq p \leq x$  into the set of fundamental discriminants  $|d| \leq x$  (since for any such prime,  $p$  or  $-p$  is a fundamental discriminant), it follows from (4-21) that

$$\sum_{L_1 \leq \ell \leq L_2} \max_{\alpha \in [0,1)} \left| \sum_{\substack{s_\ell \leq n \leq s_{\ell+1} \\ P^+(n) > y}} \frac{\psi_p(n)h(n)e(\alpha n)}{n} \right| \ll \frac{\log_4 x}{\log_2 x \log_3 x}, \tag{5-6}$$

for all primes  $p \leq x$  except for a set  $\mathcal{E}_1(x)$  of size

$$|\mathcal{E}_1(x)| \ll x \exp\left(-\frac{\log x \log_4 x}{20 \log_2 x}\right).$$

The result follows along the same exact lines of the proof of Theorem 2.1, by replacing Proposition 4.5 by Proposition 5.2.  $\square$

### 6. The distribution of $L(1, \psi_p)$ : proof of Theorem 1.10

**6A. Proof of the estimate (1-8).** In the range  $2 \leq \tau \leq (\log_2 x)/2 - 2 \log_3 x$ , we shall use [Lamzouri 2017] where we proved an asymptotic formula for complex moments of  $L(1, \psi_p)$  involving a secondary term coming from a possible Landau–Siegel exceptional discriminant. Indeed, it follows from Theorem 1.2 of [Lamzouri 2017] that for all real numbers  $2 \leq k \leq \sqrt{\log x} / (\log_2 x)^2$  we have

$$\frac{2}{\text{Li}(x)} \sum_{\substack{p \leq x \\ p \equiv 3 \pmod{4}}} L(1, \psi_p)^k = \mathbb{E}(L(1, \mathbb{X})^k) + E_1 + O\left(\exp\left(-\frac{\sqrt{\log x}}{10 \log \log x}\right)\right), \tag{6-1}$$

where

$$|E_1| \leq \left| \mathbb{E}(\mathbb{X}(|d_1|)L(1, \mathbb{X})^k) \right|,$$

and where  $d_1$  is the possible “exceptional” discriminant<sup>7</sup> with  $|d_1| \leq \exp(\sqrt{\log x})$ , and

$$L(1, \mathbb{X}) := \prod_{q \text{ prime}} \left( 1 - \frac{\mathbb{X}(q)}{q} \right)^{-1},$$

with the  $\mathbb{X}(q)$  being I.I.D. random variables taking the values  $\pm 1$  with probability  $1/2$ . We cannot rule out that the term  $E_1$  is of the same size as the main term, but we will prove that it will not heavily affect the size of the  $k$ -th moment of  $L(1, \psi_p)$  if  $k$  is large. Indeed, we observe that

$$\frac{\mathbb{E}(\mathbb{X}(|d_1|)L(1, \mathbb{X})^k)}{\mathbb{E}(L(1, \mathbb{X})^k)} = \prod_{q|d_1} \frac{\left(1 - \frac{1}{q}\right)^{-k} - \left(1 + \frac{1}{q}\right)^{-k}}{\left(1 - \frac{1}{q}\right)^{-k} + \left(1 + \frac{1}{q}\right)^{-k}} = \prod_{q|d_1} \frac{1 - \delta_q}{1 + \delta_q}, \tag{6-2}$$

where  $\delta_q = \left(1 - \frac{2}{q+1}\right)^k$ . Let  $\varepsilon > 0$  be a suitably small constant. Let  $q_1$  be the largest prime factor of  $d_1$ . Since  $d_1$  is square-free and  $d_1 > (\log x)^{10/\varepsilon}$ , if  $x$  is large enough, by Siegel’s theorem, we must have  $q_1 > (\log |d_1|)/2 > (5/\varepsilon) \log \log x$ , since otherwise  $|d_1| < \prod_{q \leq (\log |d_1|)/2} q = |d_1|^{1/2+o(1)}$  by the prime number theorem, which is a contradiction. Inserting this estimate in (6-2) gives

$$0 < \frac{\mathbb{E}(\mathbb{X}(|d_1|)L(1, \mathbb{X})^k)}{\mathbb{E}(L(1, \mathbb{X})^k)} \leq 1 - \delta_{q_1} = 1 - \left(1 - \frac{2}{q_1 + 1}\right)^k \leq 1 - \exp\left(-\varepsilon \frac{k}{\log k}\right), \tag{6-3}$$

for all real numbers  $2 \leq k \leq \sqrt{\log x}/(\log_2 x)^2$ . On the other hand, it follows from Proposition 1.2 of [Lamzouri 2010] that

$$\mathbb{E}(L(1, \mathbb{X})^k) = \exp\left(k \log_2 k + k\gamma + \frac{k}{\log k} \left(B_0 - 1 + O\left(\frac{1}{\log k}\right)\right)\right). \tag{6-4}$$

Combining this estimate with (6-1) and (6-3) we obtain

$$\frac{2}{\pi(x)} \sum_{\substack{p \leq x \\ p \equiv 3 \pmod{4}}} L(1, \psi_p)^k = \exp\left(k \log_2 k + k\gamma + \frac{k}{\log k} (B_0 - 1 + O(\varepsilon))\right). \tag{6-5}$$

Finally, the estimate for the distribution function (1-8) follows from the proof of Theorem 0.1 of [Lamzouri 2010] (which holds for general random models of this type), with the choice  $\tau = \log k + B_0$ . This completes the proof of (1-8) in the case  $a = 3$ . We also note that the case  $a = 1$  is similar, since one can derive the same estimate as (6-1) over the primes  $p \equiv 1 \pmod{4}$  using the method of [Lamzouri 2017].

---

<sup>7</sup>By the Landau–Page Theorem (see Chapter 20 of [Davenport 2000]), there is at most one square-free integer  $d_1$  such that  $|d_1| \leq \exp(\sqrt{\log x})$  and  $L(s, \chi_{d_1})$  has a zero in the region  $\text{Re}(s) > 1 - c/\sqrt{\log x}$ , for some positive constant  $c$ . If it exists, we refer to such  $d_1$  as the exceptional discriminant in the range  $|d_1| \leq \exp(\sqrt{\log x})$ .

We now show how to obtain the same estimate for the proportion of primes  $p \equiv a \pmod 4$  such that  $L(1, \psi_p \chi_{-3}) > (2e^\gamma/3)\tau$ . First, by a slight adaptation of the proof of Theorem 1.2 of [Lamzouri 2017] we get

$$\frac{2}{\text{Li}(x)} \sum_{\substack{p \leq x \\ p \equiv a \pmod 4}} L(1, \psi_p \chi_{-3})^k = \mathbb{E}(L(1, \mathbb{X} \chi_{-3})^k) + E_2 + O\left(\exp\left(-\frac{\sqrt{\log x}}{10 \log \log x}\right)\right), \tag{6-6}$$

where

$$|E_2| \leq |\mathbb{E}(\mathbb{X}(|d_1|)L(1, \mathbb{X} \chi_{-3})^k)|,$$

and where  $d_1$  is the exceptional discriminant (if it exists) with  $|d_1| \leq \exp(\sqrt{\log x})$ , and

$$L(1, \mathbb{X} \chi_{-3}) := \prod_{q \text{ prime}} \left(1 - \frac{\mathbb{X}(q) \chi_{-3}(q)}{q}\right)^{-1}.$$

By the independence of the  $\mathbb{X}(q)$ 's we observe that

$$\mathbb{E}(L(1, \mathbb{X} \chi_{-3})^k) = \prod_{q \neq 3} \mathbb{E}\left(\left(1 - \frac{\mathbb{X}(q)}{q}\right)^{-k}\right) = \left(\frac{2}{3}\right)^k \left(\frac{2}{1+2^{-k}}\right) \mathbb{E}(L(1, \mathbb{X})^k).$$

Furthermore, if we fix a suitably small constant  $\varepsilon > 0$ , then a similar argument leading to (6-3) gives

$$|E_2| \leq \left(1 - \exp\left(-\varepsilon \frac{k}{\log k}\right)\right) \mathbb{E}(L(1, \mathbb{X} \chi_{-3})^k), \tag{6-7}$$

if  $x$  is large enough. Thus by (6-4) we deduce that

$$\frac{2}{\pi(x)} \sum_{\substack{p \leq x \\ p \equiv a \pmod 4}} \left(\frac{3}{2} L(1, \psi_p \chi_{-3})\right)^k = \exp\left(k \log_2 k + k\gamma + \frac{k}{\log k} (B_0 - 1 + O(\varepsilon))\right).$$

Finally, the result follows from the proof of Theorem 0.1 of [Lamzouri 2010].

**Remark 6.1.** We observe that assuming GRH, Holmin, Jones, Kurlberg, McLeman and Petersen [Holmin et al. 2019] established the asymptotic formula (6-1) without the term  $E_1$  in the larger range  $k \leq (\log x)/(50(\log \log x)^2)$ . This justifies Remark 1.8.

**6B. Proof of (1-9).** In the range  $(\log_2 x)/2 - 2 \log_3 x \leq \tau \leq \log_2 x - \log_3 x - C_2$ , we need to use a different argument since asymptotic formulas for very large moments of  $L(1, \psi_p)$  are not known, due to the lack of strong unconditional bounds on character sums over primes. In this case, our strategy is to use Theorem 2.7 to truncate  $L(1, \psi_p)$  over  $y$ -friable integers, and then control the behaviour of  $\psi_p(q)$  over the primes  $q \leq y$ . To this end we establish the following lemma which follows from Bombieri's proof [1987, Chapter 6] of Linnik's theorem.

**Lemma 6.2.** *Let  $\{\varepsilon_q\}_q$  prime be a sequence of  $\pm 1$ , and  $a \in \{1, 3\}$ . Let  $x$  be large and  $3 \leq y \leq c_0 \log x$  be a real number, where  $c_0$  is a suitably small constant. Let  $\mathcal{P}(x, y, a, \{\varepsilon_q\})$  be the set of primes  $p \equiv a \pmod 4$  with  $p \leq x$  such that  $\psi_p(q) = \varepsilon_q$  for all primes  $q \leq y$ . Then we have*

$$|\mathcal{P}(x, y, a, \{\varepsilon_q\})| \gg \pi(x) \exp\left(-3y + O\left(\frac{y}{\log y}\right)\right).$$

*Proof.* First recall that if  $p \equiv 3 \pmod 4$  then  $\psi_p(2) = 1$  if  $p \equiv 7 \pmod 8$ , and equals  $-1$  if  $p \equiv 3 \pmod 8$ . Similarly, if  $p \equiv 1 \pmod 4$  then  $\psi_p(2) = 1$  if  $p \equiv 1 \pmod 8$ , and  $\psi_p(2) = -1$  if  $p \equiv 5 \pmod 8$ . Let  $3 \leq q \leq y$  be a prime number. By the law of quadratic reciprocity, there exists a residue class  $b_q \pmod q$  such that if  $p \equiv b_q \pmod q$  then  $\psi_p(q) = \varepsilon_q$ . We now define  $Q := 8 \prod_{3 \leq q \leq y} q$ . Then, by the Chinese remainder theorem there exists a residue class  $b \pmod Q$  such that if  $p \equiv b \pmod Q$  then  $p \equiv a \pmod 4$  and  $\psi_p(q) = \varepsilon_q$  for all primes  $q \leq y$ . By Bombieri’s proof [1987, Chapter 6] of Linnik’s theorem we deduce that there exists a small constant  $c > 0$  such that if  $x > Q^{1/c}$  then

$$|\mathcal{P}(x, y, a, \{\varepsilon_q\})| \geq \pi(x; Q, b) \gg \frac{\pi(x)}{Q^3}.$$

The result follows upon noting that  $Q = \exp(y(1 + O(1/\log y)))$  by the prime number theorem. □

*Proof of the lower bound (1-9).* Let  $Z = x^{21/40}$ . By (2-9) we have

$$L(1, \psi_p) = \sum_{n \leq Z} \frac{\psi_p(n)}{n} + O(1),$$

for all primes  $p \leq x$ . Let  $c_0$  be the constant in Lemma 6.2, and  $3 \leq y \leq c_0 \log x$  be a parameter to be chosen. Define  $\mathcal{P}(y)$  to be the set of primes  $p \leq x$  such that  $p \equiv a \pmod 4$ ,  $\psi_p(q) = 1$  for all primes  $q \leq y$  and

$$\left| \sum_{\substack{1 \leq n \leq Z \\ P^+(n) > y_1}} \frac{\psi_p(n)}{n} \right| \leq e^\gamma, \tag{6-8}$$

where  $y_1 := 10y \log y$ . Combining Lemma 6.2 with Theorem 2.7 yields

$$|\mathcal{P}(y)| \gg \pi(x) \exp(-4y). \tag{6-9}$$

On the other hand, by (2-10) and (6-8) we deduce that for every prime  $p \in \mathcal{P}(y)$  we have

$$\begin{aligned} L(1, \psi_p) &= \sum_{\substack{1 \leq n \leq Z \\ P^+(n) \leq y_1}} \frac{\psi_p(n)}{n} + O(1) = \sum_{\substack{n \geq 1 \\ P^+(n) \leq y_1}} \frac{\psi_p(n)}{n} + O(1) \\ &= \prod_{q \leq y} \left(1 - \frac{1}{q}\right)^{-1} \exp\left(\sum_{y < q \leq y_1} \frac{\psi_p(q)}{q} + O\left(\frac{1}{y}\right)\right) + O(1) \\ &\geq e^\gamma \log y \cdot \exp\left(-\sum_{y < q \leq y_1} \frac{1}{q}\right) \left(1 + O\left(\frac{1}{\log y}\right)\right) + O(1) \\ &\geq e^\gamma \log y - e^\gamma \log \log y + O(1), \end{aligned}$$

by Mertens' theorem. Combining the above estimates and choosing  $y = C\tau e^\tau$  for some large constant  $C$ , implies that  $L(1, \psi_p) > e^\gamma \tau$  whenever  $p \in \mathcal{P}(y)$ . Appealing to (6-9) completes the proof.  $\square$

We end this section by proving the upper bound of (1-9) which we deduce from Theorem 2.7.

*Proof of the upper bound of (1-9).* Let  $Z = x^{21/40}$  and  $\delta = 1/100$ . Let  $y = e^{\tau-B}$ , where  $B$  is a parameter to be chosen later. First by (2-9) we observe that

$$\begin{aligned} L(1, \psi_p) &= \sum_{n \leq Z} \frac{\psi_p(n)}{n} + O(x^{-\delta}) \leq \sum_{\substack{n \geq 1 \\ P^+(n) \leq y}} \frac{1}{n} + \left| \sum_{\substack{1 \leq n \leq Z \\ P^+(n) > y}} \frac{\psi_p(n)}{n} \right| + O(x^{-\delta}) \\ &\leq e^\gamma \tau - e^\gamma B + \left| \sum_{\substack{1 \leq n \leq Z \\ P^+(n) > y}} \frac{\psi_p(n)}{n} \right| + O\left(\frac{1}{\tau}\right). \end{aligned} \tag{6-10}$$

since  $\prod_{p \leq y} (1 - 1/p)^{-1} = e^\gamma \log y + O(1/\log y)$  by the prime number theorem. Therefore, there exists a positive constant  $c$  such that the proportion of primes  $p \leq x$  with  $p \equiv a \pmod 4$  and such that  $L(1, \psi_p) \geq e^\gamma \tau$  is bounded by the proportion of primes  $p \leq x$  such that

$$\left| \sum_{\substack{1 \leq n \leq Z \\ P^+(n) > y}} \frac{\psi_p(n)}{n} \right| \geq e^\gamma \left( B - \frac{c}{\tau} \right).$$

Thus, appealing to Theorem 2.7 with  $B = 2 + c/\tau$  completes the proof of (1-9) for  $L(1, \psi_p)$ . Finally, the analogous result for  $L(1, \psi_p \chi_{-3})$  follows along the same lines upon taking  $h(n) = \chi_{-3}(n)$  in Theorem 2.7 and noting that

$$\left| \sum_{\substack{1 \leq n \leq Z \\ P^+(n) \leq y}} \frac{\psi_p(n) \chi_{-3}(n)}{n} \right| \leq \prod_{\substack{q \neq 3 \\ q \leq y}} \left( 1 - \frac{1}{q} \right)^{-1} = \frac{2e^\gamma}{3} \log y + O\left(\frac{1}{\log y}\right),$$

by the prime number theorem.  $\square$

### 7. Positivity of sums of the Legendre symbol: Proof of Theorems 1.12 and 1.13

Let  $Z = x^{21/40}$  and  $\delta = 1/100$ . Let  $p \leq x$  be a prime number such that  $p \equiv 3 \pmod 4$ . Since  $\psi_p$  is odd and  $\tau(\psi_p) = i\sqrt{p}$ , then by (2-1) we have for any  $\alpha \in (0, 1)$

$$\sum_{n \leq \alpha p} \psi_p(n) = \frac{\sqrt{p}}{\pi} \sum_{1 \leq n \leq Z} \frac{\psi_p(n)(1 - \cos(2\pi n\alpha))}{n} + O(p^{1/2-1/40} \log p). \tag{7-1}$$

We shall now focus on the sum

$$F(\alpha, p) := \sum_{1 \leq n \leq Z} \frac{\psi_p(n)(1 - \cos(2\pi n\alpha))}{n}.$$

Our strategy for proving Theorems 1.12 and 1.13 consists in using Theorem 2.7 to bound the part of this sum over nonfriable integers uniformly over  $\alpha \in [0, 1)$ , for all primes  $p \leq x$  except for a small set of exceptions, and then prescribe the signs of  $\psi_p(q)$  for the small primes  $q$ .

*Proof of Theorem 1.12.* We start by proving (1-10). Let  $Z = x^{21/40}$  and  $2 \leq y \leq \log x$  be a parameter to be chosen. Let  $\mathcal{E}_1(x)$  be the set of primes  $\sqrt{x} \leq p \leq x$  such that  $p \equiv 3 \pmod 4$  and

$$\max_{\alpha \in [0,1)} \left| \sum_{\substack{1 \leq n \leq Z \\ P^+(n) > y}} \frac{\psi_p(n)(1 - \cos(2\pi n\alpha))}{n} \right| > 2e^y.$$

Then it follows from Theorem 2.7 that

$$|\mathcal{E}_1(x)| \ll \pi(x) \exp\left(-\frac{y}{3 \log y}\right). \tag{7-2}$$

We now define  $\mathcal{A}(y)$  to be the set of primes  $\sqrt{x} \leq p \leq x$  such that  $p \equiv 3 \pmod 4$  and  $\psi_p(q) = 1$  for all primes  $q \leq y_0$ , where  $y_0 := y/(20 \log y)$ . Moreover, we put  $\mathcal{D}(y) := \mathcal{A}(y) \setminus \mathcal{E}_1(x)$ . By Lemma 6.2 and the estimate (7-2) we have  $|\mathcal{D}(y)| \gg \pi(x) \exp(-y/(4 \log y))$ . Moreover, if  $p \in \mathcal{D}(y)$  then it follows from (2-10) that for all  $\alpha \in (0, 1)$  we have

$$\begin{aligned} F(\alpha, p) &= \sum_{\substack{1 \leq n \leq Z \\ P^+(n) \leq y}} \frac{\psi_p(n)(1 - \cos(2\pi n\alpha))}{n} + O(1) = \sum_{\substack{n \geq 1 \\ P^+(n) \leq y}} \frac{\psi_p(n)(1 - \cos(2\pi n\alpha))}{n} + O(1) \\ &= \sum_{\substack{n \geq 1 \\ P^+(n) \leq y_0}} \frac{1 - \cos(2\pi n\alpha)}{n} + O(\log_2 y) = \sum_{\substack{n \geq 1 \\ P^+(n) \leq y}} \frac{1 - \cos(2\pi n\alpha)}{n} + O(\log_2 y), \end{aligned} \tag{7-3}$$

since

$$\sum_{\substack{n \geq 1 \\ y_0 < P^+(n) \leq y}} \frac{1}{n} = \prod_{p \leq y} \left(1 - \frac{1}{p}\right)^{-1} - \prod_{p \leq y_0} \left(1 - \frac{1}{p}\right)^{-1} \ll \log_2 y,$$

by Mertens' theorem. Furthermore, by Lemma 3.4 of [Bober et al. 2018] we have

$$\left| \sum_{\substack{n \geq 1 \\ P^+(n) \leq y}} \frac{\cos(2\pi n\alpha)}{n} \right| \leq \left| \sum_{\substack{n \geq 1 \\ P^+(n) \leq y}} \frac{e(n\alpha)}{n} \right| = \sum_{\substack{n \leq 1/\|\alpha\| \\ P^+(n) \leq y}} \frac{1}{n} + O(1), \tag{7-4}$$

where  $\|\cdot\|$  denotes the distance to the nearest integer. We now write  $\|\alpha\| = y^{-u}$  for some  $u > 0$ . By Lemma 3.3 of [Bober et al. 2018] we get

$$\sum_{\substack{n \leq 1/\|\alpha\| \\ P^+(n) \leq y}} \frac{1}{n} = \sum_{\substack{n \leq y^u \\ P^+(n) \leq y}} \frac{1}{n} = (\log y) \int_0^u \rho(t) dt + O(1),$$

where  $\rho$  is the Dickman–de Bruijn function, defined by  $\rho(t)$  for  $0 \leq t \leq 1$ , and  $t\rho'(t) = -\rho(t - 1)$  for  $t > 1$ . Combining this estimate with (7-3) and (7-4) we obtain

$$F(\alpha, p) \geq \log y \left( e^\gamma - \int_0^u \rho(t) dt \right) + O(\log_2 y), \tag{7-5}$$

for all  $p \in \mathcal{D}(y)$  and all  $\alpha \in (0, 1)$ , where  $u = -\log\|\alpha\|/\log y$ . Note that  $\int_0^u \rho(t) dt$  is increasing in  $u$  and that  $\int_0^\infty \rho(t) dt = e^\gamma$ . Moreover, one has the estimate

$$e^\gamma - \int_0^u \rho(t) dt = \int_u^\infty \rho(t) dt = \frac{1}{u^{u(1+o(1))}}, \tag{7-6}$$

which follows from the standard estimate  $\rho(t) = t^{-t(1+o(1))}$ . Let  $C$  be a suitably large constant, and  $u_0$  be the solution of the equation

$$\int_{u_0}^\infty \rho(t) dt = \frac{C \log_2 y}{\log y}. \tag{7-7}$$

Then, it follows from (7-5) that for all  $p \in \mathcal{D}(y)$  and  $\|\alpha\| > y^{-u_0}$  we have  $F(\alpha, p) \geq 10$  if  $C$  is suitably large, and hence

$$\sum_{n \leq \alpha p} \psi_p(n) > \sqrt{p}$$

by (7-1) if  $x$  is large enough. To finish the proof, we choose  $y$  such that  $T = y^{u_0}/2$ , which implies that  $\lambda(p) > 1 - 1/T$  for all  $p \in \mathcal{D}(y)$ . Moreover, it follows from the estimates (7-6) and (7-7) that  $u_0 = (1 + o(1)) \log_2 y / \log_3 y$ , and hence we get

$$y = \exp\left(\frac{(1 + o(1)) \log T \log_3 T}{\log_2 T}\right),$$

as desired. □

In order to prove (1-11) for small  $T$ , we require a more precise estimate than the one provided by Lemma 6.2. To this end we prove the following result, which gives an asymptotic formula for the cardinality of the set  $\mathcal{P}(x, y, a, \{\varepsilon_q\})$  in Lemma 6.2, in the smaller range  $3 \leq y \leq \log \log x$ .

**Lemma 7.1.** *Let  $\{\varepsilon_q\}_q$  prime be a sequence of  $\pm 1$ , and  $a \in \{1, 3\}$ . Let  $x$  be large and  $3 \leq y \leq \log \log x$  be a real number. Let  $\mathcal{P}(x, y, a, \{\varepsilon_q\})$  be the set in Lemma 6.2. Then we have*

$$|\mathcal{P}(x, y, a, \{\varepsilon_q\})| = \frac{\pi(x)}{2^{\pi(y)+1}} + O(xe^{-c\sqrt{\log x}}),$$

for some positive constant  $c$ .

*Proof.* Let  $Q = \prod_{q \leq y} q$ . Note that by the prime number theorem and our assumption we have  $Q = e^{y+o(y)} \leq (\log x)^{1+o(1)}$ . We observe that for a prime  $y < p \leq x$  we have

$$\frac{1}{2^{\pi(y)+1}} (1 + \chi_{-4}(a)\chi_{-4}(p)) \prod_{q \leq y} (1 + \varepsilon_q \psi_p(q)) = \begin{cases} 1 & \text{if } p \in \mathcal{P}(x, y, a, \{\varepsilon_q\}), \\ 0 & \text{otherwise,} \end{cases} \tag{7-8}$$

where  $\chi_{-4}$  is the nonprincipal character modulo 4. We extend the sequence  $\varepsilon_q$  multiplicatively to all square-free numbers  $\ell$  by letting  $\varepsilon_\ell = \prod_{q|\ell} \varepsilon_q$ . Therefore, expanding the product on the left-hand side of (7-8) we deduce that

$$\begin{aligned} |\mathcal{P}(x, y, a, \{\varepsilon_q\})| &= \frac{1}{2^{\pi(y)+1}} \sum_{y < p \leq x} (1 + \chi_{-4}(a)\chi_{-4}(p)) \prod_{q \leq y} (1 + \varepsilon_q \psi_p(q)) + O(y) \\ &= \frac{1}{2^{\pi(y)+1}} \sum_{p \leq x} (1 + \chi_{-4}(a)\chi_{-4}(p)) \sum_{\ell|Q} \varepsilon_\ell \psi_p(\ell) + O(y) \\ &= \frac{1}{2^{\pi(y)+1}} \left( \sum_{\ell|Q} \varepsilon_\ell \sum_{p \leq x} \left(\frac{\ell}{p}\right) + \chi_{-4}(a) \sum_{\ell|Q} \varepsilon_\ell \sum_{p \leq x} \chi_{-4}(p) \left(\frac{\ell}{p}\right) \right) + O(y). \end{aligned} \tag{7-9}$$

If  $\ell \neq 1$ , then the law of quadratic reciprocity implies that  $\xi_1 = \left(\frac{\ell}{\cdot}\right)$  is a nonprincipal character of conductor  $\ell$  or  $4\ell$ . Similarly, for all  $\ell$  the character  $\xi_2 = \chi_{-4}\xi_1$  is a nonprincipal character of conductor  $4\ell$ . Furthermore, note that  $4\ell \leq 4Q \leq (\log x)^2$  if  $x$  is large enough. Thus, it follows from Corollary 11.18 of [Montgomery and Vaughan 2007] that for  $j = 1$  if  $\ell \neq 1$  and  $\ell | Q$ , and for  $j = 2$  for all  $\ell | Q$  we have

$$\sum_{p \leq x} \xi_j(p) \ll x \exp(-c\sqrt{\log x}). \tag{7-10}$$

Inserting these bounds in (7-9) completes the proof. □

*Proof of Theorem 1.13.* We start by proving (1-11). Let  $Z = x^{21/40}$  and  $\delta = 1/100$ . By (7-1) we have

$$\sum_{n \leq \alpha p} \psi_p(n) = \frac{\sqrt{p}}{\pi} F(\alpha, p) + O(p^{1/2-\delta}),$$

for all primes  $\sqrt{x} \leq p \leq x$  and all  $\alpha \in (0, 1)$ . Let  $3 \leq y \leq (\log_2 x)(\log_3 x)^2$  be a parameter to be chosen, and  $\mathcal{E}_2(x)$  be the set of primes  $\sqrt{x} \leq p \leq x$  such that  $p \equiv 3 \pmod{4}$  and

$$\max_{\alpha \in [0,1)} \left| \sum_{\substack{1 \leq n \leq Z \\ P^+(n) > y}} \frac{\psi_p(n)(1 - \cos(2\pi n\alpha))}{n} \right| > \frac{2e^\gamma}{\log y}.$$

By Theorem 2.7 we have

$$|\mathcal{E}_2(x)| \ll \pi(x) \exp\left(-\frac{y}{3(\log y)^3}\right). \tag{7-11}$$

We will use the same choice as Montgomery [1976] for the values of  $\psi_p(q)$  for small  $q$ . Let  $h(n)$  be the completely multiplicative function such that  $h(q) = \chi_{-3}(q)$  for all primes  $q \neq 3$  and  $h(3) = -1$ . Montgomery [1976] showed that for all  $\alpha \in (0, 1)$  such that  $\|\alpha\| < 1/3$  we have

$$U(\alpha) := \sum_{n=1}^{\infty} \frac{h(n) \cos(2\pi n\alpha)}{n} > \frac{\pi}{8\sqrt{3}}. \tag{7-12}$$

Let  $y_0 := y/(3 \log y)^2$  and  $3 \leq H \leq y_0$  be a parameter to be chosen. Let  $\mathcal{B}(y, H)$  be the set of primes  $\sqrt{x} \leq p \leq x$  such that  $p \equiv 3 \pmod{4}$  and

$$\psi_p(q) = \begin{cases} h(q) & \text{if } q \leq H, \\ -1 & \text{if } H < q \leq y_0. \end{cases}$$

We define  $\mathcal{T}(y) := \mathcal{B}(y, H) \setminus \mathcal{E}_2(x)$ . Then we have  $|\mathcal{T}(y)| \gg \pi(x)/2^{\pi(y_0)}$  by Lemma 7.1 and the estimate (7-11). We now let  $p$  be a prime in  $\mathcal{T}(y)$ . By (2-10) and our assumption on  $p$  we get

$$\begin{aligned} F(\alpha, p) &= \sum_{\substack{1 \leq n \leq Z \\ P^+(n) \leq y}} \frac{\psi_p(n)(1 - \cos(2\pi n\alpha))}{n} + O\left(\frac{1}{\log y}\right) \\ &= \sum_{\substack{n \geq 1 \\ P^+(n) \leq y}} \frac{\psi_p(n)(1 - \cos(2\pi n\alpha))}{n} + O\left(\frac{1}{\log y}\right). \end{aligned} \tag{7-13}$$

Furthermore, we have

$$\begin{aligned} \sum_{\substack{n \geq 1 \\ P^+(n) \leq y}} \frac{\psi_p(n)}{n} &= \prod_{q \leq y} \left(1 - \frac{\psi_p(q)}{q}\right)^{-1} \\ &= \frac{3}{4} \prod_{q \leq H} \left(1 - \frac{\chi_{-3}(q)}{q}\right)^{-1} \prod_{H < q \leq y_0} \left(1 + \frac{1}{q}\right)^{-1} \prod_{y_0 < q \leq y} \left(1 - \frac{\psi_p(q)}{q}\right)^{-1}. \end{aligned} \tag{7-14}$$

By the prime number theorem in arithmetic progressions we have

$$\sum_{q \leq t} \chi_{-3}(q) \ll t \exp(-c\sqrt{\log t}),$$

for some constant  $c > 0$ . Therefore, by partial summation we obtain

$$\sum_{q > H} \frac{\chi_{-3}(q)}{q} \ll \exp\left(-\frac{c}{2}\sqrt{\log H}\right).$$

This implies that

$$\begin{aligned} \prod_{q \leq H} \left(1 - \frac{\chi_{-3}(q)}{q}\right)^{-1} &= L(1, \chi_{-3}) + O\left(\exp\left(-\frac{c}{2}\sqrt{\log H}\right)\right) \\ &= \frac{\pi}{3\sqrt{3}} + O\left(\exp\left(-\frac{c}{2}\sqrt{\log H}\right)\right). \end{aligned}$$

Inserting this estimate in (7-14) and using Mertens' theorem we deduce that

$$0 < \sum_{\substack{n \geq 1 \\ P^+(n) \leq y}} \frac{\psi_p(n)}{n} \leq \frac{\pi}{4\sqrt{3}} \frac{\log H \log y}{(\log y_0)^2} \left(1 + O\left(\frac{1}{\log H}\right)\right).$$

We now choose  $H = \sqrt{y}/(C_1(\log y)^2)$  where  $C_1$  is a suitably large constant. This gives

$$\sum_{\substack{n \geq 1 \\ P^+(n) \leq y}} \frac{\psi_p(n)}{n} \leq \frac{\pi}{8\sqrt{3}} - \frac{\log C_1}{\log y}. \tag{7-15}$$

On the other hand, by Parseval's identity we have

$$\begin{aligned} \int_0^1 \left| \sum_{\substack{n \geq 1 \\ P^+(n) \leq y}} \frac{\psi_p(n) \cos(2\pi n\alpha)}{n} - U(\alpha) \right|^2 d\alpha &= \sum_{\substack{n \geq 1 \\ P^+(n) \leq y}} \frac{(\psi_p(n) - h(n))^2}{n^2} + \sum_{\substack{n \geq 1 \\ P^+(n) > y}} \frac{h(n)^2}{n^2} \\ &\ll \sum_{\substack{n \geq 1 \\ P^+(n) > H}} \frac{1}{n^2} \ll \frac{\log y}{\sqrt{y}}, \end{aligned}$$

since

$$\sum_{\substack{n \geq 1 \\ P^+(n) > H}} \frac{1}{n^2} = \frac{\pi^2}{6} - \sum_{\substack{n \geq 1 \\ P^+(n) \leq H}} \frac{1}{n^2} = \frac{\pi^2}{6} \left( 1 - \prod_{q > H} \left( 1 - \frac{1}{q^2} \right) \right) \ll \sum_{q > H} \frac{1}{q^2} \ll \frac{1}{H \log H}.$$

Let  $S_p$  be the set of  $\alpha \in (0, 1)$  such that

$$\left| \sum_{\substack{n \geq 1 \\ P^+(n) \leq y}} \frac{\psi_p(n) \cos(2\pi n\alpha)}{n} - U(\alpha) \right| > \frac{1}{\log y}.$$

Then

$$\mu(S_p) \leq (\log y)^2 \int_0^1 \left| \sum_{\substack{n \geq 1 \\ P^+(n) \leq y}} \frac{\psi_p(n) \cos(2\pi n\alpha)}{n} - U(\alpha) \right|^2 d\alpha \ll \frac{(\log y)^3}{\sqrt{y}}.$$

Thus, recalling the definitions of the sets  $\mathcal{T}(y)$  and  $S_p$  and using the estimate (7-12) we deduce that if  $p \in \mathcal{T}(y)$  and  $\alpha \in (0, 1/3) \cup (2/3, 1) \setminus S_p$  then

$$\sum_{\substack{n \geq 1 \\ P^+(n) \leq y}} \frac{\psi_p(n) \cos(2\pi n\alpha)}{n} > \frac{\pi}{8\sqrt{3}} - \frac{1}{\log y}.$$

Combining this estimate with (7-13) and (7-15) gives

$$F(\alpha, p) = \sum_{\substack{n \geq 1 \\ P^+(n) \leq y}} \frac{\psi_p(n)}{n} - \sum_{\substack{n \geq 1 \\ P^+(n) \leq y}} \frac{\psi_p(n) \cos(2\pi n\alpha)}{n} + O\left(\frac{1}{\log y}\right) \leq -\frac{C_2}{\log y},$$

for some positive constant  $C_2$  if  $C_1$  is suitably large. This in turn implies that

$$\sum_{n \leq \alpha p} \psi_p(n) < -\frac{\sqrt{p}}{\log_3 p}$$

by (7-1), if  $x$  is large enough. Hence for  $p \in \mathcal{T}(y)$  we have

$$\lambda(p) \leq \frac{1}{3} + \mu(\mathcal{S}_p) \leq \frac{1}{3} + C_3 \frac{(\log y)^3}{\sqrt{y}},$$

for some absolute constant  $C_3 > 0$ . Choosing  $y = C_4 T^2 (\log T)^6$  for some suitably large constant  $C_4$  completes the proof of (1-11).

To prove (1-12) we follow the exact same lines, and replace Lemma 7.1 by Lemma 6.2. In this case we make the following choices of the parameters:  $y_0 = y/(3 \log y)^3 \leq c_0 \log x$  (where  $c_0$  is the constant in Lemma 6.2),  $H = \sqrt{y}/(C_5 (\log y)^3)$ , and  $y = C_6 T^2 (\log T)^8$  for some suitably large constants  $C_5$  and  $C_6$ . This completes the proof.  $\square$

### Acknowledgments

The author would like to thank Andrew Granville for useful discussions. We would also like to thank the referee for carefully reading the paper and for comments and suggestions. This work was completed while the author was on a Délégation CNRS at the IRL3457 CRM-CNRS in Montréal. We would like to thank the CNRS for its support and the Centre de Recherches Mathématiques for its excellent working conditions.

### References

- [Baker and Montgomery 1990] R. C. Baker and H. L. Montgomery, “Oscillations of quadratic  $L$ -functions”, pp. 23–40 in *Analytic number theory* (Allerton Park, IL, 1989), edited by B. C. Berndt et al., Progr. Math. **85**, Birkhäuser, Boston, MA, 1990. MR Zbl
- [Bateman and Chowla 1950] P. T. Bateman and S. Chowla, “Averages of character sums”, *Proc. Amer. Math. Soc.* **1** (1950), 781–787. MR Zbl
- [Bober 2014] J. Bober, “Averages of character sums”, preprint, 2014. arXiv 1409.1840
- [Bober and Goldmakher 2013] J. W. Bober and L. Goldmakher, “The distribution of the maximum of character sums”, *Mathematika* **59**:2 (2013), 427–442. MR Zbl
- [Bober and Goldmakher 2016] J. W. Bober and L. Goldmakher, “Pólya–Vinogradov and the least quadratic nonresidue”, *Math. Ann.* **366**:1-2 (2016), 853–863. MR Zbl
- [Bober et al. 2018] J. Bober, L. Goldmakher, A. Granville, and D. Koukoulopoulos, “The frequency and the structure of large character sums”, *J. Eur. Math. Soc.* **20**:7 (2018), 1759–1818. MR Zbl
- [Bombieri 1987] E. Bombieri, *Le grand crible dans la théorie analytique des nombres*, Astérisque **18**, Soc. Math. France, Paris, 1987. MR Zbl
- [Davenport 2000] H. Davenport, *Multiplicative number theory*, 3rd ed., Grad. Texts in Math. **74**, Springer, 2000. MR Zbl
- [Elliott 1970] P. D. T. A. Elliott, “On the mean value of  $f(p)$ ”, *Proc. Lond. Math. Soc.* (3) **21**:1 (1970), 28–96. MR Zbl
- [Goldmakher 2012] L. Goldmakher, “Multiplicative mimicry and improvements to the Pólya–Vinogradov inequality”, *Algebra Number Theory* **6**:1 (2012), 123–163. MR Zbl
- [Granville and Mangerel 2023] A. Granville and A. P. Mangerel, “Three conjectures about character sums”, *Math. Z.* **305**:3 (2023), art. id. 49. MR Zbl
- [Granville and Soundararajan 2003] A. Granville and K. Soundararajan, “The distribution of values of  $L(1, \chi_d)$ ”, *Geom. Funct. Anal.* **13**:5 (2003), 992–1028. MR Zbl
- [Granville and Soundararajan 2006] A. Granville and K. Soundararajan, “Extreme values of  $|\zeta(1+it)|$ ”, pp. 65–80 in *The Riemann zeta function and related themes* (Bangalore, 2003), edited by R. Balasubramanian and K. Srinivas, Ramanujan Math. Soc. Lect. Notes Ser. **2**, Ramanujan Math. Soc., Mysuru, India, 2006. MR Zbl

- [Granville and Soundararajan 2007] A. Granville and K. Soundararajan, “Large character sums: pretentious characters and the Pólya–Vinogradov theorem”, *J. Amer. Math. Soc.* **20**:2 (2007), 357–384. MR Zbl
- [Heath-Brown 1995] D. R. Heath-Brown, “A mean value estimate for real character sums”, *Acta Arith.* **72**:3 (1995), 235–275. MR Zbl
- [Holmin et al. 2019] S. Holmin, N. Jones, P. Kurlberg, C. McLeman, and K. Petersen, “Missing class groups and class number statistics for imaginary quadratic fields”, *Exp. Math.* **28**:2 (2019), 233–254. MR Zbl
- [Hussain 2022a] A. Hussain, *Character studies: investigating the limiting distribution of character sums*, Ph.D. thesis, University of Bristol, 2022, available at <https://research-information.bris.ac.uk/en/studentTheses/character-studies>.
- [Hussain 2022b] A. Hussain, “The limiting distribution of character sums”, *Int. Math. Res. Not.* **2022**:20 (2022), 16292–16326. MR Zbl
- [Hussain and Lamzouri 2023] A. Hussain and Y. Lamzouri, “The limiting distribution of Legendre paths”, 2023. To appear in *J. Éc. Polytech. Math.* arXiv 2304.13025
- [Joshi 1970] P. T. Joshi, “The size of  $L(1, \chi)$  for real nonprincipal residue characters  $\chi$  with prime modulus”, *J. Number Theory* **2**:1 (1970), 58–73. MR Zbl
- [Lamzouri 2010] Y. Lamzouri, “Distribution of values of  $L$ -functions at the edge of the critical strip”, *Proc. Lond. Math. Soc.* (3) **100**:3 (2010), 835–863. MR Zbl
- [Lamzouri 2011] Y. Lamzouri, “Extreme values of  $\arg L(1, \chi)$ ”, *Acta Arith.* **146**:4 (2011), 335–354. MR Zbl
- [Lamzouri 2017] Y. Lamzouri, “The number of imaginary quadratic fields with prime discriminant and class number up to  $H$ ”, *Q. J. Math.* **68**:4 (2017), 1379–1393. MR Zbl
- [Lamzouri and Mangerel 2022] Y. Lamzouri and A. P. Mangerel, “Large odd order character sums and improvements of the Pólya–Vinogradov inequality”, *Trans. Amer. Math. Soc.* **375**:6 (2022), 3759–3793. MR Zbl
- [Littlewood 1928] J. E. Littlewood, “On the class number of the corpus  $P(\sqrt{-k})$ ”, *Proc. Lond. Math. Soc.* (2) **27**:5 (1928), 358–372. MR Zbl
- [Mangerel 2020] A. P. Mangerel, “Short character sums and the Pólya–Vinogradov inequality”, *Q. J. Math.* **71**:4 (2020), 1281–1308. MR Zbl
- [Mehkari 2005] S. Mehkari, *Distribution of sums of the Legendre symbol*, master’s thesis, Université de Montréal, 2005, available at <https://hdl.handle.net/1866/17278>.
- [Montgomery 1971] H. L. Montgomery, *Topics in multiplicative number theory*, Lecture Notes in Math. **227**, Springer, 1971. MR Zbl
- [Montgomery 1976] H. L. Montgomery, “Distribution questions concerning a character sum”, pp. 195–203 in *Topics in number theory* (Debrecen, Hungary, 1974), edited by P. Turán, Colloq. Math. Soc. János Bolyai **13**, North-Holland, Amsterdam, 1976. MR Zbl
- [Montgomery and Vaughan 1977] H. L. Montgomery and R. C. Vaughan, “Exponential sums with multiplicative coefficients”, *Invent. Math.* **43**:1 (1977), 69–82. MR Zbl
- [Montgomery and Vaughan 1979] H. L. Montgomery and R. C. Vaughan, “Mean values of character sums”, *Canadian J. Math.* **31**:3 (1979), 476–487. MR Zbl
- [Montgomery and Vaughan 1999] H. L. Montgomery and R. C. Vaughan, “Extreme values of Dirichlet  $L$ -functions at 1”, pp. 1039–1052 in *Number theory in progress, II* (Zakopane/Kościełisko, Poland, 1997), edited by K. Györy et al., de Gruyter, Berlin, 1999. MR Zbl
- [Montgomery and Vaughan 2007] H. L. Montgomery and R. C. Vaughan, *Multiplicative number theory, I: Classical theory*, Cambridge Stud. Adv. Math. **97**, Cambridge Univ. Press, 2007. MR Zbl
- [Paley 1932] R. E. A. C. Paley, “A theorem on characters”, *J. Lond. Math. Soc.* **7**:1 (1932), 28–32. MR Zbl

Communicated by Roger Heath-Brown

Received 2023-03-02    Revised 2023-09-12    Accepted 2023-11-27

youness.lamzouri@univ-lorraine.fr

*Institut Élie Cartan de Lorraine, Université de Lorraine, CNRS, Nancy, France*  
*IRL3457 CRM-CNRS, Centre de Recherches Mathématiques,*  
*Université de Montréal, Montréal, QC, Canada*



# Algebra & Number Theory

msp.org/ant

## EDITORS

MANAGING EDITOR  
Antoine Chambert-Loir  
Université Paris-Diderot  
France

EDITORIAL BOARD CHAIR  
David Eisenbud  
University of California  
Berkeley, USA

## BOARD OF EDITORS

Jason P. Bell	University of Waterloo, Canada	Philippe Michel	École Polytechnique Fédérale de Lausanne
Bhargav Bhatt	University of Michigan, USA	Martin Olsson	University of California, Berkeley, USA
Frank Calegari	University of Chicago, USA	Irena Peeva	Cornell University, USA
J.-L. Colliot-Thélène	CNRS, Université Paris-Saclay, France	Jonathan Pila	University of Oxford, UK
Brian D. Conrad	Stanford University, USA	Anand Pillay	University of Notre Dame, USA
Samit Dasgupta	Duke University, USA	Bjorn Poonen	Massachusetts Institute of Technology, USA
Hélène Esnault	Freie Universität Berlin, Germany	Victor Reiner	University of Minnesota, USA
Gavril Farkas	Humboldt Universität zu Berlin, Germany	Peter Sarnak	Princeton University, USA
Sergey Fomin	University of Michigan, USA	Michael Singer	North Carolina State University, USA
Edward Frenkel	University of California, Berkeley, USA	Vasudevan Srinivas	SUNY Buffalo, USA
Wee Teck Gan	National University of Singapore	Shunsuke Takagi	University of Tokyo, Japan
Andrew Granville	Université de Montréal, Canada	Pham Huu Tiep	Rutgers University, USA
Ben J. Green	University of Oxford, UK	Ravi Vakil	Stanford University, USA
Christopher Hacon	University of Utah, USA	Akshay Venkatesh	Institute for Advanced Study, USA
Roger Heath-Brown	Oxford University, UK	Melanie Matchett Wood	Harvard University, USA
János Kollár	Princeton University, USA	Shou-Wu Zhang	Princeton University, USA
Michael J. Larsen	Indiana University Bloomington, USA		

## PRODUCTION

production@msp.org  
Silvio Levy, Scientific Editor

---

See inside back cover or [msp.org/ant](http://msp.org/ant) for submission instructions.

---

The subscription price for 2024 is US \$525/year for the electronic version, and \$770/year (+\$65, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to MSP.

---

Algebra & Number Theory (ISSN 1944-7833 electronic, 1937-0652 printed) at Mathematical Sciences Publishers, 798 Evans Hall #3840, c/o University of California, Berkeley, CA 94720-3840 is published continuously online.

---

ANT peer review and production are managed by EditFLOW<sup>®</sup> from MSP.

PUBLISHED BY

 **mathematical sciences publishers**  
nonprofit scientific publishing

<http://msp.org/>

© 2024 Mathematical Sciences Publishers

# Algebra & Number Theory

Volume 18    No. 11    2024

---

Galois orbits of torsion points near atoral sets VESSELIN DIMITROV and PHILIPP HABEGGER	1945
Rooted tree maps for multiple $L$ -values from a perspective of harmonic algebras HIDEKI MURAHARA, TATSUSHI TANAKA and NORIKO WAKABAYASHI	2003
Terminal orders on arithmetic surfaces DANIEL CHAN and COLIN INGALLS	2027
Word measures on $GL_n(q)$ and free group algebras DANIELLE ERNST-WEST, DORON PUDER and MATAN SEIDEL	2047
The distribution of large quadratic character sums and applications YOUNESS LAMZOURI	2091