

# *Algebra & Number Theory*

Volume 19  
2025  
No. 12

**Reduction modulo  $p$  of Noether's problem**

Emiliano Ambrosi and Domenico Valloni





# Reduction modulo $p$ of Noether's problem

Emiliano Ambrosi and Domenico Valloni

Let  $R$  be a complete valuation ring of mixed characteristic  $(0, p)$  with algebraically closed fraction field  $K$  and residue field  $k$ . Let  $X/R$  be a smooth projective morphism. We show that if  $X_k$  is stably rational, then  $H_{\text{ét}}^3(X_K, \hat{\mathbb{Z}})$  is torsion-free. The proof uses the integral  $p$ -adic Hodge theory of Bhatt, Morrow and Scholze and the study of differential forms in positive characteristic. We then apply this result to study the Noether problem for finite  $p$ -groups.

1. Introduction	2359
2. Proof of Theorem 1.1.5	2362
3. A stably irrational variety reducing to a rational variety	2365
Acknowledgements	2366
References	2366

## 1. Introduction

Many techniques to study the birational geometry of varieties are based on specialization methods in equicharacteristic or in mixed characteristic (see [Voi15; CTP16], for instance), often exploiting pathological phenomena in positive characteristic [Kol95; Tot16; ABBGvB21].

In this paper we study birational invariants in mixed characteristic. Let us fix a complete valuation ring  $R$  of mixed characteristic  $(0, p)$  with algebraically closed fraction field  $K$  and residue field  $k$  (necessarily algebraically closed) and a smooth proper morphism  $X/R$ . Motivated by the different behavior of the Noether problem in positive and zero characteristic (see Section 1.2 for details) we study the following question:

**Question 1.1.** What can one say about the generic fiber  $X_K$  knowing that the special fiber  $X_k$  is stably rational? (The condition means that  $X_k \times \mathbb{P}_k^n$  is birational to  $\mathbb{P}_k^N$  for some  $n, N \in \mathbb{N}$ .)

In general,  $X_K$  need not be stably rational, as we show for instance in Section 3, where we follow a suggestion of Colliot-Thélène and adapt the constructions of [HPT18] to construct smooth proper morphisms  $X/R$  with  $X_K$  stably irrational and  $X_k$  rational. On the other hand, we show that the following vanishing holds:

**Theorem 1.2.** *Let  $X/R$  be a smooth proper morphism, and assume that the special fiber  $X_k$  is stably rational. Then the  $p$ -torsion of  $H_{\text{ét}}^3(X_K, \mathbb{Z}_p)$  vanishes.*

MSC2020: 14E08, 14F30, 14M20.

Keywords: Noether's problem, crystalline cohomology, Artin–Mumford invariant.

If  $K \subset \mathbb{C}$  is an embedding, then the vanishing of  $H_{\text{ét}}^3(X_K, \mathbb{Z}_p)[p]$  together with smooth-proper base change in étale cohomology (with  $\ell$ -adic coefficients,  $\ell \neq p$ ) implies that  $H_{\text{sing}}^3(X(\mathbb{C}), \mathbb{Z})$  is torsion free whenever  $X_k$  is stably rational. We recall that the torsion of  $H_{\text{sing}}^3(X(\mathbb{C}), \mathbb{Z})$  is a stably birational invariant of smooth proper varieties over  $\mathbb{C}$ , which is naturally isomorphic to  $\text{Br}(X)/\text{Br}(X)_{\text{div}}$ , where  $\text{Br}(X) := H_{\text{ét}}^2(X, \mathbb{G}_m)$ . This group was used by Artin and Mumford in their seminal paper [AM72] to give the first elementary example of a unirational threefold which was not rational.

**1.1. Strategy.** To explain the strategy, we begin by showing an analogous statement concerning global differential forms. We retain the notation from the previous section. By the Künneth formula and Hartogs' lemma, the vector spaces  $H^0(X_K, \Omega_{X_K}^i)$  are stably birational invariants of smooth proper  $K$ -varieties, and therefore they vanish if  $X_K$  is stably rational. If  $X/R$  is a smooth proper morphism, the semicontinuity theorem yields the inequality

$$(1.1.3) \quad \dim_k(H^0(X_k, \Omega_{X_k}^i)) \geq \dim_K(H^0(X_K, \Omega_{X_K}^i)),$$

from which it follows that if  $X_k$  is stably rational, then  $H^0(X_K, \Omega_{X_K}^i) = 0$  necessarily.

Concerning étale cohomology, as already mentioned, the proper smooth base change asserts that for a prime  $\ell \neq p$  one has

$$H_{\text{ét}}^3(X_K, \mathbb{Z}_\ell)[\ell] \simeq H_{\text{ét}}^3(X_k, \mathbb{Z}_\ell)[\ell] = 0,$$

where the last equality follows from the stable birational invariance of  $H_{\text{ét}}^3(X_k, \mathbb{Z}_\ell)[\ell]$  and the fact that  $X_k$  is stably rational. For  $p$ -adic coefficients, we can replace the smooth proper base-change with the integral  $p$ -adic Hodge theory of Bhatt, Morrow and Scholze, which will play the role of the semicontinuity theorem (1.1.3). By [BMS18, Theorem 1.1 (ii)], one has the inequality

$$(1.1.4) \quad \dim_k(H_{\text{crys}}^3(X_k/W)[p]) \geq \dim_{\mathbb{F}_p}(H_{\text{ét}}^3(X_K, \mathbb{Z}_p)[p]),$$

where  $H_{\text{crys}}^3(X_k/W)$  is the third crystalline cohomology group with integral coefficients of  $X_k$ . Thus, it would be enough to show that  $\dim_k(H_{\text{crys}}^3(X_k)[p])$  is a stably birational invariant of smooth proper varieties. It is unclear how to prove this without assuming resolution of singularities, also because crystalline cohomology behaves badly for open or singular varieties. We prove instead the following vanishing, which is enough to deduce Theorem 1.2:

**Theorem 1.1.5.** *Let  $k$  be an algebraically closed field of characteristic  $p$  and  $X$  be a smooth proper  $k$ -variety. Assume that*

- (1)  $H^i(X, \mathcal{O}_X) = 0$  for  $i = 2, 3$ ;
- (2)  $H^0(X, \Omega_X^2) = 0$ ;
- (3)  $\text{Br}(X)[p] = 0$ .

*Then  $H_{\text{crys}}^3(X/W)[p] = 0$ .*

The assumptions of Theorem 1.1.5 are satisfied if  $X$  is stably rational, since all the conditions are stably birational invariants of smooth proper varieties. (See [CR11, Theorem 1] for (1); point (2) follows from Hartog's lemma; [CTS21, Corollary 6.2.11, p. 170] gives (3).) Hence, Theorem 1.2 follows from Theorem 1.1.5 and (1.1.4).

**1.2. Applications to the Noether problem.** As mentioned, our original motivation was the Noether problem, which we now briefly recall. For a finite group  $G$  and a field  $K$ , the Noether problem asks whether  $\mathbb{P}_K^n/G$  is a stably rational variety, where  $G$  acts on  $\mathbb{P}_K^n$  in a linear and faithful way. The problem is well-posed since the stably birational class of the quotient does not depend on the chosen representation [BK85, Lemma 1.3]. For  $K = \mathbb{C}$  we can then define the Artin–Mumford invariant  $\text{AM}(G)$  of  $G$  as  $\text{Tors}(H_{\text{sing}}^3(X, \mathbb{Z}))$  where  $X/\mathbb{C}$  is any smooth proper birational model of  $\mathbb{P}_K^n/G$ .

The first counterexample of the Noether problem over  $\mathbb{C}$  was given by Saltman in [Sal84], who for any prime  $p$  produced a finite  $p$ -group  $G$  for which the quotient  $\mathbb{P}_{\mathbb{C}}^n/G$  is not stably rational, by showing that  $\text{AM}(G) \neq 0$ . A general formula for  $\text{AM}(G)$  was later given by Bogomolov [Bog88, Theorem 3.1], and this group is now known as the Bogomolov multiplier of  $G$ .

The connection to Theorem 1.2 comes from the classical observation that if  $G$  is a  $p$ -group and  $K$  has characteristic  $p$ , then  $\mathbb{P}_K^n/G$  is always rational; see, e.g., [Kun54] and [Gas59].

Keeping the notation as in the previous section, we can fix a finite  $p$ -group  $G$ , and linear faithful actions of  $G$  on  $\mathbb{P}_k^n$  and  $\mathbb{P}_K^n$ . Theorem 1.2 together with the observation above implies the following:

**Corollary 1.2.1.** *If  $\text{AM}(G) \neq 0$ , there does not exist a smooth proper  $X/R$  such that  $X_k$  is stably birational to  $\mathbb{P}_k^n/G$  and  $X_K$  is stably birational to  $\mathbb{P}_K^n/G$ .*

It follows that all the examples constructed in [Sal84; Bog88] cannot have good stably rational reduction. This also implies that if  $\text{AM}(G) \neq 0$  and  $G \rightarrow \text{PGL}_n(R)$  is a representation such that the reduction map  $G \rightarrow \text{PGL}_n(k)$  is injective, then one cannot resolve the singularities of  $\mathbb{P}_R^n/G$  relatively to  $R$ , i.e., there cannot be a smooth projective  $X/R$  with a  $R$ -morphism  $\pi : X \rightarrow \mathbb{P}_R^n/G$  which induces a resolution of singularities on both fibers.

Peyre [Pey08] constructed groups  $G$  such that  $\mathbb{P}_{\mathbb{C}}^n/G$  is not rational but  $\text{AM}(G) = 0$ . It is an interesting question at this point whether there is a  $p$ -group  $G$  for which a resolution  $X$  of  $\mathbb{P}_R^n/G$  like in the corollary above exists, but  $\mathbb{P}_K^n/G$  is not stably rational.

**Remark 1.2.2.** A related phenomenon appears in the recent work of Lazda and Skorobogatov in [LS23]. They prove that, if  $p = 2$  and  $Y \rightarrow \text{Spec}(R)$  is an abelian surface such that  $X_k$  is not supersingular, then one can resolve the singularities of  $Y/\{\pm 1\}$  to obtain a smooth proper morphism  $X \rightarrow \text{Spec}(R)$  such that the generic fiber is the Kummer variety  $\text{Kum}(Y_k)$  of  $Y_k$  and the special fiber is the Kummer variety  $\text{Kum}(Y_K)$  of  $Y_K$ .

On the other hand, if  $Y_k$  is supersingular, then  $\text{Kum}(Y_k)$  is a rational surface due to [Kat78]. Since  $H^0(\text{Kum}(Y_K), \Omega_{Y_K}^2) \neq 0$ , the argument in the beginning of Section 1.1 applies, and such  $Y$  does not exist. Since  $H^0(\text{Kum}(Y_k), \Omega_{Y_k}^2) = 0$  if and only if  $X_k$  is supersingular, the group  $H^0(\text{Kum}(Y_K), \Omega_{Y_K}^2)$  is the only obstruction to the construction of such  $Y$  in this case.

**2. Proof of Theorem 1.1.5**

**2.1. Notation.** Let  $k$  be an algebraically closed field of characteristic  $p > 0$  and let  $X/k$  be a smooth proper variety such that

$$H^2(X, \mathcal{O}_X) = H^3(X, \mathcal{O}_X) = H^0(X, \Omega_{X/k}^2) = \text{Br}(X)[p] = 0.$$

We let  $\Omega_X^\bullet$  be the de Rham complex of  $X$  and we define the following sheaves over  $X$ :

$$Z_X^i := \text{Ker}(d : \Omega_X^i \rightarrow \Omega_X^{i+1}); \quad B_X^i := \text{Im}(d : \Omega_X^{i-1} \rightarrow \Omega_X^i); \quad \mathcal{H}_X^i = \frac{Z_X^i}{B_X^i}.$$

For every complex of sheaves  $\mathcal{F}^\bullet$  over  $X$  and every  $i \in \mathbb{N}$ , we let  $\tau_{\geq i}\mathcal{F}^\bullet$  (resp.  $\tau_{\leq i}\mathcal{F}^\bullet$ ) be the upper (resp. lower) canonical truncation of  $\mathcal{F}^\bullet$  and  $\mathcal{F}^{\geq i}$  (resp.  $\mathcal{F}^{\leq i}$ ) the upper (resp. lower) naive filtration of  $\mathcal{F}^\bullet$ . Recall that, for every  $i \in \mathbb{N}$  there exists an exact triangle (see, e.g., [Stacks, Remark 08J5]):

$$\tau_{\leq i}\mathcal{F}^\bullet \rightarrow \mathcal{F}^\bullet \rightarrow \tau_{\geq i+1}\mathcal{F}^\bullet.$$

**2.2. Preliminary reductions.** The universal coefficient theorem for crystalline cohomology (see, e.g., [Ill79, (4.9.1), p. 623]), gives us an exact sequence

$$0 \rightarrow H_{\text{crys}}^2(X/W) \otimes k \rightarrow H_{\text{dR}}^2(X) \rightarrow H_{\text{crys}}^3(X/W)[p] \rightarrow 0,$$

so that, to prove Theorem 1.1.5, it is enough to show that the natural map

$$H_{\text{crys}}^2(X/W) \otimes k \rightarrow H_{\text{dR}}^2(X)$$

is surjective. In fact, we shall prove that the cycle class map  $\text{cl}_{\text{dR}} : \text{NS}(X) \otimes k \rightarrow H_{\text{dR}}^2(X/k)$  is surjective, which is enough due to the commutative diagram

$$(2.2.1) \quad \begin{array}{ccc} \text{NS}(X) \otimes W & \xrightarrow{\text{cl}_{\text{crys}}} & H_{\text{crys}}^2(X/W) \\ \downarrow & & \downarrow \\ \text{NS}(X) \otimes k & \xrightarrow{\text{cl}_{\text{dR}}} & H_{\text{dR}}^2(X), \end{array}$$

where the first horizontal arrow is the crystalline cycle class map.

**2.3. Factorization of the cycle class map.** To study  $\text{cl}_{\text{dR}} : \text{NS}(X) \otimes k \rightarrow H_{\text{dR}}^2(X/k)$ , we factorize it in three arrows. Let  $\iota : Z_X^1[-1] \rightarrow \Omega_X^\bullet$  be the natural inclusion.

Recall that, by construction, the cycle class map  $\text{cl}_{\text{dR}} : \text{NS}(X) \otimes k \rightarrow H_{\text{dR}}^2(X/k)$  is induced by the dlog map

$$\text{dlog} : \mathcal{O}_X^*[-1] \rightarrow \Omega_X^\bullet$$

(see, e.g., [Stacks, Section 0FLE]), which factors through

$$\mathcal{O}_X^*[-1] \xrightarrow{\text{dlog}} Z_X^1[-1] \xrightarrow{\iota} \Omega_X^\bullet.$$

Hence, the cycle class map factors through the induced map  $\iota : H^1(X, Z_X^1) \rightarrow H_{\text{dR}}^2(X)$ , giving a first factorization

$$\text{cl}_{\text{dR}} : \text{NS}(X) \otimes k \rightarrow H^1(X, Z_X^1) \xrightarrow{\iota} H_{\text{dR}}^2(X/k).$$

To go further, recall from [Ill79, (5.1.4), p. 626], that for every  $i$ , there is a canonical isomorphism  $H_{\text{flat}}^i(X, \mu_p) \xrightarrow{\cong} H^{i-1}(X, \mathcal{O}_X/(\mathcal{O}_X^*)^p)$  hence a canonical map  $\alpha : H^2(X, \mu_p) \rightarrow H^1(X, Z_X^1)$  induced again by  $\text{dlog} : \mathcal{O}_X/(\mathcal{O}_X^*)^p \rightarrow Z_X^1$ . In conclusion, the map  $\text{cl}_{\text{dR}} : \text{NS}(X) \otimes k \rightarrow H_{\text{dR}}^2(X)$  factorises further as

$$\text{NS}(X) \otimes k \xrightarrow{\text{cl}_{\text{flat}} \otimes k} H_{\text{flat}}^2(X, \mu_p) \otimes k \xrightarrow{\alpha} H^1(X, Z_X^1) \xrightarrow{\iota} H_{\text{dR}}^2(X),$$

where  $\text{cl}_{\text{flat}} : \text{NS}(X) \rightarrow H_{\text{flat}}^2(X, \mu_p)$  is the cycle class map in flat cohomology.

**2.4. Studying the factorization.** To show that  $\text{cl}_{\text{dR}} : \text{NS}(X) \otimes k \rightarrow H_{\text{dR}}^2(X)$  is surjective, we show that:

- (1) the map  $\text{cl}_{\text{flat}} : \text{NS}(X) \rightarrow H_{\text{flat}}^2(X, \mu_p)$  is surjective;
- (2) the map  $\alpha : H_{\text{flat}}^2(X, \mu_p) \otimes k \rightarrow H^1(X, Z_X^1)$  is an isomorphism;
- (3) the map  $\iota : H^1(X, Z_X^1) \rightarrow H_{\text{dR}}^2(X)$  is surjective.

*Proof of (1).* The map  $\text{cl}_{\text{flat}} : \text{NS}(X) \rightarrow H_{\text{flat}}^2(X, \mu_p)$  is induced by the connecting map

$$H^1(X, \mathcal{O}_X^*) = \text{Pic}(X) \rightarrow H_{\text{flat}}^2(X, \mu_p)$$

in the long exact sequence associated to the short exact sequence

$$0 \rightarrow \mu_p \rightarrow \mathcal{O}_X^* \xrightarrow{(-)^p} \mathcal{O}_X^* \rightarrow 0$$

of sheaves in flat site of  $X$ . Hence the surjectivity of  $\text{cl}_{\text{flat}} : \text{NS}(X) \rightarrow H_{\text{flat}}^2(X, \mu_p)$  follows directly from the assumption  $H_{\text{flat}}^2(X, \mathcal{O}_X^*) = H_{\text{ét}}^2(X, \mathcal{O}_X^*) = \text{Br}(X) = 0$ .

*Proof of (2).* Recall from [Ill79, (2.1.23), p. 518] the exact sequence of étale sheaves

$$0 \rightarrow \mathcal{O}_X^*/(\mathcal{O}_X^*)^p \xrightarrow{\text{dlog}} Z_X^1 \xrightarrow{i-C} \Omega_X^1 \rightarrow 0,$$

where  $i : Z_X^1 \rightarrow \Omega_X^1$  is the natural inclusion and  $C : Z_X^1 \rightarrow \Omega_X^1$  is the Cartier operator. Since étale cohomology and Zariski cohomology agree for coherent sheaves, taking the associated long exact sequence in cohomology, we get an exact sequence

$$H^0(X, Z_X^1) \xrightarrow{i-C} H^0(X, \Omega_X^1) \rightarrow H_{\text{fppf}}^2(X, \mu_p) \rightarrow H^1(X, Z_X^1) \xrightarrow{i-C} H^1(X, \Omega_X^1)$$

where  $i$  is a linear morphism and  $C$  is a Frobenius-linear map.

To prove (2), it is then enough to show that  $H^0(X, Z_X^1) \xrightarrow{i-C} H^0(X, \Omega_X^1)$  is surjective and that

$$\text{Ker}(H^1(X, Z_X^1) \xrightarrow{i-C} H^1(X, \Omega_X^1)) \otimes k \simeq H^1(X, Z_X^1).$$

Since  $H^i(X, Z_X^1)$  and  $H^i(X, \Omega_X^1)$  are finite-dimensional vector spaces, we can use the following classical lemma (see, e.g., [Mil16, Lemma 4.13, p. 128]).

**Lemma 2.4.1.** *Let  $k$  be an algebraically closed field of characteristic  $p > 0$  and let  $V$  be a finite-dimensional vector space over  $k$ . Let  $f : V \rightarrow V$  be a  $k$ -linear isomorphism and let  $C : V \rightarrow V$  be any Frobenius-linear map. Then:*

- (a)  $f - C$  is surjective.
- (b) If  $C$  is bijective, then  $V = \ker(f - C) \otimes k$ .

Hence to prove (2) it is enough to show that the following maps are isomorphisms:

$$i : H^0(X, Z_X^1) \rightarrow H^0(X, \Omega_X^1), \quad i : H^1(X, Z_X^1) \rightarrow H^1(X, \Omega_X^1), \quad C : H^1(X, Z_X^1) \rightarrow H^1(X, \Omega_X^1).$$

That the map  $H^0(X, Z_X^1) \rightarrow H^0(X, \Omega_X^1)$  is an isomorphism follows directly from the exact sequence of sheaves

$$0 \rightarrow Z_X^1 \rightarrow \Omega_X^1 \xrightarrow{d} \Omega_X^2$$

and the assumption  $H^0(X, \Omega_X^2) = 0$ .

As for  $i$  and  $C$ , by comparing dimensions we see that it is enough to prove that

- (i)  $i : H^1(X, Z_X^1) \rightarrow H^1(X, \Omega_X^1)$  is injective, and
- (ii)  $C : H^1(X, Z_X^1) \rightarrow H^1(X, \Omega_X^1)$  is surjective.

*Proof of (i).* Since  $Z_X^1[-1] = \tau_{\leq 1} \Omega_X^{\geq 1}$ , there is an exact triangle

$$Z_X^1[-1] \rightarrow \Omega_X^{\geq 1} \rightarrow \tau_{\geq 2} \Omega_X^{\geq 1},$$

so that the map  $H^1(X, Z_X^1) \rightarrow H^2(X, \Omega_X^{\geq 1})$  is injective, since  $H^1(X, \tau_{\geq 2} \Omega_X^{\geq 1}) = 0$ , because  $\tau_{\geq 2} \Omega_X^{\geq 1}$  is concentrated in degrees  $\geq 2$ . So it is enough to show that the natural map  $H^2(X, \Omega_X^{\geq 1}) \rightarrow H^2(X, \Omega_X^1[-1]) = H^1(X, \Omega_X^1)$  induced by the map  $\Omega_X^{\geq 1} \rightarrow \Omega_X^1[-1]$  is injective. But the latter fits in the short exact sequence of complexes

$$0 \rightarrow \Omega_X^{\geq 2} \rightarrow \Omega_X^{\geq 1} \rightarrow \Omega_X^1[-1] \rightarrow 0,$$

so that we just need to prove that  $H^2(X, \Omega_X^{\geq 2}) = 0$ . This follows from the inclusion

$$H^2(X, \Omega_X^{\geq 2}) = \text{Ker}(H^0(X, \Omega_X^2) \xrightarrow{d} H^0(X, \Omega_X^3)) \subseteq H^0(X, \Omega_X^2) = 0$$

by our assumption on  $X$ . Hence  $i : H^1(X, Z_X^1) \rightarrow H^1(X, \Omega_X^1)$  is injective.

*Proof of (ii).* Since  $k$  is perfect, from [Ill79, (2.1.22)] we have a short exact sequence of sheaves

$$0 \rightarrow B_X^1 \rightarrow Z_X^1 \xrightarrow{C} \Omega_X^1 \rightarrow 0,$$

where  $C$  is the Cartier operator. Hence it is enough to show that  $H^2(X, B_X^1)$  is zero. But this follows from the short exact sequence

$$0 \rightarrow \mathcal{O}_X \xrightarrow{(-)^p} \mathcal{O}_X \xrightarrow{d} B_X^1 \rightarrow 0$$

and the assumption  $H^2(X, \mathcal{O}_X) = H^3(X, \mathcal{O}_X) = 0$ .

*Proof of (3).* Since  $H^2(X, \mathcal{O}_X) = 0$  by assumption, the short exact sequence

$$0 \rightarrow Z_X^1[-1] \rightarrow \tau_{\leq 1}\Omega_X^\bullet \rightarrow \mathcal{O}_X \rightarrow 0$$

shows that the natural map  $H^1(X, Z_X^1) \rightarrow H^2(X, \tau_{\leq 1}\Omega_X^\bullet)$  is surjective. So, it is enough to show that the natural map  $H^2(X, \tau_{\leq 1}\Omega_X^\bullet) \rightarrow H^2(X, \Omega_X^\bullet) = H_{\text{dR}}^2(X)$  is surjective. Since there is an exact triangle

$$\tau_{\leq 1}\Omega_X^\bullet \rightarrow \Omega_X^\bullet \rightarrow \tau_{\geq 2}\Omega_X^\bullet,$$

it is then enough to show that  $H^2(X, \tau_{\geq 2}\Omega_X^\bullet) = 0$ . But

$$H^2(X, \tau_{\geq 2}\Omega_X^\bullet) = \ker(H^0(X, \tau_{\geq 2}\Omega_X^\bullet[2]) \rightarrow H^0(X, \Omega_X^3)) = H^0(X, \ker(\Omega_X^2/B_X^2 \rightarrow \Omega_X^3)) = H^0(X, \mathcal{H}_X^2).$$

Again, since  $k$  is perfect, the inverse Cartier operator [Ill79, (2.1.22)] gives an isomorphism  $\mathcal{H}^2(\Omega_X^\bullet) \cong \Omega_X^2$ , so

$$H^2(X, \tau_{\geq 2}\Omega_X^\bullet) \simeq H^0(X, \Omega_X^2) = 0.$$

Hence the natural map  $H^1(X, Z_X^1) \rightarrow H_{\text{dR}}^2(X/k)$  is surjective. This concludes the proof of (3) and the proof of Theorem 1.1.5.

### 3. A stably irrational variety reducing to a rational variety

Let  $R$  be the ring of integers of  $K := \mathbb{C}_p$  and  $k$  its residue field. In this last section, we show how to construct, for every  $p \gg 0$ , examples of smooth proper schemes  $X \rightarrow \text{Spec}(R)$  such that  $X_K$  is not stably rational and such that  $X_k$  is rational, as suggested to us by Colliot-Thélène. The construction uses and is based on the analogous construction in [HPT18] of a family of proper smooth varieties over the complex number with stably irrational general fiber but with some rational fiber.

**3.1. A general lemma.** We begin by reducing the construction of examples to the construction of mixed characteristic families with properties that are easier to check. Let  $B/R$  be smooth with geometrically integral fibers and  $X \rightarrow B$  a smooth proper family of varieties.

**Lemma 3.1.1.** *Assume that there exists a point  $b \in B(\mathbb{C}_p)$  such that  $X_b$  is not stably rational. Then, for every  $a \in B(k)$  there exists a lift  $b' \in B(R)$  of  $a$  such that  $X_{b'}$  is not stably rational.*

*Proof.* By [NO21, Corollary 4.1.2], the set

$$B(\mathbb{C}_p)_r := \{b \in B(\mathbb{C}_p) : X_b \text{ is stably rational}\}$$

is a countable union of closed subvarieties. Define now  $B(\mathbb{C}_p)_{nr} := B(\mathbb{C}_p) \setminus B(\mathbb{C}_p)_r$ . By the assumption on  $b$ , the set  $B(\mathbb{C}_p)_r$  is the countable union of proper closed subvarieties.

Since, by the Hensel lemma, the map  $\pi : B(R) \rightarrow B(k)$  is surjective, we can choose a lift  $b'$  of  $a$ . The set  $\pi^{-1}(a) \subseteq B(R) \subseteq B(K)$  is an open neighborhood of  $b'$  in  $B(K)$ . Since  $B(\mathbb{C}_p)_r$  is the countable union of proper closed subvarieties, we can apply [MP12, Lemma 4.29] to deduce that there exists a  $b' \in B(\mathbb{C}_p)_{nr} \cap \pi^{-1}(a)$ . This concludes the proof.  $\square$

**3.2. An example.** By Lemma 3.1.1, to construct a smooth proper scheme  $X \rightarrow \text{Spec}(R)$  such that  $X_K$  is not stably rational and such that  $X_k$  is rational, it is enough to construct a family  $X \rightarrow B$  over  $R$  such that there exist points  $b \in B(\mathbb{C}_p)$  and  $a \in B(k)$  such that  $X_b$  is stably irrational and  $X_a$  is rational. Such a family can be constructed directly using the example of Hassett, Pirutka, and Tschinkel [HPT18]; see also [CTS21, Section 12.2.2] and [Sch19]. We give some details.

Let  $X' \rightarrow Z$  be the universal family of quadric bundles over  $\mathbb{P}_{\mathbb{Q}}^2$  given in  $\mathbb{P}_{\mathbb{Q}}^2 \times \mathbb{P}_{\mathbb{Q}}^3$  by a bihomogeneous form of bidegree  $(2, 2)$ . After choosing coordinates  $x, y, z$  and  $U, V, W, T$  on  $\mathbb{P}_{\mathbb{Q}}^2 \times \mathbb{P}_{\mathbb{Q}}^3$ , the variety  $Z$  identifies with the space of bihomogeneous forms  $F = F(x, y, z, U, V, W, T)$  of bidegree  $(2, 2)$  in  $\mathbb{P}_{\mathbb{Q}}^2 \times \mathbb{P}_{\mathbb{Q}}^3$  that are symmetric quadratic forms in the variables  $U, V, W, T$ , since any such  $F$  determines a quadric bundle over  $\mathbb{P}_{\mathbb{Q}}^2$  via the projection  $\mathbb{P}_{\mathbb{Q}}^2 \times \mathbb{P}_{\mathbb{Q}}^3 \rightarrow \mathbb{P}_{\mathbb{Q}}^2$ . In turn, these forms are given by a  $4 \times 4$  symmetric matrix  $A = (a_{i,j})_{1 \leq i, j \leq 4}$  where each entry  $a_{i,j} = a_{i,j}(x, y, z)$  is a homogeneous polynomial of degree 2.

By the arguments in [Sch19] and Bertini's theorem, there exists a dense open Zariski  $B_{\mathbb{Q}} \subset Z$  such that the restriction of the family  $X_{\mathbb{Q}} \rightarrow B_{\mathbb{Q}}$  to  $B_{\mathbb{Q}}$  parametrizes quadric bundle flat over  $\mathbb{P}_{\mathbb{Q}}^2$  and with smooth total space.

By spreading out, this construction extends to give a smooth family  $X \rightarrow B$  over  $\mathbb{Z}[1/n]$  for  $n$  large enough whose base change to  $\mathbb{Q}$  identifies with  $X_{\mathbb{Q}} \rightarrow B_{\mathbb{Q}}$ . By the main result of [HPT18], the general fiber of  $X(\mathbb{C}) \rightarrow B(\mathbb{C})$  is not stably rational, hence, for every  $p$ , there exists a  $b \in B(\mathbb{C}_p)$  such that  $Y_b$  is not stably rational. So, we are left to show that for  $p \gg 0$ , there exists a  $a \in B(\overline{\mathbb{F}}_p)$  such that  $X_a$  is rational.

Using Bertini, there exists a rational point  $r \in B(\overline{\mathbb{Q}})$  such that the corresponding  $4 \times 4$  symmetric matrix  $A = (a_{i,j})_{1 \leq i, j \leq 4}$  has  $a_{1,1} = 0$ . By spreading out, we can choose  $p \gg 0$  such that  $r \in B(\overline{\mathbb{Q}})$  extends to a point  $\tilde{a} \in B(\overline{\mathbb{Z}}_p)$  whose reduction  $a$  modulo  $p$  defines a flat quadric bundle  $X_a \rightarrow \mathbb{P}_{\mathbb{F}_p}^2$  with smooth total space, whose associated matrix has  $a_{1,1} = 0$ . Since  $a_{1,1} = 0$ , for every  $x \in \mathbb{P}_{\mathbb{F}_p}^2$ , the point  $[1 : 0 : 0 : 0]$  is  $k(x)$ -rational point of the fiber of  $X_a \rightarrow \mathbb{P}_{\mathbb{F}_p}^2$  in  $x$ . In particular the morphism  $X_a \rightarrow \mathbb{P}_{\mathbb{F}_p}^2$  has a rational section, hence  $X_a$  is rational. This concludes the construction of a proper smooth scheme over  $R$  with rational special fiber and stably irrational generic fiber.

### Acknowledgements

The authors are grateful to Stefan Schreieder for having introduced Valloni to the Noether problem, to Giuseppe Ancona for many interesting discussions and comments on a preliminary version of the article, and to Colliot-Thélène for pointing out to us the example in Section 3. The authors also thank the referees for their comments and for having greatly improved both the statement and proof of our main result.

### References

- [ABBGvB21] A. Auel, A. Bigazzi, C. Böhning, and H.-C. Graf von Bothmer, “Unramified Brauer groups of conic bundle threefolds in characteristic two”, *Amer. J. Math.* **143**:5 (2021), 1601–1631. MR
- [AM72] M. Artin and D. Mumford, “Some elementary examples of unirational varieties which are not rational”, *Proc. Lond. Math. Soc.* (3) **25** (1972), 75–95. MR

- [BK85] F. A. Bogomolov and P. I. Katsylo, "Rationality of some quotient varieties", *Mat. Sb. (N.S.)* **126(168)**:4 (1985), 584–589. In Russian; translated in *Math. USSR-Sb.* **54**:2 (1986), 571–576. MR
- [BMS18] B. Bhatt, M. Morrow, and P. Scholze, "Integral  $p$ -adic Hodge theory", *Publ. Math. Inst. Hautes Études Sci.* **128** (2018), 219–397. MR
- [Bog88] F. A. Bogomolov, "The Brauer group of quotient spaces by linear group actions", *Izv. Akad. Nauk SSSR Ser. Mat.* **51**:3 (1987), 485–516. In Russian; translated in *Math. USSR-Izv.* **30**:3 (1988), 455–485. MR
- [CR11] A. Chatzistamatiou and K. Rülling, "Higher direct images of the structure sheaf in positive characteristic", *Algebra Number Theory* **5**:6 (2011), 693–775. MR
- [CTP16] J.-L. Colliot-Thélène and A. Pirutka, "Hypersurfaces quartiques de dimension 3: non-rationalité stable", *Ann. Sci. École Norm. Sup. (4)* **49**:2 (2016), 371–397. MR
- [CTS21] J.-L. Colliot-Thélène and A. N. Skorobogatov, *The Brauer–Grothendieck group*, *Ergebnisse der Math. (3)* **71**, Springer, 2021. MR
- [Gas59] W. Gaschütz, "Fixkörper von  $p$ -Automorphismengruppen rein-transzendenter Körpererweiterungen von  $p$ -Charakteristik", *Math. Z.* **71** (1959), 466–468. MR
- [HPT18] B. Hassett, A. Pirutka, and Y. Tschinkel, "Stable rationality of quadric surface bundles over surfaces", *Acta Math.* **220**:2 (2018), 341–365. MR
- [Ill79] L. Illusie, "Complexe de de Rham–Witt et cohomologie cristalline", *Ann. Sci. École Norm. Sup. (4)* **12**:4 (1979), 501–661. MR
- [Kat78] T. Katsura, "On Kummer surfaces in characteristic 2", pp. 525–542 in *Proceedings of the International Symposium on Algebraic Geometry* (Kyoto, 1977), edited by M. Nagata, Kinokuniya, Tokyo, 1978. MR
- [Kol95] J. Kollár, "Nonrational hypersurfaces", *J. Amer. Math. Soc.* **8**:1 (1995), 241–249. MR
- [Kun54] H. Kuniyoshi, "On purely-transcendence of a certain field", *Tohoku Math. J. (2)* **6** (1954), 101–108. MR
- [LS23] C. D. Lazda and A. N. Skorobogatov, "Reduction of Kummer surfaces modulo 2 in the non-supersingular case", *Épjournal Géom. Algébrique* **7** (2023), art. id. 10. MR
- [Mil16] J. S. Milne, *Étale cohomology*, Princeton Math. Ser. **33**, Princeton Univ. Press, 1980. MR
- [MP12] D. Maulik and B. Poonen, "Néron–Severi groups under specialization", *Duke Math. J.* **161**:11 (2012), 2167–2206. MR
- [NO21] J. Nicaise and J. C. Ottem, "A refinement of the motivic volume, and specialization of birational types", pp. 291–322 in *Rationality of varieties*, edited by G. Farkas et al., Progr. Math. **342**, Birkhäuser, Cham, 2021. MR
- [Pey08] E. Peyre, "Unramified cohomology of degree 3 and Noether's problem", *Invent. Math.* **171**:1 (2008), 191–225. MR
- [Sal84] D. J. Saltman, "Noether's problem over an algebraically closed field", *Invent. Math.* **77**:1 (1984), 71–84. MR
- [Sch19] S. Schreieder, "On the rationality problem for quadric bundles", *Duke Math. J.* **168**:2 (2019), 187–223. MR
- [Stacks] "The Stacks project", electronic reference, 2005–, available at <http://stacks.math.columbia.edu>.
- [Tot16] B. Totaro, "Hypersurfaces that are not stably rational", *J. Amer. Math. Soc.* **29**:3 (2016), 883–891. MR
- [Voi15] C. Voisin, "Unirational threefolds with no universal codimension 2 cycle", *Invent. Math.* **201**:1 (2015), 207–237. MR

Communicated by Jean-Louis Colliot-Thélène

Received 2023-04-02

Revised 2024-11-20

Accepted 2024-12-23

eambrosi@unistra.fr

IRMA (Université de Strasbourg), 67084 Strasbourg, France

domenico.valloni@epfl.ch

EPFL, 1015 Lausanne, Switzerland



# Algebra & Number Theory

msp.org/ant

## EDITORS

MANAGING EDITOR  
Antoine Chambert-Loir  
Université Paris-Diderot  
France

EDITORIAL BOARD CHAIR  
David Eisenbud  
University of California  
Berkeley, USA

## BOARD OF EDITORS

Jason P. Bell	University of Waterloo, Canada	Philippe Michel	École Polytechnique Fédérale de Lausanne
Bhargav Bhatt	University of Michigan, USA	Martin Olsson	University of California, Berkeley, USA
Frank Calegari	University of Chicago, USA	Irena Peeva	Cornell University, USA
J.-L. Colliot-Thélène	CNRS, Université Paris-Saclay, France	Jonathan Pila	University of Oxford, UK
Brian D. Conrad	Stanford University, USA	Anand Pillay	University of Notre Dame, USA
Samit Dasgupta	Duke University, USA	Bjorn Poonen	Massachusetts Institute of Technology, USA
Hélène Esnault	Freie Universität Berlin, Germany	Victor Reiner	University of Minnesota, USA
Gavril Farkas	Humboldt Universität zu Berlin, Germany	Peter Sarnak	Princeton University, USA
Sergey Fomin	University of Michigan, USA	Michael Singer	North Carolina State University, USA
Edward Frenkel	University of California, Berkeley, USA	Vasudevan Srinivas	SUNY Buffalo, USA
Wee Teck Gan	National University of Singapore	Shunsuke Takagi	University of Tokyo, Japan
Andrew Granville	Université de Montréal, Canada	Pham Huu Tiep	Rutgers University, USA
Ben J. Green	University of Oxford, UK	Ravi Vakil	Stanford University, USA
Christopher Hacon	University of Utah, USA	Akshay Venkatesh	Institute for Advanced Study, USA
Roger Heath-Brown	Oxford University, UK	Melanie Matchett Wood	Harvard University, USA
János Kollár	Princeton University, USA	Shou-Wu Zhang	Princeton University, USA
Michael J. Larsen	Indiana University Bloomington, USA		

## PRODUCTION

production@msp.org  
Silvio Levy, Scientific Editor

---

See inside back cover or [msp.org/ant](http://msp.org/ant) for submission instructions.

---

The subscription price for 2025 is US \$565/year for the electronic version, and \$820/year (+\$70, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to MSP.

---

Algebra & Number Theory (ISSN 1944-7833 electronic, 1937-0652 printed) at Mathematical Sciences Publishers, 2000 Allston Way # 59, Berkeley, CA 94701-4004, is published continuously online.

---

ANT peer review and production are managed by EditFLOW<sup>®</sup> from MSP.

PUBLISHED BY

 **mathematical sciences publishers**  
nonprofit scientific publishing

<http://msp.org/>

© 2025 Mathematical Sciences Publishers

# Algebra & Number Theory

Volume 19    No. 12    2025

---

Perfectoid towers and their tilts: with an application to the étale cohomology groups of local log-regular rings	2307
SHINNOSUKE ISHIRO, KEI NAKAZATO and KAZUMA SHIMOMOTO	
Reduction modulo $p$ of Noether's problem	2359
EMILIANO AMBROSI and DOMENICO VALLONI	
On the Grothendieck ring of a quasireductive Lie superalgebra	2369
MARIA GORELIK, VERA SERGANOVA and ALEXANDER SHERMAN	
Combing a hedgehog over a field	2409
ALEXEY ANANYEVSKIY and MARC LEVINE	
Self-correlations of Hurwitz class numbers	2433
ALEXANDER WALKER	
On two definitions of wave-front sets for $p$ -adic groups	2471
CHENG-CHIANG TSAI	
Irregular Hodge filtration of hypergeometric differential equations	2481
YICHEN QIN and DAXIN XU	