

# *Algebra & Number Theory*

Volume 19  
2025  
No. 5

**Syzygies of tangent-developable surfaces and K3 carpets via secant varieties**

Jinhyung Park





# Syzygies of tangent-developable surfaces and K3 carpets via secant varieties

Jinhyung Park

We give simple geometric proofs of Aprodu, Farkas, Papadima, Raicu and Weyman’s theorem on syzygies of tangent-developable surfaces of rational normal curves and Raicu and Sam’s result on syzygies of K3 carpets. As a consequence, we obtain a quick proof of Green’s conjecture for general curves of genus  $g$  over an algebraically closed field  $\mathbb{k}$  with  $\text{char}(\mathbb{k}) = 0$  or  $\text{char}(\mathbb{k}) \geq \lfloor (g - 1)/2 \rfloor$ . Our approach provides a new way to study tangent-developable surfaces in general. Along the way, we show the arithmetic normality of tangent-developable surfaces of arbitrary smooth projective curves of large degree.

## 1. Introduction

Let  $C$  be a smooth projective curve of genus  $g \geq 3$  over the field  $\mathbb{C}$  of complex numbers. If  $\text{Cliff}(C) \geq 1$ , i.e.,  $C$  is nonhyperelliptic, then  $K_C$  is very ample. In this case, Noether’s theorem says that the canonical curve  $C \subseteq \mathbb{P}^{g-1}$  is projectively normal. Petri’s theorem states that if  $\text{Cliff}(C) \geq 2$ , then the defining ideal  $I_{C|\mathbb{P}^{g-1}}$  of  $C$  in  $\mathbb{P}^{g-1}$  is generated by quadrics. To generalize classical theorems of Noether and Petri, Green [1984, Conjecture 5.1] formulated a very famous conjecture that predicts

$$K_{p,2}(C, K_C) = 0 \quad \text{for } 0 \leq p \leq \text{Cliff}(C) - 1.$$

By the Green–Lazarsfeld nonvanishing theorem [Green 1984, Appendix], we have  $K_{p,2}(C, K_C) \neq 0$  for  $\text{Cliff}(C) \leq p \leq g - 3$ . Then Green’s conjecture determines the shape of the minimal free resolution of the canonical ring  $R(C, K_C) = \bigoplus_{m \in \mathbb{Z}} H^0(C, mK_C)$  (see [Aprodu and Nagel 2010, Remark 4.19]). Although the conjecture is still open, Voisin [2002; 2005] resolved the general curve case in the early 2000s. To prove the generic Green conjecture, it suffices to exhibit one curve of each genus for which the assertion holds. If  $S \subseteq \mathbb{P}^g$  is a K3 surface of degree  $2g - 2$ , then its general hyperplane section is a canonical curve  $C \subseteq \mathbb{P}^{g-1}$  and  $K_{p,2}(S, \mathcal{O}_S(1)) = K_{p,2}(C, K_C)$ . Based on the Hilbert scheme interpretation of Koszul cohomology, Voisin accomplished sophisticated cohomology computations on Hilbert schemes of K3 surfaces to show  $K_{p,2}(S, \mathcal{O}_S(1)) = 0$ . For an introduction to Voisin’s work; see [Aprodu and Nagel 2010]. Recently, Kemeny [2021] gave a simpler proof of Voisin’s theorem for even-genus case and a streamlined version of her arguments for the odd-genus case.

---

Park was partially supported by the National Research Foundation (NRF) funded by the Korea government (MSIT) (RS-2021-NR061320 and RS-2022-NR119741).

MSC2020: 14N05, 14N07, 13D02.

Keywords: tangent-developable surfaces, K3 carpets, secant varieties, syzygies.

Prior to Voisin’s work, O’Grady and Buchweitz–Schreyer independently observed that one could use the *tangent-developable surface*  $T \subseteq \mathbb{P}^g$  of a rational normal curve of degree  $g$  to solve Green’s conjecture for general curves of genus  $g$  (see [Eisenbud 1992]). Note that  $T \subseteq \mathbb{P}^g$  is arithmetically Cohen–Macaulay as for a K3 surface. One can actually view  $T$  as a degeneration of a K3 surface (see [Aprodu et al. 2019, Remark 6.6]). A general hyperplane section of  $T$  is a canonically embedded  $g$ -cuspidal rational curve  $\bar{C} \subseteq \mathbb{P}^{g-1}$  of degree  $2g - 2$ , which is a degeneration of a general canonical curve  $C \subseteq \mathbb{P}^{g-1}$  with  $\text{Cliff}(C) = \lfloor (g - 1)/2 \rfloor$ . By the upper semicontinuity of graded Betti numbers,  $K_{p,2}(\bar{C}, \mathcal{O}_{\bar{C}}(1)) = K_{p,2}(T, \mathcal{O}_T(1)) = 0$  implies  $K_{p,2}(C, K_C) = 0$  (see [loc. cit., Section 6]). The required vanishing of  $K_{p,2}(T, \mathcal{O}_T(1))$  for confirming the generic Green conjecture was finally established by Aprodu, Farkas, Papadima, Raicu and Weyman [loc. cit.] just a few years ago. Their important result gives not only an alternative proof of the generic Green conjecture but also an extension to positive characteristic. This circle of ideas is surveyed in [Ein and Lazarsfeld 2020]. In this paper, we give a simple geometric proof of the main result of [Aprodu et al. 2019].

**Theorem 1.1.** *Let  $T \subseteq \mathbb{P}^g$  be the tangent-developable surface of a rational normal curve of degree  $g \geq 3$  over an algebraically closed field  $\mathbb{k}$  with  $\text{char}(\mathbb{k}) = 0$  or  $\text{char}(\mathbb{k}) \geq (g + 2)/2$ . Then*

$$K_{p,2}(T, \mathcal{O}_T(1)) = 0 \quad \text{for } 0 \leq p \leq \lfloor (g - 3)/2 \rfloor.$$

The proof of Theorem 1.1 in [Aprodu et al. 2019] goes as follows. We have a short exact sequence

$$0 \longrightarrow \mathcal{O}_T \longrightarrow \nu_* \mathcal{O}_{\tilde{T}} \longrightarrow \omega_{\mathbb{P}^1} \longrightarrow 0,$$

where  $\nu : \tilde{T} = \mathbb{P}^1 \times \mathbb{P}^1 \rightarrow T$  is a resolution of singularities. Then it is elementary to see that

$$K_{p,2}(T, \mathcal{O}_T(1)) = \text{coker}(K_{p,1}(T, \nu_* \mathcal{O}_{\tilde{T}}; \mathcal{O}_T(1)) \xrightarrow{\gamma} K_{p,1}(\mathbb{P}^1, \omega_{\mathbb{P}^1}; \mathcal{O}_{\mathbb{P}^1}(g))).$$

Let  $U := H^0(\mathbb{P}^1, \mathcal{O}_{\mathbb{P}^1}(1))$ ,  $V := D^{p+2}U$ ,  $W := D^{2p+2}U$ , and  $q := g - p - 3$ . The authors of [Aprodu et al. 2019] devoted considerable effort to show that  $\gamma$  arises as the composition

$$S^q V \otimes W \xrightarrow{\text{id}_{S^q V} \otimes \Delta} S^q V \otimes \wedge^2 V \xrightarrow{\delta} \ker(S^{q+1} V \otimes V \xrightarrow{\delta} S^{q+2} V), \tag{1-1}$$

where  $\Delta$  is the *co-Wahl map* and  $\delta$  is the *Koszul differential*. To achieve this, they established an explicit characteristic-free Hermite reciprocity for  $\mathfrak{sl}_2$ -representations, and they carried out complicated algebraic computations. Now,  $K_{p,2}(T, \mathcal{O}_T(1))$  is the homology of a complex

$$S^q V \otimes W \xrightarrow{\gamma} S^{q+1} V \otimes V \xrightarrow{\delta} S^{q+2} V.$$

This homology, denoted by  $W_q(V, W)$ , is the degree- $q$  piece of the *Koszul module* (or *Weyman module*) associated to  $(V, W)$ . It is enough to prove that

$$W_q(V, W) = 0 \quad \text{for } q \geq p. \tag{1-2}$$

The vanishing result (1-2) was first proved in characteristic zero in [Aprodu et al. 2022] by an application of Bott vanishing, and the argument is extended in [Aprodu et al. 2019] to positive characteristics.

Our strategy to prove Theorem 1.1 is essentially the same as that of [Aprodu et al. 2019], but our geometric approach utilizing the *secant variety*  $\Sigma \subseteq \mathbb{P}^g$  of a rational normal curve  $C$  of degree  $g$  provides a substantial simplification of the proof. The tangent surface  $T$  is a Weil divisor on  $\Sigma$ , and  $\tilde{T}$  is a Cartier divisor on  $B$ , where  $\beta : B \rightarrow \Sigma$  is the blow-up of  $\Sigma$  along  $C$  with the exceptional divisor  $Z = \mathbb{P}^1 \times \mathbb{P}^1$ . Letting  $M_H := \beta^* M_{\mathcal{O}_\Sigma(1)}$ , we realize (1-1) as maps induced in cohomology of vector bundles on  $B$  (see (3-1)):

$$H^1(\tilde{T}, \wedge^{p+2} M_H|_{\tilde{T}}) \xrightarrow{\alpha} H^2(B, \wedge^{p+2} M_H \otimes \omega_B(Z)) \xrightarrow{\delta} H^2(Z, \omega_{\mathbb{P}^1} \boxtimes \wedge^{p+2} M_L \otimes \omega_{\mathbb{P}^1}).$$

This was previously asked in [Aprodu et al. 2019, last paragraph on p. 666]. There is a rank-2 vector bundle  $E$  on  $\mathbb{P}^2$  such that  $B = \mathbb{P}(E)$ . If  $\pi : B \rightarrow \mathbb{P}^2$  is the canonical fibration, then  $Q := \pi(\tilde{T})$  is a smooth conic and  $\tilde{T} = \mathbb{P}(E|_Q)$ . It is easy to check that  $\alpha = \text{id}_{S^q V} \otimes \Delta$ , where  $\Delta$  is the dual of the restriction map  $H^0(\mathbb{P}^2, \mathcal{O}_{\mathbb{P}^2}(p+1)) \rightarrow H^0(Q, \mathcal{O}_Q(p+1))$ . Put  $M_E := \pi_* M_H$  and  $\sigma := \pi|_Z$ . The map  $\delta$  is naturally factored as

$$\begin{array}{ccc} H^2(B, \wedge^{p+2} M_H \otimes \omega_B(Z)) & \xrightarrow{\text{id}_{S^q V} \otimes \iota} & H^2(Z, \sigma^* \wedge^{p+2} M_E \otimes (\omega_{\mathbb{P}^1} \boxtimes \omega_{\mathbb{P}^1})) & \xrightarrow{m \otimes \text{id}_V} & H^2(Z, \omega_{\mathbb{P}^1} \boxtimes \wedge^{p+2} M_L \otimes \omega_{\mathbb{P}^1}) \\ \parallel & & \parallel & & \parallel \\ S^q V \otimes \wedge^2 V & & S^q V \otimes V \otimes V & & S^{q+1} V \otimes V \end{array}$$

where  $\iota$  is the canonical injection and  $m$  is the multiplication map identified with

$$H^1(\mathbb{P}^{p+2} \times \mathbb{P}^1, \mathcal{O}_{\mathbb{P}^{p+2}}(q) \boxtimes \omega_{\mathbb{P}^1}(-p-2)) \longrightarrow H^1(\mathbb{P}^{p+2} \times \mathbb{P}^1, \mathcal{O}_{\mathbb{P}^{p+2}}(q+1) \boxtimes \omega_{\mathbb{P}^1}).$$

Without any lengthy computation, we quickly obtain the key part of [Aprodu et al. 2019] — the descriptions of the maps in (1-1) (see Lemma 3.1). This provides a conceptual explanation of the difficult computation in [loc. cit.] and a geometric understanding of syzygies of  $T$  in  $\mathbb{P}^g$  as expected in [loc. cit., last paragraph on p. 666]. Next, regarding  $V = H^0(\mathbb{P}^{p+2}, \mathcal{O}_{\mathbb{P}^{p+2}}(1))$ , we give a direct proof of (1-2) using vector bundles on  $\mathbb{P}^{p+2}$ . This part of the proof is largely equivalent to the original proof in [Aprodu et al. 2019; 2022].<sup>1</sup> Put  $M_V := M_{\mathcal{O}_{\mathbb{P}^{p+2}}(1)}$ . Then (1-1) can be identified with

$$W \otimes H^0(\mathbb{P}^{p+2}, \mathcal{O}_{\mathbb{P}^{p+2}}(q)) \longrightarrow \wedge^2 V \otimes H^0(\mathbb{P}^{p+2}, \mathcal{O}_{\mathbb{P}^{p+2}}(q)) \longrightarrow H^0(\mathbb{P}^{p+2}, M_V(q+1)).$$

Under the assumption  $\text{char}(\mathbb{k}) = 0$  or  $\text{char}(\mathbb{k}) \geq (g+2)/2$ , we show that  $W \otimes \mathcal{O}_{\mathbb{P}^{p+2}} \rightarrow M_V(1)$  is surjective and its kernel is  $(p+1)$ -regular in the sense of Castelnuovo and Mumford. This implies (1-2). Here the characteristic assumption plays a crucial role (see Remark 3.3).

It is worth noting that the characteristic assumption on the field  $\mathbb{k}$  in Theorem 1.1 cannot be improved. If  $2 \leq \text{char}(\mathbb{k}) \leq (g+1)/2$ , then  $K_{\lfloor (g-3)/2 \rfloor, 2}(T, \mathcal{O}_T(1)) \neq 0$  (see [Aprodu et al. 2019, Remark 5.17]). Theorem 1.1 implies the generic Green conjecture for general curves of genus  $g$  when  $\text{char}(\mathbb{k}) = 0$  or  $\text{char}(\mathbb{k}) \geq (g+2)/2$ , but this is not optimal. Raicu and Sam [2022, Theorem 1.5] recently obtained a sharp result that Green’s conjecture holds for general curves of genus  $g$  when  $\text{char}(\mathbb{k}) = 0$  or  $\text{char}(\mathbb{k}) \geq \lfloor (g-1)/2 \rfloor$ . This confirms a conjecture of Eisenbud and Schreyer [2019, Conjecture 1.1]. For the failure of Green’s

<sup>1</sup>After writing the paper, the author learned from Claudiu Raicu [2021] that a similar argument proving (1-2) directly on projective spaces is given in his lecture notes based on Robert Lazarsfeld’s suggestion.

conjecture in small characteristic, see [Eisenbud and Schreyer 2019; Schreyer 1986]. It has long been known that the generic Green conjecture would follow from the canonical ribbon conjecture [Bayer and Eisenbud 1995]. A *canonical ribbon* is a hyperplane section of a *K3 carpet*  $X = X(a, b) \subseteq \mathbb{P}^{a+b+1}$  for integers  $b \geq a \geq 1$ , which is a unique double structure on a rational normal surface scroll  $S(a, b) \subseteq \mathbb{P}^{a+b+1}$  of type  $(a, b)$  such that  $\omega_X = \mathcal{O}_X$  and  $h^1(X, \mathcal{O}_X) = 0$  (see [Gallego and Purnaprajna 1997, Theorem 1.3]). The K3 carpet  $X$  is a degeneration of a K3 surface of degree  $2(a + b)$ , and a canonical ribbon is a degeneration of a general canonical curve  $C$  of genus  $a + b + 1$  with  $\text{Cliff}(C) = a$ . By extending algebraic arguments of [Aprodu et al. 2019], Raicu and Sam [2022, Theorem 1.1] proved  $K_{p,2}(X, \mathcal{O}_X(1)) = 0$  for  $0 \leq p \leq a - 1$  when  $\text{char}(\mathbb{k}) = 0$  or  $\text{char}(\mathbb{k}) \geq a$ . This implies the canonical ribbon conjecture and hence the generic Green conjecture (see [loc. cit., Section 6]). However, for settling Eisenbud and Schreyer conjecture, we only need to consider the case  $a = \lfloor (g - 1)/2 \rfloor$  and  $b = \lfloor g/2 \rfloor$ , and we recover [loc. cit., Theorem 1.1] for this case here.

**Theorem 1.2.** *Let  $X = X(\lfloor (g - 1)/2 \rfloor, \lfloor g/2 \rfloor) \subseteq \mathbb{P}^g$  be a K3 carpet with  $g \geq 3$  over an algebraically closed field  $\mathbb{k}$  with  $\text{char}(\mathbb{k}) = 0$  or  $\text{char}(\mathbb{k}) \geq \lfloor (g - 1)/2 \rfloor$  and  $\text{char}(\mathbb{k}) \neq 2$ . Then*

$$K_{p,2}(X, \mathcal{O}_X(1)) = 0 \quad \text{for } 0 \leq p \leq \lfloor (g - 3)/2 \rfloor.$$

*In particular, Green's conjecture holds for general curves of genus  $g$  over  $\mathbb{k}$ .*

Recall that Schreyer [1986] verified Green's conjecture for every curve of genus  $g \leq 6$  over an algebraically closed field of arbitrary characteristic. In the theorem, if  $g \geq 7$ , then the condition  $\text{char}(\mathbb{k}) \neq 2$  is redundant. Our proof of Theorem 1.2 is essentially different from that of [Raicu and Sam 2022] but surprisingly the same as that of Theorem 1.1. The key point is that the K3 carpet  $X$  in the theorem is a Weil divisor linearly equivalent to the tangent-developable surface  $T$  on the secant variety  $\Sigma$ . Thus  $X$  is a degeneration of the tangent surface  $T$ . The proof of Theorem 1.1 works for  $X$ , and the characteristic assumption on  $\mathbb{k}$  for (1-2) with another  $W$  can be improved. Consequently, we obtain a very quick proof of the generic Green conjecture and Eisenbud and Schreyer's conjecture.

In view of Theorem 1.1, it is quite natural to study syzygies of tangent-developable surfaces of smooth projective curves of genus  $g \geq 1$ . However, there was no research in this direction to the best of the author's knowledge. As a first step, we show the arithmetic normality, and compute the Castelnuovo–Mumford regularity.

**Theorem 1.3.** *Let  $C$  be a smooth projective curve of genus  $g \geq 1$  over an algebraically closed field of characteristic zero,  $L$  be a line bundle on  $C$  with  $\deg L \geq 4g + 3$ , and  $T$  be the tangent-developable surface of  $C$  embedded in  $\mathbb{P}^r$  by  $|L|$ . Then  $T \subseteq \mathbb{P}^r$  is arithmetically normal but not arithmetically Cohen–Macaulay, and  $H^i(T, \mathcal{O}_T(m)) = 0$  for  $i > 0$ ,  $m > 0$  but  $H^1(T, \mathcal{O}_T) \neq 0$ ,  $H^2(T, \mathcal{O}_T) \neq 0$ . In particular,  $\text{reg } \mathcal{O}_T = 3$ , and  $\text{reg } \mathcal{I}_{T|\mathbb{P}^r} = 4$ .*

To prove the theorem, we develop new techniques based on the methods for secant varieties in [Ein et al. 2020]. Along with our proof of Theorem 1.1, this provides a general framework for syzygies of tangent-developable surfaces. First, we show that the dualizing sheaf  $\omega_T$  is trivial (Proposition 2.5). The

hard part of the proof of Theorem 1.3 is to check the 2-normality of  $T \subseteq \mathbb{P}^r$ , which is turned out to be equivalent to  $H^1(C \times C, (L \boxtimes L)(-3D)) = 0$ , where  $D$  is the diagonal of  $C \times C$ . This cohomology vanishing was established by Bertram, Ein and Lazarsfeld [Bertram et al. 1991] when  $\deg L \geq 4g + 3$ . This degree condition is optimal: If  $\deg L = 4g + 2$ , then  $T \subseteq \mathbb{P}^r$  is arithmetically normal if and only if  $C$  is neither elliptic nor hyperelliptic (see Remark 4.4). We will discuss some conjectures on syzygies of tangent-developable surfaces at the end of Section 4.

After setting notation and presenting basic facts in Section 2, we prove Theorems 1.1 and 1.2 in Section 3 and Theorem 1.3 in Section 4. Throughout the paper, we work over an algebraically closed field  $\mathbb{k}$  of arbitrary characteristic unless otherwise stated.

## 2. Preliminaries

**2.1. Syzygies.** Let  $X$  be a projective scheme,  $B$  be a coherent sheaf on  $X$ , and  $L$  be a very ample line bundle on  $X$ . Put  $V := H^0(X, L)$ . The Koszul cohomology  $K_{p,q}(X, B; L)$  is the cohomology of the Koszul-type complex

$$\wedge^{p+1} V \otimes H^0(X, B \otimes L^{q-1}) \longrightarrow \wedge^p V \otimes H^0(X, B \otimes L^q) \longrightarrow \wedge^{p-1} V \otimes H^0(X, B \otimes L^{q+1}).$$

When  $B = \mathcal{O}_X$ , we put  $K_{p,q}(X, L) := K_{p,q}(X, \mathcal{O}_X; L)$  and  $\kappa_{p,q}(X, L) := \dim_{\mathbb{k}} K_{p,q}(X, L)$ . Let  $S := \bigoplus_{m \geq 0} S^m V$ . Then the graded  $S$ -module  $R(X, B; L) := \bigoplus_{m \in \mathbb{Z}} H^0(X, B \otimes L^m)$  admits a minimal free resolution

$$0 \longleftarrow R(X, B; L) \longleftarrow E_0 \longleftarrow E_1 \longleftarrow \dots \longleftarrow E_r \longleftarrow 0,$$

where  $E_p = \bigoplus_q K_{p,q}(X, B; L) \otimes_{\mathbb{k}} S(-p-q)$ . We may think that  $K_{p,q}(X, B; L)$  is the space of  $p$ -th syzygies of weight  $q$ . For a globally generated vector bundle  $E$  on  $X$ , we denote by  $M_E$  the kernel of the evaluation map  $H^0(X, E) \otimes \mathcal{O}_X \rightarrow E$ . The following is well known.

**Proposition 2.1** (see [Park 2022, Proposition 2.1]). *Assume that  $H^i(X, B \otimes L^m) = 0$  for  $i > 0$  and  $m > 0$ . For  $q \geq 2$ , we have  $K_{p,q}(X, B; L) = H^{q-1}(X, \wedge^{p+q-1} M_L \otimes B \otimes L)$ . If furthermore  $H^{q-1}(X, B) = H^q(X, B) = 0$ , then  $K_{p,q}(X, B; L) = H^q(X, \wedge^{p+q} M_L \otimes B)$ .*

**2.2. Castelnuovo–Mumford regularity.** A coherent sheaf  $\mathcal{F}$  on  $\mathbb{P}^n$  is said to be  $m$ -regular if

$$H^i(\mathbb{P}^n, \mathcal{F}(m-i)) = 0$$

for  $i > 0$ . By Mumford’s theorem [Lazarsfeld 2004, Theorem 1.8.3], if  $\mathcal{F}$  is  $m$ -regular, then  $\mathcal{F}$  is  $(m+1)$ -regular. The Castelnuovo–Mumford regularity  $\text{reg } \mathcal{F}$  is the minimum  $m$  such that  $\mathcal{F}$  is  $m$ -regular. If  $\mathcal{F}$  fits into an exact sequence  $\dots \rightarrow \mathcal{F}_2 \rightarrow \mathcal{F}_1 \rightarrow \mathcal{F}_0 \rightarrow \mathcal{F} \rightarrow 0$  of coherent sheaves on  $\mathbb{P}^n$  and  $\mathcal{F}_i$  is  $(m+i)$ -regular for each  $i \geq 0$ , then  $\mathcal{F}$  is  $m$ -regular [loc. cit., Example 1.8.7]. If  $\mathcal{F}$  is  $m$ -regular and  $E$  is an  $m'$ -regular vector bundle on  $\mathbb{P}^n$ , then  $\mathcal{F} \otimes E$  is  $(m+m')$ -regular [loc. cit., Proposition 1.8.9]. We can think of the regularity of a coherent sheaf  $\mathcal{G}$  on a closed subscheme  $X \subseteq \mathbb{P}^n$  by regarding  $\mathcal{G}$  as a sheaf on  $\mathbb{P}^n$ .

**2.3. Multilinear algebra.** Let  $V$  be a finite-dimensional vector space over  $\mathbb{k}$ . The symmetric group  $\mathfrak{S}_n$  naturally acts on  $V^{\otimes n}$  by permuting the factors. The *divided power*  $D^n V$  is the subspace

$$\{\omega \in V^{\otimes n} \mid \sigma(\omega) = \omega \text{ for all } \sigma \in \mathfrak{S}_n\} \subseteq T^n V,$$

and the *symmetric power*  $S^n V$  is the quotient of  $V^{\otimes n}$  by the span of  $\sigma(\omega) - \omega$  for all  $\omega \in V^{\otimes n}$  and  $\sigma \in \mathfrak{S}_n$ . We have a natural identification  $D^n V = (S^n V^\vee)^\vee$ . By composing the inclusion of  $D^n V$  into  $V^{\otimes n}$  with the projection onto  $S^n V$ , we get a natural map  $D^n V \rightarrow S^n V$ . This is an isomorphism if  $\text{char}(\mathbb{k}) = 0$  or  $\text{char}(\mathbb{k}) > n$ , but it may be neither injective nor surjective in general. The *wedge product*  $\wedge^n V$  is the quotient of  $V^{\otimes n}$  by the span of  $v_1 \otimes \cdots \otimes v_n$  for all  $v_1, \dots, v_n \in V$  with  $v_i = v_j$  for some  $i \neq j$ . We write  $v_1 \wedge \cdots \wedge v_n$  for the class of  $v_1 \otimes \cdots \otimes v_n$  in the quotient. There is a natural inclusion  $\wedge^n V \rightarrow V^{\otimes n}$  given by  $v_1 \wedge \cdots \wedge v_n \mapsto \sum_{\sigma \in \mathfrak{S}_n} \text{sgn}(\sigma) \sigma(v_1 \otimes \cdots \otimes v_n)$ . This gives a splitting of the quotient map  $V^{\otimes n} \rightarrow \wedge^n V$  if and only if  $\text{char}(\mathbb{k}) = 0$  or  $\text{char}(\mathbb{k}) > n$ . We refer to [Aprodu et al. 2019, Section 3] for more details.

**2.4. Projective spaces.** Throughout the paper, we put  $U := H^0(\mathbb{P}^1, \mathcal{O}_{\mathbb{P}^1}(1))$ , and fix a basis  $1, x$  of  $U$ . The monomials  $1, x, \dots, x^d$  form a basis of  $S^d U$ , and the divided power monomials  $x^{(0)}, x^{(1)}, \dots, x^{(d)}$  form a basis of  $D^d U$ . Let  $1, y$  be the dual basis of  $U^\vee$  to  $1, x$ . There is a natural identification  $U^\vee = \wedge^2 U \otimes U^\vee = U$  sending  $1, y$  to  $-x, 1$ . Then  $H^0(\mathbb{P}^1, \mathcal{O}_{\mathbb{P}^1}(d)) = S^d U$  and  $H^1(\mathbb{P}^1, \mathcal{O}_{\mathbb{P}^1}(-d-2)) = D^d U$ . Note that  $M_{\mathcal{O}_{\mathbb{P}^1}(d)} = S^{d-1} U \otimes \mathcal{O}_{\mathbb{P}^1}(-1)$ .

We may regard  $\mathbb{P}^n$  as the symmetric product of  $\mathbb{P}^1$ . By permuting the components,  $\mathfrak{S}_n$  acts on the ordinary product  $(\mathbb{P}^1)^n$ , and the line bundle  $\mathcal{O}_{\mathbb{P}^1}(d)^{\boxtimes n}$  on  $(\mathbb{P}^1)^n$  descends to a line bundle  $T_{n, \mathcal{O}_{\mathbb{P}^1}(d)} = \mathcal{O}_{\mathbb{P}^n}(d)$  on  $\mathbb{P}^n$  in such a way that  $q_n^* T_{n, \mathcal{O}_{\mathbb{P}^1}(d)} = \mathcal{O}_{\mathbb{P}^1}(d)^{\boxtimes n}$ , where  $q_n : (\mathbb{P}^1)^n \rightarrow \mathbb{P}^n$  is the quotient map. Then  $H^0(\mathbb{P}^n, \mathcal{O}_{\mathbb{P}^n}(d)) = D^n S^d U$ . Since  $H^0(\mathbb{P}^n, \mathcal{O}_{\mathbb{P}^n}(1)) = D^n U$ , we get  $H^0(\mathbb{P}^n, \mathcal{O}_{\mathbb{P}^n}(d)) = S^d D^n U$ . This gives *Hermite reciprocity*  $D^n S^d U = S^d D^n U$ . If the action of  $\mathfrak{S}_n$  on  $(\mathbb{P}^1)^n$  is alternating, then  $\mathcal{O}_{\mathbb{P}^1}(d)^{\boxtimes n}$  descends to  $N_{n, \mathcal{O}_{\mathbb{P}^1}(d)} = \mathcal{O}_{\mathbb{P}^n}(d-n+1)$  and  $H^0(\mathbb{P}^n, \mathcal{O}_{\mathbb{P}^n}(d-n+1)) = \wedge^n S^d U$ . This gives another *Hermite reciprocity*

$$\wedge^n S^d U = S^{d-n+1} D^n U. \tag{2-1}$$

See [Aprodu et al. 2019, Remark 3.2] and [Park 2022, Subsection 2.3]. Our Hermite reciprocity coincides with the explicit map constructed in [Aprodu et al. 2019, Section 3] (see [Raicu and Sam 2021]), but we will not use this fact.

Let  $D_n$  be the image of the injective map  $\mathbb{P}^{n-1} \times \mathbb{P}^1 \rightarrow \mathbb{P}^n \times \mathbb{P}^1$  given by  $(\xi, z) \mapsto (\xi + z, z)$ . Note that the effective divisor  $D_n$  on  $\mathbb{P}^n \times \mathbb{P}^1$  is defined by

$$\sum_{i=0}^n (-1)^i (x^0 \wedge \cdots \wedge \widehat{x^i} \wedge \cdots \wedge x^n) \otimes x^i \in \wedge^n S^n U \otimes S^n U = H^0(\mathbb{P}^n \times \mathbb{P}^1, \mathcal{O}_{\mathbb{P}^n}(1) \boxtimes \mathcal{O}_{\mathbb{P}^1}(n)).$$

Consider the short exact sequence

$$0 \longrightarrow \mathcal{O}_{\mathbb{P}^n}(d-1) \boxtimes \mathcal{O}_{\mathbb{P}^1}(-n) \xrightarrow{\cdot D_n} \mathcal{O}_{\mathbb{P}^n}(d) \boxtimes \mathcal{O}_{\mathbb{P}^1} \longrightarrow \mathcal{O}_{\mathbb{P}^{n-1}}(d) \boxtimes \mathcal{O}_{\mathbb{P}^1}(d) \longrightarrow 0. \tag{2-2}$$

Pushing forward to  $\mathbb{P}^1$  yields a short exact sequence

$$0 \longrightarrow \wedge^n M_{\mathcal{O}_{\mathbb{P}^1}(d+n-1)} \longrightarrow \wedge^n S^{d+n-1} U \otimes \mathcal{O}_{\mathbb{P}^1} \longrightarrow \wedge^{n-1} M_{\mathcal{O}_{\mathbb{P}^1}(d+n-1)} \otimes \mathcal{O}_{\mathbb{P}^1}(d+n-1) \longrightarrow 0.$$

On the other hand,  $D_n$  can be also defined by

$$\sum_{i=0}^n (-1)^i x^{(n-i)} \otimes x^i \in D^n U \otimes S^n U = H^0(\mathbb{P}^n \times \mathbb{P}^1, \mathcal{O}_{\mathbb{P}^n}(1) \boxtimes \mathcal{O}_{\mathbb{P}^1}(n)).$$

Then we see that the map

$$\begin{array}{ccc} H^1(\mathbb{P}^n \times \mathbb{P}^1, \mathcal{O}_{\mathbb{P}^n}(d-1) \boxtimes \omega_{\mathbb{P}^1}(-n)) & \xrightarrow{D_n} & H^1(\mathbb{P}^n \times \mathbb{P}^1, \mathcal{O}_{\mathbb{P}^n}(d) \boxtimes \omega_{\mathbb{P}^1}) \\ \parallel & & \parallel \\ S^{d-1} D^n U \otimes D^n U & & S^d D^n U \end{array} \tag{2-3}$$

is given by  $f \otimes x^{(i)} \mapsto (-1)^n f x^{(i)}$ . We simply regard this as the multiplication map.

**2.5. Secant varieties.** We recall the set-up of [Ein et al. 2020; 2021], and we present some preliminary results. Let  $C$  be a smooth projective curve of genus  $g \geq 0$ , and  $L$  be a line bundle on  $C$  with  $\deg L \geq 2g + 3$ .<sup>2</sup> We assume  $\text{char}(\mathbb{k}) = 0$  whenever  $g \geq 1$ . We denote by  $C^2 = C \times C$  the ordinary product of  $C$  and by  $C_2 = C^2/\mathfrak{S}_2$  the symmetric product of  $C$ . The quotient map  $\sigma : C^2 \rightarrow C_2$  is given by  $(x, y) \mapsto x + y$ . If we regard  $C_2$  as the Hilbert scheme of two points on  $C$ , then  $\sigma$  is the universal family. For any line bundle  $A$  on  $C$ , there is a line bundle  $T_A$  on  $C_2$  such that  $\sigma^* T_A = A \boxtimes A$ . Let  $D$  be the diagonal of  $C^2$ , and  $Q := \sigma(D)$ . Then  $Q \cong C$  unless  $g = 0$  and  $\text{char}(\mathbb{k}) = 2$ . We may write  $Q = 2\delta$  for some divisor  $\delta$  on  $C_2$ . Note that  $\sigma^* \delta = D$  and  $\sigma_* \mathcal{O}_{C^2} = \mathcal{O}_{C_2}(-\delta) \oplus \mathcal{O}_C$  (see [Ein et al. 2020, Lemma 3.5]). We can write  $K_{C_2} = T_{K_C}(-\delta)$ . Consider the *tautological bundle*  $E := \sigma_*(\mathcal{O}_C \boxtimes L)$  on  $C_2$ . We have  $\text{rank } E = 2$  and  $\det E = T_L(-\delta)$ . Let  $B := \mathbb{P}(E)$ , and  $\pi : B \rightarrow C_2$  be the canonical fibration. As  $H^0(C_2, E) = H^0(C, L)$  and  $E$  is globally generated,  $|\mathcal{O}_{\mathbb{P}(E)}(1)|$  induces a map  $\beta : B \rightarrow \mathbb{P}^r = \mathbb{P}H^0(C, L)$ . Then  $\Sigma := \beta(B)$  is the *secant variety* of  $C$  in  $\mathbb{P}^r$ , and  $\beta : B \rightarrow \Sigma$  is the blow-up of  $\Sigma$  along  $C$  (see [Ein et al. 2021, Theorem 1.1]). Unless  $g = 0$  and  $\deg L = 3$  (in this case  $\Sigma = \mathbb{P}^3$ ),  $\text{Sing } \Sigma = C$  and  $\beta : B \rightarrow \Sigma$  is a resolution of singularities. Let  $Z := \beta^{-1}(C) \cong C^2$ . Then  $\pi|_Z : Z \rightarrow C_2$  is just  $\sigma$ , and  $\beta|_Z : Z = C \times C \rightarrow C$  is the second projection:

$$\begin{array}{ccc} Z = C \times C & \xrightarrow{\quad} & B = \mathbb{P}(E) \xrightarrow{\beta} \Sigma \subseteq \mathbb{P}^r \\ & \searrow \sigma & \downarrow \pi \\ & & C_2 \end{array}$$

**Theorem 2.2** [Ein et al. 2020, Theorems 1.1 and 1.2].  $\Sigma \subseteq \mathbb{P}^r$  is arithmetically Cohen–Macaulay. If  $g = 0$ , then  $\Sigma$  has rational singularities and  $\text{reg } \mathcal{O}_\Sigma = 2$ . If  $g \geq 1$  and  $\text{char}(\mathbb{k}) = 0$ , then  $\Sigma$  has normal Du Bois singularities and  $\text{reg } \mathcal{O}_\Sigma = 4$ .

Pick  $H \in |\mathcal{O}_{\mathbb{P}(E)}(1)|$ . We may write  $K_B = -2H + \pi^* T_{K_C+L}(-2\delta)$  and  $Z = 2H - \pi^* T_L(-2\delta)$ . Take  $\bar{S} \in |2\delta|$ . Let  $\tilde{S} := \pi^{-1}(\bar{S}) = \mathbb{P}(E|_{\bar{S}})$ , and  $S := \beta(\tilde{S})$ . We are mostly interested in the case  $\bar{S} = Q$ . In fact,  $Q$  is a unique member in  $|2\delta|$  when  $g \geq 2$ . Note that  $\dim |2\delta| = 5$  when  $g = 0$  and  $\dim |2\delta| = 1$  when  $g = 1$ . Assume that  $\text{char}(\mathbb{k}) \neq 2$  when  $g = 0$ . Note that  $M_E|_Q = N_{C|\mathbb{P}^r}^\vee \otimes L$  and  $E|_Q = \mathcal{O}^1(L)$  is the first

<sup>2</sup>It is assumed that  $\text{char}(\mathbb{k}) = 0$  in [Ein et al. 2020; 2021], but everything works when  $g = 0$  and  $\text{char}(\mathbb{k}) \geq 0$  (see also [Raicu and Sam 2021]).

jet bundle. By [Kaji 1986, Corollary 1.18],  $\mathcal{P}^1(L)$  is the unique nontrivial extension of  $L$  by  $\omega_C \otimes L$  when  $\text{char}(\mathbb{k}) \nmid \deg L$ . Let  $\tilde{T} := \pi^{-1}(Q)$ . Then  $T := \beta(\tilde{T})$  is the *tangent-developable surface* of  $C$  in  $\mathbb{P}^r$ , and  $T$  has cuspidal singularities along  $C$ . Note that  $\nu := \beta|_{\tilde{T}} : \tilde{T} \rightarrow T$  is a resolution of singularities and  $T$  is Cohen–Macaulay.

**Proposition 2.3.**  $\deg S = 2 \deg L + 2g - 2$ .

*Proof.* We have  $\deg S = (H|_{\tilde{S}})^2 = (\det E) \cdot \bar{S} = 2 \deg L + 2g - 2$ . □

Recall from [Ein et al. 2020, Theorem 5.2] that  $\beta_* \mathcal{O}_B(-Z) = \mathcal{I}_{C|\Sigma}$  and  $R^1 \beta_* \mathcal{O}_B(-Z) = 0$ .

**Lemma 2.4.**  $\beta_* \mathcal{O}_B(-\tilde{S} - Z) = \mathcal{I}_{S|\Sigma}$  and  $R^1 \beta_* \mathcal{O}_B(-\tilde{S} - Z) = 0$ .

*Proof.* As  $\beta_* \mathcal{O}_Z(-\tilde{S}) = 0$ , we have  $\beta_* \mathcal{O}_B(-\tilde{S} - Z) = \beta_* \mathcal{O}_B(-\tilde{S}) = \mathcal{I}_{S|\Sigma}$ . For the second assertion, following [loc. cit., proof of Theorem 5.2(2)], we show that  $R^1 \beta_* \mathcal{O}_B(-\tilde{S} - Z)_x = 0$  for any  $x \in C \subseteq \Sigma$ . Let  $F := \beta^{-1}(x) \cong C$ . By the formal function theorem, it suffices to prove that

$$H^1(F, \mathcal{O}_B(-\tilde{S} - Z) \otimes \mathcal{O}_B / \mathcal{I}_{F|B}^m) = 0 \quad \text{for } m \geq 1.$$

It is enough to check that

$$H^1(F, \mathcal{O}_B(-\tilde{S} - Z) \otimes \mathcal{I}_{F|B}^m / \mathcal{I}_{F|B}^{m+1}) = 0 \quad \text{for } m \geq 0.$$

As  $(-\tilde{S} - Z)|_Z = (L \boxtimes -L)(-4D)$  and  $F$  is a fiber of the second projection  $Z = C \times C \rightarrow C$ , we have  $\mathcal{O}_B(-\tilde{S} - Z)|_F = L(-4x)$ . Note that  $\mathcal{I}_{F|B}^m / \mathcal{I}_{F|B}^{m+1} = S^m N_{F|B}^\vee$ . Recall from [Ein et al. 2020, Proposition 3.13] that  $N_{F|B}^\vee = \mathcal{O}_C \oplus L(-2x)$ . The problem is then reduced to verifying that

$$H^1(C, (m + 1)L + (-4 - 2m)x) = 0 \quad \text{for } m \geq 0.$$

This vanishing holds since  $\deg((m + 1)L + (-4 - 2m)x) \geq (m + 1)(2g + 3) - 4 - 2m \geq 2g - 1$ . □

Consider the following commutative diagram with short exact sequences:

$$\begin{array}{ccccccc}
 & & & 0 & & 0 & \\
 & & & \downarrow & & \downarrow & \\
 0 & \longrightarrow & \mathcal{O}_B(-\tilde{S} - Z) & \xrightarrow{\cdot Z} & \mathcal{O}_B(-\tilde{S}) & \longrightarrow & \mathcal{O}_Z(-2D) \longrightarrow 0 \\
 & & \parallel & & \downarrow \cdot \tilde{S} & & \downarrow \\
 0 & \longrightarrow & \mathcal{O}_B(-\tilde{S} - Z) & \xrightarrow{\cdot \tilde{S} + Z} & \mathcal{O}_B & \longrightarrow & \mathcal{O}_{\tilde{S} + Z} \longrightarrow 0 \\
 & & & & \downarrow & & \downarrow \\
 & & & & \mathcal{O}_{\tilde{S}} & \xlongequal{\quad} & \mathcal{O}_{\tilde{S}} \\
 & & & & \downarrow & & \downarrow \\
 & & & & 0 & & 0
 \end{array} \tag{2-4}$$

By Lemma 2.4, applying  $\beta_*$  to the second-row exact sequence in (2-4), we find  $\beta_* \mathcal{O}_{\tilde{S} + Z} = \mathcal{O}_S$  and  $R^1 \beta_* \mathcal{O}_{\tilde{S} + Z} = R^1 \beta_* \mathcal{O}_B = H^1(C, \mathcal{O}_C) \otimes \mathcal{O}_C$ , and we get a short exact sequence

$$0 \longrightarrow \beta_* \mathcal{O}_B(-\tilde{S} - Z) \longrightarrow \mathcal{O}_\Sigma \longrightarrow \mathcal{O}_S \longrightarrow 0. \tag{2-5}$$

Note that

$$R^1 \beta_* \mathcal{O}_{\tilde{S}+Z} = R^1 \beta_* \mathcal{O}_B = R^1 (\beta|_Z)_* \mathcal{O}_Z = R^1 (\beta|_Z)_* \mathcal{O}_Z(-D).$$

The kernel of the map  $R^1 (\beta|_Z)_* \mathcal{O}_Z(-2D) \rightarrow R^1 (\beta|_Z)_* \mathcal{O}_Z(-D)$  is  $\mathcal{O}_D(-D|_D) = \omega_C$ . By applying  $\beta_*$  to the rightmost vertical exact sequence in (2-4), we get a short exact sequence

$$0 \longrightarrow \mathcal{O}_S \longrightarrow \beta_* \mathcal{O}_{\tilde{S}} \longrightarrow \omega_C \longrightarrow 0. \tag{2-6}$$

**Proposition 2.5.** *The dualizing sheaf  $\omega_S$  is trivial.*

*Proof.* Consider two short exact sequences

$$0 \longrightarrow \omega_B \longrightarrow \omega_B(Z) \longrightarrow \omega_Z \longrightarrow 0 \quad \text{and} \quad 0 \longrightarrow \omega_B \longrightarrow \omega_B(\tilde{S}) \longrightarrow \omega_{\tilde{S}} \longrightarrow 0. \tag{2-7}$$

By Theorem 2.2 for  $g = 0$  and Grauert–Riemenschneider vanishing for  $g \geq 1$ , we have  $R^1 \beta_* \omega_B = 0$ . Then  $R^1 \beta_* \omega_B(Z) = R^1 \beta_* \omega_Z = \omega_C$  and  $R^1 \beta_* \omega_B(\tilde{S}) = 0$ . By taking  $-\otimes \mathcal{O}_B(\tilde{S})$  to the left of (2-7), we see that  $R^1 \beta_* \omega_B(\tilde{S} + Z) = 0$ . When  $g = 0$ , we have  $\beta_* \omega_B(Z) = \beta_* \omega_B = \omega_\Sigma$  by Theorem 2.2. When  $g \geq 1$ , Theorem 2.2 and [Kovács et al. 2010, Theorem 1.1] show that  $\beta_* \omega_B(Z) = \omega_\Sigma$ . As  $\omega_{\tilde{S}}(Z|_{\tilde{S}}) = \mathcal{O}_{\tilde{S}}$ , applying  $\beta_*$  to the short exact sequence with the consideration of (2-6),

$$0 \longrightarrow \omega_B(Z) \xrightarrow{\cdot \tilde{S}} \omega_B(\tilde{S} + Z) \longrightarrow \mathcal{O}_{\tilde{S}} \longrightarrow 0,$$

we obtain a short exact sequence

$$0 \longrightarrow \omega_\Sigma \longrightarrow \beta_* \omega_B(\tilde{S} + Z) \longrightarrow \mathcal{O}_S \longrightarrow 0.$$

On the other hand, applying  $\mathcal{H}om_\Sigma(-, \omega_\Sigma)$  to (2-5), we get a short exact sequence

$$0 \longrightarrow \omega_\Sigma \longrightarrow \mathcal{H}om_\Sigma(\beta_* \mathcal{O}_B(-\tilde{S} - Z), \omega_\Sigma) \longrightarrow \omega_S \longrightarrow 0$$

since  $\Sigma$  is Cohen–Macaulay. Notice that

$$\mathcal{H}om_\Sigma(\beta_* \mathcal{O}_B(-\tilde{S} - Z), \omega_\Sigma) = \mathcal{H}om_\Sigma(\beta_* \mathcal{O}_B(-\tilde{S}), \omega_\Sigma) = \beta_* \omega_B(\tilde{S} + Z).$$

Hence we conclude that  $\omega_S = \mathcal{O}_S$ . □

**Remark 2.6.** When  $g = 0$  and  $S = T$  is the tangent-developable surface, the above proposition was shown in [Aprodu et al. 2019, Corollary 5.16]. When  $g = 1$  and  $\deg L = 5$ , applying the Serre construction to the tangent-developable surface  $T$  in  $\mathbb{P}^4$ , Hulek [1986, Chapter VII] gave a new construction of the Horrocks–Mumford bundle on  $\mathbb{P}^4$ .

**2.6. Rational curve case.** Assume that  $\text{char}(\mathbb{k}) \neq 2$ . Let  $C \subseteq \mathbb{P}^g$  be a rational normal curve of degree  $g \geq 3$ , and  $L := \mathcal{O}_{\mathbb{P}^1}(g)$ . When we consider a rational normal curve,  $g$  is not the genus but the degree. Note that  $C_2 = \mathbb{P}^2$  and  $\mathcal{O}_{\mathbb{P}^2}(\delta) = \mathcal{O}_{\mathbb{P}^2}(1)$ . We have  $T_{\mathcal{O}_{\mathbb{P}^1}(d)} = \mathcal{O}_{\mathbb{P}^2}(d)$ , and put  $N_{\mathcal{O}_{\mathbb{P}^1}(d)} := T_{\mathcal{O}_{\mathbb{P}^1}(d)}(-\delta) = \mathcal{O}_{\mathbb{P}^2}(d - 1)$ .

Then  $\sigma^*T_{\mathcal{O}_{\mathbb{P}^1}(d)} = \mathcal{O}_{\mathbb{P}^1}(d) \boxtimes \mathcal{O}_{\mathbb{P}^1}(d)$  and  $\sigma^*N_{\mathcal{O}_{\mathbb{P}^1}(d)} = \mathcal{O}_{\mathbb{P}^1}(d-1) \boxtimes \mathcal{O}_{\mathbb{P}^1}(d-1)$ . Note that  $\pi_*\mathcal{O}_B(-Z) = 0$  and  $R^1\pi_*\mathcal{O}_B(-Z) = \mathcal{O}_{\mathbb{P}^2}(-\delta)$ . By applying  $\pi_*$  to the short exact sequence

$$0 \longrightarrow \mathcal{O}_B(-Z) \xrightarrow{\cdot Z} \mathcal{O}_B \longrightarrow \mathcal{O}_Z \longrightarrow 0, \tag{2-8}$$

we get a splitting short exact sequence

$$0 \longrightarrow \mathcal{O}_{\mathbb{P}^2} \longrightarrow \sigma_*\mathcal{O}_Z \longrightarrow \mathcal{O}_{\mathbb{P}^2}(-\delta) \longrightarrow 0.$$

Then we obtain the following canonically splitting short exact sequence:

$$\begin{array}{ccccccc} 0 & \longrightarrow & H^0(\mathbb{P}^2, T_{\mathcal{O}_{\mathbb{P}^1}(d)}) & \longrightarrow & H^0(\mathbb{P}^1 \times \mathbb{P}^1, \mathcal{O}_{\mathbb{P}^1}(d) \boxtimes \mathcal{O}_{\mathbb{P}^1}(d)) & \longrightarrow & H^0(\mathbb{P}^2, N_{\mathcal{O}_{\mathbb{P}^1}(d)}) \longrightarrow 0 \\ & & \parallel & & \parallel & & \parallel \\ & & D^2S^dU & & S^dU \otimes S^dU & & \wedge^2S^dU \end{array} \tag{2-9}$$

**Lemma 2.7.**  $M_E = S^{g-2}U \otimes \mathcal{O}_{\mathbb{P}^2}(-1)$ .

*Proof.* Recall that there is an injective map  $D_2 = \mathbb{P}^1 \times \mathbb{P}^1 \rightarrow \mathbb{P}^2 \times \mathbb{P}^1$ . Let  $p_1 : \mathbb{P}^2 \times \mathbb{P}^1 \rightarrow \mathbb{P}^2$  be the first projection. Then  $(p_1)|_{D_2} = \sigma$ . By applying  $p_{1,*}$  to the short exact sequence

$$0 \longrightarrow \mathcal{O}_{\mathbb{P}^2}(-1) \boxtimes \mathcal{O}_{\mathbb{P}^1}(g-2) \xrightarrow{\cdot D_2} \mathcal{O}_{\mathbb{P}^2} \boxtimes \mathcal{O}_{\mathbb{P}^1}(g) \longrightarrow \mathcal{O}_{\mathbb{P}^1} \boxtimes \mathcal{O}_{\mathbb{P}^1}(g) \longrightarrow 0,$$

we get a short exact sequence

$$0 \longrightarrow S^{g-2}U \otimes \mathcal{O}_{\mathbb{P}^2}(-1) \longrightarrow S^gU \otimes \mathcal{O}_{\mathbb{P}^2} \longrightarrow E \longrightarrow 0,$$

where  $H^0(\mathbb{P}^2, E) \otimes \mathcal{O}_{\mathbb{P}^2} = S^gU \otimes \mathcal{O}_{\mathbb{P}^2} \rightarrow E$  is the evaluation map. Thus the lemma follows. □

**Proposition 2.8.** (1)  $K_B + \tilde{S} + Z = 0$ .

(2)  $S \subseteq \mathbb{P}^g$  is arithmetically Cohen–Macaulay and  $\text{reg } \mathcal{O}_S = 3$ .

(3) The Hilbert function of  $S \subseteq \mathbb{P}^g$  is given by  $H_S(t) = (g-1)t^2 + 2$  for  $t \geq 1$ .

*Proof.* The assertion (1) follows from the direct computation

$$K_B + \tilde{S} + Z = (-2H + \pi^*\mathcal{O}_{\mathbb{P}^2}(g-4)) + \pi^*\mathcal{O}_{\mathbb{P}^2}(2) + (2H - \pi^*\mathcal{O}_{\mathbb{P}^2}(g-2)) = 0.$$

As  $\beta_*\mathcal{O}_B(-\tilde{S} - Z) = \beta_*\omega_B = \omega_\Sigma$  by Theorem 2.2, the short exact sequence (2-5) is

$$0 \longrightarrow \omega_\Sigma \longrightarrow \mathcal{O}_\Sigma \longrightarrow \mathcal{O}_S \longrightarrow 0. \tag{2-10}$$

By Theorem 2.2, we readily obtain (2). Now,  $H_T(t)$  agrees with the Hilbert polynomial  $P_S(t)$  for  $t \geq 1$ . Note that  $\det P_S(t) = 2$  and the leading coefficient of  $P_T(t)$  is  $(\deg S)/2 = g-1$ . As  $P_S(0) = \chi(\mathcal{O}_S) = 2$  and  $P_S(1) = g+1$ , we get (3). □

When  $S = T$  is the tangent-developable surface, Proposition 2.8(2) was first shown by Schreyer [1983, Proposition 6.1] (see also [Aprodu et al. 2019, Theorem 5.1]). Observe that  $S \subseteq \mathbb{P}^g$  has the same Hilbert polynomial with a K3 surface of degree  $2g-2$  in  $\mathbb{P}^g$ .

Suppose that  $\bar{S} = Q$ . Here  $Q = \sigma(D)$  is a smooth conic in  $\mathbb{P}^2$ . When  $\text{char}(\mathbb{k}) = 0$  or  $\text{char}(\mathbb{k}) \geq (g+2)/2$ , [Kaji 1986, Corollary 1.18] implies that  $E|_Q = \mathcal{O}_{\mathbb{P}^1}(g-1) \oplus \mathcal{O}_{\mathbb{P}^1}(g-1)$ . In this case,  $\tilde{T} = \mathbb{P}^1 \times \mathbb{P}^1$ , and  $\mathcal{O}_{\tilde{T}}(H|\tilde{T}) = \mathcal{O}_{\mathbb{P}^1}(g-1) \boxtimes \mathcal{O}_{\mathbb{P}^1}(1)$ . When  $\text{char}(\mathbb{k}) \mid g$ , [loc. cit., Corollary 1.18] implies that  $\tilde{T} \neq \mathbb{P}^1 \times \mathbb{P}^1$ . However, we always have  $Z = \mathbb{P}^1 \times \mathbb{P}^1$ . Now, consider the following short exact sequence:

$$\begin{array}{ccccccc}
 0 & \longrightarrow & H^0(\mathbb{P}^2, \mathcal{O}_{\mathbb{P}^2}(d-1)) & \xrightarrow{\cdot Q} & H^0(\mathbb{P}^2, \mathcal{O}_{\mathbb{P}^2}(d+1)) & \longrightarrow & H^0(Q, \mathcal{O}_Q(d+1)) \longrightarrow 0 \\
 & & \parallel & & \parallel & & \parallel \\
 & & \wedge^2 S^d U & & \wedge^2 S^{d+2} U & & S^{2d+2} U
 \end{array} \tag{2-11}$$

Since  $\sigma^*Q = 2D$  and the diagonal  $D$  of  $\mathbb{P}^1 \times \mathbb{P}^1$  is defined by

$$x \otimes 1 - 1 \otimes x \in U \otimes U = H^0(\mathbb{P}^1 \times \mathbb{P}^1, \mathcal{O}_{\mathbb{P}^1}(1) \boxtimes \mathcal{O}_{\mathbb{P}^1}(1)),$$

the inclusion  $H^0(\mathbb{P}^2, \mathcal{O}_{\mathbb{P}^2}(d-1)) \rightarrow H^0(\mathbb{P}^2, \mathcal{O}_{\mathbb{P}^2}(d+1))$  in (2-11) is

$$\wedge^2 S^d U \longrightarrow \wedge^2 S^{d+2} U, \quad x^i \wedge x^j \longmapsto x^{i+2} \wedge x^j - 2x^{i+1} \wedge x^{j+1} + x^i \wedge x^{j+2}.$$

The surjection  $H^0(\mathbb{P}^2, \mathcal{O}_{\mathbb{P}^2}(d+1)) \rightarrow H^0(Q, \mathcal{O}_Q(d+1))$  in (2-11) is

$$\mu_{d+2} : \wedge^2 S^{d+2} U \longrightarrow S^{2d+2} U, \quad x^i \wedge x^j \longmapsto (i-j)x^{i+j-1}, \tag{2-12}$$

which is the *Wahl map* (or the *Gaussian map*). See [Bertram et al. 1991; Wahl 1990] for details on Wahl maps.

Suppose that  $\bar{S} = 2\ell$ , where  $\ell \subseteq \mathbb{P}^2$  is a line meeting  $Q$  at two distinct points. We may write  $E|_\ell = \mathcal{O}_{\mathbb{P}^1}(a) \oplus \mathcal{O}_{\mathbb{P}^1}(b)$  for some integers  $b \geq a \geq 1$ . As  $\det E = N_L = \mathcal{O}_{\mathbb{P}^2}(g-1)$ , we have  $a+b = g-1$ . Note that  $\sigma^{-1}(\ell) = \mathbb{P}^1$  and  $\sigma|_{\sigma^{-1}(\ell)} : \sigma^{-1}(\ell) \rightarrow \ell$  is a two-to-one map. By restricting the short exact sequence

$$0 \longrightarrow \mathcal{O}_{\mathbb{P}^1}(g-1) \boxtimes \mathcal{O}_{\mathbb{P}^1}(-1) \longrightarrow \sigma^*E \longrightarrow \mathcal{O}_{\mathbb{P}^1} \boxtimes \mathcal{O}_{\mathbb{P}^1}(g) \longrightarrow 0$$

to  $\sigma^{-1}(\ell)$ , we get a short exact sequence

$$0 \longrightarrow \mathcal{O}_{\mathbb{P}^1}(g-2) \longrightarrow \mathcal{O}_{\mathbb{P}^1}(2a) \oplus \mathcal{O}_{\mathbb{P}^1}(2b) \longrightarrow \mathcal{O}_{\mathbb{P}^1}(g) \longrightarrow 0.$$

It follows that  $(a, b) = (\lfloor (g-1)/2 \rfloor, \lfloor g/2 \rfloor)$ . Then  $S$  is a double structure on a rational normal surface scroll  $S(a, b) = \mathbb{P}(E|_\ell) \subseteq \mathbb{P}^g$ . Recall that  $H^1(S, \mathcal{O}_S) = 0$  and  $\omega_S = \mathcal{O}_S$ . Thus  $S = X(\lfloor (g-1)/2 \rfloor, \lfloor g/2 \rfloor) \subseteq \mathbb{P}^g$  is a *K3 carpet* (see [Gallego and Purnaprajna 1997, Definition 1.2 and Theorem 1.3]<sup>3</sup>). On the other hand, consider the following short exact sequence:

$$\begin{array}{ccccccc}
 0 & \longrightarrow & H^0(\mathbb{P}^2, \mathcal{O}_{\mathbb{P}^2}(d-1)) & \xrightarrow{\cdot 2\ell} & H^0(\mathbb{P}^2, \mathcal{O}_{\mathbb{P}^2}(d+1)) & \longrightarrow & H^0(2\ell, \mathcal{O}_{2\ell}(d+1)) \longrightarrow 0 \\
 & & \parallel & & \parallel & & \parallel \\
 & & \wedge^2 S^d U & & \wedge^2 S^{d+2} U & & 
 \end{array} \tag{2-13}$$

<sup>3</sup>The proof of [Gallego and Purnaprajna 1997, Theorem 1.3] also works in positive characteristic.

We may assume that  $\sigma^{-1}(\ell)$  in  $\mathbb{P}^1 \times \mathbb{P}^1$  is defined by

$$x \otimes 1 + 1 \otimes x \in U \otimes U = H^0(\mathbb{P}^1 \times \mathbb{P}^1, \mathcal{O}_{\mathbb{P}^1}(1) \boxtimes \mathcal{O}_{\mathbb{P}^1}(1)).$$

Then the inclusion  $H^0(\mathbb{P}^2, \mathcal{O}_{\mathbb{P}^2}(d-1)) \rightarrow H^0(\mathbb{P}^2, \mathcal{O}_{\mathbb{P}^2}(d+1))$  in (2-13) is

$$\wedge^2 S^d U \longrightarrow \wedge^2 S^{d+2} U, \quad x^i \wedge x^j \longmapsto x^{i+2} \wedge x^j + 2x^{i+1} \wedge x^{j+1} + x^i \wedge x^{j+2}.$$

It is a direct summand of the inclusion

$$H^0(\mathbb{P}^1 \times \mathbb{P}^1, \mathcal{O}_{\mathbb{P}^1}(d) \boxtimes \mathcal{O}_{\mathbb{P}^1}(d)) \xrightarrow{2\sigma^{-1}(\ell)} H^0(\mathbb{P}^1 \times \mathbb{P}^1, \mathcal{O}_{\mathbb{P}^1}(d+2) \boxtimes \mathcal{O}_{\mathbb{P}^1}(d+2))$$

whose cokernel is  $S^{2d+2}U \oplus S^{2d+4}U$ . The surjection  $H^0(\mathbb{P}^2, \mathcal{O}_{\mathbb{P}^2}(d+1)) \rightarrow H^0(2\ell, \mathcal{O}_{2\ell}(d+1))$  in (2-13) factors through

$$\tau_{d+2} : \wedge^2 S^{d+2} U \longrightarrow \begin{matrix} S^{2d+2}U \\ \oplus \\ S^{2d+4}U \end{matrix}, \quad x^i \wedge x^j \longmapsto \begin{cases} ((-1)^i(i-j)x^{i+j-1}, 0) & \text{if } i \equiv j \pmod{2}, \\ (0, (-1)^i x^{i+j}) & \text{if } i \not\equiv j \pmod{2}, \end{cases} \quad (2-14)$$

in such a way that  $\text{im}(\tau_{d+2})$  injects into  $H^0(2\ell, \mathcal{O}_{2\ell}(d+1))$ .

### 3. Proofs of Theorems 1.1 and 1.2

Assume that  $\text{char}(\mathbb{k}) \neq 2$ . Let  $C \subseteq \mathbb{P}^g$  be a rational normal curve of degree  $g \geq 3$ , and  $L := \mathcal{O}_{\mathbb{P}^1}(g)$ . We use the notation in Section 2. From (2-10), we see that

$$K_{p,2}(S, \mathcal{O}_S(1)) = \text{coker}(K_{p,2}(\Sigma, K_\Sigma; \mathcal{O}_\Sigma(1)) \xrightarrow{\rho_{p+2}} K_{p,2}(\Sigma, \mathcal{O}_\Sigma(1))).$$

Thus  $K_{p,2}(S, \mathcal{O}_S(1)) = 0$  if and only if  $\rho_{p+2}$  is surjective. Recall from Theorem 2.2 that  $\Sigma$  has rational singularities. As  $\beta^* M_{\mathcal{O}_\Sigma(1)} = M_H$ , by Proposition 2.1, we find

$$\begin{aligned} K_{p,2}(\Sigma, K_\Sigma; \mathcal{O}_\Sigma(1)) &= H^2(\Sigma, \wedge^{p+2} M_{\mathcal{O}_\Sigma(1)} \otimes \omega_\Sigma) = H^2(B, \wedge^{p+2} M_H \otimes \omega_B), \\ K_{p,2}(\Sigma, \mathcal{O}_\Sigma(1)) &= H^2(\Sigma, \wedge^{p+2} M_{\mathcal{O}_\Sigma(1)}) = H^2(B, \wedge^{p+2} M_H). \end{aligned}$$

In (2-4), we have  $-\tilde{S} - Z = K_B$ ,  $-\tilde{S} = K_B + Z$ ,  $-2D = K_Z$ . Then  $\rho_{p+2}$  fits into the following commutative diagram with exact sequences induced from (2-4):

$$\begin{array}{ccccc} & & H^1(\tilde{S}, \wedge^{p+2} M_H|_{\tilde{S}}) & & \\ & & \downarrow \alpha_{p+2} & \searrow \gamma_{p+2} & \\ H^2(B, \wedge^{p+2} M_H \otimes \omega_B) & \hookrightarrow & H^2(B, \wedge^{p+2} M_H \otimes \omega_B(Z)) & \xrightarrow{\delta_{p+2}} & H^2(Z, \omega_{\mathbb{P}^1} \boxtimes \wedge^{p+2} M_L \otimes \omega_{\mathbb{P}^1}) \\ & \searrow \rho_{p+2} & \downarrow & & \\ & & H^2(B, \wedge^{p+2} M_H) & & \end{array} \quad (3-1)$$

It is easy to check that  $\rho_{p+2}$  is surjective if and only if  $\gamma_{p+2} = \delta_{p+2} \circ \alpha_{p+2}$  surjects onto  $\text{im}(\delta_{p+2})$ . We shall prove the latter for  $0 \leq p \leq \lfloor (g-3)/2 \rfloor$  when  $S = T$  is the tangent-developable surface and

$S = X(\lfloor (g-1)/2 \rfloor, \lfloor g/2 \rfloor)$  is a K3 carpet. To this end, we first describe  $\gamma_{p+2} = \delta_{p+2} \circ \alpha_{p+2}$ . Let  $q := g - p - 3$ ,  $V := D^{p+2}U$ ,  $W := H^1(\bar{S}, \mathcal{O}_{\bar{S}}(-p-2))$ .

**Lemma 3.1.** *The map  $\gamma_{p+2}$  is the composition*

$$S^q V \otimes W \xrightarrow{\alpha_{p+2}} S^q V \otimes \wedge^2 V \xrightarrow{\delta_{p+2}} S^{q+1} V \otimes V$$

such that  $\delta_{p+2}$  is the Koszul differential given by  $f \otimes (x^{(i)} \wedge x^{(j)}) \mapsto f x^{(i)} \otimes x^{(j)} - f x^{(j)} \otimes x^{(i)}$  and  $\alpha_{p+2} = \text{id}_{S^q V} \otimes \Delta_{p+2}$ , where  $\Delta_{p+2}$  fits into the short exact sequence<sup>4</sup>

$$0 \longrightarrow H^1(\bar{S}, \mathcal{O}_{\bar{S}}(-p-2)) \xrightarrow{\Delta_{p+2}} H^2(\mathbb{P}^2, \mathcal{O}_{\mathbb{P}^2}(-p-2)(-\bar{S})) \xrightarrow{\cdot \bar{S}} H^2(\mathbb{P}^2, \mathcal{O}_{\mathbb{P}^2}(-p-2)) \longrightarrow 0.$$

*Proof.* First, we give a description of  $\alpha_{p+2}$ . As  $\pi_* M_H = M_E$ , we have a short exact sequence

$$0 \longrightarrow \pi^* M_E \longrightarrow M_H \longrightarrow \mathcal{O}_B(-H) \otimes \pi^* N_L \longrightarrow 0. \tag{3-2}$$

Then we get  $\pi_* \wedge^{p+2} M_H = \wedge^{p+2} M_E$  and  $R^1 \pi_* \wedge^{p+2} M_H = 0$ . By restricting (3-2) to  $\tilde{S}$ , we also get  $(\pi|_{\tilde{S}})_* \wedge^{p+2} M_H|_{\tilde{S}} = \wedge^{p+2} M_E|_{\tilde{S}}$  and  $R^1(\pi|_{\tilde{S}})_* \wedge^{p+2} M_H|_{\tilde{S}} = 0$ . Recall from Lemma 2.7 that  $M_E = S^{g-2}U \otimes \mathcal{O}_{\mathbb{P}^2}(-1)$ . Thus we find

$$\begin{aligned} H^1(\tilde{S}, \wedge^{p+2} M_H|_{\tilde{S}}) &= H^1(\bar{S}, \wedge^{p+2} M_E|_{\bar{S}}) = \wedge^{p+2} S^{g-2}U \otimes H^1(\bar{S}, \mathcal{O}_{\bar{S}}(-p-2)), \\ H^2(B, \wedge^{p+2} M_H \otimes \omega_B(Z)) &= H^2(\mathbb{P}^2, \wedge^{p+2} M_E \otimes \mathcal{O}_{\mathbb{P}^2}(-\bar{S})) = \wedge^{p+2} S^{g-2}U \otimes H^2(\mathbb{P}^2, \mathcal{O}_{\mathbb{P}^2}(-p-2)(-\bar{S})), \\ H^2(B, \wedge^{p+2} M_H) f &= H^2(\mathbb{P}^2, \wedge^{p+2} M_E) = \wedge^{p+2} S^{g-2}U \otimes H^2(\mathbb{P}^2, \mathcal{O}_{\mathbb{P}^2}(-p-2)). \end{aligned}$$

By considering the vertical short exact sequence in (3-1), we obtain  $\alpha_{p+2} = \text{id}_{\wedge^{p+2} S^{g-2}U} \otimes \Delta_{p+2}$ . By Hermite reciprocity (2-1), we have  $\wedge^{p+2} S^{g-2}U = S^q V$ .

To describe  $\delta_{p+2}$ , we restrict (3-2) to  $Z$  to get a short exact sequence

$$\begin{array}{ccccccc} 0 & \longrightarrow & \sigma^* M_E & \longrightarrow & \mathcal{O}_{\mathbb{P}^1} \boxtimes M_L & \longrightarrow & (\mathcal{O}_{\mathbb{P}^1} \boxtimes -L) \otimes \sigma^* N_L \longrightarrow 0 \\ & & \parallel & & \parallel & & \parallel \\ & & M_{\mathcal{O}_{\mathbb{P}^1}(g-1)} \boxtimes \mathcal{O}_{\mathbb{P}^1}(-1) & & S^{g-1}U \otimes \mathcal{O}_{\mathbb{P}^1} \boxtimes \mathcal{O}_{\mathbb{P}^1}(-1) & & \mathcal{O}_{\mathbb{P}^1}(g-1) \boxtimes \mathcal{O}_{\mathbb{P}^1}(-1) \end{array}$$

where the maps are the identity on  $\mathcal{O}_{\mathbb{P}^1}(-1)$  and the map  $S^{g-1}U \otimes \mathcal{O}_{\mathbb{P}^1} \rightarrow \mathcal{O}_{\mathbb{P}^1}(g-1)$  is the evaluation map. The induced map  $H^2(Z, \sigma^* \wedge^{p+2} M_E \otimes \omega_Z) \rightarrow H^2(Z, \omega_{\mathbb{P}^1} \boxtimes \wedge^{p+2} M_L \otimes \omega_{\mathbb{P}^1})$  can be written as  $\delta'_{p+2} \otimes \text{id}_V$ , where

$$\delta'_{p+2} : H^1(\mathbb{P}^1, \wedge^{p+2} M_{\mathcal{O}_{\mathbb{P}^1}(g-1)} \otimes \omega_{\mathbb{P}^1}) \longrightarrow \wedge^{p+2} S^{g-1}U \otimes H^1(\mathbb{P}^1, \omega_{\mathbb{P}^1})$$

and  $V = D^{p+2}U = H^1(\mathbb{P}^1, \omega_{\mathbb{P}^1}(-p-2))$ . By considering the push-forward of (2-2) to  $\mathbb{P}^1$ , we can identify  $\delta'_{p+2}$  with

$$H^1(\mathbb{P}^{p+2} \times \mathbb{P}^1, \mathcal{O}_{\mathbb{P}^{p+2}}(g-p-3) \boxtimes \omega_{\mathbb{P}^1}(-p-2)) \xrightarrow{\cdot D_{p+2}} H^1(\mathbb{P}^{p+2} \times \mathbb{P}^1, \mathcal{O}_{\mathbb{P}^{p+2}}(g-p-2) \boxtimes \omega_{\mathbb{P}^1}).$$

<sup>4</sup> When  $S = T$ , one can easily confirm that the map  $\gamma$  in the Introduction coincides with the map  $\gamma_{p+2}$  by replacing the range by  $\text{im}(\delta_{p+2}) = K_{p,1}(\mathbb{P}^1, \omega_{\mathbb{P}^1}; \mathcal{O}_{\mathbb{P}^1}(g))$ . In this case,  $\Delta_{p+2}$  can be thought of as the dual of the Wahl map  $\mu_{p+2}$ .

This is the map (2-3) with  $n = p + 2$  and  $d = g - p - 2 = q + 1$ , so  $\delta'_{p+2} : S^q V \otimes V \rightarrow S^{q+1} V$  is the multiplication map, where we use Hermite reciprocity (2-1) to have that

$$\begin{aligned} H^1(\mathbb{P}^1, \wedge^{p+2} M_{\mathcal{O}_{\mathbb{P}^1}(g-1)} \otimes \omega_{\mathbb{P}^1}) &= \wedge^{p+2} S^{g-2} U \otimes D^{p+2} U = S^q V \otimes V, \\ \wedge^{p+2} S^{g-1} U \otimes H^1(\mathbb{P}^1, \omega_{\mathbb{P}^1}) &= \wedge^{p+2} S^{g-1} U = S^{q+1} V. \end{aligned}$$

Now, consider the following commutative diagram with short exact sequences:

$$\begin{array}{ccccccccc} 0 & \longrightarrow & \pi^* M_E \otimes \mathcal{O}_B(-Z) & \xrightarrow{\cdot Z} & \pi^* M_E & \longrightarrow & \sigma^* M_E & \longrightarrow & 0 \\ & & \downarrow & & \downarrow & & \downarrow & & \\ 0 & \longrightarrow & M_H \otimes \mathcal{O}_B(-Z) & \xrightarrow{\cdot Z} & M_H & \longrightarrow & \mathcal{O}_{\mathbb{P}^1} \boxtimes M_L & \longrightarrow & 0 \end{array}$$

The top row is  $(S^{g-2} U \otimes \pi^* \mathcal{O}_{\mathbb{P}^2}(-1)) \otimes (2-8)$ . The diagram induces the following commutative diagram:

$$\begin{array}{ccc} H^2(B, \pi^* \wedge^{p+2} M_E \otimes \omega_B(Z)) & \xrightarrow{\text{id}_{S^q V} \otimes \iota_V} & H^2(Z, \sigma^* \wedge^{p+2} M_E \otimes \omega_Z) \\ \parallel & & \downarrow \delta'_{p+2} \otimes \text{id}_V \\ H^2(B, \wedge^{p+2} M_H \otimes \omega_B(Z)) & \xrightarrow{\delta_{p+2}} & H^2(Z, \omega_{\mathbb{P}^1} \boxtimes \wedge^{p+2} M_L \otimes \omega_{\mathbb{P}^1}) \end{array}$$

Recalling  $\omega_B(Z) = \pi^* \omega_{\mathbb{P}^2}(1)$  and  $\wedge^{p+2} S^{g-2} U = S^q V$ , we have

$$\begin{aligned} H^2(B, \pi^* \wedge^{p+2} M_E \otimes \omega_B(Z)) &= S^q V \otimes H^0(\mathbb{P}^2, \mathcal{O}_{\mathbb{P}^2}(p+1))^\vee, \\ H^2(Z, \sigma^* \wedge^{p+2} M_E \otimes \omega_Z) &= S^q V \otimes H^0(Z, \mathcal{O}_{\mathbb{P}^1}(p+2) \boxtimes \mathcal{O}_{\mathbb{P}^1}(p+2))^\vee. \end{aligned}$$

By considering (2-8), we see that the upper horizontal injection in the above commutative diagram can be written as  $\text{id}_{S^q V} \otimes \iota_V$ , where

$$\iota_V : H^2(\mathbb{P}^2, \omega_{\mathbb{P}^2}(-p-1)) \longrightarrow H^2(Z, (\mathcal{O}_{\mathbb{P}^1}(-p-2) \boxtimes \mathcal{O}_{\mathbb{P}^1}(-p-2)) \otimes \omega_Z)$$

is the dual of the canonical surjection in (2-9) with  $d = p + 2$ . In other words, the map  $\iota_V : \wedge^2 V \rightarrow V \otimes V$  is given by  $\iota_V(x^{(i)} \wedge x^{(j)}) = x^{(i)} \otimes x^{(j)} - x^{(j)} \otimes x^{(i)}$ . Hence we conclude that  $\delta_{p+2} = (\delta'_{p+2} \otimes \text{id}_V) \circ (\text{id}_{S^q V} \otimes \iota_V) : S^q V \otimes \wedge^2 V \rightarrow S^{q+1} V \otimes V$  is the Koszul differential.  $\square$

From now on, we consider  $\gamma_{p+2} : W \otimes S^q V \rightarrow \text{im}(\delta_{p+2})$  (the target  $S^{q+1} V \otimes V$  is replaced by  $\text{im}(\delta_{p+2})$ ). Our aim is to show the surjectivity of  $\gamma_{p+2}$  for  $0 \leq p \leq \lfloor (g-3)/2 \rfloor$ .<sup>5</sup> Viewing  $V = H^0(\mathbb{P}^{p+2}, \mathcal{O}_{\mathbb{P}^{p+2}}(1))$  and putting  $M_V := M_{\mathcal{O}_{\mathbb{P}^{p+2}}(1)}$ , we have a short exact sequence

$$0 \longrightarrow \wedge^2 M_V \longrightarrow \wedge^2 V \otimes \mathcal{O}_{\mathbb{P}^{p+2}} \longrightarrow M_V(1) \longrightarrow 0.$$

Consider the composition

$$W \otimes \mathcal{O}_{\mathbb{P}^{p+2}} \xrightarrow{\Delta_{p+2} \otimes \text{id}_{\mathcal{O}_{\mathbb{P}^{p+2}}}} \wedge^2 V \otimes \mathcal{O}_{\mathbb{P}^{p+2}} \longrightarrow M_V(1). \quad (3-3)$$

<sup>5</sup> The Koszul complex  $S^q V \otimes \wedge^2 V \rightarrow S^{q+1} V \otimes V \rightarrow S^{q+2} V$  is exact, so  $\text{im}(\delta_{p+2}) = \ker(S^{q+1} V \otimes V \rightarrow S^{q+2} V)$ . Thus the surjectivity of  $\gamma_{p+2}$  for  $0 \leq p \leq \lfloor (g-3)/2 \rfloor$  is equivalent to (1-2):  $W_q(V, W) = 0$  for  $q = g - p - 3 \geq p \geq 0$ .

**Lemma 3.2.** *Assume that  $\text{char}(\mathbb{k}) = 0$  or  $\text{char}(\mathbb{k}) \geq \lfloor (g - 1)/2 \rfloor$ . If the composition (3-3) is surjective, then  $\gamma_{p+2}$  is surjective for  $0 \leq p \leq \lfloor (g - 3)/2 \rfloor$ .*

*Proof.* We can form a short exact sequence

$$0 \longrightarrow K \longrightarrow W \otimes \mathcal{O}_{\mathbb{P}^{p+2}} \xrightarrow{(3-3)} M_V(1) \longrightarrow 0, \tag{3-4}$$

where  $K$  is a vector bundle with  $\text{rank } K = p + 1$  and  $\det K = \mathcal{O}_{\mathbb{P}^{p+2}}(-p - 1)$ . Notice that  $\gamma_{p+2}$  is the  $k = q$  case of the map

$$\gamma'_k : W \otimes H^0(\mathbb{P}^{p+2}, \mathcal{O}_{\mathbb{P}^{p+2}}(k)) \longrightarrow H^0(\mathbb{P}^{p+2}, M_V(k + 1)).$$

As  $q = g - p - 3 \geq p$ , it suffices to show the surjectivity of  $\gamma'_k$  for  $k \geq p$ . To this end, consider the dual of (3-4):

$$0 \longrightarrow M_V^\vee(-1) \longrightarrow W^\vee \otimes \mathcal{O}_{\mathbb{P}^{p+2}} \longrightarrow K^\vee \longrightarrow 0.$$

Since  $M_V^\vee(-1)$  is 1-regular and  $W^\vee \otimes \mathcal{O}_{\mathbb{P}^{p+2}}$  is 0-regular, it follows that  $K^\vee$  is 0-regular. Then  $(K^\vee)^{\otimes p}$  is 0-regular, and so is  $\wedge^p K^\vee$  because  $\text{char}(\mathbb{k}) = 0$  or  $\text{char}(\mathbb{k}) \geq \lfloor (g - 1)/2 \rfloor \geq p + 1$ . Thus  $K = \wedge^p K^\vee \otimes \det K$  is  $(p + 1)$ -regular, and hence,  $\gamma'_k$  is surjective for  $k \geq p$ .  $\square$

It only remains to prove that the composition (3-3) is surjective. The second map in (3-3) is just the globalization of the map

$$\wedge^2 V \longrightarrow V_h, \quad x^{(i)} \wedge x^{(j)} \longmapsto a_j x^{(i)} - a_i x^{(j)},$$

where  $V_h$  is the kernel of a nonzero linear functional  $h := \sum_{i=0}^{p+2} a_i y^i \in V^\vee = S^{p+2}U^\vee$ . We now regard  $h = \sum_{i=0}^{p+2} b_i x^i \in S^{p+2}U$ , where  $b_i = (-1)^i a_{p+2-i}$ . The surjectivity of (3-3) is equivalent to the injectivity of the composition of the dual maps

$$V_h^\vee \longrightarrow \wedge^2 V^\vee \xrightarrow{\Delta_{p+2}^\vee} W^\vee. \tag{3-5}$$

The image of  $V_h^\vee$  in  $\wedge^2 V^\vee$  is spanned by  $v_j := x^j \wedge h$  for  $j = 0, \dots, d - 1, d + 1, \dots, p + 2$ , where  $d := \deg h$ . Recall that  $\Delta_{p+2}^\vee$  is the restriction map  $H^0(\mathbb{P}^2, \mathcal{O}_{\mathbb{P}^2}(p + 1)) \rightarrow H^0(\bar{S}, \mathcal{O}_{\bar{S}}(p + 1))$ .

*Proof of Theorem 1.1.* We consider the case  $\bar{S} = Q$ . In this case,  $\Delta_{p+2}^\vee = \mu_{p+2}$  (see (2-12)). We have  $\Delta_{p+2}^\vee(v_j) = \sum_{i=0}^d (j - i)b_i x^{i+j-1}$ . As  $\text{char}(\mathbb{k}) = 0$  or  $\text{char}(\mathbb{k}) \geq (g + 2)/2 \geq p + 3$ , we get  $\deg(\Delta_{p+2}^\vee(v_j)) = d + j - 1$ . This implies that

$$\Delta_{p+2}^\vee(v_0), \dots, \Delta_{p+2}^\vee(v_{d-1}), \Delta_{p+2}^\vee(v_{d+1}), \dots, \Delta_{p+2}^\vee(v_{p+2}) \text{ are linearly independent.}$$

Thus the composition (3-5) is injective.  $\square$

*Proof of Theorem 1.2.* We consider the case  $\bar{S} = 2\ell$ . In this case,  $\Delta_{p+2}^\vee$  factors through  $\tau_{p+2}$  (see (2-14)). Write  $\tau_{p+2}(v_j) = (u_j, w_j)$ . By Proposition 2.8 (2), we only need to deal with the case  $p \geq 1$ . As  $\text{char}(\mathbb{k}) = 0$  or  $\text{char}(\mathbb{k}) \geq \lfloor (g - 1)/2 \rfloor \geq (p + 3)/2$ , we have

$$\begin{aligned} \deg u_j &= d + j - 1 & \text{and} & & \deg w_j &\leq d + j - 1 & \text{when } d \equiv j \pmod{2}, \\ \deg u_j &\leq d + j - 2 & \text{and} & & \deg w_j &= d + j & \text{when } d \not\equiv j \pmod{2}. \end{aligned}$$

This implies

$$\tau_{p+2}(v_0), \dots, \tau_{p+2}(v_{d-1}), \tau_{p+2}(v_{d+1}), \dots, \tau_{p+2}(v_{p+2}) \text{ are linearly independent.}$$

Thus the composition (3-5) is injective. □

**Remark 3.3.** The characteristic assumption in Theorem 1.1 is used to prove the injectivity of (3-5). If  $k := \text{char}(\mathbb{k}) \leq p + 2$ , then (3-5) is not injective for  $h = 1$ . Indeed,  $x^k \wedge 1 \neq 0$  in  $\wedge^2 V^\vee$  is sent to 0 in  $W^\vee$ . The characteristic assumption in Theorem 1.2 is used in Lemma 3.2.

**Remark 3.4.** By the duality theorem,  $K_{p,q}(S, \mathcal{O}_S(1)) = K_{g-p-2,3-q}(S, \mathcal{O}_S(1))^\vee$ . In the situation of Theorem 1.1 or Theorem 1.2, we have

$$K_{p,1}(S, \mathcal{O}_S(1)) = 0 \quad \text{for } p \geq \lfloor g/2 \rfloor.$$

As  $K_{p+1,1}(S, \mathcal{O}_S(1)) = \ker(\rho_{p+2}) = \ker(\gamma_{p+2})$  and  $\gamma_{p+2}$  is surjective for  $0 \leq p \leq \lfloor (g - 3)/2 \rfloor$ , we can compute  $\kappa_{p,1}(S, \mathcal{O}_S(1))$  for  $0 \leq p \leq \lfloor (g - 1)/2 \rfloor$ . Note that  $K_{p,0}(S, \mathcal{O}_S(1)) \neq 0$  if and only if  $p = 0$ . In this case,  $\kappa_{0,0}(S, \mathcal{O}_S(1)) = 1$ . Thus we can completely determine all graded Betti numbers  $\kappa_{p,q}(S, \mathcal{O}_S(1))$ .

### 4. Proof of Theorem 1.3

Assume that  $\text{char}(\mathbb{k}) = 0$ . Let  $C \subseteq \mathbb{P}^r$  be a smooth projective curve of genus  $g \geq 1$  embedded by  $|L|$ , where  $L$  is a line bundle on  $C$  with  $\text{deg } L \geq 2g + 3$ . We use the notation in Section 2. Recall from Theorem 2.2 that  $\Sigma \subseteq \mathbb{P}^r$  is arithmetically Cohen–Macaulay and  $H^3(\Sigma, \mathcal{O}_\Sigma(m)) = 0$  for  $m > 0$ . By considering (2-5) and Lemma 2.4, we see that

$$\begin{aligned} H^i(T, \mathcal{O}_T(m)) &= H^{i+1}(B, \mathcal{O}_B(mH - \tilde{T} - Z)) \quad \text{for } i \geq 1, m \geq 1, \\ T \subseteq \mathbb{P}^r \text{ is } m\text{-normal} &\iff H^1(B, \mathcal{O}_B(mH - \tilde{T} - Z)) = 0. \end{aligned} \tag{4-1}$$

Note that  $mH - \tilde{T} - Z = (m - 2)H + \pi^*T_L(-4\delta)$ . Thus  $R^1\pi_*\mathcal{O}_B(mH - \tilde{T} - Z) = 0$  for  $m \geq 1$ . When  $m = 1$ , we have  $H^j(B, \mathcal{O}_B(H - \tilde{T} - Z)) = 0$  for  $j > 0$  since  $\pi_*\mathcal{O}_B(H - \tilde{T} - Z) = 0$ . When  $m = 2$ , we have  $H^j(B, \mathcal{O}_B(2H - \tilde{T} - Z)) = H^j(C_2, T_L(-4\delta))$  for  $j > 0$ . Recall that  $\sigma_*\mathcal{O}_Z = \mathcal{O}_{C_2} \oplus \mathcal{O}_{C_2}(-\delta)$ . Then we have  $\sigma_*(L \boxtimes L)(-kD) = T_L(-(k + 1)\delta) \oplus T_L(-k\delta)$ .

**Lemma 4.1.**  $H^i(T, \mathcal{O}_T(m)) = 0$  for  $i > 0, m > 0$ , and  $h^1(T, \mathcal{O}_T) = 2g, h^2(T, \mathcal{O}_T) = 1$ . In particular,  $\text{reg } \mathcal{O}_T = 3$ .

*Proof.* By (2-6) and Proposition 2.5,  $h^1(T, \mathcal{O}_T) = 2g$  and  $h^2(T, \mathcal{O}_T) = 1$ . It suffices to prove that  $H^i(T, \mathcal{O}_T(m)) = 0$  for  $i = 1, m = 1, 2$  and  $i = 2, m = 1$ . Indeed, this implies  $\text{reg } \mathcal{O}_T = 3$ . By (4-1), we need to check that  $H^j(B, \mathcal{O}_B(mH - \tilde{T} - Z)) = 0$  for  $j = 2, m = 1, 2$  and  $j = 3, m = 1$ . As the required vanishing is trivial when  $m = 1$ , it is enough to show that  $H^2(C^2, (L \boxtimes L)(-3D)) = 0$ . Observe that  $R^1p_*(L \boxtimes L)(-3D) = 0$ , where  $p : C \times C \rightarrow C$  is a projection map. Then

$$H^2(C^2, (L \boxtimes L)(-3D)) = H^2(C, p_*(L \boxtimes L)(-3D)) = 0. \quad \square$$

As  $T \subseteq \mathbb{P}^r$  is linearly normal and  $h^1(T, \mathcal{O}_T) = 2g$ , a general hyperplane section of the tangent-developable surface  $T \subseteq \mathbb{P}^r$  is obtained from an isomorphic projection of an  $(r+g)$ -cuspidal curve of geometric genus  $g$  canonically embedded in  $\mathbb{P}^{r+2g-1}$ .

**Lemma 4.2.** *Suppose  $\deg L \geq 3g + 2$ . Then 2-normality of  $T \subseteq \mathbb{P}^r$  is equivalent to*

$$H^1(C \times C, (L \boxtimes L)(-3D)) = 0.$$

*Proof.* By (4-1), 2-normality of  $T \subseteq \mathbb{P}^r$  is equivalent to  $H^1(C_2, T_L(-4\delta)) = 0$ . By [Bertram et al. 1991, Theorem 1(i)],  $H^1(C^2, (L \boxtimes L)(-2D)) = 0$ . It follows that  $H^1(C_2, T_L(-3\delta)) = 0$ . Then  $H^1(C_2, T_L(-4\delta)) = 0$  if and only if  $H^1(C^2, (L \boxtimes L)(-3D)) = 0$ .  $\square$

**Proposition 4.3.**  $K_{p,3}(T, \mathcal{O}_T(1)) = 0$  for  $0 \leq p \leq r - 3$ , and  $\kappa_{r-2,3}(T, \mathcal{O}_T(1)) = 1$ .

*Proof.* By Proposition 2.1 and Lemma 4.1,  $K_{p,3}(T, \mathcal{O}_T(1)) = H^2(T, \wedge^{p+2} M_{\mathcal{O}_T(1)} \otimes \mathcal{O}_T(1))$ . Note that  $\omega_T = \mathcal{O}_T$  (Proposition 2.5) and  $\wedge^{p+2} M_{\mathcal{O}_T(1)}^\vee = \wedge^{r-p-2} M_{\mathcal{O}_T(1)} \otimes \mathcal{O}_T(1)$ . By Serre duality, we have

$$H^2(T, \wedge^{p+2} M_{\mathcal{O}_T(1)} \otimes \mathcal{O}_T(1)) = H^0(T, \wedge^{p+2} M_{\mathcal{O}_T(1)}^\vee \otimes \mathcal{O}_T(-1))^\vee = H^0(T, \wedge^{r-p-2} M_{\mathcal{O}_T(1)}^\vee).$$

Then the proposition immediately follows.  $\square$

*Proof of Theorem 1.3.* By Lemma 4.1, it suffices to show that  $T \subseteq \mathbb{P}^r$  is  $m$ -normal for  $m = 1, 2, 3$ . Indeed, this together with  $\text{reg } \mathcal{O}_T = 3$  implies  $\text{reg } \mathcal{S}_{T|\mathbb{P}^r} = 4$ , and thus,  $T \subseteq \mathbb{P}^r$  is  $m$ -normal for  $m \geq 1$ . In view of (4-1),  $T \subseteq \mathbb{P}^r$  is trivially 1-normal. For the 2-normality, we assume that  $\deg L \geq 4g + 3$ . Then [Bertram et al. 1991, Theorem 1.7(i)] says that  $H^1(C \times C, (L \boxtimes L)(-3D)) = 0$ , so the 2-normality of  $T \subseteq \mathbb{P}^r$  follows from Lemma 4.2. Now, 3-normality of  $T \subseteq \mathbb{P}^r$  is equivalent to  $K_{0,3}(T, \mathcal{O}_T(1)) = 0$ , but this is a special case of Proposition 4.3.  $\square$

**Remark 4.4.** In Theorem 1.3, the degree condition  $\deg L \geq 4g + 3$  is only used to show that  $T \subseteq \mathbb{P}^r$  is 2-normal. If  $\deg L = 4g + 2$ , then by Lemma 4.2 and [Bertram et al. 1991, Theorem 1.7(ii), (iii)],  $T \subseteq \mathbb{P}^r$  is 2-normal (and arithmetically normal) if and only if  $g \geq 3$  and  $C$  is nonhyperelliptic. More generally, using the results in [Bertram et al. 1991, Section 1] and [Pareschi 1995, Theorem 3.8(b)], one can prove that if  $\deg L \geq 4g + 3 - \text{Cliff}(C)$ , then  $T \subseteq \mathbb{P}^r$  is arithmetically normal.

**Corollary 4.5.** *Suppose that  $\deg L \geq 4g + 3$ . Then the Hilbert function of  $T \subseteq \mathbb{P}^r$  is given by  $H_T(t) = (\deg L + g - 1)t^2 + 2 - 2g$  for  $t \geq 1$ . In particular,  $\kappa_{1,1}(T, \mathcal{O}_T(1)) = (r - 2)(r - 3)/2 - 6g$ .*

*Proof.* By Theorem 1.3,  $H_T(t)$  coincides with the Hilbert polynomial  $P_T(t)$  for  $t \geq 1$ . We know that  $\deg P_T(t) = 2$  and the leading coefficient of  $P_T(t)$  is  $(\deg T)/2 = \deg L + g - 1$ . As  $P_T(0) = \chi(\mathcal{O}_T) = 2 - 2g$  and  $P_T(1) = h^0(C, L) = \deg L - g + 1$ , we easily get the assertion. The last assertion follows from  $\kappa_{1,1}(T, \mathcal{O}_T(1)) = h^0(\mathbb{P}^r, \mathcal{S}_{T|\mathbb{P}^r}(2)) = \binom{r+2}{2} - H_T(2)$ .  $\square$

Assume that  $\deg L \geq 4g + 3$ . Notice that (see Proposition 4.3)

$$\begin{aligned} K_{p,0}(T, \mathcal{O}_T(1)) \neq 0 &\iff p = 0 \quad (\kappa_{0,0}(T, \mathcal{O}_T(1)) = 1), \\ K_{p,3}(T, \mathcal{O}_T(1)) \neq 0 &\iff p = r - 2 \quad (\kappa_{r-2,3}(T, \mathcal{O}_T(1)) = 1). \end{aligned}$$



(2) Let  $C \subseteq \mathbb{P}^1 \times \mathbb{P}^1$  be a smooth projective curve of genus 2 defined by

$$x^2 \otimes (x^3 + 1) + 1 \otimes (x^2 - x) \in S^2U \otimes S^3U = H^0(\mathbb{P}^1 \times \mathbb{P}^1, \mathcal{O}_{\mathbb{P}^1}(2) \boxtimes \mathcal{O}_{\mathbb{P}^1}(3)).$$

Consider the embedding  $C \subseteq \mathbb{P}^{11}$  given by  $|(\mathcal{O}_{\mathbb{P}^1}(3) \boxtimes \mathcal{O}_{\mathbb{P}^1}(2))|_C$ . Note that  $\deg C = 13$ . The Betti table of the tangent-developable surface  $T$  of  $C$  in  $\mathbb{P}^{11}$  is the following:

	0	1	2	3	4	5	6	7	8	9	10
0	1	-	-	-	-	-	-	-	-	-	-
1	-	24	48	-	-	-	-	-	-	-	-
2	-	-	153	864	1848	2304	1827	928	288	48	4
3	-	-	-	-	-	-	-	-	-	1	-

We can confirm Conjecture 4.6 and (4-2) for this case, and we see that the expectation (4-3) is not sharp when  $g \geq 2$ .

### Acknowledgements

This paper grew out of attempts to understand the algebraic arguments of [Aprodu et al. 2019; Raicu and Sam 2022] in a geometric way, so the present author wishes to express his deep gratitude to the authors of those papers, particularly to Claudiu Raicu for valuable comments. The author is grateful to Lawrence Ein, Sijong Kwak, Robert Lazarsfeld, and Wenbo Niu, he is indebted to Daniele Agostini for introducing [Gross and Popescu 1998], and he benefited from interesting discussions about Macaulay2 computations in Example 4.8 with Kiryong Chung and Jong In Han.

### References

- [Aprodu and Nagel 2010] M. Aprodu and J. Nagel, *Koszul cohomology and algebraic geometry*, Univ. Lect. Ser. **52**, Amer. Math. Soc., Providence, RI, 2010. MR Zbl
- [Aprodu et al. 2019] M. Aprodu, G. Farkas, Ş. Papadima, C. Raicu, and J. Weyman, “Koszul modules and Green’s conjecture”, *Invent. Math.* **218**:3 (2019), 657–720. MR Zbl
- [Aprodu et al. 2022] M. Aprodu, G. Farkas, Ş. Papadima, C. Raicu, and J. Weyman, “Topological invariants of groups and Koszul modules”, *Duke Math. J.* **171**:10 (2022), 2013–2046. MR Zbl
- [Bayer and Eisenbud 1995] D. Bayer and D. Eisenbud, “Ribbons and their canonical embeddings”, *Trans. Amer. Math. Soc.* **347**:3 (1995), 719–756. MR Zbl
- [Bertram et al. 1991] A. Bertram, L. Ein, and R. Lazarsfeld, “Surjectivity of Gaussian maps for line bundles of large degree on curves”, pp. 15–25 in *Algebraic geometry* (Chicago, IL, 1989), edited by S. Bloch et al., Lecture Notes in Math. **1479**, Springer, 1991. MR Zbl
- [Ein and Lazarsfeld 2020] L. Ein and R. Lazarsfeld, “Tangent developable surfaces and the equations defining algebraic curves”, *Bull. Amer. Math. Soc. (N.S.)* **57**:1 (2020), 23–38. MR Zbl
- [Ein et al. 2020] L. Ein, W. Niu, and J. Park, “Singularities and syzygies of secant varieties of nonsingular projective curves”, *Invent. Math.* **222**:2 (2020), 615–665. MR Zbl
- [Ein et al. 2021] L. Ein, W. Niu, and J. Park, “On blowup of secant varieties of curves”, *Electron. Res. Arch.* **29**:6 (2021), 3649–3654. MR Zbl
- [Eisenbud 1992] D. Eisenbud, “Green’s conjecture: an orientation for algebraists”, pp. 51–78 in *Free resolutions in commutative algebra and algebraic geometry* (Sundance, UT, 1990), edited by D. Eisenbud and C. Huneke, Res. Notes Math. **2**, Jones & Bartlett, Boston, MA, 1992. MR Zbl

- [Eisenbud and Schreyer 2019] D. Eisenbud and F.-O. Schreyer, “Equations and syzygies of K3 carpets and unions of scrolls”, *Acta Math. Vietnam.* **44**:1 (2019), 3–29. MR Zbl
- [Gallego and Purnaprajna 1997] F. J. Gallego and B. P. Purnaprajna, “Degenerations of K3 surfaces in projective space”, *Trans. Amer. Math. Soc.* **349**:6 (1997), 2477–2492. MR Zbl
- [Green 1984] M. L. Green, “Koszul cohomology and the geometry of projective varieties”, *J. Differential Geom.* **19**:1 (1984), 125–171. MR
- [Gross and Popescu 1998] M. Gross and S. Popescu, “Equations of  $(1, d)$ -polarized abelian surfaces”, *Math. Ann.* **310**:2 (1998), 333–377. MR Zbl
- [Hulek 1986] K. Hulek, *Projective geometry of elliptic curves*, Astérisque **137**, Soc. Math. France, Paris, 1986. MR Zbl
- [Kaji 1986] H. Kaji, “On the tangentially degenerate curves”, *J. Lond. Math. Soc. (2)* **33**:3 (1986), 430–440. MR Zbl
- [Kemeny 2021] M. Kemeny, “Universal secant bundles and syzygies of canonical curves”, *Invent. Math.* **223**:3 (2021), 995–1026. MR Zbl
- [Kovács et al. 2010] S. J. Kovács, K. Schwede, and K. E. Smith, “The canonical sheaf of Du Bois singularities”, *Adv. Math.* **224**:4 (2010), 1618–1640. MR Zbl
- [Lazarsfeld 2004] R. Lazarsfeld, *Positivity in algebraic geometry, I: Classical setting: line bundles and linear series*, Ergebnisse der Math. (3) **48**, Springer, 2004. MR Zbl
- [Pareschi 1995] G. Pareschi, “Gaussian maps and multiplication maps on certain projective varieties”, *Compos. Math.* **98**:3 (1995), 219–268. MR Zbl
- [Park 2022] J. Park, “Asymptotic vanishing of syzygies of algebraic varieties”, *Commun. Am. Math. Soc.* **2** (2022), 133–148. MR Zbl
- [Raicu 2021] C. Raicu, “Koszul modules”, pp. 33–57 in *Recent developments in commutative algebra* (Trento, Italy, 2019), edited by A. Conca et al., Lecture Notes in Math. **2283**, Springer, 2021. MR Zbl
- [Raicu and Sam 2021] C. Raicu and S. V. Sam, “Hermite reciprocity and Schwarzenberger bundles”, pp. 689–721 in *Commutative algebra*, edited by I. Peeva, Springer, 2021. MR Zbl
- [Raicu and Sam 2022] C. Raicu and S. V. Sam, “Bi-graded Koszul modules, K3 carpets, and Green’s conjecture”, *Compos. Math.* **158**:1 (2022), 33–56. MR Zbl
- [Rathmann 2016] J. Rathmann, “An effective bound for the gonality conjecture”, preprint, 2016. arXiv 1604.06072
- [Schreyer 1983] F. O. Schreyer, *Syzygies of curves with special pencils*, Ph.D. thesis, Brandeis University, 1983, available at <https://www.proquest.com/docview/303128625>.
- [Schreyer 1986] F.-O. Schreyer, “Syzygies of canonical curves and special linear series”, *Math. Ann.* **275**:1 (1986), 105–137. MR Zbl
- [Voisin 2002] C. Voisin, “Green’s generic syzygy conjecture for curves of even genus lying on a K3 surface”, *J. Eur. Math. Soc.* **4**:4 (2002), 363–404. MR Zbl
- [Voisin 2005] C. Voisin, “Green’s canonical syzygy conjecture for generic curves of odd genus”, *Compos. Math.* **141**:5 (2005), 1163–1190. MR Zbl
- [Wahl 1990] J. Wahl, “Gaussian maps on algebraic curves”, *J. Differential Geom.* **32**:1 (1990), 77–98. MR Zbl

Communicated by Gavril Farkas

Received 2024-04-28      Accepted 2024-07-16

parkjh13@kaist.ac.kr

Department of Mathematical Sciences, KAIST, Daejeon, South Korea

# Algebra & Number Theory

msp.org/ant

## EDITORS

MANAGING EDITOR  
Antoine Chambert-Loir  
Université Paris-Diderot  
France

EDITORIAL BOARD CHAIR  
David Eisenbud  
University of California  
Berkeley, USA

## BOARD OF EDITORS

Jason P. Bell	University of Waterloo, Canada	Philippe Michel	École Polytechnique Fédérale de Lausanne
Bhargav Bhatt	University of Michigan, USA	Martin Olsson	University of California, Berkeley, USA
Frank Calegari	University of Chicago, USA	Irena Peeva	Cornell University, USA
J.-L. Colliot-Thélène	CNRS, Université Paris-Saclay, France	Jonathan Pila	University of Oxford, UK
Brian D. Conrad	Stanford University, USA	Anand Pillay	University of Notre Dame, USA
Samit Dasgupta	Duke University, USA	Bjorn Poonen	Massachusetts Institute of Technology, USA
Hélène Esnault	Freie Universität Berlin, Germany	Victor Reiner	University of Minnesota, USA
Gavril Farkas	Humboldt Universität zu Berlin, Germany	Peter Sarnak	Princeton University, USA
Sergey Fomin	University of Michigan, USA	Michael Singer	North Carolina State University, USA
Edward Frenkel	University of California, Berkeley, USA	Vasudevan Srinivas	SUNY Buffalo, USA
Wee Teck Gan	National University of Singapore	Shunsuke Takagi	University of Tokyo, Japan
Andrew Granville	Université de Montréal, Canada	Pham Huu Tiep	Rutgers University, USA
Ben J. Green	University of Oxford, UK	Ravi Vakil	Stanford University, USA
Christopher Hacon	University of Utah, USA	Akshay Venkatesh	Institute for Advanced Study, USA
Roger Heath-Brown	Oxford University, UK	Melanie Matchett Wood	Harvard University, USA
János Kollár	Princeton University, USA	Shou-Wu Zhang	Princeton University, USA
Michael J. Larsen	Indiana University Bloomington, USA		

## PRODUCTION

production@msp.org  
Silvio Levy, Scientific Editor

---

See inside back cover or [msp.org/ant](http://msp.org/ant) for submission instructions.

---

The subscription price for 2025 is US \$565/year for the electronic version, and \$820/year (+\$70, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to MSP.

---

Algebra & Number Theory (ISSN 1944-7833 electronic, 1937-0652 printed) at Mathematical Sciences Publishers, 798 Evans Hall #3840, c/o University of California, Berkeley, CA 94720-3840 is published continuously online.

---

ANT peer review and production are managed by EditFLOW<sup>®</sup> from MSP.

PUBLISHED BY

 **mathematical sciences publishers**  
nonprofit scientific publishing

<http://msp.org/>

© 2025 Mathematical Sciences Publishers

# Algebra & Number Theory

Volume 19 No. 5 2025

---

Presentations of Galois groups of maximal extensions with restricted ramification YUAN LIU	835
Motivic distribution of rational curves and twisted products of toric varieties LOÏS FAISANT	883
Smooth cuboids in group theory JOSHUA MAGLIONE and MIMA STANOJKOVSKI	967
Malle's conjecture for fair counting functions PETER KOYMANS and CARLO PAGANO	1007
Syzygies of tangent-developable surfaces and K3 carpets via secant varieties JINHYUNG PARK	1029