

Algebra & Number Theory

Volume 20
2026
No. 2

Moments of one-level densities in families of holomorphic cusp forms in the level aspect

Peter Cohen, Justine Dell, Oscar E. González, Simran Khunger, Chung-Hang Kwan, Steven J. Miller, Alexander Shashkov, Alicia Reina Smith, Carsten Sprunger, Nicholas Triantafyllou, Nhi Truong, Roger Van Peski and Stephen Willis



Moments of one-level densities in families of holomorphic cusp forms in the level aspect

Peter Cohen, Justine Dell, Oscar E. González, Simran Khunger, Chung-Hang Kwan, Steven J. Miller, Alexander Shashkov, Alicia Reina Smith, Carsten Sprunger, Nicholas Triantafillou, Nhi Truong, Roger Van Peski and Stephen Willis

We study the n -th centered moments of the 1-level density for the low-lying zeros of L -functions attached to holomorphic cuspidal newforms of large prime level and fixed weight. Assuming the generalized Riemann hypotheses, we compute this statistic for any $n \geq 1$ and for all test functions whose Fourier transforms are supported in $(-\frac{2}{n}, \frac{2}{n})$. This is believed to be the natural limit of the current technology. Our work significantly extends beyond the trivial range $(-\frac{1}{n}, \frac{1}{n})$ and surpasses the previous record of $(-\frac{1}{n-1}, \frac{1}{n-1})$ whenever $n > 2$. The Katz–Sarnak philosophy predicts that the aforementioned statistic can be modeled by the corresponding statistic for the eigenvalues of random orthogonal matrices. We prove that this is the case for test functions with Fourier support contained in $(-\frac{2}{n}, \frac{2}{n})$. The main technical innovation is a tractable vantage to evaluate the combinatorial zoo of terms, similar to the work of Conrey, Snaith and Mason. As an application, our work provides better bounds on the order of vanishing at the central point for the L -functions in our family.

1. Introduction	237
2. Preliminaries	243
3. Proof of Theorem 1.2 assuming Propositions 3.6 and 3.7	249
4. Proof of Propositions 3.6 and 3.7	255
5. Extending support for random matrix theory: proof of Theorem 1.3	273
Appendix A. Proof of Lemma 3.1	287
Appendix B. Increasing support for the nonsplit family	291
Appendix C. Proofs of the lemmas in Section 5	293
Appendix D. Bounding the order of vanishing at the central point	295
References	296

1. Introduction

1.1. Historical perspectives. Since Montgomery and Dyson’s discovery that the two-point correlation of the zeros of the Riemann zeta function agrees with the pair correlation function for eigenvalues of the

This work was done during the 2011, 2012, 2015, 2017, and 2021 SMALL REU program in Number Theory and Probability at Williams College supported by NSF grants DMS1561945 and DMS1659037. We thank Roger Heath-Brown for his assistance on exponential sums. We also thank Geoffrey Iyer and Yingzi Yang for significant work on parts of earlier drafts of this work. C.-H. Kwan is supported by EPSRC Grant EP/W009838/1 and the Harish-Chandra Fund from the Institute for Advanced Study. *MSC2020*: primary 11M26; secondary 11M41, 15A42.

Keywords: low-lying zeros, optimal test function, n -level density, Katz–Sarnak conjecture.

Gaussian unitary ensemble (see [36]), the connection between the zeros of L -functions and the zeros of random matrices has been a major area of study. It is now widely believed that the statistical behavior of families of L -functions can be modeled by ensembles of random matrices. Based on the observation that the spacing statistics of high zeros associated with cuspidal L -functions agree with the corresponding statistics for eigenvalues of random unitary matrices under Haar measure (see [43], for example), it was originally believed that only the unitary ensemble was important to number theory. However, Katz and Sarnak [26; 27] showed that these statistics are the same for all classical compact groups. These statistics, the n -level correlations, are unaffected by finite numbers of zeros. In particular, they failed to identify differences in behavior near the central point $s = \frac{1}{2}$.

The n -level density statistic was introduced to distinguish the behavior of families of L -functions close to $s = \frac{1}{2}$. Based partially on an analogy with the function field setting, Katz and Sarnak conjectured that the *low-lying zeros* (i.e., zeros near $s = \frac{1}{2}$) of families of L -functions behave like the eigenvalues near 1 of classical compact groups (unitary, symplectic, and orthogonal). The behavior of the eigenvalues near 1 is different for each matrix group. A growing body of evidence has shown that this conjecture holds for test functions with suitably restricted support for a wide range of families of L -functions. For a nonexhaustive list, see [1; 2; 4; 9; 10; 11; 12; 15; 16; 17; 19; 21; 25; 28; 33; 35; 38; 39; 40; 41; 42; 44; 47; 48; 49]. Much of the previous work is focused on the $n = 1$ case. We study the n -th centered moment of the 1-level density for any $n \geq 1$, a higher-order statistic first introduced by Hughes and Rudnick [22] which is combinatorially simpler than the n -level density.

In this article, we consider the family of L -functions associated with holomorphic cusp forms. We prove that the Katz–Sarnak conjecture holds for the n -th centered moment of the 1-level density for this family and for all test functions ϕ with Fourier transform $\hat{\phi}$ supported in $(-\frac{2}{n}, \frac{2}{n})$, which represents the natural limit of the current analytic machinery based on the prior works of [22] on the unitary family of primitive Dirichlet characters, as well as [17], [15] and [29] on the n -level density for the symplectic family of quadratic Dirichlet L -functions. Proving the conjecture past this support likely requires new ideas or stronger hypotheses such as the ‘*Hypothesis S*’ discussed in [25, Section 10] (or more likely its generalizations; see [32]).

Previously, Iwaniec, Luo and Sarnak [25] and Hughes and Miller [21] examined the same family and statistic addressed in this article (with the former limited to the case of $n = 1$). Under the generalized Riemann hypothesis (GRH), the conjecture was proved in [25] when $n = 1$ for $\hat{\phi}$ supported in $(-2, 2)$, and in [21] for the restricted range $(-\frac{1}{n-1}, \frac{1}{n-1})$ when $n \geq 2$. In [21], the authors were unable to handle the new terms which emerge at larger supports on both the number theory and random matrix theory side (nonetheless, that work addressed some rather challenging combinatorics). Our work develops an approach to handle all the combinatorial terms that arise when the Fourier support of the test function extends up to $2/n$, thus filling in a nontrivial gap in our current knowledge of higher-order statistics for the zeros of L -functions in the orthogonal families.

To the best of the authors’ knowledge, apart from this work, the success of a *purely combinatorial venture* (i.e., without passing to function field settings in the large q -limit and using the equidistribution

theorems à la Deligne and Katz) in verifying the Katz–Sarnak philosophy through higher-order statistics is somewhat limited. [6] calculate the n -level density for a unitary family of Dirichlet L -functions. For the symplectic family, Levinson and Miller [29] calculated the n -level density of a family of Dirichlet L -functions for $n \leq 7$, extending the work of Gao [17]. See also the contributions of [7] and [31] in the realm of random matrix theory. The pioneering work of Entin and coauthors [15; 14] calculated the n -level density for the symplectic family studied in [29] and [17] and showed it agreed with random matrix theory by passing to the function field setting. However, their method has not yet been successfully applied to any of the orthogonal families. Furthermore, the success of this article seems to suggest that the n -th centered moment statistic might offer a viable means to overcome the combinatorial barriers encountered in [17] and [29] when verifying the higher-order Katz–Sarnak comparison for the symplectic family.

1.2. Number theory setup. We now describe the main objects of study. Let $H_k^*(N)$ be the set of holomorphic cusp forms of weight k and level N (with trivial nebentypus) which are *newforms*. For $f \in H_k^*(N)$, denote by $L(s, f)$ the L -function attached to f . The completed L -function is given by

$$\Lambda(s, f) := \left(\frac{\sqrt{N}}{2\pi}\right)^s \Gamma\left(s + \frac{k-1}{2}\right)L(s, f). \tag{1-1}$$

It admits an entire continuation and satisfies the functional equation

$$\Lambda(s, f) = \epsilon_f \Lambda(1 - s, f), \tag{1-2}$$

where $\epsilon_f = \pm 1$ is the root number. The family $H_k^*(N)$ splits naturally into two disjoint subfamilies:

$$H_k^+(N) := \{f \in H_k^*(N) : \epsilon_f = +1\}, \quad H_k^-(N) := \{f \in H_k^*(N) : \epsilon_f = -1\}. \tag{1-3}$$

For each $f \in H_k^*(N)$, we denote the nontrivial zeros of $L(s, f)$ by $\rho_f = \frac{1}{2} + i\gamma_f$. The generalized Riemann hypothesis (GRH) for $L(s, f)$ asserts that $\gamma_f \in \mathbb{R}$.

As in [25], we take

$$R = k^2 N \tag{1-4}$$

as the working definition of the *analytic conductor* for the families $H_k^\bullet(N)$ ($\bullet \in \{+, -, *\}$). Our analysis is greatly simplified by all forms in the family having the same analytic conductor. Varying conductors are easily handled in 1-level calculations, but cause technical difficulties through cross terms once $n \geq 2$; see [33].

The *one-level density* of $f \in H_k^*(N)$ is a weighted sum over γ_f given by

$$D(f; \phi) := \sum_{\gamma_f} \phi\left(\frac{\log R}{2\pi} \gamma_f\right), \tag{1-5}$$

where the test function $\phi : \mathbb{R} \rightarrow \mathbb{C}$ is an even Schwartz function whose Fourier transform $\hat{\phi}$ has compact support. We denote this class of test functions by $\mathcal{S}_{ec}(\mathbb{R})$ and $\phi \in \mathcal{S}_{ec}(\mathbb{R})$. Because of the rapid decay of ϕ , the *low-lying zeros* of $L(s, f)$ contribute the most to the one-level density (1-5).

As in [21], we use the following shorthand for taking averages over $H_k^\bullet(N)$:

$$\langle Q(\cdot) \rangle_\bullet = \langle Q(\cdot) \rangle_{H_k^\bullet(N)} := \frac{1}{|H_k^\bullet(N)|} \sum_{f \in H_k^\bullet(N)} Q(f), \tag{1-6}$$

where $\bullet \in \{+, -, *\}$ and Q is any complex-valued function defined on $H_k^\bullet(N)$, e.g., $f \mapsto D(f; \phi)$ defined in (1-5).

In this article, we study the n -th centered moment of the one-level density, defined by

$$\mathcal{D}_n^\bullet(N; \phi) := \left\langle \left(D(\cdot; \phi) - \langle D(\cdot; \phi) \rangle_{H_k^\bullet(N)} \right)^n \right\rangle_{H_k^\bullet(N)} \tag{1-7}$$

where $n \geq 2$ is an integer, N is prime, $k \geq 2$ is even, and $\phi \in \mathcal{S}_{ec}(\mathbb{R})$ is a test function. Readers should keep in mind that the weight k is kept fixed and the level N goes to ∞ through primes.

1.3. Random matrix theory setup. Let $\phi \in \mathcal{S}_{ec}(\mathbb{R})$. The random matrix theory counterpart of the one-level density $D(f; \phi)$ is given by

$$Z(U; \phi) := \sum_{n=1}^M \sum_{j=-\infty}^{\infty} \phi \left(\frac{M}{2\pi} (\theta_n + 2\pi j) \right), \tag{1-8}$$

where U is an $M \times M$ special orthogonal matrix with eigenvalues $\{e^{i\theta_n} : n = 1, \dots, M\}$. In fact, the eigenvalues of U always occur in complex-conjugate pairs. We have the following correspondence when comparing (1-8) to (1-5):

$$\begin{aligned} M &\longleftrightarrow \log(k^2 N) \\ U \in SO(M) &\longleftrightarrow f \in H_k^*(N) \\ \{\theta_n + 2\pi j\}_{\substack{j \in \mathbb{Z} \\ 1 \leq n \leq M}} &\longleftrightarrow \{\gamma_f\}. \end{aligned}$$

Here, it is more standard to use the notation

$$\mathbb{E}_{SO(M)} [Z(\cdot; \phi)] := \int_{SO(M)} Z(U; \phi) dU \tag{1-9}$$

instead of the bracket of (1-6). The Haar measure dU on the compact Lie group $SO(M)$ is normalized to have total measure 1. When $\text{supp}(\hat{\phi}) \subseteq [-1, 1]$, it is well-known that

$$\mu_\pm := \lim_{M \rightarrow \infty}^\pm \mathbb{E}_{SO(M)} [Z(\cdot; \phi)] = \hat{\phi}(0) + \frac{1}{2} \int_{-1}^1 \hat{\phi}(y) dy, \tag{1-10}$$

where

$$\lim^+ := \lim_{\substack{M \rightarrow \infty \\ M \equiv 0 \pmod{2}}} \quad \text{and} \quad \lim^- := \lim_{\substack{M \rightarrow \infty \\ M \equiv 1 \pmod{2}}}.$$

Similarly to (1-7), the n -th centered moment of $Z(\cdot; \phi)$ is defined as

$$\mathcal{Z}_n(M; \phi) := \mathbb{E}_{SO(M)} \left[\left(Z(\cdot; \phi) - \mathbb{E}_{SO(M)} [Z(\cdot; \phi)] \right)^n \right] \quad (M \in \mathbb{N}). \tag{1-11}$$

1.4. Main results. We are now ready to present our main result, which extends and generalizes [25, Theorem 1.1] and [21, Theorem 1.6-1.7].

Theorem 1.1. *Let $k, n \geq 2$ be positive integers with k even. Assume GRH for $L(s, f)$ for all $f \in H_k^*(N)$, where N is any prime or $N = 1$. Assume also RH for $\zeta(s)$ and GRH for all primitive Dirichlet L -functions. Then for $\phi \in \mathcal{S}_{ec}(\mathbb{R})$ with $\text{supp}(\hat{\phi}) \subset (-\frac{2}{n}, \frac{2}{n})$, we have*

$$\lim_{\substack{N \rightarrow \infty \\ N \text{ prime}}} \mathcal{D}_n^\pm(N; \phi) = \lim_{M \rightarrow \infty}^\pm \mathcal{Z}_n(M; \phi). \tag{1-12}$$

The moments $\mathcal{D}_n^\pm(N; \phi)$ and $\mathcal{Z}_n(M; \phi)$ are defined in (1-7) and (1-11).

Theorem 1.1 follows immediately from Theorems 1.2 and 1.3 below. In fact, we prove precise formulae for each of the limits in (1-12) whenever $\text{supp}(\hat{\phi}) \subset (-\frac{1}{n-a}, \frac{1}{n-a})$, where a is an integer with $0 \leq a \leq n/2$. These formulae might be of independent interest. To state these results, we need to introduce the quantities

$$\sigma_\phi^2 := 2 \int_{-\infty}^\infty |y| \hat{\phi}(y)^2 dy, \tag{1-13}$$

$$\begin{aligned} \mathcal{R}(m, i; \phi) := & 2^{m-1} (-1)^{m+1} \sum_{\ell=0}^{i-1} (-1)^\ell \binom{m}{\ell} \\ & \times \left(-\frac{1}{2} \phi(0)^m + \int_{-\infty}^\infty \cdots \int_{-\infty}^\infty \hat{\phi}(x_2) \cdots \hat{\phi}(x_{\ell+1}) \int_{-\infty}^\infty \phi(x_1)^{m-\ell} \right. \\ & \left. \times \frac{\sin(2\pi x_1(1 + |x_2| + \cdots + |x_{\ell+1}|))}{2\pi x_1} dx_1 \cdots dx_{\ell+1} \right), \end{aligned} \tag{1-14}$$

and

$$S(n, a; \phi) := \sum_{\ell=0}^{\lfloor \frac{a-1}{2} \rfloor} \frac{n!}{(n-2\ell)! \ell!} \mathcal{R}(n-2\ell, a-2\ell; \phi) \left(\frac{\sigma_\phi^2}{2} \right)^\ell, \tag{1-15}$$

where $\phi \in \mathcal{S}_{ec}(\mathbb{R})$, $1 \leq i \leq m$, and $0 \leq a \leq n/2$.

We first state the number theory result.

Theorem 1.2. *Let n, a be integers with $n \geq 2$, $0 \leq a \leq n/2$ and let $\phi \in \mathcal{S}_{ec}(\mathbb{R})$. Under the same assumptions as Theorem 1.1, if $\text{supp}(\hat{\phi}) \subset (-\frac{1}{n-a}, \frac{1}{n-a})$, then*

$$\lim_{\substack{N \rightarrow \infty \\ N \text{ prime}}} \mathcal{D}_n^\pm(N; \phi) = \mathbb{1}_{\{\text{even}\}}(n) \cdot (n-1)!! (\sigma_\phi^2)^{n/2} \pm S(n, a; \phi), \tag{1-16}$$

where $\mathbb{1}_{\{\text{even}\}}(n)$ is equal to 1 if n is even and is 0 if n is odd.

Next we state the random matrix theory result.

Theorem 1.3. *Let n, a be integers with $n \geq 2$, $0 \leq a \leq n/2$ and let $\phi \in \mathcal{S}_{ec}(\mathbb{R})$. If $\text{supp}(\hat{\phi}) \subset (-\frac{1}{n-a}, \frac{1}{n-a})$, then*

$$\lim_{M \rightarrow \infty}^\pm \mathcal{Z}_n(M; \phi) = \mathbb{1}_{\{n \text{ even}\}} \cdot (n-1)!! (\sigma_\phi^2)^{n/2} \pm S(n, a; \phi) \tag{1-17}$$

where $\mathbb{1}_{\{\text{even}\}}(n)$ is equal to 1 if n is even and is 0 if n is odd.

Hughes and Miller [21, Theorem E.1] proved an analogue of Theorem 1.1 for the *full* family $H_k^*(N)$ under the restriction $n \leq 2k$. We remove this restriction in the following theorem, whose proof is given in Appendix B.

Theorem 1.4. *Let $k, n \geq 2$ be positive integers with k even and let $\phi \in \mathcal{S}_{ec}(\mathbb{R})$. Under the same assumptions as Theorem 1.1, if $\text{supp}(\hat{\phi}) \subset \left(-\frac{1}{n-a}, \frac{1}{n-a}\right)$, then*

$$\lim_{\substack{N \rightarrow \infty \\ N \text{ prime}}} \mathcal{D}_n^*(N; \phi) = \frac{1}{2} \left[\lim_{M \rightarrow \infty}^+ \mathcal{Z}_n(M; \phi) + \lim_{M \rightarrow \infty}^- \mathcal{Z}_n(M; \phi) \right]. \quad (1-18)$$

The moments $\mathcal{D}_n^\pm(N; \phi)$ and $\mathcal{Z}_n(M; \phi)$ are defined in (1-7) and (1-11).

1.5. Applications. As noted in [21] and [34], another application of centered moments is in bounding the order of vanishing of L -functions at the central point. In Appendix D, we show how to use Theorem 1.2 to bound the probability that a newform with negative sign will have order of vanishing exceeding some r at the central point. Similar calculations may be done for the positive sign family. Our results provide the best known bounds (conditional on GRH) for order of vanishing at the central point when $r \geq 5$, surpassing [25; 21; 5; 30].

1.6. Proof sketch and structure of the paper. We evaluate the limit (1-16) using the *explicit formula* and the *Petersson trace formula*. We use the explicit formula to transform the sum over zeros (1-5) into a weighted average of products of Hecke eigenvalues over primes in Lemma 2.11. After removing many lower order subterms with Lemmas 3.1 and 3.2, we apply the Petersson trace formula and study the resulting sums of Kloosterman sums over primes; see (3-4). We assume GRH for $L(s, f)$ when applying the Petersson trace formula; see Remark 2.9.

Using Lemma 2.2, we convert the Kloosterman sums into sums over Gauss sums. Assuming GRH for Dirichlet L -functions, we show in Lemma 3.5 that the terms involving Gauss sums with nonprincipal characters contribute negligibly in the limit when $\text{supp}(\hat{\phi}) \subset \left(-\frac{2}{n}, \frac{2}{n}\right)$. This requires strong bounds for various character sums over primes, hence the need to assume GRH for Dirichlet L -functions. Hughes and Miller [21, Theorems 1.1 and 1.3] proved results without GRH for Dirichlet L -functions for $\text{supp}(\hat{\phi}) \subset \left(-\frac{1}{n}, \frac{1}{n}\right)$ (i.e., the case where $a = 0$ in Theorem 1.2-1.3); we use GRH for Dirichlet L -functions to extend the support to $\text{supp}(\hat{\phi}) \subset \left(-\frac{2}{n}, \frac{2}{n}\right)$.

We are left to handle certain smooth sums over primes (Proposition 4.1) and a convolution sum of Ramanujan sums (Proposition 4.2). We arrive at Theorem 1.2 upon very careful bookkeeping of the resulting combinatorics (see Sections 3.4 and 4).

Beyond the regime $\text{supp}(\hat{\phi}) \subset \left(-\frac{1}{n-1}, \frac{1}{n-1}\right)$ proven by [21], more complicated terms emerge as the size of $\text{supp}(\hat{\phi})$ increases. This serves as the primary obstacle to generalizing the work of [21]. The main insight of our extension lies in the observation that many of these terms actually *vanish in the limit* (see Lemma 3.1 and Proposition 3.6). This enables us to ignore the very intricate combinatorics behind these terms. The remaining terms contribute to the limit and exhibit nicer symmetries. We are able to obtain an integral representation for these terms in Proposition 3.7 upon making our way through

the combinatorial jungle. Our work features many elaborate combinatorial simplifications that result in exact matching with the calculations from the random matrix theory side. As with number theory, the key result that allows us to obtain greater support in random matrix theory is the vanishing of many of the complicated terms which emerge at larger supports (see Lemma 5.20).

The structure of this paper is as follows. In Section 2, we review the notations and conventions, and state some needed estimates. In Section 3, we work with the geometric side of the Petersson formula with the focus of locating the main contributions in the expansion. We also prove Theorem 1.2 assuming two key propositions (3.6 and 3.7). In Section 4, we evaluate the main contributions and complete the proofs of Propositions 3.6 and 3.7. In Section 5 we work on the random matrix theory side and prove Theorem 1.3.

In Appendix A, we prove Lemma 3.1 regarding the combinatorial expansion for $\mathcal{D}_n^*(N; \phi)$ which serves as the starting point of the number-theoretic calculations. In Appendix B, we give a sketch of proof for Theorem 1.4 regarding the nonsplit family. In Appendix C, we include more details for the random matrix theory calculations. In Appendix D we use Theorem 1.2 to bound the proportion of cuspidal newforms vanishing to a certain order at the central point.

2. Preliminaries

2.1. Notations and conventions. In this article, $e(x) := e^{2\pi ix}$. The Fourier transform and its inverse transform are given by

$$\hat{\phi}(y) := \int_{-\infty}^{\infty} \phi(x)e(-xy) dx, \quad \phi(x) := \int_{-\infty}^{\infty} \hat{\phi}(y)e(xy) dy \tag{2-1}$$

for $x, y \in \mathbb{R}$. The Mellin transform and its inverse transform are

$$\tilde{\psi}(s) := \int_0^{\infty} \psi(x)x^{s-1} dx, \quad \psi(x) := \int_{(\sigma)} \tilde{\psi}(s)x^{-s} \frac{ds}{2\pi i} \tag{2-2}$$

for $x > 0$ and $s \in \mathbb{C}$ in a vertical strip, provided that the integrals of (2-1) and (2-2) converge absolutely.

For $A \subset \mathbb{R}$, let

$$\mathbb{1}_{\{x \in A\}} := \begin{cases} 1 & \text{if } x \in A \\ 0 & \text{otherwise.} \end{cases} \tag{2-3}$$

We will suppress the argument of a characteristic function when it is clear from context. For $x, y \in \mathbb{R}$, let

$$\delta(x, y) := \begin{cases} 1 & \text{if } x = y \\ 0 & \text{otherwise.} \end{cases} \tag{2-4}$$

For $x, y \in \mathbb{Z}$, let (x, y) denote the greatest common divisor of x and y . Set $(x, y^\infty) = \max_{n \in \mathbb{N}}(x, y^n)$ and $(x^\infty, y) = \max_{n \in \mathbb{N}}(x^n, y)$.

We adopt the following set of conventions throughout this work.

- (1) We use ‘ i ’ to denote the imaginary unit and ‘ i ’ for indices of summation.
- (2) We use p, p_i ’s, q_j ’s to denote prime numbers.

- (3) The test function $\phi : \mathbb{R} \rightarrow \mathbb{C}$ is an even Schwartz function with its Fourier transform $\hat{\phi}$ having compact support.
- (4) The weight k is kept fixed and the level N goes to ∞ through primes
- (5) A quantity is considered to be *negligibly small* if it is $o(1)$ as $N \rightarrow \infty$.
- (6) The implicit constants for O, \ll, \gg, \asymp , etc. may depend on k, ϵ, n, a , and of course the test function ϕ (compare Theorem 1.2). We will omit such dependencies to simplify notation.
- (7) We shall frequently adopt the epsilon convention, namely that $\epsilon > 0$ is an arbitrary small quantity and $O(N^{-\epsilon}) \cdot O(N^{-C(k, \epsilon, n, a) \cdot \epsilon}) = O(N^{-\epsilon})$ (say).
- (8) We always assume GRH for $L(s, f)$ for any $f \in H_k^*(N)$ with $N = 1$ and primes N .

2.2. Analytic preliminaries. We will frequently encounter the following exponential/character sums.

Definition 2.1. Let $m, n, q \in \mathbb{Z}$ with $q \geq 1$ and $\chi \pmod{q}$ be a Dirichlet character. We define

$$G_\chi(n) := \sum_{a \pmod{q}} \chi(a)e(an/q), \tag{2-5}$$

$$R(n, q) := \sum_{a \pmod{q}}^* e(an/q), \tag{2-6}$$

$$S(m, n; q) := \sum_{a \pmod{q}}^* e\left(\frac{ma}{q} + \frac{n\bar{a}}{q}\right), \tag{2-7}$$

where $*$ restricts the summation to the reduced residue classes $a \pmod{q}$, and $a\bar{a} \equiv 1 \pmod{q}$.

The sums $G_\chi(n)$, $R(n, q)$, and $S(m, n; q)$ are known as the Gauss sum, the Ramanujan sum, and the Kloosterman sum. When $\chi = \chi_0$ (i.e., the principal character \pmod{q}), we have $G_{\chi_0}(n) = R(n, q)$. Also, $R(n, q) = S(0, n; q)$. The Ramanujan sum admits the following explicit evaluation:

$$R(n, q) = \sum_{d|(n, q)} \mu(q/d)d = \mu\left(\frac{q}{(q, n)}\right) \frac{\phi(q)}{\phi\left(\frac{q}{(q, n)}\right)}, \tag{2-8}$$

where $\mu(\cdot)$ and $\phi(\cdot)$ are the Möbius μ -function and the Euler totient function. We will need the multiplicativity of the Ramanujan sum as well:

$$R(n, q_1q_2) = R(n, q_1)R(n, q_2) \tag{2-9}$$

for $(q_1, q_2) = 1$ and $n \in \mathbb{Z}$.

The following bounds are particularly handy in showing various sums and integrals to be negligibly small. We have

$$|G_\chi(n)| \leq \sqrt{q}, \tag{2-10}$$

and

$$|S(m, n; q)| \leq (m, n, q) \sqrt{\min\left\{\frac{q}{(m, q)}, \frac{q}{(n, q)}\right\}} \tau(q), \tag{2-11}$$

where $\tau(\cdot)$ is the divisor function. The bound (2-11) is a convenient reformulation of a well-known result of A. Weil; see [25, 2.13].

We often need to handle the Kloosterman sums in more refined ways than merely applying (2-11). In the context of low-lying zeros (see [21; 25]), it is advantageous to expand Kloosterman sums in terms of Dirichlet characters and Gauss sums. The following is a generalization of [21, Lemma C.1].

Lemma 2.2. *Let N be a prime not dividing bQm . Then*

$$S(m^2, NQ; Nb) = -\frac{1}{\varphi(b)} \sum_{\chi(b)} G_\chi(m^2) G_\chi((Q, b^\infty)) \bar{\chi}\left(\frac{Q}{(Q, b^\infty)}\right) \chi(N). \tag{2-12}$$

Proof. Set $r = (Q, b^\infty)$ and $Q' = Q/r$. Then $(Q', b) = 1$. Using the orthogonality relation and opening up the Kloosterman sum by its definition, observe that

$$\begin{aligned} S(m^2, NQ; Nb) &= \frac{1}{\varphi(b)} \sum_{\chi(b)} \sum_{a(b)}^* \chi(a) \bar{\chi}(Q') S(m^2, Nra; Nb) \\ &= \frac{1}{\varphi(b)} \sum_{\chi(b)} \bar{\chi}(Q') \sum_{d(Nb)}^* e\left(\frac{m^2 d}{Nb}\right) \sum_{a(b)}^* \chi(a) \left(\frac{rad}{b}\right). \end{aligned} \tag{2-13}$$

Making a change of variables $a \rightarrow ad$ in the a -sum and breaking up the d -sum by $d = u_1 N + u_2 b$ with $(u_1, b) = 1$ and $(u_2, N) = 1$, it follows that

$$\begin{aligned} S(m^2, NQ; Nb) &= \frac{1}{\varphi(b)} \sum_{\chi(b)} \bar{\chi}(Q') G_\chi(r) \sum_{d(Nb)}^* \chi(d) e\left(\frac{m^2 d}{Nb}\right) \\ &= \frac{1}{\varphi(b)} \sum_{\chi(b)} \bar{\chi}(Q') G_\chi(r) \chi(N) \sum_{u_1(b)}^* \chi(u_1) e\left(\frac{m^2 u_1}{b}\right) \sum_{u_2(N)}^* e\left(\frac{m^2 u_2}{N}\right). \end{aligned} \tag{2-14}$$

The u_2 -sum and u_1 -sum of (2-14) evaluate to -1 and $G_\chi(m^2)$ respectively as $(m^2, N) = 1$. The result follows. □

Lemma 2.3. *Let $\chi \pmod{b}$ be a primitive Dirichlet character. Under GRH for $L(s, \chi)$, we have*

$$\sum_{n \leq x} \Lambda(n) \chi(n) n^{-it} = O_\epsilon(x^{1/2} (bxt)^\epsilon) \tag{2-15}$$

for any $x \geq 2$ and $t \in \mathbb{R}$.

Proof. This follows from a standard argument similar to the one used to prove the prime number theorem; see [24, Chapter 5], [8], or [37, Chapter 13]. As a quick sketch, we have

$$\sum_{n \leq x} \Lambda(n) \chi(n) n^{-it} = \int_{\frac{3}{2}-ix}^{\frac{3}{2}+ix} \frac{L'}{L}(s+it, \chi) \frac{x^s}{s} ds + O(x^{1/2}) = \sum_{|\gamma-t| \leq x} \frac{x^{\rho-it}}{\rho-it} + O_\epsilon(x^{1/2} (bxt)^\epsilon)$$

using GRH and the estimate

$$\frac{L'}{L}(s, \chi) \ll (\log b|s|)^2 \tag{2-16}$$

for $-1 \leq \Re(s) \leq 2$. The desired result then follows from the fact that the number of zeros satisfying $u \leq \gamma - t \leq u + 1$ is $\ll \log(b(|u| + |t|))$. □

The Bessel functions of the first kind occur in the Petersson formula (see Lemma 2.8) and hence frequently in this paper. We collect some results for them.

Lemma 2.4. *Let $k \geq 2$ be an integer. The following bounds are satisfied for $x > 0$:*

- (1) $J_{k-1}(x) \ll_k 1$.
- (2) $J_{k-1}(x) \ll_k x$.
- (3) $J_{k-1}(x) \ll_k x^{k-1}$.
- (4) $J_{k-1}(x) \ll_k x^{-\frac{1}{2}}$.

Proof. See [18; 46]. □

We will also utilize the Mellin integral representation for the Bessel function.

Lemma 2.5. *We have*

$$J_{k-1}(x) = \frac{1}{2\pi i} \int_{\Re(s)=c} G_{k-1}(s)x^{-s} ds \tag{2-17}$$

for $x > 0$ and $1 - k < c < \frac{3}{2}$, where

$$G_{k-1}(s) := 2^{s-1} \Gamma\left(\frac{k-1+s}{2}\right) / \Gamma\left(\frac{k+1-s}{2}\right) \tag{2-18}$$

Proof. See [18, (6.561.14)]. □

2.3. Automorphic preliminaries. We collect the essential results from the standard references from [24, Chapter 14], [23, Chapters 6–7], [25, Sections 2 and 3].

Let k and N be positive integers with k even and N prime. Recall that $H_k^\bullet(N)$ ($\bullet \in \{+, -, *\}$) denotes the set of holomorphic cuspidal newforms of weight k and level N , depending on the sign of the functional equation. From [25, (2.73)], we have the dimension formulae

$$|H_k^\pm(N)| = \frac{1}{24}(k-1)N + O((kN)^{5/6}), \tag{2-19}$$

$$|H_k^*(N)| = \frac{1}{12}(k-1)N + O((kN)^{5/6}). \tag{2-20}$$

Every $f \in H_k^*(N)$ has a Fourier expansion of the form

$$f(z) = \sum_{n=1}^{\infty} \lambda_f(n)n^{\frac{k-1}{2}} e(nz) \tag{2-21}$$

for $z \in \mathbb{H} := \{x + iy : x \in \mathbb{R}, y > 0\}$, where $\lambda_f(1) = 1$. The L -function associated with f is defined by the Dirichlet series

$$L(s, f) = \sum_{n=1}^{\infty} \lambda_f(n)n^{-s} \tag{2-22}$$

which converges absolutely on $\operatorname{Re} s > 1$. The completed L -function is given by

$$\Lambda(s, f) := \left(\frac{\sqrt{N}}{2\pi}\right)^s \Gamma\left(s + \frac{k-1}{2}\right) L(s, f). \tag{2-23}$$

It admits an entire continuation and satisfies the functional equation

$$\Lambda(s, f) = \epsilon_f \Lambda(1-s, f). \tag{2-24}$$

The root number ϵ_f admits a nice formula in our case.

Lemma 2.6. *If $f \in H_k^*(N)$ and N is prime, then*

$$\epsilon_f = -i^k \lambda_f(N) \sqrt{N}. \tag{2-25}$$

In particular, we have $|\lambda_f(N)| = 1/\sqrt{N}$.

Proof. See [25, equation 3.5]. □

The following Hecke relations will be crucial in demonstrating that the number theoretic combinatorics align with those of random matrix theory.

Lemma 2.7. *Let $f \in H_k^*(N)$.*

(1) *For any $m, n \geq 1$,*

$$\lambda_f(m)\lambda_f(n) = \sum_{\substack{d|(m,n) \\ (d,N)=1}} \lambda_f\left(\frac{mn}{d^2}\right). \tag{2-26}$$

In particular, if $(m, n) = 1$ then

$$\lambda_f(m)\lambda_f(n) = \lambda_f(mn). \tag{2-27}$$

(2) *For a prime $p \nmid N$, we have*

$$\lambda_f(p)^2 = \lambda_f(p^2) + 1. \tag{2-28}$$

(3) *For $n \geq 1$ and a prime $p \nmid N$, we have*

$$\lambda_f(p)^n = \sum_{\alpha=0}^{\lfloor n/2 \rfloor} \left[\binom{n}{\alpha} - \binom{n}{\alpha-1} \right] \lambda_f(p^{n-2\alpha}). \tag{2-29}$$

Proof. Only the last property is less well-known; see [20] for its proof. □

Now, consider

$$\Delta_{k,N}^\bullet(n) := \sum_{f \in H_k^*(N)} \lambda_f(n), \quad \bullet \in \{+, -, *\}. \tag{2-30}$$

Splitting by sign with Lemma 2.6, we have

$$\Delta_{k,N}^\pm(n) = \sum_{f \in H_k^*(N)} \frac{1}{2} (1 \pm \epsilon_f) \lambda_f(n) = \frac{1}{2} \Delta_{k,N}^*(n) \mp \frac{i^k \sqrt{N}}{2} \Delta_{k,N}^*(nN) \tag{2-31}$$

whenever N is a prime and $(n, N) = 1$. We have the following useful form of the Petersson formula.

Lemma 2.8. *If N is prime and $(n, N^2) \mid N$ then*

$$\Delta_{k,N}^*(n) = \Delta'_{k,N}(n) + \Delta_{k,N}^\infty(n), \tag{2-32}$$

where

$$\Delta'_{k,N}(n) = \frac{(k-1)N}{12\sqrt{n}} \delta_{n, \square_Y} + \frac{(k-1)N}{12} \sum_{\substack{(m,N)=1 \\ m \leq Y}} \frac{2\pi i^k}{m} \sum_{\substack{c \equiv 0 \pmod N \\ c \geq N}} \frac{S(m^2, n; c)}{c} J_{k-1} \left(4\pi \frac{\sqrt{m^2 n}}{c} \right) \tag{2-33}$$

with $\delta_{n, \square_Y} = 1$ only if $n = m^2$ with $m \leq Y$ and 0 otherwise, and $\Delta_{k,N}^\infty(n)$ defined in [25, Lemma 2.12].

Proof. See [25, Propositions 2.1, 2.11 and 2.15]. □

Remark 2.9. The piece $\Delta_{k,N}^\infty(n)$ is called the *complementary sum*. By [21, Lemma A.1], assuming GRH for all $L(s, f)$ with $f \in H_k^*(1) \cup H_k^*(N)$, the complementary sum does not contribute in all cases appearing in this paper.

We have the following lemma.

Lemma 2.10. *Assume $(n, N) = 1$. Then*

$$\frac{1}{|H_k^*(N)|} \Delta'_{k,N}(Nn) \ll \sqrt{n} N^{-\frac{3}{2} + \epsilon}. \tag{2-34}$$

Proof. We take $Y = N^\epsilon$ and write $c = bN$ for $c \equiv 0 \pmod N$. Using (2-19), the Weil bound (2-11) and the bound $J_{k-1}(x) \ll x$ from Lemma 2.4, the result follows immediately. □

2.4. Density and moment sums. Let $f \in H_k^*(N)$ and $\phi \in \mathcal{S}_{ec}(\mathbb{R})$. Substituting the explicit formula for $L(s, f)$ (see [25, Section 4]) into the one-level density function

$$D(f; \phi) := \sum_{\gamma_f} \phi \left(\frac{\log R}{2\pi} \gamma_f \right), \tag{2-35}$$

we have

$$D(f; \phi) = \hat{\phi}(0) + \frac{1}{2} \phi(0) - P(f; \phi) + O \left(\frac{\log \log R}{\log R} \right), \tag{2-36}$$

where $R = k^2 N$ and

$$P(f; \phi) := \sum_{p \nmid N} \lambda_f(p) \hat{\phi} \left(\frac{\log p}{\log R} \right) \frac{2 \log p}{\sqrt{p} \log R}. \tag{2-37}$$

See [25, (4.25)] and the relevant remarks of [25, pp. 88] and [21, pp. 129]. Using (2-36), [21] expresses the n -th centered moments in terms of sums over primes (see [21, Section 2.3]).

Lemma 2.11. *If $\text{supp}(\hat{\phi}) \subset (-1, 1)$, we have*

$$\lim_{\substack{N \rightarrow \infty \\ N \text{ prime}}} \mathcal{D}_n^\pm(N; \phi) = (-1)^n \lim_{\substack{N \rightarrow \infty \\ N \text{ prime}}} S_1^{(n)}(N) \pm (-1)^{n+1} \lim_{\substack{N \rightarrow \infty \\ N \text{ prime}}} S_2^{(n)}(N) \tag{2-38}$$

provided the limits on the right side exist, where

$$S_1^{(n)}(N) := \sum_{p_1 \nmid N, \dots, p_n \nmid N} \prod_{j=1}^n \left(\hat{\phi} \left(\frac{\log p_j}{\log R} \right) \frac{2 \log p_j}{\sqrt{p_j} \log R} \right) \left\langle \prod_{j=1}^n \lambda_f(p_j) \right\rangle_* \tag{2-39}$$

and

$$S_2^{(n)}(N) := i^k \sqrt{N} \sum_{p_1 \nmid N, \dots, p_n \nmid N} \prod_{j=1}^n \left(\hat{\phi} \left(\frac{\log p_j}{\log R} \right) \frac{2 \log p_j}{\sqrt{p_j} \log R} \right) \left\langle \lambda_f(N) \prod_{j=1}^n \lambda_f(p_j) \right\rangle_* \tag{2-40}$$

Proposition 2.12. *Under the same assumptions of Theorem 1.1, we have*

$$\lim_{\substack{N \rightarrow \infty \\ N \text{ prime}}} S_1^{(n)}(N) = \begin{cases} (n-1)!! (\sigma_\phi^2)^{n/2} & \text{if } n \text{ is even,} \\ 0 & \text{if } n \text{ is odd,} \end{cases} \tag{2-41}$$

for $\phi \in \mathcal{S}_{ec}(\mathbb{R})$ with $\text{supp}(\hat{\phi}) \subset (-\frac{2}{n}, \frac{2}{n})$, where σ_ϕ^2 is defined in (1-13).

Proof. See Appendix B. □

Remark 2.13. Theorem E.1 of [21] is an analogous result but with an extra restriction $n \leq 2k$.

In Sections 3 and 4, we evaluate $S_2^{(n)}(N)$. The main result is Proposition 3.8, in which we express $S_2^{(n)}(N)$ in terms of $S(n, a; \phi)$ (see (1-16)).

3. Proof of Theorem 1.2 assuming Propositions 3.6 and 3.7

In this section, we prove Theorem 1.2 assuming the key Propositions 3.6 and 3.7. We prove these propositions in Section 4. First we decompose $S_2^{(n)}(N)$ (see (2-40)) into subterms and show that many of these subterms vanish as $N \rightarrow \infty$ through primes. Then in Section 3.4, we apply Propositions 3.6 and 3.7 to complete the proof of Theorem 1.2.

3.1. Combinatorial expansion and cleaning. We rewrite the sums over primes in (2-40) as sums over powers of *distinct* primes. This facilitates the applications of the Hecke relations (Lemma 2.7) and the Petersson formula (Lemma 2.8). More precisely, suppose $p_1 \cdots p_n = q_1^{n_1} \cdots q_\ell^{n_\ell}$ in (2-40), where q_1, \dots, q_ℓ are *distinct* primes, $n \geq 2$ and $\ell \geq 1$. We have $\langle \lambda_f(N) \prod_{j=1}^n \lambda_f(p_j) \rangle_* = \langle \lambda_f(N) \prod_{j=1}^\ell \lambda_f(q_j)^{n_j} \rangle_*$. We then apply the Hecke relations (2-29) to each $\lambda_f(q_j)^{n_j}$ and then use (2-27). Then we remove the distinctness condition from the sum over primes using a delicate inclusion-exclusion process. We conclude this process with the following lemma.

Lemma 3.1. *We have*

$$S_2^{(n)}(N) = \sum_{0 \leq \omega \leq n} \sum_{0 \leq n' \leq n} \sum_{\substack{\vec{n} := (n_1, \dots, n_\omega) \\ n_j > 1 \\ n_1 + \dots + n_\omega = n'}} \sum_{\substack{\vec{m} := (m_1, \dots, m_\omega) \\ m_j \equiv n_j \pmod{2} \\ 0 \leq m_j < n_j}} C_{\vec{n}, \vec{m}} E(\vec{n}, \vec{m}) \tag{3-1}$$

for some explicit constants $C_{\vec{n}, \vec{m}}$, where¹

$$E(\vec{n}, \vec{m}) := i^k \sqrt{N} \sum_{q_1 \nmid N, \dots, q_\omega \nmid N} \prod_{j=1}^{\omega} \left(\hat{\phi} \left(\frac{\log q_j}{\log R} \right)^{n_j} \left(\frac{2 \log q_j}{\sqrt{q_j} \log R} \right)^{n_j} \right) \times \sum_{p_1 \nmid N, \dots, p_{n-n'} \nmid N} \prod_{i=1}^{n-n'} \hat{\phi} \left(\frac{\log p_i}{\log R} \right) \frac{2 \log p_i}{\sqrt{p_i} \log R} \langle \lambda_f(NQ) \rangle_* \quad (3-2)$$

and

$$Q := \prod_{j=1}^{\omega} q_j^{m_j} \cdot \prod_{i=1}^{n-n'} p_i. \quad (3-3)$$

Lemma 3.1 is quite involved combinatorially, and we prove it in Appendix A, carefully decomposing $S_2^{(n)}(N)$ into sums over distinct primes in order to establish the condition $m_j < n_j$ in (3-1).

Lemma 3.1 allows us to apply the Petersson trace formula in the following section, as we have expressed $S_2^{(n)}(N)$ in terms of the average of a single Fourier coefficient $\lambda_f(NQ)$, as opposed to the product of Fourier coefficients in the definition (2-40). The coefficients $C_{\vec{n}, \vec{m}}$ are difficult to calculate in general, but as a consequence of Proposition 3.6 we only need to determine them in specific cases (see (3-24)).

3.2. Cleaning with Weil’s bound and the prime number theorem (PNT). The following result states that $E(\vec{n}, \vec{m})$ contributes to $\lim_{N \rightarrow \infty} \sum_{N \text{ prime}} D_n^{\pm}(N; \phi)$ only if “most” of the indices satisfy $n_j = m_j$.

Lemma 3.2. *Suppose $\text{supp}(\hat{\phi}) \subset (-\frac{1}{n-a}, \frac{1}{n-a})$ for some $a \leq n/2$. If $n' \geq a$, then $E(\vec{n}, \vec{m}) = O(N^{-\epsilon})$.*

Proof. Let $\text{supp}(\hat{\phi}) \subset (-\sigma, \sigma)$ with $\sigma < 1/(n-a)$. Using Lemma 2.10 and the PNT (with partial summation), the sum over $p_1, \dots, p_{n-n'}$ in (3-2) is $\ll N^{-3/2+\epsilon} (q_1^{m_1} \dots q_\omega^{m_\omega})^{1/2} R^{\sigma(n-n')}$. For $1 \leq j \leq \omega$, we have $n_j - m_j \geq 2$ as $m_j < n_j$ and $m_j \equiv n_j \pmod{2}$. Using the PNT again for the sums over q_1, \dots, q_ω in (3-2), we have $E(\vec{n}, \vec{m}) \ll N^{-1+\epsilon} R^{\sigma(n-n')}$. This is $O(N^{-\epsilon})$ if $n' \geq a$. \square

We are now in a position to apply the Petersson formula (Lemma 2.8) to (3-2). We assume GRH for $L(s, f)$ so that the complementary sum $\Delta_{k,N}^{\infty}$ does not contribute (see Remark 2.9). Since $|H_k^*(N)| \sim N(k-1)/12$ from (2-20), it follows that

$$E(\vec{n}, \vec{m}) = \frac{2^{n+1} \pi}{\sqrt{N}} \sum_{q_1 \nmid N, \dots, q_\omega \nmid N} \prod_{j=1}^{\omega} \left(\hat{\phi} \left(\frac{\log q_j}{\log R} \right) \frac{\log q_j}{\sqrt{q_j} \log R} \right)^{n_j} \sum_{p_1 \nmid N, \dots, p_{n-n'} \nmid N} \prod_{i=1}^{n-n'} \hat{\phi} \left(\frac{\log p_i}{\log R} \right) \frac{\log p_i}{\sqrt{p_i} \log R} \times \sum_{m \leq N^\epsilon} \frac{1}{m} \sum_{b=1}^{\infty} \frac{S(m^2, NQ; Nb)}{b} J_{k-1} \left(\frac{4\pi m \sqrt{Q}}{b \sqrt{N}} \right) + O(N^{-\epsilon}), \quad (3-4)$$

where Q was defined in (3-3) and we set $Y = N^\epsilon$. Also, recall that $n' = n_1 + \dots + n_\omega$.

¹We have omitted the dependence on ω and n' in the notation $E(\vec{n}, \vec{m})$ as it is implicitly contained in \vec{n}, \vec{m} . We have also suppressed the dependence on N for ease of notation.

We impose restrictions to the b -sum of (3-4) using the following two lemmas. We will also make use of the following bounds from Section 2:

$$S(m^2, NQ; Nb) \ll_{\epsilon} m^2 b^{1/2} (bN)^{\epsilon}, \quad J_{k-1}(x) \ll_k x. \tag{3-5}$$

Lemma 3.3. *If $\text{supp}(\hat{\phi}) \subseteq \left(-\frac{5}{2(n-n')}, \frac{5}{2(n-n')}\right)$, then the contribution from the terms in (3-4) with $(b, N) > 1$ is $O(N^{-\epsilon})$.*

Proof. This is a refinement of [21, Lemma 4.4]. Let $\text{supp}(\hat{\phi}) \subset (-\sigma, \sigma)$. If the b -sum of (3-4) is restricted to $(b, N) > 1$, i.e., $b = cN$ for some $c \geq 1$, then such a sum is $\ll N^{-2+\epsilon} \sqrt{Q}$ using (3-5). Combining this with the PNT and the fact that $n_j - m_j \geq 2$, we find that the contribution to (3-4) from terms with $(b, N) > 1$ is $\ll N^{-5/2+\epsilon} R^{(n-n')\sigma}$, which is negligible when $\sigma < \frac{5}{2(n-n')}$. □

Lemma 3.4. *If $\text{supp}(\hat{\phi}) \subset \left(-\frac{1000}{n-n'}, \frac{1000}{n-n'}\right)$, then the contribution from the terms in (3-4) with $b \geq N^{2022}$ is $O(N^{-12})$.*

Proof. This is a refinement of [21, Lemma 4.5]. Let $\text{supp}(\hat{\phi}) \subset (-\sigma, \sigma)$. If the b -sum of (3-4) is restricted to $b \geq N^{2022}$, then (3-5) implies such a sum is

$$\sum_{b \geq N^{2022}} \frac{S(m^2, NQ; Nb)}{b} J_{k-1}\left(\frac{4\pi m}{b\sqrt{N}}\right) \ll N^{-1/2+\epsilon} \sum_{b \geq N^{2022}} b^{-3/2+\epsilon} \ll b^{-1011-1/2+\epsilon}. \tag{3-6}$$

When $\sigma < \frac{1000}{n-n'}$, the contribution of (3-4) with $b \geq N^{2022}$ is $\ll N^{-1012+\epsilon} R^{(n-n')\sigma} \ll N^{-12}$ using (3-6), the PNT, and the fact that $n_j - m_j \geq 2$. This completes the proof. □

3.3. Expanding the Kloosterman sums. Suppose $\text{supp}(\hat{\phi}) \subset \left(-\frac{2}{n}, \frac{2}{n}\right)$ for $n \geq 2$. In particular, the assumptions of Lemma 3.3 and 3.4 are satisfied and we may impose the relevant restrictions on the b -sum of (3-4). Also, because $R = k^2 N$ and $\text{supp}(\hat{\phi}) \subset (-1, 1)$, the conditions $q_j \nmid N$, $p_i \nmid N$ in (3-4) are automatically satisfied provided the primes N are *sufficiently large*, and thus they will be dropped subsequently. Now, Lemma 2.2 allows us to convert the Kloosterman sums in (3-4) into Gauss sums. We thus obtain

$$\begin{aligned} E(\vec{n}, \vec{m}) &= -\frac{2^{n+1}\pi}{\sqrt{N}} \sum_{q_1, \dots, q_{\omega}} \prod_{j=1}^{\omega} \left(\hat{\phi}\left(\frac{\log q_j}{\log R}\right) \frac{\log q_j}{\sqrt{q_j} \log R} \right)^{n_j} \sum_{p_1, \dots, p_{n-n'}} \prod_{i=1}^{n-n'} \hat{\phi}\left(\frac{\log p_i}{\log R}\right) \frac{\log p_i}{\sqrt{p_i} \log R} \tag{3-7} \\ &\times \sum_{m \leq N^{\epsilon}} \frac{1}{m} \sum_{\substack{(b, N)=1 \\ b < N^{2022}}} \frac{1}{b\varphi(b)} \sum_{\chi(b)} G_{\chi}(m^2) G_{\chi}((Q, b^{\infty})) \bar{\chi}\left(\frac{Q}{(Q, b^{\infty})}\right) \chi(N) J_{k-1}\left(\frac{4\pi m \sqrt{Q}}{b\sqrt{N}}\right) + O(N^{-\epsilon}). \end{aligned}$$

Denote by $E(\vec{n}, \vec{m})|_{\chi \neq \chi_0}$ the expression (3-7) but with an extra restriction $\chi \neq \chi_0$ in the sum over $\chi(b)$. The following lemma shows that $E(\vec{n}, \vec{m})|_{\chi \neq \chi_0}$ contributes negligibly as $N \rightarrow \infty$. The shape of the expansion (2-12) allows us to capture cancellations in two ways: first, from the sum over $\chi(b)$ via

$$\frac{1}{\varphi(b)} \sum_{\chi(b)} |G_{\chi}(x) G_{\chi}(y)| \ll b, \tag{3-8}$$

and second, the character sums over primes (see Lemma 2.3). The former follows simply from Cauchy-Schwarz’s inequality and the orthogonality of characters, and the uniformity of (3-8) in x, y is important. For the latter, we crucially make use of GRH for the Dirichlet L -functions, and of course the restriction $\text{supp}(\hat{\phi}) \subset (-\frac{2}{n}, \frac{2}{n})$. Additionally, the following proposition corrects an error made in [21, Lemma 4.7].

Lemma 3.5. *Assume GRH for the Dirichlet L -functions. If $\text{supp}(\hat{\phi}) \subset (-\frac{2}{n}, \frac{2}{n})$, then*

$$E(\vec{n}, \vec{m})|_{\chi \neq \chi_0} = O(N^{-\epsilon}). \tag{3-9}$$

Proof. Upon rearranging the sums and taking absolute values, we find $E(\vec{n}, \vec{m})|_{\chi \neq \chi_0}$ is bounded by

$$\begin{aligned} & \max_{0 \leq \alpha \leq n-n'} \frac{1}{\sqrt{N}} \sum_{\substack{b < N^{2022} \\ (b, N)=1}} \frac{1}{b} \sum_{q_1, \dots, q_\omega} \prod_{j=1}^\omega \left(\left| \hat{\phi} \left(\frac{\log q_j}{\log R} \right) \right| \frac{\log q_j}{\sqrt{q_j} \log R} \right)^{n_j} \sum_{p_1|b, \dots, p_\alpha|b} \prod_{i=1}^\alpha \left| \hat{\phi} \left(\frac{\log p_i}{\log R} \right) \right| \frac{\log p_i}{\sqrt{p_i} \log R} \\ & \times \sum_{m \leq N^\epsilon} \frac{1}{m} \cdot \frac{1}{\varphi(b)} \sum_{\substack{\chi(b) \\ \chi \neq \chi_0}} |G_\chi(m^2) G_\chi((Q, b^\infty))| \\ & \times \left| \sum_{p_{\alpha+1} \nmid b, \dots, p_{n-n'} \nmid b} \prod_{\ell=\alpha+1}^{n-n'} \hat{\phi} \left(\frac{\log p_\ell}{\log R} \right) \frac{\bar{\chi}(p_\ell) \log p_\ell}{\sqrt{p_\ell} \log R} J_{k-1} \left(\frac{4\pi m \sqrt{Q}}{b\sqrt{N}} \right) \right|. \end{aligned} \tag{3-10}$$

We estimate the sum over $p_{\alpha+1}, \dots, p_{n-n'}$ in (3-10) with Lemma 2.3. To separate variables, we plug in (2-17) with $s = -1 + \epsilon + it$. Interchanging the order of sums and integrals and taking absolute values, we have

$$\begin{aligned} & \sum_{p_{\alpha+1} \nmid b, \dots, p_{n-n'} \nmid b} \prod_{\ell=\alpha+1}^{n-n'} \hat{\phi} \left(\frac{\log p_\ell}{\log R} \right) \frac{\bar{\chi}(p_\ell) \log p_\ell}{\sqrt{p_\ell} \log R} J_{k-1} \left(\frac{4\pi m \sqrt{Q}}{b\sqrt{N}} \right) \\ & \ll \int_{-\infty}^\infty \left(\frac{4\pi m \sqrt{Q'}}{b\sqrt{N}} \right)^{1-\epsilon} |G_{k-1}(1 - \epsilon + it)| \left| \sum_{p \nmid b} \hat{\phi} \left(\frac{\log p}{\log R} \right) \frac{\bar{\chi}(p) \log p}{p^{(\epsilon+it)/2} \log R} \right|^{n-n'-\alpha} dt, \end{aligned} \tag{3-11}$$

where

$$Q' := \prod_{j=1}^\omega q_j^{m_j} \cdot \prod_{i=1}^\alpha p_i \tag{3-12}$$

and $G_{k-1}(\cdot)$ was given by (2-18). Suppose $\text{supp}(\hat{\phi}) \subset (-\sigma, \sigma)$. By Lemma 2.3 (with partial summation) and the fact that

$$\sum_{p|b} \frac{\log p}{\log R} \leq \frac{\log b}{\log R}, \tag{3-13}$$

we find

$$\sum_{p \nmid b} \hat{\phi} \left(\frac{\log p}{\log R} \right) \frac{\bar{\chi}(p) \log p}{p^{(\epsilon+it)/2} \log R} \ll R^{\sigma/2} (Rbt)^\epsilon. \tag{3-14}$$

By the Stirling formula $|\Gamma(x + iy)| \asymp_x (1 + |y|)^{x-1/2} e^{-\frac{\pi}{2}|y|}$, which holds for any $x \in \mathbb{R} - \{0, -1, -2, \dots\}$ and $y \in \mathbb{R}$ (see [24, equation 5.113]), we have

$$|G_{k-1}(1 - \epsilon + it)| \ll_{k,\epsilon} (1 + |t|)^{-2+\epsilon} \tag{3-15}$$

for any $t \in \mathbb{R}$. Applying (3-14) and (3-15) to (3-11), we have

$$\sum_{p_{\alpha+1} \nmid b, \dots, p_{n-n'} \nmid b} \prod_{\ell=\alpha+1}^{n-n'} \hat{\phi} \left(\frac{\log p_\ell}{\log R} \right) \frac{\bar{\chi}(p_\ell) \log p_\ell}{\sqrt{p_\ell} \log R} J_{k-1} \left(\frac{4\pi m \sqrt{Q}}{b\sqrt{N}} \right) \ll \left(\frac{m\sqrt{Q}}{b\sqrt{N}} \right)^{1-\epsilon} R^{\sigma(n-n')/2} (Rb)^\epsilon. \tag{3-16}$$

From (3-10), (3-12) and (3-16), we have that $E(\vec{n}, \vec{m})|_{\chi \neq \chi_0}$ is bounded by

$$\begin{aligned} \max_{0 \leq \alpha \leq n-n'} \frac{R^{\epsilon+\sigma(n-n'-\alpha)/2}}{N} \sum_{b < N^{2022}} \frac{1}{b^2} \sum_{q_1, \dots, q_\omega < R^\sigma} \prod_{j=1}^\omega \left(\frac{\log q_j}{\sqrt{q_j} \log R} \right)^{n_j} q_j^{m_j/2} \sum_{p_1|b, \dots, p_\alpha|b} \prod_{i=1}^\alpha \frac{\log p_i}{\log R} \\ \times \sum_{m \leq N^\epsilon} m^{-\epsilon} \cdot \frac{1}{\varphi(b)} \sum_{\substack{\chi(b) \\ \chi \neq \chi_0}} |G_\chi(m^2) G_\chi((Q, b^\infty))| \end{aligned} \tag{3-17}$$

Applying (3-8), $\sum_{m \leq N^\epsilon} m^{-\epsilon} \ll R^\epsilon$, and (3-13) in sequence, the expression above is further bounded by

$$\max_{0 \leq \alpha \leq n-n'} \frac{R^{\epsilon+\sigma(n-n'-\alpha)/2}}{N} \sum_{q_1, \dots, q_\omega < R^\sigma} \prod_{j=1}^\omega \left(\frac{\log q_j}{\sqrt{q_j} \log R} \right)^{n_j} q_j^{m_j/2} \sum_{b < N^{2022}} \frac{1}{b} \left(\frac{\log b}{\log R} \right)^{n-n'}. \tag{3-18}$$

The contribution of the b -sum is $O(R^\epsilon)$. Since $n_j - m_j \geq 2$, it follows from the PNT that

$$E(\vec{n}, \vec{m})|_{\chi \neq \chi_0} \ll N^\epsilon N^{\sigma(n-n')/2-1}, \tag{3-19}$$

which is negligible if $\sigma < 2/(n - n')$. The result follows. □

We apply Lemma 3.5 to (3-7). This leaves only the contribution from $\chi_0 \pmod{b}$ for each $b < N^{2022}$ and $(b, N) = 1$. Note that $G_{\chi_0}(x) = R(x, b)$ is the Ramanujan sum and $\chi_0(N) = 1$. Hence, we have under GRH that

$$\begin{aligned} E(\vec{n}, \vec{m}) := & -\frac{2^{n+1}\pi}{\sqrt{N}} \sum_{q_1, \dots, q_\omega} \prod_{j=1}^\omega \left(\hat{\phi} \left(\frac{\log q_j}{\log R} \right) \frac{\log q_j}{\sqrt{q_j} \log R} \right)^{n_j} \sum_{p_1, \dots, p_{n-n'}} \prod_{i=1}^{n-n'} \hat{\phi} \left(\frac{\log p_i}{\log R} \right) \frac{\log p_i}{\sqrt{p_i} \log R} \\ & \times \sum_{m \leq N^\epsilon} \frac{1}{m} \sum_{\substack{(b, N)=1 \\ b < N^{2022}}} \frac{R(m^2, b) R((Q, b^\infty), b)}{b\varphi(b)} \chi_0 \left(\frac{Q}{(Q, b^\infty)} \right) J_{k-1} \left(\frac{4\pi m \sqrt{Q}}{b\sqrt{N}} \right), \\ & + O(N^{-\epsilon}), \end{aligned} \tag{3-20}$$

provided $\text{supp}(\hat{\phi}) \subset (-\frac{2}{n}, \frac{2}{n})$.

We complete the calculation of $E(\vec{n}, \vec{m})$ with the following two propositions, which we prove in Section 4.

Proposition 3.6. *Let $E(\vec{n}, \vec{m})$ be as in (3-2) and suppose that there exists some $1 \leq j \leq \omega$ for which $n_j + m_j > 2$. Under the same assumptions of Theorem 1.1, if $\text{supp}(\hat{\phi}) \subset (-\frac{2}{n}, \frac{2}{n})$ with $n \geq 2$, then $E(\vec{n}, \vec{m}) = O(1/\log N)$.*

Proposition 3.6 shows that many of the terms in the expansion (3-1) do not contribute in the limit. We complete the proof of Proposition 3.6 in Section 4.3.

Proposition 3.7. *Let $E(\vec{n}, \vec{m})$ be as in (3-2) and suppose $\omega = n' = 0$. Under the same assumptions of Theorem 1.1, if $\text{supp}(\hat{\phi}) \subset (-\frac{1}{n-a}, \frac{1}{n-a})$ with $1 \leq a \leq n/2$, then*

$$E(\vec{n}, \vec{m}) = (-1)^{n+1} \mathcal{R}(n, a; \phi) + O\left(\frac{(\log \log N)^2}{(\log N)^{1/2}}\right), \tag{3-21}$$

where $R(n, a; \phi)$ was defined in (1-13).

We complete the proof of Proposition 3.7 in Section 4.5.

3.4. Concluding the proof of Theorem 1.2. We will now demonstrate how to use Propositions 3.6 and 3.7 to evaluate $S_2^{(n)}(N)$.

Proposition 3.8. *Let $S(n, a; \phi)$ be defined in (1-15). Under the same assumptions of Theorem 1.1, if $\text{supp}(\hat{\phi}) \subset (-\frac{1}{n-a}, \frac{1}{n-a})$ with $1 \leq a \leq n/2$, then*

$$S_2^{(n)}(N) = (-1)^{n+1} S(n, a; \phi) + O\left(\frac{(\log \log N)^2}{(\log N)^{1/2}}\right). \tag{3-22}$$

Proof. By Lemma 3.1 and Proposition 3.6, $S_2^{(n)}(N)$ is a sum of terms of the form $E(\vec{n}, \vec{m})$ with $n_j + m_j \leq 2$ for each j up to an error of $O(1/\log N)$. Since $n_j \equiv m_j \pmod{2}$, then $n_j = m_j = 1$ or $n_j = 2$ and $m_j = 0$ for each j . Let $E_\ell(N)$ denote the term $E(\vec{n}, \vec{m})$ in which $n_j = 2$ and $m_j = 0$ for exactly ℓ values of j . We have

$$E_\ell(N) = \sum_{q_1 \nmid N, \dots, q_\ell \nmid N} \prod_{j=1}^{\ell} \left(\hat{\phi} \left(\frac{\log q_j}{\log R} \right)^2 \left(\frac{2 \log q_j}{\sqrt{q_j} \log R} \right)^2 \right) \times i^k \sqrt{N} \sum_{p_1 \nmid N, \dots, p_{n-2\ell} \nmid N} \prod_{i=1}^{n-2\ell} \hat{\phi} \left(\frac{\log p_i}{\log R} \right) \left(\frac{2 \log p_i}{\sqrt{p_i} \log R} \right) \langle \lambda_f(N p_1 \cdots p_{n-2\ell}) \rangle_*. \tag{3-23}$$

By Lemma 3.2, the contribution to $S_2^{(n)}(N)$ from $E_\ell(N)$ with $\ell \geq a/2$ is negligible. We thus have

$$S_2^{(n)}(N) = \sum_{\ell=0}^{\lfloor \frac{a-1}{2} \rfloor} \frac{n!}{2^\ell (n-2\ell)! \ell!} E_\ell(N) + O(1/\log N). \tag{3-24}$$

The combinatorial factor multiplying $E_\ell(N)$ arises from choosing the indices of the primes for which $n_j = m_j = 1$, or $n_j = 2$ and $m_j = 0$. We choose the primes for which $n_j = m_j = 1$ in $\binom{n}{2\ell}$ ways, and put the remaining primes into pairs in $(2\ell - 1)!! = (2\ell)! / (\ell! 2^\ell)$ ways. Multiplying and simplifying gives the

desired combinatorial coefficient. By Proposition 3.7, we have

$$i^k \sqrt{N} \sum_{p_1 \nmid N, \dots, p_{n-2\ell} \nmid N} \prod_{i=1}^{n-2\ell} \hat{\phi}\left(\frac{\log p_i}{\log R}\right) \left(\frac{2 \log p_i}{\sqrt{p_i} \log R}\right) \langle \lambda_f(N p_1 \cdots p_{n-2\ell}) \rangle_* \\ = (-1)^{n+1} \mathcal{R}(n-2\ell, a-2\ell; \phi) + O\left(\frac{(\log \log N)^2}{(\log N)^{1/2}}\right). \tag{3-25}$$

Applying (3-25) to (3-23) and factoring the sums over q_j gives

$$E_\ell(N) = (-1)^{n+1} \left(\mathcal{R}(n-2\ell, a-2\ell; \phi) + O\left(\frac{(\log \log N)^2}{(\log N)^{1/2}}\right) \right) \left(\sum_{q \nmid N} \hat{\phi}\left(\frac{\log q}{\log R}\right)^2 \frac{4 \log^2 q}{q \log^2 R} \right)^\ell. \tag{3-26}$$

A standard partial summation argument gives

$$\sum_{q \nmid N} \hat{\phi}\left(\frac{\log q}{\log R}\right)^2 \frac{4 \log^2 q}{q \log^2 R} \sim 2 \int_{-\infty}^{\infty} |y| \hat{\phi}(y)^2 dy, \tag{3-27}$$

which is precisely the quantity σ_ϕ^2 given by (1-13). Applying (3-27) to (3-26), we have

$$E_\ell(N) = (-1)^{n+1} (\sigma_\phi^2)^\ell \mathcal{R}(n-2\ell, a-2\ell; \phi) + O\left(\frac{(\log \log N)^2}{(\log N)^{1/2}}\right). \tag{3-28}$$

The proposition follows readily upon applying this to (3-24) and comparing with (1-15). □

Proof of Theorem 1.2. The proof follows immediately from Proposition 3.8, (2-38), and (2-41). □

4. Proof of Propositions 3.6 and 3.7

Throughout Section 4, we assume $\text{supp}(\hat{\phi}) \subset (-\frac{1}{n-a}, \frac{1}{n-a})$ with $1 \leq a \leq n/2$ and we further analyze the expression (3-20) for $E(\vec{n}, \vec{m})$.

In Section 4.1, we rewrite the sums over primes in (3-20) into a more analytically tractable expression. The key quantity to be considered is

$$B(\alpha) := N^{-1/2} \sum_{p_{\alpha+1}, \dots, p_n} J_{k-1} \left(\frac{4\pi m}{b\sqrt{N}} \sqrt{c \prod_{i=\alpha+1}^n p_i} \right) \prod_{i=\alpha+1}^n \hat{\phi}\left(\frac{\log p_i}{\log R}\right) \frac{\chi_0(p_i) \log p_i}{p_i^{1/2} \log R}, \tag{4-1}$$

where the dependencies on N, m, c, b are conveniently suppressed in the notation $B(\alpha)$. We also set $\Phi_r(x) := \phi(x)^r$ for $r \geq 0$. The main result is the following.

Proposition 4.1. *Suppose $\text{supp}(\hat{\phi}) \subset (-\frac{1}{n-a}, \frac{1}{n-a})$ with $1 \leq a \leq n/2$. Under RH for $\zeta(s)$, we have*

$$B(\alpha) = \frac{b}{2\pi m \sqrt{c}} \sum_{\delta=0}^{a-\alpha-1} \binom{n-\alpha}{\delta} \sum_{i=0}^{a-\alpha-\delta-1} (-1)^i \binom{n-\alpha-\delta}{i} \sum_{p_1, \dots, p_\delta} \prod_{j=1}^{\delta} \hat{\phi}\left(\frac{\log p_j}{\log R}\right) \frac{\chi_0(p_j) \log p_j}{p_j \log R} \\ \times \int_{x=0}^{\infty} J_{k-1}(x) \hat{\Phi}_{n-\alpha-\delta} \left(\frac{2}{\log R} \log \left(\frac{bx}{4\pi m} \sqrt{N / \left(c \prod_{j=1}^{\delta} p_j \right)} \right) \right) \frac{dx}{\log R} + O(N^{-\epsilon}). \tag{4-2}$$

where the implicit constant does not depend on N, m, c, b, α .

The evaluation of $B(\alpha)$ requires a delicate inclusion-exclusion argument to convert the sums over primes of (4-1) into sums over integers which appears on the left side of (4-10). This is followed by Lemma 4.5, which expresses the sums over integers into integrals via a standard contour-shifting argument (assuming RH). Then the proof of Proposition 4.1 is completed by further combinatorial simplifications. The final step is crucial for matching the calculations with those from the random matrix theory (see Section 5).

One must also evaluate the convolution sum of *Ramanujan sums* in (3-20). This task will be carried out in Section 4.2 and the main result is stated as follows.

Proposition 4.2. *Let $\phi \in \mathcal{S}_{ec}(\mathbb{R})$. Then as $N \rightarrow \infty$, we have*

$$\begin{aligned} & \sum_{(b,M)=1} \frac{R(1,b)R(m^2,b)}{\varphi(b)} \int_0^\infty J_{k-1}(y) \hat{\phi}\left(\frac{2 \log(by\sqrt{Q}/4\pi m)}{\log R}\right) \frac{dy}{\log R} \\ &= \delta\left(\frac{m}{(m, M^\infty)}, 1\right) \frac{\varphi(M)}{M} \left(-\frac{1}{2} \int_{-\infty}^\infty \phi(x) \sin\left(2\pi x \frac{\log(k^2 Q/16\pi^2 m^2)}{\log R}\right) \frac{dx}{2\pi x} + \frac{1}{4}\phi(0)\right) \\ & \quad + O\left(m^\epsilon \frac{(\log \log M)^2}{(\log R)^{1/2}}\right), \end{aligned} \tag{4-3}$$

uniformly for $m, M, Q \geq 1$.

The contents of Sections 4.1 and 4.2 are independent of each other. The proofs of Propositions 3.6 and 3.7 crucially rely on Propositions 4.1 and 4.2. They are the subjects of Section 4.3 and Sections 4.4–4.5, respectively. It will be essential to break up the Ramanujan sums judiciously using the multiplicativity (2-9) and perform a prime-by-prime analysis with the exact evaluation (2-8). In Section 4.4, we managed to find an integral representation for $E(\vec{n}, \vec{m})$. The proof of Proposition 3.7 follows from the combinatorial simplification in Section 4.5.

4.1. Sums over primes: proof of Proposition 4.1. This subsection is dedicated to proving Proposition 4.1. We begin by defining the quantities

$$\begin{aligned} C'(\alpha, \beta) &:= N^{-1/2} \sum_{p_{\alpha+1}, \dots, p_n} \sum_{t_{\alpha+1}, \dots, t_{\alpha+\beta}=2}^\infty J_{k-1}\left(\frac{4\pi m}{b} \sqrt{c \prod_{i=\alpha+\beta+1}^n p_i} / \sqrt{N / \prod_{j=\alpha+1}^{\alpha+\beta} p_j^{t_j}}\right) \\ & \quad \times \prod_{j=\alpha+1}^{\alpha+\beta} \hat{\phi}\left(\frac{t_j \log p_j}{\log R}\right) \frac{\chi_0(p_j) \log p_j}{p_j^{t_j/2} \log R} \prod_{i=\alpha+\beta+1}^n \hat{\phi}\left(\frac{\log p_i}{\log R}\right) \frac{\chi_0(p_i) \log p_i}{p_i^{1/2} \log R} \end{aligned} \tag{4-4}$$

and

$$\begin{aligned} C(\alpha, \beta) &:= N^{-1/2} \sum_{p_{\alpha+1}, \dots, p_{\alpha+\beta}} \sum_{t_{\alpha+1}, \dots, t_{\alpha+\beta}=2}^\infty \sum_{v_{\alpha+\beta+1}, \dots, v_n=1}^\infty J_{k-1}\left(\frac{4\pi m}{b} \sqrt{c \prod_{i=\alpha+\beta+1}^n v_i} / \sqrt{N / \prod_{j=\alpha+1}^{\alpha+\beta} p_j^{t_j}}\right) df \\ & \quad \times \prod_{j=\alpha+1}^{\alpha+\beta} \hat{\phi}\left(\frac{t_j \log p_j}{\log R}\right) \frac{\chi_0(p_j) \log p_j}{p_j^{t_j/2} \log R} \prod_{i=\alpha+\beta+1}^n \hat{\phi}\left(\frac{\log v_i}{\log R}\right) \frac{\chi_0(v_i) \Lambda(v_i)}{v_i^{1/2} \log R}. \end{aligned} \tag{4-5}$$

Observe that $C'(\alpha, \beta)$ is a generalized version of $B(\alpha)$ (see (4-1)) as $B(\alpha) = C'(\alpha, 0)$. Also, the expression $C(\alpha, \beta)$ is obtained from $C'(\alpha, \beta)$ by replacing the sums over primes p_j 's, where $\alpha + \beta + 1 \leq j \leq n$, by sums over positive integers v_j 's, as well as the weight $\log p_j$ is replaced by von Mangoldt's function $\Lambda(v_j)$. We have the following relation between $C'(\alpha, \beta)$ and $C(\alpha, \beta)$.

Property 4.3.
$$C'(\alpha, \beta) = C(\alpha, \beta) - \sum_{i=1}^{a-1-\alpha-\beta} \binom{n-\alpha-\beta}{i} C'(\alpha, \beta+i) + O(N^{-\epsilon}).$$

Proof. The property follows directly from the definition of von Mangoldt's function and a partitioning argument. The sum over i in the statement of the proposition is restricted to $i \leq a-1-\alpha-\beta$ because the contribution from $i > a-1-\alpha-\beta$ is $O(N^{-\epsilon})$, which follows readily from $J_{k-1}(x) \ll 1$. \square

Property 4.3 can be applied repeatedly to obtain a relation between (4-1) and (4-5).

Property 4.4.
$$B(\alpha) = \sum_{i=0}^{a-1-\alpha} \binom{n-\alpha}{i} C(\alpha, i) (-1)^i + O(N^{-\epsilon}).$$

Proof. Define

$$B'(\alpha, \eta) := \sum_{i=0}^{\eta} (-1)^i \binom{n-\alpha}{i} C(\alpha, i) - \sum_{i=\eta+1}^{a-1-\alpha} \binom{n-\alpha}{i} C'(\alpha, i) \sum_{j=0}^{\eta} (-1)^j \binom{i}{j} + O(N^{-\epsilon}). \quad (4-6)$$

We claim that $B'(\alpha, \eta) = B(\alpha)$ and we proceed by induction on η . The base case $\eta = 0$ holds by Property 4.3 and the fact that $B(\alpha) = C'(\alpha, 0)$. For the inductive step, assume that $B'(\alpha, k) = B(\alpha)$ for some integer $k \geq 0$. This implies

$$B(\alpha) = \sum_{i=0}^k \binom{n-\alpha}{i} C(\alpha, i) (-1)^i - \sum_{i=k+1}^{a-1-\alpha} \binom{n-\alpha}{i} C'(\alpha, i) \sum_{j=0}^k (-1)^j \binom{i}{j} + O(N^{-\epsilon}). \quad (4-7)$$

We examine the term where $i = k+1$ and we have

$$-\binom{n-\alpha}{k+1} C'(\alpha, k+1) \sum_{j=0}^k (-1)^j \binom{k+1}{j} = \binom{n-\alpha}{k+1} (-1)^{k+1} C'(\alpha, k+1). \quad (4-8)$$

This follows from the identity $\sum_{j=0}^{k+1} (-1)^j \binom{k+1}{j} = 0$, an easy consequence of the binomial theorem. It follows from Property 4.3 and reindexing the sum using the change of variables $\ell = k+1+j$ that

$$\begin{aligned} & \binom{n-\alpha}{k+1} (-1)^{k+1} C'(\alpha, k+1) \\ &= \binom{n-\alpha}{k+1} (-1)^{k+1} \left(C(\alpha, \eta+1) - \sum_{j=1}^{a-\alpha-k} \binom{n-\alpha-k-1}{j} C'(\alpha, k+1+j) \right) + O(N^{-\epsilon}) \\ &= \binom{n-\alpha}{k+1} (-1)^{k+1} C(\alpha, k+1) - \sum_{j=1}^{a-\alpha-k} \binom{n-\alpha}{k+1+j} \binom{k+1+j}{k+1} (-1)^{k+1} C'(\alpha, k+1+j) + O(N^{-\epsilon}) \\ &= \binom{n-\alpha}{k+1} (-1)^{k+1} C(\alpha, k+1) - \sum_{\ell=k+2}^{a-\alpha-1} \binom{n-\alpha}{\ell} \binom{\ell}{k+1} (-1)^{k+1} C'(\alpha, \ell) + O(N^{-\epsilon}). \end{aligned} \quad (4-9)$$

Substituting the last equality into (4-7), we have $B(\alpha) = B'(\alpha, k+1)$ and this completes the induction. \square

The innermost sums of (4-5) can be rewritten as follows.

Lemma 4.5. *Under RH for $\zeta(s)$, if $\text{supp}(\hat{\phi}) \subset (-\frac{1}{n-a}, \frac{1}{n-a})$, then*

$$\begin{aligned} & \sum_{v_1, \dots, v_{n-\eta}} \left(\prod_{i=1}^{n-\eta} \hat{\phi} \left(\frac{\log v_i}{\log R} \right) \frac{\chi_0(v_i) \Lambda(v_i)}{\sqrt{v_i} \log R} \right) J_{k-1} \left(\frac{4\pi m \sqrt{c v_1 \cdots v_{n-\eta}}}{b \sqrt{N}} \right) \\ &= \frac{b \sqrt{N}}{2\pi m \sqrt{c}} \sum_{\gamma=0}^{a-\eta-1} \sum_{j=\gamma}^{a-\eta-1} (-1)^{j-\gamma} \binom{n-\eta}{j} \binom{j}{\gamma} \sum_{v_1, \dots, v_\gamma=1}^{\infty} \left(\prod_{i=1}^{\gamma} \hat{\phi} \left(\frac{\log v_i}{\log R} \right) \frac{\chi_0(v_i) \Lambda(v_i)}{v_i \log R} \right) \\ & \quad \times \int_{x=0}^{\infty} J_{k-1}(x) \hat{\Phi}_{n-\eta-\gamma} \left(\frac{2}{\log R} \log \left(\frac{bx}{4\pi m \sqrt{c v_1 \cdots v_\gamma}} \sqrt{\frac{N}{c v_1 \cdots v_\gamma}} \right) \right) \frac{dx}{\log R} + O(N^{1/2-\epsilon}), \end{aligned} \quad (4-10)$$

where the implicit constant does not depend on N, m, c, b .

Proof. This is a generalization of [21, Lemma 4.9]. Open up the J -Bessel function on the left side of (4-10) with the Mellin inversion formula (2-17):

$$\frac{1}{2\pi i} \int_{\Re(s)=1} \left(\sum_{v=1}^{\infty} \hat{\phi} \left(\frac{\log v}{\log R} \right) \frac{\chi_0(v) \Lambda(v)}{v^{(1+s)/2} \log R} \right)^{n-\eta} G_{k-1}(s) \left(\frac{4\pi m \sqrt{c}}{b \sqrt{N}} \right)^{-s} ds. \quad (4-11)$$

Under RH, a simple contour-shifting argument (or see [21, (4.34)]) gives

$$\sum_{v=1}^{\infty} \hat{\phi} \left(\frac{\log v}{\log R} \right) \frac{\chi_0(v) \Lambda(v)}{v^{(1+s)/2} \log R} = \phi \left(\frac{1-s}{4\pi i} \log R \right) + \mathcal{E}(s), \quad (4-12)$$

where

$$\mathcal{E}(s) := -\frac{1}{2\pi i} \int_{\Re(z)=3/4} \phi \left(\frac{(2z-1-s) \log R}{4\pi i} \right) \frac{L'}{L}(z, \chi_0) dz. \quad (4-13)$$

As a consequence, (4-11) becomes

$$\begin{aligned} & \frac{1}{2\pi i} \int_{\Re(s)=1} \left(\phi \left(\frac{1-s}{4\pi i} \log R \right) + \mathcal{E}(s) \right)^{n-\eta} G_{k-1}(s) \left(\frac{4\pi m \sqrt{c}}{b \sqrt{N}} \right)^{-s} ds \\ &= \sum_{\gamma=0}^{n-\eta} \binom{n-\eta}{\gamma} \frac{1}{2\pi i} \int_{\Re(s)=1} \phi \left(\frac{1-s}{4\pi i} \log R \right)^{n-\eta-\gamma} \mathcal{E}(s)^\gamma G_{k-1}(s) \left(\frac{4\pi m \sqrt{c}}{b \sqrt{N}} \right)^{-s} ds, \end{aligned} \quad (4-14)$$

where the last line follows from the binomial theorem. Now, [21, (4.43)] gives the bound

$$\frac{1}{2\pi i} \int_{\Re(s)=1} \phi \left(\frac{1-s}{4\pi i} \log R \right)^{n-\eta-\gamma} \mathcal{E}(s)^\gamma G_{k-1}(s) \left(\frac{4\pi m \sqrt{c}}{b \sqrt{N}} \right)^{-s} \ll N^{(n-\eta-\gamma)\sigma/2+\epsilon}, \quad (4-15)$$

which is in turn $O(N^{1/2-\epsilon})$ when $\gamma > a - \eta - 1$ and $\sigma < 1/(n-a)$. Hence, (4-11) is equal to

$$\sum_{\gamma=0}^{a-\eta-1} \binom{n-\eta}{\gamma} \int_{\Re(s)=1} \phi \left(\frac{1-s}{4\pi i} \log R \right)^{n-\eta-\gamma} \mathcal{E}(s)^\gamma G_{k-1}(s) \left(\frac{4\pi m \sqrt{c}}{b \sqrt{N}} \right)^{-s} \frac{ds}{2\pi i} + O(N^{1/2-\epsilon}). \quad (4-16)$$

For integers γ, n with $0 \leq \gamma < n$, we define

$$T(\gamma, n) := \int_{\Re(s)=1} \left(\sum_{v=1}^{\infty} \hat{\phi}\left(\frac{\log v}{\log R}\right) \frac{\chi_0(v)\Lambda(v)}{v^{(1+s)/2} \log R} \right)^{\gamma} \phi\left(\frac{1-s}{4\pi i} \log R\right)^{n-\gamma} G_{k-1}(s) \left(\frac{4\pi m\sqrt{c}}{b\sqrt{N}}\right)^{-s} \frac{ds}{2\pi i}. \tag{4-17}$$

It follows that (4-11) is given by

$$\begin{aligned} \sum_{\gamma=0}^{a-\eta-1} \binom{n-\eta}{\gamma} \sum_{j=0}^{\gamma} (-1)^{j-\gamma} \binom{\gamma}{j} T(j, n-\eta) + O(N^{1/2-\epsilon}) \\ = \sum_{\gamma=0}^{a-\eta-1} \sum_{j=\gamma}^{a-\eta-1} (-1)^{j-\gamma} \binom{n-\eta}{j} \binom{j}{\gamma} T(\gamma, n-\eta) + O(N^{1/2-\epsilon}), \end{aligned} \tag{4-18}$$

using the formula for $\mathcal{E}(s)$ in (4-12).

In (4-17), applying the formula $\int_0^{\infty} J_{k-1}(x)x^{s-1} dx = G_{k-1}(s)$ with the change of variable $s = 1 + it$, it follows that

$$\begin{aligned} T(\gamma, n) = \frac{b\sqrt{N}}{8\pi^2 m\sqrt{c}} \int_{t=-\infty}^{\infty} \phi\left(\frac{-t \log R}{4\pi}\right)^{n-\gamma} \left(\sum_{v_1, \dots, v_{\gamma}=1}^{\infty} \prod_{i=1}^{\gamma} \hat{\phi}\left(\frac{\log v_i}{\log R}\right) \frac{\chi_0(v_i)\Lambda(v_i)}{v_i^{it/2+1} \log R} \right) \\ \times \left(\frac{4\pi m\sqrt{c}}{b\sqrt{N}}\right)^{-it} \int_0^{\infty} J_{k-1}(x)x^{it} dx dt. \end{aligned} \tag{4-19}$$

Upon rearranging and a change of variables $u = -t \log R/(4\pi)$, we have

$$\begin{aligned} T(\gamma, n) = \frac{b\sqrt{N}}{2\pi m\sqrt{c}} \sum_{v_1, \dots, v_{\gamma}=1}^{\infty} \left(\prod_{i=1}^{\gamma} \hat{\phi}\left(\frac{\log v_i}{\log R}\right) \frac{\chi_0(v_i)\Lambda(v_i)}{v_i \log R} \right) \\ \times \int_{x=0}^{\infty} J_{k-1}(x) \hat{\Phi}_{n-\gamma} \left(\frac{2}{4\pi m} \log \left(\frac{bx}{\log R} \sqrt{\frac{N}{cv_1 \cdots v_{\gamma}}} \right) \right) \frac{dx}{\log R}. \end{aligned} \tag{4-20}$$

Now, (4-20) and (4-18) lead to the desired claim. □

Next we apply Lemma 4.5 to $C(\alpha, \beta)$.

Property 4.6. *We have*

$$\binom{n-\alpha}{\beta} C(\alpha, \beta) = \sum_{\gamma=0}^{a-\alpha-\beta-1} D(\alpha, \beta, \gamma) \left(\sum_{i=0}^{a-\alpha-\beta-\gamma-1} (-1)^i \binom{n-\alpha}{\gamma+\beta} \binom{n-\alpha-\beta-\gamma}{i} \binom{\gamma+\beta}{\gamma} \right) + O(N^{-\epsilon}),$$

where $D(\alpha, \beta, \gamma)$ is defined by

$$\begin{aligned} \frac{b}{2\pi m\sqrt{c}} \sum_{p_{\alpha+1}, \dots, p_{\alpha+\beta+\gamma}} \sum_{t_{\alpha+1}, \dots, t_{\alpha+\beta}=2}^{\infty} \sum_{t_{\alpha+\beta+1}, \dots, t_{\alpha+\beta+\gamma}=1}^{\infty} \prod_{j=\alpha+\beta+1}^{\alpha+\beta+\gamma} \hat{\phi}\left(\frac{t_j \log p_j}{\log R}\right) \frac{\chi_0(p_j) \log p_j}{p_j^{t_j} \log R} \\ \times \int_{x=0}^{\infty} J_{k-1}(x) \hat{\Phi}_{n-\alpha-\beta-\gamma} \left(\frac{2}{\log R} \log \frac{bx\sqrt{N''}}{4\pi m} \right) \frac{dx}{\log R}, \end{aligned} \tag{4-21}$$

with

$$N'' := \frac{N}{c p_{\alpha+1}^{t_{\alpha+1}} \cdots p_{\alpha+\beta+\gamma}^{t_{\alpha+\beta+\gamma}}}. \tag{4-22}$$

Proof. Applying Lemma 4.5 to (4-5), we have

$$C(\alpha, \beta) = \sum_{\gamma=0}^{a-\alpha-\beta-1} D(\alpha, \beta, \gamma) \left(\sum_{j=\gamma}^{a-\alpha-\beta-1} (-1)^{j-\gamma} \binom{n-\alpha-\beta}{j} \binom{j}{\gamma} \right) + O(N^{-\epsilon}). \tag{4-23}$$

The property follows from the reindexing $i = j - \gamma$ and simplifying. □

Remark 4.7. The three multiple sums of (4-21) can be interpreted as follows: the first α primes are those that divide b ; the next two sums involving β powers of primes are those left over from converting to sums over integers; and the last sums over γ integers are those leftover from applying Lemma 4.5.

We apply Property 4.6 to the formula for $B(\alpha)$ described in Property 4.4.

Property 4.8.

$$B(\alpha) = \sum_{\delta=0}^{a-\alpha-1} \binom{n-\alpha}{\delta} \sum_{i=0}^{a-\alpha-\delta-1} (-1)^i \binom{n-\alpha-\delta}{i} \sum_{\gamma=0}^{\delta} (-1)^{\delta-\gamma} \binom{\delta}{\gamma} D(\alpha, \delta - \gamma, \gamma) + O(N^{-\epsilon}). \tag{4-24}$$

Proof. The property from applying Property 4.6 to the equality in Property 4.4 and collecting terms with $\delta = \beta + \gamma$. □

We now eliminate the sums over prime powers in (4-21). Define

$$G(\alpha, \delta) := \frac{b}{2\pi m \sqrt{c}} \sum_{p_1, \dots, p_\delta} \int_{x=0}^{\infty} J_{k-1}(x) \hat{\Phi}_{n-\alpha-\delta} \left(\frac{2}{\log R} \log \left(\frac{bx}{4\pi m} \sqrt{\frac{N}{cp_1 \cdots p_\delta}} \right) \right) \frac{dx}{\log R} \\ \times \prod_{j=\alpha+1}^{\alpha+\delta} \hat{\phi} \left(\frac{\log p_j}{\log R} \right) \frac{\chi_0(p_j) \log p_j}{p_j \log R}, \tag{4-25}$$

which is obtained from (4-21) upon specializing all t_i 's to be 1 and reindexing. The expressions $D(\alpha, \beta, \gamma)$ and $G(\alpha, \delta)$ (with $\delta = \beta + \gamma$ as above) satisfy the following recursion:

Property 4.9.
$$G(\alpha, \delta) = \sum_{\gamma=0}^{\delta} (-1)^{\delta-\gamma} \binom{\delta}{\gamma} D(\alpha, \delta - \gamma, \gamma).$$

This can be deduced from a more general result:

Lemma 4.10. *Let $f(t_1, \dots, t_n)$ be a symmetric function taking as its input a finite sequence t_1, \dots, t_n of arbitrary length, and define a transform $\mathcal{T}(i, j)$ on f by*

$$\mathcal{T}(i, j)(f) := \sum_{t_1, \dots, t_i=2}^{\infty} \sum_{s_1, \dots, s_j=1}^{\infty} f(t_1, \dots, t_i, s_1, \dots, s_j). \tag{4-26}$$

Then, denoting by $[1]^n$ the sequence of 1's repeated n times, we have

$$\sum_{i=0}^n (-1)^i \binom{n}{i} \mathcal{T}(i, n-i)(f) = f([1]^n), \tag{4-27}$$

Proof. We proceed by induction on n . The base case $n = 1$ holds immediately. Assume the result holds up to n and define a new function $g(t_1, \dots, t_n) = f(t_1, \dots, t_n, 1)$. Then

$$\begin{aligned} f([1]^{n+1}) &= g([1]^n) = \sum_{i=0}^n (-1)^i \binom{n}{i} \sum_{t_1, \dots, t_i=2}^{\infty} \sum_{s_1, \dots, s_{n-i}=1}^{\infty} f(t_1, \dots, t_i, s_1, \dots, s_{n-i}, 1) \\ &= \sum_{i=0}^n \binom{n}{i} \left((-1)^i \mathcal{T}(i, n+1-i)(f) + (-1)^{i+1} \mathcal{T}(i+1, n+1-(i+1))(f) \right) \\ &= \sum_{i=0}^{n+1} (-1)^i \mathcal{T}(i, n+1-i)(f) \binom{n+1}{i}. \end{aligned} \tag{4-28}$$

This completes the induction. □

Proof of Property 4.9. A special case of Lemma 4.10: take $f([1]^\delta) = G(\alpha, \delta)$ and $\mathcal{T}(i, j) = D(\alpha, i, j)$. □

Applying Property 4.9 to (4-24) gives the following relation between $B(\alpha)$ and $G(\alpha, \delta)$.

Property 4.11. *We have*

$$B(\alpha) = \sum_{\delta=0}^{a-\alpha-1} \binom{n-\alpha}{\delta} \sum_{i=0}^{a-\alpha-\delta-1} (-1)^i \binom{n-\alpha-\delta}{i} G(\alpha, \delta). \tag{4-29}$$

Proof of Proposition 4.1. Substitute (4-25) into (4-29), the result follows. □

4.2. A convolution sum of Ramanujan sums: proof of Proposition 4.2. This is a generalization of [25, Section 7] and [21, Lemma 2.12]. We take this opportunity to correct typos and include more details that were omitted in previous works.

We first claim that the left side of (4-3) is equal to

$$\lim_{\epsilon \rightarrow 0} \sum_{(b, M)=1} \frac{R(1, b)R(m^2, b)}{\varphi(b)b^\epsilon} \int_0^\infty J_{k-1}(y) \hat{\phi} \left(\frac{2}{\log R} \log \frac{by\sqrt{Q}}{4\pi m} \right) \frac{dy}{\log R}. \tag{4-30}$$

Since $\hat{\phi}$ is compactly supported, the b -sum and the y -integral of (4-30) in total is bounded by

$$\sum_{b=1}^\infty \frac{m^4}{\phi(b)} \int_{y \ll \frac{mR^{O(1)}}{b\sqrt{Q}}} \frac{dy}{\log R} \ll \frac{m^5 R^{O(1)}}{\sqrt{Q} \log R} \sum_{b=1}^\infty \frac{1}{\varphi(b)b}, \tag{4-31}$$

where the bounds $|R(1, b)| = 1$, $|R(m^2, b)| \leq m^4$, $J_{k-1}(x) \ll 1$ (see (2-8) and Lemma 2.4) are used above. The last b -sum converges because $\varphi(b) \gg b/\log \log b$ (see [3, Theorem 13.14]). Our claim now readily follows from the dominated convergence theorem.

Applying the Mellin inversion formulae (2-2) and (2-18) to the integral of (4-30) gives

$$\int_0^\infty J_{k-1}(y) \hat{\phi} \left(\frac{2}{\log R} \log \frac{by\sqrt{Q}}{4\pi m} \right) \frac{dy}{\log R} = \int_{-\infty}^\infty \phi(x \log R) \left(\frac{2\pi m}{b\sqrt{Q}} \right)^{4\pi ix} \frac{\Gamma(\frac{k}{2} - 2\pi ix)}{\Gamma(\frac{k}{2} + 2\pi ix)} dx. \tag{4-32}$$

Substituting (4-32) into (4-30) and interchanging the sum and integral, we have

$$(4-30) = \lim_{\epsilon \rightarrow 0} \int_{-\infty}^\infty \phi(x \log R) \left(\frac{2\pi m}{\sqrt{Q}} \right)^{4\pi ix} \frac{\Gamma(\frac{k}{2} - 2\pi ix)}{\Gamma(\frac{k}{2} + 2\pi ix)} \chi_M(\epsilon + 4\pi ix; m) dx, \tag{4-33}$$

where the Dirichlet series

$$\chi_M(s; m) := \sum_{(b, M)=1} \frac{R(1, b)R(m^2, b)}{\varphi(b)b^s} \quad (4-34)$$

converges absolutely on the half-plane $\Re s > 0$.

We evaluate the integral of (4-33) asymptotically by breaking it into two pieces: one for $|x| \leq X$ and the other for $|x| > X$. The first piece can be handled by Laurent expansions while the second contributes negligibly due to the rapid decay of ϕ . We must carefully keep track of the dependence on m and M .

By (2-9), the Dirichlet series (4-34) can be expressed on $\Re s > 0$ in terms of an Euler product

$$\chi_M(s; m) = \prod_{(p, M)=1} \sum_{t=0}^{\infty} \frac{R(1, p^t)R(m^2, p^t)}{\varphi(p^t)p^{ts}} = \prod_{(p, M)=1} \begin{cases} \left(1 + \frac{1}{(p-1)p^s}\right) & \text{if } (p, m) = 1, \\ \left(1 - \frac{1}{p^s}\right) & \text{if } (p, m) > 1, \end{cases} \quad (4-35)$$

where the second equality follows from the fact that $R(1, b) = \mu(b)$, $R(m^2, 1) = 1$, $R(m^2, p) = -1$ if $(m, p) = 1$, and $R(m^2, p) = \varphi(p) = p - 1$ if $(p, m) > 1$. Moreover, (4-35) can be written as

$$\begin{aligned} \chi_M(s; m) &= \prod_p \left(1 + \frac{1}{(p-1)p^s}\right) \cdot \prod_{p|M} \left(1 + \frac{1}{(p-1)p^s}\right)^{-1} \cdot \prod_{p|\frac{m}{(m, M^\infty)}} \left(1 - \frac{1}{p^s}\right) \left(1 + \frac{1}{(p-1)p^s}\right)^{-1} \\ &=: \chi^{(1)}(s) \cdot \chi_M^{(2)}(s) \cdot \chi_M^{(3)}(s; m). \end{aligned} \quad (4-36)$$

This corrects a mistake made in [25, p. 99] regarding the factorization of $\chi_M(s; m)$. For this reason, the relevant Laurent expansions must be recomputed and the correct results can be obtained as follows.

For $\Re s > 0$, it is easy to verify that

$$\chi^{(1)}(s) = \frac{\zeta(1+s)}{\zeta(2+2s)} \prod_p \left(1 + \frac{1}{(p-1)(p^{1+s}+1)}\right). \quad (4-37)$$

For $s = O(1)$, we have

$$\chi^{(1)}(s) = (s^{-1} + O(1)) \left(\frac{1}{\zeta(2)} \prod_p \left(1 + \frac{1}{p^2-1}\right) + O(|s|) \right) = s^{-1} + O(1). \quad (4-38)$$

Define $\zeta_M(s) := \prod_{p|M} (1 - p^{-s})^{-1}$. In particular, $\zeta_M(1) = M/\varphi(M)$. For any $s = O(1/\log \log M)$, we have

$$\begin{aligned} \zeta_M(1+s) \cdot \frac{\varphi(M)}{M} &= \prod_{p|M} \frac{1-p^{-1}}{1-p^{-1} + O(|s|\frac{\log p}{p})} = \prod_{p|M} \left(1 + O\left(|s|\frac{\log p}{p}\right)\right)^{-1} \\ &= 1 + O\left(|s| \sum_{p|M} \frac{\log p}{p}\right) = 1 + O(|s| \log \log M). \end{aligned} \quad (4-39)$$

From this and (4-37), it follows that

$$\begin{aligned} \chi_M^{(2)}(s) &= \frac{\varphi(M)}{M} (1 + O(|s| \log \log M)) \cdot \left(\zeta_M(2) \prod_{p|M} \left(1 + \frac{1}{p^2-1}\right)^{-1} + O(|s|) \right) \\ &= \frac{\varphi(M)}{M} (1 + O(|s| \log \log M)). \end{aligned} \quad (4-40)$$

Suppose $m/(m, M^\infty) > 1$. For $s = O(1/\log m)$, we have

$$\prod_{p|\frac{m}{(m, M^\infty)}} (1 - p^{-s}) \ll \prod_{p|\frac{m}{(m, M^\infty)}} |s| \log p \ll |s| \prod_{p|\frac{m}{(m, M^\infty)}} \log p. \tag{4-41}$$

Observe that

$$\log \prod_{p|\frac{m}{(m, M^\infty)}} \log p \ll \sum_{p=O(\log m)} \log \log p \ll \frac{(\log m) \log \log \log m}{\log \log m}, \tag{4-42}$$

which implies

$$\prod_{p|\frac{m}{(m, M^\infty)}} (1 - p^{-s}) \ll |s|m^{\epsilon'} \tag{4-43}$$

for any $\epsilon' > 0$. (In this proof, we distinguish ϵ with ϵ' for clarity.)

When $m/(m, M^\infty) = 1$, we have $\prod_{p|\frac{m}{(m, M^\infty)}} (1 - p^{-s}) = 1$. In other words,

$$\chi_M^{(3)}(s; m) = \delta\left(\frac{m}{(m, M^\infty)}, 1\right) + O(|s|m^{\epsilon'}). \tag{4-44}$$

Altogether, for $|s| \ll X \ll 1/((\log m) \log \log M)$, the following estimate holds:

$$\chi_M(s; m) = \frac{\varphi(M)}{sM} \delta\left(\frac{m}{(m, M^\infty)}, 1\right) + O(m^{\epsilon'}). \tag{4-45}$$

From [18, (8.322)] we have

$$\Gamma\left(\frac{k+s}{2}\right) = \Gamma\left(\frac{k-s}{2}\right) \left(\frac{k}{2}\right)^s (1 + O(|s|/k)). \tag{4-46}$$

For $s = \epsilon + 4\pi ix$ with $\epsilon, x \ll X$, we have

$$\chi_M(s; m) \frac{\Gamma\left(\frac{k}{2} - 2\pi ix\right)}{\Gamma\left(\frac{k}{2} + 2\pi ix\right)} = \frac{\varphi(M)}{sM} \delta\left(\frac{m}{(m, M^\infty)}, 1\right) \left(\frac{k}{2}\right)^{-4\pi ix} + O(m^{\epsilon'}). \tag{4-47}$$

Making the change of variables $x \rightarrow -x$ and using the evenness of ϕ , we have

$$\begin{aligned} & \int_{-X}^X \phi(x \log R) \left(\frac{2\pi m}{\sqrt{Q}}\right)^{4\pi ix} \frac{\Gamma\left(\frac{k}{2} - 2\pi ix\right)}{\Gamma\left(\frac{k}{2} + 2\pi ix\right)} \chi_M(\epsilon + 4\pi ix; m) dx \\ &= \frac{\varphi(M)}{M} \delta\left(\frac{m}{(m, M^\infty)}, 1\right) \int_{-X}^X \phi(x \log R) \left(\frac{k\sqrt{Q}}{4\pi m}\right)^{4\pi ix} \frac{dx}{\epsilon - 4\pi ix} + O(m^{\epsilon'} X). \end{aligned} \tag{4-48}$$

We next bound each of the three products over primes in (4-36) for $s = \epsilon + 4\pi ix$ with $|x| > X$.

The last infinite product and $\zeta(2+2s)^{-1}$ in (4-37) are clearly $O(1)$. Using the bound $\zeta(1+s) \ll \log |x|$ for $|x| \geq 3$ (see [3, Theorem 13.4]) and $\zeta(1+s) = O(|s|^{-1})$ as $s \rightarrow 0$, we have

$$|\chi^{(1)}(s)| \ll |\zeta(1+s)| \ll X^{-1} \log(3 + |x|). \tag{4-49}$$

Next, observe the inequality

$$\log \prod_{\substack{p|M \\ p>2}} \left| 1 + \frac{1}{(p-1)p^s} \right|^{-1} \leq - \sum_{\substack{p|M \\ p>2}} \log \left(1 - \frac{1}{p-1} \right) = \sum_{p|M} \frac{1}{p-1} + O(1). \tag{4-50}$$

From the identity $\zeta_M(1) = M/\varphi(M)$ and Taylor’s expansion, one easily obtains

$$\log \frac{M}{\varphi(M)} = \sum_{p|M} \frac{1}{p-1} + O(1). \tag{4-51}$$

As a result, we have

$$(4-50) \leq \log \frac{M}{\varphi(M)} + O(1) \leq \log \log \log M + O(1) \tag{4-52}$$

using the bound $\varphi(M) \gg M/\log \log M$ (again from [3, Theorem 13.14]). We may deduce that

$$|\chi_M^{(2)}(s)| \ll \log \log M, \tag{4-53}$$

where the implicit constant is absolute. The estimate for $\chi_M^{(3)}(s; m)$ follows from a similar argument. Thus

$$|\chi_M(s; m)| \ll X^{-1} m^{\epsilon'} (\log \log M) \log(3 + |x|) \tag{4-54}$$

for any $\epsilon' > 0$.

Equation (4-54) and the decay of ϕ imply

$$\begin{aligned} \left| \int_X^\infty \phi(x \log R) \chi_M(\epsilon + 4\pi i x; m) \left(\frac{2\pi m}{\sqrt{Q}} \right)^{4\pi i x} \frac{\Gamma(\frac{k}{2} - 2\pi i x)}{\Gamma(\frac{k}{2} + 2\pi i x)} dx \right| \\ \ll X^{-1} m^{\epsilon'} (\log \log M) \int_X^\infty (x \log R)^{-A} \log(3 + |x|) dx \\ \ll m^{\epsilon'} (\log \log M) (X \log R)^{-A}, \end{aligned} \tag{4-55}$$

as well as

$$\left| \int_X^\infty \phi(x \log R) \left(\frac{k\sqrt{Q}}{4\pi m} \right)^{4\pi i x} \frac{dx}{\epsilon - 4\pi i x} \right| \ll \int_X^\infty \frac{1}{x} (x \log R)^{-A} dx \ll (X \log R)^{-A} \tag{4-56}$$

for any $A > 0$.

Combining the estimates (4-55), (4-56) and (4-48), it follows that

$$(4-30) = \delta\left(\frac{m}{(m, M^\infty)}, 1\right) \frac{\varphi(M)}{M} \lim_{\epsilon \downarrow 0} \int_{-\infty}^\infty \phi(x \log R) \left(\frac{k\sqrt{Q}}{4\pi m} \right)^{4\pi i x} \frac{dx}{\epsilon - 4\pi i x} + O(m^{\epsilon'} (\log \log M) (X \log R)^{-A}) + O(m^{\epsilon'} X) \tag{4-57}$$

for any $\epsilon' > 0$, $A > 0$, and $X \ll 1/((\log m) \log \log M)$.

Taking $A = 1$ and $X^{-1} := (\log m)(\log \log M)(\log R)^{1/2}$, and following the same argument of [25, p. 100], we may now conclude that (4-30) equals

$$\begin{aligned} \delta\left(\frac{m}{(m, M^\infty)}, 1\right) \frac{\varphi(M)}{M} \left(-\frac{1}{2} \int_{-\infty}^\infty \phi(x) \sin\left(\frac{2\pi x}{\log R} \log \frac{k^2 Q}{16\pi^2 m^2}\right) \frac{dx}{2\pi x} + \frac{1}{4} \phi(0) \right) \\ + O\left(m^{\epsilon'} \frac{(\log \log M)^2}{(\log R)^{1/2}}\right). \end{aligned} \tag{4-58}$$

The proof of Proposition 4.2 is complete.

4.3. Proof of Proposition 3.6. Suppose $n_j + m_j > 2$ for some $1 \leq j \leq \omega$ in (3-20). Firstly, applying Proposition 4.1 to (3-20) with $c = p_1 \cdots p_\alpha q_1^{m_1} \cdots q_\omega^{m_\omega}$, we obtain

$$\begin{aligned}
 E(\vec{n}, \vec{m}) &= \sum_{q_1, \dots, q_\omega} \prod_{j=1}^\omega \hat{\phi}\left(\frac{\log q_j}{\log R}\right)^{n_j} \frac{\log^{n_j} q_j}{q_j^{(n_j+m_j)/2} \log^{n_j} R} \sum_{p_1, \dots, p_{\alpha+\delta}} \prod_{i=1}^{\alpha+\delta} \hat{\phi}\left(\frac{\log p_i}{\log R}\right) \frac{\log p_j}{p_j \log R} \\
 &\quad \times \sum_{m \leq N^\epsilon} \frac{1}{m^2} \sum_{\substack{(b, N p_{\alpha+1} \cdots p_{\alpha+\delta} q_{\theta+1} \cdots q_\omega)=1 \\ p_1, \dots, p_\alpha, q_1, \dots, q_\theta | b}} \frac{R(m^2, b) R(p_1 \cdots p_\alpha q_1^{m_1} \cdots q_\theta^{m_\theta}, b)}{\varphi(b)} \\
 &\quad \times \int_0^\infty J_{k-1}(x) \hat{\Phi}_{n-\alpha-\delta} \left(\frac{2}{\log R} \log \left(\frac{bx}{4\pi m} \sqrt{\frac{N}{p_1 \cdots p_{\alpha+\delta} q_1^{m_1} \cdots q_\omega^{m_\omega}}} \right) \right) \frac{dx}{\log R}, \quad (4-59)
 \end{aligned}$$

up to an error $O(N^{-\epsilon})$. The condition $b < N^{2022}$ has been removed at a negligible cost.

Secondly, we convert the sums over q_j and p_i in (4-59) into sums over *distinct* primes, requiring us to break up the sums depending on whether some of the primes in the sums are equal or not, as well as the order of the primes factors of b . It follows that $E(\vec{n}, \vec{m})$ is a sum of terms of the form

$$\begin{aligned}
 F(\vec{a}, \vec{b}, \vec{c}, \vec{d}, \vec{e}) &:= \sum_{\substack{q_1, \dots, q_\ell \\ q_j \text{ distinct}}} \prod_{j=1}^\ell \hat{\phi}\left(\frac{\log q_j}{\log R}\right)^{a_j} \frac{\log^{a_j} q_j}{q_j^{b_j} \log^{a_j} R} \sum_{m \leq N^\epsilon} \frac{1}{m^2} \sum_{\substack{(b', N q_1 \dots q_\ell)=1 \\ b=b' q_1^{c_1} \dots q_\ell^{c_\ell}}} \frac{R(m^2, b) R(q_1^{d_1} \cdots q_\ell^{d_\ell}, b)}{\varphi(b)} \\
 &\quad \times \int_{x=0}^\infty J_{k-1}(x) \hat{\Phi}_{n-\nu} \left(\frac{2}{\log R} \log \left(\frac{b'x}{4\pi m} \sqrt{N q_1^{e_1} \cdots q_\ell^{e_\ell}} \right) \right) \frac{dx}{\log R}, \quad (4-60)
 \end{aligned}$$

up to an error term of size $O(N^{-\epsilon})$, where a_i, b_i, c_i , and d_i are positive integers, and the e_i 's are integers. Additionally, $\sum a_j = \nu$ and $b_i > 1$ for some i (since $n_j + m_j > 2$ for some j), as well as $b_j \geq d_j$ for all j (since $n_j \geq m_j$ in (4-59)).

As a result, the proof of Proposition 3.6 rests on the following lemma, which will also be useful in Section 4.4.

Lemma 4.12. *Let $F(\vec{a}, \vec{b}, \vec{c}, \vec{d}, \vec{e})$ be defined as in (4-60) with a_j, b_j, c_j, d_j 's being positive integers, e_j 's being integers, $b_j \geq d_j$ for all $1 \leq j \leq \kappa$. If $b_i > 1$ or $d_i < c_i$ for some i , then $F(\vec{a}, \vec{b}, \vec{c}, \vec{d}, \vec{e}) \ll 1/\log N$.*

Proof. Using the multiplicativity of the Ramanujan sums and the totient function φ , observe that

$$\frac{R(m^2, b) R(q_1^{d_1} \cdots q_\kappa^{d_\kappa}, b)}{\varphi(b)} = \frac{R(q_1^{d_1} \cdots q_\kappa^{d_\kappa}, q_1^{c_1} \cdots q_\kappa^{c_\kappa})}{\varphi(q_1^{c_1} \cdots q_\kappa^{c_\kappa})} \cdot \frac{R(m^2, b') R(m^2, q_1^{c_1} \cdots q_\kappa^{c_\kappa}) R(1, b')}{\varphi(b')}, \quad (4-61)$$

where $b = b' q_1^{c_1} \cdots q_\kappa^{c_\kappa}$ and $(b', N q_1 \cdots q_\ell) = 1$. Apply (4-61) to (4-60). This allows us to rewrite $F(\vec{a}, \vec{b}, \vec{c}, \vec{d}, \vec{e})$ (with $Q = N q_1^{e_1} \cdots q_\ell^{e_\ell}$) as

$$\begin{aligned}
 &\sum_{\substack{q_1, \dots, q_\ell \\ q_j \text{ distinct}}} \prod_{j=1}^\ell \hat{\phi}\left(\frac{\log q_j}{\log R}\right)^{a_j} \frac{\log^{a_j} q_j}{q_j^{b_j} \log^{a_j} R} \sum_{m \leq N^\epsilon} \frac{R(q_1^{d_1} \cdots q_\kappa^{d_\kappa}, q_1^{c_1} \cdots q_\kappa^{c_\kappa})}{\varphi(q_1^{c_1} \cdots q_\kappa^{c_\kappa})} \frac{R(m^2, q_1^{c_1} \cdots q_\kappa^{c_\kappa})}{m^2} \\
 &\quad \times \sum_{(b', N q_1 \dots q_\ell)=1} \frac{R(m^2, b') R(1, b')}{\varphi(b')} \int_{x=0}^\infty J_{k-1}(x) \hat{\Phi}_{n-\nu} \left(\frac{2}{\log R} \log \frac{b'x \sqrt{Q}}{4\pi m} \right) \frac{dx}{\log R}. \quad (4-62)
 \end{aligned}$$

By Proposition 4.2, one observes that the sum over b' in (4-62) is $\ll m^\epsilon$, and hence, (4-62) is

$$\ll \sum_{\substack{q_1, \dots, q_\ell \\ q_j \text{ distinct}}} \prod_{j=1}^\ell \left| \hat{\phi} \left(\frac{\log q_j}{\log R} \right)^{a_j} \frac{\log^{a_j} q_j}{q_j^{b_j} \log^{a_j} R} \right| \sum_{m \leq N^\epsilon} \frac{|R(q_1^{d_1} \cdots q_k^{d_k}, q_1^{c_1} \cdots q_k^{c_k})| |R(m^2, q_1^{c_1} \cdots q_k^{c_k})|}{\varphi(q_1^{c_1} \cdots q_k^{c_k}) m^{2-\epsilon}}.$$

The last m -sum is bounded by

$$\left(\sum_{(m', q_1 \cdots p_k)=1} \frac{1}{(m')^{2-\epsilon}} \right) \left(\prod_{i=1}^k \sum_{t \geq 0} \frac{|R(q_i^{2t}, q_i^{c_i})| |R(q_i^{d_i}, q_i^{c_i})|}{q_i^{(2-\epsilon)t} \varphi(q_i^{c_i})} \right) \ll \prod_{i=1}^k \sum_{t \geq 0} \frac{|R(q_i^{2t}, q_i^{c_i})| |R(q_i^{d_i}, q_i^{c_i})|}{q_i^{(2-\epsilon)t} \varphi(q_i^{c_i})},$$

once again due to the multiplicativity of the Ramanujan sums.

We now analyze the sum over t , primarily relying on (2-8) to bound the Ramanujan sums. When $2t < c_i - 1$, then $R(q_i^{2t}, q_i^{c_i}) = 0$. When $2t = c_i - 1$, then $R(q_i^{2t}, q_i^{c_i}) = q_i^{2t}$. When $2t \geq c_i$, we have $R(q_i^{2t}, q_i^{c_i}) = \varphi(q_i^{c_i}) \leq q_i^{c_i}$.

The sum over t is $O(q_i^{\epsilon \lfloor c_i/2 \rfloor})$ when $d_i \geq c_i$, and is $O(q_i^{-1+\epsilon \lfloor c_i/2 \rfloor})$ when $d_i < c_i$, where the bounds $|R(q_i^{d_i}, q_i^{c_i})| \leq \varphi(q_i^{c_i})$ and $|R(q_i^{d_i}, q_i^{c_i})| \leq q_i^{d_i}$ were applied respectively. Therefore,

$$\begin{aligned} F(\vec{a}, \vec{b}, \vec{c}, \vec{d}, \vec{e}) &\ll \sum_{\substack{q_1, \dots, q_\ell \\ q_j \text{ distinct}}} \prod_{j=1}^\ell \left| \hat{\phi} \left(\frac{\log q_j}{\log R} \right)^{a_j} \right| \frac{\log^{a_j} q_j}{q_j^{b_j + \eta_j - \epsilon \lfloor c_j/2 \rfloor} \log^{a_j} R} \\ &\ll \prod_{j=1}^\ell \left(\sum_p \left| \hat{\phi} \left(\frac{\log p}{\log R} \right)^{a_j} \right| \frac{\log^{a_j} p}{p^{b_j + \eta_j - \epsilon \lfloor c_j/2 \rfloor} \log^{a_j} R} \right), \end{aligned} \tag{4-63}$$

where $\eta_j = \mathbb{1}_{d_j < c_j}$. Observe the following:

- (1) Set $x_j := b_j + \eta_j - \epsilon \lfloor c_j/2 \rfloor$. The sum over p in (4-63) is $O(1/\log^{a_j} R)$ when $x_j > 1$, and is $O(1)$ when $x_j = 1$.
- (2) Suppose $d_j \geq c_j$. By assumption, we have $b_j \geq d_j$ and so $b_j \geq c_j$. If $b_j = 1$, then $x_j = 1$, and if $b_j > 1$, then $x_j > 1$. Suppose $d_j < c_j$. Then $x_j > 1$.

In particular, $x_j \geq 1$ always holds and each of the p -sums in (4-63) is $O(1)$.

By our assumption, there exists i for which either $d_i < c_i$ or $b_j > 1$ hold. In either case, we have $x_i > 1$ and the i -th factor in (4-63) is thus $\ll 1/\log^{a_i} R \ll 1/\log N$ (since $a_i > 0$ by assumption). Taking the product over all j 's, we may now conclude that $F(\vec{a}, \vec{b}, \vec{c}, \vec{d}, \vec{e}) \ll 1/\log N$. This completes the proof of the lemma. □

Proof of Proposition 3.6. By Lemma 4.12 and the arguments preceding (4-60), $E(\vec{n}, \vec{m})$ can be written as the sum of finitely many terms of sizes $O(1/\log N)$ whenever $n_j + m_j > 2$ for some j . Note that the number of such terms is independent of N . This completes the proof. □

4.4. Analytic simplification of the main contribution. Recall the expression (3-20) for $E(\vec{n}, \vec{m})$ and the fact that the main contribution comes from terms with $\omega = 0$ and $n_j = m_j = 1$ for all $1 \leq j \leq n$, i.e.,

$$\begin{aligned}
 A := & - \sum_{\alpha=0}^{a-1} \binom{n}{\alpha} 2^{n+1} \pi \sum_{p_1, \dots, p_\alpha} \prod_{j=1}^{\alpha} \hat{\phi} \left(\frac{\log p_j}{\log R} \right) \frac{\log p_j}{\sqrt{p_j} \log R} \sum_{m \leq N^\epsilon} \frac{1}{m} \sum_{\substack{(b, N)=1 \\ b < N^{2022} \\ p_1, \dots, p_\alpha | b}} \frac{1}{b \varphi(b)} R(m^2, b) R(p_1 \cdots p_\alpha, b) \\
 & \times N^{-1/2} \sum_{p_{\alpha+1}, \dots, p_n} J_{k-1} \left(\frac{4\pi m \sqrt{p_1 \cdots p_n}}{b \sqrt{N}} \right) \prod_{j=\alpha+1}^n \hat{\phi} \left(\frac{\log p_j}{\log R} \right) \frac{\chi_0(p_j) \log p_j}{\sqrt{p_j} \log R} + O(N^{-\epsilon}). \quad (4-64)
 \end{aligned}$$

Remark 4.13. (1) The coefficient $\binom{n}{\alpha}$ comes from the choices of indices for the prime factors of b .
 (2) We have truncated the α -sum in (4-64) to $0 \leq \alpha \leq a-1$ because the contribution from $\alpha > a-1$ the term is $O(N^{-\epsilon})$, which follows from the bounds $J_{k-1}(x) \ll x$, $R(m^2, b) \leq m^4$ and $R(p_1 \cdots p_\alpha, b) \leq \varphi(b)$.

Applying Proposition 4.1 to (4-64), we have, upon simplification,

$$A = -2^n \sum_{\alpha=0}^{a-1} \binom{n}{\alpha} \sum_{\delta=0}^{a-\alpha-1} \binom{n-\alpha}{\delta} \sum_{i=0}^{a-\alpha-\delta-1} (-1)^i \binom{n-\alpha-\delta}{i} H(\alpha, \delta) + O(N^{-\epsilon}), \quad (4-65)$$

where

$$\begin{aligned}
 H(\alpha, \delta) := & \sum_{p_1, \dots, p_{\alpha+\delta}} \prod_{j=1}^{\alpha+\delta} \hat{\phi} \left(\frac{\log p_j}{\log R} \right) \frac{\log p_j}{p_j \log R} \sum_{m \leq N^\epsilon} \frac{1}{m^2} \sum_{\substack{(b, N p_{\alpha+1} \cdots p_{\alpha+\delta})=1 \\ p_1, \dots, p_\alpha | b}} \frac{R(m^2, b) R(p_1 \cdots p_\alpha, b)}{\varphi(b)} \\
 & \times \int_{x=0}^{\infty} J_{k-1}(x) \hat{\Phi}_{n-\alpha-\delta} \left(\frac{2}{\log R} \log \frac{bx \sqrt{N}}{4\pi m \sqrt{p_1 \cdots p_{\alpha+\delta}}} \right) \frac{dx}{\log R}. \quad (4-66)
 \end{aligned}$$

The rest of this subsection is dedicated to proving the following lemma.

Lemma 4.14.

$$\begin{aligned}
 H(\alpha, \delta) = & -2^{-1-\alpha-\delta} (-1)^\alpha \int_{-\infty}^{\infty} \cdots \int_{-\infty}^{\infty} \hat{\phi}(x_2) \cdots \hat{\phi}(x_{\alpha+\delta+1}) \\
 & \times \left(\int_{-\infty}^{\infty} \phi^{n-\alpha-\delta}(x_1) \frac{\sin(2\pi x_1 (1 + |x_2| + \cdots + |x_{\alpha+1}| - |x_{\alpha+2}| - \cdots - |x_{\alpha+\delta+1}|))}{2\pi x_1} dx_1 - \frac{1}{2} \phi^{n-\alpha-\delta}(0) \right) \\
 & \times dx_2 \cdots dx_{\alpha+\delta+1} + O\left(\frac{(\log \log N)^2}{(\log N)^{1/2}} \right). \quad (4-67)
 \end{aligned}$$

First, we transform the sum over the primes $p_1, \dots, p_{\alpha+\delta}$ in $H(\alpha, \delta)$ to a sum over distinct primes.

Property 4.15. A distinctness condition can be added to (4-66) at the cost of an error of size $O(1/\log N)$.

Proof. The distinctness condition can be imposed to the sums over primes of (4-66) by inclusion-exclusion, depending on which primes are equal. If $p_i = p_j$ for some $1 \leq i \leq \alpha$ and $\alpha + 1 \leq j \leq \alpha + \delta$, then the corresponding term of $H(\alpha, \delta)$ is zero due to the condition on the b -sum.

Without loss of generality, we let $p_1 \cdots p_\alpha = q_1^{u_1} \cdots q_{\alpha'}^{u_{\alpha'}}$ and $p_{\alpha+1} \cdots p_{\alpha+\delta'} = q_{\alpha'+1}^{u_{\alpha'+1}} \cdots q_{\alpha'+\delta'}^{u_{\alpha'+\delta'}}$, where the primes q_i are distinct and at least one $u_i > 1$. To add in a distinctness condition to $H(\alpha, \delta)$, our inclusion-exclusion introduced terms of the form

$$\sum_{\substack{q_1, \dots, q_{\alpha'+\delta'} \\ q_i \text{ distinct}}} \prod_{j=1}^{\alpha'+\delta'} \hat{\phi} \left(\frac{\log q_j}{\log R} \right)^{u_j} \frac{\log^{u_j} q_j}{q_j^{u_j} \log^{u_j} R} \sum_{m \leq N^\epsilon} \frac{1}{m^2} \sum_{\substack{(b, Nq_{\alpha'+1} \cdots q_{\alpha'+\delta'})=1 \\ q_1^{u_1}, \dots, q_{\alpha'}^{u_{\alpha'}} | b}} \frac{R(m^2, b)R(q_1^{u_1} \cdots q_{\alpha'}^{u_{\alpha'}}, b)}{\varphi(b)} \\ \times \int_{x=0}^{\infty} J_{k-1}(x) \hat{\Phi}_{n-\alpha-\delta} \left(\frac{2}{\log R} \log \frac{bx\sqrt{N}}{4\pi m \sqrt{q_1^{u_1} \cdots q_{\alpha'+\delta'}^{u_{\alpha'+\delta'}}}} \right) \frac{dx}{\log R}. \quad (4-68)$$

After breaking up (4-68) based on the multiplicities of the prime factors of b , we appeal to Lemma 4.12 and find (4-68) is $O(1/\log N)$ since at least one $u_i > 1$. This completes the proof. \square

Upon inserting the distinctness condition to $H(\alpha, \delta)$, we break up the terms based on the multiplicities of the prime factors of b . Consider the b -sum of (4-66) over b 's of the form $b'p_1^{c_1} \cdots p_\alpha^{c_\alpha}$, where $(b', p_1 \cdots p_\alpha) = 1$. Due to Lemma 4.12, if any of the c_i 's is > 1 , then the corresponding term is $O(1/\log N)$. Thus, it suffices to consider the case when each of the c_i 's is 1, i.e.,

$$H(\alpha, \delta) = \sum_{\substack{p_1, \dots, p_{\alpha+\delta} \\ p_i \text{ distinct}}} \prod_{j=1}^{\alpha+\delta} \hat{\phi} \left(\frac{\log p_j}{\log R} \right) \frac{\log p_j}{p_j \log R} \sum_{m \leq N^\epsilon} \frac{1}{m^2} \sum_{\substack{b=b'p_1 \cdots p_\alpha \\ (b', Np_1 \cdots p_{\alpha+\delta})=1}} \frac{R(m^2, b)R(p_1 \cdots p_\alpha, b)}{\varphi(b)} \\ \times \int_{x=0}^{\infty} J_{k-1}(x) \hat{\Phi}_{n-\alpha-\delta} \left(\frac{2}{\log R} \log \frac{bx\sqrt{N}}{4\pi m \sqrt{p_1 \cdots p_{\alpha+\delta}}} \right) \frac{dx}{\log R} + O(1/\log N). \quad (4-69)$$

The b -sum can be simplified with the multiplicativity of φ and the Ramanujan sums, $R(q, q) = \varphi(q)$, and the conditions $b = b'p_1 \cdots p_\alpha$ with $(b', Np_1 \cdots p_{\alpha+\delta}) = 1$. In fact,

$$\frac{R(m^2, b)R(p_1 \cdots p_\alpha, b)}{\varphi(b)} = \frac{R(m^2, b')R(m^2, p_1 \cdots p_\alpha)R(1, b')}{\varphi(b')}. \quad (4-70)$$

Applying this to (4-69), we have, with $Q = \frac{Np_1 \cdots p_\alpha}{p_{\alpha+1} \cdots p_{\alpha+\delta}}$,

$$H(\alpha, \delta) = \sum_{\substack{p_1, \dots, p_{\alpha+\delta} \\ p_i \text{ distinct}}} \prod_{j=1}^{\alpha+\delta} \hat{\phi} \left(\frac{\log p_j}{\log R} \right) \frac{\log p_j}{p_j \log R} \sum_{m \leq N^\epsilon} \frac{1}{m^2} R(m^2, p_1 \cdots p_\alpha) \\ \times \sum_{(b', Np_1 \cdots p_{\alpha+\delta})=1} \frac{R(m^2, b')R(1, b')}{\varphi(b')} \int_{x=0}^{\infty} J_{k-1}(x) \hat{\Phi}_{n-\alpha-\delta} \left(\frac{2}{\log R} \log \frac{b'x\sqrt{Q}}{4\pi m} \right) \frac{dx}{\log R} \\ + O(1/\log N). \quad (4-71)$$

We are ready to apply Proposition 4.2 to the sum over b' in (4-71). We first show that the contribution of the error term of (4-3) to $H(\alpha, \delta)$ is $O((\log \log N)^2/(\log N)^{1/2})$. Indeed, using the multiplicativity of

the Ramanujan sums, the m -sum in this case is

$$\begin{aligned} &\ll \sum_{m \leq N^\epsilon} \frac{R(m^2, p_1 \cdots p_\alpha)}{m^2} \cdot m^\epsilon \frac{(\log \log N)^2}{(\log N)^{1/2}} \ll \frac{(\log \log N)^2}{(\log N)^{1/2}} \sum_{(m', p_1 \cdots p_\alpha)=1} \frac{1}{(m')^{2-\epsilon}} \prod_{i=1}^{\alpha} \left(\sum_{t \geq 0} \frac{R(p_i^{2t}, p_i)}{p_i^{(2-\epsilon)t}} \right) \\ &\ll \frac{(\log \log N)^2}{(\log N)^{1/2}} \prod_{i=1}^{\alpha} \left(\sum_{t \geq 0} \frac{R(p_i^{2t}, p_i)}{p_i^{(2-\epsilon)t}} \right). \end{aligned} \quad (4-72)$$

When $t = 0$, $R(p_i^{2t}, p_i) = R(1, p_i) = 1$. When $t > 0$, $R(p_i^{2t}, p_i) = \varphi(p_i) < p_i$. From this, it is clear that the sum over t is $O(1)$ (independent of p_i) and the sum over m is $O((\log \log N)^2 / (\log N)^{1/2})$. Our claim follows by also considering the sums over primes and thus, setting $M = N p_1 \cdots p_{\alpha+\delta}$,

$$\begin{aligned} H(\alpha, \delta) &= \sum_{\substack{p_1, \dots, p_{\alpha+\delta} \\ p_i \text{ distinct}}} \prod_{j=1}^{\alpha+\delta} \hat{\phi} \left(\frac{\log p_j}{\log R} \right) \frac{\log p_j}{p_j \log R} \sum_{m \leq N^\epsilon} \frac{1}{m^2} R(m^2, p_1 \cdots p_\alpha) \delta \left(\frac{m}{(m, M^\infty)}, 1 \right) \frac{\varphi(M)}{M} \\ &\times \left(-\frac{1}{2} \int_{-\infty}^{\infty} \phi(x)^{n-\alpha-\delta} \sin \left(\frac{2\pi x}{\log R} \log \frac{k^2 Q}{16\pi^2 m^2} \right) \frac{dx}{2\pi x} + \frac{1}{4} \phi(0)^{n-\alpha-\delta} \right) + O \left(\frac{(\log \log N)^2}{(\log N)^{1/2}} \right), \end{aligned} \quad (4-73)$$

The factor $\delta \left(\frac{m}{(m, M^\infty)}, 1 \right)$ and the fact that $N \nmid m$ force m in (4-73) to take the shape $p_1^{t_1} \cdots p_{\alpha+\delta}^{t_{\alpha+\delta}}$. Since

$$Q = \frac{N p_1 \cdots p_\alpha}{p_{\alpha+1} \cdots p_{\alpha+\delta}} \quad \text{and} \quad \frac{\varphi(M)}{M} = \left(1 - \frac{1}{N} \right) \prod_{j=1}^{\alpha+\delta} \left(1 - \frac{1}{p_j} \right),$$

we can then simplify (4-73) as

$$\begin{aligned} H(\alpha, \delta) &= \sum_{\substack{p_1, \dots, p_{\alpha+\delta} \\ p_i \text{ distinct}}} \prod_{j=1}^{\alpha+\delta} \hat{\phi} \left(\frac{\log p_j}{\log R} \right) \frac{\log p_j}{p_j \log R} \left(1 - \frac{1}{p_j} \right) \sum_{0 \leq t_1, \dots, t_{\alpha+\delta} \leq \epsilon \log N} \frac{R(p_1^{2t_1} \cdots p_\alpha^{2t_\alpha}, p_1 \cdots p_\alpha)}{p_1^{2t_1} \cdots p_{\alpha+\delta}^{2t_{\alpha+\delta}}} \\ &\times \left(-\frac{1}{2} \int_{-\infty}^{\infty} \phi(x)^{n-\alpha-\delta} \sin \left(2\pi x \left(1 + \frac{\log p_1 + \cdots + \log p_\alpha - \log p_{\alpha+1} - \cdots - \log p_{\alpha+\delta}}{\log R} \right) \right) \frac{dx}{2\pi x} \right. \\ &\quad \left. + \frac{1}{4} \phi(0)^{n-\alpha-\delta} \right) \\ &\quad + O \left(\frac{(\log \log N)^2}{(\log N)^{1/2}} \right). \end{aligned} \quad (4-74)$$

Next, we show that the contribution over the complement of $t_1 = \cdots = t_{\alpha+\delta} = 0$ in (4-74) is negligible. Indeed, upon inserting an absolute value, the quantity of interest is bounded by

$$\max_{1 \leq i \leq \alpha+\delta} \sum_{\substack{p_1, \dots, p_{\alpha+\delta} \\ p_j < R}} \prod_{j=1}^{\alpha+\delta} \frac{\log p_j}{p_j \log R} \sum_{\substack{0 \leq t_1, \dots, t_{\alpha+\delta} \leq \epsilon \log N \\ t_i \neq 0}} \frac{|R(p_1^{2t_1}, p_1)| \cdots |R(p_\alpha^{2t_\alpha}, p_\alpha)|}{p_1^{2t_1} \cdots p_{\alpha+\delta}^{2t_{\alpha+\delta}}}, \quad (4-75)$$

which is equal to

$$\max_{1 \leq i \leq \alpha + \delta} \prod_{j=1}^{\alpha + \delta} \left(\sum_{p_j < R} \frac{\log p_j}{p_j \log R} \sum_{\delta(i,j) \leq t_j \leq \epsilon \log N} \frac{|R(p_j^{2t_j}, p_j^{s_j})|}{p_j^{2t_j}} \right), \tag{4-76}$$

where $s_j = 1$ for $1 \leq j \leq \alpha$ and 0 for $\alpha + 1 \leq j \leq \alpha + \delta$.

We have $|R(1, p^{s_j})| = 1$ when $t_j = 0$, whereas the sum over $t_j \geq 1$ is $O(1/p_j)$. When $j \neq i$, the sums over t_j 's and p_j 's are both $O(1)$. When $j = i$, the sum over t_i 's is $O(1/p_i)$ as the term with $t_i = 0$ is absent. As a result, we have

$$\sum_{p_i < R} \frac{\log p_i}{p_i \log R} \sum_{1 \leq t_i \leq \epsilon \log N} \frac{|R(p_i^{2t_i}, p_i^{s_i})|}{p_i^{2t_i}} \ll \sum_{p_i < R} \frac{\log p_i}{p_i^2 \log R} \ll 1/\log N. \tag{4-77}$$

Upon taking the product over j , we deduce that (4-76) is $O(1/\log N)$. Our claim follows.

It remains to consider the contribution when $t_1 = \dots = t_{\alpha + \delta} = 0$ (and so $m = 1$). In this case, we approximate $\prod_j (1 - 1/p_j)$ in (4-74) by 1 as the contribution from the rest of the terms in the expansion of such a product is negligibly small. More precisely, we have

$$\begin{aligned} H(\alpha, \delta) &= (-1)^\alpha \sum_{\substack{p_1, \dots, p_{\alpha + \delta} \\ p_i \text{ distinct}}} \prod_{j=1}^{\alpha + \delta} \hat{\phi} \left(\frac{\log p_j}{\log R} \right) \frac{\log p_j}{p_j \log R} \\ &\times \left(-\frac{1}{2} \int_{-\infty}^{\infty} \phi(x)^{n - \alpha - \delta} \sin \left(2\pi x \left(1 + \frac{\log p_1 + \dots + \log p_\alpha - \log p_{\alpha + 1} - \dots - \log p_{\alpha + \delta}}{\log R} \right) \right) \frac{dx}{2\pi x} \right. \\ &\quad \left. + \frac{1}{4} \phi(0)^{n - \alpha - \delta} \right) \\ &\quad + O \left(\frac{(\log \log N)^2}{(\log N)^{1/2}} \right), \tag{4-78} \end{aligned}$$

where we use the identity $R(1, p_1 \dots p_\alpha) = \mu(p_1 \dots p_\alpha) = (-1)^\alpha$.

Property 4.16. Equation (4-78) holds with the distinctness condition in the prime sum removed.

Proof. In the process of removing the distinctness condition, we apply inclusion-exclusion and we are left to show that the terms with $p_i = p_j$ for some $i \neq j$ can be eliminated as they contribute negligibly. Indeed, suppose the condition in the sum of (4-78) is replaced by $p_1 \dots p_{\alpha + \delta} = q_1^{a_1} \dots q_\ell^{a_\ell}$, where $q_i \neq q_j$ when $i \neq j$ and where $a_j > 1$ for some j . Now, such a contribution is bounded by

$$\sum_{\substack{q_1, \dots, q_\ell \\ q_i \text{ distinct} \\ q_i < R}} \prod_{j=1}^{\ell} \frac{\log^{a_j} q_j}{q_j^{a_j} \log^{a_j} R} \ll \sum_{\substack{q_1, \dots, q_\ell \\ q_i < R}} \prod_{j=1}^{\ell} \frac{\log^{a_j} q_j}{q_j^{a_j} \log^{a_j} R} \ll \prod_{j=1}^{\ell} \left(\sum_{q_j < R} \frac{\log^{a_j} q_j}{q_j^{a_j} \log^{a_j} R} \right). \tag{4-79}$$

The last sum is $O(1)$ and $O(1/\log^{a_j} R)$ when $a_j = 1$ and $a_j > 1$ respectively. Since $a_j > 1$ for some j , we find (4-79) is $O(1/\log^2 R)$. This completes the proof. \square

To complete the proof of Lemma 4.14, we apply the prime number theorem with partial summation to each of the sums over primes in (4-78) without the distinctness condition. We find that

$$\begin{aligned}
 H(\alpha, \delta) = & -2^{-1-\alpha-\delta}(-1)^\alpha \int_{-\infty}^{\infty} \cdots \int_{-\infty}^{\infty} \hat{\phi}(x_2) \cdots \hat{\phi}(x_{\alpha+\delta+1}) \\
 & \times \left(\int_{-\infty}^{\infty} \phi^{n-\alpha-\delta}(x_1) \frac{\sin(2\pi x_1(1+|x_2|+\cdots+|x_{\alpha+1}|-|x_{\alpha+2}|-\cdots-|x_{\alpha+\delta+1}|))}{2\pi x_1} dx_1 \right. \\
 & \left. - \frac{1}{2}\phi^{n-\alpha-\delta}(0) \right) dx_2 \cdots dx_{\alpha+\delta+1} \\
 & + O\left(\frac{(\log \log N)^2}{(\log N)^{1/2}}\right),
 \end{aligned}$$

as desired.

4.5. Proof of Proposition 3.7: combinatorial simplification of the main contribution. In this section we finish the proof of Proposition 3.7 by applying Lemma 4.14 to (4-65) and simplifying. This step is mostly combinatorial, although we need the following lemma.

Lemma 4.17.
$$\int_{-\infty}^{\infty} \hat{\phi}(y) (\sin(z + 2\pi x|y|) + \sin(z - 2\pi x|y|)) dy = 2 \sin(z)\phi(x).$$

Proof. Using that $\sin(z + 2\pi x|y|) + \sin(z - 2\pi x|y|) = 2 \sin(z) \cos(2\pi xy)$ we have

$$\begin{aligned}
 \int_{-\infty}^{\infty} \hat{\phi}(y) (\sin(z + 2\pi x|y|) + \sin(z - 2\pi x|y|)) dy &= 2 \sin(z) \int_{-\infty}^{\infty} \hat{\phi}(y) \cos(2\pi xy) dy \\
 &= 2 \sin(z) \int_{-\infty}^{\infty} \hat{\phi}(y) \Re(\exp(2\pi ixy)) dy \\
 &= 2 \sin(z)\phi(x)
 \end{aligned}$$

as desired. □

Applying Lemma 4.14 to (4-65) gives

$$\begin{aligned}
 A = & \sum_{\alpha=0}^{a-1} \sum_{\delta=0}^{a-\alpha-1} \binom{n}{\alpha+\delta} \binom{\alpha+\delta}{\alpha} \sum_{i=0}^{a-\alpha-\delta-1} (-1)^i \binom{n-\alpha-\delta}{i} 2^{n-1-\alpha-\delta} (-1)^\alpha \\
 & \times \int_{-\infty}^{\infty} \cdots \int_{-\infty}^{\infty} \hat{\phi}(x_2) \cdots \hat{\phi}(x_{\alpha+\delta+1}) \\
 & \times \left(\int_{-\infty}^{\infty} \phi^{n-\alpha-\delta}(x_1) \frac{\sin(2\pi x_1(1+|x_2|+\cdots+|x_{\alpha+1}|-|x_{\alpha+2}|-\cdots-|x_{\alpha+\delta+1}|))}{2\pi x_1} dx_1 \right. \\
 & \left. - \frac{1}{2}\phi^{n-\alpha-\delta}(0) \right) dx_2 \cdots dx_{\alpha+\delta+1} \\
 & + O\left(\frac{(\log \log N)^2}{(\log N)^{1/2}}\right). \tag{4-80}
 \end{aligned}$$

Our first step is to eliminate the integral over $\phi^{n-\alpha-\delta}(0)$ in (4-80) when $\alpha + \delta > 0$. We fix some $\nu \leq a$

and collect the terms of (4-80) for which $\alpha + \delta = \nu$:

$$\left(\sum_{\alpha=0}^{\nu} \binom{\nu}{\alpha} (-1)^{\alpha}\right) \binom{n}{\nu} \sum_{i=0}^{a-\nu-1} (-1)^i \binom{n-\nu}{i} 2^{n-1-\nu} \times \int_{-\infty}^{\infty} \cdots \int_{-\infty}^{\infty} \hat{\phi}(x_2) \cdots \hat{\phi}(x_{\nu+1}) \left(-\frac{1}{2}\phi^{n-\nu}(0)\right) dx_2 \cdots dx_{\nu+1}. \quad (4-81)$$

By the binomial theorem, we have $\sum_{\alpha=0}^{\nu} \binom{\nu}{\alpha} (-1)^{\alpha} = (1-1)^{\nu} = 0$ for $\nu > 0$, so the sum over α in (4-81) is 0 unless $\nu = 0$. Thus, the terms where $\alpha + \delta = \nu$ cancel when $\nu > 0$. When $\nu = 0$, we pull out the $-\frac{1}{2}\phi^n(0)$ term and find that

$$A = \sum_{\alpha=0}^{a-1} \sum_{\delta=0}^{a-\alpha-1} \binom{n}{\alpha+\delta} \binom{\alpha+\delta}{\alpha} \sum_{i=0}^{a-\alpha-\delta-1} (-1)^i \binom{n-\alpha-\delta}{i} 2^{n-1-\alpha-\delta} (-1)^{\alpha} I(\alpha, \delta) - 2^{n-2}\phi^n(0) \sum_{i=0}^{a-1} (-1)^i \binom{n}{i} + O\left(\frac{(\log \log N)^2}{(\log N)^{1/2}}\right), \quad (4-82)$$

where

$$I(\alpha, \delta) = \int_{-\infty}^{\infty} \cdots \int_{-\infty}^{\infty} \hat{\phi}(x_2) \cdots \hat{\phi}(x_{\alpha+\delta+1}) \int_{-\infty}^{\infty} \phi^{n-\alpha-\delta}(x_1) \times \frac{\sin(2\pi x_1(1 + |x_2| + \cdots + |x_{\alpha+1}| - |x_{\alpha+2}| - \cdots - |x_{\alpha+\delta+1}|))}{2\pi x_1} dx_1 \cdots dx_{\alpha+\delta+1}. \quad (4-83)$$

We simplify the first sum over α in (4-82), which we denote by A' . By Lemma 4.17 we have $I(\alpha, \delta) = 2I(\alpha, \delta - 1) - I(\alpha + 1, \delta - 1)$. We express A' in terms of $I(\alpha, 0)$ via the following result, where $I(\alpha, \delta)$ remains defined by (4-83):

Lemma 4.18.
$$I(\alpha, \delta) = \sum_{j=0}^{\delta} 2^{\delta-j} (-1)^j \binom{\delta}{j} I(\alpha + j, 0).$$

Proof. We show by induction that

$$I(\alpha, \delta) = \sum_{j=0}^k 2^{k-j} (-1)^j \binom{k}{j} I(\alpha + j, \delta - k), \quad (4-84)$$

after which setting $k = \delta$ proves the lemma. The base case $k = 0$ holds immediately. Suppose (4-84) holds up to k . Using that $I(\alpha, \delta) = 2I(\alpha, \delta - 1) - I(\alpha + 1, \delta - 1)$ we have

$$\begin{aligned} I(\alpha, \delta) &= \sum_{j=0}^k 2^{k-j} (-1)^j \binom{k}{j} (2I(\alpha + j, \delta - k - 1) - I(\alpha + j + 1, \delta - k - 1)) \\ &= \sum_{j=0}^{k+1} 2^{k+1-j} (-1)^j \left(\binom{k+1}{j+1} - \binom{k}{j}\right) I(\alpha + j, \delta - k - 1) \\ &= \sum_{j=0}^{k+1} 2^{k+1-j} (-1)^j \binom{k+1}{j} I(\alpha + j, \delta - k - 1) \end{aligned} \quad (4-85)$$

completing the inductive hypothesis and the proof of the lemma. □

Proof of Proposition 3.7. Apply Lemma 4.18 to (4-82), we have

$$A' = \sum_{\alpha=0}^{a-1} \sum_{\delta=0}^{a-\alpha-1} \sum_{j=0}^{\delta} \binom{n}{\alpha+\delta} \binom{\alpha+\delta}{\alpha} \sum_{i=0}^{a-\alpha-\delta-1} (-1)^{\alpha+j+i} \binom{n-\alpha-\delta}{i} 2^{n-1-\alpha-j} \binom{\delta}{j} I(\alpha+j, 0). \quad (4-86)$$

The terms are collected according to the values of $\omega := \alpha + j$. Upon simplification, it follows that

$$A' = 2^{n-1} \sum_{\omega=0}^{a-1} 2^{-\omega} (-1)^{\omega} I(\omega, 0) \sum_{\alpha=0}^{\omega} \sum_{\delta=\omega-\alpha}^{a-1-\alpha} \sum_{i=0}^{a-\delta-\alpha-1} (-1)^i \binom{n}{\delta+\alpha} \binom{\delta+\alpha}{\alpha} \binom{n-\delta-\alpha}{i} \binom{\delta}{\omega-\alpha}. \quad (4-87)$$

We then make a change of variables $\delta = \ell + \omega - \alpha$ and rewrite the sums above as

$$\begin{aligned} A' &= 2^{n-1} \sum_{\omega=0}^{a-1} 2^{-\omega} (-1)^{\omega} I(\omega, 0) \sum_{\alpha=0}^{\omega} \sum_{\ell=0}^{a-1-\omega} \sum_{i=0}^{a-\ell-\omega-1} (-1)^i \binom{n}{\ell+\omega} \binom{\ell+\omega}{\alpha} \binom{n-\ell-\omega}{i} \binom{\ell+\omega-\alpha}{\omega-\alpha} \\ &= 2^{n-1} \sum_{\omega=0}^{a-1} 2^{-\omega} (-1)^{\omega} I(\omega, 0) \sum_{\alpha=0}^{\omega} \sum_{\ell=0}^{a-1-\omega} \sum_{i=0}^{a-\ell-\omega-1} (-1)^i \binom{n}{\ell+\omega+i} \binom{\ell+\omega+i}{\ell+i} \binom{\omega}{\alpha} \binom{\ell+i}{i}. \end{aligned} \quad (4-88)$$

Grouping terms based on the values $m = \ell + i$ and rearranging, we have

$$A' = 2^{n-1} \sum_{\omega=0}^{a-1} 2^{-\omega} (-1)^{\omega} I(\omega, 0) \sum_{\alpha=0}^{\omega} \binom{\omega}{\alpha} \sum_{m=0}^{a-1-\omega} \binom{n}{m+\omega} \binom{m+\omega}{m} \sum_{i=0}^m (-1)^i \binom{m}{i}. \quad (4-89)$$

As a consequence of the binomial theorem, the sum over i is zero unless $m = 0$. Thus, summing over α yields $\binom{n}{\omega} 2^{\omega}$ and so,

$$A' = 2^{n-1} \sum_{\omega=0}^{a-1} \binom{n}{\omega} (-1)^{\omega} I(\omega, 0). \quad (4-90)$$

Applying this to (4-82), we find that $A = (-1)^{n+1} \mathcal{R}(n, a; \phi) + O\left(\frac{(\log \log N)^2}{(\log N)^{1/2}}\right)$, where $\mathcal{R}(n, a; \phi)$ was defined in (1-13). This completes the proof of Proposition 3.7. \square

5. Extending support for random matrix theory: proof of Theorem 1.3

In this section, we prove Theorem 1.3. We focus on the case where $n \geq 3$ as [21, Theorem 1.7] have proved the $n = 2$ case.

In Section 5.1, we use results from [22] and [21] to reduce the proof of Theorem 1.3 to proving Proposition 5.2, which gives a closed form expression for the quantity $Q_n(\phi)$ defined in (5-4).

The rest of the section is dedicated to evaluating $Q_n(\phi)$. In Section 5.2, we define the notion of a *system of parameters* and of a *t-class*, and express $Q_n(\phi)$ as a sum over these two objects in Lemma 5.12. Lemma 5.12 splits $Q_n(\phi)$ into a combinatorial piece and an integral piece. We evaluate the combinatorial piece in Section 5.3. In particular, we calculate the contribution to $Q_n(\phi)$ from *t-classes* with $t = 1$ using Lemma 5.19, and show that the *t-classes* with $t \geq 2$ do not contribute in Lemma 5.20. Then, in Section 5.4, we evaluate the integral piece, completing the proof of Proposition 5.2.

5.1. Proof of Theorem 1.3 assuming Proposition 5.2. We calculate the n -th centered moments of $Z(U; \phi)$, denoted in Section 1.3 by $\mathcal{Z}_n(M; \phi)$, using the method of cumulants. Weyl’s explicit representation of Haar measure would allow us to compute the higher moments directly. However, to facilitate the comparison with number theory, we use the cumulants as in [22] and [21]. The cumulants $C_\ell^+(\phi)$ and $C_\ell^-(\phi)$ are defined to satisfy the following equality of formal power series:

$$\sum_{\ell=1}^{\infty} C_\ell^+(\phi) \frac{\lambda^\ell}{\ell!} = \lim_{\substack{M \text{ even} \\ M \rightarrow \infty}} \log \mathbb{E}_{\text{SO}(M)}[\exp(\lambda Z(U; \phi))], \tag{5-1}$$

$$\sum_{\ell=1}^{\infty} C_\ell^-(\phi) \frac{\lambda^\ell}{\ell!} = \lim_{\substack{M \text{ odd} \\ M \rightarrow \infty}} \log \mathbb{E}_{\text{SO}(M)}[\exp(\lambda Z(U; \phi))]. \tag{5-2}$$

Given the first n cumulants, one can compute the first n moments. In particular, for $n \geq 2$, we have

$$\lim_{M \rightarrow \infty}^\pm \mathcal{Z}_n(M; \phi) = \sum_{\substack{2k_2+3k_3+\dots+nk_n=n \\ k_j \geq 0}} \left(\frac{C_2^\pm(\phi)}{2!}\right)^{k_2} \cdots \left(\frac{C_n^\pm(\phi)}{n!}\right)^{k_n} \frac{n!}{k_2! \cdots k_n!}. \tag{5-3}$$

Now, set $S(x) := \frac{\sin(\pi x)}{\pi x}$ and define

$$Q_n(\phi) := 2^{n-1} \sum_{m=1}^n \sum_{\substack{\lambda_1+\dots+\lambda_m=n \\ \lambda_j \geq 1}} \frac{(-1)^{m+1}}{m} \frac{n!}{\lambda_1! \cdots \lambda_m!} \int_{-\infty}^{\infty} \cdots \int_{-\infty}^{\infty} \phi(x_1)^{\lambda_1} \cdots \phi(x_m)^{\lambda_m} \\ \times S(x_1 - x_2) S(x_2 - x_3) \cdots S(x_{m-1} - x_m) S(x_m + x_1) dx_1 \cdots dx_m. \tag{5-4}$$

Thanks to the next lemma, in order to prove Theorem 1.3, it suffices to calculate $Q_n(\phi)$, which we do in the subsequent proposition.

Lemma 5.1 [22, Section 2.1]. *Let $\phi \in \mathcal{S}_{ec}(\mathbb{R})$ with $\text{supp}(\hat{\phi}) \subseteq [-\frac{2}{n}, \frac{2}{n}]$. For $n \geq 3$,*

$$C_n^{\text{SO(even)}}(\phi) = Q_n(\phi) \quad \text{and} \quad C_n^{\text{SO(odd)}}(\phi) = -Q_n(\phi). \tag{5-5}$$

For $n \geq 4$,

$$C_2^{\text{SO(even)}} = C_2^{\text{SO(odd)}} = 2 \int_{-\infty}^{\infty} |y| \hat{\phi}(y)^2 dy = \sigma_\phi^2 \tag{5-6}$$

where σ_ϕ^2 is as in (1-13).

Proposition 5.2. *Let $\phi \in \mathcal{S}_{ec}(\mathbb{R})$ with $\text{supp}(\hat{\phi}) \subseteq [-\frac{1}{n-a}, \frac{1}{n-a}]$ for some integer $1 \leq a \leq \lceil n/2 \rceil$. Let $R(n, a; \phi)$ be as in (1-13). Then*

$$Q_n(\phi) = \mathcal{R}(n, a; \phi). \tag{5-7}$$

We complete the proof of Proposition 5.2 in Section 5.4. Assuming Proposition 5.2, we now prove Theorem 1.3.

Proof of Theorem 1.3. By [21, Theorem 1.4], since $\text{supp}(\hat{\phi}) \subseteq [-\frac{1}{n-a}, \frac{1}{n-a}]$, we have $C_j^\pm(\phi) = 0$ for $3 \leq j \leq n - a$, as in this case the cumulants are the same as those of the Gaussian. Hence, restricting the sum in (5-3) to those terms with $k_3 = \cdots = k_{n-a} = 0$ does not change its value. Moreover, $a \leq \lceil n/2 \rceil$

and $\sum_{\ell=2}^n \ell k_\ell = n$ imply that $k_n, k_{n-1}, \dots, k_{n-a+1} \in \{0, 1\}$ and at most one of $k_n, k_{n-1}, \dots, k_{n-a+1}$ is equal to 1. Thus, we can rewrite (5-3) as

$$\lim_{M \rightarrow \infty}^\pm \mathcal{Z}_n(M; \phi) = \mathbb{1}_{\{n \text{ even}\}} \left(\frac{C_2^\pm(\phi)}{2} \right)^{n/2} \frac{n!}{(n/2)!} + \sum_{\substack{k_2, \ell \\ 2k_2 + (n-\ell) = n \\ 0 \leq \ell \leq a-1}} \left(\frac{C_2^\pm(\phi)}{2!} \right)^{k_2} \left(\frac{C_{n-\ell}^\pm(\phi)}{(n-\ell)!} \right) \frac{n!}{k_2!} \tag{5-8}$$

where the first term is from when $k_2 = n/2$. Observing that $2k_2 + (n - \ell) = n$ forces $\ell = 2k_2$, we have

$$\lim_{M \rightarrow \infty}^\pm \mathcal{Z}_n(M; \phi) = \mathbb{1}_{\{n \text{ even}\}} (C_2^\pm(\phi))^{n/2} (2n - 1)!! + \sum_{k_2=0}^{\lfloor \frac{a-1}{2} \rfloor} \frac{n! C_{n-2k_2}^\pm(\phi)}{k_2! (n - 2k_2)!} \left(\frac{C_2^\pm(\phi)}{2} \right)^{k_2}. \tag{5-9}$$

Now, applying Lemma 5.1 and Proposition 5.2 to the right hand side of (5-9) and simplifying completes the proof of Theorem 1.3 after comparing with (1-15). □

5.2. Decomposition of $Q_n(\phi)$. In this section, we work towards Proposition 5.2 by evaluating $Q_n(\phi)$, as defined in (5-4), when $\text{supp}(\hat{\phi}) \subseteq [-\frac{1}{n-a}, \frac{1}{n-a}]$ and $a \leq \lceil n/2 \rceil$. The main result of this subsection is Lemma 5.12, which splits $Q_n(\phi)$ into a combinatorial term and an integral term which we will then evaluate separately.

Equation (5.27) of [21] gives (independently of the choice of support) that

$$Q_n(\phi) = 2^{n-2} \int_0^\infty \cdots \int_0^\infty \hat{\phi}(y_1) \cdots \hat{\phi}(y_n) K(y_1, \dots, y_n) dy_1 \cdots dy_n, \tag{5-10}$$

where

$$K(y_1, \dots, y_n) := \sum_{m=1}^n \sum_{\substack{\lambda_1 + \dots + \lambda_m = n \\ \lambda_j \geq 1}} \frac{(-1)^{m+1}}{m} \frac{n!}{\lambda_1! \cdots \lambda_m!} \sum_{\epsilon_1, \dots, \epsilon_n \in \{\pm 1\}} \prod_{\ell=1}^m \chi_{\left\{ \left| \sum_{j=1}^n \eta(\ell, j) \epsilon_j y_j \right| \leq 1 \right\}} \tag{5-11}$$

and

$$\eta(\ell, j) := \begin{cases} +1 & \text{if } j \leq \sum_{k=1}^\ell \lambda_k, \\ -1 & \text{if } j > \sum_{k=1}^\ell \lambda_k. \end{cases} \tag{5-12}$$

5.2.1. Simplifying $K(y_1, \dots, y_n)$. To evaluate $Q_n(\phi)$, we first discuss how we will interpret the expression $K(y_1, \dots, y_n)$ for $y_1, \dots, y_n \in [0, \frac{1}{n-a}]$.

Definition 5.3. Throughout this section, if $I \subseteq \{1, \dots, n\}$, we write

$$\chi_I = \chi_{\{y_1 + \dots + y_n > 1 + 2 \sum_{i \in I} y_i\}}. \tag{5-13}$$

Definition 5.4. A system of parameters, abbreviated *s.o.p.*, is an ordered tuple $(m, \lambda_1, \dots, \lambda_m, \epsilon_1, \dots, \epsilon_n)$ with $1 \leq m \leq n$, $\lambda_1 + \dots + \lambda_m = n$, $\lambda_i \geq 1$ for all $1 \leq i \leq m$, and $\epsilon_j = \pm 1$ for each $1 \leq j \leq n$.

Given a system of parameters S , we may use $\eta_S(\ell, j)$ to denote the function $\eta(\ell, j)$ where the λ_k are taken from S . When it is clear from context that the λ_k are taken from the s.o.p. S , we simply denote this

function $\eta(\ell, j)$. Fix $n \geq 2a$ and an s.o.p. $S = (m, \lambda_1, \dots, \lambda_m, \epsilon_1, \dots, \epsilon_n)$. Consider the product

$$\prod_{\ell=1}^m \chi_{\{|\sum_{j=1}^n \eta(\ell, j)\epsilon_j y_j| \leq 1\}} \tag{5-14}$$

from (5-11). Fix $1 \leq \ell_0 \leq m$. In order to study (5-14), we study the complement of the indicator functions in (5-14), given by

$$\chi_{\{|\sum_{j=1}^n \eta(\ell_0, j)\epsilon_j y_j| > 1\}}. \tag{5-15}$$

For $y_1, \dots, y_n \in [0, \frac{1}{n-a}]$, if $\sum_{j=1}^n \eta(\ell_0, j)\epsilon_j y_j > 1$ then we cannot find $y'_1, \dots, y'_n \in [0, \frac{1}{n-a}]$ such that $\sum_{j=1}^n \eta(\ell_0, j)\epsilon_j y'_j < -1$ because $a \leq \lceil n/2 \rceil$. Thus the indicator function (5-15) is identical to (5-13) for a particular choice of I . Moreover, there exists $y_i \in [0, \frac{1}{n-a}]$ such that (5-15) is nonzero if and only if one of the following mutually exclusive conditions holds:

- (i) $|\{1 \leq j \leq n : \eta(\ell_0, j)\epsilon_j = +1\}| \leq a - 1$.
- (ii) $|\{1 \leq j \leq n : \eta(\ell_0, j)\epsilon_j = -1\}| \leq a - 1$.

In case (i), we define

$$J_{\ell_0} := \{1 \leq j \leq n : \eta(\ell_0, j)\epsilon_j = +1\} \tag{5-16}$$

and say that J_{ℓ_0} has sign $\zeta_{\ell_0} = +1$. In case (ii), we define

$$J_{\ell_0} := \{1 \leq j \leq n : \eta(\ell_0, j)\epsilon_j = -1\} \tag{5-17}$$

and say that J_{ℓ_0} has sign $\zeta_{\ell_0} = -1$. If neither case holds, then J_{ℓ_0} is undefined.

Lemma 5.5. *If $S = (m, \lambda_1, \dots, \lambda_m, \epsilon_1, \dots, \epsilon_n)$ is an s.o.p. and $J \subseteq [1, n]$ is any subset, then there is at most one $\ell_0 \in [1, m]$ and $\zeta \in \{\pm 1\}$ such that $\eta(\ell_0, i)\epsilon_i = \zeta$ for $i \in J$ and $\eta(\ell_0, j)\epsilon_j = -\zeta$ for $j \notin J$.*

Proof. Suppose $\ell_1 > \ell_0$ and that both ℓ_0 and ℓ_1 have this property for some ζ_0 and ζ_1 . Without loss of generality, we assume that $J = \{i : \eta(\ell_0, i)\epsilon_i = -1\}$. It is clear that we cannot also have $I = \{i : \eta(\ell_1, i)\epsilon_i = -1\}$, so we may assume that $I = \{i : \eta(\ell_1, i)\epsilon_i = +1\}$, but then we must have $\eta(\ell_0, j) = -\eta(\ell_1, j)$ for all j , and this is clearly impossible. □

In particular, if J_{ℓ_0} and J_{ℓ_1} are both defined, then $J_{\ell_0} \neq J_{\ell_1}$.

Definition 5.6. For an s.o.p. $S = (m, \lambda_1, \dots, \lambda_m, \epsilon_1, \dots, \epsilon_n)$, let $\{\ell_1, \dots, \ell_t\} \subseteq \{1, \dots, m\}$ be the set of indices for which J_{ℓ_j} is defined. Set

$$J(S) := \{J_{\ell_1}, \dots, J_{\ell_t}\}, \tag{5-18}$$

and define

$$I(S) := \{I_1, \dots, I_r\} \tag{5-19}$$

to be the subset of elements of $J(S)$ which are minimal with respect to inclusion. That is, $I(S)$ consists of those elements of $J(S)$ which do not strictly contain any other elements of $J(S)$. By Lemma 5.5, for

each $i \in [1, r]$ there is a unique ℓ_i such that $I_i = J_{\ell_i}$. Finally, define the function

$$\sigma^S(y_1, \dots, y_n) := \sum_{i=1}^r \sum_{1 \leq j_1 < \dots < j_i \leq r} (-1)^i (\chi_{I_{j_1}} \cdots \chi_{I_{j_i}})(y_1, \dots, y_n), \tag{5-20}$$

and the quantity

$$A(S) := \frac{(-1)^{m+1}}{m} \frac{n!}{\lambda_1! \cdots \lambda_m!}. \tag{5-21}$$

The next lemma for σ^S resembles the Möbius inversion formula from elementary number theory.

Lemma 5.7. *For any s.o.p. S , we have*

$$\sigma^S(y_1, \dots, y_n) = \begin{cases} -1 & \text{if } \chi_I(y_1, \dots, y_n) = 1 \text{ for some } I \in I(S), \\ 0 & \text{otherwise.} \end{cases} \tag{5-22}$$

Proof. Fix (y_1, \dots, y_n) . Suppose there are k elements in $I(S)$ whose support contains (y_1, \dots, y_n) . If $k = 0$ the result is immediate. Now, for $k \geq 1$ and $1 \leq i \leq k$, there are $\binom{k}{i}$ terms in the i -th summand of with coefficient $(-1)^i$ and all the other terms vanish. Thus we have

$$\sigma^S(y_1, \dots, y_n) = \sum_{i=1}^k \binom{k}{i} (-1)^i = (1-1)^k - 1 = -1. \quad \square$$

We now reformulate the quantity from (5-11) in terms of $A(S)$, defined in (5-21).

Lemma 5.8. *For $(y_1, \dots, y_n) \in [0, \frac{1}{n-a}]^n$,*

$$K(y_1, \dots, y_n) = \sum_{t=1}^n (-1)^t \sum_{\substack{(I_1, \dots, I_t) \\ \text{valid}}} (\chi_{I_1} \cdots \chi_{I_t})(y_1, \dots, y_n) \sum_{\substack{\text{s.o.p. } S \text{ with} \\ I_1, \dots, I_t \in I(S)}} A(S). \tag{5-23}$$

Proof. The product (5-14) vanishes at (y_1, \dots, y_n) if and only if there is some $J \in J(S)$ such that χ_J is supported at (y_1, \dots, y_n) , (and thus) if and only if there is some $I \in I(S)$ such that χ_I is supported at (y_1, \dots, y_n) . So, by Lemma 5.7,

$$\prod_{\ell=1}^m \chi_{\{\sum_{j=1}^n \eta(\ell, j) \epsilon_j y_j \leq 1\}}(y_1, \dots, y_n) = 1 + \sigma^S(y_1, \dots, y_n). \tag{5-24}$$

Substituting (5-24) into (5-11), we have

$$K(y_1, \dots, y_n) = \sum_{m=1}^n \sum_{\substack{\lambda_1 + \dots + \lambda_m = n \\ \lambda_j \geq 1}} \frac{(-1)^{m+1}}{m} \frac{n!}{\lambda_1! \cdots \lambda_m!} \cdot 2^n + \sum_{\text{s.o.p.'s } S} A(S) \sigma^S(y_1, \dots, y_n). \tag{5-25}$$

Now, we apply an identity given by Soshnikov [45],

$$z = \log(1 + (e^z - 1)) = \sum_{n=1}^{\infty} z^n \sum_{m=1}^n \sum_{\substack{\lambda_1 + \dots + \lambda_m = n \\ \lambda_j \geq 1}} \frac{(-1)^{m+1}}{m} \frac{1}{\lambda_1! \cdots \lambda_m!}, \tag{5-26}$$

to conclude that the first sum in (5-25) is 0. Expanding the second sum using the definition of σ^S from (5-20) and rearranging completes the proof. □

5.2.2. Simplifying $Q_n(\phi)$. In this section, we simplify $Q_n(\phi)$ by applying Lemma 5.8 to the quantity $Q_n(\phi)$ as in (5-10). First, we define further notation which allows us to express $Q_n(\phi)$ (through Lemma 5.12) in terms of combinatorial quantities which we then compute in Section 5.3.1 and 5.3.2.

The symmetric group S_n acts naturally on sets of (unordered) t -tuples of subsets of $[1, n]$ by permuting the elements in each subset of each tuple. Take such a t -tuple (I_1, \dots, I_t) and some $I_j = \{i_1, \dots, i_k\}$. Given some permutation $\tau \in S_n$, we have $\tau(I_j) = \{\tau(i_1), \dots, \tau(i_k)\}$. Let $\chi_{I_1} \cdots \chi_{I_t}$ and $\chi_{J_1} \cdots \chi_{J_t}$ be elements of Ω such that there exists a permutation $\tau \in S_n$ so that for each $1 \leq \ell \leq t$, $\tau(I_\ell) = J_\ell$. Then,

$$\int_0^\infty \cdots \int_0^\infty \hat{\phi}(y_1) \cdots \hat{\phi}(y_n) (\chi_{I_1} \cdots \chi_{I_t} - \chi_{J_1} \cdots \chi_{J_t})(y_1, \dots, y_n) dy_1 \cdots dy_n = 0. \tag{5-27}$$

This motivates the following definition.

Definition 5.9. The symmetric group S_n acts naturally on sets of (unordered) t -tuples of subsets of $[1, n]$, as described above. An orbit of this action is called a t -class.

For a t -class C , we define

$$\int C dy := \int_0^\infty \cdots \int_0^\infty \hat{\phi}(y_1) \cdots \hat{\phi}(y_n) \chi(y_1, \dots, y_n) dy_1 \cdots dy_n, \tag{5-28}$$

where $\chi := \chi_{I_1} \cdots \chi_{I_t}$ with $(I_1, \dots, I_t) \in C$. Equation (5-27) shows that the integral $\int C dy$ is well-defined.

Definition 5.10. We call an unordered tuple (I_1, \dots, I_t) of subsets of $\{1, \dots, n\}$ *valid* if $I_1, \dots, I_t \in I(S)$ for some s.o.p. S and $\chi_{I_1} \cdots \chi_{I_t}$ is supported at some point in $[0, \frac{1}{n-a}]^n$.

We can extend this definition to a t -class.

Definition 5.11. We call a t -class *valid* if it contains at least one valid tuple.

We are now ready to prove the main result of the section.

Lemma 5.12. For an s.o.p. S and a t -class C , set

$$T(S, C) := \#\{(I_1, \dots, I_t) \in C : I_1, \dots, I_t \in I(S)\}. \tag{5-29}$$

We have

$$Q_n(\phi) = 2^{n-2} \sum_{t=1}^n (-1)^t \sum_{\text{valid } t\text{-classes } C} \left(\sum_{\text{s.o.p.'s } S} T(S, C) A(S) \right) \int C dy. \tag{5-30}$$

Proof. Given a valid t -class C , there is a valid tuple $(I_1, \dots, I_t) \in C$ for which $\chi_{I_1} \cdots \chi_{I_t}$ is supported at some point $(y_1, \dots, y_n) \in [-\frac{1}{n-a}, \frac{1}{n-a}]^n$. Therefore, if $\tau \in S_n$, then $\chi_{\tau(I_1)} \cdots \chi_{\tau(I_t)}$ is supported at $(y_{\tau(1)}, \dots, y_{\tau(n)})$. Since S_n acts transitively on C , this means that every tuple in C is valid. Now, applying Lemma 5.8 to (5-10) and grouping tuples into t -classes completes the proof. \square

5.3. Computing the combinatorial piece. In this section, we calculate $\sum T(S, C) A(S)$ for valid t -classes C , where $T(S, C)$ and $A(S)$ are defined as in (5-29) and (5-21), respectively. In Section 5.3.1 we find a closed form for the case $t = 1$, and then in Section 5.3.2 we show that when $t \geq 2$ the quantity vanishes.

5.3.1. Computing for valid 1-classes. In this section, we compute the terms in (5-30) for which $t = 1$. We first classify the valid 1-tuples.

Lemma 5.13. *If I and J are subsets of $[1, n]$ such that $|I \cup J| \geq a$, then $\chi_I \cdot \chi_J$ is identically zero on $[0, \frac{1}{n-a}]^n$.*

Proof. Let I and J be as in the hypotheses, and assume for contradiction that both $y_1 + \dots + y_n > 1 + 2 \sum_{i \in I} y_i$ and $y_1 + \dots + y_n > 1 + 2 \sum_{j \in J} y_j$ for some $(y_1, \dots, y_n) \in [0, \frac{1}{n-a}]^n$. Since $y_h \leq \frac{1}{n-a}$ for every h , $\sum_{h \notin I \cup J} y_h \leq 1$, so we must have that $\sum_{i \in I} y_i < \sum_{j \in J \setminus I} y_j$ and similarly $\sum_{j \in J} y_j < \sum_{i \in I \setminus J} y_i$ by our assumptions. Adding these inequalities gives

$$\sum_{i \in I} y_i + \sum_{j \in J} y_j < \sum_{i \in I \setminus J} y_i + \sum_{j \in J \setminus I} y_j, \tag{5-31}$$

which is a contradiction, as all the y_h 's are nonnegative and the terms on the right are a subset of those on the left. □

Lemma 5.14. *If I is a subset of $[1, n]$, then the 1-tuple (I) is valid if and only if $|I| \leq a - 1$.*

Proof. If $|I| > a - 1$, then (I) is not valid by Lemma 5.13, taking both subsets to be I .

Now suppose $|I| \leq a - 1$. Let $y_j = 1/(n - a)$ for each $j \notin I$ and let $y_i = 0$ for each $i \in I$. It is clear that $\chi_I(y_1, \dots, y_n) = 1$. Now consider the system of parameters $S = (m, \lambda_1, \dots, \lambda_m, \epsilon_1, \dots, \epsilon_n)$, where $m = 1$, $\lambda_1 = n$, and $\epsilon_i = -1$ if and only if $i \in I$. Clearly, $I \in I(S)$. Therefore, (I) is valid. □

It follows from Lemma 5.14 that the valid 1-classes are exactly the classes

$$C_f := \{(I) : I \subseteq [1, n], |I| = f\} \tag{5-32}$$

with $0 \leq f \leq a - 1$.

Lemma 5.15. *Let $1 \leq f \leq a - 1$. Let $S = (m, \lambda_1, \dots, \lambda_m, \epsilon_1, \dots, \epsilon_n)$ be a system of parameters with $m \geq 2$ and suppose $(I) \in C_f$ is such that, for some $1 \leq \ell \leq m$, we have $I = J_\ell \in J(S)$. Define $\Lambda_\ell := \lambda_1 + \dots + \lambda_\ell$. Then $I \in I(S)$ if and only if $[\Lambda_{\ell-1} + 1, \Lambda_\ell] \not\subseteq I$ and $[\Lambda_\ell + 1, \Lambda_{\ell+1}] \not\subseteq I$. If $\ell = m$, then we set $[\Lambda_m + 1, \Lambda_{m+1}]$ to $[1, \Lambda_1] = [1, \lambda_1]$.*

Proof. Assume without loss of generality that J_ℓ has sign $\zeta_\ell = -1$, i.e., $J_\ell = \{j : \eta(\ell, j)\epsilon_j = -1\}$.

For any $\ell' < \ell$, we have

$$\eta(\ell, j)\epsilon_j = \begin{cases} -\eta(\ell', j)\epsilon_j & \text{if } j \in [\Lambda_{\ell'} + 1, \Lambda_\ell], \\ \eta(\ell', j)\epsilon_j & \text{if } j \notin [\Lambda_{\ell'} + 1, \Lambda_\ell]. \end{cases} \tag{5-33}$$

If $[\Lambda_{\ell-1} + 1, \Lambda_\ell] \subseteq J_\ell$, then $J_{\ell-1} = J_\ell \setminus [\Lambda_{\ell-1} + 1, \Lambda_\ell] \subsetneq J_\ell$. In particular, J_ℓ is not minimal, so $J_\ell \notin I(S)$. Similarly, if $[\Lambda_\ell + 1, \Lambda_{\ell+1}] \subseteq J_\ell$ then $J_{\ell+1} = J_\ell \setminus [\Lambda_\ell + 1, \Lambda_{\ell+1}]$ so J_ℓ is not minimal.

Now assume J_ℓ is not minimal, so there exists some $J_{\ell'} \subsetneq J_\ell$.

First, suppose the sign of $J_{\ell'}$ is $\zeta_{\ell'} = -1$. Suppose that $\ell' < \ell$. By (5-33), $J_{\ell'} \setminus [\Lambda_{\ell'} + 1, \Lambda_\ell] = J_\ell \setminus [\Lambda_{\ell'} + 1, \Lambda_\ell]$, while $J_{\ell'} \cap [\Lambda_{\ell'} + 1, \Lambda_\ell]$ and $J_\ell \cap [\Lambda_{\ell'} + 1, \Lambda_\ell]$ are disjoint with union $[\Lambda_{\ell'} + 1, \Lambda_\ell]$. So, so $J_{\ell'} \subsetneq J_\ell$ implies $[\Lambda_{\ell-1} + 1, \Lambda_\ell] \subseteq [\Lambda_{\ell'} + 1, \Lambda_\ell] \subset J_\ell$. Similarly, if $\ell' > \ell$, we have $[\Lambda_\ell + 1, \Lambda_{\ell+1}] \subset J_\ell$.

Next suppose that $\text{sign } \zeta_{\ell'} = 1$. Suppose that $\ell' < \ell$. By (5-33), $J_{\ell'} \cap [\Lambda_{\ell'} + 1, \Lambda_{\ell}] = J_{\ell} \cap [\Lambda_{\ell'} + 1, \Lambda_{\ell}]$, while $J_{\ell'} \setminus [\Lambda_{\ell'} + 1, \Lambda_{\ell}]$ and $J_{\ell} \setminus [\Lambda_{\ell'} + 1, \Lambda_{\ell}]$ are disjoint with union $[1, n] \setminus [\Lambda_{\ell'} + 1, \Lambda_{\ell}]$. Since J_{ℓ} is not minimal, we must have $[\Lambda_{\ell} + 1, \Lambda_{\ell+1}] \subset [1, n] \setminus [\Lambda_{\ell-1} + 1, \Lambda_{\ell}] \subset J_{\ell}$. When $\ell' > \ell$, by the same reasoning we have $[\Lambda_{\ell-1} + 1, \Lambda_{\ell}] \subseteq J_{\ell}$. \square

Lemma 5.16. *Fix $1 \leq f \leq a - 1$. We have*

$$\sum_{\substack{\text{s.o.p.'s } S \\ \text{with } m \geq 2}} T(S, C_f) A(S) = 2n!(-1)^n \sum_{\substack{c+d \leq n \\ c, d \geq 0}} (-1)^{c+d+1} G(n, f, c, d) \frac{1}{(n-c-d)!c!d!}, \tag{5-34}$$

where

$$G(n, f, c, d) := \binom{n}{f} - \binom{n-c}{f-c} - \binom{n-d}{f-d} + \binom{n-c-d}{f-c-d}. \tag{5-35}$$

Proof. Let $S = (m, \lambda_1, \dots, \lambda_m, \epsilon_1, \dots, \epsilon_n)$ denote a variable system of parameters. By Lemma 5.5, we can rewrite $T(S, C_f)$ as

$$T(S, C_f) = \sum_{\ell=1}^m \mathbb{1}_{\{J_{\ell} \in I(S) \text{ and } \#J_{\ell} = f\}}. \tag{5-36}$$

We sum over systems of parameters by first summing over all values of m , then summing over all possible values of ℓ , then summing over all possible values of $c = \lambda_{\ell}$ and $d = \lambda_{\ell+1}$, then summing over all possible values of $\lambda_1, \dots, \lambda_m$ and finally summing over all possible choices of $\epsilon_1, \dots, \epsilon_n$. For fixed $m, \lambda_1, \dots, \lambda_m$, the $\epsilon_1, \dots, \epsilon_n$ and J_{ℓ}, ζ_{ℓ} uniquely determine each other, so we may rewrite the innermost sum as

$$\sum_{(\epsilon_j) \in \{\pm 1\}^n} A(S) \mathbb{1}_{\{J_{\ell} \in I(S) \text{ and } \#J_{\ell} = f\}} = A(S) \sum_{\zeta_{\ell} \in \{\pm 1\}} \sum_{\#J_{\ell} = f} \mathbb{1}_{\{J_{\ell} \in I(S)\}}. \tag{5-37}$$

By Lemma 5.15, the sum over J_{ℓ} is $G(n, f, c, d)$, since we can choose a general f element subset in $\binom{n}{f}$ ways, and we need to subtract off when the c element subset $[\lambda_1 + \dots + \lambda_{\ell-1} + 1, \lambda_1 + \dots + \lambda_{\ell}] \subseteq I$ or when the d element subset $[\lambda_1 + \dots + \lambda_{\ell} + 1, \lambda_1 + \dots + \lambda_{\ell+1}] \subseteq I$. Then, we add back in the case when both subsets are contained in J_{ℓ} since we have double counted it. Finally, there are two choices for ζ_{ℓ} . We have

$$\sum_{\substack{\text{s.o.p.'s } S \\ \text{with } m \geq 2}} T(S, C_f) A(S) = \sum_{m=2}^n \sum_{\ell=1}^m \sum_{\substack{c, d \geq 1 \\ c+d \leq n}} \sum_{\substack{\lambda_1 + \dots + \lambda_m = n \\ \lambda_i \geq 1, \lambda_{\ell} = c, \lambda_{\ell+1} = d}} \frac{(-1)^{m+1}}{m} \frac{n!}{\lambda_1! \dots \lambda_m!} 2 G(n, f, c, d). \tag{5-38}$$

Noting that for each value of ℓ the inner summand is the same, we can set $\ell = m - 1$ and write

$$\sum_{\substack{\text{s.o.p.'s } S \\ \text{with } m \geq 2}} T(S, C_f) A(S) = 2n! \sum_{\substack{c, d \geq 1 \\ c+d \leq n}} \frac{G(n, f, c, d)}{c!d!} \sum_{m=2}^n m \sum_{\substack{\lambda_1 + \dots + \lambda_{m-2} \\ = n-c-d}} \frac{(-1)^{m+1}}{m} \frac{1}{\lambda_1! \dots \lambda_{m-2}!}. \tag{5-39}$$

The sum over m equals $(-1)^{n+c+d+1}/(n-c-d)!$, which follows from evaluating the coefficient of z^n in

$$\sum_{n=0}^{\infty} \frac{(-1)^n}{n!} z^n = e^{-z} = \frac{1}{1+(e^z-1)} = \sum_{n=0}^{\infty} z^n \sum_{m=1}^n \sum_{\substack{\lambda_1 + \dots + \lambda_m = n \\ \lambda_j \geq 1}} \frac{(-1)^m}{\lambda_1! \dots \lambda_m!}. \tag{5-40}$$

Applying this to (5-39) gives

$$\sum_{\substack{\text{s.o.p.'s } S \\ \text{with } m \geq 2}} T(S, C_f) A(S) = 2n!(-1)^n \sum_{\substack{c+d \leq n \\ c, d \geq 1}} (-1)^{c+d+1} G(n, f, c, d) \frac{1}{(n-c-d)!c!d!}. \tag{5-41}$$

Now, we can extend the sum to include when $c = 0$ or $d = 0$ to complete the proof as in this case $G(n, f, c, d) = 0$. □

We complete our evaluation of the case when $m \geq 2$ with the following lemma, proven in Appendix C.1.

Lemma 5.17. *Fix $1 \leq f \leq a - 1$. We have*

$$\sum_{\substack{\text{s.o.p.'s } S \\ \text{with } m \geq 2}} T(S, C_f) A(S) = 2 \binom{n}{f} ((-1)^{n+f+1} - 1). \tag{5-42}$$

Now we evaluate the case when $m = 1$.

Lemma 5.18. *Fix $1 \leq f \leq a - 1$. We have*

$$\sum_{\substack{\text{s.o.p.'s } S \\ \text{with } m=1}} T(S, C_f) A(S) = 2 \binom{n}{f}. \tag{5-43}$$

Proof. We let $S = (1, \lambda_1, \epsilon_1, \dots, \epsilon_n)$ denote a variable system of parameters. Since $m = 1$, we have $\lambda_1 = n$ and $A(S) = \frac{(-1)^2 n!}{1 n!} = 1$ for all S . Now, as in (5-37), we may rewrite the sum over $\epsilon_1, \dots, \epsilon_n$ as a sum over J_1, ζ_1 . Since $m = 1$, any f -element $J_1 \in J(S)$ will be minimal. So,

$$\sum_{\substack{\text{s.o.p.'s } S \\ \text{with } m \geq 2}} T(S, C_f) A(S) = A(S) \sum_{\zeta_1 \in \{\pm 1\}} \sum_{\#J_1=f} \mathbb{1}_{\{J_1 \in I(S)\}} = \sum_{\zeta_1 \in \{\pm 1\}} \sum_{\#J_1=f} 1 = 2 \binom{n}{f}. \tag{5-43} \quad \square$$

Adding (5-42) and (5-43) gives the main result of the section.

Lemma 5.19. *Fix $1 \leq f \leq a - 1$. Then*

$$\sum_{\text{s.o.p.'s } S} T(S, C_f) A(S) = 2(-1)^{n+f+1} \binom{n}{f}. \tag{5-44}$$

5.3.2. The vanishing of valid t -classes for $t \geq 2$. In this section, we show that all terms with $t \geq 2$ in (5-30) vanish. Our main result is the following.

Lemma 5.20. *Let C be a valid t -class with $t \geq 2$. Then*

$$\sum_{\text{s.o.p.'s } S} T(S, C) A(S) = 0. \tag{5-45}$$

Throughout this section, let $S = (m, \lambda_1, \dots, \lambda_m, \epsilon_1, \dots, \epsilon_n)$ be a system of parameters, C a valid class, and $(I_1, \dots, I_t) \in C$ a tuple of subsets of $[1, n]$ such that for each $1 \leq i \leq t$, there is some ℓ_i and $\zeta_{\ell_i} \in \{\pm 1\}$ such that $I_i = \{j : \eta(\ell_i, j) \epsilon_j = \zeta_{\ell_i}\}$. I.e., $I_i = J_{\ell_i}$ with sign ζ_{ℓ_i} . Reorder the I_i so that $\ell_1 < \ell_2 < \dots < \ell_t$ and set $I'_i := I_i - \bigcap_{k=1}^t I_k$ and $j_i = \Lambda_{\ell_i} = \sum_{k=1}^{\ell_i} \lambda_k$. To begin, we prove lemmas which characterize (I_1, \dots, I_t) .

Lemma 5.21. Set $I_1 = J_{\ell_1}$ with sign ζ_{ℓ_1} and suppose there is some minimal T such that $I_T = J_{\ell_T}$ with sign $\zeta_{\ell_T} = -\zeta_{\ell_1}$. Then, for all $i \geq T$, we have $I_i = \zeta_{\ell_i}$ with sign $\zeta_{\ell_i} = -\zeta_{\ell_1}$.

Proof. Assume without loss that $\zeta_{\ell_1} = -1$ so $I_1 = \{j : \eta(\ell_1, j)\epsilon_j = -1\}$ and let T be the smallest value such that $I_T = \{j : \eta(\ell_T, j)\epsilon_j = 1\}$. Suppose there exists some $s > T$ such that $I_s = \{j : \eta(\ell_s, j)\epsilon_j = -1\}$. If $j \leq j_{T-1}$ or $j > j_s$, then $\eta(\ell_T, j) = \eta(\ell_s, j)$, so $j \in I_T \cup I_s$ so $[1, j_{T-1}] \cup [j_s + 1, n] \subseteq I_T \cup I_s$. Similarly, if $j \in [j_{T-1} + 1, j_s]$, then $\eta(\ell_{T-1}, j) = -\eta(\ell_s, j)$, so $j \in I_{T-1} \cup I_s$ so $[j_{T-1} + 1, j_s] \subseteq I_{T-1} \cup I_s$. Since $[1, j_{T-1}] \cup [j_s + 1, n] \cup [j_{T-1} + 1, j_s] = [1, n]$ and $a \leq n/2$, we must have that either $|I_T \cup I_s| \geq a$ or $|I_{T-1} \cup I_s| \geq a$. Then, by Lemma 5.13, C is not valid, a contradiction. Thus such an s cannot exist so $I_i = \{j : \eta(\ell_i, j)\epsilon_j = +1\}$ for all $i \geq T$. \square

Lemma 5.21 shows that the sign of (I_1, I_2, \dots, I_t) can switch at most once. This motivates the following definition.

Definition 5.22. The *transition point* of (I_1, \dots, I_t) is the smallest T such that $\zeta_{\ell_T} = -\zeta_{\ell_1}$. If I_1, I_2, \dots, I_t all have the same sign (so that no such T exists), then we set $T = 1$.

Lemma 5.23. Let T be the transition point of (I_1, \dots, I_t) . Then

$$\bigcup_{i=1}^t I'_i = I'_{T-1} \cup I'_T = [1, n] \setminus [j_{T-1} + 1, j_T], \quad (5-46)$$

$$I'_{T-1} \cap I'_T = \emptyset, \quad \text{and} \quad (5-47)$$

$$\bigcap_{i=1}^t I_i = I_{T-1} \cap I_T \subseteq [j_{T-1} + 1, j_T], \quad (5-48)$$

taking indices cyclically in $[1, t]$ and intervals cyclically in $[1, n]$ so that $I'_0 := I'_t$ and $I_0 := I_t$ and $[j_0 + 1, j_1] = [j_t + 1, n] \cup [1, j_1]$. Additionally, if $1 \leq i \leq t$ with $i \neq T - 1$, then

$$(I_i \cap [j_i + 1, j_{i+1}]) \cup (I_{i+1} \cap [j_i + 1, j_{i+1}]) = [j_i + 1, j_{i+1}] \quad \text{and} \quad (5-49)$$

$$(I_i \cap [j_i + 1, j_{i+1}]) \cap (I_{i+1} \cap [j_i + 1, j_{i+1}]) = \emptyset. \quad (5-50)$$

In other words, the restriction of I_i and I_{i+1} to the interval $[j_i + 1, j_{i+1}]$ forms a partition of the interval. If $i = t$ and $T \neq 1$, again taking indices and intervals cyclically, we set $I_{t+1} = I_1$ and $[j_t + 1, j_1] = [j_t + 1, n] \cup [1, j_1]$.

Proof. We consider indices and intervals cyclically in $[1, t]$ and $[1, n]$ respectively, as in Lemma 5.23.

For $j \in [j_{T-1} + 1, j_T]$, the value $\eta(\ell_i, j)\epsilon_j \zeta_{\ell_i}$ is independent of i since for any i, i' either $\zeta_{\ell_i}/\zeta_{\ell_{i'}}$ and $\eta(\ell_i, j)/\eta(\ell_{i'}, j)$ are both 1 or both -1 . So, for any $j \in [j_{T-1} + 1, j_T]$ either $j \in I_i$ for all i of $j \notin I_i$ for all i . So, $\bigcup_{i=1}^t I'_i \subset [1, n] \setminus [j_{T-1} + 1, j_T]$ and $\bigcap_{i=1}^t I_i \cap [j_{T-1} + 1, j_T] = I_{T-1} \cap I_T \cap [j_{T-1} + 1, j_T]$.

For $j \notin [j_{T-1} + 1, j_T]$, we have $\eta(\ell_{T-1}, j) = \eta(\ell_T, j)$ and so $\eta(\ell_{T-1}, j)\epsilon_j, \zeta_{\ell_{T-1}} = -\eta(\ell_T, j)\epsilon_j, \zeta_{\ell_T}$. So, every $j \notin [j_{T-1} + 1, j_T]$ belongs to exactly one of I'_{T-1} and I'_T . We conclude that $\bigcup_{i=1}^t I'_i = I'_{T-1} \cup I'_T \subset [1, n] \setminus [j_{T-1} + 1, j_T]$ and $I_{T-1} \cap I_T \subset [j_{T-1} + 1, j_T]$. So, $I_{T-1} \cap I_T \cap [j_{T-1} + 1, j_T] = I_{T-1} \cap I_T = \bigcap_{i=1}^t I_i$.

For $i \neq T - 1$, we have $\eta(\ell_i, j) = -\eta(\ell_{i+1}, j)$ if and only if $j \in [j_i + 1, j_{i+1}]$. Since $\zeta_{\ell_i} = \zeta_{\ell_{i+1}}$, this means $\eta(\ell_i, j)\epsilon_j\zeta_{\ell_i} = -\eta(\ell_{i+1}, j)\epsilon_j\zeta_{\ell_{i+1}}$ if and only if $j \in [j_i + 1, j_{i+1}]$. Hence, each $j \in [j_i + 1, j_{i+1}]$ is contained in exactly one of I_i and I_{i+1} , as desired. \square

Definition 5.24. For each $1 \leq i \leq t$, set

$$r_i := |I_i \cap [j_i + 1, j_{i+1}]| \quad \text{and} \quad s_i := |I_{i+1} \cap [j_i + 1, j_{i+1}]|. \tag{5-51}$$

We call the ordered tuple $(T, r_1, s_1, \dots, r_t, s_t)$ the *structure* of (I_1, \dots, I_t) in S where T is the transition point of (I_1, \dots, I_t) . If $(T, r_1, s_1, \dots, r_t, s_t)$ is a structure for some $(I_1, \dots, I_t) \in C$, we call it a *valid structure* for C .

By Lemma 5.23, $r_{T-1} = s_{T-1} = |\bigcap_{k=1}^t I_k|$. Lemma 5.23 also shows that when $i \neq T - 1$, $r_i + s_i = |[j_i + 1, j_{i+1}]| = j_{i+1} - j_i = \lambda_{\ell_{i+1}} + \dots + \lambda_{\ell_{i+1}}$. The following lemma shows that the two tuples with the same structure are in the same t -class.

Lemma 5.25. *Let C be a valid t -class and let $(I_1, \dots, I_t) \in C$ such that $(I_1, \dots, I_t) \in I(S)$ for some s.o.p. S . Let (J_1, \dots, J_t) be another tuple such that $(J_1, \dots, J_t) \in I(P)$ for some s.o.p. P . If the structure of (I_1, \dots, I_t) in S is the same as the structure of (J_1, \dots, J_t) in P , then $(J_1, \dots, J_t) \in C$.*

Proof. We first set notation. Set $S = (m, \lambda_1, \dots, \lambda_m, \epsilon_1, \dots, \epsilon_n)$ and $P = (m, \lambda'_1, \dots, \lambda'_m, \epsilon'_1, \dots, \epsilon'_n)$. Set $\ell_1 < \dots < \ell_t$ and $\ell'_1 < \dots < \ell'_t$ such that $I_i = \{j : \eta_S(\ell_i, j) = \zeta_{\ell_i}\}$ and $J_i = \{j : \eta_P(\ell'_i, j) = \zeta_{\ell'_i}\}$. Lastly, define $j_i = \sum_{k=1}^{\ell_i} \lambda_k$ and $j'_i = \sum_{k=1}^{\ell'_i} \lambda'_k$.

Without loss of generality, we may assume $\zeta_{\ell_i} = \zeta_{\ell'_i}$, or else we may replace each ϵ_i with $-\epsilon_i$. Since (I_1, \dots, I_t) and (J_1, \dots, J_t) have the same structure, for each i , we have

$$|I_i \cap [j_i + 1, j_{i+1}]| = |J_i \cap [j'_i + 1, j'_{i+1}]| \quad \text{and} \quad |I_{i+1} \cap [j_i + 1, j_{i+1}]| = |J_{i+1} \cap [j'_i + 1, j'_{i+1}]|. \tag{5-52}$$

Let $\tau \in S_n$ be the permutation that maps the k -th smallest element of $|I_i \cap [j_i + 1, j_{i+1}]|$ to the k -th smallest element of $|J_i \cap [j'_i + 1, j'_{i+1}]|$ and the k -th smallest element of $|I_{i+1} \cap [j_i + 1, j_{i+1}]|$ to the k -th smallest element of $|J_{i+1} \cap [j'_i + 1, j'_{i+1}]|$.

Since $\tau([j_i + 1, j_{i+1}]) = [j'_i + 1, j'_{i+1}]$, for all $i \in [1, t]$ and $j \in [1, n]$ we have $\eta(\ell_i, j) = \eta(\ell'_i, \tau(j))$. Since (I_1, \dots, I_t) and (J_1, \dots, J_t) have the same transition value T and we assumed $\zeta_{\ell_1} = \zeta_{\ell'_1}$, we have $\zeta_{\ell_i} = \zeta_{\ell'_i}$ for all i . Moreover, for $j \in [j_i + 1, j_{i+1}]$ we have $\eta(\ell_i, j)\epsilon_j\zeta_{\ell_i} = \eta(\ell'_i, \tau(j))\epsilon'_{\tau(j)}\zeta_{\ell'_i}$ so that $\epsilon_j = \epsilon'_{\tau(j)}$. But this is true for all i , so in fact $\epsilon_j = \epsilon'_{\tau(j)}$ for all $j \in [1, n]$. So, for all $i \in [1, t]$ and $j \in [1, n]$ we have $\eta(\ell_i, j)\epsilon_j\zeta_{\ell_i} = \eta(\ell'_i, \tau(j))\epsilon'_{\tau(j)}\zeta_{\ell'_i}$. It follows that $\tau(I_1, \dots, I_t) = (J_1, \dots, J_t)$ so $(J_1, \dots, J_t) \in C$. \square

Lemma 5.25 shows that if a structure is valid for C , then all tuples with that structure are in C . Thus in order to calculate $\sum T(S, C) A(S)$, we can first sum over all valid structures for C and then count tuples and s.o.p.s with that structure. All that remains is to determine when $(I_1, \dots, I_t) \in I(S)$.

Lemma 5.26. *Suppose $I_1, \dots, I_t \in J(S)$. Then, $I_1, \dots, I_t \in I(S)$ if and only if for each $1 \leq i \leq t$, $[j_i - \lambda_{\ell_i} + 1, j_i] \not\subseteq I_i$ and $[j_i + 1, j_i + \lambda_{\ell_{i+1}}] \not\subseteq I_i$.*

Proof. Note that $I_i \in J(S)$ implies $\#I_i \leq a - 1$. So, this is an immediate corollary of Lemma 5.15 which says $I_i \in I(S)$ if and only if $[j_i - \lambda_{\ell_i} + 1, j_i] \not\subseteq I_i$ and $[j_i + 1, j_i + \lambda_{\ell_i+1}] \not\subseteq I_i$. \square

Lemma 5.27. *Let C be a valid t -class with $t \geq 2$. Then*

$$\sum_{\text{s.o.p.'s } S} T(S, C) A(S) \tag{5-53}$$

is a sum of terms of the form

$$\sum_{d=1}^f \sum_{\substack{\mu_1 + \dots + \mu_d = f \\ \mu_i \geq 1}} \frac{(-1)^d}{\mu_1! \dots \mu_d!} H(f, g, \mu_1, \mu_d), \tag{5-54}$$

for some f and g , where

$$H(f, g, \mu_1, \mu_d) := \binom{f}{g} - \binom{f - \mu_1}{g - \mu_1} - \binom{f - \mu_d}{g} + \binom{f - \mu_1 - \mu_d}{g - \mu_1}. \tag{5-55}$$

Proof. Let C be a valid t -class. By Lemma 5.25, when summing over all s.o.p.s, we can first sum over all valid structures, and then over all s.o.p.s and tuples with that structure. To do this, we can sum over all m , then over all possible values of ℓ_1, \dots, ℓ_t , then over all $\lambda_1, \dots, \lambda_m$ such that $\lambda_1 + \dots + \lambda_m = n$ and $\lambda_{\ell_i+1} + \dots + \lambda_{\ell_{i+1}} = r_i + s_i$ for each $i \neq T - 1$. Now we use Lemma 5.26 to determine the summand. We can pick the elements of $\bigcap_{k=1}^t I_k$, which by Lemma 5.23 is a subset of $[j_{T-1} + 1, j_T]$, in $G(j_T - j_{T-1}, r_T, \lambda_{\ell_{T-1}}, \lambda_{\ell_T})$ ways, where G is defined as in (5-35). Next, we choose the r_i elements of I_i contained in the interval $[j_i + 1, j_{i+1}]$ in $H(r_i + s_i, r_i, \lambda_{\ell_i+1}, \lambda_{\ell_{i+1}})$ ways. Then, there are two possible choices for the sign ζ_{ℓ_i} of I_i and then the signs for the rest of the I_i 's follow because the point of transition T is fixed. The choice of ζ_{ℓ_i} and each r_i -element set $[j_i + 1, j_{i+1}] \cap I_i$ determines all ϵ_j , so they determine exactly the same data as the I_i .

Lastly, we multiply by $A(S)$. We have

$$\begin{aligned} \sum_{\text{s.o.p.'s } S} T(S, C) \cdot A(S) = & \sum_{\substack{(T, r_1, s_1, \dots, r_t, s_t) \\ \text{a valid structure for } C}} \sum_{m=1}^n \sum_{1 \leq \ell_1 < \dots < \ell_t \leq m} \sum_{\substack{\lambda_1 + \dots + \lambda_m = n \\ \lambda_{\ell_i+1} + \dots + \lambda_{\ell_{i+1}} = r_i + s_i \text{ for each } i \neq T \\ \lambda_i \geq 1}} \\ & \times 2G(u, v, \lambda_{\ell_1}, \lambda_{\ell_{i+1}}) \prod_{\substack{1 \leq i \leq t \\ i \neq T-1}} H(r_i + s_i, r_i, \lambda_{\ell_i+1}, \lambda_{\ell_{i+1}}) \frac{(-1)^{m+1}}{m} \frac{n!}{\lambda_1! \dots \lambda_m!}. \end{aligned}$$

For each structure, we can fix some $i \neq T - 1$, which exists since $t \geq 2$, to see that this is a sum of terms of the form

$$\sum_{d=1}^{r_i+s_i} \sum_{\substack{\mu_1 + \dots + \mu_d = r_i + s_i \\ \mu_i \geq 1}} \frac{(-1)^d}{\mu_1! \dots \mu_d!} H(r_i + s_i, r_i, \mu_1, \mu_d). \tag{5-56} \quad \square$$

We finish the calculation with the following combinatorial lemma, proven in Appendix C.2.

Lemma 5.28. Fix f, g and let $H(f, g, \mu_1, \mu_d)$ be as in (5-55). Then

$$\sum_{d=1}^f \sum_{\substack{\mu_1 + \dots + \mu_d = f \\ \mu_i \geq 1}} \frac{(-1)^d}{\mu_1! \dots \mu_d!} H(f, g, \mu_1, \mu_d) = 0. \tag{5-56}$$

Combining Lemmas 5.27 and 5.28 completes the proof of Lemma 5.20. □

5.4. Computing the integral piece. In this section we complete the proof of Proposition 5.2 by calculating the integral $\int C \, dy$ appearing in (5-30). Applying Lemmas 5.20 and 5.19 to 5.12 gives

$$Q_n(\phi) = 2^{n-1} (-1)^n \sum_{\ell=0}^{a-1} (-1)^\ell \binom{n}{\ell} \int_0^\infty \dots \int_0^\infty \hat{\phi}(y_1) \dots \hat{\phi}(y_n) \chi_{\{n-\ell+1, \dots, n\}} dy_1 \dots dy_n. \tag{5-57}$$

Next we define

$$\xi_\ell(\phi) := \int_0^\infty \dots \int_0^\infty \hat{\phi}(y_1) \dots \hat{\phi}(y_n) \chi_{\{y_1 + \dots + y_{n-\ell} - y_{n-\ell+1} - \dots - y_n > 1\}} dy_1 \dots dy_n, \tag{5-58}$$

$$\bar{\xi}_\ell(\phi) := \int_{-\infty}^\infty \dots \int_{-\infty}^\infty \hat{\phi}(y_1) \dots \hat{\phi}(y_n) \chi_{\{|y_1 + \dots + y_{n-\ell} - |y_{n-\ell+1}| - \dots - |y_n| > 1\}} dy_1 \dots dy_n. \tag{5-59}$$

Equation (5-57) equals

$$Q_n(\phi) = 2^{n-1} (-1)^n \sum_{\ell=0}^{a-1} (-1)^\ell \binom{n}{\ell} \xi_\ell(\phi). \tag{5-60}$$

We express $Q_n(\phi)$ in terms of $\bar{\xi}_\ell(\phi)$ with the following lemma.

Lemma 5.29. Let $\phi \in \mathcal{S}_{ec}(\mathbb{R})$ with $\text{supp}(\hat{\phi}) \subseteq [-\frac{1}{n-a}, \frac{1}{n-a}]$. Then

$$Q_n(\phi) = 2^{n-2} (-1)^n \sum_{t=0}^{a-1} (-1)^t \binom{n}{t} \bar{\xi}_t(\phi). \tag{5-61}$$

Proof. Given $\text{supp}(\hat{\phi}) \subseteq [-\frac{1}{n-a}, \frac{1}{n-a}]$ and $t \leq a-1$, if $|y_1 + \dots + y_{n-t}| - |y_{n-t+1}| - \dots - |y_n| > 1$, then either at most $i \leq a-1-t$ of the y_j 's in the first absolute value are nonnegative and the rest are negative or at most $i \leq a-1-t$ of the y_j 's in the first absolute value are nonpositive or zero and the rest are positive. Moreover, the sign of $y_1 + \dots + y_{n-t}$ matches the second group. There are $\binom{n-t}{i}$ ways to choose these indices and we introduce a factor of 2 from choosing the sign of $y_1 + \dots + y_{n-t}$. Lastly, since $\hat{\phi}$ is even, we multiply by a factor of 2^t to account for changing the limits of integration over y_{n-t+1}, \dots, y_n . Thus we have

$$\begin{aligned} \bar{\xi}_t(\phi) &= 2^{t+1} \int_0^\infty \dots \int_0^\infty \hat{\phi}(y_1) \dots \hat{\phi}(y_n) \left(\sum_{i=0}^{a-1-t} \binom{n-t}{i} \chi_{\{y_1 + \dots + y_{n-i-t} - y_{n-i-t+1} - \dots - y_n > 1\}} \right) dy_1 \dots dy_n \\ &= 2^{t+1} \sum_{i=0}^{a-1-t} \binom{n-t}{i} \xi_{i+t}(\phi). \end{aligned} \tag{5-62}$$

Applying the identity

$$\sum_{t=0}^{\ell} (-2)^t \binom{n}{t} \binom{n-t}{\ell-t} = \binom{n}{\ell} \sum_{t=0}^{\ell} (-2)^t \binom{\ell}{t} = \binom{n}{\ell} (1-2)^{\ell} = \binom{n}{\ell} (-1)^{\ell} \tag{5-63}$$

to (5-60) gives

$$Q_n(\phi) = 2^{n-1} (-1)^n \sum_{\ell=0}^{a-1} \xi_{\ell}(\phi) \sum_{t=0}^{\ell} (-2)^t \binom{n}{t} \binom{n-t}{\ell-t}. \tag{5-64}$$

Switching the order of summation and setting $i = \ell - t$ gives

$$Q_n(\phi) = 2^{n-2} (-1)^n \sum_{t=0}^{a-1} (-1)^t \binom{n}{t} \left(2^{t+1} \sum_{i=0}^{a-1-t} \binom{n-t}{i} \xi_{i+t}(\phi) \right). \tag{5-65}$$

Applying (5-62) gives the desired result. □

We complete the evaluation of $Q_n(\phi)$ by computing $\bar{\xi}_i(\phi)$.

Lemma 5.30. *Let $\phi \in \mathcal{S}_{ec}(\mathbb{R})$ with $\text{supp}(\hat{\phi}) \subseteq [-\frac{1}{n-a}, \frac{1}{n-a}]$. Then we have*

$$\begin{aligned} \bar{\xi}_{\ell}(\phi) &= \phi^n(0) \\ &\quad - 2 \int_{-\infty}^{\infty} \cdots \int_{-\infty}^{\infty} \hat{\phi}(x_{\ell+1}) \cdots \hat{\phi}(x_2) \int_{-\infty}^{\infty} \phi^{n-\ell}(x_1) \frac{\sin(2\pi x_1(1 + |x_2| + \cdots + |x_{\ell+1}|))}{2\pi x_1} dx_1 \cdots dx_{\ell+1} \end{aligned} \tag{5-66}$$

for $\ell \leq a - 1$.

Proof. We apply to (5-59) the change of variables given by

$$\begin{aligned} x_1 &= y_1 & y_1 &= x_1 \\ x_2 &= y_1 + y_2 & y_2 &= x_2 - x_1 \\ &\vdots & &\vdots \\ x_{n-\ell} &= \sum_{j=1}^{n-\ell} y_j & y_{n-\ell} &= x_{n-\ell} - x_{n-\ell-1} \\ x_{n-\ell+1} &= y_{n-\ell+1} & y_{n-\ell+1} &= x_{n-\ell+1} \\ &\vdots & &\vdots \\ x_n &= y_n & y_n &= x_n, \end{aligned} \tag{5-67}$$

obtaining

$$\begin{aligned} \bar{\xi}_{\ell}(\phi) &= \int_{-\infty}^{\infty} \cdots \int_{-\infty}^{\infty} \hat{\phi}(x_1) \hat{\phi}(x_2 - x_1) \cdots \hat{\phi}(x_{n-\ell} - x_{n-\ell-1}) \\ &\quad \times \hat{\phi}(x_{n-\ell+1}) \cdots \hat{\phi}(x_n) \chi_{\{|x_{n-\ell}| - (|x_{n-\ell+1}| + \cdots + |x_n|) > 1\}} dx_1 \cdots dx_n. \end{aligned} \tag{5-68}$$

Repeatedly applying the identity $\int_{-\infty}^{\infty} \hat{f}(v) \hat{g}(u - v) dv = \widehat{fg}(u)$ (which arises from the convolution theorem) to (5-68) gives

$$\bar{\xi}_{\ell}(\phi) = \int_{-\infty}^{\infty} \cdots \int_{-\infty}^{\infty} \widehat{\phi^{n-\ell}}(x_{n-\ell}) \hat{\phi}(x_{n-\ell+1}) \cdots \hat{\phi}(x_n) \chi_{\{|x_{n-\ell}| - (|x_{n-\ell+1}| + \cdots + |x_n|) > 1\}} dx_{n-\ell} \cdots dx_n. \tag{5-69}$$

We rename $x_{n-\ell}$ to x_1 , $x_{n-\ell+1}$ to x_2 , and so on until x_n to $x_{\ell+1}$. This and the identity

$$\chi_{\{|x_1|-(|x_2|+\dots+|x_{\ell+1}|)>1\}} = 1 - \chi_{\{|x_1|\leq 1+|x_2|+\dots+|x_{\ell+1}|\}} \quad (5-70)$$

gives

$$\bar{\xi}_\ell(\phi) = \int_{-\infty}^{\infty} \dots \int_{-\infty}^{\infty} \widehat{\phi}^{n-\ell}(x_1) \hat{\phi}(x_2) \dots \hat{\phi}(x_{\ell+1}) (1 - \chi_{\{|x_1|\leq 1+|x_2|+\dots+|x_{\ell+1}|\}}) dx_1 \dots dx_{\ell+1}. \quad (5-71)$$

Distributing and using the identity $\phi(0) = \int_{-\infty}^{\infty} \hat{\phi}(x) dx$, we have

$$\bar{\xi}_\ell(\phi) = \phi^n(0) - \int_{-\infty}^{\infty} \dots \int_{-\infty}^{\infty} \widehat{\phi}^{n-\ell}(x_1) \hat{\phi}(x_2) \dots \hat{\phi}(x_{\ell+1}) \chi_{\{|x_1|\leq 1+|x_2|+\dots+|x_{\ell+1}|\}} dx_1 \dots dx_{\ell+1}. \quad (5-72)$$

Fix $x_2, \dots, x_{\ell+1}$ and set $S_\ell(x_1) = \sin(2\pi x_1(1 + |x_2| + \dots + |x_{\ell+1}|)) / (2\pi x_1)$. We have the identity

$$\chi_{\{|x_1|\leq 1+|x_2|+\dots+|x_{\ell+1}|\}}(x_1) = 2\hat{S}_\ell(x_1), \quad (5-73)$$

which follows from the Fourier pair

$$\frac{\sin(2\pi Ax)}{2\pi x} = \int_{-\infty}^{\infty} \frac{1}{2} \chi_{\{|u|\leq A\}} e^{2\pi i x u} du. \quad (5-74)$$

Thus Plancherel's theorem gives us (5-66), as desired. \square

Applying Lemma 5.30 to (5-61) and comparing with (1-13) completes the proof of Proposition 5.2.

Appendix A. Proof of Lemma 3.1

Proof. We begin by imposing a distinctness condition on the sum over primes in (2-40). Fix a partition $\vec{n} = (n_1, \dots, n_\ell)$ of n , where $1 \leq n_1 \leq \dots \leq n_\ell$ and $n_1 + \dots + n_\ell = n$. We want to write $p_1 \dots p_n = q_1^{n_1} \dots q_\ell^{n_\ell}$. If the parts of our partition are distinguishable, then by the multinomial theorem there are

$$\binom{n}{n_1, \dots, n_\ell} = \frac{n!}{(n_1!) \dots (n_\ell!)} \quad (A-1)$$

ways to do this. Let $r_x(\vec{n})$ denote the number of values of i for which $n_i = x$ in the partition \vec{n} . Because the parts of our partition which are the same size are *indistinguishable*, the number of ways to write $p_1 \dots p_n = q_1^{n_1} \dots q_\ell^{n_\ell}$ is

$$\sigma(\vec{n}) := \binom{n}{n_1, \dots, n_\ell} \prod_{j=1}^n \frac{1}{r_j!}. \quad (A-2)$$

Thus

$$\frac{S_2^{(n)}(N)}{i^k \sqrt{N}} = \sum_{1 \leq \ell \leq n} \sum_{\substack{\vec{n} := (n_1, \dots, n_\ell) \\ 1 \leq n_1 \leq \dots \leq n_\ell \\ n_1 + \dots + n_\ell = n}} \sigma(\vec{n}) \sum_{\substack{q_1 \nmid N, \dots, q_\ell \nmid N \\ q_j \text{ distinct}}} \prod_{j=1}^{\ell} \left(\hat{\phi} \left(\frac{\log q_j}{\log R} \right)^{n_j} \left(\frac{2 \log q_j}{\sqrt{q_j} \log R} \right)^{n_j} \right) \times \langle \lambda_f(N) \lambda_f(q_1)^{n_1} \dots \lambda_f(q_\ell)^{n_\ell} \rangle_*. \quad (A-3)$$

Now that we have a distinct sum over primes, we want to apply the multiplicative properties of Fourier coefficients given in Lemma 2.7. We do some careful bookkeeping in the process. We will sum over tuples $\vec{m} = (m_1, \dots, m_\ell)$ which are *admissible* to a given partition \vec{n} . This means that $m_i \leq n_i$, $n_i \equiv m_i \pmod{2}$,

and if $n_i = n_j$ with $i < j$ then $m_i \leq m_j$. This last condition means that we order the m_i s for each fixed value of n_i .

For some fixed \vec{n}, \vec{m} with \vec{m} admissible to \vec{n} , let $s_{x,y}(\vec{n}, \vec{m})$ be the number of values of i for which $(n_i, m_i) = (x, y)$. For each value of x , the number of ways to order the indices i for which $n_i = x$ is

$$\frac{r_x(\vec{n})!}{\prod_{j=1}^x s_{x,j}(\vec{n}, \vec{m})!}. \tag{A-4}$$

Define the auxiliary function

$$\tau(\vec{n}, \vec{m}) := \prod_{i=1}^n \frac{r_i(\vec{n})!}{\prod_{j=1}^i s_{i,j}(\vec{n}, \vec{m})!} \tag{A-5}$$

and let $t_{n,m}$ be the coefficient of $\lambda_f(p^m)$ in the expansion of $\lambda_f(p)^n$ (see (2-29)), so that

$$\lambda_f(p)^n = \sum_{m=0}^n t_{n,m} \lambda_f(p^m). \tag{A-6}$$

Note that $t_{n,n} = 1$ for all n . Expanding the Fourier coefficients with (2-29) and using (2-27), we have

$$\begin{aligned} \frac{S_2^{(n)}(N)}{i^k \sqrt{N}} &= \sum_{1 \leq \ell \leq n} \sum_{\substack{\vec{n} := (n_1, \dots, n_\ell) \\ 1 \leq n_1 \leq \dots \leq n_\ell \\ n_1 + \dots + n_\ell = n}} \sum_{\substack{\vec{m} \text{ admissible to } \vec{n} \\ q_1 \nmid N, \dots, q_\ell \nmid N \\ q_j \text{ distinct}}} \sigma(\vec{n}) \tau(\vec{n}, \vec{m}) \sum_{q_1 \nmid N, \dots, q_\ell \nmid N} \\ &\times \prod_{j=1}^{\ell} \left(t_{n_j, m_j} \hat{\phi} \left(\frac{\log q_j}{\log R} \right)^{n_j} \left(\frac{2 \log q_j}{\sqrt{q_j} \log R} \right)^{n_j} \right) \langle \lambda_f(N q_1^{m_1} \cdots q_\ell^{m_\ell}) \rangle_*. \end{aligned} \tag{A-7}$$

We want to separate (\vec{n}, \vec{m}) into a part with $n_i = m_i$ and a part with $n_i < m_i$. Let

$$\vec{n}^b = (n_1^b, n_2^b, \dots, n_\omega^b)$$

denote the subpartition of \vec{n} with $n_i > m_i$, and

$$\vec{n}^\sharp = (n_1^\sharp, n_2^\sharp, \dots, n_{\ell-\omega}^\sharp)$$

denote the subpartition of \vec{n} with $n_i = m_i$. The analogous notation holds for \vec{m} . Set $\sum n_i^b = n'$, so that $\sum n_i^\sharp = n - n'$. We have

$$\begin{aligned} \sigma(\vec{n}) \tau(\vec{n}, \vec{m}) &= \frac{n!}{n_1! \cdots n_\ell!} \prod_{i=1}^n \prod_{j=1}^i \frac{1}{s_{i,j}(\vec{n}, \vec{m})!} \\ &= \frac{n!}{n'!(n-n')!} \times \left(\frac{n!}{n_1^b! \cdots n_\omega^b!} \prod_{i=1}^n \prod_{j=1}^{i-1} \frac{1}{s_{i,j}(\vec{n}, \vec{m})!} \right) \times \left(\frac{(n-n')!}{n_1^\sharp! \cdots n_{\ell-\omega}^\sharp!} \prod_{i=1}^n \frac{1}{s_{i,i}(\vec{n}, \vec{m})!} \right) \\ &= \binom{n}{n'} \times \left(\frac{n!}{n_1^b! \cdots n_\omega^b!} \prod_{i=1}^{n'} \prod_{j=1}^i \frac{1}{s_{i,j}(\vec{n}^b, \vec{m}^b)!} \right) \times \left(\frac{(n-n')!}{n_1^\sharp! \cdots n_{\ell-\omega}^\sharp!} \prod_{i=1}^{n-n'} \frac{1}{r_i(\vec{n}^\sharp)!} \right) \\ &= \binom{n}{n'} \sigma(\vec{n}^b) \tau(\vec{n}^b, \vec{m}^b) \sigma(\vec{n}^\sharp). \end{aligned} \tag{A-8}$$

Applying this identity to (A-7) and using $t_{n,n} = 1$ and $n_i^\sharp = m_i^\sharp$ gives

$$\begin{aligned}
 \frac{S_2^{(n)}(N)}{i^k \sqrt{N}} &= \sum_{0 \leq \omega \leq n} \sum_{0 \leq n' \leq n} \binom{n}{n'} \sum_{\substack{\vec{n}^b := (n_1^b, \dots, n_\omega^b) \\ 1 \leq n_1^b \leq \dots \leq n_\omega^b \\ n_1^b + \dots + n_\omega^b = n'}} \sum_{\substack{\vec{m}^b \text{ admissible to } \vec{n}^b \\ n_j^b < m_j^b}} \sigma(\vec{n}^b) \tau(\vec{n}^b, \vec{m}^b) t_{n_1^b, m_1^b} \cdots t_{n_\omega^b, m_\omega^b} \\
 &\times \sum_{\substack{q_1 \nmid N, \dots, q_\omega \nmid N \\ q_j \text{ distinct}}} \prod_{j=1}^{\omega} \left(\hat{\phi} \left(\frac{\log q_j}{\log R} \right)^{n_j^b} \left(\frac{2 \log q_j}{\sqrt{q_j} \log R} \right)^{n_j^b} \right) \\
 &\times \sum_{0 \leq \ell - \omega \leq n - n'} \sum_{\substack{\vec{n}^\sharp := (n_1^\sharp, \dots, n_{\ell - \omega}^\sharp) \\ 1 \leq n_1^\sharp \leq \dots \leq n_{\ell - \omega}^\sharp \\ n_1^\sharp + \dots + n_{\ell - \omega}^\sharp = n - n'}} \sigma(\vec{n}^\sharp) \sum_{\substack{p_1 \nmid N, \dots, p_{\ell - \omega} \nmid N \\ p_i \text{ distinct} \\ p_i \neq q_j}} \prod_{i=1}^{\ell - \omega} \left(\hat{\phi} \left(\frac{\log p_i}{\log R} \right)^{n_i^\sharp} \left(\frac{2 \log p_i}{\sqrt{p_i} \log R} \right)^{n_i^\sharp} \right) \\
 &\times \langle \lambda_f(N q_1^{m_1^b} \cdots q_\omega^{m_\omega^b} p_1^{n_1^\sharp} \cdots p_{\ell - \omega}^{n_{\ell - \omega}^\sharp}) \rangle_*. \tag{A-9}
 \end{aligned}$$

Arguing as in (A-3), we have

$$\begin{aligned}
 &\sum_{0 \leq \ell - \omega \leq n - n'} \sum_{\substack{\vec{n}^\sharp := (n_1^\sharp, \dots, n_{\ell - \omega}^\sharp) \\ 1 \leq n_1^\sharp \leq \dots \leq n_{\ell - \omega}^\sharp \\ n_1^\sharp + \dots + n_{\ell - \omega}^\sharp = n - n'}} \sigma(\vec{n}^\sharp) \sum_{\substack{p_1 \nmid N, \dots, p_{\ell - \omega} \nmid N \\ p_i \text{ distinct} \\ p_i \neq q_j}} \prod_{i=1}^{\ell - \omega} \left(\hat{\phi} \left(\frac{\log p_i}{\log R} \right)^{n_i^\sharp} \left(\frac{2 \log p_i}{\sqrt{p_i} \log R} \right)^{n_i^\sharp} \right) \\
 &\times \langle \lambda_f(N q_1^{m_1^b} \cdots q_\omega^{m_\omega^b} p_1^{n_1^\sharp} \cdots p_{\ell - \omega}^{n_{\ell - \omega}^\sharp}) \rangle_* \\
 &= \sum_{\substack{p_1 \nmid N, \dots, p_{n - n'} \nmid N \\ p_i \neq q_j}} \prod_{i=1}^{n - n'} \left(\hat{\phi} \left(\frac{\log p_i}{\log R} \right) \frac{2 \log p_i}{\sqrt{p_i} \log R} \right) \langle \lambda_f(N q_1^{m_1^b} \cdots q_\omega^{m_\omega^b} p_1 \cdots p_{n - n'}) \rangle_*. \tag{A-10}
 \end{aligned}$$

We apply (A-10) to (A-9). In doing so, we remove the condition that $1 \leq n_1^b \cdots \leq n_\omega^b$ as we are no longer concerned with the ordering of our partition. We also relax the condition that \vec{m}^b is admissible to \vec{n}^b . We also suppress the b notation (as there are no more \sharp s). Lastly, the first line of (A-9) is purely combinatorial, so we combine the combinatorial factors appearing in (A-9) into the coefficients $C''_{\vec{n}, \vec{m}}$ appearing below.

Thus

$$S_2^{(n)}(N) = \sum_{0 \leq \omega \leq n} \sum_{0 \leq n' \leq n} \sum_{\substack{\vec{n} := (n_1, \dots, n_\omega) \\ n_j > 1 \\ n_1 + \dots + n_\omega = n'}} \sum_{\substack{\vec{m} := (m_1, \dots, m_\omega) \\ m_j \equiv n_j \pmod{2} \\ 0 \leq m_j < n_j}} C''_{\vec{n}, \vec{m}} E''(\vec{n}, \vec{m}) \tag{A-11}$$

where each $C''_{\vec{n}, \vec{m}}$ is some explicit constant only dependent on \vec{n}, \vec{m} and

$$E''(\vec{n}, \vec{m}) := \sum_{\substack{q_1 \uparrow N, \dots, q_\omega \uparrow N \\ q_j \text{ distinct}}} \prod_{j=1}^{\omega} \left(\hat{\phi} \left(\frac{\log q_j}{\log R} \right)^{n_j} \left(\frac{2 \log q_j}{\sqrt{q_j} \log R} \right)^{n_j} \right) \tag{A-12}$$

$$\times \sum_{\substack{p_1 \uparrow N, \dots, p_{n-n'} \uparrow N \\ p_i \neq q_j}} \prod_{i=1}^{n-n'} \left(\hat{\phi} \left(\frac{\log p_i}{\log R} \right) \frac{2 \log p_i}{\sqrt{p_i} \log R} \right) \langle \lambda_f(N q_1^{m_1} \cdots q_\omega^{m_\omega} p_1 \cdots p_{n-n'}) \rangle_*$$

We want to remove the condition $p_i \neq q_j$ from the sum over $p_1, \dots, p_{n-n'}$ in (A-12). We apply an inclusion-exclusion process, subtracting off the terms where some $p_i = q_j$. We have

$$E''(\vec{n}, \vec{m}) = E'(\vec{n}, \vec{m}) - \sum_{a=1}^{n-n'} \sum_{\substack{\vec{a}=(a_1, \dots, a_\omega) \\ a_j \geq 0 \\ a_1 + \dots + a_\omega = a}} C''_{\vec{n}, \vec{m}, \vec{a}} E''(\vec{n} + \vec{a}, \vec{m} + \vec{a}), \tag{A-13}$$

where each $C''_{\vec{n}, \vec{m}, \vec{a}}$ is some explicit constant only dependent on $\vec{n}, \vec{m}, \vec{a}$ and

$$E'(\vec{n}, \vec{m}) := \sum_{\substack{q_1 \uparrow N, \dots, q_\omega \uparrow N \\ q_j \text{ distinct}}} \prod_{j=1}^{\omega} \left(\hat{\phi} \left(\frac{\log q_j}{\log R} \right)^{n_j} \left(\frac{2 \log q_j}{\sqrt{q_j} \log R} \right)^{n_j} \right) \tag{A-14}$$

$$\times \sum_{p_1 \uparrow N, \dots, p_{n-n'} \uparrow N} \prod_{i=1}^{n-n'} \left(\hat{\phi} \left(\frac{\log p_i}{\log R} \right) \frac{2 \log p_i}{\sqrt{p_i} \log R} \right) \langle \lambda_f(N q_1^{m_1} \cdots q_\omega^{m_\omega} p_1 \cdots p_{n-n'}) \rangle_*$$

The addition of vectors is taken componentwise, so that $\vec{n} + \vec{a} = (n_1 + a_1, \dots, n_\omega + a_\omega)$. We apply the inclusion-exclusion identity (A-13) to each term $E''(\vec{n} + \vec{a}, \vec{m} + \vec{a})$ appearing on the right side of (A-13). Repeating this process $n - n'$ times, we find that

$$E''(\vec{n}, \vec{m}) = \sum_{a=0}^{n-n'} \sum_{\substack{\vec{a}=(a_1, \dots, a_\omega) \\ a_j \geq 0 \\ a_1 + \dots + a_\omega = a}} C'_{\vec{n}, \vec{m}, \vec{a}} E'(\vec{n} + \vec{a}, \vec{m} + \vec{a}), \tag{A-15}$$

where each $C'_{\vec{n}, \vec{m}, \vec{a}}$ is some explicit constant only dependent on $\vec{n}, \vec{m}, \vec{a}$. In particular, $C'_{\vec{n}, \vec{m}, \vec{0}} = 1$. Applying (A-15) to (A-11) gives

$$S_2^{(n)}(N) = \sum_{0 \leq \omega \leq n} \sum_{0 \leq n' \leq n} \sum_{\substack{\vec{n} := (n_1, \dots, n_\omega) \\ n_j > 1 \\ n_1 + \dots + n_\omega = n'}} \sum_{\substack{\vec{m} := (m_1, \dots, m_\omega) \\ m_j \equiv n_j \pmod{2} \\ 0 \leq m_j < n_j}} C'_{\vec{n}, \vec{m}} E'(\vec{n}, \vec{m}), \tag{A-16}$$

where each $C'_{\vec{n}, \vec{m}}$ is some explicit constant only dependent on \vec{n}, \vec{m} .

We want to remove the distinctness condition from the sum over q_1, \dots, q_ω in (A-14). We again apply inclusion-exclusion, subtracting off terms where some of the q_j 's are equal. First, we define a *partition of a set* S to be a set $\vec{\pi} = \{\pi_1, \dots, \pi_\alpha\}$ where each $\pi_i \subset S$ is a nonempty subset of S , $\pi_i \cap \pi_j = \emptyset$ for every

$i \neq j$, and $\bigcap_{1 \leq i \leq \alpha} \pi_i = S$. We have

$$E'(\vec{n}, \vec{m}) = E(\vec{n}, \vec{m}) - \sum_{1 \leq \alpha \leq \omega-1} \sum_{\substack{\vec{\pi} \text{ partitions } \{1, \dots, \omega\} \\ \vec{\pi} = \{\pi_1, \dots, \pi_\alpha\}}} E'(\vec{x}, \vec{y}), \tag{A-17}$$

where $E(\vec{n}, \vec{m})$ is as in (3-2) and for each $\vec{\pi}$ we have

$$\begin{aligned} \vec{x} &= (x_1, \dots, x_\alpha), & x_i &= \sum_{j \in \pi_i} n_j & \text{and} \\ \vec{y} &= (y_1, \dots, y_\alpha), & y_i &= \sum_{j \in \pi_i} m_j. \end{aligned} \tag{A-18}$$

We apply the inclusion-exclusion identity (A-17) to each term $E'(\vec{x}, \vec{y})$ appearing on the right hand side of (A-17). Repeating this process ω times, we have

$$E'(\vec{n}, \vec{m}) = \sum_{1 \leq \alpha \leq \omega} \sum_{\substack{\vec{\pi} \text{ partitions } \{1, \dots, \omega\} \\ \vec{\pi} = \{\pi_1, \dots, \pi_\alpha\}}} C_{\vec{\pi}} E(\vec{x}, \vec{y}) \tag{A-19}$$

where each $C_{\vec{\pi}}$ is some explicit constant only depending on $\vec{\pi}$. Applying (A-19) to (A-16) gives the desired result. □

Appendix B. Increasing support for the nonsplit family

We prove Theorem 1.4. Arguing as in Appendix E of [21], we need to bound terms of the form

$$\bar{E}(\vec{n}, \vec{m}) := 2\pi i^k \sum_{\substack{q_1, \dots, q_\ell \\ q_j \text{ distinct}}} \prod_{j=1}^{\ell} \left(\hat{\phi} \left(\frac{\log q_j}{\log R} \right)^{n_j} \left(\frac{2 \log q_j}{\sqrt{q_j} \log R} \right)^{m_j} \right) \sum_{m \leq N^\epsilon} \frac{1}{m} \sum_{b=1}^{\infty} \frac{S(m^2, Q; Nb)}{Nb} J_{k-1} \left(\frac{4\pi m \sqrt{Q}}{Nb} \right), \tag{B-1}$$

where $Q = q_1^{m_1} \cdots q_\ell^{m_\ell}$ and $n_j \equiv m_j \pmod{2}$ for all j . Showing that these terms vanish as $N \rightarrow \infty$ for $\hat{\phi}$ with $\text{supp } \hat{\phi} \subset (-\frac{2}{n}, \frac{2}{n})$ completes the proof of Theorem 1.4. These terms are very similar to the $E(\vec{n}, \vec{m})$ terms introduced in Section 3 (see (3-4), for example), and we are able to evaluate them in a similar fashion. We omit proofs as they are analogous to the proofs of the corresponding lemmas in Section 3, which we refer to. We will eventually prove the following lemma.

Lemma B.1. *Let $\bar{E}(\vec{n}, \vec{m})$ be defined as in (B-1). Under GRH for Dirichlet L-functions, if $\text{supp}(\hat{\phi}) \subset (-\frac{2}{n}, \frac{2}{n})$, then $\bar{E}(\vec{n}, \vec{m}) \ll N^{-\epsilon}$ and thus does not contribute in the limit.*

First we restrict the sum over b as in Lemmas 3.3 and 3.4.

Lemma B.2. *Suppose $\text{supp}(\hat{\phi}) \subseteq (-\frac{7}{2n}, \frac{7}{2n})$. Then the subterms of $\bar{E}(\vec{n}, \vec{m})$ in (B-1) for which $(b, N) > 1$ are $O(N^{-\epsilon})$.*

Lemma B.3. *Suppose $\text{supp}(\hat{\phi}) \subset (-\frac{1000}{n}, \frac{1000}{n})$. Then the subterms of $\bar{E}(\vec{n}, \vec{m})$ in (B-1) for which $b \geq N^{2022}$ are $O(N^{-12})$.*

Applying Lemmas B.2 and B.3 to (B-1) gives

$$\begin{aligned} \bar{E}(\vec{n}, \vec{m}) &= 2\pi i^k \sum_{\substack{q_1, \dots, q_\ell \\ q_j \text{ distinct}}} \prod_{j=1}^{\ell} \left(\hat{\phi} \left(\frac{\log q_j}{\log R} \right)^{n_j} \left(\frac{2 \log q_j}{\sqrt{q_j} \log R} \right)^{n_j} \right) \sum_{m \leq N^\epsilon} \frac{1}{m} \\ &\quad \times \sum_{\substack{(b, N)=1 \\ b < N^{2022}}} \frac{S(m^2, Q; Nb)}{Nb} J_{k-1} \left(\frac{4\pi m \sqrt{Q}}{Nb} \right) + O(N^{-\epsilon}). \end{aligned} \tag{B-2}$$

We convert the Kloosterman sums to sums over Gauss sums as in Lemma 2.2.

Lemma B.4. *Let N be a prime not dividing b, Q, m . Then*

$$S(m^2, NQ; Nb) = -\frac{1}{\varphi(Nb)} \sum_{\chi(Nb)} G_\chi(m^2) G_\chi((Q, b^\infty)) \bar{\chi} \left(\frac{Q}{(Q, b^\infty)} \right). \tag{B-3}$$

Applying Lemma B.4 to (B-2) gives

$$\begin{aligned} \bar{E}(\vec{n}, \vec{m}) &= -2\pi i^k \sum_{\substack{q_1, \dots, q_\ell \\ q_j \text{ distinct}}} \prod_{j=1}^{\ell} \left(\hat{\phi} \left(\frac{\log q_j}{\log R} \right)^{n_j} \left(\frac{2 \log q_j}{\sqrt{q_j} \log R} \right)^{n_j} \right) \sum_{m \leq N^\epsilon} \frac{1}{m} \\ &\quad \times \sum_{\substack{(b, N)=1 \\ b < N^{2022}}} \frac{1}{Nb\varphi(Nb)} \sum_{\chi(Nb)} G_\chi(m^2) G_\chi((Q, b^\infty)) \bar{\chi} \left(\frac{Q}{(Q, b^\infty)} \right) J_{k-1} \left(\frac{4\pi m \sqrt{Q}}{Nb} \right) + O(N^{-\epsilon}). \end{aligned} \tag{B-4}$$

Next, subterms involving nonprincipal characters in (B-4) are negligible in the limit. This leaves only subterms involving $\chi_0 = \bar{\chi}_0 \pmod{Nb}$ for each b . It holds that $G_{\chi_0}(x) = R(x, Nb)$, a Ramanujan sum.

Lemma B.5. *Assume GRH for Dirichlet L -functions and suppose that $\text{supp}(\hat{\phi}) \subset (-\frac{2}{n}, \frac{2}{n})$. Then the sum over all nonprincipal characters in (B-4) is $O(N^{-\epsilon})$.*

This lemma corresponds to Lemma 3.5. Applying Lemma B.5 to (B-4) gives

$$\begin{aligned} \bar{E}(\vec{n}, \vec{m}) &= -2\pi i^k \sum_{\substack{q_1, \dots, q_\ell \\ q_j \text{ distinct}}} \prod_{j=1}^{\ell} \left(\hat{\phi} \left(\frac{\log q_j}{\log R} \right)^{n_j} \left(\frac{2 \log q_j}{\sqrt{q_j} \log R} \right)^{n_j} \right) \sum_{m \leq N^\epsilon} \frac{1}{m} \\ &\quad \times \sum_{\substack{(b, N)=1 \\ b < N^{2022}}} \frac{R(m^2, Nb) R((Q, b^\infty), Nb)}{Nb\varphi(Nb)} \chi_0 \left(\frac{Q}{(Q, b^\infty)} \right) J_{k-1} \left(\frac{4\pi m \sqrt{Q}}{Nb} \right) + O(N^{-\epsilon}). \end{aligned} \tag{B-5}$$

Applying the bounds $R(m^2, Nb) \leq m^4$, $R(x, Nb) \leq \varphi(Nb)$, and $J_{k-1}(x) \ll x$ to (B-5) and using the fact that $\text{supp} \hat{\phi} \subset (-\frac{2}{n}, \frac{2}{n})$, we find that the main term is absorbed by the error term, completing the proof of Lemma B.1.

Appendix C. Proofs of the lemmas in Section 5

C.1. Proof of Lemma 5.17. We will consider each term appearing in (5-35) separately. First, define

$$\begin{aligned} g_1(n, f, c, d) &:= \binom{n}{f}, & g_2(n, f, c, d) &:= \binom{n-c}{f-c}, \\ g_3(n, f, c, d) &:= \binom{n-d}{f-d}, & g_4(n, f, c, d) &:= \binom{n-c-d}{f-c-d}, \end{aligned} \quad (\text{C-1})$$

and

$$G_i(n, f) := 2(-1)^n n! \sum_{\substack{c+d \leq n \\ c, d \geq 0}} (-1)^{c+d+1} \frac{g_i(n, f, c, d)}{(n-c-d)!c!d!}. \quad (\text{C-2})$$

We want to evaluate $G_1(n, f) - G_2(n, f) - G_3(n, f) + G_4(n, f)$. We set $\ell = c + d$ to rewrite (C-2) as

$$G_i(n, f) = 2(-1)^{n+1} n! \sum_{\ell=0}^n (-1)^\ell \sum_{c=0}^{\ell} \frac{g_i(n, f, c, \ell-c)}{(n-\ell)!(\ell-c)!c!}. \quad (\text{C-3})$$

To evaluate $G_1(n, f)$, we group the binomial coefficients to find that

$$G_1(n, f) = 2(-1)^{n+1} \binom{n}{f} \sum_{\ell=0}^n (-1)^\ell \binom{n}{\ell} \sum_{c=0}^{\ell} \binom{\ell}{c} = 2(-1)^{n+1} \binom{n}{f} \sum_{\ell=0}^n (-2)^\ell \binom{n}{\ell} = -2 \binom{n}{f}. \quad (\text{C-4})$$

Next, we note that $G_2(n, f) = G_3(n, f)$. We have

$$G_2(n, f) = 2(-1)^{n+1} \binom{n}{f} \sum_{\ell=0}^n (-1)^\ell \sum_{c=0}^{\ell} \binom{f}{c} \binom{n-c}{n-\ell} = 2(-1)^{n+1} \binom{n}{f} \sum_{c=0}^n \binom{f}{c} \sum_{\ell=c}^n (-1)^\ell \binom{n-c}{n-\ell}.$$

We reindex the sum by setting $\ell' = \ell - c$. Doing so, we see that sum over ℓ' is zero unless $n - c = 0$. However, in this case we have $\binom{f}{c} = 0$ since $f \leq n/2 < n$. Thus each term vanishes and

$$G_2(n, f) = G_3(n, f) = 0. \quad (\text{C-5})$$

Lastly, again grouping terms into binomial coefficients gives

$$G_4(n, f) = 2(-1)^{n+1} \binom{n}{f} \sum_{\ell=0}^n (-1)^\ell \binom{f}{\ell} \sum_{c=0}^{\ell} \binom{\ell}{c} = 2(-1)^{n+1} \binom{n}{f} \sum_{\ell=0}^n (-2)^\ell \binom{f}{\ell}. \quad (\text{C-6})$$

We may restrict the sum in the last line to $0 \leq \ell \leq f$ since $f \leq n$ and $\binom{f}{\ell} = 0$ when $\ell > f$. Doing so, we find the sum over ℓ is $(-1)^f$ so

$$G_4(n, f) = 2(-1)^{n+f+1} \binom{n}{f}. \quad (\text{C-7})$$

Combining (C-4), (C-5) and (C-7) completes the proof of the lemma.

C.2. Proof of Lemma 5.28. We consider each term appearing in (5-55) separately. First, define

$$\begin{aligned} h_1(f, g, \mu_1, \mu_d) &:= \binom{f}{g}, & h_2(f, g, \mu_1, \mu_d) &:= \binom{f-\mu_1}{g-\mu_1} \\ h_3(f, g, \mu_1, \mu_d) &:= \binom{f-\mu_d}{g}, & h_4(f, g, \mu_1, \mu_d) &:= \binom{f-\mu_1-\mu_d}{g-\mu_1}, \end{aligned}$$

and

$$H_i(f, g) := \sum_{d=1}^f \sum_{\substack{\mu_1+\dots+\mu_d=f \\ \mu_i \geq 1}} \frac{(-1)^d}{\mu_1! \cdots \mu_d!} h_i(f, g, \mu_1, \mu_d) \tag{C-8}$$

for $i \in \{1, 2, 3, 4\}$. We will show that $H_i(f, g) = (-1)^f / g!(f - g)!$ independent of i , so that $H_1 - H_2 - H_3 + H_4 = 0$ as desired. For H_1 the result follows immediately from comparing coefficients of z^f in the identity (5-40). For H_2 , we pull out the μ_1 term to get

$$H_2(f, g) = - \sum_{\mu_1=1}^f \frac{1}{\mu_1!} \binom{f-\mu_1}{g-\mu_1} \sum_{d=1}^{f-\mu_1} \sum_{\substack{\mu_2+\dots+\mu_d=f-\mu_1 \\ \mu_i \geq 1}} \frac{(-1)^{d-1}}{\mu_2! \cdots \mu_d!}. \tag{C-9}$$

Applying (5-40) and simplifying gives

$$H_2(f, g) = - \sum_{\mu_1=1}^f \frac{1}{\mu_1!} \binom{f-\mu_1}{g-\mu_1} \frac{(-1)^{f-\mu_1}}{(f-\mu_1)!} = - \frac{(-1)^f}{(f-g)!} \sum_{\mu_1=1}^f \frac{(-1)^{\mu_1}}{\mu_1!(g-\mu_1)!} = \frac{(-1)^f}{g!(f-g)!}, \tag{C-10}$$

where the last step comes from restricting the summation to $1 \leq \mu_1 \leq g$ and using the binomial expansion of $(1 - 1)^g$. We can show the result for H_3 similarly. For H_4 , we pull out the μ_1 and μ_d terms to get

$$H_4(f, g) = \sum_{\mu_d=1}^f \sum_{\mu_1=1}^{f-\mu_d} \frac{1}{\mu_1! \mu_d!} \binom{f-\mu_1-\mu_d}{g-\mu_1} \sum_{d=1}^{f-\mu_1-\mu_d} \sum_{\substack{\mu_2+\dots+\mu_{d-1}=f-\mu_1-\mu_d \\ \mu_i \geq 1}} \frac{(-1)^{d-2}}{\mu_2! \cdots \mu_{d-1}!}. \tag{C-11}$$

Applying (5-40) and simplifying gives

$$\begin{aligned} H_4(f, g) &= \sum_{\mu_1=1}^f \sum_{\mu_d=1}^{f-\mu_1} \frac{1}{\mu_1! \mu_d!} \binom{f-\mu_1-\mu_d}{g-\mu_1} \frac{(-1)^{f-\mu_1-\mu_d}}{(f-\mu_1-\mu_d)!} \\ &= (-1)^f \sum_{\mu_d=1}^f \frac{(-1)^{\mu_d}}{(f-g-\mu_d)! \mu_d!} \sum_{\mu_1=1}^{f-\mu_d} \frac{(-1)^{\mu_1}}{\mu_1!(g-\mu_1)!} \\ &= \frac{(-1)^{f+1}}{g!} \sum_{\mu_d=1}^f \frac{(-1)^{\mu_d}}{(f-g-\mu_d)! \mu_d!} = \frac{(-1)^f}{g!(f-g)!} \end{aligned} \tag{C-12}$$

where the last two steps come from restricting the summation to $1 \leq \mu_1 \leq g$ and $1 \leq \mu_d \leq f - g$ and using the binomial expansion of $(1 - 1)^g$ and $(1 - 1)^{f-g}$.

Appendix D. Bounding the order of vanishing at the central point

In this section, we follow the arguments of Section 6 of [21] in order to bound the proportion of newforms with negative sign whose order of vanishing exceeds a certain threshold r . While they are conditional on GRH, our results surpass the best known conditional and unconditional bounds established in [25], [21], and [5] when $r \geq 5$. We focus on the case $r = 5$, however our results may be easily generalized the case when $r > 5$. Additionally, we study the fourth centered moment as it provides the best bounds for the case $r = 5$, but utilizing higher moments provides better bounds as r increases. Lastly, similar results may be obtained for the positive sign family. See [13] for a more in-depth analysis, where the results of this paper are used to find excellent bounds for vanishing to order r or more; specifically, for a fixed test function Dutta and Miller determine what level density gives the best bound.

We utilize Theorem 1.2 with $n = 4$ and

$$\phi(x) = \left(\frac{\sin \pi \sigma x}{\pi \sigma x} \right)^2, \quad \hat{\phi}(y) = \begin{cases} 1/\sigma - |y|/\sigma^2 & \text{if } |y| < \sigma \\ 0 & \text{if } |y| \geq \sigma. \end{cases} \tag{D-1}$$

This test function is likely not optimal in general for minimizing the n -th centered moment, and optimal test functions for the case $n = 1$ and $n = 2$ are found in [25] and [5]. However, they are sufficient to surpass the bounds established in those papers. While Theorem 1.2 requires $\sigma < 0.5$ when $n = 4$, we may utilize the bounds given by $\sigma = 0.5$ by setting $\sigma = 0.5 - \epsilon$ and letting $\epsilon \rightarrow 0$. Now, Theorem 1.2 gives

$$\lim_{\substack{N \rightarrow \infty \\ N \text{ prime}}} \left(\langle D(f; \phi) - \langle D(f; \phi) \rangle_- \rangle_-^4 \right) = 3(\sigma_\phi^2)^2 - R(4, 2; \phi) = \frac{31}{105}. \tag{D-2}$$

Now, if a newform f with negative sign has order of vanishing $r \geq 5$ at the central point, then by Theorem 1.4,

$$D(f; \phi) - \langle D(f; \phi) \rangle_- \geq r\phi(0) - (\hat{\phi}(0) + \frac{1}{2}\phi(0)) = r - \frac{5}{2} \geq \frac{5}{2}. \tag{D-3}$$

Let $\Pr(r \geq 5)$ be the proportion of newforms with negative sign whose order of vanishing at the central point is at least 5. Then (D-2) and (D-3) give

$$\Pr(r \geq 5) \left(\frac{5}{2} \right)^4 \leq \frac{31}{105}, \tag{D-4}$$

so

$$\Pr(r \geq 5) \leq \left(\frac{2}{5} \right)^4 \frac{31}{105} = \frac{496}{65625} \approx 0.00756.$$

In [25] and [21] the respective upper bounds $\frac{1}{32} = 0.03125$ and $\frac{1}{49} \approx 0.02040$ were obtained; our results surpass both of these. As the order of vanishing increases, our results are even better. For instance, taking $r = 19$ and $n = 20$, we find the proportion of newforms with negative sign whose order of vanishing exceeds 19 is at most $2.86 \cdot 10^{-15}$, improving the upper bound $5.77 \cdot 10^{-6}$ given in [5] and the upper bound $3.29 \cdot 10^{-3}$ implicit in [25].

References

- [1] L. Alpoge and S. J. Miller, “Low-lying zeros of Maass form L -functions”, *Int. Math. Res. Not. IMRN* **2015**:10 (2015), 2678–2701. MR
- [2] L. Alpoge, N. Amersi, G. Iyer, O. Lazarev, S. J. Miller, and L. Zhang, “Maass waveforms and low-lying zeros”, in *Analytic number theory: in honor of Helmut Maier’s 60th birthday*.
- [3] T. M. Apostol, *Introduction to analytic number theory*, Springer-Verlag, New York-Heidelberg, 1976. MR
- [4] O. Barrett, P. Burkhardt, J. DeWitt, R. Dorward, and S. J. Miller, “One-level density for holomorphic cusp forms of arbitrary level”, *Res. Number Theory* **3** (2017), art. id. 25, 21 pp. MR
- [5] E. Boldyriew, F. Chen, C. Devlin, VI, S. J. Miller, and J. Zhao, “Determining optimal test functions for 2-level densities”, *Res. Number Theory* **9**:2 (2023), art. id. 32, 17 pp. MR
- [6] V. Chandee and Y. Lee, “ n -level density of the low-lying zeros of primitive Dirichlet L -functions”, *Adv. Math.* **369** (2020), art. id. 107185, 70 pp. MR
- [7] J. B. Conrey and N. C. Snaith, “In support of n -correlation”, *Comm. Math. Phys.* **330**:2 (2014), 639–653. MR
- [8] H. Davenport, *Multiplicative number theory*, vol. 74, 2nd ed., Graduate Texts in Mathematics, Springer-Verlag, 1980. MR
- [9] L. Devin, D. Fiorilli, and A. Södergren, “Low-lying zeros in families of holomorphic cusp forms: the weight aspect”, *Q. J. Math.* **73**:4 (2022), 1403–1426. MR
- [10] S. Drappeau, K. Pratt, and M. Radziwiłł, “One-level density estimates for Dirichlet L -functions with extended support”, *Algebra Number Theory* **17**:4 (2023), 805–830. MR
- [11] E. Dueñez and S. J. Miller, “The low lying zeros of a $GL(4)$ and a $GL(6)$ family of L -functions”, *Compos. Math.* **142**:6 (2006), 1403–1425. MR
- [12] E. Dueñez and S. J. Miller, “The effect of convolving families of L -functions on the underlying group symmetries”, *Proc. Lond. Math. Soc.* (3) **99**:3 (2009), 787–820. MR
- [13] S. Dutta and S. J. Miller, “Bounding the order of vanishing of cuspidal newforms via the n^{th} centered moments”, *Res. Number Theory* **10**:4 (2024), art. id. 76, 22 pp. MR
- [14] A. Entin, “Artin-Schreier L -functions and random unitary matrices”, *J. Number Theory* **145** (2014), 340–351. MR
- [15] A. Entin, E. Roditty-Gershon, and Z. Rudnick, “Low-lying zeros of quadratic Dirichlet L -functions, hyper-elliptic curves and random matrix theory”, *Geom. Funct. Anal.* **23**:4 (2013), 1230–1261. MR
- [16] D. Fiorilli and S. J. Miller, “Surpassing the ratios conjecture in the 1-level density of Dirichlet L -functions”, *Algebra Number Theory* **9**:1 (2015), 13–52. MR
- [17] P. Gao, “ n -level density of the low-lying zeros of quadratic Dirichlet L -functions”, *Int. Math. Res. Not. IMRN* **2014**:6 (2014), 1699–1728. MR
- [18] I. S. Gradshteyn and I. M. Ryzhik, *Table of integrals, series, and products*, 8th ed., Elsevier, Amsterdam, 2015. MR
- [19] A. M. Güloğlu, “Low-lying zeroes of symmetric power L -functions”, *Int. Math. Res. Not.* **2005**:9 (2005), 517–550. MR
- [20] R. K. Guy, “Catwalks, sandsteps and Pascal pyramids”, *J. Integer Seq.* **3**:1 (2000), art. id. 00.1.6. MR
- [21] C. P. Hughes and S. J. Miller, “Low-lying zeros of L -functions with orthogonal symmetry”, *Duke Math. J.* **136**:1 (2007), 115–172. MR
- [22] C. P. Hughes and Z. Rudnick, “Mock-Gaussian behaviour for linear statistics of classical compact groups”, *J. Phys. A* **36**:12 (2003), 2919–2932. MR
- [23] H. Iwaniec, *Topics in classical automorphic forms*, Graduate Studies in Mathematics **17**, Amer. Math. Soc., 1997. MR
- [24] H. Iwaniec and E. Kowalski, *Analytic number theory*, vol. 53, AMS Colloquium Publications, Amer. Math. Soc., 2004. MR
- [25] H. Iwaniec, W. Luo, and P. Sarnak, “Low lying zeros of families of L -functions”, *Inst. Hautes Études Sci. Publ. Math.* **91** (2000), 55–131. MR

- [26] N. M. Katz and P. Sarnak, *Random matrices, Frobenius eigenvalues, and monodromy*, vol. 45, Amer. Math. Soc. Colloquium Publications, Amer. Math. Soc., Providence, RI, 1999. MR
- [27] N. M. Katz and P. Sarnak, “Zeroes of zeta functions and symmetry”, *Bull. Amer. Math. Soc. (N.S.)* **36**:1 (1999), 1–26. MR
- [28] A. Knightly and C. Reno, “Weighted distribution of low-lying zeros of $GL(2)$ L -functions”, *Canad. J. Math.* **71**:1 (2019), 153–182. MR
- [29] J. Levinson and S. J. Miller, “The n -level densities of low-lying zeros of quadratic Dirichlet L -functions”, *Acta Arith.* **161**:2 (2013), 145–182. MR
- [30] J. Li and S. J. Miller, “Bounding vanishing at the central point of cuspidal newforms”, *J. Number Theory* **244** (2023), 279–307. MR
- [31] A. M. Mason and N. C. Snaith, *Orthogonal and symplectic n -level densities*, *Memoirs of the Amer. Math. Soc.* **1194**, 2018. MR
- [32] A. Mauro, J. B. Miller, and S. J. Miller, “Extending the support of 1- and 2-level densities for cusp form L -functions under square-root cancellation hypotheses”, *Acta Arith.* **214** (2024), 289–309. MR
- [33] S. J. Miller, “One- and two-level densities for rational families of elliptic curves: evidence for the underlying group symmetries”, *Compos. Math.* **140**:4 (2004), 952–992. MR
- [34] S. J. Miller, “An orthogonal test of the L -functions ratios conjecture”, *Proc. Lond. Math. Soc. (3)* **99**:2 (2009), 484–520. MR
- [35] S. J. Miller and R. Peckner, “Low-lying zeros of number field L -functions”, *J. Number Theory* **132**:12 (2012), 2866–2891. MR
- [36] H. L. Montgomery, “The pair correlation of zeros of the zeta function”, pp. 181–193 in *Analytic number theory* (St. Louis, MO, 1972), Proc. Sympos. Pure Math. **24**, Amer. Math. Soc., 1973. MR
- [37] H. L. Montgomery and R. C. Vaughan, *Multiplicative number theory, I: Classical theory*, Cambridge Studies in Advanced Mathematics **97**, Cambridge University Press, 2007. MR
- [38] A. E. Özlük and C. Snyder, “Small zeros of quadratic L -functions”, *Bull. Austral. Math. Soc.* **47**:2 (1993), 307–319. MR
- [39] A. E. Özlük and C. Snyder, “On the distribution of the nontrivial zeros of quadratic L -functions close to the real axis”, *Acta Arith.* **91**:3 (1999), 209–228. MR
- [40] G. Ricotta and E. Royer, “Statistics for low-lying zeros of symmetric power L -functions in the level aspect”, *Forum Math.* **23**:5 (2011), 969–1028. MR
- [41] E. Royer, “Petits zéros de fonctions L de formes modulaires”, *Acta Arith.* **99**:2 (2001), 147–172. MR
- [42] M. Rubinstein, “Low-lying zeros of L -functions and random matrix theory”, *Duke Math. J.* **109**:1 (2001), 147–181. MR
- [43] Z. Rudnick and P. Sarnak, “Zeros of principal L -functions and random matrix theory”, *Duke Math. J.* **81**:2 (1996), 269–322. MR
- [44] S. W. Shin and N. Templier, “Sato-Tate theorem for families and low-lying zeros of automorphic L -functions”, *Invent. Math.* **203**:1 (2016), 1–177. MR
- [45] A. Soshnikov, “The central limit theorem for local linear statistics in classical compact groups and related combinatorial identities”, *Ann. Probab.* **28**:3 (2000), 1353–1370. MR
- [46] G. N. Watson, *A treatise on the theory of Bessel functions*, Cambridge University Press, 1944. MR
- [47] E. Waxman, “Lower order terms for the one-level density of a symplectic family of Hecke L -functions”, *J. Number Theory* **221** (2021), 447–483. MR
- [48] A. Yang, *Distribution problems associated to zeta functions and invariant theory*, Ph.D. thesis, Princeton University, 2009, available at <https://www.proquest.com/docview/304982142>. MR
- [49] M. P. Young, “Low-lying zeros of families of elliptic curves”, *J. Amer. Math. Soc.* **19**:1 (2006), 205–250. MR

Communicated by Philippe Michel

Received 2022-10-21 Revised 2024-07-19 Accepted 2025-03-03

plcohen@mit.edu	<i>Department of Mathematics, Massachusetts Institute of Technology, Cambridge, MA 02139, United States</i>
jsdell@ucsd.edu	<i>Department of Mathematics, UCSD, 9500 Gillman Drive, San Diego, CA 92093, United States</i>
oscarq39@gmail.com	<i>Department of Mathematics, University of Puerto Rico, Box 70377, San Juan 00931, Puerto Rico</i>
skhunger@umich.edu	<i>Department of Mathematics, University of Michigan, 530 Church St, Ann Arbor, MI 48109, United States</i>
chkwan@ias.edu	<i>Institute for Advanced Study, Princeton, NJ, United States</i>
sjm1@williams.edu	<i>Department of Mathematics and Statistics, Williams College, 18 Hoxsey St, Williamstown, MA 01267, United States</i>
ashashkov@berkeley.edu	<i>Department of Mathematics, University of California, Berkeley, Berkeley, CA, United States</i>
alicita.smith01@gmail.com	<i>Department of Mathematics, Williams College, Williamstown, MA, United States</i>
csprun@umich.edu	<i>Department of Mathematics, University of Michigan, Ann Arbor, MI, United States</i>
n.triantafillou@idaccr.org	<i>IDA Center for Communications Research, Princeton, NJ, United States</i>
ntruongv@stanford.edu	
rogervanpeski@gmail.com	<i>Department of Mathematics, Columbia University, New York, NY, United States</i>
sdw2@williams.edu	<i>Department of Mathematics, Williams College, Williamstown, MA, United States</i>

Algebra & Number Theory

msp.org/ant

EDITORS

MANAGING EDITOR
Antoine Chambert-Loir
Université Paris-Diderot
France

EDITORIAL BOARD CHAIR
David Eisenbud
University of California
Berkeley, USA

BOARD OF EDITORS

Jason P. Bell	University of Waterloo, Canada	Philippe Michel	École Polytechnique Fédérale de Lausanne
Bhargav Bhatt	University of Michigan, USA	Martin Olsson	University of California, Berkeley, USA
Frank Calegari	University of Chicago, USA	Irena Peeva	Cornell University, USA
J.-L. Colliot-Thélène	CNRS, Université Paris-Saclay, France	Jonathan Pila	University of Oxford, UK
Brian D. Conrad	Stanford University, USA	Anand Pillay	University of Notre Dame, USA
Samit Dasgupta	Duke University, USA	Bjorn Poonen	Massachusetts Institute of Technology, USA
Hélène Esnault	Freie Universität Berlin, Germany	Victor Reiner	University of Minnesota, USA
Gavril Farkas	Humboldt Universität zu Berlin, Germany	Peter Sarnak	Princeton University, USA
Sergey Fomin	University of Michigan, USA	Michael Singer	North Carolina State University, USA
Edward Frenkel	University of California, Berkeley, USA	Vasudevan Srinivas	SUNY Buffalo, USA
Wee Teck Gan	National University of Singapore	Shunsuke Takagi	University of Tokyo, Japan
Andrew Granville	Université de Montréal, Canada	Pham Huu Tiep	Rutgers University, USA
Ben J. Green	University of Oxford, UK	Ravi Vakil	Stanford University, USA
Christopher Hacon	University of Utah, USA	Akshay Venkatesh	Institute for Advanced Study, USA
Roger Heath-Brown	Oxford University, UK	Melanie Matchett Wood	Harvard University, USA
János Kollár	Princeton University, USA	Shou-Wu Zhang	Princeton University, USA
Michael J. Larsen	Indiana University Bloomington, USA		

PRODUCTION

production@msp.org
Silvio Levy, Scientific Editor

See inside back cover or msp.org/ant for submission instructions.

The subscription price for 2026 is US \$590/year for the electronic version, and \$865/year (+\$75, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to MSP.

Algebra & Number Theory (ISSN 1944-7833 electronic, 1937-0652 printed) at Mathematical Sciences Publishers, 2000 Allston Way # 59, Berkeley, CA 94701-4004, is published continuously online.

ANT peer review and production are managed by EditFLOW[®] from MSP.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2026 Mathematical Sciences Publishers

Algebra & Number Theory

Volume 20 No. 2 2026

On the minimal dimension of a faithful linear representation of a finite group	219
ALEXANDER MORETÓ	
Moments of one-level densities in families of holomorphic cusp forms in the level aspect	237
PETER COHEN, JUSTINE DELL, OSCAR E. GONZÁLEZ, SIMRAN KHUNGER, CHUNG-HANG KWAN, STEVEN J. MILLER, ALEXANDER SHASHKOV, ALICIA REINA SMITH, CARSTEN SPRUNGER, NICHOLAS TRIAANTAFILLOU, NHI TRUONG, ROGER VAN PESKI and STEPHEN WILLIS	
Murmurations of modular forms in the weight aspect	299
JONATHAN BOBER, ANDREW R. BOOKER, MIN LEE and DAVID LOWRY-DUDA	
The Alperin weight conjecture and the Glauberman correspondence via character triples	333
J. MIQUEL MARTÍNEZ, NOELIA RIZO and DAMIANO ROSSI	
Asymptotics of extensions of simple \mathbb{Q} -algebras	383
FABIAN GUNDLACH and BÉRANGER SEGUIN	