

Algebra & Number Theory

Volume 20
2026
No. 2

Murmurations of modular forms in the weight aspect

Jonathan Bober, Andrew R. Booker, Min Lee and David Lowry-Duda



Murmurations of modular forms in the weight aspect

Jonathan Bober, Andrew R. Booker, Min Lee and David Lowry-Duda

We prove the existence of “murmurations” in the family of holomorphic modular forms of level 1 and weight $k \rightarrow \infty$, that is, correlations between their root numbers and Hecke eigenvalues at primes growing in proportion to the analytic conductor. This is the first demonstration of murmurations in an archimedean family.

1. Introduction

Using machine learning algorithms, He, Lee, Oliver, and Pozdnyakov [HLOP22] discovered an apparent correlation between the root numbers of elliptic curve L -functions and their Dirichlet coefficients a_p at primes p varying in proportion to the conductor. They dubbed this correlation “murmurations of elliptic curves” due to the resemblance of their graphs to the swarming patterns of flocks of birds (see Figure 1). Further experimental work by the authors of [HLOP22] and Sutherland [HLOPS] has shown this to be a general phenomenon, present in many natural families of L -functions.

Zubrilina [Zub23] has provided the first theoretical confirmation, proving the existence of murmurations in the family of self-dual holomorphic newforms of a fixed weight k and squarefree conductor $N \rightarrow \infty$.




Figure 1. Left: average a_p (ordinate) over isogeny classes of elliptic curves of conductor in $[7500, 10000]$ and fixed rank (blue = rank 0, red = rank 1), for primes $p \in [2, 7919]$ (abscissa). Reprinted from [HLOP22] with permission from the authors. Right: *Starling shapes in the evening sky* © Walter Baxter (<https://www.geograph.org.uk/p/1065181>, cc-by-sa/2.0).

Bober and Booker are supported by the Heilbronn Institute for Mathematical Research. Lee is supported by a Royal Society University Research Fellowship. Lowry-Duda is supported by the Simons Collaboration in Arithmetic Geometry, Number Theory, and Computation via the Simons Foundation grant 546235.

MSC2020: primary 11F30, 11N60; secondary 11F72.

Keywords: murmurations, weight aspect, modular forms.




Figure 2. A comparison of $M_2(y)$ for $y \in [0, 2]$ (green) and the points $(p/2^{18}, r(p))$ for primes $p < 2^{19}$ (purple), where

$$r(p) = \frac{\sum_{N \in I} \sum_{f \in H_2(N)} \epsilon_f \lambda_f(p) \sqrt{p}}{\sum_{N \in I} \sum_{f \in H_2(N)} 1},$$

and I is the set of all squarefree integers in the range $[2^{18} \pm 2^{10}]$. Data computed by Andrew Sutherland.

More precisely, for $k \in 2\mathbb{Z}_{>0}$ and $N \in \mathbb{Z}_{>0}$, let $H_k(N)$ be a basis of normalized Hecke eigenforms for $S_k^{\text{new}}(\Gamma_0(N))$. For each $f \in H_k(N)$, let ϵ_f be its root number, and let $\lambda_f(n)$ be its normalized Hecke eigenvalues, so that

$$f(z) = \sum_{n=1}^{\infty} \lambda_f(n) n^{\frac{k-1}{2}} e^{2\pi i n z}.$$

Then Zubrilina showed that there exists a continuous function $M_k : \mathbb{R}_{>0} \rightarrow \mathbb{R}$ such that, for fixed $y \in \mathbb{R}_{>0}$ and $\delta \in (0, 1)$, we have

$$\lim_{\substack{p \text{ prime} \\ p \rightarrow \infty}} \frac{\sum_{\substack{N \in [p/y, p/y+p^\delta] \cap \mathbb{N} \\ N \text{ squarefree}}} \sum_{f \in H_k(N)} \epsilon_f \lambda_f(p) \sqrt{p}}{\sum_{\substack{N \in [p/y, p/y+p^\delta] \cap \mathbb{N} \\ N \text{ squarefree}}} \sum_{f \in H_k(N)} 1} = M_k(y).$$

A comparison of $M_2(y)$ with numerical data is shown in Figure 2.

In this paper, following a suggestion of Sarnak, we investigate the murmuration phenomenon in a family of L -functions with varying archimedean parameters, namely the modular forms of level 1 and weight $k \rightarrow \infty$. In this family, the root number ϵ_f is simply $(-1)^{\frac{k}{2}}$, so we expect to see biases in the $\lambda_f(p)$ values depending on the residue class of $k \pmod 4$. Following Rubinstein (unpublished, but see [Boo15] and [LMFDB]), we define the *analytic conductor* of $f \in H_k(1)$ to be the positive real number

$$\mathcal{N}(k) := \left(\frac{\exp \psi(k/2)}{2\pi} \right)^2,$$




Figure 3. A comparison of $(-1)^\delta \nu((0, t])$ and the left-hand side of (1-1) scaled by $t\sqrt{N}$, for $K = 3850$, $H = 100$, and $t \in [0, 2]$.

where $\psi(x) = \Gamma'(x)/\Gamma(x)$; by Stirling’s formula, $\mathcal{N}(k) = \left(\frac{k-1}{4\pi}\right)^2 + O(1)$. In analogy with the arithmetic conductor scaling in [HLOP22; Zub23], one might expect to see murmurations for p growing in proportion to $\mathcal{N}(k)$; our main result confirms this expectation under the generalized Riemann hypothesis (GRH):

Theorem 1.1. *Assume GRH for the L-functions of Dirichlet characters and modular forms. Fix $\varepsilon \in (0, \frac{1}{12})$, $\delta \in \{0, 1\}$, and a compact interval $E \subset \mathbb{R}_{>0}$ with $|E| > 0$. Let $K, H \in \mathbb{R}_{>0}$ with $K^{\frac{5}{6}+\varepsilon} < H < K^{1-\varepsilon}$, and set $N = \mathcal{N}(K)$. Then as $K \rightarrow \infty$, we have*

$$\frac{\sum_{p/N \in E} \log p \sum_{\substack{k \equiv 2\delta \pmod{4} \\ |k-K| \leq H}} \sum_{f \in H_k(1)} \lambda_f(p)}{\sum_{p/N \in E} \log p \sum_{\substack{k \equiv 2\delta \pmod{4} \\ |k-K| \leq H}} \sum_{f \in H_k(1)} 1} = \frac{(-1)^\delta}{\sqrt{N}} \left(\frac{\nu(E)}{|E|} + o_{E,\varepsilon}(1) \right), \tag{1-1}$$

where

$$\nu(E) = \frac{1}{\zeta(2)} \sum_{\substack{a, q \in \mathbb{Z}_{>0} \\ \gcd(a, q) = 1 \\ (a/q)^{-2} \in E}}^* \frac{\mu(q)^2}{\varphi(q)^2 \sigma(q)} \left(\frac{q}{a}\right)^3 = \frac{1}{2} \sum_{t=-\infty}^{\infty} \prod_{p \nmid t} \frac{p^2 - p - 1}{p^2 - p} \cdot \int_E \cos \frac{2\pi t}{\sqrt{y}} dy, \tag{1-2}$$

and the $*$ indicates that terms occurring at the endpoints of E are halved.

A comparison of $\nu(E)$ with numerical data is shown in Figure 3.

Remarks. (1) We have averaged the normalized $\lambda_f(p)$ values directly, which shows a correlation of size $1/\sqrt{N}$ with the root number. As in [HLOP22; Zub23; Sar23], one could instead average $\lambda_f(p)\sqrt{p}$ to boost the correlation. A similar result holds for that average, with the right-hand side becoming

$$(-1)^\delta \left(\frac{\nu^\sharp(E)}{|E|} + o_{E,\varepsilon}(1) \right),$$

where $d\nu^\sharp(y) = \sqrt{y} d\nu(y)$, i.e.,

$$\nu^\sharp(E) = \frac{1}{\zeta(2)} \sum_{\substack{a, q \in \mathbb{Z}_{>0} \\ \gcd(a, q) = 1 \\ (a/q)^{-2} \in E}}^* \frac{\mu(q)^2}{\varphi(q)^2 \sigma(q)} \left(\frac{q}{a}\right)^4 = \frac{1}{2} \sum_{t=-\infty}^{\infty} \prod_{p \nmid t} \frac{p^2 - p - 1}{p^2 - p} \cdot \int_E \sqrt{y} \cos \frac{2\pi t}{\sqrt{y}} dy.$$

(2) Zubrilina's results exhibit correlations for an individual prime p , requiring only a short average over conductors. One can carry out some of the analysis for a fixed p with our family, but the resulting correlation depends on the residue classes of p modulo small primes; at least a short average over p is needed to get a universal result depending only on the size of p/N . Sarnak [Sar23] has speculated that this is closely related to the conductor growth of the family, and that an average over p is necessary to see murmurations in any family with at most $O(x)$ L -functions of analytic conductor $\leq x$. (Note that our family contains $\asymp x$ forms of analytic conductor $\leq x$, compared to $\asymp_k x^2$ in Zubrilina's family.)

(3) Our result depends on GRH in order to compute the sums over primes with power-saving accuracy. However, GRH for modular form L -functions is only needed to handle the sharp cutoff in the sum over k in (1-1), and could be dispensed with in a smoothed version. The combination of a weak signal (only $1/\sqrt{N}$ in size) and a distribution that is not absolutely continuous makes the proof delicate, requiring many nontrivial estimates throughout, including for this truncation error. (The large sieve does not suffice here, since the number of forms involved is small relative to the size of p .)

(4) Closely related to murmurations is the 1-level density, which involves computing a_p averages in a family of L -functions for p on the scale of N^θ for some real number $\theta > 0$. For our family of holomorphic forms in the weight aspect, the 1-level density statistics for $\theta < 2$ were computed by Iwaniec, Luo, and Sarnak [ILS00], demonstrating a sharp phase transition at $\theta = 1$, in accordance with predictions from random matrix theory [KS99]. Sarnak [Sar23] has pointed out that, in retrospect, one can see murmurations in [ILS00] by examining the $\theta = 1$ transition more closely. One technical difference with our work is that [ILS00] used the Petersson trace formula (rather than Eichler–Selberg), which sums the a_p values of an L^2 -normalized basis of forms. This changes the distribution, but it shares some of the same qualitative features, including point masses at squares of squarefree integers (corresponding to the terms of (1-2) with $a = 1$).

(5) The $o(1)$ error term in Theorem 1.1 depends on Diophantine properties of the endpoints of E . More precisely, if we write $E = [u, v]$, then the theorem holds with $o_{E, \varepsilon}(1)$ replaced by

$$O_{E, \varepsilon} \left(\left(\frac{H}{K^{1-\varepsilon}} \right)^{\frac{1}{9}} + \left(\frac{H}{K^{\frac{5}{6} + \varepsilon}} \right)^{-\frac{6}{11}} \right),$$

provided that \sqrt{u} and \sqrt{v} have irrationality measures at most 27. Generally finite irrationality measure implies a power-saving error term, but for Liouville numbers the convergence can be arbitrarily slow.

(6) Our proof should carry over to the family of Maass forms of level 1 and Laplace eigenvalue $\lambda \rightarrow \infty$ with appropriate modifications. The main difference is that one would encounter discriminants $t^2 + 4n$ in

the trace formula instead of $t^2 - 4n$ (see (2-1)), but the local analysis in Lemmas 4.1–4.3 is insensitive to this change.

(7) Sutherland [Sut23] has empirically observed that the signal is less noisy if one computes correlation sums of ϵa_p rather than separating by root number, pointing to the existence of a lower-order discrepancy. Our proof suggests a possible source for this in our family, namely the terms of (3-3) with $\ell = \pm 1$, which vanish when considering the correlation. As we show in Lemma 3.1, those terms have lower order than the main term, and they are eventually swamped by other error terms in our analysis, so it may not be possible to isolate the discrepancy theoretically.

(8) Our distribution ν shows many qualitative differences from Zubrilina's M_k ; for example, it has infinitely many point masses, and no sign changes — the terms of (1-2) are all nonnegative. It is conceivable that M_2 and ν^\sharp are the margins of some joint distribution in the large analytic conductor limit, $Nk^2 \rightarrow \infty$. It would be interesting to try to merge the two results by taking a double limit; in particular, do $\lim_{N \rightarrow \infty} \lim_{k \rightarrow \infty}$ and $\lim_{k \rightarrow \infty} \lim_{N \rightarrow \infty}$ both exist, and are they equal?

(9) Although murmurations have hitherto been investigated as a pattern in a_p values at *primes*, (1-1) would continue to make sense with the sums over p replaced by sums over integers n . The analysis becomes much simpler with sums over integers, and there is still a discernible murmur: the analogue of $\nu(E)$ is

$$\sum_{\substack{a \in \mathbb{Z}_{>0} \\ a^{-2} \in E}}^* a^{-3}.$$

It may be of interest to look for integer murmurations in other families.

2. The trace formula and overview of the proof

The main tool that we use in the proof of Theorem 1.1 is the Eichler–Selberg trace formula, which allows us to evaluate the sums over $f \in H_k(1)$ in (1-1). For a positive integer n , let T_n denote the n th Hecke operator acting on $S_k(1)$, and let

$$\mathrm{Tr} T_n | S_k(1) = \sum_{f \in H_k(1)} \lambda_f(n) n^{\frac{k-1}{2}}$$

denote its trace. Then by [Chi22, Theorem 2.1] (see also [Eic57; Hij74; Zag77; Coh77; SvdV91]), we have

$$\mathrm{Tr} T_n | S_k(1) = A_1 - A_2 - A_3 + A_4,$$

where

$$A_1 = \begin{cases} \frac{1}{12} n^{\frac{k}{2}-1} (k-1) & \text{if } \sqrt{n} \in \mathbb{Z}, \\ 0 & \text{otherwise,} \end{cases}$$

$$A_2 = \sum_{\substack{t \in \mathbb{Z}, t^2 - 4n = d\ell^2 < 0 \\ \rho^2 - t\rho + n = 0, \Im \rho > 0}} \frac{\rho^{k-1} - \bar{\rho}^{k-1}}{\rho - \bar{\rho}} \frac{h(d)}{w(d)} \prod_{p|\ell} \left(p^{\text{ord}_p(\ell)} + \left(1 - \left(\frac{d}{p}\right)\right) \frac{p^{\text{ord}_p(\ell)} - 1}{p-1} \right),$$

$$A_3 = \frac{1}{2} \sum_{d|n} \min(d, n/d)^{k-1}, \quad \text{and} \quad A_4 = \begin{cases} \sigma(n) & \text{if } k = 2, \\ 0 & \text{otherwise.} \end{cases}$$

Here d is the discriminant of $\mathbb{Q}(\sqrt{t^2 - 4n})$, $h(d)$ is the class number, and $w(d)$ is the number of units in the ring of integers.

Following [BL17, §1.1], for a discriminant $D = d\ell^2$ we define $\psi_D(m) = \left(\frac{d}{m/\text{gcd}(m,\ell)}\right)$, with the convention that $\psi_0(m) = 1$. Then, by Dirichlet’s class number formula, for $D < 0$ we have

$$L(1, \psi_D) := \sum_{m=1}^{\infty} \frac{\psi_D(m)}{m} = \frac{2\pi h(d)}{w(d)\sqrt{|D|}} \prod_{p|\ell} \left(p^{\text{ord}_p(\ell)} + \left(1 - \left(\frac{d}{p}\right)\right) \frac{p^{\text{ord}_p(\ell)} - 1}{p-1} \right).$$

Thus A_2 can be written in the form

$$A_2 = \sum_{\substack{t \in \mathbb{Z}, t^2 < 4n \\ \rho^2 - t\rho + n = 0, \Im \rho > 0}} \frac{\rho^{k-1} - \bar{\rho}^{k-1}}{\rho - \bar{\rho}} \frac{\sqrt{4n - t^2}}{2\pi} L(1, \psi_{t^2 - 4n}).$$

Next, for $t \in \mathbb{Z}$ with $t^2 < 4n$, define

$$\phi_{t,n} = \arcsin \frac{t}{2\sqrt{n}} \in \left(-\frac{\pi}{2}, \frac{\pi}{2}\right).$$

Then we have $\rho = \sqrt{n}e^{i(\frac{\pi}{2} - \phi_{t,n})}$, where ρ is the solution to $\rho^2 - t\rho + n = 0$ with positive imaginary part. Thus,

$$\begin{aligned} \frac{\rho^{k-1} - \bar{\rho}^{k-1}}{\rho - \bar{\rho}} &= n^{\frac{k}{2}-1} \frac{\sin((k-1)(\frac{\pi}{2} - \phi_{t,n}))}{\sin(\frac{\pi}{2} - \phi_{t,n})} \\ &= -(-1)^{\frac{k}{2}} n^{\frac{k}{2}-1} \frac{\cos((k-1)\phi_{t,n})}{\cos(\phi_{t,n})} = -2(-1)^{\frac{k}{2}} n^{\frac{k-1}{2}} \frac{\cos((k-1)\phi_{t,n})}{\sqrt{4n - t^2}}. \end{aligned}$$

Assume now that n is prime and $k > 2$. Then, combining these expressions, we obtain

$$\sum_{f \in H_k(1)} \lambda_f(n) = n^{\frac{1-k}{2}} \text{Tr } T_n |S_k(1) = -n^{\frac{1-k}{2}} + \frac{(-1)^{\frac{k}{2}}}{\pi} \sum_{\substack{t \in \mathbb{Z} \\ t^2 < 4n}} \cos((k-1)\phi_{t,n}) L(1, \psi_{t^2 - 4n}). \quad (2-1)$$

2.1. Overview of the proof of Theorem 1.1. We conclude this section with a sketch of the proof of Theorem 1.1, with the details carried out in Sections 3–6.

For a fixed choice of δ, K, H , and E , we define $\Sigma = \Sigma(\delta, K, H, E)$ to be the numerator on the left-hand side of (1-1):

$$\Sigma = \sum_{\substack{n \text{ prime} \\ n/N \in E}} \log n \sum_{\substack{k \equiv 2\delta \pmod{4} \\ |k-K| \leq H}} \sum_{f \in H_k(1)} \lambda_f(n).$$

Having computed the sums over $f \in H_k(1)$, we see that

$$\Sigma = \sum_{\substack{n \text{ prime} \\ n/N \in E}} \log n \sum_{\substack{k \equiv 2\delta \pmod{4} \\ |k-K| \leq H}} \left(-n^{\frac{1-k}{2}} + \frac{(-1)^\delta}{\pi} \sum_{\substack{t \in \mathbb{Z} \\ t^2 < 4n}} \cos((k-1)\phi_{t,n}) L(1, \psi_{t^2-4n}) \right).$$

As $k \rightarrow \infty$, the contribution from $n^{\frac{1-k}{2}}$ is negligible, so we are left with the sum

$$\Sigma \approx \frac{(-1)^\delta}{\pi} \sum_{\substack{n \text{ prime} \\ n/N \in E}} \log n \sum_{\substack{k \equiv 2\delta \pmod{4} \\ |k-K| \leq H}} \sum_{\substack{t \in \mathbb{Z} \\ t^2 < 4n}} \cos((k-1)\phi_{t,n}) L(1, \psi_{t^2-4n}).$$

We will shortly bring the sum over k inside, but even before applying the trace formula, we take a partition of unity to split the sum over k into smooth weighted sums over shorter ranges of length $\asymp h$. For a fixed choice of smooth function $W : \mathbb{R} \rightarrow \mathbb{R}_{\geq 0}$ supported on $[-1, 1]$ and parameters h, J and K_0 (depending on H and K), we find (assuming GRH for $L(s, f)$) that

$$\Sigma = \sum_{\substack{n \text{ prime} \\ n/N \in E}} \log n \sum_{j=0}^{J-1} \sum_{k \in 2\delta + 4\mathbb{Z}} W\left(\frac{k - (K_0 + 4jh)}{4h}\right) \sum_{f \in H_k(1)} \lambda_f(n) + O_{E,\varepsilon}(hK^{2+\varepsilon}).$$

(The denominator on the left-hand side of (1-1) is of size $\asymp HK^3$, so after dividing by $\sqrt{N} \asymp K$ we will need an error term that is $o(HK^2)$ for the claimed theorem; the size of h depends on H , but it will be always be at most $H^{1-\epsilon}$. See the beginning of Section 3.)

Upon reordering the summation and applying the trace formula, this becomes

$$\Sigma = \frac{(-1)^\delta}{\pi} \sum_{j=0}^{J-1} \sum_{\substack{n \text{ prime} \\ n/N \in E}} \log n \sum_{\substack{t \in \mathbb{Z} \\ t^2 < 4n}} L(1, \psi_{t^2-4n}) \sum_{k \in 2\delta + 4\mathbb{Z}} W\left(\frac{k - (K_0 + 4jh)}{4h}\right) \cos((k-1)\phi_{t,n}) + O_{E,\varepsilon}(hK^{2+\varepsilon}).$$

Since k runs over a fixed residue class mod 4, the inner sum is effectively concentrated around values of $\phi_{t,n}$ which are close to 0 or $\pm \frac{\pi}{2}$. However, as there are relatively few integers t with t^2 close to $4n$, the terms corresponding to $\phi_{t,n} \approx \pm \frac{\pi}{2}$ do not contribute to leading order. We see this when we apply the Poisson summation formula to the innermost sum. We will have

$$\sum_{k \in 2\delta + 4\mathbb{Z}} W\left(\frac{k - k_0}{4h}\right) \cos((k-1)\phi_{t,n}) = h \cos((k_0-1)\phi_{t,n}) \sum_{\ell \in \mathbb{Z}} \widehat{W}(h\ell + 2h\phi_{t,n}/\pi),$$

and it transpires that only the term $\ell = 0$ is large enough to overcome our error terms. (See (7) after Theorem 1.1 about the terms with $\ell = \pm 1$, however.) Specifically, after Lemma 3.1 we find that

$$\Sigma = \frac{(-1)^\delta h}{\pi} \sum_{\substack{k_0 \equiv K_0 \pmod{4h} \\ 0 \leq \frac{k_0 - K_0}{4h} < J}} \sum_{\substack{n \text{ prime} \\ n/N \in E}} \log n \sum_{\substack{t \in \mathbb{Z} \\ t^2 < 4n}} L(1, \psi_{t^2-4n}) \cos((k_0 - 1)\phi_{t,n}) \widehat{W}(2h\phi_{t,n}/\pi) + O_{E,\varepsilon} \left(hK^{2+\varepsilon} + \frac{HK^3 \log K}{h^2} \right).$$

Due to the rapid decay of \widehat{W} we can now restrict the sum over t to a smaller range and put in a linear approximation for $\phi_{t,n}$, and at the end of Section 3 we arrive at

$$\Sigma = \frac{(-1)^\delta h}{\pi} \sum_{k_0} \sum_{\substack{n \text{ prime} \\ n/N \in E}} \log n \sum_{\substack{t \in \mathbb{Z} \\ |t| \leq T}} L(1, \psi_{t^2-4n}) \cos\left(\frac{(k_0 - 1)t}{2\sqrt{n}}\right) \widehat{W}\left(\frac{ht}{\pi\sqrt{n}}\right) + O_{E,\varepsilon,\varepsilon_0} \left(hK^{2+\varepsilon} + \frac{HK^3 \log K}{h^2} \right)$$

for $T = K^{1+\varepsilon_0}/h$. (Note that as our interval does not contain 0, we always have $n \gg K^2$, so that this summation over t really is a truncation.)

The next step, and the content of Section 4, is to take the sum over primes inside and apply the prime number theorem in arithmetic progressions to compute the sum over primes for each fixed t . In the process, $L(1, \psi_{t^2-4n})$ gets replaced by $L(1, \tilde{\psi}_t)$, where $\tilde{\psi}_t$ is the local average

$$\tilde{\psi}_t(m) = \frac{1}{\varphi(m^2)} \sum_{\substack{n \pmod{m^2} \\ (n,m)=1}} \psi_{t^2-4n}(m).$$

This is the content of Lemma 4.5, which, roughly, states that for any nice enough continuous function Φ , we have

$$\sum_{\substack{n \in [A, B] \\ n \text{ prime}}} L(1, \psi_{t^2-4n}) \Phi(n) \log n \approx L(1, \tilde{\psi}_t) \int_A^B \Phi(u) du.$$

Applying this lemma (after which we will be able to extend the range of summation over t to all integers) and making a change of variables in the integration we find for the interval $E = [\alpha_2^{-2}, \alpha_1^{-2}]$ that

$$\Sigma = \frac{2h(-1)^\delta}{\pi} \sum_{k_0} \left(\frac{k_0 - 1}{4\pi}\right)^2 \int_{\lambda_{k_0}\alpha_1}^{\lambda_{k_0}\alpha_2} \sum_{t \in \mathbb{Z}} L(1, \tilde{\psi}_t) \cos(2\pi\alpha t) \widehat{W}\left(\frac{t}{x_{k_0}(\alpha)}\right) \frac{d\alpha}{\alpha^3} + O_{E,\varepsilon} \left(hK^{2+\varepsilon} + \frac{HK^{3+\varepsilon}}{h^{\frac{6}{5}}} \right),$$

where $\lambda_{k_0} = (k_0 - 1)/(4\pi\sqrt{N})$ and $x_{k_0}(\alpha) = (k_0 - 1)/(4\alpha h)$.

The value $L(1, \tilde{\psi}_t)$ turns out to be a constant times a multiplicative function (this is part of Lemma 4.3), so the sum over t can be (approximately) computed by studying additive twists of the Dirichlet series $\sum_{t=1}^\infty L(1, \tilde{\psi}_t) t^{-s}$, and we find that the mass of the integrand is concentrated around rational numbers with squarefree denominators. Specifically, in Proposition 5.1 we find that for $\alpha = a/q + \theta$ (with θ small),

$$\sum_{t \in \mathbb{Z}} L(1, \tilde{\psi}_t) \cos(2\pi\alpha) \widehat{W}\left(\frac{t}{x}\right) \approx \frac{\mu(q)^2}{\varphi(q)^2 \sigma(q)} x W(x\theta).$$

It is now natural to employ the circle method, computing the contribution to the integral over $[\lambda_{k_0}\alpha_1, \lambda_{k_0}\alpha_2]$ which comes from major arcs near rational numbers with small denominators, and bounding the rest. Noticing that $\lambda_{k_0} \approx 1$, we find that the bulk of the integral comes from rational numbers in the range $[\alpha_1, \alpha_2]$. The main term will come from a sum of the integrals

$$\int x_{k_0}(\alpha)W(x_{k_0}(\alpha)(\alpha - a/q))\frac{d\alpha}{\alpha^3} \approx \left(\frac{q}{a}\right)^3.$$

Finally we can see our answer will emerge as

$$\Sigma \approx (-1)^\delta \frac{2h}{\pi} \sum_{k_0} \left(\frac{k_0 - 1}{4\pi}\right)^2 \sum_{\alpha_1 < a/q < \alpha_2} \frac{\mu(q)}{\varphi(q)^2 \sigma(q)} \left(\frac{q}{a}\right)^3 \approx (-1)^\delta \frac{HK^2}{16\pi^3} \sum_{\alpha_1 < a/q < \alpha_2} \frac{\mu(q)}{\varphi(q)^2 \sigma(q)} \left(\frac{q}{a}\right)^3,$$

though some care must be taken with the endpoints. (See also Remark (5) following Theorem 1.1.) The denominator is easily computed using dimension formulas and the prime number theorem (see (6-3)), giving Theorem 1.1.

3. The sum over weights

Let $W : \mathbb{R} \rightarrow \mathbb{R}_{\geq 0}$ be a smooth, even function supported on $[-1, 1]$ and satisfying

$$W(x) + W(1 - x) = 1 \quad \text{for } x \in [0, 1].$$

Note that these assumptions imply that $0 \leq W(x) \leq 1$ and $\int_{\mathbb{R}} W(x) dx = 1$. For instance, we may take

$$W(x) = \begin{cases} c \int_{-1}^{1-2|x|} e^{-\frac{1}{1-t^2}} dt & \text{if } |x| < 1, \\ 0 & \text{otherwise,} \end{cases} \quad \text{where } c = \left(\int_{-1}^1 e^{-\frac{1}{1-t^2}} dt\right)^{-1} = \frac{e^{\frac{1}{2}}}{K_1(\frac{1}{2}) - K_0(\frac{1}{2})}.$$

(Here $K_\nu(y)$ is the K -Bessel function.)

Define

$$h = \lceil \max(H^{\frac{10}{9}} K^{-\frac{1}{9}}, (HK)^{\frac{5}{11}}) \rceil$$

and set

$$K_0 = \min\{k \in \mathbb{Z} : k \geq K - H + 4h, k \equiv 2\delta \pmod{4}\}, \quad J = \left\lfloor \frac{K + H - K_0}{4h} \right\rfloor.$$

We assume K is sufficiently large to ensure that $4h + 2 \leq H \leq K - 2$, which implies that $K_0 \geq 4h + 2$ and $J > 0$. Write

$$\mathbf{1}_{[K-H, K+H]}(k) = \sum_{j=0}^{J-1} W\left(\frac{k - (K_0 + 4jh)}{4h}\right) + R(k). \tag{3-1}$$

Then R is supported on $[K - H, K_0] \cup [K_0 + 4(J - 1)h, K + H]$.

Assuming GRH for $L(s, f)$, the sum over primes for a single newform satisfies the bound

$$\sum_{\substack{p \text{ prime} \\ p/N \in E}} \lambda_f(p) \log p \ll_{E, \varepsilon} K^{1+\varepsilon}.$$

Combining this with the estimates $\#H_k(1) \asymp k$ and $R(k) \ll 1$, we find that

$$\sum_{\substack{k \equiv 2\delta \pmod{4} \\ |k-K| \leq H}} R(k) \sum_{f \in H_k(1)} \sum_{\substack{p \text{ prime} \\ p/N \in E}} \lambda_f(p) \log p \ll_{E,\varepsilon} hK^{2+\varepsilon}, \tag{3-2}$$

as the support of $R(k)$ is of size $O(h)$.

Next we consider the contribution from a typical term of (3-1) to (2-1), with $k_0 = K_0 + 4jh$:

$$\sum_{k \in k_0 + 4\mathbb{Z}} \cos((k-1)\phi) W\left(\frac{k-k_0}{4h}\right) = \Re \sum_{m \in \mathbb{Z}} W(m/h) e^{i(k_0-1+4m)\phi}.$$

To that end, we have

$$\begin{aligned} \int_{\mathbb{R}} W(x/h) e^{i(k_0-1+4x)\phi} e^{2\pi ixt} dx &= \int_{\mathbb{R}} W(x/h) e^{i(k_0-1)\phi + 2\pi ix(t+2\phi/\pi)} dx \\ &= h \int_{\mathbb{R}} W(x) e^{i(k_0-1)\phi + 2\pi ix(ht+2h\phi/\pi)} dx \\ &= h e^{i(k_0-1)\phi} \widehat{W}(ht + 2h\phi/\pi), \end{aligned}$$

and we note that \widehat{W} is real as W is even. By Poisson summation, we obtain

$$\sum_{k \in k_0 + 4\mathbb{Z}} \cos((k-1)\phi) W\left(\frac{k-k_0}{4h}\right) = h \cos((k_0-1)\phi) \sum_{\ell \in \mathbb{Z}} \widehat{W}(h\ell + 2h\phi/\pi).$$

Applying this to (2-1) with n prime and $k_0 \geq 4h + 2$, we have

$$\begin{aligned} \sum_{k \in k_0 + 4\mathbb{Z}} W\left(\frac{k-k_0}{4h}\right) n^{\frac{1-k}{2}} \text{Tr } T_n |S_k(1) &= - \sum_{k \in k_0 + 4\mathbb{Z}} n^{\frac{1-k}{2}} W\left(\frac{k-k_0}{4h}\right) \\ &\quad + \frac{(-1)^{\frac{k_0}{2}} h}{\pi} \sum_{\ell \in \mathbb{Z}} \sum_{\substack{t \in \mathbb{Z} \\ t^2 < 4n}} \widehat{W}\left(h\left(\ell + \frac{2\phi_{t,n}}{\pi}\right)\right) \cos((k_0-1)\phi_{t,n}) L(1, \psi_{t^2-4n}). \end{aligned} \tag{3-3}$$

Since W is smooth, only the $\ell = 0$ term contributes significantly to (3-3), as quantified by the next result.

Lemma 3.1. *We have*

$$\begin{aligned} \sum_{\substack{n \text{ prime} \\ \frac{n}{N} \in E}} (\log n) &\left[\frac{(-1)^{\frac{k_0}{2}} h}{\pi} \sum_{\ell \in \mathbb{Z} \setminus \{0\}} \sum_{\substack{t \in \mathbb{Z} \\ t^2 < 4n}} \cos((k_0-1)\phi_{t,n}) L(1, \psi_{t^2-4n}) \widehat{W}\left(h\left(\ell + \frac{2\phi_{t,n}}{\pi}\right)\right) \right. \\ &\quad \left. - \sum_{k \in k_0 + 4\mathbb{Z}} n^{\frac{1-k}{2}} W\left(\frac{k-k_0}{4h}\right) \right] \\ &\ll_E \frac{K^3 \log K}{h}. \end{aligned}$$

Proof. First note that

$$\sum_{\substack{n \text{ prime} \\ \frac{n}{N} \in E}} \log n \sum_{k \in k_0 + 4\mathbb{Z}} n^{\frac{1-k}{2}} W\left(\frac{k-k_0}{4h}\right) \ll \sum_{n \text{ prime}} \log n \sum_{k=4}^{\infty} n^{\frac{1-k}{2}} \ll 1.$$

Next, since ψ_D is periodic modulo D with mean value 0, partial summation gives the estimate

$$L(1, \psi_D) = \sum_{m=1}^{|D|} \frac{\psi_D(m)}{m} + \int_{|D|}^{\infty} x^{-2} \sum_{m \leq x} \psi_D(m) dx \ll \sum_{m=1}^{|D|} \frac{1}{m} + \int_{|D|}^{\infty} \frac{|D|}{x^2} dx \ll \log |D|.$$

Thus,

$$\sum_{\substack{t \in \mathbb{Z} \\ t^2 < 4n}} \cos((k_0 - 1)\phi_{t,n}) L(1, \psi_{t^2-4n}) \ll \sqrt{n} \log n.$$

Since W is smooth, we may apply integration by parts to $\widehat{W}(x) = \int_{\mathbb{R}} W(u) e^{-2\pi i x u} du$ to obtain

$$\widehat{W}(x) \ll_A |x|^{-A} \quad \text{for all } A \in \mathbb{Z}_{\geq 0}.$$

Since $\phi_{t,n} \in (-\frac{\pi}{2}, \frac{\pi}{2})$, we have $|\ell + 2\phi_{t,n}/\pi| > |\ell| - 1$, so that

$$\begin{aligned} \sum_{\substack{n \text{ prime} \\ n/N \in E}} \frac{h \log n}{\pi} \sum_{\substack{\ell \in \mathbb{Z} \\ |\ell| > 1}} \sum_{\substack{t \in \mathbb{Z} \\ t^2 < 4n}} \widehat{W}\left(h \left(\ell + \frac{2\phi_{t,n}}{\pi}\right)\right) \cos((k_0 - 1)\phi_{t,n}) L(1, \psi_{t^2-4n}) \\ \ll h \sum_{\substack{n \text{ prime} \\ n/N \in E}} \sqrt{n} \log^2 n \sum_{\ell=2}^{\infty} (h(\ell - 1))^{-2} \ll_E \frac{K^3 \log K}{h}. \end{aligned}$$

It remains to estimate the contributions from $\ell = \pm 1$. Since \widehat{W} is even, those terms contribute equally, so it suffices to consider $\ell = -1$. For each fixed t , we prove different bounds based on the relative sizes of t and n . When $\ell = -1$ and $t < \sqrt{n}$, $|\ell + 2\phi_{t,n}/\pi| \gg 1$, and a similar argument as above gives $O_E(K^3(\log K)/h)$.

When $t^2/(2(1 - h^{-2}))^2 < n \leq t^2$ we use

$$\frac{\pi}{2} - \phi_{t,n} = \arcsin \sqrt{1 - \frac{t^2}{4n}} \geq \sqrt{1 - \frac{t}{2\sqrt{n}}}$$

and the estimate $\widehat{W}(x) \ll_A |x|^{-A}$ to get the bound

$$\begin{aligned} &\ll_{E,A} h \log^2 K \sum_{t \asymp_E K} \int_{(t/2)^2(1-h^{-2})^{-2}}^{t^2} h^{-A} \left(1 - \frac{t}{2\sqrt{n}}\right)^{-A/2} d\pi(n) \\ &= h^{1-A} \log^2 K \sum_{t \asymp_E K} \int_{L_t}^{t^2} R_{A,t}(n) d\pi(n) \end{aligned}$$

where $L_t = (t/2)^2(1 - h^{-2})^{-2}$ and $R_{A,t}(n) = (1 - t/(2\sqrt{n}))^{-A/2}$. Applying integration by parts,

$$\begin{aligned} \int_{L_t}^{t^2} R_{A,t}(n) d\pi(n) &= (\pi(n) - \pi(t^2/4)) R_{A,t}(n) \Big|_{L_t}^{t^2} - \int_{L_t}^{t^2} (\pi(n) - \pi(t^2/4)) R'_{A,t}(n) dn \\ &\leq 2^{A/2} \pi(t^2) - \int_{L_t}^{t^2} (\pi(n) - \pi(t^2/4)) R'_{A,t}(n) dn. \end{aligned}$$

Since $L_t - t^2/4 \gg_E (K/h)^2$, [MV07, §3.2, Corollary 4] implies that $\pi(n) - \pi(t^2/4) \ll_E (n - t^2/4)/\log K$ for $n \geq L_t$. Hence, for $A \geq 3$ we have

$$\begin{aligned} \int_{L_t}^{t^2} R_{A,t}(n) d\pi(n) &\ll_{E,A} \frac{t^2}{\log K} - \frac{1}{\log K} \int_{L_t}^{t^2} (n - t^2/4) R'_{A,t}(n) dn \\ &= \frac{t^2}{\log K} - \frac{1}{\log K} \left((n - t^2/4) R_{A,t}(n) \Big|_{L_t}^{t^2} - \int_{L_t}^{t^2} R_{A,t}(n) dn \right) \\ &\leq \frac{t^2 + (L_t - t^2/4) R_{A,t}(L_t)}{\log K} + \frac{1}{\log K} \int_{L_t}^{t^2} R_{A,t}(n) dn \\ &\ll_A \frac{t^2 h^{A-2}}{\log K}. \end{aligned}$$

Summing over $t \asymp_E K$, this again gives $O_E\left(\frac{K^3 \log K}{h}\right)$.

Finally, we estimate the terms with

$$\left(\frac{t}{2}\right)^2 < n \leq \left(\frac{t}{2(1-h^{-2})}\right)^2$$

using the trivial bound $\widehat{W}(x) \ll 1$. Summing over $t \asymp_E K$, these make a total contribution

$$\ll_E h \log^2 K \sum_{t \asymp_E K} \left[\pi\left(\left(\frac{t}{2(1-h^{-2})}\right)^2\right) - \pi\left(\left(\frac{t}{2}\right)^2\right) \right].$$

Note that $\left(\frac{t}{2(1-h^{-2})}\right)^2 = \left(\frac{t}{2}\right)^2 + O_E(K^2/h^2)$. Applying [MV07, §3.2, Corollary 4] again, it follows that

$$\pi\left(\left(\frac{t}{2(1-h^{-2})}\right)^2\right) - \pi\left(\left(\frac{t}{2}\right)^2\right) \ll_E \frac{K^2}{h^2 \log K},$$

so in total we get $\ll_E (K^3 \log K)/h$. \square

Summing the bound from Lemma 3.1 over the $J \ll H/h$ choices of k_0 , and combining with the boundary error (3-2) from the partition of unity, we get that the total contribution from everything except the $\ell = 0$ term is bounded by

$$\ll_{E,\varepsilon} h K^{2+\varepsilon} + \frac{H K^3 \log K}{h^2}.$$

Next we fix a small $\varepsilon_0 > 0$ and restrict the sum over t to $|t| \leq T = K^{1+\varepsilon_0}/h$. The error in so doing is $O_{E,\varepsilon_0}(1)$, thanks to the rapid decay of \widehat{W} . Hence, we have

$$\begin{aligned} \Sigma &= O_{E,\varepsilon_0,\varepsilon} \left(h K^{2+\varepsilon} + \frac{H K^3 \log K}{h^2} \right) \\ &\quad + \frac{(-1)^\delta h}{\pi} \sum_{\substack{k_0 \equiv K_0 \pmod{4h} \\ 0 \leq \frac{k_0 - K_0}{4h} < J}} \sum_{\substack{n \text{ prime} \\ n/N \in E}} \log n \sum_{\substack{t \in \mathbb{Z} \\ |t| \leq T}} L(1, \psi_{t^2-4n}) \cos((k_0 - 1)\phi_{t,n}) \widehat{W}\left(h \frac{2\phi_{t,n}}{\pi}\right). \end{aligned}$$

Writing $A_{k_0}(\phi) = \cos((k_0 - 1)\phi) \widehat{W}(2h\phi/\pi)$, we have $A'_{k_0}(\phi) \ll K$ uniformly, and thus

$$A_{k_0}(\phi_{t,n}) = A_{k_0}\left(\frac{t}{2\sqrt{n}}\right) + O_E\left(\frac{T^3}{K^2}\right),$$

since

$$\phi_{t,n} = \frac{t}{2\sqrt{n}} + O\left(\frac{|t|^3}{n^{\frac{3}{2}}}\right).$$

Summing this error over $|t| \leq T$, $n \asymp_E K^2$, and k_0 , we get

$$\ll_E HT^4 \log K = \frac{HK^{4+4\varepsilon_0} \log K}{h^4},$$

which is dominated by the above error term for ε_0 sufficiently small. Thus, we have

$$\begin{aligned} \Sigma = O_{E,\varepsilon_0,\varepsilon} & \left(hK^{2+\varepsilon} + \frac{HK^3 \log K}{h^2} \right) \\ & + \frac{(-1)^{\delta h}}{\pi} \sum_{\substack{k_0 \equiv K_0 \pmod{4h} \\ 0 \leq \frac{k_0 - K_0}{4h} < J}} \sum_{\substack{t \in \mathbb{Z} \\ |t| \leq T}} \sum_{\substack{n \text{ prime} \\ n/N \in E}} (\log n) L(1, \psi_{t^2-4n}) \cos \frac{(k_0 - 1)t}{2\sqrt{n}} \widehat{W}\left(\frac{ht}{\pi\sqrt{n}}\right). \end{aligned} \quad (3-4)$$

4. The sum over primes

Next we turn our attention to the sum over primes n . We begin with some lemmas.

Lemma 4.1. *Let m be a positive integer, and let D_1, D_2 be discriminants with $D_1 \equiv D_2 \pmod{(2m)^2}$. Then $\psi_{D_1}(m) = \psi_{D_2}(m)$.*

Proof. Write $D_i = d_i \ell_i^2$ where d_i is a fundamental discriminant and $\ell_i \in \mathbb{Z}_{\geq 0}$. Without loss of generality, we assume that $\text{ord}_2(\ell_1) \leq \text{ord}_2(\ell_2)$. Let $r = \gcd(\ell_1, m)$. Since r^2 divides both D_1 and m^2 , it also divides D_2 . Moreover, since $d_2/\gcd(d_2, 4)$ is squarefree, we have $r \mid \ell_2$. Since $\text{ord}_2(\ell_1) \leq \text{ord}_2(\ell_2)$, it follows that $r \mid \ell_2$.

Suppose p is a prime dividing $\gcd(\ell_2/r, m/r)$. Then $p \nmid (\ell_1/r)$ since $\gcd(\ell_1/r, m/r) = 1$. From $d_1(\ell_1/r)^2 \equiv d_2(\ell_2/r)^2 \pmod{(2m/r)^2}$ it follows that $p^2 \mid d_1$, whence $p = 2$. Dividing by 4, we find that $(d_1/4)(\ell_1/r)^2 \equiv d_2(\ell_2/(2r))^2 \pmod{4}$, which is impossible since $(d_1/4)(\ell_1/r)^2 \equiv 2$ or $3 \pmod{4}$ and $d_2(\ell_2/(2r))^2 \equiv 0$ or $1 \pmod{4}$.

Hence $r = \gcd(\ell_2, m)$, and

$$\psi_{D_1}(m) = \left(\frac{d_1}{m/r}\right) = \left(\frac{d_1(\ell_1/r)^2}{m/r}\right) = \left(\frac{d_2(\ell_2/r)^2}{m/r}\right) = \left(\frac{d_2}{m/r}\right) = \psi_{D_2}(m). \quad \square$$

Lemma 4.2. *For $t, m \in \mathbb{Z}$ with $m > 0$, define*

$$\tilde{\psi}_t(m) = \frac{1}{\varphi(m^2)} \sum_{\substack{n \pmod{m^2} \\ (n,m)=1}} \psi_{t^2-4n}(m).$$

Then for fixed t , $m \mapsto \tilde{\psi}_t(m)$ is a multiplicative function of m .

Proof. Fix $t \in \mathbb{Z}$, and consider $m_1, m_2 \in \mathbb{Z}_{>0}$ with $\gcd(m_1, m_2) = 1$. For $n \in \mathbb{Z}$, we write $t^2 - 4n = d\ell^2$ where d is a fundamental discriminant and $\ell \in \mathbb{Z}_{\geq 0}$. Let $r_1 = \gcd(m_1, \ell)$ and $r_2 = \gcd(m_2, \ell)$. Since $\gcd(m_1, m_2) = 1$, we have $\gcd(r_1, r_2) = 1$ and $\gcd(m_1 m_2, \ell) = r_1 r_2$. Therefore,

$$\psi_{t^2-4n}(m_1 m_2) = \left(\frac{d}{\frac{m_1}{r_1} \frac{m_2}{r_2}} \right) = \left(\frac{d}{m_1/r_1} \right) \left(\frac{d}{m_2/r_2} \right) = \psi_{t^2-4n}(m_1) \psi_{t^2-4n}(m_2),$$

so that

$$\begin{aligned} \tilde{\psi}_t(m_1 m_2) &= \frac{1}{\varphi(m_1 m_2)} \sum_{\substack{n \bmod (m_1 m_2)^2 \\ (n, m_1 m_2) = 1}} \psi_{t^2-4n}(m_1) \psi_{t^2-4n}(m_2) \\ &= \frac{1}{\varphi(m_1 m_2)} \sum_{\substack{n_1 \bmod m_1^2 \\ (n_1, m_1) = 1}} \sum_{\substack{n_2 \bmod m_2^2 \\ (n_2, m_2) = 1}} \psi_{t^2-4(n_1(m_2 \bar{m}_2)^2 + n_2(m_1 \bar{m}_1)^2)}(m_1) \psi_{t^2-4(n_1(m_2 \bar{m}_2)^2 + n_2(m_1 \bar{m}_1)^2)}(m_2), \end{aligned}$$

where $m_1 \bar{m}_1 \equiv 1 \pmod{m_1^2}$ and $m_2 \bar{m}_2 \equiv 1 \pmod{m_2^2}$. By Lemma 4.1 we have

$$\psi_{t^2-4(n_1(m_2 \bar{m}_2)^2 + n_2(m_1 \bar{m}_1)^2)}(m_i) = \psi_{t^2-4n_i}(m_i),$$

so we get

$$\tilde{\psi}_t(m_1 m_2) = \tilde{\psi}_t(m_1) \tilde{\psi}_t(m_2). \quad \square$$

Lemma 4.3. For $t \in \mathbb{Z}$, define

$$L(s, \tilde{\psi}_t) = \sum_{m=1}^{\infty} \frac{\tilde{\psi}_t(m)}{m^s} \quad \text{for } \Re(s) > 1.$$

Then

$$L(s, \tilde{\psi}_t) = L(s, \tilde{\psi}_1) P(s, t),$$

where

$$L(s, \tilde{\psi}_1) = \zeta(2s) \zeta(s+2) \prod_p \begin{cases} 1 - (1+p^{-1})(1+p^{-s})p^{-s-1}, & p > 2, \\ (1-2^{-s})(1-2^{-s-2}), & p = 2, \end{cases} \quad (4-1)$$

and

$$P(s, t) = \prod_{p|t} \begin{cases} \frac{1-p^{-s-2}}{1-(1+p^{-1})(1+p^{-s})p^{-s-1}}, & p > 2, \\ \frac{1+2^{-s-1}-2^{-2s}}{1-2^{-s}}, & p = 2, 4|t, \\ \frac{1+2^{-s-2}-7 \cdot 2^{-2s-3}-2^{-3s-2}}{(1-2^{-s})(1-2^{-s-2})}, & p = 2, 4 \nmid t. \end{cases} \quad (4-2)$$

Thus $L(s, \tilde{\psi}_t)$ continues analytically to $\Re(s) > \frac{1}{2}$ and satisfies

$$L(1, \tilde{\psi}_t) = Cf(t),$$

where

$$C = L(1, \tilde{\psi}_1) = \prod_p \left(1 - \frac{1}{(p-1)^2(p+1)} \right) = 0.6151326573181718 \dots$$

and

$$f(t) = P(1, t) = \prod_{p|t} \left(1 + \frac{1}{p^2 - p - 1} \right). \quad (4-3)$$

In particular,

$$L(1, \tilde{\psi}_0) = Cf(0) = \prod_p \left(1 - \frac{1}{(p-1)^2(p+1)} \right) \left(1 + \frac{1}{p^2 - p - 1} \right) = \prod_p \left(1 - \frac{1}{p^2} \right)^{-1} = \zeta(2).$$

Proof. Fix $t \in \mathbb{Z}$. By Lemma 4.2, $\tilde{\psi}_t$ is multiplicative, so we have $L(s, \tilde{\psi}_t) = \prod_p \sum_{\ell=0}^{\infty} \tilde{\psi}_t(p^\ell) p^{-\ell s}$.

Consider $m = p^e$ for some $e > 0$, and put

$$S = \sum_{\substack{n \bmod p^{2e} \\ (n, p)=1}} \psi_{t^2-4n}(p^e).$$

Suppose first that $p \nmid 2t$. Given a coprime residue $n \bmod p^{2e}$, we put $j = \text{ord}_p(t^2 - 4n)$. The term $n \equiv (t/2)^2 \bmod p^{2e}$ corresponds to $j \geq 2e$ and contributes $\psi_{t^2-4n}(m) = 1$ to S .

Consider $j \in (0, 2e)$. Then n is of the form $(t/2)^2 - (a + bp)p^j$, where $a \in \{1, \dots, p-1\}$ and $b \in \{0, \dots, p^{2e-j-1} - 1\}$. We have

$$\psi_{t^2-4n}(m) = \begin{cases} \left(\frac{a}{p^{e-j/2}} \right), & j \text{ even,} \\ 0, & j \text{ odd.} \end{cases}$$

From even $j = 2e - 2r$ we get a contribution of

$$\sum_{a=1}^{p-1} \sum_{b=0}^{p^{2r-1}-1} \left(\frac{a}{p^r} \right) = \begin{cases} \varphi(p^{2r}), & r \text{ even,} \\ 0, & r \text{ odd.} \end{cases}$$

Now consider $j = 0$. As before we have $n = (t/2)^2 - (a + bp)$ with $a \in \{1, \dots, p-1\}$ and $b \in \{0, \dots, p^{2e-1} - 1\}$, but now we have the additional constraint $a \not\equiv (t/2)^2 \bmod p$, since n has to be coprime to p . We have $\psi_{t^2-4n}(m) = (a/p^e)$, so the contribution from $j = 0$ is

$$\sum_{\substack{1 \leq a \leq p-1 \\ a \not\equiv (t/2)^2 \bmod p}} \sum_{b=0}^{p^{2e-1}-1} \left(\frac{a}{p^e} \right) = \begin{cases} (p-2)p^{2e-1}, & e \text{ even,} \\ -p^{2e-1}, & e \text{ odd.} \end{cases}$$

Altogether, for $p \nmid 2t$ and $e > 0$ we have

$$S = \sum_{\substack{n \bmod p^{2e} \\ (n, p)=1}} \psi_{t^2-4n}(p^e) = -p^{2e-1} + \sum_{k=0}^{\lfloor e/2 \rfloor} \varphi(p^{4k}),$$

so that

$$\begin{aligned} \sum_{e=0}^{\infty} \frac{\tilde{\psi}_t(p^e)}{p^{es}} &= - \sum_{e=1}^{\infty} \frac{p^{2e-1}}{p^{es}\varphi(p^{2e})} + \sum_{k=0}^{\infty} \sum_{e=2k}^{\infty} \frac{\varphi(p^{4k})}{p^{es}\varphi(p^{2e})} \\ &= \frac{1 - (1 + p^{-1})(1 + p^{-s})p^{-s-1}}{(1 - p^{-2s})(1 - p^{-s-2})}. \end{aligned}$$

Next suppose that $p = 2$ and t is odd. Then $j = 0$, so we get

$$S = \sum_{\substack{n \bmod 2^{2e} \\ (n,2)=1}} \left(\frac{t^2 - 4n}{2^e} \right) = (-1)^e 2^{2e-1}.$$

Thus,

$$\sum_{e=0}^{\infty} \frac{\tilde{\psi}_t(2^e)}{2^{es}} = 1 + \sum_{e=1}^{\infty} \frac{(-1)^e 2^{2e-1}}{2^{es}\varphi(2^{2e})} = \frac{1}{1 + 2^{-s}}.$$

Hence,

$$L(s, \tilde{\psi}_1) = \frac{1}{1 + 2^{-s}} \prod_{p>2} \frac{1 - (1 + p^{-1})(1 + p^{-s})p^{-s-1}}{(1 - p^{-2s})(1 - p^{-s-2})},$$

which yields (4-1).

Now suppose that $p|t$. When $p > 2$ we have $j = 0$, so that $\tilde{\psi}_t(p^e) = (1 + (-1)^e)/2$, and the local factor at p becomes $1/(1 - p^{-2s})$. When $p = 2$ and $4|t$, we have $\tilde{\psi}_t(2^e) = (1 - (-1)^e)/4$ for $e > 0$, so that

$$\sum_{e=0}^{\infty} \tilde{\psi}_t(2^e) 2^{-es} = 1 + \frac{1}{4} \sum_{e=1}^{\infty} (1 - (-1)^e) 2^{-es} = \frac{1 + 2^{-1-s} - 2^{-2s}}{1 - 2^{-2s}}.$$

When $2|t$ but $4 \nmid t$, we have $\tilde{\psi}_t(2^e) = 2^{1-2e} (1 + \frac{16^{\lfloor e/2 \rfloor} - 1}{15})$ for $e > 0$, so that

$$\begin{aligned} \sum_{e=0}^{\infty} \tilde{\psi}_t(2^e) 2^{-es} &= 1 + \frac{14 \cdot 2}{15} \sum_{e=1}^{\infty} 2^{-2e} 2^{-es} + \frac{2}{15} \sum_{e=1}^{\infty} 2^{-2e} 2^{4\lfloor \frac{e}{2} \rfloor} 2^{-es} \\ &= \frac{1 + 2^{-s-2} - 7 \cdot 2^{-2s-3} - 2^{-3s-2}}{(1 - 2^{-s-2})(1 - 2^{-2s})}. \end{aligned}$$

Collecting these cases, we arrive at (4-2). □

Lemma 4.4. *Assume the generalized Lindelöf hypothesis (GLH) for quadratic Dirichlet L-functions. Then for any non-square discriminant D and any $x \geq 1$,*

$$L(1, \psi_D) = \sum_{m \leq x} \frac{\psi_D(m)}{m} + O_{\varepsilon} \left(\frac{|Dx|^{\varepsilon}}{\sqrt{x}} \right).$$

Proof. Let $D = d\ell^2$ with d a fundamental discriminant and $\ell \in \mathbb{Z}_{>0}$. Then we have

$$L(s, \psi_D) := \sum_{m=1}^{\infty} \frac{\psi_D(m)}{m^s} = L(s, \psi_d) \prod_{p|\ell} \left(1 + (1 - \psi_d(p)) \sum_{j=1}^{\text{ord}_p \ell} p^{-js} \right).$$

For $\Re(s) \geq \frac{1}{2}$ we have

$$\left| 1 + (1 - \psi_d(p)) \sum_{j=1}^{\text{ord}_p \ell} p^{-js} \right| \leq 1 + 2 \sum_{j=1}^{\infty} p^{-j/2} = \frac{\sqrt{p} + 1}{\sqrt{p} - 1},$$

and hence

$$L(s, \psi_D) \ll_{\varepsilon} |L(s, \psi_d)| \ell^{\varepsilon} \ll_{\varepsilon} |Ds|^{\varepsilon},$$

under GLH. By Perron's formula, for $x \geq 2$,

$$\sum_{n \leq x} \frac{\psi_D(m)}{m} = \frac{1}{2\pi i} \int_{1 + \frac{1}{\log x} - ix}^{1 + \frac{1}{\log x} + ix} L(s, \psi_D) x^{s-1} \frac{ds}{s-1} + O\left(\frac{\log x}{x}\right).$$

The lemma follows on shifting the contour to $\Re(s) = \frac{1}{2}$. □

Lemma 4.5. *Assume GRH for Dirichlet L-functions. Let $t \in \mathbb{Z}$ and $A, B \in \mathbb{R}$ with $\frac{t^2}{4} < A < B$, and let $\Phi \in C^1([A, B])$. Set $M = \max_{u \in [A, B]} |\Phi(u)|$ and $V = \int_A^B |\Phi'(u)| du$. Then*

$$\sum_{\substack{n \in [A, B] \\ n \text{ prime}}} L(1, \psi_{t^2-4n}) \Phi(n) \log n = L(1, \tilde{\psi}_t) \int_A^B \Phi(u) du + O_{\varepsilon}(M^{\frac{4}{5}}(M+V)^{\frac{1}{5}} B^{\frac{9}{10}+\varepsilon}) \quad \forall \varepsilon \in (0, \frac{1}{10}].$$

Proof. The result is trivially true if $B < 2$, so assume otherwise. Put $I = \{n \in (A, B) : n \text{ prime}\}$. (Note that I omits the left endpoint.) For each positive integer m , we have

$$\begin{aligned} \sum_{n \in I} \psi_{t^2-4n}(m) \Phi(n) \log n &= \sum_{\substack{a \bmod m^2 \\ \gcd(a,m)=1}} \sum_{\substack{n \in I \\ n \equiv a \bmod m^2}} \psi_{t^2-4n}(m) \Phi(n) \log n + \sum_{\substack{n \in I \\ n|m}} \psi_{t^2-4n}(m) \Phi(n) \log n \\ &= \sum_{\substack{a \bmod m^2 \\ \gcd(a,m)=1}} \psi_{t^2-4a}(m) \sum_{\substack{n \in I \\ n \equiv a \bmod m^2}} \Phi(n) \log n + \sum_{\substack{n \in I \\ n|m}} \psi_{t^2-4n}(m) \Phi(n) \log n. \end{aligned}$$

Here for the first piece we use Lemma 4.1, which states that $\psi_{t^2-4n}(m) = \psi_{t^2-4a}(m)$ if $a \equiv n \pmod{m^2}$. For the second piece, we observe

$$\left| \sum_{\substack{n \in I \\ n|m}} \psi_{t^2-4n}(m) \Phi(n) \log n \right| \leq M \log m \leq M \varphi(m^2).$$

By [MV07, §13.1, Theorem 8], GRH implies that for $m \geq 1$, $(a, m) = 1$ and $x \geq 2$,

$$\theta(x; m^2, a) := \sum_{\substack{p \leq x \\ p \equiv a \bmod m^2}} \log p = \frac{x}{\varphi(m^2)} + O(x^{\frac{1}{2}} \log^2 x).$$

Hence, writing $E(x; m^2, a) = \theta(x; m^2, a) - x/\varphi(m^2)$, we have

$$\sum_{\substack{n \in I \\ n \equiv a \pmod{m^2}}} \Phi(n) = \int_A^B \Phi(u) d\theta(u; m^2, a) = \int_A^B \frac{\Phi(u)}{\varphi(m^2)} du + \int_A^B \Phi(u) dE(u; m^2, a).$$

Applying integration by parts, the error term is

$$\Phi(u)E(u; m^2, a) \Big|_A^B - \int_A^B E(u; m^2, a)\Phi'(u) du \ll (M + V)B^{\frac{1}{2}} \log^2 B,$$

so that

$$\sum_{n \in I} \psi_{t^2-4n}(m)\Phi(n) \log n = \tilde{\psi}_t(m) \int_A^B \Phi(u) du + O(\varphi(m^2)(M + V)B^{\frac{1}{2}} \log^2 B).$$

By Lemma 4.4,

$$\begin{aligned} \sum_{n \in I} L(1, \psi_{t^2-4n})\Phi(n) \log n &= \sum_{m \leq x} \frac{1}{m} \sum_{n \in I} \psi_{t^2-4n}(m)\Phi(n) \log n + \sum_{n \in I} |\Phi(n) \log n| O_\varepsilon\left(\frac{|(t^2 - 4n)x|^\varepsilon}{\sqrt{x}}\right) \\ &= \sum_{m \leq x} \frac{\tilde{\psi}_t(m)}{m} \int_A^B \Phi(u) du + O\left((M + V)B^{\frac{1}{2}} \log^2 B \sum_{m \leq x} \frac{\varphi(m^2)}{m}\right) + O_\varepsilon(MB^{1+\varepsilon}x^{-\frac{1}{2}+\varepsilon}) \\ &= \sum_{m \leq x} \frac{\tilde{\psi}_t(m)}{m} \int_A^B \Phi(u) du + O(x^2(M + V)B^{\frac{1}{2}} \log^2 B) + O_\varepsilon(MB^{1+\varepsilon}x^{-\frac{1}{2}+\varepsilon}). \end{aligned}$$

From the proof of Lemma 4.3 we may observe that $|\tilde{\psi}_t(p)| \leq \frac{1}{p}$ for primes $p > 2$, which implies the estimate

$$|\tilde{\psi}_t(m)| \leq 2 \prod_{\substack{p|m \\ p^2 \nmid m}} \frac{1}{p}.$$

Given $m > x$, we may write $m = dr$, where d is squarefree, r is squarefull, and $(d, r) = 1$. Thus,

$$\sum_{m > x} \frac{|\tilde{\psi}_t(m)|}{m} \leq 2 \sum_{d=1}^\infty \frac{\mu^2(d)}{d^2} \sum_{\substack{r \text{ squarefull} \\ r > x/d}} \frac{1}{r} \ll \sum_{d=1}^\infty \frac{\min(1, \sqrt{d/x})}{d^2} \ll \frac{1}{\sqrt{x}}.$$

Thus, we have

$$\sum_{n \in I} L(1, \psi_{t^2-4n})\Phi(n) \log n = L(1, \tilde{\psi}_t) \int_A^B \Phi(u) du + O(x^2(M + V)B^{\frac{1}{2}} \log^2 B) + O_\varepsilon(MB^{1+\varepsilon}x^{-\frac{1}{2}+\varepsilon}).$$

Finally, if A is a prime number, we absorb the remaining term $L(1, \psi_{t^2-4A})\Phi(A) \log A$ into the error term. The lemma follows on choosing $x = (B^{\frac{1}{2}}M/(M + V))^{\frac{1}{5}}$. \square

With these results in place, we can now compute the sum over n in (3-4). For fixed k_0 and t , let

$$\Phi_{k_0,t}(u) = \cos \frac{(k_0 - 1)t}{2\sqrt{u}} \widehat{W}\left(\frac{ht}{\pi\sqrt{u}}\right).$$

Then $\Phi_{k_0,t}(u) \ll 1$ and

$$\int_{NE} |\Phi'_{k_0,t}(u)| du \ll_E |t|.$$

Summing the error term from Lemma 4.5 over $|t| \leq T$ and k_0 , we get

$$\ll_{E,\varepsilon_0,\varepsilon} T^{\frac{6}{5}} K^{\frac{9}{5}+\varepsilon} H = \frac{K^{3+\frac{6}{5}\varepsilon_0+\varepsilon} H}{h^{\frac{6}{5}}}. \quad (4-4)$$

The sum of the error terms from (3-4) and (4-4) is thus

$$\ll_{E,\varepsilon_0,\varepsilon} hK^{2+\varepsilon} + \frac{HK^3 \log K}{h^2} + \frac{HK^{3+\frac{6}{5}\varepsilon_0+\varepsilon}}{h^{\frac{6}{5}}} \ll hK^{2+\varepsilon} + \frac{HK^{3+\frac{6}{5}\varepsilon_0+\varepsilon}}{h^{\frac{6}{5}}},$$

so we obtain

$$\Sigma = O_{E,\varepsilon_0,\varepsilon} \left(hK^{2+\varepsilon} + \frac{HK^{3+\frac{6}{5}\varepsilon_0+\varepsilon}}{h^{\frac{6}{5}}} \right) + \frac{(-1)^\delta h}{\pi} \sum_{k_0} \int_{NE} \sum_{\substack{t \in \mathbb{Z} \\ |t| \leq T}} L(1, \tilde{\psi}_t) \cos \frac{(k_0-1)t}{2\sqrt{u}} \widehat{W} \left(\frac{ht}{\pi\sqrt{u}} \right) du.$$

Having replaced $L(1, \psi_{t^2-4n})$ by its average, we can now extend the sum over t out to $\pm\infty$, with an error of $O_{E,\varepsilon_0}(1)$. Taking ε_0 arbitrarily small, writing $E = [\alpha_2^{-2}, \alpha_1^{-2}]$ and making the substitution $u = \left(\frac{k_0-1}{4\pi\alpha}\right)^2$, we thus get

$$\Sigma = O_{E,\varepsilon} \left(hK^{2+\varepsilon} + \frac{HK^{3+\varepsilon}}{h^{\frac{6}{5}}} \right) + \frac{(-1)^\delta 2h}{\pi} \sum_{k_0} \left(\frac{k_0-1}{4\pi} \right)^2 \int_{\lambda_{k_0}\alpha_1}^{\lambda_{k_0}\alpha_2} \sum_{t \in \mathbb{Z}} L(1, \tilde{\psi}_t) \cos(2\pi\alpha t) \widehat{W} \left(\frac{t}{x_{k_0}(\alpha)} \right) \frac{d\alpha}{\alpha^3}, \quad (4-5)$$

where $\lambda_{k_0} = \frac{k_0-1}{4\pi\sqrt{N}}$ and $x_{k_0}(\alpha) = \frac{k_0-1}{4\alpha h}$.

5. The exponential sum

In this section we evaluate the sum over t in (4-5), proving the following:

Proposition 5.1. *Assume GRH for Dirichlet L -functions. Let $\alpha, \theta, x \in \mathbb{R}$ and $a, q \in \mathbb{Z}$ with $x, q \geq 1$, $\gcd(a, q) = 1$, $\alpha = a/q + \theta$, and $|\theta| \leq 1/q^2$. Then,*

$$\begin{aligned} & \sum_{t \in \mathbb{Z}} L(1, \tilde{\psi}_t) \cos(2\pi\alpha t) \widehat{W} \left(\frac{t}{x} \right) \\ &= \frac{\mu(q)^2}{\varphi(q)^2 \sigma(q)} x W(x\theta) + O(qx^{-1} \max(1, x|\theta|)) + O_\varepsilon(q^3 x^{-\frac{7}{4}+\varepsilon} \max(1, x|\theta|)^{\frac{7}{2}}). \end{aligned}$$

Recall from Lemma 4.3 that $L(1, \tilde{\psi}_t) = Cf(t)$, where $C = L(1, \tilde{\psi}_1)$ and

$$f(t) = \prod_{p|t} \left(1 - \frac{1}{(p-1)^2(p+1)} \right)$$

is multiplicative. The next lemma studies the generating function of f and its additive twists.

Lemma 5.2. *Let $q \in \mathbb{Z}_{>0}$ and $a \in \mathbb{Z}$ with $\gcd(a, q) = 1$. For $\Re(s) > 1$, define*

$$F(s; a/q) = \sum_{n=1}^{\infty} \frac{f(n)e(an/q)}{n^s} \quad \text{and} \quad F^{\pm}(s; a/q) = \frac{F(s; a/q) \pm F(s; -a/q)}{2}.$$

Then

$$F^{\pm}(s; a/q) = \sum_{d|q} \frac{f(d)}{d^s} \sum_{\substack{\chi \bmod \frac{q}{d} \\ \chi(-1) = \pm 1}} \frac{\chi(a)\tau(\bar{\chi})}{\varphi(q/d)} \frac{L(s, \chi)L(s+2, \chi\chi_0)}{L(2(s+2), \chi\chi_0)} \prod_{p|q} \left(1 + \frac{\chi(p)p^{-s-2} \frac{p+1}{p^2-p-1}}{1 + \chi(p)p^{-s-2}} \right), \quad (5-1)$$

where χ_0 denotes the trivial character modulo q .

It follows that $F^{\pm}(s; a/q)$ has meromorphic continuation to $\Re(s) > -2$ with the following additional properties:

- $F^-(s; a/q)$ and $(s^2 - 1)F^+(s; a/q)$ are analytic for $\Re(s) \geq -\frac{3}{2}$ unconditionally, and for $\Re(s) > -\frac{7}{4}$ under GRH.
- $\text{Res}_{s=1} F^+(s; a/q) = \frac{1}{C} \frac{\mu(q)^2}{\varphi(q)^2 \sigma(q)}$.
- $F^+(0; a/q) = -\frac{1}{2}f(0)$.
- $\text{Res}_{s=-1} F^+(s; a/q) = -\frac{q}{12C} \prod_{p|q} \left(1 - \frac{1}{p^3 - p} \right)$.
- $F^-(-1; a/q) = 0$.

Proof. By orthogonality of Dirichlet characters,

$$e\left(\frac{an}{q}\right) = \sum_{d|\gcd(n,q)} \frac{1}{\varphi(q/d)} \sum_{\chi \bmod \frac{q}{d}} \chi(a)\tau(\bar{\chi})\chi\left(\frac{n}{d}\right).$$

For $\Re(s) > 1$, we have

$$\begin{aligned} F(s; a/q) &= \sum_{n=1}^{\infty} \frac{f(n)e(an/q)}{n^s} \\ &= \sum_{n=1}^{\infty} \sum_{d|\gcd(n,q)} \frac{1}{\varphi(q/d)} \sum_{\chi \bmod \frac{q}{d}} \chi(a)\tau(\bar{\chi}) \frac{f(n)\chi(n/d)}{n^s} \\ &= \sum_{d|q} \frac{1}{\varphi(q/d)} \sum_{\chi \bmod \frac{q}{d}} \chi(a)\tau(\bar{\chi}) \sum_{n=1}^{\infty} \frac{f(nd)\chi(n)}{(nd)^s}. \end{aligned}$$

Recalling (4-3), we observe that $f(n)$ is multiplicative; further, for any $j \geq 1$,

$$f(p^j) = f(p) = 1 + \frac{1}{p^2 - p - 1}.$$

So for each $d|q$ and χ modulo q/d , we get

$$\begin{aligned} \sum_{n=1}^{\infty} \frac{f(nd)\chi(n)}{(nd)^s} &= d^{-s} \prod_p \sum_{j=0}^{\infty} \frac{f(p^{\text{ord}_p(d)+j})\chi(p^j)}{p^{js}} \\ &= d^{-s} \prod_{p|d} \left(f(p) \sum_{j=0}^{\infty} \frac{\chi(p^j)}{p^{js}} \right) \prod_{p \nmid d} \left(1 + f(p) \sum_{j=1}^{\infty} \frac{\chi(p^j)}{p^{js}} \right) \\ &= d^{-s} f(d)L(s, \chi) \prod_{p \nmid q} (1 + (f(p) - 1)\chi(p)p^{-s}). \end{aligned}$$

Thus we have

$$F^{\pm}(s; a/q) = \sum_{d|q} \frac{f(d)}{d^s} \frac{1}{\varphi(q/d)} \sum_{\substack{\chi \bmod \frac{q}{d} \\ \chi(-1) = \pm 1}} \chi(a)\tau(\bar{\chi})L(s, \chi) \prod_{p \nmid q} (1 + (f(p) - 1)\chi(p)p^{-s}). \tag{5-2}$$

Further, denoting by χ_0 the trivial character mod q , we can write

$$\prod_{p \nmid q} (1 + (f(p) - 1)\chi(p)p^{-s}) = \frac{L(s+2, \chi\chi_0)}{L(2(s+2), \chi\chi_0)} \prod_{p \nmid q} \left(1 + \frac{\chi(p)p^{-s-2} \frac{p+1}{p^2-p-1}}{1 + \chi(p)p^{-s-2}} \right),$$

which in turn implies (5-1). It follows that $F^{\pm}(s; a/q)$ has meromorphic continuation to $\Re(s) > -2$, with poles possible at the poles of $L(s, \chi)L(s+2, \chi\chi_0)/L(2(s+2), \chi\chi_0)$. For $\Re(s) \geq -\frac{3}{2}$ unconditionally, and $\Re(s) > -\frac{7}{4}$ under GRH, poles can only occur at $s = \pm 1$ when χ is the trivial character modulo q/d for some $d|q$. Since the trivial character is always even, these poles can only occur in F^+ .

Let $\chi_{q/d}$ denote the trivial character modulo q/d . Then

$$\tau(\bar{\chi}_{q/d}) = \sum_{\substack{u \bmod \frac{q}{d} \\ \gcd(u, \frac{q}{d})=1}} e^{2\pi i \frac{u}{q/d}} = \mu\left(\frac{q}{d}\right),$$

so that

$$\begin{aligned} \text{Res}_{s=1} F^+(s; a/q) &= \prod_{p \nmid q} (1 + (f(p) - 1)p^{-1}) \sum_{d|q} \frac{\mu(q/d)}{\varphi(q/d)} \frac{f(d)}{d} \prod_{p|q/d} (1 - p^{-1}) \\ &= \prod_{p \nmid q} (1 + (f(p) - 1)p^{-1}) \frac{1}{q} \sum_{d|q} \mu\left(\frac{q}{d}\right) f(d) \\ &= \frac{1}{q} \prod_{p \nmid q} (1 + (f(p) - 1)p^{-1}) \begin{cases} \prod_{p|q} (f(p) - 1) & \text{if } q \text{ is squarefree,} \\ 0 & \text{otherwise.} \end{cases} \end{aligned}$$

When q is squarefree, we get

$$\text{Res}_{s=1} F^+(s; a/q) = \frac{C'}{q} \prod_{p|q} (1 + (f(p) - 1)p^{-1})^{-1} (f(p) - 1) = C' \prod_{p|q} \frac{1}{(p^2 - 1)(p - 1)},$$

where

$$C' = \prod_p \left(1 + \frac{1}{p(p^2 - p - 1)} \right) = \prod_p \frac{(p^2 - 1)(p - 1)}{p(p^2 - p - 1)} = \frac{1}{C}.$$

At $s = -1$, using that $L(s, \chi_{q/d}) = \zeta(s) \prod_{p|q/d} (1 - p^{-s})$, we have

$$\begin{aligned} \operatorname{Res}_{s=-1} F^+(s; a/q) &= \prod_{p \nmid q} \left(1 + \frac{p^{-1} \frac{p+1}{p^2-p-1}}{1+p^{-1}} \right) \sum_{d|q} \frac{f(d)d}{\varphi(q/d)} \tau(\chi_{q/d}) L(-1, \chi_{q/d}) \frac{\prod_{p|q} (1-p^{-1})}{L(2, \chi_0)} \\ &= \frac{\zeta(-1)}{\zeta(2)} \prod_{p \nmid q} f(p) \sum_{d|q} \frac{f(d)d\mu(q/d)}{\varphi(q/d)} \prod_{p|\frac{q}{d}} (1-p) \frac{\prod_{p|q} (1-p^{-1})}{\prod_{p|q} (1-p^{-2})} \\ &= \frac{\zeta(-1)}{\zeta(2)} \prod_{p \nmid q} f(p) \prod_{p|q} (1+p^{-1})^{-1} \sum_{d|q} \frac{f(d)d\mu(q/d)^2 q}{\varphi(q/d) d} \prod_{p|\frac{q}{d}} (1-p^{-1}) \\ &= \frac{\zeta(-1)}{\zeta(2)} \prod_{p \nmid q} f(p) \prod_{p|q} (1+p^{-1})^{-1} \sum_{d|q} f(d)d\mu(q/d)^2 \\ &= \frac{\zeta(-1)}{\zeta(2)} \prod_{p \nmid q} f(p) \prod_{p|q} (1+p^{-1})^{-1} \prod_{p \parallel q} (1+f(p)p) \prod_{p^2|q} f(p)(p^{\operatorname{ord}_p(q)} + p^{\operatorname{ord}_p(q)-1}) \\ &= \frac{\zeta(-1)}{\zeta(2)} \prod_p f(p) \cdot q \prod_{p|q} (1+p^{-1})^{-1} \prod_{p \parallel q} \frac{1+f(p)p}{pf(p)} \prod_{p^2|q} (1+p^{-1}). \end{aligned}$$

Since $\prod_p f(p) = f(0)$, $Cf(0) = \zeta(2)$ and $\zeta(-1) = -\frac{1}{12}$, we get

$$\operatorname{Res}_{s=-1} F^+(s; a/q) = -\frac{q}{12C} \prod_{p \parallel q} \frac{1+f(p)p}{f(p)(p+1)}.$$

Next, when χ is an odd character, $L(s, \chi)$ has a trivial zero at $s = -1$, so from (5-1) we see that $F^-(-1; a/q) = 0$.

Similarly, for even χ , $L(s, \chi)$ has a trivial zero at $s = 0$ unless χ is the character of modulus 1. Thus the only nonzero term in (5-2) comes from $d = q$, so we have

$$F^+(0; a/q) = f(q)\zeta(0) \prod_{p \nmid q} f(p) = -\frac{1}{2}f(0). \quad \square$$

Lemma 5.3. *Define*

$$g^\pm(t; \theta, x) = \frac{e(\theta t) \pm e(-\theta t)}{2} \widehat{W}\left(\frac{t}{x}\right) \quad \text{and} \quad \tilde{g}^\pm(s; \theta, x) = \int_0^\infty g^\pm(t; \theta, x) t^s \frac{dt}{t}.$$

Then $\tilde{g}^\pm(s; \theta, x)$ has meromorphic continuation to \mathbb{C} , with at most simple poles in $\{s \in \mathbb{Z} : s \leq 0, (-1)^s = \pm 1\}$, and satisfies

$$\tilde{g}^\pm(s; \theta, x) \ll_j \frac{x^{\Re(s)} \max(1, x|\theta|)^j}{(\Re(s) + j)|s(s+1) \cdots (s+j-1)|} \quad \text{for } j \in \mathbb{Z}_{\geq 0} \text{ and } \Re(s) > -j.$$

Furthermore, the following identities hold:

- $\tilde{g}^+(1; \theta, x) = \frac{1}{2}xW(x\theta)$.
- $\text{Res}_{s=0} \tilde{g}^+(s; \theta, x) = 1$.
- $\tilde{g}^+(-1; \theta, x) = \frac{\pi^2}{x} \int_{\mathbb{R}} \max(|v|, x|\theta|)W(v) dv$.

Proof. For brevity we assume that θ and x are fixed throughout this proof and suppress them from the notation. Applying integration by parts, for $j \geq 0$ and $\Re(s) > -j$, we have

$$\tilde{g}^{\pm}(s) = \frac{(-1)^j}{s(s+1)\cdots(s+j-1)} \int_0^{\infty} (g^{\pm})^{(j)}(t)t^{s+j-1} dt.$$

Note that when j is odd, $(g^+)^{(j)}(0) = 0$, so this integral representation is valid for $\Re(s) > -j - 1$, and it follows that $\tilde{g}^+(s)$ is analytic at $s = -j$. Similarly, $\tilde{g}^-(s)$ is analytic at $s = -j$ for even j .

By the Leibniz rule, for any $j \geq 0$,

$$|(g^{\pm})^{(j)}(t)| \leq \sum_{b=0}^j \binom{j}{b} |2\pi\theta|^b x^{b-j} \left| \widehat{W}^{(j-b)}\left(\frac{t}{x}\right) \right| \ll_j x^{-j} \max(1, x|\theta|)^j.$$

Inserting this estimate into the integral representation above, for $\Re(s) > -j$ we have

$$\tilde{g}^{\pm}(s) \ll_j \frac{x^{-j} \max(1, x|\theta|)^j}{|s(s+1)\cdots(s+j-1)|} \int_0^x t^{\Re(s)+j-1} dt = \frac{x^{\Re(s)} \max(1, x|\theta|)^j}{(\Re(s)+j)|s(s+1)\cdots(s+j-1)|}.$$

Turning to the additional identities, we first have

$$\tilde{g}^+(1) = \int_0^{\infty} \cos(2\pi\theta t) \widehat{W}\left(\frac{t}{x}\right) dt = \frac{x}{2} \int_{\mathbb{R}} e(x\theta t) \widehat{W}(t) dt = \frac{x}{2} W(x\theta).$$

Second, using the integral representation with $j = 1$, we have

$$\text{Res}_{s=0} \tilde{g}^+(s) = - \int_0^{\infty} (g^+)'(t) dt = g^+(0) = 1.$$

Third, for $u \in \mathbb{R}$, let

$$G(u) = \int_{-\infty}^{\infty} (g^+)'(t)t^{-1}e(-ut) dt.$$

Then

$$G'(u) = -2\pi i \int_{-\infty}^{\infty} (g^+)'(t)e(-ut) dt = -4\pi^2 u \int_{-\infty}^{\infty} g^+(t)e(-ut) dt = -4\pi^2 u \widehat{g}^+(u).$$

Since $G(u) \rightarrow 0$ as $|u| \rightarrow \infty$, we have

$$G(u) = \int_{-\infty}^u G'(v) dv = -4\pi^2 \int_{-\infty}^u v \widehat{g}^+(v) dv.$$

Taking $u = 0$, we get

$$\tilde{g}^+(-1) = \frac{1}{2} \int_{-\infty}^{\infty} (g^+)'(t)t^{-1} dt = -2\pi^2 \int_{-\infty}^0 v \widehat{g}^+(v) dv = 2\pi^2 \int_0^{\infty} v \widehat{g}^+(v) dv.$$

We now compute $\widehat{g^+}(v)$. Recalling the definition of g^+ , we have

$$\begin{aligned}\widehat{g^+}(v) &= \int_{-\infty}^{\infty} g^+(u) e(-uv) du = \frac{1}{2} \left\{ \int_{-\infty}^{\infty} \widehat{W}\left(\frac{u}{x}\right) e(-u(v-\theta)) du + \int_{-\infty}^{\infty} \widehat{W}\left(\frac{u}{x}\right) e(-u(v+\theta)) du \right\} \\ &= \frac{x}{2} \{W(x(v-\theta)) + W(x(v+\theta))\}.\end{aligned}$$

Therefore

$$\begin{aligned}\tilde{g}^+(-1) &= \pi^2 x \int_0^{\infty} v \{W(x(v-\theta)) + W(x(v+\theta))\} dv \\ &= \pi^2 \left\{ \frac{1}{x} \int_{-x|\theta|}^{\infty} v W(v) dv + \frac{1}{x} \int_{x|\theta|}^{\infty} v W(v) dv + \int_{-x|\theta|}^{x|\theta|} |\theta| W(v) dv \right\} \\ &= 2\pi^2 \left\{ \frac{1}{x} \int_{x|\theta|}^{\infty} v W(v) dv + \int_0^{x|\theta|} |\theta| W(v) dv \right\} = 2\pi^2 \int_0^{\infty} \max\left(\frac{v}{x}, |\theta|\right) W(v) dv.\end{aligned}$$

□

We can now complete the proof of Proposition 5.1. Recall again from Lemma 4.3 that $L(1, \tilde{\psi}_t) = Cf(t)$. In the notation of Lemma 5.3, we have

$$\begin{aligned}\sum_{t \in \mathbb{Z}} \cos(2\pi\alpha t) f(t) \widehat{W}\left(\frac{t}{x}\right) &= f(0) + 2 \sum_{t=1}^{\infty} f(t) \cos(2\pi\alpha t) \widehat{W}\left(\frac{t}{x}\right) \\ &= f(0) + 2 \sum_{t=1}^{\infty} f(t) \cos \frac{2\pi\alpha t}{q} g^+(t; \theta, x) + 2i \sum_{t=1}^{\infty} \sin \frac{2\pi\alpha t}{q} g^-(t; \theta, x).\end{aligned}$$

By Mellin inversion, we have

$$g^{\pm}(t; \theta, x) = \frac{1}{2\pi i} \int_{(2)} \tilde{g}^{\pm}(s; \theta, x) t^{-s} ds,$$

so that

$$\sum_{t=1}^{\infty} f(t) \cos\left(\frac{2\pi\alpha t}{q}\right) g^+(t; \theta, x) = \frac{1}{2\pi i} \int_{(2)} \tilde{g}^+(s; \theta, x) F^+(s; a/q) ds.$$

Shifting the contour to $\Re(s) = \sigma$ for $-\frac{7}{4} < \sigma \leq -\frac{3}{2}$, we get

$$\begin{aligned}\tilde{g}^+(1; \theta, x) \operatorname{Res}_{s=1} F^+(s; a/q) + F^+(0; a/q) \operatorname{Res}_{s=0} \tilde{g}^+(s; \theta, x) \\ + \tilde{g}^+(-1; \theta, x) \operatorname{Res}_{s=-1} F(s; a/q) + \frac{1}{2\pi i} \int_{(\sigma)} \tilde{g}^+(s; \theta, x) F^+(s; a/q) ds \\ = \frac{1}{C} \frac{\mu(q)^2}{\varphi(q)^2 \sigma(q)} \frac{x}{2} W(x\theta) - \frac{1}{2} f(0) - \frac{1}{2} f(0) \frac{q}{x} \prod_{p|q} \left(1 - \frac{1}{p^3-p}\right) \cdot \int_{\mathbb{R}} \max(|v|, x|\theta|) W(v) dv \\ + \frac{1}{2\pi i} \int_{(\sigma)} \tilde{g}^+(s; \theta, x) F(s; a/q) ds.\end{aligned}$$

Similarly we have

$$\begin{aligned} i \sum_{t=1}^{\infty} f(t) \sin \frac{2\pi at}{q} g^-(t; \theta, x) &= \frac{1}{2\pi i} \int_{(2)} \tilde{g}^-(s; \theta, x) F^-(s; a/q) ds \\ &= \frac{1}{2\pi i} \int_{(\sigma)} \tilde{g}^-(s; \theta, x) F^-(s; a/q) ds. \end{aligned}$$

There is no residue term in this case since $F^-(-1; a/q) = 0$.

Therefore, for $-\frac{7}{4} < \sigma \leq -\frac{3}{2}$, we have

$$\begin{aligned} \sum_{t \in \mathbb{Z}} \cos(2\pi at) f(t) \widehat{W}\left(\frac{t}{x}\right) &= \frac{1}{C} \frac{\mu(q)^2}{\varphi(q)^2 \sigma(q)} x W(x\theta) - f(0) \frac{q}{x} \prod_{p|q} \left(1 - \frac{1}{p^3 - p}\right) \int_{\mathbb{R}} \max(|v|, x|\theta|) W(v) dv \\ &\quad + \frac{1}{\pi i} \int_{(\sigma)} (\tilde{g}^+(s; \theta, x) F^+(s; a/q) + \tilde{g}^-(s; \theta, x) F^-(s; a/q)) ds. \end{aligned}$$

The second line is $O(qx^{-1} \max(1, x|\theta|))$. As for the third line, assuming GRH and applying Lemma 5.2, for $\Re(s) = \sigma \in (-\frac{7}{4}, -\frac{3}{2}]$ we have

$$\begin{aligned} F^{\pm}(s; a/q) &\ll_{\sigma, \varepsilon} \sum_{d|q} \frac{1}{\varphi(q/d)} \frac{f(d)}{d^{\sigma}} \sum_{\substack{\chi \bmod \frac{q}{d} \\ \chi(-1) = \pm 1}} |\tau(\chi)| \left| \frac{q}{d} s \right|^{\frac{1}{2} - \sigma + \frac{1}{2} - (\sigma + 2) + \varepsilon} \\ &\ll |s|^{-1-2\sigma+\varepsilon} \sum_{d|q} \frac{q^{-\frac{1}{2}-2\sigma+\varepsilon}}{d^{-\frac{1}{2}-\sigma+\varepsilon}} \ll_{\varepsilon} |s|^{-1-2\sigma+\varepsilon} q^{-\frac{1}{2}-2\sigma+\varepsilon}. \end{aligned}$$

So we get

$$\frac{1}{\pi i} \int_{(\sigma)} (\tilde{g}^+(s) F^+(s; a/q) + \tilde{g}^-(s) F^-(s; a/q)) ds \ll_{\sigma, \varepsilon} \sum_{\pm} q^{-\frac{1}{2}-2\sigma+\varepsilon} \int_{(\sigma)} |\tilde{g}^{\pm}(s)| |s|^{-1-2\sigma+\varepsilon} |ds|.$$

By Lemma 5.3, for $\Re(s) = \sigma \in (-\frac{7}{4}, -\frac{3}{2}]$ and $j \geq 2$,

$$\tilde{g}^{\pm}(s) \ll_j x^{\sigma} \frac{\max(1, x|\theta|)^j}{(\sigma + j)|s(s+1) \cdots (s+j-1)|} \ll_j x^{\sigma} |s|^{-j} \max(1, x|\theta|)^j.$$

We use $j = 2$ for $|s| < \max(1, x|\theta|)$ and $j = 4$ for $|s| \geq \max(1, x|\theta|)$, obtaining

$$\ll_{\sigma, \varepsilon} q^{-\frac{1}{2}-2\sigma+\varepsilon} x^{\sigma} \max(1, x|\theta|)^{-2\sigma+\varepsilon}.$$

Choosing $\sigma = -\frac{7}{4} + \frac{\varepsilon}{2}$ gives

$$\ll_{\varepsilon} q^3 x^{-\frac{7}{4}+\varepsilon} \max(1, x|\theta|)^{\frac{7}{2}}.$$

5.1. The Fourier series. We conclude this section by proving equality of the two expressions for $\nu(E)$ appearing in Theorem 1.1. Formally this should follow from Proposition 5.1, but we give a direct proof.

Consider the function $S : \mathbb{R}_{>0} \rightarrow \mathbb{R}$ defined by

$$S(\alpha) = \frac{1}{2} - \zeta(2)\alpha + \sum_{a/q \in \mathbb{Q} \cap (0, \alpha]} \frac{\mu(q)^2}{\varphi(q)^2 \sigma(q)}.$$

(If $\alpha \in \mathbb{Q}$ then the corresponding term of the sum is counted with the full weight.) For any α , we have

$$\begin{aligned} S(\alpha + 1) - S(\alpha) &= -\zeta(2) + \sum_{q=1}^{\infty} \sum_{\substack{\alpha q < a \leq \alpha q + q \\ (a, q) = 1}} \frac{\mu(q)^2}{\varphi(q)^2 \sigma(q)} \\ &= -\zeta(2) + \sum_{q=1}^{\infty} \frac{\mu(q)^2}{\varphi(q) \sigma(q)} \\ &= -\zeta(2) + \prod_p \left(1 + \frac{1}{(p-1)(p+1)} \right) = 0, \end{aligned}$$

so $S(\alpha)$ is periodic mod 1 and has bounded variation.

It follows that the Fourier expansion of $S(\alpha)$ converges at every point. We proceed to calculate its Fourier coefficients:

$$\begin{aligned} \int_0^1 S(\alpha) e(-\alpha t) d\alpha &= \int_0^1 \left(\frac{1}{2} - \zeta(2)\alpha + \sum_{a/q \in \mathbb{Q} \cap (0, \alpha]} \frac{\mu(q)^2}{\varphi(q)^2 \sigma(q)} \right) e(-\alpha t) d\alpha \\ &= \int_0^1 \left(\frac{1}{2} - \zeta(2)\alpha \right) e(-\alpha t) d\alpha + \sum_{q=1}^{\infty} \sum_{\substack{1 \leq a \leq q \\ (a, q) = 1}} \frac{\mu(q)^2}{\varphi(q)^2 \sigma(q)} \int_{a/q}^1 e(-\alpha t) d\alpha. \end{aligned}$$

When $t = 0$ this is

$$\frac{1 - \zeta(2)}{2} + \sum_{q=1}^{\infty} \frac{\mu(q)^2}{\varphi(q)^2 \sigma(q)} \sum_{\substack{1 \leq a \leq q \\ (a, q) = 1}} \left(1 - \frac{a}{q} \right).$$

Since $\gcd(q - a, q) = \gcd(a, q)$, the inner sum is

$$\sum_{\substack{1 \leq a \leq q \\ (a, q) = 1}} \left(1 - \frac{a}{q} \right) = \begin{cases} 0 & \text{if } q = 1, \\ \frac{1}{2} \varphi(q) & \text{if } q > 1, \end{cases}$$

so we get

$$\frac{1 - \zeta(2)}{2} + \frac{1}{2} \sum_{q=2}^{\infty} \frac{\mu(q)^2}{\varphi(q) \sigma(q)} = 0.$$

For $t \neq 0$ we get

$$\begin{aligned} \frac{\zeta(2)}{2\pi it} + \sum_{q=1}^{\infty} \sum_{\substack{1 \leq a \leq q \\ (a,q)=1}} \frac{\mu(q)^2}{\varphi(q)^2 \sigma(q)} \frac{e(-at/q) - 1}{2\pi it} &= \frac{\zeta(2)}{2\pi it} + \sum_{q=1}^{\infty} \frac{\mu(q)^2}{\varphi(q)^2 \sigma(q)} \frac{c_q(-t) - \varphi(q)}{2\pi it} \\ &= \frac{1}{2\pi it} \sum_{q=1}^{\infty} \frac{\mu(q)^2 c_q(t)}{\varphi(q)^2 \sigma(q)}, \end{aligned}$$

where $c_q(t) = \sum_{a \in (\mathbb{Z}/q\mathbb{Z})^\times} e(at/q)$ is the Ramanujan sum. By [MV07, §4.1, Theorem 1], we have

$$c_q(t) = \mu\left(\frac{q}{(q,t)}\right) \frac{\varphi(q)}{\varphi(q/(q,t))},$$

so this becomes

$$\begin{aligned} \frac{1}{2\pi it} \sum_{q=1}^{\infty} \frac{\mu(q)\mu((q,t))\varphi((q,t))}{\varphi(q)^2 \sigma(q)} &= \frac{1}{2\pi it} \prod_{p|t} \left(1 - \frac{1}{(p-1)^2(p+1)}\right) \prod_{p \nmid t} \left(1 + \frac{1}{(p-1)(p+1)}\right) \\ &= \frac{L(1, \tilde{\psi}_t)}{2\pi it}. \end{aligned}$$

Therefore, $S(\alpha)$ has Fourier series

$$\sum_{t=1}^{\infty} \frac{L(1, \tilde{\psi}_t)}{\pi t} \sin(2\pi \alpha t).$$

By the Dirichlet–Dini criterion, at the jump discontinuities of $S(\alpha)$ (i.e., at every rational with squarefree denominator), the series converges to the average of the left and right limits. Thus, it equals

$$S^*(\alpha) := -\zeta(2)\alpha + \sum_{a/q \in \mathbb{Q} \cap [0, \alpha]}^* \frac{\mu(q)^2}{\varphi(q)^2 \sigma(q)},$$

where the $*$ indicates that the endpoints are weighted by $\frac{1}{2}$, as in (1-2).

Next, to relate this to ν , we define a measure λ on $\mathbb{R}_{>0}$ that assigns mass

$$\frac{1}{\zeta(2)} \frac{\mu(q)^2}{\varphi(q)^2 \sigma(q)} \left(\frac{q}{a}\right)^3$$

to each $a/q \in \mathbb{Q}_{>0}$. By Stieltjes integration, for $\alpha_2 > \alpha_1 > 0$, we have

$$\begin{aligned} \lambda((\alpha_1, \alpha_2]) &= \frac{1}{\zeta(2)} \int_{\alpha_1}^{\alpha_2} \alpha^{-3} d(S(\alpha) + \zeta(2)\alpha) = \int_{\alpha_1}^{\alpha_2} \alpha^{-3} d\alpha + \frac{1}{\zeta(2)} \int_{\alpha_1}^{\alpha_2} \alpha^{-3} dS(\alpha) \\ &= \int_{\alpha_1}^{\alpha_2} \alpha^{-3} d\alpha + \frac{S(\alpha)}{\zeta(2)\alpha^3} \Big|_{\alpha_1}^{\alpha_2} + \frac{3}{\zeta(2)} \int_{\alpha_1}^{\alpha_2} \alpha^{-4} S(\alpha) d\alpha. \end{aligned}$$

We can replace $S(\alpha)$ by $S^*(\alpha)$ in the integral, since they differ on a set of measure 0:

$$\begin{aligned} \frac{3}{\zeta(2)} \int_{\alpha_1}^{\alpha_2} \alpha^{-4} S(\alpha) d\alpha &= \frac{3}{\zeta(2)} \int_{\alpha_1}^{\alpha_2} \alpha^{-4} S^*(\alpha) d\alpha \\ &= \frac{3}{\zeta(2)} \int_{\alpha_1}^{\alpha_2} \alpha^{-4} \sum_{t=1}^{\infty} \frac{L(1, \tilde{\psi}_t)}{\pi t} \sin(2\pi\alpha t) d\alpha. \end{aligned}$$

Since $S^*(\alpha)$ is square-integrable, we are free to change the order of sum and integral:

$$\begin{aligned} \frac{3}{\zeta(2)} \int_{\alpha_1}^{\alpha_2} \alpha^{-4} S(\alpha) d\alpha &= \frac{3}{\zeta(2)} \sum_{t=1}^{\infty} \frac{L(1, \tilde{\psi}_t)}{\pi t} \int_{\alpha_1}^{\alpha_2} \sin(2\pi\alpha t) \alpha^{-4} d\alpha \\ &= \frac{1}{\zeta(2)} \sum_{t=1}^{\infty} \frac{L(1, \tilde{\psi}_t)}{\pi t} \left(-\frac{\sin(2\pi\alpha t)}{\alpha^3} \Big|_{\alpha_1}^{\alpha_2} + 2\pi t \int_{\alpha_1}^{\alpha_2} \cos(2\pi\alpha t) \alpha^{-3} d\alpha \right) \\ &= -\frac{S^*(\alpha)}{\zeta(2)\alpha^3} \Big|_{\alpha_1}^{\alpha_2} + \frac{2}{\zeta(2)} \sum_{t=1}^{\infty} L(1, \tilde{\psi}_t) \int_{\alpha_1}^{\alpha_2} \cos(2\pi\alpha t) \alpha^{-3} d\alpha. \end{aligned}$$

Substituting back into the above and using that $L(1, \tilde{\psi}_t) = \zeta(2) \prod_{p \nmid t} \frac{p^2 - p - 1}{p^2 - p}$, we get

$$\begin{aligned} \lambda((\alpha_1, \alpha_2]) &= \frac{S(\alpha) - S^*(\alpha)}{\zeta(2)\alpha^3} \Big|_{\alpha_1}^{\alpha_2} + \sum_{t=-\infty}^{\infty} \frac{L(1, \tilde{\psi}_t)}{\zeta(2)} \int_{\alpha_1}^{\alpha_2} \cos(2\pi\alpha t) \alpha^{-3} d\alpha \\ &= \lambda((\alpha_1, \alpha_2]) - \nu([\alpha_2^{-2}, \alpha_1^{-2}]) + \frac{1}{2} \sum_{t=-\infty}^{\infty} \prod_{p \nmid t} \frac{p^2 - p - 1}{p^2 - p} \cdot \int_{\alpha_2^{-2}}^{\alpha_1^{-2}} \cos \frac{2\pi t}{\sqrt{y}} dy. \end{aligned}$$

This completes the proof.

6. The circle method

Recall that $\lambda_{k_0} = \frac{k_0 - 1}{4\pi\sqrt{N}}$ and $x_{k_0}(\alpha) = \frac{k_0 - 1}{4\alpha h}$. We further define

$$\Delta = \frac{H}{4\pi\sqrt{N}}, \quad P = \frac{1}{3\sqrt{\Delta\alpha_2}}, \quad Q = \frac{H}{4\Delta\alpha_2 h P}.$$

Note that $P \asymp_E \sqrt{K/H}$, $Q \asymp_E \sqrt{HK}/h$ and $x_{k_0}(\alpha) \asymp_E P Q \asymp_E K/h$. For $a, q \in \mathbb{Z}$ with $q > 0$ and $(a, q) = 1$, define

$$\mathfrak{M}\left(\frac{a}{q}\right) = \left\{ \frac{a}{q} + \theta : |\theta| \leq \frac{1}{qQ} \right\}.$$

Since $4h < H$, we have

$$\frac{P}{Q} = \frac{4h}{9H} < \frac{1}{9} < \frac{1}{2},$$

and it follows that $\mathfrak{M}(a/q) \cap \mathfrak{M}(a'/q') = \emptyset$ when $0 < q, q' \leq P$ and $a/q \neq a'/q'$.

For a fixed k_0 , we decompose the interval $I_{k_0} := [\lambda_{k_0}\alpha_1, \lambda_{k_0}\alpha_2]$ into major arcs \mathfrak{M}_{k_0} and minor arcs \mathfrak{m}_{k_0} , defined by

$$\mathfrak{M}_{k_0} = I_{k_0} \cap \bigcup_{\substack{a, q \in \mathbb{Z} \\ 0 < q \leq P \\ (a, q) = 1}} \mathfrak{M}\left(\frac{a}{q}\right) \quad \text{and} \quad \mathfrak{m}_{k_0} = I_{k_0} \setminus \mathfrak{M}_{k_0}.$$

Note that the definitions of P , Q , and $\mathfrak{M}(a/q)$ are independent of k_0 ; only their intersection with I_{k_0} can vary. As we will show, near each endpoint there is at most one fraction a/q with $0 < q \leq P$ for which $\mathfrak{M}(a/q) \cap I_{k_0}$ varies.

To that end, since $4\pi\sqrt{N} = K - 1 + O(1/K)$, we have

$$\begin{aligned} |\lambda_{k_0} - 1| &= \frac{|k_0 - K| + O(1/K)}{4\pi\sqrt{N}} \\ &\leq \frac{H - 4h + O(1/K)}{4\pi\sqrt{N}} \leq \Delta \end{aligned}$$

for K large enough. Thus, as k_0 varies, the endpoint $\lambda_{k_0}\alpha_i$ is confined to the interval $[(1-\Delta)\alpha_i, (1+\Delta)\alpha_i]$. Next, by Dirichlet's theorem, for $i = 1, 2$ we can choose a fraction a_i/q_i such that $q_i \leq 3P$ and

$$\alpha_i = \frac{a_i}{q_i} + \theta_i, \quad \text{with } |\theta_i| \leq \frac{1}{3q_i P}.$$

Assume that K is sufficiently large to ensure that $a_i > 0$.

Let a/q be a fraction with $0 < q \leq P$ and $a/q \notin \{a_1/q_1, a_2/q_2\}$. Adding the inequalities

$$qq_i \Delta \alpha_i \leq 3P^2 \Delta \alpha_2 = \frac{1}{3}, \quad \frac{q_i}{Q} \leq \frac{3P}{Q} < \frac{1}{3}, \quad \text{and} \quad qq_i |\theta_i| \leq \frac{q}{3P} \leq \frac{1}{3},$$

we have

$$\begin{aligned} qq_i |\theta_i| + \frac{q_i}{Q} + qq_i \Delta \alpha_i < 1 &\implies |\theta_i| + \frac{1}{qQ} + \Delta \alpha_i < \frac{1}{qq_i} \leq \left| \frac{a}{q} - \frac{a_i}{q_i} \right| \\ &\implies \left| \frac{a}{q} - \alpha_i \right| > \frac{1}{qQ} + \Delta \alpha_i. \end{aligned}$$

Thus,

$$\mathfrak{M}\left(\frac{a}{q}\right) \cap [\alpha_i(1-\Delta), \alpha_i(1+\Delta)] = \emptyset.$$

Therefore, $\mathfrak{M}(a/q) \cap I_{k_0}$ does not depend on k_0 ; in fact, recalling that $E = [\alpha_2^{-2}, \alpha_1^{-2}]$, we have

$$\mathfrak{M}\left(\frac{a}{q}\right) \cap I_{k_0} = \begin{cases} \mathfrak{M}(a/q) & \text{if } (a/q)^{-2} \in E, \\ \emptyset & \text{if } (a/q)^{-2} \notin E. \end{cases}$$

We split the integral over I_{k_0} in (4-5) as

$$\begin{aligned} \int_{I_{k_0}} \sum_{t \in \mathbb{Z}} L(1, \tilde{\psi}_t) \cos(2\pi\alpha t) \widehat{W}\left(\frac{t}{x_{k_0}(\alpha)}\right) \frac{d\alpha}{\alpha^3} \\ = \int_{\mathfrak{m}_{k_0}} + \int_{\mathfrak{M}_{k_0}} = \int_{\mathfrak{m}_{k_0}} + \sum_{\substack{a \in (\mathbb{Q} \setminus \{\frac{a_1}{q_1}, \frac{a_2}{q_2}\}) \cap [\alpha_1, \alpha_2] \\ q \leq P}} \int_{\mathfrak{M}(\frac{a}{q})} + \sum_{i=1}^2 \int_{\mathfrak{M}_{k_0} \cap \mathfrak{M}(\frac{a_i}{q_i})}. \end{aligned}$$

Note that the integral over $\mathfrak{M}_{k_0} \cap \mathfrak{M}(a_i/q_i)$ vanishes if $q_i > P$.

We evaluate the terms of this sum using Proposition 5.1. By Dirichlet’s theorem, for $\alpha \in I_{k_0}$ we may choose $q \leq Q$ and a coprime to q such that

$$\left| \alpha - \frac{a}{q} \right| \leq \frac{1}{qQ}.$$

If $\alpha \in \mathfrak{m}_{k_0}$ then $q > P$, in which case $x_{k_0}(\alpha) \left| \alpha - \frac{a}{q} \right| \ll_E 1$, so by Proposition 5.1,

$$\begin{aligned} \sum_{t \in \mathbb{Z}} L(1, \tilde{\psi}_t) \cos(2\pi\alpha t) \widehat{W}\left(\frac{t}{x_{k_0}(\alpha)}\right) &\ll_{E,\varepsilon} \frac{x_{k_0}(\alpha)}{P^{3-\varepsilon}} + Qx_{k_0}(\alpha)^{-1} + Q^3x_{k_0}(\alpha)^{-\frac{7}{4}+\varepsilon} \\ &\ll_{E,\varepsilon} h^{-1}H^{\frac{3}{2}}K^{-\frac{1}{2}+\varepsilon} + H^{\frac{1}{2}}K^{-\frac{1}{2}} + h^{-\frac{5}{4}}H^{\frac{3}{2}}K^{-\frac{1}{4}+\varepsilon} \\ &\ll h^{-\frac{5}{4}}H^{\frac{3}{2}}K^{-\frac{1}{4}+\varepsilon}. \end{aligned}$$

Note that this estimate also applies to the error terms in Proposition 5.1 for α in a major arc $\mathfrak{M}(\frac{a}{q})$ with $|\alpha - \frac{a}{q}|x_{k_0}(\alpha) \leq 1$. For $\alpha \in \mathfrak{M}(\frac{a}{q})$ with $|\alpha - \frac{a}{q}|x_{k_0}(\alpha) > 1$, the error term is

$$\ll_{\varepsilon} Q^{-1} + q^{-\frac{1}{2}}Q^{-\frac{7}{2}}x_{k_0}(\alpha)^{\frac{7}{4}+\varepsilon} \ll_{E,\varepsilon} Q^{-1} + q^{-\frac{1}{2}}Q^{-\frac{7}{2}}(K/h)^{\frac{7}{4}+\varepsilon}.$$

Summing this error term times the measure of each major arc gives

$$\ll_{E,\varepsilon} PQ^{-2} + P^{\frac{1}{2}}Q^{-\frac{9}{2}}(K/h)^{\frac{7}{4}+\varepsilon} \ll_{E,\varepsilon} h^2H^{-\frac{3}{2}}K^{-\frac{1}{2}} + h^{\frac{11}{4}}H^{-\frac{5}{2}}K^{-\frac{1}{4}+\varepsilon},$$

which is dominated by the minor arc error.

Multiplying by $\frac{2h}{\pi} \left(\frac{k_0-1}{4\pi}\right)^2$, summing over the $J \ll H/h$ choices of k_0 , combining with the other error terms in (4-5), and using $h \asymp \max(H^{\frac{10}{9}}K^{-\frac{1}{9}}, (HK)^{\frac{5}{11}})$, we get

$$\ll_{E,\varepsilon} K^{\varepsilon} (hK^2 + h^{-\frac{6}{5}}HK^3 + h^{-\frac{5}{4}}H^{\frac{5}{2}}K^{\frac{7}{4}}) \ll HK^{2+\varepsilon} (H^{\frac{1}{9}}K^{-\frac{1}{9}} + H^{-\frac{6}{11}}K^{\frac{5}{11}}). \tag{6-1}$$

We turn now to the main term in Proposition 5.1 on the major arcs. Note that for $\alpha \in I_{k_0}$, we have $x_{k_0}(\alpha) \geq x_{k_0}(\lambda_{k_0}\alpha_2) = PQ$. Therefore, when $(a/q)^{-2} \in E$ and $a/q \notin \{a_1/q_1, a_2/q_2\}$, we have

$$\left\{ x_{k_0}(\alpha) \left(\alpha - \frac{a}{q} \right) : \alpha \in \mathfrak{M}\left(\frac{a}{q}\right) \right\} \supseteq [-1, 1].$$

Hence, making the change of variables $x = x_{k_0}(\alpha)(\alpha - \frac{a}{q})$, we obtain

$$\begin{aligned} \int_{\mathfrak{M}(\frac{a}{q})} x_{k_0}(\alpha) W\left(x_{k_0}(\alpha)\left(\alpha - \frac{a}{q}\right)\right) \frac{d\alpha}{\alpha^3} &= \left(\frac{a}{q}\right)^{-3} \int_{\mathbb{R}} W(x) \left(1 - \frac{4hx}{k_0-1}\right)^2 dx \\ &= \left(\frac{a}{q}\right)^{-3} \left(1 + \left(\frac{4h}{k_0-1}\right)^2 \int_{\mathbb{R}} x^2 W(x) dx\right). \end{aligned}$$

Multiplying by $\frac{2h}{\pi} \left(\frac{k_0-1}{4\pi}\right)^2$ and summing over $k_0 = K_0 + 4hj$, we get

$$\left(\frac{a}{q}\right)^{-3} \frac{2h^3 J}{\pi^3} \left\{ \left(\frac{J-1}{2} + \frac{K_0-1}{4h}\right)^2 + \frac{J^2-1}{12} + \int_{\mathbb{R}} x^2 W(x) dx \right\}.$$

Since $J = \left\lfloor \frac{K+H-K_0}{4h} \right\rfloor = \frac{H}{2h} + O(1)$ and $\frac{J-1}{2} + \frac{K_0-1}{4h} = \frac{K}{4h} + O(1)$, this is

$$\left(\frac{a}{q}\right)^{-3} \frac{1}{\pi^3} (H + O(2h)) \left\{ \frac{K^2}{16} + \frac{H^2}{48} + O(hK) + O(h^2) + h^2 \int_{\mathbb{R}} x^2 W(x) dx \right\} = \left(\frac{a}{q}\right)^{-3} \frac{HK^2 + O(hK^2)}{16\pi^3}.$$

Multiplying by $\frac{\mu(q)^2}{\varphi(q)^2 \sigma(q)^2}$ and summing over the major arcs (excluding a_i/q_i), we get

$$\frac{HK^2}{16\pi^3} \sum_{\substack{a \in (\mathbb{Q} \setminus \{\frac{a_1}{q_1}, \frac{a_2}{q_2}\}) \cap [\alpha_1, \alpha_2] \\ q \leq P}} \frac{\mu(q)^2}{\varphi(q)^2 \sigma(q)^2} \left(\frac{a}{q}\right)^{-3} + O_E(hK^2).$$

Since $h \ll H^{\frac{10}{9}} K^{-\frac{1}{9}}$ and

$$\sum_{\substack{a \in \mathbb{Q} \cap [\alpha_1, \alpha_2] \\ q > P}} \frac{\mu(q)^2}{\varphi(q)^2 \sigma(q)^2} \left(\frac{a}{q}\right)^{-3} \ll_E \frac{1}{P} \ll_E \left(\frac{H}{K}\right)^{\frac{1}{2}},$$

we can write this as

$$\frac{HK^2}{16\pi^3} \sum_{a \in (\mathbb{Q} \setminus \{\frac{a_1}{q_1}, \frac{a_2}{q_2}\}) \cap [\alpha_1, \alpha_2]} \frac{\mu(q)^2}{\varphi(q)^2 \sigma(q)^2} \left(\frac{a}{q}\right)^{-3} + O_E(H^{\frac{10}{9}} K^{\frac{17}{9}}). \tag{6-2}$$

On RH, we have

$$\frac{1}{\sqrt{N}} \sum_{\substack{p \text{ prime} \\ p/N \in E}} \log p \sum_{\substack{k \equiv 2\delta \pmod{4} \\ |k-K| \leq H}} \sum_{f \in H_k(1)} 1 = \frac{HK^2}{96\pi} |E| + O(K^2). \tag{6-3}$$

Adding the error from (6-1) to (6-2) and dividing by (6-3) gives

$$\frac{1}{\zeta(2)|E|} \sum_{a \in (\mathbb{Q} \setminus \{\frac{a_1}{q_1}, \frac{a_2}{q_2}\}) \cap [\alpha_1, \alpha_2]} \frac{\mu(q)^2}{\varphi(q)^2 \sigma(q)^2} \left(\frac{a}{q}\right)^{-3} + O_{E,\varepsilon} \left(K^\varepsilon (H^{\frac{1}{9}} K^{-\frac{1}{9}} + H^{-\frac{6}{11}} K^{\frac{5}{11}}) \right), \tag{6-4}$$

which is the expected answer, apart from a possible difference in the contributions from a_i/q_i .

When α_i is irrational, we bound the discrepancy trivially as

$$\ll_E \frac{\mu(q_i)^2}{\varphi(q_i)^2 \sigma(q_i)} \ll_\varepsilon q_i^{-3+\varepsilon}.$$

This tends to 0 as $K \rightarrow \infty$, but we have no control over the rate of convergence in general. However, if α_i has irrationality measure $\mu_i \leq 27$ then this is dominated by the other error terms.

In more detail, suppose $(a_i/q_i)^{-2} \in E$ but the major arc integral does not always cover the full support of W , which happens when

$$(-1)^i x_{k_0}(\lambda_{k_0} \alpha_i) \left(\lambda_{k_0} \alpha_i - \frac{a_i}{q_i} \right) < 1$$

for some k_0 . Rearranging this inequality, we find

$$\left| \alpha_i - \frac{a_i}{q_i} \right| < \alpha_i \left(\frac{h}{\pi \sqrt{N}} + |\lambda_{k_0} - 1| \right) < 2\Delta \alpha_i \ll_E \frac{H}{K}.$$

A similar calculation leads to the same inequality when $(a_i/q_i)^{-2} \notin E$ but a major arc overlaps the support of W for some k_0 . On the other hand, by the definition of irrationality measure, we have $|\alpha_i - a_i/q_i| \geq q_i^{-\mu_i + o(1)}$ as $K \rightarrow \infty$, whence

$$\frac{\mu(q_i)^2}{\varphi(q_i)^2 \sigma(q_i)} \leq q_i^{-3+o(1)} \leq \left(\frac{H}{K} \right)^{\frac{3}{\mu_i} - o(1)}.$$

If $\mu_i \leq 27$ this is dominated by the error term in (6-4).

When α_i is rational, we have $a_i/q_i = \alpha_i$ for sufficiently large K . Writing $x = x_{k_0}(\alpha)(\alpha - \alpha_i)$, for $\alpha \in I_{k_0}$ we have

$$\begin{aligned} (-1)^i (\lambda_{k_0} \alpha_i - \alpha) \geq 0 &\implies (-1)^i \left(\frac{(k_0 - 1)\alpha_i}{4h\alpha} - \frac{k_0 - 1}{4h\lambda_{k_0}} \right) \geq 0 \\ &\implies (-1)^i \left(-x + \frac{k_0 - 1}{4h} - \frac{k_0 - 1}{4h\lambda_{k_0}} \right) \geq 0 \\ &\implies (-1)^i x \leq (-1)^i \frac{k_0 - 1 - 4\pi\sqrt{N}}{4h}. \end{aligned}$$

Thus,

$$\int_{\mathfrak{M}(\frac{a_1}{q_1}) \cap I_{k_0}} = \int_{\frac{k_0 - 1 - 4\pi\sqrt{N}}{4h}}^{\infty} \quad \text{and} \quad \int_{\mathfrak{M}(\frac{a_2}{q_2}) \cap I_{k_0}} = \int_{-\infty}^{\frac{k_0 - 1 - 4\pi\sqrt{N}}{4h}}.$$

Recall that $k_0 = K_0 + 4hj$, where $j = 0, \dots, J - 1$. Writing $j_0 = \left\lfloor \frac{4\pi\sqrt{N} + 1 - K_0}{4h} \right\rfloor$, we have

$$\frac{(K_0 + 4jh) - 1 - 4\pi\sqrt{N}}{4h} \in (-1, 1] \iff j \in \{j_0, j_0 + 1\}.$$

Thus, for $i = 1$ we get the full integral for $j < j_0$, 0 for $j > j_0 + 1$, and something in between for $j \in \{j_0, j_0 + 1\}$; similarly, for $i = 2$ we get 0 for $j < j_0$, the full integral for $j > j_0 + 1$, and something in between for $j \in \{j_0, j_0 + 1\}$.

We calculate that $j_0 = J/2 + O(1)$. Thus, we get half the major arc contribution from a_i/q_i , up to a relative error $\ll 1/J \ll h/H \ll (H/K)^{\frac{1}{9}}$.

Acknowledgements

We thank Peter Sarnak and Nina Zubrilina for insightful comments, and Andrew Sutherland for helping to popularize this topic and sharing his data. This project began at the workshop “Murmurations in Arithmetic” and the conference “LMFDB, Computation, and Number Theory”, both held at ICERM in July 2023. We thank ICERM and the organizers for their hospitality.

References

- [BL17] A. R. Booker and M. Lee, “The Selberg trace formula as a Dirichlet series”, *Forum Math.* **29**:3 (2017), 519–542. MR
- [Boo15] A. R. Booker, “ L -functions as distributions”, *Math. Ann.* **363**:1-2 (2015), 423–454. MR
- [Chi22] K. Child, “Twist-minimal trace formula for holomorphic cusp forms”, *Res. Number Theory* **8**:1 (2022), art. id. 11, 27 pp. MR
- [Coh77] H. Cohen, “Trace des opérateurs de Hecke sur $\Gamma_0(N)$ ”, exposé 4, 9 pp. in *Séminaire de Théorie des Nombres (1976–1977)*, Centre National de Recherche Scientifique, Laboratoire de Théorie des Nombres, Talence, 1977. MR
- [Eic57] M. Eichler, “Eine Verallgemeinerung der Abelschen Integrale”, *Math. Z.* **67** (1957), 267–298. MR
- [Hij74] H. Hijikata, “Explicit formula of the traces of Hecke operators for $\Gamma_0(N)$ ”, *J. Math. Soc. Japan* **26** (1974), 56–82. MR
- [HLOP22] Y.-H. He, K.-H. Lee, T. Oliver, and A. Pozdnyakov, “Murmurations of elliptic curves”, 2022. arXiv 2204.10140
- [HLOPS] Y.-H. He, K.-H. Lee, T. Oliver, A. Pozdnyakov, and A. V. Sutherland, “Murmurations of L -functions”, 2026. In preparation.
- [ILS00] H. Iwaniec, W. Luo, and P. Sarnak, “Low lying zeros of families of L -functions”, *Inst. Hautes Études Sci. Publ. Math.* **91** (2000), 55–131. MR
- [KS99] N. M. Katz and P. Sarnak, “Zeroes of zeta functions and symmetry”, *Bull. Amer. Math. Soc. (N.S.)* **36**:1 (1999), 1–26. MR
- [LMFDB] The L -functions and modular forms database, “Analytic conductor of a classical newform”, 2023, available at https://www.lmfdb.org/knowledge/show/cmf.analytic_conductor.
- [MV07] H. L. Montgomery and R. C. Vaughan, *Multiplicative number theory, I: Classical theory*, vol. 97, Cambridge Studies in Advanced Mathematics, Cambridge University Press, Cambridge, 2007. MR
- [Sar23] P. Sarnak, Letter to Drew Sutherland and Nina Zubrilina on murmurations and root numbers, 2023, available at <https://publications.ias.edu/sarnak/paper/2726>.
- [Sut23] A. V. Sutherland, “Murmurations of arithmetic L -functions”, presentation in the Number Theory Web Seminar, 2023, available at <https://www.ntwebseminar.org/previous-talks#h.tjmodfmt53rl>.
- [SvdV91] R. Schoof and M. van der Vlugt, “Hecke operators and the weight distributions of certain codes”, *J. Combin. Theory Ser. A* **57**:2 (1991), 163–186. MR
- [Zag77] D. Zagier, “Traces des opérateurs de Hecke”, exposé 23, 12 pp. in *Séminaire Delange–Pisot–Poitou, 17e année: 1975/76, Théorie des nombres 2*, Secrétariat Math., Paris, 1977. MR
- [Zub23] N. Zubrilina, “Murmurations”, preprint, 2023. arXiv 2310.07681

Communicated by Roger Heath-Brown

Received 2024-02-01 Revised 2024-11-07 Accepted 2025-02-11

j.bober@bristol.ac.uk *School of Mathematics, University of Bristol, Bristol, BS8 1UG, United Kingdom*

andrew.booker@bristol.ac.uk *School of Mathematics, University of Bristol, Bristol, BS8 1UG, United Kingdom*

min.lee@bristol.ac.uk *School of Mathematics, University of Bristol, Bristol, BS8 1UG, United Kingdom*

Algebra & Number Theory

msp.org/ant

EDITORS

MANAGING EDITOR
Antoine Chambert-Loir
Université Paris-Diderot
France

EDITORIAL BOARD CHAIR
David Eisenbud
University of California
Berkeley, USA

BOARD OF EDITORS

Jason P. Bell	University of Waterloo, Canada	Philippe Michel	École Polytechnique Fédérale de Lausanne
Bhargav Bhatt	University of Michigan, USA	Martin Olsson	University of California, Berkeley, USA
Frank Calegari	University of Chicago, USA	Irena Peeva	Cornell University, USA
J.-L. Colliot-Thélène	CNRS, Université Paris-Saclay, France	Jonathan Pila	University of Oxford, UK
Brian D. Conrad	Stanford University, USA	Anand Pillay	University of Notre Dame, USA
Samit Dasgupta	Duke University, USA	Bjorn Poonen	Massachusetts Institute of Technology, USA
Hélène Esnault	Freie Universität Berlin, Germany	Victor Reiner	University of Minnesota, USA
Gavril Farkas	Humboldt Universität zu Berlin, Germany	Peter Sarnak	Princeton University, USA
Sergey Fomin	University of Michigan, USA	Michael Singer	North Carolina State University, USA
Edward Frenkel	University of California, Berkeley, USA	Vasudevan Srinivas	SUNY Buffalo, USA
Wee Teck Gan	National University of Singapore	Shunsuke Takagi	University of Tokyo, Japan
Andrew Granville	Université de Montréal, Canada	Pham Huu Tiep	Rutgers University, USA
Ben J. Green	University of Oxford, UK	Ravi Vakil	Stanford University, USA
Christopher Hacon	University of Utah, USA	Akshay Venkatesh	Institute for Advanced Study, USA
Roger Heath-Brown	Oxford University, UK	Melanie Matchett Wood	Harvard University, USA
János Kollár	Princeton University, USA	Shou-Wu Zhang	Princeton University, USA
Michael J. Larsen	Indiana University Bloomington, USA		

PRODUCTION

production@msp.org
Silvio Levy, Scientific Editor

See inside back cover or msp.org/ant for submission instructions.

The subscription price for 2026 is US \$590/year for the electronic version, and \$865/year (+\$75, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to MSP.

Algebra & Number Theory (ISSN 1944-7833 electronic, 1937-0652 printed) at Mathematical Sciences Publishers, 2000 Allston Way # 59, Berkeley, CA 94701-4004, is published continuously online.

ANT peer review and production are managed by EditFLOW[®] from MSP.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2026 Mathematical Sciences Publishers

Algebra & Number Theory

Volume 20 No. 2 2026

On the minimal dimension of a faithful linear representation of a finite group	219
ALEXANDER MORETÓ	
Moments of one-level densities in families of holomorphic cusp forms in the level aspect	237
PETER COHEN, JUSTINE DELL, OSCAR E. GONZÁLEZ, SIMRAN KHUNGER, CHUNG-HANG KWAN, STEVEN J. MILLER, ALEXANDER SHASHKOV, ALICIA REINA SMITH, CARSTEN SPRUNGER, NICHOLAS TRIAANTAFILLOU, NHI TRUONG, ROGER VAN PESKI and STEPHEN WILLIS	
Murmurations of modular forms in the weight aspect	299
JONATHAN BOBER, ANDREW R. BOOKER, MIN LEE and DAVID LOWRY-DUDA	
The Alperin weight conjecture and the Glauberman correspondence via character triples	333
J. MIQUEL MARTÍNEZ, NOELIA RIZO and DAMIANO ROSSI	
Asymptotics of extensions of simple \mathbb{Q} -algebras	383
FABIAN GUNDLACH and BÉRANGER SEGUIN	