

Algebra & Number Theory

Volume 20
2026
No. 5

**On the cohomology of tautological bundles
over Quot schemes of curves**

Alina Marian, Dragos Oprea and Steven V Sam



On the cohomology of tautological bundles over Quot schemes of curves

Alina Marian, Dragos Oprea and Steven V Sam

We consider tautological bundles and their exterior and symmetric powers on the Quot scheme over the projective line. We prove and conjecture several statements regarding the vanishing of their higher cohomology, and we describe their spaces of global sections via tautological constructions. To this end, we make use of the embedding of the Quot scheme as an explicit local complete intersection in the product of two Grassmannians, studied by Strømme. This allows us to construct resolutions with vanishing cohomology for the tautological bundles and their exterior and symmetric powers. We further illustrate our approach with a few additional cohomological calculations.

1. Introduction

1.1. Tautological vector bundles. We consider the Quot scheme $\text{Quot}_{\mathbb{P}^1}(\mathbb{C}^N, n, r)$ parametrizing rank- r degree- n quotients of the trivial bundle of rank N over \mathbb{P}^1 :

$$0 \rightarrow S \rightarrow \mathbb{C}^N \otimes \mathcal{O}_{\mathbb{P}^1} \rightarrow Q \rightarrow 0, \quad \text{rank } Q = r, \quad \text{deg } Q = n.$$

It is a smooth projective variety which comes equipped with the universal sequence

$$0 \rightarrow S \rightarrow \mathbb{C}^N \otimes \mathcal{O} \rightarrow Q \rightarrow 0$$

over $\mathbb{P}^1 \times \text{Quot}_{\mathbb{P}^1}(\mathbb{C}^N, n, r)$. We let p and π denote the projections to the two factors.

The Quot scheme over the projective line is an important testing ground for ideas in moduli theory. It has rich geometry and bears ties to homogeneous and quiver varieties while not being one of them. A beautiful systematic study of $\text{Quot}_{\mathbb{P}^1}(\mathbb{C}^N, n, r)$ was carried out in [Str], where the Quot scheme was shown to be rational, and the Betti numbers, generators for the Chow ring, and the nef cone were calculated. As noted in [Str], $\text{Quot}_{\mathbb{P}^1}(\mathbb{C}^N, n, r)$ is a compactification of the space of degree- d morphisms from \mathbb{P}^1 to the Grassmannian $\mathbf{G}(N, r)$ of r -dimensional quotients of \mathbb{C}^N . With this point of view, in the 1990s, the Quot scheme was used effectively to calculate the small quantum cohomology ring of the Grassmannian, leading eventually to a calculation for all flag varieties [Be; CF1; CF2; K; C]. Further progress included the description of the equivariant cohomology ring in [BCS], the calculation of the effective cone [J], and the birational study [I] which established $\text{Quot}_{\mathbb{P}^1}(\mathbb{C}^N, n, r)$ as a Mori dream space, among others.

MSC2020: 14H60, 14M99.

Keywords: Quot scheme, sheaf cohomology.

In this paper, we take up the problem of calculating the cohomology of Schur functors of tautological bundles over $\text{Quot}_{\mathbb{P}^1}(\mathbb{C}^N, n, r)$. While our results are primarily in the setting of zero quotient rank ($r = 0$), the method is available for any rank. Conjectures for any r are formulated in [Section 1.3](#) below.

To start, note that for any line bundle $L \rightarrow \mathbb{P}^1$, there is an induced tautological complex of rank $n + r(\deg L + 1)$ over $\text{Quot}_{\mathbb{P}^1}(\mathbb{C}^N, r, n)$, given by

$$L^{[n]} = R\pi_*(p^*L \otimes \mathcal{Q}). \tag{1.1.1}$$

When $r = 0$, $L^{[n]}$ is a *vector bundle* of rank n . (Note that $R^1\pi_*(p^*L \otimes \mathcal{Q}) = 0$ for $r = 0$ since the support of \mathcal{Q} is finite in each fiber of π .) We let $\text{Quot}_{\mathbb{P}^1}(\mathbb{C}^N, n)$ denote the Quot scheme in this case, and show the following results.

Theorem 1.1.2. (1) *For all line bundles $L \rightarrow \mathbb{P}^1$ with $\deg L \geq n \geq k$, we have*

$$H^0(\text{Quot}_{\mathbb{P}^1}(\mathbb{C}^N, n), \wedge^k L^{[n]}) \cong \wedge^k H^0(L^{\oplus N})$$

and the higher cohomology vanishes:

$$H^i(\text{Quot}_{\mathbb{P}^1}(\mathbb{C}^N, n), \wedge^k L^{[n]}) = 0, \quad i > 0.$$

(2) *More generally, assume $\deg L \geq n \geq k$ and let p_1, \dots, p_t be nonnegative integers, $0 \leq t \leq N - 1$.*

We have

$$\text{Ext}^i(\wedge^{p_1} L^{[n]} \otimes \dots \otimes \wedge^{p_t} L^{[n]}, \wedge^k L^{[n]}) = \begin{cases} \wedge^{k-|p|} H^0(L^{\oplus N}) & \text{if } i = 0, \\ 0 & \text{if } i > 0, \end{cases}$$

for $|p| = p_1 + \dots + p_t \leq k$. If $|p| > k$, all the above Ext groups vanish.

Theorem 1.1.3. (1) *For all line bundles $L \rightarrow \mathbb{P}^1$ with $\deg L \geq n \geq k$, we have*

$$H^0(\text{Quot}_{\mathbb{P}^1}(\mathbb{C}^N, n), \text{Sym}^k L^{[n]}) \cong \text{Sym}^k H^0(L^{\oplus N})$$

and the higher cohomology vanishes

$$H^i(\text{Quot}_{\mathbb{P}^1}(\mathbb{C}^N, n), \text{Sym}^k L^{[n]}) = 0, \quad i > 0.$$

(2) *More generally, assume $\deg L \geq n \geq k$ and let p_1, \dots, p_t be nonnegative integers, $0 \leq t \leq N - 1$.*

Then

$$\text{Ext}^i(\wedge^{p_1} L^{[n]} \otimes \dots \otimes \wedge^{p_t} L^{[n]}, \text{Sym}^k L^{[n]}) = \begin{cases} \text{Sym}^{k-|p|} H^0(L^{\oplus N}) & \text{if } i = 0 \text{ and all } p_j \in \{0, 1\}, \\ 0 & \text{otherwise,} \end{cases}$$

for $|p| \leq k$. If $|p| > k$, all the above Ext groups vanish.

We expect that the vanishing of higher cohomology of $\wedge^k L^{[n]}$ and $\text{Sym}^k L^{[n]}$ in [Theorems 1.1.2](#) and [1.1.3](#) holds whenever $\deg L \geq -1$. For arbitrary exterior powers, the above vanishings cannot be accessed by the classical theorems. In the special case of the determinant line bundle $\wedge^n L^{[n]}$, the bound in our theorems improves the bound $\deg L \geq Nn - N - n$ obtained by Kodaira vanishing. The latter can be

applied using the description of the ample cone in [Str] and a standard calculation of the canonical bundle via Grothendieck–Riemann–Roch.

It is easy to see how the sections of the bundles $\wedge^k L^{[n]}$ and $\text{Sym}^k L^{[n]}$ over $\text{Quot}_{\mathbb{P}^1}(\mathbb{C}^N, n)$ arise geometrically. Indeed, from the universal quotient

$$\mathbb{C}^N \otimes \mathcal{O} \rightarrow \mathcal{Q} \rightarrow 0$$

over $\mathbb{P}^1 \times \text{Quot}_{\mathbb{P}^1}(\mathbb{C}^N, n)$, after tensoring by L and pushing forward, we immediately obtain a map

$$H^0(L^{\oplus N}) \otimes \mathcal{O}_{\text{Quot}} \rightarrow L^{[n]}.$$

Taking exterior and symmetric powers, and taking cohomology, we obtain morphisms

$$\Phi_k : \wedge^k H^0(L^{\oplus N}) \rightarrow H^0(\wedge^k L^{[n]}), \quad \Psi_k : \text{Sym}^k H^0(L^{\oplus N}) \rightarrow H^0(\text{Sym}^k L^{[n]}). \quad (1.1.4)$$

Our proofs will show that Φ_k and Ψ_k are isomorphisms when $\deg L \geq n \geq k$. Thus all sections of $\wedge^k L^{[n]}$ and $\text{Sym}^k L^{[n]}$ are obtained tautologically, while the higher cohomology vanishes.

We further illustrate the techniques developed here by showing that:

Theorem 1.1.5. *For all line bundles $L, M \rightarrow \mathbb{P}^1$ with $\deg M \geq n$ and $0 \leq \deg M - \deg L \leq 1$, for all $p_1, \dots, p_t \geq 0$, not all zero, and $1 \leq t \leq N - 1$, we have*

$$H^i(\text{Quot}_{\mathbb{P}^1}(\mathbb{C}^N, n), (\wedge^{p_1} L^{[n]})^\vee \otimes (\wedge^{p_2} M^{[n]})^\vee \otimes \dots \otimes (\wedge^{p_t} M^{[n]})^\vee) = 0, \quad i \geq 0.$$

Of course, the case $L = M$ is contained in Theorems 1.1.2(2) and 1.1.3(2) for $k = 0$.

Although we focus for notational simplicity on the case of quotients of the trivial bundle, the same arguments easily apply to the Quot scheme $\text{Quot}_{\mathbb{P}^1}(E, n)$ of finite quotients of an arbitrary vector bundle $E \rightarrow \mathbb{P}^1$.

Let a denote the largest degree appearing in the splitting of E as a direct sum of line bundles, and set $\alpha = -\deg E + (N - 1)a$. For Theorems 1.1.2 and 1.1.3, we replace all instances of $H^0(L^{\oplus N})$ by $H^0(E \otimes L)$. For instance, Theorem 1.1.2(1) takes the form

$$H^0(\text{Quot}_{\mathbb{P}^1}(E, n), \wedge^k L^{[n]}) \cong \wedge^k H^0(E \otimes L), \quad H^i(\text{Quot}_{\mathbb{P}^1}(E, n), \wedge^k L^{[n]}) = 0, \quad i > 0,$$

whenever

$$\deg L \geq n + \alpha, \quad n \geq k \geq 0.$$

Remark 2.1.2 below explains how this bound emerges. Similarly, the analogue of Theorem 1.1.3(1) reads

$$H^0(\text{Quot}_{\mathbb{P}^1}(E, n), \text{Sym}^k L^{[n]}) \cong \text{Sym}^k H^0(E \otimes L), \quad H^i(\text{Quot}_{\mathbb{P}^1}(E, n), \text{Sym}^k L^{[n]}) = 0, \quad i > 0.$$

The more general Theorem 1.1.2(2) and Theorem 1.1.3(2) are also correct after the analogous modifications. Likewise, Theorem 1.1.5 remains true under the assumption $\deg M \geq n + \alpha$.

For a smooth projective curve C of arbitrary genus, the holomorphic Euler characteristics of $\wedge^k L^{[n]}$, $\text{Sym}^k L^{[n]}$ and $(\wedge^p L^{[n]})^\vee$ on $\text{Quot}_C(\mathbb{C}^N, n)$ were calculated in [OS] by equivariant localization. The following expectation regarding individual cohomology groups was also formulated in [OS, Question 20]:

$$H^\bullet(\text{Quot}_C(\mathbb{C}^N, n), \wedge^k L^{[n]}) \cong \wedge^k H^\bullet(L^{\oplus N}) \otimes \text{Sym}^{n-k} H^\bullet(\mathcal{O}_C). \tag{1.1.6}$$

The exterior and symmetric powers on the right are understood in the graded sense.

In the case when $C = \mathbb{P}^1$, our theorems confirm this expectation for $\text{deg } L \geq n$. We offer additional modest evidence for $k = 1$ and L of arbitrary degree:

Corollary 1.1.7. *For all line bundles $L \rightarrow \mathbb{P}^1$, we have*

$$H^i(\text{Quot}_{\mathbb{P}^1}(\mathbb{C}^N, n), L^{[n]}) = 0, \quad i \geq 2.$$

These results and formula (1.1.6) reflect a full parallelism to the cohomology of tautological bundles over the Hilbert scheme of points on a surface computed in [D; Sc1; Sc2; Kr1; A]. For instance, for all line bundles $L \rightarrow X$ over smooth projective surfaces, we have

$$H^\bullet(X^{[n]}, \wedge^k L^{[n]}) = \wedge^k H^\bullet(X, L) \otimes \text{Sym}^{n-k} H^\bullet(X, \mathcal{O}_X).$$

The Bridgeland–King–Reid correspondence plays a central role in the proof. We refer the reader to the beautiful article [Kr1] for state-of-the-art calculations in the surface case.

1.2. Proofs. To establish Theorems 1.1.2, 1.1.3 and 1.1.5, the key idea is to use the twofold Grothendieck embedding of the Quot scheme into a product of Grassmannians,

$$\iota : \text{Quot}_{\mathbb{P}^1}(\mathbb{C}^N, n) \hookrightarrow \mathbf{G}_1 \times \mathbf{G}_2,$$

so that the image of ι is the zero locus of a regular section σ of an explicit homogeneous vector bundle

$$\mathcal{E} \rightarrow \mathbf{G}_1 \times \mathbf{G}_2.$$

This embedding as an explicit local complete intersection is specific to the Quot scheme (of quotients of all ranks) over the projective line. It was considered and studied in detail by Strømme [Str], who used it to derive information about the Chow ring. Our paper widens the picture, and shows that the embedding is also well-suited to the study of the tautological bundles.

Using the Koszul resolution for σ ,

$$\dots \rightarrow \wedge^2 \mathcal{E}^\vee \rightarrow \wedge^1 \mathcal{E}^\vee \rightarrow \mathcal{O}_{\mathbf{G}_1 \times \mathbf{G}_2} \rightarrow \mathcal{O}_{\text{Quot}} \rightarrow 0,$$

we obtain resolutions

$$\dots \rightarrow \mathcal{R}_2 \rightarrow \mathcal{R}_1 \rightarrow \mathcal{R}_0 \rightarrow \iota_* \mathcal{F} \rightarrow 0$$

for each one of the tautological bundles \mathcal{F} appearing in Theorems 1.1.2, 1.1.3 and 1.1.5. The resolutions thus obtained are special. Remarkably, we show that the terms \mathcal{R}_ℓ have vanishing cohomology for all

$\ell \geq 1$, while \mathcal{R}_0 has no higher cohomology. This allows us to control the cohomology of the tautological bundles and establish our results.

The argument makes crucial use of the Borel–Weil–Bott theorem on the two Grassmannians $\mathbf{G}_1, \mathbf{G}_2$, along with several combinatorial arguments involving the Littlewood–Richardson rule. In intermediate stages, statements of independent interest are established generally over arbitrary Grassmannians; we refer the reader to [Section 3](#) for details. It takes a delicate interplay between Borel–Weil–Bott and Littlewood–Richardson vanishings to show that all higher terms $\mathcal{R}_\ell, \ell \geq 1$, of the resolution have no cohomology at all.

While the above theorems concern genus 0, we also obtain the following corollary in arbitrary genus. Let y be a variable. Setting

$$\bigwedge_y V := \sum_k y^k \bigwedge^k V, \quad \text{Sym}_y V := \sum_k y^k \text{Sym}^k V,$$

the result below recovers [Theorem 1](#), a special case of [Theorem 2](#), and [Theorem 4](#) in [\[OS\]](#).

Corollary 1.2.1. *Let $L, M_1, \dots, M_t \rightarrow C$ be line bundles over a smooth projective curve, where $1 \leq t \leq N - 1$. Then*

$$\sum_{n=0}^{\infty} q^n \chi \left(\text{Quot}_C(\mathbb{C}^N, n), \bigwedge_y L^{[n]} \right) = (1 - q)^{-\chi(\mathcal{O}_C)} (1 + qy)^{N\chi(L)}, \tag{1.2.1a}$$

$$\sum_{n=0}^{\infty} q^n \chi \left(\text{Quot}_C(\mathbb{C}^N, n), \bigotimes_{i=1}^t \bigwedge_{y_i} M_i^{[n]} \vee \right) = (1 - q)^{-\chi(\mathcal{O}_C)}. \tag{1.2.1b}$$

Furthermore, in genus 0, we have

$$\sum_{n \geq k} q^n y^k \chi \left(\text{Quot}_{\mathbb{P}^1}(\mathbb{C}^N, n), \text{Sym}^k L^{[n]} \right) = (1 - q)^{-1} (1 - qy)^{-N\chi(L)}. \tag{1.2.1c}$$

We will show how to derive [Corollary 1.2.1](#) from [Theorems 1.1.2, 1.1.3 and 1.1.5](#) in [Section 4.1](#).

Formulas [\(1.2.1a\)](#), [\(1.2.1b\)](#) and [\(1.2.1c\)](#) were previously established in [\[OS\]](#) based on reduction to genus 0 using universality statements as in [\[EGL; OS; St\]](#), and equivariant torus localization in genus 0. The localization calculation is combinatorially involved and relies on several mysterious simplifications. In the present paper, [Theorems 1.1.2, 1.1.3 and 1.1.5](#) reflect an efficient and more conceptual approach to the full cohomology, which cannot be accessed by localization.

In all genera, [Corollary 1.2.1](#) also holds for an arbitrary vector bundle E instead of the trivial bundle $\mathbb{C}^N \otimes \mathcal{O}_{\mathbb{P}^1}$, with the only modification that all instances of $N\chi(L)$ are replaced by $\chi(E \otimes L)$.

1.3. Higher rank. A natural direction is to apply the techniques of this paper to study the cohomology of the Schur functors of tautological bundles over $\text{Quot}_{\mathbb{P}^1}(\mathbb{C}^N, n, r)$, when $r > 0$. In this setting, even the numerical K -theoretic invariants of these Schur functors are largely unexplored.

Recalling the definition of the tautological complex $L^{[n]}$ from [\(1.1.1\)](#), we propose the following conjectures.

Conjecture 1.3.1. *Let $n = (N - r)a + b$ with $0 \leq b < N - r$. Then for all line bundles $L \rightarrow \mathbb{P}^1$, we have*

$$\chi(\text{Quot}_{\mathbb{P}^1}(\mathbb{C}^N, n, r), \wedge^k L^{[n]}) = \binom{N\chi(L)}{k}$$

for all $k \leq n + r(a + 1)$.

For symmetric powers, we state the following

Conjecture 1.3.2. *Let $n = (N - r)a + b$ with $0 \leq b < N - r$. Then for all line bundles $L \rightarrow \mathbb{P}^1$, we have*

$$\chi(\text{Quot}_{\mathbb{P}^1}(\mathbb{C}^N, n, r), \text{Sym}^k L^{[n]}) = \binom{N\chi(L) + k - 1}{k}$$

for all $k \leq n + r(a + 1)$.

Finally, for the dualized exterior powers, we have

Conjecture 1.3.3. *Let $r > 0$ and write $n = ar + b$ with $0 \leq b < r$. Let $1 \leq t \leq N - r - 1$ and p_1, \dots, p_t be nonnegative integers with $0 < p_1 + \dots + p_t \leq n + (N - r)(a + 1)$. Then for all line bundles $L_1, \dots, L_t \rightarrow \mathbb{P}^1$, we have*

$$\chi(\text{Quot}_{\mathbb{P}^1}(\mathbb{C}^N, n, r), (\wedge^{p_1} L_1^{[n]})^\vee \otimes \dots \otimes (\wedge^{p_t} L_t^{[n]})^\vee) = 0.$$

When $r = 0$, Conjectures 1.3.1 and 1.3.2 recover Theorems 1.1.2(1) and 1.1.3(1), while the case $r = N - 1$ can be verified by hand since the Quot scheme is a projective space. For all three conjectures, we checked the answer by computer in several other cases. The bounds on k appear to be sharp.

It is natural to expect that the three conjectures can be lifted in obvious fashion to cohomology. This suggests that Theorems 1.1.2(1), 1.1.3(1) and 1.1.5 continue to hold for higher-rank quotients subject to the appropriate bounds on k and the condition that the degree of the L 's be nonnegative. While Strømme's construction is valid for quotients of arbitrary rank, the Borel–Weil–Bott arguments become more involved and will be left for future study.

The answers predicted by the conjectures stabilize as n becomes large with respect to N, k, r . Equivalently, for each fixed k , the generating series

$$\sum_{n=0}^{\infty} q^n \chi(\text{Quot}_{\mathbb{P}^1}(\mathbb{C}^N, n, r), \wedge^k L^{[n]}), \quad \sum_{n=0}^{\infty} q^n \chi(\text{Quot}_{\mathbb{P}^1}(\mathbb{C}^N, n, r), \text{Sym}^k L^{[n]})$$

are given by rational functions with a (simple) pole only at $q = 1$. It is natural to wonder whether this statement is correct for all partitions and for the associated Schur functors of $L^{[n]}$.

1.4. Subsequent developments. After our preprint appeared on arXiv, the conjectural identity (1.1.6) was extended to cover more general Ext groups involving tensor products of wedge powers of tautological bundles; see [Kr2, Conjecture 1.1]. The extended conjecture is consistent with the Euler characteristic calculations in [OS]. Theorem 1.1.2 and Theorem 1.1.5 establish part of the conjecture. We also refer the reader to [Kr2, Theorems 1.2, 1.3, 1.5] for related results. Both equation (1.1.6) and the conjecture formulated in [Kr2] were very recently confirmed in [MN] using different methods.

1.5. Plan of the paper. We review Strømme’s embedding, construct the resolutions of the tautological bundles, and establish Theorems 1.1.2, 1.1.3 and 1.1.5 in Section 2. This relies on the Borel–Weil–Bott analysis of the resolutions which is carried out in Section 3. Corollaries 1.2.1 and 1.1.7 are proved in Section 4.

2. Grassmannian embedding of the Quot scheme and resolutions

2.1. Strømme’s embedding. We begin by describing Strømme’s construction which exhibits the Quot scheme over \mathbb{P}^1 as the zero locus of a regular section of a vector bundle over the product of two Grassmannians [Str].

For each integer $m \geq n$, the embedding takes the form

$$\iota_m : \text{Quot}_{\mathbb{P}^1}(\mathbb{C}^N, n) \hookrightarrow \mathbf{G}(V_{m-1}, n) \times \mathbf{G}(V_m, n),$$

where

$$\mathbf{G}_1 = \mathbf{G}(V_{m-1}, n), \quad \mathbf{G}_2 = \mathbf{G}(V_m, n)$$

are the Grassmannians of n -dimensional *quotients* of two vector spaces of dimensions Nm and $N(m + 1)$ respectively. We identify

$$V_{m-1} = H^0(\mathbb{P}^1, \mathcal{O}_{\mathbb{P}^1}(m - 1)^{\oplus N}), \quad V_m = H^0(\mathbb{P}^1, \mathcal{O}_{\mathbb{P}^1}(m)^{\oplus N}).$$

Explicitly, ι_m is the product of two Grothendieck embeddings. When $m \geq n$, each short exact sequence in the Quot scheme

$$0 \rightarrow S \rightarrow \mathbb{C}^N \otimes \mathcal{O}_{\mathbb{P}^1} \rightarrow Q \rightarrow 0,$$

yields two exact sequences of vector spaces

$$0 \rightarrow H^0(S(m - 1)) \rightarrow H^0(\mathcal{O}_{\mathbb{P}^1}(m - 1)^{\oplus N}) \rightarrow H^0(Q(m - 1)) \rightarrow 0,$$

$$0 \rightarrow H^0(S(m)) \rightarrow H^0(\mathcal{O}_{\mathbb{P}^1}(m)^{\oplus N}) \rightarrow H^0(Q(m)) \rightarrow 0.$$

Then ι_m is given by the assignment

$$[0 \rightarrow S \rightarrow \mathbb{C}^N \otimes \mathcal{O}_{\mathbb{P}^1} \rightarrow Q \rightarrow 0] \mapsto [V_{m-1} \rightarrow H^0(Q(m - 1))] \times [V_m \rightarrow H^0(Q(m))].$$

We write $W = H^0(\mathbb{P}^1, \mathcal{O}_{\mathbb{P}^1}(1))$. There is a natural morphism

$$V_{m-1} \rightarrow V_m \otimes W,$$

obtained from the natural section cutting out the diagonal $\Delta \hookrightarrow \mathbb{P}^1 \times \mathbb{P}^1$:

$$\mathcal{O}_{\mathbb{P}^1 \times \mathbb{P}^1} \rightarrow \mathcal{O}_{\mathbb{P}^1 \times \mathbb{P}^1}(\Delta) = \mathcal{O}_{\mathbb{P}^1}(1) \boxtimes \mathcal{O}_{\mathbb{P}^1}(1),$$

tensoring by $\mathcal{O}_{\mathbb{P}^1}(m - 1)$ on the first factor, and taking cohomology.

To describe the equations cutting out $\text{Quot}_{\mathbb{P}^1}(\mathbb{C}^N, n)$ in $\mathbf{G}_1 \times \mathbf{G}_2$, let

$$\begin{aligned} 0 \rightarrow \mathcal{A}_1 \rightarrow V_{m-1} \otimes \mathcal{O}_{\mathbf{G}_1} \rightarrow \mathcal{B}_1 \rightarrow 0, \\ 0 \rightarrow \mathcal{A}_2 \rightarrow V_m \otimes \mathcal{O}_{\mathbf{G}_2} \rightarrow \mathcal{B}_2 \rightarrow 0, \end{aligned}$$

be the tautological sequences over the two Grassmannians \mathbf{G}_1 and \mathbf{G}_2 . Let pr_1 and pr_2 be the two projections on $\mathbf{G}_1 \times \mathbf{G}_2$. The sheaf

$$\mathcal{E} = \text{pr}_1^* \mathcal{A}_1^\vee \otimes W \otimes \text{pr}_2^* \mathcal{B}_2 \rightarrow \mathbf{G}_1 \times \mathbf{G}_2 \tag{2.1.1}$$

admits a natural section σ induced by the composition

$$\sigma : \text{pr}_1^* \mathcal{A}_1 \rightarrow V_{m-1} \otimes \mathcal{O}_{\mathbf{G}_1 \times \mathbf{G}_2} \rightarrow V_m \otimes W \otimes \mathcal{O}_{\mathbf{G}_1 \times \mathbf{G}_2} \rightarrow \text{pr}_2^* \mathcal{B}_2 \otimes W.$$

Strømme shows that the section σ is regular and vanishes exactly along $\text{Quot}_{\mathbb{P}^1}(\mathbb{C}^N, n)$; see [Str, §4].

Remark 2.1.2. For an arbitrary vector bundle $E \rightarrow \mathbb{P}^1$ and m sufficiently large, we similarly have an embedding

$$\iota : \text{Quot}_{\mathbb{P}^1}(E, n) \rightarrow \mathbf{G}_1 \times \mathbf{G}_2, \quad \mathbf{G}_1 = \mathbf{G}(V_{m-1}, n), \quad \mathbf{G}_2 = \mathbf{G}(V_m, n),$$

where

$$V_{m-1} = H^0(E(m-1)), \quad V_m = H^0(E(m)).$$

To obtain a precise lower bound for m , we need to ensure the vanishing

$$H^1(S(m-1)) = H^1(S(m)) = 0,$$

for every exact sequence $0 \rightarrow S \rightarrow E \rightarrow Q \rightarrow 0$. Let

$$E = \bigoplus_{i=1}^N \mathcal{O}_{\mathbb{P}^1}(a_i),$$

and set a to be the largest of the summand degrees a_i , $1 \leq i \leq N$. If

$$S = \bigoplus_{i=1}^N \mathcal{O}_{\mathbb{P}^1}(s_i),$$

then we have $s_i \leq a$, $1 \leq i \leq N$, since there is an injection

$$\mathcal{O}_{\mathbb{P}^1}(s_i) \rightarrow S \rightarrow E \rightarrow \bigoplus_{i=1}^N \mathcal{O}_{\mathbb{P}^1}(a).$$

As $\sum_{i=1}^N s_i = \text{deg } E - n$, we also have

$$s_i \geq \text{deg } E - n - (N-1)a, \quad 1 \leq i \leq N.$$

Thus for

$$m \geq n + (N-1)a - \text{deg } E,$$

we obtain $H^1(S(m-1)) = H^1(S(m)) = 0$, as wished.

With no additional conditions on m , Strømme's arguments show that $\text{Quot}_{\mathbb{P}^1}(E, n)$ is also cut out by a regular section σ of the tautological bundle (2.1.1) over $\mathbf{G}_1 \times \mathbf{G}_2$. Consequently, the arguments below presented for the case of trivial E also carry over without change to arbitrary E .

2.2. Resolutions. As a result of the above discussion, the section σ induces a Koszul resolution

$$\cdots \rightarrow \wedge^2 \mathcal{E}^\vee \rightarrow \wedge^1 \mathcal{E}^\vee \rightarrow \mathcal{O}_{\mathbf{G}_1 \times \mathbf{G}_2} \rightarrow \mathcal{O}_{\text{Quot}} \rightarrow 0. \tag{2.2.1}$$

Note that if $\deg L = m$, by the definition of the embedding ι_m we have

$$L^{[n]} = \iota_m^* \text{pr}_2^* \mathcal{B}_2.$$

Hence, tensoring (2.2.1) with $\text{pr}_2^* \wedge^k \mathcal{B}_2$, we obtain the resolution

$$\cdots \rightarrow \wedge^2 \mathcal{E}^\vee \otimes \text{pr}_2^* \wedge^k \mathcal{B}_2 \rightarrow \wedge^1 \mathcal{E}^\vee \otimes \text{pr}_2^* \wedge^k \mathcal{B}_2 \rightarrow \text{pr}_2^* \wedge^k \mathcal{B}_2 \rightarrow (\iota_m)_* (\wedge^k L^{[n]}) \rightarrow 0$$

over $\mathbf{G}_1 \times \mathbf{G}_2$. We set

$$\mathcal{V}_\ell = \wedge^\ell \mathcal{E}^\vee \otimes \text{pr}_2^* \wedge^k \mathcal{B}_2, \quad \ell \geq 0.$$

This yields the resolution

$$\cdots \rightarrow \mathcal{V}_2 \rightarrow \mathcal{V}_1 \rightarrow \mathcal{V}_0 \rightarrow (\iota_m)_* (\wedge^k L^{[n]}) \rightarrow 0, \tag{2.2.2}$$

corresponding to Theorem 1.1.2(1).

Theorem 1.1.2(2) is conceptually analogous, but the notation becomes slightly more involved. For this reason, it may be helpful to present the simpler case (1) first. To treat case (2), we consider the bundle

$$\mathcal{F} = (\wedge^{p_1} L^{[n]})^\vee \otimes \cdots \otimes (\wedge^{p_r} L^{[n]})^\vee \otimes (\wedge^k L^{[n]}). \tag{2.2.3a}$$

By similar reasoning, we are led to the resolution

$$\cdots \rightarrow \mathcal{X}_2 \rightarrow \mathcal{X}_1 \rightarrow \mathcal{X}_0 \rightarrow (\iota_m)_* \mathcal{F} \rightarrow 0, \tag{2.2.3b}$$

where

$$\mathcal{X}_\ell = \wedge^\ell \mathcal{E}^\vee \otimes (\text{pr}_2^* \wedge^{p_1} \mathcal{B}_2^\vee \otimes \text{pr}_2^* \wedge^{p_2} \mathcal{B}_2^\vee \otimes \cdots \otimes \text{pr}_2^* \wedge^{p_r} \mathcal{B}_2^\vee \otimes \text{pr}_2^* \wedge^k \mathcal{B}_2).$$

Proposition 2.2.4. *In the resolution (2.2.2),*

$$\cdots \rightarrow \mathcal{V}_2 \rightarrow \mathcal{V}_1 \rightarrow \mathcal{V}_0 \rightarrow (\iota_m)_* (\wedge^k L^{[n]}) \rightarrow 0,$$

the sheaves \mathcal{V}_ℓ have no cohomology for $\ell \geq 1$, while the sheaf \mathcal{V}_0 has no higher cohomology.

More generally, in the resolution (2.2.3b), the sheaves \mathcal{X}_ℓ have no cohomology for $\ell \geq 1$, while the sheaf \mathcal{X}_0 has no higher cohomology.

For the symmetric products, we similarly define

$$\mathcal{W}_\ell = \wedge^\ell \mathcal{E}^\vee \otimes \text{pr}_2^* \text{Sym}^k \mathcal{B}_2,$$

and have an analogous resolution

$$\cdots \rightarrow \mathcal{W}_2 \rightarrow \mathcal{W}_1 \rightarrow \mathcal{W}_0 \rightarrow (t_m)_* (\text{Sym}^k L^{[n]}) \rightarrow 0. \tag{2.2.5}$$

This corresponds to [Theorem 1.1.3\(1\)](#). For the more general [Theorem 1.1.3\(2\)](#), we let

$$\mathcal{G} = (\wedge^{p_1} L^{[n]})^\vee \otimes \cdots \otimes (\wedge^{p_t} L^{[n]})^\vee \otimes \text{Sym}^k L^{[n]}. \tag{2.2.6a}$$

Setting

$$\mathcal{Y}_\ell = \wedge^\ell \mathcal{E}^\vee \otimes (\text{pr}_2^* \wedge^{p_1} \mathcal{B}_2^\vee \otimes \text{pr}_2^* \wedge^{p_2} \mathcal{B}_2^\vee \otimes \cdots \otimes \text{pr}_2^* \wedge^{p_t} \mathcal{B}_2^\vee \otimes \text{pr}_2^* \text{Sym}^k \mathcal{B}_2),$$

we obtain the resolution

$$\cdots \rightarrow \mathcal{Y}_2 \rightarrow \mathcal{Y}_1 \rightarrow \mathcal{Y}_0 \rightarrow (t_m)_* \mathcal{G} \rightarrow 0. \tag{2.2.6b}$$

Proposition 2.2.7. *In the resolution [\(2.2.5\)](#)*

$$\cdots \rightarrow \mathcal{W}_2 \rightarrow \mathcal{W}_1 \rightarrow \mathcal{W}_0 \rightarrow (t_m)_* (\text{Sym}^k L^{[n]}) \rightarrow 0,$$

the sheaves \mathcal{W}_ℓ have no cohomology if $\ell \geq 1$ and $\deg L \geq n \geq k$, while \mathcal{W}_0 has no higher cohomology.

More generally, under the same assumptions, in the resolution [\(2.2.6b\)](#), the sheaves \mathcal{Y}_ℓ have no cohomology for $\ell \geq 1$, while the sheaf \mathcal{Y}_0 has no higher cohomology.

A further analysis is needed for [Theorem 1.1.5](#). The case $L = M$ is already covered either by [Theorem 1.1.2\(2\)](#) or [Theorem 1.1.3\(2\)](#) for $k = 0$. Thus we may assume $\deg M = m \geq n$ and $\deg L = m - 1$. In this case, we have

$$L^{[n]} = t_m^* \text{pr}_1^* \mathcal{B}_1, \quad M^{[n]} = t_m^* \text{pr}_2^* \mathcal{B}_2.$$

Thus for the bundle

$$\mathcal{H} = (\wedge^{p_1} L^{[n]})^\vee \otimes (\wedge^{p_2} M^{[n]})^\vee \otimes \cdots \otimes (\wedge^{p_t} M^{[n]})^\vee$$

which appears in [Theorem 1.1.5](#), we obtain a resolution

$$\cdots \rightarrow \mathcal{U}_2 \rightarrow \mathcal{U}_1 \rightarrow \mathcal{U}_0 \rightarrow (t_m)_* \mathcal{H} \rightarrow 0, \tag{2.2.8}$$

where

$$\mathcal{U}_\ell = \wedge^\ell \mathcal{E}^\vee \otimes (\text{pr}_1^* \wedge^{p_1} \mathcal{B}_1^\vee \otimes \text{pr}_2^* \wedge^{p_2} \mathcal{B}_2^\vee \otimes \cdots \otimes \text{pr}_2^* \wedge^{p_t} \mathcal{B}_2^\vee).$$

Proposition 2.2.9. *For p_1, \dots, p_t not all zero, the cohomology of \mathcal{U}_ℓ vanishes for all $\ell \geq 0$.*

2.2.1. The main theorems. Before turning our attention to the proofs of the above propositions, we note that our main [Theorems 1.1.2, 1.1.3](#) and [1.1.5](#) follow immediately from them.

For [Theorem 1.1.2](#), we use [Proposition 2.2.4](#). To establish case (1) of the theorem, we make use of the resolution [\(2.2.2\)](#). The associated spectral sequence shows that the higher cohomology of $\wedge^k L^{[n]}$ vanishes, while in degree zero we have

$$H^0(\text{Quot}_{\mathbb{P}^1}(\mathbb{C}^N, n), \wedge^k L^{[n]}) = H^0(\mathbf{G}_1 \times \mathbf{G}_2, \mathcal{V}_0).$$

Recalling that $\mathcal{V}_0 = \text{pr}_2^* \wedge^k \mathcal{B}_2$, we compute

$$H^0(\mathbf{G}_1 \times \mathbf{G}_2, \mathcal{V}_0) = H^0(\mathbf{G}_2, \wedge^k \mathcal{B}_2) = \wedge^k V_m = \wedge^k H^0(\mathbb{C}^N \otimes \mathcal{O}_{\mathbb{P}^1}(m)) = \wedge^k H^0(L^{\oplus N}),$$

as needed.

For part (2) of [Theorem 1.1.2](#), we note that

$$\text{Ext}^i(\wedge^{p_1} L^{[n]} \otimes \cdots \otimes \wedge^{p_t} L^{[n]}, \wedge^k L^{[n]}) = H^i(\mathcal{F}),$$

where the bundle \mathcal{F} is defined in [\(2.2.3a\)](#). Using the resolution [\(2.2.3b\)](#) and [Proposition 2.2.4](#), we see that the only contribution to the cohomology of \mathcal{F} comes from the term

$$H^0(\mathbf{G}_1 \times \mathbf{G}_2, \mathcal{X}_0) = H^0(\mathcal{B}_2^\vee \otimes \cdots \otimes \wedge^{p_t} \mathcal{B}_2^\vee \otimes \wedge^k \mathcal{B}_2) = \wedge^{k-|p|} H^0(\mathbf{G}_2, \mathcal{B}_2) = \wedge^{k-|p|} H^0(L^{\oplus N})$$

for $|p| = p_1 + \cdots + p_t \leq k \leq n$. The second equality requires further explanation. Since we need additional notation, the argument will be presented later; see equation [\(2.3.11\)](#) in the proof of [Proposition 2.2.4](#).

Turning to [Theorem 1.1.3](#), for part (1), we make use of the resolution [\(2.2.5\)](#) and [Proposition 2.2.7](#). This time, the initial term $\mathcal{W}_0 = \text{pr}_2^* \text{Sym}^k \mathcal{B}_2$ has sections

$$H^0(\mathbf{G}_1 \times \mathbf{G}_2, \mathcal{W}_0) = H^0(\mathbf{G}_2, \text{Sym}^k \mathcal{B}_2) = \text{Sym}^k V_m = \text{Sym}^k H^0(\mathbb{C}^N \otimes \mathcal{O}_{\mathbb{P}^1}(m)) = \text{Sym}^k H^0(L^{\oplus N}).$$

For part (2), we note

$$\text{Ext}^i(\wedge^{p_1} L^{[n]} \otimes \cdots \otimes \wedge^{p_t} L^{[n]}, \text{Sym}^k L^{[n]}) = H^i(\mathcal{G}),$$

where the sheaf \mathcal{G} was defined in [\(2.2.6a\)](#). Using the resolution [\(2.2.6b\)](#) and [Proposition 2.2.7](#), the only nontrivial contribution to the cohomology of \mathcal{G} comes from the sheaf \mathcal{Y}_0 and it equals

$$H^0(\mathbf{G}_2, \wedge^{p_1} \mathcal{B}_2^\vee \otimes \cdots \otimes \wedge^{p_t} \mathcal{B}_2^\vee \otimes \text{Sym}^k \mathcal{B}_2) = \text{Sym}^{k-|p|} H^0(\mathbf{G}_2, \mathcal{B}_2) = \text{Sym}^{k-|p|} H^0(L^{\oplus N}).$$

This requires that all $p_j \in \{0, 1\}$ and $|p| \leq k$. The cohomology vanishes altogether if this condition fails. The first equality will be explained after we develop more notation; see equation [\(2.3.13\)](#) in the proof of [Proposition 2.2.7](#) below.

Finally, for [Theorem 1.1.5](#), the argument uses [Proposition 2.2.9](#). This time around, the cohomology vanishes for all terms of the resolution [\(2.2.8\)](#) and in all degrees. □

2.3. Analysis of the resolutions. We now turn to [Propositions 2.2.4, 2.2.7, 2.2.9](#) and deduce them from the Grassmannian vanishing results of [Section 3](#). We begin by making the terms of the resolutions more explicit.

2.3.1. Partitions and Cauchy's formula. We use standard terminology on partitions $\lambda = (\lambda_1, \dots, \lambda_r)$, where $\lambda_1 \geq \lambda_2 \geq \cdots \geq \lambda_r \geq 0$. We set

$$|\lambda| = \sum_{i=1}^r \lambda_i,$$

and we let λ^\dagger be the transpose partition obtained by exchanging the rows and columns of the Young diagram of λ .

Throughout, we will always assume that our base scheme is defined over a field of characteristic 0.

For each partition λ , we let \mathbf{S}_λ denote the associated Schur functor (for example, these are defined in [W, §2.1] where they are called L_{λ^\dagger}). For a partition λ and any vector bundle $V \rightarrow Y$ over a base Y , there is an associated vector bundle $\mathbf{S}_\lambda(V) \rightarrow Y$. The cases $\lambda = (1^k)$ and $\lambda = (k)$ correspond to the k -th exterior and k -th symmetric powers, respectively.

The vector bundles $\mathbf{S}_\lambda(V) \rightarrow Y$ are also defined when λ is not a partition but rather an arbitrary dominant weight $\lambda_1 \geq \dots \geq \lambda_r$, where $r = \text{rank}(V)$, and we now allow the entries to be negative. We have

$$\mathbf{S}_{-\lambda}(V) = \mathbf{S}_\lambda(V^\vee),$$

where $-\lambda$ denotes the sequence $-\lambda_r \geq \dots \geq -\lambda_1$. In addition,

$$\mathbf{S}_\lambda(V) \otimes \det V = \mathbf{S}_{\lambda+(1^r)}(V).$$

If $V, W \rightarrow Y$ are two vector bundles, Cauchy’s identity

$$\bigwedge^\ell(V \otimes W) = \bigoplus_{|\lambda|=\ell} \mathbf{S}_{\lambda^\dagger}(V) \otimes \mathbf{S}_\lambda(W)$$

holds, where the sum is over all partitions λ of size ℓ . We only need to consider those partitions λ with at most $\text{rank}(W)$ rows and at most $\text{rank}(V)$ columns since the term is 0 otherwise. Applying this formula to the bundle \mathcal{E} whose section cuts out the Quot scheme, we obtain

$$\bigwedge^\ell \mathcal{E}^\vee = \bigoplus_{|\lambda|=\ell} \mathbf{S}_{\lambda^\dagger}(\mathcal{A}_1) \boxtimes \mathbf{S}_\lambda(\mathcal{B}_2^\vee \oplus \mathcal{B}_2^\vee). \tag{2.3.1a}$$

Here, λ is a partition with at most $2n$ rows and the number of columns at most equal to

$$\text{rank}(\mathcal{A}_1) = \dim V_{m-1} - n = \dim V_m - N - n \leq \dim V_m - n - 1. \tag{2.3.1b}$$

In the discussion below, the abbreviation $d = \dim V_m$ will often be used.

Using (2.3.1a), we immediately obtain the expressions

$$\mathcal{V}_\ell = \bigoplus_{|\lambda|=\ell} \mathbf{S}_{\lambda^\dagger}(\mathcal{A}_1) \boxtimes (\mathbf{S}_\lambda(\mathcal{B}_2^\vee \oplus \mathcal{B}_2^\vee) \otimes \bigwedge^k \mathcal{B}_2), \tag{2.3.2a}$$

and

$$\mathcal{W}_\ell = \bigoplus_{|\lambda|=\ell} \mathbf{S}_{\lambda^\dagger}(\mathcal{A}_1) \boxtimes (\mathbf{S}_\lambda(\mathcal{B}_2^\vee \oplus \mathcal{B}_2^\vee) \otimes \text{Sym}^k \mathcal{B}_2) \tag{2.3.2b}$$

corresponding to Theorem 1.1.2(1) and Theorem 1.1.3(1). For the second halves of the two theorems, the expressions are slightly more complicated due to additional wedge powers:

$$\mathcal{X}_\ell = \bigoplus_{|\lambda|=\ell} \mathbf{S}_{\lambda^\dagger}(\mathcal{A}_1) \boxtimes (\mathbf{S}_\lambda(\mathcal{B}_2^\vee \oplus \mathcal{B}_2^\vee) \otimes \bigwedge^{p_1} \mathcal{B}_2^\vee \otimes \dots \otimes \bigwedge^{p_i} \mathcal{B}_2^\vee \otimes \bigwedge^k \mathcal{B}_2). \tag{2.3.2c}$$

and

$$\mathcal{Y}_\ell = \bigoplus_{|\lambda|=\ell} \mathbf{S}_{\lambda^\dagger}(\mathcal{A}_1) \boxtimes (\mathbf{S}_\lambda(\mathcal{B}_2^\vee \oplus \mathcal{B}_2^\vee) \otimes \wedge^{p_1} \mathcal{B}_2^\vee \otimes \cdots \otimes \wedge^{p_t} \mathcal{B}_2^\vee \otimes \text{Sym}^k \mathcal{B}_2). \tag{2.3.2d}$$

Finally, we have

$$\mathcal{U}_\ell = \bigoplus_{|\lambda|=\ell} (\mathbf{S}_{\lambda^\dagger}(\mathcal{A}_1) \otimes \wedge^{p_1} \mathcal{B}_1^\vee) \boxtimes (\mathbf{S}_\lambda(\mathcal{B}_2^\vee \oplus \mathcal{B}_2^\vee) \otimes \wedge^{p_2} \mathcal{B}_2^\vee \otimes \cdots \otimes \wedge^{p_t} \mathcal{B}_2^\vee). \tag{2.3.2e}$$

2.3.2. The Borel–Weil–Bott theorem. To compute the cohomology of the above bundles, we use the Borel–Weil–Bott theorem [B]. For integers $0 < n < d$, let $\mathbf{G} = \mathbf{G}(d, n)$ denote the Grassmannian of n -dimensional *quotients* of a d -dimensional vector space, and let $\mathcal{A}, \mathcal{B} \rightarrow \mathbf{G}$ denote the tautological subbundle and quotient. For a partition

$$\mu = (\mu_1, \dots, \mu_{d-n})$$

with $d - n$ rows, we form the string

$$\rho + (0, \mu) = (d - 1, d - 2, \dots, 1, 0) + \underbrace{(0, \dots, 0)}_n, \mu_1, \dots, \mu_{d-n}. \tag{2.3.3}$$

Theorem 2.3.4 (Borel–Weil–Bott). *The bundle $\mathbf{S}_\mu(\mathcal{A})$ has at most one nonzero cohomology group. Furthermore, if the string $\rho + (0, \mu)$ contains repetitions, then all cohomology groups of $\mathbf{S}_\mu(\mathcal{A})$ vanish.*

This formulation can be found in [W, Corollary 4.1.9]. We are using a few translations. First, the Weyl functors K_γ defined there are isomorphic to the Schur functors used here; see [W, Proposition 2.1.18(c)]. Moreover, the dual of the tautological subbundle \mathcal{A} is the quotient bundle on the dual Grassmannian, and on the latter space, $\mathbf{S}_\mu(\mathcal{A})$ corresponds to $\mathcal{V}(0, \mu)$ in the notation of [W].

We note from **Theorem 2.3.4** that $\mathbf{S}_\mu(\mathcal{A})$ has no cohomology provided there exists j such that

$$j \leq \mu_j \leq n + j - 1. \tag{2.3.5a}$$

Let us record the “dual” rephrasing of condition (2.3.5a) which is also useful here. For a partition ν with n rows, the bundle $\mathbf{S}_\nu(\mathcal{B}^\vee)$ over the Grassmannian $\mathbf{G}(d, n)$ has no cohomology provided that there exists j such that

$$j \leq \nu_j \leq d - n + j - 1. \tag{2.3.5b}$$

In **Proposition 2.2.9**, we also consider the bundle

$$\mathbf{S}_\mu(\mathcal{A}) \otimes \wedge^p \mathcal{B}^\vee = \mathbf{S}_{-\mu}(\mathcal{A}^\vee) \otimes \mathbf{S}_{(1^p)} \mathcal{B}^\vee$$

for $0 \leq p \leq n$. Again by the Borel–Weil–Bott theorem [W, Corollary 4.1.9], all cohomology vanishes provided that the string

$$(d - 1, \dots, 0) + (-\mu_{d-n}, \dots, -\mu_1, \underbrace{1, \dots, 1}_p, \underbrace{0, \dots, 0}_{n-p}) \tag{2.3.6a}$$




Figure 1. A partition of 2-index $i = 3$.

has repetitions. That happens when there exists j such that

$$j - 1 \leq \mu_j \leq n + j - 1 \text{ and } \mu_j \neq j + p - 1. \tag{2.3.6b}$$

Of course, the case $p = 0$ recovers (2.3.5a). In fact, for $p = 0$, the string (2.3.3) has repetitions if and only if the same holds for (2.3.6a).

2.3.3. Indices of partitions. The following definition is not standard but is crucial for our arguments.

Definition 2.3.7. Let n be a nonnegative integer. Let $\lambda \neq 0$ be a partition satisfying the condition that (*) for all j , the number of boxes in the j -th column of λ is either $< j$ or $\geq n + j$.

Let i denote the largest index j such that the j -th column has $\geq n + j$ boxes. We refer to i as the n -index of λ . If λ does not satisfy (*), we leave the n -index undefined.

It may help to visualize partitions λ of n -index i . There are i “long” columns with at least $n + i$ boxes, while the remaining columns are “short” containing at most i boxes. In Figure 1, the long columns are shown in gray, while the short columns are white. Thus, for a partition λ of n -index i , we have

$$\lambda_{i+1} = \dots = \lambda_{i+n} = i. \tag{2.3.8}$$

The following variation is needed for Proposition 2.2.9 and is connected to condition (2.3.6b) above.

Definition 2.3.9. Let $0 \leq p \leq n$ be integers. Let $\lambda \neq 0$, $\lambda \neq (1^p)$ be a partition satisfying the condition that

(**) for all j , the number of boxes in the j -th column of λ is either $< j - 1$ or $\geq n + j$ or equal to $j + p - 1$.

Let i denote the largest index j such that the j -th column has $\geq n + j$ boxes. We refer to i as the (p, n) -index of λ , when defined. The case $p = 0$ corresponds to the n -index defined above.

The partition $\lambda = (1^p)$ is not considered here. The reason is that λ satisfies (**), yet no column has at least $n + 1$ boxes, so the index is undefined.

For a partition λ of (p, n) -index i , two shapes are possible:

- (a) The partition λ has i “long” columns with $\geq n + i$ boxes, and the remaining columns are “short”, having $\leq i - 1$ boxes.
- (b) The partition λ has i “long” columns with $\geq n + i$ boxes, the $(i + 1)$ st column has $p + i$ boxes, and the remaining columns are “short”, having $\leq i$ boxes.




Figure 2. Partitions of $(1, 3)$ -index $i = 2$.

The partitions in [Figure 2](#) satisfy (a) and (b) respectively. For case (b), the long and short columns are shown in dark gray and white, while the middle $(i + 1)$ st column is lighter gray. In both cases

$$i \leq \lambda_{i+1} \leq i + 1, \quad \dots, \quad i \leq \lambda_{i+n} \leq i + 1. \tag{2.3.10}$$

Proof of [Proposition 2.2.4](#). For simplicity, we consider the case of the resolution \mathcal{V}_\bullet first. Recall from [\(2.3.2a\)](#) that

$$\mathcal{V}_\ell = \bigoplus_{|\lambda|=\ell} \mathbf{S}_{\lambda^\dagger}(\mathcal{A}_1) \boxtimes (\mathbf{S}_\lambda(\mathcal{B}_2^\vee \oplus \mathcal{B}_2^\vee) \otimes \wedge^k \mathcal{B}_2).$$

For $\ell = 0$, we must have $\lambda = 0$, and $\mathcal{V}_0 = \text{pr}^* \wedge^k \mathcal{B}_2$ has no higher cohomology.

When $\ell \geq 1$, we have $\lambda \neq 0$. For a partition $\lambda \neq 0$ appearing in the above sum, we distinguish two mutually exclusive situations:

- (\dagger) there exists j such that $j \leq \lambda_j^\dagger \leq n + j - 1$, or
- ($*$) for all j , the number of boxes in the j -th column of λ is either $< j$ or $\geq n + j$.

Of course, condition ($*$) already appeared in [Definition 2.3.7](#). In case (\dagger), we noted in [\(2.3.5a\)](#) that $\mathbf{S}_{\lambda^\dagger}(\mathcal{A}_1)$ has no cohomology. In case ($*$), [Proposition 3.1.2](#) in [Section 3](#) below shows that

$$\mathbf{S}_\lambda(\mathcal{B}_2^\vee \oplus \mathcal{B}_2^\vee) \otimes \wedge^k \mathcal{B}_2$$

has no cohomology either. Consequently, \mathcal{V}_ℓ has no cohomology when $\ell \geq 1$, establishing the first half of [Proposition 2.2.4](#).

For the second half, recall from [\(2.3.2c\)](#) that

$$\mathcal{X}_\ell = \bigoplus_{|\lambda|=\ell} \mathbf{S}_{\lambda^\dagger}(\mathcal{A}_1) \boxtimes (\mathbf{S}_\lambda(\mathcal{B}_2^\vee \oplus \mathcal{B}_2^\vee) \otimes \wedge^{p_1} \mathcal{B}_2^\vee \otimes \dots \otimes \wedge^{p_t} \mathcal{B}_2^\vee \otimes \wedge^k \mathcal{B}_2).$$

When $\ell \neq 0$, then either $\lambda \neq 0$ satisfies condition (\dagger), which guarantees cohomology vanishing for the Schur bundle on the first factor, or else $\lambda \neq 0$ satisfies ($*$). The latter situation yields vanishing on the second factor by [Proposition 3.1.4\(1\)](#) below. The hypothesis of the proposition is verified since λ was seen in [\(2.3.1b\)](#) to have at most

$$\dim V_m - N - n \leq \dim V_m - (t + 1) - n$$

columns and $t \leq N - 1$.

In the proof of [Theorem 1.1.2](#) in [Section 2.2.1](#), we also claimed that the cohomology of \mathcal{X}_0 is given by

$$H^0(\mathbf{G}_2, \wedge^{p_1} \mathcal{B}_2^\vee \otimes \cdots \otimes \wedge^{p_t} \mathcal{B}_2^\vee \otimes \wedge^k \mathcal{B}_2) = H^0(\mathbf{G}_2, \wedge^{k-|p|} \mathcal{B}_2) = \wedge^{k-|p|} H^0(\mathbf{G}_2, \mathcal{B}_2). \tag{2.3.11}$$

Here we assume $|p| = p_1 + \cdots + p_t \leq k \leq n$, while the answer is understood to be 0 if this assumption fails. We can now justify the first equality using Pieri’s rule combined with Borel–Weil–Bott. This is certainly well-known, but it appears easier to write the argument than to find a reference.

Consider an arbitrary Grassmannian $\mathbf{G}(d, n)$ with tautological quotient \mathcal{B} , and assume $t \leq d - n$. The latter is true in our setting since $t \leq N - 1 \leq \dim V_m - n$. We inspect the tensor product

$$\wedge^{p_1} \mathcal{B}^\vee \otimes \cdots \otimes \wedge^{p_t} \mathcal{B}^\vee \otimes \wedge^k \mathcal{B} = \wedge^{p_1} \mathcal{B}^\vee \otimes \cdots \otimes \wedge^{p_t} \mathcal{B}^\vee \otimes \wedge^{n-k} \mathcal{B}^\vee \otimes \det \mathcal{B}. \tag{2.3.12}$$

By Pieri’s rule, tensorization by $\wedge^p \mathcal{B}^\vee$ has the effect of adding p boxes, no two in the same row. Thus, if $\mathbf{S}_\nu(\mathcal{B}^\vee)$ is any summand of $\wedge^{p_1} \mathcal{B}^\vee \otimes \cdots \otimes \wedge^{p_t} \mathcal{B}^\vee \otimes \wedge^{n-k} \mathcal{B}^\vee$, then we create at most $t + 1$ columns. Hence

$$1 \leq \nu_1 \leq t + 1 \leq d - n + 1.$$

On the other hand, tensorization by $\det \mathcal{B}$ subtracts 1 box from all the rows. Consequently, the summands $\mathbf{S}_\mu(\mathcal{B}^\vee)$ that appear in [\(2.3.12\)](#) satisfy

$$\mu_1 = \nu_1 - 1.$$

In general, we have $1 \leq \mu_1 \leq d - n$, so Borel–Weil–Bott shows that $\mathbf{S}_\mu(\mathcal{B}^\vee)$ has no cohomology; see condition [\(2.3.5b\)](#). There is one exception corresponding to $\mu_1 = 0$. In this case, we must have $\nu_1 = 1$, which forces $\nu = (1^{|p|+n-k})$ and then $-\mu = (1^{k-|p|})$. This yields the term $\mathbf{S}_\mu(\mathcal{B}^\vee) = \mathbf{S}_{-\mu}(\mathcal{B}) = \wedge^{k-|p|} \mathcal{B}$, and justifies [\(2.3.11\)](#). □

Proof of [Proposition 2.2.7](#). We consider the simpler case of the resolution \mathcal{W}_\bullet first. By [\(2.3.2b\)](#) we have

$$\mathcal{W}_\ell = \bigoplus_{|\lambda|=\ell} \mathbf{S}_{\lambda^\dagger}(\mathcal{A}_1) \boxtimes (\mathbf{S}_\lambda(\mathcal{B}_2^\vee \oplus \mathcal{B}_2^\vee) \otimes \mathrm{Sym}^k \mathcal{B}_2).$$

The argument is similar to that of [Equation \(2.2.2\)](#). This time, to deal with case (*) we invoke [Proposition 3.1.3](#).

The analysis of the bundles \mathcal{Y}_ℓ for $\ell \geq 1$ in the general case uses [Proposition 3.1.4\(2\)](#) instead.

In [Section 2.2.1](#), we also needed the cohomology of the bundle \mathcal{Y}_0 . To this end, we show that on an arbitrary Grassmannian $\mathbf{G}(d, n)$, for $t \leq d - n$ and positive integers $p_1, \dots, p_t > 0$, the bundle $\wedge^{p_1} \mathcal{B}^\vee \otimes \cdots \otimes \wedge^{p_t} \mathcal{B}^\vee \otimes \mathrm{Sym}^k \mathcal{B}$ has no cohomology unless $p_1 = \cdots = p_t = 1$ and $k \geq t$, in which case there is only one nontrivial cohomology group

$$H^0(\wedge^{p_1} \mathcal{B}^\vee \otimes \cdots \otimes \wedge^{p_t} \mathcal{B}^\vee \otimes \mathrm{Sym}^k \mathcal{B}) = \mathrm{Sym}^{k-t} H^0(\mathcal{B}). \tag{2.3.13}$$

Indeed, let $\mathbf{S}_\mu(\mathcal{B}^\vee)$ be a summand of $\wedge^{p_1} \mathcal{B}^\vee \otimes \cdots \otimes \wedge^{p_t} \mathcal{B}^\vee \otimes \mathrm{Sym}^k \mathcal{B}$. Dualizing, we have that $\mathbf{S}_{-\mu}(\mathcal{B}^\vee)$

is a summand of

$$\wedge^{p_1} \mathcal{B} \otimes \cdots \otimes \wedge^{p_t} \mathcal{B} \otimes \text{Sym}^k \mathcal{B}^\vee = \wedge^{n-p_1} \mathcal{B}^\vee \otimes \cdots \otimes \wedge^{n-p_t} \mathcal{B}^\vee \otimes \text{Sym}^k \mathcal{B}^\vee \otimes (\det \mathcal{B})^t.$$

Let $\mathbf{S}_\nu(\mathcal{B}^\vee)$ be a summand of $\wedge^{n-p_1} \mathcal{B}^\vee \otimes \cdots \otimes \wedge^{n-p_t} \mathcal{B}^\vee \otimes \text{Sym}^k \mathcal{B}^\vee$. Applying the Pieri rules, we start with the partition (k) corresponding to $\text{Sym}^k \mathcal{B}^\vee$, to which we add $n - p_1, \dots, n - p_t$ boxes respectively, such that at each stage we do not add two boxes to the same row. Therefore

$$0 \leq \nu_n \leq t$$

(unless $n = 1$, which can be treated separately). If $\nu_n = t$, then $\nu_2 = \cdots = \nu_{n-1} = t$. In this case we can say a bit more. Note that the last $n - 1$ rows of ν each contain t boxes, and $n - p_1, \dots, n - p_t$ are all $\leq n - 1$. For this to be possible, equality must hold, hence $p_1 = \cdots = p_t = 1$. Moreover, all boxes have to be added to the last $n - 1$ rows, and so none can be added to the first row. Hence ν is the partition

$$\nu_1 = k, \quad \nu_2 = \cdots = \nu_n = t.$$

For this to make sense, we must have $\nu_1 = k \geq t = \nu_2$.

Finally, each $\mathbf{S}_{-\mu}(\mathcal{B}^\vee) = \mathbf{S}_\nu(\mathcal{B}^\vee) \otimes (\det \mathcal{B})^t$ satisfies $\mu_{n+1-i} + \nu_i = t$. Since $0 \leq \nu_n \leq t$, we also have $0 \leq \mu_1 \leq t$. Since $t \leq d - n$, it follows by condition (2.3.5b) that we only have nontrivial cohomology for $\mathbf{S}_\mu(\mathcal{B}^\vee)$ when $\mu_1 = 0$. This case corresponds to $\nu_n = t$. We have seen above this means $\nu_1 = k$ and $\nu_2 = \cdots = \nu_n = t$. This yields $\mu = (0, \dots, 0, t - k)$ and $\mathbf{S}_\mu(\mathcal{B}^\vee) = \text{Sym}^{k-t} \mathcal{B}$, which implies the claim. \square

Proof of Proposition 2.2.9. We need to inspect

$$\mathcal{U}_\ell = \bigoplus_{|\lambda|=\ell} (\mathbf{S}_{\lambda^\dagger}(\mathcal{A}_1) \otimes \wedge^{p_1} \mathcal{B}_1^\vee) \boxtimes (\mathbf{S}_\lambda(\mathcal{B}_2^\vee \oplus \mathcal{B}_2^\vee) \otimes \wedge^{p_2} \mathcal{B}_2^\vee \otimes \cdots \otimes \wedge^{p_t} \mathcal{B}_2^\vee).$$

The case $\ell = 0$ follows from either (2.3.11) or (2.3.13) with $k = 0$.

Let $\ell \neq 0$. When λ^\dagger satisfies (2.3.6b) (for $p = p_1$), the first factor $\mathbf{S}_{\lambda^\dagger}(\mathcal{A}_1) \otimes \wedge^{p_1} \mathcal{B}_1^\vee$ has no cohomology. Otherwise, λ satisfies condition (**) from Definition 2.3.9. In this case, we claim the second factor has no cohomology.

Indeed, when $\lambda = (1^{p_1})$, we have that $\mathbf{S}_\lambda = \wedge^{p_1}$ is an exterior power and

$$\wedge^{p_1} (\mathcal{B}_2^\vee \oplus \mathcal{B}_2^\vee) \cong \bigoplus_{j=0}^{p_1} \wedge^j \mathcal{B}_2^\vee \otimes \wedge^{p_1-j} \mathcal{B}_2^\vee.$$

As in the above analysis of (2.3.11), every summand $\mathbf{S}_\mu(\mathcal{B}_2^\vee)$ which appears in the second factor then satisfies $1 \leq \mu_1 \leq t + 1$ by the Pieri rule. Since $t + 1 \leq N \leq \dim V_m - n = d - n$, we get vanishing by Borel–Weil–Bott and (2.3.5b).

When $\lambda \neq (1^{p_1})$, letting i denote the (p_1, n) -index of λ , the second factor

$$\mathbf{S}_\lambda(\mathcal{B}_2^\vee \oplus \mathcal{B}_2^\vee) \otimes \wedge^{p_2} \mathcal{B}_2^\vee \otimes \cdots \otimes \wedge^{p_t} \mathcal{B}_2^\vee$$

has no cohomology by Proposition 3.1.4(3). \square

3. Cohomology vanishing on the Grassmannian

3.1. Overview. We establish the vanishing results which played a crucial role in the proofs of Propositions 2.2.4, 2.2.7 and 2.2.9.

We continue to write $\mathbf{G} = \mathbf{G}(d, n)$ for the Grassmannian of n -dimensional quotients of a d -dimensional vector space, and $\mathcal{B} \rightarrow \mathbf{G}$ for the tautological rank- n quotient.

Recall from Section 2.3.2 that $\mathbf{S}_\delta(\mathcal{B}^\vee)$ has no cohomology provided that there exists j such that

$$j \leq \delta_j \leq d - n + j - 1. \tag{3.1.1}$$

We will establish three results. The first corresponds to the resolution \mathcal{V}_\bullet , while the second pertains to the resolution \mathcal{W}_\bullet . Together, these already capture the main ideas. The third result covers the resolutions \mathcal{X}_\bullet , \mathcal{Y}_\bullet and \mathcal{U}_\bullet . Although the notation in this case is more involved, the argument does not require new ideas. We present these results separately for clarity.

Proposition 3.1.2. *Let $\lambda \neq 0$ be a partition that fits in the $(2n) \times (d - n - 1)$ rectangle and assume that λ has n -index i . For every summand $\mathbf{S}_\delta(\mathcal{B}^\vee) \subset \mathbf{S}_\lambda(\mathcal{B}^\vee \oplus \mathcal{B}^\vee) \otimes \wedge^\bullet(\mathcal{B})$, the partition δ satisfies condition (3.1.1) with $j = i$.*

In particular, all cohomology of $\mathbf{S}_\lambda(\mathcal{B}^\vee \oplus \mathcal{B}^\vee) \otimes \wedge^\bullet(\mathcal{B})$ vanishes.

Proposition 3.1.3. *Let $\lambda \neq 0$ be a partition that fits in the $(2n) \times (d - n - 1)$ rectangle and assume that λ has n -index i .*

(1) *If $i < n$, then for every summand $\mathbf{S}_\delta(\mathcal{B}^\vee) \subset \mathbf{S}_\lambda(\mathcal{B}^\vee \oplus \mathcal{B}^\vee) \otimes \text{Sym}^\bullet(\mathcal{B})$, the partition δ satisfies condition (3.1.1) with $j = i$.*

In particular, all cohomology of $\mathbf{S}_\lambda(\mathcal{B}^\vee \oplus \mathcal{B}^\vee) \otimes \text{Sym}^\bullet(\mathcal{B})$ vanishes.

(2) *If $i = n$ and $k \leq n$, then for every summand $\mathbf{S}_\delta(\mathcal{B}^\vee) \subset \mathbf{S}_\lambda(\mathcal{B}^\vee \oplus \mathcal{B}^\vee) \otimes \text{Sym}^k(\mathcal{B})$, the partition δ satisfies condition (3.1.1) with $j = i$.*

In particular, all cohomology of $\mathbf{S}_\lambda(\mathcal{B}^\vee \oplus \mathcal{B}^\vee) \otimes \text{Sym}^k(\mathcal{B})$ vanishes.

Proposition 3.1.4. *Let $\lambda \neq 0$ be a partition that fits in the $(2n) \times (d - n - t - 1)$ rectangle, for some $t \geq 0$.*

(1) *Assume that λ has n -index i . Let p_1, \dots, p_t be nonnegative integers and let $0 \leq k \leq n$. Then every summand*

$$\mathbf{S}_\delta(\mathcal{B}^\vee) \subset \mathbf{S}_\lambda(\mathcal{B}^\vee \oplus \mathcal{B}^\vee) \otimes \wedge^{p_1} \mathcal{B}^\vee \otimes \dots \otimes \wedge^{p_t} \mathcal{B}^\vee \otimes \wedge^k \mathcal{B}$$

satisfies condition (3.1.1) with $j = i$.

In particular, all cohomology of $\mathbf{S}_\lambda(\mathcal{B}^\vee \oplus \mathcal{B}^\vee) \otimes \wedge^{p_1} \mathcal{B}^\vee \otimes \dots \otimes \wedge^{p_t} \mathcal{B}^\vee \otimes \wedge^k \mathcal{B}$ vanishes.

(2) *Assume that λ has n -index i . Let p_1, \dots, p_t be nonnegative integers and let $0 \leq k \leq n$. All summands of*

$$\mathbf{S}_\lambda(\mathcal{B}^\vee \oplus \mathcal{B}^\vee) \otimes \wedge^{p_1} \mathcal{B}^\vee \otimes \dots \otimes \wedge^{p_t} \mathcal{B}^\vee \otimes \text{Sym}^k(\mathcal{B})$$

satisfy condition (3.1.1) for $j = i$, and therefore this bundle has no cohomology.

(3) Assume $\lambda \neq (1^p)$ has (p, n) -index i , for some $0 \leq p \leq n$. Let p_1, \dots, p_{t-1} be nonnegative integers. Then every summand

$$\mathbf{S}_\delta(\mathcal{B}^\vee) \subset \mathbf{S}_\lambda(\mathcal{B}^\vee \oplus \mathcal{B}^\vee) \otimes \wedge^{p_1} \mathcal{B}^\vee \otimes \dots \otimes \wedge^{p_{t-1}} \mathcal{B}^\vee$$

satisfies condition (3.1.1) with $j = i$.

In particular, all cohomology of $\mathbf{S}_\lambda(\mathcal{B}^\vee \oplus \mathcal{B}^\vee) \otimes \wedge^{p_1} \mathcal{B}^\vee \otimes \dots \otimes \wedge^{p_{t-1}} \mathcal{B}^\vee$ vanishes.

3.2. Littlewood–Richardson coefficients. For the proofs of the above propositions, we need a few preliminaries about the Littlewood–Richardson coefficients. The material below is well-known, but to establish the notation, we recall several definitions and basic facts, some of which can be found in [SS, §§2, 3].

For two partitions α and β of the same size $|\alpha| = |\beta|$, write $\alpha \geq \beta$ (α dominates β) if, for all m , we have

$$\alpha_1 + \dots + \alpha_m \geq \beta_1 + \dots + \beta_m.$$

Note that $\alpha \geq \beta$ if and only if $\alpha^\dagger \leq \beta^\dagger$.

Given partitions α, β, γ , let $c_{\alpha, \beta}^\gamma$ denote the Littlewood–Richardson coefficient, which is the multiplicity of the Schur functor \mathbf{S}_γ in the tensor product $\mathbf{S}_\alpha \otimes \mathbf{S}_\beta$.

The coefficient $c_{\alpha, \beta}^\gamma$ counts the number of Littlewood–Richardson tableaux. These are fillings of the skew tableau of shape γ/α with content β (i.e., for all i , the label i appears exactly β_i times) such that the following two properties hold:

- (semistandard) In each row, the entries are weakly increasing from left to right, and in each column, the entries are strictly increasing from top to bottom.
- (lattice word property) Let w be the word (called reading word) obtained by reading the entries in each row from right to left, starting with the top row and going down. For each i and m , let $w_i(m)$ be the number of times that i appears in the first m entries of w . Then for all m and i , we have $w_i(m) \geq w_{i+1}(m)$.

We collect a few facts about these coefficients in the next result.

Proposition 3.2.1. (1) For any complex vector bundles V, W , we have

$$\mathbf{S}_\gamma(V \oplus W) \cong \bigoplus_{\alpha, \beta} (\mathbf{S}_\alpha(V) \otimes \mathbf{S}_\beta(W))^{\oplus c_{\alpha, \beta}^\gamma},$$

where the sum is over all partitions α, β .

- (2) If $c_{\alpha, \beta}^\gamma \neq 0$, then $|\gamma| = |\alpha| + |\beta|$.
- (3) If $c_{\alpha, \beta}^\gamma \neq 0$, then γ contains both α and β , i.e., $\gamma_i \geq \max(\alpha_i, \beta_i)$ for all i .
- (4) In a Littlewood–Richardson tableau of shape γ/α and type β , all occurrences of the number i must appear in rows i and later.

As a consequence, if $c_{\alpha,\beta}^\gamma \neq 0$, then $\alpha + \beta$ dominates γ , i.e., for all m , we have

$$\sum_{i=1}^m (\alpha_i + \beta_i) \geq \sum_{i=1}^m \gamma_i.$$

(5) If $c_{\alpha,\beta}^\gamma \neq 0$, then γ dominates $\alpha \cup \beta$ (this is the partition obtained from all of the rows of α and β placed one after the other according to their lengths).

Proof. (1) See [SS, (4.5)] for a derivation.

(2) and (3) are clear from the interpretation in terms of tableaux.

(4) We prove this by induction on i . If $i = 1$, there is nothing to show. Now suppose the statement is true for i . Suppose that there is a Littlewood–Richardson tableau in which $i + 1$ appears in the first i rows. Let w be the reading word of this tableau. By the lattice word property, this instance of $i + 1$ cannot appear in a row before the earliest (relative to w) instance of i , so it must appear in row i , and it must appear to the left of i in the tableau. However, this violates the semistandard condition.

(5) This is a consequence of (4) since $\alpha \cup \beta = (\alpha^\dagger + \beta^\dagger)^\dagger$ and $c_{\alpha,\beta}^\gamma = c_{\alpha^\dagger,\beta^\dagger}^{\gamma^\dagger}$. □

3.3. Lemmas. To carry out the proofs of Propositions 3.1.2, 3.1.3, and 3.1.4, we first establish a few supporting results.

First, by Proposition 3.2.1(1), we have

$$\mathbf{S}_\lambda(\mathcal{B}^\vee \oplus \mathcal{B}^\vee) \cong \bigoplus_{\alpha,\beta} (\mathbf{S}_\alpha(\mathcal{B}^\vee) \otimes \mathbf{S}_\beta(\mathcal{B}^\vee))^{\oplus c_{\alpha,\beta}^\lambda} \cong \bigoplus_{\alpha,\beta,\gamma} \mathbf{S}_\gamma(\mathcal{B}^\vee)^{\oplus c_{\alpha,\beta}^\lambda c_{\alpha,\beta}^\gamma}. \tag{3.3.1}$$

Here, the number of rows of the partitions α, β, γ is less than or equal to n , while the number of rows in the partition λ is less than or equal to $2n$.

We assume first that λ is contained in the $(2n) \times (d - n - 1)$ rectangle as needed in Proposition 3.1.2 and 3.1.3.

We reserve i to be the n -index of λ .

Pick a triple α, β, γ such that $\mathbf{S}_\gamma(\mathcal{B}^\vee) \neq 0$, and $c_{\alpha,\beta}^\lambda c_{\alpha,\beta}^\gamma \neq 0$. We will deduce a number of restrictions on the partitions α, β, γ .

Lemma 3.3.2. *We have $\alpha_i \geq i$.*

Proof. Suppose that $\alpha_i < i$. Since λ has n -index i , recall from (2.3.8) that $\lambda_{i+1} = \dots = \lambda_{i+n} = i$ and thus $\lambda_i \geq i$. Then the i -th column of the skew shape λ/α has at least $n + 1$ boxes (in rows i through $n + i$). But then any valid Littlewood–Richardson tableau of shape λ/α needs at least $n + 1$ labels (because of the semistandard condition). This implies that $\beta_{n+1} > 0$, contradicting the fact that β has at most n rows. □

Lemma 3.3.3. *We have*

$$(\alpha_1 + \beta_1) + \dots + (\alpha_i + \beta_i) \leq i(d - n + i - 1).$$

Proof. By Proposition 3.2.1(5), we know that $\lambda \geq \alpha \cup \beta$, so that

$$\lambda_1 + \cdots + \lambda_{2i} \geq (\alpha \cup \beta)_1 + \cdots + (\alpha \cup \beta)_{2i} \geq (\alpha_1 + \beta_1) + \cdots + (\alpha_i + \beta_i). \tag{3.3.3a}$$

Since $\lambda_j \leq d - n - 1$ for $j = 1, \dots, i$ and $\lambda_j \leq i$ for $j = i + 1, \dots, 2i$, the lemma follows. \square

Lemma 3.3.4. *We have $i + 1 \leq \gamma_i \leq d - n + i - 1$.*

Proof. We know that $\alpha_i \geq i$ from Lemma 3.3.2. If in fact $\alpha_i \geq i + 1$, then we can use Proposition 3.2.1(3) to conclude that $\gamma_i \geq i + 1$.

Otherwise, we have $\alpha_i = i$. Suppose that $\gamma_i = i$. Then the i -th row of γ/α has no boxes. Since $c_{\alpha, \beta}^\lambda \neq 0$, there is a Littlewood–Richardson tableau of shape λ/α and type β . Since $\lambda_{i+n} = i$, the i -th column of λ/α has at least $i + n - \alpha_i^\dagger$ boxes, so that β has at least $i + n - \alpha_i^\dagger$ rows (by the semistandard condition).

Next, there is also a Littlewood–Richardson tableau of shape γ/α and type β . The integers in the interval $[i, i + n - \alpha_i^\dagger]$ cannot go in the first $i - 1$ rows of γ/α by Proposition 3.2.1(4), and cannot go in the i -th row since it is empty. Thus, these numbers must go in rows $i + 1$ or higher. Again, since $\gamma_i = i$, they are also constrained to the first i columns of γ/α as well. Now, in γ/α , the i -th column only has boxes in rows $\alpha_i^\dagger + 1, \dots, n$, at most. Consequently, the labels $[i, i + n - \alpha_i^\dagger]$ can only be placed in rows $\alpha_i^\dagger + 1, \dots, n$.

Suppose it is possible to do this. Consider the subdiagram D of γ/α consisting of boxes that are filled with entries $\geq i$. If we subtract $i - 1$ from every entry, we claim that the result is a valid Littlewood–Richardson tableau of shape D . Subtracting the same amount from each entry does not affect any of the semistandard inequalities. Furthermore, if w is the reading word of the Littlewood–Richardson tableau of γ/α of type β that we’re considering, and w' is the reading word of its restriction to D , then in the notation of Section 3.2, for $j \geq i$ and any m , we have $w'_j(m) = w_j(m)$. In particular, then $w'_j(m) \geq w'_{j+1}(m)$ for all $j \geq i$ and hence the result of subtracting $i - 1$ from all entries of D has the lattice word property.

But then we have too many labels: in fact, at least $n - \alpha_i^\dagger + 1$ labels and only $n - \alpha_i^\dagger$ rows to put them into. We have a contradiction to Proposition 3.2.1(4) and hence $\gamma_i \geq i + 1$.

Finally, again by Proposition 3.2.1(4), we have $\alpha + \beta \geq \gamma$. Using Lemma 3.3.3, we obtain

$$i\gamma_i \leq \gamma_1 + \cdots + \gamma_i \leq i(d - n + i - 1), \tag{3.3.4a}$$

and hence $\gamma_i \leq d - n + i - 1$.¹ \square

Lemma 3.3.5. (1) *If $i < n$, then $\gamma_{i+1} \geq i$.*

(2) *If $i = n$, then $\gamma_n \geq 2n$.*

Proof. Since $\lambda_{i+1} = i$, we have $\alpha_{i+1} \leq i$ by Proposition 3.2.1(3).

Let $c = i - \alpha_{i+1}$. Then λ/α contains the subrectangle occupying rows $i + 1, \dots, i + n$ and columns $\alpha_{i+1} + 1, \dots, i$, which has n rows and c columns. Since $c_{\alpha, \beta}^\lambda \neq 0$, we can fill λ/α with content β .

¹We could relax the condition that λ is contained in the $(2n) \times (d - n - 1)$ rectangle here. It would suffice to know that $\lambda_1 + \cdots + \lambda_i \leq i(d - n - 1) + i - 1$.

By examining the $n \times c$ subrectangle and using the semistandard property, we obtain $\beta_n \geq c$. Let β' be the result of subtracting c from all parts of β . Then

$$\mathbf{S}_\beta(\mathcal{B}^\vee) = (\det \mathcal{B}^\vee)^{\otimes c} \otimes \mathbf{S}_{\beta'}(\mathcal{B}^\vee)$$

since $\text{rank}(\mathcal{B}^\vee) = n$. Hence to compute $\mathbf{S}_\alpha(\mathcal{B}^\vee) \otimes \mathbf{S}_\beta(\mathcal{B}^\vee)$, we can first add c to all values of $(\alpha_1, \dots, \alpha_n)$ and then tensor with $\mathbf{S}_{\beta'}(\mathcal{B}^\vee)$.

In particular, if $i < n$, then $\gamma_{i+1} \geq \alpha_{i+1} + c = i$, again by [Proposition 3.2.1\(3\)](#). Otherwise, if $i = n$, since $\alpha_{n+1} = 0$ we find $c = n$. Using [Lemma 3.3.2](#), we have $\alpha_n \geq n$, and thus $\gamma_n \geq \alpha_n + c \geq 2n$. \square

3.4. Vanishing. We continue to use the notation from the previous section.

Proof of [Proposition 3.1.2](#). Assume λ fits in the $(2n) \times (d - n - 1)$ rectangle, and let $\mathbf{S}_\gamma(\mathcal{B}^\vee)$ be a summand of $\mathbf{S}_\lambda(\mathcal{B}^\vee \oplus \mathcal{B}^\vee)$. Consider the tensor product $\mathbf{S}_\gamma(\mathcal{B}^\vee) \otimes \wedge^k(\mathcal{B})$. First we use that $\wedge^k \mathcal{B} = \det(\mathcal{B}) \otimes \wedge^{n-k} \mathcal{B}^\vee$ and $\wedge^{n-k} \mathcal{B}^\vee = \mathbf{S}_{(1^{n-k})}(\mathcal{B}^\vee)$. The Pieri rule describes the outcome of tensoring with $\wedge^{n-k} \mathcal{B}^\vee$. The result is a sum over partitions where we add $n - k$ boxes, no two in the same row. Tensoring with $\det(\mathcal{B})$ is the same as subtracting 1 from all entries. Therefore, for any summand $\mathbf{S}_\delta(\mathcal{B}^\vee)$ of this tensor product, we have $\gamma_i - 1 \leq \delta_i \leq \gamma_i$. Hence we conclude from [Lemma 3.3.4](#) that

$$i \leq \delta_i \leq d - n + i - 1,$$

completing the argument in this case. \square

Proof of [Proposition 3.1.3](#). The Pieri rule applied to symmetric powers tells us that $\mathbf{S}_\nu(\mathcal{B}^\vee)$ is a summand of $\mathbf{S}_\mu(\mathcal{B}^\vee) \otimes \text{Sym}^k(\mathcal{B}^\vee)$ if and only if $|\nu| = |\mu| + k$ and the interlacing property $\nu_j \geq \mu_j \geq \nu_{j+1}$ holds for all j . In fact, it makes no difference if some entries of ν and μ are negative since we can make them nonnegative by twisting by powers of $\det(\mathcal{B}^\vee)$ and untwisting after.

If $\mathbf{S}_\delta(\mathcal{B}^\vee)$ is a summand of $\mathbf{S}_\gamma(\mathcal{B}^\vee) \otimes \text{Sym}^k(\mathcal{B})$, we obtain by dualizing that $\mathbf{S}_{-\delta}(\mathcal{B}^\vee)$ is a summand of $\mathbf{S}_{-\gamma}(\mathcal{B}^\vee) \otimes \text{Sym}^k(\mathcal{B}^\vee)$. Thus, $|\gamma| = |\delta| + k$ and the interlacing property gives

$$\gamma_{j+1} \leq \delta_j \leq \gamma_j.$$

(Thus, if $\mathbf{S}_\delta(\mathcal{B}^\vee)$ is a summand of $\mathbf{S}_\gamma(\mathcal{B}^\vee) \otimes \text{Sym}^k(\mathcal{B})$, then $\mathbf{S}_\gamma(\mathcal{B}^\vee)$ is a summand of $\mathbf{S}_\delta(\mathcal{B}^\vee) \otimes \text{Sym}^k(\mathcal{B}^\vee)$.)

Consider the n -index i of λ . If $i < n$, then [Lemma 3.3.5](#) tells us that $\gamma_{i+1} \geq i$. The interlacing property then forces $\delta_i \geq i$. If $i = n$, then $\gamma_n \geq 2n$. Since γ is obtained from δ by adding k boxes, we have

$$\delta_n \geq \gamma_n - k \geq 2n - k \geq n$$

since we assume that $k \leq n$.

In any case, under either assumption, we have shown that $\delta_i \geq i$ and also $\delta_i \leq \gamma_i \leq d - n + i - 1$ by [Lemma 3.3.4](#). This is what we set out to prove. \square

Proof of Proposition 3.1.4. Assume now that λ is contained in the $(2n) \times (d - n - t - 1)$ rectangle. For case (1), for a partition λ of n -index i , we have

$$\lambda_j \leq d - n - t - 1 \text{ for } j \leq i, \quad \lambda_j \leq i \text{ for } i + 1 \leq j \leq 2i.$$

Thus, by (3.3.3a) and (3.3.4a), we have

$$i\gamma_i \leq \gamma_1 + \cdots + \gamma_i \leq \lambda_1 + \cdots + \lambda_{2i} \leq i(d - n - t - 1) + i \cdot i \implies \gamma_i \leq d - n - t - 1 + i.$$

We also have $\gamma_i \geq i + 1$ by Lemma 3.3.4. We inspect the summands

$$\mathbf{S}_\delta(\mathcal{B}^\vee) \subset \mathbf{S}_\gamma(\mathcal{B}^\vee) \otimes \wedge^{p_1} \mathcal{B}^\vee \otimes \wedge^{p_2} \mathcal{B}^\vee \otimes \cdots \otimes \wedge^{p_t} \mathcal{B}^\vee \otimes \wedge^k \mathcal{B}.$$

We use $\wedge^k \mathcal{B} = \wedge^{n-k} \mathcal{B}^\vee \otimes \det \mathcal{B}$. By repeated application of the Pieri rule, and taking into account that tensorization by $\det \mathcal{B}$ subtracts one box from each entry, we conclude

$$\gamma_i - 1 \leq \delta_i \leq \gamma_i + t.$$

The conclusion follows since

$$i + 1 \leq \gamma_i \leq d - n - t - 1 + i \implies i \leq \delta_i \leq d - n + i - 1.$$

For (2), we saw in the proof of Proposition 3.1.3 that all summands $\mathbf{S}_\delta(\mathcal{B}^\vee)$ of $\mathbf{S}_\lambda(\mathcal{B}^\vee \oplus \mathcal{B}^\vee) \otimes \text{Sym}^k \mathcal{B}$ satisfy

$$i \leq \delta_i \leq d - n - t - 1 + i,$$

where the modified upper bound is due to the different size of the rectangle that contains λ . By the Pieri rule, tensorization by $(\wedge^\bullet \mathcal{B}^\vee)^{\otimes t}$ can only increase the lengths of rows, and if so by at most t boxes. Thus all summands $\mathbf{S}_\nu(\mathcal{B}^\vee)$ of $\mathbf{S}_\lambda(\mathcal{B}^\vee \oplus \mathcal{B}^\vee) \otimes \wedge^{p_1} \mathcal{B}^\vee \otimes \cdots \otimes \wedge^{p_t} \mathcal{B}^\vee \otimes \text{Sym}^k \mathcal{B}$ satisfy

$$i \leq \nu_i \leq d - n + i - 1,$$

as claimed.

Finally, we prove (3). If $\mathbf{S}_\gamma(\mathcal{B}^\vee)$ is a summand of $\mathbf{S}_\lambda(\mathcal{B}^\vee \oplus \mathcal{B}^\vee)$, then by the same reasoning as in Lemma 3.3.2 we have $i \leq \alpha_i \leq \gamma_i$. By (2.3.10)

$$\lambda_1, \dots, \lambda_i \leq d - n - t - 1, \quad \lambda_{i+1}, \dots, \lambda_{2i} \leq i + 1.$$

Using (3.3.3a) and (3.3.4a), we obtain

$$i\gamma_i \leq \gamma_1 + \cdots + \gamma_i \leq \lambda_1 + \cdots + \lambda_{2i} \leq i(d - n - t - 1) + i(i + 1) \implies \gamma_i \leq d - n - t + i.$$

By repeated application of the Pieri rule, all summands $\mathbf{S}_\delta(\mathcal{B}^\vee)$ of $\mathbf{S}_\gamma(\mathcal{B}^\vee) \otimes \wedge^{p_1} \mathcal{B}^\vee \otimes \cdots \otimes \wedge^{p_{t-1}} \mathcal{B}^\vee$ satisfy $\gamma_i \leq \delta_i \leq \gamma_i + (t - 1)$. Since $i \leq \gamma_i \leq d - n - t + i$, we have $i \leq \delta_i \leq d - n + i - 1$. Therefore, condition (3.1.1) is satisfied for δ and $j = i$, and all cohomology vanishes. \square

4. Corollaries

4.1. Corollary 1.2.1 and universality. We explain the universality arguments needed to derive Corollary 1.2.1 from the genus 0 computations in Theorems 1.1.2, 1.1.3, and 1.1.5.

Regarding equation (1.2.1a), we have the factorization

$$\sum_{n=0}^{\infty} q^n \chi(\text{Quot}_C(\mathbb{C}^N, n), \bigwedge_y L^{[n]}) = A^{\chi(\mathcal{O}_C)} \cdot B^{\chi(L)} \tag{4.1.1}$$

where $A, B \in 1 + q \mathbb{Q}[y][[q]]$ are two universal power series whose coefficients may depend on N but not on the pair (C, L) . This factorization is by now a standard fact; see for instance [EGL; OS; St] for various incarnations of this statement. To establish (1.2.1a), we show that

$$A = (1 - q)^{-1}, \quad B = (1 + qy)^N.$$

Specializing $C = \mathbb{P}^1$ and $\deg L = \ell \geq n$ in (4.1.1), and using Theorem 1.1.2(1) we obtain

$$[q^n]A \cdot B^{\ell+1} = \sum_{k=0}^n y^k \binom{N \chi(L)}{k},$$

where the brackets denote extracting the relevant coefficient in the q -expansion. By direct calculation, we also have

$$[q^n](1 - q)^{-1} \cdot ((1 + qy)^N)^{\ell+1} = \sum_{k=0}^n y^k \binom{N \chi(L)}{k}.$$

It remains to explain that the coefficients

$$[q^n]A \cdot B^{\ell+1} \text{ for all } \ell \geq n$$

determine the series A, B at most uniquely. We argue inductively, each coefficient at a time. Explicitly, we write

$$A = 1 + a_1q + a_2q^2 + \dots, \quad B = 1 + b_1q + b_2q^2 + \dots.$$

Then

$$[q^n]A \cdot B^{\ell+1} = a_n + (\ell + 1)b_n + \text{lower-order terms in } n.$$

The lower-order terms are determined by the induction hypothesis. The inductive step follows since the principal terms $a_n + (\ell + 1)b_n$ for all $\ell \geq n$ determine a_n, b_n at most uniquely.

For (1.2.1b) the argument is similar, using the factorization

$$\sum_{n=0}^{\infty} q^n \chi\left(\text{Quot}_C(\mathbb{C}^N, n), \bigotimes_{i=1}^t \left(\bigwedge_{y_i} M_i^{[n]}\right)^\vee\right) = A^{\chi(\mathcal{O}_C)} \cdot B_1^{\chi(M_1)} \dots B_t^{\chi(M_t)}.$$

This time, we specialize $C = \mathbb{P}^1$, and

$$M_1 = M_2 = \dots = M_t = M, \quad \deg M = m \geq n.$$

By [Theorem 1.1.5](#), we have

$$[q^n]A \cdot (B_1 \cdots B_t)^{m+1} = 1 \text{ for all } m \geq n.$$

Indeed, the only nonzero contribution appears from the free term $y_1 = \cdots = y_t = 0$ and yields the answer $\chi(\text{Quot}_{\mathbb{P}^1}(\mathbb{C}^N, n), \mathcal{O}) = 1$ since $\text{Quot}_{\mathbb{P}^1}(\mathbb{C}^N, n)$ is rational. By the above reasoning, the series

$$A = (1 - q)^{-1}, \quad B_1 \cdots B_t = 1$$

are uniquely determined. Next, we set

$$M_1 = L, \quad M_2 = \cdots = M_t = M,$$

where $\deg L = \deg M - 1 = m - 1 \geq n - 1$. This time around, [Theorem 1.1.5](#) implies

$$[q^n](A \cdot B_1^{-1}) \cdot (B_1 \cdots B_t)^{m+1} = 1 \implies A \cdot B_1^{-1} = (1 - q)^{-1}, \quad B_1 \cdots B_t = 1.$$

Therefore

$$A = (1 - q)^{-1}, \quad B_1 = \cdots = B_t = 1,$$

and [\(1.2.1b\)](#) follows.

Equation [\(1.2.1c\)](#) uses [Theorem 1.1.3\(1\)](#). Indeed, we have the factorization

$$\sum_{n=0}^{\infty} q^n \chi(\text{Quot}_C(\mathbb{C}^N, n), \text{Sym}_y L^{[n]}) = A^{\chi(\mathcal{O}_C)} \cdot B^{\chi(L)},$$

where $A, B \in 1 + q\mathbb{Q}[[y]][[q]]$. Fix $n \geq 1$. By [Theorem 1.1.3](#), if $\deg L = \ell \geq n$, we have

$$[q^n]A \cdot B^{\ell+1} = \sum_{k \geq 0} y^k \chi(\text{Quot}_{\mathbb{P}^1}(\mathbb{C}^N, n), \text{Sym}^k L^{[n]}) = \sum_{k=0}^n y^k (-1)^k \binom{-N(\ell+1)}{k} \pmod{y^{n+1}}.$$

Both sides of this identity are polynomials in $(\ell + 1)$. (On the left hand side, these polynomials depend on the first q -coefficients $a_1, \dots, a_n, b_1, \dots, b_n$ of A, B considered modulo y^{n+1} .) Consequently, the same equality holds for all values of ℓ without restrictions. Hence, for all L , we have

$$\begin{aligned} \sum_{n=0}^{\infty} \sum_{k=0}^n q^n y^k \chi(\text{Quot}_{\mathbb{P}^1}(\mathbb{C}^N, n), \text{Sym}^k L^{[n]}) &= \sum_{n=0}^{\infty} q^n \left(\sum_{k=0}^n y^k (-1)^k \binom{-N(\ell+1)}{k} \right) \\ &= (1 - q)^{-1} \cdot (1 - qy)^{-N\chi(L)} \end{aligned}$$

as stated in [\(1.2.1c\)](#). □

4.2. Corollary 1.1.7. We analyze the cohomology groups of $L^{[n]}$ for all line bundles $L \rightarrow \mathbb{P}^1$ using a few simple considerations. The corollary can also be derived by combining the methods of [\[BGS, Corollary 9.3\]](#) when adapted to the case of the projective line, followed by a calculation on the symmetric product.

Let $p \in \mathbb{P}^1$ and write $\mathcal{Q}_p = \mathcal{Q}|_{p \times \text{Quot}}$. The exact sequence

$$0 \rightarrow L(-p) \rightarrow L \rightarrow L_p \rightarrow 0$$

yields an exact sequence over Quot:

$$0 \rightarrow L(-p)^{[n]} \rightarrow L^{[n]} \rightarrow \mathcal{Q}_p \rightarrow 0. \quad (4.2.1)$$

When $\deg L \geq n + 1$, the bundles $L^{[n]}$ and $L(-p)^{[n]}$ carry no higher cohomology by Theorems 1.1.2(1) or 1.1.3(1) for $k = 1$. Taking cohomology in (4.2.1), we obtain

$$H^i(\mathcal{Q}_p) = 0, \quad i \geq 1. \quad (4.2.2)$$

We go back to (4.2.1) written for arbitrary L , not necessarily sufficiently positive. Considering cohomology again and using (4.2.2), we obtain

$$H^i(L^{[n]}) = H^i(L(-p)^{[n]}), \quad i \geq 2. \quad (4.2.3)$$

Since $H^i(L^{[n]}) = 0$ for $\deg L \geq n$ and $i \geq 2$, it follows from (4.2.3) that $H^i(L^{[n]}) = 0$ for all $i \geq 2$ and all L . \square

Acknowledgements

We thank Shubham Sinha and Jerzy Weyman for useful related conversations. A. Marian was supported by NSF grant DMS 1902310, D. Oprea was supported by NSF grant DMS 1802228, and S. V Sam was supported by NSF grant DMS 1812462.

References

- [A] N. Arbesfeld, “K-theoretic Donaldson–Thomas theory and the Hilbert scheme of points on a surface”, *Algebr. Geom.* **8**:5 (2021), 587–625. [MR](#)
- [B] R. Bott, “Homogeneous vector bundles”, *Ann. of Math. (2)* **66** (1957), 203–248. [MR](#)
- [BCS] T. Braden, L. Chen, and F. Sottile, “The equivariant Chow rings of Quot schemes”, *Pacific J. Math.* **238**:2 (2008), 201–232. [MR](#)
- [Be] A. Bertram, “Quantum Schubert calculus”, *Adv. Math.* **128**:2 (1997), 289–305. [MR](#)
- [BGS] I. Biswas, C. Gangopadhyay, and R. Sebastian, “Infinitesimal deformations of some Quot schemes”, *Int. Math. Res. Not.* **2024**:9 (2024), 8067–8100. [MR](#)
- [C] L. Chen, “Quantum cohomology of flag manifolds”, *Adv. Math.* **174**:1 (2003), 1–34. [MR](#)
- [CF1] I. Ciocan-Fontanine, “The quantum cohomology ring of flag varieties”, *Trans. Amer. Math. Soc.* **351**:7 (1999), 2695–2729. [MR](#)
- [CF2] I. Ciocan-Fontanine, “On quantum cohomology rings of partial flag varieties”, *Duke Math. J.* **98**:3 (1999), 485–524. [MR](#)
- [D] G. Danila, “Sections de la puissance tensorielle du fibré tautologique sur le schéma de Hilbert des points d’une surface”, *Bull. Lond. Math. Soc.* **39**:2 (2007), 311–316. [MR](#)
- [EGL] G. Ellingsrud, L. Göttsche, and M. Lehn, “On the cobordism class of the Hilbert scheme of a surface”, *J. Algebraic Geom.* **10**:1 (2001), 81–100. [MR](#)
- [I] A. Ito, “On birational geometry of the space of parametrized rational curves in Grassmannians”, *Trans. Amer. Math. Soc.* **369**:9 (2017), 6279–6301. [MR](#)
- [J] S.-Y. Jow, “The effective cone of the space of parametrized rational curves in a Grassmannian”, *Math. Z.* **272**:3-4 (2012), 947–960. [MR](#)
- [K] B. Kim, “Quot schemes for flags and Gromov invariants for flag varieties”, preprint, 1995. [arXiv 9512003v1](#)

- [Kr1] A. Krug, “Extension groups of tautological sheaves on Hilbert schemes”, *J. Algebraic Geom.* **23**:3 (2014), 571–598. [MR](#)
- [Kr2] A. Krug, “Extension groups of tautological bundles on punctual Quot schemes of curves”, *J. Math. Pures Appl.* (9) **189** (2024), art. id. 103600, 30 pp. [MR](#)
- [MN] A. Marian and A. Negut, “Derived categories of Quot schemes on smooth curves and tautological bundles”, preprint, 2024. [arXiv 2411.08695v1](#)
- [OS] D. Oprea and S. Sinha, “Euler characteristics of tautological bundles over Quot schemes of curves”, *Adv. Math.* **418** (2023), art. id. 108943, 45 pp. [MR](#)
- [Sc1] L. Scala, “Cohomology of the Hilbert scheme of points on a surface with values in representations of tautological bundles”, *Duke Math. J.* **150**:2 (2009), 211–267. [MR](#)
- [Sc2] L. Scala, “Higher symmetric powers of tautological bundles on Hilbert schemes of points on a surface”, preprint, 2015. [arXiv 1502.07595v2](#)
- [SS] S. V. Sam and A. Snowden, “Introduction to twisted commutative algebras”, preprint, 2012. [arXiv 1209.5122v1](#)
- [St] S. Stark, “Cosection localization and the Quot scheme $\text{Quot}^{\ell}(\mathcal{E})$ ”, *Proc. A* **478**:2268 (2022), art. id. 20220419, 16 pp. [MR](#)
- [Str] S. A. Strømme, “On parametrized rational curves in Grassmann varieties”, pp. 251–272 in *Space curves* (Rocca di Papa, Italy, 1985), Lecture Notes in Math. **1266**, Springer, 1987. [MR](#)
- [W] J. Weyman, *Cohomology of vector bundles and syzygies*, Cambridge Tracts in Mathematics **149**, Cambridge University Press, 2003. [MR](#)

Communicated by Gavril Farkas

Received 2023-06-21 Revised 2025-04-17 Accepted 2025-05-22

a.marian@northeastern.edu

Department of Mathematics, Northeastern University, Boston, MA, United States

doprea@math.ucsd.edu

Department of Mathematics, University of California San Diego, La Jolla, CA, United States

ssam@ucsd.edu

Department of Mathematics, University of California San Diego, La Jolla, CA, United States

Algebra & Number Theory

msp.org/ant

EDITORS

MANAGING EDITOR

Antoine Chambert-Loir
Université Paris-Diderot
France

EDITORIAL BOARD CHAIR

David Eisenbud
University of California
Berkeley, USA

BOARD OF EDITORS

Jason P. Bell	University of Waterloo, Canada	Michael J. Larsen	Indiana University Bloomington, USA
Olivier Benoist	Ecole Normale Supérieure, France	Philippe Michel	École Polytechnique Fédérale de Lausanne
Bhargav Bhatt	University of Michigan, USA	Martin Olsson	University of California, Berkeley, USA
Frank Calegari	University of Chicago, USA	Irena Peeva	Cornell University, USA
J-L. Colliot-Thélène	CNRS, Université Paris-Saclay, France	Jonathan Pila	University of Oxford, UK
Brian D. Conrad	Stanford University, USA	Anand Pillay	University of Notre Dame, USA
Samit Dasgupta	Duke University, USA	Bjorn Poonen	Massachusetts Institute of Technology, USA
Hélène Esnault	Freie Universität Berlin, Germany	Victor Reiner	University of Minnesota, USA
Gavril Farkas	Humboldt Universität zu Berlin, Germany	Peter Sarnak	Princeton University, USA
Sergey Fomin	University of Michigan, USA	Michael Singer	North Carolina State University, USA
Edward Frenkel	University of California, Berkeley, USA	Vasudevan Srinivas	SUNY Buffalo, USA
Wee Teck Gan	National University of Singapore	Shunsuke Takagi	University of Tokyo, Japan
Andrew Granville	Université de Montréal, Canada	Pham Huu Tiep	Rutgers University, USA
Ben J. Green	University of Oxford, UK	Ravi Vakil	Stanford University, USA
Christopher Hacon	University of Utah, USA	Akshay Venkatesh	Institute for Advanced Study, USA
Roger Heath-Brown	Oxford University, UK	Melanie Matchett Wood	Harvard University, USA
János Kollár	Princeton University, USA	Shou-Wu Zhang	Princeton University, USA

PRODUCTION

production@msp.org

Silvio Levy, Scientific Editor


See inside back cover or msp.org/ant for submission instructions.

The subscription price for 2026 is US \$590/year for the electronic version, and \$865/year (+\$75, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to MSP.

Algebra & Number Theory (ISSN 1944-7833 electronic, 1937-0652 printed) at Mathematical Sciences Publishers, 2000 Allston Way # 59, Berkeley, CA 94701-4004, is published continuously online.

ANT peer review and production are managed by EditFLOW[®] from MSP.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2026 Mathematical Sciences Publishers

Algebra & Number Theory

Volume 20 No. 5 2026

A coarse Jacquet–Zagier trace formula for $GL(n)$, with applications	861
LIYANG YANG	
On the cohomology of tautological bundles over Quot schemes of curves	943
ALINA MARIAN, DRAGOS OPREA and STEVEN V SAM	
Admissible pairs and p-adic Hodge structures, I: transcendence of the de Rham lattice	971
SEAN HOWE and CHRISTIAN KLEVDAL	
Local square mean in the hyperbolic circle problem	1029
ANDRÁS BIRÓ	