

Algebra & Number Theory

Volume 20
2026
No. 6

**The geometry of the unipotent component of the moduli space of
Weil–Deligne representations**

Daniel Funck



The geometry of the unipotent component of the moduli space of Weil–Deligne representations

Daniel Funck

We study the moduli space of unipotent Weil–Deligne representations valued in a split reductive group G and characterise which irreducible components are smooth. We apply these smoothness results to show that a certain space of ordinary automorphic forms is a locally generically free module over the corresponding global deformation ring.

1. Introduction and overview	1125
2. Considerateness and the relation to the stack of L-parameters	1128
3. Smoothness results for X_C	1132
4. Automorphic forms for unitary groups	1141
5. Galois representations and deformation rings	1145
Appendix: Weighted Dynkin diagrams for distinguished orbits in types D_n and E_n with $n \leq 7$	1156
Acknowledgements	1157
References	1157

1. Introduction and overview

Let F be a local p -adic field and let G be a connected reductive algebraic group over F and \hat{G} its Langlands dual. The local Langlands conjectures (proven for GL_n by Harris and Taylor in [HT01]) stipulate the existence of a natural map with finite fibres:

$$\frac{\{\text{smooth irreducible representations of } G(F)\}}{\{\text{isomorphism}\}} \rightarrow \frac{\{\text{L-parameters of } {}^L G\}}{\{\hat{G}\text{-conjugacy}\}}$$

Let $l \neq p$ be a prime. Let $L \subset \overline{\mathbb{Q}_l}$ be an l -adic field and \mathcal{O} its ring of integers with residue field \mathbb{F} . In recent years, through the work in [BG19; Hel23; DHKM20; Zhu25; FS25], there has been great interest in studying the properties of a moduli space of L-parameters $\text{Loc}_{\hat{G}, \mathcal{O}}$ and a closely related space, the moduli space of framed L-parameters $\text{Loc}_{\hat{G}, \mathcal{O}}^{\square}$.

The spaces $\text{Loc}_{\hat{G}, \mathcal{O}}^{\square}$ and $\text{Loc}_{\hat{G}, \mathcal{O}}$ can be respectively defined as the scheme whose R -points are the set of L-parameters with R coefficients, and the algebraic stack obtained from the stack quotient of $\text{Loc}_{\hat{G}, \mathcal{O}}^{\square}$ modulo the natural action of \hat{G} via conjugation (that is, equivalence of representations):

$$\text{Loc}_{\hat{G}, \mathcal{O}}^{\square}(R) = \{\text{L-parameters of } \hat{G} \text{ with } R\text{-coefficients}\}, \quad \text{Loc}_{\hat{G}, \mathcal{O}} = [\text{Loc}_{\hat{G}, \mathcal{O}}^{\square} / \hat{G}].$$

MSC2020: primary 11F70, 11F80; secondary 11F33.

Keywords: number theory, moduli space of Langlands parameters, Galois representation, patching, automorphic form.

In addition to this interpretation, the scheme $\mathrm{Loc}_{\hat{G}, \mathcal{O}}^{\square}$ satisfies the secondary property that the (completions of) local rings of $\mathrm{Loc}_{\hat{G}, \mathcal{O}}^{\square}$ can be interpreted as local Galois deformation rings. In this way, it is hoped to better understand these rings, which are crucial ingredients in the Taylor–Wiles–Kisin and Calegari–Geraghty patching methods.

Let W_F be the Weil group of the field F . In the case of a split reductive group G , one would want to define an L-parameter as a homomorphism $W_F \rightarrow \hat{G}(R)$ which satisfies some kind of continuity, but the problem with this naive approach is that the ring R has no topological structure in general. Historically, there have been multiple solutions to this issue, with varying degrees of usefulness. We are interested in the moduli spaces of Bellovin and Gee [BG19], defined via Weil–Deligne representations (defined below); and of Dat, Helm, Kurinczuk and Moss [DHKM20], defined through representations of a particular dense subgroup $W_F^0 \subseteq W_F$ (defined in Section 1.2 of the reference).

Definition 1.1. A Weil–Deligne representation valued in G with R -coefficients is a pair (r, N) , where $r : W_F \rightarrow G(R)$ is a homomorphism with open kernel and N is an element of the nilpotent cone $\mathcal{N}_G(R) \subseteq \mathrm{Lie}(G)(R)$ such that $\mathrm{Ad}(g)N = |g|N$ for all $g \in W_F$, where $|\cdot| : W_F \rightarrow F^\times \rightarrow \mathbb{R}^{\geq 0}$ is the valuation of W_F coming from local class field theory.

The two moduli problems of [BG19] and [DHKM20] are representable by algebraic stacks $\mathrm{Loc}_{G, \mathcal{O}}^{BG}$ and $\mathrm{Loc}_{G, \mathcal{O}}$, respectively, and are disjoint unions of quotient stacks.

It is known, as in Proposition 2.7 of [DHKM20], that these definitions give isomorphic moduli spaces over fields of characteristic 0. In positive characteristic l (or mixed characteristic), only the latter moduli space is generally well-behaved, giving the deformation rings as the completions of its local rings. However, when l is greater than the Coxeter number h_G of G , the exponential and logarithm maps of Proposition 2.7 of [DHKM20] that arise from Grothendieck’s l -adic monodromy are well defined polynomials, and we obtain an isomorphism between the two moduli spaces in this case too.

Let \mathcal{O} be a discrete valuation ring (or field) of residue characteristic (resp. characteristic) $l > h_G$ or 0. Let $\mathcal{N}_G \subseteq \mathfrak{g}$ be the nilpotent cone inside the Lie algebra \mathfrak{g} . In this paper, we seek to understand the irreducible components of the scheme studied in [Hel23]. This is a reduced affine scheme of finite type $S_{G, \mathcal{O}}$, over the ring \mathcal{O} , whose R -points (R an \mathcal{O} -algebra) are given by

$$S_{G, \mathcal{O}}(R) = \{(\Phi, N) \in G(R) \times \mathcal{N}_G(R) \mid \mathrm{Ad}(\Phi)N = qN\},$$

where $q \in \mathcal{O}^\times$ is some prime power.

The scheme $S_{G, \mathcal{O}}$ is naturally the space of framed unipotent Weil–Deligne representations over \mathcal{O} with values in G (following Definition 2.1.2 of [BG19]). We will be interested especially in the case where \mathcal{O} is the ring of integers in a finite extension of \mathbb{Q}_l because the \mathfrak{m}_R -adic completion of the local rings R of the closed points of this scheme can be interpreted as local Galois deformation rings for well behaved l (in fact, whenever the exponential and logarithm maps of Grothendieck’s l -adic monodromy theorem give an isomorphism onto a connected component, as above). Since $S_{G, \mathcal{O}}$ is a connected component of the tame parameters $Z^1(W^0/P, G)_{\mathcal{O}}$, equation 4.5 of [DHKM20] extends the description of $S_{G, \overline{\mathbb{Q}}_l}$

for various groups G to a description of the geometry of many other connected components of $\text{Loc}_{G, \overline{\mathbb{Q}_l}}^\square$ (those which correspond to the case where the action Ad_φ is trivial).

Outline. In Section 2, we collect some basic results for S_G in the mixed characteristic setting. The results of this paper depend strongly on the technical relationship between the residue characteristic l and the element $q \in \mathcal{O}$. Therefore, the section begins by defining the notions of G -banality and q -considerateness, and explaining how they are related. We then give a description of a decomposition of S_G into (unions of) irreducible components, generalising the decomposition of S_{GL_n} given in Proposition 2.1 of [Hel23], as follows: Let

$$p : S_{G, \mathcal{O}} \rightarrow \mathcal{N}_G$$

be the projection map onto the second factor. Let $C \subset \mathcal{N}_{G, L}$ be a G -conjugacy class inside $\mathcal{N}_{G, L}$. (In the case of GL_n , these can be characterised by partitions of n and in this situation we will denote the conjugacy class corresponding to λ by C_λ .) We note that because $S_{G, \mathcal{O}}$ is flat over \mathcal{O} , the irreducible components biject naturally with those of $S_{G, L}$. Then $\overline{p^{-1}(C)} \subseteq S_{G, \mathcal{O}}$ is a union of irreducible components of $S_{G, \mathcal{O}}$ (and, in the case of $G = \text{GL}_n$, is itself irreducible). In Section 3, we expand on and generalise the results of Bellovin [Bel16, Section 7.2 and Proposition 7.10] by proving our Theorems 3.2 and 3.4, which together state:

Theorem 1.2. *Assume q is considerate towards $G_{\mathcal{O}}$ (see Definition 2.1).*

- (1) *Suppose $C \subseteq \mathcal{N}_{G, L}$ is a distinguished nilpotent orbit, or the zero orbit with corresponding component $X_C = \overline{p^{-1}(C)}$. Then X_C is a disjoint union of smooth connected components.*
- (2) *Conversely, when $C \subseteq \mathcal{N}_{G, L}$ is a nontrivial nondistinguished orbit, the scheme X_C is singular.*

In Sections 4 and 5, we apply the smoothness result of Section 3 to Hida families of ordinary automorphic forms using the Taylor–Wiles–Kisin patching method in a situation very similar to that studied in [Ger19]. Let l be a prime and K be a finite extension of \mathbb{Q}_l with ring of integers \mathcal{O} . Let F^+ be a totally real global number field, and consider an imaginary quadratic extension F of F^+ . The Galois representations considered will correspond to certain Hida families of ordinary automorphic forms on a unitary algebraic group G_D/F^+ which is a unitary form of a unit group of a division algebra D/F^+ . We will define a certain space of Hida families of ordinary automorphic forms $S^{\text{ord}}(U(l^\infty), L/\mathcal{O})_m$ for G_D with Hecke operators \mathbb{T} and a corresponding deformation ring R_S^{univ} . We will then use the Taylor–Wiles patching method to deduce the following theorem.

Theorem 1.3 (Theorem 5.13). *Suppose $l > n$. The module $S^{\text{ord}}(U(l^\infty), L/\mathcal{O})_m^\vee[1/l]$ is a finite locally free $R_S^{\text{univ}}[1/l]$ -module.*

As a consequence, we can deduce that $R_S^{\text{univ}}[1/l] \cong \mathbb{T}[1/l]$, and that the multiplicity of automorphic forms with a given characteristic zero Galois representation is constant along connected components of $R_S^{\text{univ}}[1/l]$. One can then extend any such multiplicity results from the classical case to the case of nonclassical Hida families.

2. Considerateness and the relation to the stack of L-parameters

Let \mathcal{O} be a discrete valuation ring or a field with residue field \mathbb{F} of characteristic l or 0 and fraction field L . Let G be a connected reductive algebraic group over \mathcal{O} , let \mathfrak{g} be its Lie algebra, and let h_G be its Coxeter number. Throughout the paper, we assume $l > h_G$ whenever the residue characteristic of \mathcal{O} is finite.

Definition 2.1. Let h_G be the Coxeter number of G . Let $q \in \mathcal{O}^\times$ be an element of \mathcal{O} such that $q^k - 1$ is invertible in \mathcal{O} for all $k \leq h_G$. When this occurs, we say that q is *considerate* towards G over \mathcal{O} .

In applications, the ring \mathcal{O} will either be a field or the ring of integers in some field extension of \mathbb{Q}_l . Notice that, in this case, q -considerateness is equivalent to the condition that $1, q, q^2, \dots, q^{h_G}$ are all distinct in the residue field \mathbb{F} (in a sense, q “treads lightly” around G).

Definition 2.2. Let G be a split reductive group over a field L of characteristic l .

- l is called *G-banal* if l does not divide the order of the finite group $G(\mathbb{F}_q)$.
- l is called *geometrically G-banal* if, for any algebraically closed field E of characteristic l , any $\phi \in \text{Loc}_{G,E}^\square$ can be *Frobenius twisted* by some $g \in C_G(\phi(I_F))$ (that is, the centraliser of the inertia subgroup) so that ϕ^g is a smooth point of $\text{Loc}_{G,E}^\square$.

The *Frobenius twist* of a representation $\phi : W_F \rightarrow G(L)$ by $g \in C_G(\phi(I_F))$ is the representation $\phi^g : W_F \rightarrow G(L)$ whose restriction to the inertia subgroup equals ϕ , and for which $\phi^g(\text{Frob}) = \phi(\text{Frob})g$.

Remark. Let \hat{G} denote the Langlands dual group of G . We remark that the notion “ l is geometrically \hat{G} -banal” is precisely the notion that “ l is ${}^L G$ -banal”, as defined in Definition 5.27 of [DHKM20]. We introduce the notion simply to remove the extraneous Langlands dual, which is only required in the following Proposition.

Proposition 2.3. *Suppose that \mathbb{F} is a field of positive characteristic $l > h_G$ and that G is a split reductive group. Then we have the following implications.*

- *If q is considerate towards G/\mathbb{F} , then l is geometrically G -banal.*
- *If l is geometrically \hat{G} -banal (that is, ${}^L G$ -banal), then l is G -banal.*
- *If $G = \text{GL}_n$ or SL_n , then the condition “ l is G -banal” implies that q is considerate towards \hat{G}/\mathcal{O} . Thus, in these cases all three concepts are equivalent.*

Proof. By definition, q is considerate towards G/\mathbb{F} when the order of q within \mathbb{F} is greater than the Coxeter number $h = h_G$. This is equivalent to $\prod_{n \leq h} \Phi_n(q) \neq 0$ inside \mathbb{F} , where Φ_n is the n -th cyclotomic polynomial. This is the polynomial $\chi_{G,1}^*(q)$ of [DHKM20, Definition B.3]. Hence, by Theorems 5.6 and 5.7 of [DHKM20], this condition implies that l is ${}^L G$ -banal.

That l being ${}^L G$ -banal implies that l is G -banal is a consequence of the Chevalley–Steinberg formula

$$|G(\mathbb{F}_q)| = q^N \prod_d (q^d - 1)$$

(see [Ste16, Theorem 25a]), where d ranges over the fundamental degrees of the Weyl group of G . If l divides $\prod_d (q^d - 1)$, then l certainly divides $\prod_{n \leq h} \Phi_n(q)$ as the Coxeter number is the highest fundamental degree. This shows the second statement by virtue of Theorem 5.7 in [DHKM20].

In the case $G = \mathrm{SL}_n$, we get $|\mathrm{SL}_n(\mathbb{F}_q)| = q^N \prod_{i=2}^n (q^i - 1)$. Hence, if l is G -banal, the order of q in \mathbb{F} is at least n . Thus q is considerate towards $\hat{G}_{/\mathbb{F}}$. The case of GL_n is similar. \square

Remark. It is worth noting that Corollary 5.27 of [DHKM20] gives the criterion that G -banal and ${}^L G$ -banal are equivalent concepts whenever G is unramified and has no exceptional factors (where triality forms of type D_4 are also considered exceptional), but this does not hold in general (see, for example, Remark 5.22 of [DHKM20]). Outside the case of type A_n , considerateness is strictly weaker than geometric- G -banality. For example, the order of $G = \mathrm{Sp}_6(\mathbb{F}_q)$ is

$$|\mathrm{Sp}_6(\mathbb{F}_q)| = q^9 (q^2 - 1)(q^4 - 1)(q^6 - 1)$$

and the Coxeter number of $\hat{G} = \mathrm{SO}_7$ is $h = 6$. However, if $q^5 \equiv 1 \pmod{l}$, then l is G -banal, but q is not considerate towards $\hat{G}_{\mathcal{O}}$.

Definition 2.4. We define the affine scheme $S_{G,\mathcal{O}}$ over \mathcal{O} as the scheme whose R points (for R an \mathcal{O} -algebra) are $\{(\Phi, N) \in G(R) \times \mathcal{N}_G(R) : \mathrm{Ad}(\Phi)N = qN\}$.

Corollary 5.4 of [Bel16] shows that this is a reduced scheme when \mathcal{O} is a characteristic zero field, and hence, is a variety (i.e., a finite-type, separated, reduced scheme over a field). As discussed in the introduction, we may picture $S_{G,\mathcal{O}}$ as the moduli space of unipotent Weil–Deligne representations (r, N) valued in $G_{\mathcal{O}}$. In this context, “unipotent” means that the restriction of r to the inertia subgroup I_F is trivial (that is, $r(I_F) = 1$).

Proposition 2.5. (1) *Suppose q is considerate towards $G_{/\mathcal{O}}$. If $(\Phi, N) \in G \times \mathfrak{g}$ satisfies $\mathrm{Ad}(\Phi)N = qN$, then $N \in \mathcal{N}_G$. Hence,*

$$S_G(R) = \{(\Phi, N) \in G(R) \times \mathfrak{g}(R) : \mathrm{Ad}(\Phi)N = qN\},$$

and the requirement that $N \in \mathcal{N}_G(R)$ is redundant.

(2) *If G is split and $l > h_G$, then $S_{G,\mathcal{O}}$ is isomorphic to a closed subscheme of the moduli space of tame parameters $Z^1(W_F^0/P_F, G)_{\mathcal{O}}$ (see Section 1.2 of [DHKM20] for a definition of this space).*

(3) *When l is geometrically G -banal, this space is a connected component of $Z^1(W_F^0/P_F, G)_{\mathcal{O}}$.*

Proof. Because $l > h_G$, the prime l is very good in the notation of Section 2.4 of [Cot22]. Hence, by Theorem 4.13 of [Cot22], we have an isomorphism of \mathcal{O} -algebras

$$\mathcal{O}[\mathfrak{g}]^G \rightarrow \mathcal{O}[\mathfrak{t}]^W$$

given by the restriction of functions on \mathfrak{g} to \mathfrak{t} , where \mathfrak{t} is a Cartan subalgebra of \mathfrak{g} and W is the Weyl group.

By Chapter 3 of [Hum90] (see Table 1 of Section 3.7 and Table 2 of Section 3.18) the generators of $\mathcal{O}[\mathfrak{t}]^W$ are homogeneous of degree at most the Coxeter number h_G , and hence the same is true for $\mathcal{O}[\mathfrak{g}]^G$. (Although [Hum90] considers only the case of characteristic zero, the results extend to \mathcal{O} because $|W|$ is invertible inside \mathcal{O} and $\mathcal{O}[\mathfrak{t}]^W$ is a free \mathcal{O} -module.)

Let s be a generator of $\mathcal{O}[\mathfrak{g}]^G$ and suppose $(\Phi, N) \in G(R) \times \mathfrak{g}(R)$ satisfies $\text{Ad}(\Phi)N = qN$. Then as s is G -invariant and homogeneous of degree at most the Coxeter number h_G , the condition $s(\text{Ad}(\Phi)N) = s(qN)$ implies $s(N) = q^i s(N)$ for some $i \leq h_G$. As q is considerate towards G/\mathcal{O} , we see that $q^i - 1$ is a nonzero divisor in \mathcal{O} , and hence that $s(N) = 0$. Thus the image of N in the GIT quotient \mathfrak{g}/G is zero. Since l is very good, Theorem 4.12 of [Cot22] shows that N lies in the set of R -points of the nilpotent cone. Part (1) of the proposition follows.

Suppose G is a split group. As $l \neq p$, the space $Z^1 = Z^1(W_F^0/P_F, G)_{\mathcal{O}}$ has a model as a flat affine scheme over \mathcal{O} with R -points equal to

$$Z^1(W_F^0/P_F, G)_{\mathcal{O}}(R) = \{(\phi, \sigma) \in G(R)^2 : \phi\sigma\phi^{-1} = \sigma^q\}.$$

Since $l > h_G$, the exponential and logarithm maps of Section 6 of [BDP17] are well defined polynomials, and thus we have an isomorphism between the nilpotent cone in \mathcal{N}_G and unipotent cone \mathcal{U}_G . Hence, we have a map

$$S_{G,\mathcal{O}} \rightarrow Z^1(W_F^0/P_F, G)_{\mathcal{O}}, \quad (\Phi, N) \mapsto (\Phi, \exp N),$$

which is an isomorphism onto the closed subscheme of $Z^1(W_F^0/P_F, G)_{\mathcal{O}}$ given by those elements (ϕ, σ) with $\sigma \in \mathcal{U} \subseteq G$, where \mathcal{U} is the unipotent cone. This shows (2).

For (3), suppose l is geometrically G -banal. Let \mathcal{U}^+ be the scheme-theoretic image of $Z^1(W_F^0/P_F, G)_{\mathcal{O}}$ through the second projection onto G . Proposition 2.6 of [DHKM20] tells us that the underlying reduced scheme of \mathcal{U}^+ is a subscheme of $\{\sigma \in G/\mathbb{F} : \sigma^M = 1\}$ for some fixed $M \in \mathbb{N}$. Thus \mathcal{U}^+ is 0-dimensional over \mathcal{O} . As $Z^1(W_F^0/P_F, G)$ is flat and \mathcal{O} is a discrete valuation ring, \mathcal{U}^+ is a finite flat \mathcal{O} -scheme.

We claim that no two distinct \mathcal{O} -points of \mathcal{U}^+ reduce to the same \mathbb{F} -point. The preimages of the L -points (resp. \mathbb{F} -points) of \mathcal{U}^+ in $Z^1(W_F^0/P_F, G)_L$ (resp. $Z^1(W_F^0/P_F, G)_{\mathbb{F}}$) are (unions of) connected components, thus it suffices to show that no two distinct points in $\mathcal{U}^+(L)$ reduce to the same point in $\mathcal{U}^+(\mathbb{F})$. This follows in turn from the statement that no two connected components of $Z^1(W_F^0/P_F, G)_L$ reduce to the same component of $Z^1(W_F^0/P_F, G)_{\mathbb{F}}$, which follows from Proposition 5.26 of [DHKM20], which states that $Z^1(W_F^0/P_F, G)_{\mathbb{F}}$ is reduced. We can conclude that the point $\text{Spec } \mathcal{O} \hookrightarrow \mathcal{U}^+$ defined by $\sigma = 1$ is a connected component of \mathcal{U}^+ . It follows that S_G , the preimage of this point, is a connected component of $Z^1(W_F^0/P_F, G)_{\mathcal{O}}$. \square

We will also need the following results.

Proposition 2.6. (1) *The algebraic group G acts on S_G via simultaneous conjugation:*

$$g \cdot (\Phi, N) = (g\Phi g^{-1}, \text{Ad}(g)N).$$

Assume in addition that l is geometrically G -banal.

- (2) The scheme $S_{G,\mathcal{O}}$ is flat and a local complete intersection of relative dimension $\dim G$ over \mathcal{O} .
- (3) Define the second projection map $p : S_G \rightarrow \mathcal{N}_G$ as before. If C is a G/L -conjugacy class inside $\mathcal{N}_{G,L} \subseteq \mathcal{N}_G$, then the closed subscheme $X_C := \overline{p^{-1}(C)} \subset S_G$ is a union of irreducible components and $S_G = \bigcup_C X_C$.
- (4) If in addition $G = \mathrm{GL}_n$, the X_C are irreducible components of $S_{n,\mathcal{O}} := S_{\mathrm{GL}_n,\mathcal{O}}$ and these can be naturally identified with partitions of n . For a partition p , we call the corresponding component X_p .
- (5) The scheme $S_{G,\mathcal{O}}$ is reduced.

Proof. (1) This is clear.

(2) This follows from Proposition 2.5(3) and Corollary 2.5 of [DHKM20].

(3) As $S_{G,\mathcal{O}}$ is flat over \mathcal{O} , the irreducible components of $S_{G,\mathcal{O}}$ are exactly those of the open subscheme $S_{G,L}$. This then follows from the proof of part 2, after noticing that $\mathcal{N}_{G,L} = \bigcup_C C_L$ as sets.

(4) Suppose $G = \mathrm{GL}_n$. Then C is a quotient of GL_n , and so is irreducible. Because centralisers inside GL_n are irreducible, the map $p^{-1}(C) \rightarrow C$ is flat with irreducible smooth fibres, and thus is smooth and open. By [Stacks, Lemma 004Z], it follows that $p^{-1}(C)$ is irreducible, and thus so is X_C . The final claim follows from the theory of Jordan normal forms.

(5) This follows from Proposition 2.8 of [DHKM20] and Proposition 2.5(3). □

2.1. Lemmas in commutative algebra and algebraic geometry. The remainder of this section proves some lemmas from algebraic geometry and commutative algebra that we will need later.

Lemma 2.7. *Let G be a smooth algebraic group over a scheme S , and let X be an S -scheme. Suppose that we have a morphism $m : G \times_S X \rightarrow X$ defining a group action of G on X . Then m is a smooth morphism.*

Proof. The morphism $p_X : G \times_S X \rightarrow X$ obtained by the base change of $G \rightarrow S$ is smooth. The automorphism ϕ of $G \times_S X$ given by $(g, x) \mapsto (g, g.x)$ is also smooth, since it is an isomorphism. As a composition of smooth morphisms, $m = p_X \circ \phi$ is smooth. □

Lemma 2.8. *Let $f : X \rightarrow Y$ be a smooth morphism of schemes. Let $p \in X$. Then Y is regular at $f(p)$ if and only if X is regular at p .*

Proof. After reducing the problem to local ring maps on stalks, this follows from Theorem 23.7 of [Mat86]. □

Lemma 2.9. *Assume that \mathcal{O} is complete. Let R be a local \mathcal{O} -algebra, and assume that it is topologically of finite type with respect to the \mathfrak{m}_R -adic topology. Let x and \bar{x} be prime ideals of R that give rise to the commutative diagram*

$$\begin{array}{ccc}
 R & \xrightarrow{x} & \mathcal{O} \hookrightarrow L = \mathcal{O}[1/I] \\
 & \searrow \bar{x} & \downarrow \\
 & & \mathbb{F}
 \end{array}$$

Then

$$R_{\bar{x}}^{\wedge}[1/l]_x^{\wedge} \cong R_x^{\wedge}.$$

Proof. Since $R \setminus x \supseteq R \setminus \bar{x} \cup \{1/l\}$, that $R_{\bar{x}}[1/l]_x \cong R_x$. As R is Noetherian, we have $\bigcap_n \bar{x}^n = 0$, and thus we have an injection $R_{\bar{x}} \hookrightarrow R_{\bar{x}}^{\wedge}$. This gives us a local homomorphism inclusion

$$R_x = R_{\bar{x}}[1/l]_x \hookrightarrow R_{\bar{x}}^{\wedge}[1/l]_x.$$

We notice that $R_x/x \cong L$, so that

$$[R_{\bar{x}}^{\wedge}[1/l]_x]/x \cong [\varprojlim_n (R/\bar{x}^n)/x][1/l] \cong \varprojlim_n (R/(x, l^n))[1/l] \cong (\varprojlim \mathcal{O}/l^n)[1/l] = L,$$

the last equality arising because \mathcal{O} is complete. Thus, by [Stacks, Lemma 0394], $R_{\bar{x}}^{\wedge}[1/l]_x^{\wedge}$ is the completion of $R_{\bar{x}}^{\wedge}[1/l]_x$ under the x -adic topology arising from R_x , and is a finite R_x^{\wedge} -module. It follows that the map

$$R_x^{\wedge} \rightarrow R_{\bar{x}}^{\wedge}[1/l]_x^{\wedge}$$

is an injection, and induces an isomorphism modulo x . If C is the cokernel (which we now know to be a finite R_x^{\wedge} -module) then we see $xC = C$, and so Nakayama’s lemma shows us that $C = 0$, implying that the map is an isomorphism. □

3. Smoothness results for X_C

In this section we prove Theorem 1.2. Let G be a connected reductive group over \mathcal{O} and $S_{G,\mathcal{O}}$ as before. As each map $X_C \rightarrow \text{Spec}(\mathcal{O})$ is flat, we can (and do) reduce the problem to the case $\mathcal{O} = \mathbb{F}$ is a field of characteristic 0 or l (see [Stacks, Lemma 01V8]). Since smoothness is an fpqc-local property, we can assume that \mathbb{F} is algebraically closed. We make these assumptions throughout this section.

3.1. Associated cocharacters. In what follows, we will require some setup, notation, and knowledge of Bala–Carter theory. Let G be a connected reductive group over an algebraically closed field \mathbb{F} with Lie algebra \mathfrak{g} and let $C \subseteq \mathcal{N}_G$ be a nilpotent orbit. In what follows, we restrict to the case where the derived subgroup of G is (almost) simple. When G is of adjoint type, we can do this because then $G = \prod_i G_i$ for G_i almost simple, adjoint, and $S_{G,\mathcal{O}} \cong \prod_i S_{G_i,\mathcal{O}}$. If G is not of adjoint type, then $S_{G,\mathcal{O}} \rightarrow S_{G^{ad},\mathcal{O}}$ is a $Z(G)$ -torsor, and since $Z(G)$ is smooth (under our considerateness condition), any smoothness result translates between the cases for G and G^{ad} .

Let \tilde{T} be a maximal torus of G and Π the set of roots. Let $e \in C \subseteq \mathfrak{g}$. Let $L \subseteq G$ be a Levi subgroup of G , minimal subject to $e \in \text{Lie}(L)$. Let Z_L be the centre of L . Following Definition 2.8 of [FR08] (see also Section 2.3 of [Pre03]), the nilpotent element e is called distinguished in L if Z_L° is a maximal torus of the centraliser $C_L(e)$ of e in L . By Proposition 2.11 of [FR08], there is an associated cocharacter $\lambda : \mathbb{G}_m \rightarrow \tilde{T}$ such that $\text{Ad}(\lambda(t)).e = t^2e$ and $\text{im}(\lambda) \subseteq [L, L]$.

The group \mathbb{G}_m acts on $\text{Lie}(G)$ through $\text{Ad} \circ \lambda : \mathbb{G}_m \rightarrow \text{Aut}(\mathfrak{g})$, which gives a decomposition

$$\mathfrak{g} = \bigoplus_{i \in \mathbb{Z}} \mathfrak{g}(\lambda, i).$$

Through Lemma 5.6.5 of [Car93] and the preceding discussion, we choose a base of simple roots $\Delta \subseteq \Pi$ such that $\langle \alpha, \lambda \rangle \geq 0$ for all $\alpha \in \Delta$. Call the corresponding Borel subgroup B .

We define a parabolic subgroup $P_\lambda \subseteq L$ such that $\text{Lie}(P_\lambda) = \bigoplus_{i \geq 0} \mathfrak{g}_L(\lambda, i)$, where each $\mathfrak{g}_L(\lambda, i) = \mathfrak{g}(\lambda, i) \cap \text{Lie}(L)$. We note that P_λ has a Levi decomposition $P_\lambda = M_\lambda U_\lambda$ with $\text{Lie}(U_\lambda) = \bigoplus_{i > 0} \mathfrak{g}_L(\lambda, i)$ and $\text{Lie}(M_\lambda) = \mathfrak{g}_L(\lambda, 0)$. We say P_λ is a *distinguished parabolic subgroup* in L if $\dim \mathfrak{g}_L(\lambda, 0) = \dim(\mathfrak{g}_L(\lambda, 2)) + \dim(Z_L)$, and by Proposition 2.5 of [Pre03] e is distinguished if and only if P_λ is distinguished.

The primary result of Bala–Carter theory is that there is a bijection between the adjoint orbits of \mathcal{N}_G and pairs (M, P) , where M is a Levi subgroup of G , and P is a distinguished parabolic subgroup of M .

3.2. The smoothness result for irreducible components corresponding to distinguished nilpotent orbits.

This section contains the main result of the paper, concerning the smoothness of X_C (Theorem 3.2).

Lemma 3.1. *Assume $e \in C$ is a distinguished nilpotent element with λ an associated cocharacter. Let $C_G(e)$ be the centraliser of e in G with Levi decomposition $C_G(e) = MR$, where $R = R_u(C_G(e))$ is unipotent and M is reductive. Suppose that $t \in \mathbb{G}_m$ is sufficiently generic so that $\mathfrak{g}^{\text{Ad}(\lambda(t))} = \mathfrak{g}(\lambda, 0)$. Then every element of $C_G(e)\lambda(t)$ is conjugate to an element of $M\lambda(t)$.*

Proof. Since e is distinguished, Theorem A of [Pre03] tells us that $M = M_\lambda \cap C_G(e)$ and $R = U_\lambda \cap C_G(e)$. Further, following Definition 2.8 of [FR08], the maximal torus of M is Z_G° so that M/Z_G° is a rank 0 reductive group; ergo finite. Thus, all unipotent elements of $C_G(e)$ lie in R . Let $g \in \lambda(t)C_G(e) \subseteq \tilde{T}C_G(e)$ have abstract Jordan decomposition $g = su$ with s semisimple and u unipotent. As s is semisimple, it lies in a maximal torus T' of $\tilde{T}C_G(e)$ and so there is some $x \in \tilde{T}C_G(e)$ such that $xsx^{-1} \in \tilde{T}$. In fact, we can assume without loss of generality that $x \in C_G(e)$, because \tilde{T} is abelian. Then, because $\lambda(t)$ normalises $C_G(e)$, we see that $xgx^{-1} \in \lambda(t)C_G(e)$. Since u is unipotent in $\tilde{T}C_G(e)$ we obtain $u \in C_G(e)$, and thus we get

$$xsx^{-1} = xgx^{-1}(xux^{-1})^{-1} \in \lambda(t)C_G(e) \cap \tilde{T}.$$

Hence

$$xsx^{-1} \in \tilde{T} \cap \lambda(t)C_G(e) = \tilde{T} \cap \tilde{T}M \cap \lambda(t)C_G(e) = \lambda(t)\tilde{T} \cap \lambda(t)M = \lambda(t)[\tilde{T} \cap M].$$

Then $xux^{-1} \in C_G(xsx^{-1}) = C_G(\lambda(t)s')$ for some $s' \in \tilde{T} \cap M$. By the genericity condition on t , we see that $C_G(\lambda(t)) = M_\lambda$ and so, by Theorem 3.5.3 of [Car93], $C_G(\lambda(t)s') \subseteq C_G(\lambda(t)) = M_\lambda$. Hence $xux^{-1} \in M$ and $xux^{-1} = 1$. The result follows. \square

Theorem 3.2. *Let G/\mathcal{O} be a connected reductive group with centre Z , let \mathfrak{g} be the Lie algebra of G , and suppose $q \in \mathcal{O}$ is considerate towards G over \mathcal{O} . Suppose $C \subseteq \mathcal{N}_{G,L}$ is either 0 or a distinguished nilpotent adjoint orbit. Then X_C is smooth over \mathcal{O} and there is a bijection between the connected components of X_C and the set of Φ_0 -twisted conjugacy classes of the group $\pi_0(C_G(e))$.*

Proof. Consider first the case $C = 0$. Then $X_C = \{(\Phi, 0) \in S_{G,\mathcal{O}}\} \cong G$ via the map projecting to the Φ coordinate. Since G is smooth, this proves the theorem.

Keep the notation of before, with e a distinguished nilpotent element of \mathfrak{g} with associated cocharacter $\lambda : \mathbb{G}_m \rightarrow T$. After making some choice of $\sqrt{q} \in \mathbb{F}$, set $\Phi_0 = \lambda(\sqrt{q})$. The centraliser $C_G(e)$ exhibits a Levi decomposition $C_G(e) = MR$ with M reductive and R the unipotent radical. Because e is a distinguished nilpotent element in \mathfrak{g} , we see that Z° , the connected component of the identity of the centre Z of G , is a maximal torus of M by Proposition 2.11(iii) of [FR08]. Since M/Z° is a split reductive group of rank 0, it is finite; because unipotent radicals are connected, it is isomorphic to the component group $A(e)$ of the centraliser $C_G(e)$. We choose a set of representatives S such that $M = \coprod_{s \in S} sZ^\circ$.

Define $Y = M\Phi \times \mathfrak{g}(\lambda, 2)$. Since the characteristic l exceeds h , we see that M is a smooth subgroup of G making Y a smooth \mathbb{F} -scheme. It is clear that Y is a closed subscheme of S_G , because if $(m\Phi_0, N) \in M\Phi_0 \times \mathfrak{g}(\lambda, 2)$ then

$$\text{Ad}(m\Phi_0).N = \text{Ad}(m)\text{Ad}(\Phi_0).N = \text{Ad}(m).qN = qN.$$

In fact, Y lies inside the irreducible component X_C . The distinguished element e lies in the unique open dense P_λ -orbit inside $\mathfrak{g}(2, \lambda)$ (See, for example, Proposition 5.8.7b of [Car93]) and thus $P_\lambda.[M\Phi_0 \times \{e\}]$ is a dense open subscheme of Y . As $M\Phi_0 \times \{e\} \subseteq p^{-1}(C)$, we obtain $P_\lambda.[M\Phi_0 \times \{e\}] \subseteq p^{-1}(C)$ and consequently $Y \subseteq X_C$. Define the morphism

$$f : G \times Y \rightarrow X_C, \quad (g, (\Phi, N)) \mapsto (g\Phi g^{-1}, \text{Ad}(g)N).$$

As $G \times Y$ is a smooth variety, Lemma 2.8 implies the theorem provided we can show that f is a smooth surjective morphism.

Surjectivity is equivalent to the statement that every pair $(\Phi, N) \in X_C$ is conjugate to a pair in Y . To prove this, it suffices to show that Φ is conjugate to an element of $M\Phi_0$ whenever $(\Phi, N) \in X_C$. As there is some $\text{Ad}(g).e \in \mathfrak{g}$ upon which Φ acts as multiplication by q , we see that Φ is conjugate to some element of $C_G(e)\Phi_0$. By Lemma 3.1, then, Φ is conjugate to an element of $M\Phi_0$, proving surjectivity.

We now proceed to prove that f is smooth. Consider the commutative diagram

$$\begin{array}{ccc} G \times Y & \xrightarrow{f} & X_C \\ \downarrow & & \downarrow \\ G \times M\Phi_0 & \longrightarrow & G \end{array}$$

where the vertical maps come from the “forget N ” projections $(g, m\Phi_0, N) \in G \times Y \mapsto (g, m\Phi_0) \in G \times M\Phi_0$ and $(\Phi, N) \in X_C \mapsto \Phi \in G$ and the horizontal maps are defined via the conjugation action of $g \in G$ on Y , so that the diagram commutes. Choosing a set of representatives S of M/Z° , the map $G \times M\Phi_0 \rightarrow G$ factors through

$$\begin{array}{ccc} G \times M\Phi_0 & \longrightarrow & G \\ \parallel & & \uparrow \\ \coprod_{s \in S} G \times sZ^\circ\Phi_0 & \xrightarrow{m} & \coprod_s Z^\circ G_{s\Phi_0} \end{array}$$

where $G_{s\Phi_0}$ denotes the conjugacy class of $s\Phi_0$ in G . Each $Z^\circ G_{s\Phi_0}$ defines a locally closed subvariety

of G . If any two subschemes $Z^\circ G_{s\Phi_0}$ and $Z^\circ G_{t\Phi_0}$ intersect in G , say $x \in Z^\circ G_{s\Phi_0} \cap Z^\circ G_{t\Phi_0}$, then

$$z_1 g_1 s \Phi_0 g_1^{-1} = x = z_2 g_2 t \Phi_0 g_2^{-1}$$

implies that $t\Phi_0 = (z_1 z_2^{-1})(g_2^{-1} g_1) s \Phi_0 (g_2^{-1} g_1)^{-1}$; and thus leads us to $Z^\circ G_{s\Phi_0} = Z^\circ G_{t\Phi_0}$ as locally closed subschemes of G . Thus, by possibly restricting to a subset S' of the set of representatives S if necessary, we can view $\coprod_{s \in S'} Z^\circ G_{s\Phi_0}$ as a locally closed subscheme of G .

Because the map $f : G \times Y \rightarrow X_C$ is surjective, the map $X_C \rightarrow G$ also factors through $\coprod_{s \in S'} Z^\circ G_{s\Phi_0}$, giving us the commutative diagram

$$\begin{array}{ccc} G \times Y & \xrightarrow{f} & X_C \\ \downarrow & & \downarrow \\ G \times M\Phi_0 & \xrightarrow{m} & \coprod_{s \in S'} Z^\circ G_{s\Phi_0} \end{array} \tag{1}$$

We claim that this is a pullback square. Since e is distinguished, we see $\mathfrak{g}(\lambda, 1) = 0$ and thus each simple root α has its corresponding character eigenspace \mathfrak{g}_α either contained inside $\mathfrak{g}(\lambda, 0)$ or $\mathfrak{g}(\lambda, 2)$. Hence, as every positive root is the sum of at most $h - 1$ simple roots (where $h = h_G$ is the Coxeter number of G), we see that

$$\{i \in \mathbb{Z} : \mathfrak{g}(\lambda, i) \neq 0\} \subseteq 2\mathbb{Z} \cap [-2h + 2, 2h - 2].$$

Given that q is considerate towards G/\mathbb{F} , it follows that the subspace of \mathfrak{g} upon which $\Phi_0 = \lambda(\sqrt{q})$ acts as multiplication by q is precisely $\mathfrak{g}(\lambda, 2)$.

Thus, if $(g, \Phi) \in G \times M\Phi_0$ and $(\Phi', N') \in X_C$ are such that $g\Phi g^{-1} = \Phi'$, then $\text{Ad}(\Phi)(\text{Ad}(g^{-1})N') = q\text{Ad}(g^{-1})N'$ and $\text{Ad}(g^{-1})N' \in \mathfrak{g}(\lambda, 2)$ by the previous discussion. So the morphism

$$((g, \Phi), (\Phi', N')) \rightarrow (g, (\Phi, \text{Ad}(g^{-1})N'))$$

gives an inverse to the natural morphism $G \times Y \rightarrow (G \times M\Phi_0) \times_{\coprod_{s \in S'} Z^\circ G_{s\Phi_0}} X_C$. This shows that the commutative diagram (1) is a pullback square.

By the theory of homogeneous spaces, the bottom map m is flat with fibres isomorphic to $\text{Stab}_G(\Phi_0)$, which are smooth group schemes. This shows that m is smooth. Hence, since f is the base change of m to X_C , by Proposition 10.1 of [Har77] we see that f is smooth. We conclude X_C is smooth over \mathbb{F} .

The statement on the number of connected components is Theorem 2.5 of [Sho24], and is included for completeness. □

Remark. A question arises regarding the generality and necessity of the considerateness condition: When exactly is q -considerateness a necessary condition for smoothness? As one can see in the proof, we used considerateness to prove that $G \times Y$ was the pullback of the diagram

$$\begin{array}{ccc} G \times Y & \longrightarrow & X_C \\ \downarrow & & \downarrow \\ G \times M\Phi_0 & \longrightarrow & M.G_{\Phi_0} \end{array}$$

arising from the fact that $\{N \in \mathfrak{g} : \text{Ad}(\Phi_0)N = qN\} = \mathfrak{g}(\lambda, 2)$. When C is the *regular* nilpotent orbit, $\mathfrak{g}(\lambda, i) \neq 0$ for every $i \in [-2h + 2, 2h - 2] \cap 2\mathbb{Z}$, so we see that q -considerateness is precisely the condition that $\{N \in \mathfrak{g} : \text{Ad}(\Phi_0)N = qN\} = \mathfrak{g}(\lambda, 2)$. When C is distinguished and nonregular, we have $\mathfrak{g}(\lambda, 2h - 2) = 0$. Hence, there is some $r < h$, depending on the distinguished orbit C , such that if $q \in \mathcal{O}$ has the property that $1, q, \dots, q^r$ are distinct, then X_C is smooth via the above proof. This r can always be taken to be

$$r = 1 + \max\{i : \mathfrak{g}(\lambda, 2i) \neq 0\}.$$

Remark. Let X_C^s be the image of f restricted to $G \times {}^sZ^\circ\Phi_0$ (so that X_C^s is an irreducible component of X_C). Another way of interpreting the Cartesian diagram (1) is that $X_C^s \rightarrow Z^\circ G_{s\Phi_0}$ is the total space of the vector bundle on $Z^\circ G_{s\Phi_0}$ whose fibre above Φ is $\ker(\text{Ad}(\Phi) - q) \subseteq \mathfrak{g}$. In particular, we obtain:

Corollary 3.3. *If C is a distinguished orbit and s, Φ_0 are as in the proof of Theorem 3.2, then X_C^s is described as the closed subscheme of $G \times \mathfrak{g}$ cut out by the equations*

- $\text{Ad}(\Phi)N = qN$ and
- any set of equations describing the closed orbit $Z^\circ G_{s\Phi_0} \subseteq G$.

3.3. The converse nonsmoothness result. We now begin work towards the converse of Theorem 3.2. Consider the situation described at the beginning of the chapter with reductive group G over an algebraically closed field \mathbb{F} with (almost) simple derived subgroup with maximal torus T , set of roots Π , a set of simple roots $\Delta \subseteq \Pi$, and a nilpotent element e with associated cocharacter λ . Let L be the smallest Levi subgroup of G with $e \in \text{Lie}(L)$, so that e is distinguished for L . Let $\Delta_L \subseteq \Delta$ be the simple roots of L .

Theorem 3.4. *Let G be as before. Suppose $C \subseteq \mathcal{N}_G$ is a nilpotent adjoint orbit, distinguished in a proper and nontrivial Levi subgroup $T \neq L \subsetneq G$. Then $X_C \subseteq S_G$ is singular.*

Before proving this theorem, we need some terminology and a lemma. Let $D = (\Delta, \Sigma)$ be the Dynkin diagram of G , and $D_L = (\Delta_L, \Sigma_L) \subseteq D$ the maximal subdiagram containing exactly the vertices Δ_L . (Note that D is connected when its derived subgroup is (almost) simple but D_L may not be connected). Recall that, given a distinguished nilpotent e , one can attach to each vertex $\beta \in \Delta_L$ the number $\langle \alpha, \lambda \rangle \in \{0, 2\}$, and this is called the weighted Dynkin diagram $D_L(e)$.

We call a root $\alpha \in \Delta_L$ *exposed* if there is an edge in D connecting α to a root $\beta \in \Delta \setminus \Delta_L$.

Lemma 3.5. *Any exposed root $\alpha \in \Delta_L$ has $\langle \alpha, \lambda \rangle = 2$.*

Proof. In the case of types A, B, C and D , either all simple factors of the Levi subgroup are of type A , or exactly one of the almost simple factors of L is of type B, C, D , respectively. If all the simple factors of the Levi subgroup are type A , then all roots $\alpha \in \Delta_L$ have $\langle \alpha, \lambda \rangle = 2$ because all distinguished nilpotents orbits are regular in type A . In the case with one factor of type B, C or D , the only “exposed” root of this factor is on the end of the string, and one can see from the tables on pages 174 and 175 of [Car93]




Figure 1. The distinguished weighted Dynkin diagrams of Levi subgroups of F_4 .

that, independent of the choice of distinguished nilpotent, this exposed root α always has $\langle \alpha, \lambda \rangle = 2$. This proves the lemma in types A , B , C and D .

In type G_2 , the only proper Levi subgroups are of type A_1 , so all roots $\alpha \in \Delta_M$ have $\langle \alpha, \lambda \rangle = 2$.

In type F_4 , there are three possibilities for a Levi factor not of type A , these being C_2 , C_3 and B_3 . The distinguished orbits of these Levi subgroups are described on pages 174 and 175 of [Car93] and are listed in Figure 1. From this we directly see that all exposed roots have $\langle \alpha, \lambda \rangle = 2$.

In type E_l , there are three possibilities for the Levi subgroup types. Either there are only factors of type A for which the result holds, or there is a unique factor of type D_n and $n \leq 7$, or there is a factor of type E_6 or E_7 . In the case of a factor of type E_6 or E_7 , the weighted Dynkin diagrams of distinguished parabolic subgroups (of which there are 3 of type E_6 and 6 of type E_7) are listed on page 176 of [Car93], from which it is clear that all exposed roots α have the desired property. See also the figures on page 1157.

This leaves only the case that L has a factor of type D_n . All distinguished orbits of D_n with $n \leq 7$ are listed on 1156, from whose table we see that all exposed roots have the desired property. \square

Remark. In the description of nonregular distinguished parabolic subgroups in type B_l on page 175 of [Car93], one requires $k \geq 2$ (in the notation of the source) for the conditions to make sense, though this isn't explicit. In our application, the important fact is that for B_l with $l = 3$, the only distinguished orbit is the regular orbit. This can also be seen from the description of distinguished orbits in Theorem 8.2.14 of [CM93] via partitions of $2l + 1 = 7$ into distinct odd parts.

Proof of Theorem 3.4. Consider a point $P = (\Phi_0, 0) \in X_C$ with $\Phi_0 \in T$. Define four subvarieties of S_G that contain P as follows:

- (1) Let $\mathcal{O} = G.P$ be the G -orbit of P .
- (2) Let \tilde{T} be the maximal torus of G seen as a closed subvariety of S_G via the inclusion $\Phi \mapsto (\Phi, 0)$.
- (3) Let $\mathcal{N}_0 = \{N \in \mathfrak{g} : \text{Ad}(\Phi_0)N = qN\}$ viewed as a closed subvariety of S_G via the inclusion $N \mapsto (\Phi_0, N)$.
- (4) Let $U_0 = U^- \cap \text{Stab}_G(P)$ where U^- is the opposite unipotent subgroup to $[B, B]$, viewed as a closed subvariety of S_G via the inclusion $u \mapsto (\Phi_0 u, 0)$.

Claim 1. *The tangent spaces $T_P \mathcal{O}$, $T_P \tilde{T}$, $T_P \mathcal{N}_0$ and $T_P U_0$ form a direct sum inside $T_P S_G$.*

Observe that $T_P(S_G) \subseteq T_P(G \times \mathfrak{g}) = \mathfrak{g} \times \mathfrak{g}$. We briefly describe $T_P\mathcal{O}$ as a subspace of $\mathfrak{g} \times \{0\}$ (which we will conflate with \mathfrak{g} as this shouldn't cause confusion).

Consider the map $f : G \rightarrow G : g \mapsto g\Phi_0g^{-1}\Phi_0^{-1}$, comprised of the conjugation action of g on Φ_0 followed by right multiplication by Φ_0^{-1} (to ensure the identity is sent to the identity). Then \mathcal{O} is isomorphic to the set-theoretic image of f , which is a locally closed subscheme of G . The derivative of this map is $\text{id} - \text{Ad}(\Phi_0) : \mathfrak{g} \rightarrow \mathfrak{g}$, and it factors through $\mathfrak{g} \rightarrow T_PC \hookrightarrow \mathfrak{g}$. We hence see a natural identification of $T_P\mathcal{O}$ with $\text{im}(\text{id} - \text{Ad}(\Phi_0))$.

We now proceed to prove the claim. Firstly, $T_P\tilde{T} \cap T_P U_0 = \{0\}$ because $\text{Lie}(\tilde{T}) \cap \text{Lie}(U^-) = \{0\}$. Next, consider $T_P\mathcal{O} \cap (T_P\tilde{T} \oplus T_P U_0)$. Note that $U_0, \tilde{T} \subseteq \text{Stab}_G(\Phi) \subseteq G$ and $T_P\text{Stab}_G(\Phi_0) = \ker(\text{id} - \text{Ad}(\Phi_0))$. Then because Φ_0 is semisimple, the intersection of $\text{im}(\text{id} - \text{Ad}(\Phi_0))$ and $\ker(\text{id} - \text{Ad}(\Phi_0))$ is trivial (this is easily checked for GL_n , and extended to all G because G can always be embedded into some GL_N). Hence $T_P(\text{Stab}_G(\Phi_0)) \cap T_P\mathcal{O} = 0$, and thus $T_P\mathcal{O} \cap (T_P\tilde{T} \oplus T_P U_0) = 0$.

To show that $T_P\mathcal{N}_0$ intersects $T_P\mathcal{O} + T_P\tilde{T} + T_P U_0$ at the origin, it suffices to notice that an element of $T_PC + T_P\tilde{T} + T_P U_0$ takes the form $P' = (\Phi', 0)$, while an element $P' \in T_P\mathcal{N}_0$ takes the form $P' = (\Phi_0, N) \in S_G(\mathbb{F}[\epsilon]/\epsilon^2)$. For these to be equal, we must have $\Phi' = \Phi_0$ and $N = 0$, so $P' = P$. This proves the claim.

Suppose that $C \subseteq \mathcal{N}_G$ is an adjoint orbit, neither zero nor a distinguished nilpotent orbit as in the hypothesis. Define $e \in C$, a choice of maximal torus and Borel $\tilde{T} \subset B$, the associated cocharacter λ and minimal Levi $L \subset G$, all as in the general setup. Set $D = \tilde{T} \cap X_C$ and $\mathcal{N}_1 = \mathcal{N}_0 \cap X_C$, and note that $U_0, \mathcal{O} \subseteq X_C$ already. Our aim is to show that there is a point $P \in X_C$ such that

$$\dim_{\mathbb{F}}(T_P\mathcal{O}) + \dim_{\mathbb{F}}(T_P D) + \dim_{\mathbb{F}}(T_P\mathcal{N}_1) + \dim_{\mathbb{F}}(T_P U_0) > \dim(X_C) = \dim(G).$$

It is clear that whenever $z \in Z_L$ and $\mu \in \mathbb{F}$, the point $(z\lambda(\sqrt{q}), \mu e)$ lies in X_C . We choose the point $P = (\Phi_0, 0) \in X_C$ with $\Phi_0 = z\lambda(\sqrt{q})$ for some $z \in Z_L$, which we will determine momentarily.

Regardless of the choice of z for now, recall the decomposition

$$\text{Lie}(L) = \bigoplus_{i \in \mathbb{Z}} \mathfrak{g}_L(\lambda, i).$$

As $\mathfrak{g}_L(\lambda, 0)$ is a Levi subalgebra of $\text{Lie}(L)$, there is a Levi subgroup $M_0 \subset L$ with $\text{Lie}(M_0) = \mathfrak{g}_L(\lambda, 0)$. It is clear that $M_0 \subseteq \text{Stab}_G(\Phi_0)$. Observe also that $\mathfrak{g}_L(\lambda, 2) \subseteq \mathcal{N}_1$ and $Z_L\Phi_0 \subseteq D$. We define positive integers

- (1) $\epsilon_0 := \dim(\text{Stab}_G(\Phi_0)) - \dim_{\mathbb{F}}(\mathfrak{g}_L(\lambda, 0)) = \dim(G) - \dim(\mathcal{O}) - \dim_{\mathbb{F}}(\mathfrak{g}_L(\lambda, 0))$,
- (2) $\epsilon_1 := \dim_{\mathbb{F}}(T_P D) - \dim(Z_L)$,
- (3) $\epsilon_2 := \dim_{\mathbb{F}}(T_P\mathcal{N}_1) - \dim_{\mathbb{F}}(\mathfrak{g}_L(\lambda, 2))$,
- (4) $\epsilon_3 := \dim_{\mathbb{F}}(T_P U_0)$.

Putting this together, and using the fact that

$$\dim(\mathfrak{g}_L(\lambda, 0)) = \dim(\mathfrak{g}_L(\lambda, 2)) + \dim(Z_L)$$




Figure 2. An example of α in the case where G is of type E_8 and L of type $D_4 \times A_2$.

because e is distinguished inside L (this follows from a generalisation of Lemma 8.2.1 in [CM93] to reductive groups in good characteristic), we see that

$$\begin{aligned} \dim(T_P X_C) &\geq \dim(T_P \mathcal{O}) + \dim(T_P D) + \dim(T_P \mathcal{N}_1) + \dim(T_P U_0) \\ &\geq [\dim(G) - \dim(\mathfrak{g}_L(\lambda, 0)) - \epsilon_0] + [\dim(Z_L) + \epsilon_1] + [\dim(\mathfrak{g}_L(\lambda, 2)) + \epsilon_2] + \epsilon_3 \\ &= \dim(G) + \epsilon_1 + \epsilon_2 + \epsilon_3 - \epsilon_0. \end{aligned}$$

Thus, it is enough to find some choice of $z \in Z_L$ such that $\epsilon_1 + \epsilon_2 + \epsilon_3 - \epsilon_0 > 0$.

Fix some root $\alpha \in \Delta \setminus \Delta_L$ adjacent to a root in Δ_L in the Dynkin diagram. See Figure 2 for an example. There is a morphism of algebraic groups

$$B = \prod_{\beta \in \Delta} \beta : \tilde{T} \rightarrow \prod_{\beta \in \Delta} \mathbb{G}_m$$

which is surjective and has kernel $Z \subset \tilde{T}$. Because B is surjective, we can choose $z \in T$ such that $\alpha(z) = 1$ and $\beta(z) = 1$ whenever $\beta \in \Delta_L$ (so that $z \in Z_L$) and $\beta(z) = q \neq 1$ in all other cases. Consider $\Phi_0 = z\lambda(\sqrt{q})$. Whenever $\gamma \in \Pi_G^+$ is a positive root of G , it decomposes as a product of simple roots $\gamma = \prod_{\beta \in \Delta} \beta^{c_\beta} \in X(T)$ with $\sum_\beta c_\beta < h$. By design, each $\beta(\Phi_0)$ is either 1 or q , so $\gamma(\Phi_0) \in \{1, q, q^2, \dots, q^{h-1}\}$, and $\gamma(\Phi_0) = 1$ if and only if all the simple roots with $c_\beta \neq 0$ satisfy $\beta(\Phi_0) = 1$.

Claim 2. *If $\gamma \in \Pi_G^+ \setminus \Pi_L^+$ has $\gamma(\Phi_0) = 1$, then $\gamma = \alpha$.*

If γ is simple, then $\gamma = \alpha$ by our choice of Φ_0 . Suppose for contradiction that γ is not simple. Then it contains at least two simple roots in its decomposition, and one of these must be α , as otherwise all simple roots are in Δ_L and $\gamma \in \Pi_L^+$. There must also be another root $\beta \in \Delta_L$ with $c_\beta \neq 0$, and for γ to be a root, there must be a path from α to β (in the Dynkin diagram) passing through vertexes β' with $c_{\beta'} \neq 0$, and each of these as such (since $\gamma(\Phi_0) = 1$) has $\beta'(\Phi_0) = 1$. But as $\alpha \in \Delta \setminus \Delta_L$ and $\beta \in \Delta_L$, at least one of the β' is an exposed root, and thus $\beta'(\Phi_0) = q$ by Lemma 3.5. This is a contradiction. We conclude that $\gamma(\Phi_0) = 1$ implies either $\gamma = \alpha$ or $\gamma \in \Pi_L^+$. This proves the claim.

When $\beta \in \Pi$, denote the root subgroup of β by $U_\beta \leq G$. By Theorem 3.5.3 of [Car93], the (connected) centraliser of Φ_0 is

$$C_G(\Phi_0)^\circ = \langle \tilde{T}, U_\beta, U_{-\beta} : \beta(\Phi_0) = 1 \rangle.$$

The subgroup generated by \tilde{T} and all U_β with $\beta(\Phi_0) = 1$ and $\beta \in \Pi_L$ is simply M_0° , so we see that

$$C_G(\Phi_0)^\circ = \langle M_0^\circ, U_\alpha, U_{-\alpha} \rangle$$

and hence $\dim C_G(\Phi_0) = \dim(M_0) + 2$ (or, in other words, $\epsilon_0 = 2$).

The reflection $s_\alpha \in N(\tilde{T})/\tilde{T}$ acts on \tilde{T} and stabilises Φ_0 ; thus it acts on $T_P X_C$. Further, this action preserves the subspaces $T_P D$ and $T_P \mathcal{N}_1$. However, since α is adjacent to a simple root of L , the reflection s_α does not preserve the Levi subgroup L , and hence preserves neither Z_L nor $\mathfrak{g}_L(\lambda, i)$, and we see that $s_\alpha(Z_L) \neq Z_L$ and $s_\alpha(\mathfrak{g}_L(\lambda, 2)) \neq \mathfrak{g}_L(\lambda, 2)$, so that $T_P D \supseteq s_\alpha(T_P Z_L) \cup T_P Z_L$ and $T_P \mathcal{N}_1 \supseteq s_\alpha(T_P \mathfrak{g}_L(\lambda, 2)) \cup T_P \mathfrak{g}_L(\lambda, 2)$, forcing $\epsilon_1 > 0$ and $\epsilon_2 > 0$ respectively.

For ϵ_3 , consider any choice of isomorphism $u_{-\alpha} : \mathbb{G}_a \xrightarrow{\sim} U_{-\alpha}$ and the adjoint action of $u_{-\alpha}(a) \in U_{-\alpha}$ on $e = \sum_\beta e_\beta \in \text{Lie}(L)$. Because α is a simple root in $\Delta_G \setminus \Delta_L$, we see that

$$[e_{-\alpha}, \sum_\beta e_\beta] = \sum_\beta [e_{-\alpha}, e_\beta] = 0$$

and thus that $\text{Ad}(u_{-\alpha}(a))e = e$. Hence,

$$\{(\Phi_0 u_{-\alpha}(a), \mu e) : a \in \mathbb{G}_a, \mu \in \mathbb{G}_m\}$$

is a locally open subscheme of $p^{-1}(C)$, from which we see that $(\Phi_0 u_{-\alpha}(\epsilon), 0)$ is a deformation in $T_P U_0$, forcing $\epsilon_3 > 0$.

We then obtain the inequality $\epsilon_1 + \epsilon_2 + \epsilon_3 - \epsilon_0 \geq 3 - 2 = 1$, proving that $(\Phi_0, 0)$ is a singular point of X_C . □

Example 1. Consider the group

$$G = \text{GSp}_4(R) = \{M \in \text{GL}_4(R) : M\Omega M^{-1} = \lambda\Omega \text{ for some } \lambda \in \mathbb{G}_m(R)\}, \quad \Omega = \begin{pmatrix} & & & 1 \\ & & 1 & \\ & -1 & & \\ -1 & & & \end{pmatrix},$$

where Ω is chosen so that a Borel subgroup can be given by the intersection of GSp_4 with the upper triangular matrices in GL_4 . We let $L = \text{GL}_2 \subseteq G$ be the Levi subgroup corresponding to the short root. Then

$$e = \begin{pmatrix} 0 & 1 & & \\ & 0 & 0 & \\ & & 0 & -1 \\ & & & 0 \end{pmatrix}$$

is distinguished in L and the associated cocharacter is $\lambda(t) = \text{Diag}(t, t^{-1}, t, t^{-1})$. We choose $\Phi_0 = \text{Diag}(q, 1, 1, q^{-1})$ and α to be the root corresponding to the one-parameter subgroup defined as

$$U_\alpha = \left\{ \begin{pmatrix} 1 & & & \\ & 1 & * & \\ & & 1 & \\ & & & 1 \end{pmatrix} \right\}.$$

Explicitly, we now see that $\text{Stab}_G(P) = \mathbb{G}_m \times \text{GL}_2$. We can also describe the subvarieties:

$$Z_L = \left\{ \left(\begin{pmatrix} ab & & & \\ & ab & & \\ & & ab^{-1} & \\ & & & ab^{-1} \end{pmatrix}, 0 \right) : a, b \in \mathbb{G}_m \right\},$$

$$\begin{aligned}
 s_\alpha(Z_L) &= \left\{ \left(\begin{pmatrix} ab & & & \\ & ab^{-1} & & \\ & & ab & \\ & & & ab^{-1} \end{pmatrix}, 0 \right) : a, b \in \mathbb{G}_m \right\}, \\
 \mathfrak{g}_L(\lambda, 2) &= \left\{ \left(\Phi_0, \begin{pmatrix} 0 & a & & \\ & 0 & & \\ & & 0 & -a \\ & & & 0 \end{pmatrix} \right) : a \in \mathbb{G}_a \right\}, \\
 s_\alpha(\mathfrak{g}_L(\lambda, 2)) &= \left\{ \left(\Phi_0, \begin{pmatrix} 0 & a & & \\ & 0 & & \\ & & 0 & a \\ & & & 0 \end{pmatrix} \right) : a \in \mathbb{G}_a \right\}, \\
 U_0 &= \left\{ \left(\begin{pmatrix} 1 & & & \\ & a & & \\ & & 1 & \\ & & & a^{-1} \end{pmatrix}, 0 \right) : a \in \mathbb{G}_a \right\}.
 \end{aligned}$$

When we put all this together, we see the contribution from D, \mathcal{N}_1 and U_0 is 6-dimensional, and thus

$$\dim T_P X_C \geq \dim(\mathrm{GSp}_4) - \dim(\mathrm{Stab}_G(P)) + 6 = \dim(\mathrm{GSp}_4) + 1.$$

We can piece Theorems 3.2 and 3.4 together:

Corollary 3.6. *Let G be a connected reductive group over a field \mathbb{F} of characteristic 0 or l , and suppose $G^{\mathrm{ad}} = \prod_i G_i$ where each G_i has a (almost) simple derived subgroup. Suppose q is considerate towards each $G_{i, \mathbb{F}}$. Then the smooth irreducible components of $S_{G^{\mathrm{ad}}}$ are precisely those of the form $\prod_i X_i$ where each $X_i \subseteq S_{G_i}$ is a smooth irreducible component of S_{G_i} . That is, each X_i corresponds to a distinguished nilpotent orbit of G_i or the zero orbit. The smooth components of S_G are precisely preimages the smooth components of $S_{G^{\mathrm{ad}}}$ under the obvious map $S_G \rightarrow S_{G^{\mathrm{ad}}}$.*

3.4. Distinguished orbits in type D and E. For the convenience of the reader in understanding Lemma 3.5, we give in the Appendix a list of weighted Dynkin diagrams for all distinguished orbits in types D_n and E_n with $n \leq 7$.

4. Automorphic forms for unitary groups

We now turn to an application of the smoothness result found in Section 3. In this section, we define the space of ordinary automorphic forms and the Hecke algebra attached to it. We then state a freeness result, which we will prove in Section 5.4 (Theorem 5.13).

Let l be a prime. Suppose F^+ is a totally real number field with an imaginary quadratic extension F such that all primes v of F^+ above l split in F . Let S_l be the set of all primes of F^+ that lie above l . Let L be a finite extension of \mathbb{Q}_l with ring of integers \mathcal{O} and residue field \mathbb{F} . Let \bar{L} be a choice of algebraic closure. We will assume that L is large enough that all embeddings $F \hookrightarrow \bar{L}$ lie inside L . Let $c \in \mathrm{Gal}(F/F^+)$ be the unique nontrivial element, given by complex conjugation. For $a \in F$, we will denote $c(a)$ by \bar{a} when convenient.

4.1. Unitary groups. Consider a central simple algebra D/F of F -dimension n^2 , and let S_D be a finite set of primes of F^+ that split in F . Suppose that

- (1) D splits at all places w of F that do not lie above any place in S_D ;

- (2) there is an isomorphism $D^{\text{op}} \cong D \otimes_{F,c} F$ of F -algebras;
- (3) the intersection $S_D \cap S_l$ is empty;
- (4) D_w is a division algebra at all places w of F above a place in S_D ;
- (5) either n is odd, or n is even and $\frac{1}{2}n[F^+ : \mathbb{Q}] + \#S_D \equiv 0 \pmod{2}$.

Because of condition (1) (which ensures that all places in S_D split), together with (2) and (5), we can find an involution of the second kind on D by [CHT08, Section 3.3, p. 95]. That is, we may construct a map

$$* : D \rightarrow D$$

such that

- $*$ is an F^+ linear anti-automorphism of D ,
- $(a^*)^* = a$ for all $a \in D$, and
- the involution $*$ coincides with complex conjugation when restricted to F .

In addition, we assume that this involution of the second kind is positive. That is, for any $\gamma \in D \setminus \{0\}$,

$$\text{tr}_{F:\mathbb{Q}}[\text{tr}_{D/F}(\gamma\gamma^*)] > 0.$$

Such an involution gives rise to a positive Hermitian form $\langle , \rangle : D \times D \rightarrow D$ given by $\langle x, y \rangle = x^*y$.

Let \mathcal{O}_D be an order in D such that $\mathcal{O}_D^* = \mathcal{O}_D$ and such that $\mathcal{O}_{D,v}$ is a maximal order of D_v for any split prime v of F^+ , as in Section 3.3 of [CHT08]. Define the unitary group over \mathcal{O}_{F^+} whose R -points (where R is an \mathcal{O}_{F^+} -algebra) are given by $G_D(R) = \{g \in (\mathcal{O}_D \otimes_{\mathcal{O}_{F^+}} R)^\times : g^*g = 1\}$. Then G_D is an algebraic group over \mathcal{O}_{F^+} . By the positivity condition, we have $G_{D,v} \cong U(n)$ at each infinite place v of F^+ .

For each prime v of F^+ that splits in F , choose a prime \tilde{v} of F lying above v . This choice allows us to give an isomorphism $i_{\tilde{v}} : G_D(F_v^+) \rightarrow (D \otimes_F F_{\tilde{v}})^\times$ which restricts to an isomorphism $G_D(\mathcal{O}_{F^+,v}) \cong (\mathcal{O}_{D,\tilde{v}})^\times$ as in Section 3.3 of [CHT08]. Note that when $v \notin S_D$ is split in F , then G_D is split, so that $G_D(F_v^+) \cong (D \otimes_{F^+} F_v^+)^\times \cong (D \otimes_F F_{\tilde{v}})^\times \cong \text{GL}_n(F_{\tilde{v}})$. If T is a set of primes of F^+ that splits in F , set $\tilde{T} = \{\tilde{v} : v \in T\}$.

4.2. Automorphic forms of G_D . We define the automorphic forms for G_D as in [Gro99] and [CHT08]. To do this, we recall some facts from the representation theory of reductive groups.

Let G be a split reductive group defined over L , and let $T \subseteq B \subseteq G$ be a choice of split maximal torus and Borel subgroup of G . Recall that finite-dimensional simple modules of G are uniquely determined by their highest weight in the character group of the torus $X^\bullet(T) := \text{Hom}(T, \mathbb{G}_m)$, and that such a representation exists if and only if this highest weight ν lies in a dominant Weyl chamber.

In the case of GL_n and the standard upper Borel subgroup and maximal torus (defined over L), the set of weights naturally corresponds to \mathbb{Z}^n , and the set of dominant weights is $\mathbb{Z}_+^n := \{\nu = (\nu_1, \dots, \nu_n) \in \mathbb{Z}^n : \nu_i \geq \nu_{i+1} \forall i\}$. We set the L -vector space W_ν to be the irreducible representation of highest weight ν . We will need to choose a \mathcal{O} lattice of W_ν , which we do in the style of [Ger19, Section 1.1] as follows. Note

that GL_n , B , and T are defined over \mathcal{O} . For a dominant weight ν , set ξ_ν to be the induced representation $\mathrm{Ind}_B^{\mathrm{GL}_n}(w_0(\nu))_{/\mathcal{O}}$ of the algebraic group $\mathrm{GL}_n/\mathcal{O}$ defined as the functor whose R points are

$$\mathrm{Ind}_B^G(w_0(\nu)) := \{f \in R[\mathrm{GL}_n] : f(bg) = w_0(\nu)(b) \cdot f(g) \forall g \in \mathrm{GL}(R), b \in B(R)\},$$

where w_0 the longest element of the Weyl group. By Proposition II.2.2 and Corollary II.5.6 of [Jan03], the representation ξ_ν is irreducible of highest weight ν . We denote by M_ν the representation given by the \mathcal{O} -points of ξ_ν , so that $M_\nu \otimes_{\mathcal{O}} L \cong \xi_\nu(L) \cong W_\nu$.

Remark. The presence of w_0 is due to our convention that chooses B as the Borel of *upper* triangular matrices, whereas Jantzen induces from the Borel of *lower* triangular matrices. These two choices of Borel subgroup are related by w_0 .

The finite-dimensional algebraic representations in L vector spaces of $G_{D, F_l^+} \cong \prod_{w \in \tilde{S}_l} \mathrm{GL}_{n, F_w}$ are characterised by the sequence of dominant weights, one for each embedding corresponding to $w \in \tilde{S}_l$. We define the set as $W = (\mathbb{Z}_+^n)^{\mathrm{Hom}(F^+, L)}$. For each $\mu \in W$, we can now define the algebraic representation of $G_{D/\mathcal{O}_{F^+}}$ with highest weight μ by $M_\mu = \bigotimes_{\tau \in \mathrm{Hom}(F^+, L), \mathcal{O}} M_{\nu_\tau}$, and $W_\mu = M_\mu \otimes_{\mathcal{O}} L$.

For each $v \in S_D$, choose a finite-free \mathcal{O} -module representation $\rho_v : G_D(\mathcal{O}_{F^+, v}) \rightarrow \mathrm{GL}(M_v)$ with open kernel such that $M_v \otimes \bar{L}$ is irreducible. Set $M_{\{\rho_v\}} = \bigotimes_{v \in S_D} M_v$. We set $M_{\mu, \{\rho_v\}} = M_\mu \otimes M_{\{\rho_v\}}$.

Definition 4.1. Let $\lambda = (\mu, \{\rho_v\})$ be as above. We define the space of automorphic forms for G_D of weight λ with A -coefficients $S_\lambda(A)$, where A is an \mathcal{O} -module, as the space of functions

$$f : G_D(F^+) \backslash G_D(\mathbb{A}_{F^+}^\infty) \rightarrow M_\lambda \otimes_{\mathcal{O}} A$$

such that there is an open compact subgroup

$$U \subset G_D(\mathbb{A}_{F^+}^{\infty, S_l}) \times G_D(\mathcal{O}_{F^+, l})$$

with

$$u|_{S_l \cup S_D} f(gu) = f(g)$$

for all $g \in G_D(\mathbb{A}_{F^+}^\infty)$ and $u \in U$ where $u|_{S_l \cup S_D}$ denotes the action of u on M_λ factoring through $\prod_{v \in S_D \cup S_l} G_D(F_v^+)$.

Notice that $S_\lambda(A)$ is a smooth representation of $G_D(\mathbb{A}_{F^+}^\infty)$ under the action

$$(h \cdot f)(g) = h|_{S_l \cup S_D} f(gh).$$

(Again, $h|_{S_l \cup S_D}$ acts through the representation of $G_D(F_l^+) \times \prod_{v \in S_D} G_D(F_v^+)$ on M_λ .) We denote by $S_\lambda(U, A) = S_\lambda(A)^U$ the invariants under this action.

4.3. Hecke operators. For much of what remains, the argument will be a slight adaptation of that in [Ger19], the important details of which can be found in Sections 2 and 4. Let T be a finite set of places of F^+ containing $S_D \cup S_l$ such that all places in T split in F , and let \tilde{T} be a set of primes of F above those in T as defined before. Fix an open compact subgroup $U = \prod_v U_v$ of $G_D(\mathbb{A}_{F^+}^\infty)$ such that U_v is hyperspecial at all places v outside T . Suppose further that U is sufficiently small; that is, there is a place

v such that U_v contains no elements of finite order other than the identity. We define the Hecke operators on the subspace $S_\lambda(U, A)$.

Hecke operators at unramified places. Let v be a place of F^+ split in F and \tilde{v} be a place in F over v . Let $\varpi_{\tilde{v}}$ be a uniformiser. We can define the Hecke operators as the double coset operators

$$T_v^{(i)} = \left[i_v^{-1} \left(\mathrm{GL}_n(\mathcal{O}_{F, \tilde{v}}) \begin{pmatrix} \varpi_{\tilde{v}} I_i & 0 \\ 0 & I_{n-i} \end{pmatrix} \mathrm{GL}_n(\mathcal{O}_{F, \tilde{v}}) \right) \times U^v \right]$$

Hecke operators at places dividing l . At places dividing the residual characteristic of \mathcal{O} , we set

$$\alpha_{\tilde{v}}^{(i)} = \begin{pmatrix} \varpi_{\tilde{v}} I_i & 0 \\ 0 & I_{n-i} \end{pmatrix} \quad \text{and} \quad U_{\mu, \tilde{v}}^{(i)} = (w_0 \mu_v) (\alpha_{\tilde{v}}^{(i)})^{-1} [U \alpha_{\tilde{v}}^{(i)} U],$$

where w_0 is the longest element of the Weyl group of GL_n and $\mu = (\mu_v) \in W$ with μ_v the dominant weight for the corresponding embedding $F^+ \hookrightarrow L$.

We make the following adjustment to the group U .

Definition 4.2. For v a place of F^+ above l and b a positive integer, let $I^b(\tilde{v})$ be the set of matrices in $\mathrm{GL}_n(\mathcal{O}_{F_{\tilde{v}}})$ which are upper-triangular and unipotent mod \tilde{v}^b . Define $U(l^b) = \prod_{v \in S_l} i_v^{-1} (I^b(\tilde{v})) \times U^l \subseteq G_D(\mathbb{A}_{F^+}^\infty)$ where U^l denotes the product $\prod_{v \nmid l} U_v$.

In the case with the group $U(l^b)$, we further define the following diamond operators:

Definition 4.3. Let T_n be the maximal torus inside GL_n as before. For $v \in S_l$, and $u \in T_n(\mathcal{O}_{F_{\tilde{v}}})$, define $\langle u \rangle$ as the operator

$$[U(l^b)uU(l^b)]$$

on $S_\lambda(U(l^b), A)$. For $u \in T_n(\mathcal{O}_{F^+, l}) = \prod_{v \in S_l} T_n(\mathcal{O}_{F_v}) \cong \prod_{v \in S_l} T_n(\mathcal{O}_{F_{\tilde{v}}})$, define $\langle u \rangle = \prod_{v \in S_l} \langle u_{\tilde{v}} \rangle$.

Let A be an \mathcal{O} -algebra and M an A -module. Define the Hecke algebra $\mathbb{T}^T = \mathbb{T}^T(U(l^b), M)$ as the A -subalgebra of $\mathrm{End}(S_\lambda(U(l^b), M))$ generated by all the operators

$$\left\{ (T_{\tilde{v}}^{(i)}, (T_{\tilde{v}}^{(n)})^{-1}) : v \notin T \text{ split in } F \right\} \cup \left\{ U_{\mu, \tilde{v}}^{(i)} : v \in S_l \right\} \cup \left\{ \langle u \rangle : u \in T_n(\mathcal{O}_{F^+, l}) \right\}.$$

Notice that the map $u \mapsto \langle u \rangle$ defines a group homomorphism

$$T_n(\mathcal{O}_{F^+, l}) \rightarrow \mathbb{T}^T(U(l^b), M)^\times$$

which factors through $T_n(\mathcal{O}_{F^+, l}/l^b) = \prod_{v \in S_l} T_n(\mathcal{O}_{F^+, v}/v^b)$.

4.4. Big ordinary Hecke algebras and the action of Λ . From this point on, we wish to focus on the cases where $A \in \mathrm{Mod}_{\mathcal{O}}$ is one of \mathcal{O} , L/\mathcal{O} or a finite module $\mathcal{O}/\pi^n \mathcal{O}$.

Recall from Hida theory, as fully explained in Section 2.4 of [Ger19], that for any place $v \in S_l$ and any i , the operator $e_v^{(i)} := \lim_{n \rightarrow \infty} (U_{\mu, \tilde{v}}^{(i)})^{n!}$ is a projection on $S_\lambda(U, A)$. We can further define the projection $e = \prod_{v, i} e_v^{(i)}$. We define the ordinary submodule $S_\lambda^{\mathrm{ord}}(U, A) := e.S_\lambda(U, A)$ as the image of this projection. Since all the Hecke operators commute, this is a Hecke invariant submodule. We also define $\mathbb{T}^{T, \mathrm{ord}}(U(l^b), A) = e \mathbb{T}^T(U(l^b), A)$.

Definition 4.4. Let T_n be the maximal torus of GL_n as before. For $b \geq 1$, let $T_n(l^b)$ be the kernel of $T_n(\mathcal{O}_{F^+,l}) \rightarrow T_n(\mathcal{O}/l^b)$. We define Λ as the algebra

$$\Lambda = \mathcal{O}[[T_n(l)]] = \varprojlim_{b \geq 1} \mathcal{O}[T_n(l)/T_n(l^b)].$$

We denote by a_N the kernel of the map $\Lambda \rightarrow \mathcal{O}[T_n(l)/T_n(l^N)]$. Since U is sufficiently small, we see $S_\lambda^{\mathrm{ord}}(U(l^{b,c}), A)$ is a free Λ/a_b -module, through the action of $T_n(\mathcal{O}_{F^+,l})$, and hence we have an inclusion $\Lambda/a_b \hookrightarrow \mathbb{T}(U(l^b), L/\mathcal{O})$ by Proposition 2.20 of [Ger19].

4.4.1. Infinite level. We need to consider the big ordinary Hecke algebra. Set

$$\mathbb{T}^{T,\mathrm{ord}}(U(l^\infty), A) = \varprojlim_{b > 0} \mathbb{T}^{T,\mathrm{ord}}(U(l^b), A) \quad \text{and} \quad S^{\mathrm{ord}}(U(l^\infty), A) = \varinjlim_{b > 0} S^{\mathrm{ord}}(U(l^b), A).$$

Because of the inclusions $\Lambda/a_b \hookrightarrow \mathbb{T}^{T,\mathrm{ord}}(U(l^{b,c}), L/\mathcal{O})$, we get an inclusion $\Lambda \hookrightarrow \mathbb{T}^{T,\mathrm{ord}}(U(l^\infty), L/\mathcal{O})$, and we see that $S^{\mathrm{ord}}(U(l^\infty), L/\mathcal{O})$ is a discrete Λ -module, so its Pontryagin dual is a compact Λ -module. (and in fact is finite free, by Proposition 2.20 of [Ger19] since we assume $U(l)$ is sufficiently small.)

We can now give a statement of a theorem that we prove by the application of Theorem 3.2. Under certain hypotheses, to be determined in Section 5, we have:

Theorem 5.13. *The $\mathbb{T}^{T,\mathrm{ord}}(U(l^\infty), L/\mathcal{O})$ -module $S^{\mathrm{ord}}(U(l^\infty), L/\mathcal{O})^\vee$ is locally free over the generic fibre $\mathbb{T}^{T,\mathrm{ord}}(U(l^\infty), L/\mathcal{O})[1/l]$.*

Therefore the multiplicity of $S^{\mathrm{ord}}(U(l^\infty), L/\mathcal{O})^\vee$ is the same at every point of $\mathbb{T}^{T,\mathrm{ord}}(U(l^\infty), L/\mathcal{O})$ of characteristic zero, and thus we expect the multiplicity of nonclassical points (those corresponding to Hida families of ordinary automorphic forms) is the same as at classical automorphic forms in $S_\lambda(U, A)$.

5. Galois representations and deformation rings

5.1. Local deformation rings. In this section, we let G_{F^+} and G_F be the absolute Galois groups of F^+ and F , and $G_{F^+,v}$, $G_{F,w}$ be the decomposition groups at the places v, w of F^+ and F .

We now define a deformation problem. Let $v \in S_D$ with residue field of size q_v , and let $X_{\mathrm{St}} \subseteq S_{\mathrm{GL}_n}$ be the irreducible component corresponding to the regular nilpotent orbit. We say that an n -dimensional representation $\rho : G_{F^+,v} \rightarrow \mathrm{GL}_n(A)$ is Steinberg if the representation ρ lies in the A -points of this irreducible component X_{St} . When A is a field of characteristic zero and $WD(\rho) = (r, N)$ is the Weil–Deligne representation obtained from ρ , then this condition is equivalent to the condition r being unramified and the eigenvalues of $r(\mathrm{Frob}_{q_v})$ are in the ratio $q_v^{n-1} : q_v^{n-2} : \dots : q_v : 1$.

Let $\mathcal{C}_{\mathcal{O}}$ be the category of local Artinian \mathcal{O} -algebras with residue field \mathbb{F} , (that is, the category of coefficient rings as defined in Mazur’s article in [CSS97]). For each $v \in S_D$ and Steinberg representation $\bar{\rho}_v : G_{F,\tilde{v}} \rightarrow \mathrm{GL}_n(\mathbb{F})$ define a functor

$$D_{\bar{\rho}_v}^{n,\square} : \mathcal{C}_{\mathcal{O}} \rightarrow \mathfrak{Set}, \quad A \mapsto \{\text{Steinberg liftings of } \bar{\rho}_v \text{ to } A\}.$$

This functor is pro-representable by the complete Noetherian local ring $R_v^{\square,\mathrm{st}} := \mathcal{O}_{X_{\mathrm{St}},\bar{\rho}_v}^\wedge$. When we view X_{St} as a scheme over L , Theorem 3.2 tells us, since q is not a root of unity in L , and is therefore

considerate towards G_L , that any localisation of $R_v^{\square, \text{st}}[1/l]$ is a regular ring and thus that $R_v^{\square, \text{st}}[1/l]$ is regular.

We recall the definition of \tilde{r} -discrete series representations found in Section 2.4.5 in [CHT08].

Definition 5.1. Let $\tilde{r}_v : G_{F, \tilde{v}} \rightarrow \text{GL}_d(\mathcal{O})$ be a representation with the following properties:

- (1) $\tilde{r}_v \otimes \mathbb{F}$ is absolutely irreducible (\mathbb{F} the residue field of \mathcal{O}).
- (2) Every irreducible subquotient of $(\tilde{r}_v \otimes \mathbb{F})|_{I_{\tilde{v}}}$ is absolutely irreducible.
- (3) $\tilde{r} \otimes \mathbb{F} \not\cong \tilde{r} \otimes \mathbb{F}(i)$ for each $i = 0, \dots, m$, where $_ (i)$ denotes the twist by the unramified character sending Frob to q^i .

Whenever R is an \mathcal{O} algebra, we say that $\rho : G_{F, \tilde{v}} \rightarrow \text{GL}_{\text{md}}(R)$ is an \tilde{r} -discrete series representation if there is a decreasing filtration $\{\text{Fil}^i\}$ of ρ by R -direct summands such that

$$\text{gr}^i \rho \cong \text{gr}^0 \rho(i) \quad \text{for } i = 0, \dots, m - 1 \quad \text{and} \quad \text{gr}^0 \rho|_{I, \tilde{v}} \cong \tilde{r}|_{I, \tilde{v}} \otimes_{\mathcal{O}} R.$$

Proposition 5.2. Suppose $l > h_G$. Let \tilde{r} be a rank- d representation as above, and let n be an integer with $d|n$. Let $X_{\tilde{r}, n}$ be the moduli space of framed \tilde{r} -discrete series representations of rank n , defined over \mathcal{O} . Then the base change $(X_{\tilde{r}, n})_L$ is smooth over L .

Proof. Let $S_{\tilde{r}}$ be the moduli stack over \mathcal{O} of n -dimensional \tilde{r} -discrete representations, so that $S_{\tilde{r}} \cong [X_{\tilde{r}}/\text{GL}_n]$, and let $S_{\mathbb{1}}$ be the stack of $m := n/d$ -dimensional $\mathbb{1}$ -discrete series representations. Let $S_{\tilde{r}}^{\text{WD}}$ be the stack over L whose groupoid over R consists of objects (ρ', N) , where ρ' is \tilde{r} -discrete series representation of rank $n = dm$ with open kernel, and N is an element of $\text{End}_R(R^n)$ such that $\rho' N \rho'^{-1} = q^\nu N$. Define $S_{\mathbb{1}}^{\text{WD}}$ analogously. Let t_l be the homomorphism $t_l : I \rightarrow \mathbb{Z}_l$ sending any lift of the topological generator of tame inertia to $1 \in \mathbb{Z}_l$. Recall that there is a morphism $S_{\tilde{r}}^{\text{WD}} \rightarrow S_{\tilde{r}}$ given by (ρ', N) is sent to the unique representation ρ given by $\rho(g) = \rho'(g) \exp(t_l(g)N)$ for $g \in I$ and $\rho(\text{Frob}) = \rho'(\text{Frob})$. Recall that this is an isomorphism on the base change to L .

Then we have an morphism of algebraic stacks $S_{\mathbb{1}}^{\text{WD}} \rightarrow S_{\tilde{r}}^{\text{WD}}$ given by $(\rho', N) \mapsto (\rho', N) \otimes \tilde{r}$. We claim that this is an isomorphism. By an exercise in Clifford theory and by assumptions on \tilde{r} , the restriction $\tilde{r}|_I$ can be written as a direct sum of pairwise nonisomorphic absolutely irreducible I -representations $\tau \oplus \tau^{\text{Frob}} \oplus \dots \oplus \tau^{\text{Frob}^{k-1}}$ for some $k \in \mathbb{N}$. As ρ' is $\mathbb{1}$ -discrete series in characteristic zero, we see that $(\rho' \otimes \tilde{r})|_I \cong m(\tau \oplus \tau^{\text{Frob}} \oplus \dots \oplus \tau^{\text{Frob}^{k-1}})$. Let $V_{\tilde{r}}(R) = \text{End}_{R[I]}(\tilde{r}^m)$ be the space of I -equivariant maps of any representation in $S^{\text{WD}_{\tilde{r}}}(R)$, and define $V_{\mathbb{1}}(R) = \text{End}_{R[I]}(\mathbb{1}^m)$ similarly. The map

$$V_{\mathbb{1}}(R) \rightarrow V_{\tilde{r}}(R), \quad N \mapsto N \otimes \text{id}_{\tilde{r}}, \tag{2}$$

is injective, and hence is isomorphic onto its image. We claim that if $(\rho, N) \in S_{\tilde{r}}^{\text{WD}}(R)$, then N is in the image of this map.

First, note that N is I -equivariant. We calculate using Schur’s lemma that $V_{\tilde{r}}(R) \cong M_m(R)^k$, since each τ^{Frob^i} is absolutely irreducible, and we see that the above map corresponds to the diagonal map $\Delta : M_m(R) \rightarrow M_m(R)^k$.

The space $V_{\tilde{r}}(R)$ has a natural action of Frobenius on it, and under this action $N = (N_1, \dots, N_k) \in M_m(R)^k$ has $\text{Frob} \cdot (N_1, \dots, N_k) = q(N_1, \dots, N_k)$. Notice that Frob induces an isomorphism of the underlying spaces $\tau^m \rightarrow (\tau^{\text{Frob}})^m$, which gives us a commutative diagram

$$\begin{array}{ccc} \tau^m & \xrightarrow{\text{Frob}} & (\tau^{\text{Frob}})^m \\ \downarrow N_1 & & \downarrow qN_2 \\ \tau^m & \xrightarrow{\text{Frob}} & (\tau^{\text{Frob}})^m \end{array} \tag{3}$$

Hence, we see $(qN_2, \dots, qN_k, qN_1) = q(N_1, \dots, N_{k-1}, N_k)$, and thus N lies in the image of the diagonal map. This proves the claim.

Let $\chi_{\tilde{r}} = \text{Hom}_I(\tau, \tilde{r})$. This is an unramified character. We claim that $(\text{Hom}_I(\tau, _) \otimes \chi_{\tilde{r}}^{-1}, \Delta^{-1}) : S_{\tilde{r}}^{\text{WD}} \rightarrow S_{\mathbb{1}}^{\text{WD}}$ is an inverse defining the equivalence.

We first show that the composition $S_{\tilde{r}}^{\text{WD}}(R) \rightarrow S_{\mathbb{1}}^{\text{WD}}(R) \rightarrow S_{\tilde{r}}^{\text{WD}}(R)$ is the identity. For $(\Theta, N) \in S_{\tilde{r}}^{\text{WD}}(R)$, the previous claim gives us an isomorphism on the N -part of the stacks $S_{\tilde{r}}^{\text{WD}}(R)$, so we focus on the representation part. Since I acts through a finite quotient, and R is an algebra over a field of characteristic zero, we see that Θ is semisimple and hence we get a sequence of I -representation isomorphisms:

$$\begin{aligned} \Theta &\cong \bigoplus_{i=0}^{k-1} \text{Hom}_I(\tau^{\text{Frob}^i}, \Theta) \otimes \chi_{\tilde{r}}^{-1} \otimes \tau^{\text{Frob}^i} \\ &\cong \text{Hom}_I(\tau, \Theta) \otimes \chi_{\tilde{r}}^{-1} \otimes \bigoplus_{i=0}^{k-1} \tau^{\text{Frob}^i} \cong \text{Hom}_I(\tau, \Theta) \otimes \chi_{\tilde{r}}^{-1} \otimes \tilde{r}. \end{aligned}$$

To show this isomorphism also respects the W_F -action, we observe that each graded part of Θ has $\text{gr}^i(\Theta) \cong \tilde{r} \otimes \chi(i)$, where χ is some unramified character. Then we obtain

$$\text{Hom}_{L[I]}(\tau, \text{gr}^i(\Theta)) \cong \text{Hom}_{L[I]}(\tau, \tilde{r} \otimes \chi(i)) \cong \tilde{r} \otimes \chi(i),$$

so that both sides of the isomorphism are naturally $\tilde{r} \otimes \chi(i)$ as W_F -representations. Hence, the composition $S_{\tilde{r}}^{\text{WD}}(R) \rightarrow S_{\mathbb{1}}^{\text{WD}}(R) \rightarrow S_{\tilde{r}}^{\text{WD}}(R)$ is the identity.

To show that $S_{\mathbb{1}}^{\text{WD}}(R) \rightarrow S_{\tilde{r}}^{\text{WD}}(R) \rightarrow S_{\mathbb{1}}^{\text{WD}}(R)$ is the identity, let $\rho \in S_{\mathbb{1}}(R)$. Then the natural map

$$\rho \rightarrow \text{Hom}_I(\tau, \rho \otimes \tilde{r}), \quad v \mapsto \{w \mapsto v \otimes w\}, \tag{4}$$

defines an I isomorphism. So we need only check that $\rho \otimes \chi_{\tilde{r}}$ and $\text{Hom}_I(\tau, \rho \otimes \tilde{r})$ have the same action of Frobenius. This can be checked again, by looking at the character $\text{gr}^i(\rho)$. Hence, we have exhibited an equivalence of categories $S_{\mathbb{1}} \leftrightarrow S_{\tilde{r}}$.

Given a choice of Frobenius Frob and a topological generator s of the tame inertia group we can explicitly write an isomorphism of stacks

$$\begin{aligned} S_{\mathbb{1}} &\cong [X_{\text{St}}/\text{GL}_m], \\ \rho &\mapsto (\rho(\text{Frob}), \log(\rho(s))), \\ \rho_{\Phi}(\text{Frob}^n x) &= \Phi^n \exp(Nt_l(x)) \leftarrow (\Phi, N). \end{aligned}$$

As $(X_{St})_L$ is a smooth scheme by Theorem 3.2, this shows that $S_{\mathbb{1}}[1/l]$ is a smooth stack, and thus that $S_{\tilde{r}}[1/l]$ and $(X_{\tilde{r},n})_L$ are smooth. \square

In light of this proposition, if $\bar{\rho} : G_{F,\bar{v}} \rightarrow \text{GL}_n(\mathbb{F})$ is an \tilde{r} -discrete series representation, we let $R_v^{\square,\tilde{r}}$ be the universal lifting ring of \tilde{r} -discrete series representations. By the proposition, the ring $R_v^{\square,\tilde{r}}[1/l]$ is regular at every maximal ideal.

5.1.1. Deformation rings at primes above l . For $v \in S_l$, let $\bar{I}_{\bar{v}}$ be the inertia subgroup of $G_{F,\bar{v}}^{\text{ab}}$, let $\bar{I}_{\bar{v}}(l)$ be the pro- l part, and let $\Lambda_{\bar{v}} := \mathcal{O}[[\bar{I}_{\bar{v}}(l)^n]]$, which we can identify with the universal lifting algebra of an ordered set of inertial characters $\{\bar{\chi}_i : I_{\bar{v}} \rightarrow \mathbb{F}^\times\}_{i=1,\dots,n}$. Following chapter 3 of [Ger19] we can define a lifting $\Lambda_{\bar{v}}$ -algebra R_v^Δ as follows.

Take the universal lifting ring $R_v^{\square,\Lambda}$, so that a morphism $r : R_v^{\square,\Lambda} \rightarrow A$ corresponds to a pair $(\rho, \{\chi_i\}_{i=1,\dots,n})$ consisting of a representation $\rho : G_v \rightarrow \text{GL}_n(A)$ lifting $\bar{\rho}$ and a sequence of characters $\chi_i : I_{\bar{v}} \rightarrow A^\times$. Let $\mathcal{F}lag$ be the flag variety defined over \mathcal{O} . There is a subscheme \mathcal{G} of $\mathcal{F}lag \times_{\mathcal{O}} \text{Spec } R_v^{\square,\Lambda}$ whose A -points are the triples $(\text{Fil}, \rho, \{\chi_i\}) \in (\mathcal{F}lag \times_{\mathcal{O}} \text{Spec } R^\square)(A)$ such that $\rho : G_v \rightarrow \text{GL}_n(A)$ preserves the filtration Fil on A^n , and such that the action of $I_{\bar{v}}$ on the graded part $\text{Fil}_j/\text{Fil}_{j-1}$ is χ_j . Then R_v^Δ is defined as the image of the natural morphism $R_v^{\square,\Lambda} \rightarrow \Gamma(\mathcal{G}, \mathcal{O}_{\mathcal{G}})$.

By Lemma 3.3 of [Ger19], the morphism $R_v^\square \rightarrow \mathcal{O}$ corresponding to a representation $\rho : G_v \rightarrow \text{GL}_n(\mathcal{O})$ factors through R_v^Δ if and only if ρ is $\text{GL}_n(\mathcal{O})$ -conjugate to an upper triangular representation with diagonal characters equal to χ_1, \dots, χ_n when restricted to inertia.

Definition 5.3. If A is a \mathbb{Z}_l -algebra and $v \in S_l$, we call a representation $\rho : G_v \rightarrow \text{GL}_n(A)$ *ordinary* if it is $\text{GL}_n(A)$ -conjugate to an upper triangular matrix. Likewise, if $\rho : \text{Gal}(\bar{F} : F) \rightarrow \text{GL}_n(A)$ is a global Galois representation, we say ρ is ordinary if $\rho|_{G_v}$ is ordinary at all places $v \in S_l$.

In this terminology, the fact above can be restated by saying that a point $x : R_v^\square \rightarrow \mathcal{O}$ factors through R_v^Δ if and only if the corresponding representation ρ_x is ordinary.

Lemma 5.4. *Suppose that $\bar{\rho}_v : G_{F,\bar{v}} \rightarrow \text{GL}_n(\mathbb{F})$ is an ordinary Galois representation with diagonal characters $\bar{\chi}_1, \bar{\chi}_2, \dots, \bar{\chi}_n$ such that no pair $i < j$ has $\bar{\chi}_i = \varepsilon \bar{\chi}_j$, where ε is the cyclotomic character. Then $R_v^\Delta[1/l]$ is formally smooth of dimension $[F_{\bar{v}} : \mathbb{Q}_l]^{\frac{1}{2}}n(n+1) + n^2$ over L .*

Proof. This follows from Lemmas 3.17 and 3.7 of [Ger19]. (To apply Lemma 3.17 as stated there, one must note that \mathcal{G}^{ar} is a union of irreducible components of \mathcal{G} and $\bar{\rho}_v$ lies in the open subset of $\mathcal{G}[1/l]$ whose closure is defined to be \mathcal{G}^{ar} . Thus $R_v^{\Delta,ar} = R_v^\Delta$.) \square

5.2. Local-global compatibility. We start by introducing the group \mathcal{G}_n from [CHT08], defined as the group scheme that is the semidirect product of $\text{GL}_n \times \text{GL}_1$ with $C_2 = \{1, j\}$, where j acts as

$$j(g, \mu)j^{-1} = (\mu(g^{-1})^T, \mu).$$

By Lemma 2.1.1 of [CHT08], representations $r : G_{F^+} \rightarrow \mathcal{G}_n(R)$ such that $r^{-1}(\text{GL}_n(R) \times \text{GL}_1(R)) = G_F$ are in correspondence with pairs (ρ, χ) , where ρ is an n -dimensional representation of G_F and χ is a character of G_{F^+} such that $\rho^c \cong \chi\rho^\vee$ and $c \in G_{F^+}$ is sent to j .

For brevity, whenever we have a homomorphism $r : G_{F^+} \rightarrow \mathcal{G}_n(R)$ and a subgroup $H \subset G_{F^+}$, we use $r|_H$ to mean restriction to H , followed by projection to GL_n . Typically, H will be the subgroup G_F or its localisations $G_{F,w}$.

Proposition 5.5. *Suppose that $\mathfrak{m} \trianglelefteq \mathbb{T}^{T,\mathrm{ord}}(U(l^\infty), \mathcal{O})$ is a maximal ideal with residue field \mathbb{F} . Then there is a unique continuous semisimple representation*

$$\bar{r}_{\mathfrak{m}} : G_F \rightarrow \mathrm{GL}_n(\mathbb{F})$$

satisfying the following conditions:

- (1) $\bar{r}_{\mathfrak{m}}^c \cong \bar{r}_{\mathfrak{m}}^\vee \otimes \varepsilon^{1-n}$.
- (2) $\bar{r}_{\mathfrak{m}}|_w$ is unramified at all places v of F^+ outside T .
- (3) If v additionally splits as $v = ww^c$ in F , then the characteristic polynomial of $\bar{r}_{\mathfrak{m}}(\mathrm{Frob}_w)$ is

$$X^n - T_w^{(1)} X^{n-1} + \dots + (-1)^j N(w)^{\frac{1}{2}j(j-1)} T_w^{(j)} X^{n-j} + \dots + (-1)^n N(w)^{\frac{1}{2}n(n-1)} T_w^{(n)}$$

modulo \mathfrak{m} .

- (4) Let $\tilde{r}_{\tilde{v}} : G_F \rightarrow \mathrm{GL}_{m_v}(\mathcal{O})$ be constructed from the smooth representation $\rho_v : G_D(F_v^+) \rightarrow \mathrm{GL}(M_v)$ via the Jacquet–Langlands and local Langlands correspondences, as in [CHT08, Section 3.3, p. 97]. If $v \in S_D$ and $U_v = G_D(\mathcal{O}_{F^+,v})$, then $\bar{r}_{\mathfrak{m}}|_{G_{F,v}}$ is $\tilde{r}_{\tilde{v}}$ -discrete series.

Proof. We prove only (4), the other statements amounting to Proposition 2.28 in [Ger19]. By the argument in the proof of that same proposition, the maximal ideals of \mathbb{T} are in bijection with those of \mathbb{T}/m_Λ . Hence, (4) follows from the classical situation (that is, usual automorphic forms for G_D rather than Hida families of ordinary automorphic forms). The proof of this can be found in Proposition 3.4.2 of [CHT08]. \square

Proposition 5.6. *If \mathfrak{m} is non-Eisenstein (that is, if $\bar{r}_{\mathfrak{m}}$ is irreducible), then $\bar{r}_{\mathfrak{m}}$ can be extended to a representation $\bar{r}_{\mathfrak{m}} : G_{F^+} \rightarrow \mathcal{G}_n(\mathbb{F})$, which in turn can be lifted to a representation*

$$r_{\mathfrak{m}} : G_{F^+} \rightarrow \mathcal{G}_n(\mathbb{T}^{T,\mathrm{ord}}(U(l^\infty), \mathcal{O})_{\mathfrak{m}})$$

with the following properties:

- (1) If $\nu : \mathcal{G}_n \rightarrow \mathrm{GL}_1$ is the second projection, then $\nu \circ r_{\mathfrak{m}} = \varepsilon^{1-n} \delta_{F/F^+}^{\mu_{\mathfrak{m}}}$, where ε is the cyclotomic character, δ_{F/F^+} is the nontrivial character of G_{F^+}/G_F , and $\mu_{\mathfrak{m}} \in \mathbb{Z}/2$.
- (2) $\bar{r}_{\mathfrak{m}}|_{\tilde{v}}$ is unramified at all places $v \notin T$.
- (3) If v in addition splits as $v = ww^c$ in F , then the characteristic polynomial of $\bar{r}_{\mathfrak{m}}(\mathrm{Frob}_w)$ is

$$X^n - T_w^{(1)} X^{n-1} + \dots + (-1)^j N(w)^{\frac{1}{2}j(j-1)} T_w^{(j)} X^{n-j} + \dots + (-1)^n N(w)^{\frac{1}{2}n(n-1)} T_w^{(n)}.$$

- (4) If $v \in S_D$, then $r_{\mathfrak{m}}|_{G_{F,\tilde{v}}}$ is $\tilde{r}_{\tilde{v}}$ -discrete series.

Proof. As with the previous proposition, this is Proposition 2.29 in [Ger19] along with the additional (4), which we prove. By the proof of Proposition 2.29 of [Ger19], we may find a sequence of maximal ideals $\mathfrak{m}_b \subset \mathbb{T}^{T,\text{ord}}(U(l^b), \mathcal{O})$ such that $\mathbb{T}_{\mathfrak{m}} = \varprojlim_b \mathbb{T}^{T,\text{ord}}(U(l^b), \mathcal{O})_{\mathfrak{m}_b}$, and we define $r_{\mathfrak{m}} = \varprojlim_b r_{\mathfrak{m}_b}$. By Lemma 3.4.4 of [CHT08], each $r_{\mathfrak{m}_b}|_{G_{F,\tilde{v}}}$ is $\tilde{r}_{\tilde{v}}$ -discrete series, and so now it remains to show that $r_{\mathfrak{m}}|_{G_{F,\tilde{v}}}$ is, too. Since

$$r_{\mathfrak{m}_b} \otimes \mathbb{T}^{T,\text{ord}}(U(l^{c,c}), \mathcal{O})_{\mathfrak{m}_c} = r_{\mathfrak{m}_c}$$

whenever $b > c$, the filtration Fil_b^i on $r_{\mathfrak{m}_b}$ descends to a filtration $\text{Fil}_b^i \otimes \mathbb{T}^{T,\text{ord}}(U(l^{c,c}), \mathcal{O})_{\mathfrak{m}_c}$ on $r_{\mathfrak{m}_c}$, and the graded parts have

$$[\text{gr}^i(r_{\mathfrak{m},b})] \otimes \mathbb{T}^{T,\text{ord}}(U(l^{c,c}), \mathcal{O})_{\mathfrak{m}_c} \cong \text{gr}^i[r_{\mathfrak{m},b} \otimes \mathbb{T}^{T,\text{ord}}(U(l^{c,c}), \mathcal{O})_{\mathfrak{m}_c}].$$

It follows that $\text{Fil}_b^i \otimes \mathbb{T}^{T,\text{ord}}(U(l^{c,c}), \mathcal{O})_{\mathfrak{m}_c}$ is a defining filtration on $r_{\mathfrak{m}_c}$. From Lemma 2.4.25 of [CHT08], such a filtration is unique, so we have a compatible system of filtrations on the $r_{\mathfrak{m}_b}$ which lift to a filtration on $r_{\mathfrak{m}}|_{G_{F,\tilde{v}}}$. We see from this compatibility that $\text{gr}^i(r_{\mathfrak{m}}) = \varprojlim_b \text{gr}^i(r_{\mathfrak{m}_b})$, and so $r_{\mathfrak{m}}|_{G_{F,\tilde{v}}}$ is $\tilde{r}_{\tilde{v}}$ -discrete series. □

To complete the results we need for local-global compatibility, we need the following lemma:

Lemma 5.7. *Let $\tilde{v} \in \tilde{S}_l$, and let $R_{\tilde{v}}^{\Delta}$ be as before. Then there is a map $R_{\tilde{v}}^{\Delta} \rightarrow \mathbb{T}^{T,\text{ord}}(U(l^{\infty}), \mathcal{O})_{\mathfrak{m}}$ such that*

$$\begin{array}{ccc} G_{F,\tilde{v}} & \xrightarrow{\rho^{\Delta}} & \mathcal{G}_n(R_{\tilde{v}}^{\Delta}) \\ & \searrow r_{\mathfrak{m}} & \downarrow \\ & & \mathcal{G}_n(\mathbb{T}^{T,\text{ord}}(U(l^{\infty}))) \end{array}$$

commutes.

Proof. This follows directly from Corollary 4.3 of [Ger19]. □

5.3. Global deformation rings. Let F/F^+ and let $\bar{\rho} : G_F \rightarrow \text{GL}_n(\mathbb{F})$ be a representation with local representations $\rho_w = \bar{\rho}|_{G_{F,w}}$, where w is a place of F . Let R be the set of places v of F^+ such that v splits and there is a place w of F above v where ρ ramifies. Set $T = S_l \sqcup S_D \sqcup R$, and define \tilde{T} as before. We make the following assumptions:

- The representation $\bar{\rho}$ is an irreducible automorphic representation. That is, there is a non-Eisenstein maximal ideal $\mathfrak{m} \trianglelefteq \mathbb{T}^{T,\text{ord}}(U(l^{\infty}), \mathcal{O})$ such that $\bar{\rho} \cong \bar{r}_{\mathfrak{m}}$.
- The subgroup $\rho(G_{F^+(\zeta_l)}) \subseteq \mathcal{G}_n(\mathbb{F})$ is adequate in the sense of Definition 2.3 of [Tho12].
- The representation $\bar{\rho}$ is unramified outside \tilde{T} .
- At any place $v \in R$, any lift of $\bar{\rho}_v$ to $\bar{\mathbb{Q}}_l$ is nondegenerate in the sense of Section 3.3 of [Sho18]; in particular they lie on a single irreducible component of $\text{Loc}_{\text{GL}_n, \mathbb{Q}_l}^{\square}$.
- For each $v \in S_l$, we have $\text{Hom}_{G_{F,\tilde{v}}}(\bar{\rho}_{\tilde{v}}, \bar{\rho}_{\tilde{v}}\varepsilon) = 0$, for ε the cyclotomic character.

As $\bar{\rho} \cong \bar{r}_m$ is irreducible, it can be extended to a representation $\bar{\rho} : G_{F^+} \rightarrow \mathcal{G}_n(\mathbb{F})$ such that $\nu \circ \bar{\rho} = \varepsilon^{1-n} \delta_{F/F^+}^{\mu_m}$ via Proposition 5.6. We fix such an extension.

For each $v \in T$, define R_v^\square as the framed deformation ring for $\bar{\rho}_{\bar{v}}$. Set

$$R^{\text{loc}} := \left(\widehat{\bigotimes}_{v \in S_I} R_v^\Delta \right) \widehat{\otimes}_{\mathcal{O}} \left(\widehat{\bigotimes}_{v \in S_D} R_v^{\square, \tilde{r}_{\bar{v}}} \right) \widehat{\otimes}_{\mathcal{O}} \left(\widehat{\bigotimes}_{v \in R} R_v^\square \right)$$

as the local deformation ring for $\bar{\rho}$. Our first observation is that, since each R_v^Δ is a $\Lambda_{\bar{v}}$ -module, the algebra R^{loc} inherits the structure of a $\widehat{\bigotimes}_{v \in S_I} \Lambda_{\bar{v}} \cong \Lambda$ -module. The isomorphism $\widehat{\bigotimes}_{v \in S_I} \Lambda_{\bar{v}} \cong \Lambda$ is inherited from the group isomorphisms

$$T_n(\mathfrak{h}) \cong \prod_{v \in S_I} T_n \mathcal{O}_{F^+, v}(l) \cong \prod_{v \in S_I} T_n \mathcal{O}_{F, \bar{v}}(l) \cong \prod_{v \in S_I} \bar{I}_{\bar{v}}(l)^n$$

where the final isomorphism is given by the Artin map of local class field theory.

Lemma 5.8. *The ring $R^{\text{loc}}[1/l]$ is regular.*

Proof. Recall the construction of the complete ring R_v^Δ as the image of $R_v^{\square, \Lambda} = R_v^\square \widehat{\otimes}_{\mathcal{O}} \Lambda_v$ in the global sections of $\mathcal{G} \subseteq \text{Flag} \times_{\mathcal{O}} R_v^{\square, \Lambda}$. We “decomplete” R_v^Δ as follows: R_v^\square is the completion of a local ring \tilde{R}_v^\square at a closed point on a finite-type scheme over \mathcal{O} . The ring $\Lambda_v = \mathcal{O}[[\bar{I}_{\bar{v}}(l)^n]]$ is a completed group algebra of a group which is topologically finitely generated, generated by a fixed choice of generators $\{s_i\}$. So we can choose a subring $\tilde{\Lambda}_v = \mathcal{O}[s_i^{\pm 1}]/\langle \text{relations} \rangle \subseteq \Lambda_v$ of finite type over \mathcal{O} which is dense in Λ_v . Thus, there is a ring $\tilde{R}_v^{\square, \Lambda}$ of finite type over \mathcal{O} whose completion is $R_v^{\square, \Lambda}$.

We can define a closed subscheme $\tilde{\mathcal{G}} \subseteq \text{Flag} \times_{\mathcal{O}} \text{Spec}(\tilde{R}_v^{\square, \Lambda})$ cut out by the same equations for \mathcal{G} as in the definition of R_v^Δ . Of course, there is a natural commutative diagram

$$\begin{array}{ccc} \tilde{R}_v^{\square, \Lambda} & \xrightarrow{\tilde{\phi}} & \mathcal{O}(\tilde{\mathcal{G}}) \\ \downarrow & & \downarrow \\ R_v^{\square, \Lambda} & \xrightarrow{\phi} & \mathcal{O}(\mathcal{G}) \end{array}$$

We set \tilde{R}_v^Δ as the image of $\tilde{\phi}$. It is a finite-type ring over \mathcal{O} and, since the equations defining $\mathcal{G} \subseteq \text{Flag} \times \text{Spec}(R_v^{\square, \Lambda})$ are rational (that is, the defining ideal \mathcal{I} equals $\tilde{\mathcal{I}}_{\mathcal{O}} \mathcal{O}_{\text{Flag} \times \text{Spec}(R_v^{\square, \Lambda})}$ for some ideal $\tilde{\mathcal{I}} \subseteq \text{Flag} \times \text{Spec}(\tilde{R}_v^{\square, \Lambda})$), the image $\text{im}(\phi) = R_v^\Delta$ is a completion of \tilde{R}_v^Δ .

It follows that each of R_v^\square , $R_v^{\square, \tilde{r}_{\bar{v}}}$ and R_v^Δ (for $v \in R$, S_D , S_I respectively) is a completion of a local ring at a closed point P_v on a finite-type \mathcal{O} -scheme X_v . By the last two hypotheses on $\bar{\rho}$ listed above, Lemma 5.4 and Proposition 3.6 of [Sho18], we see that $R_v^\Delta[1/l]$ (for $v \in S_I$) is regular and $R_v^\square[1/l]$ (for $v \in R$) is formally smooth. Thus, the closed points P_v on X_v (where $v \in S_I \cup R$) lie on an open subscheme $U_v \subseteq X_v$ whose generic fibre $U_v[1/l]$ is smooth over L . By Proposition 5.2, the same is true for $R_v^{\square, \tilde{r}_{\bar{v}}}$ for $v \in S_D$. Set

$$\tilde{R}^{\text{loc}} := \left(\bigotimes_{\mathcal{O}, v \in S_I} \tilde{R}_v^\Delta \right) \otimes_{\mathcal{O}} \left(\bigotimes_{\mathcal{O}, v \in S_D} \tilde{R}_v^{\square, \tilde{r}_{\bar{v}}} \right) \otimes_{\mathcal{O}} \left(\bigotimes_{\mathcal{O}, v \in R} \tilde{R}_v^\square \right).$$

Then \tilde{R}^{loc} is of finite type over \mathcal{O} and has a maximal ideal m , corresponding to the closed point $(P_v)_v$, with respect to which R^{loc} is the m -adic completion. In addition, $\tilde{R}^{\text{loc}}[1/l]$ is a regular L -algebra. To show that $R^{\text{loc}}[1/l]$ is a regular ring is now a simple application of Lemma 2.9 and [Stacks, 07NY]. \square

In fact, the same argument shows that $R_\infty[1/l]$ is a regular ring whenever R_∞ is a power series ring in a finite number of variables with coefficients in R^{loc} .

Let \mathcal{S} be the tuple

$$\mathcal{S} = (F/F^+, T, \tilde{T}, \varepsilon^{1-n} \delta_{F/F^+}^{\mu_m}, \{R_v^{\Delta, ar} : v \in S_I\}, \{R_v^{\square, st} : v \in S_D\}, \{R_v^\square : v \in R\})$$

and say that $\rho : G_{F^+} \rightarrow \mathcal{G}(A)$ is a lifting of $\bar{\rho}$ to $A \in \mathcal{C}_\Lambda$ of type \mathcal{S} if it has the following properties:

- (1) $\rho|_{G_F}$ lifts \bar{r}_m .
- (2) ρ is unramified outside T .
- (3) For $v \in S_D$, the local representation ρ_v is \tilde{r} -discrete series and gives rise to the morphism $R_v^\square \rightarrow A$ which factors through $R_v^{\square, \tilde{r}}$.
- (4) For $v \in S_I$, the restriction ρ_v and the Λ -structure on A give a morphism $R_v^\square \otimes \Lambda \rightarrow A$ which factors through R_v^Δ .
- (5) $\nu \circ \rho = \varepsilon^{1-n} \delta_{F/F^+}^{\mu_m}$.

By Proposition 2.2.9 of [CHT08], we can construct the universal deformation ring $R_{\mathcal{S}}^{\text{univ}}$ and the universal lifting ring $R_{\mathcal{S}}^\square$.

Let $h_0 = [F^+ : \mathbb{Q}] \cdot \frac{1}{2}n(n-1) + [F^+ : \mathbb{Q}] \cdot \frac{1}{2}n(1 - (-1)^{\mu_m-1})$ and let h be an integer larger than both h_0 and $\dim[H_{\mathcal{L}^\perp}^1(G_{F^+, T}, \text{ad } \bar{\rho}(1))]$. (Here, the space $H_{\mathcal{L}^\perp}^1(G_{F^+, T}, \text{ad } \bar{\rho}(1))$ is a particular subspace of the cohomology group $H^1(G_{F^+, T}, \text{ad } \bar{\rho}(1))$ of the Galois group $G_{F^+, T}$ of the maximal extension of F^+ unramified outside of T , defined in Proposition 4.4 of [Tho12].)

After Thorne [Tho12], we will call a triple $(Q, \tilde{Q}, \{\bar{\psi}_v\}_{v \in Q})$ a Taylor–Wiles triple if

- (1) Q is a set of primes of F^+ which split in F ,
- (2) $l | \text{Nm}_{F^+}(v) - 1$ for each $v \in Q$,
- (3) $|Q| = h$,
- (4) \tilde{Q} is the set $\{\tilde{v} | v \in Q\}$, and
- (5) for each $v \in Q$, the representation $\bar{\rho}|_{G_v}$ splits as a direct sum into $\bar{s}_v \oplus \bar{\psi}_v$ where $\bar{\psi}_v$ is a generalised eigenspace with eigenvalue $\bar{\alpha}_v \in \mathbb{F}$ of dimension d_v .

For any Taylor–Wiles set Q we can define a deformation problem $\mathcal{S}(Q)$ that is the same as \mathcal{S} , but now we allow $\rho_{\tilde{v}}$, for $v \in Q$, to ramify in the following way: $\rho_{\tilde{v}}$ splits as a direct sum $s \oplus \psi$, and the two summands lift to \bar{s} and $\bar{\psi}$ in such a way that s is unramified and $\psi|_{I_v} : I_v \rightarrow \text{GL}_{d_v}$ factors through the scalar action on the underlying representation space. Using Proposition 2.2.9 in [CHT08] again, we can now take the universal deformation ring $R_{\mathcal{S}(Q)}^{\text{univ}}$. Because stipulating that the local deformations at

Taylor–Wiles primes are unramified is a closed condition, this presents us with a surjection $R_{S(Q)}^{\text{univ}} \rightarrow R_S^{\text{univ}}$. We also have a natural map $R^{\text{loc}} \rightarrow R_{S(Q)}^{\text{univ}}$ given by restrictions to the local subgroups at the level of functors.

Proposition 5.9. *For each $N \in \mathbb{N}$, we can find a Taylor–Wiles triple $(Q_N, \tilde{Q}_N, \{\bar{\psi}_v\}_{v \in Q})$ such that $l^N \mid |\text{Nm}_F(v) - 1|$ for all $v \in Q_N$ and the global deformation ring $R_{S(Q)}^{\text{univ}}$ can be topologically generated over R^{loc} by $h - h_0$ generators.*

Proof. This follows from Lemma 4.4 of [Tho12] applied in the case of Theorem 8.6. □

In light of this proposition, set $R_\infty = R^{\text{loc}}[[X_1, \dots, X_h]]$, set $R_N = R_{S(Q_N)}^{\text{univ}}$, and set $R_0 = R_S^{\text{univ}}$ so that we have surjections $R_\infty \twoheadrightarrow R_N$ and $R_N \twoheadrightarrow R_0$.

We now define some important subgroups of $G_D(\mathbb{A}_{F^+}^\infty)$.

Definition 5.10. For $v \in Q_N$, suppose that $\bar{r}|_v = \bar{s} \oplus \bar{\psi}$ as before, with $\bar{\psi}$ a d_v -dimensional semisimple unramified representation with all Frobenius eigenvalues equal. We take the group $U_i(\tilde{v})$ to be the subgroup of $U_v \subseteq G_D(F_v^+)$ (identified with $\text{GL}_n(F_{\tilde{v}})$ via the isomorphism $i_{\tilde{v}}$) of elements that take the form

$$\begin{pmatrix} \varpi_{\tilde{v}}^* & * \\ 0 & aI_{d_v} \end{pmatrix}$$

modulo \tilde{v} with $a \equiv 1 \pmod{\tilde{v}}$ when $i = 1$, and arbitrary when $i = 0$. Set $U_i(Q) = U^Q \times \prod_{v \in Q} U_i(\tilde{v}) \subseteq G_D(\mathbb{A}_{F^+}^\infty)$.

Let Δ_N be the maximal l -power quotient of $U_0(Q_N)/U_1(Q_N) \cong \prod_{v \in Q_N} k(\tilde{v})^\times$. We may view Δ_N as the maximal l -quotient of $\prod_{v \in Q_N} k(\tilde{v})^\times \cong (\mathbb{Z}/l^N)^q$. We claim there is an action of Δ_N on the ring $R_{S(Q)}^{\text{univ}}$. The map $\det \circ r_N^{\text{univ}} : I_{F, \tilde{v}} \rightarrow (R_{S(Q)}^{\text{univ}})^\times$ given by the determinant of the universal deformation $r_N^{\text{univ}} := r_{S(Q_N), \bar{\rho}}^{\text{univ}}$ factors through the kernel of $(R_{S(Q)}^{\text{univ}})^\times \rightarrow \mathbb{F}^\times$, which is an abelian l -power group. By local class field theory, there is an isomorphism $I_{F^{\text{ab}}, \tilde{v}} \rightarrow \mathcal{O}_{F, \tilde{v}}^\times$, and the l -power quotient of this group is the l -power quotient of $k(\tilde{v})^\times$. Hence we see that there is a map $\Delta_N \rightarrow (R_{S(Q_N)}^{\text{univ}})^\times$ and thus a ring map $\Lambda[\Delta_N] \rightarrow R_{S(Q)}^{\text{univ}}$, so that $R_{S(Q_N)}^{\text{univ}}$ inherits the structure of a finitely generated $\Lambda[\Delta_N]$ -algebra. If a_N is the augmentation ideal of $\Lambda[\Delta_N]$, then $R_{S(Q_N)}^{\text{univ}}/a_N$ is the ring of the universal deformation ring which parametrises Galois deformations of type \mathcal{S} . (These deformations are required to be unramified at places above Q_N .) Note that $\Delta_N \cong (\mathbb{Z}/l^m\mathbb{Z})^h$ by our choice of Q_N .

As in Sections 4.3 and 4.4, we can construct the Hecke operators

$$\mathbb{T}^{T \cup Q_N, \text{ord}}(U_1(Q_N)(l^\infty), \mathcal{O})$$

and, through a map $\mathbb{T}^{T \cup Q_N, \text{ord}}(U_1(Q_N)(l^\infty), \mathcal{O}) \rightarrow \mathbb{T}^{T, \text{ord}}(U(l^\infty), \mathcal{O})$, we can lift our choice of maximal ideal \mathfrak{m} to a maximal ideal $\mathfrak{m}_N \subset \mathbb{T}^{T \cup Q_N, \text{ord}}(U_1(Q_N)(l^\infty), \mathcal{O})$. Set $\mathbb{T}_{N,1} := \mathbb{T}^{T \cup Q_N, \text{ord}}(U_1(Q_N)(l^\infty), \mathcal{O})_{\mathfrak{m}_N}^\wedge$ as the \mathfrak{m}_N -adic completion. As in Proposition 5.6, we can construct a representation $r_{\mathfrak{m}_N} : G_{F^+} \rightarrow \mathcal{G}_n(\mathbb{T}_{N,1})$ which by the proof of Theorem 6.8 of [Tho12] gives us an $\mathcal{S}(Q_N)$ -lifting of $\bar{\rho}$. Hence, we get a surjection $R_{S(Q)}^{\text{univ}} \twoheadrightarrow \mathbb{T}_{N,1}$ for each N .

5.4. Patching. We now define a module H_N over $\mathbb{T}^{T \cup Q_N, \text{ord}}(U_1(Q_N)(l^\infty), \mathcal{O})_m$ for each set Q_N , and quote a patching theorem that will allow us to construct the patched “limit” module H_∞ , which we use to prove our local freeness result.

Define the space of automorphic forms $S^{\text{ord}}(U_i(Q_N)(l^\infty), L/\mathcal{O})_m$ as before and set

$$H_0 = S^{\text{ord}}(U(l^\infty), L/\mathcal{O})_m^\vee.$$

In Proposition 5.9 of [Tho12], Thorne describes a projection Pr_v on $S^{\text{ord}}(U_i(Q_N)(l^\infty), L/\mathcal{O})_m$ and, later on, modules

$$H_{i,N} := \prod_{v \in Q_N} \text{Pr}_v [S^{\text{ord}}(U_i(Q_N)(l^\infty), L/\mathcal{O})_m]^\vee$$

with the following properties:

Proposition 5.11 [Tho12, part of proof of Theorem 6.8]. (1) H_{1,Q_N} is a free $\Lambda[\Delta_{Q_N}]$ -module and restriction to $S^{\text{ord}}(U_0(Q_N)(l^\infty), L/\mathcal{O})_m$ gives an isomorphism $H_{1,Q_N}/\mathfrak{a}_N \cong H_{0,Q_N}$.

(2) The map

$$\left(\prod_{v \in Q_N} \text{Pr}_v \right)^\vee : H_{0,Q_N} \rightarrow H_0$$

is an isomorphism.

Theorem 5.12 (patching). Let $R \twoheadrightarrow \mathbb{T}$ be a surjective Λ -algebra homomorphism with \mathbb{T} a finite Λ -algebra. Define $S_N = \Lambda[(\mathbb{Z}/l^h\mathbb{Z})^h] \cong \Lambda[\Delta_{Q_N}]$ with augmentation ideal \mathfrak{a}_N and define the inverse limit $S'_\infty := \varprojlim \Lambda[\Delta_{Q_N}] \cong \Lambda[[Y_1, \dots, Y_h]]$. Set $S_\infty = S'_\infty \hat{\otimes}_{\mathcal{O}} \mathcal{T}$, where $\mathcal{T} = \mathcal{O}[[X_1, \dots, X_{|T|n^2}]]$. Suppose we have the following data:

- (1) integers $t, h \geq 1$,
- (2) a finite \mathbb{T} -module H ,
- (3) for each $N \geq 1$,
 - (a) an S_N -algebra homomorphism $R_N \twoheadrightarrow \mathbb{T}_N$ that gets reduced to $R \twoheadrightarrow \mathbb{T}$ under reduction modulo \mathfrak{a}_N , and
 - (b) a finite \mathbb{T}_N -module H_N , which is finite and free over S_N and whose S_N -rank is independent of N , and
- (4) an S_∞ -algebra R_∞ such that $R_\infty \twoheadrightarrow R_N$ with kernel $\ker(S_\infty \rightarrow S_N)R_\infty$.

Then there is an $R_\infty \otimes S_\infty$ -module H_∞ , such that $H_\infty/\mathfrak{a}H_\infty \cong H$, H_∞ is a finite free S_∞ -module, and the action of S_∞ on H_∞ factors through that of R_∞ .

Proof. The details of the Taylor–Wiles–Kisin patching method used here are as in Chapter 4.3 of [Ger19]. They can also be found in Chapter 8 of [Tho12], under the heading “another patching argument”. \square

Theorem 5.13. *The module $H_0[1/l]$ is a finite locally free $R_S^{\text{univ}}[1/l]$ -module.*

Proof. We calculate that

$$\dim(S_\infty) = \dim(\Lambda) + h + |T|n^2 = n[F^+ : \mathbb{Q}]n + h + |T|n^2$$

and that

$$\begin{aligned} \dim(R_\infty) &= 1 + \sum_{v \in S_l} ([F_v : \mathbb{Q}_l] \cdot \frac{1}{2}n(n+1) + n^2) + n^2|S_D \cup R| + h - h_0 \\ &= [F^+ : \mathbb{Q}] \cdot \frac{1}{2}n(n+1) + |T|n^2 + h - h_0 \\ &= [F^+ : \mathbb{Q}]n + |T|n^2 + h - [F^+ : \mathbb{Q}] \cdot \frac{1}{2}n(1 - (-1)^{\mu_m - n}) \end{aligned}$$

Consider the module H_∞^\square . Since H_∞^\square is a finite free S_∞ module, and since the action of S_∞ factors through R_∞ , we see that

$$\dim(S_\infty) = \text{depth}_{S_\infty}(H_\infty^\square) \leq \text{depth}_{R_\infty}(H_\infty^\square) \leq \dim(R_\infty)$$

and thus, the only possible way for this inequality to hold is if equality holds throughout. This implies $\mu_m \equiv n \pmod{2}$ and H_∞^\square is a maximal Cohen–Macaulay R_∞ module.

Now, consider the generic fibre. Let $m \subseteq R_\infty[1/l]$ be a maximal ideal. Lemma 5.8 shows that $R_\infty[1/l]_m$ is a regular local ring. Thus, any finitely generated maximal Cohen–Macaulay $R_\infty[1/l]_m$ -module has finite projective dimension, and hence any maximal Cohen–Macaulay module is projective by the Auslander–Buchsbaum formula. This shows that $H_\infty^\square[1/l]_m$ is a free $R_\infty[1/l]_m$ -module, this shows that $H_\infty^\square[1/l]$ is a locally finite free $R_\infty[1/l]$ -module. It follows that $H_0[1/l]$ is a locally finite free $R_S^{\text{univ}}[1/l]$ -module. \square

Corollary 5.14. $R_S^{\text{univ}}[1/l] = \mathbb{T}[1/l].$

Proof. Let I be the kernel of the surjection $R_S^{\text{univ}}[1/l] \rightarrow \mathbb{T}[1/l]$. Choose any maximal ideal m of $R_S^{\text{univ}}[1/l]$. Since localisation is an exact functor, we get a short exact sequence

$$0 \rightarrow I_m \rightarrow R_S^{\text{univ}}[1/l]_m \rightarrow \mathbb{T}[1/l]_m \rightarrow 0.$$

Note that the action of $R_S^{\text{univ}}[1/l]_m$ on $H_0[1/l]_m$ factors through $\mathbb{T}[1/l]_m$, so that I_m annihilates all of $H_0[1/l]_m$. Since this is a free module, this shows that I_m is trivial. Since this is true for every m , this shows that $\text{Supp}(I) = \emptyset$ and hence $I = 0$. Hence the surjection above is an isomorphism

$$R_S^{\text{univ}}[1/l] \cong \mathbb{T}[1/l]. \quad \square$$

Remark. As an application of Theorem 5.13, whenever M is a locally free coherent sheaf on a connected space X , the rank function


$$X \rightarrow \mathbb{N} \cup \{0\}, \quad x \mapsto \text{Rank}_x(M),$$

is locally constant. Therefore, the rank of a geometrically connected component can be calculated by


calculating the rank at any special point $x \in X$. In our special case, the rank of the module $H_0[1/l]$ can be interpreted as the number of distinct automorphic forms with a given set of Hecke eigenvalues, which can be interpreted as the multiplicity of the Galois representation determined by said Hecke eigenvalues inside the space of automorphic forms. We have shown that for these automorphic forms, the multiplicity is determined only by the connected component of $R_\infty[1/l]$ on which the representation ρ_m lies. By Lemma 4.2 of [Ger19], we see that the minimal primes of $R_\infty[1/l]$ biject with the minimal primes of Λ . Thus, if one could show that for each component of $\text{Spec } \Lambda$, there is an automorphic form of some classical weight had multiplicity 1, then all the Hida families of forms would also have multiplicity 1.

Appendix: Weighted Dynkin diagrams for distinguished orbits in types D_n and E_n with $n \leq 7$.


Weighted Dynkin diagrams of distinguished orbits of type D_n :



Weighted Dynkin diagrams of distinguished orbits of type E_6 :



Weighted Dynkin diagrams of distinguished orbits of type E_7 :



Acknowledgements

The research in this work is part of the author’s PhD thesis, supported by the Engineering and Physical Sciences Research Council (EPSRC). I would like to thank my supervisor, Jack Shotton, for the advice and support given me throughout this project. I also thank reviewer for the detailed comments and suggestions given.

References

[BDP17] V. Balaji, P. Deligne, and A. J. Parameswaran, “On complete reducibility in characteristic p ”, *Épjournal Géom. Algébrique* **1** (2017), art. id. 3, 27 pp. MR

[Bel16] R. Bellovin, “Generic smoothness for G -valued potentially semi-stable deformation rings”, *Ann. Inst. Fourier (Grenoble)* **66**:6 (2016), 2565–2620. MR

[BG19] R. Bellovin and T. Gee, “ G -valued local deformation rings and global lifts”, *Algebra Number Theory* **13**:2 (2019), 333–378. MR

- [Car93] R. W. Carter, *Finite groups of Lie type: conjugacy classes and complex characters*, Wiley, New York, 1985. MR
- [CHT08] L. Clozel, M. Harris, and R. Taylor, “Automorphy for some l -adic lifts of automorphic mod l Galois representations”, *Publ. Math. Inst. Hautes Études Sci.* 108 (2008), 1–181. MR
- [CM93] D. H. Collingwood and W. M. McGovern, *Nilpotent orbits in semisimple Lie algebras*, Van Nostrand, New York, 1993. MR
- [Cot22] S. Cotner, “Springer isomorphisms over a general base scheme”, preprint, 2022. arXiv 2211.08383
- [CSS97] G. Cornell, J. H. Silverman, and G. Stevens (editors), *Modular forms and Fermat’s last theorem: papers from the Instructional Conference on Number Theory and Arithmetic Geometry* (Boston, 1995), Springer, 1997. MR
- [DHKM20] J.-F. Dat, D. Helm, R. Kurinczuk, and G. Moss, “Moduli of Langlands parameters”, *J. Eur. Math. Soc. (JEMS)* 27:5 (2025), 1827–1927. MR
- [FR08] R. Fowler and G. Röhrle, “On cocharacters associated to nilpotent elements of reductive groups”, *Nagoya Math. J.* 190 (2008), 105–128. MR
- [FS25] L. Fargues, “Geometrization of the local Langlands correspondence: an overview”, *J. Math. Sci. Univ. Tokyo* 32:2 (2025), 157–240. MR
- [Ger19] D. Geraghty, “Modularity lifting theorems for ordinary Galois representations”, *Math. Ann.* 373:3-4 (2019), 1341–1427. MR
- [Gro99] B. H. Gross, “Algebraic modular forms”, *Israel J. Math.* 113 (1999), 61–93. MR
- [Har77] R. Hartshorne, *Algebraic geometry*, Graduate Texts in Mathematics 52, Springer, 1977. MR
- [Hel23] E. Hellmann, “On the derived category of the Iwahori–Hecke algebra”, *Compos. Math.* 159:5 (2023), 1042–1110. MR
- [HT01] M. Harris and R. Taylor, *The geometry and cohomology of some simple Shimura varieties*, Annals of Mathematics Studies 151, Princeton University Press, 2001. MR
- [Hum90] J. E. Humphreys, *Reflection groups and Coxeter groups*, Cambridge Studies in Advanced Mathematics 29, Cambridge University Press, 1990. MR
- [Jan03] J. C. Jantzen, *Representations of algebraic groups*, 2nd ed., Mathematical Surveys and Monographs 107, Amer. Math. Soc., 2003. MR
- [Mat86] H. Matsumura, *Commutative ring theory*, Cambridge Studies in Advanced Mathematics 8, Cambridge University Press, Cambridge, 1986. MR
- [Pre03] A. Premet, “Nilpotent orbits in good characteristic and the Kempf–Rousseau theory”, *J. Algebra* 260:1 (2003), 338–366. MR
- [Sho18] J. Shotton, “The Breuil–Mézard conjecture when $l \neq p$ ”, *Duke Math. J.* 167:4 (2018), 603–678. MR
- [Sho24] J. Shotton, “Irreducible components of the moduli space of Langlands parameters”, *Int. Math. Res. Not.* 2024:11 (2024), 9020–9035. MR
- [Stacks] “The Stacks project”, electronic reference, 2023, available at <http://stacks.math.columbia.edu>.
- [Ste16] R. Steinberg, *Lectures on Chevalley groups*, corrected ed., University Lecture Series 66, Amer. Math. Soc., 2016. MR
- [Tho12] J. Thorne, “On the automorphy of l -adic Galois representations with small residual image”, *J. Inst. Math. Jussieu* 11:4 (2012), 855–920. MR
- [Zhu25] X. Zhu, “Coherent sheaves on the stack of Langlands parameters”, pp. 39–123 in *The Langlands program, II: Geometrization of the Langlands correspondence*, Proc. Sympos. Pure Math. 112.2, Amer. Math. Soc., 2025. MR

Communicated by Frank Calegari

Received 2023-07-24 Revised 2025-03-25 Accepted 2025-06-27

danielfunckmaths@gmail.com

University of Tübingen, 72076 Tübingen, Germany

Algebra & Number Theory

msp.org/ant

EDITORS

MANAGING EDITOR

Antoine Chambert-Loir
Université Paris-Diderot
France

EDITORIAL BOARD CHAIR

David Eisenbud
University of California
Berkeley, USA

BOARD OF EDITORS

Jason P. Bell	University of Waterloo, Canada	Michael J. Larsen	Indiana University Bloomington, USA
Olivier Benoist	Ecole Normale Supérieure, France	Philippe Michel	École Polytechnique Fédérale de Lausanne
Bhargav Bhatt	University of Michigan, USA	Martin Olsson	University of California, Berkeley, USA
Frank Calegari	University of Chicago, USA	Irena Peeva	Cornell University, USA
J-L. Colliot-Thélène	CNRS, Université Paris-Saclay, France	Jonathan Pila	University of Oxford, UK
Brian D. Conrad	Stanford University, USA	Anand Pillay	University of Notre Dame, USA
Samit Dasgupta	Duke University, USA	Bjorn Poonen	Massachusetts Institute of Technology, USA
Hélène Esnault	Freie Universität Berlin, Germany	Victor Reiner	University of Minnesota, USA
Gavril Farkas	Humboldt Universität zu Berlin, Germany	Peter Sarnak	Princeton University, USA
Sergey Fomin	University of Michigan, USA	Michael Singer	North Carolina State University, USA
Edward Frenkel	University of California, Berkeley, USA	Vasudevan Srinivas	SUNY Buffalo, USA
Wee Teck Gan	National University of Singapore	Shunsuke Takagi	University of Tokyo, Japan
Andrew Granville	Université de Montréal, Canada	Pham Huu Tiep	Rutgers University, USA
Ben J. Green	University of Oxford, UK	Ravi Vakil	Stanford University, USA
Christopher Hacon	University of Utah, USA	Akshay Venkatesh	Institute for Advanced Study, USA
Roger Heath-Brown	Oxford University, UK	Melanie Matchett Wood	Harvard University, USA
János Kollár	Princeton University, USA	Shou-Wu Zhang	Princeton University, USA

PRODUCTION

production@msp.org
Silvio Levy, Scientific Editor


See inside back cover or msp.org/ant for submission instructions.

The subscription price for 2026 is US \$590/year for the electronic version, and \$865/year (+\$75, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to MSP.

Algebra & Number Theory (ISSN 1944-7833 electronic, 1937-0652 printed) at Mathematical Sciences Publishers, 2000 Allston Way # 59, Berkeley, CA 94701-4004, is published continuously online.

ANT peer review and production are managed by EditFLOW[®] from MSP.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2026 Mathematical Sciences Publishers

Algebra & Number Theory

Volume 20 No. 6 2026

Effective multiplicative independence of three singular moduli YURI BILU, SANOLI GUN and EMANUELE TRON	1073
The geometry of the unipotent component of the moduli space of Weil–Deligne representations DANIEL FUNCK	1125
Smoothness of stabilisers in generic characteristic BEN MARTIN, DAVID I. STEWART and LEWIS TOPLEY	1159
Derived isogenies and isogenies for abelian surfaces ZHIYUAN LI and HAITAO ZOU	1185
Injectivity and vanishing for the Du Bois complexes of isolated singularities MIHNEA POPA, WANCHUN SHEN and ANH DUC VO	1235