

ANALYSIS & PDE

Volume 6

No. 1

2013

TERENCE TAO

**LOCALISATION AND COMPACTNESS PROPERTIES
OF THE NAVIER-STOKES GLOBAL REGULARITY PROBLEM**



LOCALISATION AND COMPACTNESS PROPERTIES OF THE NAVIER–STOKES GLOBAL REGULARITY PROBLEM

TERENCE TAO

In this paper we establish a number of implications between various qualitative and quantitative versions of the global regularity problem for the Navier–Stokes equations in the periodic, smooth finite energy, smooth H^1 , Schwartz, and mild H^1 categories, and with or without a forcing term. In particular, we show that if one has global well-posedness in H^1 for the periodic Navier–Stokes problem with a forcing term, then one can obtain global regularity both for periodic and for Schwartz initial data (thus yielding a positive answer to both official formulations of the problem for the Clay Millennium Prize), and can also obtain global almost smooth solutions from smooth H^1 data or smooth finite energy data, although we show in this category that fully smooth solutions are not always possible. Our main new tools are localised energy and enstrophy estimates to the Navier–Stokes equation that are applicable for large data or long times, and which may be of independent interest.

1. Introduction

The purpose of this paper is to establish some implications between various formulations of the global regularity problem (either with or without a forcing term) for the Navier–Stokes system of equations, including the four formulations appearing in the Clay Millennium Prize formulation [Fefferman 2006] of the problem, and in particular to isolate a single formulation that implies these four formulations, as well as several other natural versions of the problem. In the course of doing so, we also establish some new local energy and local enstrophy estimates which seem to be of independent interest.

To describe these various formulations, we must first define properly the concept of a solution to the Navier–Stokes problem. We will need to study a number of different types of solutions, including periodic solutions, finite energy solutions, H^1 solutions, and smooth solutions; we will also consider a forcing term f in addition to the initial data u_0 . We begin in the classical regime of smooth solutions. Note that even within the category of smooth solutions, there is some choice in what decay hypotheses to place on the initial data and solution; for instance, one can require that the initial velocity u_0 be Schwartz class, or merely smooth with finite energy. Intermediate between these two will be data which is smooth and in H^1 .

More precisely, we define:

The author is supported by NSF Research Award CCF-0649473, the NSF Waterman Award and a grant from the MacArthur Foundation.

MSC2010: 35Q30, 76D05, 76N10.

Keywords: Navier–Stokes equation, global regularity.

Definition 1.1 (Smooth solutions to the Navier–Stokes system). A *smooth set of data* for the Navier–Stokes system up to time T is a triplet (u_0, f, T) , where $0 < T < \infty$ is a time, the initial velocity vector field $u_0 : \mathbb{R}^3 \rightarrow \mathbb{R}^3$ and the forcing term $f : [0, T] \times \mathbb{R}^3 \rightarrow \mathbb{R}^3$ are assumed to be smooth on \mathbb{R}^3 and $[0, T] \times \mathbb{R}^3$ respectively (thus, u_0 is infinitely differentiable in space, and f is infinitely differentiable in space-time), and u_0 is furthermore required to be divergence-free:

$$\nabla \cdot u_0 = 0. \quad (1)$$

If $f = 0$, we say that the data is *homogeneous*.

The *total energy* $E(u_0, f, T)$ of a smooth set of data (u_0, f, T) is defined by the quantity¹

$$E(u_0, f, T) := \frac{1}{2} (\|u_0\|_{L_x^2(\mathbb{R}^3)} + \|f\|_{L_t^1 L_x^2([0, T] \times \mathbb{R}^3)})^2, \quad (2)$$

and (u_0, f, T) is said to have *finite energy* if $E(u_0, f, T) < \infty$. We define the H^1 *norm* $\mathcal{H}^1(u_0, f, T)$ of the data to be the quantity

$$\mathcal{H}^1(u_0, f, T) := \|u_0\|_{H_x^1(\mathbb{R}^3)} + \|f\|_{L_t^\infty H_x^1(\mathbb{R}^3)} < \infty,$$

and say that (u_0, f, T) is H^1 if $\mathcal{H}^1(u_0, f, T) < \infty$; note that the H^1 regularity is essentially one derivative higher than the energy regularity, which is at the level of L^2 , and instead matches the regularity of the *initial enstrophy*

$$\frac{1}{2} \int_{\mathbb{R}^3} |\omega_0(t, x)|^2 dx,$$

where $\omega_0 := \nabla \times u_0$ is the initial vorticity. We say that a smooth set of data (u_0, f, T) is *Schwartz* if, for all integers $\alpha, m, k \geq 0$, one has

$$\sup_{x \in \mathbb{R}^3} (1 + |x|)^k |\nabla_x^\alpha u_0(x)| < \infty$$

and

$$\sup_{(t, x) \in [0, T] \times \mathbb{R}^3} (1 + |x|)^k |\nabla_x^\alpha \partial_t^m f(x)| < \infty.$$

Thus, for instance, the Schwartz property implies H^1 , which in turn implies finite energy. We also say that (u_0, f, T) is *periodic* with some period $L > 0$ if one has $u_0(x + Lk) = u_0(x)$ and $f(t, x + Lk) = f(t, x)$ for all $t \in [0, T]$, $x \in \mathbb{R}^3$, and $k \in \mathbb{Z}^3$. Of course, periodicity is incompatible with the Schwartz, H^1 , and finite energy properties, unless the data is zero. To emphasise the periodicity, we will sometimes write a periodic set of data (u_0, f, T) as (u_0, f, T, L) .

A *smooth solution to the Navier–Stokes system*, or a *smooth solution*, is a quintuplet (u, p, u_0, f, T) , where (u_0, f, T) is a smooth set of data, and the velocity vector field $u : [0, T] \times \mathbb{R}^3 \rightarrow \mathbb{R}^3$ and pressure field $p : [0, T] \times \mathbb{R}^3 \rightarrow \mathbb{R}$ are smooth functions on $[0, T] \times \mathbb{R}^3$ that obey the Navier–Stokes equation

$$\partial_t u + (u \cdot \nabla) u = \Delta u - \nabla p + f \quad (3)$$

and the incompressibility property

$$\nabla \cdot u = 0 \quad (4)$$

¹We will review our notation for space-time norms such as $L_t^p L_x^q$, together with sundry other notation, in [Section 2](#).

on all of $[0, T] \times \mathbb{R}^3$, and also the initial condition

$$u(0, x) = u_0(x) \tag{5}$$

for all $x \in \mathbb{R}^3$. We say that a smooth solution (u, p, u_0, f, T) has *finite energy* if the associated data (u_0, f, T) has finite energy, and in addition one has²

$$\|u\|_{L_t^\infty L_x^2([0, T] \times \mathbb{R}^3)} < \infty. \tag{6}$$

Similarly, we say that (u, p, u_0, f, T) is H^1 if the associated data (u_0, f, T) is H^1 , and in addition one has

$$\|u\|_{L_t^\infty H_x^1([0, T] \times \mathbb{R}^3)} + \|u\|_{L_t^2 H_x^2([0, T] \times \mathbb{R}^3)} < \infty. \tag{7}$$

We say instead that a smooth solution (u, p, u_0, f, T) is *periodic* with period $L > 0$ if the associated data $(u_0, f, T) = (u_0, f, T, L)$ is periodic with period L , and if $u(t, x + Lk) = u(t, x)$ for all $t \in [0, T]$, $x \in \mathbb{R}^3$, and $k \in \mathbb{Z}^3$. (Following [Fefferman 2006], however, we will not initially directly require any periodicity properties on the pressure.) As before, we will sometimes write a periodic solution (u, p, u_0, f, T) as (u, p, u_0, f, T, L) to emphasise the periodicity.

We will sometimes abuse notation and refer to a solution (u, p, u_0, f, T) simply as (u, p) or even u . Similarly, we will sometimes abbreviate a set of data (u_0, f, T) as (u_0, f) or even u_0 (in the homogeneous case $f = 0$).

Remark 1.2. In [Fefferman 2006], one considered³ smooth finite energy solutions associated to Schwartz data, as well as periodic smooth solutions associated to periodic smooth data. In the latter case, one can of course normalise the period L to equal 1 by a simple scaling argument. In this paper we will be focussed on the case when the data (u_0, f, T) is large, although we will not study the asymptotic regime when $T \rightarrow \infty$.

We recall the two standard *global regularity* conjectures for the Navier–Stokes equation, using the formulation in [Fefferman 2006]:

Conjecture 1.3 (Global regularity for homogeneous Schwartz data). Let $(u_0, 0, T)$ be a homogeneous Schwartz set of data. Then there exists a smooth finite energy solution $(u, p, u_0, 0, T)$ with the indicated data.

Conjecture 1.4 (Global regularity for homogeneous periodic data). Let $(u_0, 0, T)$ be a smooth homogeneous periodic set of data. Then there exists a smooth periodic solution $(u, p, u_0, 0, T)$ with the indicated data.

²Following [Fefferman 2006], we omit the finite energy dissipation condition $\nabla u \in L_t^2 L_x^2([0, T] \times \mathbb{R}^3)$ that often appears in the literature, particularly when discussing Leray–Hopf weak solutions. However, it turns out that this condition is actually automatic from (6) and smoothness; see Lemma 8.1. Similarly, from Corollary 11.1 we shall see that the $L_t^2 H_x^2$ condition in (7) is in fact redundant.

³The viscosity parameter ν was not normalised in [Fefferman 2006] to equal 1, as we are doing here, but one can easily reduce to the $\nu = 1$ case by a simple rescaling.

In view of these conjectures, one can naturally try to extend them to the inhomogeneous case as follows:

Conjecture 1.5 (Global regularity for Schwartz data). Let (u_0, f, T) be a Schwartz set of data. Then there exists a smooth finite energy solution (u, p, u_0, f, T) with the indicated data.

Conjecture 1.6 (Global regularity for periodic data). Let (u_0, f, T) be a smooth periodic set of data. Then there exists a smooth periodic solution (u, p, u_0, f, T) with the indicated data.

As described in [Fefferman 2006], a positive answer to either [Conjecture 1.3](#) or [Conjecture 1.4](#), or a negative answer to [Conjecture 1.5](#) or [Conjecture 1.6](#), would qualify for the Clay Millennium Prize.

However, [Conjecture 1.6](#) is not quite the “right” extension of [Conjecture 1.4](#) to the inhomogeneous setting, and needs to be corrected slightly. This is because there is a technical quirk in the inhomogeneous periodic problem as formulated in [Conjecture 1.6](#), due to the fact that the pressure p is not required to be periodic. This opens up a Galilean invariance in the problem which allows one to homogenise away the role of the forcing term. More precisely, we have:

Proposition 1.7 (Elimination of forcing term). *[Conjecture 1.6](#) is equivalent to [Conjecture 1.4](#).*

We establish this fact in [Section 6](#). We remark that this is the only implication we know of that can deduce a global regularity result for the inhomogeneous Navier–Stokes problem from a global regularity result for the homogeneous Navier–Stokes problem.

[Proposition 1.7](#) exploits the technical loophole of nonperiodic pressure. The same loophole can also be used to easily demonstrate failure of uniqueness for the periodic Navier–Stokes problem (although this can also be done by the much simpler expedient of noting that one can adjust the pressure by an arbitrary constant without affecting [\(3\)](#)). This suggests that in the nonhomogeneous case $f \neq 0$, one needs an additional normalisation to “fix” the periodic Navier–Stokes problem to avoid such loopholes. This can be done in a standard way, as follows. If one takes the divergence of [\(3\)](#) and uses the incompressibility [\(4\)](#), one sees that

$$\Delta p = -\partial_i \partial_j (u_i u_j) + \nabla \cdot f, \tag{8}$$

where we use the usual summation conventions. If (u, p, u_0, f, T) is a smooth periodic solution, then the right-hand side of [\(8\)](#) is smooth and periodic and has mean zero. From Fourier analysis, we see that given any smooth periodic mean-zero function F , there is a unique smooth periodic mean-zero function $\Delta^{-1} F$ with Laplacian equal to F . We then say that the periodic smooth solution (u, p, u_0, f, T) has *normalised pressure* if one has⁴

$$p = -\Delta^{-1} \partial_i \partial_j (u_i u_j) + \Delta^{-1} \nabla \cdot f. \tag{9}$$

We remark that this normalised pressure condition can also be imposed for smooth finite energy solutions (because $\partial_i \partial_j (u_i u_j)$ is a second derivative of an $L_x^1(\mathbb{R}^3)$ function, and $\nabla \cdot f$ is the first derivative of an $L_x^2(\mathbb{R}^3)$ function), but it will turn out that normalised pressure is essentially automatic in that setting anyway; see [Lemma 4.1](#).

⁴Up to the harmless freedom to add a constant to p , this normalisation is equivalent to requiring that the pressure be periodic with the same period as the solution u .

It is well known that once one imposes the normalised pressure condition, the periodic Navier–Stokes problem becomes locally well-posed in the smooth category (in particular, smooth solutions are now unique, and exist for sufficiently short times from any given smooth data); see [Theorem 5.1](#). Related to this, the Galilean invariance trick that allows one to artificially homogenise the forcing term f is no longer available. We can then pose a “repaired” version of [Conjecture 1.6](#):

Conjecture 1.8 (Global regularity for periodic data with normalised pressure). Let (u_0, f, T) be a smooth periodic set of data. Then there exists a smooth periodic solution (u, p, u_0, f, T) with the indicated data and with normalised pressure.

It is easy to see that the homogeneous case $f = 0$ of [Conjecture 1.8](#) is equivalent to [Conjecture 1.4](#); see, for example, [Lemma 4.1](#) below.

We now leave the category of classical (smooth) solutions for now, and turn instead to the category of *periodic H^1 mild solutions* (u, p, u_0, f, T, L) . By definition, these are functions

$$u, f : [0, T] \times \mathbb{R}^3 / L\mathbb{Z}^3 \rightarrow \mathbb{R}^3, \quad p : [0, T] \times \mathbb{R}^3 / L\mathbb{Z}^3 \rightarrow \mathbb{R}, \quad u_0 : \mathbb{R}^3 / L\mathbb{Z}^3 \rightarrow \mathbb{R}^3,$$

with $0 < T, L < \infty$, obeying the regularity hypotheses

$$\begin{aligned} u_0 &\in H_x^1(\mathbb{R}^3 / L\mathbb{Z}^3), \\ f &\in L_t^\infty H_x^1([0, T] \times (\mathbb{R}^3 / L\mathbb{Z}^3)), \\ u &\in L_t^\infty H_x^1 \cap L_t^2 H_x^2([0, T] \times (\mathbb{R}^3 / L\mathbb{Z}^3)), \end{aligned}$$

with p being given by [\(9\)](#), which obey the divergence-free conditions [\(4\)](#), [\(1\)](#) and obey the integral form

$$u(t) = e^{t\Delta} u_0 + \int_0^t e^{(t-t')\Delta} (-(u \cdot \nabla)u - \nabla p + f)(t') dt' \quad (10)$$

of the Navier–Stokes equation [\(3\)](#) with initial condition [\(5\)](#); using the Leray projection P onto divergence-free vector fields, we may also express [\(19\)](#) equivalently as

$$u(t) = e^{t\Delta} u_0 + \int_0^t e^{(t-t')\Delta} (PB(u, u) + Pf)(t') dt', \quad (11)$$

where $B(u, v)$ is the symmetric bilinear form

$$B(u, v)_i := -\frac{1}{2} \partial_j (u_i v_j + u_j v_i). \quad (12)$$

Similarly, we define *periodic H^1 data* to be a quadruplet (u_0, f, T, L) whose H^1 norm

$$\mathfrak{H}^1(u_0, f, T, L) := \|u_0\|_{H_x^1(\mathbb{R}^3 / L\mathbb{Z}^3)} + \|f\|_{L_t^\infty H_x^1(\mathbb{R}^3 / L\mathbb{Z}^3)}$$

is finite, with u_0 divergence-free.

Note from Duhamel’s formula [\(20\)](#) that every smooth periodic solution with normalised pressure is automatically a periodic H^1 mild solution.

As we will recall in [Theorem 5.1](#) below, the Navier–Stokes equation is locally well-posed in the periodic H^1 category. We can then formulate a global well-posedness conjecture in this category:

Conjecture 1.9 (Global well-posedness in periodic H^1). Let (u_0, f, T, L) be a periodic H^1 set of data. Then there exists a periodic H^1 mild solution (u, p, u_0, f, T, L) with the indicated data.

We may also phrase a quantitative variant of this conjecture:

Conjecture 1.10 (*A priori* periodic H^1 bound). There exists a function $F : \mathbb{R}^+ \times \mathbb{R}^+ \times \mathbb{R}^+ \rightarrow \mathbb{R}^+$ with the property that whenever (u, p, u_0, f, T, L) is a smooth periodic normalised-pressure solution with $0 < T < T_0 < \infty$ and

$$\mathcal{H}^1(u_0, f, T, L) \leq A < \infty,$$

we have

$$\|u\|_{L^\infty H_x^1([0, T] \times \mathbb{R}^3 / L\mathbb{Z}^3)} \leq F(A, L, T_0).$$

Remark 1.11. By rescaling, one may set $L = 1$ in this conjecture without any loss of generality; by partitioning the time interval $[0, T_0]$ into smaller subintervals, we may also simultaneously set $T_0 = 1$ if desired. Thus, the key point is that the size of the data A is allowed to be large (for small A the conjecture follows from the local well-posedness theory; see [Theorem 5.1](#)).

As we shall soon see, [Conjecture 1.9](#) and [Conjecture 1.10](#) are actually equivalent.

We now turn to the nonperiodic setting. In [Conjecture 1.5](#), the hypothesis that the initial data be Schwartz may seem unnecessarily restrictive, given that the incompressible nature of the fluid implies that the Schwartz property need not be preserved over time; also, there are many interesting examples of initial data that are smooth and finite energy (or H^1) but not Schwartz. In particular, one can consider generalising [Conjecture 1.5](#) to data that is merely smooth and H^1 , or even smooth and finite energy, rather than Schwartz⁵ of [Conjecture 1.5](#). Unfortunately, the naive generalisation of [Conjecture 1.5](#) (or even [Conjecture 1.3](#)) fails instantaneously in this case:

Theorem 1.12 (No smooth solutions from smooth H^1 data). *There exists smooth $u_0 \in H_x^1(\mathbb{R}^3)$ such that there does not exist any smooth finite energy solution $(u, p, u_0, 0, T)$ with the indicated data for any $T > 0$.*

We prove this proposition in [Section 15](#). At first glance, this proposition looks close to being a negative answer to either [Conjecture 1.5](#) or [Conjecture 1.3](#), but it relies on a technicality; for smooth H^1 data, the second derivatives of u_0 need not be square-integrable, and this can cause enough oscillation in the pressure to prevent the pressure from being C_t^2 (or the velocity field from being C_t^3) at the initial time⁶ $t = 0$. This theorem should be compared with the classical local existence theorem of Heywood [\[1980\]](#), which obtains smooth solutions for small *positive* times from smooth data with finite enstrophy, but merely obtains continuity at the initial time $t = 0$.

The situation is even worse in the inhomogeneous setting; the argument in [Theorem 1.12](#) can be used to construct inhomogeneous smooth H^1 data whose solutions will now be nonsmooth in time at all times,

⁵We are indebted to Andrea Bertozzi for suggesting these formulations of the Navier–Stokes global regularity problem.

⁶For most evolutionary PDEs, one can gain unlimited time differentiability at $t = 0$ assuming smooth initial data by differentiating the PDE in time (see the proof of the Cauchy–Kowalesky theorem). However, the problem here is that the pressure p in the Navier–Stokes equation does not obey an evolutionary PDE, but is instead determined in a nonlocal fashion from the initial data u (see [\(9\)](#)), which prevents one from obtaining much time regularity of the pressure initially.

not just at the initial time $t = 0$. Because of this, we will not attempt to formulate a global regularity problem in the inhomogeneous smooth H^1 or inhomogeneous smooth finite energy categories.

In the homogeneous setting, though, we can get around this technical obstruction by introducing the notion of an *almost smooth finite energy solution* (u, p, u_0, f, T) , which is the same concept as a smooth finite energy solution, but instead of requiring u, p to be smooth on $[0, T] \times \mathbb{R}^3$, we instead require that u, p are smooth on $(0, T] \times \mathbb{R}^3$, and for each $k \geq 0$, the functions $\nabla_x^k u, \partial_t \nabla_x^k u, \nabla_x^k p$ exist and are continuous on $[0, T] \times \mathbb{R}^3$. Thus, the only thing that almost smooth solutions lack when compared to smooth solutions is a limited amount of time differentiability at the starting time $t = 0$; informally, u is only $C_t^1 C_x^\infty$ at $t = 0$, and p is only $C_t^0 C_x^\infty$ at $t = 0$. This is still enough regularity to interpret the Navier–Stokes equation (3) in the classical sense, but is not a completely smooth solution.

The “corrected” conjectures for global regularity in the homogeneous smooth H^1 and smooth finite energy categories are then:

Conjecture 1.13 (Global almost regularity for homogeneous H^1). Let $(u_0, 0, T)$ be a smooth homogeneous H^1 set of data. Then there exists an almost smooth finite energy solution $(u, p, u_0, 0, T)$ with the indicated data.

Conjecture 1.14 (Global almost regularity for homogeneous finite energy data). Let $(u_0, 0, T)$ be a smooth homogeneous finite energy set of data. Then there exists an almost smooth finite energy solution $(u, p, u_0, 0, T)$ with the indicated data.

We carefully note that these conjectures only concern *existence* of smooth solutions, and not uniqueness; we will comment on some of the uniqueness issues later in this paper.

Another way to repair the global regularity conjectures in these settings is to abandon smoothness altogether, and work instead with the notion of mild solutions. More precisely, define a H^1 *mild solution* (u, p, u_0, f, T) to be fields $u, f : [0, T] \times \mathbb{R}^3 \rightarrow \mathbb{R}^3, p : [0, T] \times \mathbb{R}^3 \rightarrow \mathbb{R}, u_0 : \mathbb{R}^3 \rightarrow \mathbb{R}^3$ with $0 < T < \infty$, obeying the regularity hypotheses

$$\begin{aligned} u_0 &\in H_x^1(\mathbb{R}^3), \\ f &\in L_t^\infty H_x^1([0, T] \times \mathbb{R}^3), \\ u &\in L_t^\infty H_x^1 \cap L_t^2 H_x^2([0, T] \times \mathbb{R}^3), \end{aligned}$$

with p being given by (9), which obey (4), (1), and (10) (and thus (11)). Similarly, define the concept of H^1 data (u_0, f, T) .

We then have the following conjectures in the homogeneous setting:

Conjecture 1.15 (Global well-posedness in homogeneous H^1). Let $(u_0, 0, T)$ be a homogeneous H^1 set of data. Then there exists an H^1 mild solution $(u, p, u_0, 0, T)$ with the indicated data.

Conjecture 1.16 (*A priori* homogeneous H^1 bound). There exists a function $F : \mathbb{R}^+ \times \mathbb{R}^+ \rightarrow \mathbb{R}^+$ with the property that whenever $(u, p, u_0, 0, T)$ is a smooth H^1 solution with $0 < T < T_0 < \infty$ and

$$\|u_0\|_{H_x^1(\mathbb{R}^3)} \leq A < \infty,$$

we have

$$\|u\|_{L_t^\infty H_x^1([0,T] \times \mathbb{R}^3)} \leq F(A, T_0).$$

We also phrase a global-in-time variant:

Conjecture 1.17 (*A priori* global homogeneous H^1 bound). There exists a function $F : \mathbb{R}^+ \rightarrow \mathbb{R}^+$ with the property that whenever $(u, p, u_0, 0, T)$ is a smooth H^1 solution with

$$\|u_0\|_{H_x^1(\mathbb{R}^3)} \leq A < \infty,$$

then

$$\|u\|_{L_t^\infty H_x^1([0,T] \times \mathbb{R}^3)} \leq F(A).$$

In the inhomogeneous setting, we will state two slightly technical conjectures:

Conjecture 1.18 (Global well-posedness from spatially smooth Schwartz data). Let (u_0, f, T) be data obeying the bounds

$$\sup_{x \in \mathbb{R}^3} (1 + |x|)^k |\nabla_x^\alpha u_0(x)| < \infty$$

and

$$\sup_{(t,x) \in [0,T] \times \mathbb{R}^3} (1 + |x|)^k |\nabla_x^\alpha f(x)| < \infty$$

for all $k, \alpha \geq 0$. Then there exists an H^1 mild solution (u, p, u_0, f, T) with the indicated data.

Conjecture 1.19 (Global well-posedness from spatially smooth H^1 data). Let (u_0, f, T) be an H^1 set of data, such that

$$\sup_{x \in K} |\nabla_x^\alpha u_0(x)| < \infty$$

and

$$\sup_{(t,x) \in [0,T] \times K} |\nabla_x^\alpha f(x)| < \infty$$

for all $\alpha \geq 0$ and all compact K . Then there exists an H^1 mild solution (u, p, u_0, f, T) with the indicated data.

Needless to say, we do not establish⁷ any of these conjectures unconditionally in this paper. However, as the main result of this paper, we are able to establish the following implications:

Theorem 1.20 (Implications). (i) *Conjectures 1.9 and 1.10 are equivalent.*

(ii) *Conjecture 1.9 implies Conjecture 1.8 (and hence also Conjectures 1.6 and 1.4).*

(iii) *Conjecture 1.9 implies Conjecture 1.19, which is equivalent to Conjecture 1.18.*

(iv) *Conjecture 1.19 implies Conjectures 1.13 and 1.5 (and hence also Conjecture 1.3).*

(v) *Conjecture 1.13 is equivalent to Conjecture 1.14.*

(vi) *Conjectures 1.13, 1.15, 1.16, and 1.17 are all equivalent.*

⁷Indeed, the arguments here do not begin to address the main issue in any of these conjectures, namely the analysis of fine-scale (and turbulent) behaviour. The results in this paper do not prevent singularities from occurring in the Navier–Stokes flow; but they can largely localise the impact of such singularities to a bounded region of space.




Figure 1. Known implications between the various conjectures described here (existence of smooth or mild solutions, or local or global quantitative bounds in the periodic, Schwartz, H^1 , or finite energy categories, with or without normalised pressure, and with or without the $f = 0$ condition) and also in [Tao 2007] (the latter conjectures and implications occupy the far left column). A positive solution to the red problems, or a negative solution to the blue problems, qualify for the Clay Millennium prize, as stated in [Fefferman 2006].

The logical relationship between these conjectures, given by the implications above (as well as some trivial implications, and the equivalences in [Tao 2007]), is displayed in Figure 1.

Among other things, these results essentially show that in order to solve the Navier–Stokes global regularity problem, it suffices to study the periodic setting (but with the caveat that one now has to consider forcing terms with the regularity of $L_t^\infty H_x^1$).

Theorem 1.20(i) is a variant of the compactness arguments used in [Tao 2007] (see also [Gallagher 2001; Rusin and Šverák 2011]), and is proven in Section 7. Part (ii) of this theorem is a standard consequence of the periodic H^1 local well-posedness theory, which we review in Section 5. In the homogeneous $f = 0$ case it is possible to reverse this implication by the compactness arguments mentioned previously; see [Tao 2007]. However, we were unable to obtain this converse implication in the inhomogeneous case. Part (iv) is similarly a consequence of the nonperiodic H^1 local well-posedness theory, and is also proven in Section 5.

Part (vi) is also a variant of the results in [Tao 2007], with the main new ingredient being a use of concentration compactness instead of compactness in order to deal with the unboundedness of the spatial domain \mathbb{R}^3 , using the methods from [Bahouri and Gérard 1999; Gérard 1998; Gallagher 2001]. We establish these results in Section 14.

The more novel aspects of this theorem are parts (iii) and (v), which we establish in Sections 12 and 13 respectively. These results rely primarily on a new localised enstrophy inequality (Theorem 10.1) which can be viewed as a weak version of finite speed of propagation⁸ for the enstrophy $\frac{1}{2} \int_{\mathbb{R}^3} |\omega(t, x)|^2 dx$, where $\omega := \nabla \times u$ is the vorticity. We will also obtain a similar localised energy inequality for the energy $\frac{1}{2} \int_{\mathbb{R}^3} |u(t, x)|^2 dx$, but it will be the enstrophy inequality that is of primary importance to us, as the enstrophy is a subcritical quantity and can be used to obtain regularity (and local control on enstrophy can similarly be used to obtain local regularity). Remarkably, one is able to obtain local enstrophy inequalities even though the only *a priori* controlled quantity, namely the energy, is supercritical; the main difficulty is a harmonic analysis one, namely to control nonlinear effects primarily in terms of the local enstrophy and only secondarily in terms of the energy.

Remark 1.21. As one can see from Figure 1, the precise relationship between all the conjectures discussed here is rather complicated. However, if one is willing to ignore the distinction between homogeneous and inhomogeneous data, as well as the (rather technical) distinction between smooth and almost smooth solutions, then the main implications can then be informally summarised as follows:

- (Homogenisation) Without pressure normalisation, the inhomogeneity in the periodic global regularity conjecture is irrelevant: the inhomogeneous regularity conjecture is equivalent to the homogeneous one.
- (Localisation) The global regularity problem in the Schwartz, H^1 , and finite energy categories are “essentially” equivalent to each other.
- (More localisation) The global regularity problem in any of the above three categories is “essentially” a consequence of the global regularity problem in the periodic category.
- (Concentration compactness) Quantitative and qualitative versions of the global regularity problem (in a variety of categories) are “essentially” equivalent to each other.

The qualifier “essentially” here though needs to be taken with a grain of salt; again, one should consult Figure 1 for an accurate depiction of the implications.

The local enstrophy inequality has a number of other consequences, for instance allowing one to construct Leray–Hopf weak solutions whose (spatial) singularities are compactly supported in space; see Proposition 11.9.

Remark 1.22. Since the submission of this manuscript, the referee pointed out that the partial regularity theory of Caffarelli, Kohn, and Nirenberg [1982] also allows one to partially reverse the implication in

⁸Actually, in our setting, “finite distance of propagation” would be more accurate; we obtain an L_t^1 bound for the propagation velocity (see Proposition 9.1) rather than an L_t^∞ bound.

[Theorem 1.20\(iii\)](#), and more specifically to deduce [Conjecture 1.8](#) from [Conjecture 1.19](#). We sketch the referee’s argument in [Remark 12.3](#).

2. Notation and basic estimates

We use $X \lesssim Y$, $Y \gtrsim X$, or $X = O(Y)$ to denote the estimate $X \leq CY$ for an absolute constant C . If we need C to depend on a parameter, we shall indicate this by subscripts; thus for instance $X \lesssim_s Y$ denotes the estimate $X \leq C_s Y$ for some C_s depending on s . We use $X \sim Y$ as shorthand for $X \lesssim Y \lesssim X$.

We will occasionally use the Einstein summation conventions, using Roman indices i, j to range over the three spatial dimensions 1, 2, 3, though we will not bother to raise and lower these indices; for instance, the components of a vector field u will be u_i . We use ∂_i to denote the derivative with respect to the i -th spatial coordinate x_i . Unless otherwise specified, the Laplacian $\Delta = \partial_i \partial_i$ will denote the spatial Laplacian. (In [Lemma 12.1](#), though, we will briefly need to deal with the Laplace–Beltrami operator Δ_{S^2} on the sphere S^2 .) Similarly, ∇ will refer to the spatial gradient $\nabla = \nabla_x$ unless otherwise stated. We use the usual notations ∇f , $\nabla \cdot u$, $\nabla \times u$, for the gradient, divergence, or curl of a scalar field f or a vector field u .

It will be convenient (particularly when dealing with nonlinear error terms) to use *schematic notation*, in which an expression such as $\mathbb{O}(uvw)$ involving some vector- or tensor-valued quantities u, v, w denotes some constant-coefficient combination of products of the components of u, v, w respectively, and similarly for other expressions of this type. Thus, for instance, $\nabla \times \nabla \times u$ could be written schematically as $\mathbb{O}(\nabla^2 u)$, $|u \times v|^2$ could be written schematically as $\mathbb{O}(uuvv)$, and so forth.

For any centre $x_0 \in \mathbb{R}^3$ and radius $R > 0$, we use $B(x_0, R) := \{x \in \mathbb{R}^3 : |x - x_0| \leq R\}$ to denote the (closed) Euclidean ball. Much of our analysis will be localised to a ball $B(x_0, R)$, an annulus $B(x_0, R) \setminus B(x_0, r)$, or an exterior region $\mathbb{R}^3 \setminus B(x_0, R)$ (and often x_0 will be normalised to the origin 0).

We define the absolute value of a tensor in the usual Euclidean sense. Thus, for instance, if $u = u_i$ is a vector field, then $|u|^2 = u_i u_i$, $|\nabla u|^2 = (\partial_i u_j)(\partial_i u_j)$, $|\nabla^2 u|^2 = (\partial_i \partial_j u_k)(\partial_i \partial_j u_k)$, and so forth.

If E is a set, we use 1_E to denote the associated indicator function; thus $1_E(x) = 1$ when $x \in E$ and $1_E(x) = 0$ otherwise. We sometimes also use a statement in place of E ; thus for instance $1_{k \neq 0}$ would equal 1 if $k \neq 0$ and 0 when $k = 0$.

We use the usual Lebesgue spaces $L^p(\Omega)$ for various domains Ω (usually subsets of Euclidean space \mathbb{R}^3 or a torus $\mathbb{R}^3/\mathbb{Z}^3$) and various exponents $1 \leq p \leq \infty$, which will always be equipped with an obvious Lebesgue measure. We often write $L^p(\Omega)$ as $L_x^p(\Omega)$ to emphasise the spatial nature of the domain Ω . Given an absolutely integrable function $f \in L_x^1(\mathbb{R}^3)$, we define the Fourier transform $\hat{f} : \mathbb{R}^3 \rightarrow \mathbb{C}$ by the formula

$$\hat{f}(\xi) := \int_{\mathbb{R}^3} e^{-2\pi i x \cdot \xi} f(x) dx;$$

we then extend this Fourier transform to tempered distributions in the usual manner. For a function f which is periodic with period 1, and thus representable as a function on the torus $\mathbb{R}^3/\mathbb{Z}^3$, we define the

discrete Fourier transform $\hat{f} : \mathbb{Z}^3 \rightarrow \mathbb{C}$ by the formula

$$\hat{f}(k) := \int_{\mathbb{R}^3/\mathbb{Z}^3} e^{-2\pi i k \cdot x} f(x) dx$$

when f is absolutely integrable on $\mathbb{R}^3/\mathbb{Z}^3$, and extend this to more general distributions on $\mathbb{R}^3/\mathbb{Z}^3$ in the usual fashion. Strictly speaking, these two notations are not compatible with each other, but it will always be clear in context whether we are using the nonperiodic or the periodic Fourier transform.

For any spatial domain Ω (contained in either \mathbb{R}^3 or $\mathbb{R}^3/L\mathbb{Z}^3$) and any natural number $k \geq 0$, we define the classical Sobolev norms $\|u\|_{H_x^k(\Omega)}$ of a smooth function $u : \Omega \rightarrow \mathbb{R}$ by the formula

$$\|u\|_{H_x^k(\Omega)} := \left(\sum_{j=0}^k \|\nabla^j u\|_{L_x^2(\Omega)}^2 \right)^{1/2},$$

and say that $u \in H_x^k(\Omega)$ when $\|u\|_{H_x^k(\Omega)}$ is finite. Note that we do not impose any vanishing conditions at the boundary of Ω , and to avoid technical issues we will not attempt to define these norms for nonsmooth functions u in the event that Ω has a nontrivial boundary. In the domain \mathbb{R}^3 and for $s \in \mathbb{R}$, we define the Sobolev norm $\|u\|_{H_x^s(\mathbb{R}^3)}$ of a tempered distribution $u : \mathbb{R}^3 \rightarrow \mathbb{R}$ by the formula

$$\|u\|_{H_x^s(\mathbb{R}^3)} := \left(\int_{\mathbb{R}^3} (1 + |\xi|^2)^s |\hat{u}(\xi)|^2 d\xi \right)^{1/2}.$$

Strictly speaking, this conflicts slightly with the previous notation when k is a nonnegative integer, but the two norms are equivalent up to constants (and both norms define a Hilbert space structure), so the distinction will not be relevant for our purposes. For $s > -\frac{3}{2}$, we also define the homogeneous Sobolev norm

$$\|u\|_{\dot{H}_x^s(\mathbb{R}^3)} := \left(\int_{\mathbb{R}^3} |\xi|^{2s} |\hat{u}(\xi)|^2 d\xi \right)^{1/2},$$

and let $H_x^s(\mathbb{R}^3)$, $\dot{H}_x^s(\mathbb{R}^3)$ be the space of tempered distributions with finite $H_x^s(\mathbb{R}^3)$ or $\dot{H}_x^s(\mathbb{R}^3)$ norm respectively. Similarly, on the torus $\mathbb{R}^3/\mathbb{Z}^3$ and $s \in \mathbb{R}$, we define the Sobolev norm $\|u\|_{H_x^s(\mathbb{R}^3/\mathbb{Z}^3)}$ of a distribution $u : \mathbb{R}^3/\mathbb{Z}^3 \rightarrow \mathbb{R}$ by the formula

$$\|u\|_{H_x^s(\mathbb{R}^3/\mathbb{Z}^3)} := \left(\sum_{k \in \mathbb{Z}^3} (1 + |k|^2)^s |\hat{u}(k)|^2 \right)^{1/2};$$

again, this conflicts slightly with the classical Sobolev norms $H_x^k(\mathbb{R}^3/\mathbb{Z}^3)$, but this will not be a serious issue in this paper. We define $H_x^s(\mathbb{R}^3/\mathbb{Z}^3)$ to be the space of all distributions u with finite $H_x^s(\mathbb{R}^3/\mathbb{Z}^3)$ norm, and $H_x^s(\mathbb{R}^3/\mathbb{Z}^3)_0$ to be the codimension-one subspace of functions or distributions u which are mean-zero in the sense that $\hat{u}(0) = 0$.

In a similar vein, given a spatial domain Ω and a natural number $k \geq 0$, we define $C_x^k(\Omega)$ to be the space of all k times continuously differentiable functions $u : \Omega \rightarrow \mathbb{R}$ whose norm

$$\|u\|_{C_x^k(\Omega)} := \sum_{j=0}^k \|\nabla^j u\|_{L_x^\infty(\Omega)}$$

is finite⁹. Given any spatial norm $\|\cdot\|_{X_x(\Omega)}$ associated to a function space X_x defined on a spatial domain Ω , and a time interval I , we can define mixed-norms $\|u\|_{L_t^p X_x(I \times \Omega)}$ on functions $u : I \times \Omega \rightarrow \mathbb{R}$ by the formula

$$\|u\|_{L_t^p X_x(I \times \Omega)} := \left(\int_I \|u(t)\|_{X_x(\Omega)}^p dt \right)^{1/p}$$

when $1 \leq p < \infty$, and

$$\|u\|_{L_t^\infty X_x(I \times \Omega)} := \text{ess sup}_{t \in I} \|u(t)\|_{X_x(\Omega)},$$

assuming in both cases that $u(t)$ lies in $X(\Omega)$ for almost every Ω , and then let $L_t^p X_x(I \times \Omega)$ be the space of functions (or, in some cases, distributions) whose $L_t^p X_x(I \times \Omega)$ is finite. Thus, for instance, $L_t^\infty C_x^2(I \times \Omega)$ would be the space of functions $u : I \times \Omega \rightarrow \mathbb{R}$ such that for almost every $x \in I$, $u(t) : \Omega \rightarrow \mathbb{R}$ is in $C_x^2(\Omega)$, and the norm

$$\|u\|_{L_t^\infty C_x^2(I \times \Omega)} := \text{ess sup}_{t \in I} \|u(t)\|_{C_x^2(\Omega)}$$

is finite.

Similarly, for any natural number $k \geq 0$, we define $C_t^k X_x(I \times \Omega)$ to be the space of all functions $u : I \times \Omega \rightarrow \mathbb{R}$ such that the curve $t \mapsto u(t)$ from I to $X_x(\Omega)$ is k times continuously differentiable, and such that the norm

$$\|u\|_{C_t^k X_x(I \times \Omega)} := \sum_{j=0}^k \|\nabla^j u\|_{L_t^\infty X_x(I \times \Omega)}$$

is finite.

Given two normed function spaces X, Y on the same domain (in either space or space-time), we can endow their intersection $X \cap Y$ with the norm

$$\|u\|_{X \cap Y} := \|u\|_X + \|u\|_Y.$$

For us, the most common example of such hybrid norms will be the spaces

$$X^s(I \times \Omega) := L_t^\infty H_x^s(I \times \Omega) \cap L_x^2 H_x^{s+1}(I \times \Omega), \quad (13)$$

defined whenever I is a time interval, s is a natural number, and Ω is a spatial domain, or whenever I is a time interval, s is real, and Ω is either \mathbb{R}^3 or $\mathbb{R}^3/\mathbb{Z}^3$. The X^s spaces (particularly X^1) will play a prominent role in the (subcritical) local well-posedness theory for the Navier–Stokes equations; see

⁹Note that if Ω is noncompact, then it is possible for a smooth function to fail to lie in $C^k(\Omega)$ if it becomes unbounded or excessively oscillatory at infinity. One could use a notation such as $C_{x,\text{loc}}^k(\Omega)$ to describe the space of functions that are k times continuously differentiable with no bounds on derivatives, but we will not need such notation here.

Section 5. The space X^0 will also be naturally associated with energy estimates, and the space X^1 with enstrophy estimates.

All of these above function spaces can of course be extended to functions that are vector or tensor-valued without difficulty (there are multiple ways to define the norms in these cases, but all such definitions will be equivalent up to constants).

We use the Fourier transform to define a number of useful multipliers on \mathbb{R}^3 or $\mathbb{R}^3/\mathbb{Z}^3$. On \mathbb{R}^3 , we formally define the inverse Laplacian operator Δ^{-1} by the formula

$$\widehat{\Delta^{-1}f}(\xi) := \frac{-1}{4\pi^2|\xi|^2} \hat{f}(\xi), \quad (14)$$

which is well-defined for any tempered distribution $f : \mathbb{R}^3 \rightarrow \mathbb{R}$ for which the right-hand side of (14) is locally integrable. This is for instance the case if f lies in the k -th derivative of a function in $L_x^1(\mathbb{R}^3)$ for some $k \geq 0$, or the k -th derivative of a function in $L_x^2(\mathbb{R}^3)$ for some $k \geq 1$. If $f \in L_x^1(\mathbb{R}^3)$, then as is well known, one has the Newton potential representation

$$\Delta^{-1}f(x) = \frac{-1}{4\pi} \int_{\mathbb{R}^3} \frac{f(y)}{|x-y|} dy. \quad (15)$$

Note in particular that (15) implies that if $f \in L_x^1(\mathbb{R}^3)$ is supported on some closed set K , then $\Delta^{-1}f$ will be smooth away from K . Also observe from Fourier analysis (and decomposition into local and global components) that if f is smooth and is either the k -th derivative of a function in $L_x^1(\mathbb{R}^3)$ for some $k \geq 0$, or the k -th derivative of a function in $L_x^2(\mathbb{R}^3)$ for some $k \geq 1$, then $\Delta^{-1}f$ will be smooth also.

We also note that the Newton potential $-1/(4\pi|x-y|)$ is smooth away from the diagonal $x=y$. Because of this, we will often be able to obtain large amounts of regularity in space in the “far field” region when $|x|$ is large, for fields such as the velocity field u . However, it will often be significantly more challenging to gain significant amounts of regularity in *time*, because the inverse Laplacian Δ^{-1} has no smoothing properties in the time variable.

On $\mathbb{R}^3/\mathbb{Z}^3$, we similarly define the inverse Laplacian operator Δ^{-1} for distributions $f : \mathbb{R}^3/\mathbb{Z}^3 \rightarrow \mathbb{R}$ with $\hat{f}(0) = 0$ by the formula

$$\widehat{\Delta^{-1}f}(k) := \frac{-1_{k \neq 0}}{4\pi^2|k|^2} \hat{f}(k). \quad (16)$$

We define the *Leray projection* Pu of a (tempered distributional) vector field $u : \mathbb{R}^3 \rightarrow \mathbb{R}^3$ by the formula

$$Pu := \Delta^{-1}(\nabla \times \nabla \times u).$$

If u is square-integrable, then Pu is the orthogonal projection of u onto the space of square-integrable divergence-free vector fields; from Calderón–Zygmund theory, we know that the projection P is bounded on $L_x^p(\mathbb{R}^3)$ for every $1 < p < \infty$, and from Fourier analysis we see that P is also $H_x^s(\mathbb{R}^3)$ for every $s \in \mathbb{R}$. Note that if u is square-integrable and divergence-free, then $Pu = u$, and we thus have the *Biot–Savart law*

$$u = \Delta^{-1}(\nabla \times \omega), \quad (17)$$

where $\omega := \nabla \times u$.

In either \mathbb{R}^3 or $\mathbb{R}^3/L\mathbb{Z}^3$, we let $e^{t\Delta}$ for $t > 0$ be the usual heat semigroup associated to the heat equation $u_t = \Delta u$. On \mathbb{R}^3 , this takes the explicit form

$$e^{t\Delta} f(x) = \frac{1}{(4\pi t)^{3/2}} \int_{\mathbb{R}^3} e^{-|x-y|^2/4t} f(y) dy$$

for $f \in L_x^p(\mathbb{R}^3)$ for some $1 \leq p \leq \infty$. From Young's inequality, we thus record the dispersive inequality

$$\|e^{t\Delta} f\|_{L^q(\mathbb{R}^3)} \lesssim t^{3/2q-3/2p} \|f\|_{L^p(\mathbb{R}^3)} \quad (18)$$

whenever $1 \leq p \leq q \leq \infty$ and $t > 0$.

We recall *Duhamel's formula*

$$u(t) = e^{(t-t_0)\Delta} u(t_0) + \int_{t_0}^t e^{(t-t')\Delta} (\partial_t u - \Delta u)(t') dt' \quad (19)$$

whenever $u : [t_0, t] \times \Omega \rightarrow \mathbb{R}$ is a smooth tempered distribution, with Ω equal to either \mathbb{R}^3 or $\mathbb{R}^3/\mathbb{Z}^3$.

We record some linear and bilinear estimates involving Duhamel-type integrals and the spaces X^s defined in (13), which are useful in the local H^1 theory for the Navier–Stokes equation:

Lemma 2.1 (Linear and bilinear estimates). *Let $[t_0, t_1]$ be a time interval, let Ω be either \mathbb{R}^3 or $\mathbb{R}^3/\mathbb{Z}^3$, and suppose that $u : [t_0, t_1] \times \Omega \rightarrow \mathbb{R}$ and $F : [t_0, t_1] \times \Omega \rightarrow \mathbb{R}$ are tempered distributions such that*

$$u(t) = e^{(t-t_0)\Delta} u(t_0) + \int_{t_0}^t e^{(t-t')\Delta} F(t') dt'. \quad (20)$$

Then we have the standard energy estimate¹⁰

$$\|u\|_{X^s([t_0, t_1] \times \Omega)} \lesssim_s \|u(t_0)\|_{H_x^s(\Omega)} + \|F\|_{L_t^1 H_x^s([t_0, t_1] \times \Omega)} \quad (21)$$

for any $s \geq 0$, as well as the variant

$$\|u\|_{X^s([t_0, t_1] \times \Omega)} \lesssim_s \|u(t_0)\|_{H_x^s(\Omega)} + \|F\|_{L_t^2 H_x^{s-1}([t_0, t_1] \times \Omega)} \quad (22)$$

for any $s \geq 1$. We also note the further variant

$$\|u\|_{X^s([t_0, t_1] \times \Omega)} \lesssim_s \|u(t_0)\|_{H_x^s(\Omega)} + \|F\|_{L_t^4 L_x^2([t_0, t_1] \times \Omega)} \quad (23)$$

for any $s < 3/2$.

We also have the bilinear estimate

$$\|\nabla(uv)\|_{L_t^4 L_x^2([t_0, t_1] \times \Omega)} \lesssim \|u\|_{X^1([t_0, t_1] \times \Omega)} \|v\|_{X^1([t_0, t_1] \times \Omega)} \quad (24)$$

for any $u, v : [t_0, t_1] \times \mathbb{R}^3 \rightarrow \mathbb{R}$, which in particular implies (by a Hölder in time) that

$$\|\nabla(uv)\|_{L_t^2 L_x^2([t_0, t_1] \times \mathbb{R}^3)} \lesssim (t_1 - t_0)^{1/4} \|u\|_{X^1([t_0, t_1] \times \mathbb{R}^3)} \|v\|_{X^1([t_0, t_1] \times \mathbb{R}^3)}. \quad (25)$$

¹⁰We adopt the convention that an estimate is vacuously true if the right-hand side is infinite or undefined.

Proof. The estimates¹¹ (22), (23), (24) are established in [Tao 2007, Lemma 2.1, Proposition 2.2]. The estimate (21) follows from the $F = 0$ case of (21) and Minkowski's inequality. \square

Finally, we define the Littlewood–Paley projection operators on \mathbb{R}^3 . Let $\varphi(\xi)$ be a fixed bump function supported in the ball $\{\xi \in \mathbb{R}^3 : |\xi| \leq 2\}$ and equal to 1 on the ball $\{\xi \in \mathbb{R}^3 : |\xi| \leq 1\}$. Define a *dyadic number* to be a number N of the form $N = 2^k$ for some integer k . For each dyadic number N , we define the Fourier multipliers

$$\begin{aligned}\widehat{P_{\leq N} f}(\xi) &:= \varphi(\xi/N) \widehat{f}(\xi), \\ \widehat{P_{> N} f}(\xi) &:= (1 - \varphi(\xi/N)) \widehat{f}(\xi), \\ \widehat{P_N f}(\xi) &:= \psi(\xi/N) \widehat{f}(\xi) := (\varphi(\xi/N) - \varphi(2\xi/N)) \widehat{f}(\xi).\end{aligned}$$

We similarly define $P_{< N}$ and $P_{\geq N}$. Thus for any tempered distribution, we have $f = \sum_N P_N f$ in a weakly convergent sense at least, where the sum ranges over dyadic numbers. We recall the usual *Bernstein estimates*

$$\begin{aligned}\|D^s P_N f\|_{L_x^p(\mathbb{R}^3)} &\lesssim_{p,s,D^s} N^s \|P_N f\|_{L_x^p(\mathbb{R}^3)}, \\ \|\nabla^k P_N f\|_{L_x^p(\mathbb{R}^3)} &\sim_{k,s} N^k \|P_N f\|_{L_x^p(\mathbb{R}^3)}, \\ \|P_{\leq N} f\|_{L_x^q(\mathbb{R}^3)} &\lesssim_{p,q} N^{3/p-3/q} \|P_{\leq N} f\|_{L_x^p(\mathbb{R}^3)}, \\ \|P_N f\|_{L_x^q(\mathbb{R}^3)} &\lesssim_{p,q} N^{3/p-3/q} \|P_N f\|_{L_x^p(\mathbb{R}^3)},\end{aligned}\tag{26}$$

for all $1 \leq p \leq q \leq \infty$, $s \in \mathbb{R}$, $k \geq 0$, and pseudodifferential operators D^s of order s ; see, for example, [Tao 2006, Appendix A].

We recall the *Littlewood–Paley trichotomy*: an expression of the form $P_N((P_{N_1} f_1)(P_{N_2} f_2))$ vanishes unless one of the following three scenarios holds:

- (Low-high interaction) $N_2 \lesssim N_1 \sim N$.
- (High-low interaction) $N_1 \lesssim N_2 \sim N$.
- (High-high interaction) $N \lesssim N_1 \sim N_2$.

This trichotomy is useful for obtaining estimates on bilinear expressions, as we shall see in Section 9.

We have the following frequency-localised variant of (18):

Lemma 2.2. *If N is a dyadic number and $f : \mathbb{R}^3 \rightarrow \mathbb{R}$ has Fourier transform supported on an annulus $\{\xi : |\xi| \sim N\}$, then we have*

$$\|e^{t\Delta} f\|_{L^q(\mathbb{R}^3)} \lesssim t^{3/2q-3/2p} \exp(-ctN^2) \|f\|_{L^p(\mathbb{R}^3)}\tag{27}$$

for some absolute constant $c > 0$ and all $1 \leq p \leq q \leq \infty$.

¹¹Strictly speaking, the result in [Tao 2007] was stated for the torus rather than \mathbb{R}^3 , but the argument works without modification in either domain, after first truncating $u(t_0)$, F to be Schwartz to avoid technicalities at infinity, and using a standard density argument.

Proof. By Littlewood–Paley projection, it suffices to show that

$$\|e^{t\Delta} P_N f\|_{L^q(\mathbb{R}^3)} \lesssim t^{3/2q-3/2p} \exp(-ctN^2) \|f\|_{L^p(\mathbb{R}^3)}$$

for all test functions f . By rescaling, we may set $t = 1$; in view of (18) we may then set $N \geq 1$. One then verifies from Fourier analysis that $e^{t\Delta} P_N$ is a convolution operator whose kernel has an $L_x^\infty(\mathbb{R}^3)$ and an $L_x^1(\mathbb{R}^3)$ norm that are both $O(\exp(-cN^2))$ for some absolute constant $c > 0$, and the claim follows from Young’s inequality. \square

From the uniform smoothness of the heat kernel, we also observe the estimate

$$\|e^{t\Delta} f\|_{C_x^k(K)} \lesssim_{k,K,T,p} \exp(-cTr^2) \|f\|_{L_x^p(\mathbb{R}^3)} \quad (28)$$

whenever $0 \leq t \leq T$, $1 \leq p \leq \infty$, $k \geq 0$, K is a compact subset of \mathbb{R}^3 , $r \geq 1$, f is supported on the set $\{x \in \mathbb{R}^3 : \text{dist}(x, K) \geq r\}$, and some quantity $c_T > 0$ depending only on T . In practice, this estimate will be an effective substitute for finite speed of propagation for the heat equation.

3. Symmetries of the equation

In this section we review some well known symmetries of the Navier–Stokes flow that transform a given smooth solution (u, p, u_0, f, T) to another smooth solution $(\tilde{u}, \tilde{p}, \tilde{u}_0, \tilde{f}, \tilde{T})$, as these symmetries will be useful at various points in the paper.

The simplest symmetry is the *spatial translation symmetry*

$$\begin{aligned} \tilde{u}(t, x) &:= u(t, x - x_0), \\ \tilde{p}(t, x) &:= p(t, x - x_0), \\ \tilde{u}_0(x) &:= u_0(x - x_0), \\ \tilde{f}(t, x) &:= f(t, x - x_0), \\ \tilde{T} &:= T, \end{aligned} \quad (29)$$

valid for any $x_0 \in \mathbb{R}^3$; this transformation clearly maps mild, smooth, or almost smooth solutions to solutions of the same type, and also preserves conditions such as finite energy, H^1 , periodicity, pressure normalisation, or the Schwartz property. In a similar vein, we have the *time translation symmetry*

$$\begin{aligned} \tilde{u}(t, x) &:= u(t + t_0, x), \\ \tilde{p}(t, x) &:= p(t + t_0, x), \\ \tilde{u}_0(x) &:= u(t_0, x), \\ \tilde{f}(t, x) &:= f(t + t_0, x), \\ \tilde{T} &:= T - t_0, \end{aligned} \quad (30)$$

valid for any $t_0 \in [0, T]$. Again, this maps mild, smooth, or almost smooth solutions to solutions of the same type (and if $t_0 > 0$, then almost smooth solutions are even upgraded to smooth solutions). If the original solution is finite energy or H^1 , then the transformed solution will be finite energy or H^1

also. Note however that if it is only the original *data* that is assumed to be finite energy or H^1 , as opposed to the *solution*, it is not immediately obvious that the time-translated solution remains finite energy or H^1 , especially in view of the fact that the H^1 norm (or the enstrophy) is not a conserved quantity of the Navier–Stokes flow. (See however [Lemma 8.1](#) and [Corollary 11.1](#) below.) The situation is particularly dramatic in the case of Schwartz data; as remarked earlier, time translation can instantly convert¹² Schwartz data to non-Schwartz data, due to the slow decay of the Newton potential appearing in [\(9\)](#) (or of its derivatives, such as the Biot–Savart kernel in [\(17\)](#)).

Next, we record the *scaling symmetry*

$$\begin{aligned}\tilde{u}(t, x) &:= \frac{1}{\lambda} u\left(\frac{t}{\lambda^2}, \frac{x}{\lambda}\right), \\ \tilde{p}(t, x) &:= \frac{1}{\lambda^2} p\left(\frac{t}{\lambda^2}, \frac{x}{\lambda}\right), \\ \tilde{u}_0(x) &:= \frac{1}{\lambda} u\left(\frac{x}{\lambda}\right), \\ \tilde{f}(t, x) &:= \frac{1}{\lambda^3} f\left(\frac{t}{\lambda^2}, \frac{x}{\lambda}\right), \\ \tilde{T} &:= T\lambda^2,\end{aligned}\tag{31}$$

valid for any $\lambda > 0$; it also maps mild, smooth, or almost smooth solutions to solutions of the same type, and preserves properties such as finite energy, finite enstrophy, pressure normalisation, periodicity, or the Schwartz property, though note in the case of periodicity that a solution of period L will map to a solution of period λL . We will only use scaling symmetry occasionally in this paper, mainly because most of the quantities we will be manipulating will be supercritical with respect to this symmetry. Nevertheless, this scaling symmetry serves a fundamentally important conceptual purpose, by making the key distinction between subcritical, critical (or dimensionless), and supercritical quantities, which can help illuminate many of the results in this paper (and was also crucial in allowing the author to discover¹³ these results in the first place).

We record three further symmetries that impact upon the issue of pressure normalisation. The first is the *pressure shifting symmetry*

$$\begin{aligned}\tilde{u}(t, x) &:= u(t, x), \\ \tilde{p}(t, x) &:= p(t, x) + C(t), \\ \tilde{u}_0(x) &:= u_0(x), \\ \tilde{f}(t, x) &:= f(t, x), \\ \tilde{T} &:= T,\end{aligned}\tag{32}$$

¹²This can be seen for instance by noting that moments such as $\int_{\mathbb{R}^3} \omega_1(t, x)(x_2^2 - x_3^2) dx$ are not conserved in time, but must equal zero whenever $u(t)$ is Schwartz.

¹³The author also found dimensional analysis to be invaluable in checking the calculations for errors. One *could*, if one wished, exploit the scaling symmetry to normalise a key parameter (for example, the energy E , or a radius parameter r) to equal one, which would simplify the numerology slightly, but then one would lose the use of dimensional analysis to check for errors, and so we have elected to largely avoid the use of scaling normalisations in this paper.

valid for any smooth function $C : \mathbb{R} \rightarrow \mathbb{R}$. This clearly maps smooth or almost smooth solutions to solutions of the same type, and preserves properties such as finite energy, H^1 , periodicity, and the Schwartz property; however, it destroys pressure normalisation (and thus the notion of a mild solution). A slightly more sophisticated symmetry in the same spirit is the *Galilean symmetry*

$$\begin{aligned}
 \tilde{u}(t, x) &:= u\left(t, x - \int_0^t v(s) ds\right) + v(t), \\
 \tilde{p}(t, x) &:= p\left(t, x - \int_0^t v(s) ds\right) - x \cdot v'(t), \\
 \tilde{u}_0(x) &:= u_0(x) + v(0), \\
 \tilde{f}(t, x) &:= f\left(t, x - \int_0^t v(s) ds\right), \\
 \tilde{T} &:= T,
 \end{aligned} \tag{33}$$

valid for any smooth function $v : \mathbb{R} \rightarrow \mathbb{R}^3$. One can carefully check that this symmetry indeed maps mild, smooth solutions to smooth solutions and preserves periodicity (recall here that in our definition of a periodic solution, the pressure was *not* required to be periodic). On the other hand, this symmetry does not preserve finite energy, H^1 , or the Schwartz property. It also clearly destroys the pressure normalisation property.

Finally, we observe that one can absorb divergences into the forcing term via the *forcing symmetry*

$$\begin{aligned}
 \tilde{u}(t, x) &:= u(t, x), \\
 \tilde{p}(t, x) &:= p(t, x) + q(t, x), \\
 \tilde{u}_0(x) &:= u_0(x), \\
 \tilde{f}(t, x) &:= f(t, x) + \nabla \cdot q(t, x), \\
 \tilde{T} &:= T,
 \end{aligned} \tag{34}$$

valid for any smooth function $P : [0, T] \times \mathbb{R}^3 \rightarrow \mathbb{R}^3$. If the new forcing term \tilde{f} still has finite energy or is still periodic, then the normalisation of pressure is preserved. In the periodic setting, we will apply (34) with a linear term $q(t, x) := x \cdot a(t)$, allowing one to alter f by an arbitrary constant $a(t)$. In the finite energy or H^1 setting, one can use (34) and the Leray projection P to reduce to the divergence-free case $\nabla \cdot f = 0$; note, though, that this projection can destroy the Schwartz nature of f . This divergence-free reduction is particularly useful in the case of normalised pressure, since (9) then simplifies to

$$p = -\Delta^{-1} \partial_i \partial_j (u_i u_j). \tag{35}$$

One can of course compose these symmetries together to obtain a larger (semi)group of symmetries. For instance, by combining (33) and (34), we observe the symmetry

$$\begin{aligned}
\tilde{u}(t, x) &:= u\left(t, x - \int_0^t v(s) ds\right) + v(t), \\
\tilde{p}(t, x) &:= p\left(t, x - \int_0^t v(s) ds\right), \\
\tilde{u}_0(x) &:= u_0(x) + v(0), \\
\tilde{f}(t, x) &:= f\left(t, x - \int_0^t v(s) ds\right) + v'(t), \\
\tilde{T} &:= T,
\end{aligned} \tag{36}$$

for any smooth function $v : \mathbb{R} \rightarrow \mathbb{R}^3$. This symmetry is particularly useful for periodic solutions; note that it preserves both the periodicity property and the normalised pressure property. By choosing $v(t)$ appropriately, we see that we can use this symmetry to normalise periodic data (u_0, f, T, L) to be *mean-zero* in the sense that

$$\int_{\mathbb{R}^3/L\mathbb{Z}^3} u_0(x) dx = 0 \tag{37}$$

and

$$\int_{\mathbb{R}^3/L\mathbb{Z}^3} f(t, x) dx = 0 \tag{38}$$

for all $0 \leq t \leq T$. By integrating (3) over the torus $\mathbb{R}^3/L\mathbb{Z}^3$, we then conclude with this normalisation that u remains mean-zero for all times $0 \leq t \leq T$:

$$\int_{\mathbb{R}^3/L\mathbb{Z}^3} u(t, x) dx = 0. \tag{39}$$

The same conclusion also holds for periodic H^1 mild solutions.

4. Pressure normalisation

The symmetries in (32), (34) can alter the velocity field u and pressure p without affecting the data (u_0, f, T) , thus leading to a breakdown of uniqueness for the Navier–Stokes equation. In this section we investigate this loss of uniqueness, and show that (in the smooth category, at least) one can “quotient out” these symmetries by reducing to the situation (9) of normalised pressure, at which point uniqueness can be recovered (at least in the H^1 category).

More precisely, we show:

Lemma 4.1 (Reduction to normalised pressure). (i) *If (u, p, u_0, f, T) is an almost smooth finite energy solution, then for almost every time $t \in [0, T]$, one has*

$$p(t, x) = -\Delta^{-1} \partial_i \partial_j (u_i u_j)(t, x) + \Delta^{-1} \nabla \cdot f(t, x) + C(t), \tag{40}$$

for some bounded measurable function $C : [0, T] \rightarrow \mathbb{R}$.

(ii) If (u, p, u_0, f, T) is a periodic smooth solution, then there exist smooth functions $C : [0, T] \rightarrow \mathbb{R}$ and $a : [0, T] \rightarrow \mathbb{R}^3$ such that

$$p(t, x) = -\Delta^{-1} \partial_i \partial_j (u_i u_j)(t, x) + \Delta^{-1} \nabla \cdot f(t, x) + x \cdot a(t) + C(t). \quad (41)$$

In particular, after applying a Galilean transformation (33) followed by a pressure-shifting transformation (32), one can transform (u, p, u_0, f, T) into a periodic smooth solution with normalised pressure.

Remark 4.2. Morally, in (i) the function C should be smooth (at least for times $t > 0$), which would then imply that one can apply a pressure-shifting transformation (32) to convert (u, p, u_0, f, T) into a smooth solution with normalised pressure. However, there is the technical difficulty that in our definition of a finite energy smooth solution, we do not *a priori* have any control of time derivatives of u in any $L_x^p(\mathbb{R}^3)$ norms, and as such we do not have time regularity on the component $\Delta^{-1} \partial_i \partial_j (u_i u_j)$ of (40). In practice, though, this possible irregularity of $C(t)$ will not bother us, as we only need to understand the gradient ∇p of the pressure, rather than the pressure itself, in order to solve the Navier–Stokes equations (3).

Proof. We begin with the periodic case, which is particularly easy due to Liouville’s theorem (which, among other things, implies that the only harmonic periodic functions are the constants). We may normalise the period L to equal 1. Fix an almost smooth periodic solution (u, p, u_0, f, T) . Define the normalised pressure $p_0 : [0, T] \times \mathbb{R}^3 \rightarrow \mathbb{R}$ by the formula

$$p_0 := -\Delta^{-1} \partial_i \partial_j (u_i u_j) + \Delta^{-1} \nabla \cdot f. \quad (42)$$

As u, f are smooth and periodic, p_0 is smooth also, and from (8) one has $\Delta p = \Delta p_0$. Thus one has

$$p = p_0 + h,$$

where $h : [0, T] \times \mathbb{R}^3 \rightarrow \mathbb{R}$ is a smooth function with $h(t)$ harmonic in space for each time t . The function h need not be periodic; however, from (3) we have

$$\partial_t u + (u \cdot \nabla) u = \Delta u - \nabla p_0 - \nabla h + f.$$

Every term aside from ∇h is periodic, and so ∇h is periodic also. Since ∇h is also harmonic, it must therefore be constant in space by Liouville’s theorem. We therefore may write

$$h(t, x) = x \cdot a(t) + C(t)$$

for some $a(t) \in \mathbb{R}^3$ and $C(t) \in \mathbb{R}$; since h is smooth, a, C are smooth also, and the claim follows.

Now we turn to the finite energy case; thus (u, p, u_0, f, T) is now an almost smooth finite energy solution. By the time translation symmetry (30) with an arbitrarily small time shift parameter t_0 , we may assume without loss of generality that (u, p, u_0, f, T) is smooth (and not just almost smooth). We define the normalised pressure p_0 by (42) as before; then for each time $t \in [0, T]$, one sees from (8) that

$$p(t) = p_0(t) + h(t)$$

for some harmonic function $h(t) : \mathbb{R}^3 \rightarrow \mathbb{R}$. As u, f are smooth and finite energy, one sees from (42) that p_0 is bounded on compact subsets of space-time; since p is smooth, we conclude that h is bounded on compact subsets of space-time also. From harmonicity, this implies that all spatial derivatives $\nabla^k h$ are also bounded on compact subsets of space time. However, as noted previously, we cannot impose any time regularity on p_0 or h because we do not have decay estimates on time derivatives of u .

It is easy to see that h is measurable. To obtain the lemma, it suffices to show that $h(t)$ is a constant function of x for almost every time t .

Let $[t_1, t_2]$ be any interval in $[0, T]$. Integrating (3) in time on this interval, we see that

$$u(t_2, x) - u(t_1, x) + \int_{t_1}^{t_2} (u \cdot \nabla) u(t, x) dt = \int_{t_1}^{t_2} \Delta u(t, x) - \nabla p(t, x) + f(t, x) dt.$$

Next, let $\chi : \mathbb{R}^3 \rightarrow \mathbb{R}$ be a smooth compactly supported spherically symmetric function of total mass 1. We integrate the above formula against $(1/R^3)\chi(x/R)$ for some large parameter R , and conclude after some integration by parts (which is justified by the compact support of χ and the smooth (and hence C^1) nature of all functions involved) that

$$\begin{aligned} R^{-3} \int_{\mathbb{R}^3} u(t_2, x) \chi\left(\frac{x}{R}\right) dx - R^{-3} \int_{\mathbb{R}^3} u(t_1, x) \chi\left(\frac{x}{R}\right) dx - R^{-4} \int_{t_1}^{t_2} \int_{\mathbb{R}^3} u(t, x) (u(t, x) \cdot \nabla \chi)\left(\frac{x}{R}\right) dx dt \\ = R^{-5} \int_{t_1}^{t_2} \int_{\mathbb{R}^3} u(t, x) (\Delta \chi)\left(\frac{x}{R}\right) dx dt + R^{-3} \int_{t_1}^{t_2} \int_{\mathbb{R}^3} \nabla p(t, x) \chi\left(\frac{x}{R}\right) dx dt \\ + R^{-3} \int_{t_1}^{t_2} \int_{\mathbb{R}^3} f(t, x) \chi\left(\frac{x}{R}\right) dx dt. \end{aligned}$$

From the finite energy hypothesis and the Cauchy–Schwarz inequality, one easily verifies that

$$\begin{aligned} \lim_{R \rightarrow \infty} R^{-3} \int_{\mathbb{R}^3} u(t_i, x) \chi\left(\frac{x}{R}\right) dx &= 0, \\ \lim_{R \rightarrow \infty} R^{-4} \int_{t_1}^{t_2} \int_{\mathbb{R}^3} u(t, x) (u(t, x) \cdot \nabla \chi)\left(\frac{x}{R}\right) dx dt &= 0, \\ \lim_{R \rightarrow \infty} R^{-5} \int_{t_1}^{t_2} \int_{\mathbb{R}^3} u(t, x) (\Delta \chi)\left(\frac{x}{R}\right) dx dt &= 0, \\ \lim_{R \rightarrow \infty} R^{-3} \int_{t_1}^{t_2} \int_{\mathbb{R}^3} f(t, x) \chi\left(\frac{x}{R}\right) dx dt &= 0, \end{aligned}$$

and thus

$$\lim_{R \rightarrow \infty} R^{-3} \int_{t_1}^{t_2} \int_{\mathbb{R}^3} \nabla p(t, x) \chi\left(\frac{x}{R}\right) dx dt = 0. \quad (43)$$

Next, by an integration by parts and (42), we can express

$$R^{-3} \int_{t_1}^{t_2} \int_{\mathbb{R}^3} \nabla p_0(t, x) \chi\left(\frac{x}{R}\right) dx dt$$

as

$$R^{-4} \int_{t_1}^{t_2} \int_{\mathbb{R}^3} u_i u_j(t, x) (\nabla \Delta^{-1} \partial_i \partial_j \chi) \left(\frac{x}{R} \right) dx dt + R^{-3} \int_{t_1}^{t_2} \int_{\mathbb{R}^3} f_i(t, x) (\nabla \Delta^{-1} \partial_i \chi) \left(\frac{x}{R} \right) dx dt.$$

From the finite energy nature of (u, p, u_0, f, T) we see that this expression goes to zero as $R \rightarrow \infty$. Subtracting this from (43), we conclude that

$$\lim_{R \rightarrow \infty} R^{-3} \int_{t_1}^{t_2} \int_{\mathbb{R}^3} \nabla h(t, x) \chi \left(\frac{x}{R} \right) dx dt = 0. \quad (44)$$

The function $x \mapsto \int_{t_1}^{t_2} \nabla h(t, x)$ is weakly harmonic, and hence harmonic. By the mean-value property of harmonic functions (and our choice of χ), we thus have

$$R^{-3} \int_{t_1}^{t_2} \int_{\mathbb{R}^3} \nabla h(t, x) \chi \left(\frac{x}{R} \right) dx dt = \int_{t_1}^{t_2} \nabla h(t, 0) dt,$$

and thus

$$\int_{t_1}^{t_2} \nabla h(t, 0) dt = 0.$$

Since t_1, t_2 were arbitrary, we conclude from the Lebesgue differentiation theorem that $\nabla h(t, 0) = 0$ for almost every $t \in [0, T]$. Using spatial translation invariance (29) to replace the spatial origin by an element of a countable dense subset of \mathbb{R}^3 , and using the fact that harmonic functions are continuous, we conclude that $\nabla h(t)$ is identically zero for almost every $t \in [0, T]$, and so $h(t)$ is constant for almost every t as desired. \square

We note a useful corollary of Lemma 4.1(i):

Corollary 4.3 (Almost smooth H^1 solutions are essentially mild). *Let (u, p, u_0, f, T) be an almost smooth H^1 solution. Then $(u, \tilde{p}, u_0, f, T)$ is a mild H^1 solution, where*

$$\tilde{p}(t, x) := -\Delta^{-1} \partial_i \partial_j (u_i u_j)(t, x) + \Delta^{-1} \nabla \cdot f(t, x).$$

Furthermore, for almost every $t \in [0, T]$, $p(t)$ and $\tilde{p}(t)$ differ by a constant (and thus $\nabla p = \nabla \tilde{p}$).

Proof. By Lemma 4.1(i), ∇p is equal to $\nabla \tilde{p}$ almost everywhere; in particular, $\nabla p = \nabla \tilde{p}$ is a smooth tempered distribution. The claim then follows from (3) and the Duhamel formula (19). \square

5. Local well-posedness theory in H^1

In this section we review the (subcritical) local well-posedness theory for both periodic and nonperiodic H^1 mild solutions. The material here is largely standard (and in most cases has been superseded by the more powerful critical well-posedness theory); for instance the uniqueness theory already follows from [Prodi 1959] and [Serrin 1963], the blowup criterion already is present in [Leray 1934], the local existence theory follows from [Kato and Ponce 1988], regularity of mild solutions follows from [Ladyzhenskaya 1967], the stability results given here follow from the stronger stability results of [Chemin and Gallagher 2009], and the compactness results were already essentially present in [Tao 2007]. However, for the

convenience of the reader (and because we want to use the X^s function spaces defined in (13) as the basis for the theory) we shall present all this theory in a self-contained manner. There are now a number of advanced local well-posedness results at critical regularity, most notably that of [Koch and Tataru 2001], but we will not need such powerful results here.

We begin with the periodic theory. By taking advantage of the scaling symmetry (31), we may set the period L equal to 1. Using the symmetry (36), we may also restrict attention to data obeying the mean zero conditions (37), (38), and thus $u_0 \in H_x^1(\mathbb{R}^3/\mathbb{Z}^3)_0$ and $f \in L_t^\infty H_x^1([0, T] \times \mathbb{R}^3/\mathbb{Z}^3)_0$.

Theorem 5.1 (Local well-posedness in periodic H^1). *Let $(u_0, f, T, 1)$ be periodic H^1 data obeying the mean-zero conditions (37), (38).*

(i) (Strong solution). *If $(u, p, u_0, f, T, 1)$ is a periodic H^1 mild solution, then*

$$u \in C_t^0 H_x^1([0, T] \times \mathbb{R}^3/\mathbb{Z}^3).$$

In particular, one can unambiguously define $u(t)$ in $H_x^1(\mathbb{R}^3/\mathbb{Z}^3)$ for each $t \in [0, T]$.

(ii) (Local existence). *If*

$$(\|u_0\|_{H_x^1(\mathbb{R}^3/\mathbb{Z}^3)} + \|f\|_{L_t^1 H_x^1(\mathbb{R}^3/\mathbb{Z}^3)})^4 T \leq c \quad (45)$$

for a sufficiently small absolute constant $c > 0$, then there exists a periodic H^1 mild solution $(u, p, u_0, f, T, 1)$ with the indicated data with

$$\|u\|_{X^1([0, T] \times \mathbb{R}^3/\mathbb{Z}^3)} \lesssim \|u_0\|_{H_x^1(\mathbb{R}^3/\mathbb{Z}^3)} + \|f\|_{L_t^1 H_x^1(\mathbb{R}^3/\mathbb{Z}^3)}$$

and more generally

$$\|u\|_{X^k([0, T] \times \mathbb{R}^3/\mathbb{Z}^3)} \lesssim_{k, T} \|u_0\|_{H_x^k(\mathbb{R}^3/\mathbb{Z}^3)} + \|f\|_{L_t^1 H_x^k(\mathbb{R}^3/\mathbb{Z}^3)}^1$$

for each $k \geq 1$. In particular, one has local existence whenever T is sufficiently small depending on $\mathcal{H}^1(u_0, f, T, 1)$.

(iii) (Uniqueness). *There is at most one periodic H^1 mild solution $(u, p, u_0, f, T, 1)$ with the indicated data.*

(iv) (Regularity). *If $(u, p, u_0, f, T, 1)$ is a periodic H^1 mild solution, and $(u_0, f, T, 1)$ is smooth, then $(u, p, u_0, f, T, 1)$ is smooth.*

(v) (Lipschitz stability). *Let $(u, p, u_0, f, T, 1)$ be a periodic H^1 mild solution with the bounds $0 < T \leq T_0$ and*

$$\|u\|_{X^1([0, T] \times \mathbb{R}^3/\mathbb{Z}^3)} \leq M.$$

Let $(u'_0, f', T, 1)$ be another set of periodic H^1 data, and define the function

$$F(t) := e^{t\Delta}(u'_0 - u_0) + \int_0^t e^{(t-t')\Delta}(f'(t') - f(t')) dt'.$$

If the quantity $\|F\|_{X^1([0, T] \times \mathbb{R}^3/\mathbb{Z}^3)}$ is sufficiently small depending on T, M , then there exists a periodic

mild solution $(u', p', u'_0, f', T, 1)$ with

$$\|u - u'\|_{X^1([0, T] \times \mathbb{R}^3 / \mathbb{Z}^3)} \lesssim_{T, M} \|F\|_{X^1([0, T] \times \mathbb{R}^3 / \mathbb{Z}^3)}.$$

Proof. We first prove the strong solution claim (i). The linear solution

$$e^{t\Delta} u_0 + \int_0^t e^{(t-t')\Delta} P f(t') dt'$$

is easily verified to lie in $C_t^0 H_x^1([0, T] \times \mathbb{R}^3 / \mathbb{Z}^3)$, so in view of (11), it suffices to show that

$$\int_0^t e^{(t-t')\Delta} P B(u(t'), u(t')) dt'$$

also lies in $C_t^0 H_x^1([0, T] \times \mathbb{R}^3 / \mathbb{Z}^3)$. But as u is an H^1 mild solution, u lies in $X^1([0, T] \times \mathbb{R}^3 / \mathbb{Z}^3)$, so by (24), $P B(u, u)$ lies in $L_t^4 L_x^2([0, T] \times \mathbb{R}^3 / \mathbb{Z}^3)$. The claim (i) then follows easily from (22).

Now we establish local existence (ii). Let $\delta := \|u_0\|_{H_x^1(\mathbb{R}^3 / \mathbb{Z}^3)} + \|f\|_{L_t^1 H_x^1(\mathbb{R}^3 / \mathbb{Z}^3)}$; thus by (45) we have $\delta^4 T \leq c$. Using this and (25), (22), one easily establishes that the nonlinear map $u \mapsto \Phi(u)$ defined by

$$\Phi(u)(t) := e^{t\Delta} u_0 + \int_0^t e^{(t-t')\Delta} P B(u(t'), u(t')) + P f(t') dt'$$

is a contraction on the ball

$$\{u \in X^1([0, T] \times \mathbb{R}^3 / \mathbb{Z}^3) : \|u\|_{X^1([0, T] \times \mathbb{R}^3 / \mathbb{Z}^3)} \leq C\delta\}$$

if C is large enough. From the contraction mapping principle, we may then find a fixed point of Φ in this ball, and the claim (ii) follows (the estimates for higher k follow from variants of the above argument and an induction on k , and are left to the reader).

Now we establish uniqueness (iii). Suppose, in order to get a contradiction, that we have distinct solutions $(u, p, u_0, f, T, 1)$ and $(u', p', u_0, f, T, 1)$ for the same data. Then we have

$$\|u\|_{X^1([0, T] \times \mathbb{R}^3 / \mathbb{Z}^3)}, \|u'\|_{X^1([0, T] \times \mathbb{R}^3 / \mathbb{Z}^3)} \leq M.$$

To show uniqueness, it suffices to do so assuming that T is sufficiently small depending on M , as the general case then follows by subdividing $[0, T]$ into small enough time intervals and using induction. Subtracting (11) for u, u' and writing $v := u' - u$, we see that

$$v(t) = \int_0^t e^{(t-t')\Delta} P (2B(u(t'), v(t')) + B(v(t'), v(t'))) dt',$$

and thus by (22),

$$\|v\|_{X^1([0, T] \times \mathbb{R}^3 / \mathbb{Z}^3)} \lesssim M T^{1/4} \|v\|_{X^1([0, T] \times \mathbb{R}^3 / \mathbb{Z}^3)}.$$

If T is sufficiently small depending on M , this forces $\|v\|_{X^1([0, T] \times \mathbb{R}^3 / \mathbb{Z}^3)} = 0$, giving uniqueness up to time T ; iterating this argument gives the claim (iii).

Now we establish regularity (iv). To abbreviate the notation, all norms will be on $[0, T] \times \mathbb{R}^3 / \mathbb{Z}^3$. As u is an H^1 mild solution, it lies in X^1 , and hence by (25), $P B(u, u)$ lies in $L_t^4 L_x^2$. Applying (11), (23), and the smoothness of u_0, f , we conclude that $u \in X^s$ for all $s < \frac{3}{2}$. In particular, by Sobolev embedding we see that

$u \in L_t^\infty L_x^{12}$, $\nabla u \in L_t^2 L_x^{12} \cap L_t^\infty L_x^{12/5}$, and $\nabla^2 u \in L_t^2 L_x^{12/5}$, and hence $PB(u, u) \in L_t^2 H_x^1([0, T] \times \mathbb{R}^3/\mathbb{Z}^3)$. Returning to (11), (23), we now conclude that $u \in X^2([0, T] \times \mathbb{R}^3/\mathbb{Z}^3)$. One can then repeat these arguments iteratively to conclude that $u \in X^k([0, T] \times \mathbb{R}^3/\mathbb{Z}^3)$ for all $k \geq 1$, and thus $u \in L_t^\infty C^k([0, T] \times \mathbb{R}^3/\mathbb{Z}^3)$ for all $k \geq 0$. From (9) we then have $p \in L^\infty C^k([0, T] \times \mathbb{R}^3/\mathbb{Z}^3)$ for all $k \geq 0$, and then from (3) we have $\partial_t u \in L_t^\infty C^k([0, T] \times \mathbb{R}^3/\mathbb{Z}^3)$ for all $k \geq 0$. One can then obtain bounds on $\partial_t p$ and then on higher time derivatives of u and t , giving the desired smoothness, and the claim (iv) follows.

Now we establish stability (v). It suffices to establish the claim in the short-time case when T is sufficiently small depending only on M (more precisely, we take $M^4 T \leq c$ for some sufficiently small absolute constant $c > 0$), as the long-time case then follows by subdividing time and using induction. The existence of the solution $(u', p', u'_0, f'_0, T, 1)$ is then guaranteed by (ii). Evaluating (11) for u, u' and subtracting, and setting $v := u' - u$, we see that

$$v(t) = F + \int_0^t e^{(t-t')\Delta} P(2B(u, v) + B(v, v))(t') dt'$$

for all $t \in [0, T]$. Applying (22), (25), we conclude that

$$\|v\|_{X^1} \lesssim \|F\|_{X^1} + T^{1/4}(\|u\|_{X^1} + \|v\|_{X^1})\|v\|_{X^1},$$

where all norms are over $[t_0, t_1] \times \mathbb{R}^3$. Since $\|u\|_{X^1} + \|v\|_{X^1}$ is finite, we conclude (if T is small enough) that $\|v\|_{X^1([0, T] \times \mathbb{R}^3/\mathbb{Z}^3)} \lesssim \|F\|_{X^1([0, T] \times \mathbb{R}^3/\mathbb{Z}^3)}$, and the claim follows. \square

We may iterate the local well-posedness theory to obtain a dichotomy between existence and blowup. Define an *incomplete periodic mild H^1 solution* $(u, p, u_0, f, T_*^-, 1)$ from periodic H^1 data $(u_0, f, T_*, 1)$ to be fields $u : [0, T_*) \times \mathbb{R}^3/\mathbb{Z}^3 \rightarrow \mathbb{R}^3$ and $v : [0, T_*) \times \mathbb{R}^3/\mathbb{Z}^3 \rightarrow \mathbb{R}$ such that for any $0 < T < T_*$, the restriction $(u, p, u_0, f, T, 1)$ of $(u, p, u_0, f, T_*^-, 1)$ to the slab $[0, T] \times \mathbb{R}^3/\mathbb{Z}^3$ is a periodic mild H^1 solution. We similarly define the notion of an incomplete periodic smooth solution.

Corollary 5.2 (Maximal Cauchy development). *Let $(u_0, f, T, 1)$ be periodic H^1 data. Then at least one of the following two statements holds:*

- *There exists a periodic H^1 mild solution $(u, p, u_0, f, T, 1)$ with the given data.*
- *There exist a blowup time $0 < T_* < T$ and an incomplete periodic H^1 mild solution*

$$(u, p, u_0, f, T_*^-, 1)$$

up to time T_^- , which blows up in H^1 in the sense that*

$$\lim_{t \rightarrow T_*^-} \|u(t)\|_{H_x^1(\mathbb{R}^3/\mathbb{Z}^3)} = +\infty.$$

We refer to such solutions as maximal Cauchy developments.

A similar statement holds with “ H^1 data” and “ H^1 mild solution” replaced by “smooth data” and “smooth solution” respectively.

Next we establish a compactness property of the periodic H^1 flow.

Proposition 5.3 (Compactness). *If $(u_0^{(n)}, f^{(n)}, T, 1)$ is a sequence of periodic H^1 data obeying (37), (38) which is uniformly bounded in $H_x^1(\mathbb{R}^3/\mathbb{Z}^3)_0 \times L_t^\infty H_x^1([0, T] \times \mathbb{R}^3/\mathbb{Z}^3)_0$ and converges weakly¹⁴ to $(u_0, f, T, 1)$, and $(u, p, u_0, f, T, 1)$ is a periodic H^1 mild solution with the indicated data, then for n sufficiently large, there exist periodic H^1 mild solutions $(u^{(n)}, p^{(n)}, u_0^{(n)}, f^{(n)}, T, 1)$ with the indicated data, with $u^{(n)}$ converging weakly in $X^1([0, T] \times \mathbb{R}^3/\mathbb{Z}^3)$ to u . Furthermore, for any $0 < \tau < T$, $u^{(n)}$ converges strongly in $X^1([\tau, T] \times \mathbb{R}^3/\mathbb{Z}^3)$ to u .*

If $u_0^{(n)}$ converges strongly in $H_x^1(\mathbb{R}^3/\mathbb{Z}^3)_0$ to u_0 , then one can set $\tau = 0$ in the previous claim.

Proof. This result is essentially in [Tao 2007, Proposition 2.2], but for the convenience of the reader we give a full proof here.

To begin with, we assume that $u^{(n)}$ converges strongly in $H_x^1(\mathbb{R}^3/\mathbb{Z}^3)_0$ to u_0 , and relax this to weak convergence later. In view of the stability component of Theorem 5.1, it suffices to show that $F^{(n)}$ converges strongly in $X^1([0, T] \times \mathbb{R}^3/\mathbb{Z}^3)$ to zero, where

$$F^{(n)}(t) := e^{t\Delta}(u_0^{(n)} - u_0) + \int_0^t e^{(t-t')\Delta} P(f^{(n)}(t') - f(t')) dt'.$$

We have that $u_0^{(n)} - u_0$ converges strongly in $H_x^1(\mathbb{R}^3/\mathbb{Z}^3)$ to zero, while $f^{(n)} - f$ converges weakly in $L_t^\infty H_x^1([0, T] \times \mathbb{R}^3/\mathbb{Z}^3) \rightarrow 0$, and hence strongly in $L_t^2 L_x^2([0, T] \times \mathbb{R}^3/\mathbb{Z}^3)$. The claim then follows from (22).

Now we only assume that $u^{(n)}$ converges weakly in $H_x^1(\mathbb{R}^3/\mathbb{Z}^3)_0$ to u_0 . Let $0 < \tau < T$ be a sufficiently small time; then from local existence (Theorem 5.1(ii)) we see that $u^{(n)}$ and u are bounded in $X^1([0, \tau] \times \mathbb{R}^3/\mathbb{Z}^3)$ uniformly in n by some finite quantity M . Writing $v^{(n)} := u^{(n)} - u$, we obtain from (11) the difference equation

$$v^{(n)}(t) = F^{(n)}(t) + \int_0^t e^{(t-t')\Delta} P(B(u, v^{(n)}) + B(u^{(n)}, v^{(n)}))(t') dt'.$$

Since $u_0^{(n)} - u_0$ converges weakly in $H_x^1(\mathbb{R}^3/\mathbb{Z}^3)$ to zero, it converges strongly in $L_x^2(\mathbb{R}^3/\mathbb{Z}^3)$ to zero too. Using (21) as before, we see that $F^{(n)}$ converges strongly in $X^0([0, \tau] \times \mathbb{R}^3/\mathbb{Z}^3)$ to zero. From (22) we thus have

$$\|v^{(n)}\|_{X^0} \lesssim o(1) + \|B(u, v^{(n)})\|_{L_t^2 H_x^{-1}} + \|B(u^{(n)}, v^{(n)})\|_{L_t^2 H_x^{-1}},$$

where $o(1)$ goes to zero as $n \rightarrow \infty$, and all space-time norms are over $[0, \tau] \times \mathbb{R}^3/\mathbb{Z}^3$. From the form of B and Hölder's inequality, we have

$$\|B(u^{(n)}, v^{(n)})\|_{L_t^2 H_x^{-1}} \lesssim \|\mathbb{C}(u^{(n)} v^{(n)})\|_{L_t^2 L_x^2} \lesssim \tau^{1/4} \|u^{(n)}\|_{L_t^\infty L_x^6} \|v^{(n)}\|_{L_t^\infty L_x^2} \|v^{(n)}\|_{L_t^2 L_x^6}^{1/2} \lesssim M \tau^{1/4} \|v^{(n)}\|_{X^0},$$

and similarly for $B(u, v^{(n)})$, and thus

$$\|v^{(n)}\|_{X^0} \lesssim o(1) + M \tau^{1/4} \|v^{(n)}\|_{X^0}.$$

¹⁴Strictly speaking, we should use “converges in the weak-* sense” or “converges in the sense of distributions” here, in order to avoid the pathological (and irrelevant) elements of the dual space of $L_t^\infty H_x^1$ that can be constructed from the axiom of choice.

Thus, for τ small enough, one has

$$\|v^{(n)}\|_{X^0} = o(1),$$

which among other things gives weak convergence of $u^{(n)}$ to u in $[0, \tau] \times \mathbb{R}^3/\mathbb{Z}^3$. Also, by the pigeonhole principle, one can find times $\tau^{(n)}$ in $[0, \tau]$ such that

$$\|v^{(n)}(\tau^{(n)})\|_{H_x^1(\mathbb{R}^3/\mathbb{Z}^3)} = o(1).$$

Using the stability theory, and recalling that τ is small, this implies that

$$\|v^{(n)}(\tau)\|_{H_x^1(\mathbb{R}^3/\mathbb{Z}^3)} = o(1);$$

thus $u^{(n)}(\tau)$ converges strongly to $u(\tau)$. Now we can use our previous arguments to extend $u^{(n)}$ to all of $[0, T] \times \mathbb{R}^3/\mathbb{Z}^3$ and obtain strong convergence in $X^1([0, T] \times \mathbb{R}^3/\mathbb{Z}^3)$, as desired. \square

Now we turn to the nonperiodic setting. We have the following analogue of [Theorem 5.1](#):

Theorem 5.4 (Local well-posedness in H^1). *Let (u_0, f, T) be H^1 data.*

(i) (Strong solution). *If $(u, p, u_0, f, T, 1)$ is an H^1 mild solution, then*

$$u \in C_t^0 H_x^1([0, T] \times \mathbb{R}^3).$$

(ii) (Local existence and regularity). *If*

$$(\|u_0\|_{H_x^1(\mathbb{R}^3)} + \|f\|_{L_t^1 H_x^1(\mathbb{R}^3)})^4 T \leq c \tag{46}$$

for a sufficiently small absolute constant $c > 0$, then there exists a H^1 mild solution (u, p, u_0, f, T) with the indicated data, with

$$\|u\|_{X^1([0, T] \times \mathbb{R}^3)} \lesssim \|u_0\|_{H_x^1(\mathbb{R}^3)} + \|f\|_{L_t^1 H_x^1(\mathbb{R}^3)},$$

and more generally

$$\|u\|_{X^k([0, T] \times \mathbb{R}^3)} \lesssim_k \|u_0\|_{H_x^k(\mathbb{R}^3)} + \|f\|_{L_t^1 H_x^k(\mathbb{R}^3)},$$

for each $k \geq 1$. In particular, one has local existence whenever T is sufficiently small depending on $\mathcal{H}^1(u_0, f, T)$.

(iii) (Uniqueness). *There is at most one H^1 mild solution (u, p, u_0, f, T) with the indicated data.*

(iv) (Regularity). *If $(u, p, u_0, f, T, 1)$ is an H^1 mild solution, and (u_0, f, T) is Schwartz, then u and p are smooth; in fact, one has $\partial_t^j u, \partial_t^j p \in L_t^\infty H^k([0, T] \times \mathbb{R}^3)$ for all $j, k \geq 0$.*

(v) (Lipschitz stability). *Let $(u, p, u_0, f, T), (u', p', u'_0, f', T)$ be H^1 mild solutions with the bounds $0 < T \leq T_0$ and*

$$\|u\|_{X^1([0, T] \times \mathbb{R}^3)}, \|u'\|_{X^1([0, T] \times \mathbb{R}^3)} \leq M.$$

Define the function

$$F(t) := e^{t\Delta}(u'_0 - u_0) + \int_0^t e^{(t-t')\Delta}(f'(t') - f(t')) dt'.$$

If the quantity $\|F\|_{L_t^2 L_x^2([0, T] \times \mathbb{R}^3)}$ is sufficiently small depending on T, M , then

$$\|u - u'\|_{X^1([0, T] \times \mathbb{R}^3)} \lesssim_{T, M} \|F\|_{L_t^2 L_x^2([0, T] \times \mathbb{R}^3)}.$$

Proof. This proceeds by repeating the proof of [Theorem 5.1](#) verbatim. The one item which perhaps requires some care is the regularity item (iv). The arguments from [Theorem 5.1](#) yield the regularity

$$u \in X^k([0, T] \times \mathbb{R}^3)$$

for all $k \geq 0$ without difficulty. In particular, $u \in L_t^\infty H_x^k([0, T] \times \mathbb{R}^3)$ for all $k \geq 0$. From (9) and Sobolev embedding, one then has $p \in L_t^\infty H_x^k([0, T] \times \mathbb{R}^3)$ for all $k \geq 0$, and then from (3) and more Sobolev embedding, one has $\partial_t u \in L_t^\infty H_x^k([0, T] \times \mathbb{R}^3)$ for all $k \geq 0$. One can then obtain bounds on $\partial_t p$ and then on higher time derivatives of u and t , giving the desired smoothness, and the claim (iv) follows. (Note that these arguments did not require the full power of the hypothesis that (u_0, f, T) was Schwartz; it would have sufficed to have $u_0 \in H_x^k(\mathbb{R}^3)$ and $f \in C_t^j H_x^k(\mathbb{R}^3)$ for all $j, k \geq 0$.) \square

From the regularity component of the above theorem, we immediately conclude that [Conjecture 1.19](#) implies [Conjecture 1.5](#), which is one half of [Theorem 1.20\(iv\)](#).

We will also need a more quantitative version of the regularity statement in [Theorem 5.4](#).

Lemma 5.5 (Quantitative regularity). *Let (u, p, u_0, f, T) be an H^1 mild solution obeying (46) for a sufficiently small absolute constant $c > 0$, and such that*

$$\|u_0\|_{H_x^1(\mathbb{R}^3)} + \|f\|_{L_t^1 H_x^k(\mathbb{R}^3)} \leq M < \infty.$$

Then one has

$$\|u\|_{L_t^\infty H_x^k([\tau, T] \times \mathbb{R}^3)} \lesssim_{k, \tau, T, M} 1$$

for all natural numbers $k \geq 1$ and all $0 < \tau < T$.

Proof. We allow all implied constants to depend on k, T, M . From [Theorem 5.1](#) we have

$$\|u\|_{X^1([0, T] \times \mathbb{R}^3)} \lesssim 1,$$

which already gives the $k = 1$ case. Now we turn to the $k \geq 2$ case. From (25) we have

$$\|PB(u, u)\|_{L_t^4 L_x^2([0, T] \times \mathbb{R}^3)} \lesssim 1,$$

while from Fourier analysis one has

$$\|e^{t\Delta} u_0\|_{L_t^\infty H_x^k([\tau, T] \times \mathbb{R}^3)} \lesssim_\tau 1.$$

From this and (11), (21) we see that

$$\|u\|_{X^s([\tau, T] \times \mathbb{R}^3)} \lesssim_{s, \tau} 1$$

for all $s < \frac{3}{2}$. From Sobolev embedding we conclude

$$\begin{aligned} \|u\|_{L_t^\infty L_x^{12}([\tau, T] \times \mathbb{R}^3)} &\lesssim_\tau 1, \\ \|\nabla u\|_{L_t^2 L_x^{12}([\tau, T] \times \mathbb{R}^3)} &\lesssim_\tau 1, \\ \|\nabla u\|_{L_t^\infty L_x^{12/5}([\tau, T] \times \mathbb{R}^3)} &\lesssim_\tau 1, \\ \|\nabla^2 u\|_{L_t^2 L_x^{12}([\tau, T] \times \mathbb{R}^3)} &\lesssim_\tau 1, \end{aligned}$$

and hence

$$\|PB(u, u)\|_{L_t^2 H_x^1([\tau, T] \times \mathbb{R}^3)} \lesssim 1.$$

Returning to (11), (23), we now conclude that

$$\|u\|_{X^2([\tau, T] \times \mathbb{R}^3)} \lesssim_\tau 1,$$

which gives the $k = 2$ case. One can repeat these arguments iteratively to then give the higher k cases. \square

We extract a particular consequence of the above lemma:

Proposition 5.6 (Almost regularity). *Let $(u, p, u_0, 0, T)$ be a homogeneous H^1 mild solution obeying (46) for a sufficiently small absolute constant $c > 0$. Then u, p are smooth on $[\tau, T] \times \mathbb{R}^3$ for all $0 < \tau < T$; in fact, all derivatives of u, p lie in $L_t^\infty L_x^2([\tau, T] \times \mathbb{R}^3)$. If furthermore u_0 is smooth, then $(u, p, u_0, 0, T)$ is an almost smooth solution.*

Proof. From Lemma 5.5 we see that

$$u \in L_t^\infty H_x^k([\tau, T] \times \mathbb{R}^3)$$

for all $k \geq 0$ and $0 < \tau < T$. Arguing as in the proof of Theorem 5.4(iv), we conclude that u, p are smooth on $[\tau, T] \times \mathbb{R}^3$.

Now suppose that u_0 is smooth. Then (since u_0 is also in $H_x^1(\mathbb{R}^3)$) $e^{t\Delta}u_0$ is smooth¹⁵ on $[0, T] \times \mathbb{R}^3$, and in particular one has

$$\eta e^{t\Delta}u_0 \in L_t^\infty H_x^k([0, T] \times \mathbb{R}^3)$$

for any smooth, compactly supported cutoff function $\eta : \mathbb{R}^3 \rightarrow \mathbb{R}$. Meanwhile, by arguing as in Lemma 5.5 one has

$$PB(u, u) \in L_t^4 L_x^2([0, T] \times \mathbb{R}^3). \quad (47)$$

Using (11), (21), one concludes that

$$\eta u \in X^s([0, T] \times \mathbb{R}^3)$$

for all cutoff functions η and all $s < \frac{3}{2}$. Continuing the arguments from Lemma 5.5, we conclude that

$$\eta PB(u, u) \in L_t^2 H_x^1([0, T] \times \mathbb{R}^3)$$

¹⁵To obtain smoothness at a point (t_0, x_0) , one can for instance split u_0 into a smooth compactly supported component and a component that vanishes near x_0 but lies in $H_x^1(\mathbb{R}^3)$, and verify that the contribution of each component to $e^{t\Delta}u_0$ is smooth at (t_0, x_0) .

for all cutoffs η . Using (11), (23) (and using (28), (47) to deal with the far field contribution of $PB(u, u)$, and shrinking η as necessary), one then concludes that

$$\eta u \in X^2([0, T] \times \mathbb{R}^3)$$

for all cutoffs η . Repeating these arguments iteratively, one eventually concludes that

$$\eta u \in X^k([0, T] \times \mathbb{R}^3)$$

for all cutoffs η , and in particular

$$u \in L_t^\infty H_x^k([0, T] \times K)$$

for all $k \geq 0$ and all compact sets K . By Sobolev embedding, this implies that

$$u \in L_t^\infty C_x^k([0, T] \times K)$$

for all $k \geq 0$ and all compact sets K .

We also have $u \in X^1([0, T] \times \mathbb{R}^3)$, and hence

$$u \in L_t^\infty H_x^1([0, T] \times \mathbb{R}^3).$$

In particular,

$$u_i u_j \in L_t^\infty L_x^1([0, T] \times \mathbb{R}^3) \quad (48)$$

and

$$u_i u_j \in L_t^\infty C_x^k([0, T] \times K)$$

for all $k \geq 0$ and compact K . From this and (9) (splitting the inverse Laplacian Δ^{-1} smoothly into local and global components), one has

$$p \in L_t^\infty C_x^k([0, T] \times K);$$

inserting this into (3), we then see that

$$\partial_t u \in L_t^\infty C_x^k([0, T] \times K) \quad (49)$$

for all $k \geq 0$ and compact K .

This is a little weaker than what we need for an almost smooth solution, because we want $\nabla^k u$, $\nabla^k p$, $\partial_t \nabla^k p$ to extend continuously down to $t = 0$, and the above estimates merely give $L_t^\infty C_x^\infty$ control on these quantities. To upgrade the L_t^∞ control to continuity in time, we first observe¹⁶ from (49) and integration in time that we can at least make $\nabla^k u$ extend continuously to $t = 0$:

$$u \in C_t^0 C_x^k([0, T] \times K).$$

In particular,

$$u_i u_j \in C_t^0 C_x^k([0, T] \times K) \quad (50)$$

¹⁶An alternate argument here would be to approximate the initial data u_0 by Schwartz divergence-free data (using Lemma 12.1) and to use a limiting argument and the stability and regularity theory in Theorem 5.1; we omit the details.

for all $k \geq 0$ and compact K .

Now we consider $\nabla^k p$ in a compact region $[0, T] \times K$. From (9) we have

$$\nabla^k p(t, x) = \nabla^k \partial_i \partial_j \int_{\mathbb{R}^3} \frac{1}{4\pi|x-y|} u_i u_j(t, y) dy.$$

Using a smooth cutoff, we split the Newton potential $1/(4\pi|x-y|)$ into a “local” portion supported on $B(0, 2R)$ and a “global” portion supported outside of $B(0, R)$, where R is a large radius. From (50) one can verify that the contribution of the local portion is continuous on $[0, T] \times K$, while from (48) the contribution of the global portion is $O_u(1/R^3)$. Sending $R \rightarrow \infty$, we conclude that $\nabla^k p$ is continuous on $[0, T] \times K$, and thus

$$p \in C_t^0 C_x^k([0, T] \times K)$$

for all $k \geq 0$ and compact K . Inserting this into (3), we then conclude that

$$\partial_t u \in C_t^0 C_x^k([0, T] \times K)$$

for all $k \geq 0$ and compact K , and so we have an almost smooth solution as required. \square

Remark 5.7. Because u has the regularity of $L_t^\infty H_x^1$, we can continue iterating the above argument a little more, and eventually get $u \in C_t^2 C_x^k([0, T] \times K)$ and $p \in C_t^1 C_x^k([0, T] \times K)$ for all $k \geq 0$ and compact K . Using the vorticity equation (see (84) below), one can then also get $\omega \in C_t^3 C_x^k([0, T] \times K)$ as well. But without further decay conditions on higher derivatives of u (or of ω), one cannot gain infinite regularity on u, p, ω in time; see Section 15.

On the other hand, it is possible to use energy methods and the vorticity equation (84) to show (working in the homogeneous case $f = 0$ for simplicity) that if u_0 is smooth and the initial vorticity $\omega_0 := \nabla \times u_0$ is Schwartz, then the solution in Proposition 5.6 is in fact smooth, with ω remaining Schwartz throughout the lifespan of that solution; we omit the details.

As a corollary of the above proposition we see that Conjecture 1.19 implies Conjecture 1.13, thus completing the proof of Theorem 1.20(iv).

As before, we obtain a dichotomy between existence and blowup. Define an *incomplete mild H^1 solution* (u, p, u_0, f, T_*^-) from H^1 data (u_0, f, T_*) to be fields $u : [0, T_*^-) \times \mathbb{R}^3 \rightarrow \mathbb{R}^3$ and $v : [0, T_*^-) \times \mathbb{R}^3 \rightarrow \mathbb{R}$ such that for any $0 < T < T_*$, the restriction $(u, p, u_0, f, T, 1)$ of $(u, p, u_0, f, T_*^-, 1)$ to the slab $[0, T] \times \mathbb{R}^3$ is a mild H^1 solution. We similarly define the notion of an incomplete smooth H^1 solution.

Corollary 5.8 (Maximal Cauchy development). *Let (u_0, f, T) be H^1 data. Then at least one of the following two statements holds:*

- *There exists a mild H^1 solution (u, p, u_0, f, T) with the given data.*
- *There exist a blowup time $0 < T_* < T$ and an incomplete mild H^1 solution (u, p, u_0, f, T_*^-) up to time T_*^- that blows up in the enstrophy norm in the sense that*

$$\lim_{t \rightarrow T_*^-} \|u(t)\|_{H_x^1(\mathbb{R}^3)} = +\infty.$$

Remark 5.9. In the second conclusion of [Corollary 5.8](#), more information about the blowup is known. For instance, in [\[Iskauriaza et al. 2003\]](#) it was demonstrated that the $L_x^3(\mathbb{R}^3)$ norm must also blow up (in the homogeneous case $f = 0$, at least).

6. Homogenisation

In this section we prove [Proposition 1.7](#).

Fix smooth periodic data (u_0, f, T, L) ; our objective is to find a smooth periodic solution

$$(u, p, u_0, f, T, L)$$

(without pressure normalisation) with this data. By the scaling symmetry [\(31\)](#), we may normalise the period L to equal 1. Using the symmetry [\(36\)](#), we may impose the mean-zero conditions [\(37\)](#), [\(38\)](#) on this data.

By hypothesis, one can find a smooth periodic solution $(\tilde{u}, \tilde{p}, u_0, 0, T, 1)$ with data $(u_0, 0, T, 1)$. By [Lemma 4.1](#), and applying a Galilean transform [\(33\)](#) if necessary, we may assume the pressure is normalised, which in particular makes $(\tilde{u}, \tilde{p}, u_0, 0, T, 1)$ a periodic H^1 mild solution.

By the Galilean invariance [\(33\)](#) (with a linearly growing velocity $v(t) := 2wt$), it suffices to find a smooth periodic solution (u, p, u_0, f_w, T) (this time *with* pressure normalisation) for the Galilean-shifted data (u_0, f_w, T) , where

$$f_w(t, x) := f(t, x - wt^2),$$

and $w \in \mathbb{R}^3$ is arbitrary. Note that the data (u_0, f_w, T) continues to obey the mean-zero conditions [\(37\)](#), [\(38\)](#) and is bounded in $H_x^1(\mathbb{R}^3/\mathbb{Z}^3)_0 \times L_t^\infty H_x^1([0, T] \times \mathbb{R}^3/\mathbb{Z}^3)_0$ uniformly in w . We now make a key observation:

Lemma 6.1. *If $\alpha \in \mathbb{R}^3/\mathbb{Z}^3$ is irrational in the sense that $k \cdot \alpha \neq 0$ in \mathbb{R}/\mathbb{Z} for all $k \in \mathbb{Z}^3 \setminus \{0\}$, then $f_{\lambda\alpha}$ converges weakly (or more precisely, converges in the sense of space-time distributions) to zero in $L_t^\infty H_x^1([0, T] \times \mathbb{R}^3/\mathbb{Z}^3)_0$.*

Proof. It suffices to show that

$$\int_0^T \int_{\mathbb{R}^3/\mathbb{Z}^3} f_{\lambda\alpha}(t, x) \phi(t, x) dx dt \rightarrow 0$$

for all smooth functions $\phi : [0, T] \times \mathbb{R}^3/\mathbb{Z}^3 \rightarrow \mathbb{R}$. Taking the Fourier transform, the left-hand side becomes

$$\sum_{k \in \mathbb{Z}^3} \int_0^T e^{-2\pi i \lambda k t^2 \cdot \alpha} \widehat{f}(t)(k) \widehat{\phi}(t)(-k) dt,$$

with the sum being absolutely convergent due to the rapid decrease of the Fourier transform of $\phi(t)$. Because f has mean zero, we can delete the $k = 0$ term from the sum. This makes $k \cdot \alpha$ nonzero by irrationality, and so by the Riemann–Lebesgue lemma, each summand goes to zero as $\lambda \rightarrow \infty$. The claim then follows from the dominated convergence theorem. \square

Let $\alpha \in \mathbb{R}^3/\mathbb{Z}^3$ be irrational. By the above lemma, $(u_0, f_{\lambda\alpha}, T, 1)$ converges weakly to $(u_0, 0, T, 1)$ while being bounded in $H_x^1(\mathbb{R}^3/\mathbb{Z}^3)_0 \times L_t^\infty H_x^1(\mathbb{R}^3/\mathbb{Z}^3)_0$. As $(u_0, 0, T, 1)$ has a periodic mild H^1 solution $(\tilde{u}, \tilde{p}, u_0, 0, T, 1)$, we conclude from [Proposition 5.3](#) that for λ sufficiently large, $(u_0, f_{\lambda\alpha}, T, 1)$ also has a periodic mild H^1 solution, which is necessarily smooth since u_0 and $f_{\lambda\alpha}$ are smooth. The claim follows.

Remark 6.2. Suppose that $(u_0, f, \infty, 1)$ is periodic H^1 data extending over the half-infinite time interval $[0, +\infty)$. The above argument shows (assuming [Conjecture 1.4](#)) that one can, for each $0 < T < \infty$, construct a smooth periodic (but not pressure-normalised) solution $(u^{(T)}, p^{(T)}, u_0, f, T, 1)$ up to time T with the above data, by choosing a sufficiently rapidly growing linear velocity $v^{(T)} = 2w^{(T)}t$, applying a Galilean transform, and then using the compactness properties of the H^1 local well-posedness theory. As stated, this argument gives a different solution $(u^{(T)}, p^{(T)}, u_0, f, T, 1)$ for each time T (note that we do not have uniqueness once we abandon pressure normalisation). However, it is possible to modify the argument to obtain a single global smooth periodic solution $(u, p, u_0, f, \infty, 1)$ (which is still not pressure-normalised, of course), by using the ability in [\(33\)](#) to choose a *nonlinear* velocity $v(t)$ rather than a linear one. By reworking the above argument, and taking $v(t)$ to be a sufficiently rapidly growing function of t , it is then possible to obtain a global smooth periodic solution $(u, p, u_0, f, \infty, 1)$ to the indicated data; we omit the details.

7. Compactness

In this section we prove [Theorem 1.20\(i\)](#) by following the compactness arguments of [\[Tao 2007\]](#). By the scaling symmetry [\(31\)](#), we may normalise $L = 1$.

We first assume that [Conjecture 1.10](#) holds, and deduce [Conjecture 1.9](#). Suppose for contradiction that [Conjecture 1.9](#) failed. By [Corollary 5.2](#), there thus exists an incomplete periodic pressure-normalised mild H^1 solution $(u, p, u_0, f, T_*^-, 1)$ such that

$$\lim_{t \rightarrow T_*^-} \|u(t)\|_{H_x^1(\mathbb{R}^3/\mathbb{Z}^3)} = \infty. \quad (51)$$

By Galilean invariance [\(36\)](#), we may assume that u_0 and f (and hence u) have mean zero.

Let $(u_0^{(n)}, f^{(n)}, T_*, 1)$ be a sequence of periodic smooth mean-zero data converging strongly in

$$H_x^1(\mathbb{R}^3/\mathbb{Z}^3)_0 \times L_t^\infty H_x^1([0, T_*] \times \mathbb{R}^3/\mathbb{Z}^3)_0$$

to the periodic H^1 data $(u, f, T_*, 1)$. For each time $0 < T < T_*$, we see from [Theorem 5.1](#) that for n sufficiently large, we may find a smooth solution $(u^{(n)}, p^{(n)}, u_0^{(n)}, T, 1)$ with this data, with $u^{(n)}$ converging strongly in $L_t^\infty H_x^1([0, T] \times \mathbb{R}^3/\mathbb{Z}^3)$ to u . By [Conjecture 1.10](#), the $L_t^\infty H_x^1([0, T] \times \mathbb{R}^3/\mathbb{Z}^3)$ norm of $u^{(n)}$ is bounded uniformly in both T and n , so by taking limits as $n \rightarrow \infty$, we conclude that $\|u(t)\|_{H_x^1(\mathbb{R}^3/\mathbb{Z}^3)}$ is bounded uniformly for $0 \leq t < T_*$, contradicting [\(51\)](#) as desired.

Conversely, suppose that [Conjecture 1.9](#) held, but [Conjecture 1.10](#) failed. Carefully negating all the quantifiers, we conclude that there exists a time $0 < T_0 < \infty$ and a sequence $(u^{(n)}, p^{(n)}, u_0^{(n)}, f^{(n)}, T^{(n)}, 1)$ of smooth periodic data with $0 < T^{(n)} < T_0$ and $\mathcal{H}^1(u_0^{(n)}, f^{(n)}, T^{(n)}, 1)$ uniformly bounded in n , such that

$$\lim_{n \rightarrow \infty} \|u\|_{L_t^\infty H_x^1([0, T^{(n)}] \times \mathbb{R}^3/\mathbb{Z}^3)} = \infty. \quad (52)$$

Using Galilean transforms (36), we may assume that $u_0^{(n)}, f^{(n)}$ (and hence $u^{(n)}$) have mean zero. From the short-time local existence (and uniqueness) theory in Theorem 5.1, we see that $T^{(n)}$ is bounded uniformly away from zero. Thus by passing to a subsequence, we may assume that $T^{(n)}$ converges to a limit T_* with $0 < T_* \leq T_0$.

By sequential weak compactness, we may pass to a further subsequence and assume that for each $0 < T < T_*$, $(u_0^{(n)}, f^{(n)}, T, 1)$ converges weakly (or more precisely, in the sense of distributions) to a periodic H^1 limit $(u_0, f, T, 1)$; gluing these limits together, one obtains periodic H^1 data $(u_0, f, T_*, 1)$, which still has mean zero. By Conjecture 1.9, we can then find a periodic H^1 mild solution $(u, p, u_0, f, T_*, 1)$ with this data, which then necessarily also has mean zero.

By Theorem 5.1 and Proposition 5.3, we see that for every $0 < \tau < T < T_*$, $u^{(n)}$ converges strongly in $L_t^\infty H_x^1([\tau, T] \times \mathbb{R}^3/\mathbb{Z}^3)$ to u . In particular, for any $0 < T < T_*$, one has

$$\limsup_{n \rightarrow \infty} \|u^{(n)}(T)\|_{H_x^1(\mathbb{R}^3/\mathbb{Z}^3)} \leq \|u\|_{L_t^\infty H_x^1([0, T_*] \times \mathbb{R}^3/\mathbb{Z}^3)} < \infty.$$

Taking T sufficiently close to T_* and then taking n sufficiently large, we conclude from Theorem 5.1 that

$$\limsup_{n \rightarrow \infty} \|u^{(n)}\|_{L_t^\infty H_x^1([T, T^{(n)}] \times \mathbb{R}^3/\mathbb{Z}^3)} < \infty;$$

also, from the strong convergence in $L_t^\infty H_x^1([\tau, T] \times \mathbb{R}^3/\mathbb{Z}^3)$, we have

$$\limsup_{n \rightarrow \infty} \|u^{(n)}\|_{L_t^\infty H_x^1([\tau, T] \times \mathbb{R}^3/\mathbb{Z}^3)} < \infty$$

for any $0 < \tau < T$, and finally from the local existence (and uniqueness) theory in Theorem 5.1, one has

$$\limsup_{n \rightarrow \infty} \|u^{(n)}\|_{L_t^\infty H_x^1([0, \tau] \times \mathbb{R}^3/\mathbb{Z}^3)} < \infty$$

for sufficiently small τ . Putting these bounds together, we contradict (52), and the claim follows.

Remark 7.1. It should be clear to the experts that one could have replaced the H^1 regularity in the above conjectures by other subcritical regularities, such as H^k for $k > 1$, and obtained a similar result to Theorem 1.20(i).

As remarked previously, the homogeneous case $f = 0$ of Theorem 1.20(i) was established in [Tao 2007]. We recall the main results of that paper. We introduce the following homogeneous periodic conjectures:

Conjecture 7.2 (*A priori* homogeneous periodic H^1 bound). There exists a function $F : \mathbb{R}^+ \times \mathbb{R}^+ \times \mathbb{R}^+ \rightarrow \mathbb{R}^+$ with the property that whenever $(u, p, u_0, 0, T, L)$ is a smooth periodic homogeneous normalised-pressure solution with $0 < T < T_0 < \infty$ and

$$\mathcal{H}^1(u_0, 0, T, L) \leq A < \infty,$$

then

$$\|u\|_{L_t^\infty H_x^1([0, T] \times \mathbb{R}^3/L\mathbb{Z}^3)} \leq F(A, L, T_0).$$

Conjecture 7.3 (*A priori homogeneous global periodic H^1 bound*). There exists a function

$$F : \mathbb{R}^+ \times \mathbb{R}^+ \rightarrow \mathbb{R}^+$$

with the property that whenever $(u, p, u_0, 0, T, L)$ is a smooth periodic homogeneous normalised-pressure solution with

$$\mathcal{H}^1(u_0, 0, T, L) \leq A < \infty,$$

then

$$\|u\|_{L_t^\infty H_x^1([0, T] \times \mathbb{R}^3 / L\mathbb{Z}^3)} \leq F(A, L).$$

Conjecture 7.4 (*Global well-posedness in periodic homogeneous H^1*). Let $(u_0, 0, T, L)$ be a homogeneous periodic H^1 set of data. Then there exists a periodic H^1 mild solution $(u, p, u_0, 0, T, L)$ with the indicated data.

Conjecture 7.5 (*Global regularity for homogeneous periodic data with normalised pressure*). Suppose $(u_0, 0, T)$ is a smooth periodic set of data. Then there exists a smooth periodic solution $(u, p, u_0, 0, T)$ with the indicated data and with normalised pressure.

In [Tao 2007, Theorem 1.4] it was shown that Conjectures 1.4, 7.2, 7.3 are equivalent. As implicitly observed in that paper also, Conjecture 1.4 is equivalent to Conjecture 7.5 (this can be seen from Lemma 4.1), and from the local well-posedness and regularity theory (Theorem 5.1 or [Tao 2007, Proposition 2.2]), we also see that Conjecture 7.5 is equivalent to Conjecture 7.4.

8. Energy localisation

In this section we establish the energy inequality for the Navier–Stokes equation in the smooth finite energy setting. This energy inequality is utterly standard (see for example [Scheffer 1976]) for weaker notions of solutions, so long as one has regularity of $L_t^2 H_x^1$, but (somewhat ironically) requires more care in the smooth finite energy setting, because we do *not* assume *a priori* that smooth finite energy solutions lie in the space $L_t^2 H_x^1$. The methods used here are local in nature, and will also provide an energy localisation estimate for the Navier–Stokes equation (see Theorem 8.2).

We begin with the global energy inequality.

Lemma 8.1 (*Global energy inequality*). *Let (u, p, u_0, f, T) be a finite energy almost smooth solution. Then*

$$\|u\|_{L_t^\infty L_x^2([0, T] \times \mathbb{R}^3)} + \|\nabla u\|_{L_t^2 L_x^2([0, T] \times \mathbb{R}^3)} \lesssim E(u_0, f, T)^{1/2}. \quad (53)$$

In particular, u lies in the space $X^1([0, T] \times \mathbb{R}^3)$.

Proof. To abbreviate the notation, all spatial norms here will be over \mathbb{R}^3 .

Using the forcing symmetry (34), we may set f to be divergence-free, so in particular by Corollary 4.3 we have

$$\nabla p(t) = \nabla \tilde{p}(t) \quad (54)$$

for almost all times t , where

$$\tilde{p} = -\Delta^{-1} \partial_i \partial_j (u_i u_j). \quad (55)$$

As (u, p, u_0, f, T) is finite energy, we have the *a priori* hypothesis

$$\|u\|_{L_t^\infty L_x^2([0, T] \times \mathbb{R}^3)} \leq A$$

for some $A < \infty$, though recall that our final bounds are not allowed to depend on this quantity A . Because u is smooth, we see in particular from Fatou's lemma that

$$\|u(t)\|_{L_x^2} \leq A \quad (56)$$

for all $t \in [0, T]$.

Taking the inner product of the Navier–Stokes equation (3) with u and rearranging, we obtain the energy density identity

$$\partial_t \left(\frac{1}{2} |u|^2 \right) + u \cdot \nabla \left(\frac{1}{2} |u|^2 \right) = \Delta \left(\frac{1}{2} |u|^2 \right) - |\nabla u|^2 - u \cdot \nabla p + u \cdot f. \quad (57)$$

We would like to integrate this identity over all of \mathbb{R}^3 , but we do not yet have enough decay in space to achieve this, even with the normalised pressure. Instead, we will localise by integrating the identity against a cutoff η^4 , where $\eta(x) := \chi((|x| - R)/r)$, $\chi : \mathbb{R} \rightarrow \mathbb{R}^+$ is a fixed smooth function that equals 0 on $[0, +\infty]$ and 1 on $[-\infty, -1]$, and $0 < r < R/2$ are parameters to be chosen later. (The exponent 4 is convenient for technical reasons, in that η^4 and $\nabla(\eta^4)$ share a large common factor η^3 , but it should be ignored on a first reading.) Thus we see that η^4 is supported on the ball $B(0, R)$ and equals 1 on $B(0, R - r)$, with the derivative bounds

$$\nabla^j \eta = O(r^{-j}) \quad (58)$$

for $j = 0, 1, 2$. We define the localised energy

$$E_{\eta^4}(t) := \int_{\mathbb{R}^3} \frac{1}{2} |u|^2(t, x) \eta^4(x) dx. \quad (59)$$

Clearly we have the initial condition

$$E_{\eta^4}(0) \lesssim E(u_0, f). \quad (60)$$

Because η^4 is compactly supported and u is almost smooth, E_{η^4} is C_t^1 , and we may differentiate under the integral sign and integrate by parts without difficulty; using (54), we see for almost every time t that

$$\partial_t E_{\eta^4} = -X_1 + X_2 + X_3 + X_4 + X_5, \quad (61)$$

where X_1 is the dissipation term

$$X_1 := \int_{\mathbb{R}^3} |\nabla u|^2 \eta^4 dx = \|\eta^2 \nabla u\|_{L_x^2}^2, \quad (62)$$

X_2 is the heat flux term

$$X_2 := \frac{1}{2} \int_{\mathbb{R}^3} |u|^2 \Delta(\eta^4) dx,$$

X_3 is the transport term

$$X_3 := 4 \int_{\mathbb{R}^3} |u|^2 u \cdot \eta^3 \nabla \eta \, dx,$$

X_4 is the forcing term

$$X_4 := \int_{\mathbb{R}^3} u \cdot f \eta^4 \, dx,$$

and X_5 is the pressure term

$$X_5 := 4 \int_{\mathbb{R}^3} u \tilde{p} \eta^3 \nabla \eta \, dx.$$

The dissipation term X_1 is nonnegative, and will be useful in controlling some of the other terms present here. The heat flux term X_2 can be bounded using (56) and (58) by

$$X_2 \lesssim \frac{A^2}{r^2},$$

so we turn now to the transport term X_3 . Using Hölder's inequality and (58), we may bound

$$X_3 \lesssim \frac{1}{r} \|u \eta^2\|_{L_x^6}^{3/2} \|u\|_{L_x^2}^{3/2}, \quad (63)$$

and thus by (56) and Sobolev embedding

$$X_3 \lesssim \frac{A^{3/2}}{r} \|\nabla(u \eta^2)\|_{L_x^2}^{3/2}.$$

By the Leibniz rule and (62), (56), (58), one has

$$\|\nabla(u \eta^2)\|_{L_x^2} \lesssim X_1^{1/2} + \frac{A}{r},$$

and thus

$$X_3 \lesssim \frac{A^{3/2}}{r} X_1^{3/4} + \frac{A^3}{r^{5/2}}.$$

Now we move on to the forcing term X_4 . By Cauchy–Schwarz, we can bound this term by

$$X_4 \lesssim E_{\eta^4}^{1/2} a(t),$$

where $a(t) := \|f(t)\|_{L_x^2(B(0,R))}$. Note from (2) that

$$\int_0^T a(t) \, dt \lesssim E(u_0, f, T)^{1/2}. \quad (64)$$

Now we turn to the pressure term X_5 . From (55) we have

$$X_5 = \int_{\mathbb{R}^3} \mathbb{O}(u(\Delta^{-1} \nabla^2(uu)) \eta^3 \nabla \eta).$$

We will argue as in the estimation of X_4 , but we will first need to move the η^3 weight past the singular integral $\Delta^{-1} \nabla^2$. We therefore bound $X_5 = X_{5,1} + X_{5,2}$, where

$$X_{5,1} = \int_{\mathbb{R}^3} \mathbb{O}(u(\Delta^{-1} \nabla^2(uu \eta^3)) \nabla \eta)$$

and

$$X_{5,2} = \int_{\mathbb{R}^3} \mathbb{O}(u[\Delta^{-1}\nabla^2, \eta^3](uu)\nabla\eta),$$

where $[A, B] := AB - BA$ is the commutator and η^3 is interpreted as the multiplication operator $\eta^3 : u \mapsto \eta^3 u$. For $X_{6,1}$, we apply Hölder's inequality and (58) to obtain

$$X_{5,1} \lesssim \frac{1}{r} \|u\|_{L_x^2} \|\Delta^{-1}\nabla^2(uu\eta^3)\|_{L_x^2}.$$

The singular integral $\Delta^{-1}\nabla^2$ is bounded on L^2 , so it may be discarded; applying Hölder's inequality again, we conclude that

$$X_{5,1} \lesssim \frac{1}{r} \|u\|_{L_x^2}^{3/2} \|u\eta^2\|_{L_x^6}^{3/2}.$$

This is the same bound (63) used to bound X_3 , and so by repeating the X_3 analysis, we conclude that

$$X_{5,1} \lesssim \frac{A^{3/2}}{r} X_1^{3/4} + \frac{A^3}{r^{5/2}}.$$

As for $X_{5,2}$, we observe from direct computation of the integral kernel that when $r = 1$, $[\Delta^{-1}\nabla^2, \chi^3]$ is a smoothing operator of infinite order (see [Kato and Ponce 1988]), and in particular

$$\|[\Delta^{-1}\nabla^2, \eta^3]f\|_{L_x^2} \lesssim \|f\|_{L_x^1}$$

in the $r = 1$ case. In the general case, a rescaling argument then gives

$$\|[\Delta^{-1}\nabla^2, \eta^3]f\|_{L_x^2} \lesssim \frac{1}{r^{3/2}} \|f\|_{L_x^1}.$$

Applying Hölder's inequality and (56), we conclude that

$$X_{5,2} \lesssim \frac{A^3}{r^{5/2}}.$$

Putting all the estimates together, we conclude that

$$\partial_t E_{\eta^4} \leq -X_1 + O\left(\frac{A^2}{r^2} + \frac{A^{3/2}}{r} X_1^{3/4} + \frac{A^3}{r^{5/2}} + E_{\eta^4}^{1/2} a(t)\right).$$

By Young's inequality, we have

$$-\frac{1}{2}X_1 + O\left(\frac{A^{3/2}}{r} X_1^{3/4}\right) \lesssim \frac{A^6}{r^4}$$

and

$$\frac{A^3}{r^{5/2}} \lesssim \frac{A^2}{r^2} + \frac{A^6}{r^4},$$

and so we obtain

$$\partial_t E_{\eta^4} + X_1 \lesssim \frac{A^2}{r^2} + \frac{A^6}{r^4} + E_{\eta^4}^{1/2} a(t), \quad (65)$$

and hence for almost every time t ,

$$\partial_t (E_{\eta^4} + E(u_0, f, T))^{1/2} \lesssim E(u_0, f, T)^{-1/2} \left(\frac{A^2}{r^2} + \frac{A^6}{r^4} \right) + a(t).$$

By the fundamental theorem of calculus, (64), and (60), we conclude that

$$E_{\eta^4}(t)^{1/2} \lesssim E(u_0, f, T)^{1/2} + E(u_0, f, T)^{-1/2} \left(\frac{A^2}{r^2} + \frac{A^6}{r^4} \right) T$$

for all $t \in [0, T]$ and all sufficiently large R ; sending $r, R \rightarrow \infty$ and using the monotone convergence theorem, we conclude that

$$\|u\|_{L_t^\infty L_x^2([0, T] \times \mathbb{R}^3)} \lesssim E(u_0, f, T)^{1/2}.$$

In particular, we have

$$E_{\eta^4}(t) \lesssim E(u_0, f, T)$$

for all r, R ; inserting this back into (65) and integrating, we obtain that

$$\int_0^T X_1(t) dt \lesssim \left(\frac{A^2}{r^2} + \frac{A^6}{r^4} \right) T + E(u_0, f, T).$$

Sending $r, R \rightarrow \infty$ and using monotone convergence again, we conclude that

$$\|\nabla u\|_{L_t^2 L_x^2([0, T] \times \mathbb{R}^3)} \lesssim E(u_0, f, T)^{1/2},$$

and Lemma 8.1 follows. \square

We can bootstrap the proof of Lemma 8.1 as follows. *A posteriori*, we see that we may take $A \lesssim E(u_0, f, T)^{1/2}$. If we return to (65), we may then obtain

$$\partial_t (E_{\eta^4} + e)^{1/2} \lesssim e^{-1/2} \left(\frac{E(u_0, f, T)}{r^2} + \frac{E(u_0, f, T)^3}{r^4} \right) + a(t),$$

where $e > 0$ is an arbitrary parameter which we will optimise later. From the fundamental theorem of calculus, we then have

$$E_{\eta^4}^{1/2} \lesssim E_{\eta^4}(0)^{1/2} + e^{1/2} + e^{-1/2} \left(\frac{E(u_0, f, T)}{r^2} + \frac{E(u_0, f, T)^3}{r^4} \right) T + \|f\|_{L_t^1 L_x^2},$$

where the $L_t^1 L_x^2$ norm is over $[0, T] \times B(0, R)$; optimising in e , we conclude that

$$E_{\eta^4}^{1/2} \lesssim E_{\eta^4}(0)^{1/2} + \left(\frac{E(u_0, f, T)}{r^2} + \frac{E(u_0, f, T)^3}{r^4} \right)^{1/2} T^{1/2} + \|f\|_{L_t^1 L_x^2}.$$

Inserting this back into (65) and integrating, we also conclude that

$$\int_0^T X_1(t) dt \lesssim \left(E_{\eta^4}(0)^{1/2} + \left(\frac{E(u_0, f, T)}{r^2} + \frac{E(u_0, f, T)^3}{r^4} \right)^{1/2} T^{1/2} + \|f\|_{L_t^1 L_x^2} \right)^2.$$

Applying spatial translation invariance (29) to move the origin from 0 to an arbitrary point x_0 , we deduce an energy localisation result:

Theorem 8.2 (Local energy estimate). *Let (u, p, u_0, f, T) be a finite energy almost smooth solution with f divergence-free. Then for any $x_0 \in \mathbb{R}^3$ and any $0 < r < R/2$, one has*

$$\begin{aligned} & \|u\|_{L_t^\infty L_x^2([0, T] \times B(x_0, R-r))} + \|\nabla u\|_{L_t^2 L_x^2([0, T] \times B(x_0, R-r))} \\ & \lesssim \|u_0\|_{L_x^2(B(x_0, R))} + \|f\|_{L_t^1 L_x^2([0, T] \times B(x_0, R))} + \frac{E(u_0, f, T)^{1/2} T^{1/2}}{r} + \frac{E(u_0, f, T)^{3/2} T^{1/2}}{r^2}. \end{aligned} \quad (66)$$

Remark 8.3. One can verify that the estimate (66) is dimensionally consistent. Indeed, if L denotes a length scale, then $r, R, E(u_0, f)$ have the units of L , T has the units of L^2 , u has the units of L^{-1} , and all terms in (66) have the scaling of $L^{1/2}$. Note also that the global energy estimate 8.1 can be viewed as the limiting case of (66) when one sends r, R to infinity.

Remark 8.4. A minor modification of the proof of Theorem 8.2 allows one to replace the ball $B(x_0, R)$ by an annulus

$$B(x_0, R') \setminus B(x_0, R)$$

for some $0 < R < R'$ with $r < (R' - R)/2, R/2$, with the smaller ball $B(x_0, R - r)$ being replaced by the smaller annulus

$$B(x_0, R' - r) \setminus B(x_0, R + r).$$

The proof is essentially the same, except that the cutoff η has to be adapted to the two indicated annuli rather than to the two indicated balls; we omit the details. Sending $R' \rightarrow \infty$ using the monotone convergence theorem, we deduce in particular an external local energy estimate of the form

$$\begin{aligned} & \|u\|_{L_t^\infty L_x^2([0, T] \times (\mathbb{R}^3 \setminus B(x_0, R+r)))} + \|\nabla u\|_{L_t^2 L_x^2([0, T] \times (\mathbb{R}^3 \setminus B(x_0, R+r)))} \\ & \lesssim \|u_0\|_{L_x^2(\mathbb{R}^3 \setminus B(x_0, R))} + \|f\|_{L_t^1 L_x^2([0, T] \times (\mathbb{R}^3 \setminus B(x_0, R)))} + \frac{E(u_0, f, T)^{1/2} T^{1/2}}{r} + \frac{E(u_0, f, T)^{3/2} T^{1/2}}{r^2}, \end{aligned} \quad (67)$$

whenever $0 < r < R/2$.

Remark 8.5. The hypothesis that f is divergence-free can easily be removed using the symmetry (34), but then f needs to be replaced by Pf on the right-hand side of (66).

Remark 8.6. Theorem 8.2 can be extended without difficulty to the periodic setting, with the energy $E(u_0, f, T)$ being replaced by the periodic energy

$$E_L(u_0, f, T) := \frac{1}{2} \left(\|u_0\|_{L_x^2(\mathbb{R}^3/L\mathbb{Z}^3)} + \|f\|_{L_t^1 L_x^2([0, T] \times \mathbb{R}^3/L\mathbb{Z}^3)} \right)^2$$

as long as the radius R of the ball is significantly smaller than the period L of the solution, for example, $R < L/100$. The reason for this is that the analysis used to prove Theorem 8.2 takes place almost entirely inside the ball $B(x_0, R)$, and so there is almost no distinction between the finite energy and the periodic cases. The only place where there is any “leakage” outside of $B(x_0, R)$ is in the estimation of the term $X_{5,2}$, which involves the nonlocal commutator $[\Delta^{-1} \nabla^2, \eta^3]$. However, in the regime $R < L/100$, one easily verifies that the commutator essentially obeys the same sort of kernel bounds in the periodic setting as it does in the nonperiodic setting, and so the argument goes through as before. We omit the details.

Remark 8.7. [Theorem 8.2](#) asserts, roughly speaking, that if the energy of the data is small in a large ball, then the energy will remain small in a slightly smaller ball for future times T ; similarly, [\(67\)](#) asserts that if the energy of the data is small outside a ball, then the energy will remain small outside a slightly larger ball for future times T . Unfortunately, this estimate is not of major use for the purposes of establishing [Theorem 1.20](#), because energy is a supercritical quantity for the Navier–Stokes equation, and so smallness of energy (local or global) is not a particularly powerful conclusion. To achieve this goal, we will need a variant of [Theorem 8.2](#) in which the energy $\frac{1}{2} \int |u|^2$ is replaced by the *enstrophy* $\frac{1}{2} \int |\omega|^2$, which is subcritical and thus able to control the regularity of solutions effectively.

Remark 8.8. It should be possible to extend [Theorem 8.2](#) to certain classes of weak solutions, such as mild solutions or Leray–Hopf solutions, perhaps after assuming some additional regularity on the solution u . We will not pursue these matters here.

9. Bounded total speed

Let (u, p, u_0, f, T) be an almost smooth finite energy solution. Applying the Leray projection P to [\(3\)](#) (and using [Corollary 4.3](#)), we see that

$$\partial_t u = \Delta u + PB(u, u) + Pf \tag{68}$$

for almost all times t , where $B(u, v) = \mathbb{O}(\nabla(uv))$ was defined in [\(12\)](#). As all expressions here are tempered distributions, we thus have the Duhamel formula [\(11\)](#), which we rewrite here as

$$u(t) = e^{t\Delta} u_0 + \int_0^t e^{(t-t')\Delta} (P\mathbb{O}(\nabla(uu)) + Pf)(t') dt'. \tag{69}$$

One can then insert the *a priori* bounds from [Lemma 8.1](#) into [\(69\)](#) to obtain further *a priori* bounds on u in terms of the energy $E(u_0, f, T)$ (although, given that [\(53\)](#) was supercritical with respect to scaling, any further bounds obtained by this scheme must be similarly supercritical).

Many such bounds of this type already exist in the literature. For instance:¹⁷

- One can bound the vorticity $\omega := \nabla \times u$ in $L_t^\infty L_x^1$ norm [[Constantin 1990](#); [Qian 2009](#)].
- One can bound $\nabla^2 u$ in $L_{t,x}^{4/3, \infty}$ [[Constantin 1990](#); [Lions 1996](#)].
- More generally, for any $\alpha \geq 1$, one can bound $\nabla^\alpha u$ in $L_t^{4/(\alpha+1), \infty} L_x^{4/(\alpha+1), \infty}$ [[Vasseur 2010](#); [Choi and Vasseur 2011](#)].
- For any $k \geq 0$, one can bound $t^k \partial_t^k u$ in $L_{t,x}^2$ [[Chae 1992](#)].
- One can bound ∇u in $L_t^{1/2} L_x^\infty$ [[Foias et al. 1981](#)].
- For any $r \geq 0$ and $k \geq 1$, one can bound $D_t^r \nabla_x^s u$ in $L_t^{2/(4r+2k-1)} L_x^2$ [[Foias et al. 1981](#); [Doering and Foias 2002](#); [Duff 1990](#)].

¹⁷These bounds are usually localised in both time and space, or are restricted to the periodic setting, and some bounds were only established in the model case $f = 0$; some of these bounds also apply to weaker notions of solution than classical solutions. For the purposes of this exposition we will not detail these technicalities.

- For any $1 \leq m \leq \infty$, one can bound ω in $L_t^{2m/(4m-3)} L_x^{2m}$ [Gibbon 2012].
- One can bound moments of wave-number like quantities [Doering and Gibbon 2002; Cheskidov and Shvydkoy 2011].

In this section we present another *a priori* bound which will be absolutely crucial for our localisation arguments, and which (somewhat surprisingly) does not appear to be previously in the literature:

Proposition 9.1 (Bounded total speed). *Let (u, p, u_0, f, T) be a finite energy almost smooth solution. Then we have*

$$\|u\|_{L_t^1 L_x^\infty([0, T] \times \mathbb{R}^3)} \lesssim E(u_0, f, T)^{1/2} T^{1/4} + E(u_0, f, T). \quad (70)$$

We observe that the estimate (70) is dimensionally consistent with respect to the scaling (31). Indeed, if L denotes a length scale, then T scales like L^2 , u scales like L^{-1} , and E_0 scales like L , so both sides of (70) have the scaling of L .

Before we prove this proposition rigorously, let us first analyse equation (68) heuristically, using Littlewood–Paley projections, to get some feel of what kind of *a priori* estimates one can hope to establish purely from (68) and (53). For the simplicity of this exposition we shall assume $f = 0$. We consider a high-frequency component $u_N := P_N u$ of the velocity field u for some $N \gg 1$. Applying P_N to (68), and using the ellipticity of Δ to adopt the heuristic¹⁸ $P_N \Delta \sim -N^2 P_N$ and $P_N P \nabla \sim N P_N$, we arrive at the heuristic equation

$$\partial_t u_N = -N^2 u_N + \mathcal{O}(N P_N(u^2)).$$

Let us cheat even further and pretend that $P_N(u^2)$ is analogous to $u_N u_N$ (in practice, there will be more terms than this, but let us assume this oversimplification for the sake of discussion). Then we have

$$\partial_t u_N = -N^2 u_N + \mathcal{O}(N u_N^2).$$

Heuristically, this suggests that the high-frequency component u_N should quickly damp itself out into nothingness if $|u_N| \ll N$, but can exhibit nonlinear behaviour when $|u_N| \gg N$. Thus, as a heuristic, one can pretend that u_N has magnitude $\gg N$ on the regions where it is nonnegligible.

This heuristic, coupled with the energy bound (53), already can be used to informally justify many of the known *a priori* bounds on Navier–Stokes solutions. In particular, projecting (53) to the u_N component, one expects that

$$\|u_N\|_{L_t^2 L_x^2} \lesssim N^{-1} \quad (71)$$

(dropping the dependencies of constants on parameters such as E_0 and being vague about the space-time region on which the norms are being evaluated), which by Bernstein’s inequality implies that

$$\|u_N\|_{L_t^2 L_x^\infty} \lesssim N^{1/2}.$$

However, with the heuristic that $|u_N| \gg N$ on the support of u_N , we expect that

$$\|u_N\|_{L_t^1 L_x^\infty} \lesssim \frac{1}{N} \|u_N\|_{L_t^2 L_x^\infty}^2 \lesssim 1;$$

¹⁸One can informally justify this heuristic by inspecting the symbols of the Fourier multipliers appearing in these expressions.

summing in N (and ignoring the logarithmic divergence that results, which can in principle be recovered by using Bessel's inequality to improve upon (71)), we obtain a nonrigorous derivation of [Proposition 9.1](#).

We now turn to the formal proof of [Proposition 9.1](#). All space-time norms are understood to be over the region $[0, T] \times \mathbb{R}^3$ (and all spatial norms over \mathbb{R}^3) unless otherwise indicated. We abbreviate $E_0 := E(u_0, f, T)$. From (53) and (2) we have the bounds

$$\|u\|_{L_t^\infty L_x^2} \lesssim E_0^{1/2}, \quad (72)$$

$$\|\nabla u\|_{L_t^2 L_x^2} \lesssim E_0^{1/2}, \quad (73)$$

$$\|u_0\|_{L_x^2} + \|f\|_{L_t^1 L_x^2} \lesssim E_0^{1/2}. \quad (74)$$

We expand out u using (69). For the free term $e^{t\Delta}u_0$, one has by (18)

$$\|e^{t\Delta}u_0\|_{L_x^\infty} \lesssim t^{-3/4}\|u_0\|_{L_x^2}$$

for $t \in [0, T]$, so this contribution to (70) is acceptable by (74). In a similar spirit, we have

$$\|e^{(t-t')\Delta}Pf(t')\|_{L_x^\infty} \lesssim (t-t')^{-3/4}\|Pf(t')\|_{L_x^2} \lesssim (t-t')^{-3/4}\|f(t')\|_{L_x^2},$$

and so this contribution is also acceptable by the Minkowski and Young inequalities and (74).

It remains to show that

$$\left\| \int_0^t e^{(t-t')\Delta} \mathbb{O}(P\nabla(uu)(t')) dt' \right\|_{L_t^1 L_x^\infty} \lesssim E_0.$$

By Littlewood–Paley decomposition, the triangle inequality, and Minkowski's inequality, we can bound the left-hand side by

$$\lesssim \sum_N \int_0^T \int_0^t \|P_N e^{(t-t')\Delta} \mathbb{O}(P\nabla(uu)(t'))\|_{L_x^\infty} dt' dt.$$

Using (27) and bounding the first-order operator $P\nabla$ by N on the range of P_N , we may bound this by

$$\lesssim \sum_N \int_0^T \int_0^t \exp(-c(t-t')N^2) N \|P_N \mathbb{O}(uu)(t')\|_{L_x^\infty} dt' dt$$

for some $c > 0$; interchanging integrals and evaluating the t integral, this becomes

$$\lesssim \sum_N \int_0^T N^{-1} \|P_N \mathbb{O}(uu)(t')\|_{L_x^\infty} dt'. \quad (75)$$

We now apply the Littlewood–Paley trichotomy (see [Section 2](#)) and symmetry to write

$$P_N \mathbb{O}(uu) = \sum_{N_1 \sim N} \sum_{N_2 \lesssim N} P_N \mathbb{O}(u_{N_1} u_{N_2}) + \sum_{N_1 \gtrsim N} \sum_{N_2 \sim N_1} P_N \mathbb{O}(u_{N_1} u_{N_2}),$$

where $u_N := P_N u$. For N_1, N_2 in the first sum, we use Bernstein's inequality to estimate

$$\begin{aligned} \|P_N \mathbb{O}(u_{N_1} u_{N_2})\|_{L_x^\infty} &\lesssim \|u_{N_1}\|_{L_x^\infty} \|u_{N_2}\|_{L_x^\infty} \\ &\lesssim N_1^{3/2} \|u_{N_1}\|_{L_x^2} N_2^{3/2} \|u_{N_2}\|_{L_x^2} \\ &\lesssim N(N_2/N_1)^{1/2} \|\nabla u_{N_1}\|_{L_x^2} \|\nabla u_{N_2}\|_{L_x^2}. \end{aligned}$$

For N_1, N_2 in the second sum, we use Bernstein's inequality in a slightly different way to estimate

$$\begin{aligned} \|P_N \mathbb{O}(u_{N_1} u_{N_2})\|_{L_x^\infty} &\lesssim N^3 \|\mathbb{O}(u_{N_1} u_{N_2})\|_{L_x^1} \\ &\lesssim N^3 \|u_{N_1}\|_{L_x^2} \|u_{N_2}\|_{L_x^2} \\ &\lesssim N(N/N_1)^2 \|\nabla u_{N_1}\|_{L_x^2} \|\nabla u_{N_2}\|_{L_x^2}. \end{aligned}$$

Applying these bounds, we can estimate (75) by

$$\begin{aligned} &\lesssim \sum_N \sum_{N_1 \sim N} \sum_{N_2 \lesssim N} (N_2/N_1)^{1/2} \int_0^T \|\nabla u_{N_1}(t')\|_{L_x^2} \|\nabla u_{N_2}(t')\|_{L_x^2} dt' \\ &\quad + \sum_N \sum_{N_1 \gtrsim N} \sum_{N_2 \sim N_1} (N/N_1)^2 \int_0^T \|\nabla u_{N_1}(t')\|_{L_x^2} \|\nabla u_{N_2}(t')\|_{L_x^2} dt'. \end{aligned}$$

Performing the N summation first and then using Cauchy–Schwarz, one can bound this by

$$\lesssim \sum_{N_1 \gtrsim 1} \sum_{N_2 \lesssim N_1} (N_2/N_1)^{1/2} a_{N_1} a_{N_2} + \sum_{N_1 \gtrsim 1} \sum_{N_2 \sim N_1} a_{N_1} a_{N_2},$$

where

$$a_N := \|\nabla u_N\|_{L_t^2 L_x^2}.$$

But from (73) and Bessel's inequality (or the Plancherel theorem), one has

$$\sum_N a_N^2 \lesssim E_0,$$

and the claim (70) then follows from Schur's test (or Young's inequality).

Remark 9.2. An inspection of this argument reveals that the L_x^∞ norm in (70) can be strengthened to a Besov norm $(\dot{B}_1^{0,\infty})_x$, defined by

$$\|u\|_{(\dot{B}_1^{0,\infty})_x} := \sum_N \|P_N u\|_{L_x^\infty}.$$

Remark 9.3. An inspection of the proof of Proposition 9.1 reveals that the time-dependent factor $T^{1/4}$ on the right-hand side of Proposition 9.1 was only necessary in order to bound the linear components

$$e^{t\Delta} u_0 + \int_0^t e^{(t-t')\Delta} (Pf)(t') dt'$$

of the Duhamel formula (69). If one had some other means to bound these components in $L_t^1 L_x^\infty$ by a bound independent of T (for instance, if one had some further control on the decay of u_0 and f , such as

L_x^1 and $L_t^1 L_x^1$ bounds), then this would lead to a similarly time-independent bound in [Proposition 9.1](#), which could be useful for analysis of the long-time asymptotics of Navier–Stokes solutions (which is not our primary concern here).

Remark 9.4. It is worth comparing the (supercritical) control given by [Proposition 9.1](#) with the well-known (critical) Prodi–Serrin–Ladyzhenskaya regularity condition [[Prodi 1959](#); [Serrin 1963](#); [Ladyzhenskaya 1967](#); [Fabes et al. 1972](#); [Struwe 1988](#)], a special case of which (roughly speaking) asserts that smooth solutions to the Navier–Stokes system can be continued as long as u is bounded in $L_t^2 L_x^\infty$, and the equally well known (and also critical) regularity condition of Beale, Kato, and Majda [[1984](#)], which asserts that smooth solutions can be continued as long as the *vorticity*

$$\omega := \nabla \times u \tag{76}$$

stays bounded in $L_t^1 L_x^\infty$.

Remark 9.5. As pointed out by the anonymous referee, one can also obtain $L_t^1 L_x^\infty$ bounds on the velocity field u by a Gagliardo–Nirenberg type interpolation between the $L_t^{1/2} L_x^\infty$ bound on ∇u from [[Foaş et al. 1981](#)] with the $L_t^2 L_x^6$ bound on u arising from the energy inequality and Sobolev embedding.

Although we will not need it in this paper, [Proposition 9.1](#) when combined with the Picard well-posedness theorem for ODE yields the following immediate corollary, which may be of use in future applications:

Corollary 9.6 (Existence of material coordinates). *Let (u, p, u_0, f, T) be a finite energy smooth solution. Then there exists a unique smooth map $\Phi : [0, T] \times \mathbb{R}^3 \rightarrow \mathbb{R}^3$ such that*

$$\Phi(0, x) = x$$

for all $x \in \mathbb{R}^3$, and

$$\partial_t \Phi(t, x) = u(\Phi(t, x))$$

for all $(t, x) \in [0, T] \times \mathbb{R}^3$, and furthermore $\Phi(t) : \mathbb{R}^3 \rightarrow \mathbb{R}^3$ is a diffeomorphism for all $t \in [0, T]$. Finally, one has

$$|\Phi(t, x) - x| \lesssim E(u_0, f, T)^{1/2} T^{1/4} + E(u_0, f, T)$$

for all $(t, x) \in [0, T] \times \mathbb{R}^3$.

Remark 9.7. One can extend the results in this section to the periodic case, as long as one assumes normalised pressure and imposes the additional condition $T \leq L^2$, which roughly speaking ensures that the periodic heat kernel behaves enough like its nonperiodic counterpart that estimates such as (18) are maintained; we omit the details. (Without normalised pressure, the Galilean invariance (33) shows that one cannot hope to bound the $L_t^1 L_x^\infty$ norm of u by the initial data, and even energy estimates do not work any more.) When the inequality $T \leq L^2$ fails, one can still obtain estimates (but with weaker bounds) by using the crude observation that a solution which is periodic with period L is also periodic with period kL for any positive integer k , and choosing k to be the first integer such that $T \leq (kL)^2$.

10. Enstrophy localisation

The purpose of this section is to establish a subcritical analogue of [Theorem 8.2](#), in which the energy $\frac{1}{2} \int |u|^2$ is replaced by the enstrophy $\frac{1}{2} \int |\omega|^2$. Because the latter quantity is not conserved, we will need a smallness condition on the initial local enstrophy; however, the initial *global* enstrophy is allowed to be arbitrarily large (or even infinite).

Theorem 10.1 (Enstrophy localisation). *Let (u, p, u_0, f, T) be a finite energy almost smooth solution. Let $B(x_0, R)$ be a ball such that*

$$\|\omega_0\|_{L_x^2(B(x_0, R))} + \|\nabla \times f\|_{L_t^1 L_x^2([0, T] \times B(x_0, R))} \leq \delta \quad (77)$$

for some $\delta > 0$, where $\omega_0 := \nabla \times u_0$ is the initial vorticity. Assume the smallness condition

$$\delta^4 T + \delta^5 E(u_0, f, T)^{1/2} T \leq c \quad (78)$$

for some sufficiently small absolute constant $c > 0$ (independent of all parameters). Let $0 < r < R/2$ be a quantity such that

$$r > C(E(u_0, f, T) + E(u_0, f, T)^{1/2} T^{1/4} + \delta^{-2}) \quad (79)$$

for some sufficiently large absolute constant C (again independent of all parameters). Then

$$\|\omega\|_{L_x^\infty L_x^2([0, T] \times B(x_0, R-r))} + \|\nabla \omega\|_{L_t^2 L_x^2([0, T] \times B(x_0, R-r))} \lesssim \delta.$$

Remark 10.2. Once again, this theorem is dimensionally consistent (and so one could use [\(31\)](#) to normalise one of the nondimensionless parameters above to equal 1 if desired). Indeed, if L is a unit of length, then u has the units of L^{-1} , ω has the units of L^{-2} , $E(u_0, f, T)$, r , R have the units of L , T has the units of L^2 , and δ has the units of $L^{-1/2}$ (so in particular $\delta^4 T$ and $\delta^5 E(u_0, f, T)^{1/2} T$ are dimensionless).

Remark 10.3. The smallness of $\delta^4 T$ also comes up, not coincidentally, as a condition in the local well-posedness theory for the Navier–Stokes at the level of H^1 ; see [\(46\)](#). The smallness of $\delta^5 E(u_0, f, T)^{1/2} T$ is a more artificial condition, and it is possible that a more careful argument would eliminate it, but we will not need to do so for our applications. For future reference, it will be important to note the fact that δ is permitted to be large in the above theorem, so long as the time T is small.

Remark 10.4. A variant to [Theorem 10.1](#) can also be deduced from the result¹⁹ in [[Caffarelli et al. 1982](#), Theorem D]. Here, instead of assuming a small L^2 condition on the enstrophy, one needs to assume smallness of quantities such as $\int_{\mathbb{R}^3} (|u_0(x)|^2 / |x - x_0|) dx$ for all sufficiently large x_0 , and then regularity results are obtained outside of a sufficiently large ball in space-time.

We now prove the theorem. Let (u, p, u_0, f, T) , $B(x_0, R)$, δ , r be as in the theorem. We may use spatial translation symmetry [\(29\)](#) to normalise $x_0 = 0$. We assume $c > 0$ is a sufficiently small absolute constant, and then assume $C > 0$ is a sufficiently large constant (depending on c). We abbreviate $E_0 := E(u_0, f, T)$.

In principle, this is a subcritical problem, because the local enstrophy $\frac{1}{2} \int_{B(x_0, R)} |\omega|^2$ (or regularised versions thereof) is subcritical with respect to scaling [\(31\)](#). As such, standard energy methods should

¹⁹We thank the anonymous referee for this observation.

in principle suffice to keep the enstrophy small for small times (using the smallness condition (78), of course). The main difficulty is that the local enstrophy is not fully *coercive*: it controls ω (and, to a lesser extent, u) inside $B(x_0, R)$, but not outside $B(x_0, R)$; while we do have some global control of the solution thanks to the energy estimate (Lemma 8.1), this is supercritical and thus needs to be used sparingly. We will therefore expend a fair amount of effort to prevent our estimates from “leaking” outside $B(x_0, R)$; in particular, one has to avoid the use of nonlocal singular integrals (such as the Leray projection or the Biot–Savart law) and work instead with more local techniques such as integration by parts. This will inevitably lead to some factors that blow up as one approaches the boundary of $B(x_0, R)$ (actually, for technical reasons, we will be using a slightly smaller ball $B(x_0, R'(t))$ as our domain). It turns out, however, that thanks to a moderate amount of harmonic analysis, these boundary factors can (barely) be controlled if one chooses exactly the right type of weight function to define the local enstrophy (it has to be Lipschitz continuous, but no better).

We turn to the details. We will need an auxiliary initial radius $R' = R'(0)$ in the interval $[R - r/4, R]$ which we will choose later (by a pigeonholing argument). Given this R' , we then define a time-dependent radius function

$$R'(t) := R' - \frac{1}{c} \int_0^t \|u(s)\|_{L_x^\infty(\mathbb{R}^3)} ds.$$

From Proposition 9.1 one has

$$R'(t) \geq R' - O_c(E_0 + E_0^{1/2} T^{1/4}),$$

and thus (by (79)) one has

$$R'(t) \geq R - r/2$$

if the constant C in (79) is sufficiently large depending on c . The reason we introduce this rapidly shrinking radius is that we intend to “outrun” all difficulties caused by the transport component of the Navier–Stokes equation when we deploy the energy method. Note that the bounded total speed property (Proposition 9.1) prevents us from running the radius down to zero when we do this.

We introduce a time-varying Lipschitz continuous cutoff function

$$\eta(t, x) = \min(\max(0, c^{-0.1} \delta^2 (R'(t) - |x|)), 1).$$

This function is supported on the ball $B(0, R'(t))$ and equals one on $B(0, R'(t) - c^{0.1} \delta^{-2})$, and is radially decreasing; in particular, from (79), we see that η is supported on $B(0, R)$ and equals 1 on $B(0, R - r)$ if C is large enough. As t increases, this cutoff shrinks at speed $(1/c)\|u(t)\|_{L_x^\infty(\mathbb{R}^3)}$, leading to the useful pointwise estimate

$$\partial_t \eta(t, x) \leq -\frac{1}{c} \|u(t)\|_{L_x^\infty(\mathbb{R}^3)} |\nabla_x \eta(t, x)|, \quad (80)$$

which we will use later in this argument to control transport-like terms in the energy estimate (or more precisely, the enstrophy estimate).

Remark 10.5. It will be important that η is Lipschitz continuous but no better; Lipschitz is the minimal regularity for which one can still control the heat flux term (see Y_3 below), but is also the maximal regularity for which there is enough coercivity to control the nonlinear term (see Y_6 below). The argument

is in fact remarkably delicate, necessitating a careful application of harmonic analysis techniques (and in particular, a Whitney decomposition of the ball).

We introduce the localised enstrophy

$$W(t) := \frac{1}{2} \int_{\mathbb{R}^3} |\omega(t, x)|^2 \eta(t, x) dx. \quad (81)$$

From the hypothesis (77) one has the initial condition

$$W(0) \lesssim \delta^2, \quad (82)$$

and to obtain the proposition, it will suffice to show that

$$W(t) \lesssim_c \delta^2 \quad (83)$$

for all $t \in [0, T]$.

As u is almost smooth, W is C_t^1 . As in Section 8, we will compute the derivative $\partial_t W$. We first take the curl of (3) to obtain the well-known *vorticity equation*

$$\partial_t \omega + (u \cdot \nabla) \omega = \Delta \omega + \mathbb{O}(\omega \nabla u) + \nabla \times f. \quad (84)$$

This leads to the *enstrophy equation*

$$\partial_t \frac{1}{2} |\omega|^2 + (u \cdot \nabla) \frac{1}{2} |\omega|^2 = \Delta \left(\frac{1}{2} |\omega|^2 \right) - |\nabla \omega|^2 + \mathbb{O}(\omega \omega \nabla u) + \omega \cdot (\nabla \times f).$$

All terms in this equation are smooth. Integrating this equation against the Lipschitz, compactly supported η and integrating by parts as in Section 8 (interpreting derivatives of η in a distributional sense), we conclude that

$$\partial_t W = -Y_1 - Y_2 + Y_3 + Y_4 + Y_5 + Y_6, \quad (85)$$

where Y_1 is the dissipation term

$$Y_1 := \int_{\mathbb{R}^3} |\nabla \omega|^2 \eta,$$

Y_2 is the recession term

$$Y_2 := -\frac{1}{2} \int_{\mathbb{R}^3} |\omega|^2 \partial_t \eta,$$

Y_3 is the heat flux term

$$Y_3 := \frac{1}{2} \int_{\mathbb{R}^3} |\omega|^2 \Delta \eta,$$

Y_4 is the transport term

$$Y_4 := \frac{1}{2} \int_{\mathbb{R}^3} |\omega|^2 u \cdot \nabla \eta,$$

Y_5 is the forcing term

$$Y_5 := \int_{\mathbb{R}^3} \omega \cdot (\nabla \times f) \eta,$$

and Y_6 is the nonlinear term

$$Y_6 := \int_{\mathbb{R}^3} \mathbb{O}(\omega \omega \nabla u) \eta.$$

The term Y_1 is nonnegative, and will be needed to control some of the other terms. The term Y_2 is also nonnegative; by (80) we see that

$$\int_{\mathbb{R}^3} |\omega|^2 |\nabla \eta| \lesssim c \|u(t)\|_{L_x^\infty(\mathbb{R}^3)} Y_2. \quad (86)$$

We skip the heat flux term Y_3 for now and use (86) to bound the transport term Y_4 by

$$|Y_4| \lesssim c Y_2. \quad (87)$$

Now we turn to the forcing term Y_5 . By Cauchy–Schwarz and (81), we have

$$|Y_5| \lesssim W^{1/2} a(t),$$

where

$$a(t) := \|\nabla \times f\|_{L_x^2(B(0, R))}.$$

Note from (77) that

$$\int_0^T a(t) dt \lesssim \delta. \quad (88)$$

We return now to the heat flux term Y_3 . Computing the distributional Laplacian²⁰ of η in polar coordinates, we see that

$$Y_3 \lesssim b(t),$$

where $b(t) = b_{R'}(t)$ is the quantity

$$b(t) := c^{-0.1} \delta^2 R^2 \int_{S^2} |\omega(t, R'(t)\alpha)|^2 d\alpha + c^{-0.2} \delta^4 \int_{R'(t) - c^{0.1} \delta^{-2} \leq |x| \leq R'(t)} |\omega(t, x)|^2 dx,$$

and $d\alpha$ is surface measure on the unit sphere S^2 . (Note that while $\Delta \eta$ also has a component on the sphere $|x| = R'(t) - c^{0.1} \delta^{-2}$, this component is negative and thus can be discarded.)

To control $b(t)$, we take advantage of the freedom to choose R' . From Fubini's theorem and a change of variables, we see that

$$\int_{R-\frac{r}{4}}^R \int_0^T b_{R'}(t) dt dR' \lesssim c^{-0.1} \delta^2 \int_0^T \int_{\mathbb{R}^3} |\omega(t, x)|^2 dx.$$

From Lemma 8.1, the right-hand side is $O(\delta^2 E_0 / c^{0.1})$. Thus, by the pigeonhole principle, we may select a radius R' such that

$$\int_0^T b(t) dt \lesssim \frac{\delta^2 E_0}{c^{0.1} r},$$

and in particular, by (79),

$$\int_0^T b(t) dt \lesssim \delta^2 \quad (89)$$

if C is large enough.

²⁰Alternatively, if one wishes to avoid distributions, one can regularise η by a small epsilon parameter to become smooth, compute the Laplacian of the regularised term, and take limits as epsilon goes to zero. One can also rescale either R or δ (but not both) to equal 1 to simplify the computations.

Henceforth we fix R' so that (89) holds. We now turn to the most difficult term, namely the nonlinear term Y_6 . Morally speaking, the ∇u term in Y_6 has the “same strength” as ω , and so Y_6 is heuristically as strong as

$$\int_{\mathbb{R}^3} \mathbb{O}(\omega^3) \eta.$$

A standard Whitney decomposition of the support of η , followed by rescaled versions of the Sobolev inequality, bounds this latter expression by

$$O\left(\left(\int_{\mathbb{R}^3} |\omega|^2 \eta\right)^{1/2} \left(\int_{\mathbb{R}^3} |\nabla \omega|^2 \eta\right)\right).$$

If we could similarly bound Y_6 by this expression by an analogous argument, this would greatly simplify the argument below. Unfortunately, the relationship between ∇u and ω is rather delicate (especially when working relative to the weight η), and we have to perform a much more involved analysis (though still ultimately one which is inspired by the preceding argument).

We turn to the details. We fix t and work in the domain

$$\Omega := B(0, R'(t)).$$

We apply a Whitney-type decomposition, covering Ω by a boundedly overlapping collection of balls $B_i = B(x_i, r_i)$ with radius

$$r_i := \frac{1}{100} \min(\text{dist}(x_i, \partial\Omega), c^{0.1}/\delta^2).$$

In particular, we have

$$\eta \sim c^{-0.1} \delta^2 r_i \tag{90}$$

on $B(x_i, 10r_i)$. We can then bound

$$|Y_6| \lesssim c^{-0.1} \delta^2 \sum_i r_i \int_{B_i} |\omega|^2 |\nabla u|.$$

The first step is to convert ∇u into an expression that only involves ω (modulo lower-order terms), while staying inside the domain Ω . To do this, we first observe from the divergence-free nature of u that

$$\Delta u = \nabla \times \nabla \times u = \nabla \times \omega.$$

Let ψ_i be a smooth cutoff to the ball $3B_i := B(x_i, 3r_i)$ that equals 1 on $2B_i := B(x_i, 2r_i)$. On $2B_i$, we thus have the local Biot–Savart law

$$u = \mathbb{O}(\Delta^{-1} \nabla(\psi_i \omega)) + v,$$

where v is harmonic on $2B_i$. In particular, from Sobolev embedding one has

$$\|v\|_{L_x^2(2B_i)} \lesssim \|\psi_i \omega\|_{L_x^{6/5}(\mathbb{R}^3)} + \|u\|_{L_x^2(2B_i)}.$$

From Hölder’s inequality one has

$$\|\psi_i \omega\|_{L_x^{6/5}(\mathbb{R}^3)} \lesssim r_i \|\omega\|_{L_x^2(2B_i)},$$

while from the mean value principle for harmonic functions one has

$$\|\nabla v\|_{L_x^\infty(B_i)} \lesssim r_i^{-5/2} \|v\|_{L_x^2(2B_i)}.$$

We conclude that

$$\|\nabla v\|_{L_x^\infty(B_i)} \lesssim r_i^{-3/2} \|\omega\|_{L_x^2(2B_i)} + r_i^{-5/2} \|u\|_{L_x^2(2B_i)},$$

and we thus have the pointwise estimate

$$|\nabla u| \lesssim |\nabla \Delta^{-1} \nabla(\psi_i \omega)| + r_i^{-3/2} \|\omega\|_{L_x^2(2B_i)} + r_i^{-5/2} \|u\|_{L_x^2(2B_i)}$$

on B_i . We can thus bound $|Y_6| \leq Y_{6,1} + Y_{6,2}$, where

$$Y_{6,1} \lesssim c^{-0.1} \delta^2 \sum_i r_i \int_{B_i} |\omega|^2 F_i \tag{91}$$

and

$$F_i := |\nabla \Delta^{-1} \nabla(\psi_i \omega)| + r_i^{-3/2} \|\omega\|_{L_x^2(2B_i)}$$

and

$$Y_{6,2} \lesssim c^{-0.1} \delta^2 \sum_i r_i^{-3/2} \|u\|_{L_x^2(2B_i)} \int_{B_i} |\omega|^2.$$

Let us first deal with $Y_{6,2}$, which is the only term not locally controlled by the vorticity alone. If the ball B_i is contained in the annular region

$$\{x \in \Omega : |x| \geq R'(t) - c^{0.1} \delta^{-2}\},$$

which is the region where η is not constant, then we use Hölder to get the bound

$$r^{-3/2} \|u\|_{L_x^2(2B_i)} \lesssim \|u\|_{L_x^\infty(\mathbb{R}^3)}$$

and observe that $c^{-0.1} \delta^2 = |\nabla \eta|$ on B_i . Thus, by (86), the contribution of this term to $Y_{6,2}$ is $O(c^{0.9} Y_2)$. If instead the ball B_i intersects the ball $B(0, R'(t) - c^{0.1} \delta^{-2})$, then $r_i \sim c^{0.1} \delta^{-2}$ and $\eta \sim 1$ on B_i , and we use Lemma 8.1 to obtain the bound

$$r_i^{-3/2} \|u\|_{L_x^2(2B_i)} \lesssim c^{-0.15} \delta^3 E_0^{1/2},$$

and then by (81), (78) the contribution of this case is

$$O(c^{-0.25} \delta^5 E_0^{1/2} W) = O(c^{0.75} W/T);$$

and thus

$$Y_{6,2} \lesssim c^{0.9} Y_2 + c^{0.75} W/T.$$

Now we turn to $Y_{6,1}$. From Plancherel's theorem we have

$$\|\nabla \Delta^{-1} \nabla(\psi_i \omega)\|_{L_x^2(\mathbb{R}^3)} \lesssim \|\psi_i \omega\|_{L_x^2(\mathbb{R}^3)} \lesssim \|\omega\|_{L_x^2(2B_i)},$$

and thus

$$\|F_i\|_{L_x^2(B_i)} \lesssim \|\omega\|_{L_x^2(2B_i)}.$$

From Hölder’s inequality we thus have

$$Y_{6,1} \lesssim c^{-0.1} \delta^2 \sum_i r_i^{3/2} \|\omega\|_{L_x^6(B_i)}^2 \|\omega\|_{L_x^2(2B_i)}.$$

To deal with this, we let w_i denote the averages

$$w_i := \left(\frac{1}{|3B_i|} \int_{3B_i} |\omega|^2 \right)^{1/2};$$

then

$$\|\omega\|_{L_x^2(2B_i)} \lesssim r_i^{3/2} w_i.$$

Also, from the Sobolev inequality one has

$$\|\omega\|_{L_x^6(B_i)} \lesssim \|\omega\psi_i\|_{L_x^6(\mathbb{R}^3)} \lesssim \|\nabla(\omega\psi_i)\|_{L_x^2(\mathbb{R}^3)} \lesssim \|\nabla\omega\|_{L_x^2(3B_i)} + r_i^{-1} \|\omega\|_{L_x^2(3B_i)} \lesssim \|\nabla\omega\|_{L_x^2(3B_i)} + r_i^{1/2} w_i,$$

and thus

$$Y_{6,1} \lesssim c^{-0.1} \delta^2 \sum_i r_i^3 w_i \|\nabla\omega\|_{L_x^2(3B_i)}^2 + c^{-0.1} \delta^2 \sum_i r_i^4 w_i^3. \quad (92)$$

To deal with the first term of (92), observe from (81) and (90) that

$$\sum_i r_i^4 w_i^2 \lesssim c^{0.1} \delta^{-2} W, \quad (93)$$

and in particular

$$w_i \lesssim c^{0.05} \delta^{-1} W^{1/2} r_i^{-2} \quad (94)$$

for all i . We may thus bound

$$c^{-0.1} \delta^2 \sum_i r_i^3 w_i \|\nabla\omega\|_{L_x^2(3B_i)}^2 \lesssim c^{-0.05} \delta W^{1/2} \sum_i r_i \|\nabla\omega\|_{L_x^2(3B_i)}^2,$$

which by (90) and the bounded overlap of the B_i is

$$\lesssim c^{0.05} \delta^{-1} W^{1/2} \int_{\Omega} |\nabla\omega|^2 \eta \lesssim c^{0.05} \delta^{-1} W^{1/2} Y_1.$$

The second term of (92), $c^{-0.1} \delta^2 \sum_i r_i^4 w_i^3$, is trickier to handle. Call a ball “large” if its radius is at least $10^{-4} c^{-0.1} \delta^{-2}$ (say), and “small” otherwise. To deal with the small balls we use the Poincaré inequality.

From this inequality, we see in particular that

$$\left| \left(\frac{1}{|3B_i|} \int_{3B_i} |\omega|^2 \right)^{1/2} - \left(\frac{1}{|3B_j|} \int_{3B_j} |\omega|^2 \right)^{1/2} \right| \lesssim \left(r_i^{-1} \int_{10B_i} |\nabla\omega|^2 \right)^{1/2}$$

whenever B_i, B_j intersect. (Indeed, the Poincaré inequality implies that both terms in the left-hand side are within $O((r_i^{-1} \int_{10B_i} |\nabla\omega|^2)^{1/2})$ of $|(1/|10B_i|) \int_{10B_i} \omega|$.) In other words, we have

$$|w_i - w_j| \lesssim r_i^{-1/2} \left(\int_{10B_i} |\nabla\omega|^2 \right)^{1/2} \quad (95)$$

whenever B_i, B_j intersect.

Now for any small ball B_i , we may assign a “parent” ball $B_{p(i)}$ which touches the ball but has radius at least 1.001 (say) times as large as that of B_i . We may iterate this until we reach a large ball $B_{a(i)}$, and write

$$w_i \leq w_{a(i)} + \sum_{k \geq 0} |w_{p^k(i)} - w_{p^{k+1}(i)}|,$$

where the sum is over all k for which $p^{k+1}(i)$ is well-defined; note that this inequality also holds for large balls if we set $a(i) = i$. Taking cubes and using Hölder’s inequality, we obtain

$$w_i^3 \lesssim w_{a(i)}^3 + \sum_{k \geq 0} (1+k)^{10} |w_{p^k(i)} - w_{p^{k+1}(i)}|^3,$$

and so we can bound $c^{-0.1} \delta^2 \sum_i r_i^4 w_i^3$ by

$$\lesssim c^{-0.1} \delta^2 \sum_i r_i^4 w_{a(i)}^3 + c^{-0.1} \delta^2 \sum_{k \geq 0} (1+k)^{10} \sum_i r_i^4 |w_{p^k(i)} - w_{p^{k+1}(i)}|^3.$$

If one fixes a large ball B_j , one easily checks that $\sum_{i:a(i)=j} r_i^4 \lesssim r_j^4$, and thus

$$c^{-0.1} \delta^2 \sum_i r_i^4 w_{a(i)}^3 \lesssim c^{-0.1} \delta^2 \sum_{j:r_j > 10^{-4} c^{0.1} \delta^{-2}} r_j^4 w_j^3;$$

applying (94) and (93), we thus have

$$c^{-0.1} \delta^2 \sum_i r_i^4 w_{a(i)}^3 \lesssim c^{-0.25} \delta^5 W^{1/2} \sum_j r_j^4 w_j^2 \lesssim c^{-0.15} \delta^3 W^{3/2}.$$

Similarly, if one fixes a small ball B_j , one verifies that

$$\sum_{k \geq 0} (1+k)^{10} \sum_{i:p^k(i)=j} r_i^4 \lesssim r_j^4,$$

and thus

$$c^{-0.1} \delta^2 \sum_{k \geq 0} (1+k)^{10} \sum_i r_i^4 |w_{p^k(i)} - w_{p^{k+1}(i)}|^3 \lesssim c^{-0.1} \delta^2 \sum_{j:r_j \leq 10^{-4} c^{0.1} \delta^{-2}} r_j^4 |w_j - w_{p(j)}|^3.$$

From (94) (once) and (95) (twice) one has

$$|w_j - w_{p(j)}|^3 \lesssim c^{0.05} \delta^{-1} W^{1/2} r_j^{-3} \int_{10B_j} |\nabla \omega|^2,$$

and so we may bound the preceding expression by

$$\lesssim c^{-0.05} \delta W^{1/2} \sum_j r_j \int_{10B_j} |\nabla \omega|^2,$$

which by (90) and the bounded overlap of the B_j can be bounded by

$$\lesssim c^{0.05} \delta^{-1} W^{1/2} \int_{\Omega} |\nabla \omega|^2 \eta \lesssim c^{0.05} \delta^{-1} W^{1/2} Y_1.$$

Putting the $Y_{6,1}$ bounds together, we conclude that

$$Y_{6,1} \lesssim c^{-0.15} \delta^3 W^{3/2} + c^{0.05} \delta^{-1} W^{1/2} Y_1;$$

collecting the bounds for Y_1, \dots, Y_6 , we thus have

$$\partial_t W \leq -Y_1 + O(c^{0.05} \delta^{-1} W^{1/2} Y_1 + c^{-0.15} \delta^3 W^{3/2} + c^{0.75} W/T + a(t) W^{1/2} + b(t)).$$

To solve this differential inequality we use the continuity method. Suppose that $0 \leq T' \leq T$ is a time for which

$$\sup_{t \in [0, T']} W(t) \leq c^{-0.01} \delta^2. \quad (96)$$

Then, if c is small enough, we can absorb the $O(c^{0.05} \delta^{-1} W^{1/2} Y_1)$ term by the $-Y_1$ term, and can also use this bound and (78) to obtain

$$c^{-0.15} \delta^3 W^{3/2} \lesssim c^{-0.155} \delta^4 W \lesssim c^{0.75} W/T$$

and

$$a(t) W^{1/2} \lesssim c^{-0.005} \delta a(t).$$

We thus have

$$\partial_t W \lesssim c^{0.75} W/T + c^{-0.005} \delta a(t) + b(t).$$

From Gronwall's inequality and (82), (88), (89), we thus have

$$\sup_{t \in [0, T']} W(t) \lesssim c^{-0.005} \delta^2.$$

For c a small enough absolute constant, this is (slightly) better than the hypothesis (96), and so from the continuity method (and (82)), we conclude that

$$\sup_{t \in [0, T]} W(t) \lesssim c^{-0.005} \delta^2,$$

and the claim (83) follows. The proof of [Theorem 10.1](#) is now complete.

Remark 10.6. As with [Remark 8.4](#), we may adapt the proof of [Theorem 10.1](#) to an annulus, replacing the ball $B(x_0, R)$ with an annulus $B(x_0, R') \setminus B(x_0, R)$ for some $0 < R < R'$ with $0 < r < R/2, (R' - R)/2$, and replacing the smaller ball $B(x_0, R - r)$ with the smaller annulus $B(x_0, R' - r) \setminus B(x_0, R + r)$. To do this, one has to replace the cutoff η (which was shrinking inside the ball $B(x_0, R)$ towards $B(x_0, R - r)$) with a slightly more complicated cutoff (which is shrinking inside the annulus $B(x_0, R') \setminus B(x_0, R)$ towards the smaller annulus $B(x_0, R' - r) \setminus B(x_0, R + r)$). However, aside from this detail, the proof method is essentially identical and is omitted. Sending R' to infinity and using the monotone convergence theorem, we may in fact replace the annulus $B(x_0, R') \setminus B(x_0, R)$ with the exterior region $\mathbb{R}^3 \setminus B(x_0, R)$, and the annulus $B(x_0, R' - r) \setminus B(x_0, R + r)$ with $\mathbb{R}^3 \setminus B(x_0, R + r)$.

[Theorem 10.1](#) asserts, roughly speaking, that if the H_x^1 norm of the data is small on a ball, then for a quantitative amount of later time, the H_x^1 norm of the solution remains small on a slightly smaller ball.

As the H^1 norm is subcritical, we expect this sort of result to persist to higher regularities, in the spirit of [Serrin 1962]. It is therefore unsurprising that this is indeed the case:

Proposition 10.7 (Higher regularity). *Let (u, p, u_0, f, T) be a finite energy almost smooth solution with $T \leq T_*$. Let $B(x_0, R)$, η , δ , r obey the conditions (77), (78), (79) from Theorem 10.1. Then for any compact subset K in the interior of $B(x_0, R - r)$ and any $k \geq 1$, one can bound*

$$\|\nabla^k u\|_{L_t^\infty L_x^2([0, T] \times K)} + \|\nabla^{k+1} u\|_{L_t^2 L_x^2([0, T] \times K)} \lesssim_{k, K, E(u_0, f, T), \delta, T_*, R, A_k} 1,$$

where

$$A_k := \sum_{j=0}^k \|\nabla^j u_0\|_{L_x^2(B(x_0, R))} + \|\nabla^j f\|_{L_t^\infty L_x^2([0, T] \times B(x_0, R))}.$$

In particular, one has

$$\|u\|_{X^k([0, T] \times K)} \lesssim_{k, K, E(u_0, f, T), \delta, T_*, R, A_k} 1.$$

Proof. We allow all implied constants to depend on $k, K, E(u_0, f, T), \delta, T_*, R, A_k$. We introduce a compact set

$$K \subset K_1 \subset K_2 \subset K_3 \subset K_4 \subset K_5 \subset B(x_0, R - r),$$

with each set lying in the interior of the next set. Let η be a smooth function supported on K_2 that equals 1 on K_1 ; we allow implied constants to depend on η .

We begin with the $k = 1$ case. From Theorem 10.1 one already has

$$\|\omega\|_{L_t^\infty L_x^2([0, T] \times K_1)} + \|\nabla \omega\|_{L_t^2 L_x^2([0, T] \times K_1)} \lesssim 1.$$

To pass from ω to u , we use integration by parts. Since $\omega = \nabla \times u$ and u is divergence-free, a standard integration by parts shows that

$$\frac{1}{2} \int_{\mathbb{R}^3} |\omega|^2 \eta = \int_{\mathbb{R}^3} |\nabla u|^2 \eta + \int_{\mathbb{R}^3} \mathcal{O}(|u|^2 \nabla^2 \eta).$$

By Lemma 8.1, the error term is $O(1)$, and so we have

$$\int_K |\nabla u|^2 \lesssim 1.$$

Similarly, by replacing ω and u by their derivatives, we also see that

$$\frac{1}{2} \int_{\mathbb{R}^3} |\nabla \omega|^2 \eta = \int_{\mathbb{R}^3} |\nabla^2 u|^2 \eta + \int_{\mathbb{R}^3} \mathcal{O}(|\nabla u|^2 \nabla^2 \eta).$$

By Lemma 8.1, the error term is $O(1)$ after integration in time, and so we also have

$$\int_0^T \int_K |\nabla^2 u|^2 dx dt \lesssim 1$$

as desired.

We now turn to the $k = 2$ case. This is the most difficult, as we currently only control regularities that are half a derivative better than the critical regularity (which would place u in $H_x^{1/2}$), and wish to

boost this to three halves of a derivative above critical; this requires at least two iterations of the Duhamel formula. The arguments will be analogous to the regularity arguments in [Theorem 5.1](#) or [Lemma 5.5](#). By [\(68\)](#) we see that $u\eta$ obeys the truncated equation

$$\partial_t(\eta u) - \Delta(\eta u) = \eta \mathbb{O}(P\nabla(uu)) + \eta Pf + \mathbb{O}(\nabla u \nabla \eta) + \mathbb{O}(u \nabla^2 \eta) \quad (97)$$

for almost all t . Meanwhile, from the $k = 1$ case and [Lemma 8.1](#), we already have the estimates

$$\|u\|_{L_t^\infty L_x^2([0, T] \times \mathbb{R}^3)} + \|\nabla u\|_{L_t^\infty L_x^2([0, T] \times K_4)} + \|\nabla^2 u\|_{L_t^2 L_x^2([0, T] \times K_4)} \lesssim 1, \quad (98)$$

and from the definition of A_2 , we have

$$\|\nabla^j u_0\|_{L_x^2(B(x_0, R))} + \|\nabla^j f\|_{L_t^\infty L_x^2([0, T] \times B(x_0, R))} \lesssim 1 \quad (99)$$

for $j = 0, 1, 2$.

We claim that all terms on the right-hand side of [\(97\)](#) have an $L_t^4 L_x^2([0, T] \times \mathbb{R}^3)$ norm of $O(1)$. The only difficult term here is $\eta P \mathbb{O}(\nabla(uu))$; the other three terms on the right-hand side are easily estimated in $L_t^4 L_x^2$ (and even in $L_t^2 L_x^2$) using [\(98\)](#) and [\(99\)](#). We now estimate

$$\|\eta \mathbb{O}(P\nabla(uu))\|_{L_t^4 L_x^2([0, T] \times \mathbb{R}^3)}.$$

We split $uu = \tilde{\eta}uu + (1 - \tilde{\eta})uu$, where $\tilde{\eta}$ is a smooth cutoff supported on K_4 that equals 1 on K_3 . For the contribution of the nonlocal portion $(1 - \tilde{\eta})uu$, one can use the smoothness of the kernel of the operator P away from the origin to bound this contribution by $\lesssim \|\mathbb{O}(uu)\|_{L_t^4 L_x^1([0, T] \times \mathbb{R}^3)}$, which is acceptable by [\(98\)](#); for future reference, we note that this argument bounds this contribution in $L_t^2 L_x^2$ norm as well as in $L_t^4 L_x^2$ norm. For the local portion $\tilde{\eta}uu$, we discard the η and P projections and bound this by

$$\lesssim \|\mathbb{O}(\nabla(\tilde{\eta}uu))\|_{L_t^4 L_x^2([0, T] \times \mathbb{R}^3)}.$$

But this is acceptable by [\(24\)](#).

We have now placed the right-hand side of [\(97\)](#) in $L_t^4 L_x^2([0, T] \times \mathbb{R}^3)$ with norm $O(1)$. Meanwhile, from [\(99\)](#), the initial data $u_0\eta$ is in $H_x^2(\mathbb{R}^3)$ with norm $O(1)$. Applying the energy estimate [\(23\)](#), we conclude that

$$\|u\eta\|_{L_t^\infty H_x^{3/2-\sigma}([0, T] \times \mathbb{R}^3)} + \|u\eta\|_{L_t^2 H_x^{5/2-\sigma}([0, T] \times \mathbb{R}^3)} \lesssim_\sigma 1$$

for any $\sigma > 0$. A similar argument (shifting the compact sets) also gives

$$\|u\eta'\|_{L_t^\infty H_x^{3/2-\sigma}([0, T] \times \mathbb{R}^3)} + \|u\eta'\|_{L_t^2 H_x^{5/2-\sigma}([0, T] \times \mathbb{R}^3)} \lesssim_\sigma 1,$$

where η' is a smooth function supported on K_5 that equals 1 on K_4 . In particular, by Sobolev embedding, on $[0, T] \times K_4$, u is in $L_t^\infty L_x^{12}$, ∇u is in $L_t^2 L_x^{12} \cap L_t^\infty L_x^{12/5}$, and $\nabla^2 u$ is in $L_t^2 L_x^{12/5}$, which together with [\(98\)](#) and the Hölder inequality now allows one to conclude that $\mathbb{O}(\nabla(\tilde{\eta}uu))$ has an $L_t^2 H_x^1([0, T] \times \mathbb{R}^3)$ norm of $O(1)$. Repeating the previous arguments, we now conclude that the right-hand side of [\(97\)](#) lies

in $L_t^2 H_x^1([0, T] \times \mathbb{R}^3)$ with norm $O(1)$, and hence by (22),

$$\|\eta u\|_{L_t^\infty H_x^2([0, T] \times \mathbb{R}^3)} + \|\eta u\|_{L_t^2 H_x^3([0, T] \times \mathbb{R}^3)},$$

which gives the $k = 2$ case.

The higher k cases are proven by similar arguments, but are easier as we now have enough regularity to place u in $L_t^\infty L_x^\infty([0, T] \times K_5)$ with norm $O(1)$; we leave the details to the reader. (For instance, to establish the $k = 3$ case, one can verify using the estimates already obtained from the $k = 2$ case that the right-hand side of (97) has an $L_t^2 H_x^1([0, T] \times \mathbb{R}^3)$ norm of $O(1)$. \square)

Remark 10.8. As in Remark 9.7, one can extend the results here to the periodic setting so long as one has $T \leq L^2$ and $R \leq L$; we omit the details.

For our application to constructing Leray–Hopf weak solutions, we will need a generalisation of Theorem 10.1 to the case when one has hyperdissipation. More precisely, we introduce a small hyperdissipation parameter $\varepsilon > 0$, and consider solutions $(u^{(\varepsilon)}, p^{(\varepsilon)}, u_0, f, T)$ to the regularised Navier–Stokes equation, which are defined precisely as with the usual concept of a Navier–Stokes solution, but with (3) replaced by the regularised variant

$$\partial_t u^{(\varepsilon)} + (u^{(\varepsilon)} \cdot \nabla) u^{(\varepsilon)} = \Delta u^{(\varepsilon)} - \varepsilon \Delta^2 u^{(\varepsilon)} - \nabla p^{(\varepsilon)} + f. \quad (100)$$

With hyperdissipation, the global regularity problem becomes much easier (the energy is now subcritical rather than supercritical), and indeed it is not difficult to use energy methods (see, for example, [Lions 1969]) to show the existence of a unique almost smooth finite energy solution to this regularised equation $(u^{(\varepsilon)}, p^{(\varepsilon)}, u_0, f, T)$ from any given smooth finite energy data (u_0, f, T) . The energy estimate in Lemma 8.1 remains true in this case (uniformly in ε), and one easily verifies that one obtains an additional estimate

$$\varepsilon \int_0^T \int_{\mathbb{R}^3} |\nabla^2 u(t, x)|^2 dt dx \lesssim E(u_0, f, T) \quad (101)$$

in this hyperdissipative setting. One can also verify (with a some tedious effort) that Proposition 9.1 also holds in this hyperdissipative setting as long as ε is sufficiently small, basically because the hyperdissipative heat operators $e^{t(\Delta - \varepsilon \Delta^2)}$ obey essentially the same estimates (18), (27) as $e^{t\Delta}$ if $0 \leq t \leq T$ and ε is sufficiently small depending on T ; we omit the details.

One can define the vorticity $\omega^{(\varepsilon)} := \nabla \times u^{(\varepsilon)}$ of a regularised solution as before. This vorticity obeys an equation almost identical to (84), but with an additional hyperdissipative term $-\varepsilon \nabla^2 \omega^{(\varepsilon)}$ on the right-hand side. One can then repeat the proof of Theorem 10.1 with this additional term. Integrating by parts a large number of times, one obtains a similar decomposition to (85) for the derivative of the localised enstrophy, but with the addition of a negative term $-\varepsilon \int_{\mathbb{R}^3} |\nabla^2 \omega|^2 \eta$ on the right-hand side, plus some boundary terms which are bounded by $\tilde{b}(t)$, where

$$\tilde{b}(t) := \sum_{r=R'(t), R'(t)-c^{0.1}\delta^{-2}} \varepsilon c^{-0.1} \delta^2 R^2 \int_{S^2} |\nabla \omega(t, r\alpha)|^2 d\alpha + \varepsilon c^{-0.2} \delta^4 \int_{R'(t)-c^{0.1}\delta^{-2} \leq |x| \leq R'(t)} |\nabla \omega(t, x)|^2 dx$$

is a hyperdissipative analogue of $b(t)$. By using the same averaging argument used to bound $\int_0^T b(t) dt$

for typical R' , one can also simultaneously obtain a comparable bound for $\int_0^T \tilde{b}(t)dt$ (taking advantage of the additional estimate (101)). The rest of the argument in [Theorem 10.1](#) works with essentially no changes; we omit the details. The proof of [Proposition 10.7](#) is also essentially identical, after one notes that energy estimates such as (22) continue to hold in the hyperdissipative setting. Summarising, we obtain:

Proposition 10.9. *[Theorem 10.1](#) and [Proposition 10.7](#) continue to hold in the presence of hyperdissipation, uniformly in the limit $\varepsilon \rightarrow 0$.*

11. Consequences of enstrophy localisation

We now give a number of applications of the enstrophy localisation result, [Theorem 10.1](#). Many of these applications resemble existing results in the literature, but with weaker decay hypotheses on the initial data and solution (in particular, we will usually only assume either finite energy or finite H^1 norm); the main point is that the localisation afforded by [Theorem 10.1](#) can significantly reduce the need to assume any stronger decay hypotheses.

We begin with the observation that finite energy smooth solutions automatically have bounded enstrophy if the initial data has bounded enstrophy:

Corollary 11.1 (Bounded enstrophy). *Let (u, p, u_0, f, T) be an almost smooth, finite energy solution, such that the initial data (u_0, f, T) has finite H^1 norm. Then $u \in X^1([0, T] \times \mathbb{R}^3)$; in particular, (u, p, u_0, f, T) is an H^1 solution.*

Proof. Let $\delta > 0$ be small enough (depending on $E(u_0, f, T), T$) that the condition (78) holds. As (u_0, f, T) has finite H^1 norm, we have

$$\|\omega_0\|_{L_x^2(\mathbb{R}^3)} + \|\nabla \times f\|_{L_t^1 L_x^2([0, T] \times \mathbb{R}^3)} < \infty.$$

By the monotone convergence theorem, we thus have for R sufficiently large that

$$\|\omega_0\|_{L_x^2(\mathbb{R}^3 \setminus B(0, R))} + \|\nabla \times f\|_{L_t^1 L_x^2([0, T] \times (\mathbb{R}^3 \setminus B(0, R)))} \leq \delta.$$

Applying [Theorem 10.1](#) (inverted as in [Remark 10.6](#)), we conclude that

$$\|\omega\|_{L_x^\infty L_x^2([0, T] \times (\mathbb{R}^3 \setminus B(0, R+r)))} + \|\nabla \omega\|_{L_t^2 L_x^2([0, T] \times (\mathbb{R}^3 \setminus B(0, R+r)))} \lesssim \delta$$

for some finite radius r , if R is sufficiently large; in particular, ω lies in $L_t^\infty L_x^2 \cap L_t^2 H_x^1$ in the exterior region $[0, T] \times (\mathbb{R}^3 \setminus B(0, R+r))$. On the other hand, as u is almost smooth, ω also lies in $L_t^\infty L_x^2 \cap L_t^2 H_x^1$ in the interior region $[0, T] \times B(0, R+r+1)$ (say). Gluing these two bounds together, we conclude that

$$\omega \in L_t^\infty L_x^2 \cap L_t^2 H_x^1([0, T] \times \mathbb{R}^3);$$

meanwhile, from [Lemma 8.1](#) one has

$$u \in L_t^\infty L_x^2 \cap L_t^2 H_x^1([0, T] \times \mathbb{R}^3).$$

Since u is divergence-free and $\omega = \nabla \times u$, the claim then follows from Fourier analysis. \square

Remark 11.2. From [Corollary 5.8](#) we know that smooth solutions to the Navier–Stokes solutions can be continued in time as long as the H^1 norm remains bounded. However, [Corollary 11.1](#) certainly does not allow one to solve the global regularity problem for Navier–Stokes, because the proof heavily relies on the solution u being *complete* rather than *incomplete*, and thus it is (almost) smooth all the way up to the final time T , and not just smooth on $[0, T)$. Instead, what [Corollary 11.1](#) does is to show that the solution from H^1 data is well-behaved when one is sufficiently close to spatial infinity; in particular, it does not prevent turbulent behaviour in bounded regions of space-time.

Remark 11.3. If $(u, p, u_0, 0, T)$ is an almost smooth homogeneous finite energy solution, then by [Lemma 8.1](#) we see that $u(t) \in H_x^1(\mathbb{R}^3)$ for almost every time $t \in [0, T]$. Applying the time translation symmetry (30) for a small time shift t_0 , we can then convert the finite energy data to H^1 data, and then by [Corollary 11.1](#), we conclude that in fact $u(t) \in H_x^1(\mathbb{R}^3)$ for *all* nonzero times $t \in (0, T]$, and furthermore that $u(t)$ is bounded in H_x^1 as soon as t is bounded away from zero.

Since H^1 almost smooth solutions with normalised pressure are automatically H^1 mild solutions, for which uniqueness was established in [Theorem 5.4](#), we thus have uniqueness in the almost smooth finite energy category from smooth H^1 data:

Corollary 11.4 (Unconditional uniqueness). *Let (u_0, f, T) be smooth H^1 data. Then there is at most one almost smooth finite energy solution (u, p, u_0, f, T) with this data and with normalised pressure.*

This result resembles the standard “weak-strong uniqueness” results in the literature, such as those in [\[Prodi 1959; Serrin 1963; Germain 2006; 2008\]](#). The main novelty here is the lack of decay hypotheses beyond the finite energy hypothesis; note that the almost smoothness of the solution gives plenty of integrability on compact regions of space, but does not imply any global integrability in space.

Remark 11.5. We conjecture that one still retains uniqueness even if the data (u_0, f, T_*) is merely smooth and finite energy, rather than smooth and H^1 . Note from [Lemma 8.1](#) that $u(t)$ has finite $H_x^1(\mathbb{R}^3)$ norm for almost every time t , which in principle allows one to enforce uniqueness after any given positive time (in the homogeneous case $f = 0$, at least), but it is not clear to the author how to prevent instantaneous failure of uniqueness at the initial time $t = 0$ with only a smooth finite energy hypothesis on the initial data. It may however be possible to adapt the “weak-strong” uniqueness results of Germain [\[2006; 2008\]](#) to this category, perhaps in combination with the local H^1 control given by [Theorem 10.1](#).

We now use the enstrophy localisation result to study solutions as they approach a (potential) blowup time T_* .

Proposition 11.6 (Uniform smoothness outside a ball). *Let (u, p, u_0, f, T_*^-) be an incomplete almost smooth H^1 solution with normalised pressure for all times $0 < T < T_*$. Then there exists a ball $B(0, R)$ such that*

$$u, p, f, \partial_t u \in L_t^\infty C_x^k([0, T_*) \times K) \tag{102}$$

for all $k \geq 0$ and all compact subsets K of $\mathbb{R}^3 \setminus B(0, R)$.

We remark that similar results were obtained by Caffarelli, Kohn, and Nirenberg [1982] assuming additional spatial decay hypotheses on the data at infinity, and in particular that $\int_{\mathbb{R}^3} |u_0(0, x)|^2 |x| dx < \infty$. The main novelty in this proposition is that one only assumes square-integrability of u_0 and its first derivatives, without any further decay assumption.

Proof. From the argument in the proof of Corollary 11.1 (noting that the bounds are uniform for all times T in a compact set), one can already find a ball $B(0, R_0)$ for which

$$u \in X^1([0, T_*] \times (\mathbb{R}^3 \setminus B(0, R_0))).$$

Using Proposition 10.7, we then conclude the existence of a larger ball $B(0, R)$ such that

$$u \in X^k([0, T_*] \times K)$$

for all $k \geq 1$ and all compact subsets K of $\mathbb{R}^3 \setminus B(0, R)$. From this, Sobolev embedding, and (9) (using the smoothness of the kernel of $\nabla^k \Delta^{-1}$ away from the origin), we obtain (102) for u, p, f as desired. If one then applies (3) and solves for $\partial_t u$, one obtains the bound for $\partial_t u$ also. \square

Remark 11.7. From (102) one can *continuously* extend u up to the portion $\{T_*\} \times (\mathbb{R}^3 \setminus B(0, R))$ of the boundary (compare the partial regularity theory in [Caffarelli et al. 1982]). However, we were unable to demonstrate that u could be extended *smoothly* up to the boundary (or even that $\partial_t u$ is continuous in time at the boundary). The problem is due to the nonlocal effects of pressure; the solution u could be blowing up at time T_* in the interior of $B(0, R)$, leading (via (9)) to time oscillations of the pressure in K (which cannot be directly damped out by the smoothness of the Δ^{-1} kernel, which only attenuates *spatial* oscillations), which by (3) could lead to time oscillations of the solution u in K . Indeed, as Theorem 1.12 shows, these time oscillations can have a nontrivial effect on the regularity of the solution.

Remark 11.8. For future reference, we observe that Proposition 11.6 did not require the full space-time smoothness on f ; it would suffice to have $f \in L_t^\infty C_x^k([0, T_*] \times K)$ for all $k \geq 0$ and compact K in order to obtain the conclusion (102). This is because at no stage in the argument was it necessary to differentiate f in time.

In a similar spirit, we may construct Leray–Hopf weak solutions that are spatially smooth outside of a ball for any fixed time T . More precisely, define a *Leray–Hopf weak solution* (u, p, u_0, f, T) to smooth finite energy data (u_0, f, T) to be a distributional solution $u \in X^0([0, T] \times \mathbb{R}^3)$ to (3) (after expressing this equation in divergence form) which is continuous in time in the weak topology of $L_x^2(\mathbb{R}^3)$, and which obeys the energy inequality

$$\frac{1}{2} \|u(t)\|_{L_x^2(\mathbb{R}^3)}^2 + \int_0^t \|\nabla u(t)\|_{L_x^2(\mathbb{R}^3)}^2 dx \leq E(u_0, f, T). \quad (103)$$

The existence of such solutions was famously demonstrated in [Leray 1934] for arbitrary finite energy data (u_0, f, T) ; the singularities of these solutions were analysed in a vast number of papers, which are too numerous to cite here, but we will point out in particular the seminal work [Caffarelli et al. 1982].

Our main regularity result for Leray–Hopf solutions is as follows.

Proposition 11.9 (Existence of partially smooth Leray–Hopf weak solutions). *Let (u_0, f, T) be smooth H^1 data. Then there exists a Leray–Hopf weak solution (u, p, u_0, f, T) to the given data and a ball $B(0, R)$ such that u is spatially smooth in $[0, T] \times (\mathbb{R}^3 \setminus B(0, R))$ (that is, for each $t \in [0, T]$, $u(t)$ is smooth outside of $B(0, R)$).*

Again, similar results were obtained in [Caffarelli et al. 1982] under stronger decay hypotheses on the initial data. We also remark that weak solutions which were only locally of finite energy, from data of uniformly locally finite energy, were constructed in [Lemarié-Rieusset 1999]; the ability to localise the weak solution construction in this fashion is similar in spirit to the results in the proposition.

Proof. (Sketch) We use a standard hyperdissipation²¹ regularisation argument. Let $\varepsilon > 0$ be a small parameter, and consider the almost smooth finite-energy solution $(u^{(\varepsilon)}, p^{(\varepsilon)}, u_0, f, T)$ to the regularised Navier–Stokes system (100), which can be shown to exist by energy methods. By Proposition 10.9, we can extend Theorem 10.1 and Proposition 10.7 (and thence Proposition 11.6) to these regularised solutions $u^{(\varepsilon)}$, with bounds that are uniform in ε as $\varepsilon \rightarrow 0$. As a consequence, we can find a ball $B(0, R)$ independent of ε such that for every compact set K outside of $B(0, R)$ and every $k \geq 0$, $\nabla^k u^{(\varepsilon)}$ lies in $L_t^\infty L_x^\infty([0, T_*] \times K)$ uniformly in N . If we then extract a weak limit point u of the $u^{(\varepsilon)}$, then by standard arguments one verifies that u is a Leray–Hopf weak solution which is spatially smooth outside of $B(0, R)$. \square

Remark 11.10. As before, we are unable to demonstrate regularity of u in time due to potential nonlocal effects caused by the pressure, which could in principle cause singularities inside $B(0, R)$ to create time singularities outside of $B(0, R)$.

Remark 11.11. Uniqueness of Leray–Hopf solutions remains a major unsolved problem, for which we have nothing new to contribute; in particular, we do not assert that *all* Leray–Hopf solutions from smooth data obey the conclusions of Proposition 11.9. However, if (u_0, f, ∞) is globally defined smooth H^1 data, the argument above gives a single global Leray–Hopf weak solution (u, p, u_0, f, ∞) with the property that, for each finite time $T < \infty$, there exists a radius $R_T < \infty$ such that u is smooth in $[0, T] \times (\mathbb{R}^3 \setminus B(0, R))$. If we restrict to the case $f = 0$, then from (103) we see that $\|\nabla u(t)\|_{L_x^2(\mathbb{R}^3)}$ must become arbitrarily small along some sequence of times $t = t_n$ going to infinity. If $\|\nabla u(t)\|_{L_x^2(\mathbb{R}^3)}$ is small enough depending on $E(u_0, 0, \infty)$, then standard perturbation theory arguments (see, for example, [Kato 1984]) allow one to obtain a smooth, bounded enstrophy solution from the data $u(t)$ on $(t, +\infty)$, which by the uniqueness theory of Serrin [1963] must match the Leray–Hopf weak solution u on $(t, +\infty)$. As such, we conclude in the homogeneous smooth H^1 case that one can construct a global Leray–Hopf weak solution which is spatially smooth outside of a compact subset of space-time $[0, +\infty) \times \mathbb{R}^3$. Again, we emphasise that this global weak solution need not be unique.

²¹ It may also be possible to use other regularisation methods here, such as velocity regularisation, to construct the Leray–Hopf weak solution; however, due to the delicate nature of the proof of the localised enstrophy estimate (Theorem 10.1), we were not able to verify that this estimate remained true in the velocity-regularised setting, uniformly in the regularisation parameter, due to the less favourable vorticity equation in this setting.

12. Smooth H^1 solutions

The purpose of this section is to establish [Theorem 1.20\(iii\)](#). To do this, we will need the ability to localise smooth divergence-free vector fields, as follows.

Lemma 12.1 (Localisation of divergence-free vector fields). *Let $T > 0$, $0 < R_1 < R_2 < R_3 < R_4$, and let $u : [0, T] \times (B(0, R_4) \setminus B(0, R_1)) \rightarrow \mathbb{R}^3$ be spatially smooth and divergence-free, such that*

$$u, \partial_t u \in L_t^\infty C_x^k([0, T] \times (B(0, R_4) \setminus B(0, R_1)))$$

for all $k \geq 0$ and

$$\int_{|x|=r} u(t, x) \cdot n \, d\alpha(x) = 0 \quad (104)$$

for all $R_1 < r < R_4$ and $t \in [0, T]$, where n is the outward normal and $d\alpha$ is surface measure. Then there exists a spatially smooth and divergence-free vector field $\tilde{u} : [0, T] \times (B(0, R_4) \setminus B(0, R_1)) \rightarrow \mathbb{R}^3$ which agrees with u on $[0, T] \times (B(0, R_2) \setminus B(0, R_1))$ but vanishes on $[0, T] \times (B(0, R_4) \setminus B(0, R_3))$. Furthermore, we have

$$\tilde{u}, \partial_t \tilde{u} \in L_t^\infty C_x^k([0, T] \times (B(0, R_4) \setminus B(0, R_1)))$$

for all $k \geq 0$.

Finally, if we have

$$1 \leq 2R_2 \leq R_3 \lesssim R_2,$$

then we have the more quantitative bound

$$\|\tilde{u}\|_{L_t^\infty H^k([0, T] \times (B(0, R_4) \setminus B(0, R_1)))} \lesssim_k \|u\|_{L_t^\infty H^{k+1}([0, T] \times (B(0, R_4) \setminus B(0, R_1)))} \quad (105)$$

for any k . (This latter property will come in handy in the next section.)

Note that the hypothesis (104) is necessary, as can be seen from Stokes' theorem. Lemmas of this type first appear in [\[Bogovskii 1980\]](#).

Proof. One can obtain this lemma as a consequence of the machinery of compactly supported divergence-free wavelets [\[Lemarie-Rieusset 1992\]](#), but for the convenience of the reader we give a self-contained proof here.

Let X denote the vector space of all divergence-free smooth functions $u : B(0, R_4) \setminus B(0, R_1) \rightarrow \mathbb{R}^3$ obeying the mean zero condition

$$\int_{|x|=r} u(x) \cdot n \, d\alpha(x) = 0 \quad (106)$$

for all $R_1 < r < R_4$, and such that $\|u\|_{C^k([0, R_4] \setminus B(0, R_1))} < \infty$ for all k . It will suffice to construct a linear transformation $P : X \rightarrow X$ that is bounded²² from C^{k+2} to C^k , that is,

$$\|Pu\|_{C^k([0, R_4] \setminus B(0, R_1))} \lesssim_{R_1, R_2, R_3, R_4, k} \|u\|_{C^{k+2}([0, R_4] \setminus B(0, R_1))}$$

²²One can reduce this loss of regularity by working in more robust spaces than the classical C^k spaces, such as Sobolev spaces H^s or Hölder spaces $C^{k, \alpha}$, but we will not need to do so here.

for all $k \geq 0$, and such that Pu equals u on $B(0, R_2) \setminus B(0, R_1)$ and vanishes on $B(0, R_4) \setminus B(0, R_3)$, as one can then simply define $\tilde{u}(t) := P\tilde{u}(t)$ for each $t \in [0, T)$.

We now construct P . We work in polar coordinates $x = r\alpha$ with $R_1 \leq r \leq R_4$ and $\alpha \in S^2$ (thus avoiding the coordinate singularity at the origin), and decompose $u(r, \alpha)$ as the sum of a radial vector field $u_r(r, \alpha)\alpha$ for some scalar field u_r and an angular vector field $u_\alpha(r, \alpha)$ which is orthogonal to α ; thus, for fixed r , $u_\alpha(r)$ can be viewed as a smooth vector field on the unit sphere S^2 (that is, a smooth section of the tangent bundle of S^2). The divergence-free condition on u in these coordinates then reads

$$\partial_r u_r(r) + \frac{1}{r} \nabla_\alpha \cdot u_\alpha(r) = 0, \quad (107)$$

while the mean-zero condition (106) reads

$$\int_{S^2} u_r(r, \alpha) d\alpha = 0.$$

Note that either of these conditions implies that $\partial_r u_r(r)$ has mean zero on S^2 for each r . From (107) and Hodge theory, we see that

$$u_\alpha(r) = r \Delta_\alpha^{-1} \nabla_\alpha \partial_r u_r(r) + v(r),$$

where Δ_α^{-1} inverts the Laplace–Beltrami operator Δ_α on smooth mean-zero functions on S^2 and $v(r)$ is a smooth divergence-free vector field on S^2 that varies smoothly with r .

Let $\eta : [R_1, R_4] \rightarrow \mathbb{R}^+$ be a smooth function that equals 1 on $[R_1, R_2]$ and vanishes on $[R_3, R_4]$. Set

$$\tilde{u}_r := \eta(r)u_r$$

and

$$\tilde{u}_\alpha(r) = r \Delta_\alpha^{-1} \nabla_\alpha \partial_r \tilde{u}_r(r) + \eta(r)v(r)$$

and

$$Tu := \tilde{u} := \tilde{u}_r \alpha + \tilde{u}_\alpha.$$

One then easily verifies that \tilde{u} is smooth and divergence-free and obeys (106), depends linearly on u , equals u on $B(0, R_2) \setminus B(0, R_1)$, and vanishes on $B(0, R_4) \setminus B(0, R_3)$. It is also not difficult (using the fundamental solution of Δ_α^{-1}) to see that T maps C^{k+2} to C^k (with some room to spare). The claim follows.

Finally, we prove (105). It suffices to show that

$$\|Tu\|_{H^k(B(0, R_3) \setminus B(0, R_2))} \lesssim_k 1$$

whenever $k \geq 0$, and $u \in X$ is such that

$$\|u\|_{H^{k+2}(B(0, R_4) \setminus B(0, R_1))} \lesssim 1.$$

Henceforth all spatial norms will be on $B(0, R_3) \setminus B(0, R_2)$, and all implied constants may depend on k . As u has an H^{k+1} norm of $O(1)$, u_r and hence \tilde{u}_r has an H^{k+1} norm of $O(1)$ also. As for \tilde{u}_α , we observe from the Leibniz rule that

$$\tilde{u}_\alpha = \eta u_\alpha + (r \partial_r \eta(r)) \Delta_\alpha^{-1} \nabla_\alpha u_r(r).$$

As u has an H^{k+1} norm of $O(1)$, we know $r^{-i} \nabla_\alpha^i \partial_r^j u_\alpha$ has an L^2 norm of $O(1)$ whenever $i + j \leq k + 1$, which (using elliptic regularity in the angular variable) implies that $r^{-i} \nabla_\alpha^i \partial_r^j \tilde{u}_\alpha$ has an L^2 norm of $O(1)$ whenever $i + j \leq k$. This gives $\tilde{u} = \tilde{u}_r + \tilde{u}_\alpha$, an H^k norm of $O(1)$, as claimed. \square

We can now establish [Theorem 1.20\(iii\)](#):

Theorem 12.2. *Suppose [Conjecture 1.9](#) is true. Then [Conjecture 1.19](#) is true.*

Proof. In view of [Corollary 5.8](#), it suffices to show that if (u, p, u_0, f, T_*^-) is an incomplete H^1 mild solution up to time T_* , with u_0, f spatially smooth in the sense of [Conjecture 1.19](#), then u does not blow up in enstrophy norm; thus

$$\limsup_{t \rightarrow T_*^-} \|u(t)\|_{H_x^1(\mathbb{R}^3)} < \infty.$$

Let $R > 0$ be a sufficiently large radius. By arguing as in [Corollary 11.1](#), we have

$$u \in L_t^\infty H_x^1(\mathbb{R}^3 \setminus B(0, R)),$$

and thus the blowup must be localised in space:

$$\limsup_{t \rightarrow T_*^-} \|u(t)\|_{H_x^1(B(0, R))} < \infty. \quad (108)$$

By [Proposition 11.6](#) and [Remark 11.8](#) (and increasing R if necessary), we also have

$$u, p, f, \partial_t u \in L_t^\infty C_x^k([0, T_*) \times (B(0, 5R) \setminus B(0, 2R))) \quad (109)$$

for all $k \geq 0$. From Stokes' theorem and the divergence-free nature of u , we also have

$$\int_{|x|=r} u(t, x) \cdot n \, d\alpha(x) = 0$$

for all $r > 0$ and $t \in [0, T)$. Applying [Lemma 12.1](#), we can then find a spatially smooth divergence-free vector field $\tilde{u} : [0, T) \times (B(0, 5R) \setminus B(0, 2R)) \rightarrow \mathbb{R}^3$ which agrees with u on $B(0, 3R) \setminus B(0, 2R)$ and vanishes outside of $B(0, 4R)$, with

$$\tilde{u}, \partial_t \tilde{u} \in L_t^\infty C_x^k(B(0, 5R) \setminus B(0, 2R)) \quad (110)$$

for all $k \geq 0$. We then extend \tilde{u} by zero outside of $B(0, 5R)$ and by u inside of $B(0, 2R)$; then \tilde{u} is now smooth on all of $[0, T) \times \mathbb{R}^3$.

Let η be a smooth function supported on $B(0, 5R)$ that equals 1 on $B(0, 4R)$. We define a new forcing term $\tilde{f} : [0, T) \times \mathbb{R}^3 \rightarrow \mathbb{R}$ by the formula

$$\tilde{f} := \partial_t \tilde{u} + (\tilde{u} \cdot \nabla) \tilde{u} - \Delta \tilde{u} + \nabla(p\eta); \quad (111)$$

then \tilde{f} is spatially smooth and supported on $B(0, 5R)$ and agrees with f on $B(0, 3R)$. From this and [\(110\)](#), [\(109\)](#) we easily verify that

$$\tilde{f} \in L_t^\infty H_x^1([0, T_*) \times \mathbb{R}^3).$$

Note from taking divergences in (111) and using the compact support of $p\eta$, \tilde{u} , \tilde{f} that

$$p\eta = -\Delta^{-1}((\tilde{u} \cdot \nabla)\tilde{u}) + \Delta^{-1}\nabla \cdot \tilde{f}.$$

Thus, $(\tilde{u}, p\eta, \tilde{u}(0), \tilde{f}, T_*^-)$ is an incomplete H^1 pressure-normalised (and hence mild) solution with all components supported in $B(0, 5R)$. If we then choose a period L larger than $10R$, we may embed $B(0, 5R)$ inside $\mathbb{R}^3/L\mathbb{Z}^3$ and obtain an incomplete periodic smooth solution

$$(\iota(\tilde{u}), \iota(p\eta), \iota(\tilde{u}(0)), \iota(\tilde{f}), T_*^-, L),$$

where we use $\iota(f)$ to denote the extension by zero of a function f supported in $B(0, 5R)$, after embedding the latter in $\mathbb{R}^3/L\mathbb{Z}^3$. By construction, we then have

$$\iota(\tilde{f}) \in L_t^\infty H_x^1([0, T_*] \times \mathbb{R}^3/L\mathbb{Z}^3).$$

As $\{T_*\}$ has measure zero, we may arbitrarily extend \tilde{f} to $[0, T_*] \times \mathbb{R}^3/L\mathbb{Z}^3$ while staying in $L_t^\infty H_x^1$. Applying either [Conjecture 1.9](#) (and the uniqueness component to [Theorem 5.1](#)) or [Conjecture 1.10](#), we conclude that

$$\iota(\tilde{u}) \in L_t^\infty H_x^1([0, T_*] \times \mathbb{R}^3/L\mathbb{Z}^3),$$

which implies (since u and \tilde{u} agree on $B(0, R)$) that

$$u \in L_t^\infty H_x^1([0, T_*] \times B(0, R)),$$

which contradicts [\(108\)](#). The claim follows. \square

Observe that if we omit the embedding of $B(0, 5R)$ in $\mathbb{R}^3/L\mathbb{Z}^3$ in the preceding argument, we can also deduce [Conjecture 1.19](#) from [Conjecture 1.18](#). Since [Conjecture 1.19](#) clearly implies [Conjecture 1.18](#) as a special case, we obtain [Theorem 1.20\(iii\)](#).

Remark 12.3. The referee has pointed out a variant of the argument above using the partial regularity theory of Caffarelli, Kohn, and Nirenberg [1982], which allows one to partially reverse the above implications, and in particular deduce [Conjecture 1.8](#) from [Conjecture 1.19](#). We sketch the argument as follows. Assume [Conjecture 1.19](#), and assume for contradiction that [Conjecture 1.8](#) fails; thus, there is a periodic solution with smooth inhomogeneous data which first develops singularities at some finite time T , and in particular at some location (T, x_0) . We may extend the solution beyond this time as a weak solution. Applying a periodic version of the theory in [Caffarelli et al. 1982], we see that the set of singularities has zero one-dimensional parabolic measure, which among other things implies that the set of radii $r > 0$ such that the solution is singular at (T, x) for some x with $|x - x_0| = r$ has measure zero. Because of this, one can find radii $r_2 > r_1 > 0$ such that the solution is smooth in the annular region $\{(t, x) : 0 \leq t \leq T; r_1 \leq |x - x_0| \leq r_2\}$. By smoothly truncating the solution u to this annulus as in the proof of [Theorem 12.2](#), one can then create a nonperiodic H^1 mild solution to the inhomogeneous Navier–Stokes equation with spatially smooth data which develops a singularity at (T, x_0) while remaining smooth up to time T , contradicting [Conjecture 1.19](#) (when combined with standard uniqueness and regularity results, such as those in [Theorem 5.4](#)).

13. Smooth finite energy solutions

In this section we establish [Theorem 1.20\(v\)](#). It is trivial that [Conjecture 1.14](#) implies [Conjecture 1.13](#), so it suffices to establish:

Theorem 13.1. *Suppose that [Conjecture 1.13](#) is true. Then [Conjecture 1.14](#) is true.*

Proof. Let $(u_0, 0, T)$ be smooth homogeneous finite energy data. Our task is to obtain an almost smooth finite energy solution $(u, p, u_0, 0, T)$ with this data. We allow all implied constants to depend on u_0 .

We use a regularisation argument. Let N_n be a sequence of frequencies going to infinity, and set $u_0^{(n)} := P_{\leq N_n} u_0$; then $u_0^{(n)}$ converges to u_0 strongly in $L_x^2(\mathbb{R}^3)$, and $(u_0^{(n)}, 0, T)$ is smooth H^1 data for each n . Thus, by hypothesis, we may find a sequence of almost smooth finite energy solutions $(u^{(n)}, p^{(n)}, u_0^{(n)}, 0, T)$ with this data.

One could try invoking weak compactness right now to extract a solution, but as is well known, one only obtains a Leray–Hopf weak solution by doing so, which need not be smooth. So we will first work to establish some additional regularity on the sequence (after passing to a subsequence as necessary) before extracting a weakly convergent limit.

Since the $(u_0^{(n)}, 0, T)$ are uniformly bounded in energy, we see from [Lemma 8.1](#) that

$$\|u^{(n)}\|_{X^0([0, T] \times \mathbb{R}^3)} \lesssim 1. \quad (112)$$

Now let $0 < \tau_0 < T/2$ be a small time. From [\(112\)](#) and the pigeonhole principle, we may find a sequence of times $\tau^{(n)} \in [0, \tau_0]$ such that

$$\|u^{(n)}(\tau^{(n)})\|_{H_x^1(\mathbb{R}^3)} \lesssim \tau_0^{-1}.$$

Passing to a subsequence, we may assume that $\tau^{(n)}$ converges to a limit $\tau \in [0, \tau_0]$. If we then take $\tau' \in [\tau, 2\tau_0]$ sufficiently close to τ , we may apply [Lemma 5.5](#) and conclude that

$$\|u^{(n)}(\tau')\|_{H_x^{10}(\mathbb{R}^3)} \lesssim_{\tau, \tau', \tau_0} 1$$

(say) for all sufficiently large n . Passing to a further subsequence, we may then assume that $u^{(n)}(\tau')$ converges weakly in $H_x^{10}(\mathbb{R}^3)$ (and thus locally strongly in H_x^9) to a limit $u'_0 \in H_x^{10}(\mathbb{R}^3)$. By hypothesis, we may thus find an almost smooth H^1 solution $(u', p', u'_0, 0, T - \tau')$ with this data.

Meanwhile, by time translation symmetry [\(30\)](#), $(u^{(n)}(\cdot + \tau'), p^{(n)}(\cdot + \tau'), u^{(n)}(\tau'), 0, T - \tau')$ is also a sequence of almost smooth H^1 solutions. Since $u^{(n)}(\tau')$ converges locally strongly in $H_x^9(\mathbb{R}^3)$ to u'_0 , we would like to conclude that $u^{(n)}(t + \tau')$ also converges locally strongly to $u(t)$ in $H_x^1(\mathbb{R}^3)$, uniformly in $t \in [0, T - \tau']$. This does not quite follow from the standard local well-posedness theory in [Theorem 5.4](#), because this theory requires strong convergence in the *global* $H_x^1(\mathbb{R}^3)$ norm. However, we may take advantage of the local enstrophy estimates to spatially localise the local well-posedness theory, as follows.

Let $\varepsilon > 0$ be a small quantity (depending on the solution $u' = (u', p', u'_0, 0, T - \tau')$) to be chosen later, and let $R > 0$ be a sufficiently large radius (depending on ε and $(u', p', u'_0, 0, T - \tau')$) to be chosen later. Since u'_0 is in $H_x^{10}(\mathbb{R}^3)$, we see from monotone convergence that

$$\|u'_0\|_{H_x^{10}(\mathbb{R}^3 \setminus B(0, R))} \lesssim \varepsilon, \quad (113)$$

if R is sufficiently large depending on ε . Since the $u^{(n)}(\tau')$ converge locally strongly in $H_x^1(\mathbb{R}^3)$ to u'_0 , we conclude that

$$\|u^{(n)}(\tau')\|_{H_x^{10}(B(0,10R)\setminus B(0,R))} \lesssim \varepsilon,$$

if n is sufficiently large depending on R, ε . Applying [Theorem 10.1](#), we conclude (if R is large enough depending on u'_0 and $T - \tau'$) that

$$\|u^{(n)}(\cdot + \tau')\|_{X^1([0, T-\tau'] \times (B(0,9R)\setminus B(0,2R)))} \lesssim \varepsilon,$$

for n sufficiently large depending on R, ε . Using Duhamel's formula (and [Corollary 4.3](#)) repeatedly as in the proof of [Proposition 10.7](#), we may in fact conclude that

$$\|\partial_t^i u^{(n)}(\cdot + \tau')\|_{L_t^\infty H_x^6([0, T-\tau'] \times (B(0,8R)\setminus B(0,3R)))} \lesssim_{u', T} \varepsilon \quad (114)$$

(say) for $i = 0, 1$, taking R large enough depending on u', T, ε to ensure that the contributions to the Duhamel formula coming outside $B(0, 9R)$ or inside $B(0, 2R)$ are negligible, and taking n sufficiently large as always.

We let $\tilde{p}^{(n)}$ be the normalised pressure, defined by [\(9\)](#); by [Corollary 4.3](#), $\tilde{p}^{(n)}(t)$ and $p^{(n)}(t)$ differ by a constant $C(t)$ for almost every t . Using [\(9\)](#), [\(114\)](#) and [Lemma 8.1](#), we see that

$$\|\tilde{p}^{(n)}\|_{L_t^\infty H_x^2([0, T-\tau'] \times (B(0,7R)\setminus B(0,4R)))} \lesssim_{u', T} \varepsilon,$$

if R is large enough depending on u', T, ε .

Applying [Lemma 12.1](#), we may find divergence-free smooth vector fields $\tilde{u}^{(n)} : [\tau', T] \times \mathbb{R}^3 \rightarrow \mathbb{R}^3$ which agree with $u^{(n)}$ on $[\tau', T] \times B(0, 5R)$ but vanish outside of $[\tau', T] \times B(0, 6R)$, with

$$\|\partial_t^i \tilde{u}^{(n)}(\cdot + \tau')\|_{L_t^\infty H_x^5([0, T-\tau'] \times (B(0,8R)\setminus B(0,3R)))} \lesssim_{u', T} \varepsilon \quad (115)$$

(say) for n sufficiently large and $i = 0, 1$.

Let η be a smooth function that equals 1 on $B(0, 6R)$, is supported on $B(0, 7R)$, and obeys the usual derivative bounds in between. We then consider the smooth solutions

$$(\tilde{u}^{(n)}(\cdot + \tau'), \eta \tilde{p}^{(n)}(\cdot + \tau'), \tilde{u}^{(n)}(\tau'), \tilde{f}^{(n)}, T - \tau'), \quad (116)$$

where

$$\tilde{f}^{(n)} := (\partial_t \tilde{u}^{(n)} + \tilde{u}^{(n)} \cdot \nabla \tilde{u}^{(n)} - \Delta \tilde{u}^{(n)} + \nabla(\eta p^{(n)}))(\cdot + \tau').$$

By construction, \tilde{f}' and $\tilde{f}^{(n)}$ are smooth and supported on $[0, T - \tau'] \times (B(0, 7R)\setminus B(0, 5R))$, and the [\(116\)](#) are smooth, compactly supported solutions. From the preceding bounds on $\tilde{u}^{(n)}, \tilde{p}^{(n)}$, we see that

$$\|\tilde{f}^{(n)}\|_{L_t^\infty H_x^1([0, T-\tau'] \times \mathbb{R}^3)} \lesssim_{u', T} \varepsilon$$

for n sufficiently large.

Also, using [\(113\)](#), [\(115\)](#) we have

$$\|\tilde{u}^{(n)}(\tau') - u'_0\|_{H_x^1(\mathbb{R}^3)} \lesssim_{u', T} \varepsilon$$

for n sufficiently large. If ε is sufficiently small, we conclude from the local H^1 well-posedness theory (Theorem 5.4) that

$$\|\tilde{u}^{(n)}(\cdot + \tau') - u'\|_{X^1([0, T-\tau'] \times \mathbb{R}^3)} \lesssim_{u', T} \varepsilon,$$

and in particular

$$\|u^{(n)}(\cdot + \tau') - u'\|_{X^1([0, T-\tau'] \times B(0, R))} \lesssim_{u', T} \varepsilon$$

for n large enough. Sending ε to zero (and R to infinity), we conclude that $u^{(n)}(\cdot + \tau')$ converges weakly to u' . In particular, we see that any weak limit of the $u^{(n)}$ is smooth on $[\tau', T] \times \mathbb{R}^3$ (and furthermore, the weak limit is unique in this space-time region).

The above analysis was for a single choice of τ . Choosing τ to be a sequence of times going to zero (and repeatedly taking subsequences of the $u^{(n)}$ and diagonalising as necessary), we may thus arrive at a subsequence $u^{(n)}$ with the property that there is a unique weak limit u of the $u^{(n)}$, which is smooth on $(0, T] \times \mathbb{R}^3$. If we then set p by (9), we see on taking distributional limits that $(u, p, u_0, 0, T)$ is a Leray–Hopf weak solution to the initial data $(u_0, 0, T)$.

To finish the argument, we need to show that $(u, p, u_0, 0, T)$ is almost smooth at $(0, x_0)$ for every $x_0 \in \mathbb{R}^3$. Fix x_0 , and let $R > 0$ be a large radius. As u_0 is smooth, $\|u_0\|_{H^1(B(x_0, 5R))}$ is finite, and hence $\|u_0^{(n)}\|_{H^1(B(x_0, 5R))}$ is uniformly bounded. Applying Theorem 10.1 (recalling that the $u^{(n)}$ have uniformly bounded energy), we conclude (for R large enough) that there exists $0 < \tau < T$ such that $\|u^{(n)}\|_{X^1([0, \tau] \times B(x_0, 4R))}$ is uniformly bounded in n . Using Duhamel's formula as in Proposition 11.6, and noting that $u^{(n)}$ is uniformly smooth on $B(x_0, 4R)$, we conclude that $\|u^{(n)}\|_{L_t^\infty C^k([0, \tau] \times B(x_0, 3R))}$ is uniformly bounded for all $k \geq 0$. Taking weak limits, we conclude that

$$u \in L_t^\infty C^k((0, \tau] \times B(x_0, 3R))$$

for all $k \geq 0$. From this and (9) (and Lemma 8.1), we also see that

$$p \in L_t^\infty C^k((0, \tau] \times B(x_0, 2R))$$

for all $k \geq 0$. Using (3), we conclude that

$$\partial_t u \in L_t^\infty C^k((0, \tau] \times B(x_0, 2R))$$

for all $k \geq 0$. A similar argument also shows that

$$\partial_t u^{(n)} \in L_t^\infty C^k((0, \tau] \times B(x_0, 2R))$$

uniformly in n . From this, we see that the $\nabla_x^k u^{(n)}$ are uniformly Lipschitz in a neighbourhood of $(0, x_0)$. Since $\nabla_x^k u^{(n)}$ converges weakly to the smooth function $\nabla_x^k u$ in $(0, T] \times \mathbb{R}^3$, and also converges strongly at time zero in $H_x^1(\mathbb{R}^3)$ to the smooth function $\nabla_x^k u_0$, we conclude that $\nabla_x^k u$ can be extended in a locally Lipschitz continuous manner from $(0, T] \times \mathbb{R}^3$ to $[0, T] \times \mathbb{R}^3$ in such a way that it agrees with $\nabla_x^k u_0$ at time zero.

Now we consider derivatives $\nabla^k p$ of the pressure near $(0, x_0)$. Let $\varepsilon > 0$ be arbitrary. Then by the monotone convergence theorem, we see that if $R' > 0$ is a sufficiently large radius, then

$$\|u_0\|_{L_x^2(\mathbb{R}^3 \setminus B(x_0, R'))} \leq \varepsilon,$$

and thus

$$\|u_0^{(n)}\|_{L_x^2(\mathbb{R}^3 \setminus B(x_0, R'))} \lesssim \varepsilon$$

for n large enough.

By [Theorem 8.2](#), we conclude that if R' is large enough, there exists a time $0 < \tau < T$ such that

$$\|u^{(n)}\|_{L_t^\infty L_x^2([0, \tau] \times (\mathbb{R}^3 \setminus B(x_0, 2R')))} \lesssim \varepsilon,$$

and hence on taking weak limits,

$$\|u\|_{L_t^\infty L_x^2([0, \tau] \times (\mathbb{R}^3 \setminus B(x_0, 2R')))} \lesssim \varepsilon.$$

On the other hand, as $\nabla^k u$ is continuous at $t = 0$, $u(t)$ converges in $C^k(B(x_0, 2R'))$ to u_0 as $t \rightarrow 0$ for any $k \geq 0$. From this and [\(9\)](#) (and the decay of derivatives of the kernel of Δ^{-1} away from the origin), we see that

$$\limsup_{(t,x) \rightarrow (0,x_0); t>0} |\nabla^k p(t, x) - \nabla^k p_0(x_0)| \lesssim_k \varepsilon$$

for any $k \geq 0$, where p_0 is defined from u_0 using [\(9\)](#). Sending $\varepsilon \rightarrow 0$ and $R' \rightarrow \infty$, we conclude that $\nabla^k p$ extends continuously to $\nabla^k p_0(x_0)$ at $(0, x_0)$, and thus extends continuously to $\nabla^k p_0$ on all of the initial slice $\{0\} \times \mathbb{R}^3$. By [\(3\)](#) we conclude that $\partial_t \nabla^k u$ also extends continuously to the initial slice, with the Navier–Stokes equation [\(3\)](#) being obeyed both for times $t > 0$ and times $t = 0$. We have thus constructed an almost smooth finite energy solution $(u, p, u_0, 0, T)$ as desired. \square

Remark 13.2. We emphasise that [Theorem 13.1](#) only establishes *existence* of a smooth finite energy solution (assuming [Conjecture 1.13](#)), and not uniqueness; see [Remark 11.5](#). However, it is not difficult to see from the argument that one can at least ensure that the solution constructed is independent of the choice of time T , and can thus be extended to a single global smooth finite energy solution. (Alternatively, from [Lemma 8.1](#) we see that the enstrophy of the solution will become arbitrarily small for a sequence of times going to infinity, so for a sufficiently large time one can in fact construct a global smooth solution by standard perturbation theory techniques.)

Remark 13.3. One can modify the above argument to also establish [Conjecture 1.14](#) with a nonzero Schwartz forcing term f , provided of course that one also assumes [Conjecture 1.13](#) can be extended to the same class of f . We have not, however, investigated the weakest class of forcing terms f for which the argument works, though certainly finite energy seems insufficient.

14. Quantitative H^1 bounds

In this section we prove [Theorem 1.20\(vi\)](#). We begin with some easy implications. Firstly, it is trivial that [Conjecture 1.17](#) implies [Conjecture 1.16](#), and from the local well-posedness and regularity theory

in [Theorem 5.4](#) (or [Corollary 5.8](#)), we see that [Conjecture 1.16](#) implies [Conjecture 1.15](#), which in turn implies [Conjecture 1.13](#) (thanks to [Proposition 5.6](#)).

Next, we observe from [Theorem 5.1](#) and [Lemma 5.5](#) that given any H^1 data $(u_0, 0, T)$, there exists a time $0 < \tau < T$ such that one has an H^1 mild solution $(u, p, u_0, 0, \tau)$ with $u(\tau)$ smooth. If [Conjecture 1.13](#) holds, then one can then continue the solution in an almost smooth finite energy manner (and hence in an almost smooth H^1 manner, thanks to [Corollary 11.1](#)) from τ up to T . Normalising the pressure of this latter solution using [Lemma 4.1](#) and gluing the two solutions together, we obtain an H^1 mild solution up to time T . From this we see that [Conjecture 1.13](#) implies [Conjecture 1.15](#).

Now we show that [Conjecture 1.16](#) implies [Conjecture 1.17](#). Suppose that one has homogeneous H^1 data $(u_0, 0, T)$ with

$$\|u_0\|_{H_x^1(\mathbb{R}^3)} \leq A < \infty.$$

By [Conjecture 1.16](#) (which implies [Conjecture 1.15](#)), we may obtain a mild H^1 solution $(u, p, u_0, 0, T)$, which is smooth for positive times. Our objective is to show that

$$\|u\|_{L_t^\infty H_x^1([0, T] \times \mathbb{R}^3)} \lesssim_A 1.$$

Let $\varepsilon > 0$ be a quantity depending on A to be chosen later. We may assume that T is sufficiently large depending on ε, A ; otherwise the claim will follow immediately from [Conjecture 1.16](#). Using [Lemma 8.1](#) and the pigeonhole principle, we may then find a time $0 < T_1 < T$ with $T_1 \lesssim_A 1$ such that

$$\|\nabla u(T_1)\|_{L_x^2(\mathbb{R}^3)} \leq \varepsilon.$$

Meanwhile, from energy estimates, one has

$$\|u(T_1)\|_{L_x^2(\mathbb{R}^3)} \lesssim_A 1.$$

On $[T_1, T]$, we split $u = u_1 + v$, where u_1 is the linear solution $u_1(t) := e^{(t-T_1)\Delta} u(T_1)$ and $v := u - u_1$. From [\(21\)](#), one thus has

$$\|u_1\|_{X^0} \lesssim_A 1$$

and

$$\|\nabla u_1\|_{X^0} \lesssim \varepsilon.$$

From [\(11\)](#), [\(22\)](#) one has

$$\|v\|_{X^1([T_1, T] \times \mathbb{R}^3)} \lesssim \|\mathbb{O}(u_1 \nabla u_1 + u_1 \nabla v + v \nabla u_1 + v \nabla v)\|_{L_t^2 L_x^2([T_1, T] \times \mathbb{R}^3)}.$$

We now estimate various contributions to the right-hand side. We begin with the nonlinear term $\mathbb{O}(v \nabla v)$. By Hölder (and dropping the domain $[T_1, T] \times \mathbb{R}^3$ for brevity) followed by [Lemma 8.1](#), we have

$$\|\mathbb{O}(v \nabla v)\|_{L_t^2 L_x^2} \lesssim \|\nabla v\|_{L_t^2 L_x^6}^{1/2} \|\nabla v\|_{L_t^\infty L_x^2}^{1/2} \|v\|_{L_t^\infty L_x^6}^{1/2} \|v\|_{L_t^2 L_x^6}^{1/2} \lesssim \|v\|_{X^1}^{3/2} \|v\|_{X^0} \lesssim_A \|v\|_{X^1}^{3/2}.$$

A similar argument gives

$$\|\mathbb{O}(v \nabla u_1)\|_{L_t^2 L_x^2} \lesssim \|\nabla u_1\|_{X^0} \|v\|_{X^1}^{1/2} \|v\|_{X^0}^{1/2} \lesssim \varepsilon \|v\|_{X^1}$$

and

$$\|\mathbb{O}(u_1 \nabla u_1)\|_{L_t^2 L_x^2} \lesssim \|\nabla u_1\|_{X^0} \|\nabla u_1\|_{X^0}^{1/2} \|u_1\|_{X^0}^{1/2} \lesssim_A \varepsilon^{3/2}$$

and

$$\|\mathbb{O}(u_1 \nabla v)\|_{L_t^2 L_x^2} \lesssim \|\nabla v\|_{X^0} \|\nabla u_1\|_{X^0}^{1/2} \|u_1\|_{X^0}^{1/2} \lesssim_A \varepsilon^{1/2} \|v\|_{X^1}$$

and thus

$$\|v\|_{X^1} \lesssim_A \varepsilon^{3/2} + \varepsilon^{1/2} \|v\|_{X^1} + \|v\|_{X^1}^{3/2}.$$

If ε is small enough depending on A , a continuity argument in the T variable then gives

$$\|v\|_{X^1} \lesssim_A \varepsilon^{3/2}$$

and thus

$$\|u\|_{X^1([T_1, T])} \lesssim_A 1.$$

Using this and the triangle inequality, we conclude that [Conjecture 1.16](#) implies [Conjecture 1.17](#).

We now turn to the most difficult implication:

Proposition 14.1 (Concentration compactness). *If [Conjecture 1.15](#) is true, so is [Conjecture 1.16](#).*

We now prove this proposition. The methods are essentially those of [\[Gallagher 2001\]](#) (which are in turn based in [\[Bahouri and Gérard 1999; Gérard 1998\]](#)), which treated the (more difficult) critical analogue of this implication; indeed, one can view [Proposition 14.1](#) as a subcritical analogue of the critical result [\[Gallagher 2001, Corollary 1\]](#). For the convenience of the reader, though, we give a self-contained proof here, which does not need the full power of the machinery in the previously cited papers because we are now working in a subcritical regularity H^1 rather than a critical regularity such as $\dot{H}^{1/2}$, and as such one does not need to consider the role of the scaling symmetry [\(31\)](#).

We first make the remark that to prove [Conjecture 1.16](#), it suffices to do so with the condition

$$\|u_0\|_{H_x^1(\mathbb{R}^3)} \leq A \tag{117}$$

replaced by (say)

$$\|u_0\|_{H_x^{100}(\mathbb{R}^3)} \leq A. \tag{118}$$

To see this, observe that if we take data u_0 in $H_x^1(\mathbb{R}^3)$, then from [Theorem 5.4](#) and [Lemma 5.5](#) there exists a time $T_1 > 0$ depending only on A such that

$$\|u\|_{L_t^\infty H_x^1([0, \min(T, T_1)] \times \mathbb{R}^3)} \lesssim_A 1,$$

and such that

$$\|u(T_1)\|_{H_x^{100}(\mathbb{R}^3)} \lesssim_A 1$$

if $T > T_1$. From this and time translation symmetry [\(30\)](#), we see that we can deduce the $H_x^1(\mathbb{R}^3)$ version of [Conjecture 1.16](#) from the $H_x^{100}(\mathbb{R}^3)$ version.

Now suppose for contradiction that the $H_x^{100}(\mathbb{R}^3)$ version of [Conjecture 1.16](#) failed. Carefully negating the quantifiers, we can find a sequence $(u^{(n)}, p^{(n)}, u_0^{(n)}, 0, T^{(n)})$ of smooth homogeneous H^1 solutions, with $T^{(n)}$ uniformly bounded, and $u_0^{(n)}$ uniformly bounded in $H_x^{100}(\mathbb{R}^3)$, such that

$$\lim_{n \rightarrow \infty} \|u^{(n)}\|_{L_t^\infty H_x^1([0, T^{(n)}] \times \mathbb{R}^3)} = \infty. \quad (119)$$

By [Lemma 4.1](#) we may assume that these solutions have normalised pressure.

If we were working on a compact domain, such as $\mathbb{R}^3/\mathbb{Z}^3$, we could now extract a subsequence of the $u_0^{(n)}$ that converged strongly in a lower regularity space, such as $H_x^{99}(\mathbb{R}^3/\mathbb{Z}^3)$. But our domain \mathbb{R}^3 is noncompact, and in particular has the action of a noncompact symmetry group, namely the translation group $\tau_{x_0}u(x) := u(x - x_0)$. However, as is well known, we have a substitute for compactness in this setting, namely *concentration compactness*. Specifically:

Proposition 14.2 (Profile decomposition). *Let $u_0^{(n)} \in H_x^{100}(\mathbb{R}^3)$ be a sequence with*

$$\limsup_{n \rightarrow \infty} \|u_0^{(n)}\|_{H_x^{100}(\mathbb{R}^3)} \leq A,$$

and let $\varepsilon > 0$. Then, after passing to a subsequence, there exists a decomposition

$$u_0^{(n)} = \sum_{j=1}^J \tau_{x_j^{(n)}} w_{j,0} + r_0^{(n)},$$

where $|J| \lesssim_{A,\varepsilon} 1$, $w_{1,0}, \dots, w_{J,0} \in H_x^{100}(\mathbb{R}^3)$, $x_j^{(n)} \in \mathbb{R}^3$, and the remainder $r_0^{(n)}$ obeys the estimates

$$\limsup_{n \rightarrow \infty} \|r_0^{(n)}\|_{H_x^{100}(\mathbb{R}^3)} \leq A$$

and

$$\limsup_{n \rightarrow \infty} \|r_0^{(n)}\|_{L_x^\infty(\mathbb{R}^3)} \leq \varepsilon. \quad (120)$$

Furthermore, for any $1 \leq j < j' \leq J$, one has

$$|x_j^{(n)} - x_{j'}^{(n)}| \rightarrow \infty, \quad (121)$$

and for any $1 \leq j \leq J$, the sequence $\tau_{-x_j^{(n)}} r_0^{(n)}$ converges weakly in $H_x^{100}(\mathbb{R}^3)$ to zero.

Finally, if the $u_0^{(n)}$ are divergence-free, then the $w_{j,0}$ and $r_0^{(n)}$ are also divergence-free.

Proof. See, for example, [\[Gérard 1998\]](#). We sketch the (standard) proof as follows. If

$$\|u_0^{(n)}\|_{L_x^\infty(\mathbb{R}^3)} \leq \varepsilon$$

for all sufficiently large n , then there is nothing to prove (just take $J = 0$ and $r_0^{(n)} := u_0^{(n)}$). Otherwise, after passing to a subsequence, we can find a sequence $x_1^{(n)} \in \mathbb{R}^3$ such that $|u_0^{(n)}(x_1^{(n)})| \geq \varepsilon/2$ (say). The sequence $\tau_{-x_1^{(n)}} u_0^{(n)}$ is then bounded in $H_x^{100}(\mathbb{R}^3)$ and bounded away from zero at the origin; by passing to a further subsequence, we may assume that it converges weakly in $H_x^{100}(\mathbb{R}^3)$ to a limit w_1 , which

then has an $H_x^{100}(\mathbb{R}^3)$ norm of $\lesssim_{A,\varepsilon} 1$ and is asymptotically orthogonal in the Hilbert space $H_x^{100}(\mathbb{R}^3)$ to $\tau_{-x_1^{(n)}} u_0^{(n)}$. We then have the decomposition

$$u_0^{(n)} = \tau_{x_1^{(n)}} w_{1,0} + u_0^{(n),1},$$

and from an application of the cosine rule in the Hilbert space $H_x^{100}(\mathbb{R}^3)$, one can verify that

$$\limsup_{n \rightarrow \infty} \|u_0^{(n),1}\|_{H_x^{100}(\mathbb{R}^3)}^2 \leq A^2 - c$$

for some $c > 0$ depending only on ε, A . We can then iterate this procedure $O_{J,\varepsilon}(1)$ times to obtain the desired decomposition. \square

We apply this proposition with a value of $\varepsilon > 0$ depending on A, T to be chosen later. The $w_{j,0}$ lie in $H_x^{100}(\mathbb{R}^3)$, and thus by the assumption that [Conjecture 1.15](#) is true, we can find mild H^1 solutions $(w_j, p_j, w_{j,0}, 0, T)$ with this data. By [Theorem 5.1](#), we have

$$\|w_j\|_{X^{100}} < \infty$$

for each $1 \leq j \leq J$, and to abbreviate the notation, we adopt the convention that the space-time domain is understood to be $[0, T] \times \mathbb{R}^3$.

Next, we consider the remainder term $r_0^{(n)}$. From [\(21\)](#) one has

$$\|e^{t\Delta} r_0^{(n)}\|_{X^{100}} \lesssim A,$$

while from [\(120\)](#) one has

$$\|e^{t\Delta} r_0^{(n)}\|_{L_t^\infty L_x^\infty} \lesssim \varepsilon$$

for n sufficiently large. Interpolating between the two, we soon conclude that

$$\|e^{t\Delta} r_0^{(n)}\|_{X^1} \lesssim_{A,T} \varepsilon^c$$

for some absolute constant $c > 0$. If we take ε sufficiently small depending on A, T , we can use stability of the zero solution (see [Theorem 5.1](#); one could also have used here the results from [\[Chemin and Gallagher 2009\]](#)) to conclude the existence of a mild H^1 solution $(r^{(n)}, p_*^{(n)}, r_0^{(n)}, 0, T)$ with this data, with the estimates

$$\|r^{(n)}\|_{X^1} \lesssim_{A,T} \varepsilon^c; \tag{122}$$

from [Theorem 5.1](#), we then also have

$$\|r^{(n)}\|_{X^{100}} \lesssim_{A,T} 1.$$

We now form the solution

$$(\tilde{u}^{(n)}, \tilde{p}^{(n)}, u_0^{(n)}, \tilde{f}^{(n)}, T),$$

where the velocity field $\tilde{u}^{(n)}$ is given by

$$\tilde{u}^{(n)} := \sum_{j=1}^J \tau_{x_j^{(n)}} w_j + r^{(n)},$$

the pressure field $\tilde{p}^{(n)}$ is given by (9), and the forcing term $\tilde{f}^{(n)}$ is given by the formula

$$\tilde{f}^{(n)} := \partial_t \tilde{u}^{(n)} - \Delta \tilde{u}^{(n)} - PB(\tilde{u}^{(n)}, \tilde{u}^{(n)}).$$

This is clearly a mild H^1 solution, with

$$\|\tilde{u}^{(n)}\|_{X^{100}} \lesssim_{A,T,\varepsilon} 1.$$

We now estimate $\tilde{f}^{(n)}$. From (68) for the solutions $\tau_{x_j^{(n)}} w_j + r^{(n)}$, we have an expansion of $\tilde{f}^{(n)}$ purely involving nonlinear interaction terms:

$$\tilde{f}^{(n)} = \sum_{1 \leq j < j' \leq J} P\mathbb{C}(\nabla(\tau_{x_j^{(n)}} w_j, \tau_{x_{j'}^{(n)}} w_{j'})) + \sum_{1 \leq j < J} P\mathbb{C}(\nabla(\tau_{x_j^{(n)}} w_j, r^{(n)})).$$

In particular, from the triangle inequality and translation invariance we have

$$\|\tilde{f}^{(n)}\|_{L_t^2 L_x^2} \lesssim \sum_{1 \leq j < j' \leq J} \|\mathbb{C}(\nabla(w_j, \tau_{x_{j'}^{(n)} - x_j^{(n)}} w_{j'}))\|_{L_t^2 L_x^2} + \sum_{1 \leq j < J} \|\mathbb{C}(\nabla(w_j, \tau_{-x_j^{(n)}} r^{(n)}))\|_{L_t^2 L_x^2}.$$

But by (121) and Sobolev embedding,

$$\tau_{x_{j'}^{(n)} - x_j^{(n)}} w_{j'} \quad \text{and} \quad \tau_{-x_j^{(n)}} r^{(n)}$$

are bounded in $L_t^\infty L_x^\infty$ and converge locally uniformly to zero, and so we conclude that

$$\lim_{n \rightarrow \infty} \|\tilde{f}^{(n)}\|_{L_t^2 L_x^2} = 0.$$

From this and the stability theory in Theorem 5.4, we conclude that for n large enough, there is an H^1 mild solution $(u^{(n)}, p^{(n)}, u_0^{(n)}, 0, T)$ with

$$\lim_{n \rightarrow \infty} \|\tilde{u}^{(n)} - u^{(n)}\|_{X^1} = 0,$$

and in particular

$$\limsup_{n \rightarrow \infty} \|u^{(n)}\|_{L_t^\infty H_x^1([0, T] \times \mathbb{R}^3)} < \infty.$$

By the uniqueness theory in Theorem 5.4, this solution must agree with the original solutions

$$(u^{(n)}, p^{(n)}, u_0^{(n)}, 0, T^{(n)})$$

on $[0, T^{(n)}] \times \mathbb{R}^3$; but then we contradict (119). Proposition 14.1 follows.

15. Nonexistence of smooth solutions

In this section we establish Theorem 1.12. Informally, the reason for the irregularity is as follows. Assuming normalised pressure, one concludes from (9) that

$$p = \mathbb{C}(\Delta^{-1} \nabla^2 (uu)).$$

If one then differentiates this twice in time, using (3) to convert time derivatives of u into Δu plus lower-order terms and using integration by parts to redistribute derivatives, we eventually obtain (formally, at least) a formula of the form

$$\partial_t^2 p = \mathcal{O}(\Delta^{-1} \nabla^2 (\Delta u \Delta u)) + \text{lower-order terms.}$$

But if u_0 is merely assumed to be smooth and in H^1 , then Δu can grow arbitrarily fast at infinity at time $t = 0$, and this should cause p to fail to be C_t^2 at time zero.

We turn to the details. To eliminate the normalised pressure assumption, we will work with ∇p instead of p , and thus we will seek to establish bad behaviour for $\nabla \partial_t^2 p$ at time $t = 0$. For technical reasons it is convenient to work in the weak topology in space. The key quantitative step is the following:

Proposition 15.1 (Quantitative failure of regularity). *Let $u_0 : \mathbb{R}^3 \rightarrow \mathbb{R}^3$ be smooth, divergence-free, and compactly supported, and let $\psi : \mathbb{R}^3 \rightarrow \mathbb{R}$ be smooth, compactly supported, and have total mass $\int_{\mathbb{R}^3} \psi = 1$. Let $R, M, \varepsilon > 0$. Then there exists a smooth divergence-free compactly supported function u_1 which vanishes on $B(0, R)$ with*

$$\|u_1\|_{H_x^1(\mathbb{R}^3)} \lesssim \varepsilon$$

and such that if $(u, p, u_0 + u_1, 0, T)$ is a mild H^1 (and hence smooth, by Proposition 5.6) solution with data $(u_0 + u_1, 0, T)$, then

$$\left| \int_{\mathbb{R}^3} \nabla \partial_t^2 p(0, x) \psi(x) dx \right| > M. \quad (123)$$

Let us assume this proposition for now and conclude Theorem 1.12. We will use an argument reminiscent of that used to establish the Baire category theorem or the uniform boundedness principle. Let $\psi : \mathbb{R}^3 \rightarrow \mathbb{R}$ be a fixed smooth, compactly supported function with total mass 1. We will need a rapidly decreasing sequence

$$\varepsilon^{(1)} > \varepsilon^{(2)} > \dots > 0$$

of small quantities to be chosen later, with each $\varepsilon^{(n)}$ sufficiently small depending on the previous $\varepsilon^{(1)}, \dots, \varepsilon^{(n-1)}$. Applying Proposition 15.1 recursively starting with $u_0 = 0$, one can then find a sequence of smooth, divergence-free, and compactly supported functions $u_1^{(n)}$ for $n = 1, 2, \dots$ such that

$$\|u_1^{(n)}\|_{H_x^1(\mathbb{R}^3)} \lesssim \varepsilon^{(n)},$$

with $u_1^{(n)}$ vanishing on $B(0, 1/\varepsilon^{(n)})$, such that if $(u^{(n)}, p^{(n)}, u_0^{(n)}, 0, T^{(n)})$ is a mild H^1 (and hence smooth) solution with data

$$u_0^{(n)} := u_1^{(1)} + \dots + u_1^{(n)},$$

then

$$\left| \int_{\mathbb{R}^3} \nabla \partial_t^2 p^{(n)}(0, x) \psi(x) dx \right| > 1/\varepsilon^{(n)}. \quad (124)$$

Furthermore, each $u_1^{(n)}$ depends only on $\varepsilon^{(1)}, \dots, \varepsilon^{(n)}$, and in particular is independent of $\varepsilon^{(n+1)}$.

By the triangle inequality (and assuming the $\varepsilon^{(n)}$ decay fast enough), the data $u_0^{(n)}$ is strongly convergent in $H_x^1(\mathbb{R}^3)$ to a limit $u_0 = \sum_{n=1}^{\infty} u_1^{(n)} \in H_x^1(\mathbb{R}^3)$, with

$$\|u_0 - u_0^{(n)}\|_{H_x^1(\mathbb{R}^3)} \lesssim \varepsilon^{(n+1)}.$$

If we make each $\varepsilon^{(n+1)}$ sufficiently small depending on $u_0^{(n)}$, and hence on $\varepsilon^{(1)}, \dots, \varepsilon^{(n)}$, then the $u_1^{(n)}$ will have disjoint supports; as each $u_1^{(n)}$ is smooth and divergence-free, this implies that

$$u_0 = \sum_{n=1}^{\infty} u_1^{(n)}$$

is also smooth and divergence-free.

Applying [Theorem 5.1](#), we may then take the times $T^{(n)} = 1$ (if the $\varepsilon^{(n)}$ are small enough), and $(u^{(n)}, p^{(n)}, u_0^{(n)}, 0, 1)$ will converge to a mild H^1 solution $(u, p, u_0, 0, 1)$ in the sense that $u^{(n)}$ converges strongly in $X^1([0, 1] \times \mathbb{R}^3)$ to u . Indeed, from the Lipschitz stability property, we see (if the $\varepsilon^{(n)}$ decay fast enough) that

$$\|u - u^{(n)}\|_{X^1([0, 1] \times \mathbb{R}^3)} \lesssim \varepsilon^{(n+1)}.$$

Also, $u, u^{(n)}$ are bounded in $X^1([0, 1] \times \mathbb{R}^3)$ by $O(1)$. Using [\(9\)](#) and Sobolev embedding, this implies

$$\|p - p^{(n)}\|_{L_t^\infty L_x^3([0, 1] \times \mathbb{R}^3)} \lesssim \varepsilon^{(n+1)},$$

and so if one sets

$$F^{(n)}(t) := \int_{\mathbb{R}^3} \nabla p^{(n)}(t, x) \psi(x) dx$$

and

$$F(t) := \int_{\mathbb{R}^3} \nabla p(t, x) \psi(x) dx,$$

then from integration by parts, we have

$$\|F - F^{(n)}\|_{L_t^\infty([0, 1])} \lesssim \varepsilon^{(n+1)}. \quad (125)$$

Meanwhile, each $F^{(n)}$ is smooth, and F continuous, from [Proposition 5.6](#), and from [\(124\)](#) one has

$$|\partial_t^2 F^{(n)}(0)| \geq 1/\varepsilon^{(n)}.$$

In particular, if $\varepsilon^{(n+1)}$ is sufficiently small depending on $F^{(n)}$ (which in turn depends on $\varepsilon^{(1)}, \dots, \varepsilon^{(n)}$), one has from Taylor's theorem with remainder that

$$\frac{|F^{(n)}(2(\varepsilon^{(n+1)})^{0.1}) - 2F^{(n)}((\varepsilon^{(n+1)})^{0.1}) + F^{(n)}(0)|}{(\varepsilon^{(n+1)})^{0.2}} \gtrsim \frac{1}{\varepsilon^{(n)}}.$$

Applying [\(125\)](#), we conclude that

$$\frac{|F(2(\varepsilon^{(n+1)})^{0.1}) - 2F((\varepsilon^{(n+1)})^{0.1}) + F(0)|}{(\varepsilon^{(n+1)})^{0.2}} \gtrsim \frac{1}{\varepsilon^{(n)}},$$

if $\varepsilon^{(n+1)}$ is sufficiently small depending on $\varepsilon^{(1)}, \dots, \varepsilon^{(n)}$. In particular,

$$\limsup_{h \rightarrow 0^+} \frac{|F(2h) - 2F(h) + F(0)|}{h^2} = +\infty,$$

which by Taylor's theorem with remainder implies that F is not smooth at 0.

We claim that the data u_0 gives the desired counterexample to [Theorem 1.12](#). Indeed, suppose for contradiction that there was a smooth solution $(\tilde{u}, \tilde{p}, u_0, 0, T)$ for some $T > 0$. By shrinking T , we may assume $T \leq 1$. By [Lemma 4.1](#), we see that $\tilde{p}(t)$ has normalised pressure up to a constant for almost every t , and thus after adjusting $\tilde{p}(t)$ by that constant, $(\tilde{u}, \tilde{p}, u_0, 0, T)$ is a mild H^1 solution. Using the uniqueness property in [Theorem 5.1](#), we conclude that $u = \tilde{u}$, and $p(t)$ and $\tilde{p}(t)$ differ by a constant for almost every t , and hence (by continuity of both p and \tilde{p}) for every t . In particular, $\nabla p = \nabla \tilde{p}$, and so

$$F(t) = \int_{\mathbb{R}^3} \nabla \tilde{p}(t, x) \psi(x) dx.$$

But as \tilde{p} is smooth on $[0, T] \times \mathbb{R}^3$, F is smooth at 0, a contradiction.

Remark 15.2. The above argument showed that ∇p failed to be smooth at $t = 0$; by using [\(3\)](#), we conclude that the velocity field u must then also be nonsmooth at $t = 0$ (though the velocity u has one more degree of time regularity than the pressure p). Thus the failure of regularity is not just an artefact of pressure normalisation. Using the vorticity equation [\(84\)](#), one can then show a similar failure of time regularity for the vorticity, although again one gains an additional degree of time differentiability over the velocity u .

The irregularities in time stem from the unbounded growth of high derivatives of the initial data. If one assumes that *all* spatial derivatives of u_0 are in $L_x^2(\mathbb{R}^3)$, that is, that $u_0 \in H^\infty(\mathbb{R}^3)$, then one can prove iteratively²³ that all time derivatives of u and p at time zero are bounded, and also have first spatial derivatives in $H^\infty(\mathbb{R}^3)$ (basically because the first derivative of the kernel of the Leray projection is integrable at infinity). In particular, u and p now remain smooth at time 0.

It remains to establish [Proposition 15.1](#). Fix $u_0, \psi, R, M, \varepsilon$, and let u_1 be a smooth divergence-free compactly supported function u_1 vanishing on $B(0, R)$ with $H_x^1(\mathbb{R}^3)$ norm $O(\varepsilon)$ to be chosen later. Let $(u, p, u_0 + u_1, 0, T)$ be a mild H^1 solution with this given data. By [Theorem 5.1](#), this is a smooth solution, with all derivatives of u, p lying in $L_t^\infty L_x^2$. From [Lemma 4.1](#) we thus have

$$\nabla p = -\nabla \Delta^{-1} \partial_i \partial_j (u_i u_j) \tag{126}$$

for almost all times t . But both sides are smooth in $[0, T] \times \mathbb{R}^3$, so this formula is valid for all times t (and in particular at $t = 0$). In particular, we may apply a Leray projection P to [\(3\)](#) and conclude that

$$\partial_t u = \Delta u + PB(u, u). \tag{127}$$

We differentiate [\(126\)](#) once in time to obtain

$$\nabla \partial_t p = -2\nabla \Delta^{-1} \partial_i \partial_j (u_i \partial_t u_j).$$

²³We thank Richard Melrose for this observation.

Expanding out $\partial_t u_j$ using (3), we obtain

$$\nabla \partial_t p = -2\nabla \Delta^{-1} \partial_i \partial_j (u_i \Delta u_j) + \mathcal{O}(\Delta^{-1} \nabla^3 (u PB(u, u))).$$

Writing

$$\partial_i \partial_j (u_i \Delta u_j) = -2\partial_i \partial_j ((\partial_k u_i)(\partial_k u_j)) + \mathcal{O}(\nabla^4 (uu)),$$

we thus have

$$\nabla \partial_t p = 2\nabla \Delta^{-1} \partial_i \partial_j (\partial_k u_i \partial_k u_j) + \mathcal{O}(\Delta^{-1} \nabla^5 (uu)) + \mathcal{O}(\Delta^{-1} \nabla^3 (u PB(u, u))).$$

We differentiate this in time again and use (127) to obtain

$$\begin{aligned} \nabla \partial_t^2 p &= 4\nabla \Delta^{-1} \partial_i \partial_j (\partial_k u_i \partial_k \Delta u_j) \\ &\quad + \mathcal{O}(\Delta^{-1} \nabla^3 ((\nabla u) \nabla PB(u, u))) \\ &\quad + \mathcal{O}(\Delta^{-1} \nabla^5 (u \partial_t u)) \\ &\quad + \mathcal{O}(\Delta^{-1} \nabla^3 ((\partial_t u) PB(u, u))) \\ &\quad + \mathcal{O}(\Delta^{-1} \nabla^3 (u PB(u, \partial_t u))). \end{aligned}$$

We can write $\partial_k u_i \partial_k \Delta u_j = -(\Delta u_i)(\Delta u_j) + \mathcal{O}(\nabla(\nabla u \Delta u))$, so that

$$\begin{aligned} \nabla \partial_t^2 p &= -4\nabla \Delta^{-1} \partial_i \partial_j (\Delta u_i \Delta u_j) \\ &\quad + \mathcal{O}(\Delta^{-1} \nabla^4 (\nabla u \Delta u)) \\ &\quad + \mathcal{O}(\Delta^{-1} \nabla^3 ((\nabla u) \nabla PB(u, u))) \\ &\quad + \mathcal{O}(\Delta^{-1} \nabla^5 (u \partial_t u)) \\ &\quad + \mathcal{O}(\Delta^{-1} \nabla^3 ((\partial_t u) PB(u, u))) \\ &\quad + \mathcal{O}(\Delta^{-1} \nabla^3 (u PB(u, \partial_t u))). \end{aligned}$$

Integrating this against ψ , we may thus expand

$$\int_{\mathbb{R}^3} \nabla \partial_t^2 p(0, x) \psi(x) dx = 4X_0 + \sum_{i=1}^5 \mathcal{O}(X_i),$$

where

$$\begin{aligned} X_0 &:= \int_{\mathbb{R}^3} (\partial_i \partial_j \nabla \Delta^{-1} \psi) \Delta u_i \Delta u_j, & X_1 &:= \int_{\mathbb{R}^3} (\nabla^4 \Delta^{-1} \psi) \nabla u \Delta u, \\ X_2 &:= \int_{\mathbb{R}^3} (\nabla^3 \Delta^{-1} \psi) \nabla u \nabla PB(u, u), & X_3 &:= \int_{\mathbb{R}^3} (\nabla^5 \Delta^{-1} \psi) u \partial_t u, \\ X_4 &:= \int_{\mathbb{R}^3} (\nabla^3 \Delta^{-1} \psi) (\partial_t u) PB(u, u), & X_5 &:= \int_{\mathbb{R}^3} (\nabla^3 \Delta^{-1} \psi) u PB(u, \partial_t u), \end{aligned}$$

with all expressions being evaluated at time 0.

From (127) and Sobolev embedding, one has

$$\|\partial_t u(0)\|_{L_x^2(\mathbb{R}^3)} \lesssim_{u_0} 1 + \|u_1\|_{H_x^2(\mathbb{R}^3)}.$$

Meanwhile, if ε is small enough, we see that

$$\|u(0)\|_{H_x^1(\mathbb{R}^3)} \lesssim_{u_0} 1,$$

and thus from the Gagliardo–Nirenberg inequality,

$$\|u(0)\|_{L_x^\infty(\mathbb{R}^3)} \lesssim_{u_0} (1 + \|u_1\|_{H_x^2(\mathbb{R}^3)})^{1/2}.$$

From many applications of the Sobolev and Hölder inequalities (and, in the case of X_5 , an integration by parts to move the derivative off of $\partial_t u$), we conclude that

$$|X_i| \lesssim_{u_0, \psi} (1 + \|u_1\|_{H_x^2(\mathbb{R}^3)})^{3/2},$$

for $i = 1, 2, 3, 4, 5$. In a similar spirit, one has

$$X_0 = \int_{\mathbb{R}^3} (\partial_i \partial_j \nabla \Delta^{-1} \psi) \Delta u_{1,i} \Delta u_{1,j} + O_{u_0, \psi} (1 + \|u_1\|_{H_x^2(\mathbb{R}^3)}).$$

To demonstrate (123), it thus suffices to exhibit a sequence $u_1^{(n)} : \mathbb{R}^3 \rightarrow \mathbb{R}^3$ of smooth divergence-free compactly supported vector fields supported outside of $B(0, R)$ such that

$$\left| \int_{\mathbb{R}^3} (\partial_i \partial_j \nabla \Delta^{-1} \psi) \Delta u_{1,i}^{(n)} \Delta u_{1,j}^{(n)} \right| \gtrsim_{R, \psi} \|u_1^{(n)}\|_{H_x^2(\mathbb{R}^3)}^2,$$

with

$$\|u_1^{(n)}\|_{H_x^1(\mathbb{R}^3)} \rightarrow 0 \quad \text{and} \quad \|u_1^{(n)}\|_{H_x^2(\mathbb{R}^3)} \rightarrow \infty.$$

We construct $u_1^{(n)}$ explicitly as the “wave packet”

$$u_1^{(n)}(x) := n^{-5/2} \nabla \times \Psi^{(n)}(x_0),$$

where e_1, e_2, e_3 is the standard basis, $x_0 \in \mathbb{R}^3$ is a point (independent of n) outside of $B(0, R+1)$ to be chosen later, and

$$\Psi^{(n)}(x) = \chi(x) \sin(n\xi \cdot x) \eta,$$

where $\xi \in \mathbb{R}^3$ is a nonzero frequency (independent of n) to be chosen later, $\eta \in \mathbb{R}^3$ is a nonzero direction, and $\chi : \mathbb{R}^3 \rightarrow \mathbb{R}$ is a smooth bump function supported on $B(0, 1)$ to be chosen later. Note from construction that $u_1^{(n)}$ is smooth, divergence-free, and supported on $B(x_0, 1)$, and thus vanishing on $B(0, R)$ for $R_0 > R+1$. One can compute that

$$\|u_1^{(n)}\|_{H_x^1(\mathbb{R}^3)} \ll_\chi n^{-1/2} \quad \text{and} \quad \|u_1^{(n)}\|_{H_x^2(\mathbb{R}^3)} \gg_\chi n^{1/2},$$

as long as χ is not identically zero. To conclude the theorem, it thus suffices to show that

$$\left| \int_{\mathbb{R}^3} (\partial_i \partial_j \nabla \Delta^{-1} \psi) \Delta u_{1,i}^{(n)} \Delta u_{1,j}^{(n)} \right| \gg_{R_0, \psi, \chi} n$$

if R_0 and n are large enough.

Observe that

$$u_1^{(n)}(x) := n^{-3/2} \sin(n\xi \cdot (x - x_0)) \chi(x - x_0) (\xi \times \eta) + O(n^{-5/2})$$

and similarly

$$\Delta u_1^{(n)}(x) := -n^{1/2}|\xi|^2 \sin(n\xi \cdot (x - x_0))\chi(x - x_0)(\xi \times \eta) + O(n^{-1/2}),$$

and so by choosing χ appropriately and using the Riemann–Lebesgue lemma, it suffices to find $x_0, \xi, \eta \in \mathbb{R}^3$ such that

$$(\partial_i \partial_j \nabla \Delta^{-1} \psi)(\xi \times \eta)_i (\xi \times \eta)_j(x_0) \neq 0.$$

But as ψ has mean one, we see that $\nabla^3 \Delta^{-1} \psi(x_0)$ is not identically zero for x_0 large enough, and the claim follows.

Acknowledgements

The author is also indebted to Andrea Bertozzi for valuable discussions, and in particular for raising the question of whether the answer to the global regularity problem of the Navier–Stokes equation is sensitive to the decay hypotheses on the initial velocity field. We thank Xiaoyi Zhang and the anonymous referee for corrections and suggestions.

References

- [Bahouri and Gérard 1999] H. Bahouri and P. Gérard, “High frequency approximation of solutions to critical nonlinear wave equations”, *Amer. J. Math.* **121**:1 (1999), 131–175. [MR 2000i:35123](#) [Zbl 0919.35089](#)
- [Beale et al. 1984] J. T. Beale, T. Kato, and A. Majda, “Remarks on the breakdown of smooth solutions for the 3-D Euler equations”, *Comm. Math. Phys.* **94**:1 (1984), 61–66. [MR 85j:35154](#) [Zbl 0573.76029](#)
- [Bogovskii 1980] M. E. Bogovskii, “Solutions of some problems of vector analysis, associated with the operators div and grad”, pp. 5–40, 149 in *Theory of cubature formulas and the application of functional analysis to problems of mathematical physics*, Trudy Sem. S. L. Soboleva **1**, Akad. Nauk SSSR Sibirsk. Otdel. Inst. Mat., Novosibirsk, 1980. In Russian. [MR 82m:26014](#)
- [Caffarelli et al. 1982] L. Caffarelli, R. Kohn, and L. Nirenberg, “Partial regularity of suitable weak solutions of the Navier–Stokes equations”, *Comm. Pure Appl. Math.* **35**:6 (1982), 771–831. [MR 84m:35097](#) [Zbl 0509.35067](#)
- [Chae 1992] D. Chae, “Some a priori estimates for weak solutions of the 3-D Navier–Stokes equations”, *J. Math. Anal. Appl.* **167**:1 (1992), 236–244. [MR 93d:35112](#) [Zbl 0755.35085](#)
- [Chemin and Gallagher 2009] J.-Y. Chemin and I. Gallagher, “Wellposedness and stability results for the Navier–Stokes equations in \mathbf{R}^3 ”, *Ann. Inst. H. Poincaré Anal. Non Linéaire* **26**:2 (2009), 599–624. [MR 2010i:35278](#) [Zbl 1165.35038](#)
- [Cheskidov and Shvydkoy 2011] A. Cheskidov and R. Shvydkoy, “Regularity problem for the 3D Navier–Stokes equations: the use of Kolmogorov’s dissipation range”, preprint, 2011. [arXiv 1102.1944v1](#)
- [Choi and Vasseur 2011] K. Choi and A. Vasseur, “Estimates on fractional higher derivatives of weak solutions for the Navier–Stokes equations”, preprint, Oxford Centre for Nonlinear PDE, 2011, Available at <http://goo.gl/wDKRz>. [arXiv 1105.1526v1](#)
- [Constantin 1990] P. Constantin, “Navier–Stokes equations and area of interfaces”, *Comm. Math. Phys.* **129**:2 (1990), 241–266. [MR 91i:35148](#) [Zbl 0725.35080](#)
- [Doering and Foias 2002] C. R. Doering and C. Foias, “Energy dissipation in body-forced turbulence”, *J. Fluid Mech.* **467** (2002), 289–306. [MR 2003i:76048](#) [Zbl 1029.76025](#)
- [Doering and Gibbon 2002] C. R. Doering and J. D. Gibbon, “Bounds on moments of the energy spectrum for weak solutions of the three-dimensional Navier–Stokes equations”, *Phys. D* **165**:3–4 (2002), 163–175. [MR 2003j:76022](#) [Zbl 0997.35046](#)
- [Duff 1990] G. F. D. Duff, “Derivative estimates for the Navier–Stokes equations in a three-dimensional region”, *Acta Math.* **164**:3–4 (1990), 145–210. [MR 91j:35218](#) [Zbl 0728.35083](#)
- [Fabes et al. 1972] E. B. Fabes, B. F. Jones, and N. M. Rivière, “The initial value problem for the Navier–Stokes equations with data in L^p ”, *Arch. Rational Mech. Anal.* **45** (1972), 222–240. [MR 47 #5463](#) [Zbl 0254.35097](#)

- [Fefferman 2006] C. L. Fefferman, “Existence and smoothness of the Navier–Stokes equation”, pp. 57–67 in *The millennium prize problems*, edited by J. Carlson et al., Clay Math. Inst., 2006. [MR 2238274](#) [Zbl 1194.35002](#)
- [Foaïaş et al. 1981] C. Foaïaş, C. Guillopé, and R. Temam, “New a priori estimates for Navier–Stokes equations in dimension 3”, *Comm. Partial Differential Equations* **6**:3 (1981), 329–359. [MR 82f:35153](#) [Zbl 0472.35070](#)
- [Gallagher 2001] I. Gallagher, “Profile decomposition for solutions of the Navier–Stokes equations”, *Bull. Soc. Math. France* **129**:2 (2001), 285–316. [MR 2002h:35235](#) [Zbl 0987.35120](#)
- [Gérard 1998] P. Gérard, “Description du défaut de compacité de l’injection de Sobolev”, *ESAIM Control Optim. Calc. Var.* **3** (1998), 213–233. [MR 99h:46051](#)
- [Germain 2006] P. Germain, “Multipliers, paramultipliers, and weak-strong uniqueness for the Navier–Stokes equations”, *J. Differential Equations* **226**:2 (2006), 373–428. [MR 2007c:35128](#) [Zbl 1104.35033](#)
- [Germain 2008] P. Germain, “Strong solutions and weak-strong uniqueness for the nonhomogeneous Navier–Stokes system”, *J. Anal. Math.* **105** (2008), 169–196. [MR 2009j:35268](#) [Zbl 1198.35179](#)
- [Gibbon 2012] J. D. Gibbon, “A hierarchy of length scales for weak solutions of the three-dimensional Navier–Stokes equations”, *Commun. Math. Sci.* **10**:1 (2012), 131–136. [MR 2901304](#)
- [Heywood 1980] J. G. Heywood, “The Navier–Stokes equations: on the existence, regularity and decay of solutions”, *Indiana Univ. Math. J.* **29**:5 (1980), 639–681. [MR 81k:35131](#) [Zbl 0494.35077](#)
- [Iskauriaza et al. 2003] L. Iskauriaza, G. A. Serëgin, and V. Shverak, “ $L_{3,\infty}$ -solutions of Navier–Stokes equations and backward uniqueness”, *Uspekhi Mat. Nauk* **58**:2(350) (2003), 3–44. In Russian. Translated in *Russian Math. Surveys*, **58**:2 (2003), 211–250. [MR 2004m:35204](#) [Zbl 1064.35134](#)
- [Kato 1984] T. Kato, “Strong L^p -solutions of the Navier–Stokes equation in \mathbf{R}^m , with applications to weak solutions”, *Math. Z.* **187**:4 (1984), 471–480. [MR 86b:35171](#) [Zbl 0545.35073](#)
- [Kato and Ponce 1988] T. Kato and G. Ponce, “Commutator estimates and the Euler and Navier–Stokes equations”, *Comm. Pure Appl. Math.* **41**:7 (1988), 891–907. [MR 90f:35162](#) [Zbl 0671.35066](#)
- [Koch and Tataru 2001] H. Koch and D. Tataru, “Well-posedness for the Navier–Stokes equations”, *Adv. Math.* **157**:1 (2001), 22–35. [MR 2001m:35257](#) [Zbl 0972.35084](#)
- [Ladyzhenskaya 1967] O. A. Ladyzhenskaya, “Uniqueness and smoothness of generalized solutions of Navier–Stokes equations”, *Zap. Naučn. Sem. Leningrad. Otdel. Mat. Inst. Steklov. (LOMI)* **5** (1967), 169–185. In Russian. Translated in *Sem. Math. Steklov Math. Inst., Leningrad* **5** (1969), 60–67. [MR 38 #4836](#)
- [Lemarie-Rieusset 1992] P. G. Lemarie-Rieusset, “Analyses multi-résolutions non orthogonales, commutation entre projecteurs et dérivation et ondelettes vecteurs à divergence nulle”, *Rev. Mat. Iberoamericana* **8**:2 (1992), 221–237. [MR 94d:42044](#)
- [Lemarié-Rieusset 1999] P. G. Lemarié-Rieusset, “Solutions faibles d’énergie infinie pour les équations de Navier–Stokes dans \mathbb{R}^3 ”, *C. R. Acad. Sci. Paris Sér. I Math.* **328**:12 (1999), 1133–1138. [MR 2000d:35179](#) [Zbl 0930.35125](#)
- [Leray 1934] J. Leray, “Sur le mouvement d’un liquide visqueux emplissant l’espace”, *Acta. Math.* **63** (1934), 183–248. [JFM 60.0726.05](#)
- [Lions 1969] J.-L. Lions, *Quelques méthodes de résolution des problèmes aux limites non linéaires*, Dunod, 1969. [MR 41 #4326](#) [Zbl 0189.40603](#)
- [Lions 1996] P.-L. Lions, *Mathematical topics in fluid mechanics, I: Incompressible models*, Oxford Lecture Series in Mathematics and its Applications **3**, Oxford University Press, New York, 1996. [MR 98b:76001](#) [Zbl 0866.76002](#)
- [Prodi 1959] G. Prodi, “Un teorema di unicITÀ per le equazioni di Navier–Stokes”, *Ann. Mat. Pura Appl.* (4) **48** (1959), 173–182. [MR 23 #A3384](#) [Zbl 0148.08202](#)
- [Qian 2009] Z. Qian, “An estimate for the vorticity of the Navier–Stokes equation”, *C. R. Math. Acad. Sci. Paris* **347**:1-2 (2009), 89–92. [MR 2010h:35296](#) [Zbl 1155.35070](#)
- [Rusin and Šverák 2011] W. Rusin and V. Šverák, “Minimal initial data for potential Navier–Stokes singularities”, *J. Funct. Anal.* **260**:3 (2011), 879–891. [MR 2011m:35276](#) [Zbl 1206.35199](#)
- [Scheffer 1976] V. Scheffer, “Partial regularity of solutions to the Navier–Stokes equations”, *Pacific J. Math.* **66**:2 (1976), 535–552. [MR 56 #12677](#) [Zbl 0325.35064](#)

- [Serrin 1962] J. Serrin, “On the interior regularity of weak solutions of the Navier–Stokes equations”, *Arch. Rational Mech. Anal.* **9** (1962), 187–195. [MR 25 #346](#) [Zbl 0106.18302](#)
- [Serrin 1963] J. Serrin, “The initial value problem for the Navier–Stokes equations”, pp. 69–98 in *Nonlinear problems* (Madison, WI, 1962), edited by R. E. Langer, Univ. of Wisconsin Press, Madison, WI, 1963. [MR 27 #442](#) [Zbl 0115.08502](#)
- [Struwe 1988] M. Struwe, “On partial regularity results for the Navier–Stokes equations”, *Comm. Pure Appl. Math.* **41**:4 (1988), 437–458. [MR 89h:35270](#) [Zbl 0632.76034](#)
- [Tao 2006] T. Tao, *Nonlinear dispersive equations: local and global analysis*, CBMS Regional Conference Series in Mathematics **106**, Conference Board of the Mathematical Sciences, Washington, DC, 2006. [MR 2008i:35211](#) [Zbl 1106.35001](#)
- [Tao 2007] T. Tao, “A quantitative formulation of the global regularity problem for the periodic Navier–Stokes equation”, *Dyn. Partial Differ. Equ.* **4**:4 (2007), 293–302. [MR 2009d:35255](#) [Zbl 1137.35055](#)
- [Vasseur 2010] A. Vasseur, “Higher derivatives estimate for the 3D Navier–Stokes equation”, *Ann. Inst. H. Poincaré Anal. Non Linéaire* **27**:5 (2010), 1189–1204. [MR 2011k:35172](#) [Zbl 05800075](#)

Received 4 Aug 2011. Revised 7 Aug 2011. Accepted 23 May 2012.

TERENCE TAO: tao@math.ucla.edu

Department of Mathematics, University of California, Los Angeles, Los Angeles, CA 90095-1555, United States

Analysis & PDE

msp.org/apde

EDITORS

EDITOR-IN-CHIEF

Maciej Zworski

zworski@math.berkeley.edu

University of California

Berkeley, USA

BOARD OF EDITORS

Michael Aizenman	Princeton University, USA aizenman@math.princeton.edu	Nicolas Burq	Université Paris-Sud 11, France nicolas.burq@math.u-psud.fr
Luis A. Caffarelli	University of Texas, USA caffarel@math.utexas.edu	Sun-Yung Alice Chang	Princeton University, USA chang@math.princeton.edu
Michael Christ	University of California, Berkeley, USA mchrist@math.berkeley.edu	Charles Fefferman	Princeton University, USA cf@math.princeton.edu
Ursula Hamenstaedt	Universität Bonn, Germany ursula@math.uni-bonn.de	Nigel Higson	Pennsylvania State University, USA higson@math.psu.edu
Vaughan Jones	University of California, Berkeley, USA vfr@math.berkeley.edu	Herbert Koch	Universität Bonn, Germany koch@math.uni-bonn.de
Izabella Laba	University of British Columbia, Canada ilaba@math.ubc.ca	Gilles Lebeau	Université de Nice Sophia Antipolis, France lebeau@unice.fr
László Lempert	Purdue University, USA lempert@math.purdue.edu	Richard B. Melrose	Massachusetts Institute of Technology, USA rbm@math.mit.edu
Frank Merle	Université de Cergy-Pontoise, France Frank.Merle@u-cergy.fr	William Minicozzi II	Johns Hopkins University, USA minicozz@math.jhu.edu
Werner Müller	Universität Bonn, Germany mueller@math.uni-bonn.de	Yuval Peres	University of California, Berkeley, USA peres@stat.berkeley.edu
Gilles Pisier	Texas A&M University, and Paris 6 pisier@math.tamu.edu	Tristan Rivière	ETH, Switzerland riviere@math.ethz.ch
Igor Rodnianski	Princeton University, USA irod@math.princeton.edu	Wilhelm Schlag	University of Chicago, USA schlag@math.uchicago.edu
Sylvia Serfaty	New York University, USA serfaty@cims.nyu.edu	Yum-Tong Siu	Harvard University, USA siu@math.harvard.edu
Terence Tao	University of California, Los Angeles, USA tao@math.ucla.edu	Michael E. Taylor	Univ. of North Carolina, Chapel Hill, USA met@math.unc.edu
Gunther Uhlmann	University of Washington, USA gunther@math.washington.edu	András Vasy	Stanford University, USA andras@math.stanford.edu
Dan Virgil Voiculescu	University of California, Berkeley, USA dvv@math.berkeley.edu	Steven Zelditch	Northwestern University, USA zelditch@math.northwestern.edu

PRODUCTION

production@msp.org

Silvio Levy, Scientific Editor

See inside back cover or msp.org/apde for submission instructions.

The subscription price for 2013 is US \$160/year for the electronic version, and \$310/year (+\$35, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues from the last three years and changes of subscribers address should be sent to MSP.

Analysis & PDE (ISSN 1948-206X electronic, 2157-5045 printed) at Mathematical Sciences Publishers, 798 Evans Hall #3840, c/o University of California, Berkeley, CA 94720-3840, is published continuously online. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices.

APDE peer review and production are managed by EditFLOW[®] from Mathematical Sciences Publishers.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2013 Mathematical Sciences Publishers

ANALYSIS & PDE

Volume 6 No. 1 2013

Some results on scattering for log-subcritical and log-supercritical nonlinear wave equations	1
HSI-WEI SHIH	
Localisation and compactness properties of the Navier–Stokes global regularity problem	25
TERENCE TAO	
A variational principle for correlation functions for unitary ensembles, with applications	109
DORON S. LUBINSKY	
Relative Kähler–Ricci flows and their quantization	131
ROBERT J. BERMAN	
Resolvent estimates for elliptic quadratic differential operators	181
MICHAEL HITRIK, JOHANNES SJÖSTRAND and JOE VIOLA	
Bilinear Hilbert transforms along curves I: the monomial case	197
XIAOCHUN LI	
A gluing formula for the analytic torsion on singular spaces	221
MATTHIAS LESCH	