

# ANALYSIS & PDE

Volume 6

No. 2

2013

KENICHI ITO AND SHU NAKAMURA

**MICROLOCAL PROPERTIES OF SCATTERING MATRICES FOR  
SCHRÖDINGER EQUATIONS ON SCATTERING MANIFOLDS**



# MICROLOCAL PROPERTIES OF SCATTERING MATRICES FOR SCHRÖDINGER EQUATIONS ON SCATTERING MANIFOLDS

KENICHI ITO AND SHU NAKAMURA

Let  $M$  be a scattering manifold, i.e., a Riemannian manifold with an asymptotically conic structure, and let  $H$  be a Schrödinger operator on  $M$ . One can construct a natural time-dependent scattering theory for  $H$  with a suitable reference system, and a scattering matrix is defined accordingly. We show here that the scattering matrices are Fourier integral operators associated to a canonical transform on the boundary manifold generated by the geodesic flow. In particular, we learn that the wave front sets are mapped according to the canonical transform. These results are generalizations of a theorem by Melrose and Zworski, but the framework and the proof are quite different. These results may be considered as generalizations or refinements of the classical off-diagonal smoothness of the scattering matrix for two-body quantum scattering on Euclidean spaces.

## 1. Introduction

Let  $M$  be an  $n$ -dimensional smooth noncompact manifold such that  $M = M_c \cup M_\infty$ , where  $M_c$  is relatively compact, and  $M_\infty$  is diffeomorphic to  $\mathbb{R}_+ \times \partial M$ , where  $\partial M$  is a compact manifold. In the following, we often identify  $M_\infty$  with  $\mathbb{R}_+ \times \partial M$ , and we also suppose  $M_c \cap M_\infty \subset (0, 1) \times \partial M$  under this identification.

We recall the construction of the model introduced in [Ito and Nakamura 2010]. Let  $\{\varphi_\alpha : U_\alpha \rightarrow \mathbb{R}^{n-1}\}$ ,  $U_\alpha \subset \partial M$ , be a local coordinate system of  $\partial M$ . We take

$$\{\tilde{\varphi}_\alpha = I \otimes \varphi_\alpha : \tilde{U}_\alpha = \mathbb{R}_+ \times U_\alpha \rightarrow \mathbb{R} \times \mathbb{R}^{n-1}\}$$

as the local coordinate system for  $M_\infty \cong \mathbb{R}_+ \times \partial M$ , and we use  $(r, \theta) \in \mathbb{R} \times \mathbb{R}^{n-1}$  to represent a point in  $M_\infty$ .

We suppose  $\partial M$  is equipped with a smooth strictly positive density  $H = H(\theta)$  and a positive  $(2, 0)$ -tensor  $h = (h^{jk}(\theta))$  on  $\partial M$ . We let

$$Q = -\frac{1}{2} \sum_{j,k} H(\theta)^{-1} \frac{\partial}{\partial \theta_j} H(\theta) h^{jk}(\theta) \frac{\partial}{\partial \theta_k} \quad \text{on } \mathcal{H}_b = L^2(\partial M, H(\theta)d\theta).$$

$Q$  is an essentially self-adjoint operator on  $\mathcal{H}_b$ , and we denote its unique self-adjoint extension by the same symbol  $Q$ .

---

Ito was partially supported by a JSPS Research Fellowship during 2006–2007 and JSPS Wakate (B) grant number 21740090 during 2009–2012. Nakamura was partially supported by JSPS Kiban (B) grant number 17340033 during 2005–2008, and Kiban (A) grant number 21244008 during 2009–2013.

MSC2010: 35P25, 35S30, 58J40, 58J50.

Keywords: scattering matrix, scattering manifolds, Schrödinger operators, semiclassical analysis.



We let  $G$  be a smooth strictly positive density on  $M$  such that

$$G(x) dx = r^{n-1} H(\theta) dr d\theta \quad \text{on } (1, \infty) \times \partial M \subset M_\infty,$$

and we set  $\mathcal{H} = L^2(M, G(x) dx)$ . Let  $P$  be a formally self-adjoint second order elliptic operator on  $M$  such that

$$P = -\frac{1}{2} G^{-1}(\partial_r, \partial_\theta/r) G \begin{pmatrix} a_1 & a_2 \\ a_2 & a_3 \end{pmatrix} \begin{pmatrix} \partial_r \\ \partial_\theta/r \end{pmatrix} + V \quad \text{on } M_\infty,$$

where  $\begin{pmatrix} a_1 & a_2 \\ a_2 & a_3 \end{pmatrix}$  defines a real-valued smooth tensor and  $V$  is a real-valued smooth function. As in [Ito and Nakamura 2010], we introduce the following assumption:

**Assumption A.** There is  $\mu > 0$  such that for any  $\ell \in \mathbb{Z}_+$ ,  $\alpha \in \mathbb{Z}_+^{n-1}$ , there is  $C_{\ell\alpha} > 0$  and

$$\begin{aligned} |\partial_r^\ell \partial_\theta^\alpha (a_1(r, \theta) - 1)| &\leq C_{\ell\alpha} r^{-1-\mu-\ell}, & |\partial_r^\ell \partial_\theta^\alpha a_2(r, \theta)| &\leq C_{\ell\alpha} r^{-\mu-\ell}, \\ |\partial_r^\ell \partial_\theta^\alpha (a_3(r, \theta) - h(\theta))| &\leq C_{\ell\alpha} r^{-\mu-\ell}, & |\partial_r^\ell \partial_\theta^\alpha V(r, \theta)| &\leq C_{\ell\alpha} r^{-1-\mu-\ell}, \end{aligned}$$

in each local coordinate of  $M_\infty$  described above.

We may consider  $P$  as a *short range* perturbation of  $-\frac{1}{2}\partial_r^2 + \frac{1}{r^2}Q$ , but we will use different operators to construct a scattering theory. It is known that  $P$  is essentially self-adjoint, that  $\sigma_{ess}(P) = [0, \infty)$ , and that  $P$  is absolutely continuous except on a countable discrete spectrum, the only possible accumulation point being 0 (see [Ito and Nakamura 2010] and references therein). We construct a time-dependent scattering theory for  $H$  as follows: We set

$$M_f = \mathbb{R} \times \partial M, \quad \mathcal{H}_f = L^2(M_f, H(\theta) dr d\theta), \quad P_f = -\frac{1}{2} \frac{\partial^2}{\partial r^2} \quad \text{on } M_f.$$

$P_f$  is the one-dimensional free Schrödinger operator, and it is self-adjoint with  $\mathcal{D}(P_f) = H^2(\mathbb{R}) \otimes \mathcal{H}_b$ . Let  $j(r) \in C^\infty(\mathbb{R})$  such that  $j(r) = 0$  on  $(-\infty, \frac{1}{2}]$  and  $j(r) = 1$  on  $[1, \infty)$ . We define  $\mathcal{F} : \mathcal{H}_f \rightarrow \mathcal{H}$  by

$$(\mathcal{F}\varphi)(r, \theta) = r^{-(n-1)/2} j(r)\varphi(r, \theta) \quad \text{if } (r, \theta) \in M_\infty,$$

and  $(\mathcal{F}\varphi)(x) = 0$  if  $x \notin M_\infty$ . We define the wave operators by

$$W_\pm = W_\pm(P, P_f, \mathcal{F}) = \text{s-lim}_{t \rightarrow \pm\infty} e^{itP} \mathcal{F} e^{-itP_f}.$$

It is shown in [Ito and Nakamura 2010] that these operators exist and are complete in the following sense. Let  $\mathcal{F}$  be the Fourier transform in  $r$ , i.e.,

$$(\mathcal{F}\varphi)(\rho, \theta) = (2\pi)^{-1/2} \int_{-\infty}^\infty e^{-ir\rho} \varphi(r, \theta) dr \quad \text{for } \varphi \in C_0^\infty(M_f),$$

and extend it to a unitary map in  $L^2(M_f)$ . If we set

$$\mathcal{H}_{f,\pm} = \{\varphi \in \mathcal{H}_f \mid \text{supp}(\mathcal{F}\varphi) \subset \overline{\mathbb{R}_\pm} \times \partial M\},$$

then  $\mathcal{H}_f = \mathcal{H}_{f,+} \oplus \mathcal{H}_{f,-}$ . We consider  $W_\pm$  as maps from  $\mathcal{H}_{f,\pm}$  to  $\mathcal{H}$ ; they are asymptotically complete,

i.e., unitary operators from  $\mathcal{H}_{f,\pm}$  to  $\mathcal{H}_{ac}(P)$  [ibid., Theorem 2]. Then the scattering operator defined by

$$S = W_+^* W_- : \mathcal{H}_{f,-} \rightarrow \mathcal{H}_{f,+}$$

is unitary. By the intertwining property ( $P_f S = S P_f$ ), there is  $S(\lambda) \in \mathcal{B}(\mathcal{H}_b)$  for  $\lambda > 0$  such that

$$(\mathcal{F} S \mathcal{F}^{-1} \varphi)(\rho, \cdot) = S(\rho^2/2) \varphi(-\rho, \cdot) \quad \text{for } \rho > 0, \varphi \in \mathcal{F} \mathcal{H}_{f,-}.$$

$S(\lambda)$  is our scattering matrix, and we study its microlocal properties.

Let

$$q(\theta, \omega) = \frac{1}{2} \sum_{j,k} h^{jk}(\theta) \omega_j \omega_k \quad \text{for } (\theta, \omega) \in T^* \partial M$$

be the classical Hamiltonian associated to  $Q$ . We denote the Hamilton flow generated by  $b$  by  $\exp(tH_b)$  for  $t \in \mathbb{R}$ .

**Theorem 1.1.** *Suppose Assumption A holds, and let  $u \in \mathcal{H}_b$ . Then*

$$WF(S(\lambda)u) = \exp(\pi H_{\sqrt{2q}}) WF(u),$$

where  $WF(u)$  denotes the wave front set of  $u$ .

If  $\mu = 1$ , then we can show  $S(\lambda)$  is a Fourier integral operator (FIO). This is a slight extension of a theorem by Melrose and Zworski [1996].

**Theorem 1.2.** *Suppose Assumption A holds with  $\mu = 1$ . Then for each  $\lambda > 0$ ,  $S(\lambda)$  is an FIO associated to  $\exp(\pi H_{\sqrt{2q}})$ .*

If  $0 < \mu < 1$ , then  $S(\lambda)$  is not necessarily an FIO in the usual sense, but we can still show it is an FIO in a generalized sense:

**Theorem 1.3.** *Suppose Assumption A holds, and let  $S(\lambda)$  be the scattering matrix defined as above. Then for each  $\lambda > 0$ ,  $S(\lambda)$  is an FIO associated to an asymptotically homogeneous canonical transform in  $T^* \partial M$ , which is asymptotic to  $\exp(\pi H_{\sqrt{2q}})$  as  $\omega \rightarrow \infty$ .*

The exact definition of the phrase *an FIO associated to an asymptotically homogeneous canonical transform* is given in [Ito and Nakamura 2012], and we discuss it in Section 6.

**Remark 1.4.** Since we do not introduce a Riemannian metric, our model looks rather different from the scattering metric defined by Melrose [1994; Melrose and Zworski 1996]. However, as explained in [Ito and Nakamura 2010, Appendix A], the Laplacian on scattering manifolds is a special case of our model. Namely, their model corresponds to the case that  $\mu = 1$  and that each  $a_j$  has asymptotic expansion in  $r^{-1}$  as  $r \rightarrow \infty$  and  $V = 0$ .

Theorems 1.1 and 1.2 are essentially corollaries of Theorem 1.3, but they can be proved by a simpler argument than Theorem 1.3. We feel the simpler argument is interesting in itself, and we first prove Theorems 1.1 and 1.2, and then we refine the argument to prove Theorem 1.3 later.

The main idea to prove Theorems 1.1–1.3 is to consider the evolution

$$A(t) = e^{itP_f/h} \mathcal{G}^* e^{-itP/h} a(hr, D_r, \theta, hD_\theta) e^{itP/h} \mathcal{G} e^{-itP_f/h}$$

with some symbol  $a$ , and use an argument similar to Egorov's theorem for this time-dependent operator. We use a semiclassical argument, i.e., we consider the asymptotic behavior of the operator as  $h \rightarrow 0$ . We consider  $W(t) = e^{itP_f/h} g^* e^{-itP/h}$  as a time-evolution, and then construct an asymptotic solution for  $A(t)$  (with slight modifications) as a solution to a Heisenberg equation. The construction of the asymptotic solution relies on the classical Hamilton flow generated by  $p$ , the symbol of  $P$ . The dominant part of the symbol  $p$  is given by the unperturbed conic Hamiltonian:  $p_c = \frac{1}{2}\rho^2 + \frac{1}{r^2}q(\theta, \omega)$ . The classical scattering operator for the pair  $p_c$  and  $p_f = \frac{1}{2}\rho^2$  is explicitly computed, and it is  $\exp(\pi H_{\sqrt{2q}})$ , which appears in the statement of our main theorems. Thus, one may consider our results as a quantization of the classical mechanical scattering on the scattering manifold. More precisely, we show that the canonical transform appearing in [Theorem 1.3](#) is actually the classical scattering map for the pair  $p$  and  $p_f$ , which is not necessarily homogeneous, and we need to use the method of FIOs with asymptotically conic Lagrangian manifolds.

As mentioned in the beginning, [Theorem 1.2](#) is slight generalization of the Melrose–Zworski theorem [\[1996\]](#) (see also [\[Vasy 1998\]](#) for a simplification of the theory). They used the theory of Legendre distribution and the notion of scattering wave front sets, whereas we use relatively elementary pseudodifferential operator calculus with somewhat nonstandard symbol classes, and a Beals-type characterization of FIOs. We also note that our proof, as well as the setting, are time-dependent-theoretical, and we investigate the scattering phenomena directly to obtain the properties of the wave operators and scattering operators, whereas the Melrose–Zworski paper relies on the stationary, generalized eigenfunction expansion theory.

Our method is closely related to our previous works on the propagation of singularities for Schrödinger evolution equations [\[Nakamura 2009a; 2009b; Ito and Nakamura 2009; 2012\]](#). In these works, we considered singularities of solutions, which are described by their high energy behavior, whereas in the scattering phenomena we are concerned with the large  $r$  behavior (which in turn is related to the high  $|\omega|$  behavior, where  $\omega$  is the conjugate variable to  $\theta \in \partial M$ ). Thus we are forced to use different symbol classes in the calculus, and the corresponding classical mechanics look slightly different, but the general strategy is essentially the same as in these papers.

If  $M = \mathbb{R}^n$  and the Hamiltonian  $P$  is a short-range perturbation of the Laplacian  $-\frac{1}{2}\Delta$ , then the canonical map  $\exp(\pi H_{\sqrt{2q}})$  is the antipodal map on  $T^*S^{n-1}$ . In this case, the off-diagonal smoothness of the scattering cross-section is well-known (see [\[Isozaki and Kitada 1986\]](#), and [Section 9.4](#) and the references of [\[Yafaev 2000\]](#)), and our result (as well as the Melrose–Zworski theorem) may be considered as its generalizations. For such models, our result implies the scattering matrix is an FIO (associated to a canonical map which is asymptotic to the identity map), and if  $\mu = 1$  then it is in fact a pseudodifferential operator. It is also not difficult to show from our argument that the scattering matrix is a pseudodifferential operator with symbol in  $S_{\mu,0}^0(S^{n-1})$  if  $\mu \in (0, 1)$ .

The paper is organized as follows. In [Section 2](#), we discuss Hamilton flows generated by  $p_c$  and  $p$ , and their scattering theory. In [Section 3](#), we prepare the symbol calculus on the scattering manifolds. In [Section 4](#), we discuss an Egorov-type theorem and the construction of asymptotic solutions, which are sufficient to show [Theorems 1.1](#) and [1.2](#). We prove [Theorems 1.1](#) and [1.2](#) in [Section 5](#). In [Section 6](#), we discuss the modification of the argument to show [Theorem 1.3](#). We discuss a local decay estimate

necessary in the proof in [Appendix A](#). A Beals-type characterization, or an inverse of Egorov’s theorem, is discussed in [Appendix B](#), along with a technical lemma on FIOs used in the proof.

Throughout this paper, we use the following notation: For norm spaces  $X$  and  $Y$ , the space of bounded linear maps is denoted by  $B(X, Y)$ , and if  $X = Y$ , we also write  $B(X, X) = B(X)$ . More generally, if  $X$  and  $Y$  are topological linear spaces, the space of continuous linear maps is denoted by  $\mathcal{L}(X, Y)$ . For a symbol  $g$  on  $T^*X$  with a manifold  $X$ , we denote by  $\exp(tH_g)$  the Hamilton flow generated by the Hamilton vector field

$$H_g = \frac{\partial g}{\partial \xi} \cdot \frac{\partial}{\partial x} - \frac{\partial g}{\partial x} \cdot \frac{\partial}{\partial \xi}.$$

We also write  $T^*X \setminus 0 = \{(x, \xi) \in T^*X \mid \xi \neq 0\}$ .

### 2. Classical flow and scattering theory

In this section, we consider the classical mechanics, or the Hamilton flow for the Hamiltonian with conic structure on  $T^*M_\infty$ , where  $M_\infty = \mathbb{R}_+ \times \partial M$ , and then the Hamilton flow generated by the principal symbol of  $P$ .

**Exact solutions to the conic Hamilton flow.** We set

$$p_c(r, \rho, \theta, \omega) = \frac{1}{2}\rho^2 + \frac{1}{r^2}q(\theta, \omega) \quad \text{and} \quad q(\theta, \omega) = \frac{1}{2} \sum_{j,k} h^{jk}(\theta)\omega_j\omega_k$$

on  $T^*M_\infty \cong T^*\mathbb{R}_+ \times T^*\partial M$ . We consider

$$(r(t), \rho(t), \theta(t), \omega(t)) = \exp(tH_{p_c})(r_0, \rho_0, \theta_0, \omega_0),$$

with  $(r_0, \rho_0, \theta_0, \omega_0) \in T^*\mathbb{R}_+ \times (T^*\partial M \setminus 0)$ , that is, with  $\omega_0 \neq 0$ . It satisfies the Hamilton equation

$$\begin{aligned} r'(t) &= \frac{\partial p_c}{\partial \rho} = \rho(t), & \rho'(t) &= -\frac{\partial p_c}{\partial r} = \frac{2}{r(t)^3}q(\theta(t), \omega(t)), \\ \theta'(t) &= \frac{\partial p_c}{\partial \omega} = \frac{1}{r(t)^2} \frac{\partial q}{\partial \omega}(\theta(t), \omega(t)), & \omega'(t) &= \frac{\partial p_c}{\partial \theta} = -\frac{1}{r(t)^2} \frac{\partial q}{\partial \theta}(\theta(t), \omega(t)). \end{aligned}$$

The solution has two invariants: total energy  $E_0 = p_c(r_0, \rho_0, \theta_0, \omega_0)$  and angular energy  $q_0 = q(\theta_0, \omega_0)$ . (The conservation of the total energy follows from  $\{p_c, p_c\} = 0$ , and of the angular energy from  $\{q, p_c\} = \frac{1}{2}\{q, \rho^2\} + \{q, \frac{1}{r^2}\}q + \frac{1}{r^2}\{q, q\} = 0$ .) Then  $(r(t), \rho(t))$  satisfies

$$r'(t) = \rho(t), \quad \rho'(t) = \frac{2}{r(t)^3}q_0,$$

which is independent of  $(\theta(t), \omega(t))$ . Noting that  $(r^2(t))'' = 4E_0$ , we can easily solve this equation to obtain

$$r(t) = \sqrt{2E_0t^2 + 2r_0\rho_0t + r_0^2}, \quad \rho(t) = \frac{2E_0t + r_0\rho_0}{\sqrt{2E_0t^2 + 2r_0\rho_0t + r_0^2}}, \quad t \in \mathbb{R}.$$

We now set

$$\tau(t) = \int_0^t \frac{ds}{r(s)^2} = \frac{1}{\sqrt{2q_0}} \left( \tan^{-1} \frac{2E_0t + r_0\rho_0}{\sqrt{2q_0}} - \tan^{-1} \frac{r_0\rho_0}{\sqrt{2q_0}} \right).$$

Then  $(\theta(t), \omega(t))$  satisfies

$$\frac{\partial \theta}{\partial \tau} = \frac{\partial q}{\partial \omega}(\theta, \omega), \quad \frac{\partial \omega}{\partial \tau} = -\frac{\partial q}{\partial \theta}(\theta, \omega),$$

and hence we learn that

$$(\theta(t), \omega(t)) = \exp(\tau(t)H_q)(\theta_0, \omega_0).$$

Moreover, if we set  $\sigma(t) = \sqrt{2q_0} \cdot \tau(t)$ , then we learn that

$$\frac{\partial \theta}{\partial \sigma} = \frac{1}{\sqrt{2q}} \frac{\partial q}{\partial \omega} = \frac{\partial \sqrt{2q}}{\partial \omega}(\theta, \omega), \quad \frac{\partial \omega}{\partial \sigma} = -\frac{1}{\sqrt{2q}} \frac{\partial q}{\partial \theta} = -\frac{\partial \sqrt{2q}}{\partial \theta}(\theta, \omega),$$

and hence that

$$(\theta(t), \omega(t)) = \exp(\sigma(t)H_{\sqrt{2q}})(\theta_0, \omega_0).$$

Note that  $\exp(tH_{\sqrt{2q}})$  is the geodesic flow on  $\partial M$  with respect to the (co)metric  $(h^{jk}(\theta))$  on  $T^*\partial M$ .

**Classical mechanical wave operators and a scattering operator for the conic Hamilton flow.** Now we consider the asymptotics as  $t \rightarrow \pm\infty$ . We set

$$\begin{aligned} r_{\pm} &= \lim_{t \rightarrow \pm\infty} \tilde{r}(t) = \lim_{t \rightarrow \pm\infty} (r(t) - t\rho(t)) = \pm \frac{r_0\rho_0}{\sqrt{2E_0}}, \\ \rho_{\pm} &= \lim_{t \rightarrow \pm\infty} \rho(t) = \pm\sqrt{2E_0}, \\ (\theta_{\pm}, \omega_{\pm}) &= \lim_{t \rightarrow \pm\infty} (\theta(t), \omega(t)) = \exp(\sigma_{\pm}H_{\sqrt{2q}})(\theta_0, \omega_0), \end{aligned}$$

where  $\sigma_{\pm} = \pm\frac{1}{2}\pi - \tan^{-1}(r_0\rho_0/\sqrt{2q_0})$ . Note we need a modification only for  $r(t)$ .  $(r_{\pm}, \rho_{\pm}, \theta_{\pm}, \omega_{\pm})$  are the scattering data for the trajectory  $(r(t), \rho(t), \theta(t), \omega(t))$ . We also note the identities

$$E_0 = \frac{1}{2}\rho_0^2 + \frac{1}{r_0^2}q_0 = \frac{1}{2}\rho_{\pm}^2, \quad r_0\rho_0 = r_{\pm}\rho_{\pm}, \quad q_0 = q(\theta_{\pm}, \omega_{\pm}).$$

Using these, we can solve  $(r_0, \rho_0, \theta_0, \omega_0)$  for given  $(r_{\pm}, \rho_{\pm}, \theta_{\pm}, \omega_{\pm})$  if  $\pm\rho_{\pm} > 0$  and  $\omega_{\pm} \neq 0$ :

$$r_0 = \sqrt{r_{\pm}^2 + 2q_0/\rho_{\pm}^2}, \quad \rho_0 = \frac{r_{\pm}\rho_{\pm}}{\sqrt{r_{\pm}^2 + 2q_0/\rho_{\pm}^2}}, \quad (\theta_0, \omega_0) = \exp(-\sigma_{\pm}H_{\sqrt{2q}})(\theta_{\pm}, \omega_{\pm}),$$

where  $\sigma_{\pm} = \pm\frac{1}{2}\pi - \tan^{-1}(r_{\pm}\rho_{\pm}/\sqrt{2q})$ . We define the classical wave operators (for the pair  $p_c$  and  $p_f := \frac{1}{2}\rho^2$ ) by

$$w_{c,\pm} : (r_{\pm}, \rho_{\pm}, \theta_{\pm}, \omega_{\pm}) \mapsto (r_0, \rho_0, \theta_0, \omega_0).$$

We can also write

$$w_{c,\pm}(r_{\pm}, \rho_{\pm}, \theta_{\pm}, \omega_{\pm}) = \lim_{t \rightarrow \pm\infty} \exp(-tH_{p_c}) \circ \exp(tH_{p_f})(r_{\pm}, \rho_{\pm}, \theta_{\pm}, \omega_{\pm}).$$

It is easy to check that  $w_{c,\pm}$  are diffeomorphisms from  $\mathbb{R} \times \mathbb{R}_{\pm} \times (T^*\partial M \setminus 0)$  to  $\mathbb{R}_+ \times \mathbb{R} \times (T^*\partial M \setminus 0)$ . Hence the classical scattering operator

$$s_c = w_{c,+}^{-1} \circ w_{c,-} : (r_-, \rho_-, \theta_-, \omega_-) \mapsto (r_+, \rho_+, \theta_+, \omega_+)$$

is a diffeomorphism from  $\mathbb{R} \times \mathbb{R}_- \times (T^*\partial M \setminus 0)$  to  $\mathbb{R} \times \mathbb{R}_+ \times (T^*\partial M \setminus 0)$ . We can easily compute  $s_c$  explicitly, we have

$$s_c(r, \rho, \theta, \omega) = (-r, -\rho, \exp(\pi H_{\sqrt{2q}})(\theta, \rho)),$$

and this is the classical analogue of the Melrose–Zworski theorem.

We write

$$w_c(t) = \exp(-tH_{p_c}) \circ \exp(tH_{p_f}) \quad \text{so that } w_{c,\pm} = \lim_{t \rightarrow \pm\infty} w_c(t).$$

Let  $U \subset \mathbb{R}_+ \times \mathbb{R} \times (T^*\partial M \setminus 0)$  be a relatively compact domain. Then the convergence of  $w_c(t)^{-1}$  to  $w_{c,\pm}^{-1}$  (as  $t \rightarrow \pm\infty$ ) is uniform in  $U$ , along with all derivatives. Since the limit is a diffeomorphism, its inverse  $w(t)$  also has the same property (on  $w_c(t)^{-1}U$ ). In particular, all the derivatives of  $w_c(t)^{-1}$  on  $U$  are uniformly bounded in  $t$ , and all the derivatives of  $w_c(t)$  on  $w_c(t)^{-1}U$  are uniformly bounded.

We note that it is easy to check that  $w_{c,\pm}$  and hence  $s_c$  are homogeneous of order one with respect to the  $(r, \omega)$ -variables, i.e.,

$$w_{c,\pm}^{-1}(\lambda r_0, \rho_0, \theta_0, \lambda \omega_0) = (\lambda r_{\pm}, \rho_{\pm}, \theta_{\pm}, \lambda \omega) \quad \text{for } \lambda > 0.$$

This is consistent with the scaling property of  $w_c(t)$ :

$$w_c^{-1}(\lambda t)(\lambda r_0, \rho_0, \theta_0, \lambda \omega_0) = (\lambda \tilde{r}(t), \rho(t), \theta(t), \lambda \omega(t))$$

for any  $\lambda > 0, t \in \mathbb{R}$ .

**Classical flow generated by the scattering metric.** Here we discuss the Hamilton flow generated by the symbol of  $P$ :

$$p(r, \rho, \theta, \omega) = \frac{1}{2} \left( a_1(r, \theta)\rho^2 + \frac{2\rho a_2(r, \theta) \cdot \omega}{r} + \frac{\omega \cdot a_3(r, \theta)\omega}{r^2} \right) + V \tag{2-1}$$

on  $T^*M_\infty$ .

We let  $\Omega_0 \Subset T^*\mathbb{R}_+ \times (T^*\partial M \setminus 0)$ . For  $h \in (0, 1]$ , we set

$$\Omega_0^h = \{(r, \rho, \theta, \omega) \in T^*\mathbb{R}_+ \times (T^*\partial M \setminus 0) \mid (hr, \rho, \theta, h\omega) \in \Omega_0\},$$

and we consider the Hamilton flow with initial conditions in  $\Omega_0^h$ . We show that if  $h$  is sufficiently small then the classical (inverse) wave operators exist on  $\Omega_0^h$ , and they are very close to  $w_{c,\pm}^{-1}$ , the (inverse) wave operators for the conic metric.

**Theorem 2.1.** (i) *Let  $\Omega_0$  and  $\Omega_0^h$  as above. Then there is  $h_0 > 0$  such that if  $h \in (0, h_0]$ , then*

$$w_{\pm}^*(r, \rho, \theta, \omega) := \lim_{t \rightarrow \pm\infty} \exp(-tH_{p_f}) \circ \exp(tH_p)(r, \rho, \theta, \omega)$$

*exists for  $(r, \rho, \theta, \omega) \in \Omega_0^h$ , and the convergence holds in the  $C^\infty$ -topology on  $\Omega_0^h$ .*

(ii) *We write*

$$\begin{aligned} (r(t), \rho(t), \theta(t), \omega(t)) &= \exp(tH_p)(r, \rho, \theta, \omega), \\ (r_c(t), \rho_c(t), \theta_c(t), \omega_c(t)) &= \exp(tH_{p_c})(r, \rho, \theta, \omega), \end{aligned}$$



for  $(r, \rho, \theta, \omega) \in \Omega_0^h$ . Then for any indices  $\alpha, \beta, \gamma$  and  $\delta$ , there is  $C > 0$  such that

$$\begin{aligned} & |\partial_r^\alpha \partial_\rho^\beta \partial_\theta^\gamma \partial_\omega^\delta (r(t) - r_c(t))| + |\partial_r^\alpha \partial_\rho^\beta \partial_\theta^\gamma \partial_\omega^\delta (\omega(t) - \omega_c(t))| \leq Ch^{-1+\mu+|\alpha|+|\delta|}, \\ & |\partial_r^\alpha \partial_\rho^\beta \partial_\theta^\gamma \partial_\omega^\delta (\rho(t) - \rho_c(t))| + |\partial_r^\alpha \partial_\rho^\beta \partial_\theta^\gamma \partial_\omega^\delta (\theta(t) - \theta_c(t))| \leq Ch^{\mu+|\alpha|+|\delta|}, \end{aligned}$$

for  $(r, \rho, \theta, \omega) \in \Omega_0^h, t \in \mathbb{R}, 0 < h \leq h_0$ .

(iii) If we write

$$w_\pm^*(r, \rho, \theta, \omega) = (r_\pm, \rho_\pm, \theta_\pm, \omega_\pm) \quad \text{and} \quad w_{c,\pm}^*(r, \rho, \theta, \omega) = (r_{c,\pm}, \rho_{c,\pm}, \theta_{c,\pm}, \omega_{c,\pm}),$$

then

$$\begin{aligned} & |\partial_r^\alpha \partial_\rho^\beta \partial_\theta^\gamma \partial_\omega^\delta (r_\pm - r_{c,\pm})| + |\partial_r^\alpha \partial_\rho^\beta \partial_\theta^\gamma \partial_\omega^\delta (\omega_\pm - \omega_{c,\pm})| \leq Ch^{-1+\mu+|\alpha|+|\delta|}, \\ & |\partial_r^\alpha \partial_\rho^\beta \partial_\theta^\gamma \partial_\omega^\delta (\rho_\pm - \rho_{c,\pm})| + |\partial_r^\alpha \partial_\rho^\beta \partial_\theta^\gamma \partial_\omega^\delta (\theta_\pm - \theta_{c,\pm})| \leq Ch^{\mu+|\alpha|+|\delta|} \end{aligned}$$

for  $(r, \rho, \theta, \omega) \in \Omega_0^h, 0 < h \leq h_0$ .

For  $(r_0, \rho_0, \theta_0, \omega_0) \in \Omega_0$ , we define  $(r^h(t), \rho^h(t), \theta^h(t), \omega^h(t))$  so that

$$(h^{-1}r^h(t), \rho^h(t), \theta^h(t), h^{-1}\omega^h(t)) = \exp(h^{-1}tH_p)(h^{-1}r_0, \rho_0, \theta_0, h^{-1}\omega_0).$$

We also set

$$p^h(r, \rho, \theta, \omega) = p(h^{-1}r, \rho, \theta, h^{-1}\omega), \quad (r, \rho, \theta, \omega) \in T^*M_\infty.$$

Then it is easy to check that

$$(r^h(t), \rho^h(t), \theta^h(t), \omega^h(t)) = \exp(tH_{p^h})(r_0, \rho_0, \theta_0, \omega_0).$$

On the other hand, if we write

$$p^h(r, \rho, \theta, \omega) = p_c(r, \rho, \theta, \omega) + v^h(r, \rho, \theta, \omega),$$

then we learn by [Assumption A](#) that for any indices  $\alpha, \beta, \gamma, \delta$ ,

$$|\partial_r^\alpha \partial_\rho^\beta \partial_\theta^\gamma \partial_\omega^\delta v^h(r, \rho, \theta, \omega)| \leq C_{\alpha\beta\gamma\delta} h^\mu (r^{-1}\langle \rho \rangle^2 + r^{-1}\langle \rho \rangle \langle \omega \rangle + r^{-2}\langle \omega \rangle^2) r^{-\mu-|\alpha|} \langle \rho \rangle^{-|\beta|} \langle \omega \rangle^{-|\delta|}. \quad (2-2)$$

In order to prove [Theorem 2.1](#), it suffices to show:

**Theorem 2.2.** (i) *There is  $h_0 > 0$  such that if  $h \in (0, h_0]$ , then*

$$w_{\pm,h}^*(r_0, \rho_0, \theta_0, \omega_0) := \lim_{t \rightarrow \pm\infty} \exp(-tH_{p_f}) \circ \exp(tH_{p^h})(r_0, \rho_0, \theta_0, \omega_0)$$

*exists for  $(r_0, \rho_0, \theta_0, \omega_0) \in \Omega_0$ , and the convergence holds in the  $C^\infty$ -topology.*

(ii) *For any indices  $\alpha, \beta, \gamma, \delta$ , there is  $C > 0$  such that*

$$\begin{aligned} & |\partial_{r_0}^\alpha \partial_{\rho_0}^\beta \partial_{\theta_0}^\gamma \partial_{\omega_0}^\delta (r^h(t) - r_c(t))| + |\partial_{r_0}^\alpha \partial_{\rho_0}^\beta \partial_{\theta_0}^\gamma \partial_{\omega_0}^\delta (\rho^h(t) - \rho_c(t))| \\ & \quad + |\partial_{r_0}^\alpha \partial_{\rho_0}^\beta \partial_{\theta_0}^\gamma \partial_{\omega_0}^\delta (\theta^h(t) - \theta_c(t))| + |\partial_{r_0}^\alpha \partial_{\rho_0}^\beta \partial_{\theta_0}^\gamma \partial_{\omega_0}^\delta (\omega^h(t) - \omega_c(t))| \leq Ch^\mu \end{aligned}$$

for  $(r_0, \rho_0, \theta_0, \omega_0) \in \Omega_0, h \in (0, h_0], t \in \mathbb{R}$ , where

$$(r_c(t), \rho_c(t), \theta_c(t), \omega_c(t)) = \exp(tH_{p_c})(r_0, \rho_0, \theta_0, \omega_0).$$

(iii) *Writing*

$$(r_{\pm}^h, \rho_{\pm}^h, \theta_{\pm}^h, \omega_{\pm}^h) = w_{\pm, h}^*(r_0, \rho_0, \theta_0, \omega_0),$$

we have for any indices  $\alpha, \beta, \gamma, \delta$  that

$$\begin{aligned} & \left| \partial_{r_0}^{\alpha} \partial_{\rho_0}^{\beta} \partial_{\theta_0}^{\gamma} \partial_{\omega_0}^{\delta} (r_{\pm}^h - r_{c, \pm}) \right| + \left| \partial_{r_0}^{\alpha} \partial_{\rho_0}^{\beta} \partial_{\theta_0}^{\gamma} \partial_{\omega_0}^{\delta} (\rho_{\pm}^h - \rho_{c, \pm}) \right| \\ & \quad + \left| \partial_{r_0}^{\alpha} \partial_{\rho_0}^{\beta} \partial_{\theta_0}^{\gamma} \partial_{\omega_0}^{\delta} (\theta_{\pm}^h - \theta_{c, \pm}) \right| + \left| \partial_{r_0}^{\alpha} \partial_{\rho_0}^{\beta} \partial_{\theta_0}^{\gamma} \partial_{\omega_0}^{\delta} (\omega_{\pm}^h - \omega_{c, \pm}) \right| \leq Ch^{\mu}. \end{aligned}$$

*Proof of Theorem 2.2.* The proof is analogous to the arguments in [Nakamura 2009a, Section 2; Ito and Nakamura 2009, Section 2]. We only outline the proof, and we omit the details.

Step 1. By the standard virial-type argument, we learn that there is  $R > 0$  such that

$$\frac{d^2}{dt^2} (r^h(t)^2) \geq c > 0 \quad \text{if } r^h(t) \geq R,$$

if  $(r_0, \rho_0, \theta_0, \omega_0) \in \Omega_0$ . Here we use the fact that  $|\rho|$  and  $|\omega/r|$  are uniformly bounded by the conservation of energy. On the other hand, since  $v^h = O(h^{\mu})$ , we also learn that  $r^h(t) \rightarrow r_c(t)$  as  $h \downarrow 0$ , locally uniformly in  $t$ . Thus, if  $t_0$  is large and  $h$  is small enough,  $r^h(t) \geq R$ , and combining this with the above observation, we have

$$|r^h(t)| \geq \sqrt{R + c|t - t_0|^2/2} \quad \text{for } t \geq t_0.$$

Hence we learn

$$c_1(t) \leq r^h(t) \leq c_2(t) \quad \text{for } h \in (0, h_0], t > 0,$$

with some  $h_0, c_1, c_2 > 0$ . The case  $t < 0$  can be handled similarly.

Step 2. We consider the time evolution of  $q_0(t) = q(\theta^h(t), \omega^h(t))$ . By the Hamilton equation and (2-2), we have

$$\frac{d}{dt} q_0(t) = -\{p^h, q_0\} = -\{v^h, q_0\} = O(h^{\mu} r^{-1-\mu} \langle \omega \rangle^2) = O(h^{\mu} \langle t \rangle^{-1-\mu} (1 + q_0(t))).$$

Here we have used the boundedness of  $|\rho(t)|$  and  $|\omega(t)/r(t)|$  again. Then by the Duhamel formula, we learn that  $q_0(t)$  is uniformly bounded for initial conditions in  $\Omega_0$  and  $h \in (0, h_0]$ . This implies  $|\omega^h(t)|$  is also uniformly bounded.

Step 3. Combining these observations with the Hamilton equation, we learn that

$$\left| \frac{d\rho^h(t)}{dt} \right| \leq C \langle t \rangle^{-2-\mu}, \quad \left| \frac{d\theta^h(t)}{dt} \right| \leq C \langle t \rangle^{-1-\mu}, \quad \left| \frac{d}{dt} (r^h(t) - t\rho^h(t)) \right| \leq C \langle t \rangle^{-1-\mu}, \quad \left| \frac{d\omega^h(t)}{dt} \right| \leq C \langle t \rangle^{-1-\mu},$$

uniformly for  $(r_0, \rho_0, \theta_0, \omega_0) \in \Omega_0, h \in (0, h_0]$  and  $t \in \mathbb{R}$ . These imply the existence of  $w_{\pm, h}^*$  on  $\Omega_0$ . We can show the similar estimates for the derivatives, i.e.,

$$\left| \frac{d}{dt} (\partial_{r_0}^{\alpha} \partial_{\rho_0}^{\beta} \partial_{\theta_0}^{\gamma} \partial_{\omega_0}^{\delta} \rho^h(t)) \right| \leq C \langle t \rangle^{-2-\mu-|\alpha|},$$

and so on. These imply the convergence in  $C^{\infty}$ -topology, and we conclude that assertion (i) holds.

Step 4. We set

$$g^h(t) = |r^h(t) - r_c(t)| + |\rho^h(t) - \rho_c(t)| + |\theta^h(t) - \theta_c(t)| + |\omega^h(t) - \omega_c(t)|.$$

Then by the Hamilton equation, (2-2), and the estimates in Steps 1 and 2, we learn that

$$\left| \frac{d}{dt} g^h(t) \right| \leq C \langle t \rangle^{-1-\mu} g^h(t) + Ch^\mu \langle t \rangle^{-1-\mu}$$

uniformly for initial conditions in  $\Omega_0$  and  $h \in (0, h_0]$ . Then by using the Duhamel formula and noting that  $g^h(0) = 0$ , we obtain

$$|g^h(t)| \leq Ch^\mu, \quad t \in \mathbb{R}.$$

This is assertion (ii) with  $\alpha = \beta = \gamma = \delta = 0$ . The derivatives can be estimated similarly by induction. For the details of this argument, we refer to [Craig et al. 1995, Section 2; Nakamura 2009a, Section 2]. Assertion (iii) follows immediately from assertion (ii).  $\square$

By the above argument, we also learn that  $w_{\pm, h}^*$  are invertible for small  $h$ . The inverses are uniformly bounded, and their inverses

$$w_{\pm, h} = (w_{\pm, h}^*)^{-1}$$

are well-defined for  $h \in (0, h_0]$ . It follows that

$$w_{\pm} = (w_{\pm}^*)^{-1}$$

is well-defined and diffeomorphic on  $w_{\pm}^*[\Omega_0^h]$  with  $h \in (0, h_0]$ . Thus we can define the classical scattering operator by

$$s = w_+^* \circ w_-$$

on  $w_-^*[\Omega_0^h]$ , with sufficiently small  $h$ .

### 3. Symbol classes and their quantization on scattering manifolds

Here we prepare a pseudodifferential operator calculus which is used extensively in the proof of the main theorems. We refer to [Hörmander 1985; Taylor 1981, Chapter XVIII] for the standard theory of microlocal analysis.

In the following, we employ symbol calculus on  $T^*M$ , but we always suppose the symbol is supported in  $T^*M_\infty$ , and we use a local coordinate system as in Section 1. More specifically, we choose a local coordinate system on  $\partial M$ :  $\{\varphi_\alpha : U_\alpha \rightarrow \mathbb{R}^{n-1}\}$ ,  $U_\alpha \subset \partial M$ , and we use the coordinate system  $\{1 \otimes \varphi_\alpha : \mathbb{R}_+ \times U_\alpha \rightarrow \mathbb{R} \times \mathbb{R}^{n-1}\}$  on  $M_\infty$ . We also use a similar local coordinate system on  $M_f$ , defined by  $\{1 \otimes \varphi_\alpha : \mathbb{R} \times U_\alpha \rightarrow \mathbb{R} \times \mathbb{R}^{n-1}\}$ . We often identify  $U_\alpha$  (or  $\mathbb{R}_+ \times U_\alpha$ ,  $\mathbb{R} \times U_\alpha$ , respectively) with  $\text{Ran } \varphi_\alpha$  (or  $\text{Ran } (1 \otimes \varphi_\alpha)$ , respectively).

**Symbol classes.** We define a metric either on  $T^*M_\infty$  or  $T^*M_f$  by

$$g_1 = \frac{dr^2}{\langle r \rangle^2} + d\rho^2 + d\theta^2 + \frac{d\omega^2}{\langle \omega \rangle^2},$$

and consider symbols in  $S(m, g_1)$  with a weight function  $m$ , i.e.,  $a \in S(m, g_1)$  if and only if for any indices  $\alpha, \beta, \gamma, \delta$ , there is  $C$  such that

$$|\partial_r^\alpha \partial_\rho^\beta \partial_\theta^\gamma \partial_\omega^\delta a(r, \rho, \theta, \omega)| \leq Cm(r, \rho, \theta, \omega) \langle r \rangle^{-|\alpha|} \langle \omega \rangle^{-|\delta|}.$$

Later, we will consider the calculus of such symbols on sets  $\Omega^h = \{(r, \rho, \theta, \omega) \mid (hr, \rho, \theta, h\omega) \in \Omega\}$ , where  $\Omega \subset T^*M_\infty$  is some compact set (supported away from  $\{\omega = 0\}$ ) and  $h > 0$  is small. In such cases, the symbol satisfies

$$|\partial_r^\alpha \partial_\rho^\beta \partial_\theta^\gamma \partial_\omega^\delta a(h; r, \rho, \theta, \omega)| \leq Cm(h)h^{|\alpha|+|\delta|},$$

and we denote such a ( $h$ -dependent) symbol as  $a \in S_h(m, g_1^h)$ , where  $m$  is an  $h$ -dependent weight. The corresponding metric is naturally

$$g_1^h = h^2 dr^2 + d\rho^2 + d\theta^2 + h^2 d\omega^2.$$

**Weyl quantization.** Let  $\{\chi_\alpha^2\}$  be a partition of unity on  $\partial M$  compatible with our coordinate system  $\{\varphi_\alpha, U_\alpha\}$ , that is,  $\chi_\alpha \in C_0^\infty(U_\alpha)$  and  $\sum_\alpha \chi_\alpha(\theta)^2 \equiv 1$  on  $\partial M$ . We set  $\tilde{\chi}_\alpha(r, \theta) = \chi_\alpha(\theta)j(r) \in C^\infty(M_\infty)$ . Let  $a \in S(m, g_1)$  be a symbol on  $T^*M_\infty$ , and let  $u \in C_0^\infty(T^*M)$ . We denote by  $a_{(\alpha)}$  and  $G_{(\alpha)}$  the representations of  $a$  and  $G$  in the local coordinate  $(1 \otimes \varphi_\alpha, \mathbb{R} \times U_\alpha)$ , respectively. We quantize  $a$  by

$$\text{Op}^W(a)u = \sum_\alpha \tilde{\chi}_\alpha G_{(\alpha)}^{-1/2} a_{(\alpha)}^W(r, D_r, \theta, D_\theta) G_{(\alpha)}^{1/2} \tilde{\chi}_\alpha u,$$

where  $a_{(\alpha)}^W(r, D_r, \theta, D_\theta)$  denotes the usual Weyl quantization on the Euclidean space  $\mathbb{R}^n$ , and we use the identification  $\mathbb{R}_+ \times U_\alpha \cong \mathbb{R}_+ \times (\text{Ran } \varphi_\alpha)$  for each  $\alpha$ . (Strictly speaking, we should have written this as

$$\text{Op}^W(a)u = \sum_\alpha \tilde{\chi}_\alpha (\tilde{\varphi}_\alpha)_* (G_{(\alpha)}^{-1/2} a_{(\alpha)}^W(r, D_r, \theta, D_\theta) G_{(\alpha)}^{1/2} (\tilde{\varphi}_\alpha)_* (\tilde{\chi}_\alpha u)),$$

but we will omit  $(\tilde{\varphi}_\alpha)_*$ ,  $(\tilde{\varphi}_\alpha)^*$ ,  $\dots$ , when there can be no confusion.) This definition is compatible with the standard definition of pseudodifferential operators on manifolds, but we choose a specific quantization that preserves the asymptotically conic structure of  $M$ . Similarly, for a symbol  $a$  on  $T^*M_f$ , we quantize it by

$$\text{Op}^W(a)u = \sum_\alpha \chi_\alpha H_{(\alpha)}^{-1/2} a_{(\alpha)}^W(r, D_r, \theta, D_\theta) H_{(\alpha)}^{1/2} \chi_\alpha u$$

for  $u \in C_0^\infty(M_f)$ , where  $H_{(\alpha)}$  denotes the representation of  $H$  in the local coordinate  $(\varphi_\alpha, U_\alpha)$ . In this case, the linear structure in  $r$  is preserved.

In the above definition, we put weights around the locally defined pseudodifferential operators  $a_\alpha^W$  so that  $\text{Op}^W(a)$  is symmetric if  $a$  is real-valued. Moreover, by virtue of these weights, the symbol corresponding to the operator is unique, including the subprincipal symbol, though we will not take advantage of this fact in this paper.

The above definitions of quantizations also have the convenient property that if we identify a symbol  $a$  on  $T^*M_\infty$  with a symbol on  $T^*M_f$  (by the obvious identification), then we have

$$\mathcal{I} \text{Op}^W(a) \mathcal{I}^* = \text{Op}^W(a) \quad \text{on } \mathcal{H},$$

provided  $a$  is supported in  $\{r > 1\}$ , and we may identify these quantizations by using  $\mathcal{F}$ . For a symbol supported in  $\{r > 1\}$ , we may consider  $\text{Op}^W(a)$  as an operator from  $\mathcal{H}$  to  $\mathcal{H}_f$  (or from  $\mathcal{H}_f$  to  $\mathcal{H}$ ) also. We define these operators by

$$\text{Op}^W(a)u = \sum_{\alpha} \chi_{\alpha} H_{(\alpha)}^{-1/2} a_{(\alpha)}^W(r, D_r, \theta, D_{\theta}) G_{(\alpha)}^{1/2} \tilde{\chi}_{\alpha} u$$

for  $u \in C_0^{\infty}(M)$  and

$$\text{Op}^W(a)u = \sum_{\alpha} \tilde{\chi}_{\alpha} G_{(\alpha)}^{-1/2} a_{(\alpha)}^W(r, D_r, \theta, D_{\theta}) H_{(\alpha)}^{1/2} \chi_{\alpha} u$$

for  $u \in C_0^{\infty}(M_f)$ .

If  $A = \text{Op}^W(a)$ , we denote the (Weyl) symbol of  $A$  by  $a = \Sigma(A)$ .

**Hamiltonians.** Now we consider properties of our Schrödinger operators and related operators as a preparation for the next section.

We note that, as in the usual Weyl calculus on  $\mathbb{R}^n$ , if  $a(x, \xi) = \sum_{j,k} a_{jk}(x) \xi_j \xi_k$ , then

$$\text{Op}^W(a) = \sum_{j,k} D_j a_{jk}(x) D_k - \frac{1}{4} \sum_{j,k} (\partial_j \partial_k a_{jk}(x)).$$

Hence, if we let  $p$  be the symbol of  $P$  as in (2-1), we have

$$\text{Op}^W(p) = P + f,$$

where  $f \in C^{\infty}(M_f)$  is such that

$$|\partial_r^{\alpha} \partial_{\theta}^{\beta} f(r, \theta)| \leq C_{\alpha\beta} \langle r \rangle^{-2-|\alpha|}$$

for any  $\alpha, \beta$ . Thus, we can include this error term in  $V$  and we may consider  $P = \text{Op}^W(p)$ . On the other hand, it is easy to see  $P_f = \text{Op}^W(p_f)$  on  $\mathcal{H}_f$ , where  $p_f = \frac{1}{2} \rho^2$ .

#### 4. An Egorov-type theorem

Let  $(r_0, \rho_0, \theta_0, \omega_0) \in T^*(\mathbb{R}_+ \times \partial M)$ ,  $\omega_0 \neq 0$ , and suppose  $a \in C_0^{\infty}(T^*(\mathbb{R}_+ \times \partial M))$  is supported in a small neighborhood of  $(r_0, \rho_0, \theta_0, \omega_0)$  so that  $a$  is supported away from  $\{\omega = 0\}$ . We set

$$a^h(r, \rho, \theta, \omega) = a(h; hr, \rho, \theta, h\omega), \quad h > 0,$$

where  $a$  itself may depend on the parameter  $h > 0$ , but we suppose it is bounded uniformly in the  $C_0^{\infty}$ -topology, and supported in the same small neighborhood of  $(r_0, \rho_0, \theta_0, \omega_0)$ . The notation here is different from that of Section 2. We set

$$A_0 = \text{Op}^W(a^h) \quad \text{on } M.$$

We set  $\varepsilon > 0$  so small that

$$\exp(tH_{p_c})(\text{supp } a) \cap \{r \leq \varepsilon \langle t \rangle\} = \emptyset$$



for all  $t \in \mathbb{R}$ . We choose  $\eta \in C^\infty(\mathbb{R})$  such that  $\eta(r) = 1$  for  $r \geq 1$  and  $\eta(r) = 0$  for  $r \leq 1/2$ , and we set

$$Y = \eta\left(\frac{hr}{\varepsilon\langle t \rangle}\right).$$

Then we define

$$A(t) = e^{itP_f/h} \mathcal{F}^* Y e^{-itP/h} A_0 e^{itP/h} Y \mathcal{F} e^{-itP_f/h}$$

for  $t \in \mathbb{R}$ . The purpose of this section is to obtain the symbols of  $A(t)$  as a pseudodifferential operator, and to study its behavior as  $t \rightarrow \pm\infty$ .

We compute (formally) that

$$\frac{d}{dt}(e^{itP/h} Y \mathcal{F} e^{-itP_f/h}) = \frac{i}{h} e^{itP/h} T(t) e^{-itP_f/h},$$

where

$$T(t) = PY\mathcal{F} - Y\mathcal{F}P_f - \frac{h(hr)t}{i\varepsilon\langle t \rangle^3} \eta'\left(\frac{hr}{\varepsilon\langle t \rangle}\right) \mathcal{F}.$$

We further rewrite this as

$$\begin{aligned} \frac{d}{dt}(e^{itP/h} Y \mathcal{F} e^{-itP_f/h}) &= \frac{i}{h} (e^{itP/h} Y \mathcal{F} e^{-itP_f/h}) (e^{itP_f/h} \mathcal{F}^* T(t) e^{-itP_f/h}) + \frac{i}{h} e^{itP/h} (1 - Y\mathcal{F}\mathcal{F}^*) T(t) e^{-itP_f/h} \\ &= \frac{i}{h} (e^{itP/h} Y \mathcal{F} e^{-itP_f/h}) L(t) + R_1(t), \end{aligned}$$

where

$$L(t) = e^{itP_f/h} \mathcal{F}^* T(t) e^{-itP_f/h} \quad \text{and} \quad R_1(t) = \frac{i}{h} e^{itP/h} (1 - Y\mathcal{F}\mathcal{F}^*) T(t) e^{-itP_f/h}.$$

We now consider the symbols of  $T(t)$  and  $L(t)$  as pseudodifferential operators. By direct computations, it is easy to see that for any indices  $\alpha, \beta, \gamma, \delta$ ,

$$\begin{aligned} &|\partial_r^\alpha \partial_\rho^\beta \partial_\theta^\gamma \partial_\omega^\delta \Sigma(T(t))(r, \rho, \theta, \omega)| \\ &\leq C(\langle r \rangle^{-1-\mu} \langle \rho \rangle^2 + \langle r \rangle^{-1-\mu} \langle \rho \rangle \langle \omega \rangle + \langle r \rangle^{-2} \langle \omega \rangle^2) \langle r \rangle^{-|\alpha|} \langle \rho \rangle^{-|\beta|} \langle \omega \rangle^{-|\delta|}. \end{aligned} \quad (4-1)$$

Since  $T(t)$  is supported in  $\{r \geq \varepsilon\langle t \rangle/2h\}$ , we may replace  $\langle r \rangle$  by  $\langle r \rangle + \varepsilon\langle t \rangle/2h$  in the above estimate. We also have

$$\begin{aligned} &\left| \partial_r^\alpha \partial_\rho^\beta \partial_\theta^\gamma \partial_\omega^\delta \left( \Sigma(T(t)) - Y \frac{q(\theta, \omega)}{r^2} \right) \right| \\ &\leq C(\langle r \rangle^{-1-\mu} \langle \rho \rangle^2 + \langle r \rangle^{-1-\mu} \langle \rho \rangle \langle \omega \rangle + \langle r \rangle^{-2-\mu} \langle \omega \rangle^2) \langle r \rangle^{-|\alpha|} \langle \rho \rangle^{-|\beta|} \langle \omega \rangle^{-|\delta|}. \end{aligned}$$

In particular, we learn that

$$\left| \partial_r^\alpha \partial_\rho^\beta \partial_\theta^\gamma \partial_\omega^\delta \left( \Sigma(T(t)) - Y \frac{q(\theta, \omega)}{r^2} \right) \right| \leq C \langle t \rangle^{-1-\mu-|\alpha|} h^{\mu+|\alpha|+|\delta|} \quad (4-2)$$

on  $\exp(tH_{p_c})[\text{supp } a^h]$ , where the constant is independent of  $t$  and  $h$ .

Now we note, by virtue of the Weyl calculus (and our choice of the quantization), that

$$\Sigma(L(t))(r, \rho, \theta, \omega) = \Sigma(\mathcal{F}^* T(t))(r + (t/h)\rho, \rho, \theta, \omega).$$

Hence we have, by (4-1),

$$|\partial_r^\alpha \partial_\rho^\beta \partial_\theta^\gamma \partial_\omega^\delta \Sigma(L(t))| \leq C(\langle \tilde{r} \rangle^{-1-\mu} \langle \rho \rangle^2 + \langle \tilde{r} \rangle^{-1-\mu} \langle \rho \rangle \langle \omega \rangle + \langle \tilde{r} \rangle^{-2} \langle \omega \rangle^2) \langle \tilde{r} \rangle^{-|\alpha|} \langle \omega \rangle^{-|\delta|},$$

where  $\tilde{r} = r + (t/h)\rho$ . Note that we take advantage of the cut-off function  $Y$  in this estimate. We also note, along with (4-2), that

$$\left| \partial_r^\alpha \partial_\rho^\beta \partial_\theta^\gamma \partial_\omega^\delta \left( \Sigma(L(t)) - \frac{q(\theta, \omega)}{\tilde{r}^2} \right) \right| \leq C \langle t \rangle^{-1-\mu-|\alpha|} h^{\mu+|\alpha|+|\delta|} \tag{4-3}$$

on  $\exp(-tH_{p_f}) \circ \exp(tH_{p_c})[\text{supp } a^h] = \text{supp}(a^h \circ w_c(t))$ .

We then construct an asymptotic solution to the Heisenberg equation as

$$\frac{d}{dt} B(t) = -\frac{i}{h} [L(t), B(t)], \quad B(0) = \mathcal{F}^* A_0 \mathcal{F}. \tag{4-4}$$

**Lemma 4.1.** *There exists  $b^h(t; r, \rho, \theta, \omega) \in C_0^\infty(T^*M_f)$  satisfying the following conditions:*

- (i)  $b^h(0) = a^h$ .
- (ii)  $b^h(t)$  is supported in  $w_c(t/h)^{-1}[\text{supp } a^h]$ .
- (iii)  $b^h(t) \in S(1, g_1^h)$ , and it is bounded uniformly in  $t \in \mathbb{R}$ .
- (iv)  $b^h(t) - a^h \circ w_c(t/h) \in S(h^\mu, g_1^h)$ , i.e., the principal symbol of  $b^h(t)$  is given by  $a^h \circ w_c(t/h)$ , and the remainder is bounded uniformly in  $t$ .
- (v) If we set  $B(t) = \text{Op}^W(b^h(t))$ , then

$$\left\| \frac{d}{dt} B(t) + \frac{i}{h} [L(t), B(t)] \right\| \leq C_N \langle t \rangle^{-1-\mu} h^N, \quad h > 0,$$

for any  $N$ .

- (vi)  $B(t)$  converges to  $B_\pm$  as  $t \rightarrow \pm\infty$  in  $B(\mathcal{H}_f)$ , and the symbols  $b_\pm^h := \Sigma(B_\pm)$  satisfy

$$b_\pm^h - a^h \circ w_{c,\pm} \in S(h^\mu, g_1^h).$$

*Proof.* We follow the standard procedure to construct asymptotic solutions to Heisenberg equations (see [Taylor 1981, Chapter 8; Martinez 2002, Chapter 4]). We let

$$\ell_0(t; r, \rho, \theta, \omega) = \frac{q(\theta, \omega)}{(r + t\rho)^2}$$

be the principal symbol of  $L(ht)$ . If we set

$$b_0(t) = a \circ w_c(t) = a \circ \exp(-tH_{p_c}) \circ \exp(tH_{p_f}),$$

then  $b_0$  satisfies the equation

$$\frac{\partial}{\partial t} b_0(t) = -\{\ell_0(t), b_0(t)\}, \quad b_0(0) = a.$$

We set  $b_0^h(t; r, \rho, \theta, \omega) = b_0(t/h; hr, \rho, \theta, h\omega)$ , and we also set  $B_0(t) = \text{Op}^W(b_0^h(t))$ . We note that

$$|\partial_r^\alpha \partial_\rho^\beta \partial_\theta^\gamma \partial_\omega^\delta b_0^h(t; r, \rho, \theta, \omega)| \leq Ch^{|\alpha|+|\delta|}$$

uniformly in  $t$  with any  $\alpha, \beta, \gamma, \delta$ , since  $b_0(t)$  converges to  $a \circ w_{c,\pm}$  as  $t \rightarrow \pm\infty$ . We write

$$R_0^0(t) = \frac{d}{dt} B_0(t) + \frac{i}{h} [L(t), B_0(t)], \quad r_0^0(t) = \Sigma(R_0^0(t)).$$

Then by (4-3) and the symbol calculus,  $r_0^0(t)$  is supported on  $w_c(t/h)^{-1}[\text{supp } a^h]$  modulo  $O(h^\infty)$ -terms, and

$$\partial_r^\alpha \partial_\rho^\beta \partial_\theta^\gamma \partial_\omega^\delta r_0^0(t) \leq C \langle t \rangle^{-1-\mu-|\alpha|} h^{\mu+|\alpha|+|\delta|} \tag{4-5}$$

for any  $\alpha, \beta, \gamma, \delta$ . We set  $\tilde{r}_0^0(t)$  so that

$$\tilde{r}_0^0(t/h; hr, \rho, \theta, h\omega) = r_0^0(t; r, \rho, \theta, \omega),$$

and solve the transport equation

$$\frac{\partial}{\partial t} b_1(t) + \{\ell_0(t), b_1(t)\} = -\tilde{r}_0^0(t), \quad b_1(0) = 0.$$

By (4-5), it is easy to observe that  $|\partial_r^\alpha \partial_\rho^\beta \partial_\theta^\gamma \partial_\omega^\delta b_1(t; r, \rho, \theta, \omega)| \leq Ch^\mu$  uniformly in  $t$ . Moreover,  $b_1(t)$  converges to a symbol supported in  $w_{c,\pm}^{-1}[\text{supp } a]$  in the  $C_0^\infty$ -topology as  $t \rightarrow \pm\infty$ . We then set

$$B_1(t) = \text{Op}^W(b_1^h(t)), \quad b_1^h(t; r, \rho, \theta, \omega) = b_1(t/h; hr, \rho, \theta, h\omega).$$

We construct  $b_j, j = 1, 2, \dots$ , iteratively, so that  $b_j^h \in S(h^{j\mu}, g_1^h)$ , and set

$$b^h(t) \sim \sum_{j=0}^\infty b_j^h(t), \quad B(t) = \text{Op}^W(b^h(t)).$$

By construction,  $b^h(t)$  and  $B(t) = \text{Op}^W(b^h(t))$  satisfy the assertion. □

We then observe that  $A(t)$  is very close to  $B(t)$  constructed as above.

**Lemma 4.2.** *For any  $N$ , there is  $C_N > 0$  such that*

$$\|A(t) - B(t)\| \leq C_N h^N, \quad t \in \mathbb{R}.$$

*In particular,  $A_+$  and  $A_-$ , defined by*

$$A_\pm := \text{w-lim}_{t \rightarrow \pm\infty} A(t),$$

*have the symbols  $b_\pm^h$  as pseudodifferential operators.*

*Proof.* We first observe that

$$\begin{aligned} \|A(t) - B(t)\| &= \|e^{itP_f/h} \mathcal{G}^* Y e^{-itP/h} A_0 e^{itP/h} Y \mathcal{G} e^{-itP_f/h} - B(t)\| \\ &= \|\mathcal{G}^* Y e^{-itP/h} A_0 e^{itP/h} Y \mathcal{G} - e^{-itP_f/h} B(t) e^{itP_f/h}\| \\ &\leq \|Y \mathcal{G} \mathcal{G}^* Y e^{-itP/h} A_0 e^{itP/h} Y \mathcal{G} \mathcal{G}^* Y - Y \mathcal{G} e^{-itP_f/h} B(t) e^{itP_f/h} \mathcal{G}^* Y\| \\ &\leq \|e^{-itP/h} A_0 e^{itP/h} - Y \mathcal{G} e^{-itP_f/h} B(t) e^{itP_f/h} \mathcal{G}^* Y\| + R_2 \\ &= \|A_0 - \tilde{B}(t)\| + R_2, \end{aligned}$$

where

$$R_2 = 2\|(1 - Y\mathcal{G}\mathcal{G}^*Y)e^{-itP/h}A_0\| \quad \text{and} \quad \tilde{B}(t) = e^{itP/h}Y\mathcal{G}e^{-itP_f/h}B(t)e^{itP_f/h}\mathcal{G}^*Ye^{-itP/h}.$$

By [Corollary A.2](#), we learn that  $R_2 = O(\langle t \rangle^{-N}h^N)$  for any  $N$ . We then show  $\tilde{B}(t)$  is very close to  $A_0$  uniformly in  $t$ . We compute

$$\begin{aligned} \frac{d}{dt}\tilde{B}(t) &= (e^{itP/h}Y\mathcal{G}e^{-itP_f/h})\frac{d}{dt}B(t)(e^{itP_f/h}\mathcal{G}^*Ye^{-itP/h}) \\ &\quad + \frac{i}{h}(e^{itP/h}Y\mathcal{G}e^{-itP_f/h})L(t)B(t)(e^{itP_f/h}\mathcal{G}^*Ye^{-itP/h}) \\ &\quad - \frac{i}{h}(e^{itP/h}Y\mathcal{G}e^{-itP_f/h})B(t)L(t)^*(e^{itP_f/h}\mathcal{G}^*Ye^{-itP/h}) \\ &\quad + R_1(t)B(t)(e^{itP_f/h}\mathcal{G}^*Ye^{-itP/h}) - (e^{itP/h}Y\mathcal{G}e^{-itP_f/h})B(t)R_1(t)^* \\ &= (e^{itP/h}Y\mathcal{G}e^{-itP_f/h})\left(\frac{d}{dt}B(t) + \frac{i}{h}[L(t), B(t)]\right)(e^{itP_f/h}\mathcal{G}^*Ye^{-itP/h}) + R_3(t), \end{aligned}$$

where

$$\begin{aligned} R_3(t) &= R_1(t)B(t)(e^{itP_f/h}\mathcal{G}^*Ye^{-itP/h}) - (e^{itP/h}Y\mathcal{G}e^{-itP_f/h})B(t)R_1(t)^* \\ &\quad + \frac{i}{h}(e^{itP/h}Y\mathcal{G}e^{-itP_f/h})B(t)(L(t) - L(t)^*)(e^{itP_f/h}\mathcal{G}^*Ye^{-itP/h}). \end{aligned}$$

We can show that  $\|R_3(t)\| = O(\langle t \rangle^{-N}h^N)$  for any  $N$ . For example,

$$\begin{aligned} \|R_1(t)B(t)(e^{itP_f/h}\mathcal{G}^*Ye^{-itP/h})\| &\leq h^{-1}\|(1 - Y\mathcal{G}\mathcal{G}^*)T(t)e^{-itP_f/h}B(t)\| \\ &= h^{-1}\|e^{itP_f/h}(1 - Y\mathcal{G}\mathcal{G}^*)T(t)e^{-itP_f/h}B(t)\|. \end{aligned}$$

As we have seen already,  $e^{itP_f/h}(1 - Y\mathcal{G}\mathcal{G}^*)T(t)e^{-itP_f/h}$  is a pseudodifferential operator, and its support is separated from the support of  $b^h(t)$  by a distance not less than  $c\langle t \rangle h^{-1}$ , for some  $c > 0$ . Thus their product has a vanishing symbol, and its norm is  $O(\langle t \rangle^{-N}h^N)$  with any  $N$ . The other terms are estimated similarly. Combining this with [Lemma 4.1\(v\)](#), we learn that

$$\left\| \frac{d}{dt}\tilde{B}(t) \right\| \leq C_N\langle t \rangle^{-1-\mu}h^N$$

for any  $N$ , and hence  $\|\tilde{B}(t) - \tilde{B}(0)\| \leq C_Nh^N$ . We note that

$$\tilde{B}(0) = \eta\left(\frac{hr}{\varepsilon}\right)\mathcal{G}\mathcal{G}^*A_0\mathcal{G}\mathcal{G}^*\eta\left(\frac{hr}{\varepsilon}\right) = A_0 + O(h^N)$$

by the choice of  $\varepsilon > 0$ . Combining these facts, we conclude the assertion holds. □

### 5. Proofs of Theorems 1.1 and 1.2

Let  $(r_0, \rho_0, \theta_0, \omega_0) \in T^*(\mathbb{R}_+ \times \partial M)$ , and suppose  $\omega_0 \neq 0$  as in the last section. Also we let  $a$  in  $C_0^\infty(T^*(\mathbb{R}_+ \times \partial M))$  be supported in a small neighborhood of  $(r_0, \rho_0, \theta_0, \omega_0)$  and we set

$$A_0 = \text{Op}^W(a^h), \quad a^h(r, \rho, \theta, \omega) = a(hr, \rho, \theta, h\omega).$$

Let  $\varepsilon > 0$  also as in the last section. Write

$$(r_{\pm}, \rho_{\pm}, \theta_{\pm}, \omega_{\pm}) = w_{c,\pm}^{-1}(r_0, \rho_0, \theta_0, \omega_0)$$

as in Section 2, and recall that  $w_{c,\pm}$  are diffeomorphisms from  $\mathbb{R} \times \mathbb{R}_{\pm} \times (T^*\partial M \setminus 0)$  to  $\mathbb{R}_+ \times \mathbb{R} \times (T^*\partial M \setminus 0)$ .

We also note that

$$E_0 = p_c(r_0, \rho_0, \theta_0, \omega_0) = \frac{1}{2}\rho_{\pm}^2 > 0$$

by conservation of energy.

**Lemma 5.1.** *If  $\delta > 2\varepsilon^2$ , then*

$$\text{w-lim}_{t \rightarrow \pm\infty} \eta(P_f/\delta)A(t)\eta(P_f/\delta) = \eta(P_f/\delta)W_{\pm}^*A_0W_{\pm}\eta(P_f/\delta).$$

*Proof.* It is easy to show by the stationary phase method that

$$\text{s-lim}_{t \rightarrow \pm\infty} \left(1 - \eta\left(\frac{hr}{\varepsilon\langle t \rangle}\right)\right) \mathcal{G}e^{-itP_f/h}\eta(P_f/\delta) = 0$$

(for fixed  $h$ ), since the stationary points (in  $\rho$ ) satisfy  $hr = t\rho$ . This implies that

$$\text{s-lim}_{t \rightarrow \pm\infty} e^{itP/h}Y\mathcal{G}e^{-itP_f/h}\eta(P_f/\delta) = W_{\pm}\eta(P_f/\delta),$$

and the claim follows immediately. □

This implies, combined with Lemmas 4.1 and 4.2:

**Lemma 5.2.** *Let  $A_0$  as above. Then  $W_{\pm}^*A_0W_{\pm}$  are pseudodifferential operators with the symbols  $b_{\pm}^h$  given in Lemma 4.1. In particular,  $\Sigma(W_{\pm}^*A_0W_{\pm})$  are supported in  $w_{c,\pm}^{-1}[\text{supp } a^h]$  modulo  $O(h^\infty)$ -terms, and the principal symbols (modulo  $S(h^\mu, g_1^h)$ ) are given by  $a^h \circ w_{c,\pm}$ .*

For the moment, we set

$$\rho_0 = 0 \quad \text{and hence} \quad r_{\pm} = 0.$$

Then we may take  $\varepsilon = \sqrt{E_0}$  provided  $a$  is supported in a sufficiently small neighborhood of  $(r_0, 0, \theta_0, \omega_0)$ .

Now let us suppose  $(0, \rho_-, \theta_-, \omega_-)$  (with  $\omega_- \neq 0, \rho_- > 2\varepsilon$ ) is given, and  $(0, \rho_0, \theta_0, \omega_0)$  is defined by  $w_{c,-}(0, \rho_-, \theta_-, \omega_-) = (0, \rho_0, \theta_0, \omega_0)$ . The converse of Lemma 5.2 is given as follows:

**Lemma 5.3.** *Let  $\tilde{a} \in C_0^\infty(\mathbb{R} \times \mathbb{R}_- \times (T^*\partial M \setminus 0))$  be supported in a small neighborhood of  $(0, \rho_-, \theta_-, \omega_-)$ , and let*

$$\tilde{A} = \text{Op}^W(\tilde{a}^h), \quad \tilde{a}^h(r, \rho, \theta, \omega) = \tilde{a}(hr, \rho, \theta, h\omega).$$

*Then there is a symbol  $a_0^h$  supported in  $w_{c,-}[\text{supp } \tilde{a}^h]$  such that for any  $f \in C_0^\infty(\mathbb{R}_+)$ ,*

$$f(P)A_0f(P) = W_-f(P_f)\tilde{A}f(P_f)W_-^*,$$

*where  $A_0 = \text{Op}^W(a_0^h)$ . Moreover, the principal symbol (modulo  $S(h^\mu, g_1^h)$ ) is  $\tilde{a}^h \circ w_{c,-}^{-1}$ .*



*Proof.* We set  $a_{0,0}^h = \tilde{a}^h \circ w_{c,-}^{-1}$ . Then by [Lemma 5.2](#), we have

$$a_{-,1}^h := \Sigma(\tilde{A} - W_-^* \text{Op}^W(a_{0,0}^h) W_-) \in S(h^\mu, g_1^h),$$

and it is supported in  $\text{supp}[\tilde{a}^h]$  modulo  $O(h^\infty)$ -terms. Then we set  $a_{0,1} = a_{-,1}^h \circ w_{c,-}^{-1}$ , and set

$$a_{-,2}^h := \Sigma(\tilde{A} - W_-^* \text{Op}^W(a_{0,0}^h + a_{0,1}^h) W_-) \in S(h^{2\mu}, g_1^h).$$

We construct  $a_{-,j}^h$ ,  $j = 2, 3, \dots$ , iteratively by

$$a_{-,j}^h := \Sigma(\tilde{A} - W_-^* \text{Op}^W(a_{0,0}^h + \dots + a_{0,j-1}^h) W_-) \in S(h^{j\mu}, g_1^h),$$

$a_{0,j}^h = a_{-,j}^h \circ w_{c,-}^{-1}$ , and we set  $a_0^h \sim \sum_{j=0}^\infty a_{0,j}^h$  as an asymptotic sum. Then we have

$$\tilde{A} = W_-^* A_0 W_-$$

modulo  $S(h^\infty \langle r \rangle^{-\infty} \langle \omega \rangle^{-\infty}, g_1)$ -terms. Since there are no positive eigenvalues [\[Ito and Skibsted 2011; Melrose and Zworski 1996\]](#), we also have  $W_\pm f(P_f) W_\pm^* = f(P)$  by virtue of the intertwining property and asymptotic completeness [\[Ito and Nakamura 2010\]](#). These imply

$$W_- f(P_f) \tilde{A} f(P_f) W_-^* = W_- f(P_f) W_-^* A_0 W_- f(P_f) W_-^* = f(P) A_0 f(P),$$

and this implies the assertion. □

We note [Lemma 5.3](#) naturally holds for  $w_{c,+}$  instead of  $w_{c,-}$ , but we only use the above case. By [Lemma 5.3](#), we learn that

$$Sf(P_f) \tilde{A} f(P_f) S^* = W_+^* f(P) A_0 f(P) W_+ = f(P_f) (W_+^* A_0 W_+) f(P_f).$$

By [Lemma 5.2](#),  $W_+^* A_0 W_+$  is a pseudodifferential operator. By choosing  $f \in C_0^\infty(\mathbb{R}_+)$  so that  $f(\rho^2/2) = 1$  in a neighborhood of the support of  $\tilde{a}$ , we may omit  $f(P_f)$  factors up to negligible terms. Thus,  $S\tilde{A}S^*$  is a pseudodifferential operator with a symbol supported in  $s_c[\text{supp } \tilde{a}^h]$ , and the principal symbol is given by  $\tilde{a}^h \circ s_c^{-1}$ , where  $\tilde{a}$  is the symbol given in [Lemma 5.3](#), i.e.,  $\tilde{a}$  is supported in a small neighborhood of  $(0, \rho_-, \theta_-, \omega_-)$ .

We note that, by the intertwining property of the scattering operator,

$$e^{-itP_f} S = S e^{-itP_f}, \quad \forall t \in \mathbb{R}.$$

This in turn implies

$$T_\tau S = S T_\tau, \quad \forall \tau \in \mathbb{R}, \quad \text{where } T_\tau = \exp(-i\tau\sqrt{2P_f}).$$

On the other hand,  $\sqrt{2P_f} = \mp i \frac{\partial}{\partial r}$  on  $\mathcal{H}_{f,\pm}$ , and hence  $T_\tau$  are translations with respect to  $r$ . More precisely, we have

$$T_\tau u_\pm(r, \theta) = u_\pm(r \mp \tau, \theta) \quad \text{for } u_\pm \in \mathcal{H}_{f,\pm}.$$

We learn from these facts that

$$S(T_\tau \tilde{A} T_\tau^*) S^* = T_\tau (S \tilde{A} S^*) T_\tau^*,$$

and the symbols of  $T_\tau \tilde{A} T_\tau^*$  and  $T_\tau (S\tilde{A}S^*) T_\tau^*$  are given by  $\tilde{a}^h(r + \tau, \rho, \theta, \omega)$  and  $\Sigma(S\tilde{A}S^*)(r + \tau, \rho, \theta, \omega)$ , respectively. Using this observation, we may replace  $\tilde{a}$  by a symbol supported in a small neighborhood  $(r_-, \rho_-, \theta_-, \omega_-)$  with arbitrary  $r_- \in \mathbb{R}$ . Thus we have proved:

**Lemma 5.4.** *Let  $a \in C_0^\infty(\mathbb{R} \times \mathbb{R}_- \times (T^* \partial M \setminus 0))$  be supported in a small neighborhood of  $(r_-, \rho_-, \theta_-, \omega_-)$  with  $|\rho_-| \geq 2\varepsilon$ , and let*

$$\tilde{A} = \text{Op}^W(a^h), \quad a^h(r, \rho, \theta, \omega) = a(hr, \rho, \theta, h\omega).$$

*Then  $S\tilde{A}S^*$  is a pseudodifferential operator with a symbol supported in  $s_c[\text{supp } a^h]$  modulo  $O(h^\infty)$ -terms, and the principal symbol (modulo  $S(h^\mu, g_1^h)$ ) is given by  $a^h \circ s_c^{-1}$ .*

Here we have used the formula

$$s_c(r, \rho, \theta, \omega) = (-r, -\rho, \exp(\pi H_{\sqrt{2q}})(\theta, \omega)).$$

We set  $\hat{\mathcal{H}}_{f,\pm} = \mathcal{F}\mathcal{H}_{f,\pm}$ . Then  $\mathcal{F}S\mathcal{F}^{-1}$  is a unitary map from  $\hat{\mathcal{H}}_{f,-}$  to  $\hat{\mathcal{H}}_{f,+}$ . For notational simplicity, we set

$$\Pi u(r, \theta) = u(-r, \theta) \quad \text{for } u \in \mathcal{H}_{f,\pm},$$

so that  $\mathcal{F}(S\Pi)\mathcal{F}^{-1}$  is a unitary map on  $\hat{\mathcal{H}}_{f,+}$ . By the intertwining property above,  $\mathcal{F}(S\Pi)\mathcal{F}^{-1}$  commutes with functions of  $\rho$ , and hence is decomposed so that

$$\mathcal{F}(S\Pi)\mathcal{F}^{-1}u(\rho, \omega) = (S(\rho^2/2)u(\rho, \cdot))(\omega) \quad \text{on } \hat{\mathcal{H}}_{f,+} \cong L^2(\mathbb{R}_+; L^2(\partial M)),$$

where  $S(\lambda) \in B(L^2(\partial M))$  is the scattering matrix.

*Proof of Theorem 1.1.* We recall the semiclassical-type characterization of the wave front set: Let  $g(\rho, \theta) \in \mathcal{D}'(\mathbb{R}_+ \times \partial M)$ , and let  $(\rho_0, \theta_0, r_0, \omega_0) \in T^*(\mathbb{R}_+ \times \partial M)$ .  $(\rho_0, \theta_0, r_0, \omega_0) \notin WF(g)$  if and only if there is  $a \in C_0^\infty(T^*(\mathbb{R}_+ \times \partial M))$  such that  $a(\rho_0, \theta_0, r_0, \omega_0) \neq 0$  and

$$\|a(\rho, \theta, hD_\rho, hD_\theta)g\| = O(h^\infty) \quad \text{as } h \rightarrow +0.$$

We may replace  $a$  by an  $h$ -dependent symbol with a principal symbol which does not vanish at  $(\rho_0, \theta_0, r_0, \omega_0)$ .

We fix  $\lambda_0 = \rho_0^2/2$  with  $\rho_0 > 2\varepsilon$  and consider  $S(\lambda)$  where  $\lambda$  is in a small neighborhood of  $\lambda_0$ . Let  $u \in L^2(\partial M)$  and let  $v \in C_0^\infty(\mathbb{R}_+)$  be supported in a small neighborhood of  $\lambda_0$ . Then it is easy to see that

$$WF(v(\rho)u(\theta)) = \{(\rho, \theta, 0, \omega) \mid \rho \in \text{supp } v, (\theta, \omega) \in WF(u)\}.$$

Then, by Lemma 5.4 and the above characterization of the wave front set, we learn that

$$\begin{aligned} WF(\mathcal{F}(S\Pi)\mathcal{F}^{-1}v(\rho)u(\theta)) &= (1 \otimes \exp(\pi H_{\sqrt{2q}}))WF(v(\rho)u(\theta)) \\ &= \{(\rho, \theta, 0, \omega) \mid \rho \in \text{supp } v, (\theta, \omega) \in \exp(\pi H_{\sqrt{2q}})WF(u)\}; \end{aligned}$$

see [Nakamura 2009b]. By the definition of the scattering matrix, this implies

$$WF(S(\lambda)u) \subset \exp(\pi H_{\sqrt{2q}})WF(u)$$

for  $\lambda \in \text{supp } v$ . Since this argument works for  $S^{-1}$  also, the above inclusion is actually an equality, and we conclude [Theorem 1.1](#). □

*Proof of [Theorem 1.2](#).* Here we suppose  $\mu = 1$ . Then by [Lemma 5.4](#) and the Beals-type characterization of FIOs ([Theorem B.1](#)),  $\mathcal{F}(S\Pi)\mathcal{F}^{-1}$  is an FIO associated to  $1 \otimes \exp(\pi H_{\sqrt{2q}})$  on  $\{(\rho, \theta, r, \omega) \mid \omega \neq 0\}$ . Since  $\mathcal{F}(S\Pi)\mathcal{F}^{-1}$  is decomposed as  $\{S(\lambda)\}$ , this implies  $S(\lambda)$  are FIOs on  $\partial M$  associated to the canonical transform  $\exp(\pi H_{\sqrt{2q}})$  (see [Proposition B.4](#)). □

### 6. Proof of [Theorem 1.3](#)

Here we discuss how to generalize the proof of [Theorem 1.2](#) to conclude [Theorem 1.3](#).

We first modify the Egorov-type argument in [Section 4](#). Let  $(r_0, \rho_0, \theta_0, \omega_0) \in T^*M_\infty$ ,  $\omega_0 \neq 0$ , and let  $\Omega_0$  be a small neighborhood of  $(r_0, \rho_0, \theta_0, \omega_0)$ . We suppose  $a \in C_0^\infty(T^*M_\infty)$  is supported in  $\Omega_0$ , and we consider the behavior of  $A(t)$  as in [Section 4](#). We set

$$w^*(t) = \exp(-itH_{p_f}) \circ \exp(tH_p),$$

which is well-defined for  $X \in T^*M_\infty$  as long as  $\exp(tH_p)(X) \in T^*M_\infty$ . By the discussion in the proof of [Theorem 2.2](#), this condition is always satisfied if  $X = (r, \rho, \theta, \omega) \in \Omega_0^h$  and  $h$  is sufficiently small. We set

$$w(t) = w^*(t)^{-1} = \exp(-tH_p) \circ \exp(itH_{p_f})$$

on the range of  $w(t)$ . We note that

$$w_\pm^* = \lim_{t \rightarrow \pm\infty} w^*(t)$$

on  $\Omega_0^h$  with sufficiently small  $h$ , and that

$$w_\pm = \lim_{t \rightarrow \pm\infty} w(t)$$

on  $w_\pm^{-1}[\Omega_0^h]$  with sufficiently small  $h$ . Convergence of these maps holds in the  $C^\infty$ -topology.

We replace [Lemma 4.1](#) by the following slightly different statement:

**Lemma 6.1.** *There exists  $b^h(t; r, \rho, \theta, \omega) \in C_0^\infty(T^*M_f)$  satisfying the following conditions:*

- (i)  $b^h(0) = a^h$ .
- (ii)  $b^h(t)$  is supported in  $w^*(t)[\text{supp } a^h]$ .
- (iii)  $b^h(t) \in S(1, g_1^h)$ , and it is bounded uniformly in  $t \in \mathbb{R}$ .
- (iv)  $b^h(t) - a^h \circ w(t) \in S(h, g_1^h)$ , i.e., the principal symbol of  $b^h(t)$  is given by  $a^h \circ w(t)$ , and the remainder is bounded uniformly in  $t$ .
- (v) If we set  $B(t) = \text{Op}^W(b^h(t))$ , then

$$\left\| \frac{d}{dt} B(t) + \frac{i}{h} [L(t), B(t)] \right\| \leq C_N \langle t \rangle^{-1-\mu} h^N, \quad h > 0,$$

for any  $N$ .

(vi)  $B(t)$  converges to  $B_{\pm}$  as  $t \rightarrow \pm\infty$  in  $B(\mathcal{H}_f)$ , and the symbols  $b_{\pm}^h := \Sigma(B_{\pm})$  satisfy

$$b_{\pm}^h - a^h \circ w_{\pm} \in S(h, g_1^h).$$

We note that  $w(t)$  is not homogeneous in the  $(r, \omega)$ -variables, but very close to a homogeneous map when  $|(r, \omega)|$  is very large thanks to [Theorem 2.2](#).

In order to prove [Lemma 6.1](#), we set

$$b_0^h(t) = a^h \circ w(t) = a \circ \exp(-tH_p) \circ \exp(tH_{p_f}),$$

which is supported in  $w^*(t)[\Omega_0^h]$ . We have  $b_0^h(t) \in S(1, g_1^h)$  uniformly in  $t$  (for small  $h$ ) again by [Theorem 2.2](#). Moreover,  $b_0^h$  satisfies

$$\frac{\partial}{\partial t} b_0^h(t) = -h^{-1} \{\ell(t), b_0^h(t)\},$$

where  $\ell(t) = \Sigma(L(t))$ . Hence the first remainder term  $r_0^0(t)$  (as defined in [Section 4](#)) satisfies

$$|\partial_r^\alpha \partial_\rho^\beta \partial_\theta^\gamma \partial_\omega^\delta r_0^0(t)| \leq C \langle t \rangle^{-1-\mu-|\alpha|} h^{1+|\alpha|+|\delta|}$$

for any indices  $\alpha, \beta, \gamma, \delta$ . Then we construct the asymptotic solution as in the proof of [Lemma 4.1](#) by solving transport equations

$$\frac{\partial}{\partial t} b_j^h(t) + h^{-1} \{\ell(t), b_j^h(t)\} = -r_j^h(t), \quad j = 0, 1, 2, \dots,$$

and we conclude [Lemma 6.1](#). □

[Lemma 4.2](#) holds when the construction of  $B(t)$  is replaced by the one above, with no modifications. [Lemmas 5.2](#) and [5.3](#) hold in the following form. The proofs are the same.

**Lemma 6.2.** *Let  $A_0$  as above. Then  $W_{\pm}^* A_0 W_{\pm}$  are pseudodifferential operators with the symbols  $b_{\pm}^h$  given in [Lemma 6.1](#). In particular,  $\Sigma(W_{\pm}^* A_0 W_{\pm})$  are supported in  $w_{\pm}^{-1}[\text{supp } a^h]$  modulo  $O(h^\infty)$ -terms, and the principal symbols (modulo  $S(h, g_1^h)$ ) are given by  $a^h \circ w_{\pm}$ .*

**Lemma 6.3.** *Let  $\tilde{a} \in C_0^\infty(\mathbb{R} \times \mathbb{R}_- \times (T^* \partial M \setminus 0))$  be supported in a small neighborhood of  $(0, \rho_-, \theta_-, \omega_-)$ , and let*

$$\tilde{A} = \text{Op}^W(\tilde{a}^h), \quad \tilde{a}^h(r, \rho, \theta, \omega) = \tilde{a}(hr, \rho, \theta, h\omega).$$

*Then  $W_- \tilde{A} W_-^*$  is a pseudodifferential operator with a symbol supported in  $w_-[\text{supp } \tilde{a}^h]$ , and the principal symbol (modulo  $S(h, g_1^h)$ ) is given by  $\tilde{a}^h \circ w_-^*$ .*

Combining these, we learn (as in [Section 5](#)) the following assertion.

**Lemma 6.4.** *Let  $a \in C_0^\infty(\mathbb{R} \times \mathbb{R}_- \times (T^* \partial M \setminus 0))$  be supported in a small neighborhood of  $(r_-, \rho_-, \theta_-, \omega_-)$  with  $|\rho_-| \geq 2\varepsilon$ ,  $\omega_- \neq 0$ , and let*

$$\tilde{A} = \text{Op}^W(a^h), \quad a^h(r, \rho, \theta, \omega) = a(hr, \rho, \theta, h\omega).$$

*Then  $S \tilde{A} S^*$  is a pseudodifferential operator with a symbol supported in  $s[\text{supp } a^h]$  modulo  $O(h^\infty)$ -terms, and the principal symbol (modulo  $S(h, g_1^h)$ ) is given by  $a^h \circ s^{-1}$ .*

In the following, we consider  $(r, \rho, \theta, \omega) \in \Omega_0^h$  with some  $\Omega_0$  and sufficiently small  $h$ , or equivalently, when  $|\omega|$  is sufficiently large. By conservation of energy (or equivalently, by invariance under a shift in  $r$ ), the classical scattering operator has the form

$$s(r, \rho, \theta, \omega) = (-r + g(\rho, \theta, \omega), -\rho, s(\lambda)(\theta, \omega)), \tag{6-1}$$

where  $\lambda = \rho^2/2$  and  $s(\lambda)$  is a canonical transform on  $T^*\partial M$  for each  $\lambda > 0$ . (We note that without  $g(\rho, \theta, \omega)$ , the map  $s$  is not necessarily canonical.) Moreover, by [Theorem 2.1](#), we have for any indices  $\alpha, \beta, \gamma$  that

$$\begin{aligned} |\partial_\rho^\alpha \partial_\theta^\beta \partial_\omega^\gamma g(\rho, \theta, \omega)| &\leq Ch^{-1+\mu+|\gamma|}, \\ |\partial_\rho^\alpha \partial_\theta^\beta \partial_\omega^\gamma s_1(\rho, \theta, \omega)| &\leq Ch^{\mu+|\gamma|}, \\ |\partial_\rho^\alpha \partial_\theta^\beta \partial_\omega^\gamma s_2(\rho, \theta, \omega)| &\leq Ch^{-1+\mu+|\gamma|}, \end{aligned}$$

on  $\Omega_0^h$ , where  $\Omega_0$  is a small neighborhood of  $(0, \rho_-, \theta_-, \omega_-)$ , and  $s_1, s_2$  are defined by

$$(s_1(\rho, \theta, \omega), s_2(\rho, \theta, \omega)) = s(\lambda)(\theta, \omega) - \exp(\pi H_{\sqrt{2q}})(\theta, \omega),$$

i.e.,  $s_1$  denotes the  $\theta$ -components of the right-hand side terms, and  $s_2$  denotes the  $\omega$ -components. These estimates imply that  $s$  is asymptotically homogeneous (in  $(r, \omega)$ -variables) in the sense of [\[Ito and Nakamura 2012, Section 4\]](#).

In general, an operator  $U$  with distribution kernel  $u$  is called an FIO of order  $m$  associated to an asymptotically homogeneous canonical transform  $S$  if  $u$  is a Lagrangian distribution associated to

$$\Sigma_S := \{(x, y, \xi, -\eta) \mid (x, \xi) = S(y, \eta)\},$$

that is, for any  $a_1, \dots, a_N \in S_{cl}^1$  such that  $a_j$  vanishes on  $\Sigma_S$  for each  $j$ , we have that  $\text{Op}(a_1) \cdots \text{Op}(a_N)u$  is in  $B_{2,\infty}^{-m-n/2,\infty}(\mathbb{R}^{2n})$  [\[Ito and Nakamura 2012\]](#). The Beals-type characterization of FIOs discussed in [Appendix B](#) holds for such FIOs without any change.

By [Lemma 6.4](#) and the analogue of [Corollary B.2](#), we learn that  $S$  is an FIO associated to the classical scattering map  $s$ . Moreover, by [Proposition B.4](#), we learn that the scattering matrix  $S(\lambda)$  is an FIO associated to  $s(\lambda)$ , where  $s(\lambda)$  is defined by (6-1) and it is asymptotic to  $\exp(\pi H_{\sqrt{2q}})$ . Thus we have proved the following slightly more precise version of [Theorem 1.3](#):

**Theorem 6.5.** *Suppose [Assumption A](#) holds. Then for each  $\lambda > 0$ ,  $S(\lambda)$  is an FIO associated to  $s(\lambda)$  defined by (6-1). The canonical map  $s(\lambda)$  on  $T^*\partial M$  is asymptotically homogeneous in  $\omega$ , asymptotic to  $\exp(\pi H_{\sqrt{2q}})$  with the error of  $O(|\omega|^{1-\mu})$ .*

### Appendix A: Local decay estimates

Let  $P$  be as in [Section 1](#). For a symbol  $a$ , we set  $a^h(r, \rho, \theta, \omega) = a(hr, \rho, \theta, h\omega)$ . Then we have the following:

**Theorem A.1.** *Let  $(r_0, \rho_0, \theta_0, \omega_0) \in T^*M_\infty \cong T^*\mathbb{R}_+ \times T^*\partial M$ , and suppose  $\omega_0 \neq 0$ . We denote the  $\varepsilon$ -neighborhood of  $(r_0, \rho_0, \theta_0, \omega_0)$  by  $\Omega_\varepsilon$ . We suppose  $\varepsilon > 0$  so small that  $\Omega_{2\varepsilon} \Subset T^*\mathbb{R}_+ \times (T^*\partial M \setminus 0)$ .*



If  $a \in C_0^\infty(T^*M_\infty)$  is real-valued, and supported in  $\Omega_\varepsilon$ , then there is an  $h$ -dependent symbol  $b(t)$  in  $C_0^\infty(T^*M_\infty)$  for any  $t \in \mathbb{R}$  such that:

- (i)  $|a(r, \rho, \theta, \omega)| \leq c_1 b(0; r, \rho, \theta, \omega)$  with some  $c_1 > 0$ .
- (ii)  $b(t)$  is supported in  $\Omega(t) := \exp(tH_{p_c})[\Omega_{2\varepsilon}]$  for  $t \in \mathbb{R}$ .
- (iii) For any indices  $\alpha, \beta, \gamma$  and  $\delta$ , there is  $C_{\alpha\beta\gamma\delta} > 0$  such that

$$|\partial_r^\alpha \partial_\rho^\beta \partial_\theta^\gamma \partial_\omega^\delta b(t, r, \rho, \theta, \omega)| \leq C_{\alpha\beta\gamma\delta}, \quad (r, \rho, \theta, \omega) \in T^*M_\infty, t \in \mathbb{R}.$$

- (iv) There is  $R(t) \in B(L^2(M))$  such that  $\|R(t)\| \leq C_N h^N$  for any  $N$ , and

$$e^{-itP/h} \text{Op}^W(a^h) e^{itP/h} \leq c_1 \text{Op}^W(b^h(t)) + R(t)$$

for  $t > 0$ , and the reverse inequality for  $t < 0$ . Moreover,  $R(t)$  satisfies

$$\|K^N R(t) K^N\|_{B(L^2)} \leq C_N h^N, \quad t \in \mathbb{R},$$

for any  $N$ , where  $K(\cdot) = \langle \text{dist}(\cdot, \Omega^h(t)) \rangle$  with

$$\text{supp}[b^h(t)] \subset \Omega^h(t) := \{(r, \rho, \theta, \omega) \mid (hr, \rho, \theta, h\omega) \in \Omega(t)\}.$$

Before proving [Theorem A.1](#), we present a corollary which is needed in [Section 4](#).

**Corollary A.2.** Let  $\bar{\eta} \in C^\infty(\mathbb{R})$  be such that  $\bar{\eta}(r) = 0$  if  $r > 2$ , and  $\bar{\eta}(r) = 1$  if  $r \leq 1$ . We choose  $\varepsilon_1 > 0$  so small that

$$\text{dist}(\{(r, \rho, \theta, \omega) \mid |r| \leq \varepsilon_1 \langle t \rangle\}, \Omega(t)) \geq \delta \langle t \rangle$$

with some  $\delta > 0$ . Then for any  $N$  there is  $C_N > 0$  such that

$$\left\| \bar{\eta} \left( \frac{hr}{\varepsilon_1 \langle t \rangle} \right) e^{-itP/h} \text{Op}(a^h) \right\| \leq C_N h^N \langle t \rangle^{-N}, \quad t \in \mathbb{R}.$$

We note that if  $\varepsilon > 0$  is chosen sufficiently small, then we can find  $\varepsilon_1 > 0$  satisfying the property above.

*Proof of Corollary A.2.* We apply [Theorem A.1](#) with  $\tilde{a}$  such that  $\text{Op}^W(\tilde{a}) = \text{Op}^W(a) \text{Op}^W(a)^*$ , which satisfies the same condition. Then we have

$$\begin{aligned} \left\| \bar{\eta} \left( \frac{hr}{\varepsilon_1 \langle t \rangle} \right) e^{-itP/h} \text{Op}(a^h) \right\|^2 &= \bar{\eta} \left( \frac{hr}{\varepsilon_1 \langle t \rangle} \right) e^{-itP/h} \text{Op}(\tilde{a}^h) e^{itP/h} \bar{\eta} \left( \frac{hr}{\varepsilon_1 \langle t \rangle} \right) \\ &\leq c_1 \bar{\eta} \left( \frac{hr}{\varepsilon_1 \langle t \rangle} \right) \text{Op}(b^h(t)) \bar{\eta} \left( \frac{hr}{\varepsilon_1 \langle t \rangle} \right) + \bar{\eta} \left( \frac{hr}{\varepsilon_1 \langle t \rangle} \right) R(t) \bar{\eta} \left( \frac{hr}{\varepsilon_1 \langle t \rangle} \right) \\ &\leq C_N h^N \langle t \rangle^{-N}, \end{aligned}$$

where we used the fact that  $\text{supp}[b^h(t)]$  is separated from  $\Omega^h(t)$  by a distance not less than  $\delta \langle t/h \rangle$ .  $\square$

*Proof of Theorem A.1.* The proof is analogous to that of [Nakamura 2009b; Ito 2006; Ito and Nakamura 2009, Section 3], and we only sketch the main steps. We may suppose  $a$  is nonnegative without loss of generality. If we set

$$\psi(t) = a \circ \exp(tH_{p_c})^{-1},$$

then it is easy to see that

$$\frac{\partial}{\partial t} \psi = -\{p_c, \psi\}, \quad \psi(0) = a,$$

and this is a good candidate for the principal term of  $b(t)$ , but  $\psi$  does not satisfy the boundedness of the derivatives uniformly in  $t$ . We choose  $\varphi \in C_0^\infty(\mathbb{R})$  so that

$$\text{supp } \varphi \subset [-1, 1], \quad \varphi(t) \geq 0 \text{ for all } t, \quad \int_{-1}^1 \varphi(t) dt = 1,$$

and moreover,  $\pm\varphi'(t) \leq 0$  for  $\pm t \geq 0$ . We set

$$\varphi_\nu(t) = \varphi(t/\nu), \quad \text{for } \nu > 0,$$

and we denote convolution in  $t$  by  $\ast_t$ . Then we set

$$b_0(t, \cdot) = \varphi_{\delta\langle t \rangle} \ast_t \psi = \int \varphi_{\delta\langle t \rangle}(t-s)\psi(s, \cdot) ds$$

with sufficiently small  $\delta > 0$ . Then we have

$$\begin{aligned} \frac{\partial}{\partial t} b_0 &= \int \partial_t(\varphi_{\delta\langle t \rangle}(t-s))\psi(s, \cdot) ds = - \int \frac{t(t-s)}{\delta\langle t \rangle^3} \varphi'((t-s)/\delta\langle t \rangle)\psi(s, \cdot) ds + \varphi_{\delta\langle t \rangle} \ast_t (\partial_t \psi) \\ &\geq -\varphi_{\delta\langle t \rangle} \ast_t \{p_c, \psi\} = -\{p_c, b_0(t, \cdot)\} \end{aligned} \tag{A-1}$$

for  $t > 0$ , by the conditions on  $\varphi$ . We have the reverse inequality for  $t < 0$ .

We then show the derivatives of  $b_0$  satisfy the required uniform boundedness. We first note that

$$\tilde{\psi}(t; r, \rho, \theta, \omega) := \psi(t; r + t\rho, \rho, \theta, \omega) \rightarrow a \circ w_{\pm} \quad (t \rightarrow \pm\infty)$$

in the  $C_0^\infty$ -topology, by virtue of the existence of the classical scattering for  $p_c$ . Thus we have the representation

$$\psi(t; r, \rho, \theta, \omega) = \tilde{\psi}(t; r - t\rho, \rho, \theta, \omega),$$

with  $\tilde{\psi}(t)$  uniformly bounded in  $C_0^\infty(T^*M)$ . Hence we learn that the derivatives in variables other than  $\rho$  are uniformly bounded. Then this property applies also to  $b_0(t)$ . Let us consider the first derivative of  $b_0(t)$  in  $\rho$ :

$$\partial_\rho b_0(t) = - \int \varphi_{\delta\langle t \rangle}(t-s)s(\partial_r \tilde{\psi})(s, r - s\rho, \rho, \theta, \omega) ds + \int \varphi_{\delta\langle t \rangle}(t-s)(\partial_\rho \tilde{\psi})(s, r - s\rho, \rho, \theta, \omega) ds.$$

The second term is clearly uniformly bounded. We note that

$$(\partial_r \tilde{\psi})(s; r - s\rho, \rho, \theta, \omega) = -\frac{1}{\rho} \left\{ \frac{\partial}{\partial s} [\tilde{\psi}(s; r - s\rho, \rho, \theta, \omega)] - (\partial_s \tilde{\psi})(s; r - s\rho, \rho, \theta, \omega) \right\},$$

and then by integration by parts we have

$$\begin{aligned} & \int \varphi_{\delta\langle t \rangle}(t-s)s(\partial_r \tilde{\psi})(s, r-s\rho, \rho, \theta, \omega) ds \\ &= \frac{1}{\rho} \int \frac{\partial}{\partial s} \left( \varphi \left( \frac{t-s}{\delta\langle t \rangle} \right) s \right) \tilde{\psi}(s, r-s\rho, \rho, \theta, \omega) ds + \frac{1}{\rho} \int \varphi \left( \frac{t-s}{\delta\langle t \rangle} \right) s(\partial_s \tilde{\psi})(s; r-s\rho, \rho, \theta, \omega) ds \\ &= \frac{1}{\rho} \int \varphi \left( \frac{t-s}{\delta\langle t \rangle} \right) \tilde{\psi}(s, r-s\rho, \rho, \theta, \omega) ds - \frac{1}{\rho} \int \frac{s}{\delta\langle t \rangle} \varphi' \left( \frac{t-s}{\delta\langle t \rangle} \right) \tilde{\psi}(s, r-s\rho, \rho, \theta, \omega) ds \\ & \qquad \qquad \qquad + \frac{1}{\rho} \int \varphi \left( \frac{t-s}{\delta\langle t \rangle} \right) s(\partial_s \tilde{\psi})(s; r-s\rho, \rho, \theta, \omega) ds. \end{aligned}$$

Each term in the last expression is bounded uniformly in  $t$  since  $s \sim t$ , and  $\partial_s \tilde{\psi} = O(\langle s \rangle^{-2})$ . Repeating this procedure, we can show that all the derivatives of  $b_0$  are uniformly bounded. It is also easy to check that  $b_0$  satisfies the required support property provided  $a$  is supported in a sufficiently small neighborhood, and  $\delta > 0$  is chosen sufficiently small.

Now by (A-1) and the sharp Gårding inequality, we have

$$\frac{d}{dt} \text{Op}^W(b_0^h(t)) \geq -\frac{i}{h} [P, \text{Op}^W(b_0^h(t))] + \text{Op}(r_1^h(t))$$

with  $r_1(t) = O(h^\mu)$ . We set  $c_j = 7/4 - 2^{-j}$  for  $j = 1, 2, \dots$ , and set

$$a_j(r, \rho, \theta, \omega) = a\left(\frac{r}{c_j}, \frac{\rho}{c_j}, \frac{\theta}{c_j}, \frac{\omega}{c_j}\right), \quad b_j(t) = \varphi_{\delta\langle t \rangle} *_t (a_j \circ \exp(tH_{p_c})).$$

Then we set

$$b(t) \sim b_0(t) + \sum_{j=1}^{\infty} \mu_j b_j(t),$$

with appropriately chosen constants  $\mu_j > 0$  so that

$$\frac{d}{dt} \text{Op}^W(b^h(t)) \geq -\frac{i}{h} [P, \text{Op}^W(b^h(t))] + O(h^\infty),$$

and  $b(t)$  satisfies all the required properties. We refer to [Nakamura 2009b; Ito and Nakamura 2009] for the details of the construction. □

### Appendix B: Beals-type characterization of Fourier integral operators

In this appendix, we consider operators on  $\mathbb{R}^n$ , and we discuss Beals-type characterization of FIOs in terms of  $h$ -pseudodifferential operators. We use the result for scattering manifolds, but the generalization is straightforward, and we omit it. Most of the arguments here are similar to those of [Ito and Nakamura 2012, Section 2], and we mainly discuss the modifications necessary to show our results.

We let  $S$  be a canonical diffeomorphism on  $T^*\mathbb{R}^n$ , which is also supposed to be homogeneous in the  $\xi$ -variable, i.e.,

$$\text{if } (y, \eta) = S(x, \xi), \text{ then } S(x, \lambda\xi) = (y, \lambda\eta) \text{ for } \lambda > 0.$$

We also let  $U \in \mathcal{L}(\mathcal{S}, \mathcal{S}')$ , and let  $u \in \mathcal{D}'(\mathbb{R}^{2n})$  be its distribution kernel. For a symbol  $a \in C^\infty(T^*\mathbb{R}^n)$ , we write

$$a^h(x, \xi) = a(x, h\xi), \quad \text{Op}^W(a^h) = a^W(x, hD_x),$$

for  $h > 0$  as before. For  $a \in C_0^\infty(T^*\mathbb{R}^n \setminus 0)$ , we define

$$\text{Ad}_S(a^h)U = \text{Op}^W(a^h \circ S^{-1})U - U\text{Op}^W(a^h) \in \mathcal{L}(\mathcal{S}, \mathcal{S}').$$

We note that  $\text{Op}^W(a^h \circ S^{-1}) = \text{Op}^W((a \circ S^{-1})^h)$  since  $S$  is homogeneous in  $\xi$ .

**Theorem B.1.** *Let  $U \in B(L_{cpt}^2(\mathbb{R}^n), L_{loc}^2(\mathbb{R}^n))$ . Suppose for any  $a_1, a_2, \dots, a_N \in C_0^\infty(T^*\mathbb{R}^n \setminus 0)$ , there is  $C_N > 0$  such that*

$$\|\text{Ad}_S(a_1^h) \text{Ad}_S(a_2^h) \cdots \text{Ad}_S(a_N^h)U\|_{B(L^2)} \leq C_N h^N. \quad (\text{B-1})$$

*Then  $U$  is an FIO of order 0 associated to  $S$ .*

**Corollary B.2.** *Let  $S$  and  $U$  as above. If for any  $a \in C_0^\infty(T^*\mathbb{R}^n \setminus 0)$  there is an  $h$ -dependent symbol  $b \in C_0^\infty(T^*\mathbb{R}^n \setminus 0)$  such that*

$$|\partial_x^\alpha \partial_\xi^\beta b(h; x, \xi)| \leq C_{\alpha\beta} h,$$

*for any  $\alpha, \beta \in \mathbb{Z}_+^n$ ,  $h \in (0, 1]$ , and*

$$\text{Ad}_S(a^h)U = \text{Op}^W(b^h)U + R, \quad \|R\|_{B(L^2)} = O(h^\infty),$$

*then  $U$  is an FIO of order 0 associated to  $S$ .*

*Proof of Corollary B.2.* We show (B-1) follows from the above condition. The cases  $N = 0, 1$  are obvious. Let  $N = 2$  and we write

$$\text{Ad}_S(a_j^h)U = \text{Op}^W(b_j^h)U + R_j, \quad j = 1, 2.$$

Then we have

$$\begin{aligned} \text{Ad}_S(a_1^h) \text{Ad}_S(a_2^h)U &= \text{Op}^W(a_1^h \circ S^{-1})\text{Op}^W(b_2^h)U - \text{Op}^W(b_2^h)U\text{Op}^W(a_1^h) + \text{Ad}_S(a_1^h)R_2 \\ &= [\text{Op}^W(a_1^h \circ S^{-1}), \text{Op}^W(b_2^h)]U + \text{Op}^W(b_2^h)\text{Op}^W(b_1^h)U + \text{Ad}_S(a_1^h)R_2 + \text{Op}^W(b_2^h)R_1 \\ &= \text{Op}^W(b_{12}^h)U + R_{12}, \end{aligned}$$

where  $R_{12} = O(h^\infty)$  and  $b_{12} \in C_0^\infty(T^*\mathbb{R}^n \setminus 0)$  satisfies

$$|\partial_x^\alpha \partial_\xi^\beta b_{12}(h; x, \xi)| \leq C'_{\alpha\beta} h^2, \quad \text{for any } \alpha, \beta \in \mathbb{Z}_+^n, h \in (0, 1],$$

and (B-1) for  $N = 2$  follows. Iterating this procedure, we obtain (B-1) for any  $N$ .  $\square$

In order to prove Theorem B.1, we first note the semiclassical-type characterization of Besov spaces. By the standard partition-of-unity argument, it is straightforward to observe that  $u \in B_{2,loc}^{\sigma,\infty}(\mathbb{R}^m)$  if and only if for any  $(x_0, \xi_0) \in T^*\mathbb{R}^m \setminus 0$  there is  $\varphi \in C_0^\infty(T^*\mathbb{R}^m)$  such that  $\varphi(x_0, \xi_0) \neq 0$  and

$$\|\text{Op}^W(\varphi^h)u\|_{L^2} \leq Ch^\sigma, \quad h > 0.$$

Thus, in turn, we learn that  $u \in B_{2,loc}^{\sigma,\infty}(\mathbb{R}^{2n})$  if and only if for any  $(x_0, y_0, \xi_0, \eta_0), (\xi_0, \eta_0) \neq (0, 0)$ , there are  $\varphi_1, \varphi_2 \in C_0^\infty(\mathbb{R}^n)$  such that  $\varphi_1(x_0, \xi_0) \neq 0, \varphi_2(y_0, \eta_0) \neq 0$ , and

$$\|\text{Op}^W(\varphi_1^h)U\text{Op}^W(\varphi_2^h)\|_{HS} \leq Ch^\sigma, \quad h > 0,$$

where  $\|\cdot\|_{HS}$  denotes the Hilbert–Schmidt norm in  $B(L^2(\mathbb{R}^n))$ . Now we choose  $\varphi_3 \in C_0^\infty(\mathbb{R}^n)$  so that  $\varphi_3 = 1$  in a neighborhood of  $\text{supp } \varphi_2$ . We note that

$$\|\text{Op}^W(\varphi_3)\|_{HS} = (2\pi)^{-n/2} \left( \int_{\mathbb{R}^n} |\varphi_3(x, h\xi)|^2 dx d\xi \right)^{1/2} = (2\pi h)^{-n/2} \left( \int_{\mathbb{R}^n} |\varphi_3(x, \xi)|^2 dx d\xi \right)^{1/2} = Ch^{-n/2}$$

for  $h > 0$  with some  $C > 0$ . Hence we have

$$\begin{aligned} \|\text{Op}^W(\varphi_1^h)U\text{Op}^W(\varphi_2^h)\|_{HS} &\leq \|\text{Op}^W(\varphi_1^h)U\text{Op}^W(\varphi_2^h)\text{Op}^W(\varphi_3^h)\|_{HS} + R \\ &\leq Ch^{-n/2}\|\text{Op}^W(\varphi_1^h)U\text{Op}^W(\varphi_2^h)\|_{B(L^2)} + R, \end{aligned}$$

where

$$R = \|\text{Op}^W(\varphi_1^h)U\text{Op}^W(\varphi_2^h)(1 - \text{Op}^W(\varphi_3^h))\|_{HS} = O(h^\infty)$$

by the symbol calculus. Thus we have proved the following lemma:

**Lemma B.3.** *If for any  $(x_0, y_0, \xi_0, \eta_0) \in T^*\mathbb{R}^{2n}$  with  $(\xi_0, \eta_0) \neq (0, 0)$  there are  $\varphi_1, \varphi_2 \in C_0^\infty(T^*\mathbb{R}^n)$  such that  $\varphi_1(x_0, \xi_0) \neq 0, \varphi_2(y_0, \eta_0) \neq 0$  and*

$$\|\text{Op}^W(\varphi_1^h)U\text{Op}^W(\varphi_2^h)\|_{B(L^2)} \leq C, \quad h > 0,$$

then  $u \in B_{2,loc}^{-n/2,\infty}(\mathbb{R}^{2n})$ .

*Proof of Theorem B.1.* We modify the proof of Theorem 2.1 in [Ito and Nakamura 2012], to which we refer for further details.

We first note that

$$WF(u) \subset \Lambda_S = \{(x, y, \xi, -\eta) \in T^*\mathbb{R}^{2n} \mid (x, \xi) = S(y, \eta)\}.$$

We note that if  $(x_0, y_0, \xi_0, -\eta_0) \notin \Lambda_S$  with  $\eta_0 \neq 0$ , it is straightforward to show  $(x_0, y_0, \xi_0, -\eta_0) \notin WF(u)$ . If  $\xi_0 \neq 0$ , we consider  $U^*$  and we can also conclude  $(x_0, y_0, \xi_0, -\eta_0) \notin WF(u)$ .

Now we let  $a_1, a_2, \dots, a_N \in S_{cl}^1(\mathbb{R}^n)$  and let  $(x_0, \xi_0) = S(y_0, \eta_0)$ . We may assume  $a_j$  are homogeneous of order one in the  $\xi$ -variable. By Lemma B.3 and the proof just cited, it suffices to show the following to conclude  $U$  is an FIO of order 0 associated to  $S$ : There are  $\psi_1, \psi_2 \in C_0^\infty(T^*\mathbb{R}^n)$  such that  $\psi_1(x_0, \xi_0) \neq 0, \psi_2(y_0, \eta_0) \neq 0$  and

$$\|\text{Op}^W(\psi_1^h)[\text{Ad}_S(a_1) \cdots \text{Ad}_S(a_N)U]\text{Op}^W(\psi_2^h)\|_{B(L^2)} \leq C, \quad h \in (0, 1], \tag{B-2}$$

with some  $C > 0$ .

We set  $\Psi_0, \Psi_1 \in C_0^\infty(T^*\mathbb{R}^n)$  so that they are supported in a small neighborhood of  $(y_0, \eta_0)$ ,  $\Psi_j = 1$  on a neighborhood of  $(y_0, \eta_0)$ , and  $\Psi_0 = 1$  on  $\text{supp } \Psi_1$ . We then set

$$\varphi_j(x, \xi) = a_j(x, \xi)\Psi_0(x, \xi) \in C_0^\infty(T^*\mathbb{R}^n).$$



We note, since  $a_j$  are homogeneous of order one in  $\xi$ , that

$$a_j(x, \xi)\Psi_0(x, h\xi) = h^{-1}a_j(x, h\xi)\Psi_0(x, h\xi) = h^{-1}\varphi_j(x, h\xi).$$

We also set

$$\psi_1 = \Psi_1 \circ S^{-1} \quad \text{and} \quad \psi_2 = \Psi_1$$

so  $\psi_1(1 - \Psi_0 \circ S^{-1}) = 0$  and  $(1 - \Psi_0)\psi_2 = 0$ . This implies, in particular, that

$$\begin{aligned} \psi_1(x, h\xi)(a_j \circ S^{-1})(x, \xi) &= h^{-1}\psi_1(x, h\xi)(\varphi_j \circ S^{-1})(x, h\xi), \\ a_j(y, \eta)\psi_2(y, h\eta) &= h^{-1}\varphi_j(y, h\eta)\psi_2(y, h\eta). \end{aligned}$$

Using these, and applying the  $h$ -pseudodifferential operator calculus, we learn that

$$\begin{aligned} \text{Op}^W(\psi_1^h)[\text{Ad}_S(a_1) \cdots \text{Ad}_S(a_N)U]\text{Op}^W(\psi_2^h) \\ = h^{-N}\text{Op}^W(\psi_1^h)[\text{Ad}_S(\varphi_1^h) \cdots \text{Ad}_S(\varphi_N^h)U]\text{Op}^W(\psi_2^h) + O(h^\infty), \end{aligned}$$

and this implies the right-hand side is bounded by the assumption of [Theorem B.1](#). Now [\(B-2\)](#) follows from this observation, and we conclude that the assertion hold. □

We note that the conditions and the assertion of [Theorem B.1](#) are microlocal, and hence the theorem is easily extended to a statement in a conic set in  $T^*\mathbb{R}^n$ . In the next proposition, we use the extended statement on a conic set.

**Proposition B.4.** *Let  $\mathbb{R}^m = \mathbb{R}^n \times \mathbb{R}^k$ , and let  $U$  be a bounded operator on  $L^2(\mathbb{R}^m)$  and let  $S$  be a homogeneous canonical diffeomorphism on  $T^*\mathbb{R}^m$ . Suppose  $U$  commutes with multiplication operators in  $y$  so that  $U$  is decomposed as*

$$U = \int_{\mathbb{R}^k}^{\oplus} \tilde{U}(y)dy \quad \text{on } L^2(\mathbb{R}^m) \cong L^2(\mathbb{R}_y^k, L^2(\mathbb{R}_x^n)),$$

where  $\{U(y)\}$  is a family of operators on  $L^2(\mathbb{R}_x^n)$ . Suppose also that  $S$  is decomposed as

$$S : (x, \xi, y, \eta) \mapsto (\tilde{S}(y)(x, \xi), y, \eta + g(x, \xi, y))$$

for  $(x, \xi, y, \eta) \in T^*\mathbb{R}^n \cong T^*\mathbb{R}_x^n \times T^*\mathbb{R}_y^k$ , where  $\{\tilde{S}(y)\}$  is a family of canonical maps on  $T^*\mathbb{R}_x^n$ . If  $U$  is an FIO associated to  $S$  on a conic set  $\{(x, \xi, t, \eta) \mid \xi \neq 0\}$ , then for each  $y \in \mathbb{R}^k$ ,  $\tilde{U}(y)$  is an FIO of order 0 associated to  $\tilde{S}(y)$ .

**Remark B.5.** The assumption on  $S$  actually follows from the properties of  $U$ . We include it to introduce the notations.

*Proof.* Let  $a \in C_0^\infty(T^*\mathbb{R}^n \setminus 0)$ , and let  $\varphi, \psi \in C_0^\infty(\mathbb{R}^k)$  such that  $\varphi, \psi \geq 0$  and  $\int \psi(\eta) d\eta = 1$ . We also denote  $\psi_z(\eta) = \psi(\eta - z)$  for  $z \in \mathbb{R}^k$ . We consider

$$A_z = a_z(x, hD_x, y, hD_y) = a(x, hD_x)\varphi(y)\psi_z(hD_y).$$

Since  $U$  is an FIO, there is  $b_z$ , which is bounded in  $C_0^\infty(T^*\mathbb{R}^m)$  uniformly in  $h \in (0, 1]$ , such that

$$UA_z = B_zU + O(h^\infty), \quad B_z = b_z(x, hD_x, y, hD_y),$$

with the principal symbol

$$a_z \circ S^{-1} = (a \circ \tilde{S}(y)^{-1})(x, \xi)\varphi(y)\psi(\eta - g(\tilde{S}(y)^{-1}(x, \xi), y) - z).$$

Since  $U$  commutes with  $\{e^{iy \cdot z} \mid z \in \mathbb{R}^k\}$  (translations in the  $\eta$ -variable), we learn that

$$b_z(x, \xi, y, \eta) = b_0(x, \xi, y, \eta - z),$$

and the remainder term also satisfies this property. Moreover, these symbols decay rapidly outside  $S[\text{supp } a_z]$ .

It is also easy to see that

$$\int_{|z| \leq R} A_z dz \rightarrow a(x, hD_x)\varphi(y) \quad \text{and} \quad \int_{|z| \leq R} B_z dz \rightarrow \tilde{b}(x, hD_x, y)$$

strongly as  $R \rightarrow \infty$ , where  $\tilde{b}(x, \xi, y) = \int_{\mathbb{R}^k} b_0(x, \xi, y, \eta) d\eta$ . The principal symbol of  $\tilde{b}$  is given by  $(a \circ \tilde{S}(y)^{-1})(x, \xi)\varphi(y)$ . These facts imply that

$$\tilde{U}(y)a(x, hD_x)\varphi(y) = \tilde{b}(x, hD_x, y)\tilde{U}(y) + O(h^\infty),$$

where  $\tilde{b}(x, \xi, y) - (a \circ \tilde{S}(y)^{-1})(x, \xi)\varphi(y) = O(h)$ . Since  $\varphi \in C_0^\infty(\mathbb{R}^k)$  is arbitrary, for a fixed  $y \in \mathbb{R}^k$  we may replace  $\varphi(y)$  by 1, and we learn  $\tilde{U}(y)$  is an FIO of order 0 associated to  $\tilde{S}(y)$  by [Corollary B.2](#).  $\square$

### Acknowledgement

We thank Dr. S. Itozaki for pointing out errors in [Lemma 4.1](#) and [Lemma 5.3](#) in the original manuscript. We also thank the referee for constructive comments.

### References

- [Craig et al. 1995] W. Craig, T. Kappeler, and W. Strauss, “Microlocal dispersive smoothing for the Schrödinger equation”, *Comm. Pure Appl. Math.* **48**:8 (1995), 769–860. [MR 96m:35057](#) [Zbl 0856.35106](#)
- [Hörmander 1985] L. Hörmander, *The analysis of linear partial differential operators, III: Pseudodifferential operators*, Grundlehren Math. Wiss. **274**, Springer, Berlin, 1985. [MR 87d:35002a](#) [Zbl 0601.35001](#)
- [Isozaki and Kitada 1986] H. Isozaki and H. Kitada, “Scattering matrices for two-body Schrödinger operators”, *Sci. Papers College Arts Sci. Univ. Tokyo* **35**:2 (1986), 81–107. [MR 87k:35196](#) [Zbl 0615.35065](#)
- [Ito 2006] K. Ito, “Propagation of singularities for Schrödinger equations on the Euclidean space with a scattering metric”, *Comm. Partial Differential Equations* **31**:10-12 (2006), 1735–1777. [MR 2008c:35258](#) [Zbl 1117.35005](#)
- [Ito and Nakamura 2009] K. Ito and S. Nakamura, “Singularities of solutions to the Schrödinger equation on scattering manifold”, *Amer. J. Math.* **131**:6 (2009), 1835–1865. [MR 2011g:58046](#) [Zbl 1186.35004](#)
- [Ito and Nakamura 2010] K. Ito and S. Nakamura, “Time-dependent scattering theory for Schrödinger operators on scattering manifolds”, *J. Lond. Math. Soc.* (2) **81**:3 (2010), 774–792. [MR 2011g:58047](#) [Zbl 1204.58024](#)
- [Ito and Nakamura 2012] K. Ito and S. Nakamura, “Remarks on the fundamental solution to Schrödinger equation with variable coefficients”, *Ann. Inst. Fourier* **63**:3 (2012), 1091–1121. [Zbl 06093174](#)

- [Ito and Skibsted 2011] K. Ito and E. Skibsted, “Absence of embedded eigenvalues for Riemannian Laplacians”, preprint, 2011. [arXiv 1109.1928](#)
- [Martinez 2002] A. Martinez, *An introduction to semiclassical and microlocal analysis*, Springer, New York, 2002. [MR 2003b:35010](#) [Zbl 0994.35003](#)
- [Melrose 1994] R. B. Melrose, “Spectral and scattering theory for the Laplacian on asymptotically Euclidian spaces”, pp. 85–130 in *Spectral and scattering theory* (Sanda, Japan, 1992), edited by M. Ikawa, Lecture Notes in Pure and Appl. Math. **161**, Dekker, New York, 1994. [MR 95k:58168](#) [Zbl 0837.35107](#)
- [Melrose and Zworski 1996] R. Melrose and M. Zworski, “Scattering metrics and geodesic flow at infinity”, *Invent. Math.* **124**:1-3 (1996), 389–436. [MR 96k:58230](#) [Zbl 0855.58058](#)
- [Nakamura 2009a] S. Nakamura, “Semiclassical singularities propagation property for Schrödinger equations”, *J. Math. Soc. Japan* **61**:1 (2009), 177–211. [MR 2010e:35062](#) [Zbl 1176.35045](#)
- [Nakamura 2009b] S. Nakamura, “Wave front set for solutions to Schrödinger equations”, *J. Funct. Anal.* **256**:4 (2009), 1299–1309. [MR 2010h:35068](#) [Zbl 1155.35014](#)
- [Taylor 1981] M. E. Taylor, *Pseudodifferential operators*, Princeton Mathematical Series **34**, Princeton University Press, 1981. [MR 82i:35172](#) [Zbl 0453.47026](#)
- [Vasy 1998] A. Vasy, “Geometric scattering theory for long-range potentials and metrics”, *Internat. Math. Res. Notices* **1998**:6 (1998), 285–315. [MR 99b:58233](#) [Zbl 0922.58085](#)
- [Yafaev 2000] D. Yafaev, *Scattering theory: some old and new problems*, Lecture Notes in Mathematics **1735**, Springer, Berlin, 2000. [MR 2001j:81248](#) [Zbl 0951.35003](#)

Received 20 Apr 2011. Revised 28 Mar 2012. Accepted 23 May 2012.

KENICHI ITO: [ito\\_ken@math.tsukuba.ac.jp](mailto:ito_ken@math.tsukuba.ac.jp)

*Graduate School of Pure and Applied Sciences, University of Tsukuba, 1-1-1 Tennodai, Tsukuba, Ibaraki 305-8571, Japan*

SHU NAKAMURA: [shu@ms.u-tokyo.ac.jp](mailto:shu@ms.u-tokyo.ac.jp)

*Graduate School of Mathematical Sciences, University of Tokyo, 3-8-1, Komaba, Meguro, Tokyo 153-8914, Japan*

<http://www.ms.u-tokyo.ac.jp/~shu/>

# Analysis & PDE

[msp.org/apde](http://msp.org/apde)

## EDITORS

EDITOR-IN-CHIEF

Maciej Zworski  
[zworski@math.berkeley.edu](mailto:zworski@math.berkeley.edu)

University of California  
Berkeley, USA

## BOARD OF EDITORS

Michael Aizenman	Princeton University, USA <a href="mailto:aizenman@math.princeton.edu">aizenman@math.princeton.edu</a>	Nicolas Burq	Université Paris-Sud 11, France <a href="mailto:nicolas.burq@math.u-psud.fr">nicolas.burq@math.u-psud.fr</a>
Luis A. Caffarelli	University of Texas, USA <a href="mailto:caffarel@math.utexas.edu">caffarel@math.utexas.edu</a>	Sun-Yung Alice Chang	Princeton University, USA <a href="mailto:chang@math.princeton.edu">chang@math.princeton.edu</a>
Michael Christ	University of California, Berkeley, USA <a href="mailto:mchrist@math.berkeley.edu">mchrist@math.berkeley.edu</a>	Charles Fefferman	Princeton University, USA <a href="mailto:cf@math.princeton.edu">cf@math.princeton.edu</a>
Ursula Hamenstaedt	Universität Bonn, Germany <a href="mailto:ursula@math.uni-bonn.de">ursula@math.uni-bonn.de</a>	Nigel Higson	Pennsylvania State University, USA <a href="mailto:higson@math.psu.edu">higson@math.psu.edu</a>
Vaughan Jones	University of California, Berkeley, USA <a href="mailto:vfr@math.berkeley.edu">vfr@math.berkeley.edu</a>	Herbert Koch	Universität Bonn, Germany <a href="mailto:koch@math.uni-bonn.de">koch@math.uni-bonn.de</a>
Izabella Laba	University of British Columbia, Canada <a href="mailto:ilaba@math.ubc.ca">ilaba@math.ubc.ca</a>	Gilles Lebeau	Université de Nice Sophia Antipolis, France <a href="mailto:lebeau@unice.fr">lebeau@unice.fr</a>
László Lempert	Purdue University, USA <a href="mailto:lempert@math.purdue.edu">lempert@math.purdue.edu</a>	Richard B. Melrose	Massachusetts Institute of Technology, USA <a href="mailto:rbm@math.mit.edu">rbm@math.mit.edu</a>
Frank Merle	Université de Cergy-Pontoise, France <a href="mailto:Frank.Merle@u-cergy.fr">Frank.Merle@u-cergy.fr</a>	William Minicozzi II	Johns Hopkins University, USA <a href="mailto:minicozz@math.jhu.edu">minicozz@math.jhu.edu</a>
Werner Müller	Universität Bonn, Germany <a href="mailto:mueller@math.uni-bonn.de">mueller@math.uni-bonn.de</a>	Yuval Peres	University of California, Berkeley, USA <a href="mailto:peres@stat.berkeley.edu">peres@stat.berkeley.edu</a>
Gilles Pisier	Texas A&M University, and Paris 6 <a href="mailto:pisier@math.tamu.edu">pisier@math.tamu.edu</a>	Tristan Rivière	ETH, Switzerland <a href="mailto:riviere@math.ethz.ch">riviere@math.ethz.ch</a>
Igor Rodnianski	Princeton University, USA <a href="mailto:irod@math.princeton.edu">irod@math.princeton.edu</a>	Wilhelm Schlag	University of Chicago, USA <a href="mailto:schlag@math.uchicago.edu">schlag@math.uchicago.edu</a>
Sylvia Serfaty	New York University, USA <a href="mailto:serfaty@cims.nyu.edu">serfaty@cims.nyu.edu</a>	Yum-Tong Siu	Harvard University, USA <a href="mailto:siu@math.harvard.edu">siu@math.harvard.edu</a>
Terence Tao	University of California, Los Angeles, USA <a href="mailto:tao@math.ucla.edu">tao@math.ucla.edu</a>	Michael E. Taylor	Univ. of North Carolina, Chapel Hill, USA <a href="mailto:met@math.unc.edu">met@math.unc.edu</a>
Gunther Uhlmann	University of Washington, USA <a href="mailto:gunther@math.washington.edu">gunther@math.washington.edu</a>	András Vasy	Stanford University, USA <a href="mailto:andras@math.stanford.edu">andras@math.stanford.edu</a>
Dan Virgil Voiculescu	University of California, Berkeley, USA <a href="mailto:dvv@math.berkeley.edu">dvv@math.berkeley.edu</a>	Steven Zelditch	Northwestern University, USA <a href="mailto:zelditch@math.northwestern.edu">zelditch@math.northwestern.edu</a>

## PRODUCTION

[production@msp.org](mailto:production@msp.org)

Silvio Levy, Scientific Editor

---

See inside back cover or [msp.org/apde](http://msp.org/apde) for submission instructions.

---

The subscription price for 2013 is US \$160/year for the electronic version, and \$310/year (+\$35, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues from the last three years and changes of subscribers address should be sent to MSP.

---

Analysis & PDE (ISSN 1948-206X electronic, 2157-5045 printed) at Mathematical Sciences Publishers, 798 Evans Hall #3840, c/o University of California, Berkeley, CA 94720-3840, is published continuously online. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices.

---

APDE peer review and production are managed by EditFLOW<sup>®</sup> from Mathematical Sciences Publishers.

PUBLISHED BY

 **mathematical sciences publishers**  
nonprofit scientific publishing

<http://msp.org/>

© 2013 Mathematical Sciences Publishers

# ANALYSIS & PDE

Volume 6 No. 2 2013

---

Microlocal properties of scattering matrices for Schrödinger equations on scattering manifolds	257
KENICHI ITO and SHU NAKAMURA	
Local well-posedness of the viscous surface wave problem without surface tension	287
YAN GUO and IAN TICE	
Hypoellipticity and nonhypoellipticity for sums of squares of complex vector fields	371
ANTONIO BOVE, MARCO MUGHETTI and DAVID S. TARTAKOFF	
Porous media: the Muskat problem in three dimensions	447
ANTONIO CÓRDOBA, DIEGO CÓRDOBA and FRANCISCO GANCEDO	
Embeddings of infinitely connected planar domains into $\mathbb{C}^2$	499
FRANC FORSTNERIČ and ERLEND FORNÆSS WOLD	