

ANALYSIS & PDE

Volume 7

No. 3

2014

LORENZO GIACOMELLI, JOSE M. MAZÓN AND SALVADOR MOLL

**THE 1-HARMONIC FLOW WITH VALUES IN A HYPEROCTANT
OF THE N -SPHERE**

THE 1-HARMONIC FLOW WITH VALUES IN A HYPEROCTANT OF THE N -SPHERE

LORENZO GIACOMELLI, JOSE M. MAZÓN AND SALVADOR MOLL

We prove the existence of solutions to the 1-harmonic flow — that is, the formal gradient flow of the total variation of a vector field with respect to the L^2 -distance — from a domain of \mathbb{R}^m into a hyperoctant of the N -dimensional unit sphere, \mathbb{S}_+^{N-1} , under homogeneous Neumann boundary conditions. In particular, we characterize the lower-order term appearing in the Euler–Lagrange formulation in terms of the “geodesic representative” of a BV-director field on its jump set. Such characterization relies on a lower semicontinuity argument which leads to a nontrivial and nonconvex minimization problem: to find a shortest path between two points on \mathbb{S}_+^{N-1} with respect to a metric which penalizes the closeness to their geodesic midpoint.

1. Introduction	627
2. Preliminaries	631
3. Existence of solutions	639
4. A nonconvex variational problem	652
Acknowledgement	669
References	669

1. Introduction

Throughout the paper, $\Omega \subset \mathbb{R}^m$ is a bounded domain with Lipschitz continuous boundary $\partial\Omega$ and \mathbb{S}^{N-1} is the unit sphere of \mathbb{R}^N . For a smooth map $\mathbf{u} : \Omega \rightarrow \mathbb{S}^{N-1}$ and $1 \leq p < \infty$, the p -energy of \mathbf{u} is given by

$$E_p(\mathbf{u}) = \int_{\Omega} |\mathbf{D}\mathbf{u}|^p dx.$$

A critical point $\mathbf{u} \in C^1(\Omega; \mathbb{S}^{N-1})$ of the p -energy, a p -harmonic map, formally satisfies the Euler–Lagrange equation

$$-\operatorname{div}(|\mathbf{D}\mathbf{u}|^{p-2} \mathbf{D}\mathbf{u}) = |\mathbf{D}\mathbf{u}|^p \mathbf{u}. \tag{1-1}$$

The term $|\mathbf{D}\mathbf{u}|^p$ plays the role of a Lagrange multiplier corresponding to the pointwise constraint $|\mathbf{u}| = 1$.

Mazón and Moll have been partially supported by the Spanish MEC project MTM2012–31103.

MSC2010: primary 35K55, 49Q20, 53C22, 35K67, 35K92; secondary 49J45, 53C44, 58E20, 68U10.

Keywords: harmonic flows, total variation flow, nonlinear parabolic systems, lower semicontinuity and relaxation, nonconvex variational problems, geodesics, Riemannian manifolds with boundary, image processing.

One well-known method to obtain (distributional) solutions to (1-1), the so-called *heat-flow method*, introduced by J. Eells and J. H. Sampson [1964] for $p = 2$ in the general framework of Riemannian manifolds, consists in looking at long time limits of solutions to

$$\mathbf{u}_t = \operatorname{div}(|D\mathbf{u}|^{p-2}D\mathbf{u}) + \mathbf{u}|D\mathbf{u}|^p, \quad |\mathbf{u}| = 1. \quad (1-2)$$

Equation (1-2) is also a prototype for often quite complicated reaction-diffusion systems for the evolution of director fields which arise in various contexts — multigrain problems [Kobayashi et al. 2000], theory of liquid crystals [van der Hout 2001], ferromagnetism [DeSimone and Podio-Guidugli 1996], and image processing [Sapiro 2001]. For $p > 1$, (1-2) with various boundary conditions has been widely studied over the last decades; referenced discussions of the cases $p = 2$ and $p \in (1, \infty)$ may be found, for example, in [Bertsch et al. 2003; Bertsch et al. 2002; Chen 1989; Struwe 1992] and [Chen et al. 1994; Hungerbühler 2004; Misawa 2002], respectively.

Here we are interested in the case $p = 1$, for which (1-2) formally reads

$$\mathbf{u}_t = \operatorname{div}\left(\frac{D\mathbf{u}}{|D\mathbf{u}|}\right) + \mathbf{u}|D\mathbf{u}|, \quad \mathbf{u} \in \mathbb{S}^{N-1}. \quad (1-3)$$

More precisely, we focus on the homogeneous Neumann problem for (1-3) when the target space is a compact subset \mathbb{A} of \mathbb{S}^{N-1} ; that is,

$$\begin{cases} \mathbf{u}_t = \operatorname{div}\left(\frac{D\mathbf{u}}{|D\mathbf{u}|}\right) + \mathbf{u}|D\mathbf{u}|, \quad \mathbf{u} \in \mathbb{A} \subseteq \mathbb{S}^{N-1}, & \text{in } Q_T = (0, T) \times \Omega, \\ \frac{D\mathbf{u}}{|D\mathbf{u}|} \nu = 0 & \text{on } S_T = (0, T) \times \partial\Omega, \\ \mathbf{u}(0, \cdot) = \mathbf{u}_0(\cdot), \quad \mathbf{u}_0 \in \mathbb{A}, & \text{in } \Omega, \end{cases} \quad (1-4)$$

where ν denotes the outward unit normal to $\partial\Omega$. Problem (1-4) was proposed as a tool to denoise either two-dimensional image gradients and optical flows, in which case $N = 2$ and $\mathbb{A} = \mathbb{S}^1$ [Tang et al. 2000], or color images by smoothing the chromaticity data while preserving the contrast, in which case $N = 3$ and \mathbb{A} is an octant of the sphere [Tang et al. 2001].

While the scalar and unconstrained version of (1-3), that is, the so-called total variation flow, is by now well understood after the pioneering paper [Andreu et al. 2001] (see the monograph [Andreu-Vaillio et al. 2004] and the references therein or [Bonforte and Figalli 2012] for an up-to-date reference list). An existence theory for (1-3) is still open in general. Special cases considered so far have dealt with piece-wise constant data [Giga and Kobayashi 2003; Giga et al. 2005; Giga et al. 2007], initial data with “small” energy [Giga et al. 2004], and rotationally symmetric solutions [Giga and Kuroda 2004; Dal Passo et al. 2008; Giacomelli and Moll 2010]. We refer to [Giacomelli et al. 2013a] for a detailed discussion of previous attempts to obtain a solution to (1-4) given in [Barrett et al. 2008; Feng 2010].

In dealing with (1-3), the most delicate issue is of course the interpretation of the bounded matrix \mathbf{Z} , which represents $D\mathbf{u}/|D\mathbf{u}|$, and of the measure $\boldsymbol{\mu}$, which represents $\mathbf{u}|D\mathbf{u}|$, the latter being the product between a measure and a possibly discontinuous function. Very recently, an interpretation of (1-3) has

been proposed in [Giacomelli et al. 2013a]: in summary,

$$\mathbf{u}_t(t) - \operatorname{div} \mathbf{Z}(t) \in \mathbf{u}_g |D\mathbf{u}|(t), \quad \mathbf{u}(t) \in \mathbb{A} \quad \text{for a.e. } t \in [0, T] \tag{1-5}$$

in the sense of distributions, where $\mathbf{Z}(t)$ is a bounded matrix that represents $D\mathbf{u}(t)/|D\mathbf{u}(t)|$ (the precise meaning is given in Proposition 3.5) and $\mathbf{u}_g |D\mathbf{u}|(t)$ denotes a set of vector-valued measures which are oriented as $\mathbf{u}(t)^*$ (the precise representative of $\mathbf{u}(t)$) and have total variation density $|D\mathbf{u}(t)|$. For $N = 2$, this interpretation has led to the existence and uniqueness of a solution to (1-4) when \mathbb{A} is a semicircle [Giacomelli et al. 2013a, Theorems 4.1 and 5.1] together with the existence of a solution when $\mathbb{A} = \mathbb{S}^1$ and $\mathbf{u}_0 \in BV(\Omega; \mathbb{S}^1)$ has no jumps by an ‘‘angle’’ larger than π .

The aim of this paper is to prove an existence result, according to the same interpretation, for an arbitrary dimension of the target sphere. We consider (1-4) in the first hyperoctant of the N -sphere:

$$\mathbb{A} = \mathbb{S}_+^{N-1} := \{(x_1, \dots, x_N) \in \mathbb{S}^{N-1} : x_i \geq 0 \text{ for } i = 1, \dots, N\}$$

(a natural assumption in the context of image processing; see above). Note that in this case, for every pair $\mathbf{u}_-, \mathbf{u}_+ \in \mathbb{S}_+^{N-1}$ there exists a unique *geodesic midpoint*, $\mathbf{u}_g = (\mathbf{u}_+ + \mathbf{u}_-)/|\mathbf{u}_+ + \mathbf{u}_-|$ (see Definition 3.1). Hence we may define the *geodesic representative* of $\mathbf{u} \in BV(\Omega; \mathbb{S}_+^{N-1})$, $\mathbf{u}_g := \mathbf{u}^*/|\mathbf{u}^*|$ (see Definition 3.2 and Remark 3.3) and the set of measures in (1-5) reduces to the singleton $\mathbf{u}(t)_g |D\mathbf{u}(t)|$.

The complete definition of a solution and the statement of the main result are given in Definition 3.4 and Theorem 3.6, respectively. We obtain a solution as the limit of a sequence of solutions to the following approximating problems (see Proposition 3.7 and Lemma 3.8):

$$\begin{cases} \mathbf{u}_t^\varepsilon = \operatorname{div} \mathbf{Z}^\varepsilon + \boldsymbol{\mu}^\varepsilon, & \mathbf{u}^\varepsilon \in \mathbb{S}_+^{N-1} & \text{in } \Omega_T, \\ [\mathbf{Z}^\varepsilon, \nu] = 0 & & \text{on } S_T, \\ \mathbf{u}^\varepsilon(0, \cdot) = \mathbf{u}_0^\varepsilon(\cdot) & & \text{in } \Omega, \end{cases}$$

where

$$\mathbf{Z}^\varepsilon = \varepsilon^\alpha \nabla \mathbf{u}^\varepsilon + \frac{\nabla \mathbf{u}^\varepsilon}{\sqrt{|\nabla \mathbf{u}^\varepsilon|^2 + \varepsilon^2}}, \quad \boldsymbol{\mu}^\varepsilon = \varepsilon^\alpha \mathbf{u}^\varepsilon |\nabla \mathbf{u}^\varepsilon|^2 + \mathbf{u}^\varepsilon \frac{|\nabla \mathbf{u}^\varepsilon|^2}{\sqrt{|\nabla \mathbf{u}^\varepsilon|^2 + \varepsilon^2}}, \tag{1-6}$$

and the initial data suitably converge to a given $\mathbf{u}_0 \in BV(\Omega; \mathbb{S}_+^{N-1})$ (see Lemma 3.9). The strategy we follow is completely different from that in [Giacomelli et al. 2013a], where the special structure of \mathbb{S}^1 was heavily used. Its core, neglecting any technicality and concentrating on the crucial issues, may be summarized as follows (see also [Giacomelli et al. 2013b] for a slightly more detailed discussion). By fairly standard compactness arguments, we obtain convergence of \mathbf{u}^ε , \mathbf{Z}^ε , and $\boldsymbol{\mu}^\varepsilon$ to \mathbf{u} , \mathbf{Z} , and $\boldsymbol{\mu}$, respectively (see Step 1 in the proof of Theorem 3.6). The functions \mathbf{u} and \mathbf{Z} can be seen to satisfy, for a.e. $t \in [0, T]$,

$$\mathbf{u}_t(t) - \operatorname{div} \mathbf{Z}(t) = \boldsymbol{\mu}(t) \quad \text{in } \mathcal{M}(\Omega; \mathbb{R}^N).$$

Then we show, by a relatively soft argument, which nevertheless requires quite a few preliminaries, that

$$\boldsymbol{\mu} = *(*(\mathbf{Z} \wedge \mathbf{u}) \wedge D\mathbf{u}) \quad \text{and} \quad |\boldsymbol{\mu}(t)| \leq |D\mathbf{u}(t)| \quad \text{as measures for a.e. } t \in [0, T] \tag{1-7}$$

(see Step 2 in the proof of Theorem 3.6). Hence, in order to identify μ it suffices to show that

$$\mathbf{u}(t)_g \cdot \frac{\mu(t)}{|D\mathbf{u}(t)|} \geq 1 \quad \text{for a.e. } t \in [0, T], \quad (1-8)$$

where $\mu(t)/|D\mathbf{u}(t)|$ denotes the Radon–Nikodým derivative of $\mu(t)$ with respect to $|D\mathbf{u}(t)|$. Indeed, (1-7) and simple vectorial identities then imply that

$$\mu(t) = \mathbf{u}(t)_g |D\mathbf{u}(t)| \quad \text{for a.e. } t \in [0, T]$$

(see Step 6 in the proof of Theorem 3.6). In view of (1-6), the lower bound (1-8) for the diffuse part of μ follows (see Step 4 in the proof of Theorem 3.6) from a suitable modification of a relaxation result [Alicandro et al. 2007], applied to each of the components of

$$\mathcal{F}(\mathbf{v}) := \int_{\Omega} \mathbf{v}(x) |\nabla \mathbf{v}(x)| dx$$

(see Section 2F). On the other hand, the same argument would lead to a suboptimal lower bound on $\mu(t)$ over the jump set of $\mathbf{u}(t)$ (see Remark 3.10). Moreover, the results in [Alicandro et al. 2007] can not be directly applied to $\mathbf{u}(t)_g \cdot \mu^\varepsilon(t)$, since $\mathbf{u}(t)_g$ is a discontinuous function (though a very special one). For these reasons, we revisit the blow-up argument in [Fonseca and Müller 1993] and the dimensional reduction argument in [Fonseca and Rybka 1992] to conclude that

$$\mathbf{u}(t)_g \cdot \frac{\mu(t)}{|D\mathbf{u}(t)|} \geq \frac{1}{|\mathbf{u}(t)_+(x) - \mathbf{u}(t)_-(x)|} \inf_{\boldsymbol{\gamma} \in \tilde{\Gamma}_N} \int_0^1 \mathbf{u}(t)_g(x) \cdot \boldsymbol{\gamma}(s) |\boldsymbol{\gamma}'(s)| ds \quad (1-9)$$

for a.e. t and \mathcal{H}^{m-1} -a.e. $x \in J_{\mathbf{u}(t)}$, where

$$\tilde{\Gamma}_N := \{\boldsymbol{\gamma} \in W^{1,1}((0, 1); \mathbb{S}_+^{N-1}) : \boldsymbol{\gamma}(0) = \mathbf{u}(t)_-(x), \boldsymbol{\gamma}(1) = \mathbf{u}(t)_+(x)\} \quad (1-10)$$

(see Step 5 in the proof of Theorem 3.6). The minimization problem which appears on the right-hand side of (1-9) is crucial in our argument. In Section 4 we argue that

$$\min_{\boldsymbol{\gamma} \in \Gamma} \int_0^1 \mathbf{u}_g \cdot \boldsymbol{\gamma}(s) |\boldsymbol{\gamma}'(s)| ds = |\mathbf{u}_+ - \mathbf{u}_-|, \quad \text{where } \Gamma = \{\boldsymbol{\gamma} \in W^{1,1}((0, 1); \mathbb{S}_+^{N-1}) : \boldsymbol{\gamma}(0) = \mathbf{u}_-, \boldsymbol{\gamma}(1) = \mathbf{u}_+\} \quad (1-11)$$

(see Theorem 4.1). Together with (1-9), (1-11) yields the lower bound (1-8) on the jump set of $\mathbf{u}(t)$ too.

The minimization problem in (1-11) is equivalent to finding—and characterizing the length of—shortest paths between \mathbf{u}_- and \mathbf{u}_+ in a Riemannian manifold with boundary whose metric penalizes the closeness to \mathbf{u}_g . In addition, the metric may degenerate at a point of the manifold: for instance, if $N = 3$, $\mathbf{u}_- = (0, 0, 1)$, and $\mathbf{u}_+ = (0, 1, 0)$, then $\mathbf{u}_g \cdot (1, 0, 0) = 0$. In these respects, the minimization problem has a geometrical interest of its own.

It turns out that the minimum in (1-11) is achieved by the standard geodesic on \mathbb{S}_+^{N-1} connecting \mathbf{u}_- and \mathbf{u}_+ ; see Lemma 4.2. Nevertheless, the analysis of (1-11) is highly nontrivial for two reasons. Firstly, one has to characterize the length of candidate shortest paths which may in principle intersect and/or de-touch from the boundary of the manifold. Secondly, the functional in (1-11) is genuinely nonconvex:

indeed, besides the aforementioned standard geodesic, it always possesses a second smooth critical point, which we show not to be a shortest path. In addition, in the extreme cases in which \mathbf{u}_+ and \mathbf{u}_- are two distinct “vertices” of \mathbb{S}_+^{N-1} , the functional in (1-11) possesses a second shortest path which is not a critical point: it follows the boundary of \mathbb{S}_+^{N-1} and passes through the point of degeneracy. For instance, if $N = 3$, $\mathbf{u}_- = (0, 0, 1)$, and $\mathbf{u}_+ = (0, 1, 0)$, then $\mathbf{u}_g = (0, 1, 1)/\sqrt{2}$ and the curve

$$\boldsymbol{\gamma}(s) = \begin{cases} (\sin(\pi s), 0, \cos(\pi s)) & \text{if } s \in [0, 1/2], \\ (\sin(\pi s), -\cos(\pi s), 0) & \text{if } s \in (1/2, 1] \end{cases}$$

is such that

$$\int_0^1 \mathbf{u}_g \cdot \boldsymbol{\gamma}(s) |\boldsymbol{\gamma}'(s)| ds = 2 \int_0^{1/2} \frac{\cos(\pi s)}{\sqrt{2}} \pi ds = \sqrt{2} = |\mathbf{u}_+ - \mathbf{u}_-|.$$

Finally, we note that if the paths in Γ are allowed to take values in a set \mathbb{A} which contains \mathbb{S}_+^{N-1} , then in general the standard geodesic is not a minimizer and (1-11) does not hold; an example is given in Remark 4.4.

The paper is organized as follows. In Section 2 we collect the definitions and results which we need concerning multivector fields, functions of bounded variations, a generalized Green’s formula, tensor fields, and lower semicontinuity of integral functionals. In Section 3 we introduce the concept of and prove the existence of a solution to (1-4). Section 4 is devoted to the minimization problem in (1-11).

2. Preliminaries

In this section we introduce some notation and some preliminary results that we need in the sequel.

General notations. Throughout this paper \mathcal{H}^{m-1} denotes the $(m - 1)$ -dimensional Hausdorff measure and \mathcal{L}^m the m -dimensional Lebesgue measure. We denote by $\mathcal{M}(\Omega; \mathbb{R}^N)$ the space of \mathbb{R}^N -valued finite Radon measures on Ω ; see [Ambrosio et al. 2000, Definition 1.40]. We recall that $\mathcal{M}(\Omega; \mathbb{R}^N)$ is the dual space of $C_0(\Omega; \mathbb{R}^N)$. Throughout, the subscript $_0$ denotes spaces of compactly supported functions. We denote $\mathcal{D}(\Omega; \mathbb{R}^N) := C_0^\infty(\Omega; \mathbb{R}^N)$. When $N = 1$, we often do not specify the target space (for example, $\mathcal{M}(\Omega) = \mathcal{M}(\Omega; \mathbb{R})$). Finally, if $\mathbb{A} \subset \mathbb{R}^N$ is compact and $\Upsilon(\Omega; \mathbb{R}^N)$ is a space of functions, we sometimes use the notation $\Upsilon(\Omega; \mathbb{A}) := \{\mathbf{u} \in \Upsilon(\Omega; \mathbb{R}^N) : \mathbf{u}(x) \in \mathbb{A} \text{ for } \mathcal{L}^m\text{-a.e. } x \in \Omega\}$.

2A. Multivectors. Here we recall some definitions and basic properties about multivectors that we need in our analysis. We refer to, for example, [Federer 1969, Chapter 1; Darling 1994, Chapter 1] for details.

The spaces $\Lambda_0(\mathbb{R}^N)$ and $\Lambda_1(\mathbb{R}^N)$ coincide with \mathbb{R} and \mathbb{R}^N , respectively. For $2 \leq k \leq \mathbb{N}$, the k -th exterior power of \mathbb{R}^N , denoted by $\Lambda_k(\mathbb{R}^N)$, is a set spanned by elements of the form

$$\mathbf{u}_1 \wedge \cdots \wedge \mathbf{u}_k, \quad \mathbf{u}_i \in \mathbb{R}^N, \quad i = 1, \dots, k$$

(elements of this form are called “generators”) and subject to the following rules:

- (1) $(a\mathbf{v} + b\mathbf{w}) \wedge \mathbf{u}_2 \wedge \cdots \wedge \mathbf{u}_k = a(\mathbf{v} \wedge \mathbf{u}_2 \wedge \cdots \wedge \mathbf{u}_k) + b(\mathbf{w} \wedge \mathbf{u}_2 \wedge \cdots \wedge \mathbf{u}_k)$.
- (2) $\mathbf{u}_1 \wedge \cdots \wedge \mathbf{u}_k$ changes sign if two entries are transposed.

(3) For any basis $\{e_1, \dots, e_n\}$ of \mathbb{R}^N , $\{e_\alpha := e_{\alpha_1} \wedge \dots \wedge e_{\alpha_k} : \alpha \in I(k, N)\}$ is a basis for $\Lambda_k(\mathbb{R}^N)$.

Here we have used the standard notation for ordered multiindexes:

$$I(k, N) = \{\alpha = (\alpha_1, \dots, \alpha_k) : \alpha_i \in \mathbf{Z}, 1 \leq \alpha_1 < \dots < \alpha_k \leq N\}. \quad (2-1)$$

The elements of $\Lambda_k(\mathbb{R}^N)$ are called multivectors (or k -vectors), and $\Lambda_k(\mathbb{R}^N)$ is a vector space of dimension $\binom{N}{k}$. We will use the well-known equality [Darling 1994, Formula 1.68]

$$|\mathbf{a}|^2 |\mathbf{b}|^2 = (\mathbf{a} \cdot \mathbf{b})^2 + (\mathbf{a} \wedge \mathbf{b})^2 \quad \text{for all } \mathbf{a}, \mathbf{b} \in \mathbb{R}^N. \quad (2-2)$$

Given $k, p \in \{0, \dots, N\}$ with $k + p \leq N$, there exists a unique bilinear map $(\boldsymbol{\lambda}, \boldsymbol{\mu}) \rightarrow \boldsymbol{\lambda} \wedge \boldsymbol{\mu}$ from $\Lambda_k(\mathbb{R}^N) \times \Lambda_p(\mathbb{R}^N)$ to $\Lambda_{k+p}(\mathbb{R}^N)$, whose effect on generators is

$$(\mathbf{u}_1 \wedge \mathbf{u}_2 \wedge \dots \wedge \mathbf{u}_k) \wedge (\mathbf{v}_1 \wedge \mathbf{v}_2 \wedge \dots \wedge \mathbf{v}_p) = \mathbf{u}_1 \wedge \mathbf{u}_2 \wedge \dots \wedge \mathbf{u}_k \wedge \mathbf{v}_1 \wedge \mathbf{v}_2 \wedge \dots \wedge \mathbf{v}_p.$$

This map satisfies

$$\boldsymbol{\lambda} \wedge \boldsymbol{\mu} = (-1)^{-kp} (\boldsymbol{\mu} \wedge \boldsymbol{\lambda}) \quad \text{for } \boldsymbol{\lambda} \in \Lambda_k(\mathbb{R}^N), \boldsymbol{\mu} \in \Lambda_p(\mathbb{R}^N). \quad (2-3)$$

The *Hodge-star operator* is an isomorphism from $\Lambda_k(\mathbb{R}^N)$ to $\Lambda_{N-k}(\mathbb{R}^N)$, defined on the basis as

$$*(e_{\alpha_1} \wedge \dots \wedge e_{\alpha_k}) := e_{\alpha_{k+1}} \wedge \dots \wedge e_{\alpha_N}, \quad (2-4)$$

where $\{\alpha_1, \dots, \alpha_N\}$ has positive signature. In particular, in what follows we will systematically identify $\Lambda_{N-1}(\mathbb{R}^N)$ with \mathbb{R}^N . We will use the following well-known formulas:

$$*(*\boldsymbol{\lambda}) = (-1)^{k(N-k)} \boldsymbol{\lambda} \quad \text{for all } \boldsymbol{\lambda} \in \Lambda_k(\mathbb{R}^N) \quad (2-5)$$

(see, for example, [Darling 1994, (1.64)]) and

$$\mathbf{a} \wedge *(b \wedge c) = (\mathbf{a} \cdot c) *b - (\mathbf{a} \cdot b) *c \quad \text{for all } \mathbf{a}, \mathbf{b}, \mathbf{c} \in \mathbb{R}^N \quad (2-6)$$

(see, for example, [Darling 1994, Table 1.2]). It follows from (2-3), (2-6), and (2-5) that

$$|\mathbf{b}|^2 \mathbf{a} = (\mathbf{a} \cdot \mathbf{b}) \mathbf{b} - *(*(\mathbf{a} \wedge \mathbf{b}) \wedge \mathbf{b}) \quad \text{for all } \mathbf{a}, \mathbf{b} \in \mathbb{R}^N. \quad (2-7)$$

Introducing the norm

$$|\boldsymbol{\lambda}|_k = \left(\sum_{\alpha \in I(k, N)} |\lambda_\alpha|^2 \right)^{1/2}, \quad \text{where } \boldsymbol{\lambda} = \sum_{\alpha \in I(k, N)} \lambda_\alpha e_\alpha \quad (2-8)$$

and using (2-4), it is immediate to see that

$$|*\boldsymbol{\lambda}|_{N-k} = |\boldsymbol{\lambda}|_k \quad \text{for any } \boldsymbol{\lambda} \in \Lambda_k(\mathbb{R}^N). \quad (2-9)$$

Finally, we recall that, given $\boldsymbol{\lambda} \in \Lambda_k(\mathbb{R}^N)$ and $\boldsymbol{\eta} \in \Lambda_p(\mathbb{R}^N)$ such that one of them is a generator, we have

$$|\boldsymbol{\lambda} \wedge \boldsymbol{\eta}|_{k+p} \leq |\boldsymbol{\lambda}|_k |\boldsymbol{\eta}|_p; \quad (2-10)$$

see [Federer 1969, p. 32].

2B. Vector-valued functions. Let $(X, \|\cdot\|)$ be a Banach space with dual X' and let $U \subset \mathbb{R}^d$ be a bounded open set endowed with the Lebesgue measure \mathcal{L}^d . We denote by $\langle \cdot, \cdot \rangle$ the pairing between X and X' . A function $u : U \rightarrow X$ is called *simple* if there exist $x_1, \dots, x_n \in X$ and $U_1, \dots, U_n \mathcal{L}^m$ -measurable subsets of U such that $u = \sum_{i=1}^n x_i \chi_{U_i}$. The function u is called *strongly measurable* if there exists a sequence of simple functions $\{u_n\}$ such that $\|u_n(x) - u(x)\| \rightarrow 0$ as $n \rightarrow +\infty$ for almost all $x \in U$. If $1 \leq p < \infty$, then $L^p(U; X)$ stands for the space of (equivalence classes of) strongly measurable functions $u : U \rightarrow X$ with

$$\|u\|_p := \left(\int_U \|u(x)\|^p dx \right)^{1/p} < \infty.$$

Endowed with this norm, $L^p(U; X)$ is a Banach space. For $p = \infty$, the symbol $L^\infty(U; X)$ stands for the space of (equivalence classes of) strongly measurable functions $u : U \rightarrow X$ such that

$$\|u\|_\infty := \text{esssup}\{\|u(x)\| : x \in U\} < \infty.$$

If $U = (0, T)$, we write $L^p(0, T; X) = L^p((0, T); X)$. For $1 \leq p < \infty$, $L^{p'}(0, T; X')$ ($1/p + 1/p' = 1$) is isometric to a subspace of $(L^p(0, T; X))'$, with equality if and only if X' has the Radon–Nikodým property; see, for instance, [Diestel and Uhl 1977].

We consider the vector space $\mathcal{D}(U; X) := C_0^\infty(U; X)$, endowed with the topology for which a sequence $\varphi_n \rightarrow 0$ as $n \rightarrow +\infty$ if there exists $K \subset U$ compact such that $\text{supp}(\varphi_n) \subset K$ for any $n \in \mathbb{N}$ and $D^\alpha \varphi_n \rightarrow 0$ uniformly on K as $n \rightarrow +\infty$ for all multiindexes α . We denote by $\mathcal{D}'(U; X)$ the space of distributions on U with values in X , that is, the set of all linear continuous maps $T : \mathcal{D}(U; X) \rightarrow \mathbb{R}$. As is well known, $L^p(U; X) \subset \mathcal{D}'(U; X)$ through the standard continuous injection. Given $T \in \mathcal{D}'(U; X)$, the distributional derivative of T is defined by

$$\langle D_i T, \varphi \rangle := -\langle T, \partial_i \varphi \rangle \quad \text{for any } \varphi \in \mathcal{D}(U; X) \text{ and any } i \in \{1, \dots, d\}. \tag{2-11}$$

General notations for matrices. If $\mathbf{A} = (a_i^\ell)$ is an $N \times m$ matrix, we write $a^\ell = (a_1^\ell, \dots, a_m^\ell)$ for $1 \leq \ell \leq N$ and $\mathbf{a}_i = (a_i^1, \dots, a_i^N)$ for $1 \leq i \leq m$. If $\mathbf{B} = (b_i^\ell)$ is also an $N \times m$ matrix, we let

$$\mathbf{A} : \mathbf{B} = \sum_{\ell=1}^N \sum_{i=1}^m a_i^\ell b_i^\ell \quad \text{and} \quad |\mathbf{A}| = (\mathbf{A} : \mathbf{A})^{1/2} = \left(\sum_{\ell=1}^N \sum_{i=1}^m (a_i^\ell)^2 \right)^{1/2}.$$

Given $\mathbf{A} = (\mathbf{a}_1, \dots, \mathbf{a}_m) \in \mathbb{R}^{N \times m}$ and $\mathbf{b} \in \mathbb{R}^N$, we let

$$\begin{aligned} \mathbf{A} \wedge \mathbf{b} &:= (\mathbf{a}_1 \wedge \mathbf{b}, \dots, \mathbf{a}_m \wedge \mathbf{b}), \\ *(\mathbf{A} \wedge \mathbf{b}) &:= (*(\mathbf{a}_1 \wedge \mathbf{b}), \dots, *(\mathbf{a}_m \wedge \mathbf{b})). \end{aligned}$$

2C. Functions of bounded variation. A vector-field $\mathbf{u} \in L^1(\Omega; \mathbb{R}^N)$ has bounded variation, and we write $\mathbf{u} \in \text{BV}(\Omega; \mathbb{R}^N)$, if there is an $N \times m$ matrix $D\mathbf{u}$, whose components $D_i u^\ell$ are finite Radon measures, such that

$$\sum_{\ell=1}^N \int_{\Omega} u^{\ell} \operatorname{div} \varphi^{\ell} dx = - \sum_{\ell=1}^N \sum_{i=1}^m \int_{\Omega} \varphi_i^{\ell} dD_i u^{\ell} \quad \text{for all } \varphi \in (C_0^1(\Omega; \mathbb{R}^N))^m.$$

Its variation measure $|D\mathbf{u}|$ is a finite Radon measure defined on open sets $U \subseteq \Omega$ by

$$|D\mathbf{u}|(U) = \sup \left\{ \sum_{\ell=1}^N \int_U u^{\ell} \operatorname{div} \varphi^{\ell} dx : \varphi \in (C_0^1(U; \mathbb{R}^N))^m, \|\varphi\|_{\infty} \leq 1 \right\}.$$

The matrix-valued Radon measure $D\mathbf{u}$ is decomposed into three mutually orthogonal measures (see [Ambrosio et al. 2000; Evans and Gariepy 1992; Ziemer 1989]):

$$D\mathbf{u} = \nabla \mathbf{u} \mathcal{L}^m + D^c \mathbf{u} + D^j \mathbf{u},$$

where $\nabla \mathbf{u}$ denotes the Radon–Nikodým derivative of $D\mathbf{u}$ with respect to \mathcal{L}^m . The *Cantor part* $D^c \mathbf{u}$ is supported on the set of *Lebesgue points of \mathbf{u}* , $\Omega \setminus S_{\mathbf{u}}$, that is, those points $x \in \Omega$ for which there exists $\tilde{\mathbf{u}}(x) \in \mathbb{R}^N$ such that

$$\lim_{\rho \downarrow 0} \frac{1}{\mathcal{L}^m(B_{\rho}(x))} \int_{B_{\rho}(x)} |\mathbf{u}(y) - \tilde{\mathbf{u}}(x)| dy = 0.$$

The *jump part* $D^j \mathbf{u}$ is supported on the set of *approximate jump points of \mathbf{u}* , $J_{\mathbf{u}}$, that is, those points $x \in \Omega$ for which there exist $\mathbf{u}_{+}(x) \neq \mathbf{u}_{-}(x) \in \mathbb{R}^N$ and $\nu_{\mathbf{u}}(x) \in \mathbb{S}^{m-1}$ such that

$$\lim_{\rho \downarrow 0} \frac{1}{\mathcal{L}^m(B_{\rho}^{\pm}(x, \nu_{\mathbf{u}}(x)))} \int_{B_{\rho}^{\pm}(x, \nu_{\mathbf{u}}(x))} |\mathbf{u}(y) - \mathbf{u}_{\pm}(x)| dy = 0,$$

where

$$B_{\rho}^{\pm}(x, \nu_{\mathbf{u}}(x)) = \{y \in B_{\rho}(x) : \langle y - x, \nu_{\mathbf{u}}(x) \rangle \gtrless 0\}.$$

The jump set $J_{\mathbf{u}}$ is a Borel subset of $S_{\mathbf{u}}$ that satisfies $\mathcal{H}^{m-1}(S_{\mathbf{u}} \setminus J_{\mathbf{u}}) = 0$. The *precise representative* $\mathbf{u}^* : \Omega \setminus (S_{\mathbf{u}} \setminus J_{\mathbf{u}}) \rightarrow \mathbb{R}^N$ of \mathbf{u} is defined to be equal to $\tilde{\mathbf{u}}$ on $\Omega \setminus S_{\mathbf{u}}$ and equal to $(\mathbf{u}_{-} + \mathbf{u}_{+})/2$ on $J_{\mathbf{u}}$. In what follows, we identify $\mathbf{u} = \tilde{\mathbf{u}} = \mathbf{u}^*$ on $\Omega \setminus S_{\mathbf{u}}$.

2D. A generalized Green's formula. Let

$$X_{\mathcal{M}}(\Omega) = \{z \in L^{\infty}(\Omega; \mathbb{R}^m) : \operatorname{div} z \in \mathcal{M}(\Omega)\}$$

and

$$\mathcal{M}_{\mathcal{H}}(\Omega; \mathbb{R}^N) := \{\boldsymbol{\mu} \in \mathcal{M}(\Omega; \mathbb{R}^N) : |\boldsymbol{\mu}|(B) = 0 \text{ for any Borel set } B \subset \Omega : \mathcal{H}^{m-1}(B) = 0\}.$$

In [Anzellotti 1983, Theorem 1.2] (see also [Andreu-Vaillo et al. 2004; Chen and Frid 1999]), the weak trace on $\partial\Omega$ of the normal component of $z \in X_{\mathcal{M}}(\Omega)$ is defined. Namely, it is proved that there exists a linear operator $[\cdot, \nu] : X_{\mathcal{M}}(\Omega) \rightarrow L^{\infty}(\partial\Omega)$ such that $\|[z, \nu]\|_{L^{\infty}(\partial\Omega)} \leq \|z\|_{L^{\infty}(\Omega)}$ for all $z \in X_{\mathcal{M}}(\Omega)$ and $[z, \nu]$ coincides with the pointwise trace of the normal component if z is smooth:

$$[z, \nu](x) = z(x) \cdot \nu(x) \quad \text{for all } x \in \partial\Omega \text{ if } z \in C^1(\bar{\Omega}, \mathbb{R}^m).$$

It follows from [Chen and Frid 1999, Proposition 3.1] or [Ambrosio et al. 2005, Proposition 3.4] that

$$\operatorname{div} z \in \mathcal{M}_{\mathcal{H}}(\Omega) \quad \text{for all } z \in X_{\mathcal{M}}(\Omega). \tag{2-12}$$

Therefore, given $z \in X_{\mathcal{M}}(\Omega)$ and $u \in BV(\Omega) \cap L^\infty(\Omega)$, the functional $(z, Du) \in \mathcal{D}'(\Omega)$ given by

$$\langle (z, Du), \varphi \rangle := - \int_{\Omega} u^* \varphi \, d(\operatorname{div} z) - \int_{\Omega} uz \nabla \varphi \, dx \tag{2-13}$$

is well defined, and the following holds (in [Caselles 2011], see Lemma 5.1, Theorem 5.3, and the discussion after Lemma 5.4):

Lemma 2.1. *Let $z \in X_{\mathcal{M}}(\Omega)$ and $u \in BV(\Omega) \cap L^\infty(\Omega)$. Then the functional $(z, Du) \in \mathcal{D}'(\Omega)$ defined by (2-13) is a Radon measure which is absolutely continuous with respect to $|Du|$. Furthermore,*

$$\int_{\Omega} u^* \, d(\operatorname{div} z) + (z, Du)(\Omega) = \int_{\partial\Omega} [z, \nu] u \, d\mathcal{H}^{m-1}$$

and

$$\operatorname{div}(zu) = u^* \operatorname{div} z + (z, Du) \quad \text{as measures.}$$

We will use the vector-valued version of Lemma 2.1. To this aim, we introduce the space

$$X_{\mathcal{M}}^N(\Omega) = \{ \mathbf{Z} = (z^1, \dots, z^N)^T : z^\ell \in X_{\mathcal{M}}(\Omega) \text{ for } \ell = 1, \dots, N \}.$$

Given $\mathbf{Z} \in X_{\mathcal{M}}^N(\Omega)$ and $\mathbf{u} \in BV(\Omega; \mathbb{R}^N) \cap L^\infty(\Omega; \mathbb{R}^N)$, we use the notation

$$\begin{aligned} \operatorname{div} \mathbf{Z} &:= (\operatorname{div} z^1, \dots, \operatorname{div} z^N), \\ [\mathbf{Z}, \nu] &:= ([z^1, \nu], \dots, [z^N, \nu]), \\ \mathbf{Z} : D\mathbf{u} &:= \sum_{\ell=1}^N (z^\ell, Du^\ell). \end{aligned}$$

Then, as an immediate consequence of (2-12) and Lemma 2.1, we have:

Corollary 2.2. *Let $\mathbf{Z} \in X_{\mathcal{M}}^N(\Omega)$. Then*

$$\operatorname{div} \mathbf{Z} \in \mathcal{M}_{\mathcal{H}}(\Omega; \mathbb{R}^N).$$

Furthermore, for any $\mathbf{u} \in BV(\Omega; \mathbb{R}^N) \cap L^\infty(\Omega; \mathbb{R}^N)$, $\mathbf{Z} : D\mathbf{u}$ is a Radon measure which is absolutely continuous with respect to $|D\mathbf{u}|$,

$$\int_{\Omega} \mathbf{u}^* \cdot d(\operatorname{div} \mathbf{Z}) + (\mathbf{Z} : D\mathbf{u})(\Omega) = \int_{\partial\Omega} [\mathbf{Z}, \nu] \cdot \mathbf{u} \, d\mathcal{H}^{m-1} \tag{2-14}$$

and

$$\operatorname{div}(\mathbf{Z}^T \mathbf{u}) = \mathbf{u}^* \cdot \operatorname{div} \mathbf{Z} + \mathbf{Z} : D\mathbf{u} \quad \text{as measures.} \tag{2-15}$$

2E. Multivector fields. Let $U \subset \mathbb{R}^d$. A multivector distribution in U is a linear continuous map $\lambda \in \mathcal{D}'(U; \Lambda_k(\mathbb{R}^N))$ (see Section 2B). It may be expressed in terms of the basis (3) as

$$\lambda = \sum_{\alpha \in I(k, N)} \lambda_\alpha \mathbf{e}_\alpha \quad \text{with } \lambda_\alpha \in \mathcal{D}'(U; \mathbb{R}^N) \text{ for any } \alpha \in I(k, N).$$

Then, according to (2-11),

$$D_i \lambda = \sum_{\alpha \in I(k, N)} D_i \lambda_\alpha \mathbf{e}_\alpha \quad \text{for any } i \in \{1, \dots, d\}. \quad (2-16)$$

From (2-16), the following two identities are easily seen to hold for $k, p \in \mathbb{N}$ and $i \in \{1, \dots, d\}$:

$$D_i(\lambda \wedge \eta) = D_i \lambda \wedge \eta + \lambda \wedge D_i \eta \quad (2-17)$$

for any $\lambda \in L^2(U; \Lambda_k(\mathbb{R}^N))$ such that $D_i \lambda \in L^2(U; \Lambda_k(\mathbb{R}^N))$ and any $\eta \in L^2(U; \Lambda_p(\mathbb{R}^N))$ such that $D_i \eta \in L^2(U; \Lambda_p(\mathbb{R}^N))$;

$$*(D_i \lambda) = D_i(*\lambda) \quad \text{for any } \lambda \in \mathcal{D}'(U; \Lambda_k(\mathbb{R}^N)). \quad (2-18)$$

For any $k \in \mathbb{N}$, $(\Lambda_k(\mathbb{R}^N))^m$ is a Banach space. We use the norm

$$\|\mathcal{A}\| := \left(\sum_{i=1}^m |\mathcal{A}_i|_k^2 \right)^{1/2} \quad \text{for } \mathcal{A} = (\mathcal{A}_1, \dots, \mathcal{A}_m)$$

with $|\cdot|_k$ given by (2-8).

We will now state and prove the analogue of Corollary 2.2 for a multivector field

$$\mathcal{A} = (\mathcal{A}_1, \dots, \mathcal{A}_m) \in L^\infty(\Omega; (\Lambda_k(\mathbb{R}^N))^m).$$

We define

$$\operatorname{div} \mathcal{A} := \sum_{i=1}^m D_i(\mathcal{A}_i). \quad (2-19)$$

Square-integrability of $\operatorname{div} \mathcal{A}$ suffices for our purposes. Hence, we introduce the space

$$X_2(\Omega; \Lambda_{N-2}(\mathbb{R}^N)) := \{\mathcal{A} \in L^\infty(\Omega; (\Lambda_{N-2}(\mathbb{R}^N))^m) : \operatorname{div} \mathcal{A} \in L^2(\Omega; \Lambda_{N-2}(\mathbb{R}^N))\}.$$

The following holds:

Lemma 2.3. *Let $\mathcal{A} \in X_2(\Omega; \Lambda_{N-2}(\mathbb{R}^N))$ and consider $\mathbf{u} \in \operatorname{BV}(\Omega; \mathbb{R}^N) \cap L^2(\Omega; \mathbb{R}^N)$. Then the functional $*(\mathcal{A} \wedge D\mathbf{u}) : \mathcal{D}(\Omega; \mathbb{R}^N) \rightarrow \mathbb{R}$ defined by*

$$\langle *(\mathcal{A} \wedge D\mathbf{u}), \Phi \rangle := - \int_{\Omega} *(\operatorname{div} \mathcal{A} \wedge \mathbf{u}) \cdot \Phi \, dx - \sum_{i=1}^m \int_{\Omega} *(\mathcal{A}_i \wedge \mathbf{u}) \cdot \partial_i \Phi \, dx \quad (2-20)$$

is an \mathbb{R}^N -valued Radon measure on Ω , absolutely continuous with respect to $|D\mathbf{u}|$, with

$$|*(\mathcal{A} \wedge D\mathbf{u})|(B) \leq \|\mathcal{A}\|_\infty |D\mathbf{u}|(B) \quad \text{for any Borel set } B \subseteq \Omega. \quad (2-21)$$

Furthermore,

$$\operatorname{div}(*(\mathcal{A} \wedge \mathbf{u})) = *(\mathcal{A} \wedge D\mathbf{u}) + *(\operatorname{div} \mathcal{A} \wedge \mathbf{u}) \mathcal{L}^m \quad \text{as measures.} \tag{2-22}$$

Proof. Since Ω has compact Lipschitz boundary, it follows from [Ambrosio et al. 2000, Theorem 3.21, Remark 3.22, and Corollary 3.80] that the sequence $\mathbf{u}_n := (T\mathbf{u}) \star \rho_n \in C^\infty(\bar{\Omega})$ (here T denotes an extension operator) is such that $\mathbf{u}_n \rightharpoonup \mathbf{u}$ in $\operatorname{BV}(\Omega; \mathbb{R}^N)$, $\int_\Omega |\nabla \mathbf{u}_n| \, dx \rightarrow |D\mathbf{u}|(\Omega)$, and $\mathbf{u}_n \rightarrow \mathbf{u}^*$ \mathcal{H}^{m-1} -a.e. in $\bar{\Omega}$. Furthermore, by construction and since $\mathbf{u} \in L^2(\Omega; \mathbb{R}^N)$, we have $\mathbf{u}_n \rightarrow \mathbf{u}$ in $L^2(\Omega; \mathbb{R}^N)$. Then

$$\langle *(\mathcal{A} \wedge D\mathbf{u}), \Phi \rangle \stackrel{(2-20)}{=} - \lim_{n \rightarrow \infty} \left(\int_\Omega *(\operatorname{div} \mathcal{A} \wedge \mathbf{u}_n) \cdot \Phi \, dx + \sum_{i=1}^m \int_\Omega *(\mathcal{A}_i \wedge \mathbf{u}_n) \cdot \partial_i \Phi \, dx \right).$$

Integrating by parts and using (2-18), we obtain

$$\begin{aligned} \langle *(\mathcal{A} \wedge D\mathbf{u}), \Phi \rangle &= - \lim_{n \rightarrow \infty} \left(\int_\Omega *(\operatorname{div} \mathcal{A} \wedge \mathbf{u}_n) \cdot \Phi \, dx - \sum_{i=1}^m \int_\Omega *(\partial_i(\mathcal{A}_i \wedge \mathbf{u}_n)) \cdot \Phi \, dx \right) \\ &\stackrel{(2-17)}{=} \lim_{n \rightarrow \infty} \sum_{i=1}^m \int_\Omega *(\mathcal{A}_i \wedge \partial_i \mathbf{u}_n) \cdot \Phi \, dx. \end{aligned}$$

Therefore, applying the Hölder and Cauchy–Schwarz inequalities,

$$\begin{aligned} |\langle *(\mathcal{A} \wedge D\mathbf{u}), \Phi \rangle| &\stackrel{(2-9)}{\leq} \|\Phi\|_\infty \lim_{n \rightarrow \infty} \sum_{i=1}^m \int_\Omega |\mathcal{A}_i \wedge \partial_i \mathbf{u}_n|_{N-1} \, dx \\ &\stackrel{(2-10)}{\leq} \|\Phi\|_\infty \lim_{n \rightarrow \infty} \int_\Omega \sum_{i=1}^m |\partial_i \mathbf{u}_n| |\mathcal{A}_i|_{N-2} \, dx \\ &\leq \|\Phi\|_\infty \lim_{n \rightarrow \infty} \int_\Omega |\nabla \mathbf{u}_n| \left(\sum_{i=1}^m |\mathcal{A}_i|_{N-2}^2 \right)^{1/2} \, dx \\ &\leq \|\Phi\|_\infty \|\mathcal{A}\|_\infty \lim_{n \rightarrow \infty} \int_\Omega |\nabla \mathbf{u}_n| \, dx \\ &= \|\Phi\|_\infty \|\mathcal{A}\|_\infty |D\mathbf{u}|(\Omega). \end{aligned}$$

The arbitrariness of Φ completes the proof of (2-21). It follows from (2-20) and (2-19) that $\operatorname{div}(*(\mathcal{A} \wedge \mathbf{u}))$ is also an \mathbb{R}^N -valued Radon measure in Ω , and (2-22) follows from (2-19). \square

2F. Lower semicontinuity of integral functionals over $W^{1,1}(\Omega; \mathbb{S}_+^{N-1})$. Let $f : \Omega \times \mathbb{S}_+^{N-1} \rightarrow \mathbb{R}_+$ and consider the energy functional defined in $L^1(\Omega; \mathbb{S}_+^{N-1})$ by

$$\mathcal{F}_f(\mathbf{v}) := \begin{cases} \int_\Omega f(x, \mathbf{v}(x)) |\nabla \mathbf{v}(x)| \, dx & \text{if } \mathbf{v} \in W^{1,1}(\Omega; \mathbb{S}_+^{N-1}), \\ +\infty & \text{otherwise.} \end{cases}$$

The purpose of this section is to restate, to the extent we need in the present setting, a few lower semicontinuity results obtained in [Fonseca and Rybka 1992; Alicandro et al. 2007] (see also [Giaquinta

and Mucci 2006] for related results when the target space is a general manifold). We consider the following hypotheses for f :

(H1) f is continuous and nonnegative;

(H2) (uniform boundedness) a positive constant C_1 exists such that

$$|f(x, \mathbf{s})| \leq C_1 \quad \text{for all } (x, \mathbf{s}) \in \Omega \times \mathbb{S}_+^{N-1};$$

(H3) for every compact set $U \subset \Omega$, there exist a continuous function ω , with $\omega(0) = 0$, such that

$$|f(x, \mathbf{s}) - f(x', \mathbf{s}')| = \omega(|x - x'| + |\mathbf{s} - \mathbf{s}'|) \quad \text{for all } (x, \mathbf{s}), (x', \mathbf{s}') \in U \times \mathbb{S}_+^{N-1}.$$

For $\varrho \in \mathbb{R}^m$ such that $|\varrho| = 1$, we define $Q_\varrho := R_\varrho[-\frac{1}{2}, \frac{1}{2}]^m$, where R_ϱ denotes a rotation such that $R_\varrho e_m = \varrho$. Given $\mathbf{a}, \mathbf{b} \in \mathbb{S}_+^{N-1}$, we set

$$K_f(x, \mathbf{a}, \mathbf{b}, \varrho) := \inf \left\{ \int_{Q_\varrho} f(x, \mathbf{v}(y)) |\nabla \mathbf{v}(y)| dy : \mathbf{v} \in \mathcal{P}(\mathbf{a}, \mathbf{b}, \varrho) \right\}, \quad (2-23)$$

where

$$\mathcal{P}(\mathbf{a}, \mathbf{b}, \varrho) := \left\{ \mathbf{v} \in W^{1,1}(Q_\varrho; \mathbb{S}_+^{N-1}) : \mathbf{v}(x) = \mathbf{a} \text{ if } x \cdot \varrho = -\frac{1}{2}, \mathbf{v}(x) = \mathbf{b} \text{ if } x \cdot \varrho = \frac{1}{2} \right\}. \quad (2-24)$$

Lemma 2.4. *Assume (H1). Then*

$$K_f(x, \mathbf{a}, \mathbf{b}, \varrho) = \inf \left\{ \int_0^1 f(x, \boldsymbol{\gamma}(t)) |\dot{\boldsymbol{\gamma}}(t)| dt : \boldsymbol{\gamma} \in W^{1,1}((0, 1); \mathbb{S}_+^{N-1}), \boldsymbol{\gamma}(0) = \mathbf{a}, \boldsymbol{\gamma}(1) = \mathbf{b} \right\}. \quad (2-25)$$

The proof of Lemma 2.4 is identical to that of [Fonseca and Rybka 1992, Proposition 2.6], where the same result has been proved (under more general assumptions on the energy density) when the target space is \mathbb{R}^N rather than \mathbb{S}_+^{N-1} . Therefore we omit it.

In order to obtain a lower bound on the lower semicontinuous envelope of \mathcal{F}_f , in particular of its jump part, one needs an approximation lemma which relates a generic sequence in $W^{1,1}(Q_\varrho; \mathbb{S}_+^{N-1})$, converging to a step function, to a nongeneric one in $\mathcal{P}(\mathbf{a}, \mathbf{b}, \varrho)$:

Lemma 2.5. *Assume (H1) and (H2).*

Let $\mathbf{a}, \mathbf{b} \in \mathbb{S}_+^{N-1}$ and let $\mathbf{v}_n \in W^{1,1}(Q_\varrho; \mathbb{S}_+^{N-1})$ such that $\mathbf{v}_n \rightarrow \mathbf{u}_0$ in $L^1(Q_\varrho; \mathbb{S}_+^{N-1})$, where

$$\mathbf{u}_0(x) := \begin{cases} \mathbf{b} & \text{if } \langle x, \varrho \rangle \geq 0, \\ \mathbf{a} & \text{if } \langle x, \varrho \rangle < 0. \end{cases}$$

Then a sequence $\mathbf{w}_n \in \mathcal{P}(\mathbf{a}, \mathbf{b}, \varrho)$ exists such that $\mathbf{w}_n \rightarrow \mathbf{u}_0$ in $L^1(Q_\varrho; \mathbb{S}_+^{N-1})$ and

$$\liminf_{n \rightarrow \infty} \int_{Q_\varrho} f(x, \mathbf{v}_n) |\nabla \mathbf{v}_n| dx \geq \limsup_{n \rightarrow \infty} \int_{Q_\varrho} f(x, \mathbf{w}_n) |\nabla \mathbf{w}_n| dx.$$

Lemma 2.5 may be proved following line by line that of [Alicandro et al. 2007, Lemma 5.2], where the same result was proved (under more general assumptions on the energy density) when the target space is \mathbb{S}^{N-1} , and therefore we omit it. We just mention that the proof may in fact be simplified in the present

setting by using the standard projection onto \mathbb{S}_+^{N-1} (see estimate (3-23) and Lemma 3.9 below for a related approximation result).

Let \mathcal{G}_f be the functional defined in $\text{BV}(\Omega; \mathbb{S}_+^{N-1})$ by

$$\mathcal{G}_f(\mathbf{v}) := \int_{\Omega} f(x, \mathbf{v}) |\nabla \mathbf{v}| dx + \int_{J(\mathbf{v})} K_f(x, \mathbf{v}_-, \mathbf{v}_+, \nu_{\mathbf{v}}) d\mathcal{H}^{m-1} + \int_{\Omega} f(x, \mathbf{v}) d|D^c \mathbf{v}|$$

(and $+\infty$ elsewhere). Under an additional coercivity assumption on f , and when the target space is \mathbb{S}^{N-1} , in [Alicandro et al. 2007, Proposition 5.1] it is proved that \mathcal{G}_f coincides with the lower semicontinuous envelope of \mathcal{F}_f with respect to the L^1 -convergence. Of course, coercivity is crucial for the upper bound in that it guarantees that *any* sequence along which \mathcal{G}_f is bounded has a convergent subsequence. However, it may be dropped when only a lower bound is needed, provided it is a priori known that a sequence has good convergence properties:

Proposition 2.6. *Let f satisfy (H1)–(H3) and let $\mathbf{v}_n \in W^{1,1}(\Omega; \mathbb{S}_+^{N-1})$ such that $\mathbf{v}_n \rightharpoonup \mathbf{v} \in \text{BV}(\Omega; \mathbb{S}_+^{N-1})$ and $\mathbf{v}_n \rightarrow \mathbf{v}$ in $L^1(\Omega; \mathbb{S}_+^{N-1})$. Then*

$$\mathcal{G}_f(\mathbf{v}) \leq \liminf_{n \rightarrow \infty} \mathcal{F}_f(\mathbf{v}_n).$$

Given Lemma 2.5, the proof follows line by line that of [Alicandro et al. 2007, Proposition 5.1], and the difference between the target spaces (\mathbb{S}^{N-1} versus \mathbb{S}_+^{N-1}) is harmless. Therefore we omit it.

3. Existence of solutions

In this section we introduce the notion of solutions to (1-4) and we prove their existence.

As is mentioned in Section 2C, on its jump set J_u a function $\mathbf{u} \in \text{BV}(\Omega; \mathbb{R}^N)$ has a jump discontinuity between two distinct values, \mathbf{u}_+ and \mathbf{u}_- , and the value of the precise representative of \mathbf{u} is given by $(\mathbf{u}_+ + \mathbf{u}_-)/2$. Note that $(\mathbf{u}_+ + \mathbf{u}_-)/2$ is the midpoint of the segment which connects \mathbf{u}_+ and \mathbf{u}_- . In this sense, $(\mathbf{u}_+ + \mathbf{u}_-)/2$ has natural counterparts in \mathbb{S}^{N-1} endowed with the standard geodesic distance d_g on \mathbb{S}^{N-1} , the *geodesic midpoints*:

Definition 3.1. Let \mathbb{A} be a geodesically convex subset of \mathbb{S}^{N-1} and let $\mathbf{u}_-, \mathbf{u}_+ \in \mathbb{A}$. A point $\mathbf{u}_g \in \mathbb{A}$ is called a *geodesic midpoint* on \mathbb{A} between \mathbf{u}_- and \mathbf{u}_+ if

- (i) \mathbf{u}_g belongs to a greatest circle of \mathbb{S}^{N-1} passing through \mathbf{u}_- and \mathbf{u}_+ , and
- (ii) $d_g(\mathbf{u}_g, \mathbf{u}_-) = d_g(\mathbf{u}_g, \mathbf{u}_+)$.

In particular, when $\mathbb{A} = \mathbb{S}_+^{N-1}$, geodesic midpoints are uniquely determined:

$$\mathbf{u}_g = \frac{\mathbf{u}_- + \mathbf{u}_+}{|\mathbf{u}_- + \mathbf{u}_+|} \quad \text{for all } \mathbf{u}_-, \mathbf{u}_+ \in \mathbb{S}_+^{N-1}.$$

Thus we can introduce the notion of *geodesic representative* of $\mathbf{u} \in \text{BV}(\Omega; \mathbb{S}_+^{N-1})$:

Definition 3.2. Let $\mathbf{u} \in \text{BV}(\Omega; \mathbb{S}_+^{N-1})$. The *geodesic representative* $\mathbf{u}_g : \Omega \setminus (S_u \setminus J_u) \rightarrow \mathbb{S}_+^{N-1}$ of \mathbf{u} is defined by

$$\mathbf{u}_g = \begin{cases} \mathbf{u}^* & \text{on } \Omega \setminus S_u, \\ \mathbf{u}^*/|\mathbf{u}^*| & \text{on } J_u. \end{cases}$$

Note that $\mathbf{u}_g \in \text{BV}(\Omega; \mathbb{S}_+^{N-1})$ since \mathbf{u}_+ and \mathbf{u}_- are \mathcal{H}^{m-1} -measurable on J_u ; see [Ambrosio et al. 2000, Proposition 3.69]. Hence the following Radon measures are well defined:

$$|\mathbf{u}^*| |D\mathbf{u}| := |\nabla \mathbf{u}| \mathcal{L}^m + |D^c \mathbf{u}| + |\mathbf{u}^*| |\mathbf{u}_+ - \mathbf{u}_-| \mathcal{H}^{m-1} \llcorner_{J_u}, \quad (3-1)$$

$$\mathbf{u}_g |D\mathbf{u}| := \mathbf{u} (|\nabla \mathbf{u}| \mathcal{L}^m + |D^c \mathbf{u}|) + \mathbf{u}_g |\mathbf{u}_+ - \mathbf{u}_-| \mathcal{H}^{m-1} \llcorner_{J_u}. \quad (3-2)$$

Moreover, $\mathbf{u}_g |D\mathbf{u}| \in \mathcal{M}_{\mathcal{H}}(\Omega; \mathbb{R}^N)$ (see Section 2D).

Remark 3.3. As shown in the proof of Lemma 3.9, the projections onto \mathbb{S}_+^{N-1} of the mollifications of \mathbf{u} point-wise converge to \mathbf{u}_g in Ω . In this sense, the geodesic representative \mathbf{u}_g is a natural representative for BV-vector fields with values into \mathbb{S}_+^{N-1} .

We are now ready to introduce the concept of solution for (1-4).

Definition 3.4. Let $\mathbb{A} = \mathbb{S}_+^{N-1}$, $T > 0$, and $\mathbf{u}_0 \in \text{BV}(\Omega; \mathbb{S}_+^{N-1})$. A function

$$\mathbf{u} \in L^\infty(0, T; \text{BV}(\Omega; \mathbb{R}^N)) \cap C(0, T; L^1(\Omega; \mathbb{R}^N)), \quad \mathbf{u}_t \in L^2(0, T; L^2(\Omega; \mathbb{R}^N))$$

is a solution to (1-4) in Q_T if $\mathbf{u}(0) = \mathbf{u}_0$, $\mathbf{u} \in \mathbb{S}_+^{N-1}$ a.e. in Q_T , and there exists a matrix-valued function $\mathbf{Z} \in L^\infty(Q_T, \mathbb{R}^{N \times m})$, with $\|\mathbf{Z}\|_\infty \leq 1$ and $\mathbf{Z}(t) \in X_{\mathcal{M}}^N(\Omega)$ for almost all $t \in (0, T)$, such that

$$\mathbf{u}_t(t) - \text{div } \mathbf{Z}(t) = \mathbf{u}(t)_g |D\mathbf{u}(t)| \quad \text{as measures for a.e. } t \in [0, T], \quad (3-3)$$

$$\mathbf{u}_t(t) \wedge \mathbf{u}(t) = \text{div}(\mathbf{Z}(t) \wedge \mathbf{u}(t)) \quad \text{in } L^2(\Omega; \Lambda_2(\mathbb{R}^N)) \text{ for a.e. } t \in [0, T], \quad (3-4)$$

$$\mathbf{Z}^T \mathbf{u} = 0 \quad \text{a.e. in } Q_T, \quad (3-5)$$

and

$$[\mathbf{Z}(t), \nu] = 0 \quad \mathcal{H}^{m-1}\text{-a.e. on } \partial\Omega \text{ for a.e. } t \in [0, T]. \quad (3-6)$$

The next observation clarifies the concept of solution given in Definition 3.4.

Proposition 3.5. Let \mathbf{u} be a solution of (1-4) in the sense of Definition 3.4. Then

$$\mathbf{Z}(t) : D\mathbf{u}(t) = |\mathbf{u}(t)^*| |D\mathbf{u}(t)| \quad \text{as measures for a.e. } t \in (0, T). \quad (3-7)$$

Proof. We take any $\varphi \in \mathcal{D}(\Omega)$. Then

$$\begin{aligned} \int_{\Omega} \varphi d(\mathbf{Z}(t) : D\mathbf{u}(t)) &\stackrel{(2-15)}{=} - \int_{\Omega} \varphi \mathbf{u}(t)^* \cdot d(\text{div } \mathbf{Z}(t)) - \int_{\Omega} (\mathbf{Z}(t)^T \mathbf{u}(t)) \cdot \nabla \varphi dx \\ &\stackrel{(3-5)}{=} \int_{\Omega} \varphi \mathbf{u}^*(t) \cdot d(\mathbf{u}_t(t) + \mathbf{u}(t)_g |D\mathbf{u}(t)|) \\ &\stackrel{(3-3)}{=} \int_{\Omega} \varphi \mathbf{u}(t)^* \cdot d(\mathbf{u}(t)_g |D\mathbf{u}(t)|), \end{aligned}$$

where in the last line we have used the facts that $|\mathbf{u}(t)| = 1$, $\mathbf{u}_t \in L^2(Q_T; \mathbb{R}^N)$ and the fact that $\mathbf{u}(t)_g |D\mathbf{u}(t)| \in \mathcal{M}_{\mathcal{H}}(\Omega; \mathbb{R}^N)$ a.e. $t \in (0, T)$. Finally, by (3-1) we get

$$\begin{aligned} \int_{\Omega} \varphi d(\mathbf{Z}(t) : D\mathbf{u}(t)) &= \int_{\Omega} \varphi d(|\nabla \mathbf{u}(t)| \mathcal{L}^m + |D^c(\mathbf{u}(t))|) + \int_{J_{\mathbf{u}(t)}} \varphi |\mathbf{u}(t)^*| |\mathbf{u}(t)_+ - \mathbf{u}(t)_-| d\mathcal{H}^{m-1} \\ &= \int_{\Omega} \varphi d(|\mathbf{u}(t)^*| |D\mathbf{u}(t)|). \end{aligned} \quad \square$$

Our main result is the following existence theorem.

Theorem 3.6. *For any $T > 0$ and any $\mathbf{u}_0 \in \text{BV}(\Omega; \mathbb{S}_+^{N-1})$, there exists a solution \mathbf{u} to (1-4) in the sense of Definition 3.4.*

To prove Theorem 3.6 we need to recall or establish several results. The first one follows as a particular case from [Barrett et al. 2008, Theorem 4.1, (4.24), and (4.25)] (with $\lambda = \mathbf{g} = 0$ and $p = 2$).

Proposition 3.7. *Let $\varepsilon > 0$, $T > 0$, and $\alpha > 0$. If $\mathbf{u}_0^\varepsilon \in W^{1,2}(\Omega; \mathbb{S}^{N-1})$, then there exists*

$$\mathbf{u}^\varepsilon \in L^\infty(0, T; W^{1,2}(\Omega; \mathbb{R}^N)) \cap W^{1,2}(0, T; L^2(\Omega; \mathbb{R}^N))$$

such that $\mathbf{u}^\varepsilon(0, \cdot) = \mathbf{u}_0^\varepsilon$,

$$|\mathbf{u}^\varepsilon| = 1 \quad \text{a.e. in } Q_T, \tag{3-8}$$

and \mathbf{u}^ε is a weak solution to

$$\begin{cases} \mathbf{u}_t^\varepsilon = \text{div } \mathbf{Z}^\varepsilon + \boldsymbol{\mu}^\varepsilon & \text{in } Q_T, \\ [\mathbf{Z}^\varepsilon, \nu] = 0 & \text{in } S_T, \end{cases} \tag{3-9}$$

where

$$\mathbf{Z}^\varepsilon = \varepsilon^\alpha \nabla \mathbf{u}^\varepsilon + \frac{\nabla \mathbf{u}^\varepsilon}{\sqrt{|\nabla \mathbf{u}^\varepsilon|^2 + \varepsilon^2}} \quad \text{and} \quad \boldsymbol{\mu}^\varepsilon = \varepsilon^\alpha \mathbf{u}^\varepsilon |\nabla \mathbf{u}^\varepsilon|^2 + \mathbf{u}^\varepsilon \frac{|\nabla \mathbf{u}^\varepsilon|^2}{\sqrt{|\nabla \mathbf{u}^\varepsilon|^2 + \varepsilon^2}} \tag{3-10}$$

in the sense that

$$\int_0^T \int_{\Omega} (\mathbf{u}_t^\varepsilon \cdot \mathbf{v} + \mathbf{Z}^\varepsilon : \nabla \mathbf{v} - \boldsymbol{\mu}^\varepsilon \cdot \mathbf{v}) \, dx \, dt = 0 \quad \text{for all } \mathbf{v} \in C^1(\bar{Q}_T; \mathbb{R}^N). \tag{3-11}$$

Furthermore,

$$(\mathbf{Z}^\varepsilon)^T \mathbf{u}^\varepsilon = 0 \quad \text{a.e. in } Q_T, \tag{3-12}$$

$$\mathbf{u}_t^\varepsilon \cdot \mathbf{u}^\varepsilon = 0 \quad \text{a.e. in } Q_T, \tag{3-13}$$

$$\mathbf{u}_t^\varepsilon \wedge \mathbf{u}^\varepsilon = \text{div}(\mathbf{Z}^\varepsilon \wedge \mathbf{u}^\varepsilon), \tag{3-14}$$

and

$$J_\alpha^\varepsilon(\mathbf{u}^\varepsilon(t)) + \int_0^t \int_{\Omega} |\mathbf{u}_s^\varepsilon|^2 \, dx \, ds \leq J_\alpha^\varepsilon(\mathbf{u}_0) \quad \text{for a.e. } t \in [0, T], \tag{3-15}$$

where the energy functional J_α^ε is defined as

$$J_\alpha^\varepsilon(\mathbf{v}) := \varepsilon^\alpha \int_{\Omega} |\nabla \mathbf{v}(x)|^2 \, dx + \int_{\Omega} \sqrt{|\nabla \mathbf{v}(x)|^2 + \varepsilon^2} \, dx, \quad \mathbf{v} \in W^{1,2}(\Omega; \mathbb{R}^N),$$

and a positive ε -independent constant C exists such that

$$\|\operatorname{div} \mathbf{Z}^\varepsilon\|_{L^2(0,T;L^1(\Omega;\mathbb{R}^N))} \leq C, \quad (3-16)$$

$$\|\operatorname{div}(\mathbf{Z}^\varepsilon \wedge \mathbf{u}^\varepsilon)\|_{L^2(0,T;L^2(\Omega;\Lambda_2(\mathbb{R}^N)))} \leq C, \quad (3-17)$$

$$\varepsilon^{\alpha/2} \|\nabla \mathbf{u}^\varepsilon(t)\|_{L^\infty(0,T;L^2(\Omega;\mathbb{R}^{N \times m}))} \leq C. \quad (3-18)$$

We next show that if \mathbf{u}_0^ε takes values in the first hyperoctant, then \mathbf{u}^ε does too:

Lemma 3.8. *If $\mathbf{u}_0^\varepsilon \in W^{1,2}(\Omega; \mathbb{S}_+^{N-1})$, then the weak solution to Problem (3-9) given by Proposition 3.7 verifies $\mathbf{u}^\varepsilon \in \mathbb{S}_+^{N-1}$ a.e. in Q_T .*

Proof. Let $(s)^- = \max\{0, -s\}$ and let $(\mathbf{u}^\varepsilon)^- = ((u^{\varepsilon,1})^-, \dots, (u^{\varepsilon,N})^-)$. Pick a sequence of smooth functions \mathbf{v}_n such that $\mathbf{v}_n \rightarrow (\mathbf{u}^\varepsilon)^-$ in $L^2(0, T; W^{1,2}(\Omega)) \cap W^{1,2}(0, T; L^2(\Omega))$ as $n \rightarrow +\infty$. Choosing $\mathbf{v} = \mathbf{v}_n$ in (3-11) and passing to the limit as $n \rightarrow +\infty$, we obtain on the one hand

$$\int_0^T \int_\Omega (\mathbf{u}^\varepsilon)^- \cdot \mathbf{u}_t^\varepsilon \, dx \, dt = \int_0^T \int_\Omega \left(\varepsilon^\alpha + \frac{1}{\sqrt{\varepsilon^2 + |\nabla \mathbf{u}^\varepsilon|^2}} \right) |\nabla (\mathbf{u}^\varepsilon)^-|^2 (1 - |(\mathbf{u}^\varepsilon)^-|^2) \, dx \, dt \geq 0. \quad (3-19)$$

On the other hand, since $\mathbf{u}^\varepsilon \in W^{1,2}(0, T; L^2(\Omega; \mathbb{R}^N))$,

$$0 \leq \int_0^T \int_\Omega (\mathbf{u}^\varepsilon)^- \cdot \mathbf{u}_t^\varepsilon \, dx \, dt = \int_\Omega (|(\mathbf{u}_0)^-|^2 - |(\mathbf{u}^\varepsilon(T))^-|^2) \, dx = - \int_\Omega |(\mathbf{u}^\varepsilon(T))^-|^2 \, dx, \quad (3-20)$$

hence the negative part of each component remains 0 for all times. \square

Provided α is large enough, any function in $\operatorname{BV}(\Omega; \mathbb{S}_+^{N-1})$ can be approximated in $W^{1,2}(\Omega; \mathbb{S}_+^{N-1})$ in such a way that the initial energy is controlled.

Lemma 3.9. *Given $\mathbf{u}_0 \in \operatorname{BV}(\Omega; \mathbb{S}_+^{N-1})$ and $\alpha > m$, there exist $\mathbf{u}_0^\varepsilon \in W^{1,2}(\Omega; \mathbb{S}_+^{N-1})$ such that*

- (i) $\mathbf{u}_0^\varepsilon \rightarrow \mathbf{u}_0$ in $L^p(\Omega; \mathbb{R}^N)$ for all $p < \infty$ as $\varepsilon \rightarrow 0$,
- (ii) $\mathbf{u}_0^\varepsilon \rightarrow (\mathbf{u}_0)_g \mathcal{H}^{m-1}$ -a.e. in Ω as $\varepsilon \rightarrow 0$,
- (iii) $J_\alpha^\varepsilon(\mathbf{u}_0^\varepsilon) \rightarrow L < +\infty$ as $\varepsilon \rightarrow 0$.

Proof. We will construct \mathbf{u}_0^ε as the projection onto \mathbb{S}_+^{N-1} of the convolution of a suitable extension $T\mathbf{u}_0$ of \mathbf{u}_0 with a standard mollifier. In order to do this, we proceed as in [Ambrosio et al. 2000, Proposition 3.21], to which we refer for further details; see also [Brezis 2011, Theorem 9.7].

Since $\bar{\Omega}$ is compact, there exists a finite collection $\{R_i\}_{i \in I}$ of open rectangles whose union B contains $\bar{\Omega}$, which satisfies the following property: for any $i \in I$, either

- (a) $R_i \subset \Omega$ or
- (b) $\partial\Omega \cap R_i$ is the graph of a Lipschitz function defined on one face L_i of R_i and the closure of $\partial\Omega \cap R_i$ intersects neither \bar{L}_i nor the closure of the face opposite to L_i .

Let $\Omega_i = \Omega \cap R_i$. In case (b), up to a translation, a rotation, and a homothety, we have $R_i = L_i \times (-1, 1)$ with Ω_i on the upper side of R_i (that is, $\Omega_i = \{x = (y, z) : z > \phi_i(y)\}$). A vertical deformation $\varphi : R_i \rightarrow R_i$

exists such that $\varphi(\Omega_i) = R_i^+ = L_i \times (0, 1)$ and both φ and its inverse are Lipschitz. Given $u \in \text{BV}(\Omega)$, the operator $T_i : R_i \rightarrow \mathbb{R}$ is defined as the identity in case (a) and as

$$T_i(u) = T'_i(u \circ \varphi^{-1}) \circ \varphi, \quad \text{where } T'_i(u)(y, z) = u(y, |z|),$$

in case (b). Note that $|u_0| = 1$ a.e. in Ω , the maps φ and its inverse are Lipschitz, and T'_i does not change the value of u . Hence

$$U_i := \{x \in R_i : |(T_i(u_0^1), \dots, T_i(u_0^N))| \neq 1\} \quad \text{has zero measure.}$$

Let $\{\eta_i\}_{i \in I}$ be a partition of unity relative to $\{R_i\}_{i \in I}$, that is, $\text{supp}(\eta_i) \subset R_i$, $0 \leq \eta_i \leq 1$ for any $i \in I$ and there exists $r > 0$ such that $\sum_{i \in I} \eta_i \equiv 1$ in a neighborhood of $\bar{\Omega}$ containing $\Omega \oplus B_r$. We now define

$$T\mathbf{u}_0 : B = \bigcup_{i \in I} R_i \rightarrow \mathbb{R}^N, \quad T\mathbf{u}_0 := \left(\sum_{i \in I} T_i(u_0^1)\eta_i, \dots, \sum_{i \in I} T_i(u_0^N)\eta_i \right).$$

It is readily checked that $T \in \text{BV}(\Omega \oplus B_r; \mathbb{R}^N)$. Now let $k > 0$ be the cardinality of I and $U = \bigcup_{i \in I} U_i$ (a set of zero measure). We observe that

$$|T\mathbf{u}_0(x)| \geq \frac{1}{k} \quad \text{for all } x \in \Omega \oplus B_r \setminus U. \tag{3-21}$$

Indeed, for each $x \in (\Omega \oplus B_r) \setminus U$, there exists $i(x) \in I$ such that $\eta_{i(x)}(x) \geq 1/k$: since each component of \mathbf{u}_0 is nonnegative and $x \notin U_{i(x)}$,

$$|T\mathbf{u}_0(x)|^2 \geq \frac{1}{k^2} ((T_{i(x)}(u_0^1))^2 + \dots + (T_{i(x)}(u_0^N))^2) = \frac{1}{k^2}.$$

Given $\varepsilon < r$, let $\rho_\varepsilon(x) := \varepsilon^{-m} \rho(x/\varepsilon)$ be a standard mollifier. As is well known (see, for example, [Ambrosio et al. 2000, Remark 3.22]) $T\mathbf{u}_0 \star \rho_\varepsilon$ converges to $T\mathbf{u}_0$ strictly in $\text{BV}(\Omega; \mathbb{R}^N)$ and strongly in $L^1(\Omega; \mathbb{R}^N)$. Since $\|T\mathbf{u}_0 \star \rho_\varepsilon\|_\infty \leq 1$, the last convergence upgrades to

$$T\mathbf{u}_0 \star \rho_\varepsilon \rightarrow T\mathbf{u}_0 \quad \text{in } L^p(\Omega; \mathbb{R}^N) \quad \text{for all } 1 \leq p < \infty. \tag{3-22}$$

By (3-21) and since $(T(\mathbf{u}_0))^\ell \geq 0$ for $\ell = 1, \dots, N$, a direct computation shows that

$$|T\mathbf{u}_0 \star \rho_\varepsilon(x)| \geq \frac{1}{k\sqrt{N}} \quad \text{for all } x \in \Omega. \tag{3-23}$$

In addition, it follows from [Ambrosio et al. 2000, Corollary 3.80] that $T\mathbf{u}_0 \star \rho_\varepsilon \rightarrow (T\mathbf{u}_0)^* = \mathbf{u}_0^*$ pointwise in $\Omega \setminus (S_{\mathbf{u}_0} \setminus J_{\mathbf{u}_0})$. Together with (3-23), this implies that

$$\mathbf{u}_0^\varepsilon := \frac{T\mathbf{u}_0 \star \rho_\varepsilon}{|T\mathbf{u}_0 \star \rho_\varepsilon|} \rightarrow (\mathbf{u}_0)_g \quad \mathscr{H}^{m-1}\text{-a.e. in } \Omega. \tag{3-24}$$

Furthermore, (3-23) and (3-22) easily imply that $\mathbf{u}_0^\varepsilon \rightarrow \mathbf{u}_0$ in $L^p(\Omega; \mathbb{R}^N)$ for all $1 \leq p < \infty$. Finally, applying the chain rule and (3-23), [Ambrosio et al. 2000, Proposition 3.2], and [Ambrosio et al. 2000, Theorem 2.2(b)] (in this order), we see that

$$\int_\Omega |\nabla \mathbf{u}_0^\varepsilon| dx \leq C \int_\Omega |\nabla(T\mathbf{u}_0 \star \rho_\varepsilon)| dx = C \int_\Omega |(DT\mathbf{u}_0) \star \rho_\varepsilon| dx \leq C |DT\mathbf{u}_0|(\Omega \oplus B_\varepsilon). \tag{3-25}$$

Similarly,

$$\int_{\Omega} |\nabla \mathbf{u}_0^\varepsilon|^2 dx \leq C \int_{\Omega} |(DT\mathbf{u}_0) \star \rho_\varepsilon|^2 dx \leq C \|(DT\mathbf{u}_0) \star \rho_\varepsilon\|_\infty \int_{\Omega} |(DT\mathbf{u}_0) \star \rho_\varepsilon| dx,$$

and, using the definition of ρ_ε , we conclude that

$$\varepsilon^\alpha \int_{\Omega} |\nabla \mathbf{u}_0^\varepsilon|^2 dx \leq C \varepsilon^{\alpha-m} (|DT\mathbf{u}_0|(\Omega \oplus B_\varepsilon))^2. \quad (3-26)$$

Inequalities (3-25) and (3-26), together with (3-24), complete the proof. \square

Proof of Theorem 3.6. We proceed in steps. In the first step, we use the previous lemmas, together with standard compactness arguments, to identify a triplet $(\mathbf{u}, \mathbf{Z}, \boldsymbol{\mu})$. In the second step we identify $\boldsymbol{\mu}$ in terms of \mathbf{u} and \mathbf{Z} , which automatically yields an upper bound on $|\boldsymbol{\mu}|$. In the third step, collecting the information of the previous two steps, we note that \mathbf{u} satisfies all the properties in Definition 3.4 except for

$$\boldsymbol{\mu}(t) = \mathbf{u}(t)_g |D\mathbf{u}(t)| \quad \text{as measures for a.e. } t \in [0, T], \quad (3-27)$$

to which the rest of the proof is devoted. In the fourth step we use the lower semicontinuity results in Section 2F to prove a lower bound on $\boldsymbol{\mu}(t)$ over the diffuse support of $|D\mathbf{u}(t)|$. In the fifth step, we revise the blow-up argument given in [Fonseca and Müller 1993; Fonseca and Rybka 1992] to obtain a lower bound on $\boldsymbol{\mu}(t)$ over $J_{\mathbf{u}(t)}$. Finally, in the sixth step we complete the proof.

Step 1: Passage to the limit. Let \mathbf{u}_0^ε and \mathbf{u}^ε be as given by Lemma 3.9 and Proposition 3.7, respectively. By Lemma 3.8, $\mathbf{u}^\varepsilon \in \mathbb{S}_+^{N-1}$ a.e. in Q_T . By (3-8), Lemma 3.9(iii), and (3-15), a positive constant C (independent of ε) exists such that

$$\sup_{t \in (0, T)} \|\mathbf{u}^\varepsilon\|_{W^{1,1}(\Omega)} \leq C, \quad (3-28)$$

$$\|\mathbf{u}_t^\varepsilon\|_{L^2(0, T; L^2(\Omega; \mathbb{R}^N))} \leq C. \quad (3-29)$$

We recall that $BV(\Omega; \mathbb{R}^N)$ is compactly embedded in $L^1(\Omega; \mathbb{R}^N)$ [Ambrosio et al. 2000, Theorem 3.23]. Hence the Aubin–Simon compactness criterion [Simon 1987, Corollary 8.4], together with (3-28) and (3-29), implies that

$$\mathbf{u}^\varepsilon \rightarrow \mathbf{u} \quad \text{in } C(0, T; L^1(\Omega; \mathbb{R}^N)) \text{ and a.e. in } Q_T \quad (3-30)$$

for a subsequence. By the lower semicontinuity of the total variation [Ambrosio et al. 2000, Remark 3.5], (3-30) and (3-15) imply that

$$\mathbf{u} \in L^\infty(0, T; BV(\Omega; \mathbb{R}^N)). \quad (3-31)$$

From (3-30) and Lemma 3.9(i), we have

$$\mathbf{u}(0) = \mathbf{u}_0 \quad (3-32)$$

and, using also (3-8),

$$|\mathbf{u}| = 1 \quad \text{a.e. in } Q_T. \quad (3-33)$$

By a standard interpolation argument, the boundedness of \mathbf{u}^ε in $L^\infty(0, T; L^\infty(\Omega; \mathbb{R}^N))$ and (3-30) imply that

$$\mathbf{u}^\varepsilon \rightharpoonup \mathbf{u} \quad \text{in } L^p(0, T; L^q(\Omega; \mathbb{R}^N)) \text{ for all } p, q \in [1, \infty) \text{ and a.e. in } Q_T. \tag{3-34}$$

Moreover, it follows from (3-29) that

$$\mathbf{u}_t^\varepsilon \rightharpoonup \mathbf{u}_t \quad \text{in } L^2(0, T; L^2(\Omega; \mathbb{R}^N)). \tag{3-35}$$

By (3-15), Lemma 3.9(iii), and (3-18), a subsequence exists such that

$$\varepsilon^\alpha \nabla \mathbf{u}^\varepsilon \rightharpoonup 0 \quad \text{in } L^2(0, T; L^2(\Omega; \mathbb{R}^{N \times m})), \tag{3-36}$$

$$\frac{\nabla \mathbf{u}^\varepsilon}{\sqrt{|\nabla \mathbf{u}^\varepsilon|^2 + \varepsilon^2}} \xrightarrow{*} \mathbf{Z} \quad \text{in } L^\infty(Q_T; \mathbb{R}^{N \times m}). \tag{3-37}$$

Recalling the definition (3-10) of \mathbf{Z}^ε , by (3-36) and (3-37) we obtain that

$$\mathbf{Z}^\varepsilon \rightharpoonup \mathbf{Z} \quad \text{in } L^2(0, T; L^2(\Omega; \mathbb{R}^{N \times m})), \tag{3-38}$$

and from (3-37) we also obtain that

$$\|\mathbf{Z}\|_{L^\infty(Q_T)} \leq 1. \tag{3-39}$$

Since $\{\boldsymbol{\mu}^\varepsilon\}$ is bounded in $L^\infty(0, T; L^1(\Omega; \mathbb{R}^N))$ and

$$L^\infty(0, T; L^1(\Omega; \mathbb{R}^N)) \subset L^\infty(0, T; \mathcal{M}(\Omega; \mathbb{R}^N)) \subset (L^1(0, T; C_0(\Omega; \mathbb{R}^N)))'$$

(see Section 2B), we have

$$\boldsymbol{\mu}^\varepsilon \xrightarrow{*} \boldsymbol{\mu} \quad \text{in } (L^1(0, T; C_0(\Omega; \mathbb{R}^N)))'. \tag{3-40}$$

Analogously, by (3-16),

$$\operatorname{div} \mathbf{Z}^\varepsilon \xrightarrow{*} \operatorname{div} \mathbf{Z} \quad \text{in } (L^2(0, T; C_0(\Omega; \mathbb{R}^N)))'. \tag{3-41}$$

Passing to the limit as $\varepsilon \rightarrow 0$ in (3-9)₁ (using (3-35), (3-41), and (3-40)), we obtain

$$\mathbf{u}_t - \operatorname{div} \mathbf{Z} = \boldsymbol{\mu} \quad \text{in } (L^2(0, T; C_0(\Omega; \mathbb{R}^N)))'. \tag{3-42}$$

Passing to the limit as $\varepsilon \rightarrow 0$ in (3-12) (using (3-38) and (3-34)), in (3-13) (using (3-35) and (3-34)), and in (3-14) (using (3-35), (3-38), and (3-34)), we get that

$$\mathbf{Z}^T \mathbf{u} = 0 \quad \text{a.e. in } Q_T, \tag{3-43}$$

$$\mathbf{u}_t \cdot \mathbf{u} = 0 \quad \text{a.e. in } Q_T, \tag{3-44}$$

$$\mathbf{u}_t(t) \wedge \mathbf{u}(t) = \operatorname{div}(\mathbf{Z}(t) \wedge \mathbf{u}(t)) \quad \text{in } L^2(\Omega; \Lambda_2(\mathbb{R}^N)) \text{ for a.e. } t \in [0, T]. \tag{3-45}$$

Step 2: The intermediate identification of $\boldsymbol{\mu}$ and its upper bound. We claim that

$$\boldsymbol{\mu} = *((\mathbf{Z} \wedge \mathbf{u}) \wedge D\mathbf{u}) \in L^\infty(0, T; \mathcal{M}(\Omega; \mathbb{R}^N)) \tag{3-46}$$

with

$$|\boldsymbol{\mu}(t)| \leq |D\mathbf{u}(t)| \quad \text{as measures for a.e. } t \in [0, T]. \quad (3-47)$$

Let

$$\mathcal{A} = (\mathcal{A}_1, \dots, \mathcal{A}_m) := *(\mathbf{Z} \wedge \mathbf{u}) \in L^\infty(Q_T; (\Lambda_{N-2}(\mathbb{R}^N))^m). \quad (3-48)$$

We have

$$*(\mathbf{u}_t \wedge \mathbf{u}) \stackrel{(3-45)}{=} *(\operatorname{div}(\mathbf{Z} \wedge \mathbf{u})) \stackrel{(2-18)}{=} \operatorname{div}(*(\mathbf{Z} \wedge \mathbf{u})) = \operatorname{div} \mathcal{A}, \quad (3-49)$$

hence $\mathcal{A}(t) \in \mathbf{X}_2(\Omega; \Lambda_{N-2}(\mathbb{R}^N))$ for a.e. t . Therefore, by Lemma 2.3, $*(\mathcal{A}(t) \wedge D\mathbf{u}(t)) \in \mathcal{M}(\Omega; \mathbb{R}^N)$ for almost every t with

$$|*(\mathcal{A}(t) \wedge D\mathbf{u}(t))| \stackrel{(2-21)}{\leq} \|\mathcal{A}(t)\|_\infty |D\mathbf{u}(t)| \stackrel{(2-9)}{=} \|\mathbf{Z}(t) \wedge \mathbf{u}(t)\|_\infty |D\mathbf{u}(t)| \stackrel{(2-10), (3-33), (3-39)}{\leq} |D\mathbf{u}(t)|, \quad (3-50)$$

and in addition

$$*(\mathcal{A}(t) \wedge D\mathbf{u}(t)) \stackrel{(2-22)}{=} -*(\operatorname{div} \mathcal{A}(t) \wedge \mathbf{u}(t)) \mathcal{L}^m + \operatorname{div}(*(\mathcal{A}(t) \wedge \mathbf{u}(t))). \quad (3-51)$$

It follows from (3-50) and (3-31) that

$$*(\mathcal{A} \wedge D\mathbf{u}) \in L^\infty(Q_T; \mathcal{M}(\Omega; \mathbb{R}^N)). \quad (3-52)$$

Using (3-51), we see that

$$\begin{aligned} \mathbf{u}_t &\stackrel{(3-33)}{=} |\mathbf{u}|^2 \mathbf{u}_t \stackrel{(2-7)}{=} (\mathbf{u}_t \cdot \mathbf{u}) \mathbf{u} - *(*(\mathbf{u}_t \wedge \mathbf{u}) \wedge \mathbf{u}) \\ &\stackrel{(3-44)}{=} -*(\operatorname{div} \mathcal{A} \wedge \mathbf{u}) \stackrel{(3-51)}{=} *(\mathcal{A} \wedge D\mathbf{u}) - \operatorname{div}(*(\mathcal{A} \wedge \mathbf{u})). \end{aligned} \quad (3-53)$$

On the other hand,

$$\begin{aligned} -*(\mathcal{A} \wedge \mathbf{u}) &\stackrel{(3-48)}{=} -*(*(\mathbf{Z} \wedge \mathbf{u}) \wedge \mathbf{u}) \\ &= -*(*(\mathbf{z}_1 \wedge \mathbf{u}) \wedge \mathbf{u}), \dots, *(*(\mathbf{z}_m \wedge \mathbf{u}) \wedge \mathbf{u}) \\ &\stackrel{(2-7)}{=} (|\mathbf{u}|^2 \mathbf{z}_1 - (\mathbf{u} \cdot \mathbf{z}_1) \mathbf{u}), \dots, (|\mathbf{u}|^2 \mathbf{z}_m - (\mathbf{u} \cdot \mathbf{z}_m) \mathbf{u}) \\ &= \mathbf{Z} - (\mathbf{Z}^T \mathbf{u}) \mathbf{u} \stackrel{(3-43)}{=} \mathbf{Z}. \end{aligned} \quad (3-54)$$

Combining (3-53) and (3-54), we obtain

$$\mathbf{u}_t = *(\mathcal{A} \wedge D\mathbf{u}) + \operatorname{div} \mathbf{Z},$$

which together with (3-42), (3-48), and (3-52), implies (3-46). Finally, (3-47) follows immediately from (3-46) and (3-50).

Step 3: Intermediate summary. It follows from (3-42), (3-35), and (3-46) that $\operatorname{div} \mathbf{Z} \in L^2(0, T; \mathcal{M}(\Omega; \mathbb{R}^N))$. Hence (3-42) upgrades to

$$\mathbf{u}_t(t) - \operatorname{div} \mathbf{Z}(t) = \boldsymbol{\mu}(t) \quad \text{as measures for a.e. } t \in [0, T]. \quad (3-55)$$

In particular,

$$\mathbf{Z}(t) \in \mathbf{X}_{\mathcal{M}}^N(\Omega) \quad \text{for a.e. } t \in [0, T]. \tag{3-56}$$

Thus the weak trace $[\mathbf{Z}(t), \nu]$ on $\partial\Omega$ of the normal component of $\mathbf{Z}(t)$ is well defined, and for all smooth \mathbf{w} we have

$$\begin{aligned} \int_0^T \int_{\partial\Omega} [\mathbf{Z}, \nu] \cdot \mathbf{w} \, d\mathcal{H}^{m-1} \, dt &\stackrel{(2-14)}{=} \int_0^T \left(\int_{\Omega} \mathbf{w} \cdot d(\operatorname{div} \mathbf{Z}) + \mathbf{Z} : \nabla \mathbf{w} \, dx \right) dt \\ &\stackrel{(3-38)}{=} \lim_{\varepsilon \rightarrow 0} \left(\int_0^T \int_{\Omega} (\mathbf{w} \cdot \operatorname{div} \mathbf{Z}^\varepsilon + \mathbf{Z}^\varepsilon : \nabla \mathbf{w}) \, dx \, dt \right) \stackrel{(3-9)_2}{=} 0. \end{aligned}$$

Hence

$$[\mathbf{Z}(t), \nu] = 0 \quad \mathcal{H}^{m-1}\text{-a.e. on } \partial\Omega \text{ for a.e. } t \in [0, T]. \tag{3-57}$$

Collecting (3-31), (3-30), (3-35), (3-32), (3-33), (3-39), (3-56), (3-43), (3-45), and (3-57), we see that all the properties of \mathbf{u} stated in Definition 3.4 are satisfied except for (3-3). In view of (3-55), in order to prove (3-3), it remains to show (3-27).

Step 4: The lower bound on $\boldsymbol{\mu}$ over the diffuse support of $|D\mathbf{u}|$. In view of (3-47), $\boldsymbol{\mu}(t)$ can be decomposed as

$$\boldsymbol{\mu}(t) = \frac{\boldsymbol{\mu}(t)}{|D\mathbf{u}(t)|} (|\nabla \mathbf{u}(t)|_{\mathcal{L}^m} + |D^c(\mathbf{u}(t))|) + \frac{\boldsymbol{\mu}(t)}{|D\mathbf{u}(t)|} |\mathbf{u}(t)_+ - \mathbf{u}(t)_-|_{\mathcal{H}^{m-1}} \lfloor J_{\mathbf{u}(t)}, \tag{3-58}$$

where $\boldsymbol{\mu}(t)/|D\mathbf{u}(t)| \in (L^1(\Omega; |D\mathbf{u}(t)|))^N$ denotes the Radon–Nikodým derivative of $\boldsymbol{\mu}(t)$ with respect to $|D\mathbf{u}(t)|$. We claim that

$$\mathbf{u}(t) \cdot \frac{\boldsymbol{\mu}(t)}{|D\mathbf{u}(t)|} \geq 1 \quad (|\nabla \mathbf{u}(t)|_{\mathcal{L}^m} + |D^c(\mathbf{u}(t))|)\text{-a.e. in } \Omega. \tag{3-59}$$

We first notice that

$$\boldsymbol{\mu}^{\varepsilon, \ell} \stackrel{(3-10)}{\geq} u^{\varepsilon, \ell} (\sqrt{\varepsilon^2 + |\nabla \mathbf{u}^\varepsilon|^2} - \varepsilon) \geq u^{\varepsilon, \ell} (|\nabla \mathbf{u}^\varepsilon| - \varepsilon), \quad \ell = 1, \dots, N. \tag{3-60}$$

For any $\varphi \in C(\bar{\Omega}; [0, \infty))$, $0 \leq \psi \in L^1((0, T))$, and $\ell \in \{1, \dots, N\}$, we have

$$\begin{aligned} \int_0^T \psi(t) \left(\int_{\Omega} \varphi \, d\boldsymbol{\mu}^\ell(t) \right) dt &\stackrel{(3-40)}{=} \lim_{\varepsilon \rightarrow 0} \int_0^T \psi(t) \left(\int_{\Omega} \varphi \boldsymbol{\mu}^{\varepsilon, \ell}(t) \, dx \right) dt \\ &\stackrel{(3-60)}{\geq} \liminf_{\varepsilon \rightarrow 0} \int_0^T \psi(t) \left(\int_{\Omega} \varphi u^{\varepsilon, \ell}(t) |\nabla \mathbf{u}^\varepsilon(t)| \, dx \right) dt. \end{aligned}$$

We claim that, for a.e. $t \in (0, T)$,

$$\mathbf{u}^\varepsilon(t) \rightharpoonup \mathbf{u}(t) \quad \text{in } \operatorname{BV}(\Omega; \mathbb{R}^N) \text{ as } \varepsilon \rightarrow 0. \tag{3-61}$$

Indeed, in view of (3-28), for a.e. t we have $\|\mathbf{u}^\varepsilon(t)\|_{W^{1,1}(\Omega)} < \infty$. Take any such t and assume for a contradiction that (3-61) does not hold, that is, that $\mathbf{u}^\varepsilon(t) \not\rightharpoonup \mathbf{u}(t)$ for a subsequence. By (3-28), a further subsequence would exist such that $\mathbf{u}^\varepsilon(t) \rightharpoonup \tilde{\mathbf{u}}$ for some $\tilde{\mathbf{u}} \in \operatorname{BV}(\Omega; \mathbb{R}^N)$. On the other hand, because of (3-30), $\mathbf{u}^\varepsilon(t) \rightarrow \mathbf{u}(t)$ in $L^1(\Omega; \mathbb{R}^N)$: hence $\tilde{\mathbf{u}} = \mathbf{u}(t)$, a contradiction.

In view of (3-61) and (3-30), we may apply Proposition 2.6 to the right-hand side of (3-60) with $f = f_{\varphi, \ell} : \Omega \times \mathbb{R}^N \rightarrow [0, \infty)$ defined by $f_{\varphi, \ell}(x, s) := \varphi(x)s^\ell|\xi|$. This implies that

$$\int_0^T \psi(t) \left(\int_{\Omega} \varphi d\mu^\ell(t) \right) dt \geq \int_0^T \psi(t) \left(\int_{\Omega} \varphi u^\ell(t) (|\nabla \mathbf{u}(t)| dx + d|D^c \mathbf{u}(t)|) + \int_{J_{\mathbf{u}(t)}} \varphi K_t^\ell d\mathfrak{H}^{m-1} \right) dt,$$

where

$$K_t^\ell = \inf \left\{ \int_0^1 \gamma^\ell(\tau) |\dot{\gamma}(\tau)| d\tau : \gamma \in W^{1,1}((0, 1); \mathbb{S}_+^{N-1}), \gamma(0) = \mathbf{u}(t)_-, \gamma(1) = \mathbf{u}(t)_+ \right\}. \quad (3-62)$$

By the arbitrariness of ψ , we conclude that

$$\int_{\Omega} \varphi d\mu^\ell(t) \geq \int_{\Omega} \varphi u^\ell(t) (|\nabla \mathbf{u}(t)| dx + d|D^c \mathbf{u}(t)|) + \int_{J_{\mathbf{u}(t)}} \varphi K_t^\ell d\mathfrak{H}^{m-1} \quad \text{for all } \varphi \in C(\bar{\Omega}) \quad (3-63)$$

for a.e. $t \in [0, T]$ and for all $\ell \in \{1, \dots, N\}$. Recalling (3-58), (3-63) yields

$$\frac{\mu^\ell(t)}{|D\mathbf{u}(t)|} \geq u^\ell(t) \quad (|\nabla \mathbf{u}(t)| \mathcal{L}^m + |D^c(\mathbf{u}(t))|)\text{-a.e. in } \Omega$$

for a.e. $t \in [0, T]$ and all $\ell = 1, \dots, N$. Now, recalling that $|\mathbf{u}(t)| = 1$ a.e. in Ω , we obtain the inequality (3-59) at once.

Remark 3.10. On the jump set $J_{\mathbf{u}(t)}$, the above argument would yield

$$|\mathbf{u}_+(t) - \mathbf{u}_-(t)| \mathbf{u}(t)_g \cdot \frac{\boldsymbol{\mu}(t)}{|D\mathbf{u}(t)|} \geq \mathbf{u}(t)_g \cdot (K_t^1, \dots, K_t^N) \quad \mathfrak{H}^{m-1}\text{-a.e. on } J_{\mathbf{u}(t)}.$$

Unfortunately, by (3-62) and obvious properties of the infimum,

$$\begin{aligned} & \mathbf{u}(t)_g \cdot (K_t^1, \dots, K_t^N) \\ & \leq \inf \left\{ \int_0^1 \mathbf{u}(t)_g \cdot \boldsymbol{\gamma}(\tau) |\dot{\boldsymbol{\gamma}}(\tau)| d\tau : \boldsymbol{\gamma} \in W^{1,1}((0, 1); \mathbb{S}_+^{N-1}), \boldsymbol{\gamma}(0) = \mathbf{u}(t)_-, \boldsymbol{\gamma}(1) = \mathbf{u}(t)_+ \right\}, \end{aligned} \quad (3-64)$$

whilst, as we shall see, it is the right-hand side of (3-64) which yields the sharp lower bound on the jump part (cf. (3-70)–(3-73) below). On the other hand, we can not use the results in Proposition 2.6 directly on $\mathbf{u}^* \cdot \boldsymbol{\mu}^\varepsilon$, since \mathbf{u}^* is a discontinuous function (though a very special one). This motivates the discussion that follows.

Step 5: The lower bound on $\boldsymbol{\mu}$ over $J_{\mathbf{u}(t)}$. We claim that

$$\mathbf{u}(t)_g \cdot \frac{\boldsymbol{\mu}(t)}{|D\mathbf{u}(t)|} \geq 1 \quad \mathfrak{H}^{m-1}\text{-a.e. on } J_{\mathbf{u}(t)}. \quad (3-65)$$

It follows from (3-8) and (3-28) that, for a.e. $t \in [0, T]$, there exists a subsequence ε_k such that

$$\mathbf{u}^{\varepsilon_k}(t) |\nabla \mathbf{u}^{\varepsilon_k}(t)| \xrightarrow{*} \tilde{\boldsymbol{\mu}}(t) \quad \text{in } \mathcal{M}(\Omega; \mathbb{R}^N). \quad (3-66)$$

Then (3-60) and the fact that $u^{\varepsilon, \ell} \geq 0$ imply that

$$\mu^\ell(t) \geq \tilde{\mu}^\ell(t) \geq 0 \quad \text{as measures for a.e. } t \in [0, T], \quad \ell \in \{1, \dots, N\}. \quad (3-67)$$

Hereafter, we argue for a fixed t and we do not specify dependence on t for notational convenience. Using the Radon–Nikodým theorem [Ambrosio et al. 2000, Theorem 1.28], we decompose $\tilde{\mu}$ into four mutually orthogonal measures:

$$\tilde{\mu} = \frac{\tilde{\mu}}{|Du|} |\nabla u| \mathcal{L}^N + \frac{\tilde{\mu}}{|Du|} |D^c u| + \frac{\tilde{\mu}}{|Du|} |u_+ - u_-| \mathcal{H}^{m-1} \llcorner J_u + (\tilde{\mu})^o$$

with $(\tilde{\mu})^o \perp |Du|$. It follows from (3-67) and (3-58) that

$$u_g \cdot \frac{\mu}{|Du|} \geq u_g \cdot \frac{\tilde{\mu}}{|Du|} \quad \mathcal{H}^{m-1}\text{-a.e. on } J_u.$$

Therefore, (3-65) is proved once we have shown that

$$u_g \cdot \frac{\tilde{\mu}}{|Du|} \geq 1 \quad \mathcal{H}^{m-1}\text{-a.e. on } J_u. \tag{3-68}$$

To prove (3-68) we apply the same blow-up argument as in [Fonseca and Müller 1993, Section 3].

From the Besicovitch differentiation theorem [Ambrosio et al. 2000, Theorem 2.22], for \mathcal{H}^{m-1} -a.e. $x_0 \in J_u$, we have

$$\frac{\tilde{\mu}}{|Du|}(x_0) = \lim_{\delta \rightarrow 0} \frac{\tilde{\mu}(x_0 + \delta Q_{v_u(x_0)})}{|u_+ - u_-| \mathcal{H}^{m-1}(J_u \cap (x_0 + \delta Q_{v_u(x_0)}))},$$

where Q_ζ is defined in Section 2F. On the other hand, by [Fonseca and Müller 1993, Lemma 2.6], for \mathcal{H}^{m-1} a.e. $x_0 \in J_u$, we also have

$$\lim_{\delta \rightarrow 0} \frac{1}{\delta^{m-1}} \int_{(x_0 + \delta Q_{v_u(x_0)}) \cap J_u} |u_+(x) - u_-(x)| d\mathcal{H}^{m-1} = |u_+(x_0) - u_-(x_0)|.$$

Therefore, letting

$$M = |u_+(x_0) - u_-(x_0)|$$

for notational convenience, we obtain that

$$M \frac{\tilde{\mu}}{|Du|}(x_0) = \lim_{\delta \rightarrow 0} \frac{1}{\delta^{m-1}} \int_{x_0 + \delta Q_{v_u(x_0)}} d\tilde{\mu}.$$

Then, for any $\ell \in \{1, \dots, N\}$, since the function $\chi_{x_0 + \delta Q_{v_u(x_0)}}$ is upper semicontinuous with compact support in Ω if δ is sufficiently small, we have

$$\begin{aligned} M \frac{\tilde{\mu}^\ell}{|Du|}(x_0) &= \lim_{\delta \rightarrow 0} \frac{1}{\delta^{m-1}} \int_{x_0 + \delta Q_{v_u(x_0)}} d\tilde{\mu}^\ell \\ &\stackrel{(3-66)}{\geq} \limsup_{\delta \rightarrow 0} \limsup_{k \rightarrow \infty} \frac{1}{\delta^{m-1}} \int_{x_0 + \delta Q_{v_u(x_0)}} u^{\varepsilon_k, \ell} |\nabla u^{\varepsilon_k}| dx \\ &= \limsup_{\delta \rightarrow 0} \limsup_{k \rightarrow \infty} \int_{Q_{v_u(x_0)}} v_{\delta, k}^\ell(y) |\nabla v_{\delta, k}(y)| dy, \end{aligned} \tag{3-69}$$

where

$$v_{\delta, k}(y) := u^{\varepsilon_k}(x_0 + \delta y).$$

We now observe that $\mathbf{v}_{\delta,k} \in W^{1,1}(Q_{v_{\mathbf{u}(x_0)}}; \mathbb{R}^N)$ and (see [Fonseca and Müller 1993, formula (3.2)])

$$\lim_{\delta \rightarrow 0} \lim_{k \rightarrow \infty} \|\mathbf{v}_{\delta,k} - \mathbf{w}_0\|_{L^1(Q_{v_{\mathbf{u}(x_0)}}; \mathbb{R}^N)} = 0,$$

where

$$\mathbf{w}_0(y) := \begin{cases} c\mathbf{c}\mathbf{u}_+(x_0) & \text{if } y \cdot v_{\mathbf{u}(x_0)} > 0, \\ \mathbf{u}_-(x_0) & \text{if } y \cdot v_{\mathbf{u}(x_0)} < 0. \end{cases}$$

Then, by a diagonalization argument, we may extract a subsequence \mathbf{v}_k converging to \mathbf{w}_0 in $L^1(Q_{v_{\mathbf{u}(x_0)}}; \mathbb{R}^N)$.

It follows from (3-69) that

$$M \frac{\tilde{\mu}^\ell}{|D\mathbf{u}|}(x_0) \geq \lim_{k \rightarrow \infty} \int_{Q_{v_{\mathbf{u}(x_0)}}} v_k^\ell(y) |\nabla \mathbf{v}_k(y)| dy.$$

Since $(u^\ell)^* \geq 0$ for all $\ell \in \{1, \dots, N\}$, this implies that

$$M \left(\mathbf{u}_g \cdot \frac{\tilde{\mu}}{|D\mathbf{u}|} \right)(x_0) \geq \lim_{k \rightarrow \infty} \int_{Q_{v_{\mathbf{u}(x_0)}}} \mathbf{u}_g(x_0) \cdot \mathbf{v}_k(y) |\nabla \mathbf{v}_k(y)| dy.$$

The function $f(x, \mathbf{s}) = f(\mathbf{s}) = \mathbf{u}_g(x_0) \cdot \mathbf{s}$ is continuous, nonnegative, and bounded. Then, applying Lemma 2.5, we obtain a new sequence

$$\mathbf{w}_k \in \mathcal{P}(\mathbf{u}_+(x_0), \mathbf{u}_-(x_0), v_{\mathbf{u}(x_0)})$$

(with \mathcal{P} given by (2-24)) converging to \mathbf{w}_0 in $L^1(Q_{v_{\mathbf{u}(x_0)}}; \mathbb{R}^N)$ and such that

$$M \left(\mathbf{u}_g \cdot \frac{\tilde{\mu}}{|D\mathbf{u}|} \right)(x_0) \geq \limsup_{k \rightarrow \infty} \int_{Q_{v_{\mathbf{u}(x_0)}}} \mathbf{u}_g(x_0) \cdot \mathbf{w}_k(y) |\nabla \mathbf{w}_k(y)| dy.$$

We may now apply Lemma 2.4. It follows from (2-23) and (2-25) that

$$M \left(\mathbf{u}_g \cdot \frac{\tilde{\mu}}{|D\mathbf{u}|} \right)(x_0) \geq \inf_{\boldsymbol{\gamma} \in \tilde{\Gamma}_N(\mathbf{u}_+(x_0), \mathbf{u}_-(x_0))} J_N[\mathbf{u}_+(x_0), \mathbf{u}_-(x_0)](\boldsymbol{\gamma}), \quad (3-70)$$

where

$$J_N[\mathbf{v}_0, \mathbf{v}_1](\boldsymbol{\gamma}) := \int_0^1 \mathbf{v}_g \cdot \boldsymbol{\gamma}(t) |\dot{\boldsymbol{\gamma}}(t)| dt, \quad \mathbf{v}_g := \frac{\mathbf{v}_0 + \mathbf{v}_1}{|\mathbf{v}_0 + \mathbf{v}_1|}, \quad (3-71)$$

and

$$\tilde{\Gamma}_N(\mathbf{v}_0, \mathbf{v}_1) := \{\boldsymbol{\gamma} \in W^{1,1}((0, 1); \mathbb{S}_+^{N-1}) : \boldsymbol{\gamma}(0) = \mathbf{v}_0, \boldsymbol{\gamma}(1) = \mathbf{v}_1\}. \quad (3-72)$$

In view of (3-70), (3-68) and therefore (3-65) follows from

$$\inf_{\boldsymbol{\gamma} \in \tilde{\Gamma}_N(\mathbf{u}_+(x_0), \mathbf{u}_-(x_0))} J_N[\mathbf{u}_+(x_0), \mathbf{u}_-(x_0)](\boldsymbol{\gamma}) \geq M = |\mathbf{u}_+(x_0) - \mathbf{u}_-(x_0)|. \quad (3-73)$$

This last inequality will be proved in Theorem 4.1, to which the next section is devoted.

Step 6: Conclusion. Recalling (3-58), the upper bound on $|\boldsymbol{\mu}|$ given by (3-47) immediately implies that

$$\left| \frac{\boldsymbol{\mu}(t)}{|D\mathbf{u}(t)|} \right| \leq 1 \quad |D\mathbf{u}(t)|\text{-a.e. in } \Omega \quad (3-74)$$

for a.e. $t \in [0, T]$. In particular, recalling (3-33),

$$\mathbf{u}(t) \cdot \frac{\boldsymbol{\mu}(t)}{|D\mathbf{u}(t)|} \leq 1 \quad (|\nabla\mathbf{u}(t)|\mathcal{L}^m + |D^c(\mathbf{u}(t))|)\text{-a.e. in } \Omega,$$

$$\mathbf{u}(t)_g \cdot \frac{\boldsymbol{\mu}(t)}{|D\mathbf{u}(t)|} \leq 1 \quad \mathcal{H}^{m-1}\text{-a.e. on } J_{\mathbf{u}(t)}$$

for a.e. $t \in [0, T]$. Combining these inequalities with the lower bounds in (3-59) and (3-65), we obtain

$$\mathbf{u}(t) \cdot \frac{\boldsymbol{\mu}(t)}{|D\mathbf{u}(t)|} = 1 \quad (|\nabla\mathbf{u}(t)|\mathcal{L}^m + |D^c(\mathbf{u}(t))|)\text{-a.e. in } \Omega, \quad (3-75)$$

$$\mathbf{u}(t)_g \cdot \frac{\boldsymbol{\mu}(t)}{|D\mathbf{u}(t)|} = 1 \quad \mathcal{H}^{m-1}\text{-a.e. on } J_{\mathbf{u}(t)} \quad (3-76)$$

for a.e. $t \in [0, T]$. We are now ready to complete the proof. By (2-2), we have

$$\left| \frac{\boldsymbol{\mu}(t)}{|D\mathbf{u}(t)|} \wedge \mathbf{u}(t) \right|^2 = \left| \frac{\boldsymbol{\mu}(t)}{|D\mathbf{u}(t)|} \right|^2 - \left| \frac{\boldsymbol{\mu}(t)}{|D\mathbf{u}(t)|} \cdot \mathbf{u}(t) \right|^2 \quad (|\nabla\mathbf{u}(t)|\mathcal{L}^m + |D^c(\mathbf{u}(t))|)\text{-a.e. in } \Omega \quad (3-77)$$

and

$$\left| \frac{\boldsymbol{\mu}(t)}{|D\mathbf{u}(t)|} \wedge \mathbf{u}(t)_g \right|^2 = \left| \frac{\boldsymbol{\mu}(t)}{|D\mathbf{u}(t)|} \right|^2 |\mathbf{u}(t)_g|^2 - \left| \frac{\boldsymbol{\mu}(t)}{|D\mathbf{u}(t)|} \cdot \mathbf{u}(t)_g \right|^2 \quad \mathcal{H}^{m-1}\text{-a.e. on } J_{\mathbf{u}(t)}. \quad (3-78)$$

Now, from (3-74),(3-75), and (3-77), we get

$$\left| \frac{\boldsymbol{\mu}(t)}{|D\mathbf{u}(t)|} \wedge \mathbf{u}(t) \right|^2 = 0 \quad (|\nabla\mathbf{u}(t)|\mathcal{L}^m + |D^c(\mathbf{u}(t))|)\text{-a.e. in } \Omega,$$

and from (3-74),(3-76), and (3-78), we get

$$\left| \frac{\boldsymbol{\mu}(t)}{|D\mathbf{u}(t)|} \wedge \mathbf{u}(t)_g \right|^2 = 0 \quad \mathcal{H}^{m-1}\text{-a.e. on } J_{\mathbf{u}(t)}.$$

Hence the wedge products on the left-hand side are zero.

Therefore, applying (2-7) and using once more the equalities in (3-75) and (3-76), we conclude that

$$\begin{aligned} \frac{\boldsymbol{\mu}(t)}{|D\mathbf{u}(t)|} &= |\mathbf{u}(t)|^2 \frac{\boldsymbol{\mu}(t)}{|D\mathbf{u}(t)|} = \mathbf{u}(t) \quad (|\nabla\mathbf{u}(t)|\mathcal{L}^m + |D^c(\mathbf{u}(t))|)\text{-a.e. in } \Omega, \\ \frac{\boldsymbol{\mu}(t)}{|D\mathbf{u}(t)|} &= |\mathbf{u}(t)_g|^2 \frac{\boldsymbol{\mu}(t)}{|D\mathbf{u}(t)|} = \mathbf{u}(t)_g \quad \mathcal{H}^{m-1}\text{-a.e. on } J_{\mathbf{u}(t)}. \end{aligned}$$

Plugging these expressions into (3-58), we obtain (3-27), and the proof is complete. \square

4. A nonconvex variational problem

In this section we study the minimization of a nonconvex functional, and, as a result, we prove the inequality (3-73).

Theorem 4.1. *Let $\mathbf{v}_0, \mathbf{v}_1 \in \mathbb{S}^{N-1}$ and let $J_N[\mathbf{v}_0, \mathbf{v}_1](\boldsymbol{\gamma})$ and $\tilde{\Gamma}_N(\mathbf{v}_0, \mathbf{v}_1)$ be given by (3-71) and (3-72). If $\mathbf{v}_0 \cdot \mathbf{v}_1 \geq 0$, then*

$$\min_{\boldsymbol{\gamma} \in \tilde{\Gamma}_N(\mathbf{v}_0, \mathbf{v}_1)} J_N[\mathbf{v}_0, \mathbf{v}_1](\boldsymbol{\gamma}) = |\mathbf{v}_1 - \mathbf{v}_0|.$$

Of course, it suffices to consider $\mathbf{v}_0 \neq \mathbf{v}_1$. Up to a rotation, we may assume without loss of generality that

$$\mathbf{v}_g = \frac{\mathbf{v}_0 + \mathbf{v}_1}{|\mathbf{v}_0 + \mathbf{v}_1|} = \mathbf{e}_N \quad \text{and} \quad \mathbf{v}_0, \mathbf{v}_1 \in \text{span}\{\mathbf{e}_{N-1}, \mathbf{e}_N\}.$$

Since \mathbf{v}_g is the geodesic midpoint and $\mathbf{v}_0 \cdot \mathbf{v}_1 \geq 0$, there exists $\theta_0 \in (0, \pi/4]$ such that

$$\mathbf{v}_0 = (0, \dots, 0, \sin \theta_0, \cos \theta_0) \quad \text{and} \quad \mathbf{v}_1 = (0, \dots, 0, -\sin \theta_0, \cos \theta_0).$$

Then

$$|\mathbf{v}_1 - \mathbf{v}_0| = 2 \sin \theta_0. \tag{4-1}$$

A curve which attains the equality in (4-3) is easily obtained: it is just the geodesic with respect to the standard metric of \mathbb{S}^{N-1} .

Lemma 4.2. *Let $\boldsymbol{\gamma}_{\min}(t) = (0, \dots, 0, \sin((1-2t)\theta_0), \cos((1-2t)\theta_0))$. Then $J(\boldsymbol{\gamma}_{\min}) = 2 \sin \theta_0$.*

After the above-mentioned rotation, \mathbb{S}_+^{N-1} is transformed into a geodesic simplex \mathcal{T} in \mathbb{S}^{N-1} . We consider a larger set of curves: let $\mathcal{P}_N(\mathbf{v}_0, \mathbf{v}_1)$ be given by

$$\mathcal{P}_N(\mathbf{v}_0, \mathbf{v}_1) = \{\mathbf{v} \in \mathbb{S}^{N-1} : \mathbf{v} \cdot \mathbf{v}_0 \geq 0, \mathbf{v} \cdot \mathbf{v}_1 \geq 0\}$$

and let

$$\Gamma_N(\mathbf{v}_0, \mathbf{v}_1) = \{\boldsymbol{\gamma} \in W^{1,1}((0, 1); \mathcal{P}_N(\mathbf{v}_0, \mathbf{v}_1)) : \boldsymbol{\gamma}(0) = \mathbf{v}_0, \boldsymbol{\gamma}(1) = \mathbf{v}_1\}.$$

Then

$$J_N[\mathbf{v}_0, \mathbf{v}_1](\boldsymbol{\gamma}) = \int_0^1 \gamma^N(t) |\dot{\boldsymbol{\gamma}}(t)| dt \quad \text{for } \boldsymbol{\gamma} = (\gamma^1, \dots, \gamma^N) \in \Gamma_N(\mathbf{v}_0, \mathbf{v}_1).$$

Hence, recalling (4-1) and Lemma 4.2, it suffices to prove that

$$\inf_{\boldsymbol{\gamma} \in \Gamma_N(\mathbf{v}_0, \mathbf{v}_1)} J_N[\mathbf{v}_0, \mathbf{v}_1](\boldsymbol{\gamma}) \geq 2 \sin \theta_0. \tag{4-2}$$

We now show that the problem in \mathbb{S}^{N-1} may be reduced to the same problem in \mathbb{S}^2 . Let

$$\bar{\mathbf{v}}_i = (0, (-1)^i \sin \theta_0, \cos \theta_0), \quad i = 0, 1,$$

denote the projection of \mathbf{v}_i onto the three-dimensional subspace $\text{span}\{\mathbf{e}_{N-2}, \mathbf{e}_{N-1}, \mathbf{e}_N\}$.

Lemma 4.3. *Let $N \geq 4$. Then*

$$\inf_{\boldsymbol{\gamma} \in \Gamma_N(\mathbf{v}_0, \mathbf{v}_1)} J_N[\mathbf{v}_0, \mathbf{v}_1](\boldsymbol{\gamma}) \geq \inf_{\boldsymbol{\gamma} \in \Gamma_3(\bar{\mathbf{v}}_0, \bar{\mathbf{v}}_1)} J_3[\bar{\mathbf{v}}_0, \bar{\mathbf{v}}_1](\boldsymbol{\gamma}).$$

Proof. For any $\boldsymbol{\gamma} \in \Gamma_N(\mathbf{v}_0, \mathbf{v}_1)$, consider the curve

$$\tilde{\boldsymbol{\gamma}} = (0, \dots, 0, \sqrt{(\gamma^1)^2 + \dots + (\gamma^{N-2})^2}, \gamma^{N-1}, \gamma^N).$$

Clearly $\tilde{\boldsymbol{\gamma}} \in W^{1,1}((0, 1), \mathbb{S}^{N-1})$. Since \mathbf{v}_0 and \mathbf{v}_1 belong to $\text{span}\{\mathbf{e}_{N-1}, \mathbf{e}_N\}$ and the projections of $\tilde{\boldsymbol{\gamma}}$ and $\boldsymbol{\gamma}$ onto $\text{span}\{\mathbf{e}_{N-1}, \mathbf{e}_N\}$ coincide, $\tilde{\boldsymbol{\gamma}} \in W^{1,1}((0, 1); \mathcal{P}_N(\mathbf{v}_0, \mathbf{v}_1))$ and the end-point conditions are satisfied. Therefore $\tilde{\boldsymbol{\gamma}} \in \Gamma_N(\mathbf{v}_0, \mathbf{v}_1)$. In addition, letting

$$\delta = (\gamma^1, \dots, \gamma^{N-2}),$$

we may apply the chain rule [Ambrosio and Dal Maso 1990, Corollary 3.2]: since $\delta \in W^{1,1}((0, 1); \mathbb{R}^{N-2})$ and $f(x) = |x|$ is a Lipschitz function with $f(0) = 0$, we have $|\delta| = f \circ \delta \in W^{1,1}((0, 1); \mathbb{R})$, for almost every $t \in (0, 1)$ the restriction of f to the affine space

$$T_t^\delta := \{y \in \mathbb{R}^{N-2} : y = \delta(t) + \eta \dot{\delta}(t) \text{ for some } \eta \in \mathbb{R}\}$$

is differentiable at $\delta(t)$, and finally

$$\frac{d}{dt}|\delta| = \nabla(f|_{T_t^\delta})(\delta(t)) \cdot \dot{\delta}(t) \quad \text{for a.e. } t \in (0, 1).$$

Since the Lipschitz constant of f is 1, we get that $|(d/dt)|\delta|| \leq |(d/dt)\delta|$. Hence $|(d/dt)\tilde{\boldsymbol{\gamma}}| \leq |(d/dt)\boldsymbol{\gamma}|$, which implies that $J_N[\mathbf{v}_0, \mathbf{v}_1](\tilde{\boldsymbol{\gamma}}) \leq J_N[\mathbf{v}_0, \mathbf{v}_1](\boldsymbol{\gamma})$, since $\tilde{\boldsymbol{\gamma}}^N = \boldsymbol{\gamma}^N$. Arguing as above, we also see that

$$\tilde{\boldsymbol{\gamma}} = (\sqrt{(\gamma^1)^2 + \dots + (\gamma^{N-2})^2}, \gamma^{N-1}, \gamma^N)$$

belongs to $\Gamma_3(\bar{\mathbf{v}}_0, \bar{\mathbf{v}}_1)$. Since $J_N[\mathbf{v}_0, \mathbf{v}_1](\tilde{\boldsymbol{\gamma}}) = J_3[\bar{\mathbf{v}}_0, \bar{\mathbf{v}}_1](\tilde{\boldsymbol{\gamma}})$, the proof is complete. □

Hereafter we let

$$\mathbf{v}_i := \bar{\mathbf{v}}_i, \quad J := J_3[\mathbf{v}_0, \mathbf{v}_1], \quad \mathcal{P} := \mathcal{P}_3(\mathbf{v}_0, \mathbf{v}_1), \quad \Gamma := \Gamma_3(\mathbf{v}_0, \mathbf{v}_1).$$

Because of (4-2) and Lemma 4.3, it suffices to prove that

$$\inf_{\boldsymbol{\gamma} \in \Gamma} J(\boldsymbol{\gamma}) \geq 2 \sin \theta_0. \tag{4-3}$$

Proving (4-3) is far from trivial, both since the functional is genuinely nonconvex (see Lemmas 4.9 and 4.10) and since the curves are constrained to an octant of the sphere. However, it is exactly for this reason that the lower bound holds:

Remark 4.4. In the extremal case $\theta_0 = \pi/4$, there are exactly two paths $\boldsymbol{\gamma}$ such that $J(\boldsymbol{\gamma}) = 2 \sin \theta_0$: the one given in Lemma 4.2, and the one which coincides with $\partial\mathcal{P}$ (see Section 1 or Lemma 4.14 with $\varphi_0 = 0$ and $\varphi_1 = \pi/2$). If the constraint is removed, the lower bound (4-3) does not hold any more: for instance, the curve

$$\boldsymbol{\gamma}(t) := \begin{cases} (0, \sin \theta, \cos \theta), & \theta = \theta_0 + 3t(\pi/2 - \theta_0) \in (\theta_0, \pi/2) & \text{if } 0 \leq t \leq \frac{1}{3}, \\ (\sin \varphi, \cos \varphi, 0), & \varphi = 3\pi(t - \frac{1}{3}) \in (0, \pi) & \text{if } \frac{1}{3} < t \leq \frac{2}{3}, \\ (0, -\sin \theta, \cos \theta), & \theta = \pi/2 + 3(t - \frac{2}{3})(\theta_0 - \pi/2) \in (\theta_0, \pi/2) & \text{if } \frac{2}{3} < t \leq 1 \end{cases}$$

is such that

$$J(\boldsymbol{\gamma}) = 2 \int_0^{1/3} \cos \theta |\dot{\theta}| dt = 2 \int_{\theta_0}^{\pi/2} \cos \theta d\theta = 2(1 - \sin \theta_0),$$

hence $J(\boldsymbol{\gamma}) = 2(1 - \sin \theta_0) < 2 \sin \theta_0$ if $\theta_0 > \pi/6$.

We will often use spherical coordinates centered at $(0, 0, 1)$:

$$X(\varphi, \theta) := (\sin \varphi \sin \theta, \cos \varphi \sin \theta, \cos \theta). \quad (4-4)$$

In this case $\boldsymbol{v}_0 = X(0, \theta_0)$, $\boldsymbol{v}_1 = X(\pi, \theta_0)$, the functional reads

$$J(\boldsymbol{\gamma}) = \int_0^1 \cos \theta(t) \sqrt{(\dot{\theta}(t))^2 + (\dot{\varphi}(t))^2 \sin^2 \theta(t)} dt, \quad \text{where } \boldsymbol{\gamma}(t) = X(\varphi(t), \theta(t)), \quad (4-5)$$

and the constraint $\boldsymbol{\gamma}(t) \in \mathcal{P}$ is equivalent to

$$\theta(t) \in [0, \pi/2], \quad \theta(t) \leq \arctan \frac{1}{\tan \theta_0 |\cos \varphi(t)|} =: \theta^*(\varphi(t)). \quad (4-6)$$

It is convenient to cut-off from \mathcal{P} a neighborhood of $z = 0$: in this way, the new constraint has a smooth boundary and the density of J does not degenerate. Thus, let $\theta_\varepsilon^* \in C^\infty(\mathbb{R})$ be such that

$$\begin{aligned} \theta_\varepsilon^* &\text{ is } \pi\text{-periodic, even w.r.t. } \pi/2, \text{ increasing in } (0, \pi/2), \\ \theta_\varepsilon^*(\varphi) &= \theta^*(\varphi) \text{ if } |\pi/2 - \varphi| \geq \varepsilon, \theta_\varepsilon^* < \pi/2, \text{ and } |(\theta_\varepsilon^*)'| \leq C \end{aligned} \quad (4-7)$$

for some ε -independent positive constant C . Note that here and after primes denote differentiation with respect to φ , and that the latter property of θ_ε^* may be fulfilled since θ^* is Lipschitz-continuous. Now let

$$\begin{aligned} \mathcal{P}_\varepsilon &:= \{X(\varphi, \theta) : \varphi \in [0, 2\pi], 0 \leq \theta \leq \theta_\varepsilon^*(\varphi)\}, \\ \Gamma_\varepsilon &:= \{\boldsymbol{\gamma} \in W^{1,1}((0, 1); \mathcal{P}_\varepsilon) : \boldsymbol{\gamma}(0) = \boldsymbol{v}_0, \boldsymbol{\gamma}(1) = \boldsymbol{v}_1\}. \end{aligned}$$

In what follows, $\omega(\varepsilon)$ denotes a generic positive universal function which goes to zero as $\varepsilon \rightarrow 0$. The next lemma shows that we may equivalently work on \mathcal{P}_ε :

Lemma 4.5. *Assume that*

$$\inf_{\boldsymbol{\gamma} \in \Gamma_\varepsilon} J(\boldsymbol{\gamma}) \geq 2 \sin \theta_0 - \omega(\varepsilon). \quad (4-8)$$

Then (4-3) holds true, and therefore so does Theorem 4.1.

Proof. Given $\boldsymbol{\gamma} \in \Gamma$, we replace the parts of $\boldsymbol{\gamma}$ which enter into $\mathcal{P} \setminus \mathcal{P}_\varepsilon$ by arcs of $\partial \mathcal{P}_\varepsilon$. More precisely, let

$$I_\varepsilon = \{t \in (0, 1) : \boldsymbol{\gamma}(t) \in \mathcal{P} \setminus \mathcal{P}_\varepsilon\}.$$

Since the spherical coordinates (4-4) are a bijection away from the north pole $(0, 0, 1)$, in I_ε we may define $\varphi(t)$ and $\theta(t)$ through $\boldsymbol{\gamma}(t) =: X(\varphi(t), \theta(t))$. Then we let

$$\boldsymbol{\gamma}_\varepsilon(t) = \begin{cases} \boldsymbol{\gamma}(t) & \text{if } t \notin I_\varepsilon, \\ (\varphi(t), \theta_\varepsilon^*(\varphi(t))) & \text{if } t \in I_\varepsilon. \end{cases}$$

It follows from (4-7) that

$$\left| \frac{\pi}{2} - \theta_\varepsilon^*(\varphi(t)) \right| = \omega(\varepsilon). \tag{4-9}$$

We may now estimate

$$J(\boldsymbol{\gamma}) - J(\boldsymbol{\gamma}_\varepsilon) \stackrel{(4-5)}{\geq} - \int_{I_\varepsilon} \cos \theta_\varepsilon^*(\varphi) |\dot{\varphi}| \sqrt{(\theta_\varepsilon^{*\prime})^2 + \sin^2 \theta_\varepsilon^*(\varphi)} dt \stackrel{(4-7)}{\geq} -\omega(\varepsilon) \int_0^1 |\dot{\varphi}| dt. \tag{4-9}$$

Therefore

$$J(\boldsymbol{\gamma}) \stackrel{(4-8)}{\geq} 2 \sin \theta_0 - \omega(\varepsilon) \left(1 + \int_0^1 |\dot{\varphi}| dt \right).$$

Passing to the limit as $\varepsilon \rightarrow 0$, the arbitrariness of $\boldsymbol{\gamma} \in \Gamma$ yields (4-3). □

The rest of the section will be concerned with the proof of (4-8). Let

$$\Gamma_\varepsilon(\mathbf{w}_0, \mathbf{w}_1) := \{ \boldsymbol{\gamma} \in W^{1,1}((0, 1); \mathcal{P}_\varepsilon) : \boldsymbol{\gamma}(0) = \mathbf{w}_0, \boldsymbol{\gamma}(1) = \mathbf{w}_1 \} \quad \text{for } \mathbf{w}_0, \mathbf{w}_1 \in \mathcal{P}_\varepsilon.$$

Lemma 4.6. *For any $\mathbf{w}_0, \mathbf{w}_1 \in \mathcal{P}_\varepsilon$, there exists a minimizer $\boldsymbol{\gamma}$ of J in $\Gamma_\varepsilon(\mathbf{w}_0, \mathbf{w}_1)$. Furthermore $\boldsymbol{\gamma}$ lies in $W^{1,\infty}((0, 1); \mathbb{R}^3)$, satisfies $\gamma^3 |\dot{\boldsymbol{\gamma}}| = J(\boldsymbol{\gamma})$ a.e. in $[0, 1]$, and is also a minimizer of*

$$E(\boldsymbol{\gamma}) := \int_0^1 (\gamma^3(t))^2 |\dot{\boldsymbol{\gamma}}(t)|^2 dt$$

among all $\boldsymbol{\gamma} \in \Gamma_\varepsilon(\mathbf{w}_0, \mathbf{w}_1) \cap H^1((0, 1); \mathbb{R}^3)$.

Though we could appeal to general results on geodesics for Riemannian manifolds with boundary (see [Alexander et al. 1993] and the references therein), we prefer to give a self-contained proof.

Proof. We preliminarily observe that

for all $\boldsymbol{\gamma} \in \Gamma_\varepsilon(\mathbf{w}_0, \mathbf{w}_1)$, there exists $\tilde{\boldsymbol{\gamma}} \in \Gamma_\varepsilon(\mathbf{w}_0, \mathbf{w}_1) \cap W^{1,\infty}((0, 1); \mathbb{R}^3)$

$$\text{such that } \tilde{\boldsymbol{\gamma}}^3(t) |\dot{\tilde{\boldsymbol{\gamma}}}(t)| = L := J(\boldsymbol{\gamma}) \text{ for a.e. } t \in [0, 1]. \tag{4-10}$$

To see this, let

$$s(t) = \frac{1}{L} \int_0^t \gamma^3(\tau) |\dot{\boldsymbol{\gamma}}(\tau)| d\tau. \tag{4-11}$$

Obviously $s \in W^{1,1}([0, 1]; [0, 1])$, s is nondecreasing, and $s(t_1) = s(t_2)$ if and only if $\boldsymbol{\gamma}(t) = \boldsymbol{\gamma}(t_1)$ in $[t_1, t_2]$. Therefore, for any $\sigma \in [0, 1]$, either there exists a unique $t(\sigma)$ such that $s(t(\sigma)) = \sigma$, or there exists an interval I_σ such that $s(t) = \sigma$ for all $t \in I_\sigma$, and in this case we let, for example, $t(\sigma) = \inf I_\sigma$, so that again $s(t(\sigma)) = \sigma$. Now let $\tilde{\boldsymbol{\gamma}}(\sigma) := \boldsymbol{\gamma}(t(\sigma))$. By construction,

$$\boldsymbol{\gamma}(t) = \tilde{\boldsymbol{\gamma}}(s(t)) \quad \text{for all } t \in [0, 1]. \tag{4-12}$$

Note that $\tilde{\boldsymbol{\gamma}} \in W^{1,\infty}((0, 1); \mathcal{P}_\varepsilon)$. Indeed,

$$\begin{aligned} |\tilde{\boldsymbol{\gamma}}(\sigma_1) - \tilde{\boldsymbol{\gamma}}(\sigma_2)| &= |\boldsymbol{\gamma}(t(\sigma_1)) - \boldsymbol{\gamma}(t(\sigma_2))| \leq \int_{t(\sigma_1)}^{t(\sigma_2)} |\dot{\boldsymbol{\gamma}}(\tau)| d\tau \\ &\stackrel{(4-11)}{\leq} \frac{L}{\inf_{\tau \in [t(\sigma_1), t(\sigma_2)]} \gamma^3(\tau)} |s(t(\sigma_1)) - s(t(\sigma_2))| \\ &\stackrel{(4-7)}{\leq} \frac{L}{\omega(\varepsilon)} |\sigma_1 - \sigma_2|. \end{aligned} \quad (4-13)$$

Hence, it follows from (4-12) and the chain rule formula given in [Ambrosio et al. 2000, Theorem 3.101] that

$$\dot{\boldsymbol{\gamma}}(t) = \frac{d\tilde{\boldsymbol{\gamma}}}{ds}(s(t))\dot{s}(t) \quad \text{in } L^1((0, 1)). \quad (4-14)$$

Therefore,

$$L = \int_0^1 \gamma^3(t) |\dot{\boldsymbol{\gamma}}(t)| dt \stackrel{(4-14)}{\stackrel{(4-12)}{=}} \int_0^1 \tilde{\gamma}^3(s(t)) \left| \frac{d\tilde{\boldsymbol{\gamma}}}{ds}(s(t)) \right| \dot{s}(t) dt = \int_0^1 \tilde{\gamma}^3(s) \left| \frac{d\tilde{\boldsymbol{\gamma}}}{ds}(s) \right| ds. \quad (4-15)$$

On the other hand, given $s \in [0, 1]$ and $\varepsilon > 0$, let $s_1, s_2 \in [0, 1]$ with $|s_i - s| < \varepsilon$. Then

$$\tilde{\gamma}^3(s) |\tilde{\boldsymbol{\gamma}}(s_1) - \tilde{\boldsymbol{\gamma}}(s_2)| \stackrel{(4-13)}{\leq} \frac{\tilde{\gamma}^3(s)}{\inf_{\tau \in [t(s_1), t(s_2)]} \gamma^3(\tau)} L |s_2 - s_1|. \quad (4-16)$$

If $\tau \in [t(s_1), t(s_2)]$, then, by the monotonicity of s and since $s(\tau(s)) = s$, we have $s(\tau) \in [s_1, s_2]$. Hence

$$\inf_{\tau \in [t(s_1), t(s_2)]} \gamma^3(\tau) \stackrel{(4-12)}{=} \inf_{\tau \in [t(s_1), t(s_2)]} \tilde{\gamma}^3(s(\tau)) \geq \inf_{s \in [s_1, s_2]} \tilde{\gamma}^3(s). \quad (4-17)$$

Combining (4-16) and (4-17) and passing to the limit as $\varepsilon \rightarrow 0$, we obtain

$$\tilde{\gamma}^3(s) \left| \frac{d\tilde{\boldsymbol{\gamma}}}{ds}(s) \right| \leq L \quad \text{for a.e. } s \in [0, 1],$$

which together with (4-15) concludes the proof of the claim (4-10).

We consider the functional E defined on $G_\varepsilon(\mathbf{w}_0, \mathbf{w}_1) := \Gamma_\varepsilon(\mathbf{w}_0, \mathbf{w}_1) \cap H^1((0, 1); \mathbb{R}^3)$. By the Cauchy–Schwarz inequality,

$$(J(\boldsymbol{\gamma}))^2 \leq E(\boldsymbol{\gamma}) \quad \text{for all } \boldsymbol{\gamma} \in G_\varepsilon(\mathbf{w}_0, \mathbf{w}_1). \quad (4-18)$$

Hence $\inf E(\boldsymbol{\gamma}) \geq \inf (J(\boldsymbol{\gamma}))^2$. On the other hand, let $\boldsymbol{\gamma}_n$ be a minimizing sequence for J , and let $\tilde{\boldsymbol{\gamma}}_n$ be as given by (4-10): then $E(\tilde{\boldsymbol{\gamma}}_n) = (J(\boldsymbol{\gamma}_n))^2$, which means that $\inf E \leq \inf J^2$. Therefore,

$$\inf_{\boldsymbol{\gamma} \in G_\varepsilon(\mathbf{w}_0, \mathbf{w}_1)} E(\boldsymbol{\gamma}) = \inf_{\boldsymbol{\gamma} \in \Gamma_\varepsilon(\mathbf{w}_0, \mathbf{w}_1)} (J(\boldsymbol{\gamma}))^2.$$

The inf on the left-hand side is attained. Indeed, let $\boldsymbol{\gamma}_n$ be a minimizing sequence. By the coercivity of E ensured by the definition of \mathcal{P}_ε , a subsequence (not relabeled) exists such that $\boldsymbol{\gamma}_n \rightharpoonup \boldsymbol{\gamma}$ weakly in $H^1((0, 1); \mathcal{P}_\varepsilon)$ and in $C([0, 1]; \mathcal{P}_\varepsilon)$. Therefore $E(\boldsymbol{\gamma}) \leq \liminf_{n \rightarrow +\infty} E(\boldsymbol{\gamma}_n)$.

Let $\boldsymbol{\gamma}_0$ be a minimizer of E , and let $\tilde{\boldsymbol{\gamma}}_0$ be as given by (4-10). Then

$$E(\boldsymbol{\gamma}_0) \stackrel{(4-18)}{\geq} (J(\boldsymbol{\gamma}_0))^2 = (J(\tilde{\boldsymbol{\gamma}}_0))^2 = E(\tilde{\boldsymbol{\gamma}}_0),$$

that is, $\tilde{\boldsymbol{\gamma}}_0$ is also a minimizer of E , and

$$(J(\boldsymbol{\gamma}))^2 = (J(\tilde{\boldsymbol{\gamma}}))^2 = E(\tilde{\boldsymbol{\gamma}}) \geq E(\tilde{\boldsymbol{\gamma}}_0) = (J(\tilde{\boldsymbol{\gamma}}_0))^2 \quad \text{for all } \boldsymbol{\gamma} \in \Gamma_\varepsilon(\mathbf{w}_0, \mathbf{w}_1),$$

hence $\tilde{\boldsymbol{\gamma}}_0$ (or $\boldsymbol{\gamma}_0$) is a minimizer of J . Therefore J has a minimizer too. □

The rest of the section is concerned with estimating the length of a minimizer of J in Γ_ε as given by Lemma 4.6, a *shortest path* in what follows. Our first observation concerns those shortest paths which pass through the north pole:

Lemma 4.7. *If a shortest path $\boldsymbol{\gamma}$ passes through $(0, 0, 1)$, then $J(\boldsymbol{\gamma}) \geq 2 \sin \theta_0$.*

Proof. Let t_0 and t_1 be the first, respectively the last, time in which $\boldsymbol{\gamma} = (0, 0, 1)$. Then, using the spherical coordinates (4-4),

$$\begin{aligned} J(\boldsymbol{\gamma}) &\geq \int_0^{t_0} \cos \theta \sqrt{(\dot{\theta})^2 + (\dot{\varphi})^2 \sin^2 \theta} dt + \int_{t_1}^1 \cos \theta \sqrt{(\dot{\theta})^2 + (\dot{\varphi})^2 \sin^2 \theta} dt \\ &\geq \int_0^{t_0} \cos \theta |\dot{\theta}| dt + \int_{t_1}^1 \cos \theta |\dot{\theta}| dt = \int_0^{t_0} \left| \frac{d}{dt} \sin \theta \right| dt + \int_{t_1}^1 \left| \frac{d}{dt} \sin \theta \right| dt, \end{aligned}$$

and the lemma follows, since $\theta(t_0) = \theta(t_1) = 0$ and $\theta(0) = \theta(1) = \theta_0$. □

We may therefore restrict our attention to shortest paths not passing through the north pole. There, the spherical coordinates (4-4) are a diffeomorphism. In fact, we may also restrict our attention to those paths for which φ is nondecreasing and which are symmetric with respect to $\varphi = \pi/2$. In what follows, we shall call them *symmetric* shortest paths.

Lemma 4.8. *Let $\boldsymbol{\gamma} = X(\varphi, \theta)$ be a shortest path not passing through $(0, 0, 1)$. Then $\varphi \in [0, \pi]$ and φ is nondecreasing. Moreover, there exists a shortest path $\tilde{\boldsymbol{\gamma}} = X(\tilde{\varphi}, \tilde{\theta})$ not passing through $(0, 0, 1)$ such that $\tilde{\varphi}$ is symmetric with respect to $\pi/2$:*

$$\{(\tilde{\varphi}(t), \tilde{\theta}(t)) : t \in [0, 1]\} = \{(\pi - \tilde{\varphi}(t), \tilde{\theta}(t)) : t \in [0, 1]\}.$$

Proof. Without loss of generality, $\varphi(0) = 0$ and $\varphi(1) = (2k + 1)\pi$ with $k \geq 0$. It is straightforward to see that $\max\{\varphi, 0\}$ and $\min\{\varphi, \pi\}$ both decrease the value of J , hence $k = 0$. Analogously, if $t_0 < t_1 < t_2$ are such that $\varphi(t_1) < \varphi(t_2) = \varphi(t_0)$, then replacing φ with $\varphi(t_0)$ in (t_0, t_2) decreases the value of J . Therefore, φ is nondecreasing along a shortest path.

In order to construct $\tilde{\boldsymbol{\gamma}}$, we claim that

$$J_1 := \int_0^{t_*} \cos \theta \sqrt{(\dot{\theta})^2 + (\dot{\varphi})^2 \sin^2 \theta} dt = \int_{t_*}^1 \cos \theta \sqrt{(\dot{\theta})^2 + (\dot{\varphi})^2 \sin^2 \theta} dt =: J_2 \quad (4-19)$$

for any $t^* \in (0, 1)$ such that $\varphi(t^*) = \pi/2$. Suppose by contradiction that (4-19) does not hold. Then, without loss of generality, we can suppose $J_1 < J_2$. We define $\tilde{\boldsymbol{\gamma}}(t) = X(\tilde{\varphi}(t), \tilde{\theta}(t))$, where

$$\tilde{\theta}(t) = \begin{cases} \theta(t) & \text{if } t \leq t_*, \\ \theta\left(\frac{t_*(1-t)}{1-t_*}\right) & \text{if } t > t_*, \end{cases} \quad \text{and} \quad \tilde{\varphi}(t) = \begin{cases} \varphi(t) & \text{if } t \leq t_*, \\ \pi - \varphi\left(\frac{t_*(1-t)}{1-t_*}\right) & \text{if } t > t_*. \end{cases} \quad (4-20)$$

Then, by letting $\hat{t} = t_*(1-t)/(1-t_*)$ and using the 1-homogeneity of the integrands with respect to t , we see that

$$\begin{aligned} J(\tilde{\boldsymbol{\gamma}}) &= J_1 + \frac{t_*}{1-t_*} \int_{t_*}^1 \cos \theta(\hat{t}) \sqrt{(\dot{\theta}(\hat{t}))^2 + (\dot{\varphi}(\hat{t}))^2 \sin^2 \theta(\hat{t})} dt \\ &= J_1 + \int_0^{t_*} \cos \theta(\hat{t}) \sqrt{(\dot{\theta}(\hat{t}))^2 + (\dot{\varphi}(\hat{t}))^2 \sin^2 \theta(\hat{t})} d\hat{t} = 2J_1 \\ &< J_1 + J_2 = J(\boldsymbol{\gamma}), \end{aligned} \quad (4-21)$$

a contradiction, since $\boldsymbol{\gamma}$ is a shortest path. Therefore (4-19) holds. Then, defining $\tilde{\boldsymbol{\gamma}}(t) = X(\tilde{\varphi}(t), \tilde{\theta}(t))$ as in (4-20), it follows from (4-21) that $J(\tilde{\boldsymbol{\gamma}}) = 2J_1 = J_1 + J_2 = J(\boldsymbol{\gamma})$, hence $\tilde{\boldsymbol{\gamma}}$ is also a shortest path. \square

We now characterize arcs of shortest paths contained in $\mathring{\mathcal{P}}_\varepsilon$.

Lemma 4.9. *Let $\boldsymbol{\gamma}$ be a shortest path not passing through $(0, 0, 1)$ and let (t_0, t_1) be an interval in which $\boldsymbol{\gamma}|_{(t_0, t_1)} \subset \mathring{\mathcal{P}}_\varepsilon$. Then*

$$\frac{\cos \theta(t) \sin^2 \theta(t) \dot{\varphi}(t)}{\sqrt{(\dot{\theta}(t))^2 + (\dot{\varphi}(t))^2 \sin^2 \theta(t)}} = K \quad \text{for all } t \in (t_0, t_1) \quad (4-22)$$

for some $K \in [0, 1/2]$. If $K = 0$, then φ is constant. If $K > 0$, then φ is strictly increasing, the function

$$(\varphi(t_0), \varphi(t_1)) =: I \ni \varphi \mapsto \theta(t(\varphi)) \quad (4-23)$$

is a smooth solution of

$$\theta'' \sin \theta \cos \theta = (\theta')^2 (\cos^2 \theta + \cos(2\theta)) + \cos(2\theta) \sin^2 \theta \quad (4-24)$$

with

$$\sin^2 \theta (\cos^2 \theta \sin^2 \theta - K^2) = K^2 (\theta')^2, \quad (4-25)$$

and

$$J(\boldsymbol{\gamma}|_{\chi(t_0, t_1)}) = \int_{\varphi(t_0)}^{\varphi(t_1)} \cos \theta \sqrt{(\theta')^2 + \sin^2 \theta} d\varphi. \quad (4-26)$$

Proof. Up to a linear reparametrization, $\boldsymbol{\gamma}$ is also a minimizer of J in $\Gamma_\varepsilon(\boldsymbol{\gamma}(t_0), \boldsymbol{\gamma}(t_1))$. Hence, by Lemma 4.6, it is also a minimizer of E in $\Gamma_\varepsilon(\boldsymbol{\gamma}(t_0), \boldsymbol{\gamma}(t_1)) \cap H^1((0, 1); \mathbb{R}^3)$. Since it does not touch the north pole, we can write $\boldsymbol{\gamma} = X(\varphi, \theta)$ with φ and θ Lipschitz, and

$$E(\boldsymbol{\gamma}|_{\chi(t_0, t_1)}) = \int_{t_0}^{t_1} \cos^2 \theta (\dot{\theta}^2 + \sin^2 \theta \dot{\varphi}^2) dt.$$

Taking the first variation with respect to φ , we obtain $\sin^2 \theta \cos^2 \theta \dot{\varphi} = H$, and, recalling that $\gamma^3 |\dot{\boldsymbol{\gamma}}|$ is constant, (4-22) follows. Since $\sin \theta > 0$ ($\boldsymbol{\gamma}$ does not cross the north pole), $\cos \theta > 0$ ($\boldsymbol{\gamma} \in P_\varepsilon$), and

φ is nondecreasing (by Lemma 4.8), we see that $K \geq 0$. If $K = 0$, then φ is constant. If $K > 0$, then $\dot{\varphi} > 0$ in (t_0, t_1) and we may use φ as independent variable: letting θ be as in (4-23), we have $\theta' = d\theta/d\varphi = \dot{\theta}/\dot{\varphi} \in L^\infty((t_0, t_1))$ (because of (4-22)). Then (4-26) follows at once from (4-5) and the definition of K may be rewritten as

$$\frac{\cos \theta \sin^2 \theta}{\sqrt{(\theta')^2 + \sin^2 \theta}} = K, \tag{4-27}$$

which is equivalent to (4-25). From (4-25) one sees immediately that $K \leq 1/2$. Differentiating (4-27), we obtain (4-24) in the sense of distributions, and a bootstrap argument starting from $\theta \in W^{1,\infty}((t_0, t_1))$ yields smoothness. □

If $\gamma = X(\varphi, \theta(\varphi)) : (t_0, t_1) \rightarrow \mathbb{S}^2$ is a curve which does not pass through $(0, 0, 1)$ and such that $\varphi \in I := (\varphi(t_0), \varphi(t_1))$ is strictly increasing, then, following (4-26), we hereafter write (with a slight abuse of notation)

$$J(\gamma \chi_{(t_0, t_1)}) = J_I(\theta) := \int_I \cos \theta \sqrt{(\theta')^2 + \sin^2 \theta} \, d\varphi, \quad J(\theta) := J_{(0, \pi)}(\theta).$$

In view of Lemma 4.9, it is convenient to state a few properties of the solutions to (4-24), some of which are visualized in Figure 1.

Lemma 4.10. *Let θ be any solution of (4-24) such that $\theta \in (0, \pi/2)$ at some point of its domain. Then:*

- (a) θ is globally defined, periodic, and $\theta \in (0, \pi/2)$;
- (b) within a period, θ has a unique local (and therefore global) maximum, $\theta_M \geq \pi/4$, and a unique local (and therefore global) minimum, $\theta_m = \pi/2 - \theta_M$, and it is symmetric with respect to its maximum (minimum) point;
- (c) the period P is larger than π ;
- (d) the length of each interval in which $\theta \leq \pi/4$ is at least $\pi/\sqrt{2}$;
- (e) θ' has a unique local (and therefore global) maximum and a unique local (and therefore global) minimum.

Proof. (a) and (b) easily follow from (4-25) rewritten as

$$(\theta')^2 = \frac{1}{K^2} \sin^2 \theta (\sin^2 \theta \cos^2 \theta - K^2) =: f_K(\theta), \quad K \in [0, 1/2] \tag{4-28}$$

and plotted in the phase space (see Figure 1). We just observe explicitly that, since $\theta' = 0$ at the extremal values of θ , we can characterize K from (4-25) as

$$K = \cos \theta_m \sin \theta_m = \cos \theta_M \sin \theta_M, \tag{4-29}$$

which explains why $\theta_M = \pi/2 - \theta_m$. Also (e) follows immediately from (4-28), since after differentiation we see that

$$2\theta'' = f'_K(\theta),$$




Figure 1. The phase plane (θ, θ') .

whence the arrows in Figure 1.

To prove (c), we let φ_m and φ_M be a point of minimum and of maximum, respectively, chosen such that no local extremum exists in between. Then, in view of (b),

$$\frac{P}{2} = \int_{\varphi_m}^{\varphi_M} d\varphi \stackrel{(4-25)}{=} \int_{\varphi_m}^{\varphi_M} \frac{K\theta'}{\sin\theta \sqrt{\cos^2\theta \sin^2\theta - K^2}} d\varphi = \int_{\theta_m}^{\theta_M} \frac{K}{\sin\theta \sqrt{\cos^2\theta \sin^2\theta - K^2}} d\theta.$$

We now observe that

$$K \stackrel{(4-29)}{=} \cos\theta_M \sin\theta_M = \cos\theta_M \cos\theta_m \geq \cos\theta_M \cos\theta \quad \text{for all } \theta \in (\theta_m, \theta_M).$$

Therefore,

$$\frac{P}{2} \geq \cos\theta_M \int_{\theta_m}^{\theta_M} \frac{\cos\theta}{\sin\theta \sqrt{\cos^2\theta \sin^2\theta - K^2}} d\theta,$$

whose primitives may be computed explicitly:

$$\cos\theta_M \int \frac{\cos\theta}{\sin\theta \sqrt{\cos^2\theta \sin^2\theta - K^2}} d\theta = \frac{1}{2 \sin\theta_M} \arcsin \frac{\sin^2\theta - 2 \sin^2\theta_M \cos^2\theta_M}{\sin^2\theta |1 - 2 \cos^2\theta_M|}.$$

Hence

$$\frac{P}{2} \geq \frac{1}{2 \sin\theta_M} \left(\frac{\pi}{2} + \frac{\pi}{2} \right) = \frac{\pi}{2 \sin\theta_M} > \frac{\pi}{2},$$

which proves (c).

To prove (d), let φ_m be a minimum point and let φ_* be the closest point to φ_m such that $\varphi_m \leq \varphi_*$ and $\theta(\varphi_*) = \pi/4$. By (b), the length of the interval within a period where $\theta \leq \pi/4$ is exactly

$$2 \int_{\varphi_m}^{\varphi_*} d\varphi \stackrel{(4-25)}{=} 2 \int_{\theta_m}^{\pi/4} \frac{K}{\sin\theta \sqrt{\cos^2\theta \sin^2\theta - K^2}} d\theta.$$

Since $\cos \theta \geq 1/\sqrt{2}$ if $\theta \in [0, \pi/4]$,

$$2 \int_{\varphi_m}^{\varphi_*} d\varphi \geq \sqrt{2} \int_{\theta_m}^{\pi/4} \frac{K}{\sin \theta \cos \theta \sqrt{\cos^2 \theta \sin^2 \theta - K^2}} d\theta.$$

The primitives of the right-hand side may be computed explicitly (via the substitution $z = \sin^2(2\theta)$):

$$\int \frac{K}{\sin \theta \cos \theta \sqrt{\cos^2 \theta \sin^2 \theta - K^2}} d\theta = -\arctan \frac{2K \cos(2\theta)}{\sqrt{\sin^2(2\theta) - 4K^2}} + C. \tag{4-30}$$

After a substitution we get (d). □

Lemma 4.11. *Let γ be a symmetric shortest path not passing through $(0, 0, 1)$ and let $\gamma = X(\varphi, \theta)$. If $t_1 \in [0, 1]$ is such that $\theta(t_1) < \pi/6$ and $\varphi(t_1) < \pi/2$, then $\theta(t) < \pi/6$ as long as $\varphi(t) < \pi - \varphi(t_1)$.*

Proof. Let $w = \sin \theta$ and let $t_2 > t_1$ be the first time in which $\varphi(t_2) = \pi - \varphi(t_1)$. We have

$$J(\gamma \chi_{(t_1, t_2)}) = \int_{t_1}^{t_2} \sqrt{(\dot{w}(t))^2 + (\dot{\varphi}(t))^2 w(t)^2 (1 - w(t)^2)} dt.$$

By assumption, $w(t_1) < \frac{1}{2}$. If there is an interval $\tilde{I} \subset [t_1, t_2]$ where $w(t) > \frac{1}{2}$, then a symmetrization of w with respect to $\frac{1}{2}$ would strictly decrease the value of J , since

$$(1 - w)^2(1 - (1 - w)^2) - w^2(1 - w^2) = 2w(1 - w)(1 - 2w) < 0 \quad \text{if } w \in (1/2, 1).$$

This contradicts that γ is a shortest path and thus proves the lemma. □

We are now ready to exclude shortest paths which are contained in $\mathring{\mathcal{P}}_\varepsilon$:

Lemma 4.12. *There is no symmetric shortest path γ not passing through $(0, 0, 1)$ such that $\gamma((0, 1)) \subset \mathring{\mathcal{P}}_\varepsilon$.*

Proof. Assume for a contradiction that such a γ exists. We will argue that $J(\gamma) > 2 \sin \theta_0$, in contradiction with Lemma 4.2 (note that $\gamma_{\min} \subseteq \mathcal{P}_\varepsilon$ for all ε).

Since φ has to travel from 0 to π , it can not be constant in $[0, 1]$. Then, it follows from Lemma 4.9 that $\gamma(t) = X(\varphi(t), \theta(t))$, where $\varphi \mapsto \theta(t(\varphi))$ is a smooth solution of (4-24) such that $\theta(0) = \theta(\pi) = \theta_0$. Since γ is symmetric, we have $\theta'(\pi/2) = 0$. Because of (b) and (c) in Lemma 4.10, θ is monotone in $(0, \pi/2)$. Hence, letting $\theta_1 = \theta(\pi/2)$, we have

$$K \stackrel{(4-25)}{=} K(\theta_1) = \cos \theta_1 \sin \theta_1 \quad \text{and} \quad \cos^2 \theta \sin^2 \theta \geq K^2 \quad \text{for all } t \in (0, 1). \tag{4-31}$$

We claim that $\theta_1 > \pi/4$. If not, it follows from (4-31) that $\theta_1 \leq \theta_0$. Hence θ is nonincreasing in $(0, \pi/2)$, and

$$\begin{aligned} \frac{\pi}{2} &\stackrel{(4-25)}{=} \int_0^{\pi/2} \frac{K\theta'}{\sin \theta \sqrt{\cos^2 \theta \sin^2 \theta - K^2}} d\varphi = \int_{\theta_1}^{\theta_0} \frac{K}{\sin \theta \sqrt{\cos^2 \theta \sin^2 \theta - K^2}} d\theta \\ &\leq \cos(\theta_1) \int_{\theta_1}^{\pi/4} \frac{K}{\sin \theta \cos \theta \sqrt{\cos^2 \theta \sin^2 \theta - K^2}} d\theta. \end{aligned} \tag{4-32}$$

By (4-30), we would have $\pi \leq \pi \cos(\theta_1)$, a contradiction. Hence $\theta_1 > \pi/4$.

We note the obvious bound

$$\frac{1}{2}J(\theta) \geq \int_0^{\pi/2} \cos \theta \sin \theta \, d\varphi \stackrel{(4-31)}{\geq} \frac{\pi}{2} \sin \theta_1 \cos \theta_1.$$

Hence we are done if

$$\frac{\pi}{2} \sin \theta_1 \cos \theta_1 > \sin \theta_0,$$

that is, if

$$\theta_0 < \arcsin\left(\frac{\pi}{2} \sin \theta_1 \cos \theta_1\right) = \arcsin\left(\frac{\pi}{4} \sin(2\theta_1)\right). \quad (4-33)$$

We claim that (4-33) does hold. If not, recalling Lemma 4.11, we would have

$$\theta_0 \geq \max\left\{\arcsin\left(\frac{\pi}{4} \sin(2\theta_1)\right), \frac{\pi}{6}\right\} =: f(\theta_1).$$

Then, arguing as in (4-32), we write

$$\begin{aligned} \frac{\pi}{2} &= \int_{\theta_0}^{\theta_1} \frac{K}{\sin \theta \sqrt{\cos^2 \theta \sin^2 \theta - K^2}} \, d\theta \\ &< \cos \frac{\pi}{6} \int_{f(\theta_1)}^{\pi/4} \frac{K}{\sin \theta \cos \theta \sqrt{\cos^2 \theta \sin^2 \theta - K^2}} \, d\theta + \cos \frac{\pi}{4} \int_{\pi/4}^{\theta_1} \frac{K}{\sin \theta \cos \theta \sqrt{\cos^2 \theta \sin^2 \theta - K^2}} \, d\theta \\ &= \frac{\sqrt{3}}{2} \arctan \frac{\sin \theta_1 \cos \theta_1 \cos(2f(\theta_1))}{\sqrt{\sin^2(2f(\theta_1)) - 4 \sin^2 \theta_1 \cos^2 \theta_1}} + \frac{\sqrt{2}}{2} \frac{\pi}{2} =: F(\theta_1). \end{aligned}$$

It is now a calculus exercise to check that F is increasing in $(\pi/4, \bar{\theta}) := (0, \frac{1}{2}(\pi - \arcsin(2/\pi)))$ and decreasing in $(\bar{\theta}, \pi/2)$: therefore F has a global maximum at $\bar{\theta}$, with $F(\bar{\theta}) < \pi/2$. Since this is impossible, (4-33) holds and the proof is complete. \square

The rest of the section is concerned with estimating the length of candidate symmetric shortest paths which intersect $\partial\mathcal{P}_\varepsilon$ (and do not pass through the north pole). We firstly infer some properties of those candidate shortest paths which reach $\partial\mathcal{P}_\varepsilon$.

Lemma 4.13. *Let $\theta_0 < \pi/4$, let ε be sufficiently small, and let $\boldsymbol{\gamma} = X(\varphi, \theta)$ be a symmetric shortest path not intersecting the north pole. If $t_1 > 0$ exists such that $\boldsymbol{\gamma}(t_1) \in \partial\mathcal{P}_\varepsilon$ and $\boldsymbol{\gamma}(t) \in \overset{\circ}{\mathcal{P}}_\varepsilon$ in $[0, t_1)$, then:*

- (i) $\theta(t) \geq \pi/6$ for all $t \in [0, t_1)$;
- (ii) θ is increasing in $[0, t_1)$;
- (iii) $\varphi(t_1) \leq \pi/2 - \varepsilon$.

Proof. (i) follows immediately from Lemma 4.11.

To prove (ii), we note that by Lemma 4.9, (4-22) holds in $[0, t_1)$. Let $\varphi_1 = \varphi(t_1)$. By symmetry, $\varphi_1 \leq \pi/2$. If $K = 0$, we would have $\varphi(t) = \varphi_1$ in $(0, t_1)$: since $\boldsymbol{\gamma}$ does not reach the north pole, this means that $\varphi_1 = 0$ and θ is increasing from θ_0 up to $\theta(t_1) = \pi/2 - \theta_0$. If instead $K > 0$, then (4-23) holds in $(0, \varphi_1)$. We will prove that $\theta' \geq 0$ in $(0, \varphi_1)$, which implies (ii). Assume by contradiction that $\theta' < 0$ somewhere in $(0, \varphi_1)$. Then, by Lemma 4.10(b), there exists $\varphi_2 \in (0, \varphi_1)$ such that $\theta(\varphi_2) = \theta_m$. By

Lemma 4.10(d) and since $\theta(\varphi_1) > \pi/4$, we have $\varphi_1 \geq \varphi_1 - \varphi_2 \geq \pi/(2\sqrt{2})$. Then, since θ_ε^* is increasing in $(0, \pi/2)$ and provided ε is sufficiently small,

$$\theta_1 := \theta(\varphi_1) = \theta_\varepsilon^*(\varphi_1) \geq \theta_\varepsilon^*\left(\frac{\pi}{2\sqrt{2}}\right) = \theta^*\left(\frac{\pi}{2\sqrt{2}}\right) \stackrel{(4-6)}{\geq} \arctan \frac{1}{\cos(\pi/(2\sqrt{2}))} > \frac{\pi}{3}.$$

By (4-25), this implies that $\sin \theta_m \cos \theta_m \leq \sin \theta_1 \cos \theta_1 < \sqrt{3}/4$, that is, $\theta_m < \pi/6$, which is impossible in view of Lemma 4.11.

To prove (iii), assume for a contradiction that $\varphi(t_1) \in (\pi/2 - \varepsilon, \pi/2]$. We have

$$\frac{\pi}{2} - \varepsilon \leq \int_0^{\varphi(t_1)} d\varphi \stackrel{(4-25)}{=} \int_{\theta_0}^{\theta(\varphi(t_1))} \frac{K}{\sin \theta \sqrt{\cos^2 \theta \sin^2 \theta - K^2}} d\theta \leq \int_{\theta_0}^{\theta_M} \frac{K}{\sin \theta \sqrt{\cos^2 \theta \sin^2 \theta - K^2}} d\theta,$$

where in the last inequality we have used (ii). Splitting the right-hand side and applying (i), we then obtain

$$\begin{aligned} \frac{\pi}{2} - \varepsilon &\leq \int_{\pi/6}^{\pi/4} \frac{K}{\sin \theta \sqrt{\cos^2 \theta \sin^2 \theta - K^2}} d\theta + \frac{\sqrt{2}}{2} \int_{\pi/4}^{\theta_M} \frac{K}{\sin \theta \cos \theta \sqrt{\cos^2 \theta \sin^2 \theta - K^2}} d\theta \\ &\stackrel{(4-30)}{=} \int_{\pi/6}^{\pi/4} \frac{K}{\sin \theta \sqrt{\cos^2 \theta \sin^2 \theta - K^2}} d\theta + \frac{\pi\sqrt{2}}{4}. \end{aligned} \tag{4-34}$$

Furthermore, again by (ii), we have

$$K = \sin \theta_M \cos \theta_M \leq \sin \theta(\varphi(t_1)) \cos \theta(\varphi(t_1)) \leq \sin \theta^*(\varphi - \varepsilon) \cos \theta^*(\varphi - \varepsilon) \rightarrow 0 \quad \text{as } \varepsilon \rightarrow 0.$$

Therefore the integral on the right-hand side of (4-34) vanishes as $\varepsilon \rightarrow 0$, yielding a contradiction for ε sufficiently small. □

Lemma 4.14. *Let $\varphi \in I = (\varphi_0, \varphi_1) \subseteq [0, \pi/2 - \varepsilon]$. Then*

$$J_I(\theta_\varepsilon^*) = \left[\frac{\sin \theta_0 \sin \varphi}{\sqrt{1 + \tan^2 \theta_0 \cos^2 \varphi}} \right]_{\varphi=\varphi_0}^{\varphi=\varphi_1}. \tag{4-35}$$

Proof. Since $\theta_\varepsilon^* = \theta^*$ for $\varphi \leq \pi/2 - \varepsilon$, a straightforward computation shows that

$$\cos \theta_\varepsilon^* \sqrt{(\theta_\varepsilon^{*'})^2 + \sin^2 \theta_\varepsilon^*} = \frac{4 \sin \theta_0 \cos \varphi}{\sqrt{1 + \tan^2 \theta_0 \cos^2 \varphi} (3 + \cos 2\theta_0 + 2 \cos 2\varphi \sin^2 \theta_0)}.$$

An integration of this expression yields (4-35). □

We now show that if the graph of a solution to (4-24) emanates from $\partial\mathcal{P} \cap \partial\mathcal{P}_\varepsilon$ into $\mathring{\mathcal{P}}_\varepsilon$, then it does not return to $\partial\mathcal{P} \cap \partial\mathcal{P}_\varepsilon$.

Lemma 4.15. *Let $\varphi_1 \in [0, \pi/2 - \varepsilon]$ and let θ be a solution of (4-24) such that $\theta(\varphi_1) = \theta^*(\varphi_1)$ and $\theta'(\varphi_1) \leq \theta^{*'}(\varphi_1)$. Then $X(\varphi, \theta(\varphi)) \subset \mathring{\mathcal{P}}_\varepsilon$ for all $\varphi \in (\varphi_1, \pi/2 - \varepsilon]$.*

Proof. We let $\theta_1 = \theta(\varphi_1)$ and we distinguish two cases.

Case 1: $\theta'(\varphi_1) \leq 0$. If $\theta_0 = \pi/4$, $\varphi_1 = 0$, and $\theta'(\varphi_1) = 0$, then $\theta \equiv \pi/4$ and the lemma is trivially true. Else Lemma 4.10 implies that θ decreases until either $\varphi = \pi$ or it reaches its minimum. In the former case the lemma is proved. In the latter, part (d) of Lemma 4.10 implies that $\theta < \theta^*(\varphi_1)$ at least until $\varphi = \varphi_1 + \pi/\sqrt{2} > \pi/2$.

Case 2: $\theta'(\varphi_1) > 0$. It is convenient to set

$$v(\varphi) = \log \tan\left(\frac{1}{2}\theta(\varphi)\right).$$

Lengthy but straightforward computations show that

$$v'' = \frac{\cosh(2v) - 3}{\sinh(2v)}(1 + (v')^2).$$

We now observe that

$$\begin{aligned} \cosh(2v) < 3 &\iff \frac{1}{2} \log(3 - 2\sqrt{2}) < v < \frac{1}{2} \log(3 + 2\sqrt{2}) \\ &\iff \log \tan(\pi/8) < \log \tan(\theta/2) < \log \tan(3\pi/8) \\ &\iff \theta \in (\pi/4, \pi/2), \\ \sinh(2v) > 0 &\iff v > 0 \iff \theta \in (\pi/8, \pi/2). \end{aligned}$$

Hence $v'' < 0$ if $\theta > \pi/4$. On the other hand, as long as $\varphi \leq \pi/2 - \varepsilon$, we have

$$\theta < \theta_\varepsilon^* = \theta^* \iff v < v^*(\varphi) := \log \tan\left(\frac{1}{2} \arctan \frac{1}{\tan \theta_0 |\cos \varphi|}\right)$$

with

$$v^{*''} = \frac{\sin \theta_0 \cos \varphi}{(\sin^2 \theta_0 \cos^2 \varphi + \cos^2 \theta_0)^{3/2}} > 0.$$

Hence $(v - v^*)'' < 0$ as long as $\theta > \pi/4$ and $\varphi \leq \pi/2 - \varepsilon$. Since $v = v^*$ and $v' \leq v^{*'} at $\varphi = \varphi_1$, we have $v < v^*$ as long as either $\varphi = \pi/2 - \varepsilon$ or $\theta = \pi/4$. In the former case the proof is complete. In the latter case, part (d) of Lemma 4.10 implies that θ will then remain below $\pi/4$ at least in an interval of length $\pi/\sqrt{2} > \pi/2$, and the proof is complete. $\square$$

We now estimate J over a candidate symmetric shortest path which de-touches from $\partial\mathcal{P} \cap \partial\mathcal{P}_\varepsilon$ and reaches $\varphi = \pi/2 - \varepsilon$:

Lemma 4.16. *Let γ be a symmetric shortest path not passing through $(0, 0, 1)$ and let $\gamma = X(\varphi, \theta)$. If $t_1 \geq 0$ exists such that $\varphi_1 = \varphi(t_1) \in [0, \pi/2 - \varepsilon)$, $\theta(t_1) = \theta^*(\varphi_1)$, and $\gamma \not\subset \partial\mathcal{P}$ in a right-neighborhood of t_1 , then*

$$J_{(\varphi_1, \pi/2 - \varepsilon)}(\theta) > J_{(\varphi_1, \pi/2)}(\theta^*) - \frac{\varepsilon}{2}.$$

Proof. By assumption, for all $\sigma > 0$, there exists $t_\sigma \in (t_1, t_1 + \sigma)$ such that $\boldsymbol{\gamma}(t_\sigma) \in \mathring{\mathcal{P}}_\varepsilon$. By continuity, there exists $\tilde{t}_\sigma \in [t_1, t_\sigma)$ such that $\boldsymbol{\gamma}(\tilde{t}_\sigma) \in \partial\mathcal{P}_\varepsilon$ and $\boldsymbol{\gamma}(t) \in \mathring{\mathcal{P}}_\varepsilon$ for all $t \in (\tilde{t}_\sigma, t_\sigma]$. Then we may apply Lemma 4.9 in $(\tilde{t}_\sigma, t_\sigma]$.

If $K = 0$, then φ is constant, and since the curve is on $\partial\mathcal{P}_\varepsilon$ at $t = \tilde{t}_\sigma$, θ must decrease. Hence $\boldsymbol{\gamma}$ remains smooth down to $\theta = 0$, the north pole. Therefore this case is excluded.

Then $K > 0$, φ is strictly increasing, and $\theta(\varphi)$ solves (4-24) in $(\tilde{t}_\sigma, t_\sigma)$. By Lemma 4.15, we in fact have $X(\varphi, \theta(\varphi)) \subset \mathring{\mathcal{P}}_\varepsilon$ as long as $\varphi \leq \pi/2 - \varepsilon$, which in particular implies that $\tilde{t}_\sigma = t_1$ and that θ solves (4-24) as long as $\varphi \leq \pi/2 - \varepsilon$. We let

$$\theta_1 = \theta(\varphi_1) = \theta^*(\varphi_1)$$

and we distinguish two cases.

Case 1: $\theta'(\varphi_1) \leq 0$. Lemma 4.10 and the symmetry of the path imply that θ does not increase until $\pi/2$ and $\theta(\pi/2) = \theta_m > 0$. If $\theta_1 = \theta_0 = \pi/4$ and $\varphi_1 = 0$, then $\boldsymbol{\gamma}((0, 1)) \subset \mathring{\mathcal{P}}_\varepsilon$, a case which has already been ruled out in Lemma 4.12. Hence $\theta_1 > \pi/4$.

We claim that

$$\min_{\varphi \in [0, \pi/2]} \sin \theta \cos \theta = \sin \theta_m \cos \theta_m. \tag{4-36}$$

By (4-25),

$$\min_{\varphi \in [\varphi_1, \pi/2]} \sin \theta \cos \theta = \sin \theta_m \cos \theta_m. \tag{4-37}$$

In particular,

$$\sin(\theta_1) \cos(\theta_1) \geq \sin \theta_m \cos \theta_m. \tag{4-38}$$

On the other hand, by Lemma 4.13(ii), $\theta \in [\theta_0, \theta_1]$ for $\varphi \in [0, \varphi_1]$. Hence

$$\min_{\varphi \in [0, \varphi_1]} \sin \theta \cos \theta = \min\{\sin \theta_0 \cos \theta_0, \sin \theta_1 \cos \theta_1\}. \tag{4-39}$$

Since θ^* is increasing,

$$\sin(\theta_1) \cos(\theta_1) \leq \sin(\theta^*(0)) \cos(\theta^*(0)) = \sin\left(\frac{\pi}{2} - \theta_0\right) \cos\left(\frac{\pi}{2} - \theta_0\right) = \sin(\theta_0) \cos(\theta_0),$$

therefore (4-39) reads

$$\min_{\varphi \in [0, \varphi_1]} \sin \theta \cos \theta = \sin \theta_1 \cos \theta_1 \stackrel{(4-38)}{\geq} \sin \theta_m \cos \theta_m \tag{4-40}$$

and (4-36) follows from (4-37) and (4-40).

We denote by $\tilde{\varphi}_1 \in (\varphi_1, \pi/2)$ the unique point such that $\theta(\tilde{\varphi}) = \theta_0$ (recall that $\theta_1 > \pi/4$, $\theta(\pi/2) = \theta_m < \pi/4$ and θ is decreasing in $(\varphi_1, \pi/2)$), and we define (see Figure 2)

$$\tilde{\theta}(\varphi) := \begin{cases} \theta(\varphi + \tilde{\varphi}_1) & \text{if } 0 \leq \varphi \leq \frac{\pi}{2} - \tilde{\varphi}_1, \\ \theta_m & \text{if } \frac{\pi}{2} - \tilde{\varphi}_1 \leq \varphi \leq \frac{\pi}{2}. \end{cases}$$

We have

$$J_{(0, \tilde{\varphi}_1)}(\theta) > \sin \theta_m \cos \theta_m \tilde{\varphi}_1 = J_{(\pi/2 - \tilde{\varphi}_1, \pi/2)}(\tilde{\theta}) \quad \text{and} \quad J_{(0, \pi/2 - \tilde{\varphi}_1)}(\tilde{\theta}) = J_{(\tilde{\varphi}_1, \pi/2)}(\theta).$$




Figure 2. Case 1 in the proof of Lemma 4.16. The path $(\varphi, \theta(\varphi))$ (continuous) is beaten by its competitor $(\varphi, \tilde{\theta}(\varphi))$ (dashed).

Therefore γ is not a shortest path and this case is excluded.

Case 2: $\theta'(\varphi_1) > 0$. Lemma 4.10 and the symmetry of the path imply that θ increases until $\pi/2 - \varepsilon$. We now estimate its length in $I_\varepsilon = (\varphi_1, \pi/2 - \varepsilon)$. By the assumption of Case 2, and since $\gamma \in \mathcal{P}$ in I_ε ,

$$0 < \theta'(\varphi_1) \leq (\theta^*)'(\varphi_1). \tag{4-41}$$

By (4-25),

$$\sin^2 \theta (\cos^2 \theta \sin^2 \theta - K^2) = K^2 (\theta')^2 \quad \text{for some } K \in (0, 1/2). \tag{4-42}$$

Evaluating this expression at φ_1 , we have

$$K = \frac{\sin^2 \theta^*(\varphi_1) \cos \theta^*(\varphi_1)}{\sqrt{(\theta'(\varphi_1))^2 + \sin^2 \theta^*(\varphi_1)}} \stackrel{(4-41)}{\geq} \frac{\sin^2 \theta^*(\varphi_1) \cos \theta^*(\varphi_1)}{\sqrt{(\theta^*)'(\varphi_1)^2 + \sin^2 \theta^*(\varphi_1)}}.$$

Of course, we have

$$J_{I_\varepsilon}(\theta) \geq \int_{I_\varepsilon} \cos \theta \sin \theta \, d\varphi \stackrel{(4-42)}{\geq} |I_\varepsilon| K.$$

This chain of inequalities implies that

$$J_{I_\varepsilon}(\theta) > |I_\varepsilon| \frac{\sin^2 \theta^*(\varphi_1) \cos \theta^*(\varphi_1)}{\sqrt{(\theta^*)'(\varphi_1)^2 + \sin^2 \theta^*(\varphi_1)}} = \frac{|I_\varepsilon| \sin \theta_0 \cos \varphi_1}{\sqrt{1 + \tan^2 \theta_0 \cos^2 \varphi_1}}$$

(the latter equality follows from an explicit computation). On the other hand, by Lemma 4.14, the curve which just stays on the obstacle, $\gamma^* = X(\varphi, \theta^*(\varphi))$, $\varphi \in (\varphi_1, \pi/2)$, is such that

$$J_{(\varphi_1, \pi/2)}(\theta^*) = \sin \theta_0 \left(1 - \frac{\sin \varphi_1}{\sqrt{1 + \tan^2 \theta_0 \cos^2 \varphi_1}} \right).$$

Hence

$$\begin{aligned} (J_{I_\varepsilon}(\theta) - J_{(\varphi_1, \pi/2)}(\theta^*)) & \frac{\sqrt{1 + \tan^2 \theta_0 \cos^2 \varphi_1}}{\sin \theta_0} \\ & > |I_\varepsilon| \cos \varphi_1 + \sin \varphi_1 - \sqrt{1 + \tan^2 \theta_0 \cos^2 \varphi_1} \\ & > \left(\frac{\pi}{2} - \varepsilon - \varphi_1\right) \cos \varphi_1 + \sin \varphi_1 - \sqrt{1 + \cos^2 \varphi_1} =: F(\varphi_1) - \varepsilon \cos \varphi_1. \end{aligned}$$

Another calculus exercise shows that F is decreasing $[0, \pi/2]$: since $F(\pi/2) = 0$, F is positive. Therefore

$$J_{I_\varepsilon}(\theta) > J_{(\varphi_1, \pi/2)}(\theta^*) - \varepsilon \frac{\sin \theta_0 \cos \varphi_1}{\sqrt{1 + \tan^2 \theta_0 \cos^2 \varphi_1}} > J_{(\varphi_1, \pi/2)}(\theta^*) - \frac{1}{2}\varepsilon. \quad \square$$

Next we characterize the candidate shortest paths joining $X(0, \theta^*(0))$ with another point on $\partial\mathcal{P} \cap \partial\mathcal{P}_\varepsilon$ which is on the same side with respect to $\pi/2$.

Lemma 4.17. *Let $0 < \bar{\varphi} \leq \pi/2 - \varepsilon$. The shortest path which connects $X(0, \pi/2 - \theta_0)$ and $X(\bar{\varphi}, \theta^*(\bar{\varphi}))$ is (a smooth reparametrization of) $\boldsymbol{\gamma}^* = X(\varphi, \theta^*(\varphi))$, $\varphi \in [0, \bar{\varphi}]$.*

Proof. Let $I = (0, \bar{\varphi})$. We recall by Lemma 4.14 that

$$J_I(\theta^*) = \frac{\sin \theta_0 \sin \bar{\varphi}}{\sqrt{1 + \tan^2 \theta_0 \cos^2 \bar{\varphi}}} < \sin \theta_0.$$

First note that $\boldsymbol{\gamma}$ does not reach the north pole. For if it did, at a time $\bar{t} \in (0, 1)$, we would have

$$J(\boldsymbol{\gamma}) \geq \int_0^{\bar{t}} \cos \theta |\dot{\theta}| dt \geq \sin(\theta(0)) = \sin\left(\frac{\pi}{2} - \theta_0\right) = \cos \theta_0 > \sin \theta_0 > J_I(\theta^*),$$

which is impossible.

Therefore we may use the spherical coordinates (4-4), and arguing as in the proof of Lemma 4.8 we see that φ is nondecreasing.

Assume for a contradiction that $\boldsymbol{\gamma}$ does not coincide (up to a smooth reparametrization) with $\boldsymbol{\gamma}^*$. Then $t_1 > 0$ and a right-neighborhood \tilde{I} of t_1 exist such that $\varphi_1 := \varphi(t_1) < \bar{\varphi}$ and $\boldsymbol{\gamma}(\tilde{I}) \not\subset \partial\mathcal{P}$. Arguing as in the first lines of the proof of Lemma 4.16, one finds that there is $t_2 > t_1$ such that $\boldsymbol{\gamma}(t) \in \overset{\circ}{\mathcal{P}}$ for all $t \in (t_1, t_2)$. Then, arguing as in the proof of Lemma 4.9, one finds that (4-22) holds, and $K \geq 0$ since φ is nondecreasing. If $K > 0$, then $\theta(\varphi)$ would solve (4-24) in (t_1, t_2) ; but in view of Lemma 4.15, such a solution will not rehit the constraint until $\varphi = \pi/2 - \varepsilon$, hence $K > 0$ can not occur. If $K = 0$, then $\varphi \equiv \varphi_1$, and since we are on $\partial\mathcal{P}$ at time t_1 , θ must move inwards. Hence $\boldsymbol{\gamma}$ remains smooth up to $\theta = 0$, the north pole, a contradiction. \square

Proof of Theorem 4.1. First of all, we note that Lemma 4.14 implies that $J(\theta^*) = 2 \sin \theta_0$. Hence, in view of Lemma 4.5, it suffices to show that

$$\inf_{\boldsymbol{\gamma} \in \Gamma_\varepsilon} J(\boldsymbol{\gamma}) \geq J(\theta^*) - \omega(\varepsilon) = 2 \sin \theta_0 - \omega(\varepsilon), \tag{4-43}$$

where ω is a universal function which vanishes as $\varepsilon \rightarrow 0$. By Lemma 4.6, the inf on the left-hand side of (4-43) is attained. Let $\boldsymbol{\gamma}$ be one such shortest path. If $\boldsymbol{\gamma}$ passes through $(0, 0, 1)$, then (4-43) follows from




Figure 3. Case 1 in the proof of Theorem 4.1. The path $(\varphi(t), \theta(t))$ (gray) is beaten by its competitor $(\varphi(t), \tilde{\theta}(t))$ (dashed).

Lemma 4.7. If not, we let $\gamma = X(\varphi, \theta)$ and, by Lemma 4.8, we assume without loss of generality that γ is symmetric. For simplicity, we distinguish between $\theta_0 < \pi/4$ and $\theta_0 = \pi/4$.

Case 1: $\theta_0 < \pi/4$. We already know from Lemma 4.12 that γ has to intersect $\partial\mathcal{P}_\varepsilon$. Let t_0 and t_1 be, respectively, the first time in which $\theta(t) = \pi/4$ and the first time in which γ intersects $\partial\mathcal{P}_\varepsilon$:

$$t_1 := \sup\{t > 0 : \gamma \in \mathring{\mathcal{P}}_\varepsilon \text{ in } [0, t)\} \quad \text{and} \quad \varphi_1 = \varphi(t_1).$$

Provided ε is sufficiently small, by Lemma 4.13(ii), θ is increasing in $(0, t_1)$. Hence the curve

$$\tilde{\gamma}(t) := X(\varphi(t), \tilde{\theta}(t)), \quad \tilde{\theta}(t) := \begin{cases} \frac{\pi}{2} - \theta(t) & t \in [0, t_0], \\ \theta(t) & t \in [t_0, t_1] \end{cases}$$

is contained in \mathcal{P}_ε (see Figure 3). We claim that

$$J(\tilde{\gamma} \chi_{(0,t_1)}) < J(\gamma \chi_{(0,t_1)}), \tag{4-44}$$

which is equivalent to

$$J(\tilde{\gamma} \chi_{(0,t_0)}) < J(\gamma \chi_{(0,t_0)}). \tag{4-45}$$

By Lemma 4.9, γ satisfies (4-22) in $(0, t_0)$. If $K = 0$, then $\varphi \equiv 0$ and (4-45) follows from the expression (4-5) of J :

$$\cos\left(\frac{\pi}{2} - \theta\right) = \sin(\theta) < \cos \theta \quad \text{if } \theta \leq \pi/4.$$

Otherwise, by Lemma 4.9 $\varphi \mapsto \theta(\varphi)$ solves (4-24) in $(0, \varphi_0)$, where $\varphi_0 = \varphi(t_0)$. Then it follows by Lemma 4.13(iii) that $\varphi_0 < \pi/2 - \varepsilon$, and we may use the equivalent expression (4-26) for J : since

$$\cos^2 \tilde{\theta}((\tilde{\theta}')^2 + \sin^2 \tilde{\theta}) = \sin^2 \theta((\theta')^2 + \cos^2 \theta) < \cos^2 \theta((\theta')^2 + \sin^2 \theta) \quad \text{in } (0, \varphi_0),$$

(4-45) follows.

Since $\tilde{\gamma}$ is a path connecting $X(0, \pi/2 - \theta_0)$ to $X(\varphi_1, \theta^*(\varphi_1))$, Lemma 4.17 implies that $J(\tilde{\gamma} \chi_{(0,t_1)}) \geq J_{(0,\varphi_1)}(\theta^*)$. Together with (4-44), we obtain

$$J_{(0,\varphi_1)}(\theta^*) < J(\gamma \chi_{(0,t_1)}). \tag{4-46}$$

Now let $t_2 \geq t_1$ be defined by

$$t_2 := \max\{t \geq t_1 : \boldsymbol{\gamma} \in \partial\mathcal{P}_\varepsilon \text{ in } [t_1, t]\} \quad \text{and} \quad \varphi_2 = \varphi(t_2).$$

The estimate in (t_1, t_2) is trivial since $\boldsymbol{\gamma}$ coincides with $\boldsymbol{\gamma}^* := X(\varphi, \theta^*)$:

$$J(\boldsymbol{\gamma}\chi_{(t_1, t_2)}) = J_{(\varphi_1, \varphi_2)}(\theta^*).$$

On $(\varphi_2, \pi/2 - \varepsilon)$, Lemma 4.16 implies that

$$J_{(\varphi_2, \pi/2 - \varepsilon)}(\theta) > J_{(\varphi_2, \pi/2)}(\theta^*) - \frac{\varepsilon}{2} \quad \text{if } \varphi_2 < \frac{\pi}{2} - \varepsilon. \quad (4-47)$$

Finally, we just observe that

$$J_{(\pi/2 - \varepsilon, \pi/2)}(\theta^*) \stackrel{(4-7)}{\leq} \omega(\varepsilon). \quad (4-48)$$

Collecting (4-46)–(4-48) and recalling the symmetry of $\boldsymbol{\gamma}$, we obtain (4-43).

Case 2: $\theta_0 = \pi/4$. This case is simpler. We let

$$t_2 = \max\{t \geq 0 : \boldsymbol{\gamma} \in \partial\mathcal{P}_\varepsilon \text{ in } [0, t]\} \geq 0 \quad \text{and} \quad \varphi_2 = \varphi(t_2),$$

and we argue exactly as above to obtain $J(\boldsymbol{\gamma}\chi_{(0, t_2)}) = J_{(0, \varphi_2)}(\theta^*)$ and (4-47)–(4-48). \square

Acknowledgement

We thank Manuel Ritoré for fruitful discussions on the geometrical problem in Section 4.

References

- [Alexander et al. 1993] S. B. Alexander, I. D. Berg, and R. L. Bishop, “Cut loci, minimizers, and wavefronts in Riemannian manifolds with boundary”, *Michigan Math. J.* **40**:2 (1993), 229–237. MR 94c:53053 Zbl 0817.53023
- [Alicandro et al. 2007] R. Alicandro, A. Corbo Esposito, and C. Leone, “Relaxation in BV of integral functionals defined on Sobolev functions with values in the unit sphere”, *J. Convex Anal.* **14**:1 (2007), 69–98. MR 2008b:49007 Zbl 1138.49017
- [Ambrosio and Dal Maso 1990] L. Ambrosio and G. Dal Maso, “A general chain rule for distributional derivatives”, *Proc. Amer. Math. Soc.* **108**:3 (1990), 691–702. MR 90j:26019 Zbl 0685.49027
- [Ambrosio et al. 2000] L. Ambrosio, N. Fusco, and D. Pallara, *Functions of bounded variation and free discontinuity problems*, Clarendon Press, New York, 2000. MR 2003a:49002 Zbl 0957.49001
- [Ambrosio et al. 2005] L. Ambrosio, G. Crippa, and S. Maniglia, “Traces and fine properties of a BD class of vector fields and applications”, *Ann. Fac. Sci. Toulouse Math.* (6) **14**:4 (2005), 527–561. MR 2007b:35040 Zbl 1091.35007
- [Andreu et al. 2001] F. Andreu, C. Ballester, V. Caselles, and J. M. Mazón, “Minimizing total variation flow”, *Differential Integral Equations* **14**:3 (2001), 321–360. MR 2002e:35109 Zbl 1020.35037
- [Andreu-Vaillio et al. 2004] F. Andreu-Vaillio, V. Caselles, and J. M. Mazón, *Parabolic quasilinear equations minimizing linear growth functionals*, Progress in Mathematics **223**, Birkhäuser, Basel, 2004. MR 2005c:35002 Zbl 1053.35002
- [Anzellotti 1983] G. Anzellotti, “Pairings between measures and bounded functions and compensated compactness”, *Ann. Mat. Pura Appl.* (4) **135** (1983), 293–318. MR 85m:46042 Zbl 0572.46023
- [Barrett et al. 2008] J. W. Barrett, X. Feng, and A. Prohl, “On p -harmonic map heat flows for $1 \leq p < \infty$ and their finite element approximations”, *SIAM J. Math. Anal.* **40**:4 (2008), 1471–1498. MR 2009m:35202 Zbl 1182.35154
- [Bertsch et al. 2002] M. Bertsch, R. Dal Passo, and R. van der Hout, “Nonuniqueness for the heat flow of harmonic maps on the disk”, *Arch. Ration. Mech. Anal.* **161**:2 (2002), 93–112. MR 2003a:35093 Zbl 1006.35050

- [Bertsch et al. 2003] M. Bertsch, R. Dal Passo, and A. Pisante, “Point singularities and nonuniqueness for the heat flow for harmonic maps”, *Comm. Partial Differential Equations* **28**:5-6 (2003), 1135–1160. MR 2004c:53099 Zbl 1029.58008
- [Bonforte and Figalli 2012] M. Bonforte and A. Figalli, “Total variation flow and sign fast diffusion in one dimension”, *J. Differential Equations* **252**:8 (2012), 4455–4480. MR 2881044 Zbl 1242.35049
- [Brezis 2011] H. Brezis, *Functional analysis, Sobolev spaces and partial differential equations*, Springer, New York, 2011. MR 2012a:35002 Zbl 1220.46002
- [Caselles 2011] V. Caselles, “On the entropy conditions for some flux limited diffusion equations”, *J. Differential Equations* **250**:8 (2011), 3311–3348. MR 2012d:35179 Zbl 1231.35101
- [Chen 1989] Y. M. Chen, “The weak solutions to the evolution problems of harmonic maps”, *Math. Z.* **201**:1 (1989), 69–74. MR 90i:58030 Zbl 0685.58015
- [Chen and Frid 1999] G.-Q. Chen and H. Frid, “Divergence-measure fields and hyperbolic conservation laws”, *Arch. Ration. Mech. Anal.* **147**:2 (1999), 89–118. MR 2000d:35136 Zbl 0942.35111
- [Chen et al. 1994] Y. M. Chen, M. C. Hong, and N. Hungerbühler, “Heat flow of p -harmonic maps with values into spheres”, *Math. Z.* **215**:1 (1994), 25–35. MR 94k:58145 Zbl 0793.53049
- [Dal Passo et al. 2008] R. Dal Passo, L. Giacomelli, and S. Moll, “Rotationally symmetric 1-harmonic maps from D^2 to S^2 ”, *Calc. Var. Partial Differential Equations* **32**:4 (2008), 533–554. MR 2009f:58024 Zbl 1147.58015
- [Darling 1994] R. W. R. Darling, *Differential forms and connections*, Cambridge University Press, 1994. MR 95j:53038 Zbl 0822.53001
- [DeSimone and Podio-Guidugli 1996] A. DeSimone and P. Podio-Guidugli, “On the continuum theory of deformable ferromagnetic solids”, *Arch. Rational Mech. Anal.* **136**:3 (1996), 201–233. MR 98a:73051 Zbl 1002.74521
- [Diestel and Uhl 1977] J. Diestel and J. J. Uhl, Jr., *Vector measures*, Mathematical Surveys **15**, Amer. Math. Soc., Providence, RI, 1977. MR 56 #12216 Zbl 0369.46039
- [Eells and Sampson 1964] J. Eells, Jr. and J. H. Sampson, “Harmonic mappings of Riemannian manifolds”, *Amer. J. Math.* **86** (1964), 109–160. MR 29 #1603 Zbl 0122.40102
- [Evans and Gariepy 1992] L. C. Evans and R. F. Gariepy, *Measure theory and fine properties of functions*, CRC Press, Boca Raton, FL, 1992. MR 93f:28001 Zbl 0804.28001
- [Federer 1969] H. Federer, *Geometric measure theory*, Die Grundlehren der mathematischen Wissenschaften **153**, Springer, New York, 1969. MR 41 #1976 Zbl 0176.00801
- [Feng 2010] X. Feng, “Divergence- L^q and divergence-measure tensor fields and gradient flows for linear growth functionals of maps into the unit sphere”, *Calc. Var. Partial Differential Equations* **37**:1-2 (2010), 111–139. MR 2011a:35281 Zbl 1184.35171
- [Fonseca and Müller 1993] I. Fonseca and S. Müller, “Relaxation of quasiconvex functionals in $BV(\Omega, \mathbb{R}^P)$ for integrands $f(x, u, \nabla u)$ ”, *Arch. Rational Mech. Anal.* **123**:1 (1993), 1–49. MR 94h:49023 Zbl 0788.49039
- [Fonseca and Rybka 1992] I. Fonseca and P. Rybka, “Relaxation of multiple integrals in the space $BV(\Omega, \mathbb{R}^P)$ ”, *Proc. Roy. Soc. Edinburgh Sect. A* **121**:3-4 (1992), 321–348. MR 94g:49032 Zbl 0794.49012
- [Giacomelli and Moll 2010] L. Giacomelli and S. Moll, “Rotationally symmetric 1-harmonic flows from D^2 to S^2 : local well-posedness and finite time blowup”, *SIAM J. Math. Anal.* **42**:6 (2010), 2791–2817. MR 2011k:35084 Zbl 1230.35046
- [Giacomelli et al. 2013a] L. Giacomelli, J. M. Mazón, and S. Moll, “The 1-harmonic flow with values into \mathbb{S}^1 ”, *SIAM J. Math. Anal.* **45**:3 (2013), 1723–1740. MR 3063148 Zbl 06200948
- [Giacomelli et al. 2013b] L. Giacomelli, J. M. Mazón, and S. Moll, “Solutions to the 1-harmonic flow with values into a hyper-octant of the N -sphere”, *Appl. Math. Lett.* **26**:11 (2013), 1061–1064. MR 3089565
- [Giaquinta and Mucci 2006] M. Giaquinta and D. Mucci, “The BV-energy of maps into a manifold: Relaxation and density results”, *Ann. Sc. Norm. Super. Pisa Cl. Sci. (5)* **5**:4 (2006), 483–548. MR 2008k:49090 Zbl 1150.49020
- [Giga and Kobayashi 2003] Y. Giga and R. Kobayashi, “On constrained equations with singular diffusivity”, *Methods Appl. Anal.* **10**:2 (2003), 253–277. MR 2005e:58030 Zbl 1058.58006
- [Giga and Kuroda 2004] Y. Giga and H. Kuroda, “On breakdown of solutions of a constrained gradient system of total variation”, *Bol. Soc. Parana. Mat. (3)* **22**:1 (2004), 9–20. MR 2005f:35149 Zbl 1064.35084

- [Giga et al. 2004] Y. Giga, Y. Kashima, and N. Yamazaki, “Local solvability of a constrained gradient system of total variation”, *Abstr. Appl. Anal.* **8** (2004), 651–682. MR 2005k:35195 Zbl 1068.35054
- [Giga et al. 2005] Y. Giga, H. Kuroda, and N. Yamazaki, “An existence result for a discretized constrained gradient system of total variation flow in color image processing”, *Interdiscip. Inform. Sci.* **11**:2 (2005), 199–204. MR 2006h:65161
- [Giga et al. 2007] Y. Giga, H. Kuroda, and N. Yamazaki, “Global solvability of constrained singular diffusion equation associated with essential variation”, pp. 209–218 in *Free boundary problems*, edited by I. N. Figueiredo et al., Internat. Ser. Numer. Math. **154**, Birkhäuser, Basel, 2007. MR 2007m:35141 Zbl 1119.35028
- [van der Hout 2001] R. van der Hout, “Flow alignment in nematic liquid crystals in flows with cylindrical symmetry”, *Differential Integral Equations* **14**:2 (2001), 189–211. MR 2001m:76007 Zbl 1021.35028
- [Hungerbühler 2004] N. Hungerbühler, “Heat flow into spheres for a class of energies”, pp. 45–65 in *Variational problems in Riemannian geometry*, edited by P. Baird et al., Progr. Nonlinear Differential Equations Appl. **59**, Birkhäuser, Basel, 2004. MR 2005d:53106 Zbl 1063.58010
- [Kobayashi et al. 2000] R. Kobayashi, J. A. Warren, and W. C. Carter, “A continuum model of grain boundaries”, *Phys. D* **140**:1-2 (2000), 141–150. MR 2000m:74071 Zbl 0956.35123
- [Misawa 2002] M. Misawa, “On the p -harmonic flow into spheres in the singular case”, *Nonlinear Anal.* **50**:4, Ser. A: Theory Methods (2002), 485–494. MR 2003f:53119 Zbl 1163.53339
- [Sapiro 2001] G. Sapiro, *Geometric partial differential equations and image analysis*, Cambridge University Press, 2001. MR 2002a:68142 Zbl 0968.35001
- [Simon 1987] J. Simon, “Compact sets in the space $L^p(0, T; B)$ ”, *Ann. Mat. Pura Appl.* (4) **146** (1987), 65–96. MR 89c:46055 Zbl 0629.46031
- [Struwe 1992] M. Struwe, “The evolution of harmonic maps: Existence, partial regularity, and singularities”, pp. 485–491 in *Nonlinear diffusion equations and their equilibrium states, 3* (Gregynog, 1989), edited by N. G. Lloyd et al., Progr. Nonlinear Differential Equations Appl. **7**, Birkhäuser, Boston, MA, 1992. MR 1167858 Zbl 0769.58016
- [Tang et al. 2000] B. Tang, G. Sapiro, and V. Caselles, “Diffusion of general data on non-flat manifolds via harmonic maps theory: The direction diffusion case”, *Int. J. Comput. Vis.* **36**:2 (2000), 149–161.
- [Tang et al. 2001] B. Tang, G. Sapiro, and V. Caselles, “Color image enhancement via chromaticity diffusion”, *IEEE Trans. Image Process.* **10**:5 (2001), 701–707. Zbl 1037.68792
- [Ziemer 1989] W. P. Ziemer, *Weakly differentiable functions: Sobolev spaces and functions of bounded variation*, Graduate Texts in Mathematics **120**, Springer, New York, 1989. MR 91e:46046 Zbl 0692.46022

Received 18 Apr 2013. Accepted 27 Nov 2013.

LORENZO GIACOMELLI: lorenzo.giacomelli@sbai.uniroma1.it
 SBAI Department, Sapienza University of Rome, Via Scarpa, 16, I-00161 Roma, Italy

JOSE M. MAZÓN: mazon@uv.es
 Departament d’Anàlisi Matemàtica, Universitat de València, Dr. Moliner, 50, 46100 Burjassot, Spain

SALVADOR MOLL: j.salvador.moll@uv.es
 Departament d’Anàlisi Matemàtica, Universitat de València, Dr. Moliner, 50, 46100 Burjassot, Spain

Analysis & PDE

msp.org/apde

EDITORS

EDITOR-IN-CHIEF

Maciej Zworski
zworski@math.berkeley.edu
University of California
Berkeley, USA

BOARD OF EDITORS

Nicolas Burq	Université Paris-Sud 11, France nicolas.burq@math.u-psud.fr	Yuval Peres	University of California, Berkeley, USA peres@stat.berkeley.edu
Sun-Yung Alice Chang	Princeton University, USA chang@math.princeton.edu	Gilles Pisier	Texas A&M University, and Paris 6 pisier@math.tamu.edu
Michael Christ	University of California, Berkeley, USA mchrist@math.berkeley.edu	Tristan Rivière	ETH, Switzerland riviere@math.ethz.ch
Charles Fefferman	Princeton University, USA cf@math.princeton.edu	Igor Rodnianski	Princeton University, USA irod@math.princeton.edu
Ursula Hamenstaedt	Universität Bonn, Germany ursula@math.uni-bonn.de	Wilhelm Schlag	University of Chicago, USA schlag@math.uchicago.edu
Vaughan Jones	U.C. Berkeley & Vanderbilt University vaughan.f.jones@vanderbilt.edu	Sylvia Serfaty	New York University, USA serfaty@cims.nyu.edu
Herbert Koch	Universität Bonn, Germany koch@math.uni-bonn.de	Yum-Tong Siu	Harvard University, USA siu@math.harvard.edu
Izabella Laba	University of British Columbia, Canada ilaba@math.ubc.ca	Terence Tao	University of California, Los Angeles, USA tao@math.ucla.edu
Gilles Lebeau	Université de Nice Sophia Antipolis, France lebeau@unice.fr	Michael E. Taylor	Univ. of North Carolina, Chapel Hill, USA met@math.unc.edu
László Lempert	Purdue University, USA lempert@math.purdue.edu	Gunther Uhlmann	University of Washington, USA gunther@math.washington.edu
Richard B. Melrose	Massachusetts Institute of Technology, USA rbm@math.mit.edu	András Vasy	Stanford University, USA andras@math.stanford.edu
Frank Merle	Université de Cergy-Pontoise, France Frank.Merle@u-cergy.fr	Dan Virgil Voiculescu	University of California, Berkeley, USA dvv@math.berkeley.edu
William Minicozzi II	Johns Hopkins University, USA minicozz@math.jhu.edu	Steven Zelditch	Northwestern University, USA zelditch@math.northwestern.edu
Werner Müller	Universität Bonn, Germany mueller@math.uni-bonn.de		

PRODUCTION

production@msp.org
Silvio Levy, Scientific Editor

See inside back cover or msp.org/apde for submission instructions.

The subscription price for 2014 is US \$180/year for the electronic version, and \$355/year (+\$50, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues from the last three years and changes of subscribers address should be sent to MSP.

Analysis & PDE (ISSN 1948-206X electronic, 2157-5045 printed) at Mathematical Sciences Publishers, 798 Evans Hall #3840, c/o University of California, Berkeley, CA 94720-3840, is published continuously online. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices.

APDE peer review and production are managed by EditFLOW[®] from Mathematical Sciences Publishers.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2014 Mathematical Sciences Publishers

ANALYSIS & PDE

Volume 7 No. 3 2014

Prescription du spectre de Steklov dans une classe conforme PIERRE JAMMES	529
Semilinear geometric optics with boundary amplification JEAN-FRANCOIS COULOMBEL, OLIVIER GUÈS and MARK WILLIAMS	551
The 1-harmonic flow with values in a hyperoctant of the N -sphere LORENZO GIACOMELLI, JOSE M. MAZÓN and SALVADOR MOLL	627
Decomposition rank of \mathcal{L} -stable C^* -algebras AARON TIKUISIS and WILHELM WINTER	673
Scattering for a massless critical nonlinear wave equation in two space dimensions MARTIN SACK	701
Large-time blowup for a perturbation of the cubic Szegő equation HAIYAN XU	717
A geometric tangential approach to sharp regularity for degenerate evolution equations EDUARDO V. TEIXEIRA and JOSÉ MIGUEL URBANO	733
The theory of Hahn-meromorphic functions, a holomorphic Fredholm theorem, and its applications JÖRN MÜLLER and ALEXANDER STROHMAIER	745



2157-5045(2014)7:3;1-D