

ANALYSIS & PDE

Volume 7

No. 4

2014

AURÉLIEN POIRET, DIDIER ROBERT AND LAURENT THOMANN

**PROBABILISTIC GLOBAL WELL-POSEDNESS FOR THE
SUPERCRITICAL NONLINEAR HARMONIC OSCILLATOR**

PROBABILISTIC GLOBAL WELL-POSEDNESS FOR THE SUPERCRITICAL NONLINEAR HARMONIC OSCILLATOR

AURÉLIEN POIRET, DIDIER ROBERT AND LAURENT THOMANN

Thanks to an approach inspired by Burq and Lebeau [*Ann. Sci. Éc. Norm. Supér.* (4) **6:6** (2013)], we prove stochastic versions of Strichartz estimates for Schrödinger with harmonic potential. As a consequence, we show that the nonlinear Schrödinger equation with quadratic potential and any polynomial nonlinearity is almost surely locally well-posed in $L^2(\mathbb{R}^d)$ for any $d \geq 2$. Then, we show that we can combine this result with the high-low frequency decomposition method of Bourgain to prove a.s. global well-posedness results for the cubic equation: when $d = 2$, we prove global well-posedness in $\mathcal{H}^s(\mathbb{R}^2)$ for any $s > 0$, and when $d = 3$ we prove global well-posedness in $\mathcal{H}^s(\mathbb{R}^3)$ for any $s > \frac{1}{6}$, which is a supercritical regime.

Furthermore, we also obtain almost sure global well-posedness results with scattering for NLS on \mathbb{R}^d without potential. We prove scattering results for L^2 -supercritical equations and L^2 -subcritical equations with initial conditions in L^2 without additional decay or regularity assumption.

1. Introduction and results

1A. Introduction. It is known from several works that a probabilistic approach can help to give insight into the dynamics of dispersive nonlinear PDEs, even for low Sobolev regularity. This point of view was initiated by Lebowitz, Rose and Speer [1988], developed by Bourgain [1994; 1996] and Zhidkov [2001], and enhanced by Tzvetkov [2006; 2008; 2010], Burq and Tzvetkov [2008a; 2008b], Oh [2009/10; 2009], Colliander and Oh [2012] and others. In this paper we study the Cauchy problem for the nonlinear Schrödinger–Gross–Pitaevskii equation

$$\begin{cases} i \frac{\partial u}{\partial t} + \Delta u - |x|^2 u = \pm |u|^{p-1} u, & (t, x) \in \mathbb{R} \times \mathbb{R}^d, \\ u(0) = u_0, \end{cases} \quad (1-1)$$

with $d \geq 2$, $p \geq 3$ an odd integer and where u_0 is a random initial condition.

Much work has been done on dispersive PDEs with random initial conditions since the papers of Burq and Tzvetkov [2008a; 2008b]. In these articles, the authors showed that, thanks to a randomisation of the initial condition, one can prove well-posedness results even for data with supercritical Sobolev regularity. We also refer to [Burq and Tzvetkov 2014; Thomann 2009; Burq et al. 2010; Poiret 2012a; 2012b;

Robert was partly supported by the grant “NOSEVOL” ANR-2011-BS01019 01. Thomann was partly supported by the grant “HANDDY” ANR-10-JCJC 0109 and by the grant “ANAÉ” ANR-13-BS01-0010-03.

MSC2010: 35P05, 35Q55, 35R60.

Keywords: harmonic oscillator, supercritical nonlinear Schrödinger equation, random initial conditions, scattering, global solutions.

de Suzzoni 2013; Nahmod and Staffilani 2013] for strong solutions in a probabilistic sense. Concerning weak solutions, see [Burq et al. 2012; ≥ 2014; Nahmod et al. 2013].

More recently, Burq and Lebeau [2013] considered a different randomisation method, and thanks to fine spectral estimates they obtained better stochastic bounds, which enabled them to improve the previous known results for the supercritical wave equation on a compact manifold. In [Poiret et al. 2013] we extended the results of [Burq and Lebeau 2013] to the harmonic oscillator in \mathbb{R}^d . This approach enables us to prove a stochastic version of the usual Strichartz estimates with a gain of $d/2$ derivatives, which we will use here to apply to the nonlinear problem. These estimates (the result of Proposition 2.1) can be seen as a consequence of [Poiret et al. 2013, Inequality (1.6)], but we give here an alternative proof suggested by Nicolas Burq.

Consider a probability space $(\Omega, \mathcal{F}, \mathbb{P})$ and let $\{g_n\}_{n \geq 0}$ be a sequence of real random variables, which we will assume to be independent and identically distributed. We assume that the common law ν of g_n satisfies, for some $c > 0$, the bound

$$\int_{-\infty}^{+\infty} e^{\gamma x} d\nu \leq e^{c\gamma^2} \quad \text{for all } \gamma \in \mathbb{R}. \tag{1-2}$$

This condition implies in particular that the g_n are centred variables. It is easy to check that (1-2) is satisfied for centred Gauss laws and for any centred law with bounded support. Under condition (1-2), we can prove the Khinchin inequality (Lemma 2.3), which we will use in the sequel.

Let $d \geq 2$. We denote by

$$H = -\Delta + |x|^2$$

the harmonic oscillator and by $\{\varphi_j \mid j \geq 1\}$ an orthonormal basis of $L^2(\mathbb{R}^d)$ of eigenvectors of H (the Hermite functions). The eigenvalues of H are the $\{2(\ell_1 + \dots + \ell_d) + d \mid \ell \in \mathbb{N}^d\}$, and we can order them in a non-decreasing sequence $\{\lambda_j \mid j \geq 1\}$, repeated according to their multiplicities, and so that $H\varphi_j = \lambda_j\varphi_j$.

We define the harmonic Sobolev spaces for $s \geq 0, p \geq 1$ by

$$\begin{aligned} \mathring{W}^{s,p} &= \mathring{W}^{s,p}(\mathbb{R}^d) = \{u \in L^p(\mathbb{R}^d) \mid H^{s/2}u \in L^p(\mathbb{R}^d)\}, \\ \mathcal{H}^s &= \mathcal{H}^s(\mathbb{R}^d) = \mathring{W}^{s,2}. \end{aligned}$$

The natural norms are denoted by $\|u\|_{\mathring{W}^{s,p}}$ and up to equivalence of norms, for $1 < p < +\infty$, we have [Yajima and Zhang 2004, Lemma 2.4]

$$\|u\|_{\mathring{W}^{s,p}} = \|H^{s/2}u\|_{L^p} \equiv \|(-\Delta)^{s/2}u\|_{L^p} + \|\langle x \rangle^s u\|_{L^p}.$$

For $j \geq 1$, let

$$I(j) = \{n \in \mathbb{N} \mid 2j \leq \lambda_n < 2(j+1)\}.$$

Observe that, for all $j \geq d/2, I(j) \neq \emptyset$ and that $\#I(j) \sim c_d j^{d-1}$ when $j \rightarrow +\infty$.

Let $s \in \mathbb{R}$. Any $u \in \mathcal{H}^s(\mathbb{R}^d)$ can be written in a unique fashion as

$$u = \sum_{j=1}^{+\infty} \sum_{n \in I(j)} c_n \varphi_n.$$

Following a suggestion of Nicolas Burq, we introduce the condition

$$|c_k|^2 \leq \frac{C}{\#I(j)} \sum_{n \in I(j)} |c_n|^2 \quad \text{for all } j \geq 1 \text{ and } k \in I(j), \tag{1-3}$$

which means that the coefficients have almost the same size on each level of energy $I(j)$. Observe that this condition is always satisfied in dimension $d = 1$. We define the set $\mathcal{A}_s \subset \mathcal{H}^s(\mathbb{R}^d)$ by

$$\mathcal{A}_s = \left\{ u = \sum_{j=1}^{+\infty} \sum_{n \in I(j)} c_n \varphi_n \in \mathcal{H}^s(\mathbb{R}^d) \mid \text{condition (1-3) holds for some } C > 0 \right\}.$$

It is easy to check the following properties:

- If $u \in \mathcal{A}_s$, then for all $c \in \mathbb{C}$, $cu \in \mathcal{A}_s$.
- The set \mathcal{A}_s is neither closed nor open in \mathcal{H}^s .
- The set \mathcal{A}_s is invariant under the linear Schrödinger flow e^{-itH} .
- The set \mathcal{A}_s depends on the choice of the orthonormal basis $(\varphi_n)_{n \geq 1}$. Indeed, given $u \in \mathcal{H}^s$, it is easy to see that there exists a Hilbertian basis $(\tilde{\varphi}_n)_{n \geq 1}$ such that $u \in \tilde{\mathcal{A}}_s$, where $\tilde{\mathcal{A}}_s$ is the space based on $(\tilde{\varphi}_n)_{n \geq 1}$.

Let $\gamma \in \mathcal{A}_s$. We define the probability measure μ_γ on \mathcal{H}^s via the map

$$\begin{aligned} \Omega &\rightarrow \mathcal{H}^s(\mathbb{R}^d), \\ \omega &\mapsto \gamma^\omega = \sum_{j=1}^{+\infty} \sum_{n \in I(j)} c_n g_n(\omega) \varphi_n. \end{aligned}$$

In other words, μ_γ is defined by the condition, that for all measurable $F : \mathcal{H}^s \rightarrow \mathbb{R}$,

$$\int_{\mathcal{H}^s(\mathbb{R}^d)} F(v) d\mu_\gamma(v) = \int_{\Omega} F(\gamma^\omega) d\mathbb{P}(\omega).$$

In particular, we can check that μ_γ satisfies:

- If $\gamma \in \mathcal{H}^s \setminus \mathcal{H}^{s+\varepsilon}$, then $\mu_\gamma(\mathcal{H}^{s+\varepsilon}) = 0$.
- Assume that for all $j \geq 1$ such that $I(j) \neq \emptyset$ we have $c_j \neq 0$. Then for all nonempty open subsets $B \subset \mathcal{H}^s$, $\mu_\gamma(B) > 0$.

Finally, we denote by \mathcal{M}^s the set of all such measures, $\mathcal{M}^s = \bigcup_{\gamma \in \mathcal{A}_s} \{\mu_\gamma\}$.

1B. Main results. Before we state our results, let us recall some facts concerning the deterministic study of the nonlinear Schrödinger equation (1-1). We say that (1-1) is locally well-posed in $\mathcal{H}^s(\mathbb{R}^d)$ if, for any initial condition $u_0 \in \mathcal{H}^s(\mathbb{R}^d)$, there exists a unique local in time solution $u \in \mathcal{C}([-T, T]; \mathcal{H}^s(\mathbb{R}^d))$, and if the flow-map is uniformly continuous. We denote by

$$s_c = \frac{d}{2} - \frac{2}{p-1}$$

the critical Sobolev index. Then one can show that NLS is well-posed in $\mathcal{H}^s(\mathbb{R}^d)$ when $s > \max(s_c, 0)$, and ill-posed when $s < s_c$. We refer to the introduction of [Thomann 2009] for more details on this topic.

1B1. Local existence results. We are now able to state our first result on the local well-posedness of (1-1).

Theorem 1.1. *Let $d \geq 2$, let $p \geq 3$ be an odd integer, and fix $\mu = \mu_\gamma \in \mathcal{M}^0$. Then there exists $\Sigma \subset L^2(\mathbb{R}^d)$ with $\mu(\Sigma) = 1$ and such that:*

(i) *For all $u_0 \in \Sigma$ there exist $T > 0$ and a unique local solution u to (1-1) with initial data u_0 satisfying*

$$u(t) - e^{-itH} u_0 \in \mathcal{C}([-T, T]; \mathcal{H}^s(\mathbb{R}^d)), \tag{1-4}$$

for some s such that $\frac{d}{2} - \frac{2}{p-1} < s < \frac{d}{2}$.

(ii) *More precisely, for all $T > 0$, there exists $\Sigma_T \subset \Sigma$ with*

$$\mu(\Sigma_T) \geq 1 - C \exp(-cT^{-\delta} \|\gamma\|_{L^2(\mathbb{R}^2)}^{-2}), \quad C, c, \delta > 0,$$

and such that for all $u_0 \in \Sigma_T$ the lifespan of u is larger than T .

Let $\gamma = \sum_{n=0}^{+\infty} c_n \varphi_n(x)$. Then

$$u_0^\omega := \sum_{n=0}^{+\infty} g_n(\omega) c_n \varphi_n(x)$$

is a typical element in the support of μ_γ . Another way to state Theorem 1.1 is: for any $T > 0$, there exists an event $\Omega_T \subset \Omega$ such that

$$\mathbb{P}(\Omega_T) \geq 1 - C \exp(-cT^{-\delta} \|\gamma\|_{L^2(\mathbb{R}^d)}^{-2}), \quad C, c, \delta > 0,$$

and that for all $\omega \in \Omega_T$, there exists a unique solution of the form (1-4) to (1-1) with initial data u_0^ω .

We will see in Proposition 2.1 that the stochastic approach yields a gain of $d/2$ derivatives compared to the deterministic theory. To prove Theorem 1.1 we only have to gain $s_c = d/2 - 2/(p-1)$ derivatives. The solution is constructed by a fixed point argument in a Strichartz space $X_T^s \subset \mathcal{C}([-T, T]; \mathcal{H}^s(\mathbb{R}^d))$ with continuous embedding, and uniqueness holds in the class X_T^s .

The deterministic Cauchy problem for (1-1) was studied by Oh [1989] (see also [Cazenave 2003, Chapter 9] for more references). Thomann [2009] has proven an almost sure local existence result for (1-1) in the supercritical regime (with a gain of $\frac{1}{4}$ of a derivative), for any $d \geq 1$. This local existence result was improved by [Burq et al. 2010] when $d = 1$ (gain of $\frac{1}{2}$ a derivative), by [Deng 2012] when $d = 2$, and by Poiret [2012a; 2012b] in any dimension.

Remark 1.2. The results of Theorem 1.1 also hold true for any quadratic potential

$$V(x) = \sum_{1 \leq j \leq d} \alpha_j x_j^2, \quad \alpha_j > 0, \quad 1 \leq j \leq d,$$

and for more general potentials such that $V(x) \approx \langle x \rangle^2$.

1B2. Global existence and scattering results for NLS. As an application of the results of the previous part, we are able to construct global solutions to the nonlinear Schrödinger equation without potential, which scatter when $t \rightarrow \pm\infty$. Consider the equation

$$\begin{cases} i \frac{\partial u}{\partial t} + \Delta u = \pm |u|^{p-1} u, & (t, x) \in \mathbb{R} \times \mathbb{R}^d. \\ u(0) = u_0. \end{cases} \tag{1-5}$$

The well-posedness indexes for this equation are the same as for (1-1). Namely, (1-5) is well-posed in $H^s(\mathbb{R}^d)$ when $s > \max(s_c, 0)$, and ill-posed when $s < s_c$.

For the next result, we will need an additional condition on the law ν . We assume that

$$\mathbb{P}(|g_n| < \rho) > 0 \quad \text{for all } \rho > 0, \tag{1-6}$$

which ensures that the random variable can take arbitrarily small values. Then we can prove:

Theorem 1.3. *Let $d \geq 2$, let $p \geq 3$ be an odd integer, and fix $\mu = \mu_\nu \in \mathcal{M}^0$. Assume that (1-6) holds. Then there exists $\Sigma \subset L^2(\mathbb{R}^d)$ with $\mu(\Sigma) > 0$ and such that:*

(i) *For all $u_0 \in \Sigma$ there exists a unique global solution u to (1-5) with initial data u_0 satisfying*

$$u(t) - e^{it\Delta} u_0 \in \mathcal{C}(\mathbb{R}; \mathcal{H}^s(\mathbb{R}^d)),$$

for some s such that $\frac{d}{2} - \frac{2}{p-1} < s < \frac{d}{2}$.

(ii) *For all $u_0 \in \Sigma$ there exist states $f_+, f_- \in \mathcal{H}^s(\mathbb{R}^d)$ such that when $t \rightarrow \pm\infty$,*

$$\|u(t) - e^{it\Delta}(u_0 + f_\pm)\|_{H^s(\mathbb{R}^d)} \rightarrow 0.$$

(iii) *If we assume that the distribution of ν is symmetric, then*

$$\mu(u_0 \in L^2(\mathbb{R}^d) : \text{assertion (ii) holds true} \mid \|u_0\|_{L^2(\mathbb{R}^d)} \leq \eta) \rightarrow 1,$$

when $\eta \rightarrow 0$.

We can show [Poiret 2012a, Théorème 20] that for all $s > 0$, if $u_0 \notin \mathcal{H}^s(\mathbb{R}^d)$ then $\mu(H^s(\mathbb{R}^d)) = 0$. This shows that the randomisation does not yield a gain of derivative in the Sobolev scale; thus Theorem 1.3 gives results for initial conditions not covered by the deterministic theory.

There is a large literature on the deterministic local and global theory with scattering for (1-5). We refer to [Banica et al. 2008; Nakanishi and Ozawa 2002; Carles 2009] for such results and more references.

We do not give here the details of the proof of Theorem 1.3, since one can follow the main lines of the argument of Poiret [2012a; 2012b] but with different constants (see, e.g., [Poiret 2012b, Théorème 4]). The proof of (i) and (ii) is based on the use of an explicit transform, called the lens transform \mathcal{L} , which links the solutions of (1-5) to solutions of NLS with harmonic potential. The transform \mathcal{L} has been used in different contexts; see [Carles 2009] for scattering results and more references. More precisely, for $u(t, x) :]-\frac{\pi}{4}, \frac{\pi}{4}[\times \mathbb{R}^d \rightarrow \mathbb{C}$ we define

$$v(t, x) = \mathcal{L}u(t, x) = \left(\frac{1}{\sqrt{1+4t^2}} \right)^{d/2} u \left(\frac{\arctan(2t)}{2}, \frac{x}{\sqrt{1+4t^2}} \right) e^{i|x|^2 t / (1+4t^2)},$$

then u is a solution to

$$i \frac{\partial u}{\partial t} - Hu = \lambda \cos(2t)^{\frac{1}{2}d(p-1)-2} |u|^{p-1} u$$

if and only if v satisfies $i \partial v / \partial t + \Delta v = \lambda |v|^{p-1} v$. **Theorem 1.1** provides solutions with lifespan larger than $\pi/4$ for large probabilities, provided that the initial conditions are small enough.

Part (iii) is stated in [Poiret 2012a, Théorème 9], and can be understood as a small data result.

In **Theorem 1.3** we assumed that $d \geq 2$ and that $p \geq 3$ was an odd integer, so we had $p \geq 1 + 4/d$, or, in other words, we were in an L^2 -supercritical setting. Our approach also allows to get results in an L^2 -subcritical context, i.e., when $1 + 2/d < p < 1 + 4/d$.

Theorem 1.4. *Let $d = 2$ and $2 < p < 3$. Assume that (1-6) holds and fix $\mu = \mu_\gamma \in \mathcal{M}^0$. Then there exists $\Sigma \subset L^2(\mathbb{R}^2)$ with $\mu(\Sigma) > 0$ and such that for all $0 < \varepsilon < 1$:*

(i) *For all $u_0 \in \Sigma$ there exists a unique global solution u to (1-5) with initial data u_0 satisfying*

$$u(t) - e^{it\Delta} u_0 \in \mathcal{C}(\mathbb{R}; \mathcal{H}^{1-\varepsilon}(\mathbb{R}^2)).$$

(ii) *For all $u_0 \in \Sigma$ there exist states $f_+, f_- \in \mathcal{H}^{1-\varepsilon}(\mathbb{R}^2)$ such that when $t \rightarrow \pm\infty$,*

$$\|u(t) - e^{it\Delta}(u_0 + f_\pm)\|_{H^{1-\varepsilon}(\mathbb{R}^2)} \rightarrow 0.$$

(iii) *If we assume that the distribution of v is symmetric, then*

$$\mu(u_0 \in L^2(\mathbb{R}^2) : \text{assertion (ii) holds true} \mid \|u_0\|_{L^2(\mathbb{R}^2)} \leq \eta) \rightarrow 1,$$

when $\eta \rightarrow 0$.

In the case $p \leq 1 + 2/d$, Barab [1984] showed that a nontrivial solution to (1-5) never scatters; therefore even with a stochastic approach one can not have scattering in this case. When $d = 2$, the condition $p > 2$ in **Theorem 1.4** is therefore optimal. Usually, deterministic scattering results in L^2 -subcritical contexts are obtained in the space $H^1 \cap \mathcal{F}(H^1)$. Here we assume $u_0 \in L^2$, and thus we relax both the regularity and the decay assumptions (this latter point is the most striking in this context). Again we refer to [Banica et al. 2008] for an overview of scattering theory for NLS.

When $\mu \in \mathcal{M}^\sigma$ for some $0 < \sigma < 1$ we are able to prove the same result with $\varepsilon = 0$. Since the proof is much easier, we give it before the case $\sigma = 0$ (see **Section 3B**).

Finally, we point out that in **Theorem 1.4** we are only able to consider the case $d = 2$ because of the lack of regularity of the nonlinear term $|u|^{p-1} u$.

1B3. *Global existence results for NLS with quadratic potential.* We also get global existence results for defocusing Schrödinger equation with harmonic potential. For $d = 2$ or $d = 3$, consider the equation

$$\begin{cases} i \frac{\partial u}{\partial t} - Hu = |u|^2 u, & (t, x) \in \mathbb{R} \times \mathbb{R}^d, \\ u(0) = u_0, \end{cases} \tag{1-7}$$

and denote by E the energy of (1-7), namely

$$E(u) = \|u\|_{\mathcal{H}^1(\mathbb{R}^d)}^2 + \frac{1}{2} \|u\|_{L^4(\mathbb{R}^d)}^4.$$

Deterministic global existence for (1-7) has been studied by Zhang [2005] and by Carles [2011] in the case of time-dependent potentials.

When $d = 3$, our global existence result for (1-7) is the following:

Theorem 1.5. *Let $d = 3$, $\frac{1}{6} < s < 1$ and fix $\mu = \mu_\gamma \in \mathcal{M}^s$. Then there exists a set $\Sigma \subset \mathcal{H}^s(\mathbb{R}^3)$ such that $\mu(\Sigma) = 1$ and that the following holds true:*

(i) *For all $u_0 \in \Sigma$, there exists a unique global solution to (1-7), which reads*

$$u(t) = e^{-itH}u_0 + w(t), \quad w \in \mathcal{C}(\mathbb{R}, \mathcal{H}^1(\mathbb{R}^3)).$$

(ii) *The previous line defines a global flow Φ , which leaves the set Σ invariant:*

$$\Phi(t)(\Sigma) = \Sigma, \quad \text{for all } t \in \mathbb{R}.$$

(iii) *There exist $C, c_s > 0$ such that, for all $t \in \mathbb{R}$,*

$$E(w(t)) \leq C(M + |t|)^{c_s+},$$

where M is a positive random variable such that

$$\mu(u_0 \in \mathcal{H}^s(\mathbb{R}^3) : M > K) \leq C e^{-cK^\delta / \|\gamma\|_{\mathcal{H}^s(\mathbb{R}^3)}^2}.$$

Here the critical Sobolev space is $\mathcal{H}^{1/2}(\mathbb{R}^3)$; thus the local deterministic theory, combined with the conservation of the energy, immediately gives global well-posedness in $\mathcal{H}^1(\mathbb{R}^3)$. Using a kind of interpolation method due to Bourgain, one may obtain deterministic global well-posedness in $\mathcal{H}^s(\mathbb{R}^3)$ for some $1/2 < s < 1$. Instead, for the proof of Theorem 1.5, we will rely on the almost well-posedness result of Theorem 1.1, and this gives global well-posedness in a supercritical context.

The constant $c_s > 0$ can be computed explicitly (see (4-16)), and we do not think that we have obtained the optimal rate. By reversibility of the equation, it is enough to consider only positive times.

With a similar approach, in dimension $d = 2$, we can prove:

Theorem 1.6. *Let $d = 2$, $0 < s < 1$ and fix $\mu = \mu_\gamma \in \mathcal{M}^s$. Then there exists a set $\Sigma \subset \mathcal{H}^s(\mathbb{R}^2)$ such that $\mu(\Sigma) = 1$ and that, for all $u_0 \in \Sigma$, there exists a unique global solution to (1-7),*

$$u(t) = e^{-itH}u_0 + w(t), \quad w \in \mathcal{C}(\mathbb{R}, \mathcal{H}^1(\mathbb{R}^2)).$$

In addition, statements (ii) and (iii) of Theorem 1.5 are also satisfied with $c_s = \frac{1-s}{s}$.

Here the critical Sobolev space is $L^2(\mathbb{R}^2)$; thus Theorem 1.6 shows global well-posedness for any subcritical cubic nonlinear Schrödinger equations in dimension two.

Using the smoothing effect, which yields a gain of $\frac{1}{2}$ a derivative, a global well-posedness result for (1-1), in the defocusing case, was given in [Burq et al. 2010] in the case $d = 1$, for any $p \geq 3$. The global existence is proved for a typical initial condition on the support of a Gibbs measure, which is $\bigcap_{\sigma>0} \mathcal{H}^{-\sigma}(\mathbb{R})$. This result was extended by Deng [2012] in dimension $d = 2$ for radial functions. However, this approach has the drawback that it relies on the invariance of a Gibbs measure, which is a

rigid object, and is supported in rough Sobolev spaces. Therefore it seems difficult to adapt this strategy in higher dimensions.

Here instead we obtain the results of Theorems 1.5 and 1.6 as a combination of Theorem 1.1 with the high-low frequency decomposition method of [Bourgain 1999, p. 84]. This approach has been successful in different contexts, and has been first used together with probabilistic arguments by Colliander and Oh [2012] for the cubic Schrödinger below $L^2(\mathbb{S}^1)$ and later on by Burq and Tzvetkov [2014] for the wave equation.

1C. Notations and plan of the paper. In this paper $c, C > 0$ denote constants, the value of which may change from line to line. These constants will always be universal, or uniformly bounded with respect to the other parameters.

We let $L_T^p = L_{[-T,T]}^p = L^p(-T, T)$ for $T > 0$ and we write $L_x^p = L^p(\mathbb{R}^d)$. We denote the harmonic oscillator on \mathbb{R}^d by $H = -\Delta + |x|^2 = \sum_{j=1}^d (-\partial_j^2 + x_j^2)$, and for $s \geq 0$ we define the Sobolev space \mathcal{H}^s by the norm $\|u\|_{\mathcal{H}^s} = \|H^{s/2}u\|_{L^2(\mathbb{R}^d)}$. More generally, we define the spaces ${}^{\mathcal{W}}W^{s,p}$ by the norm $\|u\|_{{}^{\mathcal{W}}W^{s,p}} = \|H^{s/2}u\|_{L^p(\mathbb{R}^d)}$. If E is a Banach space and μ is a measure on E , we write $L_\mu^p = L^p(d\mu)$ and $\|u\|_{L_\mu^p E} = \|\|u\|_E\|_{L_\mu^p}$.

The rest of the paper is organised as follows. In Section 2 we recall some deterministic results on the spectral function, and prove stochastic Strichartz estimates. Section 3 is devoted to the proof of Theorem 1.1 and of the scattering results for NLS without potential. Finally, in Section 4 we study the global existence for the Schrödinger–Gross–Pitaevskii equation (1-1).

2. Stochastic Strichartz estimates

The main result of this section is the following probabilistic improvement of the Strichartz estimates.

Proposition 2.1. *Let $s \in \mathbb{R}$ and $\mu = \mu_\gamma \in \mathcal{M}^s$. Let $1 \leq q < +\infty$, $2 \leq r \leq +\infty$, and set $\alpha = d(\frac{1}{2} - \frac{1}{r})$ if $r < +\infty$ and $\alpha < d/2$ if $r = +\infty$. Then there exist $c, C > 0$ such that, for all $\tau \in \mathbb{R}$,*

$$\mu\left(u \in \mathcal{H}^s(\mathbb{R}^d) : \|e^{-i(t+\tau)H}u\|_{L_{[0,T]}^q {}^{\mathcal{W}}W^{s+\alpha,r}(\mathbb{R}^d)} > K\right) \leq Ce^{-cK^2/T^{2/q}} \|\gamma\|_{\mathcal{H}^s(\mathbb{R}^d)}^2.$$

When $r = +\infty$, this result expresses a gain μ -a.s. of $d/2$ derivatives in space compared to the deterministic Strichartz estimates (see the bound (3-2)).

Proposition 2.1 is a consequence of [Poiret et al. 2013, Inequality (1.6)], but we give here a self-contained proof suggested by Nicolas Burq.

There are two key ingredients in the proof of Proposition 2.1. The first one is a deterministic estimate on the spectral function given in Lemma 2.2, and the second is the Khinchin inequality stated in Lemma 2.3.

2A. Deterministic estimates of the spectral function. We define the spectral function π_H for the harmonic oscillator by

$$\pi_H(\lambda; x, y) = \sum_{\lambda_j \leq \lambda} \varphi_j(x) \overline{\varphi_j(y)},$$

and this definition does not depend on the choice of $\{\varphi_j \mid j \in \mathbb{N}\}$.

Let us recall some results of π_H , which were essentially obtained by Thangavelu [1993, Lemma 3.2.2, p. 70] (see also [Karadzhov 1995] and [Poiret et al. 2013, Section 3] for more details).

Thanks to the Mehler formula, we can prove

$$\pi_H(\lambda; x, x) \leq C\lambda^{d/2} \exp\left(-c\frac{|x|^2}{\lambda}\right) \quad \text{for all } x \in \mathbb{R}^d \text{ and } \lambda \geq 1. \tag{2-1}$$

One also has the following more subtle bound, which is the heart of [Karadzhov 1995]:

$$|\pi_H(\lambda + \mu; x, x) - \pi_H(\lambda; x, x)| \leq C(1 + |\mu|)\lambda^{d/2-1} \quad \text{for } \lambda \geq 1, |\mu| \leq C_0\lambda. \tag{2-2}$$

This inequality gives a bound on π_H in energy interval of size ~ 1 , which is the finest one can obtain.

Then we can prove (see [Poiret et al. 2013, Lemma 3.5]):

Lemma 2.2. *Let $d \geq 2$ and assume that $|\mu| \leq c_0, r \geq 1$ and $\theta \geq 0$. Then there exists $C > 0$ such that for all $\lambda \geq 1$*

$$\|\pi_H(\lambda + \mu; x, x) - \pi_H(\lambda; x, x)\|_{L^r(\mathbb{R}^d)} \leq C\lambda^{\frac{1}{2}d(1+1/r)-1}.$$

2B. Proof of Proposition 2.1. To begin with, recall the Khinchin inequality, which shows a smoothing property of the random series in the L^k spaces for $k \geq 2$; for example, see [Burq and Tzvetkov 2008a, Lemma 4.2].

Lemma 2.3. *There exists $C > 0$ such that for all real $k \geq 2$ and $(c_n) \in \ell^2(\mathbb{N})$*

$$\left\| \sum_{n \geq 1} g_n(\omega) c_n \right\|_{L^k_{\mathbb{P}}} \leq C\sqrt{k} \left(\sum_{n \geq 1} |c_n|^2 \right)^{\frac{1}{2}}.$$

Now we fix $\gamma = \sum_{n=0}^{+\infty} c_n \varphi_n \in \mathcal{A}_s$ and let $\gamma^\omega = \sum_{n=0}^{+\infty} g_n(\omega) c_n \varphi_n$.

Firstly, we treat the case $r < +\infty$. Set $\alpha = d(\frac{1}{2} - \frac{1}{r})$ and set $\sigma = s + \alpha$. Observe that it suffices to prove the estimation for $K \gg \|\gamma\|_{\mathcal{H}^s(\mathbb{R}^d)}$.

Let $k \geq 1$. By definition,

$$\begin{aligned} \int_{\mathcal{H}^s(\mathbb{R}^d)} \|e^{-i(t+\tau)H} u\|_{L^q_{[0,T]W^{\sigma,r}(\mathbb{R}^d)}}^k d\mu(u) &= \int_{\Omega} \|e^{-i(t+\tau)H} \gamma^\omega\|_{L^q_{[0,T]W^{\sigma,r}(\mathbb{R}^d)}}^k d\mathbb{P}(\omega) \\ &= \int_{\Omega} \|e^{-i(t+\tau)H} H^{\sigma/2} \gamma^\omega\|_{L^q_{[0,T]L^r(\mathbb{R}^d)}}^k d\mathbb{P}(\omega). \end{aligned} \tag{2-3}$$

Since $e^{-i(t+\tau)H} H^{\sigma/2} \gamma^\omega(x) = \sum_{n=0}^{+\infty} g_n(\omega) c_n \lambda_n^{\sigma/2} e^{-i(t+\tau)\lambda_n} \varphi_n(x)$, by Lemma 2.3 we get

$$\|e^{-i(t+\tau)H} H^{\sigma/2} \gamma^\omega(x)\|_{L^k_{\mathbb{P}}} \leq C\sqrt{k} \|e^{-i(t+\tau)H} H^{\sigma/2} \gamma^\omega(x)\|_{L^2_{\mathbb{P}}} = C\sqrt{k} \left(\sum_{n=0}^{+\infty} \lambda_n^\sigma |c_n|^2 |\varphi_n(x)|^2 \right)^{\frac{1}{2}}.$$

Assume that $k \geq r$. By the integral Minkowski inequality, the previous line and the triangle inequality we get

$$\begin{aligned} \|e^{-i(t+\tau)H} H^{\sigma/2} \gamma^\omega\|_{L^k_{\mathbb{P}} L^r_x} &\leq \|e^{-i(t+\tau)H} H^{\sigma/2} \gamma^\omega\|_{L^r_x L^k_{\mathbb{P}}} \\ &\leq C\sqrt{k} \left\| \sum_{k=0}^{+\infty} \lambda_k^\sigma |c_k|^2 |\varphi_k|^2 \right\|_{L^{r/2}(\mathbb{R}^d)}^{\frac{1}{2}} \\ &\leq C\sqrt{k} \left(\sum_{j=1}^{+\infty} \left\| \sum_{k \in I(j)} \lambda_k^\sigma |c_k|^2 |\varphi_k|^2 \right\|_{L^{r/2}(\mathbb{R}^d)} \right)^{\frac{1}{2}}. \end{aligned} \tag{2-4}$$

Condition (1-3) implies that for all $x \in \mathbb{R}^d$ and $k \in I(j) = \{n \in \mathbb{N} \mid 2j \leq \lambda_n < 2(j+1)\}$

$$\lambda_k^\sigma |c_k|^2 |\varphi_k(x)|^2 \leq Cj^\sigma \sum_{n \in I(j)} |c_n|^2 \frac{|\varphi_n(x)|^2}{\#I(j)},$$

and thus, by Lemma 2.2 and the fact that $\#I(j) \sim cj^{d-1}$,

$$\begin{aligned} \left\| \sum_{k \in I(j)} \lambda_k^\sigma |c_k|^2 |\varphi_k(x)|^2 \right\|_{L^{r/2}(\mathbb{R}^d)} &\leq Cj^\sigma \sum_{n \in I(j)} |c_n|^2 \frac{\left\| \sum_{k \in I(j)} |\varphi_k(x)|^2 \right\|_{L^{r/2}(\mathbb{R}^d)}}{\#I(j)} \\ &\leq Cj^{\sigma+d(1/r-1/2)} \sum_{n \in I(j)} |c_n|^2 \\ &= Cj^s \sum_{n \in I(j)} |c_n|^2. \end{aligned}$$

The latter inequality together with (2-4) gives

$$\|e^{-i(t+\tau)H} H^{\sigma/2} \gamma^\omega\|_{L^k_{\mathbb{P}} L^r_x} \leq C\sqrt{k} \|\gamma\|_{\mathcal{H}^s(\mathbb{R}^d)},$$

and for $k \geq r$, by Minkowski,

$$\|e^{-i(t+\tau)H} H^{\sigma/2} \gamma^\omega\|_{L^k_{\mathbb{P}} L^q_{[0,T]} L^r_x} \leq C\sqrt{k} T^{1/q} \|\gamma\|_{\mathcal{H}^s(\mathbb{R}^d)}.$$

Then, using (2-3) and the Bienaymé–Chebishev inequality, we obtain

$$\begin{aligned} \mu(u \in \mathcal{H}^s : \|e^{-i(t+\tau)H} u\|_{L^q_{[0,T]} \mathcal{W}^{\sigma,r}(\mathbb{R}^d)} > K) &\leq (K^{-1} \|e^{-i(t+\tau)H} H^{\sigma/2} \gamma^\omega\|_{L^k_{\mathbb{P}} L^q_{[0,T]} L^r_x})^k \\ &\leq (CK^{-1} \sqrt{k} T^{1/q} \|\gamma\|_{\mathcal{H}^s(\mathbb{R}^d)})^k. \end{aligned}$$

Finally, if $K \gg \|\gamma\|_{\mathcal{H}^s(\mathbb{R}^d)}$, we can choose $k = K^2/2CT^{2/q} \|\gamma\|_{\mathcal{H}^s(\mathbb{R}^d)}^2 \geq r$, which yields the result.

Now assume $r = +\infty$. We use the Sobolev inequality to get $\|u\|_{\mathcal{W}^{s,\infty}} \leq C\|u\|_{\mathcal{W}^{s,\tilde{r}}}$ with $\tilde{s} = s + 2d/\tilde{r}$ for $\tilde{r} \geq 1$ large enough; hence we can apply the previous result for $r < +\infty$.

Remark 2.4. A similar result to Proposition 2.1 holds, with the same gain of derivatives, when $I(\lambda)$ is replaced with the dyadic interval $J(j) = \{n \in \mathbb{N} \mid 2^j \leq \lambda_n < 2^{j+1}\}$. Then the condition (1-3) becomes

$$|c_k|^2 \leq \frac{C}{\#J(j)} \sum_{n \in J(j)} |c_n|^2 \quad \text{for all } j \geq 1 \text{ and } k \in J(j), \tag{2-5}$$

which seems more restrictive. Indeed neither condition imply the other.

Observe that if we want to prove the result under condition (2-5), the subtle estimate (2-2) is not needed; (2-1) is enough.

Remark 2.5. For $d = 1$, condition (1-3) is always satisfied but condition (2-2) is not. Instead we can use that $\|\varphi_k\|_p \leq C\lambda_k^{-\theta(p)}$ with $\theta(p) > 0$ for $p > 2$ [Koch and Tataru 2005]. For example if $p > 4$ we have $\theta(p) = \frac{1}{4} - (p - 1)/6p$. Thus we get the Proposition 2.1 with $s = p\theta(p)/4$ (see [Thomann 2009; Burq et al. 2010], where this is used).

Remark 2.6. Another approach could have been to exploit the particular basis $(\varphi_n)_{n \geq 1}$, which satisfies the good L^∞ estimates given in [Poiret et al. 2013, Theorem 1.3], and to construct the measures μ as the image measures of random series of the form

$$\gamma^\omega(x) = \sum_{n \geq 1} c_n g_n(\omega) \varphi_n(x),$$

with $c_n \in \ell^2(\mathbb{N})$ not necessarily satisfying (1-3). A direct application of the Khinchin inequality (as in [Thomann 2009, Proposition 2.3]) then gives the same bounds as in Proposition 2.1. Observe that condition (1-3) is also needed in this approach, but it directly intervenes in the construction of the φ_n .

We believe that the strategy we adopted here is slightly more general, since it seems to work even in cases where we do not have a basis of eigenfunctions that satisfy bounds analogous to [Poiret et al. 2013, Theorem 1.3], as for example in the case of the operator $-\Delta + |x|^4$.

3. Application to the local theory of the supercritical Schrödinger equation

3A. Almost sure local well-posedness. This subsection, devoted to the proof of Theorem 1.1, follows the argument of [Poiret 2012b].

Let $u_0 \in L^2(\mathbb{R}^d)$. We look for a solution to (1-1) of the form $u = e^{-itH}u_0 + v$, where v is some fluctuation term more regular than the linear profile $e^{-itH}u_0$. By the Duhamel formula, the unknown v has to be a fixed point of the operator

$$L(v) := \mp i \int_0^t e^{-i(t-s)H} |e^{-isH}u_0 + v(s)|^{p-1} (e^{-isH}u_0 + v(s)) ds, \tag{3-1}$$

in some adequate functional space, which is a Strichartz space.

To begin with, we recall the Strichartz estimates for the harmonic oscillator. A couple $(q, r) \in [2, +\infty]^2$ is called admissible if

$$\frac{2}{q} + \frac{d}{r} = \frac{d}{2} \quad \text{and} \quad (d, q, r) \neq (2, 2, +\infty),$$

and if one defines

$$X_T^s := \bigcap_{\substack{(q,r) \\ \text{admissible}}} L^q([-T, T]; \mathcal{W}^{s,r}(\mathbb{R}^d)),$$

then for all $T > 0$ there exists $C_T > 0$ such that for all $u_0 \in \mathcal{H}^s(\mathbb{R}^d)$ we have

$$\|e^{-itH} u_0\|_{X_T^s} \leq C_T \|u_0\|_{\mathcal{H}^s(\mathbb{R}^d)}. \tag{3-2}$$

We will also need the inhomogeneous version of Strichartz: For all $T > 0$, there exists $C_T > 0$ such that for all admissible couples (q, r) and functions $F \in L^{q'}([-T, T]; \mathcal{W}^{s,r'}(\mathbb{R}^d))$,

$$\left\| \int_0^t e^{-i(t-s)H} F(s) ds \right\|_{X_T^s} \leq C_T \|F\|_{L^{q'}([-T, T], \mathcal{W}^{s,r'}(\mathbb{R}^d))}, \tag{3-3}$$

where q' and r' are the Hölder conjugates of q and r . We refer to [Poiret 2012b] for a proof.

The next result is a direct application of the Sobolev embeddings and Hölder.

Lemma 3.1. *Let $(q, r) \in [2, \infty[\times [2, \infty]$, and let $s, s_0 \geq 0$ be such that $s - s_0 > \frac{d}{2} - \frac{2}{q} - \frac{d}{r}$. Then there exist $\kappa, C > 0$ such that for any $T \geq 0$ and $u \in X_T^s$,*

$$\|u\|_{L^q([-T, T], \mathcal{W}^{s_0, r}(\mathbb{R}^d))} \leq C T^\kappa \|u\|_{X_T^s}.$$

We now introduce the appropriate sets in which we can profit from the stochastic estimates of the previous section. Fix $\mu = \mu_\gamma \in M^0$ and, for $K \geq 0$ and $\varepsilon > 0$, define the set $G_d(K)$ as

$$G_d(K) = \{w \in L^2(\mathbb{R}^d) \mid \|w\|_{L^2(\mathbb{R}^d)} \leq K \text{ and } \|e^{-itH} w\|_{L^{1/\varepsilon}_{[-2\pi, 2\pi]} \mathcal{W}^{d/2-\varepsilon, \infty}(\mathbb{R}^d)} \leq K\}.$$

Then by Proposition 2.1,

$$\mu(G_d(K)^c) \leq \mu(\|w\|_{L^2(\mathbb{R}^d)} > K) + \mu(\|e^{-itH} w\|_{L^{1/\varepsilon}_{[-2\pi, 2\pi]} \mathcal{W}^{d/2-\varepsilon, \infty}(\mathbb{R}^d)} > K) \leq C e^{-cK^2/\|\gamma\|_{L^2}^2}. \tag{3-4}$$

We want to perform a fixed point argument on L with initial condition $u_0 \in G_d(K)$ for some $K > 0$ and $\varepsilon > 0$ small enough. We begin by establishing some estimates.

Lemma 3.2. *Let $s \in]\frac{d}{2} - \frac{2}{p-1}, \frac{d}{2}[$. For $\varepsilon > 0$ small enough there exist $C > 0$ and $\kappa > 0$ such that for any $0 < T \leq 1$, $u_0 \in G_d(K)$, $v \in X_T^s$ and $f_i = v$ or $f_i = e^{-itH} u_0$,*

$$\left\| H^{s/2}(v) \prod_{i=2}^p f_i \right\|_{L^1([-T, T], L^2(\mathbb{R}^d))} \leq C T^\kappa (K^p + \|v\|_{X_T^s}^p), \tag{3-5}$$

and

$$\left\| H^{s/2}(e^{-itH} u_0) \prod_{i=2}^p f_i \right\|_{L^1([-T, T], L^2(\mathbb{R}^d))} \leq C T^\kappa (K^p + \|v\|_{X_T^s}^p). \tag{3-6}$$

Proof. First we prove (3-5). Thanks to the Hölder inequality,

$$\left\| |\nabla|^s(v) \prod_{i=2}^p f_i \right\|_{L^1([-T, T], L^2(\mathbb{R}^d))} \leq \| |\nabla|^s(v) \|_{L^\infty([-T, T], L^2(\mathbb{R}^d))} \prod_{i=2}^p \|f_i\|_{L^{p-1}([-T, T], L^\infty(\mathbb{R}^d))}$$

and

$$\begin{aligned} \left\| \langle x \rangle^s v \prod_{i=2}^p f_i \right\|_{L^1([-T, T], L^2(\mathbb{R}^d))} &\leq \|\langle x \rangle^s v\|_{L^\infty([-T, T], L^2(\mathbb{R}^d))} \prod_{i=2}^p \|f_i\|_{L^{p-1}([-T, T], L^\infty(\mathbb{R}^d))} \\ &\leq \|v\|_{L^\infty([-T, T], H^s(\mathbb{R}^d))} \prod_{i=2}^p \|f_i\|_{L^{p-1}([-T, T], L^\infty(\mathbb{R}^d))}. \end{aligned}$$

If $f_i = v$, then as $s > \frac{d}{2} - \frac{2}{p-1}$, we can use [Lemma 3.1](#) to obtain

$$\|v\|_{L^{p-1}([-T, T], L^\infty(\mathbb{R}^d))} \leq CT^\kappa \|v\|_{X_T^s}.$$

If $f_i = e^{-itH} u_0$, then by definition of $G_d(K)$ we have, for $\varepsilon > 0$ small enough,

$$\|e^{-itH} u_0\|_{L^{p-1}([-T, T], L^\infty(\mathbb{R}^d))} \leq T^\kappa \|e^{-itH} u_0\|_{L^{1/\varepsilon}([-2\pi, 2\pi], \mathcal{W}^{d/2-\varepsilon, \infty}(\mathbb{R}^d))} \leq T^\kappa K.$$

We now turn to (3-6). Thanks to the Hölder inequality, we have

$$\begin{aligned} \left\| |\nabla|^s (e^{-itH} u_0) \prod_{i=2}^p f_i \right\|_{L^1([-T, T], L^2(\mathbb{R}^d))} &\leq \| |\nabla|^s (e^{-itH} u_0) \|_{L^p([-T, T], L^{2dp}(\mathbb{R}^d))} \prod_{i=2}^p \|f_i\|_{L^p([-T, T], L^{2dp(p-1)/(dp-1)}(\mathbb{R}^d))} \\ &\leq \|e^{-itH} u_0\|_{L^p([-T, T], \mathcal{W}^{s, 2dp}(\mathbb{R}^d))} \prod_{i=2}^p \|f_i\|_{L^p([-T, T], L^{2dp(p-1)/(dp-1)}(\mathbb{R}^d))}. \end{aligned}$$

If $f_i = e^{-itH} u_0$, by interpolation we obtain, for some $0 \leq \theta \leq 1$,

$$\|e^{-itH} u_0\|_{L^p([-T, T], L^{2dp(p-1)/(dp-1)}(\mathbb{R}^d))} \leq CT^\kappa \|u_0\|_{L^2(\mathbb{R}^d)}^{1-\theta} \|e^{-itH} u_0\|_{L^{1/\varepsilon}([-T, T], L^\infty(\mathbb{R}^d))}^\theta \leq CT^\kappa K.$$

If $f_i = v$, as $s > \frac{d}{2} - \frac{2}{p-1} > \frac{d}{2} - \frac{2}{p} - \frac{d(dp-1)}{2dp(p-1)}$ (because $p \geq 3$ and $d \geq 2$), then thanks to [Lemma 3.1](#) we find

$$\|v\|_{L^p([-T, T], L^{2dp(p-1)/(dp-1)}(\mathbb{R}^d))} \leq CT^\kappa \|v\|_{X_T^s}. \quad \square$$

We are now able to establish the estimates that will be useful in the application of a fixed point theorem.

Proposition 3.3. *Let $s \in]\frac{d}{2} - \frac{2}{p-1}, \frac{d}{2}[$. Then for $\varepsilon > 0$ small enough, there exist $C > 0$ and $\kappa > 0$ such that if $u_0 \in G_d(K)$ for some $K > 0$. For any $v, v_1, v_2 \in X_T^s$ and $0 < T \leq 1$,*

$$\left\| \int_0^t e^{-i(t-s)H} |e^{-isH} u_0 + v|^{p-1} (e^{-isH} u_0 + v) ds \right\|_{X_T^s} \leq CT^\kappa (K^p + \|v\|_{X_T^s}^p),$$

and

$$\begin{aligned} \left\| \int_0^t e^{-i(t-s)H} |e^{-isH} u_0 + v_1|^{p-1} (e^{-isH} u_0 + v_1) ds \right. \\ \left. - \int_0^t e^{-i(t-s)H} |e^{-isH} u_0 + v_2|^{p-1} (e^{-isH} u_0 + v_2) ds \right\|_{X_T^s} \\ \leq CT^\kappa \|v_1 - v_2\|_{X_T^s} (K^{p-1} + \|v_1\|_{X_T^s}^{p-1} + \|v_2\|_{X_T^s}^{p-1}). \end{aligned}$$

Proof. We only prove the first claim, since the proof of the second is similar. Using the Strichartz inequalities (3-3), we obtain

$$\begin{aligned} & \left\| \int_0^t e^{-i(t-s)H} |e^{-isH} u_0 + v|^{p-1} (e^{-isH} u_0 + v) ds \right\|_{X_T^s} \\ & \leq C \left\| |e^{-isH} u_0 + v|^{p-1} (e^{-isH} u_0 + v) \right\|_{L^1([-T, T], L^2(\mathbb{R}^d))}. \end{aligned}$$

Then, using Lemma 3.2, we obtain the existence of $\kappa > 0$ such that for any $u_0 \in G_d(K)$, $0 < T \leq 1$ and $v \in X_T^s$,

$$\left\| H^{s/2} (|e^{-isH} u_0 + v|^{p-1} (e^{-isH} u_0 + v)) \right\|_{L^1([-T, T], L^2(\mathbb{R}^d))} \leq CT^\kappa (K^p + \|v\|_{X_T^s}^p). \quad \square$$

Proof of Theorem 1.1. We now complete the contraction argument on L defined in (3-1) with some $u_0 \in G_d(K)$. According to Proposition 3.3, there exist $C > 0$ and $\kappa > 0$ such that

$$\begin{aligned} \|L(v)\|_{X_T^s} & \leq CT^\kappa (K^p + \|v\|_{X_T^s}^p) \\ \|L(v_1) - L(v_2)\|_{X_T^s} & \leq CT^\kappa \|v_1 - v_2\|_{X_T^s} (K^{p-1} + \|v_1\|_{X_T^s}^{p-1} + \|v_2\|_{X_T^s}^{p-1}). \end{aligned}$$

Hence, if we choose $T > 0$ such that $K = (8CT^\kappa)^{-1/(p-1)}$, then L is a contraction in the space $B_{X_T^s}(0, K)$ (the ball of radius K in X_T^s). Thus if we set $\Sigma_T = G_d(K)$, with the previous choice of K , the result follows from (3-4). \square

Proof of Theorem 1.3. We introduce

$$\begin{cases} i \frac{\partial w}{\partial t} - Hw = \pm \cos(2t)^{\frac{1}{2}d(p-1)-2} |w|^{p-1} w, & (t, x) \in \mathbb{R} \times \mathbb{R}^d, \\ v(0) = u_0, \end{cases} \quad (3-7)$$

and let $s \in]\frac{d}{2} - \frac{2}{p-1}, \frac{d}{2}[$, $T = \frac{\pi}{4}$ and $1 \gg \varepsilon > 0$. Thanks to Proposition 3.3, there exist $C > 0$ and $\kappa > 0$ such that if $u_0 \in G_d(K)$ for some $K > 0$ then, for all v ,

$$\left\| \int_0^t e^{-i(t-s)H} (\cos(2s)^{\frac{d}{2}(p-1)-2} |e^{-isH} u_0 + v|^{p-1} (e^{-isH} u_0 + v)) ds \right\|_{X_T^s} \leq CT^\kappa (K^p + \|v\|_{X_T^s}^p). \quad (3-8)$$

As in Theorem 1.1, we can choose $K = (8CT^\kappa)^{-1/(p-1)}$ to obtain, for $u_0 \in G_d(K)$, a unique local solution $w = e^{-itH} u_0 + v$ in time interval $]-\frac{\pi}{4}, \frac{\pi}{4}[$ to (3-7) with $v \in X_T^1$.

We set $u = \mathcal{L}v$. Then u is a global solution to (1-5). Thanks to [Poiret 2012b, Propositions 20 and 22], we obtain that $u = e^{it\Delta} u_0 + v'$ with $v' \in X_T^1$.

Moreover, thanks to (3-8), we have that

$$\int_0^t e^{-i(t-s)H} (\cos(2s)^{\frac{d}{2}(p-1)-2} |e^{-isH} u_0 + v|^{p-1} (e^{-isH} u_0 + v)) ds \in \mathcal{C}^0([-T, T], \mathcal{H}^s(\mathbb{R}^d)).$$

Then there exist $L \in \mathcal{H}^s$ such that

$$\lim_{t \rightarrow T} \left\| e^{-itH} \int_0^t e^{-isH} (\cos(2s)^{\frac{d}{2}(p-1)-2} |e^{-isH} u_0 + v|^{p-1} (e^{-isH} u_0 + v)) ds - L \right\|_{\mathcal{H}^s(\mathbb{R}^d)} = 0.$$

Using [Poiret 2012b, Lemma 70], we obtain that

$$\lim_{t \rightarrow T} \|u(t) - e^{it\Delta} u_0 - e^{it\Delta} (-i e^{-iT H} L)\|_{H^s(\mathbb{R}^d)} = 0.$$

Finally, to establish **Theorem 1.3**, it suffices to set $\Sigma = G_d(K)$ and to prove that $\mu(u_0 \in G_d(K)) > 0$. We can write

$$u_0 = \chi\left(\frac{H}{N}\right)u_0 + (1 - \chi)\left(\frac{H}{N}\right)u_0 := [u_0]_N + [u_0]^N,$$

where χ is a truncation function. Using the triangle inequality and independence, we obtain that

$$\mu(u_0 \in G_d(K)) \geq \mu([u_0]_N \in G_d(K/2))\mu([u_0]^N \in G_d(K/2)).$$

For all N , $\mu([u_0]_N \in G_d(K/2)) > 0$ because the hypothesis (1-6) is satisfied and thanks to **Proposition 2.1** we have

$$\mu([u_0]^N \in G_d(K/2)) \geq 1 - C e^{-cK^2/\|[u_0]^N\|_{L^2}^2} \rightarrow 1 \quad \text{as } N \rightarrow \infty,$$

and there exists N such that $\mu([u_0]^N \in G_d(K/2)) > 0$. □

3B. Almost sure local well-posedness of the time dependent equation and scattering for NLS. This section is devoted to the proof of **Theorem 1.4**. The strategy is similar to the proof of **Theorem 1.3**: we solve the equation which is mapped by \mathcal{L} to (1-5) up to time $T = \pi/4$ and we conclude as previously. The difference here is that the nonlinear term of the equation we have to solve is singular at time $T = \pi/4$. More precisely, we consider the equation

$$\begin{cases} i \frac{\partial u}{\partial t} - Hu = \pm \cos(2t)^{p-3} |u|^{p-1} u, & (t, x) \in \mathbb{R} \times \mathbb{R}^2, \\ u(0) = u_0, \end{cases} \tag{3-9}$$

when $2 < p < 3$.

Let us first consider the easier case $\sigma > 0$.

3B1. Proof of Theorem 1.4 in the case $\sigma > 0$. Let $\sigma > 0$ and $\mu = \mu_\gamma \in \mathcal{M}^\sigma$, and for $K \geq 0$ and $\varepsilon > 0$ define the set $F_\sigma(K)$ as

$$F_\sigma(K) = \{w \in \mathcal{H}^\sigma(\mathbb{R}^2) \mid \|w\|_{\mathcal{H}^\sigma(\mathbb{R}^2)} \leq K \text{ and } \|e^{-itH} w\|_{L_{[0,2\pi]}^{1/\varepsilon} \mathcal{W}^{1+\sigma-\varepsilon,\infty}(\mathbb{R}^2)} \leq K\}.$$

The parameter $\varepsilon > 0$ will be chosen small enough that we can apply **Proposition 2.1** and get

$$\mu(F_\sigma(K)^c) \leq \mu(\|w\|_{\mathcal{H}^\sigma} > K) + \mu(\|e^{-itH} w\|_{L_{[0,2\pi]}^{1/\varepsilon} \mathcal{W}^{1+\sigma-\varepsilon,\infty}} > K) \leq C e^{-cK^2/\|\gamma\|_{\mathcal{H}^\sigma}^2}.$$

The next proposition is the key in the proof of **Theorem 1.4** when $\sigma > 0$.

Proposition 3.4. *Let $\sigma > 0$. There exist $C > 0$ and $\kappa > 0$ such that if $u_0 \in F_\sigma(K)$ for some $K > 0$ then for any $v, v_1, v_2 \in X_T^1$ and $0 < T \leq 1$,*

$$\left\| \int_0^t e^{-i(t-\tau)H} (\cos(2\tau)^{p-3} |e^{-i\tau H} u_0 + v|^{p-1} (e^{-i\tau H} u_0 + v)) d\tau \right\|_{X_T^1} \leq CT^\kappa (K^p + \|v\|_{X_T^1}^p) \tag{3-10}$$

and

$$\begin{aligned} & \left\| \int_0^t e^{-i(t-\tau)H} (\cos(2\tau)^{p-3} |e^{-i\tau H} u_0 + v_1|^{p-1} (e^{-i\tau H} u_0 + v_1)) d\tau \right. \\ & \quad \left. - \int_0^t e^{-i(t-\tau)H} (\cos(2\tau)^{p-3} |e^{-i\tau H} u_0 + v_2|^{p-1} (e^{-i\tau H} u_0 + v_2)) d\tau \right\|_{X_T^1} \\ & \leq CT^\kappa \|v_1 - v_2\|_{X_T^1} (K^{p-1} + \|v_1\|_{X_T^1}^{p-1} + \|v_2\|_{X_T^1}^{p-1}). \end{aligned} \quad (3-11)$$

Proof. We first prove (3-10). Using the Strichartz inequalities (3-3), we obtain

$$\begin{aligned} & \left\| \int_0^t e^{-i(t-\tau)H} (\cos(2\tau)^{p-3} |e^{-i\tau H} u_0 + v|^{p-1} (e^{-i\tau H} u_0 + v)) d\tau \right\|_{X_T^1} \\ & \leq C \left\| \cos(2\tau)^{p-3} |e^{-i\tau H} u_0 + v|^{p-1} (e^{-i\tau H} u_0 + v) \right\|_{L^1_{[-T,T]} \mathfrak{H}^1(\mathbb{R}^2)}. \end{aligned}$$

We use the formula

$$\nabla(|u|^{p-1}u) = \frac{p+1}{2}|u|^{p-1}\nabla u + \frac{p-1}{2}|u|^{p-3}u^2\nabla\bar{u}. \quad (3-12)$$

We let $f = e^{-isH}u_0$, then

$$\begin{aligned} \left\| \nabla(|f+v|^{p-1}(f+v)) \right\|_{L^2(\mathbb{R}^2)} & \leq C \left\| |f+v|^{p-1}\nabla(f+v) \right\|_{L^2(\mathbb{R}^2)} + C \left\| |f+v|^{p-3}(f+v)^2\nabla(f+v) \right\|_{L^2(\mathbb{R}^2)} \\ & \leq C \left\| |f+v|^{p-1} \right\|_{L^\infty(\mathbb{R}^2)} \left\| \nabla v \right\|_{L^2(\mathbb{R}^2)} + C \left\| |f+v|^{p-1} \right\|_{L^{2(p-1)}(\mathbb{R}^2)} \left\| \nabla f+v \right\|_{L^\infty(\mathbb{R}^2)}. \end{aligned}$$

Therefore

$$\begin{aligned} & \left\| |f+v|^{p-1}(f+v) \right\|_{\mathfrak{H}^1(\mathbb{R}^2)} \\ & \leq C (\left\| |f|^{p-1} \right\|_{L^\infty(\mathbb{R}^2)} + \left\| |v|^{p-1} \right\|_{L^\infty(\mathbb{R}^2)}) \left\| v \right\|_{\mathfrak{H}^1(\mathbb{R}^2)} + C (\left\| |f|^{p-1} \right\|_{L^{2(p-1)}(\mathbb{R}^2)} + \left\| |v|^{p-1} \right\|_{L^{2(p-1)}(\mathbb{R}^2)}) \left\| f \right\|_{W^{1,\infty}(\mathbb{R}^2)}. \end{aligned}$$

Now observe that $\|v\|_{L^{\infty}_{[-T,T]}L^{2(p-1)}} \leq \|v\|_{X_T^1}$ as well as, for all $r < +\infty$, $\|v\|_{L^r_{[-T,T]}L^\infty} \leq \|v\|_{X_T^1}$.

Then, for all $q > 1$,

$$\begin{aligned} \left\| |f+v|^{p-1}(f+v) \right\|_{L^q_{[-T,T]} \mathfrak{H}^1(\mathbb{R}^2)} & \leq CT^\kappa \left(\left(\left\| |f|^{p-1} \right\|_{L^\infty_{[-T,T]}L^\infty(\mathbb{R}^2)} + \left\| |v|^{p-1} \right\|_{X_T^1} \right) \left\| v \right\|_{X_T^1} \right. \\ & \quad \left. + \left(\left\| |f|^{p-1} \right\|_{L^\infty_{[-T,T]}L^{2(p-1)}(\mathbb{R}^2)} + \left\| |v|^{p-1} \right\|_{X_T^1} \right) \left\| f \right\|_{L^\infty_{[-T,T]}W^{1,\infty}(\mathbb{R}^2)} \right) \\ & \leq CT^\kappa (K^p + \|v\|_{X_T^1}^p). \end{aligned} \quad (3-13)$$

Choose $q > 1$ so that $q'(3-p) < 1$. We have $\|\cos(2\tau)^{p-3}\|_{L^{q'}_{[-T,T]}} < \infty$; thus from (3-13) and Hölder, we infer

$$\begin{aligned} & \left\| \cos(2\tau)^{p-3} |f+v|^{p-1}(f+v) \right\|_{L^1_{[-T,T]} \mathfrak{H}^1(\mathbb{R}^2)} \\ & \leq C \left\| \cos(2\tau)^{p-3} \right\|_{L^{q'}_{[-T,T]}} \left\| |f+v|^{p-1}(f+v) \right\|_{L^q_{[-T,T]} \mathfrak{H}^1(\mathbb{R}^2)} \\ & \leq CT^\kappa (K^p + \|v\|_{X_T^1}^p). \end{aligned}$$

For the proof of (3-11) we can proceed similarly. Namely, we use the estimates

$$||z_1|^{p-1} - |z_2|^{p-1}| \leq C(|z_1|^{p-2} + |z_2|^{p-2})|z_1 - z_2| \tag{3-14}$$

and

$$||z_1|^{p-3}z_1^2 - |z_2|^{p-3}z_2^2| \leq C(|z_1|^{p-2} + |z_2|^{p-2})|z_1 - z_2|,$$

which are proven in [Cazenave et al. 2011, Remark 2.3] together with (3-12). □

3B2. *Proof of Theorem 1.4 in the case $\sigma = 0$.* The strategy of the proof in this case is similar, at the price of some technicalities, since the Leibniz rule (3-12) does not hold true for non-integer derivatives. Actually, when $\sigma = 0$, we will have to work in X_T^s for $s < 1$ because the probabilistic term $e^{-itH}u_0 \notin \mathcal{W}^{1,\infty}(\mathbb{R}^2)$.

Moreover, we are not able to obtain a contraction estimate in X_T^s . Therefore, we will do a fixed point in the space $\{\|v\|_{X_T^s} \leq K\}$ endowed with the weaker metric induced by X_T^0 . We can check that this space is complete. Actually, by the Banach–Alaoglu theorem, the closed balls of each component space of X_T^s are compact for the weak* topology.

For $0 < s < 1$, we use the following characterisation of the usual $H^s(\mathbb{R}^2)$ norm:

$$\|g\|_{H^s(\mathbb{R}^2)} = \left(\int_{\mathbb{R}^2 \times \mathbb{R}^2} \frac{|g(x) - g(y)|^2}{|x - y|^{2s+2}} dx dy \right)^{1/2}. \tag{3-15}$$

For $\mu = \mu_\gamma \in \mathcal{M}^0$, $K \geq 0$ and $\varepsilon > 0$, define the set $\tilde{F}_0(K)$ as

$$\begin{aligned} \tilde{F}_0(K) = \{w \in L^2(\mathbb{R}^2) \mid & \|w\|_{L^2(\mathbb{R}^2)} \leq K, \|e^{-itH}w\|_{L_{[0,2\pi]}^{1/\varepsilon} \mathcal{W}^{1-\varepsilon,\infty}(\mathbb{R}^2)} \leq K \\ & \text{and } \|(e^{-itH}w)(x) - (e^{-itH}w)(y)\|_{L_{t \in [0,2\pi]}^\infty} \leq K|x - y|^{1-\varepsilon}\}. \end{aligned}$$

The next result states that $\tilde{F}_0(K)$ is a set with large measure.

Lemma 3.5. *If $\varepsilon > 0$ is small enough,*

$$\mu((\tilde{F}_0(K))^c) \leq C e^{-cK^2/\|\gamma\|_{L^2(\mathbb{R}^2)}^2}.$$

Proof. We only have to study the contribution of the Lipschitz term in $\tilde{F}_0(K)$, since the others are controlled by Proposition 2.1.

We fix $\gamma = \sum_{n=0}^{+\infty} c_n \varphi_n \in \mathcal{A}_0$ and set $\gamma^\omega = \sum_{n=0}^{+\infty} g_n(\omega) c_n \varphi_n$. Let $k \geq 1$. By definition,

$$\begin{aligned} \int_{L^2(\mathbb{R}^2)} \|e^{-itH}u(x) - e^{-itH}u(y)\|_{L_{[0,2\pi]}^\infty}^k d\mu(u) \\ = \int_{\Omega} \|e^{-itH}\gamma^\omega(x) - e^{-itH}\gamma^\omega(y)\|_{L_{[0,2\pi]}^\infty}^k d\mathbb{P}(\omega). \end{aligned} \tag{3-16}$$

We have $e^{-itH}\gamma^\omega(x) - e^{-itH}\gamma^\omega(y) = \sum_{n=0}^{+\infty} g_n(\omega)c_n e^{-it\lambda_n}(\varphi_n(x) - \varphi_n(y))$. By Khinchin (Lemma 2.3) we get

$$\begin{aligned} \|e^{-itH}\gamma^\omega(x) - e^{-itH}\gamma^\omega(y)\|_{L^k_{\mathbb{P}}} &\leq C\sqrt{k} \left(\sum_{n=0}^{+\infty} |c_n|^2 |\varphi_n(x) - \varphi_n(y)|^2 \right)^{1/2} \\ &= C\sqrt{k} \left(\sum_{j=1}^{+\infty} \sum_{n \in I(j)} |c_n|^2 |\varphi_n(x) - \varphi_n(y)|^2 \right)^{1/2}, \end{aligned}$$

Recall that $k \in I(j) = \{n \in \mathbb{N} \mid 2j \leq \lambda_n < 2(j+1)\}$ and that $\#I(j) \sim cj$. Next, by condition (1-3), we deduce that

$$\|e^{-itH}\gamma^\omega(x) - e^{-itH}\gamma^\omega(y)\|_{L^k_{\mathbb{P}}} \leq C\sqrt{k} \left(\sum_{j=1}^{+\infty} j^{-1} \left(\sum_{\ell \in I(j)} |c_\ell|^2 \right) \sum_{n \in I(j)} |\varphi_n(x) - \varphi_n(y)|^2 \right)^{1/2}$$

Now we need the following estimate, proven in [Imekraz et al. 2014, Lemma 6.1]:

$$\sum_{n \in I(j)} |\varphi_n(y) - \varphi_n(x)|^2 \leq C|y-x|^2 j.$$

Therefore, we obtain

$$\|e^{-itH}\gamma^\omega(x) - e^{-itH}\gamma^\omega(y)\|_{L^k_{\mathbb{P}}} \leq C\sqrt{k}|x-y|\|\gamma\|_{L^2(\mathbb{R}^2)},$$

and for $k \geq q$ an integration in time and Minkowski yield

$$\|e^{-itH}\gamma^\omega(x) - e^{-itH}\gamma^\omega(y)\|_{L^k_{\mathbb{P}}L^q_{[0,2\pi]}} \leq C\sqrt{k}|x-y|\|\gamma\|_{L^2(\mathbb{R}^2)}.$$

However, since the case $q = +\infty$ is forbidden, the previous estimate is not enough to have a control on the $L^\infty_{[0,2\pi]}$ -norm. To tackle this issue, we claim that for $k \geq q$ we have

$$\|e^{-itH}\gamma^\omega(x) - e^{-itH}\gamma^\omega(y)\|_{L^k_{\mathbb{P}}W^{1,q}_{[0,2\pi]}} \leq C\sqrt{k}\|\gamma\|_{L^2(\mathbb{R}^2)}. \tag{3-17}$$

Then by a usual Sobolev embedding argument we get (by taking $q \gg 1$ large enough) that for all $\varepsilon > 0$

$$\|e^{-itH}\gamma^\omega(x) - e^{-itH}\gamma^\omega(y)\|_{L^k_{\mathbb{P}}L^\infty_{[0,2\pi]}} \leq C\sqrt{k}|x-y|^{1-\varepsilon}\|\gamma\|_{L^2(\mathbb{R}^2)},$$

which in turn by (3-16) implies that

$$\mu(u \in L^2(\mathbb{R}^2) \mid \|e^{-itH}u(x) - e^{-itH}u(y)\|_{L^\infty_{[0,2\pi]}} > K|x-y|^{1-\varepsilon}) \leq Ce^{-cK^2/\|\gamma\|_{L^2}^2},$$

as we did in the end of the proof of Proposition 2.1.

Let us now prove (3-17). We have

$$\partial_t(e^{-itH}\gamma^\omega(x) - e^{-itH}\gamma^\omega(y)) = -i \sum_{n=0}^{+\infty} g_n(\omega)\lambda_n c_n e^{-it\lambda_n}(\varphi_n(x) - \varphi_n(y)),$$

and with the previous arguments we get

$$\begin{aligned} \|\partial_t(e^{-itH}\gamma^\omega(x) - e^{-itH}\gamma^\omega(y))\|_{L^k_{\mathbb{P}}} &\leq C\sqrt{k}\left(\sum_{j=1}^{+\infty}\left(\sum_{\ell\in I(j)}|c_\ell|^2\right)\sum_{n\in I(j)}|\varphi_n(x) - \varphi_n(y)|^2\right)^{1/2} \\ &\leq C\sqrt{k}\|\gamma\|_{L^2(\mathbb{R}^2)}, \end{aligned}$$

where here we have used the Thangavelu–Karadzhov estimate (see [Poiret et al. 2013, Lemma 3.5])

$$\sup_{x\in\mathbb{R}^2}\sum_{n\in I(j)}|\varphi_n(x)|^2 \leq C.$$

We conclude the proof of (3-17) by integrating in time and using Minkowski. □

We will also need the following technical result.

Lemma 3.6. *Let $u_0 \in \tilde{F}_0(K)$ and $f(t, x) = e^{-itH}u_0(x)$. Let $2 \leq q < +\infty$ and $g \in L^q([-T, T]; L^2(\mathbb{R}^2))$. Then, if $\varepsilon > 0$ is small enough in the definition of $\tilde{F}_0(K)$,*

$$\left\|\left(\int_{\mathbb{R}^2 \times \mathbb{R}^2} \frac{|f(t, x) - f(t, y)|^2 |g(t, x)|^2}{|x - y|^{2s+2}} dx dy\right)^{1/2}\right\|_{L^q_{[-T, T]}} \leq CK\|g\|_{L^q_{[-T, T]}L^2(\mathbb{R}^2)}. \quad (3-18)$$

Proof. We consider such f, g , and we split the integral. On the one hand, we use that f is Lipschitz:

$$\begin{aligned} \int_{|x-y|\leq 1} \frac{|f(t, x) - f(t, y)|^2 |g(t, x)|^2}{|x - y|^{2s+2}} dx dy &\leq K^2 \int_{x\in\mathbb{R}^2} |g(t, x)|^2 \left(\int_{y:|x-y|\leq 1} \frac{dy}{|x - y|^{2s+2\varepsilon}}\right) dx \\ &\leq CK^2\|g(t, \cdot)\|_{L^2(\mathbb{R}^2)}^2, \end{aligned}$$

provided that $s + \varepsilon < 1$. We take the $L^q_{[-T, T]}$ -norm, and we see that this contribution is bounded by the right side of (3-18).

On the other hand

$$\begin{aligned} \int_{|x-y|\geq 1} \frac{|f(t, x) - f(t, y)|^2 |g(t, x)|^2}{|x - y|^{2s+2}} dx dy &\leq C\|f(t, \cdot)\|_{L^\infty(\mathbb{R}^2)}^2 \int_{x\in\mathbb{R}^2} |g(t, x)|^2 \left(\int_{y:|x-y|\geq 1} \frac{dy}{|x - y|^{2s+2}}\right) dx \\ &\leq C\|f(t, \cdot)\|_{L^\infty(\mathbb{R}^2)}^2 \|g(t, \cdot)\|_{L^2(\mathbb{R}^2)}^2 \end{aligned}$$

if $s > 0$. Now we take the $L^q_{[-T, T]}$ -norm, and use the fact that $\|f\|_{L^q_{[0, 2\pi]}L^\infty(\mathbb{R}^2)} \leq K$ if $\varepsilon < 1/q$. □

We now state the main estimates of this section.

Proposition 3.7. *There exist $C > 0$ and $\kappa > 0$ such that if $u_0 \in \tilde{F}_0(K)$ for some $K > 0$ then for any $v, v_1, v_2 \in X^s_T$ and $0 < T \leq 1$,*

$$\left\|\int_0^t e^{-i(t-\tau)H}(\cos(2\tau)^{p-3}|e^{-i\tau H}u_0 + v|^{p-1}(e^{-i\tau H}u_0 + v))d\tau\right\|_{X^s_T} \leq CT^\kappa(K^p + \|v\|_{X^s_T}^p), \quad (3-19)$$

and

$$\begin{aligned} & \left\| \int_0^t e^{-i(t-\tau)H} (\cos(2\tau)^{p-3} |e^{-i\tau H} u_0 + v_1|^{p-1} (e^{-i\tau H} u_0 + v_1)) d\tau \right. \\ & \quad \left. - \int_0^t e^{-i(t-\tau)H} (\cos(2\tau)^{p-3} |e^{-i\tau H} u_0 + v_2|^{p-1} (e^{-i\tau H} u_0 + v_2)) d\tau \right\|_{X_T^0} \\ & \leq CT^\kappa \|v_1 - v_2\|_{X_T^0} (K^{p-1} + \|v_1\|_{X_T^s}^{p-1} + \|v_2\|_{X_T^s}^{p-1}). \end{aligned} \tag{3-20}$$

Proof. Let $u_0 \in \tilde{F}_0(K)$ and set $f = e^{-isH} u_0$. Let $2 < p < 3$, then there exists $q \gg 1$ such that $q'(3-p) < 1$, which in turn implies $\|\cos(2s)^{p-3}\|_{L_{[-T,T]}^{q'}}$ $\leq CT^\kappa$. Next, if $s < 1$ is large enough we have, by Sobolev,

$$\|v\|_{L_{[-T,T]}^\infty L^{2(p-1)}(\mathbb{R}^2)} \leq \|v\|_{X_T^s} \quad \text{and} \quad \|v\|_{L_{[-T,T]}^{q(p-1)} L^\infty(\mathbb{R}^2)} \leq \|v\|_{X_T^s}. \tag{3-21}$$

First we prove (3-19). From Strichartz and Hölder, we get

$$\begin{aligned} & \left\| \int_0^t e^{-i(t-s)H} (\cos(2s)^{p-3} |f + v|^{p-1} (f + v)) ds \right\|_{X_T^s} \\ & \leq C \|\cos(2s)^{p-3} |f + v|^{p-1} (f + v)\|_{L_{[-T,T]}^1 \mathcal{H}^s(\mathbb{R}^2)} \\ & \leq C \|\cos(2s)^{p-3}\|_{L_{[-T,T]}^{q'}} \| |f + v|^{p-1} (f + v) \|_{L_{[-T,T]}^q \mathcal{H}^s(\mathbb{R}^2)} \\ & \leq CT^\kappa \| |f + v|^{p-1} (f + v) \|_{L_{[-T,T]}^q \mathcal{H}^s(\mathbb{R}^2)}. \end{aligned} \tag{3-22}$$

By using the characterization (3-15), we will prove that

$$\| |f + v|^{p-1} (f + v) \|_{L_{[-T,T]}^q \mathcal{H}^s(\mathbb{R}^2)} \leq C (K^p + \|v\|_{X_T^s}^p). \tag{3-23}$$

The term $\| \langle x \rangle^s |f + v|^{p-1} (f + v) \|_{L_{[-T,T]}^q L^2(\mathbb{R}^2)}$ is easily controlled; thus we only detail the contribution of the H^s norm. With (3-14), it is easy to check that, for all $x, y \in \mathbb{R}^2$,

$$\begin{aligned} & \left| |f + v|^{p-1} (f + v)(x) - |f + v|^{p-1} (f + v)(y) \right| \\ & \leq C |v(x) - v(y)| (|v(x)|^{p-1} + |v(y)|^{p-1} + |f(x)|^{p-1} + |f(y)|^{p-1}) \\ & \quad + C |f(x) - f(y)| (|v(x)|^{p-1} + |v(y)|^{p-1} + |f(x)|^{p-1} + |f(y)|^{p-1}). \end{aligned}$$

By (3-21) the contribution in $L_{[-T,T]}^q H^s(\mathbb{R}^2)$ of the first term in the previous expression is at most

$$C \left(\|f\|_{L_{[-T,T]}^{q(p-1)} L^\infty(\mathbb{R}^2)}^{p-1} + \|v\|_{L_{[-T,T]}^{q(p-1)} L^\infty(\mathbb{R}^2)}^{p-1} \right) \|v\|_{X_T^s} \leq C (K^{p-1} + \|v\|_{X_T^s}^{p-1}) \|v\|_{X_T^s}.$$

To bound the second term, we apply Lemma 3.6, which gives a contribution of at most

$$\left(\|f\|_{L_{[-T,T]}^{q(p-1)} L^{2(p-1)}(\mathbb{R}^2)}^{p-1} + \|v\|_{L_{[-T,T]}^{q(p-1)} L^{2(p-1)}(\mathbb{R}^2)}^{p-1} \right) K \leq C (K^{p-1} + \|v\|_{X_T^s}^{p-1}) K,$$

which concludes the proof of (3-23).

The proof of (3-20) is in the same spirit, and even easier. We do not write the details. □

Thanks to the estimates of [Proposition 3.7](#), for $K > 0$ small enough (see the proof of [Theorem 1.3](#) for more details) we are able to construct a unique solution $v \in \mathcal{C}([-\pi/4, \pi/4]; L^2(\mathbb{R}^2))$ such that $v \in L^\infty([-\pi/4, \pi/4]; \mathcal{H}^s(\mathbb{R}^2))$. By interpolation we deduce that $v \in \mathcal{C}([-\pi/4, \pi/4]; \mathcal{H}^{s'}(\mathbb{R}^2))$ for all $s' < s$. The end of the proof of [Theorem 1.4](#) is similar to the proof of [Theorem 1.3](#), using here [Lemma 3.5](#).

4. Global well-posedness for the cubic equation

4A. The case of dimension $d = 3$. We now turn to the proof of [Theorem 1.5](#), which is obtained thanks to the high-low frequency decomposition method of [\[Bourgain 1999, p. 84\]](#).

Let $0 \leq s < 1$ and fix $\mu = \mu_\gamma \in \mathcal{M}^s$. For $K \geq 0$ define the set $F_s(K)$ as

$$F_s(K) = \{w \in \mathcal{H}^s(\mathbb{R}^3) \mid \|w\|_{\mathcal{H}^s(\mathbb{R}^3)} \leq K, \|w\|_{L^4(\mathbb{R}^3)} \leq K \text{ and } \|e^{-itH}w\|_{L_{[0,2\pi]}^{1/\varepsilon} W^{3/2+s-\varepsilon,\infty}(\mathbb{R}^3)} \leq K\}.$$

Then, by [Proposition 2.1](#),

$$\begin{aligned} &\mu((F_s(K))^c) \\ &\leq \mu(\|w\|_{\mathcal{H}^s} > K) + \mu(\|w\|_{L^4} > K) + \mu(\|e^{-itH}w\|_{L_{[0,2\pi]}^{1/\varepsilon} W^{3/2+s-\varepsilon,\infty}} > K) \leq C e^{-cK^2/\|\gamma\|_{\mathcal{H}^s}^2}. \end{aligned} \quad (4-1)$$

Now we define a smooth version of the usual spectral projector. Let $\chi \in \mathcal{C}_0^\infty(-1, 1)$, so that $0 \leq \chi \leq 1$, with $\chi = 1$ on $[-\frac{1}{2}, \frac{1}{2}]$. We define the operators $S_N = \chi(\frac{H}{N^2})$ as

$$S_N \left(\sum_{n=0}^{+\infty} c_n \varphi_n \right) = \sum_{n=0}^{+\infty} \chi \left(\frac{\lambda_n}{N^2} \right) c_n \varphi_n,$$

and we write

$$v_N = S_N v, \quad v^N = (1 - S_N)v.$$

It is clear that for any $\sigma \geq 0$ we have $\|S_N\|_{\mathcal{H}^\sigma \rightarrow \mathcal{H}^\sigma} = 1$. Moreover, by [\[Burq et al. 2010, Proposition 4.1\]](#), for all $1 \leq r \leq +\infty$, $\|S_N\|_{L^r \rightarrow L^r} \leq C$, uniformly in $N \geq 1$.

It is straightforward to check that

$$\|v_N\|_{\mathcal{H}^1} \leq N^{1-s} \|v\|_{\mathcal{H}^s}, \quad \|v^N\|_{L^2} \leq N^{-s} \|v\|_{\mathcal{H}^s}. \quad (4-2)$$

Next, let $u_0 \in F_s(N^\varepsilon)$. By the definition of $F_s(N^\varepsilon)$ and [\(4-2\)](#), $\|u_{0,N}\|_{\mathcal{H}^1} \leq N^{1-s} \|u_0\|_{\mathcal{H}^s} \leq N^{1-s+\varepsilon}$. The nonlinear term of the energy can be controlled by the quadratic term. Indeed

$$\|u_{0,N}\|_{L^4}^4 \leq CN^\varepsilon \leq N^{2(1-s+\varepsilon)},$$

and thus

$$E(u_{0,N}) \leq 2N^{2(1-s+\varepsilon)}. \quad (4-3)$$

We also have

$$\|u_{0,N}\|_{L^2} \leq \|u_0\|_{\mathcal{H}^s} \leq N^\varepsilon.$$

For a nice description of the stochastic version of the low-high frequency decomposition method we use here, we refer to the introduction of [\[Colliander and Oh 2012\]](#). To begin with, we look for a solution u to [\(1-7\)](#) of the form $u = u^1 + v^1$, where u^1 is the solution to

$$\begin{cases} i \frac{\partial u^1}{\partial t} - H u^1 = |u^1|^2 u^1, & (t, x) \in \mathbb{R} \times \mathbb{R}^3, \\ u^1(0) = u_{0,N}, \end{cases} \tag{4-4}$$

and where $v^1 = e^{-itH} u_0^N + w^1$ satisfies

$$\begin{cases} i \frac{\partial w^1}{\partial t} - H w^1 = |w^1 + e^{-itH} u_0^N + u^1|^2 (w^1 + e^{-itH} u_0^N + u^1) - |u^1|^2 u^1, & (t, x) \in \mathbb{R} \times \mathbb{R}^3, \\ w^1(0) = 0. \end{cases} \tag{4-5}$$

Since (4-4) is \mathcal{H}^1 -subcritical, by the usual deterministic arguments there exists a unique global solution $u^1 \in \mathcal{C}(\mathbb{R}, \mathcal{H}^1(\mathbb{R}^3))$.

We now turn to (4-5), for which we have the next local existence result.

Proposition 4.1. *Let $0 < s < 1$ and $\mu = \mu_\gamma \in \mathcal{M}^s$. Set $T = N^{-4(1-s)-\varepsilon}$ with $\varepsilon > 0$. Assume that $E(u^1) \leq 4N^{2(1-s+\varepsilon)}$ and $\|u^1\|_{L^\infty_{[0,T]}L^2} \leq 2N^\varepsilon$. Then:*

(i) *There exists a set $\Sigma_T^1 \subset \mathcal{H}^s$, which only depends on T , so that*

$$\mu(\Sigma_T^1) \geq 1 - C \exp(-cT^{-\delta} \|\gamma\|_{\mathcal{H}^s(\mathbb{R}^3)}^{-2}), \quad \text{with some } \delta > 0.$$

(ii) *For all $u_0 \in \Sigma_T^1$ there exists a unique solution $w^1 \in \mathcal{C}([0, T], \mathcal{H}^1(\mathbb{R}^3))$ to (4-5), which satisfies the bounds*

$$\|w^1\|_{L^\infty_{[0,T]}\mathcal{H}^1} \leq CN^{\beta(s)+c\varepsilon}, \tag{4-6}$$

with

$$\beta(s) = \begin{cases} -\frac{5}{2} & \text{if } 0 \leq s \leq \frac{1}{2}, \\ 2s - \frac{7}{2} & \text{if } \frac{1}{2} \leq s \leq 1, \end{cases} \tag{4-7}$$

and

$$\|w^1\|_{L^\infty_{[0,T]}L^2} \leq CN^{-9/2+2s+c\varepsilon}. \tag{4-8}$$

Proof. In the next lines, we write $C^{a+} = C^{a+b\varepsilon}$, for some absolute quantity $b > 0$. Since $d = 3$, for $T > 0$ we define the space $X_T^1 = L^\infty([0, T]; \mathcal{H}^1(\mathbb{R}^3)) \cap L^2([0, T]; \mathcal{W}^{1,6}(\mathbb{R}^3))$. Let $\varepsilon > 0$, and define $\Sigma_T^1 = F_s(N^\varepsilon)$. By (4-1) and the choice $T = N^{-4(1-s)-\varepsilon}$, the set Σ_T^1 satisfies (i).

Let $u_0 \in \Sigma_T^1$. To simplify the notations in the proof, we write $w = w^1$, $u = u^1$ and $f = e^{-itH} u_0^N$. We define the map

$$L(w) = \mp i \int_0^t e^{-i(t-s)H} (|f + u + w|^2 (f + u + w) - |u|^2 u)(s) ds. \tag{4-9}$$

First we prove (4-6). By Strichartz (3-3),

$$\|L(w)\|_{X_T^1} \leq C \| |f + u + w|^2 (f + u + w) - |u|^2 u \|_{L_T^1 \mathcal{H}^1 + L^2 \mathcal{W}^{1,6/5}}. \tag{4-10}$$

By estimating the contribution of every term, we now prove that

$$\|L(w)\|_{X_T^1} \leq CN^{\beta(s)+} + N^{0-} \|w\|_{X_T^1} + N^{-2(1-s)+} \|w\|_{X_T^1}^3, \tag{4-11}$$

where $\beta(s) < (1 - s)$ is as in the statement. It is enough to prove that L maps a ball of size $CN^{\beta(s)+}$ into itself for times $T = N^{-4(1-s)-\varepsilon}$. With similar arguments one can show that L is a contraction (we do not write the details) and get w satisfying (4-6).

Observe that the complex conjugation is harmless with respect to the norms considered; thus we can forget it. By the definition of $\Sigma_T^1 = F_s(N^\varepsilon)$ and (4-2) we have the estimates used in the sequel: for all $\sigma < \frac{3}{2}$,

$$\|f\|_{L_T^\infty L^2} \leq CN^{-s+\varepsilon} \quad \text{and} \quad \|H^{\sigma/2} f\|_{L_T^{\infty-} L^\infty} \leq CN^{\sigma-3/2-s+2\varepsilon}. \quad (4-12)$$

Let us prove the second estimate in detail:

$$\begin{aligned} \|H^{\sigma/2} f\|_{L_T^{\infty-} L^\infty} &= N^\sigma \left\| \left(\frac{H}{N^2} \right)^{\sigma/2} \left(1 - \chi \left(\frac{H}{N^2} \right) \right) e^{-itH} u_0 \right\|_{L_T^{\infty-} L^\infty} \\ &\leq CN^\sigma \left\| \left(\frac{H}{N^2} \right)^{(3/2+s-\varepsilon)/2} \left(1 - \chi \left(\frac{H}{N^2} \right) \right) e^{-itH} u_0 \right\|_{L_T^{\infty-} L^\infty} \\ &\leq CN^{\sigma-3/2-s+\varepsilon} \|e^{-itH} u_0\|_{L_T^{\infty-\varrho} \mathcal{W}^{3/2+s-\varepsilon, \infty}} \\ &\leq CN^{\sigma-3/2-s+2\varepsilon}, \end{aligned}$$

where we have used that $x^{\sigma/2}(1 - \chi(x)) \leq Cx^{(3/2+s-\varepsilon)/2}(1 - \chi(x))$.

Observe also that by assumption

$$\|u\|_{L_T^\infty L^2} \leq CN^\varepsilon, \quad \|u\|_{L_T^\infty \mathcal{H}^1} \leq CN^{1-s+\varepsilon}, \quad \text{and} \quad \|u\|_{L_T^\infty L^4} \leq CN^{(1-s+\varepsilon)/2}.$$

We now estimate each term in the right side of (4-10):

• Source terms: Observe that $L_T^{4/3} \mathcal{W}^{1,3/2} \subset L_T^1 \mathcal{H}^1 + L^{2\varrho} \mathcal{W}^{1,6/5}$. By Hölder and (4-12),

$$\begin{aligned} \|f u^2\|_{L_T^1 \mathcal{H}^1 + L^{2\varrho} \mathcal{W}^{1,6/5}} &\leq C \|f u H^{1/2} u\|_{L_T^{4/3} L^{3/2}} + C \|u^2 H^{1/2} f\|_{L_T^1 L^2} \\ &\leq CT^{3/4-} \|u\|_{L_T^\infty \mathcal{H}^1} \|u\|_{L_T^\infty L^6} \|f\|_{L_T^{\infty-} L^\infty} + CT^{1-} \|u\|_{L_T^\infty L^4}^2 \|H^{1/2} f\|_{L_T^{\infty-} L^\infty} \\ &\leq CN^{-5/2+} + CN^{-7/2+2s+} \leq CN^{\beta(s)+}, \end{aligned}$$

where we have set $\beta(s) = \max(-\frac{5}{2}, -\frac{7}{2} + 2s)$, which is precisely (4-7). Similarly,

$$\begin{aligned} \|f^2 u\|_{L_T^1 \mathcal{H}^1} &\leq C \|f^2 H^{1/2} u\|_{L_T^1 L^2} + C \|u f H^{1/2} f\|_{L_T^1 L^2} \\ &\leq CT^{1-} \|u\|_{L_T^\infty \mathcal{H}^1} \|f\|_{L_T^{\infty-} L^\infty}^2 + CT^{1-} \|u\|_{L_T^\infty L^2} \|f\|_{L_T^{\infty-} L^\infty} \|H^{1/2} f\|_{L_T^{\infty-} L^\infty} \\ &\leq CT^{1-} N^{-2-3s+} + CT^{1-} N^{-2-2s+} \leq CN^{-6+2s+} \leq CN^{\beta(s)+}. \end{aligned}$$

Finally,

$$\begin{aligned} \|f^3\|_{L_T^1 \mathcal{H}^1} &\leq C \|f^2 H^{1/2} f\|_{L_T^1 L^2} \leq CT^{1-} \|H^{1/2} f\|_{L_T^{\infty-} L^\infty} \|f\|_{L_T^{\infty-} L^\infty} \|f\|_{L_T^\infty L^2} \\ &\leq CT^{1-} N^{-1/2-s+} N^{-3/2-s+} N^{-s+} \leq CN^{-6+s+} \leq CN^{\beta(s)+}. \end{aligned}$$

• Linear terms in w :

$$\begin{aligned} \|wf^2\|_{L_T^1 \mathcal{H}^1} &\leq C \|f^2 H^{1/2} w\|_{L_T^1 L^2} + C \|wf H^{1/2} f\|_{L_T^1 L^2} \\ &\leq CT^{1-} \|f\|_{L_T^\infty L^\infty}^2 \|w\|_{L_T^\infty \mathcal{H}^1} + CT^{1-} \|w\|_{L_T^\infty L^2} \|f\|_{L_T^\infty L^\infty} \|H^{1/2} f\|_{L_T^\infty L^\infty} \\ &\leq CN^{-6+2s+} \|w\|_{X_T^1} \leq CN^{0-} \|w\|_{X_T^1}. \end{aligned}$$

Using that $\|w\|_{L_T^{4/3+} L^\infty-} \leq CT^{1/2-} \|w\|_{L_T^4 L^\infty-} \leq CT^{1/2-} \|w\|_{L_T^4 \mathcal{W}^{1,3}}$ and $X_T^1 \subset L^4([0, T]; \mathcal{W}^{1,3})$,

$$\begin{aligned} \|wu^2\|_{L_T^1 \mathcal{H}^1 + L^2 \mathcal{W}^{1,6/5}} &\leq C \|u^2 H^{1/2} w\|_{L_T^1 L^2} + C \|wu H^{1/2} u\|_{L_T^{4/3+} L^{3/2-}} \\ &\leq C \|u\|_{L_T^4 L^6}^2 \|w\|_{L_T^2 \mathcal{W}^{1,6}} + C \|w\|_{L_T^{4/3+} L^\infty-} \|u\|_{L_T^\infty L^6} \|u\|_{L_T^\infty \mathcal{H}^1} \\ &\leq CT^{1/2-} \|u\|_{L_T^\infty \mathcal{H}^1}^2 \|w\|_{X_T^1} \leq CN^{0-} \|w\|_{X_T^1}. \end{aligned}$$

• The cubic term in w : by Sobolev and $X_T^1 \subset L^{4-}([0, T]; \mathcal{W}^{1,3+}) \subset L^{4-}([0, T]; L^\infty)$, we have

$$\begin{aligned} \|w^3\|_{L_T^1 \mathcal{H}^1} &\leq C \|w^2 H^{1/2} w\|_{L_T^1 L^2} \leq C \|w\|_{L_T^\infty \mathcal{H}^1} \|w\|_{L_T^2 L^\infty}^2 \\ &\leq CT^{1/2-} \|w\|_{X_T^1}^3 \leq CN^{-2(1-s)+} \|w\|_{X_T^1}^3. \end{aligned}$$

• Quadratic terms in w : with similar arguments, we check that they are controlled by the previous ones.

This completes the proof of (4-11). Hence, for all $u_0 \in \Sigma_T^1$, L has a unique fixed point w .

Let $w \in X_T^1$ be defined this way, and let us prove that $\|w\|_{X_T^0} \leq CN^{-9/2+2s+}$, which will imply (4-8). By the Strichartz inequality (3-3),

$$\|w\|_{X_T^0} \leq C \| |f+u+w|^2(f+u+w) - |u|^2 u \|_{L_T^1 L^2 + L^2 L^{6/5}}.$$

As previously, the main contribution in the source term is

$$\|fu^2\|_{L_T^1 L^2} \leq T^{1-} \|u\|_{L_T^\infty L^4}^2 \|f\|_{L_T^\infty L^\infty} \leq CN^{-4(1-s)+1-s-3/2-s+} = CN^{-9/2+2s+}.$$

For the cubic term we write

$$\begin{aligned} \|w^3\|_{L_T^1 L^2} &\leq \|w\|_{L_T^\infty L^2} \|w\|_{L_T^2 L^\infty}^2 \leq CT^{1/2-} \|w\|_{L_T^\infty L^2} \|w\|_{X_T^1}^2 \\ &\leq CN^{-2(1-s)+\beta(s)+} \|w\|_{L_T^\infty L^2} \leq CN^{0-} \|w\|_{X_T^0}, \end{aligned}$$

which gives a control by the linear term.

The other terms are controlled with similar arguments, and we leave the details to the reader. This finishes the proof of Proposition 4.1. \square

Lemma 4.2. *Under the assumptions of Proposition 4.1, for all $u_0 \in \Sigma_T^1$ we have*

$$|E(u^1(T) + w^1(T)) - E(u^1(T))| \leq CN^{1-s+\beta(s)+}.$$

Proof. Write $u = u^1$ and $w = w^1$. A direct expansion and Hölder give

$$|E(u(T) + w(T)) - E(u(T))| \leq 2\|u\|_{L_T^\infty \mathcal{H}^1} \|w\|_{L_T^\infty \mathcal{H}^1} + \|w\|_{L_T^\infty \mathcal{H}^1}^2 + C\|w\|_{L_T^\infty L^4} \|u\|_{L_T^\infty L^4}^3 + C\|w\|_{L_T^\infty L^4}^4.$$

Since $\beta(s) \leq (1 - s)$, we directly have

$$2\|u\|_{L_T^\infty \mathcal{H}^1} \|w\|_{L_T^\infty \mathcal{H}^1} + \|w\|_{L_T^\infty \mathcal{H}^1}^2 \leq CN^{1-s+\beta(s)+}.$$

By Sobolev and [Proposition 4.1](#),

$$\|w\|_{L_T^\infty L^4} \leq C\|w\|_{L_T^\infty \mathcal{H}^{3/4}} \leq C\|w\|_{L_T^\infty L^2}^{1/4} \|w\|_{L_T^\infty \mathcal{H}^1}^{3/4} \leq CN^{\eta(s)+}, \tag{4-13}$$

with $\eta(s) = \max(-3 + s/2, -15/4 + 2s) \leq (1 - s + \beta(s))/3$. Hence,

$$\|w\|_{L_T^\infty L^4}^3 \leq CN^{1-s+\beta(s)+}.$$

From the bounds $\|u\|_{L_T^\infty L^4} \leq CN^{(1-s)/2}$ and [\(4-13\)](#), we infer

$$\|w\|_{L_T^\infty L^4} \|u\|_{L_T^\infty L^4}^3 \leq CN^{\delta(s)+},$$

where $\delta(s) = \max(-3 + s/2, -15/4 + 2s) \leq 1 - s + \beta(s)$ (with equality when $0 < s \leq \frac{1}{2}$). This completes the proof. \square

With the results of [Proposition 4.1](#) and [Lemma 4.2](#), we are able to iterate the argument. At time $t = T$, write $u = u^2 + v^2$ where u^2 is the solution to

$$\begin{cases} i \frac{\partial u^2}{\partial t} - Hu^2 = |u^2|^2 u^2, & (t, x) \in \mathbb{R} \times \mathbb{R}^3, \\ u^2(T) = u^1(T) + w^1(T) \in \mathcal{H}^1(\mathbb{R}^3), \end{cases} \tag{4-14}$$

and where $v^2 = e^{-itH} u_0^N + w^2$ satisfies

$$\begin{cases} i \frac{\partial w^2}{\partial t} - Hw^2 = |w^2 + e^{-itH} u_0^N + u^2|^2 (w^2 + e^{-itH} u_0^N + u^2) - |u^2|^2 u^2, & (t, x) \in \mathbb{R} \times \mathbb{R}^3, \\ w^2(T) = 0. \end{cases}$$

By [Proposition 4.1](#), $w^1(T) \in \mathcal{H}^1(\mathbb{R}^3)$; thus [\(4-14\)](#) is globally well-posed. Then, thanks to [Lemma 4.2](#), by the conservation of the energy,

$$E(u^2) = E(u^1(T) + w^1(T)) \leq 4N^{2(1-s+\varepsilon)},$$

and, by the conservation of the mass,

$$\|u^2\|_{L_T^\infty L^2} = \|u^1(T) + w^1(T)\|_{L^2} \leq 2N^\varepsilon.$$

Therefore there exists a set $\Sigma_T^2 \subset \mathcal{H}^s$ with

$$\mu(\Sigma_T^2) \geq 1 - C \exp(-cT^{-\delta} \|\gamma\|_{\mathcal{H}^s}^{-2}),$$

and such that for all $u_0 \in \Sigma_T^2$ there exists a unique $w^2 \in \mathcal{C}([T, 2T], \mathcal{H}^1(\mathbb{R}^3))$ that satisfies the result of Proposition 4.1, with the same $T > 0$. Here we use crucially that the large deviation bounds of Proposition 2.1 are invariant under time shift τ .

Fix a time $A > 0$. We can iterate the previous argument and construct u^j, v^j and w^j for $1 \leq j \leq \lfloor A/T \rfloor$ such that the function u^j is the solution to (4-14) with initial condition

$$u^j(t = (j - 1)T) = u^{j-1}((j - 1)T) + w^{j-1}((j - 1)T),$$

then we set $v^j(t) = e^{-itH} u_0^N + w^j(t)$, where the function w^j is the solution to

$$\begin{cases} i \frac{\partial w^j}{\partial t} - Hw^j = |w^j + e^{-itH} u_0^N + u^j|^2 (w^j + e^{-itH} u_0^N + u^j) - |u^j|^2 u^j, & (t, x) \in \mathbb{R} \times \mathbb{R}^3, \\ w^j((j - 1)T) = 0. \end{cases}$$

This enables us to define a unique solution u to the initial problem (1-7) defined by $u(t) = u^j(t) + v^j(t)$ for $t \in [(j - 1)T, jT]$, with $1 \leq j \leq \lfloor A/T \rfloor$ provided that $u_0 \in \Gamma_T^A$, where $\Gamma_T^A := \bigcap_{j=1}^{\lfloor A/T \rfloor} \Sigma_T^j$.

Thanks to the exponential bounds, we have $\mu((\Gamma_T^A)^c) \leq C \exp(-cT^{-\delta/2} \|\gamma\|_{\mathcal{H}^s}^{-2})$, with $T = N^{-4(1-s)-\varepsilon}$.

For uniform bounds on the energy and the mass, it remains to check whether $E(u^j) \leq 4N^{2(1-s+\varepsilon)}$ and $\|u^j\|_{L^2(\mathbb{R}^3)} \leq 2N^\varepsilon$ for all $1 \leq j \leq \lfloor A/T \rfloor$. By Lemma 4.2, for $T = N^{-4(1-s)-}$,

$$E(u^j) \leq E(u_{0,N}) + CAT^{-1} N^{1-s+\beta(s)+} \leq 2N^{2(1-s+\varepsilon)} + CAN^{\beta(s)+5(1-s)+}, \tag{4-15}$$

which satisfies the prescribed bound if and only if $3(1 - s) + \beta(s) < 0$.

- If $\frac{1}{2} \leq s \leq 1$, the condition is $3(1 - s) + 2s - \frac{7}{2} < 0$, or equivalently $s > -\frac{1}{2}$, which is satisfied.
- If $0 \leq s \leq \frac{1}{2}$, the condition is $3(1 - s) - \frac{5}{2} < 0$, or equivalently $s > \frac{1}{6}$. The same argument applies to control $\|u^j\|_{L^2}$.

If $\frac{1}{6} < s < 1$, we optimise in (4-15) with the choice of $N \geq 1$ so that $A \sim cN^{-3(1-s)-\beta(s)}$, and get that, for $1 \leq j \leq \lfloor A/T \rfloor$,

$$E(u^j) \leq CA^{c_s+},$$

with

$$c_s = \begin{cases} \frac{2(1-s)}{6s-1} & \text{if } \frac{1}{6} < s \leq \frac{1}{2}, \\ \frac{2(1-s)}{2s+1} & \text{if } \frac{1}{2} \leq s \leq 1. \end{cases} \tag{4-16}$$

Denote by $\Gamma^A = \Gamma_T^A$ the set defined with the previous choice of N and $T = N^{-4(1-s)-\varepsilon}$.

Lemma 4.3. *Let $\frac{1}{6} < s < 1$. Then for all $A \in \mathbb{N}$ and all $u_0 \in \Gamma^A$ there exists a unique solution to (1-7) on $[0, A]$, which reads*

$$u(t) = e^{-itH} u_0 + w(t), \quad \text{with } w \in \mathcal{C}([0, A], \mathcal{H}^1(\mathbb{R}^3)), \quad \sup_{t \in [0, A]} E(w(t)) \leq CA^{c_s+}.$$

Proof. On the time interval $[(j - 1)T, jT]$ we have $u = u^j + v^j$ where $v^j = e^{-itH} u_0^N + w^j$ and $u^j = e^{-itH} u_{0,N} + z^j$, for some $z^j \in \mathcal{C}([0, +\infty[, \mathcal{H}^1(\mathbb{R}^3))$. Therefore, if we define $w \in \mathcal{C}([0, A], \mathcal{H}^1(\mathbb{R}^3))$

by $w(t) = z^j(t) + w^j(t)$ for $t \in [(j-1)T, jT]$ and $1 \leq j \leq \lfloor A/T \rfloor$, we get $u(t) = e^{-itH}u_0 + w(t)$ for all $t \in [0, A]$. Next, for $t \in [(j-1)T, jT]$,

$$E(w(t)) \leq CE(z^j) + CE(w^j) \leq CE(u^j) + CE(e^{-itH}u_{0,N}) + CE(w^j) \leq CA^{c_s+},$$

which was the claim. □

We are now able to complete the proof of [Theorem 1.5](#). Set

$$\Theta = \bigcap_{k=1}^{+\infty} \bigcup_{A \geq k} \Gamma^A \quad \text{and} \quad \Sigma = \Theta + \mathcal{H}^1.$$

We have $\mu(\Theta) = \lim_{k \rightarrow \infty} \mu(\bigcup_{A \geq k} \Gamma^A)$ and $\mu(\bigcup_{A \geq k} \Gamma^A) \geq 1 - c \exp(-k^\delta \|\gamma\|_{\mathcal{H}^s}^{-2})$. So $\mu(\Theta) = 1$, and thus $\mu(\Sigma) = 1$.

By definition, for all $u_0 \in \Theta$, there exists a unique global solution to (1-7), which reads

$$u(t) = e^{-itH}u_0 + w(t), \quad w \in \mathcal{C}(\mathbb{R}, \mathcal{H}^1(\mathbb{R}^3)).$$

Then by [Lemma 4.3](#) for all $u_0 \in \Theta$, there exists a unique $w \in \mathcal{C}([0, +\infty[, \mathcal{H}^1(\mathbb{R}^3))$, which satisfies, for all N , the bound

$$\sup_{t \in [0, N]} E(w(t)) \leq CN^{c_s+}.$$

Now, if $U_0 \in \Sigma$ then $U_0 = u_0 + v$ with $u_0 \in \Theta$, $v \in \mathcal{H}^1$ and we can use the method of [Proposition 4.1](#), [Lemma 4.2](#) and [Lemma 4.3](#) with $U_{0,N}$ replaced by $u_{0,N} + v$. And the set Σ satisfies properties (i) and (ii).

Coming back to the definition of Σ_T^j , we have $e^{-itH}(\Sigma_T^j) = \Sigma_T^j$ for all $t \in \mathbb{R}$; thus $e^{-itH}(\Theta) = \Theta$. Finally, thanks to property (i), the set Σ is invariant under the dynamics and property (iii) is satisfied.

4B. The case of dimension $d = 2$. In this section, we prove [Theorem 1.6](#). The proof is analogous to [Theorem 1.5](#) in a simpler context; that is why we only explain the key estimates.

According to [Proposition 2.1](#), we set

$$F_s(K) = \{w \in \mathcal{H}^s(\mathbb{R}^2) \mid \|w\|_{\mathcal{H}^s(\mathbb{R}^2)} \leq K, \|w\|_{L^4(\mathbb{R}^2)} \leq K \text{ and } \|e^{-itH}w\|_{L_{[0,2\pi]}^{1/\varepsilon} \mathcal{W}^{1+s-\varepsilon, \infty}(\mathbb{R}^2)} \leq K\},$$

and we fix $u_0 \in F_s(N^\varepsilon)$.

Then, if $f = e^{-itH}u_0^N$, we have

$$\|f\|_{L_{[0,2\pi]}^\infty L^2} \leq CN^{-s+\varepsilon} \quad \text{and} \quad \|H^{\sigma/2}f\|_{L_{[0,2\pi]}^\infty L^\infty} \leq CN^{\sigma-1-s+\varepsilon}.$$

In [Proposition 4.1](#) we can choose $T = N^{-2(1-s)-\varepsilon}$ to have

$$\|u^1\|_{L_T^\infty L^2} \leq CN^\varepsilon \quad \text{and} \quad \|u^1\|_{L_T^\infty \mathcal{H}^1} \leq CN^{1-s+\varepsilon}.$$

Moreover, as $u_0 \in F_s(N^\varepsilon)$, we obtain

$$\|u_{0,N}\|_{L^4} \leq CN^\varepsilon.$$

Hence, we establish

$$E(u^1) = \|u^1\|_{\mathcal{H}^1(\mathbb{R}^2)}^2 + \frac{1}{2}\|u^1\|_{L^4(\mathbb{R}^2)}^4 = \|u_{0,N}\|_{\mathcal{H}^1(\mathbb{R}^2)}^2 + \frac{1}{2}\|u_{0,N}\|_{L^4(\mathbb{R}^2)}^4 \leq N^{2(1-s+\varepsilon)} + CN^{4\varepsilon} \leq 4N^{2(1-s+\varepsilon)},$$

and

$$\|u^1\|_{L_T^\infty L^4} \leq CN^{(1-s+\varepsilon)/2}.$$

In Proposition 4.1, we obtain $\|w^1\|_{L_{[0,T]}^\infty \mathcal{H}^1} \leq CN^{-1+}$ and $\|w^1\|_{L_{[0,T]}^\infty L^2} \leq CN^{-2+}$. The proof is essentially the same. We define the map L as in (4-9). For the first estimate, we prove that

$$\|L(w)\|_{X_T^1} \leq CN^{-1+} + N^{0-}\|w\|_{X_T^1} + N^{-2(1-s)+}\|w\|_{X_T^1}^3.$$

We only give details for the source terms. First,

$$\begin{aligned} \|fu^2\|_{L_T^{1+} W^{1,2-} + L_T^1 \mathcal{H}^1} &\leq C\|fuH^{1/2}u\|_{L_T^{1+} L^{2-}} + C\|u^2H^{1/2}f\|_{L_T^1 L^2} \\ &\leq CT^{1-}\|u\|_{L_T^\infty \mathcal{H}^1}\|u\|_{L_T^\infty L^\infty} + C\|f\|_{L_T^\infty L^\infty} + CT^{1-}\|u\|_{L_T^\infty L^4}^2\|H^{1/2}f\|_{L_T^\infty L^\infty} \\ &\leq CT^{1-}\max(N^{1-3s+}, N^{1-2s+}) \leq CT^{1-}N^{1-2s+} \leq CN^{-1+}. \end{aligned}$$

Similarly,

$$\begin{aligned} \|f^2u\|_{L_T^1 \mathcal{H}^1} &\leq C\|f^2H^{1/2}u\|_{L_T^1 L^2} + C\|ufH^{1/2}f\|_{L_T^1 L^2} \\ &\leq CT^{1-}\|u\|_{L_T^\infty \mathcal{H}^1}\|f\|_{L_T^\infty L^\infty}^2 + CT^{1-}\|u\|_{L_T^\infty L^2}\|f\|_{L_T^\infty L^\infty}\|H^{1/2}f\|_{L_T^\infty L^\infty} \\ &\leq CT^{1-}\max(N^{-1-3s+}, N^{-1-2s+}) \leq CT^{1-}N^{-1-2s+} \leq CN^{-3+} \leq CN^{-1+}. \end{aligned}$$

Finally,

$$\begin{aligned} \|f^3\|_{L_T^1 \mathcal{H}^1} &\leq C\|f^2H^{1/2}f\|_{L_T^1 L^2} \leq CT^{1-}\|H^{1/2}f\|_{L_T^\infty L^\infty}\|f\|_{L_T^\infty L^\infty}\|f\|_{L_T^\infty L^2} \\ &\leq CT^{1-}N^{-s+}N^{-1-s+}N^{-s+} \leq CN^{-3-s+} \leq CN^{-1+}. \end{aligned}$$

Analogously to Lemma 4.2, we obtain $|E(u^1(T) + w^1(T)) - E(u^1(T))| \leq CN^{-s+}$, because here $\beta(s) = 1-$ and the estimates on u^1 are the same as in dimension $d = 3$.

Finally, the globalisation argument holds if (4-15) is satisfied, that is to say

$$CAT^{-1}N^{-s+} \leq 4N^{2(1-s)+},$$

which is equivalent to $2(1-s) - s < 2(1-s)$, hence $s > 0$. In this case, we set $A \sim cN^s$ and we get that, for $0 \leq t \leq A$,

$$E(w(t)) \leq CA^{c_s+}, \text{ with } c_s = \frac{1-s}{s}.$$

Theorem 1.6 follows.

Acknowledgements

We are grateful to Nicolas Burq for discussions on this subject. We thank Rémi Carles for discussions on scattering theory which led us toward Theorem 1.4.

References

- [Banica et al. 2008] V. Banica, R. Carles, and G. Staffilani, “Scattering theory for radial nonlinear Schrödinger equations on hyperbolic space”, *Geom. Funct. Anal.* **18**:2 (2008), 367–399. MR 2010h:35364 Zbl 1186.35198
- [Barab 1984] J. E. Barab, “Nonexistence of asymptotically free solutions for a nonlinear Schrödinger equation”, *J. Math. Phys.* **25**:11 (1984), 3270–3273. MR 86a:35121 Zbl 0554.35123
- [Bourgain 1994] J. Bourgain, “Periodic nonlinear Schrödinger equation and invariant measures”, *Comm. Math. Phys.* **166**:1 (1994), 1–26. MR 95k:35185 Zbl 0822.35126
- [Bourgain 1996] J. Bourgain, “Invariant measures for the 2D-defocusing nonlinear Schrödinger equation”, *Comm. Math. Phys.* **176**:2 (1996), 421–445. MR 96m:35292 Zbl 0852.35131
- [Bourgain 1999] J. Bourgain, *Global solutions of nonlinear Schrödinger equations*, American Mathematical Society Colloquium Publications **46**, Amer. Math. Soc., Providence, RI, 1999. MR 2000h:35147 Zbl 0933.35178
- [Burq and Lebeau 2013] N. Burq and G. Lebeau, “Injections de Sobolev probabilistes et applications”, *Ann. Sci. Éc. Norm. Supér.* (4) **46**:6 (2013), 917–962. MR 3134684 Zbl 06275361
- [Burq and Tzvetkov 2008a] N. Burq and N. Tzvetkov, “Random data Cauchy theory for supercritical wave equations, I: Local theory”, *Invent. Math.* **173**:3 (2008), 449–475. MR 2009k:58057 Zbl 1156.35062
- [Burq and Tzvetkov 2008b] N. Burq and N. Tzvetkov, “Random data Cauchy theory for supercritical wave equations, II: A global existence result”, *Invent. Math.* **173**:3 (2008), 477–496. MR 2010i:58025 Zbl 1187.35233
- [Burq and Tzvetkov 2014] N. Burq and N. Tzvetkov, “Probabilistic well-posedness for the cubic wave equation”, *J. Eur. Math. Soc. (JEMS)* **16**:1 (2014), 1–30. MR 3141727 Zbl 06260652
- [Burq et al. 2010] N. Burq, L. Thomann, and N. Tzvetkov, “Long time dynamics for the one dimensional non linear Schrödinger equation”, preprint, 2010. To appear in *Ann. Inst. Fourier*. arXiv 1002.4054
- [Burq et al. 2012] N. Burq, L. Thomann, and N. Tzvetkov, “Global infinite energy solutions for the cubic wave equation”, preprint, 2012. To appear in *Bull. Soc. Math. France*. arXiv 1210.2086
- [Burq et al. ≥ 2014] N. Burq, L. Thomann, and N. Tzvetkov, “Remarks on the Gibbs measures for nonlinear dispersive equations”, In preparation.
- [Carles 2009] R. Carles, “Rotating points for the conformal NLS scattering operator”, *Dyn. Partial Differ. Equ.* **6**:1 (2009), 35–51. MR 2010i:35365 Zbl 1191.35270
- [Carles 2011] R. Carles, “Nonlinear Schrödinger equation with time dependent potential”, *Commun. Math. Sci.* **9**:4 (2011), 937–964. MR 2901811 Zbl 1285.35105
- [Cazenave 2003] T. Cazenave, *Semilinear Schrödinger equations*, Courant Lecture Notes in Mathematics **10**, Courant Institute, New York, 2003. MR 2004j:35266 Zbl 1055.35003
- [Cazenave et al. 2011] T. Cazenave, D. Fang, and Z. Han, “Continuous dependence for NLS in fractional order spaces”, *Ann. Inst. H. Poincaré Anal. Non Linéaire* **28**:1 (2011), 135–147. MR 2012a:35292 Zbl 1209.35124
- [Colliander and Oh 2012] J. Colliander and T. Oh, “Almost sure well-posedness of the cubic nonlinear Schrödinger equation below $L^2(\mathbb{T})$ ”, *Duke Math. J.* **161**:3 (2012), 367–414. MR 2881226 Zbl 1260.35199
- [Deng 2012] Y. Deng, “Two-dimensional nonlinear Schrödinger equation with random radial data”, *Anal. PDE* **5**:5 (2012), 913–960. MR 3022846 Zbl 1264.35212
- [Imekraz et al. 2014] R. Imekraz, D. Robert, and L. Thomann, “On random Hermite series”, preprint, 2014. arXiv 1403.4913
- [Karadzhov 1995] G. E. Karadzhov, “Riesz summability of multiple Hermite series in L^p spaces”, *Math. Z.* **219**:1 (1995), 107–118. MR 96g:42015 Zbl 0824.42019
- [Koch and Tataru 2005] H. Koch and D. Tataru, “ L^p eigenfunction bounds for the Hermite operator”, *Duke Math. J.* **128**:2 (2005), 369–392. MR 2006d:35044 Zbl 1075.35020
- [Lebowitz et al. 1988] J. L. Lebowitz, H. A. Rose, and E. R. Speer, “Statistical mechanics of the nonlinear Schrödinger equation”, *J. Statist. Phys.* **50**:3–4 (1988), 657–687. MR 89f:82006 Zbl 1084.82506
- [Nahmod and Staffilani 2013] A. Nahmod and G. Staffilani, “Almost sure well-posedness for the periodic 3D quintic nonlinear Schrödinger equation below the energy space”, preprint, 2013. arXiv 1308.1169

- [Nahmod et al. 2013] A. R. Nahmod, N. Pavlović, and G. Staffilani, “Almost sure existence of global weak solutions for supercritical Navier–Stokes equations”, *SIAM J. Math. Anal.* **45**:6 (2013), 3431–3452. MR 3131480 Zbl 06261007
- [Nakanishi and Ozawa 2002] K. Nakanishi and T. Ozawa, “Remarks on scattering for nonlinear Schrödinger equations”, *NoDEA Nonlinear Differential Equations Appl.* **9**:1 (2002), 45–68. MR 2003a:35177 Zbl 0991.35082
- [Oh 1989] Y.-G. Oh, “Cauchy problem and Ehrenfest’s law of nonlinear Schrödinger equations with potentials”, *J. Differential Equations* **81**:2 (1989), 255–274. MR 90i:35260 Zbl 0703.35158
- [Oh 2009] T. Oh, “Invariant Gibbs measures and a.s. global well posedness for coupled KdV systems”, *Diff. Int. Eq.* **22**:7-8 (2009), 637–668. MR 2010g:35275 Zbl 1240.35477
- [Oh 2009/10] T. Oh, “Invariance of the Gibbs measure for the Schrödinger–Benjamin–Ono system”, *SIAM J. Math. Anal.* **41**:6 (2009/10), 2207–2225. MR 2011b:35467 Zbl 1205.35268
- [Poiret 2012a] A. Poiret, “Solutions globales pour des équations de Schrödinger sur-critiques en toutes dimensions”, preprint, 2012. arXiv 1207.3519
- [Poiret 2012b] A. Poiret, “Solutions globales pour l’équation de Schrödinger cubique en dimension 3”, preprint, 2012. arXiv 1207.1578
- [Poiret et al. 2013] A. Poiret, D. Robert, and L. Thomann, “Random weighted Sobolev inequalities on \mathbb{R}^d and application to Hermite functions”, preprint, 2013. To appear in *Ann. Henri Poincaré*. arXiv 1307.4976
- [de Suzzoni 2013] A.-S. de Suzzoni, “Large data low regularity scattering results for the wave equation on the Euclidean space”, *Comm. Partial Differential Equations* **38**:1 (2013), 1–49. MR 3005545 Zbl 1267.35130
- [Thangavelu 1993] S. Thangavelu, *Lectures on Hermite and Laguerre expansions*, Mathematical Notes **42**, Princeton University Press, 1993. MR 94i:42001 Zbl 0791.41030
- [Thomann 2009] L. Thomann, “Random data Cauchy problem for supercritical Schrödinger equations”, *Ann. Inst. H. Poincaré Anal. Non Linéaire* **26**:6 (2009), 2385–2402. MR 2010m:35494 Zbl 1180.35491
- [Tzvetkov 2006] N. Tzvetkov, “Invariant measures for the nonlinear Schrödinger equation on the disc”, *Dyn. Partial Differ. Equ.* **3**:2 (2006), 111–160. MR 2008f:35386 Zbl 1142.35090
- [Tzvetkov 2008] N. Tzvetkov, “Invariant measures for the defocusing nonlinear Schrödinger equation”, *Ann. Inst. Fourier (Grenoble)* **58**:7 (2008), 2543–2604. MR 2010g:35307 Zbl 1171.35116
- [Tzvetkov 2010] N. Tzvetkov, “Construction of a Gibbs measure associated to the periodic Benjamin–Ono equation”, *Probab. Theory Related Fields* **146**:3-4 (2010), 481–514. MR 2011b:35471 Zbl 1188.35183
- [Yajima and Zhang 2004] K. Yajima and G. Zhang, “Local smoothing property and Strichartz inequality for Schrödinger equations with potentials superquadratic at infinity”, *J. Differential Equations* **202**:1 (2004), 81–110. MR 2005f:35266 Zbl 1060.35121
- [Zhang 2005] J. Zhang, “Sharp threshold for blowup and global existence in nonlinear Schrödinger equations under a harmonic potential”, *Comm. Partial Differential Equations* **30**:10-12 (2005), 1429–1443. MR 2006j:35226 Zbl 1081.35109
- [Zhidkov 2001] P. E. Zhidkov, *Korteweg-de Vries and nonlinear Schrödinger equations: qualitative theory*, Lecture Notes in Mathematics **1756**, Springer, Berlin, 2001. MR 2002h:35285 Zbl 0987.35001

Received 20 Sep 2013. Revised 9 Apr 2014. Accepted 8 May 2014.

AURÉLIEN POIRET: aurelien.poiret@math.u-psud.fr

Laboratoire de Mathématiques, UMR 8628 du CNRS, Université Paris Sud, 91405 Orsay Cedex, France

DIDIER ROBERT: didier.robert@univ-nantes.fr

Laboratoire de Mathématiques J. Leray, UMR 6629 du CNRS, Université de Nantes, 2, rue de la Houssinière, 44322 Nantes Cedex 03, France

LAURENT THOMANN: laurent.thomann@univ-nantes.fr

Laboratoire de Mathématiques J. Leray, UMR 6629 du CNRS, Université de Nantes, 2, rue de la Houssinière, 44322 Nantes Cedex 03, France

Analysis & PDE

msp.org/apde

EDITORS

EDITOR-IN-CHIEF

Maciej Zworski
zworski@math.berkeley.edu

University of California
Berkeley, USA

BOARD OF EDITORS

Nicolas Burq	Université Paris-Sud 11, France nicolas.burq@math.u-psud.fr	Yuval Peres	University of California, Berkeley, USA peres@stat.berkeley.edu
Sun-Yung Alice Chang	Princeton University, USA chang@math.princeton.edu	Gilles Pisier	Texas A&M University, and Paris 6 pisier@math.tamu.edu
Michael Christ	University of California, Berkeley, USA mchrist@math.berkeley.edu	Tristan Rivière	ETH, Switzerland riviere@math.ethz.ch
Charles Fefferman	Princeton University, USA cf@math.princeton.edu	Igor Rodnianski	Princeton University, USA irod@math.princeton.edu
Ursula Hamenstaedt	Universität Bonn, Germany ursula@math.uni-bonn.de	Wilhelm Schlag	University of Chicago, USA schlag@math.uchicago.edu
Vaughan Jones	U.C. Berkeley & Vanderbilt University vaughan.f.jones@vanderbilt.edu	Sylvia Serfaty	New York University, USA serfaty@cims.nyu.edu
Herbert Koch	Universität Bonn, Germany koch@math.uni-bonn.de	Yum-Tong Siu	Harvard University, USA siu@math.harvard.edu
Izabella Laba	University of British Columbia, Canada ilaba@math.ubc.ca	Terence Tao	University of California, Los Angeles, USA tao@math.ucla.edu
Gilles Lebeau	Université de Nice Sophia Antipolis, France lebeau@unice.fr	Michael E. Taylor	Univ. of North Carolina, Chapel Hill, USA met@math.unc.edu
László Lempert	Purdue University, USA lempert@math.purdue.edu	Gunther Uhlmann	University of Washington, USA gunther@math.washington.edu
Richard B. Melrose	Massachusetts Institute of Technology, USA rbm@math.mit.edu	András Vasy	Stanford University, USA andras@math.stanford.edu
Frank Merle	Université de Cergy-Pontoise, France Frank.Merle@u-cergy.fr	Dan Virgil Voiculescu	University of California, Berkeley, USA dvv@math.berkeley.edu
William Minicozzi II	Johns Hopkins University, USA minicozz@math.jhu.edu	Steven Zelditch	Northwestern University, USA zelditch@math.northwestern.edu
Werner Müller	Universität Bonn, Germany mueller@math.uni-bonn.de		

PRODUCTION

production@msp.org

Silvio Levy, Scientific Editor

See inside back cover or msp.org/apde for submission instructions.

The subscription price for 2014 is US \$180/year for the electronic version, and \$355/year (+\$50, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues from the last three years and changes of subscribers address should be sent to MSP.

Analysis & PDE (ISSN 1948-206X electronic, 2157-5045 printed) at Mathematical Sciences Publishers, 798 Evans Hall #3840, c/o University of California, Berkeley, CA 94720-3840, is published continuously online. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices.

APDE peer review and production are managed by EditFLOW[®] from Mathematical Sciences Publishers.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2014 Mathematical Sciences Publishers

ANALYSIS & PDE

Volume 7 No. 4 2014

- The global stability of the Minkowski spacetime solution to the Einstein-nonlinear system in wave coordinates 771
JARED SPECK
- Dispersion for the Schrödinger equation on the line with multiple Dirac delta potentials and on delta trees 903
VALERIA BANICA and LIVIU I. IGNAT
- The Cuntz semigroup and stability of close C^* -algebras 929
FRANCESC PERERA, ANDREW TOMS, STUART WHITE and WILHELM WINTER
- Wave and Klein–Gordon equations on hyperbolic spaces 953
JEAN-PHILIPPE ANKER and VITTORIA PIERFELICE
- Probabilistic global well-posedness for the supercritical nonlinear harmonic oscillator 997
AURÉLIEN POIRET, DIDIER ROBERT and LAURENT THOMANN