

ANALYSIS & PDE

Volume 7

No. 5

2014

ALAIN GRIGIS AND ANDRÉ MARTINEZ

**RESONANCE WIDTHS FOR THE MOLECULAR
PREDISSOCIATION**



RESONANCE WIDTHS FOR THE MOLECULAR PREDISSOCIATION

ALAIN GRIGIS AND ANDRÉ MARTINEZ

We consider a semiclassical 2×2 matrix Schrödinger operator of the form

$$P = -h^2 \Delta \mathbf{I}_2 + \text{diag}(V_1(x), V_2(x)) + hR(x, hD_x),$$

where V_1, V_2 are real-analytic, V_2 admits a nondegenerate minimum at 0 with $V_2(0) = 0$, V_1 is nontrapping at energy 0, and $R(x, hD_x) = (r_{j,k}(x, hD_x))_{1 \leq j, k \leq 2}$ is a symmetric 2×2 matrix of first-order pseudodifferential operators with analytic symbols. We also assume that $V_1(0) > 0$. Then, denoting by e_1 the first eigenvalue of $-\Delta + \langle V_2''(0)x, x \rangle / 2$, and under some ellipticity condition on $r_{1,2}$ and additional generic geometric assumptions, we show that the unique resonance ρ_1 of P such that $\rho_1 = (e_1 + r_{2,2}(0, 0))h + \mathcal{O}(h^2)$ (as $h \rightarrow 0_+$) satisfies

$$\text{Im } \rho_1 = -h^{n_0 + (1 - n_\Gamma)/2} f\left(h, \ln \frac{1}{h}\right) e^{-2S/h},$$

where $f(h, \ln \frac{1}{h}) \sim \sum_{0 \leq m \leq \ell} f_{\ell, m} h^\ell (\ln \frac{1}{h})^m$ is a symbol with $f_{0,0} > 0$, $S > 0$ is the so-called Agmon distance associated with the degenerate metric $\max(0, \min(V_1, V_2)) dx^2$, between 0 and $\{V_1 \leq 0\}$, and $n_0 \geq 1, n_\Gamma \geq 0$ are integers that depend on the geometry.

1. Introduction

The theory of predissociation goes back to the very first years of quantum mechanics (see [Kronig 1928; Landau 1932a; 1932b; Zener 1932; Stückelberg 1932], for example). Roughly speaking, it describes the possibility for a molecule to dissociate spontaneously (after a sufficiently large time) into several submolecules, for energies below the crossing of the corresponding energy surfaces of the initial molecule and the final dissociated state. From a physical point of view, one naturally expects that this (typically quantum) phenomenon occurs with extremely small (but nonzero) probability.

Despite the fact that statements concerning this problem are present in the physics literature for more than 70 years, the first mathematically rigorous result is due to M. Klein [1987], where an upper bound on the time of predissociation is given in the framework of the Born–Oppenheimer approximation. More precisely, denoting by h the square root of the ratio of electronic to nuclear mass, Klein proves the existence of resonances ρ with real part below the crossing of the energy surfaces and with exponentially small imaginary part; that is

$$|\text{Im } \rho| = \mathcal{O}(e^{-2(1-\varepsilon)S/h}),$$

Grigis partly supported by the GDR INdAM-CNRS GrefiMefi. Martinez partly supported by Università di Bologna, Funds for Selected Research Topics.

MSC2010: primary 35P15, 35C20; secondary 35S99, 47A75.

Keywords: resonances, Born–Oppenheimer approximation, eigenvalue crossing, microlocal analysis.

where $S > 0$ is a geometric constant, $\varepsilon > 0$ is fixed arbitrarily, and the estimate holds uniformly as h goes to zero.

In terms of probabilities, this result corresponds to an upper bound on the transition probability between the initial molecule and the dissociated state. The purpose of this article is to obtain more complete information on this quantity, in particular, a lower bound. More precisely, under suitable conditions, we prove that the imaginary part of the lowest resonance admits a complete asymptotic expansion of the type

$$\operatorname{Im} \rho_1 = -h^{n_0+(1-n_\Gamma)/2} e^{-2S/h} \sum_{0 \leq m \leq \ell} f_{\ell,m} h^\ell \left(\ln \frac{1}{h} \right)^m,$$

in the sense that, for any $N \geq 1$, one has

$$\left| \operatorname{Im} \rho_1 + h^{n_0+(1-n_\Gamma)/2} e^{-2S/h} \sum_{0 \leq m \leq \ell \leq N} f_{\ell,m} h^\ell \left(\ln \frac{1}{h} \right)^m \right| = \mathcal{O}(h^{n_0+(1-n_\Gamma)/2+N} e^{-2S/h}),$$

where $S > 0$, $n_0 \geq 1$ and $n_\Gamma \geq 0$ are all geometric constants, and where the leading coefficient $f_{0,0}$ is positive.

As is well-known, the quantity $\operatorname{Im} \rho$ is closely related to the oscillatory behavior of the corresponding resonant state in the unbounded classically allowed region. Hence, the main issue will be to know sufficiently well this behavior.

The strategy of the proof consists in starting from the WKB construction at the bottom of the well and then trying to extend it as much as possible, at least up to the classically allowed unbounded region. This is mainly the same strategy used in [Helffer and Sjöstrand 1986] for the study of shape resonances.

However, from a technical point of view, several new problems are encountered, because of the crossing of the electronic levels.

The first one is that, at the crossing, the only reference on WKB constructions is that of [Pettersson 1997], which has been done for a special type of matrix Schrödinger operators. In particular, it strongly uses the fact that only differential operators are involved. In our case, since our operator comes from a Born–Oppenheimer reduction, it is necessarily of pseudodifferential kind (see [Klein et al. 1992; Martinez and Sordoni 2009], for example). As a consequence, our first step will consist in extending Pettersson’s method to pseudodifferential operators. Unfortunately, this extension is far from being straightforward, and needs a specific formal calculus adapted to expressions involving the Weber functions.

The second one is that, after having overcome the crossing, the symbols of the resulting WKB expansions do not anymore satisfy analytic estimates (usually needed in order to resum them, up to exponentially small error terms). In particular, this prevents us from using directly the constructions of [Helffer and Sjöstrand 1986] near the classically allowed unbounded region. Instead, we have to adapt the method of Fujiié, Lahmar-Benbernou and Martinez [Fujiié et al. 2011], which, without analyticity, allows us to extend the WKB constructions into the classically allowed unbounded region up to a distance of order $(h \ln |h|)^{2/3}$ from the barrier. This is not much, but it is enough to have sufficient control in this region on the difference between the true solution and the WKB one. This is actually done by adapting the specific arguments of propagation introduced in [loc. cit.], where the propagation takes place in h -dependent domains.

In the next section, we describe in details the geometrical context and the assumptions.

In [Section 3](#), we state our main result.

[Section 4](#) is devoted to the WKB constructions, starting from the well and proceeding away along some minimal geodesics, until crossing the boundary of the classically forbidden region. It is in this section that we develop a formal pseudodifferential calculus adapted to expressions involving the Weber functions.

Next, in [Section 5](#), we extend the well-known Agmon estimates to our pseudodifferential context. In this case, the main feature is that, since we cannot use general Lipschitz weight functions, we replace them by h -dependent smooth functions with bounded gradient, but with derivatives of higher order that can grow to infinity as $h \rightarrow 0$.

In [Section 6](#), we use these estimates in order to obtain a bound for the difference between the WKB solutions and a solution of a modified problem, and this permits us to define an asymptotic solution in a whole neighborhood of the classically forbidden region (but only up to a distance of order $(h \ln |h|)^{1/3}$ from this region).

[Section 7](#) contains the a priori estimates and the propagation arguments that lead to a good control on the difference between the asymptotic solution and the actual one.

Finally, [Section 8](#) makes the link with the width of the resonance. Even if the idea is standard (practically an application of the Green formula; see [\[Helffer and Sjöstrand 1986\]](#), for example), here we have to be careful with the double problem that, on the one hand, we deal with pseudodifferential (not differential) operators and, on the other hand, the magnitude of freedom outside the classically forbidden region is of order $(h \ln |h|)^{1/3}$ as $h \rightarrow 0$.

2. Geometrical assumptions

We consider the semiclassical 2×2 matrix Schrödinger operator

$$P = \begin{pmatrix} P_1 & 0 \\ 0 & P_2 \end{pmatrix} + hR(x, hD_x), \tag{2-1}$$

with

$$P_j := -h^2 \Delta + V_j(x), \quad j = 1, 2,$$

where $x = (x_1, \dots, x_n)$ is the current variable in \mathbb{R}^n ($n \geq 1$), $h > 0$ denotes the semiclassical parameter, and $R(x, hD_x) = (r_{j,k}(x, hD_x))_{1 \leq j,k \leq 2}$ is a formally self-adjoint 2×2 matrix of first-order semiclassical pseudodifferential operators, in the sense that, for all $\alpha \in \mathbb{N}^{2n}$, $\partial^\alpha r_{j,k}(x, \xi) = \mathcal{O}(1 + |\xi|)$ uniformly on \mathbb{R}^{2n} .


Let us observe that this is typically the kind of operator one obtains in the Born–Oppenheimer approximation, after reduction to an effective Hamiltonian [\[Klein et al. 1992; Martinez and Sordoni 2009\]](#). In that case, the quantity h^2 stands for the inverse of the mass of the nuclei.

Assumption 1: The potentials V_1 and V_2 are smooth and bounded on \mathbb{R}^n , and satisfy:

$$V_1(0) > 0 \text{ and } E = 0 \text{ is a nontrapping energy for } V_1, \tag{2-2}$$

$$V_1 \text{ has a strictly negative limit as } |x| \rightarrow \infty, \tag{2-3}$$

$$V_2 \geq 0, \quad V_2^{-1}(0) = \{0\}, \quad \text{Hess } V_2(0) > 0, \quad \liminf_{|x| \rightarrow \infty} V_2 > 0. \tag{2-4}$$



In particular, we assume that V_2 has a unique nondegenerate well at $x = 0$. We define the island \ddot{O} as the bounded open set

$$\ddot{O} = \{x \in \mathbb{R}^n : V_1(x) > 0\}, \tag{2-5}$$

and the sea as the set where $V_1(x) < 0$. With (2-2) and (2-4), the well $\{x = 0\}$ for V_2 is included in the island.

The fact that 0 is a nontrapping energy for V_1 means that, for any $(x, \xi) \in p_1^{-1}(0)$, one has that $|\exp t H_{p_1}(x, \xi)| \rightarrow +\infty$ as $t \rightarrow \infty$, where we let $p_1(x, \xi) := \xi^2 + V_1(x)$ be the symbol of P_1 and $H_{p_1} := (\nabla_\xi p_1, -\nabla_x p_1)$ be the Hamilton field of p_1 .

Conditions (2-2)–(2-4) correspond to *molecular predissociation*, as described in [Klein 1987].

Since we plan to study the resonances of P near the energy level $E = 0$, we also assume:

Assumption 2: The potentials V_1 and V_2 extend to bounded holomorphic functions near a complex sector of the form $\mathcal{S}_{R_0, \delta} := \{x \in \mathbb{C}^n : |\operatorname{Re} x| \geq R_0, |\operatorname{Im} x| \leq \delta |\operatorname{Re} x|\}$, with $R_0, \delta > 0$. Moreover V_1 tends to its limit at ∞ in this sector and $\operatorname{Re} V_2$ stays away from 0 in this sector.

Assumption 3: The symbols $r_{j,k}(x, \xi)$ for $(j, k) = (1, 1), (1, 2), (2, 2)$ extend to holomorphic functions in (x, ξ) near

$$\tilde{\mathcal{F}}_{R_0, \delta} := \mathcal{S}_{R_0, \delta} \times \{\xi \in \mathbb{C}^n : |\operatorname{Im} \xi| \leq \max(\delta \langle \operatorname{Re} x \rangle, \sqrt{M_0})\},$$

with

$$M_0 > \sup_{x \in \mathbb{R}^n} \min(V_1(x), V_2(x)),$$

and, for real x , $r_{j,k}$ is a smooth function of x with values in the set of holomorphic functions of ξ near $\{|\operatorname{Im} \xi| \leq \sqrt{M_0}\}$. Moreover we assume that, for any $\alpha \in \mathbb{N}^{2n}$, they satisfy

$$\partial^\alpha r_{j,k}(x, \xi) = \mathcal{O}(\langle \operatorname{Re} \xi \rangle) \quad \text{uniformly on } \tilde{\mathcal{F}}_{R_0, \delta} \cup (\mathbb{R}^n \times \{|\operatorname{Im} \xi| \leq \sqrt{M_0}\}). \tag{2-6}$$

Now we define the cirque Ω as

$$\Omega = \{x \in \mathbb{R}^n : V_2(x) < V_1(x)\}. \tag{2-7}$$

Hence, the well is in the cirque and the cirque is in the island.

We also consider the Agmon distance associated to the pseudometric

$$(\min(V_1, V_2))_+ dx^2;$$

see [Pettersson 1997]. There are three places where this metric is not a standard one.

The first is near the well 0, but this case is well-known. It was treated in [loc. cit.] and also in [Helffer and Sjöstrand 1984]. The Agmon distance,

$$\varphi(x) := d(x, 0), \tag{2-8}$$

is smooth at 0. The point $(x, \xi) = (0, 0)$ is a hyperbolic singular point of the Hamilton vector field H_{q_2} , where $q_2 = \xi^2 - V_2(x)$, and the stable and unstable manifold near this point are respectively the Lagrangian manifolds $\{\xi = \nabla\varphi(x)\}$ and $\{\xi = -\nabla\varphi(x)\}$.

Secondly, on $\partial\Omega$, precisely at the points where $V_1 = V_2$. This case has been also considered by Pettersson. At such a point, if one assume that $\nabla V_1 \neq \nabla V_2$, then any geodesic which is transversal to the hypersurface $\{V_1 = V_2\}$ is C^1 .

Finally there is the boundary of the island $\partial\ddot{O}$, where $V_1 = 0$. This situation was considered in [Helffer and Sjöstrand 1986]. We will follow this work in the next assumption.

Now we consider the distance from the well to the sea, that is, to $\partial\ddot{O}$:

$$S := d(0, \partial\ddot{O}). \tag{2-9}$$

Setting $B_S := \{x \in \ddot{O} : \varphi(x) < S\}$ and denoting by \bar{B}_S its closure, we also consider the set $\bar{B}_S \cap \partial\ddot{O}$ that consists of the points of the boundary of the island that are joined to the well by a minimal d -geodesic included in the island. These points are called points of type 1 in [loc. cit.], and we denote by G the set of minimal geodesics joining such a point to 0 in \ddot{O} .

We make the following assumption:

Assumption 4: For all $\gamma \in G$, γ intersects $\partial\Omega$ at a finite number of points and the intersection is transversal at each of these points. Moreover, $\nabla V_1 \neq \nabla V_2$ on $\gamma \cap \partial\Omega$.

Let us recall that the assumption that 0 is a nontrapping energy for V_1 implies that $\nabla V_1 \neq 0$ on $\partial\ddot{O}$, and therefore that $\partial\ddot{O}$ is a smooth hypersurface.

We define the caustic set \mathcal{C} as the union of the set of points of type 1 and the set of points $x \in \ddot{O}$ with $\varphi(x) = S + d(x, \partial\ddot{O})$. As in [loc. cit.] we assume:

Assumption 5: The points of type 1 form a submanifold Γ , and \mathcal{C} has a contact of order exactly two with $\partial\ddot{O}$ along Γ .

We denote by n_Γ the dimension of Γ . Moreover, for any $\gamma \in G$, we denote by $N_\gamma := \#(\gamma \cap \partial\Omega)$ the number of points where γ crosses the boundary of the cirque, and we set

$$n_0 := \min_{\gamma \in G} N_\gamma, \quad G_0 := \{\gamma \in G : N_\gamma = n_0\}.$$

Then, we make an assumption that somehow insures that an interaction between the two Schrödinger operators does exist.

Assumption 6: There exists at least one $\gamma \in G_0$ for which the ellipticity condition $r_{12}(x, i\nabla\varphi(x)) \neq 0$ holds at every point $x \in \gamma \cap \partial\Omega$.

3. Main result

Under the previous assumption we plan to study the resonances of the operator P given in (2-1), where $R(x, hD_x)$ is defined as

$$R(x, hD_x) := \begin{pmatrix} \text{Op}_h^L(r_{1,1}) & \text{Op}_h^L(r_{1,2}) \\ \text{Op}_h^R(\overline{r_{1,2}}) & \text{Op}_h^L(r_{2,2}) \end{pmatrix},$$

where for any symbol $a(x, \xi)$ we use the quantizations

$$\begin{aligned} \text{Op}_h^L(a)u(x) &= \frac{1}{(2\pi h)^n} \int e^{i(x-y)\xi/h} a(x, \xi)u(y) dy d\xi, \\ \text{Op}_h^R(a)u(x) &= \frac{1}{(2\pi h)^n} \int e^{i(x-y)\xi/h} a(y, \xi)u(y) dy d\xi. \end{aligned}$$

In order to define the resonances we consider the distortion given as follows. Let $F(x) \in C^\infty(\mathbb{R}^n, \mathbb{R}^n)$ be such that $F(x) = 0$ for $|x| \leq R_0$ and $F(x) = x$ for $|x|$ large enough. For $\theta > 0$ small enough, we define the distorted operator P_θ as the value at $\nu = i\theta$ of the extension to the complex numbers of the operator $U_\nu P U_\nu^{-1}$, which is defined for ν real small enough and analytic in ν , where we have set

$$U_\nu \phi(x) = \det(1 + \nu dF(x))^{1/2} \phi(x + \nu F(x)). \tag{3-1}$$

Since we have a pseudodifferential operator $R(x, hD)$, the fact that $U_\nu P U_\nu^{-1}$ is analytic in ν is not completely standard but can be done without problem (thanks to [Assumption 3](#)), and by using the Weyl perturbation theorem, one can also see that there exists $\varepsilon_0 > 0$ such that for any $\theta > 0$ small enough, the spectrum of P_θ is discrete in $[-\varepsilon_0, \varepsilon_0] - i[0, \varepsilon_0\theta]$. The eigenvalues of P_θ are called the resonances of P [[Hunziker 1986](#); [Helffer and Sjöstrand 1986](#); [Helffer and Martinez 1987](#)].

We will need another small parameter $k > 0$ related to the semiclassical parameter $h > 0$, defined as

$$k := h \ln \frac{1}{h}. \tag{3-2}$$

In the sequel, we will study the resonances in the domain $[-\varepsilon_0, Ch] - i[0, Ck]$, where $C > 0$ is arbitrarily large. In this case, we can adapt the WKB constructions near the well made in [[Helffer and Sjöstrand 1984](#)] and show that these resonances form a finite set $\{\rho_1, \dots, \rho_m\}$, with asymptotic expansions as $h \rightarrow 0$ of the form

$$\rho_j \sim h \sum_{\ell \geq 0} \rho_{j,\ell} h^{\ell/2},$$

where $\rho_{j,\ell} \in \mathbb{R}$ and $\rho_{j,0} = e_j + r_{2,2}(0, 0)$, e_j being the j -th eigenvalue of the harmonic oscillator $-\Delta + \langle V_2''(0)x, x \rangle / 2$ (actually, to be more precise, one must also assume that the arbitrarily large constant C does not coincide with one of the e_j).

In this paper we are interested in the imaginary part of these resonances. We have:

Theorem 3.1. *Under Assumptions 1 to 6, the first resonance ρ_1 of P is such that*

$$\text{Im } \rho_1 = -h^{n_0+(1-n_\Gamma)/2} f\left(h, \ln \frac{1}{h}\right) e^{-2S/h},$$

where $f\left(h, \ln \frac{1}{h}\right)$ admits an asymptotic expansion of the form

$$f\left(h, \ln \frac{1}{h}\right) \sim \sum_{0 \leq m \leq \ell} f_{\ell,m} h^\ell \left(\ln \frac{1}{h}\right)^m \quad \text{as } h \rightarrow 0,$$

with $f_{0,0} > 0$ and $S > 0$ as defined in (2-9).

Moreover the other resonances in $[-\varepsilon_0, Ch] - i[0, Ck]$ verify

$$\text{Im } \rho_j = \mathcal{O}(h^{\beta_j} e^{-2S/h}),$$

for some real β_j , uniformly as $h \rightarrow 0$.

4. WKB constructions

In this section, we fix some minimal d -geodesic $\gamma \in G$ and we denote by $x^{(1)}, \dots, x^{(N_\gamma)}$ the sequence of points that constitute $\gamma \cap \partial\Omega$, ordered from the closest to 0 up to the closest to \check{O} (note that N_γ is necessarily an odd number). We also denote by $\gamma^{(1)}, \gamma^{(2)}, \dots, \gamma^{(N_\gamma+1)}$ the portions of $\gamma \setminus \partial\Omega$ that are in-between 0 and $x^{(1)}, x^{(1)}$ and $x^{(2)}, \dots, x^{(N_\gamma)}$ and \check{O} , respectively, in such a way that we have

$$\gamma = \gamma^{(1)} \cup \{x^{(1)}\} \cup \gamma^{(2)} \cup \dots \cup \{x^{(N_\gamma)}\} \cup \gamma^{(N_\gamma+1)},$$

where the union is disjoint (in particular, by convention we assume that $0 \in \gamma^{(1)}$). Moreover, we start by considering the first resonance ρ_1 only.

In the cirque. As in [Pettersson 1997], the starting point of the construction consists of the WKB asymptotics given near the well $x = 0$ by a method due to Helffer and Sjöstrand [1984]. More precisely, because of the matricial nature of the operator and the fact that p_1 is elliptic above $x = 0$, one finds a formal solution w_1 of $Pw_1 = \rho_1 w_1$ of the form

$$w_1(x; h) = \begin{pmatrix} ha_1(x, h) \\ a_2(x, h) \end{pmatrix} e^{-\varphi(x)/h}, \tag{4-1}$$

where φ is defined in (2-8) and a_j ($j = 1, 2$) is a classical symbol of order 0 in h , that is, a formal series in h of the form

$$a_j(x, h) = \sum_{k=0}^{\infty} h^k a_{j,k}(x), \tag{4-2}$$

with $a_{j,k}$ smooth near 0 (here no half-powers of h appear since we consider the first resonance ρ_1 only). Moreover, a_2 is elliptic in the sense that $a_{2,0}$ never vanishes. Note that the generalization of the constructions of [Helffer and Sjöstrand 1984] to the case of pseudodifferential operators is done by the use of a so-called *formal semiclassical pseudodifferential calculus*, which in our case is based on the following result.

Lemma 4.1. *Let $\tilde{\varphi} = \tilde{\varphi}(x)$ be a real bounded C^∞ function on \mathbb{R}^n and let $p = p(x, \xi) \in S(1)$ extend to a bounded function, holomorphic with respect to ξ in a neighborhood of the set*

$$\{(x, \xi) \in \text{supp } \nabla \tilde{\varphi} \times \mathbb{C}^n : |\text{Im } \xi| \leq |\nabla \tilde{\varphi}(x)|\}.$$

Then, denoting by Op_h^L the left (or standard) semiclassical quantization of symbols, the operator $e^{\tilde{\varphi}/h} \text{Op}_h^L(p)e^{-\tilde{\varphi}/h}$ is uniformly bounded on $L^2(\mathbb{R}^n)$ and, for any $a \in C_0^\infty(\mathbb{R}^n)$ and $N \geq 1$, one has, with $\Phi(x, y) := \tilde{\varphi}(x) - \tilde{\varphi}(y) - (x - y)\nabla\tilde{\varphi}(x)$,

$$(e^{\tilde{\varphi}/h} \text{Op}_h^L(p)e^{-\tilde{\varphi}/h}a)(x; h) = \sum_{|\alpha| \leq N} \frac{1}{\alpha!} \left(\frac{h}{i}\right)^{|\alpha|} \partial_\xi^\alpha p(x, i\nabla\tilde{\varphi}(x)) \partial_y^\alpha (a(y)e^{\Phi(x,y)/h}) \Big|_{y=x} + \mathcal{O}(h^{N/2}), \tag{4-3}$$

locally uniformly with respect to x , and uniformly with respect to h small enough.

The proof of this lemma is rather standard, e.g., [Martinez 1987] and we omit it.

Then, the construction can be performed by using the formal series given in (4-3) in order to define the formal action of $R(x, hD_x)$ on w_1 . Afterwards, these constructions can be continued along the integral curves of the vector field $\nabla_\xi p_2(x, i\nabla\varphi(x))D_x = 2\nabla\varphi(x) \cdot \nabla_x$ (that is, along the minimal geodesic of d starting at 0), as long as $p_1(x, i\nabla\varphi(x))$ does not vanish (that is, as long as these minimal geodesics stay inside the cirque Ω). In that way, after resummation and multiplication by a cutoff function, we obtain a function w_1 of the form (4-1) that satisfies

$$Pw_1 - \rho_1 w_1 = \mathcal{O}(h^\infty e^{-\varphi/h}) \tag{4-4}$$

locally uniformly in $\bigcup \gamma$, where the union is taken over all the minimal d -geodesics γ coming from the well 0 and staying in Ω . In particular, (4-4) is satisfied in a neighborhood \mathcal{N}_1 of $\gamma^{(1)}$.

At the boundary of the cirque. Now, we study the situation near the point $x^{(1)} \in \partial\Omega$. By Theorem 2.14 of [Pettersson 1997], we know that there exist a neighborhood \mathcal{V}_1 of $x^{(1)}$ and two positive functions $\varphi_1, \varphi_2 \in C^\infty(\mathcal{V}_1)$ such that

$$\begin{aligned} \varphi_1 &= \varphi && \text{on } \mathcal{V}_1 \cap \{V_1 < V_2\}; \\ \varphi_2 &= \varphi && \text{on } \mathcal{V}_1 \cap \{V_2 < V_1\}; \\ |\nabla\varphi_j(x)|^2 &= V_j(x), && j = 1, 2; \\ \varphi_1 &= \varphi_2 \text{ and } \nabla\varphi_1 = \nabla\varphi_2 && \text{on } \mathcal{V}_1 \cap \partial\Omega; \\ \varphi_2(x) - \varphi_1(x) &\sim d(x, \partial\Omega)^2. \end{aligned}$$

Actually, φ_2 is just $d_2(0, x)$, where d_2 is the Agmon distance associated with the metric $V_2(x) dx^2$ and φ_1 is the phase function of the Lagrangian manifold obtained as the flow-out of $\{(x, \nabla\varphi_2(x)) : x \in \mathcal{V}_1 \cap \partial\Omega\}$ under the Hamilton flow of $q_1(x, \xi) := \xi^2 - V_1(x)$.

Then, we set

$$\psi := \frac{1}{2}(\varphi_1 + \varphi_2), \tag{4-5}$$

and we consider the smooth function $z(x)$ defined for $x \in \mathcal{V}_1$ by

$$\begin{aligned} z(x)^2 &= 2(\varphi_2(x) - \varphi_1(x)) \\ z(x) &< 0 \quad \text{on } \mathcal{V}_1 \cap \{V_2 < V_1\}. \end{aligned} \tag{4-6}$$

In order to extend the WKB construction (4-1) across $\partial\Omega$ near $x^{(1)}$, we follow Pettersson and try a formal ansatz,

$$w_2(x; h) = \sum_{k \geq 0} h^k \left(\alpha_k(x, h) Y_{k,0} \left(\frac{z(x)}{\sqrt{h}} \right) + \sqrt{h} \beta_k(x, h) Y_{k,1} \left(\frac{z(x)}{\sqrt{h}} \right) \right) e^{-\psi(x)/h}, \tag{4-7}$$

where

$$\alpha_k(x, h) = \begin{pmatrix} h\alpha_{k,1}(x, h) \\ \alpha_{k,2}(x, h) \end{pmatrix}, \quad \beta_k(x, h) = \begin{pmatrix} \beta_{k,1}(x, h) \\ h\beta_{k,2}(x, h) \end{pmatrix}. \tag{4-8}$$

Here $\alpha_{k,j}$ and $\beta_{k,j}$ are formal symbols of the type

$$\sum_{l \geq 0} \sum_{m=0}^l h^l (\ln h)^m \gamma^{l,m}(x) \tag{4-9}$$

(with $\gamma^{l,m}$ smooth in ω_1) and, for any $k \geq 0$ and $\varepsilon \in \mathbb{C}$, the function $Y_{k,\varepsilon}$ is the so-called Weber function, defined by

$$Y_{k,\varepsilon}(z) = \partial_\varepsilon^k Y_{0,\varepsilon}(z), \tag{4-10}$$

where $Y_{0,\varepsilon}$ is the unique entire function with respect to ε and z that is a solution of the Weber equation,

$$Y''_{0,\varepsilon} + \left(\frac{1}{2} - \varepsilon - \frac{z^2}{4} \right) Y_{0,\varepsilon} = 0, \tag{4-11}$$

such that, for $\varepsilon > 0$, one has

$$Y_{0,\varepsilon}(z) \sim e^{-z^2/4} z^{-\varepsilon} \quad \text{as } z \rightarrow -\infty. \tag{4-12}$$

(Then, one also has $Y_{0,\varepsilon}(z) \sim (\sqrt{2\pi}/\Gamma(\varepsilon)) e^{z^2/4} z^{\varepsilon-1}$ as $z \rightarrow +\infty$, by Proposition A.2 of [Pettersson 1997].) As is shown in Pettersson’s Theorem 4.3, a resummation of (4-7) is possible up to an error of order $\mathcal{O}(h^\infty e^{-\varphi/h})$.

Now, since φ is not C^∞ (but only C^1) near $x^{(1)}$, we need to find some generalization of Lemma 4.1. For technical reasons, in the rest of this section we prefer to work with the *right* semiclassical quantization of symbols, which we denote by Op_h^R .

For $\nu_0 > 0$ and $g \in C^\infty(\mathbb{R}^{2n}; \mathbb{R}_+)$, we denote by $S_{\nu_0}(g(x, \xi))$ the set of (possibly h -dependent) functions $p \in C^\infty(\mathbb{R}^{2n})$ that extend to holomorphic functions with respect to ξ in the strip

$$\mathcal{A}_{\nu_0} := \{(x, \xi) \in \mathbb{R}^n \times \mathbb{C}^n : |\text{Im } \xi| < \nu_0\}$$

such that, for all $\alpha \in \mathbb{N}^{2n}$, one has

$$\partial^\alpha p(x, \xi) = \mathcal{O}(g(x, \text{Re } \xi)), \tag{4-13}$$

uniformly with respect to $(x, \xi) \in \mathcal{A}_{\nu_0}$ and $h > 0$ small enough. We also denote by $S_0(g)$ the analogous space of smooth symbols obtained by switching \mathbb{R}^{2n} to \mathcal{A}_{ν_0} and “smooth” to “holomorphic”.

Lemma 4.2. *Let $\nu_0 > 0$, $m \in \mathbb{R}$, $p = p(x, \xi) \in S_{\nu_0}(\langle \xi \rangle^m)$, and let $\phi = \phi(x)$ be a real bounded Lipschitz function on \mathbb{R}^n such that*

$$\|\nabla \phi(x)\|_{L^\infty} < \nu_0.$$

Let also $a = a(x; h) \in C^\infty(\mathbb{R}^n)$ be such that, for all $\alpha \in \mathbb{N}^n$,

$$(hD_x)^\alpha a(x; h) = \mathcal{O}(e^{-\phi(x)/h}),$$

uniformly with respect to h small enough and $x \in \mathbb{R}^n$. Then

$$(\text{Op}_h^R(p)a)(x; h) = \mathcal{O}(e^{-\phi(x)/h})$$

uniformly with respect to h small enough and $x \in \mathbb{R}^n$.

Proof. We write

$$e^{\phi(x)/h} \text{Op}_h^R(p)a(x; h) = \frac{1}{(2\pi h)^n} \int e^{i(x-y)\xi/h + \phi(x)/h} p(y, \xi) a(y; h) dy d\xi \tag{4-14}$$

and, following [Sjöstrand 1982], we make the change of contour of integration in ξ ,

$$\mathbb{R}^n \ni \xi \mapsto \xi + i\nu_1 \frac{x-y}{|x-y|}, \tag{4-15}$$

where $\|\nabla\phi(x)\|_{L^\infty} < \nu_1 < \nu_0$. We obtain

$$e^{\phi(x)/h} \text{Op}_h^R(p)a(x; h) = \frac{1}{(2\pi h)^n} \int e^{i(x-y)\xi/h} p\left(y, \xi + i\nu_1 \frac{x-y}{|x-y|}\right) \theta(x, y; h) dy d\xi, \tag{4-16}$$

with

$$\theta(x, y; h) = a(y; h) e^{(\phi(x) - \nu_1|x-y|)/h} = \mathcal{O}(e^{\phi(x) - \phi(y) - \nu_1|x-y|/h}).$$

Therefore

$$\theta(x, y; h) = \mathcal{O}(e^{-\delta|x-y|/h}), \tag{4-17}$$

with $\delta = \nu_1 - \|\nabla\phi\|_{L^\infty} > 0$.

Then, in the case $m < -n$, the result follows immediately from (4-16)–(4-17) (and standard estimates on oscillatory integrals). In the general case, we just write

$$\text{Op}_h^R(p) = \text{Op}_h^R(p)(2\nu_0 - h^2\Delta_x)^{-k} (2\nu_0 - h^2\Delta_x)^k, \tag{4-18}$$

with k an integer large enough (e.g., $k = 1 + |[m]| + n$) and, since $\text{Op}_h^R(p)(2\nu_0 - h^2\Delta_x)^{-k}$ is a semiclassical pseudodifferential operator with (h -dependent) symbol in $S_{\nu_0}(\langle \xi \rangle^{m-2k}) \subset S_{\nu_0}(\langle \xi \rangle^{-n-1})$, the result follows by applying the previous case with a replaced by $(2\nu_0 - h^2\Delta_x)^k a$. □

Now, as preparation for defining a formal pseudodifferential calculus acting on expressions such as (4-7), for $j = 1, \dots, n$ and $x \in \omega_1$, we set

$$A_j(x) := \begin{pmatrix} \frac{\partial\varphi_2(x)}{\partial x_j} & 0 \\ 0 & \frac{\partial\varphi_1(x)}{\partial x_j} \end{pmatrix} \in \mathcal{M}_2(\mathbb{R}). \tag{4-19}$$

Then, for any $k \geq 0$, we have (see [Pettersson 1997, (4.18)])

$$(hD_{x_j} - iA_j(x)) \begin{pmatrix} Y_{k,0} \left(\frac{z(x)}{\sqrt{h}} \right) \\ Y_{k,1} \left(\frac{z(x)}{\sqrt{h}} \right) \end{pmatrix} e^{-\psi(x)/h} = \frac{\sqrt{h}}{i} (\partial_{x_j} z(x)) \begin{pmatrix} kY_{k-1,1} \left(\frac{z(x)}{\sqrt{h}} \right) \\ Y_{k,0} \left(\frac{z(x)}{\sqrt{h}} \right) \end{pmatrix} e^{-\psi(x)/h}. \tag{4-20}$$

If a and b are (scalar) formal symbols of the type (4-9) and $k \in \mathbb{N}$, we set

$$I_k(a, b)(x; h) = a(x; h)Y_{k,0} \left(\frac{z(x)}{\sqrt{h}} \right) + b(x; h)Y_{k,1} \left(\frac{z(x)}{\sqrt{h}} \right), \tag{4-21}$$

and we plan to exploit (4-20) in order to define a formal action of a pseudodifferential operator on $I_k(a, b)e^{-\psi/h}$. Using (4-20), we see that we have

$$\begin{aligned} (hD_x - i\nabla\varphi_2(x))(I_k(a, 0)e^{-\psi/h}) &= (I_k(hD_x a, 0) + I_{k-1}(0, k\sqrt{h}aD_x z))e^{-\psi/h}, \\ (hD_x - i\nabla\varphi_1(x))(I_k(0, b)e^{-\psi/h}) &= I_k(\sqrt{h}bD_x z, hD_x b)e^{-\psi/h}. \end{aligned} \tag{4-22}$$

Now, for any $p \in S_{v_0}(\langle \xi \rangle^m)$, $N \geq 1$ and $j = 1, 2$, Taylor’s formula gives

$$p(x, \xi) = \sum_{|\alpha| \leq N} \frac{1}{\alpha!} \partial_\xi^\alpha p(x, i\nabla\varphi_j(x)) (\xi - i\nabla\varphi_j(x))^\alpha + \sum_{|\alpha| = N+1} p_{j,\alpha}(x, \xi) (\xi - i\nabla\varphi_j(x))^\alpha, \tag{4-23}$$

where the $p_{j,\alpha}$ are in $S_{v_0}(\langle \xi \rangle^m)$. Moreover, we have:

Lemma 4.3. *Let $v_0 > \sup_{x \in \omega_1} \min(\sqrt{V_1(x)}, \sqrt{V_2(x)})$ and $m \in \mathbb{R}$. Then, for any $q = q(x, \xi) \in S_{v_0}(\langle \xi \rangle^m)$, $k \geq 0$, a in $C_0^\infty(\omega_1)$ and $\alpha \in \mathbb{N}^n$, one has*

$$\begin{aligned} \text{Op}_h^R(q(x, \xi)(\xi - i\nabla\varphi_2(x))^\alpha)(I_k(a, 0)e^{-\psi(x)/h}) &= \mathcal{O}(|\ln h|^k h^{|\alpha|/2} e^{-\varphi(x)/h}), \\ \text{Op}_h^R(q(x, \xi)(\xi - i\nabla\varphi_1(x))^\alpha)(I_k(0, a)e^{-\psi(x)/h}) &= \mathcal{O}(|\ln h|^k h^{|\alpha|/2} e^{-\varphi(x)/h}), \end{aligned} \tag{4-24}$$

where the estimates hold uniformly for h small enough and $x \in \mathbb{R}^n$.

Proof. We prove both estimates together, by induction on $|\alpha|$. We first notice that, by Lemma 4.6 of [Pettersson 1997], for $\beta \in \mathbb{N}^n$ and $j \in \{0, 1\}$, one has

$$(hD_x)^\beta \left(Y_{k,j} \left(\frac{z(x)}{\sqrt{h}} \right) e^{-\psi(x)/h} \right) = \mathcal{O}(|\ln h|^k e^{-\varphi(x)/h}). \tag{4-25}$$

As a consequence, the result for $\alpha = 0$ follows directly from Lemma 4.2.

Now, assume it is true for $|\alpha| \leq N$ ($N \in \mathbb{N}$ fixed arbitrarily), and let $\gamma \in \mathbb{N}^n$, $|\gamma| = 1$. Using the notation

$$I_{k,2}(a) := I_k(a, 0)e^{-\psi/h}, \quad I_{k,1}(a) := I_k(0, a)e^{-\psi/h}, \tag{4-26}$$

we write (for $|\alpha| \leq N$ and $j = 1, 2$)

$$\begin{aligned} \text{Op}_h^R(q(x, \xi)(\xi - i\nabla\varphi_j(x))^{\alpha+\gamma}) I_{k,j}(a) e^{-\psi(x)/h} \\ = \frac{1}{(2\pi h)^n} \int e^{i(x-y)\xi/h} f_\alpha(y, \xi) (\xi - i\nabla\varphi_j(y))^\gamma I_{k,j}(a)(y) e^{-\psi(y)/h} dy d\xi, \end{aligned}$$

with $f_\alpha(y, \xi) := q(y, \xi)(\xi - i\nabla\varphi_j(y))^\alpha$. Now, assuming without loss of generality that $\gamma = (1, 0, \dots, 0)$ and using the fact that

$$\xi_1 e^{i(x-y)\xi/h} = -h D_{y_1} (e^{i(x-y)\xi/h}),$$

we obtain

$$\begin{aligned} \text{Op}_h^R(q(x, \xi)(\xi - i\nabla\varphi_j(x))^{\alpha+\gamma}) I_{k,j}(a) e^{-\psi(x)/h} \\ = \frac{1}{(2\pi h)^n} \int e^{i(x-y)\xi/h} \left(h D_{y_1} - i \frac{\partial\varphi_j}{\partial x_1}(y) \right) I_{k,j}(f_\alpha(y, \xi) a(y)) e^{-\psi(y)/h} dy d\xi, \end{aligned}$$

and therefore, by (4-22),

$$\text{Op}_h^R(q(x, \xi)(\xi - i\nabla\varphi_j(x))^{\alpha+\gamma}) I_{k,j}(a) e^{-\psi(x)/h} = \frac{1}{(2\pi h)^n} \int e^{i(x-y)\xi/h} \tilde{I}_{k,j}(f_\alpha(y, \xi) a(y)) e^{-\psi(y)/h} dy d\xi,$$

with

$$\tilde{I}_{k,2}(a) := h I_{k,2}(D_{x_1} a) + k\sqrt{h} I_{k-1,1}(a D_{x_1} z), \quad \tilde{I}_{k,1}(a) := \sqrt{h} I_{k,2}(a D_{x_1} z) + h I_{k,1}(D_{x_1} a).$$

Then, applying the induction hypothesis (and using the fact that $D_{y_1} f_\alpha$ is a sum of terms of the type $g(y, \xi)(\eta - iA(y))^\beta$ with $|\beta| \geq |\alpha| - 1$) this gives

$$\begin{aligned} \text{Op}_h^R(q(x, \xi)(\xi - i\nabla\varphi_j(x))^{\alpha+\gamma}) I_{k,j}(a) e^{-\psi(x)/h} \\ = \mathcal{O}(|\ln h|^k h^{1+(|\alpha|-1)/2} + |\ln h|^{k-1} h^{(1+|\alpha|)/2} + |\ln h|^k h^{(1+|\alpha|)/2}) e^{-\varphi/h} = \mathcal{O}(|\ln h|^k h^{(1+|\alpha|)/2}) e^{-\varphi/h}, \end{aligned}$$

and the proof is complete. □

Using Lemma 4.3 and (4-23), for any a in $C_0^\infty(\omega_1)$, we obtain (with the notation (4-26))

$$\begin{aligned} \text{Op}_h^R(p)(I_{k,j}(a) e^{-\psi/h}) \\ = \sum_{|\alpha| \leq N} \frac{1}{\alpha!} \text{Op}_h^R(\partial_\xi^\alpha p(x, i\nabla\varphi_j(x))(\xi - i\nabla\varphi_j(x))^\alpha)(I_{k,j}(a) e^{-\psi(x)/h}) + \mathcal{O}(h^{N/2} e^{-\varphi/h}) \\ = \sum_{\substack{|\alpha| \leq N \\ \beta \leq \alpha}} \frac{1}{i^{|\beta|} \beta! (\alpha - \beta)!} \text{Op}_h^R(\partial_\xi^\alpha p(x, i\nabla\varphi_j(x)) (\nabla\varphi_j)^\beta \xi^{\alpha-\beta})(I_{k,j}(a) e^{-\psi(x)/h}) + \mathcal{O}(h^{N/2} e^{-\varphi/h}), \end{aligned}$$

and thus, as before, writing down the corresponding oscillatory integral, in the same way we deduce

$$\begin{aligned} \text{Op}_h^R(p)(I_{k,j}(a) e^{-\psi/h}) \\ = \sum_{\substack{|\alpha| \leq N \\ \beta \leq \alpha}} \frac{1}{i^{|\beta|} \beta! (\alpha - \beta)!} (h D_x)^{\alpha-\beta} [(\nabla\varphi_j)^\beta \partial_\xi^\alpha p(x, i\nabla\varphi_j) I_{k,j}(a) e^{-\psi/h}] + \mathcal{O}(h^{N/2} e^{-\varphi/h}). \quad (4-27) \end{aligned}$$

Now, for an integer $M \leq 0$ and $\Omega \subset \mathbb{R}^n$ open, we consider the space of sequences of formal symbols,

$$S^M(\omega_1) := \left\{ a = (a_k)_{k \in \mathbb{N}} : a_k(x, h) = \sum_{l=-M}^\infty \sum_{m=0}^l h^l (\ln h)^m \gamma_k^{l,m}(x), \gamma_k^{l,m} \in C^\infty(\omega_1) \right\},$$

and, for $a, b \in S^M(\omega_1)$, we set

$$I(a, b) := \sum_{k \geq 0} h^k I_k(a_k, \sqrt{h}b_k). \tag{4-28}$$

Using (4-22), we see that, for $j = 1, \dots, n$, we have

$$hD_{x_j} I(a, b)e^{-\psi/h} = I((iA_j + L_j)(a, b))e^{-\psi/h}, \tag{4-29}$$

where $iA_j(a, b) = (i(\partial_{x_j}\varphi_2)a, i(\partial_{x_j}\varphi_1)b)$ and L_j is the operator

$$L_j : S^M \times S^M \rightarrow S^{M-1} \times S^{M-1}, \\ (a, b) \mapsto (\tilde{a}^j, \tilde{b}^j),$$

defined by, for $k \in \mathbb{N}$,

$$\tilde{a}_k^j := hD_{x_j}a_k + hb_kD_{x_j}z, \\ \tilde{b}_k^j := hD_{x_j}b_k + (k+1)ha_{k+1}D_{x_j}z. \tag{4-30}$$

In particular, using the notation $L = (L_1, \dots, L_n)$ and $L^\alpha = L_1^{\alpha_1} \dots L_n^{\alpha_n}$, for all $\alpha \in \mathbb{N}^n$ we have

$$L^\alpha \text{ maps } S^M(\omega_1) \times S^M(\omega_1) \text{ into } S^{M-|\alpha|}(\omega_1) \times S^{M-|\alpha|}(\omega_1). \tag{4-31}$$

For any smooth diagonal $\mathcal{M}_2(\mathbb{C})$ -valued function $B(x) = \text{diag}(B_1(x), B_2(x))$, we let it act on $S^M \times S^M$ by setting

$$B(a, b) = (B_1a, B_2b), \tag{4-32}$$

and we define the formal action of a pseudodifferential operator with symbol $p \in S_{v_0}(\langle \xi \rangle^m)$ on expressions of the type $I(a, b)e^{-\psi/h}$ by the formula

$$\text{Op}_h^F(p)(I(a, b)e^{-\psi/h}) := \sum_{\substack{\alpha \in \mathbb{N}^n \\ \beta \leq \alpha}} \frac{1}{i^{|\beta|}\beta!(\alpha - \beta)!} I((iA(x) + L)^{\alpha - \beta} A(x)^\beta \partial_\xi^\alpha p(x, iA(x))(a, b))e^{-\psi/h}, \tag{4-33}$$

where we have also set $A := (A_1, \dots, A_n)$ and

$$\partial_\xi^\alpha p(x, iA(x))(a, b) := (\partial_\xi^\alpha p(x, i(\nabla\varphi_2))a, \partial_\xi^\alpha p(x, i(\nabla\varphi_1))b).$$

Then, in view of Lemma 4.3 and (4-27), we immediately obtain:

Proposition 4.4. *Let $a, b \in S^M(\omega_1)$ and denote by $\tilde{I}(a, b)e^{-\psi/h}$ any resummation of $I(a, b)e^{-\psi/h}$ up to an $\mathcal{O}(h^\infty e^{-\varphi/h})$ error term. Then, for any $\chi \in C_0^\infty(\omega_1)$, the quantity $\text{Op}_h^R(p)(\chi \tilde{I}(a, b)e^{-\psi/h})$ is a resummation of $\text{Op}_h^F(p)(I(\chi a, \chi b)e^{-\psi/h})$, up to an $\mathcal{O}(h^\infty e^{-\varphi/h})$ error term.*

In particular, the operator P naturally acts (up to $\mathcal{O}(h^\infty e^{-\varphi/h})$ error terms) on expressions of the type

$$w_2 = \begin{pmatrix} I(h\alpha_1, \beta_1) \\ I(\alpha_2, h\beta_2) \end{pmatrix} e^{-\psi/h}, \tag{4-34}$$

where $\alpha_j = (\alpha_{j,k})_{k \geq 0}$ and $\beta_j = (\beta_{j,k})_{k \geq 0}$ are in $S^0(\omega_1)$ ($j = 1, 2$).

Writing down the equation $\tilde{P}w_2 = \rho_1 w_2$, setting

$$\alpha_{j,k} = \sum_{l \geq 0} \sum_{m=0}^l h^l (\ln h)^m \alpha_{j,k}^{l,m}(x),$$

and the analogous formula for $\beta_{j,k}$, and identifying the coefficients of $h^l (\ln h)^m$ for $0 \leq m \leq l \leq 1$, we find (denoting by $p = \begin{pmatrix} p_1 + hr_{1,1} & hr_{1,2} \\ hr_{2,1} & p_2 + hr_{2,2} \end{pmatrix}$ the right symbol of P),

$$p_1(x, i\nabla\varphi_2)\alpha_{1,0}^{0,0} + r_{1,2}(x, i\nabla\varphi_2)\alpha_{2,0}^{0,0} + \left[\frac{1}{i} \nabla_\xi p_1(x, i\nabla\varphi_1)(\nabla z) + \frac{1}{2} \langle (\text{Hess}_\xi p_1)(x, i\nabla\varphi_1) \nabla z, \nabla(\varphi_2 - \varphi_1) \rangle \right] \beta_{1,0}^{0,0} = 0; \tag{4-35}$$

$$[\partial_\xi p_1(x, i\nabla\varphi_1) D_x - i(\nabla_x \cdot \nabla_\xi p_1)(x, i\nabla\varphi_1) + r_{1,1}(x, i\nabla\varphi_1) - \rho_1] \beta_{1,0}^{0,0} = 0; \tag{4-36}$$

$$p_2(x, i\nabla\varphi_1)\beta_{2,0}^{0,0} + r_{2,1}(x, i\nabla\varphi_1)\beta_{1,0}^{0,0} + \left[\frac{1}{i} \partial_\xi p_2(x, i\nabla\varphi_2)(\nabla z) + \frac{1}{2} \langle (\text{Hess}_\xi p_2)(x, i\nabla\varphi_2) \nabla z, \nabla(\varphi_1 - \varphi_2) \rangle \right] \alpha_{2,1}^{0,0} = 0; \tag{4-37}$$

$$(\partial_\xi p_2(x, i\nabla\varphi_2) D_x - i(\nabla_x \cdot \nabla_\xi p_2)(x, i\nabla\varphi_2) + r_{2,2}(x, i\nabla\varphi_2) - \rho_1) \alpha_{2,0}^{0,0} = 0. \tag{4-38}$$

Here we also have used the fact that $\rho \sim \sum_{k \geq 1} h^k \rho_k$ as $h \rightarrow 0$.

Identifying the other coefficients, one obtains a series of equations that (in a way similar to [Pettersson 1997, Section 4]) can be solved in \mathcal{V}_1 (possibly after having shrunk it a little bit around $x^{(1)}$), and in such a way that one also has

$$\tilde{w}_2 - \tilde{w}_1 = \mathcal{O}(h^\infty e^{-\varphi/h}) \quad \text{locally uniformly in } \mathcal{V}_1 \cap \{V_2 < V_1\}, \tag{4-39}$$

where w_1 is defined in (4-1) and \tilde{w}_1 and \tilde{w}_2 are resummations of w_1 and w_2 . Among other things, this implies

$$\alpha_{2,0}^{0,0} = a_{2,0} \quad \text{in } \mathcal{V}_1 \cap \{V_2 < V_1\}. \tag{4-40}$$

Moreover, we see in (4-36) and (4-38) that $\beta_{1,0}^{0,0}$ (respectively $\alpha_{2,0}^{0,0}$) is a solution of a differential equation of order 1 on each integral curve of the real vector field $\nabla\varphi_1(y) \cdot \nabla_y$ (respectively $\nabla\varphi_2(y) \cdot \nabla_y$). In particular, because of the ellipticity of $a_{2,0}$, we deduce from (4-38) and (4-40) that we have that

$$\alpha_{2,0}^{0,0} \quad \text{never vanishes in } \mathcal{V}_1. \tag{4-41}$$

Now, Assumption 6 implies that, if $\gamma \in G_0$, then

$$r_{1,2}(x, i\nabla\varphi_2) \neq 0 \quad \text{on } \mathcal{V}_1. \tag{4-42}$$

Since $p_1(y, i\nabla\varphi_2) = p_1(y, i\nabla\varphi_1) = 0$ on $\omega_1 \cap \partial\Omega$, we deduce from (4-35) and (4-41) that, if $\gamma \in G_0$, then $\beta_{1,0}^{0,0}$ does not vanish on $\omega_1 \cap \partial\Omega$. As before, because of (4-36) (and the fact that $R(x, hD_x)$ is formally self-adjoint), this implies:

$$\text{if } \gamma \in G_0, \text{ then } \beta_{1,0}^{0,0} \text{ never vanishes in } \mathcal{V}_1. \tag{4-43}$$

In the island, outside the cirque. Now, we look at what happens on $\gamma^{(2)}$ and, at first, near $x^{(1)}$. Using the asymptotics of $Y_{k,\varepsilon}(z/\sqrt{h})$ given in [Pettersson 1997, Section 4], one also finds that, in $\mathcal{V}_1 \cap \{V_1 < V_2\}$, w_2 can be formally identified with

$$w_3(x, h) = \sqrt{2\pi h} \begin{pmatrix} b_1(x, h) \\ hb_2(x, h) \end{pmatrix} e^{-\varphi(x)/h}, \tag{4-44}$$

where b_1, b_2 are symbols of the form

$$b_j(x; h) = \sum_{l \geq 0} \sum_{m=0}^l h^l (\ln h)^m b_j^{l,m}(x) \quad (j = 1, 2), \tag{4-45}$$

with $b_j^{l,m} \in C^\infty(\mathcal{V}_1 \cap \{V_1 < V_2\})$, in the sense that, for any resummations \tilde{w}_2 and \tilde{w}_3 of w_2 and w_3 ,

$$\tilde{w}_2 - \tilde{w}_3 = \mathcal{O}(h^\infty e^{-\varphi/h}) \quad \text{locally uniformly in } \Omega \cap \Gamma_+. \tag{4-46}$$

Moreover, one also has

$$b_1^{0,0} = \beta_{1,0}^{0,0}, \tag{4-47}$$

which, by (4-43), shows that, when $\gamma \in G_0$, b_1 is elliptic in $\mathcal{V}_1 \cap \{V_1 < V_2\}$.

Since $p_2(x, i\nabla\varphi(x)) \neq 0$ in $\{V_1 < V_2\}$, we can formally solve the equation $Pw_3 = \rho_1 w_3$, and we see again that b_1 and b_2 can be continued along the integral curves of $\nabla\varphi$, as long as these curves stay inside $\{V_1 < V_2\}$ and φ_1 does not develop caustics. In particular, they can be continued in a neighborhood \mathcal{N}_2 of $\gamma^{(2)}$, and the continuation of b_1 remains elliptic in Ω_2 .

Clearly, the previous steps can be repeated near $x^{(2)}, x^{(3)}$, etc. (in the case $N_\gamma \geq 3$), up to $x^{(N_\gamma+1)}$, obtaining in that way (after having pasted everything in a standard way by using a partition of unity) a function $\mathbf{w}(x, h)$, smooth on a neighborhood $\mathcal{N}(\gamma)$ of γ in \ddot{O} , satisfying

$$(P - \rho_1)\mathbf{w} = \mathcal{O}(h^\infty e^{-\varphi/h}),$$

locally uniformly in $\mathcal{N}(\gamma)$. Moreover, $\mathcal{N}(\gamma)$ can be decomposed into

$$\mathcal{N}(\gamma) = \mathcal{N}_1 \cup \mathcal{V}_1 \cup \dots \cup \mathcal{V}_{N_\gamma} \cup \mathcal{N}_{N_\gamma+1},$$

where, for all j , \mathcal{V}_j is a neighborhood of $x^{(j)}$ and \mathcal{N}_j is a neighborhood of $\gamma^{(j)}$, in such a way that, in each \mathcal{N}_j , \mathbf{w} admits a WKB asymptotics of the form,

$$\mathbf{w}(x; h) \sim h^{(j-1)/2} \begin{pmatrix} h^{(1-(-1)^j)/2} a_1^{(j)}(x, h) \\ h^{(1+(-1)^j)/2} a_2^{(j)}(x, h) \end{pmatrix} e^{-\varphi(x)/h}, \tag{4-48}$$

where $a_1^{(j)}$ and $a_2^{(j)}$ are symbols of the same form as in (4-45), and $a_1^{(j)}$ is elliptic if j is even, while $a_2^{(j)}$ is elliptic if j is odd (in particular, $a_1^{(N_\gamma+1)}$ is elliptic). On the other hand, in each \mathcal{V}_j , \mathbf{w} can be represented by means of the Weber function, in a way similar to that of (4-7).

At and after the boundary of the island. Let us denote by $x_\gamma \in \gamma \cap \partial \ddot{O}$ the point of type 1 where γ touches the boundary of the island. When $x \in \gamma \cap \ddot{O}$ is close enough to x_γ , we know from the previous subsection that the asymptotic solution w is of the form

$$w(x; h) \sim h^{N_\gamma/2} \begin{pmatrix} b_1(x, h) \\ hb_2(x, h) \end{pmatrix} e^{-\varphi(x)/h}, \tag{4-49}$$

where b_1, b_2 are smooth symbols on $\mathcal{N}_{N_\gamma+1}$ of the same form as in (4-45), and b_1 is elliptic. Moreover, as x approaches x_γ , b_1 and b_2 (together with φ) develop singularities on some set \mathcal{C} (called the caustic set). However, following an idea of [Helffer and Sjöstrand 1986], we can represent $h^{-N_\gamma/2}e^{S/h}w$ in the integral (Airy) form

$$I[c_1, c_2](x, h) = h^{-\frac{1}{2}} \int_{\gamma(x)} \begin{pmatrix} c_1(x', \xi_n, h) \\ hc_2(x', \xi_n, h) \end{pmatrix} e^{-(x_n \xi_n + g(x', \xi_n))/h} d\xi_n, \tag{4-50}$$

where we have used local Euclidean coordinates $(x', x_n) \in \mathbb{R}^{n-1} \times \mathbb{R}$ centered at $\gamma \cap \partial \ddot{O}$, such that $V_1(x) = -C_0 x_n + \mathcal{O}(x^2)$ near this point. For x in \ddot{O} close to $\gamma \cap \partial \ddot{O}$, the phase function $\xi_n \mapsto x_n \xi_n + g(x', \xi_n)$ admits two real critical points that are close to 0. Then, choosing conveniently the x -dependent interval $\gamma(x)$, the steepest descent method at one of these points gives us the asymptotic expansion of $I[c_1, c_2]$. Comparing this with the symbols b_1 and b_2 , one can determine c_1 and c_2 so that the asymptotic expansion of $h^{-N_\gamma/2}e^{S/h}w$ coincides with that of $I[c_1, c_2]$ in \ddot{O} . In particular, when $\gamma \in G_0$, one finds that c_1 remains elliptic near 0.

At this point, since we did not assume any analyticity of the potentials near \ddot{O} , we have to follow the methods of [Fujiié et al. 2011] — a reference we will henceforth abbreviate as [FLM] — where a similar situation is considered. Indeed, following the constructions of [FLM, Section 4] (that are made in the scalar case, but can be generalized without problem to our vectorial case), we see that there exists a constant $\delta > 0$ such that, for any $N \geq 1$, one can construct a (vectorial) function w_N , smooth on the set

$$\mathfrak{W}_N(\gamma) := \{|x - x_\gamma| < \varepsilon\} \cap \{\text{dist}(x, \ddot{O}) < 2(Nk)^{2/3}\} \tag{4-51}$$

with $\varepsilon > 0$ small enough (recall from (3-2) that $k = |h \ln h|$), such that [FLM, Propositions 4.5 and 4.6]:

- $(P - \rho_1)w_N = \mathcal{O}(h^{\delta N} e^{-\text{Re} \tilde{\varphi}_N/h})$ uniformly in $\mathfrak{W}_N(\gamma)$.
- For any $\alpha \in \mathbb{Z}_+^n$, there exists $m_\alpha \geq 0$ independent of N such that

$$\partial_x^\alpha w_N = \mathcal{O}(h^{-m_\alpha} e^{-\text{Re} \tilde{\varphi}_N/h})$$

uniformly in $\mathfrak{W}_N(\gamma)$.

- w_N can be represented by an integral of the form (4-50) (with $\gamma(x) = \gamma_N(x)$ depending on N) in all of $\mathfrak{W}_N(\gamma)$.
- $w_N = w$ in $\mathcal{N}_{N_\gamma+1} \cap \mathfrak{W}_N(\gamma)$.

- For any large enough L , there exist $C_L > 0$ and $\delta_L > 0$, both independent of N , such that, uniformly in ${}^{\mathfrak{W}}\mathcal{W}_N(\gamma) \cap \{\text{dist}(x, \ddot{O}) \geq (Nk)^{2/3}\}$, one has

$$w_N(x, h) = h^{N_\gamma/2} \left(\sum_{\substack{\ell=0 \\ 0 \leq m \leq \ell}}^{L+[Nk/C_L h]} h^\ell (\ln h)^m \begin{pmatrix} f_{1,N}^{\ell,m}(x) \\ h f_{2,N}^{\ell,m}(x) \end{pmatrix} + \mathcal{O}(h^{\delta_L N} + h^L) \right) e^{-\tilde{\varphi}_N(x)/h} \quad \text{as } h \rightarrow 0, \quad (4-52)$$

with $f_{1,N}^{\ell,m}(x), f_{2,N}^{\ell,m}(x)$ independent of h and of the form

$$\tilde{f}_{j,N}^{\ell,m}(x) = (\text{dist}(x, \mathcal{C}))^{-3\ell/2-1/4} \beta_{j,N}^{\ell,m}(x, \text{dist}(x, \mathcal{C})), \quad j = 1, 2, \quad (4-53)$$

where $\beta_{j,N}^{\ell,m}$ is smooth near $(x_\gamma, 0)$, and $\beta_1^{\ell,m}(x_\gamma, 0) \neq 0$ in the case $\gamma \in G_0$.

Here, $\tilde{\varphi}_N$ is a (complex-valued) C^1 function on ${}^{\mathfrak{W}}\mathcal{W}_N(\gamma)$, smooth on ${}^{\mathfrak{W}}\mathcal{W}_N(\gamma) \setminus \mathcal{C}$, such that [FLM, Lemma 4.1]:

- $\tilde{\varphi}_N = \varphi + \mathcal{O}(h^\infty)$ uniformly in $\mathcal{N}_{N_\gamma+1} \cap {}^{\mathfrak{W}}\mathcal{W}_N(\gamma)$.
- $(\nabla \tilde{\varphi}_N)^2 = V_1(x) + \mathcal{O}(h^\infty)$ uniformly in ${}^{\mathfrak{W}}\mathcal{W}_N(\gamma)$.
- There exists $\varepsilon(h) = \mathcal{O}(h^\infty)$ real such that, for $x \in {}^{\mathfrak{W}}\mathcal{W}_N(\gamma) \setminus \ddot{O}$, one has

$$\text{Re } \tilde{\varphi}_N(x) \geq S - \varepsilon(h). \quad (4-54)$$

- One has

$$\text{Im } \nabla \varphi_N(x) = -\nu_N(x) \sqrt{\text{dist}(x, \mathcal{C})} \nabla \text{dist}(x, \mathcal{C}) + \mathcal{O}(\text{dist}(x, \mathcal{C})),$$

uniformly with respect to $h > 0$ small enough and $x \in {}^{\mathfrak{W}}\mathcal{W}_N(\gamma) \setminus \ddot{O}$ with $\nu_N(x) \geq \delta$.

The previous results show that we can extend \mathbf{w} by taking w_N in ${}^{\mathfrak{W}}\mathcal{W}_N(\gamma)$, and we obtain in that way a function \mathbf{w}_N smooth on $\mathcal{N}(\gamma) \cup {}^{\mathfrak{W}}\mathcal{W}_N(\gamma)$, such that $(P - \rho_1)\mathbf{w}_N = \mathcal{O}(h^{\delta N} e^{-\text{Re } \tilde{\varphi}_N/h})$ uniformly in $\mathcal{N}(\gamma) \cup {}^{\mathfrak{W}}\mathcal{W}_N(\gamma)$. Note that, thanks to Assumption 4, the number N_γ is constant on each connected component of Γ .

5. Agmon estimates

Preliminaries. In order to perform Agmon estimates in the same spirit as in [Helffer and Sjöstrand 1984], we need some preliminary results because of the fact that we have to deal with pseudodifferential operators (and not only Schrödinger operators). For this reason, we prefer to work with C^∞ weight functions (instead of Lipschitz ones), and the idea is to take h -dependent regularizations of Lipschitz weights.

At first, we need:

Proposition 5.1. *Let $\nu_0 > 0, m \geq 0, a = a(x, \xi) \in S_{\nu_0}(\langle \xi \rangle^{2m})$. For $h > 0$ small enough, let also $\Phi_h \in C^\infty(\mathbb{R}^n)$ be real-valued, such that*

$$\sup |\nabla \Phi_h| < \nu_0 \quad (5-1)$$

and, for any multi-index $\alpha \in \mathbb{N}^n$ with $|\alpha| \geq 2$,

$$\partial^\alpha \Phi_h(x) = \mathcal{O}(h^{1-|\alpha|}), \quad (5-2)$$

uniformly for $x \in \mathbb{R}^n$ and $h > 0$ small enough. Then, for any $\tilde{\Sigma} \subset \mathbb{R}^n$ with $\text{dist}(\Sigma, \mathbb{R}^n \setminus \tilde{\Sigma}) > 0$, the operator $e^{\Phi_h/h} A e^{-\Phi_h/h} := e^{\Phi_h/h} \text{Op}_h^W(a) e^{-\Phi_h/h}$ satisfies

$$\|e^{\Phi_h/h} A e^{-\Phi_h/h} u\|_{L^2} \leq C_1 \| \langle h D_x \rangle^m u \|_{L^2}, \tag{5-3}$$

uniformly for all $h > 0$ small enough and $u \in H^m(\mathbb{R}^n)$.

Proof. For $u \in C_0^\infty(\mathbb{R}^n)$, we write

$$e^{\Phi/h} A e^{-\Phi/h} u(x) = \frac{1}{(2\pi h)^n} \int e^{i(x-y)\xi/h + (\Phi(x) - \Phi(y))/h} a\left(\frac{x+y}{2}, \xi\right) u(y) dy d\xi,$$

and the property (5-1) shows that we can make the change of contour of integration given by

$$\mathbb{R}^n \ni \xi \mapsto \xi + i\Psi(x, y),$$

where $\Psi(x, y) := \int_0^1 \nabla \Phi((1-t)x + ty) dt$ (in particular, one has $\Phi(x) - \Phi(y) = (x - y)\Psi(x, y)$). Then, denoting by Op_h the semiclassical quantization of symbols depending on $3n$ variables (see, e.g., [Martinez 2002, Section 2.5]), we obtain

$$e^{\Phi/h} A e^{-\Phi/h} = \text{Op}_h\left(a\left(\frac{x+y}{2}, \xi + i\Psi(x, y)\right)\right),$$

and, using (5-2), we see that, for any $\alpha, \beta, \gamma \in \mathbb{Z}_+^n$, we have

$$\partial_x^\alpha \partial_y^\beta \partial_\xi^\gamma \left(a\left(\frac{x+y}{2}, \xi + i\Psi(x, y)\right) \right) = \mathcal{O}(h^{-|\alpha+\beta|} \langle \xi \rangle^m). \tag{5-4}$$

Then, the result is an easy consequence of the Calderón–Vaillancourt Theorem; see [Martinez 2002, Exercise 2.10.15], for example. □

Proposition 5.2. *Let ϕ and V be two bounded real-valued Lipschitz functions on \mathbb{R}^n with $|\nabla \phi(x)|^2 \leq V(x)$ almost everywhere. Let also $\chi_1 \in C_0^\infty(\mathbb{R}^n; [0, 1])$ be supported in the ball $\{|x| \leq 1\}$, with $\int \chi_1(x) dx = 1$. For any $h > 0$, we set $\chi_h(x) = h^{-n} \chi(x/h)$. Then, the smooth function*

$$\phi_h := \chi_h * \phi$$

(where $*$ stands for the standard convolution) satisfies:

- $\phi_h = \phi + \mathcal{O}(h)$ uniformly for $h > 0$ small enough and $x \in \mathbb{R}^n$.
- For all $x \in \mathbb{R}^n$, one has $|\nabla \phi_h(x)|^2 \leq V(x) + h \|\nabla V\|_{L^\infty}$.
- For all $\alpha \in \mathbb{Z}_+^n$ with $|\alpha| \geq 1$, one has $\partial^\alpha \phi_h = \mathcal{O}(h^{1-|\alpha|})$.

The proof of this proposition is very standard and almost obvious, and we leave it to the reader. Observe that, in particular, ϕ_h satisfies the estimates (5-2).

Agmon estimates. As a corollary of the two previous propositions, we have:

Corollary 5.3. *Let ϕ and ϕ_h be as in Proposition 5.2, with $V = \min(V_1, V_2)_+$. Then one has, for any $u = (u_1, u_2) \in H^2(\mathbb{R}^n) \oplus H^2(\mathbb{R}^n)$,*

$$\begin{aligned} & \operatorname{Re}\langle e^{\phi_h/h} P u, e^{\phi_h/h} u \rangle \\ & \geq \|h\nabla(e^{\phi_h/h} u)\|^2 + \sum_{j=1}^2 \langle (V_j - |\nabla\phi_h|^2)e^{\phi_h/h} u_j, e^{\phi_h/h} u_j \rangle - C_R h (\|e^{\phi_h/h} u\|^2 + \|h\nabla(e^{\phi_h/h} u)\|^2), \end{aligned}$$

where $C_R > 0$ is a constant that depends on $R(x, hD_x)$, χ_1 and $\sup |\nabla\phi|$ only.

Proof. It is standard (and elementary) to show that

$$\operatorname{Re}\langle e^{\phi_h/h} (-h^2\Delta + V_j)u_j, e^{\phi_h/h} u_j \rangle = \|h\nabla(e^{\phi_h/h} u_j)\|^2 + \langle (V_j - |\nabla\phi_h|^2)e^{\phi_h/h} u_j, e^{\phi_h/h} u_j \rangle.$$

Therefore, it is enough to estimate $\langle e^{\phi_h/h} R(x, hD_x)u, e^{\phi_h/h} u \rangle$. Applying Proposition 5.1, we see that the operator $e^{\phi_h/h} R(x, hD_x)e^{-\phi_h/h} \langle hD_x \rangle^{-1}$ is uniformly bounded on L^2 .

Moreover, since the constants appearing in the estimates (5-4) depend on a, α , and on the estimates on the $\partial^\beta \Phi$ only, we see that the norm of $e^{\phi_h/h} R(x, hD_x)e^{-\phi_h/h} \langle hD_x \rangle^{-1}$ depends on r and on estimates on $\partial^\beta (\chi_h * \nabla\phi) = (\partial^\beta \chi_h) * \nabla\phi$ ($|\beta| \leq |\alpha|$) only. Since the latter depend on α, χ_1 and $\sup |\nabla\phi|$ only, the result follows. \square

6. Global asymptotic solution

The constructions of Section 4 can be done in a neighborhood of any minimal geodesic $\gamma \in G$, and give rise (after having pasted them together with a partition of unity) to an asymptotic solution (still denoted by w_N) on a neighborhood of $\bigcup_{\gamma \in G} \gamma$. Now, we plan to extend this solution to a whole (h -dependent) neighborhood of $\{V_1 \geq 0\}$, by using a modified self-adjoint operator with discrete spectrum near 0.

At first, we fix $\varepsilon_0 > 0$ sufficiently small, and a cutoff function $\chi_0 \in C_0^\infty(\ddot{O}; [0, 1])$ such that

$$\chi_0(x) = 1 \quad \text{if } V_1(x) \geq 2\varepsilon_0, \quad \chi_0(x) = 0 \quad \text{if } V_1(x) \leq \varepsilon_0,$$

and we set

$$\tilde{V}_1 := \chi_0 V_1 + \varepsilon_0(1 - \chi_0). \tag{6-1}$$

In particular, \tilde{V}_1 coincides with V_1 on $\{V_1 \geq 2\varepsilon_0\}$, and we have $\tilde{V}_1 \geq \varepsilon_0$ everywhere. Then, we define $\tilde{P}_1 := -h^2\Delta + \tilde{V}_1$, and we consider the self-adjoint operator

$$\tilde{P} = \begin{pmatrix} \tilde{P}_1 & 0 \\ 0 & P_2 \end{pmatrix} + hR(x, hD_x). \tag{6-2}$$

By construction, for all $C > 0$ and h small enough, the spectrum of \tilde{P} is discrete in $[-Ch, Ch]$, and a straightforward adaptation of the arguments used in [Helffer and Sjöstrand 1984] shows that its first eigenvalue E_1 admits the same asymptotics as ρ_1 as $h \rightarrow 0_+$. We denote by v its first normalized eigenfunction, and by $\mathcal{N}_0 \subset \{V_1 > 2\varepsilon_0\}$ some fixed neighborhood of $\bigcup_{\gamma \in G} \gamma \cap \{V_1 > 2\varepsilon_0\}$ where the asymptotic solution w_N is well-defined. We have:

Proposition 6.1. *There exists $\theta_0 \in \mathbb{R}$ independent of h such that, for any compact subset K of \mathcal{N}_0 , and for any $\alpha \in \mathbb{Z}_+^n$, one has*

$$\|e^{\varphi/h} \partial^\alpha (e^{i\theta_0} \mathbf{v} - h^{n/4} \mathbf{w}_N)\|_K = \mathcal{O}(h^\infty).$$

Proof. The existence of θ_0 such that $\partial^\alpha (e^{i\theta_0} \mathbf{v} - h^{n/4} \mathbf{w}_N) = \mathcal{O}(h^\infty)$ uniformly near 0 is a consequence of [Helffer and Sjöstrand 1984, Proposition 2.5] and standard Sobolev estimates. Let $\chi \in C_0^\infty(\mathcal{N}_0; [0, 1])$, with $\chi = 1$ in a neighborhood of $K \cup \{0\}$. Following [Helffer and Sjöstrand 1984; Pettersson 1997], we plan to apply Corollary 5.3 to $u := \chi (e^{i\theta_0} \mathbf{v} - h^{n/4} \mathbf{w}_N)$, with a suitable weight function ϕ . Let us first observe that, using Corollary 5.3, for any $\varepsilon > 0$ one has

$$\|e^{(1-\varepsilon)\tilde{\varphi}/h} \langle hD_x \rangle \mathbf{v}\|_{H^1} = \mathcal{O}(1), \tag{6-3}$$

where $\tilde{\varphi}(x) \geq \varphi(x)$ is the Agmon distance associated with $\min(\tilde{V}_1, V_2)$ between 0 and x . Now, for $C \geq 1$ arbitrarily large, we define

$$\phi(x) := \min(\phi_1, \phi_2),$$

where

$$\phi_1(x) := \begin{cases} \varphi(x) - Ch \ln\left(\frac{\varphi(x)}{h}\right) & \text{if } \varphi(x) \geq Ch, \\ \varphi(x) - Ch \ln C & \text{if } \varphi(x) \leq Ch, \end{cases}$$

$$\phi_2(x) := \begin{cases} \inf_{\chi(y) \neq 1} (1 - 2\varepsilon)(\varphi(y) + d(y, x)) & \text{if } x \in \text{supp } \chi, \\ (1 - 2\varepsilon)\varphi(x) & \text{if } x \notin \text{supp } \chi. \end{cases}$$

Here, $\varepsilon > 0$ is taken sufficiently small to have $\phi_2(x) > \varphi(x)$ when $x \in K$. Then, ϕ is Lipschitz continuous, and one has $\phi = \phi_1$ on K and $\phi = \phi_2$ on $\mathbb{R}^n \setminus \{\chi = 1\}$. Moreover, one sees as in the proof of [Pettersson 1997, Theorem 5.5] that, if we set $V := \min(V_1, V_2)$, ϕ satisfies

$$|\nabla\phi|^2 = V \quad \text{in } \{\varphi \leq Ch\}, \quad |\nabla\phi|^2 \leq V - \delta_0 Ch \quad \text{in } \{\varphi \geq Ch\},$$

where $\delta_0 = \inf_{x \in \text{supp } \chi, x \neq 0} (V(x)/\varphi(x)) > 0$. As a consequence, by Proposition 5.2, the regularized ϕ_h of ϕ satisfies

$$|\nabla\phi_h|^2 \leq V + h\|V\|_{L^\infty} \quad \text{in } \{\varphi \leq Ch\}, \quad |\nabla\phi_h|^2 \leq V - (\delta_0 C - \|V\|_{L^\infty})h \quad \text{in } \{\varphi \geq Ch\}.$$

Then, choosing C sufficiently large and setting $u := \chi (e^{i\theta_0} \mathbf{v} - h^{n/4} \mathbf{w}_N)$, we see that Corollary 5.3 implies

$$\|h\nabla(e^{\phi_h/h} u)\|^2 + C'h\|e^{\phi_h/h} u\|_{\{\varphi \geq Ch\}}^2 \leq \langle e^{\phi_h/h} (\tilde{P} - E_1)u, e^{\phi_h/h} u \rangle, \tag{6-4}$$

with $C' = C'(C)$ arbitrarily large. Moreover, if $\tilde{\chi} \in C_0^\infty(\mathcal{N}_0)$ is such that $\tilde{\chi}\chi = \chi$, we have

$$(\tilde{P} - E_1)u = [\tilde{P}, \tilde{\chi}]\tilde{\chi}u + \mathcal{O}(h^\infty e^{-\varphi/h}),$$

and since $\phi_h = (1 - 2\varepsilon)\varphi + \mathcal{O}(h)$ on $\text{supp } \nabla\chi$, $\min_{\text{supp } \nabla\chi} \varphi =: \delta_1 > 0$ and, by Proposition 5.1, the operator $e^{\phi_h/h} [R, \chi] e^{-\phi_h/h}$ is uniformly bounded, we obtain, using also (6-3),

$$\langle e^{\phi_h/h} (\tilde{P} - E_1)u, e^{\phi_h/h} u \rangle = \mathcal{O}(\|e^{(1-\varepsilon)\varphi/h} \langle hD_x \rangle \tilde{\chi}u\|_{\text{supp } \nabla\chi}^2 + h\|e^{\phi_h/h} u\|^2) = \mathcal{O}(e^{-2\varepsilon\delta_1/h} + h\|e^{\phi_h/h} u\|^2).$$

Inserting this estimate into (6-4) and taking C sufficiently large, this permits us to obtain

$$\|h\nabla(e^{\phi_h/h}u)\|^2 + h\|e^{\phi_h/h}u\|^2 = \mathcal{O}(e^{-2\varepsilon\delta_1/h} + \|e^{\phi_h/h}u\|_{\{\varphi \leq Ch\}}^2).$$

In particular, since $\phi_h = \phi_1 + \mathcal{O}(h)$ on K and $\phi_h = (1 - 2\varepsilon)\varphi \leq Ch$ on $\{\varphi \leq Ch\}$,

$$\|h^C \varphi^{-C} e^{\varphi/h} h \nabla u\|_K^2 + \|h^C \varphi^{-C} e^{\varphi/h} u\|_K^2 = \mathcal{O}(e^{-2\varepsilon\delta_1/h} + \|u\|_{\{\varphi \leq Ch\}}^2).$$

Therefore,

$$\|e^{\varphi/h} \nabla u\|_K^2 + \|e^{\varphi/h} u\|_K^2 = \mathcal{O}(h^\infty),$$

and the result follows by standard Sobolev estimates. □

Now, following [FLM, Section 4.3], we observe that, if ε_0 has been taken small enough, the asymptotic solution \mathbf{w}_N is $\mathcal{O}(h^{\delta N} e^{-S/h})$ uniformly on the set

$$\left\{ \text{dist}\left(x, \bigcup_{\gamma \in G} \gamma\right) \geq \varepsilon_0 \right\} \cap \{V_1 \leq 2\varepsilon_0\} \cap \left(\bigcup_{\gamma \in G} \mathcal{N}(\gamma) \cup \mathcal{W}_N(\gamma) \right)$$

Moreover, by (6-3), the same is true for \mathbf{v} on $\{\text{dist}(x, \bigcup_{\gamma \in G} \gamma) \geq \varepsilon_0\} \cap \{V_1 \leq 2\varepsilon_0\}$. Therefore, using also Proposition 6.1, we can paste together $e^{i\theta_0} \mathbf{v}$ and $h^{-n/4} \mathbf{w}_N$ in order to obtain a function \mathbf{u}_N that satisfies the properties of the following proposition; see also [FLM, Proposition 4.6].

Proposition 6.2. *There exists a function \mathbf{u}_N , smooth on $\ddot{O}_N := \{\text{dist}(x, \ddot{O}) < 2(Nk)^{2/3}\}$, such that*

$$(P - \rho)\mathbf{u}_N = \mathcal{O}(h^{\delta N} e^{-\text{Re } \varphi_N/h}), \quad \partial^\alpha \mathbf{u}_N = \mathcal{O}(h^{-m_\alpha} e^{-\text{Re } \varphi_N/h}),$$

uniformly on \ddot{O}_N , where $\tilde{\varphi}_N$ is as in (4-54). Moreover, in $\bigcup_{\gamma \in G} \mathcal{W}_N(\gamma) \cap \{\text{dist}(x, \ddot{O}) \geq (Nk)^{2/3}\}$ one can write \mathbf{u}_N as in (4-52) (with $\beta_1^{\ell,m}(x,\gamma, 0) \neq 0$), while away from $\bigcup_{\gamma \in G} \mathcal{W}_N(\gamma) \cap \{x \notin \ddot{O}\}$, \mathbf{u}_N is $\mathcal{O}(h^{\delta N} e^{-\text{Re } \varphi_N/h})$.

7. Comparison between asymptotic and true solution

A priori estimates. In the same spirit as in [FLM, Theorem 2.2], we start with an a priori estimate for the resonant state of P . From now on, we denote by \mathbf{u} the outgoing solution of

$$P\mathbf{u} = \rho_1 \mathbf{u}, \tag{7-1}$$

normalized in the following way: we fix some analytic distorted space (also more recently introduced, in the context of computational physics, under the name of perfectly matched layer; see [Berenger 1994], for example) of the form

$$\tilde{\mathbb{R}}_\theta^n := \{x + i\theta F(x) : x \in \mathbb{R}^n\}, \tag{7-2}$$

where $F \in C^\infty(\mathbb{R}^n; \mathbb{R}^n)$, $F(x) = 0$ if $|x| \leq R_0$, $F(x) = x$ for $|x|$ large enough, and where $\theta > 0$ is sufficiently small and may also tend to 0 with h , but not too rapidly (here, we take $\theta = h|\ln h| = k$). Then, by definition, the fact that ρ_1 is a resonance of P means that (7-1) admits a solution in $L^2(\tilde{\mathbb{R}}_\theta^n)$, and here we take \mathbf{u} such that

$$\|\mathbf{u}\|_{L^2(\tilde{\mathbb{R}}_\theta^n)} = 1. \tag{7-3}$$

As before, d stands for the Agmon distance associated with the pseudometric $\min(V_1, V_2)_+ dx^2$, and we denote by $B_d(S) := \{x \in \mathbb{R}^n : d(0, x) < S\}$ the corresponding open ball of radius $S = d(0, \partial\mathring{O})$. Then, we first have:

Proposition 7.1. *For any compact subset $K \subset \mathbb{R}^n$, there exists $N_K \geq 0$ such that*

$$\|e^{s(x)/h} \mathbf{u}\|_{H^1(K)} = \mathcal{O}(h^{-N_K}),$$

uniformly as $h \rightarrow 0$, where $s(x) = \varphi(x)$ if $x \in B_d(S)$ and $s(x) = S$ otherwise.

Proof. The proof is very similar to that of [FLM, Theorem 2.2], with the only difference that here we have to deal with pseudodifferential operators, forbidding us to use Dirichlet realizations and nonsmooth weight functions. Instead, we modify V_1 in a way similar to (6-1), and we regularize the weights as in Proposition 5.2.

We consider a cutoff function $\hat{\chi}$ (dependent on h) such that

$$\hat{\chi}(x) = 1 \quad \text{if } V_1(x) \geq 2k^{2/3}, \quad \hat{\chi}(x) = 0 \quad \text{if } V_1(x) \leq k^{2/3}, \quad \partial^\alpha \hat{\chi} = \mathcal{O}(k^{-2|\alpha|/3}),$$

and we set

$$\hat{V}_1 := \hat{\chi} V_1 + k^{2/3}(1 - \hat{\chi}), \quad \hat{P}_1 := -h^2 \Delta + \hat{V}_1, \quad \hat{P} = \begin{pmatrix} \hat{P}_1 & 0 \\ 0 & P_2 \end{pmatrix} + hR(x, hD_x). \quad (7-4)$$

We denote by \hat{E} the first eigenvalue of \hat{P} , and by \hat{v} its first normalized eigenfunction. Moreover, we consider the Agmon distance \hat{d} associated with the pseudometric $(\min(V_1, V_2)_+ - \hat{E}) dx^2$, and we set $\hat{\varphi}(x) := \hat{d}(0, x)$. Then, the same proof as in [FLM, Lemma 3.1] shows the existence of a constant $C_1 > 0$ such that

$$s(x) - C_1 k \leq \hat{\varphi}(x) \leq \varphi(x) \quad (x \in \mathbb{R}^n). \quad (7-5)$$

Moreover, an adaptation of the proof of [FLM, Lemma 3.2] (obtained by using Proposition 5.2 in order to regularize the Lipschitz weight) gives

$$\|e^{\hat{\varphi}/h} \hat{v}\|_{H^1(\mathbb{R}^n)} = \mathcal{O}(h^{-N_0}), \quad (7-6)$$

for some $N_0 \geq 0$. Then, the result follows by considering the function $\hat{\chi} \hat{v}$ and by observing that, thanks to (7-6), one has [FLM, Lemma 3.3 and Formula (3.20)]

$$\left\| \hat{\chi} \hat{v} - \frac{1}{2i\pi} \int_\gamma (z - P_\theta)^{-1} \hat{\chi} \hat{v} dz \right\|_{H^1} = \mathcal{O}(h^{-N_1} e^{-S/h}).$$

Here, γ is the oriented complex circle $\{z \in \mathbb{C} : |z - \hat{E}| = h^2\}$ and P_θ is a convenient distortion of P . The previous estimate actually shows that the distorted \mathbf{u}_θ of \mathbf{u} coincides — up to $\mathcal{O}(h^{-N_1} e^{-S/h})$ — with $\mu \hat{\chi} \hat{v}$, where μ is a complex constant satisfying $|\mu| = 1 + \mathcal{O}(e^{-\delta/h})$, for some $\delta > 0$. □

Remark 7.2. The previous proof also gives a global estimate on \mathbf{u}_θ ,

$$\|e^{s(x)/h} \mathbf{u}_\theta\|_{H^1(\mathbb{R}^n)} = \mathcal{O}(h^{-N'_1}),$$

for some constant $N'_1 \geq 0$. See [FLM, Lemma 3.3 and Formula (3.20)].

Now, we plan to give an even better a priori estimate on the difference $\mathbf{u} - \mathbf{u}_N$ near the boundary of the island. Here again, we follow the arguments given in [FLM, Section 5]. For any $N \geq 1$, we set

$$U_N := \{x \in \mathbb{R}^n : \text{dist}(x, \partial \ddot{O}) < 2(Nk)^{2/3}\}.$$

We have [FLM, Propositions 5.1 and 5.2]:

Proposition 7.3. *There exist $N_1 \geq 0$ and $C \geq 1$ such that, for any $N \geq 1$ large enough, one has*

$$\|\mathbf{u} - \mathbf{u}_{CN}\|_{H^1(U_N)} \leq h^{-N_2} e^{-S/h}.$$

Proof. We just recall the main lines of the proof in [FLM]. At first, thanks to Proposition 7.1 and the particular form of \mathbf{u}_{CN} , we immediately see that the estimate is true on the set $\{\varphi(x) \geq S - 2k\}$. Then, we take a cutoff function $\tilde{\chi} \in C_0^\infty(\varphi(x) < S - k)$ such that $\tilde{\chi} = 1$ on $\{\varphi(x) \geq S - 2k\}$ and $\partial^\alpha \tilde{\chi} = \mathcal{O}(h^{-N_\alpha})$ for some $N_\alpha \geq 0$. We also consider the Lipschitz weight

$$\phi_N(x) = \min(\varphi(x) + C_1 Nk + k(S - \varphi(x))^{1/3}, S + (1 - k^{1/3})(S - \varphi(x)))$$

and, by using Propositions 7.1 and 6.2, we see that, if C is large enough, we have

$$\|e^{\phi_N/h} (P - \rho_1) \tilde{\chi}(\mathbf{u} - \mathbf{u}_{CN})\|_{L^2(\mathbb{R}^n)} = \mathcal{O}(h^{-M_1}),$$

for some $M_1 \geq 0$ independent of N . Then, regularizing ϕ_N as in Proposition 5.2, we can perform Agmon estimates as in the proof of [FLM, Proposition 5.1], and we find

$$\|e^{\phi_N/h} \tilde{\chi}(\mathbf{u} - \mathbf{u}_{CN})\|_{L^2(\mathbb{R}^n)} = \mathcal{O}(h^{-M_2}),$$

for some $M_1 \geq 0$ independent of N , and the result follows. □

Propagation. Now, we plan to prove (see [FLM, Proposition 6.1]):

Theorem 7.4. *For any $L > 0$ and for any $\alpha \in \mathbb{Z}_+^n$, there exists $N_{L,\alpha} \geq 1$ such that, for any $N \geq N_{L,\alpha}$,*

$$\partial_x^\alpha (\mathbf{u} - \mathbf{u}_{CN})(x, h) = \mathcal{O}(h^L e^{-S/h}) \quad \text{as } h \rightarrow 0, \tag{7-7}$$

uniformly in U_N .

Proof. As in [FLM], the proof relies on three different types of microlocal propagation arguments. We fix some $\hat{x} \in \partial \ddot{O}$ and we define the Fourier–Bors–Iagolnitzer transform T (see [Sjöstrand 1982; Martinez 2002], for example) as

$$Tu(x, \xi; h) := \int_{\mathbb{R}^n} e^{i(x-y)\xi/h - (x-y)^2/2h} u(y) dy.$$

(1) *Standard C^∞ propagation.* Since \mathbf{u} is outgoing (that is, it becomes L^2 when restricted to the distorted space or the perfectly matched layer defined in (7-2)), one can see as in [FLM, Lemma 6.2] that, if $t_0 > 0$ is large enough, one has

$$T\mathbf{u}(x, \xi) = \mathcal{O}(h^\infty e^{-S/h}),$$

uniformly near $\exp(-t_0 H_{p_1})(\hat{x}, 0)$. Moreover, by Proposition 7.1, we know that $e^{S/h} \mathbf{u}$ remains $\mathcal{O}(h^{-N_0})$ (for some $N_0 \geq 0$) on a neighborhood of the x -projection of $\{\exp(-t H_{p_1})(\hat{x}, 0) : 0 < t \leq t_0\}$.

Then, the standard C^∞ propagation of the frequency set for the solution to a real principal type operator (see, e.g., [Martinez 2002]) shows that the previous estimate remains valid near $\exp(-tH_{p_1})(\hat{x}, 0)$ for any $t > 0$.

(2) *Nonstandard propagation in h -dependent domains.* Thanks to the previous result, we can concentrate our attention on a sufficiently small neighborhood of \hat{x} . As before, we choose local Euclidean coordinates $(x', x_n) \in \mathbb{R}^{n-1} \times \mathbb{R}$ centered at \hat{x} , such that $V_1(x) = -C_0x_n + \mathcal{O}(|x - \hat{x}|^2)$. We also set $\mu_N := (Nk)^{-\frac{1}{3}}$, and we consider the modified Fourier–Bors–Iagolnitzer transform T_N defined by

$$T_N u(x, \xi; h) := \int_{\mathbb{R}^n} e^{i(x-y)\xi/h - (x'-y')^2/2h - \mu_N(x_n - y_n)^2/2h} u(y) dy. \tag{7-8}$$

Then, using the previous result it is elementary to show that, for any (fixed) $t > 0$ small enough, one has [FLM, Lemma 6.3]

$$T_N \mathbf{1}_{K_1} u(x, \xi) = \mathcal{O}(h^\infty e^{-S/h}),$$

uniformly near $\exp(-tH_{p_1})(\hat{x}, 0)$. Here K_1 is of the form $K_1 = K \setminus B_d(S)$, where K is any compact neighborhood of the closure of \ddot{O} . The interest of the latter property is that, as shown in [FLM], it can be propagated up h -dependent times t of order $(Nk)^{1/3}$. More precisely, setting

$$\exp(tH_{p_1})(\hat{x}, 0) = (x'(t), x_n(t); \xi'(t), \xi_n(t)) \quad (t \in \mathbb{R}),$$

we have [FLM, Lemma 6.4]:

Lemma 7.5. *There exists $\delta_0 > 0$ such that, for any $\delta \in (0, \delta_0]$, for all $N \geq 1$ large enough, and for $t_{N,\delta} := \delta^{-1}(Nk)^{1/3}$, one has*

$$\mathbf{T}_N \mathbf{1}_{K_1} u = \mathcal{O}(h^{\delta N} e^{-S/h}) \quad \text{uniformly in } \mathcal{W}(t_N, h),$$

where

$$\mathcal{W}_\delta(N, h) := \left\{ |x_n - x_n(-t_{N,\delta})| \leq \delta(Nk)^{2/3}, |\xi_n - \xi_n(-t_{N,\delta})| \leq \delta(Nk)^{1/3}, |x' - x'(-t_{N,\delta})| \leq \delta(Nk)^{1/3}, |\xi' - \xi'(-t_{N,\delta})| \leq \delta(Nk)^{1/3} \right\}.$$

Proof. The proof is based on the refined exponential weighted estimates (in the same spirit as in [Martinez 2002]) given in [FLM, Proposition 8.3], which we apply here to the operator P_1 . Since the proof is very similar to that of [FLM, Lemma 6.4], we omit the details. □

On the other hand, using the explicit form of u_{CN} given in (4-52), one also sees that, for any L large enough, there exists $\delta_L > 0$ such that, for any $N \geq 1$, one has [FLM, Lemma 6.7]:

$$T_N \mathbf{1}_{K_1} u_{CN} = \mathcal{O}((h^{\delta_L N} + h^L) e^{-S/h}) \quad \text{uniformly in } \mathcal{W}_\delta(N, h).$$

In particular, taking $N = L/\delta_L$ with $L \gg 1$, we obtain a sequence $N = N_L$ along which

$$T_N \mathbf{1}_{K_1} u_{CN} = \mathcal{O}(h^{\delta_L N} e^{-S/h}) \quad \text{uniformly in } \mathcal{W}_\delta(N, h),$$

and with both N and $\delta_L N$ arbitrarily large.

As a consequence, along the same sequence, we also obtain

$$T_N \mathbf{1}_{K_1}(\mathbf{u} - \mathbf{u}_{CN}) = \mathcal{O}(h^{\delta'_L N} e^{-S/h}) \quad \text{uniformly in } \mathcal{W}_\delta(N, h),$$

with $\delta'_L = \min(\delta, \delta_L)$.

Moreover we see that, when $y \in U_N \cap B_d(S)$ and $x \in \Pi_x \mathcal{W}_\delta(N, h)$ (where Π_x stands for the natural projection onto the x -space), we have

$$\mu_N(x_n - y_n)^2 + s(x) - S \geq C_\delta Nk,$$

with $C_\delta > 0$ constant (and actually $C_\delta \rightarrow \infty$ as $\delta \rightarrow 0$). Therefore, using [Proposition 7.3](#) and the expression (7-8) for T_N , we also obtain

$$T_N \mathbf{1}_{U_N \cap B_d(S)}(\mathbf{u} - \mathbf{u}_{CN}) = \mathcal{O}(h^{\delta N} e^{-S/h}) \quad \text{uniformly in } \mathcal{W}_\delta(N, h).$$

As a consequence, if we set

$$\chi_N(x) := \chi_0\left(\frac{|x_n - \hat{x}_n|}{(Nk)^{2/3}}\right) \chi_0\left(\frac{|x' - \hat{x}'|}{(Nk)^{1/2}}\right), \tag{7-9}$$

where the function $\chi_0 \in C_0^\infty(\mathbb{R}_+; [0, 1])$ satisfies $\chi_0 = 1$ in a sufficiently large neighborhood of 0, and is fixed in such a way that $\chi_N(x) = 1$ in

$$\{|x_n - \hat{x}_n| \leq |x_n(-t_N) - \hat{x}_n| + 2\delta(Nk)^{2/3}, |x' - \hat{x}'| \leq |x'(-t_N) - \hat{x}'| + 2\delta(Nk)^{1/2}\}$$

(here, t_N and δ are those of [Lemma 7.5](#)), then the function

$$v_N := \chi_N e^{S/h}(\mathbf{u} - \mathbf{u}_{CN})$$

is such that

$$T_N v_N = \mathcal{O}(h^{\delta'_L N} e^{-S/h}) \quad \text{uniformly in } \mathcal{W}_\delta(N, h). \tag{7-10}$$

Moreover, we have [\[FLM, Section 6.2\]](#)

$$(P - \rho_1)v_N = [P, \chi_N]e^{S/h}(\mathbf{u} - \mathbf{u}_{CN}) + \mathcal{O}(h^{\delta N}),$$

and thus, on $\{d_N(x, \text{supp } \nabla \chi_N) \geq \varepsilon\} \times \mathbb{R}^n$, where $\varepsilon > 0$ is fixed small enough and d_N is the distance associated with the metric $(Nk)^{-1}(dx')^2 + (Nk)^{-\frac{4}{3}} dx_n^2$,

$$T_N(P - \rho_1)v_N = \mathcal{O}(h^{\delta' N}),$$

for some $\delta' = \delta'(\varepsilon) > 0$.

(3) *(Almost) standard analytic propagation.* Although we are in a region where no analytic assumption is made, a rescaling of the problem gives estimates similar to those encountered in the analytic context. Indeed, setting

$$\tilde{h} = \tilde{h}_N := \frac{h}{Nk} = \left(N \ln \frac{1}{h}\right)^{-1},$$

and performing the change of variables (still working in the same coordinates, for which $\hat{x} = 0$)

$$\begin{aligned} x &\mapsto \tilde{x} = (\tilde{x}', \tilde{x}_n) := ((Nk)^{-\frac{1}{2}}x', (Nk)^{-\frac{2}{3}}x_n), \\ \xi &\mapsto \tilde{\xi} = (\tilde{\xi}', \tilde{\xi}_n) := ((Nk)^{-\frac{1}{2}}\xi', (Nk)^{-\frac{2}{3}}\xi_n), \end{aligned}$$

we see that the estimate (7-10) implies (see [FLM, Formula (6.43)])

$$T\tilde{v}_N(\tilde{x}, \tilde{\xi}; \tilde{h}_N) = \mathcal{O}(e^{-\delta'_L/2\tilde{h}_N})$$

uniformly in the tubular domain

$$\begin{aligned} \tilde{W}(\tilde{h}) := \{ &|\tilde{x}_n - \tilde{x}_n(-\delta^{-1})| \leq \delta, |\tilde{\xi}_n - \tilde{\xi}_n(-\delta^{-1})| \leq \delta, |\tilde{x}' - \tilde{x}'(-\delta^{-1})| \leq \delta(Nk)^{-\frac{1}{6}}, \\ &|\tilde{\xi}' - \tilde{\xi}'(-\delta^{-1})| \leq \delta(Nk)^{-\frac{1}{6}} \}, \end{aligned} \quad (7-11)$$

where

$$\begin{aligned} \tilde{v}_N(\tilde{x}) &:= (Nk)^{(n-1)/4+\frac{1}{3}}v_N((Nk)^{1/2}\tilde{x}', (Nk)^{2/3}\tilde{x}_n), \\ (\tilde{x}(\tilde{t}), \tilde{\xi}(\tilde{t})) &:= \exp \tilde{t}H_{\tilde{p}_1}(0, 0), \\ \tilde{p}_1(\tilde{x}, \tilde{\xi}) &:= (Nk)^{1/3}|\tilde{\xi}'|^2 + \tilde{\xi}_n^2 + W_1(\tilde{x}, \tilde{h}), \\ W_1(\tilde{x}, \tilde{h}) &:= (Nk)^{-\frac{2}{3}}V_1((Nk)^{1/2}\tilde{x}', (Nk)^{2/3}\tilde{x}_n) - (Nk)^{-\frac{2}{3}}\rho_1. \end{aligned}$$

Moreover, setting

$$\tilde{P} := -(Nk)^{1/3}\tilde{h}^2\Delta_{\tilde{x}'} - \tilde{h}^2\partial_{\tilde{x}_n}^2 + W_1(\tilde{x}),$$

then, for any $N \geq 1$ large enough, we also have

$$T\tilde{P}\tilde{v}_N(\tilde{x}, \tilde{\xi}; \tilde{h}_N) = \mathcal{O}(e^{-\delta'/2\tilde{h}_N}),$$

uniformly with respect to $h > 0$ small enough and $(\tilde{x}, \tilde{\xi}) \in \mathbb{R}^{2n}$ satisfying

$$d_N(((Nk)^{1/2}\tilde{x}', (Nk)^{2/3}\tilde{x}_n), \text{supp } \nabla \chi_N) \geq \varepsilon.$$

Finally, by Proposition 7.1 and Proposition 7.3, we have the a priori estimate

$$\|\tilde{v}_N\|_{H^1} = \mathcal{O}(h^{-N_1}) = \mathcal{O}(e^{N_1/(N\tilde{h})}),$$

for some $N_1 \geq 0$ independent of N , and we observe that, for $N = L/\delta_L$, one has $N_1/(\delta_L N) \rightarrow 0$ as $L \rightarrow +\infty$.

At this point, a small refinement of the propagation of the microsupport (see [FLM, Proposition 6.8]) gives the existence of a constant $\delta_1 > 0$ independent of L such that, for all L large enough and $N = L/\delta_L$, one has

$$T\tilde{v}_N(\tilde{x}, \tilde{\xi}; \tilde{h}) = \mathcal{O}(e^{-\delta_1\delta_L/\tilde{h}}), \quad (7-12)$$

uniformly in $V(\delta_1) = \{\tilde{x} : |\tilde{x}| \leq \delta_1\} \times \{\tilde{\xi} : (Nk)^{1/6}|\tilde{\xi}'| + |\tilde{\xi}_n| \leq \delta_1\}$.

Then, using an ellipticity property of \tilde{p}_1 away from $\{\tilde{\xi} : (Nk)^{1/6}|\tilde{\xi}'| + |\tilde{\xi}_n| \leq \delta_1\}$ and reconstructing \tilde{v}_N from $T\tilde{v}_N$, one finally finds

$$\|\tilde{v}_N\|_{H^m(|\tilde{x}| \leq \delta_2)} = \mathcal{O}(e^{-\delta_2\delta_L/\tilde{h}}),$$

with $m \geq 0$ arbitrary, $\delta_2 > 0$ independent of L , $N = L/\delta_L$, and L arbitrarily large. Therefore, turning back to the original coordinates x and parameter h and making \hat{x} vary on all of $\partial\ddot{O}$, [Theorem 7.4](#) follows. \square

8. Asymptotics of the width

As before, we denote by P_θ the distorted operator obtained from P by means of a complex distortion as in (7-2), with R_0 sufficiently large in order to have $\ddot{O} \subset \{|x| \leq R_0/2\}$. We also denote by \mathbf{u}_θ the corresponding distorted state obtain from \mathbf{u} by applying the same distortion (see, e.g., [\[FLM\]](#) for more details).

Let $\psi_0 \in C_0^\infty([0, 2]; [0, 1])$ with $\psi_0 = 1$ near $[0, 1]$, and set

$$\psi_N(x) := \psi_0\left(\frac{\text{dist}(x, \ddot{O})}{(Nk)^{2/3}}\right),$$

where, as before, $N = L/\delta_L$ with $L \geq 1$ arbitrarily large.

Then, since $\psi_N \mathbf{u} = \psi_N \mathbf{u}_\theta$, $P_\theta \mathbf{u}_\theta = \rho_1 \mathbf{u}_\theta$ and $\psi_N P_\theta \psi_N \mathbf{u}_\theta = \psi_N P \psi_N \mathbf{u}$, we have

$$\text{Im } \rho_1 \|\psi_N \mathbf{u}\|^2 = \text{Im} \langle \psi_N P_\theta \mathbf{u}_\theta, \psi_N \mathbf{u} \rangle = \text{Im} \langle [\psi_N, P_\theta] \mathbf{u}_\theta, \psi_N \mathbf{u} \rangle,$$

and thus

$$\text{Im } \rho_1 = \text{Im} \frac{\langle 2h^2(\nabla \psi_N) \nabla \mathbf{u} + h^2(\Delta \psi_N) \mathbf{u}, \psi_N \mathbf{u} \rangle + h \langle [\psi_N, R_\theta] \mathbf{u}_\theta, \psi_N \mathbf{u} \rangle}{\|\psi_N \mathbf{u}\|^2}. \tag{8-1}$$

Moreover, we know that $\|\psi_N \mathbf{u}\| = 1 + \mathcal{O}(e^{-\delta/h})$ with $\delta > 0$ and, by [Theorem 7.4](#), on $\text{supp } \tilde{\psi}_N$ we can replace \mathbf{u} by \mathbf{u}_{CN} , up to an error $\mathcal{O}(h^L e^{-S/h})$. Also, using [Proposition 7.1](#) we deduce

$$\text{Im } \rho_1 = \text{Im} \langle 2h^2(\nabla \psi_N) \nabla \mathbf{u}_{CN} + h^2(\Delta \psi_N) \mathbf{u}, \psi_N \mathbf{u}_{CN} \rangle + h \langle [\psi_N, R_\theta] \mathbf{u}_\theta, \psi_N \mathbf{u} \rangle + \mathcal{O}(h^{L-N_0}) e^{-2S/h} \tag{8-2}$$

for some fixed $N_0 \geq 0$ independent of L .

Now, we let $\tilde{\psi}_0 \in C_0^\infty((1, 2); [0, 1])$ with $\tilde{\psi}_0 = 1$ near $\text{supp } \nabla \psi_0$ and set $\tilde{\psi}_N(x) = \tilde{\psi}_0\left(\frac{\text{dist}(x, \ddot{O})}{(Nk)^{2/3}}\right)$.

Lemma 8.1. *One has*

$$\langle [\psi_N, R_\theta] \mathbf{u}_\theta, \psi_N \mathbf{u} \rangle = \langle \psi_N [\psi_N, R] \tilde{\psi}_N \mathbf{u}, \tilde{\psi}_N \mathbf{u} \rangle + \mathcal{O}(h^\infty e^{-2S/h}). \tag{8-3}$$

Proof. Thanks to [Assumption 3](#), in $[\psi_N, R_\theta]$, we can make the (complex) change of contour of integration

$$\mathbb{R}^n \ni \xi \mapsto \xi + i\sqrt{M_0} \frac{x-y}{\sqrt{(x-y)^2 + h^2}}.$$

We obtain

$$[\psi_N, R_\theta] \mathbf{u}_\theta(x) = \frac{1}{(2\pi h)^n} \int e^{i(x-y)\xi/h - \Phi/h} (\psi_N(x) - \psi_N(y)) \tilde{r}_\theta \bar{\mathbf{u}}_\theta(y) dy d\xi,$$

with

$$\Phi := \sqrt{M_0} \frac{(x-y)^2}{\sqrt{(x-y)^2 + h^2}}, \quad \partial_{x,y}^\alpha \partial_\xi^\beta \tilde{r}_\theta(x, y, \xi) = \mathcal{O}(h^{-|\alpha|}(\xi)).$$

By construction, on the set

$$A_N := \text{supp}(\psi_N(x) - \psi_N(y)) \cap \{\tilde{\psi}_N(x) \neq 1 \text{ or } \tilde{\psi}_N(y) \neq 1\},$$

we have $|x - y| \geq c(Nk)^{2/3}$ for some constant $c > 0$. As a consequence, on this set, the quantity $|x - y|/\sqrt{(x - y)^2 + h^2}$ tends to 1 uniformly as $h \rightarrow 0$. Moreover, still on this set, we have either $s(x) = S$ or $s(y) = S$, and since $|s(x) - s(y)| \leq \mu|x - y|$ with $0 < \mu < \sqrt{M_0}$, we deduce the existence of a constant $c_0 > 0$ such that for $(x, y) \in A_N$, one has $s(x) + s(y) + \Phi \geq 2S + c_0(Nk)^{2/3}$.

Therefore, by the Calderón–Vaillancourt theorem (and also using Proposition 5.2 in order to regularize the function $s(x)$), we obtain

$$\|e^{-s/h}[\psi_N, R_\theta]e^{-s/h}(1 - \tilde{\psi}_N)\langle hD_n \rangle^{-1}\| + \|(1 - \tilde{\psi}_N)e^{-s/h}[\psi_N, R_\theta]e^{-s/h}\langle hD_n \rangle^{-1}\| = \mathcal{O}(h^\infty e^{-2S/h}).$$

Then, writing

$$\langle [\psi_N, R_\theta]\mathbf{u}_\theta, \psi_N \mathbf{u} \rangle = \langle e^{-s/h}[\psi_N, R_\theta]e^{-s/h}(e^{s/h}\mathbf{u}_\theta), \psi_N e^{s/h}\mathbf{u} \rangle$$

and using Proposition 7.1 and Remark 7.2, the result follows. \square

Inserting (8-3) into (8-2) and approaching $\tilde{\psi}_N \mathbf{u}$ by $\tilde{\psi}_N \mathbf{u}_{CN}$, we obtain

$$\begin{aligned} \operatorname{Im} \rho_1 &= \operatorname{Im} \langle 2h^2(\nabla \psi_N) \nabla \mathbf{u}_{CN} \\ &\quad + h^2(\Delta \psi_N) \mathbf{u}, \psi_N \mathbf{u}_{CN} \rangle + h \langle \psi_N [\psi_N, R] \tilde{\psi}_N \mathbf{u}_{CN}, \tilde{\psi}_N \mathbf{u}_{CN} \rangle + \mathcal{O}(h^{L-N_0})e^{-2S/h}. \end{aligned} \quad (8-4)$$

Finally, using Proposition 6.2 (in particular the expression (4-52) of \mathbf{u}_{CN} in $\bigcup_{\gamma \in G} {}^q W_N(\gamma) \cap \operatorname{supp} \tilde{\psi}_N$), we can perform a stationary-phase expansion in (8-4) (as in [FLM, Section 7]) and, for L large enough, we obtain

$$\operatorname{Im} \rho_1 = -h^{(1-n_\Gamma)/2} \sum_{j=n_0}^L \sum_{0 \leq m \leq \ell \leq L} f_{j,\ell,m} h^{j+\ell} |\ln h|^m e^{-2S/h} + \mathcal{O}(h^{L/2})e^{-2S/h},$$

with $f_{n_0,0,0} > 0$. In particular, the result for ρ_1 follows.

The result for ρ_j , $j \geq 2$, can be done along the same lines, by using a representation of $\operatorname{Im} \rho_j$ analogous to (8-1) and by approaching \mathbf{u} by a linear combination of WKB expressions similar to \mathbf{u}_{CN} , where the number of terms depends on the asymptotic multiplicity of the resonance; see [Helffer and Sjöstrand 1986, Section 10].

References

- [Berenger 1994] J.-P. Berenger, “A perfectly matched layer for the absorption of electromagnetic waves”, *J. Comput. Phys.* **114**:2 (1994), 185–200. MR 95e:78002 Zbl 0814.65129
- [Fujiié et al. 2011] S. Fujiié, A. Lahmar-Benbernou, and A. Martinez, “Width of shape resonances for non globally analytic potentials”, *J. Math. Soc. Japan* **63**:1 (2011), 1–78. MR 2012i:81100 Zbl 1210.81037
- [Helffer and Martinez 1987] B. Helffer and A. Martinez, “Comparaison entre les diverses notions de résonances”, *Helv. Phys. Acta* **60**:8 (1987), 992–1003. MR 89h:35083
- [Helffer and Sjöstrand 1984] B. Helffer and J. Sjöstrand, “Multiple wells in the semiclassical limit, I”, *Commun. Part. Diff. Eq.* **9**:4 (1984), 337–408. MR 86c:35113 Zbl 0546.35053
- [Helffer and Sjöstrand 1986] B. Helffer and J. Sjöstrand, *Résonances en limite semi-classique*, Mém. Soc. Math. France (N.S.) **24-25**, Société Mathématique de France, Paris, 1986. MR 88i:81025 Zbl 0631.35075
- [Hunziker 1986] W. Hunziker, “Distortion analyticity and molecular resonance curves”, *Ann. Inst. H. Poincaré Phys. Théor.* **45**:4 (1986), 339–358. MR 88h:81241 Zbl 0619.46068

- [Klein 1987] M. Klein, “On the mathematical theory of predissociation”, *Ann. Physics* **178**:1 (1987), 48–73. [MR 89f:81278](#) [Zbl 0649.35024](#)
- [Klein et al. 1992] M. Klein, A. Martinez, R. Seiler, and X. P. Wang, “On the Born–Oppenheimer expansion for polyatomic molecules”, *Comm. Math. Phys.* **143**:3 (1992), 607–639. [MR 93b:81069](#) [Zbl 0754.35099](#)
- [Kronig 1928] R. de L. Kronig, “Zur Deutung der Bandenspektren, II”, *Z. Phys.* **50**:5-6 (1928), 347–362.
- [Landau 1932a] L. Landau, “Zur Theorie der Energieübertragung bei Stößen”, *Phys. Z. Sowjetunion* **1** (1932), 88–98. [Zbl 0004.13704](#)
- [Landau 1932b] L. Landau, “Zur Theorie der Energieübertragung, II”, *Phys. Z. Sowjetunion* **2** (1932), 46–51. [Zbl 0005.27801](#)
- [Martinez 1987] A. Martinez, “Estimations de l’effet tunnel pour le double puits, I”, *J. Math. Pures Appl.* (9) **66**:2 (1987), 195–215. [MR 88i:35041](#) [Zbl 0564.35028](#)
- [Martinez 2002] A. Martinez, *An introduction to semiclassical and microlocal analysis*, Springer, New York, 2002. [MR 2003b:35010](#) [Zbl 0994.35003](#)
- [Martinez and Sordoni 2009] A. Martinez and V. Sordoni, *Twisted pseudodifferential calculus and application to the quantum evolution of molecules*, Mem. Amer. Math. Soc. **936**, Amer. Math. Soc., Providence, RI, 2009. [MR 2011e:35323](#) [Zbl 1179.35357](#)
- [Pettersson 1997] P. Pettersson, “WKB expansions for systems of Schrödinger operators with crossing eigenvalues”, *Asymptot. Anal.* **14**:1 (1997), 1–48. [MR 98b:35126](#) [Zbl 0885.35104](#)
- [Sjöstrand 1982] J. Sjöstrand, “Singularités analytiques microlocales”, pp. 1–166 *Astérisque* **95**, Soc. Math. France, Paris, 1982. [MR 84m:58151](#) [Zbl 0524.35007](#)
- [Stückelberg 1932] E. C. G. Stückelberg, “Theorie der unelastischen Stösse zwischen Atomen”, *Helv. Phys. Acta* **5**:6 (1932), 369–422. [Zbl 0006.09006](#)
- [Zener 1932] C. Zener, “Non-adiabatic crossing of energy levels”, *Proc. R. Soc. London Ser. A* **137**:833 (1932), 696–702. [Zbl 0005.18605](#)

Received 28 Oct 2011. Revised 23 May 2014. Accepted 30 Jun 2014.

ALAIN GRIGIS: grigis@math.univ-paris13.fr

Département de Mathématiques, Université Paris 13 - Institut Galilée, Avenue Jean-Baptiste Clement, 93430 Villetaneuse, France

ANDRÉ MARTINEZ: andre.martinez@unibo.it

Dipartimento di Matematica, Università di Bologna, Piazza di Porta San Donato, I-40127 Bologna, Italy

Analysis & PDE

msp.org/apde

EDITORS

EDITOR-IN-CHIEF

Maciej Zworski
zworski@math.berkeley.edu
University of California
Berkeley, USA

BOARD OF EDITORS

Nicolas Burq	Université Paris-Sud 11, France nicolas.burq@math.u-psud.fr	Yuval Peres	University of California, Berkeley, USA peres@stat.berkeley.edu
Sun-Yung Alice Chang	Princeton University, USA chang@math.princeton.edu	Gilles Pisier	Texas A&M University, and Paris 6 pisier@math.tamu.edu
Michael Christ	University of California, Berkeley, USA mchrist@math.berkeley.edu	Tristan Rivière	ETH, Switzerland riviere@math.ethz.ch
Charles Fefferman	Princeton University, USA cf@math.princeton.edu	Igor Rodnianski	Princeton University, USA irod@math.princeton.edu
Ursula Hamenstaedt	Universität Bonn, Germany ursula@math.uni-bonn.de	Wilhelm Schlag	University of Chicago, USA schlag@math.uchicago.edu
Vaughan Jones	U.C. Berkeley & Vanderbilt University vaughan.f.jones@vanderbilt.edu	Sylvia Serfaty	New York University, USA serfaty@cims.nyu.edu
Herbert Koch	Universität Bonn, Germany koch@math.uni-bonn.de	Yum-Tong Siu	Harvard University, USA siu@math.harvard.edu
Izabella Laba	University of British Columbia, Canada ilaba@math.ubc.ca	Terence Tao	University of California, Los Angeles, USA tao@math.ucla.edu
Gilles Lebeau	Université de Nice Sophia Antipolis, France lebeau@unice.fr	Michael E. Taylor	Univ. of North Carolina, Chapel Hill, USA met@math.unc.edu
László Lempert	Purdue University, USA lempert@math.purdue.edu	Gunther Uhlmann	University of Washington, USA gunther@math.washington.edu
Richard B. Melrose	Massachusetts Institute of Technology, USA rbm@math.mit.edu	András Vasy	Stanford University, USA andras@math.stanford.edu
Frank Merle	Université de Cergy-Pontoise, France Frank.Merle@u-cergy.fr	Dan Virgil Voiculescu	University of California, Berkeley, USA dvv@math.berkeley.edu
William Minicozzi II	Johns Hopkins University, USA minicozz@math.jhu.edu	Steven Zelditch	Northwestern University, USA zelditch@math.northwestern.edu
Werner Müller	Universität Bonn, Germany mueller@math.uni-bonn.de		

PRODUCTION

production@msp.org

Silvio Levy, Scientific Editor

See inside back cover or msp.org/apde for submission instructions.

The subscription price for 2014 is US \$180/year for the electronic version, and \$355/year (+\$50, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues from the last three years and changes of subscribers address should be sent to MSP.

Analysis & PDE (ISSN 1948-206X electronic, 2157-5045 printed) at Mathematical Sciences Publishers, 798 Evans Hall #3840, c/o University of California, Berkeley, CA 94720-3840, is published continuously online. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices.

APDE peer review and production are managed by EditFLOW[®] from Mathematical Sciences Publishers.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2014 Mathematical Sciences Publishers

ANALYSIS & PDE

Volume 7 No. 5 2014

Resonance widths for the molecular predissociation	1027
ALAIN GRIGIS and ANDRÉ MARTINEZ	
Quasimodes and a lower bound on the uniform energy decay rate for Kerr–AdS spacetimes	1057
GUSTAV HOLZEGEL and JACQUES SMULEVICI	
Cylindrical estimates for hypersurfaces moving by convex curvature functions	1091
BEN ANDREWS and MAT LANGFORD	
Uniform L^p -improving for weighted averages on curves	1109
BETSY STOVALL	
Propagation of singularities for rough metrics	1137
HART F. SMITH	
Well-posedness of Lagrangian flows and continuity equations in metric measure spaces	1179
LUIGI AMBROSIO and DARIO TREVISAN	
Erratum to “Poisson statistics for eigenvalues of continuum random Schrödinger operators”	1235
JEAN-MICHEL COMBES, FRANÇOIS GERMINET and ABEL KLEIN	