

ANALYSIS & PDE

Volume 8

No. 5

2015

TOMÁS LUNGENSTRASS AND GEORGI RAIKOV

**LOCAL SPECTRAL ASYMPTOTICS
FOR METRIC PERTURBATIONS OF THE LANDAU
HAMILTONIAN**

LOCAL SPECTRAL ASYMPTOTICS FOR METRIC PERTURBATIONS OF THE LANDAU HAMILTONIAN

TOMÁS LUNGENSTRASS AND GEORGI RAIKOV

We consider metric perturbations of the Landau Hamiltonian. We investigate the asymptotic behavior of the discrete spectrum of the perturbed operator near the Landau levels, for perturbations of compact support, and of exponential or power-like decay at infinity.

1. Introduction

Let

$$H_0 := (-i\nabla - A_0)^2$$

with $A_0 = (A_{0,1}, A_{0,2}) := \frac{1}{2}b(-x_2, x_1)$ be the Landau Hamiltonian, self-adjoint in $L^2(\mathbb{R}^2)$, and essentially self-adjoint on $C_0^\infty(\mathbb{R}^2)$. In other words, H_0 is the two-dimensional Schrödinger operator with constant scalar magnetic field $b > 0$, that is, the Hamiltonian of a two-dimensional, spinless, nonrelativistic quantum particle subject to a constant magnetic field. As is well known, the spectrum $\sigma(H_0)$ consists of infinitely degenerate eigenvalues $\Lambda_q := b(2q + 1)$, $q \in \mathbb{Z}_+ := \{0, 1, 2, \dots\}$, called *Landau levels* (see, e.g., [Fock 1928; Landau 1930]).

In the present article we consider metric perturbations of H_0 . Namely, let

$$m(x) = \{m_{jk}(x)\}_{j,k=1,2}, \quad x \in \mathbb{R}^2,$$

be a Hermitian 2×2 matrix such that $m(x) \geq 0$ for all $x \in \mathbb{R}^2$. Throughout the article we assume that $m_{jk} \in C_b^\infty(\mathbb{R}^2)$, $j, k = 1, 2$, i.e., $m_{jk} \in C^\infty(\mathbb{R}^2)$, and the entries m_{jk} together with all their derivatives are bounded on \mathbb{R}^2 . Set

$$\Pi_j := -i \frac{\partial}{\partial x_j} - A_{0,j}, \quad j = 1, 2, \tag{1-1}$$

so that $H_0 = \Pi_1^2 + \Pi_2^2$. On $\text{Dom } H_0$, define the operators

$$H_\pm := \sum_{j,k=1,2} \Pi_j (\delta_{jk} \pm m_{jk}) \Pi_k = H_0 \pm W,$$

where $W := \sum_{j,k=1,2} \Pi_j m_{jk} \Pi_k$; in the case of H_- , we suppose additionally that $\sup_{x \in \mathbb{R}^2} |m(x)| < 1$. Thus, the matrices $g_\pm(x) = \{g_{jk}^\pm(x)\}_{j,k=1,2}$ with $g_{jk}^\pm := \delta_{jk} \pm m_{jk}$ are positive definite for each $x \in \mathbb{R}^2$. Under these assumptions, the operators H_\pm are self-adjoint in $L^2(\mathbb{R}^2)$, and essentially self-adjoint on $C_0^\infty(\mathbb{R}^2)$ (see the Appendix).

MSC2010: 35J10, 35P20, 47G30, 81Q10.

Keywords: Landau Hamiltonian, metric perturbations, position-dependent mass, spectral asymptotics.

From a mathematical physics point of view, the operators H_{\pm} are special cases of Schrödinger operators with *position-dependent mass*, which have a long history (see, e.g., [Bastard et al. 1975; von Roos 1983]), but have received increased attention during the last decade (see, e.g., [Midya et al. 2010; Gadella and Smolyanov 2008; Killingbeck 2011]). We would like to mention especially [de Souza Dutra and de Oliveira 2009], where the model considered is quite close to the operators H_{\pm} discussed here.

The operators H_{\pm} admit also a geometric interpretation, since they are related to the Bochner Laplacians corresponding to connections with constant nonvanishing curvature (see, e.g., [Rosenberg 1997; Colin de Verdière 1986]); we discuss this relation in more detail at the end of Section 2. Further, assume that

$$\lim_{|x| \rightarrow \infty} m_{jk}(x) = 0, \quad j, k = 1, 2. \quad (1-2)$$

Thus m models a localized perturbation with respect to a reference medium. Under condition (1-2), the resolvent difference $H_{\pm}^{-1} - H_0^{-1}$ is a compact operator (see the Appendix), and therefore the essential spectra of H_{\pm} and H_0 coincide:

$$\sigma_{\text{ess}}(H_{\pm}) = \sigma_{\text{ess}}(H_0) = \sigma(H_0) = \bigcup_{q=0}^{\infty} \{\Lambda_q\}.$$

The spectrum $\sigma(H_{\pm})$ on $\mathbb{R} \setminus \bigcup_{q=0}^{\infty} \{\Lambda_q\}$ may consist of discrete eigenvalues whose only possible accumulation points are the Landau levels. Moreover, taking into account that $W \geq 0$, and applying [Birman and Solomjak 1987, Section 9.4, Theorem 7], we find that the eigenvalues of H_+ (resp. H_-) may accumulate to a given Landau level Λ_q only from above (resp. from below). Fix $q \in \mathbb{Z}_+$. Let $\{\lambda_{k,q}^-\}$ be the eigenvalues of H_- lying on the interval $(\Lambda_{q-1}, \Lambda_q)$ with $\Lambda_{-1} := -\infty$, counted with multiplicities and enumerated in increasing order. Similarly, let $\{\lambda_{k,q}^+\}$ be the eigenvalues of H_+ lying on the interval $(\Lambda_q, \Lambda_{q+1})$, counted with multiplicities and enumerated in decreasing order.

The aim of the article is to investigate the rate of convergence of $\lambda_{k,q}^{\pm} - \Lambda_q$ as $k \rightarrow \infty$, with $q \in \mathbb{Z}_+$ fixed, for perturbations m of compact support, of exponential decay, or of power-like decay at infinity.

The properties of the discrete spectrum generated by perturbative second-order differential operators with decaying coefficients have been considered also in [Alama et al. 1994; Boyarchenko and Levendorskii 1997; Briet et al. 2009; Raikov 2015].

The article is organized as follows. In Section 2 we formulate our main results and briefly comment on them. In Section 3 we reduce our analysis to the study of operators of Berezin–Toeplitz type, and in Section 4 we establish several useful unitary equivalences for these operators. Section 5 contains the proofs of our results in the case of rapid decay, i.e., of compact support or exponential decay, while the proofs for slow, i.e., power-like decay, can be found in Section 6. Finally, in the Appendix we address some standard issues concerning the domain of the operators H_{\pm} and the compactness of the resolvent difference $H_0^{-1} - H_{\pm}^{-1}$.

2. Main results

First, we formulate our results concerning perturbations m of compact support. Denote by $m_{<}(x)$ and $m_{>}(x)$, with $m_{<}(x) \leq m_{>}(x)$, the two eigenvalues of the matrix $m(x)$, $x \in \mathbb{R}^2$.

Theorem 2.1. *Assume that the support of the matrix m is compact, and its smaller eigenvalue $m_<$ does not vanish identically. Fix $q \in \mathbb{Z}_+$. Then we have*

$$\ln(\pm(\lambda_{k,q}^\pm - \Lambda_q)) = -k \ln k + O(k), \quad k \rightarrow \infty. \tag{2-1}$$

Remarks. (i) Under additional technical hypotheses on m_\geq , we could make the asymptotic relation (2-1) more precise. Namely, assume that there exists a nonincreasing sequence $\{s_j\}_{j \in \mathbb{N}}$ such that $s_j > 0$, $j \in \mathbb{N}$, $\lim_{j \rightarrow \infty} s_j = 0$, and the level lines

$$\{x \in \mathbb{R}^2 \mid m_<(x) = s_j\}, \quad j \in \mathbb{N},$$

are bounded Lipschitz curves. In particular, the existence of such a sequence follows from the Sard lemma (see, e.g., [Sternberg 1964, Chapter 2, Theorem 3.1]) if we assume that $m_< \in C^2(\mathbb{R}^2)$. Further, denote by \mathcal{C}_\geq the logarithmic capacities (see, e.g., [Landkof 1972, Chapter II, Section 4]) of $\text{supp } m_\geq$. Then we have

$$\left(1 + \ln\left(\frac{b\mathcal{C}_<^2}{2}\right)\right)k + o(k) \leq \ln(\pm(\lambda_{k,q}^\pm - \Lambda_q)) + k \ln k \leq \left(1 + \ln\left(\frac{b\mathcal{C}_>^2}{2}\right)\right)k + o(k) \tag{2-2}$$

as $k \rightarrow \infty$. We omit the details of the proof of (2-2), which is inspired by [Filonov and Pushnitski 2006].

(ii) For $q \in \mathbb{Z}_+$ and $\lambda > 0$, set

$$\mathcal{N}_q^\pm(\lambda) := \#\{k \in \mathbb{Z}_+ \mid \pm(\lambda_{k,q}^\pm - \Lambda_q) > \lambda\}. \tag{2-3}$$

Then, a less precise version of (2-1), namely

$$\ln(\pm(\lambda_{k,q}^\pm - \Lambda_q)) = -k \ln k (1 + o(1)), \quad k \rightarrow \infty,$$

is equivalent to

$$\mathcal{N}_q^\pm(\lambda) = \frac{|\ln \lambda|}{\ln |\ln \lambda|} (1 + o(1)), \quad \lambda \downarrow 0. \tag{2-4}$$

Further, we state our results concerning perturbations of exponential decay. Assume that there exist constants $\beta > 0$ and $\gamma > 0$ such that

$$\ln m_\geq(x) = -\gamma |x|^{2\beta} + O(\ln |x|), \quad |x| \rightarrow \infty. \tag{2-5}$$

Remark. In (2-5), we suppose that the values of γ and β are the same for $m_<$ and $m_>$. Of course, the remainder $O(\ln |x|)$ could be different for $m_<$ and $m_>$.

Given $\beta > 0$ and $\gamma > 0$, set $\mu := \gamma(2/b)^\beta$, $b > 0$ being the constant magnetic field.

Theorem 2.2. *Let m_\geq satisfy (2-5). Fix $q \in \mathbb{Z}_+$.*

(i) *If $\beta \in (0, 1)$, then there exist constants $f_j = f_j(\beta, \mu)$, $j \in \mathbb{N}$, with $f_1 = \mu$, such that*

$$\ln(\pm(\lambda_{k,q}^\pm - \Lambda_q)) = - \sum_{1 \leq j < 1/(1-\beta)} f_j k^{(\beta-1)j+1} + O(\ln k), \quad k \rightarrow \infty. \tag{2-6}$$

(ii) If $\beta = 1$, then

$$\ln(\pm(\lambda_{k,q}^\pm - \Lambda_q)) = -(\ln(1 + \mu))k + O(\ln k), \quad k \rightarrow \infty. \tag{2-7}$$

(iii) If $\beta \in (1, \infty)$, then there exist constants $g_j = g_j(\beta, \mu)$, $j \in \mathbb{N}$, such that

$$\begin{aligned} &\ln(\pm(\lambda_{k,q}^\pm - \Lambda_q)) \\ &= -\frac{\beta-1}{\beta}k \ln k + \left(\frac{\beta-1-\ln(\mu\beta)}{\beta}\right)k - \sum_{1 \leq j < \beta/(\beta-1)} g_j k^{(1/\beta-1)j+1} + O(\ln k), \quad k \rightarrow \infty. \end{aligned} \tag{2-8}$$

Remarks. (i) Let us describe explicitly the coefficients f_j and g_j , $j \in \mathbb{N}$, appearing in (2-6) and (2-8) respectively. Assume first that $\beta \in (0, 1)$. For $s > 0$ and $\epsilon \in \mathbb{R}$, $|\epsilon| \ll 1$, introduce the function

$$F(s; \epsilon) := s - \ln s + \epsilon \mu s^\beta. \tag{2-9}$$

Denote by $s_<(\epsilon)$ the unique positive solution of the equation $s = 1 - \epsilon \beta \mu s^\beta$, so that $\partial F(s_<(\epsilon); \epsilon)/\partial s = 0$. Set

$$f(\epsilon) := F(s_<(\epsilon); \epsilon). \tag{2-10}$$

Note that f is a real analytic function for small $|\epsilon|$. Then $f_j := (1/j!) d^j f(0)/d\epsilon^j$, $j \in \mathbb{N}$.

Let now $\beta \in (1, \infty)$. For $s > 0$ and $\epsilon \in \mathbb{R}$, $|\epsilon| \ll 1$, introduce the function

$$G(s; \epsilon) := \mu s^\beta - \ln s + \epsilon s. \tag{2-11}$$

Denote by $s_>(\epsilon)$ the unique positive solution of the equation $\beta \mu s^\beta = 1 - \epsilon s$, so that $\partial G(s_>(\epsilon); \epsilon)/\partial s = 0$. Define

$$g(\epsilon) := G(s_>(\epsilon); \epsilon), \tag{2-12}$$

which is a real analytic function for small $|\epsilon|$. Then $g_j := (1/j!) d^j g(0)/d\epsilon^j$, $j \in \mathbb{N}$.

(ii) If, instead of (2-5), we assume that

$$\ln m_{\geq}(x) = -\gamma|x|^{2\beta}(1 + o(1)), \quad |x| \rightarrow \infty, \tag{2-13}$$

then we can prove less precise versions of (2-6), (2-7), and (2-8), namely

$$\ln(\pm(\lambda_{k,q}^\pm - \Lambda_q)) = \begin{cases} -\mu k^\beta(1 + o(1)) & \text{if } \beta \in (0, 1), \\ -(\ln(1 + \mu))k(1 + o(1)) & \text{if } \beta = 1, \\ -\frac{\beta-1}{\beta}k \ln k(1 + o(1)) & \text{if } \beta \in (1, \infty), \end{cases} \quad k \rightarrow \infty,$$

which are equivalent to

$$\mathcal{N}_q^\pm(\lambda) = \begin{cases} \mu^{-1/\beta} |\ln \lambda|^{1/\beta} (1 + o(1)) & \text{if } \beta \in (0, 1), \\ \frac{1}{\ln(1 + \mu)} |\ln \lambda| (1 + o(1)) & \text{if } \beta = 1, \\ \frac{\beta}{\beta-1} \frac{|\ln \lambda|}{\ln |\ln \lambda|} (1 + o(1)) & \text{if } \beta \in (1, \infty), \end{cases} \quad \lambda \downarrow 0. \tag{2-14}$$

Note that in (2-13), similarly to (2-5), we assume that the values of γ and β are the same for $m_<$ and $m_>$. However, since the coefficient in (2-14) with $\beta > 1$ does not depend on γ , in this case we could assume different values of $\gamma > 0$ for $m_<$ and $m_>$.

Finally, we consider perturbations m which admit a power-like decay at infinity. For $\rho > 0$ recall the definition of the Hörmander class

$$\mathcal{S}^{-\rho}(\mathbb{R}^2) := \{\psi \in C^\infty(\mathbb{R}^2) \mid |D^\alpha \psi(x)| \leq c_\alpha \langle x \rangle^{-\rho-|\alpha|}, x \in \mathbb{R}^2, \alpha \in \mathbb{Z}_+^2\},$$

where $\langle x \rangle := (1 + |x|^2)^{1/2}$, $x \in \mathbb{R}^2$. Let $\psi : \mathbb{R}^2 \rightarrow \mathbb{R}$ satisfy $\lim_{|x| \rightarrow \infty} \psi(x) = 0$. Set

$$\Phi_\psi(\lambda) := |\{x \in \mathbb{R}^2 \mid \psi(x) > \lambda\}|, \quad \lambda > 0, \tag{2-15}$$

where $|\cdot|$ denotes the Lebesgue measure. Fix $q \in \mathbb{Z}_+$, and introduce the function

$$\mathcal{T}_q(x) := \frac{1}{2}(\Lambda_q \operatorname{Tr} m(x) - 2b \operatorname{Im} m_{12}(x)), \quad x \in \mathbb{R}^2. \tag{2-16}$$

Note that $\mathcal{T}_q(x) \geq 0$ for any $x \in \mathbb{R}^2$ and $q \in \mathbb{Z}_+$.

Theorem 2.3. *Let $m_{jk} \in \mathcal{S}^{-\rho}(\mathbb{R}^2)$, $j, k = 1, 2$, with $\rho > 0$. Fix $q \in \mathbb{Z}_+$. Suppose that there exists a function $0 < \tau_q \in C^\infty(\mathbb{S}^1)$ such that*

$$\lim_{|x| \rightarrow \infty} |x|^\rho \mathcal{T}_q(x) = \tau_q\left(\frac{x}{|x|}\right).$$

Then we have

$$\mathcal{N}_q^\pm(\lambda) = \frac{b}{2\pi} \Phi_{\mathcal{T}_q}(\lambda)(1 + o(1)) \asymp \lambda^{-2/\rho}, \quad \lambda \downarrow 0, \tag{2-17}$$

which is equivalent to

$$\lim_{\lambda \downarrow 0} \lambda^{2/\rho} \mathcal{N}_q^\pm(\lambda) = \mathcal{C}_q := \frac{b}{4\pi} \int_0^{2\pi} \tau_q(\cos \theta, \sin \theta)^{2/\rho} d\theta, \tag{2-18}$$

or to

$$\pm(\lambda_{k,q}^\pm - \Lambda_q) = \mathcal{C}_q^{\rho/2} k^{-\rho/2}(1 + o(1)), \quad k \rightarrow \infty. \tag{2-19}$$

Remarks. (i) Relation (2-17) could be regarded as a semiclassical one, although here the semiclassical interpretation is somewhat implicit. In Propositions 4.1 and 4.3 below, we show that the effective Hamiltonian, which governs the asymptotics of $\mathcal{N}_q^\pm(\lambda)$ as $\lambda \downarrow 0$, is a pseudodifferential operator with anti-Wick symbol $w_{q,b} := w_q \circ \mathcal{R}_b$ defined by (4-8) and (4-31). Under the assumptions of Theorem 2.3, $\mathcal{T}_{q,b} := \mathcal{T}_q \circ \mathcal{R}_b$ (see (2-16) and (4-31)) can be considered as the principal part of the symbol $w_{q,b}$, while the difference between the anti-Wick and the Weyl quantization is negligible. Then $(1/2\pi)\Phi_{\mathcal{T}_{q,b}}(\lambda) = (b/2\pi)\Phi_{\mathcal{T}_q}(\lambda)$ is just the main semiclassical asymptotic term for the eigenvalue counting function for a compact pseudodifferential operator with Weyl symbol $\mathcal{T}_{q,b}$.

(ii) There exists an extensive family of alternative sets of assumptions for Theorem 2.3 (see, e.g., [Ivrii 1998; Dauge and Robert 1987]). We have chosen here hypotheses which, for certain, are not the most general ones, but are quite explicit and, hopefully, easy to absorb.

Let us comment briefly on our results. Nowadays, there exists a relatively wide literature on the local spectral asymptotics for various magnetic quantum Hamiltonians. Let us concentrate here on three types of perturbations of H_0 which are considered to be of particular interest (see, e.g., [Ivrii 1998; Mao 2012]):

- Electric perturbations $H_0 + Q$ where $Q : \mathbb{R}^2 \rightarrow \mathbb{R}$ plays the role of the perturbative *electric potential*.
- Magnetic perturbations $(-i\nabla - A_0 - A)^2$, where $A = (A_1, A_2)$, and $B := \partial A_2/\partial x_1 - \partial A_1/\partial x_2$ is the perturbative *magnetic field*.
- Metric perturbations $\sum_{j,k=1,2} \Pi_j (\delta_{jk} + m_{jk}) \Pi_k$, where $m = \{m_{jk}\}_{j,k=1,2}$ is an appropriate perturbative matrix-valued function.

Typically, the perturbations Q , B , or m are supposed to decay in a suitable sense at infinity. Slowly decaying Q , for example $Q \in \mathcal{S}^{-\rho}(\mathbb{R}^2)$ with $\rho > 0$, were considered in [Raïkov 1990], and the main asymptotic terms of the corresponding counting functions $\mathcal{N}_q^\pm(\lambda)$ as $\lambda \downarrow 0$ were found, utilizing, in particular, anti-Wick pseudodifferential operators. In [Ivrii 1998, Theorem 11.3.17], the case of *combined* electric, magnetic, and metric slowly decaying perturbations was investigated; the main asymptotic terms of $\mathcal{N}_q^\pm(\lambda)$ as $\lambda \downarrow 0$, as well as certain remainder estimates were obtained. The semiclassical microlocal analysis applied in [Ivrii 1998] imposed restrictions on the symbols involved, which, in some sense or another, had to decay at infinity less rapidly than their derivatives. These restrictions excluded some rapidly decaying perturbations, e.g., those of compact support, or of exponential decay with $\beta \geq \frac{1}{2}$ (see (2-5)).

Raikov and Warzel [2002] used a different approach based on the spectral analysis of Berezin–Toeplitz operators and obtained the main asymptotic terms of $\mathcal{N}_q^\pm(\lambda)$ as $\lambda \downarrow 0$ in the case of potential perturbations Q of exponential decay or of compact support. In particular, in [Raikov and Warzel 2002], formulas of the type (2-4) or (2-14) appeared for the first time. Here, we essentially improve the methods developed in [Raikov and Warzel 2002]. These improvements lead also to more precise results for certain rapidly decaying electric perturbations. Namely, assume that $Q \geq 0$ admits a decay at infinity which is compatible in a suitable sense with the decay of m . Then the results of the article extend quite easily to operators of the form

$$H_\pm \pm Q, \tag{2-20}$$

so that $H_\pm \pm Q$ are perturbations of H_0 having a definite sign. We do not include these generalizations just in order to avoid an unreasonable increase of the size of the article due to results which do not require any really new arguments.

Combined perturbations of H_0 by compactly supported B and Q were considered in [Rozenblum and Tashchiyan 2008], where the main asymptotic terms of $\mathcal{N}_q^\pm(\lambda)$ as $\lambda \downarrow 0$ were found. Note that the magnetic perturbations of H_0 are never of fixed sign, which creates specific difficulties, successfully overcome in [Rozenblum and Tashchiyan 2008].

To our best knowledge, no results on the spectral asymptotics for rapidly decaying *metric* perturbations of H_0 appeared before in the literature. We also included in the article our result on slowly decaying metric perturbations (see Theorem 2.3), since it is coherent with the unified approach of the article and is proved by methods quite different from those in [Ivrii 1998].

Finally, let us discuss briefly the relation of H_{\pm} to the Bochner Laplacians. Assume that the elements of m are real. In \mathbb{R}^2 introduce a Riemannian metric generated by the inverse of g^{\pm} , and the connection 1-form $\sum_{j=1,2} A_{0,j} dx_j$. Set $\gamma_{\pm} := (\det g^{\pm})^{-1/2}$. Then the standard positive-definite Bochner Laplacian, self-adjoint in $L^2(\mathbb{R}^2; \gamma_{\pm} dx)$, is written in local coordinates as

$$\mathcal{L}_{\pm} := \gamma_{\pm}^{-1} \sum_{j,k=1,2} \Pi_j g_{jk}^{\pm} \gamma_{\pm} \Pi_k.$$

Let $U_{\pm} : L^2(\mathbb{R}^2; \gamma_{\pm} dx) \rightarrow L^2(\mathbb{R}^2; dx)$ be the unitary operator defined by $U_{\pm} f := \gamma_{\pm}^{1/2} f$. Then we have

$$U_{\pm} \mathcal{L}_{\pm} U_{\pm}^* = H_{\pm} + Q_{\pm}, \tag{2-21}$$

where

$$Q_{\pm} := \frac{1}{4} \sum_{j,k=1,2} \left(g_{jk}^{\pm} \frac{\partial \ln \gamma_{\pm}}{\partial x_k} \frac{\partial \ln \gamma_{\pm}}{\partial x_j} + 2 \frac{\partial}{\partial x_j} \left(g_{jk}^{\pm} \frac{\partial \ln \gamma_{\pm}}{\partial x_k} \right) \right).$$

Generally speaking, the functions Q_{\pm} do not have a definite sign coinciding with the sign of the operators $H_{\pm} - H_0$; hence, the operators on the right-hand side of (2-21) are not exactly of the form of (2-20). The fact that the symbol of a Toeplitz operator does not have a definite sign may cause considerable difficulties in the study of the spectral asymptotics of this operator if the symbol decays rapidly, and, in particular, when its support is compact (see, e.g., [Pushnitski and Rozenblum 2011]). Hopefully, we will overcome these difficulties in a future work, where we would consider the local spectral asymptotics of \mathcal{L}_{\pm} .

3. Reduction to Berezin–Toeplitz operators

In this section we reduce the analysis of the functions $\mathcal{N}_q^{\pm}(\lambda)$ as $\lambda \downarrow 0$ to the spectral asymptotics for certain compact operators of Berezin–Toeplitz type. To this end, we will need some more notations, and several auxiliary results from the abstract theory of compact operators in Hilbert space.

In what follows, we denote by $\mathbb{1}_M$ the characteristic function of the set M . Let T be a self-adjoint operator in a Hilbert space,¹ and $\mathcal{I} \subset \mathbb{R}$ be an interval. Set

$$N_{\mathcal{I}}(T) := \text{rank } \mathbb{1}_{\mathcal{I}}(T),$$

where, in accordance with our general notations, $\mathbb{1}_{\mathcal{I}}(T)$ is the spectral projection of T corresponding to \mathcal{I} . Thus, if $\mathcal{I} \cap \sigma_{\text{ess}}(T) = \emptyset$, then $N_{\mathcal{I}}(T)$ is just the number of the eigenvalues of T lying on \mathcal{I} and counted with their multiplicities. In particular,

$$\mathcal{N}_q^-(\lambda) = N_{(\Lambda_{q-1}, \Lambda_q - \lambda)}(H_-), \quad q \in \mathbb{Z}_+, \lambda \in (0, 2b), \tag{3-1}$$

$$\mathcal{N}_q^+(\lambda) = N_{(\Lambda_q + \lambda, \Lambda_{q+1})}(H_+), \quad q \in \mathbb{Z}_+, \lambda \in (0, 2b), \tag{3-2}$$

the functions \mathcal{N}_q^{\pm} being defined in (2-3). Let $T = T^*$ be a linear compact operator in a Hilbert space. For $s > 0$, set

$$n_{\pm}(s; T) := N_{(s, \infty)}(\pm T);$$

¹All the Hilbert spaces considered in the article are assumed to be separable.

thus, $n_+(s; T)$ (resp. $n_-(s; T)$) is just the number of the eigenvalues of the operator T larger than s (resp. smaller than $-s$), counted with multiplicities. If $T_j = T_j^*$, $j = 1, 2$, are two linear compact operators acting in a given Hilbert space, then the Weyl inequalities

$$n_{\pm}(s_1 + s_2; T_1 + T_2) \leq n_{\pm}(s_1; T_1) + n_{\pm}(s_2; T_2) \tag{3-3}$$

hold for $s_j > 0$ (see, e.g., [Birman and Solomjak 1987, Section 9.2, Theorem 9]).

Fix $q \in \mathbb{Z}_+$ and denote by P_q the orthogonal projection onto $\text{Ker}(H_0 - \Lambda_q)$. Since the operator $H_0^{-1}WH_0^{-1}$ is compact, the operator $P_qWP_q = \Lambda_q^2P_qH_0^{-1}WH_0^{-1}P_q$ is compact as well. Similarly, the operators $H_0^{-1}WH_{\pm}^{-1/2}$ are compact, and hence the operators

$$P_qWH_{\pm}^{-1}WP_q = \Lambda_q^2P_q(H_0^{-1}WH_{\pm}^{-1/2})(H_{\pm}^{-1/2}WH_0^{-1})P_q$$

are compact as well.

Proposition 3.1. *Under the general assumptions of the article we have*

$$\begin{aligned} n_+((1 + \varepsilon)\lambda; P_qWP_q \mp P_qWH_{\pm}^{-1}WP_q) + O(1) \\ \leq N_q^{\pm}(\lambda) \leq n_+((1 - \varepsilon)\lambda; P_qWP_q \mp P_qWH_{\pm}^{-1}WP_q) + O(1), \quad \lambda \downarrow 0, \end{aligned} \tag{3-4}$$

for each $\varepsilon \in (0, 1)$.

Proof. The argument is close in spirit to the proof of [Raikov and Warzel 2002, Proposition 4.1], and is based again on the (generalized) Birman–Schwinger principle. However, since the operator $H_0^{-1/2}WH_0^{-1/2}$ is only bounded but not compact, we cannot apply the Birman–Schwinger principle to the operator pair (H_0, H_{\pm}) , and apply it instead to the resolvent pair (H_0^{-1}, H_{\pm}^{-1}) . First of all, note that there exist Λ_- and Λ_+ with $\Lambda_- \in (0, \Lambda_0)$ if $q = 0$, $\Lambda_- \in (\Lambda_{q-1}, \Lambda_q)$ if $q \in \mathbb{N}$, and $\Lambda_+ \in (\Lambda_q, \Lambda_{q+1})$ if $q \in \mathbb{Z}_+$, such that

$$N_q^-(\lambda) = N_{(\Lambda_-, \Lambda_q - \lambda)}(H_-), \quad \lambda \in (0, \Lambda_q - \Lambda_-), \tag{3-5}$$

$$N_q^+(\lambda) = N_{(\Lambda_q + \lambda, \Lambda_+)}(H_+), \quad \lambda \in (0, \Lambda_+ - \Lambda_q). \tag{3-6}$$

Further, evidently,

$$N_{(\Lambda_-, \Lambda_q - \lambda)}(H_-) = N_{((\Lambda_q - \lambda)^{-1}, \Lambda_-^{-1})}(H_-^{-1}) = N_{((\Lambda_q - \lambda)^{-1}, \Lambda_-^{-1})}(H_0^{-1} + T_-), \tag{3-7}$$

$$N_{(\Lambda_q + \lambda, \Lambda_+)}(H_+) = N_{(\Lambda_+^{-1}, (\Lambda_q + \lambda)^{-1})}(H_+^{-1}) = N_{(\Lambda_+^{-1}, (\Lambda_q + \lambda)^{-1})}(H_0^{-1} - T_+), \tag{3-8}$$

with $T_- := H_-^{-1} - H_0^{-1}$ and $T_+ := H_0^{-1} - H_+^{-1}$. Note that the operators T_{\pm} are nonnegative and compact. By the generalized Birman–Schwinger principle (see, e.g., [Alama et al. 1989, Theorem 1.3]) we have

$$\begin{aligned} N_{((\Lambda_q - \lambda)^{-1}, \Lambda_-^{-1})}(H_0^{-1} + T_-) \\ = n_+(1; T_-^{1/2}((\Lambda_q - \lambda)^{-1} - H_0^{-1})^{-1}T_-^{1/2}) - n_+(1; T_-^{1/2}(\Lambda_-^{-1} - H_0^{-1})^{-1}T_-^{1/2}) - \dim \text{Ker}(H_- - \Lambda_-), \end{aligned} \tag{3-9}$$

and

$$\begin{aligned}
 & N_{(\Lambda_+^{-1}, (\Lambda_q + \lambda)^{-1})}(H_0^{-1} - T_+) \\
 &= n_+(1; T_+^{1/2}(H_0^{-1} - (\Lambda_q + \lambda)^{-1})^{-1}T_+^{1/2}) - n_+(1; T_+^{1/2}(H_0^{-1} - \Lambda_+^{-1})^{-1}T_+^{1/2}) - \dim \text{Ker}(H_+ - \Lambda_+). \quad (3-10)
 \end{aligned}$$

Since the operators T_{\pm} are compact and $\Lambda_{\pm} \notin \sigma(H_0)$, we find that the two last terms on the right-hand side of (3-9) and (3-10), which are independent of λ , are finite. Next, the Weyl inequalities (3-3) imply

$$\begin{aligned}
 & n_+(1 + \varepsilon; T_-^{1/2}((\Lambda_q - \lambda)^{-1} - H_0^{-1})^{-1}P_q T_-^{1/2}) - n_-(\varepsilon; T_-^{1/2}((\Lambda_q - \lambda)^{-1} - H_0^{-1})^{-1}(I - P_q)T_-^{1/2}) \\
 & \leq n_+(1; T_-^{1/2}((\Lambda_q - \lambda)^{-1} - H_0^{-1})^{-1}T_-^{1/2}) \\
 & \leq n_+(1 - \varepsilon; T_-^{1/2}((\Lambda_q - \lambda)^{-1} - H_0^{-1})^{-1}P_q T_-^{1/2}) \\
 & \qquad \qquad \qquad + n_+(\varepsilon; T_-^{1/2}((\Lambda_q - \lambda)^{-1} - H_0^{-1})^{-1}(I - P_q)T_-^{1/2}) \quad (3-11)
 \end{aligned}$$

for any $\varepsilon \in (0, 1)$. The operator $T_-^{1/2}((\Lambda_q - \lambda)^{-1} - H_0^{-1})^{-1}(I - P_q)T_-^{1/2}$ tends in norm as $\lambda \downarrow 0$ to the compact operator

$$T_-^{1/2} \left(\sum_{j \in \mathbb{Z}_+ \setminus \{q\}} (\Lambda_q^{-1} - \Lambda_j^{-1})^{-1} P_j \right) T_-^{1/2}.$$

Therefore,

$$n_{\pm}(\varepsilon; T_-^{1/2}((\Lambda_q - \lambda)^{-1} - H_0^{-1})^{-1}(I - P_q)T_-^{1/2}) = O(1), \quad \lambda \downarrow 0, \quad (3-12)$$

for any $\varepsilon > 0$. Next, for any $s > 0$ we have

$$\begin{aligned}
 n_+(s; T_-^{1/2}((\Lambda_q - \lambda)^{-1} - H_0^{-1})^{-1}P_q T_-^{1/2}) &= n_+(s; ((\Lambda_q - \lambda)^{-1} - \Lambda_q^{-1})^{-1}T_-^{1/2}P_q T_-^{1/2}) \\
 &= n_+(s\lambda(\Lambda_q - \lambda)^{-1}\Lambda_q^{-1}; P_q T_- P_q). \quad (3-13)
 \end{aligned}$$

Hence, (3-9) and (3-11)–(3-13) yield

$$\begin{aligned}
 & n_+((1 + \varepsilon)\lambda(\Lambda_q - \lambda)^{-1}\Lambda_q^{-1}; P_q T_- P_q) + O(1) \\
 & \leq N_{((\Lambda_q - \lambda)^{-1}, \Lambda_+^{-1})}(H_0^{-1} + T_-) \leq n_+((1 - \varepsilon)\lambda(\Lambda_q - \lambda)^{-1}\Lambda_q^{-1}; P_q T_- P_q) + O(1), \quad \lambda \downarrow 0, \quad (3-14)
 \end{aligned}$$

for any $\varepsilon \in (0, 1)$. Similarly, (3-10) and the analogues of (3-11)–(3-13) for positive perturbations imply

$$\begin{aligned}
 & n_+((1 + \varepsilon)\lambda(\Lambda_q + \lambda)^{-1}\Lambda_q^{-1}; P_q T_+ P_q) + O(1) \\
 & \leq N_{(\Lambda_+^{-1}, (\Lambda_q + \lambda)^{-1})}(H_0^{-1} - T_+) \leq n_+((1 - \varepsilon)\lambda(\Lambda_q + \lambda)^{-1}\Lambda_q^{-1}; P_q T_+ P_q) + O(1), \quad \lambda \downarrow 0. \quad (3-15)
 \end{aligned}$$

By the resolvent identity, we have $T_{\pm} = H_0^{-1}WH_0^{-1} \mp H_0^{-1}WH_{\pm}^{-1}WH_0^{-1}$, so that

$$P_q T_{\pm} P_q = \Lambda_q^{-2}(P_q W P_q \mp P_q W H_{\pm}^{-1} W P_q).$$

Thus,

$$n_+(s; P_q T_{\pm} P_q) = n_+(s\Lambda_q^2; P_q W P_q \mp P_q W H_{\pm}^{-1} W P_q), \quad s > 0. \quad (3-16)$$

Putting together (3-5)–(3-8) and (3-14)–(3-16), we easily obtain (3-4). □

4. Unitary equivalence for Berezin–Toeplitz operators

Our first goal in this section is to show that, under certain regularity conditions on the matrix m , the operator $P_q W P_q$, $q \in \mathbb{Z}_+$, with domain $P_q L^2(\mathbb{R}^2)$, is unitarily equivalent to $P_0 w_q P_0$ with domain $P_0 L^2(\mathbb{R}^2)$, where w_q is the multiplier by a suitable function $w_q : \mathbb{R}^2 \rightarrow \mathbb{C}$. In fact, we will need a slightly more general result, and that is why we introduce first the appropriate notations.

As usual, for $x = (x_1, x_2) \in \mathbb{R}^2$ we set $z := x_1 + ix_2$ and $\bar{z} := x_1 - ix_2$, so that

$$\frac{\partial}{\partial z} = \frac{1}{2} \left(\frac{\partial}{\partial x_1} - i \frac{\partial}{\partial x_2} \right), \quad \frac{\partial}{\partial \bar{z}} = \frac{1}{2} \left(\frac{\partial}{\partial x_1} + i \frac{\partial}{\partial x_2} \right).$$

Introduce the magnetic annihilation operator

$$a := -2ie^{-b|x|^2/4} \frac{\partial}{\partial \bar{z}} e^{b|x|^2/4} = -2i \left(\frac{\partial}{\partial \bar{z}} + \frac{bz}{4} \right)$$

and the magnetic creation operator

$$a^* := -2ie^{b|x|^2/4} \frac{\partial}{\partial z} e^{-b|x|^2/4} = -2i \left(\frac{\partial}{\partial z} - \frac{b\bar{z}}{4} \right)$$

with common domain $\text{Dom } a = \text{Dom } a^* = \text{Dom } H_0^{1/2}$. The operators a and a^* are closed and mutually adjoint in $L^2(\mathbb{R}^2)$. On $\text{Dom } H_0$ we have $[a, a^*] = 2b$ and

$$H_0 = a^* a + b = a a^* - b = \frac{1}{2}(a a^* + a^* a). \tag{4-1}$$

Moreover, on $\text{Dom } H_0^{1/2}$ we have

$$\Pi_1 = \frac{1}{2}(a + a^*), \quad \Pi_2 = \frac{1}{2i}(a - a^*), \tag{4-2}$$

the operators Π_j , $j = 1, 2$, being introduced in (1-1). Next, define the operator $\mathbb{A} : \text{Dom } H_0^{1/2} \rightarrow L^2(\mathbb{R}^2; \mathbb{C}^2)$ by

$$\mathbb{A}u := \begin{pmatrix} a^* u \\ au \end{pmatrix}, \quad u \in \text{Dom } H_0^{1/2}.$$

Then, (4-1) implies that $H_0 = \frac{1}{2}\mathbb{A}^*\mathbb{A}$. Further, introduce the Hermitian matrix-valued function

$$\Omega := \begin{pmatrix} \omega_{11} & \omega_{12} \\ \omega_{21} & \omega_{22} \end{pmatrix}$$

with $\omega_{jk} \in L^\infty(\mathbb{R}^2)$, $j, k = 1, 2$. Fix $q \in \mathbb{Z}_+$ and define the operator

$$P_q \mathbb{A}^* \Omega \mathbb{A} P_q = \Lambda_q P_q H_0^{-1/2} \mathbb{A}^* \Omega \mathbb{A} H_0^{-1/2} P_q, \tag{4-3}$$

which is bounded and self-adjoint in $P_q L^2(\mathbb{R}^2)$. Utilizing (4-2), we easily find that

$$P_q W P_q = \frac{1}{2} P_q \mathbb{A}^* U \mathbb{A} P_q, \tag{4-4}$$

where

$$U := \mathbb{O}^* m \mathbb{O}, \quad \mathbb{O} := \frac{1}{\sqrt{2}} \begin{pmatrix} 1 & 1 \\ i & -i \end{pmatrix}, \tag{4-5}$$

so that

$$U = \begin{pmatrix} u_{11} & u_{12} \\ u_{21} & u_{22} \end{pmatrix} \quad \text{with} \quad \begin{aligned} u_{11} &:= \frac{1}{2}(\text{Tr } m - 2 \text{Im } m_{12}), \\ u_{22} &:= \frac{1}{2}(\text{Tr } m + 2 \text{Im } m_{12}), \\ u_{12} = \bar{u}_{21} &:= \frac{1}{2}(m_{11} - m_{22} - 2i \text{Re } m_{12}). \end{aligned}$$

Introduce the Laguerre polynomials

$$L_q^{(m)} := \sum_{j=0}^q \binom{q+m}{q-j} \frac{(-t)^j}{j!}, \quad t \in \mathbb{R}, \quad q \in \mathbb{Z}_+, \quad m \in \mathbb{Z}_+; \tag{4-6}$$

as usual, we write $L_q^{(0)} = L_q$, and for notational convenience we set $qL_{q-1} = 0$ for $q = 0$. By [Gradshteyn and Ryzhik 1965, Equation 8.974.3] we have

$$\sum_{j=0}^q L_j^{(m)}(t) = L_q^{(m+1)}(t), \quad t \in \mathbb{R}, \quad q \in \mathbb{Z}_+, \quad m \in \mathbb{Z}_+. \tag{4-7}$$

Proposition 4.1. *Let Ω be a Hermitian 2×2 matrix-valued function with entries $\omega_{jk} \in C_b^\infty(\mathbb{R}^2)$, $j, k = 1, 2$. Fix $q \in \mathbb{Z}_+$. Then the operator $P_q \mathbb{A}^* \Omega \mathbb{A} P_q$ with domain $P_q L^2(\mathbb{R}^2)$ is unitarily equivalent to the operator $P_0 w_q P_0$ with domain $P_0 L^2(\mathbb{R}^2)$, where*

$$\begin{aligned} w_q &= w_q(\Omega) \\ &:= \begin{cases} 2b(q+1)L_{q+1}\left(-\frac{\Delta}{2b}\right)\omega_{11} + 2bqL_{q-1}\left(-\frac{\Delta}{2b}\right)\omega_{22} - 8 \text{Re } L_{q-1}^{(2)}\left(-\frac{\Delta}{2b}\right)\frac{\partial^2 \omega_{12}}{\partial \bar{z}^2} & \text{if } q \geq 1, \\ 2bL_1\left(-\frac{\Delta}{2b}\right)\omega_{11} & \text{if } q = 0, \end{cases} \end{aligned} \tag{4-8}$$

and $\Delta = \sum_{j=1,2} \partial^2 / \partial x_j^2$, so that, in accordance with (4-6), $L_s^{(m)}(-\Delta/(2b))$ with $s \in \mathbb{Z}_+$ and $m \in \mathbb{Z}_+$ is just the differential operator $\sum_{j=0}^s \binom{s+m}{s-j} \Delta^j / (j!(2b)^j)$ of order $2s$ with constant coefficients.

Proof. Set

$$\begin{aligned} \varphi_{0,k}(x) &:= \sqrt{\frac{b}{2\pi k!}} \left(\frac{b}{2}\right)^{k/2} z^k e^{-b|x|^2/4}, \quad x \in \mathbb{R}^2, \quad k \in \mathbb{Z}_+, \\ \varphi_{q,k}(x) &:= \sqrt{\frac{1}{(2b)^q q!}} (a^*)^q \varphi_{0,k}(x), \quad x \in \mathbb{R}^2, \quad k \in \mathbb{Z}_+, \quad q \in \mathbb{N}. \end{aligned}$$

Then $\{\varphi_{q,k}\}_{k \in \mathbb{Z}_+}$ is an orthonormal basis of $P_q L^2(\mathbb{R}^2)$, sometimes called *the angular momentum basis* (see, e.g., [Raikov and Warzel 2002] or [Bruneau et al. 2004, Section 9.1]). Evidently, for $k \in \mathbb{Z}_+$ we have

$$a^* \varphi_{q,k} = \sqrt{2b(q+1)} \varphi_{q+1,k}, \quad q \in \mathbb{Z}_+, \quad a \varphi_{q,k} = \begin{cases} \sqrt{2bq} \varphi_{q-1,k}, & q \geq 1, \\ 0, & q = 0. \end{cases} \tag{4-9}$$

Define the unitary operator $\mathcal{W} : P_q L^2(\mathbb{R}^2) \rightarrow P_0 L^2(\mathbb{R}^2)$ by $\mathcal{W} : u \mapsto v$, where

$$u = \sum_{k \in \mathbb{Z}_+} c_k \varphi_{q,k}, \quad v = \sum_{k \in \mathbb{Z}_+} c_k \varphi_{0,k}, \quad \{c_k\}_{k \in \mathbb{Z}_+} \in \ell^2(\mathbb{Z}_+). \tag{4-10}$$

We will show that

$$P_q \mathbb{A}^* \Omega \mathbb{A} P_q = \mathcal{W}^* P_0 w_q P_0 \mathcal{W}. \tag{4-11}$$

For $V \in C_b^\infty(\mathbb{R}^2)$, $m, s \in \mathbb{Z}_+$, and $k, \ell \in \mathbb{Z}_+$, set

$$\Xi_{m,s}(V; k, \ell) := \langle V \varphi_{m,k}, \varphi_{s,\ell} \rangle,$$

where $\langle \cdot, \cdot \rangle$ denotes the scalar product in $L^2(\mathbb{R}^2)$. Taking into account (4-9) and (4-10), we easily find that

$$\begin{aligned} \langle P_q \mathbb{A}^* \Omega \mathbb{A} P_q u, u \rangle &= 2b \sum_{k \in \mathbb{Z}_+} \sum_{\ell \in \mathbb{Z}_+} ((q+1) \Xi_{q+1,q+1}(\omega_{11}; k, \ell) + q \Xi_{q-1,q-1}(\omega_{22}; k, \ell)) c_k \bar{c}_\ell \\ &\quad + 2b \sqrt{q(q+1)} 2 \operatorname{Re} \sum_{k \in \mathbb{Z}_+} \sum_{\ell \in \mathbb{Z}_+} \Xi_{q+1,q-1}(\omega_{21}; k, \ell) c_k \bar{c}_\ell \end{aligned} \tag{4-12}$$

if $q \geq 1$, and

$$\langle P_0 \mathbb{A}^* \Omega \mathbb{A} P_0 u, u \rangle = 2b \sum_{k \in \mathbb{Z}_+} \sum_{\ell \in \mathbb{Z}_+} \Xi_{1,1}(\omega_{11}; k, \ell) c_k \bar{c}_\ell. \tag{4-13}$$

Moreover,

$$\langle P_0 w_q P_0 v, v \rangle = \sum_{k \in \mathbb{Z}_+} \sum_{\ell \in \mathbb{Z}_+} \Xi_{0,0}(w_q; k, \ell) c_k \bar{c}_\ell, \quad q \in \mathbb{Z}_+. \tag{4-14}$$

In [Bruneau et al. 2004, Lemma 9.2] (see also the remark after Equation (2.2) in [Bony et al. 2014]), it was shown that

$$\Xi_{m,m}(V; k, \ell) = \Xi_{0,0} \left(L_m \left(-\frac{\Delta}{2b} \right) V; k, \ell \right), \quad m \in \mathbb{Z}_+. \tag{4-15}$$

Now (4-13), (4-15) with $m = 1$ and $V = \omega_{11}$, and (4-14) with $q = 0$ imply (4-11) in the case $q = 0$. Assume $q \geq 1$. By (4-15), we have

$$\Xi_{q+1,q+1}(\omega_{11}; k, \ell) = \Xi_{0,0} \left(L_{q+1} \left(-\frac{\Delta}{2b} \right) \omega_{11}; k, \ell \right), \tag{4-16}$$

$$\Xi_{q-1,q-1}(\omega_{22}; k, \ell) = \Xi_{0,0} \left(L_{q-1} \left(-\frac{\Delta}{2b} \right) \omega_{22}; k, \ell \right). \tag{4-17}$$

Let us now consider the quantity $\Xi_{q+1,q-1}(V; k, \ell)$. Using (4-9), we easily find that, for $q \geq 2$, we have

$$\Xi_{q+1,q-1}(V; k, \ell) = \frac{1}{\sqrt{2b(q+1)}} \Xi_{q,q-1}([V, a^*]; k, \ell) + \sqrt{\frac{q-1}{q+1}} \Xi_{q,q-2}(V; k, \ell), \tag{4-18}$$

$$\Xi_{q,q-1}([V, a^*]; k, \ell) = \frac{1}{\sqrt{2bq}} \Xi_{q-1,q-1}([[V, a^*], a^*]; k, \ell) + \sqrt{\frac{q-1}{q}} \Xi_{q-1,q-2}([V, a^*]; k, \ell). \tag{4-19}$$

Moreover, $[V, a^*] = 2i \partial V / \partial z$, and

$$[[V, a^*], a^*] = -4 \frac{\partial^2 V}{\partial z^2}. \tag{4-20}$$

Using (4-19), it is not difficult to prove by induction that

$$\mathfrak{E}_{q,q-1}([V, a^*]; k, \ell) = \frac{1}{\sqrt{2bq}} \sum_{j=0}^{q-1} \mathfrak{E}_{j,j}([V, a^*], a^*; k, \ell), \quad q \geq 1. \tag{4-21}$$

Now (4-15), (4-20), and (4-7) imply

$$\begin{aligned} \sum_{j=0}^{q-1} \mathfrak{E}_{j,j}([V, a^*], a^*; k, \ell) &= \sum_{j=0}^{q-1} \mathfrak{E}_{0,0} \left(-4L_j \left(-\frac{\Delta}{2b} \right) \frac{\partial^2 V}{\partial z^2}; k, \ell \right) \\ &= \mathfrak{E}_{0,0} \left(-4L_{q-1}^{(1)} \left(-\frac{\Delta}{2b} \right) \frac{\partial^2 V}{\partial z^2}; k, \ell \right). \end{aligned} \tag{4-22}$$

Setting

$$\mathfrak{D}_q := -4L_{q-1}^{(1)} \left(-\frac{\Delta}{2b} \right) \frac{\partial^2}{\partial z^2}, \quad q \in \mathbb{N}, \tag{4-23}$$

we find that (4-21) and (4-22) imply

$$\mathfrak{E}_{q,q-1}([V, a^*]; k, \ell) = \frac{1}{\sqrt{2bq}} \mathfrak{E}_{0,0}(\mathfrak{D}_q V; k, \ell). \tag{4-24}$$

Bearing in mind (4-18), (4-15), and (4-24), it is not difficult to prove by induction that

$$\mathfrak{E}_{q+1,q-1}(V; k, \ell) = \frac{1}{2b\sqrt{q(q+1)}} \sum_{s=1}^q \mathfrak{E}_{0,0}(\mathfrak{D}_s V; k, \ell). \tag{4-25}$$

Note that (4-7) and (4-25) imply

$$\sum_{s=1}^q \mathfrak{D}_s = -4L_{q-1}^{(2)} \left(-\frac{\Delta}{2b} \right) \frac{\partial^2}{\partial z^2}. \tag{4-26}$$

Now, (4-25) and (4-26) entail

$$2b\sqrt{q(q+1)} \mathfrak{E}_{q+1,q-1}(\omega_{21}; k, \ell) = \mathfrak{E}_{0,0} \left(-4L_{q-1}^{(2)} \left(-\frac{\Delta}{2b} \right) \frac{\partial^2 \omega_{21}}{\partial z^2}; k, \ell \right). \tag{4-27}$$

Finally, (4-12) and (4-14) combined with (4-16), (4-17), and (4-27) yield (4-11) with $q \geq 1$. □

In the rest of the section we establish two other suitable representations for the operators $P_q V P_q$, $q \in \mathbb{Z}_+$, with $V : \mathbb{R}^2 \rightarrow \mathbb{C}$.

Proposition 4.2. (i) [Fernández and Raikov 2004, Lemma 3.1; Bony et al. 2014, Section 2.3] *Let $V \in L^1_{\text{loc}}(\mathbb{R}^2)$ satisfy $\lim_{|x| \rightarrow \infty} V(x) = 0$. Then, for each $q \in \mathbb{Z}_+$, the operator $P_q V P_q$ is compact.*

(ii) [Raikov and Warzel 2002, Lemma 3.3] Assume in addition that V is radially symmetric, i.e., there exists $v : [0, \infty) \rightarrow \mathbb{C}$ such that $V(x) = v(|x|)$, $x \in \mathbb{R}^2$. Then the eigenvalues of the operator $P_q V P_q$ with domain $P_q L^2(\mathbb{R}^2)$, counted with multiplicities, coincide with the set

$$\{\langle V \varphi_{q,k}, \varphi_{q,k} \rangle\}_{k \in \mathbb{Z}_+}. \tag{4-28}$$

In particular, the eigenvalues of $P_0 V P_0$ coincide with

$$\frac{1}{k!} \int_0^\infty v\left(\left(\frac{2t}{b}\right)^{\frac{1}{2}}\right) e^{-t} t^k dt, \quad k \in \mathbb{Z}_+. \tag{4-29}$$

Remarks. (i) Let us recall that, if f is, say, a bounded function of exponential decay, then

$$(\mathcal{M}f)(z) := \int_0^\infty f(t) t^{z-1} dt, \quad z \in \mathbb{C}, \operatorname{Re} z > 0,$$

is sometimes called *the Mellin transform* of f . Some of the asymptotic properties as $k \rightarrow \infty$ of the integrals (4-29), which we will later obtain and use in the proofs of Theorems 2.1 and 2.2, could possibly be deduced from the general theory of the Mellin transform.

(ii) Combining Propositions 4.1 and 4.2, we find that, if the matrix-valued function Ω is radially symmetric and diagonal, then the operator $P_q \mathbb{A}^* \Omega \mathbb{A} P_q$ acting in $P_q L^2(\mathbb{R}^2)$ is unitarily equivalent to a *diagonal* operator in $\ell^2(\mathbb{Z}_+)$. If Ω is just radially symmetric, then $P_q \mathbb{A}^* \Omega \mathbb{A} P_q$ is unitarily equivalent to a *tridiagonal* operator acting in $\ell^2(\mathbb{Z}_+)$.

The last proposition in this section concerns the unitary equivalence between the Berezin–Toeplitz operator $P_0 W P_0$ and a certain Weyl pseudodifferential operator. Let us recall the definition of Weyl pseudodifferential operators acting in $L^2(\mathbb{R})$. Denote by $\Gamma(\mathbb{R}^2)$ the set of functions $\psi : \mathbb{R}^2 \rightarrow \mathbb{C}$ such that

$$\|\psi\|_{\Gamma(\mathbb{R}^2)} := \sup_{(y,\eta) \in \mathbb{R}^2} \sup_{\ell,m=0,1} \left| \frac{\partial^{\ell+m} \psi(y, \eta)}{\partial y^\ell \partial \eta^m} \right| < \infty.$$

Then the operator $\operatorname{Op}^w(\psi)$, defined initially as a mapping between the Schwartz class $\mathcal{S}(\mathbb{R})$ and its dual class $\mathcal{S}'(\mathbb{R})$ by

$$(\operatorname{Op}^w(\psi)u)(y) = \frac{1}{2\pi} \int_{\mathbb{R}} \int_{\mathbb{R}} \psi\left(\frac{y+y'}{2}, \eta\right) e^{i(y-y')\eta} u(y') dy' d\eta, \quad y \in \mathbb{R},$$

extends uniquely to an operator bounded in $L^2(\mathbb{R})$. Moreover, there exists a constant c such that

$$\|\operatorname{Op}^w(\psi)\| \leq c \|\psi\|_{\Gamma(\mathbb{R}^2)} \tag{4-30}$$

(see, e.g., [Boulkhemair 1999, Corollary 2.5(i)]).

Remark. Inequalities of the type (4-30) are known as *Calderón–Vaillancourt estimates*.

Put

$$\mathcal{R}_b := -b^{-1/2} \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \tag{4-31}$$

and, for $V : \mathbb{R}^2 \rightarrow \mathbb{C}$, define

$$V_b(x) := V(\mathcal{R}_b x), \quad x \in \mathbb{R}^2, \quad b > 0.$$

Moreover, set $\mathcal{G}(x) := e^{-|x|^2}/\pi, x \in \mathbb{R}^2$.

Proposition 4.3 [Pushnitski et al. 2013, Theorem 2.11, Corollary 2.8]. *Let $V \in L^1(\mathbb{R}^2) + L^\infty(\mathbb{R}^2)$. Then the operator $P_0 V P_0$ with domain $P_0 L^2(\mathbb{R}^2)$ is unitarily equivalent to the operator $\text{Op}^w(V_b * \mathcal{G})$.*

Remark. The operator $\text{Op}^{\text{aw}}(\psi) := \text{Op}^w(\psi * \mathcal{G})$ is called a pseudodifferential operator with *anti-Wick symbol* ψ (see, e.g., [Shubin 2001, Section 24]).

5. Proofs of Theorems 2.1 and 2.2

In this section we complete the proofs of Theorems 2.1 and 2.2, concerning perturbations of compact support and of exponential decay.

Let $T = T^*$ be a compact operator in a Hilbert space such that $\text{rank } \mathbb{1}_{(0,\infty)}(T) = \infty$. Denote by $\{v_k(T)\}_{k=0}^\infty$ the nonincreasing sequence of the positive eigenvalues of T , counted with multiplicities.

Recall that $m_<(x) \leq m_>(x)$ are the eigenvalues of the matrix $m(x)$ for $x \in \mathbb{R}^2$. Since the matrix U (see (4-5)) is unitarily equivalent to m , m_\geq are also the eigenvalues of U . Next, we check that Proposition 3.1 implies the following:

Corollary 5.1. *Under the general assumptions of the article, there exist constants $0 < c_{<}^\pm \leq c_{>}^\pm < \infty$ and $k_0 \in \mathbb{Z}_+$ such that*

$$c_{<}^\pm v_{k+k_0}(P_q \mathbb{A}^* m_{<} \mathbb{A} P_q) \leq \pm(\lambda_{k,q}^\pm - \Lambda_q) \leq c_{>}^\pm v_{k-k_0}(P_q \mathbb{A}^* m_{>} \mathbb{A} P_q) \tag{5-1}$$

for sufficiently large $k \in \mathbb{N}$.

Proof. It is easy to see that

$$0 \leq P_q W H_\pm^{-1} W P_q \leq c_\pm P_q W P_q \tag{5-2}$$

with

$$c_\pm := \|H_\pm^{-1/2} W H_\pm^{-1/2}\| \leq \sup_{x \in \mathbb{R}^2} |m(x)(I \pm m(x))^{-1}|.$$

Note that $0 \leq c_- < \infty$ and $0 \leq c_+ < 1$. Moreover, by (4-4) and the mini-max principle,

$$n_+(2s; P_q \mathbb{A}^* m_{<} \mathbb{A} P_q) \leq n_+(s; P_q W P_q) \leq n_+(2s; P_q \mathbb{A}^* m_{>} \mathbb{A} P_q), \quad s > 0. \tag{5-3}$$

Now, (3-4), (5-2), and (5-3) imply that, for any $\varepsilon \in (0, 1)$, we have

$$n_+(2\lambda(1 + \varepsilon); P_q \mathbb{A}^* m_{<} \mathbb{A} P_q) + O(1) \leq \mathcal{N}_q^-(\lambda) \leq n_+(2\lambda(1 - \varepsilon); (1 + c_-) P_q \mathbb{A}^* m_{>} \mathbb{A} P_q) + O(1), \tag{5-4}$$

$$n_+(2\lambda(1 - \varepsilon); P_q \mathbb{A}^* m_{>} \mathbb{A} P_q) + O(1) \geq \mathcal{N}_q^+(\lambda) \geq n_+(2\lambda(1 + \varepsilon); (1 - c_+) P_q \mathbb{A}^* m_{<} \mathbb{A} P_q) + O(1) \tag{5-5}$$

as $\lambda \downarrow 0$, and estimates (5-4)–(5-5) yield (5-1) with

$$c_{<}^- = \frac{1}{2(1 + \varepsilon)}, \quad c_{>}^- = \frac{1 + c_-}{2(1 - \varepsilon)}, \quad c_{<}^+ = \frac{1 - c_+}{2(1 + \varepsilon)}, \quad c_{>}^+ = \frac{1}{2(1 - \varepsilon)},$$

and sufficiently large $k_0 \in \mathbb{N}$. □

Let us now complete the proof of Theorem 2.1. Let $\zeta_1 \in C_0^\infty(\mathbb{R}^2)$, $\zeta_1 \geq 0$, $\zeta_1 = 1$ on $\text{supp } m_>$. Set $\zeta_2(x) := (\max_{y \in \mathbb{R}^2} m_>(y))\zeta_1(x)$, $x \in \mathbb{R}^2$. Evidently, $m_> \leq \zeta_2$ on \mathbb{R}^2 , so that

$$\nu_k(P_q \mathbb{A}^* m_> \mathbb{A} P_q) \leq \nu_k(P_q \mathbb{A}^* \zeta_2 \mathbb{A} P_q), \quad k \in \mathbb{Z}_+. \tag{5-6}$$

Further, by Proposition 4.1, the operator $P_q \mathbb{A}^* \zeta_2 \mathbb{A} P_q$ is unitarily equivalent to the operator $P_0 \zeta_3 P_0$, where

$$\zeta_3 := 2b \left((q+1) L_{q+1} \left(-\frac{\Delta}{2b} \right) + q L_{q-1} \left(-\frac{\Delta}{2b} \right) \right) \zeta_2.$$

Therefore,

$$\nu_k(P_q \mathbb{A}^* \zeta_2 \mathbb{A} P_q) = \nu_k(P_0 \zeta_3 P_0), \quad k \in \mathbb{Z}_+. \tag{5-7}$$

Let $R_> > 0$ be so large that the disk $B_{R_>}(0)$ of radius $R_>$ centered at the origin contains the support of ζ_3 . Then,

$$\nu_k(P_0 \zeta_3 P_0) \leq \max_{x \in \mathbb{R}^2} |\zeta_3(x)| \nu_k(P_0 \mathbb{1}_{B_{R_>}(0)} P_0), \quad k \in \mathbb{Z}_+. \tag{5-8}$$

Putting together (5-6), (5-7), and (5-8), we find that there exists a constant $K_> < \infty$ such that

$$\nu_k(P_q \mathbb{A}^* m_> \mathbb{A} P_q) \leq K_> \nu_k(P_0 \mathbb{1}_{B_{R_>}(0)} P_0), \quad k \in \mathbb{Z}_+. \tag{5-9}$$

On the other hand,

$$\nu_k(P_q \mathbb{A}^* m_< \mathbb{A} P_q) \geq \nu_k(P_q a m_< a^* P_q). \tag{5-10}$$

Applying (4-9), we easily find that the operators $P_q a m_< a^* P_q$ and $2b(q+1) P_{q+1} m_< P_{q+1}$ are unitarily equivalent. Hence,

$$\nu_k(P_q a m_< a^* P_q) = 2b(q+1) \nu_k(P_{q+1} m_< P_{q+1}), \quad k \in \mathbb{Z}_+. \tag{5-11}$$

Further, since $m_<$ is nonnegative, continuous, and does not vanish identically, there exist $c_0 > 0$, $R_< \in (0, \infty)$, and $x_0 \in \mathbb{R}^2$ such that $m_<(x) \geq c_0 \mathbb{1}_{B_{R_<}(x_0)}(x)$, $x \in \mathbb{R}^2$. Therefore,

$$\nu_k(P_{q+1} m_< P_{q+1}) \geq c_0 \nu_k(P_{q+1} \mathbb{1}_{B_{R_<}(x_0)} P_{q+1}), \quad k \in \mathbb{Z}_+. \tag{5-12}$$

The operators $P_{q+1} \mathbb{1}_{B_{R_<}(x_0)} P_{q+1}$ and $P_{q+1} \mathbb{1}_{B_{R_<}(0)} P_{q+1}$ are unitarily equivalent under the magnetic translation which maps x_0 into 0 (see, e.g., [Raikov and Warzel 2002, Equation (4.21)]). Therefore,

$$\nu_k(P_{q+1} \mathbb{1}_{B_{R_<}(x_0)} P_{q+1}) = \nu_k(P_{q+1} \mathbb{1}_{B_{R_<}(0)} P_{q+1}), \quad k \in \mathbb{Z}_+. \tag{5-13}$$

Combining (5-10)–(5-13), we find that there exists a constant $K_<$ such that

$$K_< \nu_k(P_{q+1} \mathbb{1}_{B_{R_<}(0)} P_{q+1}) \leq \nu_k(P_q \mathbb{A}^* m_< \mathbb{A} P_q), \quad k \in \mathbb{Z}_+. \tag{5-14}$$

By (5-9) and (5-14), it remains to study the asymptotic behavior as $k \rightarrow \infty$ of $\nu_k(P_m \mathbb{1}_{B_R(0)} P_m)$, with $m \in \mathbb{Z}_+$ and $R \in (0, \infty)$ fixed. This asymptotic analysis relies on the representation (4-28), and results sufficient for our purposes are available in the literature. Namely, we have:

Lemma 5.2 [Combes et al. 2004, Section 4, Corollary 2]. *Let $m \in \mathbb{Z}_+$, $R \in (0, \infty)$, $b \in (0, \infty)$. Set $\varrho := bR^2/2$. Then*

$$v_k(P_m \mathbb{1}_{B_R(0)} P_m) = \frac{e^{-\varrho} \varrho^{-m+1} k^{2m-1} \varrho^k}{m!k!} (1 + o(1)), \quad k \rightarrow \infty. \tag{5-15}$$

Now, asymptotic relation (2-1) follows from (5-1), (5-9), (5-14), (5-15), and the elementary fact that $\ln k! = k \ln k + O(k)$ as $k \rightarrow \infty$.

In the remaining part of this section we prove Theorem 2.2 concerning perturbations m of exponential decay. Assume that m satisfies (2-5). Then there exist $\delta_{\geq} \in \mathbb{R}$, $\delta_{<} \leq \delta_{>}$, and $r > 1$ such that

$$|x|^{\delta_{<}} e^{-\gamma|x|^{2\beta}} \mathbb{1}_{\mathbb{R}^2 \setminus B_r(0)}(x) \leq m_{<}(x) \leq m_{>}(x) \leq |x|^{\delta_{>}} e^{-\gamma|x|^{2\beta}} \mathbb{1}_{\mathbb{R}^2 \setminus B_r(0)}(x) + \max_{y \in \mathbb{R}^2} m_{>}(y) \mathbb{1}_{B_r(0)}(x), \quad x \in \mathbb{R}^2. \tag{5-16}$$

Let $\eta_{\geq,0} \in C^\infty(\mathbb{R}^2; [0, 1])$ be two radially symmetric functions such that $\eta_{<,0} = 1$ on $\mathbb{R}^2 \setminus B_{r+1}(0)$, $\eta_{<,0} = 0$ on $B_r(0)$, and $\eta_{>,0} = 1$ on $\mathbb{R}^2 \setminus B_r(0)$, $\eta_{>,0} = 0$ on $B_{r-1}(0)$. For $x \in \mathbb{R}^2$ set

$$\begin{aligned} \eta_{<,1}(x) &:= |x|^{\delta_{<}} e^{-\gamma|x|^{2\beta}} \eta_{<,0}(x), \\ \eta_{>,1}(x) &:= |x|^{\delta_{>}} e^{-\gamma|x|^{2\beta}} \eta_{>,0}(x) + \max_{y \in \mathbb{R}^2} m_{>}(y) (1 - \eta_{<,0}(x)). \end{aligned}$$

Evidently, $\eta_{\geq,1} \in C_b^\infty(\mathbb{R}^2)$, and by (5-16),

$$\eta_{<,1}(x) \leq m_{<}(x), \quad m_{>}(x) \leq \eta_{>,1}(x), \quad x \in \mathbb{R}^2.$$

Therefore, for $k \in \mathbb{Z}_+$, we have

$$\begin{aligned} v_k(P_q \mathbb{A}^* m_{<} \mathbb{A} P_q) &\geq v_k(P_q \mathbb{A}^* \eta_{<,1} \mathbb{A} P_q), \\ v_k(P_q \mathbb{A}^* m_{>} \mathbb{A} P_q) &\leq v_k(P_q \mathbb{A}^* \eta_{>,1} \mathbb{A} P_q). \end{aligned} \tag{5-17}$$

Further, set

$$\eta_{\geq,2} := 2b \left((q+1) L_{q+1} \left(-\frac{\Delta}{2b} \right) + q L_{q-1} \left(-\frac{\Delta}{2b} \right) \right) \eta_{\geq,1}.$$

According to Proposition 4.1, the operators $P_q \mathbb{A}^* \eta_{\geq,1} \mathbb{A} P_q$, $q \in \mathbb{Z}_+$, and $P_0 \eta_{\geq,2} P_0$ are unitarily equivalent. Therefore,

$$v_k(P_q \mathbb{A}^* \eta_{\geq,1} \mathbb{A} P_q) = v_k(P_0 \eta_{\geq,2} P_0), \quad k \in \mathbb{Z}_+. \tag{5-18}$$

Next, a tedious but straightforward calculation shows that

$$\eta_{\geq,2}(x) = \eta_{\geq,3}(x) (1 + o(1)), \quad |x| \rightarrow \infty, \tag{5-19}$$

where

$$\eta_{\geq,3}(x) := C_{q,\beta} |x|^{\delta_{\geq}} e^{-\gamma|x|^{2\beta}} \begin{cases} 1 & \text{if } \beta \in (0, \frac{1}{2}], \\ |x|^{2(q+1)(2\beta-1)} & \text{if } \beta \in (\frac{1}{2}, \infty), \end{cases} \quad x \in \mathbb{R}^2 \setminus \{0\},$$

and $C_{q,\beta} > 0$ are some constants. Even though the exact values of $C_{q,\beta}$ will not play any role in the sequel, we indicate here these values for the sake of the completeness of the exposition:

$$C_{q,\beta} = \begin{cases} 2\Lambda_q & \text{if } \beta \in (0, \frac{1}{2}), \\ 2b\left((q+1)L_{q+1}\left(-\frac{(2\beta\gamma)^2}{2b}\right) + qL_{q-1}\left(-\frac{(2\beta\gamma)^2}{2b}\right)\right) & \text{if } \beta = \frac{1}{2}, \\ \frac{(2\beta\gamma)^{2(q+1)}}{(2b)^q q!} & \text{if } \beta \in (\frac{1}{2}, \infty). \end{cases}$$

Hence, by (5-19), there exists $R \in (0, \infty)$ such that for $x \in \mathbb{R}^2$ we have

$$\eta_{<,2} \geq \frac{1}{2}\eta_{<,3}\mathbb{1}_{\mathbb{R}^2 \setminus B_R(0)} - c_{<}\mathbb{1}_{B_R(0)} =: \eta_{<,4}(x), \tag{5-20}$$

$$\eta_{>,2} \leq \frac{3}{2}\eta_{>,3}\mathbb{1}_{\mathbb{R}^2 \setminus B_R(0)} + c_{>}\mathbb{1}_{B_R(0)} =: \eta_{>,4}(x) \tag{5-21}$$

with $c_{\geq} := \max_{y \in \mathbb{R}^2} |\eta_{\geq,2}(y)|$. Thus, for any admissible $k \in \mathbb{Z}_+$, we have

$$\nu_k(P_0\eta_{<,2}P_0) \geq \nu_k(P_0\eta_{<,4}P_0), \quad \nu_k(P_0\eta_{>,2}P_0) \leq \nu_k(P_0\eta_{>,4}P_0). \tag{5-22}$$

In order to complete the proof of Theorem 2.2, we need a couple of auxiliary results. For $\beta > 0, \mu > 0$, and $\varrho > 0$, set

$$\mathcal{F}_{\beta,\mu}(k) := \int_0^\infty e^{-\mu t^\beta - t} t^k dt, \quad \mathcal{E}_\varrho(k) := \int_0^\varrho e^{-t} t^k dt, \quad k > -1, \tag{5-23}$$

and, for $\delta \in \mathbb{R}, c_0 > 0$ and $c_1 \in \mathbb{R}$, put

$$\mathcal{L}(k) = \mathcal{L}_{\beta,\mu,\varrho,\delta}(k; c_0, c_1) := \frac{c_0\mathcal{F}_{\beta,\mu}(k+\delta) + c_1\mathcal{E}_\varrho(k-\delta_-)}{\Gamma(k+1)}, \quad k > \max\{-1, -\delta-1\},$$

where $\delta_- := \max\{0, -\delta\}$.

Lemma 5.3. *Let $\beta > 0, \mu > 0, \varrho > 0, c_0 > 0$, and $\delta \in \mathbb{R}, c_1 \in \mathbb{R}$.*

(i) *The asymptotic relations*

$$\ln \mathcal{L}(k) = \begin{cases} -\sum_{1 \leq j < 1/(1-\beta)} f_j k^{(\beta-1)j+1} + O(\ln k) & \text{if } \beta \in (0, 1), \\ -\ln(1+\mu)k + O(\ln k) & \text{if } \beta = 1, \\ -\frac{\beta-1}{\beta}k \ln k + k\left(\frac{\beta-1-\ln(\mu\beta)}{\beta}\right) & \\ \quad -\sum_{1 \leq j < \beta/(\beta-1)} g_j k^{(1/\beta-1)j+1} + O(\ln k) & \text{if } \beta \in (1, \infty), \end{cases} \tag{5-24}$$

hold true as $k \rightarrow \infty$, the coefficients f_j and g_j being introduced in the statement of Theorem 2.2.

(ii) *We have $\mathcal{L}'(k) < 0$ for sufficiently large k .*

Proof. First, let $\delta = 0$. Assume $\beta \in (0, 1), k > 0$, and make the change of variable $t \mapsto ks$ in the first integral in (5-23). Thus we find that

$$\mathcal{F}_{\beta,\mu}(k) = k^{k+1} \int_0^\infty e^{-kF(s;k^{\beta-1})} ds. \tag{5-25}$$

The function $F(s; k^{\beta-1})$ defined in (2-9) attains its unique minimum at $s_{<}(k^{\beta-1})$, and we have $\partial^2 F(s_{<}(k^{\beta-1}); k^{\beta-1})/\partial s^2 = 1 + o(1)$, $k \rightarrow \infty$. Therefore, applying a standard argument close to the usual Laplace method for asymptotic evaluation of integrals depending on a large parameter, we easily find that

$$\int_0^\infty e^{-kF(s; k^{\beta-1})} ds = (2\pi)^{1/2} e^{-kF(s_{<}(k^{\beta-1}); k^{\beta-1})} k^{-1/2} (1 + o(1)), \quad k \rightarrow \infty. \tag{5-26}$$

Bearing in mind that $F(s_{<}(k^{\beta-1}); k^{\beta-1}) = f(k^{\beta-1})$ (see (2-10)), $f(0) = 1$, and

$$\ln \Gamma(k + 1) = k \ln k - k + \frac{1}{2} \ln k + O(1), \quad k \rightarrow \infty, \tag{5-27}$$

(see, e.g., [Abramowitz and Stegun 1964, Equation 6.1.40]), we find that (5-25)–(5-26) imply

$$\begin{aligned} \ln \left(\frac{\mathcal{F}_{\beta, \mu}(k)}{\Gamma(k + 1)} \right) &= k - kf(k^{\beta-1}) + O(\ln k) \\ &= k - k \sum_{0 \leq j < 1/(1-\beta)} \frac{1}{j!} \frac{d^j f}{d\epsilon^j}(0) k^{(\beta-1)j} + O(\ln k) \\ &= - \sum_{1 \leq j < 1/(1-\beta)} \frac{1}{j!} \frac{d^j f}{d\epsilon^j}(0) k^{(\beta-1)j+1} + O(\ln k) \\ &= - \sum_{1 \leq j < 1/(1-\beta)} f_j k^{(\beta-1)j+1} + O(\ln k), \quad k \rightarrow \infty. \end{aligned} \tag{5-28}$$

In the case $\beta = 1$, we simply have

$$\frac{\mathcal{F}_{\beta, \mu}(k)}{\Gamma(k + 1)} = \frac{1}{\Gamma(k + 1)} \int_0^\infty e^{-(\mu+1)t} t^k dt = (\mu + 1)^{-k-1},$$

that is,

$$\ln \left(\frac{\mathcal{F}_{\beta, \mu}(k)}{\Gamma(k + 1)} \right) = -(\ln(1 + \mu))k + O(1), \quad k \rightarrow \infty. \tag{5-29}$$

Now let $\beta \in (1, \infty)$. Making the change of variable $t \mapsto k^{1/\beta} s$ with $k > 0$ in (5-23), we find

$$\mathcal{F}_{\beta, \mu}(k) := k^{(k+1)/\beta} \int_0^\infty e^{-kG(s; k^{1/\beta-1})} ds. \tag{5-30}$$

The function $G(s; k^{1/\beta-1})$ defined in (2-11), attains its unique minimum at $s_{>}(k^{1/\beta-1})$, and we have

$$\frac{\partial^2 G}{\partial s^2}(s_{>}(k^{1/\beta-1}), k^{1/\beta-1}) = \beta(\mu\beta)^{2/\beta} (1 + o(1)), \quad k \rightarrow \infty.$$

Arguing as in the derivation of (5-26), we obtain

$$\int_0^\infty e^{-kG(s; k^{1/\beta-1})} ds = \sqrt{2\pi\beta} (\mu\beta)^{-1/\beta} e^{-kG(s_{>}(k^{1/\beta-1}); k^{1/\beta-1})} k^{-1/2} (1 + o(1)), \quad k \rightarrow \infty. \tag{5-31}$$

Bearing in mind that $G(s_{>}(k^{1/\beta-1}); k^{1/\beta-1}) = g(k^{1/\beta-1})$ (see (2-12)), and $g(0) = (1 + \ln(\mu\beta))/\beta$, we find that (5-30), (5-31), and (5-27) imply

$$\begin{aligned} \ln\left(\frac{\mathcal{F}_{\beta,\mu}(k)}{\Gamma(k+1)}\right) &= -\frac{\beta-1}{\beta}k \ln k + k - kg(k^{1/\beta-1}) + O(\ln k) \\ &= -\frac{\beta-1}{\beta}k \ln k + k - k \sum_{0 \leq j < \beta/(\beta-1)} \frac{1}{j!} \frac{d^j g}{d\epsilon^j}(0) k^{(1/\beta-1)j} + O(\ln k) \\ &= -\frac{\beta-1}{\beta}k \ln k + k(1 - g(0)) - \sum_{1 \leq j < \beta/(\beta-1)} \frac{1}{j!} \frac{d^j g}{d\epsilon^j}(0) k^{(1/\beta-1)j+1} + O(\ln k) \\ &= -\frac{\beta-1}{\beta}k \ln k + k \left(\frac{\beta-1 - \ln(\mu\beta)}{\beta}\right) - \sum_{1 \leq j < \beta/(\beta-1)} g_j k^{(1/\beta-1)j+1} + O(\ln k), \end{aligned} \tag{5-32}$$

as $k \rightarrow \infty$. Let us now consider general $\delta \in \mathbb{R}$. By (5-27),

$$\ln\left(\frac{\Gamma(k+\delta+1)}{\Gamma(k+1)}\right) = \delta \ln k + O(1), \quad k \rightarrow \infty. \tag{5-33}$$

Putting together (5-28), (5-29), (5-32), and (5-33), we find that

$$\ln\left(\frac{\mathcal{F}_{\beta,\mu}(k+\delta)}{\Gamma(k+1)}\right) - \ln\left(\frac{\mathcal{F}_{\beta,\mu}(k)}{\Gamma(k+1)}\right) = O(\ln k), \quad k \rightarrow \infty. \tag{5-34}$$

Finally, by (5-15), we easily find that, for each fixed $\delta \in \mathbb{R}$, we have

$$\frac{\mathcal{E}_\rho(k-\delta_-)}{\Gamma(k+1)} = o\left(\frac{\mathcal{F}_{\beta,\mu}(k+\delta)}{\Gamma(k+1)}\right), \quad k \rightarrow \infty. \tag{5-35}$$

The combination of (5-28), (5-29), (5-32), (5-34), and (5-35) implies (5-24).

For (ii), we have

$$\begin{aligned} \mathcal{L}'(k) &= c_0 \left(\frac{\mathcal{F}'_{\beta,\mu}(k+\delta)}{\Gamma(k+1)} - \frac{\Gamma'(k+1)}{\Gamma(k+1)^2} \mathcal{F}_{\beta,\mu}(k+\delta) \right) + c_1 \left(\frac{\mathcal{E}'_\rho(k-\delta_-)}{\Gamma(k+1)} - \frac{\Gamma'(k+1)}{\Gamma(k+1)^2} \mathcal{E}_\rho(k-\delta_-) \right), \\ \mathcal{F}'_{\beta,\mu}(k) &= \int_0^\infty e^{-\mu t^\beta - t} t^k \ln t \, dt, \quad \mathcal{E}'_\rho(k) = \int_0^\rho e^{-t} t^k \ln t \, dt, \end{aligned} \tag{5-36}$$

and

$$\frac{\Gamma'(k+1)}{\Gamma(k+1)} = \ln k + \frac{1}{2k} + O(k^{-2}), \quad k \rightarrow \infty,$$

(see, e.g., [Abramowitz and Stegun 1964, Equation 6.3.18]). Performing an asymptotic analysis similar to the one in the proof of the first part of the lemma, we find that there exists a function $\Psi = \Psi_{\beta,\mu,\delta}$ such that $\Psi(k) < 0$ for k large enough, and

$$\frac{\mathcal{F}'_{\beta,\mu}(k+\delta)}{\Gamma(k+1)} - \frac{\Gamma'(k+1)}{\Gamma(k+1)^2} \mathcal{F}_{\beta,\mu}(k+\delta) = \Psi(k)(1 + o(1)), \tag{5-37}$$

$$\frac{\mathcal{E}'_\rho(k-\delta_-)}{\Gamma(k+1)} - \frac{\Gamma'(k+1)}{\Gamma(k+1)^2} \mathcal{E}_\rho(k-\delta_-) = o(\Psi(k)) \tag{5-38}$$

as $k \rightarrow \infty$. Putting together (5-36), (5-37), and (5-38), we conclude that $\mathcal{L}'(k) < 0$ for sufficiently large k . \square

Taking into account the definition of the functions $\eta_{\geq,4}$ in (5-20)–(5-21), the mini-max principle, representation (4-29), as well as Lemma 5.3(ii), we find that there exist constants $c_{j,\geq} > 0$, $j = 0, 1$, $\tilde{\delta}_{\geq} \in \mathbb{R}$, and $k_0 \in \mathbb{Z}_+$ such that

$$\begin{aligned} v_k(P_0\eta_{<,4}P_0) &\geq \mathcal{L}_{\beta,\mu,\varrho,\tilde{\delta}_<}(k+k_0; c_{0,<}, -c_{1,<}) \\ v_k(P_0\eta_{>,4}P_0) &\leq \mathcal{L}_{\beta,\mu,\varrho,\tilde{\delta}_>}(k; c_{0,>}, c_{1,>}), \end{aligned} \tag{5-39}$$

for $\mu = \gamma(2/b)^\beta$, $\varrho = bR^2/2$, and sufficiently large $k \in \mathbb{Z}_+$.

Putting together (5-1), (5-17), (5-18), (5-22), (5-39), and (5-24), we obtain (2-6)–(2-8).

6. Proof of Theorem 2.3

Estimates (3-4) combined with the Weyl inequalities (3-3) and the mini-max principle entail

$$\begin{aligned} n_+(\lambda(1+\varepsilon); P_q W P_q) + O(1) \\ \leq N_q^-(\lambda) \leq n_+(\lambda(1-\varepsilon)^2; P_q W P_q) + n_+(\lambda\varepsilon(1-\varepsilon); P_q W H_-^{-1} W P_q) + O(1), \end{aligned} \tag{6-1}$$

and

$$\begin{aligned} n_+(\lambda(1+\varepsilon)^2; P_q W P_q) - n_+(\lambda\varepsilon(1+\varepsilon); P_q W H_+^{-1} W P_q) + O(1) \\ \leq N_q^+(\lambda) \leq n_+(\lambda(1-\varepsilon); P_q W P_q) + O(1) \end{aligned} \tag{6-2}$$

as $\lambda \downarrow 0$. It is easy to check that we have

$$P_q W H_\pm^{-1} W P_q \leq C_{1,\pm} P_q \mathbb{A}^* \langle \cdot \rangle^{-2\rho} \mathbb{A} P_q$$

with

$$C_{1,\pm} := \|H_0^{1/2} H_\pm^{-1/2}\|^2 \left(\sup_{x \in \mathbb{R}^2} \langle x \rangle^\rho m_\pm(x) \right)^2.$$

Therefore, for any $s > 0$,

$$n_+(s; P_q W H_\pm^{-1} W P_q) \leq n_+(s; C_{1,\pm} P_q \mathbb{A}^* \langle \cdot \rangle^{-2\rho} \mathbb{A} P_q). \tag{6-3}$$

Further, by Proposition 4.1, the operator $P_q W P_q$ (resp. $P_q \mathbb{A}^* \langle \cdot \rangle^{-2\rho} \mathbb{A} P_q$) is unitarily equivalent to $\frac{1}{2} P_0 w_q(U) P_0$ (resp. $P_0 w_q(\langle \cdot \rangle^{-2\rho} I) P_0$). Hence, for any $s > 0$,

$$n_+(s; P_q W P_q) = n_+(2s; P_0 w_q(U) P_0), \tag{6-4}$$

$$n_+(s; P_q \mathbb{A}^* \langle \cdot \rangle^{-2\rho} \mathbb{A} P_q) = n_+(s; P_0 w_q(\langle \cdot \rangle^{-2\rho} I) P_0) \leq n_+(s; C_2 P_0 \langle \cdot \rangle^{-2\rho} P_0) \tag{6-5}$$

with $C_2 := \sup_{x \in \mathbb{R}^2} \langle x \rangle^{2\rho} |w_q(\langle x \rangle^{-2\rho} I)|$. Now, write

$$\frac{1}{2} w_q(U) = \mathcal{T}_q + \tilde{\mathcal{T}}_q,$$

the symbol \mathcal{T}_q being defined in (2-16), and note the crucial circumstance that $\tilde{\mathcal{T}}_q \in \mathcal{S}^{-\rho-2}(\mathbb{R}^2)$. Then the Weyl inequalities (3-3) entail

$$\begin{aligned} n_+(s(1 + \varepsilon); P_0\mathcal{T}_q P_0) - n_-(s\varepsilon; P_0\tilde{\mathcal{T}}_q P_0) &\leq n_+(2s; P_0w_q(U)P_0) \\ &\leq n_+(s(1 - \varepsilon); P_0\mathcal{T}_q P_0) + n_+(s\varepsilon; P_0\tilde{\mathcal{T}}_q P_0) \end{aligned} \tag{6-6}$$

for any $s > 0$ and $\varepsilon \in (0, 1)$. Evidently,

$$n_{\pm}(s; P_0\tilde{\mathcal{T}}_q P_0) \leq n_+(s; C_3 P_0\langle \cdot \rangle^{-\rho-2} P_0), \quad s > 0, \tag{6-7}$$

with $C_3 := \sup_{x \in \mathbb{R}^2} \langle x \rangle^{\rho+2} |\tilde{\mathcal{T}}_q(x)|$. Recalling Proposition 4.3, we find that we have reduced the asymptotic analysis of $N_q^{\pm}(\lambda)$ as $\lambda \downarrow 0$ to the eigenvalue asymptotics for a pseudodifferential operator with elliptic anti-Wick symbol of negative order. The spectral asymptotics for operators of this type has been extensively studied in the literature since the 1970s. In particular, we have the following:

Proposition 6.1. *Let $0 \leq \psi \in \mathcal{S}^{-\rho}(\mathbb{R}^2)$, $\rho > 0$. Assume that there exists $0 < \psi_0 \in C^\infty(\mathbb{S}^1)$ such that $\lim_{|x| \rightarrow \infty} |x|^\rho \psi(x) = \psi_0(x/|x|)$. Then we have*

$$n_+(\lambda; \text{Op}^{\text{aw}}(\psi)) = (2\pi)^{-1} \Phi_\psi(\lambda)(1 + o(1)), \quad \lambda \downarrow 0, \tag{6-8}$$

which is equivalent to

$$\lim_{\lambda \downarrow 0} \lambda^{2/\rho} n_+(\lambda; \text{Op}^{\text{aw}}(\psi)) = \mathcal{C}(\psi_0) := \frac{1}{4\pi} \int_0^{2\pi} \psi_0(\cos \theta, \sin \theta)^{2/\rho} d\theta.$$

Proof. Evidently, for each $\varepsilon \in (0, 1)$ there exist real functions $\psi_{\pm, \varepsilon} \in C^\infty(\mathbb{R}^2)$ such that

$$\begin{aligned} \psi_{-, \varepsilon}(x) &\leq \psi(x) \leq \psi_{+, \varepsilon}(x), \quad x \in \mathbb{R}^2, \\ \psi_{\pm, \varepsilon}(x) &= (1 \mp \varepsilon)^{-1} |x|^{-\rho} \psi_0\left(\frac{x}{|x|}\right), \quad x \in \mathbb{R}^2, |x| \geq R, \end{aligned}$$

for some $R \in (0, \infty)$. Applying the monotonicity of the anti-Wick quantization with respect to the symbol (see, e.g., [Shubin 2001, Proposition 24.1]), the mini-max principle, and the Weyl inequalities, we obtain

$$\begin{aligned} n_+((1 + \varepsilon)\lambda; \text{Op}^{\text{w}}(\psi_{-, \varepsilon})) - n_-(\varepsilon\lambda; (\text{Op}^{\text{aw}}(\psi_{-, \varepsilon}) - \text{Op}^{\text{w}}(\psi_{-, \varepsilon}))) \\ \leq n_+(\lambda; \text{Op}^{\text{aw}}(\psi)) \leq n_+((1 - \varepsilon)\lambda; \text{Op}^{\text{w}}(\psi_{+, \varepsilon})) + n_+(\varepsilon\lambda; (\text{Op}^{\text{aw}}(\psi_{+, \varepsilon}) - \text{Op}^{\text{w}}(\psi_{+, \varepsilon}))). \end{aligned} \tag{6-9}$$

By [Dauge and Robert 1987], we have the semiclassical result

$$n_+(\lambda; \text{Op}^{\text{w}}(\psi_{\pm, \varepsilon})) = (2\pi)^{-1} \Phi_{\psi_{\pm, \varepsilon}}(\lambda)(1 + o(1)), \quad \lambda \downarrow 0. \tag{6-10}$$

Further, by [Shubin 2001, Theorem 24.1] the differences $\text{Op}^{\text{aw}}(\psi_{\pm, \varepsilon}) - \text{Op}^{\text{w}}(\psi_{\pm, \varepsilon})$ are pseudodifferential operators of lower order than $\text{Op}^{\text{w}}(\psi_{\pm, \varepsilon})$, so that we easily obtain

$$\lim_{\lambda \downarrow 0} \lambda^{2/\rho} n_{\pm}(\varepsilon\lambda; (\text{Op}^{\text{aw}}(\psi_{\pm, \varepsilon}) - \text{Op}^{\text{w}}(\psi_{\pm, \varepsilon}))) = 0, \quad \varepsilon > 0. \tag{6-11}$$

Now, (6-9)–(6-11) imply

$$(1 + \varepsilon)^{-4/\rho} \mathcal{C}(\psi_0) \leq \liminf_{\lambda \downarrow 0} \lambda^{2/\rho} n_+(\lambda; \text{Op}^{\text{aw}}(\psi)) \leq \limsup_{\lambda \downarrow 0} \lambda^{2/\rho} n_+(\lambda; \text{Op}^{\text{aw}}(\psi)) \leq (1 - \varepsilon)^{-4/\rho} \mathcal{C}(\psi_0)$$

for $\varepsilon \in (0, 1)$. Letting $\varepsilon \downarrow 0$, we obtain (6-8). □

By Propositions 4.3 and 6.1, we have

$$n_+(\lambda; P_0 \mathcal{T}_q P_0) = n_+(\lambda; \text{Op}^{\text{aw}}(\mathcal{T}_{q,b})) = \frac{1}{2\pi} \Phi_{\mathcal{T}_{q,b}}(\lambda)(1+o(1)) = \frac{b}{2\pi} \Phi_{\mathcal{T}_q}(\lambda)(1+o(1)), \quad \lambda \downarrow 0, \quad (6-12)$$

with $\mathcal{T}_{q,b} = \mathcal{T}_q \circ \mathcal{R}_b$, \mathcal{R}_b being defined in (4-31). Finally, for $\rho_0 > \rho$, we have

$$n_+(\lambda; P_0 \langle \cdot \rangle^{-\rho_0} P_0) = O(\lambda^{-2/\rho_0}) = o(\Phi_{\mathcal{T}_q}(\lambda)), \quad \lambda \downarrow 0. \quad (6-13)$$

Now, (2-17) easily follows from (6-1)–(6-8), (6-12), and (6-13). The equivalence of (2-18) and (2-19) can be checked by arguing as in the proof of [Shubin 2001, Proposition 13.1].

Appendix: Compactness of the resolvent differences

A priori, the operators H_0 and H_{\pm} , self-adjoint in $L^2(\mathbb{R}^2)$, could be defined as the Friedrichs extensions of the operators $\sum_{j=1,2} \Pi_j^2$ and $\sum_{j,k=1,2} \Pi_j g_{jk}^{\pm} \Pi_k$ defined on $C_0^\infty(\mathbb{R}^2)$. Such a definition implies immediately that

$$\text{Dom } H_0^{1/2} = \text{Dom } H_{\pm}^{1/2} = \{u \in L^2(\mathbb{R}^2) \mid \Pi_j u \in L^2(\mathbb{R}^2), j = 1, 2\},$$

and that the operators $H_{\pm}^{1/2} H_0^{-1/2}$ and $H_0^{1/2} H_{\pm}^{-1/2}$ are bounded. By [Gérard et al. 1991, Proposition A.2], the operators H_0 and H_{\pm} are essentially self-adjoint on $C_0^\infty(\mathbb{R}^2)$ and have a common domain

$$\text{Dom } H_0 = \text{Dom } H_{\pm} = \{u \in L^2(\mathbb{R}^2) \mid \Pi_j \Pi_k u \in L^2(\mathbb{R}^2), j, k = 1, 2\}.$$

Let us now prove the compactness of the operator $H_0^{-1} - H_{\pm}^{-1}$ in $L^2(\mathbb{R}^2)$. Since we have

$$H_0^{-1} - H_{\pm}^{-1} = \pm H_0^{-1} W H_{\pm}^{-1} = \pm H_0^{-1} W H_0^{-1} H_0 H_{\pm}^{-1},$$

it suffices to prove the compactness of $H_0^{-1} W H_0^{-1}$. The operators $H_0^{-1} W H_0^{-1} = \frac{1}{2} H_0^{-1} \mathbb{A}^* U \mathbb{A} H_0^{-1}$ and $\frac{1}{2} H_0^{-1} \mathbb{A}^* m_{>} \mathbb{A} H_0^{-1}$ are bounded, self-adjoint, and positive. Moreover,

$$H_0^{-1} \mathbb{A}^* U \mathbb{A} H_0^{-1} \leq H_0^{-1} \mathbb{A}^* m_{>} \mathbb{A} H_0^{-1}. \quad (A-1)$$

On the other hand,

$$H_0^{-1} \mathbb{A}^* m_{>} \mathbb{A} H_0^{-1} = H_0^{-1} a^* m_{>} a H_0^{-1} + H_0^{-1} a m_{>} a^* H_0^{-1}. \quad (A-2)$$

By (A-1) and (A-2), it suffices to prove the compactness of the operator $m_{>}^{1/2} a^* H_0^{-1}$. We have

$$m_{>}^{1/2} a^* H_0^{-1} = m_{>}^{1/2} H_0^{-1/2} (H_0^{-1/2} a^* + 2b H_0^{-1/2} a^* H_0^{-1}).$$

The operator $H_0^{-1/2} a^* + 2b H_0^{-1/2} a^* H_0^{-1}$ is bounded, so that it suffices to prove the compactness of $m_{>}^{1/2} H_0^{-1/2}$ which follows from $m_{>} \in L^\infty(\mathbb{R}^2)$, $\lim_{|x| \rightarrow \infty} m_{>}(x) = 0$, and the diamagnetic inequality (see, e.g., [Avron et al. 1978, Theorem 2.5]).

Acknowledgements

The final version of this work has been done during the authors' visit to the Isaac Newton Institute, Cambridge, UK, in January 2015. The authors thank the Newton Institute for financial support and hospitality. The partial support by the Chilean Scientific Foundation *Fondecyt* under Grant 1130591, by *Núcleo Milenio de Física Matemática* RC120002, and by the Faculty of Mathematics, PUC, Santiago de Chile, is gratefully acknowledged as well.

References

- [Abramowitz and Stegun 1964] M. Abramowitz and I. A. Stegun, *Handbook of mathematical functions with formulas, graphs, and mathematical tables*, National Bureau of Standards Applied Mathematics Series **55**, U.S. Government Printing Office, Washington, D.C., 1964. MR 29 #4914 Zbl 0171.38503
- [Alama et al. 1989] S. Alama, P. A. Deift, and R. Hempel, "Eigenvalue branches of the Schrödinger operator $H - \lambda W$ in a gap of $\sigma(H)$ ", *Comm. Math. Phys.* **121**:2 (1989), 291–321. MR 90e:35046 Zbl 0676.47032
- [Alama et al. 1994] S. Alama, M. Avellaneda, P. A. Deift, and R. Hempel, "On the existence of eigenvalues of a divergence-form operator $A + \lambda B$ in a gap of $\sigma(A)$ ", *Asymptotic Anal.* **8**:4 (1994), 311–344. MR 95g:47069 Zbl 0806.47042
- [Avron et al. 1978] J. Avron, I. Herbst, and B. Simon, "Schrödinger operators with magnetic fields, I: General interactions", *Duke Math. J.* **45**:4 (1978), 847–883. MR 80k:35054 Zbl 0399.35029
- [Bastard et al. 1975] G. Bastard, J. K. Furdyna, and J. Mycielski, "Landau levels and cyclotron resonance in graded mixed semiconductors", *Phys. Rev. B* **12** (1975), 4356–4359.
- [Birman and Solomjak 1987] M. S. Birman and M. Z. Solomjak, *Spectral theory of selfadjoint operators in Hilbert space*, Reidel, Dordrecht, 1987. MR 93g:47001
- [Bony et al. 2014] J.-F. Bony, V. Bruneau, and G. Raikov, "Resonances and spectral shift function singularities for magnetic quantum Hamiltonians", pp. 77–100 in *Spectral and scattering theory and related topics*, edited by F. Hiroshima, RIMS Kôkyûroku Bessatsu **B45**, Res. Inst. Math. Sci. (RIMS), Kyoto, 2014. MR 3287673 Zbl 1301.35074
- [Boulkhemair 1999] A. Boulkhemair, " L^2 estimates for Weyl quantization", *J. Functional Anal.* **165**:1 (1999), 173–204. MR 2001f:35454 Zbl 0934.35217
- [Boyarchenko and Levendorskiĭ 1997] S. I. Boyarchenko and S. Z. Levendorskiĭ, "An asymptotic formula for the number of eigenvalue branches of a divergence form operator $A + \lambda B$ in a spectral gap of A ", *Comm. Partial Differential Equations* **22**:11–12 (1997), 1771–1786. MR 99e:35165 Zbl 0899.35070
- [Briet et al. 2009] P. Briet, H. Kovařík, G. Raikov, and E. Soccorsi, "Eigenvalue asymptotics in a twisted waveguide", *Comm. Partial Differential Equations* **34**:7–9 (2009), 818–836. MR 2010j:35089 Zbl 1195.35119
- [Bruneau et al. 2004] V. Bruneau, A. Pushnitski, and G. Raikov, "Spectral shift function in strong magnetic fields", *Algebra i Analiz* **16**:1 (2004), 207–238. Reprinted in *St. Petersburg Math. J.* **16** (2005), 181–209. MR 2005h:81118 Zbl 1082.35115
- [Colin de Verdière 1986] Y. Colin de Verdière, "L'asymptotique de Weyl pour les bouteilles magnétiques", *Comm. Math. Phys.* **105**:2 (1986), 327–335. MR 87k:58273 Zbl 0612.35102
- [Combes et al. 2004] J.-M. Combes, P. D. Hislop, F. Klopp, and G. Raikov, "Global continuity of the integrated density of states for random Landau Hamiltonians", *Comm. Partial Differential Equations* **29**:7–8 (2004), 1187–1213. MR 2005g:81091 Zbl 1085.47050
- [Dauge and Robert 1987] M. Dauge and D. Robert, "Weyl's formula for a class of pseudodifferential operators with negative order on $L^2(\mathbb{R}^n)$ ", pp. 91–122 in *Pseudodifferential operators* (Oberwolfach, 1986), edited by H. O. Cordes et al., Lecture Notes in Math. **1256**, Springer, Berlin, 1987. MR 88k:35150 Zbl 0629.35098
- [Fernández and Raikov 2004] C. Fernández and G. Raikov, "On the singularities of the magnetic spectral shift function at the Landau levels", *Ann. Henri Poincaré* **5**:2 (2004), 381–403. MR 2005g:81092 Zbl 1062.81043

- [Filonov and Pushnitski 2006] N. Filonov and A. Pushnitski, “Spectral asymptotics of Pauli operators and orthogonal polynomials in complex domains”, *Comm. Math. Phys.* **264**:3 (2006), 759–772. MR 2007h:81064 Zbl 1106.81040
- [Fock 1928] V. Fock, “Bemerkung zur Quantelung des harmonischen Oszillators im Magnetfeld”, *Z. Physik* **47**:5–6 (1928), 446–448. JFM 54.0966.02
- [Gadella and Smolyanov 2008] M. Gadella and O. G. Smolyanov, “Feynman formulas for particles with position-dependent mass”, *Dokl. Akad. Nauk* **418**:6 (2008), 727–730. In Russian; translated in *Dokl. Math.* **77**:1 (2008), 120–123. MR 2009m:81125 Zbl 1159.35426
- [Gérard et al. 1991] C. Gérard, A. Martinez, and J. Sjöstrand, “A mathematical approach to the effective Hamiltonian in perturbed periodic problems”, *Comm. Math. Phys.* **142**:2 (1991), 217–244. MR 93a:47079 Zbl 0753.35057
- [Gradshteyn and Ryzhik 1965] I. S. Gradshteyn and I. M. Ryzhik, *Table of integrals, series, and products*, Academic Press, New York, 1965. Zbl 1300.65001
- [Ivrii 1998] V. Ivrii, *Microlocal analysis and precise spectral asymptotics*, Springer, Berlin, 1998. MR 99e:58193 Zbl 0906.35003
- [Killingbeck 2011] J. P. Killingbeck, “The Schrödinger equation with position-dependent mass”, *J. Phys. A* **44**:28 (2011), 285208, 6. MR 2012i:81090 Zbl 1222.81163
- [Landau 1930] L. Landau, “Diamagnetismus der Metalle”, *Z. Physik* **64**:9–10 (1930), 629–637.
- [Landkof 1972] N. S. Landkof, *Foundations of modern potential theory*, Grundlehren der Math. Wissenschaften **180**, Springer, New York, 1972. MR 50 #2520 Zbl 0253.31001
- [Mao 2012] S. Mao, “Singularities for solutions to Schrödinger equations with asymptotically constant magnetic fields”, *J. Math. Phys.* **53**:7 (2012), Art. ID #073707. MR 2985266 Zbl 1276.81054
- [Midya et al. 2010] B. Midya, B. Roy, and R. Roychoudhury, “Position dependent mass Schrödinger equation and isospectral potentials: intertwining operator approach”, *J. Math. Phys.* **51**:2 (2010), Art. ID #022109. MR 2011g:81089 Zbl 1309.81082
- [Pushnitski and Rozenblum 2011] A. Pushnitski and G. Rozenblum, “On the spectrum of Bargmann–Toeplitz operators with symbols of a variable sign”, *J. Anal. Math.* **114** (2011), 317–340. MR 2837088 Zbl 1253.47018
- [Pushnitski et al. 2013] A. Pushnitski, G. Raikov, and C. Villegas-Blas, “Asymptotic density of eigenvalue clusters for the perturbed Landau Hamiltonian”, *Comm. Math. Phys.* **320**:2 (2013), 425–453. MR 3053767 Zbl 1277.47059
- [Raikov 1990] G. D. Raikov, “Eigenvalue asymptotics for the Schrödinger operator with homogeneous magnetic potential and decreasing electric potential, I: Behaviour near the essential spectrum tips”, *Comm. Partial Differential Equations* **15**:3 (1990), 407–434. MR 91k:35188 Zbl 0739.35055
- [Raikov 2015] G. D. Raikov, “Spectral asymptotics for waveguides with perturbed periodic twisting”, preprint, 2015. To appear in *J. Spectral Theory*. arXiv 1407.0757
- [Raikov and Warzel 2002] G. D. Raikov and S. Warzel, “Quasi-classical versus non-classical spectral asymptotics for magnetic Schrödinger operators with decreasing electric potentials”, *Rev. Math. Phys.* **14**:10 (2002), 1051–1072. MR 2003m:81069 Zbl 1033.81038
- [von Roos 1983] O. von Roos, “Position-dependent effective masses in semiconductor theory”, *Phys. Rev. B* **27** (1983), 7547–7552.
- [Rosenberg 1997] S. Rosenberg, *The Laplacian on a Riemannian manifold: an introduction to analysis on manifolds*, London Mathematical Society Student Texts **31**, Cambridge University Press, 1997. MR 98k:58206 Zbl 0868.58074
- [Rozenblum and Tashchiyan 2008] G. Rozenblum and G. Tashchiyan, “On the spectral properties of the perturbed Landau Hamiltonian”, *Comm. Partial Differential Equations* **33**:4–6 (2008), 1048–1081. MR 2009e:81073 Zbl 1158.47029
- [Shubin 2001] M. A. Shubin, *Pseudodifferential operators and spectral theory*, 2nd ed., Springer, Berlin, 2001. MR 2002d:47073 Zbl 0980.35180
- [de Souza Dutra and de Oliveira 2009] A. de Souza Dutra and J. A. de Oliveira, “Two-dimensional position-dependent massive particles in the presence of magnetic fields”, *J. Phys. A* **42**:2 (2009), 025304, 13. MR 2010i:82200 Zbl 1154.82030
- [Sternberg 1964] S. Sternberg, *Lectures on differential geometry*, Prentice-Hall, Englewood Cliffs, N.J., 1964. MR 33 #1797 Zbl 0129.13102

Received 3 Dec 2014. Revised 30 Jan 2015. Accepted 15 Apr 2015.

TOMÁS LUNGENSTRASS: tlungens@mat.puc.cl

Facultad de Matemáticas, Pontificia Universidad Católica de Chile, Vicuña Mackenna 4860, 7820436 Santiago, Chile

GEORGI RAIKOV: graikov@mat.puc.cl

Facultad de Matemáticas, Pontificia Universidad Católica de Chile, Vicuña Mackenna 4860, 7820436 Santiago, Chile

Analysis & PDE

msp.org/apde

EDITORS

EDITOR-IN-CHIEF

Maciej Zworski
zworski@math.berkeley.edu
University of California
Berkeley, USA

BOARD OF EDITORS

Nicolas Burq	Université Paris-Sud 11, France nicolas.burq@math.u-psud.fr	Yuval Peres	University of California, Berkeley, USA peres@stat.berkeley.edu
Sun-Yung Alice Chang	Princeton University, USA chang@math.princeton.edu	Gilles Pisier	Texas A&M University, and Paris 6 pisier@math.tamu.edu
Michael Christ	University of California, Berkeley, USA mchrist@math.berkeley.edu	Tristan Rivière	ETH, Switzerland riviere@math.ethz.ch
Charles Fefferman	Princeton University, USA cf@math.princeton.edu	Igor Rodnianski	Princeton University, USA irod@math.princeton.edu
Ursula Hamenstaedt	Universität Bonn, Germany ursula@math.uni-bonn.de	Wilhelm Schlag	University of Chicago, USA schlag@math.uchicago.edu
Vaughan Jones	U.C. Berkeley & Vanderbilt University vaughan.f.jones@vanderbilt.edu	Sylvia Serfaty	New York University, USA serfaty@cims.nyu.edu
Herbert Koch	Universität Bonn, Germany koch@math.uni-bonn.de	Yum-Tong Siu	Harvard University, USA siu@math.harvard.edu
Izabella Laba	University of British Columbia, Canada ilaba@math.ubc.ca	Terence Tao	University of California, Los Angeles, USA tao@math.ucla.edu
Gilles Lebeau	Université de Nice Sophia Antipolis, France lebeau@unice.fr	Michael E. Taylor	Univ. of North Carolina, Chapel Hill, USA met@math.unc.edu
László Lempert	Purdue University, USA lempert@math.purdue.edu	Gunther Uhlmann	University of Washington, USA gunther@math.washington.edu
Richard B. Melrose	Massachusetts Institute of Technology, USA rbm@math.mit.edu	András Vasy	Stanford University, USA andras@math.stanford.edu
Frank Merle	Université de Cergy-Pontoise, France Frank.Merle@u-cergy.fr	Dan Virgil Voiculescu	University of California, Berkeley, USA dvv@math.berkeley.edu
William Minicozzi II	Johns Hopkins University, USA minicozz@math.jhu.edu	Steven Zelditch	Northwestern University, USA zelditch@math.northwestern.edu
Werner Müller	Universität Bonn, Germany mueller@math.uni-bonn.de		

PRODUCTION

production@msp.org
Silvio Levy, Scientific Editor

See inside back cover or msp.org/apde for submission instructions.

The subscription price for 2015 is US \$205/year for the electronic version, and \$390/year (+\$55, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues from the last three years and changes of subscribers address should be sent to MSP.

Analysis & PDE (ISSN 1948-206X electronic, 2157-5045 printed) at Mathematical Sciences Publishers, 798 Evans Hall #3840, c/o University of California, Berkeley, CA 94720-3840, is published continuously online. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices.

APDE peer review and production are managed by EditFLOW[®] from MSP.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2015 Mathematical Sciences Publishers

ANALYSIS & PDE

Volume 8 No. 5 2015

Partial collapsing and the spectrum of the Hodge–de Rham operator COLETTE ANNÉ and JUNYA TAKAHASHI	1025
Sharp L^p bounds for the wave equation on groups of Heisenberg type DETLEF MÜLLER and ANDREAS SEEGER	1051
Global well-posedness on the derivative nonlinear Schrödinger equation YIFEI WU	1101
On the boundary value problem for the Schrödinger equation: compatibility conditions and global existence CORENTIN AUDIARD	1113
On estimates for fully nonlinear parabolic equations on Riemannian manifolds BO GUAN, SHUJUN SHI and ZHENAN SUI	1145
Concentration phenomena for the nonlocal Schrödinger equation with Dirichlet datum JUAN DÁVILA, MANUEL DEL PINO, SERENA DIPIERRO and ENRICO VALDINOCI	1165
Local spectral asymptotics for metric perturbations of the Landau Hamiltonian TOMÁS LUNGENSTRASS and GEORGI RAIKOV	1237
Hilbert transform along measurable vector fields constant on Lipschitz curves: L^2 boundedness SHAOMING GUO	1263



2157-5045(2015)8:5;1-9