

ANALYSIS & PDE

Volume 9

No. 2

2016

PHAN THÀNH NAM, NICOLAS ROUGERIE AND ROBERT SEIRINGER

**GROUND STATES OF LARGE BOSONIC SYSTEMS:
THE GROSS-PITAEVSKII LIMIT REVISITED**

GROUND STATES OF LARGE BOSONIC SYSTEMS: THE GROSS–PITAEVSKII LIMIT REVISITED

PHAN THÀNH NAM, NICOLAS ROUGERIE AND ROBERT SEIRINGER

We study the ground state of a dilute Bose gas in a scaling limit where the Gross–Pitaevskii functional emerges. This is a repulsive nonlinear Schrödinger functional whose quartic term is proportional to the scattering length of the interparticle interaction potential. We propose a new derivation of this limit problem, with a method that bypasses some of the technical difficulties that previous derivations had to face. The new method is based on a combination of Dyson’s lemma, the quantum de Finetti theorem and a second moment estimate for ground states of the effective Dyson Hamiltonian. It applies equally well to the case where magnetic fields or rotation are present.

1. Introduction	459
2. Overall strategy	462
3. Second moment estimate	467
4. Energy lower bound and convergence of states	475
Acknowledgments	484
References	484

1. Introduction

The rigorous derivation of effective nonlinear theories from many-body quantum mechanics has been studied extensively in recent years, motivated in part by experiments in cold atom physics. For bosons, the emergence of the limit theories can be interpreted as due to most of the particles occupying the same quantum state: this is the Bose–Einstein condensation phenomenon, observed first in dilute alkali vapors some twenty years ago.

The parameter regime most relevant for the description of the actual physical setup is the Gross–Pitaevskii limit. It is also the most mathematically demanding regime considered in the literature so far; see [Lieb and Yngvason 1998; Lieb et al. 2000; Lieb and Seiringer 2002; 2006] for the derivation of equilibrium states and [Erdős et al. 2009; 2010; Benedikter et al. 2015; Pickl 2015] for dynamics (more extensive lists of references may be found in [Lieb et al. 2005b; Rougerie 2015; Benedikter et al. 2016]). The main reason for this sophistication is the fact that interparticle correlations due to two-body scattering play a leading-order role in this regime. The goal of this paper is to present a method for the derivation of Gross–Pitaevskii theory at the level of the ground state that is conceptually and technically simpler

MSC2010: 35Q40, 81V70.

Keywords: many-body quantum mechanics, mean-field limits, Bose–Einstein condensates.

than existing proofs, in particular that of [Lieb and Seiringer 2006], which was so far the only method applicable when an external magnetic field is present.

Our setting is as follows: we consider N interacting bosons in the three-dimensional space \mathbb{R}^3 , described by the many-body Schrödinger Hamiltonian

$$H_N = \sum_{j=1}^N h_j + \sum_{1 \leq j < k \leq N} w_N(x_j - x_k) \tag{1-1}$$

acting on the space $\mathfrak{H}^N = \bigotimes_{\text{sym}}^N L^2(\mathbb{R}^3)$ of permutation-symmetric square integrable functions. The one-body operator is given by

$$h := (-i\nabla + A(x))^2 + V(x),$$

with a magnetic (or a rotation) field A satisfying

$$A \in L^3_{\text{loc}}(\mathbb{R}^3, \mathbb{R}^3), \quad \lim_{|x| \rightarrow \infty} |A(x)|e^{-b|x|} = 0, \tag{1-2}$$

for some constant $b > 0$ and an external potential V satisfying

$$0 \leq V \in L^1_{\text{loc}}(\mathbb{R}^3), \quad \lim_{|x| \rightarrow \infty} V(x) = +\infty. \tag{1-3}$$

We thus consider nonrelativistic particles in a trapping potential, possibly under the influence of an effective magnetic field, which might be due to rotation of the sample or the interaction with optical fields.

The particles interact pairwise via a repulsive potential w_N given by

$$w_N(x) = N^2 w(Nx), \tag{1-4}$$

where w is a fixed function which is nonnegative, radial and of finite range, i.e., $\mathbb{1}(|x| > R_0)w(x) \equiv 0$ for some constant $R_0 > 0$. Different scalings of the interaction potential of the form

$$w_{\beta,N}(x) = \frac{1}{N} N^{3\beta} w(N^\beta x), \tag{1-5}$$

with $0 \leq \beta \leq 1$, have been considered in the literature. The N^{-1} prefactor makes the interaction energy in (1-1) of the same order as the one-particle energy. Indeed, if $\beta > 0$, then

$$N^{3\beta} w(N^\beta x) \xrightarrow{N \rightarrow \infty} \left(\int w \right) \delta_0 \tag{1-6}$$

weakly and thus the interaction potential $w_{\beta,N}$ should be thought of as leading to a bounded interaction energy per pair of particles. Generally speaking, the larger the parameter β , the faster the potential converges to a point interaction, and thus the harder the analysis. Note that the cases $\beta < \frac{1}{3}$ and $\beta > \frac{1}{3}$ correspond to two physically rather different scenarios: in the former, the range of the potential is much larger than the typical interparticle distance $N^{-1/3}$, and we should expect many weak collisions; while in the latter, we rather have very few but very strong collisions. In this paper, we consider the most interesting case $\beta = 1$, where the naive approximation (1-6) does *not* capture the leading-order behavior of the physical system. In fact, the strong correlations at short distances $O(N^{-1})$ yield a nonlinear correction, which essentially amounts to replacing the coupling constant $\int w$ by $(8\pi) \times$ (the scattering length of w).

Let us quickly recall the definition of the scattering length; a more complete discussion can be found in [Lieb et al. 2005b, Appendix C]. Under our assumption on w , the zero-energy scattering equation

$$(-2\Delta + w(x))f(x), \quad \lim_{|x| \rightarrow \infty} f(x) = 1,$$

has a unique solution and it satisfies

$$f(x) = 1 - \frac{a}{|x|} \quad \forall |x| > R_0$$

for some constant $a \geq 0$ which is called the *scattering length* of w . In particular, if w is the potential for hard spheres, namely $w(x) \equiv +\infty$ when $|x| < R_0$ and $w(x) \equiv 0$ when $|x| \geq R_0$, then the scattering length of w is exactly R_0 . In a dilute gas, the scattering length can be interpreted as an effective range of the interaction: a quantum particle far from the others is felt by them as a hard sphere of radius a . A useful variational characterization of a is

$$8\pi a = \inf \left\{ \int_{\mathbb{R}^3} 2|\nabla f|^2 + w|f|^2, \quad \lim_{|x| \rightarrow \infty} f(x) = 1 \right\}. \quad (1-7)$$

Consequently, $8\pi a$ is smaller than $\int w$ (the strict inequality can be seen by taking the trial function $1 - \lambda g$ with $g \in C_c^2(\mathbb{R}^3, \mathbb{R})$ satisfying $g(x) \equiv 1$ when $|x| < R_0$, and $\lambda > 0$ sufficiently small). Moreover, a simple scaling shows that the scattering length of $w_N = N^2 w(N \cdot)$ is a/N .

We are going to prove that the ground-state energy and ground states of H_N converge to those of the Gross–Pitaevskii functional

$$\mathcal{E}_{\text{GP}}(u) := \langle u, hu \rangle + 4\pi a \int_{\mathbb{R}^3} |u(x)|^4 dx \quad (1-8)$$

in a suitable sense. Note that the occurrence of the scattering length in (1-8) is subtle: this functional is *not* obtained by testing H_N with factorized states of the form $u^{\otimes N}$ (which would lead to a functional with $4\pi a$ replaced by $\frac{1}{2} \int w$). Taking into account the short-range correlation structure which gives rise to (1-8) is the main difficulty in the proof of the following theorem, which is our main result.

Theorem 1.1 (Derivation of the Gross–Pitaevskii functional).

Under conditions (1-2), (1-3) and (1-4), we have

$$\lim_{N \rightarrow \infty} \inf_{\|\Psi\|_{\mathfrak{S}_N} = 1} \frac{\langle \Psi, H_N \Psi \rangle}{N} = \inf_{\|u\|_{L^2(\mathbb{R}^3)} = 1} \mathcal{E}_{\text{GP}}(u) =: e_{\text{GP}}. \quad (1-9)$$

Moreover, if Ψ_N is an approximate ground state for H_N , namely

$$\lim_{N \rightarrow \infty} \frac{\langle \Psi_N, H_N \Psi_N \rangle}{N} = e_{\text{GP}},$$

then there exists a subsequence Ψ_{N_ℓ} and a Borel probability measure μ supported on the set of minimizers of $\mathcal{E}_{\text{GP}}(u)$ such that

$$\lim_{\ell \rightarrow \infty} \text{Tr} \left| \gamma_{\Psi_{N_\ell}}^{(k)} - \int |u^{\otimes k} \rangle \langle u^{\otimes k} | d\mu(u) \right| = 0 \quad \forall k \in \mathbb{N}, \quad (1-10)$$

where $\gamma_{\Psi_N}^{(k)} = \text{Tr}_{k+1 \rightarrow N} |\Psi_N\rangle\langle\Psi_N|$ is the k -particle reduced density matrix of Ψ_N . In particular, if $\mathcal{E}_{\text{GP}}(u)$ subject to $\|u\|_{L^2} = 1$ has a unique minimizer u_0 (up to a complex phase), then there is complete Bose–Einstein condensation

$$\lim_{N \rightarrow \infty} \text{Tr} |\gamma_{\Psi_N}^{(k)} - |u_0^{\otimes k}\rangle\langle u_0^{\otimes k}|| = 0 \quad \forall k \in \mathbb{N}. \tag{1-11}$$

The energy upper bound in (1-9) was proved in [Lieb et al. 2000; Seiringer 2003] (see also [Benedikter et al. 2016, Appendix A] for an alternative approach). The energy lower bound in (1-9) and the convergence of one-particle density matrices were proved in [Lieb and Seiringer 2006]. The simpler case $A \equiv 0$ has been treated before in [Lieb et al. 2000] (ground-state energy) and [Lieb and Seiringer 2002] (condensation). In this case, the uniqueness of the Gross–Pitaevskii minimizer u_0 follows from a simple convexity argument. The result in Theorem 1.1 is thus not new, but the existing proofs are fairly difficult, in particular that of [Lieb and Seiringer 2006] which deals with the case $A \neq 0$.

In the present paper, we will provide alternative proofs of the energy lower bound and the convergence of states using the quantum de Finetti theorem in the same spirit as in [Lewin et al. 2014; 2015a]. Our proofs are conceptually and technically simpler than those provided in [Lieb and Seiringer 2006]. The overall strategy will be explained in the next section.

Our result covers the case of a rotating gas where the minimizers of the Gross–Pitaevskii functional can develop quantized vortices. This corresponds to taking $A(x) = \Omega \wedge x$, with Ω being the angular velocity vector. In this case, V should be interpreted as the trapping potential minus $\frac{1}{2}(\Omega \wedge x)^2$. The assumption $V(x) \rightarrow \infty$ as $|x| \rightarrow \infty$ is to ensure that all particles are confined to the system. Here our conditions on A and V are slightly more general than those of [Lieb and Seiringer 2006], where A is assumed to grow at most polynomially and V is assumed to grow at least logarithmically.

The finite range assumption on w is not a serious restriction because we can always restrict the support of w to a finite ball without changing the scattering length significantly. In fact, it is sufficient to assume that w is integrable at infinity, in which case the scattering length is well-defined. We can also work with a more general interaction $w_N \geq 0$ (with scattering length a_N) rather than the specific choice (1-4), as long as its range goes to zero and $\lim_{N \rightarrow \infty} Na_N$ exists; then the result in Theorem 1.1 still holds with a replaced by $\lim_{N \rightarrow \infty} Na_N$. In particular, if w_N is chosen as in (1-6) for some $0 < \beta < 1$, then $Na_N \rightarrow (8\pi)^{-1} \int w$. The critical case $\beta = 1$ considered in this paper is much more interesting because in the limit, the true scattering length appears instead of its first-order Born approximation $(8\pi)^{-1} \int w$.

2. Overall strategy

In this section we give an outline of the proofs of our main results, in order to better emphasize the key new points for the energy lower bound and the convergence of states.

We shall use the following notation: Let $\theta : \mathbb{R}^3 \rightarrow \mathbb{R}$ be a radial smooth Heaviside-like function; i.e.,

$$0 \leq \theta \leq 1, \quad \theta(x) \equiv 0 \quad \text{for } |x| \leq 1 \quad \text{and} \quad \theta(x) \equiv 1 \quad \text{for } |x| \geq 2.$$

Let $U : \mathbb{R}^3 \rightarrow \mathbb{R}$ be a radial smooth function supported on the annulus $\frac{1}{2} \leq |x| \leq 1$ such that

$$U(x) \geq 0 \quad \text{and} \quad \int_{\mathbb{R}^3} U = 4\pi a.$$

For every $R > 0$, define

$$\theta_R(x) = \theta\left(\frac{x}{R}\right), \quad U_R(x) = \frac{1}{R^3} U\left(\frac{x}{R}\right).$$

The smooth cut-off function θ_R will be used to perform cut-offs in both space and momentum variables, the latter being always denoted by

$$p = -i\nabla.$$

The potential U_R will be used to replace the original one. The important points will be that the integral of U_R yields the correct physical scattering length, and that we will have some freedom in choosing the range R of U_R .

Step 1 (Dyson’s lemma). The main difficulty in dealing with the Gross–Pitaevskii limit is that an ansatz $u^{\otimes N}$ does *not* give the correct energy asymptotics. In this regime, correlations between particles *do* matter, and one should rather think of an ansatz of the form

$$\prod_{i=1}^N u(x_i) \prod_{1 \leq i < j \leq N} f(x_i - x_j), \tag{2-1}$$

or a close variant, where f is linked to the two-body scattering process. We shall follow the approach of [Lieb and Seiringer 2006], relying on a generalization of an idea due to Dyson [1957]. The following lemma, proved in [Lieb et al. 2005a], allows us to bound our Hamiltonian from below by an effective one which is much less singular, but still encodes the scattering length of the original interaction potential.

Lemma 2.1 (Generalized Dyson lemma).

For all $s > 0$, $1 > \varepsilon > 0$ and $R > 2R_0/N$, we have

$$H_N \geq \sum_{j=1}^N (h_j - (1 - \varepsilon)p_j^2 \theta_s(p_j)) + \frac{(1 - \varepsilon)^2}{N} W_N - C \frac{N^2 R^2 s^5}{\varepsilon}, \tag{2-2}$$

where

$$W_N := \sum_{i \neq j}^N U_R(x_i - x_j) \prod_{k \neq i, j} \theta_{2R}(x_j - x_k). \tag{2-3}$$

Here and in the sequel, C stands for a generic positive constant.

Proof. Recall that the scattering length of w_N is a/N . Therefore, from equation (50) and the first estimate in (52) in [Lieb et al. 2005a], with (v, a, χ, s) replaced by $(w_N, a/N, \theta_s, s^{-1})$, one has

$$p^2 \theta_s(p) + \frac{1}{2} \sum_{j=1}^{N-1} w_N(x - y_j) \geq \frac{1 - \varepsilon}{N} \sum_{j=1}^{N-1} U_R(x - y_j) - \frac{CaR^2s^5}{\varepsilon}$$

on $L^2(\mathbb{R}^3)$ for all given points y_j satisfying $\min_{j \neq k} |y_j - y_k| \geq 2R$. Since the left side is nonnegative, we can relax the condition $\min_{j \neq k} |y_j - y_k| \geq 2R$ by multiplying the right side by $\prod_{k \neq j} \theta_{2R}(y_j - y_k)$.

Thus for every $i = 1, 2, \dots, N$,

$$p_i^2 \theta_s(p_i) + \frac{1}{2} \sum_{j \neq i}^N w_N(x_i - x_j) \geq \frac{1-\varepsilon}{N} \sum_{j \neq i} U_R(x_i - x_j) \prod_{k \neq i, j} \theta_{2R}(x_j - x_k) - \frac{CaR^2s^5}{\varepsilon}.$$

Multiplying both sides by $1 - \varepsilon$ and summing over i , we obtain (2-2). □

Clarification. The reader should keep in mind that we will choose $R = R(N) \rightarrow 0$ (actually $N^{-1/2} \gg R \gg N^{-2/3}$), then $s \rightarrow \infty$ and $\varepsilon \rightarrow 0$.

The main point of Dyson’s lemma is that we can replace the hard interaction potential w_N by a softer one U_R which encodes the scattering length conveniently as $\int U_R = 4\pi a$. The price we have to pay for this advantage is twofold, however: first, we have to use all the high-momentum part of the kinetic energy (note that $\theta_s(p) = 1$ when $p \geq 2s$); and second, the new potential $U_R(x_i - x_j)$ comes with the cut-off $\prod_{k \neq i, j} \theta(x_j - x_k)$. Together they really describe a “nearest neighbor” potential instead of an ordinary two-body potential. While the first problem is not too annoying, as the low part of the momentum is sufficient to recover the full energy in the limit, the second problem is much more serious.

Step 2 (Second moment estimate). The lower bound (2-2) leads us to consider the effective Hamiltonian

$$\tilde{H}_N := \sum_{j=1}^N \tilde{h}_j + \frac{(1-\varepsilon)^2}{N} W_N, \tag{2-4}$$

where

$$\tilde{h} := h - (1 - \varepsilon)p^2\theta_s(p) - \kappa_{\varepsilon,s}, \quad \kappa_{\varepsilon,s} := \inf \sigma(h - (1 - \varepsilon)p^2\theta_s(p) - 1). \tag{2-5}$$

Here we use the freedom to add and remove the constant $N\kappa_{\varepsilon,s}$ to the Hamiltonian to reduce to the case $\tilde{h} \geq 1$. In order to ensure that $\kappa_{\varepsilon,s}$ is finite, we need the extra condition

$$\lim_{|x| \rightarrow \infty} \frac{|A(x)|^2}{V(x)} = 0, \tag{2-6}$$

which can be removed at a later stage, as we shall explain below.

We will now seek a lower bound to the ground-state energy of (2-4). The philosophy, as in the previous work [Lieb and Seiringer 2006], is that if Ψ_N is the ground state of the original Hamiltonian, then roughly

$$\Psi_N \approx \tilde{\Psi}_N \prod_{1 \leq i < j \leq N} f(x_i - x_j),$$

where f encodes the two-body scattering process and $\tilde{\Psi}_N$ is a ground state for (2-4). Thus the Dyson lemma allows to extract the short-range correlation structure, and we now want to justify that $\tilde{\Psi}_N$ can be approximated by a tensor power $u^{\otimes N}$; that is, we want to justify the mean-field approximation at the level of the ground state of (2-4).

There are two key difficulties left:

- The effective Hamiltonian is genuinely many-body. It can be bounded below by a three-body Hamiltonian, but obviously one will ultimately have to show that the three-body contribution can be neglected.

- To recover the correct energy in the limit, we need to take $R \ll N^{-1/3}$ in order to be able to neglect the three-body contribution in the effective Hamiltonian. We thus still have to deal with the mean-field approximation in the “rare but strong collisions” limit. In other words, even though the effective Hamiltonian is much less singular than the original one, we do not have the freedom to reduce the singularity as much as we would like.

It is in treating these two difficulties that our new method significantly departs from the previous works [Lewin et al. 2015a; Lieb and Seiringer 2006]. We shall rely on a strong a priori estimate for ground states of (2-4). In Lemma 3.1, we assume (2-6) and show that (provided $R \gg N^{-2/3}$, which is sufficient for our purpose)

$$(\tilde{H}_N)^2 \geq \frac{1}{3} \left(\sum_{j=1}^N \tilde{h}_j \right)^2. \quad (2-7)$$

Note that a bound of this kind is not available for the original H_N due to the singularity of its interaction potential. In particular, (2-7) implies that every ground state $\tilde{\Psi}_N$ of \tilde{H}_N satisfies the strong a priori estimate

$$\langle \tilde{\Psi}_N, \tilde{h}_1 \tilde{h}_2 \tilde{\Psi}_N \rangle \leq C_{\varepsilon, s}. \quad (2-8)$$

This second moment estimate is the key point in our analysis in the next steps. It is reminiscent of similar estimates used in the literature for the time-dependent problem [Erdős et al. 2007; 2009; 2010; Erdős and Yau 2001].

Notation. We always denote by C_ε (or $C_{\varepsilon, s}$) a (generic) constant independent of s , N and R (or independent of N and R , respectively).

Step 3 (Three-body estimate). Next we have to remove the cut-off

$$\prod_{k \neq i, j} \theta(x_j - x_k)$$

in W_N to obtain a lower bound in terms of a two-body Hamiltonian. Using the elementary inequality (see [Lieb and Seiringer 2006, equation (22)])

$$\prod_{k: k \neq i, j} \theta_{2R}(x_j - x_k) \geq 1 - \sum_{k: k \neq i, j} (1 - \theta_{2R}(x_j - x_k)),$$

we have

$$W_N \geq \sum_{i \neq j}^N U_R(x_i - x_j) - \sum_{k \neq i \neq j \neq k} U_R(x_i - x_j)(1 - \theta_{2R}(x_j - x_k)), \quad (2-9)$$

and we thus have only a three-body term to estimate. Since the summand in this term is zero except when $|x_i - x_j| \leq R$ and $|x_j - x_k| \leq 4R$, the last sum of (2-9) can be removed if the probability of having three or more particles in a region of diameter $O(R)$ is small enough. This should be the case if R is much smaller than $N^{-1/3}$, the average distance between particles, but it is rather difficult to confirm this intuition rigorously.

In [Lieb and Seiringer 2006], a three-body estimate was established using a subtle argument based on path integrals (the Trotter product formula). In this paper, we will follow a different, simpler approach. Instead of working directly with a ground state of H_N as in [Lieb and Seiringer 2006], we will consider a ground state $\tilde{\Psi}_N$ of the effective Hamiltonian \tilde{H}_N . Thanks to the second moment estimate (2-7), we can show that (see Lemma 3.4)

$$\sum_{k=3}^N \langle \tilde{\Psi}_N, U_R(x_1 - x_2)\theta_{2R}(x_2 - x_k)\tilde{\Psi}_N \rangle \leq C_{\varepsilon,s}NR^2. \tag{2-10}$$

The right side of (2-10) is small with our choice $N^{-1/2} \gg R$.

Step 4 (Mean-field approximation). With the cut-off in W_N removed, \tilde{H}_N turns into the two-body Hamiltonian

$$K_N := \sum_{j=1}^N \tilde{h}_j + \frac{(1-\varepsilon)^2}{N} \sum_{i \neq j} U_R(x_i - x_j)$$

for which we can validate the *mean-field approximation*. This is the simplest approximation for Bose gases where one restricts the many-body wave functions to the pure tensor products $u^{\otimes N}$. Since U_R converges to the delta-interaction with mass $\int U_R = 4\pi a$, we formally obtain the following approximation for the ground-state energy

$$e_{\text{NL}}(\varepsilon, s) := \inf_{\|u\|_{L^2}=1} \left(\langle u, \tilde{h}u \rangle + (1-\varepsilon)^2 4\pi a \int |u|^4 \right).$$

In Section 4A, we will show that

$$\lim_{N \rightarrow \infty} \frac{\inf \sigma(K_N)}{N} = e_{\text{NL}}(\varepsilon, s). \tag{2-11}$$

A similar result was proved in [Lieb and Seiringer 2006] using a coherent state method, which is a generalization of the c -number substitution in [Lieb et al. 2005c]. In the present paper, we will provide an alternative proof of (2-11) using the quantum de Finetti theorem of Størmer [1969] and Hudson and Moody [1976]. We note that this theorem has proved useful also in the derivation of the Gross–Pitaevskii equation in the dynamical case; see [Chen et al. 2015]. The following formulation is taken from [Lewin et al. 2014, Corollary 2.4] (see [Rougerie 2015] for a general discussion and more references):

Theorem 2.2 (Quantum de Finetti).

Let \mathfrak{K} be an arbitrary separable Hilbert space and let $\Psi_N \in \bigotimes_{\text{sym}}^N \mathfrak{K}$ with $\|\Psi_N\| = 1$. Assume that the sequence of one-particle density matrices $\gamma_{\Psi_N}^{(1)}$ converges strongly in trace class when $N \rightarrow \infty$. Then, up to a subsequence, there exists a (unique) Borel probability measure μ on the unit sphere $S\mathfrak{K}$, invariant under the group action of S^1 , such that

$$\lim_{N \rightarrow \infty} \text{Tr} \left| \gamma_{\Psi_N}^{(k)} - \int |u^{\otimes k}\rangle \langle u^{\otimes k}| d\mu(u) \right| = 0 \quad \forall k \in \mathbb{N}. \tag{2-12}$$

This theorem validates the mean-field approximation for a large class of trapped Bose gases, in particular (see [Lewin et al. 2014] and references therein) when the strength of the interaction is proportional to the

inverse of the particle number, case $\beta = 0$ in (1-6). However, when the interaction becomes stronger, the mean-field approximation is harder to justify. The convergence (2-11) with $R \gg N^{-2/15}$ was proved in [Lewin et al. 2015a] by using a quantitative version of Theorem 2.2 valid for finite-dimensional spaces [Christandl et al. 2007; Chiribella 2011; Lewin et al. 2015b]. However, this range of R is too small for our purpose because we are forced to choose $R \ll N^{-1/2}$ in the previous steps.

In this paper, thanks to the strong a priori estimate (2-8), we are able to prove (2-11) for the larger range $R \gg N^{-2/3}$. As in [Lewin et al. 2015a; Lieb and Seiringer 2006], we localize the problem onto energy levels of the one-body Hamiltonian \tilde{h} lying below a chosen cut-off Λ . At fixed Λ , it turns out that the projected Hamiltonian is bounded proportionally to N . We are thus in a usual mean-field scaling if we are allowed to take $N \rightarrow \infty$ first, and then $\Lambda \rightarrow \infty$ later. Taking limits in this order demands a very strong control on the localization error made by projecting the Hamiltonian, however. This control is provided again by the moment estimate (2-8).

Combining the arguments in Steps 1–4, we can pass to the limit $N \rightarrow \infty$; then $s \rightarrow \infty$ and $\varepsilon \rightarrow 0$ to obtain the energy convergence (1-9) under the extra condition (2-6). In Section 4B, we remove this technical assumption using a concavity argument from [Lieb and Seiringer 2006] and a binding inequality which goes back to an idea in [Lieb 1984].

Step 5 (Convergence of ground states). In Section 4C, we prove the convergence of (approximate) ground states using the convergence of the ground state energy of a perturbed Hamiltonian and the Feynman–Hellmann principle. A similar approach was used in [Lieb and Seiringer 2006] to prove the convergence of the 1-particle density matrix. However, the quantum de Finetti theorem helps us to avoid the complicated convex analysis in [Lieb and Seiringer 2006], simplifying the proof significantly and giving access to higher-order density matrices.

3. Second moment estimate

In this section, we consider the effective Hamiltonian obtained after applying the generalized Dyson lemma to the original problem, namely

$$\tilde{H}_N = \sum_{j=1}^N \tilde{h}_j + \frac{(1-\varepsilon)^2}{N} W_N,$$

where \tilde{h} and W_N are defined in (2-5) and (2-3), respectively. We will work under the extra assumption (2-6). Since $A \in L^3_{\text{loc}}(\mathbb{R}^3, \mathbb{R}^3)$ and V grows faster than $|A|^2$ at infinity, for every $\varepsilon > 0$, we have

$$\left(\frac{V}{2} - 2\varepsilon^{-1}|A|^2 \right)_- \in L^{3/2}(\mathbb{R}^3),$$

and hence

$$\left(\frac{\varepsilon}{4} \right) p^2 + \frac{V}{2} - 2\varepsilon^{-1}|A|^2 \geq -C_\varepsilon.$$

In combination with the Cauchy–Schwarz inequality, we get

$$h - (1-\varepsilon)p^2 \theta_s(p) \geq \frac{\varepsilon}{2} p^2 - 2\varepsilon^{-1}|A|^2 + V \geq \frac{\varepsilon}{4} p^2 + \frac{V}{2} - C_\varepsilon.$$

Therefore, $\inf \sigma(h) - 1 \geq \kappa_{\varepsilon,s} \geq -C_\varepsilon$ and

$$\tilde{h} \geq C_\varepsilon(-\Delta + V + 1). \tag{3-1}$$

The key estimate in this section is the following:

Lemma 3.1 (Second moment estimate).

Assume that (2-6) holds. For every $1 > \varepsilon > 0$ and $s > 0$, if

$$R = R(N) \gg N^{-2/3}$$

when $N \rightarrow \infty$, then for N large enough, we have the operator bound

$$(\tilde{H}_N)^2 \geq \frac{1}{3} \left(\sum_{j=1}^N \tilde{h}_j \right)^2. \tag{3-2}$$

We will show in Section 3C that a convenient lower bound to Dyson’s potential W_N in terms of truly two-body operators follows from Lemma 3.1.

Before proving Lemma 3.1 in Section 3B, we first collect some useful inequalities on a generic translation-invariant interaction operator $W(x - y)$ that will be used throughout the paper.

3A. Operator inequalities for interaction potentials. We state several useful inequalities in the following lemma. In fact, (3-3) is well-known, and (3-4) with $\delta = 0$ was proved earlier in [Erdős and Yau 2001, Lemma 5.3]. In the sequel, we will crucially rely on the improvement to $\delta > 0$, and on (3-5), which seem to be new.

Lemma 3.2 (Inequalities for a repulsive interaction potential).

For every $0 \leq W \in L^1 \cap L^2(\mathbb{R}^3)$, the multiplication operator $W(x - y)$ on $L^2((\mathbb{R}^3)^2)$ satisfies

$$0 \leq W(x - y) \leq C \|W\|_{L^{3/2}(\mathbb{R}^3)}(-\Delta_x), \tag{3-3}$$

and, for any $0 \leq \delta < \frac{1}{4}$,

$$0 \leq W(x - y) \leq C_\delta \|W\|_{L^1(\mathbb{R}^3)}(1 - \Delta_x)^{1-\delta}(1 - \Delta_y)^{1-\delta}. \tag{3-4}$$

Moreover, for all $1 > \varepsilon > 0$, $s > 0$, $A \in L^3_{\text{loc}}(\mathbb{R}^3, \mathbb{R}^3)$ and $0 \leq V \in L^1_{\text{loc}}(\mathbb{R}^3)$,

$$\tilde{h}_x W(x - y) + W(x - y) \tilde{h}_x \geq -C (\|W\|_{L^2} + (1 + s^2) \|W\|_{L^{3/2}})(1 - \Delta_x)(1 - \Delta_y). \tag{3-5}$$

Proof of Lemma 3.2. We will prove this in several steps.

Proof of (3-3). From Hölder’s and Sobolev’s inequalities, we have

$$\begin{aligned} \langle f, W(x - y)f \rangle &= \iint W(x - y) |f(x, y)|^2 dx dy \\ &\leq \int \left(\int W(x - y)^{3/2} dx \right)^{2/3} \left(\int |f(x, y)|^6 dx \right)^{1/3} dy \\ &\leq C \|W\|_{L^{3/2}(\mathbb{R}^3)} \int \left(\int |\nabla_x f(x, y)|^2 dx \right) dy \end{aligned}$$

for every function $f \in H^1((\mathbb{R}^3)^2)$. Therefore, (3-3) follows immediately.

Proof of (3-4). The estimate (3-4) with $\delta = 0$ was first derived in [Erdős and Yau 2001]. The following proof is adapted from the proof (again for $\delta = 0$) in [Lieb and Seiringer 2006]. Note that for every operator K , we have $K^*K \leq 1$ if and only if $KK^* \leq 1$. Therefore, (3-4) is equivalent to

$$\sqrt{W(x-y)}(1-\Delta_x)^{\delta-1}(1-\Delta_y)^{\delta-1}\sqrt{W(x-y)} \leq C_\delta \|W\|_{L^1}. \tag{3-6}$$

Let G be the Green function of $(1-\Delta)^{\delta-1}$ whose Fourier transform is given by

$$\widehat{G}(k) := \int_{\mathbb{R}^3} e^{-2\pi i x \cdot k} G(x) dx = \frac{1}{(1+4\pi^2|k|^2)^{1-\delta}}.$$

For every function $f \in L^2((\mathbb{R}^3)^2)$, one has

$$\begin{aligned} & \langle f, \sqrt{W(x-y)}(1-\Delta_x)^{\delta-1}(1-\Delta_y)^{\delta-1}\sqrt{W(x-y)}f \rangle \\ &= \int \overline{f(x,y)}\sqrt{W(x-y)}G(x-x')G(y-y')\sqrt{W(x'-y')}f(x',y') dx dy dx' dy' \\ &\leq \int \frac{W(x-y)|G(x-x')|^2|f(x',y')|^2 + W(x'-y')|G(y-y')|^2|f(x,y)|^2}{2} \\ &= C_\delta \|W\|_{L^1} \langle f, f \rangle, \end{aligned}$$

where

$$C_\delta := \int |G|^2 = \int |\widehat{G}|^2 = \int_{\mathbb{R}^3} \frac{dk}{(1+4\pi^2|k|^2)^{2(1-\delta)}},$$

which is finite for all $0 \leq \delta < \frac{1}{4}$. Thus (3-6), and hence (3-4), holds true.

Simpler version of (3-5). We are going to deduce (3-5) from the inequality

$$(-\Delta_x)W(x-y) + W(x-y)(-\Delta_x) \geq -C(\|W\|_{L^{3/2}} + \|W\|_{L^2})(1-\Delta_x)(1-\Delta_y). \tag{3-7}$$

By an approximation argument, one can assume that W is smooth. For every $f \in H^2(\mathbb{R}^3 \times \mathbb{R}^3)$, a straightforward calculation using integration by parts, and the identity $\nabla_x(W(x-y)) = -\nabla_y(W(x-y))$ gives us

$$\begin{aligned} & \langle f, ((-\Delta_x)W(x-y) + W(x-y)(-\Delta_x))f \rangle \\ &= 2\Re \iint \nabla_x \overline{f(x,y)} \nabla_x (W(x-y)) f(x,y) dx dy \\ &= 2 \iint |\nabla_x f(x,y)|^2 W(x-y) + 2\Re \iint \nabla_x \overline{f(x,y)} \nabla_x (W(x-y)) f(x,y) dx dy \\ &\geq -2\Re \iint \nabla_x \overline{f(x,y)} \nabla_y (W(x-y)) f(x,y) dx dy \\ &= 2\Re \iint \nabla_y ((\nabla_x \overline{f(x,y)}) f(x,y)) W(x-y) dx dy \\ &= 2\Re \iint (\nabla_x \overline{f(x,y)} \nabla_y f(x,y) + \nabla_y (\nabla_x \overline{f(x,y)}) f(x,y)) W(x-y) dx dy. \end{aligned}$$

Using Cauchy–Schwarz and Sobolev’s inequality (3-3), we get

$$\begin{aligned} \left| \iint \nabla_x \overline{f(x, y)} \nabla_y f(x, y) W(x - y) dx dy \right| &\leq \iint \frac{|\nabla_x f(x, y)|^2 + |\nabla_y f(x, y)|^2}{2} |W(x - y)| dx dy \\ &\leq C \|W\|_{L^{3/2}} \langle f, (-\Delta_x)(-\Delta_y)f \rangle. \end{aligned}$$

Moreover, by the Cauchy–Schwarz inequality again and (3-4) (with $\delta = 0$ and W replaced by W^2),

$$\begin{aligned} \left| \iint (\nabla_y \nabla_x \overline{f(x, y)}) f(x, y) W(x - y) dx dy \right| \\ \leq \left(\iint |\nabla_y \nabla_x f(x, y)|^2 dx dy \right)^{1/2} \left(\iint |f(x, y)|^2 |W(x - y)|^2 dx dy \right)^{1/2} \\ \leq C \|W\|_{L^2} \langle f, (1 - \Delta_x)(1 - \Delta_y)f \rangle. \end{aligned}$$

Thus we obtain

$$\langle f, ((-\Delta_x)W(x - y) + W(x - y)(-\Delta_x))f \rangle \geq -C(\|W\|_{L^{3/2}} + \|W\|_{L^2}) \langle f, (1 - \Delta_x)(1 - \Delta_y)f \rangle$$

for all $f \in H^2(\mathbb{R}^3 \times \mathbb{R}^3)$. This proves (3-7).

Proof of (3-5). From the commutator relation

$$p_x W(x - y) = W(x - y)p_x + (-i \nabla_x W)(x - y),$$

we find that

$$\begin{aligned} (p_x A(x) + A(x)p_x + |A(x)|^2)W(x - y) + W(x - y)(p_x A(x) + A(x)p_x + |A(x)|^2) \\ = 2(p_x W(x - y)A(x) + A(x)W(x - y)p_x + |A(x)|^2 W(x - y)) \\ = 2(p_x + A(x))W(x - y)(p_x + A(x)) - 2p_x W(x - y)p_x. \end{aligned}$$

Using

$$(p_x + A(x))W(x - y)(p_x + A(x)) \geq 0$$

and estimating $p_x W(x - y)p_x$ by Sobolev’s inequality (3-3), we get

$$\begin{aligned} (p_x A(x) + A(x)p_x + |A(x)|^2)W(x - y) + W(x - y)(p_x A(x) + A(x)p_x + |A(x)|^2) \\ \geq -C \|W\|_{L^{3/2}} (-\Delta_x)(-\Delta_y). \end{aligned} \tag{3-8}$$

Finally, by (3-3) again and the Cauchy–Schwarz inequality for operators

$$\pm(XY + Y^* X^*) \leq \delta X X^* + \delta^{-1} Y^* Y \quad \forall \delta > 0, \tag{3-9}$$

we obtain

$$\begin{aligned} p_x^2(1 - \theta_s(p_x))W(x - y) + W(x - y)p_x^2(1 - \theta_s(p_x)) \\ \geq -\delta p_x^2(1 - \theta_s(p_x))W(x - y)p_x^2(1 - \theta_s(p_x)) + \delta^{-1} W(x - y) \\ \geq -C \|W\|_{L^{3/2}} (\delta p_x^4(1 - \theta_s(p_x))^2 + \delta^{-1})(-\Delta_x) \end{aligned}$$

for all $\delta > 0$. Using $1 - \theta_s(p) \leq \mathbb{1}(|p| \leq 2s)$ and choosing $\delta \sim s^{-2}$ gives

$$p_x^2(1 - \theta_s(p_x))W(x - y) + W(x - y)p_x^2(1 - \theta_s(p_x)) \geq -Cs^2 \|W\|_{L^{3/2}} (-\Delta_x). \tag{3-10}$$

From (3-7), (3-8) and (3-10), the bound (3-5) follows. □

3B. Proof of Lemma 3.1. Before completing the proof of Lemma 3.1, we make a remark on the simpler case with the Dyson potential W_N replaced by a truly two-body interaction.

Remark 3.3 (Second moment estimate with two-body interactions).

Consider the model case

$$K_N := \sum_{j=1}^N \tilde{h}_j + \frac{(1-\varepsilon)^2}{N} \sum_{i \neq j} U_R(x_i - x_j).$$

By expanding K_N^2 and using the fact that $\tilde{h}_i \geq 0$ commutes with $U_R(x_j - x_k) \geq 0$ when $i \neq j$ and $i \neq k$, and then using (3-5) to estimate terms of the form

$$\tilde{h}_i U_R(x_i - x_j) + U(x_i - x_j) \tilde{h}_i,$$

we obtain

$$K_N^2 \geq \frac{1}{3} \sum_{1 \leq i \neq j \leq N} \tilde{h}_i \tilde{h}_j \quad (3-11)$$

provided that $R = R(N) \gg N^{-2/3}$. A similar estimate also holds when \tilde{h} is replaced by the original kinetic operator h .

We stress once again that we do *not* expect (3-11) to hold for our original Hamiltonian H_N , which is in the more singular regime $R \sim N^{-1}$. We thus need to work with the Dyson Hamiltonian, and its rather intricate nature makes the actual proof of Lemma 3.1 more difficult than the one we have sketched for (3-11). We now proceed with this proof.

Proof of Lemma 3.1. We have

$$(\tilde{H}_N)^2 - \left(\sum_{j=1}^N \tilde{h}_j \right)^2 = \frac{(1-\varepsilon)^2}{N} \sum_{\ell=1}^N (\tilde{h}_\ell W_N + W_N \tilde{h}_\ell) + \frac{(1-\varepsilon)^4}{N^2} W_N^2. \quad (3-12)$$

As in Remark 3.3, the goal is to bound $\tilde{h}_1 W_N + W_N \tilde{h}_1$ from below. We first decompose the interaction operator as

$$W_N = W_a + W_b,$$

where

$$W_a = \sum_{1 \in \{i, j\}} U_R(x_i - x_j) \prod_{k \neq i, j} \theta_{2R}(x_j - x_k),$$

$$W_b = \sum_{i, j \geq 2} U_R(x_i - x_j) \prod_{k \neq i, j} \theta_{2R}(x_j - x_k).$$

Estimate of W_a . By the Cauchy–Schwarz inequality (3-9), we get

$$\pm (\tilde{h}_1 W_a + W_a \tilde{h}_1) \leq N^{-1} \tilde{h}_1 W_a \tilde{h}_1 + N W_a. \quad (3-13)$$

Let us show that

$$W_a \leq \frac{C}{R^3}. \quad (3-14)$$

Indeed, for every given $(x_1, x_2, \dots, x_N) \in (\mathbb{R}^3)^N$, the product

$$U_R(x_1 - x_j) \prod_{k \neq 1, j} \theta_{2R}(x_j - x_k)$$

is bounded by $\|U_R\|_{L^\infty} \leq CR^{-3}$ and it is zero except in the case

$$|x_1 - x_j| < R < 2R < \min_{k \neq 1, j} |x_j - x_k|.$$

By the triangle inequality, the latter condition implies that

$$|x_1 - x_j| < R < \min_{k \neq 1, j} |x_1 - x_k|,$$

and it is satisfied by at most one index $j \neq 1$. Therefore,

$$\sum_{j \geq 2} U_R(x_1 - x_j) \prod_{k \neq 1, j} \theta_{2R}(x_j - x_k) \leq \frac{C}{R^3}.$$

Similarly, we have

$$\sum_{i \geq 2} U_R(x_i - x_1) \prod_{k \neq 1, i} \theta_{2R}(x_1 - x_k) \leq \frac{C}{R^3},$$

and hence (3-14) holds true. From (3-13) and (3-14), we obtain

$$\pm(\tilde{h}_1 W_a + W_a \tilde{h}_1) \leq \frac{C}{NR^3} (\tilde{h}_1)^2 + 2N \sum_{1 \in \{i, j\}} U_R(x_i - x_j) \prod_{k \neq i, j} \theta_{2R}(x_j - x_k). \tag{3-15}$$

Here we do not need to estimate the second term on the right side of (3-15) because this term is part of W_N , which will be controlled by W_N^2 in \tilde{H}_N^2 .

Estimate of W_b . We need a further decomposition

$$W_b = \sum_{i, j \geq 2} U_R(x_i - x_j) \prod_{k \neq i, j} \theta_{2R}(x_j - x_k) = W_c - W_d,$$

where

$$W_c := \sum_{i, j \geq 2} U_R(x_i - x_j) \prod_{k \neq 1, i, j} \theta_{2R}(x_j - x_k),$$

$$W_d := \sum_{i, j \geq 2} U_R(x_i - x_j) (1 - \theta_{2R}(x_j - x_1)) \prod_{k \neq 1, i, j} \theta_{2R}(x_j - x_k).$$

Note that

$$W_c \geq 0, \quad W_d \geq 0 \quad \text{and} \quad \tilde{h}_1 W_c = W_c \tilde{h}_1 \geq 0.$$

On the other hand, by the Cauchy–Schwarz inequality (3-9) again,

$$\pm(\tilde{h}_1 W_d + W_d \tilde{h}_1) \leq \delta \tilde{h}_1 W_d \tilde{h}_1 + \delta^{-1} W_d. \tag{3-16}$$

We have two different ways to bound W_d . First, by (3-3) and (3-1),

$$(1 - \theta_{2R}(x_j - x_1)) \leq C \|1 - \theta_{2R}\|_{L^{3/2}} (1 - \Delta_1) \leq C_\varepsilon R^2 \tilde{h}_1.$$

Since here $i, j \geq 2$, both sides of the latter estimate commute with

$$U_R(x_i - x_j) \prod_{k \neq 1, i, j} \theta_{2R}(x_j - x_k),$$

and we deduce that

$$(1 - \theta_{2R}(x_j - x_1))U_R(x_i - x_j) \prod_{k \neq 1, i, j} \theta_{2R}(x_j - x_k) \leq C_\varepsilon R^2 \tilde{h}_1 U_R(x_i - x_j) \prod_{k \neq 1, i, j} \theta_{2R}(x_j - x_k).$$

Taking the sum over $i, j \geq 2$, we obtain

$$W_d \leq C_\varepsilon R^2 \tilde{h}_1 W_c. \quad (3-17)$$

Second, let us show that

$$W_d \leq \frac{C}{R^3}. \quad (3-18)$$

Indeed, for every given $(x_1, x_2, \dots, x_N) \in (\mathbb{R}^3)^N$, the product

$$U_R(x_i - x_j)(1 - \theta_{2R}(x_j - x_1)) \prod_{k \neq 1, i, j} \theta_{2R}(x_j - x_k)$$

is zero except in the case

$$|x_i - x_j| < R, \quad |x_j - x_1| < 4R, \quad \min_{k \neq 1, i, j} |x_j - x_k| > 2R. \quad (3-19)$$

By the triangle inequality, (3-19) implies that the ball $B(x_1, 5R)$ contains $B(x_i, \frac{1}{2}R)$, $B(x_j, \frac{1}{2}R)$, and the balls $B(x_i, \frac{1}{2}R)$, $B(x_j, \frac{1}{2}R)$ do not intersect with $B(x_k, \frac{1}{2}R)$ for all $k \neq 1, i, j$. Since $B(x_1, 5R)$ can contain only a finite number of disjoint balls of radius $\frac{1}{2}R$, we see that there are only a finite number of pairs (i, j) satisfying (3-19). Thus we can conclude that

$$W_d \leq C \|U_R\|_{L^\infty} \leq CR^{-3}.$$

From (3-16), (3-17) and (3-18), we obtain

$$\tilde{h}_1 W_b + W_b \tilde{h}_1 = \tilde{h}_1 W_d + W_d \tilde{h}_1 + 2\tilde{h}_1 W_c \geq -\frac{C\delta}{R^3}(\tilde{h}_1)^2 + \left(2 - \frac{C_\varepsilon R^2}{\delta}\right)\tilde{h}_1 W_c.$$

Choosing $\delta \sim R^2$, we get

$$\tilde{h}_1 W_b + W_b \tilde{h}_1 \geq -\frac{C_\varepsilon}{R}(\tilde{h}_1)^2. \quad (3-20)$$

Conclusion. From (3-15) and (3-20), we get

$$\tilde{h}_1 W_N + W_N \tilde{h}_1 \geq -\left(\frac{C}{NR^3} + \frac{C_\varepsilon}{R}\right)(\tilde{h}_1)^2 - 2N \sum_{1 \in \{i, j\}} U_R(x_i - x_j) \prod_{k \neq i, j} \theta_{2R}(x_j - x_k).$$

Summing the similar estimates with 1 replaced by ℓ and using

$$\sum_{\ell=1}^N \sum_{\ell \in \{i, j\}} U_R(x_i - x_j) \prod_{k \neq i, j} \theta_{2R}(x_j - x_k) = 2W_N,$$

we find that

$$\sum_{\ell=1}^N (\tilde{h}_\ell W_N + W_N \tilde{h}_\ell) \geq -\left(\frac{C}{NR^3} + \frac{C_\varepsilon}{R}\right) \sum_{\ell=1}^N (\tilde{h}_\ell)^2 - 2NW_N.$$

Therefore, coming back to (3-12), we conclude that (completing a square in the last inequality)

$$\begin{aligned} (\tilde{H}_N)^2 - \left(\sum_{j=1}^N \tilde{h}_j\right)^2 &= \frac{(1-\varepsilon)^2}{N} \sum_{\ell=1}^N (\tilde{h}_\ell W_N + W_N \tilde{h}_\ell) + \frac{(1-\varepsilon)^4}{N^2} W_N^2 \\ &\geq -\left(\frac{C}{N^2 R^3} + \frac{C_\varepsilon}{NR}\right) \sum_{\ell=1}^N (\tilde{h}_\ell)^2 - 2(1-\varepsilon)^2 W_N + \frac{(1-\varepsilon)^4}{N^2} W_N^2 \\ &\geq -\left(\frac{C}{N^2 R^3} + \frac{C_\varepsilon}{NR}\right) \sum_{\ell=1}^N (\tilde{h}_\ell)^2 - N^2. \end{aligned}$$

When $R \gg N^{-2/3}$, we have

$$\frac{C}{N^2 R^3} + \frac{C_\varepsilon}{NR} \ll 1,$$

and hence

$$(\tilde{H}_N)^2 \geq 2 \sum_{1 \leq i < j \leq N} \tilde{h}_i \tilde{h}_j + (1 - o(1)) \sum_{\ell=1}^N (\tilde{h}_\ell)^2 - N^2,$$

which yields the result, recalling that in our convention, $\tilde{h} \geq 1$. □

3C. Three-body estimate. A first consequence of the second moment estimate in Lemma 3.1 is that we can conveniently bound Dyson’s Hamiltonian from below by a two-body Hamiltonian. This is done by first using a simple bound in terms of a three-body Hamiltonian, and then bounding the unwanted three-body part.

Lemma 3.4 (Three-body estimate).

Assume the extra condition (2-6) holds. For every $1 > \varepsilon > 0$ and $s > 0$, if $R = R(N) \gg N^{-2/3}$, then

$$\sum_{i \neq j} U_R(x_i - x_j) \sum_{k \neq i, j} (1 - \theta_{2R}(x_j - x_k)) \leq C_{\varepsilon, s} \frac{R^2}{N} (\tilde{H}_N)^4. \tag{3-21}$$

Consequently,

$$\tilde{H}_N \geq \sum_{j=1}^N \tilde{h}_j + \frac{(1-\varepsilon)^2}{N} \sum_{i \neq j} U_R(x_i - x_j) - C_{\varepsilon, s} \frac{R^2}{N^2} (\tilde{H}_N)^4. \tag{3-22}$$

Note the error term involving $(\tilde{H}_N)^4$, which is well under control since we are interested in its expectation value in a ground state.

Proof. By (3-3) and (3-1), we have

$$(1 - \theta_{2R}(x_2 - x_k)) \leq C_{\varepsilon, s} R^2 \tilde{h}_k \quad \text{for } k \geq 3.$$

Since $U_R(x_1 - x_2)$ commutes with both sides, we get

$$\begin{aligned}
U_R(x_1 - x_2) \sum_{k \geq 3} (1 - \theta_{2R}(x_2 - x_k)) &\leq C_{\varepsilon, s} R^2 U_R(x_1 - x_2) \sum_{k \geq 3} \tilde{h}_3 \\
&= \frac{1}{2} C_{\varepsilon, s} R^2 (\tilde{H}_N - \tilde{h}_1 - \tilde{h}_2 - (1 - \varepsilon)^2 N^{-1} W_N) U_R(x_1 - x_2) \\
&\quad + \frac{1}{2} C_{\varepsilon, s} R^2 U_R(x_1 - x_2) (\tilde{H}_N - \tilde{h}_1 - \tilde{h}_2 - (1 - \varepsilon)^2 N^{-1} W_N) \\
&\leq \frac{1}{2} C_{\varepsilon, s} R^2 (\tilde{H}_N U_R(x_1 - x_2) + U_R(x_1 - x_2) \tilde{H}_N) \\
&\quad + \frac{1}{2} C_{\varepsilon, s} R^2 \sum_{j=1}^2 (\tilde{h}_j U_R(x_1 - x_2) + U_R(x_1 - x_2) \tilde{h}_j). \tag{3-23}
\end{aligned}$$

In the last estimate, we have used $W_N \geq 0$. Thanks to (3-5) and (3-1), we get for all $j = 1, 2$,

$$\tilde{h}_j U_R(x_1 - x_2) + U_R(x_1 - x_2) \tilde{h}_j \geq -C_{\varepsilon, s} R^{-3/2} (1 - \Delta_1)(1 - \Delta_2) \geq -C_{\varepsilon, s} R^{-3/2} \tilde{h}_1 \tilde{h}_2. \tag{3-24}$$

On the other hand, by the Cauchy–Schwarz inequality (3-9) and (3-4) (with $\delta = 0$ and $W = U_R$),

$$\begin{aligned}
\tilde{H}_N U_R(x_1 - x_2) + U_R(x_1 - x_2) \tilde{H}_N &\leq \delta \tilde{H}_N U_R(x_1 - x_2) \tilde{H}_N + \delta^{-1} U_R(x_1 - x_2) \\
&\leq C_{\varepsilon, s} \delta \tilde{H}_N \tilde{h}_1 \tilde{h}_2 \tilde{H}_N + C_{\varepsilon, s} \delta^{-1} \tilde{h}_1 \tilde{h}_2 \tag{3-25}
\end{aligned}$$

for all $\delta > 0$. Choosing $\delta = N^{-1}$ and using $R^{-3/2} \leq N$, we deduce from (3-23), (3-24) and (3-25) that

$$U_R(x_1 - x_2) \sum_{k \geq 3} (1 - \theta_{2R}(x_2 - x_k)) \leq C_{\varepsilon, s} R^2 (N^{-1} \tilde{H}_N \tilde{h}_1 \tilde{h}_2 \tilde{H}_N + N \tilde{h}_1 \tilde{h}_2).$$

By symmetrization with respect to the indices, we find that

$$\sum_{i \neq j} U_R(x_1 - x_2) \sum_{k \neq i, j} (1 - \theta_{2R}(x_j - x_k)) \leq C_{\varepsilon, s} R^2 \left(N^{-1} \tilde{H}_N \sum_{i \neq j} \tilde{h}_i \tilde{h}_j \tilde{H}_N + N \sum_{i \neq j} \tilde{h}_i \tilde{h}_j \right).$$

Combining with the second moment estimate (3-2), we obtain (3-21). From the three-body estimate (3-21) and the elementary inequality (2-9), the operator bound (3-22) follows. \square

4. Energy lower bound and convergence of states

4A. Mean-field approximation. We are now reduced to justifying the mean-field approximation for a new Hamiltonian with the two-body interaction $U_R(x - y)$, which converges to a Dirac delta much slower than the original one. The analysis in this section provides an alternative to the coherent states method of [Lieb and Seiringer 2006].

Proposition 4.1 (Mean-field approximation).

Assume that (2-6) holds. For every $1 > \varepsilon > 0$ and $s > 0$, if

$$N^{-1/2} \gg R = R(N) \gg N^{-2/3}$$

then

$$\lim_{N \rightarrow \infty} \frac{\inf \sigma(\tilde{H}_N)}{N} = \inf_{\|u\|_{L^2}=1} \left(\langle u, \tilde{h}u \rangle + (1 - \varepsilon)^2 4\pi a \int |u|^4 \right) =: e_{\text{NL}}(\varepsilon, s). \tag{4-1}$$

Proof. The upper bound in (4-1) can be obtained easily using trial states of the form $u^{\otimes N}$. For the lower bound, let us consider a ground state $\tilde{\Psi}_N$ of \tilde{H}_N (which exists because \tilde{h} has compact resolvent). Using the ground-state equation, we find that

$$\langle \tilde{\Psi}_N, (\tilde{H}_N)^k \tilde{\Psi}_N \rangle = (\inf \sigma(\tilde{H}_N))^k \leq (C_{\varepsilon, s} N)^k \tag{4-2}$$

for all $k \in \mathbb{N}$. In particular, the second moment estimate (3-2) implies that

$$\langle \tilde{\Psi}_N, \tilde{h}_1 \tilde{h}_2 \tilde{\Psi}_N \rangle \leq C_{\varepsilon, s}, \tag{4-3}$$

and the operator estimate (3-22) implies that

$$\liminf_{N \rightarrow \infty} \frac{\langle \tilde{\Psi}_N, \tilde{H}_N \tilde{\Psi}_N \rangle}{N} \geq \liminf_{N \rightarrow \infty} (\text{Tr}(\tilde{h} \gamma_{\tilde{\Psi}_N}^{(1)}) + (1 - \varepsilon)^2 \text{Tr}(U_R \gamma_{\tilde{\Psi}_N}^{(2)})). \tag{4-4}$$

Here $\gamma_{\tilde{\Psi}_N}^{(k)}$ is the k -particle density matrix of $\tilde{\Psi}_N$ and U_R is understood as the multiplication operator $U_R(x - y)$ on \mathfrak{H}^2 . Since $\text{Tr}(\tilde{h} \gamma_{\tilde{\Psi}_N}^{(1)})$ is bounded uniformly in N and \tilde{h} has compact resolvent, up to a subsequence we can assume that $\gamma_{\tilde{\Psi}_N}^{(1)}$ converges strongly in trace class. By the quantum de Finetti theorem, Theorem 2.2, up to a subsequence we can find a Borel probability measure $\tilde{\mu}$ on the unit sphere $S\mathfrak{H}$ such that

$$\lim_{N \rightarrow \infty} \text{Tr} \left| \gamma_{\tilde{\Psi}_N}^{(k)} - \int |u^{\otimes k} \rangle \langle u^{\otimes k}| d\tilde{\mu}(u) \right| = 0 \quad \forall k \in \mathbb{N}. \tag{4-5}$$

We will show that

$$\liminf_{N \rightarrow \infty} (\text{Tr}(\tilde{h} \gamma_{\tilde{\Psi}_N}^{(1)}) + (1 - \varepsilon)^2 \text{Tr}(U_R \gamma_{\tilde{\Psi}_N}^{(2)})) \geq \int \left(\langle u, \tilde{h}u \rangle + (1 - \varepsilon)^2 4\pi a \int |u|^4 \right) d\tilde{\mu}(u), \tag{4-6}$$

and then the lower bound in (4-1) follows immediately. Since \tilde{h} is positive and independent of N , (4-5) and Fatou's lemma imply

$$\liminf_{N \rightarrow \infty} \text{Tr}(\tilde{h} \gamma_{\tilde{\Psi}_N}^{(1)}) \geq \int \langle u, \tilde{h}u \rangle d\tilde{\mu}(u). \tag{4-7}$$

It remains to prove

$$\liminf_{N \rightarrow \infty} \text{Tr}(U_R \gamma_{\tilde{\Psi}_N}^{(2)}) \geq 4\pi a \int \|u\|_{L^4}^4 d\tilde{\mu}(u). \tag{4-8}$$

Note that (4-8) does not follow from (4-5) and Fatou's lemma easily because U_R depends on $R = R(N)$. We need to replace U_R by an operator bounded independently of N . Since \tilde{h} has compact resolvent, for every $\Lambda \geq 1$, the projection

$$P_\Lambda := \mathbb{1}(\tilde{h} \leq \Lambda)$$

has finite rank. Let us denote

$$\Pi := \mathbb{1}_{\mathfrak{H}^2} - P_\Lambda^{\otimes 2}.$$

Since $U_R \geq 0$, we can apply the Cauchy–Schwarz inequality (3-9) with

$$X = P_\Lambda^{\otimes 2} U_R^{1/2} \quad \text{and} \quad Y = U_R^{1/2} \Pi$$

to obtain

$$\begin{aligned} U_R &= (P_\Lambda^{\otimes 2} + \Pi) U_R (P_\Lambda^{\otimes 2} + \Pi) \\ &= P_\Lambda^{\otimes 2} U_R P_\Lambda^{\otimes 2} + \Pi U_R \Pi + P_\Lambda^{\otimes 2} U_R \Pi + \Pi U_R P_\Lambda^{\otimes 2} \\ &\geq P_\Lambda^{\otimes 2} U_R P_\Lambda^{\otimes 2} - \delta^{-1} \Pi U_R \Pi - \delta P_\Lambda^{\otimes 2} U_R P_\Lambda^{\otimes 2} \end{aligned}$$

for all $\delta > 0$. Using the operator bound (3-4) and the fact that the $\frac{4}{5}$ -th power is operator monotone [Bhatia 1997], we have

$$U_R(x_1 - x_2) \leq C \|U_R\|_{L^1} (1 - \Delta_1)^{4/5} (1 - \Delta_2)^{4/5} \leq C_{\varepsilon,s} (\tilde{h}_1)^{4/5} (\tilde{h}_2)^{4/5}. \quad (4-9)$$

Therefore,

$$P_\Lambda^{\otimes 2} U_R P_\Lambda^{\otimes 2} \leq C_{\varepsilon,s} \tilde{h}_1 \tilde{h}_2 \quad \text{and} \quad \Pi U_R \Pi \leq C_{\varepsilon,s} \Lambda^{-1/5} \tilde{h}_1 \tilde{h}_2.$$

Here in the second estimate, we have used $\mathbb{1}_{\mathcal{J}} - P_\Lambda \leq \Lambda^{-1/5} (\tilde{h})^{1/5}$, which is a consequence of the definition of P_Λ . Thus

$$U_R - P_\Lambda^{\otimes 2} U_R P_\Lambda^{\otimes 2} \geq -C_{\varepsilon,s} (\delta^{-1} + \delta \Lambda^{-1/5}) \tilde{h}_1 \tilde{h}_2.$$

If we choose $\delta = \Lambda^{-1/10}$ and take the trace against $\gamma_{\tilde{\Psi}_N}^{(2)}$, then by the a priori estimate (4-3), we find

$$\text{Tr}(U_R \gamma_{\tilde{\Psi}_N}^{(2)}) - \text{Tr}(P_\Lambda^{\otimes 2} U_R P_\Lambda^{\otimes 2} \gamma_{\tilde{\Psi}_N}^{(2)}) \geq -C_{\varepsilon,s} \Lambda^{-1/10}. \quad (4-10)$$

On the other hand, from (4-9) and the definition of P_Λ , it follows that the operator norm $\|P_\Lambda^{\otimes 2} U_R P_\Lambda^{\otimes 2}\|$ is bounded uniformly in N for fixed Λ . Therefore, the strong convergence (4-5) implies that

$$\lim_{N \rightarrow \infty} \left(\text{Tr}(P_\Lambda^{\otimes 2} U_R P_\Lambda^{\otimes 2} \gamma_{\tilde{\Psi}_N}^{(2)}) - \int \langle (P_\Lambda u)^{\otimes 2}, U_R (P_\Lambda u)^{\otimes 2} \rangle d\tilde{\mu}(u) \right) = 0. \quad (4-11)$$

Since the left side of (4-7) is finite, every function u in the support of $d\tilde{\mu}$ belongs to the quadratic form domain $Q(\tilde{h})$ of \tilde{h} , and hence $P_\Lambda u \rightarrow u$ strongly in $Q(\tilde{h})$. Using the continuous embeddings $Q(\tilde{h}) \subset H^1 \subset L^4$, we get

$$\lim_{\Lambda \rightarrow \infty} \lim_{R \rightarrow 0} \langle (P_\Lambda u)^{\otimes 2}, U_R (P_\Lambda u)^{\otimes 2} \rangle = \lim_{\Lambda \rightarrow \infty} \|P_\Lambda u\|_{L^4}^4 = \|u\|_{L^4}^4.$$

By Fatou's lemma,

$$\liminf_{\Lambda \rightarrow \infty} \liminf_{N \rightarrow \infty} \int \langle (P_\Lambda u)^{\otimes 2}, U_R (P_\Lambda u)^{\otimes 2} \rangle d\tilde{\mu}(u) \geq 4\pi a \int \|u\|_{L^4}^4 d\tilde{\mu}(u). \quad (4-12)$$

The desired convergence (4-8) follows from (4-10), (4-11) and (4-12). \square

Remark 4.2 (Mean-field approximation with two-body interactions).

From the preceding proposition, we obtain easily the convergence (2-11) mentioned in Section 2 because $\tilde{H}_N \leq K_N$. In fact, K_N satisfies the second moment estimate (3-11) (see Remark 3.3), and hence (2-11) can be proved directly. In particular, the method can be used to derive the energy asymptotics when the interaction potential is given by (1-5); for $\beta < \frac{2}{3}$, Step 1 (and thus also Step 3) in the proof are not needed.

One can also obtain some explicit error estimate in Proposition 4.1 and (2-11) by using a quantitative version of the quantum de Finetti theorem as in [Lewin et al. 2015a, Lemma 3.4].

4B. Convergence of ground-state energy. We now conclude the proof of the convergence of the ground-state energy. There are two things left to do: remove the high momentum cut-off in the final effective functional, and relax the additional assumption (2-6).

Proof of energy convergence (1-9). The upper bound in (1-9) was proved in [Seiringer 2003]. The proof of the lower bound is divided into three steps.

Step 1. We start with the simple case when the extra condition (2-6) holds true. Recall that we are choosing

$$N^{-1/2} \gg R = R(N) \gg N^{-2/3}.$$

From Lemma 2.1 and Proposition 4.1, it follows that for every $1 > \varepsilon > 0$ and $s > 0$,

$$\liminf_{N \rightarrow \infty} \frac{\inf \sigma(H_N)}{N} \geq \liminf_{N \rightarrow \infty} \left(\frac{\inf \sigma(\tilde{H}_N)}{N} + \kappa_{\varepsilon,s} \right) = e_{\text{NL}}(\varepsilon, s) + \kappa_{\varepsilon,s}.$$

Thus to obtain the lower bound in (1-9), it remains to show that

$$\lim_{\varepsilon \rightarrow 0} \lim_{s \rightarrow \infty} (e_{\text{NL}}(\varepsilon, s) + \kappa_{\varepsilon,s}) = e_{\text{GP}}. \tag{4-13}$$

The upper bound in (4-13) is trivial as $\mathcal{E}_{\text{NL}}(u) + \kappa_{\varepsilon,s} \leq \mathcal{E}_{\text{GP}}(u)$. The lower bound in (4-13) can be done by a standard compactness argument provided in [Lieb and Seiringer 2006]. We recall this here for the reader’s convenience. Let $u_{\varepsilon,s}$ be a ground state for $e_{\text{NL}}(\varepsilon, s)$, namely

$$e_{\text{NL}}(\varepsilon, s) = \langle u_{\varepsilon,s}, \tilde{h}u_{\varepsilon,s} \rangle + (1 - \varepsilon)^2 4\pi a \int |u_{\varepsilon,s}|^4.$$

From (3-1), it follows that $\langle u_{\varepsilon,s}, (-\Delta + V)u_{\varepsilon,s} \rangle$ is bounded uniformly in s . Since $-\Delta + V$ has compact resolvent, for every given $\varepsilon > 0$, there exists a subsequence $s_j \rightarrow \infty$ such that u_{ε,s_j} converges strongly in L^2 and pointwise (in both p -space and x -space) to a function u_ε . By Fatou’s lemma, we have

$$\begin{aligned} \liminf_{j \rightarrow \infty} \int |u_{\varepsilon,s_j}(x)|^4 dx &\geq \int |u_\varepsilon(x)|^4 dx, \\ \liminf_{j \rightarrow \infty} \int p^2(1 - \theta_{s_j}(p))|\hat{u}_{\varepsilon,s_j}(p)|^2 dp &\geq \int p^2|\hat{u}_\varepsilon(p)|^2 dp. \end{aligned}$$

Next, using (2-6) as before, we have

$$\varepsilon p^2 + pA + Ap + |A|^2 + V + C_\varepsilon \geq 0$$

for some $C_\varepsilon \geq 0$. Using Fatou’s lemma again and the strong convergence in L^2 , we deduce

$$\liminf_{j \rightarrow \infty} \langle u_{\varepsilon,s_j}, (\varepsilon p^2 + pA + Ap + |A|^2 + V + \kappa_{\varepsilon,s})u_{\varepsilon,s_j} \rangle \geq \langle u_\varepsilon, (\varepsilon p^2 + pA + Ap + |A|^2 + V)u_\varepsilon \rangle.$$

Combining these estimates, we get

$$\liminf_{j \rightarrow \infty} (e_{\text{NL}}(\varepsilon, s_j) + \kappa_{\varepsilon,s_j}) \geq \langle u_\varepsilon, hu_\varepsilon \rangle + (1 - \varepsilon)^2 4\pi a \int |u_\varepsilon|^4 \geq (1 - \varepsilon)^2 e_{\text{GP}}.$$

Taking $\varepsilon \rightarrow 0$, we obtain the lower bound in (4-13).

Step 2. From now on we do not assume (2-6). Let us introduce the Hamiltonian

$$H_{M,N} := \sum_{j=1}^M h_j + \sum_{1 \leq i < j \leq M} w_N(x_i - x_j)$$

and denote by $E(M, N)$ its (bosonic) ground-state energy. In this step, we will prove the lower bound in (1-9) using the additional assumption

$$E(N, N) - E(N - 1, N) \leq C. \tag{4-14}$$

We will find a function $f : \mathbb{R}^3 \rightarrow \mathbb{R}_+$ growing faster than $|A|$, namely

$$\lim_{|x| \rightarrow \infty} \frac{|A(x)|}{f(x)} = 0, \tag{4-15}$$

such that for a ground state Ψ_N for H_N , we have

$$\langle \Psi_N, f^2(x_1) \Psi_N \rangle \leq C. \tag{4-16}$$

Once this is achieved, we get

$$\frac{\inf \sigma(H_N)}{N} \geq \frac{\inf \sigma(H_N + \eta \sum_{j=1}^N f^2(x_j))}{N} - C\eta$$

for every $\eta > 0$. Since the growth condition (2-6) holds true with V replaced by $V + \eta f^2$, we can apply the result in Step 1 to the Hamiltonian

$$H_N + \eta \sum_{j=1}^N f^2(x_j)$$

for every given $\eta > 0$. Then the lower bound in (1-9) follows by taking $\eta \rightarrow 0$.

Now we find such a function f . We will establish a simple binding inequality using an idea in [Lieb 1984]. From the ground-state equation $H_{N,N} \Psi_N = E(N, N) \Psi_N$, it follows that

$$E(N, N) \langle \Psi_N, f^2(x_N) \Psi_N \rangle = \Re \langle \Psi_N, f^2(x_N) H_{N,N} \Psi_N \rangle. \tag{4-17}$$

By the variational principle and (4-14), we have

$$H_{N,N} - h_N \geq H_{N-1,N} \geq E(N - 1, N) \geq E(N, N) - C.$$

Note that $f^2(x_N)$ commutes with all terms in the latter inequality. If f is bounded and sufficiently regular, we have the IMS-type formula

$$\frac{1}{2}(f^2 h + h f^2) = f h f - |\nabla f|^2 \geq V f^2 - |\nabla f|^2, \tag{4-18}$$

and we deduce from (4-17) that

$$\langle \Psi_N, (V(x_N) f^2(x_N) - |\nabla f(x_N)|^2 - C f^2(x_N)) \Psi_N \rangle \leq 0. \tag{4-19}$$

Note that if we choose $f(x) = e^{b|x|}$ for some constant $b > 0$, then (4-15) follows from the assumption (1-2). Moreover, heuristically, (4-16) follows from (4-19) as Vf^2 grows faster than $|\nabla f|^2 + Cf^2$. To make this idea rigorous, let us apply (4-19) with $f(x)$ replaced by

$$g_r(x) = \exp[b[r - ||x| - r|]_+].$$

Note that $g_r(x) = e^{b|x|}$ when $|x| \leq r$ and $g_r(x) = 1$ when $|x| \geq 2r$. We can thus apply (4-18) to g_r .

Moreover,

$$\begin{aligned} Vg_r^2 - |\nabla g_r|^2 - Cg_r^2 &\geq (V - b^2 - C)g_r^2 \\ &\geq g_r^2 - (b^2 + C + 1)g_r^2 \mathbb{1}(V \leq b^2 + C + 1) \\ &\geq g_r^2 - C_0 \end{aligned}$$

for some constant C_0 independent of $r > 0$. Here we have used the fact that $g_r^2 \mathbb{1}(V \leq b^2 + C + 1)$ is bounded independently of $r > 0$, which follows from the assumption $\lim_{|x| \rightarrow \infty} V(x) = +\infty$. Thus (4-19) gives us

$$\langle \Psi_N, g_r(x_N) \Psi_N \rangle \leq C_0$$

for all $r > 0$. Taking $r \rightarrow \infty$, we obtain (4-16) with $f(x) = e^{b|x|}$.

Step 3. Now we explain how to remove the additional assumption (4-14). This can be done by following the strategy in [Lieb and Seiringer 2006], which we recall quickly below for the reader’s convenience.

By choosing trial states $u^{\otimes N}$, we get the upper bound

$$E(N, N) \leq C_0 N$$

for some constant $C_0 > 2e_{\text{GP}}$. For every $N \in \mathbb{N}$, we denote by $M = M(N)$ the largest integer $\leq N$ such that

$$E(M(N), N) - E(M(N) - 1, N) \leq C_0. \tag{4-20}$$

Then by the choice of $M(N)$, we obtain

$$E(N, N) - E(M(N), N) \geq (N - M(N))C_0. \tag{4-21}$$

We can find a subsequence $N_j \rightarrow \infty$ such that $M(N_j)/N_j \rightarrow \lambda \in [0, 1]$. Since (4-20) holds with $M = M(N_j)$, we can apply the result in Step 2 with w replaced by λw and find that

$$\liminf_{j \rightarrow \infty} \frac{E(M(N_j), N_j)}{M_j} \geq e_{\text{GP}}(\lambda a) \geq \lambda e_{\text{GP}}(a). \tag{4-22}$$

Here $e_{\text{GP}}(\lambda a)$ is the Gross–Pitaevskii energy with a replaced by λa and the last inequality in (4-22) is obtained by simply ignoring part of the one-body energy in the corresponding Gross–Pitaevskii functional. From (4-21) and (4-22), it follows that

$$\begin{aligned} e_{\text{GP}}(a) &\geq \liminf_{j \rightarrow \infty} \frac{E(N_j, N_j)}{N_j} \geq \liminf_{j \rightarrow \infty} \left(\frac{E(M(N_j), N_j)}{N_j} + C_0 \frac{N_j - M(N_j)}{N_j} \right) \\ &\geq \lambda^2 e_{\text{GP}}(a) + C_0(1 - \lambda). \end{aligned}$$

Since

$$e_{\text{GP}}(a) \leq \lambda^2 e_{\text{GP}}(a) + 2(1 - \lambda)e_{\text{GP}}(a)$$

and $C_0 > 2e_{\text{GP}}(a)$, we must have $\lambda = 1$. Thus $M(N)/N \rightarrow 1$ for the whole sequence and

$$\liminf_{N \rightarrow \infty} \frac{E(N, N)}{N} = \liminf_{j \rightarrow \infty} \frac{E(N_j, N_j)}{N_j} \geq e_{\text{GP}}(a).$$

This completes the proof of the energy convergence (1-9). □

4C. Convergence of density matrices. Now we prove the convergence of ground states in (1-10) by means of the Feynman–Hellmann principle. For $v \in L^2(\mathbb{R}^3)$ and $\ell \in \mathbb{N}$, we will perturb H_N by

$$S_{v,\ell} := \frac{\ell!}{N^{\ell-1}} \sum_{1 \leq i_1 < \dots < i_\ell \leq N} |v^{\otimes \ell}\rangle \langle v^{\otimes \ell}|_{i_1, \dots, i_\ell}.$$

Here $|v^{\otimes \ell}\rangle \langle v^{\otimes \ell}|_{i_1, \dots, i_\ell}$ denotes the operator $|v^{\otimes \ell}\rangle \langle v^{\otimes \ell}|$ acting on the ℓ -body Hilbert space of the i_1 -th, ..., i_ℓ -th variables. We have the following extension of (1-9).

Lemma 4.3 (Energy lower bound for perturbed Hamiltonians).

We assume (1-2), (1-3) and (1-4). For every $v \in L^2(\mathbb{R}^3)$ and $\ell \in \mathbb{N}$, we have

$$\liminf_{N \rightarrow \infty} \frac{\inf \sigma(H_N - S_{v,\ell})}{N} \geq \inf_{\|u\|_{L^2}=1} (\mathcal{E}_{\text{GP}}(u) - |\langle v, u \rangle|^{2\ell}). \tag{4-23}$$

Proof. We first work under the extra condition (2-6), and then explain how to remove it at the end. Let $1 > \varepsilon > 0$ and $s > 0$ and

$$N^{-1/2} \gg R = R(N) \gg N^{-2/3}.$$

Recall that from (2-2), we have

$$H_N - S_{v,\ell} \geq \tilde{H}_N - S_{v,\ell} + N\kappa_{\varepsilon,s} - C_{\varepsilon,s}NR^2. \tag{4-24}$$

Let Φ_N be a ground state for $\tilde{H}_N - S_{v,\ell}$. Since $\|S_{v,\ell}\|/N$ is bounded uniformly in N , (4-2) still holds true with $\tilde{\Psi}_N$ replaced by Φ_N , namely

$$\langle \Phi_N, (\tilde{H}_N)^k \Phi_N \rangle \leq (C_{\varepsilon,s}N)^k \tag{4-25}$$

for all $k \in \mathbb{N}$. Combining (4-25) with the three-body estimate in Lemma 3.4, we get the following analogue of (4-4):

$$\begin{aligned} \liminf_{N \rightarrow \infty} \frac{\inf \sigma(\tilde{H}_N - S_{v,\ell})}{N} &= \liminf_{N \rightarrow \infty} \frac{\langle \Phi_N, (\tilde{H}_N - S_{v,\ell}) \Phi_N \rangle}{N} \\ &\geq \liminf_{N \rightarrow \infty} (\text{Tr}(\tilde{h}\gamma_{\tilde{\Psi}_N}^{(1)}) + (1 - \varepsilon)^2 \text{Tr}(U_R\gamma_{\tilde{\Psi}_N}^{(2)}) - \text{Tr}(|v^{\otimes \ell}\rangle \langle v^{\otimes \ell}| \gamma_{\Phi_N}^{(\ell)})). \end{aligned} \tag{4-26}$$

Moreover, (4-25) and the second moment estimate (3-2) imply the a priori estimate $\langle \Phi_N, \tilde{h}_1 \tilde{h}_2 \Psi_N \rangle \leq C_{\varepsilon,s}$. Therefore, we can estimate the right side of (4-26) by proceeding exactly as in the proof of Proposition 4.1. More precisely, by the quantum de Finetti theorem, Theorem 2.2, we can find a Borel probability measure μ_Φ on the unit sphere $S\mathfrak{H}$ such that, up to a subsequence,

$$\lim_{N \rightarrow \infty} \text{Tr} \left| \gamma_{\Phi_N}^{(k)} - \int |u^{\otimes k}\rangle \langle u^{\otimes k}| d\mu_\Phi(u) \right| = 0 \quad \forall k \in \mathbb{N}.$$

Using (4-6) with $\tilde{\Psi}_N$ replaced by Φ_N and employing the fact that $|v^{\otimes \ell}\rangle\langle v^{\otimes \ell}|$ is bounded, we obtain

$$\begin{aligned} \liminf_{N \rightarrow \infty} (\text{Tr}(\tilde{h}\gamma_{\tilde{\Psi}_N}^{(1)}) + (1 - \varepsilon)^2 \text{Tr}(U_R\gamma_{\tilde{\Psi}_N}^{(2)}) - \text{Tr}(|v^{\otimes \ell}\rangle\langle v^{\otimes \ell}| \gamma_{\Phi_N}^{(\ell)})) \\ \geq \int \left(\langle u, \tilde{h}u \rangle + (1 - \varepsilon)^2 4\pi a \int |u|^4 - |\langle v, u \rangle|^{2\ell} \right) d\mu_{\Phi}(u). \end{aligned} \quad (4-27)$$

From (4-24), (4-26) and (4-27), it follows that

$$\liminf_{N \rightarrow \infty} \frac{\inf \sigma(H_N - S_{v,\ell})}{N} \geq \inf_{\|u\|_{L^2}=1} \left(\langle u, \tilde{h}u \rangle + (1 - \varepsilon)^2 4\pi a \int |u|^4 - |\langle v, u \rangle|^{2\ell} \right) + \kappa_{\varepsilon,s}.$$

The lower bound (4-23) follows by passing to the limits $s \rightarrow 0$ and then $\varepsilon \rightarrow 0$ as in the proof of (4-13).

To remove the assumption (2-6), we may use the argument in Section 4B. The only extra difficulty is that when dealing with the analogue of (4-17) with $H_{N,N}$ replaced by $H_{N,N} - S_{v,\ell}$, we have to take care of the operator $f^2|v\rangle\langle v| = |f^2v\rangle\langle v|$, which may be unbounded as $f(x) = e^{b|x|}$ with $b > 0$ and v is merely in $L^2(\mathbb{R}^3)$. However, we can still proceed with all functions v in $L^2(\mathbb{R}^3)$ which have compact support. Then after obtaining the lower bound (4-23) with those nice functions v , we can extend the lower bound to all functions v in $L^2(\mathbb{R}^3)$ by a standard density argument. \square

Now we are able to prove the convergence of density matrices.

Proof of state convergence (1-10). Let Ψ_N be an approximate ground state for H_N as in Theorem 1.1. For every $v \in L^2(\mathbb{R}^3)$ and $\ell \in \mathbb{N}$, from the upper bound in (1-9) and the lower bound in Lemma 4.3, we have

$$\begin{aligned} \limsup_{N \rightarrow \infty} \text{Tr}(|v^{\otimes \ell}\rangle\langle v^{\otimes \ell}| \gamma_{\Psi_N}^{(\ell)}) &= \limsup_{N \rightarrow \infty} \left(\frac{\langle \Psi_N, H_N \Psi_N \rangle}{N} - \frac{\langle \Psi_N, (H_N - S_{v,\ell}) \Psi_N \rangle}{N} \right) \\ &\leq \limsup_{N \rightarrow \infty} \left(\frac{\inf \sigma(H_N)}{N} - \frac{\inf \sigma(H_N - S_{v,\ell})}{N} \right) \\ &\leq e_{\text{GP}} - \inf_{\|u\|_{L^2}=1} (\mathcal{E}_{\text{GP}}(u) - |\langle v, u \rangle|^{2\ell}). \end{aligned}$$

Here v is not necessarily normalized. Therefore, we can replace v by $\lambda^{1/(2\ell)}v$ with $\lambda > 0$ and obtain

$$\limsup_{N \rightarrow \infty} \text{Tr}(|v^{\otimes \ell}\rangle\langle v^{\otimes \ell}| \gamma_{\Psi_N}^{(\ell)}) \leq \frac{1}{\lambda} \left(e_{\text{GP}} - \inf_{\|u\|_{L^2}=1} (\mathcal{E}_{\text{GP}}(u) - \lambda |\langle v, u \rangle|^{2\ell}) \right). \quad (4-28)$$

With given v and ℓ , for every $\lambda > 0$, let u_λ be a (normalized) minimizer for $u \mapsto \mathcal{E}_{\text{GP}}(u) - \lambda |\langle v, u \rangle|^{2\ell}$. Since $\langle u_\lambda, hu_\lambda \rangle$ is bounded and h has compact resolvent, there exists a subsequence $\lambda_j \rightarrow 0$ such that u_{λ_j} converges to u_0 in L^2 . By Fatou's lemma, u_0 is a minimizer of $\mathcal{E}_{\text{GP}}(u)$. Moreover,

$$\begin{aligned} \limsup_{j \rightarrow \infty} \frac{1}{\lambda_j} \left(e_{\text{GP}} - \inf_{\|u\|_{L^2}=1} (\mathcal{E}_{\text{GP}}(u) - \lambda_j |\langle v, u \rangle|^{2\ell}) \right) \\ \leq \limsup_{j \rightarrow \infty} \frac{1}{\lambda_j} \left(\mathcal{E}_{\text{GP}}(u_{\lambda_j}) - (\mathcal{E}_{\text{GP}}(u_{\lambda_j}) - \lambda_j |\langle v, u_{\lambda_j} \rangle|^{2\ell}) \right) = |\langle v, u_0 \rangle|^{2\ell}. \end{aligned} \quad (4-29)$$

From (4-28) and (4-29), we conclude that for every $v \in L^2(\mathbb{R}^3)$ and $\ell \in \mathbb{N}$,

$$\limsup_{N \rightarrow \infty} \text{Tr}(|v^{\otimes \ell}\rangle\langle v^{\otimes \ell}| \gamma_{\Psi_N}^{(\ell)}) \leq \sup_{u \in \mathcal{M}_{\text{GP}}} |\langle v, u \rangle|^{2\ell}, \tag{4-30}$$

where \mathcal{M}_{GP} is the set of minimizers of $\mathcal{E}_{\text{GP}}(u)$.

Note that also in [Lieb and Seiringer 2006], the upper bound (4-30) with $\ell = 1$ was proved, and from it, the convergence of the one-particle density matrices was deduced using an abstract argument of convex analysis. In the following, we will provide a simpler way to conclude the convergence of density matrices from (4-30), using the quantum de Finetti theorem. Indeed, by Theorem 2.2 as before, up to a subsequence of Ψ_N , there exists a Borel probability measure μ on the unit sphere $S\mathfrak{H}$ such that

$$\lim_{N \rightarrow \infty} \text{Tr} \left| \gamma_{\Psi_N}^{(k)} - \int |u^{\otimes k}\rangle\langle u^{\otimes k}| d\mu(u) \right| = 0 \quad \forall k \in \mathbb{N}. \tag{4-31}$$

We will show that μ is supported on \mathcal{M}_{GP} . From (4-30) and (4-31), we get

$$\int |\langle v, u \rangle|^{2k} d\mu(u) \leq \sup_{u \in \mathcal{M}_{\text{GP}}} |\langle v, u \rangle|^{2k} \quad \forall v \in L^2(\mathbb{R}^3), k \in \mathbb{N}. \tag{4-32}$$

We assume for contradiction that there exists v_0 in the support of μ and $v_0 \notin \mathcal{M}_{\text{GP}}$. We claim that we could then find $\delta \in (0, \frac{1}{2})$ such that

$$|\langle v, u \rangle| \leq 1 - 3\delta^2 \quad \forall u \in \mathcal{M}_{\text{GP}}, \forall v \in B, \tag{4-33}$$

where B is the set of all points in the support of μ within an L^2 -distance less than δ from v_0 . Indeed, if that were not the case, we would have two sequences strongly converging in L^2 ,

$$v_n \rightarrow v_0, \quad u_n \rightarrow u_0 \in \mathcal{M}_{\text{GP}},$$

with $\|u_n - v_n\| \rightarrow 0$, and thus $v_0 \in \mathcal{M}_{\text{GP}}$. Here we have used that \mathcal{M}_{GP} is a compact subset of $L^2(\mathbb{R}^3)$.

On the other hand, by the triangle inequality,

$$|\langle v, u \rangle| \geq \frac{\|u\|^2 + \|v\|^2 - \|u - v\|^2}{2} \geq 1 - 2\delta^2 \quad \forall u, v \in B. \tag{4-34}$$

Combining (4-32), (4-33) and (4-34), we find that

$$\begin{aligned} (\mu(B))^2 (1 - 2\delta^2)^{2k} &\leq \int_B \int_B |\langle v, u \rangle|^{2k} d\mu(u) d\mu(v) \\ &\leq \int_B \sup_{u \in \mathcal{M}_{\text{GP}}} |\langle v, u \rangle|^{2k} d\mu(v) \leq \mu(B) (1 - 3\delta^2)^{2k} \end{aligned} \tag{4-35}$$

for all $k \in \mathbb{N}$, and hence, taking $k \rightarrow \infty$, we have $\mu(B) = 0$. This, however, is a contradiction to the fact that v_0 belongs to the support of μ and μ is a Borel measure. Thus we conclude that μ is supported on \mathcal{M}_{GP} and the proof is complete. □

Acknowledgments

We thank Mathieu Lewin for helpful discussions. Phan Thành Nam is supported partially by the European Union's Seventh Framework Programme under REA grant agreement no. 291734. Nicholas Rougerie acknowledges financial support from the ANR (MaThoStaQ project, ANR-13-JS01-0005-01).

References

- [Benedikter et al. 2015] N. Benedikter, G. de Oliveira, and B. Schlein, “Quantitative derivation of the Gross–Pitaevskii equation”, *Comm. Pure Appl. Math.* **68**:8 (2015), 1399–1482. [MR 3366749](#) [Zbl 1320.35318](#)
- [Benedikter et al. 2016] N. Benedikter, M. Porta, and B. Schlein, *Effective evolution equations from quantum dynamics*, SpringerBriefs in Mathematical Physics **7**, Springer, Cham, 2016. [MR 3382225](#) [Zbl 06492716](#)
- [Bhatia 1997] R. Bhatia, *Matrix analysis*, Graduate Texts in Mathematics **169**, Springer, New York, 1997. [MR 98i:15003](#) [Zbl 0863.15001](#)
- [Chen et al. 2015] T. Chen, C. Hainzl, N. Pavlović, and R. Seiringer, “Unconditional uniqueness for the cubic Gross–Pitaevskii hierarchy via quantum de Finetti”, *Comm. Pure Appl. Math.* **68**:10 (2015), 1845–1884. [MR 3385343](#) [Zbl 1326.35332](#)
- [Chiribella 2011] G. Chiribella, “On quantum estimation, quantum cloning and finite quantum de Finetti theorems”, pp. 9–25 in *Theory of quantum computation, communication, and cryptography*, edited by W. van Dam et al., Lecture Notes in Comput. Sci. **6519**, Springer, Berlin, 2011. [MR 2012i:81019](#) [Zbl 1310.81034](#)
- [Christandl et al. 2007] M. Christandl, R. König, G. Mitchison, and R. Renner, “One-and-a-half quantum de Finetti theorems”, *Comm. Math. Phys.* **273**:2 (2007), 473–498. [MR 2008m:81100](#) [Zbl 1126.81032](#)
- [Dyson 1957] F. J. Dyson, “Ground-state energy of a hard-sphere gas”, *Phys. Rev.* **106**:1 (1957), 20–26. [Zbl 0077.23503](#)
- [Erdős and Yau 2001] L. Erdős and H.-T. Yau, “Derivation of the nonlinear Schrödinger equation from a many body Coulomb system”, *Adv. Theor. Math. Phys.* **5**:6 (2001), 1169–1205. [MR 2004c:82075](#) [Zbl 1014.81063](#)
- [Erdős et al. 2007] L. Erdős, B. Schlein, and H.-T. Yau, “Derivation of the cubic non-linear Schrödinger equation from quantum dynamics of many-body systems”, *Invent. Math.* **167**:3 (2007), 515–614. [MR 2007m:81258](#) [Zbl 1123.35066](#)
- [Erdős et al. 2009] L. Erdős, B. Schlein, and H.-T. Yau, “Rigorous derivation of the Gross–Pitaevskii equation with a large interaction potential”, *J. Amer. Math. Soc.* **22**:4 (2009), 1099–1156. [MR 2010g:82041](#) [Zbl 1207.82031](#)
- [Erdős et al. 2010] L. Erdős, B. Schlein, and H.-T. Yau, “Derivation of the Gross–Pitaevskii equation for the dynamics of Bose–Einstein condensate”, *Ann. of Math. (2)* **172**:1 (2010), 291–370. [MR 2011g:82068](#) [Zbl 1204.82028](#)
- [Hudson and Moody 1976] R. L. Hudson and G. R. Moody, “Locally normal symmetric states and an analogue of de Finetti’s theorem”, *Z. Wahrscheinlichkeitstheorie und Verw. Gebiete* **33**:4 (1976), 343–351. [MR 53 #1280](#) [Zbl 0304.60001](#)
- [Lewin et al. 2014] M. Lewin, P. T. Nam, and N. Rougerie, “Derivation of Hartree’s theory for generic mean-field Bose systems”, *Adv. Math.* **254** (2014), 570–621. [MR 3161107](#) [Zbl 1316.81095](#)
- [Lewin et al. 2015a] M. Lewin, P. T. Nam, and N. Rougerie, “The mean-field approximation and the non-linear Schrödinger functional for trapped Bose gases”, *Trans. Amer. Math. Soc.* (online publication October 2015).
- [Lewin et al. 2015b] M. Lewin, P. T. Nam, and N. Rougerie, “Remarks on the quantum de Finetti theorem for bosonic systems”, *Appl. Math. Res. Express. AMRX* **1** (2015), 48–63. [MR 3335056](#) [Zbl 1310.81169](#)
- [Lieb 1984] E. H. Lieb, “Bound on the maximum negative ionization of atoms and molecules”, *Phys. Rev. A* **29**:6 (1984), 3018–3028.
- [Lieb and Seiringer 2002] E. H. Lieb and R. Seiringer, “Proof of Bose–Einstein condensation for dilute trapped gases”, *Phys. Rev. Lett.* **88**:17 (2002), paper ID 170409.
- [Lieb and Seiringer 2006] E. H. Lieb and R. Seiringer, “Derivation of the Gross–Pitaevskii equation for rotating Bose gases”, *Commun. Math. Phys.* **264**:2 (2006), 505–537. [Zbl 1233.82004](#)
- [Lieb and Yngvason 1998] E. H. Lieb and J. Yngvason, “Ground state energy of the low density Bose gas”, *Phys. Rev. Lett.* **80**:12 (1998), 2504–2507.

- [Lieb et al. 2000] E. H. Lieb, R. Seiringer, and J. Yngvason, “Bosons in a trap: a rigorous derivation of the Gross–Pitaevskii energy functional”, *Phys. Rev. A* **61**:4 (2000), paper ID 043602.
- [Lieb et al. 2005a] E. H. Lieb, R. Seiringer, and J. P. Solovej, “Ground-state energy of the low-density Fermi gas”, *Phys. Rev. A* **71**:5 (2005), paper ID 053605.
- [Lieb et al. 2005b] E. H. Lieb, R. Seiringer, J. P. Solovej, and J. Yngvason, *The mathematics of the Bose gas and its condensation*, Oberwolfach Seminars **34**, Birkhäuser, Basel, 2005. MR 2006e:82001 Zbl 1104.82012
- [Lieb et al. 2005c] E. H. Lieb, R. Seiringer, and J. Yngvason, “Justification of c -number substitutions in bosonic Hamiltonians”, *Phys. Rev. Lett.* **94**:8 (2005), paper ID 080401.
- [Pickl 2015] P. Pickl, “Derivation of the time dependent Gross–Pitaevskii equation with external fields”, *Rev. Math. Phys.* **27**:1 (2015), paper ID 1550003. MR 3317556 Zbl 1320.35328
- [Rougerie 2015] N. Rougerie, “De Finetti theorems, mean-field limits and Bose–Einstein condensation”, lecture notes, 2015. arXiv 1506.05263
- [Seiringer 2003] R. Seiringer, “Ground state asymptotics of a dilute, rotating gas”, *J. Phys. A* **36**:37 (2003), 9755–9778. MR 2004h:82016 Zbl 1151.82401
- [Størmer 1969] E. Størmer, “Symmetric states of infinite tensor products of C^* -algebras”, *J. Functional Analysis* **3** (1969), 48–68. MR 39 #3327 Zbl 0167.43403

Received 30 Apr 2015. Revised 8 Sep 2015. Accepted 6 Jan 2016.

PHAN THÀNH NAM: pnam@ist.ac.at

Institute of Science and Technology Austria, Am Campus 1, 3400 Klosterneuburg, Austria

NICOLAS ROUGERIE: nicolas.rougerie@grenoble.cnrs.fr

Université Grenoble 1 and Centre National de la Recherche Scientifique, Laboratoire de Physique et Modélisation des Milieux Condensés (UMR 5493), BP 166, 38042 Grenoble, France

ROBERT SEIRINGER: robert.seiringer@ist.ac.at

Institute of Science and Technology Austria, Am Campus 1, 3400 Klosterneuburg, Austria

Analysis & PDE

msp.org/apde

EDITORS

EDITOR-IN-CHIEF

Patrick Gérard

patrick.gerard@math.u-psud.fr

Université Paris Sud XI

Orsay, France

BOARD OF EDITORS

Nicolas Burq	Université Paris-Sud 11, France nicolas.burq@math.u-psud.fr	Werner Müller	Universität Bonn, Germany mueller@math.uni-bonn.de
Massimiliano Berti	Scuola Intern. Sup. di Studi Avanzati, Italy berti@sissa.it	Yuval Peres	University of California, Berkeley, USA peres@stat.berkeley.edu
Sun-Yung Alice Chang	Princeton University, USA chang@math.princeton.edu	Gilles Pisier	Texas A&M University, and Paris 6 pisier@math.tamu.edu
Michael Christ	University of California, Berkeley, USA mchrist@math.berkeley.edu	Tristan Rivière	ETH, Switzerland riviere@math.ethz.ch
Charles Fefferman	Princeton University, USA cf@math.princeton.edu	Igor Rodnianski	Princeton University, USA irod@math.princeton.edu
Ursula Hamenstaedt	Universität Bonn, Germany ursula@math.uni-bonn.de	Wilhelm Schlag	University of Chicago, USA schlag@math.uchicago.edu
Vaughan Jones	U.C. Berkeley & Vanderbilt University vaughan.f.jones@vanderbilt.edu	Sylvia Serfaty	New York University, USA serfaty@cims.nyu.edu
Vadim Kaloshin	University of Maryland, USA vadim.kaloshin@gmail.com	Yum-Tong Siu	Harvard University, USA siu@math.harvard.edu
Herbert Koch	Universität Bonn, Germany koch@math.uni-bonn.de	Terence Tao	University of California, Los Angeles, USA tao@math.ucla.edu
Izabella Laba	University of British Columbia, Canada ilaba@math.ubc.ca	Michael E. Taylor	Univ. of North Carolina, Chapel Hill, USA met@math.unc.edu
Gilles Lebeau	Université de Nice Sophia Antipolis, France lebeau@unice.fr	Gunther Uhlmann	University of Washington, USA gunther@math.washington.edu
László Lempert	Purdue University, USA lempert@math.purdue.edu	András Vasy	Stanford University, USA andras@math.stanford.edu
Richard B. Melrose	Massachusetts Inst. of Tech., USA rbb@math.mit.edu	Dan Virgil Voiculescu	University of California, Berkeley, USA dvv@math.berkeley.edu
Frank Merle	Université de Cergy-Pontoise, France Frank.Merle@u-cergy.fr	Steven Zelditch	Northwestern University, USA zelditch@math.northwestern.edu
William Minicozzi II	Johns Hopkins University, USA minicozz@math.jhu.edu	Maciej Zworski	University of California, Berkeley, USA zworski@math.berkeley.edu
Clément Mouhot	Cambridge University, UK c.mouhot@dpms.cam.ac.uk		

PRODUCTION

production@msp.org

Silvio Levy, Scientific Editor

See inside back cover or msp.org/apde for submission instructions.

The subscription price for 2016 is US \$235/year for the electronic version, and \$430/year (+\$55, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues from the last three years and changes of subscribers address should be sent to MSP.

Analysis & PDE (ISSN 1948-206X electronic, 2157-5045 printed) at Mathematical Sciences Publishers, 798 Evans Hall #3840, c/o University of California, Berkeley, CA 94720-3840, is published continuously online. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices.

APDE peer review and production are managed by EditFlow[®] from MSP.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2016 Mathematical Sciences Publishers

ANALYSIS & PDE

Volume 9 No. 2 2016

Resonances for large one-dimensional “ergodic” systems FRÉDÉRIC KLOPP	259
On characterization of the sharp Strichartz inequality for the Schrödinger equation JIN-CHENG JIANG and SHUANGLIN SHAO	353
Future asymptotics and geodesic completeness of polarized T^2 -symmetric spacetimes PHILIPPE G. LEFLOCH and JACQUES SMULEVICI	363
Obstacle problem with a degenerate force term KAREN YERESSIAN	397
A counterexample to the Hopf–Oleinik lemma (elliptic case) DARYA E. APUSHKINSKAYA and ALEXANDER I. NAZAROV	439
Ground states of large bosonic systems: the Gross–Pitaevskii limit revisited PHAN THÀNH NAM, NICOLAS ROUGERIE and ROBERT SEIRINGER	459
Nontransversal intersection of free and fixed boundaries for fully nonlinear elliptic operators in two dimensions EMANUEL ÎNDREI and ANDREAS MINNE	487
Correction to the article Scattering threshold for the focusing nonlinear Klein–Gordon equation SLIM IBRAHIM, NADER MASMOUDI and KENJI NAKANISHI	503