

ANALYSIS & PDE

Volume 9

No. 4

2016

KIRIL DATCHEV

**RESONANCE FREE REGIONS FOR NONTRAPPING MANIFOLDS
WITH CUSPS**

RESONANCE FREE REGIONS FOR NONTRAPPING MANIFOLDS WITH CUSPS

KIRIL DATCHEV

We prove resolvent estimates for nontrapping manifolds with cusps which imply the existence of arbitrarily wide resonance free strips, local smoothing for the Schrödinger equation, and resonant wave expansions. We obtain lossless limiting absorption and local smoothing estimates, but the estimates on the holomorphically continued resolvent exhibit losses. We prove that these estimates are optimal in certain respects.

1. Introduction

Resonance free regions near the essential spectrum have been extensively studied since the foundational work of Lax and Phillips and of Vainberg. Their size is related to the dynamical structure of the set of trapped classical trajectories. More trapping typically results in a smaller region, and the largest resonance free regions exist when there is no trapping.

Example. Let \mathbb{H}^2 be the hyperbolic upper half plane. Let (X, g) be a nonpositively curved, compactly supported, smooth, metric perturbation of the quotient space $(z \mapsto z + 1) \backslash \mathbb{H}^2$. As we show in [Section 2D](#), such a surface has no trapped geodesics (that is, all geodesics are unbounded).

Let (X, g) be as in the example above, or as in [Section 2A](#), with dimension $n + 1$ and Laplacian $\Delta \geq 0$. The resolvent $(\Delta - \frac{1}{4}n^2 - \sigma^2)^{-1}$ is holomorphic for $\text{Im } \sigma > 0$, except at any $\sigma \in i\mathbb{R}$ such that $\sigma^2 + \frac{1}{4}n^2$ is an eigenvalue, and has essential spectrum $\{\text{Im } \sigma = 0\}$; see [Figure 1](#).

Theorem. For all $\chi \in C_0^\infty(X)$, there exists $M_0 > 0$ such that for all $M_1 > 0$ there exists $M_2 > 0$ such that the cutoff resolvent $\chi(\Delta - \frac{1}{4}n^2 - \sigma^2)^{-1}\chi$ continues holomorphically to $\{|\text{Re } \sigma| \geq M_2, \text{Im } \sigma \geq -M_1\}$, where it obeys the estimate

$$\|\chi(\Delta - \frac{1}{4}n^2 - \sigma^2)^{-1}\chi\|_{L^2(X) \rightarrow L^2(X)} \leq M_2 |\sigma|^{-1+M_0|\text{Im } \sigma|}. \quad (1-1)$$

In the example above, and in many of the examples in [Section 2D](#), $\chi(\Delta - \frac{1}{4}n^2 - \sigma^2)^{-1}\chi$ is meromorphic in \mathbb{C} . The poles of the meromorphic continuation are called *resonances*.

Logarithmically large resonance free regions go back to work of Regge [\[1958\]](#) on potential scattering. In the setting of obstacle scattering they go back to work of Lax and Phillips [\[1989\]](#) and Vainberg [\[1989\]](#), whose results were generalized by Morawetz, Ralston and Strauss [\[1977\]](#) and Melrose and Sjöstrand [\[1982\]](#). When X is Euclidean outside of a compact set, they have been established for very general nontrapping perturbations of the Laplacian by Sjöstrand and Zworski in [\[2007, Theorem 1\]](#), which extends earlier work of Martinez [\[2002\]](#) and Sjöstrand [\[1990\]](#). More recently, Baskin and Wunsch [\[2013\]](#),

MSC2010: 58J50.

Keywords: cusp, resonances, resolvent, scattering, waves.




Figure 1. We prove that the cutoff resolvent continues holomorphically to arbitrarily wide strips and obeys polynomial bounds.

Galkowski and Smith [2015], and Galkowski [2015; 2016] have weakened slightly the sense in which the perturbation must be nontrapping. These works give a larger resonance free region and a stronger resolvent estimate than the Theorem above, but require asymptotically Euclidean geometry near infinity. On the other hand, as shown in recent work of Datchev, Kang and Kessler [2015], nontrapping manifolds with cusps which are merely $C^{1,1}$ (and not C^∞) do *not* have arbitrarily wide resonance free strips as in the Theorem.

The manifolds considered in this paper are nontrapping, but the cusp makes them not uniformly so: for a sufficiently large compact set $K \subset X$, we have

$$\sup_{\gamma \in \Gamma} \text{diam } \gamma^{-1}(K) = +\infty,$$

where Γ is the set of unit-speed geodesics in X . This is because geodesics may travel arbitrarily far into the cusp before escaping down the funnel; this dynamical peculiarity makes it difficult to separate the analysis in the cusp from the analysis in the funnel and is the reason for the relatively involved resolvent estimate gluing procedure we use below.

Resonance free strips also exist in some trapping situations, with width determined by dynamical properties of the trapped set. These go back to work of Ikawa [1982], with recent progress by Nonnenmacher and Zworski [2009; 2015], Petkov and Stoyanov [2010], Alexandrova and Tamura [2011], Wunsch and Zworski [2011], Dyatlov [2015b], and Dyatlov and Zahl [2015]. Resonance free regions and resolvent estimates have applications to evolution equations, and this is an active area: examples include resonant wave expansions and wave decay, local smoothing estimates, Strichartz estimates, geometric control, wave damping, and radiation fields [Burq 2004; Burq and Zworski 2004; Bony and Häfner 2008; Guillarmou and Naud 2009; Christianson 2009; Burq, Guillarmou and Hassell 2010; Dyatlov 2012; 2015a; Melrose, Sá Barreto and Vasy 2014; Christianson, Schenck, Vasy and Wunsch 2014; Wang 2014]; see also [Wunsch 2012] for a recent survey and more references. In Section 7 we apply (1-1) to local smoothing and resonant wave expansions.

If (X, g) is evenly asymptotically hyperbolic (in the sense of Mazzeo and Melrose [1987] and Guillarmou [2005]) and nontrapping, then for any $M_1 > 0$ there is $M_2 > 0$ such that

$$\|\chi(\Delta - \frac{1}{4}n^2 - \sigma^2)^{-1}\chi\|_{L^2(X) \rightarrow L^2(X)} \leq M_2|\sigma|^{-1}, \quad |\text{Re } \sigma| \geq M_2, \text{Im } \sigma \geq -M_1, \quad (1-2)$$

by work of Vasy [2013, (1.1)] (see also the analogous estimate for asymptotically Euclidean spaces by Sjöstrand and Zworski [2007, Theorem 1’], and related but slightly weaker estimates for more general asymptotically hyperbolic and conformally compact manifolds by Wang [2014] and Sá Barreto and Wang [2015]).

The bound (1-1) is weaker than (1-2) due to the presence of a cusp. Indeed, by studying low angular frequencies (which correspond to geodesics which travel far into the cusp before escaping down the funnel) in Proposition 8.1 we show that if $(X, g) = \langle z \mapsto z + 1 \rangle \backslash \mathbb{H}^2$, then

$$\| \chi (\Delta - \frac{1}{4}n^2 - \sigma^2)^{-1} \chi \|_{L^2(X) \rightarrow L^2(X)} \geq e^{-C|\text{Im}\sigma|} |\sigma|^{-1+2|\text{Im}\sigma|} / C \tag{1-3}$$

for σ in the lower half-plane and near, but bounded away from, the real axis.

The lower bound (1-3) gives a sense in which (1-1) is optimal, but finding the maximal resonance free region remains an open problem. The only known explicit example of this type is $(X, g) = \langle z \mapsto z + 1 \rangle \backslash \mathbb{H}^2$, for which Borthwick [2007, §5.3] expresses the resolvent in terms of Bessel functions and shows there is only one resonance and it is simple (see also Proposition 8.1). On the other hand, Guillopé and Zworski [1997] study more general surfaces, and prove that if the 0-volume is not zero, then there are infinitely many resonances and optimal lower and upper bounds hold on their number in disks. We apply their result to our setting in Section 2D, giving a family of surfaces with infinitely many resonances to which our Theorem applies, but it is not clear even in this case whether or not the resonance free region given by the Theorem is optimal. The delicate nature of this question is indicated by the result in [Datchev, Kang and Kessler 2015] showing that nontrapping manifolds with cusps which are merely $C^{1,1}$ (and not C^∞) do *not* have arbitrarily wide resonance free strips.

Cardoso and Vodev [2002, Corollary 1.2], extending work of Burq [1998; 2002], proved resolvent estimates for very general infinite-volume manifolds (including the ones studied here; note that the presence of a funnel implies that the volume is infinite) which imply an exponentially small resonance free region. Our Theorem gives the first large resonance free region for a family of manifolds with cusps.

For $\text{Im } \sigma = 0$, (1-1) is lossless; that is to say it agrees with the result for general nontrapping operators on asymptotically Euclidean or hyperbolic manifolds (see [Cardoso, Popov and Vodev 2004, (1.6)] and references therein). However, if (X, g) is asymptotically Euclidean or hyperbolic in the sense of [Datchev and Vasy 2012a, §4], then the gluing methods of that paper show that such a lossless estimate for $\text{Im } \sigma = 0$ implies (1-2) for some $M_1 > 0$; see [Datchev 2012]. In this sense it is due to the cusp that $\mathcal{O}(|\sigma|^{-1})$ bounds hold for $\text{Im } \sigma = 0$ but not in any strip containing the real axis.

The Theorem also provides a first step in support of the following:

Conjecture (fractal Weyl upper bound). *Let Γ be a geometrically finite discrete group of isometries of \mathbb{H}^{n+1} such that $X = \Gamma \backslash \mathbb{H}^{n+1}$ is a smooth noncompact manifold. Let $R(X)$ denote the set of eigenvalues and resonances of X included according to multiplicity, let $K \subset T^*X$ be the set of maximally extended, bounded, unit speed geodesics, and let m be the Hausdorff dimension of K . Then for any $C_0 > 0$ there is $C_1 > 0$ such that, for $r \in \mathbb{R}$,*

$$\#\{\sigma \in R(X) : |\sigma - r| \leq C_0\} \leq C_1(1 + |r|)^{(m-1)/2}.$$

This statement is a partial generalization to the case of resonances of the Weyl asymptotic for eigenvalues of a compact manifold; such results go back to work of Sjöstrand [1990]. If $\Gamma \backslash \mathbb{H}^{n+1}$ has funnels but no cusps, this is proved in [Datchev and Dyatlov 2013] (generalizing earlier results of Zworski [1999] and Guillopé, Lin and Zworski [2004]); if $X = \Gamma \backslash \mathbb{H}^2$ has cusps but no funnels, this follows from work of Selberg [1990]. When $n = 1$ the remaining case is $\Gamma \backslash \mathbb{H}^2$ having both cusps and funnels. The methods of the present paper, combined with those of [Sjöstrand and Zworski 2007; Datchev and Dyatlov 2013], provide a possible approach to the conjecture in this case. When $n \geq 2$, cusps can have mixed rank, and in this case even meromorphic continuation of the resolvent was proved only recently by Guillarmou and Mazzeo [2012].

In Section 2 we give the general assumptions on (X, g) under which the Theorem holds, and deduce consequences for the geodesic flow and for the spectrum of the Laplacian. We then give examples of manifolds which satisfy the assumptions, including examples with infinitely many resonances and examples with at least one eigenvalue.

In Section 3 we use a resolvent gluing method, based on one developed in [Datchev and Vasy 2012a], to reduce the Theorem to proving resolvent estimates and propagation of singularities results for three model operators. The first model operator is semiclassically elliptic outside of a compact set, and we analyze it in Section 4 following [Sjöstrand and Zworski 2007] and [Datchev and Vasy 2012a].

In Section 5 we study the second model operator, the model in the cusp. We use a separation of variables, a semiclassically singular rescaling, and an elliptic variant of the gluing method of Section 3 to reduce its study to that of a family of one-dimensional Schrödinger operators for which uniform resolvent estimates and propagation of singularities results hold. The rescaling causes losses for the resolvent estimate on the real axis, and we remove these by a noncompact variant of the method of propagation of singularities through trapped sets developed in [Datchev and Vasy 2012b]. The lower bound (1-3) shows that these losses cannot be removed for the continued resolvent; see also [Bony and Petkov 2013] for related and more general lower bounds in Euclidean scattering.

In Section 6 we study the third model operator, the model in the funnel, and we again reduce to a family of one-dimensional Schrödinger operators. To obtain uniform estimates we use a variant of the method of complex scaling of Aguilar and Combes [1971] and Simon [1972], following the geometric approach of Sjöstrand and Zworski [1991]. The method of complex scaling was first adapted to such families of operators by Zworski [1999], but we use here the approach of [Datchev 2010], which is slightly simpler and is adapted to nonanalytic manifolds. The analysis in this section could be replaced by that of [Vasy 2013], which avoids separating variables; the advantage of our approach is that it gives an estimate in a logarithmically large neighborhood of the real axis (although this does not make a difference here) and also requires less preliminary setup.

In Section 7 we apply (1-1) to local smoothing and resonant wave expansions. For the latter we need the additional assumption, satisfied in the example above and in many of the examples in Section 2D, that $\chi(\Delta - \frac{1}{4}n^2 - \sigma^2)^{-1}\chi$ is meromorphic in \mathbb{C} . In Section 8 we prove (1-3) using Bessel function asymptotics.

2. Preliminaries

Throughout the paper $C > 0$ is a large constant which may change from line to line, and estimates are always uniform for $h \in (0, h_0]$, where $h_0 > 0$ may change from line to line.

2A. Assumptions. Let S be a compact manifold (without boundary) of dimension n , and let

$$X := \mathbb{R}_r \times S.$$

Let $R_g > 0$, and let g be a Riemannian metric on X such that

$$g|_{\{\pm r > R_g\}} = dr^2 + e^{2(r+\beta(r))} dS_{\pm}, \tag{2-1}$$

where dS_+ and dS_- are metrics on S , $R_g > 0$ and $\beta \in C^\infty(\mathbb{R})$. We call the region $\{r < -R_g\}$ the *cusp*, and the region $\{r > R_g\}$ the *funnel*; see [Figure 2](#).

Suppose there is $\theta_0 \in (0, \frac{\pi}{4})$ such that β is holomorphic and bounded in the sectors where $|z| > R_g$ and $\min\{|\arg z|, |\arg(-z)|\} < 2\theta_0$. By Cauchy estimates, for all $k \in \mathbb{N}$ there are $C, C_k > 0$, such that if $|z| > R_g$ and $\min\{|\arg z|, |\arg(-z)|\} \leq \theta_0$, then

$$|\beta^{(k)}(z)| \leq C_k |z|^{-k}, \quad |\operatorname{Im} \beta(z)| \leq C |\operatorname{Im} z|/|z|.$$

In particular, after possibly redefining R_g to be larger, we may assume without loss of generality that, for all $r \in \mathbb{R}$,

$$|\beta'(r)| + |\beta''(r)| \leq \frac{1}{4}. \tag{2-2}$$

In the example at the beginning of the paper $\beta \equiv 0$. When the funnel end is an exact hyperbolic funnel, $\beta(r) = C + \log(1 + e^{-2r})$ for $r > R_g$.

We make two dynamical assumptions: if $\gamma : \mathbb{R} \rightarrow X$ is a maximally extended geodesic, assume $\gamma(\mathbb{R})$ is not bounded and $\gamma^{-1}(\{r < -R_g\})$ is connected. See [Section 2D](#) for examples.




Figure 2. The manifold X .

2B. Dynamics near infinity. Let $p + 1 \in C^\infty(T^*X)$ be the geodesic Hamiltonian; that is,

$$p = \rho^2 + e^{-2(r+\beta(r))} \sigma_\pm - 1$$

in the region $\{\pm r > R_g\}$, where ρ is dual to r , and σ_\pm is the geodesic Hamiltonian of (S, dS_\pm) . From this we conclude that, along geodesic flow lines, we have

$$\dot{r}(t) = H_p \rho = 2\rho(t), \quad \dot{\rho}(t) = -H_p r = 2[1 + \beta'(r(t))]e^{-2(r+\beta(r(t)))} \sigma_\pm,$$

so long as the trajectory remains within $\{\pm r > R_g\}$. In particular,

$$\ddot{r}(t) = 4[1 + \beta'(r(t))]e^{-2(r+\beta(r(t)))} \sigma_\pm \geq 0. \tag{2-3}$$

Dividing the equation for $\dot{\rho}$ by $p + 1 - \rho^2$, putting $\hat{\rho} = \rho/\sqrt{p + 1}$, and integrating we find

$$\tanh^{-1} \hat{\rho}(t) - \tanh^{-1} \hat{\rho}(0) = 2\sqrt{p + 1} \left(t + \int_0^t \beta'(r(s)) ds \right) \geq \frac{3}{4} \frac{r(t) - r(0)}{\max\{\hat{\rho}(s) : s \in [0, t]\}}, \tag{2-4}$$

where the equality holds so long as the trajectory remains in $\{\pm r > R_g\}$, and the inequality (which follows from (2-2) and the equation for \dot{r}) holds when additionally $t \geq 0, \rho(0) \geq 0$.

2C. The essential spectrum and semiclassical formulation of the problem. The nonnegative Laplacian is given by

$$\Delta|_{\{\pm r > R_g\}} = D_r^2 - in(1 + \beta'(r))D_r + e^{-2(r+\beta(r))} \Delta_{S_\pm},$$

where $D_r = -i\partial_r$, and Δ_{S_\pm} is the Laplacian on (S, dS_\pm) . Fix $\varphi \in C^\infty(X)$ such that

$$\varphi|_{\{|r| > R_g\}} = \frac{1}{2}n(r + \beta(r)). \tag{2-5}$$

Then

$$(e^\varphi \Delta e^{-\varphi})|_{\{\pm r > R_g\}} = D_r^2 + e^{-2(r+\beta(r))} \Delta_{S_\pm} + \frac{1}{4}n^2 + V(r), \tag{2-6}$$

where

$$V(r) = \varphi'' + \varphi'^2 - \frac{1}{4}n^2 = \frac{1}{2}n\beta'' + \frac{1}{2}n^2\beta' + \frac{1}{4}n^2\beta'^2.$$

This shows that the essential spectrum of Δ is $[\frac{1}{4}n^2, \infty)$ (see for example [Reed and Simon 1978, Theorem XIII.14, Corollary 3]); the potential perturbation V is relatively compact since β' and β'' tend to zero at infinity (see for example Rellich's criterion [ibid., Theorem XIII.65]).

In this paper we study

$$P := h^2(e^\varphi \Delta e^{-\varphi} - \frac{1}{4}n^2) - 1. \tag{2-7}$$

This is an unbounded selfadjoint operator on $L_\varphi^2(X) := \{e^\varphi u : u \in L^2(X)\}$ with domain

$$H_\varphi^2(X) := \{u \in L_\varphi^2(X) : e^\varphi \Delta e^{-\varphi} u \in L_\varphi^2(X)\} = \{e^\varphi u : u \in H^2(X)\}.$$

Over the course of Sections 3–6 we will prove the following:

Proposition 2.1. *For every $\chi \in C_0^\infty(X)$, $E \in (0, 1)$ there exists $C_0 > 0$ such that for every $\Gamma > 0$ there exist $C, h_0 > 0$ such that the cutoff resolvent $\chi(P - \lambda)^{-1}\chi$ continues holomorphically from $\{\text{Im } \lambda > 0\}$ to $[-E, E] - i[0, \Gamma h]$ and satisfies*

$$\|\chi(P - \lambda)^{-1}\chi\|_{L^2_\varphi(X) \rightarrow L^2_\varphi(X)} \leq Ch^{-1-C_0|\text{Im } \lambda|/h} \tag{2-8}$$

uniformly for $\lambda \in [-E, E] - i[0, \Gamma h]$ and $h \in (0, h_0]$.

This implies the [Theorem](#).

2D. Examples. In this section we give a family of examples of manifolds satisfying the assumptions of [Section 2A](#). I am very grateful to John Lott for suggesting this family of examples. In this section $d_g(p, q)$ denotes the distance between p and q with respect to the Riemannian metric g , and $L_g(c)$ denotes the length of a curve c with respect to g .

Let (\mathbb{H}^{n+1}, g_h) be hyperbolic space with coordinates

$$(r, y) \in \mathbb{R} \times \mathbb{R}^n, \quad g_h := dr^2 + e^{2r} dy^2.$$

Let (X, g_h) be a parabolic cylinder obtained by quotienting the y variables to a torus:

$$X := \mathbb{R} \times ((y \mapsto y + c_1, \dots, y \mapsto y + c_n) \backslash \mathbb{R}^n),$$

where the c_j are linearly independent vectors in \mathbb{R}^n . Let $R_g > 0$, put $dS_+ = dS_- = dy^2$, and take $\beta \in C^\infty(\mathbb{R})$ satisfying all assumptions of [Section 2A](#), including (2-2). On $\{|r| > R_g\}$ define g by (2-1), and on $\{|r| \leq R_g\}$ let g be any metric with all sectional curvatures nonpositive. The calculation in the [Appendix](#) shows that the sectional curvatures in $\{|r| > R_g\}$ are nonpositive so long as (2-2) holds.

The two dynamical assumptions in the last paragraph of [Section 2A](#) will follow from the following classical theorem (see for example [\[Bridson and Haefliger 1999, Theorem III.H.1.7\]](#)).

Proposition 2.2 (stability of quasigeodesics). *Let (\mathbb{H}^{n+1}, g_h) be the $(n+1)$ -dimensional hyperbolic space, let $p, q \in \mathbb{H}^{n+1}$, and let $\gamma_h : [t_1, t_2] \rightarrow \mathbb{H}^{n+1}$ be the unit-speed geodesic from p to q . Suppose $c : [t_1, t_2] \rightarrow \mathbb{H}^{n+1}$ satisfies $c(t_1) = p, c(t_2) = q$, and there is $C_1 > 0$ such that*

$$\frac{1}{C_1}|t - t'| \leq d_{g_h}(c(t), c(t')) \leq C_1|t - t'| \tag{2-9}$$

for all $t, t' \in [t_1, t_2]$. Then

$$\max_{t \in [t_1, t_2]} d_{g_h}(\gamma_h(t), c(t)) \leq C_2, \tag{2-10}$$

where C_2 depends only on C_1 .

To apply this theorem, observe first that just as g_h descends to a metric on X , so g lifts to a metric on \mathbb{H}^{n+1} ; call the lifted metric g as well. Observe there is C_g such that

$$\frac{1}{C_g}g_h(u, u) \leq g(u, u) \leq C_g g_h(u, u), \quad u \in T_x X, \quad x \in X. \tag{2-11}$$

Indeed, for x varying in a compact set this is true for any pair of metrics, and on $\{|r| > R_g\}$ it suffices if $C_g \geq e^{2 \max |\beta|}$. We will show that if c is a unit-speed g -geodesic in \mathbb{H}^n , then (2-9) holds with a constant C_1 depending only on C_g . Since both g and g_h have nonnegative curvature and hence distance-minimizing geodesics, it is equivalent to show that

$$\frac{1}{C_1} d_g(p, q) \leq d_{g_h}(p, q) \leq C_1 d_g(p, q) \tag{2-12}$$

holds for all $p, q \in \mathbb{H}^{n+1}$, with a constant C_1 which depends only on C_g . For this last we compute as follows: let γ be a unit-speed g -geodesic from p to q . Then

$$d_{g_h}(p, q) \leq L_{g_h}(\gamma) = \int_{t_1}^{t_2} \sqrt{g_h(\dot{\gamma}, \dot{\gamma})} dt \leq \int_{t_1}^{t_2} \sqrt{C_g g(\dot{\gamma}, \dot{\gamma})} dt = \sqrt{C_g} L_g(\gamma) = \sqrt{C_g} d_g(p, q).$$

This proves the second inequality of (2-12), and the first follows from the same calculation since (2-11) is unchanged if we switch g and g_h .

Let $\gamma : \mathbb{R} \rightarrow X$ be a g -geodesic and $\gamma_h : \mathbb{R} \rightarrow X$ a g_h -geodesic. For any $x \in X$ we have

$$\lim_{t \rightarrow \infty} d_{g_h}(\gamma_h(t), x) = \lim_{t \rightarrow \infty} d_g(\gamma(t), x) = \infty,$$

and by (2-10) the same holds if γ_h is replaced by γ . In particular $\gamma(\mathbb{R})$ is not bounded.

We check finally that $\gamma^{-1}(\{r < -R_g\})$ is connected. It suffices to check that if instead $\gamma : \mathbb{R} \rightarrow \mathbb{H}^{n+1}$ is a g -geodesic, then $\gamma^{-1}(\{r < -N\})$ is connected for N large enough, with N independent of γ . We then conclude by redefining R_g to be larger than N .

We argue by way of contradiction. From (2-3) we see that $\dot{r}(t)$ is nondecreasing along γ in $\{r < -R_g\}$. Hence, if $\gamma^{-1}(\{r < -N\})$ is to contain at least two intervals for some $N > R_g$, there must exist times $t_1 < t_2 < t_3$ such that $r(\gamma(t_1)), r(\gamma(t_3)) < -N$ and $r(\gamma(t_2)) = -R_g$. Now the g_h -geodesic $\gamma_h : [t_1, t_3] \rightarrow \mathbb{H}^n$ joining $\gamma(t_1)$ to $\gamma(t_3)$ has $r(\gamma_h(t)) < -N$ for all $t \in [t_1, t_3]$. It follows that $d_{g_h}(\gamma_h(t_2), \gamma(t_2)) \geq N - R_g$, and if N is large enough this violates (2-10).

2D1. Examples with infinitely many resonances. In this subsection we specialize to the case $n = 1$, $\beta(r) = 0$ for $r < -R_g$, $\beta(r) = \beta_0 + \log(1 + e^{-2r})$ for $r > R_g$ and for some $\beta_0 \in \mathbb{R}$. Then the cusp and funnel of X are isometric to the standard cusp and funnel obtained by quotienting \mathbb{H}^2 by a *nonelementary* Fuchsian subgroup (see, e.g., [Borthwick 2007, §2.4]; note that the funnel end is slightly different here than in the example at the beginning of the paper).

In particular there is $l > 0$ such that

$$X = \mathbb{R}_r \times (\mathbb{R}/l\mathbb{Z})_t, \quad g|_{\{r > R_g\}} = dr^2 + \cosh^2 r dt^2.$$

If $(X_0, g_0) = [0, \infty) \times (\mathbb{R}/l\mathbb{Z})$, $g_0 = dr^2 + \cosh^2 r dt^2$, then the 0-volume of X is

$$0\text{-vol}(X) \stackrel{\text{def}}{=} \text{vol}_g(X \cap \{r < R_g\}) - \text{vol}_{g_0}(X_0 \cap \{r < R_g\}).$$

Let $R_\chi(\sigma)$ denote the meromorphic continuation of $\chi(\Delta - \frac{1}{4} - \sigma^2)^{-1} \chi$. In this case, $R_\chi(\sigma)$ is meromorphic in \mathbb{C} [Mazzeo and Melrose 1987; Guillopé and Zworski 1997], and near each pole σ_0 we

have

$$R_\chi(\sigma) = \chi \left(\sum_{j=1}^k \frac{A_j}{(\sigma - \sigma_0)^j} + A(\sigma) \right) \chi,$$

where the $A_j : L^2_{\text{comp}}(X) \rightarrow L^2_{\text{loc}}(X)$ are finite rank and $A(\sigma)$ is holomorphic near σ_0 . The multiplicity of a pole, $m(\sigma_0)$ is given by

$$m(\sigma) \stackrel{\text{def}}{=} \text{rank} \left(\sum_{j=1}^k A_j \right).$$

Proposition 2.3 [Guillopé and Zworski 1997, Theorem 1.3]. *If $0\text{-vol}(X) \neq 0$, then there exists a constant C such that*

$$\lambda^2/C \leq \sum_{|\sigma| \leq \lambda} m(\sigma) \leq C\lambda^2, \quad \lambda > C.$$

We can ensure that $0\text{-vol}(X) \neq 0$ by adding, if necessary, a small compactly supported metric perturbation to g . Then, as $\lambda \rightarrow \infty$, the meromorphic continuation of R_χ will have $\sim \lambda^2$ -many poles in a disk of radius λ , but none of them will be in the strips (1-1).

2D2. Examples with at least one eigenvalue. In this subsection we consider examples of the form

$$X := \mathbb{R} \times (\mathbb{R}^n / \mathbb{Z}^n), \quad g := dr^2 + \exp\left(2r + 2 \int_{-\infty}^r b\right) dy^2, \quad b \in C_0^\infty(\mathbb{R}). \quad (2-13)$$

As in (2-3), we have $\ddot{r} = 4(1 + b(r))e^{-2(r + \int^r b)}\sigma$, and this is nonnegative as long as $b \geq -1$; consequently, as long as $b \geq -1$ the assumptions of Section 2A hold. We will give a sufficient condition on b such that X has at least one eigenvalue, and also infinitely many resonances.

By the calculation in Section 2C, if $\varphi(r) := \frac{1}{2}(r + \int_{-\infty}^r b)$ for all $r \in \mathbb{R}$, then

$$e^{-\varphi} \Delta e^\varphi = D_r^2 + e^{-2(r + \int^r b)} \Delta_{\mathbb{R}^n / \mathbb{Z}^n} + \frac{1}{4}n^2 + V(r), \quad V(r) := \frac{1}{2}nb'(r) + \frac{1}{4}n^2b(r)^2 + \frac{1}{2}n^2b(r).$$

Note $V \in C_0^\infty(\mathbb{R})$, and consequently (see for example [Reed and Simon 1978, Theorem XIII.110]) $D_r^2 + V(r)$ has a negative eigenvalue provided $V \not\equiv 0$ and $\int V \leq 0$; it suffices for example to take $b \leq 0$. But Zworski [1987, Theorem 2] has shown that if $V \not\equiv 0$, then $D_r^2 + V(r)$ has infinitely many resonances: indeed, the number in a disk of radius λ is given by

$$\frac{2}{\pi}(\text{diam supp } V)\lambda + o(\lambda), \quad \lambda \rightarrow \infty.$$

This eigenvalue and these resonances correspond to an eigenvalue and resonances for Δ : one multiplies the eigenfunction and resonant states by e^φ and regards them as functions on X which depend on r only.

In summary, if (X, g) is given by (2-13), then the assumptions of Section 2A hold if $b \geq -1$. It has infinitely many resonances and at least one eigenvalue if additionally $b \not\equiv 0, b \leq 0$.

2E. Pseudodifferential operators. In this section we review some facts about semiclassical pseudodifferential operators, following [Dimassi and Sjöstrand 1999; Zworski 2012; Dyatlov and Zworski 2016].

2E1. *Pseudodifferential operators on \mathbb{R}^n .* For $m \in \mathbb{R}$, $\delta \in [0, \frac{1}{2})$, let $S_\delta^m(\mathbb{R}^n)$ be the symbol class of functions $a = a_h(x, \xi) \in C^\infty(T^*\mathbb{R}^n)$ satisfying

$$|\partial_x^\alpha \partial_\xi^\beta a| \leq C_{\alpha,\beta} h^{-\delta(|\alpha|+|\beta|)} (1 + |\xi|^2)^{(m-|\beta|)/2} \tag{2-14}$$

uniformly in $T^*\mathbb{R}^n$. The *principal symbol* of a is its equivalence class in $S_\delta^m(\mathbb{R}^n)/hS_\delta^{m-1}(\mathbb{R}^n)$. Let $S^m(\mathbb{R}^n) = S_0^m(\mathbb{R}^n)$.

We quantize $a \in S_\delta^m(\mathbb{R}^n)$ to an operator $\text{Op}(a)$ using the formula

$$(\text{Op}(a)u)(x) = \frac{1}{(2\pi h)^n} \iint e^{i(x-y)\cdot\xi/h} a_h(x, \xi) u(y) dy d\xi, \tag{2-15}$$

and put $\Psi_\delta^m(\mathbb{R}^n) = \{\text{Op}(a) : a \in S_\delta^m(\mathbb{R}^n)\}$, $\Psi^m(\mathbb{R}^n) = \Psi_0^m(\mathbb{R}^n)$. If $A = \text{Op}(a)$ then a is the *full symbol* of A , and the principal symbol of A is the principal symbol of a . If $A \in \Psi_\delta^m(\mathbb{R}^n)$, then for any $s \in \mathbb{R}$ we have $\|A\|_{H_h^{s+m}(\mathbb{R}^n) \rightarrow H_h^s(\mathbb{R}^n)} \leq C$, where (if $\Delta \geq 0$)

$$\|u\|_{H_h^s(\mathbb{R}^n)} = \|(1 + h^2 \Delta)^{s/2} u\|_{L^2(\mathbb{R}^n)}.$$

If $A \in \Psi_\delta^m(\mathbb{R}^n)$ and $B \in \Psi_\delta^{m'}(\mathbb{R}^n)$, then $AB \in \Psi_\delta^{m+m'}(\mathbb{R}^n)$ and $[A, B] = AB - BA \in h^{1-2\delta} \Psi_\delta^{m+m'-1}(\mathbb{R}^n)$. If a and b are the principal symbols of A and B , then the principal symbol of $h^{2\delta-1}[A, B]$ is $iH_b a$, where H_b is the Hamiltonian vector field of b .

If $K \subset T^*\mathbb{R}^n$ has either K or $T^*\mathbb{R}^n \setminus K$ bounded in ξ , then $a \in S_\delta^m(\mathbb{R}^n)$ is *elliptic* on K if

$$|a| \geq (1 + |\xi|^2)^{m/2} / C \tag{2-16}$$

uniformly for $(x, \xi) \in K$. We say that $A \in \Psi_\delta^m(\mathbb{R}^n)$ is elliptic on K if its principal symbol is. For such K , we say A is *microsupported* in K if the full symbol a of A obeys

$$|\partial_x^\alpha \partial_\xi^\beta a| \leq C_{\alpha,\beta,N} h^N (1 + |\xi|^2)^{-N} \tag{2-17}$$

uniformly on $T^*\mathbb{R}^n \setminus K$, for any α, β, N . If A_1 is microsupported in K_1 and A_2 is microsupported in K_2 , then $A_1 A_2$ is microsupported in $K_1 \cap K_2$.

If $A \in \Psi_\delta^m(\mathbb{R}^n)$ is elliptic on K , then it is invertible there in the following sense: there exists $G \in \Psi_\delta^{-m}(\mathbb{R}^n)$ such that $AG - \text{Id}$ and $GA - \text{Id}$ are both microsupported in $T^*\mathbb{R}^n \setminus K$. Hence if $B \in \Psi_\delta^{m'}(\mathbb{R}^n)$ is microsupported in K and A is elliptic in an ε -neighborhood of K for some $\varepsilon > 0$, then, for any $s, N \in \mathbb{R}$,

$$\|Bu\|_{H_h^{s+m}(\mathbb{R}^n)} \leq C \|ABu\|_{H_h^s(\mathbb{R}^n)} + \mathcal{O}(h^\infty) \|u\|_{H_h^{-N}(\mathbb{R}^n)}. \tag{2-18}$$

The *sharp Gårding inequality* says that if the principal symbol of $A \in \Psi_\delta^m(\mathbb{R}^n)$ is nonnegative near K and $B \in \Psi_\delta^{m'}(\mathbb{R}^n)$ is microsupported in K , then

$$\langle ABu, Bu \rangle_{L^2(\mathbb{R}^n)} \geq -Ch^{1-2\delta} \|Bu\|_{H^{(m-1)/2}(\mathbb{R}^n)}^2 - \mathcal{O}(h^\infty) \|u\|_{H_h^{-N}(\mathbb{R}^n)}. \tag{2-19}$$

2E2. Pseudodifferential operators on a manifold. These results extend to the case of a noncompact manifold X , provided we require our estimates to be uniform only on compact subsets of X . For convenience we work in the setting of Section 2A, with the notation of Section 2C, but the discussion below applies to any manifold; see also the discussions in [Datchev and Dyatlov 2013, §3.1] and [Dyatlov and Zworski 2016, Appendix E]. Note that we take care to quantize a symbol which is compactly supported in space to an operator which is compactly supported in space.

Write $S_\delta^m(X)$ for the symbol class of functions $a \in C^\infty(T^*X)$ satisfying (2-14) on coordinate patches (note that this condition is invariant under change of coordinates). The principal symbol of a is its equivalence class in $S_\delta^m(X)/hS_\delta^{m-1}(X)$, and let $S^m(X) = S_0^m(X)$.

Let $h^\infty\Psi^{-\infty}(X)$ be the set of linear operators R such that for any $\chi \in C_0^\infty(X)$, we have

$$\|\chi R\|_{H_{\varphi,h}^{-N}(X) \rightarrow H_{\varphi,h}^N(X)} + \|R\chi\|_{H_{\varphi,h}^{-N}(X) \rightarrow H_{\varphi,h}^N(X)} \leq C_N h^N$$

for any N , where

$$\|u\|_{H_{\varphi,h}^s(X)} := \|(2 + P)^{s/2} u\|_{L_\varphi^2(X)}. \tag{2-20}$$

We quantize $a \in S_\delta^m(X)$ to an operator $\text{Op}(a)$ by using a partition of unity and the formula (2-15) in coordinate patches. Let $\Psi_\delta^m(X) = \{\text{Op}(a) + R : a \in S_\delta^m(X), R \in h^\infty\Psi^{-\infty}(X)\}$. The quantization Op depends on the choices of coordinates and partition of unity, but the class $\Psi_\delta^m(X)$ does not. If $A \in \Psi_\delta^m(X)$ and $\chi \in C_0^\infty(X)$, then χA and $A\chi$ are bounded as operators $H_{\varphi,h}^{s+m}(X) \rightarrow H_{\varphi,h}^s(X)$, uniformly in h . If $A \in \Psi_\delta^m(X)$ and $B \in \Psi_\delta^{m'}(X)$, then

$$AB \in \Psi_\delta^{m+m'}(X) \quad \text{and} \quad h^{2\delta-1}[A, B] \in \Psi_\delta^{m+m'-1}(X).$$

If a and b are the principal symbols of A and B (the principal symbol is invariantly defined, although the total symbol is not), then the principal symbol of $h^{2\delta-1}[A, B]$ is $iH_b a$, where H_b is the Hamiltonian vector field of b .

Let $K \subset T^*X$ have either $K \cap T^*U$ bounded for every bounded $U \subset X$, or $T^*U \setminus K$ bounded for every bounded $U \subset X$. We say $a \in S_\delta^m(X)$ is *elliptic* on K if (2-16) holds uniformly on $T^*U \cap K$ for every bounded $U \subset X$. We say that $A \in \Psi_\delta^m(X)$ is elliptic on K if its principal symbol is. We say A is *microsupported* in K if a full symbol a of A obeys (2-17) uniformly on $T^*U \setminus K$ for every bounded $U \subset X$ and for any α, β, N (note that if this holds for one full symbol of A , it also does for all the others).

If $B \in \Psi_\delta^{m'}(X)$ is microsupported in K and A is elliptic in an ε -neighborhood of K for some $\varepsilon > 0$, then, for any $s, N \in \mathbb{R}$ and $\chi \in C_0^\infty(X)$,

$$\|B\chi u\|_{H_{\varphi,h}^{s+m}(X)} \leq C \|AB\chi u\|_{H_{\varphi,h}^s(X)} + \mathcal{O}(h^\infty) \|\chi u\|_{H_{\varphi,h}^{-N}(X)}. \tag{2-21}$$

The *sharp Gårding inequality* says that if the principal symbol of $A \in \Psi_\delta^m(X)$ is nonnegative near K and $B \in \Psi_\delta^{m'}(X)$ is microsupported in K , then for every $\chi \in C_0^\infty(X)$, $N \in \mathbb{R}$,

$$\langle AB\chi u, B\chi u \rangle_{L_\varphi^2(X)} \geq -Ch^{1-2\delta} \|B\chi u\|_{H_{\varphi,h}^{(m-1)/2}(X)}^2 - \mathcal{O}(h^\infty) \|\chi u\|_{H_{\varphi,h}^{-N}(X)}. \tag{2-22}$$

2E3. Exponentiation of operators. For $q \in C_0^\infty(T^*X)$, $Q = \text{Op}(q)$, and $\varepsilon \in [0, C_0 h \log(1/h)]$, we will be interested in operators of the form $e^{\varepsilon Q/h}$. By the discussion above, since $q \in S^m(X)$ for every $m \in \mathbb{R}$, we have $\|Q\|_{H_{\varphi,h}^{-N} \rightarrow H_{\varphi,h}^N} \leq C_N$ for every $N \in \mathbb{R}$.

We write

$$e^{\varepsilon Q/h} := \sum_{j=0}^{\infty} \frac{(\varepsilon/h)^j}{j!} Q^j,$$

with the sum converging in the $H_{\varphi,h}^s(X) \rightarrow H_{\varphi,h}^s(X)$ norm operator topology, but the convergence is not uniform as $h \rightarrow 0$. Beals’s characterization [Zworski 2012, Theorem 9.12] can be used to show that $e^{\varepsilon Q/h} \in \Psi_\delta^0(X)$ for any $\delta > 0$, but we will not need this. Let $s \in \mathbb{R}$. Then

$$\|e^{\varepsilon Q/h}\| \leq \sum_{j=0}^{\infty} \frac{(C_0 \log(1/h))^j}{j!} \|Q\|^j = e^{C_0 \log(1/h) \|Q\|} = h^{-C_0 \|Q\|}, \tag{2-23}$$

where all norms are $H_{\varphi,h}^s(X) \rightarrow H_{\varphi,h}^s(X)$.

If $A \in \Psi_\delta^m(X)$ is bounded as an operator $H_{\varphi,h}^{s+m}(X) \rightarrow H_{\varphi,h}^s(X)$, uniformly in h , (without needing to be multiplied by a cutoff), then, by (2-23),

$$\|e^{\varepsilon Q/h} A e^{-\varepsilon Q/h}\|_{H_{\varphi,h}^{s+m}(X) \rightarrow H_{\varphi,h}^s(X)} \leq C h^{-N} \tag{2-24}$$

for any $s \in \mathbb{R}$, where

$$N = C_0(\|Q\|_{H_{\varphi,h}^{s+m}(X) \rightarrow H_{\varphi,h}^{s+m}(X)} + \|Q\|_{H_{\varphi,h}^s(X) \rightarrow H_{\varphi,h}^s(X)}).$$

But, writing $\text{ad}_Q A = [Q, A]$ and $e^{\varepsilon Q/h} A e^{-\varepsilon Q/h} = e^{\varepsilon \text{ad}_Q/h} A$, for any $J \in \mathbb{N}$ we have the Taylor expansion

$$e^{\varepsilon Q/h} A e^{-\varepsilon Q/h} = \sum_{j=0}^J \frac{\varepsilon^j}{j!} \left(\frac{\text{ad}_Q}{h}\right)^j A + \frac{\varepsilon^{J+1}}{J!} \int_0^1 (1-t)^J e^{-\varepsilon t \text{ad}_Q/h} \left(\frac{\text{ad}_Q}{h}\right)^{J+1} A dt. \tag{2-25}$$

For any $M \in \mathbb{N}$, the integrand maps $H_{\varphi,h}^M(X)$ to $H_{\varphi,h}^{-M}(X)$ with norm $\mathcal{O}(h^{-2\delta(J+1)-N})$, where

$$N = C_0(\|Q\|_{H_{\varphi,h}^M(X) \rightarrow H_{\varphi,h}^M(X)} + \|Q\|_{H_{\varphi,h}^{-M}(X) \rightarrow H_{\varphi,h}^{-M}(X)}).$$

Hence applying (2-25) with J sufficiently large we see that (2-24) can be improved to

$$\|e^{\varepsilon Q/h} A e^{-\varepsilon Q/h}\|_{H_{\varphi,h}^{s+m}(X) \rightarrow H_{\varphi,h}^s(X)} \leq C,$$

and the integrand in (2-25) maps $H_{\varphi,h}^M(X)$ to $H_{\varphi,h}^{-M}(X)$ with norm $\mathcal{O}(1)$. Applying (2-25) with $J \rightarrow \infty$ shows that $e^{\varepsilon Q/h} A e^{-\varepsilon Q/h} \in \Psi_\delta^m(X)$, and applying (2-25) with $J = 1$ we find

$$e^{\varepsilon Q/h} A e^{-\varepsilon Q/h} = A - \varepsilon[A, Q/h] + \varepsilon^2 h^{-4\delta} R, \tag{2-26}$$

where $R \in \Psi_\delta^{-\infty}(X)$.

3. Reduction to estimates for model operators

3A. Resolvent gluing. In Section 2 we showed that the Theorem follows from (2-8). In this section, we reduce (2-8) to several estimates for model operators using a variant of the gluing method of [Datchev and Vasy 2012a], adapted to the dynamics on X .

We will use the following open cover of X :

$$\Omega_C := \{r < -R_g\}, \quad \Omega_K := \{|r| < R_g + 3\}, \quad \Omega_F := \{r > R_g\}.$$

Let P_C, P_K, P_F be differential operators on X which are *model operators* for P , with respect to this open cover, in the sense that they satisfy

$$P_j|_{\Omega_j} = P|_{\Omega_j}, \quad j \in \{C, K, F\}. \tag{3-1}$$

So P_C is a model in the cusp, P_F is a model in the funnel, and P_K is a model in a neighborhood of the remaining region (see Figure 2).

More specifically, let $W_K \in C^\infty(X; [0, 1])$ be 0 near $\{|r| \leq R_g + 3\}$, and 1 near $\{|r| \geq R_g + 4\}$, and let

$$P_K = P - iW_K;$$

let $W_C \in C^\infty(\mathbb{R}; [0, 1])$ be 0 near $\{r \leq -R_g\}$, and 1 near $\{r \geq 0\}$, and let

$$P_C = h^2 D_r^2 + h^2 e^{-2(r+\beta(r))} \Delta_{S_-} + h^2 V(r) - 1 - iW_C(r);$$

let $W_F \in C^\infty(\mathbb{R}; [0, 1])$ be 0 near $\{r \geq R_g\}$, and 1 near $\{r \leq 0\}$, nonincreasing, and let

$$P_F = h^2 D_r^2 + h^2 (1 - W_F(r)) e^{-2(r+\beta(r))} \Delta_{S_+} + h^2 V(r) - 1 - iW_F(r).$$

The functions W_j for $j \in \{C, K, F\}$, are called *complex absorbing barriers* and they make each P_j semiclassically elliptic in the region where $W_j = 1$. Note that we have also chosen P_C and P_F so that we can separate variables, and so that P_F has no exponentially growing term.

Now observe that $P_j + iW_j$ is selfadjoint on L_j^2 , where

$$L_K^2 := L_\phi^2(X), \quad L_C^2 := L^2(X, dr dS_-), \quad L_F^2 := L^2(X, dr dS_+).$$

Moreover, $W_j \geq 0$ implies $\langle \text{Im } P_j u, u \rangle_{L_j^2} \leq 0$, and hence

$$\|u\|_{L_j^2} \leq (\text{Im } \lambda)^{-1} \|(P_j - \lambda)u\|_{L_j^2}, \quad \text{Im } \lambda > 0,$$

and, consequently (since W_j is bounded on L_j^2), when $\text{Im } \lambda > 0$, we can define the resolvents

$$R_j(\lambda) := (P_j - \lambda)^{-1} : L_j^2 \rightarrow L_j^2, \quad j \in \{C, K, F\}.$$

Using (2-20) and (3-1) gives, for any $\chi_j \in C^\infty(X)$, bounded with all derivatives, and satisfying $\text{supp } \chi_j \subset \Omega_j$,

$$\max_{j \in \{C, K, F\}} \|\chi_j R_j(\lambda) \chi_j\|_{L_\phi^2(X) \rightarrow H_{\phi, h}^2(X)} \leq C(|\lambda| + (\text{Im } \lambda)^{-1}), \quad \text{Im } \lambda > 0. \tag{3-2}$$

Below we will show that for every $\chi_j \in C_0^\infty(X)$ with $\text{supp } \chi_j \subset \Omega_j$, $E \in (0, 1)$, there is $C_0 > 0$ such that for all $\Gamma > 0$ the cutoff resolvents $\chi_j R_j(\lambda)\chi_j$ continue holomorphically to $\lambda \in [-E, E] + i[-\Gamma h, \infty)$, where they satisfy

$$\max_{j \in \{C, K, F\}} \|\chi_j R_j(\lambda)\chi_j\|_{L_\varphi^2(X) \rightarrow H_{\varphi, h}^2(X)} \leq C h^{-1 - C_0 |\text{Im } \lambda| / 5h}. \tag{3-3}$$

Here E , C_0 , and Γ are the same as in (2-8), but as elsewhere in the paper the constant C and the implicit constant h_0 may be different.

We will also show that the $R_j(\lambda)$ propagate singularities forward along bicharacteristics, in the following limited sense. Let $\chi_1 \in C_0^\infty(X)$ and let $\chi_2, \chi_3 \in \Psi^1(X)$ be compactly supported differential operators.

- Suppose $\text{supp } \chi_1 \subset \Omega_K$, $\text{supp } \chi_2 \subset \Omega_K \cap \Omega_F$, and $\text{supp } \chi_3 \subset \Omega_F$. If further $\text{supp } \chi_1 \cup \text{supp } \chi_3 \subset \{r < R_g + 2\}$ and $\text{supp } \chi_2 \subset \{r > R_g + 2\}$, then, for any $N \in \mathbb{N}$,

$$\|\chi_3 R_F(\lambda)\chi_2 R_K(\lambda)\chi_1\|_{L_\varphi^2(X) \rightarrow L_\varphi^2(X)} = \mathcal{O}(h^\infty) \tag{3-4}$$

uniformly for $|\text{Re } \lambda| \leq E$, $\text{Im } \lambda \in [-\Gamma h, h^{-N}]$.

- Suppose $\text{supp } \chi_1 \subset \Omega_C$, $\text{supp } \chi_2 \subset \Omega_C \cap \Omega_K$, and $\text{supp } \chi_3 \subset \Omega_K$. If further $\text{supp } \chi_1 \cup \text{supp } \chi_3 \subset \{r < -R_g - 2\}$ and $\text{supp } \chi_2 \subset \{r > -R_g - 2\}$, then, for any $N \in \mathbb{N}$,

$$\|\chi_3 R_K(\lambda)\chi_2 R_C(\lambda)\chi_1\|_{L_\varphi^2(X) \rightarrow L_\varphi^2(X)} = \mathcal{O}(h^\infty) \tag{3-5}$$

uniformly for $|\text{Re } \lambda| \leq E$, $\text{Im } \lambda \in [-\Gamma h, h^{-N}]$.

Note that in either case there can exist no bicharacteristic passing through $T^* \text{supp } \chi_1$, $T^* \text{supp } \chi_2$, $T^* \text{supp } \chi_3$ in that order. In the first case this is implied by (2-3), and in the second by (2-3) together with the assumption that $\gamma^{-1}(\{r < -R_g\})$ is connected for any geodesic $\gamma : \mathbb{R} \rightarrow X$. We will use these facts in the proofs of (3-4) and (3-5) below. Before doing that, however, we will show that these estimates imply the [Theorem](#).

Proposition 3.1. *The estimate (2-8) follows from (3-3), (3-4), and (3-5).*

Proof. Let $\chi_C, \chi_K, \chi_F \in C^\infty(\mathbb{R})$ satisfy $\chi_C + \chi_K + \chi_F = 1$, $\text{supp } \chi_F \subset (R_g + 1, \infty)$, $\text{supp}(1 - \chi_F) \subset (R_g + 2, \infty)$, and $\chi_C(r) = \chi_F(-r)$ for all $r \in \mathbb{R}$. Then define a parametrix for $P - \lambda$ by

$$G = \chi_C(r - 1)R_C(\lambda)\chi_C(r) + \chi_K(|r - 1|)R_K(\lambda)\chi_K(|r|) + \chi_F(r + 1)R_F(\lambda)\chi_F(r).$$

Then G is defined for $\text{Im } \lambda > 0$ and $\chi G \chi$ continues holomorphically to $\lambda \in [-E, E] - i[0, \Gamma h]$. Define operators A_C, A_K, A_F by

$$\begin{aligned} (P - \lambda)G &= \text{Id} + [\chi_C(r - 1), h^2 D_r^2]R_C(\lambda)\chi_C(r) + [\chi_K(|r - 1|), h^2 D_r^2]R_K(\lambda)\chi_K(|r|) \\ &\quad + [\chi_F(r + 1), h^2 D_r^2]R_F(\lambda)\chi_F(r) \\ &= \text{Id} + A_C + A_K + A_F; \end{aligned}$$




Figure 3. The remainders A_C , A_K , and A_F are localized on the right in the region to the back of the arrows, and on the left near the tips of the arrows (A_C is localized on the right at the support of χ_C and on the left at the support of $\chi'_C(\cdot - 1)$, and so on), and this implies (3-6). They are microlocalized on the left in the indicated directions, and this implies (3-7) (since, by (2-3), no geodesic can follow one of the A_K arrows and then the A_F arrow, and so on).

see Figure 3. The estimates (3-2) and (3-3) only allow us to remove the remainders A_C, A_K, A_F by Neumann series for a narrow range of λ . To obtain a parametrix with improved remainders, observe that the support properties of the χ_j imply that

$$A_C^2 = A_K^2 = A_F^2 = A_C A_F = A_F A_C = 0; \tag{3-6}$$

so, solving away using G , we obtain

$$(P - \lambda)G(\text{Id} - A_C - A_K - A_F) = \text{Id} - A_K A_C - A_C A_K - A_F A_K - A_K A_F.$$

Now the propagation of singularities estimates (3-4) and (3-5) imply

$$\|A_F A_K\|_{L^2_\varphi(X) \rightarrow L^2_\varphi(X)} + \|A_C A_K A_C A_K\|_{L^2_\varphi(X) \rightarrow L^2_\varphi(X)} = \mathcal{O}(h^\infty). \tag{3-7}$$

In this sense the $A_F A_K$ remainder term is negligible. We again use (3-6) to write

$$\begin{aligned} (P - \lambda)G(\text{Id} - A_C - A_K - A_F + A_K A_C + A_C A_K + A_K A_F) \\ = \text{Id} - A_F A_K + A_C A_K A_C + A_F A_K A_C + A_K A_C A_K + A_C A_K A_F + A_K A_F A_K. \end{aligned}$$

Now all remainders but $A_C A_K A_C, A_K A_C A_K,$ and $A_C A_K A_F$ are negligible in the sense of (3-7). Solving away again gives

$$\begin{aligned} (P - \lambda)G(\text{Id} - A_C - A_K - A_F + A_K A_C + A_C A_K + A_K A_F - A_C A_K A_C - A_K A_C A_K - A_C A_K A_F) \\ = \text{Id} - A_F A_K + A_F A_K A_C + A_K A_F A_K - A_K A_C A_K A_C \\ - A_C A_K A_C A_K - A_F A_K A_C A_K - A_K A_C A_K A_F. \end{aligned}$$

Now all remainders but $A_K A_C A_K A_C$ are negligible. Solving away one last time gives

$$\begin{aligned} (P - \lambda)G(\text{Id} - A_C - A_K - A_F + A_K A_C + A_C A_K + A_K A_F \\ - A_C A_K A_C - A_K A_C A_K - A_C A_K A_F + A_K A_C A_K A_C) \\ = \text{Id} - A_F A_K + A_C A_K A_C + A_F A_K A_C + A_K A_F A_K - A_C A_K A_C A_K \\ - A_F A_K A_C A_K - A_K A_C A_K A_F + A_C A_K A_C A_K A_C + A_F A_K A_C A_K A_C =: \text{Id} + R, \end{aligned}$$

where R is defined by the equation, and $\|R\|_{L^2_\varphi(X) \rightarrow L^2_\varphi(X)} = \mathcal{O}(h^\infty)$. So for h small enough we may write, for $\text{Im } \lambda > 0$,

$$\begin{aligned} (P - \lambda)^{-1} = G(\text{Id} - A_C - A_K - A_F + A_K A_C + A_C A_K + A_K A_F \\ - A_C A_K A_C - A_K A_C A_K - A_C A_K A_F + A_K A_C A_K A_C) \sum_{k=0}^\infty (-R)^k. \end{aligned}$$

Combining this equation with (3-3), we see that $\chi(P - \lambda)^{-1} \chi$ continues to holomorphically to $|\text{Re } \lambda| \leq E$, $\text{Im } \lambda \geq -\Gamma h$ and obeys

$$\|\chi(P - \lambda)^{-1} \chi\|_{L^2_\varphi(X) \rightarrow H^2_{\varphi,h}(X)} \leq C h^{-1 - C_0 |\text{Im } \lambda|/h}. \quad \square$$

In summary, to prove (2-8) (and hence (1-1)), it remains to prove (3-3), (3-4) and (3-5).

3B. Statements of estimates for model operators. In this subsection we state six propositions: a resolvent estimate and a propagation of singularities estimate, for each of R_K , R_C , and R_F . Propositions 3.2, 3.4, and 3.6 imply (3-3) for $j = K, C$, and F , respectively. As we discuss after the statements, Propositions 3.3, 3.5, and 3.7 imply (3-4) and (3-5). The first two propositions concern R_K , and we prove them in Section 4. The next two concern R_C , and we prove them in Section 5. The last two concern R_F , and we prove them in Section 6. Hence at the end of Section 6 the proof of the Theorem will be complete.

Proposition 3.2. *For any $E \in (0, 1)$ there is $C_0 > 0$ such that for any $M > 0$ there are $C, h_0 > 0$ such that*

$$\|R_K(\lambda)\|_{L^2_\varphi(X) \rightarrow H^2_{\varphi,h}(X)} \leq C \begin{cases} h^{-1} + |\lambda|, & \text{Im } \lambda > 0, \\ h^{-1} e^{C_0 |\text{Im } \lambda|/h}, & \text{Im } \lambda \leq 0, \end{cases} \quad (3-8)$$

for $|\text{Re } \lambda| \leq E, \text{Im } \lambda \geq -M h \log(1/h), h \in (0, h_0]$.

Proposition 3.3. *Let $\Gamma \in \mathbb{R}, E \in (0, 1)$. Let $A, B \in \Psi^0(X)$ have full symbols a and b with the projections to X of $\text{supp } a$ and $\text{supp } b$ compact and suppose that*

$$\text{supp } a \cap \left[\text{supp } b \cup \bigcup_{t \geq 0} \exp(tH_p)[p^{-1}([-E, E]) \cap \text{supp } b] \right] = \emptyset, \quad (3-9)$$

where $\exp(tH_p)$ is the bicharacteristic flow of p . Then, for any $N \in \mathbb{N}$,

$$\|AR_K(\lambda)B\|_{L^2_\varphi(X) \rightarrow H^2_{\varphi,h}(X)} = \mathcal{O}(h^\infty) \quad (3-10)$$

for $|\text{Re } \lambda| \leq E, -\Gamma h \leq \text{Im } \lambda \leq h^{-N}$.

Proposition 3.4. *For every $\chi \in C_0^\infty(X)$, $E \in (0, 1)$, there is $C_0 > 0$ such that, for any $M > 0$, there are $h_0, C > 0$ such that the cutoff resolvent $\chi R_C(\lambda)\chi$ continues holomorphically from $\{\text{Im } \lambda > 0\}$ to $\{|\text{Re } \lambda| \leq E, \text{Im } \lambda \geq -Mh\}$, $h \in (0, h_0]$, and obeys*

$$\|\chi R_C(\lambda)\chi\|_{L_\varphi^2(X) \rightarrow H_{\varphi,h}^2(X)} \leq C \begin{cases} h^{-1} + |\lambda|, & \text{Im } \lambda > 0, \\ h^{-1-C_0|\text{Im } \lambda|/h}, & \text{Im } \lambda \leq 0. \end{cases} \tag{3-11}$$

Proposition 3.5. *Let $r_0 < 0$, $\chi_- \in C_0^\infty((-\infty, r_0))$, $\chi_+ \in C_0^\infty((r_0, \infty))$, $\varphi \in C^\infty(\mathbb{R})$ supported in $(-\infty, 0)$ and bounded with all derivatives, $E \in (0, 1)$, $\Gamma > 0$ be given. Then there exists $h_0 > 0$ such that*

$$\|\varphi(hD_r)\chi_+(r)R_C(\lambda)\chi_-(r)\|_{L_\varphi^2(X) \rightarrow H_{\varphi,h}^2(X)} = \mathcal{O}(h^\infty) \tag{3-12}$$

for $|\text{Re } \lambda| \leq E$, $-\Gamma h \leq \text{Im } \lambda \leq h^{-N}$, $h \in (0, h_0]$.

Proposition 3.6. *For every $\chi \in C_0^\infty(X)$, $E \in (0, 1)$, there is $C_0 > 0$ such that, for any $M > 0$, there are $h_0, C > 0$ such that the cutoff resolvent $\chi R_F(\lambda)\chi$ continues holomorphically from $\{\text{Im } \lambda > 0\}$ to $\{|\text{Re } \lambda| \leq E, \text{Im } \lambda \geq -Mh \log(1/h)\}$, $h \in (0, h_0]$, where it satisfies*

$$\|\chi R_F(\lambda)\chi\|_{L_\varphi^2(X) \rightarrow H_{\varphi,h}^2(X)} \leq C \begin{cases} h^{-1} + |\lambda|, & \text{Im } \lambda > 0, \\ h^{-1}e^{C_0|\text{Im } \lambda|/h}, & \text{Im } \lambda \leq 0. \end{cases} \tag{3-13}$$

Proposition 3.7. *Let $r_0 > R_g$, $\chi_- \in C_0^\infty((-\infty, r_0))$, $\chi_+ \in C_0^\infty((r_0, \infty))$, $\varphi \in C^\infty(\mathbb{R})$ supported in $(0, \infty)$ and bounded with all derivatives, $E \in (0, 1)$, $\Gamma > 0$ be given. Then there exists $h_0 > 0$ such that*

$$\|\chi_+(r)R_F(\lambda)\chi_-(r)\varphi(hD_r)\|_{L_\varphi^2(X) \rightarrow H_{\varphi,h}^2(X)} = \mathcal{O}(h^\infty) \tag{3-14}$$

for $|\text{Re } \lambda| \leq E$, $-\Gamma h \leq \text{Im } \lambda \leq h^{-N}$, $h \in (0, h_0]$.

We conclude the subsection by deducing (3-4) and (3-5) from the above propositions.

Take $\varphi \in C^\infty(\mathbb{R})$, bounded with all derivatives and supported in $(0, \infty)$, and take $\tilde{\chi}_2, \tilde{\chi}_3 \in C_0^\infty(X)$ such that $\text{supp } \tilde{\chi}_2 \subset \{r > R_g + 2\}$ and $\tilde{\chi}_3 \subset \{r < R_g + 2\}$, and such that $\tilde{\chi}_2\chi_2 = \chi_2\tilde{\chi}_2 = \chi_2$ and $\tilde{\chi}_3\chi_3 = \chi_3\tilde{\chi}_3 = \chi_3$. Then (3-4) follows from

$$\|\tilde{\chi}_3 R_F \tilde{\chi}_2 \varphi(hD_r)\|_{L_\varphi^2(X) \rightarrow H_{\varphi,h}^2(X)} + \|\tilde{\chi}_2(\text{Id} - \varphi(hD_r))R_K \chi_1\|_{L_\varphi^2(X) \rightarrow H_{\varphi,h}^2(X)} = \mathcal{O}(h^\infty). \tag{3-15}$$

The estimate on the first term follows from (3-14), while the estimate on the second term follows from (3-10) if $\text{supp}(1 - \varphi)$ is contained in a sufficiently small neighborhood of $(-\infty, 0]$; it suffices to take a neighborhood small enough that no bicharacteristic in $p^{-1}([-E, E])$ goes from $T^* \text{supp } \chi_1$ to $(T^* \text{supp } \tilde{\chi}_2) \cap \text{supp}(1 - \varphi(\rho))$, where ρ is the dual variable to r in T^*X , and such a neighborhood exists by (2-4) because when a bicharacteristic leaves $T^* \text{supp } \chi_1$ it has $\rho \geq 0$, and (2-4) gives a minimum amount by which ρ must grow in the time it takes the bicharacteristic to reach $T^* \text{supp } \tilde{\chi}_2$. An analogous argument reduces (3-5) to (3-12): the analog of (3-15) is

$$\|\tilde{\chi}_3 R_K(\text{Id} - \varphi(hD_r))\tilde{\chi}_2\|_{L_\varphi^2(X) \rightarrow H_{\varphi,h}^2(X)} + \|\varphi(hD_r)\tilde{\chi}_2 R_C \chi_1\|_{L_\varphi^2(X) \rightarrow H_{\varphi,h}^2(X)} = \mathcal{O}(h^\infty),$$

where $\varphi \in C^\infty(\mathbb{R})$ is bounded with all derivatives and supported in $(-\infty, 0)$, and $\tilde{\chi}_2, \tilde{\chi}_3 \in C_0^\infty(X)$ have $\text{supp } \tilde{\chi}_2 \subset \{r > -R_g - 2\}$ and $\tilde{\chi}_3 \subset \{r < -R_g - 2\}$, and such that $\tilde{\chi}_2 \chi_2 = \chi_2 \tilde{\chi}_2 = \chi_2$ and $\tilde{\chi}_3 \chi_3 = \chi_3 \tilde{\chi}_3 = \chi_3$.

4. Model operator in the nonsymmetric region

In this section we prove Propositions 3.2 and 3.3. Although the techniques involved are all essentially well known, we go over them in some detail here because they are important in the more complicated analysis of P_C and P_F below.

4A. Proof of Proposition 3.2. This is similar to the argument in [Sjöstrand and Zworski 2007, §4]. Fix

$$E_0 \in (E, 1), \quad \varepsilon = 10Mh \log(1/h).$$

We will use the assumption that the flow is nontrapping to construct an *escape function* $q \in C_0^\infty(T^*X)$, that is to say a function such that

$$H_p q \leq -1 \quad \text{near } T^* \text{supp}(1 - W_K) \cap p^{-1}([-E_0, E_0]). \tag{4-1}$$

The construction will be given below. Then let $Q \in \Psi^{-\infty}(X)$ be a quantization of q , and

$$P_{K,\varepsilon} = e^{\varepsilon Q/h} P_K e^{-\varepsilon Q/h} = P_K - \varepsilon[P_K, Q/h] + \varepsilon^2 R,$$

where $R \in \Psi^{-\infty}(X)$ (see (2-26)). We will prove that

$$\|(P_{K,\varepsilon} - E')^{-1}\|_{L_\varphi^2(X) \rightarrow H_{\varphi,h}^2(X)} \leq 5/\varepsilon, \quad E' \in [-E_0, E_0], \tag{4-2}$$

from which it follows, using first the openness of the resolvent set and then (2-23), that

$$\|(P_K - \lambda)^{-1}\|_{L_\varphi^2(X) \rightarrow H_{\varphi,h}^2(X)} \leq \frac{h^{-N}}{M \log(1/h)}, \quad |\text{Re } \lambda| \leq E_0, \quad |\text{Im } \lambda| \leq Mh \log(1/h), \tag{4-3}$$

where

$$N = 10M(\|Q\|_{H_{\varphi,h}^2(X) \rightarrow H_{\varphi,h}^2(X)} + \|Q\|_{L_\varphi^2(X) \rightarrow L_\varphi^2(X)}) + 1.$$

Then we will show how to use complex interpolation to improve (4-3) to (3-8).

*Construction of $q \in C_0^\infty(T^*X)$ satisfying (4-1).* As in [Vasy and Zworski 2000, §4], we take q of the form

$$q = \sum_{j=1}^J q_j, \tag{4-4}$$

where each q_j is supported near a bicharacteristic in $T^* \text{supp}(1 - W_K) \cap p^{-1}([-E_0, E_0])$.

First, for each $\varphi \in T^* \text{supp}(1 - W_K) \cap p^{-1}([-E_0, E_0])$, define the following *escape time*:

$$T_\varphi = \inf\{T \in \mathbb{R} : |t| \geq T - 1 \implies \exp(tH_p)\varphi \notin T^* \text{supp}(1 - W_K)\}.$$

Then put

$$T = \max\{T_\varphi : \varphi \in T^* \text{supp}(1 - W_K) \cap p^{-1}([-E_0, E_0])\}.$$

Note that the nontrapping assumption in Section 2A implies that $T < \infty$. Let S_φ be a hypersurface through φ , transversal to H_p near φ . If U_φ is a small enough neighborhood of φ , then

$$V_\varphi = \{\exp(tH_p)\varphi' : \varphi' \in U_\varphi \cap S_\varphi, |t| < T + 1\}$$

is diffeomorphic to $\mathbb{R}^{2n-1} \times (-T - 1, T + 1)$ with φ mapped to $(0, 0)$. Denote this diffeomorphism by (y_φ, t_φ) . Further shrinking U_φ if necessary, we may assume the inverse image of $\mathbb{R}^{2n-1} \times \{|t| \geq T\}$ is disjoint from $T^* \text{supp}(1 - W_K)$. Then take $\varphi \in C_0^\infty(\mathbb{R}^{2n-1}; [0, 1])$ identically 1 near 0, and $\chi \in C_0^\infty((-T - 1, T + 1))$ with $\chi' = -1$ near $[-T, T]$, and put

$$q_\varphi = \varphi(y_\varphi)\chi(t_\varphi), \quad H_p q_\varphi = \varphi(y_\varphi)\chi'(t_\varphi).$$

Note $H_p q_\varphi \leq 0$ on $T^* \text{supp}(1 - W_K)$ because $\chi' = -1$ there. Let V'_φ be the interior of $\{H_p q_\varphi = -1\}$, note that the V'_φ cover $T^*(1 - W_K) \cap p^{-1}([-E_0, E_0])$, and extract a finite subcover $\{V'_{\varphi_1}, \dots, V'_{\varphi_J}\}$. Then put $q_j = q_{\varphi_j}$ and define q by (4-4), so that

$$H_p q = \sum_{j=1}^J \varphi(y_{\varphi_j})\chi'_{\varphi_j}(t_{\varphi_j}).$$

Then $H_p q \leq -1$ near $T^*(1 - W_K) \cap p^{-1}([-E_0, E_0])$ because at each point at least one summand is, and the other summands are nonpositive. □

Proof of (4-2). Let $\chi_0 \in C_0^\infty(X; [0, 1])$ be identically 1 on a large enough set that $\chi_0 Q = Q \chi_0 = Q$. In particular we have $(1 - \chi_0)W_K = 1 - \chi_0$, allowing us to write

$$\|(1 - \chi_0)u\|_{L^2_\varphi(X)}^2 = -\text{Im}\langle (P_{K,\varepsilon} - E')(1 - \chi_0)u, (1 - \chi_0)u \rangle_{L^2_\varphi(X)}.$$

Hence

$$\|(1 - \chi_0)u\|_{L^2_\varphi(X)} \leq \|(P_{K,\varepsilon} - E')u\|_{L^2_\varphi(X)} + \|[P_{K,\varepsilon}, \chi_0]u\|_{L^2_\varphi(X)}.$$

To estimate $\|\chi_0 u\|_{L^2_\varphi(X)}$ and the remainder term $\|[P_{K,\varepsilon}, \chi_0]u\|_{L^2_\varphi(X)}$ we introduce a microlocal cutoff $\phi \in C_0^\infty(T^*X)$ which is identically 1 near $T^* \text{supp}(1 - W_K) \cap p^{-1}([-E_0, E_0])$ and is supported in the interior of the set where $H_p q \leq -1$. Since the principal symbol of $P_{K,\varepsilon} - E'$ is

$$p_{K,\varepsilon} - E' = p - iW_K - E' - i\varepsilon\{p - iW_K, q\},$$

we have

$$|p_{K,\varepsilon} - E'| \geq 1 - E_0 \quad \text{near } \text{supp}(1 - \phi)$$

for $|E'| \leq E_0$, provided h (and hence ε) is sufficiently small. Then if $\Phi \in \Psi^{-\infty}(X)$ is a quantization of ϕ , we find using the semiclassical elliptic estimate (2-21) that

$$\|(\text{Id} - \Phi)\chi_0 u\|_{H^2_{\varphi,h}(X)} \leq C(\|(P_{K,\varepsilon} - E')u\|_{L^2_\varphi(X)} + h\|u\|_{H^1_{\varphi,h}(X)}).$$




Figure 4. Bounds on f used in the complex interpolation argument.

Since $H_p q \leq -1$ near $\text{supp } \phi$ we see that

$$\text{Im } p_{K,\varepsilon} - E' = -W_K - \varepsilon\{p, q\} \leq -\varepsilon \quad \text{near } \text{supp } \phi.$$

Then, using the sharp Gårding inequality (2-22), we find that

$$\begin{aligned} \|(P_{K,\varepsilon} - E')\Phi\chi_0 u\|_{L^2_\varphi(X)} \|\Phi\chi_0 u\|_{L^2_\varphi(X)} &\geq -\langle \text{Im}(P_{K,\varepsilon} - E')\Phi\chi_0 u, \Phi\chi_0 u \rangle_{L^2_\varphi(X)} \\ &\geq \varepsilon \|\Phi\chi_0 u\|_{L^2_\varphi(X)}^2 - Ch \|u\|_{H^{1/2}_{\varphi,h}(X)}^2. \end{aligned}$$

This implies that

$$\begin{aligned} \|u\|_{L^2_\varphi(X)} &\leq \|(1 - \chi_0)u\|_{L^2_\varphi(X)} + \|\Phi\chi_0 u\|_{L^2_\varphi(X)} + \|(\text{Id} - \Phi)\chi_0 u\|_{L^2_\varphi(X)} \\ &\leq C \|(P_{K,\varepsilon} - E')u\|_{L^2_\varphi(X)} + \varepsilon^{-1} \|(P_{K,\varepsilon} - E')u\|_{L^2_\varphi(X)} + Ch^{1/2} \|u\|_{H^{1/2}_{\varphi,h}(X)}. \end{aligned}$$

As in the proof of (3-2), combining this with

$$\begin{aligned} \|u\|_{H^2_{\varphi,h}(X)} &\leq 3\|u\|_{L^2_\varphi(X)} + \|(P - E')u\|_{L^2_\varphi(X)} \\ &\leq 4\|u\|_{L^2_\varphi(X)} + \|(P_{K,\varepsilon} - E')u\|_{L^2_\varphi(X)} + C\varepsilon \|u\|_{L^2_\varphi(X)}, \end{aligned} \tag{4-5}$$

we obtain (4-2) for h sufficiently small. □

Proof that (4-3) implies (3-8). We follow the approach of [Tang and Zworski 1998] as presented in [Nakamura, Stefanov and Zworski 2003, Lemma 3.1]. Observe first that (3-2) implies (3-8) for $\text{Im } \lambda \geq C_\Omega h$ for any $C_\Omega > 0$.

Let $f(\lambda, h)$ be holomorphic in λ for $\lambda \in \Omega = [-E_0, E_0] + i[-Mh \log(1/h), C_\Omega h]$ and bounded uniformly in h there. Suppose further that, for $\lambda \in \Omega$,

$$|\text{Re } \lambda| \leq E \implies |f| \geq 1, \quad |\text{Re } \lambda| \in \left[\frac{1}{2}(E + E_0), E_0\right] \implies |f| \leq h^N.$$

For example, we may take f to be a characteristic function convolved with a gaussian:

$$f(\lambda, h) = \frac{2}{\sqrt{\pi}} \log(1/h) \int_{-\tilde{E}}^{\tilde{E}} \exp(-\log^2(1/h)(\lambda - y)^2) dy$$

$$= \operatorname{erfc}(\log(1/h)(\lambda - \tilde{E})) - \operatorname{erfc}(\log(1/h)(\lambda + \tilde{E})),$$

where $\tilde{E} = \frac{1}{4}(3E + E_0)$, $\operatorname{erfc} z = 2 \int_z^\infty e^{-t^2} dt / \sqrt{\pi}$. We bound $|f|$ using the identity $\operatorname{erfc}(z) + \operatorname{erfc}(-z) = 2$ and the fact that $\operatorname{erfc} z = \pi^{-1/2} z^{-1} e^{-z^2} (1 + \mathcal{O}(z^{-2}))$ for $|\arg z| < \frac{3\pi}{4}$.

Then the subharmonic function

$$g(\lambda, h) = \log \|(P_K - \lambda)^{-1}\|_{L^2_\varphi(X) \rightarrow H^2_{\varphi,h}(X)} + \log |f(\lambda, h)| + \frac{N \operatorname{Im} \lambda}{Mh}$$

obeys

$$g \leq C \quad \text{on } \partial\Omega \cap (\{|\operatorname{Re} \lambda| = E_0\} \cup \{\operatorname{Im} \lambda = -Mh \log(1/h)\})$$

and

$$g \leq C + \log(1/h) \quad \text{on } \partial\Omega \cap \{\operatorname{Im} \lambda = C_\Omega h\}.$$

From the maximum principle and the lower bound on $|f|$ we obtain

$$\log \|(P_K - \lambda)^{-1}\|_{L^2_\varphi(X) \rightarrow H^2_{\varphi,h}(X)} + \frac{N \operatorname{Im} \lambda}{Mh} \leq C + \log(1/h),$$

for $\lambda \in \Omega$, $|\operatorname{Re} \lambda| \leq E$, from which (3-8) follows for $\lambda \in \Omega$. □

4B. Proof of Proposition 3.3. This is similar to [Datchev and Vasy 2012a, Lemma 5.1]. By (2-21), without loss of generality we may assume that a is supported in a neighborhood of $p^{-1}([-E, E]) \cap \operatorname{supp}(1 - W_K)$ which is as small as we please (but independent of h). In particular we may assume $\operatorname{supp} a$ is compact.

We will show that if $(P_K - \lambda)u = Bf$ with $\|f\|_{L^2_\varphi(X)} = 1$, and if $\|A_0 u\| \leq Ch^k$ for some $A_0 \in \Psi^0(X)$ with full symbol a_0 such that

$$a_0 = 1 \quad \text{near } \operatorname{supp} a \cap p^{-1}([-E, E]), \quad \operatorname{supp} a_0 \cap \bigcup_{t \geq 0} \exp(tH_p) \operatorname{supp} b = \emptyset,$$

then $\|A_1 u\| \leq Ch^{k+1/2}$ for each $A_1 \in \Psi^0(X)$ with full symbol a_1 satisfying $a_0 = 1$ near $\operatorname{supp} a_1$. Then the conclusion (3-10) follows by induction; the base step is given by (3-8).

Let $q \in C_0^\infty(T^*X; [0, \infty))$ such that

$$a_0 = 1 \quad \text{near } \operatorname{supp} q, \quad H_p(q^2) \leq -(2\Gamma + 1)q^2 \quad \text{near } \operatorname{supp} a_1, \tag{4-6}$$

$$H_p q \leq 0 \quad \text{on } T^* \operatorname{supp}(1 - W_K). \tag{4-7}$$

The construction of q is very similar to that of the function q used in the proof of Proposition 3.2 above, and is also given in [loc. cit.]. Write

$$H_p(q^2) = -\ell^2 + r,$$

where $\ell, r \in C_0^\infty(T^*X)$ satisfy

$$\ell^2 \geq (2\Gamma + 1)q^2, \quad \text{supp } r \subset \{W_K = 1\}. \tag{4-8}$$

Let $Q, L, R \in \Psi^{-\infty}(X)$ have principal symbols q, ℓ, r respectively. Then

$$i[P, Q^*Q] = -hL^*L + hR + h^2F + R_\infty,$$

where $F \in \Psi^{-\infty}(X)$ has full symbol supported in $\text{supp } q$ and $R_\infty \in h^\infty\Psi^{-\infty}(X)$. From this we conclude that

$$\begin{aligned} \|Lu\|_{L_\phi^2(X)}^2 &= -\frac{2}{h} \text{Im}\langle Q^*QPu, u \rangle_{L_\phi^2(X)} + \langle Ru, u \rangle_{L_\phi^2(X)} + h\langle Fu, u \rangle_{L_\phi^2(X)} + \mathcal{O}(h^\infty)\|u\|_{L_\phi^2(X)}^2 \\ &= -\frac{2}{h} \text{Im}\langle Q^*Q(P_K - \lambda)u, u \rangle_{L_\phi^2(X)} - \text{Re}\langle Q^*QW_Ku, u \rangle_{L_\phi^2(X)} - \frac{2}{h} \text{Im } \lambda \|Qu\|_{L_\phi^2(X)}^2 \\ &\quad + \langle Ru, u \rangle_{L_\phi^2(X)} + h\langle Fu, u \rangle_{L_\phi^2(X)} + \mathcal{O}(h^\infty)\|u\|_{L_\phi^2(X)}^2. \end{aligned} \tag{4-9}$$

We now estimate the right-hand side of (4-9) term by term to prove that

$$\|Lu\|_{L_\phi^2(X)}^2 \leq 2\Gamma\|Qu\|_{L_\phi^2(X)}^2 + Ch\|A_0u\|_{L_\phi^2(X)}^2 + \mathcal{O}(h^\infty)\|u\|_{L_\phi^2(X)}^2. \tag{4-10}$$

Indeed, since $\text{supp } q \cap \text{supp } b = \emptyset$ and since $(P_K - \lambda)u = Bf$ it follows that

$$\langle Q^*Q(P_K - \lambda)u, u \rangle_{L_\phi^2(X)} = \mathcal{O}(h^\infty)\|u\|_{L_\phi^2(X)}^2.$$

Next, we write

$$-\text{Re}\langle Q^*QW_Ku, u \rangle_{L_\phi^2(X)} = -\text{Re}\langle W_KQu, Qu \rangle_{L_\phi^2(X)} + \langle Q^*[W_K, Q]u, u \rangle_{L_\phi^2(X)},$$

and observe that the first term is nonpositive because $W_K \geq 0$, and the second term is bounded by $Ch\|A_0u\|_{L_\phi^2(X)}^2$. Since $\text{Im } \lambda \geq -\Gamma h$ we have

$$-\frac{2}{h} \text{Im } \lambda \|Qu\|_{L_\phi^2(X)}^2 \leq 2\Gamma\|Qu\|_{L_\phi^2(X)}^2,$$

while since $W_K = 1$ on $\text{supp } r$ we have the elliptic estimate

$$\langle Ru, u \rangle_{L_\phi^2(X)} = C\|R(P_K - \lambda)u\|_{L_\phi^2(X)}\|u\|_{L_\phi^2(X)} + Ch\|A_0u\|_{L_\phi^2(X)}^2,$$

and the first term is $\mathcal{O}(h^\infty)\|u\|_{L_\phi^2(X)}^2$ since $\text{supp } r \cap \text{supp } b = \emptyset$. Finally $h\langle Fu, u \rangle_{L_\phi^2(X)} \leq Ch\|A_0u\|^2$ by the inductive hypothesis, giving (4-10).

But by (4-8) and the sharp Gårding inequality we have

$$\langle (D^*D - (2\Gamma + 1)Q^*Q)u, u \rangle \geq -Ch\|A_0u\|^2 - \mathcal{O}(h^\infty)\|u\|^2.$$

Hence by the inductive hypothesis we have

$$\|Qu\|^2 \leq Ch^{2k+1}\|u\|^2,$$

completing the inductive step.

5. Model operator in the cusp

In this section we prove Propositions 3.4 and 3.5. We begin by separating variables over the eigenspaces of Δ_{S_-} , writing

$$P_C = \bigoplus_{m=0}^{\infty} h^2 D_r^2 + (h\lambda_m)^2 e^{-2(r+\beta(r))} + h^2 V(r) - 1 - i W_C(r),$$

where $0 = \lambda_0 < \lambda_1 \leq \dots$ are square roots of the eigenvalues of Δ_{S_-} . Roughly speaking, it suffices to prove (3-11), (3-12) with P_C replaced by $P(\alpha)$, with estimates uniform in $\alpha \in \{0\} \cup [h\lambda_1, \infty)$, where

$$P(\alpha) = h^2 D_r^2 + \alpha^2 e^{-2(r+\beta(r))} + h^2 V(r) - 1 - i W_C(r).$$

The precise estimates for these operators which imply Propositions 3.4 and 3.5 are stated in Lemmas 5.1, 5.2, and 5.3 below.

5A. The case $\alpha = 0$. The analysis of $(P(0) - \lambda)^{-1}$ is very similar to that of R_K in Section 4. The only additional technical ingredient is the method of complex scaling, which for this operator works just as in [Sjöstrand and Zworski 1991; 2007].

Lemma 5.1. *For every $\chi \in C_0^\infty(X)$, $E \in (0, 1)$, there is $C_0 > 0$ such that, for any $M > 0$, there exist $h_0, C > 0$ such that the cutoff resolvent $\chi(P(0) - \lambda)^{-1}\chi$ continues holomorphically from $\{\text{Im } \lambda > 0\}$ to $\{|\text{Re } \lambda| \leq E, \text{Im } \lambda \geq -Mh \log(1/h)\}$, $h \in (0, h_0]$, and obeys*

$$\|\chi(P(0) - \lambda)^{-1}\chi\|_{L^2(\mathbb{R}) \rightarrow H_h^2(\mathbb{R})} \leq Ch^{-1}e^{C_0|\text{Im } \lambda|/h}. \tag{5-1}$$

Let $r_0 \in \mathbb{R}$, $\chi_- \in C_0^\infty((-\infty, r_0))$, $\chi_+ \in C_0^\infty((r_0, \infty))$, $\varphi \in C^\infty(\mathbb{R})$ supported in $(-\infty, 0)$ and bounded with all derivatives, $\Gamma > 0$ be given. Then there exists $h_0 > 0$ such that

$$\|\varphi(hD_r)\chi_+(r)(P(0) - \lambda)^{-1}\chi_-(r)\|_{L^2(\mathbb{R}) \rightarrow H_h^2(\mathbb{R})} = \mathcal{O}(h^\infty) \tag{5-2}$$

for $|\text{Re } \lambda| \leq E$, $-\Gamma h \leq \text{Im } \lambda \leq h^{-N}$, $h \in (0, h_0]$.

Proof of (5-1). We use complex scaling to replace $P(0)$ by the complex scaled operator $P_\delta(0)$, defined below. As we will see, $P_\delta(0)$ is semiclassically elliptic for $|r|$ sufficiently large and obeys (5-1) without cutoffs.

We have

$$P(0) = h^2 D_r^2 + h^2 V(r) - 1 - i W_C(r).$$

Fix $R > R_g$ sufficiently large that

$$\text{supp } \chi \cup \text{supp } \chi_+ \cup \text{supp } \chi_- \subset (-R, \infty). \tag{5-3}$$

Let $\gamma \in C^\infty(\mathbb{R})$ be nondecreasing and obey $\gamma(r) = 0$ for $r \geq -R$, $\gamma'(r) = \tan \theta_0$ for $r \leq -R - 1$ (here θ_0 is as in Section 2A), and impose further that $\beta(r)$ is holomorphic near $r + i\delta\gamma(r)$ for every $r < -R$, $\delta \in (0, 1)$. Below we will take $\delta \ll 1$ independent of h .

Now put

$$P_\delta(0) = \frac{h^2 D_r^2}{(1 + i\delta\gamma'(r))^2} - h \frac{\delta\gamma''(r)hD_r}{(1 + i\delta\gamma'(r))^3} + h^2 V(r + i\delta\gamma(r)) - 1 - iW_C(r + i\delta\gamma(r)).$$

If we define the differential operator with complex coefficients

$$\tilde{P}(0) = h^2 D_z^2 + h^2 V(z) - 1 - iW_C(z),$$

where z varies in $\{z = r + i\delta\gamma(r) : r \in \mathbb{R}, \delta \in (0, 1)\}$, and where $W_C(z) := 0$ whenever $\text{Im } z \neq 0$, then we have

$$P(0) = \tilde{P}(0)|_{\{z=r:r \in \mathbb{R}\}}, \quad P_\delta(0) = \tilde{P}(0)|_{\{z=r+i\delta\gamma(r):r \in \mathbb{R}\}}. \tag{5-4}$$

We will show that if $\chi_0 \in C^\infty(\mathbb{R})$ has $\text{supp } \chi_0 \cap \text{supp } \gamma = \emptyset$, then

$$\chi_0(P(0) - \lambda)^{-1} \chi_0 = \chi_0(P_\delta(0) - \lambda)^{-1} \chi_0, \quad \text{Im } \lambda > 0. \tag{5-5}$$

From this it follows that if one of these operators has a holomorphic continuation to any domain, then so does the other, and the continuations agree, so that it suffices to prove (5-1) and (5-2) with $P(0)$ replaced by $P_\delta(0)$. To prove (5-5) we will prove that if

$$(P(0) - \lambda)u = v \quad \text{and} \quad (P_\delta(0) - \lambda)u_\delta = v$$

for $v \in L^2(\mathbb{R})$ with $\text{supp } v \subset \{r : \gamma(r) = 0\}$, and $u, u_\delta \in L^2(\mathbb{R})$, then

$$u|_{\{r:\gamma(r)=0\}} = u_\delta|_{\{r:\gamma(r)=0\}}.$$

Thanks to (5-4), it suffices to show that if \tilde{u} solves $(\tilde{P}(0) - \lambda)\tilde{u} = v$ with $\tilde{u}|_{\{z=r:r \in \mathbb{R}\}} \in L^2(\mathbb{R})$, then $\tilde{u}|_{\{z=r+i\delta\gamma(r):r \in \mathbb{R}\}} \in L^2(\mathbb{R})$. For the proof of this statement we may take λ fixed with $\text{Re } \lambda = 0$ since the general statement follows by holomorphic continuation.

Observe that for $\text{Re } z < -R$, we have

$$(\tilde{P}(0) - \lambda)\tilde{u}(z) = 0. \tag{5-6}$$

We will use the WKB method to construct solutions u_\pm to (5-6) which are exponentially growing or decaying as $\text{Re } z \rightarrow -\infty$. Define

$$f(z) = V(z) - (1 + \lambda)/h^2, \quad \varphi(z) = (4f(z)f''(z) - 5f'(z)^2)(16f(z))^{-5/2}.$$

Now (see, e.g., [Olver 1974, Chapter 6, Theorem 11.1]) there exist two solutions to (5-6) given by

$$u_\pm(z) = f(z)^{-1/4} e^{\pm \int_{\gamma_{z,-R}} \sqrt{f(z')} dz'} (1 + b_\pm(z)), \quad \text{Re } z < -R,$$

taking principal branches of the roots and with the contour of integration $\gamma_{z,-R}$ taken from z to $-R$ such that $\sqrt{\text{Re } z'}$ is monotonic along $\gamma_{z,-R}$. The functions b_\pm obey

$$|b_\pm(z)| \leq \exp(\max(|\varphi(z')| : z' \in \gamma_\pm)) - 1 \leq Ch$$

when $\operatorname{Re} z > R$, where γ_+ and γ_- are contours from $-\infty$ to z and from z to $-R$, respectively, such that $\sqrt{\operatorname{Re} z'}$ is monotonic along the contour. It follows that, for fixed h sufficiently small,

$$|u_+(z)| \leq C e^{\operatorname{Re} z/C}, \quad |u_-(z)| \geq C e^{-\operatorname{Re} z/C}$$

for $\operatorname{Re} z < -R$. Hence $\tilde{u}|_{\{z=r:r \in \mathbb{R}\}} \in L^2(\mathbb{R})$ implies that \tilde{u} is proportional to u_+ . This implies that $\tilde{u}|_{\{z=r+i\delta\gamma(r):r \in \mathbb{R}\}} \in L^2(\mathbb{R})$, completing the proof of (5-5).

Fix

$$E_0 \in (E, 1), \quad \varepsilon = 10Mh \log(1/h).$$

The semiclassical principal symbol of $P_\delta(0)$ is

$$p_\delta(0) = \frac{\rho^2}{(1 + i\delta\gamma'(r))^2} - 1 = \rho^2(1 + \mathcal{O}(\delta)) - 1. \tag{5-7}$$

In this case the escape function can be made more explicit: we take $q \in C_0^\infty(T^*\mathbb{R})$ with

$$q(r, \rho) = -4r\rho(1 - E_0)^{-2}, \quad H_{p_\delta(0)}q = -8\rho^2(1 - E_0)^{-2}(1 + \mathcal{O}(\delta)) \tag{5-8}$$

on $\{|r| \leq R + 1, |\rho| \leq 2\}$. Let $Q \in \Psi^{-\infty}(\mathbb{R})$ be a quantization of q and put

$$P_{\delta,\varepsilon}(0) = e^{\varepsilon Q/h} P_\delta(0) e^{-\varepsilon Q/h} = P_\delta(0) - \varepsilon [P_\delta(0), Q/h] + \varepsilon^2 R,$$

where $R \in \Psi^{-\infty}(\mathbb{R})$ (see (2-26)). We will prove

$$\|(P_{\delta,\varepsilon}(0) - E')^{-1}\|_{L^2(\mathbb{R}) \rightarrow H_h^2(\mathbb{R})} \leq 5/\varepsilon, \quad E' \in [-E_0, E_0], \tag{5-9}$$

from which it follows by (2-23) that

$$\|(P_\delta(0) - \lambda)^{-1}\|_{L^2(\mathbb{R}) \rightarrow H_h^2(\mathbb{R})} \leq \frac{h^{-N}}{M \log(1/h)}, \quad |\operatorname{Re} \lambda| \leq E_0, \quad |\operatorname{Im} \lambda| \leq Mh \log(1/h), \tag{5-10}$$

where $N = 10M(\|Q\|_{H_h^2(\mathbb{R}) \rightarrow H_h^2(\mathbb{R})} + \|Q\|_{L^2(\mathbb{R}) \rightarrow L^2(\mathbb{R})}) + 1$. As before we will use complex interpolation to improve (5-10) to

$$\|(P_\delta(0) - \lambda)^{-1}\|_{L^2(\mathbb{R}) \rightarrow H_h^2(\mathbb{R})} \leq Ch^{-1} e^{C|\operatorname{Im} \lambda|/h} \tag{5-11}$$

for $-E \leq \operatorname{Re} \lambda \leq E, \operatorname{Im} \lambda > -Mh \log(1/h)$. Combining (5-5) and (5-11) gives (5-1).

Let $\phi \in C_0^\infty(\mathbb{R}; [0, 1])$ have $\phi(\rho) = 1$ for $|\rho|$ near $[1 - E_0, 1 + E_0]$ and $\operatorname{supp} \phi \subset \{\frac{1}{2}(1 - E_0) < |\rho| < 2\}$. By (5-7), if δ is small enough and h is small enough depending on δ , then on $\operatorname{supp}(1 - \phi(\rho))$ we have $|p_{\delta,\varepsilon}(0) - E'| \geq \delta(1 + \rho^2)/C$, uniformly in $E' \in [-E_0, E_0]$ and in h , where $p_{\delta,\varepsilon}(0)$ is the semiclassical principal symbol of $P_{\delta,\varepsilon}(0)$. Hence, by the semiclassical elliptic estimate (2-18),

$$\|(\operatorname{Id} - \phi(hD_r))u\|_{H_h^2(\mathbb{R})} \leq C\delta^{-1} \|(P_{\delta,\varepsilon}(0) - E')(\operatorname{Id} - \phi(hD_r))u\|_{L^2(\mathbb{R})} + \mathcal{O}(h^\infty) \|u\|_{H_h^{-N}(\mathbb{R})}.$$

On $\operatorname{supp} \phi(\rho)$ we use the negativity of the imaginary part of the principal symbol of $P_{\delta,\varepsilon}(0)$. Indeed, on $\{(r, \rho) : \rho \in \operatorname{supp} \phi, |r| \leq R + 1\}$ we have, using (5-8),

$$\operatorname{Im} p_{\delta,\varepsilon}(0) = \operatorname{Im} p_\delta(0) + \operatorname{Im} i\varepsilon H_{p_{\delta,\varepsilon}(0)}q = \frac{-2\delta\gamma'(r)\rho^2}{|1 + i\delta\gamma'(r)|^4} - \frac{8\varepsilon\rho^2}{(1 - E_0)^2}(1 + \mathcal{O}(\delta)) \leq -\varepsilon,$$

provided δ is sufficiently small. Meanwhile, on $\{(r, \rho) : \rho \in \text{supp } \phi, |r| \geq R + 1\}$ we have

$$\text{Im } p_{\delta, \varepsilon}(0) = \text{Im } p_{\delta}(0) + \text{Im } i \varepsilon H_{p_{\delta, \varepsilon}(0)} q = \frac{-2\delta \tan \theta_0 \rho^2}{|1 + i \delta \tan \theta_0|^4} + \mathcal{O}(\varepsilon) \leq -\delta/C,$$

provided h (and hence ε) is sufficiently small.

Then, using the sharp Gårding inequality (2-19), we have, for h sufficiently small,

$$\begin{aligned} \|\varphi(hD_r)u\|_{L^2(\mathbb{R})} \|(P_{\delta, \varepsilon}(0) - E')\varphi(hD_r)u\|_{L^2(\mathbb{R})} &\geq -\langle \text{Im}(P_{\delta, \varepsilon}(0) - E')\varphi(hD_r)u, \varphi(hD_r)u \rangle_{L^2(\mathbb{R})} \\ &\geq \varepsilon \|\varphi(hD_r)u\|_{L^2(\mathbb{R})}^2 - Ch \|u\|_{H_h^{1/2}(\mathbb{R})}^2. \end{aligned}$$

We deduce (5-9) from this just as we did (4-2) above.

To improve (5-10) to (5-11) we use almost the same complex interpolation argument as we did to improve (4-3) to (3-8). The only difference is that in the first step we note that

$$\text{Im } p_{\delta}(0) = \frac{-2\delta \gamma'(r)}{|1 + i \delta \gamma'(r)|^4} \leq 0,$$

so by the sharp Gårding inequality (2-19) we have, for some $C_{\Omega} > 0$,

$$\langle \text{Im } P_{\delta}(0)u, u \rangle_{L^2(\mathbb{R})} \geq -C_{\Omega} h \|u\|_{L^2(\mathbb{R})}^2,$$

so that $\|(P_{\delta}(0) - \lambda)^{-1}\|_{L^2(\mathbb{R})} \leq 1/C_{\Omega} h$, when $\text{Im } \lambda \geq 2C_{\Omega} h$. □

Proof of (5-2). Let $(P_{\delta}(0) - \lambda)u = f$, where $\|f\|_{L^2(\mathbb{R})} = 1$, $\text{supp } f \subset \text{supp } \chi_-$ and $P_{\delta}(0)$ is as in the proof of (5-1). We must show that

$$\|\varphi(hD_r)\chi_+(r)u\|_{H_h^2(\mathbb{R})} = \mathcal{O}(h^{\infty}); \tag{5-12}$$

recall that the replacement of $P(0)$ by $P_{\delta}(0)$ is justified by (5-5). To prove (5-12) we use an argument by induction based on a nested sequence of escape functions.

More specifically, take

$$q = \varphi_r(r)\varphi_{\rho}(\rho), \quad H_{p_{\delta}(0)}q = 2\rho\varphi'_r(r)\varphi_{\rho}(\rho) + \mathcal{O}(\delta),$$

where $\varphi_r \in C_0^{\infty}(\mathbb{R}; [0, \infty))$ with $\text{supp } \varphi_r \subset (r_0, \infty)$, $\varphi'_r \geq 0$ near $[r_0, R + 1]$ (here R is as in (5-3)), $\varphi'_r > 0$ near $\text{supp } \chi_+$. Take $\varphi_{\rho} \in C_0^{\infty}(\mathbb{R}; [0, \infty))$ with $\text{supp } \varphi_{\rho} \subset (-\infty, 0)$, $\varphi'_{\rho} \leq 0$ near $[-2, 0]$, $\varphi_{\rho} \neq 0$ near $\text{supp } \varphi \cap [-2, 0]$. Impose further that $\sqrt{\varphi_r}, \sqrt{\varphi_{\rho}} \in C_0^{\infty}(\mathbb{R})$, and that $\varphi'_r \geq c\varphi_r$ for $r \leq R + 1$, where $c > 0$ is chosen large enough that $H_{p_0(\delta)}q \leq -(2\Gamma + 1)q$ on $\{r \leq R + 1, \rho \geq -2\}$; see Figure 5.

We will show that if $\|A_0u\|_{L^2(\mathbb{R})} \leq Ch^k$ for $A_0 \in \Psi^0(\mathbb{R})$ with full symbol supported sufficiently near $\text{supp } q$ and for some $k \in \mathbb{R}$, then $\|A_1u\|_{L^2(\mathbb{R})} \leq Ch^{k+1/2}$ for $A_1 \in \Psi^0(\mathbb{R})$ with full symbol supported sufficiently near $\{r \in \text{supp } \chi_+, \rho \in \text{supp } \varphi\}$. The conclusion (5-12) then follows by induction. (The base step of the induction follows from (5-11) or even from (5-10).)

In the remainder of the proof all norms and inner products are in $L^2(\mathbb{R})$ and we omit the subscript for brevity.

We write

$$H_{p_{\delta}(0)}q^2 = -b^2 + e,$$




Figure 5. The escape function q used to prove propagation of singularities (5-2) in the case $\alpha = 0$. The derivative along the flow lines $H_{p_\delta(0)}q$ is negative and provides ellipticity for our positive commutator argument near $\{r \in \text{supp } \chi_+, \rho \in \text{supp } \varphi\}$. We allow $H_{p_\delta(0)}q > 0$ (the unfavorable sign for us) only in $\{r > R + 1\}$ and in $\{\rho < -2\}$, because in this region $p_\delta(0)$ is elliptic.

where $b, e \in C_0^\infty(T^*\mathbb{R})$, $b > 0$ near $\{r \in \text{supp } \chi_+, \rho \in \text{supp } \varphi, -2 \leq \rho\}$, $b^2 \geq (2\Gamma + 1)q^2$ everywhere, and $\text{supp } e \cap (\{r \leq R + 1, \rho \geq -2\} \cup \{r \leq r_0\}) = \emptyset$. Let Q, B, E be quantizations of q, b, e respectively. Then

$$i[P_\delta(0), Q^*Q] = -hB^*B + hE + h^2F,$$

where $F \in \Psi^0(\mathbb{R})$ has full symbol supported in $\text{supp } q$. From this we conclude that

$$\|Bu\|^2 = -\frac{2}{h} \text{Im} \langle Q^*Q(P_\delta(0) - \lambda)u, u \rangle - \frac{2}{h} \text{Im} \lambda \|Qu\|^2 + \langle Eu, u \rangle + h \langle Fu, u \rangle + \mathcal{O}(h^\infty) \|u\|^2.$$

From $(P_\delta(0) - \lambda)u = f$ and $\text{WF}'_h Q \cap T^* \text{supp } f = \emptyset$ it follows that the first term is $\mathcal{O}(h^\infty) \|u\|^2$. Similarly $\text{WF}'_h E \cap (\text{supp } f \cup p_\delta^{-1}(0)) = \emptyset$ implies by (2-18) that the third term is $\mathcal{O}(h^\infty) \|u\|^2$. The fourth term is bounded by $Ch^{2k+1} \|u\|^2$ by the inductive hypothesis, giving

$$\|Bu\|^2 \leq 2\Gamma \|Qu\|^2 + Ch^{2k+1} \|u\|^2.$$

By (2-19) we have

$$\langle (B^*B - (2\Gamma + 1)Q^*Q)u, u \rangle \geq -Ch \|Ru\|^2,$$

where $R \in \Psi_0^{0,0}(\mathbb{R})$ is microsupported in an arbitrarily small neighborhood of $\text{WF}'_h Q$. Hence $\|Ru\| \leq Ch^k \|u\|$ and we have

$$\|Qu\|^2 \leq Ch^{2k+1} \|u\|^2,$$

completing the inductive step and also the proof. □

5B. The case $\alpha \geq \lambda_1 h$. Propositions 3.4 and 3.5 follow from (5-1), (5-2) and the following two lemmas.

Lemma 5.2. *For any $E \in (0, 1)$ there is $C_0 > 0$ such that for any $M, \lambda_1 > 0$ there are $h_0, C > 0$ such that if $h \in (0, h_0], \alpha \geq \lambda_1 h, \lambda \in [-E, E] + i[-Mh, \infty)$, then*

$$\|(P(\alpha) - \lambda)^{-1}\|_{L^2(\mathbb{R}) \rightarrow H_h^2(\mathbb{R})} \leq C \log(1/h) h^{-1-C_0|\text{Im } \lambda|/h}. \tag{5-13}$$

If $\chi \in C^\infty(\mathbb{R})$ has $\chi' \in C_0^\infty(\mathbb{R})$ and $\chi(r) = 0$ for r sufficiently negative, then

$$\|\chi(P(\alpha) - \lambda)^{-1}\chi\|_{L^2(\mathbb{R}) \rightarrow H_h^2(\mathbb{R})} \leq Ch^{-1-2C_0|\text{Im } \lambda|/h} \tag{5-14}$$

in the same range of h, α, λ , and with the same C_0 and h_0 (but with different C).

Lemma 5.3. *Let $r_0 < 0, \chi_- \in C_0^\infty((-\infty, r_0)), \chi_+ \in C_0^\infty((r_0, \infty)), \varphi \in C_0^\infty((-\infty, 0)), E \in (0, 1), \Gamma, \lambda_1, N > 0$ be given. Then there exists $h_0 > 0$ such that*

$$\|\varphi(hD_r)\chi_+(r)(P(\alpha) - \lambda)^{-1}\chi_-(r)\|_{L^2(\mathbb{R}) \rightarrow H_h^2(\mathbb{R})} = \mathcal{O}(h^\infty) \tag{5-15}$$

uniformly for $\alpha \geq \lambda_1 h, \text{Re } \lambda \in [-E, E], -\Gamma h \leq \text{Im } \lambda \leq h^{-N}, h \in (0, h_0]$.

Take $\alpha_0 > 0$ such that if $\alpha \geq \alpha_0$ and $r \leq 0$ then $\alpha^2 e^{-2(r+\beta(r))} \geq 3$. We consider the cases $\lambda_1 h \leq \alpha \leq \alpha_0$ and $\alpha_0 \leq \alpha$ separately.

Proof of (5-13), (5-14), and (5-15) for $\alpha_0 \leq \alpha$. In this case $P(\alpha)$ is ‘‘elliptic’’ (although not pseudodifferential in the usual sense because of the exponentially growing term $\alpha^2 e^{-2(r+\beta(r))}$) and better estimates hold. Use the fact that $W_C \geq 0$ and $\alpha^2 e^{-2(r+\beta(r))} \geq 3$ for $r \leq 0$ to write

$$\begin{aligned} \int_{-\infty}^0 |u|^2 dr &\leq \frac{1}{3} \int_{-\infty}^{\infty} \alpha^2 e^{-2(r+\beta(r))} |u|^2 dr \leq \frac{1}{3} \text{Re} \langle P(\alpha)u, u \rangle_{L^2(\mathbb{R})} + \left(\frac{1}{3} + \mathcal{O}(h^2)\right) \|u\|_{L^2(\mathbb{R})}^2, \\ \int_0^{\infty} |u|^2 dr &= \int_0^{\infty} W_C |u|^2 dr \leq \int_{-\infty}^{\infty} W_C |u|^2 dr = -\text{Im} \langle P(\alpha)u, u \rangle_{L^2(\mathbb{R})}. \end{aligned}$$

Adding the inequalities gives

$$\|u\|_{L^2(\mathbb{R})}^2 \leq 2\|(P(\alpha) - \lambda)u\|_{L^2(\mathbb{R})} \|u\|_{L^2(\mathbb{R})} + \left(\frac{1}{3} \text{Re } \lambda - \text{Im } \lambda + \frac{1}{3} + \mathcal{O}(h^2)\right) \|u\|_{L^2(\mathbb{R})}^2.$$

So long as $\text{Im } \lambda - \frac{1}{3} \text{Re } \lambda + \frac{2}{3} \geq \epsilon$ for some $\epsilon > 0$, it follows that

$$\|u\|_{L^2(\mathbb{R})} \leq C \|(P(\alpha) - \lambda)u\|_{L^2(\mathbb{R})}. \tag{5-16}$$

To obtain (5-13) we observe that

$$\begin{aligned} &\|h^2 D_r^2 u\|_{L^2(\mathbb{R})}^2 \\ &= \|(h^2 D_r^2 + \alpha^2 e^{-2(r+\beta(r))})u\|_{L^2(\mathbb{R})}^2 - \|\alpha^2 e^{-2(r+\beta(r))} u\|_{L^2(\mathbb{R})}^2 - 2 \text{Re} \langle h^2 D_r^2 u, \alpha^2 e^{-2(r+\beta(r))} u \rangle_{L^2(\mathbb{R})}, \end{aligned}$$

while

$$\begin{aligned} &-\text{Re} \langle h^2 D_r^2 u, \alpha^2 e^{-2(r+\beta(r))} u \rangle_{L^2(\mathbb{R})} \\ &= -\|\alpha e^{-(r+\beta(r))} h D_r u\|_{L^2(\mathbb{R})}^2 + 2 \text{Im} \langle h D_r u, (1 + \beta'(r)) h \alpha^2 e^{-2(r+\beta(r))} u \rangle_{L^2(\mathbb{R})}, \end{aligned}$$

so that

$$\|h^2 D_r^2 u\|_{L^2(\mathbb{R})} \leq 2\|(h^2 D_r^2 + \alpha^2 e^{-2(r+\beta(r))})u\|_{L^2(\mathbb{R})} \leq 2\|(P(\alpha) - \lambda)u\|_{L^2(\mathbb{R})} + C|\lambda|\|u\|_{L^2(\mathbb{R})}.$$

Together with (5-16), this implies (5-13) (and hence (5-14)) with the right-hand side replaced by $C(1 + |\lambda|)$. The estimate (5-15) follows from the stronger Agmon estimate

$$\|\chi_+(r)(P(\alpha) - \lambda)^{-1}\chi_-(r)\|_{L^2(\mathbb{R}) \rightarrow H_h^2(\mathbb{R})} = \mathcal{O}(e^{-1/(Ch)});$$

see for example [Zworski 2012, Theorems 7.3 and 7.1]. □

Proof of (5-13) for $\lambda_1 h \leq \alpha \leq \alpha_0$. For this range of α we use the following rescaling (I'm very grateful to Nicolas Burq for suggesting this rescaling):

$$\tilde{r} = r / \log(2\alpha_0/\alpha), \quad \tilde{h} = h / \log(2\alpha_0/\alpha). \tag{5-17}$$

In these variables we have

$$P(\alpha) = (\tilde{h} D_{\tilde{r}})^2 + 4\alpha_0^2 e^{-2[(1+\tilde{r})\log(2\alpha_0/\alpha) + \tilde{\beta}(\tilde{r})]} + \tilde{h}^2 \tilde{V}(\tilde{r}) - 1 - i \tilde{W}_C(\tilde{r}),$$

where

$$\tilde{\beta}(\tilde{r}) = \beta(r), \quad \tilde{V}(\tilde{r}) = \log(2\alpha_0/\alpha)^2 V(r), \quad \tilde{W}_C(\tilde{r}) = W_C(r).$$

We will show that

$$\|(P(\alpha) - \lambda)^{-1}\|_{L_{\tilde{r}}^2 \rightarrow H_{\tilde{h}, \tilde{r}}^2} \leq C \tilde{h}^{-1} e^{C_0 |\operatorname{Im} \lambda| / \tilde{h}} \tag{5-18}$$

for $|\operatorname{Re} \lambda| \leq E$, $\operatorname{Im} \lambda \geq -M \tilde{h} \log(1/\tilde{h})$, from which (5-13) follows.

We now use a variant of the gluing argument in Section 3A to replace the exponentially growing term $4\alpha_0^2 e^{-2[(1+\tilde{r})\log(\alpha_0/\alpha) + \tilde{\beta}(\tilde{r})]}$ with a bounded one. Fix $\tilde{R} > 0$ such that

$$\tilde{r} \leq -\tilde{R}, \quad \alpha \leq \alpha_0 \quad \implies \quad \alpha_0^2 e^{-2[(1+\tilde{r})\log(2\alpha_0/\alpha) + \tilde{\beta}(\tilde{r})]} > 1.$$

Take $\tilde{V}_B, \tilde{V}_E \in C^\infty(\mathbb{R}, [0, \infty))$ such that

$$\tilde{V}_E(\tilde{r}) = 4\alpha_0^2 e^{-2[(1+\tilde{r})\log(2\alpha_0/\alpha) + \tilde{\beta}(\tilde{r})]} \quad \text{for } \tilde{r} \leq -\tilde{R}$$

and $\tilde{V}_E(\tilde{r}) \geq 4$ for all \tilde{r} , while

$$\tilde{V}_B(\tilde{r}) = 4\alpha_0^2 e^{-2[(1+\tilde{r})\log(2\alpha_0/\alpha) + \tilde{\beta}(\tilde{r})]} \quad \text{for } \tilde{r} \geq -\tilde{R} - 3$$

and \tilde{V}_B is decreasing in \tilde{r} and bounded together with all derivatives, uniformly in α (see Figure 6).

Let

$$P_E(\alpha) = (\tilde{h} D_{\tilde{r}})^2 + \tilde{V}_E(\tilde{r}) + \tilde{h}^2 \tilde{V}(\tilde{r}) - 1 - i \tilde{W}_C(\tilde{r}),$$

$$P_B(\alpha) = (\tilde{h} D_{\tilde{r}})^2 + \tilde{V}_B(\tilde{r}) + \tilde{h}^2 \tilde{V}(\tilde{r}) - 1 - i \tilde{W}_C(\tilde{r}),$$

and let $R_E = (P_E(\alpha) - \lambda)^{-1}$, $R_B = (P_B(\alpha) - \lambda)^{-1}$. Note that

$$\|R_E\|_{L_{\tilde{r}}^2 \rightarrow H_{\tilde{h}, \tilde{r}}^2} \leq C$$




Figure 6. The model potentials \tilde{V}_E and \tilde{V}_B . The former agrees with the function $4\alpha_0^2 e^{-2[(1+\tilde{r}) \log(2\alpha_0/\alpha) + \tilde{\beta}(\tilde{r})]}$ for $\tilde{r} \leq -\tilde{R}$, and \tilde{V}_B agrees with the same function for $\tilde{r} \geq -\tilde{R} - 3$.

by the same proof as that of (5-13) for $\alpha \geq \alpha_0$. We will show that (5-18) follows from

$$\|R_B\|_{L_{\tilde{r}}^2 \rightarrow H_{h,\tilde{r}}^2} \leq C\tilde{h}^{-1} e^{C_0|\text{Im } \lambda|/\tilde{h}} \tag{5-19}$$

for $|\text{Re } \lambda| \leq E, \text{Im } \lambda \geq -M\tilde{h} \log(1/\tilde{h})$. Indeed, let $\chi_E \in C^\infty(\mathbb{R}; \mathbb{R})$ have $\chi_E(\tilde{r}) = 1$ near $\tilde{r} \leq -\tilde{R} - 2$ and $\chi_E(\tilde{r}) = 0$ near $\tilde{r} \geq -\tilde{R} - 1$, and let $\chi_B = 1 - \chi_E$. Let

$$G = \chi_E(\tilde{r} - 1)R_E\chi_E(\tilde{r}) + \chi_B(\tilde{r} + 1)R_B\chi_B(\tilde{r}).$$

Then

$$(P(\alpha) - \lambda)G = \text{Id} + [\tilde{h}^2 D_{\tilde{r}}^2, \chi_E(\tilde{r} - 1)]R_E\chi_E(\tilde{r}) + [\tilde{h}^2 D_{\tilde{r}}^2, \chi_B(\tilde{r} + 1)]R_B\chi_B(\tilde{r}) = \text{Id} + A_E + A_B.$$

As in Section 3A we have $A_E^2 = A_B^2 = 0$. We also have the Agmon estimate

$$\|A_E\|_{L_{\tilde{r}}^2 \rightarrow L_{\tilde{r}}^2} \leq e^{-1/(C\tilde{h})};$$

see for example [Zworski 2012, Theorems 7.3 and 7.1]. Solving away A_B using G we find that

$$(P(\alpha) - \lambda)G(\text{Id} - A_B) = \text{Id} + \mathcal{O}_{L_{\tilde{r}}^2 \rightarrow L_{\tilde{r}}^2}(e^{-1/(C\tilde{h})}), \tag{5-20}$$

and since $\|G(\text{Id} - A_B)\|_{L_{\tilde{r}}^2 \rightarrow H_{h,\tilde{r}}^2} \leq C\tilde{h}^{-1} e^{C|\text{Im } \lambda|/\tilde{h}}$, this implies (5-18).

The proof of (5-19) follows that of (5-1) with these differences: the $-i\tilde{W}_C(\tilde{r})$ term removes the need for complex scaling, and the $\tilde{V}_B(\tilde{r})$ term puts P_B in a mildly exotic operator class and leads to a slightly modified escape function q and microlocal cutoff ϕ . Fix

$$E_0 \in (E, 1), \quad \varepsilon = 10M\tilde{h} \log(1/\tilde{h}). \tag{5-21}$$

The \tilde{h} -semiclassical principal symbol of P_B (note that $P_B \in \Psi_\delta^2(\mathbb{R})$ for any $\delta > 0$) is

$$p_B = \tilde{\rho}^2 + \tilde{V}_B(\tilde{r}) - 1 - i\tilde{W}_C(\tilde{r}), \tag{5-22}$$

where $\tilde{\rho}$ is dual to \tilde{r} . Take $q \in C_0^\infty(T^*\mathbb{R})$ such that on $\{-\tilde{R} \leq \tilde{r} \leq 0, |\tilde{\rho}| \leq 2\}$ we have

$$q(\tilde{r}, \tilde{\rho}) = -C_q(\tilde{r} + \tilde{R} + 1)\tilde{\rho},$$

$$\operatorname{Re} H_{p_B} q = -2C_q \tilde{\rho}^2 + C_q(\tilde{r} + \tilde{R} + 1)\tilde{V}'_B(\tilde{r}) \leq -C_q(\operatorname{Re} p_B + 1),$$

where $C_q > 0$ is a large constant which will be specified below, and where for the inequality we used (2-2). Let $Q \in \Psi^{-\infty}(\mathbb{R})$ be a quantization of q with \tilde{h} as semiclassical parameter and put

$$P_{B,\varepsilon} = e^{\varepsilon Q/\tilde{h}} P_B e^{-\varepsilon Q/\tilde{h}} = P_B - \varepsilon[P_B, Q/\tilde{h}] + \varepsilon^2 \tilde{h}^{-4\delta} R, \tag{5-23}$$

where $R \in \Psi_\delta^{-\infty}(\mathbb{R})$ by (2-26). The \tilde{h} -semiclassical principal symbol of $P_{B,\varepsilon}$ is

$$p_{B,\varepsilon} = \tilde{\rho}^2 + V_B(\tilde{r}) - 1 - i\tilde{W}_C(\tilde{r}) + i\varepsilon H_{p_B} q.$$

We will prove

$$\|(P_{B,\varepsilon} - E')^{-1}\|_{L_{\tilde{r}}^2 \rightarrow H_{\tilde{h},\tilde{r}}^2} \leq 5/\varepsilon, \quad E' \in [-E_0, E_0], \tag{5-24}$$

from which it follows by (2-23) that

$$\|(P_{B,\varepsilon} - \lambda)^{-1}\|_{L_{\tilde{r}}^2 \rightarrow H_{\tilde{h},\tilde{r}}^2} \leq \frac{\tilde{h}^{-N}}{M \log(1/\tilde{h})}, \quad |\operatorname{Re} \lambda| \leq E_0, \quad |\operatorname{Im} \lambda| \leq M\tilde{h} \log(1/\tilde{h}), \tag{5-25}$$

where

$$N = 10M(\|Q\|_{H_{\tilde{h},\tilde{r}}^2 \rightarrow H_{\tilde{h},\tilde{r}}^2} + \|Q\|_{L_{\tilde{r}}^2 \rightarrow L_{\tilde{r}}^2}) + 1.$$

The proof that (5-25) implies (5-19) is the same as the proof that (4-3) implies (3-8).

Let $\phi \in C_0^\infty(T^*\mathbb{R})$ be identically 1 near $\{(\tilde{r}, \tilde{\rho}) : -\tilde{R} \leq \tilde{r} \leq 0, |\tilde{\rho}| \leq 2, |\operatorname{Re} p_B(\tilde{r}, \tilde{\rho})| \leq E_0\}$ and be supported such that $\operatorname{Re} H_{p_B} q < 0$ on $\operatorname{supp} \phi$. Let Φ be the quantization of ϕ with \tilde{h} as semiclassical parameter. For h (and hence \tilde{h} and ε) small enough, we have $|p_{B,\varepsilon} - E'| \geq (1 + \tilde{\rho}^2)/C$ on $\operatorname{supp}(1 - \phi)$, uniformly in $E' \in [-E_0, E_0]$, in $\alpha \leq \alpha_0$ and in h . Hence, by the semiclassical elliptic estimate (2-18),

$$\|(\operatorname{Id} - \Phi)u\|_{H_{\tilde{h},\tilde{r}}^2} \leq C\|(P_{B,\varepsilon} - E')(\operatorname{Id} - \Phi)u\|_{L_{\tilde{r}}^2} + \mathcal{O}(h^\infty)\|u\|_{H_{\tilde{h},\tilde{r}}^{-N}}.$$

Using the fact that $\operatorname{Re} H_{p_B} q < 0$ on $\operatorname{supp} \phi$, fix C_q large enough that on $\operatorname{supp} \phi$ we have

$$\operatorname{Im} p_{B,\varepsilon} = -\tilde{W}_C(\tilde{r}) + \varepsilon \operatorname{Re} H_{p_B} q \leq -\varepsilon.$$

Then, using the sharp Gårding inequality (2-19), we have, for h sufficiently small,

$$\begin{aligned} \|\Phi u\|_{L_{\tilde{r}}^2(\mathbb{R})} \|(P_{B,\varepsilon} - E')\Phi u\|_{L_{\tilde{r}}^2(\mathbb{R})} &\geq -\langle \operatorname{Im}(P_{B,\varepsilon} - E')\Phi u, \Phi u \rangle_{L_{\tilde{r}}^2(\mathbb{R})} \\ &\geq \varepsilon \|\Phi u\|_{L_{\tilde{r}}^2(\mathbb{R})}^2 - C\tilde{h}^{1-2\delta} \|u\|_{H_{\tilde{h},\tilde{r}}^{1/2}(\mathbb{R})}^2. \end{aligned}$$

We deduce (5-24) from this just as we did (4-2) above. □

Proof of (5-14) for $\lambda_1 h \leq \alpha \leq \alpha_0$. It suffices to show that

$$\|\chi R_B \chi\|_{L_{\tilde{r}}^2 \rightarrow H_{\tilde{h},\tilde{r}}^2} \leq C/h \tag{5-26}$$

when $|\operatorname{Re} \lambda| \leq E_0$, $\operatorname{Im} \lambda \geq 0$, with R_B as in the proof of (5-13) for $\lambda_1 h \leq \alpha \leq \alpha_0$, E_0 as in (5-21).¹ Then $\|\chi(P(\alpha) - \lambda)^{-1} \chi\|_{L^2_{\tilde{r}} \rightarrow H^2_{\tilde{h},r}} \leq C/h$ (for the same range of parameters) follows by the same argument that reduced (5-13) to (5-19) above. After this, (5-14) follows by complex interpolation as in the proof that (4-3) implies (3-8) above. Indeed, take $f(\lambda, h)$ holomorphic in λ , bounded uniformly for $\lambda \in \Omega = [-E_0, E_0] + i[-Mh \log \log(1/h), 0]$, and satisfying

$$|\operatorname{Re} \lambda| \leq E \implies |f| \geq 1, \quad |\operatorname{Re} \lambda| \leq \left[\frac{1}{2}(E + E_0), E_0\right] \implies |f| \leq h^2$$

for $\lambda \in \Omega$. Then define the subharmonic function

$$g(\lambda, h) = \log \|\chi(P(\alpha) - \lambda)^{-1} \chi\|_{L^2_{\tilde{r}} \rightarrow H^2_{\tilde{h},r}} + \log |f(\lambda, h)| + 2C_0 \frac{\operatorname{Im} \lambda}{h} \log(1/h),$$

and apply the maximum principle to g on Ω , observing that $g \leq C + \log(1/h)$ on $\partial\Omega$.

It now remains to prove (5-26), which we do using a “noncompact” variant of the positive commutator method of [Datchev and Vasy 2012b]. Fix $-R_0 < \inf \operatorname{supp} \chi$ and take $f \in L^2_r$ with $\operatorname{supp} f \subset (-R_0, \infty)$. Let $u = R_B f$. We will show that $\|\chi u\|_{H^2_{\tilde{h},r}} \leq C \|f\|_{L^2_r}/h$.

As an escape function take $q \in S^0(\mathbb{R})$ with $q \geq 0$ everywhere and such that

$$q(r, \rho) = \begin{cases} 1 + 2R_0 e^{-1/R_0}, & -R_0 \geq r, \\ 1 + 2R_0 e^{-1/R_0} - \rho(r + R_0 + 1)e^{-1/(r+R_0)}, & -R_0 < r \leq 0 \text{ and } |\rho| \leq 2. \end{cases}$$

We do not prescribe additional conditions on q outside of this range of (r, ρ) , as P_B is semiclassically elliptic there. The h -semiclassical principal symbol of P_B is (see (5-22))

$$p_B = \rho^2 + V_B(r) - 1 - iW_C(r),$$

where $V_B(r) = \tilde{V}_B(\tilde{r})$. Making $-\tilde{R}$ more negative if necessary, we may suppose without loss of generality that

$$r \geq -R_0 \implies V_B(r) = \alpha^2 e^{-2(r+\beta(r))}.$$

For $r \leq -R_0$ we have $H_{p_B} q = 0$, and for $-R_0 < r \leq 0$, $|\rho| \leq 2$ we have

$$\begin{aligned} \operatorname{Re} H_{p_B} q(r, \rho) &= [-2\rho^2(1 + 1/(r + R_0)) + V'_B(r)(r + R_0 + 1)]e^{-1/(r+R_0)} \\ &\leq -(\operatorname{Re} p_B + 1)e^{-1/(r+R_0)}. \end{aligned}$$

Consequently, we may write

$$\operatorname{Re} H_{p_B}(q^2) = -b^2 + a,$$

where $a, b \in C^\infty_0(T^*\mathbb{R})$ and $\operatorname{supp} a$ is disjoint from $\{r \leq -R_0\}$ and from $\{-R_0 < r \leq 0\} \cap \{|\rho| \leq 2\}$. Note that

$$b \neq 0 \quad \text{on } \{|p_B| \leq E_0\} \cap T^*(-R_0, 0). \tag{5-27}$$

Let $Q = \operatorname{Op}(q)$ as in (2-15). Then

$$i[P_B, Q^* Q] = -hB^* B + hA + [W_C, Q^* Q] + h^2 Y, \tag{5-28}$$

¹Note that for this proof we do not use the variables \tilde{r} and \tilde{h} .

where $B, A, Y \in \Psi^{-\infty}(\mathbb{R})$ and B, A have semiclassical principal symbols b, a . Note that if $\chi_0 \in C_0^\infty((-R_0, \infty))$, then by (5-27) and (2-18) we have

$$\|\chi_0 u\|_{H_{h,r}^2}^2 \leq C(\|Bu\|_{L_r^2}^2 + \log^2(1/h)\|f\|_{L_r^2}^2), \tag{5-29}$$

so it suffices to show that

$$\|Bu\|_{L_r^2}^2 \leq Ch^{-2}\|f\|_{L_r^2}^2. \tag{5-30}$$

Combining (5-28) with

$$\langle i[P_B, Q^*Q]u, u \rangle_{L_r^2} = -2 \operatorname{Im} \langle Q^*Qu, f \rangle_{L_r^2} + 2 \langle W_C Q^*Qu, u \rangle_{L_r^2} + 2 \operatorname{Im} \lambda \|Qu\|_{L_r^2}^2$$

gives

$$\begin{aligned} \|Bu\|_{L_r^2}^2 &= \langle Au, u \rangle_{L_r^2} + \frac{2}{h} \operatorname{Im} \langle Q^*Qu, f \rangle_{L_r^2} - \frac{1}{h} \langle (W_C Q^*Q + Q^*QW_C)u, u \rangle_{L_r^2} \\ &\quad - \frac{2 \operatorname{Im} \lambda}{h} \|Qu\|_{L_r^2}^2 + h \langle Yu, u \rangle_{L_r^2}. \end{aligned} \tag{5-31}$$

We now estimate the right-hand side term by term to obtain (5-30). Since $P_B - \lambda$ is semiclassically elliptic on $\operatorname{supp} a$, by (2-18) followed by (5-13) we have

$$|\langle Au, u \rangle_{L_r^2}| \leq C\|f\|_{L_r^2}^2 + Ch^2\|u\|_{L_r^2}^2 \leq C \log^2(1/h)\|f\|_{L_r^2}^2.$$

For any $\epsilon > 0$ and $\chi_1 \in C_0^\infty(\mathbb{R})$ with $\chi_1 = 1$ near $\operatorname{supp} f$ we have

$$\frac{2}{h} \operatorname{Im} \langle Q^*Qu, f \rangle_{L_r^2} \leq \epsilon \|\chi_1 u\|_{L_r^2}^2 + \frac{C}{h^2 \epsilon} \|f\|_{L_r^2}^2.$$

By (5-27) and the elliptic estimate (2-18), if further $\inf \operatorname{supp} \chi_1 > -R_0$, then (5-29) gives

$$\frac{2}{h} \operatorname{Im} \langle Q^*Qu, f \rangle_{L_r^2} \leq C\epsilon \|Bu\|_{L_r^2}^2 + \frac{C}{h^2 \epsilon} \|f\|_{L_r^2}^2.$$

Next we have, using $W_C \geq 0$ and the fact that $h^{-1}[W_C, Q^*]Q$ has imaginary principal symbol, followed by (5-13),

$$\begin{aligned} -\frac{1}{h} \langle (W_C Q^*Q + Q^*QW_C)u, u \rangle_{L_r^2} &= -\frac{2}{h} \langle W_C Qu, Qu \rangle_{L_r^2} + \frac{2}{h} \operatorname{Re} \langle [W_C, Q^*]Qu, u \rangle_{L_r^2} \\ &\leq Ch\|u\|_{L_r^2}^2 \leq C \frac{\log^2(1/h)}{h} \|f\|_{L_r^2}^2. \end{aligned}$$

Finally we observe that $-2 \operatorname{Im} \lambda \|Qu\|_{L_r^2}^2 / h \leq 0$ since $\operatorname{Im} \lambda \geq 0$, while (5-13) implies

$$h \langle Yu, u \rangle_{L_r^2} \leq C \frac{\log^2(1/h)}{h} \|f\|_{L_r^2}^2.$$

This completes the estimation of (5-31) term by term, giving (5-30). □

Proof of (5-15) for $\lambda_1 h \leq \alpha \leq \alpha_0$. We begin this proof with the same rescaling to \tilde{r} and \tilde{h} , and the same parametrix construction as for the proof of (5-13) for $\lambda_1 h \leq \alpha \leq \alpha_0$ above, but with the additional requirement that

$$-\tilde{R} \leq r_0 / \log 2.$$

Then if we put

$$\tilde{\chi}_+(\tilde{r}) = \chi_+(r), \quad \tilde{\chi}_-(\tilde{r}) = \chi_-(r),$$

we have

$$\text{supp } \tilde{\chi}_+ \subset (r_0 / \log(2\alpha_0/\alpha), \infty) \subset (r_0 / \log 2, \infty), \quad \text{supp } \chi_E \subset (-\infty, -\tilde{R} - 1),$$

and hence

$$\tilde{\chi}_+(\tilde{r})\chi_E(\tilde{r} - 1) = 0. \tag{5-32}$$

Then, noting that (5-20) implies

$$(P(\alpha) - \lambda)^{-1} = G(\text{Id} - A_B)(\text{Id} + \mathcal{O}_{L_{\tilde{r}}^2 \rightarrow L_{\tilde{r}}^2}(e^{-1/(C\tilde{h})})),$$

we use (5-32) to write

$$\tilde{\chi}_+(\tilde{r})(P(\alpha) - \lambda)^{-1}\tilde{\chi}_-(\tilde{r}) = \tilde{\chi}_+(\tilde{r})R_B\tilde{\chi}_-(\tilde{r}) + \mathcal{O}_{L_{\tilde{r}}^2 \rightarrow H_{\tilde{h},\tilde{r}}^2}(e^{-1/(C\tilde{h})}).$$

Returning to the r and h variables, we see that it suffices to show that

$$\|\varphi(hD_r)\chi_+(r)R_B\chi_-(r)\|_{L_r^2 \rightarrow H_{h,r}^2} = \mathcal{O}(h^\infty). \tag{5-33}$$

The proof of (5-33) is almost the same as that of (5-2). There are two differences.

The first difference is that as an escape function we use

$$q = \varphi_r(r)\varphi_\rho(\rho), \quad \text{Re } H_{p_B}q = 2\rho\varphi'_r(r)\varphi_\rho(\rho) - V'_C(r)\varphi'_r(r)\varphi'_\rho(\rho),$$

where $\varphi_r \in C_0^\infty(\mathbb{R}; [0, \infty))$ with $\text{supp } \varphi_r \subset (r_0, \infty)$, $\varphi'_r \geq 0$ near $[r_0, 0]$, $\varphi'_r > 0$ near $\text{supp } \chi_+$. Take $\varphi_\rho \in C_0^\infty(\mathbb{R}; [0, \infty))$ with $\text{supp } \varphi_\rho \subset (-\infty, 0)$, $\varphi'_\rho \leq 0$ near $[-2, 0]$, $\varphi_\rho \neq 0$ near $\text{supp } \varphi \cap [-2, 0]$. Impose further that $\sqrt{\varphi_r}, \sqrt{\varphi_\rho} \in C_0^\infty(\mathbb{R})$, and that $\varphi'_r \geq c\varphi_r$ for $r \leq 0$, where $c > 0$ is chosen large enough that $\text{Re } H_{p_B}q \leq -(2\Gamma + 1)q$ on $\{r \leq 0, \rho \geq -2\}$.

The second difference is that the complex absorbing barrier W_C produces a remainder term in the positive commutator estimate, analogous to the one in the proof of (5-14) for $\lambda_1 h \leq \alpha \leq \alpha_0$ above. The same argument removes the remainder term in this case. \square

6. Model operator in the funnel

In this section we prove Propositions 3.6 and 3.7. As in Section 5, we begin by separating variables over the eigenspaces of Δ_{S_+} , writing

$$P_F = \bigoplus_{m=0}^{\infty} h^2 D_r^2 + (1 - W_F(r))(h\lambda_m)^2 e^{-2(r+\beta(r))} + h^2 V(r) - 1 - iW_F(r),$$

where $0 = \lambda_0 < \lambda_1 \leq \dots$ are square roots of the eigenvalues of Δ_{S^+} . Roughly speaking, it suffices to prove (3-13), (3-14) with P_F replaced by $P(\alpha)$, with estimates uniform in $\alpha \geq 0$, where

$$P(\alpha) = h^2 D_r^2 + (1 - W_F(r))\alpha^2 e^{-2(r+\beta(r))} + h^2 V(r) - 1 - i W_F(r).$$

More specifically, with notation as in those two propositions, (3-13) follows from

$$\|\chi(P(\alpha) - \lambda)^{-1} \chi\|_{L^2(\mathbb{R}) \rightarrow H_h^2(\mathbb{R})} \leq C \begin{cases} h^{-1} + |\lambda|, & \text{Im } \lambda > 0, \\ h^{-1} e^{C_0 |\text{Im } \lambda|/h}, & \text{Im } \lambda \leq 0, \end{cases} \tag{6-1}$$

and (3-14) follows from

$$\|\chi_+(r)(P(\alpha) - \lambda)^{-1} \chi_-(r)\varphi(h D_r)\|_{L^2(\mathbb{R}) \rightarrow H_h^2(\mathbb{R})} = \mathcal{O}(h^\infty), \tag{6-2}$$

so in this section we will prove (6-1) and (6-2).

To do that we use a variant of the method of complex scaling presented in the proof of Lemma 5.1, but with contours γ depending on α in such a way as to give estimates uniform in α ; the α -dependence is needed because the term $\alpha^2(1 - W_F(r))e^{-2(r+\beta(r))}$, although exponentially decaying, is not uniformly exponentially decaying as $\alpha \rightarrow \infty$. Such contours were first used in [Zworski 1999, §4]; here we present a simplified approach based on that in [Datchev 2010, §5.2].

Fix $R > R_g$ sufficiently large that

$$\text{supp } \chi \cup \text{supp } \chi_+ \cup \text{supp } \chi_- \subset (-\infty, R)$$

and that

$$\text{Re } z \geq R, \quad 0 \leq \arg z \leq \theta_0 \quad \implies \quad |\text{Im } \beta(z)| \leq \frac{1}{2} |\text{Im } z|, \tag{6-3}$$

where θ_0 is as in Section 2A. Let $\gamma = \gamma_\alpha(r)$ be real-valued, smooth in r with $\gamma'(r) \geq 0$ for all r , and obey $\gamma(r) = 0$ for $r \leq R$ (here and below $\gamma' = \partial_r \gamma$). Suppose $\gamma'' \in C_0^\infty(\mathbb{R})$ for each α , but not necessarily uniformly in α . Now put

$$P_\gamma(\alpha) = \frac{h^2 D_r^2}{(1 + i\gamma'(r))^2} - h \frac{\gamma''(r)h D_r}{(1 + i\gamma'(r))^3} + \alpha^2(1 - W_F(r))e^{-2(r+i\gamma(r)+\beta(r+i\gamma(r)))} + h^2 V(r + i\gamma(r)) - 1 - i W_F(r).$$

If we define the differential operator with complex coefficients

$$\tilde{P}(\alpha) = h^2 D_z^2 + \alpha^2(1 - W_F(z))e^{-2(z+\beta(z))} + h^2 V(z) - 1 - i W_F(z),$$

where z varies in $\{z = r + i\delta\gamma(r) : r \in \mathbb{R}, \delta \in (0, 1)\}$, and where $W_F(z) := 0$ whenever $\text{Im } z \neq 0$, then we have

$$P(\alpha) = \tilde{P}(\alpha)|_{\{z=r:r \in \mathbb{R}\}}, \quad P_\gamma(\alpha) = \tilde{P}(\alpha)|_{\{z=r+i\gamma(r):r \in \mathbb{R}\}}.$$

If $\chi_0 \in C^\infty(\mathbb{R})$ has $\text{supp } \chi_0 \cap \text{supp } \gamma = \emptyset$, then

$$\chi_0(P(\alpha) - \lambda)^{-1} \chi_0 = \chi_0(P_\gamma(\alpha) - \lambda)^{-1} \chi_0, \quad \text{Im } \lambda > 0,$$

by an argument almost identical to that used to prove (5-5); the only difference is we construct WKB solutions which are exponentially growing and decaying as $\text{Re } z \rightarrow +\infty$ rather than $-\infty$, and we take $f(z) = (\alpha^2 e^{-2(z+\beta(z))} + h^2 V(z) - 1 - \lambda)/h^2$.

Consequently, to prove (6-1) and (6-2), it is enough to show that

$$\|(P_\gamma(\alpha) - \lambda)^{-1}\|_{L^2(\mathbb{R}) \rightarrow H_h^2(\mathbb{R})} \leq C e^{C_0 |\text{Im } \lambda|/h} \tag{6-4}$$

and

$$\|\chi_+(r)(P_\gamma(\alpha) - \lambda)^{-1}\chi_-(r)\varphi(hD_r)\|_{L^2(\mathbb{R}) \rightarrow H_h^2(\mathbb{R})} = \mathcal{O}(h^\infty) \tag{6-5}$$

for a suitably chosen γ , with estimates uniform in $\alpha \geq 0$.

Fix $R_- > R$ such that

$$|\text{Im } \beta(z)| \leq \frac{1}{2} \text{Im } z \tag{6-6}$$

for $\text{Re } z \geq R_-$, $0 \leq \arg z \leq \theta_0$, with θ_0 as in Section 2A. Take $\alpha_0 > 0$ such that

$$\alpha_0^2 e^{-2(R+1)} e^{-2 \max |\text{Re } \beta|} = 8, \tag{6-7}$$

where $\max |\text{Re } \beta|$ is taken over $\mathbb{R} \cup \{|z| > R_g, 0 \leq \arg z \leq \theta_0\}$. We consider the cases $\alpha \leq \alpha_0$ and $\alpha \geq \alpha_0$ separately.

Proof of (6-4) for $0 \leq \alpha \leq \alpha_0$. Fix

$$E_0 \in (E, 1), \quad \varepsilon = 10Mh \log(1/h).$$

We use the same complex scaling as in the proof of Lemma 5.1. In this range γ is independent of α and we put $\gamma = \delta\gamma_-$, where $0 < \delta \ll 1$ will be specified later, and we require $\gamma_-(r) = 0$ for $r \leq R_-$, $\gamma'_-(r) \geq 0$ for all r , and $\gamma'_-(r) = \tan \theta_0$ for $r \geq R_- + 1$.

The semiclassical principal symbol of $P_\gamma(\alpha)$ is

$$\begin{aligned} p_\gamma(\alpha) &= \frac{\rho^2}{(1 + i\gamma'(r))^2} + \alpha^2(1 - W_F(r))e^{-2(r+i\gamma(r)+\beta(r+i\gamma(r)))} - 1 - iW_F(r) \\ &= \rho^2 + \alpha^2(1 - W_F(r))e^{-2(r+\beta(r))} - 1 - iW_F(r) + \mathcal{O}(\delta), \end{aligned}$$

where the implicit constant in \mathcal{O} is uniform in compact subsets of $T^*\mathbb{R}$. Moreover,

$$\text{Re } p_\gamma(\alpha) + 1 \geq \rho^2 - \mathcal{O}(\delta),$$

and, using (6-6),

$$\begin{aligned} \text{Im } p_\gamma(\alpha) &\leq -\alpha^2(1 - W_F(r))e^{-2(r+\text{Re } \beta(r+i\gamma(r)))} \sin(2(\gamma(r) + \text{Im } \beta(r + i\gamma(r)))) \\ &\leq -\alpha^2(1 - W_F(r))e^{-2(r+\text{Re } \beta(r+i\gamma(r)))} \sin \gamma(r) \\ &= -\alpha^2(1 - W_F(r))e^{-2(r+\text{Re } \beta(r+i\gamma(r)))} \gamma(r)(1 + \mathcal{O}(\delta^2)), \end{aligned} \tag{6-8}$$

again uniformly on compact subsets of $T^*\mathbb{R}$. Take $q \in C_0^\infty(T^*\mathbb{R})$ such that on $\{0 \leq r \leq R_- + 1, |\rho| \leq 2\}$ we have

$$\begin{aligned} q &= -C_q(r + 1)\rho, \\ \frac{\operatorname{Re} H_{p_\gamma} q}{C_q} &= -2\rho^2 - (W'_F(r) + 2(1 + \beta'(r))(r + 1)\alpha^2 e^{-2(r+\beta(r))}) + \mathcal{O}(\delta) \\ &\leq -(\operatorname{Re} p_\gamma + 1) \leq -\rho^2 + \mathcal{O}(\delta), \end{aligned}$$

where $C_q > 0$ will be specified later, and provided δ is sufficiently small. Let $Q = \operatorname{Op}(q)$ and put

$$P_{\gamma,\varepsilon}(\alpha) = e^{\varepsilon Q/h} P_\gamma(\alpha) e^{-\varepsilon Q/h} = P_\gamma(\alpha) - \varepsilon [P_\gamma(\alpha), Q/h] + \varepsilon^2 R,$$

where $R \in \Psi^{-\infty}(\mathbb{R})$ (see (2-26)). As in the proof of Lemma 5.1, (6-4) follows from

$$\|(P_{\gamma,\varepsilon}(\alpha) - E')^{-1}\|_{L^2(\mathbb{R}) \rightarrow H_h^2(\mathbb{R})} \leq 5/\varepsilon \tag{6-9}$$

for $E' \in [-E_0, E_0]$.

The proof of (6-9) combines elements of the proofs of (5-9) and (5-24). Let $\phi \in C_0^\infty(T^*\mathbb{R})$ be identically 1 near $\{0 \leq r \leq R_- + 1, |\rho| \leq 2, |\operatorname{Re} p_\gamma| \leq E_0\}$ and be supported such that $\operatorname{Re} H_{p_\gamma} q < 0$ on $\operatorname{supp} \phi$. Let Φ be the quantization of ϕ . For δ small enough, and h (and hence ε) small enough depending on δ , we have $|p_{\gamma,\varepsilon} - E'| \geq \delta(1 + \rho^2)/C$ on $\operatorname{supp}(1 - \phi)$, uniformly in $E' \in [-E_0, E_0]$, in $\alpha \leq \alpha_0$ and in h , where $p_{\gamma,\varepsilon}(\alpha)$ is the semiclassical principal symbol of $P_{\gamma,\varepsilon}(\alpha)$. Hence, by the semiclassical elliptic estimate (2-18),

$$\|(\operatorname{Id} - \Phi)u\|_{H_h^2(\mathbb{R})} \leq C\delta^{-1} \|(P_{\gamma,\varepsilon} - E')(\operatorname{Id} - \Phi)u\|_{L^2(\mathbb{R})} + \mathcal{O}(h^\infty) \|u\|_{H_h^{-N}(\mathbb{R})}.$$

Using (6-8) and $\operatorname{supp} \phi \subset \{\operatorname{Re} H_{p_c} q < 0\}$, fix C_q large enough that on $\operatorname{supp} \phi$ we have

$$\operatorname{Im} p_{\gamma,\varepsilon} = \operatorname{Im} p_\gamma + \varepsilon \operatorname{Re} H_{p_c} q \leq -\alpha^2(1 - W_F)e^{-2(r+\operatorname{Re} \beta)} \gamma(1 + \mathcal{O}(\delta^2)) + \varepsilon \operatorname{Re} H_{p_c} q \leq -\varepsilon.$$

Then, using the sharp Gårding inequality (2-19), we have, for h sufficiently small,

$$\begin{aligned} \|\Phi u\|_{L^2(\mathbb{R})} \|(P_{C,\varepsilon} - E')\Phi u\|_{L^2(\mathbb{R})} &\geq -\langle \operatorname{Im}(P_{C,\varepsilon} - E')\Phi u, \Phi u \rangle_{L^2(\mathbb{R})} \\ &\geq \varepsilon \|\Phi u\|_{L^2(\mathbb{R})}^2 - Ch \|u\|_{L^2(\mathbb{R})}^2. \end{aligned}$$

This implies (6-9) just as in the proofs of (5-9) and (5-24). □

Proof of (6-4) for $\alpha \geq \alpha_0$. Define contours $\gamma = \gamma_\alpha(r)$ as follows. Take R_α such that

$$\alpha^2 e^{-2R_\alpha} e^{2 \max |\operatorname{Re} \beta|} = \min\left\{\frac{1}{4}, \frac{1}{2} \tan \theta_0\right\}, \tag{6-10}$$

where $\max |\operatorname{Re} \beta|$ is taken over $\mathbb{R} \cup \{|z| > R_g, 0 \leq \arg z \leq \theta_0\}$. Note that $R_\alpha > R + 1$ by (6-7). Take γ smooth and supported in (R, ∞) , with $0 \leq \gamma'(r) \leq \frac{1}{2}$, and such that

$$\begin{cases} \gamma(r) \leq \frac{\pi}{9}, & r \leq R + 1, \\ \frac{\pi}{18} \leq \gamma(r) \leq \frac{\pi}{6}, & R + 1 \leq r \leq R_\alpha, \\ \gamma'(r) = \min\left\{\frac{1}{2}, \tan \theta_0\right\}, & r \geq R_\alpha. \end{cases}$$

We prove that

$$|p_\gamma(\alpha) - E'| \geq (1 + \rho^2)/C \tag{6-11}$$

uniformly for $-E \leq E' \leq E$ and $\alpha \geq \alpha_0$, by considering each range of r individually. By (2-18) this implies (6-4) for $\alpha \geq \alpha_0$.

(1) For $r \leq R + 1$ we have

$$\begin{aligned} \operatorname{Re} p_\gamma(\alpha) + 1 &= \frac{\rho^2(1 - \gamma'(r)^2)}{|1 + i\gamma'(r)|^4} + \alpha^2(1 - W_F(r)) \operatorname{Re} e^{-2(r+i\gamma(r)+\beta(r+i\gamma(r)))} \\ &\geq \frac{1}{3}\rho^2 + \alpha^2(1 - W_F(r))e^{-2(r+\operatorname{Re} \beta(r+i\gamma(r)))} \cos(3\gamma(r)) \\ &\geq \frac{1}{3}\rho^2 + 4(1 - W_F(r)), \end{aligned} \tag{6-12}$$

where for the first inequality we used $\gamma' \leq \frac{1}{2}$ and (6-6), and for the second (6-7) and $\gamma \leq \frac{\pi}{9}$. Since $\operatorname{Im} p_\gamma = -W_F$ whenever $W_F \neq 0$, this gives (6-11) for $r \leq R + 1$.

(2) For $R + 1 \leq r \leq R_\alpha$ we have $\operatorname{Re} p_\gamma(\alpha) \geq \frac{1}{3}\rho^2 - 1$ by the same argument as in (6-12). This gives (6-11) for $R + 1 \leq r \leq R_\alpha$ once we note that (6-6) and (6-10) imply

$$\begin{aligned} -\operatorname{Im} p_\gamma(\alpha) &= \frac{2\rho^2\gamma'(r)}{|1 + i\gamma'(r)|^4} - \alpha^2 \operatorname{Im} e^{-2(r+i\gamma(r)+\beta(r+i\gamma(r)))} \\ &\geq e^{-2 \max|\operatorname{Re} \beta|} \sin\left(\frac{\pi}{18}\right) \min\left\{\frac{1}{2}, \frac{1}{2} \tan \theta_0\right\}. \end{aligned}$$

(3) For $r \geq R_\alpha$, note that $\alpha^2|e^{-2(r+i\gamma(r)+\beta(r+i\gamma(r)))}| \leq \gamma'(r)$. We again deduce (6-11) by considering two ranges of ρ individually. When $\rho^2/|1 + i\gamma'(r)|^4 \leq \frac{1}{2}$ we have

$$\begin{aligned} \operatorname{Re} p_\gamma(\alpha) &= \frac{\rho^2(1 - \gamma'(r)^2)}{|1 + i\gamma'(r)|^4} + \alpha^2 \operatorname{Re} e^{-2(r+i\gamma(r)+\beta(r+i\gamma(r)))} - 1 \\ &\leq \frac{1}{2} + \frac{1}{4} - 1 = -\frac{1}{4}. \end{aligned}$$

When $\rho^2/|1 + i\gamma'(r)|^4 \geq \frac{1}{2}$ we have

$$\begin{aligned} \operatorname{Im} p_\gamma(\alpha) &= \frac{-2\rho^2\gamma'(r)}{|1 + i\gamma'(r)|^4} + \alpha^2 \operatorname{Im} e^{-2(r+i\gamma(r)+\beta(r+i\gamma(r)))} \\ &\leq \frac{-2\rho^2\gamma'(r)}{|1 + i\gamma'(r)|^4} + \frac{1}{2}\gamma'(r) \leq -\frac{3}{2}\gamma'(r) = -\min\left\{\frac{3}{4}, \frac{3}{2} \tan \theta_0\right\}. \quad \square \end{aligned}$$

For $\alpha \geq \alpha_0$, (6-5) follows from an Agmon estimate just as in the proof of (5-15) for $\alpha \geq \alpha_0$ above. For $\alpha \leq \alpha_0$, (6-5) follows from the same positive commutator argument as was used for the proof of (5-33).

7. Applications

In this section we give applications of the Theorem to solutions to Schrödinger and wave equations. Since such applications are well-known, we only sketch the arguments below, giving references to sources with further details.

We use the notation

$$\|u\|_s := \|(1 + \Delta)^{s/2}u\|_{L^2(X)}, \quad \|A\|_{s \rightarrow s'} := \sup_{\|u\|_s=1} \|Au\|_{s'}, \quad s, s' \in \mathbb{R}.$$

We begin by using (1-1) to deduce polynomial bounds on the resolvent between Sobolev spaces. If $\chi, \tilde{\chi} \in C_0^\infty(X)$ satisfy $\tilde{\chi}\chi = \chi$, then for any $s \in \mathbb{R}$, we have

$$\|\Delta\chi u\|_s \leq C(\|\tilde{\chi}u\|_s + \|\tilde{\chi}\Delta u\|_s).$$

Hence, for any $s, s' \in \mathbb{R}$, we have, letting $R_\chi(\sigma) := \chi(\Delta - \frac{1}{4}n^2 - \sigma^2)^{-1}\chi$,

$$\begin{aligned} \|R_\chi(\sigma)\|_{s \rightarrow s} &\leq C\|R_{\tilde{\chi}}(\sigma)\|_{s' \rightarrow s'}, \\ \|R_\chi(\sigma)\|_{s \rightarrow s'+2} &\leq C(1 + |\sigma|^2)(\|R_{\tilde{\chi}}(\sigma)\|_{s \rightarrow s} + \|R_{\tilde{\chi}}(\sigma)\|_{s \rightarrow s'}), \\ \|R_\chi(\sigma)\|_{s \rightarrow s'} &\leq C(1 + |\sigma|^2)^{-1}(\|R_{\tilde{\chi}}(\sigma)\|_{s \rightarrow s'+2} + \|R_{\tilde{\chi}}(\sigma)\|_{s \rightarrow s'}). \end{aligned}$$

Consequently, (1-1) implies that for any $\chi \in C_0^\infty(X)$, there is $M_0 > 0$ such that for any $M_1 > 0, s \in \mathbb{R}, s' \leq s + 2$, there is $M_2 > 0$ such that

$$\|R_\chi(\sigma)\|_{s \rightarrow s'} \leq M_2|\sigma|^{M_0|\text{Im}\sigma|+s'-s-1} \tag{7-1}$$

when $|\text{Re}\sigma| \geq M_2, \text{Im}\sigma \geq -M_1$.

7A. Local smoothing. By the self-adjoint functional calculus of Δ , the Schrödinger propagator is unitary on all Sobolev spaces: for any $s, t \in \mathbb{R}$, if $u \in H^s(X)$,

$$\|e^{-it\Delta}u\|_s = \|u\|_s.$$

The Kato local smoothing effect says that if we localize in space and average in time, then Sobolev regularity improves by half a derivative: for any $\chi \in C_0^\infty(X), T > 0, s \in \mathbb{R}$ there is $C > 0$ such that if $u \in H^s(X)$,

$$\int_0^T \|\chi e^{-it\Delta}u\|_{s+1/2}^2 dt \leq C\|u\|_s^2. \tag{7-2}$$

This follows by a TT^* argument from (7-1) applied with $\text{Im}\sigma = s = 0, s' = 1$ (see, e.g., [Burq 2004, p. 424]); note that in this case the right-hand side of (7-1) is independent of σ .

7B. Resonant wave expansions. Suppose $\chi(\Delta - \frac{1}{4}n^2 - \sigma^2)^{-1}\chi$ is meromorphic for $\sigma \in \mathbb{C}$. For example we may take (X, g) as in Section 2D1. More generally, if the funnel end is evenly asymptotically hyperbolic as in [Guillarmou 2005, Definition 1.2] then this follows as in the proof of Theorem 1.1 in [Sjöstrand and Zworski 1991, p. 747], but in the interest of brevity we do not pursue this here.

Then (7-1) implies that, when the initial data is compactly supported, solutions to the wave equation $(\partial_t^2 + \Delta - \frac{1}{4}n^2)u = 0$ can be expanded into a superposition of eigenstates and resonant states, with a remainder which decays exponentially on compact sets:

Let $\chi \in C_0^\infty(X)$. There is $M_0 > 0$ such that for any $s \in \mathbb{R}$, $f \in H^{s+1}(X)$, $g \in H^s(X)$ satisfying $\chi f = f$, $\chi g = g$, and for any $M_1 > 0$ and

$$s' < s - M_0 M_1, \tag{7-3}$$

there are $C, T > 0$ such that if $t \geq T$, $H = \sqrt{\Delta - \frac{1}{4}n^2}$, then

$$\left\| \chi \left(\cos(tH) f + \frac{\sin(tH)}{H} g - \sum_{\text{Im } \sigma_j > -M_1} \sum_{m=1}^{M(\sigma_j)} e^{-i\sigma_j t} t^{m-1} w_{j,m} \right) \right\|_{s'} \leq C e^{-M_1 t},$$

where the sum is taken over poles of $R_\chi(\sigma)$ (and is finite by the [Theorem](#)), $M(\sigma_j)$ is the rank of the residue of the pole at σ_j , and each $w_{j,m}$ is a linear combination of the projections of f and g onto the m -th eigenstate or resonant state at σ_j . This follows from (7-1) by an argument of [[Lax and Phillips 1989](#); [Vainberg 1989](#)]; see also [[Tang and Zworski 2000](#), Theorem 3.3] or [[Datchev and Vasy 2012a](#), Corollary 6.1].

Remark. The local smoothing estimate (7-2) is lossless in the sense that the result is the same if (X, g) is nontrapping and asymptotically Euclidean or hyperbolic (see [[Cardoso, Popov and Vodev 2004](#), (1.6)] for a general result). This is because the resolvent estimates (1-1) and (1-2) agree when $\text{Im } \sigma = 0$. The resonant wave expansion exhibits a loss in the Sobolev spaces in which the remainder is controlled: the improvement from (1-1) to (1-2) for $\text{Im } \sigma < 0$ means that, when (1-2) holds, we can replace (7-3) with $s' < s$.

8. Lower bounds

In this section we prove that, in the setting of an exact quotient, the holomorphic continuation of the resolvent grows polynomially. As in [[Borthwick 2007](#), §5.3], we use the fact that in this case the integral kernel of the resolvent can be written in terms of modified Bessel functions.

Proposition 8.1. *Let (X, g) be given by*

$$X = \mathbb{R} \times S, \quad g = dr^2 + e^{2r} dS,$$

where (S, dS) is a compact Riemannian manifold without boundary of dimension n . Then for any $\chi \in C_0^\infty(X)$ which is not identically 0, the cutoff resolvent $\chi(\Delta - \frac{1}{4}n^2 - \sigma^2)^{-1} \chi$ continues holomorphically from $\{\text{Im } \sigma > 0\}$ to $\mathbb{C} \setminus 0$, with a simple pole of rank 1 at $\sigma = 0$.

Moreover, if $\chi \neq 0$ in a neighborhood of $\{r = 0\}$, for any $\varepsilon > 0$ there exists $C > 0$ such that

$$\left\| \chi(\Delta - \frac{1}{4}n^2 - \sigma^2)^{-1} \chi \right\|_{L^2(X) \rightarrow L^2(X)} \geq e^{-C|\text{Im } \sigma|} |\sigma|^{2|\text{Im } \sigma| - 1} / C \tag{8-1}$$

when $\text{Im } \sigma \leq -\varepsilon$, $\text{Re } \sigma \geq C$, $|\text{Im } \sigma| \leq C|\text{Re } \sigma|^{2/3}$.

Proof. As in [Section 2C](#) a conjugation and separation of variables reduce this to the study of the following family of ordinary differential operators:

$$P_m = D_r^2 + \lambda_m^2 e^{-2r},$$

where $0 = \lambda_0 < \lambda_1 \leq \lambda_2 \leq \dots$ are square roots of the eigenvalues of Δ . We will show that $\chi(P_m - \sigma^2)^{-1}\chi$ is entire in σ for $m > 0$, and that it is holomorphic in $\mathbb{C} \setminus 0$ with a simple pole of rank 1 at $\sigma = 0$ for $m = 0$. We will further show that

$$\|\chi(P_1 - \sigma^2)^{-1}\chi\|_{L^2(\mathbb{R}) \rightarrow L^2(\mathbb{R})} \geq e^{-C|\operatorname{Im}\sigma|}|\sigma|^{2|\operatorname{Im}\sigma|-1}/C \tag{8-2}$$

when $\operatorname{Im}\sigma \leq -\varepsilon$, $\operatorname{Re}\sigma \geq C$, $|\operatorname{Im}\sigma| \leq |\operatorname{Re}\sigma|^{2/3}$.

We write the integral kernel of the resolvent of each P_m using the following variation of parameters formula:

$$R_m(r, r') = -\psi_1(\max\{r, r'\})\psi_2(\min\{r, r'\})/W(\psi_1, \psi_2), \tag{8-3}$$

where ψ_1 and ψ_2 are linearly independent solutions to $(P_m - \sigma^2)u = 0$ and $W(\psi_1, \psi_2)$ is their Wronskian.

If $m = 0$ we take $\psi_1(r) = e^{ir\sigma}$ and $\psi_2(r) = e^{-ir\sigma}$ (this is the choice for which the resolvent maps L^2 to L^2 for $\operatorname{Im}\sigma > 0$), so that $W(\psi_1, \psi_2) = 2i\sigma$. Now the asserted continuation is immediate from the formula (8-3).

To study $m > 0$ we use, as in [Borthwick 2007, §5.3], the Bessel functions

$$\psi_1(r) = I_\nu(\lambda_m e^{-r}), \quad \psi_2(r) = K_\nu(\lambda_m e^{-r}), \quad \nu = -i\sigma. \tag{8-4}$$

We recall the definitions:

$$I_\nu(z) := \frac{z^\nu}{2^\nu} \sum_{k=0}^\infty \frac{(z/2)^{2k}}{k!\Gamma(\nu+k+1)}, \tag{8-5}$$

$$K_\nu(z) := \frac{\pi}{2\sin(\pi\nu)} (I_{-\nu}(z) - I_\nu(z)). \tag{8-6}$$

This pair solves the desired equation (see for example [Olver 1974, Chapter 7, (8.01)]) and has Wronskian $W = 1$ (see for example [ibid., Chapter 7, (8.07)]). When $\operatorname{Im}\sigma > 0$, we have $\operatorname{Re}\nu > 0$ and this resolvent maps L^2 to L^2 thanks to the asymptotic

$$I_\nu(z) = \frac{z^\nu}{2^\nu\Gamma(\nu+1)} \left(1 + \mathcal{O}\left(\frac{z^2}{\nu}\right)\right), \tag{8-7}$$

which is a consequence of (8-5), and thanks to the fact that $K_\nu(z) \sim e^{-z}\sqrt{\pi/2z}$ as $z \rightarrow \infty$ (see for example [ibid., Chapter 7, (8.04)]). Because I and K are entire in ν , we have the desired holomorphic continuation of the resolvent for all $m > 0$.

To estimate the resolvent we use (8-6) and (8-7) to write

$$K_\nu(z) = \frac{\pi}{2\sin(\pi\nu)} \left(\frac{z^{-\nu}}{2^{-\nu}\Gamma(-\nu+1)} - \frac{z^\nu}{2^\nu\Gamma(\nu+1)}\right) \left(1 + \mathcal{O}\left(\frac{z^2}{\nu}\right)\right).$$

Using Euler’s reflection formula for the gamma function (see for example [ibid., Chapter 2, (1.07)]),

$$\frac{\pi}{\sin(\pi\nu)\Gamma(\nu+1)} = -\Gamma(-\nu) = \frac{\Gamma(-\nu+1)}{\nu},$$

it follows that

$$\begin{aligned} K_\nu(z) &= \frac{\Gamma(\nu+1)}{2^\nu} \left(\frac{z^{-\nu}}{2^{-\nu}} - \frac{z^\nu \Gamma(-\nu+1)}{2^\nu \Gamma(\nu+1)} \right) \left(1 + \mathcal{O}\left(\frac{z^2}{\nu}\right) \right) \\ &= \frac{\Gamma(\nu+1)}{2^\nu} \left(\frac{z^{-\nu}}{2^{-\nu}} + \frac{\nu z^\nu \sin(\pi\nu) \Gamma(-\nu)^2}{2^\nu \pi} \right) \left(1 + \mathcal{O}\left(\frac{z^2}{\nu}\right) \right). \end{aligned} \tag{8-8}$$

To prove (8-1) we assume (without loss of generality) that there is $a > 0$ such that $\chi \geq 1$ on $[-a, a]$, and fix such an a . Let f be the characteristic function of $[0, a]$, and let

$$u(r) := (P_1 - \sigma^2)^{-1} f(r) = - \int_0^a R_1(r, r') dr' = K_\nu(\lambda_1 e^{-r}) \int_0^a I_\nu(\lambda_1 e^{-r'}) dr'.$$

Then $\|\chi(P_1 - \sigma^2)^{-1} \chi\|_{L^2(\mathbb{R}) \rightarrow L^2(\mathbb{R})} \geq \|\chi u\|_{L^2(\mathbb{R})} / \|f\|_{L^2(\mathbb{R})}$ and hence

$$\begin{aligned} \|\chi(P_1 - \sigma^2)^{-1} \chi\|_{L^2(\mathbb{R}) \rightarrow L^2(\mathbb{R})}^2 &\geq \frac{1}{a} \int_{-a}^a |u(r)|^2 dr \geq \frac{1}{a} \int_{-a}^a \left| K_\nu(\lambda_1 e^{-r}) \int_0^a I_\nu(\lambda_1 e^{-r'}) dr' \right|^2 dr \\ &= \frac{1}{a} \left| \int_0^a I_\nu(\lambda_1 e^{-r'}) dr' \right|^2 \int_{-a}^0 |K_\nu(\lambda_1 e^{-r})|^2 dr. \end{aligned}$$

Using (8-7) and (8-8) we obtain

$$\begin{aligned} \|\chi(P_1 - \sigma^2)^{-1} \chi\|_{L^2(\mathbb{R}) \rightarrow L^2(\mathbb{R})}^2 &\geq \frac{1}{8a|\nu|^2} \left| \int_0^a \frac{(\lambda_1 e^{-r'})^\nu}{2^\nu} dr' \right|^2 \int_{-a}^0 \left| \frac{(\lambda_1 e^{-r})^{-\nu}}{2^{-\nu}} + \frac{\nu(\lambda_1 e^{-r})^\nu \sin(\pi\nu) \Gamma(-\nu)^2}{2^\nu \pi} \right|^2 dr, \end{aligned} \tag{8-9}$$

provided $|\nu|$ is sufficiently large.

We now bound the two integrals from below one by one. First,

$$\left| \int_0^a \frac{(\lambda_1 e^{-r'})^\nu}{2^\nu} dr' \right| = \frac{\lambda_1^{\operatorname{Re} \nu}}{2^{\operatorname{Re} \nu} |\nu|} |e^{-a\nu} - 1| \geq e^{-C|\operatorname{Re} \nu|} / C|\nu|, \tag{8-10}$$

since $\operatorname{Re} \nu = \operatorname{Im} \sigma \leq -\varepsilon$. Second, using Stirling’s formula (see for example [ibid., Chapter 8, (4.04)])

$$\Gamma(-\nu) = e^\nu (-\nu)^{-\nu} \sqrt{-2\pi/\nu} (1 + \mathcal{O}(\nu^{-1})),$$

with

$$\arg(-\nu) := \frac{\pi}{2} - \arctan \frac{|\operatorname{Re} \nu|}{|\operatorname{Im} \nu|}$$

taking values in $(0, \frac{\pi}{2})$, and where the branch of $(-\nu)^{-\nu}$ is real and positive when $-\nu$ is, we write

$$\begin{aligned} |\nu \sin(\pi\nu) \Gamma(-\nu)^2| &= \pi e^{\pi |\operatorname{Im} \nu|} e^{-2|\operatorname{Re} \nu|} |\nu|^{2|\operatorname{Re} \nu|} e^{-2|\operatorname{Im} \nu| \arg(-\nu)} (1 + \mathcal{O}(|\operatorname{Im} \nu|^{-1})), \\ &= \pi e^{-2|\operatorname{Re} \nu|} |\nu|^{2|\operatorname{Re} \nu|} e^{2|\operatorname{Im} \nu| \arctan |\operatorname{Re} \nu / \operatorname{Im} \nu|} (1 + \mathcal{O}(|\operatorname{Im} \nu|^{-1})) \\ &= \pi |\nu|^{2|\operatorname{Re} \nu|} e^{-\frac{2}{3}|\operatorname{Re} \nu|^3 / |\operatorname{Im} \nu|^2} (1 + \mathcal{O}(|\operatorname{Re} \nu|^5 |\operatorname{Im} \nu|^{-4} + |\operatorname{Im} \nu|^{-1})). \end{aligned}$$

Hence, as long as $|\operatorname{Re} \nu|^{-3} |\operatorname{Im} \nu|^2$ is bounded and $|\nu|$ is sufficiently large, and using $\operatorname{Re} \nu \leq -\varepsilon$,

$$\begin{aligned} \left| \frac{(\lambda_1 e^{-r})^{-\nu}}{2^{-\nu}} + \frac{\nu(\lambda_1 e^{-r})^\nu \sin(\pi \nu) \Gamma(-\nu)^2}{2^\nu \pi} \right| &\geq \frac{1}{2} |\nu|^{-2 \operatorname{Re} \nu} e^{\frac{2}{3}(\operatorname{Re} \nu)^3 / (\operatorname{Im} \nu)^2} \frac{(\lambda_1 e^{-r})^{\operatorname{Re} \nu}}{2^{\operatorname{Re} \nu}} - \frac{2^{\operatorname{Re} \nu}}{(\lambda_1 e^{-r})^{\operatorname{Re} \nu}} \\ &\geq \frac{1}{C} |\nu|^{2|\operatorname{Re} \nu|} \left(\frac{2e^r}{\lambda_1} \right)^{|\operatorname{Re} \nu|} \end{aligned}$$

for $|r| \leq a$. This implies

$$\begin{aligned} \int_a^0 \left| \frac{(\lambda_1 e^{-r})^{-\nu}}{2^{-\nu}} + \frac{\nu(\lambda_1 e^{-r})^\nu \sin(\pi \nu) \Gamma(-\nu)^2}{2^\nu \pi} \right|^2 dr &\geq \frac{1}{C} |\nu|^{4|\operatorname{Re} \nu|} \left(\frac{2}{\lambda_1} \right)^{2|\operatorname{Re} \nu|} \int_{-a}^0 e^{2|\operatorname{Re} \nu|r} dr \\ &\geq |\nu|^{4|\operatorname{Re} \nu|} e^{-C \operatorname{Re} \nu} / C. \end{aligned}$$

Combining this with (8-9) and (8-10), and using $\nu = -i\sigma$, gives (8-2) and hence (8-1). □

Appendix: The curvature of a warped product

The result of this calculation is used in the examples in Section 2D, and although it is well known, we include the details for the convenience of the reader. For this section only, let (S, \tilde{g}) be a compact Riemannian manifold, and let $X = \mathbb{R} \times S$ have the metric

$$g = dr^2 + f(r)^2 \tilde{g},$$

where $f \in C^\infty(\mathbb{R}; (0, \infty))$. Let $p \in X$, let P be a two-dimensional subspace of $T_p X$, and let $K(P)$ be the sectional curvature of P with respect to g . We will show that if $\partial_r \in P$, then

$$K(P) = -f''(r)/f(r),$$

while if $P \subset T_p S$ and $\tilde{K}(P)$ is the sectional curvature of P with respect to \tilde{g} , then

$$K(P) = (\tilde{K}(P) - f'(r)^2)/f(r)^2.$$

We work in coordinates $(x^0, \dots, x^n) = (r, x^1, \dots, x^n)$, and write

$$g = g_{\alpha\beta} dx^\alpha dx^\beta = dr^2 + g_{ij} dx^i dx^j = dr^2 + f(r)^2 \tilde{g}_{ij} dx^i dx^j,$$

using the Einstein summation convention. We use Greek letters for indices which include 0, that is indices which include r , and Latin letters for indices which do not. Then

$$\partial_\alpha g_{r\alpha} = 0, \quad \partial_r g_{jk} = 2f^{-1} f' g_{jk}, \quad \partial_i g_{jk} = f^2 \partial_i \tilde{g}_{jk}.$$

We write Γ for the Christoffel symbols of g , and $\tilde{\Gamma}$ for those of \tilde{g} . These are given by

$$\Gamma^r_{r\alpha} = \Gamma^\alpha_{rr} = 0, \quad \Gamma^r_{jk} = -f^{-1} f' g_{jk}, \quad \Gamma^i_{jr} = f^{-1} f' \delta_j^i, \quad \Gamma^i_{jk} = \tilde{\Gamma}^i_{jk}.$$

Let R be the Riemann curvature tensor of g :

$$R_{\alpha\beta\gamma}{}^\delta = \partial_\alpha \Gamma^\delta_{\beta\gamma} + \Gamma^\varepsilon_{\beta\gamma} \Gamma^\delta_{\alpha\varepsilon} - \partial_\beta \Gamma^\delta_{\alpha\gamma} - \Gamma^\varepsilon_{\alpha\gamma} \Gamma^\delta_{\beta\varepsilon}.$$

Now if $P \subset T_p X$ is spanned by a pair of orthogonal unit vectors $V^\alpha \partial_\alpha$ and $W^\alpha \partial_\alpha$, then $K(P) = R_{\alpha\beta\gamma\delta} V^\alpha W^\beta W^\gamma V^\delta$, and similarly for \tilde{R} and \tilde{K} . Then

$$\begin{aligned} R_{ijk}{}^l &= \tilde{R}_{ijk}{}^l + \Gamma^r{}_{jk} \Gamma^l{}_{ir} - \Gamma^r{}_{ik} \Gamma^l{}_{jr} = \tilde{R}_{ijk}{}^l + (f^{-1})^2 (f')^2 (-\delta_i^l g_{jk} + \delta_j^l g_{ik}), \\ R_{rjk}{}^r &= \partial_r \Gamma^r{}_{jk} - \Gamma^m{}_{rk} \Gamma^r{}_{jm} = -(f^{-1} f' g_{jk})' + (f^{-1} f')^2 g_{jk} = -f^{-1} f'' g_{jk}. \end{aligned}$$

If $\partial_r \in P$ we take $V = \partial_r$ and $W = W^j \partial_j$ any unit vector in $T_p X$ orthogonal to V . Then

$$K(P) = R_{rjkr} W^j W^k = -f^{-1} f'' g_{jk} W^j W^k = -f^{-1} f''.$$

Meanwhile, if $\partial_r \perp P$, we may write $V = V^j \partial_j$ and $W = W^j \partial_j$. Then

$$K(P) = (f^2 \tilde{R}_{ijkl} + (f^{-1})^2 (f')^2 (-g_{li} g_{jk} + g_{lj} g_{ik})) V^i W^j W^k V^l.$$

Using the fact that fV and fW are orthogonal unit vectors for \tilde{g} , we see that

$$K(P) = f^{-2} \tilde{K}(P) - (f^{-1})^2 (f')^2.$$

Acknowledgements

I am indebted especially to Maciej Zworski for his generous guidance, advice, and unflinching encouragement throughout the course of this project. Thanks also to András Vasy, Nicolas Burq, John Lott, David Borthwick, Colin Guillarmou, Hamid Hezari, Semyon Dyatlov, and Richard Melrose for their interest and for their many very helpful ideas, comments, and suggestions. I am also grateful to the several anonymous referees for their careful reading, and for pointing out corrections and suggesting a number of improvements to the presentation, and to Matt Tucker-Simmons for his meticulous editing.

Thanks finally to the National Science Foundation and the Simons Foundation for partial support (under NSF grant DMS-0654436, under an NSF MSPRF grant, and under a Simons Collaboration Grant for Mathematicians), and to the Mathematical Sciences Research Institute and the Université Paris 13 for their hospitality while I was a visitor.

References

- [Aguilar and Combes 1971] J. Aguilar and J. M. Combes, “A class of analytic perturbations for one-body Schrödinger Hamiltonians”, *Comm. Math. Phys.* **22** (1971), 269–279. [MR 0345551](#) [Zbl 0219.47011](#)
- [Alexandrova and Tamura 2011] I. Alexandrova and H. Tamura, “Resonance free regions in magnetic scattering by two solenoidal fields at large separation”, *J. Funct. Anal.* **260**:6 (2011), 1836–1885. [MR 2754895](#) [Zbl 1211.81093](#)
- [Baskin and Wunsch 2013] D. Baskin and J. Wunsch, “Resolvent estimates and local decay of waves on conic manifolds”, *J. Differential Geom.* **95**:2 (2013), 183–214. [MR 3128982](#) [Zbl 1296.53075](#)
- [Bony and Häfner 2008] J.-F. Bony and D. Häfner, “Decay and non-decay of the local energy for the wave equation on the de Sitter–Schwarzschild metric”, *Comm. Math. Phys.* **282**:3 (2008), 697–719. [MR 2426141](#) [Zbl 1159.35007](#)
- [Bony and Petkov 2013] J.-F. Bony and V. Petkov, “Semiclassical estimates of the cut-off resolvent for trapping perturbations”, *J. Spectr. Theory* **3**:3 (2013), 399–422. [MR 3073417](#) [Zbl 1290.47013](#)
- [Borthwick 2007] D. Borthwick, *Spectral theory of infinite-area hyperbolic surfaces*, Progress in Mathematics **256**, Birkhäuser, Boston, 2007. [MR 2344504](#) [Zbl 1130.58001](#)

- [Bridson and Haefliger 1999] M. R. Bridson and A. Haefliger, *Metric spaces of non-positive curvature*, Grundlehren der Math. Wissenschaften **319**, Springer, Berlin, 1999. MR 1744486 Zbl 0988.53001
- [Burq 1998] N. Burq, “Décroissance de l’énergie locale de l’équation des ondes pour le problème extérieur et absence de résonance au voisinage du réel”, *Acta Math.* **180**:1 (1998), 1–29. MR 1618254 Zbl 0918.35081
- [Burq 2002] N. Burq, “Lower bounds for shape resonances widths of long range Schrödinger operators”, *Amer. J. Math.* **124**:4 (2002), 677–735. MR 1914456 Zbl 1013.35019
- [Burq 2004] N. Burq, “Smoothing effect for Schrödinger boundary value problems”, *Duke Math. J.* **123**:2 (2004), 403–427. MR 2066943 Zbl 1061.35024
- [Burq and Zworski 2004] N. Burq and M. Zworski, “Geometric control in the presence of a black box”, *J. Amer. Math. Soc.* **17**:2 (2004), 443–471. MR 2051618 Zbl 1050.35058
- [Burq, Guillarmou and Hassell 2010] N. Burq, C. Guillarmou, and A. Hassell, “Strichartz estimates without loss on manifolds with hyperbolic trapped geodesics”, *Geom. Funct. Anal.* **20**:3 (2010), 627–656. MR 2720226 Zbl 1206.58009
- [Cardoso and Vodev 2002] F. Cardoso and G. Vodev, “Uniform estimates of the resolvent of the Laplace–Beltrami operator on infinite volume Riemannian manifolds, II”, *Ann. Henri Poincaré* **3**:4 (2002), 673–691. MR 1933365 Zbl 1021.58016
- [Cardoso, Popov and Vodev 2004] F. Cardoso, G. Popov, and G. Vodev, “Semi-classical resolvent estimates for the Schrödinger operator on non-compact complete Riemannian manifolds”, *Bull. Braz. Math. Soc. (N.S.)* **35**:3 (2004), 333–344. MR 2106308 Zbl 1159.58308
- [Christianson 2009] H. Christianson, “Applications of cutoff resolvent estimates to the wave equation”, *Math. Res. Lett.* **16**:4 (2009), 577–590. MR 2525026 Zbl 1189.58012
- [Christianson, Schenck, Vasy and Wunsch 2014] H. Christianson, E. Schenck, A. Vasy, and J. Wunsch, “From resolvent estimates to damped waves”, *J. Anal. Math.* **122** (2014), 143–162. MR 3183526 Zbl 1301.35191
- [Datchev 2010] K. Datchev, *Distribution of resonances for manifolds with hyperbolic ends*, Ph.D. thesis, University of California, Berkeley, 2010, available at <http://search.proquest.com/docview/748836620>.
- [Datchev 2012] K. Datchev, “Extending cutoff resolvent estimates via propagation of singularities”, *Comm. Partial Differential Equations* **37**:8 (2012), 1456–1461. MR 2957548 Zbl 1254.35034
- [Datchev and Dyatlov 2013] K. Datchev and S. Dyatlov, “Fractal Weyl laws for asymptotically hyperbolic manifolds”, *Geom. Funct. Anal.* **23**:4 (2013), 1145–1206. MR 3077910 Zbl 1297.58006
- [Datchev and Vasy 2012a] K. Datchev and A. Vasy, “Gluing semiclassical resolvent estimates via propagation of singularities”, *Int. Math. Res. Not.* **2012**:23 (2012), 5409–5443. MR 2999147 Zbl 1262.58019
- [Datchev and Vasy 2012b] K. Datchev and A. Vasy, “Propagation through trapped sets and semiclassical resolvent estimates”, *Ann. Inst. Fourier (Grenoble)* **62**:6 (2012), 2347–2377. MR 3060760 Zbl 1271.58014
- [Datchev, Kang and Kessler 2015] K. R. Datchev, D. D. Kang, and A. P. Kessler, “Non-trapping surfaces of revolution with long-living resonances”, *Math. Res. Lett.* **22**:1 (2015), 23–42. MR 3342177 Zbl 1328.58028
- [Dimassi and Sjöstrand 1999] M. Dimassi and J. Sjöstrand, *Spectral asymptotics in the semi-classical limit*, London Mathematical Society Lecture Note Series **268**, Cambridge University Press, 1999. MR 1735654 Zbl 0926.35002
- [Dyatlov 2012] S. Dyatlov, “Asymptotic distribution of quasi-normal modes for Kerr–de Sitter black holes”, *Ann. Henri Poincaré* **13**:5 (2012), 1101–1166. MR 2935116 Zbl 1246.83111
- [Dyatlov 2015a] S. Dyatlov, “Asymptotics of linear waves and resonances with applications to black holes”, *Comm. Math. Phys.* **335**:3 (2015), 1445–1485. MR 3320319 Zbl 1315.83022
- [Dyatlov 2015b] S. Dyatlov, “Resonance projectors and asymptotics for r -normally hyperbolic trapped sets”, *J. Amer. Math. Soc.* **28**:2 (2015), 311–381. MR 3300697 Zbl 06394348
- [Dyatlov and Zahl 2015] S. Dyatlov and J. Zahl, “Spectral gaps, additive energy, and a fractal uncertainty principle”, preprint, 2015. arXiv 1504.06589
- [Dyatlov and Zworski 2016] S. Dyatlov and M. Zworski, “Mathematical theory of scattering resonances”, book project in progress, 2016, available at <http://math.mit.edu/~dyatlov/res/>.
- [Galkowski 2015] J. Galkowski, “Distribution of resonances in scattering by thin barriers”, preprint, 2015. arXiv 1404.3709

- [Galkowski 2016] J. Galkowski, “A quantitative Vainberg method for black box scattering”, preprint, 2016. To appear in *Comm. Math. Phys.* [arXiv 1511.05894](#)
- [Galkowski and Smith 2015] J. Galkowski and H. F. Smith, “Restriction bounds for the free resolvent and resonances in lossy scattering”, *Int. Math. Res. Not.* **2015**:16 (2015), 7473–7509. [MR 3428971](#) [Zbl 06486428](#)
- [Guillarmou 2005] C. Guillarmou, “Meromorphic properties of the resolvent on asymptotically hyperbolic manifolds”, *Duke Math. J.* **129**:1 (2005), 1–37. [MR 2153454](#) [Zbl 1099.58011](#)
- [Guillarmou and Mazzeo 2012] C. Guillarmou and R. Mazzeo, “Resolvent of the Laplacian on geometrically finite hyperbolic manifolds”, *Invent. Math.* **187**:1 (2012), 99–144. [MR 2874936](#) [Zbl 1252.58015](#)
- [Guillarmou and Naud 2009] C. Guillarmou and F. Naud, “Wave decay on convex co-compact hyperbolic manifolds”, *Comm. Math. Phys.* **287**:2 (2009), 489–511. [MR 2481747](#) [Zbl 1196.58011](#)
- [Guillopé and Zworski 1997] L. Guillopé and M. Zworski, “Scattering asymptotics for Riemann surfaces”, *Ann. of Math.* (2) **145**:3 (1997), 597–660. [MR 1454705](#) [Zbl 0898.58054](#)
- [Guillopé, Lin and Zworski 2004] L. Guillopé, K. K. Lin, and M. Zworski, “The Selberg zeta function for convex co-compact Schottky groups”, *Comm. Math. Phys.* **245**:1 (2004), 149–176. [MR 2036371](#) [Zbl 1075.11059](#)
- [Ikawa 1982] M. Ikawa, “Decay of solutions of the wave equation in the exterior of two convex obstacles”, *Osaka J. Math.* **19**:3 (1982), 459–509. [MR 676233](#) [Zbl 0498.35008](#)
- [Lax and Phillips 1989] P. D. Lax and R. S. Phillips, *Scattering theory*, 2nd ed., Pure and Applied Mathematics **26**, Academic Press, Boston, 1989. [MR 1037774](#) [Zbl 0697.35004](#)
- [Martinez 2002] A. Martinez, “Resonance free domains for non globally analytic potentials”, *Ann. Henri Poincaré* **3**:4 (2002), 739–756. Erratum published in *Ann. Henri Poincaré* **8**:7 (2007), 1425–1431. [MR 1933368](#) [Zbl 1026.81012](#)
- [Mazzeo and Melrose 1987] R. R. Mazzeo and R. B. Melrose, “Meromorphic extension of the resolvent on complete spaces with asymptotically constant negative curvature”, *J. Funct. Anal.* **75**:2 (1987), 260–310. [MR 916753](#) [Zbl 0636.58034](#)
- [Melrose and Sjöstrand 1982] R. B. Melrose and J. Sjöstrand, “Singularities of boundary value problems, II”, *Comm. Pure Appl. Math.* **35**:2 (1982), 129–168. [MR 644020](#) [Zbl 0546.35083](#)
- [Melrose, Sá Barreto and Vasy 2014] R. Melrose, A. Sá Barreto, and A. Vasy, “Asymptotics of solutions of the wave equation on de Sitter–Schwarzschild space”, *Comm. Partial Differential Equations* **39**:3 (2014), 512–529. [MR 3169793](#) [Zbl 1286.35145](#)
- [Morawetz, Ralston and Strauss 1977] C. S. Morawetz, J. V. Ralston, and W. A. Strauss, “Decay of solutions of the wave equation outside nontrapping obstacles”, *Comm. Pure Appl. Math.* **30**:4 (1977), 447–508. [MR 0509770](#) [Zbl 0372.35008](#)
- [Nakamura, Stefanov and Zworski 2003] S. Nakamura, P. Stefanov, and M. Zworski, “Resonance expansions of propagators in the presence of potential barriers”, *J. Funct. Anal.* **205**:1 (2003), 180–205. [MR 2020213](#) [Zbl 1037.35064](#)
- [Nonnenmacher and Zworski 2009] S. Nonnenmacher and M. Zworski, “Quantum decay rates in chaotic scattering”, *Acta Math.* **203**:2 (2009), 149–233. [MR 2570070](#) [Zbl 1226.35061](#)
- [Nonnenmacher and Zworski 2015] S. Nonnenmacher and M. Zworski, “Decay of correlations for normally hyperbolic trapping”, *Invent. Math.* **200**:2 (2015), 345–438. [MR 3338007](#) [Zbl 06442708](#)
- [Olver 1974] F. W. J. Olver, *Asymptotics and special functions*, Academic Press, New York, 1974. [MR 0435697](#) [Zbl 0303.41035](#)
- [Petkov and Stoyanov 2010] V. Petkov and L. Stoyanov, “Analytic continuation of the resolvent of the Laplacian and the dynamical zeta function”, *Anal. PDE* **3**:4 (2010), 427–489. [MR 2718260](#) [Zbl 1251.37031](#)
- [Reed and Simon 1978] M. Reed and B. Simon, *Methods of modern mathematical physics, IV: Analysis of operators*, Academic Press, New York, 1978. [MR 0493421](#) [Zbl 0401.47001](#)
- [Regge 1958] T. Regge, “Analytic properties of the scattering matrix”, *Nuovo Cimento* (10) **8** (1958), 671–679. [MR 0095702](#) [Zbl 0080.41903](#)
- [Sá Barreto and Wang 2015] A. Sá Barreto and Y. Wang, “High energy resolvent estimates on conformally compact manifolds with variable curvature at infinity”, preprint, 2015. [arXiv 1511.05891](#)
- [Selberg 1990] A. Selberg, “Remarks on the distribution of poles of Eisenstein series”, pp. 251–278 in *Festschrift in honor of I. I. Piatetski-Shapiro on the occasion of his sixtieth birthday, Part II* (Ramat Aviv, 1989), edited by S. Gelbart et al., Israel Math. Conf. Proc. **3**, Weizmann, Jerusalem, 1990. [MR 1159119](#) [Zbl 0712.11034](#)

- [Simon 1972] B. Simon, “Quadratic form techniques and the Balslev–Combes theorem”, *Comm. Math. Phys.* **27** (1972), 1–9. [MR 0321456](#) [Zbl 0237.35025](#)
- [Sjöstrand 1990] J. Sjöstrand, “Geometric bounds on the density of resonances for semiclassical problems”, *Duke Math. J.* **60**:1 (1990), 1–57. [MR 1047116](#) [Zbl 0702.35188](#)
- [Sjöstrand and Zworski 1991] J. Sjöstrand and M. Zworski, “Complex scaling and the distribution of scattering poles”, *J. Amer. Math. Soc.* **4**:4 (1991), 729–769. [MR 1115789](#) [Zbl 0752.35046](#)
- [Sjöstrand and Zworski 2007] J. Sjöstrand and M. Zworski, “Fractal upper bounds on the density of semiclassical resonances”, *Duke Math. J.* **137**:3 (2007), 381–459. [MR 2309150](#) [Zbl 1201.35189](#)
- [Tang and Zworski 1998] S.-H. Tang and M. Zworski, “From quasimodes to resonances”, *Math. Res. Lett.* **5**:3 (1998), 261–272. [MR 1637824](#) [Zbl 0913.35101](#)
- [Tang and Zworski 2000] S.-H. Tang and M. Zworski, “Resonance expansions of scattered waves”, *Comm. Pure Appl. Math.* **53**:10 (2000), 1305–1334. [MR 1768812](#) [Zbl 1032.35148](#)
- [Vainberg 1989] B. R. Vainberg, *Asymptotic methods in equations of mathematical physics*, Gordon & Breach, New York, 1989. [MR 1054376](#) [Zbl 0743.35001](#)
- [Vasy 2013] A. Vasy, “Microlocal analysis of asymptotically hyperbolic and Kerr–de Sitter spaces”, *Invent. Math.* **194**:2 (2013), 381–513. [MR 3117526](#) [Zbl 1315.35015](#)
- [Vasy and Zworski 2000] A. Vasy and M. Zworski, “Semiclassical estimates in asymptotically Euclidean scattering”, *Comm. Math. Phys.* **212**:1 (2000), 205–217. [MR 1764368](#) [Zbl 0955.58023](#)
- [Wang 2014] Y. Wang, “Resolvent and radiation fields on non-trapping asymptotically hyperbolic manifolds”, preprint, 2014. [arXiv 1410.6936](#)
- [Wunsch 2012] J. Wunsch, “Resolvent estimates with mild trapping”, pp. Exposé 13 in *Journées équations aux dérivées partielles* (Biarritz, France, 2012), Cedram, 2012.
- [Wunsch and Zworski 2011] J. Wunsch and M. Zworski, “Resolvent estimates for normally hyperbolic trapped sets”, *Ann. Henri Poincaré* **12**:7 (2011), 1349–1385. [MR 2846671](#) [Zbl 1228.81170](#)
- [Zworski 1987] M. Zworski, “Distribution of poles for scattering on the real line”, *J. Funct. Anal.* **73**:2 (1987), 277–296. [MR 899652](#) [Zbl 0662.34033](#)
- [Zworski 1999] M. Zworski, “Dimension of the limit set and the density of resonances for convex co-compact hyperbolic surfaces”, *Invent. Math.* **136**:2 (1999), 353–409. [MR 1688441](#) [Zbl 1016.58014](#)
- [Zworski 2012] M. Zworski, *Semiclassical analysis*, Graduate Studies in Mathematics **138**, American Mathematical Society, Providence, RI, 2012. [MR 2952218](#) [Zbl 1252.58001](#)

Received 16 Dec 2015. Accepted 26 Feb 2016.

KIRIL DATCHEV: kdatchev@purdue.edu

Mathematics Department, Purdue University, 150 N. University Street, West Lafayette, IN 47907, United States

Analysis & PDE

msp.org/apde

EDITORS

EDITOR-IN-CHIEF

Patrick Gérard

patrick.gerard@math.u-psud.fr

Université Paris Sud XI

Orsay, France

BOARD OF EDITORS

Nicolas Burq	Université Paris-Sud 11, France nicolas.burq@math.u-psud.fr	Werner Müller	Universität Bonn, Germany mueller@math.uni-bonn.de
Massimiliano Berti	Scuola Intern. Sup. di Studi Avanzati, Italy berti@sissa.it	Yuval Peres	University of California, Berkeley, USA peres@stat.berkeley.edu
Sun-Yung Alice Chang	Princeton University, USA chang@math.princeton.edu	Gilles Pisier	Texas A&M University, and Paris 6 pisier@math.tamu.edu
Michael Christ	University of California, Berkeley, USA mchrist@math.berkeley.edu	Tristan Rivière	ETH, Switzerland riviere@math.ethz.ch
Charles Fefferman	Princeton University, USA cf@math.princeton.edu	Igor Rodnianski	Princeton University, USA irod@math.princeton.edu
Ursula Hamenstaedt	Universität Bonn, Germany ursula@math.uni-bonn.de	Wilhelm Schlag	University of Chicago, USA schlag@math.uchicago.edu
Vaughan Jones	U.C. Berkeley & Vanderbilt University vaughan.f.jones@vanderbilt.edu	Sylvia Serfaty	New York University, USA serfaty@cims.nyu.edu
Vadim Kaloshin	University of Maryland, USA vadim.kaloshin@gmail.com	Yum-Tong Siu	Harvard University, USA siu@math.harvard.edu
Herbert Koch	Universität Bonn, Germany koch@math.uni-bonn.de	Terence Tao	University of California, Los Angeles, USA tao@math.ucla.edu
Izabella Laba	University of British Columbia, Canada ilaba@math.ubc.ca	Michael E. Taylor	Univ. of North Carolina, Chapel Hill, USA met@math.unc.edu
Gilles Lebeau	Université de Nice Sophia Antipolis, France lebeau@unice.fr	Gunther Uhlmann	University of Washington, USA gunther@math.washington.edu
László Lempert	Purdue University, USA lempert@math.purdue.edu	András Vasy	Stanford University, USA andras@math.stanford.edu
Richard B. Melrose	Massachusetts Inst. of Tech., USA rbb@math.mit.edu	Dan Virgil Voiculescu	University of California, Berkeley, USA dvv@math.berkeley.edu
Frank Merle	Université de Cergy-Pontoise, France Frank.Merle@u-cergy.fr	Steven Zelditch	Northwestern University, USA zelditch@math.northwestern.edu
William Minicozzi II	Johns Hopkins University, USA minicozz@math.jhu.edu	Maciej Zworski	University of California, Berkeley, USA zvorski@math.berkeley.edu
Clément Mouhot	Cambridge University, UK c.mouhot@dpms.cam.ac.uk		

PRODUCTION

production@msp.org

Silvio Levy, Scientific Editor


See inside back cover or msp.org/apde for submission instructions.

The subscription price for 2016 is US \$235/year for the electronic version, and \$430/year (+\$55, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues from the last three years and changes of subscribers address should be sent to MSP.

Analysis & PDE (ISSN 1948-206X electronic, 2157-5045 printed) at Mathematical Sciences Publishers, 798 Evans Hall #3840, c/o University of California, Berkeley, CA 94720-3840, is published continuously online. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices.

APDE peer review and production are managed by EditFlow® from MSP.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2016 Mathematical Sciences Publishers

ANALYSIS & PDE

Volume 9 No. 4 2016

Peierls substitution for magnetic Bloch bands	773
SILVIA FREUND and STEFAN TEUFEL	
Dispersive estimates in \mathbb{R}^3 with threshold eigenstates and resonances	813
MARIUS BECEANU	
Interior nodal sets of Steklov eigenfunctions on surfaces	859
JIUYI ZHU	
Some counterexamples to Sobolev regularity for degenerate Monge–Ampère equations	881
CONNOR MOONEY	
Mean ergodic theorem for amenable discrete quantum groups and a Wiener-type theorem for compact metrizable groups	893
HUICHI HUANG	
Resonance free regions for nontrapping manifolds with cusps	907
KIRIL DATCHEV	
Characterizing regularity of domains via the Riesz transforms on their boundaries	955
DORINA MITREA, MARIUS MITREA and JOAN VERDERA	