

ANALYSIS & PDE

Volume 11 No. 3 2018

JOACHIM KRIEGER AND YANNICK SIRE

SMALL DATA GLOBAL REGULARITY FOR HALF-WAVE MAPS

SMALL DATA GLOBAL REGULARITY FOR HALF-WAVE MAPS

JOACHIM KRIEGER AND YANNICK SIRE

We formulate the half-wave maps problem with target S^2 and prove global regularity in sufficiently high spatial dimensions for a class of small critical data in Besov spaces.

1. The problem

Let $u : \mathbb{R}^{n+1} \rightarrow S^2 \hookrightarrow \mathbb{R}^3$ be smooth, and assume that it converges to some $p \in S^2$ at spatial infinity. Further, assume that on each fixed time slice $\nabla_{t,x} u \in L^p(\mathbb{R}^n)$ for some $p \in (1, \infty)$. Denote by \times the standard vectorial product in three dimensions. We call this a *half-wave map*, provided it satisfies the relation

$$\partial_t u = u \times (-\Delta)^{\frac{1}{2}} u. \tag{1-1}$$

Here we define the operator $(-\Delta)^{\frac{1}{2}}$ via

$$(-\Delta)^{\frac{1}{2}} u = - \sum_{j=1}^n (-\Delta)^{-\frac{1}{2}} \partial_j (\partial_j u),$$

a specification necessary on account of the fact that u does not vanish at infinity, but instead approaches some $p \in S^2$, while ∇u does vanish at infinity. In fact, the expression $(-\Delta)^{\frac{1}{2}} u$ under our current definition is then well-defined since $\nabla_{t,x} u(t, \cdot) \in L^p(\mathbb{R}^n)$ for some $p \in (1, \infty)$, for all t .

We note that the model (1-1) appears formally related to the much-studied Schrödinger maps problem, which can be written in the form

$$\partial_t u = u \times \Delta u,$$

and moreover, we shall see shortly that (1-1) also appears closely related to the classical wave maps problem with target S^2 . We also note that we have a formally conserved quantity

$$E(t) := \int_{\mathbb{R}^n} |(-\Delta)^{\frac{1}{4}} u|^2 dx, \tag{1-2}$$

where we let $(-\Delta)^{\frac{1}{4}} u = - \sum_{j=1}^n (-\Delta)^{-\frac{3}{4}} \partial_j (\partial_j u)$. Such kinds of quantities have been considered in the works of Da Lio and Rivière in the study of fractional harmonic maps; see for instance [Da Lio and Rivière 2011a; 2011b; Da Lio 2013]. We also note that on account of the results on fractional harmonic maps previously mentioned, this model moreover displays a very rich class of static solutions; see also [Millot and Sire 2015].

MSC2010: 35L05, 35B40, 35A01.

Keywords: wave equation, fractional wave maps, half-wave maps, Besov spaces, global regularity.

On the other hand, (1-1) scales just like wave maps, which means that in all dimensions $n \geq 2$ the problem (1-1) is formally supercritical.

We formulated the model (1-1) as a nonlocal wave analogue to Schrödinger maps in 2014, but have since learned from E. Lenzmann¹ that it already exists in the physics literature. We learned from Lenzmann that the half-wave map equation arises as the continuum version of the so-called integrable spin Calogero–Moser systems, which in turn comes from the completely integrable quantum spin systems called Haldane–Shastry systems.² Recent work by Schikorra and Lenzmann [2017] completely classifies the travelling wave solutions for this model in the critical case $n = 1$.

In the present note, our goal is to approach the issue of global solutions corresponding to small data, attempting to parallel the developments in [Tataru 1998; Tao 2001a]. We will see that (1-1) can be reformulated as a nonlinear wave-type equation of the schematic form

$$\square u = F(u) \nabla_{t,x} u \cdot \nabla_{t,x} u, \quad (1-3)$$

although this is an oversimplification as the true underlying wave equation displays nonlocal expressions. It has been known now for a while, see [Sterbenz 2004], that (1-3) admits global solutions corresponding to initial data of small critical, i.e., scaling invariant, Besov $\dot{B}_2^{\frac{n}{2},1}$ norms, provided one restricts oneself to spatial dimensions $n \geq 6$, and that passing to lower dimensions appears to require some sort of null-structure. Here, we show that (1-1) does have enough of an intrinsic null-structure to allow for the following.

Theorem 1.1. *Let $n \geq 5$. Let*

$$u[0] = (u(0, \cdot), u_t(0, \cdot)) = (u_0, u_1) : \mathbb{R}^n \rightarrow S^2 \times TS^2$$

be a smooth data pair with $u_1 = u_0 \times (-\Delta)^{\frac{1}{2}} u_0$, and such that u_0 is constant outside of a compact subset of \mathbb{R}^n (this condition in particular ensures that $(-\Delta)^{\frac{1}{2}} u_0$ is well-defined). Also, assume the smallness condition

$$\|u[0]\|_{\dot{B}_2^{n/2,1} \times \dot{B}_2^{n/2-1,1}} < \epsilon,$$

where $\epsilon \ll 1$ is sufficiently small. Then problem (1-1) admits a global smooth solution.

To prove this theorem, we shall have to reformulate (1-1) as a wave equation, which we do next.

Remark 1.2. We note that the restriction $n \geq 5$ comes from the fact that we use the $L_t^2 L_x^4$ -Strichartz estimate, which is not available in spatial dimension $n = 4$. However, it is quite likely that this can be circumvented, and that the structures exhibited in this paper suffice to push the result to $n = 4$. However, both the issue of passing to the critical space $\dot{H}^{\frac{n}{2}}$, as well as going to lower spatial dimensions $n \leq 3$, appear nontrivial, as there are novel trilinear terms which no longer seem to have the same strong null-structure as the leading term coming from the wave maps equation.

¹The name of half-wave map was suggested by Lenzmann.

²Lenzmann provided us with the references [Haldane 1988; Shastry 1988; Hikami and Wadati 1993; Blom and Langmann 1998] and we refer to his work for an account on the passage from the physics to the mathematical model.

2. Passage to a wave equation

Departing from (1-1), we compute

$$\begin{aligned}\partial_t^2 u &= \partial_t u \times (-\Delta)^{\frac{1}{2}} u + u \times (-\Delta)^{\frac{1}{2}} \partial_t u \\ &= (u \times (-\Delta)^{\frac{1}{2}} u) \times (-\Delta)^{\frac{1}{2}} u + u \times (-\Delta)^{\frac{1}{2}} (u \times (-\Delta)^{\frac{1}{2}} u).\end{aligned}$$

Then using the basic formula $a \times (b \times c) = b(a \cdot c) - c(a \cdot b)$, $a, b, c \in \mathbb{R}^3$, we rewrite the first term on the right as

$$(u \times (-\Delta)^{\frac{1}{2}} u) \times (-\Delta)^{\frac{1}{2}} u = -u((-\Delta)^{\frac{1}{2}} u \cdot (-\Delta)^{\frac{1}{2}} u) + (-\Delta)^{\frac{1}{2}} u(u \cdot (-\Delta)^{\frac{1}{2}} u).$$

For the second term on the right above, introducing a commutator term, we write it in the form

$$\begin{aligned}u \times (-\Delta)^{\frac{1}{2}} (u \times (-\Delta)^{\frac{1}{2}} u) &= u \times (-\Delta)^{\frac{1}{2}} (u \times (-\Delta)^{\frac{1}{2}} u) - u \times (u \times (-\Delta)u) + u \times (u \times (-\Delta)u) \\ &= u \times (-\Delta)^{\frac{1}{2}} (u \times (-\Delta)^{\frac{1}{2}} u) - u \times (u \times (-\Delta)u) + u(u \cdot (-\Delta)u) + \Delta u.\end{aligned}$$

Using the fact that $u \cdot u = 1$, whence

$$u \cdot \Delta u + \nabla u \cdot \nabla u = 0,$$

we arrive at the equation

$$\begin{aligned}(\partial_t^2 - \Delta)u &= -u((-\Delta)^{\frac{1}{2}} u \cdot (-\Delta)^{\frac{1}{2}} u) + (-\Delta)^{\frac{1}{2}} u(u \cdot (-\Delta)^{\frac{1}{2}} u) + u \times (-\Delta)^{\frac{1}{2}} (u \times (-\Delta)^{\frac{1}{2}} u) \\ &\quad - u \times (u \times (-\Delta)u) + u(\nabla u \cdot \nabla u).\end{aligned}$$

Carefully note that ∇u here only involves the spatial derivatives. In order to make this appear closer to the wave maps equation and introduce better null-structure, we have to also make the time derivatives visible on the right-hand side, for which the first line on the right-hand side is pivotal. In fact, we get

$$(-u((-\Delta)^{\frac{1}{2}} u \cdot (-\Delta)^{\frac{1}{2}} u) + (-\Delta)^{\frac{1}{2}} u(u \cdot (-\Delta)^{\frac{1}{2}} u)) \cdot u = -|u \times (-\Delta)^{\frac{1}{2}} u|^2 = -|\partial_t u|^2,$$

and so the equation becomes

$$\begin{aligned}(\partial_t^2 - \Delta)u &= u(\nabla u \cdot \nabla u - \partial_t u \cdot \partial_t u) + \Pi_{u^\perp} ((-\Delta)^{\frac{1}{2}} u)(u \cdot (-\Delta)^{\frac{1}{2}} u) \\ &\quad + u \times (-\Delta)^{\frac{1}{2}} (u \times (-\Delta)^{\frac{1}{2}} u) - u \times (u \times (-\Delta)u), \quad (2-1)\end{aligned}$$

where Π_{u^\perp} denotes projection onto the orthogonal complement of u . Thus we see that formally the nonlinearity involves the precise wave maps source term, as well as two error terms, which formally behave like

$$u \nabla u \cdot \nabla u.$$

3. Technical preliminaries

Our main tools shall be the classical Strichartz estimates, combined with some $X^{s,b}$ -space technology. Specifically, we let P_k , $k \in \mathbb{Z}$, be standard Littlewood–Paley multipliers on \mathbb{R}^n (acting on the spatial variables), and furthermore, we denote by Q_j , $j \in \mathbb{Z}$, multipliers which localise a space-time function $F(t, x)$

to dyadic distance $\sim 2^j$ from the light cone $|\tau| = |\xi|$ on the Fourier side. Specifically, letting $\tilde{F}(\tau, \xi)$ denote the space time Fourier transform of F , while $\hat{f}(\xi)$ denotes the Fourier transform with respect to the spatial variables, and letting $\chi \in C_0^\infty(\mathbb{R}_+)$ be a smooth cutoff satisfying

$$\sum_{k \in \mathbb{Z}} \chi\left(\frac{x}{2^k}\right) = 1 \quad \text{for all } x \in \mathbb{R}_+,$$

we set

$$\widehat{P_k f}(\xi) = \chi\left(\frac{|\xi|}{2^k}\right) \hat{f}(\xi), \quad \widehat{Q_j F} = \chi\left(\frac{||\tau| - |\xi||}{2^j}\right) \tilde{F}(\tau, \xi).$$

Using these ingredients one can then define the norms

$$\|u\|_{\dot{X}^{n/2, 1/2, \infty}} := \sup_{j \in \mathbb{Z}} 2^{\frac{j}{2}} \|\nabla_x^{\frac{n}{2}} Q_j u\|_{L_{t,x}^2}, \quad \|F\|_{\dot{X}^{n/2-1, -1/2, 1}} := \sum_{j \in \mathbb{Z}} 2^{-\frac{j}{2}} \|\nabla_x^{\frac{n}{2}-1} Q_j F\|_{L_{t,x}^2}.$$

In addition to these, we rely on the classical Strichartz norms, which are the mixed-type Lebesgue norms $\|\cdot\|_{L_t^p L_x^q}$, $\frac{1}{p} + \frac{n-1}{2q} \leq \frac{n-1}{4}$, $p \geq 2$, where we shall always restrict to $n \geq 5$. Call such pairs (p, q) admissible.

We shall freely use the fact that Fourier localisers of the form $P_k Q_j$ act in bounded fashion on spaces of the form $L_t^p L_x^2$, $1 \leq p \leq \infty$; see e.g., [Tao 2001b]. We can now define a norm controlling our solutions as follows:

$$\|u\|_S := \sum_{k \in \mathbb{Z}} \sup_{(p,q) \text{ admissible}} 2^{\left(\frac{1}{p} + \frac{n}{q} - 1\right)k} \|\nabla_{t,x} P_k u\|_{L_t^p L_x^q} + \|\nabla_{t,x} P_k u\|_{\dot{X}^{n/2-1, 1/2, \infty}} =: \sum_{k \in \mathbb{Z}} \|P_k u\|_{S_k}. \tag{3-1}$$

We also introduce

$$\|F\|_N := \sum_{k \in \mathbb{Z}} \|P_k F\|_{L_t^1 \dot{H}^{n/2-1} + \dot{X}^{n/2-1, -1/2, 1}}, \tag{3-2}$$

as well as the norms

$$\|u\|_{\dot{B}_2^{r, 1}} := \sum_{k \in \mathbb{Z}} \|P_k u\|_{\dot{H}^r}.$$

Then the following inequality is by now completely standard; see e.g., [Krieger 2008; Tao 2001a; Tataru 1998]:

Proposition 3.1. $\|u\|_S \lesssim \|u[0]\|_{\dot{B}_2^{n/2, 1} \times \dot{B}_2^{n/2-1, 1}} + \|\square u\|_N. \tag{3-3}$

Sketch of proof. The fact that

$$\sum_{k \in \mathbb{Z}} \sup_{(p,q) \text{ admissible}} 2^{\left(\frac{1}{p} + \frac{n}{q} - 1\right)k} \|\nabla_{t,x} P_k u\|_{L_t^p L_x^q} \lesssim \sum_{k \in \mathbb{Z}} [\|P_k u[0]\|_{\dot{H}^{n/2} \times \dot{H}^{n/2-1}} + \|\square P_k u\|_{L_t^1 \dot{H}^{n/2-1}}]$$

is a direct consequence of the Strichartz estimates; see, e.g., [Shatah and Struwe 1998]. The fact that

$$\|\nabla_{t,x} P_k u\|_{\dot{X}^{n/2-1, 1/2, \infty}} \lesssim \|\square P_k u\|_{L_t^1 \dot{H}^{n/2-1}}$$

follows by localising the modulation and applying Holder’s inequality: setting $k = 0$, as we may by scaling invariance,

$$\begin{aligned} 2^{\frac{j}{2}} \|\nabla_{t,x} P_0 Q_j u\|_{L^2_{t,x}} &\sim 2^{\frac{j}{2}} (1 + 2^j) \|\chi(|\xi|) \widehat{Q_j u}(t, \xi)\|_{L^2_t L^2_\xi} \\ &\lesssim 2^j (1 + 2^j) \|\chi(|\xi|) \widehat{Q_j u}(t, \xi)\|_{L^2_\xi L^1_t} \lesssim \|\square P_0 Q_j u\|_{L^1_t L^2_x}. \end{aligned}$$

Furthermore, the fact that

$$\sup_{(p,q) \text{ admissible}} 2^{(\frac{1}{p} + \frac{n}{q} - 1)k} \|\nabla_{t,x} P_k Q_j u\|_{L^p_t L^q_x} \lesssim \|\square P_k Q_j u\|_{\dot{X}^{n/2-1, -1/2, 1}} \tag{3-4}$$

is a consequence of the fact that the function $P_k Q_j u$ may be represented as a weighted average of free waves, in conjunction with the standard Strichartz estimates: putting $k = 0$ as we may, write

$$\begin{aligned} \widehat{P_0 Q_j u}(\tau, \xi) &= \chi(|\xi|) \chi\left(\frac{|\tau| - |\xi|}{2^j}\right) \tilde{u}(\tau, \xi) \\ &= \chi(|\xi|) \sum_{\pm, \pm} \int \chi\left(\pm \frac{a}{2^j}\right) \tilde{u}(|\xi| + a, \xi) \delta(\pm\tau - |\xi| - a) da \\ &= \chi(|\xi|) \sum_{\pm, \pm} \int_{a \sim \pm 2^j} \widehat{e^{\pm ita} u_a^{\pm, \pm}}(\tau, \xi) da. \end{aligned}$$

Here each $u_a^{\pm, \pm}$ is a free wave and we have

$$\sum_{\pm, \pm} \int_{a \sim \pm 2^j} \|u_a^{\pm, \pm}\|_{L^\infty_t L^2_x} da \lesssim 2^{\frac{j}{2}} \sum_{\pm, \pm} \left(\int_{a \sim \pm 2^j} \|u_a^{\pm, \pm}\|_{L^\infty_t L^2_x}^2 da \right)^{\frac{1}{2}} \lesssim 2^{\frac{j}{2}} \|P_0 Q_j u\|_{L^2_{t,x}},$$

where we have used Plancherel’s theorem in the last step. This gives the case $j \leq k$ in (3-4) as a direct consequence of the Strichartz estimates, while the case $j \geq k$ follows directly from Bernstein’s inequality.

The fact that

$$\|\nabla_{t,x} P_k u\|_{\dot{X}^{n/2-1, 1/2, \infty}} \lesssim \|\square P_k u\|_{\dot{X}^{n/2-1, -1/2, 1}}$$

is immediate. This concludes our sketch of the proof of the proposition. \square

In order to deal with the nonlocal expressions such as $(-\Delta)^{\frac{1}{2}}(u \times (-\Delta)^{\frac{1}{2}}u)$, the following simple lemma shall be useful:

Lemma 3.2. Consider the bilinear expression (where $\chi_{k_j}(\cdot)$ smoothly localises to the annulus $|\xi| \sim 2^{k_j}$)

$$F(u, v)(x) := \int_{\mathbb{R}^n} \int_{\mathbb{R}^n} m(\xi, \eta) e^{ix \cdot (\xi + \eta)} \chi_{k_1}(\xi) \hat{u}(\xi) \chi_{k_2}(\eta) \hat{v}(\eta) d\xi d\eta$$

where the multiplier $m(\xi, \eta)$ is C^∞ with respect to the coordinates on the support of $\chi_{k_1}(\xi) \cdot \chi_{k_2}(\eta)$, and satisfies the pointwise bounds

$$|m(\xi, \eta)| \leq \gamma \lesssim 1, \quad |(2^{k_1} \nabla_\xi)^i (2^{k_2} \nabla_\eta)^j m(\xi, \eta)| \lesssim_{i,j} 1 \quad \text{for all } i, j.$$

Then if $\|\cdot\|_Z, \|\cdot\|_Y, \|\cdot\|_X$ are translation invariant norms with the property that

$$\|u \cdot v\|_Z \leq \|u\|_X \cdot \|v\|_Y,$$

then it follows that

$$\|F(u, v)\|_Z \lesssim \gamma^{(1-)} \|P_{k_1} u\|_X \|P_{k_2} v\|_Y,$$

where the implied constant only depends on the size of finitely many derivatives of m .

Proof. This follows by Fourier expansion of the multiplier $m(\xi, \eta)$: write

$$m(\xi, \eta) \chi_{k_1}(\xi) \chi_{k_2}(\eta) = \sum_{m \in \mathbb{Z}^n} \sum_{p \in \mathbb{Z}^n} a_{mp} e^{i(2^{-k_1} m \cdot \xi + 2^{-k_2} p \cdot \eta)},$$

where we have

$$\begin{aligned} |a_{m_1 m_2}| &\leq (2^{-k_1} |m| + 2^{-k_2} |p|)^{-Mn} \|\nabla_{\xi, \eta}^{Mn} [m(\xi, \eta) \chi_{k_1}(\xi) \chi_{k_2}(\eta)]\|_{L_{\xi, \eta}^\infty} \\ &\lesssim_{M, n} [|m| + |p|]^{-Mn}, \end{aligned}$$

while we also get the trivial bound $|a_{m_1 m_2}| \leq \gamma$. It follows that

$$\begin{aligned} F(u, v)(x) &= \sum_{\substack{m, p \in \mathbb{Z}^n \\ |m| + |p| < \gamma^{-1/(nM)}}} a_{mp} \int_{\mathbb{R}^n} \int_{\mathbb{R}^n} \hat{u}(\xi) \hat{v}(\eta) e^{i([2^{-k_1} m + x] \cdot \xi + [2^{-k_2} p + x] \cdot \eta)} d\xi d\eta \\ &\quad + \sum_{\substack{m, p \in \mathbb{Z}^n \\ |m| + |p| \geq \gamma^{-1/(nM)}}} a_{mp} \int_{\mathbb{R}^n} \int_{\mathbb{R}^n} \hat{u}(\xi) \hat{v}(\eta) e^{i([2^{-k_1} m + x] \cdot \xi + [2^{-k_2} p + x] \cdot \eta)} d\xi d\eta \end{aligned}$$

and so

$$\|F(u, v)\|_Z \lesssim \|u\|_X \|v\|_Y \left[\sum_{\substack{m, p \in \mathbb{Z}^n \\ |m| + |p| < \gamma^{-1/(nM)}}} \gamma + \sum_{\substack{m, p \in \mathbb{Z}^n \\ |m| + |p| \geq \gamma^{-1/(nM)}}} (|m| + |p|)^{-Mn} \right] \lesssim \|u\|_X \|v\|_Y \gamma^{1 - \frac{1}{M}}.$$

Here the constant M may be chosen arbitrarily large (with implied constant depending on M). □

4. Multilinear estimates

Here we gather the multilinear estimates which allow us to obtain a solution for (2-1) by means of a suitable iteration scheme:

Proposition 4.1. *Assume that u takes values in S^2 and converges to $p \in S^2$ at spatial infinity. Then using the norms $\|\cdot\|_S, \|\cdot\|_N$ introduced in the previous section, we have the bounds*

$$\|P_k [u(\nabla u \cdot \nabla u - \partial_t u \cdot \partial_t u)]\|_N \lesssim (1 + \|u\|_S) \|u\|_S \left(\sum_{k_1 \in \mathbb{Z}} 2^{-\sigma|k-k_1|} \|P_{k_1} u\|_{S_{k_1}} \right). \quad (4-1)$$

Furthermore, if \tilde{u} maps into a small neighbourhood of S^2 and $\|\tilde{u}\|_S \lesssim 1$, we have the similar bound

$$\|P_k (\Pi_{\tilde{u}_\perp} ((-\Delta)^{\frac{1}{2}} u)(u \cdot (-\Delta)^{\frac{1}{2}} u))\|_N \lesssim \prod_{v=u, \tilde{u}} (1 + \|v\|_S) \|u\|_S \left(\sum_{k_1 \in \mathbb{Z}} 2^{-\sigma|k-k_1|} \|P_{k_1} u\|_{S_{k_1}} \right), \quad (4-2)$$

as well as

$$\begin{aligned} & \| P_k (\Pi_{\tilde{u}_\perp} [u \times (-\Delta)^{\frac{1}{2}} (u \times (-\Delta)^{\frac{1}{2}} u) - u \times (u \times (-\Delta) u)]) \|_N \\ & \lesssim \prod_{v=u, \tilde{u}} (1 + \|v\|_S) \|u\|_S \left(\sum_{k_1 \in \mathbb{Z}} 2^{-\sigma|k-k_1|} \|P_{k_1} u\|_{S_{k_1}} \right). \end{aligned} \quad (4-3)$$

We also have corresponding difference estimates: assuming that $u^{(j)}$, $j = 1, 2$, map into S^2 , while $\tilde{u}^{(j)}$ map into a small neighbourhood of S^2 , then using the notation

$$\Delta_{1,2} F^{(j)} = F^{(1)} - F^{(2)}$$

we have

$$\begin{aligned} & \| \Delta_{1,2} P_k [u^{(j)} (\nabla u^{(j)} \cdot \nabla u^{(j)} - \partial_t u^{(j)} \cdot \partial_t u^{(j)})] \|_N \\ & \lesssim (1 + \max_j \|u^{(j)}\|_S) (\max_j \|u^{(j)}\|_S) \left(\sum_{k_1 \in \mathbb{Z}} 2^{-\sigma|k-k_1|} \|P_{k_1} u^{(1)} - P_{k_1} u^{(2)}\|_{S_{k_1}} \right) \\ & \quad + (1 + \max_j \|u^{(j)}\|_S) (\|u^{(1)} - u^{(2)}\|_S) \left(\max_j \sum_{k_1 \in \mathbb{Z}} 2^{-\sigma|k-k_1|} \|P_{k_1} u^{(j)}\|_{S_{k_1}} \right), \end{aligned} \quad (4-4)$$

and similarly

$$\begin{aligned} & \| P_k \Delta_{1,2} (\Pi_{\tilde{u}_\perp} ((-\Delta)^{\frac{1}{2}} u^{(j)}) (u^{(j)} \cdot (-\Delta)^{\frac{1}{2}} u^{(j)})) \|_N \\ & \lesssim \max_j \prod_{v=u^{(j)}, \tilde{u}^{(j)}} (1 + \|v\|_S) \|u^{(j)}\|_S \left(\sum_{k_1 \in \mathbb{Z}} 2^{-\sigma|k-k_1|} \|P_{k_1} u^{(1)} - P_{k_2} u^{(2)}\|_{S_{k_1}} \right) \\ & \quad + \max_j \prod_{v=u^{(j)}, \tilde{u}^{(j)}} (1 + \|v\|_S) \|u^{(1)} - u^{(2)}\|_S \left(\max_j \sum_{k_1 \in \mathbb{Z}} 2^{-\sigma|k-k_1|} \|P_{k_1} u^{(j)}\|_{S_{k_1}} \right) \\ & \quad + \max_j (1 + \|u^{(j)}\|_S) \|\tilde{u}^{(1)} - \tilde{u}^{(2)}\|_S \left(\max_j \sum_{k_1 \in \mathbb{Z}} 2^{-\sigma|k-k_1|} \|P_{k_1} u^{(j)}\|_{S_{k_1}} \right). \end{aligned} \quad (4-5)$$

The analogous difference estimate for (4-3) is similar. In fact, in all these estimates the choice $\sigma = 1$ works.

Proof. We shall only deal in detail with the case $n = 5$, since the case $n \geq 6$ is simpler due to the better decay with respect to large frequencies. Also, we note that then the desired estimates follow for a slightly different functional framework from [Sterbenz 2004]. We observe that the proof of (4-1) is really quite standard and follows for example from [Tataru 1998]. For completeness's sake, we include a simple version here.

Before giving the proof, we note that the $X^{s,b}$ -type components of our spaces are only used to prove (4-1), and not (4-2), (4-3). The key fact behind the proof of (4-1) is the identity

$$2[u_t \cdot v_t - \nabla_x u \cdot \nabla_x v] = \square(uv) - \square uv - u \square v;$$

see for example [Krieger 2008] for further discussion and earlier references.

On the other hand, the estimates (4-2), (4-3) only use Strichartz norms; there the idea is to move a derivative from a high- to a low-frequency factor, using algebraic relations such as

$$a \times (b \times c) = b(a \cdot c) - c(a \cdot b), \quad a, b, c \in \mathbb{R}^3.$$

Proof of (4-1) To achieve it, we localise the second and third factors to frequencies $\sim 2^{k_1}, 2^{k_2}$, respectively, and we shall restrict the output logarithmic frequency k to size 0. This is possible on account of the scaling invariance of the estimate. We shall obtain exponential gains in terms of these frequencies in certain cases, and summation over all allowed frequencies will result in the desired bound (4-1).

(1) *High-high interactions* $\max\{k_1, k_2\} > 10$. This is schematically written as

$$P_0[u \nabla_{t,x} u_{k_1} \nabla_{t,x} u_{k_2}].$$

Then if $k_1 = k_2 + O(1)$, we estimate this by

$$\|P_0[u \nabla_{t,x} u_{k_1} \nabla_{t,x} u_{k_2}]\|_{L_t^1 L_x^2} \lesssim \|\nabla_{t,x} u_{k_1}\|_{L_t^2 L_x^4} \|\nabla_{t,x} u_{k_2}\|_{L_t^2 L_x^4} \lesssim 2^{-\frac{3}{2}k_1} \prod_{j=1,2} \|u_{k_j}\|_{S_{k_j}}.$$

If $k_2 > k_1 + 10$, say then we estimate it by

$$\begin{aligned} \|P_0[u \nabla_{t,x} u_{k_1} \nabla_{t,x} u_{k_2}]\|_{L_t^1 L_x^2} &= \|P_0[P_{k_2+O(1)} u \nabla_{t,x} u_{k_1} \nabla_{t,x} u_{k_2}]\|_{L_t^1 L_x^2} \\ &\lesssim \|P_{k_2+O(1)} u\|_{L_t^2 L_x^4} \|\nabla_{t,x} u_{k_2}\|_{L_t^2 L_x^4} \|\nabla_{t,x} u_{k_1}\|_{L_t^\infty L_x^2 + L_{t,x}^\infty} \\ &\lesssim 2^{-\frac{5}{2}k_2} \|u_{k_2}\|_{S_{k_2}} \|u_{k_2+O(1)}\|_{S_{k_2+O(1)}} \|u_{k_1}\|_{S_{k_1}}. \end{aligned}$$

The case $k_1 > k_2 + 10$ is of course the same. Summation over the suitable ranges of k_1, k_2 implies (4-1) in this case with $\sigma = \frac{3}{2}$.

(2) *High-low interactions* $\max\{k_1, k_2\} < -10$. Here one places $\nabla_{t,x} u_{k_j}$, $j = 1, 2$, into $L_t^2 L_x^\infty$ and $u = P_{O(1)} u$ into $L_t^\infty L_x^2$.

(3) *Low-high interactions* $\max\{k_1, k_2\} \in [-10, 10]$. This is the most delicate case. We may assume that $k_1 < k_2 - 10$, since else we argue as in (1). Thus $k_2 \in [-10, 10]$. Note that then

$$\begin{aligned} \|P_0[u_{\geq k_1-10} \nabla_{t,x} u_{k_1} \nabla_{t,x} u_{k_2}]\|_{L_t^1 L_x^2} &\lesssim \|u_{\geq k_1-10}\|_{L_t^2 L_x^\infty} \|\nabla_{t,x} u_{k_1}\|_{L_t^2 L_x^\infty} \|\nabla_{t,x} u_{k_2}\|_{L_t^\infty L_x^2} \\ &\lesssim \|u\|_S \|u_{k_2}\|_{S_{k_2}} \|u_{k_1}\|_{S_{k_1}}, \end{aligned}$$

which can be summed over $k_1 < k_2 - 10$. One similarly estimates

$$P_0[Q_{\geq k_1-10} u_{< k_1-10} \nabla_{t,x} u_{k_1} \nabla_{t,x} u_{k_2}].$$

We have now reduced to estimating

$$P_0[Q_{< k_1-10} u_{< k_1-10} \partial_\alpha u_{k_1} \partial^\alpha u_{k_2}].$$

Here note that

$$\begin{aligned} \|P_0[Q_{< k_1-10} u_{< k_1-10} \partial_\alpha u_{k_1} \partial^\alpha Q_{> k_1-10} u_{k_2}]\|_{L_t^1 L_x^2} &\lesssim \|\partial_\alpha u_{k_1}\|_{L_t^2 L_x^\infty} \|\partial^\alpha Q_{> k_1-10} u_{k_2}\|_{L_{t,x}^2} \\ &\lesssim \|u_{k_1}\|_{S_{k_1}} \|u_{k_2}\|_{S_{k_2}}, \end{aligned}$$

again summable over $k_1 < k_2 - 10$. Also, we get

$$\begin{aligned} & \|P_0[Q_{<k_1-10} u_{<k_1-10} \partial_\alpha Q_{\geq k_1+10} u_{k_1} \partial^\alpha Q_{<k_1-10} u_{k_2}]\|_{X^{\frac{n}{2}-1, -\frac{1}{2}, 1}} \\ & \lesssim \sum_{j \geq k_1+10} 2^{-\frac{j}{2}} \|Q_j \nabla_{t,x} u_{k_1}\|_{L_t^2 L_x^\infty} \|\partial^\alpha Q_{<k_1-10} u_{k_2}\|_{L_t^\infty L_x^2} \lesssim \|u_{k_1}\|_{S_{k_1}} \|u_{k_2}\|_{S_{k_2}}, \end{aligned}$$

and hence it is summable over $k_1 < k_2 - 10$. Finally, we expand the expression out using its null-structure:

$$\begin{aligned} & 2P_0[Q_{<k_1-10} u_{<k_1-10} \partial_\alpha Q_{<k_1+10} u_{k_1} \partial^\alpha Q_{<k_1-10} u_{k_2}] \\ & = P_0[Q_{<k_1-10} u_{<k_1-10} \square(Q_{<k_1+10} u_{k_1} Q_{<k_1-10} u_{k_2})] \\ & \quad - P_0[Q_{<k_1-10} u_{<k_1-10} \square Q_{<k_1+10} u_{k_1} Q_{<k_1-10} u_{k_2}] \\ & \quad - P_0[Q_{<k_1-10} u_{<k_1-10} Q_{<k_1+10} u_{k_1} \square Q_{<k_1-10} u_{k_2}]. \end{aligned}$$

Then we bound each of these:

$$\begin{aligned} & P_0[Q_{<k_1-10} u_{<k_1-10} \square(Q_{<k_1+10} u_{k_1} Q_{<k_1-10} u_{k_2})] \\ & = \square P_0[Q_{<k_1-10} u_{<k_1-10} (Q_{<k_1+10} u_{k_1} Q_{<k_1-10} u_{k_2})] \\ & \quad - P_0[\nabla_{t,x} Q_{<k_1-10} u_{<k_1-10} \nabla_{t,x} (Q_{<k_1+10} u_{k_1} Q_{<k_1-10} u_{k_2})] \\ & \quad - P_0[\nabla_{t,x}^2 Q_{<k_1-10} u_{<k_1-10} (Q_{<k_1+10} u_{k_1} Q_{<k_1-10} u_{k_2})]. \end{aligned}$$

The last two terms on the right can be easily placed into $L_t^1 L_x^2$ using the $L_t^2 L_x^\infty$ norm for the low-frequency factors, while for the first term on the right, we get

$$\begin{aligned} & \|\square P_0[Q_{<k_1-10} u_{<k_1-10} (Q_{<k_1+10} u_{k_1} Q_{<k_1-10} u_{k_2})]\|_{X^{\frac{n}{2}-1, -\frac{1}{2}, 1}} \\ & \lesssim \sum_{j < k_1+20} 2^{\frac{j}{2}} \|P_0 Q_j [Q_{<k_1-10} u_{<k_1-10} (Q_{<k_1+10} u_{k_1} Q_{<k_1-10} u_{k_2})]\|_{L_{t,x}^2} \\ & \lesssim 2^{\frac{k_1}{2}} \|Q_{<k_1+10} u_{k_1}\|_{L_t^2 L_x^\infty} \|Q_{<k_1-10} u_{k_2}\|_{L_t^\infty L_x^2} \\ & \lesssim \|u_{k_1}\|_{S_{k_1}} \|u_{k_2}\|_{S_{k_2}}. \end{aligned}$$

Further, we get

$$\begin{aligned} & \|P_0[Q_{<k_1-10} u_{<k_1-10} \square Q_{<k_1+10} u_{k_1} Q_{<k_1-10} u_{k_2}]\|_{L_t^1 L_x^2} \\ & \lesssim \|\square Q_{<k_1+10} u_{k_1}\|_{L_t^2 L_x^4} \|Q_{<k_1-10} u_{k_2}\|_{L_t^2 L_x^4} \\ & \lesssim 2^{\frac{k_1}{4}} \|u_{k_1}\|_{S_{k_1}} \|u_{k_2}\|_{S_{k_2}}. \end{aligned}$$

To close things, we also get

$$\begin{aligned} & \|P_0[Q_{<k_1-10} u_{<k_1-10} Q_{<k_1+10} u_{k_1} \square Q_{<k_1-10} u_{k_2}]\|_{L_t^1 L_x^2} \\ & \lesssim \|Q_{<k_1+10} u_{k_1}\|_{L_t^2 L_x^\infty} \|\square Q_{<k_1-10} u_{k_2}\|_{L_{t,x}^2} \\ & \lesssim \|u_{k_1}\|_{S_{k_1}} \|u_{k_2}\|_{S_{k_2}} \end{aligned}$$

and the desired bound follows again by summing over $k_1 < k_2 - 10$.

Proof of (4-2) Here we shall be able to get by only using Strichartz-type norms, by taking advantage of the condition $u \cdot u = 1$. Using the standard Littlewood–Paley trichotomy, we have

$$0 = u \cdot u - p \cdot p = \sum_{|k_1 - k_2| \leq 10} u_{k_1} u_{k_2} + 2 \sum_{k_1} u_{k_1} \cdot u_{<k_1-10}. \quad (4-6)$$

This implies

$$0 = \sum_{|k_1 - k_2| < 10} (-\Delta)^{\frac{1}{2}} (u_{k_1} u_{k_2}) + 2 \sum_{k_1} (-\Delta)^{\frac{1}{2}} (u_{k_1} \cdot u_{<k_1-10}). \quad (4-7)$$

Here the first term is better, since the outer derivative falls on the low-frequency output. We shall use this to replace the second term on the right by the first. Write

$$\begin{aligned} \Pi_{\tilde{u}_\perp} ((-\Delta)^{\frac{1}{2}} u) (u \cdot (-\Delta)^{\frac{1}{2}} u) &= \sum_{|k_1 - k_2| \leq 10} \Pi_{\tilde{u}_\perp} ((-\Delta)^{\frac{1}{2}} u) (u_{k_1} \cdot (-\Delta)^{\frac{1}{2}} u_{k_2}) \\ &+ \sum_{k_1} \Pi_{\tilde{u}_\perp} ((-\Delta)^{\frac{1}{2}} u) (u_{k_1} \cdot (-\Delta)^{\frac{1}{2}} u_{<k_1-10}) \\ &+ \sum_{k_2} \Pi_{\tilde{u}_\perp} ((-\Delta)^{\frac{1}{2}} u) (u_{<k_2-10} \cdot (-\Delta)^{\frac{1}{2}} u_{k_2}). \end{aligned} \quad (4-8)$$

Then for the first term on the right we infer

$$\begin{aligned} &\left\| P_0 \left[\sum_{|k_1 - k_2| \leq 10} \Pi_{\tilde{u}_\perp} ((-\Delta)^{\frac{1}{2}} u) (u_{k_1} \cdot (-\Delta)^{\frac{1}{2}} u_{k_2}) \right] \right\|_{L_t^1 L_x^2} \\ &\lesssim \sum_{\substack{|k_1 - k_2| \leq 10 \\ k_1 < -20}} \| P_{[-20, 20]} (\Pi_{\tilde{u}_\perp} ((-\Delta)^{\frac{1}{2}} u)) \|_{L_t^\infty L_x^2} \| u_{k_1} \|_{L_t^2 L_x^\infty} \| (-\Delta)^{\frac{1}{2}} u_{k_2} \|_{L_t^2 L_x^\infty} \\ &+ \sum_{\substack{|k_1 - k_2| \leq 10 \\ k_1 \geq -20}} \| \Pi_{\tilde{u}_\perp} ((-\Delta)^{\frac{1}{2}} u) \|_{L_t^\infty L_x^2 + L_{t,x}^\infty} \| u_{k_1} \|_{L_t^2 L_x^4} \| (-\Delta)^{\frac{1}{2}} u_{k_2} \|_{L_t^2 L_x^4}. \end{aligned}$$

Then using a further elementary frequency decomposition it is easy to see (see the Appendix) that

$$\begin{aligned} \| P_{[-20, 20]} (\Pi_{\tilde{u}_\perp} ((-\Delta)^{\frac{1}{2}} u)) \|_{L_t^\infty L_x^2} &\lesssim \sum_{k_3 \in \mathbb{Z}} 2^{-|k_3|} \| P_{k_3} u \|_{S_{k_3}} (1 + \|\tilde{u}\|_S), \\ \| (\Pi_{\tilde{u}_\perp} ((-\Delta)^{\frac{1}{2}} u)) \|_{L_t^\infty L_x^2 + L_{t,x}^\infty} &\lesssim \sum_{k_3 \in \mathbb{Z}} 2^{-|k_3|} \| P_{k_3} u \|_{S_{k_3}} (1 + \|\tilde{u}\|_S), \end{aligned}$$

and so we obtain that

$$\begin{aligned} &\sum_{\substack{|k_1 - k_2| \leq 10 \\ k_1 < -20}} \| P_{[-20, 20]} (\Pi_{\tilde{u}_\perp} ((-\Delta)^{\frac{1}{2}} u)) \|_{L_t^\infty L_x^2} \| u_{k_1} \|_{L_t^2 L_x^\infty} \| (-\Delta)^{\frac{1}{2}} u_{k_2} \|_{L_t^2 L_x^\infty} \\ &\lesssim \sum_{\substack{|k_1 - k_2| \leq 10 \\ k_1 < -20}} 2^{\frac{k_2 - k_1}{2}} \prod_{j=1,2} \| P_{k_j} u \|_{S_{k_j}} \left(\sum_{k_3 \in \mathbb{Z}} 2^{-|k_3|} \| P_{k_3} u \|_{S_{k_3}} (1 + \|\tilde{u}\|_S) \right) \\ &\lesssim \left(\sum_{k_3 \in \mathbb{Z}} 2^{-|k_3|} \| P_{k_3} u \|_{S_{k_3}} \right) \| u \|_S^2 (1 + \|\tilde{u}\|_S), \end{aligned}$$

as well as

$$\begin{aligned} \sum_{\substack{|k_1-k_2|\leq 10 \\ k_1\geq -20}} \|\Pi_{\tilde{u}_\perp}((-\Delta)^{\frac{1}{2}}u)\|_{L_t^\infty L_x^2+L_{t,x}^\infty} \|u_{k_1}\|_{L_t^2 L_x^4} \|(-\Delta)^{\frac{1}{2}}u_{k_2}\|_{L_t^2 L_x^4} \\ \lesssim \left(\sum_{\substack{|k_1-k_2|\leq 10 \\ k_1\geq -20}} 2^{-\frac{5}{2}k_1} \|u_{k_1}\|_{S_{k_1}} \|u_{k_2}\|_{S_{k_2}} \right) \sum_{k_3\in\mathbb{Z}} 2^{-|k_3|} \|P_{k_3}u\|_{S_{k_3}} (1 + \|\tilde{u}\|_S) \\ \lesssim \left(\sum_{k_3\in\mathbb{Z}} 2^{-|k_3|} \|P_{k_3}u\|_{S_{k_3}} \right) \|u\|_S^2 (1 + \|\tilde{u}\|_S). \end{aligned}$$

This concludes the required bound for the first term on the right-hand side of (4-8).

Now we pass to the second term. We write it as a sum of three terms:

$$\begin{aligned} \sum_{k_1} \Pi_{\tilde{u}_\perp}((-\Delta)^{\frac{1}{2}}u)(u_{k_1} \cdot (-\Delta)^{\frac{1}{2}}u_{<k_1-10}) &= \sum_{k_1\geq 5} \Pi_{\tilde{u}_\perp}((-\Delta)^{\frac{1}{2}}u)(u_{k_1} \cdot (-\Delta)^{\frac{1}{2}}u_{<k_1-10}) \\ &+ \sum_{k_1\in[-5,5]} \Pi_{\tilde{u}_\perp}((-\Delta)^{\frac{1}{2}}u)(u_{k_1} \cdot (-\Delta)^{\frac{1}{2}}u_{<k_1-10}) \\ &+ \sum_{k_1<-5} \Pi_{\tilde{u}_\perp}((-\Delta)^{\frac{1}{2}}u)(u_{k_1} \cdot (-\Delta)^{\frac{1}{2}}u_{<k_1-10}). \end{aligned}$$

Then we get

$$\begin{aligned} \left\| P_0 \left(\sum_{k_1\geq 5} \Pi_{\tilde{u}_\perp}((-\Delta)^{\frac{1}{2}}u)(u_{k_1} \cdot (-\Delta)^{\frac{1}{2}}u_{<k_1-10}) \right) \right\|_{L_t^1 L_x^2} \\ \lesssim \sum_{k_1\geq 5} \|P_{[k_1-5, k_1+5]}(\Pi_{\tilde{u}_\perp}((-\Delta)^{\frac{1}{2}}u))\|_{L_t^\infty L_x^2} \|u_{k_1}\|_{L_t^2 L_x^\infty} \|(-\Delta)^{\frac{1}{2}}u_{<k_1-10}\|_{L_t^2 L_x^\infty} \\ \lesssim \sum_{k_1\geq 5, k_2<k_1-10} \sum_{k_3} 2^{-\frac{3}{2}|k_3|} \|P_{k_3}u\|_{S_{k_3}} (1 + \|\tilde{u}\|_S) \|u_{k_1}\|_{S_{k_1}} \|u_{k_2}\|_{S_{k_2}} \\ \lesssim \left(\sum_{k_3} 2^{-\frac{3}{2}|k_3|} \|P_{k_3}u\|_{S_{k_3}} \right) (1 + \|\tilde{u}\|_S) \|u\|_S^2. \end{aligned}$$

Similarly, for the term of intermediate k_1 , we have

$$\begin{aligned} \left\| P_0 \left[\sum_{k_1\in[-5,5]} \Pi_{\tilde{u}_\perp}((-\Delta)^{\frac{1}{2}}u)(u_{k_1} \cdot (-\Delta)^{\frac{1}{2}}u_{<k_1-10}) \right] \right\|_{L_t^1 L_x^2} \\ \lesssim \sum_{k_1\in[-5,5]} \|P_{<10}\Pi_{\tilde{u}_\perp}((-\Delta)^{\frac{1}{2}}u)\|_{L_t^2 L_x^\infty} \|u_{k_1}\|_{S_{k_1}} \|(-\Delta)^{\frac{1}{2}}u_{<k_1-10}\|_{L_t^2 L_x^\infty}, \end{aligned}$$

and one closes by observing (see the Appendix for the first bound) that

$$\|P_{<10}\Pi_{\tilde{u}_\perp}((-\Delta)^{\frac{1}{2}}u)\|_{L_t^2 L_x^\infty} \lesssim (\|\tilde{u}\|_S + 1) \|u\|_S, \quad \|(-\Delta)^{\frac{1}{2}}u_{<k_1-10}\|_{L_t^2 L_x^\infty} \lesssim \|u\|_S.$$

Finally, for the range of low $k_1 < -5$, we place both u_{k_1} and $(-\Delta)^{\frac{1}{2}}u_{<k_1-10}$ into $L_t^2 L_x^\infty$ and observe that

$$\|u_{k_1}\|_{L_t^2 L_x^\infty} \|(-\Delta)^{\frac{1}{2}}u_{<k_1-10}\|_{L_t^2 L_x^\infty} \lesssim \|u_{k_1}\|_{S_{k_1}} \|u\|_S.$$

Then we close by using that

$$P_0\left(\sum_{k_1 < -5} \Pi_{\tilde{u}_\perp}((-\Delta)^{\frac{1}{2}}u)(u_{k_1} \cdot (-\Delta)^{\frac{1}{2}}u_{<k_1-10})\right) = P_0\left(\sum_{k_1 < -5} P_{[-2,2]}(\Pi_{\tilde{u}_\perp}((-\Delta)^{\frac{1}{2}}u))(u_{k_1} \cdot (-\Delta)^{\frac{1}{2}}u_{<k_1-10})\right),$$

as well as

$$\|P_{[-2,2]}(\Pi_{\tilde{u}_\perp}((-\Delta)^{\frac{1}{2}}u))\|_{L_t^\infty L_x^2} \lesssim (1 + \|\tilde{u}\|_S) \left(\sum_{k_3} 2^{-\frac{3}{2}|k_3|} \|P_{k_3}u\|_{S_{k_3}}\right).$$

This concludes the required bound for the second term on the right in (4-8).

Finally, the third term in (4-8) is the most delicate, as the derivative $(-\Delta)^{\frac{1}{2}}$ lands on the higher-frequency term u_{k_2} . To deal with it, we note, using Lemma 3.2, that the difference

$$\sum_{k_2} \Pi_{\tilde{u}_\perp}((-\Delta)^{\frac{1}{2}}u)(u_{<k_2-10} \cdot (-\Delta)^{\frac{1}{2}}u_{k_2}) - \sum_{k_2} \Pi_{\tilde{u}_\perp}((-\Delta)^{\frac{1}{2}}u)(-\Delta)^{\frac{1}{2}}(u_{<k_2-10} \cdot u_{k_2})$$

can be estimated like the second term on the right in (4-8), and hence it suffices to bound

$$\sum_{k_2} \Pi_{\tilde{u}_\perp}((-\Delta)^{\frac{1}{2}}u)(-\Delta)^{\frac{1}{2}}(u_{<k_2-10} \cdot u_{k_2}) = - \sum_{|k_3-k_4| < 10} \frac{1}{2} \Pi_{\tilde{u}_\perp}((-\Delta)^{\frac{1}{2}}u)(-\Delta)^{\frac{1}{2}}(u_{k_3} \cdot u_{k_4}),$$

where we have used (4-6). This term is again straightforward to estimate: we have

$$\begin{aligned} & \left\| P_0 \left[\sum_{\substack{|k_3-k_4| < 10 \\ k_3 < -20}} \frac{1}{2} \Pi_{\tilde{u}_\perp}((-\Delta)^{\frac{1}{2}}u)(-\Delta)^{\frac{1}{2}}(u_{k_3} \cdot u_{k_4}) \right] \right\|_{L_t^1 L_x^2} \\ & \lesssim \sum_{\substack{|k_3-k_4| < 10 \\ k_3 < -20}} \|P_{[-10,10]}[\Pi_{\tilde{u}_\perp}((-\Delta)^{\frac{1}{2}}u)]\|_{L_t^\infty L_x^2} \|(-\Delta)^{\frac{1}{2}}(u_{k_3} \cdot u_{k_4})\|_{L_t^1 L_x^\infty}, \end{aligned}$$

and we close for the case $k_3 < -20$ by observing that

$$\sum_{\substack{|k_3-k_4| < 10 \\ k_3 < -20}} \|(-\Delta)^{\frac{1}{2}}(u_{k_3} \cdot u_{k_4})\|_{L_t^1 L_x^\infty} \lesssim \sum_{\substack{|k_3-k_4| < 10 \\ k_3 < -20}} 2^{k_3} \|u_{k_3}\|_{L_t^2 L_x^\infty} \|u_{k_4}\|_{L_t^2 L_x^\infty} \lesssim \|u\|_S^2,$$

as well as

$$\|P_{[-10,10]}[\Pi_{\tilde{u}_\perp}((-\Delta)^{\frac{1}{2}}u)]\|_{L_t^\infty L_x^2} \lesssim (1 + \|\tilde{u}\|_S) \sum_{k_3} 2^{-\frac{3}{2}|k_3|} \|P_{k_3}u\|_{S_{k_3}}.$$

On the other hand, if $k_3 > -20$, we place both $u_{k_3,4}$ into $L_t^2 L_x^4$. We omit the simple details. This finally concludes the bound of estimate (4-2).

Proof of (4-3) We commence by observing that we may in fact get rid of the outer operator $\Pi_{\tilde{u}_\perp}$, since one easily checks that

$$\|P_0[\Pi_{\tilde{u}_\perp} F]\|_{L_t^1 L_x^2} \lesssim (1 + \|\tilde{u}\|_S) \sum_{k_1} 2^{-|k_1|} \|P_{k_1} F\|_{L_t^1 \dot{H}^{n/2-1}}.$$

Then assuming that we have proved the bound

$$\|P_{k_1} F\|_{L_t^1 \dot{H}^{n/2-1}} \lesssim \sum_{k_2} 2^{-\sigma|k_1-k_2|} \|P_{k_2} u\|_{S_{k_2}}$$

for some $\sigma > 1$, we then infer the bound

$$\|P_0[\Pi_{\tilde{u}^\perp} F]\|_{L_t^1 L_x^2} \lesssim (1 + \|\tilde{u}\|_S) \sum_{k_2} 2^{-|k_2|} \|P_{k_2} u\|_{S_{k_2}}.$$

Next, localising the last two factors to dyadic frequencies, and the output to frequency ~ 1 as we may, we arrive at the expression

$$P_0[u \times (-\Delta)^{\frac{1}{2}}(u_{k_1} \times (-\Delta)^{\frac{1}{2}}u_{k_2}) - u \times (u_{k_1} \times (-\Delta)u_{k_2})].$$

Then we first dispose of the easy cases:

Both frequencies large: $\max\{k_1, k_2\} > 10$. If $k_1 = k_2 + O(1)$, we simply place both high-frequency factors into $L_t^2 L_x^4$, resulting in

$$\begin{aligned} \|P_0[u \times (-\Delta)^{\frac{1}{2}}(u_{k_1} \times (-\Delta)^{\frac{1}{2}}u_{k_2}) - u \times (u_{k_1} \times (-\Delta)u_{k_2})]\|_{L_t^1 L_x^2} &\lesssim 2^{2k_1} \|P_{k_1} u\|_{L_t^2 L_x^4} \|u_{k_2}\|_{L_t^2 L_x^4} \\ &\lesssim 2^{2k_1 - (1 + \frac{5}{2})k_1} \prod_{j=1,2} \|u_{k_j}\|_{S_{k_j}}, \end{aligned}$$

whence we have

$$\begin{aligned} &\left\| \sum_{k_1=k_2+O(1)>10} P_0[u \times (-\Delta)^{\frac{1}{2}}(u_{k_1} \times (-\Delta)^{\frac{1}{2}}u_{k_2}) - u \times (u_{k_1} \times (-\Delta)u_{k_2})] \right\|_{L_t^1 L_x^2} \\ &\lesssim \sum_{k_1=k_2+O(1)>10} 2^{-\frac{3}{2}|k_1|} \prod_{j=1,2} \|u_{k_j}\|_{S_{k_j}} \lesssim \left(\sum_{k_1} 2^{-\frac{3}{2}|k_1|} \|P_{k_1} u\|_{S_{k_1}} \right) \|u\|_S. \end{aligned}$$

On the other hand, if $k_2 \gg k_1$, we use

$$\begin{aligned} &P_0[u \times (-\Delta)^{\frac{1}{2}}(u_{k_1} \times (-\Delta)^{\frac{1}{2}}u_{k_2}) - u \times (u_{k_1} \times (-\Delta)u_{k_2})] \\ &= P_0[P_{k_2+O(1)}u \times (-\Delta)^{\frac{1}{2}}(u_{k_1} \times (-\Delta)^{\frac{1}{2}}u_{k_2}) - P_{k_2+O(1)}u \times (u_{k_1} \times (-\Delta)u_{k_2})]. \end{aligned}$$

Then place the first and third factors into $L_t^2 L_x^4$ and the middle factor into $L_t^\infty L_x^2 + L_{t,x}^\infty$. The case $k_2 \ll k_1$ is similar.

Both frequencies small: $\max\{k_1, k_2\} < -10$. Here we observe that Lemma 3.2 allows us to place one derivative $(-\Delta)^{\frac{1}{2}}$ onto the factor u_{k_1} , even if $k_1 < k_2 - 10$. Thus we reduce to bounding the schematic expression

$$P_0[P_{[-5,5]}u \nabla_x u_{k_1} \nabla_x u_{k_2}],$$

which is straightforward since we can place the second and third factors into $L_t^2 L_x^\infty$. We omit the simple details.

One frequency intermediate, the other small: $\max\{k_1, k_2\} \in [-10, 10]$. This case is a bit more difficult, and we shall exploit the geometric structure of the expression. We split this further into two cases:

(i) $k_1 \in [-10, 10]$, $k_2 < 10$. Here the difference structure inherent in the term is not helpful. In fact, we can immediately estimate

$$\begin{aligned} \|P_0[u \times (u_{k_1} \times (-\Delta)u_{k_2})]\|_{L_t^1 L_x^2} &\lesssim \|u_{k_1}\|_{L_t^2 L_x^4} \|(-\Delta)u_{k_2}\|_{L_t^2 L_x^4} \\ &\lesssim 2^{\frac{k_2}{4}} \|u_{k_2}\|_{S_{k_2}} \|u_{k_1}\|_{S_{k_1}}, \end{aligned}$$

and here of course we can sum over $k_2 < 10$ to infer the desired bound. Next, using Lemma 3.2 allows us to replace the term

$$P_0[u \times (-\Delta)^{\frac{1}{2}}(u_{k_1} \times (-\Delta)^{\frac{1}{2}}u_{k_2})]$$

by

$$P_0[u \times ((-\Delta)^{\frac{1}{2}}u_{k_1} \times (-\Delta)^{\frac{1}{2}}u_{k_2})]$$

up to a term which is estimated like $P_0[u \times (u_{k_1} \times (-\Delta)u_{k_2})]$. Before exploiting the algebraic structure of the term above, we reduce the first factor u to frequency $< 2^{k_2-10}$, which we can on account of

$$\begin{aligned} \|P_0[u_{\geq k_2-10} \times ((-\Delta)^{\frac{1}{2}}u_{k_1} \times (-\Delta)^{\frac{1}{2}}u_{k_2})]\|_{L_t^1 L_x^2} &\lesssim \|u_{\geq k_2-10}\|_{L_t^2 L_x^\infty} \|(-\Delta)^{\frac{1}{2}}u_{k_1}\|_{L_t^\infty L_x^2} \|(-\Delta)^{\frac{1}{2}}u_{k_2}\|_{L_t^2 L_x^\infty} \\ &\lesssim \|u_{k_1}\|_{S_{k_1}} \|u_{k_2}\|_{S_{k_2}} \|u\|_S. \end{aligned}$$

Summing over $k_2 < 10$ and recalling that $k_1 \in [-10, 10]$ leads to the desired bound.

Consider now the expression

$$P_0[u_{<k_2-10} \times ((-\Delta)^{\frac{1}{2}}u_{k_1} \times (-\Delta)^{\frac{1}{2}}u_{k_2})].$$

Write this as

$$\begin{aligned} &P_0[u_{<k_2-10} \times ((-\Delta)^{\frac{1}{2}}u_{k_1} \times (-\Delta)^{\frac{1}{2}}u_{k_2})] \\ &= P_0[(-\Delta)^{\frac{1}{2}}u_{k_1}(u_{<k_2-10} \cdot (-\Delta)^{\frac{1}{2}}u_{k_2}) - (-\Delta)^{\frac{1}{2}}u_{k_2}(u_{<k_2-10} \cdot (-\Delta)^{\frac{1}{2}}u_{k_1})]. \end{aligned}$$

In order to estimate this, we use a frequency-localised version of (4-6). Specifically, we have

$$0 = 2u_k \cdot u_{<k-10} + \sum_{k_1=k_2+O(1)} P_k(u_{k_1} \cdot u_{k_2}) + 2^{-k} L(u_k, \nabla_x u_{<k-10}), \quad (4-9)$$

where L is a bilinear operator of the form used in Lemma 3.2 with a bounded kernel $m(\xi, \eta)$. We conclude the schematic relation

$$(-\Delta)^{\frac{1}{2}}u_k \cdot u_{<k-10} = -\frac{1}{2}(-\Delta)^{\frac{1}{2}} \sum_{k_1=k_2+O(1)} P_k(u_{k_1} \cdot u_{k_2}) + L(u_k, \nabla_x u_{<k-10}).$$

It follows that we can write

$$\begin{aligned} &P_0[(-\Delta)^{\frac{1}{2}}u_{k_1}(u_{<k_2-10} \cdot (-\Delta)^{\frac{1}{2}}u_{k_2})] \\ &= -\frac{1}{2}P_0\left[(-\Delta)^{\frac{1}{2}}u_{k_1} \sum_{k_3=k_4+O(1)} (-\Delta)^{\frac{1}{2}}P_{k_2}(u_{k_3} \cdot u_{k_4})\right] + P_0[(-\Delta)^{\frac{1}{2}}u_{k_1} L(u_{k_2}, \nabla_x u_{<k_2-10})], \end{aligned}$$

and here we have (keeping in mind that $k_1 \in [-10, 10]$)

$$\begin{aligned} & \left\| P_0 \left[(-\Delta)^{\frac{1}{2}} u_{k_1} \sum_{k_3=k_4+O(1)} (-\Delta)^{\frac{1}{2}} P_{k_2}(u_{k_3} \cdot u_{k_4}) \right] \right\|_{L_t^1 L_x^2} \\ & \lesssim 2^{k_2} \sum_{k_3=k_4+O(1) \geq k_2} \|(-\Delta)^{\frac{1}{2}} u_{k_1}\|_{L_t^\infty L_x^2} \|u_{k_3}\|_{L_t^2 L_x^\infty} \|u_{k_4}\|_{L_t^2 L_x^\infty} \\ & \lesssim \|(-\Delta)^{\frac{1}{2}} u_{k_1}\|_{L_t^\infty L_x^2} \sum_{k_3=k_4+O(1) \geq k_2} 2^{k_2-k_3} \|u_{k_3}\|_{S_{k_3}} \|u_{k_4}\|_{S_{k_4}}, \end{aligned}$$

and here we can sum over $k_2 < 10$ to arrive at an upper bound of $\lesssim \|u_{k_1}\|_{S_{k_1}} \|u\|_S^2$, as desired. We also have the simple bound

$$\begin{aligned} & \left\| P_0 \left[(-\Delta)^{\frac{1}{2}} u_{k_1} L(u_{k_2}, \nabla_x u_{<k_2-10}) \right] \right\|_{L_t^1 L_x^2} \lesssim \|(-\Delta)^{\frac{1}{2}} u_{k_1}\|_{L_t^\infty L_x^2} \|u_{k_2}\|_{L_t^2 L_x^\infty} \|\nabla_x u_{<k_2-10}\|_{L_t^2 L_x^\infty} \\ & \lesssim \|(-\Delta)^{\frac{1}{2}} u_{k_1}\|_{L_t^\infty L_x^2} \|u_{k_2}\|_{S_{k_2}} \|u\|_S, \end{aligned}$$

and summing over $k_2 < 10$, we arrive again at the bound

$$\lesssim \|u_{k_1}\|_{S_{k_1}} \|u\|_S^2.$$

This concludes the case (i).

(ii) $k_2 \in [-10, 10], k_1 < 10$. Proceeding in analogy to case (i), we immediately reduce to the expression

$$P_0[u_{<k_1-10} \times (-\Delta)^{\frac{1}{2}}(u_{k_1} \times (-\Delta)^{\frac{1}{2}}u_{k_2}) - u_{<k_1-10} \times (u_{k_1} \times (-\Delta)u_{k_2})].$$

Here we first note that on account of Lemma 3.2 we have

$$\begin{aligned} & \left\| P_0 \left[u_{<k_1-10} \times (-\Delta)^{\frac{1}{2}}(u_{k_1} \times (-\Delta)^{\frac{1}{2}}u_{k_2}) - (-\Delta)^{\frac{1}{2}}(u_{<k_1-10} \times (u_{k_1} \times (-\Delta)^{\frac{1}{2}}u_{k_2})) \right] \right\|_{L_t^1 L_x^2} \\ & \lesssim \|(-\Delta)^{\frac{1}{2}} u_{<k_1-10}\|_{L_t^2 L_x^\infty} \|u_{k_1}\|_{L_t^2 L_x^\infty} \|(-\Delta)^{\frac{1}{2}} u_{k_2}\|_{L_t^\infty L_x^2} \\ & \lesssim \|u\|_S \|u_{k_1}\|_{S_{k_1}} \|u_{k_2}\|_{S_{k_2}}. \end{aligned}$$

Then summation over $k_1 < 10$ gives the required bound.

Next, we expand out

$$\begin{aligned} & P_0 \left[(-\Delta)^{\frac{1}{2}} (u_{<k_1-10} \times (u_{k_1} \times (-\Delta)^{\frac{1}{2}}u_{k_2})) - u_{<k_1-10} \times (u_{k_1} \times (-\Delta)u_{k_2}) \right] \\ & = P_0 (-\Delta)^{\frac{1}{2}} (u_{k_1} (u_{<k_1-10} \cdot (-\Delta)^{\frac{1}{2}}u_{k_2}) - (-\Delta)^{\frac{1}{2}}u_{k_2} (u_{<k_1-10} \cdot u_{k_1})) \\ & \quad - P_0 (u_{k_1} (u_{<k_1-10} \cdot (-\Delta)u_{k_2}) - (-\Delta)u_{k_2} (u_{<k_1-10} \cdot u_{k_1})). \quad (4-10) \end{aligned}$$

Then pairing up these last four terms suitably, we have

$$\begin{aligned} & P_0 (-\Delta)^{\frac{1}{2}} (u_{k_1} (u_{<k_1-10} \cdot (-\Delta)^{\frac{1}{2}}u_{k_2})) - P_0 (u_{k_1} (u_{<k_1-10} \cdot (-\Delta)u_{k_2})) \\ & = P_0 (-\Delta)^{\frac{1}{2}} (u_{k_1} (-\Delta)^{\frac{1}{2}}(u_{<k_1-10} \cdot u_{k_2})) - P_0 (u_{k_1} (-\Delta)(u_{<k_1-10} \cdot u_{k_2})) + u_{k_1} L(\nabla_x u_{<k_1-10}, u_{k_2}) \\ & = L((-\Delta)^{\frac{1}{2}}u_{k_1}, (-\Delta)^{\frac{1}{2}}(u_{<k_1-10} \cdot u_{k_2})) + u_{k_1} L(\nabla_x u_{<k_1-10}, u_{k_2}). \end{aligned}$$

The last term is straightforward since

$$\begin{aligned} \|u_{k_1} L(\nabla_x u_{<k_1-10}, u_{k_2})\|_{L_t^1 L_x^2} &\lesssim \|u_{k_1}\|_{L_t^2 L_x^\infty} \|\nabla_x u_{<k_1-10}\|_{L_t^2 L_x^\infty} \|u_{k_2}\|_{L_t^\infty L_x^2} \\ &\lesssim \|u\|_S \|u_{k_1}\|_{S_{k_1}} \|u_{k_2}\|_{S_{k_2}}, \end{aligned}$$

and we can sum over $k_1 < 10$. Further, we see that

$$L((-\Delta)^{\frac{1}{2}} u_{k_1}, (-\Delta)^{\frac{1}{2}} (u_{<k_1-10} \cdot u_{k_2})) = L((-\Delta)^{\frac{1}{2}} u_{k_1}, (-\Delta)^{\frac{1}{2}} (u_{<k_2-10} \cdot u_{k_2})) + \text{error},$$

where the term error here is estimated exactly like the previous term. But then taking advantage of (4-9), we find

$$\begin{aligned} L((-\Delta)^{\frac{1}{2}} u_{k_1}, (-\Delta)^{\frac{1}{2}} (u_{<k_2-10} \cdot u_{k_2})) \\ = -\frac{1}{2} \sum_{k_3=k_4+O(1)>k_2} L((-\Delta)^{\frac{1}{2}} u_{k_1}, (-\Delta)^{\frac{1}{2}} P_{k_2}(u_{k_3} \cdot u_{k_4})) \\ + 2^{-k_2} L((-\Delta)^{\frac{1}{2}} u_{k_1}, (-\Delta)^{\frac{1}{2}} L(\nabla_x u_{<k_2-10}, u_{k_2})). \end{aligned}$$

Then we have

$$\begin{aligned} \left\| -\frac{1}{2} \sum_{k_3=k_4+O(1)>k_2} L((-\Delta)^{\frac{1}{2}} u_{k_1}, (-\Delta)^{\frac{1}{2}} P_{k_2}(u_{k_3} \cdot u_{k_4})) \right\|_{L_t^1 L_x^2} \\ \lesssim \sum_{k_3=k_4+O(1)>k_2} 2^{k_2} \|(-\Delta)^{\frac{1}{2}} u_{k_1}\|_{L_t^2 L_x^\infty} \|u_{k_3}\|_{L_t^2 L_x^\infty} \|u_{k_4}\|_{L_t^\infty L_x^2}. \end{aligned}$$

The preceding sum can be further bounded by

$$\begin{aligned} &\lesssim \sum_{k_3=k_4+O(1)>k_2} 2^{k_2} 2^{\frac{k_1-k_3}{2}} 2^{-\frac{5}{2}k_4} \|u_{k_1}\|_{S_{k_1}} \|u_{k_3}\|_{S_{k_3}} \|u_{k_4}\|_{S_{k_4}} \\ &\lesssim \left(\sum_{k_1} 2^{-|k_4-k_2|} \|u_{k_4}\|_{S_{k_4}} \|u\|_S \right) \|u_{k_1}\|_{S_{k_1}}. \end{aligned}$$

This can be summed over $k_1 < 10$ to yield the desired kind of bound.

Finally, we have the simpler bound

$$\begin{aligned} \|2^{-k_2} L((-\Delta)^{\frac{1}{2}} u_{k_1}, (-\Delta)^{\frac{1}{2}} L(\nabla_x u_{<k_2-10}, u_{k_2}))\|_{L_t^1 L_x^2} \\ \lesssim \|(-\Delta)^{\frac{1}{2}} u_{k_1}\|_{L_t^2 L_x^\infty} \|\nabla_x u_{<k_2-10}\|_{L_t^2 L_x^\infty} \|u_{k_2}\|_{L_t^\infty L_x^2}, \end{aligned}$$

which after summation over $k_1 < 10$ is again bounded by $\lesssim \|u\|_S^2 \|u_{k_2}\|_{S_{k_2}}$.

Returning to (4-10), it remains to bound the difference

$$\begin{aligned} P_0(-\Delta)^{\frac{1}{2}} [(-\Delta)^{\frac{1}{2}} u_{k_2} (u_{<k_1-10} \cdot u_{k_1})] - P_0 [(-\Delta) u_{k_2} (u_{<k_1-10} \cdot u_{k_1})] \\ = -L \left((-\Delta)^{\frac{1}{2}} u_{k_2}, \sum_{k_3=k_4+O(1)>k_1} (-\Delta)^{\frac{1}{2}} P_{k_1}(u_{k_3} \cdot u_{k_4}) \right) \\ + L \left((-\Delta)^{\frac{1}{2}} u_{k_2}, (-\Delta)^{\frac{1}{2}} 2^{-k_1} L(\nabla_x u_{<k_1-10}, u_{k_1}) \right). \end{aligned}$$

Then the first term is bounded by

$$\begin{aligned} & \left\| \frac{1}{2} L \left((-\Delta)^{\frac{1}{2}} u_{k_2}, \sum_{k_3=k_4+O(1)>k_1} (-\Delta)^{\frac{1}{2}} P_{k_1}(u_{k_3} \cdot u_{k_4}) \right) \right\|_{L_t^1 L_x^2} \\ & \lesssim 2^{k_1} \sum_{k_3=k_4+O(1)>k_1} \| (-\Delta)^{\frac{1}{2}} u_{k_2} \|_{L_t^\infty L_x^2} \| u_{k_3} \|_{L_t^2 L_x^\infty} \| u_{k_4} \|_{L_t^2 L_x^\infty} \\ & \lesssim \| u_{k_2} \|_{S_{k_2}} \sum_{k_3=k_4+O(1)>k_1} 2^{k_1-k_3} \| u_{k_3} \|_{S_{k_3}} \| u_{k_4} \|_{S_{k_4}}. \end{aligned}$$

This expression can be summed over k_1 to give the desired bound. Similarly, we get

$$\begin{aligned} & \left\| L \left((-\Delta)^{\frac{1}{2}} u_{k_2}, (-\Delta)^{\frac{1}{2}} 2^{-k_1} L(\nabla_x u_{<k_1-10}, u_{k_1}) \right) \right\|_{L_t^1 L_x^2} \\ & \lesssim \| (-\Delta)^{\frac{1}{2}} u_{k_2} \|_{L_t^\infty L_x^2} \| \nabla_x u_{<k_1-10} \|_{L_t^2 L_x^\infty} \| u_{k_1} \|_{L_t^2 L_x^\infty} \\ & \lesssim \| u_{k_2} \|_{S_{k_2}} \| u_{k_1} \|_{S_{k_1}} \| u \|_S, \end{aligned}$$

and summation over $k_1 < 10$ yields the desired bound. This concludes case (ii), and thereby of (4-3).

The estimates (4-4), (4-5) are proved similarly, after passing to the differences. One only needs to make sure to reformulate the terms as in the preceding using (4-6), (4-9), before passing to the differences. \square

5. The iteration scheme

Here we solve (2-1). Specifically, we prove the following.

Theorem 5.1. *Let $n \geq 5$. Let $u[0] = (u, u_t) : \mathbb{R}^n \rightarrow S^2 \times TS^2$ be a smooth data pair with $u \cdot u_t = 0$ pointwise, and such that u is constant outside of a compact subset of \mathbb{R}^n . Also, assume the smallness condition*

$$\|u[0]\|_{\dot{B}_2^{n/2,1} \times \dot{B}_2^{n/2-1,1}} < \epsilon,$$

where $\epsilon \ll 1$ is sufficiently small. Then problem (2-1) admits a global smooth solution with these data.

Proof. We do this by means of a suitable iteration scheme: first, let $u^{(0)} = p$, where $p \in S^2$ is the limit of the initial data $u|_{t=0}$ at spatial infinity. Then let $u^{(1)}$ be the wave map into S^2 with the given data (which is possible since $u_t(0, \cdot) \cdot u(0, \cdot) = 0$ from our assumption), thus solving

$$(\partial_t^2 - \Delta)u^{(1)} = u^{(1)}(\nabla u^{(1)} \cdot \nabla u^{(1)} - \partial_t u^{(1)} \cdot \partial_t u^{(1)}).$$

It is given by $u^{(1)} = p + \sum_{k \in \mathbb{Z}} u_k^{(1)}$, and its existence follows via simple iteration from (4-1) and the corresponding difference estimate. Then we define the higher iterates $u^{(j)}$, $j \geq 2$, via the following iterative scheme:

$$\begin{aligned} & (\partial_t^2 - \Delta)u^{(j)} \\ & = u^{(j)}(\nabla u^{(j)} \cdot \nabla u^{(j)} - \partial_t u^{(j)} \cdot \partial_t u^{(j)}) \\ & \quad + \Pi_{u_\perp^{(j)}} \left((-\Delta)^{\frac{1}{2}} u^{(j-1)} \right) (u^{(j-1)} \cdot (-\Delta)^{\frac{1}{2}} u^{(j-1)}) \\ & \quad + \Pi_{u_\perp^{(j)}} \left[u^{(j-1)} \times (-\Delta)^{\frac{1}{2}} (u^{(j-1)} \times (-\Delta)^{\frac{1}{2}} u^{(j-1)}) - u^{(j-1)} \times (u^{(j-1)} \times (-\Delta)u^{(j-1)}) \right]. \end{aligned} \tag{5-1}$$

This equation defines $u^{(j)}$ implicitly, and so to actually compute it, we have to run a subiteration

$$\begin{aligned}
& (\partial_t^2 - \Delta)u^{(j,i)} \\
&= u^{(j,i-1)}(\nabla u^{(j,i-1)} \cdot \nabla u^{(j,i-1)} - \partial_t u^{(j,i-1)} \cdot \partial_t u^{(j,i-1)}) \\
&\quad + \Pi_{u_\perp} u^{(j,i-1)} ((-\Delta)^{\frac{1}{2}} u^{(j-1)})(u^{(j-1)} \cdot (-\Delta)^{\frac{1}{2}} u^{(j-1)}) \\
&\quad + \Pi_{u_\perp} u^{(j,i-1)} [u^{(j-1)} \times (-\Delta)^{\frac{1}{2}} (u^{(j-1)} \times (-\Delta)^{\frac{1}{2}} u^{(j-1)}) - u^{(j-1)} \times (u^{(j-1)} \times (-\Delta) u^{(j-1)})] \quad (5-2)
\end{aligned}$$

for $i \geq 1$, while $u^{(j,0)}$ is the free wave evolution of the data $u[0]$. Then we again have $u^{(j,i)} = p + \sum_k u_k^{(j,i)}$, and in particular each $u^{(j,i)}$ is close to S^2 with respect to the L^∞ norm, while convergence with respect to $\|\cdot\|_S$ follows from Proposition 4.1. We also get higher regularity of each $u^{(j,i)}$ and $u^{(j)}$ by differentiating the equation.

Our choice of iterative scheme (5-1) implies

$$\square(u^{(j)} \cdot u^{(j)} - 1) = (u^{(j)} \cdot u^{(j)} - 1)(\nabla u^{(j)} \cdot \nabla u^{(j)} - \partial_t u^{(j)} \cdot \partial_t u^{(j)}),$$

as well as $(u^{(j)} \cdot u^{(j)} - 1)[0] = (0, 0)$, which inductively gives that $u^{(j)}$ maps into S^2 for all j . Finally, convergence of the $u^{(j)}$ with respect to $\|\cdot\|_S$ follows again via Proposition 4.1. Differentiating (5-1) then also gives higher regularity of the limit function u . The latter is then easily seen to solve (2-1). For later purposes, we also note that Proposition 4.1 in conjunction with the assumptions that $(u - p)|_{t=0} \in C_0^\infty$ and $u_t|_{t=0} = u \times (-\Delta)^{\frac{1}{2}}|_{t=0}$ imply that we have improved control over low frequencies: $u(t, \cdot) \in \dot{H}^{\frac{n}{2}-\frac{1}{2}}$, $u_t(t, \cdot) \in \dot{H}^{\frac{n}{2}-\frac{3}{2}}$ for all t . \square

6. Proof of Theorem 1.1

It remains to show that the solution $u(t, x)$ obtained in Theorem 5.1 actually solves (1-1). For this introduce the quantity

$$X := u_t - u \times (-\Delta)^{\frac{1}{2}} u,$$

as well as the energy type functional

$$\tilde{E}(t) := \frac{1}{2} \int_{\mathbb{R}^n} |(-\Delta)^{\frac{n}{4}-\frac{3}{4}} X(t, \cdot)|^2 dx.$$

Note that we have $\nabla_{t,x} u \in \dot{H}^{\frac{n}{2}-\frac{3}{2}}$ as observed previously, and hence $\tilde{E}(t)$ is well defined and also continuously differentiable (on account of the higher-regularity properties of u). Retracing the steps that led to the final wave equation (2-1), we deduce

$$\partial_t X = -X \times (-\Delta)^{\frac{1}{2}} u - u \times (-\Delta)^{\frac{1}{2}} X - u(X \cdot (u \times (-\Delta)^{\frac{1}{2}} u + u_t)),$$

and so we deduce

$$\begin{aligned}
\frac{d}{dt} \tilde{E}(t) &= - \int_{\mathbb{R}^n} (-\Delta)^{\frac{n}{4}-\frac{3}{4}} (X \times (-\Delta)^{\frac{1}{2}} u + u \times (-\Delta)^{\frac{1}{2}} X) \cdot (-\Delta)^{\frac{n}{4}-\frac{3}{4}} X dx \\
&\quad - \int_{\mathbb{R}^n} (-\Delta)^{\frac{n}{4}-\frac{3}{4}} (u(X \cdot (u \times (-\Delta)^{\frac{1}{2}} u + u_t))) \cdot (-\Delta)^{\frac{n}{4}-\frac{3}{4}} X dx.
\end{aligned}$$

Then we note that³

$$\begin{aligned} & \|(-\Delta)^{\frac{n}{4}-\frac{3}{4}}(X \times (-\Delta)^{\frac{1}{2}}u + u \times (-\Delta)^{\frac{1}{2}}X) - u \times (-\Delta)^{\frac{n}{4}-\frac{1}{4}}X\|_{L_x^2} \\ & \lesssim \|(-\Delta)^{\frac{n}{4}-\frac{3}{4}}X\|_{L_x^2} \|(-\Delta)^{\frac{1}{2}}u\|_{L_x^\infty} + \|X\|_{L_x^{2n/3}} \|(-\Delta)^{\frac{n}{4}-\frac{1}{4}}u\|_{L_x^{2n/(n-3)}} \\ & \quad + \|(-\Delta)^{\frac{n}{4}-\frac{3}{4}}u\|_{L_x^{2n/(n-5)}} \|(-\Delta)^{\frac{1}{2}}X\|_{L_x^{2n/5}} \lesssim u \|(-\Delta)^{\frac{n}{4}-\frac{3}{4}}X\|_{L_x^2}^2 \end{aligned}$$

on account of Sobolev’s embedding and higher regularity of u , and further, we observe that

$$\begin{aligned} & \int_{\mathbb{R}^n} (u \times (-\Delta)^{\frac{n}{4}-\frac{1}{4}}X) \cdot (-\Delta)^{\frac{n}{4}-\frac{3}{4}}X \, dx \\ & = \int_{\mathbb{R}^n} (-\Delta)^{\frac{1}{4}}(u \times (-\Delta)^{\frac{n}{4}-\frac{2}{4}}X) \cdot (-\Delta)^{\frac{n}{4}-\frac{3}{4}}X \, dx + O(\|(-\Delta)^{\frac{n}{4}-\frac{3}{4}}X\|_{L_x^2}^2 \|\nabla_x u\|_{L_x^\infty}) \\ & = O(\|(-\Delta)^{\frac{n}{4}-\frac{3}{4}}X\|_{L_x^2}^2 \|\nabla_x u\|_{L_x^\infty}). \end{aligned}$$

Similarly, we infer

$$\left| \int_{\mathbb{R}^n} (-\Delta)^{\frac{n}{4}-\frac{3}{4}}(u(X \cdot (u \times (-\Delta)^{\frac{1}{2}}u + u_t))) \cdot (-\Delta)^{\frac{n}{4}-\frac{3}{4}}X \, dx \right| \lesssim u \|(-\Delta)^{\frac{n}{4}-\frac{3}{4}}X\|_{L_x^2}^2.$$

But then the preceding implies that

$$\frac{d}{dt} \tilde{E}(t) \leq C(u) \tilde{E}(t)$$

and furthermore $\tilde{E}(0) = 0$, which implies $\tilde{E}(t) = 0$ throughout. It follows that $X = 0$ identically, which completes the proof of Theorem 1.1.

Appendix

Here we prove some bounds related to the projection operator $\Pi_{\tilde{u}_\perp}$ used in the proof of Proposition 4.1.

Lemma A.1. *Assume that $\tilde{u} : \mathbb{R}^{5+1} \rightarrow S^2$ maps into a small neighbourhood of S^2 with $\|\tilde{u}\|_S \lesssim 1$. Then for any $a \in \mathbb{Z}$ we have the bounds*

$$\|P_{[-a,a]}(\Pi_{\tilde{u}_\perp}((-\Delta)^{\frac{1}{2}}u))\|_{L_t^\infty L_x^2} \lesssim a \sum_{k_3 \in \mathbb{Z}} 2^{-|k_3|} \|P_{k_3}u\|_{S_{k_3}} (1 + \|\tilde{u}\|_S), \tag{A-1}$$

$$\|(\Pi_{\tilde{u}_\perp}((-\Delta)^{\frac{1}{2}}u))\|_{L_t^\infty L_x^2 + L_{t,x}^\infty} \lesssim \sum_{k_3 \in \mathbb{Z}} 2^{-|k_3|} \|P_{k_3}u\|_{S_{k_3}} (1 + \|\tilde{u}\|_S), \tag{A-2}$$

$$\|P_{<a} \Pi_{\tilde{u}_\perp}((-\Delta)^{\frac{1}{2}}u)\|_{L_t^2 L_x^\infty} \lesssim a (\|\tilde{u}\|_S + 1) \|u\|_S. \tag{A-3}$$

Proof of (A-1). Note that we can write

$$\Pi_{\tilde{u}_\perp}((-\Delta)^{\frac{1}{2}}u) = (-\Delta)^{\frac{1}{2}}u - F(\tilde{u}) \cdot (-\Delta)^{\frac{1}{2}}u$$

³Here $2n/(n-5)$ gets replaced by ∞ if $n = 5$.

for a suitable C^∞ function $F : \mathbb{R}^3 \rightarrow \mathbb{R}^3$, which in addition to all its derivatives is bounded. Then since

$$\|P_{[-a,a]}(-\Delta)^{\frac{1}{2}}u\|_{L_t^\infty L_x^2} \lesssim a \sum_{k_3 \in \mathbb{Z}} 2^{-|k_3|} \|P_{k_3}u\|_{S_{k_3}},$$

it suffices to consider $\|P_{[-a,a]}[F(\tilde{u}) \cdot (-\Delta)^{\frac{1}{2}}u]\|_{L_t^\infty L_x^2}$. To deal with this expression, observe first that

$$\begin{aligned} \|P_l F(\tilde{u})\|_{L_t^\infty L_x^2} &\lesssim 2^{-l} \|P_l [P_{<l-20}(\nabla \tilde{u}) F'(\tilde{u})]\|_{L_t^\infty L_x^2} + 2^{-\frac{5}{2}l} \|\tilde{u}\|_{\dot{H}^{n/2}} \\ &\lesssim 2^{-2l} \|P_l [P_{<l-20}(\nabla \tilde{u}) P_{<l-20}(\nabla \tilde{u}) F''(\tilde{u})]\|_{L_t^\infty L_x^2} + 2^{-\frac{5}{2}l} \|\tilde{u}\|_{\dot{H}^{n/2}} \\ &\lesssim 2^{-3l} \|P_l [P_{<l-20}(\nabla \tilde{u}) P_{<l-20}(\nabla \tilde{u}) P_{<l-20}(\nabla \tilde{u}) F'''(\tilde{u})]\|_{L_t^\infty L_x^2} + 2^{-\frac{5}{2}l} \|\tilde{u}\|_{\dot{H}^{n/2}} \end{aligned}$$

and we can estimate the last term by

$$\begin{aligned} 2^{-3l} \|P_l [P_{<l-20}(\nabla \tilde{u}) P_{<l-20}(\nabla \tilde{u}) P_{<l-20}(\nabla \tilde{u}) F'''(\tilde{u})]\|_{L_t^\infty L_x^2} \\ \lesssim 2^{-3l} \|P_{<l-20}(\nabla \tilde{u})\|_{L_t^\infty L_x^6}^3 \lesssim 2^{-\frac{5}{2}l} \|\tilde{u}\|_S^3 \lesssim 2^{-\frac{5}{2}l} \|\tilde{u}\|_S, \end{aligned}$$

whence in summary $\|P_l F(\tilde{u})\|_{L_t^\infty L_x^2} \lesssim 2^{-\frac{5}{2}l} \|\tilde{u}\|_S$. To conclude, we estimate

$$\begin{aligned} \|P_{[-a,a]}[F(\tilde{u}) \cdot (-\Delta)^{\frac{1}{2}}u]\|_{L_t^\infty L_x^2} &\leq \|P_{[-a,a]}[P_{<-a-10}[F(\tilde{u})] \cdot (-\Delta)^{\frac{1}{2}}u]\|_{L_t^\infty L_x^2} \\ &\quad + \|P_{[-a,a]}[P_{[-a-10,a+10]}[F(\tilde{u})] \cdot (-\Delta)^{\frac{1}{2}}u]\|_{L_t^\infty L_x^2} \\ &\quad + \|P_{[-a,a]}[P_{>a+10}[F(\tilde{u})] \cdot (-\Delta)^{\frac{1}{2}}u]\|_{L_t^\infty L_x^2}, \end{aligned}$$

and we have

$$\begin{aligned} \|P_{[-a,a]}[P_{<-a-10}[F(\tilde{u})] \cdot (-\Delta)^{\frac{1}{2}}u]\|_{L_t^\infty L_x^2} &\lesssim \|P_{[-a-20,a+20]}(-\Delta)^{\frac{1}{2}}u\|_{L_t^\infty L_x^2} \\ &\lesssim a \sum_{k_3 \in \mathbb{Z}} 2^{-|k_3|} \|P_{k_3}u\|_{S_{k_3}}, \end{aligned}$$

$$\begin{aligned} \|P_{[-a,a]}[P_{[-a-10,a+10]}[F(\tilde{u})] \cdot (-\Delta)^{\frac{1}{2}}u]\|_{L_t^\infty L_x^2} &\lesssim \|P_{[-a-10,a+10]}[F(\tilde{u})]\|_{L_t^\infty L_x^2} \|P_{<a+10}(-\Delta)^{\frac{1}{2}}u\|_{L_t^\infty L_x^2} \\ &\lesssim a \|\tilde{u}\|_S \sum_{k_3 \in \mathbb{Z}} 2^{-|k_3|} \|P_{k_3}u\|_{S_{k_3}}, \end{aligned}$$

where we have used the preceding bound for $P_l F(\tilde{u})$ to control $\|P_{[-a-10,a+10]}[F(\tilde{u})]\|_{L_t^\infty L_x^2}$. Finally, we get

$$\begin{aligned} \|P_{[-a,a]}[P_{>a+10}[F(\tilde{u})] \cdot (-\Delta)^{\frac{1}{2}}u]\|_{L_t^\infty L_x^2} &\leq \sum_{k_1=k_2+O(1)>a+10} \|P_{[-a,a]}[P_{k_1}[F(\tilde{u})] \cdot P_{k_2}(-\Delta)^{\frac{1}{2}}u]\|_{L_t^\infty L_x^2} \\ &\lesssim a \sum_{k_1=k_2+O(1)>a+10} \|P_{k_1}[F(\tilde{u})]\|_{L_t^\infty L_x^2} \|P_{k_2}(-\Delta)^{\frac{1}{2}}u\|_{L_t^\infty L_x^2} \\ &\lesssim a \|\tilde{u}\|_S \sum_{k_3 \in \mathbb{Z}} 2^{-|k_3|} \|P_{k_3}u\|_{S_{k_3}}, \end{aligned}$$

where we have used Bernstein's and Holder's inequalities as well as the preceding bound for $P_l F(\tilde{u})$. \square

Proof of (A-2). This is similar to the preceding bound; one places

$$P_{<0}[(\Pi_{\tilde{u}_\perp}((-\Delta)^{\frac{1}{2}}u))]$$

into $L_{t,x}^\infty$ and

$$P_{\geq 0}[(\Pi_{\tilde{u}_\perp}((-\Delta)^{\frac{1}{2}}u))]$$

into $L_t^\infty L_x^2$. □

Proof of (A-3). We use the preceding bounds, and reduce to bounding $\|P_{<a}(F(\tilde{u})(-\Delta)^{\frac{1}{2}}u)\|_{L_t^2 L_x^\infty}$. Then

$$\begin{aligned} \|P_{<a}(F(\tilde{u})(-\Delta)^{\frac{1}{2}}u)\|_{L_t^2 L_x^\infty} &\lesssim \|P_{<a}(F(\tilde{u})P_{<a+10}(-\Delta)^{\frac{1}{2}}u)\|_{L_t^2 L_x^\infty} + \|P_{<a}(F(\tilde{u})P_{\geq a+10}(-\Delta)^{\frac{1}{2}}u)\|_{L_t^2 L_x^\infty} \end{aligned}$$

and we can bound

$$\begin{aligned} \|P_{<a}(F(\tilde{u})P_{<a+10}(-\Delta)^{\frac{1}{2}}u)\|_{L_t^2 L_x^\infty} &\lesssim \sum_{k < a+10} \|P_k(-\Delta)^{\frac{1}{2}}u\|_{L_t^2 L_x^\infty} \\ &\lesssim \sum_{k < a+10} 2^{\frac{k}{2}} \|P_k u\|_{S_k} \lesssim_a \|u\|_S, \end{aligned}$$

as well as

$$\begin{aligned} \|P_{<a}(F(\tilde{u})P_{\geq a+10}(-\Delta)^{\frac{1}{2}}u)\|_{L_t^2 L_x^\infty} &\lesssim \sum_{k_1=k_2+O(1)\geq a+10} \|P_{<a}(P_{k_2}F(\tilde{u})P_{k_1}(-\Delta)^{\frac{1}{2}}u)\|_{L_t^2 L_x^\infty} \\ &\lesssim 2^{\frac{5}{2}a} \sum_{k_1=k_2+O(1)\geq a+10} \|P_{k_2}F(\tilde{u})\|_{L_t^\infty L_x^2} \|P_{k_1}(-\Delta)^{\frac{1}{2}}u\|_{L_t^2 L_x^\infty} \\ &\lesssim 2^{\frac{5}{2}a} \sum_{k_1=k_2+O(1)\geq a+10} 2^{-\frac{5}{2}k_2} \|\tilde{u}\|_S \cdot 2^{\frac{k_1}{2}} \|P_{k_1}u\|_{S_{k_1}} \\ &\lesssim_a \|\tilde{u}\|_S \|u\|_S. \end{aligned} \quad \square$$

References

[Blom and Langmann 1998] J. Blom and E. Langmann, “Novel integrable spin-particle models from gauge theories on a cylinder”, *Phys. Lett. B* **429**:3-4 (1998), 336–342.

[Da Lio 2013] F. Da Lio, “Fractional harmonic maps into manifolds in odd dimension $n > 1$ ”, *Calc. Var. Partial Differential Equations* **48**:3-4 (2013), 421–445. MR Zbl

[Da Lio and Rivière 2011a] F. Da Lio and T. Rivière, “Sub-criticality of non-local Schrödinger systems with antisymmetric potentials and applications to half-harmonic maps”, *Adv. Math.* **227**:3 (2011), 1300–1348. MR Zbl

[Da Lio and Rivière 2011b] F. Da Lio and T. Rivière, “Three-term commutator estimates and the regularity of $\frac{1}{2}$ -harmonic maps into spheres”, *Anal. PDE* **4**:1 (2011), 149–190. MR Zbl

[Haldane 1988] F. D. M. Haldane, “Exact Jastrow–Gutzwiller resonating-valence-bond ground state of the spin- $\frac{1}{2}$ antiferromagnetic Heisenberg chain with $1/r^2$ exchange”, *Phys. Rev Lett.* **60**:7 (1988), 635–638.

[Hikami and Wadati 1993] K. Hikami and M. Wadati, “Integrability of Calogero–Moser spin system”, *J. Phys. Soc. Japan* **62**:2 (1993), 469–472. MR Zbl

- [Krieger 2008] J. Krieger, “Global regularity and singularity development for wave maps”, pp. 167–201 in *Surveys in differential geometry, XII: Geometric flows*, edited by H.-D. Cao and S.-T. Yau, Surv. Differ. Geom. **12**, International Press, Somerville, MA, 2008. MR Zbl
- [Millot and Sire 2015] V. Millot and Y. Sire, “On a fractional Ginzburg–Landau equation and $\frac{1}{2}$ -harmonic maps into spheres”, *Arch. Ration. Mech. Anal.* **215**:1 (2015), 125–210. MR Zbl
- [Schikorra and Lenzmann 2017] A. Schikorra and E. Lenzmann, “On energy-critical half-wave maps into \mathbb{S}^2 ”, preprint, 2017. arXiv
- [Shastry 1988] B. S. Shastry, “Exact solution of an $S = \frac{1}{2}$ Heisenberg antiferromagnetic chain with long-ranged interactions”, *Phys. Rev. Lett.* **60**:7 (1988), 639–642.
- [Shatah and Struwe 1998] J. Shatah and M. Struwe, *Geometric wave equations*, Courant Lecture Notes in Mathematics **2**, New York University, 1998. MR Zbl
- [Sterbenz 2004] J. Sterbenz, “Global regularity for general non-linear wave equations, I: $(6 + 1)$ and higher dimensions”, *Comm. Partial Differential Equations* **29**:9-10 (2004), 1505–1531. MR Zbl
- [Tao 2001a] T. Tao, “Global regularity of wave maps, I: Small critical Sobolev norm in high dimension”, *Internat. Math. Res. Notices* **6** (2001), 299–328. MR Zbl
- [Tao 2001b] T. Tao, “Global regularity of wave maps, II: Small energy in two dimensions”, *Comm. Math. Phys.* **224**:2 (2001), 443–544. MR Zbl
- [Tataru 1998] D. Tataru, “Local and global results for wave maps, I”, *Comm. Partial Differential Equations* **23**:9-10 (1998), 1781–1793. MR Zbl

Received 22 Oct 2016. Revised 17 May 2017. Accepted 23 Sep 2017.

JOACHIM KRIEGER: joachim.krieger@epfl.ch
Bâtiment des Mathématiques, EPFL, Lausanne, Switzerland

YANNICK SIRE: sire@math.jhu.edu
Department of Mathematics, Johns Hopkins University, Baltimore, MD, United States

Analysis & PDE

msp.org/apde

EDITORS

EDITOR-IN-CHIEF

Patrick Gérard
patrick.gerard@math.u-psud.fr
Université Paris Sud XI
Orsay, France

BOARD OF EDITORS

| | | | |
|----------------------|---|-----------------------|--|
| Nicolas Burq | Université Paris-Sud 11, France nicolas.burq@math.u-psud.fr | Werner Müller | Universität Bonn, Germany mueller@math.uni-bonn.de |
| Massimiliano Berti | Scuola Intern. Sup. di Studi Avanzati, Italy berti@sissa.it | Gilles Pisier | Texas A&M University, and Paris 6 pisier@math.tamu.edu |
| Sun-Yung Alice Chang | Princeton University, USA chang@math.princeton.edu | Tristan Rivière | ETH, Switzerland riviere@math.ethz.ch |
| Michael Christ | University of California, Berkeley, USA mchrist@math.berkeley.edu | Igor Rodnianski | Princeton University, USA irod@math.princeton.edu |
| Charles Fefferman | Princeton University, USA cf@math.princeton.edu | Wilhelm Schlag | University of Chicago, USA schlag@math.uchicago.edu |
| Ursula Hamenstaedt | Universität Bonn, Germany ursula@math.uni-bonn.de | Sylvia Serfaty | New York University, USA serfaty@cims.nyu.edu |
| Vaughan Jones | U.C. Berkeley & Vanderbilt University vaughan.f.jones@vanderbilt.edu | Yum-Tong Siu | Harvard University, USA siu@math.harvard.edu |
| Vadim Kaloshin | University of Maryland, USA vadim.kaloshin@gmail.com | Terence Tao | University of California, Los Angeles, USA tao@math.ucla.edu |
| Herbert Koch | Universität Bonn, Germany koch@math.uni-bonn.de | Michael E. Taylor | Univ. of North Carolina, Chapel Hill, USA met@math.unc.edu |
| Izabella Laba | University of British Columbia, Canada ilaba@math.ubc.ca | Gunther Uhlmann | University of Washington, USA gunther@math.washington.edu |
| Gilles Lebeau | Université de Nice Sophia Antipolis, France lebeau@unice.fr | András Vasy | Stanford University, USA andras@math.stanford.edu |
| Richard B. Melrose | Massachusetts Inst. of Tech., USA rbm@math.mit.edu | Dan Virgil Voiculescu | University of California, Berkeley, USA dvv@math.berkeley.edu |
| Frank Merle | Université de Cergy-Pontoise, France Frank.Merle@u-cergy.fr | Steven Zelditch | Northwestern University, USA zelditch@math.northwestern.edu |
| William Minicozzi II | Johns Hopkins University, USA minicozz@math.jhu.edu | Maciej Zworski | University of California, Berkeley, USA zworski@math.berkeley.edu |
| Clément Mouhot | Cambridge University, UK c.mouhot@dpms.cam.ac.uk | | |

PRODUCTION

production@msp.org
Silvio Levy, Scientific Editor


See inside back cover or msp.org/apde for submission instructions.

The subscription price for 2018 is US \$275/year for the electronic version, and \$480/year (+\$55, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues from the last three years and changes of subscriber address should be sent to MSP.

Analysis & PDE (ISSN 1948-206X electronic, 2157-5045 printed) at Mathematical Sciences Publishers, 798 Evans Hall #3840, c/o University of California, Berkeley, CA 94720-3840, is published continuously online. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices.

APDE peer review and production are managed by EditFlow® from MSP.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2018 Mathematical Sciences Publishers

ANALYSIS & PDE

Volume 11 No. 3 2018

| | |
|---|-----|
| The endpoint perturbed Brascamp–Lieb inequalities with examples RUIXIANG ZHANG | 555 |
| Square function estimates for discrete Radon transforms MARIUSZ MIREK | 583 |
| On the Kato problem and extensions for degenerate elliptic operators DAVID CRUZ-URIBE, JOSÉ MARÍA MARTELL and CRISTIAN RIOS | 609 |
| Small data global regularity for half-wave maps JOACHIM KRIEGER and YANNICK SIRE | 661 |
| The semigroup generated by the Dirichlet Laplacian of fractional order TSUKASA IWABUCHI | 683 |
| Klein’s paradox and the relativistic δ -shell interaction in \mathbb{R}^3 ALBERT MAS and FABIO PIZZICHILLO | 705 |
| Dimension-free L^p estimates for vectors of Riesz transforms associated with orthogonal ex- pansions BŁAŻEJ WRÓBEL | 745 |
| Reducibility of the quantum harmonic oscillator in d -dimensions with polynomial time-dependent perturbation DARIO BAMBUSI, BENOÎT GRÉBERT, ALBERTO MASPERO and DIDIER ROBERT | 775 |
| Eigenfunction scarring and improvements in L^∞ bounds JEFFREY GALKOWSKI and JOHN A. TOTH | 801 |



2157-5045(2018)11:3;1-R