

ANALYSIS & PDE

Volume 11 No. 5 2018

PIERRE-YVES BIENAIMÉ AND ABDESSLAM BOULKHEMAIR

**WELL-POSEDNESS AND SMOOTHING EFFECT FOR
GENERALIZED NONLINEAR SCHRÖDINGER EQUATIONS**

WELL-POSEDNESS AND SMOOTHING EFFECT FOR GENERALIZED NONLINEAR SCHRÖDINGER EQUATIONS

PIERRE-YVES BIENAIMÉ AND ABDESSLAM BOULKHEMAIR

We improve the result obtained by one of the authors, Bienaimé (2014), and establish the well-posedness of the Cauchy problem for some nonlinear equations of Schrödinger type in the usual Sobolev space $H^s(\mathbb{R}^n)$ for $s > \frac{n}{2} + 2$ instead of $s > \frac{n}{2} + 3$. We also improve the smoothing effect of the solution and obtain the optimal exponent.

1. Introduction

Consider the nonlinear Cauchy problem

$$\begin{cases} \partial_t u = i\mathcal{L}u + F(u, \nabla_x u, \bar{u}, \nabla_x \bar{u}), & t \in \mathbb{R}, x \in \mathbb{R}^n, \\ u(x, 0) = u_0(x) \in H^s(\mathbb{R}^n), \end{cases} \quad (1)$$

where the function F is sufficiently regular in $\mathbb{C} \times \mathbb{C}^n \times \mathbb{C} \times \mathbb{C}^n$, the operator \mathcal{L} has the form

$$\mathcal{L} = \sum_{j \leq j_0} \partial_{x_j}^2 - \sum_{j > j_0} \partial_{x_j}^2,$$

with a fixed $j_0 \in \{1, 2, \dots, n\}$, and $H^s(\mathbb{R}^n)$, $s \in \mathbb{R}$, is the usual Sobolev space on \mathbb{R}^n . Thus, \mathcal{L} generalizes the Laplace operator but is not elliptic unless $j_0 = n$. Hence, such equations are generalizations of the nonlinear Schrödinger (NLS) equations.

In this paper, we continue the work undertaken in [Bienaimé 2014] and study the local existence and the smoothing effect of the solutions of the Cauchy problem (1) with essentially the following goal: to obtain the optimal index s of regularity for which (1) is well-posed. In fact, since the partial differential equation is of second order and is semilinear, the optimal condition on s should be $s > \frac{n}{2} + 1$. Unfortunately, up to now and due to issues that occur when estimating the remainder obtained after the linearization of the nonlinear equation, we have not been able to prove the desired result under such a condition. In any case, we shall return to this question in a future work. In this paper, we establish the following:

Theorem 1.1. *Assume that F vanishes to the third order at 0; that is, F and its partial derivatives up to the second order vanish at 0. Then, for every $s > \frac{n}{2} + 2$ and every initial data $u_0 \in H^s(\mathbb{R}^n)$, there exists a real number $T > 0$ such that the Cauchy problem (1) has a unique solution u which is defined on the interval $[0, T]$ and satisfies*

$$u \in C([0, T]; H^s(\mathbb{R}^n))$$

MSC2010: 47G20, 47G30.

Keywords: Cauchy problem, well-posedness, smoothing effect, nonlinear equation, Schrödinger, paradifferential, pseudodifferential, operator, parilinearization.

and

$$\| \| J^{s+\frac{1}{2}} u \| \|_T \stackrel{\text{def}}{=} \sup_{\mu \in \mathbb{Z}^n} \left(\int_0^T \int_{\mathbb{R}^n} |\langle x - \mu \rangle^{-\sigma_0} J^{s+\frac{1}{2}} u(x, t)|^2 dx dt \right)^{\frac{1}{2}} < \infty,$$

where $J = (1 - \Delta)^{\frac{1}{2}}$, $\Delta = \sum_{k=1}^{k=n} \partial_{x_k}^2$ and $\sigma_0 > \frac{1}{2}$ is fixed. Moreover, given a bounded subset B of $H^s(\mathbb{R}^n)$, there exists a real number $T > 0$ such that, for every $u_0 \in B$, the associated solution u of (1) exists on the interval $[0, T]$ and the map which associates u to u_0 is Lipschitz continuous from B into the space

$$\{w \in C([0, T]; H^s(\mathbb{R}^n)) : \| \| J^{s+\frac{1}{2}} w \| \|_T < \infty\}.$$

In [Bienaimé 2014], this theorem is proved under the assumption $s > \frac{n}{2} + 3$. We also improve the result with respect to the smoothing effect of the solution since $\sigma_0 = 2$ there. Note that the assumption $\sigma_0 > \frac{1}{2}$ in the above theorem seems to be sharp; we refer for example to the survey article [Robbiano 2013] on the subject of Kato’s smoothing effect. Recall that at the origin of [Bienaimé 2014] was the significant work of C. E. Kenig, G. Ponce and L. Vega [Kenig et al. 1998], who first studied (1) with such a nonelliptic \mathcal{L} and established the local existence and the smoothing effect of the solutions assuming that F is a polynomial and $s \geq s_0$, the index s_0 being sufficiently large. Note that these authors did not give an idea about the value of s_0 , but by going back to the details of their proof, one can see that s_0 is of the order of $\frac{n}{2} + 10n + 1$. These authors also studied the case where F (is a polynomial and) vanishes to the second order at 0. However, it seems that in that case we need to work in weighted Sobolev spaces.

The Cauchy problem (1) was extensively studied in the 90s mainly when $\mathcal{L} = \Delta$, that is, in the case of the Schrödinger equation. See the introduction of [Kenig et al. 1998]. The case $\mathcal{L} \neq \Delta$ is less well-known. Nevertheless, it is motivated by several equations coming from the applications such as Ishimori-type equations or Davey–Stewartson-type systems. For more details, we refer the reader to the instructive introduction of [Kenig et al. 1998]. Let us now quote some papers which are more or less related to this subject. In [Kenig et al. 2004], the authors extended their results of 1998 to the quasilinear case assuming essentially that the corresponding dispersive operator \mathcal{L} is elliptic and nontrapping. The nonelliptic case is treated in [Kenig et al. 2006; 2005]. In [Bejenaru and Tataru 2008], the authors solved the Cauchy problem (1) for $s > \frac{n}{2} + 1$ in modified Sobolev spaces and assuming $F(u, \nabla_x u, \bar{u}, \nabla_x \bar{u})$ bilinear. More recently, in [Marzuola et al. 2012; 2014], the authors considered the quasilinear Schrödinger equation

$$i \partial_t u + \sum_{j,k} g^{j,k}(u, \nabla_x u) \partial_j \partial_k u = F(u, \nabla_x u)$$

and obtained the local well-posedness of the associated Cauchy problem for $s > \frac{n}{2} + 3$ in the quadratic case (with modified Sobolev spaces) and for $s > \frac{n}{2} + \frac{5}{2}$ in the nonquadratic case. However, they assume the smallness of the data and they do not seem to obtain the smoothing effect of the solutions.

The proof of Theorem 1.1 follows the same ideas as that of [Kenig et al. 1998; Bienaimé 2014]. Of course, the general plan is unoriginal: linearization of the nonlinear equation, then, establishing energy estimates for solutions of the linear equation, and finally, solving the nonlinear equation by means of an appropriate fixed-point theorem. Like [Bienaimé 2014], we start by applying a parilinearization, that is, a

linearization in the sense of [Bony 1981] instead of the classical linearization. This leads us to the use of the paradifferential calculus whose main interest lies in the fact that it eliminates the usual losses of regularity due to commutators. One obtains a parilinear equation and most of the proof of the theorem is concerned with the study of such an equation, that is, the well-posedness in the Sobolev spaces of the associated Cauchy problem by means of energy and smoothing effect estimates. As did Kenig, Ponce and Vega, we establish the smoothing effect estimate by using Doi's argument [1994] via Gårding's inequality, and we prove the energy estimates by following an idea of [Takeuchi 1992], that is, by constructing a nonclassical invertible pseudodifferential operator C which allows estimates for Cu if u is a solution of the parilinear equation. Finally, we solve the nonlinear Cauchy problem (1) by applying these estimates to an integrodifferential equation which is equivalent to (1) and obtain the solution as the fixed point of an appropriate contraction in an appropriate complete metric space.

Now, in order to give a more precise idea about our proof, let us indicate the differences with that given in [Bienaimé 2014]. In fact, there are three main differences:

- We simplify certain arguments of that paper; for example, we no longer need to use the general Hörmander symbol spaces $S_{\rho,\delta}^m$; we only use $S_{1,0}^m$ and $S_{0,0}^m$. Also, we only use the original paradifferential operators (see Section 2) and not the variant introduced in [Bienaimé 2014].
- The linear theorem, that is, Theorem 3.1 (see Section 3), is proved for general paradifferential operators T_{b_1} and T_{b_2} of order 0 instead of paramultiplication operators. Note also that we allow the operators C_1 and C_2 to be abstract bounded operators.
- The third difference lies in the nonlinear part (see Section 4) and is crucial for our improvement of the result of [Bienaimé 2014]: we use anisotropic Sobolev spaces and an interpolation inequality (see Proposition A.5) to estimate the remainder of the parilinearized equation.

2. Notations and preliminary results

Some notation used in the paper:

- $J^s = (1 - \Delta)^{\frac{s}{2}} = \langle D \rangle^s$ is the operator whose symbol is $\langle \xi \rangle^s = (1 + \xi^2)^{\frac{s}{2}}$.
- $D_{x_k} = -i \partial_{x_k}$, $D_x = -i \partial_x$.
- $|\alpha| = \sum_{j=1}^n \alpha_j$ if $\alpha \in \mathbb{N}^n$.
- $\Delta v = (\Delta v_1, \dots, \Delta v_n)$ and $\nabla v = (\nabla v_1, \dots, \nabla v_n)$ if $v = (v_1, \dots, v_n)$.
- $\mathcal{S}(\mathbb{R}^n)$ denotes the Schwartz space of rapidly decreasing functions in \mathbb{R}^n .
- $\mathcal{D}(\mathbb{R}^n)$ denotes the space of smooth functions with compact support in \mathbb{R}^n .
- $\mathcal{D}'(\mathbb{R}^n)$ denotes the space of distributions in \mathbb{R}^n .
- $\mathcal{S}'(\mathbb{R}^n)$ denotes the space of tempered distributions in \mathbb{R}^n .
- \hat{u} or $\mathcal{F}(u)$ denotes the Fourier transform of u .
- $H^s(\mathbb{R}^n) = \{u \in \mathcal{S}'(\mathbb{R}^n) : \langle \xi \rangle^s \hat{u} \in L^2(\mathbb{R}^n)\}$ is the usual Sobolev space of regularity s .

- $\|u\|_s = \left(\int_{\mathbb{R}^n} \langle \xi \rangle^{2s} |\hat{u}(\xi)|^2 d\xi\right)^{\frac{1}{2}}$ denotes the norm of u in $H^s(\mathbb{R}^n)$.
- $\|u\|_E$ denotes the norm of u in the space E .
- Hörmander’s classes of symbols: if $m \in \mathbb{R}$ and $\gamma, \delta \in [0, 1]$,

$$S_{\gamma,\delta}^m = \{a \in C^\infty(\mathbb{R}^n \times \mathbb{R}^n) : |\partial_x^\alpha \partial_\xi^\beta a(x, \xi)| \leq A_{\alpha,\beta} \langle \xi \rangle^{m-\gamma|\beta|+\delta|\alpha|} \text{ for all } \alpha, \beta \in \mathbb{N}^n\}.$$

- If $\varrho > 0$ is an integer, $C^\varrho(\mathbb{R}^n)$ denotes the set of functions in \mathbb{R}^n which are bounded, of class C^m and their derivatives up to m are bounded. If $\varrho > 0$ is not an integer, $C^\varrho(\mathbb{R}^n)$ denotes the Hölder class, that is, the set of u in $C^{[\varrho]}(\mathbb{R}^n)$ such that

$$\exists C \in \mathbb{R}, \forall (x, y) \in \mathbb{R}^n \times \mathbb{R}^n, \quad |\partial^\alpha u(x) - \partial^\alpha u(y)| \leq C|x - y|^{\varrho-[\varrho]}.$$

- $\text{Op } S$ denotes the set of pseudodifferential operators whose symbols belong to S .

The following statement summarizes the pseudodifferential calculus associated to Hörmander’s classes of symbols $S_{\gamma,\delta}^m$:

Theorem 2.1. *If $a \in S_{\gamma,\delta}^m$, $b \in S_{\gamma,\delta}^{m'}$, $m, m' \in \mathbb{R}$, and $0 \leq \delta < \gamma \leq 1$ or $0 \leq \delta \leq \gamma < 1$, then:*

- (i) $a(x, D)b(x, D) = c(x, D)$ with $c \in S_{\gamma,\delta}^{m+m'}$. Moreover,

$$\begin{aligned} c(x, \xi) &= \int e^{-iy \cdot \eta} a(x, \xi + \eta) b(x + y, \xi) \frac{dy d\eta}{(2\pi)^n} \\ &= \sum_{|v| < N} \frac{1}{v!} \partial_\xi^v a(x, \xi) D_x^v b(x, \xi) + \sum_{|v|=N} \frac{1}{v!} \int_0^1 (1-\theta)^{N-1} r_{v,\theta}(x, \xi) d\theta, \end{aligned}$$

where

$$r_{v,\theta}(x, \xi) = \int e^{-iy \cdot \eta} \partial_\xi^v a(x, \xi + \theta\eta) D_x^v b(x + y, \xi) \frac{dy d\eta}{(2\pi)^n},$$

and the $S_{\gamma,\delta}^{m+m'-N(\gamma-\delta)}$ seminorms of $r_{v,\theta}$ are bounded by products of seminorms of a and b uniformly in $\theta \in [0, 1]$.

- (ii) $a(x, D)^* = a^*(x, D)$ with $a^* \in S_{\gamma,\delta}^m$. Moreover,

$$\begin{aligned} a^*(x, \xi) &= \int e^{-iy \cdot \eta} \bar{a}(x + y, \xi + \eta) \frac{dy d\eta}{(2\pi)^n} \\ &= \sum_{|v| < N} \frac{1}{v!} \partial_\xi^v D_x^v \bar{a}(x, \xi) + \sum_{|v|=N} \frac{1}{v!} \int_0^1 (1-\theta)^{N-1} r_{v,\theta}^*(x, \xi) d\theta, \end{aligned}$$

where

$$r_{v,\theta}^*(x, \xi) = \int e^{-iy \cdot \eta} \partial_\xi^v D_x^v \bar{a}(x + y, \xi + \theta\eta) \frac{dy d\eta}{(2\pi)^n},$$

and the $S_{\gamma,\delta}^{m-N(\gamma-\delta)}$ seminorms of $r_{v,\theta}^*$ are bounded by seminorms of a uniformly in $\theta \in [0, 1]$.

See [Taylor 1991], for instance, for the proof. We shall also often need the following version of the Calderón–Vaillancourt theorem:

Theorem 2.2. *Let $a : \mathbb{R}^n \times \mathbb{R}^n \rightarrow \mathbb{C}$ be a bounded function. Assume that, for all $\alpha, \beta \in \mathbb{N}^n$ such that $|\alpha| + |\beta| \leq n + 1$, there exists a constant $C_{\alpha,\beta} > 0$ such that $|\partial_x^\alpha \partial_\xi^\beta a(x, \xi)| \leq C_{\alpha,\beta} \in \mathbb{R}^{2n}$. Then, the pseudodifferential operator $a(x, D)$ is bounded in $L^2(\mathbb{R}^n)$ and its operator norm is estimated by*

$$\sup_{|\alpha|+|\beta|\leq n+1} \|\partial_x^\alpha \partial_\xi^\beta a\|_{L^\infty}.$$

See [Coifman and Meyer 1978] for the proof.

The following technical lemma, which is a consequence of Theorem 2.1, will be very useful in many of our proofs:

Lemma 2.3. *Let $a \in S_{0,0}^m$, $m, \sigma \in \mathbb{R}$ and $\mu \in \mathbb{R}^n$. Then:*

- (i) *We have $\langle x - \mu \rangle^\sigma a(x, D) \langle x - \mu \rangle^{-\sigma} = a_\mu(x, D)$, where $a_\mu \in S_{0,0}^m$ and the seminorms of a_μ are bounded by seminorms of a uniformly in μ .*
- (ii) *If $\sigma \geq 0$ and if, in addition, $a(x, \xi)$ is rapidly decreasing with respect to $x - \mu$, then we have $\langle x - \mu \rangle^\sigma a(x, D) \langle x - \mu \rangle^{-\sigma} = b_\mu(x, D)$, where $b_\mu \in S_{0,0}^m$, b_μ is also rapidly decreasing in $x - \mu$ and the seminorms of b_μ are estimated uniformly in μ by expressions of the form*

$$\sup_{|\alpha|+|\beta|\leq N} \|\langle x - \mu \rangle^{2\sigma} \langle \xi \rangle^{-m} \partial_x^\alpha D_\xi^\beta a\|_{L^\infty}.$$

Here, the fact that the symbol $a(x, \xi)$ is rapidly decreasing with respect to $x - \mu$ means that, for every integer N and all multi-indices α, β , the function $\langle x - \mu \rangle^N \langle \xi \rangle^{-m} \partial_x^\alpha D_\xi^\beta a$ is bounded in $\mathbb{R}^n \times \mathbb{R}^n$, and we shall often meet such symbols in this paper.

Proof. (i) When $\sigma \geq 0$, we can use Theorem 2.1(i) and integrations by parts to obtain

$$\begin{aligned} a_\mu(x, \xi) &= \langle x - \mu \rangle^\sigma (2\pi)^{-n} \int e^{-iy \cdot \eta} a(x, \xi + \eta) \langle x + y - \mu \rangle^{-\sigma} dy d\eta \\ &= \langle x - \mu \rangle^\sigma (2\pi)^{-n} \int e^{-iy \cdot \eta} J_\eta^N [\langle \eta \rangle^{-N} a(x, \xi + \eta)] \langle y \rangle^{-N} J_y^N [\langle x + y - \mu \rangle^{-\sigma}] dy d\eta, \end{aligned}$$

where N is a large and even integer. Hence, by taking derivatives and bounding, and next by applying Peetre’s inequality,

$$\begin{aligned} |a_\mu(x, \xi)| &\leq C \|\langle \xi \rangle^{-m} a\|_{C^N} \langle x - \mu \rangle^\sigma \int \langle \eta \rangle^{-N} \langle \xi + \eta \rangle^m \langle y \rangle^{-N} \langle x + y - \mu \rangle^{-\sigma} dy d\eta \\ &\leq 2^{\frac{\sigma+|m|}{2}} C \|\langle \xi \rangle^{-m} a\|_{C^N} \langle \xi \rangle^m \int \langle \eta \rangle^{|m|-N} \langle y \rangle^{\sigma-N} dy d\eta = C' \langle \xi \rangle^m \|\langle \xi \rangle^{-m} a\|_{C^N}, \end{aligned}$$

where C and C' are constants which are independent of μ , and N is taken for example such that $N \geq |m| + \sigma + n + 1$. Of course, the derivatives of a_μ are treated in the same manner.

The case $\sigma < 0$ follows from the preceding case by considering the adjoint

$$a_\mu(x, D)^* = \langle x - \mu \rangle^{-\sigma} a(x, D)^* \langle x - \mu \rangle^\sigma$$

and by applying Theorem 2.1(ii).

(ii) By using the formula in Theorem 2.1(ii) once more, it is easy to see that, if a is rapidly decreasing with respect to $x - \mu$, then the symbol a^* is also rapidly decreasing with respect to $x - \mu$ and that, for all

$N \in \mathbb{N}$, $\alpha, \beta \in \mathbb{N}^n$, there exist $M \in \mathbb{N}$ and a nonnegative constant $C_{N,\alpha,\beta}$ which does not depend on μ such that

$$\|\langle x - \mu \rangle^N \langle \xi \rangle^{-m} \partial_x^\alpha D_\xi^\beta a^* \|_{L^\infty} \leq C_{N,\alpha,\beta} \sup_{|\alpha'|+|\beta'| \leq M} \|\langle x - \mu \rangle^N \langle \xi \rangle^{-m} \partial_x^{\alpha'} D_\xi^{\beta'} a \|_{L^\infty}.$$

Now, by following the same argument as that used in the first part, one can check that the same claim holds exactly when we replace a^* by a_μ in the above assertion; in particular, we have the estimate

$$\|\langle x - \mu \rangle^N \langle \xi \rangle^{-m} \partial_x^\alpha D_\xi^\beta a_\mu \|_{L^\infty} \leq C_{N,\alpha,\beta} \sup_{|\alpha'|+|\beta'| \leq M} \|\langle x - \mu \rangle^N \langle \xi \rangle^{-m} \partial_x^{\alpha'} D_\xi^{\beta'} a \|_{L^\infty},$$

and since we can write obviously $b_\mu(x, \xi) = \langle x - \mu \rangle^{2\sigma} a_\mu(x, \xi)$, this achieves the proof of the lemma. \square

When studying the nonlinear equation, the following result is important in order to explain the assumption made on the nonlinearity F .

Lemma 2.4. *For all $s \geq 0$ and all $\sigma > \frac{n}{2}$, there exists a constant $C > 0$ such that, for all $v \in H^s(\mathbb{R}^n)$, the sequence $\mu \mapsto \|\langle x - \mu \rangle^{-\sigma} v\|_s$ is in $\ell^2(\mathbb{Z}^n)$ and*

$$\sum_{\mu} \|\langle x - \mu \rangle^{-\sigma} v\|_s^2 \leq C \|v\|_s^2.$$

In particular, if $s > \frac{n}{2}$, $u, v \in H^s(\mathbb{R}^n)$ and χ is a smooth and rapidly decreasing function, then, $\mu \mapsto \|\chi(x - \mu)uv\|_s$ is in $\ell^1(\mathbb{Z}^n)$ and

$$\sum_{\mu} \|\chi(x - \mu)uv\|_s \leq C \|u\|_s \|v\|_s.$$

Proof. The case $s = 0$ is obvious and follows from the fact that $\sum_{\mu} \langle x - \mu \rangle^{-2\sigma}$ is a bounded function. The case where s is a positive integer reduces to the case $s = 0$ by taking derivatives via Leibniz formula. The general case is obtained by interpolation. Indeed, since the map $v \mapsto \langle x - \mu \rangle^{-\sigma} v$ is linear and bounded from H^s into $\ell^2(\mathbb{Z}^n, H^s)$ for integral indices $s = s_1, s_2$, it will be also bounded from $H^{s'}$ into $\ell^2(\mathbb{Z}^n, H^{s'})$ for any real s' between s_1 and s_2 . This follows from the fact that

$$[\ell^2(\mathbb{Z}^n, H^{s_1}), \ell^2(\mathbb{Z}^n, H^{s_2})]_{\theta} = \ell^2(\mathbb{Z}^n, [H^{s_1}, H^{s_2}]_{\theta})$$

for $0 < \theta < 1$. See for example [Bergh and Löfström 1976, Theorem 5.1.2, page 107].

The second part is a consequence of the first one and the fact that $H^s(\mathbb{R}^n)$ is an algebra if $s > \frac{n}{2}$. \square

Let us now recall some results on paradifferential operators.

Definition 2.5. We define the class Σ_{ρ}^m where $m \in \mathbb{R}$ and $\rho \geq 0$ to be the class of symbols $a(x, \xi)$ defined on $\mathbb{R}^n \times \mathbb{R}^n$ which are C^∞ in ξ and C^ρ in x , in the sense that

$$\text{for all } \alpha \in \mathbb{N}^n, \quad |\partial_\xi^\alpha a(x, \xi)| \langle \xi \rangle^{-m+|\alpha|} \in C^\rho(\mathbb{R}^n \times \mathbb{R}^n),$$

C^ϱ being replaced by L^∞ when $\varrho = 0$. If $a \in \Sigma_\varrho^m$, then m is the order of a and ϱ is its regularity. Following J.-M. Bony, we associate to a symbol a in Σ_ϱ^m the paradifferential operator $T_{a,\chi}$ defined by the expression

$$\widehat{T_{a,\chi}u}(\xi) = (2\pi)^{-n} \int_{\mathbb{R}^n} \chi(\xi - \eta, \eta) \mathcal{F}_1(a)(\xi - \eta, \eta) \hat{u}(\eta) d\eta,$$

where χ is what one calls a paratruncature, that is, a C^∞ function in $\mathbb{R}^n \times \mathbb{R}^n$ satisfying the following properties:

- (i) There exists $\varepsilon > 0$ such that $\varepsilon < 1$ and $\chi(\xi, \eta) = 0$ if $|\xi| \geq \varepsilon|\eta|$, $\xi, \eta \in \mathbb{R}^n$.
- (ii) There exist $\varepsilon' > 0$, $\varepsilon'' > 0$ such that $\varepsilon' < \varepsilon$ and $\chi(\xi, \eta) = 1$ if $|\xi| \leq \varepsilon'|\eta|$ and $|\eta| \geq \varepsilon''$.
- (iii) For all $\alpha \in \mathbb{N}^{2n}$, there exists $A_\alpha > 0$ such that for all $\zeta \in \mathbb{R}^{2n}$, we have $\langle \zeta \rangle^{|\alpha|} |\partial^\alpha \chi(\zeta)| \leq A_\alpha$.

The first important result on paradifferential operators is that, even if one can show that $T_{a,\chi} = \tilde{a}(x, D)$ with some $\tilde{a} \in S_{1,1}^m$, they are bounded in the Sobolev spaces in the usual manner. In fact, we have:

Theorem 2.6. *Assume that χ satisfies only the first and third properties among the above ones. Then, for every real s , the operator $T_{a,\chi}$ is bounded from $H^s(\mathbb{R}^n)$ into $H^{s-m}(\mathbb{R}^n)$ and its operator norm is estimated by a seminorm of a in Σ_ϱ^m . In particular, if $a = a(x) \in L^\infty(\mathbb{R}^n)$, then, for every real s , the operator $T_{a,\chi}$ is bounded in $H^s(\mathbb{R}^n)$ with an operator norm bounded by a constant times $\|a\|_{L^\infty}$.*

Proof. See [Bony 1981; Meyer 1981; Taylor 1991]. □

Concerning the dependence with respect to the paratruncature χ , one can say the following:

Theorem 2.7. *If $\varrho > 0$ and χ_1, χ_2 are paratruncatures, then the operator $T_{a,\chi_1} - T_{a,\chi_2}$ is bounded from $H^s(\mathbb{R}^n)$ into $H^{s-m+\varrho}(\mathbb{R}^n)$ and its operator norm is estimated by a seminorm of a in Σ_ϱ^m .*

Proof. See [Bony 1981; Meyer 1981; Taylor 1991]. □

This result shows that the dependence of $T_{a,\chi}$ on χ is less important than that on a . It also explains why the remainders in the paradifferential theory are only ϱ -regularizing. From now on, we shall write T_a instead of $T_{a,\chi}$ unless it is needed.

Note also that a possible choice of the paratruncature that we shall often use in the sequel is given by

$$\chi(\xi, \eta) = \chi_1(\xi/|\eta|)(1 - \psi_1(\eta)),$$

where $\psi_1, \chi_1 \in C^\infty(\mathbb{R}^n)$, $\psi_1 = 1$ in a neighbourhood of 0, $\psi_1 = 0$ out of $B(0, \varepsilon'')$, and $\chi_1 = 1$ on $B(0, \varepsilon')$, $\text{supp}(\chi) \subset B(0, \varepsilon)$, with ε and ε' satisfying $0 < \varepsilon' < \varepsilon < 1$. In this case, $T_{a,\chi} = \tilde{a}(x, D)$ with the following expression of \tilde{a} :

$$\tilde{a}(x, \xi) = (1 - \psi_1(\xi)) |\xi|^n \int_{\mathbb{R}^n} \mathcal{F}^{-1}(\chi_1)(|\xi|(x - y)) a(y, \xi) dy. \tag{2}$$

The following lemma gives some properties of \tilde{a} which will be needed in the sequel and often used implicitly.

Lemma 2.8. *Let $\varrho \geq 0$ and $a \in \Sigma_\varrho^m$. Then, \tilde{a} is smooth and*

$$|\partial_\xi^\beta \partial_x^\alpha \tilde{a}(x, \xi)| \leq A_{\alpha, \beta} \langle \xi \rangle^{m-|\beta|} \quad \text{if } |\alpha| \leq \varrho, \tag{3}$$

$$|\partial_\xi^\beta \partial_x^\alpha \tilde{a}(x, \xi)| \leq A_{\alpha, \beta} \langle \xi \rangle^{m-|\beta|+|\alpha|-\varrho} \quad \text{if } |\alpha| > \varrho, \tag{4}$$

where $A_{\alpha, \beta}$ are nonnegative constants; more precisely, the $A_{\alpha, \beta}$ can be estimated by seminorms of a in Σ_ϱ^m . In particular, $\tilde{a} \in S_{1,1}^m$.

Moreover, if θ is a smooth function with support in some compact subset of \mathbb{R}^n and $\theta_\mu(x) = \theta(x - \mu)$, $\mu \in \mathbb{Z}^n$, then, for all $N \in \mathbb{N}$, we have

$$\langle x - \mu \rangle^N |\partial_\xi^\beta \partial_x^\alpha \widetilde{\theta_\mu a}(x, \xi)| \leq A_{\alpha, \beta, N} \langle \xi \rangle^{m-|\beta|} \quad \text{if } |\alpha| \leq \varrho, \tag{5}$$

$$\langle x - \mu \rangle^N |\partial_\xi^\beta \partial_x^\alpha \widetilde{\theta_\mu a}(x, \xi)| \leq A_{\alpha, \beta, N} \langle \xi \rangle^{m-|\beta|+|\alpha|-\varrho} \quad \text{if } |\alpha| > \varrho, \tag{6}$$

where the $A_{\alpha, \beta, N}$ do not depend on μ and are estimated by seminorms of a in Σ_ϱ^m .

Proof. For the first part we refer to [Meyer 1981; Taylor 1991]. The second part follows from the first one by using, for example, for even N the decomposition

$$\langle x - \mu \rangle^N = \sum_\alpha \frac{(x - y)^\alpha}{\alpha!} \partial_y^\alpha \langle y - \mu \rangle^N$$

together with the expression (2). □

When dealing with nonlinear terms, we shall frequently use the following classical result:

Proposition 2.9. *If F is a C^∞ (or sufficiently regular) function in \mathbb{C}^m , $F(0) = 0$ and u_1, \dots, u_m are functions in $H^s(\mathbb{R}^n)$, $s > \frac{n}{2}$, then, $F(u_1, \dots, u_m) \in H^s(\mathbb{R}^n)$ and we have precisely*

$$\|F(u_1, \dots, u_m)\|_s \leq C(\|(u_1, \dots, u_m)\|_{L^\infty}) \|(u_1, \dots, u_m)\|_s,$$

where $\xi \mapsto C(\xi)$ is a nonnegative and nondecreasing function.

An important property of the paradifferential operators consists in the fact that they are necessary to write down Bony’s linearization formula, a formula that we recall here.

Theorem 2.10 (Bony’s linearization formula). *For all real functions $u_1, \dots, u_m \in H^{\frac{n}{2}+\varrho}(\mathbb{R}^n)$, $\varrho > 0$, and every function F of m real variables which is C^∞ (or sufficiently regular) and vanishes in 0, we have*

$$F(u_1, \dots, u_m) = \sum_{i=1}^{i=m} T_{\partial_{u_i}} F u_i + r \quad \text{with } r \in H^{\frac{n}{2}+2\varrho}(\mathbb{R}^n).$$

Proof. See [Bony 1981; Meyer 1981; Meyer 1982]. □

The remainder r in the above formula depends of course on (u_1, \dots, u_m) . The following result essentially shows that r is a locally Lipschitz function of (u_1, \dots, u_m) . More precisely:

Theorem 2.11. *If $u = (u_1, \dots, u_m) \in H^s(\mathbb{R}^n, \mathbb{R}^m)$, $s = \frac{n}{2} + \varrho$, $\varrho > 0$, let us denote by $r(u)$ the remainder in Bony’s formula. For all $u, v \in H^s(\mathbb{R}^n, \mathbb{R}^m)$, we have then*

$$\|r(u) - r(v)\|_{s+\varrho} \leq \theta(\|u\|_s, \|v\|_s)\|u - v\|_s,$$

where $\theta(\|u\|_s, \|v\|_s)$ is bounded if u and v vary in a bounded subset of $H^s(\mathbb{R}^n, \mathbb{R}^m)$.

Proof. See [Bienaimé 2014]. □

Remark. In the case of our equation, that is (1), even if u has complex values, we shall be able to apply Bony’s formula to the nonlinear expression $F(u, \nabla u, \bar{u}, \nabla \bar{u})$ where $u \in H^{\frac{n}{2}+1+\varrho}(\mathbb{R}^n)$. Indeed, we can write

$$F(u, \nabla u, \bar{u}, \nabla \bar{u}) = G(\operatorname{Re}(u), \nabla \operatorname{Re}(u), \operatorname{Im}(u), \nabla \operatorname{Im}(u))$$

where $G(x_1, x_2, y_1, y_2) = F(x_1 + iy_1, x_2 + iy_2, x_1 - iy_1, x_2 - iy_2)$ which is a function from \mathbb{R}^{2n+2} into \mathbb{C} . We apply then Bony’s formula to G and obtain that

$$F(u, \nabla u, \bar{u}, \nabla \bar{u}) = T_{\partial_{x_1}G} \operatorname{Re}(u) + T_{\partial_{x_2}G} \nabla \operatorname{Re}(u) + T_{\partial_{y_1}G} \operatorname{Im}(u) + T_{\partial_{y_2}G} \nabla \operatorname{Im}(u) + r(u).$$

At last, by using the fact that $\operatorname{Re}(u) = \frac{u+\bar{u}}{2}$, $\operatorname{Im}(u) = \frac{u-\bar{u}}{2i}$, $\partial_z = \frac{1}{2}(\partial_x - i\partial_y)$ and $\partial_{\bar{z}} = \frac{1}{2}(\partial_x + i\partial_y)$, and then the linearity of T_b with respect to b , we obtain the formula used in this paper:

$$F(u, \bar{u}, \nabla u, \nabla \bar{u}) = T_{\partial_u}Fu + T_{\partial_{\bar{u}}}F\bar{u} + T_{\partial_{\nabla u}}F\nabla u + T_{\partial_{\nabla \bar{u}}}F\nabla \bar{u} + r(u)$$

with $r(u) \in H^{\frac{n}{2}+2\varrho}(\mathbb{R}^n)$ if $u \in H^{\frac{n}{2}+1+\varrho}(\mathbb{R}^n)$.

We shall also often need the following result similar to Lemma 2.3:

Lemma 2.12. *Let $a \in \Sigma_0^0(\mathbb{R}^n)$, $\theta \in \mathcal{D}(\mathbb{R}^n)$, $\theta_\mu(x) = \theta(x - \mu)$, $\mu \in \mathbb{R}^n$ and $s \in \mathbb{R}$, and consider the paradifferential operator $T_{\theta_\mu a} = T_{\theta_\mu a, \chi}$ (where the paratruncature χ does not necessarily satisfy the second property of Definition 2.5). Then, for all $\sigma \geq 0$, the operator $\langle x - \mu \rangle^\sigma T_{\theta_\mu a} \langle x - \mu \rangle^\sigma$ is bounded in $H^s(\mathbb{R}^n)$ and there exist $N \in \mathbb{N}$ and a nonnegative constant C such that, for every $\mu \in \mathbb{R}^n$,*

$$\|\langle x - \mu \rangle^\sigma T_{\theta_\mu a} \langle x - \mu \rangle^\sigma\|_{\mathcal{L}(H^s)} \leq C \sup_{|\alpha| \leq N} \|\langle \xi \rangle^{|\alpha|} \partial_\xi^\alpha a\|_{L^\infty}.$$

Proof. First, one can assume that σ is an integer and even an even integer. Let us denote by a_μ the symbol $\theta_\mu a$ and consider first the operator $T_{a_\mu} \langle x - \mu \rangle^\sigma$. Recall that $T_{a_\mu} = \tilde{a}_\mu(x, D)$ with

$$\tilde{a}_\mu(x, \xi) = (1 - \psi_1(\xi))|\xi|^n \int_{\mathbb{R}^n} \mathcal{F}^{-1}(\chi_1)(|\xi|(x - y))a_\mu(y, \xi) dy. \tag{7}$$

where $\psi_1, \chi_1 \in C^\infty(\mathbb{R}^n)$, $\psi_1 = 1$ in a neighbourhood of 0, $\psi_1 = 0$ out of $B(0, \varepsilon'')$, and $\chi_1 = 1$ on $B(0, \varepsilon')$, $\operatorname{supp}(\chi) \subset B(0, \varepsilon)$, with ε and ε' satisfying $0 < \varepsilon' < \varepsilon < 1$. Hence, we can write for arbitrary

$u \in \mathcal{S}(\mathbb{R}^n)$,

$$\begin{aligned} T_{a_\mu} \langle x - \mu \rangle^\sigma u(x) &= (2\pi)^{-n} \int e^{ix\xi} \tilde{a}_\mu(x, \xi) \mathcal{F}(\langle x - \mu \rangle^\sigma u)(\xi) d\xi \\ &= (2\pi)^{-n} \int e^{ix\xi} \tilde{a}_\mu(x, \xi) \langle D_\xi + \mu \rangle^\sigma \hat{u}(\xi) d\xi \\ &= (2\pi)^{-n} \int \langle D_\xi - \mu \rangle^\sigma [e^{ix\xi} \tilde{a}_\mu(x, \xi)] \hat{u}(\xi) d\xi \\ &= (2\pi)^{-n} \sum_\alpha \frac{1}{\alpha!} D_x^\alpha [\langle x - \mu \rangle^\sigma] \int e^{ix\xi} \partial_\xi^\alpha \tilde{a}_\mu(x, \xi) \hat{u}(\xi) d\xi, \end{aligned}$$

where we have applied integrations by parts and the Leibniz formula. So, we have proved that

$$T_{a_\mu} \langle x - \mu \rangle^\sigma = \sum_\alpha \frac{1}{\alpha!} D_x^\alpha [\langle x - \mu \rangle^\sigma] (\partial_\xi^\alpha \tilde{a}_\mu)(x, D),$$

where the sum is of course finite. Now, let us consider the operator $(\partial_\xi^\alpha \tilde{a}_\mu)(x, D)$ and let us remark that, for example,

$$\begin{aligned} \partial_{\xi_k} \tilde{a}_\mu(x, \xi) &= (1 - \psi_1(\xi)) |\xi|^n \int_{\mathbb{R}^n} \mathcal{F}^{-1}(\chi_1)(|\xi|(x - y)) \partial_{\xi_k} a_\mu(y, \xi) dy \\ &\quad - (1 - \psi_1(\xi)) |\xi|^n \int_{\mathbb{R}^n} \mathcal{F}^{-1}(\chi_2)(|\xi|(x - y)) a_\mu(y, \xi) \frac{\xi_k}{|\xi|^2} dy \\ &\quad - \partial_k \psi_1(\xi) |\xi|^n \int_{\mathbb{R}^n} \mathcal{F}^{-1}(\chi_1)(|\xi|(x - y)) a_\mu(y, \xi) dy, \end{aligned}$$

where $\chi_2(\eta) = \sum_{j=1}^n \eta_j \partial_j \chi_1(\eta)$. This shows that

$$(\partial_{\xi_k} \tilde{a}_\mu)(x, D) = \sum_{l=1}^3 T_{\theta_\mu a^l, \chi^l},$$

where the a^l are symbols in Σ_0^{-1} and the χ^l are paratruncatures which satisfy the first and third properties of Definition 2.5. By induction, $(\partial_\xi^\alpha \tilde{a}_\mu)(x, D)$ is then a finite sum of operators of the same form as $T_{a_\mu} = T_{a_\mu, \chi}$ (of order $\leq -|\alpha|$), and note also that the seminorms of the associated symbols are bounded uniformly in μ by a seminorm of a . Hence, $T_{a_\mu} \langle x - \mu \rangle^\sigma$ is a finite sum of operators of the form $P(x - \mu) T_{a_\mu}$, where P is a polynomial (of degree $\leq \sigma$), and consequently the problem is reduced to the study of the operator $\langle x - \mu \rangle^\sigma T_{a_\mu}$ only. Now, the symbol of the latter can be written as

$$\begin{aligned} \langle x - \mu \rangle^\sigma \tilde{a}_\mu(x, \xi) &= \sum_{|\alpha| \leq \sigma} \frac{1}{\alpha!} (1 - \psi_1(\xi)) |\xi|^n \int_{\mathbb{R}^n} (x - y)^\alpha \mathcal{F}^{-1}(\chi_1)(|\xi|(x - y)) \partial_y^\alpha [\langle y - \mu \rangle^\sigma] a_\mu(y, \xi) dy \\ &= \sum_{|\alpha| \leq L} \frac{1}{\alpha!} (1 - \psi_1(\xi)) |\xi|^n \int_{\mathbb{R}^n} \mathcal{F}^{-1}(\chi_1^\alpha)(|\xi|(x - y)) \theta^\alpha(y - \mu) a^\alpha(y, \xi) dy, \end{aligned}$$

where χ_1^α and θ^α are similar to χ_1 and θ respectively, and $a^\alpha \in \Sigma_0^{-|\alpha|}$ with seminorms bounded by those of a . Hence, $\langle x - \mu \rangle^\sigma T_{a_\mu}$ is a finite sum of operators of the same form as T_{a_μ} whose symbols have seminorms bounded uniformly in μ by a seminorm of a . Eventually, the lemma follows from Theorem 2.6. □

Let us also recall the Gårding inequality which will be used crucially to prove the smoothing effect estimate.

Theorem 2.13 (sharp Gårding inequality for systems). *Let $a(x, \xi)$ be a $k \times k$ matrix whose elements are in $S_{1,0}^m$ and which satisfies*

$$\langle (a(x, \xi) + a^*(x, \xi))\zeta, \zeta \rangle \geq 0$$

for all $\zeta \in \mathbb{C}^k$ and all (x, ξ) such that $|\xi| \geq A_0$, where a^* denotes the adjoint matrix of a and $\langle \cdot, \cdot \rangle$ is the usual hermitian scalar product of \mathbb{C}^k . Then, there exist a nonnegative constant A and an integer N such that, for all $u \in \mathcal{S}(\mathbb{R}^n, \mathbb{C}^k)$, we have

$$\operatorname{Re} \langle a(x, D)u, u \rangle \geq -A \sup_{|\alpha|+|\beta| \leq N} \|\langle \xi \rangle^{|\beta|-m} \partial_x^\alpha \partial_\xi^\beta a\|_{L^\infty} \|u\|_{\frac{m-1}{2}}^2,$$

where A depends only on n, k and A_0 .

Proof. See [Taylor 1991; Tataru 2002] for example. □

3. The parilinear equation

In this section, we solve the Cauchy problem for the parilinear equation, that is, the linear equation obtained from (1) by applying Bony’s linearization formula (Theorem 2.10).

Recall that Q_μ is the cube $\mu + [0, 1]^n$, $\mu \in \mathbb{Z}^n$ and that Q_μ^* is a larger cube with side length 2, for example, $\mu + [-\frac{1}{2}, \frac{3}{2}]^n$.

Theorem 3.1. *Given $s \in \mathbb{R}$, consider the following linear Cauchy problem:*

$$\begin{cases} \partial_t u = i\mathcal{L}u + T_{b_1} \cdot \nabla_x u + T_{b_2} \cdot \nabla_x \bar{u} + C_1 u + C_2 \bar{u} + f(x, t), \\ u(x, 0) = u_0 \in H^s(\mathbb{R}^n). \end{cases} \tag{8}$$

We assume that C_1 and C_2 are bounded operators in $H^s(\mathbb{R}^n)$ and in $H^{s+2}(\mathbb{R}^n)$, that $b_1, b_2 \in \Sigma_\varrho^m$, $\varrho > 0$, and more precisely that

$$\begin{aligned} b_k(x, \xi) &= \sum_{\mu \in \mathbb{Z}^n} \alpha_{k,\mu} \varphi_{k,\mu}(x, \xi), & \sum_{\mu} |\alpha_{k,\mu}| &\leq A_k, \quad k = 1, 2, \\ \operatorname{supp}(x \mapsto \varphi_{k,\mu}(x, \xi)) &\subseteq Q_\mu^*, & \sup_{|\beta| \leq N_0} \|\langle \xi \rangle^{|\beta|} \partial_\xi^\beta \varphi_{k,\mu}\|_{C^0} &\leq 1, \end{aligned} \tag{9}$$

and $\|C_k\|_{\mathcal{L}(H^s)}, \|C_k\|_{\mathcal{L}(H^{s+2})} \leq A_k$, $k = 1, 2$, N_0 being a large and fixed integer. We further assume that $b_2(x, \xi)$ is even in ξ and that $f \in L_{\text{loc}}^1(\mathbb{R}, H^s(\mathbb{R}^n))$. Then, problem (8) has a unique solution u which is in $C(\mathbb{R}, H^s(\mathbb{R}^n))$ and satisfies, for all $T > 0$,

$$\sup_{-T \leq t \leq T} \|u(t)\|_s^2 \leq A(\|u_0\|_s^2 + I_T(J^s f, J^s u)), \tag{10}$$

$$\|J^{s+\frac{1}{2}} u\|_T^2 \leq A(\|u_0\|_s^2 + I_T(J^s f, J^s u)), \tag{11}$$

where the constant A depends only on $n, s, \varrho, T, A_1,$ and $A_2,$ and the expression $I_T(v, w)$ is a finite sum of terms of the form

$$\sup_{\mu \in \mathbb{Z}^n} \int_{-T}^T |\langle G_\mu v, w \rangle| dt$$

with $G_\mu \in \text{Op } S_{0,0}^0$ and the seminorms of its symbol (up to N_0) are uniformly bounded by a constant that depends only on s, n, ϱ, A_1 and $A_2.$

Recall that $\|u\|_T^2 = \sup_{\mu} \int_{-T}^T \int_{\mathbb{R}^n} \langle x - \mu \rangle^{-2\sigma_0} |u(x, t)|^2 dt dx,$ where $\sigma_0 > \frac{1}{2}$ is fixed.

Proof. Let us start by noting that the uniqueness is an obvious matter. Indeed, if u_1 and u_2 are solutions of (8), then, $u_1 - u_2$ is a solution of (8) with $u_0 = 0$ and $f = 0,$ and the conclusion follows from (10).

As for the existence, as is customary with linear differential equations, it will follow from the a priori estimates (10) and (11) by using more or less standard arguments of functional analysis, and the proof of Theorem 3.1 will consist essentially in establishing them.

Another useful remark is that it will be sufficient to prove the theorem in $C(\mathbb{R}_+, H^s(\mathbb{R}^n))$ instead of $C(\mathbb{R}, H^s(\mathbb{R}^n))$ and the estimates (10) and (11) on $[0, T]$ instead of $[-T, T].$ In fact, if the theorem is proved on $\mathbb{R}_+,$ one can apply it to $v(t) = u(-t),$ which satisfies a Cauchy problem of the same type as (8). The result is then that $v(-t)$ will extend u to \mathbb{R}_- and satisfy (8) on $\mathbb{R}_-,$ in addition to the fact that the estimates (10) and (11) are also extended to $[-T, 0].$

So, let us assume that $u \in C([0, T]; H^s(\mathbb{R}^n))$ is a solution of the Cauchy problem (8).

In what follows, it will be quite convenient to use the notation

$$v_N(\varphi) = \sup_{1 \leq j \leq N} \sup_{|\alpha|+|\beta| \leq N} \|\langle \xi \rangle^{|\beta|} \partial_x^\alpha \partial_\xi^\beta \varphi\|_{L^\infty}^j,$$

and note that such a quantity is not a norm in general but it is well-defined for $\varphi \in S_{1,0}^0.$ Note also that, if $M \geq 1, v_N(\varphi)^M \leq v_{NM}(\varphi),$ a remark that will be often used implicitly.

In fact, the inequalities (10) and (11) will be deduced from the following ones:

Proposition 3.2. *Assume that the functions $\varphi_{k,\mu}$ defining the b_k are $C^\infty;$ that is, $\varphi_{k,\mu} \in S_{1,0}^0, k = 1, 2.$ Then, there exist a positive real number A and an integer N such that, for all $R \geq 1,$ there exists a pseudodifferential operator $C \in \text{Op } S_{0,0}^0$ such that, for all $T > 0,$ any solution $u \in C([0, T]; H^s(\mathbb{R}^n))$ of the Cauchy problem (8) satisfies*

$$\begin{aligned} & \sup_{0 \leq t \leq T} \|Cu(t)\|_s^2 \\ & \leq \|Cu_0\|_s^2 + 2 \int_0^T |\langle C J^s f, C J^s u \rangle| dt + A \sup_{k,\mu} v_N(\varphi_{k,\mu}) \left(RT \sup_{0 \leq t \leq T} \|u(t)\|_s^2 + \frac{1}{R} \|J^{s+\frac{1}{2}} u\|_T^2 \right). \end{aligned}$$

Moreover, regarding the operator $C,$ we have the following precise bounds for $v \in H^s(\mathbb{R}^n) :$

$$\begin{aligned} \|Cv\|_s & \leq A \sup_{\mu} v_N(\varphi_{1,\mu}) \|v\|_s, \\ \|v\|_s & \leq A \sup_{\mu} v_N(\varphi_{1,\mu}) \|Cv\|_s + \frac{A}{R} \sup_{\mu} v_N(\varphi_{1,\mu})^2 \|v\|_s. \end{aligned}$$

Proposition 3.3. *Under the same assumptions as above and with the same elements A, R, C and N , there exist also pseudodifferential operators $\psi_j(x, D) \in \text{Op } S_{1,0}^0$, $j = 1, 2, 3, 4$, such that, for all $T > 0$, any solution $u \in C([0, T]; H^s(\mathbb{R}^n))$ of the Cauchy problem (8) satisfies*

$$\begin{aligned} \| \| J^{s+\frac{1}{2}} u \| \|_T^2 &\leq A(1 + T + T \sup_{\mu,k} v_N(\varphi_{k,\mu})) \sup_{[0,T]} \| u \|_s^2 + A \sum_{j=1}^4 \sup_{\mu \in \mathbb{Z}^n} \int_0^T |\langle \psi_j(x - \mu, D) J^s f, J^s u \rangle| dt, \\ \| \| J^{s+\frac{1}{2}} C u \| \|_T^2 &\leq A(1 + T + T \sup_{\mu,k} v_N(\varphi_{k,\mu})) \sup_{[0,T]} \| C u \|_s^2 + A \sum_{j=1}^4 \sup_{\mu} \int_0^T |\langle \psi_j(x - \mu, D) C J^s f, C J^s u \rangle| dt \\ &\quad + A \sup_{k,\mu} v_N(\varphi_{k,\mu}) \left(RT \sup_{0 \leq t \leq T} \| u(t) \|_s^2 + \frac{1}{R} \| \| J^{s+\frac{1}{2}} u \| \|_T^2 \right). \end{aligned}$$

Admitting these propositions (see Sections 5 and 6 for their proofs), let us go on and finish the proof of Theorem 3.1. In order to apply the above inequalities we have to regularize the b_k , $k = 1, 2$, by setting

$$\varphi_{k,\mu,m}(x, \xi) = m^n \int_{\mathbb{R}^n} \chi(m(x - y)) \varphi_{k,\mu}(y, \xi) dy \quad \text{and} \quad b_{k,m} = \sum_{\mu} \alpha_{k,\mu} \varphi_{k,\mu,m},$$

where χ is a nonnegative C^∞ function with support in the unit ball and whose integral is equal to 1. Note that $\varphi_{k,\mu,m}$ has its support (with respect to x) in a compact set which is slightly larger than Q_μ^* but this has no effect on the proofs. Since we can write

$$\partial_t u = i \mathcal{L} u + T_{b_{1,m}} \cdot \nabla_x u + T_{b_{2,m}} \cdot \nabla_x \bar{u} + C_1 u + C_2 \bar{u} + f_m,$$

where

$$f_m = f + T_{b_1 - b_{1,m}} \cdot \nabla u + T_{b_2 - b_{2,m}} \cdot \nabla \bar{u},$$

we can apply Proposition 3.2 to obtain

$$\begin{aligned} \sup_{[0,T]} \| C_m u \|_s^2 &\leq \| C_m u_0 \|_s^2 + 2 \int_0^T |\langle C_m J^s f_m, C_m J^s u \rangle| dt \\ &\quad + A \sup_{k,\mu} v_N(\varphi_{k,\mu,m}) \left(RT \sup_{[0,T]} \| u \|_s^2 + \frac{1}{R} \| \| J^{s+\frac{1}{2}} u \| \|_T^2 \right), \end{aligned}$$

where the operator C is denoted here by C_m to indicate its dependence on m . Now, clearly, we have

$$v_N(\varphi_{k,\mu,m}) \leq Am^{N^2} \sup_{1 \leq j \leq N} \sup_{|\beta| \leq N} \| \langle \xi \rangle^{|\beta|} \partial_\xi^\beta \varphi_{k,\mu} \|_{L^\infty}^j \leq Am^{N^2}.$$

Hence,

$$\begin{aligned} \sup_{[0,T]} \| C_m u \|_s^2 &\leq \| C_m u_0 \|_s^2 + 2 \int_0^T |\langle C_m J^s f, C_m J^s u \rangle| dt + 2 \int_0^T |\langle C_m J^s T_{b_1 - b_{1,m}} \nabla u, C_m J^s u \rangle| dt \\ &\quad + 2 \int_0^T |\langle C_m J^s T_{b_2 - b_{2,m}} \nabla \bar{u}, C_m J^s u \rangle| dt + Am^{N^2} \left(RT \sup_{[0,T]} \| u \|_s^2 + \frac{1}{R} \| \| J^{s+\frac{1}{2}} u \| \|_T^2 \right), \end{aligned}$$

and the problem now is to estimate the third and fourth terms in the right-hand side of this inequality. This is done in the following lemma.

Lemma 3.4. *Let \tilde{u} stand for u or \bar{u} , and $\sigma = \inf\{\varrho, 1\}$. Then, there exists a constant A such that, for all $k \in \{1, 2\}$, $m \geq 1$, $R \geq 1$ and $m' \geq m$,*

$$\int_0^T |\langle \mathbf{C}_m J^s T_{b_k - b_{k,m}} \nabla \tilde{u}, \mathbf{C}_m J^s u \rangle| dt \leq \left(\frac{Am^{2N^2}}{m'^\sigma} + \frac{Am'^{3N^2}}{R} \right) \| \| J^{s+\frac{1}{2}} u \| \|_T^2 + Am'^{3N^2} T \sup_{[0,T]} \|u\|_s^2 + \frac{A}{m^\sigma} \| \| J^{s+\frac{1}{2}} \mathbf{C}_m u \| \|_T^2.$$

See the Appendix for the proof of this lemma. Applying this lemma yields

$$\begin{aligned} \sup_{[0,T]} \| \mathbf{C}_m u \|_s^2 &\leq \| \mathbf{C}_m u_0 \|_s^2 + 2 \int_0^T |\langle \mathbf{C}_m J^s f, \mathbf{C}_m J^s u \rangle| dt + \frac{A}{m^\sigma} \| \| J^{s+\frac{1}{2}} \mathbf{C}_m u \| \|_T^2 \\ &\quad + \left(\frac{Am^{2N^2}}{m'^\sigma} + \frac{Am'^{3N^2}}{R} \right) \| \| J^{s+\frac{1}{2}} u \| \|_T^2 + Am'^{3N^2} RT \sup_{[0,T]} \|u\|_s^2, \end{aligned}$$

an inequality that we can improve, thanks to Proposition 3.3, as follows:

$$\begin{aligned} \sup_{[0,T]} \| \mathbf{C}_m u \|_s^2 &\leq \| \mathbf{C}_m u_0 \|_s^2 + 2 \int_0^T |\langle \mathbf{C}_m J^s f, \mathbf{C}_m J^s u \rangle| dt \\ &\quad + \frac{A}{m^\sigma} \sum_{j=1}^4 \sup_{\mu} \int_0^T |\langle \psi_j(x - \mu, D) \mathbf{C}_m J^s f, \mathbf{C}_m J^s u \rangle| dt + \frac{A(1 + Tm^N)}{m^\sigma} \sup_{[0,T]} \| \mathbf{C}_m u \|_s^2 \\ &\quad + \left(\frac{Am^{2N^2}}{m'^\sigma} + \frac{Am'^{3N^2}}{R} \right) \| \| J^{s+\frac{1}{2}} u \| \|_T^2 + Am'^{3N^2} RT \sup_{[0,T]} \|u\|_s^2 \\ &\leq \| \mathbf{C}_m u_0 \|_s^2 + 2 \int_0^T |\langle \mathbf{C}_m J^s f, \mathbf{C}_m J^s u \rangle| dt \\ &\quad + \frac{A}{m^\sigma} \sum_{j=1}^4 \sup_{\mu} \int_0^T |\langle \psi_j(x - \mu, D) \mathbf{C}_m J^s f, \mathbf{C}_m J^s u \rangle| dt + \frac{A(1 + Tm^N)}{m^\sigma} \sup_{[0,T]} \| \mathbf{C}_m u \|_s^2 \\ &\quad + \left(\frac{Am^{2N^2}}{m'^\sigma} + \frac{Am'^{3N^2}}{R} \right) \sum_{j=1}^4 \sup_{\mu} \int_0^T |\langle \psi_j(x - \mu, D) J^s f, J^s u \rangle| dt \\ &\quad + \left(\frac{Am^{2N^2}}{m'^\sigma} + \frac{Am'^{3N^2}}{R} \right) (1 + Tm^N) \sup_{[0,T]} \|u\|_s^2 + Am'^{3N^2} RT \sup_{[0,T]} \|u\|_s^2. \end{aligned}$$

Next, by taking m such that, for example, $m^\sigma \geq 4A$ and T such that $Tm^N \leq 1$, we get

$$\begin{aligned} \sup_{[0,T]} \| \mathbf{C}_m u \|_s^2 &\leq 2 \| \mathbf{C}_m u_0 \|_s^2 + 4 \int_0^T |\langle \mathbf{C}_m J^s f, \mathbf{C}_m J^s u \rangle| dt + \sum_{j=1}^4 \sup_{\mu} \int_0^T |\langle \psi_j(x - \mu, D) \mathbf{C}_m J^s f, \mathbf{C}_m J^s u \rangle| dt \\ &\quad + \left(\frac{2Am^{2N^2}}{m'^\sigma} + \frac{2Am'^{3N^2}}{R} \right) \sum_{j=1}^4 \sup_{\mu} \int_0^T |\langle \psi_j(x - \mu, D) J^s f, J^s u \rangle| dt \\ &\quad + \left(\frac{Am^{2N^2}}{m'^\sigma} + \frac{Am'^{3N^2}}{R} + Am'^{3N^2} RT \right) \sup_{[0,T]} \|u\|_s^2, \end{aligned}$$

and by using the second part of Proposition 3.2, we obtain

$$\begin{aligned} & \sup_{[0, T]} \|u\|_s^2 \\ & \leq Am^{2N^2} \left(m^{2N^2} \|u_0\|_s^2 + \int_0^T |\langle C_m J^s f, C_m J^s u \rangle| dt + \sum_{j=1}^4 \sup_{\mu} \int_0^T |\langle \psi_j(x-\mu, D) C_m J^s f, C_m J^s u \rangle| dt \right) \\ & \quad + \left(\frac{Am^{4N^2}}{m'^{\sigma}} + \frac{Am'^{5N^2}}{R} \right) \sum_{j=1}^4 \sup_{\mu} \int_0^T |\langle \psi_j(x-\mu, D) J^s f, J^s u \rangle| dt + C(m, m', R, T) \sup_{[0, T]} \|u\|_s^2, \end{aligned}$$

where

$$C(m, m', R, T) = \frac{Am^{4N^2}}{m'^{\sigma}} + \frac{Am'^{5N^2}}{R} + Am'^{5N^2} RT + \frac{Am^{4N^2}}{R^2}.$$

Finally, since m is fixed (and depends only on A), we take m' such that $Am^{4N^2}/m'^{\sigma} \leq \frac{1}{8}$, then we take R such that $Am'^{5N^2}/R \leq \frac{1}{8}$ and $Am^{4N^2}/R^2 \leq \frac{1}{8}$, and last we take T such that $Am'^{5N^2} RT \leq \frac{1}{8}$. With these choices, we have of course $C(m, m', R, T) \leq \frac{1}{2}$, which allows to bound $\sup_{[0, T]} \|u\|_s^2$ and to get (10) (and also (11), thanks to Proposition 3.3) with

$$\begin{aligned} I_T(v, w) = & \int_0^T |\langle C^* C v, w \rangle| dt + \sum_{j=1}^4 \sup_{\mu} \int_0^T |\langle C^* \psi_j(x-\mu, D) C v, w \rangle| dt \\ & + \sup_{\mu} \int_0^T |\langle \psi_j(x-\mu, D) v, w \rangle| dt. \end{aligned}$$

In fact, we have proved (10) and (11) only for $T = T_0$ and T_0 is sufficiently small. Let us show, if $T_0 < T$, that they hold true in the whole interval $[0, T]$ where the solution u is defined. Indeed, note first that the T_0 as determined above depends only on the constant A (so, only on n, s, ϱ, A_1 and A_2) but not on the given function (or distribution) f . Next, take a $T_1 \leq T_0$ such that $T_1 = T/n_1$, with some integer $n_1 \geq 2$. Then, if we consider the function $v(x, t) = u(x, t + T_1)$, we note that v is a solution (defined at least in $[0, T - T_1]$) of (8) with $v(0) = u(T_1)$ and $g(x, t) = f(x, t + T_1)$ instead of $f(x, t)$. It follows from the above arguments that v satisfies (10) and (11) for $T = T_0$ and hence for $T = T_1$. Since

$$\begin{aligned} \sup_{[T_1, 2T_1]} \|u\|_s^2 &= \sup_{[0, T_1]} \|v\|_s^2 \leq A(\|u(T_1)\|_s^2 + I_{T_1}(J^s g, J^s v)) \leq A(\|u(T_1)\|_s^2 + I_{2T_1}(J^s f, J^s u)) \\ &\leq A(A\|u_0\|_s^2 + AI_{T_1}(J^s f, J^s u) + I_{2T_1}(J^s f, J^s u)) \\ &\leq (A^2 + A)(\|u_0\|_s^2 + I_{2T_1}(J^s f, J^s u)), \end{aligned}$$

we obtain that u satisfies (10) and (11) for $T = 2T_1$ and with the constant $A^2 + A$ instead of A . Repeating this argument, we obtain that u satisfies (10) and (11) on $[0, n_1 T_1] = [0, T]$ and with the constant $\sum_{j=1}^{n_1} A^j \simeq A^{T/T_1}$ instead of A .

As for the existence, let us consider the approximating Cauchy problem

$$\begin{cases} \partial_t u = i\mathcal{L}u + T_{b_1} \nabla h(\varepsilon D)u + T_{b_2} \nabla h(\varepsilon D)\bar{u} + C_1 u + C_2 \bar{u} + f(x, t), \\ u(x, 0) = u_0 \in H^s(\mathbb{R}^n), \end{cases} \tag{12}$$

where h is a nonnegative C^∞ function on \mathbb{R}^n which is equal to 1 near 0 and has a compact support. It is easy to see, if $\int_0^T \|f\|_s dt < +\infty$, that the above problem has a unique solution, denoted by u_ε , which is

in $C([0, T]; H^s(\mathbb{R}^n))$. Indeed, the Cauchy problem (12) is clearly equivalent to the integral equation

$$u = e^{it\mathcal{L}}u_0 + \int_0^t e^{i(t-t')\mathcal{L}}(T_{b_1}\nabla h(\varepsilon D)u + T_{b_2}\nabla h(\varepsilon D)\bar{u} + C_1u + C_2\bar{u} + f(x, t')) dt'$$

and one can easily show that the map defined by the right-hand side of this equation is a contraction in $C([0, T_\varepsilon]; H^s(\mathbb{R}^n))$ with some $T_\varepsilon > 0$ sufficiently small, which allows one to apply the fixed-point theorem and to get a solution u_ε . Now, since T_ε does not depend on the data u_0 and f , one can extend u_ε to a solution of (12) on the whole interval $[0, T]$.

The idea now is to let ε tend to 0. This is possible because u_ε satisfies the estimates (10) and (11) and even uniformly with respect to ε . Indeed, it is sufficient to remark that the Cauchy problem (12) is of the same type as (8) because we can write

$$T_{b_k}\nabla h(\varepsilon D) = T_{b_{k,\varepsilon}}\nabla,$$

where $b_{k,\varepsilon}(x, \xi) = b_k(x, \xi)h(\varepsilon\xi)$ and $b_{k,\varepsilon}$ satisfies the assumptions of Theorem 3.1 uniformly in ε . Hence, we have in particular

$$\sup_{[0,T]} \|u_\varepsilon\|_s^2 \leq A\|u_0\|_s^2 + AI_T(J^s f, J^s u_\varepsilon),$$

and it follows from the Calderón–Vaillancourt theorem that

$$AI_T(J^s f, J^s u_\varepsilon) \leq AA' \sup_{[0,T]} \|u_\varepsilon\|_s \int_0^T \|f\|_s dt \leq \frac{1}{2} \sup_{[0,T]} \|u_\varepsilon\|_s^2 + \frac{1}{2}(AA')^2 \left(\int_0^T \|f\|_s dt \right)^2,$$

so that,

$$\sup_{[0,T]} \|u_\varepsilon\|_s \leq A\|u_0\|_s + A \int_0^T \|f\|_s dt. \tag{13}$$

Next, to check the convergence of u_ε , let us consider $v = u_\varepsilon - u_{\varepsilon'}$. It is clear that v is the solution of (12) with $u_0 = 0$ and

$$f = T_{b_1}\nabla(h(\varepsilon D) - h(\varepsilon' D))u_{\varepsilon'} + T_{b_2}\nabla(h(\varepsilon D) - h(\varepsilon' D))\bar{u}_{\varepsilon'}.$$

Therefore, it follows from (13) that

$$\sup_{[0,T]} \|v\|_s \leq A \int_0^T \|T_{b_1}\nabla(h(\varepsilon D) - h(\varepsilon' D))u_{\varepsilon'} + T_{b_2}\nabla(h(\varepsilon D) - h(\varepsilon' D))\bar{u}_{\varepsilon'}\|_s dt, \tag{14}$$

and from the boundedness of the T_{b_k} in the Sobolev spaces that

$$\sup_{[0,T]} \|v\|_s \leq A|\varepsilon - \varepsilon'| \int_0^T \|u_{\varepsilon'}\|_{s+2} dt \leq A|\varepsilon - \varepsilon'|T \sup_{[0,T]} \|u_{\varepsilon'}\|_{s+2}, \tag{15}$$

that is, thanks to (13),

$$\sup_{[0,T]} \|u_\varepsilon - u_{\varepsilon'}\|_s \leq A|\varepsilon - \varepsilon'| \left(\|u_0\|_{s+2} + \int_0^T \|f\|_{s+2} dt \right), \tag{16}$$

which proves that (u_ε) is a Cauchy sequence in $C([0, T]; H^s(\mathbb{R}^n))$ if one assumes that $u_0 \in H^{s+2}(\mathbb{R}^n)$ and $f \in L^1([0, T]; H^{s+2}(\mathbb{R}^n))$. In this case, $u_\varepsilon \rightarrow u$ in $C([0, T]; H^s(\mathbb{R}^n))$ when $\varepsilon \rightarrow 0$, and by passing to the limit in (12), we obtain that u is a solution of (8). Moreover, by passing to the limit in (13), we get

$$\sup_{[0, T]} \|u\|_s \leq A \left(\|u_0\|_s + \int_0^T \|f\|_s dt \right). \tag{17}$$

Now, if we have only $u_0 \in H^s(\mathbb{R}^n)$ and $f \in L^1([0, T]; H^s(\mathbb{R}^n))$, by density of the smooth functions, we can take sequences (u_0^j) in $H^{s+2}(\mathbb{R}^n)$ and (f^j) in $L^1([0, T]; H^{s+2}(\mathbb{R}^n))$ such that $\|u_0^j - u_0\|_s \rightarrow 0$ and $\int_0^T \|f^j - f\|_s dt \rightarrow 0$, and we can consider the solution u^j of (8) associated to the data u_0^j and f^j . Then, $u^j - u^k$ is the solution of (8) associated to the data $u_0^j - u_0^k$ and $f^j - f^k$. Hence, thanks to (17),

$$\sup_{[0, T]} \|u^j - u^k\|_s \leq A \left(\|u_0^j - u_0^k\|_s + \int_0^T \|f^j - f^k\|_s dt \right),$$

which shows that (u^j) is a Cauchy sequence in $C([0, T]; H^s(\mathbb{R}^n))$ which is then convergent to some $u \in C([0, T]; H^s(\mathbb{R}^n))$. Of course, u is a solution of (8) and satisfies the estimates (10), (11) and also (17). This achieves the proof of Theorem 3.1. □

4. The nonlinear equation

Consider the nonlinear Cauchy problem

$$\begin{cases} \partial_t u = i \mathcal{L}u + F(u, \nabla_x u, \bar{u}, \nabla_x \bar{u}), & t \in \mathbb{R}, x \in \mathbb{R}^n, \\ u(x, 0) = u_0(x) \in H^s(\mathbb{R}^n), \end{cases} \tag{18}$$

where the function $F(u, v, \bar{u}, \bar{v})$ is sufficiently regular in $\mathbb{C} \times \mathbb{C}^n \times \mathbb{C} \times \mathbb{C}^n$ and vanishes to the third order at 0, the operator \mathcal{L} has the form

$$\mathcal{L} = \sum_{j \leq k} \partial_{x_j}^2 - \sum_{j > k} \partial_{x_j}^2,$$

with a fixed $k \in \{1, 2, \dots, n\}$, $H^s(\mathbb{R}^n)$ is the usual Sobolev space on \mathbb{R}^n , and $s = \frac{n}{2} + 2 + \varrho$, $\varrho > 0$. Using Bony’s linearization formula, (18) is equivalent to

$$\begin{cases} \partial_t u = i \mathcal{L}u + T_{b_1} \nabla_x u + T_{b_2} \nabla_x \bar{u} + T_{a_1} u + T_{a_2} \bar{u} + R(u, \nabla_x u, \bar{u}, \nabla_x \bar{u}), \\ u(x, 0) = u_0(x) \in H^s(\mathbb{R}^n), \end{cases} \tag{19}$$

where $R(u, \nabla_x u, \bar{u}, \nabla_x \bar{u})$ is Bony’s remainder and

$$\begin{aligned} b_1 &= \partial_v F(u, \nabla_x u, \bar{u}, \nabla_x \bar{u}), & b_2 &= \partial_{\bar{v}} F(u, \nabla_x u, \bar{u}, \nabla_x \bar{u}), \\ a_1 &= \partial_u F(u, \nabla_x u, \bar{u}, \nabla_x \bar{u}), & a_2 &= \partial_{\bar{u}} F(u, \nabla_x u, \bar{u}, \nabla_x \bar{u}). \end{aligned}$$

Recall that $R(u, \nabla_x u, \bar{u}, \nabla_x \bar{u}) \in H^{2(s-1)-\frac{n}{2}}(\mathbb{R}^n)$ if $u \in H^s(\mathbb{R}^n)$, $s > \frac{n}{2} + 1$. Note also that it follows from Proposition 2.9 that the b_j and a_j , $j = 1$ or 2 , are in $H^{s-1}(\mathbb{R}^n)$ if $u \in H^s(\mathbb{R}^n)$, $s > \frac{n}{2} + 1$, and that

$$\|b_j\|_{s-1} \leq C(\|u\|_{L^\infty}, \|\nabla u\|_{L^\infty}) \|u\|_s, \quad \|a_j\|_{s-1} \leq C(\|u\|_{L^\infty}, \|\nabla u\|_{L^\infty}) \|u\|_s, \quad j = 1, 2.$$

Moreover, by introducing the notation

$$\begin{aligned} b_1^0 &= \partial_v F(u_0, \nabla_x u_0, \bar{u}_0, \nabla_x \bar{u}_0), & b_2^0 &= \partial_{\bar{v}} F(u_0, \nabla_x u_0, \bar{u}_0, \nabla_x \bar{u}_0), \\ a_1^0 &= \partial_u F(u_0, \nabla_x u_0, \bar{u}_0, \nabla_x \bar{u}_0), & a_2^0 &= \partial_{\bar{u}} F(u_0, \nabla_x u_0, \bar{u}_0, \nabla_x \bar{u}_0), \end{aligned}$$

the above Cauchy problem is in fact equivalent to

$$\begin{cases} \partial_t u = i\mathcal{L}u + T_{b_1^0} \nabla_x u + T_{b_2^0} \nabla_x \bar{u} + T_{a_1^0} u + T_{a_2^0} \bar{u} + \tilde{R}(u, \nabla_x u, \bar{u}, \nabla_x \bar{u}), \\ u(x, 0) = u_0(x) \in H^s(\mathbb{R}^n), \end{cases} \tag{20}$$

where

$$\tilde{R}(u, \nabla_x u, \bar{u}, \nabla_x \bar{u}) = T_{b_1 - b_1^0} \nabla_x u + T_{b_2 - b_2^0} \nabla_x \bar{u} + T_{a_1 - a_1^0} u + T_{a_2 - a_2^0} \bar{u} + R(u, \nabla_x u, \bar{u}, \nabla_x \bar{u}). \tag{21}$$

Clearly, the last Cauchy problem is of the same type as (8), which is studied in Theorem 3.1, and in fact we are going to apply that theorem to

$$\begin{cases} \partial_t u = i\mathcal{L}u + T_{b_1^0} \nabla_x u + T_{b_2^0} \nabla_x \bar{u} + T_{a_1^0} u + T_{a_2^0} \bar{u} + f, \\ u(x, 0) = u_0(x) \in H^s(\mathbb{R}^n). \end{cases} \tag{22}$$

This is possible because b_1^0 and b_2^0 satisfy the assumptions of Theorem 3.1. Indeed, it follows from the Taylor formula and the assumptions on F that one can write for example

$$b_1^0 = \partial_v F(z_0) = u_0 G_1(z_0) + \nabla_x u_0 G_2(z_0) + \bar{u}_0 G_3(z_0) + \nabla_x \bar{u}_0 G_4(z_0), \tag{23}$$

where $z_0 = (u_0, \nabla_x u_0, \bar{u}_0, \nabla_x \bar{u}_0)$ and G_1, G_2, G_3 and G_4 are sufficiently regular and vanish at 0. Since $s - 1 > \frac{n}{2}$, we know that the $G_i(z_0)$ are in $H^{s-1}(\mathbb{R}^n)$ and it follows from (23) and Lemma 2.4 that b_1^0 satisfies the assumption (9) of Theorem 3.1; that is, one can write

$$b_1^0 = \sum_{\mu} \alpha_{1,\mu} \varphi_{1,\mu},$$

where $\alpha_{1,\mu} = \|q_{\mu} b_1^0\|_{H^{s-1}}$, $\varphi_{1,\mu} = q_{\mu} b_1^0 / \alpha_{1,\mu}$, and $\sum_{\mu} q_{\mu} = 1$ is a smooth partition of unity with $q_{\mu}(x) = q(x - \mu)$ and $\text{supp}(q) \subset Q_0^*$. Note that we have precisely the bound

$$\begin{aligned} \sum_{\mu} \|q_{\mu} b_1^0\|_{H^{s-1}} &\leq C (\|u_0\|_{H^{s-1}} \|G_1(z_0)\|_{H^{s-1}} + \|\nabla_x u_0\|_{H^{s-1}} \|G_2(z_0)\|_{H^{s-1}} \\ &\quad + \|\bar{u}_0\|_{H^{s-1}} \|G_3(z_0)\|_{H^{s-1}} + \|\nabla_x \bar{u}_0\|_{H^{s-1}} \|G_4(z_0)\|_{H^{s-1}}), \end{aligned}$$

with some positive constant C . Of course, the same is true for b_2^0 . Moreover, since a_1^0 and a_2^0 are bounded (they are in $H^{s-1}(\mathbb{R}^n)$), the paramultiplication operators $T_{a_1^0}$ and $T_{a_2^0}$ are bounded in $H^s(\mathbb{R}^n)$.

Now, by application of Theorem 3.1 to (22), let us consider the unique solution of (22) with $f = 0$ and denote it by $U(t)u_0$.

Next, for $T > 0$, let us define the norms $\lambda_1(w)$, $\lambda_2(w)$, $\lambda_3(w)$ and $\lambda(w)$ by

$$\lambda_1(w) = \sup_{[0,T]} \|w\|_s, \quad \lambda_2(w) = \|J^{s+\frac{1}{2}} w\|_T, \quad \lambda_3(w) = \sup_{[0,T]} \|\partial_t w\|_{s-2}, \quad \lambda(w) = \max_{1 \leq i \leq 3} \lambda_i(w),$$

the space Z by

$$Z = \{w \in C([0, T]; H^s(\mathbb{R}^n)) : w(x, 0) = u_0(x) \text{ and } \lambda(w) \leq K\},$$

where the positive constant K is to be determined later, and, for $w \in C([0, T]; H^s(\mathbb{R}^n))$, the operator Υ by

$$\Upsilon w(t) = U(t)u_0 + \int_0^t U(t-t')\tilde{R}(w(t'), \nabla_x w(t'), \bar{w}(t'), \nabla_x \bar{w}(t')) dt'.$$

Let us first remark that Υw satisfies

$$\begin{cases} \partial_t \Upsilon w = i\mathcal{L}\Upsilon w + T_{b_1^0} \nabla_x \Upsilon w + T_{b_2^0} \nabla_x \overline{\Upsilon w} + T_{a_1^0} \Upsilon w + T_{a_2^0} \overline{\Upsilon w} + \tilde{R}(w, \nabla_x w, \bar{w}, \nabla_x \bar{w}), \\ \Upsilon w(0) = u_0, \end{cases} \tag{24}$$

and that a fixed point of Υ will be a solution of (20), hence, a solution of (18). So, in what follows, we are going to study $\lambda(\Upsilon w)$ in order to prove that Υ has a fixed point in the complete metric space (Z, λ) . Let us also note that since the life time T will be small, we can assume from now on that $T \leq 1$.

We start by applying Theorem 3.1 to (24). It follows from (10) and (11) that

$$\max\{\lambda_1(\Upsilon w)^2, \lambda_2(\Upsilon w)^2\} \leq A(\|u_0\|_s^2 + I_T(J^s \tilde{R}, J^s \Upsilon w)), \tag{25}$$

where, for simplicity, $\tilde{R} = \tilde{R}(w, \nabla_x w, \bar{w}, \nabla_x \bar{w})$ and $I_T(u, v)$ is a finite sum of terms of the form

$$\sup_{\mu \in \mathbb{Z}^n} \int_0^T |\langle G_\mu u, v \rangle| dt,$$

where $G_\mu \in \text{Op } S_{0,0}^0$ and the seminorms of its symbol are uniformly bounded with respect to μ . Recall that the constant A depends only on n, s and u_0 and we remark right now a fact that will be useful later: if we let u_0 vary in a bounded subset of $H^s(\mathbb{R}^n)$, it follows from the linear theory that we can take the constant A in the above inequality that depends only on that bounded set. The same remark holds for $\sup_\mu \|G_\mu\|_{\mathcal{L}(L^2)}$ or the seminorms of the operators G_μ uniformly in μ .

Thus, we have to estimate uniformly in μ the sum

$$\begin{aligned} & \int_0^T |\langle G_\mu J^s T_{b_1-b_1^0} \nabla_x w, J^s \Upsilon w \rangle| dt + \int_0^T |\langle G_\mu J^s T_{b_2-b_2^0} \nabla_x \bar{w}, J^s \Upsilon w \rangle| dt \\ & + \int_0^T |\langle G_\mu J^s T_{a_1-a_1^0} w, J^s \Upsilon w \rangle| dt + \int_0^T |\langle G_\mu J^s T_{a_2-a_2^0} \bar{w}, J^s \Upsilon w \rangle| dt \\ & + \int_0^T |\langle G_\mu J^s R(w, \nabla_x w, \bar{w}, \nabla_x \bar{w}), J^s \Upsilon w \rangle| dt. \end{aligned} \tag{26}$$

First, let us consider the third term. It follows from the preceding remark, the Cauchy–Schwarz inequality, the Calderón–Vaillancourt theorem and Theorem 2.6 that

$$\int_0^T |\langle G_\mu J^s T_{a_1-a_1^0} w, J^s \Upsilon w \rangle| dt \leq A \|a_1 - a_1^0\|_{L^\infty} \int_0^T \|w\|_s \|\Upsilon w\|_s dt,$$

and from Proposition 2.9 that

$$\|a_1 - a_1^0\|_{L^\infty} \leq C(\|w\|_s)\|w\|_s + C(\|u_0\|_s)\|u_0\|_s \leq C(K)K + C(\|u_0\|_s)\|u_0\|_s \leq 2C(K)K.$$

Hence,

$$\int_0^T |(G_\mu J^s T_{a_1 - a_1^0} w, J^s \Upsilon w)| dt \leq ATC(K)\lambda_1(w)\lambda_1(\Upsilon w) \leq ATC(K)\lambda(w)\lambda(\Upsilon w), \quad (27)$$

with a modified constant $C(K)$.

The fourth term of (26) is treated in the same manner.

Now, let us estimate the first term of (26). Using a smooth partition of unity $1 = \sum_{v \in \mathbb{Z}^n} \chi_v$, with $\chi_v(x) = \chi(x - v)$ and χ having a compact support, we can write

$$\begin{aligned} &\langle G_\mu J^s T_{b_1 - b_1^0} \nabla_x w, J^s \Upsilon w \rangle \\ &= \sum_v \langle J^{-\frac{1}{2}} G_\mu J^s T_{\chi_v(b_1 - b_1^0)} \nabla_x w, J^{s+\frac{1}{2}} \Upsilon w \rangle \\ &= \sum_v \langle G_{\mu,v} \langle x - v \rangle^{\sigma_0} T_{\chi_v(b_1 - b_1^0)} \langle x - v \rangle^{\sigma_0} H_\mu \langle x - v \rangle^{-\sigma_0} J^{s+\frac{1}{2}} w, \langle x - v \rangle^{-\sigma_0} J^{s+\frac{1}{2}} \Upsilon w \rangle, \end{aligned}$$

where

$$G_{\mu,v} = \langle x - v \rangle^{\sigma_0} J^{-\frac{1}{2}} G_\mu J^s \langle x - v \rangle^{-\sigma_0}, \quad H_v = \langle x - v \rangle^{-\sigma_0} J^{-s-\frac{1}{2}} \nabla \langle x - v \rangle^{\sigma_0}.$$

Next, it follows from the pseudodifferential composition formula and from Lemma 2.3 that $G_{\mu,v}$ is in $\text{Op } S_{0,0}^{s-\frac{1}{2}}$, H_v is in $\text{Op } S_{1,0}^{\frac{1}{2}-s}$, and that their seminorms are uniformly bounded with respect to μ and v . Going back to the first term of (26), these considerations in addition to Lemma 2.12 allow us to estimate it as follows:

$$\begin{aligned} &\int_0^T |\langle G_\mu J^s T_{b_1 - b_1^0} \nabla_x w, J^s \Upsilon w \rangle| dt \\ &\leq \sum_v \int_0^T \|G_{\mu,v} \langle x - v \rangle^{\sigma_0} T_{\chi_v(b_1 - b_1^0)} \langle x - v \rangle^{\sigma_0} H_v\|_{\mathcal{L}(L^2)} \left\| \frac{J^{s+\frac{1}{2}} w}{\langle x - v \rangle^{\sigma_0}} \right\|_0 \left\| \frac{J^{s+\frac{1}{2}} \Upsilon w}{\langle x - v \rangle^{\sigma_0}} \right\|_0 dt \\ &\leq A \sum_v \int_0^T \|\langle x - v \rangle^{\sigma_0} T_{\chi_v(b_1 - b_1^0)} \langle x - v \rangle^{\sigma_0}\|_{\mathcal{L}(H^{s-\frac{1}{2}})} \left\| \frac{J^{s+\frac{1}{2}} w}{\langle x - v \rangle^{\sigma_0}} \right\|_0 \left\| \frac{J^{s+\frac{1}{2}} \Upsilon w}{\langle x - v \rangle^{\sigma_0}} \right\|_0 dt \\ &\leq A \sum_v \int_0^T \|\chi_v(b_1 - b_1^0)\|_{L^\infty} \left\| \frac{J^{s+\frac{1}{2}} w}{\langle x - v \rangle^{\sigma_0}} \right\|_0 \left\| \frac{J^{s+\frac{1}{2}} \Upsilon w}{\langle x - v \rangle^{\sigma_0}} \right\|_0 dt \\ &\leq A \sum_v \sup_{[0,T]} \|\chi_v(b_1 - b_1^0)\|_{L^\infty} \|J^{s+\frac{1}{2}} w\|_T \|J^{s+\frac{1}{2}} \Upsilon w\|_T. \end{aligned}$$

Now, it follows from the Taylor formula that we can write

$$\begin{aligned} b_1 - b_1^0 &= \partial_v F(z) - \partial_v F(z_0) \\ &= (w - u_0)G_1(z_0, z) + \nabla_x(w - u_0)G_2(z_0, z) + (\bar{w} - \bar{u}_0)G_3(z_0, z) + \nabla_x(\bar{w} - \bar{u}_0)G_4(z_0, z), \end{aligned}$$

where, for simplicity, $z_0 = (u_0, \nabla_x u_0, \bar{u}_0, \nabla_x \bar{u}_0)$ and $z = (w, \nabla_x w, \bar{w}, \nabla_x \bar{w})$, and the G_k are functions of the form

$$\int_0^1 F_k(z_0 + \tau(z - z_0)) d\tau,$$

where F_k is a second-order partial derivative of F . Next, it follows from the assumption on F that $G_k(0, 0) = 0$ for all k , from which one deduces easily that

$$\|\chi_\nu(b_1 - b_1^0)\|_{L^\infty} \leq C(\|(z_0, z)\|_{L^\infty}) \|\chi_\nu(z_0, z)\|_{L^\infty} \tilde{\chi}_\nu(z_0, z)\|_{L^\infty},$$

where $\tilde{\chi}_\nu$ is similar to χ_ν , and, by using the Sobolev injection, that is, Proposition A.5(i), that

$$\begin{aligned} \|\chi_\nu(b_1 - b_1^0)\|_{L^\infty} &\leq C(\|(z_0, z)\|_{L^\infty}) \|\chi_\nu(z_0, z)\|_{H^\sigma([0, T]; H^{s'})} \|\tilde{\chi}_\nu(z_0, z)\|_{H^\sigma([0, T]; H^{s'})} \\ &\leq C(K) \|\chi_\nu(u_0, w)\|_{H^\sigma([0, T]; H^{s'+1})} \|\tilde{\chi}_\nu(u_0, w)\|_{H^\sigma([0, T]; H^{s'+1})}, \end{aligned}$$

where $\sigma > \frac{1}{2}$ and $s' > \frac{n}{2}$. Thus, to obtain the summability in ν of $\|\chi_\nu(b_1 - b_1^0)\|_{L^\infty}$, it is sufficient to prove that $\|\chi_\nu(u_0, w)\|_{H^\sigma([0, T]; H^{s'+1})}$ is square summable in ν . To this end and to get an explicit bound for the sum, let us apply the interpolation inequality of Proposition A.5. This yields, by taking $\frac{1}{2} < \sigma < 1$,

$$\begin{aligned} \|\chi_\nu(u_0, w)\|_{H^\sigma([0, T]; H^{s'+1})} &\leq A \|\chi_\nu(u_0, w)\|_{L^2([0, T]; H^{s'+2})}^{1-\sigma} \|\chi_\nu(u_0, w)\|_{H^1([0, T]; H^{s''})}^\sigma \\ &\leq A (\|\chi_\nu(u_0, w)\|_{L^2([0, T]; H^{s'+2})} + \|\chi_\nu(u_0, w)\|_{L^2([0, T]; H^{s'+2})}^{1-\sigma} \|\chi_\nu \partial_t w\|_{L^2([0, T]; H^{s''})}^\sigma), \end{aligned}$$

where s'' is such that $(1 - \sigma)(s' + 2) + \sigma s'' = s' + 1$, that is, $s'' = s' + 2 - \frac{1}{\sigma}$. One can choose s' and σ such that $s'' = s - 2$, that is, such that $s' = s - 4 + \frac{1}{\sigma}$. In fact, if $\sigma = \frac{1}{2} + \varepsilon$, then $s' = \frac{n}{2} + \varrho - 4\varepsilon/(1 + 2\varepsilon)$, which is larger than $\frac{n}{2}$ if ε is small enough. With such a choice, we also have $s' + 2 < s$, so, the expressions $\|\chi_\nu(u_0, w)\|_{L^2([0, T]; H^{s'+2})}$ and $\|\chi_\nu \partial_t w\|_{L^2([0, T]; H^{s''})}$ are both square summable in ν , which shows that $\|\chi_\nu(u_0, w)\|_{H^\sigma([0, T]; H^{s'+1})}$ is itself square summable in ν and that, by applying Hölder's inequality,

$$\begin{aligned} \sum_\nu \|\chi_\nu(u_0, w)\|_{H^\sigma([0, T]; H^{s'+1})}^2 &\leq A \sum_\nu \|\chi_\nu(u_0, w)\|_{L^2([0, T]; H^{s'})}^2 + A \left(\sum_\nu \|\chi_\nu(u_0, w)\|_{L^2([0, T]; H^{s'})}^2 \right)^{1-\sigma} \left(\sum_\nu \|\chi_\nu \partial_t w\|_{L^2([0, T]; H^{s-2})}^2 \right)^\sigma \\ &\leq A (T\lambda_1(w)^2 + (T\lambda_1(w)^2)^{1-\sigma} (T\lambda_3(w)^2)^\sigma) \leq AT\lambda(w)^2, \end{aligned}$$

where, of course, the constant A changes from one inequality to the other. Consequently,

$$\sum_\nu \|\chi_\nu(b_1 - b_1^0)\|_{L^\infty} \leq AC(K)T\lambda(w)^2,$$

which allows us finally to bound the first term of (26) as follows:

$$\begin{aligned} \int_0^T |(G_\mu J^s T_{b_1 - b_1^0} \nabla_x w, J^s \Upsilon w)| dt &\leq AC(K)T\lambda(w)^2 \lambda_2(w) \lambda_2(\Upsilon w) \\ &\leq AC(K)K^2 T\lambda(w) \lambda(\Upsilon w). \end{aligned} \tag{28}$$

The second term of (26) is treated in the same manner.

Let us now consider the last term of (26). As above, let z stand for $(w, \nabla_x w, \bar{w}, \nabla_x \bar{w})$. As $z \in H^{s-1}(\mathbb{R}^n) = H^{\frac{n}{2}+1+\varrho}(\mathbb{R}^n)$, it follows from Bony’s formula, that is, Theorem 2.10, that $R(z) \in H^{2(s-1)-\frac{n}{2}}(\mathbb{R}^n) = H^{s+\varrho}(\mathbb{R}^n)$ and that

$$\|R(z)\|_{s+\varrho} \leq C(K)\|z\|_{s-1} \leq C(K)\|w\|_s.$$

Hence,

$$\begin{aligned} \int_0^T |\langle G_\mu J^s R(z), J^s \Upsilon w \rangle| dt &\leq A \int_0^T \|R(z)\|_s \|\Upsilon w\|_s dt \leq AC(K) \int_0^T \|w\|_s \|\Upsilon w\|_s dt \\ &\leq AC(K)T\lambda_1(w)\lambda_1(\Upsilon w) \leq AC(K)T\lambda(w)\lambda(\Upsilon w). \end{aligned} \tag{29}$$

Thus, we have bounded all the terms of (26), which leads to the estimate

$$\max\{\lambda_1(\Upsilon w), \lambda_2(\Upsilon w)\} \leq A\|u_0\|_s + \sqrt{AC(K)T\lambda(w)\lambda(\Upsilon w)}, \tag{30}$$

where the constants A and $C(K)$ have changed of course.

It remains to estimate $\lambda_3(\Upsilon w)$. Recall that Υw satisfies the Cauchy problem (24). Hence, applying Theorem 2.6 yields

$$\begin{aligned} \|\partial_t \Upsilon w\|_{s-2} &\leq \|\Upsilon w\|_s + A(\|b_1^0\|_{L^\infty} + \|b_2^0\|_{L^\infty})\|\Upsilon w\|_{s-1} \\ &\quad + A(\|a_1^0\|_{L^\infty} + \|a_2^0\|_{L^\infty})\|\Upsilon w\|_{s-2} + A(\|b_1 - b_1^0\|_{L^\infty} + \|b_2 - b_2^0\|_{L^\infty})\|w\|_{s-1} \\ &\quad + A(\|a_1 - a_1^0\|_{L^\infty} + \|a_2 - a_2^0\|_{L^\infty})\|w\|_{s-2} + \|R(z)\|_{s-2} \\ &\leq A\|\Upsilon w\|_s + A(\|b_1 - b_1^0\|_{L^\infty} + \|b_2 - b_2^0\|_{L^\infty} \\ &\quad + \|a_1 - a_1^0\|_{L^\infty} + \|a_2 - a_2^0\|_{L^\infty})\|w\|_s + \|R(z)\|_{s-2}. \end{aligned} \tag{31}$$

Now, as before, it follows from Proposition A.5 that

$$\|b_j - b_j^0\|_{L^\infty} \leq A\|b_j - b_j^0\|_{H^\sigma([0,T];H^{s'})} \leq A\|b_j - b_j^0\|_{L^2([0,T];H^{s'+1})}^{1-\sigma} \|b_j - b_j^0\|_{H^1([0,T];H^{s''})}^\sigma,$$

where $j = 1, 2$, $\sigma > \frac{1}{2}$, $s' > \frac{n}{2}$ and s'' is such that $(1 - \sigma)(s' + 1) + \sigma s'' = s'$. In fact, we can take $s'' = s - 3$, which corresponds to $s' = s + \frac{1}{\sigma} - 4 = \frac{n}{2} + \varrho + \frac{1}{\sigma} - 2$; so, $s' < s - 2$ and if σ is close enough to $\frac{1}{2}$, then, $s' > \frac{n}{2}$. Therefore, with such a choice, we have

$$\|b_j - b_j^0\|_{L^\infty} \leq A\|b_j - b_j^0\|_{L^2([0,T];H^{s-1})} + A\|b_j - b_j^0\|_{L^2([0,T];H^{s-1})}^{1-\sigma} \|\partial_t b_j\|_{L^2([0,T];H^{s-3})}^\sigma.$$

Next, applying Proposition 2.9 yields

$$\begin{aligned} \|b_j - b_j^0\|_{L^2([0,T];H^{s-1})}^2 &= \int_0^T \|b_j - b_j^0\|_{s-1}^2 dt \\ &\leq \int_0^T (C(\|z\|_{L^\infty})\|z\|_{s-1} + C(\|z_0\|_{L^\infty})\|z_0\|_{s-1})^2 dt \leq C(K)^2 T\lambda_1(w)^2, \end{aligned}$$

and

$$\begin{aligned} \|\partial_t b_j\|_{L^2([0,T];H^{s-3})}^2 &= \int_0^T \|(\partial_v F)'(z)\partial_t z\|_{s-3}^2 dt \leq A \int_0^T \|(\partial_v F)'(z)\|_{s-2}^2 \|\partial_t z\|_{s-3}^2 dt \\ &\leq A \int_0^T \|(\partial_v F)'(z)\|_{s-2}^2 \|\partial_t w\|_{s-2}^2 dt \leq ATC(K)^2 \lambda_3(w)^2, \end{aligned}$$

which imply that

$$\|b_j - b_j^0\|_{L^\infty} \leq AC(K)\sqrt{T}\lambda_1(w) + AC(K)\sqrt{T}\lambda_1(w)^{1-\sigma}\lambda_3(w)^\sigma \leq AC(K)\sqrt{T}\lambda(w).$$

Of course, the same inequality holds for $\|a_j - a_j^0\|_{L^\infty}$, $j = 1, 2$. Note that we have applied the following classical lemma:

Lemma 4.1. *If $s > \frac{n}{2}$ and $|r| \leq s$, then $H^r(\mathbb{R}^n) \cdot H^s(\mathbb{R}^n) \subset H^r(\mathbb{R}^n)$ with continuous injection.*

Finally, it follows from Theorem 2.10 and Theorem 2.11 that

$$\begin{aligned} \|R(z)\|_{s-2} &= \|R(z)\|_{\frac{n}{2}+\varrho} \leq \|R(z) - R(z_0)\|_{\frac{n}{2}+\varrho} + \|R(z_0)\|_{\frac{n}{2}+\varrho} \\ &\leq C_1(\|z\|_{\frac{n}{2}+\varrho}, \|z_0\|_{\frac{n}{2}+\varrho})\|z - z_0\|_{\frac{n}{2}+\varrho} + C_2(\|z_0\|_{\frac{n}{2}+\varrho})\|z_0\|_{\frac{n}{2}+\varrho} \\ &\leq C_1(\|w\|_{\frac{n}{2}+\varrho+1}, \|u_0\|_{\frac{n}{2}+\varrho+1})\|w - u_0\|_{\frac{n}{2}+\varrho+1} + C_2(\|u_0\|_{\frac{n}{2}+\varrho+1})\|u_0\|_{\frac{n}{2}+\varrho+1} \\ &\leq C(K)\|w - u_0\|_{\frac{n}{2}+\varrho+1} + A\|u_0\|_{\frac{n}{2}+\varrho+1}, \end{aligned}$$

and, using once again Proposition A.5, we obtain

$$\begin{aligned} \sup_{[0,T]} \|w - u_0\|_{s'} &\leq A\|w - u_0\|_{H^\sigma([0,T];H^{s'})} \leq A\|w - u_0\|_{L^2([0,T];H^{s'+1})}^{1-\sigma} \|w - u_0\|_{H^1([0,T];H^{s''})}^\sigma \\ &\leq A\|w - u_0\|_{L^2([0,T];H^{s'+1})} + A\|w - u_0\|_{L^2([0,T];H^{s'+1})}^{1-\sigma} \|\partial_t w\|_{L^2([0,T];H^{s''})}^\sigma \\ &\leq A\|w - u_0\|_{L^2([0,T];H^s)} + A\|w - u_0\|_{L^2([0,T];H^s)}^{1-\sigma} \|\partial_t w\|_{L^2([0,T];H^{s-2})}^\sigma \\ &\leq A\sqrt{T}\lambda_1(w) + A\sqrt{T}\lambda_1(w)^{1-\sigma}\lambda_3(w)^\sigma \leq A\sqrt{T}\lambda(w), \end{aligned}$$

where $s' = \frac{n+\varrho}{2} + 1 < s$, $\sigma > \frac{1}{2}$, $s'' = \frac{n+\varrho}{2} + 2 - \frac{1}{\sigma}$ and $s'' \leq s - 2$ if σ is close to $\frac{1}{2}$. Hence,

$$\sup_{[0,T]} \|R(z)\|_{s-2} \leq A\|u_0\|_{\frac{n+\varrho}{2}+1} + AC(K)\sqrt{T}\lambda(w).$$

Thus, we have bounded all the terms of (31) and the result is that

$$\begin{aligned} \lambda_3(\Upsilon w) &\leq A\lambda_1(\Upsilon w) + AC(K)\sqrt{T}\lambda(w)\lambda_1(w) + A\|u_0\|_{\frac{n+\varrho}{2}+1} + AC(K)\sqrt{T}\lambda(w) \\ &\leq A\|u_0\|_s + \sqrt{AC(K)T\lambda(w)\lambda(\Upsilon w)} + AC(K)\sqrt{T}\lambda(w), \end{aligned} \tag{32}$$

where, of course, we have used (30). Therefore,

$$\begin{aligned} \lambda(\Upsilon w) &\leq A\|u_0\|_s + \sqrt{AC(K)T\lambda(w)\lambda(\Upsilon w)} + AC(K)\sqrt{T}\lambda(w) \\ &\leq A\|u_0\|_s + \frac{1}{2}AC(K)T\lambda(w) + \frac{1}{2}\lambda(\Upsilon w) + AC(K)\sqrt{T}\lambda(w), \end{aligned}$$

which leads to

$$\lambda(\Upsilon w) \leq 2A\|u_0\|_s + AC(K)T\lambda(w) + 2AC(K)\sqrt{T}\lambda(w),$$

that is, an estimate which is, by changing the constants and taking $T \leq 1$, of the form

$$\lambda(\Upsilon w) \leq A\|u_0\|_s + AC(K)\sqrt{T}\lambda(w). \tag{33}$$

This is the main nonlinear estimate. In fact, when $u_0 \neq 0$, by taking $K = 2A\|u_0\|_s$ for example, and then, $T > 0$ such that

$$T \leq \left(\frac{A\|u_0\|_s}{AC(K)K}\right)^2 = \left(\frac{1}{2AC(K)}\right)^2,$$

it follows from (33) that $\lambda(\Upsilon w) \leq K$ when $\lambda(w) \leq K$, that is, $\Upsilon(Z) \subset Z$. When $u_0 = 0$, it suffices to take $K > 0$ and $T \leq 1/A^2C(K)^2$ to obtain the same result.

Let us now show that $\Upsilon : Z \rightarrow Z$ is a contraction mapping. In fact, the arguments are similar to the above ones and we shall be brief. If $w_1, w_2 \in Z$, then $W = \Upsilon w_1 - \Upsilon w_2$ satisfies the Cauchy problem

$$\begin{cases} \partial_t W = i\mathcal{L}W + T_{b_1^0}\nabla_x W + T_{b_2^0}\nabla_x \bar{W} + T_{a_1^0}W + T_{a_2^0}\bar{W} + \tilde{R}(z_1) - \tilde{R}(z_2), \\ W(0) = 0, \end{cases} \tag{34}$$

where, as before, $z_j = (w_j, \nabla_x w_j, \bar{w}_j, \nabla_x \bar{w}_j)$, $j = 1, 2$. Applying Theorem 3.1 to (34) gives

$$\max\{\lambda_1(W)^2, \lambda_2(W)^2\} \leq AI_T(J^s(\tilde{R}(z_1) - \tilde{R}(z_2)), J^s W), \tag{35}$$

and, consequently, we have to estimate uniformly in μ the integral

$$\int_0^T |(G_\mu J^s(\tilde{R}(z_1) - \tilde{R}(z_2)), J^s W)| dt.$$

It follows from (21) that

$$\begin{aligned} \tilde{R}(z_1) - \tilde{R}(z_2) &= T_{b_1(z_1)-b_1^0}\nabla(w_1 - w_2) + T_{b_1(z_1)-b_1(z_2)}\nabla w_2 \\ &\quad + T_{b_2(z_1)-b_2^0}\nabla(\bar{w}_1 - \bar{w}_2) + T_{b_2(z_1)-b_2(z_2)}\nabla \bar{w}_2 \\ &\quad + T_{a_1(z_1)-a_1^0}(w_1 - w_2) + T_{a_1(z_1)-a_1(z_2)}w_2 \\ &\quad + T_{a_2(z_1)-a_2^0}(\bar{w}_1 - \bar{w}_2) + T_{a_2(z_1)-a_2(z_2)}\bar{w}_2 \\ &\quad + R(z_1) - R(z_2), \end{aligned} \tag{36}$$

and we have to estimate the integral corresponding to each term of the above sum. Let us first consider the terms of the third line in (36). By the same argument as that used to obtain (27), we have

$$\int_0^T |(G_\mu J^s(T_{a_1(z_1)-a_1^0}(w_1 - w_2) + T_{a_1(z_1)-a_1(z_2)}w_2), J^s W)| dt \leq ATC(K)\lambda(w_1 - w_2)\lambda(W),$$

where we also applied Proposition 2.9 for the second term. Of course, we have the same estimate for the integral corresponding to the terms of the fourth line in (36).

As for the terms of the first line in (36), applying an argument similar to that yielding (28), one obtains

$$\begin{aligned} \int_0^T & \left| \langle G_\mu J^s (T_{b_1(z_1)-b_1^0} \nabla(w_1 - w_2) + T_{b_1(z_1)-b_1(z_2)} \nabla w_2), J^s W \rangle \right| dt \\ & \leq A, C(K)T(\lambda(w_1)^2 \lambda_2(w_1 - w_2) + \lambda(w_1 - w_2)(\lambda(w_1) + \lambda(w_2)) \lambda_2(w_2)) \lambda_2(W) \\ & \leq AC(K)T(\lambda(w_1)^2 + \lambda(w_1)\lambda(w_2) + \lambda(w_2)^2)\lambda(w_1 - w_2)\lambda(W) \\ & \leq AC(K)K^2T\lambda(w_1 - w_2)\lambda(W), \end{aligned}$$

and the same estimate holds for the terms of the second line in (36).

Last, for the terms of the fifth line in (36), applying Theorem 2.11 and estimating as in (29), we obtain

$$\begin{aligned} \int_0^T & \left| \langle G_\mu J^s (R(z_1) - R(z_2)), J^s W \rangle \right| dt \leq A \int_0^T \|z_1 - z_2\|_{s-1} \|W\|_s dt \leq AC(K) \int_0^T \|w_1 - w_2\|_s \|W\|_s dt \\ & \leq AC(K)T\lambda_1(w_1 - w_2)\lambda_1(W) \\ & \leq AC(K)T\lambda(w_1 - w_2)\lambda(W). \end{aligned}$$

Summing up and going back to (35), we can conclude that

$$\max\{\lambda_1(W)^2, \lambda_2(W)^2\} \leq AC(K)T\lambda(w_1 - w_2)\lambda(W).$$

It remains to estimate $\lambda_3(W)$. Using the fact that W satisfies the Cauchy problem (34) and an argument similar to that yielding (32), we obtain

$$\begin{aligned} \lambda_3(W) & \leq A\lambda_1(W) + AC(K)\sqrt{T}(\lambda(w_1)\lambda_1(w_1 - w_2) + \lambda(w_1 - w_2)\lambda_1(w_2)) + AC(K)\sqrt{T}\lambda(w_1 - w_2) \\ & \leq A\lambda_1(W) + AC(K)\sqrt{T}(\lambda(w_1) + \lambda(w_2))\lambda(w_1 - w_2) + AC(K)\sqrt{T}\lambda(w_1 - w_2) \\ & \leq \sqrt{AC(K)T\lambda(w_1 - w_2)\lambda(W)} + AC(K)\sqrt{T}\lambda(w_1 - w_2). \end{aligned}$$

Summing up, we have obtained

$$\lambda(W) \leq \sqrt{AC(K)T\lambda(w_1 - w_2)\lambda(W)} + AC(K)\sqrt{T}\lambda(w_1 - w_2).$$

Hence,

$$\lambda(W) \leq \frac{1}{2}AC(K)T\lambda(w_1 - w_2) + \frac{1}{2}\lambda(W) + AC(K)\sqrt{T}\lambda(w_1 - w_2);$$

that is,

$$\lambda(W) = \lambda(\Upsilon w_1 - \Upsilon w_2) \leq AC(K)\sqrt{T}\lambda(w_1 - w_2),$$

with modified constants. This clearly implies, if T is taken small enough, that $\Upsilon : Z \rightarrow Z$ is a contraction mapping and, thus, it has a unique fixed point u in Z which is a solution of (18). In fact, this is the solution of (18) in $C([0, T], H^s(\mathbb{R}^n))$ because the above method gives the local uniqueness and we obtain eventually the full uniqueness by applying a classical bootstrap argument. This proves the first part of Theorem 1.1.

The second part of Theorem 1.1 concerns the continuity of the solution operator $u_0 \mapsto u$ and we start its proof by remarking that this operator maps bounded subsets of $H^s(\mathbb{R}^n)$ into bounded subsets of $C([0, T], H^s(\mathbb{R}^n))$. In fact, if B is a bounded subset of $H^s(\mathbb{R}^n)$, as remarked at the beginning of this

section, the constant A and the bounds of the seminorms of the operators G_μ can be taken to depend only on B ; that is, if $u_0 \in B$, the estimates proven above and satisfied by Υ can be rewritten as

$$\lambda(\Upsilon w) \leq A(B)\|u_0\|_s + A(B)C(K)\sqrt{T}\lambda(w), \tag{37}$$

$$\lambda(\Upsilon w_1 - \Upsilon w_2) \leq A(B)C(K)\sqrt{T}\lambda(w_1 - w_2), \tag{38}$$

where $A(B)$ depends only on n, s and B , which implies that the constants K and T can be chosen depending only on B . Hence, for all $u_0 \in B$, the associated solutions u are all defined on the same interval $[0, T]$ and are all in the ball of radius K . As for the continuity, let B be a bounded subset of $H^s(\mathbb{R}^n)$, $u_0, u_0^* \in B$, u, u^* the respective associated solutions and $w = u - u^*$. Then, w satisfies the Cauchy problem

$$\begin{cases} \partial_t w = i\mathcal{L}w + Du - D^*u^* + \tilde{R} - \tilde{R}^* = i\mathcal{L}w + Dw + (D - D^*)u^* + \tilde{R} - \tilde{R}^*, \\ w(x, 0) = u_0(x) - u_0^*(x), \end{cases} \tag{39}$$

where

$$\begin{aligned} Dw &= T_{b_1^0}\nabla w + T_{b_2^0}\nabla\bar{w} + T_{a_1^0}w + T_{a_2^0}\bar{w}, & D^*w &= T_{b_1^{0,*}}\nabla w + T_{b_2^{0,*}}\nabla\bar{w} + T_{a_1^{0,*}}w + T_{a_2^{0,*}}\bar{w}, \\ \tilde{R} &= \tilde{R}(u, \nabla u, \bar{u}, \nabla\bar{u}) & \tilde{R}^* &= \tilde{R}(u^*, \nabla u^*, \bar{u}^*, \nabla\bar{u}^*). \end{aligned}$$

Of course, the b_j^0, a_j^0 correspond to u_0 whereas the $b_j^{0,*}, a_j^{0,*}$ correspond to u_0^* . Applying Theorem 3.1 gives us the inequality

$$\max\{\lambda_1(w)^2, \lambda_2(w)^2\} \leq A(B)\|u_0 - u_0^*\|_s^2 + A(B)I_T(J^s((D - D^*)u^* + \tilde{R} - \tilde{R}^*), J^s w). \tag{40}$$

As it can be seen easily by going back to (21), we can write

$$\begin{aligned} \tilde{R} - \tilde{R}^* &= T_{b_1(u)-b_1^0}\nabla w + T_{b_1(u)-b_1(u^*)}\nabla u^* + T_{b_1^{0,*}-b_1^0}\nabla u^* \\ &\quad + T_{b_2(u)-b_2^0}\nabla\bar{w} + T_{b_2(u)-b_2(u^*)}\nabla\bar{u}^* + T_{b_2^{0,*}-b_2^0}\nabla\bar{u}^* \\ &\quad + T_{a_1(u)-a_1^0}w + T_{a_1(u)-a_1(u^*)}u^* + T_{a_1^{0,*}-a_1^0}u^* \\ &\quad + T_{a_2(u)-a_2^0}\bar{w} + T_{a_2(u)-a_2(u^*)}\bar{u}^* + T_{a_2^{0,*}-a_2^0}\bar{u}^* \\ &\quad + R(u, \nabla u, \bar{u}, \nabla\bar{u}) - R(u^*, \nabla u^*, \bar{u}^*, \nabla\bar{u}^*), \end{aligned} \tag{41}$$

and we also have

$$(D - D^*)u^* = T_{b_1^0-b_1^{0,*}}\nabla u^* + T_{b_2^0-b_2^{0,*}}\nabla\bar{u}^* + T_{a_1^0-a_1^{0,*}}u^* + T_{a_2^0-a_2^{0,*}}\bar{u}^*.$$

Using the same arguments as before to estimate the integrals corresponding to each of the above terms yields

$$\max\{\lambda_1(w)^2, \lambda_2(w)^2\} \leq A(B)\|u_0 - u_0^*\|_s^2 + A_1(B)C_1(K)T(\lambda(w)\|u_0 - u_0^*\|_s + \lambda(w)^2), \tag{42}$$

which becomes, after a change of the constants and assuming $T \leq 1$,

$$\max\{\lambda_1(w), \lambda_2(w)\} \leq A(B)\|u_0 - u_0^*\|_s + A(B)C(K)\sqrt{T}\lambda(w). \tag{43}$$

Next, using (39) and similar arguments, one can easily get

$$\lambda_3(w) \leq A(B)C(K)(\|u_0 - u_0^*\|_s + \lambda_1(w)),$$

which becomes, after use of (43) and a possible change of the constants,

$$\lambda_3(w) \leq A(B)C(K)(\|u_0 - u_0^*\|_s + \sqrt{T}\lambda(w)).$$

Hence,

$$\lambda(w) \leq A(B)C(K)\|u_0 - u_0^*\|_s + A(B)C(K)\sqrt{T}\lambda(w), \tag{44}$$

which, by taking $T \leq (1/2A(B)C(K))^2$ (for example), leads to the Lipschitz estimate

$$\lambda(w) = \lambda(u - u^*) \leq 2A(B)C(K)\|u_0 - u_0^*\|_s, \tag{45}$$

and this achieves the proof of Theorem 1.1.

5. Proof of Proposition 3.2

We shall only give the main steps for the convenience of the reader and refer to [Bienaimé 2014] for the full details.

Let us start by remarking that it is sufficient to treat the case $s = 0$. Indeed, if $v = J^s u$ and $v_0 = J^s u_0$, it is easy to see that u is a solution of (8) if and only if v satisfies

$$\begin{cases} \partial_t v = i\mathcal{L}v + T_{b_1} \cdot \nabla_x v + T_{b_2} \cdot \nabla_x \bar{v} + \tilde{C}_1 v + \tilde{C}_2 \bar{v} + \tilde{f}(x, t), \\ v(x, 0) = v_0 \in L^2(\mathbb{R}^n), \end{cases} \tag{46}$$

where $\tilde{f} = J^s f$ and $\tilde{C}_k = J^s C_k J^{-s} + [J^s, T_{b_k} \cdot \nabla_x] J^{-s}$, $k = 1$ or 2 , and, thanks to the paradifferential calculus, the \tilde{C}_k are bounded operators in $L^2(\mathbb{R}^n)$.

The idea of proof is that of [Kenig et al. 1998], inspired by [Takeuchi 1992], and consists in constructing a pseudodifferential operator C which is bounded and invertible in $L^2(\mathbb{R}^n)$ and estimating $\sup_{[0, T]} \|Cu\|_0$ instead of estimating directly $\sup_{[0, T]} \|u\|_0$. Since $\frac{d}{dt} \langle Cu, Cu \rangle = \langle C \partial_t u, Cu \rangle + \langle Cu, C \partial_t u \rangle$ and u is a solution of (8), we obtain that

$$\begin{aligned} \frac{d}{dt} \|Cu\|_0^2 &= \langle iC \mathcal{L}u, Cu \rangle + \langle CT_{b_1} \nabla u, Cu \rangle + \langle CT_{b_2} \nabla \bar{u}, Cu \rangle \\ &\quad + \langle CC_1 u, Cu \rangle + \langle CC_2 \bar{u}, Cu \rangle + \langle Cf, Cu \rangle \\ &\quad + \langle Cu, iC \mathcal{L}u \rangle + \langle Cu, CT_{b_1} \nabla u \rangle + \langle Cu, CT_{b_2} \nabla \bar{u} \rangle \\ &\quad + \langle Cu, CC_1 u \rangle + \langle Cu, CC_2 \bar{u} \rangle + \langle Cu, Cf \rangle, \end{aligned} \tag{47}$$

and since

$$\langle i\mathcal{L}Cu, Cu \rangle + \langle Cu, i\mathcal{L}Cu \rangle = 0,$$

we have finally

$$\begin{aligned} \frac{d}{dt} \|Cu\|_0^2 &= 2 \operatorname{Re} \langle (i[C, \mathcal{L}] + CT_{b_1} \nabla)u, Cu \rangle + 2 \operatorname{Re} \langle CT_{b_2} \nabla \bar{u}, Cu \rangle \\ &\quad + 2 \operatorname{Re} \langle Cu, Cf \rangle + 2 \operatorname{Re} (\langle CC_1 u, Cu \rangle + \langle CC_2 \bar{u}, Cu \rangle). \end{aligned}$$

The idea of [Kenig et al. 1998] is precisely to choose C so that the operator $i[C, \mathcal{L}] + CT_{b_1} \nabla$ will be small in some sense. Here, we will make a refinement by writing $b_1 = b'_1 + ib''_1$ with real b'_1, b''_1 , and by considering the operator $i[C, \mathcal{L}] + iCT_{b'_1} \nabla$ instead. This has been already used by [Bienaimé 2014] and essentially allows one to construct a *real* operator C , that is, with the property $\overline{Cu} = C\bar{u}$, which will be convenient in certain arguments. Now, clearly,

$$|2 \operatorname{Re}(\langle C C_1 u, C u \rangle + \langle C C_2 \bar{u}, C u \rangle)| \leq 2(A_1 + A_2) \|C\|_{\mathcal{L}(L^2)}^2 \|u\|_0^2,$$

and integrating on $[0, T']$, $T' \leq T$, yields

$$\begin{aligned} \|Cu(T')\|_2^2 &\leq \|Cu_0\|_0^2 + 2 \left| \operatorname{Re} \int_0^{T'} \langle (i[C, \mathcal{L}] + iCT_{b'_1} \nabla)u, Cu \rangle dt \right| + 2 \left| \operatorname{Re} \int_0^{T'} \langle CT_{b'_1} \nabla u, Cu \rangle dt \right| \\ &\quad + 2 \left| \operatorname{Re} \int_0^{T'} \langle CT_{b_2} \nabla \bar{u}, Cu \rangle dt \right| + 2 \left| \operatorname{Re} \int_0^{T'} \langle Cu, Cf \rangle dt \right| \\ &\quad + 2(A_1 + A_2) \|C\|_{\mathcal{L}(L^2)}^2 \int_0^{T'} \|u(t)\|_0^2 dt, \end{aligned} \tag{48}$$

and our task will be to estimate appropriately each of the terms in the right-hand side of this inequality. The most difficult one is

$$\left| \int_0^T \langle (i[C, \mathcal{L}] + iCT_{b'_1} \nabla)u, Cu \rangle dt \right|$$

and C will be constructed so that this term will be small with respect to some parameters to be defined later. To this end, let us denote by c the symbol of C and define

$$p(x, \xi) = -2\xi^\sharp \cdot \nabla_x c(x, \xi) - c(x, \xi) \tilde{b}''_1(x, \xi) \cdot \xi, \tag{49}$$

where $\xi^\sharp = (\xi_1, \dots, \xi_{j_0}, -\xi_{j_0+1}, \dots, -\xi_n)$ and \tilde{b}''_1 is such that $T_{b'_1} = \tilde{b}''_1(x, D)$; see (2). The problem lies essentially in the fact that $p(x, \xi)$ is not the true principal symbol of the pseudodifferential (or paradifferential) operator $i[C, \mathcal{L}] + iCT_{b'_1} \nabla$ since C will be merely in the class $\operatorname{Op}S_{0,0}^0$. Nevertheless, the constructed C will allow us to obtain good estimates.

Set $c(x, \xi) = \exp(\gamma(x, \xi))$ and $\gamma(x, \xi) = \sum_{\mu \in \mathbb{Z}^n} \alpha_{1,\mu} \gamma_\mu(x, \xi)$, where the $\alpha_{1,\mu}$ are the coefficients of b_1 in its decomposition with respect to the $\varphi_{1,\mu}$, see (9), and the $\gamma_\mu(x, \xi)$ are defined a little later. Note here that one can assume the $\alpha_{1,\mu}$ real (and even nonnegative) without loss of generality. We can then write

$$p(x, \xi) = c(x, \xi) \sum_{\mu} \alpha_{1,\mu} (-2\xi^\sharp \cdot \nabla_x \gamma_\mu(x, \xi) - \tilde{\varphi}_{1,\mu}(x, \xi) \cdot \xi),$$

and this suggests considering the function

$$\eta_\mu(x, \xi) = \frac{1}{2} \int_0^\infty \operatorname{Im}(\tilde{\varphi}_{1,\mu})(x + s\xi^\sharp, \xi) \cdot \xi \, ds.$$

One can show that such a function is smooth and satisfies, for all multi-indices α, β ,

$$|\partial_x^\alpha \partial_\xi^\beta \eta_\mu(x, \xi)| \leq A_{\alpha,\beta} \sup_{\beta' \leq \beta} \|\langle \xi \rangle^{|\beta'|} \partial_x^\alpha \partial_\xi^{\beta'} \varphi_{1,\mu}\|_{L^\infty} \langle x - \mu \rangle^{|\beta|} \langle \xi \rangle^{-|\beta|}, \tag{50}$$

and, moreover,

$$-2\xi^\sharp \cdot \nabla_x \eta_\mu(x, \xi) - \text{Im}(\tilde{\varphi}_{1,\mu})(x, \xi) \cdot \xi = 0. \tag{51}$$

See [Kenig et al. 1998; Bienaimé 2014] for the proof. To get an even function, we replace η_μ by

$$\zeta_\mu(x, \xi) = \frac{1}{2}(\eta_\mu(x, \xi) + \eta_\mu(x, -\xi)),$$

which satisfies the same properties as η_μ , and then set

$$\gamma_\mu(x, \xi) = \theta\left(\frac{|\xi|}{R}\right) \psi\left(\frac{R(x - \mu)}{\langle \xi \rangle}\right) \zeta_\mu(x, \xi),$$

where θ and ψ are smooth (real) functions on \mathbb{R} such that $\theta(t) = 1$ if $t \geq 2$, $\theta(t) = 0$ if $t \leq 1$, $\psi(x) = 1$ if $|t| \leq 1$, $\psi = 0$ outside some compact set and R is a large parameter that will be fixed later. One can easily check that $\gamma_\mu \in S_{0,0}^0$ and that its seminorms are uniformly bounded with respect to μ and R . The following lemma gives the main properties of the operator C and its symbol

$$c(x, \xi) = \exp(\gamma(x, \xi)) = \exp\left(\sum_\mu \alpha_{1,\mu} \gamma_\mu(x, \xi)\right).$$

Lemma 5.1. (i) *The symbol $c(x, \xi)$ is real and even in ξ .*

(ii) *The symbol $c(x, \xi)$ is in the class $S_{0,0}^0$. More precisely, for all $\alpha, \beta \in \mathbb{N}^n$,*

$$|\partial_x^\alpha \partial_\xi^\beta c(x, \xi)| \leq \frac{A_{\alpha,\beta}}{R^{|\beta|}} \sup_{1 \leq j \leq |\alpha|+|\beta|} \sup_\mu \sup_{\alpha' \leq \alpha; \beta' \leq \beta} \|\langle \xi \rangle^{|\beta'|} \partial_x^{\alpha'} \partial_\xi^{\beta'} \varphi_{1,\mu}\|_{L^\infty}^j \leq \frac{A_{\alpha,\beta}}{R^{|\beta|}} \sup_\mu v_{|\alpha|+|\beta|}(\varphi_{1,\mu}).$$

(iii) *There exist $N \in \mathbb{N}$ and $A > 0$ such that, for all $R \geq 1$ and all $v \in L^2(\mathbb{R}^n)$,*

$$\begin{aligned} \|Cv\|_0 &\leq A \sup_\mu v_N(\varphi_{1,\mu}) \|v\|_0, \\ \|v\|_0 &\leq A \sup_\mu v_N(\varphi_{1,\mu}) \|Cv\|_s + \frac{A}{R} \sup_\mu v_N(\varphi_{1,\mu})^2 \|v\|_s. \end{aligned}$$

(iv) *The symbol*

$$p(x, \xi) = -2\xi^\sharp \cdot \nabla_x c(x, \xi) - c(x, \xi) \tilde{b}_1''(x, \xi) \cdot \xi$$

is in $S_{0,0}^0$ and its seminorms (of order $\leq M$) are estimated by $AR \sup_\mu v_{M+1}(\varphi_{1,\mu})$.

Even if here the function $\varphi_{1,\mu}$ is more general, the proof follows the same lines as that of [Bienaimé 2014, Lemmas 3.5 and 3.6] and we refer to it. These properties are sufficient to allow us to get the following estimates:

Lemma 5.2. *Let $b(x, \xi)$ be a symbol satisfying*

$$b(x, \xi) = \sum_{\mu \in \mathbb{Z}^n} \alpha_\mu \varphi_\mu(x, \xi), \quad \varphi_\mu \in S_{1,0}^0, \quad \sum_\mu |\alpha_\mu| \leq A_0, \tag{52}$$

$x \mapsto \varphi_\mu(x, \xi)$ is rapidly decreasing in $x - \mu$,

and let \tilde{u} stand for u or \bar{u} . Then, there exist $N \in \mathbb{N}$ and $A > 0$ such that, for all $T > 0$, $T' \in [0, T]$, $R \geq 1$ and every $H = h(x, D)$ in $\text{Op } S_{0,0}^0$, the following estimates hold true:

- (i) $\int_0^{T'} | \langle (CT_b \nabla - (c\tilde{b})(x, D)\nabla)\tilde{u}, Hu \rangle | dt \leq \frac{A}{R} \|h\|_{C^N} \sup_{\mu} \nu_N(\varphi_{1,\mu}) \sup_{\mu} \|\varphi_{\mu}\|_{C^N} \|J^{\frac{1}{2}}u\|_T^2.$
- (ii) $\int_0^{T'} | \langle (i[C, \mathcal{L}] + iCT_{b'_1} \nabla)u, Hu \rangle | dt \leq A \|h\|_{C^N} \sup_{\mu} \nu_N(\varphi_{1,\mu}) \left(RT \sup_{[0,T]} \|u\|_0^2 + \frac{1}{R} \|J^{\frac{1}{2}}u\|_T^2 \right).$
- (iii) $\int_0^{T'} | \langle [C, J^s T_b J^{-s} \nabla]\tilde{u}, Hu \rangle | dt \leq A \|h\|_{C^N} \sup_{\mu} \nu_N(\varphi_{1,\mu}) \sup_{\mu} \|\varphi_{\mu}\|_{C^N} \left(T \sup_{[0,T]} \|u\|_0^2 + \frac{\|J^{\frac{1}{2}}u\|_T^2}{R} \right).$

Remark. The case $s \neq 0$ in (iii) is needed in the Appendix.

Proof. Using the pseudodifferential calculus, we can write the symbol $e(x, \xi)$ of the operator $E = CT_b \nabla - (c\tilde{b})(x, D)\nabla$ as $e = \sum_{\mu} \alpha_{\mu} e_{\mu}$, where e_{μ} is given by

$$e_{\mu}(x, \xi) = \frac{1}{(2\pi)^n} \sum_{j=1}^n \int_0^1 \int e^{-iy\eta} \partial_{\xi_j} c(x, \xi + t\eta) \partial_{x_j} \tilde{\varphi}_{\mu}(x + y, \xi) \cdot \xi dy d\eta dt, \tag{53}$$

and we first remark that $e_{\mu} \in \text{Op } S_{0,0}^1$ and that using the fast decrease of $\tilde{\varphi}_{\mu}(x, \xi)$ in $x - \mu$ and integrations by parts yields the fact that $e_{\mu}(x, \xi)$ is itself rapidly decreasing in $x - \mu$. Next, setting $E_{\mu} = e_{\mu}(x, D)$, we can write

$$\begin{aligned} \langle E\tilde{u}, Hu \rangle &= \sum_{\mu} \alpha_{\mu} \langle E_{\mu}\tilde{u}, Hu \rangle = \sum_{\mu} \alpha_{\mu} \langle H^* E_{\mu}\tilde{u}, u \rangle \\ &= \sum_{\mu} \alpha_{\mu} \langle \langle x - \mu \rangle^{\sigma_0} \tilde{H} \langle x - \mu \rangle^{-\sigma_0} \langle x - \mu \rangle^{\sigma_0} \tilde{E}_{\mu} \langle x - \mu \rangle^{\sigma_0} \tilde{u}, u_{\mu} \rangle, \end{aligned}$$

where $\tilde{H} = J^{-\frac{1}{2}} H^* J^{\frac{1}{2}}$, $\tilde{E}_{\mu} = J^{-\frac{1}{2}} E_{\mu} J^{-\frac{1}{2}}$ and $u_{\mu} = \langle x - \mu \rangle^{-\sigma_0} J^{\frac{1}{2}} u$. Now, it follows from the pseudodifferential calculus (Theorem 2.1) that \tilde{H} and \tilde{E}_{μ} are in $\text{Op } S_{0,0}^0$ and that we can estimate the seminorms of \tilde{H} and \tilde{E}_{μ} by those of H and E_{μ} respectively. Moreover, it is easy to see that the symbol of \tilde{E}_{μ} inherits the fast decrease in $x - \mu$ which implies, by virtue of Lemma 2.3, that the operator $\langle x - \mu \rangle^{\sigma_0} \tilde{E}_{\mu} \langle x - \mu \rangle^{\sigma_0}$ is also in $\text{Op } S_{0,0}^0$ and that its seminorms are estimated by those of E_{μ} uniformly in μ . The same property holds for the operator $\langle x - \mu \rangle^{\sigma_0} \tilde{H} \langle x - \mu \rangle^{-\sigma_0}$, as it follows also from Lemma 2.3. This allows us to apply the Calderón–Vaillancourt theorem to obtain

$$\begin{aligned} \int_0^{T'} | \langle E\tilde{u}, Hu \rangle | dt &\leq \sum_{\mu} |\alpha_{\mu}| \int_0^{T'} \| \langle x - \mu \rangle^{\sigma_0} \tilde{H} \langle x - \mu \rangle^{-\sigma_0} \|_{\mathcal{L}(L^2)} \| \langle x - \mu \rangle^{\sigma_0} \tilde{E}_{\mu} \langle x - \mu \rangle^{\sigma_0} \|_{\mathcal{L}(L^2)} \| u_{\mu} \|_0^2 dt \\ &\leq A \|h\|_{C^{N_1}} \sup_{\mu} \sum_{|\alpha|+|\beta| \leq N_1} \| \langle x - \mu \rangle^{2\sigma_0} \partial_x^{\alpha} \partial_{\xi}^{\beta} e_{\mu} \|_{L^{\infty}} \| J^{\frac{1}{2}} u \|_T^2 \\ &\leq \frac{A}{R} \|h\|_{C^{N_1}} \sup_{\mu} \nu_{N_2}(\varphi_{1,\mu}) \sup_{\mu} \|\varphi_{\mu}\|_{C^{N_2}} \| J^{\frac{1}{2}} u \|_T^2, \tag{54} \end{aligned}$$

which proves (i).

To prove (ii), note first that the symbol of $i[\mathbf{C}, \mathcal{L}]$ is given by

$$-2\xi^\# \cdot \nabla_x \mathbf{c}(x, \xi) + (\mathcal{L}_x \mathbf{c})(x, \xi)$$

and that of $i\mathbf{C}T_{b_1'}\nabla$ can be written as

$$i\mathbf{c}(x, \xi)\tilde{b}_1''(x, \xi).i\xi + \frac{1}{(2\pi)^n} \sum_{j=1}^n \int_0^1 \int e^{-iy\eta} \partial_{\xi_j} \mathbf{c}(x, \xi + t\eta) \partial_{x_j} \tilde{b}_1''(x + y, \xi).i\xi \, dy \, d\eta \, dt.$$

Thus, the symbol of the operator $i[\mathbf{C}, \mathcal{L}] + i\mathbf{C}T_{b_1'}\nabla$ is given by

$$(\mathcal{L}_x \mathbf{c})(x, \xi) + p(x, \xi) + ie(x, \xi),$$

where $p(x, \xi)$ is given by (49), $e = \sum_{\mu} \alpha_{\mu} e_{\mu}$ and $e_{\mu}(x, \xi)$ is given by (53) with $\alpha_{\mu} = \alpha_{1,\mu}$ and $\varphi_{\mu} = \text{Im}(\varphi_{1,\mu})$. Hence, applying Lemma 5.1 and the Calderón–Vaillancourt theorem yields the estimate

$$\int_0^{T'} |((\mathcal{L}_x \mathbf{c})(x, D) + p(x, D))u, Hu| \, dt \leq ART \|h\|_{C^{N_1}} \sup_{\mu} \nu_{N_1}(\varphi_{1,\mu})^2 \sup_{[0,T]} \|u\|_0^2,$$

and applying part (i) gives the estimate

$$\int_0^{T'} |(ie(x, D)u, Hu)| \, dt \leq \frac{A}{R} \|h\|_{C^{N_2}} \sup_{\mu} \nu_{N_2}(\varphi_{1,\mu})^2 \|J^{\frac{1}{2}}u\|_T^2,$$

which proves (ii).

To prove (iii), we first treat the case $s = 0$ and note that the symbol of $[\mathbf{C}, T_b \nabla] = \mathbf{C}T_b \nabla - T_b \nabla \mathbf{C}$ can be written simply as $e(x, \xi) - e_0(x, \xi)$, where $e(x, \xi)$ is the symbol of the operator E studied in (i) and

$$e_0(x, \xi) = \frac{1}{(2\pi)^n} \sum_{j=1}^n \int_0^1 \int e^{-iy\eta} \partial_{\xi_j} (\tilde{b}(x, \xi + t\eta).(\xi + t\eta)) \partial_{x_j} \mathbf{c}(x + y, \xi) \, dy \, d\eta \, dt.$$

Since $\partial_{\xi_j} (\tilde{b}(x, \xi).\xi)$ is of order 0, the symbol $e_0(x, \xi)$ is in fact in $S_{0,0}^0$ and the seminorms of e_0 are estimated by a product of seminorms of \tilde{b} and \mathbf{c} . Hence, by using the decomposition of b as above, we get

$$\int_0^{T'} |\langle e_0(x, D)\tilde{u}, Hu \rangle| \, dt \leq AT \|h\|_{C^{N_1}} \sup_{\mu} \|\varphi_{\mu}\|_{C^{N_2}} \sup_{\mu} \nu_{N_2}(\varphi_{1,\mu}) \sup_{[0,T]} \|u\|_0^2,$$

which, together with (54), yields (iii) in the case $s = 0$. If $s \neq 0$, it follows from the pseudodifferential and paradifferential calculi that $J^s T_b J^{-s} = T_{b^\#}$, where $b^\# = \sum_{\mu} \alpha_{\mu} \psi_{\mu}$ and ψ_{μ} is given by

$$\psi_{\mu}(x, \xi) = \frac{1}{(2\pi)^n} \int e^{-iy\eta} \langle \xi + \eta \rangle^s \varphi_{\mu}(x + y, \xi) \langle \xi \rangle^{-s} \, dy \, d\eta,$$

which implies that ψ_{μ} is also rapidly decreasing in $x - \mu$ and that it is in $S_{1,0}^0$ with seminorms estimated by those of φ_{μ} . This shows that the case $s \neq 0$ follows from the case $s = 0$ and achieves the proof of Lemma 5.2. □

Lemma 5.3. *Let b be as in the preceding lemma. Then, there exist $N \in \mathbb{N}$ and $A > 0$ such that, for all $T > 0$, $T' \in [0, T]$ and $R \geq 1$, the following estimates hold true:*

(i) *If $b(x, \xi)$ is even in ξ , then*

$$\int_0^{T'} |\langle \mathbf{C} T_b \nabla \bar{u}, \mathbf{C} u \rangle| dt \leq A \sup_{\mu} v_N(\varphi_{1,\mu}) \sup_{\mu} \|\varphi_{\mu}\|_{C^N} \left(T \sup_{0 \leq t \leq T} \|u\|_0^2 + \frac{1}{R} \|J^{\frac{1}{2}} u\|_T \right).$$

(ii) *If b is real, then*

$$\left| \operatorname{Re} \int_0^{T'} \langle \mathbf{C} T_b \nabla u, \mathbf{C} u \rangle dt \right| \leq A \sup_{\mu} v_N(\varphi_{1,\mu}) \sup_{\mu} \|\varphi_{\mu}\|_{C^N} \left(T \sup_{0 \leq t \leq T} \|u\|_0^2 + \frac{1}{R} \|J^{\frac{1}{2}} u\|_T \right).$$

Proof. Since \mathbf{C} is real, we can write

$$\langle \mathbf{C} T_b \nabla \bar{u}, \mathbf{C} u \rangle = \langle T_b \nabla \mathbf{C} \bar{u}, \mathbf{C} u \rangle + \langle [\mathbf{C}, T_b \nabla] \bar{u}, \mathbf{C} u \rangle = \langle T_b \nabla \overline{\mathbf{C} u}, \mathbf{C} u \rangle + \langle [\mathbf{C}, T_b \nabla] \bar{u}, \mathbf{C} u \rangle.$$

Now, the integral corresponding to $\langle [\mathbf{C}, T_b \nabla] \bar{u}, \mathbf{C} u \rangle$ is treated by Lemma 5.2(iii). As for the other term, we note that it is of the form $\langle T_b \nabla \bar{v}, v \rangle$, so it suffices to study such a term. Since $b(x, \xi)$ is even in ξ , we have

$$\langle T_b \nabla \bar{v}, v \rangle = \overline{\langle v, T_b \nabla \bar{v} \rangle} = \langle \bar{v}, T_{\bar{b}} \nabla v \rangle = \langle (T_{\bar{b}} \nabla)^* \bar{v}, v \rangle,$$

and it follows from the pseudodifferential (or paradifferential) calculus that

$$(T_{\bar{b}} \nabla)^* = -T_b \nabla + E_1, \tag{55}$$

where E_1 is of type $S_{1,0}^0$ and its seminorms (up to some finite order) are estimated by those of b . Hence,

$$\langle T_b \nabla \bar{v}, v \rangle = -\langle T_b \nabla \bar{v}, v \rangle + \langle E_1 \bar{v}, v \rangle,$$

and $\langle T_b \nabla \bar{v}, v \rangle = \frac{1}{2} \langle E_1 \bar{v}, v \rangle$, that is, $\langle T_b \nabla \overline{\mathbf{C} u}, \mathbf{C} u \rangle = \frac{1}{2} \langle E_1 \overline{\mathbf{C} u}, \mathbf{C} u \rangle$, and (i) follows just by applying the Calderón–Vaillancourt theorem and Lemma 5.1.

To prove (ii), we write as before

$$\langle \mathbf{C} T_b \nabla u, \mathbf{C} u \rangle = \langle T_b \nabla \mathbf{C} u, \mathbf{C} u \rangle + \langle [\mathbf{C}, T_b \nabla] u, \mathbf{C} u \rangle,$$

and then apply Lemma 5.2(iii) to reduce the problem to the study of $\operatorname{Re} \langle T_b \nabla \mathbf{C} u, \mathbf{C} u \rangle$. Now, it follows from (55) and the fact that b is real that we have

$$2 \operatorname{Re} \langle T_b \nabla \mathbf{C} u, \mathbf{C} u \rangle = \langle T_b \nabla \mathbf{C} u, \mathbf{C} u \rangle + \langle \mathbf{C} u, T_b \nabla \mathbf{C} u \rangle = \langle (T_b \nabla + (T_b \nabla)^*) \mathbf{C} u, \mathbf{C} u \rangle = \langle E_1 \mathbf{C} u, \mathbf{C} u \rangle$$

and the proof ends like that of (i). The lemma is thus proved. □

It is clear now that applying Lemmas 5.1, 5.2 and 5.3 to the inequality (48) yields Proposition 3.2.

6. Proof of Proposition 3.3

By the same argument as that used in the beginning of the proof of Proposition 3.2, it is sufficient to establish the first estimate in the case $s = 0$.

The proof follows the same ideas as that of [Kenig et al. 1998; Bienaimé 2014]. The difference is that here the T_{b_k} , $k = 1, 2$, are general paradifferential operators of order 0 instead of merely multiplication or paramultiplication operators.

Since

$$\begin{aligned} \partial_t u &= i\mathcal{L}u + T_{b_1} \cdot \nabla u + T_{b_2} \cdot \nabla \bar{u} + C_1 u + C_2 \bar{u} + f, \\ \partial_t \bar{u} &= -i\mathcal{L}\bar{u} + T_{\bar{b}_1} \cdot \nabla \bar{u} + T_{\bar{b}_2} \cdot \nabla u + \bar{C}_1 \bar{u} + \bar{C}_2 u + \bar{f}, \end{aligned}$$

where the operators \bar{C}_k are defined by $\bar{C}_k u = \overline{C_k \bar{u}}$, one starts by remarking that the vector unknown $w = \begin{pmatrix} u \\ \bar{u} \end{pmatrix}$ satisfies the system

$$\partial_t w = iHw + Bw + Cw + F, \tag{56}$$

where

$$H = \begin{pmatrix} \mathcal{L} & 0 \\ 0 & -\mathcal{L} \end{pmatrix}, \quad B = \begin{pmatrix} T_{b_1} \nabla & T_{b_2} \nabla \\ T_{\bar{b}_2} \nabla & T_{\bar{b}_1} \nabla \end{pmatrix}, \quad C = \begin{pmatrix} C_1 & C_2 \\ \bar{C}_1 & \bar{C}_2 \end{pmatrix}, \quad F = \begin{pmatrix} f \\ \bar{f} \end{pmatrix},$$

and the idea then is to estimate the expression $\langle \Psi w, w \rangle$ by means of Gårding’s inequality for systems via Doi’s argument. Here,

$$\Psi = \begin{pmatrix} \Psi_0 & 0 \\ 0 & -\Psi_0 \end{pmatrix},$$

and Ψ_0 is an appropriate pseudodifferential operator in $\text{OpS}_{1,0}^0$ to be chosen a little later. By using (56), one gets easily

$$\begin{aligned} \partial_t \langle \Psi w, w \rangle &= \langle \Psi \partial_t w, w \rangle + \langle \Psi w, \partial_t w \rangle \\ &= \langle (i[\Psi, H] + B^* \Psi + \Psi B + C^* \Psi + \Psi C)w, w \rangle + \langle \Psi F, w \rangle + \langle \Psi w, F \rangle, \end{aligned} \tag{57}$$

and, as one can check also easily, the principal symbol of the first-order operator

$$i[\Psi, H] + B^* \Psi + \Psi B + C^* \Psi + \Psi C$$

is given by

$$M(x, \xi) = \begin{pmatrix} 2\xi^\sharp \cdot \nabla_x \psi_0(x, \xi) - 2\xi \cdot \text{Im}(\tilde{b}_1)(x, \xi) \psi_0(x, \xi) & 2i\xi \cdot \tilde{b}_2(x, \xi) \psi_0(x, \xi) \\ -2i\xi \cdot \tilde{b}_2(x, \xi) \psi_0(x, \xi) & 2\xi^\sharp \cdot \nabla_x \psi_0(x, \xi) - 2\xi \cdot \text{Im}(\tilde{b}_1)(x, \xi) \psi_0(x, \xi) \end{pmatrix},$$

where ψ_0 denotes the symbol of Ψ_0 . Now, for ψ_0 , we shall make the following choice which follows the idea of [Doi 1994]. Define

$$\begin{aligned} p(x, \xi) &= \langle \xi \rangle^{-1} \sum_{j=1}^n \xi_j^\sharp h(x_j) \quad \text{with } h(t) = \int_0^t \langle s \rangle^{-2\sigma_0} ds, \\ p_\mu(x, \xi) &= p(x - \mu, \xi) + A_0 \sum_{\mu' \in \mathbb{Z}^n} (|\alpha_{1, \mu'}| + |\alpha_{2, \mu'}|) p(x - \mu', \xi), \\ \psi_0(x, \xi) &= \psi_\mu(x, \xi) = \exp(-p_\mu(x, \xi)). \end{aligned}$$

Here, the $\alpha_{1, \mu'}$ and $\alpha_{2, \mu'}$ are the coefficients of b_1 and b_2 in their decompositions with respect to the $\varphi_{1, \mu'}$ and $\varphi_{2, \mu'}$ respectively, see (9), A_0 is a large constant that will be determined later and $\mu \in \mathbb{Z}^n$

is fixed for the moment. However, from now on, we shall write Ψ_μ and ψ_μ instead of Ψ_0 and ψ_0 to emphasize the dependance on μ . First, note that p_μ and ψ_μ are in $S_{1,0}^0$ and that their seminorms are uniformly bounded with respect to μ . Next, with these notations, the symbol $M(x, \xi)$ can be rewritten as

$$M(x, \xi) = 2\psi_\mu(x, \xi) \begin{pmatrix} -\xi^\# \cdot \nabla_x p_\mu(x, \xi) - \xi \cdot \text{Im}(\tilde{b}_1)(x, \xi) & i\xi \cdot \tilde{b}_2(x, \xi) \\ -i\xi \cdot \tilde{b}_2(x, \xi) & -\xi^\# \cdot \nabla_x p_\mu(x, \xi) - \xi \cdot \text{Im}(\tilde{b}_1)(x, \xi) \end{pmatrix}.$$

Consider now the matrix $Z(x, \xi) = -M(x, \xi) - V(x, \xi)$, where

$$V(x, \xi) = \frac{2\psi_\mu(x, \xi)|\xi|^2}{\langle \xi \rangle \langle x - \mu \rangle^{2\sigma_0}} \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}.$$

$Z(x, \xi)$ is a matrix of symbols in $S_{1,0}^1$ and, in order to apply Gårding's inequality, we are going to show that, for large ξ , it is a nonnegative matrix, that is, $\langle Z(x, \xi)v, v \rangle \geq 0$ for all $v \in \mathbb{C}^2$. In fact, $Z(x, \xi)$ is of the form

$$2\psi_\mu(x, \xi) \begin{pmatrix} \alpha & \beta \\ \bar{\beta} & \alpha \end{pmatrix},$$

where

$$\alpha = \xi^\# \cdot \nabla_x p_\mu(x, \xi) - \frac{|\xi|^2}{\langle \xi \rangle \langle x - \mu \rangle^{2\sigma_0}} + \xi \cdot \text{Im}(\tilde{b}_1)(x, \xi) \quad \text{and} \quad \beta = -i\xi \cdot \tilde{b}_2(x, \xi),$$

and it is sufficient to show that the two eigenvalues $\alpha \pm |\beta|$ of $\begin{pmatrix} \alpha & \beta \\ \bar{\beta} & \alpha \end{pmatrix}$ are nonnegative, or, equivalently, that $\alpha \geq |\beta|$, that is,

$$\xi^\# \cdot \nabla_x p_\mu(x, \xi) - \frac{|\xi|^2}{\langle \xi \rangle \langle x - \mu \rangle^{2\sigma_0}} + \xi \cdot \text{Im}(\tilde{b}_1)(x, \xi) \geq |-i\xi \cdot \tilde{b}_2(x, \xi)|. \tag{58}$$

Now, the main reason for the choice of the symbol p_μ is that it allows to get the following inequality:

$$\begin{aligned} \xi^\# \cdot \nabla_x p_\mu(x, \xi) &= \xi^\# \cdot \nabla_x p(x - \mu, \xi) + A_0 \sum_{\mu' \in \mathbb{Z}^n} (|\alpha_{1,\mu'}| + |\alpha_{2,\mu'}|) \xi^\# \cdot \nabla_x p(x - \mu', \xi) \\ &= \sum_{j=1}^n \frac{\xi_j^2}{\langle \xi \rangle \langle x_j - \mu_j \rangle^{2\sigma_0}} + A_0 \sum_{\mu' \in \mathbb{Z}^n} (|\alpha_{1,\mu'}| + |\alpha_{2,\mu'}|) \sum_{j=1}^n \frac{\xi_j^2}{\langle \xi \rangle \langle x_j - \mu'_j \rangle^{2\sigma_0}} \\ &\geq \frac{|\xi|^2}{\langle \xi \rangle \langle x - \mu \rangle^{2\sigma_0}} + A_0 \sum_{\mu' \in \mathbb{Z}^n} (|\alpha_{1,\mu'}| + |\alpha_{2,\mu'}|) \frac{|\xi|^2}{\langle \xi \rangle \langle x - \mu' \rangle^{2\sigma_0}}; \end{aligned} \tag{59}$$

that is,

$$\xi^\# \cdot \nabla_x p_\mu(x, \xi) - \frac{|\xi|^2}{\langle \xi \rangle \langle x - \mu \rangle^{2\sigma_0}} \geq A_0 \sum_{\mu' \in \mathbb{Z}^n} (|\alpha_{1,\mu'}| + |\alpha_{2,\mu'}|) \frac{|\xi|^2}{\langle \xi \rangle \langle x - \mu' \rangle^{2\sigma_0}}. \tag{60}$$

Besides, we have

$$\tilde{b}_k(x, \xi) = \sum_{\mu' \in \mathbb{Z}^n} \alpha_{k,\mu'} \tilde{\varphi}_{k,\mu'}(x, \xi), \quad k = 1, 2,$$

and it follows from Lemma 2.8 that

$$\langle x - \mu' \rangle^{2\sigma_0} |\tilde{\varphi}_{k,\mu'}(x, \xi)| \leq A(n),$$

with a constant $A(n)$ which depends only on the dimension. Hence,

$$|i\xi \tilde{b}_k(x, \xi)| \leq A(n) \sum_{\mu' \in \mathbb{Z}^n} |\alpha_{k,\mu'}| \frac{|\xi|}{\langle x - \mu' \rangle^{2\sigma_0}} \leq \sqrt{2}A(n) \sum_{\mu' \in \mathbb{Z}^n} |\alpha_{k,\mu'}| \frac{|\xi|^2}{\langle \xi \rangle \langle x - \mu' \rangle^{2\sigma_0}}, \quad k = 1, 2,$$

if $|\xi| \geq 1$, which, together with (60), implies (58) by taking $A_0 \geq \sqrt{2}A(n)$. Thus, the matrix symbol $Z(x, \xi)$ is nonnegative, and since it is also hermitian, $Z(x, \xi) + Z(x, \xi)^*$ is also nonnegative and we can apply Gårding’s inequality for systems:

$$\operatorname{Re}\langle Z(x, D)w, w \rangle \geq -A\left(1 + \sup_{|\alpha|+|\beta|\leq N} \sup_{k,\mu'} \|\langle \xi \rangle^{|\beta|} \partial_x^\alpha \partial_\xi^\beta \varphi_{k,\mu'}\|_{L^\infty}\right) \|w\|_0^2, \tag{61}$$

where the constant A depends only on A_1, A_2 and the dimension n and the integer N depends only on the dimension n . Now, going back to (57), we can rewrite it as

$$\partial_t \langle \Psi w, w \rangle = \langle (-Z(x, D) - V(x, D) + E)w, w \rangle + \langle \Psi F, w \rangle + \langle \Psi w, F \rangle,$$

where E is a bounded operator in $L^2(\mathbb{R}^n)$, and integrating it on $[0, T]$ yields

$$\begin{aligned} \int_0^T \langle V(x, D)w, w \rangle dt &= \langle \Psi w(0), w(0) \rangle - \langle \Psi w(T), w(T) \rangle \\ &\quad - \int_0^T \langle Z(x, D)w, w \rangle dt + \int_0^T \langle Ew, w \rangle dt + \int_0^T \langle \Psi F, w \rangle dt + \int_0^T \langle \Psi w, F \rangle dt. \end{aligned}$$

Taking the real part, using (61) and estimating, we obtain

$$\begin{aligned} \operatorname{Re} \int_0^T \langle V(x, D)w, w \rangle dt &\leq A \sup_{[0,T]} \|w\|_0^2 + AT\left(1 + \sup_{k,\mu'} \nu_N(\varphi_{k,\mu'})\right) \sup_{[0,T]} \|w\|_0^2 \\ &\quad + \left| \int_0^T \langle \Psi F, w \rangle dt \right| + \left| \int_0^T \langle \Psi w, F \rangle dt \right|, \end{aligned}$$

and since $\psi_\mu(x, \xi) \geq \exp(-A)$ and, for $|\xi| \geq 1$,

$$V(x, \xi) \geq e^{-A} \frac{\langle \xi \rangle}{\langle x - \mu \rangle^{2\sigma_0}} \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix},$$

a second application of Gårding’s inequality gives us

$$\begin{aligned} \operatorname{Re} \int_0^T \langle J^{\frac{1}{2}} \langle x - \mu \rangle^{-2\sigma_0} J^{\frac{1}{2}} w, w \rangle dt &\leq A \sup_{[0,T]} \|w\|_0^2 (1 + T + T \sup_{k,\mu'} \nu_N(\varphi_{k,\mu'})) \\ &\quad + \left| \int_0^T \langle \Psi F, w \rangle dt \right| + \left| \int_0^T \langle \Psi w, F \rangle dt \right|, \end{aligned}$$

with a modified constant A . Since we can write

$$\langle \Psi F, w \rangle = \langle \Psi_\mu f, u \rangle - \overline{\langle \Psi_\mu f, u \rangle}$$

and a similar expression for $\langle \Psi w, F \rangle$, by going back to u , we get eventually

$$\int_0^T \|\langle x-\mu \rangle^{-\sigma_0} J^{\frac{1}{2}} u\|_0^2 dt \leq A \sup_{[0,T]} \|u\|_0^2 (1+T+T \sup_{k,\mu'} \nu_N(\varphi_{k,\mu'})) + \int_0^T |\langle \Psi_\mu f, u \rangle| dt + \int_0^T |\langle \bar{\Psi}_\mu f, u \rangle| dt + \int_0^T |\langle \Psi_\mu^* f, u \rangle| dt + \int_0^T |\langle \bar{\Psi}_\mu^* f, u \rangle| dt,$$

which yields the first part of Proposition 3.3 by taking the supremum over all $\mu \in \mathbb{Z}^n$.

As for the second estimate of Proposition 3.3, we first remark that, since C is real, Cu satisfies

$$\partial_t Cu = i\mathcal{L}Cu + T_{b_1'} \cdot \nabla Cu + T_{b_2} \cdot \nabla \bar{C}u + C_1 Cu + C_2 \bar{C}u + \tilde{f},$$

where $k = 1, 2$, $b_1 = b_1' + ib_1''$ with real b_1', b_1'' , and

$$\tilde{f} = (i[C, \mathcal{L}] + CT_{ib_1''} \nabla)u + [C, T_{b_1'} \cdot \nabla]u + [C, T_{b_2} \cdot \nabla]\bar{u} + [C, C_1]u + [C, C_2]\bar{u} + Cf.$$

Hence, we can apply the first estimate of Proposition 3.3 to Cu obtaining

$$\| \| J^{s+\frac{1}{2}} Cu \| \|_T^2 \leq A(1+T+T \sup_{k,\mu} \nu_N(\varphi_{k,\mu})) \sup_{[0,T]} \|Cu\|_s^2 + \sum_{j=1}^4 \sup_{\mu} \int_0^T |\langle \Psi_{j,\mu} J^s \tilde{f}, J^s Cu \rangle| dt, \tag{62}$$

where $\Psi_{j,\mu} = \psi_j(x - \mu, D)$. Thus, we are led to estimate essentially the terms

$$\int_0^T \left| \langle J^s (i[C, \mathcal{L}] + CT_{ib_1''} \nabla)u, \Psi_{j,\mu}^* J^s Cu \rangle \right| dt + \int_0^T \left| \langle J^s [C, T_{b_1'} \cdot \nabla]u, \Psi_{j,\mu}^* J^s Cu \rangle \right| dt + \int_0^T \left| \langle J^s [C, T_{b_2} \cdot \nabla]\bar{u}, \Psi_{j,\mu}^* J^s Cu \rangle \right| dt.$$

Indeed, since the operators $\Psi_{j,\mu} J^s [C, C_1] J^{-s}$ and $\Psi_{j,\mu} J^s [C, C_2] J^{-s}$ are bounded in L^2 (and so is $J^s C J^{-s}$), the corresponding terms are easily estimated by

$$AT \sup_{\mu} \nu_N(\varphi_{1,\mu}) \sup_{0 \leq t \leq T} \|u(t)\|_s^2.$$

We need now for the other terms the following simple lemma:

Lemma 6.1. *If $a \in S_{0,0}^m$, then, for any real s ,*

$$J^s a(x, D) J^{-s} = a(x, D) + e(x, D),$$

where $e \in S_{0,0}^{m-1}$ and the seminorms of e are bounded by those of a .

Proof. It suffices to apply the pseudodifferential calculus and to remark that

$$e(x, \xi) = \frac{1}{(2\pi)^n} \sum_{j=1}^n \int_0^1 \int e^{-iy\eta} \partial_{\xi_j} (\langle \xi + t\eta \rangle^s) \partial_{x_j} a(x+y, \xi) \langle \xi \rangle^{-s} dy d\eta dt. \quad \square$$

We apply the lemma successively with

$$\begin{aligned} a(x, D) &= i[C, \mathcal{L}] + CT_{ib_1'}\nabla, \\ a(x, D) &= [C, T_{b_1'}\cdot\nabla], \\ a(x, D) &= [C, T_{b_2}\cdot\nabla]. \end{aligned}$$

Since here $m = 1$, we obtain that at each time the operator $e(x, D)$ is bounded in L^2 and that its operator norm is estimated by the seminorms of a . Next, it follows from the pseudodifferential calculus that $\Psi_{j,\mu}^* \in \text{Op}S_{1,0}^0$ and their seminorms are uniformly bounded with respect to μ , and, consequently, also that $\Psi_{j,\mu}^* J^s C J^{-s} \in \text{Op}S_{0,0}^0$ and their seminorms are uniformly estimated by those of C . Hence, the integrals corresponding to the operators $e(x, D)$ are easily estimated by

$$ART \sup_{k,\mu} v_N(\varphi_{k,\mu}) \sup_{0 \leq t \leq T} \|u(t)\|_S^2.$$

Thus, it remains to estimate the sum

$$\begin{aligned} &\int_0^T \left| \langle (i[C, \mathcal{L}] + CT_{ib_1'}\nabla) J^s u, \Psi_{j,\mu}^* J^s C u \rangle \right| dt \\ &\quad + \int_0^T \left| \langle [C, T_{b_1'}\cdot\nabla] J^s u, \Psi_{j,\mu}^* J^s C u \rangle \right| dt + \int_0^T \left| \langle [C, T_{b_2}\cdot\nabla] J^s \bar{u}, \Psi_{j,\mu}^* J^s C u \rangle \right| dt, \end{aligned}$$

to which we apply Lemma 5.2 with $S = \Psi_{j,\mu}^* J^s C J^{-s}$. We obtain eventually

$$\begin{aligned} &\sum_{j=1}^4 \sup_{\mu} \int_0^T \left| \langle \Psi_{j,\mu} J^s \tilde{f}, J^s C u \rangle \right| dt \\ &\quad \leq \sum_{j=1}^4 \sup_{\mu} \int_0^T \left| \langle \Psi_{j,\mu} J^s C f, J^s C u \rangle \right| dt + A \sup_{k,\mu} v_N(\varphi_{k,\mu}) \left(RT \sup_{[0,T]} \|u\|_S^2 + \frac{1}{R} \|J^{s+\frac{1}{2}} u\|_T^2 \right), \end{aligned}$$

which, together with (62), implies the second estimate of Proposition 3.3.

Appendix

Proof of Lemma 3.4. We need the following general estimate:

Lemma A.1. *Let b satisfy*

$$\begin{aligned} b(x, \xi) &= \sum_{\mu \in \mathbb{Z}^n} \alpha_{\mu} \varphi_{\mu}(x, \xi), \quad \sum_{\mu} |\alpha_{\mu}| \leq A_0, \\ \text{supp}(x \mapsto \varphi_{\mu}(x, \xi)) &\subseteq Q_{\mu}^*, \quad \sup_{\mu} \sup_{|\beta| \leq N_0} \|\langle \xi \rangle^{|\beta|} \partial_{\xi}^{\beta} \varphi_{\mu}\|_{L^{\infty}} < \infty, \end{aligned} \tag{63}$$

where N_0 is a sufficiently large integer, and let \tilde{u} stand for u or \bar{u} . Then, there exist $N \in \mathbb{N}$ and $A > 0$ such that, for all $T > 0$ and every $S_1 = s_1(x, D)$, $S_2 = s_2(x, D)$ in $\text{Op} S_{0,0}^0$, we have

$$\int_0^T \left| \langle S_1 J^s T_b J^{-s} \nabla \tilde{u}, S_2 u \rangle \right| dt \leq A \|s_1\|_{C^N} \|s_2\|_{C^N} \sup_{\mu} \sup_{|\beta| \leq N} \|\langle \xi \rangle^{|\beta|} \partial_{\xi}^{\beta} \varphi_{\mu}\|_{L^{\infty}} \|J^{\frac{1}{2}} u\|_T^2.$$

Proof. One can write

$$\begin{aligned} & \langle S_1 J^s T_b J^{-s} \nabla \tilde{u}, S_2 u \rangle \\ &= \sum_{\mu} \alpha_{\mu} \langle S_1 J^s T_{\varphi_{\mu}} J^{-s} \nabla \tilde{u}, S_2 u \rangle = \sum_{\mu} \alpha_{\mu} \langle S_2^* S_1 J^s T_{\varphi_{\mu}} J^{-s} \nabla \tilde{u}, u \rangle \\ &= \sum_{\mu} \alpha_{\mu} \langle \langle x - \mu \rangle^{\sigma_0} J^{-\frac{1}{2}} S_2^* S_1 J^s T_{\varphi_{\mu}} J^{-s} \nabla J^{-\frac{1}{2}} \langle x - \mu \rangle^{\sigma_0} \langle x - \mu \rangle^{-\sigma_0} J^{\frac{1}{2}} \tilde{u}, \langle x - \mu \rangle^{-\sigma_0} J^{\frac{1}{2}} u \rangle \\ &= \sum_{\mu} \alpha_{\mu} \langle S_{\mu} \langle x - \mu \rangle^{\sigma_0} T_{\varphi_{\mu}} \langle x - \mu \rangle^{\sigma_0} J_{\mu} \tilde{u}_{\mu}, u_{\mu} \rangle \end{aligned}$$

where

$$S_{\mu} = \langle x - \mu \rangle^{\sigma_0} J^{-\frac{1}{2}} S_2^* S_1 J^s \langle x - \mu \rangle^{-\sigma_0}, \quad J_{\mu} = \langle x - \mu \rangle^{-\sigma_0} J^{-s} \nabla J^{-\frac{1}{2}} \langle x - \mu \rangle^{\sigma_0}, \quad u_{\mu} = \langle x - \mu \rangle^{-\sigma_0} J^{\frac{1}{2}} u.$$

Now, it follows from the pseudodifferential calculus (Theorem 2.1) and from Lemma 2.3 that S_{μ} and J_{μ} are in $\text{Op } S_{0,0}^{s-\frac{1}{2}}$ and $\text{Op } S_{0,0}^{\frac{1}{2}-s}$ respectively, and that we can estimate their seminorms uniformly in μ . Next, it follows from Lemma 2.12 that the operator norm of $\langle x - \mu \rangle^{\sigma_0} T_{\varphi_{\mu}} \langle x - \mu \rangle^{\sigma_0}$ acting in $H^{s-\frac{1}{2}}(\mathbb{R}^n)$ is estimated by $\sup_{|\beta| \leq N} \|\langle \xi \rangle^{|\beta|} \partial_{\xi}^{\beta} \varphi_{\mu}\|_{L^{\infty}}$ uniformly in μ . Hence, the application of the Cauchy–Schwarz inequality and the Calderón–Vaillancourt theorem allows us to obtain

$$\begin{aligned} & \int_0^T |\langle S_1 J^s T_b J^{-s} \nabla \tilde{u}, S_2 u \rangle| dt \\ & \leq \sum_{\mu} |\alpha_{\mu}| \|S_{\mu}\|_{\mathcal{L}(H^{s-1/2}, L^2)} \|\langle x - \mu \rangle^{\sigma_0} T_{\varphi_{\mu}} \langle x - \mu \rangle^{\sigma_0}\|_{\mathcal{L}(H^{s-1/2})} \|J_{\mu}\|_{\mathcal{L}(L^2, H^{s-1/2})} \int_0^T \|u_{\mu}\|_0^2 dt \\ & \leq A \|s_1\|_{C^N} \|s_2\|_{C^N} \sup_{\mu} \sup_{|\beta| \leq N} \|\langle \xi \rangle^{|\beta|} \partial_{\xi}^{\beta} \varphi_{\mu}\|_{L^{\infty}} \|J^{\frac{1}{2}} u\|_T^2, \end{aligned}$$

which proves the lemma. □

Now, let us write $T_{b_k - b_{k,m}} = T_{b_k - b_{k,m'}} + T_{b_{k,m'} - b_{k,m}}$ and apply Lemma A.1 first to $b = b_k - b_{k,m'}$ with $S_1 = S_2 = C_m$. We obtain

$$\begin{aligned} & \int_0^T |\langle C_m J^s T_{b_k - b_{k,m'}} \nabla \tilde{u}, C_m J^s u \rangle| dt \\ &= \int_0^T |\langle C_m J^s T_{b_k - b_{k,m'}} J^{-s} \nabla \tilde{v}, C_m v \rangle| dt \\ &\leq A \sup_{\mu} v_N(\varphi_{1,\mu,m})^2 \sup_{\mu} \sup_{|\beta| \leq N} \|\langle \xi \rangle^{|\beta|} \partial_{\xi}^{\beta} (\varphi_{k,\mu} - \varphi_{k,\mu,m'})\|_{L^{\infty}} \|J^{\frac{1}{2}} v\|_T^2, \\ &\leq A \frac{m^{2N^2}}{m^{\sigma}} \sup_{\mu} \sup_{|\beta| \leq N} \|\langle \xi \rangle^{|\beta|} \partial_{\xi}^{\beta} \varphi_{k,\mu}\|_{C^{\sigma}} \|J^{\frac{1}{2}+s} u\|_T^2 \leq A \frac{m^{2N^2}}{m^{\sigma}} \|J^{\frac{1}{2}+s} u\|_T^2, \end{aligned}$$

where $v = J^s u$ and $\sigma = \inf\{\varrho, 1\}$. As for the study of the other term, we write

$$\begin{aligned} & \langle C_m J^s T_{b_{k,m'} - b_{k,m}} \nabla \tilde{u}, C_m J^s u \rangle \\ &= \langle C_m J^s T_{b_{k,m'} - b_{k,m}} J^{-s} \nabla \tilde{v}, C_m v \rangle \\ &= \langle J^s T_{b_{k,m'} - b_{k,m}} J^{-s} \nabla C_m \tilde{v}, C_m v \rangle + \langle [C_m, J^s T_{b_{k,m'} - b_{k,m}} J^{-s} \nabla] \tilde{v}, C_m v \rangle, \quad (64) \end{aligned}$$

and then apply Lemma 5.2(iii) to the second term in (64) to obtain

$$\begin{aligned} \int_0^T | \langle [C_m, J^s T_{b_{k,m'}-b_{k,m}} J^{-s} \nabla] \tilde{v}, C_m v \rangle | dt & \\ & \leq A \sup_{\mu} v_N(\varphi_{1,\mu,m})^2 \sup_{\mu} \|\varphi_{k,\mu,m'} - \varphi_{k,\mu,m}\|_{C^N} \left(T \sup_{[0,T]} \|v\|_0^2 + \frac{1}{R} \| \| J^{\frac{1}{2}} v \| \|_T^2 \right) \\ & \leq A m'^{2N^2} (m'^N + m^N) \left(T \sup_{[0,T]} \|u\|_s^2 + \frac{1}{R} \| \| J^{s+\frac{1}{2}} u \| \|_T^2 \right) \\ & \leq A m'^{2N^2+N} \left(T \sup_{[0,T]} \|u\|_s^2 + \frac{1}{R} \| \| J^{s+\frac{1}{2}} u \| \|_T^2 \right). \end{aligned}$$

Finally, recalling that $C_m \bar{u} = \overline{C_m u}$ and applying Lemma A.1 to the first term in (64) with $S_1 = S_2 = \text{Id}$, we get

$$\begin{aligned} \int_0^T | \langle J^s T_{b_{k,m'}-b_{k,m}} J^{-s} \nabla C_m \tilde{v}, C_m v \rangle | dt & \\ & \leq A \sup_{\mu} \sup_{|\beta| \leq N} \| \langle \xi \rangle^{|\beta|} \partial_{\xi}^{\beta} (\varphi_{k,\mu,m'} - \varphi_{k,\mu,m}) \|_{L^{\infty}} \| \| J^{\frac{1}{2}} C_m v \| \|_T^2 \\ & \leq A \left(\sup_{\mu} \sup_{|\beta| \leq N} \| \langle \xi \rangle^{|\beta|} \partial_{\xi}^{\beta} (\varphi_{k,\mu,m'} - \varphi_{k,\mu,m}) \|_{L^{\infty}} + \sup_{\mu} \sup_{|\beta| \leq N} \| \langle \xi \rangle^{|\beta|} \partial_{\xi}^{\beta} (\varphi_{k,\mu} - \varphi_{k,\mu,m}) \|_{L^{\infty}} \right) \| \| J^{\frac{1}{2}} C_m v \| \|_T^2 \\ & \leq \left(\frac{A}{m'^{\sigma}} + \frac{A}{m^{\sigma}} \right) \sup_{\mu} \sup_{|\beta| \leq N} \| \langle \xi \rangle^{|\beta|} \partial_{\xi}^{\beta} \varphi_{k,\mu} \|_{C^{\sigma}} \| \| J^{\frac{1}{2}} C_m v \| \|_T^2 \\ & \leq \left(\frac{A}{m'^{\sigma}} + \frac{A}{m^{\sigma}} \right) \| \| J^{\frac{1}{2}} C_m v \| \|_T^2 \leq \frac{A}{m^{\sigma}} \| \| J^{\frac{1}{2}} C_m v \| \|_T^2. \end{aligned}$$

It remains to compare $\| \| J^{\frac{1}{2}} C_m v \| \|_T^2 = \| \| J^{\frac{1}{2}} C_m J^s u \| \|_T^2$ with $\| \| J^{s+\frac{1}{2}} C_m u \| \|_T^2$. Of course, one can write $J^{\frac{1}{2}} C_m J^s u = J^{s+\frac{1}{2}} J^{-s} C_m J^s u$ and it follows from Lemma 6.1 that $J^{-s} C_m J^s - C_m = E_m$ is in $\text{Op } S_{0,0}^{-1}$ and the seminorms of E_m are bounded by those of C_m . Hence, since $J^{s+\frac{1}{2}} E_m J^{-s}$ is in $\text{Op } S_{0,0}^{-\frac{1}{2}}$,

$$\begin{aligned} \| \| J^{s+\frac{1}{2}} E_m u \| \|_T^2 & = \sup_{\mu} \int_0^T \int | \langle x - \mu \rangle^{-\sigma_0} J^{s+\frac{1}{2}} E_m u |^2 dx dt \\ & \leq \int_0^T \int | J^{s+\frac{1}{2}} E_m u |^2 dx dt \\ & \leq A \sup_{\mu} v_N(\varphi_{1,\mu,m})^2 \int_0^T \int | J^s u |^2 dx dt \leq A T m^{2N^2} \sup_{[0,T]} \|u\|_s^2 \end{aligned}$$

and

$$\| \| J^{\frac{1}{2}} C_m v \| \|_T^2 \leq 2 \| \| J^{s+\frac{1}{2}} C_m u \| \|_T^2 + 2 A T m^{2N^2} \sup_{[0,T]} \|u\|_s^2,$$

which implies that

$$\int_0^T | \langle J^s T_{b_{k,m'}-b_{k,m}} J^{-s} \nabla C_m \tilde{v}, C_m v \rangle | dt \leq \frac{A}{m^{\sigma}} \| \| J^{s+\frac{1}{2}} C_m u \| \|_T^2 + A T m^{2N^2} \sup_{[0,T]} \|u\|_s^2,$$

where, of course, the constant A has changed. Summing up, we have proven that

$$\int_0^T | \langle C_m J^s T_{b_k - b_{k,m}} \nabla \tilde{u}, C_m J^s u \rangle | dt \leq \frac{Am^{2N^2}}{m'^\sigma} \| J^{s+\frac{1}{2}} u \|_T^2 + Am'^{2N^2+N} \left(T \sup_{[0,T]} \| u \|_s^2 + \frac{1}{R} \| J^{s+\frac{1}{2}} u \|_T^2 \right) + \frac{A}{m^\sigma} \| J^{s+\frac{1}{2}} C_m u \|_T^2; \tag{65}$$

that is, we have proven Lemma 3.4.

Anisotropic Sobolev spaces. There are several notions of anisotropic Sobolev space in the literature. However, we have not been able to find a reference with the results we need in this paper. Therefore, we are going to define our spaces and next prove the results we need.

We denote by (x, y) the variable in $\mathbb{R}^n \times \mathbb{R}^{n'}$ and by (ξ, η) its Fourier dual variable.

Definition A.2. If $s, s' \in \mathbb{R}$, we denote by $H^{s,s'}(\mathbb{R}^n \times \mathbb{R}^{n'})$ the space of tempered distributions u in $\mathbb{R}^n \times \mathbb{R}^{n'}$ such that the integral

$$\int_{\mathbb{R}^n \times \mathbb{R}^{n'}} \langle \xi \rangle^{2s} \langle \eta \rangle^{2s'} |\hat{u}(\xi, \eta)|^2 d\xi d\eta \tag{66}$$

is finite.

We call this space an anisotropic Sobolev space. Note that this is different, for example, from the classical space $H^{r,s}$ of [Lions and Magenes 1968]. Clearly, $H^{s,s'}(\mathbb{R}^n \times \mathbb{R}^{n'})$ is a Hilbert space when it is provided with the obvious scalar product. We also denote by $\|u\|_{s,s'}$ the norm of u in this space and, of course, it is equal to the square root of (66).

Additionally, note that the space $H^{s,s'}(\mathbb{R}^n \times \mathbb{R}^{n'})$ in the above definition coincides with the space $H^s(\mathbb{R}^n, H^{s'}(\mathbb{R}^{n'}))$ and, by symmetry, with $H^{s'}(\mathbb{R}^{n'}, H^s(\mathbb{R}^n))$.

In this paper, we need the following two results on anisotropic Sobolev spaces. The first one is the Sobolev injection:

Proposition A.3. *If $s > \frac{n}{2}$ and $s' > \frac{n'}{2}$, then $H^{s,s'}(\mathbb{R}^n \times \mathbb{R}^{n'}) \subset L^\infty(\mathbb{R}^n \times \mathbb{R}^{n'})$ with continuous injection.*

Proof. If $u \in H^{s,s'}$, then

$$\hat{u}(\xi, \eta) = \langle \xi \rangle^{-s} \langle \eta \rangle^{-s'} \cdot \langle \xi \rangle^s \langle \eta \rangle^{s'} \hat{u}(\xi, \eta);$$

hence, $\hat{u} \in L^2 \cdot L^2 \subset L^1$ and $\|u\|_{L^\infty} \leq C \|\hat{u}\|_{L^1} \leq C' \|u\|_{s,s'}$, where C and C' are constants which are independent of u . □

The other result is an interpolation inequality:

Proposition A.4. *If $s = (1 - \theta)s_1 + \theta s_2$ and $s' = (1 - \theta)s'_1 + \theta s'_2$, where $\theta \in [0, 1]$, $s_1, s_2, s'_1, s'_2 \in \mathbb{R}$, then, for any $u \in H^{s_1, s'_1}(\mathbb{R}^n \times \mathbb{R}^{n'}) \cap H^{s_2, s'_2}(\mathbb{R}^n \times \mathbb{R}^{n'})$, we have*

$$\|u\|_{s,s'} \leq \|u\|_{s_1, s'_1}^{1-\theta} \|u\|_{s_2, s'_2}^\theta.$$

Proof. Indeed, we have

$$\begin{aligned} \|u\|_{s,s'}^2 &= \int_{\mathbb{R}^n \times \mathbb{R}^{n'}} \langle \xi \rangle^{2(1-\theta)s_1 + 2\theta s_2} \langle \eta \rangle^{2(1-\theta)s'_1 + 2\theta s'_2} |\hat{u}(\xi, \eta)|^2 d\xi d\eta \\ &= \int_{\mathbb{R}^n \times \mathbb{R}^{n'}} \left(\langle \xi \rangle^{s_1} \langle \eta \rangle^{s'_1} |\hat{u}(\xi, \eta)| \right)^{2(1-\theta)} \left(\langle \xi \rangle^{s_2} \langle \eta \rangle^{s'_2} |\hat{u}(\xi, \eta)| \right)^{2\theta} d\xi d\eta \\ &\leq \left(\int_{\mathbb{R}^n \times \mathbb{R}^{n'}} \langle \xi \rangle^{2s_1} \langle \eta \rangle^{2s'_1} |\hat{u}(\xi, \eta)|^2 d\xi d\eta \right)^{1-\theta} \left(\int_{\mathbb{R}^n \times \mathbb{R}^{n'}} \langle \xi \rangle^{2s_2} \langle \eta \rangle^{2s'_2} |\hat{u}(\xi, \eta)|^2 d\xi d\eta \right)^\theta \\ &= \|u\|_{s_1, s'_1}^{2(1-\theta)} \|u\|_{s_2, s'_2}^{2\theta}, \end{aligned}$$

where we have applied Hölder’s inequality. □

Actually, we need the above results for anisotropic Sobolev spaces on domains Ω in $\mathbb{R}^n \times \mathbb{R}^{n'}$, and since the theory of such spaces is less simple, we shall restrict ourselves to the case that arises in this paper, that is, the case $\Omega = I \times \mathbb{R}^n$ where I is a bounded interval in \mathbb{R} , and only to the case $s \geq 0$. First, let us set, by definition,

$$H^{s,s'}(\Omega) = H^s(I, H^{s'}(\mathbb{R}^n)),$$

in the sense that $u(x, y)$ is in $H^{s,s'}(\Omega)$ if and only if

$$\partial_x^\alpha J_y^{s'} u \in L^2(\Omega) \quad \text{for } |\alpha| \leq s$$

and

$$\int_{I \times I \times \mathbb{R}^n} \frac{|\partial_x^\alpha J_y^{s'} u(x, y) - \partial_x^\alpha J_y^{s'} u(x', y)|^2}{|x - x'|^{1+2\sigma}} dx dx' dy < \infty \quad \text{if } 0 < \sigma = s - [s] < 1.$$

Of course, the norm in this space is defined by

$$\|u\|_{s,s',\Omega}^2 = \sum_{|\alpha| \leq s} \|\partial_x^\alpha J_y^{s'} u\|_{L^2(\Omega)}^2 \quad \text{if } s \in \mathbb{N},$$

and

$$\|u\|_{s,s',\Omega}^2 = \sum_{|\alpha| \leq [s]} \|\partial_x^\alpha J_y^{s'} u\|_{L^2(\Omega)}^2 + \int_{I \times I \times \mathbb{R}^n} \frac{|\partial_x^\alpha J_y^{s'} u(x, y) - \partial_x^\alpha J_y^{s'} u(x', y)|^2}{|x - x'|^{1+2\sigma}} dx dx' dy \quad \text{otherwise.}$$

Now, we can prove for $H^{s,s'}(\Omega)$ the results analogous to the above ones.

Proposition A.5. (i) *If $s > \frac{1}{2}$ and $s' > \frac{n}{2}$, then $H^{s,s'}(\Omega) \subset L^\infty(\Omega)$ with continuous injection.*

(ii) *If $s = (1 - \theta)s_1 + \theta s_2$ and $s' = (1 - \theta)s'_1 + \theta s'_2$, where $\theta \in [0, 1]$, $s_1 \geq 0$, $s_2 \geq 0$, $s'_1, s'_2 \in \mathbb{R}$, then there exists a constant C such that, for any $u \in H^{s_1, s'_1}(\Omega) \cap H^{s_2, s'_2}(\Omega)$, we have*

$$\|u\|_{s,s',\Omega} \leq C \|u\|_{s_1, s'_1, \Omega}^{1-\theta} \|u\|_{s_2, s'_2, \Omega}^\theta.$$

Proof. Since we cannot use directly the Fourier transformation, the idea is to construct a bounded linear extension operator

$$P_\Omega : H^{s,s'}(\Omega) \rightarrow H^{s,s'}(\mathbb{R} \times \mathbb{R}^n), \tag{67}$$

that is, it satisfies $P_\Omega u|_\Omega = u$, for all $u \in H^{s,s'}(\Omega)$. Indeed, assume that such a P_Ω exists. Then, for $u \in H^{s,s'}(\Omega)$ with $s > \frac{1}{2}$ and $s' > \frac{n}{2}$,

$$\|u\|_{L^\infty(\Omega)} = \|P_\Omega u\|_{L^\infty(\Omega)} \leq \|P_\Omega u\|_{L^\infty(\mathbb{R} \times \mathbb{R}^n)} \leq C \|P_\Omega u\|_{s,s'} \leq C' \|u\|_{s,s',\Omega},$$

where we have applied Proposition A.3 and the boundedness of P_Ω , and this proves (i).

Furthermore, under the assumptions of (ii), we have

$$\|u\|_{s,s',\Omega} = \|P_\Omega u\|_{s,s',\Omega} \leq \|P_\Omega u\|_{s,s',\mathbb{R} \times \mathbb{R}^n},$$

and it is a classical fact that there exists a constant C such that, for all $v \in H^s(\mathbb{R}^d)$,

$$\sum_{|\alpha| \leq [s]} \|\partial^\alpha v\|_{L^2(\mathbb{R}^d)}^2 + \int_{\mathbb{R}^d \times \mathbb{R}^d} \frac{|\partial^\alpha v(x) - \partial^\alpha v(x')|^2}{|x - x'|^{d+2\sigma}} dx dx' \leq C \|v\|_s^2;$$

now, applying this inequality to $v(x) = J_y^{s'} P_\Omega u(x, y)$, with $d = 1$, and integrating with respect to y gives

$$\|P_\Omega u\|_{s,s',\mathbb{R} \times \mathbb{R}^n}^2 \leq C \|P_\Omega u\|_{s,s'}^2.$$

Finally, applying Proposition A.4 and the boundedness of P_Ω yields

$$\|u\|_{s,s',\Omega} \leq \sqrt{C} \|P_\Omega u\|_{s,s'} \leq \sqrt{C} \|P_\Omega u\|_{s_1,s'_1}^{1-\theta} \|P_\Omega u\|_{s_2,s'_2}^\theta \leq C' \|u\|_{s_1,s'_1,\Omega}^{1-\theta} \|u\|_{s_2,s'_2,\Omega}^\theta,$$

which establishes (ii).

It remains to construct P_Ω as in (67). In fact, the classical theory of Sobolev spaces already provides a bounded linear extension operator

$$P_I : H^s(I) \rightarrow H^s(\mathbb{R}) \tag{68}$$

such that $P_I u|_I = u$ for all $u \in H^s(I)$. If $u \in H^{s,s'}(\Omega)$, let us set

$$P_\Omega u(x, y) = (P_I)_x u(x, y).$$

Clearly, this defines a linear operator such that $P_\Omega u|_\Omega = u$. Let us show the boundedness of $P_\Omega : H^{s,s'}(\Omega) \rightarrow H^{s,s'}(\mathbb{R} \times \mathbb{R}^n)$. It follows from the definition that $x \mapsto J_y^{s'} u(x, y)$ is in the Sobolev space $H^s(I)$ for almost all $y \in \mathbb{R}^n$. Hence, $x \mapsto (P_I)_x J_y^{s'} u(x, y)$ is in $H^s(\mathbb{R})$ for almost all $y \in \mathbb{R}^n$ and there exists a constant C which depends neither on u nor on y such that

$$\|(P_I)_x J_y^{s'} u(x, y)\|_{H^s(\mathbb{R})} \leq C \|J_y^{s'} u(x, y)\|_{H^s(I)} \quad \text{for a.e. } y \in \mathbb{R}^n.$$

Since $(P_I)_x J_y^{s'} u = J_y^{s'} P_\Omega u$, this inequality can be written more explicitly as

$$\begin{aligned} \int_{\mathbb{R}} |J_x^s J_y^{s'} P_\Omega u(x, y)|^2 dx &\leq C^2 \sum_{|\alpha| \leq [s]} \int_I |\partial_x^\alpha J_y^{s'} u(x, y)|^2 dx \\ &+ C^2 \int_{I \times I} \frac{|\partial_x^\alpha J_y^{s'} u(x, y) - \partial_x^\alpha J_y^{s'} u(x', y)|^2}{|x - x'|^{1+2\sigma}} dx dx' \quad \text{for a.e. } y \in \mathbb{R}^n. \end{aligned}$$

Integrating over \mathbb{R}^n with respect to y gives

$$\|P_{\Omega}u\|_{s,s'}^2 \leq C^2 \|u\|_{s,s',\Omega}^2,$$

which proves the boundedness of P_{Ω} and achieves the proof of the proposition. \square

References

- [Bejenaru and Tataru 2008] I. Bejenaru and D. Tataru, “Large data local solutions for the derivative NLS equation”, *J. Eur. Math. Soc. (JEMS)* **10**:4 (2008), 957–985. MR Zbl
- [Bergh and Löfström 1976] J. Bergh and J. Löfström, *Interpolation spaces: an introduction*, Grundlehren der Mathematischen Wissenschaften **223**, Springer, 1976. MR Zbl
- [Bienaimé 2014] P.-Y. Bienaimé, “Existence locale et effet régularisant précisés pour des équations non linéaires de type Schrödinger”, *Rev. Mat. Iberoam.* **30**:3 (2014), 751–798. MR Zbl
- [Bony 1981] J.-M. Bony, “Calcul symbolique et propagation des singularités pour les équations aux dérivées partielles non linéaires”, *Ann. Sci. École Norm. Sup. (4)* **14**:2 (1981), 209–246. MR Zbl
- [Coifman and Meyer 1978] R. R. Coifman and Y. Meyer, *Au delà des opérateurs pseudo-différentiels*, Astérisque **57**, Société Mathématique de France, Paris, 1978. MR Zbl
- [Doi 1994] S.-i. Doi, “On the Cauchy problem for Schrödinger type equations and the regularity of solutions”, *J. Math. Kyoto Univ.* **34**:2 (1994), 319–328. MR Zbl
- [Kenig et al. 1998] C. E. Kenig, G. Ponce, and L. Vega, “Smoothing effects and local existence theory for the generalized nonlinear Schrödinger equations”, *Invent. Math.* **134**:3 (1998), 489–545. MR Zbl
- [Kenig et al. 2004] C. E. Kenig, G. Ponce, and L. Vega, “The Cauchy problem for quasi-linear Schrödinger equations”, *Invent. Math.* **158**:2 (2004), 343–388. MR Zbl
- [Kenig et al. 2005] C. E. Kenig, G. Ponce, C. Rolvung, and L. Vega, “Variable coefficient Schrödinger flows for ultrahyperbolic operators”, *Adv. Math.* **196**:2 (2005), 373–486. MR Zbl
- [Kenig et al. 2006] C. E. Kenig, G. Ponce, C. Rolvung, and L. Vega, “The general quasilinear ultrahyperbolic Schrödinger equation”, *Adv. Math.* **206**:2 (2006), 402–433. MR Zbl
- [Lions and Magenes 1968] J.-L. Lions and E. Magenes, *Problèmes aux limites non homogènes et applications, II*, Travaux et Recherches Mathématiques **18**, Dunod, Paris, 1968. MR Zbl
- [Marzuola et al. 2012] J. L. Marzuola, J. Metcalfe, and D. Tataru, “Quasilinear Schrödinger equations, I: Small data and quadratic interactions”, *Adv. Math.* **231**:2 (2012), 1151–1172. MR Zbl
- [Marzuola et al. 2014] J. L. Marzuola, J. Metcalfe, and D. Tataru, “Quasilinear Schrödinger equations, II: Small data and cubic nonlinearities”, *Kyoto J. Math.* **54**:3 (2014), 529–546. MR Zbl
- [Meyer 1981] Y. Meyer, “Remarques sur un théorème de J.-M. Bony”, pp. 1–20 in *Proceedings of the Seminar on Harmonic Analysis* (Pisa, 1980), 1981. Issue as a supplement to *Rend. Circ. Mat. Palermo (2)*. MR Zbl
- [Meyer 1982] Y. Meyer, “Nouvelles estimations pour les solutions d’équations aux dérivées partielles non linéaires”, exposé 6, pp. 1–12 in *Goulaouic–Meyer–Schwartz Seminar*, 1981/1982, École Polytech., Palaiseau, 1982. MR Zbl
- [Robbiano 2013] L. Robbiano, “Kato smoothing effect for Schrödinger operator”, pp. 355–369 in *Studies in phase space analysis with applications to PDEs*, edited by M. Cicognani et al., Progr. Nonlinear Differential Equations Appl. **84**, Springer, 2013. MR Zbl
- [Takeuchi 1992] J. Takeuchi, “Le problème de Cauchy pour certaines équations aux dérivées partielles du type de Schrödinger, VIII: Symétrisations indépendantes du temps”, *C. R. Acad. Sci. Paris Sér. I Math.* **315**:10 (1992), 1055–1058. MR Zbl
- [Tataru 2002] D. Tataru, “On the Fefferman–Phong inequality and related problems”, *Comm. Partial Differential Equations* **27**:11–12 (2002), 2101–2138. MR Zbl
- [Taylor 1991] M. E. Taylor, *Pseudodifferential operators and nonlinear PDE*, Progress in Mathematics **100**, Birkhäuser, Boston, 1991. MR Zbl

Received 20 Jun 2017. Revised 26 Nov 2017. Accepted 2 Jan 2018.

PIERRE-YVES BIENAIMÉ: pierre-yves.bienaime@univ-nantes.fr

Laboratoire de Mathématiques Jean Leray, CNRS UMR6629, Université de Nantes, Nantes, France

ABDESSLAM BOULKHEMAIR: boulkhemair-a@univ-nantes.fr

Laboratoire de Mathématiques Jean Leray, CNRS UMR6629, Université de Nantes, Nantes, France

Analysis & PDE

msp.org/apde

EDITORS

EDITOR-IN-CHIEF

Patrick Gérard
patrick.gerard@math.u-psud.fr
Université Paris Sud XI
Orsay, France

BOARD OF EDITORS

Massimiliano Berti	Scuola Intern. Sup. di Studi Avanzati, Italy berti@sissa.it	Clément Mouhot	Cambridge University, UK c.mouhot@dpmms.cam.ac.uk
Sun-Yung Alice Chang	Princeton University, USA chang@math.princeton.edu	Werner Müller	Universität Bonn, Germany mueller@math.uni-bonn.de
Michael Christ	University of California, Berkeley, USA mchrist@math.berkeley.edu	Gilles Pisier	Texas A&M University, and Paris 6 pisier@math.tamu.edu
Alessio Figalli	ETH Zurich, Switzerland alessio.figalli@math.ethz.ch	Tristan Rivière	ETH, Switzerland riviere@math.ethz.ch
Charles Fefferman	Princeton University, USA cf@math.princeton.edu	Igor Rodnianski	Princeton University, USA irod@math.princeton.edu
Ursula Hamenstaedt	Universität Bonn, Germany ursula@math.uni-bonn.de	Sylvia Serfaty	New York University, USA serfaty@cims.nyu.edu
Vaughan Jones	U.C. Berkeley & Vanderbilt University vaughan.f.jones@vanderbilt.edu	Yum-Tong Siu	Harvard University, USA siu@math.harvard.edu
Vadim Kaloshin	University of Maryland, USA vadim.kaloshin@gmail.com	Terence Tao	University of California, Los Angeles, USA tao@math.ucla.edu
Herbert Koch	Universität Bonn, Germany koch@math.uni-bonn.de	Michael E. Taylor	Univ. of North Carolina, Chapel Hill, USA met@math.unc.edu
Izabella Laba	University of British Columbia, Canada ilaba@math.ubc.ca	Gunther Uhlmann	University of Washington, USA gunther@math.washington.edu
Gilles Lebeau	Université de Nice Sophia Antipolis, France lebeau@unice.fr	András Vasy	Stanford University, USA andras@math.stanford.edu
Richard B. Melrose	Massachusetts Inst. of Tech., USA rbm@math.mit.edu	Dan Virgil Voiculescu	University of California, Berkeley, USA dvv@math.berkeley.edu
Frank Merle	Université de Cergy-Pontoise, France Frank.Merle@u-cergy.fr	Steven Zelditch	Northwestern University, USA zelditch@math.northwestern.edu
William Minicozzi II	Johns Hopkins University, USA minicozz@math.jhu.edu	Maciej Zworski	University of California, Berkeley, USA zworski@math.berkeley.edu

PRODUCTION

production@msp.org
Silvio Levy, Scientific Editor


See inside back cover or msp.org/apde for submission instructions.

The subscription price for 2018 is US \$275/year for the electronic version, and \$480/year (+\$55, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues from the last three years and changes of subscriber address should be sent to MSP.

Analysis & PDE (ISSN 1948-206X electronic, 2157-5045 printed) at Mathematical Sciences Publishers, 798 Evans Hall #3840, c/o University of California, Berkeley, CA 94720-3840, is published continuously online. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices.

APDE peer review and production are managed by EditFlow® from MSP.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2018 Mathematical Sciences Publishers

ANALYSIS & PDE

Volume 11 No. 5 2018

Large sets avoiding patterns	1083
ROBERT FRASER and MALABIKA PRAMANIK	
On minimizers of an isoperimetric problem with long-range interactions under a convexity constraint	1113
MICHAEL GOLDMAN, MATTEO NOVAGA and BERARDO RUFFINI	
Nonautonomous maximal L^p -regularity under fractional Sobolev regularity in time	1143
STEPHAN FACKLER	
Transference of bilinear restriction estimates to quadratic variation norms and the Dirac–Klein–Gordon system	1171
TIMOTHY CANDY and SEBASTIAN HERR	
Well-posedness and smoothing effect for generalized nonlinear Schrödinger equations	1241
PIERRE-YVES BIENAIMÉ and ABDESSLAM BOULKHEMAIR	
The shape of low energy configurations of a thin elastic sheet with a single disclination	1285
HEINER OLBERMANN	
The thin-film equation close to self-similarity	1303
CHRISTIAN SEIS	



2157-5045(2018)11:5;1-P