

ANALYSIS & PDE

Volume 11

No. 5

2018

CHRISTIAN SEIS

THE THIN-FILM EQUATION CLOSE TO SELF-SIMILARITY

THE THIN-FILM EQUATION CLOSE TO SELF-SIMILARITY

CHRISTIAN SEIS

We study well-posedness and regularity of the multidimensional thin-film equation with linear mobility in a neighborhood of the self-similar Smyth–Hill solutions. To be more specific, we perform a von Mises change of dependent and independent variables that transforms the thin-film free boundary problem into a parabolic equation on the unit ball. We show that the transformed equation is well-posed and that solutions are smooth and even analytic in time and angular direction. The latter gives the analyticity of level sets of the original equation, and thus, in particular, of the free boundary.

1. Introduction and main results

1.1. The background. We are concerned with a thin-film equation in arbitrary space dimensions. Our interest is in the simplest case of linear mobility; that is, we consider the partial differential equation

$$\partial_t u + \nabla \cdot (u \nabla \Delta u) = 0 \quad (1)$$

in \mathbb{R}^N . In this model, u describes the thickness of a viscous thin liquid film on a flat substrate. We will focus on what is usually referred to as the complete wetting regime, in which the liquid–solid contact angle at the film boundary is presumed to be zero. Notice that in the three-dimensional physical space, the dimension N of the substrate is 2.

Equation (1) belongs to the following family of thin-film equations:

$$\partial_t u + \nabla \cdot (m(u) \nabla \Delta u) = 0, \quad (2)$$

where the mobility factor is given by $m(u) = u^3 + \beta^{n-3} u^n$ with β being the slippage length. The nonlinearity exponent $n > 0$ depends on the slip condition at the solid–liquid interface: $n = 3$ models no-slip conditions and $n = 2$ models Navier–slip conditions. The case $n = 1$ is a further relaxation and the linear mobility considered here is obtained to leader order in the limit $u \rightarrow 0$.

The evolution described in (2) was originally derived as a long-wave approximation from the free-surface problem related to the Navier–Stokes equations and suitable model reductions; see, e.g., [Oron et al. 1997]. At the same time, it can be obtained as the Wasserstein gradient flow of the surface-tension energy [Otto 1998; Giacomelli and Otto 2001; Matthes et al. 2009] and serves thus as the natural dissipative model for surface-tension-driven transport of viscous liquids over solid substrates.

The analytical treatment of the equation is challenging and the mathematical understanding is far from being satisfactory. As a fourth-order problem, the thin-film equation lacks a maximum principle.

MSC2010: primary 35K30; secondary 76A20.

Keywords: thin-film equation, fourth-order equation, self-similar solution, well-posedness.

Moreover, the parabolicity degenerates where u vanishes and, as a consequence, for compactly supported initial data (“droplets”), the solution remains compactly supported [Bernis 1996; Bertsch et al. 1998]. The thin-film equation features thus a free boundary given by $\partial\{u > 0\}$, which in physical terms is the contact line connecting the phases liquid, solid and vapor. Nonetheless, by using estimates for the surface energy and compactness arguments Bernis and Friedman [1990] established the existence of weak nonnegative solutions over a quarter of a century ago. The regularity of these solutions could be slightly improved with the help of certain entropy-type estimates [Beretta et al. 1995; Bertozzi and Pugh 1996; Dal Passo et al. 1998], but this regularity is not sufficient for proving general uniqueness results. To gain an understanding of the thin-film equation and its qualitative features, it is thus natural to find and study special solutions first. In the past ten years, most of the attention has been focused on the one-dimensional setting, for instance, near stationary solutions [Giacomelli et al. 2008; Giacomelli and Knüpfer 2010], traveling waves [Giacomelli et al. 2014; Gnann 2016], and self-similar solutions [Gnann 2015; Belgacem et al. 2016]. The only regularity and well-posedness result in higher dimensions available so far is due to John [2015], who analyzed the equation around stationary solutions. For completeness, we remark the thin-film equation is also studied with nonzero contact angles; see, e.g., [Otto 1998; Giacomelli and Otto 2001; 2003; Knüpfer 2011; 2015; Knüpfer and Masmoudi 2015; Belgacem et al. 2016; Degtyarev 2017]. The latter of these works is particularly interesting as it deals with the multidimensional situation.

In the present paper, we will conduct a study similar to John’s and investigate the qualitative behavior of solutions close to self-similarity. A family of self-similar solutions to (1) is given by

$$u_*(t, x) = \frac{\alpha_N}{t^{N/(N+4)}} \left(\sigma_M - \frac{|x|^2}{t^{2/(N+4)}} \right)_+^2,$$

where $\alpha_N = 1/(8(N+4)(N+2))$ and σ_M is a positive number that is determined by the mass constraint

$$\int u_* dx = M,$$

and the subscript plus sign denotes the positive part of a quantity; i.e., $(\cdot)_+ = \max\{0, \cdot\}$. These solutions were first found by Smyth and Hill [1988] in the one-dimensional case and then rediscovered in [Ferreira and Bernis 1997]. As in related parabolic settings, the Smyth–Hill solutions play a distinguished role in the theory of the thin-film equation as they are believed to describe the leading-order large-time asymptotic behavior of *any* solution—a fact that is currently known only for strong [Carrillo and Toscani 2002] and minimizing movement [Matthes et al. 2009] solutions. Besides that, these particular solutions are considered to feature the same regularity properties as any “typical” solution, at least for large times. Thus, under suitable assumptions on the initial data, we expect the solutions of (1) to be smooth up to the boundary of their support. (Notice that this behavior is exclusive for the linear-mobility thin-film equation; see [Giacomelli et al. 2013].)

In the present work we consider solutions that are in some suitable sense close to the self-similar Smyth–Hill solution. Instead of working with (1) directly, we will perform a certain von Mises change of dependent and independent variables, which has the advantage that it freezes the free boundary $\partial\{u > 0\}$ to the unit ball. We will mainly address the following four questions:

- (1) Is there some uniqueness principle available for the transformed equation?
- (2) Are solutions smooth?
- (3) Can we deduce some regularity for the moving interface $\partial\{u > 0\}$?
- (4) Do solutions depend smoothly on the initial data?

We will provide positive answers to all of these questions. In fact, applying a perturbation procedure we will show that the transformed equation is well-posed in a sufficiently small neighborhood of u_* . We will furthermore show that the unique solution is smooth in time and space. In fact, our results show that solutions to the transformed equation are analytic in time and in direction tangential to the free boundary. The latter in particular implies that all level sets and thus also the free boundary line corresponding to the original solutions are analytic. We finally prove analytic dependence on the initial data.

The fact that solutions depend differentiably (or even better) on the initial data will be of great relevance in a companion study on fine large-time asymptotic expansions. Indeed, in [Seis \geq 2018], we investigate the rates at which solutions converge to the self-similarity at any order. Optimal rates of convergence were already found by Carrillo and Toscani [2002] and Matthes, McCann and Savaré [Matthes et al. 2009], and these rates are saturated by spatial translations of the Smyth–Hill solutions. Jointly with McCann [McCann and Seis 2015] we diagonalized the differential operator obtained after formal linearization around the self-similar solution. The goal of [Seis \geq 2018] is to translate the spectral information obtained in [McCann and Seis 2015] into large-time asymptotic expansions for the nonlinear problem. For this, it is necessary to rigorously linearize the equation, the framework for which is obtained in the current paper. This strategy was recently successfully applied to the porous-medium equation near the self-similar Barenblatt solutions [Seis 2014; 2015]. The present work parallels in parts [Seis 2015] as well as the pioneering work by Koch [1999] and the further developments by Kienzler [2016] and John [2015].

1.2. Global transformation onto fixed domain. In this subsection it is our goal to transform the thin-film equation (1) into a partial differential equation that is posed on a fixed domain and that appears to be more suitable for a regularity theory than the original equation. The first step is a customary change of variables that translates the self-similarly spreading Smyth–Hill solution into a stationary solution. This is, for instance, achieved by setting

$$\hat{x} = \frac{1}{\sqrt{\sigma_M}} \frac{x}{t^{1/(N+4)}}, \quad \hat{t} = \frac{1}{N+4} \log t, \quad \hat{u} = \frac{N+4}{\sigma_M^2} t^{N/(N+4)} u,$$

with the effect that the Smyth–Hill solution becomes

$$\gamma \hat{u}_*(\hat{x})^{1/2} = \frac{1}{2}(1 - |\hat{x}|^2)_+, \quad \gamma = \sqrt{2(N+2)}, \tag{3}$$

and the thin-film equation (1) turns into the *confined* thin-film equation

$$\partial_{\hat{t}} \hat{u} + \hat{\nabla} \cdot (\hat{u} \hat{\nabla} \hat{\Delta} \hat{u}) - \hat{\nabla} \cdot (\hat{x} \hat{u}) = 0. \tag{4}$$

It is easily checked that \hat{u}_* is indeed a stationary solution to (4) and mass is no longer spreading over all of \mathbb{R}^N . Instead, the confinement term pushes all mass towards the stationary \hat{u}_* at the origin. To simplify the notation in the following, we will drop the hats immediately!




Figure 1. The definition of the (z, w) -coordinates.

Now that the Smyth–Hill solution has become stationary, we will perform a change of dependent and independent variables that parametrizes the solution as a graph over u_* . This type of a change of variables is sometimes referred to as a von Mises transformation. It is convenient to temporarily introduce the variable $v = \gamma u^{1/2}$, so that $\sqrt{2v_*}$ maps the unit ball onto the upper half-sphere. The new variables are obtained by projecting the point $(x, \sqrt{2v(x)})$ orthogonally onto the graph of $\sqrt{2v_*}$: noting that $\sqrt{2v(x) + |x|^2}$ is the hypotenuse of the triangle with the edges $(0, 0)$, $(0, |x|)$ and $(|x|, \sqrt{2v(x)})$, the projection point has the coordinates $(z, \sqrt{2v_*(z)})$ with

$$z = \frac{x}{\sqrt{2v(x) + |x|^2}}.$$

We define the new dependent variable w as the distance of the point $(x, \sqrt{2v(x)})$ from the sphere; that is,

$$1 + w = \sqrt{2v + |x|^2}, \tag{5}$$

which gives that $x = (1 + w)z$. This change of variables is illustrated in [Figure 1](#).

The transformation of the thin-film equation (4) under this change of variables leads to straightforward but tedious computations that we conveniently defer to the [Appendix](#). The new variable w obeys the equation

$$\partial_t w + \mathcal{L}^2 w + N \mathcal{L} w = f[w] \tag{6}$$

on $B_1(0)$, where

$$\mathcal{L} w = -\rho^{-1} \nabla \cdot (\rho^2 \nabla w) = -\rho \Delta w + 2z \cdot \nabla w$$

is precisely the linear operator that is obtained by linearizing the porous-medium equation $\partial_t u - \Delta u^{3/2} = 0$ about the Barenblatt solution; see, e.g., [[Seis 2014; 2015](#)]. Before specifying the particular form of the nonlinearity $f[w]$, let us notice that the linear operator $\mathcal{L}^2 + N \mathcal{L}$ corresponding to the thin-film dynamics was previously found by McCann and the author [[McCann and Seis 2015](#)] by a formal computation. Its relation to the porous-medium linear operator \mathcal{L} is not surprising but reflects the deep relation between

both equations. Indeed, as first exploited by Carrillo and Toscani [2002], the dissipation rate of the porous-medium entropy is just the surface energy that drives the thin-film dynamics. On a more abstract level, this observation can be expressed by the so-called energy-information relation, first formulated by Matthes, McCann and Savaré [Matthes et al. 2009], which connects the Wasserstein gradient flow structures of both equations [Otto 1998; 2001; Giacomelli and Otto 2001].

Let us finally discuss the right-hand side of (6). We can split $f[w]$ according to $f[w] = f^1[w] + f^2[w] + f^3[w]$, where

$$\begin{aligned} f^1[w] &= p \star R_1[w] \star ((\nabla w)^{2\star} + \nabla w \star \nabla^2 w), \\ f^2[w] &= p \star R_1[w] \star \rho((\nabla^2 w)^{2\star} + \nabla w \star \nabla^3 w), \\ f^3[w] &= p \star R_2[w] \star \rho^2((\nabla^2 w)^{3\star} + \nabla \tilde{w} \star \nabla^2 \tilde{w} \star \nabla^3 w + (\nabla w)^{2\star} \star \nabla^4 w), \end{aligned}$$

and

$$R_i[w] = \frac{(\nabla w)^{k\star}}{(1 + w + z \cdot \nabla w)^{k+i}}$$

for some $k \in \mathbb{N}_0$. We will see in Section 3 below that the particular form of the nonlinearity is irrelevant for the perturbation argument. We have thus introduced a slightly condensed notation to simplify the terms in the nonlinearity: we write $f \star g$ to denote any arbitrary linear combination of the tensors (vectors, matrices) f and g . For instance, $\nabla^{m_1} \tilde{w} \star \nabla^{m_2} \tilde{w}$ is an arbitrary linear combination of products of derivatives of orders m_1 and m_2 . The iterated application of the \star is abbreviated as $f^{j\star} = f \star \dots \star f$ if the latter product has j factors. The conventions $f^{1\star} = 1 \star f$ and $f^{0\star} = 1$ apply. We furthermore use p as an arbitrary representative of a (tensor-valued) polynomial in z . We have only kept track of those ρ prefactors, which will be of importance later on.

1.3. The intrinsic geometry and function spaces. In our analysis of the linear equation associated to (6), i.e.,

$$\partial_t w + \mathcal{L}^2 w + N \mathcal{L} w = f \tag{7}$$

for some general f , we will make use of the framework developed earlier in [Koch 1999; Seis 2015] for the second-order equation

$$\partial_t w + \mathcal{L} w = f. \tag{8}$$

The underlying point of view in there is the fact that the previous equation can be interpreted as a heat flow on a weighted manifold, i.e., a Riemannian manifold to which a new volume element (typically a positive multiple of the one induced by the metric tensor) is assigned. The theories for heat flows on a weighted manifold parallel those on Riemannian manifolds in many respects; see [Grigor'yan 2006]. For instance, a Calderón–Zygmund theory is available for (8). The crucial idea in [Koch 1999; Seis 2015] is to trade the Euclidean distance on $B_1(0)$ for the geodesic distance induced by the heat flow interpretation. In this way, we equip the unit ball with a non-Euclidean Carnot–Carathéodory metric, see, e.g., [Bellaïche and Risler 1996], which has the advantage that the parabolicity of the linear equation can be restored. The same strategy has been applied in similar settings in [Daskalopoulos and Hamilton 1998; Koch 1999; Kienzler 2016; John 2015; Denzler et al. 2015; Seis 2015].

We define

$$d(z, z') = \frac{|z - z'|}{\sqrt{\rho(z)} + \sqrt{\rho(z')} + \sqrt{|z - z'|}}$$

for any $z, z' \in B_1(0)$. Notice that d is not a metric as it lacks a proper triangle inequality. This semidistance is in fact equivalent to the geodesic distance induced on the (weighted) Riemannian manifold associated with the heat flow (8); see [Seis 2015, Proposition 4.2]. We define open balls with respect to the metric d by

$$B_r^d(z) = \{z' \in \overline{B_1(0)} : d(z, z') < r\},$$

and set $Q_r^d(z) = (\frac{1}{2}r^4, r^4) \times B_r^d(z)$ and also $Q(T) = (T, T + 1) \times B_1(0)$. Properties of intrinsic balls and volumes will be cited in Section 2.1 below.

With these preparations, we are in the position to introduce the (semi)norms

$$\begin{aligned} \|w\|_{X(p)} &= \sum_{(\ell, k, |\beta|) \in \mathcal{E}} \sup_{\substack{z \in \overline{B_1(0)} \\ 0 < r \leq 1}} \frac{r^{4k + |\beta| - 1}}{\theta(r, z)^{2\ell - |\beta| + 1}} |Q_r^d(z)|^{-1/p} \|\rho^\ell \partial_t^k \partial_z^\beta w\|_{L^p(Q_r^d(z))} \\ &\quad + \sum_{(\ell, k, |\beta|) \in \mathcal{E}} \sup_{T \geq 1} T \|\rho^\ell \partial_t^k \partial_z^\beta w\|_{L^p(Q(T))}, \\ \|f\|_{Y(p)} &= \sup_{\substack{z \in \overline{B_1(0)} \\ 0 < r \leq 1}} \frac{r^3}{\theta(r, z)} |Q_r^d(z)|^{-1/p} \|f\|_{L^p(Q_r^d(z))} + \sup_{T \geq 1} T \|f\|_{L^p(Q(T))} \end{aligned}$$

for $p \geq 1$, where

$$\mathcal{E} = \{(0, 1, 0), (0, 0, 2), (1, 0, 3), (2, 0, 4)\}.$$

These norms (or Whitney measures) induce the function spaces $X(p)$ and $Y(p)$, respectively, in the obvious way.

1.4. Statement of the results. In view of the particular form of the nonlinearity it is apparent that any well-posedness theory for the transformed equation (6) requires an appropriate control of w and ∇w to prevent the denominators in $R_i[w]$ from degenerating. This is achieved when the Lipschitz norm $\|w\|_{W^{1,\infty}} = \|w\|_{L^\infty} + \|\nabla w\|_{L^\infty}$ is sufficiently small. A suitable function space for existence and uniqueness is provided by $X(p) \cap L^\infty(W^{1,\infty})$ under minimal assumptions on the initial data. Here we have used the convention that $L^q(X) = L^q((0, T); X)$ for some (possibly infinite) $T > 0$. Our first main result is:

Theorem 1 (existence and uniqueness). *Let $p > N + 4$ be given. There exists $\varepsilon, \varepsilon_0 > 0$ such that for every $g \in W^{1,\infty}$ with*

$$\|g\|_{W^{1,\infty}} \leq \varepsilon_0$$

there exists a solution w to the nonlinear equation (6) with initial datum g and w is unique among all solutions with $\|w\|_{L^\infty(W^{1,\infty})} + \|w\|_{X(p)} \leq \varepsilon$. Moreover, this solution satisfies the estimate

$$\|w\|_{L^\infty(W^{1,\infty})} + \|w\|_{X(p)} \lesssim \|g\|_{W^{1,\infty}}.$$

Theorem 1 contains the first (conditional) uniqueness result for the multidimensional thin-film equation in a neighborhood of self-similar solutions. Since any solution to the thin-film equation is expected to converge towards the self-similar Smyth–Hill solution, our result can be considered as a uniqueness result for large times. Notice that the smallness of the Lipschitz norm of w can be translated back into the closeness of u to the stationary u_* . Indeed, $\|w\|_{W^{1,\infty}} \ll 1$ can be equivalently expressed as

$$\|u - u_*\|_{L^\infty(\mathcal{P}(u))} + \|\nabla u + x\|_{L^\infty(\mathcal{P}(v))} \ll 1$$

if $\mathcal{P}(u) = \{u > 0\}$ is the positivity set of u . Recall that $\nabla u_* = -x$ inside $B_1(0)$.

Our second result addresses the regularity of the unique solution found above.

Theorem 2. *Let w be the solution from [Theorem 1](#). Then w is smooth and analytic in time and angular direction.*

It is clear that the smoothness of w immediately translates into the smoothness of u up to the boundary of its support. Moreover, the analyticity result particularly implies the analyticity of the level sets of u . Indeed, the level set of u at height $\lambda \geq 0$ is given by

$$\left\{ (t, x) : w(t, r, \phi) = \sqrt{r^2 + \frac{2\sqrt{\lambda}}{\gamma}} - 1 \right\}$$

if r and ϕ are the radial and angular coordinates, respectively. As a consequence, the temporal and tangential analyticity of w translates into the analyticity of the level sets of u . Notice that the zero-level set is nothing but the free boundary $\partial\{u = 0\}$, and thus, [Theorem 2](#) proves the analyticity of the free boundary of solutions near self-similarity.

In the forthcoming paper [[Seis \$\geq\$ 2018](#)], we will use the gained regularity in time for a construction of invariant manifolds that characterize the large-time asymptotic behavior at any order.

The proof of existence and uniqueness in [Theorem 1](#) follows from a fixed-point argument and a maximal regularity theory for the linear equation. Analyticity and regularity are essentially consequences of an argument first introduced by Angenent [[1990](#)] and later improved by Koch and Lamm [[2012](#)].

1.5. Notation. One word about constants. In the major part of the subsequent analysis, we will not keep track of constants in inequalities but prefer to use the sloppy notation $a \lesssim b$ if $a \leq Cb$ for some universal constant C . Sometimes, however, we have to include constants like $e^{\pm Ct}$ when dealing with exponential growth or decay rates. In such cases, C will always be a positive constant which is generic in the sense that it will not depend on t or z , for instance. This constant might change from line to line, which allows us to write things like $e^{2Ct} \lesssim e^{Ct}$ even for large t .

2. The linear problem

Our goal in this section is the study of the initial value problem for the linear equation ([7](#)). In fact, our analysis also applies to the slightly more general equation

$$\begin{cases} \partial_t w + \mathcal{L}_\sigma^2 w + \mathcal{L}_\sigma w = f & \text{in } (0, \infty) \times B_1(0), \\ w(0, \cdot) = g & \text{in } B_1(0), \end{cases} \tag{9}$$

where \mathcal{L}_σ corresponds to the linearized porous-medium equation considered in [Seis 2014; 2015], defined by

$$\mathcal{L}_\sigma w = -\rho^{-\sigma} \nabla \cdot (\rho^{\sigma+1} \nabla w) = -\rho \Delta w + (\sigma + 1)z \cdot \nabla w$$

for any smooth function $w \in C^\infty(\bar{B})$, and $n \geq 0$ is arbitrary. The constant σ is originally chosen greater than -1 , but we will restrict our attention to the case $\sigma > 0$ for convenience.

Our notion of a weak solution is the following:

Definition 3 (weak solution). Let $0 < T \leq \infty$ and $f \in L^2((0, T); L^2_\sigma)$, $g \in L^2_\sigma$. We call w a weak solution to (9) if $w \in L^2((0, T); L^2_\sigma)$ with $\mathcal{L}w \in L^2((0, T); L^2_\sigma)$ solves

$$\begin{aligned} - \int_0^T \int \partial_t \zeta w \, d\mu_\sigma \, dt + \int_0^T \int \mathcal{L}_\sigma \zeta \mathcal{L}_\sigma w \, d\mu_\sigma \, dt + n \int_0^T \int \nabla \zeta \cdot \nabla w \, d\mu_{\sigma+1} \, dt \\ = \int_0^T \int \zeta f \, d\mu_\sigma \, dt + \int \zeta(0, \cdot) g \, d\mu_\sigma \end{aligned} \tag{10}$$

for all $\zeta \in C^\infty([0, \infty) \times \overline{B_1(0)})$ with $\text{spt } \zeta \subset [0, T) \times \overline{B_1(0)}$.

Here we have used the notation L^2_σ for Lebesgue space $L^2(\mu_\sigma)$ if μ_σ is the absolute continuous measure defined by

$$d\mu_\sigma = \rho^\sigma \, dx.$$

The Hilbert-space theory for (9) is relatively easy and will be developed in Section 2.3 below. In order to perform a perturbation argument on the nonlinear equation, however, we need to control the solution w in the Lipschitz norm. The function spaces $X(p)$ and $Y(p)$ introduced earlier are suitable for such an argument. In fact, our objective in this section is the following result for the linear equation (9).

Theorem 4. *Let $p > N + 4$ be given. Assume that $g \in W^{1,\infty}$. Then there exists a unique weak solution w to (9), and this solution satisfies the a priori estimate*

$$\|w\|_{W^{1,\infty}} + \|w\|_{X(p)} \lesssim \|f\|_{Y(p)} + \|g\|_{W^{1,\infty}}.$$

As mentioned earlier, a change from the Euclidean distance to a Carnot–Carathéodory distance suitable for the second-order operator \mathcal{L}_σ will be crucial for our subsequent analysis. In the following subsection we will recall some basic properties of the corresponding intrinsic volumes and balls, which were derived earlier in [Seis 2015]. Section 2.2 intends to provide some tools that allow us to switch from the spherical setting to the Cartesian one. Energy estimates are established in Section 2.3. In Section 2.4 we treat the homogeneous problem and derive Gaussian estimates. A bit of Calderón–Zygmund theory is provided in Section 2.5. Finally, Section 2.6 contains the theory for the inhomogeneous equation.

2.1. Intrinsic balls and volumes. In the following, we will collect some definitions and properties that are related to our choice of geometry and that will become relevant in the subsequent analysis of the linearized equation. Details and derivations can be found in [Seis 2015, Chapter 4].

It can be shown that the intrinsic balls $B_r^d(z)$ are equivalent to Euclidean balls in the sense that there exists a constant $C < \infty$ such that

$$B_{C^{-1}r\theta(r,z)}(z) \subset B_r^d(z) \subset B_{Cr\theta(r,z)}(z) \tag{11}$$

for any $z \in \overline{B_1(0)}$. Here θ is defined by

$$\theta(r, z) = r \vee \sqrt{\rho(z)}.$$

For the local estimates, it will be crucial to notice that

$$\sqrt{\rho(z_0)} \lesssim r \implies \sqrt{\rho(z)} \lesssim r \quad \text{for all } z \in B_r^d(z_0), \tag{12}$$

$$\sqrt{\rho(z_0)} \gg r \implies \rho(z) \sim \rho(z_0) \quad \text{for all } z \in B_r^d(z_0). \tag{13}$$

In particular, it holds that $\theta(r, \cdot) \sim \theta(r, z_0)$ in $B_r^d(z_0)$. Moreover, if $\sqrt{\rho(z_0)} \lesssim r$, and $z_0 \neq 0$, then (11) implies

$$B_{C^{-1}r^2}\left(\frac{z_0}{|z_0|}\right) \cap \overline{B_1(0)} \subset B_r^d(z_0) \subset B_{Cr^2}\left(\frac{z_0}{|z_0|}\right) \cap \overline{B_1(0)}. \tag{14}$$

We will sometimes write $|A|_\sigma = \mu_\sigma(A)$ for measurable sets A . The volume of an intrinsic ball can be calculated as follows:

$$|B_r^d(z)|_\sigma \sim r^N \theta(r, z)^{N+2\sigma}. \tag{15}$$

In particular, it holds that

$$\frac{|B_r^d(z)|_\sigma}{|B_r^d(z')|_\sigma} \lesssim \left(1 + \frac{d(z, z')}{r}\right)^{2N+2\sigma}. \tag{16}$$

2.2. Preliminary results. By the symmetry of \mathcal{L}_σ and the Cauchy–Schwarz inequality in L^2_σ , we have the interpolation

$$\|\nabla \zeta\|_{\sigma+1}^2 = \int \zeta \mathcal{L}_\sigma \zeta \, d\mu_\sigma \leq \|\zeta\|_\sigma \|\mathcal{L}_\sigma \zeta\|_\sigma. \tag{17}$$

For further reference, we also quote the maximal regularity estimate

$$\|\nabla w\|_\sigma + \|\nabla^2 w\|_{\sigma+2} \lesssim \|\mathcal{L}_\sigma w\|_\sigma; \tag{18}$$

see [Seis 2015, Lemma 4.6].

Close to the boundary, the operator \mathcal{L}_σ can be approximated with the linear operator studied in [Kienzler 2016],

$$\tilde{\mathcal{L}}_\sigma w = -z_N^{-\sigma} \nabla \cdot (z_N^{\sigma+1} \nabla w) = -z_N \Delta w - (\sigma + 1) \partial_N w.$$

This operator is considered on the half-space $\mathbb{R}_+^N = \{z \in \mathbb{R}^N : z > 0\}$. Defining $\tilde{\mu}_\sigma = z_N^\sigma dz$ for any $\sigma > -1$ and using the notation $\|\cdot\|_\sigma$ for the norm on $L^2(\tilde{\mu}_\sigma)$ (slightly abusing notation), we have, analogously to (18), that

$$\|\nabla w\|_\sigma + \|\nabla^2 w\|_{\sigma+2} \lesssim \|\tilde{\mathcal{L}}_\sigma w\|_\sigma. \tag{19}$$

Our first lemma shows how the second-order elliptic equation can be transformed onto a problem on the half-space.

Lemma 5. *Suppose that*

$$\mathcal{L}_\sigma w = \xi$$

for some w such that $\text{spt}(w) \subset \overline{B_1(0)} \cap B_\varepsilon(e_N)$ for some $\varepsilon > 0$. Let $\Phi(z) = \sqrt{1 - |z'|^2}e_N - z$ for $z \in \overline{B_1(0)}$. If ε is sufficiently small, then Φ is a diffeomorphism on $\overline{B_1(0)} \cap B_\varepsilon(e_N)$. Moreover, \tilde{w} defined by $\tilde{w}(\Phi(z)) = w(z)$ solves the equation

$$\tilde{\mathcal{L}}_\sigma \tilde{w} = \tilde{\xi},$$

where $A = A(\tilde{z}) \in \mathbb{R}^{N \times N}$ and $b = b(\tilde{z}) \in \mathbb{R}^N$ are smooth functions with

$$\tilde{\xi} = \xi \circ \Phi^{-1} + A : \tilde{\nabla}^2 \tilde{w} + b \cdot \tilde{\nabla} \tilde{w}$$

with $|A(\tilde{z})| \lesssim |\tilde{z}|^2$, $|b(\tilde{z})| \lesssim |\tilde{z}|$.

Proof. It is clear that Φ is a diffeomorphism from a small ball around $z = e_N$ into a small neighborhood around the origin in \mathbb{R}_+^N . Moreover, a direct calculation and Taylor expansion show that

$$\begin{aligned} \Delta w &= \tilde{\Delta} \tilde{w} + \frac{2\tilde{z}'}{\sqrt{1 - |\tilde{z}'|^2}} \cdot \tilde{\nabla}' \tilde{\partial}_N \tilde{w} + \frac{|\tilde{z}'|^2}{1 - |\tilde{z}'|^2} \tilde{\partial}_N^2 \tilde{w} - \left(\frac{N - 1}{\sqrt{1 - |\tilde{z}'|^2}} + \frac{|\tilde{z}'|^2}{(1 - |\tilde{z}'|^2)^{3/2}} \right) \tilde{\partial}_N \tilde{w} \\ &= \tilde{\Delta} \tilde{w} + A_1(\tilde{z}) : \tilde{\nabla}^2 \tilde{w} + b_1(\tilde{z}) \cdot \tilde{\nabla} \tilde{w}, \end{aligned}$$

$$\begin{aligned} z \cdot \nabla w &= -\sqrt{1 - |\tilde{z}'|^2} \tilde{\partial}_N \tilde{w} + \tilde{z}' \cdot \tilde{\nabla}' \tilde{w} + \left(\tilde{z}_N - \frac{|\tilde{z}'|^2}{\sqrt{1 - |\tilde{z}'|^2}} \right) \tilde{\partial}_N \tilde{w} \\ &= -\tilde{\partial}_N \tilde{w} + b_2(\tilde{z}) \cdot \tilde{\nabla} \tilde{w}, \end{aligned}$$

$$\rho(z) = \sqrt{1 - |\tilde{z}'|^2} \tilde{z}_N - \frac{1}{2} \tilde{z}_N^2 = \tilde{z}_N + c(\tilde{z}),$$

where $|A_1(\tilde{z})| \lesssim |\tilde{z}|$, $|b_1(\tilde{z})| \lesssim 1$, $|b_2(\tilde{z})| \lesssim |\tilde{z}|$, $|c(\tilde{z})| \lesssim |\tilde{z}|^2$. We easily infer the statement. □

A helpful tool in the derivation of the L_σ^2 maximal regularity estimates for our parabolic problem (9) will be the following estimate for the Cartesian problem.

Lemma 6. *Suppose \tilde{w} is a smooth solution of the equation*

$$\tilde{\mathcal{L}}_\sigma \tilde{w} = \tilde{\xi}$$

for some smooth $\tilde{\xi}$. Then

$$\|\tilde{\nabla}^2 \tilde{w}\|_\sigma + \|\tilde{\nabla}^3 \tilde{w}\|_{\sigma+2} + \|\tilde{\nabla}^4 \tilde{w}\|_{\sigma+4} \lesssim \|\tilde{\nabla} \tilde{\xi}\|_\sigma + \|\tilde{\nabla}^2 \tilde{\xi}\|_{\sigma+2}.$$

Proof. We start with the derivation of higher-order tangential regularity. Since $\tilde{\mathcal{L}}_\sigma$ commutes with $\tilde{\partial}_i$ for any $i \in \{1, \dots, N - 1\}$, differentiation yields $\tilde{\mathcal{L}}_\sigma \tilde{\partial}_i \tilde{w} = \tilde{\partial}_i \tilde{\xi}$, and thus via (19),

$$\|\tilde{\nabla} \tilde{\partial}_i \tilde{w}\|_\sigma + \|\tilde{\nabla}^2 \tilde{\partial}_i \tilde{w}\|_{\sigma+2} \lesssim \|\tilde{\nabla} \tilde{\xi}\|_\sigma.$$

We take second-order derivatives in tangential direction and rewrite the resulting equation as $\tilde{\mathcal{L}}_{\sigma+2}\tilde{\partial}_{ij}\tilde{w} = \tilde{\partial}_{ij}\tilde{\xi} - 2\tilde{\partial}_{ijN}\tilde{w}$, where $i, j \in \{1, \dots, N-1\}$. From the above estimate and (19) we obtain

$$\|\tilde{\nabla}\tilde{\partial}_{ij}\tilde{w}\|_{\sigma+2} + \|\tilde{\nabla}^2\tilde{\partial}_{ij}\tilde{w}\|_{\sigma+4} \lesssim \|\tilde{\nabla}\tilde{\xi}\|_{\sigma} + \|\tilde{\nabla}^2\tilde{\xi}\|_{\sigma+2}.$$

Transversal derivatives do not commute with $\tilde{\mathcal{L}}_{\sigma}$. Instead, it holds that $\tilde{\partial}_N\tilde{\mathcal{L}}_{\sigma} = \tilde{\mathcal{L}}_{\sigma+1}\tilde{\partial}_N - \tilde{\Delta}'$. A double differentiation in transversal direction yields thus $\tilde{\mathcal{L}}_{\sigma+2}\tilde{\partial}_N^2\tilde{w} = \tilde{\partial}_N^2\tilde{\xi} + 2\tilde{\Delta}'\tilde{\partial}_N\tilde{w}$. We invoke (19) and obtain with the help of the previous estimate

$$\|\tilde{\nabla}\tilde{\partial}_N^2\tilde{w}\|_{\sigma+2} + \|\tilde{\nabla}^2\tilde{\partial}_N^2\tilde{w}\|_{\sigma+4} \lesssim \|\tilde{\nabla}\tilde{\xi}\|_{\sigma} + \|\tilde{\nabla}^2\tilde{\xi}\|_{\sigma+2}.$$

Finally, the control of $\tilde{\partial}_N^2\tilde{w}$ follows by using the transversally differentiated equation in the sense of $(\sigma+2)\tilde{\partial}_N^2\tilde{w} = -\tilde{\partial}_N\tilde{\xi} - \tilde{z}_N\tilde{\Delta}\tilde{\partial}_N\tilde{w} - \tilde{\Delta}'\tilde{w}$ and the previous bounds. \square

2.3. Energy estimates. In this subsection, we derive the basic well-posedness result, maximal regularity estimates and local estimates in the Hilbert-space setting. We start with existence and uniqueness.

Lemma 7. *Let $0 < T \leq \infty$ and $f \in L^2((0, T); L^2_{\sigma})$, $g \in L^2_{\sigma}$. Then there exists a unique weak solution to (9). Moreover, it holds that*

$$\sup_{(0, T)} \|w\|_{\sigma}^2 + \int_0^T \|\nabla w\|_{\sigma}^2 dt + \int_0^T \|\nabla^2 w\|_{\sigma+2}^2 dt \lesssim \int_0^T \|f\|_{\sigma}^2 dt + \|g\|_{\sigma}^2.$$

Proof. Existence of weak solutions can be proved, for instance, by using an implicit Euler scheme. Indeed, thanks to (18), it is easily seen that for any $h > 0$ and $f \in L^2_{\sigma}$ the elliptic problem

$$\frac{1}{h}w + (\mathcal{L}_{\sigma}^2 + n\mathcal{L}_{\sigma})w = f$$

has a unique solution w satisfying $\mathcal{L}_{\sigma}w \in L^2_{\sigma}$; see [Seis 2014, Appendix] for the analogous second-order problem. This solution satisfies the a priori estimate

$$\frac{1}{\sqrt{h}}\|w\|_{\sigma} + \|\mathcal{L}_{\sigma}w\|_{\sigma} \lesssim \|f\|_{\sigma}.$$

With these insights, it is an exercise to construct time-discrete solutions to (9), and standard compactness arguments allow for passing to the limit, both in the equation and in the estimate. In view of the linearity of the equation, uniqueness follows immediately. \square

Our next result is a maximal regularity estimate for the homogeneous problem.

Lemma 8. *Let w be a solution to the initial value problem (9) with $g = 0$ and $f \in L^2((0, T); L^2_{\sigma})$ for some $0 < T \leq \infty$. Then the mappings $t \mapsto \|w(t)\|_{\sigma}$ and $t \mapsto \|\nabla w(t)\|_{\sigma+1}$ are continuous on $[0, T]$ and $\nabla^2 w, \nabla^3 w, \rho\nabla^4 w \in L^2((0, T); L^2_{\sigma})$ with*

$$\|\partial_t w\|_{L^2(L^2_{\sigma})} + \|\nabla^2 w\|_{L^2(L^2_{\sigma})} + \|\nabla^3 w\|_{L^2(L^2_{\sigma+2})} + \|\nabla^4 w\|_{L^2(L^2_{\sigma+4})} \lesssim \|f\|_{L^2(L^2_{\sigma})}.$$

In the statement of the lemma, we have written $L^2(L^2_{\sigma})$ for $L^2((0, T); L^2_{\sigma})$.

Proof. We perform a quite formal argument that can be made rigorous by using the customary approximation procedures. Choosing $\zeta = \chi_{[t_1, t_2]} \partial_t w$ as a test function in (10), we have the identity

$$\int_{t_1}^{t_2} \|\partial_t w\|_\sigma^2 dt + \frac{1}{2} \|\mathcal{L}_\sigma w(t_2)\|_\sigma^2 + \frac{1}{2} n \|\nabla w(t_2)\|_{\sigma+1} = \int_{t_1}^{t_2} \int f \partial_t w d\mu_\sigma dt + \frac{1}{2} \|\mathcal{L}_\sigma w(t_1)\|_\sigma^2 + \frac{1}{2} n \|\nabla w(t_1)\|_{\sigma+1}.$$

Combining this bound with the estimate from Lemma 7, we deduce that the mappings $t \mapsto \|w(t)\|_\sigma$ and $t \mapsto \|\nabla w(t)\|_{\sigma+1}$ are continuous. Moreover,

$$\int_0^T \|\partial_t w\|_\sigma^2 dt \lesssim \int_0^T \|f\|_\sigma^2 dt$$

because $g = 0$. A similar estimate holds for $\mathcal{L}_\sigma w$ by virtue of Lemma 7, and the statement thus follows upon proving

$$\|\nabla^2 w\|_\sigma + \|\nabla^3 w\|_{\sigma+2} + \|\nabla^4 w\|_{\sigma+4} \lesssim \|\xi\|_\sigma + \|\nabla \xi\|_\sigma + \|\nabla^2 \xi\|_{\sigma+2} \tag{20}$$

for any solution of the elliptic problem $\mathcal{L}_\sigma w = \xi$, because the right-hand side is bounded by $\|f\|_\sigma$ thanks to (18) and Lemma 7. It is not difficult to obtain estimates in the interior of $B_1(0)$. For instance, since

$$\mathcal{L}_{\sigma+2} \partial_i w = \partial_i \xi - z_i \Delta w + 2z \cdot \nabla \partial_i w - (\sigma + 1) \partial_i w$$

is bounded in $L^2_{\sigma+2}$ and because $L^2_\sigma \subset L^2_{\sigma+2}$, an application of (18) yields

$$\|\nabla^2 w\|_{\sigma+2} + \|\nabla^3 w\|_{\sigma+4} \lesssim \|\partial_i \xi\|_\sigma + \|\nabla w\|_\sigma + \|\nabla^2 w\|_{\sigma+2} \lesssim \|\xi\|_\sigma + \|\nabla \xi\|_\sigma. \tag{21}$$

Since $\rho \sim 1$ in the interior of $B_1(0)$, this estimate gives the desired control of the second- and third-order derivatives in the interior of $B_1(0)$. Fourth-order derivatives can be estimated similarly.

To derive estimates at the boundary of $B_1(0)$, it is convenient to locally flatten the boundary. For this purpose, we localize the equation with the help of a smooth cut-off function η that is supported in a small ball centered at a given boundary point, say e_N ,

$$\mathcal{L}_\sigma(\eta w) = \eta \xi - 2\rho \nabla \eta \cdot \nabla w - \rho \Delta \eta w + (\sigma + 1) z \cdot \nabla \eta w =: \tilde{\xi}.$$

A short computation shows that

$$\|\tilde{\xi}\|_\sigma + \|\nabla \tilde{\xi}\|_\sigma + \|\nabla^2 \tilde{\xi}\|_{\sigma+2} \lesssim \|\xi\|_\sigma + \|\nabla \xi\|_\sigma + \|\nabla^2 \xi\|_{\sigma+2},$$

where we have used (18) and (21) and the Hardy–Poincaré inequality $\|w\|_\sigma \lesssim \|\nabla w\|_{\sigma+1}$ from [Seis 2014, Lemma 3]. (For this, notice that we can assume that w has zero average because solutions to $\mathcal{L}_\sigma w = \xi$ are unique up to constants.) Establishing (20) for this localized equation is now a straightforward calculation based on the transformation from Lemma 5 and the a priori estimate from Lemma 6. A covering argument concludes the proof. \square

A crucial step in the derivation of the Gaussian estimates is the following local estimate.

Lemma 9. *Let $0 < \hat{\varepsilon} < \varepsilon < 1$ and $0 < \delta < \hat{\delta} < 1$ be given. Let w be a solution to the inhomogeneous equation (9). Then the following holds for any $z_0 \in \overline{B_1(0)}$, $\tau \geq 0$ and $0 < r \lesssim 1$:*

$$\begin{aligned} & \iint_Q (\partial_t w)^2 d\mu_\sigma dt + \frac{\theta(r, z_0)^4}{r^4} \iint_Q |\nabla^2 w|^2 d\mu_\sigma dt \\ & \quad + \frac{\theta(r, z_0)^2}{r^2} \iint_Q |\nabla^3 w|^2 d\mu_{\sigma+2} dt + \iint_Q |\nabla^4 w|^2 d\mu_{\sigma+4} dt \\ & \lesssim \iint_{\hat{Q}} f^2 d\mu_\sigma dt + \frac{1}{r^8} \iint_{\hat{Q}} w^2 + r^2 \theta(r, z_0)^2 |\nabla w|^2 d\mu_\sigma dt, \end{aligned}$$

where $Q = (\tau + \varepsilon r^2, \tau + r^2) \times B_{\delta r}^d(z_0)$ and $\hat{Q} = (\tau + \hat{\varepsilon} r^2, \tau + r^2) \times B_{\hat{\delta} r}^d(z_0)$.

Proof. Because (9) is invariant under time shifts, we may set $\tau = 0$. We start by recalling that

$$\mathcal{L}_\sigma(\eta w) = \eta \mathcal{L}_\sigma w - 2\rho \nabla \eta \cdot \nabla w + (\mathcal{L}_\sigma \eta) w$$

for any two functions η and w , and thus, via iteration,

$$\mathcal{L}_\sigma^2(\eta w) = \eta \mathcal{L}_\sigma^2 w - 2\rho \nabla \eta \cdot \nabla \mathcal{L}_\sigma w + \mathcal{L}_\sigma \eta \mathcal{L}_\sigma w - 2\mathcal{L}_\sigma(\rho \nabla \eta \cdot \nabla w) + \mathcal{L}_\sigma((\mathcal{L}_\sigma \eta) w).$$

In the sequel, we will choose η as a smooth cut-off function that is supported in the intrinsic space-time cylinder \hat{Q} , and is constantly 1 in the smaller cylinder Q . For such cut-off functions, it holds that $|\partial_t^k \partial_z^\beta \eta| \lesssim r^{-2k-|\beta|} \theta(r, z_0)^{-|\beta|}$. (Here and in the following, the dependency on $\varepsilon, \hat{\varepsilon}, \delta$ and $\hat{\delta}$ is neglected in the inequalities.) Then ηw solves the equation

$$\begin{aligned} \partial_t(\eta w) + \mathcal{L}_\sigma^2(\eta w) + n \mathcal{L}_\sigma(\eta w) &= \eta f + \partial_t \eta w - 4\mathcal{L}_\sigma(\rho \nabla \eta \cdot \nabla w) - 4\rho \nabla(\rho \nabla \eta) : \nabla^2 w \\ &\quad - 2\rho z \cdot \nabla \eta \Delta w - 2(\sigma + 1 + n)\rho \nabla \eta \cdot \nabla w + 2\mathcal{L}_\sigma(\rho \nabla \eta) \cdot \nabla w \\ &\quad + \mathcal{L}_\sigma((\mathcal{L}_\sigma \eta) w) + \mathcal{L}_\sigma \eta \mathcal{L}_\sigma w + n(\mathcal{L}_\sigma \eta) w. \end{aligned}$$

For simplicity, we denote the right-hand side by \tilde{f} . Testing against ηw and using the symmetry and nonnegativity properties of \mathcal{L}_σ and the fact that $\eta w = 0$ initially, we obtain the estimate

$$\iint (\mathcal{L}_\sigma(\eta w))^2 d\mu_\sigma dt \leq \iint \eta w \tilde{f} d\mu_\sigma dt.$$

A tedious but straightforward computation then yields

$$\begin{aligned} & \left| \int \eta w \tilde{f} d\mu_\sigma \right| \\ & \lesssim \left(r^2 \|\chi f\|_\sigma + \|\mathcal{L}_\sigma(\eta w)\|_\sigma + \frac{1}{r^2} \|\chi w\|_\sigma + \frac{\theta_0}{r} \|\chi \nabla w\|_\sigma \right) \left(\frac{1}{r^2} \|\chi w\|_\sigma + \frac{\theta_0}{r} \|\chi \nabla w\|_\sigma \right), \end{aligned} \tag{22}$$

where $\chi = \chi_{\text{spt}(\eta)}$ and $\theta_0 = \theta(r, z_0)$, which in turn implies

$$\begin{aligned} & \iint_Q |\nabla w|^2 d\mu_\sigma dt + \iint_Q |\nabla^2 w|^2 d\mu_{\sigma+2} dt \\ & \lesssim r^4 \iint_{\hat{Q}} f^2 d\mu_\sigma dt + \frac{1}{r^4} \iint_{\hat{Q}} w^2 + r^2 \theta_0^2 |\nabla w|^2 d\mu_\sigma dt \end{aligned} \tag{23}$$

via (18) and Young’s inequality. We will show the argument for (22) for the leading-order terms only. For instance, from the symmetry of \mathcal{L}_σ and the fact that $|\rho \nabla \eta| \lesssim \theta_0/r$, we deduce that

$$\left| \int \eta w \mathcal{L}_\sigma(\rho \nabla \eta \cdot \nabla w) \, d\mu_\sigma \right| \lesssim \frac{\theta_0}{r} \|\mathcal{L}_\sigma(\eta w)\|_\sigma \|\chi \nabla w\|_\sigma.$$

Similarly, by integration by parts we calculate

$$\left| \int \eta w \rho \nabla(\rho \nabla \eta) : \nabla^2 w \, d\mu_\sigma \right| \lesssim \frac{\theta_0^2}{r^2} \int |\nabla(\eta w)| |\nabla w| \, d\mu_\sigma + \frac{1}{r^2} \int \eta |w| |\nabla w| \, d\mu_\sigma,$$

and conclude observing that $|\nabla(\eta w)| \lesssim \theta_0^{-1} r^{-1} |w| + |\nabla w|$. The remaining terms of \tilde{f} can be estimated similarly.

To gain control over the third-order derivatives of ηw , we test the equation with $\mathcal{L}_\sigma(\eta w)$. With the help of the symmetry and nonnegativity properties of \mathcal{L}_σ , we obtain the estimate

$$\iint |\nabla \mathcal{L}_\sigma(\eta w)|^2 \, d\mu_{\sigma+1} \, dt \leq \iint \mathcal{L}_\sigma(\eta w) \tilde{f} \, d\mu_\sigma \, dt.$$

We have to find suitable estimates for the inhomogeneity term. Because we have to make use of the previous bound (23), we have to shrink the cylinders Q and \hat{Q} such that the new function η is supported in the set where the old η was constantly 1. We claim that

$$\iint_Q |\nabla \mathcal{L}_\sigma w|^2 \, d\mu_{\sigma+1} \, dt \lesssim r^2 \iint_{\hat{Q}} f^2 \, d\mu_\sigma \, dt + \frac{1}{r^6} \iint_{\hat{Q}} w^2 + r^2 \theta_0^2 |\nabla w| \, d\mu_\sigma \, dt. \tag{24}$$

Again, we will provide the argument for the leading-order terms only. We use the symmetry of \mathcal{L}_σ , the bounds on derivatives of η and the scaling of ρ , see (12) and (13), to estimate

$$\begin{aligned} \left| \int \mathcal{L}_\sigma(\eta w) \mathcal{L}_\sigma(\rho \nabla \eta \cdot \nabla w) \, d\mu_\sigma \right| &= \left| \int \nabla \mathcal{L}_\sigma(\eta w) \cdot \nabla(\rho \nabla \eta \cdot \nabla w) \, d\mu_{\sigma+1} \right| \\ &\lesssim \|\nabla \mathcal{L}_\sigma(\eta w)\|_{\sigma+1} \left(\frac{\theta_0}{r^2} \|\chi \nabla w\|_\sigma + \frac{1}{r} \|\chi \nabla^2 w\|_{\sigma+2} \right). \end{aligned}$$

Similarly,

$$\left| \int \mathcal{L}_\sigma(\eta w) \rho \nabla(\rho \nabla \eta) : \nabla^2 w \, d\mu_\sigma \right| \lesssim \frac{1}{r^2} \|\mathcal{L}_\sigma(\eta w)\|_\sigma \|\chi \nabla^2 w\|_{\sigma+2}.$$

The estimates of the remaining terms have a similar flavor. We deduce (24) with the help of Young’s inequality and (23).

The estimate (24) is beneficial as it allows us to estimate \tilde{f} in L^2_σ . This time, it is enough to study the term that involves the third-order derivatives of w . We rewrite

$$\begin{aligned} \mathcal{L}_\sigma(\rho \partial_i \eta \partial_i w) &= \rho \partial_i \eta \mathcal{L}_\sigma \partial_i w - 2\rho \nabla(\rho \partial_i \eta) \cdot \nabla \partial_i w + \mathcal{L}_\sigma(\rho \partial_i \eta) \partial_i w, \\ \mathcal{L}_\sigma \partial_i w &= \partial_i \mathcal{L}_\sigma w - z_i \Delta w - (\sigma + 1) \partial_i w, \end{aligned}$$

and estimate

$$\begin{aligned} \|\mathcal{L}(\rho\partial_i\eta\partial_i w)\|_\sigma &\lesssim \frac{1}{r\theta_0}\|\chi\mathcal{L}_\sigma\partial_i w\|_{\sigma+2} + \frac{1}{r^2}\|\chi\nabla^2 w\|_{\sigma+2} + \frac{\theta_0}{r^3}\|\chi\nabla w\|_\sigma \\ &\lesssim \frac{1}{r}\|\chi\partial_i\mathcal{L}_\sigma w\|_{\sigma+1} + \frac{1}{r^2}\|\chi\nabla^2 w\|_{\sigma+2} + \frac{\theta_0}{r^3}\|\chi\nabla w\|_\sigma. \end{aligned}$$

Upon redefining Q and \hat{Q} as in the derivation of (24), an application of (23) and (24) then yields

$$\int_{\varepsilon r^2}^{r^2} \|\mathcal{L}(\rho\partial_i\eta\partial_i w)\|_\sigma^2 dt \lesssim \iint_{\hat{Q}} f^2 d\mu_\sigma dt + \frac{1}{r^8} \iint_{\hat{Q}} w^2 + r^2\theta_0^2|\nabla w|^2 d\mu_\sigma dt.$$

The remaining terms of \tilde{f} can be estimated in a similar way. Applying the energy estimate from Lemma 8 to the evolution equation for ηw , we thus deduce

$$\begin{aligned} \iint_Q (\partial_t w)^2 d\mu_\sigma dt + \iint_Q |\nabla^2 w|^2 d\mu_\sigma dt + \iint_Q |\nabla^3 w|^2 d\mu_{\sigma+2} dt + \iint_Q |\nabla^4 w|^2 d\mu_{\sigma+4} dt \\ \lesssim \iint_{\hat{Q}} f^2 d\mu_\sigma dt + \frac{1}{r^8} \iint_{\hat{Q}} w^2 + r^2\theta_0^2|\nabla w|^2 d\mu_\sigma dt. \end{aligned} \quad (25)$$

Notice that the above bound on the second-order derivatives and (23) together imply

$$\frac{\theta_0^4}{r^4} \iint_Q |\nabla^2 w|^2 d\mu_\sigma dt \lesssim \iint_{\hat{Q}} f^2 d\mu_\sigma dt + \frac{1}{r^8} \iint_{\hat{Q}} w^2 + r^2\theta_0^2|\nabla w|^2 d\mu_\sigma dt.$$

Similarly, we can produce the factor θ_0^2/r^2 in front of the integral containing the third-order derivatives. Indeed, because $\nabla\mathcal{L}_\sigma w = \mathcal{L}_{\sigma+1}\nabla w + z\Delta w - \nabla^2 w z + (\sigma + 1)\nabla w$, the bound (24) yields

$$\begin{aligned} \iint_Q |\mathcal{L}_{\sigma+1}\nabla w|^2 d\mu_{\sigma+1} dt \\ \lesssim r^2 \iint_{\hat{Q}} f^2 d\mu_\sigma dt + \frac{1}{r^6} \iint_{\hat{Q}} w^2 + r^2\theta_0^2|\nabla w|^2 d\mu_\sigma dt + \iint_{\hat{Q}} |\nabla^2 w|^2 d\mu_{\sigma+1} dt + \iint_{\hat{Q}} |\nabla w|^2 d\mu_{\sigma+1} dt. \end{aligned}$$

The second-order term on the right-hand side is controlled with the help of the Cauchy–Schwarz inequality, (23) and (25). The first-order term is of higher order as a consequence of (23). It remains to invoke (18) to the effect that

$$\iint_Q |\nabla^3 w|^2 d\mu_{\sigma+3} dt \lesssim r^2 \iint_{\hat{Q}} f^2 d\mu_\sigma dt + \frac{1}{r^6} \iint_{\hat{Q}} w^2 + r^2\theta_0^2|\nabla w|^2 d\mu_\sigma dt.$$

Combining the latter with (25) yields the statement of the lemma. □

2.4. Estimates for the homogeneous equation. In this subsection, we study the initial value problem for the homogeneous equation

$$\begin{cases} \partial_t w + \mathcal{L}_\sigma^2 w + n\mathcal{L}_\sigma w = 0 & \text{in } (0, \infty) \times B_1(0), \\ w(0, \cdot) = g & \text{in } B_1(0). \end{cases} \quad (26)$$

Our first goal is a pointwise higher-order regularity estimate.

Lemma 10. *Let $0 < \varepsilon < 1$ and $0 < \delta < 1$ be given. Let w be a solution to the homogeneous equation (26). If $\varepsilon, \delta \in (0, 1)$ and δ is sufficiently small, then the following holds for any $z_0 \in \overline{B_1(0)}$, $\tau \geq 0$ and $0 < r \lesssim 1$:*

$$|\partial_t^k \partial_z^\beta w(t, z)|^2 \lesssim \frac{r^{-8k-2|\beta|} \theta(r, z_0)^{-2|\beta|}}{r^4 |B_r^d(z_0)|_\sigma} \int_\tau^{\tau+r^4} \int_{B_r^d(z_0)} w^2 + r^2 \theta(r, z_0)^2 |\nabla w|^2 d\mu_\sigma dt$$

for any $(t, z) \in (\tau + \varepsilon r^4, \tau + r^4] \times B_{\delta r}^d(z_0)$.

Proof. The lemma is a consequence of the local higher-order regularity estimate

$$\iint_Q (\partial_t^k \partial_z^\beta w)^2 d\mu_\sigma dt \lesssim r^{-8k-2|\beta|} \theta(r, z_0)^{-2|\beta|} \iint_{\hat{Q}} w^2 + r^2 \theta(r, z_0)^2 |\nabla w|^2 d\mu_\sigma dt, \tag{27}$$

where Q and \hat{Q} are defined as in Lemma 9, and a Morrey estimate in the weighted space $L^2(\mu_\sigma)$; see, e.g., [Seis 2015, Lemma 4.9]. Notice that (27) is trivial for $(k, |\beta|) \in \{(0, 1), (0, 2)\}$. In the following, we write $\theta_0 = \theta(r, z_0)$.

To prove (27) for general choices of k and β , it is convenient to consider separately the two cases $\sqrt{\rho(z_0)} \lesssim r$ and $\sqrt{\rho(z_0)} \gg r$. The second case is relatively simple: Since $\rho \sim \rho(z_0)$ by (13) in both Q and \hat{Q} , we deduce (27) in the cases $(k, |\beta|) \in \{(1, 0), (0, 2), (0, 3), (0, 4)\}$ directly from Lemma 9 (with $f = 0$). To gain control on higher-order derivatives, we differentiate with respect to z_i ,

$$\begin{aligned} \partial_t \partial_i w + \mathcal{L}_\sigma^2 \partial_i w + n \mathcal{L}_\sigma \partial_i w \\ = -z_i \Delta \mathcal{L}_\sigma w - (\sigma + 1) \partial_i \mathcal{L}_\sigma w - \mathcal{L}_\sigma (z_i \Delta w) - (\sigma + 1) \mathcal{L}_\sigma \partial_i w - n z_i \Delta w - n(\sigma + 1) \partial_i w. \end{aligned}$$

Denoting by \tilde{f} the right-hand side of this identity, applying Lemma 9 yields the estimate

$$\iint_Q |\nabla^4 \partial_i w|^2 d\mu_{\sigma+4} dt \lesssim \iint_{\hat{Q}} \tilde{f}^2 d\mu_\sigma dt + \frac{1}{r^8} \iint_{\hat{Q}} (\partial_i w)^2 + r^2 \theta_0^2 |\nabla \partial_i w|^2 d\mu_\sigma dt.$$

We invoke the previously derived bound and the fact that $\rho \sim \rho(z_0)$ to conclude the statement in the case $(k, |\beta|) = (0, 5)$. Higher-order derivatives are controlled similarly via iteration.

The proof in the case $\rho(z_0) \lesssim r$ is lengthy and tedious. As similar results have been recently obtained in [John 2015; Kienzler 2016; Seis 2015] and most the involved tools have been already applied earlier in this paper, we will only outline the argument in the following. Thanks to (14), it is enough to study the situation where $z_0 \in \partial B_1(0)$, and upon shrinking δ , we may assume that Φ constructed in Lemma 5 is a diffeomorphism from $B_{\delta r}^2(z_0)$ onto a subset of the half-space. Under Φ , the homogeneous equation (26) transforms into

$$\partial_t \tilde{w} + \tilde{\mathcal{L}}_\sigma^2 \tilde{w} + n \tilde{\mathcal{L}}_\sigma \tilde{w} = \tilde{f},$$

where \tilde{f} is of higher-order at the boundary. Because $\tilde{\mathcal{L}}_\sigma$ commutes with tangential derivatives $\tilde{\partial}_i$ for $i \in \{1, \dots, N - 1\}$, control on higher-order tangential derivatives is deduced from Lemma 9. To obtain control on vertical derivatives, we recall that $\tilde{\partial}_N \tilde{\mathcal{L}}_\sigma = \tilde{\mathcal{L}}_{\sigma+1} \tilde{\partial}_N - \tilde{\Delta}'$. Arguing as in the proof of Lemma 6 gives the desired estimates. Again, bounds on higher-order derivatives and mixed derivatives are obtained by iteration. □

For the proof of the Gaussian estimates and the Whitney-measure estimates for the homogeneous problem, it is convenient to introduce a family of auxiliary functions $\chi_{a,b} : \overline{B_1(0)} \times \overline{B_1(0)} \rightarrow \mathbb{R}$, given by

$$\chi_{a,b}(z, z_0) = \frac{a\hat{d}(z, z_0)^2}{\sqrt{b^2 + \hat{d}(z, z_0)^2}},$$

where $a, b \in \mathbb{R}$ are given parameters, and

$$\hat{d}(z, z_0)^2 = \frac{|z - z_0|^2}{\sqrt{\rho(z)^2 + \rho(z_0)^2 + |z - z_0|^2}} \sim d(z, z_0)^2.$$

It can be verified by a short computation that $\rho|\nabla_z \hat{d}^2|^2 \lesssim \hat{d}$ and $\rho|\nabla_z^2 \hat{d}^2| \lesssim 1$, with the consequence that

$$\sqrt{\rho(z)}|\nabla_z \chi_{a,b}(z, z_0)| \lesssim |a|, \tag{28}$$

$$\rho(z)|\nabla_z^2 \chi_{a,b}(z, z_0)| \lesssim \frac{|a|}{|b|} \tag{29}$$

uniformly in $z, z_0 \in \overline{B_1(0)}$. Because g is conformally flat with $g \sim \rho^{-1}(dz)^2$, the gradient ∇_g on (\mathcal{M}, g) obeys the scaling $\nabla_g \sim \rho\nabla$, and thus (28) can be rewritten as $\sqrt{g(\nabla_g \chi, \nabla_g \chi)} \lesssim |a|$ (where we have dropped the indices and z_0). The latter implies that $\chi = \chi_{a,b}(\cdot, z_0)$ is Lipschitz with respect to the intrinsic topology; that is,

$$|\chi_{a,b}(z, z_0) - \chi_{a,b}(z', z_0)| \lesssim |a| d(z, z'). \tag{30}$$

We derive some new weighted energy estimates.

Lemma 11. *Let w be the solution to the homogeneous equation (26). Let $a, b \in \mathbb{R}$ and $z_0 \in \overline{B_1(0)}$ be given. Define $\chi = \chi_{a,b}(\cdot, z_0)$. Then there exists a constant $C > 0$ such that for any $T > 0$ it holds that*

$$\sup_{[0, T]} \int e^{2\chi} w^2 d\mu_\sigma + \int_0^T \int e^{2\chi} |\nabla w|^2 + (\mathcal{L}_\sigma(e^\chi w))^2 d\mu_\sigma dt \lesssim e^{C(a^2/b^2 + a^4)T} \int e^{2\chi} g^2 d\mu_\sigma.$$

Proof. The quantity $e^\chi w$ evolves according to

$$\begin{aligned} & \partial_t(e^\chi w) + \mathcal{L}_\sigma^2(e^\chi w) + n\mathcal{L}_\sigma(e^\chi w) \\ &= -2\rho\nabla e^\chi \cdot \nabla \mathcal{L}_\sigma w + \mathcal{L}_\sigma e^\chi \mathcal{L}_\sigma w - 2\mathcal{L}_\sigma(\rho\nabla e^\chi \cdot \nabla w) + \mathcal{L}_\sigma((\mathcal{L}_\sigma e^\chi)w) - 2n\rho\nabla e^\chi \cdot \nabla w + n(\mathcal{L}_\sigma e^\chi)w. \end{aligned}$$

Denoting the right-hand side by \tilde{f} and testing with $e^\chi w$ yields

$$\frac{d}{dt} \frac{1}{2} \int (e^\chi w)^2 d\mu_\sigma + \int (\mathcal{L}_\sigma(e^\chi w))^2 d\mu_\sigma + n \int |\nabla(e^\chi w)|^2 d\mu_{\sigma+1} = \int e^\chi w \tilde{f} d\mu_\sigma, \tag{31}$$

where we have used once more the symmetry of \mathcal{L}_σ . We claim that the term on the right can be estimated as follows:

$$\int e^\chi w \tilde{f} d\mu_\sigma \lesssim \varepsilon(\|\mathcal{L}_\sigma(e^\chi w)\|_\sigma^2 + \|\nabla(e^\chi w)\|_{\sigma+1}) + \left(1 + \frac{a^2}{b^2} + a^4\right) \|e^\chi w\|_\sigma^2, \tag{32}$$

where ε is some small constant that allows us to absorb the first two terms in the left-hand side of the energy estimate above. Indeed, multiple integrations by parts and the bounds (28) and (29) yield that the left-hand side of (31) is bounded by

$$\begin{aligned} &|a| \|\mathcal{L}_\sigma \zeta\|_\sigma \|\nabla \zeta\|_{\sigma+1} + a^2 \|\nabla \zeta\|_{\sigma+1}^2 + \left(|a| + \frac{a^2}{|b|} + |a|^3\right) \|\nabla \zeta\|_{\sigma+1} \|\zeta\|_\sigma \\ &+ a^2 \|\nabla \zeta\|_{\sigma+1} \|\zeta\|_{\sigma-1} + \left(\frac{|a|}{|b|} + a^2\right) \|\mathcal{L}_\sigma \zeta\|_\sigma \|\zeta\|_\sigma + \left(1 + \frac{a^2}{b^2} + a^4\right) \|\zeta\|_\sigma^2 \\ &+ \left(\frac{a^2}{|b|} + |a|^3\right) \|\zeta\|_{\sigma-1} \|\zeta\|_\sigma + |a| \|\mathcal{L}_\sigma \zeta\|_\sigma \|\zeta\|_{\sigma-1} + a^2 \|\zeta\|_{\sigma-1}^2, \end{aligned}$$

where we have set $\zeta = e^\chi w$. We next claim that

$$\|\zeta\|_{\sigma-1} \lesssim \|\zeta\|_\sigma + \|\nabla \zeta\|_{\sigma+1}. \tag{33}$$

Indeed, recall the Hardy–Poincaré inequality

$$\left\| \zeta - \int \zeta \, d\mu_{\tilde{\sigma}-1} \right\|_{\tilde{\sigma}-1} \lesssim \|\nabla \zeta\|_{\sigma+1},$$

see [Seis 2014, Lemma 3], which holds true for any $\tilde{\sigma} \geq \sigma$, because $\sigma > 0$. In particular, $\|\zeta\|_{\sigma-1} \lesssim \|\nabla \zeta\|_{\sigma+1} + \left| \int \zeta \, d\mu_{\sigma-1} \right|$. Notice that for any $\alpha \in (0, \tilde{\sigma})$, it holds that

$$\left| \int \zeta \, d\mu_{\tilde{\sigma}-1} \right| = \left| \int \zeta \rho^\alpha \, d\mu_{\tilde{\sigma}-1-\alpha} \right| \lesssim \left(\int \zeta^2 \rho \, d\mu_{\tilde{\sigma}-1-\alpha} \right)^{1/2} \lesssim \|\zeta\|_{\tilde{\sigma}-1+\alpha}$$

by Jensen’s inequality because $\mu_{\tilde{\sigma}-1+\alpha}$ is a finite measure. Applying the previous two estimates iteratively yields (33). Hence, combining (33) and the interpolation inequality (17) with the bound on the inhomogeneity and using Young’s inequality yields (32).

Now (31) and (32) imply for ε sufficiently small that

$$\frac{d}{dt} \int (e^\chi w)^2 \, d\mu_\sigma + \int (\mathcal{L}_\sigma(e^\chi w))^2 \, \mu_\sigma \lesssim \left(1 + \frac{a^2}{b^2} + a^4\right) \int (e^\chi w)^2 \, d\mu_\sigma.$$

In view of the bound (18) we have the estimate $\|\nabla(e^\chi w)\|_\sigma \lesssim \|\mathcal{L}_\sigma(e^\chi w)\|_\sigma$. Therefore, invoking the product rule of differentiation

$$\|e^\chi \nabla w\|_\sigma \leq \|\nabla(e^\chi w)\|_\sigma + \|e^\chi w \nabla \chi\|_\sigma \lesssim \|\mathcal{L}_\sigma(e^\chi w)\|_\sigma + |a| \|e^\chi w\|_{\sigma-1}.$$

Observe that (33) and (17) imply

$$|a| \|e^\chi w\|_{\sigma-1} \lesssim (|a| + a^2) \|e^\chi w\|_\sigma + \|\mathcal{L}_\sigma(e^\chi w)\|_\sigma.$$

Combining the previous estimates with a Grönwall argument yields the statement of the lemma. □

The following estimate is a major step towards Gaussian estimates.

Lemma 12. *Let w be the solution to the homogeneous equation (26). Let $a, b \in \mathbb{R}$ be given. Then there exists a constant $C > 0$ such that for all $z, z_0 \in \overline{B_1(0)}$, $0 < r \lesssim 1$, $t \in (\frac{1}{2}r^4, r^4)$, $k \in \mathbb{N}_0$ and $\beta \in \mathbb{N}_0^N$ it holds that*

$$|\partial_t^k \partial_z^\beta w(t, z)| \lesssim \frac{r^{-4k-|\beta|} |\theta(r, z)|^{-|\beta|}}{|B_r^d(z)|_\sigma^{1/2}} e^{C(a^2/b^2+a^4)t-\chi_{a,b}(z,z_0)} \|e^{\chi_{a,b}(\cdot, z_0)} g\|_\sigma.$$

Proof. For simplicity, we write $\chi = \chi_{a,b}(\cdot, z_0)$ and $\theta = \theta(r, z)$. From Lemma 10 (with $z_0 = z$ and $\tau = 0$) we deduce the estimate

$$|\partial_t^k \partial_z^\beta w(t, z)|^2 \lesssim \frac{r^{-8k-2|\beta|} |\theta(r, z)|^{-2|\beta|}}{r^4 |B_r^d(z)|_\sigma} \sup_{B_r^d(z)} e^{-2\chi} \int_0^{r^4} \int_{B_r^d(z)} e^{2\chi} w^2 + r^2 \theta^2 e^{2\chi} |\nabla w|^2 d\mu_\sigma dt \quad (34)$$

for all $t \in (\frac{1}{2}r^4, r^4)$. We first observe that the Lipschitz estimate (30) implies

$$\sup_{B_r^d(z)} e^{-\chi} \lesssim e^{-\chi(z)+a^4 r^4}.$$

To estimate the integral expression in (34), we distinguish the cases $\sqrt{\rho(z)} \leq r$ and $\sqrt{\rho(z)} \geq r$. In the first case, we apply Lemma 11 and obtain

$$r^4 \left(\sup_{[0, r^4]} \int e^{2\chi} w^2 d\mu_\sigma + \int_0^{r^4} \int e^{2\chi} |\nabla w|^2 d\mu_\sigma dt \right) \lesssim r^4 e^{C(a^2/b^2+a^4)r^4} \int e^{2\chi} g^2 d\mu_\sigma$$

for some $C > 0$. In the second case, we only focus on second term, i.e., the gradient term. The argument for the first term remains unchanged. Because $\rho \sim \rho(z)$ in the domain of integration, see (13), it holds that

$$\int_0^{r^4} \int_{B_r^d(z)} r^2 \theta^2 e^{2\chi} |\nabla w|^2 d\mu_\sigma dt \lesssim r^2 a^2 \int_0^{r^4} \int e^{2\chi} w^2 d\mu_\sigma dt + r^2 \int_0^{r^4} \int |\nabla(e^\chi w)|^2 d\mu_{\sigma+1} dt,$$

where we have used (28). By using (17) and Young’s inequality, we further estimate

$$r^2 \int |\nabla(e^\chi w)|^2 d\mu_{\sigma+1} \lesssim r^4 \int (\mathcal{L}_\sigma(e^\chi w))^2 d\mu_\sigma + \int (e^\chi w)^2 d\mu_\sigma,$$

which in turn yields

$$\int_0^{r^4} \int_{B_r^d(z)} r^2 \theta^2 e^{2\chi} |\nabla w|^2 d\mu_\sigma dt \lesssim r^4 (1 + r^2 a^2) e^{C(a^2/b^2+a^4)r^4} \int e^{2\chi} g^2 d\mu_\sigma$$

via Lemma 11. Notice that we can eliminate the factor $r^2 a^2$ in the previous expression upon enlarging the constant C . Substituting the previous bounds into (34) yields the statement of the lemma. \square

For large times, we have exponential decay as established in the lemma that follows.

Lemma 13. *Let w be the solution of the initial value problem (9) with $f = 0$. Then for any $k \in \mathbb{N}_0$, $\beta \in \mathbb{N}_0^N$, $t \geq \frac{1}{2}$ and $z \in \overline{B_1(0)}$ it holds that*

$$\left| \partial_t^k \partial_z^\beta \left(w(t, z) - \int g \mu_\sigma \right) \right| \lesssim e^{-\lambda_1 t} \|\nabla g\|_{\sigma+1}.$$

Proof. The proof is an easy consequence of [Lemma 10](#) and a spectral gap estimate for \mathcal{L}_σ . Indeed, applying [Lemma 10](#) with $t = \tau + \frac{1}{4}$, $\varepsilon = \frac{1}{4}$, $r = 1$ and $\tau \geq \frac{1}{4}$ to $w - c$, where $c = \int w \, d\mu_\sigma$ is a constant of the evolution, we obtain the estimate

$$|\partial_t^k \partial_z^\beta (w(t, z) - c)| \lesssim \int_{t-1/4}^{t+3/4} \int (w - c)^2 + |\nabla w|^2 \, d\mu_\sigma \, dt.$$

Thanks to the Hardy–Poincaré inequality [[Seis 2014](#), Lemma 3] and because $\mu_{\sigma+1} \lesssim \mu_\sigma$, we can drop the term $(w - c)^2$ in the integrand. To prove the statement of the lemma, we thus have to establish the estimate

$$\int_{t-1/4}^{t+3/4} |\nabla w|^2 \, d\mu_\sigma \, dt \lesssim e^{-2\lambda_1 t} \int |\nabla g|^2 \, d\mu_{\sigma+1}. \tag{35}$$

For this purpose, we test the homogeneous equation with w and invoke the symmetry and nonnegativity properties of \mathcal{L}_σ and obtain the energy estimate

$$\frac{d}{dt} \frac{1}{2} \int |\nabla w|^2 \, d\mu_{\sigma+1} + \int (\mathcal{L}_\sigma w)^2 \, d\mu_\sigma \leq 0.$$

On the one hand, integration in time over $[t - \frac{1}{4}, t + \frac{3}{4}]$ and the a priori estimate (18) yield

$$\int_{t-1/4}^{t+3/4} |\nabla w|^2 \, d\mu_\sigma \, dt \lesssim \int |\nabla w(t - \frac{1}{4})|^2 \, d\mu_{\sigma+1}. \tag{36}$$

On the other hand, the smallest nonzero eigenvalue λ_1 of \mathcal{L}_σ yields the spectral gap estimate

$$\int (\mathcal{L}_\sigma w)^2 \, d\mu_\sigma = \int \nabla w \cdot \nabla \mathcal{L}_\sigma w \, d\mu_{\sigma+1} \geq \lambda_1 \int |\nabla w|^2 \, d\mu_{\sigma+1},$$

which we combine with the energy estimate from above to get

$$\int |\nabla w(t - \frac{1}{4})|^2 \, d\mu_{\sigma+1} \lesssim e^{-2\lambda_1 t} \int |\nabla g|^2 \, d\mu_{\sigma+1}.$$

Plugging this estimate into (36) yields (35) as desired. □

We are now in the position to prove the desired maximal regularity estimate for the homogeneous problem. Let us start with the latter.

Proposition 14. *Let w be the solution to the homogeneous equation (26). Then*

$$\begin{aligned} \|w\|_{L^\infty} &\lesssim \|g\|_{L^\infty}, \\ \|w\|_{X(p)} + \|\nabla w\|_{L^\infty} &\lesssim \|\nabla g\|_{L^\infty}. \end{aligned}$$

Proof. Thanks to the exponential decay estimates from [Lemma 13](#), it is enough to focus on the norms for small times, $T \leq 1$. We fix $z_0 \in \overline{B_1(0)}$ for a moment and let $r \lesssim 1$ and $t \in (\frac{1}{2}r^4, r^4)$ be arbitrarily given. As before, we set $\theta_0 = \theta(r, z_0)$. Because $w - g(z_0)$ is a solution to the homogeneous equation with initial value $g - g(z_0)$, an application of [Lemma 12](#) with $a = -\frac{1}{r}$ and $b = r$ yields the estimate

$$|\partial_t^k \partial_z^\beta|_{z=z_0} (w(t, z) - g(z_0))| \lesssim \frac{r^{-4k-|\beta|} \theta_0^{-|\beta|}}{|B_r^d(z_0)|_\sigma^{1/2}} \|e^{\chi_{-1/r, r}(\cdot, z_0)} (g - g(z_0))\|_\sigma. \tag{37}$$

Notice that the function χ drops out in the exponential prefactor because $\chi(z_0, z_0) = 0$. We claim that

$$\|e^{\chi_{-1/r,r}(\cdot, z_0)}(g - g(z_0))\|_\sigma \lesssim \min\{\|g\|_{L^\infty}, r\theta_0\|\nabla g\|_{L^\infty}\}|B_r^d(z_0)|_\sigma^{1/2}. \tag{38}$$

The proof of this estimate has been already displayed earlier; see, e.g., the proof of Proposition 4.2 in [Seis 2014]. For the convenience of the reader, we recall the simple argument. Notice first that $|g(z) - g(z_0)| \lesssim \min\{\|g\|_{L^\infty}, |z - z_0|\|\nabla g\|_{L^\infty}\}$. On every annulus

$$A_j = B_{jr}^{\hat{d}}(z_0) \setminus B_{(j-1)r}^{\hat{d}}(z_0)$$

it holds that $\chi_{-1/r,r}(z, z_0) \leq -(j - 1)/\sqrt{2}$ as can be verified by an elementary computation, and thus, for $s \in \{0, 1\}$, we have

$$\int_{A_j} e^{2\chi_{-1/r,r}(z, z_0)} |z - z_0|^{2s} d\mu_\sigma(z) \lesssim j^{2s} r^{2s} \theta(jr, z_0)^{2s} e^{-\sqrt{2}j} |A_j|_\sigma$$

as a consequence of (11). Clearly, $\theta(jr, z_0) \leq j\theta_0$. We notice that $A_j = \emptyset$ for each $j \gg \frac{1}{r}$. On the other hand, thanks to the volume formula (15), it holds that

$$|A_j|_\sigma \lesssim j^{2(N+\sigma)} |B_r^d(z_0)|_\sigma.$$

It remains to notice that the annuli $\{A_j\}_{j \in \mathbb{N}}$ cover $\overline{B_1(0)}$ and deduce that

$$\|e^{\chi_{-1/r,r}(\cdot, z_0)}|\cdot - z_0|^s\|_\sigma \lesssim (r\theta_0)^s |B_r^d(z_0)|_\sigma^{1/2} \left(\sum_{j \in \mathbb{N}} e^{-\sqrt{2}j} j^\kappa\right)^{1/2}$$

for some $\kappa = \kappa(s) > 0$. Because the series is convergent, we have thus proved the bound in (38).

We now combine (37) and (38) to the effect of

$$\begin{aligned} r^{4k+|\beta|} \theta_0^{|\beta|} |\partial_t^k \partial_z^\beta|_{z=z_0}(w(t, z) - g(z_0)) &\lesssim \|g\|_{L^\infty}, \\ r^{4k+|\beta|-1} \theta_0^{|\beta|-1} |\partial_t^k \partial_z^\beta|_{z=z_0}(w(t, z) - g(z_0)) &\lesssim \|\nabla g\|_{L^\infty}. \end{aligned}$$

We obtain the uniform bounds on w and ∇w in the time interval $[0, 1]$ by setting $(k, |\beta|) = (0, 0)$ in the first and $(k, |\beta|) = (0, 1)$ in the second estimate. (Recall that we use Lemma 13 to extend the estimates to times $t \geq 1$.) To control in $X(p)$, we choose $(k, |\beta|) \in \{(1, 0), (0, 2), (0, 3), (0, 4)\}$, raise the second of the above estimates to the power p and average over $Q_r(z)$. For instance, if $(k, |\beta|) = (0, 2)$, this leads to

$$\frac{r^p}{|Q_r^d(z)|} \iint_{Q_r^d(z)} \theta(r, z_0)^p |\nabla^2 w(t, z_0)|^p dz_0 dt \lesssim \|\nabla g\|_{L^\infty}^p.$$

If view of (12) and (13), it holds that $\theta(r, z_0) \sim \theta(r, z)$ uniformly in $B_r^d(z)$, and thus, from maximizing in r and z we obtain

$$\sup_{\substack{z \in \overline{B_1(0)} \\ 0 < r \lesssim 1}} r \theta(r, z) |Q_r^d(z)|^{-1/p} \|\nabla^2 w\|_{L^p(Q_r^d(z))} \lesssim \|\nabla g\|_{L^\infty}.$$

Higher-order derivatives are bounded analogously. □

Gaussian estimates are contained in the following statement.

Proposition 15. *There exists a unique function $G : (0, \infty) \times \overline{B_1(0)} \times \overline{B_1(0)} \rightarrow \mathbb{R}$ with the following properties:*

- (1) *If w is the solution to the homogeneous equation (26), then for any $k \in \mathbb{N}_0$, $\beta \in \mathbb{N}_0^N$ and $(t, z) \in (0, \infty) \times \overline{B_1(0)}$*

$$\partial_t^k \partial_z^\beta w(t, z) = \int \partial_t^k \partial_z^\beta G(t, z, z') g(z') d\mu_\sigma.$$

- (2) *The function G is symmetric in the last two variables; that is,*

$$G(t, z, z') = G(t, z', z)$$

for all $(t, z, z') \in (0, \infty) \times \overline{B_1(0)} \times \overline{B_1(0)}$.

- (3) *For any $z' \in B_1(0)$, $G' = G(\cdot, \cdot, z')$ solves the homogeneous equation*

$$\partial_t G' + \mathcal{L}_\sigma^2 G' + \mathcal{L}_\sigma G' = 0.$$

Moreover,

$$\rho^\sigma G' \xrightarrow{t \downarrow 0} \delta_{z'} \quad \text{in the sense of distributions.}$$

- (4) *It holds that*

$$|\partial_t^k \partial_z^\beta G(t, z, z')| \lesssim \frac{\sqrt[4]{t}^{-4k-|\beta|} \theta(\sqrt[4]{t}, z)^{-|\beta|}}{|B_{\sqrt[4]{t}}^d(z)|_\sigma^{1/2} |B_{\sqrt[4]{t}}^d(z')|_\sigma^{1/2}} e^{-C(d(z, z')/\sqrt[4]{t})^{4/3}}$$

for all $(t, z, z') \in (0, 1] \times \overline{B_1(0)} \times \overline{B_1(0)}$ and any $k \in \mathbb{N}_0$ and $\beta \in \mathbb{N}_0^N$.

- (5) *It holds that*

$$|\partial_t^k \partial_z^\beta (G(t, z, z') - |B_1(0)|_\sigma^{-1})| \lesssim e^{-\lambda_1 t}$$

for all $(t, z, z') \in [1, \infty) \times \overline{B_1(0)} \times \overline{B_1(0)}$ and any $k \in \mathbb{N}_0$ and $\beta \in \mathbb{N}_0^N$.

The estimates in the fourth statement are usually referred to as ‘‘Gaussian estimates’’.

Remark 16. In the fourth statement we may freely interchange the balls centered at z by balls centered at z' and vice versa. Likewise, we can substitute $\theta(\sqrt[4]{t}, z)$ by $\theta(\sqrt[4]{t}, z')$. This is a consequence of (16).

The proof of this proposition is (almost) exactly the one of [Seis 2014, Proposition 4.3]. We display the argument for completeness and the convenience of the reader.

Proof. We first notice that the linear mapping $L_\sigma^2 \ni g \mapsto \partial_t^k \partial_z^\beta w(t, z) \in \mathbb{R}$ is bounded for every fixed $(t, z) \in (0, \infty) \times \overline{B_1(0)}$ and $(k, \beta) \in \mathbb{N}_0 \times \mathbb{N}_0^N$. Indeed, for small times, boundedness is a consequence of Lemma 12 (with $a = 0$), and for large times, boundedness follows from successively applying Lemmas 13 and 12 (with $a = 0$), namely $|\partial_t^k \partial_z^\beta w(t, z)| \lesssim \|w(\frac{1}{2})\|_\sigma + \|\nabla w(\frac{1}{2})\|_{\sigma+1} \lesssim \|g\|_\sigma$. Riesz’ representation theorem thus provides us with the existence of a unique function $G_{k,\beta}(t, z, \cdot) \in L_\sigma^2$ such that

$$\partial_t^k \partial_z^\beta w(t, z) = \int G_{k,\beta}(t, z, z') g(z') d\mu_\sigma(z').$$

Setting $G = G_{0,0}$, uniqueness implies $G_{k,\beta} = \partial_t^k \partial_z^\beta G$. Notice that G inherits the symmetry in z and z' from the symmetry of the linear operator $\mathcal{L}_\sigma^2 + \mathcal{L}_\sigma$ via the symmetry of the associated semigroup operator $e^{(\mathcal{L}_\sigma^2 + \mathcal{L}_\sigma)t}$.

We now turn to the proof of the Gaussian estimates. We shall write $\chi = \chi_{a,b}(\cdot, z_0)$ for some fixed $z_0 \in \overline{B_1(0)}$ and set $\theta = \theta(r, z)$. We first notice that by Lemma 12, for $r \sim \sqrt[4]{t}$, we have

$$|B_r^d(z)|_\sigma^{1/2} e^{\chi(z)} |w(\frac{1}{2}t, z)| \lesssim e^{C(a^2/b^2 + a^4)t} \|e^\chi g\|_\sigma,$$

and thus, the mapping \mathcal{A} defined by

$$(\mathcal{A}h)(z) = |B_r^d(z)|_\sigma^{1/2} e^{\chi(z)} \int G(\frac{1}{2}t, z, z') e^{-\chi(z')} h(z') d\mu_\sigma(z')$$

for $z \in \overline{B_1(0)}$, is a bounded linear mapping from L_σ^2 to L^∞ with

$$\|\mathcal{A}\|_{L_\sigma^2 \rightarrow L^\infty} \lesssim e^{C(a^2/b^2 + a^4)t}.$$

By the symmetry of the Green's function, it holds that

$$\begin{aligned} \int \mathcal{A}h \xi d\mu_\sigma &= \iint |B_r^d(z)|_\sigma^{1/2} e^{\chi(z)} G(\frac{1}{2}t, z', z) e^{-\chi(z')} h(z') d\mu_\sigma(z') d\mu_\sigma(z) \\ &= \int e^{-\chi} w_\xi(\frac{1}{2}t) h d\mu_\sigma \end{aligned}$$

if w_ξ denotes the solution to the homogeneous equation with initial value $g_\xi = |B_r^d(\cdot)|_\sigma^{1/2} e^\chi \xi$, and if $\xi \in L_\sigma^1$ is such that $g_\xi \in L_\sigma^2$. In particular, the action of the dual $\mathcal{A}^* : (L^\infty)^* \rightarrow L_\sigma^2$ on such functions ξ is given by $\mathcal{A}^* \xi = e^{-\chi} w_\xi(\frac{1}{2}t)$. Because $\|\mathcal{A}\|_{L_\sigma^2 \rightarrow L^\infty} = \|\mathcal{A}^*\|_{(L^\infty)^* \rightarrow L_\sigma^2}$, we then have the estimate

$$\|e^{-\chi} w_\xi(\frac{1}{2}t)\|_\sigma \lesssim e^{C(a^2/b^2 + a^4)t} \|\xi\|_{L_\sigma^1}.$$

An application of Lemma 12 with a replaced by $-a$ then yields that

$$\left| \int \partial_t^k \partial_z^\beta G(t, z, \cdot) |B_r^d(\cdot)|_\sigma^{1/2} e^\chi \xi d\mu_\sigma \right| \lesssim \frac{r^{-4k - |\beta| \theta - |\beta|}}{|B_r^d(z)|_\sigma^{1/2}} e^{C(a^2/b^2 + a^4)t + \chi(z)} \|\xi\|_{L_\sigma^1}.$$

By approximation, it is clear that this estimate holds for any $\xi \in L_\sigma^1$. Thanks to the duality $L^\infty = (L_\sigma^1)^*$, we thus have

$$|\partial_t^k \partial_z^\beta G(t, z, z')| \lesssim \frac{r^{-4k - |\beta| \theta - |\beta|}}{|B_r^d(z)|_\sigma^{1/2} |B_r^d(z')|_\sigma^{1/2}} e^{C(a^2/b^2 + a^4)t + \chi(z) - \chi(z')}.$$

The term $-\chi(z')$ drops out of the exponent upon choosing $z' = z_0$. To conclude the argument for the Gaussian estimates, we distinguish two cases: First, if $\sqrt[4]{t} \geq d(z, z_0)$, then

$$1 \lesssim e^{-C(d(z, z_0)/\sqrt[4]{t})^{4/3}},$$

and thus the statement follows with $a = 0$. Otherwise, if $\sqrt[4]{t} \leq d(z, z_0)$, we choose $a = -\ell$ for some $\ell > 0$ and $b \sim d = d(z, z_0)$ so that the exponent becomes

$$\left(\frac{\ell^2}{d^2} + \ell^4\right)t - \ell d$$

modulo constant prefactors. We optimize the last two terms in ℓ by choosing $\ell \sim (d/t)^{1/3}$. It is easily checked that the exponent is bounded by an expression of the form $1 - (d/\sqrt[4]{t})^{4/3}$, which yields the desired result.

The remaining properties are immediate consequences of the preceding analysis. □

2.5. Calderón–Zygmund estimates. We will see at the beginning of the next subsection that the kernel representation of solutions of the homogeneous problem carries over to the ones of the inhomogeneous problem. This observation is commonly referred to as Duhamel’s principle. To study regularity in the inhomogeneous problem, the detailed knowledge of the Gaussian kernel provided by Proposition 15 is very helpful. A major step in the analysis of Whitney measures is the translation of the energy estimates from weighted L^2 to standard L^p spaces. We are thus led to the study of singular integrals in the spirit of Calderón and Zygmund and the theory of Muckenhoupt weights.

Out of the Euclidean setting, a good framework for these studies is provided by spaces of homogeneous type, see [Coifman and Weiss 1971], which are metric measure spaces, i.e., metric spaces endowed with a doubling Borel measure.¹ The theory of singular integrals in spaces of homogeneous type was elaborated by Koch [1999; 2004; 2008]. For the Euclidean theory, we refer to [Stein 1970; 1993].

Let us recall some pieces of the abstract theory. Let (X, D) be a metric space endowed with a doubling Borel measure μ . A linear operator T on $L^q(X, \mu)$ with $q \in (1, \infty)$ is called a *Calderón–Zygmund operator* if T can be written as

$$Tf(x) = \int_X K(x, y)f(y) d\mu(y)$$

for all $x \in (\text{spt } f)^c$ and $f \in L^\infty(X, \mu) \cap L^q(X, \mu)$, where $K : X \times X \rightarrow \mathbb{R}$ is a measurable kernel such that

$$\begin{aligned} y \mapsto K(x, y) &\in L^1_{\text{loc}}(X \setminus \{x\}, \mu), \\ x \mapsto K(x, y) &\in L^1_{\text{loc}}(X \setminus \{y\}, \mu), \end{aligned}$$

satisfying the following boundedness and *Calderón–Zygmund cancellation conditions*:

$$\sup_{x \neq y} V(x, y)|K(x, y)| \lesssim 1 \tag{39}$$

and

$$\sup_{x \neq y} \sup_{x' \neq y'} V(x, y) \wedge V(x', y') |K(x, y) - K(x', y')| \lesssim \left(\frac{D(x, x') + D(y, y')}{D(x, y) + D(x', y')}\right)^\delta \tag{40}$$

¹In fact, Coifman and Weiss introduced the notion of spaces of homogeneous type with quasimetrics instead of metrics.

for some $\delta \in (0, 1]$. Here we have used the notation

$$V(x, y) = \mu\left(B_{D(x,y)}^D\left(\frac{x+y}{2}\right)\right).$$

It is worth noting that the doubling property of μ implies that we could equivalently have chosen to center the above balls at x or y .

Finally, we call ω a p -Muckenhoupt weight if

$$\sup_B \left(\frac{1}{\mu(B)} \int_B \omega \, d\mu\right) \left(\frac{1}{\mu(B)} \int_B \omega^{-1/(p-1)} \, d\mu\right)^{p-1} < \infty.$$

The class of p -Muckenhoupt weights is denoted by $A_p(X, D, \mu)$.

The theory of singular integrals asserts that any Calderón–Zygmund operator T extends to a bounded operator on any $L^p(X, \mu)$ with $p \in (1, \infty)$; i.e.,

$$\|Tf\|_{L^p(\mu)} \lesssim \|f\|_{L^p(\mu)}.$$

Moreover, if $\omega \in A_p$ is a Muckenhoupt weight, then T is also bounded on $L^p(\mu \llcorner \omega)$, where $d(\mu \llcorner \omega) = \omega d\mu$.

In order to establish L^p maximal regularity estimates for our problem at hand, we have to study singular integrals of the form

$$T_{\ell,k,\beta} f(t, z) = \int_0^\infty \int K_{\ell,k,\beta}((t, z), (t', z')) f(t', z') \, d\mu_\sigma(z') \, dt',$$

where $K_{\ell,k,\beta}((t, z), (t', z')) = \chi_{(0,t)}(t') \rho(z)^\ell \partial_t^k \partial_z^\beta G(t - t', z, z')$. In fact, we will see that $T_{\ell,k,\beta}$ is a Calderón–Zygmund operator on the product space $(0, \infty) \times B_1(0)$ provided that ℓ, k , and β are such that

$$(\ell, k, |\beta|) \in \mathcal{E} = \{(0, 1, 0), (0, 0, 2), (1, 0, 3), (2, 0, 4)\}. \tag{41}$$

We will accordingly refer to any tuple (ℓ, k, β) in the above class as a *Calderón–Zygmund exponent*.

The product space $X = (0, \infty) \times B_1(0)$ will be endowed with the metric

$$D((t, z), (t', z')) = \sqrt[4]{|t - t'| + d(z, z')^4},$$

which reflects the parabolic scaling of the linear differential operator, and the product measure $\mu = \lambda^1 \otimes \mu_\sigma$, with λ^1 denoting the one-dimensional Lebesgue. Because d is doubling, so is D , and thus the metric measure space (X, D, μ) is of *homogeneous type* in the sense of [Coifman and Weiss 1971] and is thus suitable for Calderón–Zygmund theory. Notice also that the volume tensor $V((t, z), (t', z'))$ simplifies to

$$V((t, z), (t', z')) \sim D((t, z), (t', z'))^4 \left| B_{D((t,z),(t',z'))}^d\left(\frac{z+z'}{2}\right) \right|_\sigma. \tag{42}$$

Without proof, we state the following lemma:

Lemma 17. *If (ℓ, k, β) is such that (41) holds, then $T_{\ell,k,\beta}$ is a Calderón–Zygmund operator.*

The proof is almost identical to the one in the porous-medium setting; see Lemmas 4.20 and 4.21 in [Seis 2015]. We will thus refrain from repeating the argument and refer the interested reader to the quoted paper.

2.6. The inhomogeneous problem. In this subsection, we consider the inhomogeneous problem with zero initial datum,

$$\begin{cases} \partial_t w + \mathcal{L}_\sigma^2 w + n\mathcal{L}_\sigma w = f & \text{in } (0, \infty) \times B_1(0), \\ w(0, \cdot) = 0 & \text{in } B_1(0). \end{cases} \tag{43}$$

Our first observation is that the kernel representation from [Proposition 15](#) carries over to the inhomogeneous setting.

Lemma 18 (Duhamel’s principle). *If $f \in L^2(L_\sigma^2)$ and w is the solution to the initial value problem (43), then*

$$w(t, z) = \int_0^t \int G(t - t', z, z') f(t', z') d\mu_\sigma(z') dt'$$

for all $(t, z) \in (0, \infty) \times \overline{B_1(0)}$.

Proof. The statement follows from the fact that G is a fundamental solution, see [Proposition 15\(3\)](#). \square

Proposition 19. *Let w be the solution to the initial value problem (43). Then, for any $p \in (1, \infty)$ it holds that*

$$\|\partial_t w\|_{L^p} + \|\nabla w\|_{L^p} + \|\nabla^2 w\|_{L^p} + \|\rho \nabla^3 w\|_{L^p} + \|\rho^2 \nabla^4 w\|_{L^p} \lesssim \|f\|_{L^p}. \tag{44}$$

Proof. The purpose of this lemma is to carry the energy estimates from [Lemma 8](#) over to the standard L^p setting. This is achieved by applying the abstract theory recalled in the previous subsection. In fact, as a consequence of [Lemma 18](#), any function $\rho^\ell \partial_t^k \partial_z^\beta w$ has the kernel representation

$$T_{\ell,k,\beta} f(t, z) = \int_0^\infty \int K_{\ell,k,\beta}((t, z), (t', z')) f(t', z') d\mu_\sigma(z') dt',$$

where

$$K_{\ell,k,\beta}((t, z), (t', z')) = \chi_{(0,t)}(t') \rho(z)^\ell \partial_t^k \partial_z^\beta G(t - t', z, z').$$

If (ℓ, k, β) are Calderón–Zygmund exponents [\(41\)](#), by [Lemma 17](#), the energy estimates from [Lemma 8](#) carry over to any $L^p(L^p(\mu_\sigma))$ space with $p \in (1, \infty)$. Moreover, if ν is a Muckenhoupt weight in $A_p(B_1, d, \mu_\sigma)$, then the operators $T_{\ell,k,\beta}$ are bounded on $L^p(L^p(\mu_\sigma \upharpoonright \nu))$. Notice that this is the case for weights of the form $\nu = \rho^\gamma$ precisely if $-(\sigma + 1) < \gamma < (p - 1)(\sigma + 1)$. In particular, choosing $\gamma = -\sigma$, we see that $T_{\ell,k,\beta}$ is bounded on $L^p = L^p(L^p)$ for any $p \in (1, \infty)$ because $\sigma > 0$. This is the statement of the proposition apart from the term $\|\nabla w\|_{L^p}$. The control of this term can be deduced, for instance, from the analogous estimates for the porous-medium equation, see [Proposition 4.23](#) in [\[Seis 2015\]](#), applied to $\partial_t w + n\mathcal{L}_\sigma w = f - \mathcal{L}_\sigma^2 w$. \square

In the following, we consider the larger cylinders

$$\widehat{Q}_r^d(z_0) := \left(\frac{1}{4}r^4, r^4\right) \times B_{2r}^d(z_0) \quad \text{and} \quad \widehat{Q}(T) = \left(\frac{1}{4}T, T\right) \times \overline{B_1(0)}.$$

Lemma 20. (1) *Suppose that $\text{spt } f \subset \widehat{Q}_r^d(z_0)$ for some $z_0 \in \overline{B_1(0)}$ and $0 < r \lesssim 1$. Then for any (ℓ, k, β) satisfying [\(41\)](#) and any $p \in (1, \infty)$, it holds that*

$$r^4 |\widehat{Q}_r^d(z_0)|^{-1/p} \|\rho^\ell \partial_t^k \partial_z^\beta w\|_{L^p(\widehat{Q}_r^d(z_0))} \lesssim \|f\|_{Y(p)}.$$

(2) Suppose that $\text{spt } f \subset \widehat{Q}(T)$ for some $T \geq 1$. Then it holds for any $p \in (1, \infty)$ that

$$\sum_{(\ell, k, |\beta|) \in \mathcal{E}} T \|\rho^\ell \partial_t^k \partial_z^\beta w\|_{L^p(Q(T))} \lesssim \|f\|_{Y(p)}.$$

Proof. We will only prove the first statement. The argument for the second one is very similar. The desired estimate is an immediate consequence of Proposition 19. Indeed, the latter implies

$$\|\rho^\ell \partial_t^k \partial_z^\beta w\|_{L^p(Q_r^d(z_0))} \lesssim \|f\|_{L^p(\widehat{Q}_r^d(z_0))}.$$

If now $\{Q_{r_i}^d(z_i)\}_{i \in I}$ is a finite cover of $\widehat{Q}_r^d(z_0)$ with radii $r_i \sim r$ and such that $\sum_i |Q_{r_i}^d(z_i)| \lesssim |\widehat{Q}_r^d(z_0)|$, then

$$\|f\|_{L^p(\widehat{Q}_r^d(z_0))} \leq \sum_{i \in I} \|f\|_{L^p(Q_{r_i}^d(z_i))} \lesssim \frac{1}{r^4} |\widehat{Q}_r^d(z_0)|^{1/p} \|f\|_{Y(p)}.$$

Notice that $|\widehat{Q}_r^d(z_0)| \lesssim |Q_r^d(z_0)|$, because $\mu_\sigma \otimes \lambda^1$ is doubling, which concludes the proof. \square

In view of the definition of the $X(p)$ norm, the estimates on the second- and third-order spatial derivative derived in the previous lemma are not strong enough for balls $B_r^d(z_0)$ that are relatively far away from the boundary in the sense that $\sqrt{\rho(z_0)} \gg r$. Estimates in such balls, as well as uniform bounds on w and ∇w , are derived in the lemma that follows.

Lemma 21. (1) Suppose that $\text{spt } f \subset \widehat{Q}_r^d(z_0)$ for some $z_0 \in \overline{B_1(0)}$ and $0 < r \lesssim 1$ and let $p > N + 4$. Then it holds for any $0 < t \lesssim r^4$ that

$$|w(t, z_0)| + |\nabla w(t, z_0)| \lesssim \|f\|_{Y(p)}.$$

If moreover $\sqrt{\rho(z_0)} \gg r$, then it holds that

$$r\theta(r, z_0) |Q_r^d(z_0)|^{-1/p} \|\nabla^2 w\|_{L^p(Q_r^d(z_0))} + r^2\theta(r, z_0)^2 |Q_r^d(z_0)|^{-1/p} \|\nabla^3 w\|_{L^p(Q_r^d(z_0))} \lesssim \|f\|_{Y(p)}.$$

(2) Suppose that $\text{spt } f \subset \widehat{Q}(T)$ for some $T \geq 1$. Then it holds for any $p > 1 + \frac{1}{2}N$ that

$$\|w\|_{L^\infty(Q(T))} + \|\nabla w\|_{L^\infty(Q(T))} \lesssim \|f\|_{Y(p)}.$$

Proof. (1) As a consequence of Lemma 18 and Hölder’s inequality, we have

$$|\partial_z^\beta w(t, z)| \leq \left(\int_0^{r^4} \|\partial_z^\beta G(\tau, z, \cdot)\|_{L_{q\sigma}^q}^q d\tau \right)^{1/q} \|f\|_{L^p}, \tag{45}$$

where q is such that $\frac{1}{p} + \frac{1}{q} = 1$ and $\beta \in \mathbb{N}_0^N$. The statements thus follow from suitable estimates for the kernel functions. From Proposition 15 we recall that

$$|\partial_z^\beta G(\tau, z, z')| \lesssim \sqrt[4]{\tau}^{-|\beta|} \theta(\sqrt[4]{\tau}, z)^{-|\beta|} |B_{\sqrt[4]{\tau}}^d(z)|_\sigma^{-1} e^{-C(d(z, z')/\sqrt[4]{\tau})^{4/3}}. \tag{46}$$

Let $\{B_j^d(\sqrt[4]{\tau}(z))\}_{j \in J}$ be a finite cover of $\overline{B_1}$. Then

$$\int e^{-qC(d(z, z')/\sqrt[4]{\tau})^{4/3}} d\mu_{q\sigma}(z') \leq \sum_{j \in J} e^{-qC(j-1)^{4/3}} |B_j^d(\sqrt[4]{\tau}(z))|_{q\sigma}.$$

Notice that by the virtue of (15),

$$|B_{j\sqrt[4]{\tau}}^d(z)|_{q\sigma} \lesssim j^{2N} |B_{\sqrt[4]{\tau}}^d(z)|_{q\sigma} \sim j^{2N} |B_{\sqrt[4]{\tau}}^d(z)|^{1-q} |B_{\sqrt[4]{\tau}}^d(z)|_{\sigma}^q,$$

which in turn implies

$$\int e^{-qC(d(z,z')/\sqrt[4]{\tau})^{4/3}} d\mu_{q\sigma}(z') \lesssim \left(\sum_{j \in J} e^{-q(j-1)^{4/3}} j^{2N} \right) |B_{\sqrt[4]{\tau}}^d(z)|^{1-q} |B_{\sqrt[4]{\tau}}^d(z)|_{\sigma}^q.$$

The sum is converging and can thus be absorbed in the (suppressed) constant. We now integrate (46) over time and space and obtain

$$\int_0^{r^4} \|\partial_z^\beta G(\tau, z, \cdot)\|_{L_{q\sigma}^q}^q d\tau \lesssim \int_0^{r^4} \sqrt[4]{\tau}^{-|\beta|q} \theta(\sqrt[4]{\tau}, z)^{-|\beta|q} |B_{\sqrt[4]{\tau}}^d(z)|^{1-q} d\tau \tag{47}$$

for any $z \in \bar{B}_1$.

First, if $\sqrt{\rho(z)} \lesssim r$, then by (12), estimate (47) turns into

$$\int_0^{r^4} \|\partial_z^\beta G(\tau, z, \cdot)\|_{L_{q\sigma}^q}^q d\tau \lesssim \int_0^{r^4} \sqrt[4]{\tau}^{-q|\beta|-(q-1)N} d\tau \lesssim r^{4-2q|\beta|-2(q-1)N},$$

provided that $N + 2 < (2 - |\beta|)p$, which is consistent with the assumptions in the lemma only if $|\beta| \in \{0, 1\}$.

It remains to notice that

$$r^{4-2(q-1)N} \lesssim r^{4q} |Q_r^d(z)|^{1-q}$$

by virtue of (15). From this and (45), we easily derive the first estimate in the first part of the lemma in the case where $\sqrt{\rho(z)} \lesssim r$.

Second, if $\sqrt{\rho(z)} \gg r$, then (47) becomes

$$\begin{aligned} \int_0^{r^4} \|\partial_z^\beta G(\tau, z, \cdot)\|_{L_{q\sigma}^q}^q d\tau &\lesssim \sqrt{\rho(z)}^{-|\beta|q-(q-1)N} \int_0^{r^4} \sqrt[4]{\tau}^{-|\beta|q-N(q-1)} d\tau \\ &\lesssim \sqrt{\rho(z)}^{-|\beta|q-(q-1)N} r^{4-|\beta|q-(q-1)N}, \end{aligned}$$

provided that $N + 4 < (4 - |\beta|)p$, which is consistent with the assumptions only if $|\beta| \in \{0, 1, 2, 3\}$. Now we notice that

$$\sqrt{\rho(z)}^{-|\beta|q-(q-1)N} r^{4-|\beta|q-(q-1)N} \lesssim r^{(4-|\beta|)q} \theta^{-|\beta|q} |Q_r^d(z)|^{1-q},$$

using (15) again. It is not difficult to see that the latter estimates in combination with (45) imply remaining estimates in the first part of the lemma.

(2) By Duhamel’s principle in Lemma 18 and the fact that f is concentrated on $\hat{Q}(T)$, we have for any $(t, z) \in Q(T)$ and $\beta \in \mathbb{N}_0^N$ that

$$\begin{aligned} |\partial_z^\beta w(t, z)| &\leq \int_{T/4}^{t-1} \|\partial_z^\beta G(t-t', z, z')\| |f(t', z')| d\mu_\sigma(z') dt' \\ &\quad + \int_{t-1}^t \|\partial_z^\beta G(t-t', z, z')\| |f(t', z')| d\mu_\sigma(z') dt', \end{aligned}$$

with the convention that the first integral is zero if $\frac{1}{4}T \geq t - 1$. If it is nonzero, we use Proposition 15(5) and estimate the latter by

$$\int_{T/4}^{t-1} \int |f| d\mu_\sigma dt' \lesssim T^{1-1/p} \|f\|_{L^p(\hat{Q}(T))} \leq T \|f\|_{L^p(\hat{Q}(T))}.$$

Similarly, applying the same strategy as in part (1) above, we bound the second term by

$$\|\partial_z^\beta G(\cdot, z, \cdot)\|_{L^q((0,1);L^q_\sigma)} \|f\|_{L^p(\hat{Q}(T))} \lesssim \|f\|_{L^p(\hat{Q}(T))}.$$

The statement thus follows by choosing $|\beta| \in \{0, 1\}$. □

We need some estimates for the off-diagonal parts.

Lemma 22. (1) *Suppose that $\text{spt } f \subset [0, r^4] \times \overline{B_1(0)} \setminus \hat{Q}_r^d(z_0)$ for some $z_0 \in \overline{B_1(0)}$ and $0 < r \lesssim 1$. Then it holds for any $p \in (1, \infty)$ that*

$$\|w\|_{L^\infty(Q_r^d(z_0))} + \|\nabla w\|_{L^\infty(Q_r^d(z_0))} + \sum_{(\ell,k,|\beta|) \in \mathcal{E}} \frac{r^{4k+|\beta|}}{\theta(r, z_0)^{2\ell-|\beta|}} \|\rho^\ell \partial_t^k \partial_z^\beta w\|_{L^p(Q_r^d(z_0))} \lesssim \|f\|_{Y(p)}.$$

(2) *Suppose that $\text{spt } f \subset [\frac{1}{2}, \frac{1}{4}T] \times \overline{B_1(0)}$ for some $T \geq 2$. Then it holds for any $p \in (1, \infty)$ that*

$$\|w\|_{L^\infty(Q(T))} + \|\nabla w\|_{L^\infty(Q(T))} + \sum_{(\ell,k,|\beta|) \in \mathcal{E}} T \|\rho^\ell \partial_t^k \partial_z^\beta w\|_{L^p(Q(T))} \lesssim \|f\|_{Y(p)}.$$

Proof. (1) We begin our proof with a helpful elementary estimate: if θ and C are given positive constants, then there exists a new constant \tilde{C} such that

$$\sqrt[4]{t-t'}^{-\theta} e^{-C(d(z,z')/\sqrt[4]{t-t'})^{4/3}} \lesssim r^{-\theta} e^{-\tilde{C}(d(z,z')/r)^{4/3}} \tag{48}$$

for all $(t, z) \in Q_r^d(z_0)$ and $(t', z') \in [0, r^4] \times \overline{B_1} \setminus \hat{Q}_r^d(z_0)$. The argument for (48) runs as follows: To simplify the notation slightly, we write $\tau = t - t'$ and $d = d(z, z')$. If $z' \in B_{2r}^d(z_0)$, then necessarily $t' \notin (\frac{1}{4}r^4, r^4)$, and therefore $\tau \gtrsim r^4$. It follows that

$$\sqrt[4]{\tau}^{-\theta} e^{-C(d/\sqrt[4]{\tau})^{4/3}} \leq \sqrt[4]{\tau} \lesssim r^{-\theta} e^{-C(d/r)^{4/3}},$$

because $d(z, z') \leq d(z, z_0) + d(z_0, z') \leq 3r$. Otherwise, if $z' \notin B_{2r}^d(z_0)$, it holds that $2r \leq d(z', z_0) \leq d(z, z') + r$, and thus $\sqrt[4]{\tau} \lesssim r \lesssim d$. Using the fact that $\tau \mapsto \sqrt[4]{\tau}^{-\theta} e^{-C(d/\sqrt[4]{\tau})^{4/3}}$ is increasing for $0 < \tau \lesssim d^4$, we then estimate

$$\sqrt[4]{\tau}^{-\theta} e^{-C(d/\sqrt[4]{\tau})^{4/3}} \lesssim r^{-\theta} e^{-\tilde{C}(d/r)^{4/3}}.$$

This completes the proof of (48).

In the following, C will be a uniform constant whose value may change from line to line.

Because $f = 0$ in $Q_r^d(z_0)$, Duhamel’s principle (Lemma 18) and the Gaussian estimates from Proposition 15 imply

$$\begin{aligned} & |\partial_t^k \partial_z^\beta w(t, z)| \\ & \lesssim \int_0^t \int \frac{\sqrt[4]{\tau}^{-4k-|\beta|} \theta(\sqrt[4]{\tau}, z)^{-|\beta|}}{|B_{\sqrt[4]{\tau}}^d(z)|_\sigma} e^{-C(d(z, z')/\sqrt[4]{\tau})^{4/3}} |f(t - \tau, z')| d\mu_\sigma(z') d\tau \\ & \lesssim \int_0^{r^4} \int \left(\frac{\sqrt[4]{\tau}}{\sqrt[4]{\tau} + \sqrt{\rho(z)}} \right)^{|\beta|+N+2\sigma} \sqrt[4]{\tau}^{-4k-2|\beta|-2N-2\sigma} e^{-C(d(z, z')/\sqrt[4]{\tau})^{4/3}} |f(t - \tau, z')| d\mu_\sigma(z') d\tau. \end{aligned}$$

As a consequence of (48), Remark 16 and the monotonicity of the function $s \mapsto s/(s + c)$ for any fixed $c > 0$, we may substitute any $\sqrt[4]{\tau}$ by r and find

$$\begin{aligned} & \frac{r^{4k+|\beta|-1}}{\theta(r, z)^{2\ell-|\beta|+1}} |\rho(z)^\ell \partial_t^k \partial_z^\beta w(t, z)| \\ & \lesssim \frac{1}{|B_r^d(z)|_\sigma} \int_0^{r^4} \int e^{-C(d(z, z')/r)^{4/3}} r^{-1} \theta(r, z')^{-1} |f(t', z')| d\mu_\sigma(z') dt'. \end{aligned}$$

We consider now a finite family of balls $\{B_r^d(z_0)\}_{i \in I}$ covering $\overline{B_1(0)}$. Since $d(z, z_i) \leq d(z, z') + r$ for any $z' \in B_r^d(z_i)$ and

$$\sum_{i \in I} e^{-C(d(z, z_i)/r)^{4/3}} < \infty$$

uniformly in r and z , we further estimate the right-hand side of the last inequality by

$$\sup_{\tilde{z} \in B_1} \frac{1}{|B_r^d(\tilde{z})|_\sigma} \int_0^{r^4} \int_{B_r^d(\tilde{z})} r^{-1} \theta(r, z')^{-1} |f(t', z')| d\mu_\sigma(z') dt'. \tag{49}$$

We claim that this term is controlled by $\|f\|_{Y(p)}$. To see this, we fix $\tilde{z} \in B_1(0)$ and let $r_j = (\sqrt{2/3})^j r$. Applying a non-Euclidean version of Vitali’s covering lemma, see Lemma 2.2.2 in [Koch 1999], we find a finite family of balls $\{B_{r_j}^d(z_{ij})\}_{i \in I_j}$ covering $B_r^d(\tilde{z})$ and such that

$$\sum_{i \in I_j} |B_{r_j}^d(z_{ij})|_\sigma \lesssim |B_r^d(\tilde{z})|_\sigma \tag{50}$$

uniformly in j, r , and \tilde{z} . Then $(0, r^4) \times B_r^d(\tilde{z})$ is contained in the countable union $\bigcup_{j \in \mathbb{N}_0} \bigcup_{i \in I_j} Q_{r_j}^d(z_{ij})$. Invoking Hölder’s inequality we thus find

$$\begin{aligned} & \int_0^{r^4} \int_{B_r^d(\tilde{z})} r^{-1} \theta(r, z')^{-1} |f(t', z')| d\mu_\sigma(z') dt' \\ & \leq \sum_{j \in \mathbb{N}_0} \sum_{i \in I_j} (\sqrt{2/3})^j r_j^{-1} \|\theta(r, \cdot)^{-1}\|_{L_{q\sigma}^q(Q_{r_j}^d(z_{ij}))} \|f\|_{L^p(Q_{r_j}^d(z_{ij}))}, \end{aligned}$$

where, as usual, $\frac{1}{p} + \frac{1}{q} = 1$. Notice that $\mu_{q\sigma}$ is a finite measure for any $q \in (1, \infty)$. From (12) and (13) we deduce that $\theta(r_j, z_{ij}) \sim \theta(r_j, z') \leq \theta(r, z')$ for any $z' \in B_{r_j}^d(z_{ij})$, and thus, as a consequence of (15),

$$\|\theta(r, \cdot)^{-1}\|_{L_{q\sigma}^q(Q_{r_j}^d(z_{ij}))} \lesssim r_j^4 \theta(r_j, z_{ij})^{-1} |Q_{r_j}^d(z_{ij})|^{-1/p} |B_{r_j}^d(z_{ij})|_\sigma.$$

Combining the previous two estimates, using (50) and the convergence of the geometric series finally yields that the term in (49) is bounded by $\|f\|_{Y(p)}$. We have thus proved that

$$\frac{r^{4k+|\beta|-1}}{\theta(r, z)^{2\ell-|\beta|+1}} |\rho(z)^\ell \partial_t^k \partial_z^\beta w(t, z)| \lesssim \|f\|_{Y(p)}.$$

We easily deduce the statement of the lemma.

(2) To prove the second statement, we use Lemma 18 and Proposition 15(5) to estimate

$$|\partial_t^k \partial_z^\beta w(t, z)| \lesssim \int_{1/2}^{T/4} e^{-(t-t')\lambda_1} \int |f(t', z')| d\mu_\sigma(z') dt'$$

for any $(t, z) \in Q(T)$ and any $k \in \mathbb{N}_0$ and $\beta \in \mathbb{N}_0^N$ with $k + |\beta| \geq 1$. Let $M \in \mathbb{Z}$ be such that $2^M \leq \frac{1}{4}T < 2^{M+1}$. We then split and compute

$$\begin{aligned} |\partial_t^k \partial_z^\beta w(t, z)| &\lesssim \sum_{m=0}^M \int_{2^{m-1}}^{2^m} e^{-(t-t')\lambda_1} \int |f| d\mu_\sigma dt + \int_{T/8}^{T/4} e^{-(t-t')\lambda_1} \int |f| d\mu_\sigma dt' \\ &\lesssim \sum_{m=0}^M e^{-(t-2^m)\lambda_1} \|f\|_{L^p(Q(2^m))} + e^{-(T/4)\lambda_1} \|f\|_{L^p(Q(T/4))} \\ &\lesssim e^{-(T/4)\lambda_1} \|f\|_{Y(p)}. \end{aligned}$$

We easily infer all estimates but the uniform bound on w . To gain control on $\|w\|_{L^\infty}$, we argue similarly and get

$$|w(t, z)| \lesssim \sum_{m=0}^{M+1} \int_{2^{m-1}}^{2^m} \int |f| d\mu_\sigma dt' \lesssim \left(\sum_{m=0}^{M+1} \frac{1}{(2^p)^m} \right) \|f\|_{Y(p)}.$$

The desired estimate follows from the convergence of the geometric series. □

A combination of the results in this subsection yields the maximal regularity estimate for the inhomogeneous problem (43).

Proposition 23. *Suppose that $p > N + 4$. Let w be a solution to the homogeneous problem (43). Then*

$$\|w\|_{L^\infty(W^{1,\infty})} + \|w\|_{X(p)} \lesssim \|f\|_{Y(p)}.$$

Proof. The statement follows immediately from Lemmas 20, 21 and 22 and the superposition principle for linear equations: For small times, we split f into $\eta f + (1 - \eta)f$ with η being a smooth cut-off function such that $\eta = 1$ on $Q_r^d(z_0)$ and $\eta = 0$ outside $\widehat{Q}_r^d(z_0)$ for some arbitrarily fixed $r \lesssim 1$ and $z_0 \in \overline{B_1(0)}$. For large times, we make a hard temporal cut-off by splitting f into $\chi f + (1 - \chi)f$, where χ is the characteristic function on $\widehat{Q}(T)$. Notice that to estimate the large times, it is enough to study such f 's that are zero in the initial time interval $(0, \frac{1}{2})$. For details, we refer to [Seis 2015]. □

3. The nonlinear problem

Our goal in this section is the derivation of Theorems 1 and 2. The existence of a unique solution to the nonlinear problem is a consequence of a fixed-point argument. We need the following lemma:

Lemma 24. *Let w_1 and w_2 be two functions satisfying*

$$\|w_i\|_{L^\infty(W^{1,\infty})} + \|w_i\|_{X(p)} \leq \varepsilon, \quad i = 1, 2, \tag{51}$$

for some small $\varepsilon > 0$. Then

$$\|f[w_1] - f[w_2]\|_{Y(p)} \lesssim \varepsilon (\|w_1 - w_2\|_{L^\infty(W^{1,\infty})} + \|w_1 - w_2\|_{X(p)}).$$

Proof. For notational convenience, we write $f_i^j = f^j[w_i]$ for any $i \in \{1, 2\}$ and $j \in \{1, 2, 3\}$. We will also just write w instead of w_1 or w_2 if the index doesn't matter. We remark that by the virtue of (51), it holds that

$$\begin{aligned} |R_k[w_1] - R_k[w_2]| &\lesssim \|w_1 - w_2\|_{L^\infty(W^{1,\infty})}, \\ |R_k[w]| &\lesssim 1 \end{aligned}$$

for any value of k .

The estimates of the differences of the f_i^j is very similar. We focus on the leading-order terms, i.e., $f_1^3 - f_2^3$. Using (51) and the previous bounds on the R_k , we first notice that

$$\begin{aligned} |f_1^3 - f_2^3| &\lesssim \rho^2 \|w_1 - w_2\|_{L^\infty(W^{1,\infty})} (|\nabla^2 w|^3 + |\nabla^2 w| |\nabla^3 w| + |\nabla w| |\nabla^4 w|) \\ &\quad + \rho^2 |\nabla^2 w_1 - \nabla^2 w_2| (|\nabla^2 w|^2 + |\nabla w| |\nabla^3 w|) \\ &\quad + \rho^2 |\nabla^3 w_1 - \nabla^3 w_2| |\nabla w| |\nabla^2 w| + \rho^2 |\nabla^4 w_1 - \nabla^4 w_2| |\nabla w|^2. \end{aligned}$$

The control of the individual terms is derived very similarly. There are a few obvious cases; for instance the last term, which is simply controlled by using (51):

$$\|\rho^2 |\nabla^4 w_1 - \nabla^4 w_2| |\nabla w|^2\|_{Y(p)} \leq \|w\|_{L^\infty(W^{1,\infty})}^2 \|w_1 - w_2\|_{X(p)} \leq \varepsilon \|w_1 - w_2\|_{X(p)}.$$

For most of the remaining terms, we have to make use of the following interpolation inequality

$$\|\nabla^i \zeta\|_{L_r^m}^m \lesssim \|\zeta\|_{L^\infty}^{m-i} \|\nabla^m \zeta\|_{L_r^p}^i,$$

provided that $mp = ir$ for some integers $i < m$, which has been proved in Appendix A of [Seis 2015]. For instance, setting $\zeta = \eta \nabla w$ for some smooth cut-off function η satisfying $\eta = 1$ in $B_r^d(z_0)$ and $\eta = 0$ outside $B_{2r}^d(z_0)$, we have

$$\|\rho^2 |\nabla^2 w|^3\|_{L^p(B_r^d(z_0))} = \|\nabla^2 w\|_{L_{2r}^{3p}(B_r^d(z_0))}^3 \leq \|\nabla \zeta\|_{L_{2r}^{3p}}^3.$$

Applying the above interpolation inequality and using the fact that η varies on the scale $r\theta(r, z_0)$ and $\rho \lesssim \theta(r, z_0)^2$ in $B_{2r}^d(z_0)$, see (12) and (13), we then get

$$\begin{aligned} \|\rho^2 |\nabla^2 w|^3\|_{L^p(B_r^d(z_0))} &\lesssim \|\nabla w\|_{L^\infty}^2 \left(\|\rho^2 \nabla^4\|_{L^p(B_{2r}^d(z_0))} + \frac{\theta}{r} \|\rho \nabla^3 w\|_{L^p(B_{2r}^d(z_0))} \right. \\ &\quad \left. + \frac{\theta^2}{r^2} \|\nabla^2 w\|_{L^p(B_{2r}^d(z_0))} + \frac{\theta}{r^3} \|\nabla w\|_{L^p(B_{2r}^d(z_0))} \right), \end{aligned}$$

where $\theta = \theta(r, z_0)$. Integrating in time over $(\frac{1}{2}r^4, r^4)$, multiplying by r^3/θ and using (51) then yields

$$\sup_{r, z_0} \frac{r^3}{\theta} |Q_r^d(z_0)|^{-1/p} \|\rho^2 |\nabla^2 w|^3\|_{L^p(Q_r^d(z_0))} \lesssim \varepsilon \|w\|_{X(p)}.$$

This type of estimate can be used, for instance, to bound the first term in the above estimate for $f_1^3 - f_2^3$ for small times. The remaining terms and the large-time parts of the $Y(p)$ norm can be controlled in a similar way. □

We are now in the position to prove Theorems 1 and 2.

Proof of Theorems 1 and 2. To simplify the notation in the following, we denote by $\bar{X}(p)$ the intersection $X(p) \cap L^\infty(W^{1,\infty})$ and set $\|\cdot\|_{\bar{X}(p)} = \|\cdot\|_{X(p)} + \|\cdot\|_{L^\infty(W^{1,\infty})}$. Let ε and ε_0 be two positive constants. We denote by M_ε the set of all functions w in $\bar{X}(p)$ such that $\|w\|_{\bar{X}(p)} \leq \varepsilon$ and by N_{ε_0} the set of all functions g such that $\|g\|_{W^{1,\infty}} \leq \varepsilon_0$. We divide the proof into several steps.

Step 1: existence and uniqueness. For $w \in M_\varepsilon$ and $g \in N_{\varepsilon_0}$ given, we denote by $\tilde{w} := I(w, g)$ the unique solution to the linear problem (43) with inhomogeneity $f = f[w]$. By Theorem 4, we have the estimate $\|\tilde{w}\|_{\bar{X}(p)} \lesssim \|f[w]\|_{Y(p)} + \|g\|_{W^{1,\infty}}$. Applying Lemma 24 with $w_1 = w$ and $w_2 = 0$ and using the assumptions on w and g , we find that $\|\tilde{w}\|_{\bar{X}(p)} \leq C(\varepsilon^2 + \varepsilon_0)$ for some positive constant C . We choose ε and ε_0 small enough so that $C\varepsilon^2 \leq \frac{1}{2}\varepsilon$ and $C\varepsilon_0 \leq \frac{1}{2}\varepsilon$, with the consequence that $\tilde{w} \in M_\varepsilon$. This reasoning implies that for any fixed $g \in N_{\varepsilon_0}$, the function $\tilde{w}(\cdot, g)$ maps the set M_ε into itself. Moreover, given w_1 and w_2 in M_ε , we find by linearity and Lemma 24 that

$$\|I(w_1, g) - I(w_2, g)\|_{\bar{X}(p)} \lesssim \|f[w_1] - f[w_2]\|_{Y(p)} \lesssim \varepsilon \|w_1 - w_2\|_{\bar{X}(p)}.$$

Thus, choosing ε even smaller, if necessary, the previous estimate shows that $I(\cdot, g)$ is a contraction on M_ε . By Banach's fixed-point argument, there exists thus a unique $w^* \in M_\varepsilon$ such that $w^* = I(w^*, g)$. In particular, w^* solves the nonlinear equation. From the previous choice of ε , we moreover deduce that $\|w^*\|_{\bar{X}(p)} \lesssim \|g\|_{W^{1,\infty}}$.

Step 2: analytic dependence on initial data. In order to show that w^* depends analytically on g , we will apply the analytic implicit function theorem; see [Deimling 1985, Theorem 15.3]. Because the nonlinearity $f = f[w]$ is a rational function of w and ∇w , and thus analytic away from its poles, the contraction map I is an analytic function on $M_\varepsilon \times N_{\varepsilon_0}$. We consider the map $J : M_\varepsilon \times N_{\varepsilon_0} \rightarrow M_\varepsilon$ defined by $J(w, g) = w - I(w, g)$. Because I is analytic, so is J . It holds that $I(0, 0) = 0$ and $D_w I(0, 0) = \text{id}$. From the analytic implicit function theorem we deduce the existence of two constants $\tilde{\varepsilon} < \varepsilon$ and $\tilde{\varepsilon}_0 < \varepsilon_0$ and of an analytic map $A : N_{\tilde{\varepsilon}_0} \rightarrow M_{\tilde{\varepsilon}}$ with $A(0) = 0$ and such that $J(w, g) = 0$ if and only if $A(g) = w$. From the uniqueness of the fixed point and the definition of J we then conclude that the map $g \mapsto w^*$ is analytic from $N_{\tilde{\varepsilon}_0}$ to $M_{\tilde{\varepsilon}}$.

Step 3: analytic dependence on time and tangential coordinates. Let us now change from Euclidean to spherical coordinates. For $z = (z_1, \dots, z_N)^T \in \bar{B}_1(0)$, we find radius $s \in [0, 1]$ and an angle vector $\phi = (\phi_1, \dots, \phi_{N-1})^T \in \mathcal{A}_{N-1} := [0, \pi]^{N-2} \times [0, 2\pi]$ such that $z_n = s(\prod_{i=1}^{n-1} \sin \phi_i) \cos \phi_n$ for $n \leq N-1$

and $z_N = s \prod_{i=1}^{N-1} \sin \phi_i$. By a slight abuse of notation, we write $w(t, z) = w(t, s, \phi)$. For $\lambda \in \mathbb{R}$ and $\psi \in \mathcal{A}_{N-1}$ we define

$$w_{\lambda, \psi}^* := w^* \circ \Xi_{\lambda, \psi}, \quad \Xi_{\lambda, \psi}(t, s, \phi) := (\lambda t, s, \phi + t \psi).$$

A short computation reveals that $w_{\lambda, \psi}$ solves the equation

$$\partial_t w_{\lambda, \psi}^* + \mathcal{H}_\sigma w_{\lambda, \psi}^* = f_{\lambda, \psi}[w_{\lambda, \psi}^*],$$

where

$$f_{\lambda, \psi}[w] := \lambda f[w] + (1 - \lambda)\mathcal{H}_\sigma w + \psi \cdot \nabla_\phi w, \quad \mathcal{H}_\sigma = \mathcal{L}_\sigma^2 + n\mathcal{L}_\sigma.$$

Clearly, $f_{1,0} = f$. Similarly as above, we denote by $I_{\lambda, \psi}(w, g)$ the solution to the linear equation with inhomogeneity $f_{\lambda, \psi}[w]$ and initial datum g . We furthermore set $J_{\lambda, \psi}(w, g) := w - I_{\lambda, \psi}(w, g)$. It is obvious that $J_{1,0}(0, 0) = 0$ and $D_w J_{1,0}(0, 0) = \text{id}$. Applying the analytic implicit function theorem once more, we find positive constants $\delta, \hat{\varepsilon} < \tilde{\varepsilon}, \hat{\varepsilon}_0 < \tilde{\varepsilon}_0$ and an analytic function $A_{\lambda, \psi}(g) = A(\lambda, \psi, g)$ from $B_\delta^{\mathbb{R}}(1) \times B_\delta^{\mathbb{R}^{N-1}}(0) \times N_{\hat{\varepsilon}_0}$ to $M_{\hat{\varepsilon}}$ such that $J_{\lambda, \psi}(A_{\lambda, \psi}(g), g) = 0$. In particular, the above uniqueness result gives that $A_{\lambda, \psi}(g) = A(g) \circ \Xi_{\lambda, \psi}$. We conclude that $w_{\lambda, \psi} \in \bar{X}(p)$ depends analytically on λ and ψ in a neighborhood of $(1, 0) \in \mathbb{R} \times \mathbb{R}^{N-1}$. In particular, there exists a constant Λ dependent only on N such that for any $k \in \mathbb{N}_0$ and $\beta' \in \mathbb{N}_0^{N-1}$, it holds that

$$\|\partial_\lambda^k \partial_\psi^{\beta'}|_{(\lambda, \psi)=(1,0)} w_{\lambda, \psi}\|_{\bar{X}(p)} \lesssim \Lambda^{-k-|\beta'|} k! \beta'! \|g\|_{W^{1,\infty}}.$$

It remains to notice that

$$\partial_\lambda^k \partial_\psi^{\beta'}|_{(\lambda, \psi)=(1,0)} w_{\lambda, \psi}(t, z) = t^{k+|\beta'|} \partial_t^k \partial_\phi^{\beta'} w(t, r, \phi)$$

to deduce

$$t^{k+|\beta'|} |\partial_t^k \partial_\phi^{\beta'} \nabla w(t, r, \phi)| \lesssim \Lambda^{-k-|\beta'|} k! \beta'! \|g\|_{W^{1,\infty}}. \tag{52}$$

Step 4: regularity in transversal direction. The derivation of the transversal regularity relies on the analyticity bounds established above together with the Morrey estimate

$$\begin{aligned} \|v\|_{L^\infty(Q_r^d(z))} &\lesssim |Q_r^d(z)|_\sigma^{-1/p} \|v\|_{L_\sigma^p(Q_r^d(z))} \\ &\quad + r\theta |Q_r^d(z)|_\sigma^{-1/p} \|\nabla v\|_{L_\sigma^p(Q_r^d(z))} + r^4 |Q_r^d(z)|_\sigma^{-1/p} \|\partial_t v\|_{L_\sigma^p(Q_r^d(z))}, \end{aligned} \tag{53}$$

which holds for any $p > N$ uniformly in r and z . The proof of this estimate proceeds analogously to the Euclidean case; see, e.g., [Evans and Gariepy 1992, Chapter 4.5]. We omit the argument.

In the following discussion, we keep r and z fixed and we set $\theta = \theta(r, z)$. For $b \in \{2, 3\}$, we choose $\sigma = (b - 1)p$ and apply (53) to the effect that

$$\begin{aligned} \|\nabla_\psi^{4-b} \partial_s^b w\|_{L^\infty(Q_r^d(z))} &\lesssim |Q_r^d(z)|_{(b-1)p}^{-1/p} \|\rho^{b-1} \nabla_\psi^{4-b} \partial_s^b w\|_{L^p(Q_r^d(z))} \\ &\quad + r\theta |Q_r^d(z)|_{(b-1)p}^{-1/p} \|\rho^{b-1} \nabla_\psi^{4-b} \partial_r^b \nabla w\|_{L^p(Q_r^d(z))} \\ &\quad + r^4 |Q_r^d(z)|_{(b-1)p}^{-1/p} \|\rho^{b-1} \nabla_\psi^{4-b} \partial_r^b \partial_t w\|_{L^p(Q_r^d(z))}. \end{aligned}$$

We recall from (15) that $|Q_r^d(z)|_\sigma \sim \theta^{2\sigma} |Q_r^d(z)|$ and that $\sqrt{\rho(\tilde{z})} \lesssim \theta$ for any $\tilde{z} \in B_r^d(z)$ by virtue of (13). Therefore,

$$\begin{aligned} \|\nabla_\psi^{4-b} \partial_s^b w\|_{L^\infty(Q_r^d(z))} &\lesssim \theta^{4-2b} |Q_r^d(z)|^{-1/p} \|\rho^{b-2} \nabla_\psi^{4-b} \partial_s^b w\|_{L^p(Q_r^d(z))} \\ &\quad + r \theta^{3-2b} |Q_r^d(z)|^{-1/p} \|\rho^{b-1} \nabla_\psi^{4-b} \partial_r^b \nabla w\|_{L^p(Q_r^d(z))} \\ &\quad + r^4 \theta^{4-2b} |Q_r^d(z)|^{-1/p} \|\rho^{b-2} \nabla_\psi^{4-b} \partial_r^b \partial_t w\|_{L^p(Q_r^d(z))}. \end{aligned}$$

With the help of the analyticity estimates (52), we easily deduce that

$$r \theta \|t^2 \nabla_\psi^2 \partial_s^2 w\|_{L^\infty(Q_r^d(z))} + (r \theta)^2 \|t \nabla_\psi \partial_s^3 w\|_{L^\infty(Q_r^d(z))} \lesssim \|g\|_{W^{1,\infty}}. \tag{54}$$

An analogous argument yields the corresponding control of the time derivatives, namely

$$\frac{r^3}{\theta} \|\partial_t w\|_{L^\infty(Q_r^d(z))} \lesssim \|g\|_{W^{1,\infty}}. \tag{55}$$

In order to deduce control over the fourth-order vertical derivatives, we rewrite the nonlinear equation (6) as

$$\rho \partial_r^2 (\rho \partial_r^2 w) = f[w] - \partial_t w + \text{l.o.t.}$$

The terms on the right-hand side are all uniformly controlled thanks to (52), (54) and (55). Similarly, we may write

$$-\rho^{-1} \partial_r (\rho^2 \partial_r v) = h$$

for some h such that $t^\kappa h \in L^\infty$ for some $\kappa > 0$, and where $v = -\rho^{-1} \partial_r (\rho^2 \partial_r w)$. This identity can be integrated so that

$$\partial_r v = \rho^{-2} \int_r^1 \rho h \, d\tilde{r}.$$

The expression on the right is differentiable with

$$\partial_r^2 v = 2\rho^{-3} r \int_r^1 \rho h \, d\tilde{r} - \rho^{-1} h.$$

We deduce that $\rho t^\kappa \partial_r^2 v \in L^\infty$ and thus $\rho^2 t^\kappa \partial_r^4 w \in L^\infty$.

This argument can be iterated and yields the smoothness of w . □

Appendix: Derivation of the transformed equation

Let us write $z = \Phi_t(x)$. We will first verify that Φ defines a diffeomorphism. For this purpose, we compute the derivatives of Φ in terms of x and v ,

$$\partial_i \Phi^j = \frac{\delta_{ij}}{(2v + |x|^2)^{1/2}} - \frac{x_j (\partial_i v + x_i)}{(2v + |x|^2)^{3/2}}.$$

Recalling the elementary formula $\det(I - a \otimes b) = I - a \cdot b$ for any two vectors a and b , we compute that

$$\det \nabla \phi(x) = \frac{2v - x \cdot \nabla v}{(2v + |x|^2)^{N/2+1}}.$$

If v is close to the Smyth–Hill solution in the sense that

$$\|v - v_*\|_{L^\infty(\mathcal{P}(v))} + \|\nabla v + x\|_{L^\infty(\mathcal{P}(v))} \leq \varepsilon$$

for some small ε , we find that $2v - x \cdot \nabla v \geq 1 - 3\varepsilon$ and $2v + |x|^2 \geq 1 - 2\varepsilon$, which implies that the Jacobi determinant is finite if ε is sufficiently small.

Let us express the derivative of Φ in terms of the new variables z and w . Differentiating (5) yields

$$\partial_i v + x_i = (1 + w)\nabla w \cdot \partial_i \Phi = \partial_i w - \frac{z \cdot \nabla w}{1 + w}(\partial_i v + x_i),$$

and thus

$$\partial_i v + x_i = \frac{1 + w}{1 + w + z \cdot \nabla w} \partial_i w.$$

Plugging this and (5) into the expression for the derivatives of Φ , we find

$$\partial_i \Phi^j = \frac{\delta_{ij}}{1 + w} - \frac{z_j \partial_i w}{(1 + w)(1 + w + z \cdot \nabla w)}.$$

Under the assumption that w is such that

$$\|w\|_{L^\infty} + \|\nabla w\|_{L^\infty} \leq \varepsilon$$

for some small ε , we see by a calculation similar as the one above that Φ is a diffeomorphism.

We will now compute how the change of variables acts on the confined thin-film equation (3). For notational convenience, we set

$$\rho(z) = \frac{1}{2}(1 - |z|^2),$$

and $\tilde{w} = 1 + w$, with the effect that

$$\rho \tilde{w}^2 = v = \gamma u^{1/2}. \tag{56}$$

For an arbitrary function $f = f(z)$, it thus holds that

$$\partial_i (f(\Phi)) = \frac{\partial_i f}{\tilde{w}} - \frac{(z \cdot \nabla f) \partial_i \tilde{w}}{\tilde{w}(\tilde{w} + z \cdot \nabla \tilde{w})}. \tag{57}$$

Now, differentiating (56) with respect to x_i yields

$$\begin{aligned} \gamma^2 \partial_i u &= \frac{1}{\tilde{w}} \partial_i (\rho^2 \tilde{w}^4) - \frac{\partial_i \tilde{w}}{\tilde{w}(\tilde{w} + z \cdot \nabla \tilde{w})} z \cdot \nabla (\rho^2 \tilde{w}^4) \\ &= -2\rho \tilde{w}^3 z_i + 2 \frac{\rho \tilde{w}^3 \partial_i \tilde{w}}{\tilde{w} + z \cdot \nabla \tilde{w}}. \end{aligned}$$

Differentiating with respect to x_i again, we obtain that

$$\begin{aligned} \frac{\gamma^2}{2} \partial_i^2 u &= -(\rho - z_i^2) \tilde{w}^2 + \frac{\tilde{w}^2}{\tilde{w} + z \cdot \nabla \tilde{w}} \rho^{-1} \partial_i (\rho^2 \partial_i \tilde{w}) \\ &\quad - \frac{\tilde{w}^2}{\tilde{w} + z \cdot \nabla \tilde{w}} \rho z \cdot \nabla \left(\frac{(\partial_i \tilde{w})^2}{\tilde{w} + z \cdot \nabla \tilde{w}} \right) + (\rho + |z|^2) \frac{\tilde{w}^2 (\partial_i \tilde{w})^2}{(\tilde{w} + z \cdot \nabla \tilde{w})^2}. \end{aligned}$$

Hence, summing over i and rearranging terms yields

$$\frac{\gamma^2}{2} \Delta u \times \frac{\tilde{w} + z \cdot \nabla \tilde{w}}{\tilde{w}^2} = (1 - (N + 2)\rho)(\tilde{w} + z \cdot \nabla \tilde{w}) - \mathcal{L}\tilde{w} + (1 - \rho) \frac{|\nabla \tilde{w}|^2}{\tilde{w} + z \cdot \nabla \tilde{w}} - \rho z \cdot \nabla \left(\frac{|\nabla \tilde{w}|^2}{\tilde{w} + z \cdot \nabla \tilde{w}} \right).$$

With the help of the \star -notation, the (nonlinear) term in the second line of the above identity can be rewritten as

$$p \star \sum_{k=1}^2 \frac{(\nabla \tilde{w})^{(k-1)\star}}{(\tilde{w} + z \cdot \nabla \tilde{w})^k} ((\nabla \tilde{w})^{2\star} + \rho \nabla \tilde{w} \star \nabla^2 \tilde{w}).$$

In what follows, it should become clear why this way of writing drastically simplifies the notation.

With the help of (57), we compute

$$\partial_i \left(\left(\frac{\tilde{w}^2}{\tilde{w} + z \cdot \nabla \tilde{w}} f \right) (\Phi) \right) = \frac{\tilde{w}}{\tilde{w} + z \cdot \nabla \tilde{w}} \left(\partial_i f - z \cdot \nabla \left(\frac{\partial_i \tilde{w} f}{\tilde{w} + z \cdot \nabla \tilde{w}} \right) \right)$$

for any function $f = f(z)$, and thus

$$\begin{aligned} \frac{\gamma^2}{2} (\partial_i \Delta u - x_i) \times \frac{\tilde{w} + z \cdot \nabla \tilde{w}}{\tilde{w}} &= -N \partial_i \tilde{w} - \partial_i \mathcal{L}\tilde{w} \\ &+ p \star \frac{(\nabla \tilde{w})^{2\star} + \nabla \tilde{w} \star \nabla^2 \tilde{w} + \rho (\nabla^2 \tilde{w})^{2\star} + \rho \nabla \tilde{w} \star \nabla^3 \tilde{w}}{\tilde{w} + z \cdot \nabla \tilde{w}} \\ &+ p \star \frac{(\nabla \tilde{w})^{3\star} + (\nabla \tilde{w})^{2\star} \star \nabla^2 \tilde{w} + \rho \nabla \tilde{w} \star (\nabla^2 \tilde{w})^{2\star} + \rho (\nabla \tilde{w})^{2\star} \star \nabla^3 \tilde{w}}{(\tilde{w} + z \cdot \nabla \tilde{w})^2} \\ &+ p \star \frac{(\nabla \tilde{w})^{4\star} + (\nabla \tilde{w})^{3\star} \star \nabla^2 \tilde{w} + \rho (\nabla \tilde{w})^{2\star} \star (\nabla^2 \tilde{w})^{2\star} + \rho (\nabla \tilde{w})^{3\star} \star \nabla^3 \tilde{w}}{(\tilde{w} + z \cdot \nabla \tilde{w})^3} \\ &+ p \star \frac{(\nabla \tilde{w})^{5\star} + \rho (\nabla \tilde{w})^{4\star} \star \nabla^2 \tilde{w} + \rho (\nabla \tilde{w})^{3\star} \star (\nabla^2 \tilde{w})^{2\star}}{(\tilde{w} + z \cdot \nabla \tilde{w})^4}. \end{aligned}$$

We notice that the nonlinearity belongs to the class

$$p \star \sum_{k=1}^4 \frac{(\nabla \tilde{w})^{(k-1)\star}}{(\tilde{w} + z \cdot \nabla \tilde{w})^k} \star ((\nabla \tilde{w})^{2\star} + \nabla \tilde{w} \star \nabla^2 \tilde{w} + \rho (\nabla^2 \tilde{w})^{2\star} + \rho \nabla \tilde{w} \star \nabla^3 \tilde{w}).$$

Similarly to the above, we compute for an arbitrary function $f = f(z)$ that

$$\partial_i \left(\left(\frac{\tilde{w}^5}{\tilde{w} + z \cdot \nabla \tilde{w}} f \right) (\Phi) \right) = \frac{\tilde{w}^4}{\tilde{w} + z \cdot \nabla \tilde{w}} \left(\partial_i f + 3 \frac{\partial_i \tilde{w} f}{\tilde{w} + z \cdot \nabla \tilde{w}} - z \cdot \nabla \left(\frac{\partial_i \tilde{w} f}{\tilde{w} + z \cdot \nabla \tilde{w}} \right) \right),$$

and thus

$$\begin{aligned} \frac{\gamma^4}{2} \nabla \cdot (u \nabla \Delta u - u x) \times \frac{\tilde{w} + z \cdot \nabla \tilde{w}}{\rho \tilde{w}^4} &= (N + \mathcal{L}) \mathcal{L}\tilde{w} \\ &+ p \star \tilde{R}_{-1}[\tilde{w}] \star ((\nabla \tilde{w})^{2\star} + \nabla \tilde{w} \star \nabla^2 \tilde{w}) \\ &+ p \star \tilde{R}_{-1}[\tilde{w}] \star \rho ((\nabla^2 \tilde{w})^{2\star} + \nabla \tilde{w} \star \nabla^3 \tilde{w}) \\ &+ p \star \tilde{R}_{-2}[\tilde{w}] \star \rho^2 ((\nabla^2 \tilde{w})^{3\star} + \nabla \tilde{w} \star \nabla^2 \tilde{w} \star \nabla^3 \tilde{w} + (\nabla \tilde{w})^{2\star} \star \nabla^4 \tilde{w}), \end{aligned}$$

where $\tilde{R}_i[\tilde{w}] = r_i(\nabla\tilde{w}, \tilde{w} + z \cdot \nabla\tilde{w})$ for some rational functions r_i that are homogeneous of degree i , i.e., $r_i(sa, sb) = s^i r_i(a, b)$.

We finally turn to the computation of the time derivative. For this notice first that

$$\partial_t \Phi_t(x) = -\frac{\gamma^2}{2} \frac{z}{\rho\tilde{w}^4} \partial_t u,$$

and thus, a short computation shows that

$$\frac{\gamma^2}{2} \partial_t u = \frac{\rho\tilde{w}^4}{\tilde{w} + z \cdot \nabla\tilde{w}} \partial_t \tilde{w}.$$

After a rescaling of time $t \rightarrow \gamma^2 t$, and recalling that $\tilde{w} = 1 + w$, we find the transformed equation (6).

Acknowledgements

The author thanks Herbert Koch for helpful discussions. The present work was done when the author was affiliated with the Universität Bonn.

References

- [Angenent 1990] S. B. Angenent, “Nonlinear analytic semiflows”, *Proc. Roy. Soc. Edinburgh Sect. A* **115**:1-2 (1990), 91–107. [MR](#) [Zbl](#)
- [Belgacem et al. 2016] F. B. Belgacem, M. V. Gnann, and C. Kuehn, “A dynamical systems approach for the contact-line singularity in thin-film flows”, *Nonlinear Anal.* **144** (2016), 204–235. [MR](#) [Zbl](#)
- [Bellaïche and Risler 1996] A. Bellaïche and J.-J. Risler (editors), *Sub-Riemannian geometry*, Progress in Mathematics **144**, Birkhäuser, Basel, 1996. [MR](#) [Zbl](#)
- [Beretta et al. 1995] E. Beretta, M. Bertsch, and R. Dal Passo, “Nonnegative solutions of a fourth-order nonlinear degenerate parabolic equation”, *Arch. Rational Mech. Anal.* **129**:2 (1995), 175–200. [MR](#) [Zbl](#)
- [Bernis 1996] F. Bernis, “Finite speed of propagation and continuity of the interface for thin viscous flows”, *Adv. Differential Equations* **1**:3 (1996), 337–368. [MR](#) [Zbl](#)
- [Bernis and Friedman 1990] F. Bernis and A. Friedman, “Higher order nonlinear degenerate parabolic equations”, *J. Differential Equations* **83**:1 (1990), 179–206. [MR](#) [Zbl](#)
- [Bertozzi and Pugh 1996] A. L. Bertozzi and M. Pugh, “The lubrication approximation for thin viscous films: regularity and long-time behavior of weak solutions”, *Comm. Pure Appl. Math.* **49**:2 (1996), 85–123. [MR](#) [Zbl](#)
- [Bertsch et al. 1998] M. Bertsch, R. Dal Passo, H. Garcke, and G. Grün, “The thin viscous flow equation in higher space dimensions”, *Adv. Differential Equations* **3**:3 (1998), 417–440. [MR](#) [Zbl](#)
- [Carrillo and Toscani 2002] J. A. Carrillo and G. Toscani, “Long-time asymptotics for strong solutions of the thin film equation”, *Comm. Math. Phys.* **225**:3 (2002), 551–571. [MR](#) [Zbl](#)
- [Coifman and Weiss 1971] R. R. Coifman and G. Weiss, *Analyse harmonique non-commutative sur certains espaces homogènes*, Lecture Notes in Mathematics **242**, Springer, 1971. [MR](#) [Zbl](#)
- [Dal Passo et al. 1998] R. Dal Passo, H. Garcke, and G. Grün, “On a fourth-order degenerate parabolic equation: global entropy estimates, existence, and qualitative behavior of solutions”, *SIAM J. Math. Anal.* **29**:2 (1998), 321–342. [MR](#) [Zbl](#)
- [Daskalopoulos and Hamilton 1998] P. Daskalopoulos and R. Hamilton, “Regularity of the free boundary for the porous medium equation”, *J. Amer. Math. Soc.* **11**:4 (1998), 899–965. [MR](#) [Zbl](#)
- [Degtyarev 2017] S. Degtyarev, “Classical solvability of the multidimensional free boundary problem for the thin film equation with quadratic mobility in the case of partial wetting”, *Discrete Contin. Dyn. Syst.* **37**:7 (2017), 3625–3699. [MR](#) [Zbl](#)

- [Deimling 1985] K. Deimling, *Nonlinear functional analysis*, Springer, 1985. [MR](#) [Zbl](#)
- [Denzler et al. 2015] J. Denzler, H. Koch, and R. J. McCann, *Higher-order time asymptotics of fast diffusion in Euclidean space: a dynamical systems approach*, Mem. Amer. Math. Soc. **1101**, Amer. Math. Soc., Providence, RI, 2015. [MR](#) [Zbl](#)
- [Evans and Gariepy 1992] L. C. Evans and R. F. Gariepy, *Measure theory and fine properties of functions*, CRC Press, Boca Raton, FL, 1992. [MR](#) [Zbl](#)
- [Ferreira and Bernis 1997] R. Ferreira and F. Bernis, “Source-type solutions to thin-film equations in higher dimensions”, *European J. Appl. Math.* **8**:5 (1997), 507–524. [MR](#) [Zbl](#)
- [Giacomelli and Knüpfer 2010] L. Giacomelli and H. Knüpfer, “A free boundary problem of fourth order: classical solutions in weighted Hölder spaces”, *Comm. Partial Differential Equations* **35**:11 (2010), 2059–2091. [MR](#) [Zbl](#)
- [Giacomelli and Otto 2001] L. Giacomelli and F. Otto, “Variational formulation for the lubrication approximation of the Hele-Shaw flow”, *Calc. Var. Partial Differential Equations* **13**:3 (2001), 377–403. [MR](#) [Zbl](#)
- [Giacomelli and Otto 2003] L. Giacomelli and F. Otto, “Rigorous lubrication approximation”, *Interfaces Free Bound.* **5**:4 (2003), 483–529. [MR](#) [Zbl](#)
- [Giacomelli et al. 2008] L. Giacomelli, H. Knüpfer, and F. Otto, “Smooth zero-contact-angle solutions to a thin-film equation around the steady state”, *J. Differential Equations* **245**:6 (2008), 1454–1506. [MR](#) [Zbl](#)
- [Giacomelli et al. 2013] L. Giacomelli, M. V. Gnann, and F. Otto, “Regularity of source-type solutions to the thin-film equation with zero contact angle and mobility exponent between $\frac{3}{2}$ and 3”, *European J. Appl. Math.* **24**:5 (2013), 735–760. [MR](#) [Zbl](#)
- [Giacomelli et al. 2014] L. Giacomelli, M. V. Gnann, H. Knüpfer, and F. Otto, “Well-posedness for the Navier-slip thin-film equation in the case of complete wetting”, *J. Differential Equations* **257**:1 (2014), 15–81. [MR](#) [Zbl](#)
- [Gnann 2015] M. V. Gnann, “Well-posedness and self-similar asymptotics for a thin-film equation”, *SIAM J. Math. Anal.* **47**:4 (2015), 2868–2902. [MR](#) [Zbl](#)
- [Gnann 2016] M. V. Gnann, “On the regularity for the Navier-slip thin-film equation in the perfect wetting regime”, *Arch. Ration. Mech. Anal.* **222**:3 (2016), 1285–1337. [MR](#) [Zbl](#)
- [Grigor’yan 2006] A. Grigor’yan, “Heat kernels on weighted manifolds and applications”, pp. 93–191 in *The ubiquitous heat kernel*, edited by J. Jorgenson and L. Walling, Contemp. Math. **398**, Amer. Math. Soc., Providence, RI, 2006. [MR](#) [Zbl](#)
- [John 2015] D. John, “On uniqueness of weak solutions for the thin-film equation”, *J. Differential Equations* **259**:8 (2015), 4122–4171. [MR](#) [Zbl](#)
- [Kienzler 2016] C. Kienzler, “Flat fronts and stability for the porous medium equation”, *Comm. Partial Differential Equations* **41**:12 (2016), 1793–1838. [MR](#) [Zbl](#)
- [Knüpfer 2011] H. Knüpfer, “Well-posedness for the Navier slip thin-film equation in the case of partial wetting”, *Comm. Pure Appl. Math.* **64**:9 (2011), 1263–1296. [MR](#) [Zbl](#)
- [Knüpfer 2015] H. Knüpfer, “Well-posedness for a class of thin-film equations with general mobility in the regime of partial wetting”, *Arch. Ration. Mech. Anal.* **218**:2 (2015), 1083–1130. [MR](#) [Zbl](#)
- [Knüpfer and Masmoudi 2015] H. Knüpfer and N. Masmoudi, “Darcy’s flow with prescribed contact angle: well-posedness and lubrication approximation”, *Arch. Ration. Mech. Anal.* **218**:2 (2015), 589–646. [MR](#) [Zbl](#)
- [Koch 1999] H. Koch, *Non-Euclidean singular integrals and the porous medium equation*, habilitation thesis, Universität Heidelberg, 1999, available at www.math.uni-bonn.de/people/koch/habil.ps.gz.
- [Koch 2004] H. Koch, “Partial differential equations and singular integrals”, pp. 59–122 in *Dispersive nonlinear problems in mathematical physics*, edited by P. D’Ancona and V. Georgev, Quad. Mat. **15**, Dept. Math., Seconda Univ. Napoli, Caserta, 2004. [MR](#) [Zbl](#)
- [Koch 2008] H. Koch, “Partial differential equations with non-Euclidean geometries”, *Discrete Contin. Dyn. Syst. Ser. S* **1**:3 (2008), 481–504. [MR](#) [Zbl](#)
- [Koch and Lamm 2012] H. Koch and T. Lamm, “Geometric flows with rough initial data”, *Asian J. Math.* **16**:2 (2012), 209–235. [MR](#) [Zbl](#)
- [Matthes et al. 2009] D. Matthes, R. J. McCann, and G. Savaré, “A family of nonlinear fourth order equations of gradient flow type”, *Comm. Partial Differential Equations* **34**:10–12 (2009), 1352–1397. [MR](#) [Zbl](#)

- [McCann and Seis 2015] R. J. McCann and C. Seis, “The spectrum of a family of fourth-order nonlinear diffusions near the global attractor”, *Comm. Partial Differential Equations* **40**:2 (2015), 191–218. [MR](#) [Zbl](#)
- [Oron et al. 1997] A. Oron, S. H. Davis, and S. G. Bankoff, “Long-scale evolution of thin liquid films”, *Rev. Mod. Phys.* **69** (1997), 931–980.
- [Otto 1998] F. Otto, “Lubrication approximation with prescribed nonzero contact angle”, *Comm. Partial Differential Equations* **23**:11-12 (1998), 2077–2164. [MR](#) [Zbl](#)
- [Otto 2001] F. Otto, “The geometry of dissipative evolution equations: the porous medium equation”, *Comm. Partial Differential Equations* **26**:1-2 (2001), 101–174. [MR](#) [Zbl](#)
- [Seis 2014] C. Seis, “Long-time asymptotics for the porous medium equation: the spectrum of the linearized operator”, *J. Differential Equations* **256**:3 (2014), 1191–1223. [MR](#) [Zbl](#)
- [Seis 2015] C. Seis, “Invariant manifolds for the porous medium equation”, preprint, 2015. [arXiv](#)
- [Seis \geq 2018] C. Seis, “Invariant manifolds for the thin film equation”, work in preparation.
- [Smyth and Hill 1988] N. F. Smyth and J. M. Hill, “High-order nonlinear diffusion”, *IMA J. Appl. Math.* **40**:2 (1988), 73–86. [MR](#) [Zbl](#)
- [Stein 1970] E. M. Stein, *Singular integrals and differentiability properties of functions*, Princeton Mathematical Series **30**, Princeton University Press, 1970. [MR](#) [Zbl](#)
- [Stein 1993] E. M. Stein, *Harmonic analysis: real-variable methods, orthogonality, and oscillatory integrals*, Princeton Mathematical Series **43**, Princeton University Press, 1993. [MR](#) [Zbl](#)

Received 5 Sep 2017. Accepted 2 Jan 2018.

CHRISTIAN SEIS: seis@wwu.de

Institut für Analysis und Numerik, Westfälische Wilhelms-Universität Münster, Münster, Germany

Analysis & PDE

msp.org/apde

EDITORS

EDITOR-IN-CHIEF

Patrick Gérard

patrick.gerard@math.u-psud.fr

Université Paris Sud XI

Orsay, France

BOARD OF EDITORS

Massimiliano Berti	Scuola Intern. Sup. di Studi Avanzati, Italy berti@sissa.it	Clément Mouhot	Cambridge University, UK c.mouhot@dpmms.cam.ac.uk
Sun-Yung Alice Chang	Princeton University, USA chang@math.princeton.edu	Werner Müller	Universität Bonn, Germany mueller@math.uni-bonn.de
Michael Christ	University of California, Berkeley, USA mchrist@math.berkeley.edu	Gilles Pisier	Texas A&M University, and Paris 6 pisier@math.tamu.edu
Alessio Figalli	ETH Zurich, Switzerland alessio.figalli@math.ethz.ch	Tristan Rivière	ETH, Switzerland riviere@math.ethz.ch
Charles Fefferman	Princeton University, USA cf@math.princeton.edu	Igor Rodnianski	Princeton University, USA irod@math.princeton.edu
Ursula Hamenstaedt	Universität Bonn, Germany ursula@math.uni-bonn.de	Sylvia Serfaty	New York University, USA serfaty@cims.nyu.edu
Vaughan Jones	U.C. Berkeley & Vanderbilt University vaughan.f.jones@vanderbilt.edu	Yum-Tong Siu	Harvard University, USA siu@math.harvard.edu
Vadim Kaloshin	University of Maryland, USA vadim.kaloshin@gmail.com	Terence Tao	University of California, Los Angeles, USA tao@math.ucla.edu
Herbert Koch	Universität Bonn, Germany koch@math.uni-bonn.de	Michael E. Taylor	Univ. of North Carolina, Chapel Hill, USA met@math.unc.edu
Izabella Laba	University of British Columbia, Canada ilaba@math.ubc.ca	Gunther Uhlmann	University of Washington, USA gunther@math.washington.edu
Gilles Lebeau	Université de Nice Sophia Antipolis, France lebeau@unice.fr	András Vasy	Stanford University, USA andras@math.stanford.edu
Richard B. Melrose	Massachusetts Inst. of Tech., USA rbb@math.mit.edu	Dan Virgil Voiculescu	University of California, Berkeley, USA dvv@math.berkeley.edu
Frank Merle	Université de Cergy-Pontoise, France Frank.Merle@u-cergy.fr	Steven Zelditch	Northwestern University, USA zelditch@math.northwestern.edu
William Minicozzi II	Johns Hopkins University, USA minicozz@math.jhu.edu	Maciej Zworski	University of California, Berkeley, USA zvorski@math.berkeley.edu

PRODUCTION

production@msp.org

Silvio Levy, Scientific Editor


See inside back cover or msp.org/apde for submission instructions.

The subscription price for 2018 is US \$275/year for the electronic version, and \$480/year (+\$55, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues from the last three years and changes of subscriber address should be sent to MSP.

Analysis & PDE (ISSN 1948-206X electronic, 2157-5045 printed) at Mathematical Sciences Publishers, 798 Evans Hall #3840, c/o University of California, Berkeley, CA 94720-3840, is published continuously online. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices.

APDE peer review and production are managed by EditFlow[®] from MSP.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2018 Mathematical Sciences Publishers

ANALYSIS & PDE

Volume 11 No. 5 2018

Large sets avoiding patterns	1083
ROBERT FRASER and MALABIKA PRAMANIK	
On minimizers of an isoperimetric problem with long-range interactions under a convexity constraint	1113
MICHAEL GOLDMAN, MATTEO NOVAGA and BERARDO RUFFINI	
Nonautonomous maximal L^p -regularity under fractional Sobolev regularity in time	1143
STEPHAN FACKLER	
Transference of bilinear restriction estimates to quadratic variation norms and the Dirac–Klein–Gordon system	1171
TIMOTHY CANDY and SEBASTIAN HERR	
Well-posedness and smoothing effect for generalized nonlinear Schrödinger equations	1241
PIERRE-YVES BIENAIMÉ and ABDESSLAM BOULKHEMAIR	
The shape of low energy configurations of a thin elastic sheet with a single disclination	1285
HEINER OLBERMANN	
The thin-film equation close to self-similarity	1303
CHRISTIAN SEIS	