

ANALYSIS & PDE

Volume 11

No. 7

2018

SYLVIA SERFATY AND JOAQUIM SERRA

**QUANTITATIVE STABILITY OF THE FREE BOUNDARY
IN THE OBSTACLE PROBLEM**

QUANTITATIVE STABILITY OF THE FREE BOUNDARY IN THE OBSTACLE PROBLEM

SYLVIA SERFATY AND JOAQUIM SERRA

We prove some detailed quantitative stability results for the contact set and the solution of the classical obstacle problem in \mathbb{R}^n ($n \geq 2$) under perturbations of the obstacle function, which is also equivalent to studying the variation of the equilibrium measure in classical potential theory under a perturbation of the external field.

To do so, working in the setting of the whole space, we examine the evolution of the free boundary Γ^t corresponding to the boundary of the contact set for a family of obstacle functions h^t . Assuming that $h = h^t(x) = h(t, x)$ is $C^{k+1, \alpha}$ in $[-1, 1] \times \mathbb{R}^n$ and that the initial free boundary Γ^0 is regular, we prove that Γ^t is twice differentiable in t in a small neighborhood of $t = 0$. Moreover, we show that the “normal velocity” and the “normal acceleration” of Γ^t are respectively $C^{k-1, \alpha}$ and $C^{k-2, \alpha}$ scalar fields on Γ^t . This is accomplished by deriving equations for this velocity and acceleration and studying the regularity of their solutions via single- and double-layer estimates from potential theory.

1. Introduction

Motivation of the problem. Consider the classical obstacle problem; see for instance [Kinderlehrer and Nirenberg 1977; Caffarelli 1998]. If the obstacle h is perturbed into $h + t\xi$ with t small and ξ regular enough, how much does the contact set (or coincidence set) move? The best known answer to this question is in [Blank 2001], where it is proved that the new contact set is $O(t)$ -close to the old one in Hausdorff distance, in the setting of a bounded domain with Dirichlet boundary condition. Some results are also proved in [Schaeffer 1975] in an analytic setting, by Nash–Moser inversion.

Our paper is concerned with getting stronger and more quantitative stability estimates, in particular obtaining closeness of the contact sets in $C^{k, \alpha}$ norms with explicitly described first and second derivatives with respect to t , which come together with an explicit asymptotic expansion of the solution itself. We believe that such results are of natural and independent interest for the obstacle problem. They are also for us motivated by an application on the analysis of Coulomb systems in statistical mechanics from [Leblé and Serfaty 2018], which relies on the present paper.

Let us get into more detail on this aspect. In potential theory, the so-called (Frostman) “equilibrium measure” for Coulomb interactions with an external “field” Q is the unique probability measure μ on \mathbb{R}^n which minimizes

$$\int_{\mathbb{R}^n \times \mathbb{R}^n} P(x - y) d\mu(x) d\mu(y) + \int_{\mathbb{R}^n} Q(x) d\mu(x), \quad (1-1)$$

MSC2010: 35R35, 31B35, 49K99.

Keywords: obstacle problem, contact set, coincidence set, stability, equilibrium measure, potential theory.

where P is the Newtonian potential in dimension n . If Q grows fast enough at infinity, then setting

$$u(x) = \int_{\mathbb{R}^n} P(x-y) d\mu(y), \quad (1-2)$$

the equilibrium measure μ is compactly supported and uniquely characterized by the fact that there exists a constant c such that

$$u \geq c - \frac{1}{2}Q \quad \text{and} \quad u = c - \frac{1}{2}Q \quad \mu - \text{a.e.};$$

see for instance [Saff and Totik 1997]. We thus find that $\mu = -\Delta u$, where u solves the classical obstacle problem in the whole space

$$\min\{-\Delta u, u - h\} = 0$$

with obstacle $h = c - \frac{1}{2}Q$ —the two problems (identifying the equilibrium measure and solving the obstacle problem) are in fact convex dual minimization problems, as seen in [Ekeland and Temam 1976]; see for instance [Serfaty 2015, Chapter 2] for a description of this correspondence. Thus, the support of the equilibrium measure is equal to the contact set wherever the obstacle is “active”.

The understanding of the dependence of the equilibrium measure on the external field—which is thus equivalent to the understanding of the dependence of the solution and its contact set on the obstacle function—is crucial for the analysis of systems of particles with logarithmic or Coulomb interactions; in particular it allows one to show that the linear statistics of fluctuations of such systems converge to Gaussians. Following the method first introduced by [Johansson 1998], this relies on the computation of the Laplace transform of the fluctuations, which directly leads to considering the same system but with perturbed external field. Previously, the analysis of the perturbation of the equilibrium measure, as done in [Ameur et al. 2011], relied on Sakai’s theory [1991], a complex-analytic approach which is thus only valid in two dimensions and imposed analyticity assumptions on the external field and the boundary of the coincidence set.

In that context, the evolution of the contact sets sometimes goes by the name “Laplacian growth” or “Hele-Shaw flow” or the “Hele-Shaw equation”, see [Hedenmalm and Makarov 2004; 2013], and seems related to the quantum Hele-Shaw flow introduced by the physicists Wiegmann [2002] and Zabrodin. It has only been examined in dimension 2.

Setting of the study. Both for simplicity and for the applications we have in mind mentioned above, we consider global solutions to the obstacle problem in \mathbb{R}^n , $n \geq 2$. We note that the setting in \mathbb{R}^2 is slightly different than the setting in \mathbb{R}^n for $n \geq 3$ due to the fact that the logarithmic Newtonian potential does not decay to zero at infinity, and this will lead us to often making parallel statements about the two. We also note that the potential u associated to the equilibrium measure in (1-2) behaves like P at infinity, since μ is a compactly supported probability measure, i.e., tends to 0 if $n \geq 3$ and behaves like $-\frac{1}{2\pi} \log |x|$ if $n = 2$. Specifying the total mass of $-\Delta u$ is equivalent to specifying the ratio of $u/-\log |x|$ at infinity in dimension 2, or to adding an appropriate¹ constant to u in dimension $n \geq 3$.

¹Let u^t be defined as (1-3). For $n \geq 3$ there is a nonlinear (but monotone and continuous) relation between the mass $\int_{\mathbb{R}^n} \Delta u^t$ and value of the constant c^t . For c^t large enough, the mass is 0, and when c^t decreases, the mass increases continuously. This allows us to solve the equation with prescribed mass by varying the constant c^t .

With the above motivation, in order to consider the perturbations of the obstacle, we thus consider for each $t \in [-1, 1]$, given c^t a function of t , the function u^t solving the obstacle problem

$$\min\{-\Delta u^t, u^t - h^t\} = 0 \quad \text{in } \mathbb{R}^n, \quad \begin{cases} \lim_{|x| \rightarrow \infty} u^t(x) = c^t & (n \geq 3), \\ \lim_{|x| \rightarrow \infty} \frac{u^t(x)}{-\log|x|} = c^t & (n = 2). \end{cases} \quad (1-3)$$

We assume $\Delta h^0 < 0$ on $\{u^0 = h^0\}$, i.e., the obstacle must be “active” in the contact set, and

$$\begin{cases} \lim_{|x| \rightarrow \infty} h^t(x) < c^t & (n \geq 3), \\ \lim_{|x| \rightarrow \infty} \frac{h^t(x)}{-\log|x|} < c^t & (n = 2), \end{cases} \quad (1-4)$$

$$h = h^t(x) = h(t, x) \in C^{k-1, \alpha}([-1, 1] \times B_R), \quad (1-5)$$

while

$$c = c^t = c(t) \in C^2([-1, 1]). \quad (1-6)$$

For $n = 2$ we assume $c > 0$.

In addition, we assume

$$\Delta(h^t - h^0) \quad \text{is compactly supported in } B_R \quad (1-7)$$

and

$$h^t - h^0 \rightarrow 0 \quad \text{as } |x| \rightarrow \infty \quad (n \geq 3), \quad (1-8a)$$

$$\frac{h^t - h^0}{-\log|x|} \rightarrow 0 \quad \text{as } |x| \rightarrow \infty \quad (n = 2). \quad (1-8b)$$

In particular, letting $\dot{\cdot}$ denote the derivative with respect to t , this implies

$$\dot{h}^t \rightarrow 0 \quad \text{as } |x| \rightarrow \infty \quad (n \geq 3), \quad (1-9a)$$

$$\frac{\dot{h}^t}{-\log|x|} \rightarrow 0 \quad \text{as } |x| \rightarrow \infty \quad (n = 2). \quad (1-9b)$$

Let us denote by

$$\Omega^t := \{u^t - h^t > 0\} \quad \text{and} \quad \Gamma^t := \partial\Omega^t$$

the complement of the contact set and the free boundary, respectively.

We will assume that all points of the “initial” free boundary Γ^0 are regular points in the sense of Caffarelli [1977; 1998]. In particular we assume that Ω^0 is an open set with smooth boundary.

For the analysis of the paper it is convenient to identify precisely the quantities on which the (constants in the) estimates depend. To this aim, let us fix $\rho > 0$ and make the following quantitative assumptions.

First, we assume that, for some $U \subset B_R$, we have

$$\Delta h^0 \leq -\rho \quad \text{in } \bar{U} \quad \text{and} \quad \begin{cases} u^0 - h^0 \geq \rho & \text{in } \mathbb{R}^n \setminus U \quad (n \geq 3), \\ \frac{u^0 - h^0}{-\log|x|} \geq \rho & \text{in } \mathbb{R}^n \setminus U \quad (n = 2), \end{cases} \quad (1-10)$$

where $U \subset B_R$ is some open set containing $\{u^0 = 0\}$.

Second, we assume

$$\text{all points of } \Gamma^0 \text{ can be touched from both sides by balls of radius } \rho. \tag{1-11}$$

This is a quantitative version of the assumption that all points of Γ^0 are regular points.

Throughout the paper, if \mathcal{C} is a set of parameters of the problem, we denote by $C(\mathcal{C})$ a constant depending only on \mathcal{C} . We define

$$\mathcal{C} := \{n, k, \alpha, \mathbf{R}, \mathbf{U}, \rho, \|h\|_{C^{k+1,\alpha}([-1,1] \times \bar{U})}, \|c\|_{C^2([-1,1])}\}, \tag{1-12}$$

$$\mathcal{C}^0 := \{n, k, \alpha, \mathbf{R}, \mathbf{U}, \rho, \|h^0\|_{C^{k+1,\alpha}(\bar{U})}, c^0\}. \tag{1-13}$$

For $n = 2$ we also add to \mathcal{C} the constant $\inf_{[-1,1]} c > 0$.

Main result. Let $t_o > 0$ and let $\Psi = \Psi^t(x) = \Psi(t, x)$ be a 1-parameter family of diffeomorphisms $\Psi : (-t_o, t_o) \times \mathbb{R}^n \rightarrow \mathbb{R}^n$. We say that Ψ fixes the complement of \mathbf{U} if $\Psi(x) = x$ for all $x \in \mathbb{R}^n \setminus \mathbf{U}$.

We say Ψ is continuously differentiable if for all $t \in (-t_o, t_o)$ there exists $\dot{\Psi}^t \in C^0(\mathbb{R}^n; \mathbb{R}^n)$ such that

$$\begin{aligned} \|\Psi^{t+s}(x) - \Psi^t(x) - s\dot{\Psi}^t(x)\|_{C^0(\mathbb{R}^n; \mathbb{R}^n)} &= o(s), \\ \|\dot{\Psi}^{t+s}(x) - \dot{\Psi}^t(x)\|_{C^0(\mathbb{R}^n; \mathbb{R}^n)} &= o(1) \end{aligned}$$

as $s \rightarrow 0$.

We say Ψ is twice continuously differentiable if, in addition, for all $t \in (-t_o, t_o)$ there exists $\ddot{\Psi}^t \in C^0(\mathbb{R}^n; \mathbb{R}^n)$ such that

$$\begin{aligned} \|\Psi^{t+s}(x) - \Psi^t(x) - s\dot{\Psi}^t(x) - \frac{1}{2}s^2\ddot{\Psi}^t(x)\|_{C^0(\mathbb{R}^n; \mathbb{R}^n)} &= o(s^2), \\ \|\ddot{\Psi}^{t+s}(x) - \ddot{\Psi}^t(x)\|_{C^0(\mathbb{R}^n; \mathbb{R}^n)} &= o(1) \end{aligned}$$

as $s \rightarrow 0$.

Throughout the paper, given a function $f : (-t_o, t_o) \times Y \rightarrow \mathbb{R}$ we use the notation $f = f^t(x) = f(t, x)$,

$$\delta_t f^s := \frac{f^{s+t} - f^s}{t} \quad \text{and} \quad \dot{f}^s := \lim_{t \downarrow 0} \delta_t f^s = \partial_t f(s, y).$$

The main result of the paper is the following. In its statement, and throughout the paper, we denote by

$$v^t : \Gamma^t \rightarrow S^{n-1}$$

the unit normal vector to Γ^t pointing towards Ω^t .

Theorem 1.1. *Let $n \geq 2$, $k \geq 1$, $\alpha \in (0, 1)$, and u^t satisfy (1-3), with h and c satisfying (1-4)–(1-8). Assume (1-10) and (1-11) hold.*

Then, there exists $t_o > 0$ and a 1-parameter differentiable family of diffeomorphisms $\Psi^t \in C^{k,\alpha}(\mathbb{R}^n; \mathbb{R}^n)$ that fixes the complement of \mathbf{U} and which satisfies, for every $t \in (-t_o, t_o)$,

$$\begin{aligned} \Psi^t(\Omega^0) &= \Omega^t, \quad \Psi^t(\Gamma^0) = \Gamma^t, \\ \|\dot{\Psi}^t\|_{C^{k-1,\alpha}(\mathbb{R}^n)} &\leq C \quad \text{and} \quad (\dot{\Psi}^t \circ (\Psi^t)^{-1}) \cdot v^t = \frac{\partial_{v^t} V^t}{\Delta h^t} \quad \text{on } \Gamma^t, \end{aligned} \tag{1-14}$$

where $V^t := \dot{u}^t - \dot{h}^t$ is the solution² to

$$\begin{cases} \Delta V^t = -\Delta \dot{h}^t & \text{in } \Omega^t, \\ V^t = 0 & \text{on } \Gamma^t, \\ \lim_{x \rightarrow \infty} V^t(x) = \dot{c}^t & (n \geq 3), \end{cases} \quad \begin{cases} \Delta V^t = -\Delta \dot{h}^t & \text{in } \Omega^t, \\ V^t = 0 & \text{on } \Gamma^t, \\ \lim_{x \rightarrow \infty} \frac{V^t(x)}{-\log|x|} = \dot{c}^t & (n = 2). \end{cases} \quad (1-15)$$

In addition, we have

$$\dot{u}^t = \dot{h}^t + V^t \chi_{\Omega^t} \quad \text{in all } \mathbb{R}^n.$$

If moreover $k \geq 2$ then Ψ is twice differentiable and we have

$$\|\ddot{\Psi}^t\|_{C^{k-2,\alpha}(\mathbb{R}^n)} \leq C_\circ \quad (1-16)$$

and

$$\|\ddot{u}^t\|_{L^\infty(\mathbb{R}^n)} + \|\nabla \ddot{u}^t\|_{C^{k-2,\alpha}((\Omega^0 \cup \Omega^t)^c \cup \overline{\Omega^0 \cap \Omega^t})} \leq C_\circ. \quad (1-17)$$

The constants t_\circ and C_\circ depend only on³ \mathcal{C} .

An informal rephrasing of [Theorem 1.1](#) is as follows. If the moving obstacle $h(t, x)$ is $C^{k+1,\alpha}$ and $c(t)$ is C^2 , then Γ^t is “twice differentiable” for t in a small neighborhood of 0. Moreover, the “normal velocity” of Γ^t and the “normal acceleration” of Γ^t are respectively $C^{k-1,\alpha}$ and $C^{k-2,\alpha}$ scalar fields on Γ^t , with the normal velocity precisely identified via a Dirichlet-to-Neumann transformation: to compute it, one finds the solution V^t to the Dirichlet problem in a exterior domain (1-15) and the normal velocity at a point of Γ^t is given by the normal derivative of V^t divided by the Laplacian of the obstacle at that point.

Open questions. It is of course natural to ask whether similar results hold for more general obstacle problems, such as those associated to fully nonlinear operators or to fractional Laplacians.

In view of our results,⁴ a natural open question, which we believe to be delicate, is whether one can improve [Theorem 1.1](#) to

$$\Psi(t, x) \in C^{k,\alpha} \quad (\text{jointly in } t \text{ and } x).$$

Structure of the proof and organization of the paper. For the proof, we first reduce to a situation where the contact set is growing, i.e., $\Omega^t \subset \Omega^0$. We then define a coordinate system near the free boundary Γ^0 , and express the “height” η^t of Γ^t in these coordinates.

In [Section 3](#), assuming that an expansion of the type $\eta^t = \eta^0 + \dot{\eta}^0 t + \frac{1}{2} \ddot{\eta}^0 t^2 + \dots$ holds as $t \rightarrow 0$, we derive equations for $\dot{\eta}^0$ and $\ddot{\eta}^0$, which allow us to obtain explicit formulae and Hölder regularity for these quantities via single- and double-layer potential-theoretic estimates. These regularity estimates are delicate to obtain because the relations characterizing $\dot{\eta}^0$ and $\ddot{\eta}^0$ are at first implicit and one needs to show they can be “closed” for regularity.

²Since we assumed \dot{h}^t tends to 0 (resp. is $\ll |\log|x||$ if $n = 2$) at ∞ , we have V^t is the unique solution such that $V^t + \dot{h}^t$ is bounded, coincides with h^t in the complement of Ω^t and is harmonic in Ω^t . In fact, $V^t + \dot{h}^t$ is the unique bounded harmonic extension of \dot{h}^t outside of $(\Omega^t)^c$.

³The set of constants of the problem \mathcal{C} was defined in (1-12).

⁴We establish that if $h \in C^{k+1,\alpha}$ then $\Psi^t \in C^{k,\alpha}$, $\dot{\Psi}^t \in C^{k,\alpha}$ and $\ddot{\Psi}^t \in C^{k-2,\alpha}$.

In [Section 4](#), we show that the existence of an expansion in t for η^t , which was previously assumed, does hold. This is done by using a second set of adapted coordinates near Γ^0 (a sort of hodograph transform) and again single- and double-layer potential estimates.

Finally, in [Section 5](#) we prove the main result by showing how to treat the general case where the contact set is not necessarily growing. In the [Appendix](#), we collect the potential-theoretic estimates we need and some additional proofs.

2. Preliminaries

Known results. Throughout the paper it is useful to quantify the smoothness of the (boundaries of the) domains Ω^t . Let us introduce some more notation with that aim. Let U be some open set and $r > 0$. We write $\partial U \in C_r^{k,\alpha}$ if for all $x_o \in \partial U$ there are some orthonormal coordinates y_i , $1 \leq i \leq n$, with origin at x_o (these coordinates may vary from point to point), and a function $F_{x_o} \in C^{k,\alpha}(\bar{B}'_r)$ such that

$$U \cap \{|y'| < r, |y_n| < r\} = \{y_n < F_{x_o}(y')\} \cap \{|y'| < r, |y_n| < r\},$$

where $y' = (y_1, y_2, \dots, y_{n-1})$.

In this framework we define

$$\|\partial U\|_{C_r^{k,\alpha}} := \sup_{x_o \in \partial U} \|F_{x_o}\|_{C^{k,\alpha}(\bar{B}'_r)} < \infty, \tag{2-1}$$

where $B'_r = \{|y'| < r\} \subset \mathbb{R}^{n-1}$.

With the previous assumptions we have in our notation:

Proposition 2.1 [[Caffarelli 1977; 1998; Kinderlehrer and Nirenberg 1977; Blank 2001](#)]. *There exist universal constants $t_o > 0$ and C_o depending only on \mathcal{C} such that the following hold:*

(i) *We have*

$$\|\Gamma^t\|_{C_{\rho/4}^{k,\alpha}} \leq C_o \quad \text{for all } t \in (-t_o, t_o).$$

(ii) *For every pair $t, s \in (-t_o, t_o)$, the Hausdorff distance between Γ^t and Γ^s satisfies*

$$d_{\text{Hausdorff}}(\Gamma^t, \Gamma^s) \leq C_o |t - s|.$$

[Proposition 2.1](#) is contained in the results of [[Blank 2001](#)]. However, for the sake of completeness, we briefly sketch the proof in the [Appendix](#). This is done by combining the classical results for the obstacle problem in [[Caffarelli 1977; 1998; Kinderlehrer and Nirenberg 1977](#)] and the key sharp estimate $|\Omega^t \Delta \Omega^s| \leq C|t - s|$ for the symmetric difference of the positivity sets (or of the contact sets) from [[Blank 2001](#)].

Scalar parametrization of deformations (definition of η^t). By [Proposition 2.1](#) the free boundaries Γ^t are “uniformly” $C^{k,\alpha}$ for $|t|$ small and the difference between Γ^t and Γ^s is bounded by $C|t' - t|$ in the L^∞ norm. A goal of the paper is to prove that the difference is bounded $C|t' - t|$ also in a $C^{k-1,\alpha}$ norm. To prove this type of result it is convenient to have a scalar function representing the “difference” between Γ^t and Γ^s . This has a clear meaning locally —since both Γ^t and Γ^s are graphs, and one can simply subtract the two functions that define these graphs. We next give a global analogue of this.

In an open neighborhood U_o of Γ^0 we define coordinates

$$(z, s) : U_o \rightarrow \mathcal{Z} \times (-s_o, s_o),$$

where $s_o > 0$ and \mathcal{Z} is some smooth approximation of Γ^0 .

We assume the vector field

$$N := \partial_s$$

is a smooth approximation of ν^0 on Γ^0 . More precisely, we assume

$$N \in C^\infty(U_o; \mathbb{R}^n), \quad |N| = 1 \quad \text{and} \quad N \cdot \nu^t \geq (1 - \varepsilon_o) \quad \text{for } t \in (-t_o, t_o), \quad (2-2)$$

where ε_o is a constant that in the sequel will be chosen to be small enough — depending only on \mathcal{C} .

In this framework, [Proposition 2.1](#) implies that for all $t \in (-t_o, t_o)$ with t_o small enough there exists $\eta^t \in C^{k,\alpha}(\mathcal{Z})$ such that

$$\Gamma^t = \{s = \eta^t(z)\} \subset U_o. \quad (2-3)$$

Remark 2.2. From the data of Γ^0 we may always construct \mathcal{Z} and (z, s) satisfying the previous properties — for ε_o arbitrarily small — by taking \mathcal{Z} to be a smooth approximation of Γ^0 and N a smooth approximation of ν^0 . Once \mathcal{Z} and N are chosen, the coordinates (z, s) are then defined respectively as the projection on \mathcal{Z} and the signed distance to \mathcal{Z} along integral curves of N .

3. A priori estimates

Roughly speaking, the goal of this section is to show that if an expansion of the type

$$\eta^t = \eta^0 + \dot{\eta}^0 t + \frac{1}{2} \ddot{\eta}^0 t^2 + \dots$$

holds, where

$$\frac{\eta^t - \eta^0}{t} \rightarrow \dot{\eta}^0 \quad \text{and} \quad \frac{\eta^t - \eta^0 - \dot{\eta}^0 t}{t^2} \rightarrow \frac{\ddot{\eta}^0}{2} \quad \text{as } t \rightarrow 0, \quad \text{in } C^0(\mathcal{Z}),$$

then $\dot{\eta}^0$ and $\ddot{\eta}^0$ must satisfy certain equations that have uniqueness of solution and a priori estimates. From these equations we obtain conditional (or a priori) estimates for $\|\dot{\eta}^0\|_{C^{k-1,\alpha}(\mathcal{Z})}$ and $\|\ddot{\eta}^0\|_{C^{k-2,\alpha}(\mathcal{Z})}$.

In the next sections, let us provisionally assume

$$\Delta(h^t - h^0) \geq 0 \quad \text{and} \quad c^t - c^0 \leq 0 \quad (3-1)$$

for all $t \geq 0$, which is not essential but simplifies the analysis: Assumption (3-1) guarantees that $\Omega^t \subset \Omega^0$ for all $t \geq 0$. Indeed, this is an immediate consequence of the characterization of

$$\tilde{u}^t := u^t - h^t$$

as the infimum of all nonnegative supersolutions with the same right-hand side and appropriate condition at infinity. More precisely, we have the following lemma, whose proof is standard in dimension $n \geq 3$ and which we sketch in dimension $n = 2$ in the [Appendix](#).

Lemma 3.1. *The function \tilde{u}^t can be defined as the infimum of all f satisfying $f \geq 0$, $\Delta f \leq -\Delta h^t$,*

$$\lim_{x \rightarrow \infty} (f + h^t) \geq c^t \quad (n \geq 3), \tag{3-2a}$$

$$\lim_{x \rightarrow \infty} \frac{f + h^t}{-\log |x|} \geq c^t \quad (n = 2). \tag{3-2b}$$

Note that in particular $f = \tilde{u}^0$ is included since $\Delta \tilde{u}^0 = -\Delta h^0 \leq -\Delta h^t$, and

$$\lim_{x \rightarrow \infty} (\tilde{u}^0 + h^t) = \lim_{x \rightarrow \infty} (\tilde{u}^0 + h^0) + \lim_{x \rightarrow \infty} (h^t - h^0) \geq c^0 \geq c^t \quad (n \geq 3),$$

$$\lim_{x \rightarrow \infty} \frac{(\tilde{u}^0 + h^t)}{-\log |x|} \geq c(t) \quad (n = 2).$$

Therefore, applying Lemma 3.1 we obtain $\tilde{u}^0 \geq \tilde{u}^t$ and

$$\Omega^t = \{\tilde{u}^t > 0\} \subset \{\tilde{u}^0 > 0\} = \Omega^0$$

for all $t > 0$. Equivalently (3-1) implies $\eta^t \geq 0$ on \mathcal{Z} for $t > 0$.

Later, when we prove Theorem 1.1, we will reduce to this case by decomposing h^t as a sum of two functions, one with nonnegative Laplacian and one with nonpositive Laplacian.

Let us define

$$v^t := \delta_t \tilde{u}^0 = \frac{1}{t}(\tilde{u}^t - \tilde{u}^0). \tag{3-3}$$

The function v^t is a solution of

$$\begin{cases} \Delta v^t = -\Delta \delta_t h^0 & \text{in } \Omega^t, \\ v^t = -\frac{1}{t} \tilde{u}^0 & \text{on } \Gamma^t, \\ \lim_{x \rightarrow \infty} v^t = \delta_t c^0 \quad (n \geq 3), \end{cases} \quad \begin{cases} \Delta v^t = -\Delta \delta_t h^0 & \text{in } \Omega^t, \\ v^t = -\frac{1}{t} \tilde{u}^0 & \text{on } \Gamma^t, \\ \lim_{x \rightarrow \infty} \frac{v^t(x)}{-\log |x|} = \delta_t c^0 \quad (n = 2). \end{cases} \tag{3-4}$$

Since $\tilde{u}^0 = |\nabla \tilde{u}^0| = 0$ on Γ^0 , using the classical estimate⁵

$$\|u^0\|_{C^{1,1}(\mathbb{R}^n)} \leq (n - 1) \|h^0\|_{C^{1,1}(\mathbb{R}^n)},$$

we obtain

$$|\tilde{u}^0| \leq C \|h\|_{C^{1,1}(\mathbb{R}^n)} d_{\text{Hausdorff}}^2(\Gamma^t, \Gamma^0) \leq Ct^2 \quad \text{on } \Gamma^t.$$

Then, using that Ω^t grows to Ω^0 as $t \downarrow 0$ and uniform estimates for v^t we find that $v^t \rightarrow v$ as $t \downarrow 0$, where v is the solution of

$$\begin{cases} \Delta v = -\Delta \dot{h}^0 & \text{in } \Omega^0, \\ v = 0 & \text{on } \Gamma^0, \\ v(\infty) = \dot{c}^0 \quad (n \geq 3), \end{cases} \quad \begin{cases} \Delta v = -\Delta \dot{h}^0 & \text{in } \Omega^0, \\ v = 0 & \text{on } \Gamma^0, \\ \lim_{x \rightarrow \infty} \frac{v(x)}{-\log |x|} = \dot{c}^0 \quad (n = 2). \end{cases} \tag{3-5}$$

Here $\Delta \dot{h}^0 = \lim_{t \downarrow 0} \Delta \delta_t h^0 = (\Delta \partial_t h)(0, x)$.

⁵Since u^0 is a solution of the obstacle problem in the whole \mathbb{R}^n with a semiconcave obstacle h^0 , we know u^0 is semiconcave with $D^2 u^0 \geq -\|h\|_{C^{1,1}(\mathbb{R}^n)} \text{Id}$ and the estimate follows using $\Delta u^0 = 0$, where $u^0 > h^0$.

Equation and estimate for $\dot{\eta}^0$. We first prove the following

Proposition 3.2. *Let $k \geq 1$. Assume that for some $t_m \downarrow 0$ there exists $\dot{\eta}^0 \in C^0(\mathcal{Z})$ such that*

$$\delta_{t_m} \eta^0 \rightarrow \dot{\eta}^0 \quad \text{in } C^0(\mathcal{Z}) \quad \text{as } m \rightarrow \infty.$$

Then, the limit $\dot{\eta}^0$ is given by

$$\dot{\eta}^0(z) = \left(\frac{\partial_N v}{(N \cdot \nu^0)^2 \Delta h^0} \right)(z, \eta^0(z)), \tag{3-6}$$

with v as in (3-5). As a consequence, $\dot{\eta}^0$ is independent of the sequence t_m and we have $\dot{\eta} \in C^{k-1,\alpha}(\mathcal{Z})$ with the estimate

$$\|\dot{\eta}^0\|_{C^{k-1,\alpha}(\mathcal{Z})} \leq C(\mathcal{C}^0)(\|\dot{h}^0\|_{C^{k,\alpha}(B_R)} + |c^0|). \tag{3-7}$$

Proof. We split the proof into two steps.

Step 1. We prove (3-6). Recall that since \tilde{u}^t is a solution of a zero obstacle problem we have

$$\tilde{u}^t = |\nabla \tilde{u}^t| = 0 \quad \text{on } \Gamma^t.$$

Thus,

$$\partial_s v^t = \frac{1}{t}(\partial_s \tilde{u}^t - \partial_s \tilde{u}^0) = -\frac{\partial_s \tilde{u}^0}{t} \quad \text{on } \Gamma^t. \tag{3-8}$$

From (3-8) we deduce that

$$\partial_s v^{t_m}(z, \eta^{t_m}) = -\frac{1}{t_m} \partial_s \tilde{u}^0(z, \eta^{t_m}) = -\frac{1}{t_m} (\partial_s \tilde{u}^0(z, \eta^0) + \partial_{ss} \tilde{u}^0(z, \eta^0)(\eta^{t_m} - \eta^0) + o(t_m)), \tag{3-9}$$

where η^0 and η^{t_m} are evaluated at z (although we omit this in the notation) and where $\partial_{ss} \tilde{u}^0(z, \eta^0)$ is understood as the limit from the Ω^0 -side. To justify the validity of the previous Taylor expansion we use that $\tilde{u}^0 \in C^{2,\alpha}(\bar{\Omega}^0)$; see Lemma 3.6.

Since $\tilde{u}^0 = |\tilde{\nabla} u^0| = 0$ on Γ^0 we obtain

$$\partial_{ee} \tilde{u}^0 = (e \cdot \nu)^2 \partial_{\nu\nu} \tilde{u}^0 = (e \cdot \nu)^2 \Delta \tilde{u}^0 = -(e \cdot \nu)^2 \Delta h^0 \quad \text{on } \Gamma^0$$

for every vector e , where $\nu = \nu^0$ is the normal vector to Γ^0 (pointing towards Ω^0). Again, the previous second derivatives on Γ^0 mean the limits from the Ω^0 -side. Hence, we have

$$\partial_s \tilde{u}^0(z, \eta^0(z)) = 0 \quad \text{and} \quad \partial_{ss} \tilde{u}^0(z, \eta^0(z)) = -((N \cdot \nu^0)^2 \Delta h^0)(z, \eta^0(z)), \tag{3-10}$$

where $\partial_{ss} \tilde{u}^0(z, \eta^0(z))$ is from the Ω^0 -side. Dividing (3-10) by t_m and taking the limit as $t_m \downarrow 0$ in (3-9) using the assumption, we obtain

$$\partial_s v(z, \eta^0(z)) = -\partial_{ss} \tilde{u}^0(z, \eta^0(z)) \dot{\eta}^0(z) = ((N \cdot \nu^0)^2 \Delta h^0)(z, \eta^0(z)) \dot{\eta}^0(z), \tag{3-11}$$

where $\partial_s v(z, \eta^0(z))$ and $\partial_{ss} \tilde{u}^0(z, \eta^0(z))$ are from the Ω^0 -side. When computing the limit that yields (3-11) we must check that

$$\partial_s v^{t_m}(z, \eta^{t_m}(z)) \rightarrow \partial_s v(z, \eta^0(z)), \tag{3-12}$$

where $\partial_s v(z, \eta^0)$ is from the Ω^0 -side. To prove this, note that the equation (3-4) for v^t , since we have uniform $C^{1,\alpha}$ estimates for the boundary Γ^t , implies that $\|\nabla v^t\|_{C^{0,\alpha}(\bar{\Omega}^t)}$ is uniformly bounded (for $t > 0$ small). This implies that ∇v^t converges uniformly to ∇v in every compact set of Ω^0 . Then using the uniform continuity of the derivatives of v on $\bar{\Omega}^0$ we show that

$$\lim \nabla v^{t_p}(x_p) \rightarrow \nabla v(x) \quad \text{as } p \rightarrow \infty \quad \text{whenever } t_p \downarrow 0, \quad x_p \rightarrow x \text{ and } x_p \in \bar{\Omega}^{t_p}.$$

This establishes (3-12) and (3-11). Then, (3-6) follows immediately from (3-11), after recalling that $N = \partial_s$.

Step 2. We prove (3-7). Indeed, from (3-5), and using that $\Gamma^0 = \partial\Omega^0 \in C_{\rho/4}^{k,\alpha}$ with norm universally bounded, we obtain

$$\|v\|_{C^{k,\alpha}(\Omega^0)} \leq C(\mathcal{C}^0)(\|\Delta \dot{h}^0\|_{C^{k-2,\alpha}(\Omega^0)} + |\dot{c}^0|) \leq C(\mathcal{C}^0)(\|\dot{h}^0\|_{C^{k,\alpha}(\Omega^0)} + |\dot{c}^0|). \tag{3-13}$$

Now recalling that N is smooth, that $\|v^0\|_{C^{k-1,\alpha}(\Gamma^0)} \leq C\|\Gamma^0\|_{C_{\rho/4}^{k,\alpha}} \leq C$, that $-\Delta h^0 \geq \rho$, and that $\|\eta^0\|_{C^{k,\alpha}(\mathcal{Z})} \leq C$, (3-6) and (3-13) imply (3-7). □

Equation and estimate for $\ddot{\eta}^0$. In this section we estimate the second derivative in t of η at $t = 0$. It is convenient to introduce here the following notation, which we shall use throughout the paper. Given a function $f : (-t_o, t_o) \times Y \rightarrow \mathbb{R}$, recall the notation $f = f^t(y) = f(t, x)$. Let us also define

$$\delta_t^2 f^s := 2 \frac{\delta_t f^s - f^s}{t} \quad \text{and} \quad \ddot{f}^s := \lim_{t \downarrow 0} \delta_t^2 f^s = \partial_{tt} f(y, 0).$$

From now on let us consider v to be defined in all of \mathbb{R}^n by extending the solution of (3-5) by 0 in $\mathbb{R}^n \setminus \Omega^0$. Note that this is consistent with $v = \lim_{t \downarrow 0} v^t$ and $v^t = \delta_t \tilde{u}^0 = 0$ in $\mathbb{R}^n \setminus \Omega^0$ (since both \tilde{u}^t and \tilde{u}^0 vanish there).

We now introduce the function, defined in all of \mathbb{R}^n ,

$$w^t := \delta_t v^0 = \frac{1}{t}(v^t - v) = \frac{1}{2} \delta_t^2 \tilde{u}^0.$$

Using (3-5) and the identity

$$\Delta v^t = \frac{1}{t} \Delta(\tilde{u}^t - \tilde{u}^0) = -\frac{1}{t} \Delta \tilde{u}^0 = \frac{1}{t} \Delta h^0 \quad \text{in } \Omega^0 \setminus \Omega^t,$$

we find, in the distributional sense,

$$\begin{cases} \Delta w^t = \frac{1}{t}((\partial_N v / (N \cdot v^0)) \mathcal{H}^{n-1} \upharpoonright_{\Gamma^0} + (\frac{1}{t} \Delta h^0 - \Delta \dot{h}^0) \chi_{\Omega^0 \setminus \Omega^t}) - \frac{1}{2} \Delta \delta_t^2 h^0 \chi_{\Omega^t} & \text{in } \mathbb{R}^n, \\ w^t(\infty) = \frac{1}{2} \delta_t^2 c^0 \quad (n \geq 3), \end{cases} \tag{3-14a}$$

$$\begin{cases} \Delta w^t = \frac{1}{t}((\partial_N v / (N \cdot v^0)) \mathcal{H}^{n-1} \upharpoonright_{\Gamma^0} + (\frac{1}{t} \Delta h^0 - \Delta \dot{h}^0) \chi_{\Omega^0 \setminus \Omega^t}) - \frac{1}{2} \Delta \delta_t^2 h^0 \chi_{\Omega^t} & \text{in } \mathbb{R}^n, \\ \lim_{x \rightarrow \infty} \frac{w^t}{-\log|x|} = \frac{1}{2} \delta_t^2 c^0 \quad (n = 2), \end{cases} \tag{3-14b}$$

where \mathcal{H} denotes the Hausdorff measure. Indeed, note also that for $v = v^0$ we have

$$\partial_N v = (N \cdot v^0) \partial_\nu v \quad \text{on } \Gamma_{\text{out}}^0, \quad \text{while} \quad \partial_\nu v = 0 \quad \text{on } \Gamma_{\text{in}}^0.$$

Here, “ Γ_{out}^0 ” refers to the limit from the Ω^0 -side, while “ Γ_{in}^0 ” refers to the limit from the $(\mathbb{R}^n \setminus \bar{\Omega}^0)$ -side. Therefore, Δw^t has some mass concentrated on Γ^0 which is given by the jump in the normal derivative of v , namely,

$$\frac{1}{t} \frac{\partial_N v}{N \cdot \nu^0} \mathcal{H}^{n-1} \upharpoonright_{\Gamma^0}.$$

In the following lemma, and throughout the paper, P denotes the Newtonian potential in dimension n , namely,

$$P(x) = \frac{1}{n(n-2)|B_1|} |x|^{2-n} \quad (n \geq 3)$$

or

$$P(x) = -\frac{1}{2\pi} \log |x| \quad (n = 2).$$

Recall that $-\Delta P = \delta_{x=0}$ in the sense of distributions.

We also need to introduce the Jacobian

$$J(z, s) := |\det D(z, s)^{-1}|$$

of the coordinates (z, s) defined by

$$\int_A f(x) dx = \int_{(z,s)(A)} f(z, s) J(z, s) dz ds.$$

We use the following abuse of notation:

- When $f = f(x)$ we denote by $f(z, s)$ the composition $f \circ (z, s)^{-1}$.
- Conversely, when $g = g(z, s)$ we will denote by $g(x)$ the composition $g \circ (z, s)$.

Finally, let us denote by

$$\pi_1 : U_\circ \rightarrow \mathcal{Z}$$

the projection map along N , which is defined in the coordinates (z, s) by

$$(z, s) \mapsto (z, 0).$$

We will need the following:

Lemma 3.3. *Given $f : \Gamma^0 \rightarrow \mathbb{R}$ continuous we have*

$$\int_{\Gamma^0} (N \cdot \nu^0)(x) f(x) d\mathcal{H}^{n-1}(x) = \int_{\mathcal{Z}} f(z, \eta^0(z)) J(z, \eta^0(z)) dz.$$

Proof. Let us assume without loss of generality that f is defined and continuous in the neighborhood U_\circ of Γ^0 . Given $\varepsilon > 0$ let

$$A^\varepsilon := \{x \in U_\circ : \eta^0(z(x)) \leq s(x) \leq \eta^0(z(x)) + \varepsilon\}.$$

Recalling that $N = \partial_s$ and that $|N| = 1$, we have

$$\int_{\Gamma^0} (N \cdot \nu^0)(x) f(x) d\mathcal{H}^{n-1}(x) = \lim_{\varepsilon \downarrow 0} \frac{1}{\varepsilon} \int_{A^\varepsilon} f(x) d\mathcal{H}^n(x).$$

On the other hand, for $(z, s)(A^\varepsilon) := \{(z, s) \in \mathcal{Z} \times (-s_o, s_o) : \eta^0(z) \leq s \leq \eta^0(z) + \varepsilon\}$ we have, by the definition of J ,

$$\begin{aligned} \frac{1}{\varepsilon} \int_{A^\varepsilon} f(x) d\mathcal{H}^n(x) &= \frac{1}{\varepsilon} \int_{(z,s)(A^\varepsilon)} f(z, s) J(z, s) dz ds \\ &= \int_{\mathcal{Z}} dz \frac{1}{\varepsilon} \int_0^\varepsilon d\bar{s} f(z, \eta^0(z) + \bar{s}) J(z, \eta^0(z) + \bar{s}) \\ &= \int_{\mathcal{Z}} f(z, \eta^0(z)) J(z, \eta^0(z)) dz + o(1) \end{aligned}$$

as $\varepsilon \downarrow 0$ and the lemma follows. □

Lemma 3.4. *Let $k \geq 2$. Assume that for some $t_m \downarrow 0$ there exist $\dot{\eta}, \ddot{\eta} \in C^0(\mathcal{Z})$ such that*

$$\delta_{t_m}^2 \eta^0 = 2 \frac{\eta^{t_m} - \eta^0 - \dot{\eta}^0 t_m}{t_m^2} \rightarrow \ddot{\eta}^0 \quad \text{in } C^0(\mathcal{Z})$$

as $t_m \downarrow 0$. Then,

$$w^{t_m} \xrightarrow{\text{weakly}} w \quad \text{in } \mathbb{R}^n,$$

where w can be decomposed as

$$w = w_{\text{solid}} + w_{\text{single}} + w_{\text{double}} + w_{\text{implicit}} + \text{constant} \tag{3-15}$$

for

$$w_{\text{solid}}(x) := \int_{\mathbb{R}^n} d\mathcal{H}^n(y) (\Delta \dot{h}^0 \chi_{\Omega_0})(y) P(x-y), \tag{3-16}$$

$$w_{\text{single}}(x) := \int_{\Gamma_0} d\mathcal{H}^{n-1}(y) \left((N \cdot \nu^0)(\dot{\eta}^0 \circ \pi_1) \Delta \dot{h}^0 - \frac{1}{2} (\dot{\eta}^0 \circ \pi_1)^2 \frac{N \cdot \nu^0}{J} \partial_N (Jh^0) \right)(y) P(x-y), \tag{3-17}$$

$$w_{\text{double}}(x) := \int_{\Gamma_0} d\mathcal{H}^{n-1}(y) \left(\frac{1}{2} (\dot{\eta}^0 \circ \pi_1)^2 \frac{N \cdot \nu^0}{J} (J \Delta h^0) \right)(y) \partial_N P(x-y), \tag{3-18}$$

$$w_{\text{implicit}}(x) := \int_{\Gamma_0} d\mathcal{H}^{n-1}(y) \frac{\Theta}{(N \cdot \nu)}(y) P(x-y), \tag{3-19}$$

where $\Theta : \Gamma^0 \rightarrow \mathbb{R}$,

$$\Theta := \frac{1}{2} (N \cdot \nu^0)^2 \Delta h^0 (\ddot{\eta}^0 \circ \pi_1). \tag{3-20}$$

Proof. Define

$$\mathcal{D}^t := \Delta w^t = \frac{1}{t} \frac{\partial_N \nu}{N \cdot \nu^0} \mathcal{H}^{n-1} \upharpoonright_{\Gamma^0} - \left(\frac{1}{t^2} \Delta h^0 + \frac{1}{t} \Delta \dot{h}^0 \right) \chi_{\Omega^0 \setminus \Omega^t} - \frac{1}{2} \Delta \delta_t^2 h^0 \chi_{\Omega^t}.$$

Let us show that $\mathcal{D}^{t_m} \rightarrow \mathcal{D}$ in the sense of distributions, for some distribution \mathcal{D} that we compute.

Let us first write

$$\mathcal{D}^t = \mathcal{D}'_1 + \mathcal{D}'_2,$$

where

$$\mathcal{D}'_1 := -\frac{1}{t} \Delta \dot{h}^0 \chi_{\Omega^0 \setminus \Omega^t} - \frac{1}{2} \Delta \delta_t^2 h^0 \chi_{\Omega^t} \tag{3-21}$$

and

$$\mathcal{D}_2^t := \frac{1}{t} \left(\frac{\partial_N v}{N \cdot \nu^0} \mathcal{H}^{n-1} \upharpoonright_{\Gamma^0} - \frac{1}{t} \Delta h^0 \chi_{\Omega^0 \setminus \Omega^t} \right). \tag{3-22}$$

First we clearly have, for $\phi \in C_c^\infty(\mathbb{R}^n)$,

$$\begin{aligned} \int \phi(x) \left(\frac{1}{t} \Delta \dot{h}^0 \chi_{\Omega^0 \setminus \Omega^t} \right)(x) dx &= \frac{1}{t} \int_{\mathcal{Z}} \int_{\eta^0}^{\eta^t} J(z, s) (\Delta \dot{h}^0 \phi)(z, s) dz ds \\ &\rightarrow \int_{\mathcal{Z}} \dot{\eta}^0(z) J(z, \eta^0) (\Delta \dot{h}^0 \phi)(z, \eta^0) dz = \int_{\Gamma^0} (N \cdot \nu^0) (\dot{\eta}^0 \circ \pi_1) \Delta \dot{h}^0 \phi d\mathcal{H}^{n-1} \end{aligned}$$

as $t = t_m \downarrow 0$, where we have used [Lemma 3.3](#), and hence

$$\mathcal{D}_1^{t_m} \xrightarrow{\text{weakly}} -(N \cdot \nu^0) (\dot{\eta}^0 \circ \pi_1) \Delta \dot{h}^0 \mathcal{H}^{n-1} \upharpoonright_{\Gamma^0} - \frac{1}{2} \Delta \dot{h}^0 \chi_{\Omega^0}. \tag{3-23}$$

Next, using [\(3-6\)](#), we compute, for $\phi \in C_c^\infty(\mathbb{R}^n)$,

$$\begin{aligned} \int \phi \mathcal{D}_2^t &= \frac{1}{t} \left(\int_{\mathcal{Z}} dz \frac{J(z, \eta^0)}{(N \cdot \nu^0)^2(z, \eta^0)} \partial_N v(z) \phi(z, \eta^0) - \int_{\mathcal{Z}} dz \frac{1}{t} \int_{\eta^0}^{\eta^t} ds (J \Delta h^0 \phi)(z, s) \right) \\ &= \frac{1}{t} \int_{\mathcal{Z}} dz \left((J \Delta h^0 \phi)(z, \eta^0) \dot{\eta}^0 - \frac{1}{t} \int_{\eta^0}^{\eta^t} ds (J \Delta h^0 \phi)(z, s) \right) \\ &= I_1 + I_2, \end{aligned} \tag{3-24}$$

where

$$I_1 := \frac{1}{t} \int_{\mathcal{Z}} dz \left((J \Delta h^0 \phi)(z, \eta^0) \dot{\eta}^0 - \frac{1}{t} \int_{\eta^0}^{\eta^0 + \dot{\eta}^0 t} ds (J \Delta h^0 \phi)(z, s) \right)$$

and

$$I_2 := -\frac{1}{t^2} \int_{\mathcal{Z}} dz \int_{\eta^0 + \dot{\eta}^0 t}^{\eta^t} ds (J \Delta h^0 \phi)(z, s).$$

On one hand, letting $s = \eta^0 + \dot{\eta}^0 t \bar{s}$,

$$\begin{aligned} I_1 &= \int_{\mathcal{Z}} dz \int_0^1 \dot{\eta}^0(z) t d\bar{s} \frac{\bar{s}}{t} \left(\frac{(J \Delta h^0 \phi)(z, \eta^0) - (J \Delta h^0 \phi)(z, \eta^0 + \dot{\eta}^0 t \bar{s})}{\bar{s} t} \right) \\ &= \int_{\mathcal{Z}} dz \int_0^1 (\dot{\eta}^0)^2(z) \bar{s} d\bar{s} \partial_s (J \Delta h^0 \phi)(z, \eta^0) + o(1) \\ &= \int_0^1 \bar{s} d\bar{s} \int_{\mathcal{Z}} dz (\dot{\eta}^0)^2(z) \partial_s (J \Delta h^0 \phi)(z, \eta^0) + o(1) \\ &= \frac{1}{2} \int_{\Gamma^0} (N \cdot \nu^0) d\mathcal{H}^{n-1} \frac{1}{J} (\dot{\eta}^0 \circ \pi_1)^2 \partial_N (J \Delta h^0 \phi) + o(1). \end{aligned} \tag{3-25}$$

as $t = t_m \downarrow 0$, where for the last relation we used [Lemma 3.3](#) with

$$f(x) = \left(\frac{1}{J} (\dot{\eta}^0 \circ \pi_1)^2 \partial_N (J \Delta h^0 \phi) \right)(x),$$

noting also that $\partial_s = \partial_N$ and $(\dot{\eta}^0 \circ \pi_1)^2(z, \eta^0(z)) = (\dot{\eta}^0)^2(z)$. On the other hand,

$$\begin{aligned} I_2 &= -\frac{1}{t^2} \int_{\mathcal{Z}} dz \int_{\eta^0 + \dot{\eta}^0 t}^{\eta^0 + \dot{\eta}^0 t + \frac{1}{2} \ddot{\eta}^0 t^2} ds (J \Delta h^0 \phi)(z, s) + o(1) \\ &= -\frac{1}{2} \int_{\mathcal{Z}} dz \ddot{\eta}^0 (J \Delta h^0 \phi)(z, \eta^0) + o(1) \\ &= -\frac{1}{2} \int_{\Gamma^0} d\mathcal{H}^{n-1} (N \cdot \nu^0) (\ddot{\eta}^0 \circ \pi_1) \Delta h^0 \phi + o(1) \end{aligned} \tag{3-26}$$

as $t = t_m \downarrow 0$. Therefore, $\mathcal{D}_2^{t_m} \rightarrow \mathcal{D}_2$, where

$$\int \phi \mathcal{D}_2 = \frac{1}{2} \int_{\Gamma^0} d\mathcal{H}^{n-1} (\dot{\eta}^0 \circ \pi_1)^2 \frac{N \cdot \nu^0}{J} \partial_N (J \Delta h^0 \phi) - \frac{1}{2} \int_{\Gamma^0} d\mathcal{H}^{n-1} (N \cdot \nu^0) (\ddot{\eta}^0 \circ \pi_1) \Delta h^0 \phi. \tag{3-27}$$

In dimension $n \geq 3$ we have

$$w^{t_m}(\infty) = \frac{1}{2} \lim_{x \rightarrow \infty} \delta_{t_m}^2 \tilde{u}^0(\infty) = \frac{1}{2} \delta_{t_m}^2 c^0 \rightarrow \frac{1}{2} \ddot{c}^0,$$

and thus

$$w(\infty) = \frac{1}{2} \ddot{c}^0 = \text{constant}.$$

In dimension $n = 2$ we have instead

$$\lim_{x \rightarrow \infty} \frac{w(x)}{-\log|x|} = \frac{1}{2} \ddot{c}^0$$

and this implies $2\pi \frac{1}{2} \ddot{c}^0 = \int_{\mathbb{R}^2} \Delta w$ and that w can be obtained (up to an additive constant) by convolving the Newtonian potential P with Δw .

Therefore, combining (3-23) and (3-27), we obtain that (3-15)–(3-19) hold. □

We may now state the final result of this section.

Proposition 3.5. *Let $k \geq 2$. Assume that for some $t_m \downarrow 0$ there exist $\dot{\eta}, \ddot{\eta} \in C^0(\mathcal{Z})$ such that*

$$\delta_{t_m}^2 \eta^0 = 2 \frac{\eta^{t_m} - \eta^0 - \dot{\eta}^0 t_m}{t_m^2} \rightarrow \ddot{\eta}^0 \quad \text{in } C^0(\mathcal{Z})$$

as $t_m \downarrow 0$. Assume $w \in C^1(\bar{\Omega}^0)$ and

$$\lim \nabla w^{t_m}(x_m) \rightarrow \nabla w(x) \quad \text{as } m \rightarrow \infty \quad \text{for all } x_m \rightarrow x \text{ such that } x_m \in \bar{\Omega}^{t_m}. \tag{3-28}$$

Then, $\Theta : \Gamma^0 \rightarrow \mathbb{R}$ defined by (3-20) satisfies

$$\Theta - \frac{1}{2} \partial_{sss} u^0 (\dot{\eta}^0 \circ \pi_1)^2 = \partial_{ss} v \dot{\eta}^0 + \partial_s w \quad \text{on } \Gamma_{\text{out}}^0. \tag{3-29}$$

Moreover, $\ddot{\eta}^0$ does not depend on (t_m) and

$$\|\ddot{\eta}^0\|_{C^{k-2,\alpha}(\mathcal{Z})} \leq C(\mathcal{C}^0) \mathcal{Q}, \tag{3-30}$$

where

$$\mathcal{Q} := \|\dot{h}^0\|_{C^{k-1,\alpha}(\mathbb{R}^n)} + |\dot{c}^0| + (\|\dot{h}^0\|_{C^{k,\alpha}(\mathbb{R}^n)} + |\dot{c}^0|)(\|\dot{h}^0\|_{L^\infty(\mathbb{R}^n)} + |\dot{c}^0|).$$

As for η , the independence of t_m and regularity of η will be consequences of the fact that Θ solves (3-29), for which regularity estimates and uniqueness hold. However, note that (3-29) is an implicit equation for Θ since w_{implicit} depends on Θ , which makes the analysis more involved.

To prove Proposition 3.5, we will need two auxiliary lemmas with standard proofs.

Lemma 3.6. *We have*

$$\|\tilde{u}^0\|_{C^{k+1,\alpha}(\overline{B_R \cap \Omega^t})} \leq C(\mathcal{C}^0).$$

More generally, for $t \in [0, t_o]$, where $t_o = t_o(\mathcal{C})$, we have $\tilde{u}^t \in C^{k+1,\alpha}(\overline{\Omega^t})$ with

$$\|\tilde{u}^t\|_{C^{k+1,\alpha}(\overline{B_R \cap \Omega^t})} \leq C(\mathcal{C}).$$

Proof. Note that $\partial_i \tilde{u}^t$ solves

$$\Delta(\partial_i \tilde{u}^t) = -\Delta(\partial_i h^t) \quad \text{in } \Omega^t, \quad \text{with } \partial_i \tilde{u}^t = 0 \quad \text{on } \Gamma^t = \partial \Omega^t.$$

Since $-\Delta(\partial_i h^t) \in C^{k-2,\alpha}(\mathbb{R}^n)$ and Γ^t belongs to $C_\rho^{k,\alpha}$, using standard Schauder estimates up to the boundary we obtain

$$\partial_i \tilde{u}^t \in C^{k,\alpha}(\overline{B_R \cap \Omega^t}),$$

and hence

$$\tilde{u}^t \in C^{k+1,\alpha}(\overline{B_R \cap \Omega^t}). \quad \square$$

Lemma 3.7. *Let $U \subset \overline{B_R} \subset \mathbb{R}^n$ be bounded with ∂U belonging to $C_r^{m+2,\alpha}$ for some $r > 0$ and $f \in C_c^{m,\alpha}(B_{2R})$, where $m \geq 0$. Let W be the solution of*

$$\begin{cases} \Delta W = f \chi_{\mathbb{R}^n \setminus U} & \text{in } \mathbb{R}^n, \\ W(\infty) = 0 & (n \geq 3), \end{cases} \quad \begin{cases} \Delta W = f \chi_{\mathbb{R}^n \setminus U} & \text{in } \mathbb{R}^n, \\ \lim_{x \rightarrow \infty} \frac{W(x)}{-\log|x|} = 2\pi \int_{\mathbb{R}^2} f \chi_{\mathbb{R}^n \setminus U} & (n = 2), \end{cases}$$

which is given in dimension 2 by convolution with the logarithmic Newtonian potential.

Then,

$$\|W\|_{C^{m+2,\alpha}(\overline{B_{2R} \setminus U})} + \|W\|_{C^{m+2,\alpha}(\overline{U})} \leq C\|f\|_{C^{m,\alpha}(\overline{B_{2R}})},$$

where $C = C(n, m, \alpha, R, r, \|\partial U\|_{C_r^{m+2,\alpha}})$.

Proof. Let \tilde{W} be the solution of

$$\begin{cases} \Delta \tilde{W} = f & \text{in } \mathbb{R}^n \setminus U, \\ \tilde{W} = 0 & \text{on } \partial U, \\ \tilde{W}(\infty) = 0 & (n \geq 3), \end{cases} \quad \begin{cases} \Delta \tilde{W} = f & \text{in } \mathbb{R}^n \setminus U, \\ \tilde{W} = 0 & \text{on } \partial U, \\ \lim_{x \rightarrow \infty} \frac{\tilde{W}(x)}{-\log|x|} = 0 & (n = 2). \end{cases}$$

We consider \tilde{W} defined in all of \mathbb{R}^n by extending it by 0 in U .

Note that by standard Schauder estimates up to the boundary we have

$$\|\tilde{W}\|_{C^{m+2,\alpha}(\overline{B_{2R} \setminus U})} \leq C\|f\|_{C^{m,\alpha}(\overline{B_{2R}})}. \tag{3-31}$$

On the other hand, the difference $(\tilde{W} - W)$ solves, in all of \mathbb{R}^n ,

$$\begin{cases} \Delta(\tilde{W} - W) = \partial_{v, \text{out}} \tilde{W} H^{n-1} \upharpoonright_{\partial U} & \text{in } \mathbb{R}^n, \\ (\tilde{W} - W)(\infty) = 0 & (n \geq 3), \end{cases}$$

$$\begin{cases} \Delta(\tilde{W} - W) = \partial_{v, \text{out}} \tilde{W} H^{n-1} \upharpoonright_{\partial U} & \text{in } \mathbb{R}^n, \\ \lim_{x \rightarrow \infty} \frac{(\tilde{W} - W)(x)}{-\log|x|} = -2\pi \int_{\mathbb{R}^2} f \chi_{\mathbb{R}^n \setminus U} = 2\pi \int_{\partial U} \partial_{v, \text{out}} \tilde{W} & \text{at } \infty \quad (n = 2). \end{cases}$$

Therefore, $\tilde{W} - W$ is a single-layer potential and using [Theorem A.1](#) we obtain

$$\begin{aligned} \|(\tilde{W} - W)\|_{C^{m+2,\alpha}(\bar{B}_{2R} \setminus U)} + \|(\tilde{W} - W)\|_{C^{m+2,\alpha}(\bar{U})} &\leq C \|\partial_{v, \text{out}} \tilde{W}\|_{C^{m+1,\alpha}(\partial U)} \\ &\leq C \|\tilde{W}\|_{C^{m+2,\alpha}(\bar{B}_{2R} \setminus U)} \leq C \|f\|_{C^{m,\alpha}(\bar{B}_4)}. \end{aligned}$$

Using [\(3-31\)](#) and recalling that by definition $\tilde{W} \equiv 0$ in U we obtain

$$\|W\|_{C^{m+2,\alpha}(\bar{B}_{2R} \setminus U)} + \|W\|_{C^{m+2,\alpha}(\bar{U})} \leq C \|f\|_{C^{m,\alpha}(\bar{B}_{2R})}. \quad \square$$

Proof of Proposition 3.5. Step 1. We first prove [\(3-29\)](#).

Expanding [\(3-8\)](#) as in [\(3-9\)](#) but up to the next order, we find

$$\partial_s v^t(z, \eta^t) = -\partial_{ss} \tilde{u}^0(z, \eta^0)(\dot{\eta}^0 + \frac{1}{2} \ddot{\eta}^0 t + o(t)) - \frac{1}{2} \partial_{sss} \tilde{u}^0(z, \eta^0)(\dot{\eta}^0)^2 t + o(t) \tag{3-32}$$

as $t = t_m \downarrow 0$.

Here $\eta, \dot{\eta}$ and $\ddot{\eta}$ are evaluated at z (although we omit this in the notation) and $\partial_{ss} \tilde{u}^0(z, \eta^0)$ and $\partial_{sss} \tilde{u}^0(z, \eta^0)$ mean the limits from Ω^0 . To obtain the Taylor expansion up to the third order of \tilde{u}^0 we are using that, by [Lemma 3.6](#), $u^0 \in C^{k+1,\alpha}(\bar{B}_R \cap \Omega^0)$ where $k \geq 2$. Recall here that $\{u^0 = 0\} = \mathbb{R}^n \setminus \Omega^0 \subset U \subset B_R$.

Subtracting from both sides of [\(3-32\)](#) the quantity

$$\partial_s v(z, \eta^0) = -\partial_{ss} \tilde{u}^0(z, \eta^0) \dot{\eta}^0 \tag{3-33}$$

and dividing by t , we obtain

$$\frac{\partial_s v^t(z, \eta^t) - \partial_s v(z, \eta^0)}{t} = -\frac{1}{2} \partial_{ss} \tilde{u}^0(z, \eta^0) \ddot{\eta}^0 - \frac{1}{2} \partial_{sss} u^0(z, \eta^0) (\dot{\eta}^0)^2 + o(1). \tag{3-34}$$

Recall that by [Lemma 3.4](#) we have $w^t \rightarrow w$ in the sense of distributions with w given by [\(3-15\)](#)–[\(3-19\)](#). Then, the assumption [\(3-28\)](#) allows us to compute the limit of the left-hand side in [\(3-34\)](#), namely,

$$\begin{aligned} \lim_{t=t_m \downarrow 0} \frac{\partial_s v^t(z, \eta^t) - \partial_s v(z, \eta^0)}{t} &= \lim_{t=t_m \downarrow 0} \frac{\partial_s v(z, \eta^t) - \partial_s v(z, \eta^0)}{t} + \frac{\partial_s v^t(z, \eta^t) - \partial_s v(z, \eta^t)}{t} \\ &= \partial_{ss} v(z, \eta^0) \dot{\eta} + \lim_{t=t_m \downarrow 0} (N(z, \eta^t) \cdot \nabla w^t(z, \eta^t)) \\ &= \partial_{ss} v(z, \eta^0) \dot{\eta} + \partial_s w^t(z, \eta^0), \end{aligned} \tag{3-35}$$

where we have used the assumption [\(3-28\)](#).

Taking $t = t_m \downarrow 0$ in (3-34) and using (3-35) we obtain

$$-\frac{1}{2}\partial_{ss}\tilde{u}^0(z, \eta^0)\ddot{\eta}^0 - \frac{1}{2}\partial_{sss}u^0(z, \eta^0)(\dot{\eta}^0)^2 = \partial_{ss}v(z, \eta^0)\dot{\eta} + \partial_s w^t(z, \eta^0).$$

Recalling the definition of Θ in (3-20) and the fact that $\partial_{ss}\tilde{u}^0 = -\Delta h^0$ on Γ^0 —and in particular at (z, η^0) —we obtain (3-29).

Step 2. We use (3-29) to prove uniqueness and regularity of $\ddot{\eta}$. Recall that

$$\partial_s w = \partial_N w = \partial_N w_{\text{solid}} + \partial_N w_{\text{single}} + \partial_N w_{\text{double}} + \partial_N w_{\text{implicit}},$$

and while $\partial_N w_{\text{solid}}$, $\partial_N w_{\text{single}}$, $\partial_N w_{\text{double}}$ depend only on “known” functions—see (3-16), (3-17), (3-18)—the term $\partial_N w_{\text{implicit}}$ introduces a “implicit” dependence on Θ —see (3-19). We therefore need to “solve for Θ ” in (3-29) in order to prove the uniqueness and regularity of its solutions Θ .

For this, we write

$$\partial_N w_{\text{implicit}} = (N \cdot \nu)\partial_\nu w_{\text{implicit}} + (N - (N \cdot \nu)\nu) \cdot \nabla w_{\text{implicit}} \quad \text{on } \Gamma_{\text{out}}^0,$$

where $\nu = \nu^0$. Recall that by a standard result on single layer potentials—see Theorem A.1—we have

$$(N \cdot \nu)\partial_\nu w_{\text{implicit}}(x) = \frac{1}{2}\Theta(x) + \tilde{\Theta}(x) \quad \text{on } \Gamma_{\text{out}}^0, \tag{3-36}$$

where

$$\tilde{\Theta}(x) := \int_{\Gamma_0} d\mathcal{H}^{n-1}(y) \left(-\frac{\Theta}{(N \cdot \nu)} \right)(y) \nu(x) \cdot \nabla P(x - y). \tag{3-37}$$

Note that the first term in the right-hand side of (3-36) is exactly the half of the first (and main) term in the left-hand side of (3-29). Using this and defining

$$\omega(x) := (N - (N \cdot \nu)\nu)(x) \quad \text{for } x \text{ on } \Gamma^0$$

we obtain

$$\frac{1}{2}\Theta = \frac{1}{2}\partial_{sss}u^0(\dot{\eta}^0 \circ \pi_1)^2 + \dot{\eta}^0 \partial_s N \cdot \nabla v + \partial_s(w_{\text{solid}} + w_{\text{single}} + w_{\text{double}}) + \omega \cdot \nabla w_{\text{implicit}} + \tilde{\Theta} \quad \text{on } \Gamma_{\text{out}}^0. \tag{3-38}$$

Step 3. From (3-38), we may deduce optimal regularity estimates for Θ , and hence for $\ddot{\eta}^0$. To do so we will bound each of the five terms in the right-hand side of (3-38) separately.

From here on, the constant C means $C = C(n, k, \alpha, \rho, \|h^0\|_{C^{k+1, \alpha}(\mathbb{R}^n)})$.

For the first term, we use that $h^0 \in C^{k+1, \alpha}$, and we obtain that $\Gamma^0 \in C_{\rho/4}^{k, \alpha}$, that $\nu^0 \in C^{k-1, \alpha}(\Gamma^0)$, and that $\eta^0 \in C^{k, \alpha}(\mathcal{Z})$ with estimates—here we are using the regularity estimates on Γ^0 from Proposition 2.1. In particular,

$$\|\pi_1\|_{C^{k, \alpha}(\Gamma^0)} + \|\nu^0\|_{C^{k-1, \alpha}(\Gamma^0)} \leq C. \tag{3-39}$$

Observe also that the vector field N is smooth and hence $\partial_{sss}u^0$ —the third derivative of u^0 along an integral curve of N —is regular as D^3u^0 .

Therefore,

$$\begin{aligned} & \left\| \frac{1}{2} \partial_{s s s} u^0 (\dot{\eta}^0 \circ \pi_1)^2 \right\|_{C^{k-2,\alpha}(\Gamma^0)} \\ & \leq C \left(\|u^0\|_{C^{k+1,\alpha}(\overline{B_R \cap \Omega^0})} \|(\dot{\eta}^0 \circ \pi_1)^2\|_{L^\infty(\Gamma^0)} + \|u^0\|_{L^\infty(B_R \cap \Omega^0)} \|(\dot{\eta}^0 \circ \pi_1)^2\|_{C^{k-2,\alpha}(\Gamma^0)} \right) \\ & \leq C \|(\dot{\eta}^0)^2\|_{C^{k-2,\alpha}(\mathcal{Z})} \\ & \leq C \mathcal{Q}. \end{aligned} \tag{3-40}$$

For the second term, we use again that N is smooth and recalling the estimate (3-13) for v and the estimate $\dot{\eta}$ in (3-7), we obtain

$$\begin{aligned} \|(\dot{\eta}^0 \circ \pi_1) \partial_{s s} v\|_{C^{k-2,\alpha}(\Omega^0)} & \leq C \left(\|\dot{\eta}^0\|_{C^{k-2,\alpha}(\mathcal{Z})} \|v\|_{L^\infty(B_R \cap \Omega^0)} + \|\dot{\eta}^0\|_{L^\infty(\mathcal{Z})} \|v\|_{C^{k,\alpha}(\overline{B_R \cap \Omega^0})} \right) \\ & \leq C \mathcal{Q}, \end{aligned} \tag{3-41}$$

where we used (3-7) and (3-13).

For the third term, we proceed as follows. From Lemma 3.7 we obtain that

$$\|\nabla w_{\text{solid}}\|_{C^{k-2,\alpha}(B_R \cap \Omega^0)} \leq C \|\Delta \dot{h}^0\|_{C^{k-2,\alpha}} \leq C \mathcal{Q}.$$

Next, since N and J are smooth, $\Delta h^0 \in C^{k-1,\alpha}$, $\Gamma^0 \in C^{k,\alpha}$, and $v^0 \in C^{k-1,\alpha}$ we obtain by Theorem A.1(i) that

$$\|w_{\text{single}}\|_{C^{k-1,\alpha}(\Omega^0)} \leq C \left(\|(\dot{\eta}^0 \circ \pi_1) \Delta \dot{h}^0\|_{C^{k-2,\alpha}(\Gamma^0)} + \|(\dot{\eta}^0 \circ \pi_1)^2\|_{C^{k-2,\alpha}(\Gamma^0)} \right) \leq C \mathcal{Q}$$

and by Theorem A.1(iii)

$$\|w_{\text{double}}\|_{C^{k-1,\alpha}(\Omega^0)} \leq C \|(\dot{\eta}^0 \circ \pi_1)^2\|_{C^{k-1,\alpha}(\Gamma^0)} \leq C \mathcal{Q}.$$

Hence,

$$\|\partial_s (w_{\text{solid}} + w_{\text{single}} + w_{\text{double}})\|_{C^{k-2,\alpha}(\Gamma^0)} \leq C \mathcal{Q}. \tag{3-42}$$

For the term $\omega \cdot \nabla w_{\text{implicit}}$ we use that Theorem A.1(i) yields

$$\|w_{\text{implicit}}\|_{C^{k-1,\alpha}(B_R \cap \Omega^0)} \leq C \|\Theta\|_{C^{k-2,\alpha}(\Gamma^0)},$$

and thus

$$\|\omega \cdot \nabla w_{\text{implicit}}\|_{C^{k-2,\alpha}(\Gamma^0)} \leq C \|\omega\|_{C^{k-2,\alpha}(\Gamma^0)} \|\Theta\|_{C^{k-2,\alpha}(\Gamma^0)}. \tag{3-43}$$

Also, recalling the definition of $\tilde{\Theta}$ in (3-37) and using Theorem A.1(iii) we obtain

$$\|\tilde{\Theta}\|_{C^{k-2,\alpha}(\Gamma^0)} \leq C \|\Theta\|_{C^{k-3,\alpha}(\Gamma^0)}. \tag{3-44}$$

Inserting (3-40)–(3-44) into (3-38), we obtain

$$\|\Theta\|_{C^{k-2,\alpha}(\Gamma^0)} \leq C \left(\mathcal{Q} + \|\omega\|_{C^{k-2,\alpha}(\Gamma^0)} \|\Theta\|_{C^{k-2,\alpha}(\Gamma^0)} + \|\Theta\|_{C^{k-3,\alpha}(\Gamma^0)} \right).$$

Note that we may take $\|\omega\|_{C^{k-2,\alpha}(\Gamma^0)}$ arbitrarily small by taking ε_o in (2-2) small enough. Then, by a standard interpolation argument we obtain

$$\|\Theta\|_{C^{k-2,\alpha}(\Gamma^0)} \leq C \mathcal{Q}. \tag{3-45}$$

Finally we recall the definition of Θ in (3-20), use that $v^0 \in C^{k-1,\alpha}$, $-\Delta h^0 \geq \rho$ and $\Delta h^0 \in C^{k-1,\alpha}$, and observe that $\pi_0^{-1} : \mathcal{Z} \rightarrow \Gamma^0$ satisfies $\|\pi_0^{-1}\|_{C^{k,\alpha}(\mathcal{Z})} \leq C$ with C universal, to obtain

$$\|\ddot{\eta}^0\|_{C^{k-2,\alpha}(\mathcal{Z})} \leq C\mathcal{Q}. \quad \square$$

4. Removing the a priori assumptions

In Section 3 we assumed the existence of the limits

$$\frac{\eta^{t_m} - \eta^0}{t_m} \rightarrow \dot{\eta}^0 \quad \text{and} \quad 2\frac{\eta^{t_m} - \eta^0 - \dot{\eta}^0 t_m}{t_m^2} \rightarrow \ddot{\eta}^0 \quad \text{in } C^0(\mathcal{Z}) \quad (4-1)$$

and we have shown that $\dot{\eta}^0$ and $\ddot{\eta}^0$ must then satisfy certain equations for which uniqueness and regularity estimates were proven.

The purpose of the next section is to prove that under our assumptions, (4-1) indeed holds for every sequence $t_m \downarrow 0$.

The setup. We start by introducing a new system of coordinates in $U_o \cap \bar{\Omega}^0$ that are adapted to u^0 .

Let us define

$$\sigma = \sigma(x) := \partial_N \tilde{u}_0(x). \quad (4-2)$$

Note that σ is defined in $U_o \cap \bar{\Omega}^0$ and takes positive values in that neighborhood of Γ^0 if U_o is chosen small enough. An application of the implicit function theorem gives that (z, σ) are $C^{k,\alpha}$ coordinates in $U_o \cap \bar{\Omega}^0$ (up to taking a smaller neighborhood U_o). Indeed, for $v = v^0$

$$\frac{\partial \sigma}{\partial s} = \partial_{ss} \tilde{u}^0 = (N \cdot v)^2 \partial_{vv} \tilde{u}^0 = (N \cdot v)^2 \Delta \tilde{u}^0 = -(N \cdot v)^2 \Delta h^0 \quad (4-3)$$

on Γ_{out}^0 and where by assumption $-\Delta h^0 \geq \rho > 0$ in a neighborhood of Γ^0 . Note in addition that the new coordinates (z, σ) are indeed $C^{k,\alpha}$ since $\tilde{u}^0 \in C^{k+1,\alpha}(\bar{\Omega}^0)$.

Let us also introduce

$$\bar{\pi}_1 : U_o \cap \Omega^0 \rightarrow \mathcal{Z}$$

to be the projection defined in the coordinates (z, σ) by

$$(z, \sigma) \mapsto (z, 0).$$

These coordinates are clearly related to the hodograph transform of the obstacle problem introduced in [Kinderlehrer and Nirenberg 1977]. Note also that for the case of the model solution to the obstacle problem $\frac{1}{2}(x_n)_+^2$, and with $N = e_n$ the coordinate σ would simply be x_n .

In view of Proposition 2.1 there exists $\lambda^t \in C^{k,\alpha}(\mathcal{Z})$ such that

$$\Gamma^t = \{\sigma = \lambda^t(z)\} \quad \text{for } t \in (0, t_o). \quad (4-4)$$

In the coordinates (z, σ) we have

$$\lambda^0 \equiv 0 \quad (4-5)$$

since $\sigma = \partial_N u^0 \equiv 0$ on Γ^0 . In addition, from (3-8) and the definition of the coordinate σ we have

$$\partial_N v^t = -\frac{\partial_N \tilde{u}^0}{t} = -\frac{\sigma}{t} = -\frac{\lambda^t}{t} \circ \bar{\pi}_1 \quad \text{on } \Gamma^t;$$

hence

$$\frac{\lambda^t}{t}(z) = -\partial_N v^t(z, \lambda^t(z)). \tag{4-6}$$

Indeed to prove (4-6) we use (3-8) and the definition of the coordinate σ to obtain

$$\partial_N v^t = -\frac{\partial_N \tilde{u}^0}{t} = -\frac{\sigma}{t} = -\frac{\lambda^t}{t} \circ \bar{\pi}_1 \quad \text{on } \Gamma^t.$$

The relation (4-6) will allow us to prove uniform $C^{k-1,\alpha}$ estimates for λ^t/t , leading to the existence of the limit as $t \downarrow 0$ of λ^t/t , which will be denoted by $\dot{\lambda}^0$. Later on, we will prove uniform $C^{k-2,\alpha}$ estimates for

$$\frac{1}{2} \frac{\lambda^t - \dot{\lambda}^0 t}{t^2} = \frac{1}{2} \frac{\lambda^t/t - \dot{\lambda}^0}{t},$$

which will lead to the existence of its limit as $t \rightarrow 0$, denoted by $\ddot{\lambda}^0$. These estimates will be deduced from the equation

$$\frac{\lambda^t/t - \dot{\lambda}^0}{t} = -\frac{\partial_N v(z, \lambda^t(z)) - \partial_N v(z, 0)}{t} - \partial_N w^t(z, \lambda^t(z)), \tag{4-7}$$

obtained from (4-6) by subtracting $\dot{\lambda}^0(z) = -\partial_N v(z, 0)$ from both sides, dividing by t on both sides, and recalling that by definition $w^t = (v^t - v)/t$.

Estimate on λ^t/t . The goal of this subsection is to prove a regularity result (without a priori assumptions) on λ^t/t . We state it next.

Proposition 4.1. *For $t \in (0, t_0)$ we have*

$$\left\| \frac{\lambda^t}{t} \right\|_{C^{k-1,\alpha}(\mathcal{Z})} \leq C(\mathcal{C}).$$

Before proving Proposition 4.1, let us state its main corollary

Corollary 4.2. *There exist $\dot{\eta}^0$ and $\dot{\lambda}^0$ such that*

$$\frac{\eta^t - \eta^0}{t} \rightarrow \dot{\eta}^0 \quad \text{and} \quad \frac{\lambda^t}{t} \rightarrow \dot{\lambda}^0 \quad \text{in } C^0(\mathcal{Z})$$

as $t \downarrow 0$.

Proof. Let $t_p \downarrow 0$. Note that both coordinate systems (z, s) and (z, σ) are $C^{k,\alpha}$. Hence, the estimate in Proposition 4.1 implies

$$\left\| \frac{\eta^t - \eta^0}{t} \right\|_{C^{k-1,\alpha}(\mathcal{Z})} \leq C$$

and by Arzelà–Ascoli there is a subsequence t_m such that

$$\frac{\eta^{t_m} - \eta^0}{t_m} \rightarrow \ell_1 \quad \text{and} \quad \frac{\lambda^{t_m}}{t_m} \rightarrow \ell_2 \quad \text{in } C^0(\mathcal{Z})$$

for certain limit functions ℓ_1 and ℓ_2 in $C^{k-1,\alpha}(\mathcal{Z})$. Applying [Proposition 3.2](#), we must have $\ell_1 = \dot{\eta}^0$, where $\dot{\eta}^0$ is the function given by [\(3-6\)](#). Then, either using the change of variables between s and σ or passing to the limit in [\(4-6\)](#) we obtain

$$\ell_2(z) = \dot{\lambda}^0(z) := \partial_N v(z, \sigma = 0).$$

Therefore, we have proven that each sequence has a subsequence converging to a limit that is independent of the sequence. In other words the limits as $t \downarrow 0$ exist and are given by $\dot{\eta}^0$ and $\dot{\lambda}^0$. \square

In view of [\(4-6\)](#), [Proposition 4.1](#) is an immediate consequence of the following:

Lemma 4.3. *For $t \in (0, t_0)$ we have*

$$\left\| \partial_N v^t(\cdot, \lambda^t(\cdot)) - \frac{1}{2} \frac{\lambda^t}{t} \right\|_{C^{k-1,\alpha}(\mathcal{Z})} \leq C(\mathcal{C}) + \frac{1}{100} \left\| \frac{\lambda^t}{t} \right\|_{C^{k-1,\alpha}(\mathcal{Z})}.$$

Next we state a sequence of lemmas aimed at proving [Lemma 4.3](#). To study the regularity of $\partial_N v^t$, let us write down (for the first time) the equation for $v^t = \frac{1}{t}(\tilde{u}^t - \tilde{u}^0)$ in all of \mathbb{R}^n . We have

$$\begin{cases} \Delta v^t = -\frac{\Delta h^0}{t} \chi_{\Omega^0 \setminus \Omega^t} + \Delta \delta_t h^0 \chi_{\Omega^t} & \text{in } \mathbb{R}^n, \\ v^t(\infty) = \delta_t c^0 \quad (n \geq 3), \end{cases} \quad \begin{cases} \Delta v^t = -\frac{\Delta h^0}{t} \chi_{\Omega^0 \setminus \Omega^t} + \Delta \delta_t h^0 \chi_{\Omega^t} & \text{in } \mathbb{R}^n, \\ \lim_{x \rightarrow \infty} \frac{v^t(x)}{-\log|x|} = \delta_t c^0 \quad (n = 2). \end{cases} \quad (4-8)$$

Hence, we may decompose v^t as

$$v^t = v_1^t + v_2^t + \text{constant},$$

where

$$v_1^t(x) := - \int_{\mathbb{R}^n} dy \left(\frac{\Delta h^0}{t} \chi_{\Omega^0 \setminus \Omega^t} \right)(y) P(x - y), \quad (4-9)$$

$$v_2^t(x) := - \int_{\mathbb{R}^n} dy \Delta \delta_t h^0 \chi_{\Omega^t}(y) P(x - y). \quad (4-10)$$

To prove [Lemma 4.3](#) we will deal separately with the two contributions $\partial_N v_1$ and $\partial_N v_2$ to $\partial_N v$.

Note that $\partial_N v_1$ is an ‘‘approximate single-layer potential’’. To study its regularity we need the next lemma. Before giving its statement, we need to introduce some notation.

We denote by

$$\bar{J}(z, \sigma) := |\det D(z, \sigma)|^{-1}$$

the Jacobian of the coordinates (z, σ) defined by

$$\int_A f(x) dx = \int_{(z,\sigma)(A)} f(z, \sigma) \bar{J}(z, \sigma) dz d\sigma. \quad (4-11)$$

Also, for $\theta \in (0, 1)$ we define

$$\begin{aligned} \Omega_\theta^t &:= \{x \in U \cap \Omega^0 : \sigma(x) > \theta \lambda^t(z(x))\} \cup (\Omega^0 \setminus U), \\ \Gamma_\theta^t &:= \partial \Omega_\theta^t = \{\sigma = \theta \lambda^t(z)\}, \end{aligned}$$

and denote by v_θ^t the unit normal to Γ_θ^t towards Ω_θ^t . Although the following lemma will be used in this subsection for $F \equiv -\Delta h^0$, we write it for general F for later use.

Lemma 4.4. *Let V be the single-layer potential*

$$V(x) = \int_{\mathbb{R}^n} dy \left(\frac{1}{t} F \chi_{\Omega^t \setminus \Omega^0} \right)(y) P(x - y). \tag{4-12}$$

We may write

$$V = \int_0^1 V^\theta d\theta, \tag{4-13}$$

where

$$V^\theta = \int_{\Gamma_\theta^t} \mathcal{H}^{n-1}(y) \left(F \frac{\lambda^t}{t} \circ \bar{\pi}_1 \frac{(N \cdot v_\theta^t)}{\partial_{ss} u^0} \right)(y) P(x - y)$$

and for all $\theta \in (0, 1)$ we have

$$\|\nabla V^\theta\|_{C^{k-1,\alpha}(\bar{\Omega}^t)} \leq C(\mathcal{C}) \left\| F \frac{\lambda^t}{t} \circ \bar{\pi}_1 \right\|_{C^{k-1,\alpha}(\Gamma_\theta^t)}. \tag{4-14}$$

Before giving the proof of the previous lemma let us give the analogue to [Lemma 3.3](#) in the present context.

Lemma 4.5. *Given $f : \Gamma_\theta^t \rightarrow \mathbb{R}$ continuous we have*

$$\int_{\Gamma_\theta^t} \frac{(N \cdot v_\theta^t)}{\partial\sigma/\partial s}(x) f(x) d\mathcal{H}^{n-1}(x) = \int_Z f(z, \theta\lambda^t(z)) \bar{J}(z, \theta\lambda^t(z)) dz.$$

Proof. Let us assume without loss of generality that f is continuously extended in a neighborhood of Γ_θ^t contained in $U_\circ \cap \bar{\Omega}^0$. Given $\varepsilon > 0$ let

$$A^\varepsilon := \{x \in U_\circ : \theta\lambda^t(z(x)) \leq \sigma(x) \leq \theta\lambda^t(z(x)) + \varepsilon\}.$$

Recalling that $(\partial\sigma/\partial s)\partial_\sigma = N = \partial_s$ and that $|N| = 1$, we have

$$\int_{\Gamma^0} \frac{(N \cdot v_\theta^t)}{\partial\sigma/\partial s}(x) f(x) d\mathcal{H}^{n-1}(x) = \lim_{\varepsilon \downarrow 0} \frac{1}{\varepsilon} \int_{A^\varepsilon} f(x) d\mathcal{H}^n(x).$$

On the other hand, for

$$(z, \sigma)(A^\varepsilon) := \{(z, \sigma) \in \mathcal{Z} \times (-\sigma_\circ, \sigma_\circ) : \theta\lambda^t(z) \leq \sigma \leq \theta\lambda^t(z) + \varepsilon\}$$

we have, by the definition of \bar{J} ,

$$\begin{aligned} \frac{1}{\varepsilon} \int_{A^\varepsilon} f(x) d\mathcal{H}^n(x) &= \frac{1}{\varepsilon} \int_{(z,s)(A^\varepsilon)} f(z, s) \bar{J}(z, s) dz ds \\ &= \int_{\mathcal{Z}} dz \frac{1}{\varepsilon} \int_0^\varepsilon d\bar{s} f(z, \theta\lambda^t(z) + \bar{s}) \bar{J}(z, \theta\lambda^t(z) + \bar{s}) \\ &= \int_{\mathcal{Z}} f(z, \theta\lambda^t(z)) \bar{J}(z, \theta\lambda^t(z)) dz + o(1) \end{aligned}$$

as $\varepsilon \downarrow 0$ and the lemma follows. □

Proof of Lemma 4.4. The key idea in the proof is to think of an approximate single-layer potential as an average (or integral) of exact single-layer potentials. More precisely, using (4-11) we may write

$$\begin{aligned} \int \phi \Delta V &:= \frac{1}{t} \int_{\mathcal{Z}} dz \int_0^{\lambda^t(z)} d\sigma (F\phi\bar{J})(z, \sigma) \\ &= \int_{\mathcal{Z}} dz \frac{1}{t} \int_0^1 d\theta \lambda^t(z) (F\phi\bar{J})(z, \theta\lambda^t(z)) \\ &= \int_0^1 d\theta \int_{\{\sigma=\theta\lambda^t(z)\}} \frac{\lambda^t}{t} (F\phi)(y) \frac{(N \cdot v_\theta^t)(y)}{\partial\sigma/\partial s(y)} d\mathcal{H}^{n-1}(y), \end{aligned}$$

where we used Lemma 4.5.

Recalling that $\sigma = \partial_s u^0$, this proves (4-13).

To prove (4-14) we use that V^θ is a single-layer potential on the surface Γ_θ^t , with charge density

$$\left(\frac{\lambda^t}{t} \circ \bar{\pi}_1\right) F \frac{(N \cdot v_\theta^t)}{u_{ss}^0}.$$

Note that Proposition 2.1 yields $\|\lambda^t\|_{C^{k,\alpha}(\mathcal{Z})} \leq C$ and hence $\{\sigma = \theta\lambda^t(x)\}$ is $C^{k,\alpha}$ and its normal vector v_θ^t is $C^{k-1,\alpha}$. Recall also that $u^0 \in C^{k+1,\alpha}(\bar{\Omega}^0)$ and that $u_{ss}^0 \approx -(N \cdot v^0)^2 \Delta h^0 > 0$ in a neighborhood of Γ^0 . Then, if

$$F \frac{\lambda^t}{t} \circ \bar{\pi}_1 \in C^{k-1,\alpha},$$

it follows from Theorem A.1 that V^θ is $C^{k,\alpha}(\bar{\Omega}_\theta^t)$ and in particular V^θ is $C^{k,\alpha}(\bar{\Omega}^t)$ with the estimate (4-14). □

Recalling (4-9), and using Lemma 4.4 with $F = -\Delta h^0$, we may now write

$$v_1^t(x) = \int_0^1 V^\theta(x) d\theta, \tag{4-15}$$

where

$$V^\theta(x) := \int_{\Gamma_\theta^t} \left(-\Delta h^0 \frac{\lambda^t}{t} \circ \bar{\pi}_1 \frac{(N \cdot v_\theta^t)}{\partial_{ss} u^0}\right)(y) P(x - y) dy. \tag{4-16}$$

The following lemma is a straightforward consequence of Theorem A.1 in the Appendix.

Lemma 4.6. *Let V^θ be as in (4-16). We have*

$$-\partial_{v_\theta^t, \text{out}} V^\theta = \frac{1}{2} (N \cdot v_\theta^t) \frac{-\Delta h^0}{\partial_{ss} u^0} \frac{\lambda^t}{t} \circ \bar{\pi}_1 + \partial_{v_\theta^t, 0} V^\theta \quad \text{on } \Gamma_\theta^t, \tag{4-17}$$

where

$$\|\partial_{v_\theta^t, 0} V^\theta\|_{C^{k-1,\alpha}(\Gamma_\theta^t)} \leq C(\mathcal{C}) \left\| \frac{\lambda^t}{t} \right\|_{C^{k-2,\alpha}(\mathcal{Z})}.$$

Proof. We recall that $(N \cdot v_\theta^t)$, $-\Delta h^0$, $\partial_{ss} u^0 > \rho/2 > 0$, and $\bar{\pi}_1^{-1} : \mathcal{Z} \rightarrow \Gamma_\theta^t$, $-\Delta h^0$ are all $C^{k-1,\alpha}$ functions. Then, the lemma follows from Theorem A.1(ii)–(iii). □

The next lemma will be used to control the “difference”

$$-\partial_N V^\theta(z, \eta^t(z)) - \frac{1}{2} \frac{\lambda^t}{t}(z).$$

Lemma 4.7. *Let V^θ be as in (4-16). We have*

$$\left\| -\partial_N V^\theta(\cdot, \eta^t(\cdot)) - \frac{1}{2} \frac{\lambda^t}{t} \right\|_{C^{k-1,\alpha}(\mathcal{Z})} \leq C(\mathcal{C}) + \frac{1}{100} \left\| \frac{\lambda^t}{t} \right\|_{C^{k-1,\alpha}(\mathcal{Z})}.$$

Proof. Step 1. We estimate the $C^{k-1,\alpha}(\mathcal{Z})$ norm of

$$I_1(z) := \partial_N V^\theta(z, \lambda^t(z)) - \partial_{N,\text{out}} V^\theta(z, \theta \lambda^t(z)).$$

To do it we write this difference as

$$I_1 = t \int_\theta^1 d\bar{\theta} \partial_\sigma \partial_N V^s(x', \bar{\theta} \lambda^t(x')) \frac{\lambda^t}{t}(x').$$

Then, using Lemma 4.4 we obtain

$$\begin{aligned} \|I_1\|_{C^{k-1,\alpha}(\mathcal{Z})} &\leq Ct \left(\|V^\theta\|_{C^{k+1,\alpha}(\Omega'_\theta)} \left\| \frac{\lambda^t}{t} \right\|_{L^\infty(\mathcal{Z})} + \|V^\theta\|_{L^\infty(\Omega'_\theta)} \left\| \frac{\lambda^t}{t} \right\|_{C^{k-1,\alpha}(\mathcal{Z})} \right) \\ &\leq Ct \left\| \frac{\lambda^t}{t} \right\|_{C^{k,\alpha}(\mathcal{Z})} \left\| \frac{\lambda^t}{t} \right\|_{L^\infty(\mathcal{Z})} \leq C \|\lambda^t\|_{C^{k,\alpha}(\mathcal{Z})} \left\| \frac{\lambda^t}{t} \right\|_{L^\infty(\mathcal{Z})} \leq C, \end{aligned} \tag{4-18}$$

where $C = C(\mathcal{C})$. Here we have used the fact that $\|\lambda^t/t\|_{L^\infty(\mathcal{Z})} \leq C$ and information that follows from Proposition 2.1.

Step 2. We next estimate the $C^{k-1,\alpha}(\mathcal{Z})$ norm of

$$I_2(z) := \partial_{N,\text{out}} V^\theta(z, \theta \lambda^t(z)) - \frac{1}{2} \frac{\lambda^t}{t}(z).$$

Using (4-17) we have

$$I_2(z) = (N - v_\theta^t) \cdot \nabla_{\text{out}} V^\theta(z, \theta \lambda^t(z)) + \frac{1}{2} \left((N \cdot v_\theta^t) \frac{-\Delta h^0}{\partial_{ss} u^0} - 1 \right) \frac{\lambda^t}{t} \circ \bar{\pi}_1 + \partial_{v_\theta^t, 0} V^\theta.$$

Using the estimates from Lemma 4.6 and 4.4 we have

$$\|\nabla V^\theta\|_{C^{k-1,\alpha}(\Gamma'_\theta)} \leq \|V^\theta\|_{C^{k,\alpha}(\bar{\Omega}'_\theta)} \leq C \left\| \frac{\lambda^t}{t} \right\|_{C^{k-1,\alpha}(\mathcal{Z})}.$$

In addition,

$$|N - v_\theta^t| \approx 0, \quad (N \cdot v_\theta^t) \approx 1, \quad \text{and} \quad \frac{-\Delta h^0}{\partial_{ss} u^0} \approx 1 \quad \text{on} \quad \Gamma'_\theta$$

for $t \in (0, t_o)$, where $X \approx Y$ means that “ X is arbitrarily close to Y ” provided that t_o and ε_o are chosen small enough depending only of \mathcal{C} .

Therefore, using the estimate in Lemma 4.6 and an interpolation inequality, we obtain

$$\begin{aligned} \|I_2\|_{C^{k-1,\alpha}(\mathcal{Z})} &\leq \frac{\varepsilon}{2} \left\| \frac{\lambda^t}{t} \right\|_{C^{k-1,\alpha}(\mathcal{Z})} + C \left\| \frac{\lambda^t}{t} \right\|_{C^{k-2,\alpha}(\mathcal{Z})} \\ &\leq \varepsilon \left\| \frac{\lambda^t}{t} \right\|_{C^{k-1,\alpha}(\mathcal{Z})} + C \left\| \frac{\lambda^t}{t} \right\|_{L^\infty(\mathcal{Z})} \leq \varepsilon \left\| \frac{\lambda^t}{t} \right\|_{C^{k-1,\alpha}(\mathcal{Z})} + C, \end{aligned} \tag{4-19}$$

where $\varepsilon > 0$ can be taken arbitrarily small by decreasing, if necessary, t_o and ε_o .

Step 3. We conclude by the triangle inequality that

$$\left\| \partial_N V^\theta(\cdot, \eta^t(\cdot)) - \frac{1}{2} \frac{\lambda^t}{t} \right\|_{C^{k-1,\alpha}(\mathcal{Z})} \leq \|I_1\|_{C^{k-1,\alpha}(\mathcal{Z})} + \|I_2\|_{C^{k-1,\alpha}(\mathcal{Z})}$$

and the lemma follows from (4-18) and (4-19), setting $\varepsilon = \frac{1}{100}$. □

Lemmas 4.4, 4.6, and 4.7 will be used to treat the term $\partial_N v_1$. As a counterpart, the next lemma will be used to treat the term $\partial_N v_2$.

Lemma 4.8. *We have*

$$\|v_2^t\|_{C^{k,\alpha}(\bar{\Omega}^t)} \leq C(\mathcal{C}).$$

Proof. Recalling that $\Delta v_2^t = \Delta \delta_t h^0 \chi_{\Omega^t}$ and that $\Gamma^t = \partial \Omega^t$ are (uniformly) $C^{k,\alpha}$, it follows from Lemma 3.7 that

$$\|v_2^t\|_{C^{k,\alpha}(\bar{\Omega}^t)} \leq C(\mathcal{C}) \|\Delta \delta_t h^0\|_{C^{k-2,\alpha}(\mathbb{R}^n)}.$$

Using the trivial estimate

$$\|\Delta \delta_t h^0\|_{C^{k-2,\alpha}(\mathbb{R}^n)} \leq \|h\|_{C^{k+1,\alpha}([-1,1] \times \mathbb{R}^n)}$$

the lemma follows. □

Proof of Lemma 4.3. We have

$$\partial_N v^t(z, \lambda^t(z)) = (\partial_N v_1^t + \partial_N v_2^t)(z, \lambda^t(z)),$$

and by (4-15)–(4-16) we have

$$\partial_N v_1^t(z, \lambda^t(z)) = \int_0^1 \partial_N V^\theta(z, \lambda^t(z)) d\theta.$$

Hence, by the triangle inequality, and using Lemmas 4.7 and 4.8,

$$\begin{aligned} \left\| \partial_N v^t(\cdot, \lambda^t) - \frac{1}{2} \frac{\lambda^t}{t}(z) \right\|_{C^{k-1,\alpha}(\mathcal{Z})} &\leq \int_0^1 d\theta \left\| \partial_N V^\theta(\cdot, \lambda^t) - \frac{1}{2} \frac{\lambda^t}{t} \right\|_{C^{k-1,\alpha}(\mathcal{Z})} + \|\partial_N v_2^t(\cdot, \lambda^t)\|_{C^{k-1,\alpha}(\mathcal{Z})} \\ &\leq C + \frac{1}{100} \left\| \frac{\lambda^t}{t} \right\|_{C^{k-1,\alpha}(\mathcal{Z})} + C \|\partial_N v_2^t\|_{C^{k-1,\alpha}(\Gamma^t)} \\ &\leq C + \frac{1}{100} \left\| \frac{\lambda^t}{t} \right\|_{C^{k-1,\alpha}(\mathcal{Z})}, \end{aligned}$$

where $C = C(\mathcal{C})$. □

Proof of Proposition 4.1. Recall (4-6), that is, $(\lambda^t/t)(z) = -\partial_N v^t(z, \lambda^t(z))$. Subtracting $\frac{1}{2}(\lambda^t/t)(z)$ from both sides and using Lemma 4.3 we obtain

$$\frac{1}{2} \left\| \frac{\lambda^t}{t} \right\|_{C^{k-1,\alpha}(\mathcal{Z})} \leq \left\| -\partial_N v^t(\cdot, \lambda^t(\cdot)) - \frac{1}{2} \frac{\lambda^t}{t} \right\|_{C^{k-1,\alpha}(\mathcal{Z})} \leq C(\mathcal{C}) + \frac{1}{100} \left\| \frac{\lambda^t}{t} \right\|_{C^{k-1,\alpha}(\mathcal{Z})}$$

as desired. □

Estimate on $\frac{1}{t}(\lambda^t/t - \dot{\lambda}^0)$. The goal of this subsection is to prove the following regularity result (without a priori assumptions).

Proposition 4.9. *We have*

$$\left\| \frac{1}{t} \left(\frac{\lambda^t}{t} - \dot{\lambda}^0 \right) \right\|_{C^{k-2,\alpha}(\mathcal{Z})} \leq C(\mathcal{C}).$$

Before proving [Proposition 4.9](#), let us give its main corollary.

Corollary 4.10. *There exist $\ddot{\eta}^0$ and $\ddot{\lambda}^0$ such that*

$$2 \frac{\eta^t - \eta^0 - t\dot{\eta}^0}{t^2} \rightarrow \ddot{\eta}^0 \quad \text{and} \quad 2 \frac{\lambda^t - t\dot{\lambda}^0}{t^2} \rightarrow \ddot{\lambda}^0 \quad \text{in } C^0(\mathcal{Z})$$

as $t \downarrow 0$.

Proof. Let $t_p \downarrow 0$. Note that since both coordinate systems (z, s) and (z, σ) are $C^{k,\alpha}$, the estimate of [Proposition 4.9](#) yields

$$\left\| \frac{\eta^t - \eta^0 - t\dot{\eta}^0}{t^2} \right\|_{C^{k-2,\alpha}(\mathcal{Z})} \leq C(\mathcal{C}).$$

Hence, by Arzelà–Ascoli there is a subsequence t_m such that

$$2 \frac{\eta^{t_m} - \eta^0 - t_m \dot{\eta}^0}{t_m^2} \rightarrow \ell_1 \quad \text{and} \quad 2 \frac{\lambda^{t_m} - t_m \dot{\lambda}^0}{t_m^2} \rightarrow \ell_2 \quad \text{in } C^0(\mathcal{Z})$$

for certain limit functions ℓ_1 and ℓ_2 in $C^{k-2,\alpha}(\mathcal{Z})$.

Applying [Proposition 3.5](#) the limit ℓ_1 must be $\ddot{\eta}^0$, the unique solution to (3-20)–(3-29). Using the change of variables between s and σ we obtain that there is also a unique possible limit $\ell_2(z) = \dot{\lambda}^0(z)$ which is independent of the subsequence.

In other words, the limits as $t \downarrow 0$ exist and they are denoted by $\ddot{\eta}^0$ and $\ddot{\lambda}^0$. □

In view of (4-7) and the regularity of $\partial_N v$, [Proposition 4.9](#) is a consequence of the following:

Lemma 4.11. *We have*

$$\left\| \partial_N w^t(\cdot, \lambda^t(\cdot)) - \frac{1}{2} \frac{\lambda^t/t - \dot{\lambda}^0}{t} \right\|_{C^{k-2,\alpha}(\mathcal{Z})} \leq C(\mathcal{C}) + \frac{1}{100} \left\| \frac{\lambda^t/t - \dot{\lambda}^0}{t} \right\|_{C^{k-2,\alpha}(\mathcal{Z})}.$$

Let us state a sequence of lemmas which will prove [Lemma 4.11](#). To study the regularity of $\partial_N w^t$ we will use the equation for w^t in all of \mathbb{R}^n that was obtained in (3-14).

As in Step 2 of the proof of [Proposition 3.5](#) we take the decomposition

$$w^t = w_1^t + w_2^t + \text{constant},$$

where, for $n \geq 3$,

$$w_1^t(x) = \int \left(\frac{1}{t} \Delta h^0 \chi_{\Omega^0 \setminus \Omega^t} - \frac{1}{2} \Delta \delta_t^2 h^0 \chi_{\Omega^t} \right) (dy) P(x - y), \quad w_1^t = 0,$$

and

$$w_2^t(x) = - \int \frac{1}{t} \left(\frac{\partial_N v}{N \cdot \nu^0} \mathcal{H}^{n-1} \upharpoonright_{\Gamma^0} - \frac{1}{t} \Delta h^0 \chi_{\Omega^0 \setminus \Omega^t} \right) (dy) P(x - y).$$

Respectively, for $n = 2$ we define w_1^t and w_2^t as the potentials of the previous Laplacians.

The analysis of the regularity in $\bar{\Omega}^t$ of w_1^t is done using Lemmas 4.4 and 3.7, which straightforwardly imply:

Lemma 4.12. *We have*

$$\|\nabla w_1^t\|_{C^{k-2,\alpha}(\bar{\Omega}^t)} \leq C(\mathcal{C}).$$

To study w_2^t let us further split it as

$$w_2^t = w_{21}^t + w_{22}^t + \text{constant},$$

where

$$w_{21}^t(x) = \int \frac{1}{t(N \cdot \nu^0)} \left(\partial_N v + (N \cdot \nu^0)^2 \frac{\Delta h^0}{\partial \sigma / \partial s} \frac{\lambda^t}{t} \right) \mathcal{H}^{n-1} \upharpoonright_{\Gamma^0}(dy) P(x-y),$$

$$w_{22}^t(x) = - \int \frac{1}{t} \left((N \cdot \nu^0) \frac{\Delta h^0}{\partial \sigma / \partial s} \frac{\lambda^t}{t} \mathcal{H}^{n-1} \upharpoonright_{\Gamma^0} - \frac{1}{t} \Delta h^0 \chi_{\Omega^0 \setminus \Omega^t} \right) (dy) P(x-y).$$

The study of $\partial_N w_{21}^t$ is done by observing that w_{21}^t is a single-layer potential and using Theorem A.1. Indeed we have:

Lemma 4.13. *We have*

$$\left\| \partial_N w_{21}^t(\cdot, \sigma = 0) - \frac{1}{2} \frac{\lambda^t/t - \dot{\lambda}^0}{t} \right\|_{C^{k-2,\alpha}(\mathcal{Z})} \leq C(\mathcal{C}) + \frac{1}{100} \left\| \frac{\lambda^t/t - \dot{\lambda}^0}{t} \right\|_{C^{k-2,\alpha}(\mathcal{Z})}.$$

Proof. Let

$$f(x) := \frac{1}{t(N \cdot \nu^0)} \left(\partial_N v + (N \cdot \nu^0)^2 \frac{\Delta h^0}{\partial \sigma / \partial s} \frac{\lambda^t}{t} \right) (x) = \frac{1}{t(N \cdot \nu^0)} \left(\dot{\lambda}^0 - \frac{\lambda^t}{t} \right) (x)$$

for $x \in \Gamma^0$. Here we have used that $\partial_N v = -\dot{\lambda} \circ \bar{\pi}_1$ and (4-3).

On one hand, by Theorem A.1(iii) we have

$$\partial_{\nu^0, \text{out}} w_{21}^t = \frac{1}{2} f + \partial_{\nu^0, 0} w_{21}^t \quad \text{on } \Gamma^0,$$

with

$$\|\partial_{\nu^0, 0} w_{21}^t\|_{C^{k-2,\alpha}(\Gamma^0)} \leq C \|f\|_{C^{k-3,\alpha}(\Gamma^0)}$$

and

$$\|w_{21}^t\|_{C^{k-1,\alpha}(\bar{\Omega}^0)} \leq C \|f\|_{C^{k-2,\alpha}(\Gamma^0)} \quad (n \geq 3),$$

$$\|\nabla w_{21}^t\|_{C^{k-2,\alpha}(\bar{\Omega}^0)} \leq C \|f\|_{C^{k-2,\alpha}(\Gamma^0)} \quad (n = 2),$$

where $C = C(\mathcal{C})$. Therefore, using that $|N - \nu^0| \leq \varepsilon$ we have

$$\begin{aligned} \|\partial_N w_{21}^t - \frac{1}{2} f\|_{C^{k-2,\alpha}(\Gamma^0)} &\leq C\varepsilon \|w_{21}^t\|_{C^{k-1,\alpha}(\bar{\Omega}^0)} + \|\partial_{\nu^0, 0} w_{21}^t\|_{C^{k-2,\alpha}(\Gamma^0)} \\ &\leq C\varepsilon \|f\|_{C^{k-2,\alpha}(\Gamma^0)} + C \|f\|_{C^{k-3,\alpha}(\Gamma^0)}, \end{aligned}$$

and the lemma follows using interpolation and choosing ε small enough. □

It thus remains to study the regularity of w_{22}^t , which we treat as an approximate double layer.

Lemma 4.14. *We have*

$$\|\nabla w_{22}^t\|_{C^{k-2,\alpha}(\bar{\Omega}^t)} \leq C(\mathcal{C}).$$

Proof. We will first write our approximate double layer as an average of double layers and we will then use the regularity results for the single layers to deduce the regularity of double layers.

Let us compute

$$\begin{aligned} -\int \phi \Delta w_{22}^t &= \int \phi(x) \frac{1}{t} \left((N \cdot \nu^0) \frac{\Delta h^0}{\partial \sigma / \partial s} \frac{\lambda^t}{t} \mathcal{H}^{n-1} \upharpoonright_{\Gamma^0} - \frac{1}{t} \Delta h^0 \chi_{\Omega^0 \setminus \Omega^t} \right) (x) dx \\ &= \frac{1}{t} \int_{\mathcal{Z}} dz \frac{\lambda^t}{t}(z) (\bar{J} \Delta h^0 \phi)(z, 0) - \frac{1}{t^2} \int_{\mathcal{Z}} dz \int_0^{\lambda^t(z)} d\sigma (\bar{J} \Delta h^0 \phi)(z, \sigma) \\ &= \int_0^1 d\theta \int_{\mathcal{Z}} dz \frac{\lambda^t}{t}(z) \frac{1}{t} \left((\bar{J} \Delta h^0 \phi)(z, 0) - (\bar{J} \Delta h^0 \phi)(z, \theta \lambda^t) \right) \\ &= - \int_0^1 d\theta \int_0^\theta d\theta' \int_{\mathcal{Z}} \left(\frac{\lambda^t}{t} \right)^2 (z) dz \partial_\sigma (\bar{J} \Delta h^0 \phi)(z, \theta' \lambda^t) \\ &= - \int_0^1 d\theta \int_0^\theta d\theta' \int_{\Gamma_{\theta'}^t} \left(\frac{\lambda^t}{t} \right)^2 \circ \bar{\pi}_1 \partial_\sigma (\bar{J} \Delta h^0 \phi) \frac{(N \cdot \nu_{\theta'}^t)}{\partial \sigma / \partial s}, \end{aligned}$$

where we have used Lemma 4.5. Changing the order of integration we find

$$\begin{aligned} -\int \phi \Delta w_{22}^t &= - \int_0^1 (1 - \theta) d\theta \int_{\Gamma_\theta^t} \left(\frac{\lambda^t}{t} \right)^2 \circ \bar{\pi}_1 \partial_\sigma (\bar{J} \Delta h^0 \phi) \frac{(N \cdot \nu_\theta^t)}{\partial \sigma / \partial s} \\ &= - \int_0^1 (1 - \theta) d\theta \int_{\Gamma_\theta^t} \left(\frac{\lambda^t}{t} \right)^2 \circ \bar{\pi}_1 \partial_s (\bar{J} \Delta h^0 \phi) \frac{(N \cdot \nu_\theta^t)}{(\partial \sigma / \partial s)^2}. \end{aligned}$$

Therefore, we have

$$w_{22}^t(x) = - \int_0^1 (1 - \theta) d\theta I_\theta(x) \tag{4-20}$$

for

$$I_\theta(x) := - \int_{\Gamma_\theta^t} d\mathcal{H}^{n-1}(y) \left(\frac{\lambda^t}{t} \right)^2 \circ \bar{\pi}_1(y) \partial_N ((\bar{J} \Delta h^0)(y) P(x - y)) \frac{(N \cdot \nu_\theta^t)}{(\partial \sigma / \partial s)^2}(y).$$

Note that

$$\begin{aligned} I_\theta(x) = I_1^\theta(x) + I_2^\theta(x) &=: \int_{\Gamma_\theta^t} d\mathcal{H}^{n-1}(y) \left(\left(\frac{\lambda^t}{t} \right)^2 \circ \bar{\pi}_1 \partial_N (\bar{J} \Delta h^0) \frac{(N \cdot \nu_\theta^t)}{(\partial \sigma / \partial s)^2} \right) (y) P(x - y) \\ &\quad + \operatorname{div}_x \left(\int_{\Gamma_\theta^t} d\mathcal{H}^{n-1}(y) \left(\left(\frac{\lambda^t}{t} \right)^2 \circ \bar{\pi}_1(y) (\bar{J} \Delta h^0) \frac{(N \cdot \nu_\theta^t)}{(\partial \sigma / \partial s)^2} N \right) (y) P(x - y) \right). \end{aligned}$$

Therefore, recalling that

$$\Gamma_\theta^t \in C^{k,\alpha}, \quad \frac{\lambda^t}{t} \in C^{k-1,\alpha}(\mathcal{Z}), \quad \bar{\pi}_1 \in C^{k,\alpha}(\Gamma_\theta^t), \quad \nu_\theta^t \in C^{k-1,\alpha}(\Gamma_\theta^t), \quad J \Delta h^0 \in C^{k-1,\alpha},$$

and $\partial\sigma/\partial s = u_{ss}^0$ positive and $C^{k-1,\alpha}$ and using [Theorem A.1](#) we obtain

$$\|\nabla I_1\|_{C^{k-2,\alpha}(\bar{\Omega}_\theta^t)} + \|\nabla I_2\|_{C^{k-2,\alpha}(\bar{\Omega}_\theta^t)} \leq C(\mathcal{C}).$$

The estimate of the lemma then follows from [\(4-20\)](#) observing that $\bar{\Omega}^t \subset \bar{\Omega}_\theta^t$ for all $\theta \in (0, 1)$. □

[Lemma 4.11](#) is now an immediate consequence of [Lemmas 4.12, 4.13, and 4.14](#), and [Proposition 4.9](#) follows.

5. Proof of the main result

In this section we conclude the proof of [Theorem 1.1](#). If one assumes $h^{\tau+t} - h^\tau$ satisfies $\Delta(h^{\tau+t} - h^\tau) \geq 0$ and $c^{\tau+t} - c^\tau \leq 0$ for $\tau, t \in (0, t_o)$ then [Theorem 1.1](#) is a straightforward consequence of the results developed in [Sections 2–5](#). Hence, the main issue that needs to be addressed is how to remove these technical sign assumptions. This is done by using a decomposition of the form

$$h^t - h^0 = \xi_+^t + \xi_-^t, \tag{5-1}$$

where $\Delta(\xi_+^{\tau+t} - \xi_+^\tau) \geq 0$ and $\lim_{x \rightarrow \infty} (\xi_+^{\tau+t} - \xi_+^\tau) \geq 0$ and the same with ξ_+ replaced by ξ_- and \geq replaced by \leq . This decomposition is defined as follows. We let

$$\phi_+(z) := 1 + \frac{1 + ze^z}{e^z + e^{-z}} \quad \text{and} \quad \phi_-(z) = -1 + \frac{-1 + ze^{-z}}{e^z + e^{-z}}$$

and note that

$$\phi_+ + \phi_- = z \tag{5-2}$$

and that ϕ_+ is similar to x^+ (the positive part), while ϕ_- is similar to $-x^-$ (minus the negative part) at large scales.

Let ζ be a radial smooth cutoff function with $\zeta \equiv 1$ in B_R and $\zeta \equiv 0$ outside of B_{2R} . For $t \in (-t_o, t_o)$ and $x \in \mathbb{R}^n$ let us define

$$\begin{aligned} \xi_+^t(x) &:= - \int_{\mathbb{R}^n} P(x-y) t \phi_+ \left(\frac{1}{t} \Delta(h^t - h^0) \right) \zeta(y), \\ \xi_-^t(x) &:= - \int_{\mathbb{R}^n} P(x-y) t \phi_- \left(\frac{1}{t} \Delta(h^t - h^0) \right) \zeta(y) \end{aligned}$$

Note that by definition we have, for τ and t small,

$$\begin{aligned} \Delta(\xi_+^{\tau+t} - \xi_+^\tau) &= \left((\tau+t)\phi_+ \left(\frac{1}{\tau+t} \Delta(h^{\tau+t} - h^0) \right) - \tau\phi_+ \left(\frac{1}{\tau} \Delta(h^\tau - h^0) \right) \right) \zeta \\ &= \left(\frac{d}{dt'} \Big|_{t'=\tau} \{t'\phi_+(\Delta\delta_{t'}h^0)\} t + O(t^{1+\alpha}) \right) \zeta \\ &= \left(\phi_+(\Delta\delta_\tau h^0)t + \tau\dot{\phi}_+(\Delta\delta_\tau h^0) \frac{d}{dt'} \Big|_{t'=\tau} (\Delta\delta_{t'}h^0)t + O(t^{1+\alpha}) \right) \zeta \\ &= \left(\phi_+(\Delta\delta_\tau h^0)t + \tau\dot{\phi}_+(\Delta\delta_\tau h^0) \frac{1}{\tau} O(\tau^\alpha) + O(t^{1+\alpha}) \right) \zeta \\ &\geq t(1 - C\tau^\alpha - Ct^\alpha)\zeta \geq 0, \end{aligned} \tag{5-3}$$

where in the passage from the third to the fourth line we have used that, since $h \in C^{3,\alpha}$,

$$\begin{aligned} \frac{d}{dt'} \Big|_{t'=\tau} (\Delta \delta_{t'} h^0) &= \Delta \frac{d}{dt'} \Big|_{t'=\tau} \left(\frac{h^{t'} - h^0}{t'} \right) = \Delta \left(-\frac{h^\tau - h^0}{\tau^2} + \frac{\dot{h}^\tau}{\tau} \right) \\ &= \left(-\frac{\Delta \dot{h}^\tau + O(\tau^{1+\alpha})}{\tau^2} + \frac{\Delta \dot{h}^\tau}{\tau} \right) = \frac{1}{\tau} O(\tau^\alpha). \end{aligned}$$

A similar inequality (with opposite sign) holds when $+$ is replaced by $-$. Moreover, by (5-2),

$$\Delta(\xi_+^t + \xi_-^t) = t\phi_+ \left(\frac{1}{t} \Delta(h^t - h^0) \right) \zeta + t\phi_- \left(\frac{1}{t} \Delta(h^t - h^0) \right) \zeta = \Delta(h^t - h^0)$$

since $\Delta(h^{s+t} - h^s) = 0$ outside of B_R and $\zeta = 1$ in B_R . Therefore (5-1) follows.

Next, for $t, \bar{t} \in (-t_o, t_o)$ we consider the two-parameter family of solutions to obstacle problems $u^{t,\bar{t}}$ defined as

$$\min\{-\Delta u^{t,\bar{t}}, u^{t,\bar{t}} - h^{t,\bar{t}}\} = 0 \quad \text{in } \mathbb{R}^n, \quad \begin{cases} \lim_{|x| \rightarrow \infty} u^{t,\bar{t}}(x) = c^{t,\bar{t}} & (n \geq 3), \\ \lim_{|x| \rightarrow \infty} \frac{u^{t,\bar{t}}(x)}{-\log|x|} = c^{t,\bar{t}} & (n = 2), \end{cases} \quad (5-4)$$

where

$$h^{t,\bar{t}} := h^0 + \xi_+^t + \xi_-^{\bar{t}}$$

and

$$c^{t,\bar{t}} := t\phi_- \left(\frac{1}{t} (c^t - c^0) \right) + \bar{t}\phi_+ \left(\frac{1}{\bar{t}} (c^{\bar{t}} - c^0) \right).$$

Note that

$$u^t = u^{t,t} \quad \text{and} \quad \eta^t = \eta^{t,t}.$$

Let us define

$$\Omega^{t,\bar{t}} := \{u^{t,\bar{t}} - h^{t,\bar{t}} > 0\} \quad \text{and} \quad \Gamma^{t,\bar{t}} := \partial\Omega^{t,\bar{t}}$$

and let $\eta^{t,\bar{t}} \in C^{k,\alpha}(\mathcal{Z})$ be defined by

$$\Gamma^{t,\bar{t}} = \{s = \eta^{t,\bar{t}}(z)\} \subset U_o. \quad (5-5)$$

In the proof of Theorem 1.1 the following observation will be useful.

Remark 5.1. For $e = (e^1, e^2) \in S^1$ making a small enough angle with $(1, 0)$ a computation similar to (5-3) shows that

$$\Delta(h^{t+e^1\bar{t}, \bar{t}+e^2\bar{t}} - h^{t,\bar{t}}) \geq 0 \quad \text{and} \quad c^{t+e^1\bar{t}, \bar{t}+e^2\bar{t}} - c^{t,\bar{t}} \leq 0 \quad (5-6)$$

for (t, \bar{t}) in a small neighborhood of $(0, 0)$. Thanks to this observation, the results developed in Sections 2–5 can be applied to obtain, in a neighborhood of $(0,0)$, estimates for the derivatives of $u^{t,\bar{t}}$ and $\eta^{t,\bar{t}}$ in a cone of directions (t, \bar{t}) . As a consequence, we obtain estimates for all the first and second derivatives $\partial_t, \partial_{\bar{t}}, \partial_{tt}, \partial_{\bar{t}\bar{t}}, \partial_{t\bar{t}}$ of $u^{t,\bar{t}}$ and $\eta^{t,\bar{t}}$ in a neighborhood of $(0, 0)$. In particular we obtain estimates in the direction $(1, 1)$ which are equivalent to estimates for $u^t = u^{t,t}$ and $\eta^t = \eta^{t,t}$.

Proof of Theorem 1.1. Step 1. Assuming that $k \geq 1$ we prove that $\eta^{t,\bar{t}}$ is once differentiable (jointly) in the two variables (t, \bar{t}) in a neighborhood of $(0, 0)$ with the estimate

$$\|\partial_e \eta^{t,\bar{t}}\|_{C^{k-1,\alpha}(\mathcal{Z})} \leq |e|C(\mathcal{C}) \tag{5-7}$$

and the formula

$$\partial_e \eta^{t,\bar{t}} = \left(\frac{\partial_N(\partial_e u^{t,\bar{t}})}{(N \cdot \nu^{t,\bar{t}})^2 \Delta h^{t,\bar{t}}} \right) (z, \eta^{t,\bar{t}}(x)), \tag{5-8}$$

which holds true for every vector e in the (t, \bar{t}) -plane.

Indeed, let $e_1 = (1, 0)$ and let e_2 be some different unit vector making a small enough angle with e_1 as in Remark 5.1.

By Remark 5.1, for fixed (t, \bar{t}) in a small enough neighborhood of $(0, 0)$ and for $i = 1, 2$, the one parameter family $(u^{t+e_i^1 \bar{t}, \bar{t}+e_i^2 \bar{t}})_{\bar{t}}$ satisfies the assumptions of Sections 2–4. Applying Corollary 4.2 to it, we find that

$$\partial_{e_i} \eta^{t,\bar{t}} := \left. \frac{d}{d\bar{t}} \right|_{\bar{t}=0} \eta^{t+e_i^1 \bar{t}, \bar{t}+e_i^2 \bar{t}}$$

exists in the sense that the limit defining this derivative exists in $C^0(\mathcal{Z})$.

Then, Proposition 3.2 yields the estimate

$$\|\partial_{e_i} \eta^{t,\bar{t}}\|_{C^{k-1,\alpha}(\mathcal{Z})} \leq C(\mathcal{C})$$

and the formula

$$\partial_{e_i} \eta^{t,\bar{t}} = \left(\frac{\partial_N \partial_{e_i} (u^{t,\bar{t}} - h^{t,\bar{t}})}{(N \cdot \nu^{t,\bar{t}})^2 \Delta h^{t,\bar{t}}} \right) (z, \eta^{t,\bar{t}}(x)).$$

Since $\partial_t = \partial_{e_1}$ and $\partial_{\bar{t}}$ is a linear combination of ∂_{e_i} we obtain that $\eta^{t,\bar{t}}$ is continuously differentiable (jointly) in the two variables (t, \bar{t}) in a neighborhood of $(0, 0)$ with the estimate (5-7) and formula (5-8).

Step 2. Applying (5-7) and formula (5-8) for (t, \bar{t}) restricted to the “diagonal” $t = \bar{t}$ (still in a neighborhood of $(0, 0)$) — i.e., with $e = (1, 1)$ — we obtain that η^t is differentiable with respect to t , with the estimate

$$\|\dot{\eta}^t\|_{C^{k-1,\alpha}(\mathcal{Z})} \leq C(\mathcal{C}) \tag{5-9}$$

and the formula

$$\dot{\eta}^t = \left(\frac{\partial_N(\dot{u}^t - \dot{h}^t)}{(N \cdot \nu^t)^2 \Delta h^t} \right) (z, \eta^t(x)). \tag{5-10}$$

Note that (5-9) and (5-10) are identical to those of Proposition 3.2 but now they are valid under more general assumptions (we do not need to assume the sign condition that implies that the contact sets are ordered).

Step 3. Similarly we obtain

$$\|\partial_{ee} \eta^{t,\bar{t}}\|_{C^{k-1,\alpha}(\mathcal{Z})} \leq |e|^2 C(\mathcal{C}). \tag{5-11}$$

Indeed, let e_1 and e_2 be as in Step 1 and let e_3 be a third vector such that the e_i are pairwise linearly independent and the angle of e_3 with $(1, 0)$ is small enough.

Using again [Remark 5.1](#), for fixed (t, \bar{t}) in a small enough neighborhood of $(0, 0)$ and for $i = 1, 2, 3$, the one parameter family $(u^{t+e_i^1\bar{t}, \bar{t}+e_i^2\bar{t}})_{\bar{t}}$ satisfies the assumptions of Sections 2–4. Applying [Corollary 4.10](#) we find that

$$\partial_{e_i e_i} \eta^{t, \bar{t}} := \left. \frac{d^2}{d\bar{t}^2} \right|_{\bar{t}=0} \eta^{t+e_i^1\bar{t}, \bar{t}+e_i^2\bar{t}}$$

exists in the sense that the limit defining this derivative exists in $C^0(\mathcal{Z})$.

Then, [Proposition 3.2](#) yields the estimate

$$\|\partial_{e_i e_i} \eta^{t, \bar{t}}\|_{C^{k-1, \alpha}(\mathcal{Z})} \leq C(\mathcal{C}).$$

Since for all e in the (t, \bar{t}) -plane ∂_{ee} is a linear combination of $\{\partial_{e_i e_i}\}_{i=1,2,3}$, we obtain that $\eta^{t, \bar{t}}$ is twice differentiable (jointly) in the two variables (t, \bar{t}) in a neighborhood of $(0, 0)$ with the estimate [\(5-11\)](#).

Step 4. Applying [\(5-11\)](#) or (t, \bar{t}) restricted to the “diagonal” $t = \bar{t}$ (still in a neighborhood of $(0, 0)$) — i.e., with $e = (1, 1)$ — we obtain that η^t is twice differentiable with respect to t , with the estimate

$$\|\ddot{\eta}^t\|_{C^{k-1, \alpha}(\mathcal{Z})} \leq C(\mathcal{C}). \tag{5-12}$$

Again note that [\(5-11\)](#) is identical to that of [Proposition 3.5](#) but now it is valid under more general assumptions.

Step 5. Finally, we complete the proof of [Theorem 1.1](#) by defining the diffeomorphisms Ψ^t from the coordinates (z, s) and the function η^t . Let $\phi \in C_c^\infty(U_\circ)$ be some function such that $\phi \equiv 1$ in a neighborhood of Γ^0 . Let us define

$$\Psi^t(x) = \begin{cases} (z, s)^{-1}(z(x), s(x) + \eta^0(z(x)) + \phi(x)\{\eta^t(z(x)) - \eta^0(z(x))\}), & x \in U_\circ, \\ x, & x \in \mathbb{R}^n \setminus U_\circ. \end{cases}$$

Since we may take $U_\circ \subset U$ we have that Ψ^t fixes the complement of U . By the definition of η^t we easily show that $\Psi^t(\omega^0) = \Omega^t$ — and thus $\Psi^t(\Gamma^0) = \Gamma^t$.

It not difficult to check that [\(5-9\)](#), [\(5-10\)](#), and [\(5-12\)](#) yield [\(1-14\)](#), [\(1-15\)](#) and [\(1-16\)](#) when rewritten in terms of Ψ . On the other hand, estimate [\(1-17\)](#) follows from the estimates for w obtained in Step 3 of the proof of [Proposition 3.5](#). □

Appendix: Single-layer potentials and auxiliary proofs

We recall here classical regularity properties and the formula for the jump in the normal derivative for a single-layer potential.

Theorem A.1. *Let $U \subset B_R \subset \mathbb{R}^n$ be a domain such that $\partial U \in C_r^{m, \alpha}$ for some $r > 0$, $m \in \mathbb{N}$ and $\alpha \in (0, 1)$. Given $f \in C^{m-1, \alpha}(\partial U)$ let us define*

$$w(x) := \int_{\partial U} d\mathcal{H}^{n-1}(y) f(y) P(x - y),$$

where P is the Newtonian potential.

We then have:

(i) $w \in C^0(\mathbb{R}^n)$, $w \in C^{m,\alpha}(\bar{U})$ and $w \in C^{m,\alpha}(\overline{\mathbb{R}^n \setminus U})$ with the estimate

$$\|w\|_{C^{m,\alpha}(\bar{U})} + \|w\|_{C^{m,\alpha}(\overline{\mathbb{R}^n \setminus U})} \leq C \|f\|_{C^{m-1,\alpha}(\partial U)},$$

where C depends only on n, m, α, r , and $\|\partial U\|_{C_r^{m,\alpha}}$.

(ii) Denoting by $\partial_{v, \text{out}} w$ and $\partial_{v, \text{in}} w$ the (outward) normal derivatives of w from outside and inside U respectively we have, for all $x \in \partial U$,

$$\begin{aligned} \partial_{v, \text{out}} w(x) &= \partial_{v,0} w(x) - \frac{1}{2} f(x), \\ \partial_{v, \text{in}} w(x) &= \partial_{v,0} w(x) + \frac{1}{2} f(x), \end{aligned}$$

where

$$\partial_{v,0} w(x) := \int_{\partial U} d\mathcal{H}^{n-1}(y) f(y) \nu(x) \cdot \nabla P(x - y).$$

(iii) The linear operator $T : f \mapsto \partial_{v,0} w$ maps continuously $C^{m-2,\alpha}(\partial U)$ to $C^{m-1,\alpha}(\partial U)$. More precisely,

$$\|\partial_{v,0} w\|_{C^{m-1,\alpha}(\partial U)} \leq C \|f\|_{C^{m-2,\alpha}(\partial U)},$$

where C depends only on n, m, α, R, r , and $\|\partial U\|_{C_r^{m,\alpha}}$. In particular T is compact in Hölder spaces.

We provide the following for completeness.

Bibliographic references and sketch of the proof of Theorem A.1. Properties of single-layer potentials in the spirit of (i)–(iii) — and related ones for double-layer potentials — are very classical results in potential theory. They are key tools in proving the existence of solutions for the Dirichlet and Neumann problems in $C^{1,\alpha}$ domains by the method of boundary potentials (by solving in Hölder spaces Fredholm integral equations on the boundary of the domain). For more information on the topic, see for instance the classical books [Sobolev and Dawson 1964; Dautray and Lions 1990].

The proofs of (i), (ii) are given in [Dautray and Lions 1990, Section II.3]. The proof of (i) is given in full detail only for $m = 1$ but the proof for general m is similar. The result for all m is stated in [Dautray and Lions 1990, p. 303].

The compactness property of T in (iii) is in the core of the theory for solving the Dirichlet and Neumann problems by the method of boundary potentials. Indeed, by (ii), the Neumann problem $\Delta w = 0$ in U , $\partial_\nu w = g$ on ∂U is equivalent to $Tf + \frac{1}{2}f = g$, where f is the charge on the boundary. Since T is compact, this equation can be solved by Fredholm’s alternative;⁶ see [Sobolev and Dawson 1964, Lectures 15–19].

Roughly speaking, the reason why Tf increases the order of differentiability of f by 1 is that the integral kernel ($x \in \partial U$) satisfies

$$k(x, y) := \nu(x) \cdot \nabla P(x - y) = c_n \nu(x) \frac{x - y}{|x - y|^n} = O(|x - y|^{n-2})$$

⁶In this case the orthogonality condition of Fredholm’s alternative requires $\int_{\partial U} g = 0$.

as $y \rightarrow x$, $y \in \partial U$, while ∂U is an $(n-1)$ -dimensional surface. The extra factor $|x - y|$ comes from $v(x) \cdot (x - y) = O(|x - y|^2)$ since ∂U is smooth enough. Thus, Tf behaves similarly to

$$f \mapsto \int_{\mathbb{R}^d} f(y) \frac{e \cdot y}{|y|^d} dy,$$

which maps $C_c^{k-1,\alpha}(\mathbb{R}^d)$ to $C^{k,\alpha}(\mathbb{R}^d)$.

Since it is not easy to find complete references for (iii), although these types of estimates are very classical, for the sake of completeness we provide next a detailed proof of a nearly optimal estimate like (iii) in the case $m = 2$ (the proof for other m is more involved but similar). For the purposes of this paper the optimal estimate is not necessary — we just state the optimal result for the convenience of the reader. In our proofs, we do not need to gain a full derivative but just obtain a control in a finer Hölder norm to control the corresponding term by interpolation. Let us prove that if $\partial U \in C_r^{2,\alpha}$ then, for all $\beta \in (0, 1)$,

$$\|Tf\|_{C^{0,\beta}(\partial U)} \leq C \|f\|_{C^{0,\alpha}(\partial U)} \tag{A-1}$$

(note that the optimal estimate would be with $C^{1,\alpha}$ instead of $C^{0,\beta}$).

As a matter of fact we will prove the stronger (and almost sharp) estimate

$$\|Tf\|_{C^{0,\beta}(\partial U)} \leq C \|f\|_{L^\infty(\partial U)}, \tag{A-2}$$

which clearly yields (A-1).

Indeed, we start by showing that

$$k(x, y) := v(x) \frac{x - y}{|x - y|^n} \tag{A-3}$$

satisfies

$$|k(x, y) - k(\bar{x}, y)| \leq C|x - \bar{x}| |\xi - y|^{-n+1}, \tag{A-4}$$

where ξ is a point of a curve on ∂U joining x and \bar{x} .

Indeed, if $\gamma \subset \partial U$ is a smooth curve joining x and \bar{x} and of length comparable to $|x - \bar{x}|$ we have, at $\xi = \gamma(t)$,

$$\frac{d}{dt} k(\gamma(t), y) = v'(\xi) \frac{\xi - y}{|\xi - y|^n} + v_i(\xi) \gamma'_j(t) \frac{|z|^2 \delta_{ij} - n z_i z_j}{|z|^{n+2}} \quad \text{for } z = \xi - y.$$

Choosing an appropriate frame, we may assume $v_i(\xi) = \delta_{1i}$ and $\gamma'_j(t) = C \delta_{2j}$ — since the former vector is normal to ∂U and the latter is tangent. Therefore

$$\left| v_i(\xi) \gamma'_j(t) \frac{|z|^2 \delta_{ij} - n z_i z_j}{|z|^{n+2}} \right| = C \frac{|z_1 z_2|}{|z|^{n+2}} \leq C \frac{|z|^2 |z|}{|z|^{n+2}} \leq C |z|^{1-n} = C |\xi - y|^{1-n},$$

where we have used that the first axis is normal to ∂U and hence we have $|z_1| \leq |z|^2$ — by C^2 regularity of ∂U and recalling that $z = \xi - y$ with both ξ and y on ∂U . Therefore, an application of the mean value theorem gives

$$|k(x, y) - k(\bar{x}, y)| \leq C|x - \bar{x}| \left| \frac{d}{dt} k(\gamma(t), y) \right| \leq C|x - \bar{x}| |\xi - y|^{1-n}$$

and proves (A-4).

Finally, recalling that

$$|k(x, y)| \leq C|x - y|^{2-n} \quad \text{and} \quad |k(\bar{x}, y)| \leq C|\bar{x} - y|^{2-n}$$

and combining this with (A-4), we obtain

$$|k(x, y) - k(\bar{x}, y)| \leq C|x - \bar{x}|^\beta |\xi - y|^{(1-n)\beta} (|x - y|^{(2-n)(1-\beta)} + |\bar{x} - y|^{(2-n)(1-\beta)}).$$

Therefore

$$\begin{aligned} |Tf(x) - Tf(\bar{x})| &= \left| \int_{\partial U} f(y)(k(x, y) - k(\bar{x}, y)) d\mathcal{H}^{n-1}(z) \right| \\ &\leq \int_{\partial U} |f(y)| |k(x, y) - k(\bar{x}, y)| d\mathcal{H}^{n-1}(z) \\ &\leq C\|f\|_{L^\infty(\partial U)} \int_{\partial U} |x - \bar{x}|^\beta |\xi - y|^{(1-n)\beta} (|x - y|^{(2-n)(1-\beta)} + |\bar{x} - y|^{(2-n)(1-\beta)}) \\ &\leq C\|f\|_{L^\infty(\partial U)} |x - \bar{x}|^\beta, \end{aligned}$$

which proves (A-2). □

Sketch of the proof of Proposition 2.1. For the sake of clarity we give a proof assuming that, for $t > 0$, we have $\Delta(h^t - h^0) \geq 0$ and $h^t - h^0 \geq 0$ and thus $\Omega^t \subset \omega^0$. We give the proof in dimension $n = 2$. The proof for $n \geq 3$ is similar; see [Blank 2001].

Step 1. We show that for some $t_o > 0$ and C_o depending only on \mathcal{C} we have

$$|\Omega^0 \setminus \Omega^t| \leq C(\mathcal{C})t. \tag{A-5}$$

Indeed, from (4-8) we know that (recall that $v^t := \delta_t \tilde{u}^0$)

$$\begin{cases} \Delta v^t = -(\Delta h^0/t)\chi_{\Omega^0 \setminus \Omega^t} + \Delta \delta_t h^0 \chi_{\Omega^t} & \text{in } \mathbb{R}^2, \\ \lim_{x \rightarrow \infty} \frac{v^t(x)}{-\log|x|} = \delta_t c^0. \end{cases}$$

Note that by (1-10) we have $\Gamma^t \subset B_R$ for $t \in [0, t_o)$, where $t_o > 0$ is a small enough constant depending only on \mathcal{C} . Recalling that by assumption $\Delta \delta_t h^0$ is supported in B_R , we have

$$\delta_t c^0 = \int_{\mathbb{R}^2} \Delta v^t = \int_{\mathbb{R}^2} -\frac{\Delta h^0}{t} \chi_{\Omega^0 \setminus \Omega^t} + \Delta \delta_t h^0 \chi_{\Omega^t}.$$

Therefore, since $-\Delta h^0 \geq \rho$, we find

$$\frac{\rho}{t} |\Omega^0 \setminus \Omega^t| \leq |\delta_t c^0| + \int_{B_R} |\Delta \delta_t h^0| \leq C(\mathcal{C}).$$

Step 2. We first show (i); that is, we prove that for t_o small enough we have

$$\|\Gamma^t\|_{C_{\rho/4}^{k,\alpha}} \leq C_o \quad \text{for all } t \in [0, t_o). \tag{A-6}$$

Indeed, by Step 1, $|\Omega^0 \setminus \Omega^t| \downarrow 0$ as $t \rightarrow 0$ and hence, for t small enough, all points of Γ^t are regular points. More precisely, for all $p \in \Gamma^t$,

$$B_\rho(p) \cap \{\tilde{u}^t = 0\} \geq c_o(\mathcal{C}) > 0.$$

Then, we apply:

- (1) $C^{1,\alpha}$ free boundary estimates near regular points [Caffarelli 1977; 1998].
- (2) $C^{1,\alpha} \Rightarrow C^{k,\alpha}$ estimates for obstacle $h \in C^{k+1,\alpha}$ [Kinderlehrer and Nirenberg 1977].

We thus obtain (A-6).

Step 3. From (A-5) and (A-6) deduce that for $t \in (0, t_o)$, the Hausdorff distance between Γ^t and Γ^s satisfies

$$d_{\text{Hausdorff}}(\Gamma^t, \Gamma^0) \leq C_o t. \quad \square$$

Sketch of the proof of Lemma 3.1. The lemma for $n \geq 3$ is very standard. Let us prove it in the case $n = 2$.

Assume $n = 2$. We want to prove that $u^t = f_*$, where

$$f_*(x) := \inf \left\{ f(x) : f \in C(\mathbb{R}^2), f \geq h^t, \Delta f \leq 0, \lim_{x \rightarrow \infty} \frac{f}{-\log|x|} = c^t \right\}. \quad (\text{A-7})$$

The admissible class in (A-7) is nonempty since the function

$$f_1(x) := c^t \min\{0, -\log|x|\} + C_1$$

is a member, provided we take $C_1 > 0$ large enough that $\log|x| + C > h^t(x)$ for all $x \in \mathbb{R}^2$ — here we are using (1-4). Hence, $f_*(x) \in [h^t(x), +\infty)$ is finite for all x .

We now check that $u^t = f_*$ is a solution of (1-3) ($n = 2$). First, as an infimum of superharmonic functions, it is superharmonic. To check that it is a subsolution of the obstacle problem, we argue by contradiction. Suppose on the contrary that there exist $r, \varepsilon, \delta > 0$ (as small as we like) and $x_o \in \mathbb{R}^2$ such that $f_* > \varepsilon + h^t$ in $B_r(x_o)$ and $f_*(x_o) > \delta + \int_{\partial B_r(x_o)} f_*$. By changing (slightly) x_o and making r and δ smaller, if necessary, we may assume $\delta < \varepsilon$ and

$$\text{osc}_{B_r(x_o)} h^t \leq \varepsilon \implies f_* > \sup_{B_r(x_o)} h^t.$$

Let $\tilde{f} \in C(B_r(x_o))$ be the unique harmonic function in $B_r(x_o)$ with Dirichlet boundary condition $\tilde{f} = \frac{1}{2}\delta + f_*$ on $\partial B_r(x_o)$. Note that $\tilde{f} > h^t$ in $B_r(x_o)$, and set

$$f(x) := \begin{cases} f_*(x), & x \notin B_r(x_o), \\ \min\{\tilde{f}, f_*(x)\}, & x \in B_r(x_o). \end{cases}$$

Then f is admissible in (A-7) and hence $f_* \leq f$. But then by the mean value formula for \tilde{f} we have

$$f_*(x_o) \leq f(x_o) \leq \tilde{f}(x_o) = \frac{1}{2}\delta + \int_{B_r(x_o)} f_* \leq \frac{1}{2}\delta + f_*(x_o) - \varepsilon < f_*(x_o),$$

a contradiction. □

Acknowledgments

Serfaty was supported by the Institut Universitaire de France and NSF grant DMS-1700278. Serra is supported by the ERC Grant “Regularity and stability in partial differential equations (RSPDE)”.

References

- [Ameur et al. 2011] Y. Ameur, H. Hedenmalm, and N. Makarov, “Fluctuations of eigenvalues of random normal matrices”, *Duke Math. J.* **159**:1 (2011), 31–81. [MR](#) [Zbl](#)
- [Blank 2001] I. Blank, “Sharp results for the regularity and stability of the free boundary in the obstacle problem”, *Indiana Univ. Math. J.* **50**:3 (2001), 1077–1112. [MR](#) [Zbl](#)
- [Caffarelli 1977] L. A. Caffarelli, “The regularity of free boundaries in higher dimensions”, *Acta Math.* **139**:3-4 (1977), 155–184. [MR](#) [Zbl](#)
- [Caffarelli 1998] L. A. Caffarelli, “The obstacle problem revisited”, *J. Fourier Anal. Appl.* **4**:4-5 (1998), 383–402. [MR](#) [Zbl](#)
- [Dautray and Lions 1990] R. Dautray and J.-L. Lions, *Mathematical analysis and numerical methods for science and technology, I: Physical origins and classical methods*, Springer, 1990. [MR](#) [Zbl](#)
- [Ekeland and Temam 1976] I. Ekeland and R. Temam, *Convex analysis and variational problems*, Studies in Mathematics and its Applications **1**, North-Holland, Amsterdam, 1976. [MR](#) [Zbl](#)
- [Hedenmalm and Makarov 2004] H. Hedenmalm and N. Makarov, “Quantum Hele-Shaw flow”, preprint, 2004. [arXiv](#)
- [Hedenmalm and Makarov 2013] H. Hedenmalm and N. Makarov, “Coulomb gas ensembles and Laplacian growth”, *Proc. Lond. Math. Soc.* (3) **106**:4 (2013), 859–907. [MR](#) [Zbl](#)
- [Johansson 1998] K. Johansson, “On fluctuations of eigenvalues of random Hermitian matrices”, *Duke Math. J.* **91**:1 (1998), 151–204. [MR](#) [Zbl](#)
- [Kinderlehrer and Nirenberg 1977] D. Kinderlehrer and L. Nirenberg, “Regularity in free boundary problems”, *Ann. Scuola Norm. Sup. Pisa Cl. Sci.* (4) **4**:2 (1977), 373–391. [MR](#) [Zbl](#)
- [Leblé and Serfaty 2018] T. Leblé and S. Serfaty, “Fluctuations of two dimensional Coulomb gases”, *Geom. Funct. Anal.* **28**:2 (2018), 443–508. [MR](#)
- [Saff and Totik 1997] E. B. Saff and V. Totik, *Logarithmic potentials with external fields*, Grundlehren der Mathematischen Wissenschaften **316**, Springer, 1997. [MR](#) [Zbl](#)
- [Sakai 1991] M. Sakai, “Regularity of a boundary having a Schwarz function”, *Acta Math.* **166**:3-4 (1991), 263–297. [MR](#) [Zbl](#)
- [Schaeffer 1975] D. G. Schaeffer, “A stability theorem for the obstacle problem”, *Advances in Math.* **17**:1 (1975), 34–47. [MR](#) [Zbl](#)
- [Serfaty 2015] S. Serfaty, *Coulomb gases and Ginzburg–Landau vortices*, European Mathematical Society, Zürich, 2015. [MR](#) [Zbl](#)
- [Sobolev and Dawson 1964] S. L. Sobolev and E. R. Dawson, *Partial differential equations of mathematical physics*, Pergamon Press, Oxford, 1964. [MR](#) [Zbl](#)
- [Wiegmann 2002] P. B. Wiegmann, “Aharonov–Bohm effect in the quantum Hall regime and Laplacian growth problems”, pp. 337–349 in *Statistical field theories* (Como, 2001), edited by A. Cappelli and G. Mussardo, NATO Sci. Ser. II Math. Phys. Chem. **73**, Kluwer Acad., Dordrecht, 2002. [MR](#) [Zbl](#)

Received 10 Aug 2017. Revised 13 Feb 2018. Accepted 9 Apr 2018.

SYLVIA SERFATY: serfaty@cims.nyu.edu

Courant Institute of Mathematical Sciences, New York University, New York, NY, United States

JOAQUIM SERRA: joaquim.serra@math.ethz.ch

Department of Mathematics, ETH Zürich, Zürich, Switzerland

Analysis & PDE

msp.org/apde

EDITORS

EDITOR-IN-CHIEF

Patrick Gérard

patrick.gerard@math.u-psud.fr

Université Paris Sud XI

Orsay, France

BOARD OF EDITORS

Massimiliano Berti	Scuola Intern. Sup. di Studi Avanzati, Italy berti@sissa.it	Clément Mouhot	Cambridge University, UK c.mouhot@dpmms.cam.ac.uk
Sun-Yung Alice Chang	Princeton University, USA chang@math.princeton.edu	Werner Müller	Universität Bonn, Germany mueller@math.uni-bonn.de
Michael Christ	University of California, Berkeley, USA mchrist@math.berkeley.edu	Gilles Pisier	Texas A&M University, and Paris 6 pisier@math.tamu.edu
Alessio Figalli	ETH Zurich, Switzerland alessio.figalli@math.ethz.ch	Tristan Rivière	ETH, Switzerland riviere@math.ethz.ch
Charles Fefferman	Princeton University, USA cf@math.princeton.edu	Igor Rodnianski	Princeton University, USA irod@math.princeton.edu
Ursula Hamenstaedt	Universität Bonn, Germany ursula@math.uni-bonn.de	Sylvia Serfaty	New York University, USA serfaty@cims.nyu.edu
Vaughan Jones	U.C. Berkeley & Vanderbilt University vaughan.f.jones@vanderbilt.edu	Yum-Tong Siu	Harvard University, USA siu@math.harvard.edu
Vadim Kaloshin	University of Maryland, USA vadim.kaloshin@gmail.com	Terence Tao	University of California, Los Angeles, USA tao@math.ucla.edu
Herbert Koch	Universität Bonn, Germany koch@math.uni-bonn.de	Michael E. Taylor	Univ. of North Carolina, Chapel Hill, USA met@math.unc.edu
Izabella Laba	University of British Columbia, Canada ilaba@math.ubc.ca	Gunther Uhlmann	University of Washington, USA gunther@math.washington.edu
Gilles Lebeau	Université de Nice Sophia Antipolis, France lebeau@unice.fr	András Vasy	Stanford University, USA andras@math.stanford.edu
Richard B. Melrose	Massachusetts Inst. of Tech., USA rbb@math.mit.edu	Dan Virgil Voiculescu	University of California, Berkeley, USA dvv@math.berkeley.edu
Frank Merle	Université de Cergy-Pontoise, France Frank.Merle@u-cergy.fr	Steven Zelditch	Northwestern University, USA zelditch@math.northwestern.edu
William Minicozzi II	Johns Hopkins University, USA minicozz@math.jhu.edu	Maciej Zworski	University of California, Berkeley, USA zvorski@math.berkeley.edu

PRODUCTION

production@msp.org

Silvio Levy, Scientific Editor


See inside back cover or msp.org/apde for submission instructions.

The subscription price for 2018 is US \$275/year for the electronic version, and \$480/year (+\$55, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues from the last three years and changes of subscriber address should be sent to MSP.

Analysis & PDE (ISSN 1948-206X electronic, 2157-5045 printed) at Mathematical Sciences Publishers, 798 Evans Hall #3840, c/o University of California, Berkeley, CA 94720-3840, is published continuously online. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices.

APDE peer review and production are managed by EditFlow[®] from MSP.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2018 Mathematical Sciences Publishers

ANALYSIS & PDE

Volume 11 No. 7 2018

On the global stability of a beta-plane equation FABIO PUSATERI and KLAUS WIDMAYER	1587
Airy-type evolution equations on star graphs DELIO MUGNOLO, DIEGO NOJA and CHRISTIAN SEIFERT	1625
On s -harmonic functions on cones SUSANNA TERRACINI, GIORGIO TORTONE and STEFANO VITA	1653
Weighted little bmo and two-weight inequalities for Journé commutators IRINA HOLMES, STEFANIE PETERMICHL and BRETT D. WICK	1693
Estimates for eigenvalues of Aharonov–Bohm operators with varying poles and non-half-integer circulation LAURA ABATANGELO, VERONICA FELLI, BENEDETTA NORIS and MANON NYS	1741
Complex rotation numbers: bubbles and their intersections NATALIYA GONCHARUK	1787
Quantitative stability of the free boundary in the obstacle problem SYLVIA SERFATY and JOAQUIM SERRA	1803