

ANALYSIS & PDE

Volume 12 No. 2 2019

ORAN GANNOT

**RESOLVENT ESTIMATES FOR SPACETIMES
BOUNDED BY KILLING HORIZONS**

RESOLVENT ESTIMATES FOR SPACETIMES BOUNDED BY KILLING HORIZONS

ORAN GANNOT

We show that the resolvent grows at most exponentially with frequency for the wave equation on a class of stationary spacetimes which are bounded by nondegenerate Killing horizons, without any assumptions on the trapped set. Correspondingly, there exists an exponentially small resonance-free region, and solutions of the Cauchy problem exhibit logarithmic energy decay.

1. Introduction

1A. Statement of results. Let (M, g) be a connected $(n+1)$ -dimensional Lorentzian manifold of signature $(1, n)$ with connected boundary ∂M , satisfying the following assumptions:

- (1) ∂M is a Killing horizon generated by a complete Killing vector field T , whose surface gravity is a positive constant $\kappa > 0$ (see Section 2C for details).
- (2) M is stationary in the sense that there is a compact spacelike hypersurface X with boundary such that each integral curve of T intersects X exactly once.
- (3) T is timelike in M° .

Consider a formally self-adjoint (with respect to the volume density) operator $L \in \text{Diff}^2(M)$ commuting with T , such that $L - \square_g \in \text{Diff}^1(M)$. Thus we can write

$$L = \square_g + \mathcal{W} + \mathcal{V},$$

where \mathcal{W} is a smooth vector field and $\mathcal{V} \in \mathcal{C}^\infty(M)$. In addition, assume that \mathcal{W} is tangent to ∂M .

Identify $M = \mathbb{R}_t \times X$ under the flow of T . Since T commutes with L , the composition

$$\mathbf{P}(\omega) = e^{i\omega t} L e^{-i\omega t} \tag{1-1}$$

descends to a differential operator on X depending on $\omega \in \mathbb{C}$. Fredholm properties of $\mathbf{P}(\omega)$ were first examined in a robust fashion by Vasy [2013] using methods of microlocal analysis, and subsequently by Warnick [2015] via physical space arguments; see also [Gannot 2018].

Here we summarize a simple version of these results, which applies in any strip of fixed width near the real axis. For $k \in \mathbb{N}$, let

$$\mathcal{X}^k = \{u \in H^{k+1}(X) : \mathbf{P}(0)u \in H^k(X)\}, \tag{1-2}$$

MSC2010: 35L05, 35P25, 35R01, 83C57.

Keywords: logarithmic decay, resolvent bounds, Killing horizons, Carleman estimates.

equipped with the graph norm. Since $\mathbf{P}(\omega) - \mathbf{P}(0) \in \text{Diff}^1(X)$, the operator $\mathbf{P}(\omega)$ is bounded $\mathcal{X}^k \rightarrow H^k(X)$ for each $\omega \in \mathbb{C}$.

Proposition 1.1 [Vasy 2013; Warnick 2015]. *The operator $\mathbf{P}(\omega) : \mathcal{X}^k \rightarrow H^k(X)$ is Fredholm of index 0 in the half-plane $\{\text{Im } \omega > -\kappa(k + \frac{1}{2})\}$ and is invertible for $\text{Im } \omega > 0$ sufficiently large.*

The inverse $\mathbf{P}(\omega)^{-1} : H^k(X) \rightarrow \mathcal{X}^k$ forms a meromorphic family of operators in $\{\text{Im } \omega > -\kappa(k + \frac{1}{2})\}$, called the resolvent family, which is independent of k in a suitable sense [Vasy 2013, Remark 2.9]. Its complex poles in $\{\text{Im } \omega > -\kappa(k + \frac{1}{2})\}$ are known as resonances, and correspond to nontrivial mode solutions $v = e^{-i\omega t}u$ of the equation $\square_g v = 0$, where $u \in C^\infty(M)$ satisfies $Tu = 0$. Thus mode solutions with $\text{Im } \omega > 0$ grow exponentially in time, whereas those with $\text{Im } \omega < 0$ exhibit exponential decay.

Given $\omega_0, C_0 > 0$, define the region

$$\Omega = \{|\text{Im } \omega| \leq e^{-C_0|\text{Re } \omega|}\} \cap \{|\omega| > \omega_0\}.$$

These parameters are fixed in the next theorem, which is the main result of this paper.

Theorem 1. *There exist $\omega_0, C_0 > 0$ such that $\mathbf{P}(\omega)$ has no resonances in Ω . Furthermore, there exists $C > 0$ such that if $\omega \in \Omega$, then*

$$\|\mathbf{P}(\omega)^{-1} f\|_{H^{k+1}} \leq e^{C|\text{Re } \omega|} \|f\|_{H^k} \quad (1-3)$$

for each $k \in \mathbb{N}$ and $f \in H^k(X)$.

Theorem 1 is also true when ∂M consists of several Killing horizons generated by T , each of which has a positive, constant surface gravity. In particular, Theorem 1 applies to any stationary perturbation of the Schwarzschild–de Sitter spacetime (which is bounded by two nondegenerate Killing horizons [Vasy 2013, Section 6]) that preserves the timelike nature of T , and for which the horizons remain nondegenerate Killing horizons. Other examples are even asymptotically hyperbolic spaces in the sense of [Guillarmou 2005].

1B. Energy decay. Theorem 1 can be used to prove logarithmic decay to constants for solutions of the Cauchy problem

$$\square_g v = 0, \quad v|_X = v_0, \quad T v|_X = v_1. \quad (1-4)$$

Given initial data $(v_0, v_1) \in H^{k+1}(X) \times H^k(X)$, the equation (1-4) admits a unique solution

$$v \in \mathcal{C}^0(\mathbb{R}_+; H^{k+1}(X)) \cap \mathcal{C}^1(\mathbb{R}_+; H^k(X)).$$

If N denotes the future-pointing unit normal to the level sets of t and $Q[v]$ is the stress-energy tensor (see Section 4C) associated to v , define the energy

$$\mathcal{E}[v](s) = \int_{\{t=s\}} Q[v](N, N) dS_X.$$

Here dS_X is the induced volume density on $X = \{t = 0\}$, which is isometric to each time slice $\{t = s\}$. Since N is timelike, it is well known that $\mathcal{E}[v](s)$ is positive definite in dv . One consequence of the

positivity of κ is the energy-boundedness statement

$$\mathcal{E}[v](t) \leq C\mathcal{E}[v](0); \quad (1-5)$$

see for instance [Warnick 2015, Corollary 3.9]. One can also define an energy $\mathcal{E}_k[v]$ controlling all derivatives up to order k , with $\mathcal{E}[v] = \mathcal{E}_1[v]$, which is similarly uniformly bounded. This can be improved to a logarithmic energy-decay statement uniformly up to the horizon, with a loss of derivatives.

Corollary 1.2. *Given $k \in \mathbb{N}$, there exists $C > 0$ such that*

$$\mathcal{E}_k[v](t)^{1/2} \leq \frac{C}{\log(2+t)} \|(v_0, v_1)\|_{\mathcal{X}^k \times H^{k+1}}$$

for each $v \in \mathcal{C}^0(\mathbb{R}_+; H^{k+1}(X)) \cap \mathcal{C}^1(\mathbb{R}_+; H^k(X))$ solving the Cauchy problem (1-4) with initial data $(v_0, v_1) \in \mathcal{X}^k \times H^{k+1}(X)$.

We can also improve Corollary 1.2 by showing that v decays logarithmically to a constant as follows. Given $(v_0, v_1) \in \mathcal{X}^k \times H^{k+1}(X)$, define the constant

$$v_\infty = \text{vol}(\partial X)^{-1} \int_X (A^{-2}v_1 - 2A^{-2}Wv_0 - \text{div}_g(A^{-2}W)v_0) A dS_X.$$

Here $A > 0$ is the lapse function and W is the shift vector as described in Section 2D.

Corollary 1.3. *Given $k \in \mathbb{N}$, there exists $C > 0$ such that*

$$\|v(t) - v_\infty\|_{H^{k+1}} + \|\partial_t v(t)\|_{H^k} \leq \frac{C}{\log(2+t)} \|(v_0, v_1)\|_{\mathcal{X}^k \times H^{k+1}}$$

for each $v \in \mathcal{C}^0(\mathbb{R}_+; H^{k+1}(X)) \cap \mathcal{C}^1(\mathbb{R}_+; H^k(X))$ solving the Cauchy problem (1-4) with initial data $(v_0, v_1) \in \mathcal{X}^k \times H^{k+1}(X)$.

By Sobolev embedding, Corollary 1.3 can be used to deduce pointwise decay estimates as well.

1C. Relationship with previous work. The analogue of Theorem 1 was first established for compactly supported perturbations of the Euclidean Laplacian in the landmark paper [Burq 1998]. There have been subsequent improvements and simplifications in the asymptotically Euclidean setting [Burq 2002; Vodev 2000; Datchev 2014], while Rodnianski and Tao [2015] considered asymptotically conic spaces. In a different direction, Holzegel and Smulevici [2013] established logarithmic energy decay on slowly rotating Kerr–AdS spacetimes, which contain a Killing horizon of the type described here in addition to a conformally timelike boundary. However, their approach made heavy use of the symmetries of Kerr–AdS, and is not adaptable to our setting.

Most relevant to the setting considered here are [Moschidis 2016; Cardoso and Vodev 2002]. The former reference shows logarithmic energy decay on Lorentzian spacetimes which may contain Killing horizons, but importantly also contain at least one asymptotically flat end. There, the mechanism of decay is radiation into the asymptotically flat region. In contrast, asymptotically flat ends are not considered in the present paper, but we do allow spacetimes which contain Killing horizons as their only boundary components. We therefore stress that the results of [Moschidis 2016] are disjoint from those of this paper.

Meanwhile, [Cardoso and Vodev 2002] applies to a wide class of Riemannian metrics, including those with hyperbolic ends. There is a close connection between asymptotically hyperbolic manifolds and black holes spacetimes, first exploited in the study of resonances by Sá Barreto and Zworski [1997]. This relationship has attracted a great deal of interest, especially following the paper [Vasy 2013]; for a survey of recent developments, see [Zworski 2017].

Common to the works described above is the use of Carleman estimates in the interior of the geometry, which is then combined with some other (typically more complicated) analysis near infinity. Although the proof of Theorem 1 adopts techniques from [Burq 1998; Moschidis 2016; Rodnianski and Tao 2015], one novelty (and simplifying feature) is that the Carleman estimate employed here is valid up to and including the horizon. In particular, this avoids the use of separation of variables and special function methods [Burq 1998; Holzegel and Smulevici 2013; Vodev 2000], Mourre-type estimates [Burq 2002], and spherical energies [Cardoso and Vodev 2002; Datchev 2014; Moschidis 2016; Rodnianski and Tao 2015].

2. Preliminaries

2A. Semiclassical rescaling. It is conceptually convenient to rescale the operator by

$$P(z) = h^2 \mathbf{P}(h^{-1}z). \quad (2-1)$$

Thus $\omega = h^{-1}z$, and uniform bounds on $P(z)$ for $\pm z$ in a compact set $[a, b] \subset (0, \infty)$ give high-frequency bounds for $\mathbf{P}(\omega)$ as $|\omega| \rightarrow \infty$. Theorem 1 is easily seen to be equivalent to the following.

Theorem 1'. *Given $[a, b] \subset (0, \infty)$, there exist $C, C_1 > 0$ such that*

$$\|u\|_{H_h^{k+1}} \leq e^{C/h} \|P(z)u\|_{H_h^k} \quad (2-2)$$

for each $u \in \mathcal{X}^k$ and $\pm z \in [a, b] + ie^{-C_1/h}[-1, 1]$.

The norms in (2-2) are semiclassically rescaled Sobolev norms. For detailed expositions on semiclassical analysis, the reader is referred to [Zworski 2012] and [Dyatlov and Zworski 2018, Appendix E].

2B. Stationarity. A tensor on M will be called stationary if it is annihilated by the Lie derivative \mathcal{L}_T . The definition of stationarity can be extended to T^*M by observing that T lifts to a vector field on T^*M via the identification

$$T^*M = T^*\mathbb{R} \oplus T^*X.$$

Any covector $\varpi \in T_q^*M$ at a point $q = (t, x)$ can be decomposed as $\varpi = \xi + \tau dt$, where $\xi \in T_x^*X$ and $\tau dt \in T_t^*\mathbb{R}$. Thus a function $F \in \mathcal{C}^\infty(T^*M)$ is stationary if it depends only on $\xi \in T_x^*X$ and $\tau \in \mathbb{R}$, which we sometimes denote by $F(x, \xi, \tau)$. Furthermore, if $\tau = \tau_0$ is fixed, then F induces a function $F(\cdot, \tau_0)$ on T^*X . This is compatible with the Poisson bracket in the sense that for stationary $F_1, F_2 \in \mathcal{C}^\infty(T^*M)$, there is the equality

$$\{F_1, F_2\}(x, \xi, \tau_0) = \{F_1(\cdot, \tau_0), F_2(\cdot, \tau_0)\}(x, \xi). \quad (2-3)$$

On the left is the Poisson bracket on T^*M , and on the right the Poisson bracket on T^*X .

In particular, this discussion applies to the dual metric function $G \in \mathcal{C}^\infty(T^*M)$, whose value at $\varpi \in T_q^*M$ is given by

$$G(x, \varpi) = g_x^{-1}(\varpi, \varpi) = g^{\alpha\beta}(x) \varpi_\alpha \varpi_\beta.$$

The semiclassical principal symbol $p = \sigma_h(P(z))$ is given by $p(x, \xi; z) = -G(x, \xi - z \, dt)$.

Lemma 2.1. *The quadratic form $(x, \xi) \mapsto G(x, \xi)$ is negative definite on T^*X° .*

Proof. The condition $\tau = 0$ implies that $\varpi = \xi + 0 \, dt$ is orthogonal to T^\flat . But T^\flat is timelike on M° , whence the result follows. \square

If $\tau_0 \in \mathbb{R}$ is fixed and $K \subset X^\circ$ is compact, then by Lemma 2.1 there exist $c, R > 0$ such that if $G(x, \xi) \geq R$, then

$$G(x, \xi + \tau_0 \, dt) \geq cG(x, \xi)$$

for each $\xi \in T_K^*X^\circ$, where the constants c, R are locally uniform in τ_0 . In particular, given a compact interval $I \subset \mathbb{R}$, the set

$$\{\xi \in T_K^*X^\circ : G(\xi + \tau \, dt) = 0 \text{ for some } \tau \in I\}$$

is a compact subset of T^*X° . This also implies that if Q is a stationary quadratic form on T^*M , then there exists $C > 0$ such that

$$|Q(x, \xi + \tau \, dt)| \leq C(1 + |G(x, \xi + \tau \, dt)|)$$

for each $\xi \in T_K^*X^\circ$ and $\tau \in I$.

2C. Killing horizons and surface gravity. Recall the hypotheses on (M, g) described in Section 1A and set

$$\mu = g(T, T).$$

The key property of (M, g) is that ∂M is a Killing horizon generated by T . By definition, this means that ∂M is a null hypersurface which agrees with a connected component of the set $\{\mu = 0, T \neq 0\}$. Of course in this case T is nowhere-vanishing. Since orthogonal null vectors are collinear, there is a smooth function $\kappa : \partial M \rightarrow \mathbb{R}$, called the surface gravity, such that

$$\nabla_g \mu = -2\kappa T \tag{2-4}$$

on ∂M . The nondegeneracy assumption means that $\kappa > 0$, and for simplicity it is assumed that κ is in fact constant along ∂M .

2D. Properties of the metric. Let N denote the future-pointing unit normal to the level sets of t , and define the lapse function $A > 0$ by $A^{-2} = g^{-1}(dt, dt)$. The shift vector is given by the formula

$$W = T - AN,$$

which by construction is tangent to the level sets of t . Let k denote the induced (positive definite) metric on X . If (x^i) are local coordinates on X , then

$$g = (A^2 - k_{ij} W^i W^j) dt^2 - 2k_{ij} W^i dx^j dt - k_{ij} dx^i dx^j.$$

Inverting this form of the metric gives

$$g^{-1} = A^{-2}(\partial_t - W^i \partial_i)^2 - k^{ij} \partial_i \partial_j. \quad (2-5)$$

Note that $k(W, W) = A^2 - \mu$, and hence $W \neq 0$ near ∂M .

Now use the condition that ∂M is a Killing horizon generated by T . The covariant form of (2-4) reads

$$\partial_i \mu = 2\kappa W^j k_{ij}. \quad (2-6)$$

By assumption $\kappa > 0$, so W is a nonzero inward-pointing normal to X along ∂X whose length with respect to k is A .

Introduce geodesic normal coordinates (r, y^A) on X near ∂X , so r is the distance to ∂X (uppercase indices will always range over $A = 2, \dots, n$). By construction, ∂_r is an inward-pointing unit normal along ∂X , so

$$W^r = A, \quad W^A = 0 \quad (2-7)$$

along the boundary. Also by construction, the components of the induced metric in (r, y^A) -coordinates satisfy $k^{rr} = 1$ and $k^{rA} = 0$.

Lemma 2.2. *The function r satisfies $g^{-1}(dr, dr) = -2\kappa A^{-1}r + r^2 \mathcal{C}^\infty(M)$.*

Proof. First observe that $k_{AB} W^A W^B \in r^2 \mathcal{C}^\infty(M)$ by (2-7), and since $k(W, W) = A^2 - \mu$,

$$A^2 - \mu = (W^r)^2 + k_{AB} W^A W^B.$$

Now μ and r are both boundary-defining functions, so $\mu = fr$ for some $f \in C^\infty(M)$, and hence $d\mu = f dr$ on ∂X . But on the boundary $\langle d\mu \rangle = 2\kappa A^2$ from (2-6), while $\langle W, dr \rangle = W^r = A$ from (2-7). Thus

$$\mu = fr = 2\kappa Ar + r^2 \mathcal{C}^\infty(M).$$

Plugging this back into the equation for $k(W, W)$ yields

$$(W^r)^2 = A^2 - 2\kappa Ar + r^2 \mathcal{C}^\infty(M),$$

and therefore $g^{-1}(dr, dr) = -k^{rr} + A^{-2}(W^r)^2 = -2\kappa A^{-1}r + r^2 \mathcal{C}^\infty(M)$ as desired. \square

Observe that the surface gravity depends on the choice of null generator T . Consider the rescaled vector field

$$\hat{T} = \frac{T}{2\kappa},$$

which changes the time coordinate by the transformation $\hat{t} = 2\kappa t$. If $\hat{P}(\hat{\omega})$ is now defined as in (1-1) but with t replacing \hat{t} , then

$$P(\omega) = \hat{P}\left(\frac{\omega}{2\kappa}\right).$$

It suffices to prove Theorem 1 for $\hat{P}(\omega)$ then, since rescaling the frequency only changes the constants ω_0, C_0, C . Dropping the hat notation, it will henceforth be assumed that $\kappa = \frac{1}{2}$.

Next, consider a conformal change $g = f\tilde{g}$, where $f > 0$ is stationary. The operator L can then be written as

$$L = f^{-1}\square_{\tilde{g}} + (n-1)f^{-2}\nabla_{\tilde{g}}f + \mathcal{W} + \mathcal{V}. \quad (2-8)$$

Thus we can write $L = f^{-1}\tilde{L}$, where \tilde{L} has the same form as L but with \tilde{g} replacing g , provided that the vector field $\nabla_{\tilde{g}}f$ is tangent to ∂M . But this follows from the stationarity of f , since

$$g(T, \nabla_g f) = 0$$

and T is normal to ∂M . Thus it suffices to prove Theorem 1 with \tilde{L} replacing L . Observe that ∂M remains a Killing horizon generated by T with respect to \tilde{g} , and the surface gravity is unchanged.

By making a conformal change and dropping the tilde notation, it will also be assumed that

$$g^{-1}(dr, dr) = -r. \quad (2-9)$$

If (τ, ρ, η_A) are dual variables to (t, r, y^A) , define a stationary quadratic form $G_0 \in \mathcal{C}^\infty(T^*M)$ by

$$G_0 = -r\rho^2 - 2\rho\tau - k_0^{AB}\eta_A\eta_B. \quad (2-10)$$

Here k_0 is the restriction of k to ∂M , which is then extended to a neighborhood of ∂M by requiring that $\mathcal{L}_{\partial_r}k_0 = 0$. In the next section, the difference $G - G_0$ will be analyzed.

2E. Negligible tensors. We now define a class of tensors which will arise as errors throughout the proof of Theorem 1'.

Definition 2.3. (1) A stationary 1-tensor $F^\alpha \partial_\alpha$ is said to be negligible if its components in a coordinate system (t, r, y^A) satisfy

$$F^t \in r\mathcal{C}^\infty(M), \quad F^r \in r^2\mathcal{C}^\infty(M), \quad F^A \in r\mathcal{C}^\infty(M).$$

(2) A stationary 2-tensor $H^{\alpha\beta} \partial_\alpha \partial_\beta$ is said to be negligible if its components in a coordinate system (t, r, y^A) satisfy

$$\begin{aligned} H^{tt} &\in \mathcal{C}^\infty(M), & H^{rr} &\in r^2\mathcal{C}^\infty(M), & H^{AB} &\in r\mathcal{C}^\infty(M), \\ H^{tA} &\in \mathcal{C}^\infty(M), & H^{tr} &\in r\mathcal{C}^\infty(M), & H^{rA} &\in r\mathcal{C}^\infty(M). \end{aligned}$$

Observe that negligibility is invariant under those coordinate changes which leave (t, r) invariant. Denote by \mathcal{N}_1 and \mathcal{N}_2 all $\mathcal{C}^\infty(T^*M)$ functions of the forms $F^\alpha \varpi_\alpha$ and $H^{\alpha\beta} \varpi_\alpha \varpi_\beta$, respectively.

Recall the definition of G_0 in (2-10). The notion of negligibility is motivated by the fact that

$$G = G_0 + \mathcal{N}_2.$$

This follows directly from (2-5), (2-7), and (2-9). We will also repeatedly reference the auxiliary functions

$$Y = (r\rho)^2 + \tau^2, \quad Z = r\rho^2 + k^{AB}\eta_A\eta_B. \quad (2-11)$$

It follows immediately from the Cauchy–Schwarz inequality $2ab < \delta a^2 + b^2/\delta$ that there exists $C > 0$ satisfying

$$Z \leq C \left(|G_0| + \frac{\tau^2}{r} \right). \quad (2-12)$$

The next two lemmas also follow from judicious applications of the Cauchy–Schwarz inequality and the trivial observation that $(r\rho)^2 = r(r\rho^2)$ is small relative to $r\rho^2$ for small values of r .

Lemma 2.4. *Let $F \in \mathcal{N}_1$. Then, for each $\gamma > 0$ there exists C_γ such that*

$$r^{-1}|\tau||F| \leq C_\gamma \tau^2 + \gamma Z.$$

Furthermore, $\rho\mathcal{N}_1 \subset \mathcal{N}_2$ and $\mathcal{N}_1 \cdot \mathcal{N}_1 \subset r\mathcal{N}_2$.

Lemma 2.5. *Let $H \in \mathcal{N}_2$. Then, for each $\gamma > 0$ there exist $C_\gamma, r_\gamma > 0$ such that*

$$|H| \leq C_\gamma Y + \gamma k^{AB} \eta_A \eta_B, \quad |H| \leq C_\gamma \tau^2 + \gamma Z$$

for $r \in [0, r_\gamma]$.

Now combine Lemma 2.5 with the bound (2-12) and the relation $G = G_0 + \mathcal{N}_2$. Thus there exists $R > 0$ and $C > 0$ such that

$$Z \leq C \left(|G| + \frac{\tau^2}{r} \right) \quad (2-13)$$

for $r \in [0, R]$.

The next goal is to compute the Poisson brackets $\{G, r\}$ and $\{G, \{G, r\}\}$. To begin, observe that

$$\{G_0, r\} = -2(r\rho + \tau), \quad \{G_0, \{G_0, r\}\} = 2(r\rho^2 + 2\tau\rho). \quad (2-14)$$

In order to replace G_0 with G we also need to consider the Poisson brackets of functions in \mathcal{N}_1 and \mathcal{N}_2 .

Lemma 2.6. *The Poisson bracket satisfies $\{\mathcal{N}_2, r\} \subset \mathcal{N}_1$ and $\{\mathcal{N}_2, \mathcal{N}_1\} \subset \mathcal{N}_2$, as well as $\{G_0, \mathcal{N}_1\} \subset \mathcal{N}_2$ and $\{\{G_0, r\}, \mathcal{N}_2\} \subset \mathcal{N}_2$. Therefore,*

$$\{G, r\} = -2(r\rho + \tau) + \mathcal{N}_1, \quad \{G, \{G, r\}\} = 2(r\rho^2 + 2\tau\rho) + \mathcal{N}_2. \quad (2-15)$$

Furthermore, $\{G, \{G, r\}\} = -2r\rho^2 + \mathcal{N}_2$ whenever $\{G, r\} = 0$.

Proof. The first part is a direct calculation, while (2-15) follows from the first part and (2-14). The last statement follows from the inclusion $\rho\mathcal{N}_1 \subset \mathcal{N}_2$. \square

3. Carleman estimates in the interior

3A. Statement of result. We now prove a Carleman estimate valid in the interior X° , but with uniform control over the exponential weight near ∂X .

Recall that r denotes the distance on X to the boundary with respect to the induced metric. Although this function is only well-defined in a small neighborhood of ∂X , for notational convenience we will assume that $[0, 3]$ is contained in the range of r (otherwise it is just a matter of replacing 3 with 3ε for an appropriate $\varepsilon > 0$).

Proposition 3.1. *Given $[a, b] \subset (0, \infty)$, there exists $r_1 \in (0, 1)$ and $\varphi_1, \varphi_2 \in \mathcal{C}^\infty(X)$ such that*

- *on $\{r \leq 1\}$ the functions φ_1, φ_2 are equal and depend only on r ,*
- *$\varphi'_i(r) < 0$ is constant on $\{r \leq r_1\}$ for $i = 1, 2$,*

with the following property: given a compact set $K \subset X^\circ$ there exists $C > 0$ such that

$$\|(e^{\varphi_1/h} + e^{\varphi_2/h})u\|_{H_h^2(X)} \leq Ch^{-1/2} \|(e^{\varphi_1/h} + e^{\varphi_2/h})P(z)u\|_{L^2(X)}$$

for each $u \in \mathcal{C}_c^\infty(K^\circ)$ and $\pm z \in [a, b]$.

It clearly suffices to prove Proposition 3.1 for the operator $L = \square_g$, since the lower-order terms can be absorbed as errors. In order to prove Theorem 1', an additional estimate is needed near the boundary; this is achieved in Section 4 below.

3B. The conjugated operator. Given $\varphi \in \mathcal{C}^\infty(X)$, define the conjugated operator

$$P_\varphi(z) = e^{\varphi/h} P(z) e^{-\varphi/h}.$$

Let $p_\varphi(z)$ denote its semiclassical principal symbol. Define $L^2(X)$ with respect to the density $A \cdot dS_X$, where recall dS_X is the induced volume density on X , and $A > 0$ is the lapse function as in Section 2C. Defining $\operatorname{Re} P_\varphi(z)$ and $\operatorname{Im} P_\varphi(z)$ with respect to this inner product, integrate by parts to find

$$\|P_\varphi(\omega)u\|_{L^2(X)}^2 = \langle P_\varphi(\omega)P_\varphi(\omega)^*u, u \rangle_{L^2(X)} + i \langle [\operatorname{Re} P_\varphi(\omega), \operatorname{Im} P_\varphi(\omega)]u, u \rangle_{L^2(X)} \quad (3-1)$$

for $u \in \mathcal{C}_c^\infty(X^\circ)$. The idea is to find φ which satisfies Hörmander's hypoellipticity condition

$$\{\operatorname{Re} p_\varphi, \operatorname{Im} p_\varphi\} > 0 \quad (3-2)$$

on the characteristic set $\{p_\varphi = 0\}$.

In order to apply the results of Section 2E without introducing additional notation, it is convenient to work with the dual metric function G directly. Define

$$G_\varphi(x, \varpi) = G(x, \varpi + i d\varphi),$$

so since we are assuming that τ is real, $\operatorname{Re} G_\varphi(x, \varpi) = G(x, \varpi) - G(x, d\varphi)$, and $\operatorname{Im} G_\varphi(x, \varpi) = (H_G \varphi)(x, \varpi)$. We will then construct φ (viewed as a stationary function on M) such that

$$\{\operatorname{Re} G_\varphi, \operatorname{Im} G_\varphi\}(x, \varpi) = (H_G^2 \varphi)(x, \varpi) + (H_G^2 \varphi)(x, d\varphi) > 0 \quad (3-3)$$

on $\{G_\varphi = 0\} \cap \{a \leq \pm \tau \leq b\}$. This will imply the original hypoellipticity condition from the discussion surrounding (2-3) and the identifications

$$p_\varphi(x, \xi; z) = -G_\varphi(x, \xi - z dt), \quad z = -\tau.$$

Note that the dual variable τ is now playing the role of a *rescaled* time frequency.

3C. Constructing the phase in a compact set. To avoid any undue topological restrictions, we will actually construct two weights φ_1, φ_2 in the interior which agree outside a large compact set. This appears already in [Burq 1998], but we will follow the closely related presentation in [Moschidis 2016; Rodnianski and Tao 2015].

Lemma 3.2. *There exist positive functions $\psi_1, \psi_2 \in C^\infty(X)$ with the following properties:*

- (1) ψ_1, ψ_2 have finitely many nondegenerate critical points, all of which are contained in $\{r > 2\}$.
- (2) $\psi_2 > \psi_1$ on $\{d\psi_1 = 0\}$, and $\psi_1 > \psi_2$ on $\{d\psi_2 = 0\}$.
- (3) The functions ψ_1, ψ_2 are equal and depend only on r in $\{r \leq 2\}$. Furthermore $\partial_r \psi_1$ and $\partial_r \psi_2$ are negative in this region.

Proof. Let $\zeta \in C^\infty(\{r \geq 2\})$ solve the boundary value problem

$$\Delta_k \zeta = 1, \quad \zeta|_{\{r=2\}} = 1.$$

Here Δ_k is the nonpositive Laplacian with respect to the induced metric k . Since $\Delta_k \zeta > 0$, none of the critical points of ζ in $\{r > 2\}$ are local maxima. In addition, since ζ clearly achieves its maximum at each point of $\{r = 2\}$, its outward-pointing normal derivative is strictly positive by Hopf's lemma [Gilbarg and Trudinger 1983, Lemma 3.4]. By construction, the outward-pointing unit normal is $-\partial_r$; hence $\zeta' < 0$ near $\{r = 2\}$ (for the remainder of the proof, prime will denote differentiation with respect to r).

The first step is to replace ζ by a Morse function. We may for instance embed $\{r \geq 2\}$ into a compact manifold X_0 without boundary, and approximate an arbitrary smooth extension of ζ to X_0 by a Morse function in the $C^\infty(X_0)$ topology. Restricting to $\{r \geq 2\}$ and again calling this replacement ζ , we still have that ζ has no local maximum in $\{r > 2\}$ and $\zeta' < 0$ near $\{r = 2\}$. In particular, all critical points of ζ are nondegenerate and lie in a compact subset of $\{r > 2\}$.

Now fix any function $\bar{\zeta} = \bar{\zeta}(r) \in C^\infty(\{r < 3\})$ such that $\bar{\zeta}' < 0$ everywhere, and $\bar{\zeta} \geq \zeta$ on their common domain of definition $\{2 \leq r < 3\}$. Choose a cutoff $H = H(r) \in C^\infty(X; [0, 1])$ such that

$$H = 1 \quad \text{for } r < 2 + \gamma, \quad \text{supp } H \subset \{r \leq 2 + 2\gamma\},$$

and $H' \leq 0$. Set $\psi_1 = H\bar{\zeta} + (1 - H)\zeta$, and compute $\psi_1' = H'(\bar{\zeta} - \zeta) + H\bar{\zeta}' + (1 - H)\zeta'$. If $\gamma > 0$ is sufficiently small, then $\psi_1' < 0$ in a neighborhood of $\text{supp } H$, since the sum of the last two terms is strictly positive on $\text{supp } H$. On the other hand, outside of such a neighborhood the only critical points of ψ_1 are those of ζ .

Let p_1, \dots, p_n enumerate the necessarily finite number of critical points of ψ_1 , and choose $\gamma > 0$ such that the closed geodesic balls $B(p_1, \gamma), \dots, B(p_n, \gamma)$ are mutually disjoint and $B(p_j, \gamma) \subset \{r > 2\}$ for each j . Since p_j is not a local maximum, for each j there is a point $q_j \in B(p_j, \gamma)$ such that

$$\psi_1(q_j) > \psi_1(p_j).$$

Now choose a diffeomorphism $g : X \rightarrow X$ which is the identity outside the union of the $B(q_j, r)$ and exchanges p_j with q_j . Then, set $\psi_2 = \psi_1 \circ g$. By construction the only critical points of ψ_2 are q_1, \dots, q_n ,

and furthermore

$$\psi_2(p_j) > \psi_1(p_j), \quad \psi_1(q_j) > \psi_2(q_j)$$

for each j . Since outside of $\{r > 2\}$ the functions $\psi_1 = \psi_2$ depend on r only, the proof is complete, adding an appropriate constant if necessary to ensure that both functions are positive. \square

Let $B_1 \subset \{r > 2\}$ be a closed neighborhood of $\{d\psi_1 = 0\}$ such that $\psi_2 > \psi_1$ on B_1 , and likewise for B_2 , exchanging the roles of ψ_1 and ψ_2 . Also, let $U_i \subset B_i$ be additional neighborhoods of $\{d\psi_i = 0\}$. Now define

$$\varphi_i = \exp(\alpha\psi_i), \quad i = 1, 2, \quad (3-4)$$

where $\alpha > 0$ is a parameter. The following lemma is a standard computation which is included for the sake of completeness.

Lemma 3.3. *Given $\varepsilon > 0$ and $\tau_0 > 0$, there exists $\alpha_0 > 0$ such that if $\alpha \geq \alpha_0$, then*

$$\{\operatorname{Re} G_{\varphi_i}, \operatorname{Im} G_{\varphi_i}\} > 0$$

on $(\{G_{\varphi_i} = 0\} \cap \{r \geq \varepsilon\} \cap \{|\tau| \leq \tau_0\}) \setminus T_{U_i}^* M$ for $i = 1, 2$.

Proof. The subscript $i = 1, 2$ will be suppressed. Use the definition (3-4) to compute

$$H_G \varphi = \alpha e^{\alpha\psi} H_G \psi, \quad H_G^2 \varphi = \alpha^2 e^{\alpha\psi} (H_G \psi)^2 + \alpha e^{\alpha\psi} H_G^2 \psi.$$

Assume that $G_\varphi(x, \varpi) = 0$. It follows from $\operatorname{Im} G_\varphi(x, \varpi) = 0$ that $(H_G \varphi)(x, \varpi) = 0$, and hence $(H_G \psi)(x, \varpi) = 0$. Therefore by (3-3),

$$\{G - G(x, d\psi), H_G \varphi\}(x, \varpi) = \alpha e^{\alpha\psi} (H_G^2 \psi)(x, \varpi) + \alpha^3 e^{3\alpha\psi} (H_G^2 \psi)(x, d\psi) + \alpha^4 e^{3\alpha\psi} |G(x, d\psi)|^2.$$

Next, use the condition $(\operatorname{Re} G_\varphi)(x, \varpi) = 0$, which implies $G(x, \varpi) = \alpha^2 e^{2\alpha\psi} G(x, d\psi)$. By the discussion following Lemma 2.1, there exists $C > 0$ such that

$$|(H_G^2 \psi)(x, \varpi)| \leq C(1 + |G(x, \varpi)|)$$

on $\{r \geq \varepsilon\} \cap \{|\tau| \leq \tau_0\}$. Thus on the set $\{G_\varphi = 0\} \cap \{r \geq \varepsilon\} \cap \{|\tau| \leq \tau_0\}$,

$$|\alpha e^{\alpha\psi} (H_G^2 \psi)(x, \varpi)| + |\alpha^3 e^{3\alpha\psi} (H_G^2 \psi)(x, d\psi)| \leq C \alpha^3 e^{3\alpha\psi}.$$

On the other hand, as soon as $d\psi \neq 0$ the third term $\alpha^4 e^{3\alpha\psi} |G(x, d\psi)|^2$ is positive by Lemma 2.1, and dominates the previous two terms for large $\alpha > 0$. Since $d\psi \neq 0$ away from B , the proof is complete. \square

3D. Constructing the phase outside of a compact set. The most delicate part of the argument is the construction of the phase outside of a compact set. Since $g^{-1}(dr, dr) = -r$ and φ is a function only of r in this region,

$$G_\varphi = G + r(\varphi')^2 + i\varphi' H_G r.$$

Now compute the Poisson bracket

$$\begin{aligned}\{\operatorname{Re} G_\varphi, \operatorname{Im} G_\varphi\} &= \{G + r(\varphi')^2, \varphi' H_G r\} \\ &= \varphi' H_G^2 r + \varphi'' (H_G r)^2 - ((\varphi')^3 + 2r(\varphi')^2 \varphi'') \partial_\rho H_G r.\end{aligned}$$

Assume that $\varphi' < 0$, in which case $\operatorname{Im} G_\varphi = 0$ is equivalent to $H_G r = 0$. The goal is then to arrange negativity of the term

$$H_G^2 r - ((\varphi')^2 + 2r\varphi'\varphi'') \partial_\rho H_G r \quad (3-5)$$

on the set $\{\operatorname{Re} G_\varphi = 0\}$. Recall the definition of Z from (2-11).

Lemma 3.4. *There exists $C > 0$ and $R > 0$ such that $Z \leq C(r(\phi')^2 + \tau^2/r)$ on $\{\operatorname{Re} G_\varphi = 0\} \cap \{0 < r \leq R\}$.*

Proof. Apply (2-13), using that $\operatorname{Re} G_\varphi = 0$ implies $G = -r(\varphi')^2$. \square

Putting everything together, it is now easy compute $H_G^2 r$ on $\{G_\varphi = 0\}$ near the boundary.

Lemma 3.5. *For each $\delta > 0$ there exists $R_\delta > 0$ such that*

$$\left| H_G^2 r + \frac{2\tau^2}{r} \right| \leq \delta \left(r(\phi')^2 + \frac{\tau^2}{r} \right)$$

on $\{G_\varphi = 0\} \cap \{0 < r \leq R_\delta\}$.

Proof. From the expression (2-15) for $H_G^2 r$ and Lemma 2.5, find $C_\gamma > 0$ and $r_\gamma > 0$ such that

$$|H_G^2 r + 2r\rho^2| < C_\gamma |\tau|^2 + \gamma Z \quad (3-6)$$

for $r \in (0, r_\gamma)$. Now multiply $H_G r$ by ρ , and use that $\rho \mathcal{N}_1 \subset \mathcal{N}_2$. Therefore by Lemma 2.5, there exist $C'_\gamma > 0$ and $r'_\gamma > 0$ such that

$$|2r\rho^2 + 2\tau\rho| < C'_\gamma |\tau|^2 + \gamma Z \quad (3-7)$$

for $r \in (0, r'_\gamma)$. On the other hand, from $H_G r = 0$, deduce that $-\tau\rho = \tau^2/r + \tau r^{-1} \mathcal{N}_1$. By Lemma 2.4, there exists $C''_\gamma > 0$ such that

$$\left| 2\tau\rho + \frac{2\tau^2}{r} \right| < C''_\gamma |\tau|^2 + \gamma Z. \quad (3-8)$$

Combine (3-6), (3-7), and (3-8) via the triangle inequality with Lemma 3.4 to find that

$$\left| H_G^2 r + \frac{2\tau^2}{r} \right| < 3\gamma C \left(r(\phi')^2 + \frac{\tau^2}{r} \right) + (C_\gamma + C'_\gamma + C''_\gamma) \tau^2$$

for $r \in (0, \min\{r_\gamma, r'_\gamma, R\})$; here $C > 0$ and $R > 0$ are provided by Lemma 3.4. Finally, choose γ sufficiently small depending on δ and a corresponding $R_\delta > 0$ such that the conclusion of the lemma holds for $r \in (0, R_\delta)$. \square

Next, observe that $-\partial_\rho H_G r = 2r + r^2 \mathcal{C}^\infty(M)$. Given $a > 0$, it follows from (3-5) and Lemma 3.5 that there exists $R_1 > 0$ such that

$$(\varphi')^{-1} \{\operatorname{Re} G_\varphi, \operatorname{Im} G_\varphi\} < -\frac{3a^2}{2r} + 3r(\varphi')^2 + 3r^2 \varphi' \varphi'' \quad (3-9)$$

on $\{G_\varphi = 0\} \cap \{0 < r \leq R_1\} \cap \{|\tau| \geq a\}$, provided that $\varphi'' \geq 0$.

Shrinking R_1 if necessary, it may be assumed that $\psi = \psi_i$ as in Lemma 3.2 satisfies $\psi' < 0$ on $[0, R_1 + 1]$. Recalling that $\varphi_i = \exp(\alpha\psi_i)$, choose $\alpha > 0$ satisfying the conclusion of Lemma 3.3 with $\varepsilon = R_1$. By further increasing α , (but keeping $a > 0$ fixed), it may also be assumed that $\varphi = \varphi_i$ satisfies

$$3(\varphi'(R_1)R_1)^2 > a^2, \quad \varphi''(r) \geq -\frac{\varphi'(r)}{r} \quad \text{for } r \in [R_1, R_1 + 1]. \quad (3-10)$$

Although φ is already defined on all of X , the following lemma allows one to redefine φ on $\{r < R_1 + 1\}$ in such a way that its derivative is controlled; this new extension will still be denoted by φ . The idea comes from [Burq 1998, Section 3.1.2], but of course the form of the operator there is quite different.

Lemma 3.6. *There exists an extension of $\varphi = \varphi_i$ from $\{r \geq R_1 + 1\}$ to $\{r < R_1 + 1\}$ such that*

$$\{\operatorname{Re} G_\varphi, \operatorname{Im} G_\varphi\} > 0$$

on $\{G_\varphi = 0\} \cap \{0 < r \leq R_1\} \cap \{|\tau| \geq a\}$. Furthermore, there exists $r_1 \in (0, R_1)$ such that $\varphi'(r) < 0$ is constant for $r \in [0, r_1]$.

Proof. Motivated by (3-9), consider the differential equation

$$-\frac{a^2}{r} + 3rk^2 + 3r^2kk' = 0, \quad k(R_1) = \varphi'(R_1) < 0.$$

This is a Bernoulli equation whose solution is given by

$$k(r) = -r^{-1} \left((\varphi'(R_1)R_1)^2 + \frac{2}{3}a^2 \log\left(\frac{r}{R_1}\right) \right)^{1/2}.$$

The solution is certainly meaningful for $r \in [R_0, R_1]$, where we define R_0 by

$$R_0 = R_1 \exp\left(\frac{1}{2} - \frac{3}{2}\left(\varphi'(R_1)\frac{R_1}{a}\right)^2\right).$$

Note that we indeed have $R_0 < R_1$ by the assumption (3-10). The value R_0 was chosen such that $k'(R_0) = 0$, and it is easy to see that $k'(r) > 0$ for $r \in (R_0, R_1]$. In addition, $k(R_0) < 0$. Let $\theta = \theta(r)$ be defined on $[0, R_1 + 1]$ by

$$\theta(r) = \begin{cases} \varphi'(r), & r \in [R_1, R_1 + 1], \\ k(r), & r \in [R_0, R_1], \\ k(R_0), & r \in [0, R_0]. \end{cases}$$

The function θ is strictly negative, and the piecewise continuous function θ' satisfies

$$-\frac{a^2}{r} + 3r\theta^2 + 3r^2\theta\theta' \leq 0$$

for $r \in (0, R_1 + 1]$. Indeed, by the construction of k and R_0 , the inequality holds for $r \in (0, R_1)$, and it is also true for $r \in (R_1, R_1 + 1]$ by (3-10). Rearranging,

$$\theta' \geq \frac{a^2}{3r^3\theta} - \frac{\theta}{r} \quad (3-11)$$

for $r \in (0, R_1 + 1]$.

We now proceed to mollify θ in such a way that the hypotheses of the lemma hold. Let $\eta_\varepsilon(r) = (1/\varepsilon)\eta(r/\varepsilon)$ denote a standard mollifier, where $\eta \in \mathcal{C}_c^\infty((-1, 1))$ has integral one. In addition, choose a cutoff $H = H(r) \in \mathcal{C}^\infty(X; [0, 1])$ such that

$$H = 1 \quad \text{for } r < R_1 + \frac{1}{4}, \quad H = 0 \quad \text{for } r > R_1 + \frac{1}{2},$$

and $H' \leq 0$. Now define

$$\theta_\varepsilon = (1 - H)\theta + \eta_\varepsilon * (H\theta).$$

Clearly θ_ε is smooth, and $\theta_\varepsilon \rightarrow \theta$ uniformly for $r \in [0, R_1 + 1]$. Furthermore, there exists $\varepsilon_0 > 0$ such that if $\varepsilon \in (0, \varepsilon_0)$, then the following properties are satisfied:

- $\theta_\varepsilon(r) < 0$ and $\theta'_\varepsilon(r) \geq 0$ for $r \in [0, R_1 + 1]$.
- $\theta_\varepsilon(r) = \varphi'(r)$ for $r \in [R_1 + \frac{3}{4}, R_1 + 1]$,
- There exists $r_1 \in (0, R_0]$ such that $\theta_\varepsilon(r) = k(R_0)$ for $r \in [0, r_1]$.

Since θ is continuous and piecewise smooth,

$$\theta'_\varepsilon = (1 - H)\theta' - H'\theta + \eta_\varepsilon * (H'\theta + H\theta'). \quad (3-12)$$

Therefore by (3-11),

$$\theta'_\varepsilon \geq -H'\theta + \eta_\varepsilon * (H'\theta) + (1 - H)\left(\frac{a^2}{3r^3\theta} - \frac{\theta}{r}\right) + \eta_\varepsilon * \left(H\left(\frac{a^2}{3r^3\theta} - \frac{\theta}{r}\right)\right)$$

for $r \in (0, R_1 + 1]$. The right-hand side converges uniformly to $a^2/(3r^3\theta) - \theta/r$ for $r \in [r_1, R_1 + 1]$ since the latter function is continuous there. Since $\theta_\varepsilon \rightarrow \theta$ uniformly for $r \in [r_1, R_1 + 1]$ as well, there exists $\varepsilon \in (0, \varepsilon_0)$ such that

$$-\frac{3a^2}{2r} + 3r\theta_\varepsilon^2 + 3r^2\theta_\varepsilon\theta'_\varepsilon \leq 0$$

for $r \in [r_1, R_1 + 1]$. This inequality is also true for $r \in (0, r_1)$, since $\theta_\varepsilon = k(R_0)$ on that interval. Now extend φ from $\{r \geq R_1 + 1\}$ to $\{r < R_1 + 1\}$ by the formula

$$\varphi(r) = \varphi(R_1 + 1) + \int_{R_1+1}^r \theta_\varepsilon(s) ds.$$

This completes the proof according to (3-9) by observing that the φ just constructed satisfies $\varphi''(r) \geq 0$. \square

As a remark, if $\tau \neq 0$, then the hypoellipticity condition also holds along $\{r = 0\}$, simply because $\text{Im } G_\varphi \neq 0$ in that case. However, since $(x, \xi) \mapsto G(x, \xi)$ is not elliptic along $\{r = 0\}$, the hypoellipticity condition alone, stated here in the semiclassical setting, is not sufficient to prove a Carleman estimate; see [Hörmander 1963, Section 8.4].

Now that the phases φ_1, φ_2 have been constructed globally, we are ready to finish the proof of Proposition 3.1. Here we come back to the operator $P_\varphi(z)$ on X . Fix a norm $|\cdot|$ on the fibers of T^*X (for instance using the induced metric k) and let $\langle \xi \rangle = (1 + |\xi|^2)^{1/2}$.

Proof of Proposition 3.1. Recall that we are given $[a, b] \subset (0, \infty)$ and a compact set $K \subset X^\circ$. Without loss, we may assume that $K = \{r \geq \varepsilon\}$ for some $\varepsilon > 0$. Let B_i, U_i be as in the discussion preceding Lemma 3.3. In particular,

$$\{\operatorname{Re} p_{\varphi_i}, \operatorname{Im} p_{\varphi_i}\} > 0$$

on $(\{p_{\varphi_i} = 0\} \cap \{r \geq \varepsilon/2\}) \setminus T_{U_i}^* X$. Let $\chi_i \in C_c^\infty(B_i^\circ)$ be such that $\chi_i = 1$ near U_i . If $\varphi = \varphi_i$, then

$$|p_\varphi|^2 + \chi^2 + h\{\operatorname{Re} p_\varphi, \operatorname{Im} p_\varphi\} \geq h(M|p_\varphi|^2 + M\chi^2 + \{\operatorname{Re} p_\varphi, \operatorname{Im} p_\varphi\})$$

for any $M > 0$, provided that $h > 0$ is sufficiently small. On the other hand, the set $\{\operatorname{Re} p_\varphi = 0\} \cap \{r \geq \varepsilon/2\}$ is compact by Lemma 2.1, uniformly for $\pm z \in [a, b]$. Therefore,

$$\langle \xi \rangle^{-4} (M|p_\varphi|^2 + M\chi^2 + \{\operatorname{Re} p_\varphi, \operatorname{Im} p_\varphi\}) > 0$$

near $T^*X \cap \{r \geq \varepsilon/2\}$ for $M > 0$ sufficiently large. By (3-1) and the semiclassical Gårding inequality applied to $e^{\varphi_i/h} u$,

$$h\|e^{\varphi_i/h} u\|_{H_h^2(X)}^2 \leq C\|e^{\varphi_i/h} P(z)u\|_{L^2(X)}^2 + C\|e^{\varphi_i/h} u\|_{L^2(B_i)}^2 \quad (3-13)$$

for $u \in C_c^\infty(K^\circ)$ and $i = 1, 2$. Since $\varphi_1 > \varphi_2$ on B_2 and $\varphi_2 > \varphi_1$ on B_1 , there is $\gamma > 0$ such that

$$e^{\varphi_i/h} \leq e^{-\gamma/h} (e^{\varphi_1/h} + e^{\varphi_2/h})$$

on B_i . Now add (3-13) for $i = 1, 2$ to absorb the integral over $B_1 \cup B_2$ into the left-hand side. \square

4. Degenerate Carleman estimates near the boundary

4A. Statement of result. We now complement Proposition 3.1 with a result valid up to the boundary. Recall that the phases φ_1, φ_2 are equal on $\{r \leq 1\}$. Since we are working near ∂X , we will thus drop the subscript and simply write φ .

Proposition 4.1. *Given $[a, b] \subset (0, \infty)$ there exists $r_0 > 0$ and $C > 0$ such that*

$$\|e^{\varphi/h} u\|_{H_{b,h}^1} \leq C(h^{-1/2}\|e^{\varphi/h} P(z)u\|_{L^2} + e^{\varphi(0)/h}\|u\|_{L^2(\partial X)}) \quad (4-1)$$

for $u \in C_c^\infty(\{r < r_0\})$ and $\pm z \in [a, b]$.

The Sobolev space appearing on the left-hand side of (4-1) is modeled on the space of vector fields $\mathcal{V}_b(X)$ which are tangent to the boundary; see [Melrose 1993]. Thus $u \in H_b^1(X)$ if $u \in L^2(X)$ and $Ku \in L^2(X)$ for any $K \in \mathcal{V}_b(X)$. If $u \in H_b^1(X)$ and $\operatorname{supp} u \subset \{r < 1\}$, we can set

$$\|u\|_{H_{b,h}^1}^2 = \int_X |u|^2 + h^2 |r \partial_r u|^2 + h^2 k^{AB} (\partial_A u \cdot \partial_B \bar{u}) dS_X.$$

Of course away from ∂X this is equivalent to the full H_h^1 norm. Observe that it is enough to prove Proposition 4.1 for the operator $L = \square_g$, since the estimate (4-1) is stable under perturbations $B \in h\operatorname{Diff}_h^1(X)$ provided that the vector field part of B is tangent to ∂X . The latter condition is satisfied by the hypothesis that \mathcal{W} is tangent to ∂M made in the Introduction.

Proposition 4.1 is proved through integration by parts. A convenient way of carrying out this procedure is by constructing an appropriate multiplier for the wave operator and applying the divergence theorem. This approach to Carleman estimates for certain geometric operators is partly inspired by [Alexakis and Shao 2015; Ionescu and Klainerman 2009].

4B. The divergence theorem. We will use the divergence theorem in the time-differentiated form

$$\frac{d}{dt} \int_X g(K, N) dS_X + \int_{\partial X} g(K, T) dS_{\partial X} = \int_X (\operatorname{div}_g K) A dS_X, \quad (4-2)$$

valid for any vector field K , see [Warnick 2015, Lemma 3.1] for instance, where recall $X = \{t = 0\}$. Thus the first term on the left-hand side of (4-2) is short-hand for

$$\frac{d}{ds} \int_{\{t=s\}} g(K, N) dS_X \text{ evaluated at } s = 0.$$

Here $dS_{\partial X}$ is the volume density on ∂X induced by k (the latter is Riemannian, and hence the induced volume density is well-defined).

4C. Stress-energy tensor. Given $v \in C^\infty(M)$, let $Q = Q[v]$ denote the usual stress-energy tensor associated to v with components

$$Q_{\alpha\beta} = \operatorname{Re}(\partial_\alpha v \cdot \partial_\beta \bar{v}) - \frac{1}{2} g^{-1}(dv, d\bar{v}) g_{\alpha\beta}.$$

This tensor has the property that $(\nabla^\beta Q_{\alpha\beta}) S^\alpha = \operatorname{Re}(\square v \cdot S \bar{v})$ for any vector field S . Given such a vector field and a function w , define the modified vector field $J = J[v]$ with components

$$J^\alpha = Q^\alpha_\beta S^\beta + \frac{1}{2} w \cdot \partial^\alpha (|v|^2) - \frac{1}{2} (\partial^\alpha w) |v|^2.$$

The relevant choices in this context are

$$S = \nabla_g r, \quad w = \lambda + \frac{1}{2} \square_g r, \quad (4-3)$$

where $\lambda = \lambda(r)$ is an undetermined function to be chosen in Lemma 4.4 below. Also, introduce the tensor Π with components

$$\Pi^{\alpha\beta} = -\nabla^{\alpha\beta} r - \lambda g^{\alpha\beta}.$$

The divergence of J satisfies

$$\operatorname{Re}(\square_g u \cdot (S \bar{v} + w \bar{v})) = \operatorname{div}_g J + \Pi(dv, d\bar{v}) + \frac{1}{2} (\square_g w) |v|^2, \quad (4-4)$$

which is verified by a direct calculation.

4D. The conjugated operator. Near ∂M , consider the conjugated operator $L_\Phi = e^\Phi \square_g e^{-\Phi}$, where $\Phi = \Phi(r)$. Then, L_Φ has the expression

$$\begin{aligned} L_\Phi &= \square_g - 2\Phi' S + ((\Phi')^2 - \Phi'') g^{-1}(dr, dr) - \Phi' \square_g r \\ &= \square_g - 2\Phi' S + V_0. \end{aligned}$$

Now $g^{-1}(dr, dr) = -r$ by assumption, and consequently the potential term V_0 satisfies

$$V_0 = r(\Phi'' - (\Phi')^2) - \Phi' \square_g r.$$

Set $V_1 = V_0 - 2\Phi'w$, multiply $L_\Phi v$ by $S\bar{v} + w\bar{v}$, and take the real part to find that

$$\operatorname{Re}(L_\Phi v \cdot (S\bar{v} + w\bar{v})) = \operatorname{Re}(\square_g v \cdot (S\bar{v} + w\bar{v})) - 2\Phi'|Sv|^2 + \operatorname{Re} V_1 v \cdot S\bar{v} + V_0 w|v|^2. \quad (4-5)$$

It is also convenient to write $\operatorname{Re}(V_1 v \cdot S\bar{v})$ as a divergence,

$$\operatorname{Re}(V_1 v \cdot S\bar{v}) = \frac{1}{2} \operatorname{div}_g(V_1 |v|^2 S) - \frac{1}{2} (S(V_1) + V_1 \square_g r) |v|^2.$$

In view of this expression, define the vector field $K = J + \frac{1}{2} V_1 |v|^2 S$. For future use, also define the modified potential V by

$$V = \frac{1}{2} (\square_g w) + V_0 w - \frac{1}{2} S(V_1) - \frac{1}{2} V_1 \square_g r + \Phi' w^2. \quad (4-6)$$

On one hand, integrating the divergence of K yields boundary integrals; the following special case of this will suffice.

Lemma 4.2. *Let $v \in C^\infty(M)$ be given by $v = e^{-izt/h} u$, where u is stationary and $z \in \mathbb{R}$. Then,*

$$\int_X (\operatorname{div}_g K) A dS_X = - \left| \frac{z}{h} \right|^2 \int_{\partial X} |u|^2 dS_{\partial X}.$$

Proof. Apply the divergence theorem (4-2). Since $z \in \mathbb{R}$, the vector field K is stationary, and hence there is no contribution from the time derivative. As for the integral over ∂M , observe that T is null and $S = -T$ on the horizon. Since $Tv = -i(z/h)v$, it follows that $g(T, K) = -|Tv|^2 = -|z/h|^2 |u|^2$ on ∂M . \square

Note that the boundary contribution from Lemma 4.2 has an unfavorable sign, which will account for the boundary term in Proposition 4.1. On the other hand, the divergence of K can also be expressed in terms of (4-5).

Lemma 4.3. *If $\Phi' < 0$, then the divergence of K satisfies*

$$(2|\Phi'|)^{-1} |L_\Phi v|^2 \geq \operatorname{div}_g K + \Pi(dv, d\bar{v}) - \Phi'|Sv|^2 + V|v|^2, \quad (4-7)$$

where V is given by (4-6).

Proof. Combine (4-5) with (4-4), and then use the Cauchy–Schwarz inequality to find

$$\operatorname{Re}(L_\Phi v \cdot (S\bar{v} + w\bar{v})) \leq (2|\Phi'|)^{-1} |L_\Phi v|^2 - \Phi'(|Sv|^2 + w^2|v|^2),$$

recalling that $\Phi' < 0$. \square

4E. Pseudoconvexity. To examine positivity properties of $\Pi(dv, d\bar{v}) - \Phi'|Sv|^2$, we establish a certain pseudoconvexity condition. A criterion of this type first appeared in work of Alinhac [1984] on unique continuation, and was also employed in [Ionescu and Klainerman 2009; Alexakis and Shao 2015]. Recall that the Poisson bracket is related to the Hessian via the formula

$$\{G, \{G, f\}\}(x, \varpi) = 4\varpi_\alpha \varpi_\beta \nabla^{\alpha\beta} f, \quad (4-8)$$

valid for any $f \in C^\infty(M)$.

Lemma 4.4. *There exists $M, c, R_0 > 0$, and a function $\lambda = \lambda(r)$ such that*

$$M\{G, r\}^2 - \{G, \{G, r\}\} - 4\lambda G \geq c((r\rho)^2 + \tau^2 + k^{AB}\eta_A\eta_B) \quad (4-9)$$

for $r \in [0, R_0]$.

Proof. Throughout, assume that $M \geq 1$. Let $r \leq (4M)^{-1}$, and define the function λ by

$$\lambda = \frac{1}{2} - (1 - \delta)rM,$$

where $\delta > 0$ will be chosen sufficiently small. Observe that $\frac{1}{4} \leq \lambda \leq \frac{1}{2}$ uniformly in $M \geq 1$ for $r \leq (4M)^{-1}$. Denote the left-hand side of (4-9) by $4\mathcal{E}$, and the corresponding quantity by $4\mathcal{E}_0$ if G is replaced with G_0 . Dividing through by 4,

$$\mathcal{E}_0 = M((r\rho)^2 + 2r\rho\tau + \tau^2) - \frac{1}{2}(r\rho^2 + 2\rho\tau) - \lambda G_0. \quad (4-10)$$

Use the expression for λG_0 and the lower bound $\lambda \geq \frac{1}{4}$ on $\{r \leq (4M)^{-1}\}$ to find that

$$\mathcal{E}_0 \geq M\delta((r\rho)^2 + 2r\rho\tau) + M\tau^2 + \frac{1}{4}k_0^{AB}\eta_A\eta_B.$$

Therefore $\mathcal{E}_0 \geq c(MY + k^{AB}\eta_A\eta_B)$ if $\delta > 0$ is sufficiently small, where recall $Y = (r\rho)^2 + \tau^2$.

Now consider the error $\mathcal{E} - \mathcal{E}_0$ incurred by replacing G with G_0 . Replacing $M\{G, r\}^2$ with $M\{G_0, r\}^2$ produces an error

$$2M\{G_0, r\}\{G - G_0, r\} + M\{G - G_0, r\}^2.$$

Using Cauchy–Schwarz on the first term to absorb a small multiple of $M\{G_0, r\}^2$ into \mathcal{E}_0 (in other words, changing the constant $c > 0$ in the lower bound for \mathcal{E}_0 above) leaves an overall error of the form

$$M(\mathcal{N}_1 \cdot \mathcal{N}_1) \subset (rM)\mathcal{N}_2.$$

The factor of rM is harmless since $rM \leq \frac{1}{4}$; thus the right-hand side is certainly in \mathcal{N}_2 uniformly in $M \geq 1$. Using that λ is uniformly bounded in $M \geq 1$ on $\{r \leq (4M)^{-1}\}$, the remaining errors $\lambda(G - G_0)$ and

$$\{G - G_0, \{G - G_0, r\}\} + \{G - G_0, \{G_0, r\}\} + \{G_0, \{G - G_0, r\}\}$$

are also in \mathcal{N}_2 by Lemma 2.6, uniformly in $M \geq 1$. Now apply the first bound in Lemma 2.5, choosing $\gamma > 0$ sufficiently small but independent of M so that $\gamma k^{AB}\eta_A\eta_B$ can be absorbed by $ck^{AB}\eta_A\eta_B$ on the right-hand side for $r \in [0, r_\gamma]$. This leaves a large multiple of Y , which is then absorbed by MY on the right-hand side by taking M sufficiently large. It then suffices to take $R_0 = \min\{(4M)^{-1}, r_\gamma\}$. \square

Fix $M > 0$ such that Lemma 4.4 is valid. This fixes the function λ , and therefore the function w in (4-3). Lemma 4.3 will be applied with the weight $\Phi = \varphi_i/h$, viewed as a stationary function on M . In particular, $\Phi' = -C/h$ on $\{r \leq r_1\}$ for some constant $C > 0$ (recall the statement of Proposition 3.1).

Before proceeding, consider the potential term V from Lemma 4.3. Instead of analyzing its sign, we more simply note that for $F' = -C/h$ one has

$$V = f_0 + h^{-1}f_1 + h^{-2}f_2, \quad (4-11)$$

where $f_0, f_1 \in C^\infty(M)$ and $f_2 \in rC^\infty(M)$. The small coefficient of f_2 means V can be treated as an error. To be precise, we have the following positivity result for the bulk terms.

Lemma 4.5. *Given $a > 0$, there exists $c, r_0 \geq 0$ such that if $|z| \geq a$, then*

$$\Pi(dv, d\bar{v}) - \Phi'|Sv|^2 + V|v|^2 \geq c(h^{-2}|u|^2 + |r \partial_r u|^2 + k^{AB} \partial_A u \partial_B \bar{u}) \quad (4-12)$$

on $\{r \leq r_0\}$ for each $v \in C^\infty(M)$ of the form $v = e^{-izt/h}u$, where u is stationary.

Proof. Since $\Phi' = -C/h$, an inequality of the form (4-12) is true for sufficiently small $h > 0$ if the term $V|v|^2$ is dropped from the left-hand side; this follows from Lemma 4.4 and (4-8). On the other hand, for a potential V satisfying (4-11), there is clearly $r_0 > 0$ such that $V|v|^2$ can be absorbed by $ch^{-2}|v|^2$ for $r \in [0, r_0]$ and $h > 0$ sufficiently small. \square

The proof of Proposition 4.1 is now immediate:

Proof of Proposition 4.1. Given $[a, b] \subset (0, \infty)$, apply Lemmas 4.2, 4.3, and 4.5 to functions of the form $v = e^{-izt/h}e^{\varphi/h}u$, where $\pm z \in [a, b]$ and $\text{supp } u \subset \{r < r_0\}$. \square

5. Proof of Theorem 1

We prove the equivalent Theorem 1'. Assume that $[a, b] \subset (0, \infty)$ has been fixed. Choose a cutoff function $\chi \in C^\infty(X)$ such that

$$\text{supp } \chi \subset \{r < r_0\}, \quad \chi = 1 \quad \text{near } \{r \leq r_0/2\},$$

where r_0 is provided by Lemma 4.5. Then, apply Proposition 4.1 to χu and Proposition 3.1 to $(1 - \chi)u$, where $u \in C^\infty(X)$. Since the commutator $[P(z), \chi]$ is supported away from ∂X , the error terms can be absorbed even though the left-hand side is only estimated in the $H_{b,h}^1$ norm. Bounding $e^{\varphi_1/h} + e^{\varphi_2/h}$ from below on the left and from above on the right yields

$$\|u\|_{H_{b,h}^1} \leq e^{C/h} (\|P(z)u\|_{L^2} + \|u\|_{L^2(\partial X)}) \quad (5-1)$$

for $u \in C^\infty(X)$ and $\pm z \in [a, b]$.

Next, we remove the boundary term on the right-hand side of (5-1). In order to estimate the boundary term, we use that L is formally self-adjoint and that \mathcal{W} is tangent to ∂M . Apply the divergence theorem (4-2) to the vector field $\bar{v} \nabla_g v - v \nabla_g \bar{v} + |v|^2 \cdot \mathcal{W}$ with $v = e^{-izt/h}u$. Since L is formally self-adjoint, we obtain Green's formula

$$(hz) \int_{\partial X} |u|^2 dS_{\partial X} = -\text{Im} \int_X P(z)u \cdot \bar{u} A dS_X.$$

There is no boundary contribution coming from \mathcal{W} since we assumed $g(T, \mathcal{W})$ vanishes on ∂M . Applying Cauchy-Schwarz to the right-hand side implies

$$e^{C/h} \|u\|_{L^2(\partial X)} \leq C_\varepsilon h^{-1} e^{2C/h} \|P(z)u\|_{L^2} + \varepsilon \|u\|_{L^2}$$

for some C_ε and every $\pm z \in [a, b]$. Therefore the boundary term on the right-hand side of (5-1) can be absorbed into the left-hand side by taking ε sufficiently small, at the expense of increasing the constant in the exponent $e^{C/h}$. We then have

$$\|u\|_{H_{b,h}^1} \leq e^{C/h} \|P(z)u\|_{L^2}.$$

The final step is to apply a bound of the form

$$\|u\|_{H_h^{k+1}} \leq Ch^{-1} (\|P(z)u\|_{H_h^k} + \|u\|_{L^2}) \quad (5-2)$$

for $u \in \mathcal{C}^\infty(X)$ and $\pm z \in [a, b]$. The most conceptual way of understanding this estimate is in terms of the semiclassical trapping present in the interior of M . For an appropriate pseudodifferential complex absorbing operator $Q \in \Psi_h^{-\infty}(X^\circ)$ with compact support in X° , the nontrapping framework of [Vasy 2013, Section 2.8] shows that $P(z) - iQ$ satisfies the nontrapping bound

$$\|u\|_{H_h^k} \leq Ch^{-1} \|(P(z) - iQ)u\|_{H_h^k}$$

for $z \in [a, b]$. Here Q is chosen to be elliptic (with the correct choice of sign) on the trapped set. In this case Q can be chosen to have compact microsupport in X° , and hence $Q : \mathcal{C}^{-\infty}(X) \rightarrow \mathcal{C}^\infty(X)$, and in particular

$$\|Qu\|_{H_h^k} \leq C\|u\|_{L^2}.$$

This clearly implies (5-2) for $z \in [a, b]$, with a similar argument when $-z \in [a, b]$.

This completes the proof of Theorem 1' in the case when $u \in \mathcal{C}^\infty(X)$ and $\pm z \in [a, b]$. By perturbation, this extends to a region $\pm z \in [a, b] + ie^{-C_1/h}[-1, 1]$. Simply write

$$P(z) - P(\operatorname{Re} z) = \operatorname{Im} z \cdot B(z),$$

where $B(z) \in \operatorname{Diff}_h^1(X)$ is bounded $H_h^{k+1}(X) \rightarrow H_h^k(X)$ uniformly for $z \in [a, b]$ (although $B(z)$ is not holomorphic in z). Thus the difference can be absorbed into the left-hand side if $|\operatorname{Im} z| \leq e^{-C_1/h}$ for $C_1 > 0$ sufficiently large. Finally, $\mathcal{C}^\infty(X)$ is dense in \mathcal{X}^k , see [Dyatlov and Zworski 2018, Lemma E.47], so (2-2) is valid for $u \in \mathcal{X}^k$ as well, thus completing the proof of Theorem 1'.

6. Logarithmic energy decay

6A. A semigroup formulation. In this section we outline how Corollary 1.2 can be deduced from the resolvent estimate (1-3) via semigroup theory. The starting point is that the Cauchy problem (1-4) is associated with a \mathcal{C}^0 semigroup $U(t) = e^{-itB}$ on $\mathcal{H}^k = H^{k+1}(X) \times H^k(X)$ satisfying

$$\|U(t)\|_{\mathcal{H}^k \rightarrow \mathcal{H}^k} \leq Ce^{\nu t} \quad (6-1)$$

for some $C, \nu > 0$ [Warnick 2015, Corollary 3.14]. Recalling the lapse function $A = g^{-1}(dt, dt)^{-1/2}$, write

$$\square_g = L_2 + L_1 \partial_t + A^{-2} \partial_t^2,$$

where L_j is identified with a differential operator on X of order j . Thus $L_2 = \mathbf{P}(0)$ and $L_1 = i \partial_\omega \mathbf{P}(0)$. More explicitly,

$$L_1 = -2A^{-2}W - \operatorname{div}_g(A^{-2}W),$$

where W is the shift vector from Section 2D. The infinitesimal generator is then given by

$$-iB = \begin{pmatrix} 0 & 1 \\ -A^2 L_2 & -A^2 L_1 \end{pmatrix}. \quad (6-2)$$

Indeed, applying $U(t)$ to initial data in $\mathcal{C}_c^\infty(X^\circ)$ shows that $-iB$ is given by (6-2) in the sense of distributions. Now the resolvent set of B is nonempty, and indeed $\sigma(B) \subset \{\operatorname{Im} \omega \leq \nu\}$ by (6-1). Therefore the domain $D(B)$ of B is characterized as those distributions $(v_0, v_1) \in \mathcal{H}^k$ such that

$$v_1 \in H^{k+1}(X), \quad L_2 v_0 + L_1 v_1 \in H^k(X).$$

Since $L_2 = \mathbf{P}(0)$ and $L_1 \in \operatorname{Diff}^1(X)$, this shows that the domain of B is

$$D(B) = \mathcal{X}^k \times H^{k+1}(X),$$

where \mathcal{X}^k is defined by (1-2). It is also easy to see that the graph norm on $D(B)$ satisfies

$$\|B(v_0, v_1)\|_{\mathcal{H}^k} + \|(v_0, v_1)\|_{\mathcal{H}^k} \leq C \|(v_0, v_1)\|_{\mathcal{X}^k \times H^{k+1}};$$

hence the two norms on $D(B)$ are equivalent by the open mapping theorem. Furthermore, the spectrum of B in $\{\operatorname{Im} \omega > -\kappa(k + \frac{1}{2})\}$ coincides with poles of $\mathbf{P}(\omega)^{-1}$, and the resolvent estimate (1-3) translates into the bound $\|(B - \omega)^{-1}\|_{\mathcal{H}^k \rightarrow \mathcal{H}^k} \leq e^{C|\operatorname{Re} \omega|}$ for $\omega \in \Omega$.

6B. Logarithmic stabilization of semigroups. The goal now is to apply a theorem on the logarithmic stabilization of certain bounded semigroups:

Theorem 2 [Burq 1998, Theorem 3, Batty and Duyckaerts 2008, Theorem 1.5]. *Let $U(t) = e^{-itB}$ be a bounded \mathcal{C}^0 semigroup on a Hilbert space \mathcal{H} . If $\sigma(B) \cap \mathbb{R} = \emptyset$ and $\|(B - \omega)^{-1}\|_{\mathcal{H} \rightarrow \mathcal{H}} \leq e^{C|\omega|}$ for $\omega \in \mathbb{R}$, then there exists $C > 0$ such that*

$$\|U(t)v\|_{\mathcal{H}} \leq \frac{C}{\log(2+t)} \|(B - i)v\|_{\mathcal{H}}$$

for each $v \in D(B)$.

A priori the semigroup $U(t)$ from Section 6A is not uniformly bounded in time on \mathcal{H}^k , since the energy $\mathcal{E}_k[v](t)$ does not control the L^2 norm of $v(t)$. Instead, observe that $\operatorname{span}\{(1, 0)\} \subset \mathcal{H}^k$ is invariant under $U(t)$, which therefore descends to a semigroup $\widehat{U}(t)$ on the quotient space

$$\widehat{\mathcal{H}}^k = \mathcal{H}^k / \operatorname{span}\{(1, 0)\}.$$

If $\pi : \mathcal{H}^k \rightarrow \widehat{\mathcal{H}}^k$ is the natural projection, then, the infinitesimal generator of $\widehat{U}(t)$ is simply the operator \widehat{B} induced by B on $\pi(D(B))$. It follows from (1-5) and the Poincaré inequality that $\widehat{U}(t)$ is a bounded \mathcal{C}^0 semigroup.

Since $\text{span}\{(0, 1)\}$ is finite-dimensional, the spectrum of \widehat{B} is contained in the spectrum of B , and furthermore the bound

$$\|(\widehat{B} - \omega)^{-1}\|_{\widehat{\mathcal{H}} \rightarrow \widehat{\mathcal{H}}} \leq e^{C|\text{Re } \omega|}$$

also holds for $\omega \in \Omega$. The final step is to show $\sigma(\widehat{B}) \cap \mathbb{R} = \emptyset$. If $\omega \in \mathbb{R} \setminus 0$, this follows from the fact that $P(\omega)^{-1}$ has no nonzero real poles [Warnick 2015, Lemma A.1].

Finally, consider the spectrum at $\omega = 0$. If ω_0 is a pole of $(B - \omega)^{-1}$ acting on \mathcal{H}^k with $\text{Im } \omega_0 > -\kappa(k + \frac{1}{2})$, then its Laurent coefficients all map into $\mathcal{C}^\infty(X) \times \mathcal{C}^\infty(X)$ [Vasy 2013, Section 2.6]. Thus $\ker B \subset \mathcal{X}^k \times H^{k+1}(X)$ is in one-to-one correspondence with smooth stationary solutions of $\square_g v = 0$. If $\square_g v = 0$ for v smooth and stationary, then (4-2) applied to the vector field $\bar{v} \nabla_g v + v \nabla_g \bar{v}$ shows that $g^{-1}(dv, d\bar{v}) = 0$ on X . Again using that v is stationary, Lemma 2.1 implies $dv = 0$, and hence v is constant. Thus $\ker B = \text{span}\{(1, 0)\}$, so $0 \notin \sigma(\widehat{B})$.

The hypotheses of Theorem 2 are therefore satisfied by $\widehat{U}(t)$, which yields the bound

$$\|\widehat{U}(t) \circ \pi(v_0, v_1)\|_{\widehat{\mathcal{H}}^k} \leq \frac{C}{\log(2+t)} \|(B - i)(v_0, v_1)\|_{\mathcal{H}^k} \quad (6-3)$$

for each $(v_0, v_1) \in \mathcal{X}^k \times H^{k+1}(X)$. This establishes Corollary 1.2, since the norm on the left-hand side of (6-3) is equivalent to $\mathcal{E}_k[v](t)^{1/2}$, where v solves the Cauchy problem (1-4) with initial data (v_0, v_1) .

6C. Decay to a constant. To prove Corollary 1.3, consider the Laurent expansion of $(B - \omega)^{-1}$ about $\omega = 0$. The range of the corresponding residue Π_0 consists of all generalized eigenvectors, and contains $\text{span}\{(1, 0)\}$.

If the algebraic multiplicity of $\omega = 0$ was greater than 1, then there would exist a solution of $\square_g v = 0$ of the form

$$v(t, x) = u(x) + t,$$

where $u \in \mathcal{C}^\infty(M)$ is stationary. This is compatible with energy boundedness, but not with the logarithmic energy decay established above. Thus $\omega = 0$ is a simple pole with algebraic multiplicity 1.

By standard spectral theory, Π_0 is the projection onto $\text{span}\{(1, 0)\}$ along $\text{range}(B)$, so

$$\Pi_0 = \langle \cdot, \psi \rangle (1, 0)$$

for some $\psi \in (\ker B)'$, which we identify with $(\mathcal{H}^k)' / \text{range}(B^*) = \ker(B^*)$. Furthermore, ψ is uniquely determined by requiring that $\langle (1, 0), \psi \rangle = 1$. Here the duality between \mathcal{H}^k and

$$(\mathcal{H}^k)' = \dot{H}^{-k-1}(X) \times \dot{H}^{-k}(X)$$

is induced by the $L^2(X)$ inner product described in Section 3B, where $\dot{H}^s(X)$ is the Sobolev space of supported distributions in the sense of [Hörmander 1985, Appendix B.2].

The domain of B^* consists of all $w \in \dot{H}^{-k-1}(X) \times \dot{H}^{-k}(X)$ for which there exists $v \in \dot{H}^{-k-1}(X) \times \dot{H}^{-k}(X)$ satisfying $(w, Bu) = (v, u)$ for every $u \in D(B) = \mathcal{X}^k \times H^{k+1}(X)$. Thus

$$D(B^*) = \dot{H}^{-k-1}(X) \times \dot{\mathcal{X}}^{-k},$$

where we define

$$\dot{\mathcal{H}}^{-k} = \{u \in \dot{H}^{-k}(X) : P(0) \in \dot{H}^{-k-1}(X)\}.$$

The action of B^* is given by

$$iB^* = \begin{pmatrix} 0 & -L_2 A^2 \\ 1 & L_1 A^2 \end{pmatrix},$$

using that L_1 is skew-adjoint.

Now we compute the kernel of B^* , which again by abstract spectral theory is one-dimensional. Let $\psi_1 = \text{vol}(\partial X)^{-1} A^{-2} \in L^2(X)$, viewed as an element of $\dot{H}^{-k}(X)$ via the $L^2(X)$ inner product, and then set

$$\psi_0 = -\text{vol}(\partial X)^{-1} L_1(1) \in \dot{H}^{-k-1}(X)$$

in the sense of supported distributions. If we set $\psi = (\psi_0, \psi_1)$, then $B^*\psi = 0$. Furthermore,

$$\begin{aligned} \text{vol}(\partial X) \langle 1, \psi_0 \rangle &= \langle L_1(1), 1 \rangle = - \int_X \text{div}_g(A^{-2}W) A \, dS_X \\ &= - \int_{\partial X} A^{-2} g(W, T) \, dS_{\partial X} = \int_{\partial X} dS_{\partial X} = \text{vol}(\partial X), \end{aligned}$$

since $g(W, T) = -g(AN, T) = -A^2$ on ∂X . Thus $\psi \in \ker B^*$ has the appropriate normalization.

Finally, let $E = \text{range}(I - \Pi_0)$, which is thus invariant under $U(t)$, and $U(t)|_E = U(t)(I - \Pi_0)$. Since

$$\mathcal{H}^k = E \dot{+} \text{span}\{(1, 0)\},$$

with $\dot{+}$ denoting a topological direct sum, it follows that E is isomorphic to the quotient $\widehat{\mathcal{H}}^k$ as a Banach space. Given $(v_0, v_1) \in D(B)$, define the constant $v_\infty = \langle v_0, \psi_0 \rangle + \langle v_1, \psi_1 \rangle$. Then

$$\|U(t)(v_0 - v_\infty, v_1)\|_{\mathcal{H}^k} = \|U(t)(v_0, v_1) - (v_\infty, 0)\|_{\mathcal{H}^k} \leq C \|\widehat{U}(t) \circ \pi(v_0, v_1)\|_{\widehat{\mathcal{H}}^k},$$

which completes the proof of Corollary 1.3.

Acknowledgments

This work was supported by the NSF grant DMS-1502632. The author would like to thank Georgios Moschidis for carefully explaining parts of his paper [Moschidis 2016], and Daniel Tataru for a discussion on Carleman estimates.

References

- [Alexakis and Shao 2015] S. Alexakis and A. Shao, “Global uniqueness theorems for linear and nonlinear waves”, *J. Funct. Anal.* **269**:11 (2015), 3458–3499. MR Zbl
- [Alinhac 1984] S. Alinhac, “Unicité du problème de Cauchy pour des opérateurs du second ordre à symboles réels”, *Ann. Inst. Fourier (Grenoble)* **34**:2 (1984), 89–109. MR Zbl
- [Batty and Duyckaerts 2008] C. J. K. Batty and T. Duyckaerts, “Non-uniform stability for bounded semi-groups on Banach spaces”, *J. Evol. Equ.* **8**:4 (2008), 765–780. MR Zbl
- [Burq 1998] N. Burq, “Décroissance de l’énergie locale de l’équation des ondes pour le problème extérieur et absence de résonance au voisinage du réel”, *Acta Math.* **180**:1 (1998), 1–29. MR Zbl

- [Burq 2002] N. Burq, “Lower bounds for shape resonances widths of long range Schrödinger operators”, *Amer. J. Math.* **124**:4 (2002), 677–735. MR Zbl
- [Cardoso and Vodev 2002] F. Cardoso and G. Vodev, “Uniform estimates of the resolvent of the Laplace–Beltrami operator on infinite volume Riemannian manifolds, II”, *Ann. Henri Poincaré* **3**:4 (2002), 673–691. MR Zbl
- [Datchev 2014] K. Datchev, “Quantitative limiting absorption principle in the semiclassical limit”, *Geom. Funct. Anal.* **24**:3 (2014), 740–747. MR Zbl
- [Dyatlov and Zworski 2018] S. Dyatlov and M. Zworski, “Mathematical theory of scattering resonances”, preprint, 2018, available at http://math.mit.edu/~dyatlov/res/res_20180406.pdf.
- [Gannot 2018] O. Gannot, “The null-geodesic flow near horizons”, *Trans. Amer. Math. Soc.* (online publication 2018).
- [Gilbarg and Trudinger 1983] D. Gilbarg and N. S. Trudinger, *Elliptic partial differential equations of second order*, 2nd ed., Grundlehren der Math. Wissenschaften **224**, Springer, 1983. MR Zbl
- [Guillarmou 2005] C. Guillarmou, “Meromorphic properties of the resolvent on asymptotically hyperbolic manifolds”, *Duke Math. J.* **129**:1 (2005), 1–37. MR Zbl
- [Holzegel and Smulevici 2013] G. Holzegel and J. Smulevici, “Decay properties of Klein–Gordon fields on Kerr–AdS space-times”, *Comm. Pure Appl. Math.* **66**:11 (2013), 1751–1802. MR Zbl
- [Hörmander 1963] L. Hörmander, *Linear partial differential operators*, Grundlehren der Math. Wissenschaften **116**, Springer, 1963. MR Zbl
- [Hörmander 1985] L. Hörmander, *The analysis of linear partial differential operators, III: Pseudo-differential operators*, Grundlehren der Math. Wissenschaften **274**, Springer, 1985. MR Zbl
- [Ionescu and Klainerman 2009] A. D. Ionescu and S. Klainerman, “On the uniqueness of smooth, stationary black holes in vacuum”, *Invent. Math.* **175**:1 (2009), 35–102. MR Zbl
- [Melrose 1993] R. B. Melrose, *The Atiyah–Patodi–Singer index theorem*, Research Notes in Math. **4**, A. K. Peters, Wellesley, MA, 1993. MR Zbl
- [Moschidis 2016] G. Moschidis, “Logarithmic local energy decay for scalar waves on a general class of asymptotically flat spacetimes”, *Ann. PDE* **2**:1 (2016), art. id. 5. MR
- [Rodnianski and Tao 2015] I. Rodnianski and T. Tao, “Effective limiting absorption principles, and applications”, *Comm. Math. Phys.* **333**:1 (2015), 1–95. MR Zbl
- [Sá Barreto and Zworski 1997] A. Sá Barreto and M. Zworski, “Distribution of resonances for spherical black holes”, *Math. Res. Lett.* **4**:1 (1997), 103–121. MR Zbl
- [Vasy 2013] A. Vasy, “Microlocal analysis of asymptotically hyperbolic and Kerr–de Sitter spaces”, *Invent. Math.* **194**:2 (2013), 381–513. MR Zbl
- [Vodev 2000] G. Vodev, “Exponential bounds of the resolvent for a class of noncompactly supported perturbations of the Laplacian”, *Math. Res. Lett.* **7**:3 (2000), 287–298. MR Zbl
- [Warnick 2015] C. M. Warnick, “On quasinormal modes of asymptotically anti-de Sitter black holes”, *Comm. Math. Phys.* **333**:2 (2015), 959–1035. MR Zbl
- [Zworski 2012] M. Zworski, *Semiclassical analysis*, Graduate Studies in Math. **138**, Amer. Math. Soc., Providence, RI, 2012. MR Zbl
- [Zworski 2017] M. Zworski, “Mathematical study of scattering resonances”, *Bull. Math. Sci.* **7**:1 (2017), 1–85. MR Zbl

Received 20 Feb 2018. Revised 27 Apr 2018. Accepted 30 May 2018.

ORAN GANNOT: gannot@northwestern.edu

Department of Mathematics, Northwestern University, Evanston, IL, United States

Analysis & PDE

msp.org/apde

EDITORS

EDITOR-IN-CHIEF

Patrick Gérard
patrick.gerard@math.u-psud.fr
Université Paris Sud XI
Orsay, France

BOARD OF EDITORS

| | | | |
|----------------------|---|-----------------------|--|
| Massimiliano Berti | Scuola Intern. Sup. di Studi Avanzati, Italy berti@sissa.it | Clément Mouhot | Cambridge University, UK c.mouhot@dpmms.cam.ac.uk |
| Sun-Yung Alice Chang | Princeton University, USA chang@math.princeton.edu | Werner Müller | Universität Bonn, Germany mueller@math.uni-bonn.de |
| Michael Christ | University of California, Berkeley, USA mchrist@math.berkeley.edu | Gilles Pisier | Texas A&M University, and Paris 6 pisier@math.tamu.edu |
| Alessio Figalli | ETH Zurich, Switzerland alessio.figalli@math.ethz.ch | Tristan Rivière | ETH, Switzerland riviere@math.ethz.ch |
| Charles Fefferman | Princeton University, USA cf@math.princeton.edu | Igor Rodnianski | Princeton University, USA irod@math.princeton.edu |
| Ursula Hamenstaedt | Universität Bonn, Germany ursula@math.uni-bonn.de | Sylvia Serfaty | New York University, USA serfaty@cims.nyu.edu |
| Vaughan Jones | U.C. Berkeley & Vanderbilt University vaughan.f.jones@vanderbilt.edu | Yum-Tong Siu | Harvard University, USA siu@math.harvard.edu |
| Vadim Kaloshin | University of Maryland, USA vadim.kaloshin@gmail.com | Terence Tao | University of California, Los Angeles, USA tao@math.ucla.edu |
| Herbert Koch | Universität Bonn, Germany koch@math.uni-bonn.de | Michael E. Taylor | Univ. of North Carolina, Chapel Hill, USA met@math.unc.edu |
| Izabella Laba | University of British Columbia, Canada ilaba@math.ubc.ca | Gunther Uhlmann | University of Washington, USA gunther@math.washington.edu |
| Gilles Lebeau | Université de Nice Sophia Antipolis, France lebeau@unice.fr | András Vasy | Stanford University, USA andras@math.stanford.edu |
| Richard B. Melrose | Massachusetts Inst. of Tech., USA rbm@math.mit.edu | Dan Virgil Voiculescu | University of California, Berkeley, USA dvv@math.berkeley.edu |
| Frank Merle | Université de Cergy-Pontoise, France Frank.Merle@u-cergy.fr | Steven Zelditch | Northwestern University, USA zelditch@math.northwestern.edu |
| William Minicozzi II | Johns Hopkins University, USA minicozz@math.jhu.edu | Maciej Zworski | University of California, Berkeley, USA zworski@math.berkeley.edu |

PRODUCTION

production@msp.org
Silvio Levy, Scientific Editor


See inside back cover or msp.org/apde for submission instructions.

The subscription price for 2019 is US \$310/year for the electronic version, and \$520/year (+\$60, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues from the last three years and changes of subscriber address should be sent to MSP.

Analysis & PDE (ISSN 1948-206X electronic, 2157-5045 printed) at Mathematical Sciences Publishers, 798 Evans Hall #3840, c/o University of California, Berkeley, CA 94720-3840, is published continuously online. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices.

APDE peer review and production are managed by EditFlow® from MSP.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2019 Mathematical Sciences Publishers

ANALYSIS & PDE

Volume 12 No. 2 2019

| | |
|---|-----|
| A unified flow approach to smooth, even L_p -Minkowski problems | 259 |
| PAUL BRYAN, MOHAMMAD N. IVAKI and JULIAN SCHEUER | |
| The Muskat problem in two dimensions: equivalence of formulations, well-posedness, and regularity results | 281 |
| BOGDAN-VASILE MATIOC | |
| Maximal gain of regularity in velocity averaging lemmas | 333 |
| DIOGO ARSÉNIO and NADER MASMOUDI | |
| On the existence and stability of blowup for wave maps into a negatively curved target | 389 |
| ROLAND DONNINGER and IRFAN GLOGIĆ | |
| Fracture with healing: A first step towards a new view of cavitation | 417 |
| GILLES FRANCFORT, ALESSANDRO GIACOMINI and OSCAR LOPEZ-PAMIES | |
| General Clark model for finite-rank perturbations | 449 |
| CONSTANZE LIAW and SERGEI TREIL | |
| On the maximal rank problem for the complex homogeneous Monge–Ampère equation | 493 |
| JULIUS ROSS and DAVID WITT NYSTRÖM | |
| A viscosity approach to the Dirichlet problem for degenerate complex Hessian-type equations | 505 |
| SŁAWOMIR DINEW, HOANG-SON DO and TAT DAT TÔ | |
| Resolvent estimates for spacetimes bounded by Killing horizons | 537 |
| ORAN GANNOT | |
| Interpolation by conformal minimal surfaces and directed holomorphic curves | 561 |
| ANTONIO ALARCÓN and ILDEFONSO CASTRO-INFANTES | |



2157-5045(2019)12:2;1-O