

# ANALYSIS & PDE

Volume 12      No. 3      2019

ALESSIO FIGALLI AND MOON-JIN KANG

**A RIGOROUS DERIVATION FROM THE KINETIC  
CUCKER-SMALE  
MODEL TO THE PRESSURELESS EULER SYSTEM  
WITH NONLOCAL ALIGNMENT**



# A RIGOROUS DERIVATION FROM THE KINETIC CUCKER–SMALE MODEL TO THE PRESSURELESS EULER SYSTEM WITH NONLOCAL ALIGNMENT

ALESSIO FIGALLI AND MOON-JIN KANG

We consider the kinetic Cucker–Smale model with local alignment as a mesoscopic description for the flocking dynamics. The local alignment was first proposed by Karper, Mellet and Trivisa (2014), as a singular limit of a normalized nonsymmetric alignment introduced by Motsch and Tadmor (2011). The existence of weak solutions to this model was obtained by Karper, Mellet and Trivisa (2014), and in the same paper they showed the time-asymptotic flocking behavior. Our main contribution is to provide a rigorous derivation from a mesoscopic to a macroscopic description for the Cucker–Smale flocking models. More precisely, we prove the hydrodynamic limit of the kinetic Cucker–Smale model with local alignment towards the pressureless Euler system with nonlocal alignment, under a regime of strong local alignment. Based on the relative entropy method, a main difficulty in our analysis comes from the fact that the entropy of the limit system has no strict convexity in terms of density variable. To overcome this, we combine relative entropy quantities with the 2-Wasserstein distance.

## 1. Introduction

This article is mainly devoted to providing a rigorous justification of the hydrodynamic limit of the kinetic Cucker–Smale model to the pressureless Euler system with nonlocal alignment force. Cucker and Smale [2007] introduced an agent-based model capturing a flocking phenomenon observed within complex systems, such as flocks of birds, schools of fish and swarms of insects. The Cucker–Smale (CS) model has received extensive attention in the mathematical community, as well as physics, biology, engineering and social science, etc.; see for instance [Carlen et al. 2015; Cañizo et al. 2011; Carrillo et al. 2010; Duan et al. 2010; Fornasier et al. 2011; Ha et al. 2014c; 2017; Ha and Tadmor 2008; Poyato and Soler 2017; Zavlanos et al. 2011]. Motsch and Tadmor [2011] proposed a modified Cucker–Smale model by replacing the original CS alignment by a normalized nonsymmetric alignment. Karper, Mellet, and Trivisa [Karper et al. 2014] proposed a new kinetic flocking model as a combination of the CS alignment and a local alignment interaction, where the latter was obtained as a singular limit of the nonsymmetric alignment introduced by Motsch and Tadmor.

---

The work of Figalli is supported by the ERC grant “Regularity and stability in partial differential equations (RSPDE)”. The work of Kang was supported by the NRF grant NRF-2017R1C1B5076510 and Sookmyung Women’s University Research Grant (1-1703-2045).

*MSC2010:* primary 35Q70; secondary 35B25.

*Keywords:* hydrodynamic limit, kinetic Cucker–Smale model, local alignment, pressureless Euler system, relative entropy, Wasserstein distance.

We consider the kinetic flocking model without Brownian noise, proposed by Karper, Mellet and Trivisa [Karper et al. 2013] on  $\mathbb{T}^d \times \mathbb{R}^d$ :

$$\begin{aligned} \partial_t f + v \cdot \nabla_x f + \nabla_v \cdot (L[f]f) + \nabla_v \cdot ((u - v)f) &= 0, \\ L[f](t, x, v) &= \int_{\mathbb{T}^d} \int_{\mathbb{R}^d} \psi(x - y) f(t, y, w) (w - v) dw dy, \\ u(t, x) &= \frac{\int_{\mathbb{R}^d} v f dv}{\int_{\mathbb{R}^d} f dv}, \quad \|f(0)\|_{L^1(\mathbb{T}^d \times \mathbb{R}^d)} = 1. \end{aligned} \tag{1-1}$$

Here  $\psi : \mathbb{T}^d \rightarrow \mathbb{R}^d$  is a Lipschitz communication weight that is positive and symmetric, i.e.,  $\psi(x - y) = \psi(y - x)$ . The term  $\nabla_v \cdot (L[f]f)$  describes a nonlocal alignment due to the original Cucker–Smale flocking mechanism, while the last term  $\nabla_v \cdot ((u - v)f)$  describes a local alignment interaction, because of the averaged local velocity  $u$ . The global existence of weak solutions to (1-1) was proved in [Karper et al. 2013]. The flocking behaviors of (1-1), however, have not been studied so far. We here provide its time-asymptotic behavior.

As a mesoscopic description, the kinetic model (1-1) is posed in  $(t, x, v) \in \mathbb{R} \times \mathbb{T}^d \times \mathbb{R}^d$ , i.e., in  $2d + 1$  dimensions. This feature provides an accurate description for a significant number of particles. However, its numerical test is very costly with respect to an associated macroscopic description. Hence, it is very important to find a suitable parameter regime on which the complexity of (1-1) is reduced.

The main goal of this article is to show a singular limit of (1-1) in a regime of strong local alignment:

$$\begin{aligned} \partial_t f^\varepsilon + v \cdot \nabla_x f^\varepsilon + \nabla_v \cdot (L[f^\varepsilon]f^\varepsilon) + \frac{1}{\varepsilon} \nabla_v \cdot ((u^\varepsilon - v)f^\varepsilon) &= 0, \\ L[f^\varepsilon](t, x, v) &= \int_{\mathbb{T}^d} \int_{\mathbb{R}^d} \psi(x - y) f^\varepsilon(t, y, w) (w - v) dw dy, \\ u^\varepsilon &= \frac{\int_{\mathbb{R}^d} v f^\varepsilon dv}{\int_{\mathbb{R}^d} f^\varepsilon dv}, \\ f^\varepsilon|_{t=0} &= f_0^\varepsilon, \quad \|f_0^\varepsilon\|_{L^1(\mathbb{T}^d \times \mathbb{R}^d)} = 1. \end{aligned} \tag{1-2}$$

As  $\varepsilon \rightarrow 0$ , it is expected that the solution  $f^\varepsilon$  of (1-2) converges, in some weak sense, to a monokinetic distribution

$$\delta_{v=u(t,x)} \otimes \rho(t, x); \tag{1-3}$$

see Remark 1.1. Here,  $\delta_{v=u(t,x)}$  denotes a Dirac mass in  $v$  centered on  $u(t, x)$ . Also, as we shall explain later, at least formally  $\rho$  and  $u$  should solve the associated limit system given by the pressureless Euler system with nonlocal flocking dissipation:

$$\begin{aligned} \partial_t \rho + \nabla \cdot (\rho u) &= 0, \\ \partial_t (\rho u) + \nabla \cdot (\rho u \otimes u) &= \int_{\mathbb{T}^d} \psi(x - y) \rho(t, x) \rho(t, y) (u(t, y) - u(t, x)) dy, \\ \rho|_{t=0} &= \rho_0, \quad u|_{t=0} = u_0, \quad \|\rho_0\|_{L^1(\mathbb{T}^d)} = 1. \end{aligned} \tag{1-4}$$

The main difficulty in the justification of this limit comes from the singularity of the monokinetic distribution. To the best of our knowledge, there is no general method to handle the hydrodynamic limit

from some kinetic equations to the pressureless Euler systems, no matter what regime is considered. Indeed, there are few results on this kinds of limit; see [Jabin and Rey 2017; Kang 2018; Kang and Vasseur 2015] (see also [Jabin 2000] for a general treatment of similar regimes that lead to the Dirac formation and pressureless gases equations).

**Remark 1.1.** In this paper we will use the symbol  $\otimes$  in two different contexts: if  $\mu$  is a measure on a complete metric space  $X$ , and  $\{\nu_x\}_{x \in X}$  is a family of measures on a complete metric space  $Y$ , then  $\nu_x \otimes \mu$  denotes the measure on  $X \times Y$  defined as

$$\int_{X \times Y} \varphi d[\nu_x \otimes \mu] = \int_X \left( \int_Y \varphi(x, y) d\nu_x(y) \right) d\mu(x) \quad \text{for all } \varphi \in C_c(X \times Y).$$

When  $\nu_x$  is independent of  $x$  (that is,  $\nu_x = \nu$  for all  $x$ ), we use the more standard notation  $\mu \otimes \nu$  (instead of  $\nu \otimes \mu$ , as done before) to denote the product measure:

$$\int_{X \times Y} \varphi d[\mu \otimes \nu] = \int_X \left( \int_Y \varphi(x, y) d\nu(y) \right) d\mu(x) \quad \text{for all } \varphi \in C_c(X \times Y).$$

Finally, if  $a, b \in \mathbb{R}^d$  are vectors, then  $a \otimes b$  denotes the  $(d \times d)$ -matrix with entries

$$(a \otimes b)_{ij} = a_i b_j \quad \text{for all } i, j = 1, \dots, d.$$

The meaning will always be clear from the context.

It is worth mentioning that the pressureless Euler system without the nonlocal alignment has been used for the formation of large-scale structures in astrophysics and the aggregation of sticky particles [Silk et al. 1983; Zeldovich 1970]. For more theoretical studies on the pressureless gases, we for example refer to [Bouchut 1994; Bouchut and James 1999; Boudin 2000; Brenier and Grenier 1998; Huang and Wang 2001; Poupaud and Rascle 1997; Weinan et al. 1996].

The macroscopic flocking model (1-4) or its variants have been formally derived under a monokinetic ansatz (1-3), and studied in various topics; see for example [Do et al. 2018; Ha et al. 2014a; 2014b; 2015; Tadmor and Tan 2014]. In [Ha et al. 2014b], the authors showed the global well-posedness of (1-4) with suitably smooth and small initial data, and the time-asymptotic flocking behavior. In [Ha et al. 2015], the authors dealt with a moving boundary problem of (1-4) with compactly supported initial density. We also refer to [Ha et al. 2014a] for a reformulation of (1-4) into hyperbolic conservation laws with damping in one dimension.

In [Karper et al. 2015], the authors showed the hydrodynamic limit of the kinetic flocking model (1-1) with Brownian motion, that is, a Vlasov-Fokker-Planck-type equation, under the regime of strong local alignment and strong Brownian motion:

$$\partial_t f^\varepsilon + v \cdot \nabla_x f^\varepsilon + \nabla_v \cdot (L[f^\varepsilon] f^\varepsilon) + \frac{1}{\varepsilon} \nabla_v \cdot ((u^\varepsilon - v) f^\varepsilon) - \frac{1}{\varepsilon} \Delta_v f^\varepsilon = 0. \tag{1-5}$$

In this case, as  $\varepsilon \rightarrow 0$ ,  $f^\varepsilon$  converges to a smooth local equilibrium given by a local Maxwellian, contrary to (1-3). There, the authors used the relative entropy method, heavily relying on a strict convexity of the

entropy of the isothermal Euler system (as a limit system of (1-5)):

$$\begin{aligned} \partial_t \rho + \nabla \cdot (\rho u) &= 0, \\ \partial_t(\rho u) + \nabla \cdot (\rho u \otimes u) + \nabla \rho &= \int_{\mathbb{T}^d} \psi(x-y) \rho(t,x) \rho(t,y) (u(t,y) - u(t,x)) dy. \end{aligned}$$

The relative entropy method based on a strict convex entropy has been successfully used to prove the hydrodynamic limit of Vlasov–Fokker–Planck-type equations; we refer to [Berthelin and Vasseur 2005; Carrillo et al. 2016; Goudon et al. 2004; Mellet and Vasseur 2008; Vasseur 2008].

On the other hand, the pressureless Euler system (1-4) has a convex entropy given by

$$\eta(\rho, \rho u) = \rho \frac{1}{2}(|u|^2), \quad (1-6)$$

which is not strictly convex with respect to  $\rho$ . For this reason, the associated relative entropy (1-6) is not enough to control the convergence of the nonlocal alignment term (compare with [Kang and Vasseur 2015], where the nonlocal alignment is not present). To overcome this difficulty, we first estimate an  $L^2$ -distance of characteristics generated by vector fields  $u^\varepsilon$  and  $u$  that controls the 2-Wasserstein distance of densities, and then combine the estimates of the relative entropy and the  $L^2$ -distance of characteristics.

As a related work on (1-5), we refer to [Carrillo et al. 2016], where the authors studied the flocking behavior and hydrodynamic limit of a coupled system of (1-5) and fluid equations via drag force.

The rest of this paper is organized as follows. In Section 2, we mention different scales of Cucker–Smale models from a microscopic level to a macroscopic level, and then specify some known existence results on the two descriptions (1-1) and (1-4). In Section 3, we present our main theorem on the hydrodynamic limit, and collect some useful results on the relative entropy method and the optimal transportation theory that are used in the proof of the main theorem. In Section 4, we present some structural hypotheses to guarantee the hydrodynamic limit in a general setting. Then we apply the general result to our systems by verifying the hypotheses in Section 5. In the Appendix, we provide the proof of the long time-asymptotic flocking dynamics and the existence of monokinetic solutions for the kinetic model (1-1).

## 2. Various scales of Cucker–Smale models

We first present various scales of Cucker–Smale models, from a microscopic level to a macroscopic level. Then we state some known results on global existence of weak solutions to the kinetic description (1-1), and local existence of smooth solutions to the limit system (1-4). Those results are crucially used in the proof of the main theorem. Finally, in Theorem 2.2, we present the time-asymptotic flocking behavior of the kinetic model (1-1).

**Variants of Cucker–Smale models.** We briefly present the kinetic CS model and its variants. Cucker and Smale [2007] proposed a mathematical model to explain the flocking phenomenon:

$$\begin{aligned} \frac{dx_i}{dt} &= v_i, \quad i = 1, \dots, N, \\ \frac{dv_i}{dt} &= \frac{1}{N} \sum_{j=1}^N \psi(x_j - x_i)(v_j - v_i), \end{aligned} \quad (2-1)$$

where  $x_i, v_i \in \mathbb{R}^d$  denote the spatial position and velocity of the  $i$ -th particle for an ensemble of  $N$  self-propelled particles. The kernel  $\psi(|x_j - x_i|)$  is a communication weight given by

$$\psi(x_j - x_i) = \frac{\lambda}{(1 + |x_j - x_i|^2)^\beta}, \quad \beta \geq 0, \lambda > 0. \quad (2-2)$$

The system (2-1) with (2-2) was used as an analytical description of the Vicsek model [Vicsek et al. 1995] without resorting to the first principle of physics.

When the number of particles is sufficiently large, the ensemble of particles can be described by the one-particle density function  $f = f(t, x, v)$  at the spatial-velocity position  $(x, v) \in \mathbb{R}^d \times \mathbb{R}^d$  at time  $t$ . Then, the evolution of  $f$  is governed by the following Vlasov-type equation:

$$\begin{aligned} \partial_t f + v \cdot \nabla_x f + \nabla_v \cdot (L[f]f) &= 0, \\ L[f](t, x, v) &= \int_{\mathbb{R}^{2d}} \psi(x - y) f(t, y, w) (w - v) dw dy. \end{aligned} \quad (2-3)$$

This was first introduced by Ha and Tadmor [2008] using the BBGKY hierarchy from the particle CS model (2-1). A rigorous mean-field limit was given in [Ha and Liu 2009].

Motsch and Tadmor [2011] recognized a drawback of the CS model (2-1), which is due to the normalization factor  $1/N$ . More precisely, when a small group of agents are located far away from a much larger group of agents, the internal dynamics of the small group is almost halted since the total number of agents is relatively very large. To solve this issue, they replaced the nonlocal alignment  $L[f]$  by a normalized nonsymmetric alignment operator:

$$\bar{L}[f](t, x, v) := \frac{\int_{\mathbb{R}^{2d}} K^r(x - y) f(t, y, w) (w - v) dw dy}{\int_{\mathbb{R}^{2d}} K^r(x - y) f(t, y, w) dw dy},$$

where the kernel  $K^r$  is a communication weight and  $r$  denotes the radius of influence of  $K^r$ .

In [Karper et al. 2014], the authors considered the case when the communication weight is extremely concentrated near each agent, so that the alignment term  $\bar{L}[f]$  corresponds to a short-range interaction. More precisely, they rigorously justified the singular limit  $r \rightarrow 0$ , i.e., as  $K^r$  converges to the Dirac distribution  $\delta_0$ , in which case  $\bar{L}[f]$  converges to a local alignment term:

$$\bar{L}[f](t, x, v) \rightarrow \frac{\int_{\mathbb{R}^d} f(t, x, w) (w - v) dw}{\int_{\mathbb{R}^d} f(t, x, w) dw} = u(t, x) - v,$$

where  $u(t, x)$  denotes the averaged local velocity defined as

$$u(t, x) = \frac{\int_{\mathbb{R}^d} v f(t, x, v) dv}{\int_{\mathbb{R}^d} f(t, x, v) dv}.$$

Hence, their new model became (1-1), which consists of two kinds of alignment force: a nonlocal alignment due to the original CS model, plus a local alignment.

**Existence of weak solutions to (1-2).** In [Karper et al. 2013], the authors showed the existence of weak solutions to the kinetic Cucker–Smale model with local alignment, noise, self-propulsion, and friction:

$$\begin{aligned} \partial_t f + v \cdot \nabla_x f + \nabla_v \cdot (L[f]f) + \nabla_v \cdot ((u - v)f) &= \sigma \Delta_v f - \nabla_v \cdot ((a - b|v|^2)vf), \\ L[f] &= \int_{\mathbb{R}^{2d}} \psi(x - y)f(t, y, w)(w - v) dw dy, \end{aligned} \tag{2-4}$$

where the kernel  $\psi$  is the same as (1-2) and  $a, b$ , and  $\sigma$  are nonnegative constants. By their result applied with  $a = b = \sigma = 0$  inside the periodic domain  $\mathbb{T}^d$ , we obtain existence of solutions for (1-2). To precisely state such an existence result, we need to define a (mathematical) entropy  $\mathcal{F}(f^\varepsilon)$  and kinetic dissipation  $\mathcal{D}_1(f^\varepsilon), \mathcal{D}_2(f^\varepsilon)$  for (1-2):

$$\begin{aligned} \mathcal{F}(f^\varepsilon) &:= \int_{\mathbb{R}^d} \frac{|v|^2}{2} f^\varepsilon dv, \\ \mathcal{D}_1(f^\varepsilon) &:= \int_{\mathbb{T}^d \times \mathbb{R}^d} f^\varepsilon |u^\varepsilon - v|^2 dv dx, \\ \mathcal{D}_2(f^\varepsilon) &:= \frac{1}{2} \int_{\mathbb{T}^{2d} \times \mathbb{R}^{2d}} \psi(x - y) f^\varepsilon(x, v) f^\varepsilon(y, w) |v - w|^2 dx dy dv dw. \end{aligned} \tag{2-5}$$

**Proposition 2.1.** *For any  $\varepsilon > 0$ , assume that  $f_0^\varepsilon$  satisfies*

$$f_0^\varepsilon \geq 0, \quad f_0^\varepsilon \in L^1 \cap L^\infty(\mathbb{R}^{2d}), \quad |v|^2 f_0^\varepsilon \in L^1(\mathbb{R}^{2d}). \tag{2-6}$$

*Then there exists a weak solution  $f^\varepsilon \geq 0$  of (1-2) such that*

$$\begin{aligned} f^\varepsilon &\in C(0, T; L^1(\mathbb{R}^{2d})) \cap L^\infty((0, T) \times \mathbb{R}^{2d}), \\ |v|^2 f^\varepsilon &\in L^\infty(0, T; L^1(\mathbb{R}^{2d})), \end{aligned} \tag{2-7}$$

*and (1-2) holds in the sense of distributions, that is, for any  $\varphi \in C_c^\infty([0, T) \times \mathbb{R}^{2d})$ , the weak formulation holds:*

$$\int_0^t \int_{\mathbb{R}^{2d}} f^\varepsilon \left( \partial_t \varphi + v \cdot \nabla_x \varphi + L[f^\varepsilon] \cdot \nabla_v \varphi + \frac{1}{\varepsilon} (u^\varepsilon - v) \cdot \nabla_v \varphi \right) dv dx ds + \int_{\mathbb{R}^{2d}} f_0^\varepsilon \varphi(0, \cdot) dv dx = 0. \tag{2-8}$$

*Moreover,  $f^\varepsilon$  preserves the total mass and satisfies the entropy inequality*

$$\int_{\mathbb{T}^d} \mathcal{F}(f^\varepsilon)(t) dx + \frac{1}{\varepsilon} \int_0^t \mathcal{D}_1(f^\varepsilon)(s) ds + \int_0^t \mathcal{D}_2(f^\varepsilon)(s) ds \leq \int_{\mathbb{T}^d} \mathcal{F}(f_0^\varepsilon) dx. \tag{2-9}$$

The entropy inequality (2-9) is crucially used in the proof of Theorem 3.1.

**Flocking behavior of the kinetic model (1-1).** We now present the time-asymptotic flocking behavior of solutions to the kinetic model (1-1). For that, we define the following two Lyapunov functionals:

$$\begin{aligned} \mathcal{E}_1(t) &:= \int_{\mathbb{T}^d \times \mathbb{R}^d} f(t, x, v) |u(t, x) - v|^2 dv dx, \\ \mathcal{E}_2(t) &:= \int_{\mathbb{T}^{2d}} \rho(t, x) \rho(t, y) |u(t, x) - u(t, y)|^2 dx dy, \end{aligned}$$



where  $\rho(t, x) = \int_{\mathbb{R}^d} f(t, x, v) dv$ . We remark that  $\mathcal{E}_1$  measures a local alignment, and  $\mathcal{E}_2$  measures alignment of the averaged local velocities. Then, for the flocking estimate, we combine the two functionals as follows:

$$\mathcal{E}(t) := \mathcal{E}_1(t) + \frac{1}{2}\mathcal{E}_2(t). \tag{2-10}$$

**Theorem 2.2.** *Let  $f$  be a solution to (1-1). Then, we have the time-asymptotic flocking estimate*

$$\mathcal{E}(t) \leq \mathcal{E}(0) \exp(-2 \min\{1, \psi_m\}t), \quad t > 0, \tag{2-11}$$

where  $\psi_m$  is the minimum communication weight:

$$\psi_m := \min_{x, y \in \mathbb{T}^d} \psi(x - y) > 0.$$

In addition, if  $u$  is uniformly Lipschitz continuous on a time interval  $[0, T]$ , namely

$$\ell_T := \sup_{t \in [0, T]} \|\nabla_x u\|_{L^\infty(\mathbb{T}^d)} < \infty,$$

then

$$\mathcal{E}_1(t) \leq \mathcal{E}_1(0)e^{2(\ell_T-1)t} \quad \text{for all } t \in [0, T]. \tag{2-12}$$

*Proof.* We postpone the proof to the Appendix. □

**Remark 2.3.** As an interesting consequence of (2-12) one obtains that, for smooth solutions,  $\mathcal{E}_1(0) = 0$  implies that  $\mathcal{E}_1(t) = 0$  for all  $t \in [0, T]$ . In other words, monokinetic initial conditions remain monokinetic as long as the velocity field is Lipschitz. One can note that monokinetic solutions to (1-1) simply correspond to solutions of the pressureless Euler system (1-4); hence the short time existence of Lipschitz solutions is guaranteed by Proposition 2.4 and Remark 2.5 below.

**Formal derivation of the hydrodynamic Cucker–Smale system (1-4).** We consider the hydrodynamic variables  $\rho^\varepsilon := \int_{\mathbb{R}^d} f^\varepsilon dv$  and  $\rho^\varepsilon u^\varepsilon := \int_{\mathbb{R}^d} v f^\varepsilon dv$ .

First of all, integrating (1-2) with respect to  $v$ , we get the continuity equation

$$\partial_t \rho^\varepsilon + \nabla_x \cdot (\rho^\varepsilon u^\varepsilon) = 0.$$

Multiplying (1-2) by  $v$ , and then integrating it with respect to  $v$ , we have

$$\partial_t (\rho^\varepsilon u^\varepsilon) + \nabla_x \cdot \left( \int_{\mathbb{R}^d} v \otimes v f^\varepsilon dv \right) = \int_{\mathbb{T}^d} \psi(x - y) \rho^\varepsilon(t, x) \rho^\varepsilon(t, y) (u^\varepsilon(t, y) - u^\varepsilon(t, x)) dy,$$

where we used

$$u^\varepsilon = \frac{\int_{\mathbb{R}^d} v f^\varepsilon dv}{\int_{\mathbb{R}^d} f^\varepsilon dv}.$$

Then, we rewrite the system for  $\rho^\varepsilon$  and  $u^\varepsilon$  as

$$\begin{aligned} \partial_t \rho^\varepsilon + \nabla_x \cdot (\rho^\varepsilon u^\varepsilon) &= 0, \\ \partial_t (\rho^\varepsilon u^\varepsilon) + \nabla_x \cdot (\rho^\varepsilon u^\varepsilon \otimes u^\varepsilon + P^\varepsilon) &= \int_{\mathbb{T}^d} \psi(x - y) \rho^\varepsilon(t, x) \rho^\varepsilon(t, y) (u^\varepsilon(t, y) - u^\varepsilon(t, x)) dy, \end{aligned} \tag{2-13}$$

where  $P^\varepsilon$  is the stress tensor given by

$$P^\varepsilon := \int_{\mathbb{R}^d} (v - u^\varepsilon) \otimes (v - u^\varepsilon) f^\varepsilon dv.$$

If we take  $\varepsilon \rightarrow 0$  in (1-2), the local alignment term  $\nabla_v \cdot ((u^\varepsilon - v) f^\varepsilon)$  converges to 0. Hence, if  $\rho^\varepsilon \rightarrow \rho$  and  $\rho^\varepsilon u^\varepsilon \rightarrow \rho u$  for some limiting functions  $\rho$  and  $u$ , we have that  $f^\varepsilon \rightarrow \delta_{v=u} \otimes \rho$  (in some suitable sense). Hence, the stress tensor  $P^\varepsilon$  should vanish in the limit, since

$$\int_{\mathbb{R}^d} (v - u) \otimes (v - u) \delta_{v=u} \rho dv = 0.$$

Therefore, at least formally, the limit quantities  $\rho$  and  $u$  satisfy the pressureless Euler system with nonlocal alignment:

$$\begin{aligned} \partial_t \rho + \nabla_x \cdot (\rho u) &= 0, \\ \partial_t (\rho u) + \nabla_x \cdot (\rho u \otimes u) &= \int_{\mathbb{T}^d} \psi(x - y) \rho(t, x) \rho(t, y) (u(t, y) - u(t, x)) dy. \end{aligned}$$

**Existence of classical solutions to (1-4).** We present here the local existence of classical solutions to the pressureless Euler system (1-4).

**Proposition 2.4.** *Assume that*

$$\rho_0 > 0 \quad \text{in } \mathbb{T}^d \quad \text{and} \quad (\rho_0, u_0) \in H^s(\mathbb{T}^d) \times H^{s+1}(\mathbb{T}^d) \quad \text{for } s > \frac{1}{2}d + 1. \quad (2-14)$$

*Then, there exists  $T_* > 0$  such that (1-4) has a unique classical solution  $(\rho, u)$  satisfying*

$$\begin{aligned} \rho &\in C^0([0, T_*]; H^s(\mathbb{T}^d)) \cap C^1([0, T_*]; H^{s-1}(\mathbb{T}^d)), \\ u &\in C^0([0, T_*]; H^{s+1}(\mathbb{T}^d)) \cap C^1([0, T_*]; H^s(\mathbb{T}^d)). \end{aligned} \quad (2-15)$$

**Remark 2.5.** Since  $s > \frac{1}{2}d + 1$ , by the Sobolev inequality it follows that  $(\rho, u) \in C^1([0, T_*] \times \mathbb{T}^d)$ .

Proposition 2.4 has been proven in [Ha et al. 2014b]. There, the authors obtained also a global well-posedness of classical solutions, provided an initial datum is suitably smooth and small.

### 3. Main result and preliminaries

We first present our main result on the hydrodynamic limit of (1-2). We next present useful results on the relative entropy method and the optimal transportation theory, which are used as main tools in the next section.

**Main result.** For the hydrodynamic limit, we consider a well-prepared initial data  $f_0^\varepsilon$  satisfying (2-6) and

$$(A1) \quad \int_{\mathbb{T}^d} \int_{\mathbb{R}^d} (f_0^\varepsilon \frac{1}{2}|v|^2 - \rho_0 \frac{1}{2}|u_0|^2) dv dx = \mathcal{O}(\varepsilon),$$

$$(A2) \quad \|\rho_0^\varepsilon - \rho_0\|_{L^1(\mathbb{T}^d)} = \mathcal{O}(\varepsilon),$$

$$(A3) \quad \|u_0^\varepsilon - u_0\|_{L^\infty(\mathbb{T}^d)} = \mathcal{O}(\varepsilon).$$

We now specify our main result on the hydrodynamic limit.

**Theorem 3.1.** *Assume that the initial data  $f_0^\varepsilon$  and  $(\rho_0, u_0)$  satisfy (2-6), (2-14), and (A1)–(A3). Let  $f^\varepsilon$  be a weak solution to (1-2) satisfying (2-9), and  $(\rho, u)$  be a local-in-time smooth solution to (1-4) satisfying (2-15) up to the time  $T_*$ . Then, there exists a positive constant  $C_*$  (depending on  $T_*$ ) such that, for all  $t \leq T_*$ ,*

$$\int_{\mathbb{T}^d} \rho^\varepsilon(t) |(u^\varepsilon - u)|^2(t) dx + W_2^2(\rho^\varepsilon(t), \rho(t)) \leq C_* \varepsilon, \tag{3-1}$$

where  $\rho^\varepsilon = \int_{\mathbb{R}^d} f^\varepsilon dv$ ,  $\rho^\varepsilon u^\varepsilon = \int_{\mathbb{R}^d} v f^\varepsilon dv$ , and  $W_2$  denotes the 2-Wasserstein distance.

Therefore, we have

$$f^\varepsilon \rightharpoonup \delta_{v=u(t,x)} \otimes \rho(t, x) \quad \text{in } \mathcal{M}((0, T_*) \times \mathbb{T}^d \times \mathbb{R}^d), \tag{3-2}$$

where  $\mathcal{M}((0, T_*) \times \mathbb{T}^d \times \mathbb{R}^d)$  is the space of nonnegative Radon measures on  $(0, T_*) \times \mathbb{T}^d \times \mathbb{R}^d$ .

The proof of this result is postponed to Section 5. In the next subsections we collect some preliminary facts that will be used later in the proof.

**Relative entropy method.** First of all, we rewrite the limit system (1-4) in an abstract form using the notation

$$\begin{aligned} P &= \rho u, & U &= \begin{pmatrix} \rho \\ P \end{pmatrix}, & A(U) &= \begin{pmatrix} P^T \\ (P \otimes P)/\rho \end{pmatrix}, \\ F(U) &= \begin{pmatrix} 0 \\ \int_{\mathbb{T}^d} \psi(x-y) \rho(t, x) \rho(t, y) (u(t, y) - u(t, x)) dy \end{pmatrix}. \end{aligned}$$

Then we can rewrite (1-4) as the balance law

$$\partial_t U + \operatorname{div}_x A(U) = F(U). \tag{3-3}$$

We consider the relative entropy and relative flux

$$\begin{aligned} \eta(V | U) &= \eta(V) - \eta(U) - D\eta(U) \cdot (V - U), \\ A(V | U) &= A(V) - A(U) - DA(U) \cdot (V - U), \end{aligned} \tag{3-4}$$

where  $DA(U) \cdot (V - U)$  is a matrix defined as

$$(DA(U) \cdot (V - U))_{ij} = \sum_{k=1}^{d+1} \partial_{U_k} A_{ij}(U) (V_k - U_k), \quad 1 \leq i \leq d+1, \quad 1 \leq j \leq d.$$

By the theory of conservation laws, the system (3-3) has a convex entropy  $\eta(U) = \rho \frac{1}{2} |u|^2$  with entropy flux  $G$  given by the identity

$$\partial_{U_i} G_j(U) = \sum_{k=1}^{d+1} \partial_{U_k} \eta(U) \partial_{U_i} A_{kj}(U), \quad 1 \leq i \leq d+1, \quad 1 \leq j \leq d.$$

Since  $\eta(U) = |P|^2/(2\rho)$ , and

$$D\eta(U) = \begin{pmatrix} D_\rho \eta \\ D_P \eta \end{pmatrix} = \begin{pmatrix} -|P|^2/(2\rho^2) \\ P/\rho \end{pmatrix} = \begin{pmatrix} -|u|^2/2 \\ u \end{pmatrix}, \tag{3-5}$$

for given  $V = \begin{pmatrix} q \\ qw \end{pmatrix}$ ,  $U = \begin{pmatrix} \rho \\ \rho u \end{pmatrix}$ , we have

$$\begin{aligned} \eta(V | U) &= \frac{1}{2}q|w|^2 - \frac{1}{2}\rho|u|^2 + \frac{1}{2}|u|^2(q - \rho) - u(qw - \rho u) \\ &= \frac{1}{2}q|u - w|^2. \end{aligned} \quad (3-6)$$

The next proposition provides a cornerstone to verify the hydrodynamic limit through the relative entropy method. For its proof, we refer to the proof of Proposition 4.2 in [Karper et al. 2015]; see also [Vasseur 2008].

**Proposition 3.2.** *Let  $U$  be a strong solution to balance law (3-3) and  $V$  be any smooth function. Then, the following holds:*

$$\begin{aligned} \frac{d}{dt} \int_{\mathbb{T}^d} \eta(V | U) dx &= \frac{d}{dt} \int_{\mathbb{T}^d} \eta(V) dx - \int_{\mathbb{T}^d} \nabla_x (D\eta(U)) : A(V | U) dx \\ &\quad - \int_{\mathbb{T}^d} D\eta(U) \cdot [\partial_t V + \operatorname{div}_x A(V) - F(V)] dx \\ &\quad - \int_{\mathbb{T}^d} [D^2 \eta(U) F(U)(V - U) + D\eta(U) F(V)] dx. \end{aligned}$$

**Wasserstein distance and representation formulae for solutions of the continuity equation.** For  $p \geq 1$ , the  $p$ -Wasserstein distance between two probability measures  $\mu_1$  and  $\mu_2$  on  $\mathbb{R}^d$  is defined by

$$W_p^p(\mu_1, \mu_2) := \inf_{\nu \in \Lambda(\mu_1, \mu_2)} \int_{\mathbb{R}^{2d}} |x - y|^p d\nu(x, y),$$

where  $\Lambda(\mu_1, \mu_2)$  denotes the set of all probability measures  $\nu$  on  $\mathbb{R}^{2d}$  with marginals  $\mu_1$  and  $\mu_2$ , i.e.,

$$\pi_{1\#}\nu = \mu_1, \quad \pi_{2\#}\nu = \mu_2,$$

where  $\pi_1 : (x, y) \mapsto x$  and  $\pi_2 : (x, y) \mapsto y$  are the natural projections from  $\mathbb{R}^d \times \mathbb{R}^d$  to  $\mathbb{R}^d$ , and  $\pi_{\#}\nu$  denotes the push forward of  $\nu$  through a map  $\pi$ , i.e.,  $\pi_{\#}\nu(B) := \nu(\pi^{-1}(B))$  for any Borel set  $B$ . This same definition can be extended to measures on the torus  $\mathbb{T}^d$  with the understanding that  $|x - y|$  denotes the distance on the torus.

To make a connection between the  $L^2$ -distance of velocities and the 2-Wasserstein distance of densities (see Lemma 5.2), we will use two different representation formulas for solutions to the continuity equation

$$\partial_t \mu_t + \operatorname{div}_x (u_t \mu_t) = 0. \quad (3-7)$$

Let us recall that, if the velocity field  $u_t : \mathbb{R}^d \rightarrow \mathbb{R}^d$  is Lipschitz with respect to  $x$ , uniformly in  $t$ , then for any  $x$  there exists a global-in-time unique characteristic  $X$  generated by  $u_t$  starting from  $x$ ,

$$\dot{X}(t, x) = u_t(X(t, x)), \quad X(0, x) = x,$$

and the solution  $\mu_t$  of (3-7) is the push forward of the initial data  $\mu_0$  through  $X(t)$ , i.e.,

$$\mu_t = X(t)_{\#}\mu_0; \quad (3-8)$$

e.g., see [Ambrosio et al. 2005, Proposition 8.1.8]. On the other hand, if the velocity field  $u_t$  is not Lipschitz with respect to  $x$ , the uniqueness of the characteristics is not guaranteed anymore. Still, a probabilistic representation formula for solutions to (3-7) holds (recall that a curve of probability measures in  $\mathbb{R}^d$  is called narrowly continuous if it is continuous in the duality with continuous bounded functions):

**Proposition 3.3.** *For a given  $T > 0$ , let  $\mu_t : [0, T] \rightarrow \mathcal{P}(\mathbb{R}^d)$  be a narrowly continuous solution of (3-7) for a Borel vector field  $u_t$  satisfying*

$$\int_0^T \int_{\mathbb{R}^d} |u_t(x)|^p d\mu_t(x) dt < \infty \quad \text{for some } p > 1.$$

Let  $\Gamma_T$  denote the space of continuous curves from  $[0, T]$  into  $\mathbb{R}^d$ . Then, there exists a probability measure  $\eta$  on  $\Gamma_T \times \mathbb{R}^d$  satisfying the following properties:

- (i)  $\eta$  is concentrated on the set of pairs  $(\gamma, x)$  such that  $\gamma$  is an absolutely continuous curve solving the ODE

$$\dot{\gamma}(t) = u_t(\gamma(t)) \quad \text{for a.e. } t \in (0, T), \text{ with } \gamma(0) = x.$$

- (ii)  $\mu_t$  satisfies

$$\int_{\mathbb{R}^d} \varphi(x) d\mu_t(x) = \int_{\Gamma_T \times \mathbb{R}^d} \varphi(\gamma(t)) d\eta(\gamma, x) \quad \text{for all } \varphi \in C_b^0(\mathbb{R}^d), t \in [0, T].$$

Again, this result readily extends on the torus.

Note that, in the case when  $u_t$  is Lipschitz, there exists a unique curve  $\gamma$  solving the ODE and starting from  $x$  (i.e.,  $\gamma = X(\cdot, x)$ ), so the measure  $\eta$  is given by the formula

$$d\eta(\gamma, x) = \delta_{\gamma=X(\cdot, x)} \otimes d\mu_0(x).$$

We refer to [Ambrosio et al. 2005, Theorem 8.2.1] for more details and a proof.

**Useful inequality.** We here present a standard inequality that is used in the proof of Lemma 5.2, for the convenience of the reader:

**Lemma 3.4.** *Let  $\rho_1, \rho_2 : \mathbb{T}^d \rightarrow \mathbb{R}$  be two probability densities. Then*

$$W_2^2(\rho_1, \rho_2) \leq \frac{1}{8}d \|\rho_1 - \rho_2\|_{L^1(\mathbb{T}^d)}.$$

*Proof.* The idea is simple: to estimate the transportation cost from  $\rho_1$  to  $\rho_2$  it suffices to consider a transport plan that keeps at rest all the mass in common between  $\rho_1$  and  $\rho_2$  (namely  $\min\{\rho_1, \rho_2\}$ ) and sends  $\rho_1 - \min\{\rho_1, \rho_2\}$  onto  $\rho_2 - \min\{\rho_1, \rho_2\}$  in an arbitrary way. For instance, assuming without loss of generality that  $\rho_1 \neq \rho_2$  (otherwise the result is trivial), we set

$$m := \|\rho_1 - \min\{\rho_1, \rho_2\}\|_{L^1(\mathbb{T}^d)} = \|\rho_2 - \min\{\rho_1, \rho_2\}\|_{L^1(\mathbb{T}^d)} = \frac{1}{2} \|\rho_1 - \rho_2\|_{L^1(\mathbb{T}^d)} > 0.$$

Then, a possible choice of transport plan between  $\rho_1$  and  $\rho_2$  is given by

$$\begin{aligned} \pi(dx, dy) := & \delta_{x=y}(dy) \otimes \min\{\rho_1(x), \rho_2(x)\}dx \\ & + \frac{1}{m} [\rho_1(x) - \min\{\rho_1(x), \rho_2(x)\}]dx \otimes [\rho_2(y) - \min\{\rho_1(y), \rho_2(y)\}]dy. \end{aligned}$$

Since the diameter of  $\mathbb{T}^d$  is bounded by  $\frac{1}{2}\sqrt{d}$ , we deduce that the  $W_2^2$ -cost to transport  $\rho_1 - \min\{\rho_1, \rho_2\}$  onto  $\rho_2 - \min\{\rho_1, \rho_2\}$  is at most

$$\begin{aligned} \int_{\mathbb{T}^{2d}} |x - y|^2 d\pi(x, y) &= \frac{1}{m} \int_{\mathbb{T}^{2d}} |x - y|^2 (\rho_1(x) - \min\{\rho_1(x), \rho_2(x)\})(\rho_2(y) - \min\{\rho_1(y), \rho_2(y)\}) dx dy \\ &\leq \frac{1}{4}d \|\rho_1 - \min\{\rho_1, \rho_2\}\|_{L^1(\mathbb{T}^d)} = \frac{1}{8}d \|\rho_1 - \rho_2\|_{L^1(\mathbb{T}^d)}, \end{aligned}$$

as desired. □

### 4. Structural lemma

In a general system, we first present some structural hypotheses to provide a Gronwall-type inequality on the relative entropy that is also controlled by 2-Wasserstein distance.

**Hypotheses.** Let  $f^\varepsilon$  be a solution to a given kinetic equation  $\text{KE}_\varepsilon$  scaled with  $\varepsilon > 0$  corresponding to initial data  $f_0^\varepsilon$ . Let  $U^\varepsilon$  and  $U_0^\varepsilon$  consist of hydrodynamic variables of  $f^\varepsilon$  and  $f_0^\varepsilon$  respectively.

Let  $U$  be a solution to a balance law (as a limit system of  $\text{KE}_\varepsilon$ ):

$$\partial_t U + \text{div}_x A(U) = F(U), \quad U|_{t=0} = U_0.$$

(H1) The kinetic equation  $\text{KE}_\varepsilon$  has a kinetic entropy  $\mathcal{F}$  such that  $\int \mathcal{F}(f^\varepsilon)(t) dx \geq 0$  and

$$\int \mathcal{F}(f^\varepsilon)(t) dx + \frac{1}{\varepsilon} \int_0^t D_1(f^\varepsilon)(s) ds + \int_0^t D_2(f^\varepsilon)(s) ds \leq \int_{\mathbb{T}^d} \mathcal{F}(f_0^\varepsilon) dx,$$

where  $D_1, D_2 \geq 0$  are some dissipations.

(H2) There exists a constant  $C > 0$  (independent of  $\varepsilon$ ) such that

$$\int \eta(U_0^\varepsilon | U_0) dx \leq C\varepsilon, \quad \int (\mathcal{F}(f_0^\varepsilon) - \eta(U_0^\varepsilon)) dx \leq C\varepsilon, \quad \int_{\mathbb{T}^d} \mathcal{F}(f_0^\varepsilon) dx \leq C.$$

(H3) The balance law has a convex entropy  $\eta$ , and the following minimization property holds:

$$\eta(U^\varepsilon) \leq \mathcal{F}(f^\varepsilon).$$

(H4) There exists a constant  $C > 0$  (independent of  $\varepsilon$ ) such that

$$\left| \int \nabla_x (D\eta(U)) : A(U^\varepsilon | U) dx \right| \leq C \int \eta(U^\varepsilon | U) dx.$$

(H5) There exists a constant  $C > 0$  (independent of  $\varepsilon$ ) such that

$$\left| \int D\eta(U) \cdot [\partial_t U^\varepsilon + \text{div}_x A(U^\varepsilon) - F(U^\varepsilon)] dx \right| \leq C D_1(f^\varepsilon).$$

(H6) Let  $\rho^\varepsilon$  be the hydrodynamic variable of  $f^\varepsilon$  as the local mass, and  $\rho$  be the corresponding variable for the balance law. Then,

$$- \int [D^2 \eta(U) F(U)(U^\varepsilon - U) + D\eta(U) F(U^\varepsilon)] dx \leq D_2(f^\varepsilon) + C W_2^2(\rho^\varepsilon, \rho) + C \int \eta(U^\varepsilon | U) dx.$$

(H7) There exists a constant  $C > 0$  (independent of  $\varepsilon$ ) such that

$$W_2^2(\rho^\varepsilon, \rho)(t) \leq C \int_0^t \int \eta(U^\varepsilon | U) dx ds + C\varepsilon.$$

**Remark 4.1.** (1) The hypotheses (H1)–(H5) provide a basic structure in applying the relative entropy method to hydrodynamic limits as in previous results, for example, [Kang and Vasseur 2015; Karper et al. 2015; Mellet and Vasseur 2008]. On the other hand, the hypotheses (H6)–(H7) provide a crucial connection between the relative entropy and Wasserstein distance.

(2) The (kinetic) entropy inequality (H1) plays an important role in controlling the dissipations  $D_1, D_2$  in (H5) and (H6).

(3) (H2) is related to a kind of well-prepared initial data.

**Lemma 4.2.** *Assume the hypotheses (H1)–(H7). Then, for a given  $T > 0$ , there exists a constant  $C > 0$  such that*

$$\int \eta(U^\varepsilon | U)(t) dx + W_2^2(\rho^\varepsilon, \rho)(t) \leq C\varepsilon, \quad t \leq T.$$

*Proof.* First of all, using Proposition 3.2, we have

$$\int_{\mathbb{T}^d} \eta(U^\varepsilon | U)(t) dx \leq I_1 + I_2 + I_3 + I_4 + I_5,$$

where

$$I_1 := \int_{\mathbb{T}^d} \eta(U_0^\varepsilon | U_0) dx,$$

$$I_2 := \int_{\mathbb{T}^d} (\eta(U^\varepsilon)(t) - \eta(U_0^\varepsilon)) dx,$$

$$I_3 := - \int_0^t \int_{\mathbb{T}^d} \nabla_x (D\eta(U)) : A(U^\varepsilon | U) dx ds,$$

$$I_4 := - \int_0^t \int_{\mathbb{T}^d} D\eta(U) \cdot [\partial_t U^\varepsilon + \operatorname{div}_x A(U^\varepsilon) - F(U^\varepsilon)] dx ds,$$

$$I_5 := - \int_0^t \int_{\mathbb{T}^d} [D^2 \eta(U) F(U) (U^\varepsilon - U) + D\eta(U) F(U^\varepsilon)] dx ds.$$

It follows from (H2) that  $I_1 \leq C\varepsilon$ .

We decompose  $I_2$  as

$$I_2 = \underbrace{\int_{\mathbb{T}^d} (\eta(U^\varepsilon)(t) - \mathcal{F}(f^\varepsilon)(t)) dx}_{=: I_2^1} + \underbrace{\int_{\mathbb{T}^d} (\mathcal{F}(f^\varepsilon)(t) - \mathcal{F}(f_0^\varepsilon)) dx}_{=: I_2^2} + \underbrace{\int_{\mathbb{T}^d} (\mathcal{F}(f_0^\varepsilon) - \eta(U_0^\varepsilon)) dx}_{=: I_2^3}. \quad (4-1)$$

First,  $I_2^1 \leq 0$  by (H3).

Since (H1) yields

$$I_2^2 \leq - \int_0^t D_2(f^\varepsilon) ds,$$

it follows from (H6) that

$$I_2^2 + I_5 \leq C \int_0^t W_2^2(\rho^\varepsilon, \rho) ds + C \int_0^t \int_{\mathbb{T}^d} \eta(U^\varepsilon | U) dx ds.$$

By (H2),  $I_2^3 \leq C\varepsilon$ .

It follows from (H4) that

$$I_3 \leq C \int_0^t \int_{\mathbb{T}^d} \eta(U^\varepsilon | U) dx ds.$$

Since (H1) and (H2) imply

$$\int_0^t D_1(f^\varepsilon)(s) ds \leq C\varepsilon,$$

we have  $I_4 \leq C\varepsilon$ .

Therefore, we have

$$\int \eta(U^\varepsilon | U)(t) dx \leq C\varepsilon + C \int_0^t \left[ \int \eta(U^\varepsilon | U)(s) dx ds + W_2^2(\rho^\varepsilon, \rho) \right] ds.$$

Hence, combining it with (H7), and using Gronwall's inequality, we have the desired result.  $\square$

## 5. Proof of Theorem 3.1

The main part of the proof consists in proving the estimate (3-1).

**Proof of (3-1).** This will be done by verifying the hypotheses (H1)–(H7), and then completed by Lemma 4.2.

*Verification of (H1).* (H1) is satisfied thanks to Lemma 5.1 below. There we show that one can replace the nonlocal dissipation  $\mathcal{D}_2$  in the kinetic entropy inequality (2-9) by another dissipation  $\tilde{\mathcal{D}}_2$  defined in terms of the hydrodynamic variables  $\rho^\varepsilon$  and  $u^\varepsilon$ .

**Lemma 5.1.** *For any  $\varepsilon > 0$ , assume that  $f_0^\varepsilon$  satisfies*

$$f_0^\varepsilon \in L^1 \cap L^\infty(\mathbb{T}^d \times \mathbb{R}^d), \quad |v|^2 f_0^\varepsilon \in L^1(\mathbb{T}^d \times \mathbb{R}^d).$$

*Then the weak solution  $f^\varepsilon$  in Proposition 2.1 also satisfies*

$$\int_{\mathbb{T}^d} \mathcal{F}(f^\varepsilon)(t) dx + \frac{1}{\varepsilon} \int_0^t \mathcal{D}_1(f^\varepsilon)(s) ds + \int_0^t \tilde{\mathcal{D}}_2(f^\varepsilon)(s) ds \leq \int_{\mathbb{T}^d} \mathcal{F}(f_0^\varepsilon) dx, \quad (5-1)$$

where  $\mathcal{F}$  and  $\mathcal{D}_1$  are as in (2-5), and

$$\tilde{\mathcal{D}}_2(f^\varepsilon) := \frac{1}{2} \int_{\mathbb{T}^{2d}} \psi(x-y) \rho^\varepsilon(t, x) \rho^\varepsilon(t, y) |u^\varepsilon(t, x) - u^\varepsilon(t, y)|^2 dx dy.$$



*Proof.* Recalling (2-9), it is enough to show  $\tilde{\mathcal{D}}_2(f^\varepsilon) \leq \mathcal{D}_2(f^\varepsilon)$ . We first rewrite  $\tilde{\mathcal{D}}_2(f^\varepsilon)$  in terms of the mesoscopic variables as follows: using  $\psi(x-y) = \psi(y-x)$ , we have

$$\begin{aligned} \tilde{\mathcal{D}}_2(f^\varepsilon) &= \frac{1}{2} \int_{\mathbb{T}^{2d} \times \mathbb{R}^{2d}} \psi(x-y) f^\varepsilon(t, x, v) f^\varepsilon(t, y, w) (v-w) \cdot (u^\varepsilon(t, x) - u^\varepsilon(t, y)) dv dw dx dy \\ &= \int_{\mathbb{T}^{2d} \times \mathbb{R}^{2d}} \psi(x-y) f^\varepsilon(t, x, v) f^\varepsilon(t, y, w) (v-w) \cdot u^\varepsilon(t, x) dv dw dx dy \\ &= \underbrace{\int_{\mathbb{T}^{2d} \times \mathbb{R}^{2d}} \psi(x-y) f^\varepsilon(t, x, v) f^\varepsilon(t, y, w) (v-w) \cdot v dv dw dx dy}_{=: \mathcal{I}_1} \\ &\quad + \underbrace{\int_{\mathbb{T}^{2d} \times \mathbb{R}^{2d}} \psi(x-y) f^\varepsilon(t, x, v) f^\varepsilon(t, y, w) (v-w) \cdot (u^\varepsilon(t, x) - v) dv dw dx dy}_{=: \mathcal{I}_2}. \end{aligned}$$

First, we have

$$\mathcal{I}_1 = \frac{1}{2} \int_{\mathbb{T}^{2d} \times \mathbb{R}^{2d}} \psi(x-y) f^\varepsilon(t, x, v) f^\varepsilon(t, y, w) |v-w|^2 dx dy dv dw = \mathcal{D}_2(f^\varepsilon).$$

We next claim  $\mathcal{I}_2 \leq 0$ .

Indeed, since

$$\rho^\varepsilon |u^\varepsilon|^2 = \frac{(\int_{\mathbb{R}^d} v f^\varepsilon dv)^2}{\int_{\mathbb{R}^d} f^\varepsilon dv} \leq \int_{\mathbb{R}^d} |v|^2 f^\varepsilon dv, \quad (5-2)$$

we have

$$\begin{aligned} \int_{\mathbb{T}^{2d} \times \mathbb{R}^{2d}} \psi(x-y) f^\varepsilon(t, x, v) f^\varepsilon(t, y, w) |v|^2 dv dw dx dy \\ \geq \int_{\mathbb{T}^{2d}} \psi(x-y) \rho^\varepsilon(t, y) \rho^\varepsilon(t, x) |u^\varepsilon(t, x)|^2 dx dy. \end{aligned}$$

Then, since

$$\begin{aligned} \int_{\mathbb{T}^{2d} \times \mathbb{R}^{2d}} \psi(x-y) f^\varepsilon(t, x, v) f^\varepsilon(t, y, w) u^\varepsilon(t, x) \cdot w dv dw dx dy \\ = \int_{\mathbb{T}^{2d}} \psi(x-y) \rho^\varepsilon(t, x) \rho^\varepsilon(t, y) u^\varepsilon(t, x) \cdot u^\varepsilon(t, y) dx dy, \\ \int_{\mathbb{T}^{2d} \times \mathbb{R}^{2d}} \psi(x-y) f^\varepsilon(t, x, v) f^\varepsilon(t, y, w) u^\varepsilon(t, x) \cdot v dv dw dx dy \\ = \int_{\mathbb{T}^{2d}} \psi(x-y) \rho^\varepsilon(t, x) \rho^\varepsilon(t, y) |u^\varepsilon(t, x)|^2 dx dy, \\ \int_{\mathbb{T}^{2d} \times \mathbb{R}^{2d}} \psi(x-y) f^\varepsilon(t, x, v) f^\varepsilon(t, y, w) v \cdot w dv dw dx dy \\ = \int_{\mathbb{T}^{2d}} \psi(x-y) \rho^\varepsilon(t, x) \rho^\varepsilon(t, y) u^\varepsilon(t, x) \cdot u^\varepsilon(t, y) dx dy, \end{aligned}$$

we conclude that  $\mathcal{I}_2 \leq 0$ , as desired.  $\square$

*Verification of (H2).* We show that the assumptions (A1)–(A3) for initial data imply (H2). Using (3-6) and assumption (A3), we have

$$\int_{\mathbb{T}^d} \eta(U_0^\varepsilon | U_0) dx = \frac{1}{2} \int_{\mathbb{T}^d} \rho_0^\varepsilon |u_0^\varepsilon - u_0|^2 dx \leq C\varepsilon^2 \int_{\mathbb{T}^d} \rho_0^\varepsilon dx \leq C\varepsilon^2.$$

Since it follows from (A1)–(A3) that

$$\int_{\mathbb{T}^d} (\mathcal{F}(f_0^\varepsilon) - \eta(U_0)) dx = \mathcal{O}(\varepsilon),$$

and

$$\begin{aligned} \int_{\mathbb{T}^d} (\eta(U_0) - \eta(U_0^\varepsilon)) dx &= \frac{1}{2} \int_{\mathbb{R}^d} (\rho_0 |u_0|^2 - \rho_0^\varepsilon |u_0^\varepsilon|^2) \\ &\leq \frac{1}{2} \int_{\mathbb{T}^d} |\rho_0 - \rho_0^\varepsilon| |u_0|^2 + \frac{1}{2} \int_{\mathbb{T}^d} \rho_0^\varepsilon ||u_0^\varepsilon|^2 - |u_0|^2| = \mathcal{O}(\varepsilon), \end{aligned}$$

we have

$$\int_{\mathbb{T}^d} (\mathcal{F}(f_0^\varepsilon) - \eta(U_0^\varepsilon)) dx = \mathcal{O}(\varepsilon).$$

It is obvious that (A1) implies

$$\int_{\mathbb{T}^d} \mathcal{F}(f_0^\varepsilon) dx \leq C.$$

*Verification of (H3).* It follows from (5-2) that

$$\eta(U^\varepsilon) = \rho^\varepsilon \frac{1}{2} |u^\varepsilon|^2 \leq \int_{\mathbb{R}^d} \frac{1}{2} |v|^2 f^\varepsilon dv = \mathcal{F}(f^\varepsilon). \tag{5-3}$$

*Verification of (H4).* Since

$$A(U) = \begin{pmatrix} P^T \\ (P \otimes P)/\rho \end{pmatrix},$$

we have

$$\begin{aligned} DA(U) \cdot (U^\varepsilon - U) &= D_\rho A(U)(\rho^\varepsilon - \rho) + D_{P_i} A(U)(P_i^\varepsilon - P_i) \\ &= \begin{pmatrix} (P^\varepsilon - P)^T \\ -((\rho^\varepsilon - \rho)/\rho^2)P \otimes P + (1/\rho)P \otimes (P^\varepsilon - P) + (1/\rho)(P^\varepsilon - P) \otimes P \end{pmatrix}, \end{aligned}$$

which yields

$$\begin{aligned} A(U^\varepsilon | U) &= \begin{pmatrix} 0 \\ (1/\rho^\varepsilon)P^\varepsilon \otimes P^\varepsilon - (1/\rho)P \otimes P + ((\rho^\varepsilon - \rho)/\rho^2)P \otimes P - (1/\rho)P \otimes (P^\varepsilon - P) - (1/\rho)(P^\varepsilon - P) \otimes P \end{pmatrix} \\ &= \begin{pmatrix} 0 \\ \rho^\varepsilon (u^\varepsilon - u) \otimes (u^\varepsilon - u) \end{pmatrix}. \end{aligned}$$

Therefore, using (3-5) and (3-6), we have

$$\begin{aligned} \left| \int \nabla_x (D\eta(U)) : A(U^\varepsilon | U) dx \right| &= \left| \int_0^t \int_{\mathbb{T}^d} \rho^\varepsilon (u^\varepsilon - u) \otimes (u^\varepsilon - u) : \nabla_x u dx ds \right| \\ &\leq C \|\nabla_x u\|_{L^\infty((0, T_*) \times \mathbb{T}^d)} \int_0^t \int_{\mathbb{T}^d} \eta(U^\varepsilon | U) dx ds. \end{aligned}$$

*Verification of (H5).* For a weak solution  $f^\varepsilon$  to (1-2), it follows from (2-13) that  $U^\varepsilon = \left(\frac{\rho^\varepsilon}{p^\varepsilon}\right)$  solves the system

$$\partial_t U^\varepsilon + \operatorname{div}_x A(U^\varepsilon) - F(U^\varepsilon) = \operatorname{div}_x \begin{pmatrix} 0 \\ -\int_{\mathbb{R}^d} (v - u^\varepsilon) \otimes (v - u^\varepsilon) f^\varepsilon dv \end{pmatrix}, \tag{5-4}$$

where the equality holds in the sense of distributions; see (2-8). Therefore, we have

$$\begin{aligned} & \left| \int D\eta(U) \cdot [\partial_t U^\varepsilon + \operatorname{div}_x A(U^\varepsilon) - F(U^\varepsilon)] dx \right| \\ &= \left| \int_{\mathbb{T}^d} \nabla_x u : \left( \int_{\mathbb{R}^d} (v - u^\varepsilon) \otimes (v - u^\varepsilon) f^\varepsilon dv \right) dx \right| \\ &\leq C \|\nabla_x u\|_{L^\infty((0, T_*) \times \mathbb{T}^d)} \int_{\mathbb{T}^d \times \mathbb{R}^d} |v - u^\varepsilon|^2 f^\varepsilon dv dx = C \|\nabla_x u\|_{L^\infty((0, T_*) \times \mathbb{T}^d)} \mathcal{D}_1(f^\varepsilon). \end{aligned}$$

*Verification of (H6).* From the proof of Proposition 4.2 in [Karper et al. 2015], we see

$$-\int_{\mathbb{T}^d} [D^2 \eta(U) F(U)(U^\varepsilon - U) + D\eta(U) F(U^\varepsilon)] dx = K_1 + K_2 + K_3,$$

where

$$\begin{aligned} K_1 &:= -\frac{1}{2} \int_{\mathbb{T}^{2d}} \psi(x - y) \rho^\varepsilon(x) \rho^\varepsilon(t, y) |(u^\varepsilon(x) - u(x)) - (u^\varepsilon(y) - u(y))|^2 dx dy, \\ K_2 &:= \frac{1}{2} \int_{\mathbb{T}^{2d}} \psi(x - y) \rho^\varepsilon(x) \rho^\varepsilon(y) |u^\varepsilon(x) - u^\varepsilon(y)|^2 dx dy, \\ K_3 &:= \int_{\mathbb{T}^{2d}} \psi(x - y) \rho^\varepsilon(x) (\rho^\varepsilon(y) - \rho(y)) (u(y) - u(x)) (u^\varepsilon(x) - u(x)) dx dy. \end{aligned}$$

Notice that  $K_1 \leq 0$ , and  $K_2 = \tilde{\mathcal{D}}_2(f^\varepsilon)$  where  $\tilde{\mathcal{D}}_2(f^\varepsilon)$  is in Lemma 5.1.

To estimate  $K_3$ , we separate it into two parts:

$$\begin{aligned} K_3 &= \int_{\mathbb{T}^d} \left( \int_{\mathbb{T}^d} \psi(x - y) u(y) (\rho^\varepsilon(y) - \rho(y)) dy \right) \rho^\varepsilon(x) (u^\varepsilon(x) - u(x)) dx \\ &\quad - \int_{\mathbb{T}^d} \left( \int_{\mathbb{T}^d} \psi(x - y) (\rho^\varepsilon(y) - \rho(y)) dy \right) u(x) \rho^\varepsilon(x) (u^\varepsilon(x) - u(x)) dx. \end{aligned}$$

Since  $\psi$  and  $u$  are Lipschitz, we use the Kantorovich–Rubinstein theorem, see [Villani 2009, Theorem 5.10 and Particular Case 5.16], to estimate

$$\begin{aligned} K_3 &\leq W_1(\rho^\varepsilon, \rho) \left( \sup_{x \in \mathbb{T}^d} \|\psi(x - \cdot) u\|_{L^\infty(0, T_*; W^{1, \infty}(\mathbb{T}^d))} + \|\psi\|_{L^\infty(0, T_*; W^{1, \infty}(\mathbb{T}^d))} \|u\|_{L^\infty((0, T_*) \times \mathbb{T}^d)} \right) \\ &\quad \times \int_{\mathbb{T}^d} \rho^\varepsilon(x) |u^\varepsilon(x) - u(x)| dx. \end{aligned}$$

Therefore, since  $W_1(\rho^\varepsilon, \rho) \leq W_2(\rho^\varepsilon, \rho)$ , we obtain

$$K_3 \leq C \left( W_2^2(\rho^\varepsilon, \rho) + \int_{\mathbb{T}^d} \rho^\varepsilon(x) |u^\varepsilon(x) - u(x)|^2 dx \right).$$

Hence we have verified (H6).

*Verification of (H7).* This will be shown by Lemma 5.2 below. We first derive some estimates for the characteristics generated by the velocity fields  $u^\varepsilon$  and  $u$ .

For the velocity  $u$  in the limit system (1-4), let  $X$  be a characteristic generated by it, that is,

$$\dot{X}(t, x) = u(t, X(t, x)), \quad X(0, x) = x. \quad (5-5)$$

Then, thanks to the smoothness of  $u$ , it follows from (3-8) that

$$X(t)_\# \rho_0(x) dx = \rho(t, x) dx.$$

On the other hand, since  $u^\varepsilon$  is not Lipschitz with respect to  $x$ , we use a probabilistic representation for  $\rho^\varepsilon$  as a solution of the continuity equation in (3-3). More precisely, (5-3) and (2-9) imply

$$\int_{\mathbb{T}^d} |u^\varepsilon(t)|^2 \rho^\varepsilon(t) dx \leq \int_{\mathbb{T}^d} \mathcal{F}(f^\varepsilon)(t) dx \leq \int_{\mathbb{T}^d} \mathcal{F}(f_0^\varepsilon) dx < \infty,$$

so it follows from Proposition 3.3 that there exists a probability measure  $\eta^\varepsilon$  in  $\Gamma_{T_*} \times \mathbb{T}^d$  that is concentrated on the set of pairs  $(\gamma, x)$  such that  $\gamma$  is a solution of the ODE

$$\dot{\gamma}(t) = u^\varepsilon(\gamma(t)), \quad \gamma(0) = x, \quad (5-6)$$

and

$$\int_{\mathbb{T}^d} \varphi(x) \rho^\varepsilon(t, x) dx = \int_{\Gamma_{T_*} \times \mathbb{T}^d} \varphi(\gamma(t)) d\eta^\varepsilon(\gamma, x) \quad \text{for all } \varphi \in C^0(\mathbb{T}^d), \quad t \in [0, T_*]. \quad (5-7)$$

In particular, this says that the time marginal of the measure  $\eta^\varepsilon$  at time 0 is given by  $\rho^\varepsilon(0) = \rho_0^\varepsilon$ . Hence, by the disintegration theorem of measures, see for instance [Ambrosio et al. 2005, Theorem 5.3.1] and the comments at the end of Section 8.2 in [Ambrosio et al. 2005], we can write

$$d\eta^\varepsilon(\gamma, x) = \eta_x^\varepsilon(d\gamma) \otimes \rho_0^\varepsilon(x) dx,$$

where  $\{\eta_x^\varepsilon\}_{x \in \mathbb{T}^d}$  is a family of probability measures on  $\Gamma_{T_*}$  concentrated on solutions of (5-6).

For the flow  $X$  in (5-5), we also consider the densities  $\tilde{\rho}^\varepsilon(t)$  defined as

$$\tilde{\rho}^\varepsilon(t, x) dx = X(t)_\# \rho_0^\varepsilon(x) dx. \quad (5-8)$$

Note that, since

$$\begin{aligned} \|\rho(t) - \tilde{\rho}^\varepsilon(t)\|_{L^1(\mathbb{T}^d)} &= \sup_{\|\varphi\|_\infty \leq 1} \int_{\mathbb{T}^d} \varphi(x) [\rho(t, x) - \tilde{\rho}^\varepsilon(t, x)] dx \\ &= \sup_{\|\varphi\|_\infty \leq 1} \int_{\mathbb{T}^d} \varphi(X(t, x)) [\rho_0(x) - \rho_0^\varepsilon(x)] dx \leq \|\rho_0^\varepsilon - \rho_0\|_{L^1(\mathbb{T}^d)}, \end{aligned}$$

we have

$$\|\rho(t) - \tilde{\rho}^\varepsilon(t)\|_{L^1(\mathbb{T}^d)} \leq \|\rho_0^\varepsilon - \rho_0\|_{L^1(\mathbb{T}^d)}. \quad (5-9)$$

We now consider the measure  $\nu^\varepsilon$  on  $\Gamma_{T_*} \times \Gamma_{T_*} \times \mathbb{T}^d$  defined as

$$d\nu^\varepsilon(\gamma, \sigma, x) = \eta_x^\varepsilon(d\gamma) \otimes \delta_{X(\cdot, x)}(d\sigma) \otimes \rho_0^\varepsilon(x) dx.$$

If we consider the evaluation map

$$E_t : \Gamma_{T_*} \times \Gamma_{T_*} \times \mathbb{T}^d \rightarrow \mathbb{T}^d \times \mathbb{T}^d, \quad E_t(\gamma, \sigma, x) = (\gamma(t), \sigma(t)),$$

it follows that the measure  $\pi_t^\varepsilon := (E_t)_\# \nu^\varepsilon$  on  $\mathbb{T}^d \times \mathbb{T}^d$  has marginals  $\rho^\varepsilon(t, x) dx$  and  $\tilde{\rho}^\varepsilon(t, y) dy$  for all  $t \geq 0$ . Therefore, we have

$$\begin{aligned} \int_{\Gamma_{T_*} \times \mathbb{T}^d} |\gamma(t) - X(t, x)|^2 \eta_x^\varepsilon(d\gamma) \otimes \rho_0^\varepsilon(x) dx &= \int_{\Gamma_{T_*} \times \Gamma_{T_*} \times \mathbb{T}^d} |\gamma(t) - \sigma(t)|^2 d\nu^\varepsilon(\gamma, \sigma, x) \\ &= \int_{\mathbb{T}^{2d}} |x - y|^2 d\pi_t^\varepsilon(x, y) \\ &\geq W_2^2(\rho^\varepsilon(t), \tilde{\rho}^\varepsilon(t)). \end{aligned} \tag{5-10}$$

We now use the above results to prove the following lemma.

**Lemma 5.2.** *Under the same assumptions as in Theorem 3.1, we have*

$$W_2^2(\rho^\varepsilon(t), \rho(t)) \leq C e^{T_*} \int_0^t \int_{\mathbb{T}^d} |u^\varepsilon(s, x) - u(s, x)|^2 \rho^\varepsilon(s, x) dx ds + \mathcal{O}(\varepsilon), \quad t \leq T_*. \tag{5-11}$$

*Proof.* Let  $\tilde{\rho}^\varepsilon$  be defined as in (5-8). We begin by observing that, thanks to Lemma 3.4, (5-9), and assumption (A2), it follows that

$$W_2^2(\tilde{\rho}^\varepsilon(t), \rho(t)) \leq \mathcal{O}(\varepsilon).$$

Hence, to prove (5-11), it is enough to bound  $W_2^2(\rho^\varepsilon(t), \tilde{\rho}^\varepsilon(t))$ .

To this aim, we try to get a Gronwall-type inequality on

$$\int_{\Gamma_{T_*} \times \mathbb{T}^d} |\gamma(t) - X(t, x)|^2 \eta_x^\varepsilon(d\gamma) \otimes \rho_0^\varepsilon(x) dx.$$

Since

$$\dot{\gamma}(t) - \dot{X}(t, x) = (u^\varepsilon(\gamma(t)) - u(\gamma(t))) + (u(\gamma(t)) - u(X(t, x)))$$

by (5-5) and (5-6), we have

$$\begin{aligned} \frac{1}{2} \frac{d}{dt} \int_{\Gamma_{T_*} \times \mathbb{T}^d} |\gamma(t) - X(t, x)|^2 d\eta_x^\varepsilon(\gamma) \otimes \rho_0^\varepsilon(x) dx &\leq \int_{\Gamma_{T_*} \times \mathbb{T}^d} |u^\varepsilon(\gamma(t)) - u(\gamma(t))|^2 d\eta_x^\varepsilon(\gamma) \otimes \rho_0^\varepsilon(x) dx \\ &\quad + \int_{\Gamma_{T_*} \times \mathbb{T}^d} |u(\gamma(t)) - u(X(t, x))|^2 d\eta_x^\varepsilon(\gamma) \otimes \rho_0^\varepsilon(x) dx \\ &\quad + 2 \int_{\Gamma_{T_*} \times \mathbb{T}^d} |\gamma(t) - X(t, x)|^2 d\eta_x^\varepsilon(\gamma) \otimes \rho_0^\varepsilon(x) dx. \end{aligned}$$

Notice that, thanks to (5-7),

$$\int_{\Gamma_{T_*} \times \mathbb{T}^d} |u^\varepsilon(\gamma(t)) - u(\gamma(t))|^2 d\eta_x^\varepsilon(\gamma) \otimes \rho_0^\varepsilon(x) dx = \int_{\mathbb{T}^d} |u^\varepsilon(t, x) - u(t, x)|^2 \rho^\varepsilon(t, x) dx.$$

Moreover, since

$$\begin{aligned} \int_{\Gamma_{T_*} \times \mathbb{T}^d} |u(\gamma(t)) - u(X(t, x))|^2 d\eta_x^\varepsilon(\gamma) \otimes \rho_0^\varepsilon(x) dx \\ \leq \|u\|_{L^\infty(0, T_*; W^{1, \infty}(\mathbb{T}^d))} \int_{\Gamma_{T_*} \times \mathbb{T}^d} |\gamma(t) - X(t, x)|^2 d\eta_x^\varepsilon(\gamma) \otimes \rho_0^\varepsilon(x) dx, \end{aligned}$$

we have

$$\begin{aligned} \frac{d}{dt} \int_{\Gamma_{T_*} \times \mathbb{T}^d} |\gamma(t) - X(t, x)|^2 d\eta_x^\varepsilon(\gamma) \otimes \rho_0^\varepsilon(x) dx \\ \leq C \int_{\Gamma_{T_*} \times \mathbb{T}^d} |\gamma(t) - X(t, x)|^2 d\eta_x^\varepsilon(\gamma) \otimes \rho_0^\varepsilon(x) dx + \int_{\mathbb{T}^d} |u^\varepsilon(t, x) - u(t, x)|^2 \rho^\varepsilon(t, x) dx. \end{aligned}$$

Therefore, using Gronwall’s inequality together with  $\gamma(0) = X(0, x) = x$  for  $\eta_x^\varepsilon$ -a.e.  $\gamma$ , we obtain

$$\int_{\Gamma_{T_*} \times \mathbb{T}^d} |\gamma(t) - X(t, x)|^2 d\eta_x^\varepsilon(\gamma) \otimes \rho_0^\varepsilon(x) dx \leq C e^{T_*} \int_0^t \int_{\mathbb{T}^d} |u^\varepsilon(s, x) - u(s, x)|^2 \rho^\varepsilon(s, x) dx ds, \quad t \leq T_*.$$

Hence, using (5-10) we get the desired control on  $W_2^2(\rho^\varepsilon(t), \tilde{\rho}^\varepsilon(t))$ , which concludes the proof.  $\square$

**Proof of (3-2).** Here we use the estimate (3-1) to show the convergence (3-2).

First, since (5-1) and (A1) imply

$$\int_0^t \mathcal{D}_1(f^\varepsilon)(s) ds \leq C\varepsilon,$$

using (3-1), we have

$$\int_0^{T_*} \int_{\mathbb{T}^d \times \mathbb{R}^d} f^\varepsilon |v - u|^2 dx dv ds \leq 2 \int_0^{T_*} \int_{\mathbb{T}^d \times \mathbb{R}^d} f^\varepsilon (|v - u^\varepsilon|^2 + |u^\varepsilon - u|^2) dx dv ds \leq C(1 + T_*)\varepsilon. \quad (5-12)$$

Then, for any  $\varphi \in C_c^1((0, T_*) \times \mathbb{T}^d \times \mathbb{R}^d)$ ,

$$\begin{aligned} \int_0^{T_*} \int_{\mathbb{T}^d \times \mathbb{R}^d} \varphi(s, x, v) f^\varepsilon dx dv ds - \int_0^{T_*} \int_{\mathbb{T}^d \times \mathbb{R}^d} \varphi(s, x, v) \rho \delta_u(dv) dx ds \\ = \int_0^{T_*} \int_{\mathbb{T}^d \times \mathbb{R}^d} \varphi(s, x, v) f^\varepsilon dx dv ds - \int_0^{T_*} \int_{\mathbb{T}^d} \varphi(s, x, u) \rho dx ds \\ = \underbrace{\int_0^{T_*} \int_{\mathbb{T}^d \times \mathbb{R}^d} f^\varepsilon (\varphi(s, x, v) - \varphi(s, x, u)) dx dv ds}_{=: I_1^\varepsilon} + \underbrace{\int_0^{T_*} \int_{\mathbb{T}^d} \varphi(s, x, u) (\rho^\varepsilon - \rho) dx ds}_{=: I_2^\varepsilon}. \end{aligned}$$

Using (5-12), we have

$$\begin{aligned} I_1^\varepsilon &\leq \|\nabla_v \varphi\|_\infty \int_0^{T_*} \int_{\mathbb{T}^d \times \mathbb{R}^d} f^\varepsilon |v - u| dx dv ds \\ &= \|\nabla_v \varphi\|_\infty \left( \int_0^{T_*} \int_{|v-u| \leq \sqrt{\varepsilon}} f^\varepsilon |v - u| dx dv ds + \int_0^{T_*} \int_{|v-u| > \sqrt{\varepsilon}} f^\varepsilon |v - u| dx dv ds \right) \end{aligned}$$

$$\begin{aligned} &\leq \|\nabla_v \varphi\|_\infty \left( \sqrt{\varepsilon} T_* + \frac{1}{\sqrt{\varepsilon}} \int_0^{T_*} \int_{|v-u| > \sqrt{\varepsilon}} f^\varepsilon |v-u|^2 dv dx ds \right) \\ &\leq C(1 + T_*)\sqrt{\varepsilon}. \end{aligned}$$

Since  $W_1(\rho^\varepsilon, \rho) \leq W_2(\rho^\varepsilon, \rho) \rightarrow 0$  by (3-1), we also have  $I_2^\varepsilon \rightarrow 0$  as  $\varepsilon \rightarrow 0$ .

Hence, this completes the proof of (3-2).

### Appendix: Proof of Theorem 2.2

We first estimate  $(d/dt)\mathcal{E}_1$  as follows:

$$\frac{d}{dt} \mathcal{E}_1 = 2 \int_{\mathbb{T}^d \times \mathbb{R}^d} f(u-v) \partial_t u dv dx + \int_{\mathbb{T}^d \times \mathbb{R}^d} \partial_t f |u-v|^2 dv dx := I_1 + I_2.$$

First of all, by the definition of  $u$ , we have  $\int f(u-v) dv = 0$ ; hence  $I_1 = 0$ .

Concerning  $I_2$ , it follows from (1-1) that

$$\begin{aligned} I_2 &= \int_{\mathbb{T}^d \times \mathbb{R}^d} |u-v|^2 (-\nabla_x \cdot (vf) - \nabla_v \cdot (L[f]f) - \nabla_v \cdot ((u-v)f)) dv dx \\ &= \underbrace{2 \int_{\mathbb{T}^d \times \mathbb{R}^d} \nabla_x u (u-v) \cdot vf dv dx}_{=: I_{21}} - \underbrace{2 \int_{\mathbb{T}^d \times \mathbb{R}^d} (u-v) \cdot L[f]f dv dx}_{=: I_{22}} - \underbrace{2 \int_{\mathbb{T}^d \times \mathbb{R}^d} |u-v|^2 f dv dx}_{=: -2\mathcal{E}_1}. \end{aligned}$$

Then, we use the stress tensor  $P = \int_{\mathbb{R}^d} (v-u) \otimes (v-u) f dv$  to rewrite  $I_{21}$  as

$$I_{21} = 2 \int_{\mathbb{T}^d \times \mathbb{R}^d} \nabla_x u (u-v) \cdot (v-u) f dv dx = 2 \int_{\mathbb{T}^d} (\nabla_x \cdot P) \cdot u dx.$$

Thanks to the estimate on  $\mathcal{I}_2$  in the proof of Lemma 5.1, we see that

$$I_{22} = -2 \int_{\mathbb{T}^{2d} \times \mathbb{R}^{2d}} \psi(x-y) f(t,x,v) f(t,y,w) (u(t,x)-v) \cdot (w-v) dv dw dx dy \leq 0.$$

Therefore, we have

$$\frac{d}{dt} \mathcal{E}_1 \leq 2 \int_{\mathbb{T}^d} (\nabla_x \cdot P) \cdot u dx - 2\mathcal{E}_1. \tag{A-1}$$

We next estimate  $(d/dt)\mathcal{E}_2$  as follows:

$$\begin{aligned} \frac{d}{dt} \mathcal{E}_2 &= 2 \int_{\mathbb{T}^{2d}} \partial_t \rho(t,x) \rho(t,y) |u(t,x) - u(t,y)|^2 dx dy \\ &\quad + 2 \int_{\mathbb{T}^{2d}} \rho(t,x) \rho(t,y) (u(t,x) - u(t,y)) \partial_t (u(t,x) - u(t,y)) dx dy \\ &:= J_1 + J_2. \end{aligned}$$

Since it follows from (2-13) with  $\varepsilon = 1$  that

$$\begin{aligned} \partial_t \rho + \nabla_x \cdot (\rho u) &= 0, \\ \rho \partial_t u + \rho u \cdot \nabla_x u + \nabla_x \cdot P &= \int_{\mathbb{R}^d} L[f]f dv, \end{aligned}$$

we obtain (recall that  $\|\rho\|_{L^1(\mathbb{T}^d)} = 1$ )

$$\begin{aligned} J_1 &= -2 \int_{\mathbb{T}^{2d}} \nabla_x \cdot (\rho u)(t, x) \rho(t, y) |u(t, x) - u(t, y)|^2 dx dy \\ &= 4 \int_{\mathbb{T}^d} \rho u \cdot \nabla_x u \cdot u dx - 4 \int_{\mathbb{T}^d} \rho u \cdot \nabla_x u dx \cdot \int_{\mathbb{T}^d} \rho u dx, \end{aligned}$$

and

$$\begin{aligned} J_2 &= 4 \int_{\mathbb{T}^{2d}} \rho(t, y) u(t, x) \rho(t, x) \partial_t u(t, x) dx dy - 4 \int_{\mathbb{T}^{2d}} \rho(t, y) u(t, y) \rho(t, x) \partial_t u(t, x) dx dy \\ &= -4 \int_{\mathbb{T}^d} \rho u \cdot \nabla_x u \cdot u dx - 4 \int_{\mathbb{T}^d} \nabla_x \cdot P \cdot u dx + 4 \underbrace{\int_{\mathbb{T}^d \times \mathbb{R}^d} u \cdot L[f] f dx dv}_{:= J_{21}} + 4 \int_{\mathbb{T}^d} \rho u \cdot \nabla_x u dx \cdot \int_{\mathbb{T}^d} \rho u dx \\ &\quad + 4 \underbrace{\int_{\mathbb{T}^d} \nabla_x \cdot P dx}_{=0} \cdot \int_{\mathbb{T}^d} \rho u dx - 4 \underbrace{\int_{\mathbb{T}^d \times \mathbb{R}^d} L[f] f dx dv}_{:= J_{22}} \cdot \int_{\mathbb{T}^d} \rho u dx. \end{aligned}$$

Now, we compute the above terms  $J_{21}$  and  $J_{22}$  as follows:

$$\begin{aligned} J_{21} &= \int_{\mathbb{T}^{2d} \times \mathbb{R}^{2d}} \psi(x-y) f(t, x, v) f(t, y, w) (w-v) \cdot u(t, x) dv dw dx dy \\ &= \int_{\mathbb{T}^{2d}} \psi(x-y) \rho(t, x) \rho(t, y) (u(t, y) - u(t, x)) \cdot u(t, x) dx dy \\ &= -\frac{1}{2} \int_{\mathbb{T}^{2d}} \psi(x-y) \rho(t, x) \rho(t, y) |u(t, x) - u(t, y)|^2 dx dy, \\ J_{22} &= \int_{\mathbb{T}^{2d} \times \mathbb{R}^{2d}} \psi(x-y) f(t, x, v) f(t, y, w) (w-v) dv dw dx dy = 0. \end{aligned}$$

Therefore, we have

$$\frac{d}{dt} \mathcal{E}_2 = -4 \int_{\mathbb{T}^d} \nabla_x \cdot P \cdot u dx - 2 \int_{\mathbb{T}^{2d}} \psi(x-y) \rho(t, x) \rho(t, y) |u(t, x) - u(t, y)|^2 dx dy.$$

Recalling (A-1), proves that

$$\begin{aligned} \frac{d}{dt} \mathcal{E} &\leq -2\mathcal{E}_1 - \int_{\mathbb{T}^{2d}} \psi(x-y) \rho(t, x) \rho(t, y) |u(t, x) - u(t, y)|^2 dx dy \\ &\leq -2\mathcal{E}_1 - \psi_m \mathcal{E}_2 \leq -2 \min\{1, \psi_m\} \mathcal{E}, \end{aligned}$$

which completes the proof of (2-11).

To show the second bound (2-12), note that if  $\ell_T := \sup_{t \in [0, T]} \|\nabla_x u\|_{L^\infty(\mathbb{T}^d)} < \infty$  then (A-1) yields

$$\frac{d}{dt} \mathcal{E}_1(t) \leq -2 \int_{\mathbb{T}^d} \nabla_x u : P dx - 2\mathcal{E}_1 \leq 2\ell_T \int_{\mathbb{T}^d \times \mathbb{R}^d} |u-v|^2 f dv dx - 2\mathcal{E}_1(t) = 2(\ell_T - 1)\mathcal{E}_1(t),$$

which proves (2-12).  $\square$



## References

- [Ambrosio et al. 2005] L. Ambrosio, N. Gigli, and G. Savaré, *Gradient flows in metric spaces and in the space of probability measures*, Birkhäuser, Basel, 2005. MR Zbl
- [Berthelin and Vasseur 2005] F. Berthelin and A. Vasseur, “From kinetic equations to multidimensional isentropic gas dynamics before shocks”, *SIAM J. Math. Anal.* **36**:6 (2005), 1807–1835. MR Zbl
- [Bouchut 1994] F. Bouchut, “On zero pressure gas dynamics”, pp. 171–190 in *Advances in kinetic theory and computing*, edited by B. Perthame, Ser. Adv. Math. Appl. Sci. **22**, World Sci., River Edge, NJ, 1994. MR Zbl
- [Bouchut and James 1999] F. Bouchut and F. James, “Duality solutions for pressureless gases, monotone scalar conservation laws, and uniqueness”, *Comm. Partial Differential Equations* **24**:11-12 (1999), 2173–2189. MR Zbl
- [Boudin 2000] L. Boudin, “A solution with bounded expansion rate to the model of viscous pressureless gases”, *SIAM J. Math. Anal.* **32**:1 (2000), 172–193. MR Zbl
- [Brenier and Grenier 1998] Y. Brenier and E. Grenier, “Sticky particles and scalar conservation laws”, *SIAM J. Numer. Anal.* **35**:6 (1998), 2317–2328. MR Zbl
- [Cañizo et al. 2011] J. A. Cañizo, J. A. Carrillo, and J. Rosado, “A well-posedness theory in measures for some kinetic models of collective motion”, *Math. Models Methods Appl. Sci.* **21**:3 (2011), 515–539. MR Zbl
- [Carlen et al. 2015] E. Carlen, M. C. Carvalho, P. Degond, and B. Wennberg, “A Boltzmann model for rod alignment and schooling fish”, *Nonlinearity* **28**:6 (2015), 1783–1803. MR Zbl
- [Carrillo et al. 2010] J. A. Carrillo, M. Fornasier, J. Rosado, and G. Toscani, “Asymptotic flocking dynamics for the kinetic Cucker–Smale model”, *SIAM J. Math. Anal.* **42**:1 (2010), 218–236. MR Zbl
- [Carrillo et al. 2016] J. A. Carrillo, Y.-P. Choi, and T. K. Karper, “On the analysis of a coupled kinetic–fluid model with local alignment forces”, *Ann. Inst. H. Poincaré Anal. Non Linéaire* **33**:2 (2016), 273–307. MR Zbl
- [Cucker and Smale 2007] F. Cucker and S. Smale, “Emergent behavior in flocks”, *IEEE Trans. Automat. Control* **52**:5 (2007), 852–862. MR Zbl
- [Do et al. 2018] T. Do, A. Kiselev, L. Ryzhik, and C. Tan, “Global regularity for the fractional Euler alignment system”, *Arch. Ration. Mech. Anal.* **228**:1 (2018), 1–37. MR Zbl
- [Duan et al. 2010] R. Duan, M. Fornasier, and G. Toscani, “A kinetic flocking model with diffusion”, *Comm. Math. Phys.* **300**:1 (2010), 95–145. MR Zbl
- [Fornasier et al. 2011] M. Fornasier, J. Haskovec, and G. Toscani, “Fluid dynamic description of flocking via the Povzner–Boltzmann equation”, *Phys. D* **240**:1 (2011), 21–31. MR Zbl
- [Goudon et al. 2004] T. Goudon, P.-E. Jabin, and A. Vasseur, “Hydrodynamic limit for the Vlasov–Navier–Stokes equations, II: Fine particles regime”, *Indiana Univ. Math. J.* **53**:6 (2004), 1517–1536. MR Zbl
- [Ha and Liu 2009] S.-Y. Ha and J.-G. Liu, “A simple proof of the Cucker–Smale flocking dynamics and mean-field limit”, *Commun. Math. Sci.* **7**:2 (2009), 297–325. MR Zbl
- [Ha and Tadmor 2008] S.-Y. Ha and E. Tadmor, “From particle to kinetic and hydrodynamic descriptions of flocking”, *Kinet. Relat. Models* **1**:3 (2008), 415–435. MR Zbl
- [Ha et al. 2014a] S.-Y. Ha, F. Huang, and Y. Wang, “A global unique solvability of entropic weak solution to the one-dimensional pressureless Euler system with a flocking dissipation”, *J. Differential Equations* **257**:5 (2014), 1333–1371. MR Zbl
- [Ha et al. 2014b] S.-Y. Ha, M.-J. Kang, and B. Kwon, “A hydrodynamic model for the interaction of Cucker–Smale particles and incompressible fluid”, *Math. Models Methods Appl. Sci.* **24**:11 (2014), 2311–2359. MR Zbl
- [Ha et al. 2014c] S.-Y. Ha, Z. Li, M. Slemrod, and X. Xue, “Flocking behavior of the Cucker–Smale model under rooted leadership in a large coupling limit”, *Quart. Appl. Math.* **72**:4 (2014), 689–701. MR Zbl
- [Ha et al. 2015] S.-Y. Ha, M.-J. Kang, and B. Kwon, “Emergent dynamics for the hydrodynamic Cucker–Smale system in a moving domain”, *SIAM J. Math. Anal.* **47**:5 (2015), 3813–3831. MR Zbl
- [Ha et al. 2017] S.-Y. Ha, J. Jeong, S. E. Noh, Q. Xiao, and X. Zhang, “Emergent dynamics of Cucker–Smale flocking particles in a random environment”, *J. Differential Equations* **262**:3 (2017), 2554–2591. MR Zbl

- [Huang and Wang 2001] F. Huang and Z. Wang, “Well posedness for pressureless flow”, *Comm. Math. Phys.* **222**:1 (2001), 117–146. MR Zbl
- [Jabin 2000] P.-E. Jabin, *Équations de transport modélisant des particules en interaction dans un fluide et comportement asymptotiques*, Ph.D. thesis, Université Paris VI, 2000.
- [Jabin and Rey 2017] P.-E. Jabin and T. Rey, “Hydrodynamic limit of granular gases to pressureless Euler in dimension 1”, *Quart. Appl. Math.* **75**:1 (2017), 155–179. MR Zbl
- [Kang 2018] M.-J. Kang, “From the Vlasov–Poisson equation with strong local alignment to the pressureless Euler–Poisson system”, *Appl. Math. Lett.* **79** (2018), 85–91. MR
- [Kang and Vasseur 2015] M.-J. Kang and A. F. Vasseur, “Asymptotic analysis of Vlasov-type equations under strong local alignment regime”, *Math. Models Methods Appl. Sci.* **25**:11 (2015), 2153–2173. MR Zbl
- [Karper et al. 2013] T. K. Karper, A. Mellet, and K. Trivisa, “Existence of weak solutions to kinetic flocking models”, *SIAM J. Math. Anal.* **45**:1 (2013), 215–243. MR Zbl
- [Karper et al. 2014] T. K. Karper, A. Mellet, and K. Trivisa, “On strong local alignment in the kinetic Cucker–Smale model”, pp. 227–242 in *Hyperbolic conservation laws and related analysis with applications*, edited by G.-Q. G. Chen et al., Springer Proc. Math. Stat. **49**, Springer, 2014. MR Zbl
- [Karper et al. 2015] T. K. Karper, A. Mellet, and K. Trivisa, “Hydrodynamic limit of the kinetic Cucker–Smale flocking model”, *Math. Models Methods Appl. Sci.* **25**:1 (2015), 131–163. MR Zbl
- [Mellet and Vasseur 2008] A. Mellet and A. Vasseur, “Asymptotic analysis for a Vlasov–Fokker–Planck/compressible Navier–Stokes system of equations”, *Comm. Math. Phys.* **281**:3 (2008), 573–596. MR Zbl
- [Motsch and Tadmor 2011] S. Motsch and E. Tadmor, “A new model for self-organized dynamics and its flocking behavior”, *J. Stat. Phys.* **144**:5 (2011), 923–947. MR Zbl
- [Poupaud and Rasclé 1997] F. Poupaud and M. Rasclé, “Measure solutions to the linear multi-dimensional transport equation with non-smooth coefficients”, *Comm. Partial Differential Equations* **22**:1-2 (1997), 337–358. MR Zbl
- [Poyato and Soler 2017] D. Poyato and J. Soler, “Euler-type equations and commutators in singular and hyperbolic limits of kinetic Cucker–Smale models”, *Math. Models Methods Appl. Sci.* **27**:6 (2017), 1089–1152. MR Zbl
- [Silk et al. 1983] J. Silk, A. Szalay, and Y. B. Zeldovich, “Large-scale structure of the universe”, *Sci. Amer.* **249**:4 (1983), 72–80.
- [Tadmor and Tan 2014] E. Tadmor and C. Tan, “Critical thresholds in flocking hydrodynamics with non-local alignment”, *Philos. Trans. R. Soc. Lond. Ser. A Math. Phys. Eng. Sci.* **372**:2028 (2014), art. id. 20130401. MR Zbl
- [Vasseur 2008] A. F. Vasseur, “Recent results on hydrodynamic limits”, pp. 323–376 in *Handbook of differential equations: evolutionary equations, IV*, edited by C. M. Dafermos and M. Pokorný, Elsevier/North-Holland, Amsterdam, 2008. MR Zbl
- [Vicsek et al. 1995] T. Vicsek, A. Czirók, E. Ben-Jacob, I. Cohen, and O. Shochet, “Novel type of phase transition in a system of self-driven particles”, *Phys. Rev. Lett.* **75**:6 (1995), 1226–1229. MR
- [Villani 2009] C. Villani, *Optimal transport: old and new*, Grundlehren der Mathematischen Wissenschaften **338**, Springer, 2009. MR Zbl
- [Weinan et al. 1996] E. Weinan, Y. G. Rykov, and Y. G. Sinai, “Generalized variational principles, global weak solutions and behavior with random initial data for systems of conservation laws arising in adhesion particle dynamics”, *Comm. Math. Phys.* **177**:2 (1996), 349–380. MR Zbl
- [Zavlanos et al. 2011] M. Zavlanos, M. Egerstedt, and G. J. Pappas, “Graph-theoretic connectivity control of mobile robot networks”, *Proc. IEEE* **99**:9 (2011), 1525–1540.
- [Zeldovich 1970] Y. B. Zeldovich, “Gravitational instability: an approximate theory for large density perturbations”, *Astron. Astrophys.* **5**:1 (1970), 84–89.

Received 22 Jan 2018. Revised 23 Apr 2018. Accepted 29 Jun 2018.

ALESSIO FIGALLI: alessio.figalli@math.ethz.ch  
 Department of Mathematics, ETH Zürich, Zürich, Switzerland

MOON-JIN KANG: moonjinkang@sookmyung.ac.kr  
 Department of Mathematics & Research Institute of Natural Sciences, Sookmyung Women’s University, Seoul, South Korea

# Analysis & PDE

msp.org/apde

## EDITORS

EDITOR-IN-CHIEF

Patrick Gérard

patrick.gerard@math.u-psud.fr

Université Paris Sud XI

Orsay, France

## BOARD OF EDITORS

Massimiliano Berti	Scuola Intern. Sup. di Studi Avanzati, Italy berti@sissa.it	Clément Mouhot	Cambridge University, UK c.mouhot@dpmms.cam.ac.uk
Sun-Yung Alice Chang	Princeton University, USA chang@math.princeton.edu	Werner Müller	Universität Bonn, Germany mueller@math.uni-bonn.de
Michael Christ	University of California, Berkeley, USA mchrist@math.berkeley.edu	Gilles Pisier	Texas A&M University, and Paris 6 pisier@math.tamu.edu
Alessio Figalli	ETH Zurich, Switzerland alessio.figalli@math.ethz.ch	Tristan Rivière	ETH, Switzerland riviere@math.ethz.ch
Charles Fefferman	Princeton University, USA cf@math.princeton.edu	Igor Rodnianski	Princeton University, USA irod@math.princeton.edu
Ursula Hamenstaedt	Universität Bonn, Germany ursula@math.uni-bonn.de	Sylvia Serfaty	New York University, USA serfaty@cims.nyu.edu
Vaughan Jones	U.C. Berkeley & Vanderbilt University vaughan.f.jones@vanderbilt.edu	Yum-Tong Siu	Harvard University, USA siu@math.harvard.edu
Vadim Kaloshin	University of Maryland, USA vadim.kaloshin@gmail.com	Terence Tao	University of California, Los Angeles, USA tao@math.ucla.edu
Herbert Koch	Universität Bonn, Germany koch@math.uni-bonn.de	Michael E. Taylor	Univ. of North Carolina, Chapel Hill, USA met@math.unc.edu
Izabella Laba	University of British Columbia, Canada ilaba@math.ubc.ca	Gunther Uhlmann	University of Washington, USA gunther@math.washington.edu
Gilles Lebeau	Université de Nice Sophia Antipolis, France lebeau@unice.fr	András Vasy	Stanford University, USA andras@math.stanford.edu
Richard B. Melrose	Massachusetts Inst. of Tech., USA rbm@math.mit.edu	Dan Virgil Voiculescu	University of California, Berkeley, USA dvv@math.berkeley.edu
Frank Merle	Université de Cergy-Pontoise, France Frank.Merle@u-cergy.fr	Steven Zelditch	Northwestern University, USA zelditch@math.northwestern.edu
William Minicozzi II	Johns Hopkins University, USA minicozz@math.jhu.edu	Maciej Zworski	University of California, Berkeley, USA zworski@math.berkeley.edu

## PRODUCTION

production@msp.org

Silvio Levy, Scientific Editor

---

See inside back cover or [msp.org/apde](http://msp.org/apde) for submission instructions.

---

The subscription price for 2019 is US \$310/year for the electronic version, and \$520/year (+\$60, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues from the last three years and changes of subscriber address should be sent to MSP.


---

Analysis & PDE (ISSN 1948-206X electronic, 2157-5045 printed) at Mathematical Sciences Publishers, 798 Evans Hall #3840, c/o University of California, Berkeley, CA 94720-3840, is published continuously online. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices.

---

APDE peer review and production are managed by EditFlow<sup>®</sup> from MSP.

PUBLISHED BY

 **mathematical sciences publishers**  
nonprofit scientific publishing

<http://msp.org/>

© 2019 Mathematical Sciences Publishers

# ANALYSIS & PDE

Volume 12 No. 3 2019

---

The BMO-Dirichlet problem for elliptic systems in the upper half-space and quantitative characterizations of VMO	605
JOSÉ MARÍA MARTELL, DORINA MITREA, IRINA MITREA and MARIUS MITREA	
Convergence of the Kähler–Ricci iteration	721
TAMÁS DARVAS and YANIR A. RUBINSTEIN	
Concentration of ground states in stationary mean-field games systems	737
ANNALISA CESARONI and MARCO CIRANT	
Generalized crystalline evolutions as limits of flows with smooth anisotropies	789
ANTONIN CHAMBOLLE, MASSIMILIANO MORINI, MATTEO NOVAGA and MARCELLO PONSIGLIONE	
Global weak solutions of the Teichmüller harmonic map flow into general targets	815
MELANIE RUPFLIN and PETER M. TOPPING	
A rigorous derivation from the kinetic Cucker–Smale model to the pressureless Euler system with nonlocal alignment	843
ALESSIO FIGALLI and MOON-JIN KANG	



2157-5045(2019)12:3;1-N