

ANALYSIS & PDE

Volume 12

No. 4

2019

STEPHEN CAMERON

**GLOBAL WELL-POSEDNESS
FOR THE TWO-DIMENSIONAL MUSKAT PROBLEM
WITH SLOPE LESS THAN 1**

GLOBAL WELL-POSEDNESS FOR THE TWO-DIMENSIONAL MUSKAT PROBLEM WITH SLOPE LESS THAN 1

STEPHEN CAMERON

We prove the existence of global, smooth solutions to the two-dimensional Muskat problem in the stable regime whenever the product of the maximal and minimal slope is less than 1. The curvature of these solutions decays to 0 as t goes to infinity, and they are unique when the initial data is $C^{1,\epsilon}$. We do this by getting a priori estimates using a nonlinear maximum principle first introduced in a paper by Kiselev, Nazarov, and Volberg (2007), where the authors proved global well-posedness for the surface quasigeostrophic equation.

1. Introduction

The Muskat problem was originally introduced in [Muskat 1934] in order to model the interface between water and oil in tar sands. In general, it describes the interface between two incompressible, immiscible fluids of different constant densities in a porous media. The fluids evolve according to Darcy's law, giving an evolution of the interface (see [Córdoba and Gancedo 2007] for derivation of equations), and the problem in two dimensions is analogous to the two-phase Hele-Shaw cell (see [Saffman and Taylor 1958]). In the case that the two fluids are of equal viscosity and the interface is given by the graph $y = f(t, x)$ with the denser fluid on bottom (i.e., the stable regime), the function f satisfies

$$f_t(t, x) = \int_{\mathbb{R}} \frac{(f_x(t, y) - f_x(t, x))(y - x)}{(f(t, y) - f(t, x))^2 + (y - x)^2} dy, \quad (1-1)$$

after the appropriate renormalization. By making a change of variables, see the proof of Lemma 5.1 of [Córdoba and Gancedo 2009], we get the equivalent system

$$f_t(t, x) = \int_{\mathbb{R}} \frac{f(t, y) - f(t, x) - (y - x)f_x(t, x)}{(f(t, y) - f(t, x))^2 + (y - x)^2} dy, \quad (1-2)$$

which will be more useful for our purposes. Since the function f is Lipschitz, the above integral can be viewed as a nonlinear perturbation of the half Laplacian. In fact, it is easy to see that linearizing around a flat solution gives

$$f_t(t, x) = -c(-\Delta)^{1/2} f(t, x), \quad (1-3)$$

demonstrating the natural parabolicity of the problem.

MSC2010: 35K55, 35Q35, 35R09.

Keywords: Muskat problem, porous media, fluid interface, global well-posedness.

The Muskat problem is known to be locally well-posed in H^k for $k \geq 3$ with solutions satisfying L^∞ and L^2 maximum principles, but neither imply any gain of derivatives; see [Córdoba and Gancedo 2009; Constantin et al. 2013].

Under the assumption $\|f'_0\|_{L^\infty} < 1$, there have been a number of positive results. In [Constantin et al. 2013] the authors proved an L^∞ maximal principle for the slope f_x along with the existence of global weak Lipschitz solutions using a regularized system. Recently, [Gancedo 2017] improved the L^2 energy estimate of [Constantin et al. 2013] (which holds for any solution) to one analogous with the energy estimate from the linear equation under this assumption on the slope. When the initial data f_0 is in $H^2(\mathbb{R})$ with $\|f_0\|_1 = \|\xi|\hat{f}_0(\xi)\|_{L^1_\xi}$ less than some explicit constant $\approx \frac{1}{3}$ (which implies slope less than 1), [Constantin et al. 2016] proved that a unique global strong solution exists. In this case [Patel and Strain 2017] proved optimal decay estimates on the norms $\|f(t, \cdot)\|_s = \|\xi|^s \hat{f}(t, \xi)\|_{L^1_\xi}$, matching the estimates for the linear equation.

Recently, [Deng et al. 2017] was also able to prove the existence of global weak solutions for arbitrarily large monotonic initial data. They did this using the regularized system from [Constantin et al. 2013] to prove that both f and f_x still obey the maximum principle under this monotonicity assumption.

Because solutions to (1-2) have the natural scaling $(1/r)f(rt, rx)$, we see that L^∞ or sign bounds on the slope f_x are scale-invariant properties. We fit these two types of assumptions into the same framework by showing that the critical quantity is in fact the product of the maximal and minimal slopes,

$$\beta(f'_0) := \left(\sup_x f'_0(x)\right) \left(\sup_y -f'_0(y)\right). \quad (1-4)$$

As we shall see in Section 3, the derivative f_x obeys the equation

$$(f_x)_t(t, x) = f_{xx}(t, x) \int_{\mathbb{R}} \frac{-h}{\delta_h f(t, x)^2 + h^2} dh + \int_{\mathbb{R}} \delta_h f_x(t, x) K(t, x, h) dh, \quad (1-5)$$

where $\delta_h f(t, x) := f(t, x+h) - f(t, x)$ and the kernel K is uniformly elliptic of order 1 whenever $\beta(f'_0) < 1$. Thus we naturally get regularizing effects from the equation whenever the initial data satisfies this bound. It's clear that $\|f'_0\|_{L^\infty} < 1$ implies $\beta(f'_0) < 1$, and for bounded monotonic data we get $\beta(f'_0) = 0$ since either $\sup f'_0 = 0$ or $\inf f'_0 = 0$. Thus this $\beta(f'_0) < 1$ provides a natural interpolation between these two types of assumptions.

In contrast to the positive results, [Castro et al. 2012] showed that there is an open subset of initial data in H^4 such that the Rayleigh–Taylor condition breaks down in finite time. That is,

$$\lim_{t \rightarrow t_0-} \|f_x(t, \cdot)\|_{L^\infty} = \infty$$

for some time t_0 , after which the interface between the fluids can no longer be described by a graph.

The authors of [Constantin et al. 2017] made great progress towards proving global regularity. They proved that if the initial data $f_0 \in H^k$, then the solution f will exist and remain in H^k so long as the slope $f_x(t, \cdot)$ remains bounded and uniformly continuous. Thus the natural next step is to prove the generation of a modulus of continuity for f_x .

Theorem 1.1. *Let $f_0 \in W^{1,\infty}(\mathbb{R})$ with*

$$\beta(f'_0) := \left(\sup_x f'_0(x) \right) \left(\sup_y -f'_0(y) \right) < 1. \quad (1-6)$$

Then there exists a classical solution

$$f \in C([0, \infty) \times \mathbb{R}) \cap C_{\text{loc}}^{1,\alpha}((0, \infty) \times \mathbb{R}) \cap L_{\text{loc}}^\infty((0, \infty); C^{1,1}) \quad (1-7)$$

to (1-2) with f_x satisfying both the maximum principle and

$$f_x(t, x) - f_x(t, y) \leq \rho \left(\frac{|x - y|}{t} \right), \quad t > 0, \quad x \neq y \in \mathbb{R}, \quad (1-8)$$

for some Lipschitz modulus of continuity ρ depending solely on $\beta(f'_0)$, $\|f'_0\|_{L^\infty}$. In the case that $f_0 \in C^{1,\epsilon}(\mathbb{R})$ for some $\epsilon > 0$, the solution f is unique with $f \in L^\infty([0, \infty); C^{1,\epsilon})$.

The uniqueness statement follows essentially from the uniqueness theorem of [Constantin et al. 2017]. We note in the Appendix the few small changes needed to their proof in order to apply it here.

The most vital part of Theorem 1.1 is the spontaneous generation of the modulus $\rho(\cdot/t)$, as everything else will follow from that. The spontaneous generation/propagation of a general modulus of continuity has old roots as classical Hölder estimates, but it's only recently that the idea to tailor-make moduli for specific equations emerged. The technique first appeared in [Kiselev et al. 2007], where the authors used it to prove global well-posedness for the surface quasigeostrophic equation. It has had great success at proving regularity for a number of active scalar equations, that is, equations of the form

$$\theta_t + (u \cdot \nabla)\theta + \mathcal{L}\theta = 0, \quad (1-9)$$

where u is a flow depending on θ and $\mathcal{L}$ is some diffusive operator. See [Kiselev 2010; Dabkowski et al. 2014] for a good overview of results using this method.

To date, these tailor-made moduli have only been applied to cases where all the nonlinearity is in the flow velocity u , and the diffusive term $\mathcal{L}$ is rather nice (typically $(-\Delta)^\alpha$, or at least a Fourier multiplier). We will be applying this method to f_x , which solves the active scalar equation (1-5). Note that in this equation, the kernel K defined in (3-4) is a highly nonlinear function of f , f_x . Thus this is the first time the method has been applied in a fully nonlinear equation.

We prove Theorem 1.1 by deriving a priori estimates for smooth solutions to (1-2) with initial data $f_0 \in C_c^\infty(\mathbb{R})$ depending primarily on $\beta(f'_0)$, $\|f'_0\|_{L^\infty}$. We prove enough estimates that by approximating in $W_{\text{loc}}^{1,\infty}$ with smooth compactly supported initial data, we get solutions f^ϵ which will converge along subsequences in C_{loc}^1 to a solution f solving (1-2) for arbitrary initial data $f_0 \in W^{1,\infty}(\mathbb{R})$ with $\beta(f'_0) < 1$.

The rest of the paper is organized as follows. We begin by repeating the breakthrough argument of [Kiselev et al. 2007] in Section 2. In Section 3, we differentiate (1-2) to derive the equation for f_x , showing that it satisfies the maximum principle when $\beta(f'_0) < 1$. In Section 4, we state how a modulus of continuity ω interacts with the equation in our main technical lemma. In Sections 5 and 6 we then derive the bounds on the drift and diffusion terms necessary to prove that lemma. In Section 7, we apply our main technical lemma to a specific modulus of continuity, and finally in Section 8 we complete the proof

of (1-8) by choosing the correct modulus ρ . In Section 9, we then use (1-8) to prove a few estimates on regularity in time, guaranteeing enough compactness to prove that there are classical solutions for rough initial data. Finally in the Appendix, we give a quick outline for how to modify the uniqueness proof of [Constantin et al. 2017] to work for initial data $f_0 \in C^{1,\epsilon}(\mathbb{R})$ with $\beta(f'_0) < 1$.

2. Breakthrough scenario

Assume that $f_0 \in C_c^\infty(\mathbb{R})$, with $\beta(f'_0) < 1$, so that there exists a solution $f \in C^1((0, T_+); H^k)$ for k arbitrarily large and some $T_+ > 0$ by [Córdoba and Gancedo 2009]. Note that under the assumption that $\beta(f'_0) < 1$, we will show that the maximum principle holds (see Proposition 3.1) and hence $\|f_x\|_{L^\infty([0, T_+] \times \mathbb{R})} \leq \|f'_0\|_{L^\infty}$ is uniformly bounded. Fix a Lipschitz modulus ρ which we will define later. For sufficiently small times, $f_x(t, \cdot)$ will have modulus $\rho(\cdot/t)$ since it is smooth and bounded. It then follows by the main theorem of [Constantin et al. 2017] that as long as $f_x(t, \cdot)$ continues to have modulus $\rho(\cdot/t)$, the solution f will exist with $T_+ > t$.

So, we proceed as in the proof for the quasigeostrophic equation in [Kiselev et al. 2007]. Suppose that $f_x(t, \cdot)$ satisfies (1-8) for all $t < T$. Then by continuity,

$$f_x(T, x) - f_x(T, y) \leq \rho\left(\frac{|x - y|}{T}\right) \quad \text{for all } x \neq y \in \mathbb{R}. \quad (2-1)$$

We first prove that if we have the strict inequality $f_x(T, x) - f_x(T, y) < \rho(|x - y|/T)$, then $f_x(t, \cdot)$ will have modulus $\rho(\cdot/t)$ for $t \leq T + \epsilon$.

Lemma 2.1. *Let $f \in C([0, T_+]; C_0^3(\mathbb{R}))$ and $T \in (0, T_+)$. Suppose that $f(T, \cdot)$ satisfies*

$$f_x(T, x) - f_x(T, y) < \rho\left(\frac{|x - y|}{T}\right) \quad \text{for all } x \neq y \in \mathbb{R}, \quad (2-2)$$

for some Lipschitz modulus of continuity ρ with $\rho''(0) = -\infty$. Then

$$f_x(T + \epsilon, x) - f_x(T + \epsilon, y) < \rho\left(\frac{|x - y|}{T + \epsilon}\right) \quad \text{for all } x \neq y \in \mathbb{R}, \quad (2-3)$$

for all $\epsilon > 0$ sufficiently small.

Proof. To begin, note that for any compact subset $K \subset \mathbb{R}^2 \setminus \{(x, x) : x \in \mathbb{R}\}$,

$$\begin{aligned} f_x(T, x) - f_x(T, y) &< \rho\left(\frac{|x - y|}{T}\right) \quad \text{for all } (x, y) \in K \\ \Rightarrow f_x(T + \epsilon, x) - f_x(T + \epsilon, y) &< \rho\left(\frac{|x - y|}{T + \epsilon}\right) \quad \text{for all } (x, y) \in K, \end{aligned} \quad (2-4)$$

for $\epsilon > 0$ sufficiently small by uniform continuity. So, we only need to focus on pairs (x, y) that are either close to the diagonal, or that are large.

To handle (x, y) near the diagonal, we start by noting that $f(T, \cdot) \in C^3(\mathbb{R})$ and $\rho''(0) = -\infty$. Thus for every x we get

$$|f_{xx}(T, x)| < \frac{\rho'(0)}{T}. \quad (2-5)$$

Since $f \in C([0, T_+); C_0^3(\mathbb{R}))$, we have $f_{xx}(T, x) \rightarrow 0$ as $x \rightarrow \infty$. Thus we can take the point where $\max_x |f_{xx}(T, x)|$ is achieved to get

$$\|f_{xx}(T, \cdot)\|_{L^\infty} < \frac{\rho'(0)}{T}. \quad (2-6)$$

By the continuity of f_{xx} , we thus have $\|f_{xx}(T + \epsilon, \cdot)\|_{L^\infty} < \rho'(0)/(T + \epsilon)$ for $\epsilon > 0$ sufficiently small. Hence,

$$f_x(T + \epsilon, x) - f_x(T + \epsilon, y) < \rho\left(\frac{|x - y|}{T + \epsilon}\right), \quad |x - y| < \delta, \quad (2-7)$$

for ϵ, δ sufficiently small.

Now let $R_1, R_2 > 0$ be such that

$$\rho(R_1/(T + \epsilon)) > \text{osc}_{\mathbb{R}} f_x(T + \epsilon, \cdot) \quad (2-8)$$

and that $|x| > R_2$ implies

$$|f_x(T + \epsilon, x)| < \frac{1}{2}\rho\left(\frac{\delta}{T + \epsilon}\right) \quad (2-9)$$

for $\epsilon > 0$ sufficiently small. Taking $R = R_1 + R_2$, it's easy to check that $|x| > R$ implies

$$|f_x(T + \epsilon, x) - f_x(T + \epsilon, y)| < \rho\left(\frac{|x - y|}{T + \epsilon}\right) \quad \text{for all } y \neq x. \quad (2-10)$$

Finally, taking

$$K = \{(x, y) \in \mathbb{R}^2 : |x - y| \geq \delta, x, y \in \bar{B}_R\},$$

we're done. □

Thus by [Lemma 2.1](#), if f_x was to lose its modulus after time T , we must have that there exist $x \neq y \in \mathbb{R}$ with

$$f_x(T, x) - f_x(T, y) = \rho\left(\frac{|x - y|}{T}\right). \quad (2-11)$$

We will show for a smooth solution f of (1-2) and the correct choice of ρ that in this case

$$\frac{d}{dt}(f_x(t, x) - f_x(t, y))\Big|_{t=T} < \frac{d}{dt}\left(\rho\left(\frac{|x - y|}{t}\right)\right)\Big|_{t=T}, \quad (2-12)$$

contradicting the fact that f_x had modulus $\rho(\cdot/t)$ for time $t < T$.

Thus we just need to prove (2-12) to complete the proof of the generation of modulus of continuity (1-8) of [Theorem 1.1](#).

3. Equation for f_x

To begin proving (2-12), we need to examine the equation that f_x solves. Since everything we will be doing is for some fixed time $T > 0$, we will suppress the time variable from now on. Differentiating (1-2),

we see that f_x solves

$$(f_x)_t(x) = f_{xx}(x) \int_{\mathbb{R}} \frac{x-y}{(f(y)-f(x))^2+(y-x)^2} dy + \int_{\mathbb{R}} (f(y)-f(x)-(y-x)f_x(x)) \frac{2((f(y)-f(x))f_x(x)+(y-x))}{((f(y)-f(x))^2+(y-x)^2)^2} dy. \quad (3-1)$$

To simplify notation, we reparametrize (3-1) by taking $y = x + h$, and letting

$$\delta_h f(x) := f(x+h) - f(x),$$

we get

$$(f_x)_t(x) = f_{xx}(x) \int_{\mathbb{R}} \frac{-h}{(\delta_h f(x))^2 + h^2} dh + \int_{\mathbb{R}} (\delta_h f(x) - hf_x(x)) \frac{2(\delta_h f(x)f_x(x) + h)}{(\delta_h f(x)^2 + h^2)^2} dh. \quad (3-2)$$

Note that

$$\delta_h f(x) - hf_x(x) = \begin{cases} \int_0^h \delta_s f_x(x) ds & \text{for } h > 0, \\ -\int_h^0 \delta_s f_x(x) ds & \text{for } h < 0. \end{cases}$$

With that in mind, define

$$k(x, s) = \frac{2(\delta_s f(x)f_x(x) + s)}{(\delta_s f(x)^2 + s^2)^2}, \quad (3-3)$$

and

$$K(x, h) = \begin{cases} \int_h^\infty k(x, s) ds, & h > 0, \\ \int_{-\infty}^h -k(x, s) ds, & h < 0. \end{cases} \quad (3-4)$$

Then integrating (3-2) by parts, we have that f_x solves the equation

$$(f_x)_t(x) = f_{xx}(x) \int_{\mathbb{R}} \frac{-h}{\delta_h f(x)^2 + h^2} dh + \int_{\mathbb{R}} \delta_h f_x(x) K(x, h) dh. \quad (3-5)$$

As

$$\frac{-\beta(f_x)}{s} \leq \frac{f_x(x)\delta_s f(x)}{s} \leq \frac{\|f_x\|_{L^\infty}^2}{s}, \quad (3-6)$$

we see that

$$\frac{2(1-\beta(f_x))}{(1+\|f_x\|_{L^\infty}^2)^2} \frac{1}{|s|^3} \leq \text{sgn}(s)k(x, s) \leq \frac{2(1+\|f_x\|_{L^\infty}^2)}{|s|^3},$$

and hence

$$\frac{1-\beta(f_x)}{(1+\|f_x\|_{L^\infty}^2)^2} \frac{1}{h^2} \leq K(x, h) \leq \frac{1+\|f_x\|_{L^\infty}^2}{h^2}. \quad (3-7)$$

Thus in the case that $\beta(f_x) \leq 1$, we then have that the kernel K is a nonnegative, from which we get immediately:

Proposition 3.1 (maximum principle). *Let f_x be a sufficiently smooth solution to (3-5) with $\beta(f'_0) \leq 1$. Then for any $0 \leq s \leq t$, we have*

$$\inf_y f_x(s, y) \leq \inf_y f_x(t, y) \leq \sup_y f_x(t, y) \leq \sup_y f_x(s, y). \quad (3-8)$$

In particular, since $\beta(f'_0) < 1$, the maximum principle tells us that

$$\beta(f_x) \leq \beta(f'_0) < 1, \quad \|f_x\|_{L^\infty} \leq \|f'_0\|_{L^\infty} < \infty. \quad (3-9)$$

Thus we get

$$0 < \frac{\lambda}{h^2} \leq K(x, h) \leq \frac{\Lambda}{h^2}, \quad (3-10)$$

where

$$\lambda = \frac{1 - \beta(f'_0)}{(1 + \|f'_0\|_{L^\infty}^2)^2}, \quad \Lambda = 1 + \|f'_0\|_{L^\infty}^2. \quad (3-11)$$

Thus K is comparable to the kernel for $(-\Delta)^{1/2}$, so f_x solves the uniformly elliptic equation (3-5). Note that the sole reason we require $\beta(f'_0) < 1$ is to ensure this ellipticity of K .

4. Moduli estimates

Our goal is to show that if $f_x(T, \cdot)$ has modulus $\rho(\cdot/T)$ and equality is achieved at two points (2-11), then (2-12) must hold, contradicting the assumptions of the breakthrough argument (see Section 2). To that end, we first need to understand how a modulus of continuity interacts with the equation for f_x (3-5). Hence:

Lemma 4.1. *Let $f : [0, \infty) \times \mathbb{R} \rightarrow \mathbb{R}$ be a bounded smooth solution to (1-2) with $\beta(f'_0) < 1$, and $\omega : [0, \infty) \rightarrow [0, \infty)$ be some fixed modulus of continuity. Assume that at some fixed time T*

$$\begin{aligned} \delta_h f_x(T, x) &\leq \omega(|h|), \\ f_x(T, \xi/2) - f_x(T, -\xi/2) &= \omega(\xi) \end{aligned} \quad (4-1)$$

for all $h \in \mathbb{R}$ and for some $\xi > 0$. Then

$$\begin{aligned} &\frac{d}{dt}(f_x(t, \xi/2) - f_x(t, -\xi/2))\big|_{t=T} \\ &\leq A\omega'(\xi) \left(\int_0^\xi \frac{\omega(h)}{h} dh + \xi \int_\xi^\infty \frac{\omega(h)}{h^2} dh + \ln(M+1)\omega(\xi) \right) \\ &\quad + A\omega(\xi) \int_{M\xi}^\infty \frac{\omega(h)}{h^2} dh + 2(\Lambda - \lambda) \int_\xi^{M\xi} \frac{(\omega(h - \xi) - \omega(\xi))_+}{h^2} dh \\ &\quad + 2\lambda \int_0^\xi \frac{\delta_h \omega(\xi) + \delta_{-h} \omega(\xi)}{h^2} dh + 2\lambda \int_\xi^\infty \frac{\omega(h + \xi) - \omega(h) - \omega(\xi)}{h^2} dh \end{aligned} \quad (4-2)$$

for any $M \geq 1$, where A depends only on $\|f'_0\|_{L^\infty}$ and λ, Λ are as in (3-11).

This is the main technical lemma that we need. Since solutions to (1-2) are closed under translation and sign change, it suffices to consider the above situation for our proof of (2-12).

Note that (4-2) holds for any value of the parameter $M \geq 1$. Later in Lemma 6.1, we will essentially use two different values of M depending on the size of ξ . In the small ξ regime we can simply take $M = 1$, but in the large ξ regime we will need to take M to be a sufficiently large constant depending only on initial data (but not on the exact size of ξ) in order to control the size of the error term

$$\omega(\xi) \int_{M\xi}^{\infty} \frac{\omega(h)}{h^2} dh.$$

The proof for Lemma 4.1 is essentially a nondivergence form argument; our function f_x is touched from above at $\xi/2$ by our modulus ω , and it is touched from below at $-\xi/2$ by $-\omega$. Specifically,

$$\begin{aligned} \delta_h f_x(\xi/2) &\leq \delta_h \omega(\xi) & \text{for all } h > -\xi, \\ \delta_h f_x(-\xi/2) &\geq -\delta_{-h} \omega(\xi) & \text{for all } h < \xi. \end{aligned} \quad (4-3)$$

From (4-3), we want to derive as much information as we can and bound $\frac{d}{dt}(f_x(\xi/2) - f_x(-\xi/2))$. To that end, by dividing (4-3) through by h and taking the limit as $h \rightarrow 0$, we then get

$$f_{xx}(\xi/2) = f_{xx}(-\xi/2) = \omega'(\xi). \quad (4-4)$$

Hence by our equation for f_x (3-5), we have

$$\begin{aligned} &\frac{d}{dt}(f_x(\xi/2) - f_x(-\xi/2)) \\ &= \omega'(\xi) \int_{\mathbb{R}} \left(\frac{-h}{\delta_h f(\xi/2)^2 + h^2} - \frac{-h}{\delta_h f(-\xi/2)^2 + h^2} \right) dh \\ &\quad + \int_{\mathbb{R}} \delta_h f_x(\xi/2) K(\xi/2, h) - \delta_h f_x(-\xi/2) K(-\xi/2, h) dh \\ &= \omega'(\xi) \int_{\mathbb{R}} \left(\frac{-h}{\delta_h f(\xi/2)^2 + h^2} - \frac{-h}{\delta_h f(-\xi/2)^2 + h^2} \right) dh + \omega'(\xi) \int_{-M\xi}^{M\xi} (hK(\xi/2, h) - hK(-\xi/2, h)) dh \\ &\quad + \int_{-M\xi}^{M\xi} (\delta_h f_x(\xi/2) - h\omega'(\xi)) K(\xi/2, h) - (\delta_h f_x(-\xi/2) - h\omega'(\xi)) K(-\xi/2, h) dh \\ &\quad + \int_{|h| > M\xi} \delta_h f_x(\xi/2) K(\xi/2, h) - \delta_h f_x(-\xi/2) K(-\xi/2, h) dh \end{aligned} \quad (4-5)$$

for any $M \geq 1$. The first two terms of the far right-hand side of (4-5) act as a drift, giving rise to the first two error terms of (4-2). The latter two terms of (4-5) act as a diffusion, giving rise to both the helpful (negative) terms in (4-2), as well as additional error terms (the middle terms of (4-2)) arising from the difference in the kernels, $|K(\xi/2, h) - K(-\xi/2, h)|$.

5. Bounds on drift terms

We begin proving [Lemma 4.1](#) by bounding the drift terms of (4-5), starting with:

Lemma 5.1. *Under the assumptions of [Lemma 4.1](#),*

$$\omega'(\xi) \left| \int_{\mathbb{R}} \frac{-h}{\delta_h f(\xi/2)^2 + h^2} - \frac{-h}{\delta_h f(-\xi/2)^2 + h^2} dh \right| \lesssim \omega'(\xi) \left(\int_0^\xi \frac{\omega(h)}{h} dh + \xi \int_\xi^\infty \frac{\omega(h)}{h^2} dh \right). \quad (5-1)$$

Proof. We want to bound (5-1) by symmetrizing the kernels for $|h| < \xi$ and then using the continuity in the first variable for $|h| > \xi$. To that end,

$$\begin{aligned} & \omega'(\xi) \int_{\mathbb{R}} \left(\frac{-h}{\delta_h f(\xi/2)^2 + h^2} - \frac{-h}{\delta_h f(-\xi/2)^2 + h^2} \right) dh \\ & \leq \omega'(\xi) \int_0^\xi h \left| \frac{\delta_h f(\xi/2)^2 - \delta_{-h} f(\xi/2)^2}{(\delta_h f(\xi/2)^2 + h^2)(\delta_{-h} f(\xi/2)^2 + h^2)} + \frac{\delta_h f(-\xi/2)^2 - \delta_{-h} f(-\xi/2)^2}{(\delta_h f(-\xi/2)^2 + h^2)(\delta_{-h} f(-\xi/2)^2 + h^2)} \right| dh \\ & \quad + \omega'(\xi) \int_{|h| > \xi} |h| \left| \frac{\delta_h f(\xi/2)^2 - \delta_{-h} f(-\xi/2)^2}{(\delta_h f(\xi/2)^2 + h^2)(\delta_{-h} f(-\xi/2)^2 + h^2)} \right| dh. \end{aligned} \quad (5-2)$$

We bound the first integral using

$$\begin{aligned} & |\delta_h f(x)| \lesssim |h|, \\ & |\delta_h f(x) + \delta_{-h} f(x)| = \left| \int_0^h f_x(x+s) - f_x(x+s-h) ds \right| \leq \omega(h)h. \end{aligned} \quad (5-3)$$

Thus we get, for $0 \leq h < \xi$,

$$\left| \frac{\delta_h f(x)^2 - \delta_{-h} f(x)^2}{(\delta_h f(x)^2 + h^2)(\delta_{-h} f(x)^2 + h^2)} \right| \lesssim \frac{\omega(h)}{h^2}, \quad (5-4)$$

and hence

$$\int_0^\xi h \left| \frac{\delta_h f(\xi/2)^2 - \delta_{-h} f(\xi/2)^2}{(\delta_h f(\xi/2)^2 + h^2)(\delta_{-h} f(\xi/2)^2 + h^2)} \right| dh \lesssim \int_0^\xi \frac{\omega(h)}{h} dh. \quad (5-5)$$

For $|h| \geq \xi$, we bound $|\delta_h f(\xi/2) + \delta_h f(-\xi/2)| \lesssim |h|$ and

$$\begin{aligned} & |\delta_h f(\xi/2) - \delta_h f(-\xi/2)| = \left| \int_0^h f_x(\xi/2+s) - f_x(-\xi/2+s) ds \right| \\ & = \left| \int_0^\xi f_x(h-\xi/2+s) - f_x(-\xi/2+s) ds \right| \leq \xi \omega(|h|), \end{aligned} \quad (5-6)$$

in order to get

$$\int_{|h|>\xi} |h| \left| \frac{\delta_h f(\xi/2)^2 - \delta_h f(-\xi/2)^2}{(\delta_h f(\xi/2)^2 + h^2)(\delta_h f(-\xi/2)^2 + h^2)} \right| dh \lesssim \xi \int_{\xi}^{\infty} \frac{\omega(h)}{h^2} dh. \quad (5-7)$$

Putting (5-5) and (5-7) together, we thus have

$$\omega'(\xi) \int_{\mathbb{R}} \left(\frac{-h}{\delta_h f(\xi/2)^2 + h^2} - \frac{-h}{\delta_h f(-\xi/2)^2 + h^2} \right) dh \lesssim \omega'(\xi) \left(\int_0^{\xi} \frac{\omega(h)}{h} dh + \xi \int_{\xi}^{\infty} \frac{\omega(h)}{h^2} dh \right), \quad (5-8)$$

completing the proof. $\square$

That leaves us with the second drift term of (4-5):

Lemma 5.2. *Under the assumptions of Lemma 4.1, for any $M \geq 1$*

$$\omega'(\xi) \left| \int_{-M\xi}^{M\xi} h K(\xi/2, h) - h K(-\xi/2, h) dh \right| \lesssim \omega'(\xi) \left(\int_0^{\xi} \frac{\omega(h)}{h} dh + \xi \int_{\xi}^{\infty} \frac{\omega(h)}{h^2} dh + \ln(M+1) \omega(\xi) \right). \quad (5-9)$$

Proof. To begin, we note

$$\begin{aligned} \omega'(\xi) \left| \int_{-M\xi}^{M\xi} h K(\xi/2, h) - h K(-\xi/2, h) dh \right| \\ \leq \omega'(\xi) \int_0^{M\xi} h \left| K(\xi/2, h) - K(\xi/2, -h) - K(-\xi/2, h) + K(-\xi/2, -h) \right| dh. \end{aligned} \quad (5-10)$$

Recall the definition of K , (3-4),

$$K(x, h) = \begin{cases} \int_h^{\infty} k(x, s) ds, & h > 0, \\ \int_{-\infty}^h -k(x, s) ds, & h < 0, \end{cases} \quad k(x, s) = \frac{2(\delta_s f(x) f_x(x) + s)}{(\delta_s f(x)^2 + s^2)^2}. \quad (5-11)$$

So, to control (5-10) we first need to bound $|k(x, s) + k(x, -s)|$ for $0 \leq s < \xi$, and $|k(\xi/2, s) - k(-\xi/2, s)|$ for $|s| > \xi$. For the first, using the bounds (5-3) we see that

$$\begin{aligned} & |k(x, s) + k(x, -s)| \\ &= \left| \frac{2(\delta_s f(x) f_x(x) + s)}{(\delta_s f(x)^2 + s^2)^2} + \frac{2(\delta_{-s} f(x) f_x(x) - s)}{(\delta_{-s} f(x)^2 + s^2)^2} \right| \\ &\leq \frac{2|\delta_s f(x) + \delta_{-s} f(x)| |f_x(x)|}{(\delta_{-s} f(x)^2 + s^2)^2} + 2|\delta_s f(x) f_x(x) + s| \left| \frac{(\delta_s f(x)^2 + s^2)^2 - (\delta_{-s} f(x)^2 + s^2)^2}{(\delta_s f(x)^2 + s^2)^2 (\delta_{-s} f(x)^2 + s^2)^2} \right| \\ &\lesssim \frac{\omega(s)}{s^3} + s \left| \frac{\delta_s f(x)^4 - \delta_{-s} f(x)^4 + 2s^2(\delta_s f(x)^2 - \delta_{-s} f(x)^2)}{s^8} \right| \\ &\lesssim \frac{\omega(s)}{s^3}. \end{aligned} \quad (5-12)$$

For the second, using (5-3), (5-6), and (4-1) we get

$$\begin{aligned}
 |k(\xi/2, s) - k(-\xi/2, s)| &= \left| \frac{2(\delta_s f(\xi/2) f_x(\xi/2) + s)}{(\delta_s f(\xi/2)^2 + s^2)^2} - \frac{2(\delta_s f(-\xi/2) f_x(-\xi/2) + s)}{(\delta_s f(-\xi/2)^2 + s^2)^2} \right| \\
 &\leq 2 \frac{|\delta_s f(\xi/2) f_x(\xi/2) - \delta_s f(-\xi/2) f_x(-\xi/2)|}{(\delta_s f(-\xi/2)^2 + s^2)^2} \\
 &\quad + 2 |\delta_s f(\xi/2) f_x(\xi/2) + s| \left| \frac{(\delta_s f(\xi/2)^2 + s^2)^2 - (\delta_s f(-\xi/2)^2 + s^2)^2}{(\delta_s f(\xi/2)^2 + s^2)^2 (\delta_s f(-\xi/2)^2 + s^2)^2} \right| \\
 &\lesssim \frac{|\delta_s f(\xi/2) - \delta_s f(-\xi/2)| |f_x(\xi/2)|}{s^4} + \frac{|\delta_s f(-\xi/2)| |f_x(\xi/2) - f_x(-\xi/2)|}{s^4} \\
 &\quad + |s| \left| \frac{\delta_s f(\xi/2)^4 - \delta_s f(-\xi/2)^4 + s^2(\delta_s f(\xi/2)^2 - \delta_s f(-\xi/2)^2)}{s^8} \right| \\
 &\lesssim \frac{\xi \omega(s)}{s^4} + \frac{\omega(\xi)}{s^3}. \tag{5-13}
 \end{aligned}$$

So using (5-12) and (5-13), we can first bound

$$\begin{aligned}
 \int_0^\xi h |K(\xi/2, h) - K(\xi/2, -h) - K(-\xi/2, h) + K(-\xi/2, -h)| dh \\
 \lesssim \int_0^\xi h \int_h^\xi \frac{\omega(s)}{s^3} ds dh + \int_0^\xi h \int_\xi^\infty \frac{\xi \omega(s)}{s^4} + \frac{\omega(\xi)}{s^3} ds dh \\
 \lesssim \int_0^\xi \frac{\omega(s)}{s^3} \int_0^s h dh ds + \int_\xi^\infty \frac{\xi^3 \omega(s)}{s^4} + \frac{\xi^2 \omega(\xi)}{s^3} ds \\
 \lesssim \int_0^\xi \frac{\omega(s)}{s} ds + \xi \int_\xi^\infty \frac{\omega(s)}{s^2} ds + \omega(\xi). \tag{5-14}
 \end{aligned}$$

For the rest of (5-10), we use (5-13) again to also bound

$$\begin{aligned}
 \int_{M\xi > |h| > \xi} |h| |K(\xi/2, h) - K(-\xi/2, h)| dh &\lesssim \int_\xi^{M\xi} h \int_h^\infty \frac{\omega(\xi)}{s^3} + \frac{\xi \omega(s)}{s^4} ds dh \\
 &\lesssim \omega(\xi) \int_\xi^{M\xi} \frac{1}{h} dh + \xi \int_\xi^{M\xi} \frac{\omega(h)}{h^2} dh \\
 &\lesssim \ln(M) \omega(\xi) + \xi \int_\xi^\infty \frac{\omega(h)}{h^2} dh, \tag{5-15}
 \end{aligned}$$

completing the proof. $\square$

6. Bounds on diffusive terms

Now we move on to proving an upper bound for the diffusive terms of (4-5). We can rewrite them as

$$\begin{aligned}
 & \int_{-M\xi}^{M\xi} (\delta_h f_x(\xi/2) - h\omega'(\xi))K(\xi/2, h) - (\delta_h f_x(-\xi/2) - h\omega'(\xi))K(-\xi/2, h) dh \\
 & \quad + \int_{|h|>M\xi} \delta_h f_x(\xi/2)K(\xi/2, h) - \delta_h f_x(-\xi/2)K(-\xi/2, h) dh \\
 & = \int_{-M\xi}^{M\xi} (\delta_h f_x(\xi/2) - h\omega'(\xi))K(\xi/2, h) - (\delta_h f_x(-\xi/2) - h\omega'(\xi))K(-\xi/2, h) dh \\
 & \quad + \int_{|h|>M\xi} [\delta_h f_x(\xi/2) - \delta_h f_x(-\xi/2)]K(\xi/2, h) dh \\
 & \quad + \int_{|h|>M\xi} \delta_h f_x(-\xi/2)[K(\xi/2, h) - K(-\xi/2, h)] dh. \tag{6-1}
 \end{aligned}$$

We begin by bounding the last term, which is an error term.

Lemma 6.1. *Under the assumptions of Lemma 4.1,*

$$\left| \int_{|h|>M\xi} \delta_h f_x(-\xi/2)[K(\xi/2, h) - K(-\xi/2, h)] dh \right| \lesssim \omega(\xi) \int_{M\xi}^{\infty} \frac{\omega(h)}{h^2} dh + \omega'(\xi)\xi \int_{\xi}^{\infty} \frac{\omega(h)}{h^2} dh. \tag{6-2}$$

Proof. Using the fact that f_x has modulus ω and the bounds (5-13), it follows that

$$\begin{aligned}
 & \int_{|h|>M\xi} \delta_h f_x(-\xi/2)[K(\xi/2, h) - K(-\xi/2, h)] dh \\
 & \lesssim \int_{M\xi}^{\infty} \omega(h) \int_h^{\infty} \frac{\omega(\xi)}{s^3} + \frac{\xi\omega(s)}{s^4} ds dh \\
 & \lesssim \omega(\xi) \int_{M\xi}^{\infty} \frac{\omega(h)}{h^2} dh + \int_{M\xi}^{\infty} \omega(h) \int_h^{\infty} \frac{\xi\omega(\xi) + \xi\omega'(\xi)(s-\xi)}{s^4} ds dh \\
 & \lesssim \omega(\xi) \int_{M\xi}^{\infty} \frac{\omega(h)}{h^2} dh + \omega(\xi) \int_{M\xi}^{\infty} \frac{\xi\omega(h)}{h^3} dh + \omega'(\xi)\xi \int_{M\xi}^{\infty} \frac{\omega(h)}{h^2} dh \\
 & \lesssim \omega(\xi) \int_{M\xi}^{\infty} \frac{\omega(h)}{h^2} dh + \omega'(\xi)\xi \int_{\xi}^{\infty} \frac{\omega(h)}{h^2} dh, \tag{6-3}
 \end{aligned}$$

completing the proof. □

For the other two terms in (6-1), we bound them in two stages.

Lemma 6.2. *Under the assumptions of Lemma 4.1,*

$$\begin{aligned}
 & \int_{-M\xi}^{M\xi} (\delta_h f_x(\xi/2) - h\omega'(\xi))K(\xi/2, h) - (\delta_h f_x(-\xi/2) - h\omega'(\xi))K(-\xi/2, h) dh \\
 & \quad + \int_{|h| > M\xi} [\delta_h f_x(\xi/2) - \delta_h f_x(-\xi/2)]K(\xi/2, h) dh \\
 & \leq \lambda \int_{\mathbb{R}} \frac{\delta_h f_x(\xi/2) - \delta_h f_x(-\xi/2)}{h^2} dh + 2(\Lambda - \lambda) \int_{\xi}^{M\xi} \frac{(\omega(h - \xi) - \omega(\xi))_+}{h^2} dh \\
 & \quad + \omega'(\xi) \int_{\xi < |h| < M\xi} |h[K(\xi/2, h) - K(-\xi/2, h)]| dh. \quad (6-4)
 \end{aligned}$$

Proof. We can bound the second term of (6-4) rather easily. Since

$$\delta_h f_x(\xi/2) - \delta_h f_x(-\xi/2) = (f_x(h + \xi/2) - f_x(h - \xi/2)) - \omega(\xi) \leq 0, \quad (6-5)$$

by the uniform ellipticity of K ,

$$\int_{|h| > M\xi} [\delta_h f_x(\xi/2) - \delta_h f_x(-\xi/2)]K(\xi/2, h) dh \leq \lambda \int_{|h| > M\xi} \frac{\delta_h f_x(\xi/2) - \delta_h f_x(-\xi/2)}{h^2} dh. \quad (6-6)$$

To bound the first term, we first define

$$G(\xi, h) = (\delta_h f_x(\xi/2) - h\omega'(\xi))K(\xi/2, h) - (\delta_h f_x(-\xi/2) - h\omega'(\xi))K(-\xi/2, h). \quad (6-7)$$

Note that since ω is concave and touches f_x from above, see (4-3), it follows that

$$\begin{aligned}
 \delta_h f_x(\xi/2) - \omega'(\xi)h & \leq \delta_h \omega(\xi) - \omega'(\xi)h \leq 0, & h \geq -\xi, \\
 \delta_h f_x(-\xi/2) - \omega'(\xi)h & \geq -\delta_{-h}\omega(\xi) - h\omega'(\xi) \geq 0, & h \leq \xi.
 \end{aligned} \quad (6-8)$$

Thus for $|h| \leq \xi$, by the uniform ellipticity of K we have the bound

$$G(\xi, h) \leq \lambda \frac{\delta_h f_x(\xi/2) - \delta_h f_x(-\xi/2)}{h^2}. \quad (6-9)$$

That just leaves us with the case $\xi \leq |h| \leq M\xi$ to analyze. Note that we can write G in two distinct ways:

$$\begin{aligned}
 G(\xi, h) & = (\delta_h f_x(\xi/2) - \delta_h f_x(-\xi/2))K(\xi/2, h) + (\delta_h f_x(-\xi/2) - h\omega'(\xi))(K(\xi/2, h) - K(-\xi/2, h)) \\
 & = (\delta_h f_x(\xi/2) - \delta_h f_x(-\xi/2))K(-\xi/2, h) + (\delta_h f_x(\xi/2) - h\omega'(\xi))(K(\xi/2, h) - K(-\xi/2, h)). \quad (6-10)
 \end{aligned}$$

By (6-8), $\delta_h f_x(\xi/2) - h\omega'(\xi) \leq 0$ for all $h > \xi$. Thus if $K(\xi/2, h) - K(-\xi/2, h) \geq 0$, then

$$G(\xi, h) \leq \lambda \frac{\delta_h f_x(\xi/2) - \delta_h f_x(-\xi/2)}{h^2} \quad \text{if } K(\xi/2, h) - K(-\xi/2, h) \geq 0. \quad (6-11)$$

On the other hand, since

$$\delta_h f_x(-\xi/2) = \delta_{h-\xi} f(\xi/2) + \omega(\xi) \geq -\omega(h-\xi) + \omega(\xi) \quad (6-12)$$

for $h \geq \xi$, we see that

$$\begin{aligned} G(\xi, h) &\leq \lambda \frac{\delta_h f_x(\xi/2) - \delta_h f_x(-\xi/2)}{h^2} + (\delta_h f_x(-\xi/2) - h\omega'(\xi))(K(\xi/2, h) - K(-\xi/2, h)) \\ &\leq \lambda \frac{\delta_h f_x(\xi/2) - \delta_h f_x(-\xi/2)}{h^2} + (\Lambda - \lambda) \frac{(\omega(h-\xi) - \omega(\xi))_+}{h^2} \\ &\quad + h\omega'(\xi) |K(\xi/2, h) - K(-\xi/2, h)| \quad \text{if } K(\xi/2, h) - K(-\xi/2, h) \leq 0. \end{aligned} \quad (6-13)$$

Putting these two together, we get

$$\begin{aligned} G(\xi, h) &\leq \lambda \frac{\delta_h f_x(\xi/2) - \delta_h f_x(-\xi/2)}{h^2} + (\Lambda - \lambda) \frac{(\omega(h-\xi) - \omega(\xi))_+}{h^2} \\ &\quad + h\omega'(\xi) |K(\xi/2, h) - K(-\xi/2, h)| \end{aligned} \quad (6-14)$$

for $h \geq \xi$. A similar argument can be made in the case that $h \leq -\xi$.

Putting this all together,

$$\begin{aligned} &\int_{-M\xi}^{M\xi} G(\xi, h) dh + \int_{|h| > M\xi} [\delta_h f_x(\xi/2) - \delta_h f_x(-\xi/2)] K(\xi/2, h) dh \\ &\leq \lambda \int_{\mathbb{R}} \frac{\delta_h f_x(\xi/2) - \delta_h f_x(-\xi/2)}{h^2} dh + 2(\Lambda - \lambda) \int_{\xi}^{M\xi} \frac{(\omega(h-\xi) - \omega(\xi))_+}{h^2} dh \\ &\quad + \omega'(\xi) \int_{\xi < |h| < M\xi} |h[K(\xi/2, h) - K(-\xi/2, h)]| dh, \end{aligned} \quad (6-15)$$

completing the proof. $\square$

It's clear that we can bound $\int_{\xi < |h| < M\xi} |h[K(\xi/2, h) - K(-\xi/2, h)]| dh$ as in (5-15). Thus the only thing remaining to prove (4-2) is:

Lemma 6.3. *Under the assumptions of Lemma 4.1,*

$$\begin{aligned} &\lambda \int_{\mathbb{R}} \frac{\delta_h f_x(\xi/2) - \delta_h f_x(-\xi/2)}{h^2} dh \\ &\leq 2\lambda \int_0^{\xi} \frac{\delta_h \omega(\xi) + \delta_{-h} \omega(\xi)}{h^2} dh + 2\lambda \int_{\xi}^{\infty} \frac{\omega(\xi+h) - \omega(h) - \omega(\xi)}{h^2} dh. \end{aligned} \quad (6-16)$$

Proof. To see this, note that formally we should have

$$\int_{\mathbb{R}} \frac{\delta_h f_x(\xi/2) - \delta_h f_x(-\xi/2)}{h^2} dh = \int_{\mathbb{R}} f_x(y) \left(\frac{1}{(y-\xi/2)^2} - \frac{1}{(y+\xi/2)^2} \right) - \frac{\omega(\xi)}{y^2} dy. \quad (6-17)$$

Thus in order to get an upper bound on (6-17), we should be taking an upper bound on $f_x(y)$ when $y > 0$ and a lower bound when $y < 0$. Note by (4-3) that

$$\begin{aligned} f_x(y) &\leq f_x(\xi/2) + \omega(y + \xi/2) - \omega(\xi) = f_x(-\xi/2) + \omega(y + \xi/2), & y > -\xi/2, \\ f_x(y) &\geq f_x(-\xi/2) - \omega(-y + \xi/2) + \omega(\xi) = f_x(\xi/2) - \omega(-y + \xi/2), & y < \xi/2. \end{aligned} \quad (6-18)$$

In particular, using the upper bounds on $\delta_h f_x(\pm\xi/2)$ for $h > 0$ and the lower bounds for $\delta_h f_x(\pm\xi/2)$ for $h < 0$ gives the result. To rigorously justify this though, we will bound

$$\int_{\epsilon}^{\infty} \frac{\delta_h f_x(\xi/2) - \delta_h f_x(-\xi/2)}{h^2} dh$$

from above. Taking $\epsilon \rightarrow 0$, we'll get

$$\int_0^{\infty} \frac{\delta_h f_x(\xi/2) - \delta_h f_x(-\xi/2)}{h^2} dh \leq \int_0^{\xi} \frac{\delta_h \omega(\xi) + \delta_{-h} \omega(\xi)}{h^2} dh + \int_{\xi}^{\infty} \frac{\omega(\xi + h) - \omega(h) - \omega(\xi)}{h^2} dh. \quad (6-19)$$

The bound for $\int_{-\infty}^0$ follows from identical arguments.

So, fix some $\epsilon \ll \xi$. By splitting the integral into a several pieces and reparametrizing, we get

$$\begin{aligned} &\int_{\epsilon}^{\infty} \frac{\delta_h f_x(\xi/2) - \delta_h f_x(-\xi/2)}{h^2} dh \\ &= \int_{\epsilon+\xi/2}^{\infty} \frac{f_x(y)}{(y - \xi/2)^2} dy - \int_{\epsilon-\xi/2}^{\infty} \frac{f_x(y)}{(y + \xi/2)^2} dy - \int_{\epsilon}^{\infty} \frac{\omega(\xi)}{y^2} dy \\ &= \int_{\epsilon+\xi/2}^{\infty} f_x(y) \left(\frac{1}{(y - \xi/2)^2} - \frac{1}{(y + \xi/2)^2} \right) dy - \int_{\epsilon}^{\infty} \frac{\omega(\xi)}{y^2} dy - \int_{\epsilon-\xi/2}^{\epsilon+\xi/2} \frac{f_x(y)}{(y + \xi/2)^2} dy. \end{aligned} \quad (6-20)$$

In the first integral of the third line, since $y > \xi/2$ we have $(y - \xi/2)^{-2} > (y + \xi/2)^{-2}$. So applying the upper bound in (6-18) gives an upper bound on the integral,

$$\begin{aligned} &\int_{\epsilon+\xi/2}^{\infty} f_x(y) \left(\frac{1}{(y - \xi/2)^2} - \frac{1}{(y + \xi/2)^2} \right) dy \\ &\leq \int_{\epsilon+\xi/2}^{\infty} (f_x(\xi/2) + \omega(y + \xi/2) - \omega(\xi)) \left(\frac{1}{(y - \xi/2)^2} - \frac{1}{(y + \xi/2)^2} \right) dy \\ &= \int_{\epsilon+\xi/2}^{\infty} \frac{f_x(\xi/2) + \omega(y + \xi/2) - \omega(\xi)}{(y - \xi/2)^2} dy - \int_{\epsilon+\xi/2}^{\infty} \frac{f_x(\xi/2) + \omega(y + \xi/2) - \omega(\xi)}{(y + \xi/2)^2} dy. \end{aligned} \quad (6-21)$$

By reparametrizing back, we get

$$\begin{aligned} \int_{\epsilon+3\xi/2}^{\infty} \frac{f_x(\xi/2) + \omega(y+\xi/2) - \omega(\xi)}{(y-\xi/2)^2} dy - \int_{\epsilon+\xi/2}^{\infty} \frac{f_x(\xi/2) + \omega(y+\xi/2) - \omega(\xi)}{(y+\xi/2)^2} dy - \int_{\epsilon+\xi}^{\infty} \frac{\omega(\xi)}{y^2} dy \\ = \int_{\epsilon+\xi}^{\infty} \frac{\omega(\xi+h) - \omega(h) - \omega(\xi)}{h^2} dh. \end{aligned} \quad (6-22)$$

Hence combining (6-20), (6-21), and (6-22) gives us

$$\begin{aligned} \int_{\epsilon}^{\infty} \frac{\delta_h f_x(\xi/2) - \delta_h f_x(-\xi/2)}{h^2} dh \\ \leq \int_{\epsilon+\xi}^{\infty} \frac{\omega(\xi+h) - \omega(h) - \omega(\xi)}{h^2} dh + \int_{\epsilon}^{\epsilon+\xi} \frac{f_x(\xi/2) + \omega(\xi+h) - \omega(\xi)}{h^2} dh - \int_{\epsilon}^{\epsilon+\xi} \frac{\omega(\xi)}{h^2} dh - \int_{\epsilon}^{\epsilon+\xi} \frac{f_x(h-\xi/2)}{h^2} dh \\ = \int_{\epsilon+\xi}^{\infty} \frac{\omega(\xi+h) - \omega(h) - \omega(\xi)}{h^2} dh + \int_{\epsilon}^{\epsilon+\xi} \frac{\delta_h \omega(\xi) + f_x(\xi/2) - f_x(h-\xi/2) - \omega(\xi)}{h^2} dh. \end{aligned} \quad (6-23)$$

Now for $h < \xi$, we have $f_x(\xi/2) - f_x(h-\xi/2) \leq \omega(\xi-h)$, and thus

$$\int_{\epsilon}^{\xi} \frac{\delta_h \omega(\xi) + f_x(\xi/2) - f_x(h-\xi/2) - \omega(\xi)}{h^2} dh \leq \int_{\epsilon}^{\xi} \frac{\delta_h \omega(\xi) + \delta_{-h} \omega(\xi)}{h^2} dh. \quad (6-24)$$

Taking the limit as $\epsilon \rightarrow 0$, we then get

$$\int_0^{\infty} \frac{\delta_h f_x(\xi/2) - \delta_h f_x(-\xi/2)}{h^2} dh \leq \int_0^{\xi} \frac{\delta_h \omega(\xi) + \delta_{-h} \omega(\xi)}{h^2} dh + \int_{\xi}^{\infty} \frac{\omega(\xi+h) - \omega(h) - \omega(\xi)}{h^2} dh. \quad \square$$

7. Modulus inequality

Combining all the estimates from the previous two sections, we get a proof of [Lemma 4.1](#). Thus under the assumptions (4-1), we have

$$\begin{aligned} \frac{d}{dt} (f_x(\xi/2) - f_x(-\xi/2)) &\leq A\omega'(\xi) \left(\int_0^{\xi} \frac{\omega(h)}{h} dh + \xi \int_{\xi}^{\infty} \frac{\omega(h)}{h^2} dh + \ln(M+1)\omega(\xi) \right) \\ &\quad + A\omega(\xi) \int_{M\xi}^{\infty} \frac{\omega(h)}{h^2} dh + 2(\Lambda - \lambda) \int_{\xi}^{M\xi} \frac{(\omega(h-\xi) - \omega(\xi))_+}{h^2} dh \\ &\quad + 2\lambda \int_0^{\xi} \frac{\delta_h \omega(\xi) + \delta_{-h} \omega(\xi)}{h^2} dh + 2\lambda \int_{\xi}^{\infty} \frac{\omega(h+\xi) - \omega(h) - \omega(\xi)}{h^2} dh \end{aligned} \quad (7-1)$$

for any $M \geq 1$, where A is a constant depending only on $\|f'_0\|_{L^\infty}$.

In [Kiselev et al. 2007], the authors showed that the modulus

$$\begin{aligned}\omega(\xi) &= \xi - \xi^{3/2}, & 0 \leq \xi \leq \delta, \\ \omega'(\xi) &= \frac{\gamma}{\xi(4 + \log(\xi/\delta))}, & \xi \geq \delta\end{aligned}\quad (7-2)$$

satisfies

$$\begin{aligned}A\omega'(\xi) \left(\int_0^\xi \frac{\omega(h)}{h} dh + \xi \int_\xi^\infty \frac{\omega(h)}{h^2} dh \right) + \lambda \int_0^\xi \frac{\delta_h \omega(\xi) + \delta_{-h} \omega(\xi)}{h^2} dh \\ + \lambda \int_\xi^\infty \frac{\omega(h + \xi) - \omega(h) - \omega(\xi)}{h^2} dh < 0\end{aligned}\quad (7-3)$$

for all $\xi \in \mathbb{R}$ so long as δ, γ are sufficiently small.

With that in mind, we will show:

Lemma 7.1. *Under the assumptions of Lemma 4.1 for the modulus ω defined in (7-2),*

$$\frac{d}{dt}(f_x(\xi/2) - f_x(-\xi/2)) < -\omega'(\xi)\omega(\xi), \quad (7-4)$$

as long as δ, γ are taken sufficiently small depending on $\beta(f'_0), \|f'_0\|_{L^\infty}$.

Proof. By Lemma 4.1 and (7-3) which was proven in [Kiselev et al. 2007], it suffices to show

$$\begin{aligned}A\omega'(\xi) \ln(M+1)\omega(\xi) + A\omega(\xi) \int_{M\xi}^\infty \frac{\omega(h)}{h^2} dh \\ + 2(\Lambda - \lambda) \int_\xi^{M\xi} \frac{(\omega(h - \xi) - \omega(\xi))_+}{h^2} dh + \lambda \int_0^\xi \frac{\delta_h \omega(\xi) + \delta_{-h} \omega(\xi)}{h^2} dh \\ + \lambda \int_\xi^\infty \frac{\omega(h + \xi) - \omega(h) - \omega(\xi)}{h^2} dh \leq -\omega'(\xi)\omega(\xi)\end{aligned}\quad (7-5)$$

for the correct choices of M , and δ, γ sufficiently small.

We proceed very similarly to [Kiselev et al. 2007]. To begin, for $\xi \leq \delta$ we take $M = 1$. Then we just need to show that

$$\begin{aligned}A\omega'(\xi)\omega(\xi) + A\omega(\xi) \int_\xi^\infty \frac{\omega(h)}{h^2} dh + \lambda \int_0^\xi \frac{\delta_h \omega(\xi) + \delta_{-h} \omega(\xi)}{h^2} dh \\ + \lambda \int_\xi^\infty \frac{\omega(h + \xi) - \omega(h) - \omega(\xi)}{h^2} dh \leq -\omega'(\xi)\omega(\xi).\end{aligned}\quad (7-6)$$

In this regime, note that we have the bounds

$$\begin{aligned} \int_{\xi}^{\delta} \frac{\omega(h)}{h^2} dh &\leq \log(\delta/\xi), \\ \int_{\delta}^{\infty} \frac{\omega(h)}{h^2} dh &= \frac{\omega(\delta)}{\delta} + \gamma \int_{\delta}^{\infty} \frac{1}{h^2(4 + \log(h/\delta))} dh \leq 1 + \frac{\gamma}{4\delta} \leq 2 \quad \text{if we take } \gamma < 4\delta, \\ \omega'(\xi) &\leq 1, \quad \omega(\xi) \leq \xi, \\ \int_0^{\xi} \frac{\omega(\xi+h) + \omega(\xi-h) - 2\omega(\xi)}{h^2} dh &\leq \xi \omega''(\xi) = -\frac{3}{2} \xi \xi^{-1/2}. \end{aligned} \quad (7-7)$$

Putting this all together, we get

$$\begin{aligned} (A+1)\omega'(\xi)\omega(\xi) + A\omega(\xi) \int_{\xi}^{\infty} \frac{\omega(h)}{h^2} dh \\ + \lambda \int_0^{\xi} \frac{\omega(\xi+h) + \omega(\xi-h) - 2\omega(\xi)}{h^2} dh + \lambda \int_{\xi}^{\infty} \frac{\omega(\xi+h) - \omega(h) - \omega(\xi)}{h^2} dh \\ \leq \xi \left((A+1)(3 + \log(\delta/\xi)) - \frac{3}{2} \lambda \xi^{-1/2} \right) < 0, \end{aligned} \quad (7-8)$$

assuming that δ is sufficiently small.

Now assume that $\xi \geq \delta$. Then what we need to show is

$$\begin{aligned} A\omega'(\xi) \ln(M+1)\omega(\xi) + A\omega(\xi) \int_{M\xi}^{\infty} \frac{\omega(h)}{h^2} dh + 2(\Lambda - \lambda) \int_{\xi}^{M\xi} \frac{(\omega(h-\xi) - \omega(\xi))_+}{h^2} dh \\ + \lambda \int_0^{\xi} \frac{\delta_h \omega(\xi) + \delta_{-h} \omega(\xi)}{h^2} dh + \lambda \int_{\xi}^{\infty} \frac{\omega(h+\xi) - \omega(h) - \omega(\xi)}{h^2} dh \leq -\omega'(\xi)\omega(\xi). \end{aligned} \quad (7-9)$$

We first bound our new error terms. Using the definition of ω and integrating by parts, we see that

$$\begin{aligned} 2(\Lambda - \lambda) \int_{2\xi}^{M\xi} \frac{\omega(h-\xi) - \omega(\xi)}{h^2} dh &\leq 2(\Lambda - \lambda) \int_{\xi}^{\infty} \frac{\omega(h) - \omega(\xi)}{h^2} dh \leq 2(\Lambda - \lambda) \int_{\xi}^{\infty} \frac{\gamma}{h^2(4 + \log(h/\delta))} dh \\ &\leq \frac{2(\Lambda - \lambda)\gamma}{\xi} \leq \frac{\lambda}{4} \frac{\omega(\delta)}{\xi} \leq \frac{\lambda}{4} \frac{\omega(\xi)}{\xi}, \end{aligned} \quad (7-10)$$

assuming $\gamma \leq (\lambda/(8(\Lambda - \lambda)))\omega(\delta)$.

In order to bound our other new error term, we will be taking M sufficiently large and then γ sufficiently small depending on M, δ . Noting that $\omega(\xi) \leq 2\|f'_0\|_{L^\infty}$, we can bound our other new error term by

integrating by parts,

$$\begin{aligned}
 A\omega(\xi) \int_{M\xi}^{\infty} \frac{\omega(h)}{h^2} dh &\leq \frac{2A\|f'_0\|_{L^\infty}}{M} \frac{\omega(M\xi)}{\xi} + 2A\|f'_0\|_{L^\infty} \int_{M\xi}^{\infty} \frac{\gamma}{h^2(4+\log(h/\delta))} dh \\
 &\leq \frac{2A\|f'_0\|_{L^\infty}}{M} \frac{\omega(M\xi)}{\xi} + \frac{2A\|f'_0\|_{L^\infty}}{M} \frac{\gamma}{\xi} \\
 &\leq \frac{\lambda}{16} \frac{\omega(M\xi)}{\xi} + \frac{\lambda}{8} \frac{\omega(\xi)}{\xi},
 \end{aligned} \tag{7-11}$$

assuming that

$$M \geq \frac{32A\|f'_0\|_{L^\infty}}{\lambda},$$

and then γ is sufficiently small so that

$$\frac{2\|f'_0\|_{L^\infty}A}{M}\gamma \leq \frac{\lambda}{8}\omega(\delta) \leq \frac{\lambda}{8}\omega(\xi).$$

Note that this is where we set a value for M , and that γ is taken sufficiently small depending on M . Now that the value for M is fixed, we can also control the value $\omega(M\xi)$ by taking γ sufficiently small that

$$\begin{aligned}
 \omega(M\xi) &= \omega(\xi) + \int_{\xi}^{M\xi} \frac{\gamma}{h(4+\log(h/\delta))} dh \leq \omega(\xi) + \gamma \ln(M) \leq \omega(\xi) + \omega(\delta) \\
 &\leq 2\omega(\xi).
 \end{aligned} \tag{7-12}$$

Hence,

$$A\omega(\xi) \int_{M\xi}^{\infty} \frac{\omega(h)}{h^2} dh \leq \frac{\lambda}{16} \frac{\omega(M\xi)}{\xi} + \frac{\lambda}{8} \frac{\omega(\xi)}{\xi} \leq \frac{\lambda}{4} \frac{\omega(\xi)}{\xi}. \tag{7-13}$$

Using the same integration-by-parts tricks, we can also show

$$\lambda \int_{\xi}^{\infty} \frac{\omega(h+\xi) - \omega(h) - \omega(\xi)}{h^2} dh \leq -\frac{3}{4}\lambda \frac{\omega(\xi)}{\xi} \tag{7-14}$$

for γ sufficiently small.

So combining these, we get

$$\begin{aligned}
 A\omega(\xi) \int_{M\xi}^{\infty} \frac{\omega(h)}{h^2} dh + 2(\Lambda - \lambda) \int_{2\xi}^{M\xi} \frac{\omega(h-\xi) - \omega(\xi)}{h^2} dh \\
 + \lambda \int_{\xi}^{\infty} \frac{\omega(h+\xi) - \omega(h) - \omega(\xi)}{h^2} dh \leq -\frac{\lambda}{4} \frac{\omega(\xi)}{\xi}.
 \end{aligned} \tag{7-15}$$

Since $\omega'(\xi)\omega(\xi) \leq \gamma\omega(\xi)/\xi$, we finally get

$$(A \ln(M+1) + 1)\omega'(\xi)\omega(\xi) - \frac{\lambda}{4} \frac{\omega(\xi)}{\xi} \leq \frac{\omega(\xi)}{\xi} \left((A \ln(M+1) + 1)\gamma - \frac{\lambda}{4} \right) < 0, \quad (7-16)$$

if γ is taken sufficiently small. $\square$

8. Our choice for the modulus ρ

We've now shown that for the modulus defined in (7-2) if the assumptions (4-1) hold then

$$\frac{d}{dt}(f_x(t, \xi/2) - f_x(t, -\xi/2))\big|_{t=T} < -\omega'(\xi)\omega(\xi). \quad (8-1)$$

We claim that in fact (8-1) will hold for any rescaling $\omega_r(h) = \omega(rh)$ as well. To see this, fix some $r > 0$, and suppose that $f(t, x)$ satisfies the conditions of Lemma 4.1 for ω_r at time T and distance ξ . Take $\tilde{f}(t, x) = rf(t/r, x/r)$, which is also a solution of (1-2). Then $\tilde{f}_x$ is a solution of (3-5) with $\beta(\tilde{f}_0') = \beta(f_0')$, $\|\tilde{f}_0'\|_{L^\infty} = \|f_0'\|_{L^\infty}$, and satisfying the conditions of Lemma 4.1 for ω at time rT and distance $r\xi$. Hence by Lemma 7.1

$$\begin{aligned} \frac{d}{dt}(f_x(t, \xi/2) - f_x(t, -\xi/2))\big|_{t=T} &= r \frac{d}{dt}(\tilde{f}_x(t, r\xi/2) - \tilde{f}_x(t, -r\xi/2))\big|_{t=rT} \\ &< -r\omega'(r\xi)\omega(r\xi) = -\omega'_r(\xi)\omega_r(\xi). \end{aligned} \quad (8-2)$$

So, (8-1) will hold for any rescaling ω_r . Also note that for $f_x(T, \xi/2) - f_x(T, -\xi/2) = \omega(\xi)$ to hold, we must necessarily have $\omega(\xi) \leq 2\|f_x(T, \cdot)\|_{L^\infty} < 2\|f_0'\|_{L^\infty}$. Thus taking

$$C = \sup_{0 < h < \omega^{-1}(2\|f_0'\|_{L^\infty})} \frac{h}{\omega(h)} = \frac{\omega^{-1}(2\|f_0'\|_{L^\infty})}{2\|f_0'\|_{L^\infty}}, \quad (8-3)$$

we see that

$$\omega(h) \geq \frac{h}{C} \quad (8-4)$$

for all relevant h . Define

$$\rho(h) := \omega(Ch) \quad (8-5)$$

so that

$$\rho(h) \geq h \quad (8-6)$$

for all $h \in [0, \rho^{-1}(2\|f_0'\|_{L^\infty})]$.

Now, suppose that at time T , f satisfies the assumptions (4-1) for $\rho(\cdot/T)$. Then since $\rho(\cdot/T)$ is a rescaling of ω , we have

$$\begin{aligned} \frac{d}{dt}(f_x(T, \xi/2) - f_x(T, -\xi/2)) &< -\frac{d}{dh}\rho(h/T)\big|_{h=\xi}\rho(\xi/T) \\ &= -\frac{1}{T}\rho'(\xi/T)\rho(\xi/T) \leq -\frac{\xi}{T^2}\rho'(\xi/T) = \frac{d}{dt}\rho(\xi/t)\big|_{t=T}. \end{aligned} \quad (8-7)$$

Thus we've constructed a modulus ρ which satisfies (2-12), completing the proof of the generation of a Lipschitz modulus of continuity (1-8) in our main theorem.

9. Regularity in time

With the construction of the modulus ρ , we get universal Lipschitz bounds in space for $f_x(t, \cdot)$. By the structure of (1-2), we also get regularity in space for f_t .

Proposition 9.1. *Let $f : (0, T) \times \mathbb{R} \rightarrow \mathbb{R}$ be a classical solution to (1-2) with $\|f(t, \cdot)\|_{W^{1,\infty}}$ bounded and $\|f_{xx}(t, \cdot)\|_{L^\infty} \lesssim 1/t$. Then $f_t(t, \cdot)$ is log-Lipschitz in space with*

$$\begin{aligned} |f_t(t, \cdot)| &\lesssim \max\{-\log(t), 1\}, \\ |f_t(t, x) - f_t(t, y)| &\lesssim -\log(|x - y|)|x - y| \left(1 + \frac{1}{t}\right), \quad 0 < |x - y| < \frac{1}{2}. \end{aligned} \quad (9-1)$$

Proof. For $t < 1$, we have

$$\begin{aligned} |f_t(t, x)| &= \left| \int_{\mathbb{R}} \frac{\delta_h f(t, x) - h f_x(t, x)}{\delta_h f(t, x)^2 + h^2} dh \right| \\ &\leq \left| \int_0^\infty \frac{\delta_h f(t, x) + \delta_{-h} f(t, x)}{\delta_{-h} f(t, x)^2 + h^2} dh \right| + \left| \int_0^\infty \frac{(\delta_h f(t, x) - h f_x(t, x))(\delta_h f(t, x)^2 - \delta_{-h} f(t, x)^2)}{(\delta_h f(t, x)^2 + h^2)(\delta_{-h} f(t, x)^2 + h^2)} dh \right| \\ &\lesssim \int_0^t \frac{1}{t} dh + \int_t^1 \frac{1}{h} dh + \int_1^\infty \frac{1}{h^2} + \frac{1}{h^3} dh \\ &\lesssim -\log(t) + 1. \end{aligned} \quad (9-2)$$

For $t > 1$, we can similarly show $|f_t(t, x)| \lesssim 1$, proving the first bound.

For regularity in space, we see that

$$\begin{aligned} &f_t(t, x) - f_t(t, y) \\ &= \int_{\mathbb{R}} \frac{\delta_h f(t, x) - h f_x(t, x)}{\delta_h f(t, x)^2 + h^2} - \frac{\delta_h f(t, y) - h f_x(t, y)}{\delta_h f(t, y)^2 + h^2} dh \\ &= \int_{\mathbb{R}} \frac{\delta_h f(t, x) - h f_x(t, x) - (\delta_h f(t, y) - h f_x(t, y))}{\delta_h f(t, y)^2 + h^2} + \frac{(\delta_h f(t, x) - h f_x(t, x))(\delta_h f(t, x)^2 - \delta_h f(t, y)^2)}{(\delta_h f(t, x)^2 + h^2)(\delta_h f(t, y)^2 + h^2)} dh \\ &\leq \left| \int_{|h| < |x-y|} \right| + \left| \int_{|x-y| < |h| < 1} \right| + \left| \int_{|h| > 1} \right|. \end{aligned} \quad (9-3)$$

For $|h| < |x - y|$, we can bound much as before to get

$$\left| \int_{|h| < |x-y|} \right| \lesssim \int_0^{|x-y|} \frac{1}{t} dh = \frac{|x-y|}{t}. \quad (9-4)$$

For midsize $|x - y| < |h| < 1$, we have

$$\begin{aligned} |\delta_h f(t, x) - hf_x(t, x) - (\delta_h f(t, y) - hf_x(t, y))| &= \left| \int_0^h \delta_s f_x(t, x) - \delta_s f_x(t, y) ds \right| \lesssim \frac{|x - y|h}{t}, \\ |\delta_h f(t, x) - \delta_h f(t, y)| &= \left| \int_0^h f_x(t, x + s) - f_x(t, y + s) ds \right| \lesssim \frac{|x - y|h}{t}. \end{aligned} \quad (9-5)$$

Thus

$$\left| \int_{|x-y| < |h| < 1} \right| \lesssim \frac{|x - y|}{t} \int_{|x-y|}^1 \frac{1}{h} dh = \frac{-\ln(|x - y|)|x - y|}{t}. \quad (9-6)$$

Finally, we use L^∞ bounds on f to get

$$\begin{aligned} \left| \int_{|h| > 1} \right| &\leq \left| \int_{|h| > 1} \frac{\delta_h f(t, x) - \delta_h f(t, y)}{\delta_h f(t, y)^2 + h^2} + \frac{(\delta_h f(t, x) - hf_x(t, x))(\delta_h f(t, x)^2 - \delta_h f(t, y)^2)}{(\delta_h f(t, x)^2 + h^2)(\delta_h f(t, y)^2 + h^2)} dh \right| \\ &\quad + |f_x(t, x) - f_x(t, y)| \left| \int_{|h| > 1} \frac{-h}{\delta_h f(t, y)^2 + h^2} dh \right| \\ &\lesssim |x - y| \int_1^\infty \frac{1}{h^2} + \frac{1}{h^3} dh + \frac{|x - y|}{t} \int_1^\infty \frac{1}{h^3} dh \lesssim \left(1 + \frac{1}{t}\right) |x - y|. \end{aligned} \quad (9-7)$$

Putting this all together, we thus have

$$|f_t(t, x) - f_t(t, y)| \lesssim -\ln(|x - y|)|x - y| \left(1 + \frac{1}{t}\right), \quad (9-8)$$

completing the proof. $\square$

Recall that in [Section 2](#), we assumed that our initial data f_0 was in $C_c^\infty(\mathbb{R})$ so that, by the local existence results of [\[Córdoba and Gancedo 2009\]](#), there was a unique solution $f \in C^1((0, T_+); H^k)$ for k arbitrarily large and some $T_+ > 0$. We were then able to prove the existence of the modulus ρ as in [Theorem 1.1](#) depending only on $\beta(f'_0)$, $\|f'_0\|_{L^\infty}$, and hence with the solution f existing for all time by the main theorem of [\[Constantin et al. 2017\]](#). For an arbitrary $f_0 \in W^{1,\infty}(\mathbb{R})$ with $\beta(f'_0) < 1$, the same result holds true by compactness. Let $\eta \in C_c^\infty(\mathbb{R})$ be a smooth mollifier and $\phi \in C_c^\infty(\mathbb{R})$ be a smooth cutoff function. For $f_0 \in W^{1,\infty}(\mathbb{R})$ with $\beta(f'_0) < 1$, take $f_0^{(\epsilon)}(x) := (f_0 * \eta_\epsilon)(x)\phi(\epsilon x)$. Then $f_0^{(\epsilon)} \rightarrow f_0$ in $W_{\text{loc}}^{1,\infty}$, with $\beta(f_0^{(\epsilon)'}) \rightarrow \beta(f'_0)$ and $\|f_0^{(\epsilon)}\|_{W^{1,\infty}(\mathbb{R})} \rightarrow \|f_0\|_{W^{1,\infty}(\mathbb{R})}$ as $\epsilon \rightarrow 0$. Thus for ϵ sufficiently small, $\beta(f_0^{(\epsilon)'}) < 1$ and the results of the previous section hold for the solution to the mollified problem $f^{(\epsilon)}$. The L^∞ bound on $f_t^{(\epsilon)}$ proven above along with the maximum principle for $f_x^{(\epsilon)}$ is enough to ensure that there a subsequence $f^{(\epsilon_k)}$ converging in $C_{\text{loc}}([0, \infty) \times \mathbb{R})$ to a Lipschitz (weak) solution f to the original problem. In order to get a classical C^1 solution, we need regularity estimates for $f_x^{(\epsilon)}$, $f_t^{(\epsilon)}$ in both time and space. The modulus ρ and [Proposition 9.1](#) give the regularity in space that we need for f_x , f_t . All that leaves is to prove regularity in time.

Proposition 9.2. *Let f be a sufficiently smooth solution to (1-2) with $\beta(f'_0) < 1$. Then $f_x, f_t \in C_{\text{loc}}^\alpha((0, \infty) \times \mathbb{R})$ with*

$$\|f_x\|_{C^\alpha(Q_{t/4}(t,x))}, \|f_t\|_{C^\alpha(Q_{t/4}(t,x))} \leq C(\beta(f'_0), \|f\|_{L_t^\infty((t/2, 3t/2); W_x^{2,\infty}(\mathbb{R}))}) \max\{t^{-\alpha}, 1\}, \quad (9-9)$$

where $Q_r(s, y) = (s - r, s] \times B_r(y)$, and $\alpha > 0$ depends only on $\beta(f'_0), \|f'_0\|_{L^\infty}$.

Proof. We have that f_x solves

$$(f_x)_t(t, x) = f_{xx}(t, x) \int_{\mathbb{R}} \frac{-h}{\delta_h f(t, x)^2 + h^2} dh + \int_{\mathbb{R}} \delta_h f_x(t, x) K(t, x, h) dh, \quad (9-10)$$

where $\lambda/h^2 \leq K(t, x, h) \leq \Lambda/h^2$ is uniformly elliptic with ellipticity constants λ, Λ depending on $\beta(f'_0), \|f'_0\|_{L^\infty}$. Rewriting this, we have that f_x satisfies

$$\begin{aligned} (f_x)_t - \int_{\mathbb{R}} \delta_h f_x(t, x) \left(\frac{K(t, x, h) + K(t, x, -h)}{2} \right) dh \\ = f_{xx}(t, x) \int_{\mathbb{R}} \frac{-h}{\delta_h f(t, x)^2 + h^2} dh + \int_{\mathbb{R}} \delta_h f_x(t, x) \left(\frac{K(t, x, h) - K(t, x, -h)}{2} \right) dh. \end{aligned} \quad (9-11)$$

Let $F(t, x)$ denote the right-hand side of (9-11). Then $F(t, x)$ is locally bounded with $|F(t, x)|$ controlled by $\|f(t, \cdot)\|_{W^{2,\infty}}$. Then since $(K(t, x, h) + K(t, x, -h))/2$ is a symmetric uniformly elliptic kernel, it follows that we have local C^α bounds for $\alpha \leq \alpha_0$ for some α_0 depending on ellipticity constants; see [Silvestre 2011].

So, all we have to do is give bounds on $F(t, x)$ depending only on $\|f(t, \cdot)\|_{W^{2,\infty}}$. Similar to the proof of Lemma 5.1,

$$\int_{\mathbb{R}} \frac{-h}{\delta_h f(t, x)^2 + h^2} dh = \int_0^\infty h \frac{\delta_h f(t, x)^2 - \delta_{-h} f(t, x)^2}{(\delta_h f(t, x)^2 + h^2)(\delta_{-h} f(t, x)^2 + h^2)} dh \lesssim \int_0^1 1 dh + \int_1^\infty \frac{1}{h^3} dh \lesssim 1. \quad (9-12)$$

Also similar to the proof of Lemma 5.2, specifically (5-12), we have

$$|K(t, x, h) - K(t, x, -h)| \lesssim \min\left\{\frac{1}{h}, \frac{1}{h^3}\right\}, \quad (9-13)$$

so

$$\left| \int_{\mathbb{R}} \delta_h f_x(t, x) \left(\frac{K(t, x, h) - K(t, x, -h)}{2} \right) dh \right| \lesssim \int_0^1 1 dh + \int_1^\infty \frac{1}{h^3} dh \lesssim 1. \quad (9-14)$$

Thus since we've bounded the right-hand side of (9-11) depending only on $\|f(t, \cdot)\|_{W^{2,\infty}}$, we have our local C^α bounds for f_x for all α sufficiently small. A C^α bound that is uniform in x for f_x then gives a log C^α estimate for f_t , similar to the proof for regularity in space in Proposition 9.1. Thus we have C^α estimates for both f_x, f_t . $\square$

Appendix: Uniqueness

We now prove that if our initial data f_0 is in $C^{1,\epsilon}(\mathbb{R})$ with $\beta(f'_0) < 1$, then the solution f given by [Theorem 1.1](#) is unique with $f \in L^\infty([0, \infty); C^{1,\epsilon})$. As mentioned before, this essentially follows from the uniqueness theorem given in [\[Constantin et al. 2017\]](#), which under our assumptions simplifies to:

Theorem A.1 [\[Constantin et al. 2017\]](#). *Let $f \in L^\infty([0, T]; W^{1,\infty})$ be a classical, C^1 solution to (1-2) with initial data $f(0, x) = f_0(x)$. Assume that $\lim_{x \rightarrow \infty} f(t, x) = 0$, and that there is some modulus of continuity $\tilde{\rho}$ such that*

$$f_x(t, x) - f_x(t, y) \leq \tilde{\rho}(|x - y|) \quad \text{for all } 0 \leq t \leq T, \ x \neq y \in \mathbb{R}. \quad (\text{A-1})$$

Then the solution f is unique.

The authors of [\[Constantin et al. 2017\]](#) note that the uniform continuity assumption should be the only real assumption; the decay is assumed for convenience in their proof. So, we start by proving that if $f_0 \in C^{1,\epsilon}(\mathbb{R})$, then the solution f is in $L^\infty([0, \infty); C^{1,\epsilon})$. To begin, suppose that $f_0 \in C^{1,1}(\mathbb{R})$. Then necessarily f'_0 has modulus $\rho(\cdot/\delta)$ for some $\delta > 0$ sufficiently small. The same proof for the instantaneous generation of the modulus ρ will give that $f_x(t, \cdot)$ has modulus $\rho(\cdot/t + \delta)$. Hence $f_x(t, \cdot)$ has modulus $\rho(\cdot/\delta)$ for all $t \geq 0$.

If $f_0 \in C^{1,\epsilon}(\mathbb{R})$, we can make the same essential argument by changing the definitions of ρ, ω . You can repeat the arguments of [Sections 7 and 8](#) for the modulus

$$\begin{aligned} \omega^{(\epsilon)}(\xi) &= \xi^\epsilon, & 0 \leq \xi \leq \delta, \\ \omega^{(\epsilon)'}(\xi) &= \frac{\gamma}{\xi(4 + \log(\xi/\delta))}, & \xi \geq \delta. \end{aligned} \quad (\text{A-2})$$

All the error terms for $\xi \leq \delta$ are of order $\xi^{2\epsilon-1}$, while the diffusion term is of order $\xi^{\epsilon-1}$, so there are no problems as long as δ is sufficiently small. The argument for $\xi \geq \delta$ is identical to the original. Taking $\rho^{(\epsilon)}$ to be some suitable rescaling of $\omega^{(\epsilon)}$, we then have that if f'_0 has modulus $\rho^{(\epsilon)}(\cdot/\delta)$, then $f_x(t, \cdot)$ will have modulus $\rho^{(\epsilon)}(\cdot/t + \delta)$.

Thus if $f_0 \in C^{1,\epsilon}(\mathbb{R})$, then the solution f given by [Theorem 1.1](#) will satisfy the main uniform continuity assumption of [Theorem A.1](#). Our solution f will not decay as $x \rightarrow \infty$, but that assumption isn't truly necessary.

Let f_1, f_2 be two uniformly continuous, classical solutions to (1-2) with the same initial data, and let $M(t) = \|f_1(t, \cdot) - f_2(t, \cdot)\|_{L^\infty}$. With the decay assumption, the authors of [\[Constantin et al. 2017\]](#) are able to assume that for almost every t , there is a point $x(t) \in \mathbb{R}$ such that

$$M(t) = |f_1(t, x(t)) - f_2(t, x(t))|, \quad \frac{d}{dt}M(t) = \left(\frac{d}{dt}|f_1 - f_2|\right)(t, x(t)). \quad (\text{A-3})$$

They then bound $\frac{d}{dt}|f_1(t, x(t)) - f_2(t, x(t))|$ using equation (1-2), $\tilde{\rho}$, and $W^{1,\infty}$ bounds.

Without the decay assumption, we instead use that

$$\frac{d}{dt}M(t) \leq \sup \left\{ \frac{d}{dt}|f_1(t, x) - f_2(t, x)| : |f_1(t, x) - f_2(t, x)| \geq M(t) - \delta \right\}, \quad (\text{A-4})$$

where $\delta > 0$ is arbitrary. When we go to bound $\frac{d}{dt}|f_1(t, x) - f_2(t, x)|$, we then get new error terms which can be bounded by

$$C(\tilde{\rho}, \max_i \|f_i(t, \cdot)\|_{W^{1,\infty}}, M(t))(\delta + |f_{1,x}(t, x) - f_{2,x}(t, x)|). \quad (\text{A-5})$$

Since $f_{i,x}(t, x)$ is bounded and has modulus $\tilde{\rho}$, it then follows that

$$|f_{1,x}(t, x) - f_{2,x}(t, x)| = o_\delta(1). \quad (\text{A-6})$$

Thus by taking δ sufficiently small depending on $\tilde{\rho}, \max_i \|f_i(t, \cdot)\|_{W^{1,\infty}}, M(t)$, we can guarantee that the new error terms $\lesssim M(t)$. Then the original proof of [Constantin et al. 2017] goes through.

Acknowledgement

I would like to thank my advisor Luis Silvestre for suggesting the problem, pointing me towards good resources, and just giving good advice in general.

References

- [Castro et al. 2012] A. Castro, D. Córdoba, C. Fefferman, F. Gancedo, and M. López-Fernández, “Rayleigh–Taylor breakdown for the Muskat problem with applications to water waves”, *Ann. of Math.* (2) **175**:2 (2012), 909–948. [MR](#) [Zbl](#)
- [Constantin et al. 2013] P. Constantin, D. Córdoba, F. Gancedo, and R. M. Strain, “On the global existence for the Muskat problem”, *J. Eur. Math. Soc. (JEMS)* **15**:1 (2013), 201–227. [MR](#) [Zbl](#)
- [Constantin et al. 2016] P. Constantin, D. Córdoba, F. Gancedo, L. Rodríguez-Piazza, and R. M. Strain, “On the Muskat problem: global in time results in 2D and 3D”, *Amer. J. Math.* **138**:6 (2016), 1455–1494. [MR](#) [Zbl](#)
- [Constantin et al. 2017] P. Constantin, F. Gancedo, R. Shvydkoy, and V. Vicol, “Global regularity for 2D Muskat equations with finite slope”, *Ann. Inst. H. Poincaré Anal. Non Linéaire* **34**:4 (2017), 1041–1074. [MR](#) [Zbl](#)
- [Córdoba and Gancedo 2007] D. Córdoba and F. Gancedo, “Contour dynamics of incompressible 3-D fluids in a porous medium with different densities”, *Comm. Math. Phys.* **273**:2 (2007), 445–471. [MR](#) [Zbl](#)
- [Córdoba and Gancedo 2009] D. Córdoba and F. Gancedo, “A maximum principle for the Muskat problem for fluids with different densities”, *Comm. Math. Phys.* **286**:2 (2009), 681–696. [MR](#) [Zbl](#)
- [Dabkowski et al. 2014] M. Dabkowski, A. Kiselev, L. Silvestre, and V. Vicol, “Global well-posedness of slightly supercritical active scalar equations”, *Anal. PDE* **7**:1 (2014), 43–72. [MR](#) [Zbl](#)
- [Deng et al. 2017] F. Deng, Z. Lei, and F. Lin, “On the two-dimensional Muskat problem with monotone large initial data”, *Comm. Pure Appl. Math.* **70**:6 (2017), 1115–1145. [MR](#) [Zbl](#)
- [Gancedo 2017] F. Gancedo, “A survey for the Muskat problem and a new estimate”, *SeMA J.* **74**:1 (2017), 21–35. [MR](#) [Zbl](#)
- [Kiselev 2010] A. Kiselev, “Regularity and blow up for active scalars”, *Math. Model. Nat. Phenom.* **5**:4 (2010), 225–255. [MR](#) [Zbl](#)
- [Kiselev et al. 2007] A. Kiselev, F. Nazarov, and A. Volberg, “Global well-posedness for the critical 2D dissipative quasi-geostrophic equation”, *Invent. Math.* **167**:3 (2007), 445–453. [MR](#) [Zbl](#)
- [Muskat 1934] M. Muskat, “Two fluid systems in porous media: the encroachment of water into an oil sand”, *J. Appl. Phys.* **5**:9 (1934), 250–264. [Zbl](#)
- [Patel and Strain 2017] N. Patel and R. M. Strain, “Large time decay estimates for the Muskat equation”, *Comm. Partial Differential Equations* **42**:6 (2017), 977–999. [MR](#) [Zbl](#)
- [Saffman and Taylor 1958] P. G. Saffman and G. Taylor, “The penetration of a fluid into a porous medium or Hele-Shaw cell containing a more viscous liquid”, *Proc. Roy. Soc. London. Ser. A* **245** (1958), 312–329. [MR](#) [Zbl](#)
- [Silvestre 2011] L. Silvestre, “On the differentiability of the solution to the Hamilton–Jacobi equation with critical fractional diffusion”, *Adv. Math.* **226**:2 (2011), 2020–2039. [MR](#) [Zbl](#)

Received 9 May 2017. Revised 14 Jan 2018. Accepted 30 Jul 2018.

STEPHEN CAMERON: scameron@math.uchicago.edu

Department of Mathematics, University of Chicago, Chicago, IL, United States

Analysis & PDE

msp.org/apde

EDITORS

EDITOR-IN-CHIEF

Patrick Gérard

patrick.gerard@math.u-psud.fr

Université Paris Sud XI

Orsay, France

BOARD OF EDITORS

Massimiliano Berti	Scuola Intern. Sup. di Studi Avanzati, Italy berti@sissa.it	Clément Mouhot	Cambridge University, UK c.mouhot@dpmms.cam.ac.uk
Sun-Yung Alice Chang	Princeton University, USA chang@math.princeton.edu	Werner Müller	Universität Bonn, Germany mueller@math.uni-bonn.de
Michael Christ	University of California, Berkeley, USA mchrist@math.berkeley.edu	Gilles Pisier	Texas A&M University, and Paris 6 pisier@math.tamu.edu
Alessio Figalli	ETH Zurich, Switzerland alessio.figalli@math.ethz.ch	Tristan Rivière	ETH, Switzerland riviere@math.ethz.ch
Charles Fefferman	Princeton University, USA cf@math.princeton.edu	Igor Rodnianski	Princeton University, USA irod@math.princeton.edu
Ursula Hamenstaedt	Universität Bonn, Germany ursula@math.uni-bonn.de	Sylvia Serfaty	New York University, USA serfaty@cims.nyu.edu
Vaughan Jones	U.C. Berkeley & Vanderbilt University vaughan.f.jones@vanderbilt.edu	Yum-Tong Siu	Harvard University, USA siu@math.harvard.edu
Vadim Kaloshin	University of Maryland, USA vadim.kaloshin@gmail.com	Terence Tao	University of California, Los Angeles, USA tao@math.ucla.edu
Herbert Koch	Universität Bonn, Germany koch@math.uni-bonn.de	Michael E. Taylor	Univ. of North Carolina, Chapel Hill, USA met@math.unc.edu
Izabella Laba	University of British Columbia, Canada ilaba@math.ubc.ca	Gunther Uhlmann	University of Washington, USA gunther@math.washington.edu
Gilles Lebeau	Université de Nice Sophia Antipolis, France lebeau@unice.fr	András Vasy	Stanford University, USA andras@math.stanford.edu
Richard B. Melrose	Massachusetts Inst. of Tech., USA rbbm@math.mit.edu	Dan Virgil Voiculescu	University of California, Berkeley, USA dvv@math.berkeley.edu
Frank Merle	Université de Cergy-Pontoise, France Frank.Merle@u-cergy.fr	Steven Zelditch	Northwestern University, USA zelditch@math.northwestern.edu
William Minicozzi II	Johns Hopkins University, USA minicozz@math.jhu.edu	Maciej Zworski	University of California, Berkeley, USA zworski@math.berkeley.edu

PRODUCTION

production@msp.org

Silvio Levy, Scientific Editor

See inside back cover or msp.org/apde for submission instructions.

The subscription price for 2019 is US \$310/year for the electronic version, and \$520/year (+\$60, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues from the last three years and changes of subscriber address should be sent to MSP.

Analysis & PDE (ISSN 1948-206X electronic, 2157-5045 printed) at Mathematical Sciences Publishers, 798 Evans Hall #3840, c/o University of California, Berkeley, CA 94720-3840, is published continuously online. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices.

APDE peer review and production are managed by EditFlow® from MSP.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2019 Mathematical Sciences Publishers

ANALYSIS & PDE

Volume 12 No. 4 2019

Quantum dynamical bounds for ergodic potentials with underlying dynamics of zero topological entropy	867
RUI HAN and SVETLANA JITOMIRSKAYA	
Two-dimensional gravity water waves with constant vorticity, I: Cubic lifespan	903
MIHAELA IFRIM and DANIEL TATARU	
Absolute continuity and α -numbers on the real line	969
TUOMAS ORPONEN	
Global well-posedness for the two-dimensional Muskat problem with slope less than 1	997
STEPHEN CAMERON	
Global well-posedness and scattering for the radial, defocusing, cubic wave equation with initial data in a critical Besov space	1023
BENJAMIN DODSON	
Nonexistence of Wente's L^∞ estimate for the Neumann problem	1049
JONAS HIRSCH	
Global geometry and C^1 convex extensions of 1-jets	1065
DANIEL AZAGRA and CARLOS MUDARRA	
Classification of positive singular solutions to a nonlinear biharmonic equation with critical exponent	1101
RUPERT L. FRANK and TOBIAS KÖNIG	
Optimal multilinear restriction estimates for a class of hypersurfaces with curvature	1115
IOAN BEJENARU	