

ANALYSIS & PDE

Volume 12

No. 8

2019

JONAS AZZAM AND MIHALIS MOURGOLOU

**TANGENT MEASURES OF ELLIPTIC MEASURE AND
APPLICATIONS**



TANGENT MEASURES OF ELLIPTIC MEASURE AND APPLICATIONS

JONAS AZZAM AND MIHALIS MOURGOLOU

Tangent measure and blow-up methods are powerful tools for understanding the relationship between the infinitesimal structure of the boundary of a domain and the behavior of its harmonic measure. We introduce a method for studying tangent measures of elliptic measures in arbitrary domains associated with (possibly nonsymmetric) elliptic operators in divergence form whose coefficients have vanishing mean oscillation at the boundary. In this setting, we show the following for domains $\omega \subset \mathbb{R}^{n+1}$, $n \geq 2$:

- (1) We extend the results of Kenig, Preiss, and Toro (*J. Amer. Math. Soc.* **22**:3 (2009), 771–796) by showing mutual absolute continuity of interior and exterior elliptic measures for *any* domains implies the tangent measures are a.e. flat and the elliptic measures have dimension n .
- (2) We generalize the work of Kenig and Toro (*J. Reine Agnew. Math.* **596** (2006), 1–44) and show that VMO equivalence of doubling interior and exterior elliptic measures for general domains implies the tangent measures are always supported on the zero sets of elliptic polynomials.
- (3) In a uniform domain that satisfies the capacity density condition and whose boundary is locally finite and has a.e. positive lower n -Hausdorff density, we show that if the elliptic measure is absolutely continuous with respect to n -Hausdorff measure then the boundary is rectifiable. This generalizes the work of Akman, Badger, Hofmann, and Martell (*Trans. Amer. Math. Soc.* **369**:8 (2017), 5711–5745).

1. Introduction	1891
2. Tangent measures	1900
3. Elliptic measures	1905
4. Harmonic polynomial measures	1917
5. Proof of Theorem I	1921
6. BMO, VMO and vanishing A_∞	1924
7. Proofs of Theorems II and III	1929
8. Proof of Theorem IV	1932
9. Proof of Proposition III	1936
Acknowledgements	1938
References	1939

1. Introduction

1A. Background. In this paper, we study how the relationships between the elliptic measures of two complementary domains in \mathbb{R}^{n+1} , for $n \geq 2$, dictate the geometry of their common boundaries. We shall denote those domains by Ω^+ and Ω^- and the respective elliptic measures by ω^+ and ω^- . Bishop,

MSC2010: 31A15, 28A75, 28A78, 28A33.
Keywords: harmonic measure, elliptic measure, capacity density condition, Δ -regular domains, tangent measures, absolute continuity, rectifiability.

Carleson, Garnett and Jones [Bishop et al. 1989] showed that, for disjoint simply connected planar domains with mutually absolutely continuous harmonic measures, the boundary has tangents on a set of positive measure. Kenig, Preiss, and Toro [Kenig et al. 2009] showed that if Ω^\pm are both nontangentially accessible (or NTA) domains in \mathbb{R}^{n+1} , with $n \geq 2$, and the interior and exterior harmonic measures are mutually absolutely continuous, then at every point of the common boundary except for a set of harmonic measure zero, $\partial\Omega^+$ looks flatter and flatter as we zoom in. We will not define NTA but refer the reader to its inception in [Jerison and Kenig 1982]. Recently, the authors of the current paper, along with Tolsa [Azzam et al. 2017b], as well as with Tolsa and Volberg [Azzam et al. 2016c], showed that additionally the boundary is n -rectifiable in the sense that, off a set of harmonic measure zero, the boundary is a union of Lipschitz images of \mathbb{R}^{n+1} , and in fact Ω^+ and Ω^- need not be NTA but just connected.

These are, however, almost everywhere phenomena, so it is interesting to ask what assumptions we need on ω^\pm to guarantee some nice limiting behavior of our blow-ups at *every* point. Kenig and Toro [2006] showed that if Ω^+ is 2-sided NTA and $\log(d\omega^-/d\omega^+) \in \text{VMO}(d\omega^+)$, then as we zoom in on any point of the boundary for a particular sequence of scales, $\partial\Omega^+$ begins to look more and more like the zero set of a harmonic polynomial (see Section 6 for the definition of VMO). In [Badger 2011], it is further shown that these harmonic polynomials are always homogeneous, and [Badger 2013] investigates the topological properties of sets where the boundary is approximated by zero sets of harmonic polynomials in this way.

To explain these results in more detail, we need to discuss what we mean by “blow-ups” and what it means for these to look like not necessarily one object but any one of a class of objects as we zoom in on harmonic measure. There are two ways we can consider this. Firstly, we can look at the Hausdorff convergence of rescaled copies of the support of a measure as we zoom in. To do this, we follow the framework of [Badger and Lewis 2015].

Definition 1.1. Let $A \subset \mathbb{R}^{n+1}$ be a set. For $x \in A$, $r > 0$, and \mathcal{S} a collection of sets, define

$$\Theta_A^{\mathcal{S}}(x, r) = \inf_{S \in \mathcal{S}} \max \left\{ \sum_{a \in A \cap B(x, r)} \frac{\text{dist}(a, x + S)}{r}, \sum_{z \in (x + S) \cap B(x, r)} \frac{\text{dist}(z, A)}{r} \right\}.$$

We say $x \in A$ is a \mathcal{S} point of A if $\lim_{r \rightarrow 0} \Theta_A^{\mathcal{S}}(x, r) = 0$. We say A is *locally bilaterally well approximated* by \mathcal{S} (or simply $\text{LBWA}(\mathcal{S})$) if, for all $\varepsilon > 0$ and all compact sets $K \subset A$, there is $r_{\varepsilon, K} > 0$ such that $\Theta_A^{\mathcal{S}}(x, r) < \varepsilon$ for all $x \in K$ and $0 < r < r_{\varepsilon, K}$.

Thus, for $x \in A$ to be an \mathcal{S} -point means that, as we zoom in on A at the point x , the set A resembles more and more an element of \mathcal{S} (though that element may change as we zoom in).

Secondly, we can look at the weak convergence of rescaled copies of the measure itself. To do this, we follow the framework of [Preiss 1987]. For $a \in \mathbb{R}^{n+1}$ and $r > 0$, set

$$T_{a,r}(x) = \frac{x - a}{r}.$$

Note that $T_{a,r}(B(a, r)) = B(0, 1)$. Given a Radon measure μ , the notation $T_{a,r}[\mu]$ is the image measure of μ by $T_{a,r}$; that is,

$$T_{a,r}[\mu](A) = \mu(rA + a), \quad A \subset \mathbb{R}^{n+1}.$$

Here and later, for a function f and a measure μ , we write $f[\mu]$ to denote the push-forward measure $f[\mu](A) = \mu(f^{-1}(A))$.

Definition 1.2. We say that ν is a *tangent measure* of μ at a point $a \in \mathbb{R}^{n+1}$ if ν is a nonzero Radon measure on \mathbb{R}^{n+1} and there are sequences $c_i > 0$ and $r_i \downarrow 0$ so that $c_i T_{a, r_i}[\mu]$ converges weakly to ν as $i \rightarrow \infty$ and write $\nu \in \text{Tan}(\mu, a)$.

That is, ν is a tangent measure of μ at a point ξ if, as we zoom in on μ at ξ for a sequence of scales, the rescaled μ converges weakly to ν .

The collections of measures and sets that we will consider are associated to zero sets of harmonic functions. Let H denote the set of harmonic functions vanishing at the origin, $P(k)$ denote the set of harmonic polynomials h of degree k such that $h(0) = 0$ and $F(k)$ denote the set of homogeneous polynomials of degree k . For $h \in H$, we define

$$\Sigma_h = \{h=0\}, \quad \Omega_h = \{h>0\},$$

and

$$\mathcal{H} = \{\omega_h : h \in H\}, \quad \mathcal{P}(k) = \{\omega_h : h \in P(k)\}, \quad \mathcal{F}(k) = \{\omega_h : h \in F(k)\},$$

where

$$\omega_h = -\nu_{\Omega_h} \cdot \nabla h \, d\sigma_{\Sigma_h}.$$

Also set

$$\mathcal{P}_\Sigma(k) = \{\Sigma_h : h \in P(1) \cup \dots \cup P(k)\}, \quad \mathcal{F}_\Sigma(k) = \{\Sigma_h : h \in F(k)\}$$

and

$$\mathcal{H}_\Sigma = \{\Sigma_h : h \in H\}.$$

Here $\nu_{\Omega_h}(x)$ stands for the measure-theoretic unit outward normal of Ω_h at $x \in \partial^* \Omega_h$, the reduced boundary of Ω_h . Now h is a harmonic function and thus, real analytic, which implies that Σ_h is an n -dimensional real analytic variety; hence, Ω_h is a set of locally finite perimeter and one can prove that $\mathcal{H}^n(\partial \Omega_h \setminus \partial^* \Omega_h) = 0$, where \mathcal{H}^n stands for the n -Hausdorff measure. Notice now that $\nu_{\Omega_h}(x)$ is defined at \mathcal{H}^n -almost every point of Σ_h and σ_{Σ_h} is the usual surface measure. For a detailed proof of this see [Azzam et al. 2017b, p. 21].

In the rest of the paper we will be dealing with unbounded domains, i.e., open and connected sets in \mathbb{R}^{n+1} , with $n \geq 2$.

We summarize the best results to date. We first mention a result by the authors, Tolosa, and Volberg.

Theorem 1.3 [Azzam et al. 2016c; 2017b]. *Let $\Omega^\pm \subset \mathbb{R}^{n+1}$ be two disjoint domains and $\omega^\pm = \omega_{\Omega^\pm}^{x_\pm}$ for some $x_\pm \in \Omega^\pm$. If ω^\pm are mutually absolutely continuous on E , then for ω^\pm -a.e. $\xi \in E$ we have $\text{Tan}(\omega^\pm, \xi) \subset \mathcal{F}(1)$ and $\omega^+|_E$ can be covered up to a set of ω^+ -measure zero by n -dimensional Lipschitz graphs. Furthermore, if $\partial \Omega^\pm$ are CDC, then $\lim_{r \rightarrow 0} \Theta_{\partial \Omega^+}^{\mathcal{F}_\Sigma(1)}(\xi, r) = 0$ for ω^+ -a.e. $\xi \in E$.*

This was originally shown by Bishop, Carleson, Garnett, and Jones [Bishop et al. 1989] for simply connected planar domains. Later, Kenig, Preiss and Toro showed that, under the same assumptions, provided that the domain is also 2-sided locally NTA, it holds that $\dim \omega^+ = n$ (but not that ω^+ is rectifiable).

Below we summarize the results so far in the situation when Ω is 2-sided NTA and the interior and exterior harmonic measures are VMO equivalent, which brings together results and techniques from Badger [2011; 2013] and Kenig and Toro [2006].

Theorem 1.4. *Let $\Omega^+ \subset \mathbb{R}^{n+1}$ and $\Omega^- = \text{ext}(\Omega^+)$ be NTA domains, and let ω^\pm be the harmonic measure in Ω^\pm with pole $x^\pm \in \Omega^\pm$. Assume that ω^+ and ω^- are mutually absolutely continuous and $f := d\omega^-/d\omega^+$ satisfies $\log f \in \text{VMO}(d\omega^+)$. Then, there exists $d \in \mathbb{N}$ (depending on n and the NTA constants) such that the boundary $\partial\Omega^+$ is LBWA($\mathcal{P}_\Sigma(d)$) and may be decomposed into sets $\Gamma_1, \dots, \Gamma_d$ satisfying the following:*

- (1) For $1 \leq k \leq d$, $\Gamma_k = \{\xi \in \partial\Omega^+ : \text{Tan}(\omega^+, \xi) \subset \mathcal{F}(k)\}$.
- (2) $\Gamma_1 \cup \dots \cup \Gamma_d = \partial\Omega^+$.
- (3) $\lim_{r \rightarrow 0} \Theta_{\partial\Omega^+}^{\mathcal{F}_\Sigma(k)}(\xi, r) = 0$ for $\xi \in \Gamma_k$.

The work of [Badger et al. 2017] studies the geometric structure of the set as well as the tangent measure structure using the conclusions of the results above. We refer to their work for more details.

1B. Blowups of elliptic measures. In this paper, our objective is to recreate some parts of these results for a class of elliptic measures. Admittedly, there are more results that could be generalized to this setting, like Tsirelson's theorem (using the method of [Tolsa and Volberg 2018]), but we content ourselves with the present results to convey the flexibility of the method.

Let $\Omega \subset \mathbb{R}^{n+1}$ be open and $A = A(\cdot) = (a_{ij}(\cdot))_{1 \leq i, j \leq n+1}$ be a matrix with real measurable coefficients in Ω . We say that A is a *uniformly elliptic matrix* in Ω with constant $\Lambda \geq 1$ and write $A \in \mathcal{A}$ if it satisfies the following conditions:

$$\Lambda^{-1}|\xi|^2 \leq \langle A(x)\xi, \xi \rangle \quad \text{for a.e. } x \in \Omega \text{ and for all } \xi \in \mathbb{R}^{n+1}, \quad (1-1)$$

$$\langle A(x)\xi, \eta \rangle \leq \Lambda|\xi||\eta| \quad \text{for a.e. } x \in \Omega \text{ and for all } \xi, \eta \in \mathbb{R}^{n+1}. \quad (1-2)$$

Notice that the matrix is *possibly nonsymmetric* and has variable coefficients. If $A \in \mathcal{A}$, we define a *uniformly elliptic operator* associated with A by

$$L_A = -\text{div}(A(\cdot)\nabla).$$

We will let $\omega_{\Omega}^{A,x}$ denote the L_A -harmonic measure in Ω with pole at x (see Section 11 in [Heinonen et al. 1993] for the definition), which we also call *elliptic measure*. It is clear that the transpose matrix of A , which we denote by A^T , is also uniformly elliptic in Ω . Finally, a function $u : \Omega \rightarrow \mathbb{R}$ that satisfies the equation $L_A u = 0$ in the weak sense is called L_A -harmonic. We will denote by \mathcal{C} the subclass of \mathcal{A} consisting of matrices with constant entries.

To make sense of tangent measures of an elliptic measure at a point ξ in its support, we need to assume that the coefficients A do not oscillate too much there on small scales.

Definition 1.5. Let $\Omega \subset \mathbb{R}^{n+1}$ and let L_A be an elliptic operator on Ω . For a compact set $K \subset \partial\Omega$, we will say that the coefficients of L_A have *vanishing mean oscillation on K* with respect to Ω (or just

$L_A \in \text{VMO}(\Omega, K)$) if

$$\lim_{r \rightarrow 0} \sup_{\xi \in K} \frac{1}{r^{n+1}} \inf_{C \in \mathcal{C}} \int_{B(\xi, r) \cap \Omega} |A(x) - C| dx = 0. \quad (1-3)$$

We also say the coefficients of L_A have VMO at $\xi \in \partial\Omega$ if

$$\lim_{r \rightarrow 0} \frac{1}{r^{n+1}} \inf_{C \in \mathcal{C}} \int_{B(\xi, r) \cap \Omega} |A(x) - C| dx = 0. \quad (1-4)$$

Much like the harmonic case, the tangent measures we will obtain are supported on zero sets of elliptic polynomials associated with an elliptic operator with constant coefficients. For a constant-coefficient matrix A with real entries, we will denote by H_A the set of L_A -harmonic functions u vanishing at zero, i.e., those functions u for which

$$\int A \nabla u \nabla \varphi dx = 0 \quad \text{for all } \varphi \in C_c^\infty(\mathbb{R}^{n+1}) \quad \text{and} \quad u(0) = 0.$$

We also let $P_A(k)$ denote the set of L_A -harmonic polynomials of degree k vanishing at the origin, and $F_A(k) \subset P_A(k)$ the subset of homogeneous L_A -harmonic polynomials of degree k . When $A = I$, we will simply write $F(k)$, $P(k)$ and H in place of $F_A(k)$, $P_A(k)$ and H_A .

For $h \in H_A$, we will write

$$d\omega_h^A = -\nu_{\Omega_h} \cdot A \nabla h d\sigma_{\Sigma_h},$$

where σ_S stands for the surface measure on a surface S and ν is the outward normal vector at $x \in \partial^* \Omega_h$, the reduced boundary of Ω_h . Once more, we used that h is real analytic since A has constant coefficients and $L_A h = 0$; see, e.g., Proposition 11.3 in [Mitrea 2013]. Again, when A is the identity, we will drop the superscripts and, for example, write ω_h in place of ω_h^A . For $\mathcal{S} \subset \mathcal{C}$, we write

$$\begin{aligned} \mathcal{H}_{\mathcal{S}} &= \{\omega_h^A : h \in H_A, A \in \mathcal{S}\}, \quad \mathcal{P}_{\mathcal{S}}(k) = \{\omega_h^A : h \in P_A(k), A \in \mathcal{S}\}, \quad \mathcal{F}_{\mathcal{S}}(k) = \{\omega_h^A : h \in F_A(k), A \in \mathcal{S}\}, \\ \mathcal{H}_A &= \mathcal{H}_{\{A\}}, \quad \mathcal{P}_A = \mathcal{P}_{\{A\}}, \quad \mathcal{F}_A = \mathcal{F}_{\{A\}}, \end{aligned}$$

and define $\mathcal{H}_{\mathcal{S}, \Sigma}$, $\mathcal{P}_{\mathcal{S}, \Sigma}$, and $\mathcal{F}_{\mathcal{S}, \Sigma}$ as we did before. Observe that $\mathcal{F}_{\mathcal{C}}(1) = \mathcal{F}_A(1) = \mathcal{F}(1)$ for any $A \in \mathcal{C}$.

Our results also recover some LBWA properties implied in previous results if we consider domains satisfying the capacity density condition (CDC), whose complements also satisfy the CDC (see Definition 3.3 below) and whose associated elliptic measures are doubling. Examples of domains satisfying these conditions are NTA domains and, by [Martio 1979, Theorem 3.1], any uniform domain Ω for which there is $s > n - 1$ such that $\mathcal{H}_{\infty}^s(B(\xi, r) \cap \partial\Omega)/r^s \geq c > 0$ for all $\xi \in \partial\Omega$ and $r > 0$ is a CDC domain.

Our first result extends the work of [Kenig et al. 2009] to the elliptic case, and for domains beyond NTA. First, recall the dimension of a measure μ .

For a Borel measure μ in \mathbb{R}^{n+1} , we define the Hausdorff dimension of μ by

$$\dim(\mu) = \inf\{\dim(Z) : \mu(\mathbb{R}^{n+1} \setminus Z) = 0\}.$$

In practice, it is easier to compute this dimension as follows. Define *lower* and *upper pointwise dimension* at a point $x \in \text{supp } \mu$ to be

$$\underline{d}_\mu(x) = \liminf_{r \rightarrow 0} \frac{\log \mu(B(x, r))}{\log r} \quad \text{and} \quad \bar{d}_\mu(x) = \limsup_{r \rightarrow 0} \frac{\log \mu(B(x, r))}{\log r}.$$

We call the common value $\underline{d}_\mu(x) = \bar{d}_\mu(x) = d_\mu(x)$, if it exists, the *pointwise dimension* of μ at $x \in \text{supp } \mu$. It is shown in [Barreira and Wolf 2006, Proposition 3] that

$$\dim(\mu) = \text{ess sup}\{\underline{d}_\mu(x) : x \in \Lambda\}.$$

Theorem I. *Let $\Omega^\pm \subset \mathbb{R}^{n+1}$ be two disjoint domains and let L_A be a uniformly elliptic operator on $\Omega^+ \cup \Omega^-$. Let also $\omega^\pm = \omega_{\Omega^\pm}^{L_A, x^\pm}$ for some $x_\pm \in \Omega^\pm$ be the L_A -harmonic measures in the respective domains and L_A be in $\text{VMO}(\Omega^+ \cup \Omega^-, \xi)$ at ω^+ -almost every $\xi \in E \subset \partial\Omega^+ \cap \partial\Omega^-$ with respect to either Ω^\pm . If ω^\pm are mutually absolutely continuous on E , then for ω^\pm -a.e. $\xi \in E$ we have $\text{Tan}(\omega^\pm, \xi) \subset \mathcal{F}(1)$ and $\dim \omega^\pm|_E = n$. Furthermore, if $\partial\Omega^\pm$ are CDC, then $\lim_{r \rightarrow 0} \Theta_{\partial\Omega^+}^{\mathcal{F}_{\Sigma}(1)}(\xi, r) = 0$ for ω^+ -a.e. $\xi \in E$.*

Kenig, Preiss, and Toro originally showed this if Ω^\pm were both NTA domains, and the dimension was computed by estimating the Hausdorff dimension directly from above and then using the monotonicity formula of Alt, Caffarelli, and Friedman [Alt et al. 1984] to estimate it from below. The latter is not available for L -harmonic functions when L satisfies the VMO condition above. For this reason, we use instead the fact that the tangent measures are all flat, which forces ω^\pm to decay like a planar n -dimensional Hausdorff measure on small scales.

Assuming a VMO condition on the interior and exterior elliptic measures, we can also obtain the results of [Kenig and Toro 2006] and [Badger 2011] for elliptic measures on domains that do not have to be NTA. We first state a pointwise version of these.

Theorem II. *Let Ω^+ be a domain in \mathbb{R}^{n+1} , let $\Omega^- := \text{ext}(\Omega^+)$ be its exterior, and let L_A be a uniformly elliptic operator in $\Omega^+ \cup \Omega^-$. Denote by ω^\pm the L_A -harmonic measures of Ω^\pm with poles at some points $x^\pm \in \Omega^\pm$, and assume that ω^\pm are mutually absolutely continuous with $f = d\omega^-/d\omega^+$. If for a fixed $\xi \in \partial\Omega^+ \cap \partial\Omega^-$ it holds that $L_A \in \text{VMO}(\Omega^+ \cup \Omega^-, \xi)$,*

$$\lim_{r \rightarrow 0} \left(\int_{B(\xi, r)} f d\omega^+ \right) \exp \left(- \int_{B(\xi, r)} \log f d\omega^+ \right) = 1, \quad (1-5)$$

and $\text{Tan}(\omega^+, \xi) \neq \emptyset$, then $\text{Tan}(\omega^+, \xi) \subset \mathcal{F}_\mathcal{C}(k)$ for some k and

$$\limsup_{r \rightarrow 0} \frac{\omega^+(B(\xi, 2r))}{\omega^+(B(\xi, r))} < \infty. \quad (1-6)$$

If Ω^\pm have the CDC, then additionally

$$\lim_{r \rightarrow 0} \Theta_{\partial\Omega^+}^{\mathcal{F}_\mathcal{C}, \Sigma(k)}(\xi, r) = 0.$$

It is well known that $\text{Tan}(\omega^+, \xi) \neq \emptyset$ whenever ω^+ satisfies the pointwise doubling condition (1-6). In our situation, however, we do not assume that, but we get it for free since $\mathcal{F}_\mathcal{C}(k)$ is compact (see [Badger 2011, Lemma 4.10] for the harmonic case and Theorem 2.4 below).

One might have guessed that a pointwise version of [Theorem 1.4](#) would have assumed instead that

$$\lim_{r \rightarrow 0} \int_{B(\xi, r)} \left| f - \int_{B(\xi, r)} \log f \, d\omega^+ \right| d\omega^+ = 0,$$

but we were not able to show that this implies [Theorem II](#). However, under certain conditions they are equivalent. We will discuss this matter in depth in [Section 6](#) below.

Next, we state a global version.

Theorem III. *Let $\Omega^\pm \subset \mathbb{R}^{n+1}$ be two disjoint domains in \mathbb{R}^{n+1} with common boundary, and let L_A be a uniformly elliptic operator in $\Omega^+ \cup \Omega^-$ such that $L_A \in \text{VMO}(\Omega^+ \cup \Omega^-, \xi)$ at every $\xi \in \partial\Omega^+ \cap \partial\Omega^-$. Denote by ω^\pm the L_A -harmonic measures of Ω^\pm with poles at some points $x^\pm \in \Omega^\pm$. If ω^+ is C -doubling, ω^\pm are mutually absolutely continuous, and $\log f = \log(d\omega^-/d\omega^+) \in \text{VMO}(d\omega^+)$, then there is d depending on n and the doubling constant so that, for every compact subset $K \subseteq \partial\Omega^+$,*

$$\lim_{r \rightarrow 0} \sup_{\xi \in K} d_1(T_{\xi, r}[\omega^+], \mathcal{P}_\mathcal{C}(d)) = 0. \quad (1-7)$$

If additionally Ω^\pm are CDC domains, then for any compact set $K \subseteq \partial\Omega$

$$\lim_{r \rightarrow 0} \sup_{\xi \in K} \Theta_{\partial\Omega^+}^{\mathcal{P}_\mathcal{C}, \Sigma(d)}(\xi, r) = 0.$$

That is, $\partial\Omega^+ \in \text{LBWA}(\mathcal{P}_\mathcal{C}, \Sigma(d))$.

See [Section 2](#) for the definition of $d_1(\cdot, \mathcal{P}_\mathcal{C}(d))$, which is essentially a distance between measures and the set $\mathcal{P}_\mathcal{C}(d)$.

The proof of [Theorem II](#) involves some useful lemmas about tangent measures that may be of independent interest. Specifically, we refer the reader to [Lemma 2.10](#).

Over the course of working on this manuscript, we also resolved a question left open in [\[Badger 2011\]](#) (see the discussion on page 861 of that work).

Proposition I. *The d -cone $\mathcal{P}(k)$ has compact basis for each $k \in (0, n]$.*

See [Section 2](#) for the definition of compact bases. A consequence of this result is that we can improve on the following theorem of Badger.

Theorem 1.6 [\[Badger 2011, Theorem 1.1\]](#). *Let $\Omega \subset \mathbb{R}^{n+1}$ be an NTA domain with harmonic measure ω and let $\xi \in \partial\Omega$. If $\text{Tan}(\omega, \xi) \subset \mathcal{P}(d)$, then $\text{Tan}(\omega, \xi) \subset \mathcal{F}(k)$ for some $k \leq d$.*

In the proof of this result, Badger relied on the NTA assumption to conclude that $\text{Tan}(\omega, \xi)$ was compact. By using [Proposition I](#) (whose proof is rather short), the compactness of $\mathcal{F}(k)$ (to which much of the proof of [Theorem 1.6](#) is dedicated), and a connectivity theorem of Preiss, we can improve this by showing that, to get the same conclusion, no a priori information about the geometry of ω is needed; it need not have been a harmonic measure, let alone one for an NTA domain:

Proposition II. *Let ω be a Radon measure in \mathbb{R}^{n+1} and $\xi \in \mathbb{R}^{n+1}$ such that $\text{Tan}(\omega, \xi) \subset \mathcal{P}(k)$ for some integer k . If $\text{Tan}(\omega, \xi) \cap \mathcal{F}(k) \neq \emptyset$ for some integer k , then $\text{Tan}(\omega, \xi) \subset \mathcal{F}(k)$.*

1C. Rectifiability and elliptic measure for uniform domains. The blow-up arguments we use also have an application to studying the relationship between rectifiability and harmonic measure, a subject in which there have been a flurry of results in the last few years. For simply connected planar domains, the problem of when harmonic measure is absolutely continuous with respect to \mathcal{H}^1 is classical. Bishop and Jones [1990] showed that, if Ω is simply connected, $\omega_\Omega^x \ll \mathcal{H}^1$ on the subset of any Lipschitz curve intersecting $\partial\Omega$. Conversely, Pommerenke [1986] showed that if $\omega_\Omega \ll \mathcal{H}^1$ on a subset $E \subset \partial\Omega$, then that set can be covered by Lipschitz graphs up to a set of harmonic measure zero. In fact, a much earlier result of the Riesz brothers says that any Jordan domain has harmonic measure and is \mathcal{H}^1 mutually absolutely continuous if and only if the boundary is rectifiable; see [Riesz and Riesz 1920] or [Garnett and Marshall 2005, Chapter VI.1].

In higher dimensions, the problem is more delicate. There are some examples of simply connected domains $\Omega \subset \mathbb{R}^{n+1}$ with n -rectifiable boundaries of finite \mathcal{H}^n -measure so that either $\omega_\Omega \not\ll \mathcal{H}^n$ or $\mathcal{H}^n \not\ll \omega_\Omega$; see [Wu 1986; Ziemer 1974]. David and Jerison [1990] showed that mutual absolute continuity occurs for NTA domains with Ahlfors–David regular boundaries. Building on that, Badger [2012] showed that $\mathcal{H}^n \ll \omega_\Omega$ if Ω is an NTA domain whose boundary simply has locally finite \mathcal{H}^n -measure, although we showed with Tolsa that the converse relation $\omega_\Omega \ll \mathcal{H}^n$ could be false for such domains [Azzam et al. 2017c].

However, in [Azzam et al. 2016b], along with Hofmann, Martell, Mayboroda, Tolsa, and Volberg, we showed that for *any* domain $\Omega \subset \mathbb{R}^{n+1}$ and $E \subset \partial\Omega$ with $\omega_\Omega(E) > 0$ and $\mathcal{H}^n(E) < \infty$, if $\omega_\Omega \ll \mathcal{H}^n$ on E , then E may be covered up to ω_Ω -measure zero by Lipschitz graphs. By a theorem of Wolff, harmonic measure in the plane lies on a set of σ -finite \mathcal{H}^1 -measure, and so the assumption that $\mathcal{H}^1(E) < \infty$ is unnecessary in this case (although very necessary in higher dimensions due to the existence of Wolff snowflakes). With Akman, we developed a converse for domains $\Omega \subset \mathbb{R}^{n+1}$ with *big complements*, meaning

$$\mathcal{H}_\infty^n(B(\xi, r) \setminus \Omega) \geq cr^n \quad \text{for all } \xi \in \partial\Omega \text{ and } 0 < r < \text{diam } \partial\Omega. \quad (1-8)$$

We showed that, for such domains, $\omega_\Omega \ll \mathcal{H}^n$ on the subset of any n -dimensional Lipschitz graph [Akman et al. 2019], and hence, for these domains, we know that absolute continuity is equivalent to rectifiability of harmonic measure (versus rectifiability of the boundary).

There are fewer positive results concerning absolute continuity and rectifiability of *elliptic* measures. Even in the case of the half-plane, without some extra assumptions on the behavior of the elliptic coefficients, elliptic measure can be singular [Caffarelli et al. 1981; Sweezy 1992; Wu 1994], and some sort of Dini condition on the coefficients near the boundary is needed [Fabes et al. 1984; Fefferman et al. 1991]. For example, Kenig and Pipher [2001], considered the following condition.

Definition 1.7. Let $\delta(x) = \text{dist}(x, \partial\Omega)$. We will say that an elliptic operator $L = -\text{div } A \nabla$ satisfies the *Kenig–Pipher condition* (or *KP-condition*) if $A = (a_{ij}(x))$ is a uniformly elliptic real matrix that has distributional derivatives such that

$$\varepsilon_\Omega^L(z) := \sup\{|\delta(x)|^2 |\nabla a_{ij}(x)|^2 : x \in \tfrac{1}{2}B(z, \delta(z)), 1 \leq i, j \leq n+1\} \quad (1-9)$$

is a Carleson measure in Ω , by which we mean that for all $x \in \partial\Omega$ and $r \in (0, \text{diam } \partial\Omega)$,

$$\int_{B(x,r) \cap \Omega} \varepsilon_{\Omega}^L(z) dz \leq Cr^n.$$

In [Kenig and Pipher 2001], they showed that for Lipschitz domains in \mathbb{R}^{n+1} , elliptic operators satisfying the KP-condition give rise to elliptic measures which are A_{∞} -equivalent to surface measure. In fact, it was proved in [Hofmann et al. 2017] that the same result can be obtained under the following more general assumptions on the coefficients:

$$(\widetilde{\text{KP}}) = \begin{cases} \nabla a_{ij} \in \text{Lip}_{\text{loc}}(\Omega), \\ \|\delta_{\Omega} |\nabla a_{ij}|\|_{L^{\infty}(\Omega)} < \infty, \\ \delta(x) |\nabla a_{ij}(x)|^2 \text{ is a Carleson measure} \end{cases} \quad (1-10)$$

for $1 \leq i, j \leq n+1$. Akman, Badger, Hofmann, and Martell observed in [Akman et al. 2017, Section 3.2] that, using the same arguments in [David and Jerison 1990], this result can be extended to NTA domains with Ahlfors–David regular boundaries. They used this fact to show that, on a *uniform domain* Ω (see Definition 8.1 below) with Ahlfors–David regular boundary, if L_A is a *symmetric* elliptic operator satisfying a local L^1 version of (1-9), i.e., $A \in \text{Lip}_{\text{loc}}(\Omega)$ and $\sup\{|\nabla a_{ij}(x)| : x \in \frac{1}{2}B(z, \delta(z)), 1 \leq i, j \leq n+1\}$ is a Carleson measure with Carleson constant depending on the ball, then $\mathcal{H}^n \ll \omega_{\Omega}^L$ implies n -rectifiability of the boundary.

Using our blow-up arguments, we can obtain the following improvement.

Theorem IV. *Let $\Omega \subset \mathbb{R}^{n+1}$ be a uniform CDC domain so that $\mathcal{H}^n|_{\partial\Omega}$ is locally finite. Let $\omega_{\Omega}^{L_A}$ be the L_A -harmonic measure associated to a (possibly nonsymmetric) elliptic operator satisfying (1-1) and (1-2). Let $E \subseteq \partial\Omega$ be a set with $\mathcal{H}^n(E) > 0$ such that $\mathcal{H}^n \ll \omega_{\Omega}^{L_A}$ on E and for \mathcal{H}^n -a.e. $\xi \in E$*

$$\theta_{\partial\Omega,*}^n(\xi, r) := \liminf_{r \rightarrow 0} \frac{\mathcal{H}^n(B(\xi, r) \cap \partial\Omega)}{(2r)^n} > 0$$

and A has vanishing mean oscillation at ξ . Then E is n -rectifiable.

Surprisingly, to get this improvement requires a very different set of techniques than originally considered in [Akman et al. 2017]. Let us point out that the argument therein uses the symmetry hypothesis on the coefficients in a significant way and does not seem easy to extend to the nonsymmetric case unless one additionally assumes that $\mathcal{H}^n \ll \omega_{\Omega}^{L_{A^T}}$.

Having VMO coefficients \mathcal{H}^n -a.e. on $\partial\Omega$ is natural as it is implied by the Carleson condition considered in [Akman et al. 2017; Kenig and Pipher 2001] by the following proposition:

Proposition III. *Let $\Omega \subset \mathbb{R}^{n+1}$ be a uniform domain and suppose that A is an elliptic matrix satisfying (1-1) and (1-2) such that $A \in \text{Lip}_{\text{loc}}(\Omega)$ and, for some ball B_0 centered on $\partial\Omega$,*

$$\int_{B_0} \delta(x) |\nabla a_{ij}(x)|^2 dx < \infty. \quad (1-11)$$

Then $L_A \in \text{VMO}(\Omega, \xi)$ for \mathcal{H}^n -a.e. $\xi \in B_0 \cap \partial\Omega$.

Discussion of related results. Near the completion of this work, we learned that Toro and Zhao [2017] simultaneously proved that $\mathcal{H}^n \ll \omega_\Omega$ implies rectifiability of the boundary if $\Omega \subseteq \mathbb{R}^{n+}$ is a uniform domain with Ahlfors–David n -regular boundary and the elliptic coefficients are in $W^{1,1}(\Omega)$. They also exploit the vanishing oscillation of the coefficients at almost every boundary point (which they show is implied by the $W^{1,1}$ condition) in the context of uniform domains, though, their proof is distinct by their use of pseudotangents and stopping-time arguments.

1D. Notation. We will write $a \lesssim b$ if there is $C > 0$ so that $a \leq Cb$ and $a \lesssim_t b$ if the constant C depends on the parameter t . We write $a \approx b$ to mean $a \lesssim b \lesssim a$ and define $a \approx_t b$ similarly.

2. Tangent measures

2A. Cones and compactness. Given two Radon measures μ and σ , we set

$$F_B(\mu, \sigma) = \sup_f \int f d(\mu - \sigma),$$

where the supremum is taken over all the nonnegative 1-Lipschitz functions supported on B . For $r > 0$, we write

$$F_r(\mu, \nu) = F_{B(0,r)}, \quad F_r(\mu) = F_r(\mu, 0) = \int (r - |z|)_+ d\mu.$$

A set of Radon measures \mathcal{M} is a d -cone if $cT_{0,r}[\mu] \in \mathcal{M}$ for all $\mu \in \mathcal{M}$, $c > 0$ and $r > 0$. We say a d -cone has *closed (resp. compact) basis* if its basis $\{\mu \in \mathcal{M} : F_1(\mu) = 1\}$ is closed (resp. compact) with respect to the weak topology.

For a d -cone \mathcal{M} , $r > 0$, and μ a Radon measure with $0 < F_r(\mu) < \infty$, we define the *distance* between μ and \mathcal{M} as

$$d_r(\mu, \mathcal{M}) = \inf \left\{ F_r \left(\frac{\mu}{F_r(\mu)}, \nu \right) : \nu \in \mathcal{M}, F_r(\nu) = 1 \right\}.$$

Lemma 2.1 [Kenig et al. 2009, Section 2]. *Let μ be a Radon measure in \mathbb{R}^{n+1} and \mathcal{M} a d -cone. For $\xi \in \mathbb{R}^{n+1}$ and $r > 0$:*

- (1) $T_{\xi,r}[\mu](B(0,s)) = \mu(B(\xi, sr))$.
- (2) $\int f dT_{\xi,r}[\mu] = \int f \circ T_{\xi,r} d\mu$.
- (3) $F_{B(\xi,r)}(\mu) = rF_1(T_{\xi,r}[\mu])$.
- (4) $F_{B(\xi,r)}(\mu, \nu) = rF_1(T_{\xi,r}[\mu], T_{\xi,r}[\nu])$.
- (5) $\mu_i \rightarrow \mu$ weakly if and only if $F_r(\mu_i, \mu) \rightarrow 0$ for all $r > 0$.
- (6) $d_r(\mu, \mathcal{M}) \leq 1$.
- (7) $d_r(\mu, \mathcal{M}) = d_1(T_{0,r}[\mu], \mathcal{M})$.
- (8) If $\mu_i \rightarrow \mu$ weakly and $F_r(\mu) > 0$, then $d_r(\mu_i, \mathcal{M}) \rightarrow d_r(\mu, \mathcal{M})$.

Lemma 2.2 [Kenig et al. 2009, Remark 2.13]. *A d -cone \mathcal{M} of Radon measures in \mathbb{R}^{n+1} has a closed basis if and only if it is a relatively closed subset of the nonzero Radon measures in \mathbb{R}^{n+1} .*

Proof. One direction is obvious, so suppose \mathcal{M} has closed basis and $\mu_i \in \mathcal{M}$ converges weakly to some nonzero Radon measure μ . Then $F_r(\mu) > 0$ for some $r > 0$. The set $\{v \in \mathcal{M} : F_1(v) = 1\}$ is closed by assumption, and since \mathcal{M} is a d -cone, the set $\{v \in \mathcal{F} : F_r(v) = 1\}$ is also closed. Hence, since $\mu_i / F_r(\mu_i) \rightarrow \mu / F_r(\mu)$, we know $\mu / F_r(\mu) \in \mathcal{M}$, and thus $\mu \in \mathcal{M}$. \square

Lemma 2.3. *If μ is a nonzero Radon measure and \mathcal{M} is a d -cone with closed basis, then $\mu \in \mathcal{M}$ if and only if $d_r(\mu, \mathcal{M}) = 0$ for all $r > 0$ for which $F_r(\mu) > 0$.*

Proof. Suppose $d_r(\mu, \mathcal{M}) = 0$ for all $r > 0$ for which $F_r(\mu) > 0$. For $j \in \mathbb{N}$ large enough, we can find a sequence $\mu_{j,k} \in \mathcal{M}$ such that

$$F_j(\mu_{j,k}) = 1 \quad \text{and} \quad \lim_{k \rightarrow \infty} F_j\left(\frac{\mu}{F_j(\mu)}, \mu_{j,k}\right) = 0. \quad (2-1)$$

In particular, we can pass to a subsequence so that $\mu_{j,k}$ converges weakly in $B(0, j)$ to a measure μ_j supported in $B(0, j)$ with $F_j(\mu_j) = 1$. In view of (2-1), the latter implies $\mu = F_j(\mu)\mu_j$ in $B(0, j)$, and thus

$$F_j(\mu)\mu_j \rightharpoonup \mu.$$

Since $\mu_{j,k} \rightharpoonup \mu_j$ and $F_j(\mu) \neq 0$ for j large, we can pick k_j so that

$$F_j(\mu_{j,k_j}, \mu_j) < \frac{1}{j F_j(\mu)}.$$

In particular, for any $r > 0$ and $j > r$,

$$\begin{aligned} F_r(\mu_{j,k_j} F_j(\mu), \mu) &\leq F_r(\mu_{j,k_j} F_j(\mu), \mu_j F_j(\mu)) + F_r(\mu_j F_j(\mu), \mu) \\ &\leq F_j(\mu_{j,k_j} F_j(\mu), \mu_j F_j(\mu)) + F_r(\mu_j F_j(\mu), \mu) \\ &< \frac{1}{j} + F_r(\mu_j F_j(\mu), \mu) \rightarrow 0. \end{aligned}$$

Thus, $\mu_{j,k_j} F_j(\mu) \rightharpoonup \mu$. By Lemma 2.2, \mathcal{M} is closed, and since we have $\mu_{j,k_j} F_j(\mu) \in \mathcal{M}$ for all j , this implies $\mu \in \mathcal{M}$. The other implication is trivial. \square

Theorem 2.4 [Preiss 1987, Corollary 2.7]. *Let μ be a Radon measure on \mathbb{R}^{n+1} and $\xi \in \text{supp } \mu$. Then $\text{Tan}(\mu, \xi)$ has compact basis if and only if*

$$\limsup_{r \rightarrow 0} \frac{\mu(B(\xi, 2r))}{\mu(B(\xi, r))} < \infty. \quad (2-2)$$

In this case, for any $v \in \text{Tan}(\mu, \xi)$, it holds that $0 \in \text{supp } v$ and

$$\frac{v(B(0, 2r))}{v(B(0, r))} \leq \limsup_{\rho \rightarrow 0} \frac{\mu(B(\xi, 2\rho))}{\mu(B(\xi, \rho))} \quad \text{for all } r > 0.$$

Lemma 2.5 [Mattila 1995, Theorem 14.3]. *Let μ be a Radon measure on \mathbb{R}^{n+1} . If $\xi \in \mathbb{R}^{n+1}$ and (2-2) holds, then every sequence $r_i \downarrow 0$ contains a subsequence such that*

$$\frac{T_{\xi, r_j}[\mu]}{\mu(B(\xi, r_j))} \rightharpoonup v \quad (2-3)$$

for some measure $v \in \text{Tan}(\mu, \xi)$.

Having tangent measures that arise as limits of the form (2-3) is very convenient, but this limit does not always converge weakly to something. This may happen if μ is not pointwise doubling at the point a . However, all tangent measures are at least dilations of tangent measures arising in this way.

Lemma 2.6 [Mattila 1995, Remark 14.4(1)]. *Let μ be a nonzero Radon measure, $\xi \in \text{supp } \mu$, and $\nu \in \text{Tan}(\mu, \xi)$. Then there are $\rho_j \downarrow 0$ and $\rho, c > 0$ so that*

$$\frac{T_{\xi, \rho_j}[\mu]}{\mu(B(\xi, \rho_j))} \rightharpoonup cT_{0, \rho}[\nu] \quad \text{and} \quad cT_{0, \rho}[\nu](\mathbb{B}) > 0.$$

Proposition 2.7 [Preiss 1987, Proposition 2.2]. *Let \mathcal{M} be a d -cone. Then \mathcal{M} has compact basis if and only if for every $\lambda > 1$ there is $\tau > 1$ such that*

$$F_{\tau r}(\Psi) \leq \lambda F_r(\Psi) \quad \text{for every } \Psi \in \mathcal{M} \text{ and } r > 0. \quad (2-4)$$

In this case, $0 \in \text{supp } \Psi$ for all $\Psi \in \mathcal{M}$.

Theorem 2.8 [Mattila 1995, Theorem 14.16]. *Let μ be a Radon measure on \mathbb{R}^{n+1} . For μ -almost every $x \in \mathbb{R}^{n+1}$, if $\nu \in \text{Tan}(\mu, x)$, the following hold:*

- (1) $T_{y, r}[\nu] \in \text{Tan}(\mu, x)$ for all $y \in \text{supp } \nu$ and $r > 0$.
- (2) $\text{Tan}(\nu, y) \subset \text{Tan}(\mu, x)$ for all $y \in \text{supp } \nu$.

Lemma 2.9 [Badger 2011, Lemma 2.6]. *Let μ be a nonzero Radon measure on \mathbb{R}^{n+1} and $x \in \text{supp}(\mu)$. If $\nu \in \text{Tan}(\mu, x)$, then $\text{Tan}(\nu, 0) \subset \text{Tan}(\mu, x)$.*

2B. Connectivity of cones. The main tool from [Kenig et al. 2009; Badger 2011] is the following “connectivity” lemma, which was originally shown in [Kenig et al. 2009, Corollary 2.16] under the assumption that \mathcal{M} had compact basis. For our purposes, we need to remove this assumption.

Lemma 2.10. *Let \mathcal{F} and \mathcal{M} be d -cones and assume \mathcal{F} has compact basis. Furthermore, suppose that there is $\varepsilon_0 > 0$ such that for $\mu \in \mathcal{M}$, if there is $r_0 > 0$ so that $d_r(\mu, \mathcal{F}) \leq \varepsilon$ for all $r \geq r_0$, then $\mu \in \mathcal{F}$. For a Radon measure η and $x \in \text{supp } \eta$, if $\text{Tan}(\eta, x) \subset \mathcal{M}$ and $\text{Tan}(\eta, x) \cap \mathcal{F} \neq \emptyset$, then $\text{Tan}(\eta, x) \subset \mathcal{F}$.*

We will first require some lemmas.

Lemma 2.11. *Let \mathcal{F} be a d -cone with compact basis. There is $\beta > 0$ depending only on \mathcal{F} so that the following holds. Suppose ω is a Radon measure in \mathbb{R}^{n+1} , $\xi \in \text{supp } \omega$, $\text{Tan}(\omega, \xi) \cap \mathcal{F} \neq \emptyset$ and*

$$\limsup_{r \rightarrow 0} d_{r_0}(T_{\xi, r}[\omega], \mathcal{F}) \geq \varepsilon_0 > 0 \quad \text{for some } r_0 > 0.$$

Then for $\varepsilon < \varepsilon_0$ small enough, we may find $\mu \in \text{Tan}(\omega, \xi) \setminus \mathcal{F}$ so that

- (1) $d_{r_0}(\mu, \mathcal{F}) = \varepsilon$,
- (2) $d_r(\mu, \mathcal{F}) \leq \varepsilon$ for all $r > r_0$, and
- (3) $\mu(B(0, r)) \leq r^\beta \mu(B(0, 4r_0))$ for all $r \geq r_0$.

This is an adaptation of the proof of [Kenig et al. 2009, Corollary 2.16], but with some extra care.

Proof. Without loss of generality, we will assume $r_0 = 1$. Let $c_j > 0$ and $r_j \downarrow 0$ be such that $c_j T_{\xi, r_j}[\omega] \rightarrow \nu \in \mathcal{F}$. Since \mathcal{F} is compact, by [Proposition 2.7](#), $0 \in \text{supp } \nu$ and so $\nu(\mathbb{B}) > 0$. Thus, by [Lemma 2.1\(5\)](#), $c_j T_{\xi, r_j}[\omega](\mathbb{B}) > 0$ for j large. By [Lemma 2.1\(8\)](#), we have that, given $\varepsilon > 0$, for j large enough,

$$d_1(T_{\xi, r_j}[\omega], \mathcal{F}) = d_1(c_j T_{\xi, r_j}[\omega], \mathcal{F}) < \varepsilon. \quad (2-5)$$

Note that $0 \in \text{supp } T_{\xi, r_j}[\omega]$ since $\xi \in \text{supp } \omega$, and so there is no accidental dividing by zero in the definition of d_1 . By assumption, there is also $s_j \downarrow 0$ so that

$$d_1(T_{\xi, s_j}[\omega], \mathcal{F}) > \varepsilon. \quad (2-6)$$

We can assume $s_j < r_j$ by passing to a subsequence. Then by (2-5) and (2-6), let $\rho_j \in (s_j, r_j)$ be the maximal number such that

$$d_1(T_{\xi, \rho_j}[\omega], \mathcal{F}) = \varepsilon. \quad (2-7)$$

Then, by the maximality of ρ_j ,

$$\sup_{t \in [\rho_j, r_j]} d_1(T_{\xi, t}[\omega], \mathcal{F}) \leq \varepsilon. \quad (2-8)$$

We claim $\rho_j / r_j \rightarrow 0$. If not, then since $\rho_j / r_j \leq 1$, we may pass to a subsequence so that $\rho_j / r_j \rightarrow t \in (0, 1)$, and so

$$c_j T_{\xi, \rho_j}[\omega] = T_{0, \rho_j / r_j}[c_j T_{\xi, r_j}[\omega]] \rightarrow T_{0, t}[\nu] \in \mathcal{F},$$

which contradicts (2-7). Thus, $\rho_j / r_j \rightarrow 0$, and so (2-8) implies that for $\alpha \geq 1$, if j is large enough, we have $1 \leq \alpha < r_j / \rho_j$. If $\omega_j = T_{\xi, \rho_j}[\omega]$, then by [Lemma 2.1\(7\)](#), it holds that

$$d_\alpha(\omega_j, \mathcal{F}) = d_\alpha(T_{\xi, \rho_j}[\omega], \mathcal{F}) = d_1(T_{\xi, \alpha \rho_j}[\omega], \mathcal{F}) \stackrel{(2-8)}{\leq} \varepsilon, \quad (2-9)$$

which by (2-7) implies

$$d_1(\omega_j, \mathcal{F}) = \varepsilon > 0 \quad \text{and} \quad \limsup_{j \rightarrow \infty} d_r(\omega_j, \mathcal{F}) \leq \varepsilon \quad \text{for } r > 1. \quad (2-10)$$

For $r \geq 1$, let $\mu_{j,r} \in \mathcal{F}$ be such that $F_{\tau r}(\mu_{j,r}) = 1$ and

$$F_{\tau r}\left(\frac{\omega_j}{F_{\tau r}(\omega_j)}, \mu_{j,r}\right) < \frac{3}{2} d_{\tau r}(\omega_j, \mathcal{F}).$$

By (2-10), for j large enough,

$$F_r\left(\frac{\omega_j}{F_{\tau r}(\omega_j)}, \mu_{j,r}\right) \leq F_{\tau r}\left(\frac{\omega_j}{F_{\tau r}(\omega_j)}, \mu_{j,r}\right) < \frac{3}{2} d_{\tau r}(\omega_j, \mathcal{F}) < 2\varepsilon. \quad (2-11)$$

Since \mathcal{F} has compact basis, by [Proposition 2.7](#) with $\lambda = 2$, there is $\tau > 1$ depending only on \mathcal{F} so that (2-4) holds for $\mathcal{M} = \mathcal{F}$. Thus, if $\varepsilon < \frac{1}{8}$, by the triangle inequality for F_r and (2-11),

$$\frac{F_r(\omega_j)}{F_{\tau r}(\omega_j)} \geq F_r(\mu_{j,r}) - 2\varepsilon \geq \frac{1}{2} F_{\tau r}(\mu_{j,r}) - 2\varepsilon = \frac{1}{2} - 2\varepsilon > \frac{1}{4}. \quad (2-12)$$

Hence, for any $r \geq 1$,

$$F_{\tau r}(\omega_j) \leq 4F_r(\omega_j).$$

Set $\mu_j = \omega_j / F_1(\omega_j)$. Then iterating the above inequality and letting $j \rightarrow \infty$, we get that for all $\ell \in \mathbb{N}$,

$$\limsup_{j \rightarrow \infty} F_{\tau^\ell}(\mu_j) \leq 4^\ell.$$

This implies that we can pass to a subsequence so that μ_j converges weakly to a measure $\mu \in \text{Tan}(\omega, \xi)$. In particular, for $r \geq 1$, since $F_1(\mu_j) = 1$, we may compute

$$d_1(\mu, \mathcal{F}) = \lim_{j \rightarrow \infty} d_1(\mu_j, \mathcal{F}) = \lim_{j \rightarrow \infty} d_1(\omega_j, \mathcal{F}) \stackrel{(2-10)}{=} \varepsilon,$$

$$d_r(\mu, \mathcal{F}) = \lim_{j \rightarrow \infty} d_r(\mu_j, \mathcal{F}) = \lim_{j \rightarrow \infty} d_r(\omega_j, \mathcal{F}) \stackrel{(2-10)}{\leq} \varepsilon,$$

and

$$\tau^\ell \mu(B(0, \tau^\ell)) \leq F_{2\tau^\ell}(\mu) \leq 4^\ell F_2(\mu) \quad \text{for all } \ell \in \mathbb{N}. \quad (2-13)$$

Since $\tau > 1$, for any $r \geq 1$, there exists $\ell > 0$ such that $\tau^{\ell-1} < r \leq \tau^\ell$. If $\tau \in (1, 4)$, then (2-13) implies

$$\tau^\ell \mu(B(0, \tau^\ell)) \leq \tau^\alpha r^\alpha \mu(\overline{B(0, 2)}),$$

where $\alpha = 1/\log_4 \tau \in (1, \infty)$ and we used that $4^\ell = \tau^{\ell\alpha}$. Therefore,

$$\mu(B(0, r)) \leq \tau^{\alpha-\ell} r^\alpha \mu(\overline{B(0, 2)}),$$

and notice that $\tau^{\alpha-\ell} \leq 1$ whenever $\tau^\ell \geq 4$; i.e., the constant is independent of τ . In the case that $1 \leq r \leq \tau^\ell < 4$, we simply use that $B(0, r) \subset B(0, 4)$ to conclude that

$$\mu(B(0, r)) \leq \mu(B(0, 4)).$$

If $\tau \geq 4$, then (2-13) trivially gives

$$\tau^\ell \mu(B(0, \tau^\ell)) \leq 4^\ell \mu(\overline{B(0, 2)}) \leq \tau^\ell \mu(\overline{B(0, 2)}),$$

which can only be true if $r \leq \tau^\ell \leq 2$. Thus, $B(0, r) \subset B(0, 2)$ and (3) readily follows. \square

Corollary 2.12. *Let \mathcal{F} be a d -cone with compact basis. There is $\beta > 0$ so that the following holds. Suppose μ is a Radon measure in \mathbb{R}^{n+1} so that*

- (1) $\text{Tan}(\mu, \xi) \cap \mathcal{F} \neq \emptyset$ and
- (2) $\text{Tan}(\mu, \xi) \setminus \mathcal{F} \neq \emptyset$.

Then there is $r_0 > 0$ so that for any $\varepsilon > 0$ sufficiently small, the conclusion of Lemma 2.11 holds.

Proof. Let $v \in \text{Tan}(\mu, \xi) \setminus \mathcal{F}$. By Lemma 2.3, there exists $r_0 > 0$ so that $F_{r_0}(v) > 0$ and $d_{r_0}(v, \mathcal{F}) > 0$. Let $c_j > 0$ and $r_j \downarrow 0$ be so that $c_j T_{\xi, r_j}[\mu] \rightarrow v$. Then, for j large enough, $d_{r_0}(T_{\xi, r_j}[\mu], \mathcal{F}) > \frac{1}{2} d_{r_0}(v, \mathcal{F}) > 0$. The corollary now follows from Lemma 2.11 with $\varepsilon_0 = \frac{1}{2} d_{r_0}(v, \mathcal{F})$. \square

Proof of Lemma 2.10. If $\text{Tan}(\eta, x) \setminus \mathcal{F} \neq \emptyset$, then, by Corollary 2.12, we may find $\mu \in \text{Tan}(\eta, x) \setminus \mathcal{F}$ and $\varepsilon, r_0 > 0$ so that $d_{r_0}(\mu, \mathcal{F}) = \varepsilon$ and $d_r(\mu, \mathcal{F}) \leq \varepsilon$ for all $r > r_0$. By assumption, this implies $\mu \in \mathcal{F}$, which is a contradiction. Thus, $\text{Tan}(\eta, x) \subset \mathcal{F}$. \square

3. Elliptic measures

3A. Uniformly elliptic operators in divergence form. Let A be a real matrix with measurable coefficients that satisfies (1-1) and (1-2). We consider the second-order elliptic operator $L = -\operatorname{div} A \nabla$ and we say that a function $u \in W_{\operatorname{loc}}^{1,2}(\Omega)$ is a *weak solution* of the equation $Lu = 0$ in Ω (or just *L-harmonic*) if

$$\int A \nabla u \cdot \nabla \varphi = 0 \quad \text{for all } \varphi \in C_0^\infty(\Omega). \quad (3-1)$$

We also say that $u \in W_{\operatorname{loc}}^{1,2}(\Omega)$ is a *supersolution* (resp. *subsolution*) for L in Ω or just *L-superharmonic* (resp. *L-subharmonic*) if $\int A \nabla u \nabla \varphi \geq 0$ (resp. $\int A \nabla u \nabla \varphi \leq 0$) for all nonnegative $\varphi \in C_0^\infty(\Omega)$.

In this section, we assume $n \geq 2$.

3B. Regularity of the domain and Dirichlet problem. We say that a point $x_0 \in \partial\Omega$ is *Sobolev L-regular* if, for each function $\varphi \in W^{1,2}(\Omega) \cap C(\bar{\Omega})$, the L -harmonic function h in Ω with $h - \varphi \in W_0^{1,2}(\Omega)$ satisfies

$$\lim_{x \rightarrow x_0} h(x) = \varphi(x_0).$$

Theorem 3.1 [Heinonen et al. 1993, Theorem 6.27]. *If for $x_0 \in \partial\Omega$ it holds that*

$$\int_0^1 \frac{\operatorname{cap}(B(x_0, r) \cap \Omega^c, B(x_0, 2r))}{\operatorname{cap}(B(x_0, r), B(x_0, 2r))} \frac{dr}{r} = +\infty,$$

then x_0 is Sobolev L-regular. Here $\operatorname{cap}(\cdot, \cdot)$ stands for the variational 2-capacity of the condenser (\cdot, \cdot) (see, e.g., [Heinonen et al. 1993, p. 27]).

We say that a point $x_0 \in \partial\Omega$ is *Wiener regular* if, for each function $f \in C(\partial\Omega; \mathbb{R})$, the L -harmonic function H_f constructed by the Perron's method satisfies

$$\lim_{x \rightarrow x_0} H_f(x) = f(x_0).$$

See [Heinonen et al. 1993, Chapter 9].

Lemma 3.2 [Heinonen et al. 1993, Theorem 9.20]. *Suppose that $x_0 \in \partial\Omega$. If x_0 is Sobolev L-regular then it is also Wiener regular.*

The aforementioned result from [Heinonen et al. 1993] is only stated for Ω bounded but in fact it holds for unbounded domains, since the only part of the proof that requires the domain to be bounded is the existence of a unique solution of the Dirichlet problem with Sobolev Dirichlet data in bounded domains. This is true though in the unbounded case as well. See, e.g., on p. 11 in [Azzam et al. 2016a] where this is shown. Moreover, ∞ is also a Wiener regular point for each unbounded $\Omega \subset \mathbb{R}^{n+1}$, if $n \geq 2$; see, e.g., Theorem 9.22 in [Heinonen et al. 1993].

We say that Ω is *Sobolev L-regular* (resp. *Wiener regular*) if all the points in $\partial\Omega$ are Sobolev L -regular (resp. Wiener regular).

Definition 3.3. A domain $\Omega \subset \mathbb{R}^{n+1}$ is called *regular* if every point of $\partial\Omega$ is regular (i.e., if the classical Dirichlet problem is solvable in Ω for the elliptic operator \mathcal{L}), where $\partial\Omega$ denotes the boundary of Ω . For $K \subset \partial\Omega$, we say that Ω has the *capacity density condition (CDC)* if, for all $x \in \partial\Omega$ and $0 < r < \text{diam } \partial\Omega$,

$$\text{cap}(B(x, r) \cap \Omega^c, B(x, 2r)) \gtrsim r^{n-1}.$$

Note that if $n \geq 2$, by Wiener's criterion, domains satisfying the CDC are both Wiener regular and L -Sobolev regular.

Let $\Omega \subset \mathbb{R}^{n+1}$ be Wiener regular and $x \in \Omega$. If $f \in C(\partial\Omega)$, then the map $f \mapsto \bar{H}_f(x)$ is a bounded linear functional on $C(\partial\Omega)$. Therefore, by the Riesz representation theorem and the maximum principle, there exists a probability measure ω^x on $\partial\Omega$ (associated to L and the point $x \in \Omega$) defined on Borel subsets of $\partial\Omega$ so that

$$\bar{H}_f(x) = \int_{\partial\Omega} f d\omega^x \quad \text{for all } x \in \Omega.$$

We call ω^x the *elliptic measure* or *L -harmonic measure* associated to L and x .

3C. Green's function and PDE estimates.

Lemma 3.4. Let $\Omega \subset \mathbb{R}^{n+1}$, $n \geq 2$, be an open, connected set so that $\partial\Omega$ is Sobolev L -regular. There exists a Green's function $G : \Omega \times \Omega \setminus \{(x, y) : x = y\} \rightarrow \mathbb{R}$ associated with L which satisfies the following. For $0 < a < 1$, there are positive constants C and c depending on a, n and Λ such that for all $x, y \in \Omega$ with $x \neq y$, it holds that

$$\begin{aligned} 0 &\leq G(x, y) \leq C|x - y|^{1-n}, \\ G(x, y) &\geq c|x - y|^{1-n} \quad \text{if } |x - y| \leq a\delta_\Omega(x), \\ G(x, \cdot) &\in C(\bar{\Omega} \setminus \{x\}) \cap W_{\text{loc}}^{1,2}(\Omega \setminus \{x\}) \quad \text{and} \quad G(x, \cdot)|_{\partial\Omega} \equiv 0, \\ G(x, y) &= G^T(y, x), \end{aligned}$$

where G^T is the Green's function associated with the operator L_{A^T} , and for every $\varphi \in C_c^\infty(\mathbb{R}^{n+1})$

$$\int_{\partial\Omega} \varphi d\omega^x - \varphi(x) = - \int_{\Omega} A^T(y) \nabla_y G(x, y) \cdot \nabla \varphi(y) dy \quad \text{for a.e. } x \in \Omega. \quad (3-2)$$

In the statement of (3-2), one should understand that the integral on right-hand side is absolutely convergent for a.e. $x \in \Omega$ and a proof of it can be found in Lemma 2.6 in [Azzam et al. 2016a]. The rest were proved in [Grüter and Widman 1982; Hofmann and Kim 2007].

The lemma below is frequently called Bourgain's lemma, as he proved a similar estimate for harmonic measure in [Bourgain 1987].

Lemma 3.5 [Heinonen et al. 1993, Lemma 11.21]. Let $\Omega \subset \mathbb{R}^{n+1}$ be any domain satisfying the CDC condition, $x_0 \in \partial\Omega$, and $r > 0$ so that $\Omega \setminus B(x_0, 2r) \neq \emptyset$. Then

$$\omega_{\Omega}^{L,x}(B(x_0, 2r)) \geq c > 0 \quad \text{for all } x \in \Omega \cap B(x_0, r), \quad (3-3)$$

where c depends on d and the constant in the CDC.

Lemma 3.6. For $\Omega \subset \mathbb{R}^{n+1}$, $n \geq 2$, and the assumptions of [Lemma 3.4](#), if B is centered on $\partial\Omega$, then

$$G(x, y)r_B^{n-1} \inf_{z \in 2B} \omega^{L, z}(4B) \lesssim \omega^{L, y}(4B) \quad \text{for } x \in B \cap \Omega \text{ and } y \in \Omega \setminus 2B. \quad (3-4)$$

In particular, for a CDC domain, we have

$$G(x, y)r_B^{n-1} \lesssim \omega^{L, y}(4B) \quad \text{for } x \in B \cap \Omega \text{ and } y \in \Omega \setminus 2B.$$

Proof. This was originally shown for harmonic measure in [\[Azzam et al. 2016b\]](#), but we cover the details here.

By Bourgain's estimate, $\omega^{L, y}(4B) \gtrsim 1$ for $y \in 2B \cap \Omega$, and so for $y \in \Omega \setminus 2B$ and $x \in B \cap \Omega$

$$\inf_{z \in 2B} \omega^{L, z}(4B)G(x, y)r_B^{n-1} \lesssim \frac{\inf_{z \in 2B} \omega^{L, z}(4B)}{|x - y|^{n-1}}r_B^{n-1} \lesssim \inf_{z \in 2B} \omega^{L, z}(4B)$$

and since $G(x, \cdot)$ vanishes on $\partial\Omega$, we thus have that, for some constant $C > 0$,

$$\limsup_{y \rightarrow \xi} C\omega^{L, y}(4B) - \inf_{z \in 2B} \omega^{L, z}(4B)G(x, y)r_B^{n-1} \geq 0 \quad \text{for all } \xi \in \partial(\Omega \setminus 2B)$$

and so (3-4) follows from the maximum principle [\[Heinonen et al. 1993, Theorem 11.9\]](#). \square

By an iteration argument using [Lemma 3.5](#), one can obtain the following lemma.

Lemma 3.7. Let $\Omega \subsetneq \mathbb{R}^{n+1}$ be open with the CDC. Let $x \in \partial\Omega$ and $0 < r < \text{diam } \Omega$. Let u be a nonnegative L -harmonic function in $B(x, 4r) \cap \Omega$ and continuous in $B(x, 4r) \cap \bar{\Omega}$ so that $u \equiv 0$ in $\partial\Omega \cap B(x, 4r)$. Then extending u by 0 in $B(x, 4r) \setminus \bar{\Omega}$, there exists a constant $\alpha > 0$ such that

$$u(y) \leq C \left(\frac{\delta_\Omega(y)}{r} \right)^\alpha \sup_{B(x, 2r)} u \quad \text{for all } y \in B(x, r), \quad (3-5)$$

where C and α depend on n , Λ and the CDC constant, and $\delta_\Omega(y) = \text{dist}(y, \Omega^c)$. In particular, u is α -Hölder continuous in $B(x, r)$.

The following lemma is standard but we provide a proof for the sake of completeness.

Lemma 3.8. Let $\Omega \subset \mathbb{R}^{n+1}$ be an open set, and assume that A is an elliptic matrix and $\Phi : \mathbb{R}^{n+1} \rightarrow \mathbb{R}^{n+1}$ is a bi-Lipschitz map. Set

$$\tilde{A} := |\det D_\Phi| D_{\Phi^{-1}}(A \circ \Phi) D_{\Phi^{-1}}^T.$$

Then u is a weak solution of $L_A u = 0$ in $\Phi(\Omega)$ if and only if $\tilde{u} = u \circ \Phi$ is a weak solution of $L_{\tilde{A}} \tilde{u} = 0$ in Ω .

Proof. Let $\varphi \in C_c^\infty(\mathbb{R}^{n+1})$ and $\varphi = \psi \circ \Phi$. Then by change of variables and the chain rule

$$\begin{aligned} \int_{\Phi(\Omega)} A \nabla u \cdot \nabla \psi &= \int_{\Omega} (A \circ \Phi) \nabla u \circ \Phi \cdot \nabla \psi \circ \Phi |\det D_\Phi| \\ &= \int_{\Omega} (A \circ \Phi) D_{\Phi^{-1}}^T \nabla(u \circ \Phi) \cdot D_{\Phi^{-1}}^T \nabla(\psi \circ \Phi) |\det D_\Phi| \\ &= \int_{\Omega} |\det D_\Phi| D_{\Phi^{-1}}(A \circ \Phi) D_{\Phi^{-1}}^T \nabla(u \circ \Phi) \cdot \nabla(\psi \circ \Phi) = \int_{\Omega} \tilde{A} \nabla \tilde{u} \cdot \nabla \varphi. \end{aligned}$$

The lemma readily follows. \square

We will usually apply the above lemma when $\Phi(x) = Sx$ for some matrix S , in which case

$$\tilde{A} = (\det S)S^{-1}(A \circ S)(S^{-1})^T. \quad (3-6)$$

Lemma 3.9. *With the same assumptions as [Lemma 3.8](#), and assuming Ω is a Wiener regular domain, we have that for any set $E \subset \Phi(\partial\Omega) = \partial\Phi(\Omega)$ and $x \in \Omega$*

$$\omega_{\Phi(\Omega)}^{L_A, \Phi(x)}(E) = \omega_{\Omega}^{L_{\tilde{A}}, x}(\Phi^{-1}(E)). \quad (3-7)$$

Proof. Let $\varphi \in C_c^\infty(\mathbb{R}^{n+1})$. Since the function

$$v(x) = \int \varphi d\omega_{\Phi(\Omega)}^{L, x}$$

is L_A -harmonic for $x \in \Phi(\Omega)$, by the previous lemma we know that the function

$$\tilde{v}(x) = \int \varphi d\omega_{\Phi(\Omega)}^{L, \Phi(x)}$$

is $L_{\tilde{A}}$ -harmonic for $x \in \Omega$. If $\xi \in \partial\Omega$, then as $x \rightarrow \xi$ in Ω , $\Phi(x) \rightarrow \Phi(\xi)$ in $\Phi(\Omega)$, and so

$$\tilde{v}(x) = \int \varphi d\omega_{\Phi(\Omega)}^{L, \Phi(x)} \rightarrow \varphi(\Phi(\xi)).$$

Thus, \tilde{v} is the $L_{\tilde{A}}$ -harmonic extension of $(\varphi \circ \Phi)|_{\partial\Omega}$ to Ω , and so

$$\int_{\partial\Phi(\Omega)} \varphi d\omega_{\Phi(\Omega)}^{L_A, \Phi(x)} = \int_{\partial\Omega} \varphi \circ \Phi d\omega_{\Omega}^{L_{\tilde{A}}, x} \quad \text{for all } x \in \Omega.$$

Since this holds for all such φ , we get that for any set $E \subset \partial\Phi(\Omega) = \Phi(\partial\Omega)$,

$$\omega_{\Phi(\Omega)}^{L_A, \Phi(x)}(E) = \omega_{\Omega}^{L_{\tilde{A}}, x}(\Phi^{-1}(E)),$$

which gives the lemma. \square

The following lemma will help us relate measures generated by elliptic polynomials to just measures generated by harmonic polynomials. In particular, if A is an elliptic matrix with constant and real coefficients, by the change of variables described below (which is just a linear transformation), if h is a harmonic polynomial solution in an open set Ω and $S = \sqrt{A_s}$ (where A_s is the symmetric part of A), then $\tilde{h} = h \circ S^{-1}$ is a polynomial solution of $-\operatorname{div} A \nabla u = 0$ in $S(\Omega)$. So, there is a bijection between the set of harmonic polynomials and the set of polynomial solutions of $-\operatorname{div} A \nabla u = 0$ in $S(\Omega)$ (for a fixed constant elliptic matrix A). Recall also that p is a harmonic polynomial in an open set if and only if it is a harmonic polynomial in \mathbb{R}^{n+1} . So, if A is as above, there is an abundance of nontrivial polynomial solutions of $-\operatorname{div} A \nabla u = 0$ in any open subset of \mathbb{R}^{n+1} (including \mathbb{R}^{n+1} itself). In fact, Theorem 2 in [\[Abramov and Petkovšek 2012\]](#) states that for such L_A , for any $k \in \mathbb{N}$, there exists a polynomial solution of $L_A h = 0$ of degree k .

Lemma 3.10. *Let A be an elliptic constant matrix, $A_s = \frac{1}{2}(A + A^T)$, and $S = \sqrt{A_s}$. Let $h \in H_A$ and $\tilde{h} = h \circ S$. Then $\tilde{A} = (\det S)I$, $\tilde{h} \in H$ and*

$$\omega_{\tilde{h}} = (\det S)^{-1} S^{-1} [\omega_h^A]. \quad (3-8)$$

Proof. Note that since L_A has constant coefficients, $L_{A_s} = L_A$ by the fact that for $u \in C^2$

$$\begin{aligned} L_A u &= \sum_{i,j} a_{ij} \partial_i \partial_j u = \frac{1}{2} \sum_{i,j} a_{ij} \partial_i \partial_j u + \frac{1}{2} \sum_{i,j} a_{ij} \partial_j \partial_i u \\ &= \sum_{i,j} \frac{(a_{ij} + a_{ji})}{2} \partial_i \partial_j u = L_{A_s} u. \end{aligned}$$

Thus, if h is an L_A -harmonic function, it is also an L_{A_s} -harmonic function. Moreover, for any $\psi \in C_c^\infty(\mathbb{R}^{n+1})$

$$\int \psi d\omega_h^{A_s} = \int_{\Omega_h} h L_{A_s}(\psi) = \int_{\Omega_h} h L_A(\psi) = \int \psi d\omega_h^A.$$

In fact, without loss of generality, we may assume that $A = A_s$.

Recall now that since A_s is a symmetric, positive definite and invertible matrix with constant real entries, then it has a unique real symmetric positive definite square root $S = \sqrt{A_s}$ which is also invertible. Hence, by [Lemma 3.8](#) and (3-6) with $A = A_s$, we have that $\tilde{A} = (\det S)I$ and \tilde{h} is $L_{(\det S)I}$ -harmonic, and thus just harmonic.

Let now $\varphi \in C_c^\infty(\mathbb{R}^{n+1})$ and $\psi \circ S = \varphi$. By Green's formula and the fact that S is also symmetric, we have

$$\begin{aligned} (\det S) \int \varphi d\omega_{\tilde{h}} &= (\det S) \int_{\Omega_{\tilde{h}}} \tilde{h} \Delta \varphi = -(\det S) \int_{\Omega_{\tilde{h}}} \nabla \tilde{h} \cdot \nabla \varphi \\ &= -(\det S) \int_{\Omega_{\tilde{h}}} S^T \nabla h \circ S \cdot S^T \nabla \psi \circ S \\ &= - \int_{S^{-1}(\Omega_h)} S S^T \nabla h \circ S \cdot \nabla \psi \circ S \\ &= - \int_{\Omega_h} A_s \nabla h \cdot \nabla \psi = \int_{\Omega_h} h L_{A_s}(\psi) \\ &= \int_{\Omega_h} h L_A(\psi) = \int \psi d\omega_h^A = \int \varphi dS^{-1}[\omega_h^A]. \quad \square \end{aligned}$$

Let us recall some simple facts from linear algebra which help us understand how the geometry of Ω is affected by the linear transformation above. Note that S is orthogonally diagonalizable since it is symmetric, which means that it represents a linear transformation with scaling in mutually perpendicular directions. Hence S^{-1} is a special bi-Lipschitz change of variables that takes balls to ellipsoids, where eigenvectors determine directions of semiaxes, eigenvalues determine lengths of semiaxes and its maximum eccentricity is given by $\sqrt{(\lambda_{\max}/\lambda_{\min})}$ (where λ_{\max} are λ_{\min} are the maximal and minimal eigenvalues of S^{-1}), which is in turn bounded below by $\sqrt{\Lambda}^{-1}$ and above by $\sqrt{\Lambda}$. In particular, $S^{-1}(\partial\Omega) = \partial(S^{-1}(\Omega))$, $\Lambda^{-1/2} \leq \|S^{-1}\| \leq \Lambda^{1/2}$; i.e., S^{-1} distorts distances by at most a constant depending on ellipticity.

3D. The main blow-up lemma. We now introduce the main tool of this paper, which is a variant of previous blow-up arguments, first introduced by Kenig and Toro [\[2006\]](#) for NTA domains, then extended to CDC domains in [\[Azzam et al. 2017b\]](#). Both these cases apply to harmonic measure but can be extended to elliptic measures with a VMO condition on the coefficients.

Lemma 3.11. *Let $\Omega^+ \subset \mathbb{R}^{n+1}$ be a CDC domain, $K \subset \partial\Omega^+$ a compact set, $\xi_j \in K$ a sequence of points, and $L = -\operatorname{div} A \nabla$ a uniformly elliptic operator in Ω^+ such that*

$$\lim_{r \rightarrow 0} \sup_{\xi \in K} \frac{1}{r^{n+1}} \inf_{C \in \mathcal{C}} \int_{B(\xi, r) \cap \Omega^+} |A(x) - C| dx = 0. \quad (3-9)$$

Let ω^+ be the elliptic measure for Ω^+ and $c_j \geq 0$, and $r_j \rightarrow 0$ such that $\omega_j^+ = c_j T_{\xi_j, r_j}[\omega^+] \rightarrow \omega_\infty^+$ for some nonzero measure ω_∞^+ . Let $\Omega_j^+ = T_{\xi_j, r_j}(\Omega^+)$. Then there is a subsequence and a closed set $\Sigma \subset \mathbb{R}^{n+1}$ such that:

- (a) *For all $R > 0$ sufficiently large, $B(0, R) \cap \partial\Omega_j^+ \neq \emptyset$ and $\partial\Omega_j^+ \cap \overline{B(0, R)} \rightarrow \Sigma \cap \overline{B(0, R)}$ in the Hausdorff metric.*
- (b) *$\Sigma^c = \Omega_\infty^+ \cup \Omega_\infty^-$, where Ω_∞^+ is a nonempty open set and Ω_∞^- is also open but possibly empty. Further, they satisfy that for any ball B with $\bar{B} \subset \Omega_\infty^\pm$, a neighborhood of \bar{B} is contained in Ω_j^\pm for all j large enough.*
- (c) *$\operatorname{supp} \omega_\infty^+ \subset \Sigma$.*
- (d) *Let $u^+(x) = G_{\Omega^+}(x, x^+)$ on Ω^+ and $u^+(x) = 0$ on $(\Omega^+)^c$. Set*

$$u_j^+(x) = c_j u^+(x r_j + \xi_j) r_j^{n-1}.$$

Then u_j^+ converges locally uniformly in \mathbb{R}^{n+1} and in $W_{\operatorname{loc}}^{1,2}(\mathbb{R}^{n+1})$ to a nonzero function u_∞^+ which is continuous in \mathbb{R}^{n+1} , vanishes in $(\Omega_\infty^+)^c$, and satisfies

$$u_\infty^+(y) \lesssim \omega_\infty^+(\overline{B(x, 4r)}) r^{1-n} \quad (3-10)$$

for $x \in \Sigma$, $r > 0$, and $y \in B(x, r) \cap \Omega_\infty^+$. Moreover, there is A_0^+ a constant elliptic matrix so that if $L_0^+ = -\operatorname{div} A_0^+ \nabla$, then

$$\int \varphi d\omega_\infty^+ = \int_{\mathbb{R}^{n+1}} u_\infty^+ L_0^+ \varphi \quad \text{for any } \varphi \in C_c^\infty(\mathbb{R}^{n+1}). \quad (3-11)$$

Suppose now that $\Omega^- = \mathbb{R}^{n+1} \setminus \bar{\Omega}^+$, so that $\partial\Omega^+ = \partial\Omega^-$ and Ω^- is also connected and has the CDC. Define analogously ω_j^- , u^- , u_j^- , and u_∞^- . Assume that A is uniformly elliptic in $\Omega^+ \cup \Omega^-$, (3-9) holds for $\Omega^+ \cup \Omega^-$ in place of Ω^+ and ω_j^- converges weakly to $\omega_\infty^- = c\omega_\infty^+$ for some number $c \in (0, \infty)$. Then $\Omega_\infty^- \neq \emptyset$ and for a suitable subsequence, (d) holds for u_j^- , u_∞^- , and Ω_∞^- . Furthermore, if we set $u_\infty = u_\infty^+ - c^{-1}u_\infty^-$, then:

- (e) *u_∞ extends to a continuous function on \mathbb{R}^{n+1} which satisfies $L_0 u_\infty = 0$ in \mathbb{R}^{n+1} .*
- (f) *$\Sigma = \{u_\infty = 0\}$, with $u_\infty > 0$ on Ω_∞^+ and $u_\infty < 0$ on Ω_∞^- . Further, Σ is a real analytic variety of dimension n .*
- (g) *$d\omega_\infty^+ = -(\partial u_\infty / \partial \nu_{A_0}) d\sigma_{\partial\Omega_\infty^+}$, where σ_S stands for the surface measure on a surface S and $\partial / \partial \nu_{A_0} = \nu \cdot A_0 \nabla$ is the outward conormal derivative.*

Proof. The proof of this lemma can be found in [Azzam et al. 2017b] for harmonic measure for the case that $K = \{\xi\}$ (i.e., so that (1-4) holds). The proof for general K is essentially the same in this setting with

minor changes. Here we shall only record the required modifications (some of which are quite substantial) for the $K = \{\xi\}$ case in order for the same proof to work for any elliptic measure as well. In this case, $\xi_j = \xi$ for all j . We set

$$A_j(x) := A(r_j x + \xi), \quad u_j^\pm(x) := c_j r_j^{n-1} u^\pm(r_j x + \xi), \quad \varphi_j(x) := \varphi\left(\frac{x - \xi}{r_j}\right).$$

Without loss of generality we can only work with u^+ since the results for u^- can be proved analogously.

Notice now that for j large enough, the pole x^+ is not in $\text{supp}(\varphi_j)$. In fact, for any ball B centered at the boundary of Ω_j , we can find $j_0 \in \mathbb{N}$ such that for all $j \geq j_0$, we have $x^+ \notin T_{\xi, r_j}(B)$. Moreover, for $x \in B \cap \Omega_j$ and j large enough,

$$u_j^+(x) = c_j r_j^{n-1} u^+(r_j x + \xi) \stackrel{(3-4)}{\lesssim} c_j r_j^{n-1} (r_j r_B)^{1-n} \omega^+(4r_j B + \xi) = r_B^{1-n} \omega_j^+(4B). \quad (3-12)$$

Proof of (b): We only need to prove the existence of $B \subset \Omega_j^+$ for large $j \in \mathbb{N}$. Suppose there is no such ball. Let φ be any continuous compactly supported nonnegative function for which $\int \varphi d\omega_\infty^+ \neq 0$, and let $M > 0$ be so that $\text{supp } \varphi \subset B(0, M)$. Thus, there must be $x_0 \in B(0, M) \cap \text{supp } \omega_\infty^+$. We set

$$\delta_j := \sup\{\text{dist}(x, (\Omega_j^+)^c) : x \in B(0, 2M)\},$$

which goes to zero by assumption. For $x \in B(0, 2M)$ and $j \in \mathbb{N}$, let $\zeta_j(x) \in (\Omega_j^+)^c$ be closest to x so that $|x - \zeta_j(x)| \leq \delta_j \leq 2M$ (the second inequality holds because $0 \in \partial\Omega_j^+$). It also holds that for all $x \in B(0, 2M)$, we have $|x - x_0| \leq |x| + |x_0| < 3M$.

Notice now that for any j big enough, u_j^+ is a solution in $B(0, 2M) \cap \Omega_j^+$ and a subsolution in $B(0, 2M)$. Moreover, if $x \in \Omega_j^+$, then $\zeta_j(x) \in \partial\Omega_j^+$. Thus, for j large, by Cauchy–Schwarz, Caccioppoli’s inequality in $B(0, M)$ (which also holds for subsolutions), and the fact that u_j^+ and φ are supported in Ω_j^+ and $B(0, M)$ respectively,

$$\begin{aligned} 0 &< \int \varphi d\omega_j^+ = \int_{\Omega_j^+} A_j \nabla u_j^+ \cdot \nabla \varphi \lesssim_{\lambda, \Lambda, n, M} \|\nabla \varphi\|_\infty \left(\int_{B(0, 2M)} |u_j^+|^2 \right)^{1/2} \\ &\stackrel{(3-5)}{\lesssim} \left(\int_{\Omega_j^+ \cap B(0, 2M)} \left(\sup_{B(\zeta_j(x), 2M)} u_j^+ \right)^2 \left(\frac{x - \zeta_j(x)}{2M} \right)^{2\alpha} dx \right)^{1/2} \\ &\stackrel{(3-12)}{\lesssim} \left(\int_{\Omega_j^+ \cap B(0, 2M)} [\omega_j^+(B(\zeta_j(x), 8M))(2M)^{1-n}]^2 dx \right)^{1/2} \left(\frac{\delta_j}{2M} \right)^\alpha \\ &\lesssim (2M)^{(n+1)/2} \omega_j^+(B(x_0, 13M))(2M)^{1-n} \left(\frac{\delta_j}{2M} \right)^\alpha, \end{aligned}$$

and thus

$$\begin{aligned} 0 &< \int \varphi d\omega_\infty^+ \lesssim_{\lambda, \Lambda, n, M, \varphi} \left(\limsup_{j \rightarrow \infty} \omega_j^+(B(x_0, 13M)) \right) \lim_j \delta_j^\alpha \\ &\leq \omega_\infty^+(\overline{B(x_0, 13M)}) \cdot 0 = 0, \end{aligned}$$

which is a contradiction. Thus, there is $B \subset \Omega_j$ for all large j (after passing to a subsequence).

Proof of (d): Arguing as in [Azzam et al. 2017b], there exists u_∞^+ which is continuous in \mathbb{R}^{n+1} and vanishes on $(\Omega_\infty^+)^c$ such that (after passing to a subsequence) $u_j^+ \rightarrow u_\infty^+$ uniformly on compact sets of \mathbb{R}^{n+1} . Moreover, it is not hard to see that $u_j^+ \in W^{1,2}(B)$ for large j . Indeed, by (3-12), it is clear that

$$\|u_j^+\|_{L^2(B)}^2 \lesssim r_B^{3-n} [\omega_j^+(4B)]^2, \quad (3-13)$$

while by Caccioppoli's inequality and (3-12),

$$\int_B |\nabla u_j^+|^2 \lesssim r_B^{-2} \int_B |u_j^+|^2 \lesssim r_B^{-2} [r_B^{1-n} \omega_j^+(4B)]^2 r_B^{n+1} = r_B^{1-n} [\omega_j^+(4B)]^2. \quad (3-14)$$

In view of (3-13) and (3-14) we have

$$\begin{aligned} \limsup_{j \rightarrow \infty} \|u_j^+\|_{W^{1,2}(B)} &\lesssim r_B^{(1-n)/2} (1 + r_B) \limsup_{j \rightarrow \infty} \omega_j^+(4B) \\ &\leq r_B^{(1-n)/2} (1 + r_B) \omega_\infty^+(\overline{4B}) < \infty. \end{aligned}$$

Therefore, by [Heinonen et al. 1993, Theorem 1.32], $u_\infty^+ \in W_{\text{loc}}^{1,2}(\mathbb{R}^{n+1})$ and there exists a further subsequence of u_j^+ that converges weakly to u_∞^+ in $W_{\text{loc}}^{1,2}(\mathbb{R}^{n+1})$.

Notice that

$$-\int_{\Omega_j^+} A_j \nabla u_j^+ \cdot \nabla \varphi = \int \varphi d\omega_j^+.$$

Indeed, by a change of variables, and letting $\varphi_j = \varphi \circ T_{\xi, r_j}$ and $\varphi_j = \varphi \circ T_{\xi, r_j}$,

$$\begin{aligned} \int \varphi d\omega_j^+ &= c_j \int \varphi_j d\omega^+ = \int_{\Omega^+} A \nabla u^+ \cdot \nabla \varphi_j \\ &= c_j r_j^n \int_{\Omega_j^+} A(r_j x + \xi) \nabla u^+(r_j x + \xi) \cdot \nabla \varphi(x) dx \\ &= \int_{\Omega_j^+} A_j \nabla u_j^+ \cdot \nabla \varphi. \end{aligned}$$

Let $C_{j,k}$ be a constant elliptic matrix so that

$$\lim_j (kr_j)^{-1-n} \int_{B(\xi, kr_j) \cap \Omega^+} |A - C_{j,k}| = 0.$$

By a diagonalization argument and compactness, we may pass to a subsequence so that for each k , $C_{j,k}$ converges to a uniformly elliptic matrix C_k with constant coefficients. It is not hard to check that we must in fact have that $C_k = A_0^+$ for some fixed matrix A_0^+ (using the fact that $\inf \delta_j > 0$). Thus, we have

$$\lim_j (Mr_j)^{-1-n} \int_{B(\xi, Mr_j) \cap \Omega^+} |A - A_0^+| = 0 \quad \text{for all } M \geq 1. \quad (3-15)$$

To see the ellipticity of A_0^+ is pretty easy but we show the details for completeness. Note that since A is uniformly elliptic for a.e. $x \in \Omega^+$, for $\xi \in \mathbb{R}^{n+1}$

$$\Lambda^{-1} |\xi|^2 \leq A(x) \xi \cdot \xi = (A(x) - A_0^+) \xi \cdot \xi + A_0^+ \xi \cdot \xi.$$

Then, if we take averages over $B(\xi, Mr_j) \cap \Omega$, use the existence of corkscrew balls in Ω_j for large j proved in (b), and then take limits as $j \rightarrow \infty$, by (3-15) we have

$$\Lambda^{-1}|\xi|^2 \leq A_0^+ \xi \cdot \xi.$$

The upper bound follows by a similar argument and the proof is omitted.

We will now estimate the difference

$$\int_{\Omega_j^+} A_j \nabla u_j^+ \cdot \nabla \varphi - \int_{\Omega_\infty^+} A_0^+ \nabla u_\infty^+ \cdot \nabla \varphi \quad (3-16)$$

for sufficiently large j .

To this end, let $\text{supp}(\varphi) \subset B(0, M)$. Note that

$$|(3-16)| \leq \left| \int_{\Omega_j^+} (A(r_j x + \xi) - A_0^+) \nabla u_j^+ \cdot \nabla \varphi \right| + \left| \int_{B(0, M)} (\nabla u_j^+ \mathbb{1}_{\Omega_j} - \nabla u_\infty^+ \mathbb{1}_{\Omega_\infty}) \cdot A_0^{+,T} \nabla \varphi \right| \leq I_1 + I_2.$$

Note that $u_j^+, u_\infty^+ \in W^{1,2}(\mathbb{R}^{n+1})$, $u_j^+ > 0$ only in Ω_j^+ , and $u_\infty^+ > 0$ only in Ω_∞^+ . Since the extension of the gradient of a function $f \in W_0^{1,2}(\Omega)$ by zero to \mathbb{R}^{n+1} (where Ω is any domain) is the same as the gradient of the extension of f by zero,¹ we have that in $W^{1,2}(B(0, M))$

$$\nabla u_j^+ \mathbb{1}_{\Omega_j^+} = \nabla(u_j^+ \mathbb{1}_{\Omega_j^+}) = \nabla u_j^+ \rightharpoonup \nabla u_\infty^+ = \nabla(u_\infty^+ \mathbb{1}_{\Omega_\infty^+}) = \nabla u_\infty^+ \mathbb{1}_{\Omega_\infty^+},$$

so we have $I_2 \rightarrow 0$. On the other hand, since A and $A_0^+ \in L^\infty(\Omega)$,

$$\begin{aligned} I_1 &\leq \|\nabla u_j^+\|_{L^2(B(0, M))} \|\nabla \varphi\|_\infty \left(\int_{B(0, M) \cap \Omega_j^+} |A(r_j x + \xi) - A_0^+|^2 dx \right)^{1/2} \\ &\stackrel{(3-14)}{\lesssim_\Lambda} M^{(1-n)/2} \omega_\infty^+(\overline{B(0, 4M)}) \left(\frac{1}{r_j^{1+n}} \int_{B(0, Mr_j) \cap \Omega_j^+} |A(x) - A_0^+|^2 dx \right)^{1/2} \\ &\stackrel{(3-15)}{\rightarrow} 0. \end{aligned}$$

Thus, combining the above estimates and taking $j \rightarrow \infty$, we infer that

$$- \int_{\Omega_\infty^+} A_0^+ \nabla u_\infty^+ \cdot \nabla \varphi = \int \varphi d\omega_\infty^+.$$

In particular, u_∞^+ is a continuous weak solution of

$$L_0^+ w = -\text{div } A_0^+ \nabla w = 0 \quad \text{in } \Omega_\infty^+.$$

Since L_0^+ is a second-order elliptic operator with constant coefficients, u_∞^+ is real analytic in Ω_∞^+ . Thus, by the definition of u_∞^+ and since the gradient of its extension by zero is the extension by zero of the gradient, we have

$$\int_{\Omega_\infty^+} A_0^+ \nabla u_\infty^+ \cdot \nabla \varphi = \int_{\mathbb{R}^{n+1}} A_0^+ \nabla u_\infty^+ \cdot \nabla \varphi.$$

¹ See Proposition 9.18 in [Brezis 2011]. It is stated for C^1 -domains, but the direction we need holds for general Ω .

We now use the divergence theorem along with the fact that $\text{supp}(\nabla\varphi) \subset B(0, M)$ and obtain (writing $L_0^{+,T} = L_{A_0^{+,T}}$)

$$\int \varphi d\omega_\infty^+ = - \int_{\mathbb{R}^{n+1}} \text{div}[u_\infty^+ A_0^{+,T} \nabla \varphi] + \int_{\mathbb{R}^{n+1}} u_\infty^+ L_0^{+,T} \varphi = -0 + \int_{\mathbb{R}^{n+1}} u_\infty^+ L_0^{+,T} \varphi,$$

which finishes the proof of (d). The rest of the proof is almost identical since one only uses that u_∞ is real analytic in \mathbb{R}^{n+1} and Liouville's theorem for positive solutions of uniformly elliptic equations; see, e.g., [Heinonen et al. 1993, Corollary 6.11].

One may argue similarly in the case of u_j^- . Notice that in this case, we will obtain a constant-coefficient uniformly elliptic matrix A_0 such that

$$\lim_j (Mr_j)^{-1-n} \int_{B(\xi, Mr_j) \cap (\Omega^+ \cup \Omega^-)} |A - A_0| = 0 \quad \text{for all } M \geq 1. \quad \square$$

Now we prove a slightly weaker version of this result in the next two lemmas. Again, this is based on the details in the proof of [Azzam et al. 2016c, Lemma 5.3], but with some adjustments for elliptic measure.

Lemma 3.12. *Let $\Omega \subset \mathbb{R}^{n+1}$ be a domain. Let $\xi_j \in \partial\Omega$ and $L = -\text{div } A \nabla$ be a uniformly elliptic operator in Ω such that (1-3) holds with $K = \{\xi_j\}$ and, if $\omega = \omega_\Omega^{L_A, x_0}$ is its L_A -harmonic measure with pole at $x_0 \in \Omega$, there is $r_j \rightarrow 0$ and $c_j > 0$ so that*

$$\omega_j := c_j T_{\xi_j, r_j}[\omega] \rightarrow \omega_\infty, \quad (3-17)$$

$$\liminf_j \frac{|\Omega \cap B(\xi_j, r_j)|}{r_j^{n+1}} > 0, \quad (3-18)$$

$$\omega^z(B(\xi_j, 2r_j)) \gtrsim 1 \quad \text{for all } j \text{ and } z \in B(\xi_j, r_j) \cap \Omega. \quad (3-19)$$

Then there is a subsequence such that the following hold: If $u(x) = G_\Omega(x, x_0)$ on Ω and $u(x) = 0$ on Ω^c , and

$$u_j(x) = c_j u(xr_j + \xi_j) r_j^{n-1},$$

then u_j converges in $L_{\text{loc}}^2(\frac{1}{2}\mathbb{B})$ to a nonzero function u_∞ which is L_{A_0} -harmonic in $\{x : u_\infty > 0\} \cap (\frac{1}{2}\mathbb{B})$ for constant uniformly elliptic matrix A_0 and such that

$$\|u_\infty\|_{L^2(\mathbb{B}/2)} \lesssim \omega_\infty(\overline{B(0, 2)}), \quad (3-20)$$

and for any $\varphi \in C_c^\infty(\frac{1}{2}\mathbb{B})$

$$\int \varphi d\omega_\infty = \int_{\mathbb{R}^{n+1}} u_\infty L_{A_0} \varphi. \quad (3-21)$$

If $\xi = \xi_j$ and A is continuous at ξ , then A_0 is just the value of A at ξ .

Proof. Recall that we let $\mathbb{B} = B(0, 1)$. Again, to simplify notation, we'll just prove the case when $\xi_j = \xi \in \partial\Omega$.

By (3-19), without loss of generality, we can scale the c_j so that

$$\omega_\infty(\tfrac{1}{4}\mathbb{B}) = 1. \quad (3-22)$$

Let $\Omega_j = T_{\xi, r_j}(\Omega)$. By (3-19) and (3-4),

$$\omega(B(\xi, 2r_j)) \gtrsim r_j^{n-1} u(x) \quad \text{for all } x \in B(\xi, r_j) \cap \Omega_1, \quad (3-23)$$

and so,

$$\omega_j(2\mathbb{B}) \gtrsim u_j(x) \quad \text{for all } x \in \mathbb{B} \cap \Omega_1^j, \quad (3-24)$$

By Caccioppoli's inequality for L -subharmonic functions and the uniform boundedness of u in \mathbb{B} , we deduce that, for $i = 1, 2$,

$$\limsup_{j \rightarrow \infty} \|\nabla u_j\|_{L^2(\mathbb{B}/2)} \lesssim \limsup_{j \rightarrow \infty} \|u_j\|_{L^2(\mathbb{B})} \lesssim \limsup_{j \rightarrow \infty} \omega_j(2\mathbb{B}) \leq \omega_\infty(\overline{2\mathbb{B}}).$$

By the Rellich–Kondrachov theorem, the unit ball of the Sobolev space $W^{1,2}(\frac{1}{2}\mathbb{B})$ is relatively compact in $L^2(\frac{1}{2}\mathbb{B})$, and thus there exists a subsequence of the functions u_j which converges *strongly* in $L^2(\frac{1}{2}\mathbb{B})$ to another function $u_\infty \in L^2(\frac{1}{2}\mathbb{B})$. This and the above inequality imply (3-20).

By the same diagonalization argument as in the proof of the previous lemma (although using (3-18) instead of $\inf \delta_j > 0$ that we used in the previous lemma), we can pass to a subsequence so that, for some uniformly elliptic matrix A_0 with constant coefficients,

$$\lim_j (Mr_j)^{-1-n} \int_{B(\xi, Mr_j) \cap \Omega} |A(x) - A_0| = 0 \quad \text{for all } M \geq 1. \quad (3-25)$$

It easy to check that

$$\int \varphi d\omega_j = \int A_j \nabla u_j \cdot \nabla \varphi dx$$

for any C^∞ function φ compactly supported in $\frac{1}{2}\mathbb{B}$. Then passing to a limit, it follows that

$$\int \varphi d\omega_\infty = \int A_0 \nabla u_\infty \cdot \nabla \varphi dx, \quad \text{for any } \varphi \in C_c^\infty(\tfrac{1}{2}\mathbb{B}). \quad \square$$

Theorem 3.13. *Let $\Omega^\pm \subset \mathbb{R}^{n+1}$ be disjoint domains. Let $\xi_j \in \partial\Omega^+ \cap \partial\Omega^-$ and $L = -\operatorname{div} A \nabla$ be a uniformly elliptic operator in $\Omega^+ \cup \Omega^-$ such that (1-3) holds with $K = \{\xi_j\}$ with respect to $\Omega^+ \cup \Omega^-$. If $\omega^\pm = \omega_{\Omega^\pm}^{L_A, x^\pm}$ is the L_A -harmonic measure with pole at $x^\pm \in \Omega^\pm$, and if there is $r_j \rightarrow 0$ and $c_j > 0$ so that*

$$\begin{aligned} \omega_j^+ &:= c_j T_{\xi_j, r_j}[\omega^+] \rightarrow \omega_\infty, \\ \omega_j^- &:= c_j T_{\xi_j, r_j}[\omega^-] \rightarrow c\omega_\infty \end{aligned}$$

for some constant $c > 0$, then there is a subsequence such that the following hold. If $u^\pm(x) = G_{\Omega^\pm}(x, x^\pm)$ on Ω^\pm , $u(x) = 0$ on $(\Omega^\pm)^c$ and

$$u_j^\pm(x) = c_j u^\pm(xr_j + \xi_j) r_j^{n-1}, \quad (3-26)$$

then $u_j := u_j^+ - c^{-1}u_j^-$ converges in $L^2(\frac{1}{2}\mathbb{B})$ to a nonzero function u_∞ , which is L_{A_0} -harmonic in $\frac{1}{2}\mathbb{B}$ for some constant uniformly elliptic matrix A_0 , and moreover,

$$\tfrac{1}{2}\mathbb{B} \cap \operatorname{supp} \omega_\infty = \{u_\infty = 0\} \cap \tfrac{1}{2}\mathbb{B} \quad (3-27)$$

and (3-20) and (3-21) hold. If $\xi_j = \xi$ and A is continuous at ξ , then A_0 is just the value of A at ξ .

By applying this result to the sequences $c_j T_{\xi_j, ar_j}[\omega^\pm]$ for all $a > 0$, we see that u_∞ extends to an L_{A_0} -harmonic function on \mathbb{R}^{n+1} so that for $r > 0$

$$\|u_\infty\|_{L^2(B(0,r))} \lesssim r^{1-n} \omega_\infty(\overline{B(0,4r)}), \quad (3-28)$$

and for any $\varphi \in C_c^\infty(\mathbb{R}^{n+1})$

$$\int \varphi d\omega_\infty = \int_{\mathbb{R}^{n+1}} u_\infty L_{A_0} \varphi. \quad (3-29)$$

Proof. The proof is mostly the same as the proof of [Azzam et al. 2016c, Lemma 5.3], but we provided some of the details here to show the differences. Again, we assume $\xi_j = \xi$. Note that since Ω^+ and Ω^- are disjoint, we may assume without loss of generality that

$$|B(\xi, \tfrac{1}{8}r_j) \setminus \Omega^+| \geq \tfrac{1}{2} |B(\xi, \tfrac{1}{8}r_j)|$$

and so Bourgain's estimate implies

$$\omega_\infty^{+,z}(B(\xi, 2r_j)) \gtrsim 1 \quad \text{for all } z \in B(\xi, r_j).$$

Hence, the conclusions of Lemma 3.12 apply to $\omega = \omega^+$, $\Omega = \Omega^+$, and $u = u^+$. In particular, (3-24) in our scenario is

$$\omega_j^+(2\mathbb{B}) \gtrsim u_j^+(x) \quad \text{for all } x \in \mathbb{B} \cap \Omega_1^j. \quad (3-30)$$

Again, by rescaling, we can assume that $\omega_\infty(\frac{1}{4}\mathbb{B}) = 1$.

Observe now that for any nonnegative $\varphi \in C_c^\infty(\frac{1}{2}\mathbb{B})$ with $\varphi = 1$ in $\frac{1}{4}\mathbb{B}$, by Cauchy–Schwarz and Caccioppoli's inequality (since u_j^\pm is positive and L_{A_j} -harmonic in $\mathbb{B} \cap \Omega_j^\pm$ and zero in $\mathbb{B} \setminus \Omega_j^\pm$) we have

$$\begin{aligned} 1 &= \omega_\infty(\tfrac{1}{4}\mathbb{B}) \leq \int \varphi d\omega_\infty = \int A_0 \nabla u_\infty^+ \cdot \nabla \varphi dx \\ &= \lim_j \int_{\Omega_j^+} A_j \nabla u_j^+ \cdot \nabla \varphi dx \\ &\leq \|A\|_{L^\infty} \|\nabla \varphi\|_{L^\infty(\mathbb{B})} \lim_j \int_{\Omega_j^+ \cap \mathbb{B}/2} |\nabla u_j^+| \\ &\lesssim \|A\|_{L^\infty} \|\nabla \varphi\|_{L^\infty(\mathbb{B})} \lim_j \left(\int_{\Omega_j^+ \cap \mathbb{B}} |u_j^+|^2 \right)^{1/2} \\ &\lesssim \lim_j \left(\int_{\mathbb{B} \cap \Omega_j^+ \cap \{u_j^+ > t\}} |u_j^+|^2 dx + \int_{\mathbb{B} \cap \Omega_j^+ \cap \{u_j^+ \leq t\}} |u_j^+|^2 dx \right)^{1/2} \\ &\lesssim \liminf_j (|\{x \in \mathbb{B} \cap \Omega_j^+ : u_j^+ > t\}|^{1/2} \cdot \|u_j^+\|_{L^\infty(\mathbb{B} \cap \Omega_j^+)} + t) \\ &\stackrel{(3-30)}{\lesssim} \liminf_j (|\{x \in \mathbb{B} \cap \Omega_j^+ : u_j^+ > t\}|^{1/2} \omega_\infty(\overline{2\mathbb{B}}) + t), \end{aligned}$$

and so, for t small enough,

$$|\mathbb{B} \cap \Omega_j^+| \geq |\{x \in \mathbb{B} \cap \Omega_j^+ : u_j^+(x) > t\}| \gtrsim \omega_\infty(\overline{2\mathbb{B}})^{-2}.$$

In particular,

$$|B(\xi, r_j) \setminus \Omega^-| \geq |B(\xi, r_j) \cap \Omega^+| \gtrsim r_j^{n+1} \omega_\infty(\overline{2\mathbb{B}})^{-2}. \quad (3-31)$$

Thus, by the same arguments as earlier in proving (3-24), we have that for j large

$$\omega_j^-(B(\xi, 2r_j)) \gtrsim u_j^-(x) \omega_\infty(\overline{2\mathbb{B}})^{-2} \quad \text{for all } x \in B(\xi, r_j) \cap \Omega^-. \quad (3-32)$$

Thus, we can apply Lemma 3.12 and can pass to a subsequence so that u_j^- converges in $L^2(\frac{1}{2}\mathbb{B})$ to a function u_∞^- . Hence, $u_j^+ - c^{-1}u_j^- \rightarrow u_\infty^+ - c^{-1}u_\infty^- =: u_\infty$ and

$$c \int \varphi d\omega_\infty = \int L_{A_0^*} \varphi u_\infty^- dx \quad \text{for any } \varphi \in C_c^\infty(\tfrac{1}{2}\mathbb{B}). \quad (3-33)$$

In particular, we can show that u_∞ is L_{A_0} -harmonic in $\frac{1}{2}\mathbb{B}$, and the rest of the proof is exactly as in [Azzam et al. 2016c] starting from equation (5.15). \square

4. Harmonic polynomial measures

4A. Preliminaries. We now review and collect some lemmas that will help us work with the quantities ω_h^A .

Lemma 4.1. *Let $h \in H_A$ and $r > 0$. Then*

$$T_{0,r}[\omega_h^A] = r^{n-1} \omega_{h \circ T_{0,r}^{-1}}^A, \quad (4-1)$$

$$F_r(\omega_h^A) = r^n F_1(\omega_{h \circ T_{0,r}^{-1}}^A). \quad (4-2)$$

Proof. By Lemma 3.10, it suffices to prove this in the case that $h \in H$. Note that if h is a harmonic function and $\varphi \in C_c^\infty(\mathbb{R}^{n+1})$, then

$$\begin{aligned} \int \varphi dT_{0,r}[\omega_h] &= \int \varphi \circ T_{0,r} d\omega_h \\ &= \int h \Delta(\varphi \circ T_{0,r}) dx = r^{-2} \int h \Delta \varphi \circ T_{0,r} dx \\ &= r^{n-1} \int h \circ T_{0,r}^{-1} \Delta \varphi dx = r^{n-1} \int \varphi d\omega_{h \circ T_{0,r}^{-1}}, \end{aligned} \quad (4-3)$$

and so (4-1) follows. Moreover, by Lemma 2.1(3),

$$F_r(\omega_h) = r F_1(T_{0,r}[\omega_h]) \stackrel{(4-1)}{=} r^n F_1(\omega_{h \circ T_{0,r}^{-1}}). \quad \square$$

Lemma 4.2. *Let $h \in F_A(k)$ and $r > 0$. Then*

$$F_r(\omega_h^A) = r^{n+k} F_1(\omega_h^A). \quad (4-4)$$

Proof. Note that since h is homogeneous of degree k ,

$$h \circ T_{0,r}^{-1}(x) = h(rx) = r^k h(x),$$

and thus, by (4-2),

$$F_r(\omega_h^A) = r^n F_1(\omega_{h \circ T_{0,r}^{-1}}^A) = r^n F_1(\omega_{r^k h}^A) = r^{n+k} F_1(\omega_h^A). \quad \square$$

The following is an immediate consequence of [Lemma 4.1](#).

Lemma 4.3 [[Badger 2011](#), Lemma 4.1]. *Since $\mathcal{F}_A(k)$, $\mathcal{P}_A(k)$, and \mathcal{H}_A are d -cones, so are $\mathcal{F}_{\mathcal{S}}(k)$, $\mathcal{P}_{\mathcal{S}}(k)$, and $\mathcal{H}_{\mathcal{S}}$ for any $\mathcal{S} \subset \mathcal{C}$.*

Lemma 4.4. *Let $A_j \in \mathcal{C}$ converge to a matrix $A \in \mathcal{C}$ and let $h_j \in H_{A_j}$ converge uniformly on compact subsets to some $h \in H_A$. Then $\omega_{h_j}^{A_j} \rightarrow \omega_h^A$ weakly.*

Proof. First we will deal with the case that $A_j = A = I$ for all j .

We first claim that, since h and h_j are harmonic, $\mathbb{1}_{\Omega_{h_j}} \rightarrow \mathbb{1}_{\Omega_h}$ a.e. Indeed, if $\mathbb{1}_{\Omega_h}(x) = 1$, then $h(x) > 0$, and by uniform convergence, $h_j(x) > 0$ for all large j , and so $\mathbb{1}_{\Omega_{h_j}}(x) = 1$ for all large j ; similarly, if $\mathbb{1}_{\Omega_h}(x) = 0$, then either $x \in \partial\Omega_h$ (which has measure zero) or $h_j(x) < 0$ for all large j , in which case $\mathbb{1}_{\Omega_{h_j}}(x) = 0$ for all large j . Thus, $\mathbb{1}_{\Omega_{h_j}} \rightarrow \mathbb{1}_{\Omega_h}$ pointwise everywhere in $(\partial\Omega_h)^c$ and thus a.e. in \mathbb{R}^{n+1} . In particular, $h_j \mathbb{1}_{\Omega_{h_j}} \rightarrow h \mathbb{1}_{\Omega_h}$ a.e. Hence, for $\varphi \in C_c^\infty(\mathbb{R}^{n+1})$, by the dominated convergence theorem,

$$\lim_{j \rightarrow \infty} \int \varphi d\omega_{h_j} = \lim_{j \rightarrow \infty} \int_{\Omega_{h_j}} h_j \Delta \varphi = \int_{\Omega_h} h \Delta \varphi = \int \varphi d\omega_h,$$

which implies $\omega_{h_j} \rightharpoonup \omega_h$ as $j \rightarrow \infty$.

Now we handle the general case. Let $A_{j,s} = \frac{1}{2}(A_j + A_j^T)$, and $S_j = \sqrt{A_{j,s}}$, and define A_s and S similarly. Let \tilde{A}_j and \tilde{A} be defined as in (3-6), and let $\tilde{h} = h \circ S$ and $\tilde{h}_j = h_j \circ S_j$. Since $\sqrt{\cdot}$ is continuous on the set of real symmetric matrices, $\tilde{h}_j \rightarrow \tilde{h}$ uniformly on compact subsets and both are harmonic. Thus, $\omega_{\tilde{h}_j} \rightharpoonup \omega_{\tilde{h}}$, and so

$$\lim_{j \rightarrow \infty} \omega_{h_j}^A \stackrel{(3-8)}{=} \lim_{j \rightarrow \infty} (\det S_j) S_j [\omega_{\tilde{h}_j}] = (\det S) S [\omega_{\tilde{h}}] \stackrel{(3-8)}{=} \omega_h^A. \quad \square$$

Lemma 4.5. *If $A \in \mathcal{C}$ and $h \in P_A(k)$ for some $k \in \mathbb{N}$, then*

$$\|h\|_{L^\infty(\mathbb{B})} \lesssim_{k,\Lambda} F_1(\omega_h^A). \quad (4-5)$$

Proof. Suppose instead that there exist $A_j \in \mathcal{C}$ and $h_j \in P_{A_j}(k)$ for which $\|h_j\|_{L^\infty(\mathbb{B})} > j F_1(\omega_{h_j}^{A_j})$. Without loss of generality, we may assume $\|h_j\|_{L^\infty(\mathbb{B})} = 1$, and thus $F_1(\omega_{h_j}^{A_j}) \rightarrow 0$. Using Cauchy estimates (see, e.g., Proposition 11.3 [[Mitrea 2013](#)]), $\{h_j\}_{j=1}^\infty$ forms a normal family in \mathbb{B} , and thus we can pass to a subsequence so that h_j converges uniformly on compact subsets of \mathbb{B} and so that A_j converges to some $A \in \mathcal{C}$. Since all h_j are polynomials of order k , we know that the coefficients of h_j converge, which, in turn, implies that h_j converges to some function $h \in \mathcal{P}_{\mathcal{C}}(k)$ uniformly on compact subsets of \mathbb{R}^{n+1} . By [Lemma 4.4](#), $\omega_{h_j}^{A_j} \rightarrow \omega_h^A$. In particular,

$$F_1(\omega_h^A) = \lim_{j \rightarrow \infty} F_1(\omega_{h_j}^{A_j}) = 0.$$

Thus, $\omega(B(0, r)) = 0$ for all $r < 1$, and so $0 \notin \text{supp } \omega_h$. We will now show that in fact $0 \in \text{supp } \omega_h^A$ in order to get a contradiction.

First, by [Lemma 3.10](#), we can assume without loss of generality that $A = I$ and $\omega_h^A = \omega_h$. Secondly, notice that as $h_j \in \mathcal{P}_{\mathcal{C}}(k)$, we have $h \in \mathcal{P}(k)$ and so $h(0) = 0$. By Lojasiewicz's structure theorem for real analytic varieties (see, e.g., [[Krantz and Parks 1992](#), Theorem 6.3.3, p. 168]), if U is a small enough

neighborhood of a point $0 \in \Sigma_h$, we have

$$U \cap \Sigma_h = V^n \cup V^{n-1} \cup \dots \cup V^0,$$

where V^0 is either the empty set or the singleton $\{0\}$ and for each $i \in \{1, \dots, n\}$, we may write V^i as a finite, disjoint union $V^i = \bigcup_{j=1}^{N_k} \Gamma_j^i$ of i -dimensional real analytic submanifolds. Further, for each $1 \leq i \leq n-1$,

$$U \cap \bar{V}^i \supset V^{i-1} \cup \dots \cup V^0.$$

Moreover, for $1 \leq k \leq n$ and $1 \leq j \leq N_k$, $U \cap \partial \Gamma_j^i$ is a union of sets of the form Γ_m^ℓ for $1 \leq \ell < i$ and $1 \leq m \leq N_\ell$ and possibly V^0 .

By the main result in [Cheeger et al. 2015], $\dim\{\nabla h=0\} \leq n-1$, and thus $V^n \cap \{\nabla h=0\}$ is a closed set of relatively empty interior in V^n , so in particular

$$\overline{V^n \setminus \{\nabla h=0\}} \cap U = \bar{V}^n \cap U = \Sigma_h \cap U \ni 0.$$

For $\zeta \in U \cap V^n \setminus \{\nabla h=0\}$, the derivative of h at ζ tangent to V^n is always zero, as h is zero on V^n , which forces ∇h to be perpendicular to V^n . Since the normal derivative is nonzero,

$$U \cap V^n \setminus \{\nabla h=0\} \subset \left\{ \zeta \in U \cap V^n : \frac{\partial h}{\partial \nu} \neq 0 \right\} \subset U \cap V^n \cap \text{supp } \omega_h.$$

Thus, $0 \in U \cap \overline{V^n \setminus \{\nabla h=0\}} \subset \text{supp } \omega_h$, which gives us the contradiction and concludes the proof. \square

4B. Proof of Proposition I. Proposition I is a consequence of the following more general result.

Lemma 4.6. *Let $\mathcal{S} \subset \mathcal{C}$ be closed (hence compact). Then $P_{\mathcal{S}}(k)$ and $\mathcal{F}_{\mathcal{S}}(k)$ have compact bases*

Proof. Let $h_j \in P_{A_j}(k)$ with $A_j \in \mathcal{S}$ and assume $\mathcal{F}(\omega_{h_j}^{A_j}) = 1$. Then by (4-5) and Cauchy estimates, we can bound each coefficient of the polynomials h_j uniformly, and then pass to a subsequence so that $A_j \rightarrow A \in \mathcal{S}$ and h_j converges on compact subsets of \mathbb{R}^{n+1} to a function $h \in P_A(k) \subset P_{\mathcal{S}}(k)$. By Lemma 4.4, we have $\omega_{h_j} \rightarrow \omega_h$, which implies that $\mathcal{P}_{\mathcal{S}}(k)$ has compact basis. The proof for $\mathcal{F}_{\mathcal{S}}(k)$ is similar. \square

As a corollary, we show the following stronger version of (4-5).

Corollary 4.7. *For $h \in P_{\mathcal{C}}(k)$ and $r > 0$,*

$$\|h\|_{L^\infty(r\mathbb{B})} \approx_k r^{-n} F_r(\omega_h). \quad (4-6)$$

Proof. Let $h \in P_{\mathcal{C}}(k)$ and $\varphi \in C_c^\infty(\mathbb{R}^{n+1})$ be such that $\mathbb{1}_{\mathbb{B}/2} \leq \varphi \leq \mathbb{1}_{\mathbb{B}}$. Since $\mathcal{P}_{\mathcal{C}}(k)$ has compact basis by Lemma 4.6, we can estimate

$$F_1(\omega_h) \stackrel{(2-4)}{\lesssim} F_{1/2}(\omega_h) \leq \int \varphi d\omega_h = \int_{\Omega_h} h \Delta \varphi \leq \|\Delta \varphi\|_\infty \int_{\mathbb{B}} |h| \lesssim \|h\|_{L^\infty(\mathbb{B})} \stackrel{(4-5)}{\lesssim} F_1(\omega_h).$$

For $r \neq 1$, by the previous inequalities we have

$$F_r(\omega_h) \stackrel{(4-2)}{=} r^n F_1(\omega_{h \circ T_{0,r}^{-1}}) \approx r^n \|h \circ T_{0,r}^{-1}\|_{L^\infty(\mathbb{B})} \approx r^n \|h\|_{L^\infty(r\mathbb{B})}. \quad \square$$

4C. Proof of Proposition II.

Lemma 4.8. *Let $h \in H_A$, $A \in \mathcal{C}$, and*

$$h(x) = \sum_{j=m}^{\infty} \sum_{|\alpha|=j} \frac{D^\alpha h(0)}{\alpha!} x^\alpha = \sum_{j=m}^{\infty} h_j(x)$$

be its Taylor series (where $m > 0$ and $h_m \neq 0$), which converges uniformly to h on compact subsets of \mathbb{R}^{n+1} . Then $\text{Tan}(\omega_h^A, 0) = \{c\omega_{h_m}^A : c > 0\}$.

Proof. For notational convenience, we will just consider the case $A = I$; the general case is identical. Note that as $r \rightarrow 0$, we have $r^{-m}h \circ T_{0,r}^{-1} \rightarrow h_m$ uniformly on compact subsets of \mathbb{R}^{n+1} . Indeed, fix $R > 0$. Then the series

$$r^{-m} \sum_{j=m}^{\infty} \sum_{|\alpha|=j} \frac{D^\alpha h(0)}{\alpha!} (rx)^\alpha = \sum_{j=m}^{\infty} \sum_{|\alpha|=j} \frac{D^\alpha h(0)}{\alpha!} x^\alpha r^{|\alpha|-m}$$

converges uniformly to $r^{-m}h \circ T_{0,r}^{-1}$ on compact subsets of $B(0, R)$, provided r is small enough. In fact, by Cauchy estimates,

$$|D^\alpha h(0)| \lesssim_n |\alpha|^{|\alpha|},$$

and since there exists a constant $C > 1$ such that $k^k/k! \lesssim C^k$, for $x \in B(0, R)$ and $r \in (0, 1/(CR))$ we have

$$\begin{aligned} |r^{-m}h \circ T_{0,r}^{-1}(x) - h_m(x)| &\leq \sum_{j=m+1}^{\infty} \sum_{|\alpha|=j} \left| \frac{D^\alpha h(0)}{\alpha!} \right| R^{|\alpha|} r^{|\alpha|-m} \\ &\lesssim_{n,m} \sum_{j=m+1}^{\infty} C^j R^j r^{j-m} \lesssim r^{-m} (CRr)^{m+1} = (CR)^{m+1} r \xrightarrow{r \downarrow 0} 0. \end{aligned}$$

Let now

$$v_r := r^{-m-n+1} T_{0,r}[\omega_h] \stackrel{(4-1)}{=} r^{-m} \omega_{h \circ T_{0,r}^{-1}} = \omega_{r^{-m}h \circ T_{0,r}^{-1}}.$$

By Lemma 4.4, $v_r \rightarrow \omega_{h_m} \in \mathcal{F}(m)$. In particular, every tangent measure of ω_h at zero must be a multiple of this one. \square

We now state an interesting consequence of these results: if a portion of tangent measures of an arbitrary Radon measure are in $\mathcal{P}(k)$, then in fact they are all in $\mathcal{F}(k)$ (that is, we did not have to assume the original measure was special like harmonic measure).

Lemma 4.9. *Let ω be a Radon measure, $\xi \in \text{supp } \omega$, and k be the minimal integer such that $\text{Tan}(\omega, \xi) \cap \mathcal{P}(k) \neq \emptyset$; then $\text{Tan}(\omega, \xi) \cap \mathcal{P}(k) \subset \mathcal{F}(k)$.*

We follow the proof in [Badger 2011, Lemma 5.9], which originally supposed that ω was a harmonic measure for an NTA domain.

Proof. If $k = 1$, then $\mathcal{P}(1) = \mathcal{F}(1)$. Now suppose $k > 1$ and there is $h \in \mathcal{P}(k)$ nonhomogeneous such that $\omega_h \in \text{Tan}(\omega, \xi) \cap \mathcal{P}(k)$. Since $h \in \mathcal{P}(k)$, we may write

$$h(x) = \sum_{j=m}^k \sum_{|\alpha|=j} \frac{D^\alpha h(0)}{\alpha!} x^\alpha = \sum_{j=m}^k h_m(x),$$

where $m < k$ since $h \in \mathcal{P}(k)$ is not homogeneous. By [Lemma 4.8](#), $\text{Tan}(\omega_h, 0) = \{c\omega_{h_m} : c > 0\} \subset \mathcal{F}(m)$, and since $\text{Tan}(\omega_h, 0) \subset \text{Tan}(\omega, \xi)$ by [Lemma 2.9](#), $\text{Tan}(\omega, \xi) \cap \mathcal{F}(m) \neq \emptyset$, contradicting the minimality of k . Thus, $\text{Tan}(\omega, \xi) \cap \mathcal{P}(k) \subset \mathcal{F}(k)$. \square

We will also need the following result.

Lemma 4.10 [[Badger 2011](#), Lemma 4.7]. *Suppose $h \in P(m)$ for some m . There exist $\varepsilon = \varepsilon(n, m, k) > 0$ and $r_0 > 0$ so that if $d_r(\omega_h, \mathcal{F}(k)) < \varepsilon$ for all $r \geq r_0$, then $m = k$.*

Proof of Proposition II. Suppose $\text{Tan}(\omega, \xi) \subset \mathcal{P}(k)$. Let m be the minimal integer for which $\text{Tan}(\omega, \xi) \cap \mathcal{P}(m) \neq \emptyset$, so $m \leq k$. Then, by [Lemma 4.9](#), $\text{Tan}(\omega, \xi) \cap \mathcal{P}(m) \subset \mathcal{F}(m)$. In particular, $\text{Tan}(\omega, \xi) \cap \mathcal{F}(m) \neq \emptyset$. Since, by [Proposition I](#), $\mathcal{P}(k)$ has compact basis, we can use [Lemmas 4.10](#) and [2.10](#) to conclude $\text{Tan}(\omega, \xi) \subset \mathcal{F}(m)$. \square

5. Proof of Theorem I

Lemma 5.1. *Let $\mathcal{S} \subset \mathcal{C}$ be closed and $\omega = \omega_{\Omega}^{A,x}$ be an L_A -harmonic measure where $A \in \mathcal{A}$ and $L_A \in \text{VMO}(\Omega, \xi)$ at $\xi \in \text{supp } \omega$. Also assume we have $\text{Tan}(\omega, \xi) \subset \mathcal{H}_{\mathcal{S}}$. Let k be the smallest integer for which $\text{Tan}(\omega, \xi) \cap \mathcal{F}_{\mathcal{S}}(k) \neq \emptyset$. Then $\text{Tan}(\omega, \xi) \subset \mathcal{F}_{\mathcal{S}}(k)$. In particular,*

$$\lim_{r \rightarrow 0} \frac{\log \omega(B(\xi, r))}{\log r} = n + k - 1; \quad (5-1)$$

i.e., the pointwise dimension of harmonic measure at the point ξ is $n + k - 1$.

Proof. If $\text{Tan}(\omega, \xi) \not\subset \mathcal{F}_{\mathcal{S}}(k)$, then by [Corollary 2.12](#), there is $r_0 > 0$ so that for any $\varepsilon > 0$ small we may find $v \in \text{Tan}(\omega, \xi) \setminus \mathcal{F}_{\mathcal{S}}(k)$ so that $d_{r_0}(v, \mathcal{F}_{\mathcal{S}}(k)) = \varepsilon$ and $d_r(v, \mathcal{F}_{\mathcal{S}}(k)) \leq \varepsilon$ for all $r \geq r_0$. Without loss of generality, we can assume $r_0 = 1$. For each $r > 1$, choose $\mu_r \in \mathcal{F}_{\mathcal{S}}(k)$ such that $F_r(\mu_r) = 1$ and

$$F_r\left(\frac{v}{F_r(v)}, \mu_r\right) < 2\varepsilon.$$

Then for $r \geq 1$,

$$\begin{aligned} \frac{F_r(v)}{F_{2r}(v)} &= \int (r - |x|)_+ d\frac{v}{F_{2r}(v)} < 2\varepsilon + \int (r - |x|)_+ d\mu_{2r} = 2\varepsilon + F_r(\mu_{2r}) \\ &\stackrel{(4-4)}{=} 2\varepsilon + 2^{-n-k} F_{2r}(\mu_{2r}) = 2\varepsilon + 2^{-n-k} = 2^{-n-k+\beta} \end{aligned}$$

for some $\beta > 0$ that goes to zero as $\varepsilon \rightarrow 0$. Similarly,

$$\frac{F_r(v)}{F_{2r}(v)} \geq 2^{-n-k-\beta}.$$

Hence, for $\ell \in \mathbb{N}$,

$$2^{\ell(n+k-\beta)} \leq \frac{F_{2^\ell r}(v)}{F_r(v)} \leq 2^{\ell(n+k+\beta)}. \quad (5-2)$$

Note that $v = \omega_h^A$ for some $h \in \mathcal{H}_A$ by [Theorem 3.13](#) and $A \in \mathcal{S}$, and so

$$\|h\|_{L^\infty(2^\ell \mathbb{B})} \stackrel{(3-28)}{\lesssim} 2^{\ell(1-n)} \omega_h(B(0, 2^{\ell+1})) \leq 2^{-\ell n-1} F_{2^{\ell+2}}(\omega_h) \stackrel{(5-2)}{\leq} 2^{\ell(k+\beta)-1} F_{2^2}(\omega_h). \quad (5-3)$$

Let α be a multi-index of length $|\alpha| > k$. Then we can pick $\varepsilon > 0$ small enough so that β is so small that $|\alpha| - k - \beta > 0$ holds. Thus, by Cauchy estimates,

$$|\partial^\alpha h(0)| \lesssim_\alpha 2^{-\ell|\alpha|} \|h\|_{L^\infty(2^\ell \mathbb{B})} \stackrel{(5-3)}{\lesssim} 2^{-\ell(|\alpha| - k - \beta)} F_{2^2}(\omega_h) \rightarrow 0$$

as $\ell \rightarrow \infty$, and so $h \in \mathcal{P}_A(k)$.

Suppose $h = \sum_{j=1}^k h_j$. If $\omega_h \notin \mathcal{F}_A(k)$, then there exists $j < k$ such that $h_j \neq 0$, and by Lemma 4.8, we infer that $\text{Tan}(\omega_h^A, 0)$ contains an element of $\mathcal{F}_A(j)$. Since $\omega_h^A \in \text{Tan}(\omega, \xi)$, we know that $\text{Tan}(\omega_h^A, 0) \subset \text{Tan}(\omega, \xi)$ by Lemma 2.9 and thus, $\text{Tan}(\omega, \xi) \cap \mathcal{F}_A(j) \neq \emptyset$. Hence $\text{Tan}(\omega, \xi) \cap \mathcal{F}_{\mathcal{J}}(j) \neq \emptyset$, contradicting the minimality of k . This proves $\text{Tan}(\omega, \xi) \subset \mathcal{F}_{\mathcal{J}}(k)$.

For the final equality, note that $\text{Tan}(\omega, \xi) \subset \mathcal{F}_{\mathcal{J}}(k)$ and so $\text{Tan}(\omega, \xi)$ has compact basis. In particular, by Lemma 2.11,

$$\lim_{r \rightarrow 0} d_1(T_{\xi, r}[\omega], \mathcal{F}_{\mathcal{J}}(k)) = 0.$$

Thus, for $\varepsilon > 0$, there is $r_0 > 0$ such that for each $r \leq r_0$ there exists $\mu_r \in \mathcal{F}_{\mathcal{J}}(k)$ so that $F_1(\mu_r) = 1$ and

$$F_1\left(\frac{T_{\xi, r}[\omega]}{F_1(T_{\xi, r}[\omega])}, \mu_r\right) < \varepsilon.$$

Setting $\nu_r = r^{-1}T_{\xi, r}^{-1}[\mu_r]$, this gives $F_r(\nu_r) = 1$ and

$$F_r\left(\frac{\omega}{F_r(\omega)}, \nu_r\right) < \varepsilon.$$

By the same arguments as earlier, we can show that there exists $\gamma > 0$, which goes to zero as $\varepsilon \rightarrow 0$, so that for all $\ell \geq 0$ and $r < 2^{-\ell-1}r_0$

$$2^{\ell(n+k-\gamma)} \leq \frac{F_{2^\ell r}(\omega)}{F_r(\omega)} \leq 2^{\ell(n+k+\gamma)}. \quad (5-4)$$

Hence, if we set $d = n + k - 1$, we get

$$\begin{aligned} \omega(B(\xi, 2^\ell r)) &= T_{\xi, r}[\omega](B(0, 2^\ell)) \leq 2^{-\ell} F_{2^{\ell+1}}(T_{\xi, r}[\omega]) \\ &\leq 2^{(\ell+1)(n+k+\gamma)-\ell} F_1(T_{\xi, r}[\omega]) \\ &\leq 2^{\ell(d+\gamma)+n+k+\gamma} T_{\xi, r}[\omega](B(0, 1)) \\ &= 2^{\ell(d+\gamma)+n+k+\gamma} \omega(B(\xi, r)). \end{aligned}$$

Similarly,

$$\begin{aligned} \omega(B(\xi, r)) &= T_{\xi, r}[\omega](B(0, 1)) \leq F_2(T_{\xi, r}[\omega]) \\ &\leq 2^{-(\ell-1)(n+k-\gamma)} F_{2^\ell}(T_{\xi, r}[\omega]) \\ &\leq 2^{-(\ell-1)(n+k-\gamma)+\ell} \omega(B(\xi, 2^\ell r)) \\ &= 2^{-\ell(d-\gamma)+n+k-\gamma} \omega(B(\xi, 2^\ell r)). \end{aligned}$$

For $r < \frac{1}{2}r_0$, let $\ell \in \mathbb{N}$ be so that $2^{-\ell-1}r_0 \leq r \leq 2^{-\ell}r_0$. Then

$$\begin{aligned} \omega(B(\xi, r)) &\leq \omega(B(\xi, 2^{-\ell}r_0)) \leq 2^{-\ell(d-\gamma)+n+k-\gamma} \omega(B(\xi, r_0)) \\ &\leq 2^{1+(n+k-\gamma)} r^{d-\gamma} \omega(B(\xi, r_0)). \end{aligned}$$

Hence, recalling that these logs are negative, we conclude

$$\liminf_{r \rightarrow 0} \frac{\log \omega(B(\xi, r))}{\log r} \geq \liminf_{r \rightarrow 0} \frac{\log(2^{1+(n+k-\gamma)} \omega(B(\xi, r_0)))}{\log r} + d - \gamma = d - \gamma.$$

A similar estimate gives

$$\limsup_{r \rightarrow 0} \frac{\log \omega(B(\xi, r))}{\log r} \leq d + \gamma.$$

If we let $\gamma \rightarrow 0$, then (5-1) follows. □

Proof of Theorem I. We set

$$E^* = \left\{ \xi \in E : \lim_{r \rightarrow 0} \frac{\omega^+(E \cap B(\xi, r))}{\omega^+(B(\xi, r))} = \lim_{r \rightarrow 0} \frac{\omega^-(E \cap B(\xi, r))}{\omega^-(B(\xi, r))} = 1 \right\},$$

$$E^{**} = \{\xi \in E^* : (1-4) \text{ holds}\}.$$

Notice that by [Mattila 1995, Corollary 2.14(1)] and because ω_1 and ω_2 are mutually absolutely continuous on E ,

$$\omega^+(E \setminus E^{**}) = \omega^-(E \setminus E^{**}) = 0.$$

Also, set

$$\Lambda_1 = \left\{ \xi \in E^{**} : 0 < h(\xi) := \frac{d\omega^-}{d\omega^+}(\xi) = \lim_{r \rightarrow 0} \frac{\omega^-(B(\xi, r))}{\omega^+(B(\xi, r))} = \lim_{r \rightarrow 0} \frac{\omega^-(E \cap B(\xi, r))}{\omega^+(E \cap B(\xi, r))} < \infty \right\},$$

$$\Gamma = \{\xi \in \Lambda_1 : \xi \text{ is a Lebesgue point for } h \text{ with respect to } \omega^+\}.$$

Again, by Lebesgue differentiation for measures (see [Mattila 1995, Corollary 2.14(2) and Remark 2.15(3)]), Γ has full measure in E^{**} and hence in E .

Next, we record a lemma which was proven in [Azzam et al. 2017b, Lemma 5.8] (which in turn is based on the work of [Kenig et al. 2009]) in the case of the harmonic functions in domains that satisfy the CDC condition, but its proof goes through unchanged for L -harmonic functions in general domains.

Lemma 5.2. *Let $\xi \in \Gamma$, $c_j \geq 0$, and $r_j \rightarrow 0$ be so that $\omega_j^+ = c_j T_{\xi, r_j}[\omega^+] \rightarrow \omega_\infty$. Then $\omega_j^- = c_j T_{\xi, r_j}[\omega^-] \rightarrow h(\xi)\omega_\infty$.*

We define

$$\mathcal{F} := \{c\mathcal{H}^n|_V : c > 0, V \text{ a } d\text{-dimensional plane containing the origin}\}.$$

It is not hard to show that \mathcal{F} has compact basis.

Lemma 5.3. *For ω^+ -a.e. $\xi \in \Gamma$,*

$$\text{Tan}(\omega^+, \xi) \cap \mathcal{F} \neq \emptyset.$$

Proof. We can pick $\xi \in \Gamma$ so that $\text{Tan}(\omega^+, \xi) \neq \emptyset$, let $\omega_\infty \in \text{Tan}(\omega^+, \xi)$, so there is $c_j > 0$ and $r_j \downarrow 0$ so that $c_j T_{\xi, r_j}[\omega^+] \rightarrow \omega_\infty$. By Lemma 5.2, we also have $c_j T_{\xi, r_j}[\omega^-] \rightarrow h(\xi)\omega_\infty$. By Theorem 3.13, (3-27) holds.

In particular, $\frac{1}{2}\mathbb{B} \cap \text{supp } \omega_\infty$ is a smooth real analytic variety, and arguing as in [Azzam et al. 2016c], for example, one deduces that

$$d\omega_\infty|_{\mathbb{B}/2} = -c_n(\nu_{\Omega_\infty^+} \cdot A_0 \nabla u_\infty) d\mathcal{H}^n|_{\partial^* \Omega_\infty^+ \cap \mathbb{B}/2},$$

where A_0 is the matrix from [Theorem 3.13](#), $\partial^* \Omega_\infty^+$ is the reduced boundary of $\Omega_\infty^+ = \{u_\infty > 0\}$ and $\nu_{\Omega_\infty^+}$ is the measure-theoretic outer unit normal. Hence, ω_∞ is absolutely continuous with respect to surface measure of $\partial \Omega_\infty^+$ in $\frac{1}{2}\mathbb{B}$. Thus, since the tangent measure at \mathcal{H}^n -almost every point of $\partial \Omega_\infty^+$ is contained in \mathcal{F} , we can take another tangent measure of ω_∞ that is in \mathcal{F} and apply [Theorem 2.8](#) to conclude the proof. \square

By [Lemmas 5.1](#) and [5.3](#), we also have that $\dim \omega^+|_E = n$. It remains to show that if Ω^\pm both have the CDC, then $\lim_{r \rightarrow 0} \Theta_{\partial \Omega^+}^{\mathcal{F}}(\xi, r) = 0$ for ω^+ -a.e. $\xi \in E$. But this follows almost immediately because, for almost every $\xi \in \Gamma$ and any $r_j \downarrow 0$, we may pass to a subsequence so that, by [Lemma 3.11\(a\)](#) and (f), $\lim_{j \rightarrow \infty} \Theta_{\partial \Omega^+}^{\mathcal{F}}(\xi, r_j) = 0$. This concludes the proof of [Theorem I](#). \square

6. BMO, VMO and vanishing A_∞

In this section, we will prove some estimates relating the logarithm of a Radon–Nikodym derivative to the mutual absolute continuity properties of two measures. We will apply them to the specific case of elliptic measure, but we will prove them for general measures.

Definition 6.1. Let μ be a Radon measure on a metric space X . We say that a function $f \in L^1_{\text{loc}}(\mu)$ is of *bounded mean oscillation* and write $f \in \text{BMO}(\mu)$ if there exists a constant $C > 0$ such that

$$\sup_{r \in (0, \infty)} \sup_{x \in \text{supp } \mu} \int_{B(x, r)} |f - f_{B(x, r)}| d\mu \leq C, \quad (6-1)$$

where $f_A := \int_A f d\mu := \mu(A)^{-1} \int_A f d\mu$ for any $A \subset X$ with $\mu(A) > 0$. We define the space of *vanishing mean oscillation* $\text{VMO}(\mu)$ to be the closure in the $\text{BMO}(\mu)$ norm of the set of bounded uniformly continuous functions defined on X . Equivalently, we say $f \in \text{VMO}(\mu)$ if $f \in L^1_{\text{loc}}(\mu)$ and

$$\lim_{r \rightarrow 0} \sup_{x \in \text{supp } \mu} \int_{B(x, r)} |f - f_{B(x, r)}| d\mu = 0. \quad (6-2)$$

Definition 6.2. For two measures μ and ν on a metric space X , we will say $\nu \in A_\infty(\mu)$ if $\mu \ll \nu$ and there is $K = K(\mu, \nu)$ so that for any ball B centered on the support of μ

$$\int_B \frac{d\nu}{d\mu} d\mu \exp\left(-\int_B \log \frac{d\nu}{d\mu} d\mu\right) \leq K(\mu, \nu). \quad (6-3)$$

We will say $\nu \in A'_\infty(\mu)$ if there are $\varepsilon, \delta \in (0, 1)$ so that for all $B \subseteq X$ and $E \subseteq B$

$$\frac{\mu(E)}{\mu(B)} < \delta \implies \frac{\nu(E)}{\nu(B)} < \varepsilon. \quad (6-4)$$

We will say $\nu \in \text{VA}_\infty(\mu)$ (or *vanishing A_∞ with respect to μ*) if

$$\lim_{r \rightarrow 0} \sup_{\xi \in \text{supp } \mu} \int_B \frac{d\nu}{d\mu} d\mu \exp\left(-\int_B \log \frac{d\nu}{d\mu} d\mu\right) = 1 \quad (6-5)$$

and $\nu \in \text{VA}'_\infty(\mu)$ if for all $r > 0$ there is $\varepsilon_r \in (0, 1)$ so that $\lim_{r \rightarrow 0} \varepsilon_r = 0$ and $\delta_r > 0$ so that for all balls $B \subset X$ with $r_B < r$ and $E \subset B$

$$\frac{\mu(E)}{\mu(B)} < \delta_r \implies \frac{\nu(E)}{\nu(B)} < \varepsilon_r. \quad (6-6)$$

In the case that $X = \mathbb{R}^{n+1}$ and μ is equal to the $(n+1)$ -dimensional Lebesgue measure, A_∞ -equivalence is the same as A'_∞ -equivalence, and this is from [Reimann and Rychener 1975], although it was also shown later in [Khrushchev 1984; García-Cuerva and Rubio de Francia 1985].

We recall a notion introduced in [Korey 1998].

Definition 6.3. A probability space (X, μ) is *halving* if every subset $E \subset X$ of positive measure has a subset $F \subset E$ so that $\mu(F) = \frac{1}{2}\mu(E)$.

We will first focus on proving the following after a series of other lemmas.

Lemma 6.4. Let (X, μ) be a metric measure space, $\nu \ll \mu$, and $f = d\nu/d\mu$:

- (1) If $\nu \in A'_\infty(\mu)$ and $\log f \in \text{BMO}(\mu)$, then $\nu \in A_\infty(\mu)$. If X is also halving, then $\nu \in A_\infty(\mu)$ implies $\nu \in A'_\infty(\mu)$ and $\log f \in \text{BMO}(\mu)$.
- (2) If $\nu \in VA'_\infty(\mu)$ and $\log f \in \text{VMO}(\mu)$, then $\nu \in VA_\infty(\mu)$. If X is also halving, then $\nu \in VA_\infty(\mu)$ implies $\nu \in VA'_\infty(\mu)$ and $\log f \in \text{VMO}(\mu)$.

The first implication of the second half of (1) of the lemma is a consequence of the following theorem.

Theorem 6.5 [Khrushchev 1984, Theorem 1]. Suppose $\nu \ll \mu$, B is a ball centered on $\text{supp } \mu$, and

$$\int_B \frac{d\nu}{d\mu} d\mu \exp\left(-\int_B \log \frac{d\nu}{d\mu} d\mu\right) \leq C.$$

Then there are $\varepsilon, \delta > 0$ so that, for any $F \subset B \cap \text{supp } \mu$,

$$\frac{\mu(F)}{\mu(B)} < \delta \implies \frac{\nu(F)}{\nu(B)} < \varepsilon. \quad (6-7)$$

Moreover, there is $\delta > 0$ so that

$$\frac{\mu(F)}{\mu(B)} < \delta \implies \frac{\nu(F)}{\nu(B)} < 2(C-1). \quad (6-8)$$

In particular, if $\nu \in A_\infty(\mu)$, then $\nu \in A'_\infty(\nu)$, and if $\nu \in VA_\infty(\mu)$, then $\nu \in VA'_\infty(\mu)$.

Proof. We follow the proof from [Khrushchev 1984, Theorem 1], since he proves (6-7) but not (6-8). Let $\delta \in (0, 1)$ to be chosen later, let $F \subseteq B$ and suppose $\mu(F) = \delta\mu(B)$; we will pick δ later. Let $f = d\nu/d\mu$, $E = B \setminus F$, and set

$$t = \frac{\nu(E)}{\nu(F)}.$$

Let $g_B = \int_B f d\mu$. Then

$$\log C \geq (\log f^{-1})_B + \log f_B = \frac{\mu(E)}{\mu(B)} (\log f^{-1})_E + \frac{\mu(F)}{\mu(B)} (\log f^{-1})_F + \log f_B. \quad (6-9)$$

By Jensen's inequality, for any set S

$$(\log f^{-1})_S = -(\log f)_S \geq -\log f_S,$$

and applying this to $S = E, F$, we have

$$\begin{aligned} \log C &\geq -\frac{\mu(E)}{\mu(B)} \log f_E - \frac{\mu(F)}{\mu(B)} \log f_F + \log f_B \\ &\geq -\frac{\mu(E)}{\mu(B)} \log f_E - \frac{\mu(F)}{\mu(B)} \log f_E + \frac{\mu(F)}{\mu(B)} \log \frac{\mu(F)}{\mu(E)} + \frac{\mu(F)}{\mu(B)} \log t + \log f_B \\ &= -\log f_E + \frac{\mu(F)}{\mu(B)} \log \frac{\mu(F)}{\mu(E)} + \frac{\mu(F)}{\mu(B)} \log t + \log f_B. \end{aligned}$$

Now observe that

$$-\log f_E = \log \left(\frac{\mu(E)}{\mu(B)} \frac{\mu(B)}{v(B)} \frac{v(B)}{v(E)} \right) = \log \frac{\mu(E)}{\mu(B)} - \log f_B + \log \left(1 + \frac{1}{t} \right)$$

and so we have

$$\begin{aligned} \log C &\geq \log \frac{\mu(E)}{\mu(B)} + \log \left(1 + \frac{1}{t} \right) + \frac{\mu(F)}{\mu(B)} \log \frac{\mu(F)}{\mu(E)} + \frac{\mu(F)}{\mu(B)} \log t \\ &= \frac{\mu(F)}{\mu(B)} \log \frac{\mu(F)}{\mu(B)} + \frac{\mu(E)}{\mu(B)} \log \frac{\mu(E)}{\mu(B)} + \log(1+t) + \frac{\mu(E)}{\mu(B)} \log \frac{1}{t} \\ &= \underbrace{\delta \log \delta + (1-\delta) \log(1-\delta)}_{=: \varphi(\delta)} + \log(1+t) + \frac{\mu(E)}{\mu(B)} \log \frac{1}{t}. \end{aligned}$$

Note that $\lim_{\delta \rightarrow 0} \varphi(\delta) = 0$. Let $\alpha > 0$ and pick $\delta > 0$ so that $|\varphi(\delta)| < \alpha \log C$. Then

$$(1+\alpha) \log C \geq \log(1+t) + \frac{\mu(E)}{\mu(B)} \log \frac{1}{t}. \quad (6-10)$$

We restrict δ further so that $\delta < \alpha$. If $t > 1$, then

$$\frac{\mu(E)}{\mu(B)} \log \frac{1}{t} \geq \log \frac{1}{t};$$

otherwise,

$$\frac{\mu(E)}{\mu(B)} \log \frac{1}{t} \geq (1-\alpha) \log \frac{1}{t}$$

since $\mu(E)/\mu(B) = 1-\delta > 1-\alpha$. Thus, in any case, we have

$$\frac{1+\alpha}{1-\alpha} \log C > \log \frac{1}{t}. \quad (6-11)$$

This implies $t \geq c = C^{-(1+\alpha)/(1-\alpha)}$, and so

$$v(F) = \frac{v(F)}{1+t} + \frac{tv(F)}{1+t} = \frac{v(F) + v(E)}{1+t} = \frac{v(B)}{1+t} \leq \frac{v(B)}{1+c}.$$

This proves (6-7) with $\varepsilon = (1+c)^{-1}$. To prove (6-8), we go back to (6-10) with the same bound on δ .

Then, since $t \geq c$,

$$\begin{aligned} (1+\alpha) \log C &\geq \log(1+t) + \frac{\mu(E)}{\mu(B)} \log \frac{1}{t} = \log \left(1 + \frac{1}{t} \right) + \frac{\mu(F)}{\mu(B)} \log t \\ &\geq \log \left(1 + \frac{1}{t} \right) - \delta \frac{1+\alpha}{1-\alpha} \log C. \end{aligned}$$

Since $\delta < \alpha$, this implies

$$\log\left(1 + \frac{1}{t}\right) < \left(1 + \alpha + \delta \frac{1+\alpha}{1-\alpha}\right) \log C = (1+\alpha) \left(1 + \frac{\delta}{1-\alpha}\right) \log C < \frac{1+\alpha}{1-\alpha} \log C,$$

and so

$$C^{(1+\alpha)/(1-\alpha)} - 1 > \frac{1}{t}.$$

We now pick α so that $C^{(1+\alpha)/(1-\alpha)} - 1 = 2(C - 1)$, and we are done. \square

Korey showed that VA_∞ implies the logarithm of the density is VMO.

Theorem 6.6 [Korey 1998, Theorem 4 and Section 3.5]. *There is a universal constant $c > 0$ so that the following holds. Let (X, μ) be a halving probability space, and suppose that*

$$\frac{(\int_X \exp g \, d\mu)}{\exp(\int_X g \, d\mu)} \leq K. \quad (6-12)$$

Then

$$\int_X \left| g - \int_X g \, d\mu \right| d\mu \leq \log 2K \quad (6-13)$$

and as $K \rightarrow 1$,

$$\int_X \left| g - \int_X g \, d\mu \right| d\mu \leq c\sqrt{K-1}. \quad (6-14)$$

Lemma 6.7. *Let (X, μ) be a metric probability space and suppose $\nu \ll \mu$. Let $\varepsilon, \delta \in (0, 1)$ be so that for any $E \subset X$*

$$\mu(E) < \delta\mu(X) \implies \nu(E) < \varepsilon\nu(X). \quad (6-15)$$

Set $f = d\nu/d\mu$ and assume

$$\int_X \left| \log f - \int_X \log f \, d\mu \right| d\mu < \eta. \quad (6-16)$$

Then

$$1 \leq \int_X f \, d\mu \exp\left(-\int_X \log f \, d\mu\right) \leq \frac{e^{\eta/\delta}}{1-\varepsilon}. \quad (6-17)$$

Proof. Without loss of generality, we may assume $\mu(X) = \nu(X) = 1$. Let $\varepsilon > 0$ and pick δ so that (6-15) holds.

Let $c = \int_X \log f \, d\mu$ and

$$G = \{|\log f - c| < \rho := \eta\delta^{-1}\}, \quad F = G^c. \quad (6-18)$$

Then, by Chebyshev's inequality and (6-16), we infer that $\mu(F) < \delta$, which, in turn, by (6-15), implies

$$\nu(F) < \varepsilon. \quad (6-19)$$

Moreover, on the set G ,

$$\frac{\eta}{\delta} > |\log f - c|$$

and so

$$f \leq e^{c+\eta/\delta} \quad \text{on } G. \quad (6-20)$$

Then,

$$\begin{aligned} 1 = \frac{\nu(X)}{\mu(X)} &= \int_X f \, d\mu \stackrel{(6-20)}{\leq} \left(\int_G e^{c+\eta/\delta} \, d\mu + \int_F f \, d\mu \right) \\ &\leq e^{c+\eta/\delta} + \nu(F) \stackrel{(6-19)}{<} e^{c+\eta/\delta} + \varepsilon. \end{aligned}$$

Thus,

$$(1 - \varepsilon) \int_X f \, d\mu = 1 - \varepsilon < e^{c+\eta/\delta}$$

and so

$$\int_X f \, d\mu < \frac{e^{c+\eta/\delta}}{1 - \varepsilon}. \quad (6-21)$$

This and Jensen's inequality imply

$$1 \leq e^{-c} \int_X f \, d\mu < e^{-c} \frac{1}{1 - \varepsilon} e^{c+\eta/\delta} = \frac{1}{1 - \varepsilon} e^{\eta/\delta}. \quad \square$$

Corollary 6.8. *Let (X, μ) be a metric measure space. Set $f = dv/d\mu$ and assume that for some sequence of balls B_j in X*

$$\lim_j \int_{B_j} \left| \log f - \int_{B_j} \log f \, d\mu \right| d\mu = 0 \quad (6-22)$$

and for all $\varepsilon > 0$ there is $\delta > 0$ so that for j sufficiently large

$$\frac{\mu(E)}{\mu(B_j)} < \delta \implies \frac{\nu(E)}{\nu(B_j)} < \varepsilon. \quad (6-23)$$

Then

$$\lim_{j \rightarrow \infty} \int_{B_j} f \, d\mu \exp\left(-\int_{B_j} \log f \, d\mu\right) = 1. \quad (6-24)$$

In particular, if $\log f \in \text{VMO}(d\mu)$ and $v \in VA'_\infty(\mu)$, then $v \in VA_\infty(\mu)$.

Proof. Let $\varepsilon, \eta > 0$ and let $\delta > 0$ be so that (6-23) holds for j large enough. Then (6-16) holds (with B_j in place of X and $\mu|_{B_j}$ in place of μ). Then (6-17) must hold. In particular,

$$\limsup_{j \rightarrow \infty} \int_{B_j} f \, d\mu \exp\left(-\int_{B_j} \log f \, d\mu\right) \leq \frac{e^{\eta/\delta}}{1 - \varepsilon}.$$

As ε and δ did not depend on η , we can send $\eta \rightarrow 0$, and then $\varepsilon \rightarrow 0$ since δ now vanishes from the inequality, and then we obtain (6-24). \square

Proof of Lemma 6.4. The second halves of (1) and (2) follow from Theorems 6.5 and 6.6. The first half of (1) follows from Lemma 6.7, and the first half of (2) is from Corollary 6.8. \square

Lemma 6.9. *Let $\Omega \subset \mathbb{R}^{n+1}$ be any connected domain and $\omega = \omega_\Omega^{L_{A,x}}$ where $A \in \mathcal{A}(\Omega)$. Then ω is halving.*

Proof. Suppose there is $E \subset \partial\Omega$ with $\omega(E) > 0$ that is not halving. For $t \in \mathbb{R}$ and $v \in \mathbb{S}^{n-1}$, let $H_{t,v} = \{x \in \mathbb{R}^{n+1} : x \cdot v \geq t\}$. Then $t \mapsto \omega(H_{t,v} \cap E)$ is not continuous for any $v \in \mathbb{S}^n$, and so there is t_v so that $\omega(\partial H_{t_v,v} \cap E) > 0$. Let $V_v = \partial H_{t_v,v}$, which is an n -dimensional plane. Since \mathbb{S}^n is uncountable, there is $\varepsilon > 0$ so that $\omega(V_v \cap E) > \varepsilon > 0$ for all v in some uncountable set $A \subset \mathbb{S}^n$. Let $A' \subset A$ be

countable. Note that for any $u, v \in A'$ distinct, $V_u \cap V_v$ is an $(n-1)$ -dimensional subspace. This implies $V_u \cap V_v$ has 2-capacity zero [Heinonen et al. 1993, Theorem 2.27]; hence it is a polar set for ω [loc. cit., Theorem 10.1] and polar sets have L_A -harmonic measure zero [loc. cit., Theorem 11.15]. Thus, if we set

$$W_u := V_u \setminus \bigcup_{\substack{v \in A' \\ v \neq u}} V_v,$$

we have $\omega(W_u \cap E) = \omega(V_u \cap E) \geq \varepsilon$ and W_u are mutually disjoint. But since A' is infinite, this implies $\omega(E) = \infty$, which is a contradiction. \square

Lemma 6.10. *Let $\Omega^+ \subset \mathbb{R}^{n+1}$ be a connected domain with connected complement $\Omega^- = \text{ext}(\Omega^+)$ and let L_A be a uniformly elliptic operator with real coefficients. If ω^\pm denote the L_A -harmonic measures of Ω^\pm with fixed poles $x^\pm \in \Omega^\pm$, then $\omega^- \in A_\infty(\omega^+)$ if and only if $\omega^- \in A'_\infty(\omega^+)$ and $\log(d\omega^-/d\omega^+) \in \text{BMO}(d\omega^+)$. Moreover, $\omega^- \in VA_\infty(\omega^+)$ if and only if $\omega^- \in VA'_\infty(\omega^+)$ and $\log(d\omega^-/d\omega^+) \in \text{VMO}(d\omega^+)$.*

Proof. This follows from Lemmas 6.4 and 6.9. \square

7. Proofs of Theorems II and III

Lemma 7.1. *Let ω^\pm be two halving Radon measures with equal supports and set $f = \log(d\omega^-/d\omega^+)$. Suppose there are $r_j \downarrow 0$ and $\xi_j \in \partial\Omega^+$ so that $\omega_j^+ = T_{\xi_j, r_j}[\omega^+]/\omega(B(\xi_j, r_j))$ converges weakly to some measure ω with $\omega(\mathbb{B}) > 0$. Further assume that for all $M > 0$*

$$\lim_j \int_{B(\xi_j, Mr_j)} f d\omega^+ \exp\left(-\int_{B(\xi_j, Mr_j)} \log f d\omega^+\right) = 1. \quad (7-1)$$

Then $\omega_j^- \rightarrow \omega$ as well.

The proof is similar to that of [Kenig and Toro 2006, Theorem 4.4], though using the techniques of the previous section, we no longer require the doubling assumption.

Proof. Let $B_j = B(\xi_j, r_j)$ and for a ball B set $c_B = \int_B \log f$. By assumption, for each $M > 0$,

$$e^{-c_{MB_j}} \frac{\omega^-(MB_j)}{\omega^+(MB_j)} \rightarrow 1 \quad \text{as } j \rightarrow \infty. \quad (7-2)$$

Let $\varphi \in C_c^\infty(\mathbb{R}^{n+1})$ with support in $B(0, M)$ for some $M > 0$ and let $\varphi_j = \varphi \circ T_{\xi_j, r_j}$. Then $\text{supp } \varphi_j \subset MB_j$. Let $\varepsilon > 0$. By (7-2), for j large enough, we have

$$0 \leq e^{-c_{B_j}} \frac{\omega^-(B_j)}{\omega^+(B_j)} - 1 < \varepsilon \quad \text{and} \quad 0 \leq e^{-c_{MB_j}} \frac{\omega^-(MB_j)}{\omega^+(MB_j)} - 1 < \varepsilon. \quad (7-3)$$

Let now $\eta = c\sqrt{1-\varepsilon}$, where c is the constant in (6-14). For j large enough, Theorem 6.6 and (7-2) imply

$$\int_{B_j} |\log f - c_{B_j}| d\omega^+ < \eta \quad \text{and} \quad \int_{MB_j} |\log f - c_{MB_j}| d\omega^+ < \eta. \quad (7-4)$$

Note that ε is independent of η . For fixed $\delta > 0$ and for a ball B , we set

$$G_B = \{\xi \in B \cap \partial\Omega^+ : |\log f(\xi) - c_B| \leq \eta/\delta\}, \quad F_B = B \setminus G_B.$$

Then, Chebyshev's inequality and (7-4) imply

$$\omega^+(F_{B_j}) < \delta \omega^+(B_j) \quad \text{and} \quad \omega^+(F_{MB_j}) < \delta \omega^+(MB_j), \quad (7-5)$$

and for $\delta > 0$ small enough and j large enough Theorem 6.5 and (7-2) imply

$$\omega^-(F_{B_j}) < \varepsilon, \quad \omega^-(B_j) \quad \text{and} \quad \omega^-(F_{MB_j}) < \varepsilon \omega^-(MB_j). \quad (7-6)$$

Let $C = 2\omega(\overline{M\mathbb{B}})/\omega(\mathbb{B})$. Since $\omega(\mathbb{B}) > 0$, we know

$$\limsup_{j \rightarrow \infty} \frac{\omega^+(MB_j)}{\omega^+(B_j)} = \limsup_{j \rightarrow \infty} \frac{\omega_j^+(M\mathbb{B})}{\omega_j^+(\mathbb{B})} \leq \frac{\omega(\overline{M\mathbb{B}})}{\omega(\mathbb{B})} = \frac{1}{2}C,$$

and so for j large enough,

$$\omega^+(MB_j) \leq C\omega^+(B_j). \quad (7-7)$$

Also, note that for j large enough,

$$\begin{aligned} |c_{B_j} - c_{MB_j}| &= \left| \int_{B_j} (c_{B_j} - c_{MB_j}) \right| d\omega^+ \\ &\leq \int_{B_j} |c_{B_j} - \log f| d\omega^+ + \int_{B_j} |\log f - c_{MB_j}| d\omega^+ \\ &\stackrel{(7-4)}{<} \eta + \frac{\omega^+(MB_j)}{\omega^+(B_j)} \int_{MB_j} |\log f - c_{MB_j}| d\omega^+ \stackrel{(7-7)}{<} (1+C)\eta. \end{aligned} \quad (7-8)$$

Hence,

$$\begin{aligned} \omega^-(MB_j) &\stackrel{(7-3)}{\leq} \omega^+(MB_j)(1+\varepsilon)e^{c_{MB_j}} \stackrel{(7-8)}{<} C\omega^+(B_j)(1+\varepsilon)e^{c_{B_j}+(1+C)\eta} \\ &\stackrel{(7-3)}{\leq} C\omega^-(B_j)(1+\varepsilon)e^{(1+C)\eta} \leq 2Ce^{(1+C)\eta}\omega^-(B_j) \lesssim_C \omega^-(B_j). \end{aligned} \quad (7-9)$$

Then

$$\begin{aligned} \int \varphi d\omega_j^- - \int \varphi d\omega_j^+ &= \frac{1}{\omega^-(B_j)} \int_{MB_j} \varphi_j d\omega^- - \frac{1}{\omega^+(B_j)} \int_{MB_j} \varphi_j d\omega^+ \\ &= \underbrace{\frac{1}{\omega^-(B_j)} \int_{MB_j \cap F_{MB_j}} \varphi_j f d\omega^+}_{=:I_1} + \underbrace{\frac{1}{\omega^-(B_j)} \int_{MB_j \cap G_{MB_j}} (f - e^{c_{MB_j}}) \varphi_j d\omega^+}_{=:I_2} \\ &\quad - \underbrace{\frac{e^{c_{MB_j}}}{\omega^-(B_j)} \int_{MB_j \cap F_{MB_j}} \varphi_j d\omega^+}_{=:I_3} + \underbrace{\frac{e^{c_{MB_j}}}{\omega^-(B_j)} \int_{MB_j} \varphi_j d\omega^+ - \frac{1}{\omega^+(B_j)} \int_{MB_j} \varphi_j d\omega^+}_{=:I_4} \\ &= I_1 + I_2 - I_3 + I_4. \end{aligned}$$

We will estimate each of these terms separately, with the understanding that j is large enough (depending on M and η):

$$|I_1| \leq \frac{\|\varphi\|_\infty}{\omega^-(B_j)} \int_{MB_j} \mathbb{1}_{F_{MB_j}} f d\omega^+ = \frac{\|\varphi\|_\infty \omega^-(F_{MB_j})}{\omega^-(B_j)} = \frac{\omega^-(MB_j)}{\omega^-(B_j)} \frac{\|\varphi\|_\infty \omega^-(F_{MB_j})}{\omega^-(MB_j)} \stackrel{(7-6)}{\stackrel{(7-9)}}{\lesssim_{C,M,\|\varphi\|_\infty}} \varepsilon.$$

Next, for points in G_{MB_j} ,

$$e^{-\eta/\delta} e^{cMB_j} \leq f \leq e^{\eta/\delta} e^{cMB_j}$$

and so

$$e^{cMB_j} (e^{-\eta/\delta} - 1) \leq f - e^{cMB_j} \leq e^{cMB_j} (e^{\eta/\delta} - 1).$$

Thus, for $\eta > 0$ small enough (i.e., for j large enough), we can make

$$|f - e^{cMB_j}| < \delta e^{cMB_j} \quad \text{on } G_{MB_j}.$$

Therefore,

$$\begin{aligned} |I_2| &\leq \frac{\delta e^{cMB_j} \|\varphi\|_\infty}{\omega^-(B_j)} \omega^+(G_{MB_j}) \leq \frac{\delta e^{cMB_j} \|\varphi\|_\infty}{\omega^-(B_j)} \omega^+(MB_j) \\ &= e^{cMB_j} \frac{\omega^+(MB_j)}{\omega^-(MB_j)} \frac{\delta \|\varphi\|_\infty \omega^-(MB_j)}{\omega^-(B_j)} \stackrel{(7-9)}{\lesssim} \|\varphi\|_\infty, C, M \delta, \\ |I_3| &\leq \frac{e^{cMB_j} \|\varphi\|_\infty}{\omega^-(B_j)} \omega^+(F_{MB_j}) \stackrel{(7-5)}{<} \delta \frac{e^{cMB_j} \|\varphi\|_\infty}{\omega^-(B_j)} \omega^+(MB_j) \\ &= \delta \frac{e^{cMB_j} \|\varphi\|_\infty \omega^-(MB_j)}{\omega^-(B_j)} \frac{\omega^+(MB_j)}{\omega^-(MB_j)} \stackrel{(7-3)}{\lesssim} C, M, \|\varphi\|_\infty \delta. \end{aligned}$$

Finally,

$$|I_4| \leq \left(e^{cMB_j} \frac{\omega^+(B_j)}{\omega^-(B_j)} - 1 \right) \frac{\omega^+(MB_j)}{\omega^+(B_j)} \int_{MB_j} \varphi_j d\omega^+ \stackrel{(7-3)}{\lesssim} C, \|\varphi\|_\infty, M \varepsilon.$$

Since these estimates hold for all j large enough, we can conclude

$$\limsup_{j \rightarrow \infty} \left| \int \varphi d\omega_j^- - \int \varphi_j d\omega_j^+ \right| \lesssim_{C, M, \|\varphi\|_\infty} \varepsilon + \delta.$$

Now send δ to zero since it only had to be small enough depending on ε . Finally, ε was arbitrarily chosen, which implies that the above limit is zero. Since this holds for all φ , we get that ω_j^\pm have the same weak limit. \square

Proof of Theorem II. Let $\omega \in \text{Tan}(\omega^+, \xi)$. We claim that $\omega \in \mathcal{H}_\ell$. By Lemma 2.6, $\omega = cT_{0,r}(\mu)$ for some constants $c, r > 0$ and some measure μ of the form $\mu = \lim_{j \rightarrow 0} T_{\xi, r_j}[\omega^+]/\omega^+(B(\xi, r_j))$ for some $r_j \downarrow 0$, where $\mu(\mathbb{B}) > 0$. By Lemma 7.1, $\mu = \lim_{j \rightarrow 0} T_{\xi, r_j}[\omega^-]/\omega^-(B(\xi, r_j))$ as well. By Theorem 3.13 (or Lemma 3.11(g) if Ω^\pm have the CDC), $\mu \in \mathcal{H}_\ell$, and since \mathcal{H}_ℓ is a d -cone by Lemma 4.3, we also have that $\omega \in \mathcal{H}_\ell$, which proves the claim.

Hence, $\omega = \omega_u$ for some $u \in H_A$ and some $A \in \mathcal{C}$. By Lemma 4.8, for some $k > 0$,

$$\text{Tan}(\omega_u, 0) = \{c\omega_{u_k} : c > 0\} \subset \mathcal{F}_A(k) \subset \mathcal{F}_\ell(k),$$

and since $\text{Tan}(\omega_u, 0) \subset \text{Tan}(\omega^+, \xi)$ by Lemma 2.9, we now know that $\text{Tan}(\omega^+, \xi) \cap \mathcal{F}_\ell(k) \neq \emptyset$ as well. By Lemma 5.1, $\text{Tan}(\omega^+, \xi) \subset \mathcal{F}_\ell(k)$. The proof that $\Theta_{\partial\Omega^+}^{\mathcal{F}_\Sigma, \mathcal{C}}(k)(\xi, r) \rightarrow 0$ if Ω^\pm have the CDC is similar to the proof of Theorem I. \square

Proof of Theorem III. Let K be any compact subset of $\partial\Omega^+$. Suppose there is a sequence of radii $r_j \downarrow 0$ and $\xi_j \in K$ so that

$$d_1(T_{\xi_j, r_j}[\omega^+], \mathcal{P}_{\mathcal{E}}(d)) \geq \varepsilon > 0, \quad (7-10)$$

where d will be chosen later, but it will depend only on n and the doubling constant of ω^+ .

Since ω^+ is doubling, we may pass to a subsequence so that $\omega_j^+ := T_{\xi_j, r_j}[\omega^+]/\omega^+(B(\xi_j, r_j))$ converges weakly to some measure ω .

If $f = d\omega^-/d\omega^+$ satisfies $\log f \in \text{VMO}(\omega^-)$, then doubling also implies that $\omega^- \in VA'_\infty(\omega^+)$. Indeed, if ω^+ is doubling, then the John–Nirenberg theorem holds, and the VMO condition tells us that on small enough balls, f is a traditional A_p -weight (see [Garnett 2007, Chapter 6.2]). This easily implies $f d\omega^+ = d\omega^- \in VA'_\infty(\omega^+)$. Thus, by Corollary 6.8, we know $\omega^- \in VA_\infty(\omega^+)$ and that (7-1) holds for every $M > 0$. By Lemma 7.1, $\omega_j^- \rightharpoonup \omega$ as well. Thus, we can pass to a subsequence so that the conclusions of Theorem 3.13 hold. In particular, $\omega = \omega_h$ for some L_0 -harmonic function h , where L_0 is a uniformly elliptic operator with constant coefficients, and also, for any $\varphi \in C_c^\infty(\mathbb{R}^{n+1})$, (3-21) holds.

Now we apply the same standard trick from [Kenig and Toro 2006]. Notice that since ω^+ is doubling, so is ω_h , which combined with Cauchy estimates implies that there exists $\beta > 0$ such that for any $\ell \in \mathbb{N}$ and any multi-index α

$$\begin{aligned} |\partial_\alpha h(0)| &\lesssim 2^{-|\alpha|\ell} \|h\|_{L^\infty(2^\ell \mathbb{B})} \stackrel{(3-28)}{\lesssim} 2^{\ell(-|\alpha|+1-n)} \omega_h(B(0, 2^{\ell+1})) \\ &\lesssim 2^{\ell(-|\alpha|+1-n+\beta)} \omega_h(B(0, 2)). \end{aligned} \quad (7-11)$$

Hence, if $|\alpha| > 1 - n + \beta$, letting $\ell \rightarrow \infty$ gives $|\partial_\alpha h(0)| = 0$, which implies h is a polynomial of degree at most $1 - n + \beta$. Setting $d = \lceil 1 - n + \beta \rceil$ gives a contradiction to (7-10). The proof of (1-7) is similar to the proof of Theorem I, where we use instead Lemma 3.11 instead of Theorem 3.13. \square

8. Proof of Theorem IV

All elliptic operators in this section will be assumed to satisfy (1-1) and (1-2). We will require a few lemmas about elliptic measures in uniform domains as well as some new notation.

Definition 8.1. Let $\Omega \subseteq \mathbb{R}^{n+1}$:

- We say Ω satisfies the *corkscrew condition* if, for some uniform constant $c > 0$ and every ball B centered on $\partial\Omega$ with $0 < r_B < \text{diam}(\partial\Omega)$, there is a ball $B(x_B, cr_B) \subseteq \Omega \cap B$. The point x_B is called a *corkscrew point relative to B* .
- We say Ω satisfies the *Harnack chain condition* if there is a uniform constant C such that for every $\rho > 0$, $\Lambda \geq 1$, and every pair of points $x, y \in \Omega$ with $\delta(x), \delta(y) \geq \rho$ and $|x - y| < \Lambda\rho$ there is a chain of open balls $B_1, \dots, B_N \subset \Omega$, $N \leq C(\Lambda)$, with $x \in B_1$, $y \in B_N$, $B_k \cap B_{k+1} \neq \emptyset$ and $C^{-1} \text{diam}(B_k) \leq \text{dist}(B_k, \partial\Omega) \leq C \text{diam}(B_k)$. The chain of balls is called a *Harnack chain*.

Definition 8.2. If Ω satisfies both the corkscrew and the Harnack chain conditions, then we say that Ω is a *uniform domain*.

Theorem 8.3. *Let $\Omega \subset \mathbb{R}^{n+1}$ be a uniform domain with the CDC and u a nonnegative L_A -elliptic function vanishing on $2B \cap \partial\Omega$, where B is a ball with $r_B < \text{diam } \partial\Omega$ and $A \in \mathcal{A}(\Omega)$. Then*

$$\sup_{x \in B \cap \Omega} u(x) \lesssim u(x_B). \quad (8-1)$$

This was originally shown in Section 4 of [Jerison and Kenig 1982] for NTA domains, but the proof only uses the Hölder continuity of u at the boundary and the fact that NTA domains are uniform, and so the proof of the above result is exactly the same.

Theorem 8.4. *Let $\Omega \subset \mathbb{R}^{n+1}$ be a uniform domain with the CDC and L_A an elliptic operator satisfying (1-1) and (1-2). Then, for all B centered on $\partial\Omega$,*

$$\omega^{L_A, x}(B) \approx r_B^{n-1} G_\Omega(x, x_B) \quad \text{for all } x \in \Omega \setminus 2B. \quad (8-2)$$

This follows from [Aikawa and Hirata 2008]. Their proof is originally for harmonic measures, but an inspection of the proof shows that it carries through for elliptic measure as well.

Theorem 8.5. *Let $\Omega \subset \mathbb{R}^{n+1}$ be a uniform domain with the CDC. If L_A is an elliptic operator satisfying (1-1) and (1-2), B is a ball centered on $\partial\Omega$, and $E \subset B \cap \partial\Omega$ is Borel, then*

$$\omega_\Omega^{L_A, x_B}(E) \approx \frac{\omega_\Omega^{L_A, x}(E)}{\omega_\Omega^{L_A, x}(B)}. \quad (8-3)$$

Again, this is [Jerison and Kenig 1982, Lemma 4.11], and since the previous two lemmas are available, the proof is exactly the same for elliptic measures modulo the proof of [loc. cit., Lemma 4.10]. The latter can also be proved by building a subuniform domain as in [loc. cit.], and then showing as in [Akman et al. 2019, Lemma 2.26] that the resulting domain is also CDC (all of this instead of a geometric localization theorem due to Jones, which only works for NTA domains).

Lemma 8.6. *Let $\Omega \subset \mathbb{R}^{n+1}$ be a uniform domain with the CDC and L_A an elliptic operator satisfying (1-1) and (1-2), and also (1-4) at ξ . If $\xi \in \partial\Omega$ and $\omega_j = \omega^{L_A, x_0}(B(\xi, r_j))^{-1} T_{\xi, r_j}(\omega^{L_A, x_0})$ converges weakly to a tangent measure $\omega_\infty \in \text{Tan}(\omega^{L_A, x_0}, \xi)$, then there is a uniform domain Ω_∞ and a constant matrix $A_0 \in \mathcal{C}$ such that, for each $x \in \Omega_\infty$, $\omega_{\Omega_j}^x \rightharpoonup \omega_{\Omega_\infty}^x$ and for all balls $B' \subset B$ centered on $\partial\Omega_\infty$, if x_B is a corkscrew point in $\Omega_\infty \cap B$,*

$$\omega_{\Omega_\infty}^{L_{A_0}, x_B}(B') \approx \frac{\omega_\infty(B')}{\omega_\infty(B)}. \quad (8-4)$$

This was originally shown in [Azzam and Mourgoglou 2018] for harmonic measure. In our situation, the proof is much shorter, so we provide it here.

Proof. By Lemma 3.11, there is $A_0 \in \mathcal{C}$ so that we can pass to a subsequence so that $u_j(x) = c_j u(xr_j + \xi)r_j^{n-1}$ converges uniformly in \mathbb{R}^{n+1} to a nonzero L_{A_0} -elliptic function u_∞ and also so that, if $\Omega_j = T_{\xi, r_j}(\Omega)$, then $\partial\Omega_j$ converges in the Hausdorff metric on compact subsets. Let $\Omega_\infty = \{u_\infty > 0\}$.

Claim. Ω_∞ is uniform. If $x, y \in \Omega_\infty$ with $\text{dist}(\{x, y\}, \partial\Omega) \geq \varepsilon|x - y|$, then they are contained in Ω_j and $\text{dist}(\{x, y\}, \partial\Omega_j) \geq \frac{1}{2}\varepsilon|x - y|$ for sufficiently large j . Since the Ω_j are uniform, for each j we can find a Harnack chain of length $N = N(\varepsilon)$ contained in Ω_j . By passing to a subsequence, we can assume the

length of this chain is constant and their centers and radii are converging, and hence the chain converges to a Harnack chain in Ω_∞ of length no more than N . A similar proof shows that Ω_∞ is a corkscrew domain. Hence, Ω_∞ is uniform.

Suppose $B' \subset \mathbb{B}$ are centered on $\partial\Omega_\infty$. Let

$$\omega_{\Omega_j}^{T_{\xi,r_j}(x)} = T_{\xi,r_j}[\omega_{\Omega_j}^{L_A,x}].$$

If $x_j = T_{\xi,r_j}(x_0)$, then

$$\omega_{\Omega_j}^{x_B}(B') \approx \frac{\omega_{\Omega_j}^{x_j}(B')}{\omega_{\Omega_j}^{x_j}(B)} = \frac{\omega_{\Omega_j}^{x_j}(\mathbb{B})}{\omega_{\Omega_j}^{x_j}(B)} \frac{\omega_{\Omega_j}^{x_j}(B')}{\omega_{\Omega_j}^{x_j}(\mathbb{B})} = \frac{\omega_j(B')}{\omega_j(B)}.$$

Since ω_j and ω_{Ω_j} are doubling measures, we have

$$\omega_{\Omega_\infty}^{x_B}(B') \leq \liminf_{j \rightarrow \infty} \omega_{\Omega_j}^{x_B}(B') \lesssim \limsup_{j \rightarrow \infty} \frac{\omega_j(B')}{\omega_j(B)} \leq \frac{\omega_\infty(\bar{B}')}{\omega_\infty(B)} \lesssim \frac{\omega_\infty(B')}{\omega_\infty(B)}.$$

A similar estimate gives the reverse inequality, and hence proves (8-4). \square

We will use the following criterion for uniform rectifiability due to Hofmann, Martell, and Uriarte-Tuero. See Theorem 1.23, equation 1.22, and Remark 1.25 in [Hofmann et al. 2014]; for a local version see Corollary 11.2 in [Mourgoglou and Tolsa 2017].

Theorem 8.7. *Let $\Omega \subset \mathbb{R}^{n+1}$ be a uniform domain with n -regular boundary and let ω_Ω be the harmonic measure defined in Ω . Suppose there is $q > 1$ so that, for any balls $B' \subset B$ centered on $\partial\Omega$, if $k_B = d\omega_\Omega^{x_B}/(d\mathcal{H}^n|_{\partial\Omega})$, then*

$$\left(\int_{B' \cap \partial\Omega} k_B^q d\mathcal{H}^n \right)^{1/q} \lesssim \int_{2B' \cap \partial\Omega} k_B d\mathcal{H}^n.$$

Then $\partial\Omega$ is uniformly rectifiable.

Recall that, by the main result of [Aikawa and Hirata 2008], harmonic measure is doubling in uniform domains satisfying the CDC, and thus, by (8-3), the right side of this inequality is comparable to $\int_{B' \cap \partial\Omega} k_B d\mathcal{H}^n$ (that is, with B' instead of $2B'$), which we will use below.

Remark 8.8. This result still holds for constant coefficients. Indeed, it is easy to see that the A_∞ -property is preserved under linear transformations that map balls to ellipsoids, as is the one in Lemma 3.10 (see the paragraph after the proof of this lemma), using that such weights are doubling. Thus, by Lemma 3.10 and the fact that being a uniformly rectifiable set, by its very definition, is invariant under bi-Lipschitz maps, $\partial\Omega_\infty$ is uniformly rectifiable.

Recall that an Ahlfors n -regular set E is *uniformly rectifiable* if there are $c, L > 0$ so that, for every ball B centered on E with $r_B < \text{diam } E$, there is an L -Lipschitz map $f : B(0, r_B) \cap \mathbb{R}^n \rightarrow \mathbb{R}^{n+1}$ so that

$$\mathcal{H}^n(f(B(0, r_B)) \cap E) \geq cr_B^n.$$

Now we prove Theorem IV. Let $\Omega \subset \mathbb{R}^{n+1}$ be a uniform CDC domain so that $\mathcal{H}^n|_{\partial\Omega}$ is locally finite. Let $\omega = \omega_\Omega^{L_A}$ be the L_A -harmonic measure associated to a (possibly nonsymmetric) elliptic operator

satisfying (1-1) and (1-2). Let $E \subseteq \partial\Omega$ be a set with $\mathcal{H}^n(E) > 0$ such that $\mathcal{H}^n \ll \omega_\Omega^{L_A}$ on E and for \mathcal{H}^n -a.e. $\xi \in E$,

$$\theta_{\partial\Omega,*}^n(\xi, r) := \liminf_{r \rightarrow 0} \frac{\mathcal{H}^n(B(\xi, r) \cap \partial\Omega)}{(2r)^n} > 0$$

and A has vanishing mean oscillation at ξ .

Assume $\mathcal{H}^n(E) > 0$ (otherwise the theorem is trivial). Then we may find a subset E' of full \mathcal{H}^n -measure, where ω and \mathcal{H}^n are mutually absolutely continuous (in particular, $\mathcal{H}^n = g\omega$ for some function g , so we pick $E' = \{x : g(x) > 0\}$). For $\mathcal{H}^n|_{\partial\Omega}$ -a.e. $\xi \in E'$, we also have

$$0 < \theta_*^n(\mathcal{H}^n|_{\partial\Omega}, \xi) \leq \theta^{n,*}(\mathcal{H}^n|_{\partial\Omega}, \xi) < \infty. \quad (8-5)$$

The lower bound is by assumption, and the upper bound is from [Mattila 1995, Theorem 6.2]. By [loc. cit., Theorem 14.7], for $\mathcal{H}^n|_{\partial\Omega}$ -a.e. $\xi \in E'$, $\text{Tan}(\mathcal{H}^n|_{\partial\Omega}, \xi)$ consists of Ahlfors–David n -regular measures. By [loc. cit., Lemmas 14.5 and 14.6], for $\mathcal{H}^n|_{\partial\Omega}$ -a.e. $\xi \in E'$,

$$\text{Tan}(\mathcal{H}^n|_{\partial\Omega}, \xi) = \text{Tan}(\mathcal{H}^n|_{E'}, \xi) = \text{Tan}(\omega, \xi)$$

and $\text{Tan}(\omega, \xi)$ consists only of Ahlfors–David n -regular measures. Let $E'' \subset E'$ be the set of points where this holds.

By the Besicovitch decomposition theorem, we can split E'' into two sets F_1 and F_2 , where F_1 is n -rectifiable and F_2 is purely n -unrectifiable. Suppose $\mathcal{H}^n(F_2) > 0$. Let $\xi \in F_2$ be a point of density of F_2 with respect to \mathcal{H}^n .

Let $r_j \downarrow 0$ be so that $\omega_j := \omega_\Omega^{L_A, x_0}(B(\xi, r_j))^{-1} T_{\xi, r_j}(\omega_\Omega^{L_A, x_0})$ converges weakly to some Ahlfors–David n -regular measure $\omega_\infty \in \text{Tan}(\omega, \xi)$. By Lemma 8.6, we may find a uniform domain Ω_∞ so that $\text{supp } \omega_\infty = \partial\Omega_\infty$ and, for any balls $B' \subset B$ centered on $\partial\Omega$,

$$\omega_{\Omega_\infty}^{L_{A_0}, x_B}(B') \approx \frac{\omega_\infty(B')}{\omega_\infty(B)} \approx \frac{r_{B'}^n}{r_B^n}$$

for some $A_0 \in \mathcal{C}$. If $\sigma = \mathcal{H}^n|_{\partial\Omega_\infty}$, then σ is Ahlfors–David n -regular and so if we set

$$k_B := \frac{d\omega_{\Omega_\infty}^{L_{A_0}, x_B}}{d\sigma},$$

then we have that for σ -a.e. $x \in B \cap \partial\Omega$

$$k_B(x) = \lim_{r \rightarrow 0} \frac{\omega_{\Omega_\infty}^{L_{A_0}, x_B}(B(x, r))}{\sigma(B(x, r))} \approx \frac{r^n/r_B^n}{r^n} = r_B^{-n}.$$

Hence, if $B' \subset B$ is centered on $\partial\Omega$,

$$\left(\int_{B'} k_B^2 d\sigma \right)^{1/2} \approx r_B^{-n} \approx \int_{B'} k_B d\sigma.$$

Thus, in light of Remark 8.8, $\partial\Omega_\infty$ is uniformly rectifiable. By the main result of [Azzam et al. 2017a], Ω_∞ is an NTA domain. In particular, we can find corkscrew balls $B_1 \subset \mathbb{B} \cap \Omega_\infty$ and $B_2 \subseteq \mathbb{B} \setminus \Omega_\infty$. We claim that, for all j sufficiently large, $\frac{1}{2}B_1 \subset \Omega_j \cap \mathbb{B}$ and $\frac{1}{2}B_2 \subset \mathbb{B} \setminus \Omega_j$. Indeed, if $\frac{1}{2}B_i \cap \partial\Omega_j \neq \emptyset$ for

infinitely many j , then since ω_j is doubling, $\omega_j(\frac{2}{3}B_i) \sim \omega_j(\mathbb{B}) = 1$ for all j , and so $\omega_\infty(B_i) > 0$, and in particular $\partial\Omega_\infty \cap B_i \neq \emptyset$, which is a contradiction. Thus, B_1 and B_2 do not intersect $\partial\Omega_j$ for sufficiently large j . They cannot both be in Ω_j for all large j , since otherwise, if they were both in Ω_j for infinitely many j then in each such Ω_j , they would be connected by a Harnack chain in Ω_j of bounded length; passing to a subsequence, this implies there is a Harnack chain connecting B_1 to B_2 , and since $B_1 \subseteq \Omega_\infty$, the whole chain, including B_2 , must be in Ω_∞ , which is a contradiction. Thus, at least one of these balls is in Ω_j^c for all j large. By the proof of [Lemma 8.6](#), $\Omega_\infty = \{u_\infty > 0\}$, and since $u_j \rightarrow u_\infty$ uniformly on compact subsets of Ω_∞ and $u_\infty > 0$ on B_1 , we have $B_1 \subset \Omega_j$ for j large, and so $B_2 \subset \Omega_j^c$ for j large. This proves the claim.

Now there is a small angle of directions around the vector parallel to the line between the centers of B_1 and B_2 where the orthogonal projection of $\partial\Omega_j \cap \mathbb{B}$ has Lebesgue measure comparable to 1. By the Besicovitch–Federer projection theorem, the purely unrectifiable part of $\partial\Omega_j$ has zero Lebesgue measure projection in almost all of these directions, and so $\partial\Omega_j \cap \mathbb{B}$ contains an n -rectifiable set of \mathcal{H}^n -measure $\gtrsim 1$ (with constant depending on the sizes of B_1 and B_2). Thus,

$$\liminf_{j \rightarrow \infty} \frac{\mathcal{H}^n(B(\xi, r_j) \cap \partial\Omega \setminus F_2)}{\mathcal{H}^n(B(\xi, r_j) \cap \partial\Omega)} \gtrsim \liminf_{j \rightarrow \infty} \frac{r_j^n}{\mathcal{H}^n(B(\xi, r_j) \cap \partial\Omega)} \stackrel{(8-5)}{>} 0.$$

But this contradicts that ξ is a point of density for F_2 . Therefore, $\mathcal{H}^n(F_2) = 0$, and we have now shown that \mathcal{H}^n -almost all of E' is rectifiable, and thus ω^{x_0} -almost all of E is contained in a countable union of Lipschitz graphs. This finishes the proof of [Theorem IV](#).

9. Proof of [Proposition III](#)

Assume the conditions of the proposition. We recall the following result.

Theorem 9.1 [[Hurri-Syrjänen 1994](#), Theorem 1.3]. *Suppose that $\Omega \subset \mathbb{R}^{n+1}$ is a bounded C -uniform² domain. If*

$$p \leq q \leq \frac{(n+1)p}{n+1-p(1-\delta)} \quad \text{and} \quad p(1-\delta) < n+1,$$

then for all $u \in L^1_{\text{loc}}(\Omega)$ such that $\nabla u(x)d(x, \partial\Omega)^\delta \in L^p(\Omega)$,

$$\inf_{a \in \mathbb{R}} \|u(x) - a\|_{L^q(\Omega)} \lesssim_{n,p,q,\delta,C} |\Omega|^{(1-\delta)/(n+1)+1/q-1/p} \|\nabla u \operatorname{dist}(\cdot, \Omega^c)^\delta\|_{L^p(\Omega)}. \quad (9-1)$$

(The explicit constant in (9-1) is written at the end of the proof on page 218 of [[Hurri-Syrjänen 1994](#)].) We will use this in the case that $\delta = \frac{1}{2}$ and $p = q = 2$, so (9-1) becomes

$$\inf_{a \in \mathbb{R}} \|u(x) - a\|_{L^2(\Omega)} \lesssim_{n,p,q,\delta,C} |\Omega|^{1/(2(n+1))} \|\nabla u \operatorname{dist}(\cdot, \Omega^c)^{1/2}\|_{L^2(\Omega)}. \quad (9-2)$$

Lemma 9.2. *Suppose $E \subset \mathbb{R}^{n+1}$ is a closed set and $\varepsilon : E^c \rightarrow [0, \infty]$ is a function such that for some ball B_0 centered on E*

$$\int_{E^c \cap B_0} \varepsilon(z) dz < \infty.$$

²In fact it holds for John domains.

Then for \mathcal{H}^n -a.e. $x \in E \cap B_0$,

$$\lim_{r \rightarrow 0} r^{-n} \int_{E^c \cap B(x, r)} \varepsilon(z) dz = 0.$$

Proof. Without loss of generality, we can assume $E \subset B_0$. Let $d\mu(z) = \varepsilon(z) dz|_{E^c}$. For $x \in E$ and $r > 0$, set

$$a(x, r) = \frac{\mu(B(x, r))}{r^n} = r^{-n} \int_{E^c \cap B(x, r)} \varepsilon(z) dz.$$

Suppose there is $F \subset E$ with $\mathcal{H}^n(F) > 0$ such that

$$\limsup_{r \rightarrow 0} a(x, r) > 0.$$

Then there is $t > 0$ and a compact set $G \subset F$ with $\mathcal{H}_\infty^n(G) > 0$ and

$$\limsup_{r \rightarrow 0} a(x, r) > t > 0 \quad \text{for all } x \in G.$$

For each $x \in G$, pick $r_{x,1} > 0$ so that $B(x, r_{x,1}) \subset B_0$ and $a(x, r_{x,1}) > t$. Let B_j^1 be a Besicovitch subcovering from $\mathcal{G}_1 := \{B(x, r_x^1) : x \in G\}$, that is, a countable collection of balls in \mathcal{G}_1 so that

$$\mathbb{1}_G \leq \sum_j \mathbb{1}_{B_j^1} \lesssim_n 1.$$

Since the B_j^1 come from \mathcal{G} , we have that for all j

$$\frac{\mu(B_j^1)}{r_{B_j^1}^n} = a(x_{B_j^1}, r_{B_j^1}) > t.$$

Let

$$L_1 = \bigcup B_j^1 \setminus E.$$

Then since the B_j^1 have bounded overlap and come from \mathcal{G}_1 ,

$$\mu(L_1) = \int_{L_1} d\mu \gtrsim \int_{L_1} \sum_j \mathbb{1}_{B_j^1} d\mu = \sum_j \mu(B_j^1) > t \sum_j r_{B_j^1}^n \geq t \mathcal{H}_\infty^n(G).$$

Since $\mu(G) = 0$, there is $\delta_1 > 0$ so that if $G_{\delta_1} = \{x \in \mathbb{R}^n : \text{dist}(x, G) < \delta_1\}$ and $L^1 = L_1 \setminus G_{\delta_1}$, then

$$\mu(L^1) > \frac{1}{2} \mu(L_1) \geq \frac{1}{2} t \mathcal{H}_\infty^n(G).$$

Now inductively, suppose we have constructed disjoint sets $L^1, \dots, L^k \subset B_0$, where

$$\mu(L^j) \gtrsim t \mathcal{H}_\infty^n(G) \quad \text{for all } j = 1, 2, \dots, k,$$

and there is $\delta_k > 0$ so that $L^1 \cup \dots \cup L^k \cap G_{\delta_k} = \emptyset$.

For each $x \in G$, we may find $r_{x,k+1} \in (0, \delta_k)$ so that $B(x, r_{x,k+1}) \subset B_0$ and $a(x, r_{x,k+1}) > t$. Let $\{B_j^{k+1}\}$ be a Besicovitch subcovering of the collection $\mathcal{G}_{k+1} = \{B(x, r_{x,k+1}) : x \in G\}$, so

$$\mathbb{1}_G \leq \sum_j \mathbb{1}_{B_j^{k+1}} \lesssim_n \mathbb{1}_{L_{k+1}},$$

where $L_{k+1} = \bigcup_j B_j^{k+1}$. Since G has $\mu(G) = 0$, there is $\delta_{k+1} \in (0, \delta_k)$ so that $L^{k+1} = L_{k+1} \setminus G_{\delta_{k+1}}$ has

$$\mu(L^{k+1}) \geq \frac{\mu(L_{k+1})}{2} = \frac{1}{2} \int \mathbb{1}_{L_{k+1}} d\mu \gtrsim \int \sum_j \mu(B_j^{k+1}) \geq t \sum_j r_{B_j^{k+1}}^n \gtrsim t \mathcal{H}_\infty^n(G).$$

Also note that by our induction hypothesis

$$L^{k+1} \subset L_{k+1} \subset G_{\delta_k} \subset (L^1 \cup \dots \cup L^k)^c.$$

Thus, by induction, we can come up with a sequence of disjoint sets $L^k \subset B_0$ so that $\mu(L^k) \gtrsim t \mathcal{H}_\infty^n(G)$ for all k , which contradicts the finiteness of μ since ε is locally integrable. \square

Now we finish the proof of [Proposition III](#). By the previous lemma, for $\varepsilon(z) = |\nabla A(z)|^2 \text{dist}(z, \Omega^c)$ and $E = \partial\Omega$, we have that for \mathcal{H}^n -a.e. $\xi \in B_0 \cap \partial\Omega$

$$\lim_{r \rightarrow 0} r^{-n} \int_{B(\xi, r) \cap \Omega} |\nabla A|^2 \text{dist}(z, \Omega^c) dz = 0. \quad (9-3)$$

Let $\xi \in B_0 \cap \partial\Omega$ be such a point. There is a universal constant M depending on the uniformity constants so that, for all $r > 0$, there is an MC -uniform domain Ω_r such that

$$\Omega \cap B(\xi, r) \subset \Omega_r \subset \Omega \cap B(\xi, Mr).$$

This follows from the proof of [\[Hofmann and Martell 2014, Lemma 3.61\]](#). See also [\[Azzam 2016, Lemma 4.1; Jerison and Kenig 1982, Lemma 6.3\]](#).

Hence, by Cauchy–Schwarz inequality,

$$\begin{aligned} \inf_C r^{-(n+1)} \int_{B(\xi, r) \cap \Omega} |A - C| &\lesssim \inf_C \left(r^{-(n+1)} \int_{B(\xi, r) \cap \Omega} |A - C|^2 \right)^{1/2} \\ &\leq \inf_C \left(r^{-(n+1)} \int_{\Omega_r} |A - C|^2 \right)^{1/2} \\ &\stackrel{(9-2)}{\lesssim} |\Omega_r|^{1/(2(n+1))} \left(\frac{1}{r^{n+1}} \int_{\Omega_r} |\nabla A|^2 \text{dist}(z, \Omega_r^c) dz \right)^{1/2} \\ &\lesssim \left(r^{-n} \int_{\Omega \cap B(\xi, Mr)} |\nabla A|^2 \text{dist}(z, \Omega^c) dz \right)^{1/2} \rightarrow 0 \quad \text{as } r \rightarrow 0. \end{aligned}$$

Acknowledgements

Both authors were supported by the ERC grant 320501 of the European Research Council (FP7/2007-2013). Mourgoglou was supported by Ikerbasque and partially supported by the grant MTM-2017-82160-C2-2-P of the Ministerio de Economía y Competitividad (Spain) and by IT-641-13 (Basque Government).

This project was started while we were postdocs under Tolsa's ERC grant 320501 of the European Research Council (FP7/2007-2013) between 2015 and 2016, and also during mutual visits between our home institutions of Universitat Autònoma de Barcelona (UAB), the Basque Center for Applied Mathematics, Universidad del País Vasco and the University of Edinburgh. We are grateful to these

institutions for their hospitality, to Xavier Tolsa for his helpful discussions and encouragement on the project, José Conde for his clear presentation of [Badger 2011] in our working seminar at the UAB, and to Tatiana Toro for her comments on our paper. We would like to thank the anonymous referee for reading our paper carefully and providing us with many useful comments.

References

- [Abramov and Petkovšek 2012] S. A. Abramov and M. Petkovšek, “On polynomial solutions of linear partial differential and (q) -difference equations”, pp. 1–11 in *Computer algebra in scientific computing* (Maribor, Slovenia, 2012), edited by V. P. Gerdt et al., Lecture Notes in Comput. Sci. **7442**, Springer, 2012. [MR](#) [Zbl](#)
- [Aikawa and Hirata 2008] H. Aikawa and K. Hirata, “Doubling conditions for harmonic measure in John domains”, *Ann. Inst. Fourier (Grenoble)* **58**:2 (2008), 429–445. [MR](#) [Zbl](#)
- [Akman et al. 2017] M. Akman, M. Badger, S. Hofmann, and J. M. Martell, “Rectifiability and elliptic measures on 1-sided NTA domains with Ahlfors–David regular boundaries”, *Trans. Amer. Math. Soc.* **369**:8 (2017), 5711–5745. [MR](#) [Zbl](#)
- [Akman et al. 2019] M. Akman, J. Azzam, and M. Mouroglou, “Absolute continuity of harmonic measure for domains with lower regular boundaries”, *Adv. Math.* **345** (2019), 1206–1252. [MR](#) [Zbl](#)
- [Alt et al. 1984] H. W. Alt, L. A. Caffarelli, and A. Friedman, “Variational problems with two phases and their free boundaries”, *Trans. Amer. Math. Soc.* **282**:2 (1984), 431–461. [MR](#) [Zbl](#)
- [Azzam 2016] J. Azzam, “Sets of absolute continuity for harmonic measure in NTA domains”, *Potential Anal.* **45**:3 (2016), 403–433. [MR](#) [Zbl](#)
- [Azzam and Mouroglou 2018] J. Azzam and M. Mouroglou, “Tangent measures and absolute continuity of harmonic measure”, *Rev. Mat. Iberoam.* **34**:1 (2018), 305–330. [MR](#) [Zbl](#)
- [Azzam et al. 2016a] J. Azzam, J. Garnett, M. Mouroglou, and X. Tolsa, “Uniform rectifiability, elliptic measure, square functions, and ε -approximability via an ACF monotonicity formula”, preprint, 2016. [arXiv](#)
- [Azzam et al. 2016b] J. Azzam, S. Hofmann, J. M. Martell, S. Mayboroda, M. Mouroglou, X. Tolsa, and A. Volberg, “Rectifiability of harmonic measure”, *Geom. Funct. Anal.* **26**:3 (2016), 703–728. [MR](#) [Zbl](#)
- [Azzam et al. 2016c] J. Azzam, M. Mouroglou, X. Tolsa, and A. Volberg, “On a two-phase problem for harmonic measure in general domains”, preprint, 2016. To appear in *Amer. J. Math.* [arXiv](#)
- [Azzam et al. 2017a] J. Azzam, S. Hofmann, J. M. Martell, K. Nyström, and T. Toro, “A new characterization of chord-arc domains”, *J. Eur. Math. Soc.* **19**:4 (2017), 967–981. [MR](#) [Zbl](#)
- [Azzam et al. 2017b] J. Azzam, M. Mouroglou, and X. Tolsa, “Mutual absolute continuity of interior and exterior harmonic measure implies rectifiability”, *Comm. Pure Appl. Math.* **70**:11 (2017), 2121–2163. [MR](#) [Zbl](#)
- [Azzam et al. 2017c] J. Azzam, M. Mouroglou, and X. Tolsa, “Singular sets for harmonic measure on locally flat domains with locally finite surface measure”, *Int. Math. Res. Not.* **2017**:12 (2017), 3751–3773. [MR](#) [Zbl](#)
- [Badger 2011] M. Badger, “Harmonic polynomials and tangent measures of harmonic measure”, *Rev. Mat. Iberoam.* **27**:3 (2011), 841–870. [MR](#) [Zbl](#)
- [Badger 2012] M. Badger, “Null sets of harmonic measure on NTA domains: Lipschitz approximation revisited”, *Math. Z.* **270**:1-2 (2012), 241–262. [MR](#) [Zbl](#)
- [Badger 2013] M. Badger, “Flat points in zero sets of harmonic polynomials and harmonic measure from two sides”, *J. Lond. Math. Soc. (2)* **87**:1 (2013), 111–137. [MR](#) [Zbl](#)
- [Badger and Lewis 2015] M. Badger and S. Lewis, “Local set approximation: Mattila–Vuorinen type sets, Reifenberg type sets, and tangent sets”, *Forum Math. Sigma* **3** (2015), art. id. e24. [MR](#) [Zbl](#)
- [Badger et al. 2017] M. Badger, M. Engelstein, and T. Toro, “Structure of sets which are well approximated by zero sets of harmonic polynomials”, *Anal. PDE* **10**:6 (2017), 1455–1495. [MR](#) [Zbl](#)
- [Barreira and Wolf 2006] L. Barreira and C. Wolf, “Pointwise dimension and ergodic decompositions”, *Ergodic Theory Dynam. Systems* **26**:3 (2006), 653–671. [MR](#) [Zbl](#)

- [Bishop and Jones 1990] C. J. Bishop and P. W. Jones, “Harmonic measure and arclength”, *Ann. of Math.* (2) **132**:3 (1990), 511–547. [MR](#) [Zbl](#)
- [Bishop et al. 1989] C. J. Bishop, L. Carleson, J. B. Garnett, and P. W. Jones, “Harmonic measures supported on curves”, *Pacific J. Math.* **138**:2 (1989), 233–236. [MR](#) [Zbl](#)
- [Bourgain 1987] J. Bourgain, “On the Hausdorff dimension of harmonic measure in higher dimension”, *Invent. Math.* **87**:3 (1987), 477–483. [MR](#) [Zbl](#)
- [Brezis 2011] H. Brezis, *Functional analysis, Sobolev spaces and partial differential equations*, Springer, 2011. [MR](#) [Zbl](#)
- [Caffarelli et al. 1981] L. A. Caffarelli, E. B. Fabes, and C. E. Kenig, “Completely singular elliptic-harmonic measures”, *Indiana Univ. Math. J.* **30**:6 (1981), 917–924. [MR](#) [Zbl](#)
- [Cheeger et al. 2015] J. Cheeger, A. Naber, and D. Valtorta, “Critical sets of elliptic equations”, *Comm. Pure Appl. Math.* **68**:2 (2015), 173–209. [MR](#) [Zbl](#)
- [David and Jerison 1990] G. David and D. Jerison, “Lipschitz approximation to hypersurfaces, harmonic measure, and singular integrals”, *Indiana Univ. Math. J.* **39**:3 (1990), 831–845. [MR](#) [Zbl](#)
- [Fabes et al. 1984] E. B. Fabes, D. S. Jerison, and C. E. Kenig, “Necessary and sufficient conditions for absolute continuity of elliptic-harmonic measure”, *Ann. of Math.* (2) **119**:1 (1984), 121–141. [MR](#) [Zbl](#)
- [Fefferman et al. 1991] R. A. Fefferman, C. E. Kenig, and J. Pipher, “The theory of weights and the Dirichlet problem for elliptic equations”, *Ann. of Math.* (2) **134**:1 (1991), 65–124. [MR](#) [Zbl](#)
- [García-Cuerva and Rubio de Francia 1985] J. García-Cuerva and J. L. Rubio de Francia, *Weighted norm inequalities and related topics*, North-Holland Math. Studies **116**, North-Holland, Amsterdam, 1985. [MR](#) [Zbl](#)
- [Garnett 2007] J. B. Garnett, *Bounded analytic functions*, Graduate Texts in Math. **236**, Springer, 2007. [MR](#) [Zbl](#)
- [Garnett and Marshall 2005] J. B. Garnett and D. E. Marshall, *Harmonic measure*, New Math. Monographs **2**, Cambridge Univ. Press, 2005. [MR](#) [Zbl](#)
- [Grüter and Widman 1982] M. Grüter and K.-O. Widman, “The Green function for uniformly elliptic equations”, *Manuscripta Math.* **37**:3 (1982), 303–342. [MR](#) [Zbl](#)
- [Heinonen et al. 1993] J. Heinonen, T. Kilpeläinen, and O. Martio, *Nonlinear potential theory of degenerate elliptic equations*, Oxford Univ. Press, 1993. [MR](#) [Zbl](#)
- [Hofmann and Kim 2007] S. Hofmann and S. Kim, “The Green function estimates for strongly elliptic systems of second order”, *Manuscripta Math.* **124**:2 (2007), 139–172. [MR](#) [Zbl](#)
- [Hofmann and Martell 2014] S. Hofmann and J. M. Martell, “Uniform rectifiability and harmonic measure, I: Uniform rectifiability implies Poisson kernels in L^p ”, *Ann. Sci. Éc. Norm. Supér.* (4) **47**:3 (2014), 577–654. [MR](#) [Zbl](#)
- [Hofmann et al. 2014] S. Hofmann, J. M. Martell, and I. Uriarte-Tuero, “Uniform rectifiability and harmonic measure, II: Poisson kernels in L^p imply uniform rectifiability”, *Duke Math. J.* **163**:8 (2014), 1601–1654. [MR](#) [Zbl](#)
- [Hofmann et al. 2017] S. Hofmann, J. M. Martell, and T. Toro, “ A_∞ implies NTA for a class of variable coefficient elliptic operators”, *J. Differential Equations* **263**:10 (2017), 6147–6188. [MR](#) [Zbl](#)
- [Hurri-Syrjänen 1994] R. Hurri-Syrjänen, “An improved Poincaré inequality”, *Proc. Amer. Math. Soc.* **120**:1 (1994), 213–222. [MR](#) [Zbl](#)
- [Jerison and Kenig 1982] D. S. Jerison and C. E. Kenig, “Boundary behavior of harmonic functions in nontangentially accessible domains”, *Adv. in Math.* **46**:1 (1982), 80–147. [MR](#) [Zbl](#)
- [Kenig and Pipher 2001] C. E. Kenig and J. Pipher, “The Dirichlet problem for elliptic equations with drift terms”, *Publ. Mat.* **45**:1 (2001), 199–217. [MR](#) [Zbl](#)
- [Kenig and Toro 2006] C. Kenig and T. Toro, “Free boundary regularity below the continuous threshold: 2-phase problems”, *J. Reine Angew. Math.* **596** (2006), 1–44. [MR](#) [Zbl](#)
- [Kenig et al. 2009] C. Kenig, D. Preiss, and T. Toro, “Boundary structure and size in terms of interior and exterior harmonic measures in higher dimensions”, *J. Amer. Math. Soc.* **22**:3 (2009), 771–796. [MR](#) [Zbl](#)
- [Khrushchev 1984] S. V. Hruščev, “A description of weights satisfying the A_∞ condition of Muckenhoupt”, *Proc. Amer. Math. Soc.* **90**:2 (1984), 253–257. [MR](#) [Zbl](#)

- [Korey 1998] M. B. Korey, “Ideal weights: asymptotically optimal versions of doubling, absolute continuity, and bounded mean oscillation”, *J. Fourier Anal. Appl.* **4**:4-5 (1998), 491–519. [MR](#) [Zbl](#)
- [Krantz and Parks 1992] S. G. Krantz and H. R. Parks, *A primer of real analytic functions*, Basler Lehrbücher **4**, Birkhäuser, Basel, 1992. [MR](#) [Zbl](#)
- [Martio 1979] O. Martio, “Capacity and measure densities”, *Ann. Acad. Sci. Fenn. Ser. A I Math.* **4**:1 (1979), 109–118. [MR](#) [Zbl](#)
- [Mattila 1995] P. Mattila, *Geometry of sets and measures in Euclidean spaces*, Cambridge Studies in Adv. Math. **44**, Cambridge Univ. Press, 1995. [MR](#) [Zbl](#)
- [Mitrea 2013] D. Mitrea, *Distributions, partial differential equations, and harmonic analysis*, Springer, 2013. [MR](#) [Zbl](#)
- [Mourgoglou and Tolsa 2017] M. Mourgoglou and X. Tolsa, “Harmonic measure and Riesz transform in uniform and general domains”, *J. Reine Angew. Math.* (online publication October 2017).
- [Pommerenke 1986] C. Pommerenke, “On conformal mapping and linear measure”, *J. Analyse Math.* **46** (1986), 231–238. [MR](#) [Zbl](#)
- [Preiss 1987] D. Preiss, “Geometry of measures in \mathbb{R}^n : distribution, rectifiability, and densities”, *Ann. of Math. (2)* **125**:3 (1987), 537–643. [MR](#) [Zbl](#)
- [Reimann and Rychener 1975] H. M. Reimann and T. Rychener, *Funktionen beschränkter mittlerer Oszillation*, Lecture Notes in Math. **487**, Springer, 1975. [MR](#) [Zbl](#)
- [Riesz and Riesz 1920] F. Riesz and M. Riesz, “Über die Randwerte einer analytischen Funktion”, pp. 27–44 in *Quatrième Congrès des Mathématiciens Scandinaves* (Stockholm, 1916), edited by M. G. Mittag-Leffler, Almqvist & Wiksell, Uppsala, 1920. [JFM](#)
- [Sweezy 1992] C. Sweezy, “The Hausdorff dimension of elliptic measure: a counterexample to the Oksendahl conjecture in \mathbb{R}^2 ”, *Proc. Amer. Math. Soc.* **116**:2 (1992), 361–368. [MR](#) [Zbl](#)
- [Tolsa and Volberg 2018] X. Tolsa and A. Volberg, “On Tsirelson’s theorem about triple points for harmonic measure”, *Int. Math. Res. Not.* **2018**:12 (2018), 3671–3683. [MR](#)
- [Toro and Zhao 2017] T. Toro and Z. Zhao, “Boundary rectifiability and elliptic operators with $W^{1,1}$ coefficients”, *Adv. Calc. Var.* (online publication October 2017).
- [Wu 1986] J.-M. Wu, “On singularity of harmonic measure in space”, *Pacific J. Math.* **121**:2 (1986), 485–496. [MR](#) [Zbl](#)
- [Wu 1994] J.-M. Wu, “Porous sets and null sets for elliptic harmonic measures”, *Trans. Amer. Math. Soc.* **346**:2 (1994), 455–473. [MR](#) [Zbl](#)
- [Zierner 1974] W. P. Zierner, “Some remarks on harmonic measure in space”, *Pacific J. Math.* **55** (1974), 629–637. [MR](#) [Zbl](#)

Received 15 Aug 2017. Revised 19 Oct 2018. Accepted 30 Nov 2018.

JONAS AZZAM: j.azzam@ed.ac.uk

School of Mathematics, University of Edinburgh, Edinburgh, United Kingdom

MIHALIS MOURGOLOU: michail.mourgoglou@ehu.eus

Departamento de Matemáticas, Universidad del País Vasco, Leioa, Spain

and

Ikerbasque, Basque Foundation for Science, Bilbao, Spain

Analysis & PDE

msp.org/apde

EDITORS

EDITOR-IN-CHIEF

Patrick Gérard

patrick.gerard@math.u-psud.fr

Université Paris Sud XI

Orsay, France

BOARD OF EDITORS

Massimiliano Berti	Scuola Intern. Sup. di Studi Avanzati, Italy berti@sissa.it	Gilles Pisier	Texas A&M University, and Paris 6 pisier@math.tamu.edu
Michael Christ	University of California, Berkeley, USA mchrist@math.berkeley.edu	Tristan Rivière	ETH, Switzerland riviere@math.ethz.ch
Charles Fefferman	Princeton University, USA cf@math.princeton.edu	Igor Rodnianski	Princeton University, USA irod@math.princeton.edu
Ursula Hamenstaedt	Universität Bonn, Germany ursula@math.uni-bonn.de	Yum-Tong Siu	Harvard University, USA siu@math.harvard.edu
Vadim Kaloshin	University of Maryland, USA vadim.kaloshin@gmail.com	Terence Tao	University of California, Los Angeles, USA tao@math.ucla.edu
Herbert Koch	Universität Bonn, Germany koch@math.uni-bonn.de	Michael E. Taylor	Univ. of North Carolina, Chapel Hill, USA met@math.unc.edu
Izabella Laba	University of British Columbia, Canada ilaba@math.ubc.ca	Gunther Uhlmann	University of Washington, USA gunther@math.washington.edu
Richard B. Melrose	Massachusetts Inst. of Tech., USA rhm@math.mit.edu	András Vasy	Stanford University, USA andras@math.stanford.edu
Frank Merle	Université de Cergy-Pontoise, France Frank.Merle@u-cergy.fr	Dan Virgil Voiculescu	University of California, Berkeley, USA dvv@math.berkeley.edu
William Minicozzi II	Johns Hopkins University, USA minicozz@math.jhu.edu	Steven Zelditch	Northwestern University, USA zelditch@math.northwestern.edu
Clément Mouhot	Cambridge University, UK c.mouhot@dpmms.cam.ac.uk	Maciej Zworski	University of California, Berkeley, USA zworski@math.berkeley.edu
Werner Müller	Universität Bonn, Germany mueller@math.uni-bonn.de		

PRODUCTION

production@msp.org

Silvio Levy, Scientific Editor

See inside back cover or msp.org/apde for submission instructions.

The subscription price for 2019 is US \$310/year for the electronic version, and \$520/year (+\$60, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues from the last three years and changes of subscriber address should be sent to MSP.

Analysis & PDE (ISSN 1948-206X electronic, 2157-5045 printed) at Mathematical Sciences Publishers, 798 Evans Hall #3840, c/o University of California, Berkeley, CA 94720-3840, is published continuously online. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices.

APDE peer review and production are managed by EditFlow[®] from MSP.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2019 Mathematical Sciences Publishers

ANALYSIS & PDE

Volume 12 No. 8 2019

Tangent measures of elliptic measure and applications	1891
JONAS AZZAM and MIHALIS MOURGOGLOU	
Discretely self-similar solutions to the Navier–Stokes equations with data in L^2_{loc} satisfying the local energy inequality	1943
ZACHARY BRADSHAW and TAI-PENG TSAI	
Continuity properties for divergence form boundary data homogenization problems	1963
WILLIAM M. FELDMAN and YUMING PAUL ZHANG	
Dynamics of one-fold symmetric patches for the aggregation equation and collapse to singular measure	2003
TAOUFIK HMIDI and DONG LI	
Coupled Kähler–Ricci solitons on toric Fano manifolds	2067
JAKOB HULTGREN	
Carleson measure estimates and the Dirichlet problem for degenerate elliptic equations	2095
STEVE HOFMANN, PHI LE and ANDREW J. MORRIS	