

ANALYSIS & PDE

Volume 12 No. 8 2019

STEVE HOFMANN, PHI LE AND ANDREW J. MORRIS

**CARLESON MEASURE ESTIMATES AND THE DIRICHLET
PROBLEM
FOR DEGENERATE ELLIPTIC EQUATIONS**

CARLESON MEASURE ESTIMATES AND THE DIRICHLET PROBLEM FOR DEGENERATE ELLIPTIC EQUATIONS

STEVE HOFMANN, PHI LE AND ANDREW J. MORRIS

We prove that the Dirichlet problem for degenerate elliptic equations $\operatorname{div}(A\nabla u) = 0$ in the upper half-space $(x, t) \in \mathbb{R}_+^{n+1}$ is solvable when $n \geq 2$ and the boundary data is in $L_\mu^p(\mathbb{R}^n)$ for some $p < \infty$. The coefficient matrix A is only assumed to be measurable, real-valued and t -independent with a degenerate bound and ellipticity controlled by an A_2 -weight μ . It is not required to be symmetric. The result is achieved by proving a Carleson measure estimate for all bounded solutions in order to deduce that the degenerate elliptic measure is in A_∞ with respect to the μ -weighted Lebesgue measure on \mathbb{R}^n . The Carleson measure estimate allows us to avoid applying the method of ϵ -approximability, which simplifies the proof obtained recently in the case of uniformly elliptic coefficients. The results have natural extensions to Lipschitz domains.

1. Introduction	2095
2. Preliminaries	2098
3. Estimates for maximal operators	2105
4. The Carleson measure estimate	2110
5. Solvability of the Dirichlet problem	2129
References	2144

1. Introduction

We consider the Dirichlet boundary value problem for the degenerate elliptic equation $\operatorname{div}(A\nabla u) = 0$ in the upper half-space \mathbb{R}_+^{n+1} when $n \geq 2$ and which we make precise below. The boundary $\mathbb{R}^n \times \{0\}$ is identified with \mathbb{R}^n and we adopt the notation $X = (x, t)$ for points $X \in \mathbb{R}_+^{n+1}$ with coordinates $x \in \mathbb{R}^n$ and $t \in (0, \infty)$. The gradient $\nabla := (\nabla_x, \partial_t)$ and divergence $\operatorname{div} := \operatorname{div}_x + \partial_t$ are with respect to all $(n+1)$ -coordinates. The coefficient A denotes an $(n+1) \times (n+1)$ matrix of measurable, real-valued and t -independent functions on \mathbb{R}_+^{n+1} . The matrix $A(x) := A(x, t)$ is not required to be symmetric. We suppose that there exist constants $0 < \lambda \leq \Lambda < \infty$ and an A_2 -weight μ on \mathbb{R}^n such that the degenerate bound and ellipticity

$$|\langle A(x)\xi, \zeta \rangle| \leq \Lambda \mu(x) |\xi| |\zeta| \quad \text{and} \quad \langle A(x)\xi, \xi \rangle \geq \lambda \mu(x) |\xi|^2 \quad (1.1)$$

hold for all $\xi, \zeta \in \mathbb{R}^{n+1}$ and almost every $x \in \mathbb{R}^n$. We use $\langle \cdot, \cdot \rangle$ and $|\cdot|$ to denote the Euclidean inner product and norm. An A_2 -weight μ on \mathbb{R}^n refers to a nonnegative locally integrable function

MSC2010: 35J25, 35J70, 42B20, 42B25.

Keywords: square functions, nontangential maximal functions, harmonic measure, Radon–Nikodym derivative, Carleson measure, divergence form elliptic equations, Dirichlet problem, A_2 Muckenhoupt weights, reverse Hölder inequality.

$\mu : \mathbb{R}^n \rightarrow [0, \infty]$ such that

$$[\mu]_{A_2(\mathbb{R}^n)} := \sup_Q \left(\frac{1}{|Q|} \int_Q \mu(x) dx \right) \left(\frac{1}{|Q|} \int_Q \frac{1}{\mu(x)} dx \right) < \infty,$$

where \sup_Q denotes the supremum over all cubes Q in \mathbb{R}^n with volume $|Q|$. We also use μ to denote the measure $\mu(Q) := \int_Q \mu(x) dx$ and consider the Lebesgue space $L_\mu^p(\mathbb{R}^n)$ with the norm $\|f\|_{L_\mu^p(\mathbb{R}^n)} := (\int_{\mathbb{R}^n} |f|^p d\mu)^{1/p}$ for all $p \in [1, \infty)$. There is also the notation $f_Q f d\mu := \mu(Q)^{-1} \int_Q f d\mu$, whilst $f_Q f := |Q|^{-1} \int_Q f(x) dx$.

If μ is identically 1, then A is called uniformly elliptic. The solvability of the Dirichlet problem for general nonsymmetric coefficients in that case was obtained only recently by Hofmann, Kenig, Mayboroda and Pipher [Hofmann et al. 2015a]. The result in dimension $n = 1$ had been obtained previously by Kenig, Koch, Pipher and Toro [Kenig et al. 2000]. These results assert that for each uniformly elliptic coefficient matrix A there exists some $p < \infty$ for which the Dirichlet problem is solvable for L^p -boundary data. Conversely, counterexamples in [Kenig et al. 2000] show that for each $p < \infty$, there exists a uniformly elliptic coefficient matrix A for which the Dirichlet problem is not solvable for L^p -boundary data. In contrast, solvability of the Dirichlet problem for symmetric coefficients in the uniformly elliptic case is well understood, and we mention only that it was obtained by Jerison and Kenig [1981] for L^p -boundary data when $2 \leq p < \infty$.

The solvability of the Dirichlet problem in the uniformly elliptic case has also been established for a variety of complex coefficient structures; see, for instance, [Auscher and Stahlhut 2014; Hofmann et al. 2015a; 2015b]. A significant portion of that theory was recently extended to the degenerate elliptic case by Auscher, Rosén and Rule [Auscher et al. 2015] for L^2 -boundary data. That extension did not include, however, the results for general nonsymmetric coefficients in [Hofmann et al. 2015a]. This paper complements the progress made in [Auscher et al. 2015] by extending the solvability obtained for the Dirichlet problem in [Hofmann et al. 2015a] to the degenerate elliptic case.

For solvability on the upper half-space \mathbb{R}_+^{n+1} , the A_2 -weight μ on \mathbb{R}^n is extended to the t -independent A_2 -weight $\mu(x, t) := \mu(x)$ on \mathbb{R}^{n+1} (and $[\mu]_{A_2(\mathbb{R}^{n+1})} = [\mu]_{A_2(\mathbb{R}^n)}$). We then say that u is a *solution* of the equation $\operatorname{div}(A \nabla u) = 0$ in an open set $\Omega \subseteq \mathbb{R}^{n+1}$ when $u \in W_{\mu, \text{loc}}^{1,2}(\Omega)$ and $\int_{\mathbb{R}_+^{n+1}} \langle A \nabla u, \nabla \Phi \rangle = 0$ for all smooth compactly supported functions $\Phi \in C_c^\infty(\Omega)$. The solution space is the local μ -weighted Sobolev space $W_{\mu, \text{loc}}^{1,2}$ defined in Section 2. The convergence of solutions to boundary data is afforded by estimates for the nontangential maximal function $N_* u$ of solutions u , defined by

$$(N_* u)(x) := \sup_{(y,t) \in \Gamma(x)} |u(y,t)| \quad \text{for all } x \in \mathbb{R}^n,$$

where $\Gamma(x)$ is the cone $\{(y,t) \in \mathbb{R}_+^{n+1} : |y-x| < t\}$. If $p \in (1, \infty)$, then the Dirichlet problem for $L_\mu^p(\mathbb{R}^n)$ -boundary data, or simply $(D)_{p,\mu}$, is said to be *solvable* when for each $f \in L_\mu^p(\mathbb{R}^n)$ there exists a solution u such that

$$\begin{cases} \operatorname{div}(A \nabla u) = 0 & \text{in } \mathbb{R}_+^{n+1}, \\ N_* u \in L_\mu^p(\mathbb{R}^n), \\ \lim_{t \rightarrow 0} u(\cdot, t) = f, \end{cases} \quad (D)_{p,\mu}$$

where the limit is required to converge in $L^p_\mu(\mathbb{R}^n)$ -norm and in the nontangential sense whereby $\lim_{\Gamma(x) \ni (y,t) \rightarrow (x,0)} u(y,t) = f(x)$ for almost every $x \in \mathbb{R}^n$. Note that this definition of solvability is distinct from *well-posedness*, which requires that such solutions are unique. We are able to obtain a uniqueness result for solutions that converge uniformly to 0 at infinity, but the question of well-posedness more generally remains open (see Theorem 5.34 and the preceding discussion).

A nonnegative Borel measure ω on a cube Q_0 in \mathbb{R}^n is said to be in the A_∞ -class with respect to μ , written $\omega \in A_\infty(\mu)$, when there exist constants $C, \theta > 0$, which we call the $A_\infty(Q_0)$ -constants, such that

$$\omega(E) \leq C \left(\frac{\mu(E)}{\mu(Q)} \right)^\theta \omega(Q)$$

for all cubes $Q \subseteq Q_0$ and all Borel sets $E \subseteq Q$. This is a scale-invariant version of the absolute continuity of ω with respect to μ . It is well known, at least in the uniformly elliptic case, that solvability of the Dirichlet problem for L^p -boundary data for some $p < \infty$ is equivalent to the property that an adapted harmonic measure (elliptic measure) belongs to A_∞ with respect to the Lebesgue measure on \mathbb{R}^n ; see Theorem 1.7.3 in [Kenig 1994]. In the degenerate case, an adapted harmonic measure ω^X , which we call degenerate elliptic measure, can also be defined at each $X \in \mathbb{R}_+^{n+1}$ (see Section 5). We prove that this degenerate elliptic measure is in A_∞ with respect to μ and then deduce the solvability of $(D)_{p,\mu}$ stated in the theorem below. This requires the notation associated with cubes Q in \mathbb{R}^n , where x_Q and $\ell(Q)$ denote the centre and side length of Q , respectively, and $X_Q := (x_Q, \ell(Q))$ denotes the corkscrew point in \mathbb{R}_+^{n+1} relative to Q .

Theorem 1.2. *If $n \geq 2$ and the t -independent coefficient matrix A satisfies the degenerate bound and ellipticity in (1.1) for some constants $0 < \lambda \leq \Lambda < \infty$ and an A_2 -weight μ on \mathbb{R}^n , then there exists $p \in (1, \infty)$ such that $(D)_{p,\mu}$ is solvable. Moreover, on each cube Q in \mathbb{R}^n , the degenerate elliptic measure $\omega := \omega^{X_Q} \llcorner Q$ satisfies $\omega \in A_\infty(\mu)$ with $A_\infty(Q)$ -constants that depend only on n, λ, Λ and $[\mu]_{A_2}$.*

In contrast to the proof of solvability in the uniformly elliptic case in [Hofmann et al. 2015a], we avoid the need to apply the method of ϵ -approximability by first establishing the Carleson measure estimate in the theorem below. This crucial estimate facilitates the main results of the paper. The connection between the Carleson measure estimate and solvability was first established in the uniformly elliptic case by Kenig, Kirchheim, Pipher and Toro [Kenig et al. 2016], and we follow their approach here, adapting it to the degenerate elliptic setting (see Lemma 5.24 below). In particular, the A_∞ -property of degenerate elliptic measure is obtained by combining the Carleson measure estimate (1.4) with the notion of good ϵ -coverings introduced in [Kenig et al. 2000].

Theorem 1.3. *If $n \geq 2$ and the t -independent coefficient matrix A satisfies the degenerate bound and ellipticity in (1.1) for some constants $0 < \lambda \leq \Lambda < \infty$ and an A_2 -weight μ on \mathbb{R}^n , then any solution $u \in L^\infty(\mathbb{R}_+^{n+1})$ of $\operatorname{div}(A \nabla u) = 0$ in \mathbb{R}_+^{n+1} satisfies the Carleson measure estimate*

$$\sup_Q \frac{1}{\mu(Q)} \int_0^{\ell(Q)} \int_Q |t \nabla u(x, t)|^2 d\mu(x) \frac{dt}{t} \leq C \|u\|_\infty^2, \quad (1.4)$$

where C depends only on n, λ, Λ and $[\mu]_{A_2}$.

Using the Carleson measure estimate in this way allows us to bypass the need to establish norm-equivalences between the nontangential maximal function N_*u and the square function Su of solutions u , defined by

$$(Su)(x) := \left(\iint_{\Gamma(x)} |t \nabla u(y, t)|^2 \frac{d\mu(y)}{\mu(\Delta(x, t))} \frac{dt}{t} \right)^{1/2} \quad \text{for all } x \in \mathbb{R}^n,$$

where $\Delta(x, t)$ is the surface ball $\{y \in \mathbb{R}^n : |y - x| < t\}$. It was shown by Dahlberg, Jerison and Kenig [Dahlberg et al. 1984], however, that such estimates are a consequence of the A_∞ -property of degenerate elliptic measure, which provides the following result.

Theorem 1.5. *If $n \geq 2$ and the t -independent coefficient matrix A satisfies the degenerate bound and ellipticity in (1.1) for some constants $0 < \lambda \leq \Lambda < \infty$ and an A_2 -weight μ on \mathbb{R}^n , then any solution of $\operatorname{div}(A \nabla u) = 0$ in \mathbb{R}_+^{n+1} satisfies*

$$\|Su\|_{L_\mu^p(\mathbb{R}^n)} \leq C \|N_*u\|_{L_\mu^p(\mathbb{R}^n)} \quad \text{for all } p \in (0, \infty),$$

and if, in addition, $u(X_0) = 0$ for some $X_0 \in \mathbb{R}_+^{n+1}$, then

$$\|N_*u\|_{L_\mu^p(\mathbb{R}^n)} \leq C \|Su\|_{L_\mu^p(\mathbb{R}^n)} \quad \text{for all } p \in (0, \infty),$$

where C depends only on X_0 , p , n , λ , Λ and $[\mu]_{A_2}$.

The paper is structured as follows. Technical preliminaries concerning weights and degenerate elliptic operators are in Section 2, whilst estimates for weighted maximal operators are in Section 3. The Carleson measure estimate in Theorem 1.3 is obtained in Section 4. The degenerate elliptic measure is constructed in Section 5 and then the A_∞ -estimates in Theorem 1.2 are deduced as part of Theorem 5.30. The square function and nontangential maximal function estimates in Theorem 1.5 are included in the more general result in Theorem 5.31, whilst the solvability of the Dirichlet problem in Theorem 1.2 is finally deduced in Theorem 5.34, where a uniqueness result is also obtained.

We state and prove our results in the upper half-space, but we note that they extend immediately to the case that the domain is the region above a Lipschitz graph, by a well-known pull-back technique which preserves the t -independence of the coefficients. In turn, our results concerning the A_∞ -property of degenerate elliptic measure may then be extended to the case of a bounded star-like Lipschitz domain, with radially independent coefficients, by a standard localization argument using the maximum principle.

The convention is adopted whereby C denotes a finite positive constant that may change from one line to the next. For $a, b \in \mathbb{R}$, the notation $a \lesssim b$ means that $a \leq Cb$, whilst $a \asymp b$ means that $a \lesssim b \lesssim a$. We write $a \lesssim_p b$ when $a \leq Cb$ and we wish to emphasise that C depends on a specified parameter p .

2. Preliminaries

We dispense with some technical preliminaries concerning general A_p -weights μ for $p \in (1, \infty)$ and degenerate elliptic operators on \mathbb{R}^n for $n \in \mathbb{N}$. All cubes Q and balls B in \mathbb{R}^n are assumed to be open (except in Section 5D where the standard dyadic cubes S in $\mathbb{D}(\mathbb{R}^n)$ are assumed to be closed to provide genuine coverings of \mathbb{R}^n). For $\alpha > 0$, let αQ and αB denote the concentric dilates of Q

and B respectively. For $x \in \mathbb{R}^n$ and $r > 0$, define the ball $B(x, r) := \{y \in \mathbb{R}^n : |y - x| < r\}$. An A_p -weight refers to a nonnegative locally integrable function μ on \mathbb{R}^n with the property that $[\mu]_{A_p(\mathbb{R}^n)} := \sup_Q (f_Q \mu)(f_Q \mu^{-1/(p-1)})^{p-1} < \infty$. The measure associated with such a weight satisfies the doubling property

$$\mu(\alpha B) \leq [\mu]_{A_p} \alpha^{np} \mu(B) \quad (2.1)$$

for all $\alpha \geq 1$; see, for instance, Section 1.5 in Chapter V of [Stein 1993].

For an open set $\Omega \subseteq \mathbb{R}^n$, the Sobolev space $W_\mu^{1,p}(\Omega)$ is defined as the completion, in the ambient space $L_\mu^p(\Omega)$, of the normed space of all $f \in C^\infty(\Omega)$ with finite norm

$$\|f\|_{W_\mu^{1,p}(\Omega)}^p := \int_\Omega |f|^p d\mu + \int_\Omega |\nabla f|^p d\mu < \infty. \quad (2.2)$$

The embedding of the completion $W_\mu^{1,p}(\Omega)$ in $L_\mu^p(\Omega)$ relies on the A_p -property of the weight (to the extent that it implies both μ and $\mu^{-1/(p-1)}$ are in $L_{\text{loc}}^1(\Omega)$), which ensures that if $(f_j)_j$ is a $W_\mu^{1,p}(\Omega)$ -Cauchy sequence in $C^\infty(\Omega)$ converging to 0 in $L_\mu^p(\Omega)$, then $(f_j)_j$ converges to 0 in $W_\mu^{1,p}(\Omega)$ -norm; see Section 2.1 in [Fabes et al. 1982b]. Therefore, since $C^\infty(\Omega)$ is dense in $W_\mu^{1,p}(\Omega)$, the gradient extends to a bounded operator $\nabla : W_\mu^{1,p}(\Omega) \rightarrow L_\mu^p(\Omega, \mathbb{R}^n)$, thereby extending (2.2) to all $f \in W_\mu^{1,p}(\Omega)$. The Sobolev space $W_{0,\mu}^{1,p}(\Omega)$ is defined as the closure of $C_c^\infty(\Omega)$ in $W_\mu^{1,p}(\Omega)$. It can be shown that $W_{0,\mu}^{1,p}(\mathbb{R}^n) = W_\mu^{1,p}(\mathbb{R}^n)$ by following the proof in the unweighted case from Proposition 1 of Chapter V in [Stein 1970] but instead using Lemma 2.2 in [Auscher et al. 2015] to deduce the convergence of the regularization in $L_\mu^p(\mathbb{R}^n)$. The local space $W_{\mu,\text{loc}}^{1,p}(\Omega)$ is then defined as the set of all $f \in L_{\mu,\text{loc}}^p(\Omega)$ such that $f \in W_\mu^{1,p}(\Omega')$ for all open sets Ω' with compact closure $\bar{\Omega}' \subset \Omega$ (henceforth denoted by $\Omega' \Subset \Omega$). Finally, the weighted Sobolev and Poincaré inequalities obtained for continuous functions in Theorems 1.2 and 1.5 in [Fabes et al. 1982b] have the following immediate extensions.

Theorem 2.3. *Let $n \geq 2$ and suppose that $B \subset \mathbb{R}^n$ denotes a ball with radius $r(B)$. If $p \in (1, \infty)$ and μ is an A_p -weight on \mathbb{R}^n , then there exists $\delta > 0$ such that*

$$\left(\int_B |f|^{p(\frac{n}{n-1} + \delta)} d\mu \right)^{1/(p(\frac{n}{n-1} + \delta))} \lesssim r(B) \left(\int_B |\nabla f|^p d\mu \right)^{1/p} \quad (2.4)$$

for all $f \in W_{0,\mu}^{1,p}(B)$, and

$$\left(\int_B |f(x) - c_B|^p d\mu \right)^{1/p} \lesssim r(B) \left(\int_B |\nabla f|^p d\mu \right)^{1/p} \quad (2.5)$$

for all $f \in W_\mu^{1,p}(B)$ and $c_B \in \{ \int_B f d\mu, \int_B f \}$, where the implicit constants depend only on n , p and $[\mu]_{A_p}$. The estimates also hold when the ball B and the radius $r(B)$ are replaced by a cube Q and the side length $\ell(Q)$.

For $n \in \mathbb{N}$, constants $0 < \lambda \leq \Lambda < \infty$ and an A_2 -weight μ on \mathbb{R}^n , let $\mathcal{E}(n, \lambda, \Lambda, \mu)$ denote the set of all $n \times n$ matrices \mathcal{A} of measurable real-valued functions on \mathbb{R}^n satisfying the degenerate bound and

ellipticity

$$|\langle \mathcal{A}(x)\xi, \zeta \rangle| \leq \Lambda \mu(x) |\xi| |\zeta| \quad \text{and} \quad \langle \mathcal{A}(x)\xi, \xi \rangle \geq \lambda \mu(x) |\xi|^2 \quad (2.6)$$

for all $\xi, \zeta \in \mathbb{R}^n$ and almost every $x \in \mathbb{R}^n$. These properties allow us to define

$$\mathcal{L}_{\mu, \Omega} : \text{Dom}(\mathcal{L}_{\mu, \Omega}) \subseteq L^2_{\mu}(\Omega) \rightarrow L^2_{\mu}(\Omega)$$

as the maximal accretive operator in $L^2_{\mu}(\Omega)$ associated with the bilinear form defined by

$$\mathfrak{a}_{\Omega}(f, g) := \int_{\Omega} \langle \mathcal{A} \nabla f, \nabla g \rangle = \int_{\Omega} \left\langle \frac{1}{\mu} \mathcal{A} \nabla f, \nabla g \right\rangle d\mu \quad (2.7)$$

for all $f, g \in W^{1,2}_{0,\mu}(\Omega)$. The domain of $\mathcal{L}_{\mu, \Omega}$ is dense in $L^2_{\mu}(\Omega)$, and in particular

$$\text{Dom}(\mathcal{L}_{\mu, \Omega}) = \left\{ f \in W^{1,2}_{0,\mu}(\Omega) : \sup_{g \in C^{\infty}_c(\Omega)} \frac{|\mathfrak{a}_{\Omega}(f, g)|}{\|g\|_{L^2_{\mu}(\Omega)}} < \infty \right\},$$

with

$$\int_{\Omega} (\mathcal{L}_{\mu, \Omega} f) g d\mu = \mathfrak{a}_{\Omega}(f, g) \quad (2.8)$$

for all $f \in \text{Dom}(\mathcal{L}_{\mu, \Omega})$ and $g \in W^{1,2}_{0,\mu}(\Omega)$. It is equivalent to define $\mathcal{L}_{\mu, \Omega}$ as the composition $-\text{div}_{\mu, \Omega}((1/\mu)\mathcal{A}\nabla)$ of unbounded operators, where $-\text{div}_{\mu, \Omega}$ is the adjoint ∇^* of the closed densely defined operator $\nabla : W^{1,2}_{0,\mu}(\Omega) \subseteq L^2_{\mu}(\Omega) \rightarrow L^2_{\mu}(\Omega, \mathbb{R}^n)$, that is,

$$\int_{\Omega} (-\text{div}_{\mu, \Omega} f) g d\mu = \int_{\Omega} \langle f, \nabla g \rangle d\mu \quad (2.9)$$

for all $f \in \text{Dom}(\text{div}_{\mu, \Omega}) := \text{Dom}(\nabla^*)$ and $g \in W^{1,2}_{0,\mu}(\Omega)$. In view of (2.7) and (2.8), we have the formal identities $\text{div}_{\mu, \Omega} = (1/\mu) \text{div}_{\Omega} \mu$ and $\mathcal{L}_{\mu, \Omega} = -(1/\mu) \text{div}_{\Omega}(\mathcal{A}\nabla)$.

Now let $\Omega = Q$ for some cube $Q \subset \mathbb{R}^n$ and denote the space of bounded linear functionals on $W^{1,2}_{0,\mu}(Q)$ by $W^{-1,2}_{0,\mu}(Q)$. The inclusions $W^{1,2}_{0,\mu}(Q) \subseteq L^2_{\mu}(Q) \subseteq W^{-1,2}_{0,\mu}(Q)$ are interpreted in the standard way by identifying $f \in L^2_{\mu}(Q)$ with the functional ℓ_f defined by $\ell_f(g) := \int_Q f g d\mu$ for all $g \in W^{1,2}_{0,\mu}(Q)$. Thus, setting

$$\mathcal{L}_{\mu, Q} f(g) := \mathfrak{a}_Q(f, g) \quad \text{and} \quad -\text{div}_{\mu, Q} f(g) := \int_Q \langle f, \nabla g \rangle d\mu$$

for all $f, g \in W^{1,2}_{0,\mu}(Q)$ and $f \in L^2(Q, \mathbb{R}^n)$, we obtain an extension of $\mathcal{L}_{\mu, Q}$ from (2.8) to a bounded invertible operator from $W^{1,2}_{0,\mu}(Q)$ onto $W^{-1,2}_{0,\mu}(Q)$, and an extension of $\text{div}_{\mu, Q}$ from (2.9) to a bounded operator from $L^2_{\mu}(Q)$ into $W^{-1,2}_{0,\mu}(Q)$. The surjectivity of $\mathcal{L}_{\mu, Q}$ relies on (2.4) and the Lax–Milgram theorem. These definitions imply

$$\|\nabla \mathcal{L}_{\mu, Q}^{-1} \text{div}_{\mu, Q} f\|_{L^2_{\mu}(Q, \mathbb{R}^n)} \lesssim \|f\|_{L^2_{\mu}(Q, \mathbb{R}^n)}$$

for all $f \in L^2_{\mu}(Q, \mathbb{R}^n)$. The topological direct sum or $W^{1,2}_{0,\mu}(Q)$ -Hodge decomposition

$$L^2_{\mu}(Q, \mathbb{R}^n) = \left\{ \frac{1}{\mu} \mathcal{A} \nabla g : g \in W^{1,2}_{0,\mu}(Q) \right\} \oplus \{h \in L^2_{\mu}(Q, \mathbb{R}^n) : \text{div}_{\mu, Q} h = 0\} \quad (2.10)$$

follows by writing

$$f = -\frac{1}{\mu} \mathcal{A} \nabla \mathcal{L}_{\mu, Q}^{-1} \operatorname{div}_{\mu, Q} f + \left(f + \frac{1}{\mu} \mathcal{A} \nabla \mathcal{L}_{\mu, Q}^{-1} \operatorname{div}_{\mu, Q} f \right) =: \frac{1}{\mu} \mathcal{A} \nabla g + h,$$

since then $\operatorname{div}_{\mu, Q} h = \operatorname{div}_{\mu, Q} f - \mathcal{L}_{\mu, Q} \mathcal{L}_{\mu, Q}^{-1} \operatorname{div}_{\mu, Q} f = 0$. This decomposition also extends to $L_{\mu}^p(Q, \mathbb{R}^n)$ for all $p \in [2, 2 + \epsilon)$ and some $\epsilon > 0$ by recent work of Le [2015], although we do not need it here.

Now let $\Omega = \mathbb{R}^n$ and consider $\operatorname{div}_{\mu} := \operatorname{div}_{\mu, \mathbb{R}^n}$ as in (2.9) so $\mathcal{L}_{\mu} := -\operatorname{div}_{\mu}((1/\mu)\mathcal{A}\nabla)$ is maximal accretive, thus having a maximal accretive square root $\mathcal{L}_{\mu}^{1/2}$, in $L_{\mu}^2(\mathbb{R}^n)$. The solution of the Kato square root problem in [Auscher et al. 2002] was recently extended to degenerate elliptic equations by Cruz-Uribe and Rios [2015]. This shows that $\|\mathcal{L}_{\mu}^{1/2} f\|_{L_{\mu}^2(\mathbb{R}^n)} \approx \|\nabla f\|_{L_{\mu}^2(\mathbb{R}^n, \mathbb{R}^n)}$ for all $f \in W_{\mu}^{1,2}(\mathbb{R}^n)$; hence $\operatorname{Dom}(\mathcal{L}_{\mu}^{1/2}) = W_{\mu}^{1,2}(\mathbb{R}^n)$.

The operator \mathcal{L}_{μ} is also injective and type- $S_{\omega+}$ in $L_{\mu}^2(\mathbb{R}^n)$ for some $\omega \in (0, \frac{\pi}{2})$, so it has a bounded $H^{\infty}(S_{\theta+}^o)$ -functional calculus in $L_{\mu}^2(\mathbb{R}^n)$ for each $\theta \in (\omega, \pi)$, where $S_{\theta+}^o := \{z \in \mathbb{C} \setminus \{0\} : |\arg z| < \theta\}$. See Section 2.2 of [Auscher 2007] for the uniformly elliptic case and Theorems F and G in [Albrecht et al. 1996] for the general theory. An equivalent property is the validity of the quadratic estimate

$$\int_0^{\infty} \|\psi(t\mathcal{L}_{\mu}^{-\frac{1}{2}})f\|_{L_{\mu}^2(\mathbb{R}^n)}^2 \frac{dt}{t} \approx \|f\|_{L_{\mu}^2(\mathbb{R}^n)}^2 \quad \text{for all } f \in L_{\mu}^2(\mathbb{R}^n), \quad (2.11)$$

for each holomorphic ψ on $S_{\theta+}^o$ satisfying $|\psi(z)| \lesssim \min\{|z|^{\alpha}, |z|^{-\beta}\}$ for some $\alpha, \beta > 0$, where the bounded operator $\psi(t\mathcal{L}_{\mu}^{-\frac{1}{2}})$ on $L_{\mu}^2(\mathbb{R}^n)$ is defined by a Cauchy integral. More generally, the relationship between bounded holomorphic functional calculi and quadratic estimates is developed in the seminal articles [McIntosh 1986; Cowling et al. 1996].

The functional calculus then defines a bounded operator $\varphi(\mathcal{L}_{\mu})$ on $L_{\mu}^2(\mathbb{R}^n)$ for each bounded holomorphic function φ on $S_{\theta+}^o$ and $\|\varphi(\mathcal{L}_{\mu})\|_{L_{\mu}^2(\mathbb{R}^n) \rightarrow L_{\mu}^2(\mathbb{R}^n)} \lesssim_{\theta} \|\varphi\|_{\infty}$. Another consequence is that $-\mathcal{L}_{\mu}$ generates a holomorphic contraction semigroup $(e^{-\zeta \mathcal{L}_{\mu}})_{\zeta \in S_{\pi/2-\omega}^o \cup \{0\}}$ on $L_{\mu}^2(\mathbb{R}^n)$; thus $e^{-t\mathcal{L}_{\mu}} f \in \operatorname{Dom}(\mathcal{L}_{\mu})$ and $\partial_t(e^{-t\mathcal{L}_{\mu}} f) = \mathcal{L}_{\mu} e^{-t\mathcal{L}_{\mu}} f$ for all $f \in L_{\mu}^2(\mathbb{R}^n)$ and $t > 0$. The functional calculus also extends to define an unbounded operator $\phi(\mathcal{L}_{\mu})$ on $L_{\mu}^2(\mathbb{R}^n)$ for each holomorphic function ϕ on $S_{\theta+}^o$ satisfying $|\phi(z)| \lesssim \max\{|z|^{\alpha}, |z|^{-\beta}\}$ for some $\alpha, \beta > 0$, but the algebra homomorphism property of the functional calculus $(\phi_1(\mathcal{L}_{\mu})\phi_2(\mathcal{L}_{\mu})) = (\phi_1\phi_2)(\mathcal{L}_{\mu})$ must then be interpreted in the sense of unbounded linear operators. This allows us to interpret both the semigroup and the square root of \mathcal{L}_{μ} in terms of the functional calculus in order to justify some otherwise formal manipulations, beginning with (2.15) in the proof of the following corollary of the solution of the Kato problem in [Cruz-Uribe and Rios 2015].

Theorem 2.12. *Let $n \geq 1$ and suppose that $\mathcal{A} \in \mathcal{E}(n, \lambda, \Lambda, \mu)$ for some constants $0 < \lambda \leq \Lambda < \infty$ and an A_2 -weight μ on \mathbb{R}^n . The operator $\mathcal{L}_{\mu} := -\operatorname{div}_{\mu}((1/\mu)\mathcal{A}\nabla)$ satisfies*

$$\int_0^{\infty} \|t\mathcal{L}_{\mu} e^{-t^2\mathcal{L}_{\mu}} f\|_{L_{\mu}^2(\mathbb{R}^n)}^2 \frac{dt}{t} \approx \|\nabla f\|_{L_{\mu}^2(\mathbb{R}^n, \mathbb{R}^n)}^2, \quad (2.13)$$

$$\int_0^{\infty} \|t^2 \nabla_{x,t} \mathcal{L}_{\mu} e^{-t^2\mathcal{L}_{\mu}} f\|_{L_{\mu}^2(\mathbb{R}^n, \mathbb{R}^{n+1})}^2 \frac{dt}{t} \lesssim \|\nabla f\|_{L_{\mu}^2(\mathbb{R}^n, \mathbb{R}^n)}^2 \quad (2.14)$$

for all $f \in W_{\mu}^{1,2}(\mathbb{R}^n)$, where the implicit constants depend only on n, λ, Λ and $[\mu]_{A_2}$.

Proof. The functional calculus of \mathcal{L}_μ justifies the identity

$$\mathcal{L}_\mu e^{-t^2 \mathcal{L}_\mu} f = \mathcal{L}_\mu^{1/2} e^{-t^2 \mathcal{L}_\mu} \mathcal{L}_\mu^{1/2} f = e^{-(t^2/2) \mathcal{L}_\mu} \mathcal{L}_\mu e^{-(t^2/2) \mathcal{L}_\mu} f \quad (2.15)$$

for all $f \in \text{Dom}(\mathcal{L}_\mu^{1/2})$ and $t > 0$. The first equality in (2.15), the quadratic estimate in (2.11) and the solution of the Kato problem in [Cruz-Urbe and Rios 2015] imply

$$\begin{aligned} \int_0^\infty \|t \mathcal{L}_\mu e^{-t^2 \mathcal{L}_\mu} f\|_{L_\mu^2(\mathbb{R}^n)}^2 \frac{dt}{t} &= \int_0^\infty \|(\tau \mathcal{L}_\mu)^{1/2} e^{-\tau \mathcal{L}_\mu} \mathcal{L}_\mu^{1/2} f\|_{L_\mu^2(\mathbb{R}^n)}^2 \frac{d\tau}{\tau} \\ &\approx \|\mathcal{L}_\mu^{1/2} f\|_{L_\mu^2(\mathbb{R}^n)}^2 \\ &\approx \|\nabla f\|_{L_\mu^2(\mathbb{R}^n, \mathbb{R}^n)}^2 \end{aligned}$$

for all $f \in \text{Dom}(\mathcal{L}_\mu^{1/2}) = W_\mu^{1,2}(\mathbb{R}^n)$, which proves (2.13).

The bounded $H^\infty(S_{\theta+}^o)$ -functional calculus of \mathcal{L}_μ implies the uniform estimate

$$\begin{aligned} \|t \nabla_{x,t} e^{-t^2 \mathcal{L}_\mu} g\|_{L_\mu^2(\mathbb{R}^n, \mathbb{R}^{n+1})}^2 &= \|t \partial_t e^{-t^2 \mathcal{L}_\mu} g\|_{L_\mu^2(\mathbb{R}^n)}^2 + \|t \nabla_x e^{-t^2 \mathcal{L}_\mu} g\|_{L_\mu^2(\mathbb{R}^n, \mathbb{R}^n)}^2 \\ &\lesssim \|t^2 \mathcal{L}_\mu e^{-t^2 \mathcal{L}_\mu} g\|_{L_\mu^2(\mathbb{R}^n)}^2 + \int_{\mathbb{R}^n} t^2 \langle \mathcal{A} \nabla_x e^{-t^2 \mathcal{L}_\mu} g, \nabla_x e^{-t^2 \mathcal{L}_\mu} g \rangle \\ &\lesssim \|g\|_{L_\mu^2(\mathbb{R}^n)}^2 + \|t^2 \mathcal{L}_\mu e^{-t^2 \mathcal{L}_\mu} g\|_{L_\mu^2(\mathbb{R}^n)} \|e^{-t^2 \mathcal{L}_\mu} g\|_{L_\mu^2(\mathbb{R}^n)} \\ &\lesssim \|g\|_{L_\mu^2(\mathbb{R}^n)}^2 \end{aligned}$$

for all $g \in L_\mu^2(\mathbb{R}^n)$ and $t > 0$. Thus, the second equality in (2.15) and the vertical square function estimate in (2.13), which we have already proved, imply

$$\int_0^\infty \|t^2 \nabla_{x,t} \mathcal{L}_\mu e^{-t^2 \mathcal{L}_\mu} f\|_{L_\mu^2(\mathbb{R}^n, \mathbb{R}^{n+1})}^2 \frac{dt}{t} \lesssim \int_0^\infty \|t \mathcal{L}_\mu e^{-(t^2/2) \mathcal{L}_\mu} f\|_{L_\mu^2(\mathbb{R}^n)}^2 \frac{dt}{t} \lesssim \|\nabla f\|_{L_\mu^2(\mathbb{R}^n, \mathbb{R}^n)}^2$$

for all $f \in W_\mu^{1,2}(\mathbb{R}^n)$, which proves (2.14). \square

Now let us return to the case when $\Omega \subseteq \mathbb{R}^n$ is an arbitrary open set and suppose that $\mathbf{f} : \Omega \rightarrow \mathbb{R}^n$ is a measurable function for which $(1/\mu)\mathbf{f} \in L^\infty(\Omega)$. A *solution* of the inhomogeneous equation $\text{div}(\mathcal{A} \nabla u) = \text{div} \mathbf{f}$ in $\Omega \subseteq \mathbb{R}^n$ refers to any function $u \in W_{\mu, \text{loc}}^{1,2}(\Omega)$ such that $\int_{\mathbb{R}^n} \langle \mathcal{A} \nabla u - \mathbf{f}, \nabla \Phi \rangle = 0$ for all $\Phi \in C_c^\infty(\Omega)$. All solutions u of the homogeneous equation $\text{div}(\mathcal{A} \nabla u) = 0$ in Ω are locally bounded and Hölder continuous in the sense that

$$\|u\|_{L^\infty(B)} \lesssim \left(\int_{2B} |u|^2 d\mu \right)^{1/2} \quad (2.16)$$

and there exists $\alpha > 0$ such that

$$|u(x) - u(y)| \lesssim \left(\frac{|x - y|}{r(B)} \right)^\alpha \left(\int_{2B} |u|^2 d\mu \right)^{1/2} \quad \text{for all } x, y \in B, \quad (2.17)$$

and if, in addition, $u \geq 0$ almost everywhere on Ω , there is the Harnack inequality

$$\sup_B u \lesssim \inf_B u \quad (2.18)$$

for all balls B of radius $r(B)$ such that $2B \subseteq \Omega$, where α and the implicit constants depend only on n , λ , Λ and $[\mu]_{A_2}$. These properties follow from Corollary 2.3.4, Lemma 2.3.5 and Theorem 2.3.12 in [Fabes et al. 1982b] by observing that the proofs do not use the assumption therein that \mathcal{A} is symmetric. The estimates also hold when the balls B are replaced by (open) cubes Q , and also when the dilate $2B$ is replaced by $C_0 B$ for any $C_0 > 1$, provided the implicit constants are understood to depend on C_0 .

The following local boundedness estimate for solutions of the inhomogeneous equation is needed in Lemma 4.3, although only for $p = 2$. This is a simpler version of Theorem 8.17 in [Gilbarg and Trudinger 1977], which we have adapted to degenerate elliptic equations. In fact, the result for $p \geq 2$ is already proven in [Fabes et al. 1982b] by combining Corollary 2.3.4 with estimates (2.3.7) and (2.3.13) therein. The proof is included here for the reader's convenience and since it implies (2.16) as a special case, which in turn is the well-known starting point for establishing (2.17).

Theorem 2.19. *Let $n \geq 2$ and suppose that $\mathcal{A} \in \mathcal{E}(n, \lambda, \Lambda, \mu)$ for some constants $0 < \lambda \leq \Lambda < \infty$ and an A_2 -weight μ on \mathbb{R}^n . Let $\Omega \subseteq \mathbb{R}^n$ denote an open set and suppose that $\mathbf{f} : \Omega \rightarrow \mathbb{R}^n$ is a measurable function such that $(1/\mu)\mathbf{f} \in L^\infty(\Omega)$. If $p \in (1, \infty)$ and $\operatorname{div}(\mathcal{A}\nabla u) = \operatorname{div} \mathbf{f}$ in Ω , then*

$$\|u\|_{L^\infty(B)} \lesssim \left(\int_{2B} |u|^p d\mu \right)^{1/p} + r(B) \left\| \frac{1}{\mu} \mathbf{f} \right\|_{L^\infty(\Omega)} \quad (2.20)$$

for all balls B of radius $r(B) > 0$ such that $2B \subseteq \Omega$, where the implicit constant depends only on p , n , λ , Λ and $[\mu]_{A_2}$.

Proof. Suppose that $\operatorname{div}(\mathcal{A}\nabla u) = \operatorname{div} \mathbf{f}$ in Ω and consider a ball B such that $2B \subseteq \Omega$. First, assume that u is nonnegative and in $L^\infty(2B)$. Let $\epsilon > 0$, set $k = r(B) \|(1/\mu)\mathbf{f}\|_{L^\infty(\Omega)}$ and $\bar{u}_\epsilon := u + k + \epsilon$. Let B_r denote the ball concentric to B with radius $r > 0$ and recall the index $\delta > 0$ from the Sobolev inequality in Theorem 2.3. We claim that if $\gamma \in [p, \infty)$ and $r(B) \leq r_1 < r_2 \leq 2r(B)$, then

$$\left(\int_{B_{r_1}} \bar{u}_\epsilon^{\gamma(\frac{n}{n-1}+\delta)} d\mu \right)^{1/(\gamma(\frac{n}{n-1}+\delta))} \lesssim \left(\gamma \frac{r_1}{r_2 - r_1} \right)^{2/\gamma} \left(\int_{B_{r_2}} \bar{u}_\epsilon^\gamma d\mu \right)^{1/\gamma}, \quad (2.21)$$

where the implicit constant depends only on p , n , λ , Λ and $[\mu]_{A_2}$. To prove (2.21), fix $\eta \in C_c^\infty(\Omega)$ such that $\eta : \Omega \rightarrow [0, 1]$, $\eta \equiv 1$ on B_{r_1} , $\eta \equiv 0$ on $\Omega \setminus B_{r_2}$ and $\|\nabla \eta\|_\infty \leq 2/(r_2 - r_1)$. Set $\beta := \gamma - 1$ and $v := \eta^2 \bar{u}_\epsilon^\beta$. Note that $v \in W_{0,\mu}^{1,2}(\Omega)$ with

$$\nabla v = 2\eta \nabla \eta \bar{u}_\epsilon^\beta + \beta \eta^2 \bar{u}_\epsilon^{\beta-1} \nabla u,$$

since $0 < \epsilon \leq \bar{u}_\epsilon(x) \leq \|u\|_{L^\infty(2B)} + k + \epsilon < \infty$ for almost every $x \in 2B$; thus

$$\int_{\mathbb{R}^n} \langle \mathcal{A}\nabla u - \mathbf{f}, 2\eta \nabla \eta \bar{u}_\epsilon^\beta \rangle = - \int_{\mathbb{R}^n} \langle \mathcal{A}\nabla u - \mathbf{f}, \beta \eta^2 \bar{u}_\epsilon^{\beta-1} \nabla u \rangle.$$

We then use this identity and Cauchy's inequality with $\sigma > 0$ to obtain

$$\begin{aligned}
\int_{\mathbb{R}^n} \eta^2 \bar{u}_\epsilon^{\beta-1} |\nabla u|^2 d\mu &\lesssim_\lambda \int_{\mathbb{R}^n} \eta^2 \bar{u}_\epsilon^{\beta-1} \langle \mathcal{A} \nabla u, \nabla u \rangle \\
&= -2\beta^{-1} \int_{\mathbb{R}^n} \eta \bar{u}_\epsilon^\beta \langle \mathcal{A} \nabla u - \mathbf{f}, \nabla \eta \rangle + \int_{\mathbb{R}^n} \eta^2 \bar{u}_\epsilon^{\beta-1} \langle \mathbf{f}, \nabla u \rangle \\
&\lesssim_\Lambda (p-1)^{-1} \int_{\mathbb{R}^n} \eta \bar{u}_\epsilon^\beta \left(|\nabla u| + \left| \frac{1}{\mu} \mathbf{f} \right| \right) |\nabla \eta| d\mu + \int_{\mathbb{R}^n} \eta^2 \bar{u}_\epsilon^{\beta-1} \left| \frac{1}{\mu} \mathbf{f} \right| |\nabla u| d\mu \\
&\lesssim_p \sigma \int_{\mathbb{R}^n} \eta^2 \bar{u}_\epsilon^{\beta-1} |\nabla u|^2 d\mu + \sigma^{-1} \int_{\mathbb{R}^n} \bar{u}_\epsilon^{\beta+1} |\nabla \eta|^2 d\mu \\
&\quad + \int_{\mathbb{R}^n} \bar{u}_\epsilon^{\beta+1} |\nabla \eta|^2 d\mu + \int_{\mathbb{R}^n} \left(\frac{\eta}{r(B)} \right)^2 \bar{u}_\epsilon^{\beta+1} d\mu \\
&\quad + \sigma \int_{\mathbb{R}^n} \eta^2 \bar{u}_\epsilon^{\beta-1} |\nabla u|^2 d\mu + \sigma^{-1} \int_{\mathbb{R}^n} \left(\frac{\eta}{r(B)} \right)^2 \bar{u}_\epsilon^{\beta+1} d\mu,
\end{aligned}$$

where in the second inequality we used the assumption that $\beta := \gamma - 1 \geq p - 1$ and in the final inequality we used the fact that $|(1/\mu)\mathbf{f}| \leq k/r(B) \leq \bar{u}_\epsilon/r(B)$ on Ω . Next, choose $\sigma > 0$ small enough, depending only on p, λ and Λ , to deduce that

$$\int_{B_{r_1}} \bar{u}_\epsilon^{\beta-1} |\nabla u|^2 d\mu \lesssim_{p, \lambda, \Lambda} \int_{\mathbb{R}^n} \bar{u}_\epsilon^{\beta+1} \left(|\nabla \eta|^2 + \left(\frac{\eta}{r(B)} \right)^2 \right) d\mu \lesssim \frac{1}{(r_2 - r_1)^2} \int_{B_{r_2}} \bar{u}_\epsilon^{\beta+1} d\mu,$$

where in the final inequality we used the fact that $r(B) \geq r_2 - r_1$. Now combine this estimate with the Sobolev inequality (2.4) and recall that $\beta := \gamma - 1$ to obtain

$$\begin{aligned}
\left(\int_{B_{r_1}} \bar{u}_\epsilon^{\gamma(\frac{n}{n-1} + \delta)} d\mu \right)^{1/(\frac{n}{n-1} + \delta)} &\lesssim r_1^2 \int_{B_{r_1}} |\nabla(\bar{u}_\epsilon^{(\beta+1)/2})|^2 d\mu \\
&\lesssim ((\beta+1)r_1)^2 \int_{B_{r_1}} \bar{u}_\epsilon^{\beta-1} |\nabla u|^2 d\mu \\
&\lesssim \left(\gamma \frac{r_1}{r_2 - r_1} \right)^2 \int_{B_{r_2}} \bar{u}_\epsilon^\gamma d\mu,
\end{aligned}$$

where the implicit constants depend only on p, n, λ, Λ and $[\mu]_{A_2}$, proving (2.21).

We now apply the Moser iteration technique to prove (2.20). Set $\chi := n/(n-1) + \delta$ and define $\Phi(q, r) := \left(\int_{B_r} \bar{u}_\epsilon^q d\mu \right)^{1/q}$ for $q, r > 0$. Estimate (2.21) implies

$$\Phi(\gamma\chi, r_1) \leq \left(C\gamma \frac{r_1}{r_2 - r_1} \right)^{2/\gamma} \Phi(\gamma, r_2),$$

where C depends only on p, n, λ, Λ and $[\mu]_{A_2}$, and it follows by induction that

$$\Phi(p\chi^m, (1 + 2^{-m})r(B)) \leq (4Cp)^{(2/p) \sum_{k=0}^{m-1} \chi^{-k}} (2\chi)^{(2/p) \sum_{k=0}^{m-1} k\chi^{-k}} \Phi(p, 2r(B)) \lesssim \Phi(p, 2r(B))$$

for all $m \in \mathbb{N}$. This shows that

$$\|\bar{u}_\epsilon\|_{L^\infty(B)} = \lim_{m \rightarrow \infty} \Phi(p\chi^m, r(B)) \lesssim \Phi(p, 2r(B)) = \left(\int_{2B} \bar{u}_\epsilon^p d\mu \right)^{1/p}$$

and therefore

$$\|u\|_{L^\infty(B)} \leq \|\bar{u}_\epsilon\|_{L^\infty(B)} \lesssim \left(\int_{2B} \bar{u}_\epsilon^p d\mu \right)^{1/p} \lesssim \left(\int_{2B} u^p d\mu \right)^{1/p} + r(B) \left\| \frac{1}{\mu} f \right\|_{L^\infty(\Omega)} + \epsilon$$

for all $\epsilon > 0$, which implies (2.20).

Finally, it remains to remove the assumption that u is nonnegative and bounded. This is achieved by setting $\bar{u}_\epsilon := \max\{u, 0\} + k + \epsilon$ and $\bar{u}_\epsilon := -\min\{u, 0\} + k + \epsilon$ respectively and in each case adjusting the proof above to incorporate the truncated test function $v := \eta^2 h_N(\bar{u}_\epsilon) \bar{u}_\epsilon$, where

$$h_N(x) := \begin{cases} x^{\beta-1}, & x \leq N + k + \epsilon, \\ (N + k + \epsilon)^{\beta-1}, & x > N + k + \epsilon. \end{cases}$$

We leave the standard details to the reader. \square

The following self-improvement property for Carleson measures will be used in conjunction with the local Hölder continuity estimate for solutions in (2.17). The result is proved in the unweighted case in Lemma 2.14 in [Auscher et al. 2001]. In that proof, the Lebesgue measure on \mathbb{R}^n can in fact be replaced by any doubling measure, since the Whitney decomposition of open sets can be adapted to any such measure; see, for instance, Lemma 2 in Chapter I of [Stein 1993]. The result below then follows.

Lemma 2.22. *Let $n \geq 1$ and suppose that μ is an A_2 -weight on \mathbb{R}^n . Let $\alpha, \beta_0 > 0$ and suppose that $(v_t)_{t>0}$ is a collection of Hölder continuous functions on a cube $Q \subset \mathbb{R}^n$ satisfying*

$$0 \leq v_t(x) \leq \beta_0 \quad \text{and} \quad |v_t(x) - v_t(y)| \leq \beta_0 \left(\frac{|x - y|}{t} \right)^\alpha$$

for all $x, y \in Q$. If there exists $\eta \in (0, 1]$, $\beta > 0$ and, for each cube $Q' \subseteq Q$, a measurable set $F' \subset Q'$ such that

$$\mu(F') \geq \eta \mu(Q') \quad \text{and} \quad \frac{1}{\mu(Q')} \int_0^{\ell(Q')} \int_{F'} v_t(x) d\mu(x) \frac{dt}{t} \leq \beta,$$

then

$$\frac{1}{\mu(Q)} \int_0^{\ell(Q)} \int_Q v_t(x) d\mu(x) \frac{dt}{t} \lesssim_{\alpha, \eta} \beta + \beta_0,$$

where the implicit constant depends only on α, η, n and $[\mu]_{A_2}$.

3. Estimates for maximal operators

We obtain estimates for a variety of maximal operators $(M_\mu, D_{*,\mu}, N_*^\eta$ and $\tilde{N}_{*,\mu}^\eta)$ adapted to an A_2 -weight μ and degenerate elliptic operators $\mathcal{L}_\mu := -\operatorname{div}_\mu((1/\mu)\mathcal{A}\nabla)$ on \mathbb{R}^n for $n \geq 2$. These will be used

to prove the Carleson measure estimate from Theorem 1.3 in Section 4. We first define the maximal operators M_μ and $D_{*,\mu}$ by

$$M_\mu f(x) := \sup_{r>0} \int_{B(x,r)} |f(y)| d\mu(y),$$

$$D_{*,\mu} g(x) := \sup_{r>0} \left(\int_{B(x,r)} \left(\frac{|g(x) - g(y)|}{|x - y|} \right)^2 d\mu(y) \right)^{1/2}$$

for all $f \in L^1_{\mu,\text{loc}}(\mathbb{R}^n)$, $g \in W^{1,2}_{\mu,\text{loc}}(\mathbb{R}^n)$ and $x \in \mathbb{R}^n$. The usual unweighted and centred Hardy–Littlewood maximal operator is abbreviated by M . The maximal operator M_μ is bounded on $L^p_\mu(\mathbb{R}^n)$ for all $p \in (1, \infty)$ and satisfies the weak-type estimate

$$\mu(\{x \in \mathbb{R}^n : |M_\mu f(x)| > \kappa\}) \lesssim \kappa^{-1} \|f\|_{L^1_\mu(\mathbb{R}^n)} \quad \text{for all } \kappa > 0, \quad (3.1)$$

for all $f \in L^1_\mu(\mathbb{R}^n)$; see, for instance, Theorem 1 in Chapter I of [Stein 1993]. There is also the following weak-type estimate for the maximal operator $D_{*,\mu}$.

Lemma 3.2. *Let $n \geq 2$. If μ is an A_2 -weight on \mathbb{R}^n , then*

$$\mu(\{x \in \mathbb{R}^n : |D_{*,\mu} f(x)| > \kappa\}) \lesssim \kappa^{-2} \|\nabla f\|_{L^2_\mu(\mathbb{R}^n, \mathbb{R}^n)}^2 \quad \text{for all } \kappa > 0, \quad (3.3)$$

for all $f \in W^{1,2}_\mu(\mathbb{R}^n)$, where the implicit constant depends only on n and $[\mu]_{A_2}$.

Proof. If $f \in C_c^\infty(\mathbb{R}^n)$, then a version of Morrey’s inequality [1966, Theorem 3.5.2] shows that

$$\frac{|f(x) - f(y)|}{|x - y|} \lesssim M(\nabla f)(x) + M(\nabla f)(y)$$

for almost every $x, y \in \mathbb{R}^n$; hence

$$D_{*,\mu} f(x) \lesssim M(\nabla f)(x) + (M_\mu[M(\nabla f)]^2(x))^{1/2}.$$

Estimate (3.3) then follows from the weak-type bound for M_μ in (3.1), the fact that M is bounded on $L^2_\mu(\mathbb{R}^n)$ (see, for instance, Theorem 1 in Chapter V of [Stein 1993]) and the density of $C_c^\infty(\mathbb{R}^n)$ in $W^{1,2}_\mu(\mathbb{R}^n)$. \square

We now define the nontangential maximal operators N_*^η and $\tilde{N}_{*,\mu}^\eta$, for $\eta > 0$, by

$$N_*^\eta u(x) := \sup_{(y,t) \in \Gamma_\eta(x)} |u(y,t)|, \quad \tilde{N}_{*,\mu}^\eta v(x) := \sup_{(y,t) \in \Gamma_\eta(x)} \left(\int_{B(y,tat)} |v(z,t)|^2 d\mu(z) \right)^{1/2}$$

for all measurable functions u, v on \mathbb{R}_+^{n+1} (such that $v(\cdot, t) \in L^2_{\mu,\text{loc}}(\mathbb{R}^n)$ for a.e. $t > 0$) and $x \in \mathbb{R}^n$, where $\Gamma_\eta(x) := \{(y,t) \in \mathbb{R}_+^{n+1} : |y - x| < \eta t\}$ is the conical nontangential approach region in \mathbb{R}_+^{n+1} with vertex at x and aperture η .

Now suppose that $\mathcal{A} \in \mathcal{E}(n, \lambda, \Lambda, \mu)$, as defined by (2.6). In particular, since \mathcal{A} has real-valued coefficients, there exists an integral kernel $W_t(x, y)$ such that

$$e^{-t\mathcal{L}\mu} f(x) = \int_{\mathbb{R}^n} W_t(x, y) f(y) d\mu(y) \quad (3.4)$$

for all $f \in L^2_\mu(\mathbb{R}^n)$, and there exists constants $C_1, C_2 > 0$ such that

$$|W_t(x, y)| \leq \frac{C_1}{\mu(B(x, \sqrt{t}))} \exp\left(-C_2 \frac{|x - y|^2}{t}\right) \quad (3.5)$$

for all $t > 0$ and $x, y \in \mathbb{R}^n$. This was proved by Cruz-Uribe and Rios for $f \in C_c^\infty(\mathbb{R}^n)$ under the assumption that \mathcal{A} is symmetric; see Theorem 1 and Remark 3 in [Cruz-Uribe and Rios 2014]. The symmetry assumption can be removed, however, by following their proof and applying the Harnack inequality for degenerate parabolic equations obtained by Ishige [1999, Theorem A], which does not require symmetric coefficients, instead of the version recorded in Proposition 3.8 of [Cruz-Uribe and Rios 2008]. The results also extend to $f \in L^2_\mu(\mathbb{R}^n)$ by density, Schur's lemma and the doubling property of μ .

We now consider the semigroup generated by $\mathcal{L}_\mu := -\operatorname{div}_\mu((1/\mu)\mathcal{A}\nabla)$ with elliptic homogeneity (t replaced by t^2) and denoted by $\mathcal{P}_t := e^{-t^2\mathcal{L}_\mu}$ in the estimates below.

Lemma 3.6. *Let $n \geq 2$ and suppose that $\mathcal{A} \in \mathcal{E}(n, \lambda, \Lambda, \mu)$ for some constants $0 < \lambda \leq \Lambda < \infty$ and an A_2 -weight μ on \mathbb{R}^n . Let $p \in (1, \infty)$ and suppose that μ is also an A_p -weight on \mathbb{R}^n . If $x \in \mathbb{R}^n$, $\eta > 0$ and $\alpha \geq 1$, then*

$$\sup_{(y,t) \in \Gamma_\eta(x)} |(\eta t)^{-1} [\mathcal{P}_{\eta t}(f - c_{B(x, \alpha \eta t)})](y)|^2 \lesssim_\alpha [M_\mu(|\nabla f|^p)(x)]^{2/p} \quad (3.7)$$

for all $f \in W_\mu^{1,p}(\mathbb{R}^n)$ and $c_{B(x, \alpha \eta t)} \in \{f_{B(x, \alpha \eta t)} f \, d\mu, \tilde{f}_{B(x, \alpha \eta t)} f\}$, and

$$|N_*^\eta(\partial_t \mathcal{P}_t f)(x)|^2 \lesssim_\eta [M_\mu(|\nabla f|^p)(x)]^{2/p}, \quad (3.8)$$

$$|\eta^{-1} N_*^\eta(\partial_t \mathcal{P}_{\eta t} f)(x)|^2 \lesssim [M_\mu(|\nabla f|^p)(x)]^{2/p}, \quad (3.9)$$

$$|\tilde{N}_{*,\mu}^\eta(\nabla_x \mathcal{P}_{\eta t} f)(x)|^2 \lesssim M_\mu([M_\mu(|\nabla f|^p)]^{2/p})(x) + M_\mu(|\nabla f|^2)(x) \quad (3.10)$$

for all $f \in W_\mu^{1,2}(\mathbb{R}^n) \cap W_{\mu,\text{loc}}^{1,p}(\mathbb{R}^n)$, where the implicit constants depend only on $n, \lambda, \Lambda, p, [\mu]_{A_2}$ and $[\mu]_{A_p}$, as well as on α in (3.7) and on η in (3.8).

Proof. Let $x \in \mathbb{R}^n$, $(y, t) \in \Gamma_\eta(x)$, $f \in W_\mu^{1,2}(\mathbb{R}^n) \cap W_{\mu,\text{loc}}^{1,p}(\mathbb{R}^n)$, $f_{B(x,t)} := f_{B(x,t)} f$ and $\tilde{f}_{B(x,t)} := \tilde{f}_{B(x,t)} f \, d\mu$. To prove (3.7), it suffices to assume that $\eta = 1$ and $\alpha \geq 1$. We set $C_0(t) := B(x, \alpha t)$ and define the dyadic annulus $C_j(t) := B(x, 2^j \alpha t) \setminus B(x, 2^{j-1} \alpha t)$ for all $j \in \mathbb{N}$. The Gaussian kernel estimates in (3.4) and (3.5) imply

$$\begin{aligned} |t^{-1} [\mathcal{P}_t(f - f_{B(x, \alpha t)})](y)| &= t^{-1} \left| \int_{\mathbb{R}^n} W_{t^2}(y, z) [f(z) - f_{B(x, \alpha t)}] \, d\mu(z) \right| \\ &\leq \sum_{j=0}^{\infty} t^{-1} \frac{C_1}{\mu(B(y, t))} \int_{C_j(t)} \exp\left(-C_2 \frac{|y - z|^2}{t^2}\right) |f(z) - f_{B(x, \alpha t)}| \, d\mu(z) \\ &=: \sum_{j=0}^{\infty} I_j. \end{aligned}$$

To estimate I_0 , note that $B(x, \alpha t) \subseteq B(y, (1 + \alpha)t)$ and apply the doubling property of μ , followed by the L_μ^p -Poincaré inequality in (2.5) with $c_B = f_{B(x, \alpha t)} f$, to obtain

$$I_0 \lesssim_\alpha t^{-1} \int_{B(x, \alpha t)} |f(z) - f_{B(x, \alpha t)}| d\mu(z) \lesssim \left(\int_{B(x, \alpha t)} |\nabla f|^p d\mu \right)^{1/p} \lesssim [M_\mu(|\nabla f|^p)(x)]^{1/p}.$$

To estimate I_j , for each $j \in \mathbb{N}$, expand $f(z) - f_{B(x, \alpha t)}$ as a telescoping sum to write

$$\begin{aligned} I_j &\leq C_1 e^{-C_2(2^{j-1}\alpha-1)^2} \frac{\mu(B(x, 2^j \alpha t))}{\mu(B(y, t))} t^{-1} \\ &\quad \times \left(\int_{B(x, 2^j \alpha t)} |f - \tilde{f}_{B(x, 2^j \alpha t)}| d\mu + \sum_{i=1}^j |\tilde{f}_{B(x, 2^i \alpha t)} - \tilde{f}_{B(x, 2^{i-1} \alpha t)}| + |\tilde{f}_{B(x, \alpha t)} - f_{B(x, \alpha t)}| \right) \\ &\lesssim e^{-C_2(2^{j-1}\alpha-1)^2} \frac{\mu(B(y, (1 + 2^j \alpha)t))}{\mu(B(y, t))} \sum_{i=0}^j t^{-1} \int_{B(x, 2^i \alpha t)} |f - \tilde{f}_{B(x, 2^i \alpha t)}| d\mu \\ &\lesssim e^{-C_2(2^{j-1}\alpha-1)^2} (1 + 2^j \alpha)^{2n} \sum_{i=0}^j 2^i \alpha \left(\int_{B(x, 2^i \alpha t)} |\nabla f|^p d\mu \right)^{1/p} \\ &\lesssim_\alpha e^{-C_4 j} 4^{nj} [M_\mu(|\nabla f|^p)(x)]^{1/p}, \end{aligned}$$

where the second inequality relies on the inclusion $B(x, 2^j \alpha t) \subseteq B(y, (1 + 2^j \alpha)t)$, whilst the third inequality uses the doubling property of μ in (2.1) with $p = 2$, and the L_μ^p -Poincaré inequality in (2.5) with $c_B = f_{B(x, 2^j \alpha t)} f d\mu$. Altogether, we have

$$|t^{-1}[\mathcal{P}_t(f - f_{B(x, \alpha t)})](y)| \lesssim_\alpha \left(\sum_{j=0}^{\infty} e^{-C_4 j} 4^{nj} \right) [M_\mu(|\nabla f|^p)(x)]^{1/p} \lesssim [M_\mu(|\nabla f|^p)(x)]^{1/p},$$

which proves (3.7) when $c_{B(x, \alpha t)} = f_{B(x, \alpha t)} f$. The proof when $c_{B(x, \alpha t)} = f_{B(x, \alpha t)} f d\mu$ follows as above by replacing $f_{B(x, \alpha t)}$ with $\tilde{f}_{B(x, \alpha t)}$, since (2.5) can still be applied.

To prove (3.8) and (3.9), suppose that $\eta > 0$. The Gaussian kernel estimate for $e^{-t\mathcal{L}_\mu}$ in (3.5) implies that $t\partial_t \mathcal{P}_t f(y)$ has an integral kernel $\tilde{W}_{t^2}(y, z)$ satisfying

$$|\tilde{W}_{t^2}(y, z)| \leq \frac{C_1}{\mu(B(y, t))} \exp\left(-C_2 \frac{|y - z|^2}{t^2}\right)$$

and the conservation property $\int_{\mathbb{R}^n} \tilde{W}_{t^2}(y, z) d\mu(y) = 0$ for all $z \in \mathbb{R}^n$ and $t > 0$. This follows from Theorem 5 in [Cruz-Urbe and Rios 2014], where the assumption that \mathcal{A} is symmetric can be removed as per the remarks preceding this lemma. Therefore, we may write

$$|\partial_t \mathcal{P}_t f(y)| = t^{-1} \left| \int_{\mathbb{R}^n} \tilde{W}_{t^2}(y, z) [f(z) - f_{B(x, \eta t)}] d\mu(z) \right|$$

and a change of variables implies

$$\sup_{(y, t) \in \Gamma^n(x)} |\partial_t \mathcal{P}_t f(y)| = \sup_{(y, t) \in \Gamma(x)} t^{-1} \left| \int_{\mathbb{R}^n} \eta \tilde{W}_{(t/\eta)^2}(y, z) [f(z) - f_{B(x, t)}] d\mu(z) \right|.$$

We can then obtain (3.8) by following the proof of (3.7) with $\alpha = 1$ in order to show that this is bounded by $[M_\mu(|\nabla f|^p)(x)]^{1/p}$, since the doubling property of μ ensures that

$$|\eta \tilde{W}_{(t/\eta)^2}(y, z)| \leq \frac{C_{1,\eta}}{\mu(B(y, t))} \exp\left(-C_{2,\eta} \frac{|y-z|^2}{t^2}\right)$$

for some positive constants $C_{1,\eta}$ and $C_{2,\eta}$ that depend on η . We obtain (3.9) as an immediate consequence of (3.8) and the fact that $\eta^{-1} \partial_t \mathcal{P}_{\eta t} = (\partial_s \mathcal{P}_s)|_{s=\eta t}$.

To prove (3.10), let $\eta > 0$, set $u_{\eta t} := \mathcal{P}_{\eta t} f$ and choose a nonnegative function $\Phi \in C_c^\infty(B(y, 2\eta t))$ such that $\Phi \equiv 1$ on $B(y, \eta t)$ and $|\nabla_x \Phi| \lesssim (\eta t)^{-1}$. Let $c > 0$ denote a constant that will be chosen later. The definition of \mathcal{L}_μ implies

$$\begin{aligned} & \int_{B(y, \eta t)} |\nabla_x \mathcal{P}_{\eta t} f|^2 d\mu \\ & \leq \frac{1}{\mu(B(y, \eta t))} \int_{\mathbb{R}^n} |\nabla_x u_{\eta t}|^2 \Phi^2 d\mu \\ & \lesssim \frac{1}{\mu(B(y, \eta t))} \int_{\mathbb{R}^n} \langle \mathcal{A} \nabla_x u_{\eta t}, \nabla_x (u_{\eta t} - c) \rangle \Phi^2 \\ & = \frac{1}{\mu(B(y, \eta t))} \int_{\mathbb{R}^n} \{ \langle \mathcal{A} \nabla_x u_{\eta t}, \nabla_x [(u_{\eta t} - c) \Phi^2] \rangle - 2 \langle \mathcal{A} \nabla_x u_{\eta t}, \nabla_x \Phi (u_{\eta t} - c) \rangle \Phi \} \\ & \lesssim \frac{1}{\mu(B(y, \eta t))} \int_{\mathbb{R}^n} \{ (\mathcal{L}_\mu u_{\eta t})(u_{\eta t} - c) \Phi^2 + |\nabla_x u_{\eta t}| |\nabla_x \Phi| |u_{\eta t} - c| \Phi \} d\mu \\ & \leq \frac{1}{\mu(B(y, \eta t))} \int_{B(y, 2\eta t)} \left(\frac{1}{2\eta^2 t} |\partial_t u_{\eta t}| |u_{\eta t} - c| \Phi^2 + |\nabla_x u_{\eta t}| |\nabla_x \Phi| |u_{\eta t} - c| \Phi \right) d\mu \\ & =: I + II. \end{aligned}$$

Now fix $c := \tilde{f}_{B(x, 3\eta t)}$. To estimate I , we use Cauchy's inequality and the doubling property of μ , combined with the fact that $B(x, \eta t) \subseteq B(y, 2\eta t) \subseteq B(x, 3\eta t)$, to obtain

$$I \lesssim \int_{B(x, 3\eta t)} (|\eta^{-1} \partial_t u_{\eta t}|^2 + (\eta t)^{-2} |u_{\eta t} - f|^2 + (\eta t)^{-2} |f - \tilde{f}_{B(x, 3\eta t)}|^2) d\mu =: I_1 + I_2 + I_3.$$

It is immediate that $I_1 \leq M_\mu(|\eta^{-1} N_*^\eta(\partial_t \mathcal{P}_{\eta t} f)|^2)(x)$, whilst the semigroup property

$$|u_{\eta t}(z) - f(z)| = \left| \int_0^{\eta t} \partial_s u_s(z) ds \right| \leq \eta t N_*(\partial_s u_s)(z)$$

implies that $I_2 \lesssim M_\mu(|N_*(\partial_s u_s)|^2)(x)$, and the L_μ^2 -Poincaré inequality in (2.5) shows that $I_3 \lesssim M_\mu(|\nabla f|^2)(x)$; hence

$$I \leq M_\mu(|\eta^{-1} N_*^\eta(\partial_t \mathcal{P}_{\eta t} f)|^2)(x) + M_\mu(|N_*(\partial_s u_s)|^2)(x) + M_\mu(|\nabla f|^2)(x).$$

To estimate II , we use Cauchy's inequality with $\epsilon > 0$ to obtain

$$II \lesssim \frac{\epsilon}{\mu(B(y, \eta t))} \int_{\mathbb{R}^n} |\nabla_x u_{\eta t}|^2 \Phi^2 d\mu + \epsilon^{-1} (I_2 + I_3).$$

A sufficiently small choice of $\epsilon > 0$ allows the ϵ -term to be subtracted, yielding

$$\int_{B(y, \eta t)} |\nabla_x \mathcal{P}_{\eta t} f|^2 d\mu \lesssim I + II \lesssim M_\mu(|\eta^{-1} N_*^\eta(\partial_t \mathcal{P}_{\eta t} f)|^2 + |N_*(\partial_t \mathcal{P}_t f)|^2 + |\nabla f|^2)(x),$$

which, combined with (3.8) and (3.9), implies (3.10). \square

The pointwise estimates in Lemma 3.6 have the following corollary.

Corollary 3.11. *Let $n \geq 2$ and suppose that $\mathcal{A} \in \mathcal{E}(n, \lambda, \Lambda, \mu)$ for some constants $0 < \lambda \leq \Lambda < \infty$ and an A_2 -weight μ on \mathbb{R}^n . If $\eta > 0$, then*

$$\mu(\{x \in \mathbb{R}^n : |N_*^\eta(\partial_t \mathcal{P}_t f)(x)| > \kappa\}) \lesssim_\eta \kappa^{-2} \|\nabla f\|_{L_\mu^2(\mathbb{R}^n, \mathbb{R}^n)}^2, \quad (3.12)$$

$$\mu(\{x \in \mathbb{R}^n : |\eta^{-1} N_*^\eta(\partial_t \mathcal{P}_{\eta t} f)(x)| > \kappa\}) \lesssim \kappa^{-2} \|\nabla f\|_{L_\mu^2(\mathbb{R}^n, \mathbb{R}^n)}^2, \quad (3.13)$$

$$\mu(\{x \in \mathbb{R}^n : |\tilde{N}_{*, \mu}^\eta(\nabla_x \mathcal{P}_{\eta t} f)(x)| > \kappa\}) \lesssim \kappa^{-2} \|\nabla f\|_{L_\mu^2(\mathbb{R}^n, \mathbb{R}^n)}^2 \quad (3.14)$$

for all $\kappa > 0$ and $f \in W_\mu^{1,2}(\mathbb{R}^n)$, where the implicit constants depend only on n, λ, Λ and $[\mu]_{A_2}$, as well as on η in (3.12).

Proof. Estimates (3.12) and (3.13) follow respectively from (3.8) and (3.9), in the case $p = 2$, since M_μ satisfies the weak-type estimate in (3.1). To prove (3.14), note that there exists $1 < q < 2$ such that μ is an A_q -weight on \mathbb{R}^n ; see, for instance, Section 3 in Chapter V of [Stein 1993]. Therefore, combining (3.10) in the case $p = q$ with (3.1) and noting that $2/q > 1$, we obtain

$$\begin{aligned} \mu(\{x \in \mathbb{R}^n : |\tilde{N}_{*, \mu}^\eta(\nabla_x \mathcal{P}_{\eta t} f)(x)| > \kappa\}) &\lesssim \kappa^{-2} (\|M_\mu(|\nabla f|^q)\|_{L_\mu^{2/q}(\mathbb{R}^n)}^{2/q} + \|\nabla f\|_{L_\mu^2(\mathbb{R}^n, \mathbb{R}^n)}^2) \\ &\lesssim \kappa^{-2} \|\nabla f\|_{L_\mu^2(\mathbb{R}^n, \mathbb{R}^n)}^2 \end{aligned}$$

for all $\kappa > 0$ and $f \in W_\mu^{1,2}(\mathbb{R}^n)$ (since $W_\mu^{1,2}(\mathbb{R}^n) \subseteq W_{\mu, \text{loc}}^{1,q}(\mathbb{R}^n)$), as required. \square

4. The Carleson measure estimate

The purpose of this section is to prove the Carleson measure estimate (1.4) in Theorem 1.3. We adopt the strategy outlined at the end of Section 3.1 in [Hofmann et al. 2015a], although the crucial technical estimate, stated here as Theorem 4.10, is not at all an obvious extension of the uniformly elliptic case. Moreover, establishing the Carleson measure estimate directly allows us to avoid “good- λ ” inequalities and thus apply a change of variables based on the $W_{0, \mu}^{1,2}$ -Hodge decomposition in (2.10), instead of the $W_0^{1,2+\epsilon}$ -version (for a sufficiently small $\epsilon > 0$) required in [Hofmann et al. 2015a].

The technical result in Theorem 4.10 establishes (1.4) on certain “big pieces” of all cubes. The passage to the general estimate ultimately follows from the self-improvement property for Carleson measures in Lemma 2.22. This requires, however, that the Carleson measure estimate on the full gradient ∇u of a solution u can be controlled by the same estimate on its transversal derivative $\partial_t u$, which is the content of Lemma 4.2. We briefly postpone the statement and proof of Lemma 4.2 and Theorem 4.10, however, in order to deduce Theorem 1.3 from those results below.

In contrast to the previous two sections, the results here concern solutions of the equation $\operatorname{div}(A\nabla u) = 0$ in open sets $\Omega \subseteq \mathbb{R}_+^{n+1}$ when $n \geq 2$ and A is a t -independent coefficient matrix that satisfies (1.1) for some $0 < \lambda \leq \Lambda < \infty$ and an A_2 -weight μ on \mathbb{R}^n . In particular, in Section 2, weighted Sobolev spaces were defined on open sets in \mathbb{R}^d and matrix coefficients $A \in \mathcal{E}(d, \lambda, \Lambda, \mu)$ were considered for all $d \in \mathbb{N}$. Those results also hold here on open sets in the upper half-space with the weight $\mu(x, t) := \mu(x)$ and the coefficients $A(x, t) := A(x)$ for all $(x, t) \in \mathbb{R}_+^{n+1}$, since then $[\mu]_{A_2(\mathbb{R}^{n+1})} = [\mu]_{A_2(\mathbb{R}^n)}$ and $A \in \mathcal{E}(n+1, \lambda, \Lambda, \mu)$. In particular, the solution space $W_{\mu, \text{loc}}^{1,2}(\Omega)$ is defined and the regularity estimates in (2.16), (2.17) and (2.18) hold when $\Omega \subseteq \mathbb{R}_+^{n+1}$.

We will also use, without reference, the well-known fact that if u is a solution of $\operatorname{div}(A\nabla u) = 0$ in $\Omega \subseteq \mathbb{R}_+^{n+1}$, then $\partial_t u$ is also a solution in Ω . In particular, to see that $\partial_t u$ is in $W_{\mu, \text{loc}}^{1,2}(\Omega)$, a Whitney decomposition of Ω reduces matters to showing that $\partial_t u$ is in $W_\mu^{1,2}(R)$ for all cubes $R \subset \Omega$ satisfying $\ell(R) < \frac{1}{2} \operatorname{dist}(R, \partial\Omega)$. To this end, define the difference quotients $D_i^h u(X) := (1/h)[u(X + he_i) - u(X)]$ for all $X \in R$ and $h < \operatorname{dist}(R, \partial\Omega)$, where e_i is the unit vector in the i -th coordinate direction in \mathbb{R}^{n+1} . The t -independence of the coefficients implies that $D_{n+1}^h u$ is a solution in R , so we use the identity $D_{n+1}^h(\partial_t u) = \partial_t(D_{n+1}^h u)$ and Caccioppoli's inequality to obtain

$$\begin{aligned} \iint_R |D_{n+1}^h(\partial_t u)|^2 d\mu &\leq \iint_R |\nabla(D_{n+1}^h u)|^2 d\mu \lesssim \ell(R)^2 \iint_{2R} |D_{n+1}^h u|^2 d\mu \\ &\leq \ell(R)^2 \iint_{2R} |\partial_t u|^2 d\mu =: K \quad \text{for all } h < \operatorname{dist}(R, \partial\Omega), \end{aligned}$$

where the implicit constant depends only on n, λ, Λ and $[\mu]_{A_2}$, and the final bound holds uniformly in h because u is in $W_\mu^{1,2}(R)$; see Lemma 7.23 in [Gilbarg and Trudinger 1977]. We can then use Lemma 7.24 in the same reference to deduce that $\partial_t u$ is in $W_\mu^{1,2}(R)$ with the estimate

$$\|\partial_i \partial_t u\|_{L_\mu^2(R)}^2 = \|\partial_t \partial_i u\|_{L_\mu^2(R)}^2 \leq K$$

for all $i \in \{1, \dots, n+1\}$, as required. Note that the proofs of Lemmas 7.23 and 7.24 in [Gilbarg and Trudinger 1977] extend immediately to the weighted context considered here because $C^\infty(R)$ is still dense in $W_\mu^{1,2}(R)$.

Proof of Theorem 1.3 from Lemma 4.2 and Theorem 4.10. Let $Q \subset \mathbb{R}^n$ denote a cube and suppose that $u \in L^\infty(\mathbb{R}_+^{n+1})$ solves $\operatorname{div}(A\nabla u) = 0$ in \mathbb{R}_+^{n+1} . It follows a fortiori from Theorem 4.10 that there exist constants $C, c_0 > 0$ and, for each cube $Q' \subseteq Q$, a measurable set $F' \subset Q'$ such that $\mu(F') \geq c_0 \mu(Q')$ and

$$\frac{1}{\mu(Q')} \int_0^{\ell(Q')} \int_{F'} |t \partial_t u(x, t)|^2 d\mu(x) \frac{dt}{t} \leq C \|u\|_\infty^2,$$

where C and c_0 depend only on n, λ, Λ and $[\mu]_{A_2}$.

The coefficient matrix A is t -independent, so $\partial_t u$ is also a solution and thus the degenerate version of Moser's estimate in (2.16), followed by Caccioppoli's inequality, shows that $\|t \partial_t u\|_\infty \lesssim \|u\|_\infty$. Moreover,

the degenerate version of the de Giorgi–Nash Hölder regularity for solutions in (2.17) shows that

$$|t \partial_t u(x, t) - t \partial_t u(y, t)| \lesssim \left(\frac{|x - y|}{t} \right)^\alpha \|t \partial_t u\|_\infty \lesssim \|u\|_\infty \left(\frac{|x - y|}{t} \right)^\alpha$$

for all $x, y \in Q$ and $t > 0$, where all of the implicit constants and the exponent $\alpha > 0$ depend only on n, λ, Λ and $[\mu]_{A_2}$. Therefore, we may apply Lemma 2.22 with

$$\{v_t, \alpha, \beta_0, \eta, \beta\} := \{(t \partial_t u)^2, \alpha, C \|u\|_\infty^2, c_0, C \|u\|_\infty^2\}$$

to obtain

$$\frac{1}{\mu(Q)} \int_0^{\ell(Q)} \int_Q |t \partial_t u(x, t)|^2 d\mu(x) \frac{dt}{t} \lesssim \|u\|_\infty^2, \quad (4.1)$$

where the implicit constant depends only on n, λ, Λ and $[\mu]_{A_2}$. This estimate holds for all cubes Q , so by Lemma 4.2, we conclude that (1.4) holds. \square

We now dispense with the following lemma, which was used in the proof of Theorem 1.3 above to reduce to a Carleson measure estimate on the transversal derivative of solutions. The proof is adapted from Section 3.1 of [Hofmann et al. 2015a].

Lemma 4.2. *Let $n \geq 2$ and consider a cube $Q \subset \mathbb{R}^n$. If A is a t -independent coefficient matrix that satisfies the degenerate bound and ellipticity in (1.1) for some constants $0 < \lambda \leq \Lambda < \infty$ and an A_2 -weight μ on \mathbb{R}^n , then any solution $u \in L^\infty(4Q \times (0, 4\ell(Q)))$ of $\operatorname{div}(A \nabla u) = 0$ in $4Q \times (0, 4\ell(Q))$ satisfies*

$$\int_0^{\ell(Q)} \int_Q |t \nabla u(x, t)|^2 d\mu(x) \frac{dt}{t} \lesssim \int_0^{4\ell(Q)} \int_{4Q} |t \partial_t u(x, t)|^2 d\mu(x) \frac{dt}{t} + \mu(Q) \|u\|_\infty^2,$$

where the implicit constant depends only on n, λ, Λ and $[\mu]_{A_2}$.

Proof. Let $0 < \delta < \frac{1}{2}$ and set $\Phi_Q(t) := \Phi(t/\ell(Q))$, where $\Phi: \mathbb{R} \rightarrow [0, 1]$ denotes a C^∞ -function such that $\Phi(t) = 1$ for all $2\delta \leq t \leq 1$, whilst $\Phi(t) = 0$ for all $t \leq \delta$ and $t \geq 2$. Integrating by parts with respect to the t -variable and noting that $\|\partial_t \Phi\|_{L^\infty([1, 2])} \lesssim 1$, whilst $\|\partial_t \Phi\|_{L^\infty([\delta, 2\delta])} \lesssim 1/\delta$, we obtain

$$\begin{aligned} I &:= \int_Q \int_0^{2\ell(Q)} |\nabla u(x, t)|^2 \Phi_Q(t) t dt d\mu(x) \\ &\approx \int_Q \int_0^{2\ell(Q)} \partial_t (|\nabla u(x, t)|^2 \Phi_Q(t)) t^2 dt d\mu(x) \\ &\lesssim \int_Q \int_0^{2\ell(Q)} \langle \nabla \partial_t u(x, t), \nabla u(x, t) \rangle \Phi_Q(t) t^2 dt d\mu(x) \\ &\quad + \int_Q \int_{\ell(Q)}^{2\ell(Q)} |\nabla u(x, t)|^2 t^2 dt d\mu(x) + \int_Q \int_{\delta\ell(Q)}^{2\delta\ell(Q)} |\nabla u(x, t)|^2 t^2 dt d\mu(x) \\ &=: I' + I'' + I'''. \end{aligned}$$

For the term I' , we apply Cauchy's inequality with an arbitrary $\epsilon > 0$ to obtain

$$I' \leq \epsilon I + \frac{1}{\epsilon} \int_Q \int_0^{2\ell(Q)} |\nabla \partial_t u(x, t)|^2 t^3 dt d\mu(x).$$

For the term I'' , we apply Caccioppoli's inequality, the doubling property of μ and the fact that $t \approx \ell(Q)$ in the domain of the integration to obtain

$$\begin{aligned} I'' &\lesssim \ell(Q) \int_Q \int_{\ell(Q)}^{2\ell(Q)} |\nabla u(x, t)|^2 dt d\mu(x) \\ &\lesssim \frac{1}{\ell(Q)} \int_{2Q} \int_{\ell(Q)/2}^{5\ell(Q)/2} |u(x, t)|^2 dt d\mu(x) \lesssim \mu(Q) \|u\|_\infty^2. \end{aligned}$$

For the term I''' , the same reasoning shows that $I''' \lesssim \mu(Q) \|u\|_\infty^2$. We now fix $\epsilon > 0$, depending only on allowable constants, such that altogether

$$I \lesssim \int_Q \int_0^{2\ell(Q)} |\nabla \partial_t u(x, t)|^2 t^3 dt d\mu(x) + \mu(Q) \|u\|_\infty^2,$$

which is justified since $I < \infty$ by Caccioppoli's inequality and the support of Φ_Q .

To complete the estimate, we let $\{W_j : j \in J\}$ denote a collection of Whitney boxes (from a Whitney decomposition of \mathbb{R}_+^{n+1}) such that $W_j \cap (Q \times (0, 2\ell(Q))) \neq \emptyset$ and $\sum_{j \in J} \mathbb{1}_{2W_j}(x, t) \lesssim 1$. The coefficient matrix A is t -independent, so $\partial_t u$ is also a solution of $\operatorname{div}(A \nabla u) = 0$ in each set W_j ; hence we may apply Caccioppoli's inequality in combination with the fact that $t \approx l(W_j)$ in W_j to obtain

$$\begin{aligned} \int_{2\delta\ell(Q)}^{\ell(Q)} \int_Q |t \nabla u(x, t)|^2 d\mu(x) \frac{dt}{t} &\lesssim \sum_{j \in J} \iint_{W_j} |\nabla \partial_t u(x, t)|^2 t^3 dt d\mu(x) + \mu(Q) \|u\|_\infty^2 \\ &\lesssim \sum_{j \in J} l(W_j) \iint_{2W_j} |\partial_t u(x, t)|^2 dt d\mu(x) + \mu(Q) \|u\|_\infty^2 \\ &\lesssim \int_0^{4\ell(Q)} \int_{4Q} |t \partial_t u(x, t)|^2 d\mu(x) \frac{dt}{t} + \mu(Q) \|u\|_\infty^2, \end{aligned}$$

where the implicit constants do not depend on δ . The final result is then obtained by applying Fatou's lemma to estimate the limit as δ approaches 0. \square

The remainder of this section is dedicated to the proof of the crucial technical estimate, Theorem 4.10, that was used to prove Theorem 1.3. The proof adapts the change of variables from Section 3.2 of [Hofmann et al. 2015a] to the degenerate elliptic case. This is used to pull back solutions to certain sawtooth domains where the Carleson measure estimate can be verified by reducing matters to the vertical square function estimates in Theorem 2.12, which we recall were obtained from the solution of the Kato problem in [Cruz-Urbe and Rios 2015]. The following technical lemma, which reprises the notation $\mathcal{P}_t := e^{-t^2 \mathcal{L}_\mu}$ for $\mathcal{L}_\mu := -\operatorname{div}_\mu((1/\mu)A \nabla)$ and $\mathcal{A} \in \mathcal{E}(n, \lambda, \Lambda, \mu)$ as in (2.6) and Lemma 3.6, will be used to justify these changes of variables.

Lemma 4.3. *Let $n \geq 2$ and suppose that $\mathcal{A} \in \mathcal{E}(n, \lambda, \Lambda, \mu)$ for some constants $0 < \lambda \leq \Lambda < \infty$ and an A_2 -weight μ on \mathbb{R}^n . Let $Q \subset \mathbb{R}^n$ denote a cube and suppose that $\mathbf{f} : 5Q \rightarrow \mathbb{R}^n$ is a measurable function such that $(1/\mu)\mathbf{f} \in L^\infty(5Q)$. Let $\phi \in W_{0,\mu}^{1,2}(5Q)$ and suppose that $\operatorname{div}(A \nabla \phi) = \operatorname{div} \mathbf{f}$ in $5Q$. If $\kappa_0 > 0$,*

$0 < \eta < \frac{1}{2}$ and $x_0 \in Q$ satisfy $\Lambda(\eta, \phi, \mathcal{A})(x_0) \leq \kappa_0$, where

$$\Lambda(\eta, \phi, \mathcal{A}) := \eta^{-1} N_*^\eta(\partial_t \mathcal{P}_{\eta t} \phi) + N_*(\partial_t \mathcal{P}_t \phi) + [M_\mu(|\nabla_x \phi|^2)]^{1/2} + D_{*,\mu} \phi, \quad (4.4)$$

then

$$|\partial_t \mathcal{P}_{\eta t} \phi(x)| \leq \eta \kappa_0 \quad \text{for all } (x, t) \in \Gamma_\eta(x_0) \quad (4.5)$$

and

$$|(I - \mathcal{P}_{\eta t})\phi(x)| \lesssim \eta \left(\kappa_0 + \left\| \frac{1}{\mu} \mathbf{f} \right\|_\infty \right) t \quad \text{for all } (x, t) \in \Gamma_\eta(x_0) \cap (2Q \times (0, 4\ell(Q))), \quad (4.6)$$

where the implicit constant depends only on n, λ, Λ and $[\mu]_{A_2}$.

Proof. Suppose that $\kappa_0 > 0$, $0 < \eta < \frac{1}{2}$ and $x_0 \in Q$ satisfy $\Lambda(\eta, \phi, \mathcal{A})(x_0) \leq \kappa_0$. It follows a fortiori that $\eta^{-1} N_*^\eta(\partial_t \mathcal{P}_{\eta t} \phi)(x_0) \leq \kappa_0$, so (4.5) holds for all $(x, t) \in \Gamma_\eta(x_0)$.

To prove (4.6), first note that the properties of the semigroup imply

$$|(I - \mathcal{P}_{\eta t})\phi(x_0)| = \left| \int_0^{\eta t} \partial_s \mathcal{P}_s \phi(x_0) ds \right| \leq \eta t \kappa_0 \quad (4.7)$$

for all $t > 0$, since $N_*(\partial_s \mathcal{P}_s \phi)(x_0) \leq \kappa_0$. Now let $(x, t) \in \Gamma_\eta(x_0) \cap (2Q \times (0, 4\ell(Q)))$. We set $\phi_{x_0, \eta t} := \int_{B(x_0, 2\eta t)} \phi(y) dy$ and apply estimate (3.7) with $\alpha = 2$ to obtain

$$|\mathcal{P}_{\eta t}(\phi - \phi_{x_0, \eta t})(x)| \lesssim \eta t [M_\mu(|\nabla_x \phi|^2)(x_0)]^{1/2} \leq \eta t \kappa_0. \quad (4.8)$$

Next, since $\operatorname{div}(\mathcal{A} \nabla(\phi - \phi(x_0))) = \operatorname{div}(\mathcal{A} \nabla \phi) = \operatorname{div} \mathbf{f}$ in $5Q$, and since $0 < \eta < \frac{1}{2}$ ensures that $B(x_0, 2\eta t) \subseteq 5Q$, we may apply the degenerate version of Moser's estimate for inhomogeneous equations in (2.20) to obtain

$$\begin{aligned} |\phi(x) - \phi(x_0)| &\lesssim \left(\int_{B(x_0, 2\eta t)} |\phi(y) - \phi(x_0)|^2 d\mu(y) \right)^{1/2} + \eta t \left\| \frac{1}{\mu} \mathbf{f} \right\|_\infty \\ &\lesssim \eta t \left(D_{*,\mu} \phi(x_0) + \left\| \frac{1}{\mu} \mathbf{f} \right\|_\infty \right) \\ &\lesssim \eta t \left(\kappa_0 + \left\| \frac{1}{\mu} \mathbf{f} \right\|_\infty \right). \end{aligned} \quad (4.9)$$

Combining estimates (4.7), (4.8) and (4.9), we obtain

$$\begin{aligned} |(I - \mathcal{P}_{\eta t})\phi(x)| &\leq |\phi(x) - \phi(x_0)| + |(I - \mathcal{P}_{\eta t})\phi(x_0)| + |\mathcal{P}_{\eta t}(\phi - \phi_{x_0, \eta t})(x_0)| + |\mathcal{P}_{\eta t}(\phi - \phi_{x_0, \eta t})(x)| \\ &\lesssim \eta \left(\kappa_0 + \left\| \frac{1}{\mu} \mathbf{f} \right\|_\infty \right) t, \end{aligned}$$

which proves (4.6), as the implicit constant depends only on n, λ, Λ and $[\mu]_{A_2}$. \square

We now present the main technical result of this section. The proof is adapted from Section 3.2 of [Hofmann et al. 2015a], although some arguments have been simplified as detailed at the beginning of this section, and the additional justification required in the degenerate elliptic case has been emphasised.

The strategy of the original proof in [Hofmann et al. 2015a] was motivated in part by the fact that integration by parts is sufficient to establish the required estimate in the case when A has a certain block

upper-triangular structure. A key idea in that paper was to account for the presence of lower-triangular coefficients \mathbf{c} (and upper-triangular coefficients) by decomposing them according to a $W_0^{1,2+\epsilon}$ -Hodge decomposition. This was done locally on a given cube Q and the idea has been adapted here. First, the $W_{0,\mu}^{1,2}$ -Hodge decomposition $\mathbf{c}\mathbb{1}_Q = \mu\mathbf{h} - A_{\parallel}^*\nabla\varphi$ is introduced in (4.13), where A_{\parallel} is the $n \times n$ submatrix of A shown in (4.12). After integrating by parts, the divergence-free component $\mu\mathbf{h}$ provides valuable cancellation, whilst the adapted gradient vector field $A_{\parallel}^*\nabla\varphi$ facilitates a reduction to the square function estimates in Theorem 2.12, which are implied by the solution to the Kato problem in [Cruz-Uribe and Rios 2015], for the boundary operator $L_{\parallel,\mu}^* := -\operatorname{div}_{\mu}((1/\mu)A_{\parallel}^*\nabla_x)$.

The latter estimates, however, require that $L_{\parallel,\mu}^*$ acts on the range of $P_t^* := e^{-t^2 L_{\parallel,\mu}^*}$ and this is arranged by initially making the Dahlberg–Kenig–Stein-type pull-back $t \mapsto t - (I - P_{\eta t}^*)\varphi(x)$ so that the lower-triangular coefficients become $\mu\mathbf{h} - A_{\parallel}^*\nabla_x P_{\eta t}^*\varphi$. This change of variables is justified by choosing $\eta > 0$ small enough so that the pull-back is bi-Lipschitz in t . Once this is in place, a set F is introduced that contains a “big piece” of Q and on which the various maximal functions in Lemma 4.3 are bounded. The integration on $F \times (0, \ell(Q))$ is then performed by introducing a smooth test function Ψ_{δ} that equals 1 on $F \times (2\delta\ell(Q), 2\ell(Q))$ and is supported on a certain truncated sawtooth domain $\Omega_{\eta/8,Q,\delta}$ over F , where $\delta > 0$ is an arbitrary (small) parameter that provides for a smooth truncation in the t -direction near the boundary of \mathbb{R}_+^{n+1} . The main integration by parts is then performed in (4.32). The two principal terms \mathcal{S}_1 and \mathcal{S}_2 arise from the tangential and transversal integration by parts, respectively, where the former is taken with respect to the measure μ and thus requires additional justification from the uniformly elliptic case. These and numerous error terms are then shown to be appropriately under control.

Theorem 4.10. *Let $n \geq 2$ and consider a cube $Q \subset \mathbb{R}^n$. If A is a t -independent coefficient matrix that satisfies the degenerate bound and ellipticity in (1.1) for some constants $0 < \lambda \leq \Lambda < \infty$ and an A_2 -weight μ on \mathbb{R}^n , then for any solution $u \in L^\infty(4Q \times (0, 4\ell(Q)))$ that solves $\operatorname{div}(A\nabla u) = 0$ in $4Q \times (0, 4\ell(Q))$, there exist constants $C, c_0 > 0$ and a measurable set $F \subset Q$ such that $\mu(F) \geq c_0\mu(Q)$ and*

$$\frac{1}{\mu(Q)} \int_0^{\ell(Q)} \int_F |t\nabla u(x, t)|^2 d\mu(x) \frac{dt}{t} \leq C \|u\|_\infty^2, \quad (4.11)$$

where C and c_0 depend only on n, λ, Λ and $[\mu]_{A_2}$.

Proof. We begin by expressing the matrix A and its adjoint A^* (which is just the transpose A^t , since the matrix coefficients are real-valued) in the form

$$A = \left[\begin{array}{c|c} A_{\parallel} & \mathbf{b} \\ \hline \mathbf{c}^t & d \end{array} \right], \quad A^* = \left[\begin{array}{c|c} A_{\parallel}^* & \mathbf{c} \\ \hline \mathbf{b}^t & d \end{array} \right], \quad (4.12)$$

where A_{\parallel} denotes the $n \times n$ submatrix of A with entries $(A_{\parallel})_{i,j} := A_{i,j}$, $1 \leq i, j \leq n$, whilst $\mathbf{b} := (A_{i,n+1})_{1 \leq i \leq n}$ is a column vector, $\mathbf{c}^t := (A_{n+1,j})_{1 \leq j \leq n}$ is a row vector and $d := A_{n+1,n+1}$ is a scalar.

Now consider a cube $Q \subset \mathbb{R}^n$. The aim is to construct a set $F \subset Q$ with the required properties. To this end, we apply the Hodge decomposition from (2.10) to the space $L_\mu^2(5Q, \mathbb{R}^n)$ in order to write

$$\frac{1}{\mu} \mathbf{c}\mathbb{1}_Q = -\frac{1}{\mu} A_{\parallel}^*\nabla\varphi + \mathbf{h}, \quad \frac{1}{\mu} \mathbf{b}\mathbb{1}_Q = -\frac{1}{\mu} A_{\parallel}\nabla\tilde{\varphi} + \tilde{\mathbf{h}}, \quad (4.13)$$

where $\varphi, \tilde{\varphi} \in W_{\mu,0}^{1,2}(5Q)$ and $\mathbf{h}, \tilde{\mathbf{h}} \in L_{\mu}^2(5Q, \mathbb{R}^n)$ are such that $\operatorname{div}_{\mu} \mathbf{h} = \operatorname{div}_{\mu} \tilde{\mathbf{h}} = 0$ and

$$\int_{5Q} (|\nabla \varphi(x)|^2 + |\mathbf{h}(x)|^2) d\mu(x) \lesssim \int_{5Q} \left| \frac{\mathbf{c}(x)}{\mu} \right|^2 d\mu(x) \lesssim 1, \quad (4.14)$$

$$\int_{5Q} (|\nabla \tilde{\varphi}(x)|^2 + |\tilde{\mathbf{h}}(x)|^2) d\mu(x) \lesssim \int_{5Q} \left| \frac{\mathbf{b}(x)}{\mu} \right|^2 d\mu(x) \lesssim 1. \quad (4.15)$$

We extend each of $\varphi, \tilde{\varphi}, \mathbf{h}, \tilde{\mathbf{h}}$ to functions on \mathbb{R}^n by setting them equal to 0 on $\mathbb{R}^n \setminus 5Q$.

In Sections 2 and 3, we investigated the operators $\mathcal{L}_{\mu} := -\operatorname{div}_{\mu}((1/\mu)\mathcal{A}\nabla)$ and $\mathcal{P}_t := e^{-t^2\mathcal{L}_{\mu}}$ for arbitrary coefficient matrices \mathcal{A} in $\mathcal{E}(n, \lambda, \Lambda, \mu)$. We now set

$$\begin{aligned} L_{\parallel, \mu} &:= -\operatorname{div}_{\mu} \left(\frac{1}{\mu} A_{\parallel} \nabla_x \right), & P_t &:= e^{-t^2 L_{\parallel, \mu}}, \\ L_{\parallel, \mu}^* &:= -\operatorname{div}_{\mu} \left(\frac{1}{\mu} A_{\parallel}^* \nabla_x \right), & P_t^* &:= e^{-t^2 L_{\parallel, \mu}^*} \end{aligned} \quad (4.16)$$

in order to apply those results in the cases $\mathcal{A} = A_{\parallel}$ and $\mathcal{A} = A_{\parallel}^*$.

We now introduce two constants $\kappa_0, \eta > 0$, which will be fixed shortly, and recall the function $\Lambda(\eta, \phi, \mathcal{A})$ from (4.4) to define the set $F \subset Q$ by

$$F := \{x \in Q : \Lambda(\eta, \varphi, A_{\parallel}^*)(x) + \Lambda(\eta, \tilde{\varphi}, A_{\parallel})(x) + \tilde{N}_{*, \mu}^{\eta}(\nabla_x P_{\eta t}^* \varphi)(x) + \tilde{N}_{*, \mu}^{\eta}(\nabla_x P_{\eta t} \tilde{\varphi})(x) \leq \kappa_0\}. \quad (4.17)$$

Applying the weak-type bounds in (3.1), (3.3), (3.13) and (3.14) followed by the estimates from the Hodge decomposition in (4.14) and (4.15), we obtain

$$\mu(Q \setminus F) \lesssim \kappa_0^{-2} (\|\nabla \varphi\|_{L_{\mu}^2(\mathbb{R}^n, \mathbb{R}^n)}^2 + \|\nabla \tilde{\varphi}\|_{L_{\mu}^2(\mathbb{R}^n, \mathbb{R}^n)}^2) \lesssim \kappa_0^{-2} \mu(Q),$$

where the implicit constants depend only on n, λ, Λ and $[\mu]_{A_2}$. This allows us to now fix $\kappa_0 > 1$ and some constant $c_0 > 0$ such that $\mu(F) \geq c_0 \mu(Q)$, where both κ_0 and c_0 depend only on the allowed constants, and thus are independent of η .

We now fix the value of η as follows. First, for $0 \leq \alpha \leq 4$ and $\beta > 0$, let

$$\Omega_{\beta} := \bigcup_{x \in F} \Gamma_{\beta}(x), \quad \Omega_{\beta, Q, \alpha} := \Omega_{\beta} \cap (2Q \times (\alpha \ell(Q), 4\ell(Q))) \quad \text{and} \quad \Omega_{\beta, Q} := \Omega_{\beta, Q, 0}$$

denote the sawtooth domains in \mathbb{R}_+^{n+1} spanned by cones centred on F of aperture β . Next, note that the properties of the Hodge decomposition in (4.13) imply $-\operatorname{div}(A_{\parallel}^* \nabla \varphi) = \operatorname{div}(\mathbf{c} \mathbb{1}_{5Q})$ and $-\operatorname{div}(A_{\parallel} \nabla \tilde{\varphi}) = \operatorname{div}(\mathbf{b} \mathbb{1}_{5Q})$ in $5Q$. Therefore, we now fix $0 < \eta < \frac{1}{2}$ in accordance with (4.5) and (4.6) such that

$$\max\{|\partial_t P_{\eta t}^* \varphi(x)|, |\partial_t P_{\eta t} \tilde{\varphi}(x)|\} \leq \eta \kappa_0 < \frac{1}{8} \quad \text{for all } (x, t) \in \Omega_{\eta} \quad (4.18)$$

and

$$\begin{aligned} &\max\{|(I - P_{\eta t}^*)\varphi(x)|, |(I - P_{\eta t})\tilde{\varphi}(x)|\} \\ &\lesssim \eta \left(\kappa_0 + \max \left\{ \left\| \frac{1}{\mu} \mathbf{c} \right\|_{\infty}, \left\| \frac{1}{\mu} \mathbf{b} \right\|_{\infty} \right\} \right) t \lesssim \eta \kappa_0 t < \frac{1}{8} t \quad \text{for all } (x, t) \in \Omega_{\eta, Q}, \end{aligned} \quad (4.19)$$

where η and the implicit constants depend only on n, λ, Λ and $[\mu]_{A_2}$.

It remains to prove (4.11). We will achieve this by changing variables in the transversal direction using the mapping $t \mapsto \tau(x, t)$, with $x \in \mathbb{R}^n$ fixed, defined by

$$\tau(x, t) := t - (I - P_{\eta t}^*)\varphi(x)$$

and having Jacobian denoted by

$$J(x, t) := \partial_t \tau(x, t) = 1 + \partial_t P_{\eta t}^* \varphi(x). \quad (4.20)$$

In order to justify such changes of variables, we note from (4.18) and (4.19) that

$$\frac{7}{8}t < \tau(x, t) < \frac{9}{8}t \quad \text{and} \quad \frac{7}{8} < J(x, t) < \frac{9}{8} \quad \text{for all } (x, t) \in \Omega_{\eta, Q}. \quad (4.21)$$

In particular, for each $x \in F$ and $0 \leq \alpha \leq \frac{1}{8}$, this implies that the mapping $t \mapsto \tau(x, t)$ is bi-Lipschitz in t on $(2\alpha\ell(Q), 2\ell(Q))$ with range

$$(4\alpha\ell(Q), \ell(Q)) \subseteq \tau(x, \cdot)((2\alpha\ell(Q), 2\ell(Q))) \subseteq (\alpha\ell(Q), 4\ell(Q)). \quad (4.22)$$

Moreover, for each $0 < \beta \leq \eta$, the mapping $(x, t) \mapsto \rho(x, t)$ defined by

$$\rho(x, t) := (x, \tau(x, t)) = (x, t + P_{\eta t}^* \varphi(x) - \varphi(x))$$

is bi-Lipschitz in t on $\Omega_{\beta, Q}$ with range

$$\Omega_{8\beta/9, Q} \subseteq \rho(\Omega_{\beta, Q}) \subseteq \Omega_{8\beta/7, Q}. \quad (4.23)$$

Now consider a bounded solution u satisfying $\operatorname{div}(A\nabla u) = 0$ in $4Q \times (0, 4\ell(Q))$. The pull-back $u_1 := u \circ \rho$ is in $L^\infty(\Omega_{\eta, Q})$ and $\operatorname{div}(A_1 \nabla u_1) = 0$ in $\Omega_{\eta, Q}$, where

$$A_1 := \left[\begin{array}{c|c} JA_{\parallel} & \mathbf{b} + A_{\parallel} \nabla_x \varphi - A_{\parallel} \nabla_x P_{\eta t}^* \varphi \\ \hline (\mu \mathbf{h} - A_{\parallel}^* \nabla_x P_{\eta t}^* \varphi)^t & \langle A \mathbf{p}, \mathbf{p} \rangle / J \end{array} \right]$$

and

$$\mathbf{p}(x, t) := \begin{bmatrix} \nabla_x \tau(x, t) \\ -1 \end{bmatrix} = \begin{bmatrix} \nabla_x P_{\eta t}^* \varphi(x) - \nabla_x \varphi(x) \\ -1 \end{bmatrix}. \quad (4.24)$$

Our statement that $\operatorname{div}(A_1 \nabla u_1) = 0$ in $\Omega_{\eta, Q}$ does not mean that A_1 satisfies (1.1), only that $u_1 \in W_{\mu, \text{loc}}^{1,2}(\Omega_{\eta, Q})$ and that $\int_{\mathbb{R}^{n+1}_+} \langle A_1 \nabla u_1, \nabla \Phi \rangle = 0$ for all $\Phi \in C_c^\infty(\Omega_{\eta, Q})$. To prove this, we combine the pointwise identity

$$\langle A((\nabla u) \circ \rho), (\nabla v) \circ \rho \rangle J = \langle A_1 \nabla(u \circ \rho), \nabla(v \circ \rho) \rangle \quad \text{for all } v \in W_{0, \mu}^{1,2}(\rho(\Omega_{\eta, Q})) \quad (4.25)$$

with the change of variables $(x, t) \mapsto \rho(x, t)$ on $\Omega_{\eta, Q}$, which is justified because ρ is bi-Lipschitz in t on $\Omega_{\eta, Q}$ with range $\rho(\Omega_{\eta, Q}) \subset 4Q \times (0, 4\ell(Q))$ by (4.23). Also, we note for later use that $\|\mathbb{1}_{\Omega_{\eta, Q}} u_1\|_\infty \leq \|u\|_\infty$ and, using (4.21), that

$$|\nabla u_1| \lesssim \left| \begin{bmatrix} \nabla_x u_1 - (\nabla_x \tau)(\partial_t u_1)/J \\ (\partial_t u_1)/J \end{bmatrix} \right| + |\nabla_x \tau| |\partial_t u_1| = |(\nabla u) \circ \rho| + |\nabla_x \tau| |\partial_t u_1| \quad (4.26)$$

on $\Omega_{\eta, Q}$.

Next, in order to work with the pull-back solution u_1 , we consider an arbitrary constant $0 < \delta \leq \frac{1}{8}$ and define a smooth cut-off function Ψ_δ adapted to $\Omega_{\eta, Q}$ as follows. Let $\delta_F(x) := \text{dist}(x, F)$, fix a C^∞ -function $\Phi : \mathbb{R} \rightarrow [0, 1]$ satisfying $\Phi(t) = 1$ when $t < \frac{1}{16}$ and $\Phi(t) = 0$ when $t \geq \frac{1}{8}$, and then define

$$\Psi_\delta(x, t) := \Phi\left(\frac{\delta_F(x)}{\eta t}\right) \Phi\left(\frac{t}{32\ell(Q)}\right) \left(1 - \Phi\left(\frac{t}{16\delta\ell(Q)}\right)\right) \quad \text{for all } (x, t) \in \mathbb{R}_+^{n+1}.$$

This function is designed so that $\Psi_\delta \equiv 1$ on $F \times (2\delta\ell(Q), 2\ell(Q))$, and since $\eta < \frac{1}{2}$, we have $\text{supp } \Psi_\delta \subseteq \Omega_{\eta/8, Q, \delta}$ and

$$|\nabla_{x,t} \Psi_\delta(x, t)| \lesssim \frac{\mathbb{1}_{E_1}(x, t)}{t} + \frac{\mathbb{1}_{E_2}(x, t)}{\ell(Q)} + \frac{\mathbb{1}_{E_3}(x, t)}{\delta\ell(Q)} \quad \text{for all } (x, t) \in \Omega_{\eta/8, Q, \delta}, \quad (4.27)$$

where

$$\begin{aligned} E_1 &:= \{(x, t) \in 2Q \times (0, 4\ell(Q)) : \frac{1}{16}\eta t \leq \delta_F(x) \leq \frac{1}{8}\eta t\}, \\ E_2 &:= 2Q \times (2\ell(Q), 4\ell(Q)), \\ E_3 &:= 2Q \times (\delta\ell(Q), 2\delta\ell(Q)). \end{aligned}$$

In contrast to Section 3.2 in [Hofmann et al. 2015a], the cut-off function Ψ_δ introduced here incorporates an additional truncation in the t -direction at the boundary. This is done to simplify subsequent integration-by-parts arguments, since it ensures that Ψ_δ vanishes on the boundary of \mathbb{R}_+^{n+1} . For later purposes, it is also convenient to isolate the following general fact here.

Remark 4.28. For each $k \in \mathbb{Z}$, let \mathbb{D}_k^η denote the grid of dyadic cubes $Q' \subset \mathbb{R}^n$ such that

$$\frac{1}{64}\eta 2^{-k} \leq \text{diam } Q' < \frac{1}{32}\eta 2^{-k}.$$

If $C_0 > 0$ and $(v_t)_{t>0}$ is a collection of nonnegative measurable functions such that

$$\sup_{t \in [2^{-k}, 2^{-k+1}]} \int_{Q'} v_t(x) d\mu(x) \leq C_0 \quad \text{for all } k \in \mathbb{Z}, \text{ for all } Q' \in \mathbb{D}_k^\eta,$$

then

$$\iint_{\mathbb{R}_+^{n+1}} \left(\frac{\mathbb{1}_{E_1}(x, t)}{t} + \frac{\mathbb{1}_{E_2}(x, t)}{\ell(Q)} + \frac{\mathbb{1}_{E_3}(x, t)}{\delta\ell(Q)} \right) v_t(x) d\mu(x) dt \lesssim C_0 \mu(Q), \quad (4.29)$$

where the implicit constant depends only on n, λ, Λ and $[\mu]_{A_2}$. To see this, first observe that since δ_F is a Lipschitz mapping with constant 1, we have

$$\begin{aligned} Q^{(1)} \times [2^{-k}, 2^{-k+1}] &\subseteq \tilde{E}_1 := \left\{ (x, t) \in 4Q \times (0, 4\ell(Q)) : \frac{\eta t}{C} \leq \delta_F(x) \leq C\eta t \right\}, \\ Q^{(2)} \times [2^{-k}, 2^{-k+1}] &\subseteq 4Q \times (\ell(Q), 8\ell(Q)), \\ Q^{(3)} \times [2^{-k}, 2^{-k+1}] &\subseteq 4Q \times \left(\frac{1}{2}\delta\ell(Q), 4\delta\ell(Q)\right) \end{aligned}$$

whenever $E_i \cap (Q^{(i)} \times [2^{-k}, 2^{-k+1}]) \neq \emptyset$ and $i \in \{1, 2, 3\}$. The estimate in (4.27) and the doubling property of μ then imply that the left side of (4.29) is bounded by

$$\begin{aligned}
C_0 \left(\sum_{k \in \mathbb{Z}} \sum_{Q' \in \mathbb{D}_k^\eta} \int_{2^{-k}}^{2^{-k+1}} \int_{Q'} \mathbb{1}_{\tilde{E}_1} d\mu \frac{dt}{t} + C \int_{\ell(Q)}^{8\ell(Q)} \mu(Q) dt + C \int_{\frac{1}{2}\delta\ell(Q)}^{4\delta\ell(Q)} \mu(Q) dt \right) \\
\lesssim C_0 \left(\int_{4Q} \int_{(1/(C\eta))\delta_F(x)}^{(C/\eta)\delta_F(x)} \frac{d\mu(x)}{t} + \mu(Q) \right) \lesssim C_0 \mu(Q),
\end{aligned}$$

as required.

We now proceed to prove (4.11). First, note that it suffices to show that

$$\sup_{0 < \delta \leq 1/8} \int_{4\delta\ell(Q)}^{\ell(Q)} \int_F |t \nabla u(x, t)|^2 d\mu(x) \frac{dt}{t} \lesssim \|u\|_\infty^2 \mu(Q), \quad (4.30)$$

since we may then obtain (4.11) by using Fatou's lemma to pass to the limit as δ approaches 0. To this end, we use (4.22), followed by the bi-Lipschitz-in- t change of variables $t \mapsto \tau(x, t)$ on $(\delta\ell(Q), 2\ell(Q))$ for each $x \in F$, estimate (4.21) and identity (4.25) to obtain

$$\begin{aligned}
\int_{4\delta\ell(Q)}^{\ell(Q)} \int_F |t \nabla u(x, t)|^2 d\mu(x) \frac{dt}{t} &\lesssim \int_F \int_{4\delta\ell(Q)}^{\ell(Q)} \langle A \nabla u, \nabla u \rangle t dt dx \\
&\lesssim \int_F \int_{2\delta\ell(Q)}^{2\ell(Q)} \langle A_1 \nabla u_1, \nabla u_1 \rangle t dt dx \\
&\leq \iint_{\mathbb{R}_+^{n+1}} \langle A_1 \nabla u_1, \nabla u_1 \rangle \Psi_\delta^2 t dx dt.
\end{aligned}$$

Thus, in order to prove (4.30) and ultimately (4.11), it suffices to show that

$$\iint_{\mathbb{R}_+^{n+1}} \langle A_1 \nabla u_1, \nabla u_1 \rangle \Psi_\delta^2 t dx dt \lesssim \|u\|_\infty^2 \mu(Q) \quad \text{for all } 0 < \delta \leq \frac{1}{8}, \quad (4.31)$$

where the implicit constant depends only on n, λ, Λ and $[\mu]_{A_2}$.

Next, we recall that $\operatorname{div}(A_1 \nabla u_1) = 0$ in $\Omega_{\eta, Q}$, noting that $u_1 \Psi_\delta^2 t \in W_{0, \mu}^{1,2}(\Omega_{\eta, Q})$, and then integrate by parts to obtain

$$\begin{aligned}
&\iint_{\mathbb{R}_+^{n+1}} \langle A_1 \nabla u_1, \nabla u_1 \rangle \Psi_\delta^2 t dx dt \\
&= -\frac{1}{2} \iint_{\mathbb{R}_+^{n+1}} \langle A_1 \nabla(u_1^2), \nabla(\Psi_\delta^2 t) \rangle dx dt \\
&= -\frac{1}{2} \iint_{\mathbb{R}_+^{n+1}} \left\langle \nabla(u_1^2), \frac{1}{\mu} A_1^* e_{n+1} \right\rangle \Psi_\delta^2 d\mu dt - \frac{1}{2} \iint_{\mathbb{R}_+^{n+1}} \langle A_1 \nabla(u_1^2), \nabla(\Psi_\delta^2) \rangle t dx dt \\
&= \frac{1}{2} \iint_{\mathbb{R}_+^{n+1}} u_1^2 (L_{\|\cdot, \mu}^* P_{\eta t}^* \varphi) \Psi_\delta^2 d\mu dt + \frac{1}{2} \iint_{\mathbb{R}_+^{n+1}} u_1^2 \partial_t \left(\frac{\langle A P, P \rangle}{J} \right) \Psi_\delta^2 dx dt \\
&\quad - \frac{1}{2} \iint_{\mathbb{R}_+^{n+1}} \langle A_1 \nabla(u_1^2), \nabla(\Psi_\delta^2) \rangle t dx dt + \frac{1}{2} \iint_{\mathbb{R}_+^{n+1}} u_1^2 \langle e_{n+1}, A_1 \nabla(\Psi_\delta^2) \rangle dx dt \\
&=: S_1 + S_2 + E_1 + E_2,
\end{aligned} \quad (4.32)$$

where $e_{n+1} := (0, \dots, 0, 1)$ denotes the unit vector in the t -direction. In particular, note that the tangential integration by parts

$$\int_{\mathbb{R}^n} \left\langle \nabla_x(u_1^2), \mathbf{h} - \frac{1}{\mu} A_{\parallel}^* \nabla_x P_{\eta t}^* \varphi \right\rangle \Psi_{\delta}^2 d\mu = \int_{\mathbb{R}^n} u_1^2 \operatorname{div}_{\mu} \left[\left(\mathbf{h} - \frac{1}{\mu} A_{\parallel}^* \nabla_x P_{\eta t}^* \varphi \right) \Psi_{\delta}^2 \right] d\mu,$$

with respect to the measure μ , is justified by the definition of the operator div_{μ} , since $P_{\eta t}^* \varphi \in \operatorname{Dom}(L_{\parallel, \mu}^*)$ and $\operatorname{div}_{\mu} \mathbf{h} = 0$ imply $(\mathbf{h} - (1/\mu) A_{\parallel}^* \nabla_x P_{\eta t}^* \varphi) \Psi_{\delta}^2 \in \operatorname{Dom}(\operatorname{div}_{\mu})$ (recall (2.8), (2.9) and (4.16)). Meanwhile, the transversal integration by parts

$$\int_0^{\infty} \partial_t(u_1^2) \left(\frac{\langle A\mathbf{p}, \mathbf{p} \rangle}{J} \right) \Psi_{\delta}^2 dt = - \int_0^{\infty} u_1^2 \partial_t \left[\left(\frac{\langle A\mathbf{p}, \mathbf{p} \rangle}{J} \right) \Psi_{\delta}^2 \right] dt$$

is justified because Ψ_{δ} vanishes on the boundary of \mathbb{R}_+^{n+1} .

We proceed to prove that, for all $\sigma \in (0, 1)$, each term in (4.32) is controlled by

$$\mathbf{S}_1 + \mathbf{S}_2 + \mathbf{E}_1 + \mathbf{E}_2 \lesssim \sigma \iint_{\mathbb{R}_+^{n+1}} \langle A_1 \nabla u_1, \nabla u_1 \rangle \Psi_{\delta}^2 t dx dt + \sigma^{-1} \|u\|_{\infty}^2 \mu(Q), \quad (4.33)$$

where the implicit constant depends only on n, λ, Λ and $[\mu]_{A_2}$. Estimate (4.31) will then follow by fixing a sufficiently small $\sigma \in (0, 1)$, depending only on allowed constants, to move the integral in (4.33) to the left side of (4.32). This is justified because the integral in (4.33) is finite by Caccioppoli's inequality and the fact that Ψ_{δ} vanishes in a neighbourhood of the boundary of \mathbb{R}_+^{n+1} ($\operatorname{supp} \Psi_{\delta} \subseteq \Omega_{\eta/8, Q, \delta}$).

We now prove (4.33) in three steps to complete the proof.

Step 1: estimates for the error terms \mathbf{E}_1 and \mathbf{E}_2 in (4.32).

We first apply Cauchy's inequality with σ to write

$$\begin{aligned} \mathbf{E}_1 &\leq \left| \frac{1}{2} \iint_{\mathbb{R}_+^{n+1}} \langle A_1 \nabla(u_1^2), \nabla(\Psi_{\delta}^2) \rangle t dx dt \right| \\ &= 2 \left| \iint_{\mathbb{R}_+^{n+1}} \langle A_1 \nabla u_1, \nabla \Psi_{\delta} \rangle u_1 \Psi_{\delta} t dx dt \right| \\ &\lesssim \sigma \iint_{\mathbb{R}_+^{n+1}} \langle A_1 \nabla u_1, \nabla u_1 \rangle \Psi_{\delta}^2 t dx dt + \sigma^{-1} \iint_{\mathbb{R}_+^{n+1}} u_1^2 \langle A_1 \nabla \Psi_{\delta}, \nabla \Psi_{\delta} \rangle t dx dt \\ &=: \sigma \iint_{\mathbb{R}_+^{n+1}} \langle A_1 \nabla u_1, \nabla u_1 \rangle \Psi_{\delta}^2 t dx dt + \sigma^{-1} \mathbf{E}'_1. \end{aligned}$$

We then use $\mu \mathbf{h} = c \mathbb{1}_{5Q} + A_{\parallel}^* \nabla \varphi$ from (4.13), the degenerate bound in (1.1) for A , the bound $\|\mathbb{1}_{\Omega_{\eta, Q}} u_1\|_{\infty} \lesssim \|u\|_{\infty}$ and the estimate for $\nabla \Psi_{\delta}$ from (4.27) to obtain

$$\mathbf{E}'_1 + \mathbf{E}_2 \lesssim \|u\|_{\infty}^2 \iint_{\Omega_{\eta/8, Q}} \left(\frac{\mathbb{1}_{E_1}}{t} + \frac{\mathbb{1}_{E_2}}{\ell(Q)} + \frac{\mathbb{1}_{E_3}}{\delta \ell(Q)} \right) (1 + |\nabla_x(I - P_{\eta t}^*) \varphi|^2) d\mu dt,$$

where (4.27) ensures that $|\nabla(\Psi_\delta^2)|$ and $|\nabla\Psi_\delta|^2 t$ can be controlled in the same manner. In order to apply Remark 4.28 with $v_t = \mathbb{1}_{\Omega_{\eta/8,Q}}(1 + |\nabla_x(I - P_{\eta t}^*)\varphi|^2)$, we observe that if $k \in \mathbb{Z}$, $Q' \in \mathbb{D}_k^\eta$ and $\Omega_{\eta/8,Q,\delta} \cap (Q' \times [2^{-k}, 2^{-k+1}]) \neq \emptyset$, then there exists $x_0 \in F$ such that $Q' \subseteq \Delta(x_0, \eta 2^{-k}) \subseteq CQ'$, where Δ is used to denote balls in \mathbb{R}^n ; hence

$$Q' \times [2^{-k}, 2^{-k+1}] \subseteq \Omega_{\eta,2Q,\delta/4} \quad (4.34)$$

and the doubling property of μ implies

$$\begin{aligned} \int_{Q'} |\nabla_x(I - P_{\eta t}^*)\varphi|^2 d\mu &\lesssim \int_{\Delta(x_0, \eta t)} |\nabla_x P_{\eta t}^*\varphi|^2 d\mu + \int_{\Delta(x_0, \eta 2^{-k})} |\nabla_x \varphi|^2 d\mu \\ &\lesssim [\tilde{N}_{*,\mu}^\eta(\nabla_x P_{\eta t}^*\varphi)(x_0)]^2 + M_\mu(|\nabla_x \varphi|^2)(x_0) \\ &\lesssim \kappa_0^2 \lesssim 1 \quad \text{for all } t \in [2^{-k}, 2^{-k+1}], \end{aligned} \quad (4.35)$$

where in the last line we used the definition of the set F in (4.17) and the weighted maximal operators $\tilde{N}_{*,\mu}$ and M_μ from Section 3. It thus follows from (4.29) that $E'_1 + E_2 \lesssim \|u\|_\infty^2 \mu(Q)$, so altogether we have

$$E_1 + E_2 \lesssim \sigma \iint_{\mathbb{R}_+^{n+1}} \langle A_1 \nabla u_1, \nabla u_1 \rangle \Psi_\delta^2 t \, dx \, dt + \sigma^{-1} \|u\|_\infty^2 \mu(Q) \quad \text{for all } \sigma \in (0, 1). \quad (4.36)$$

Step 2: estimates for the term S_1 in (4.32).

We note that $\partial_t P_{\eta t}^* = -2\eta^2 t L_{\parallel,\mu}^* P_{\eta t}^*$ on $L_\mu^2(\mathbb{R}^n)$ and integrate by parts in t to write

$$\begin{aligned} S_1 &= \frac{1}{2} \iint_{\mathbb{R}_+^{n+1}} u_1^2 (L_{\parallel,\mu}^* P_{\eta t}^* \varphi) \Psi_\delta^2 d\mu \, dt \\ &= -\frac{1}{2} \iint_{\mathbb{R}_+^{n+1}} u_1^2 \partial_t (L_{\parallel,\mu}^* P_{\eta t}^* \varphi) \Psi_\delta^2 t \, d\mu \, dt \\ &\quad + \frac{1}{2\eta^2} \iint_{\mathbb{R}_+^{n+1}} (u_1 \partial_t u_1) (\partial_t P_{\eta t}^* \varphi) \Psi_\delta^2 d\mu \, dt + \frac{1}{2\eta^2} \iint_{\mathbb{R}_+^{n+1}} u_1^2 (\partial_t P_{\eta t}^* \varphi) \Psi_\delta \partial_t \Psi_\delta d\mu \, dt \\ &=: S'_1 + S''_1 + S'''_1, \end{aligned}$$

where there is no boundary term because Ψ_δ vanishes on the boundary of \mathbb{R}_+^{n+1} .

To estimate S'''_1 , we use the definition of the set F in (4.17), the estimate for $|\nabla\Psi_\delta|$ from (4.27), and Remark 4.28 in the case $v_t \equiv 1$, to obtain

$$\begin{aligned} S'''_1 &\lesssim \|u\|_\infty^2 \iint_{\Omega_{\eta/8,Q}} N_*^\eta(\partial_t P_{\eta t}^* \varphi) |\partial_t \Psi_\delta| d\mu \, dt \\ &\lesssim \eta \kappa_0 \|u\|_\infty^2 \mu(Q) \lesssim \|u\|_\infty^2 \mu(Q). \end{aligned}$$

To estimate S'_1 , we observe that $\partial_t (L_{\parallel,\mu}^* P_{\eta t}^* \varphi) = L_{\parallel,\mu}^* (\partial_t P_{\eta t}^* \varphi)$, since $\varphi \in W_{\mu,0}^{1,2}(\mathbb{R}^n)$ and $\partial_t P_{\eta t}^* = -2\eta^2 t P_{\eta t}^* L_{\parallel,\mu}^*$ on the dense subset $\text{Dom}(L_{\parallel,\mu}^*)$ of $W_{0,\mu}^{1,2}(\mathbb{R}^n)$ (note also that $t \nabla_x P_{\eta t}^*$ and hence its adjoint are bounded operators on L_μ^2 , as can be seen from the proof of Theorem 2.12). We then apply Cauchy's

inequality with σ to write

$$\begin{aligned}
S'_1 &\leq \left| \iint_{\mathbb{R}_+^{n+1}} L_{\parallel, \mu}^* (\partial_t P_{\eta t}^* \varphi) u_1^2 \Psi_\delta^2 t \, d\mu \, dt \right| \\
&\lesssim \left| \iint_{\mathbb{R}_+^{n+1}} \left\langle \frac{1}{\mu} A_\parallel^* \nabla_x (\partial_t P_{\eta t}^* \varphi), \nabla_x u_1 \right\rangle u_1 \Psi_\delta^2 t \, d\mu \, dt \right| \\
&\quad + \left| \iint_{\mathbb{R}_+^{n+1}} \left\langle \frac{1}{\mu} A_\parallel^* \nabla_x (\partial_t P_{\eta t}^* \varphi), \nabla_x \Psi_\delta \right\rangle u_1^2 \Psi_\delta t \, d\mu \, dt \right| =: \mathbf{J} + \mathbf{K} \\
&\lesssim \sigma \iint_{\mathbb{R}_+^{n+1}} |\nabla_x u_1|^2 \Psi_\delta^2 t \, d\mu \, dt + (\sigma^{-1} + 1) \iint_{\mathbb{R}_+^{n+1}} u_1^2 |\nabla_x \partial_t P_{\eta t}^* \varphi|^2 \Psi_\delta^2 t \, d\mu \, dt \\
&\quad + \iint_{\mathbb{R}_+^{n+1}} u_1^2 |\nabla_x \Psi_\delta|^2 t \, d\mu \, dt =: \sigma S'_{11} + (\sigma^{-1} + 1) S'_{12} + S'_{13}, \quad (4.37)
\end{aligned}$$

where the integration by parts in x , with respect to the measure μ , is justified by the definition of the operator $L_{\parallel, \mu}^*$ (recall (2.8), (2.9) and (4.16)). The terms \mathbf{J} and \mathbf{K} are highlighted above for reference in Step 3.

To estimate S'_{13} , we use the estimate for $|\nabla \Psi_\delta|$ from (4.27) and Remark 4.28 in the case $v_t \equiv 1$ to obtain $S'_{13} \lesssim \|u\|_\infty^2 \mu(Q)$.

To estimate S'_{12} , we observe that $\nabla_x \partial_t P_{\eta t}^* = -2\eta^2 t \nabla_x L_{\parallel, \mu}^* P_{\eta t}^*$ on $L_\mu^2(\mathbb{R}^n)$ and then apply the vertical square function estimate from (2.14) followed by the $W_{0, \mu}^{1,2}(5Q)$ -Hodge estimate for φ from (4.14) to obtain

$$\begin{aligned}
S'_{12} &\lesssim \iint_{\mathbb{R}_+^{n+1}} u_1^2 |\nabla_x \partial_t P_{\eta t}^* \varphi|^2 \Psi_\delta^2 t \, d\mu \, dt \lesssim \|u\|_\infty^2 \iint_{\mathbb{R}_+^{n+1}} |t^2 \nabla_x L_{\parallel, \mu}^* P_{\eta t}^* \varphi|^2 \, d\mu \, \frac{dt}{t} \\
&\lesssim \|u\|_\infty^2 \|\nabla \varphi\|_{L_\mu^2(\mathbb{R}^n, \mathbb{R}^n)}^2 \lesssim \|u\|_\infty^2 \mu(Q).
\end{aligned}$$

The terms S'_{11} and S''_1 will now be estimated together. We again apply Cauchy's inequality with σ , followed by the vertical square function estimate from (2.13) with $\mathcal{L}_\mu = L_{\parallel, \mu}^*$ and the $W_{0, \mu}^{1,2}(5Q)$ -Hodge estimate for φ from (4.14) to obtain

$$\begin{aligned}
\sigma S'_{11} + S''_1 &\lesssim \sigma \iint_{\mathbb{R}_+^{n+1}} |\nabla_x u_1|^2 \Psi_\delta^2 t \, d\mu \, dt + \left| \iint_{\mathbb{R}_+^{n+1}} (u_1 \partial_t u_1) (\partial_t P_{\eta t}^* \varphi) \Psi_\delta^2 \, d\mu \, dt \right| \\
&\lesssim \sigma \iint_{\mathbb{R}_+^{n+1}} |\nabla u_1|^2 \Psi_\delta^2 t \, d\mu \, dt + \sigma^{-1} \|u\|_\infty^2 \iint_{\mathbb{R}_+^{n+1}} |\partial_t P_{\eta t}^* \varphi|^2 \, d\mu \, \frac{dt}{t} \\
&\lesssim \sigma \iint_{\mathbb{R}_+^{n+1}} \langle A_1 \nabla u_1, \nabla u_1 \rangle \Psi_\delta^2 t \, dx \, dt + \sigma \iint_{\mathbb{R}_+^{n+1}} |\nabla_x \tau|^2 |\partial_t u_1|^2 \Psi_\delta^2 t \, d\mu \, dt + \sigma^{-1} \|u\|_\infty^2 \mu(Q),
\end{aligned}$$

where we combined the pointwise estimates for ∇u_1 and J from (4.26) and (4.21) with identity (4.25) and the ellipticity of A to deduce the final inequality.

We use the dyadic decomposition from Remark 4.28 to write

$$\iint_{\mathbb{R}_+^{n+1}} |\nabla_x \tau|^2 |\partial_t u_1|^2 \Psi_\delta^2 t \, d\mu \, dt \leq \sum_{k \in \mathbb{Z}} \sum_{Q' \in \mathbb{D}_k^n} \int_{2^{-k}}^{2^{-k+1}} \int_{Q'} \mathbb{1}_{\Omega_{n, Q, \delta}} |\nabla_x \tau|^2 |\partial_t u_1|^2 t \, d\mu \, dt. \quad (4.38)$$

Observe that if $k \in \mathbb{Z}$, $Q' \in \mathbb{D}_k^\eta$ and $\Omega_{\eta/8, Q, \delta} \cap (Q' \times [2^{-k}, 2^{-k+1}]) \neq \emptyset$, then, as in (4.34) and (4.35), it holds that $Q' \times [2^{-k}, 2^{-k+1}] \subseteq \Omega_{\eta, 2Q, \delta/4}$ and

$$\int_{Q'} |\nabla_x \tau(x, t)|^2 d\mu(x) \lesssim \kappa_0^2 \quad \text{for all } t \in [2^{-k}, 2^{-k+1}].$$

Also, we have $\frac{7}{8}t < \tau(x, t) < \frac{9}{8}t$ and $J(x, t) \approx 1$ on $Q' \times [2^{-k}, 2^{-k+1}]$ by (4.21), so the degenerate version of Moser's estimate in (2.16) and t -independence show that

$$\sup_{x \in Q'} |\partial_t u_1(x, t)|^2 = \sup_{x \in Q'} |J(x, t) \partial_\tau u(x, \tau(x, t))|^2 \lesssim \int_{2Q'} \int_{t/2}^{2t} |\partial_s u(y, s)|^2 ds d\mu(y)$$

for all $t \in [2^{-k}, 2^{-k+1}]$. In particular, note that

$$2Q' \times [2^{-k-1}, 2^{-k+2}] \subseteq \Omega^* := \{(y, s) \in \mathbb{R}_+^{n+1} : \delta_F(y) < \frac{5}{8}\eta s, \frac{1}{2}\delta\ell(Q) < s < 8\ell(Q)\},$$

since there exists $(x_0, t_0) \in Q' \times [2^{-k}, 2^{-k+1}]$ satisfying $\delta_F(x_0) < \frac{1}{8}\eta t_0$, whence

$$\delta_F(y) < \text{diam}(2Q') + \frac{1}{8}\eta t_0 \leq \frac{5}{16}\eta 2^{-k} \leq \frac{5}{8}\eta s \quad \text{for all } y \in 2Q' \text{ and } s \geq 2^{-k-1},$$

whilst $\delta\ell(Q) < t_0 < 4\ell(Q)$ implies $[2^{-k}, 2^{-k+1}] \subseteq (\frac{1}{2}\delta\ell(Q), 8\ell(Q))$.

The observations in the preceding paragraph show that (4.38) is bounded by

$$\begin{aligned} & \sum_{k \in \mathbb{Z}} \sum_{Q' \in \mathbb{D}_k^\eta} \int_{2^{-k}}^{2^{-k+1}} \left(\int_{Q'} |\nabla_x \tau|^2 d\mu \right) \left(\int_{2Q'} \int_{t/2}^{2t} |\partial_s u(y, s)|^2 \mathbb{1}_{\Omega^*}(y, s) ds d\mu(y) \right) dt \\ & \lesssim \sum_{k \in \mathbb{Z}} \sum_{Q' \in \mathbb{D}_k^\eta} \int_{2^{-k-1}}^{2^{-k+2}} \int_{2Q'} |\partial_s u(y, s)|^2 \mathbb{1}_{\Omega^*}(y, s) s d\mu(y) ds \\ & \lesssim \left(\iint_{\Omega^{**}} |\partial_s u(y, s)|^2 s d\mu(y) ds + \iint_{\Omega^* \setminus \Omega^{**}} |\partial_s u(y, s)|^2 s d\mu(y) ds \right) := \mathbf{M} + \mathbf{E}, \end{aligned}$$

where we used the fact that $\sum_{k \in \mathbb{Z}} \sum_{Q' \in \mathbb{D}_k^\eta} \mathbb{1}_{2Q' \times [2^{-k-1}, 2^{-k+2}]} \lesssim \mathbb{1}_{\mathbb{R}_+^{n+1}}$ and introduced

$$\Omega^{**} := \{(y, s) \in \mathbb{R}_+^{n+1} : \delta_F(y) < \frac{1}{18}\eta s, 4\delta\ell(Q) < s < \ell(Q)\}.$$

To estimate the main term \mathbf{M} , we use (4.21)–(4.23) to observe that

$$\rho^{-1}(\Omega^{**}) \subseteq \Omega_{\eta/16} \cap (2Q \times (2\delta\ell(Q), 2\ell(Q))).$$

Thus, since $\Psi_\delta \equiv 1$ on these sets, the change of variables $(y, s) \mapsto \rho(y, s)$ gives

$$\mathbf{M} \lesssim \iint_{\mathbb{R}_+^{n+1}} |(\partial_t u) \circ \rho|^2 J \Psi_\delta^2 t d\mu dt \lesssim \iint_{\mathbb{R}_+^{n+1}} \langle A_1 \nabla u_1, \nabla u_1 \rangle \Psi_\delta^2 t dx dt,$$

where we used identity (4.25) and the ellipticity of A to deduce the final inequality.

To estimate the error term \mathbf{E} , recall that the degenerate version of Moser's estimate in (2.16), followed by Caccioppoli's inequality, ensures that $\|s \partial_s u\|_\infty \lesssim \|u\|_\infty$. Thus, by the definition of $\Omega^* \setminus \Omega^{**}$ and

the doubling property of μ , we obtain

$$E \lesssim \|u\|_\infty^2 \int_{2Q} \left(\int_{(8/5\eta)\delta_F(y)}^{(18/\eta)\delta_F(y)} \frac{ds}{s} + \int_{\ell(Q)}^{8\ell(Q)} \frac{ds}{s} + \int_{(\delta/2)\ell(Q)}^{4\delta\ell(Q)} \frac{ds}{s} \right) d\mu(y) \lesssim \|u\|_\infty^2 \mu(Q).$$

This shows that

$$\sigma S'_{11} + S''_1 \lesssim \sigma \iint_{\mathbb{R}_+^{n+1}} \langle A_1 \nabla u_1, \nabla u_1 \rangle \Psi_\delta^2 t \, dx \, dt + \sigma^{-1} \|u\|_\infty^2 \mu(Q);$$

hence

$$S_1 \lesssim \sigma \iint_{\mathbb{R}_+^{n+1}} \langle A_1 \nabla u_1, \nabla u_1 \rangle \Psi_\delta^2 t \, dx \, dt + \sigma^{-1} \|u\|_\infty^2 \mu(Q) \quad \text{for all } \sigma \in (0, 1). \quad (4.39)$$

Step 3: estimates for the term S_2 in (4.32).

We observe that since A is t -independent it is possible to write

$$\begin{aligned} 2S_2 &= \iint_{\mathbb{R}_+^{n+1}} u_1^2 \partial_t \left(\frac{\langle A\mathbf{p}, \mathbf{p} \rangle}{J} \right) \Psi_\delta^2 \, dx \, dt \\ &= \iint_{\mathbb{R}_+^{n+1}} u_1^2 \partial_t \left(\frac{1}{J} \right) \langle A\mathbf{p}, \mathbf{p} \rangle \Psi_\delta^2 \, dx \, dt \\ &\quad + \iint_{\mathbb{R}_+^{n+1}} \left(\frac{u_1^2}{J} \right) \langle \partial_t \mathbf{p}, A^* \mathbf{p} \rangle \Psi_\delta^2 \, dx \, dt + \iint_{\mathbb{R}_+^{n+1}} \left(\frac{u_1^2}{J} \right) \langle A\mathbf{p}, \partial_t \mathbf{p} \rangle \Psi_\delta^2 \, dx \, dt \\ &=: I + II + III. \end{aligned}$$

To estimate I , we recall the Jacobian $J(x, t) = 1 + \partial_t P_{\eta t}^* \varphi(x)$ from (4.20) and then integrate by parts in t to write

$$\begin{aligned} I &= - \iint_{\mathbb{R}_+^{n+1}} u_1^2 \frac{\partial_t^2 P_{\eta t}^* \varphi}{J^2} \langle A\mathbf{p}, \mathbf{p} \rangle \Psi_\delta^2 \, dx \, dt \\ &= \iint_{\mathbb{R}_+^{n+1}} \partial_t (u_1^2) \frac{\partial_t P_{\eta t}^* \varphi}{J^2} \langle A\mathbf{p}, \mathbf{p} \rangle \Psi_\delta^2 \, dx \, dt + \iint_{\mathbb{R}_+^{n+1}} u_1^2 \frac{\partial_t P_{\eta t}^* \varphi}{J^2} \partial_t (\langle A\mathbf{p}, \mathbf{p} \rangle) \Psi_\delta^2 \, dx \, dt \\ &\quad + \iint_{\mathbb{R}_+^{n+1}} u_1^2 \partial_t P_{\eta t}^* \varphi \partial_t (J^{-2}) \langle A\mathbf{p}, \mathbf{p} \rangle \Psi_\delta^2 \, dx \, dt + \iint_{\mathbb{R}_+^{n+1}} u_1^2 \frac{\partial_t P_{\eta t}^* \varphi}{J^2} \langle A\mathbf{p}, \mathbf{p} \rangle \partial_t (\Psi_\delta^2) \, dx \, dt \\ &=: I_1 + I_2 + I_3 + I_4, \end{aligned}$$

where there is no boundary term because Ψ_δ vanishes on the boundary of \mathbb{R}_+^{n+1} .

To estimate I_1 , we recall that $J \approx 1$ on $\text{supp } \Psi_\delta \subseteq \Omega_{\eta/8, Q, \delta}$ by (4.21) and then apply Cauchy's inequality with σ to obtain

$$\begin{aligned} |I_1| &\lesssim \sigma \iint_{\mathbb{R}_+^{n+1}} |\partial_t u_1|^2 |\mathbf{p}|^2 \Psi_\delta^2 t \, d\mu \, dt + \sigma^{-1} \iint_{\mathbb{R}_+^{n+1}} u_1^2 |\partial_t P_{\eta t} \varphi|^2 |\mathbf{p}|^2 \Psi_\delta^2 \, d\mu \, \frac{dt}{t} \\ &=: \sigma I'_1 + \sigma^{-1} I''_1. \end{aligned} \quad (4.40)$$

To estimate I'_1 , recall that $|\mathbf{p}|^2 = 1 + |\nabla_x \tau|^2$ by the definition of \mathbf{p} in (4.24), so we follow the treatment of (4.38) above to obtain

$$I'_1 \lesssim \iint_{\mathbb{R}_+^{n+1}} \langle A_1 \nabla u_1, \nabla u_1 \rangle \Psi_\delta^2 t \, dx \, dt + \|u\|_\infty^2 \mu(Q).$$

To estimate I''_1 , recall that $\|\mathbb{1}_{\Omega_{\eta,Q}} u_1\|_\infty \lesssim \|u\|_\infty$ and use the dyadic decomposition from Remark 4.28 to obtain

$$\begin{aligned} I''_1 &\lesssim \|u\|_\infty^2 \sum_{k \in \mathbb{Z}} \sum_{Q' \in \mathbb{D}_k^\eta} \|\partial_t P_{\eta t}^* \varphi\|_{L^\infty(Q' \times [2^{-k}, 2^{-k+1}])}^2 \int_{2^{-k}}^{2^{-k+1}} \int_{Q'} |\mathbf{p}|^2 \Psi_\delta^2 \, d\mu \, \frac{dt}{t} \\ &\lesssim \|u\|_\infty^2 \sum_{k \in \mathbb{Z}} \sum_{Q' \in \mathbb{D}_k^\eta} \mu(Q') \|\partial_t P_{\eta t}^* \varphi\|_{L^\infty(Q' \times [2^{-k}, 2^{-k+1}])}^2 \\ &\lesssim \|u\|_\infty^2 \sum_{k \in \mathbb{Z}} \sum_{Q' \in \mathbb{D}_k^\eta} \mu(Q') \int_{2^{-k-1}}^{2^{-k+2}} \int_{2Q'} |\partial_t P_{\eta t}^* \varphi|^2 \, d\mu \, dt \\ &\lesssim \|u\|_\infty^2 \sum_{k \in \mathbb{Z}} \sum_{Q' \in \mathbb{D}_k^\eta} \int_{2^{-k-1}}^{2^{-k+2}} \int_{2Q'} |\partial_t P_{\eta t}^* \varphi|^2 \, d\mu \, \frac{dt}{t} \\ &\lesssim \|u\|_\infty^2 \iint_{\mathbb{R}_+^{n+1}} |t L_{\parallel, \mu}^* e^{-t^2 L_{\parallel, \mu}^*} \varphi|^2 \, d\mu \, \frac{dt}{t} \\ &\lesssim \|u\|_\infty^2 \|\nabla \varphi\|_{L_\mu^2(\mathbb{R}^n, \mathbb{R}^n)}^2 \lesssim \|u\|_\infty^2 \mu(Q), \end{aligned} \quad (4.41)$$

where the second line uses the pointwise bound $|\mathbf{p}|^2 \Psi_\delta^2 \leq \mathbb{1}_{\Omega_{\eta/8, Q, \delta}} (1 + |\nabla_x (I - P_{\eta t}^*) \varphi|^2)$ and estimate (4.35), the third line uses the parabolic version of the degenerate Moser-type estimate in (2.16) (see Theorem B in [Fernandes 1991]), noting that $v := \partial_t (e^{-t^2 L_{\parallel, \mu}^*} \varphi)$ solves $\partial_t v = -L_{\parallel, \mu}^* v$, whilst $|\partial_t P_{\eta t}^* \varphi(x)| \lesssim |t v(x, \eta^2 t^2)|$, and the final line uses the vertical square function estimate from (2.13) with $\mathcal{L}_\mu = L_{\parallel, \mu}^*$ and the $W_{0, \mu}^{1,2}(5Q)$ -Hodge estimate for φ from (4.14).

To estimate I_2 , we again use the bound $J \approx 1$ on $\text{supp } \Psi_\delta \subseteq \Omega_{\eta/8, Q, \delta}$ from (4.21), and then recall the definition $\mathbf{p} := (\nabla_x (P_{\eta t}^* - I) \varphi, -1)$ from (4.24) to obtain

$$|I_2| \lesssim \iint_{\mathbb{R}_+^{n+1}} u_1^2 |\nabla_x \partial_t P_{\eta t}^* \varphi|^2 \Psi_\delta^2 t \, d\mu \, dt + \iint_{\mathbb{R}_+^{n+1}} u_1^2 |\partial_t P_{\eta t}^* \varphi|^2 |\mathbf{p}|^2 \Psi_\delta^2 \, d\mu \, \frac{dt}{t}. \quad (4.42)$$

The first integral in (4.42) is the same as S'_{12} from (4.37), whilst the second integral is the same as I''_1 from (4.40); hence $|I_2| \lesssim \|u\|_\infty^2 \mu(Q)$.

To estimate I_3 , we use the bound $|\partial_t P_{\eta t}^* \varphi| < \frac{1}{8}$ guaranteed by (4.18) to deduce that

$$|\partial_t (J^{-2})| = |\partial_t (1 + \partial_t P_{\eta t}^* \varphi)^{-2}| \lesssim |\partial_t^2 P_{\eta t}^* \varphi|$$

on $\text{supp } \Psi_\delta \subseteq \Omega_{\eta/8, Q, \delta}$ and write

$$|I_3| \lesssim \iint_{\mathbb{R}_+^{n+1}} u_1^2 |\partial_t P_{\eta t}^* \varphi|^2 |\mathbf{p}|^2 \Psi_\delta^2 \, d\mu \, \frac{dt}{t} + \iint_{\mathbb{R}_+^{n+1}} u_1^2 |\partial_t^2 P_{\eta t}^* \varphi|^2 |\mathbf{p}|^2 \Psi_\delta^2 t \, d\mu \, dt =: I'_3 + I''_3.$$

To estimate I'_3 , we note that it is the same as I''_1 from (4.40); thus $I'_3 \lesssim \|u\|_\infty^2 \mu(Q)$.

To estimate I''_3 , we follow the estimates and justification provided for (4.41), noting in addition that $\partial_t v = \partial_t^2(e^{-tL_{\parallel,\mu}^*} \varphi)$ solves $\partial_t(\partial_t v) = -L_{\parallel,\mu}^*(\partial_t v)$, to obtain

$$\begin{aligned}
I''_3 &\lesssim \|u\|_\infty^2 \sum_{k \in \mathbb{Z}} \sum_{Q' \in \mathbb{D}_k^\eta} \|\partial_t^2 P_{\eta t}^* \varphi\|_{L^\infty(Q' \times [2^{-k}, 2^{-k+1}])}^2 \int_{2^{-k}}^{2^{-k+1}} \int_{Q'} |p|^2 \Psi_\delta^2 d\mu \frac{dt}{t} \\
&\lesssim \|u\|_\infty^2 \sum_{k \in \mathbb{Z}} \sum_{Q' \in \mathbb{D}_k^\eta} \mu(Q') \| |t L_{\parallel,\mu}^* P_{\eta t}^* \varphi| + |t^2 \partial_t(L_{\parallel,\mu}^* P_{\eta t}^* \varphi)| \|_{L^\infty(Q' \times [2^{-k}, 2^{-k+1}])}^2 \\
&\lesssim \|u\|_\infty^2 \sum_{k \in \mathbb{Z}} \sum_{Q' \in \mathbb{D}_k^\eta} \mu(Q') \int_{2^{-k-1}}^{2^{-k+2}} \int_{2Q'} (|t L_{\parallel,\mu}^* P_{\eta t}^* \varphi|^2 + |t^2 \partial_t(L_{\parallel,\mu}^* P_{\eta t}^* \varphi)|^2) d\mu dt \\
&\lesssim \|u\|_\infty^2 \iint_{\mathbb{R}_+^{n+1}} |t L_{\parallel,\mu}^* P_{\eta t}^* \varphi|^2 d\mu \frac{dt}{t} + \|u\|_\infty^2 \iint_{\mathbb{R}_+^{n+1}} |t^2 \nabla_{x,t}(L_{\parallel,\mu}^* P_{\eta t}^* \varphi)|^2 d\mu \frac{dt}{t} \\
&\lesssim \|u\|_\infty^2 \|\nabla \varphi\|_{L_\mu^2(\mathbb{R}^n, \mathbb{R}^n)}^2 \lesssim \|u\|_\infty^2 \mu(Q),
\end{aligned}$$

where the second line uses $|\partial_t^2 P_{\eta t}^* \varphi| \lesssim |\partial_t(t L_{\parallel,\mu}^* P_{\eta t}^* \varphi)| \lesssim |L_{\parallel,\mu}^* P_{\eta t}^* \varphi| + |t \partial_t(L_{\parallel,\mu}^* P_{\eta t}^* \varphi)|$, the third line uses $|L_{\parallel,\mu}^* P_{\eta t}^* \varphi(x)| = |v(x, \eta^2 t^2)|$ and $|\partial_t(L_{\parallel,\mu}^* P_{\eta t}^* \varphi)(x)| \lesssim |t(\partial_t v)(x, \eta^2 t^2)|$, and the final line uses the vertical square function estimates from (2.13) and (2.14) with $\mathcal{L}_\mu = L_{\parallel,\mu}^*$; hence $|I_3| \lesssim \|u\|_\infty^2 \mu(Q)$.

To estimate I_4 , we use $|\partial_t P_{\eta t}^* \varphi| \lesssim 1$, $J \approx 1$ and $|p|^2 \leq (1 + |\nabla_x(I - P_{\eta t}^*)\varphi|^2)$, which hold on $\text{supp } \Psi_\delta \subseteq \Omega_{\eta/8, Q, \delta}$ by (4.18), (4.21) and (4.24), to reduce to the estimate obtained for $E'_1 + E_2$; hence $|I_4| \lesssim \|u\|_\infty^2 \mu(Q)$.

To estimate II , we use the definition $p := (\nabla_x(P_{\eta t}^* - I)\varphi, -1)$ from (4.24) to note that $\partial_t p = (\nabla_x \partial_t P_{\eta t}^* \varphi, 0)$ and use the Hodge decomposition from (4.13) to write

$$\langle \partial_t p, A^* p \rangle = \langle \nabla_x \partial_t P_{\eta t}^* \varphi, A_{\parallel}^* \nabla_x (P_{\eta t}^* - I)\varphi - c \rangle = \langle \nabla_x \partial_t P_{\eta t}^* \varphi, A_{\parallel}^* \nabla_x P_{\eta t}^* \varphi - \mu h \rangle \quad (4.43)$$

for all $x \in 5Q$ and $t > 0$. Using this and recalling that $\text{div}_\mu h = 0$, it follows that

$$\begin{aligned}
II &= \iint_{\mathbb{R}_+^{n+1}} \left(\frac{u_1^2}{J} \right) \langle \nabla_x \partial_t P_{\eta t}^* \varphi, A_{\parallel}^* \nabla_x P_{\eta t}^* \varphi - \mu h \rangle \Psi_\delta^2 dx dt \\
&= \iint_{\mathbb{R}_+^{n+1}} \left(\frac{u_1^2}{J} \right) (\partial_t P_{\eta t}^* \varphi) (L_{\parallel,\mu}^* P_{\eta t}^* \varphi) \Psi_\delta^2 d\mu dt \\
&\quad - \iint_{\mathbb{R}_+^{n+1}} \partial_t P_{\eta t}^* \varphi \langle \nabla_x \left(\frac{u_1^2}{J} \right), A_{\parallel}^* \nabla_x P_{\eta t}^* \varphi - \mu h \rangle \Psi_\delta^2 dx dt \\
&\quad - \iint_{\mathbb{R}_+^{n+1}} \left(\frac{u_1^2}{J} \right) \partial_t P_{\eta t}^* \varphi \langle \nabla_x (\Psi_\delta^2), A_{\parallel}^* \nabla_x P_{\eta t}^* \varphi - \mu h \rangle dx dt \\
&=: II_1 + II_2 + II_3,
\end{aligned} \quad (4.44)$$

where the integration by parts in x , with respect to the measure μ , is justified by the definition of the operator $L_{\parallel,\mu}^*$ (recall (2.8), (2.9) and (4.16)).

To estimate \mathbf{II}_1 , we use $J \approx 1$ and $L_{\parallel, \mu}^* P_{\eta t}^* \varphi = -(2\eta^2 t)^{-1} \partial_t P_{\eta t}^* \varphi$ to show that it can be treated the same way as \mathbf{I}_1'' in (4.40), without $|\mathbf{p}|^2$; hence $|\mathbf{II}_1| \lesssim \|u\|_\infty^2 \mu(Q)$.

To estimate \mathbf{II}_2 , we use $J \approx 1$,

$$|\nabla_x(J^{-1})| = |\nabla_x(1 + \partial_t P_{\eta t}^* \varphi)^{-1}| \lesssim |\nabla_x \partial_t P_{\eta t}^* \varphi|$$

and apply Cauchy's inequality with σ to obtain

$$\begin{aligned} |\mathbf{II}_2| &\lesssim \sigma \iint_{\mathbb{R}_+^{n+1}} |\nabla_x u_1|^2 \Psi_\delta^2 t \, d\mu \, dt + \iint_{\mathbb{R}_+^{n+1}} u_1^2 |\nabla_x \partial_t P_{\eta t}^* \varphi|^2 \Psi_\delta^2 t \, d\mu \, dt \\ &\quad + (\sigma^{-1} + 1) \iint_{\mathbb{R}_+^{n+1}} u_1^2 |\partial_t P_{\eta t}^* \varphi|^2 (|\nabla_x P_{\eta t}^* \varphi|^2 + |\mathbf{h}|^2) \Psi_\delta^2 \, d\mu \, \frac{dt}{t}. \end{aligned} \quad (4.45)$$

The first integral is the same as \mathbf{S}'_{11} from (4.37), whilst the remaining two integrals are the same as those that bound \mathbf{I}_2 in (4.42), except $(|\nabla_x P_{\eta t}^* \varphi|^2 + |\mathbf{h}|^2)$ replaces $|\mathbf{p}|^2$. This factor is controlled in the same way, however, since the Hodge decomposition in (4.13) implies

$$|\mathbf{h}|^2 = \left| \frac{1}{\mu} \mathbf{c} \mathbb{1}_{5Q} + \frac{1}{\mu} A_\parallel^* \nabla_x \varphi \right|^2 \lesssim 1 + |\nabla_x \varphi|^2;$$

hence by (4.35) we obtain

$$|\mathbf{II}_2| \lesssim \sigma \iint_{\mathbb{R}_+^{n+1}} \langle A_1 \nabla u_1, \nabla u_1 \rangle \Psi_\delta^2 t \, dx \, dt + \sigma^{-1} \|u\|_\infty^2 \mu(Q).$$

To estimate \mathbf{II}_3 , we use $J \approx 1$ and Cauchy's inequality to write

$$|\mathbf{II}_3| \lesssim \iint_{\mathbb{R}_+^{n+1}} u_1^2 |\nabla_x \Psi_\delta|^2 t \, d\mu \, dt + \iint_{\mathbb{R}_+^{n+1}} u_1^2 |\partial_t P_{\eta t}^* \varphi|^2 (|\nabla_x P_{\eta t}^* \varphi|^2 + |\mathbf{h}|^2) \Psi_\delta^2 \, d\mu \, \frac{dt}{t}.$$

The first term above is the same as \mathbf{S}'_{13} in (4.37), whilst the remaining term is the same as the last integral in (4.45); hence $|\mathbf{II}_3| \lesssim \|u\|_\infty^2 \mu(Q)$.

To estimate \mathbf{III} , we observe by analogy with (4.43) that

$$\begin{aligned} \langle A\mathbf{p}, \partial_t \mathbf{p} \rangle &= \langle A_\parallel \nabla_x (P_{\eta t}^* - I)\varphi - \mathbf{b}, \nabla_x \partial_t P_{\eta t}^* \varphi \rangle \\ &= \langle A_\parallel \nabla_x (P_{\eta t}^* \varphi - \varphi) + A_\parallel \nabla_x \tilde{\varphi} - \mu \tilde{\mathbf{h}}, \nabla_x \partial_t P_{\eta t}^* \varphi \rangle \\ &= \langle A_\parallel \nabla_x [(P_{\eta t}^* \varphi - \varphi) - (P_{\eta t} \tilde{\varphi} - \tilde{\varphi})] + A_\parallel \nabla_x P_{\eta t} \tilde{\varphi} - \mu \tilde{\mathbf{h}}, \nabla_x \partial_t P_{\eta t}^* \varphi \rangle \end{aligned}$$

for all $x \in 5Q$ and $t > 0$ and then write

$$\begin{aligned} \mathbf{III} &= \iint_{\mathbb{R}_+^{n+1}} \left(\frac{u_1^2}{J} \right) \langle \nabla_x [(P_{\eta t}^* \varphi - \varphi) - (P_{\eta t} \tilde{\varphi} - \tilde{\varphi})], A_\parallel^* \nabla_x \partial_t P_{\eta t}^* \varphi \rangle \Psi_\delta^2 \, dx \, dt \\ &\quad + \iint_{\mathbb{R}_+^{n+1}} \left(\frac{u_1^2}{J} \right) \langle A_\parallel \nabla_x P_{\eta t} \tilde{\varphi} - \mu \tilde{\mathbf{h}}, \nabla_x \partial_t P_{\eta t}^* \varphi \rangle \Psi_\delta^2 \, dx \, dt \\ &=: \mathbf{III}_1 + \mathbf{III}_2. \end{aligned}$$

To estimate \mathbf{III}_1 , we integrate by parts in x with respect to the measure μ to write

$$\begin{aligned} \mathbf{III}_1 &= \iint_{\mathbb{R}_+^{n+1}} \left(\frac{u_1^2}{J} \right) [(P_{\eta t}^* \varphi - \varphi) - (P_{\eta t} \tilde{\varphi} - \tilde{\varphi})] (L_{\parallel, \mu}^* \partial_t P_{\eta t}^* \varphi) \Psi_\delta^2 d\mu dt \\ &\quad - \iint_{\mathbb{R}_+^{n+1}} [(P_{\eta t}^* \varphi - \varphi) - (P_{\eta t} \tilde{\varphi} - \tilde{\varphi})] \langle \nabla_x \left(\frac{u_1^2 \Psi_\delta^2}{J} \right), A_\parallel^* \nabla_x \partial_t P_{\eta t}^* \varphi \rangle dx dt \\ &=: \mathbf{III}'_1 + \mathbf{III}''_1, \end{aligned}$$

which is justified by the definition of $L_{\parallel, \mu}^*$ (recall (2.8), (2.9) and (4.16)).

To estimate \mathbf{III}'_1 , we use Hardy's inequality (see, for instance, page 272 in [Stein 1970]) to observe, for the semigroups $\mathcal{P}_t \in \{e^{-t^2 L_{\parallel, \mu}^*}, e^{-t^2 L_{\parallel, \mu}}\}$, the estimate

$$\int_0^\infty |\mathcal{P}_{\eta t} f - f|^2 \frac{dt}{t^3} \leq \int_0^\infty \left(\int_0^{\eta t} |\partial_s \mathcal{P}_s f| ds \right)^2 \frac{dt}{t^3} \lesssim \int_0^\infty |\partial_t \mathcal{P}_t f|^2 \frac{dt}{t} \quad \text{for all } f \in L_\mu^2(\mathbb{R}^n).$$

We then recall that $\|\mathbb{1}_{\Omega_{\eta, Q}} u_1\|_\infty \lesssim \|u\|_\infty$ and $J \approx 1$ on $\text{supp } \Psi_\delta \subseteq \Omega_{\eta/8, Q, \delta}$ to obtain

$$\begin{aligned} |\mathbf{III}'_1| &\lesssim \|u\|_\infty^2 \int_{\mathbb{R}^n} \int_0^\infty (|P_{\eta t}^* \varphi - \varphi| + |P_{\eta t} \tilde{\varphi} - \tilde{\varphi}|) |L_{\parallel, \mu}^* \partial_t P_{\eta t}^* \varphi| dt d\mu \\ &\lesssim \|u\|_\infty^2 \int_{\mathbb{R}^n} \left(\int_0^\infty |P_{\eta t}^* \varphi - \varphi|^2 + |P_{\eta t} \tilde{\varphi} - \tilde{\varphi}|^2 \frac{dt}{t^3} \right)^{1/2} \left(\int_0^\infty |t^2 L_{\parallel, \mu}^* \partial_t P_{\eta t}^* \varphi|^2 \frac{dt}{t} \right)^{1/2} d\mu \\ &\lesssim \|u\|_\infty^2 \left(\iint_{\mathbb{R}_+^{n+1}} |\partial_t P_{\eta t}^* \varphi|^2 + |\partial_t P_{\eta t} \tilde{\varphi}|^2 d\mu \frac{dt}{t} \right)^{1/2} \left(\iint_{\mathbb{R}_+^{n+1}} |t^2 \partial_t L_{\parallel, \mu}^* P_{\eta t}^* \varphi|^2 d\mu \frac{dt}{t} \right)^{1/2} \\ &\lesssim \|u\|_\infty^2 (\|\nabla \varphi\|_{L_\mu^2(\mathbb{R}^n, \mathbb{R}^n)}^2 + \|\nabla \tilde{\varphi}\|_{L_\mu^2(\mathbb{R}^n, \mathbb{R}^n)}^2)^{1/2} \|\nabla \varphi\|_{L_\mu^2(\mathbb{R}^n, \mathbb{R}^n)} \lesssim \|u\|_\infty^2 \mu(Q), \end{aligned}$$

where the final line uses the vertical square function estimates from (2.13)–(2.14) for $\mathcal{L}_\mu \in \{L_{\parallel, \mu}^*, L_{\parallel, \mu}\}$ and the $W_{0, \mu}^{1,2}(5Q)$ -Hodge estimates for $\varphi, \tilde{\varphi}$ from (4.14)–(4.15).

To estimate \mathbf{III}''_1 , recall that $|P_{\eta t}^* \varphi - \varphi| \lesssim t$ and $|P_{\eta t} \tilde{\varphi} - \tilde{\varphi}| \lesssim t$ on $\text{supp } \Psi_\delta \subseteq \Omega_{\eta/8, Q, \delta}$ by (4.19), whilst $J \approx 1$ and $|\nabla_x (J^{-1})| \lesssim |\nabla_x \partial_t P_{\eta t}^* \varphi|$, so distributing ∇_x over u_1^2 , Ψ_δ^2 and $1/J$ yields terms that can be controlled in the same way as \mathbf{J} , \mathbf{K} and \mathbf{S}'_{12} in (4.37).

To estimate \mathbf{III}_2 , note that the estimates used to control φ and $P_{\eta t} \varphi$ also hold for $\tilde{\varphi}$ and $P_{\eta t} \tilde{\varphi}$ by (4.14)–(4.15) and (4.18)–(4.19), whilst $\text{div}_\mu \mathbf{h} = \text{div}_\mu \tilde{\mathbf{h}} = 0$ by (4.13); hence \mathbf{III}_2 can be estimated in the same way as \mathbf{II} in (4.44).

This gives

$$|\mathbf{III}''_1| + |\mathbf{III}_2| \lesssim \sigma \iint_{\mathbb{R}_+^{n+1}} \langle A_1 \nabla u_1, \nabla u_1 \rangle \Psi_\delta^2 t dx dt + \sigma^{-1} \|u\|_\infty^2 \mu(Q);$$

hence

$$\mathbf{S}_2 \lesssim \sigma \iint_{\mathbb{R}_+^{n+1}} \langle A_1 \nabla u_1, \nabla u_1 \rangle \Psi_\delta^2 t dx dt + \sigma^{-1} \|u\|_\infty^2 \mu(Q) \quad \text{for all } \sigma \in (0, 1). \quad (4.46)$$

We combine (4.36), (4.39) and (4.46) to obtain (4.33), as required. \square

5. Solvability of the Dirichlet problem

This section is dedicated to the proof of Theorem 1.2. We first consider the construction and properties of a degenerate elliptic measure ω^X for degenerate elliptic equations $\operatorname{div}(A\nabla u) = 0$ in the upper half-space, where $X = (x, t) \in \mathbb{R}_+^{n+1}$ and $n \geq 2$. The t -independent coefficient matrix A is assumed throughout to satisfy the degenerate bound and ellipticity in (1.1) for some constants $0 < \lambda \leq \Lambda < \infty$ and an A_2 -weight μ on \mathbb{R}^n . This is necessary as the literature only seems to treat bounded domains, whilst the passage to unbounded domains in the uniformly elliptic case (see Section 10 in [Littman et al. 1963] and [Hofmann and Kim 2007]) relies on a global version of the Sobolev embedding in (2.4), which is not known for A_2 -weights in general. The degenerate elliptic measure is then shown to be in the A_∞ -class with respect to μ on the boundary \mathbb{R}^n in Theorem 5.30 and the solvability of the Dirichlet problem follows in Theorem 5.34. These results together prove Theorem 1.2.

5A. Boundary estimates for solutions. We require some estimates for solutions near the boundary $\partial\Sigma$ of a bounded Lipschitz domain $\Sigma \subset \mathbb{R}^n$ (see Section 2 of [Caffarelli et al. 1981] for the standard definition). These estimates require some regularity on the domain boundary but no attempt is made here to obtain the minimal such regularity, as the focus is to define and analyse a degenerate elliptic measure on \mathbb{R}^n .

The Lipschitz regularity of the boundary $\partial\Sigma$ ensures that the smooth class $C^\infty(\bar{\Sigma})$ and the Lipschitz class $C^{0,1}(\bar{\Sigma})$ are both dense in $W_\mu^{1,2}(\Sigma)$; see Theorem 3.4.1 in [Morrey 1966] and page 29 in [Kinderlehrer and Stampacchia 1980]. This allows the usual definition, for $E \subseteq \partial\Sigma$ and $u \in W_\mu^{1,2}(\Sigma)$, whereby $u \geq 0$ on E in the $W_\mu^{1,2}(\Sigma)$ -sense means there exists a sequence u_j in $C^{0,1}(\bar{\Sigma})$ that converges to u in $W_\mu^{1,2}(\Sigma)$ with $u_j(x) \geq 0$ for all $x \in E$. This induces definitions for inequalities \leq , \geq and $=$, between functions and/or constants, on E in the $W_\mu^{1,2}(\Sigma)$ -sense; see, for instance, Definition 5.1 in [Kinderlehrer and Stampacchia 1980]. Moreover, with $\sup_{\partial\Sigma} u := \inf\{k \in \mathbb{R} : u \leq k \text{ on } \partial\Sigma \text{ in the } W_\mu^{1,2}(\Sigma)\text{-sense}\}$ and $\inf_{\partial\Sigma} := -\sup_{\partial\Sigma}(-u)$, the weak maximum principle holds [Fabes et al. 1982b, Theorem 2.2.2], and the strong version follows by the Harnack inequality in (2.18) [Fabes et al. 1982b, Corollary 2.3.10].

We can now state a Hölder continuity estimate and a Harnack inequality for certain solutions near the boundary. For a cube $Q \subset \mathbb{R}^n$, recall the corkscrew point $X_Q := (x_Q, \ell(Q))$ and denote the Carleson box in \mathbb{R}_+^{n+1} by $T_Q := Q \times (0, \ell(Q))$. Also, recall that $\mu(x, t) := \mu(x)$, so $d\mu(x, t) = \mu(x) dx dt$, for $(x, t) \in \mathbb{R}^{n+1}$. If $u \in W_\mu^{1,2}(T_{2Q})$ is a solution of $\operatorname{div}(A\nabla u) = 0$ in T_{2Q} , and $u = 0$ on $2Q$ in the $W_\mu^{1,2}(T_{2Q})$ -sense, then

$$|u(x, t)| \lesssim \left(\frac{t}{\ell(Q)}\right)^\alpha \left(\int_{T_{2Q}} |u|^2 d\mu\right)^{1/2} \quad \text{for all } (x, t) \in T_Q, \quad (5.1)$$

and if, in addition, $u \geq 0$ almost everywhere on T_{2Q} , then

$$u(X) \lesssim u(X_Q) \quad \text{for all } X \in T_Q, \quad (5.2)$$

where α is from (2.17) and the implicit constants depend only on n , λ , Λ and $[\mu]_{A_2}$. Estimate (5.1) follows from standard reflection arguments and the interior Hölder continuity estimate in (2.17), as observed on page 102 in [Fabes et al. 1982b]. Estimate (5.2) can then be deduced from (5.1) and the interior Harnack

inequality in (2.18), as in the uniformly elliptic case; see the proof of Theorem 1.1 in [Caffarelli et al. 1981], which does not use the assumption therein that A is symmetric.

5B. Definition and properties of degenerate elliptic measure. For $X \in \mathbb{R}^{n+1}$, $x \in \mathbb{R}^n$ and $r > 0$, we use $B(X, r) := \{Y \in \mathbb{R}^{n+1} : |Y - X| < r\}$ to denote balls in \mathbb{R}^{n+1} and $\Delta(x, r) := \{y \in \mathbb{R}^n : |x| < r\}$ to denote balls in \mathbb{R}^n , where $\Delta(x, r)$ is identified with the surface ball $B((x, 0), r) \cap \partial\mathbb{R}_+^{n+1}$ in \mathbb{R}^{n+1} . For each $R > 0$, consider the bounded Lipschitz domain $\Sigma_R := B(0, R) \cap \mathbb{R}_+^{n+1}$ with Lipschitz constant at most 1. For each $X \in \Sigma_R$, the degenerate elliptic measure ω_R^X is the measure on $\partial\Sigma_R$, as defined on page 583 in [Fabes et al. 1983], such that $u(X) = \int_{\partial\Sigma_R} h d\omega_R^X$ solves the Dirichlet problem for continuous boundary data $h \in C(\partial\Sigma_R)$ in the sense that $\operatorname{div}(A\nabla u) = 0$ in Σ_R and $u \in C(\bar{\Sigma}_R)$ with $u|_{\partial\Sigma_R} = h$.

We now define the degenerate elliptic measure on \mathbb{R}^n . If $f \in C_c(\mathbb{R}^n)$, fix $R_0 > 0$ such that $\operatorname{supp} f \subseteq \Delta(0, R_0)$ and set f equal to zero on \mathbb{R}_+^{n+1} , so then $f^\pm \in C(\partial\Sigma_R)$ for all $R \geq R_0$, where $f^\pm(X) := \max\{\pm f(X), 0\}$; thus

$$u_R^\pm(X) := \int_{\partial\Sigma_R} f^\pm d\omega_R^X \quad \text{for all } X \in \Sigma_R$$

solve the Dirichlet problem as above in Σ_R for all $R \geq R_0$. The maximum principle then implies that $u_{R_1}^\pm(X) \leq u_{R_2}^\pm(X)$, whenever $R_0 \leq R_1 \leq R_2$ and $X \in \Sigma_{R_1}$, and that $\sup_{R>0} \|u_R^\pm\|_\infty \leq \|f\|_\infty$. This allows us to define

$$u(X) := \lim_{R \rightarrow \infty} [u_R^+(X) - u_R^-(X)] \quad \text{for all } X \in \mathbb{R}_+^{n+1}, \quad (5.3)$$

and since the mapping $f \mapsto u(X)$ is a positive linear functional on $C_c(\mathbb{R}^n)$, the Riesz representation theorem implies that there exists a regular Borel probability measure (the degenerate elliptic measure) ω^X on \mathbb{R}^n such that $u(X) = \int_{\mathbb{R}^n} f d\omega^X$.

The function u from (5.3) solves $\operatorname{div}(A\nabla u) = 0$ in \mathbb{R}_+^{n+1} . To prove this, note that $\|u\|_\infty \leq \|f\|_\infty$, so for each compact set $K \subset \mathbb{R}_+^{n+1}$, the Hölder continuity of solutions in (2.17) ensures the equicontinuity required to apply the Arzelà–Ascoli theorem and extract a subsequence u_{R_j} that converges to u uniformly on K . This combined with Caccioppoli’s inequality shows that u_{R_j} converges to u in $W_{\mu}^{1,2}(K)$; hence $u \in W_{\mu, \text{loc}}^{1,2}(\mathbb{R}_+^{n+1})$. Moreover, if $\varphi \in C_c^\infty(\mathbb{R}_+^{n+1})$ and $K = \operatorname{supp} \varphi \subset \Sigma_R$, then

$$\left| \int_K \langle A\nabla(u - u_R), \nabla\varphi \rangle \right| \leq \Lambda \|\nabla\varphi\|_\infty \mu(K)^{1/2} \|u - u_R\|_{W_{\mu}^{1,2}(K)}, \quad (5.4)$$

from which it follows that $\int_{\mathbb{R}_+^{n+1}} \langle A\nabla u, \nabla\varphi \rangle = 0$, as required.

We note by (5.3) that, when restricted to any bounded Borel subset of \mathbb{R}^n , the measures ω_R^X converge weakly to ω^X , so Theorem 1 on page 54 of [Evans and Gariepy 1992] shows that

$$\omega^X(U) \leq \liminf_{R \rightarrow \infty} \omega_R^X(U), \quad \omega^X(K) \geq \limsup_{R \rightarrow \infty} \omega_R^X(K), \quad \omega^X(B) = \lim_{R \rightarrow \infty} \omega_R^X(B) \quad (5.5)$$

for all bounded open sets $U \subset \mathbb{R}^n$, all compact sets $K \subset \mathbb{R}^n$, and all bounded Borel sets $B \subset \mathbb{R}^n$ such that $\omega^X(\partial B) = 0$. This construction of the degenerate elliptic measure also provides for the following expected properties.

Lemma 5.6. *If $X_0, X_1 \in \mathbb{R}_+^{n+1}$ and $E \subseteq \mathbb{R}^n$ is a Borel set, then $\omega^{X_0}(E) = 0$ if and only if $\omega^{X_1}(E) = 0$. Moreover, the nonnegative function $u(X) := \omega^X(E)$ is a solution of $\operatorname{div}(A \nabla u) = 0$ in \mathbb{R}_+^{n+1} and the boundary Hölder continuity estimate*

$$|u(x, t)| \lesssim \left(\frac{t}{\ell(Q)} \right)^\alpha u(X_Q) \quad \text{for all } (x, t) \in T_Q \quad (5.7)$$

holds on all cubes Q such that $2Q \subseteq \mathbb{R}^n \setminus E$, where α is from (2.17) and the implicit constants depend only on n, λ, Λ and $[\mu]_{A_2}$,

Proof. The proof follows that of Lemma 1.2.7 in [Kenig 1994], except we must account for the fact that the solution to the Dirichlet problem in \mathbb{R}_+^{n+1} defined by (5.3) requires boundary data to have compact support, which is easily done as we now show. Suppose that $\omega^{X_0}(E) = 0$ and that $K \subseteq E$ is a compact set. The regularity of the measure implies that $\omega^{X_0}(K) = 0$ and, for each $\epsilon > 0$, there exists a bounded open set $U \supset K$ such that $\omega^{X_0}(U) < \epsilon$. In particular, we may assume that U is bounded because K is compact, so by Urysohn's lemma there exists $g \in C_c(\mathbb{R}^n)$ such that $g(x) = 1$ on K , $0 \leq g(x) \leq 1$ on U , and $\operatorname{supp} g \subset U$. It follows that $u(X) = \int_{\mathbb{R}^n} g d\omega^X$ is the solution to the Dirichlet problem in \mathbb{R}_+^{n+1} defined by (5.3) with boundary data g . Applying the Harnack inequality from (2.18) and connecting X_0 with X_1 via a Harnack chain then shows that there exists $C > 0$, depending on X_0 and X_1 , such that

$$\omega^{X_1}(K) \leq u(X_1) \leq C u(X_0) \leq C \omega^{X_0}(U) \leq C\epsilon \quad \text{for all } \epsilon > 0;$$

hence $\omega^{X_1}(K) = 0$ for all compact sets $K \subseteq E$, and so $\omega^{X_1}(E) = 0$ by regularity.

The proof that $u(X) := \omega^X(E)$ is a solution of $\operatorname{div}(A \nabla u) = 0$ in \mathbb{R}_+^{n+1} also follows that of Lemma 1.2.7 in [Kenig 1994]. It remains to prove that the boundary Hölder continuity estimate holds on all cubes Q such that $2Q \subseteq \mathbb{R}^n \setminus E$. We first consider when E is bounded. In that case, let U_δ denote the open δ -neighbourhood of E and set $\chi_{\epsilon, \delta} := \varphi_\epsilon * \mathbb{1}_{U_\delta}$ for all $\delta > \epsilon > 0$, where $\varphi_\epsilon(x) := \epsilon^{-n} \varphi(x/\epsilon)$ and $\varphi \in C_c^\infty(\Delta(0, 1))$ is a fixed nonnegative function with $\int_{\mathbb{R}^n} \varphi = 1$. In particular, since U_δ is open, we have $\mathbb{1}_E \leq \mathbb{1}_{U_\delta} \leq \liminf_{\epsilon \rightarrow 0} \chi_{\epsilon, \delta}$. Consequently, if $X = (x, t) \in \mathbb{R}_+^{n+1}$, then

$$u(X) = \omega^X(E) \leq \omega^X(U_\delta) \leq \int_{\mathbb{R}^n} \liminf_{\epsilon \rightarrow 0} \chi_{\epsilon, \delta} d\omega^X \leq \liminf_{\epsilon \rightarrow 0} \int_{\mathbb{R}^n} \chi_{\epsilon, \delta} d\omega^X. \quad (5.8)$$

The function $\chi_{\epsilon, \delta}$ belongs to $C_c^\infty(\mathbb{R}^n)$ and thus extends to a function in $C_c^\infty(\mathbb{R}^{n+1})$. The construction of the degenerate elliptic measure (see pages 580–583 in [Fabes et al. 1983], which was the starting point for our extension to the upper half-space above) thus implies $v_\epsilon(X) := \int_{\mathbb{R}^n} \chi_{\epsilon, \delta} d\omega^X$ is in $W^{1,2}(T_{(3/2)Q})$ and vanishes on $\frac{3}{2}Q$ whenever $0 < \epsilon < \delta < \frac{1}{4}\ell(Q)$, so estimate (5.8) combined with the boundary Hölder continuity estimate in (5.1) and the boundary Harnack inequality in (5.2) shows that

$$u(x, t) \leq \liminf_{\epsilon \rightarrow 0} v_\epsilon(x, t) \lesssim \left(\frac{t}{\ell(Q)} \right)^\alpha \liminf_{\epsilon \rightarrow 0} v_\epsilon(X_Q) \quad \text{for all } (x, t) \in T_Q. \quad (5.9)$$

We now let $U_{\delta, \epsilon}$ denote the open ϵ -neighbourhood of U_δ , in which case $\chi_{\epsilon, \delta} \leq \mathbb{1}_{U_{\delta, \epsilon}}$ and $v_\epsilon(X) \leq \omega^X(U_{\delta, \epsilon})$, so by (5.9) and the regularity of the degenerate elliptic measure we have

$$u(x, t) \lesssim \left(\frac{t}{\ell(Q)} \right)^\alpha \liminf_{\epsilon \rightarrow 0} \omega^{X_Q}(U_{\delta, \epsilon}) \lesssim \left(\frac{t}{\ell(Q)} \right)^\alpha \omega^{X_Q}(U_\delta) \quad \text{for all } (x, t) \in T_Q.$$

This proves (5.7) if E is bounded, since the regularity of the measure also implies that $\omega^{X_Q}(U_\delta)$ approaches $\omega^{X_Q}(E) = u(X_Q)$ as δ approaches 0. If E is not bounded, then applying (5.7) on the bounded sets $E_k := \mathbb{1}_{2^{k+1}Q \setminus 2^k Q} E$, for $k \in \mathbb{N}$, shows that

$$u(x, t) = \sum_{k=1}^{\infty} \omega^X(E_k) \lesssim \sum_{k=1}^{\infty} \left(\frac{t}{\ell(Q)} \right)^\alpha \omega^{X_Q}(E_k) = \left(\frac{t}{\ell(Q)} \right)^\alpha \omega^{X_Q}(E) \quad \text{for all } (x, t) \in T_Q,$$

as required. \square

5C. Preliminary estimates for degenerate elliptic measure. In the uniformly elliptic case, there is a rich theory for the Green's function on bounded domains, and specifically, estimates and connections with elliptic measure; see, for instance, Theorem 1.2.8 and Corollary 1.3.6 in [Kenig 1994]. This theory also extends to unbounded domains; see Section 10 in [Littman et al. 1963] and [Hofmann and Kim 2007]. In the degenerate elliptic case, the theory was developed on bounded domains in [Fabes et al. 1982a; 1982b; 1983], but it is not clear if there is always such a Green's function on unbounded domains. In particular, the construction in [Hofmann and Kim 2007] for the uniformly elliptic case relies on the (unweighted) global version of the Sobolev embedding in (2.4), which is not known for a general A_2 -weight. In what follows, we combine the properties of the Green's function on the bounded domain $\Sigma_R := B(0, R) \cap \mathbb{R}_+^{n+1}$ with the limit properties in (5.5) to deduce estimates for degenerate elliptic measure on \mathbb{R}^n . These will be used to prove Lemma 5.24 and ultimately Theorem 5.30.

For each $R > 0$, the Green's function $g_R : \bar{\Sigma}_R \times \bar{\Sigma}_R \mapsto [0, \infty]$ is constructed by following Proposition 2.4 in [Fabes et al. 1982a]. In particular, for each $Y \in \Sigma_R$, the mapping $X \mapsto g_R(X, Y)$ is the Hölder continuous function in $\bar{\Sigma}_R \setminus \{Y\}$ that vanishes on $\partial \Sigma_R$ and satisfies $\int_{\Sigma_R} \langle A \nabla g_R(\cdot, Y), \nabla \Phi \rangle = \Phi(Y)$ for all $\Phi \in C_c^\infty(\Sigma_R)$. As explained on page 583 in [Fabes et al. 1983], these properties are valid on any NTA domain, hence a fortiori on Σ_R . The proofs do not rely on the assumption therein that A is symmetric, although the symmetry property $g_R(X, Y) = g_R(Y, X)$ is no longer guaranteed, as $g_R^*(X, Y) := g_R(Y, X)$ is the Green's function for the adjoint operator $-\operatorname{div}(A^* \nabla)$. We will rely on the following two lemmas, which are immediate from Theorem 4 and Lemma 3 in [Fabes et al. 1983], respectively, to estimate the Green's function g_R and the degenerate elliptic measure ω_R on Σ_R .

Lemma 5.10. *If $X, Y \in \Sigma_R$ and $|X - Y| < \frac{1}{2} \operatorname{dist}(Y, \partial \Sigma_R)$, then*

$$g_R(X, Y) \approx \int_{|X-Y|}^{\operatorname{dist}(Y, \partial \Sigma_R)} \frac{s^2}{\mu(B(Y, s))} \frac{ds}{s},$$

where the implicit constants depend only on n, λ, Λ and $[\mu]_{A_2}$.

Lemma 5.11. *If $R > 0$ and Q is a cube in \mathbb{R}^n such that $T_2 Q \subset \Sigma_R$, then*

$$\frac{g_R(X_Q, Y)}{\ell(Q)} \approx \omega_R^Y(Q) \frac{\ell(Q)}{\mu(T_Q)} = \frac{\omega_R^Y(Q)}{\mu(Q)} \quad \text{for all } Y \in \Sigma_R \setminus T_2 Q,$$

where the implicit constants depend only on n, λ, Λ and $[\mu]_{A_2}$.

The degenerate elliptic measure ω_R^X satisfies the doubling property $\omega_R^X(2Q) \leq C_0 \omega_R^X(Q)$ for all cubes Q in \mathbb{R}^n such that $T_2Q \subset \Sigma_R$ and all $X \in \Sigma_R \setminus T_2Q$, where the doubling constant $C_0 > 0$ depends only on n, λ, Λ and $[\mu]_{A_2}$. This is proved in Lemma 1 on page 584 of [Fabes et al. 1983] by using the estimates in Lemma 5.11, the Harnack inequality in (2.18), and the doubling property of μ . The doubling constant C_0 does not depend on R , which allows us to use the inequalities in (5.5) to show that the degenerate elliptic measure ω^X is locally doubling on \mathbb{R}^n , in the sense that

$$\omega^X(2Q) \leq \liminf_{R \rightarrow \infty} \omega_R^X(2Q) \lesssim \liminf_{R \rightarrow \infty} \omega_R^X\left(\frac{1}{2}Q\right) \leq \limsup_{R \rightarrow \infty} \omega_R^X\left(\frac{1}{2}\bar{Q}\right) \leq \omega^X(Q) \quad (5.12)$$

for all cubes $Q \subset \mathbb{R}^n$ and all $X \in \mathbb{R}_+^{n+1} \setminus T_2Q$, where the implicit constant is C_0^2 . In particular, the doubling property implies $\omega^X(\partial Q) = 0$ for all cubes $Q \subset \mathbb{R}^n$ (see page 403 in [García-Cuerva and Rubio de Francia 1985] or Proposition 6.3 in [Hofmann and Martell 2014]), so (5.12) actually improves to $\omega^X(2Q) \leq C_0 \omega^X(Q)$, since by the equality in (5.5) we now have

$$\omega^X(Q) = \lim_{R \rightarrow \infty} \omega_R^X(Q) \quad (5.13)$$

for all cubes $Q \subset \mathbb{R}^n$ and all $X \in \mathbb{R}_+^{n+1} \setminus T_2Q$. This provides the following estimate for degenerate elliptic measure.

Lemma 5.14. *If Q is a cube in \mathbb{R}^n , then $\omega^{X_Q}(Q) \gtrsim 1$, where the implicit constant depends only on n, λ, Λ and $[\mu]_{A_2}$.*

Proof. Let Q denote a cube in \mathbb{R}^n and fix $R_0 > 0$ such that $T_2Q \subset \Sigma_{R_0}$. The Hölder continuity at the boundary in (5.1) and the Harnack inequality in (2.18) imply (see the proof of Lemma 3 on page 585 in [Fabes et al. 1983]) that

$$\omega_R^{X_Q}(Q) \gtrsim 1 \quad \text{for all } R \geq R_0,$$

where the implicit constant depends only on n, λ, Λ and $[\mu]_{A_2}$, and so does not depend on R . The result follows by using Harnack's inequality to shift the pole (from X_{2Q} to X_Q) in (5.12)–(5.13) to obtain $\omega^{X_Q}(Q) = \lim_{R \rightarrow \infty} \omega_R^{X_Q}(Q) \gtrsim 1$. \square

The estimates in Lemma 5.11 also imply the following comparison principle. The result is stated on page 585 in [Fabes et al. 1983] and the proof is the same as in the uniformly elliptic case; see Theorem 1.4 in [Caffarelli et al. 1981] or Lemma 1.3.7 in [Kenig 1994], neither of which use the assumption therein that A is symmetric.

Lemma 5.15 (comparison principle). *Let Q denote a cube in \mathbb{R}^n and suppose that $u, v \in W_\mu^{1,2}(T_2Q) \cap C(\bar{T}_2Q)$ with $u, v \geq 0$ on T_2Q . If $\operatorname{div}(A\nabla u) = \operatorname{div}(A\nabla v) = 0$ in T_2Q and $u = v = 0$ on $2Q$, then*

$$\frac{u(X)}{v(X)} \approx \frac{u(X_Q)}{v(X_Q)} \quad \text{for all } X \in T_Q,$$

where the implicit constants depend only on n, λ, Λ and $[\mu]_{A_2}$.

The following corollary of these preliminaries will be used in Proposition 5.18 to estimate Radon–Nikodym derivatives of the degenerate elliptic measure.

Lemma 5.16. *If Q_0 and Q are cubes in \mathbb{R}^n such that $Q \subseteq Q_0$, then*

$$\omega^{X_{Q_0}}(Q) \asymp \frac{\omega^X(Q)}{\omega^X(Q_0)} \quad \text{for all } X \in \mathbb{R}_+^{n+1} \setminus T_{2Q_0},$$

where the implicit constants depend only on n, λ, Λ and $[\mu]_{A_2}$.

Proof. Let $Q \subset Q_0$ be cubes in \mathbb{R}^n , suppose that $X \in \mathbb{R}_+^{n+1} \setminus T_{2Q_0}$ and consider $R > 0$ large enough so that $X \in \Sigma_R$ and $T_{4Q_0} \subset \Sigma_R$. Lemma 5.11 shows that

$$\begin{aligned} \omega_R^X(Q_0) \ell(Q_0) &\asymp \mu(Q_0) g_R(X_{Q_0}, X), \\ \omega_R^X(Q) \ell(Q) &\asymp \mu(Q) g_R(X_Q, X) \\ \omega_R^{X_{3Q_0}}(Q) \ell(Q) &\asymp \mu(Q) g_R(X_Q, X_{3Q_0}). \end{aligned}$$

If $u(Y) = g_R(Y, X)$ and $v(Y) = g_R(Y, X_{3Q_0})$, then $\operatorname{div}(A \nabla u) = \operatorname{div}(A \nabla v) = 0$ in T_{2Q_0} and $u = v = 0$ on $2Q_0$, so the comparison principle in Lemma 5.15 shows that

$$\frac{g_R(X_Q, X)}{g_R(X_Q, X_{3Q_0})} = \frac{u(X_Q)}{v(X_Q)} \asymp \frac{u(X_{Q_0})}{v(X_{Q_0})} = \frac{g_R(X_{Q_0}, X)}{g_R(X_{Q_0}, X_{3Q_0})}.$$

Also, Lemma 5.10 shows that $g_R(X_{Q_0}, X_{3Q_0}) \asymp \ell(Q_0)/\mu(Q_0)$, so together we obtain

$$\frac{\omega_R^X(Q)}{\omega_R^X(Q_0)} \asymp \frac{g_R(X_Q, X)}{g_R(X_{Q_0}, X)} \frac{\mu(Q)}{\mu(Q_0)} \frac{\ell(Q_0)}{\ell(Q)} \asymp \frac{g_R(X_Q, X_{3Q_0})}{g_R(X_{Q_0}, X_{3Q_0})} \frac{\mu(Q)}{\mu(Q_0)} \frac{\ell(Q_0)}{\ell(Q)} \asymp \omega_R^{X_{3Q_0}}(Q).$$

The Harnack inequality from (2.18) then shows that $\omega_R^X(Q) \asymp \omega_R^X(Q_0) \omega_R^{X_{Q_0}}(Q)$ and the result follows by using (5.13) to estimate the limit as R approaches infinity. \square

If $X, X_0 \in \mathbb{R}_+^{n+1}$, then Lemma 5.6 shows that ω^X and ω^{X_0} are mutually absolutely continuous, so the Lebesgue differentiation theorem for the locally doubling measure ω^{X_0} implies that the Radon–Nikodym derivative of ω^X satisfies

$$K(X_0, X, y) := \frac{d\omega^X}{d\omega^{X_0}}(y) = \lim_{s \rightarrow 0} \frac{\omega^X(Q(y, s))}{\omega^{X_0}(Q(y, s))}, \quad \omega^{X_0}\text{-a.e. } y \in \mathbb{R}^n, \quad (5.17)$$

where $Q(y, s)$ denotes the cube in \mathbb{R}^n with centre y and side length s . The following decay estimate for the kernel function K extends Lemma 2 on page 584 in [Fabes et al. 1983]. It is the final property of degenerate elliptic measure needed to prove Lemma 5.24.

Proposition 5.18. *If Q_0 and Q are cubes in \mathbb{R}^n such that $Q \subseteq Q_0$, then*

$$K(X_{Q_0}, X_Q, y) \lesssim \frac{1}{\omega^{X_{Q_0}}(Q)} \max \left\{ \frac{|y - x_Q|}{\ell(Q)}, 1 \right\}^{-\alpha}, \quad \omega^{X_{Q_0}}\text{-a.e. } y \in Q_0,$$

where $\alpha > 0$ from (2.17) and the implicit constant depend only on n, λ, Λ and $[\mu]_{A_2}$.

Proof. Let $Q \subseteq Q_0$ denote cubes in \mathbb{R}^n and fix $J \in \mathbb{N}$ such that $2^{J-1}Q \subseteq Q_0 \subseteq 2^JQ$. If $y \in Q$, then Lemma 5.16 and the Harnack inequality in (2.18) show that

$$\omega^{X_Q}(Q(y, s)) \approx \frac{\omega^{X_{2Q_0}}(Q(y, s))}{\omega^{X_{2Q_0}}(Q)} \approx \frac{\omega^{X_{Q_0}}(Q(y, s))}{\omega^{X_{Q_0}}(Q)}$$

whenever $0 < s < \text{dist}(y, \mathbb{R}^n \setminus Q)$. If $y \in 2^jQ \setminus 2^{j-1}Q$ for some $j \in \{1, \dots, J\}$, then the boundary Hölder continuity estimate in (5.7) combined with Lemma 5.16 and the Harnack inequality in (2.18) show that

$$\omega^{X_Q}(Q(y, s)) \lesssim \left(\frac{\ell(Q)}{2^{j-2}\ell(Q)} \right)^\alpha \omega^{X_{2^{j-2}Q}}(Q(y, s)) \approx \left(\frac{\ell(Q)}{|y - x_Q|} \right)^\alpha \frac{\omega^{X_{Q_0}}(Q(y, s))}{\omega^{X_{Q_0}}(2^jQ)}$$

whenever $0 < s < \text{dist}(y, \mathbb{R}^n \setminus (2^jQ \setminus 2^{j-2}Q))$, where $\alpha > 0$ from (2.17) and the implicit constants depend only on n, λ, Λ and $[\mu]_{A_2}$. The result follows by using these two estimates to bound the limit as s approaches zero in (5.17). \square

5D. The A_∞ -estimate for degenerate elliptic measure. We now combine the properties of degenerate elliptic measure with good ϵ_0 -coverings for sets, as introduced in [Kenig et al. 2000] and defined below (see also [Kenig et al. 2016]), to construct bounded solutions that satisfy the truncated square function estimate in Lemma 5.24. This result, combined with the Carleson measure estimate from Theorem 1.3, allows us to prove the A_∞ -estimate for the degenerate elliptic measure in Theorem 5.30. This avoids the need to apply the method of ϵ -approximability, as was done in [Hofmann et al. 2015a], and so simplifies the proof in the uniformly elliptic case.

Let $\mathbb{D}(\mathbb{R}^n)$ denote the standard collection $\{2^k(j + [0, 1]^n) : k \in \mathbb{Z}, j \in \mathbb{Z}^n\}$ of all *closed* dyadic cubes S in \mathbb{R}^n . For each $S \in \mathbb{D}(\mathbb{R}^n)$ and $\eta = 2^{-K}$, where $K \in \mathbb{N}$, define $\mathbb{D}(S) := \{S' \in \mathbb{D}(\mathbb{R}^n) : S' \subseteq S\}$ and

$$\mathbb{D}^\eta(S) := \{S' \in \mathbb{D}(S) : \ell(S') = 2^{-K}\ell(S)\}, \quad (5.19)$$

so $\mathbb{D}^\eta(S)$ is precisely the set of all *dyadic descendants* of S at scale $2^{-K}\ell(S)$.

Definition 5.20. Suppose that Q_0 is a cube in \mathbb{R}^n . If $\epsilon_0 > 0$, $k \in \mathbb{N}$, $Q \subseteq Q_0$ is a cube and $E \subseteq Q$, then a *good ϵ_0 -cover of E of length k in Q* is a collection $\{O_l\}_{l=1}^k$ of nested open sets that satisfy $E \subseteq O_k \subseteq O_{k-1} \subseteq \dots \subseteq O_1 \subseteq Q$ and each of which has a decomposition $O_l = \bigcup_{i=1}^\infty S_i^l$ given by a collection $\{S_i^l\}_{i \in \mathbb{N}} \subseteq \mathbb{D}(\mathbb{R}^n)$ of dyadic cubes with pairwise disjoint interiors such that

$$\omega^{X_{2Q_0}}(O_l \cap S_i^{l-1}) \leq \epsilon_0 \omega^{X_{2Q_0}}(S_i^{l-1}) \quad \text{for all } i \in \mathbb{N}, \text{ for all } l \in \{2, \dots, k\}. \quad (5.21)$$

Let us record a few important consequences of this definition that will be needed. It is proved on page 243 in [Kenig et al. 2000] that for each $i \in \mathbb{N}$ and $l \in \{2, \dots, k\}$, there exists a unique $j \in \mathbb{N}$ such that S_i^l is a *proper* subset of S_j^{l-1} ; thus $\ell(S_i^l) \leq \frac{1}{2}\ell(S_j^{l-1})$. Also, for $m \in \{2, \dots, k\}$, iterating (5.21) as in Lemma 2.5 of [Kenig et al. 2000] shows that

$$\omega^{X_{2Q_0}}(O_l \cap S_i^m) \leq \epsilon_0^{l-m} \omega^{X_{2Q_0}}(S_i^m) \quad \text{for all } i \in \mathbb{N}, \text{ for all } l \in \{m, \dots, k\}. \quad (5.22)$$

In the uniformly elliptic case, the following result is Lemma 2.3 from [Kenig et al. 2016]. The proof extends to the degenerate elliptic case, since it only relies on the fact that the degenerate elliptic measure $\omega^{X_2 Q_0}$ is doubling when restricted to the cube Q_0 .

Lemma 5.23. *Suppose that Q_0 is a cube in \mathbb{R}^n . If $\epsilon_0 > 0$, then there exists $\delta_0 > 0$, depending only on ϵ_0 , n , λ , Λ and $[\mu]_{A_2}$, such that the following property holds:*

If $Q \subseteq Q_0$ is a cube and $E \subseteq Q$ such that $\omega^{X_2 Q_0}(E) \leq \delta_0$, then there exists a good ϵ_0 -cover of E of length k in Q for some natural number $k \approx \log(\omega^{X_2 Q_0}(E))/\log \epsilon_0$, where the implicit constants depend only on n , λ , Λ and $[\mu]_{A_2}$.

We can now prove the following lemma by adapting the proof in [Kenig et al. 2016] to the degenerate elliptic case. The original argument has also been somewhat modified.

Lemma 5.24. *Suppose that Q_0 is a cube in \mathbb{R}^n . If $M \geq 1$, then there exists $\delta_M > 0$, depending only on M , n , λ , Λ and $[\mu]_{A_2}$, such that the following property holds:*

If $Q \subseteq Q_0$ is a cube and $E \subseteq Q$ and $\omega^{X_2 Q_0}(E) \leq \delta_M$, then there is a Borel subset \mathcal{B} of \mathbb{R}^n such that the solution $u(X) := \omega^X(\mathcal{B})$ of $\operatorname{div}(A \nabla u) = 0$ in \mathbb{R}_+^{n+1} satisfies

$$M \leq \int_0^{\gamma \ell(Q)} \int_{\Delta(x, \gamma t)} |t \nabla u(y, t)|^2 \frac{d\mu(y)}{\mu(\Delta(x, t))} \frac{dt}{t} \quad \text{for all } x \in E,$$

where $\gamma > 0$ is a constant that depends only on n , λ , Λ and $[\mu]_{A_2}$.

Proof. We introduce three constants $\epsilon_0, \delta, \eta \in (0, 1)$ that will be chosen with $\delta \leq \delta_0$, where δ_0 is determined by ϵ_0 as in Lemma 5.23, and $\eta = 2^{-K}$ for some $K \in \mathbb{N}$. Therefore, if $E \subseteq Q \subseteq Q_0$ and $\omega^{X_2 Q_0}(E) \leq \delta$, then there exists a good ϵ_0 -cover of E of length k in Q such that $k \approx \log(\omega^{X_2 Q_0}(E))/\log \epsilon_0$. This cover is denoted by $\{O_l\}_{l=1}^k$ with $O_l = \bigcup_{i=1}^\infty S_i^l$ as in Definition 5.20, and for each such cube S_i^l , a dyadic descendant \tilde{S}_i^l in $\mathbb{D}^\eta(S_i^l)$ that contains the centre of S_i^l is now fixed and

$$\tilde{O}_l := \bigcup_{i=1}^\infty \tilde{S}_i^l, \quad (5.25)$$

where we note that $\ell(\tilde{S}_i^l) = \eta \ell(S_i^l)$ in accordance with (5.19).

We claim that there exists a Borel subset \mathcal{B} of \mathbb{R}^n such that $\mathbb{1}_{\mathcal{B}} = \sum_{l=2}^k \mathbb{1}_{\tilde{O}_{l-1} \setminus O_l}$. To see this, suppose that $\sum_{l=2}^k \mathbb{1}_{\tilde{O}_{l-1} \setminus O_l}(x) \neq 0$ and let l_0 denote the smallest integer $l \in [2, k]$ such that $\mathbb{1}_{\tilde{O}_{l-1} \setminus O_l}(x) = 1$. It must hold that $x \in \tilde{O}_{l_0-1} \setminus O_{l_0}$, so then $x \notin O_{l_0}$, which implies $x \notin O_l$ and $x \notin \tilde{O}_l$ for all $l \geq l_0$; hence $\mathbb{1}_{\tilde{O}_{l-1} \setminus O_l}(x) = 0$ for all $l > l_0$ and the claim follows.

We now aim to choose $\epsilon_0, \eta \in (0, 1)$ such that $u(X) := \omega^X(\mathcal{B})$ on \mathbb{R}_+^{n+1} satisfies

$$|u(X_{\eta S_i^l}) - u(X_{\eta \hat{S}_i^l})| \gtrsim 1 \quad \text{for all } \hat{S}_i^l \in \mathbb{D}^\eta(S_i^l), \text{ for all } i \in \mathbb{N}, \text{ for all } l \in \{1, \dots, k\}, \quad (5.26)$$

where the implicit constant depends only on the allowed constants n, λ, Λ and $[\mu]_{A_2}$, and if x_i^l and \hat{x}_i^l denote the centres of S_i^l and \hat{S}_i^l , then the relevant corkscrew points are precisely $X_{\eta S_i^l} = (x_i^l, \eta \ell(S_i^l))$ and $X_{\eta \hat{S}_i^l} = (\hat{x}_i^l, \eta^2 \ell(S_i^l))$. To this end, we proceed to obtain estimates for $u(X_{\eta S_i^l})$ and $u(X_{\eta \hat{S}_i^l})$.

To estimate $u(X_{\eta S_i^l})$, write

$$u(X_{\eta S_i^l}) = \int_{\mathbb{R}^n \setminus S_i^l} \mathbb{1}_B d\omega^{X_{\eta S_i^l}} + \int_{S_i^l} \mathbb{1}_B d\omega^{X_{\eta S_i^l}} =: I + II.$$

The boundary Hölder continuity in (5.7) shows that $I \leq \omega^{X_{\eta S_i^l}}(\mathbb{R}^n \setminus S_i^l) \leq C_0 \eta^\alpha$, where $C_0, \alpha > 0$ depend only on the allowed constants. To estimate II , write

$$\begin{aligned} II &= \sum_{j=2}^l \int_{S_i^l} \mathbb{1}_{\tilde{O}_{j-1} \setminus O_j} d\omega^{X_{\eta S_i^l}} + \sum_{j=l+2}^k \int_{S_i^l} \mathbb{1}_{\tilde{O}_{j-1} \setminus O_j} d\omega^{X_{\eta S_i^l}} + \int_{S_i^l} \mathbb{1}_{\tilde{O}_l \setminus O_{l+1}} d\omega^{X_{\eta S_i^l}} \\ &=: II_1 + II_2 + II_3. \end{aligned}$$

First, observe that $II_1 = 0$, since if $m \in \{2, \dots, l\}$, then $S_i^l \subseteq O_l \subseteq O_j$ and so $(\tilde{O}_{j-1} \setminus O_j) \cap S_i^l = \emptyset$. To estimate II_2 , the kernel function representation in (5.17) and estimates in Proposition 5.18, the local doubling property of the degenerate elliptic measure in (5.12) and property (5.22) of the good ϵ_0 -covering, show that

$$\begin{aligned} II_2 &= \sum_{j=l+2}^k \int_{(\tilde{O}_{j-1} \setminus O_j) \cap S_i^l} K(X_{2Q_0}, X_{\eta S_i^l}, y) d\omega^{X_{2Q_0}}(y) \\ &\leq \frac{C_\eta}{\omega^{X_{2Q_0}}(S_i^l)} \sum_{j=l+2}^k \omega^{X_{2Q_0}}((\tilde{O}_{j-1} \setminus O_j) \cap S_i^l) \\ &\leq \frac{C_\eta}{\omega^{X_{2Q_0}}(S_i^l)} \sum_{j=l+2}^k \omega^{X_{2Q_0}}(O_{j-1} \cap S_i^l) \\ &\leq \frac{C_\eta}{\omega^{X_{2Q_0}}(S_i^l)} \sum_{j=l+2}^k \epsilon_0^{j-1-l} \omega^{X_{2Q_0}}(S_i^l) \leq \frac{C_\eta \epsilon_0}{1 - \epsilon_0}, \end{aligned}$$

where the constant $C_\eta > 0$ depends only on η and the allowed constants.

To estimate II_3 , observe that $S_i^l \cap \tilde{O}_l = \tilde{S}_i^l$ by the definition of \tilde{O}_l in (5.25); hence

$$II_3 = \int_{\tilde{S}_i^l} d\omega^{X_{\eta S_i^l}} - \int_{\tilde{S}_i^l \cap O_{l+1}} d\omega^{X_{\eta S_i^l}} =: II_3' - II_3''.$$

The term II_3'' is estimated in the same way as II_2 above to show that

$$II_3'' \leq \frac{C_\eta}{\omega^{X_{2Q_0}}(S_i^l)} \omega^{X_{2Q_0}}(O_{l+1} \cap \tilde{S}_i^l) \leq \frac{C_\eta}{\omega^{X_{2Q_0}}(S_i^l)} \omega^{X_{2Q_0}}(O_{l+1} \cap S_i^l) \leq C_\eta \epsilon_0.$$

We estimate II_3' from above and below. First, note that $X_{\eta S_i^l} = (x_i^l, \eta \ell(S_i^l))$, $x_i^l \in \tilde{S}_i^l$ and $\ell(\tilde{S}_i^l) = \eta \ell(S_i^l)$, so $\omega^{X_{\eta S_i^l}}(\tilde{S}_i^l) \approx \omega^{X_{\tilde{S}_i^l}}(\tilde{S}_i^l)$ by the Harnack inequality in (2.18), whilst $\omega^{X_{\tilde{S}_i^l}}(\tilde{S}_i^l) \gtrsim 1$ by Lemma 5.14. Thus, there exists $c_0 \in (0, 1)$ depending only on the allowed constants such that $II_3' = \omega^{X_{\eta S_i^l}}(\tilde{S}_i^l) \geq c_0$. Next, choose a different dyadic descendant $\underline{S}_i^l \neq \tilde{S}_i^l$ in $\mathbb{D}^\eta(S_i^l)$ that contains the centre of S_i^l . The

preceding argument shows that $\omega^{X_{\eta S_i^l}}(\mathcal{S}_i^l) \geq c_0$, whilst $\omega^{X_{\eta S_i^l}}(\mathcal{S}_i^l \cap \tilde{\mathcal{S}}_i^l) \leq \omega^{X_{\eta S_i^l}}(\partial \tilde{\mathcal{S}}_i^l) = 0$; hence

$$c_0 \leq II_3' = \omega^{X_{\eta S_i^l}}(\tilde{\mathcal{S}}_i^l) = 1 - \omega^{X_{\eta S_i^l}}(\mathbb{R}^n \setminus \tilde{\mathcal{S}}_i^l) \leq 1 - \omega^{X_{\eta S_i^l}}(\mathcal{S}_i^l) \leq 1 - c_0.$$

The above estimates together show that if $\epsilon_0 \in (0, \frac{1}{2})$, then

$$c_0 \leq u(X_{\eta S_i^l}) \leq C_0 \eta^\alpha + 3C_\eta \epsilon_0 + 1 - c_0. \quad (5.27)$$

To estimate $u(X_{\eta \hat{S}_i^l})$, write

$$u(X_{\eta \hat{S}_i^l}) = \int_{\mathbb{R}^n \setminus \hat{S}_i^l} \mathbb{1}_B d\omega^{X_{\eta \hat{S}_i^l}} + \int_{\hat{S}_i^l} \mathbb{1}_B d\omega^{X_{\eta \hat{S}_i^l}} =: \hat{I} + \hat{\Pi},$$

as well as

$$\begin{aligned} \hat{\Pi} &= \sum_{j=2}^l \int_{\hat{S}_i^l} \mathbb{1}_{\tilde{O}_{j-1} \setminus O_j} d\omega^{X_{\eta \hat{S}_i^l}} + \sum_{j=l+2}^k \int_{\hat{S}_i^l} \mathbb{1}_{\tilde{O}_{j-1} \setminus O_j} d\omega^{X_{\eta \hat{S}_i^l}} + \int_{\hat{S}_i^l} \mathbb{1}_{\tilde{O}_l \setminus O_{l+1}} d\omega^{X_{\eta \hat{S}_i^l}} \\ &=: \hat{\Pi}_1 + \hat{\Pi}_2 + \hat{\Pi}_3. \end{aligned}$$

The arguments used to estimate I , II_1 and II_2 show that $\hat{I} \leq \omega^{X_{\eta \hat{S}_i^l}}(\mathbb{R}^n \setminus \hat{S}_i^l) \leq C_0 \eta^\alpha$, $\hat{\Pi}_1 = 0$ and $\hat{\Pi}_2 \leq C_\eta \epsilon_0 / (1 - \epsilon_0)$. To estimate $\hat{\Pi}_3$, observe that

$$\hat{S}_i^l \cap (\tilde{O}_l \setminus O_{l+1}) = (\hat{S}_i^l \cap \tilde{S}_i^l) \setminus O_{l+1},$$

where either $\omega^{X_{\eta \hat{S}_i^l}}(\hat{S}_i^l \cap \tilde{S}_i^l) = 0$ and $\hat{\Pi}_3 = 0$, or $\hat{S}_i^l = \tilde{S}_i^l$ and

$$\hat{\Pi}_3 = \int_{\hat{S}_i^l} d\omega^{X_{\eta \hat{S}_i^l}} - \int_{\hat{S}_i^l \cap O_{l+1}} d\omega^{X_{\eta \hat{S}_i^l}} =: \hat{\Pi}_3' - \hat{\Pi}_3''.$$

The boundary Hölder continuity estimate in (5.7) shows that

$$\hat{\Pi}_3' = \omega^{X_{\eta \hat{S}_i^l}}(\hat{S}_i^l) = 1 - \omega^{X_{\eta \hat{S}_i^l}}(\mathbb{R}^n \setminus \hat{S}_i^l) \geq 1 - C_0 \eta^\alpha,$$

whilst repeating the arguments used to estimate II_3'' shows that

$$\hat{\Pi}_3'' \leq \frac{C_\eta}{\omega^{X_{2Q_0}}(\hat{S}_i^l)} \omega^{X_{2Q_0}}(O_{l+1} \cap \hat{S}_i^l) \leq \frac{C_\eta}{\omega^{X_{2Q_0}}(S_i^l)} \omega^{X_{2Q_0}}(O_{l+1} \cap S_i^l) \leq C_\eta \epsilon_0.$$

These estimates together show that if $\epsilon_0 \in (0, \frac{1}{2})$, then either

$$0 \leq u(X_{\eta \hat{S}_i^l}) \leq C_0 \eta^\alpha + 3C_\eta \epsilon_0 \quad \text{or} \quad u(X_{\eta \hat{S}_i^l}) \geq 1 - (C_0 \eta^\alpha + C_\eta \epsilon_0). \quad (5.28)$$

The estimates (5.27) and (5.28) together imply

$$|u(X_{\eta S_i^l}) - u(X_{\eta \hat{S}_i^l})| \geq c_0 - 2C_0 \eta^\alpha - 4C_\eta \epsilon_0.$$

We thus obtain (5.26) by first choosing $\eta \in (0, 1)$ so that $2C_0 \eta^\alpha \leq \frac{1}{4}c_0$ and then choosing $\epsilon_0 \in (0, \frac{1}{2})$ (depending on η) so that $4C_\eta \epsilon_0 \leq \frac{1}{4}c_0$. These choices of η and ϵ_0 , which depend only on the allowed constants, are now fixed.

To complete the proof, suppose that $M \geq 1$ and $x \in E$, and recall that $\delta \in (0, \delta_0)$ remains to be chosen, where δ_0 is now fixed by our choice of ϵ_0 as in Lemma 5.23. First, fix a cube S^k in $\{S_i^k\}_{i \in \mathbb{N}}$ such that $x \in S^k$. The remarks after Definition 5.20 then imply that for each $l \in \{1, \dots, k-1\}$, there exists a unique cube S^l in $\{S_i^l\}_{i \in \mathbb{N}}$ such that $x \in S^l$ and $S^{l+1} \subset S^l$; thus $\ell(S^{l+1}) \leq \frac{1}{2}\ell(S^l)$. Next, for each $l \in \{1, \dots, k\}$, fix a dyadic descendant \hat{S}^l in $\mathbb{D}^\eta(S^l)$ such that $x \in \hat{S}^l$.

Observe that, for some $\tau \in (0, 1)$ sufficiently close to 1 and depending only on η , the corkscrew points $X_{\eta S^l}$ and $X_{\eta \hat{S}^l}$ both belong to the dilate τQ_η^l of the cube

$$Q_\eta^l := \{(y, t) \in \mathbb{R}_+^{n+1} : |y - x|_\infty < (\tfrac{1}{2} + \tfrac{1}{4}\eta^2)\ell(S^l), \tfrac{1}{2}\eta^2\ell(S^l) < t < (1 + \eta^2)\ell(S^l)\},$$

with $\ell(Q_\eta^l) = (1 + \frac{1}{2}\eta^2)\ell(S^l)$. Therefore, if $c^l := f_{Q_\eta^l} u$, then the Moser-type estimate in (2.16), the Poincaré inequality in (2.5) and the doubling property of μ show that

$$\begin{aligned} |u(X_{\eta S^l}) - u(X_{\eta \hat{S}^l})|^2 &\lesssim |u(X_{\eta S^l}) - c^l|^2 + |u(X_{\eta \hat{S}^l}) - c^l|^2 \lesssim \|u - c^l\|_{L^\infty(\tau Q_\eta^l)}^2 \\ &\lesssim_\eta \int_{Q_\eta^l} |u - c^l|^2 d\mu \lesssim \ell(Q_\eta^l)^2 \int_{Q_\eta^l} |\nabla u|^2 d\mu \\ &\lesssim \frac{\ell(S^l)}{\mu(\Delta(x, (1 + \frac{1}{2}\eta^2)\ell(S^l)))} \int_{Q_\eta^l} |\nabla u|^2 d\mu \\ &\lesssim \iint_{Q_\eta^l} |t \nabla u(y, t)|^2 \frac{d\mu(y)}{\mu(\Delta(x, t))} \frac{dt}{t}. \end{aligned} \quad (5.29)$$

Iterating the bound $\ell(S^{l+1}) \leq \frac{1}{2}\ell(S^l)$ shows that $\ell(S^{l'}) \leq 2^{l-l'}\ell(S^l)$ when $l' \geq l$. This implies that the collection $\{Q_\eta^1, \dots, Q_\eta^k\}$ has the bounded intersection property whereby, for each $l \in \{1, \dots, k\}$, there are at most $3 + 2\log_2(1/\eta^2 + 1)$ such cubes $Q_\eta^{l'}$ satisfying $Q_\eta^{l'} \cap Q_\eta^l \neq \emptyset$. This allows us to sum estimate (5.29) over $l \in \{1, \dots, k\}$ and then apply (5.26) to obtain

$$k \lesssim_\eta \iint_{\bigcup_{l=1}^k Q_\eta^l} |t \nabla u(y, t)|^2 \frac{d\mu(y)}{\mu(\Delta(x, t))} \frac{dt}{t} \lesssim \int_0^{\gamma \ell(Q)} \int_{\Delta(x, \gamma t)} |t \nabla u(y, t)|^2 \frac{d\mu(y)}{\mu(\Delta(x, t))} \frac{dt}{t}$$

for some $\gamma > 0$ that depends only on $\eta > 0$ and thus only on the allowed constants.

To conclude, recall that $k \approx \log(\omega^{X_2 Q_0}(E)^{-1})/\log(1/\epsilon_0) \geq \log(1/\delta)/\log(1/\epsilon_0)$, since $\omega^{X_2 Q_0}(E) \leq \delta < 1$. Therefore, the result follows by choosing $\delta \in (0, \delta_0]$ such that $M \leq \log(1/\delta)$, since $\delta_M := \delta$ depends only on M and the allowed constants. \square

We now combine the above technical lemma with the Carleson measure estimate from Theorem 1.3 to prove the main A_∞ -estimate for degenerate elliptic measure.

Theorem 5.30. *Suppose that Q_0 is a cube in \mathbb{R}^n . If $X \in \mathbb{R}_+^{n+1} \setminus T_{Q_0}$ and $\omega := \omega^X \lfloor Q_0$ denotes the degenerate elliptic measure restricted to Q_0 , then $\omega \in A_\infty(\mu)$ and the following equivalent properties hold:*

- (1) *For each $\epsilon \in (0, 1)$, there exists $\delta \in (0, 1)$, depending only on $\epsilon, n, \lambda, \Lambda$ and $[\mu]_{A_2}$, such that the following property holds: if $Q \subseteq Q_0$ is a cube and $E \subseteq Q$ such that $\omega(E) \leq \delta \omega(Q)$, then $\mu(E) \leq \epsilon \mu(Q)$.*

- (2) *The measure ω is absolutely continuous with respect to μ and there exists $q \in (1, \infty)$ such that the Radon–Nikodym derivative $k := d\omega/d\mu$ satisfies, on all surface balls $\Delta \subseteq Q_0$, the reverse Hölder estimate*

$$\left(\int_{\Delta} k^q d\mu \right)^{1/q} \lesssim \int_{\Delta} k d\mu,$$

where q and the implicit constant depend only on n, λ, Λ and $[\mu]_{A_2}$.

- (3) *There exist $C, \theta > 0$, depending only on n, λ, Λ and $[\mu]_{A_2}$, such that*

$$\omega(E) \leq C \left(\frac{\mu(E)}{\mu(Q)} \right)^{\theta} \omega(Q)$$

for all cubes $Q \subseteq Q_0$ and all Borel sets $E \subseteq Q$.

Proof. It is well known that (1)–(3) are equivalent; see Theorem 1.4.13 in [Kenig 1994]. Moreover, by Lemma 5.16, it suffices to prove (1) when $X = X_{2Q_0}$. In that case, by Lemma 5.24, the Carleson measure estimate in Theorem 1.3, Fubini’s theorem and the doubling property of μ , it follows that for each $M \geq 1$, there exists $\delta_M > 0$, depending only on M and the allowed constants, such that the following property holds: if $Q \subseteq Q_0$ is a cube and $E \subseteq Q$ such that $\omega(E) \leq \delta_M \omega(Q)$, then there exists a solution u of the equation $\operatorname{div}(A\nabla u) = 0$ in \mathbb{R}_+^{n+1} with $\|u\|_{\infty} \leq 1$ such that

$$\begin{aligned} M\mu(E) &\leq \int_E \int_0^{\gamma\ell(Q)} \int_{\Delta(x, \gamma t)} |t\nabla u(y, t)|^2 \frac{d\mu(y)}{\mu(\Delta(x, t))} \frac{dt}{t} d\mu(x) \\ &\lesssim \int_0^{\tilde{\gamma}\ell(Q)} \int_{\tilde{\gamma}Q} |t\nabla u(y, t)|^2 d\mu(y) \frac{dt}{t} \lesssim \mu(Q), \end{aligned}$$

where the implicit constants and $\tilde{\gamma} > \gamma > 0$ depend only on the allowed constants. Therefore, if $\epsilon \in (0, 1)$, we choose $M(\epsilon) \geq 1$ and thus $\delta_{M(\epsilon)} \in (0, 1)$, depending only on ϵ and the allowed constants, such that $\mu(E) \leq \epsilon\mu(Q)$, as required. \square

5E. The square function and nontangential maximal function estimates. The $L_{\mu}^p(\mathbb{R}^n)$ -norm equivalence between the square function Su and the nontangential maximal function $N_{*}u$ of solutions u in Theorem 1.5 is now a corollary of the main A_{∞} -estimate for the degenerate elliptic measure in Theorem 5.30. This was proved by Dahlberg, Jerison and Kenig in Theorem 1 of [Dahlberg et al. 1984], which actually provides the more general result in Theorem 5.31 below. In particular, the degenerate elliptic case is treated on page 106 of the same paper, noting that the normalisation $u(X_0) = 0$ assumed therein is actually only required for the so-called $N \lesssim S$ -estimate.

Theorem 5.31. *Suppose that $\Phi : [0, \infty) \rightarrow [0, \infty)$ is an unbounded, nondecreasing, continuous function with $\Phi(0) = 0$ and $\Phi(2t) \leq C\Phi(t)$ for all $t > 0$ and some $C > 0$. If $\operatorname{div}(A\nabla u) = 0$ in \mathbb{R}_+^{n+1} , then*

$$\int_{\mathbb{R}^n} \Phi(Su) d\mu \lesssim \int_{\mathbb{R}^n} \Phi(N_{*}u) d\mu,$$

and if, in addition, $u(X_0) = 0$ for some $X_0 \in \mathbb{R}_+^{n+1}$, then

$$\int_{\mathbb{R}^n} \Phi(N_* u) d\mu \lesssim \int_{\mathbb{R}^n} \Phi(Su) d\mu,$$

where the implicit constants depend only on X_0 , Φ , n , λ , Λ and $[\mu]_{A_2}$.

The next result is also a consequence of the main A_∞ -estimate in Theorem 5.30. It will allow us to construct solutions to the Dirichlet problem $(D)_{p,\mu}$ as integrals of $L_\mu^p(\mathbb{R}^n)$ -boundary data with respect to degenerate elliptic measure.

Lemma 5.32. *Suppose that $1/p + 1/q = 1$, where $q \in (1, \infty)$ is the reverse Hölder exponent from Theorem 5.30. If $X = (x, t) \in \mathbb{R}_+^{n+1}$, then the Radon–Nikodym derivative $k(X, \cdot) := d\omega^X/d\mu$ is in $L_\mu^q(\mathbb{R}^n)$ and*

$$\int_{\mathbb{R}^n} k((x, t), y)^q d\mu(y) \lesssim \mu(\Delta(x, t))^{1-q}.$$

Moreover, if $f \in L_\mu^p(\mathbb{R}^n)$ and $u(X) := \int_{\mathbb{R}^n} f(y) d\omega^X$, then $\|N_* u\|_{L_\mu^p(\mathbb{R}^n)} \lesssim \|f\|_{L_\mu^p(\mathbb{R}^n)}$. The implicit constant in each estimate depends only on n , λ , Λ and $[\mu]_{A_2}$.

Proof. Suppose that $X = (x, t) \in \mathbb{R}_+^{n+1}$. The proof of Proposition 5.18 shows that

$$k((x, t), y) \lesssim 2^{-j\alpha} \frac{k((x, 2^j t), y)}{\omega(x, 2^j t)(\Delta(x, 2^j t))} \quad \text{for all } y \in \Delta(x, 2^j t) \setminus \Delta(x, 2^{j-1} t), \text{ for all } j \in \mathbb{N}.$$

Applying the reverse Hölder estimate from Theorem 5.30 then shows that

$$\begin{aligned} \int_{\mathbb{R}^n} k((x, t), y)^q d\mu(y) &= \int_{\Delta(x, t)} k((x, t), y)^q d\mu(y) + \sum_{j=1}^{\infty} \int_{\Delta(x, 2^j t) \setminus \Delta(x, 2^{j-1} t)} k((x, t), y)^q d\mu(y) \\ &\lesssim \mu(\Delta(x, t))^{1-q} + \sum_{j=1}^{\infty} 2^{-j\alpha q} \mu(\Delta(x, 2^j t))^{1-q} \lesssim \mu(\Delta(x, t))^{1-q}. \end{aligned}$$

To obtain the nontangential maximal function estimate, it suffices to consider the case when $f \geq 0$, since in general we may then decompose $f = f^+ - f^-$ into its positive and negative parts f^+ , $f^- \geq 0$. To this end, suppose that $x_0 \in \mathbb{R}^n$ and that $X = (x, t) \in \mathbb{R}_+^{n+1}$ in order to write

$$f = f \mathbb{1}_{\Delta(x_0, 2t)} + \sum_{j=1}^{\infty} f \mathbb{1}_{\Delta(x_0, 2^{j+1}t) \setminus \Delta(x_0, 2^j t)} =: \sum_{j=0}^{\infty} f_j$$

and define

$$u_j(X) := \int_{\mathbb{R}^n} f_j(y) d\omega^X(y) = \int_{\mathbb{R}^n} f_j(y) k(X, y) d\mu(y).$$

The self-improvement property of the reverse Hölder estimate from Theorem 5.30 (see Theorem 1.4.13 in [Kenig 1994]) implies that there exists an exponent $r > q$ such that

$$\left(\int_{\Delta} k((x, t), y)^r d\mu(y) \right)^{1/r} \lesssim \int_{\Delta} k((x, t), y) d\mu(y) \leq \frac{1}{\mu(\Delta)} \quad (5.33)$$

for all surface balls $\Delta \subseteq \Delta(x, \frac{1}{2}t)$.

Now suppose that $X = (x, t) \in \Gamma(x_0)$. To estimate u_0 , we apply the interior Harnack inequality in (2.18) followed by Hölder's inequality and (5.33) to obtain

$$\begin{aligned} u_0(x, t) &\approx u_0(x, 6t) \leq \int_{\Delta(x_0, 2t)} f(y) k((x, 6t), y) d\mu(y) \\ &\leq \left(\int_{\Delta(x_0, 2t)} |k((x, 6t), y)|^r d\mu(y) \right)^{1/r} \left(\int_{\Delta(x_0, 2t)} f(y)^{r'} d\mu(y) \right)^{1/r'} \\ &\lesssim \mu(\Delta(x_0, 2t))^{-1/r'} \left(\int_{\Delta(x_0, 2t)} f(y)^{r'} d\mu(y) \right)^{1/r'} \\ &\leq [M_\mu(f^{r'})(x_0)]^{1/r'}. \end{aligned}$$

To estimate u_j when $j \in \mathbb{N}$, we apply the boundary Hölder continuity estimate from (5.7) and then proceed as in the estimate above to obtain

$$\begin{aligned} u_j(x, t) &\lesssim \left(\frac{t}{2^j t} \right)^\alpha u_j(x_0, 2^j t) \approx 2^{-j\alpha} u_j(x_0, 2^{j+2}t) \\ &\leq 2^{-j\alpha} \int_{\Delta(x_0, 2^{j+1}t)} f(y) k((x_0, 2^{j+2}t), y) d\mu(y) \\ &\leq 2^{-j\alpha} \left(\int_{\Delta(x_0, 2^{j+1}t)} k((x_0, 2^{j+2}t), y)^r d\mu(y) \right)^{1/r} \left(\int_{\Delta(x_0, 2^{j+1}t)} f(y)^{r'} d\mu(y) \right)^{1/r'} \\ &\lesssim 2^{-j\alpha} \left(\int_{\Delta(x_0, 2^{j+1}t)} f(y)^{r'} d\mu(y) \right)^{1/r'} \\ &\leq 2^{-j\alpha} [M_\mu(f^{r'})(x_0)]^{1/r'}. \end{aligned}$$

The above estimates together show that

$$N_* u(x_0) \lesssim [M_\mu(f^{r'})(x_0)]^{1/r'}$$

for all $x_0 \in \mathbb{R}^n$, and since $r' < q' = p$, it follows that $\|N_* u\|_{L_\mu^p} \lesssim \|f\|_{L_\mu^p}$, as required. \square

We conclude the paper by using the preceding lemma to obtain solvability of the Dirichlet problem $(D)_{p,\mu}$. A uniqueness result is also obtained but only for solutions that converge uniformly to 0 at infinity. This restriction does not appear in the uniformly elliptic case; see Theorem 1.7.7 in [Kenig 1994]. It arises here because of the absence of a Green's function for degenerate elliptic equations on unbounded domains (see Section 5C) and it is not clear to us whether this can be improved.

Theorem 5.34. *Suppose that $1/p + 1/q = 1$, where $q \in (1, \infty)$ is the reverse Hölder exponent from Theorem 5.30. The Dirichlet problem for $L_\mu^p(\mathbb{R}^n)$ -boundary data is solvable in the sense that for each $f \in L_\mu^p(\mathbb{R}^n)$, there exists a solution u such that*

$$\begin{cases} \operatorname{div}(A\nabla u) = 0 & \text{in } \mathbb{R}_+^{n+1}, \\ N_* u \in L_\mu^p(\mathbb{R}^n), \\ \lim_{t \rightarrow 0} u(\cdot, t) = f, \end{cases} \quad (D)_{p,\mu}$$

where the limit converges in $L^p_\mu(\mathbb{R}^n)$ -norm and in the nontangential sense whereby

$$\lim_{\Gamma(x) \ni (y,t) \rightarrow (x,0)} u(y,t) = f(x)$$

for almost every $x \in \mathbb{R}^n$. Moreover, if f has compact support, then there is a unique solution u of $(D)_{p,\mu}$ that converges uniformly to 0 at infinity in the sense that $\lim_{R \rightarrow \infty} \|u\|_{L^\infty(\mathbb{R}^{n+1}_+ \setminus B(0,R))} = 0$.

Proof. Suppose that $f \in L^p_\mu(\mathbb{R}^n)$ and define $u(X) := \int_{\mathbb{R}^n} f d\omega^X$ for all $X \in \mathbb{R}^{n+1}_+$. We first prove that $\operatorname{div}(A\nabla u) = 0$ in \mathbb{R}^{n+1}_+ . Let $(f_j)_j$ denote a sequence in $C_c(\mathbb{R}^n)$ that converges to f in $L^p_\mu(\mathbb{R}^n)$ and consider the solutions $u_j(X) := \int_{\mathbb{R}^n} f_j d\omega^X$. The $L^q(\mathbb{R}^n)$ -estimate for the Radon–Nikodym derivative $d\omega^X/d\mu$ from Lemma 5.32 and the doubling property of μ show that

$$\|u_j - u\|_{L^\infty(K)} \lesssim_{\mu,K} \|f_j - f\|_{L^p_\mu(\mathbb{R}^n)}$$

for all $j \in \mathbb{N}$ and any compact set $K \subset \mathbb{R}^{n+1}_+$, so u_j converges to u in $L^2_{\mu,\text{loc}}(\mathbb{R}^n)$. Moreover, Cacioppoli's inequality and the arguments preceding (5.4) show that u_j converges to a solution v in $W^{1,2}_{\mu,\text{loc}}(\mathbb{R}^n)$, so then $u = v$ is a solution in \mathbb{R}^{n+1}_+ as required.

The nontangential maximal function estimate $\|N_* u\|_{L^p_\mu(\mathbb{R}^n)} \lesssim \|f\|_{L^p_\mu(\mathbb{R}^n)}$ is given by Lemma 5.32. To prove the nontangential convergence to the boundary datum, first recall that $u_j \in C(\bar{\mathbb{R}^{n+1}_+})$ with $u_j|_{\mathbb{R}^n} := f_j$, so $\lim_{\Gamma(x) \ni (y,t) \rightarrow (x,0)} u_j(y,t) = f_j(x)$ (see Section 5B). We combine this fact with the bound

$$|u(y,t) - f(x)| \leq |u(y,t) - u_j(y,t)| + |u_j(y,t) - f_j(x)| + |(f_j - f)(x)|$$

to obtain

$$\limsup_{\Gamma(x) \ni (y,t) \rightarrow (x,0)} |u(y,t) - f(x)| \leq |N_*(u - u_j)(x)| + |(f - f_j)(x)|$$

for all $x \in \mathbb{R}^n$. For any $\eta > 0$, we then apply Chebyshev's inequality and the nontangential maximal function estimate from Lemma 5.32 to show that

$$\begin{aligned} \mu(\{x \in \mathbb{R}^n : \limsup_{\Gamma(x) \ni (y,t) \rightarrow (x,0)} |u(y,t) - f(x)| > \eta\}) \\ \leq \mu(\{x \in \mathbb{R}^n : N_*(u - u_j)(x) > \tfrac{1}{2}\eta\}) + \mu(\{x \in \mathbb{R}^n : |(f - f_j)(x)| > \tfrac{1}{2}\eta\}) \\ \lesssim \eta^{-p} (\|N_*(u - u_j)\|_{L^p_\mu(\mathbb{R}^n)}^p + \|f - f_j\|_{L^p_\mu(\mathbb{R}^n)}^p) \\ \lesssim \eta^{-p} \|f - f_j\|_{L^p_\mu(\mathbb{R}^n)}^p. \end{aligned}$$

It follows, since f_j converges to f in $L^p_\mu(\mathbb{R}^n)$, that

$$\lim_{\Gamma(x) \ni (y,t) \rightarrow (x,0)} u(y,t) = f(x)$$

for almost every $x \in \mathbb{R}^n$, as required. The norm convergence $\lim_{t \rightarrow 0} \|u(\cdot, t) - f\|_{L^p_\mu(\mathbb{R}^n)} = 0$ then follows by Lebesgue's dominated convergence theorem.

It remains to prove that u is the unique solution satisfying $\lim_{|X| \rightarrow \infty} \|u(X)\|_\infty = 0$ when f has compact support. In that case, fix $R_0 > 0$ such that f is supported in the surface ball $\Delta(0, R_0)$. If

$X \in \mathbb{R}_+^{n+1}$ and $|X| > 2R_0$, then the reverse Hölder estimate in Theorem 5.30 shows that

$$\begin{aligned} |u(X)| &\leq \int_{\Delta(0, R_0)} |f(y)| k(X, y) d\mu(y) \\ &\leq \|f\|_{L_\mu^p(\mathbb{R}^n)} \left(\int_{\Delta(0, |X|/2)} k(X, y)^q d\mu(y) \right)^{1/q} \\ &\lesssim \|f\|_{L_\mu^p(\mathbb{R}^n)} \mu(\Delta(0, \tfrac{1}{2}|X|))^{1/q} \int_{\Delta(0, |X|/2)} k(X, y) d\mu(y) \\ &\leq \|f\|_{L_\mu^p(\mathbb{R}^n)} \mu(\Delta(0, \tfrac{1}{2}|X|))^{-1/p}, \end{aligned}$$

whilst $\lim_{R \rightarrow \infty} \mu(\Delta(0, R)) = \infty$, since μ is in the A_∞ -class with respect to Lebesgue measure on \mathbb{R}^n ; thus $\lim_{R \rightarrow \infty} \|u\|_{L^\infty(\mathbb{R}_+^{n+1} \setminus B(0, R))} = 0$. The maximum principle allows us to conclude that any solution of $(D)_{p, \mu}$ with this decay must be unique. \square

References

- [Albrecht et al. 1996] D. Albrecht, X. Duong, and A. McIntosh, “Operator theory and harmonic analysis”, pp. 77–136 in *Instructional workshop on analysis and geometry, III* (Canberra, 1995), edited by T. Cranny and J. Hutchinson, Proc. Centre Math. Appl. Austral. Nat. Univ. **34**, Austral. Nat. Univ., Canberra, 1996. MR Zbl
- [Auscher 2007] P. Auscher, *On necessary and sufficient conditions for L^p -estimates of Riesz transforms associated to elliptic operators on \mathbb{R}^n and related estimates*, Mem. Amer. Math. Soc. **871**, Amer. Math. Soc., Providence, RI, 2007. MR Zbl
- [Auscher and Stahlhut 2014] P. Auscher and S. Stahlhut, “A priori estimates for boundary value elliptic problems via first order systems”, preprint, 2014. arXiv
- [Auscher et al. 2001] P. Auscher, S. Hofmann, J. L. Lewis, and P. Tchamitchian, “Extrapolation of Carleson measures and the analyticity of Kato’s square-root operators”, *Acta Math.* **187**:2 (2001), 161–190. MR Zbl
- [Auscher et al. 2002] P. Auscher, S. Hofmann, M. Lacey, A. McIntosh, and P. Tchamitchian, “The solution of the Kato square root problem for second order elliptic operators on \mathbb{R}^n ”, *Ann. of Math. (2)* **156**:2 (2002), 633–654. MR Zbl
- [Auscher et al. 2015] P. Auscher, A. Rosén, and D. Rule, “Boundary value problems for degenerate elliptic equations and systems”, *Ann. Sci. Éc. Norm. Sup. (4)* **48**:4 (2015), 951–1000. MR Zbl
- [Caffarelli et al. 1981] L. Caffarelli, E. Fabes, S. Mortola, and S. Salsa, “Boundary behavior of nonnegative solutions of elliptic operators in divergence form”, *Indiana Univ. Math. J.* **30**:4 (1981), 621–640. MR Zbl
- [Cowling et al. 1996] M. Cowling, I. Doust, A. McIntosh, and A. Yagi, “Banach space operators with a bounded H^∞ functional calculus”, *J. Austral. Math. Soc. Ser. A* **60**:1 (1996), 51–89. MR Zbl
- [Cruz-Uribe and Rios 2008] D. Cruz-Uribe and C. Rios, “Gaussian bounds for degenerate parabolic equations”, *J. Funct. Anal.* **255**:2 (2008), 283–312. MR Zbl
- [Cruz-Uribe and Rios 2014] D. Cruz-Uribe and C. Rios, “Corrigendum to ‘Gaussian bounds for degenerate parabolic equations’”, *J. Funct. Anal.* **267**:9 (2014), 3507–3513. MR Zbl
- [Cruz-Uribe and Rios 2015] D. Cruz-Uribe and C. Rios, “The Kato problem for operators with weighted ellipticity”, *Trans. Amer. Math. Soc.* **367**:7 (2015), 4727–4756. MR Zbl
- [Dahlberg et al. 1984] B. E. J. Dahlberg, D. S. Jerison, and C. E. Kenig, “Area integral estimates for elliptic differential operators with nonsmooth coefficients”, *Ark. Mat.* **22**:1 (1984), 97–108. MR Zbl
- [Evans and Gariepy 1992] L. C. Evans and R. F. Gariepy, *Measure theory and fine properties of functions*, CRC Press, Boca Raton, FL, 1992. MR Zbl
- [Fabes et al. 1982a] E. Fabes, D. Jerison, and C. Kenig, “The Wiener test for degenerate elliptic equations”, *Ann. Inst. Fourier (Grenoble)* **32**:3 (1982), 151–182. MR Zbl

- [Fabes et al. 1982b] E. B. Fabes, C. E. Kenig, and R. P. Serapioni, “The local regularity of solutions of degenerate elliptic equations”, *Comm. Partial Differential Equations* **7**:1 (1982), 77–116. MR Zbl
- [Fabes et al. 1983] E. B. Fabes, C. E. Kenig, and D. Jerison, “Boundary behavior of solutions to degenerate elliptic equations”, pp. 577–589 in *Conference on harmonic analysis in honor of Antoni Zygmund, II* (Chicago, 1981), edited by W. Beckner et al., Wadsworth, Belmont, CA, 1983. MR Zbl
- [Fernandes 1991] J. C. Fernandes, “Mean value and Harnack inequalities for a certain class of degenerate parabolic equations”, *Rev. Mat. Iberoamericana* **7**:3 (1991), 247–286. MR Zbl
- [García-Cuerva and Rubio de Francia 1985] J. García-Cuerva and J. L. Rubio de Francia, *Weighted norm inequalities and related topics*, North-Holland Math. Studies **116**, North-Holland, Amsterdam, 1985. MR Zbl
- [Gilbarg and Trudinger 1977] D. Gilbarg and N. S. Trudinger, *Elliptic partial differential equations of second order*, Grundlehren der Math. Wissenschaften **224**, Springer, 1977. MR Zbl
- [Hofmann and Kim 2007] S. Hofmann and S. Kim, “The Green function estimates for strongly elliptic systems of second order”, *Manuscripta Math.* **124**:2 (2007), 139–172. MR Zbl
- [Hofmann and Martell 2014] S. Hofmann and J. M. Martell, “Uniform rectifiability and harmonic measure, I: Uniform rectifiability implies Poisson kernels in L^p ”, *Ann. Sci. Éc. Norm. Sup. (4)* **47**:3 (2014), 577–654. MR Zbl
- [Hofmann et al. 2015a] S. Hofmann, C. Kenig, S. Mayboroda, and J. Pipher, “Square function/non-tangential maximal function estimates and the Dirichlet problem for non-symmetric elliptic operators”, *J. Amer. Math. Soc.* **28**:2 (2015), 483–529. MR Zbl
- [Hofmann et al. 2015b] S. Hofmann, M. Mitrea, and A. J. Morris, “The method of layer potentials in L^p and endpoint spaces for elliptic operators with L^∞ coefficients”, *Proc. Lond. Math. Soc. (3)* **111**:3 (2015), 681–716. MR Zbl
- [Ishige 1999] K. Ishige, “On the behavior of the solutions of degenerate parabolic equations”, *Nagoya Math. J.* **155** (1999), 1–26. MR Zbl
- [Jerison and Kenig 1981] D. S. Jerison and C. E. Kenig, “The Dirichlet problem in nonsmooth domains”, *Ann. of Math. (2)* **113**:2 (1981), 367–382. MR Zbl
- [Kenig 1994] C. E. Kenig, *Harmonic analysis techniques for second order elliptic boundary value problems*, CBMS Regional Conf. Series in Math. **83**, Amer. Math. Soc., Providence, RI, 1994. MR Zbl
- [Kenig et al. 2000] C. Kenig, H. Koch, J. Pipher, and T. Toro, “A new approach to absolute continuity of elliptic measure, with applications to non-symmetric equations”, *Adv. Math.* **153**:2 (2000), 231–298. MR Zbl
- [Kenig et al. 2016] C. Kenig, B. Kirchheim, J. Pipher, and T. Toro, “Square functions and the A_∞ property of elliptic measures”, *J. Geom. Anal.* **26**:3 (2016), 2383–2410. MR Zbl
- [Kinderlehrer and Stampacchia 1980] D. Kinderlehrer and G. Stampacchia, *An introduction to variational inequalities and their applications*, Pure Appl. Math. **88**, Academic Press, New York, 1980. MR Zbl
- [Le 2015] P. Le, “ L^p bounds of Riesz transform and vertical square functions for degenerate elliptic operators”, preprint, 2015, available at <https://tinyurl.com/pleriesz>.
- [Littman et al. 1963] W. Littman, G. Stampacchia, and H. F. Weinberger, “Regular points for elliptic equations with discontinuous coefficients”, *Ann. Scuola Norm. Sup. Pisa (3)* **17** (1963), 43–77. MR Zbl
- [McIntosh 1986] A. McIntosh, “Operators which have an H_∞ functional calculus”, pp. 210–231 in *Miniconference on operator theory and partial differential equations* (North Ryde, Australia, 1986), edited by B. Jefferies et al., Proc. Centre Math. Anal. Austral. Nat. Univ. **14**, Austral. Nat. Univ., Canberra, 1986, 1986. MR Zbl
- [Morrey 1966] C. B. Morrey, Jr., *Multiple integrals in the calculus of variations*, Grundlehren der Math. Wissenschaften **130**, Springer, 1966. MR Zbl
- [Stein 1970] E. M. Stein, *Singular integrals and differentiability properties of functions*, Princeton Math. Series **30**, Princeton Univ. Press, 1970. MR Zbl
- [Stein 1993] E. M. Stein, *Harmonic analysis: real-variable methods, orthogonality, and oscillatory integrals*, Princeton Math. Series **43**, Princeton Univ. Press, 1993. MR Zbl

STEVE HOFMANN: hofmanns@missouri.edu

Department of Mathematics, University of Missouri, Columbia, MO, United States

PHI LE: ple101@syr.edu

Mathematics Department, Syracuse University, Syracuse, NY, United States

ANDREW J. MORRIS: a.morris.2@bham.ac.uk

School of Mathematics, University of Birmingham, Birmingham, United Kingdom

Analysis & PDE

msp.org/apde

EDITORS

EDITOR-IN-CHIEF

Patrick Gérard
patrick.gerard@math.u-psud.fr
Université Paris Sud XI
Orsay, France

BOARD OF EDITORS

Massimiliano Berti	Scuola Intern. Sup. di Studi Avanzati, Italy berti@sissa.it	Gilles Pisier	Texas A&M University, and Paris 6 pisier@math.tamu.edu
Michael Christ	University of California, Berkeley, USA mchrist@math.berkeley.edu	Tristan Rivière	ETH, Switzerland riviere@math.ethz.ch
Charles Fefferman	Princeton University, USA cf@math.princeton.edu	Igor Rodnianski	Princeton University, USA irod@math.princeton.edu
Ursula Hamenstaedt	Universität Bonn, Germany ursula@math.uni-bonn.de	Yum-Tong Siu	Harvard University, USA siu@math.harvard.edu
Vadim Kaloshin	University of Maryland, USA vadim.kaloshin@gmail.com	Terence Tao	University of California, Los Angeles, USA tao@math.ucla.edu
Herbert Koch	Universität Bonn, Germany koch@math.uni-bonn.de	Michael E. Taylor	Univ. of North Carolina, Chapel Hill, USA met@math.unc.edu
Izabella Laba	University of British Columbia, Canada ilaba@math.ubc.ca	Gunther Uhlmann	University of Washington, USA gunther@math.washington.edu
Richard B. Melrose	Massachusetts Inst. of Tech., USA rbm@math.mit.edu	András Vasy	Stanford University, USA andras@math.stanford.edu
Frank Merle	Université de Cergy-Pontoise, France Frank.Merle@u-cergy.fr	Dan Virgil Voiculescu	University of California, Berkeley, USA dvv@math.berkeley.edu
William Minicozzi II	Johns Hopkins University, USA minicozz@math.jhu.edu	Steven Zelditch	Northwestern University, USA zelditch@math.northwestern.edu
Clément Mouhot	Cambridge University, UK c.mouhot@dpms.cam.ac.uk	Maciej Zworski	University of California, Berkeley, USA zworski@math.berkeley.edu
Werner Müller	Universität Bonn, Germany mueller@math.uni-bonn.de		

PRODUCTION

production@msp.org
Silvio Levy, Scientific Editor

See inside back cover or msp.org/apde for submission instructions.

The subscription price for 2019 is US \$310/year for the electronic version, and \$520/year (+\$60, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues from the last three years and changes of subscriber address should be sent to MSP.

Analysis & PDE (ISSN 1948-206X electronic, 2157-5045 printed) at Mathematical Sciences Publishers, 798 Evans Hall #3840, c/o University of California, Berkeley, CA 94720-3840, is published continuously online. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices.

APDE peer review and production are managed by EditFlow® from MSP.

PUBLISHED BY



mathematical sciences publishers

nonprofit scientific publishing

<http://msp.org/>

© 2019 Mathematical Sciences Publishers

ANALYSIS & PDE

Volume 12 No. 8 2019

Tangent measures of elliptic measure and applications	1891
JONAS AZZAM and MIHALIS MOURGOGLOU	
Discretely self-similar solutions to the Navier–Stokes equations with data in L^2_{loc} satisfying the local energy inequality	1943
ZACHARY BRADSHAW and TAI-PENG TSAI	
Continuity properties for divergence form boundary data homogenization problems	1963
WILLIAM M. FELDMAN and YUMING PAUL ZHANG	
Dynamics of one-fold symmetric patches for the aggregation equation and collapse to singular measure	2003
TAOUFIK HMIDI and DONG LI	
Coupled Kähler–Ricci solitons on toric Fano manifolds	2067
JAKOB HULTGREN	
Carleson measure estimates and the Dirichlet problem for degenerate elliptic equations	2095
STEVE HOFMANN, PHI LE and ANDREW J. MORRIS	



2157-5045(2019)12:8;1-I