

ANALYSIS & PDE

Volume 13

No. 1

2020

MARK ALLEN AND DENNIS KRIVENTSOV

A SPIRAL INTERFACE WITH
POSITIVE ALT-CAFFARELLI-FRIEDMAN LIMIT AT THE ORIGIN

A SPIRAL INTERFACE WITH POSITIVE ALT-CAFFARELLI-FRIEDMAN LIMIT AT THE ORIGIN

MARK ALLEN AND DENNIS KRIVENTSOV

We give an example of a pair of nonnegative subharmonic functions with disjoint support for which the Alt–Caffarelli–Friedman monotonicity formula has strictly positive limit at the origin, and yet the interface between their supports lacks a (unique) tangent there. This clarifies a remark of Caffarelli and Salsa (*A geometric approach to free boundary problems*, 2005) that the positivity of the limit of the ACF formula implies unique tangents; this is true under some additional assumptions, but false in general. In our example, blow-ups converge to the expected piecewise linear two-plane function along subsequences, but the limiting function depends on the subsequence due to the spiraling nature of the interface.

1. Introduction

The Alt–Caffarelli–Friedman monotonicity formula (hereafter denoted ACF formula) has been and continues to be a powerful tool in the study of free boundary problems. It was introduced in [Alt et al. 1984] in order to prove that the solutions to a two-phase Bernoulli free boundary problem are Lipschitz continuous. The formula was then adapted to treat more general two-phase problems, and a discussion of the formula, its proof, and its applications to two-phase free boundary problems may be found in [Caffarelli and Salsa 2005]. The ACF formula has also been effective in studying obstacle-type problems, and applications of the formula for obstacle-type problems are found in [Petrosyan et al. 2012]. Further applications also include the study of segregation problems in [Caffarelli et al. 2009]. While the most typical use of the formula is to prove the optimal regularity of solutions or flatness of the free boundary, it can also be used for other purposes, such as to show the separation of phases in free boundary problems; see [Allen and Petrosyan 2012; Allen et al. 2015; Allen and Shi 2016].

The key property of the ACF formula (1-1) is given in the following proposition:

Proposition 1.1. *Let $u_1, u_2 \geq 0$ be two continuous subharmonic functions in B_R with $u_1 \cdot u_2 = 0$ and $u_1(0) = u_2(0) = 0$. Then*

$$\Phi(r, u_1, u_2) := \frac{1}{r^4} \int_{B_r(0)} \frac{|\nabla u_1|^2}{|x|^{n-2}} \int_{B_r(0)} \frac{|\nabla u_2|^2}{|x|^{n-2}} \quad (1-1)$$

is nondecreasing for $0 < r < R$. Consequently, the limit

$$\Phi(0+, u_1, u_2) := \lim_{r \searrow 0} \Phi(r, u_1, u_2)$$

is well defined.

MSC2010: 35R35, 35J05.

Keywords: ACF monotonicity formula, spiral interface, free boundary, monotonicity formula.

Our paper is motivated by the following claim, which appears as Lemma 12.9 in [Caffarelli and Salsa 2005]:

Claim 1.2. *Let $u \geq 0$ be continuous in B_1 and harmonic in $\{u > 0\}$. Let Ω_1 be a connected component of $\{u > 0\}$ and let $0 \in \partial\Omega_1$. If $u_1 = u|_{\Omega_1}$ and $u_2 = u - u_1$, then if $\Phi(0+, u_1, u_2) > 0$, exactly two connected components Ω_1 and Ω_2 of $\{u > 0\}$ are tangent at 0, and in a suitable system of coordinates,*

$$u(x) = \alpha x_1^+ + \beta x_1^- + o(|x|), \quad (1-2)$$

with $\alpha, \beta > 0$.

As no proof of this Lemma 12.9 is provided in [Caffarelli and Salsa 2005] (it is followed only by some general remarks), it is not entirely clear whether it is meant to be taken at face value. We note, for example, that if u is also assumed to satisfy a two-phase free boundary problem of the type treated in [Caffarelli and Salsa 2005], then the claim is valid, but requires heavy use of the free boundary relation to prove.

Claim 1.2, and in particular the question of whether it is true in the generality stated, drew the authors' interest when the second author was tempted to use it while working on certain eigenvalue optimization problems [Kriventsov and Lin 2019] but was unable to write down a proof. Typically, a monotonicity formula is applied together with other tools making explicit use of the free boundary relation in order to prove regularity of an interface; however, Claim 1.2 would imply that the ACF monotonicity formula, on its own, yields some regularity of the interface. This makes the claim very powerful and useful, especially in problems where the free boundary condition is difficult to exploit, such as the vector-valued free boundary problems arising from spectral optimization [Kriventsov and Lin 2018; 2019].

Unfortunately, it is also not true: the main result of this paper is to provide a counterexample to Claim 1.2.

Theorem 1.3. *For any dimension $n \geq 2$, there exist two continuous subharmonic functions $u, \tilde{u} \geq 0$ with u, \tilde{u} both harmonic in their respective positivity sets and $u \cdot \tilde{u} = 0$. Furthermore, $\Phi(0+, u, \tilde{u}) > 0$. However, $\partial\{u > 0\}$ and $\partial\{\tilde{u} > 0\}$ (which are given by a piecewise smooth, connected hypersurface when restricted to any annulus $B_1 \setminus B_r$) do not admit tangents (or approximate tangents) at the origin, nor do there exist numbers $\alpha, \beta > 0$ and a change of coordinates such that $u + \tilde{u} = \alpha x_1^+ + \beta x_1^- + o(|x|)$.*

In the above, the boundary of a measurable set A is said to *admit a tangent (plane)* at the origin if

$$0 < \liminf_{r \searrow 0} \frac{|B_r \cap A|}{|B_r|} \leq \limsup_{r \searrow 0} \frac{|B_r \cap A|}{|B_r|} < 1 \quad (1-3)$$

and there is a unit vector v such that

$$\lim_{r \searrow 0} \frac{1}{r} \max_{x \in \partial A \cap B_r} |x \cdot v| = 0.$$

The boundary is said to admit an *approximate tangent (plane)* if (1-3) holds and

$$\lim_{r \searrow 0} \frac{1}{r^{n+1}} \int_{B_r \cap \partial A} |x \cdot v|^2 d\mathcal{H}^{n-1} = 0.$$

Here \mathcal{H}^{n-1} denotes $(n-1)$ -dimensional Hausdorff measure. Note that if u, \tilde{u} are as in Claim 1.2 and A is either $\{u > 0\}$ or $\{\tilde{u} > 0\}$, then (1-3) holds; see Corollary 12.4 in [Caffarelli and Salsa 2005].

It seems that the notion of approximate tangent above (or another similar measure-theoretic notion) is the more meaningful one in this context. Indeed, there are simpler constructions which produce functions u, \tilde{u} as in Claim 1.2 for which $\partial\{u > 0\}$ does not admit a tangent at 0 but does admit an approximate tangent.

If one only considers functions u for which $\partial\{u > 0\}$ is, say, given by a 1-Lipschitz graph over some plane π_r on every annulus $B_{2r} \setminus B_r$, these two notions of tangent plane are equivalent. This property holds for the example constructed in the proof of Theorem 1.3.

The functions u, \tilde{u} we construct in proving the theorem have $\partial\{u > 0\}$ a spiral: while $u + \tilde{u}$ looks more and more like $\alpha(x \cdot \nu)_+ + \beta(x \cdot \nu)_-$ on progressively smaller balls B_r , the choice of ν cannot be made uniformly in r , and the optimal ν rotates (slowly) as r decreases. Some free boundary problems are known to exhibit spiraling patterns for the interface; see [Blank 2001; Terracini et al. 2019] for examples, although the spirals produced there have different properties from ours. We also remark that an example of nonunique tangents for an energy minimization problem is given in [White 1992].

Further questions. Before turning to the proof of Theorem 1.3 we would like to offer some discussion of the further questions raised by this theorem and speculate on what “optimal” results, both positive and negative, might look like.

A standard argument with the ACF formula shows that if u, \tilde{u} are as in Claim 1.2, then for every sequence $r_k \rightarrow 0$, there is a subsequence r_{k_j} such that

$$\lim_{j \rightarrow \infty} \frac{1}{r_{k_j}^{n+2}} \int_{B_{r_{k_j}}} |u(x) - \alpha(x \cdot \nu)_+ - \beta(x \cdot \nu)_-|^2 = 0,$$

where α, β, ν depend on the subsequence. Let us refer to any such subsequence r_{k_j} as a *blow-up subsequence*. We are interested in whether or not these parameters may be chosen independent of the blow-up subsequence.

In the example constructed below, the functions u and \tilde{u} are rotations of one another around the origin; in particular, this means that for all of the blow-up subsequences, $\alpha = \beta = c\sqrt{\Phi(0+, u, \tilde{u})}$ are the same, while ν depends on the particular subsequence.

This example gives one way for (1-2) to fail. There could, in principle, be another way: say that $\partial\{u > 0\} = \partial\{\tilde{u} > 0\}$ is given by a C^1 hypersurface (including up to the origin, so that it admits a tangent there), and that u, \tilde{u} are as in Claim 1.2. Can one find a pair u, \tilde{u} like this for which (1-2) fails? This would mean that between the various blow-up subsequences, ν would remain fixed, while α and β would vary. Note that if the hypersurface is more regular near the origin (in particular, if it is a Lyapunov–Dini surface), then this is impossible.

Another set of questions is related to optimality in Theorem 1.3. To clarify the discussion, define, for each r , $\nu(r)$ to be the best approximating normal vector:

$$\int_{B_r \cap \partial\{u > 0\}} |x \cdot \nu(r)|^2 d\mathcal{H}^{n-1} = \min_{\nu \in S^{n-1}} \int_{B_r \cap \partial\{u > 0\}} |x \cdot \nu|^2 d\mathcal{H}^{n-1}.$$

It may be verified that $v(r)$ is uniquely determined from this relation and depends in a Lipschitz manner on r . The property of having an approximate tangent, then, can be reformulated as saying that $v(r)$ has a limit as $r \rightarrow 0$, while Theorem 1.3 gives an example where

$$\int_0^1 \left| \frac{dv(r)}{dr} \right| = \infty. \quad (1-4)$$

What restrictions on the change in $v(r)$, one may ask then, are implied by the conditions in Claim 1.2? We conjecture that under those conditions, one must have

$$\int_0^1 r \left| \frac{dv(r)}{dr} \right|^2 < \infty; \quad (1-5)$$

on the other hand, for any $v_0(r)$ satisfying (1-4) and (1-5), there is a pair of functions u, \tilde{u} as in Claim 1.2 with $v_0(r)$ with

$$\left| \frac{dv(r)}{dr} \right| \geq \left| \frac{dv_0(r)}{dr} \right|.$$

To explain the source of (1-5), let us point out that in Section 2, we construct a pair of functions u, \tilde{u} for which

$$\int_0^\infty \left| \frac{dv(r)}{dr} \right| = \theta \quad \text{and} \quad \frac{\Phi(0+, u, \tilde{u})}{\Phi(\infty, u, \tilde{u})} \geq 1 - \theta^2$$

(and this dependence on θ seems to be sharp up to constants). By gluing truncated and scaled versions of this construction, one might hope to attain functions u, \tilde{u} satisfying the hypotheses of Claim 1.2, and with

$$\int_{2^j}^{2^{j+1}} \left| \frac{dv(r)}{dr} \right| \approx \theta_i$$

for any sequence θ_i for which $\prod_i (1 - \theta_i^2) > 0$. This restriction is equivalent to (1-5) for such a construction. In the actual proof of Theorem 1.3, we are unable to perform the truncation and gluing steps uniformly in θ , and so do not obtain such a quantitative result.

Finally, over the past two decades enormous progress has been made in understanding the relationship between the behavior of positive harmonic functions with zero Dirichlet condition near the boundaries of domains and the geometric measure-theoretic properties of the boundary; we do not attempt to provide a summary here, but refer the reader to the introduction and references in [Azzam et al. 2016]. We suggest that the questions above can be thought of as a continuation, or extension, of this program, with the goal of relating (finer) geometric properties of a boundary to the simultaneous behavior of positive harmonic functions on a domain and its compliment, using the ACF formula as a crucial tool.

Outline of proof. To prove Theorem 1.3 we will construct a subharmonic function $u \geq 0$ in \mathbb{R}^2 such that u is harmonic in its positivity set and $u(0) = 0$. Furthermore, $\partial\{u > 0\}$ will be invariant under a rotation of π . Consequently, if $\tilde{u}(z) := u(-z)$, then the pair u, \tilde{u} will satisfy the assumptions of the ACF formula in Proposition 1.1. Before explaining the construction of u and the outline of the paper, we first give two definitions.

We define the class of functions in \mathbb{R}^2

$$\mathcal{K} := \left\{ u \in C(B_1) : u \geq 0 \text{ in } B_1, \Delta u = 0 \text{ in } \{u > 0\}, u(0) = 0, u(z) \cdot u(-z) = 0, \text{ and } \partial\{u(z) > 0\} = \partial\{u(-z) > 0\} \right\}.$$

By working in the class \mathcal{K} , we may consider using a one-sided rescaled version of the ACF formula. If $u \in \mathcal{K}$, then

$$J(r, u) := \left(\frac{2}{\pi r^2} \int_{B_r} |\nabla u|^2 \right)^{1/2}$$

is monotonically nondecreasing in r since $J(r, u) = \left(\frac{2}{\pi}\right)^2 \sqrt{\Phi(r, u(z), u(-z))}$. Furthermore, if u is C^1 up to $\partial\{u > 0\}$ near the origin, then $J(0+, u) = |\nabla u(0)|$.

In order to prove Theorem 1.3 we first show in Section 2, working on unbounded domains, that it is possible to turn $\partial\{u > 0\}$ so that its asymptotic behavior at infinity differs from its tangent near the origin by an angle of θ while arranging that $J(\infty, u) - J(0+, u) < 1 - \theta^2$ (for small θ). In Section 3 we transfer this result to a bounded domain. In Section 4 we inductively construct a sequence of functions in \mathcal{K} and take a limit to obtain the u in Theorem 1.3. Heuristically, the value of $J(0+, u)$ should be $\prod(1 - \theta_i^2)$, and this is strictly positive if, say, $\theta_i = i^{-1}$. On successively smaller balls, the interface $\{u = 0\}$ will have turned a total amount of $\sum i^{-1} \rightarrow \infty$, which implies that the interface spirals towards the origin and therefore lacks a unique tangent there. We make these heuristic ideas rigorous, and then we show how the pair u, \tilde{u} also provide a counterexample in higher dimensions.

2. Conformal mapping

We utilize the Schwarz–Christoffel formula to obtain a conformal mapping. For a fixed angle $0 < \theta < \frac{\pi}{2}$, we map the upper half-plane to the domain Ω_θ (see Figure 1) by the conformal mapping f_θ with derivative

$$f'_\theta(z) = (z - (-1))^{(\pi+\theta)/\pi-1} (z - 1)^{(\pi-\theta)/\pi-1} = \left(\frac{z+1}{z-1} \right)^{\theta/\pi}. \quad (2-1)$$

We translate f_θ by a constant z_0 , so that the midpoint of the line segment in the image is the origin $0 + 0i$. We define $t_\theta \in (-1, 1) \subset \mathbb{R}$ to be $t_\theta = f_\theta^{-1}(0 + 0i)$. Clearly, $t_\theta \rightarrow 0$ as $\theta \rightarrow 0$. What is of importance is how quickly $t_\theta \rightarrow 0$. In order to determine this decay rate we use the following result:




Figure 1. Conformal map.

Lemma 2.1. *Let $f, g > 0$ be integrable functions on an interval I . If f/g is an increasing function, then for any $x_1 < x_2 < x_3 < x_4$ with each $x_i \in I$, we have*

$$\frac{\int_{x_1}^{x_2} f}{\int_{x_1}^{x_2} g} \leq \frac{\int_{x_3}^{x_4} f}{\int_{x_3}^{x_4} g}.$$

Proof. Since f/g is increasing we have

$$\int_{x_1}^{x_2} f(x) dx \leq \int_{x_1}^{x_2} \frac{f(x_2)}{g(x_2)} g(x) dx.$$

Consequently, we have

$$\frac{\int_{x_1}^{x_2} f(x) dx}{\int_{x_1}^{x_2} g(x) dx} \leq \frac{f(x_2)}{g(x_2)}.$$

By the same argument, we have

$$\frac{f(x_3)}{g(x_3)} \leq \frac{\int_{x_3}^{x_4} f(x) dx}{\int_{x_3}^{x_4} g(x) dx},$$

and so the conclusion follows. \square

We will also need the following:

Lemma 2.2. *Let $f \geq g > 0$ be integrable and continuous on $[0, 1]$ with $f \geq g$ and f/g increasing, and*

$$\int_0^1 f > M \quad \text{and} \quad \int_0^1 g > M.$$

Let x_1, x_2 be the unique values such that

$$M + \int_0^{x_1} g = \int_{x_1}^1 g \quad \text{and} \quad M + \int_0^{x_2} f = \int_{x_2}^1 f. \quad (2-2)$$

Then $x_1 \leq x_2$.

Proof. We have

$$\frac{M + \int_0^{x_1} f}{M + \int_0^{x_1} g} \leq \frac{\int_0^{x_1} f}{\int_0^{x_1} g} \leq \frac{\int_{x_1}^1 f}{\int_{x_1}^1 g},$$

where the second inequality is due to Lemma 2.1. Since x_1 is chosen so that (2-2) holds, we have that the denominator in the inequality above is the same so that

$$M + \int_0^{x_1} f \leq \int_{x_1}^1 f.$$

Then $x_1 \leq x_2$. \square

The two lemmas above allow us to prove:

Lemma 2.3. *Let f_θ be defined as in (2-1) and let $t_\theta = f_\theta^{-1}(0 + 0i)$. Then there exists $\theta_0 > 0$ such that $0 < t_\theta \leq 2\theta/\pi$ as long as $0 < \theta \leq \theta_0$.*

Proof. To determine the midpoint of a line segment it suffices to find the x -value. Consequently, we focus on the real part of the mapping f_θ . If $t \in (-1, 1)$, then

$$f'(t) = \left((-1) \frac{1+t}{1-t} \right)^{\theta/\pi} = \left(\frac{1+t}{1-t} \right)^{\theta/\pi} e^{i\theta}.$$

Thus, t_θ is the unique value in $(-1, 1)$ such that

$$\int_{-1}^{t_\theta} \left(\frac{1+t}{1-t} \right)^{\theta/\pi} dt = \int_{t_\theta}^1 \left(\frac{1+t}{1-t} \right)^{\theta/\pi} dt.$$

We now note that

$$\left(\frac{1+t}{1-t} \right)^{\theta/\pi} \geq \left(\frac{1+t}{2} \right)^{\theta/\pi} \quad \text{if } -1 \leq t \leq 0.$$

Then $t_\theta \leq \xi_\theta$, where ξ_θ is the unique value such that

$$\int_{-1}^0 \left(\frac{1+t}{2} \right)^{\theta/\pi} dt + \int_0^{\xi_\theta} \left(\frac{1+t}{1-t} \right)^{\theta/\pi} dt = \int_{\xi_\theta}^1 \left(\frac{1+t}{1-t} \right)^{\theta/\pi} dt.$$

We also have

$$\left(\frac{1+t}{1-t} \right)^{\theta/\pi} \leq \left(\frac{1}{1-t} \right)^{2\theta/\pi} \quad \text{if } 0 \leq t \leq 1,$$

and

$$\frac{(1/(1-t))^{2\theta/\pi}}{((1+t)/(1-t))^{\theta/\pi}} = \left(\frac{1}{1-t^2} \right)^{\theta/\pi}$$

is an increasing function on $(0, 1)$. If we let

$$M = \int_{-1}^0 \left(\frac{1+t}{2} \right)^{\theta/\pi} dt,$$

then we may apply Lemma 2.2 and conclude that $t_\theta \leq \xi_\theta \leq \tau_\theta$, where τ_θ is given by

$$\int_{-1}^0 \left(\frac{1+t}{2} \right)^{\theta/\pi} dt + \int_0^{\tau_\theta} \left(\frac{1}{1-t} \right)^{2\theta/\pi} dt = \int_{\tau_\theta}^1 \left(\frac{1}{1-t} \right)^{2\theta/\pi} dt.$$

The integrals above have elementary antiderivatives. In order to show that $\tau_\theta \leq 2\theta/\pi$ for small θ , we choose $2\theta/\pi$ as the point of integration. By taking explicit antiderivatives and simplifying, it suffices to show that for small enough θ ,

$$\frac{\left(\frac{1}{2}\right)^{\theta/\pi}}{1+\theta/\pi} + \frac{1-2(1-2\theta/\pi)^{1-2\theta/\pi}}{1-2\theta/\pi} \geq 0. \quad (2-3)$$

The expression on the left of (2-3) approaches zero as $\theta \rightarrow 0$. If we take the derivative of the left side of (2-3) with respect to θ and let $\theta \rightarrow 0$ we obtain $(1+\ln(\frac{1}{2}))/\pi > 0$. Then (2-3) is true as long as $0 < \theta \leq \theta_0$ for $\theta_0 > 0$ chosen small enough. Hence we conclude that $t_\theta \leq \tau_\theta \leq 2\theta/\pi$ for any $0 < \theta \leq \theta_0$. \square

From (2-1) we have $|f'_\theta(z)| \rightarrow 1$ as $|z| \rightarrow \infty$. We let ϕ_θ be the harmonic function in Ω_θ defined by

$$y^+ = \phi_\theta(u, v),$$

where $f_\theta = u + iv$. Since $1 = |\nabla\phi_\theta||f'(z)|$, we have $|\nabla\phi_\theta| \rightarrow 1$ as $|z| \rightarrow \infty$. By a rotation of $\frac{\pi}{2}$ of ϕ_θ we have a complementary harmonic function $\tilde{\phi}_\theta$ and can thus apply the ACF monotonicity formula. We have $J(\infty, \phi_\theta, \tilde{\phi}_\theta) = 1$. To find $J(0+, \phi_\theta, \tilde{\phi}_\theta)$ we find $|\nabla\phi_\theta(0)|$. This is given by

$$1 = |\nabla\phi_\theta(0)||f'(t_\theta)|.$$

Thus

$$1 \geq |\nabla\phi_\theta(0)| = \left(\frac{1-t_\theta}{1+t_\theta}\right)^{\theta/\pi},$$

so $|\nabla\phi_\theta(0)|$ is an increasing function of θ , and

$$1 \geq |\nabla\phi_\theta(0)| \geq \left(\frac{1-2\theta/\pi}{1+2\theta/\pi}\right)^{\theta/\pi}.$$

Using L'Hospital's rule we conclude that

$$\lim_{\theta \rightarrow 0} \frac{1 - ((1-2\theta/\pi)/(1+2\theta/\pi))^{\theta/\pi}}{(\theta/\pi)^2} = 4 > 0.$$

As a consequence we have the following result:

Lemma 2.4. *There exists θ_0 such that if $0 < \theta \leq \theta_0$, then*

$$0 < 1 - \theta^2 < |\nabla\phi_\theta(0)| \leq 1. \quad (2-4)$$

Since $J(\infty, \phi_\theta) = 1$ and $J(0+, \phi_\theta) = |\nabla\phi_\theta(0)|$, Lemma 2.4 shows that

$$J(\infty, \phi_\theta) - J(0+, \phi_\theta) < 1 - \theta^2.$$

3. Bounded domain

The aim of this section is to transfer the inequality in (2-4) to a harmonic function on a bounded domain. We approximate Ω_θ with domains $\Omega_{\theta,M}$; see Figure 2. If $f_{\theta,M}$ is the conformal mapping of the upper half-plane onto $\Omega_{\theta,M}$, then

$$f'_{\theta,M}(z) = \left(\frac{z+1}{z-1}\right)^{\theta/\pi} \left(\frac{z-z_2}{z+z_2}\right)^{1/2} \left(\frac{z+z_1}{z-z_1}\right)^{1/2}, \quad (3-1)$$

where $z_1, z_2 \in \mathbb{R}$ and $1 < z_1 < z_2$. We again translate $f_{\theta,M}$ by a constant so that the domain is centered on the origin as in Figure 2. The points z_1, z_2 are chosen so that $f_{\theta,M}(z_2) = M + 0i$. We point out that $|f'_{\theta,M}| \rightarrow 1$ as $|z| \rightarrow \infty$. We define $\phi_{\theta,M}(u, v) = y^+$, where $f_{\theta,M} = u + iv$.

Lemma 3.1. *Fix $\theta \leq \theta_0$. There exists $M > 0$, possibly depending on θ , such that $J(\infty, \phi_{\theta,M}) = 1$ and $J(0+, \phi_{\theta,M}) > 1 - \theta^2$.*

Proof. That $J(\infty, \phi_{\theta,M}) = 1$ follows from the definition of $\phi_{\theta,M}$ and (3-1). Now from the explicit formulas given for $f'_\theta(z)$ and $f_{\theta,M}$ in (2-1) and (3-1) respectively, we have $\phi_{\theta,M} \rightarrow \phi_\theta$ in C^1 up to the boundary in a neighborhood of the origin. Since $|\nabla\phi_\theta(0)| > 1 - \theta^2$, the conclusion follows. \square




Figure 2. Domain $\Omega_{\theta,M}$.

Remark 3.2. Since $J(r, \phi_{\theta,M})$ is monotonically increasing in r , it follows from Lemma 3.1 that

$$J(\infty, \phi_{\theta,M}) - J(0+, \phi_{\theta,M}) < 1 - \theta^2.$$

For any $\theta \leq \theta_0$, we fix an M that satisfies Lemma 3.1. We now transfer the decrease in energy to a finite domain.

Lemma 3.3. *Let θ and $\phi_{\theta,M}$ be as in Lemma 3.1. Let $\Omega_{\theta,M}$ be defined as before. If we define w_R to be such that*

$$\begin{cases} \Delta w_R = 0 & \text{in } B_R \cap \Omega_{\theta,M}, \\ w_R = 0 & \text{on } \partial \Omega_{\theta,M} \cap B_R, \\ w_R = y & \text{on } (\partial B_R)^+, \end{cases}$$

then $w_R \rightarrow \phi_{\theta,M}$ locally uniformly in $\Omega_{\theta,M}$ and in C^1 in $B_\rho \cap \Omega_{\theta,M}$ for small enough ρ .

Proof. Using the rescaling

$$\phi_R := \frac{\phi_{\theta,M}(Rx, Ry)}{R},$$

we have $\phi_R \rightarrow y^+$ in C^1 on $(\partial B_1)^+$. Thus, for any $\eta > 0$, there exists $R_0 > 0$ such that if $R \geq R_0$, then

$$(1 - \eta)y^+ \leq \phi_R \leq (1 + \eta)y^+ \quad \text{on } (\partial B_1)^+.$$

Then rescaling back we obtain that $(1 - \eta)y^+ \leq \phi_{\theta,M} \leq (1 + \eta)y^+$ on $(\partial B_R)^+$ if $R \geq R_0$. From the maximum principle we then have

$$(1 - \eta)w_R \leq \phi_{\theta,M} \leq (1 + \eta)w_R \quad \text{for any } R \geq R_0.$$

Then as $R \rightarrow \infty$, we have $w_R \rightarrow w_\infty$ locally uniformly in $\Omega_{\theta,M}$ and in C^1 in a neighborhood of the origin. Furthermore, we have $(1 - \eta)w_\infty \leq \phi_{\theta,M} \leq (1 + \eta)w_\infty$. Since η can be taken to be arbitrarily small, we conclude that $w_\infty = \phi_{\theta,M}$. \square

We end this section by defining a θ -turn. If $u \in \mathcal{K}$ and for some $\rho > 0$ we have $\partial\{u > 0\} \cap B_\rho$ is a line segment with inward unit normal ν , then a θ -turn in B_ρ gives a new function v with

- (i) $v \in \mathcal{K}$,
- (ii) $v = u$ on ∂B_1 ,




Figure 3. θ -turn when $v = i$.

- (iii) $\partial\{v > 0\} \cap (B_1 \setminus \bar{B}_\rho) = \partial\{u > 0\} \cap (B_1 \setminus \bar{B}_\rho)$,
- (iv) $\partial\{v > 0\} \cap B_\rho = \partial\{\phi_{\theta, M}(e^{i(v-\theta)}(2M/\rho)z) > 0\} \cap B_\rho$.

The idea of property (iv) is to shrink $\phi_{\theta, M}$ on B_{2M} to B_ρ and give v the same positivity set; see Figure 3 for when $v = i$.

4. Construction of counterexample

As before we let θ_0 be as in Lemma 2.4. This next lemma shows how to apply a θ -turn to a function that is almost linear at the origin.

Lemma 4.1. *Fix $\epsilon > 0$. Assume $u \in \mathcal{K}$, and that there is $s < r_0 < 1$ with*

- (1) $B_s \cap \partial\{u > 0\} = B_s \cap \{y_n = 0\}$,
- (2) $|u| < 2J(1, u)r_0$ on B_{r_0} .

If $\theta \leq \theta_0$, then there exists r, ρ with $s > r > \rho > 0$ with a θ -turn in B_ρ such that if v is the redefined function, then v satisfies

- (A) $|v| < 2J(1, v)r$ on B_r ,
- (B) $|v| \leq (1 + \theta^2) \sup_{B_t} |u|$ on B_t for $t \in [r_0, 1]$,
- (C) $J(1, v) \leq (1 + \theta^2)J(1, u)$,
- (D) $J(0+, v) > (1 - \theta^2)^2 J(0+, u)$.

Proof. We choose $r < s$ small enough so that

$$\left\| \frac{u(rx)}{r} - J(0+, u)y^+ \right\|_{C^1((\partial B_1)^+)} < \delta, \quad (4-1)$$

and so that $|u| < 2J(1, u)r$. We now apply a θ -turn in B_ρ with $0 < \rho < r$. As $\rho \rightarrow 0$, we have $v \rightarrow u$ uniformly away from the origin, so that by choosing ρ small enough, v satisfies (B).

We now let $\eta > 0$ be small and use a cut-off function and obtain in the standard way the Caccioppoli inequality

$$\int_{B_1 \setminus B_\eta} |\nabla v - u|^2 \leq C(\eta) \int_{B_1 \setminus B_{\eta/2}} |v - u|^2.$$

Then as $\rho \rightarrow 0$, we have $v \rightarrow u$ in $H^1(B_1 \setminus B_\eta)$ for any $\eta > 0$. We now use the monotonicity of $J(r, v)$ to prove that $v \rightarrow u$ in $H^1(B_1)$ as $\rho \rightarrow 0$. We have

$$\int_{B_\eta} |\nabla v|^2 \leq \eta^2 \int_{B_1} |\nabla v|^2 = \delta^2 \int_{B_1 \setminus B_\eta} |\nabla v|^2 + \int_{B_\eta} |\nabla v|^2,$$

so that

$$\int_{B_\eta} |\nabla v|^2 \leq \frac{\eta^2}{1 - \eta^2} \int_{B_1 \setminus B_\eta} |\nabla v|^2,$$

and we conclude that

$$\int_{B_1} |\nabla v|^2 \leq \frac{1}{1 - \eta^2} \int_{B_1 \setminus B_\eta} |\nabla v|^2.$$

Then $\|v\|_{H^1(B_1)}$ is bounded as $\rho \rightarrow 0$, so that $v \rightarrow u$ in $H^1(B_1)$ as $\rho \rightarrow 0$. We now have

$$\int_{B_1} |\nabla u|^2 \leq \lim_{\rho \rightarrow 0} \int_{B_1} |\nabla v|^2 \leq \lim_{\rho \rightarrow 0} \frac{1}{1 - \eta^2} \int_{B_1 \setminus B_\eta} |\nabla v|^2 = \frac{1}{1 - \eta^2} \int_{B_1 \setminus B_\eta} |\nabla u|^2.$$

Since η can be chosen arbitrarily small, we have $\nabla v \rightarrow \nabla u$ in $L^2(B_1)$ and thus conclude that $v \rightarrow u$ in $H^1(B_1)$ as $\rho \rightarrow 0$. Consequently, we may choose ρ even smaller so that properties (A) and (C) hold.

From (4-1), if ρ is chosen small enough we have

$$\left\| \frac{v(rz)}{r} - J(0+, u)y^+ \right\|_{C^1((\partial B_1)^+)} < \delta,$$

so that $(1 - \delta)J(0+, u)y^+ \leq v(rz)/r$ on $(\partial B_1)^+$. We now define w to be the solution to

$$\begin{cases} \Delta w = 0 & \text{in } \{v(rz)/r > 0\} \cap B_1, \\ w = 0 & \text{on } \partial\{v(rz)/r > 0\} \cap B_1, \\ w = (1 - \delta)J(0+)y^+ & \text{on } (\partial B_1)^+. \end{cases}$$

We have $w \leq v$ in B_1 , so that $|\nabla w(0)| \leq |\nabla v(0)|$ or $J(0+, w) \leq J(0+, v)$. We may rescale w and apply Lemma 3.3 to obtain that for small enough ρ , we have

$$J(0+, w) > (1 - \theta^2)(1 - \delta)J(0+, u).$$

By choosing $\delta < \theta^2$ we obtain (D). \square

Proof of Theorem 1.3 in dimension $n = 2$. We now use Lemma 4.1 to construct a sequence $u_k \in \mathcal{K}$ with $\lim u_k \rightarrow u$. The pair u and $\tilde{u}(z) := u(-z)$ will be a counterexample to Claim 1.2. The sequence u_k is constructed inductively as follows. We choose $\theta_k = 1/(k + N_0)$, where $N_0 \in \mathbb{N}$ is chosen large enough so that $\theta_k \leq \theta_0$. We then let $u_0 = y^+$ on B_1 . By Lemma 4.1 there exists $\rho_1 < r_1$ such that if a θ_1 -turn is applied in B_{ρ_1} to obtain u_1 , then u_1 will satisfy properties (A)–(D). We now suppose that u_k has been constructed for some $k \geq 1$. By rotating u_k it will satisfy assumption (1) of Lemma 4.1. Assumption (2) will also be satisfied because u_k satisfies (A) for $r = r_k$. By Lemma 4.1 there exists $\rho_{k+1} < r_{k+1}$ with $r_{k+1} < \rho_k$ so that if we apply a θ_{k+1} -turn to u_k to obtain u_{k+1} we have

(i) $|u_{k+1}| < 2J(1, u_{k+1})r$ on B_r ,

- (ii) $|u_{k+1}| \leq \prod_{j=1}^k (1 + \theta_j^2) \sup_{B_t} |u_0|$ on B_t for $t \in [r_k, 1]$,
- (iii) $J(1, u_{k+1}) \leq \prod_{j=1}^k (1 + \theta_j^2) J(1, u_0) = \prod_{j=1}^k (1 + \theta_j^2)$,
- (iv) $J(0+, u_{k+1}) > \prod_{j=1}^k (1 - \theta_j^2)^2 J(0+, u_0) = \prod_{j=1}^k (1 - \theta_j^2)^2$.

From the same arguments involving the Caccioppoli inequality as in the proof of Lemma 4.1, there exists u such that $u_k \rightarrow u$ in $H^1(B_1)$ and locally uniformly away from the origin. Then u is continuous away from the origin. From (i) we obtain that $|u| \leq Cr$ on B_r for $0 < r \leq 1$, so that u is continuous up to the origin, and $u(0) = 0$.

Now $0 < \prod_{k=1}^{\infty} (1 - \theta_k^2)^2$ if and only if $0 < \prod_{k=1}^{\infty} (1 - \theta_k^2)$ if and only if

$$\sum_{k=1}^{\infty} (k + N_0)^{-2} = \sum_{k=1}^{\infty} \theta_k^2 < \infty.$$

Since the inequality above is true, we conclude that

$$0 < \prod_{k=1}^{\infty} (1 - \theta_k^2)^2 < \prod_{k=1}^{\infty} (1 - \theta_k^2) < 1.$$

The last inequality above is due to the fact that all the terms are less than 1. Since $u_k \rightarrow u$ in $H^1(B_1)$ and from properties (ii) and (iii), we conclude that

$$0 < \prod_{k=1}^{\infty} (1 - \theta_k^2)^2 \leq J(r, u) \leq C J(1, u) < \infty \quad \text{for all } 0 < r \leq 1,$$

so that $J(0+, u) > 0$.

If we let $\tilde{u}_k(z) = u_k(-z)$, then $\tilde{u}_k \rightarrow \tilde{u}$, where $\tilde{u}(z) = u(-z)$. Furthermore, $u \cdot \tilde{u} = 0$ in B_1 . Since also u, \tilde{u} are nonnegative, continuous, and harmonic when positive, they satisfy the assumptions of the ACF monotonicity formula in Proposition 1.1. We now show that u, \tilde{u} are a counterexample to Claim 1.2. We assume by way of contradiction that $\{u > 0\}$ and $\{\tilde{u} > 0\}$ are tangent at the origin and after a rotation $u(z) + \tilde{u}(z) = \alpha x_1^+ + \beta x_1^- + o(|z|)$. Then for any small $\delta > 0$, there exists r_0 such that if $r \leq r_0$ and $|z| > \frac{1}{2}$ and $|\text{Arg}(z)| < \delta$, then

$$\frac{u(rz) + \tilde{u}(rz)}{r} > \frac{\alpha x_1^+}{2} > 0. \quad (4-2)$$

We now recall that from the construction

$$\partial\{u > 0\} \cap (B_{r_k} \setminus B_{\rho_k}) = \{z : z = te^{-i \sum_{j=1}^k \theta_j} \text{ and } \rho_k \leq |t| < r_k\}. \quad (4-3)$$

Since $\sum \theta_k = \infty$ and $\theta_k \rightarrow 0$, we obtain from (4-3) there exist infinitely many z_k with $|z_k| \rightarrow 0$ and $|\text{Arg}(z)| < \delta$ such that $u(z_k) + \tilde{u}(z_k) = 0$. This contradicts (4-2), and so Claim 1.2 is not true. \square

We now show that the pair u and \tilde{u} are also a counterexample in higher dimensions.

Proof of Theorem 1.3 in dimension $n > 2$. For u as in the proof for dimension 2, we let $w_n(x_1, x_2, \dots, x_n) = u(x_1, x_2)$. Since in dimension $n = 2$ we have

$$\frac{1}{r^2} \int_{B_r} |\nabla u|^2 \geq C > 0,$$

it follows that in dimension n ,

$$\frac{1}{r^n} \int_{B_r} |\nabla w|^2 \geq C.$$

Then

$$\frac{1}{r^2} \int_{B_r} \frac{|\nabla w|^2}{|x|^{n-2}} \geq \frac{1}{r^2} \int_{B_r} \frac{|\nabla w|^2}{r^{n-2}} = \frac{1}{r^n} \int_{B_r} |\nabla w|^2 \geq C > 0,$$

so that $\Phi(r, w, \tilde{w}) > 0$. We have already shown that $u + \tilde{u}$ cannot satisfy the conclusions in Claim 1.2; consequently, $w + \tilde{w}$ also do not satisfy those conclusions. \square

Acknowledgments

Kriventsov was supported by the NSF MSPRF fellowship DMS-1502852. The authors would like to thank Luis Caffarelli for helpful conversations regarding this problem.

References

- [Allen and Petrosyan 2012] M. Allen and A. Petrosyan, “A two-phase problem with a lower-dimensional free boundary”, *Interfaces Free Bound.* **14**:3 (2012), 307–342. MR Zbl
- [Allen and Shi 2016] M. Allen and W. Shi, “The two-phase parabolic Signorini problem”, *Indiana Univ. Math. J.* **65**:2 (2016), 727–742. MR Zbl
- [Allen et al. 2015] M. Allen, E. Lindgren, and A. Petrosyan, “The two-phase fractional obstacle problem”, *SIAM J. Math. Anal.* **47**:3 (2015), 1879–1905. MR Zbl
- [Alt et al. 1984] H. W. Alt, L. A. Caffarelli, and A. Friedman, “Variational problems with two phases and their free boundaries”, *Trans. Amer. Math. Soc.* **282**:2 (1984), 431–461. MR Zbl
- [Azzam et al. 2016] J. Azzam, S. Hofmann, J. M. Martell, S. Mayboroda, M. Mourgoglou, X. Tolsa, and A. Volberg, “Rectifiability of harmonic measure”, *Geom. Funct. Anal.* **26**:3 (2016), 703–728. MR Zbl
- [Blank 2001] I. Blank, “Sharp results for the regularity and stability of the free boundary in the obstacle problem”, *Indiana Univ. Math. J.* **50**:3 (2001), 1077–1112. MR Zbl
- [Caffarelli and Salsa 2005] L. Caffarelli and S. Salsa, *A geometric approach to free boundary problems*, Graduate Studies in Math. **68**, Amer. Math. Soc., Providence, RI, 2005. MR Zbl
- [Caffarelli et al. 2009] L. A. Caffarelli, A. L. Karakhanian, and F.-H. Lin, “The geometry of solutions to a segregation problem for nondivergence systems”, *J. Fixed Point Theory Appl.* **5**:2 (2009), 319–351. MR Zbl
- [Kriventsov and Lin 2018] D. Kriventsov and F. Lin, “Regularity for shape optimizers: the nondegenerate case”, *Comm. Pure Appl. Math.* **71**:8 (2018), 1535–1596. MR Zbl
- [Kriventsov and Lin 2019] D. Kriventsov and F. Lin, “Regularity for shape optimizers: the degenerate case”, *Comm. Pure Appl. Math.* **72**:8 (2019), 1678–1721. MR
- [Petrosyan et al. 2012] A. Petrosyan, H. Shahgholian, and N. Uraltseva, *Regularity of free boundaries in obstacle-type problems*, Graduate Studies in Math. **136**, Amer. Math. Soc., Providence, RI, 2012. MR Zbl
- [Terracini et al. 2019] S. Terracini, G. Verzini, and A. Zilio, “Spiraling asymptotic profiles of competition-diffusion systems”, *Comm. Pure Appl. Math.* (online publication March 2019).
- [White 1992] B. White, “Nonunique tangent maps at isolated singularities of harmonic maps”, *Bull. Amer. Math. Soc. (N.S.)* **26**:1 (1992), 125–129. MR Zbl

Received 26 Feb 2018. Revised 13 Sep 2018. Accepted 19 Dec 2018.

MARK ALLEN: allen@mathematics.byu.edu

Department of Mathematics, Brigham Young University, Provo, UT, United States

DENNIS KRIVENTSOV: dennisk@cims.nyu.edu

Courant Institute of Mathematical Sciences, New York University, New York, NY, United States

Analysis & PDE

msp.org/apde

EDITORS

EDITOR-IN-CHIEF

Patrick Gérard

patrick.gerard@math.u-psud.fr

Université Paris Sud XI

Orsay, France

BOARD OF EDITORS

Massimiliano Berti	Scuola Intern. Sup. di Studi Avanzati, Italy berti@sissa.it	Gilles Pisier	Texas A&M University, and Paris 6 pisier@math.tamu.edu
Michael Christ	University of California, Berkeley, USA mchrist@math.berkeley.edu	Tristan Rivière	ETH, Switzerland riviere@math.ethz.ch
Charles Fefferman	Princeton University, USA cf@math.princeton.edu	Igor Rodnianski	Princeton University, USA irod@math.princeton.edu
Ursula Hamenstädt	Universität Bonn, Germany ursula@math.uni-bonn.de	Yum-Tong Siu	Harvard University, USA siu@math.harvard.edu
Vadim Kaloshin	University of Maryland, USA vadim.kaloshin@gmail.com	Terence Tao	University of California, Los Angeles, USA tao@math.ucla.edu
Herbert Koch	Universität Bonn, Germany koch@math.uni-bonn.de	Michael E. Taylor	Univ. of North Carolina, Chapel Hill, USA met@math.unc.edu
Izabella Laba	University of British Columbia, Canada ilaba@math.ubc.ca	Gunther Uhlmann	University of Washington, USA gunther@math.washington.edu
Richard B. Melrose	Massachusetts Inst. of Tech., USA rbm@math.mit.edu	András Vasy	Stanford University, USA andras@math.stanford.edu
Frank Merle	Université de Cergy-Pontoise, France Frank.Merle@u-cergy.fr	Dan Virgil Voiculescu	University of California, Berkeley, USA dvv@math.berkeley.edu
William Minicozzi II	Johns Hopkins University, USA minicozz@math.jhu.edu	Steven Zelditch	Northwestern University, USA zelditch@math.northwestern.edu
Clément Mouhot	Cambridge University, UK c.mouhot@dpmms.cam.ac.uk	Maciej Zworski	University of California, Berkeley, USA zworski@math.berkeley.edu
Werner Müller	Universität Bonn, Germany mueller@math.uni-bonn.de		

PRODUCTION

production@msp.org

Silvio Levy, Scientific Editor

See inside back cover or msp.org/apde for submission instructions.

The subscription price for 2020 is US \$340/year for the electronic version, and \$550/year (+\$60, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues from the last three years and changes of subscriber address should be sent to MSP.

Analysis & PDE (ISSN 1948-206X electronic, 2157-5045 printed) at Mathematical Sciences Publishers, 798 Evans Hall #3840, c/o University of California, Berkeley, CA 94720-3840, is published continuously online. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices.

APDE peer review and production are managed by EditFlow® from MSP.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2020 Mathematical Sciences Publishers

ANALYSIS & PDE

Volume 13 No. 1 2020

Absence of Cartan subalgebras for right-angled Hecke von Neumann algebras MARTIJN CASPERS	1
A vector field method for radiating black hole spacetimes JESÚS OLIVER and JACOB STERBENZ	29
Stable ODE-type blowup for some quasilinear wave equations with derivative-quadratic nonlinearities JARED SPECK	93
Asymptotic expansions of fundamental solutions in parabolic homogenization JUN GENG and ZHONGWEI SHEN	147
Capillary surfaces arising in singular perturbation problems ARAM L. KARAKHANYAN	171
A spiral interface with positive Alt–Caffarelli–Friedman limit at the origin MARK ALLEN and DENNIS KRIVENTSOV	201
Infinite-time blow-up for the 3-dimensional energy-critical heat equation MANUEL DEL PINO, MONICA MUSSO and JUNCHENG WEI	215
A well-posedness result for viscous compressible fluids with only bounded density RAPHAËL DANCHIN, FRANCESCO FANELLI and MARIUS PAICU	275



2157-5045(2020)13:1;1-D