

ANALYSIS & PDE

Volume 13 No. 2 2020

GIANMARCO BROCCHI, DIOGO OLIVEIRA E SILVA
AND RENÉ QUILODRÁN

**SHARP STRICHARTZ INEQUALITIES FOR FRACTIONAL AND
HIGHER-ORDER SCHRÖDINGER EQUATIONS**

SHARP STRICHARTZ INEQUALITIES FOR FRACTIONAL AND HIGHER-ORDER SCHRÖDINGER EQUATIONS

GIANMARCO BROCCHI, DIOGO OLIVEIRA E SILVA AND RENÉ QUILODRÁN

We investigate a class of sharp Fourier extension inequalities on the planar curves $s = |y|^p$, $p > 1$. We identify the mechanism responsible for the possible loss of compactness of nonnegative extremizing sequences, and prove that extremizers exist if $1 < p < p_0$ for some $p_0 > 4$. In particular, this resolves the dichotomy of Jiang, Pausader, and Shao concerning the existence of extremizers for the Strichartz inequality for the fourth-order Schrödinger equation in one spatial dimension. One of our tools is a geometric comparison principle for n -fold convolutions of certain singular measures in \mathbb{R}^d , developed in the companion paper of Oliveira e Silva and Quilodrán (*Math. Proc. Cambridge Philos. Soc.*, (2019)). We further show that any extremizer exhibits fast L^2 -decay in physical space, and so its Fourier transform can be extended to an entire function on the whole complex plane. Finally, we investigate the extent to which our methods apply to the case of the planar curves $s = y|y|^{p-1}$, $p > 1$.

1. Introduction	477
2. Bilinear estimates and cap refinements	483
3. Existence versus concentration	491
4. Existence of extremizers	497
5. Superexponential L^2 -decay	507
6. The case of odd curves	510
Appendix A. Concentration-compactness	520
Appendix B. Revisiting Brézis–Lieb	522
Acknowledgements	524
References	524

1. Introduction

Gaussians are known to extremize certain Strichartz estimates in low dimensions. Consider, for instance, the Strichartz inequality for the homogeneous Schrödinger equation in d spatial dimensions,

$$\|e^{-it\Delta} f\|_{L_{x,t}^{2+4/d}(\mathbb{R}^{d+1})} \leq S(d) \|f\|_{L^2(\mathbb{R}^d)}, \quad (1-1)$$

Oliveira e Silva was supported by the Hausdorff Center for Mathematics and DFG grant CRC 1060.

MSC2010: 35B38, 35Q41, 42B37.

Keywords: sharp Fourier restriction theory, extremizers, Strichartz inequalities, fractional Schrödinger equation, convolution of singular measures.

with optimal constant given by

$$S(d) := \sup_{0 \neq f \in L^2} \frac{\|e^{-it\Delta} f\|_{L^{2+4/d}_{x,t}(\mathbb{R}^{d+1})}}{\|f\|_{L^2(\mathbb{R}^d)}}. \quad (1-2)$$

That $S(d) < \infty$ is of course due to the original work of Strichartz [1977], which in turn had precursors in [Tomas 1975; Segal 1976]. If $d \in \{1, 2\}$, then Gaussians extremize (1-1), and therefore $S(1) = 12^{-1/12}$ and $S(2) = 2^{-1/2}$. This was originally established in [Foschi 2007; Hundertmark and Zharnitsky 2006], and alternative proofs were subsequently given in [Bennett et al. 2009; 2015; Gonçalves 2019]. All of these approaches ultimately rely on the fact that the Strichartz exponent $2 + \frac{4}{d}$ is an even integer if $d \in \{1, 2\}$, which in turn allows us to recast inequality (1-1) in convolution form. This is a powerful technique that has proved very successful in tackling a number of problems in sharp Fourier restriction theory; see the recent survey [Foschi and Oliveira e Silva 2017].

In the recent work [Oliveira e Silva and Quilodrán 2018], we explored the convolution structure of a family of Strichartz inequalities for higher-order Schrödinger equations in two spatial dimensions in order to answer a question concerning the existence of extremizers that had appeared in the previous literature. Our purpose with the present work is three-fold. Firstly, we resolve the dichotomy from [Jiang et al. 2010] concerning the existence of extremizers for the Strichartz inequality for the fourth-order Schrödinger equation in one spatial dimension. This is related to the Fourier extension problem on the planar curve $s = y^4$. Secondly, we study similar questions in the more general setting of the Fourier extension problem on the curve $s = |y|^p$ for arbitrary $p > 1$. We also consider *odd* curves $s = y|y|^{p-1}$, $p > 1$, the case $p = 3$ relating to the Airy–Strichartz inequality [Farah and Versieux 2018; Frank and Sabin 2018; Shao 2009]. Lastly, we study superexponential decay and analyticity of the corresponding extremizers and their Fourier transform via a bootstrapping procedure.

Jiang, Pausader, and Shao [Jiang et al. 2010] considered the fourth-order Schrödinger equation with L^2 initial datum in one spatial dimension,

$$\begin{cases} i\partial_t u - \mu\partial_x^2 u + \partial_x^4 u = 0, & (x, t) \in \mathbb{R} \times \mathbb{R}, \\ u(\cdot, 0) = f \in L_x^2(\mathbb{R}), \end{cases} \quad (1-3)$$

where $u : \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{C}$, and $\mu \geq 0$. By scaling, one may restrict attention to $\mu \in \{0, 1\}$. The solution of the Cauchy problem (1-3) can be written in terms of the propagator

$$u(x, t) = e^{it(\partial_x^4 - \mu\partial_x^2)} f(x) = \frac{1}{2\pi} \int_{\mathbb{R}} e^{ix\xi} e^{it(\xi^4 + \mu\xi^2)} \hat{f}(\xi) d\xi,$$

where the spatial Fourier transform is defined as¹

$$\hat{f}(\xi) := \int_{\mathbb{R}} e^{-ix\xi} f(x) dx.$$

¹The Fourier transform will occasionally be denoted by $\mathcal{F}(f) = \hat{f}$.

The solution disperses as $|t| \rightarrow \infty$, and consequently the following Strichartz inequality due to Kenig, Ponce, and Vega [Kenig et al. 1991, Theorem 2.1] holds:²

$$\|D_\mu^{\frac{1}{3}} e^{it(\partial_x^4 - \mu \partial_x^2)} f\|_{L_{x,t}^6(\mathbb{R}^{1+1})} \lesssim \|f\|_{L^2(\mathbb{R})}. \quad (1-4)$$

The main result of [Jiang et al. 2010] is a linear profile decomposition for (1-3), which uses a refinement of the Strichartz inequality (1-4) in the scale of Besov spaces, together with improved localized Fourier restriction estimates. As a consequence, the authors of [Jiang et al. 2010] establish a dichotomy result for the existence of extremizers for (1-4) when $\mu = 0$, which can be summarized as follows: Consider the sharp inequality in multiplier form

$$\|D_0^{\frac{1}{3}} e^{it\partial_x^4} f\|_{L_{x,t}^6(\mathbb{R}^{1+1})} \leq M \|f\|_{L^2(\mathbb{R})}, \quad (1-5)$$

with optimal constant given by

$$M := \sup_{0 \neq f \in L^2} \frac{\|D_0^{\frac{1}{3}} e^{it\partial_x^4} f\|_{L_{x,t}^6(\mathbb{R}^{1+1})}}{\|f\|_{L^2(\mathbb{R})}}. \quad (1-6)$$

Then [Jiang et al. 2010, Theorem 1.8] states that either an extremizer for (1-5) exists, or there exist a sequence $\{a_n\} \subset \mathbb{R}$ satisfying $|a_n| \rightarrow \infty$ as $n \rightarrow \infty$ and a function $f \in L^2$ such that

$$M = \lim_{n \rightarrow \infty} \frac{\|D_0^{\frac{1}{3}} e^{it\partial_x^4} (e^{ia_n x} f)\|_{L_{x,t}^6(\mathbb{R}^{1+1})}}{\|f\|_{L^2(\mathbb{R})}}.$$

In the latter case, one necessarily has $M = S(1)$, where $S(1)$ denotes the optimal constant defined in (1-2). Our first main result resolves this dichotomy.

Theorem 1.1. *There exists an extremizer for (1-5).*

Theorem 1.1 will follow from a more general result which we now introduce. As noted in [Kenig et al. 1991, §2], the operator $D_0^{1/3} e^{it\partial_x^4}$ is nothing but a constant multiple of the Fourier transform at the point $(-x, -t) \in \mathbb{R}^2$ of the singular measure

$$d\sigma_4(y, s) = \delta(s - y^4) |y|^{\frac{1}{3}} dy ds \quad (1-7)$$

defined on the curve $s = y^4$. As in [Oliveira e Silva and Quilodrán 2018, §6.4], one is naturally led to consider generic power curves $s = |y|^p$. The corresponding inequality is

$$\|\mathcal{M}_p(f)\|_{L_{x,t}^6(\mathbb{R}^{1+1})} \leq M_p \|f\|_{L^2(\mathbb{R})}, \quad (1-8)$$

where the multiplier operator \mathcal{M}_p is defined as

$$\mathcal{M}_p(f)(x, t) = D_0^{\frac{p-2}{6}} e^{it|\partial_x|^p} f(x).$$

²Given $\mu \in \{0, 1\}$ and $\alpha \in \mathbb{R}$, we follow the notation from [Jiang et al. 2010] and denote by D_μ^α the differentiation operator $D_\mu^\alpha f(x) := \frac{1}{2\pi} \int_{\mathbb{R}} e^{ix\xi} (\mu + 6\xi^2)^{\alpha/2} \hat{f}(\xi) d\xi$.

Inequality (1-8) can be equivalently restated as a Fourier extension inequality,

$$\|\mathcal{E}_p(f)\|_{L^6(\mathbb{R}^2)} \leq E_p \|f\|_{L^2(\mathbb{R})}, \quad (1-9)$$

or in convolution form as

$$\|f\sigma_p * f\sigma_p * f\sigma_p\|_{L^2(\mathbb{R}^2)} \leq C_p^3 \|f\|_{L^2(\mathbb{R})}^3. \quad (1-10)$$

Here, the singular measure σ_p is defined in accordance with (1-7) by

$$d\sigma_p(y, s) = \delta(s - |y|^p) |y|^{\frac{p-2}{6}} dy ds, \quad (1-11)$$

and the Fourier extension operator $\mathcal{E}_p(f) = \mathcal{F}(f\sigma_p)(-\cdot)$ is given by

$$\mathcal{E}_p(f)(x, t) = \int_{\mathbb{R}} e^{ixy} e^{it|y|^p} |y|^{\frac{p-2}{6}} f(y) dy, \quad (1-12)$$

so that

$$6^{\frac{p-2}{12}} \mathcal{E}_p(\hat{f}) = 2\pi \mathcal{M}_p(f).$$

If f is an extremizer for (1-9), then f is likewise an extremizer for (1-10), and $\mathcal{F}^{-1}(f)$ is an extremizer for (1-8). Thus these three existence problems are essentially equivalent. The convolution form (1-10) also shows that the search for extremizers can be restricted to the class of *nonnegative* functions. An application of Plancherel's theorem further reveals that the corresponding optimal constants satisfy

$$E_p^6 = (2\pi)^2 C_p^6 = (2\pi)^3 6^{1-\frac{p}{2}} M_p^6.$$

Our next result extends the dichotomy proved in [Jiang et al. 2010, Theorem 1.8] to the case of arbitrary exponents $p > 1$. It states that one of two possible scenarios occurs, *compactness* or *concentration* at a point. We make the latter notion precise.

Definition 1.2. A sequence of functions $\{f_n\} \subset L^2(\mathbb{R})$ *concentrates at a point* $y_0 \in \mathbb{R}$ if, for every $\varepsilon, \rho > 0$, there exists $N \in \mathbb{N}$ such that, for every $n \geq N$,

$$\int_{|y-y_0| \geq \rho} |f_n(y)|^2 dy < \varepsilon \|f_n\|_{L^2(\mathbb{R})}^2.$$

We choose to phrase our second main result in terms of the convolution inequality (1-10) because, as we shall see, condition (1-13) has a very simple geometric meaning in terms of the boundary value of the relevant 3-fold convolution measure.

Theorem 1.3. *Let $p > 1$. If*

$$C_p^6 > \frac{2\pi}{\sqrt{3p(p-1)}}, \quad (1-13)$$

then any extremizing sequence of nonnegative functions in $L^2(\mathbb{R})$ for (1-10) is precompact, after normalization and scaling. In this case, extremizers for (1-10) exist. If instead equality holds in (1-13) then, given any $y_0 \in \mathbb{R}$, there exists an extremizing sequence for (1-10) which concentrates at y_0 .

A few remarks may help to further orient the reader. Firstly, if $p = 1$, then the curve $s = |y|$ has no curvature, and no nontrivial Fourier extension estimate can hold. Secondly, if equality holds in (1-13), then **Theorem 1.3** does *not* guarantee the nonexistence of extremizers. Indeed, $C_2^6 = \pi/\sqrt{3}$, and Gaussians are known to extremize (1-10) when $p = 2$. Various results of a similar flavor to that of **Theorem 1.3** have appeared in the recent literature. They are typically derived from a sophisticated application of concentration-compactness techniques [Christ and Shao 2012a; Shao 2016a], a full profile decomposition [Jiang et al. 2010; 2014; Shao 2009], or the missing mass method as in [Frank et al. 2016; Frank and Sabin 2018]. We introduce a new variant which follows the spirit of the celebrated works [Brézis and Lieb 1983; Lieb 1983; Lions 1984a; 1984b]. It seems more elementary and may be easier to adapt to other manifolds. The proof of **Theorem 1.3** involves a variant of Lions' concentration-compactness lemma [1984a], a variant of the corollary of the Brézis–Lieb lemma from [Fanelli et al. 2011], bilinear extension estimates, and a refinement of inequality (1-9) over a suitable *cap space*.

In a range of exponents that includes the case $p = 4$, we are able to resolve the dichotomy posed by **Theorem 1.3**.

Theorem 1.4. *There exists $p_0 > 4$ such that, for every $p \in (1, p_0) \setminus \{2\}$, the strict inequality (1-13) holds. In particular, if $p \in (1, p_0)$, then there exists an extremizer for (1-10).*

Our method yields $p_0 \approx 4.803$ with three decimal places, and effectively computes arbitrarily good lower bounds for the ratio of L^2 -norms in (1-10) via expansions of suitable trial functions in the orthogonal basis of Legendre polynomials. We remark that the value $p_0 \approx 4.803$ is suboptimal, in the sense that a natural refinement of our argument allows us to increase this value to ≈ 5.485 ; see **Section 4C** below.

Once the existence of extremizers has been established, their properties are typically deduced from the study of the associated Euler–Lagrange equation. Following this paradigm, we show that any extremizer of (1-9) decays superexponentially fast in L^2 , which reflects the analyticity of its Fourier transform. This is the content of our next result.

Theorem 1.5. *Let $p > 1$. If f is an extremizer for (1-9), then there exists $\mu_0 > 0$ such that*

$$x \mapsto e^{\mu_0|x|^p} f(x) \in L^2(\mathbb{R}).$$

In particular, its Fourier transform \hat{f} can be extended to an entire function on \mathbb{C} .

Note that the exponent μ_0 necessarily depends on the extremizer itself; see the discussion in [Christ and Shao 2012b, p. 964]. The proof relies on a bootstrapping argument that found similar applications in [Christ and Shao 2012b; Erdogan et al. 2011; Hundertmark and Shao 2012; Shao 2016b].

To some extent, our methods are able to handle the case of the planar *odd* curves $s = y|y|^{p-1}$, $p > 1$. Define the singular measure

$$d\mu_p(y, s) = \delta(s - y|y|^{p-1})|y|^{\frac{p-2}{6}} dy ds. \quad (1-14)$$

The associated Fourier extension operator $\mathcal{S}_p(f) = \mathcal{F}(f\mu_p)(-\cdot)$, defined in (6-2) below, satisfies the estimate $\|\mathcal{S}_p(f)\|_{L^6} \lesssim \|f\|_{L^2}$. In sharp convolution form, this can be rewritten as

$$\|f\mu_p * f\mu_p * f\mu_p\|_{L^2(\mathbb{R}^2)} \leq Q_p^3 \|f\|_{L^2(\mathbb{R})}^3, \quad (1-15)$$

where Q_p denotes the optimal constant. Odd curves are of independent interest, in particular because a new phenomenon emerges: caps centered around points with parallel tangents interact strongly, regardless of separation between the points. This mechanism was discovered in [Christ and Shao 2012a], and further explored in [Carneiro et al. 2017; Foschi 2015; Frank et al. 2016; Frank and Sabin 2018; Shao 2016a]. Some of these works include a symmetrization step which relies on the convolution structure of the underlying inequality. In the present case, we also show that the search for extremizers can be further restricted to the class of even functions, but interestingly our symmetrization argument does not depend on the convolution structure. This may be of independent interest since it applies to other Fourier extension inequalities where some additional symmetry is present, as we indicate in [Section 6A](#) below.

The following versions of Theorems 1.3 and 1.4 hold for odd curves.

Theorem 1.6. *Let $p > 1$. If*

$$Q_p^6 > \frac{5\pi}{\sqrt{3}p(p-1)}, \quad (1-16)$$

then any extremizing sequence of nonnegative, even functions in $L^2(\mathbb{R})$ for (1-15) is precompact, after normalization and scaling. In this case, extremizers for (1-15) exist. If instead equality holds in (1-16) then, given any $y_0 \in \mathbb{R}$, there exists an extremizing sequence for (1-15) which concentrates at the pair $\{-y_0, y_0\}$.

The case $p = 3$ of [Theorem 1.6](#) coincides with a special case of [Frank and Sabin 2018, Theorem 1], which was obtained by different methods.

Theorem 1.7. *If $p \in (1, 2)$, then the strict inequality (1-16) holds and, in particular, there exists an extremizer for (1-15).*

We believe that extremizers do not exist if $p \geq 2$; see [Conjecture 6.6](#) below.

Overview. The paper is organized as follows. [Section 2](#) is devoted to the technical preliminaries for the dichotomy statement concerning the existence of extremizers: bilinear estimates and cap bounds. We then prove [Theorem 1.3](#) in [Section 3](#). Existence of extremizers is the subject of [Section 4](#), where we establish [Theorem 1.4](#). [Theorem 1.5](#) addresses the regularity of extremizers and is established in [Section 5](#). Odd curves are treated in [Section 6](#), where Theorems 1.6 and 1.7 are proved. In the [Appendix](#), we establish useful variants of Lions' concentration-compactness lemma ([Proposition A.1](#)) and of a corollary of the Brézis–Lieb lemma ([Proposition B.1](#)).

Notation. If x, y are real numbers, we write $x = O(y)$ or $x \lesssim y$ if there exists a finite absolute constant C such that $|x| \leq C|y|$. If we want to make explicit the dependence of the constant C on some parameter α , we write $x = O_\alpha(y)$ or $x \lesssim_\alpha y$. We write $x \gtrsim y$ if $y \lesssim x$, and $x \simeq y$ if $x \lesssim y$ and $x \gtrsim y$. Finally, the indicator function of a set $E \subset \mathbb{R}^d$ will be denoted by $\mathbb{1}_E$, and the complement of E will at times be denoted by E^c .

2. Bilinear estimates and cap refinements

In this section, we prove the bilinear extension estimates and cap refinements which will be needed in the next section. Bilinear extension estimates are usually deep [Tao 2003; Wolff 2001], but in the one-dimensional case one may rely on the classical Hausdorff–Young inequality. Throughout this section, we shall consider the dyadic regions

$$I_k := [2^k, 2^{k+1}) \quad \text{and} \quad I_k^\bullet := (-2^{k+1}, -2^k] \cup [2^k, 2^{k+1}) \quad (k \in \mathbb{Z}).$$

2A. Bilinear estimates. Recall the definitions (1-11) and (1-12) of the measure σ_p and the Fourier extension operator \mathcal{E}_p , respectively. Our first result quantifies the principle that distant caps interact weakly.

Proposition 2.1. *Let $p > 1$ and $k, k' \in \mathbb{Z}$. Then*

$$\|\mathcal{E}_p(f)\mathcal{E}_p(g)\|_{L^3(\mathbb{R}^2)} \lesssim_p 2^{-|k-k'|\frac{p-1}{6}} \|f\|_{L^2(\mathbb{R})} \|g\|_{L^2(\mathbb{R})} \quad (2-1)$$

for every $f, g \in L^2(\mathbb{R})$ satisfying $\text{supp } f \subseteq I_k^\bullet$ and $\text{supp } g \subseteq I_{k'}^\bullet$.

Proof. Setting $\psi = |\cdot|^p$ and $w = |\cdot|^{\frac{p-2}{3}}$, we have

$$(\mathcal{E}_p(f)\mathcal{E}_p(g))(x, t) = \int_{\mathbb{R}^2} e^{ix(y+y')} e^{it(\psi(y)+\psi(y'))} f(y) g(y') w(y)^{\frac{1}{2}} w(y')^{\frac{1}{2}} dy dy'.$$

Change variables $(y, y') \mapsto (u, v) = (y + y', \psi(y) + \psi(y'))$. Except for null sets, this is a 2-to-1 map from \mathbb{R}^2 onto the region $\{(u, v): v \geq 2\psi(u/2)\}$. Its Jacobian is given by

$$J^{-1}(y, y') = \frac{\partial(u, v)}{\partial(y, y')} = \det \begin{pmatrix} 1 & \psi'(y) \\ 1 & \psi'(y') \end{pmatrix} = \psi'(y') - \psi'(y) = p(y'|y'|^{p-2} - y|y|^{p-2}) \quad (2-2)$$

and satisfies $|J^{-1}(y, y')| \geq p|y|^{p-1} - |y'|^{p-1}$, with equality if and only if $yy' \geq 0$. Thus

$$(\mathcal{E}_p(f)\mathcal{E}_p(g))(x, t) = 2 \int e^{ixu} e^{itv} f(y) g(y') w(y)^{\frac{1}{2}} w(y')^{\frac{1}{2}} J(u, v) du dv, \quad (2-3)$$

where the integral is taken over the region $\{(u, v): v \geq 2\psi(u/2)\}$. Note that this implies

$$(f\sigma_p * g\sigma_p)(u, v) = 2f(y)g(y')w(y)^{\frac{1}{2}}w(y')^{\frac{1}{2}}J(u, v) \quad (2-4)$$

for every (u, v) satisfying $v > 2\psi(u/2)$, where (y, y') is related to (u, v) via the change of variables described above.

By symmetry, we can and will restrict attention to $|y'| \leq |y|$. Taking the L^3 -norm of (2-3), invoking the Hausdorff–Young inequality, and then changing variables back to (y, y') ,

$$\begin{aligned} \|\mathcal{E}_p(f)\mathcal{E}_p(g)\|_{L^3(\mathbb{R}^2)} &\lesssim \|f(y)g(y')w(y)^{\frac{1}{2}}w(y')^{\frac{1}{2}}J(u, v)\|_{L_{u,v}^{3/2}(\mathbb{R}^{1+1})} \\ &= \|f(y)g(y')w(y)^{\frac{1}{2}}w(y')^{\frac{1}{2}}|J(y, y')|^{\frac{1}{3}}\|_{L_{y,y'}^{3/2}(\mathbb{R}^{1+1})}. \end{aligned}$$

If $2^k \leq |y| < 2^{k+1}$, $2^{k'} \leq |y'| < 2^{k'+1}$, and $k \geq k' + 2$, then

$$\frac{|yy'|^{\frac{p-2}{4}}}{p^{\frac{1}{2}}||y|^{p-1} - |y'|^{p-1}|^{\frac{1}{2}}} \lesssim \frac{2^{(k+k')\frac{p-2}{4}}}{2^{k\frac{p-1}{2}}(1-2^{-(k-k'-1)(p-1)})^{\frac{1}{2}}} \lesssim 2^{(k'-k)\frac{p}{4} - \frac{k'}{2}}. \quad (2-5)$$

It follows that

$$\begin{aligned} \|\mathcal{E}_p(f)\mathcal{E}_p(g)\|_{L^3}^{\frac{3}{2}} &\lesssim \int_{\mathbb{R}^2} |f(y)g(y')|^{\frac{3}{2}} w(y)^{\frac{3}{4}} w(y')^{\frac{3}{4}} |J(y, y')|^{\frac{1}{2}} dy dy' \\ &\leq \int_{\mathbb{R}^2} |f(y)g(y')|^{\frac{3}{2}} \frac{|yy'|^{\frac{p-2}{4}}}{p^{\frac{1}{2}}||y|^{p-1} - |y'|^{p-1}|^{\frac{1}{2}}} dy dy' \\ &\lesssim 2^{(k'-k)\frac{p}{4} - \frac{k'}{2}} 2^{\frac{k}{4}} 2^{\frac{k'}{4}} \|f\|_{L^2}^{\frac{3}{2}} \|g\|_{L^2}^{\frac{3}{2}} \\ &= 2^{-|k-k'|\frac{p-1}{4}} \|f\|_{L^2}^{\frac{3}{2}} \|g\|_{L^2}^{\frac{3}{2}}. \end{aligned} \quad (2-6)$$

If $k \in \{k', k' + 1\}$, then we can simply use the estimate $\|\mathcal{E}_p(f)\mathcal{E}_p(g)\|_{L^3} \lesssim \|f\|_{L^2} \|g\|_{L^2}$. \square

Corollary 2.2. *Let $p > 1$ and $k, k' \in \mathbb{Z}$ be such that $k' \leq k$. Then*

$$\|\mathcal{E}_p(f)\mathcal{E}_p(g)\|_{L^3(\mathbb{R}^2)} \lesssim_p 2^{-|k-k'|\frac{p-1}{6}} \|f\|_{L^2(\mathbb{R})} \|g\|_{L^2(\mathbb{R})} \quad (2-7)$$

for every $f, g \in L^2(\mathbb{R})$ satisfying $\text{supp } f \subseteq \{|y| \geq 2^k\}$ and $\text{supp } g \subseteq \{|y'| \leq 2^{k'}\}$.

Proof. Write $f = \sum_{j \geq k} f_j$ and $g = \sum_{j' < k'} g_{j'}$, where $f_j := f \mathbb{1}_{I_j^\bullet}$ and $g_{j'} := g \mathbb{1}_{I_{j'}^\bullet}$. Then

$$\begin{aligned} \|\mathcal{E}_p(f)\mathcal{E}_p(g)\|_{L^3(\mathbb{R}^2)} &\leq \sum_{j \geq k, j' < k'} \|\mathcal{E}_p(f_j)\mathcal{E}_p(g_{j'})\|_{L^3} \lesssim \sum_{j \geq k, j' < k'} 2^{-|j-j'|\frac{p-1}{6}} \|f_j\|_{L^2} \|g_{j'}\|_{L^2} \\ &\leq \left(\sum_{j \geq k, j' < k'} 2^{-|j-j'|\frac{p-1}{3}} \right)^{\frac{1}{2}} \left(\sum_{j \geq k, j' < k'} \|f_j\|_{L^2}^2 \|g_{j'}\|_{L^2}^2 \right)^{\frac{1}{2}} \\ &\simeq \left(\sum_{j \geq k} 2^{-|j-k'|\frac{p-1}{3}} \right)^{\frac{1}{2}} \|f\|_{L^2} \|g\|_{L^2} \\ &\simeq 2^{-|k-k'|\frac{p-1}{6}} \|f\|_{L^2} \|g\|_{L^2}, \end{aligned}$$

where we used the triangle inequality, Proposition 2.1, the Cauchy–Schwarz inequality, L^2 -orthogonality, and the fact that a geometric series is comparable to its largest term. \square

When studying concentration at points different from the origin, it will be useful to consider dyadic decompositions of the real line with arbitrary centers. By reflection and scaling, it suffices to consider decompositions centered at 1. Define the dyadic regions

$$\mathcal{I}_k := \{2^k \leq y - 1 < 2^{k+1}\} \quad \text{and} \quad \mathcal{I}_k^\bullet := \{2^k \leq |y - 1| < 2^{k+1}\} \quad (k \in \mathbb{Z})$$

so that $\mathcal{I}_k = 1 + I_k$ and $\mathcal{I}_k^\bullet = 1 + I_k^\bullet$. The following analogue of Proposition 2.1 holds.

Proposition 2.3. *Let $p > 1$ and $k, k' \in \mathbb{Z}$. Let $\beta = \min\{\frac{1}{6}, \frac{p-1}{6}\}$. Then*

$$\|\mathcal{E}_p(f)\mathcal{E}_p(g)\|_{L^3(\mathbb{R}^2)} \lesssim_p 2^{-\beta|k-k'|} \|f\|_{L^2(\mathbb{R})} \|g\|_{L^2(\mathbb{R})} \quad (2-8)$$

for every $f, g \in L^2(\mathbb{R})$ satisfying $\text{supp } f \subseteq \mathcal{I}_k^\bullet$ and $\text{supp } g \subseteq \mathcal{I}_{k'}^\bullet$.

Before embarking on the proof, let us take a closer look at the factor $|yy'|^{(p-2)/4} |J(y, y')|^{1/2}$ that appears after applying the Hausdorff–Young inequality in (2-6). We have already seen that

$$|J^{-1}(y, y')| = p|y|y|^{p-2} - y'y|y'|^{p-2}|. \quad (2-9)$$

In (2-5) we observed that, if y, y' are *separated* (say, $|y'| \leq \frac{1}{2}|y|$), then

$$\frac{|yy'|^{\frac{p-2}{4}}}{|y|y|^{p-2} - y'y|y'|^{p-2}|^{\frac{1}{2}}} \lesssim \frac{|yy'|^{\frac{p-2}{4}}}{|y|^{\frac{p-1}{2}}} = |y|^{-\frac{p}{4}} |y'|^{\frac{p-2}{4}}. \quad (2-10)$$

In order to obtain a useful bound in the case when both y, y' are close to 1, invoke the mean value theorem and write

$$|y|^{p-1} - |y'|^{p-1} = (p-1)s^{p-2}(|y| - |y'|)$$

for some $s \in [|y'|, |y|]$. Then, for $0 \leq y' \leq y$, we have

$$|y|y|^{p-2} - y'y|y'|^{p-2}| = |y^{p-1} - y'^{p-1}| \gtrsim \begin{cases} |y - y'|y^{p-2} & \text{if } p \in (1, 2], \\ |y - y'|y'^{p-2} & \text{if } p \in [2, \infty). \end{cases}$$

Thus the following estimate holds for every $\frac{1}{2} \leq y, y' \leq \frac{3}{2}$:

$$\frac{|yy'|^{\frac{p-2}{4}}}{|y|y|^{p-2} - y'y|y'|^{p-2}|^{\frac{1}{2}}} \lesssim |y - y'|^{-\frac{1}{2}}. \quad (2-11)$$

Proof of Proposition 2.3. Without loss of generality, assume $|k - k'| \geq 2$. We start by considering the situation when 0 is an endpoint of $\mathcal{I}_{k'}^\bullet$, i.e., $k' \in \{-1, 0\}$. Let $k' = -1$, so that $\mathcal{I}_{k'}^\bullet = (0, \frac{1}{2}] \cup [\frac{3}{2}, 2)$, split $g = g_\ell + g_r$, with $g_\ell := g\mathbb{1}_{(0, \frac{1}{2}]}$ and $g_r := g\mathbb{1}_{[\frac{3}{2}, 2)}$, and take the dyadic decomposition

$$g_\ell = \sum_{j \geq 1} g_j, \quad \text{with } g_j := g\mathbb{1}_{(2^{-(j+1)}, 2^{-j}]}.$$

If $k \leq -3$, then (2-10) implies

$$\begin{aligned} \|\mathcal{E}_p(f)\mathcal{E}_p(g_\ell)\|_{L^3} &\lesssim \sum_{j \geq 1} \left(\int_{\mathbb{R}^2} |f(y)g_j(y')|^{\frac{3}{2}} \frac{|yy'|^{\frac{p-2}{4}}}{|y|^{p-1} - |y'|^{p-1}|^{\frac{1}{2}}} dy dy' \right)^{\frac{2}{3}} \\ &\lesssim \sum_{j \geq 1} \left(2^{-j\frac{p-2}{4}} \int_{\mathbb{R}^2} |f(y)g_j(y')|^{\frac{3}{2}} dy dy' \right)^{\frac{2}{3}} \lesssim \sum_{j \geq 1} \left(2^{-j\frac{p-2}{4}} 2^{\frac{k}{4}} 2^{-\frac{j}{4}} \|f\|_{L^2}^{\frac{3}{2}} \|g_j\|_{L^2}^{\frac{3}{2}} \right)^{\frac{2}{3}} \\ &= 2^{\frac{k}{6}} \|f\|_{L^2} \sum_{j \geq 1} 2^{-j\frac{p-1}{6}} \|g_j\|_{L^2} \lesssim 2^{\frac{k}{6}} \|f\|_{L^2} \|g_\ell\|_{L^2} \lesssim 2^{-\frac{|k-k'|}{6}} \|f\|_{L^2} \|g\|_{L^2}. \end{aligned}$$

If $k \geq 1$, then [Corollary 2.2](#) applies, and directly yields

$$\|\mathcal{E}_p(f)\mathcal{E}_p(g_\ell)\|_{L^3} \lesssim 2^{-|k-k'| \frac{p-1}{6}} \|f\|_{L^2} \|g\|_{L^2}.$$

A similar analysis applies to g_r . Setting $\beta := \min\{\frac{1}{6}, \frac{p-1}{6}\}$, we conclude that, if $k' = -1$ and $|k-k'| \geq 2$, then

$$\|\mathcal{E}_p(f)\mathcal{E}_p(g)\|_{L^3} \lesssim 2^{-\beta|k-k'|} \|f\|_{L^2} \|g\|_{L^2}.$$

The case $k' = 0$ admits a similar treatment. If $k, k' \leq -2$ and $k - k' \geq 2$, then [\(2-11\)](#) implies

$$\|\mathcal{E}_p(f)\mathcal{E}_p(g)\|_{L^3} \lesssim \frac{2^{\frac{k}{6}} 2^{\frac{k'}{6}}}{2^{\frac{k}{3}}} \|f\|_{L^2} \|g\|_{L^2} = 2^{-\frac{|k-k'|}{6}} \|f\|_{L^2} \|g\|_{L^2}.$$

Finally, the remaining cases can be handled in a similar way by [Corollary 2.2](#). \square

Corollary 2.4. *Let $p > 1$ and $k, k' \in \mathbb{Z}$ be such that $k' \leq k$. Let $\beta = \min\{\frac{1}{6}, \frac{p-1}{6}\}$. Then*

$$\|\mathcal{E}_p(f)\mathcal{E}_p(g)\|_{L^3(\mathbb{R}^2)} \lesssim_p 2^{-\beta|k-k'|} \|f\|_{L^2(\mathbb{R})} \|g\|_{L^2(\mathbb{R})} \quad (2-12)$$

for every $f, g \in L^2(\mathbb{R})$ satisfying $\text{supp } f \subseteq \{|y-1| \geq 2^k\}$ and $\text{supp } g \subseteq \{|y'-1| \leq 2^{k'}\}$.

We finish this subsection by taking yet another look at the Jacobian factor [\(2-9\)](#). This will be useful in [Section 2B](#) below. Let $p \geq 2$. If $yy' \leq 0$, then $|J^{-1}(y, y')| = p(|y|^{p-1} + |y'|^{p-1})$, in which case

$$\frac{|yy'|^{\frac{p-2}{4}}}{(|y|^{p-1} + |y'|^{p-1})^{\frac{1}{2}}} \lesssim (|y| + |y'|)^{-\frac{1}{2}} = |y - y'|^{-\frac{1}{2}}$$

uniformly in y, y' . To handle the complementary case $yy' > 0$, note that, if $p \geq 2$ and $0 \leq a \leq b$, then

$$b^{p-1} - a^{p-1} \simeq (b-a)b^{p-2}. \quad (2-13)$$

It follows that, if $p \geq 2$ and $yy' > 0$, then

$$|J^{-1}(y, y')| = p||y|^{p-1} - |y'|^{p-1}| \simeq |y - y'| \max\{|y|, |y'|\}^{p-2},$$

and so if additionally $|y| \geq |y'|$, then

$$\frac{|yy'|^{\frac{p-2}{4}}}{||y|^{p-1} - |y'|^{p-1}|^{\frac{1}{2}}} \lesssim \frac{|yy'|^{\frac{p-2}{4}}}{|y|^{\frac{p-2}{2}} |y - y'|^{\frac{1}{2}}} \leq |y - y'|^{-\frac{1}{2}}.$$

Therefore the estimate

$$\|\mathcal{E}_p(f)\mathcal{E}_p(g)\|_{L^3(\mathbb{R}^2)}^{\frac{3}{2}} \lesssim \int_{\mathbb{R}^2} \frac{|f(y)g(y')|^{\frac{3}{2}}}{|y - y'|^{\frac{1}{2}}} dy dy' \quad (2-14)$$

holds as long as $p \geq 2$. We cannot hope for such a bound if $1 < p < 2$ since [\(2-13\)](#) fails in that case. However, if $|y| \simeq |y'|$, then one can check in a similar way that the estimate

$$\|\mathcal{E}_p(f_k)\mathcal{E}_p(g_k)\|_{L^3(\mathbb{R}^2)}^{\frac{3}{2}} \lesssim \int_{\mathbb{R}^2} \frac{|f_k(y)g_k(y')|^{\frac{3}{2}}}{|y - y'|^{\frac{1}{2}}} dy dy' \quad (2-15)$$

holds for any $p > 1$ and functions f_k, g_k which are both supported on I_k^\bullet .

2B. Cap bounds. An inspection of the proof of [Proposition 2.1](#) reveals that if $\text{supp } f \subseteq I_k^\bullet$ and $\text{supp } g \subseteq I_{k'}^\bullet$, for some $k, k' \in \mathbb{Z}$ satisfying $k - k' \geq 2$, then

$$\begin{aligned} \|\mathcal{E}_p(f)\mathcal{E}_p(g)\|_{L^3(\mathbb{R}^2)} &\lesssim 2^{-|k-k'|\frac{p-1}{6}} \left(|I_k|^{-\frac{1}{4}} \int_{I_k^\bullet} |f|^{\frac{3}{2}} \right)^{\frac{2}{3}} \left(|I_{k'}|^{-\frac{1}{4}} \int_{I_{k'}^\bullet} |g|^{\frac{3}{2}} \right)^{\frac{2}{3}} \\ &\lesssim 2^{-|k-k'|\frac{p-1}{6}} \Lambda(f)^{\frac{2}{9}} \Lambda(g)^{\frac{2}{9}} \|f\|_{L^2(\mathbb{R})}^{\frac{2}{3}} \|g\|_{L^2(\mathbb{R})}^{\frac{2}{3}}, \end{aligned} \quad (2-16)$$

where the quantity $\Lambda(f)$ is defined via

$$\Lambda(f) := \sup_{k \in \mathbb{Z}} |I_k|^{-\frac{1}{4}} \int_{I_k^\bullet} |f|^{\frac{3}{2}}. \quad (2-17)$$

The purpose of this subsection is to develop on this observation. Given $f \in L^2(\mathbb{R})$, write $f = \sum_{k \in \mathbb{Z}} f_k$, with $f_k := f \mathbb{1}_{I_k^\bullet}$. Our first result is the following.

Proposition 2.5. *Let $p > 1$. Then the following estimates hold for every $f \in L^2(\mathbb{R})$:*

$$\|\mathcal{E}_p(f)\|_{L^6(\mathbb{R}^2)}^3 \lesssim_p \sum_{k \in \mathbb{Z}} \|f_k\|_{L^2(\mathbb{R})}^3, \quad (2-18)$$

$$\|\mathcal{E}_p(f)\|_{L^6(\mathbb{R}^2)}^3 \lesssim_p \sum_{k \in \mathbb{Z}} \|\mathcal{E}_p(f_k)\|_{L^6(\mathbb{R}^2)}^3 + \Lambda(f)^{\frac{4}{9}} \left(\sum_{k \in \mathbb{Z}} \|f_k\|_{L^2(\mathbb{R})}^3 \right)^{\frac{1}{3}} \|f\|_{L^2(\mathbb{R})}^{\frac{4}{3}}. \quad (2-19)$$

Proof. By the triangle inequality,

$$\|\mathcal{E}_p(f)\|_{L^6}^3 \leq \sum_{(i,j,k) \in \mathbb{Z}^3} \|\mathcal{E}_p(f_i)\mathcal{E}_p(f_j)\mathcal{E}_p(f_k)\|_{L^2}.$$

For each triple (i, j, k) in the previous sum, we lose no generality in assuming that

$$|j - k| = \max\{|i' - j'| : i', j' \in \{i, j, k\}\}. \quad (2-20)$$

Hölder's inequality and [Proposition 2.1](#) then imply

$$\|\mathcal{E}_p(f_i)\mathcal{E}_p(f_j)\mathcal{E}_p(f_k)\|_{L^2} \lesssim 2^{-|j-k|\frac{p-1}{6}} \|f_i\|_{L^2} \|f_j\|_{L^2} \|f_k\|_{L^2}.$$

By the maximality of $|j - k|$, we have $|j - k| \geq \frac{1}{3}|i - j| + \frac{1}{3}|j - k| + \frac{1}{3}|k - i|$, and hence

$$\|\mathcal{E}_p(f)\|_{L^6}^3 \lesssim \sum_{(i,j,k) \in \mathbb{Z}^3} 2^{-|i-j|\frac{p-1}{18}} 2^{-|j-k|\frac{p-1}{18}} 2^{-|k-i|\frac{p-1}{18}} \|f_i\|_{L^2} \|f_j\|_{L^2} \|f_k\|_{L^2}.$$

A final application of Hölder's inequality yields [\(2-18\)](#). Estimate [\(2-19\)](#) follows from similar considerations which we now detail. Let $S := \{(i, j, k) \in \mathbb{Z}^3 : \max\{|i - j|, |j - k|, |k - i|\} \leq 1\}$ and $S^C := \mathbb{Z}^3 \setminus S$. Split the sum into diagonal and off-diagonal contributions,

$$\|\mathcal{E}_p(f)\|_{L^6}^3 \leq \left\| \sum_{(i,j,k) \in S} \mathcal{E}_p(f_i)\mathcal{E}_p(f_j)\mathcal{E}_p(f_k) \right\|_{L^2} + \left\| \sum_{(i,j,k) \in S^C} \mathcal{E}_p(f_i)\mathcal{E}_p(f_j)\mathcal{E}_p(f_k) \right\|_{L^2},$$

and analyze the two terms separately. For the diagonal term, note that

$$\begin{aligned} & \left\| \sum_{(i,j,k) \in S} \mathcal{E}_p(f_i) \mathcal{E}_p(f_j) \mathcal{E}_p(f_k) \right\|_{L^2} \\ & \leq \sum_{k \in \mathbb{Z}} (3 \|\mathcal{E}_p(f_k) \mathcal{E}_p(f_k) \mathcal{E}_p(f_{k+1})\|_{L^2} + 3 \|\mathcal{E}_p(f_{k-1}) \mathcal{E}_p(f_k) \mathcal{E}_p(f_k)\|_{L^2} + \|\mathcal{E}_p(f_k) \mathcal{E}_p(f_k) \mathcal{E}_p(f_k)\|_{L^2}) \\ & \leq \sum_{k \in \mathbb{Z}} (3 \|\mathcal{E}_p(f_k)\|_{L^6}^2 \|\mathcal{E}_p(f_{k+1})\|_{L^6} + 3 \|\mathcal{E}_p(f_{k-1})\|_{L^6} \|\mathcal{E}_p(f_k)\|_{L^6}^2 + \|\mathcal{E}_p(f_k)\|_{L^6}^3) \lesssim \sum_{k \in \mathbb{Z}} \|\mathcal{E}_p(f_k)\|_{L^6}^3. \end{aligned}$$

To handle the off-diagonal term, note that estimate (2-16) implies

$$\begin{aligned} \left\| \sum_{(i,j,k) \in S^C} \mathcal{E}_p(f_i) \mathcal{E}_p(f_j) \mathcal{E}_p(f_k) \right\|_{L^2} & \lesssim \sum'_{(i,j,k): |j-k| \geq 2} \|f_i\|_{L^2} \|\mathcal{E}_p(f_j) \mathcal{E}_p(f_k)\|_{L^3} \\ & \lesssim \Lambda(f)^{\frac{4}{9}} \sum'_{(i,j,k): |j-k| \geq 2} 2^{-|j-k|\frac{p-1}{6}} \|f_i\|_{L^2} \|f_j\|_{L^2}^{\frac{2}{3}} \|f_k\|_{L^2}^{\frac{2}{3}}, \end{aligned}$$

where the sum Σ' is taken over triples $(i, j, k) \in S^C$ for which (j, k) satisfies the maximality assumption (2-20). It follows that

$$\begin{aligned} \left\| \sum_{(i,j,k) \in S^C} \mathcal{E}_p(f_i) \mathcal{E}_p(f_j) \mathcal{E}_p(f_k) \right\|_{L^2} & \lesssim \Lambda(f)^{\frac{4}{9}} \sum_{i,j,k} 2^{-(|i-j|+|j-k|+|k-i|)\frac{p-1}{18}} \|f_i\|_{L^2} \|f_j\|_{L^2}^{\frac{2}{3}} \|f_k\|_{L^2}^{\frac{2}{3}} \\ & \lesssim \Lambda(f)^{\frac{4}{9}} \left(\sum_{k \in \mathbb{Z}} \|f_k\|_{L^2}^3 \right)^{\frac{1}{3}} \left(\sum_{k \in \mathbb{Z}} \|f_k\|_{L^2}^2 \right)^{\frac{2}{3}}. \end{aligned}$$

This implies (2-19) at once, and concludes the proof of the proposition. \square

The following L^2 dyadic cap estimate is a direct consequence of (2-18).

Corollary 2.6. *Let $p > 1$. Then, for every $f \in L^2(\mathbb{R})$,*

$$\|\mathcal{E}_p(f)\|_{L^6(\mathbb{R}^2)}^3 \lesssim_p \left(\sup_{k \in \mathbb{Z}} \|f_k\|_{L^2(\mathbb{R})} \right) \|f\|_{L^2(\mathbb{R})}^2.$$

We now derive a cap bound similar to [Jiang et al. 2010, Lemma 1.2] and [Shao 2009, Lemma 1.2].

Proposition 2.7. *Let $p > 1$. Then the following estimate holds:*

$$\|\mathcal{E}_p(f)\|_{L^6(\mathbb{R}^2)}^3 \lesssim_p \left(\sup_{k \in \mathbb{Z}} \sup_{I \subseteq I_k^\bullet} |I|^{-\frac{1}{6}} \|f\|_{L^{3/2}(I)} \right)^{\frac{2}{3}} \|f\|_{L^2(\mathbb{R})}^{\frac{7}{3}} \quad (2-21)$$

for every $f \in L^2(\mathbb{R})$, where the inner supremum is taken over all subintervals $I \subseteq I_k^\bullet$.

Proof. We start by considering the case when $f = f_k (= f \mathbb{1}_{I_k^\bullet})$. From (2-15), we have

$$\|\mathcal{E}_p(f_k)\|_{L^6}^3 \lesssim \int_{\mathbb{R}^2} \frac{|f_k(y) f_k(y')|^{\frac{3}{2}}}{|y - y'|^{\frac{1}{2}}} dy dy'. \quad (2-22)$$

Arguing as in as in [Jiang et al. 2010; Shao 2009] we obtain, for every $q > 1$, that

$$\|\mathcal{E}_p(f_k)\|_{L^6} \lesssim \left(\sup_{I \subseteq I_k^\bullet} |I|^{\frac{1}{2} - \frac{1}{q}} \|f_k\|_{L^q(I)} \right)^{\frac{1}{3}} \|f_k\|_{L^2(\mathbb{R})}^{\frac{2}{3}}. \quad (2-23)$$

For the convenience of the reader, we provide the details. In light of (2-22), we may assume $f_k \geq 0$. Normalizing the supremum in (2-23) to equal 1, we may further assume that

$$\int_I f_k^q \leq |I|^{1 - \frac{q}{2}} \quad \text{for every subinterval } I \subseteq I_k^\bullet. \quad (2-24)$$

Denote the collection of dyadic intervals of length 2^j by $\mathcal{D}_j := \{2^j[k, k+1) : k \in \mathbb{Z}\}$, and set $\mathcal{D} := \bigcup_{j \in \mathbb{Z}} \mathcal{D}_j$. We perform a Whitney decomposition of $\mathbb{R}^2 \setminus \{(y, y) : y \in \mathbb{R}\}$ in the following manner; see for instance [Dodson et al. 2018, Lemma 10] and [Bégout and Vargas 2007, Proof of Theorem 1.2]. Given distinct $y, y' \in \mathbb{R}$, there exists a unique pair of maximal dyadic intervals I, I' satisfying

$$(y, y') \in I \times I', \quad |I| = |I'|, \quad \text{and} \quad \text{dist}(I, I') \geq 4|I|.$$

Let \mathfrak{I} denote the collection of all such pairs as $y \neq y'$ ranges over $\mathbb{R} \times \mathbb{R}$. Then

$$\sum_{(I, I') \in \mathfrak{I}} \mathbb{1}_I(y) \mathbb{1}_{I'}(y') = 1 \quad \text{for every } (y, y') \in \mathbb{R}^2 \text{ with } y \neq y',$$

and therefore

$$f_k(y) f_k(y') = \sum_{(I, I') \in \mathfrak{I}} f_{k,I}(y) f_{k,I'}(y') \quad \text{for a.e. } (y, y') \in \mathbb{R}^2,$$

where $f_{k,I} := f_k \mathbb{1}_I$. Clearly, if $(y, y') \in I \times I'$ and $(I, I') \in \mathfrak{I}$, then $|y - y'| \simeq |I|$. From this and (2-22), we may choose a slightly larger dyadic interval containing $I \cup I'$ but of length comparable to $|I|$ (still denoted by I), and it suffices to show that

$$\sum_{I \in \mathcal{D}} \frac{1}{|I|^{\frac{1}{2}}} \left(\int f_{k,I}^{\frac{3}{2}} \right)^2 \lesssim \int f_k^2.$$

We further decompose $f_{k,I}$ as

$$f_{k,I} = \sum_{n \in \mathbb{Z}} f_{k,I,n}, \quad \text{where } f_{k,I,n} := f_k \mathbb{1}_{\{y \in I : \frac{2^n}{|I|^{1/2}} \leq f_k(y) < \frac{2^{n+1}}{|I|^{1/2}}\}},$$

and note that it suffices to establish

$$\sum_{I \in \mathcal{D}} \frac{1}{|I|^{\frac{1}{2}}} \left(\int f_{k,I,n}^{\frac{3}{2}} \right)^2 \lesssim 2^{-|n|\varepsilon} \int f_k^2 \quad (2-25)$$

for some $\varepsilon > 0$ and every $n \in \mathbb{Z}$. By the Cauchy–Schwarz inequality,

$$\left(\int f_{k,I,n}^{\frac{3}{2}} \right)^2 \leq \left(\int f_{k,I,n}^2 \right) \left(\int f_{k,I,n} \right).$$

By construction of $f_{k,I,n}$, Chebyshev's inequality, and normalization (2-24),

$$\int f_{k,I,n} \leq \frac{2^{n+1}}{|I|^{\frac{1}{2}}} \left| \left\{ y \in I : f_k(y) \geq \frac{2^n}{|I|^{\frac{1}{2}}} \right\} \right| \leq \frac{2^{n+1}}{|I|^{\frac{1}{2}}} \frac{\int_I f_k^q}{2^{nq} |I|^{-\frac{q}{2}}} \lesssim 2^{-|n|(q-1)} |I|^{\frac{1}{2}} \quad (2-26)$$

for every $q > 1$ and $n \geq 0$. If $n < 0$, then the following simpler estimate suffices:

$$\int f_{k,I,n} \lesssim \frac{2^n}{|I|^{\frac{1}{2}}} |I| = 2^{-|n|} |I|^{\frac{1}{2}}. \quad (2-27)$$

Combining (2-26) and (2-27), we conclude

$$\sum_{I \in \mathcal{D}} \frac{1}{|I|^{\frac{1}{2}}} \left(\int f_{k,I,n}^{\frac{3}{2}} \right)^2 \lesssim 2^{-|n|\varepsilon} \sum_{I \in \mathcal{D}} \int f_{k,I,n}^2$$

for some $\varepsilon > 0$, from which we get the desired (2-25) by noting that

$$\sum_{I \in \mathcal{D}} \int f_{k,I,n}^2 = \sum_{j \in \mathbb{Z}} \sum_{I \in \mathcal{D}_j} \int f_k^2 \mathbb{1}_{\{f_k \simeq 2^{n-j/2}\}} = \int_{\mathbb{R}} \left(\sum_{\substack{j \in \mathbb{Z}: \\ f_k(y) \simeq 2^{n-j/2}}} f_k^2(y) \right) dy \lesssim \int f_k^2.$$

This concludes the verification of (2-23). Recalling inequality (2-19), and specializing (2-23) to $q = \frac{3}{2}$, yields

$$\begin{aligned} \|\mathcal{E}_p(f)\|_{L^6}^3 &\lesssim \left(\sup_{k, I \subseteq I_k^\bullet} |I|^{-\frac{1}{6}} \|f_k\|_{L^{3/2}(I)} \right) \sum_{k \in \mathbb{Z}} \|f_k\|_{L^2}^2 + \left(\sup_{k \in \mathbb{Z}} |I_k|^{-\frac{1}{6}} \|f_k\|_{L^{3/2}} \right)^{\frac{2}{3}} \|f\|_{L^2}^{\frac{7}{3}} \\ &\lesssim \left(\sup_{k \in \mathbb{Z}} \sup_{I \subseteq I_k^\bullet} |I|^{-\frac{1}{6}} \|f_k\|_{L^{3/2}(I)} \right)^{\frac{2}{3}} \|f\|_{L^2}^{\frac{7}{3}}, \end{aligned}$$

where the last line follows from Hölder's inequality. \square

In the next section, it will be useful to have the L^1 version of (2-21) at our disposal, and this is the content of the following result.

Proposition 2.8. *Let $p > 1$. Then there exist $\gamma \in (0, 1)$ such that*

$$\|\mathcal{E}_p(f)\|_{L^6(\mathbb{R}^2)} \lesssim_{p,\gamma} \left(\sup_{k \in \mathbb{Z}} \sup_{I \subseteq I_k^\bullet} |I|^{-\frac{1}{2}} \|f\|_{L^1(I)} \right)^\gamma \|f\|_{L^2(\mathbb{R})}^{1-\gamma} \quad (2-28)$$

for every $f \in L^2(\mathbb{R})$, where the inner supremum is taken over all subintervals $I \subseteq I_k^\bullet$.

The proof below yields $\gamma = \frac{2}{45}$ and is inspired by [Christ and Shao 2012a, Proposition 2.9].

Proof of Proposition 2.8. Set $\delta := \|\mathcal{E}_p(f)\|_{L^6} \|f\|_{L^2}^{-1}$. From (2-21) we have

$$\sup_{k \in \mathbb{Z}} \sup_{I \subseteq I_k^\bullet} |I|^{-\frac{1}{6}} \|f\|_{L^{3/2}(I)} \gtrsim \delta^{\frac{9}{2}} \|f\|_{L^2(\mathbb{R})}.$$

Then there exist $k \in \mathbb{Z}$ and an interval $I \subseteq I_k^\bullet$ such that

$$\int_I |f|^{\frac{3}{2}} \geq c_0 \delta^{\frac{27}{4}} |I|^{\frac{1}{4}} \|f\|_{L^2(\mathbb{R})}^{\frac{3}{2}}$$

for a universal constant c_0 (independent of f, δ). Given $R \geq 1$, define the set $E := \{y \in I : |f(y)| \leq R\}$. Set $g := f \mathbb{1}_E$ and $h := f - g$. Then g and h have disjoint supports, and $\|g\|_{L^\infty} \leq R$. Since $|h(y)| \geq R$ for almost every $y \in I$ for which $h(y) \neq 0$, we have

$$\int_I |h|^{\frac{3}{2}} \leq R^{-\frac{1}{2}} \int_I |h|^2 \leq R^{-\frac{1}{2}} \|f\|_{L^2(\mathbb{R})}^2.$$

Choose R satisfying $R^{-\frac{1}{2}} = \frac{1}{2}c_0\delta^{\frac{27}{4}}|I|^{\frac{1}{4}}\|f\|_{L^2(\mathbb{R})}^{-\frac{1}{2}}$. Then

$$\int_I |g|^{\frac{3}{2}} = \int_I |f|^{\frac{3}{2}} - \int_I |h|^{\frac{3}{2}} \geq \frac{c_0}{2}\delta^{\frac{27}{4}}|I|^{\frac{1}{4}}\|f\|_{L^2(\mathbb{R})}^{\frac{3}{2}}.$$

Since g is supported on I , Hölder's inequality implies

$$\|g\|_{L^2} \geq |I|^{-\frac{1}{6}}\|g\|_{L^{3/2}} \geq c_1\delta^{\frac{9}{2}}\|f\|_{L^2}, \quad (2-29)$$

where c_1 is universal. Since $\|g\|_{L^\infty} \leq R$, we have (by the definition of R) that

$$|g(y)| \leq c_2\delta^{-\frac{27}{2}}|I|^{-\frac{1}{2}}\|f\|_{L^2(\mathbb{R})}\mathbb{1}_I(y) \quad \text{for almost every } y \in \mathbb{R},$$

where c_2 is universal. Together with (2-29), this implies the lower bound

$$\int_I |g| \geq \int_I |g| \frac{|g|}{c_2\delta^{-\frac{27}{2}}|I|^{-\frac{1}{2}}\|f\|_{L^2}} = c_2^{-1}\delta^{\frac{27}{2}}|I|^{\frac{1}{2}}\frac{\|g\|_{L^2}^2}{\|f\|_{L^2}} \geq c_3\delta^{\frac{45}{2}}|I|^{\frac{1}{2}}\|f\|_{L^2},$$

where c_3 is universal. Since $|g| \leq |f|$, it follows that

$$c_3\delta^{\frac{45}{2}}|I|^{\frac{1}{2}}\|f\|_{L^2(\mathbb{R})} \leq \|g\|_{L^1(I)} \leq \|f\|_{L^1(I)}.$$

Recalling the definition of δ , we obtain (2-28) with $\gamma = \frac{2}{45}$. □

3. Existence versus concentration

This section is devoted to the proof of [Theorem 1.3](#). Start by observing the scale invariance of (1-10), or equivalently that of (1-9). Indeed, if $f_\lambda(y) := f(\lambda y)$, then $\|f_\lambda\|_{L^2(\mathbb{R})} = \lambda^{-1/2}\|f\|_{L^2(\mathbb{R})}$. On the other hand, $\mathcal{E}_p(f_\lambda)(x, t) = \lambda^{-(p+4)/6}\mathcal{E}_p(f)(x/\lambda, t/\lambda^p)$, and so

$$\|\mathcal{E}_p(f_\lambda)\|_{L^6(\mathbb{R}^2)} = \lambda^{-\frac{p+4}{6} + \frac{p+1}{6}}\|\mathcal{E}_p(f)\|_{L^6(\mathbb{R}^2)} = \lambda^{-\frac{1}{2}}\|\mathcal{E}_p(f)\|_{L^6(\mathbb{R}^2)}.$$

In particular, given any sequence $\{a_n\} \subset \mathbb{R} \setminus \{0\}$, if $\{f_n\}$ is an L^2 -normalized extremizing sequence for (1-9), then so is $\{|a_n|^{1/2}f_n(a_n \cdot)\}$.

We come to the first main result of this section.

Proposition 3.1. *Let $\{f_n\} \subset L^2(\mathbb{R})$ be an L^2 -normalized extremizing sequence of nonnegative functions for (1-9). Then there exist a subsequence $\{f_{n_k}\}$ and a sequence $\{a_k\} \subset \mathbb{R} \setminus \{0\}$ such that the rescaled sequence $\{g_k\}$, $g_k := |a_k|^{1/2}f_{n_k}(a_k \cdot)$ satisfies one of the following conditions:*

- (i) *There exists $g \in L^2(\mathbb{R})$ such that $g_k \rightarrow g$ in $L^2(\mathbb{R})$ as $k \rightarrow \infty$.*
- (ii) *$\{g_k\}$ concentrates at $y_0 = 1$.*

Theorem 1.3 follows at once from Proposition 3.1 and the following result.

Lemma 3.2. *Let $p > 1$. Given $y_0 \in \mathbb{R} \setminus \{0\}$, let $\{f_n\} \subset L^2(\mathbb{R})$ be a sequence concentrating at y_0 . Then*

$$\limsup_{n \rightarrow \infty} \frac{\|f_n \sigma_p * f_n \sigma_p * f_n \sigma_p\|_{L^2(\mathbb{R}^2)}^2}{\|f_n\|_{L^2(\mathbb{R})}^6} \leq \frac{2\pi}{\sqrt{3}p(p-1)}. \quad (3-1)$$

If we set $f_n(y) = e^{-n(|y|^p - |y_0|^p - p y_0 |y_0|^{p-2} (y - y_0))} |y|^{(p-2)/6}$, then the sequence $\{f_n\|f_n\|_{L^2}^{-1}\}$ concentrates at y_0 , and equality holds in (3-1).

Convolution of singular measures is treated in much greater generality in the companion paper [Oliveira e Silva and Quilodrán 2019]. Lemma 3.2 is almost contained in [Oliveira e Silva and Quilodrán 2018; 2019], and we just indicate the necessary changes.

Proof sketch of Lemma 3.2. Once the boundary value for $|\cdot|^{(p-2)/6} \sigma_p * |\cdot|^{(p-2)/6} \sigma_p * |\cdot|^{(p-2)/6} \sigma_p$ given in (4-3) below is known to equal the right-hand side of (3-1), the proof for $p \geq 2$ follows the exact same lines as that of [Oliveira e Silva and Quilodrán 2018, Lemmas 4.1 and 4.2]. We omit the details.

If $1 < p < 2$, then the function $|\cdot|^{(p-2)/6}$ fails to be continuous at the origin, and an additional argument is needed. We show how to reduce matters to the analysis of projection measure. Let $\{f_n\} \subset L^2(\mathbb{R})$ concentrate at $y_0 \neq 0$. Then

$$\limsup_{n \rightarrow \infty} \frac{\|f_n \sigma_p * f_n \sigma_p * f_n \sigma_p\|_{L^2}^2}{\|f_n\|_{L^2}^6} = |y_0|^{p-2} \limsup_{n \rightarrow \infty} \frac{\|f_n \nu_p * f_n \nu_p * f_n \nu_p\|_{L^2}^2}{\|f_n\|_{L^2}^6}, \quad (3-2)$$

where ν_p denotes the projection measure $d\nu_p = \delta(s - |y|^p) dy ds$. To verify (3-2), consider the interval $J := [y_0/2, 3y_0/2]$. Then

$$\begin{aligned} \limsup_{n \rightarrow \infty} \frac{\|f_n \sigma_p * f_n \sigma_p * f_n \sigma_p\|_{L^2}^2}{\|f_n\|_{L^2}^6} &= \limsup_{n \rightarrow \infty} \frac{\|f_n \mathbb{1}_J \sigma_p * f_n \mathbb{1}_J \sigma_p * f_n \mathbb{1}_J \sigma_p\|_{L^2}^2}{\|f_n \mathbb{1}_J\|_{L^2}^6} \\ &= |y_0|^{p-2} \limsup_{n \rightarrow \infty} \frac{\|f_n \nu_p * f_n \nu_p * f_n \nu_p\|_{L^2}^2}{\|f_n\|_{L^2}^6}. \end{aligned}$$

Here, to justify the first equality, invoke the continuity of the operator \mathcal{E}_p , and the fact that the sequence $\{f_n\}$ concentrates at y_0 . For the second equality, additionally note that

$$\frac{\|f_n \mathbb{1}_J |\cdot|^{\frac{p-2}{6}} - f_n \mathbb{1}_J |y_0|^{\frac{p-2}{6}}\|_{L^2}}{\|f_n \mathbb{1}_J\|_{L^2}} \rightarrow 0 \quad \text{as } n \rightarrow \infty.$$

From [Oliveira e Silva and Quilodrán 2019, Proposition 2.1], the measure $\nu_p * \nu_p * \nu_p$ defines a continuous function in the interior of its support, with continuous extension to the boundary except at $(0, 0)$. Moreover, for any $y_0 \neq 0$,

$$(\nu_p * \nu_p * \nu_p)(3y_0, 3|y_0|^p) = \frac{2\pi}{\sqrt{3}p(p-1)|y_0|^{p-2}}.$$

The result now follows as in [Oliveira e Silva and Quilodrán 2018, Lemmas 4.1 and 4.2]. \square

The proof of [Proposition 3.1](#) relies on the bilinear extension estimates and cap bounds from [Section 2](#), together with a suitable variant of Lions' concentration-compactness lemma, which is formulated in the [Appendix](#) as [Proposition A.1](#). This has two important consequences for the present context, the first of which is the following.

Proposition 3.3. *Let $\{f_n\} \subset L^2(\mathbb{R})$ be an L^2 -normalized extremizing sequence for (1-9). Let $\{r_n\}$ be a sequence of nonnegative numbers satisfying $r_n \rightarrow 0$ as $n \rightarrow \infty$ and*

$$\inf_{n \in \mathbb{N}} \int_{1-r_n}^{1+r_n} |f_n(y)|^2 dy > 0.$$

Then the sequence $\{f_n\}$ concentrates at $y_0 = 1$.

Proof. Consider the intervals $J_n := [1 - r_n, 1 + r_n]$, $n \in \mathbb{N}$, and define the pseudometric

$$\varrho: \mathbb{R} \setminus \{1\} \times \mathbb{R} \setminus \{1\} \rightarrow [0, \infty), \quad \varrho(x, y) := |k - k'|, \quad (3-3)$$

where k, k' are such that $|x - 1| \in [2^k, 2^{k+1})$ and $|y - 1| \in [2^{k'}, 2^{k'+1})$. Let R be an integer. Then the ball centered at $x \neq 1$ of radius R defined by ϱ is given by

$$B(x, R) = \{y \in \mathbb{R} \setminus \{1\}: 2^{k-R} \leq |y - 1| < 2^{k+R+1}\}.$$

Let $\{f_n\}$ be as in the statement of the proposition. Apply [Proposition A.1](#) to the sequence $\{|f_n|^2\}$ with $X = \mathbb{R}$ equipped with Lebesgue measure, $\bar{x} = 1$, the function ϱ defined as in (3-3), and $\lambda = 1$. Passing to a subsequence, also denoted by $\{|f_n|^2\}$, one of three cases arises.

Case 1: The sequence $\{|f_n|^2\}$ satisfies *compactness*. In this case, there exists $\{x_n\} \subset \mathbb{R} \setminus \{1\}$ with the property that for any $\varepsilon > 0$ there exists $R < \infty$ such that, for every $n \geq 1$,

$$\int_{B(x_n, R)} |f_n|^2 \geq 1 - \varepsilon. \quad (3-4)$$

Suppose that $\limsup_{n \rightarrow \infty} |x_n - 1| > 0$. Then, possibly after extraction of a subsequence, $\{x_n\}$ is eventually far from 1; i.e., there exist $N_0 \in \mathbb{N}$, $\ell^* \in \mathbb{Z}$ such that $|x_n - 1| > 2^{\ell^*}$ for every $n \geq N_0$. Let $\varepsilon := \frac{1}{2} \inf_n \|f_n\|_{L^2(J_n)}^2 > 0$, and choose an integer R such that (3-4) holds. Now,

$$B(x_n, R) = \{y \in \mathbb{R} \setminus \{1\}: 2^{k_n-R} \leq |y - 1| < 2^{k_n+R+1}\},$$

where k_n is such that $|x_n - 1| \in [2^{k_n}, 2^{k_n+1})$, and hence $B(x_n, R) \subseteq \{y \neq 1: |y - 1| \geq 2^{\ell^*-R}\}$. Let $N_1 \geq N_0$ be such that $r_n < 2^{\ell^*-R}$ for every $n \geq N_1$. In this case, we have $J_n \cap B(x_n, R) = \emptyset$, which is impossible because our choice of ε would then force

$$1 = \int_{\mathbb{R}} |f_n|^2 \geq \int_{J_n} |f_n|^2 + \int_{B(x_n, R)} |f_n|^2 > 1.$$

It follows that $x_n \rightarrow 1$ as $n \rightarrow \infty$ and consequently the sequence $\{f_n\}$ concentrates at $y_0 = 1$. Indeed, given $\varepsilon > 0$, choose an integer R such that (3-4) holds. Then $B(x_n, R) \subseteq [1 - 2^{k_n+R+1}, 1 + 2^{k_n+R+1}] \setminus \{1\}$,

where $|x_n - 1| \in [2^{k_n}, 2^{k_n+1})$ and $k_n \rightarrow -\infty$, as $n \rightarrow \infty$, so that $2^{k_n+R+1} \rightarrow 0$, as $n \rightarrow \infty$. This forces

$$\int_{1-2^{k_n+R+1}}^{1+2^{k_n+R+1}} |f_n(y)|^2 dy \geq 1 - \varepsilon$$

for every $n \geq 1$, which implies concentration of the sequence $\{f_n\}$ at $y_0 = 1$.

Case 2: The sequence $\{|f_n|^2\}$ satisfies *dichotomy*. Let $\alpha \in (0, 1)$ be as in the dichotomy condition. Given $\varepsilon > 0$, consider the corresponding data R , k_0 , $\rho_{n,j} = |f_{n,j}|^2$, $j \in \{1, 2\}$, $\{x_n\} \subset \mathbb{R} \setminus \{1\}$, $\{R_n\} \subset [0, \infty)$. In particular,

$$\text{supp}(f_{n,1}) \subset B(x_n, R) \quad \text{and} \quad \text{supp}(f_{n,2}) \subset B(x_n, R_n)^c.$$

Since $R_n - R \rightarrow \infty$ as $n \rightarrow \infty$, by [Corollary 2.4](#) we obtain

$$\|\mathcal{E}_p(f_{n,1})\mathcal{E}_p(f_{n,2})\|_{L^3} \leq C_n \|f_{n,1}\|_{L^2} \|f_{n,2}\|_{L^2}, \quad (3-5)$$

where $C_n = C_n(\varepsilon) \lesssim 2^{-\beta(R_n - R)}$ for some $\beta > 0$. In particular, given $\varepsilon > 0$, we have $C_n \rightarrow 0$ as $n \rightarrow \infty$. Aiming at a contradiction, consider

$$\|\mathcal{E}_p(f_n - f_{n,1} - f_{n,2})\|_{L^6} \leq E_p \|f_n - (f_{n,1} + f_{n,2})\|_{L^2} \leq E_p \varepsilon^{\frac{1}{2}}. \quad (3-6)$$

The latter inequality requires a short justification which boils down to the pointwise estimate

$$(|f_n| - (|f_{n,1}| + |f_{n,2}|))^2 \leq ||f_n|^2 - (|f_{n,1}| + |f_{n,2}|)^2| = ||f_n|^2 - (|f_{n,1}|^2 + |f_{n,2}|^2)|. \quad (3-7)$$

This, in turn, follows from the disjointness of the supports of $f_{n,1}$ and $f_{n,2}$, together with the trivial estimate $||f_n| - (|f_{n,1}| + |f_{n,2}|)| \leq |f_n| + (|f_{n,1}| + |f_{n,2}|)$. In this way, [\(3-7\)](#) and [Proposition A.1](#) imply

$$\|(|f_n| - (|f_{n,1}| + |f_{n,2}|))^2\|_{L^1} \leq ||f_n|^2 - (|f_{n,1}|^2 + |f_{n,2}|^2)||_{L^1} \leq \varepsilon.$$

Coming back to [\(3-6\)](#), we have as an immediate consequence that

$$\|\mathcal{E}_p(f_n)\|_{L^6} \leq E_p \varepsilon^{\frac{1}{2}} + \|\mathcal{E}_p(f_{n,1} + f_{n,2})\|_{L^6}.$$

Expanding the binomial, using $\|f_{n,1}\|_{L^2}, \|f_{n,2}\|_{L^2} \leq 1$, and Hölder's inequality together with [\(3-5\)](#), we find that there exists c independent of n such that, for sufficiently large n ,

$$\begin{aligned} \|\mathcal{E}_p(f_{n,1} + f_{n,2})\|_{L^6}^6 &\leq \|\mathcal{E}_p(f_{n,1})\|_{L^6}^6 + \|\mathcal{E}_p(f_{n,2})\|_{L^6}^6 + c C_n \\ &\leq E_p^6 (\|f_{n,1}\|_{L^2}^6 + \|f_{n,2}\|_{L^2}^6) + c C_n \\ &\leq E_p^6 ((\alpha + \varepsilon)^3 + (1 - \alpha + \varepsilon)^3) + c C_n. \end{aligned} \quad (3-8)$$

This implies, for every sufficiently large n ,

$$\|\mathcal{E}_p(f_n)\|_{L^6} \leq E_p \varepsilon^{\frac{1}{2}} + (E_p^6 ((\alpha + \varepsilon)^3 + (1 - \alpha + \varepsilon)^3) + c C_n)^{\frac{1}{6}}.$$

Taking $n \rightarrow \infty$, and recalling that $\{f_n\}$ is an L^2 -normalized extremizing sequence for [\(1-9\)](#), we find that

$$E_p \leq E_p \varepsilon^{\frac{1}{2}} + E_p ((\alpha + \varepsilon)^3 + (1 - \alpha + \varepsilon)^3)^{\frac{1}{6}}$$

for every $\varepsilon > 0$. Taking $\varepsilon \rightarrow 0$ yields $1 \leq \alpha^3 + (1 - \alpha)^3$, which is impossible since $\alpha \in (0, 1)$. Hence dichotomy does not arise.

Case 3: The sequence $\{|f_n|^2\}$ satisfies *vanishing*. In this case,

$$\lim_{n \rightarrow \infty} \sup_{k \in \mathbb{Z}} \int_{2^{k-R} \leq |y-1| \leq 2^{k+R+1}} |f_n(y)|^2 dy = 0$$

for every integer $R < \infty$. In particular, for fixed $k \in \mathbb{N}$, we have

$$\lim_{n \rightarrow \infty} \int_{2^{-k} \leq |y-1| \leq 2^k} |f_n(y)|^2 dy = 0. \quad (3-9)$$

Set $f_{n,1} := f_n \mathbb{1}_{[1-2^{-k}, 1+2^{-k}]}$ and $f_{n,2} := f_n \mathbb{1}_{\{|y-1| \geq 2^k\}}$. Since $\|f_n - f_{n,1} - f_{n,2}\|_{L^2} \rightarrow 0$ as $n \rightarrow \infty$ it follows that $\{f_{n,1} + f_{n,2}\}_n$ is also an extremizing sequence for (1-9) for each $k \in \mathbb{N}$. This new sequence splits the mass into two separated regions, and so we expect to reach a contradiction if $\limsup_{n \rightarrow \infty} \|f_{n,2}\|_{L^2} > 0$, just as in Case 2. Set $\alpha_k := \limsup_{n \rightarrow \infty} \|f_{n,2}\|_{L^2}^2$ (recall that $f_{n,2}$ depends on k), and note that $\{\alpha_k\}$ is a constant sequence. Indeed,

$$\int_{|y-1| \geq 2^k} |f_n(y)|^2 dy = \int_{|y-1| \geq 2^{k+1}} |f_n(y)|^2 dy + \int_{2^k \leq |y-1| \leq 2^{k+1}} |f_n(y)|^2 dy \quad (3-10)$$

and from (3-9) with $k+1$ instead of k we have

$$\lim_{n \rightarrow \infty} \int_{2^k \leq |y-1| \leq 2^{k+1}} |f_n(y)|^2 dy = 0.$$

Taking $\limsup_{n \rightarrow \infty}$ in (3-10) yields $\alpha_{k+1} = \alpha_k$ for every $k \in \mathbb{N}$. An argument analogous to that of Case 2 (starting from (3-8)) shows that there exist $\beta > 0$ and a sequence $\{C_k\}$, $0 \leq C_k \lesssim 2^{-\beta k} \rightarrow 0$ as $k \rightarrow \infty$ such that

$$1 \leq \alpha_k^3 + (1 - \alpha_k)^3 + C_k \quad \text{for every } k \in \mathbb{N}.$$

Since $\alpha_k \equiv \alpha$ is constant, we may take $k \rightarrow \infty$ in the previous inequality and obtain $1 \leq \alpha^3 + (1 - \alpha)^3$. Since $\alpha \in [0, 1]$, necessarily $\alpha \in \{0, 1\}$. We claim that $\alpha = 0$. For any $k \geq 1$, the support of $f_{n,2}$ is disjoint from the interval J_n if n large enough. Thus

$$\|f_{n,2}\|_{L^2}^2 \leq 1 - \int_{J_n} |f_n|^2 \leq 1 - \inf_{n \in \mathbb{N}} \int_{J_n} |f_n|^2,$$

and therefore

$$\alpha \leq 1 - \inf_{n \in \mathbb{N}} \int_{J_n} |f_n|^2 < 1.$$

We conclude that $\alpha = 0$, as claimed. Finally, we show that vanishing implies concentration at $y = 1$. Since $1 = \|f_n\|_{L^2}^2 = \|f_{n,1}\|_{L^2}^2 + \|f_{n,2}\|_{L^2}^2 + o_n(1) = \|f_{n,1}\|_{L^2}^2 + o_n(1) = \|f_n \mathbb{1}_{[1-2^{-k}, 1+2^{-k}]}\|_{L^2}^2 + o_n(1)$, we find that, for every $k \in \mathbb{N}$,

$$\lim_{n \rightarrow \infty} \int_{1-2^{-k}}^{1+2^{-k}} |f_n(y)|^2 dy = 1.$$

This implies that the sequence $\{f_n\}$ concentrates at $y_0 = 1$.

To sum up, we proved that any sequence $\{f_n\}$ as in the statement of the proposition does not satisfy dichotomy, and that if it satisfies compactness or vanishing, then it concentrates at $y_0 = 1$. Thus the proof is complete. \square

As a second application of [Proposition A.1](#), we prove dyadic localization of extremizing sequences, after rescaling. We take $X = \mathbb{R}$, $\bar{x} = 0$, and use the dyadic pseudometric

$$\varrho: \mathbb{R} \setminus \{0\} \times \mathbb{R} \setminus \{0\} \rightarrow [0, \infty), \quad \varrho(x, y) := |k - k'|, \quad (3-11)$$

where this time $|x| \in [2^k, 2^{k+1})$ and $|y| \in [2^{k'}, 2^{k'+1})$. In this case, if R is an integer, then

$$B(x, R) = \{y \in \mathbb{R} \setminus \{0\}: 2^{k-R} \leq |y| < 2^{k+R+1}\}.$$

Proposition 3.4. *Let $\{f_n\} \subset L^2(\mathbb{R})$ be an L^2 -normalized extremizing sequence for (1-9). Then there exist a subsequence $\{f_{n_k}\}$, a sequence $\{a_k\} \subset \mathbb{R} \setminus \{0\}$, and a function $\Theta: [1, \infty) \rightarrow (0, \infty)$, $\Theta(R) \rightarrow 0$ as $R \rightarrow \infty$ such that the rescaled sequence $\{g_k\}$, $g_k := |a_k|^{1/2} f_{n_k}(a_k \cdot)$, satisfies*

$$\|g_k\|_{L^2([-R, R]^\mathbb{C})} \leq \Theta(R) \quad \text{for every } k \geq 1 \text{ and } R \geq 1. \quad (3-12)$$

This proposition will provide the input for the suitable application of the Brézis–Lieb lemma, which is formulated in the [Appendix](#) as [Proposition B.1](#).

Proof of [Proposition 3.4](#). Let $\{f_n\}$ be as in the statement of the proposition. In view of [Corollary 2.6](#), there exists $\ell_n \in \mathbb{Z}$ such that $\|f_n\|_{L^2(I_{\ell_n}^\bullet)} \gtrsim_p 1$, if n is large enough. Setting $g_n := 2^{\ell_n/2} f_n(2^{\ell_n} \cdot)$, we then have

$$\|g_n\|_{L^2(I_0^\bullet)} \gtrsim_p 1 \quad (3-13)$$

for every sufficiently large n . Using [Proposition A.1](#) with the pseudometric (3-11), we obtain a subsequence $\{|g_{n_k}|^2\}$ that satisfies one of three possibilities. Because of (3-13), vanishing does not occur. The argument given in Case 2 of the proof of [Proposition 3.3](#) can be used in conjunction with [Corollary 2.2](#) to show that the sequence $\{|g_{n_k}|^2\}$ does not satisfy dichotomy either. Therefore it must satisfy compactness. Thus, there exists a sequence $\{N_k\} \subset \mathbb{Z}$ such that, for every $k \geq 1$ and $\varepsilon > 0$, there exists an integer $r = r(\varepsilon)$ for which

$$\int_{2^{N_k-r} \leq |y| \leq 2^{N_k+r+1}} |g_k(y)|^2 dy \geq 1 - \varepsilon.$$

Because of (3-13), the sequence $\{N_k\}$ is bounded, $\sup_{k \geq 1} |N_k| =: r_0 < \infty$. By redefining r as $r + r_0 + 1$, it follows that

$$\int_{2^{-r} \leq |y| \leq 2^r} |g_k(y)|^2 dy \geq 1 - \varepsilon \quad \text{for every } k \geq 1. \quad (3-14)$$

Defining the function

$$\theta(R) := \sup_{k \geq 1} \int_{\{R^{-1} \leq |y| \leq R\}^\mathbb{C}} |g_k(y)|^2 dy,$$

$R \mapsto \theta(R)$ is a nonincreasing function of R which is bounded by 1 and, in view of (3-14), satisfies $\theta(R) \rightarrow 0$ as $R \rightarrow \infty$. By construction,

$$\int_{\{R^{-1} \leq |y| \leq R\}^{\mathbb{C}}} |g_k(y)|^2 dy \leq \theta(R) \quad \text{for every } k \geq 1, R \geq 1,$$

which implies (3-12) at once by taking $\Theta := \theta^{1/2}$. \square

We are finally ready to prove [Proposition 3.1](#).

Proof of Proposition 3.1. Let $\{f_n\}$ be as in the statement of the proposition. Apply [Proposition 3.4](#) to $\{f_n\}$, and denote the resulting rescaled subsequence by $\{g_n\}$. From the L^1 cap estimate (2-28) we know that, for each sufficiently large n , there exists an interval $J_n = [s_n - r_n, s_n + r_n]$, contained in a dyadic interval³ $[2^{k_n}, 2^{k_n+1}]$, such that

$$\int_{J_n} |g_n| \geq c |J_n|^{\frac{1}{2}}$$

for some $c > 0$ which is independent of n . By the Cauchy–Schwarz inequality,

$$\|g_n\|_{L^2(J_n)} \geq c, \quad (3-15)$$

and so estimate (3-12) implies the existence of $C > 0$ independent of n , such that $C^{-1} \leq |s_n| \leq C$. Rescaling again, we may assume $s_n = 1$ for every n .

If $r^* := \liminf_{n \rightarrow \infty} |J_n| > 0$, then passing to the relevant subsequence that realizes the limit inferior we have

$$\int_{1-2r^*}^{1+2r^*} g_n(y) dy = \int_{1-2r^*}^{1+2r^*} |g_n(y)| dy \geq \int_{J_n} |g_n| \gtrsim \sqrt{r^*},$$

provided n is large enough to ensure $J_n \subseteq [1-2r^*, 1+2r^*]$. Therefore any L^2 -weak limit of the sequence $\{g_n\}$ is nonzero. Here we used the nonnegativity of the sequence $\{g_n\}$. By [Proposition B.1](#), we conclude that there exists $0 \neq g \in L^2(\mathbb{R})$, such that possibly after a further extraction, $g_n \rightarrow g$ in $L^2(\mathbb{R})$, as $n \rightarrow \infty$. In other words, (i) holds.

It remains to consider the case when $|J_n| \rightarrow 0$, as $n \rightarrow \infty$. In view of (3-15), [Proposition 3.3](#) applies, and the sequence $\{g_n\}$ concentrates at $y_0 = 1$, i.e., (ii) holds. This finishes the proof of [Proposition 3.1](#) (and therefore of [Theorem 1.3](#)). \square

4. Existence of extremizers

In this section, we prove [Theorem 1.4](#). The basic strategy is to choose an appropriate trial function f for which the ratio from (1-10),

$$\Phi_p(f) := \frac{\|f \sigma_p * f \sigma_p * f \sigma_p\|_{L^2(\mathbb{R}^2)}^2}{\|f\|_{L^2(\mathbb{R})}^6}, \quad (4-1)$$

³Or its negative, but in that case we replace f_n by its reflection around the origin.

can be estimated via a simple lower bound. We will give different arguments depending on whether $1 < p < 2$ or $p > 2$, which rely on distinct choices of trial functions. This can be explained by the different qualitative nature of the 3-fold convolutions $wv_p * wv_p * wv_p$ in the two regimes of p ; see [Figure 1](#). Here, and throughout this section, $dv_p = \delta(s - |y|^p) dy ds$ denotes projection measure on the curve $s = |y|^p$, and the weight is given by $w = |\cdot|^{(p-2)/3}$. Note that $d\sigma_p = \sqrt{w} dv_p$.

The following analogue of [\[Oliveira e Silva and Quilodrán 2018, Proposition 6.4\]](#) holds for 3-fold convolutions in \mathbb{R}^2 .

Proposition 4.1. *Given $p > 1$, the following assertions hold for $wv_p * wv_p * wv_p$:*

- (a) *It is absolutely continuous with respect to Lebesgue measure on \mathbb{R}^2 .*
- (b) *Its support, denoted by E_p , is given by*

$$E_p = \{(\xi, \tau) \in \mathbb{R}^2 : \tau \geq 3^{1-p} |\xi|^p\}. \quad (4-2)$$

- (c) *If $p \geq 2$, then its Radon–Nikodym derivative, also denoted by $wv_p * wv_p * wv_p$, defines a bounded, continuous function in the interior of the set E_p . If $1 < p < 2$, then $wv_p * wv_p * wv_p$ defines a continuous function on the set*

$$\tilde{E}_p := \{(\xi, \tau) \in \mathbb{R}^2 : 3^{1-p} |\xi|^p < \tau < 2^{1-p} |\xi|^p\}.$$

- (d) *It is even in ξ , that is,*

$$(wv_p * wv_p * wv_p)(-\xi, \tau) = (wv_p * wv_p * wv_p)(\xi, \tau)$$

for every $\xi \in \mathbb{R}$, $\tau > 0$, and is homogeneous of degree zero in the sense that

$$(wv_p * wv_p * wv_p)(\lambda \xi, \lambda^p \tau) = (wv_p * wv_p * wv_p)(\xi, \tau) \quad \text{for every } \lambda > 0.$$

- (e) *It extends continuously to the boundary of E_p , except at the point $(\xi, \tau) = (0, 0)$, with values given by*

$$(wv_p * wv_p * wv_p)(\xi, 3^{1-p} |\xi|^p) = \frac{2\pi}{\sqrt{3}p(p-1)} \quad \text{if } \xi \neq 0. \quad (4-3)$$

Proof. For $p \geq 2$, the result follows from Proposition 2.1 and Remark 2.3 of [\[Oliveira e Silva and Quilodrán 2019\]](#). If $1 < p < 2$, then the weight w is singular at the origin, and an additional argument is required in order to establish parts (c) and (e) (as the others follow from [\[loc. cit.\]](#)). Note that part (e) also follows from [\[loc. cit.\]](#) after we verify (c), and so it suffices to show the latter.

Let $\psi = |\cdot|^p$. From [\[loc. cit., Remark 2.3\]](#), the formula

$$(wv_p * wv_p * wv_p)(\xi, \tau) = \int_{\mathbb{S}^1} \frac{\left(\left| \frac{1}{3}\xi + \alpha(\omega_1 + \omega_2) \right| \left| \frac{1}{3}\xi - \alpha\omega_1 \right| \left| \frac{1}{3}\xi - \alpha\omega_2 \right| \right)^{\frac{p-2}{3}}}{\langle \omega_1, \frac{\mathfrak{W}_1}{\alpha} \rangle + \langle \omega_2, \frac{\mathfrak{W}_2}{\alpha} \rangle} d\mu_{(\omega_1, \omega_2)}, \quad (4-4)$$

where

$$\mathfrak{W}_i(\xi, \tau, \omega_1, \omega_2) = \nabla \psi \left(\frac{1}{3}\xi + \alpha\omega_1 + \alpha\omega_2 \right) - \nabla \psi \left(\frac{1}{3}\xi - \alpha\omega_i \right), \quad i = 1, 2,$$

holds on \tilde{E}_p , provided that the function W defined by

$$W(\xi, \omega_1, \omega_2) := \left(\left| \frac{1}{3}\xi + \alpha(\omega_1 + \omega_2) \right| \left| \frac{1}{3}\xi - \alpha\omega_1 \right| \left| \frac{1}{3}\xi - \alpha\omega_2 \right| \right)^{\frac{p-2}{3}} \quad (4-5)$$

is continuous in the domain of integration. Here $\omega_1^2 + \omega_2^2 = 1$, arc-length measure on the unit circle \mathbb{S}^1 is denoted by μ , and the function $\alpha = \alpha(\xi, \tau, \omega_1, \omega_2)$ is implicitly defined by

$$\left| \frac{1}{3}\xi + \alpha(\omega_1 + \omega_2) \right|^p + \left| \frac{1}{3}\xi - \alpha\omega_1 \right|^p + \left| \frac{1}{3}\xi - \alpha\omega_2 \right|^p = \tau;$$

see [Oliveira e Silva and Quilodrán 2019] for details. It follows that

$$\left| \frac{1}{3}\xi + \alpha(\omega_1 + \omega_2) \right|^p + \left| \frac{1}{3}\xi - \alpha\omega_1 \right|^p + \left| \frac{1}{3}\xi - \alpha\omega_2 \right|^p < 2^{1-p} |\xi|^p,$$

provided $(\xi, \tau) \in \tilde{E}_p$. On the other hand, if $\frac{1}{3}\xi - \alpha\omega_1 = 0$, then convexity of ψ implies

$$\left| \frac{2}{3}\xi + \alpha\omega_2 \right|^p + \left| \frac{1}{3}\xi - \alpha\omega_2 \right|^p \geq 2^{1-p} |\xi|^p,$$

and similarly if $\frac{1}{3}\xi - \alpha\omega_2 = 0$, while if $\frac{1}{3}\xi + \alpha(\omega_1 + \omega_2) = 0$, then

$$\left| \frac{1}{3}\xi - \alpha\omega_1 \right|^p + \left| \frac{1}{3}\xi - \alpha\omega_2 \right|^p \geq 2^{1-p} \left| \frac{2}{3}\xi - \alpha(\omega_1 + \omega_2) \right|^p = 2^{1-p} |\xi|^p.$$

It follows that none of these three terms can vanish in a neighborhood of any point $(\xi, \tau) \in \tilde{E}_p$, and therefore W is continuous there. Thus identity (4-4) holds, and this concludes the verification of part (c). \square

The boundedness of $w\nu_p * w\nu_p * w\nu_p$ provides an alternative way towards estimate (1-10) via the usual application of the Cauchy–Schwarz inequality, at least in the restricted range $p \geq 2$. Moreover, identity (4-3) and the argument in Lemma 3.2 together imply that the corresponding optimal constant C_p satisfies

$$C_p^6 \geq \frac{2\pi}{\sqrt{3p(p-1)}},$$

which should be compared to (1-13).

4A. Effective lower bounds for C_p . We start by examining a simple lower bound, which is the analogue of [Oliveira e Silva and Quilodrán 2018, Lemma 6.1] for 3-fold convolutions in \mathbb{R}^2 .

Lemma 4.2. *Given a strictly convex function $\Psi : \mathbb{R} \rightarrow \mathbb{R}$ and a nonnegative function $w : \mathbb{R} \rightarrow [0, \infty)$, consider the measures $dv(y, s) = \delta(s - \Psi(y)) dy ds$ and $d\sigma = \sqrt{w} dv$. Let E denote the support of the convolution measure $\nu * \nu * \nu$. Given $\lambda > 0$, $a \in \mathbb{R}$, let $f_{\lambda, a}(y) := e^{-\lambda(\Psi(y) + ay)} \sqrt{w(y)}$. Then*

$$\frac{\|f_{\lambda, a}\sigma * f_{\lambda, a}\sigma * f_{\lambda, a}\sigma\|_{L^2(\mathbb{R}^2)}^2}{\|f_{\lambda, a}\|_{L^2(\mathbb{R})}^6} \geq \frac{\|f_{\lambda, a}\|_{L^2(\mathbb{R})}^6}{\int_E e^{-2\lambda(\tau + a\xi)} d\xi d\tau} \quad (4-6)$$

for every $f_{\lambda, a} \in L^2(\mathbb{R})$ such that $f_{\lambda, a}\sigma * f_{\lambda, a}\sigma * f_{\lambda, a}\sigma \in L^2(\mathbb{R}^2)$.

The proof is entirely parallel to that of [Oliveira e Silva and Quilodrán 2018, Lemma 6.1]. Note that (4-6) implies

$$\sup_{0 \neq f \in L^2(\mathbb{R})} \frac{\|f\sigma * f\sigma * f\sigma\|_{L^2(\mathbb{R}^2)}^2}{\|f\|_{L^2(\mathbb{R})}^6} \geq \sup_{\lambda > 0, a \in \mathbb{R}} \frac{\|f_{\lambda, a}\|_{L^2(\mathbb{R})}^6}{\int_E e^{-2\lambda(\tau + a\xi)} d\xi d\tau}. \quad (4-7)$$

Specializing [Lemma 4.2](#) to the case of the measure σ_p with the natural choice of trial function $f(y) = e^{-|y|^p} |y|^{(p-2)/6}$, a quick computation yields

$$\Phi_p(f) \geq \frac{4\Gamma\left(\frac{p+1}{3p}\right)^3}{3^{1-\frac{1}{p}} p^2 \Gamma\left(\frac{1}{p}\right)}. \quad (4-8)$$

This lower bound is good enough to establish the strict inequality [\(1-13\)](#) in a range of p that includes the cubic case $p = 3$ but *not* the quartic case $p = 4$, so we have to refine it. For the above choice of trial function, the corresponding ratio [\(4-1\)](#) can be expanded as an infinite series with nonnegative terms, whose coefficients are given in terms of the Gamma function and whose first term equals the expression on the right-hand side of [\(4-8\)](#).

Proposition 4.3. *Let $p > 1$ and $f(y) = e^{-|y|^p} |y|^{(p-2)/6} \in L^2(\mathbb{R})$. Then*

$$\Phi_p(f) = \frac{3^{1-\frac{1}{p}} p^2 \Gamma\left(\frac{1}{p}\right)}{2^3 \Gamma\left(\frac{p+1}{3p}\right)^3} \sum_{n=0}^{\infty} (4n+1) 2^{4n-1} \left(\sum_{k=0}^n \binom{2n}{2k} \binom{n+k-\frac{1}{2}}{2n} I_{2k}(p) \right)^2, \quad (4-9)$$

where the coefficients $\{I_{2k}(p)\}_{k \geq 0}$ are given by expression [\(4-15\)](#) below.

The proof will make use of the classical Legendre polynomials, denoted by $\{P_n\}_{n \geq 0}$, which constitute a family of orthogonal polynomials with respect to the L^2 -norm on the interval $[-1, 1]$. Explicitly, they are given by⁴

$$P_n(t) = 2^n \sum_{k=0}^n \binom{n}{k} \binom{\frac{n+k-1}{2}}{n} t^k, \quad -1 \leq t \leq 1, \quad (4-10)$$

from where one checks that $\langle P_m, P_n \rangle_{L^2} = (2/(2n+1)) \delta(n=m)$; see [\[Stein and Weiss 1971, Corollary 2.16, Chapter 4\]](#). See also [\[Carneiro and Oliveira e Silva 2015; Christ and Shao 2012a; Foschi 2015; Gonçalves 2019; Negro 2018\]](#) for earlier appearances of Legendre and other families of orthogonal polynomials in sharp Fourier restriction theory.

Proof of [Proposition 4.3](#). Start by noting that the function $f(y) = e^{-|y|^p} |y|^{(p-2)/6}$ coincides with $e^{-\tau} \sqrt{w(\xi)}$ on the support of σ_p . Using this together with parts (b) and (d) of [Proposition 4.1](#), we obtain

$$\begin{aligned} \|f\sigma_p * f\sigma_p * f\sigma_p\|_{L^2}^2 &= \|e^{-\tau} (w\nu_p * w\nu_p * w\nu_p)\|_{L^2}^2 \\ &= \int_0^\infty \int_{-3^{1-1/p} \tau^{1/p}}^{3^{1-1/p} \tau^{1/p}} e^{-2\tau} (w\nu_p * w\nu_p * w\nu_p)^2(\xi, \tau) d\xi d\tau \\ &= \int_0^\infty \int_{-3^{1-1/p}}^{3^{1-1/p}} \tau^{\frac{1}{p}} e^{-2\tau} (w\nu_p * w\nu_p * w\nu_p)^2(\tau^{\frac{1}{p}} \lambda, \tau) d\lambda d\tau \\ &= \left(\int_0^\infty \tau^{\frac{1}{p}} e^{-2\tau} d\tau \right) \int_{-3^{1-1/p}}^{3^{1-1/p}} (w\nu_p * w\nu_p * w\nu_p)^2(\lambda, 1) d\lambda \\ &= \frac{3^{1-\frac{1}{p}} \Gamma\left(\frac{1}{p}\right)}{p 2^{1+\frac{1}{p}}} \int_{-1}^1 (w\nu_p * w\nu_p * w\nu_p)^2(3^{1-\frac{1}{p}} t, 1) dt. \end{aligned} \quad (4-11)$$

⁴Recall that the binomial coefficient $\binom{\alpha}{n} := \alpha(\alpha-1)\cdots(\alpha-n+1)/n!$ is also defined when $\alpha \notin \mathbb{Z}$.

On the other hand,

$$\|f\|_{L^2}^2 = \int_{\mathbb{R}} e^{-2|y|^p} |y|^{\frac{p-2}{3}} dy = 2 \int_0^\infty e^{-2y^p} y^{\frac{p-2}{3}} dy = \frac{2^{\frac{2}{3}-\frac{1}{3p}}}{p} \Gamma\left(\frac{p+1}{3p}\right). \quad (4-12)$$

Given $t \in [-1, 1]$, define $g_p(t) := (w\nu_p * w\nu_p * w\nu_p)(3^{1-1/p}t, 1)$. Expanding g_p in the basis of Legendre polynomials,

$$\begin{aligned} \|g_p\|_{L^2([-1, 1], dt)}^2 &= \sum_{n=0}^{\infty} \frac{1}{\|P_n\|_{L^2}^2} \left(\int_{-1}^1 g_p(t) P_n(t) dt \right)^2 \\ &= \sum_{n=0}^{\infty} (4n+1) 2^{4n-1} \left(\sum_{k=0}^n \binom{2n}{2k} \binom{n+k-\frac{1}{2}}{2n} \int_{-1}^1 g_p(t) t^{2k} dt \right)^2, \end{aligned}$$

where the last identity follows from (4-10), the normalization $\|P_n\|_{L^2}^2 = 2/(2n+1)$, and the fact that g_p is an even function of t . We proceed to find an explicit expression for the moments $I_n(p) := \int_{-1}^1 g_p(t) t^n dt$. Given $b \in \mathbb{R}$, we compute

$$\begin{aligned} &\int_{\mathbb{R}^2} e^{-(\tau-b\xi)} (w\nu_p * w\nu_p * w\nu_p)(\xi, \tau) d\xi d\tau \\ &= \int_0^\infty \int_{-3^{1-1/p}}^{3^{1-1/p}} \tau^{\frac{1}{p}} e^{-\tau} e^{b\tau^{1/p}\lambda} (w\nu_p * w\nu_p * w\nu_p)(\lambda, 1) d\lambda d\tau \\ &= \sum_{n=0}^{\infty} \frac{3^{(1-\frac{1}{p})(2n+1)} b^{2n}}{(2n)!} \left(\int_0^\infty e^{-\tau} \tau^{\frac{2n+1}{p}} d\tau \right) \int_{-1}^1 t^{2n} (w\nu_p * w\nu_p * w\nu_p)(3^{1-\frac{1}{p}}t, 1) dt \\ &= \sum_{n=0}^{\infty} \frac{3^{(1-\frac{1}{p})(2n+1)} b^{2n}}{(2n)!} \frac{2n+1}{p} \Gamma\left(\frac{2n+1}{p}\right) I_{2n}(p). \end{aligned} \quad (4-13)$$

This Laplace transform can be alternatively computed as

$$\begin{aligned} &\int_{\mathbb{R}^2} e^{-(\tau-b\xi)} (w\nu_p * w\nu_p * w\nu_p)(\xi, \tau) d\xi d\tau = \left(\int_{\mathbb{R}} e^{-|y|^p} e^{by} |y|^{\frac{p-2}{3}} dy \right)^3 \\ &= \left(\sum_{n=0}^{\infty} \frac{2b^{2n}}{(2n)!} \int_0^\infty e^{-y^p} y^{\frac{p-2}{3}+2n} dy \right)^3 \\ &= \left(\sum_{n=0}^{\infty} \frac{2b^{2n}}{p(2n)!} \Gamma\left(\frac{p+1+6n}{3p}\right) \right)^3. \end{aligned} \quad (4-14)$$

Equating coefficients of the same degree, we obtain

$$I_{2n}(p) = \frac{2^3 (2n)!}{3^{(1-\frac{1}{p})(2n+1)} p^2 (2n+1) \Gamma\left(\frac{2n+1}{p}\right)} \sum_{k=0}^n \sum_{m=0}^{n-k} \frac{\Gamma\left(\frac{p+1+6k}{3p}\right) \Gamma\left(\frac{p+1+6m}{3p}\right) \Gamma\left(\frac{p+1+6(n-k-m)}{3p}\right)}{(2k)! (2m)! (2(n-k-m))!}. \quad (4-15)$$

Identity (4-9) follows at once, and the proof is complete. \square




Figure 1. Plots of the functions $g_{p,N}(t)$, appropriately normalized so that they are close to 1 at $t = 1$ for $p \in \{3, 4, 5, 10, 11, 12\}$ and $p \in \{\frac{4}{3}, \frac{3}{2}, \frac{5}{3}\}$. For $p \in \{3, 4, 5\}$ and $p \in \{\frac{4}{3}, \frac{3}{2}, \frac{5}{3}\}$ we used $N = 10$, for $p \in \{10, 11, 12\}$ we used $N = 15$.

Remark 4.4. From the preceding proof, we have the following approximating sequence $\{g_{p,N}\}_{N \geq 0}$ for g_p :

$$g_{p,N}(t) := \sum_{n=0}^N (4n+1) 2^{2n-1} \left(\sum_{k=0}^n \binom{2n}{2k} \binom{n+k-\frac{1}{2}}{2n} I_{2k}(p) \right) P_{2n}(t), \quad -1 \leq t \leq 1.$$

This was used to construct Figure 1. They correspond to approximate graphs of $wv_p * wv_p * wv_p$ on the region $\{(\xi, 1) : 0 \leq \xi \leq 3^{1-1/p}\}$ for different values of p . By homogeneity, the full picture on \mathbb{R}^2 can be obtained from these graphs. Figure 1 (top) indicates that, for large p , the function $g_p(t)$ becomes small as $t \rightarrow 0$. The function $(wv_p * wv_p * wv_p)(\xi, \tau)$ should then be small near the τ -axis, unlike the case of small values of p . This suggests that extremizing sequences may concentrate at the boundary if p is large enough.

4B. Proof of Theorem 1.4. We consider the case $p > 2$ first. From Theorem 1.3 and Proposition 4.3, it suffices to show that there exists $N \in \mathbb{N}$ such that

$$\frac{3^{1-\frac{1}{p}} p^2 \Gamma(\frac{1}{p})}{2^3 \Gamma(\frac{p+1}{3p})^3} \sum_{n=0}^N (4n+1) 2^{4n-1} \left(\sum_{k=0}^n \binom{2n}{2k} \binom{n+k-\frac{1}{2}}{2n} I_{2k}(p) \right)^2 > \frac{2\pi}{\sqrt{3p(p-1)}}, \quad (4-16)$$

where the coefficients $I_{2k}(p)$ are given by (4-15). The range of validity of (4-16) can be estimated by performing an accurate numerical calculation. Taking $N = 15$, one checks that inequality (4-16) holds

for every $p \in (2, p_0)$, where $p_0 \in [4, 5]$ and can be numerically estimated by $p_0 \approx 4.803$, with three decimal places. Increasing the value of N does not seem to substantially increase p_0 .

If $1 < p < 2$, then inequality (4-16) fails (for every $N \in \mathbb{N}$). Incidentally, note that if $p = 2$, then the left- and right-hand sides of (4-16) are equal (for every $N \in \mathbb{N}$) since the 3-fold convolution of projection measure on the parabola is constant inside its support; see [Foschi 2007, Lemma 4.1]. We are thus led to a different trial function. For $n \in \mathbb{N}$, define

$$f_n(y) = e^{-\frac{n}{2}(|y|^p - py)} |y|^{-\frac{2-p}{6}}. \quad (4-17)$$

In light of Lemma 3.2, the sequence $\{f_n\|f_n\|_{L^2}^{-1}\}$ concentrates at $y_0 = 1$. Passing to a continuous parameter $\lambda > 0$, Lemma 4.2 yields the lower bound

$$\Phi_p(f_\lambda) \geq \frac{\|f_\lambda\|_{L^2(\mathbb{R})}^6}{\int_{E_p} e^{-\lambda(\tau-p\xi)} d\xi d\tau} =: \phi_p(\lambda),$$

which we proceed to analyze. Since

$$\begin{aligned} \|f_\lambda\|_{L^2(\mathbb{R})}^2 &= \int_{-\infty}^{\infty} e^{-\lambda(|y|^p - py)} |y|^{-\frac{2-p}{3}} dy, \\ \int_{E_p} e^{-\lambda(\tau-p\xi)} d\xi d\tau &= \int_{-\infty}^{\infty} e^{\lambda p \xi} \left(\int_{3^{1-p}|\xi|^p}^{\infty} e^{-\lambda \tau} d\tau \right) d\xi = \frac{1}{\lambda} \int_{-\infty}^{\infty} e^{-\lambda(3^{1-p}|\xi|^p - p\xi)} d\xi, \end{aligned}$$

we have

$$\phi_p(\lambda) = \lambda \frac{\left(\int_{-\infty}^{\infty} e^{-\lambda(|y|^p - py)} |y|^{-\frac{2-p}{3}} dy \right)^3}{\int_{-\infty}^{\infty} e^{-\lambda(3^{1-p}|\xi|^p - p\xi)} d\xi}.$$

In view of (4-7), we have $C_p^6 \geq \phi_p(\lambda)$ for every $\lambda > 0$. Therefore it suffices to show that $\phi_p(\lambda) > 2\pi/(\sqrt{3}p(p-1))$, provided λ is large enough. This is the content of the following lemma, which we choose to formulate in terms of the function $\varphi_p(\lambda) := \phi_p(\lambda^{-1})$.

Lemma 4.5. *Let $p \in (1, 2)$. Then*

$$\lim_{\lambda \rightarrow 0^+} \varphi_p(\lambda) = \frac{2\pi}{\sqrt{3}p(p-1)}, \quad (4-18)$$

$$\lim_{\lambda \rightarrow 0^+} \varphi'_p(\lambda) = \frac{\pi(2-p)(2p-1)}{9\sqrt{3}p^2(p-1)^2}, \quad (4-19)$$

In particular, if $\lambda > 0$ is small enough, then $\varphi_p(\lambda) > 2\pi/(\sqrt{3}p(p-1))$.

Note that (4-18) follows from Lemma 3.2, but we choose to present a unified approach that establishes both (4-18) and (4-19).

Proof of Lemma 4.5. Rewrite ϕ_p in the equivalent form

$$\phi_p(\lambda) = \lambda \frac{\left(\int_{-\infty}^{\infty} e^{-\lambda(|y|^p - 1 - p(y-1))} |y|^{-\frac{2-p}{3}} dy \right)^3}{\int_{-\infty}^{\infty} e^{-\lambda 3^{1-p}(|y|^p - 3^p - p 3^{p-1}(y-3))} dy}.$$

Define real-valued functions $y \mapsto \alpha(y)$ and $y \mapsto \beta(y)$ via⁵

$$|y|^p - 1 - p(y-1) = \binom{p}{2}((y-1)^2 + \alpha(y-1)), \quad (4-20)$$

$$|y|^p - 3^p - p3^{p-1}(y-3) = 3^{p-2}\binom{p}{2}((y-3)^2 + \beta(y-3)).$$

By the binomial series expansion, if $|y| < 1$, then

$$\alpha(y) = \frac{p-2}{3}y^3 + \frac{(p-2)(p-3)}{12}y^4 + \dots, \quad (4-21)$$

$$\beta(y) = \frac{p-2}{3 \cdot 3}y^3 + \frac{(p-2)(p-3)}{12 \cdot 3^2}y^4 + \dots. \quad (4-22)$$

One easily checks that $|\alpha(y)| \rightarrow \infty$ and $|\beta(y)| \rightarrow \infty$ as $|y| \rightarrow \infty$, and

$$\lim_{\lambda \rightarrow \infty} \lambda \alpha(\lambda^{-\frac{1}{2}}y) = \lim_{\lambda \rightarrow \infty} \lambda \beta(\lambda^{-\frac{1}{2}}y) = 0 \quad (4-23)$$

for each $y \in \mathbb{R}$. We also have

$$\int_{\mathbb{R}} \exp\left(-\lambda \frac{|y|^p - 1 - p(y-1)}{\binom{p}{2}}\right) |y|^{-\frac{2-p}{3}} dy = \lambda^{-\frac{1}{2}} \int_{\mathbb{R}} e^{-y^2} e^{-\lambda \alpha(\lambda^{-1/2}y)} |1 + \lambda^{-\frac{1}{2}}y|^{-\frac{2-p}{3}} dy,$$

$$\int_{\mathbb{R}} \exp\left(-\lambda \frac{3^{1-p}(|y|^p - 3^p - p3^{p-1}(y-3))}{\binom{p}{2}}\right) dy = 3^{\frac{1}{2}} \lambda^{-\frac{1}{2}} \int_{\mathbb{R}} e^{-y^2} e^{-\frac{\lambda}{3}\beta((\frac{3}{\lambda})^{1/2}y)} dy,$$

and consequently

$$\phi_p\left(\frac{2\lambda}{p(p-1)}\right) = \frac{2}{\sqrt{3}p(p-1)} \frac{\left(\int_{\mathbb{R}} e^{-y^2} e^{-\lambda \alpha(\lambda^{-1/2}y)} |1 + \lambda^{-\frac{1}{2}}y|^{-\frac{2-p}{3}} dy\right)^3}{\int_{\mathbb{R}} e^{-y^2} e^{-\frac{\lambda}{3}\beta((\frac{3}{\lambda})^{1/2}y)} dy}.$$

For bookkeeping purposes, set

$$A_p(\lambda) := \left(\int_{\mathbb{R}} e^{-y^2} e^{-\lambda \alpha(\lambda^{-1/2}y)} |1 + \lambda^{-\frac{1}{2}}y|^{-\frac{2-p}{3}} dy \right)^3 \quad \text{and} \quad B_p(\lambda) := \int_{\mathbb{R}} e^{-y^2} e^{-\frac{\lambda}{3}\beta((\frac{3}{\lambda})^{1/2}y)} dy.$$

We now analyze each expression. Recalling (4-22), the numerator $A_p(\lambda)$ is seen to satisfy

$$A_p(\lambda) = \pi^{\frac{3}{2}} \left(1 - \frac{(p-2)(2p-1)}{144\lambda} + O(\lambda^{-\frac{3}{2}}) \right)^3 \quad \text{as } \lambda \rightarrow \infty. \quad (4-24)$$

Since binomial series expansions are only valid inside the unit ball, this step requires some care which we now briefly describe. Split the integral defining $A_p(\lambda)$ into three regions,

$$A_p^{\frac{1}{3}}(\lambda) = \left(\int_{-\infty}^{-\frac{\sqrt{\lambda}}{2}} + \int_{-\frac{\sqrt{\lambda}}{2}}^{\frac{\sqrt{\lambda}}{2}} + \int_{\frac{\sqrt{\lambda}}{2}}^{\infty} \right) e^{-y^2} e^{-\lambda \alpha(\lambda^{-1/2}y)} |1 + \lambda^{-\frac{1}{2}}y|^{-\frac{2-p}{3}} dy =: \text{I} + \text{II} + \text{III},$$

⁵Note that $\alpha(y) = 3^{-2}\beta(3y)$.

and estimate each of them separately. The main contribution comes from the integral $\text{II} = \text{II}(\lambda)$. Appealing to (4-21) and to the binomial series expansion, we have

$$\begin{aligned}\exp(-\lambda\alpha(\lambda^{-\frac{1}{2}}y)) &= 1 - \frac{p-2}{3}\lambda^{-\frac{1}{2}}y^3 - \frac{(p-2)(p-3)}{12}\lambda^{-1}y^4 + \frac{(p-2)^2}{18}\lambda^{-1}y^6 + O_y(\lambda^{-\frac{3}{2}}), \\ |1 + \lambda^{-\frac{1}{2}}y|^{-\frac{2-p}{3}} &= 1 + \frac{p-2}{3}\lambda^{-\frac{1}{2}}y + \frac{(p-2)(p-5)}{18}\lambda^{-1}y^2 + O_y(\lambda^{-\frac{3}{2}})\end{aligned}$$

uniformly in $y \in [-\sqrt{\lambda}/2, \sqrt{\lambda}/2]$. From this one easily checks that

$$\text{II}(\lambda) = \pi^{\frac{1}{2}} + \pi^{\frac{1}{2}} \frac{(p-2)(2p-1)}{144}\lambda^{-1} + O(\lambda^{-\frac{3}{2}}).$$

Matters are thus reduced to verifying that the contributions from I and III become negligible as $\lambda \rightarrow \infty$. On the region of integration of $\text{I} = \text{I}(\lambda)$, the factor $|1 + \lambda^{-1/2}y|^{-(2-p)/3}$ has an integrable singularity at $y = -\lambda^{1/2}$. Recalling the definition (4-20) of the function α , and changing variables $\lambda^{-1/2}y \rightsquigarrow x$, we have

$$\text{I}(\lambda) = \lambda^{\frac{1}{2}} \int_{-\infty}^{-\frac{1}{2}} e^{-\frac{2\lambda}{p(p-1)}(|1+x|^p - 1 - px)} |1+x|^{-\frac{2-p}{3}} dx.$$

Invoking the elementary inequality $|1 + x|^p - 1 - px \gtrsim_p |x|^p$, which is valid for every $x \leq -\frac{1}{2}$ and $1 < p < 2$, we may use Hölder's inequality together with the local integrability of $x \mapsto |1 + x|^{-(2-p)/3}$ in order to bound

$$\text{I}(\lambda) = O_p(\lambda^{\frac{1}{2}} \exp(-C_p \lambda))$$

for some $C_p > 0$. The contribution of $\text{III}(\lambda)$ is easier to handle because no singularity occurs on the corresponding region of integration. This concludes the verification of (4-24), which can then be differentiated term by term because there is sufficient decay. Therefore

$$\lim_{\lambda \rightarrow \infty} A_p(\lambda) = \pi^{\frac{3}{2}} \quad \text{and} \quad \lim_{\lambda \rightarrow \infty} -\lambda^2 A'_p(\lambda) = -\frac{3(p-2)(2p-1)\pi^{\frac{3}{2}}}{144}.$$

On the other hand, using the binomial series expansion (4-22) we obtain

$$\exp\left(-\frac{\lambda}{3}\beta\left(\left(\frac{3}{\lambda}\right)^{\frac{1}{2}}y\right)\right) = 1 - \frac{p-2}{3^{\frac{3}{2}}}\lambda^{-\frac{1}{2}}y^3 - \frac{(p-2)(p-3)}{36}\lambda^{-1}y^4 + \frac{(p-2)^2}{54}\lambda^{-1}y^6 + O_y(\lambda^{-\frac{3}{2}})$$

uniformly in $y \in \left[-\frac{1}{2}\left(\frac{1}{3}\lambda\right)^{1/2}, \frac{1}{2}\left(\frac{1}{3}\lambda\right)^{1/2}\right]$, so that an argument similar to that for $A_p(\lambda)$ gives

$$B_p(\lambda) = \pi^{\frac{1}{2}} + \frac{(p-2)(2p-1)\pi^{\frac{1}{2}}}{144\lambda} + O(\lambda^{-\frac{3}{2}}),$$

$$\lim_{\lambda \rightarrow \infty} B_p(\lambda) = \pi^{\frac{1}{2}} \quad \text{and} \quad \lim_{\lambda \rightarrow \infty} -\lambda^2 B'_p(\lambda) = \frac{(p-2)(2p-1)\pi^{\frac{1}{2}}}{144}.$$

We conclude

$$\lim_{\lambda \rightarrow 0^+} \varphi_p(\lambda) = \lim_{\lambda \rightarrow \infty} \phi_p(\lambda) = \lim_{\lambda \rightarrow \infty} \phi_p\left(\frac{2\lambda}{p(p-1)}\right) = \frac{2\pi}{\sqrt{3}p(p-1)}.$$

To address (4-19), note that

$$\varphi'_p(\lambda) = -\lambda^{-2}\phi'_p(\lambda^{-1}), \quad \text{and so } \lim_{\lambda \rightarrow 0^+} \varphi'_p(\lambda) = \lim_{\lambda \rightarrow \infty} -\lambda^2\phi'_p(\lambda).$$

Therefore

$$\lim_{\lambda \rightarrow \infty} -\lambda^2 \frac{d}{d\lambda} \left(\phi_p \left(\frac{2\lambda}{p(p-1)} \right) \right) = \frac{2\pi}{\sqrt{3}p(p-1)} \left(-\frac{3(p-2)(2p-1)}{144} - \frac{(p-2)(2p-1)}{144} \right) = \frac{\pi(2-p)(2p-1)}{18\sqrt{3}p(p-1)},$$

which readily implies (4-19). This completes the proof of the lemma (and therefore of Theorem 1.4). \square

4C. Improving p_0 . In view of the results from the last subsection, it is natural to let the functional Φ_p defined on (4-1) act on trial functions $f(y) = e^{-|y|^p} |y|^{(p-2)/6+a}$ for different choices of a .⁶ By doing so, the value $p_0 \approx 4.803$ can be improved. We turn to the details.

Set $\kappa := |\cdot|^{(p-2)/3+a}$, and note that

$$(\kappa \nu_p * \kappa \nu_p * \kappa \nu_p)(\lambda \xi, \lambda^p \tau) = \lambda^{3a} (\kappa \nu_p * \kappa \nu_p * \kappa \nu_p)(\xi, \tau) \quad \text{for every } \lambda > 0.$$

Reasoning as in (4-11) and (4-12), one checks that

$$\begin{aligned} \|f \sigma_p * f \sigma_p * f \sigma_p\|_{L^2(\mathbb{R}^2)}^2 &= \frac{3^{1-\frac{1}{p}} \Gamma\left(\frac{1+6a}{p}\right)}{p 2^{1+\frac{1+6a}{p}}} (1+6a) \int_{-1}^1 (\kappa \nu_p * \kappa \nu_p * \kappa \nu_p)^2(3^{1-\frac{1}{p}}t, 1) dt, \\ \|f\|_{L^2(\mathbb{R})}^2 &= \frac{2^{\frac{2}{3}-\frac{1+6a}{3p}}}{p} \Gamma\left(\frac{p+1+6a}{3p}\right). \end{aligned}$$

Given $t \in [-1, 1]$, define $h_p(t) := (\kappa \nu_p * \kappa \nu_p * \kappa \nu_p)(3^{1-1/p}t, 1)$. Expanding h_p in the basis of Legendre polynomials,

$$\|h_p\|_{L^2([-1, 1], dt)}^2 = \sum_{n=0}^{\infty} (4n+1) 2^{4n-1} \left(\sum_{k=0}^n \binom{2n}{2k} \binom{n+k-\frac{1}{2}}{2n} \int_{-1}^1 h_p(t) t^{2k} dt \right)^2.$$

We proceed to find explicit expressions for the moments $I_n(p, a) := \int_{-1}^1 h_p(t) t^n dt$. Given $b \in \mathbb{R}$, we compute as in (4-13) and (4-14)

$$\begin{aligned} \int_{\mathbb{R}^2} e^{-(\tau-b\xi)} (\kappa \nu_p * \kappa \nu_p * \kappa \nu_p)(\xi, \tau) d\xi d\tau &= \sum_{n=0}^{\infty} \frac{3^{(1-\frac{1}{p})(2n+1)} b^{2n}}{(2n)!} \frac{2n+1+3a}{p} \Gamma\left(\frac{2n+1+3a}{p}\right) I_{2n}(p, a) \\ &= \left(\sum_{n=0}^{\infty} \frac{2b^{2n}}{p(2n)!} \Gamma\left(\frac{p+1+6n+3a}{3p}\right) \right)^3. \end{aligned}$$

Equating coefficients as before, we find that the moment $I_{2n}(p, a)$ equals

$$\frac{3^{-(1-\frac{1}{p})(2n+1)} 2^3 (2n)!}{p^2 (2n+1+3a) \Gamma\left(\frac{2n+1+3a}{p}\right)} \sum_{k=0}^n \sum_{m=0}^{n-k} \frac{\Gamma\left(\frac{p+1+6k+3a}{3p}\right) \Gamma\left(\frac{p+1+6m+3a}{3p}\right) \Gamma\left(\frac{p+1+6(n-k-m)+3a}{3p}\right)}{(2k)! (2m)! (2(n-k-m))!}.$$

⁶Note that L^2 -integrability forces $a > -(p+1)/6$.

This implies

$$\Phi_p(f) = \frac{3^{1-\frac{1}{p}} p^2 \Gamma\left(\frac{1+6a}{p}\right)}{2^3 \Gamma\left(\frac{p+1+6a}{3p}\right)^3} (1+6a) \sum_{n=0}^{\infty} (4n+1) 2^{4n-1} \left(\sum_{k=0}^n \binom{2n}{2k} \binom{n+k-\frac{1}{2}}{2n} I_{2k}(p, a) \right)^2,$$

and consequently the following lower bound holds for every $N \geq 0$:

$$\Phi_p(f) \geq \frac{3^{1-\frac{1}{p}} p^2 \Gamma\left(\frac{1+6a}{p}\right)}{2^3 \Gamma\left(\frac{p+1+6a}{3p}\right)^3} (1+6a) \sum_{n=0}^N (4n+1) 2^{4n-1} \left(\sum_{k=0}^n \binom{2n}{2k} \binom{n+k-\frac{1}{2}}{2n} I_{2k}(p, a) \right)^2.$$

By numerically evaluating this sum with $N = 15$ and $a = \frac{7}{15}$, one can establish a lower bound that beats the critical threshold $2\pi/(\sqrt{3}p(p-1))$ for every $p \in (2, p_1)$, where $p_1 \approx 5.485$ with three decimal places. One further observes that the lower bound for small values of $a > 0$ is larger than that for $a = 0$, strongly suggesting that the original trial function $y \mapsto e^{-|y|^p} |y|^{(p-2)/6}$ might *not* be an extremizer in that range of exponents.

5. Superexponential L^2 -decay

This section is devoted to the proof of [Theorem 1.5](#). We follow the outline of [\[Erdoğan et al. 2011; Hundertmark and Shao 2012\]](#), and shall sometimes be brief. The Euler–Lagrange equation associated to [\(1-9\)](#) is

$$\mathcal{E}_p^*(\mathcal{E}_p(f)(\cdot, t) |\mathcal{E}_p(f)(\cdot, t)|^4) = \lambda f; \quad (5-1)$$

see [\[Christ and Quilodrán 2014, Proposition 2.4\]](#) for the variational derivation in a related context. The following 6-linear form will play a prominent role in the analysis:

$$Q(f_1, f_2, f_3, f_4, f_5, f_6) := \int_{\mathbb{R}^2} \prod_{j=1}^3 \mathcal{E}_p(f_j)(x, t) \overline{\mathcal{E}_p(f_{j+3})(x, t)} \, dx \, dt.$$

An immediate consequence of [\(1-9\)](#) is the basic estimate

$$|Q(f_1, f_2, f_3, f_4, f_5, f_6)| \lesssim \prod_{j=1}^6 \|f_j\|_{L^2(\mathbb{R})}. \quad (5-2)$$

The form Q can be rewritten as

$$Q(f_1, f_2, f_3, f_4, f_5, f_6) = \int_{\mathbb{R}^6} \prod_{j=1}^3 f_j(y_j) |y_j|^{\frac{p-2}{6}} \overline{f_{j+3}(y_{j+3})} |y_{j+3}|^{\frac{p-2}{6}} \delta(\alpha(y)) \delta(\beta(y)) \, dy,$$

where $y = (y_1, \dots, y_6) \in \mathbb{R}^6$ and

$$\begin{aligned} \alpha(y) &:= |y_1|^p + |y_2|^p + |y_3|^p - |y_4|^p - |y_5|^p - |y_6|^p, \\ \beta(y) &:= y_1 + y_2 + y_3 - y_4 - y_5 - y_6. \end{aligned}$$

We will also consider the associated form

$$K(f_1, f_2, f_3, f_4, f_5, f_6) := Q(|f_1|, |f_2|, |f_3|, |f_4|, |f_5|, |f_6|),$$

which is sublinear in each entry. Clearly,

$$|Q(f_1, f_2, f_3, f_4, f_5, f_6)| \leq K(f_1, f_2, f_3, f_4, f_5, f_6), \quad (5-3)$$

$$K(f_1, f_2, f_3, f_4, f_5, f_6) \lesssim \prod_{j=1}^6 \|f_j\|_{L^2(\mathbb{R})}. \quad (5-4)$$

Let us now introduce a parameter $s \geq 1$, which will typically be large. If there exist $j \neq k$ such that f_j is supported on $[-s, s]$ and f_k is supported outside of $[-Cs, Cs]$ for some $C > 1$, then estimate (5-4) can be improved to

$$K(f_1, f_2, f_3, f_4, f_5, f_6) \lesssim C^{-\frac{p-1}{6}} \prod_{j=1}^6 \|f_j\|_{L^2(\mathbb{R})}, \quad (5-5)$$

in accordance with the bilinear estimates of [Corollary 2.2](#). Introducing the weighted variant

$$K_G(f_1, f_2, f_3, f_4, f_5, f_6) := \int_{\mathbb{R}^6} e^{G(y_1) - \sum_{j=2}^6 G(y_j)} \prod_{j=1}^6 |f_j(y_j)| |y_j|^{\frac{p-2}{6}} \delta(\alpha(y)) \delta(\beta(y)) dy,$$

one checks at once that

$$K(e^G f_1, e^{-G} f_2, e^{-G} f_3, e^{-G} f_4, e^{-G} f_5, e^{-G} f_6) = K_G(f_1, f_2, f_3, f_4, f_5, f_6). \quad (5-6)$$

Given $\mu, \varepsilon \geq 0$, define the function

$$G_{\mu, \varepsilon}(y) := \frac{\mu|y|^p}{1 + \varepsilon|y|^p}. \quad (5-7)$$

The same proof as [\[Hundertmark and Shao 2012, Proposition 4.5\]](#) yields

$$K_{G_{\mu, \varepsilon}}(f_1, f_2, f_3, f_4, f_5, f_6) \leq K(f_1, f_2, f_3, f_4, f_5, f_6); \quad (5-8)$$

see also Remark 4.6 of that paper. Split $f = f_< + f_>$ with $f_> := f \mathbb{1}_{[-s^2, s^2]^\mathbb{C}}$, and define

$$\|f\|_{\mu, s, \varepsilon} := \|e^{G_{\mu, \varepsilon}} f_>\|_{L^2}.$$

Definition 5.1. A function $f \in L^2(\mathbb{R})$ is said to be a *weak solution* of (5-1) if there exists $\lambda > 0$ such that

$$Q(g, f, f, f, f, f) = \lambda \langle g, f \rangle_{L^2} \quad \text{for every } g \in L^2(\mathbb{R}). \quad (5-9)$$

Note that if f extremizes (1-9), then f satisfies (5-9) with $\lambda = E_p^6 \|f\|_{L^2}^4$. The following key step shows that for some positive μ , the quantity $\|f\|_{\mu, s, \varepsilon}$ is bounded in $\varepsilon > 0$.

Proposition 5.2. Given $p > 1$, let f be a weak solution of the Euler–Lagrange equation (5-1) with $\|f\|_{L^2} = 1$. If $s \geq 1$ is sufficiently large, then there exists $C < \infty$ such that

$$\lambda \|f\|_{s^{-2p}, s, \varepsilon} \leq o_1(1) \|f\|_{s^{-2p}, s, \varepsilon} + C \sum_{\ell=2}^5 \|f\|_{s^{-2p}, s, \varepsilon}^\ell + o_2(1), \quad (5-10)$$

where for $j \in \{1, 2\}$ we have $o_j(1) \rightarrow 0$ as $s \rightarrow \infty$ uniformly in ε . Moreover the constant C is independent of s and ε .

Proof. We start by introducing some notation. Let $G := G_{\mu, \varepsilon}$ be as in (5-7). Let $h := e^G f$, $h_> := e^G f_>$, and $h_< := h - h_>$. Further split $f_< = f_{\ll} + f_{\sim}$ and $h_< = h_{\ll} + h_{\sim}$, where $f_{\ll} := f \mathbb{1}_{[-s, s]}$ and $h_{\ll} := e^G f_{\ll}$. Since f satisfies (5-9), we have

$$\begin{aligned} \lambda \|e^G f_>\|_{L^2}^2 &= \lambda \langle e^{2G} f_>, f_> \rangle_{L^2} = \lambda \langle e^{2G} f_>, f \rangle_{L^2} = Q(e^{2G} f_>, f, f, f, f, f) \\ &= Q(e^G h_>, f, f, f, f, f) = Q(e^G h_>, e^{-G} h, e^{-G} h, e^{-G} h, e^{-G} h, e^{-G} h) =: Q_G. \end{aligned}$$

It follows from (5-3), (5-6), and (5-8) that $|Q_G| \lesssim K(h_>, h, h, h, h, h)$. Writing $h = h_< + h_>$, the sublinearity of K implies

$$|Q_G| \lesssim K(h_>, h_<, h_<, h_<, h_<, h_<) + \left(\sum' + \sum'' \right) K(h_>, h_{j_2}, h_{j_3}, h_{j_4}, h_{j_5}, h_{j_6}),$$

where the first sum, denoted by B_1 , is taken over indices $j_2, \dots, j_6 \in \{>, <\}$ with exactly one of the j_k equal to $>$, and the second sum, denoted by B_2 , is taken over indices $j_2, \dots, j_6 \in \{>, <\}$ with two or more of the j_k equal to $>$. We estimate the three terms separately. For the first one,

$$\begin{aligned} A &:= K(h_>, h_<, h_<, h_<, h_<, h_<) \leq K(h_>, h_{\ll}, h_<, h_<, h_<, h_<) + K(h_>, h_{\sim}, h_<, h_<, h_<, h_<) \\ &\lesssim \|h_>\|_{L^2} (s^{-\frac{p-1}{6}} \|h_{\ll}\|_{L^2} + \|h_{\sim}\|_{L^2}) \|h_<\|_{L^2}^4, \end{aligned}$$

where we made use of the support separation of $h_>$ and h_{\ll} via (5-5). Since $\|f\|_{L^2} = 1$, the estimates

$$\|h_<\|_{L^2} \lesssim e^{\mu s^{2p}}, \quad \|h_{\ll}\|_{L^2} \lesssim e^{\mu s^p}, \quad \text{and} \quad \|h_{\sim}\|_{L^2} \lesssim e^{\mu s^{2p}} \|f_{\sim}\|_{L^2}$$

hold and therefore

$$A \lesssim \|h_>\|_{L^2} (s^{-\frac{p-1}{6}} e^{\mu(s^p - s^{2p})} + \|f_{\sim}\|_{L^2}) e^{5\mu s^{2p}}.$$

The terms B_1, B_2 can be estimated in a similar way. One obtains

$$B_1 \lesssim \|h_>\|_{L^2}^2 (s^{-\frac{p-1}{6}} e^{\mu(s^p - s^{2p})} + \|f_{\sim}\|_{L^2}) e^{4\mu s^{2p}} \quad \text{and} \quad B_2 \lesssim \|h_>\|_{L^2} \left(\sum_{\ell=2}^5 \|h_>\|_{L^2}^{\ell} \right) e^{3\mu s^{2p}}.$$

The result follows by choosing $\mu = s^{-2p}$ and noting that $\|f_{\sim}\|_{L^2} \rightarrow 0$, as $s \rightarrow \infty$. \square

We are finally ready to prove that extremizers decay superexponentially fast.

Proof of Theorem 1.5. Let $f \in L^2$ be an extremizer of (1-9), normalized so that $\|f\|_{L^2} = 1$. Then f satisfies (5-9) with $\lambda = E_p^6$. Note that the function $(s, \varepsilon) \mapsto \|f\|_{s^{-2p}, s, \varepsilon}$ is continuous in $(s, \varepsilon) \in (0, \infty)^2$ and, for each fixed $\varepsilon > 0$,

$$\|f\|_{s^{-2p}, s, \varepsilon} = \|e^{G_{s^{-2p}, \varepsilon}} f \mathbb{1}_{[-s^2, s^2]} \|_{L^2} \rightarrow 0 \quad \text{as } s \rightarrow \infty. \quad (5-11)$$

Let C be the constant promised by Proposition 5.2, and consider the function

$$H(v) := \frac{1}{2} \lambda v - C(v^2 + v^3 + v^4 + v^5).$$

In (5-10) choose s sufficiently large so that $o_1(1) \leq \frac{1}{2}\lambda$ for every $\varepsilon > 0$. This is possible since $o_1(1) \rightarrow 0$ as $s \rightarrow \infty$ uniformly in $\varepsilon > 0$. Consequently,

$$H(\|f\|_{s-2p,s,\varepsilon}) \leq o_2(1) \quad \text{for every } \varepsilon > 0.$$

In view of (5-11), and the facts that $H(0) = 0$, $H'(0) > 0$, and H is concave on $[0, \infty)$, we may choose s sufficiently large so that $\sup_{\varepsilon > 0} o_2(1) < H(v_0)$ and $\|f\|_{s-2p,s,1} \leq v_0$, where $0 < v_0 < v_1$ are the two unique positive solutions of the equation

$$H(v_j) = \frac{1}{2} \max\{H(v) : v \geq 0\}.$$

By continuity, $\|f\|_{s-2p,s,\varepsilon} \leq v_0$ for every $\varepsilon > 0$. The monotone convergence theorem then implies $\|f\|_{s-2p,s,0} \leq v_0 < \infty$, which translates into

$$e^{s-2p} |\cdot|^p f \in L^2(\mathbb{R}).$$

Letting $\mu_0 := s^{-2p}$, where s is large enough so that all of the above steps hold, we have thus proved the first part. For the second part, note that, for every $\mu \in \mathbb{R}$, the function

$$e^{\mu|x|} f(x) = e^{\mu|x| - \mu_0|x|^p} \cdot e^{\mu_0|x|^p} f(x)$$

belongs to $L^2(\mathbb{R})$, since the first factor is bounded (here we use $p > 1$) and the second factor is, as we have just seen, square integrable. The result then follows from the Paley–Wiener theorem as in [Reed and Simon 1975, Theorem IX.13]. \square

We finish with two concluding remarks. Firstly, the argument can be adapted to the case of extremizers for odd curves treated in the next section. Secondly, an interesting problem is whether extremizers are smooth (and not only their Fourier transforms). This question has been addressed in the context of the Fourier extension operator on low-dimensional spheres in [Christ and Shao 2012b; Shao 2016b], but we have not investigated the extent to which their analysis can be adapted to the present case.

6. The case of odd curves

In this section we discuss the necessary modifications to establish analogues of Theorems 1.3 and 1.4 for odd curves. In general terms, the analysis is similar, but the existence of parallel tangents requires an extra symmetrization step. Estimate (1-15) can be rewritten as

$$\|\mathcal{S}_p(f)\|_{L^6(\mathbb{R}^2)} \leq \mathcal{O}_p \|f\|_{L^2(\mathbb{R})}, \quad (6-1)$$

where the Fourier extension operator on the curve $s = y|y|^{p-1}$ is given by

$$\mathcal{S}_p(f)(x, t) = \int_{\mathbb{R}} e^{ixy} e^{ity|y|^{p-1}} |y|^{\frac{p-2}{6}} f(y) dy. \quad (6-2)$$

Given a real-valued function $f \in L^2(\mathbb{R})$, denote the reflection of f with respect to the origin by $\tilde{f} := f(-\cdot)$. One easily checks that

$$\mathcal{S}_p(\tilde{f})(x, t) = \mathcal{S}_p(f)(-x, -t) = \overline{\mathcal{S}_p(f)(x, t)},$$

where the bar denotes complex conjugation. In particular,

$$\|\mathcal{S}_p(f)\mathcal{S}_p(g)\|_{L^3} = \|\mathcal{S}_p(f)\mathcal{S}_p(\tilde{g})\|_{L^3},$$

and so functions f, g supported on intervals I and $-I$, respectively, are seen to interact in the same way as if they were both supported on I , unlike the case of even curves. In this way, one is led to symmetrize with respect to reflection. This has already been observed in the case of the spheres \mathbb{S}^1 [Shao 2016a] and \mathbb{S}^2 [Christ and Shao 2012a]. Symmetrization on \mathbb{S}^2 has been efficiently handled via δ -calculus in [Foschi 2015]. The same method can be applied to the present case, but we choose to present a different argument which does not rely on the underlying convolution structure.

Lemma 6.1. *Let $p > 1$ and $f \in L^2(\mathbb{R})$. Then*

$$\frac{\|\mathcal{S}_p(f)\|_{L^6(\mathbb{R}^2)}}{\|f\|_{L^2(\mathbb{R})}} \leq \sup_{\substack{0 \neq g \in L^2(\mathbb{R}) \\ g \text{ even}}} \frac{\|\mathcal{S}_p(g)\|_{L^6(\mathbb{R}^2)}}{\|g\|_{L^2(\mathbb{R})}}. \quad (6-3)$$

If equality holds in (6-3), then f is necessarily an even function.

Proof. Given $f \in L^2(\mathbb{R})$, $f \neq 0$, take the decomposition $f = f_e + f_o$, where f_e is an even function, $f_e = \tilde{f}_e$ a.e. in \mathbb{R} , and f_o is odd, $f_o = -\tilde{f}_o$ a.e. in \mathbb{R} . Then $\|f\|_{L^2}^2 = \|f_e\|_{L^2}^2 + \|f_o\|_{L^2}^2$, and $\mathcal{S}_p(f_e)$ is real-valued, while $\mathcal{S}_p(f_o)$ is purely imaginary. Thus

$$|\mathcal{S}_p(f)(x, t)|^2 = |\mathcal{S}_p(f_e)(x, t)|^2 + |\mathcal{S}_p(f_o)(x, t)|^2 \quad \text{for almost every } (x, t) \in \mathbb{R}^2, \quad (6-4)$$

and so, by the triangle inequality for the L^3 -norm, $\|\mathcal{S}_p(f)\|_{L^6}^2 \leq \|\mathcal{S}_p(f_e)\|_{L^6}^2 + \|\mathcal{S}_p(f_o)\|_{L^6}^2$. It follows that

$$\frac{\|\mathcal{S}_p(f)\|_{L^6}^2}{\|f\|_{L^2}^2} \leq \frac{\|\mathcal{S}_p(f_e)\|_{L^6}^2 + \|\mathcal{S}_p(f_o)\|_{L^6}^2}{\|f_e\|_{L^2}^2 + \|f_o\|_{L^2}^2} \leq \max \left\{ \frac{\|\mathcal{S}_p(f_e)\|_{L^6}^2}{\|f_e\|_{L^2}^2}, \frac{\|\mathcal{S}_p(f_o)\|_{L^6}^2}{\|f_o\|_{L^2}^2} \right\},$$

where we set either ratio on the right-hand side of this chain of inequalities to zero whenever the corresponding function f_e or f_o happens to vanish identically. Therefore we may restrict attention to functions which are either even or odd. On the other hand, the equivalent convolution form (1-15) of the inequality implies $\|\mathcal{S}_p(g)\|_{L^6} \leq \|\mathcal{S}_p(|g|)\|_{L^6}$, with equality if and only if $g = |g|$ a.e. in \mathbb{R} . Thus

$$\frac{\|\mathcal{S}_p(f)\|_{L^6}^2}{\|f\|_{L^2}^2} \leq \max \left\{ \frac{\|\mathcal{S}_p(f_e)\|_{L^6}^2}{\|f_e\|_{L^2}^2}, \frac{\|\mathcal{S}_p(|f_o|)\|_{L^6}^2}{\|f_o\|_{L^2}^2} \right\} \leq \sup_{\substack{0 \neq g \in L^2 \\ g \text{ even}}} \frac{\|\mathcal{S}_p(g)\|_{L^6}}{\|g\|_{L^2}}, \quad (6-5)$$

where we used that both f_e and $|f_o|$ are even functions. In order for equality to hold in (6-3), both inequalities in (6-5) must be equalities. Inspection of the chain of inequalities leading to (6-5) shows that, if there is equality in the first inequality, then necessarily one of the following alternatives must hold:

- $\|f_o\|_{L^2} = 0$, in which case $f = f_e$, and so f is even; or
- $\|f_e\|_{L^2} = 0$ and $f_o = |f_o|$ a.e. in \mathbb{R} , which implies that $f_o \equiv 0$, and so $f \equiv 0$ which does not hold by assumption; or

- $\|f_e\|_{L^2}\|f_o\|_{L^2} \neq 0$ and $\|\mathcal{S}_p(f_e)\|_{L^6}\|f_e\|_{L^2}^{-1} = \|\mathcal{S}_p(f_o)\|_{L^6}\|f_o\|_{L^2}^{-1} = \|\mathcal{S}_p(|f_o|)\|_{L^6}\|f_o\|_{L^2}^{-1}$, which again forces $f_o = |f_o|$ a.e. in \mathbb{R} , so that $f_o = 0$ which is absurd.

Therefore equality in (6-3) forces f to be an even function, as desired. \square

For the remainder of this section, we restrict attention to nonnegative, even functions f . To prove the analogue of [Proposition 3.1](#), we need bilinear estimates as in [Propositions 2.1](#) and [2.3](#), and an L^1 cap bound as in [Proposition 2.8](#). These can be obtained in exactly the same way as for the case of even curves, since the Jacobian factor corresponding to (2-2) is now equal to $p||y'|^{p-1} - |y|^{p-1}|$, which amounts to the bound we used before. We also need an analogue of [Proposition A.1](#) with two points removed; i.e., consider $X_{\bar{x}, \bar{y}} := X \setminus \{\bar{x}, \bar{y}\}$ equipped with a pseudometric $\varrho : X_{\bar{x}, \bar{y}} \times X_{\bar{x}, \bar{y}} \rightarrow [0, \infty)$. The statement is analogous so we omit it. Next, defining the dyadic pseudometric centered at zero as in (3-11) and invoking the appropriate bilinear estimates, we obtain an analogue of [Proposition 3.4](#), the statement again being identical (omitted). The analogue of [Proposition 3.3](#) requires the pseudometric

$$\varrho : \mathbb{R} \setminus \{-1, 1\} \times \mathbb{R} \setminus \{-1, 1\} \rightarrow [0, \infty), \quad \varrho(x, y) := |k - k'|,$$

where $k, k' \in \mathbb{Z}$ are such that $||x| - 1| \in [2^k, 2^{k+1})$ and $||y| - 1| \in [2^{k'}, 2^{k'+1})$. It handles concentration at a pair of opposite points, which we now define.

Definition 6.2. Let $y_0 \in \mathbb{R}$. A sequence of even functions $\{f_n\} \subset L^2(\mathbb{R})$ *concentrates at the pair $\{-y_0, y_0\}$* if, for every $\varepsilon, \rho > 0$, there exists $N \in \mathbb{N}$ such that, for every $n \geq N$,

$$\int_{\substack{|y+y_0| \geq \rho \\ |y-y_0| \geq \rho}} |f_n(y)|^2 dy < \varepsilon \|f_n\|_{L^2(\mathbb{R})}^2.$$

The following analogue of [Proposition 3.3](#) holds for odd curves.

Proposition 6.3. Let $\{f_n\} \subset L^2(\mathbb{R})$ be an L^2 -normalized extremizing sequence of even functions for (6-1). Let $\{r_n\}$ be a sequence of nonnegative numbers satisfying $r_n \rightarrow 0$ as $n \rightarrow \infty$, and

$$\inf_{n \in \mathbb{N}} \int_{1-r_n}^{1+r_n} |f_n(y)|^2 dy > 0.$$

Then the sequence $\{f_n\}$ concentrates at the pair $\{-1, 1\}$.

As in the case of even curves, this can be used to prove the analogue of [Proposition 3.1](#).

Proposition 6.4. Let $\{f_n\} \subset L^2(\mathbb{R})$ be an L^2 -normalized extremizing sequence of nonnegative, even functions for (6-1). Then there exist a subsequence $\{f_{n_k}\}$ and a sequence $\{a_k\} \subset \mathbb{R} \setminus \{0\}$ such that the rescaled sequence $\{g_k\}$, $g_k := |a_k|^{1/2} f_{n_k}(a_k \cdot)$, satisfies one of the following conditions:

- (i) There exists $g \in L^2(\mathbb{R})$ such that $g_k \rightarrow g$ in $L^2(\mathbb{R})$ as $k \rightarrow \infty$.
- (ii) $\{g_k\}$ concentrates at the pair $\{-1, 1\}$.

Let $\{f_n\} \subset L^2(\mathbb{R})$ be an L^2 -normalized sequence of nonnegative, even functions concentrating at the pair $\{-1, 1\}$. Write $f_n = g_n + \tilde{g}_n$, where $g_n := f_n \mathbb{1}_{[0, \infty)}$. In particular, $\|g_n\|_{L^2} = 2^{-1/2}$, and the

sequence $\{g_n\}$ concentrates at $y_0 = 1$. The left-hand side of (1-15) can be expanded into

$$\begin{aligned}
\|f_n \mu_p * f_n \mu_p * f_n \mu_p\|_{L^2}^2 &= \|g_n \mu_p * g_n \mu_p * g_n \mu_p\|_{L^2}^2 + \|\tilde{g}_n \mu_p * \tilde{g}_n \mu_p * \tilde{g}_n \mu_p\|_{L^2}^2 \\
&\quad + 9\|g_n \mu_p * g_n \mu_p * \tilde{g}_n \mu_p\|_{L^2}^2 + 9\|g_n \mu_p * \tilde{g}_n \mu_p * \tilde{g}_n \mu_p\|_{L^2}^2 \\
&\quad + 6\langle g_n \mu_p * g_n \mu_p * g_n \mu_p, g_n \mu_p * g_n \mu_p * \tilde{g}_n \mu_p \rangle_{L^2} \\
&\quad + 6\langle g_n \mu_p * \tilde{g}_n \mu_p * \tilde{g}_n \mu_p, \tilde{g}_n \mu_p * \tilde{g}_n \mu_p * \tilde{g}_n \mu_p \rangle_{L^2} \\
&\quad + 18\langle g_n \mu_p * \tilde{g}_n \mu_p * \tilde{g}_n \mu_p, g_n \mu_p * g_n \mu_p * \tilde{g}_n \mu_p \rangle_{L^2} \\
&\quad + 6\langle g_n \mu_p * g_n \mu_p * g_n \mu_p, g_n \mu_p * \tilde{g}_n \mu_p * \tilde{g}_n \mu_p \rangle_{L^2} \\
&\quad + 6\langle g_n \mu_p * g_n \mu_p * \tilde{g}_n \mu_p, \tilde{g}_n \mu_p * \tilde{g}_n \mu_p * \tilde{g}_n \mu_p \rangle_{L^2} \\
&\quad + 2\langle g_n \mu_p * g_n \mu_p * g_n \mu_p, \tilde{g}_n \mu_p * \tilde{g}_n \mu_p * \tilde{g}_n \mu_p \rangle_{L^2}. \quad (6-6)
\end{aligned}$$

The last three summands vanish since the corresponding supports intersect on a Lebesgue null set. The symmetry of the inner products then implies

$$\begin{aligned}
\|f_n \mu_p * f_n \mu_p * f_n \mu_p\|_{L^2}^2 &= 20\|g_n \mu_p * g_n \mu_p * g_n \mu_p\|_{L^2}^2 + 30\langle g_n \mu_p * g_n \mu_p * g_n \mu_p * g_n \mu_p, g_n \mu_p * g_n \mu_p \rangle_{L^2}.
\end{aligned}$$

Note that $\mu_p = \sigma_p$ on the support of g_n , where σ_p was defined in (1-11). It follows that

$$\begin{aligned}
\frac{\|f_n \mu_p * f_n \mu_p * f_n \mu_p\|_{L^2}^2}{\|f_n\|_{L^2}^6} &= \frac{5}{2} \frac{\|g_n \sigma_p * g_n \sigma_p * g_n \sigma_p\|_{L^2}^2}{\|g_n\|_{L^2}^6} + \frac{15}{4} \frac{\langle g_n \sigma_p * g_n \sigma_p * g_n \sigma_p * g_n \sigma_p, g_n \sigma_p * g_n \sigma_p \rangle_{L^2}}{\|g_n\|_{L^2}^6}. \quad (6-7)
\end{aligned}$$

Since the sequence $\{g_n\}$ concentrates at $y_0 = 1$, we have

$$\lim_{n \rightarrow \infty} \langle g_n \sigma_p * g_n \sigma_p * g_n \sigma_p * g_n \sigma_p, g_n \sigma_p * g_n \sigma_p \rangle_{L^2} = 0.$$

Heuristically, $g_n \sigma_p * g_n \sigma_p$ is supported near the point $(2, 2)$, while $(g_n \sigma_p)^{(4)}$ is supported near the point $(4, 4)$, and so in the limit there is no contribution of the inner product. More precisely, given $\varepsilon > 0$, write $g_n = h_n + \kappa_n$, where $h_n := g_n \mathbb{1}_{[1-\varepsilon, 1+\varepsilon]}$ and $\|\kappa_n\|_{L^2}^2 \rightarrow 0$ as $n \rightarrow \infty$. If ε is small enough, then support considerations force

$$\langle h_n \sigma_p * h_n \sigma_p * h_n \sigma_p * h_n \sigma_p, h_n \sigma_p * h_n \sigma_p \rangle_{L^2} = 0 \quad \text{for every } n,$$

whereas the cross terms involve κ_n , whose L^2 -norm tends to zero as $n \rightarrow \infty$. We conclude

$$\limsup_{n \rightarrow \infty} \frac{\|f_n \mu_p * f_n \mu_p * f_n \mu_p\|_{L^2}^2}{\|f_n\|_{L^2}^6} = \frac{5}{2} \limsup_{n \rightarrow \infty} \frac{\|g_n \sigma_p * g_n \sigma_p * g_n \sigma_p\|_{L^2}^2}{\|g_n\|_{L^2}^6}, \quad (6-8)$$

and similarly for the limit inferior. [Lemma 3.2](#) applied to the sequence $\{g_n\}$ implies

$$\limsup_{n \rightarrow \infty} \frac{\|f_n \mu_p * f_n \mu_p * f_n \mu_p\|_{L^2(\mathbb{R}^2)}^2}{\|f_n\|_{L^2}^6} \leq \frac{5\pi}{\sqrt{3}p(p-1)}.$$

Moreover, equality holds if we take $f_n = g_n + \tilde{g}_n$, with $g_n := 2^{-1/2} h_n \|h_n\|_{L^2}^{-1}$, and

$$h_n(y) := e^{-n(|y|^p - 1 - p(y-1))} |y|^{\frac{p-2}{6}} \mathbb{1}_{[0,\infty)}(y).$$

Theorem 1.6 is now proved.

Remark 6.5. The invariant form of condition (1-16) in **Theorem 1.6** is

$$\left(\frac{Q_p}{C_2}\right)^6 > \frac{5}{p(p-1)}, \quad (6-9)$$

where $C_2^6 = \pi/\sqrt{3}$ is the best constant for the parabola in convolution form. In the case $p = 3$, a similar condition appears in [Shao 2009] on the Airy–Strichartz inequality, which translates into $(Q_3/C_2)^6 > \frac{1}{3}$. This is of course incompatible with (6-9) but, as was recently pointed out in [Frank and Sabin 2018, Remark 2.7], there is a problem in [Shao 2009, Lemma 6.1] in the passage from equation (89) to equation (90), as the argument presented there disregards the effect of symmetrization. On the other hand, the case $p = 3$ of (6-9) agrees with [Frank and Sabin 2018, Case $p = q = 6$ of Theorem 1], once the proper normalization is considered.

We now come to the question of whether extremizers for (1-15) actually exist, and discuss the case $1 < p < 2$ first. Just as in (4-17), set $g_n(y) := e^{-(u/n)n2(|y|^p - py)} |y|^{-(2-p)/6}$. Its even extension,

$$f_n := \frac{g_n \mathbb{1}_{[0,\infty)} + \tilde{g}_n \mathbb{1}_{(-\infty,0]}}{2^{\frac{1}{2}} \|g_n\|_{L^2(0,\infty)}},$$

can be used to establish the strict inequality in (1-16). One simply uses (6-8) together with the fact that the sequence $\{g_n \|g_n\|_{L^2}^{-1}\}_{s>0}$ concentrates at $y_0 = 1$, so that an argument similar to **Lemma 4.5** can be applied to the present case. Therefore, extremizers for (1-15) exist if $1 < p < 2$, and **Theorem 1.7** is now proved.

The case $p \geq 2$ seems harder. In view of (6-8), it is natural to use the methods of **Section 4** in order to find the series expansion for the trial functions $f = 2^{-1/2}(g + \tilde{g})$, where $g(y) = e^{-|y|^p} |y|^{(p-2)/6+a} \mathbb{1}_{[0,\infty)}(y)$ for different choices of a . By doing so, we find that we cannot reach the critical threshold $5\pi/(\sqrt{3}p(p-1))$, but that we can approach it from below by varying the value of a . We are led to the following conjecture.

Conjecture 6.6. For every $p \geq 2$,

$$\left(\frac{Q_p}{C_2}\right)^6 = \frac{5}{p(p-1)}.$$

Moreover, extremizers for (1-15) do not exist.

6A. On symmetric complex- and real-valued extremizers. The proof of **Lemma 6.1** merits some further remarks which we attempt to insert within a broader context.

First of all, identity (6-4) holds thanks to the symmetry with respect to the origin of both the curve $s = y|y|^{p-1}$ and the measure $d\mu_p = \delta(t - y|y|^{p-1})|y|^{(p-2)/6} dy ds$. In fact, the proof of **Lemma 6.1** immediately generalizes to the Fourier extension operator associated to any *antipodally symmetric pair* (Σ, μ) . By this we mean a set $\Sigma \subseteq \mathbb{R}^d$ (usually a smooth submanifold) together with a Borel measure

μ supported on Σ , both symmetric with respect to the origin in the sense that $T(\Sigma) = \Sigma$ and $T^* \mu = \mu$, where T denotes the antipodal map $T(y) = -y$ and $T^* \mu$ denotes the pushforward measure.

Secondly, the Lebesgue exponent 6 can be replaced with any finite exponent $r \geq 2$. More precisely, in the general context of an antipodally symmetric pair (Σ, μ) , if an estimate

$$\|\widehat{f\mu}\|_{L^r(\mathbb{R}^d)} \lesssim \|f\|_{L^2(\Sigma, \mu)} \quad (6-10)$$

does hold for some $r \in [2, \infty)$, then necessarily⁷

$$\sup_{\substack{0 \neq f \in L^2(\Sigma, \mu) \\ f \text{ real-valued}}} \frac{\|\widehat{f\mu}\|_{L^r(\mathbb{R}^d)}}{\|f\|_{L^2(\Sigma, \mu)}} = \sup_{\substack{0 \neq g \in L^2(\Sigma, \mu) \\ g \text{ real-valued, } g \text{ even or } g \text{ odd}}} \frac{\|\widehat{g\mu}\|_{L^r(\mathbb{R}^d)}}{\|g\|_{L^2(\Sigma, \mu)}}.$$

Thirdly, the discussion extends to the more general situation of complex-valued functions. For concreteness, let us specialize to the case of the unit sphere $\Sigma = \mathbb{S}^{d-1} \subseteq \mathbb{R}^d$, $d \geq 2$, equipped with its natural surface measure μ . Given an exponent $p \geq p_d := 2(d+1)/(d-1)$, the Tomas–Stein inequality states that

$$\|\widehat{(u\mu)}\|_{L^p(\mathbb{R}^d)} \lesssim_{p, d} \|u\|_{L^2(\mathbb{S}^{d-1})} \quad (6-11)$$

for every complex-valued function $u \in L^2(\mathbb{S}^{d-1})$. It is known [Fanelli et al. 2011; Frank et al. 2016] that complex-valued extremizers for (6-11) exist in the full range $p \geq p_d$, the endpoint existence in dimensions $d \geq 4$ being conditional on a celebrated conjecture concerning (1-2). Moreover, if $p \geq p_d$ is an even integer, then real-valued, even, nonnegative extremizers for (6-11) exist, by virtue of the equivalent convolution form; see [Christ and Shao 2012a; Foschi 2015; Shao 2016a]. Finally, if $p = \infty$, then one easily checks that the unique extremizers for (6-11) are the constant functions. For general $p \geq p_d$, $p \neq \infty$, we argue that the search for extremizers of (6-11) can be restricted to the class of complex-valued, symmetric functions. Indeed, write $u = f + ig$, with $f = \Re u$, $g = \Im u$. By reorganizing the summands, we may write $u = F + iG$, where $F = f_e + ig_o$ and $G = g_e - if_o$. The functions F, G are complex-valued and symmetric, in the sense that $F(y) = \overline{F(-y)}$ and $G(y) = \overline{G(-y)}$, for every $y \in \mathbb{S}^{d-1}$. Moreover, one easily checks that

$$F(y) = \frac{1}{2}(u(y) + \overline{u(-y)}), \quad G(y) = \frac{1}{2i}(u(y) - \overline{u(-y)}), \quad \|u\|_{L^2}^2 = \|F\|_{L^2}^2 + \|G\|_{L^2}^2,$$

and that, in view of the antipodal symmetry of the pair (\mathbb{S}^{d-1}, μ) , the functions $\widehat{F\mu}$, $\widehat{G\mu}$ are real-valued. Following the proof of Lemma 6.1, we are thus led to the following result.

Proposition 6.7. *Let $d \geq 2$ and $2(d+1)/(d-1) \leq p \leq \infty$. Then for every complex-valued $u \in L^2(\mathbb{S}^{d-1})$, $u \neq 0$, the following inequality holds:*

$$\frac{\|\widehat{(u\mu)}\|_{L^p(\mathbb{R}^d)}}{\|u\|_{L^2(\mathbb{S}^{d-1})}} \leq \sup_{0 \neq F \in L^2_{\text{sym}}(\mathbb{S}^{d-1})} \frac{\|\widehat{F\mu}\|_{L^p(\mathbb{R}^d)}}{\|F\|_{L^2(\mathbb{S}^{d-1})}}, \quad (6-12)$$

⁷Here, a real-valued function $g : \Sigma \rightarrow \mathbb{R}$ is naturally defined to be *even* (resp. *odd*) if $g(y) = g(-y)$ (resp. $g(y) = -g(-y)$) for μ -almost every point $y \in \Sigma$.

where $L_{\text{sym}}^2(\mathbb{S}^{d-1}) := \{F \in L^2(\mathbb{S}^{d-1}) : F(y) = \overline{F(-y)} \text{ for } \mu\text{-a.e. } y \in \mathbb{S}^{d-1}\}$. Moreover, if u realizes equality in (6-12), then there exist $F \in L_{\text{sym}}^2(\mathbb{S}^{d-1})$ and a constant $\kappa \in \mathbb{C}$ such that $u = \kappa F$, μ -a.e.

Proof. In light of the previous discussion, we can assume $p < \infty$, and only the last statement merits further justification. Suppose that u realizes equality in (6-12). In particular, u is a complex-valued extremizer for (6-11). Decompose $u = F + iG$ as before, with $F(y) = \frac{1}{2}(u(y) + \overline{u(-y)})$, $G = \frac{1}{2i}(u(y) - \overline{u(-y)})$, so that $F, G \in L_{\text{sym}}^2(\mathbb{S}^{d-1})$. If either $F \equiv 0$ or $G \equiv 0$, then there is nothing to prove, and so in what follows we assume F, G not to be identically zero. Following the proof of Lemma 6.1, we note that equality occurs in the application of the triangle inequality with respect to the $L^{p/2}(\mathbb{R}^d)$ -norm (recall that $p/2 > 1$ is finite) only if there exists $\lambda > 0$ such that⁸

$$|\widehat{F\mu}(\xi)| = \lambda |\widehat{G\mu}(\xi)| \quad \text{for every } \xi \in \mathbb{R}^d. \quad (6-13)$$

Subsequent cases of equality further imply

$$\frac{\|\widehat{(u\mu)}\|_{L^p(\mathbb{R}^d)}}{\|u\|_{L^2(\mathbb{S}^{d-1})}} = \frac{\|\widehat{F\mu}\|_{L^p(\mathbb{R}^d)}}{\|F\|_{L^2(\mathbb{S}^{d-1})}} = \frac{\|\widehat{G\mu}\|_{L^p(\mathbb{R}^d)}}{\|G\|_{L^2(\mathbb{S}^{d-1})}},$$

and so the functions F, G are also extremizers for (6-11). It suffices to show that $F = \kappa G$, where $\kappa \in \{-\lambda, \lambda\}$. Recall that $\widehat{F\mu}, \widehat{G\mu}$ are real-valued functions, since $F, G \in L_{\text{sym}}^2(\mathbb{S}^{d-1})$. Let $\xi_0 \in \mathbb{R}^d$ be such that $|\widehat{F\mu}(\xi_0)| \neq 0$. We lose no generality in assuming that $\widehat{F\mu}(\xi_0) > 0$ and $\widehat{G\mu}(\xi_0) > 0$, for otherwise we could replace F by $-F$ or G by $-G$. By continuity, there exists $r_0 > 0$ such that

$$\widehat{F\mu}(\xi + \xi_0) = \lambda \widehat{G\mu}(\xi + \xi_0) \quad \text{for every } |\xi| < r_0. \quad (6-14)$$

On the other hand, $\widehat{F\mu}(\xi + \xi_0) = \widehat{(e^{-iy \cdot \xi_0} F\mu)}(\xi)$ and $\widehat{G\mu}(\xi + \xi_0) = \widehat{(e^{-iy \cdot \xi_0} G\mu)}(\xi)$. The functions $e^{-iy \cdot \xi_0} F$ and $e^{-iy \cdot \xi_0} G$ belong to $L_{\text{sym}}^2(\mathbb{S}^{d-1})$, and may be expanded in the basis of spherical harmonics,

$$e^{-iy \cdot \xi_0} F = \sum_{n=0}^{\infty} \sum_{k=1}^{\gamma(d,n)} a_{n,k} Y_{n,k} \quad \text{and} \quad e^{-iy \cdot \xi_0} G = \sum_{n=0}^{\infty} \sum_{k=1}^{\gamma(d,n)} b_{n,k} Y_{n,k}. \quad (6-15)$$

Here, $\{Y_{n,k}\}_{k=1}^{\gamma(d,n)}$ denotes a basis for the space of spherical harmonics of degree n in the sphere \mathbb{S}^{d-1} , which has dimension $\gamma(d, n) := \binom{d+n-1}{n} - \binom{d+n-3}{n-2}$; see [Stein and Weiss 1971, Chapter IV]. The coefficients $a_{n,k}, b_{n,k}$ are complex numbers. Applying the Fourier transform to (6-15), we find that

$$\begin{aligned} \widehat{F\mu}(\xi + \xi_0) &= (2\pi)^{\frac{d}{2}} \sum_{n=0}^{\infty} \sum_{k=1}^{\gamma(d,n)} a_{n,k} i^{-n} |\xi|^{-\frac{d}{2}+1} J_{\frac{d}{2}-1+n}(|\xi|) Y_{n,k}\left(\frac{\xi}{|\xi|}\right), \\ \widehat{G\mu}(\xi + \xi_0) &= (2\pi)^{\frac{d}{2}} \sum_{n=0}^{\infty} \sum_{k=1}^{\gamma(d,n)} b_{n,k} i^{-n} |\xi|^{-\frac{d}{2}+1} J_{\frac{d}{2}-1+n}(|\xi|) Y_{n,k}\left(\frac{\xi}{|\xi|}\right). \end{aligned} \quad (6-16)$$

⁸As Fourier transforms of compactly supported distributions, both sides of (6-13) coincide with the absolute value of real-valued, *smooth* functions, so that the pointwise equality occurs at every point, and not just almost everywhere.

Using (6-14) and (6-16) together with the orthogonality of the functions $\{Y_{n,k}\}$ in $L^2(\mathbb{S}^{d-1})$, we obtain

$$a_{n,k} r^{-\frac{d}{2}+1} J_{\frac{d}{2}-1+n}(r) = \lambda b_{n,k} r^{-\frac{d}{2}+1} J_{\frac{d}{2}-1+n}(r) \quad \text{for every } r \in (0, r_0).$$

In particular, $a_{n,k} = \lambda b_{n,k}$. This and (6-15) together imply $F = \lambda G$. \square

A similar result to [Proposition 6.7](#) holds for a broader class of antipodally symmetric pairs (Σ, μ) . Indeed, let $r \in [2, \infty)$ be such that the extension estimate (6-10) holds. Then

$$\sup_{0 \neq u \in L^2(\Sigma, \mu)} \frac{\|\widehat{u\mu}\|_{L^r(\mathbb{R}^d)}}{\|u\|_{L^2(\Sigma, \mu)}} = \sup_{0 \neq F \in L^2_{\text{sym}}(\Sigma, \mu)} \frac{\|\widehat{F\mu}\|_{L^r(\mathbb{R}^d)}}{\|F\|_{L^2(\Sigma, \mu)}}, \quad (6-17)$$

with the obvious definition of $L^2_{\text{sym}}(\Sigma, \mu)$. Moreover, if μ is compactly supported and finite, then any complex extremizer u for (6-10) necessarily coincides with a multiple of a symmetric extremizer $F \in L^2_{\text{sym}}(\Sigma, \mu)$. Regarding the second part of [Proposition 6.7](#), the previous proof used the particular geometry of the sphere, but it can be modified to handle this more general situation. The crux of the matter is the fact that the Fourier transform of a compactly supported finite measure is real analytic. Indeed, if μ is a positive, compactly supported finite measure, and $F \in L^2(\Sigma, \mu)$, then, for every $\xi_0 \in \mathbb{R}^d$,

$$\begin{aligned} \widehat{F\mu}(\xi) &= \int_{\Sigma} e^{-i\xi \cdot y} F(y) d\mu(y) = \int_{\Sigma} e^{-i(\xi - \xi_0) \cdot y} e^{-i\xi_0 \cdot y} F(y) d\mu(y) \\ &= \sum_{k=0}^{\infty} \frac{(-i)^k}{k!} \int_{\Sigma} ((\xi - \xi_0) \cdot y)^k e^{-i\xi_0 \cdot y} F(y) d\mu(y), \end{aligned} \quad (6-18)$$

where the convergence is locally uniform. To see this, note the tail estimate

$$\left\| \sum_{k=K}^{\infty} \frac{(-i)^k}{k!} \int_{\Sigma} ((\xi - \xi_0) \cdot y)^k e^{-i\xi_0 \cdot y} F(y) d\mu(y) \right\|_{L_{\xi}^{\infty}(\Omega)} \leq \mu(\Sigma)^{\frac{1}{2}} \|F\|_{L^2(\Sigma, \mu)} \sum_{k=K}^{\infty} \frac{s^k}{k!},$$

which holds for every compact subset $\Omega \subseteq \mathbb{R}^d$ and every $K \in \mathbb{N}$. Here, $s = \sup_{\xi \in \Omega, y \in \Sigma} |\xi - \xi_0| |y| < \infty$. Therefore, the analogue of (6-13) in this setting leads to the corresponding (6-14), which by analyticity of (6-18) implies $\widehat{F\mu} = \lambda \widehat{G\mu}$, and therefore $F = \lambda G$.

These observations can be of interest when combined with the main result of [\[Fanelli et al. 2011\]](#), which states that complex-valued extremizers exist in the nonendpoint setting, provided μ is a positive, compactly supported finite measure. Important cases of antipodally symmetric pairs (Σ, μ) which have attracted recent attention include the aforementioned case of spheres, together with ellipsoids equipped with surface measure, and the double cone, the one- and the two-sheeted hyperboloids equipped with their natural Lorentz invariant measures; see [\[Foschi and Oliveira e Silva 2017\]](#).

We end this section with a final remark on the multiplier form of inequality (6-1). Consider the Cauchy problem

$$\begin{cases} \partial_t u - |\partial_x|^{p-1} \partial_x u = 0, & (x, t) \in \mathbb{R} \times \mathbb{R}, \\ u(\cdot, 0) = f \in L_x^2(\mathbb{R}), \end{cases} \quad (6-19)$$

whose solution can be written in terms of the propagator

$$u(x, t) = e^{t|\partial_x|^{p-1}\partial_x} f(x) = \frac{1}{2\pi} \int_{\mathbb{R}} e^{ix\xi} e^{it\xi|\xi|^{p-1}} \hat{f}(\xi) d\xi. \quad (6-20)$$

In view of (6-1), and more generally of [Kenig et al. 1991, Theorem 2.1], this satisfies the mixed norm estimate

$$\| |D|^{\frac{p-2}{r}} e^{t|\partial_x|^{p-1}\partial_x} f \|_{L_t^r L_x^s(\mathbb{R}^{1+1})} \lesssim_{r,s} \| f \|_{L^2(\mathbb{R})},$$

whenever the Lebesgue exponents r, s are such that $\frac{2}{r} + \frac{1}{s} = \frac{1}{2}$.

In this context, as noted in [Frank and Sabin 2018; Shao 2009] for the case $p = 3$, it makes sense to distinguish between real-valued and general complex-valued L^2 initial data. This is because the evolution $e^{t|\partial_x|^{p-1}\partial_x}$ preserves real-valuedness. In other words, if f is real-valued, then so is $e^{t|\partial_x|^{p-1}\partial_x} f$ for every $t \in \mathbb{R}$. In fact, if f is real-valued, then $\hat{f}(-\xi) = \hat{f}(\xi)$, and so taking the complex conjugate of (6-20) reveals that $\overline{u(x, t)} = u(x, t)$. The operator $|D|^{(p-2)/r} e^{t|\partial_x|^{p-1}\partial_x}$ is seen to preserve real-valuedness in a similar way.

It is then natural to consider the following family of sharp inequalities, for real- and complex-valued initial data and admissible Lebesgue exponents r, s :

$$\| |D|^{\frac{p-2}{r}} e^{t|\partial_x|^{p-1}\partial_x} u \|_{L_t^r L_x^s(\mathbb{R}^{1+1})} \leq M_{p,r,s}(\mathbb{C}) \| u \|_{L^2(\mathbb{R})}, \quad (6-21)$$

$$\| |D|^{\frac{p-2}{r}} e^{t|\partial_x|^{p-1}\partial_x} f \|_{L_t^r L_x^s(\mathbb{R}^{1+1})} \leq M_{p,r,s}(\mathbb{R}) \| f \|_{L^2(\mathbb{R})}, \quad (6-22)$$

where $u : \mathbb{R} \rightarrow \mathbb{C}$ is complex-valued and $f : \mathbb{R} \rightarrow \mathbb{R}$ is real-valued. The study of extremizers for (6-21)–(6-22) in the Airy–Strichartz case $p = 3$ has been considered in [Farah and Versieux 2018; Frank and Sabin 2018; Hundertmark and Shao 2012; Shao 2009]. It would be interesting to determine whether the methods developed in the present paper can be adapted to the study of extremizers for (6-21)–(6-22) in the mixed norm case $r \neq s$, so as to obtain an alternative approach to profile decomposition or the missing mass method. We do not pursue these matters here. However, we would still like to point out two interesting features of this problem which are easily derived from our previous analysis, and are the content of the following result.

Proposition 6.8. *Let $p > 1$, and $r, s \in (2, \infty)$ be such that $M_{p,r,s}(\mathbb{C})$ and $M_{p,r,s}(\mathbb{R})$ are finite. Then $M_{p,r,s}(\mathbb{C}) = M_{p,r,s}(\mathbb{R})$. Moreover, if a complex-valued extremizer u for $M_{p,r,s}(\mathbb{C})$ exists, then there exist $\kappa \in \mathbb{C}$ and a real-valued extremizer f for $M_{p,r,s}(\mathbb{R})$ such that $u = \kappa f$.*

The problem of the relationship between arbitrary complex-valued extremizers and real-valued extremizers has been considered in the literature; see, e.g., [Christ and Shao 2012b] for the case of the Tomas–Stein inequality on the sphere \mathbb{S}^2 . Note the duality with the second statement of Proposition 6.7 above.

Proof of Proposition 6.8. The equality $M_{p,r,s}(\mathbb{C}) = M_{p,r,s}(\mathbb{R})$ follows the same lines as the proof of Lemma 6.1. To see why this is the case, let $u \in L^2(\mathbb{R})$ and write $u = f + ig$, where f and g are the real and imaginary parts of u , and hence real-valued. Therefore

$$\|u\|_{L^2}^2 = \|f\|_{L^2}^2 + \|g\|_{L^2}^2, \quad (6-23)$$

$$\| |D|^{\frac{p-2}{r}} e^{t|\partial_x|^{p-1}\partial_x} u(x) \|^2 = \| |D|^{\frac{p-2}{r}} e^{t|\partial_x|^{p-1}\partial_x} f(x) \|^2 + \| |D|^{\frac{p-2}{r}} e^{t|\partial_x|^{p-1}\partial_x} g(x) \|^2 \quad (6-24)$$

for every $(x, t) \in \mathbb{R}^2$. If $r, s \geq 2$, then we can use the triangle inequality for the $L_x^{s/2}$ - and the $L_t^{r/2}$ -norms applied to (6-24), and obtain

$$\| |D|^{\frac{p-2}{r}} e^{t|\partial_x|^{p-1}\partial_x} u \|_{L_t^r L_x^s}^2 \leq \| |D|^{\frac{p-2}{r}} e^{t|\partial_x|^{p-1}\partial_x} f \|_{L_t^r L_x^s}^2 + \| |D|^{\frac{p-2}{r}} e^{t|\partial_x|^{p-1}\partial_x} g \|_{L_t^r L_x^s}^2. \quad (6-25)$$

Without loss of generality, assume that f, g are not identically zero. Reasoning as in the proof of Lemma 6.1 yields

$$\frac{\| |D|^{\frac{p-2}{r}} e^{t|\partial_x|^{p-1}\partial_x} u \|_{L_t^r L_x^s}^2}{\| u \|_{L^2}^2} \leq \max \left\{ \frac{\| |D|^{\frac{p-2}{r}} e^{t|\partial_x|^{p-1}\partial_x} f \|_{L_t^r L_x^s}^2}{\| f \|_{L^2}^2}, \frac{\| |D|^{\frac{p-2}{r}} e^{t|\partial_x|^{p-1}\partial_x} g \|_{L_t^r L_x^s}^2}{\| g \|_{L^2}^2} \right\} \quad (6-26)$$

and therefore $M_{p,r,s}(\mathbb{C}) \leq M_{p,r,s}(\mathbb{R})$. The reverse inequality is immediate. We gratefully acknowledge recent personal communication with R. Frank and J. Sabin [2018], who independently arrived at a similar conclusion.

We proceed to show that an arbitrary complex-valued extremizer for $M_{p,r,s}(\mathbb{C})$ necessarily coincides with a constant multiple of a real-valued extremizer for $M_{p,r,s}(\mathbb{R})$. Let $r, s \in (2, \infty)$, and suppose that u is a complex-valued extremizer for $M_{p,r,s}(\mathbb{C})$, which we express as the sum of its real and imaginary parts, $u = f + ig$. An inspection of the chain of inequalities leading to (6-26) shows that one of the following alternatives must hold:

- $g = 0$ and $u = f$ is a real-valued extremizer.
- $f = 0$, $u = ig$, and g is a real-valued extremizer.
- f, g are both not identically zero, and

$$\frac{\| |D|^{\frac{p-2}{r}} e^{t|\partial_x|^{p-1}\partial_x} f \|_{L_t^r L_x^s}^2}{\| f \|_{L^2}^2} = \frac{\| |D|^{\frac{p-2}{r}} e^{t|\partial_x|^{p-1}\partial_x} g \|_{L_t^r L_x^s}^2}{\| g \|_{L^2}^2} = M_{p,r,s}(\mathbb{R}), \quad (6-27)$$

so that f, g are real-valued extremizers.

It suffices to analyze the latter case. An inspection of the chain of inequalities leading to (6-25) shows that equality must hold in both applications of the triangle inequality. Since $r, s \in (2, \infty)$, this implies the existence of $\lambda > 0$ such that

$$\| |D|^{\frac{p-2}{r}} e^{t|\partial_x|^{p-1}\partial_x} f(x) \| = \lambda \| |D|^{\frac{p-2}{r}} e^{t|\partial_x|^{p-1}\partial_x} g(x) \| \quad \text{for almost every } (x, t) \in \mathbb{R}^2. \quad (6-28)$$

Equality in (6-27) then implies $\|f\|_{L^2} = \lambda \|g\|_{L^2}$. By squaring (6-28), and applying the Fourier transform, the equality of the resulting convolutions can be recast as

$$\begin{aligned} & \int_{\mathbb{R}^2} \hat{f}(y_1) \hat{f}(y_2) \delta(t - \psi(y_1) - \psi(y_2)) \delta(x - y_1 - y_2) |y_1 y_2|^{\frac{p-2}{r}} dy_1 dy_2 \\ &= \lambda^2 \int_{\mathbb{R}^2} \hat{g}(y_1) \hat{g}(y_2) \delta(t - \psi(y_1) - \psi(y_2)) \delta(x - y_1 - y_2) |y_1 y_2|^{\frac{p-2}{r}} dy_1 dy_2, \end{aligned} \quad (6-29)$$

where $(x, t) \in \mathbb{R}^2$ and $\psi(y) := y|y|^{p-1}$. Considering points (x, t) in the interior of the support of the convolution measure $\mu_p * \mu_p$, i.e., satisfying $t > 2\psi(\frac{1}{2}x)$ for $x > 0$, and $t < 2\psi(\frac{1}{2}x)$ for $x < 0$, we see

that there exists a unique positive solution $\alpha = \alpha(x, t) > 0$ of

$$t = \psi\left(\frac{1}{2}x - \alpha(x, t)\right) + \psi\left(\frac{1}{2}x + \alpha(x, t)\right), \quad (6-30)$$

and hence that the system of equations $t = \psi(y_1) + \psi(y_2)$, $x = y_1 + y_2$ has unique solutions

$$(y_1, y_2) \in \left\{ \left(\frac{1}{2}x - \alpha(x, t), \frac{1}{2}x + \alpha(x, t) \right), \left(\frac{1}{2}x + \alpha(x, t), \frac{1}{2}x - \alpha(x, t) \right) \right\}.$$

From (6-29) and a similar reasoning to that of [Oliveira e Silva and Quilodrán 2019, Proposition 2.1 and Remark 2.3], it then follows that

$$\hat{f}\left(\frac{1}{2}x - \alpha(x, t)\right)\hat{f}\left(\frac{1}{2}x + \alpha(x, t)\right) = \lambda^2 \hat{g}\left(\frac{1}{2}x - \alpha(x, t)\right)\hat{g}\left(\frac{1}{2}x + \alpha(x, t)\right)$$

for almost every $(x, t) \in \text{supp}(\mu_p * \mu_p)$. Alternatively, the latter identity follows by considering the analogue of formula (2-4) obtained in the case of even curves, which by the previous discussion applies to the present scenario as well. This yields

$$\hat{f}(x)\hat{f}(x') = \lambda^2 \hat{g}(x)\hat{g}(x') \quad (6-31)$$

for almost every $(x, x') \in \mathbb{R}^2$. As \hat{f}, \hat{g} belong to $L^2(\mathbb{R})$, we may integrate over any compact subset $I \subset \mathbb{R}$ in both variables x, x' and obtain

$$\left(\int_I \hat{f}(x) \, dx \right)^2 = \lambda^2 \left(\int_I \hat{g}(x) \, dx \right)^2. \quad (6-32)$$

Choose a compact subset $J \subset \mathbb{R}$ for which $\int_J \hat{g}(x) \, dx \neq 0$. From (6-32), we have

$$\int_J \hat{f}(x) \, dx = \lambda \int_J \hat{g}(x) \, dx \quad \text{or} \quad \int_J \hat{f}(x) \, dx = -\lambda \int_J \hat{g}(x) \, dx. \quad (6-33)$$

Integrating both sides of (6-31) over $x' \in J$, one infers from (6-33) that either $\hat{f} = \lambda \hat{g}$ or $\hat{f} = -\lambda \hat{g}$, and therefore that either $f = \lambda g$ or $f = -\lambda g$. The conclusion is that there exists $\lambda > 0$ such that either $u = (\lambda + i)g$ or $u = (-\lambda + i)g$, and so u is a constant multiple of a real-valued extremizer, as desired. \square

Appendix A: Concentration-compactness

This appendix consists of a useful observation regarding Lions' concentration-compactness lemma [1984a]. Let us start with some general considerations. Let (X, \mathcal{B}, μ) be a measure space with a *distinguished* point $\bar{x} \in X$ such that $\{\bar{x}\} \in \mathcal{B}$ and $\mu(\{\bar{x}\}) = 0$. Set $X_{\bar{x}} := X \setminus \{\bar{x}\}$. Let $\varrho : X_{\bar{x}} \times X_{\bar{x}} \rightarrow [0, \infty)$ be a pseudometric on $X_{\bar{x}}$, i.e., a measurable function on $X_{\bar{x}} \times X_{\bar{x}}$ satisfying $\varrho(x, x) = 0$, $\varrho(x, y) = \varrho(y, x)$, and $\varrho(x, y) \leq \varrho(x, z) + \varrho(z, y)$ for every $x, y, z \in X_{\bar{x}}$. Define the ball of center $x \in X_{\bar{x}}$ and radius $r \geq 0$, $B(x, r) := \{y \in X_{\bar{x}} : \varrho(x, y) \leq r\}$, and its complement $B(x, r)^c := X \setminus B(x, r)$. It is clear that

$$X_{\bar{x}} = \bigcup_{r \geq 0} B(x, r)$$

for every $x \neq \bar{x}$. We have the following concentration-compactness result, which should be compared to [Lions 1984a, Lemma I.1].

Proposition A.1. *Let $(X, \mathcal{B}, \mu), \bar{x} \in X, \varrho : X_{\bar{x}} \times X_{\bar{x}} \rightarrow [0, \infty)$ be as above. Let $\{\rho_n\}$ be a sequence in $L^1(X, \mu)$ satisfying*

$$\rho_n \geq 0 \quad \text{in } X, \quad \int_X \rho_n \, d\mu = \lambda,$$

where $\lambda > 0$ is fixed. Then there exists a subsequence $\{\rho_{n_k}\}$ satisfying one of the following three possibilities:

(i) (compactness) *There exists $\{x_k\} \subset X_{\bar{x}}$ such that $\rho_{n_k}(\cdot + x_k)$ is tight; i.e.,*

$$\text{for all } \varepsilon > 0, \text{ there exists } R < \infty \text{ such that } \int_{B(x_k, R)} \rho_{n_k} \, d\mu \geq \lambda - \varepsilon.$$

(ii) (vanishing) $\lim_{k \rightarrow \infty} \sup_{y \in X_{\bar{x}}} \int_{B(y, R)} \rho_{n_k} \, d\mu = 0$ for all $R < \infty$;

(iii) (dichotomy) *There exists $\alpha \in (0, \lambda)$ with the following property. For every $\varepsilon > 0$, there exist $R \in [0, \infty)$, $k_0 \geq 1$, and nonnegative functions $\rho_{k,1}, \rho_{k,2} \in L^1(X, \mu)$ such that, for every $k \geq k_0$,*

$$\|\rho_{n_k} - (\rho_{k,1} + \rho_{k,2})\|_{L^1(X)} \leq \varepsilon, \quad \left| \int_X \rho_{k,1} \, d\mu - \alpha \right| \leq \varepsilon, \quad \left| \int_X \rho_{k,2} \, d\mu - (\lambda - \alpha) \right| \leq \varepsilon,$$

$$\text{supp}(\rho_{k,1}) \subseteq B(x_k, R) \quad \text{and} \quad \text{supp}(\rho_{k,2}) \subseteq B(x_k, R_k)^C$$

for certain sequences $\{x_k\} \subset X_{\bar{x}}$, $\{R_k\} \subset [0, \infty)$, with $R_k \rightarrow \infty$ as $k \rightarrow \infty$.

The proof of [Proposition A.1](#) parallels that of [\[Lions 1984a, Lemma I.1\]](#) and proceeds via analysis of the sequence of *concentration functions*

$$Q_n : [0, \infty) \rightarrow \mathbb{R}, \quad Q_n(t) := \sup_{x \in X_{\bar{x}}} \int_{B(x, t)} \rho_n \, d\mu.$$

The sequence $\{Q_n\}$ consists of nondecreasing, nonnegative, uniformly bounded functions on $[0, \infty)$ which satisfy $Q_n(t) \rightarrow \lambda$ as $t \rightarrow \infty$, since $\mu(\{\bar{x}\}) = 0$. Very briefly, the argument goes as follows. By the Helly selection principle, there exists a subsequence $\{n_k\} \subset \mathbb{N}$ and a nondecreasing, nonnegative function $Q : [0, \infty) \rightarrow \mathbb{R}$ such that $Q_{n_k}(t) \rightarrow Q(t)$ as $k \rightarrow \infty$ for every $t \geq 0$. Set $\alpha := \lim_{t \rightarrow \infty} Q(t) \in [0, \lambda]$, and note that:

- If $\alpha = 0$, then $Q \equiv 0$. This translates into the vanishing condition at once.
- If $\alpha = \lambda$, then compactness occurs.
- If $0 < \alpha < \lambda$, then dichotomy occurs. In this case, the functions $\rho_{k,1}, \rho_{k,2}$ are given by $\rho_{k,1} = \rho_{n_k} \mathbb{1}_{B(x_k, R)}$ and $\rho_{k,2} = \rho_{n_k} \mathbb{1}_{B(x_k, R_k)^C}$.

We omit further details and refer the interested reader to [\[Lions 1984a\]](#).

When applying [Proposition A.1](#) to the study of extremizing sequences for (1-9), the desirable outcome (with a view towards obtaining concentration at a point under the hypotheses of [Proposition 3.3](#)) is *compactness* or *vanishing*. Therefore the possibility of *dichotomy* needs to be discarded. To this end,

Lions proposes the *strict superadditivity condition* [Lions 1984a, Section I.2], which in the present setting can be recast as follows. Define

$$I_\lambda := \sup\{\|\mathcal{E}_p(f)\|_{L^6(\mathbb{R}^2)}^6 : \|f\|_{L^2(\mathbb{R})}^2 = \lambda\}. \quad (\text{A-1})$$

The quantity I_λ is said to satisfy the strict superadditivity condition if, for every $\lambda > 0$,

$$I_\lambda > I_\alpha + I_{\lambda-\alpha} \quad \text{for every } \alpha \in (0, \lambda). \quad (\text{A-2})$$

In our case, \mathcal{E}_p is a linear operator, and so $I_\lambda = \lambda^3 I_1 = \lambda^3 E_p^6$. Thus (A-2) translates into the elementary numerical inequality $\lambda^3 > \alpha^3 + (\lambda - \alpha)^3$, which holds for every $\lambda > 0$ and $\alpha \in (0, \lambda)$. As seen in the proof of Proposition 3.3, it is condition (A-2) (applied with $\lambda = 1$) which ensures that dichotomy does not occur. A similar condition in a more general context is used in [Lieb 1983, Lemma 2.7].

Appendix B: Revisiting Brézis–Lieb

In this appendix, we prove a useful variant of [Fanelli et al. 2011, Proposition 1.1], which in turn relies on the Brézis–Lieb lemma [1983]. Proposition 1.1 of [Fanelli et al. 2011] states that, in the compact setting, the only obstruction to the strong convergence of an extremizing sequence is weak convergence to zero. In the noncompact setting, it is in general nontrivial to verify condition (iv) of [Fanelli et al. 2011, Proposition 1.1]. To overcome this difficulty, various arguments using Sobolev embeddings and the Rellich–Kondrachov compactness theorem have been employed in [Carneiro et al. 2019; Fanelli et al. 2012; Quilodrán 2013]. In our case, it is not clear how such an argument would go. Instead we take a different route, and argue that condition (iv) from [Fanelli et al. 2011, Proposition 1.1] can be replaced by uniform decay of the L^2 -norm, in a sense compactifying the space in question. The following is a precise formulation of this idea.

Proposition B.1. *Given $p > 1$, consider the Fourier extension operator $\mathcal{E}_p: L^2(\mathbb{R}) \rightarrow L^6(\mathbb{R}^2)$ defined in (1-12). Let $\{f_n\} \subset L^2(\mathbb{R})$, and let $\Theta: [1, \infty) \rightarrow (0, \infty)$ with $\Theta(R) \rightarrow 0$, as $R \rightarrow \infty$, be such that*

- (i) $\|f_n\|_{L^2(\mathbb{R})} = 1$ for every $n \in \mathbb{N}$,
- (ii) $\lim_{n \rightarrow \infty} \|\mathcal{E}_p(f_n)\|_{L^6(\mathbb{R}^2)} = E_p$,
- (iii) $f_n \rightharpoonup f \neq 0$ as $n \rightarrow \infty$,
- (iv) $\|f_n\|_{L^2([-R, R]^2)} \leq \Theta(R)$ for every $n \in \mathbb{N}$ and $R \geq 1$.

Then $f_n \rightarrow f$ in $L^2(\mathbb{R})$, as $n \rightarrow \infty$. In particular, $\|f\|_{L^2(\mathbb{R})} = 1$ and $\|\mathcal{E}_p(f)\|_{L^6(\mathbb{R}^2)} = E_p$, and so f is an extremizer of (1-9).

This variant was already observed in [Quilodrán 2012, Proposition 2.31] for the case of the cone, and the proof follows similar lines to that of [Fanelli et al. 2011, Proposition 1.1]. Note that the function Θ may depend on the sequence $\{f_n\}$, but *not* on n . The following proof is inspired by [Frank et al. 2016, Proposition 2.2].

Proof of Proposition B.1. Set $r_n := f_n - f$. Then $r_n \rightarrow 0$ as $n \rightarrow \infty$, and thus $m := \lim_{n \rightarrow \infty} \|r_n\|_{L^2}^2$ exists and satisfies $1 = \|f\|_{L^2}^2 + m$. Given $R > 0$, take the decomposition

$$r_n = r_n \mathbb{1}_{[-R, R]} + r_n \mathbb{1}_{[-R, R]^C} =: r_{n,1} + r_{n,2}.$$

Since the support of $r_{n,1}$ is compact and $r_{n,1} \rightarrow 0$ as $n \rightarrow \infty$, we know $\mathcal{E}_p(r_{n,1}) \rightarrow 0$ pointwise a.e. in \mathbb{R}^2 as $n \rightarrow \infty$. On the other hand, from condition (iv) we have

$$\|\mathcal{E}_p(r_{n,2})\|_{L^6} \leq E_p(\Theta(R) + \|f\|_{L^2([-R, R]^C)}) \quad (\text{B-1})$$

for every $R \geq 1$. This upper bound is independent of n , and tends to 0 as $R \rightarrow \infty$. We have $\mathcal{E}_p(f_n - r_{n,2}) = \mathcal{E}_p(f) + \mathcal{E}_p(r_{n,1})$, and $\|\mathcal{E}_p(f_n - r_{n,2})\|_{L^6} \leq E_p(1 + \Theta(R) + \|f\|_{L^2([-R, R]^C)})$ is uniformly bounded in n . Since $\mathcal{E}_p(f_n - r_{n,2}) \rightarrow \mathcal{E}_p(f)$ pointwise a.e. in \mathbb{R}^2 as $n \rightarrow \infty$, we can invoke the Brézis–Lieb lemma [1983] and obtain

$$\|\mathcal{E}_p(f_n - r_{n,2})\|_{L^6}^6 = \|\mathcal{E}_p(f)\|_{L^6}^6 + \|\mathcal{E}_p(r_{n,1})\|_{L^6}^6 + o(1) \quad \text{as } n \rightarrow \infty.$$

It follows that $\mu := \limsup_{n \rightarrow \infty} \|\mathcal{E}_p(r_{n,1})\|_{L^6}^6$ and $\lambda := \limsup_{n \rightarrow \infty} \|\mathcal{E}_p(f_n - r_{n,2})\|_{L^6}^6$ satisfy

$$\lambda = \|\mathcal{E}_p(f)\|_{L^6}^6 + \mu.$$

Since $\|\mathcal{E}_p(r_{n,1})\|_{L^6}^6 \leq E_p^6 \|r_{n,1}\|_{L^2}^6 \leq E_p^6 \|r_n\|_{L^2}^6$, we have $\mu \leq E_p^6 m^3$. Therefore

$$\lambda = \|\mathcal{E}_p(f)\|_{L^6}^6 + \mu \leq \|\mathcal{E}_p(f)\|_{L^6}^6 + E_p^6 (1 - \|f\|_{L^2}^2)^3.$$

Thus, replacing the definition of λ , we have proved

$$\limsup_{n \rightarrow \infty} \|\mathcal{E}_p(f_n - r_{n,2})\|_{L^6}^6 \leq \|\mathcal{E}_p(f)\|_{L^6}^6 + E_p^6 (1 - \|f\|_{L^2}^2)^3 \quad (\text{B-2})$$

for every $R \geq 1$. Now, $\|\mathcal{E}_p(f_n - r_{n,2})\|_{L^6} \geq \|\mathcal{E}_p(f_n)\|_{L^6} - \|\mathcal{E}_p(r_{n,2})\|_{L^6}$ and $\|\mathcal{E}_p(r_{n,2})\|_{L^6}$ is bounded above as quantified by (B-1). Thus

$$\limsup_{n \rightarrow \infty} \|\mathcal{E}_p(f_n - r_{n,2})\|_{L^6} \geq E_p - E_p(\Theta(R) + \|f\|_{L^2([-R, R]^C)})$$

for every $R \geq 1$. Using this together with (B-2), and letting $R \rightarrow \infty$, yields

$$E_p^6 \leq \|\mathcal{E}_p(f)\|_{L^6}^6 + E_p^6 (1 - \|f\|_{L^2}^2)^3.$$

By the elementary inequality $(1 - t)^3 \leq 1 - t^3$, valid for every $t \in [0, 1]$, we then have

$$E_p^6 \leq \|\mathcal{E}_p(f)\|_{L^6}^6 + E_p^6 (1 - \|f\|_{L^2}^6).$$

Since the reverse inequality holds by definition, we conclude that f is an extremizer. Moreover, since $f \neq 0$ and the elementary inequality is strict unless $t \in \{0, 1\}$, we conclude that $\|f\|_{L^2} = 1$. This completes the proof of the proposition. \square

Acknowledgements

We are grateful to Rupert Frank and Julien Sabin for discussions surrounding symmetric complex- and real-valued extremizers, and to Rupert Frank for clarifying some historical remarks.

References

- [Bégout and Vargas 2007] P. Bégout and A. Vargas, “Mass concentration phenomena for the L^2 -critical nonlinear Schrödinger equation”, *Trans. Amer. Math. Soc.* **359**:11 (2007), 5257–5282. [MR](#) [Zbl](#)
- [Bennett et al. 2009] J. Bennett, N. Bez, A. Carbery, and D. Hundertmark, “Heat-flow monotonicity of Strichartz norms”, *Anal. PDE* **2**:2 (2009), 147–158. [MR](#) [Zbl](#)
- [Bennett et al. 2015] J. Bennett, N. Bez, and M. Iliopoulou, “Flow monotonicity and Strichartz inequalities”, *Int. Math. Res. Not.* **2015**:19 (2015), 9415–9437. [MR](#) [Zbl](#)
- [Brézis and Lieb 1983] H. Brézis and E. Lieb, “A relation between pointwise convergence of functions and convergence of functionals”, *Proc. Amer. Math. Soc.* **88**:3 (1983), 486–490. [MR](#) [Zbl](#)
- [Carneiro and Oliveira e Silva 2015] E. Carneiro and D. Oliveira e Silva, “Some sharp restriction inequalities on the sphere”, *Int. Math. Res. Not.* **2015**:17 (2015), 8233–8267. [MR](#) [Zbl](#)
- [Carneiro et al. 2017] E. Carneiro, D. Foschi, D. Oliveira e Silva, and C. Thiele, “A sharp trilinear inequality related to Fourier restriction on the circle”, *Rev. Mat. Iberoam.* **33**:4 (2017), 1463–1486. [MR](#) [Zbl](#)
- [Carneiro et al. 2019] E. Carneiro, D. Oliveira e Silva, and M. Sousa, “Extremizers for Fourier restriction on hyperboloids”, *Ann. Inst. H. Poincaré Anal. Non Linéaire* **36**:2 (2019), 389–415. [MR](#) [Zbl](#)
- [Christ and Quilodrán 2014] M. Christ and R. Quilodrán, “Gaussians rarely extremize adjoint Fourier restriction inequalities for paraboloids”, *Proc. Amer. Math. Soc.* **142**:3 (2014), 887–896. [MR](#) [Zbl](#)
- [Christ and Shao 2012a] M. Christ and S. Shao, “Existence of extremals for a Fourier restriction inequality”, *Anal. PDE* **5**:2 (2012), 261–312. [MR](#) [Zbl](#)
- [Christ and Shao 2012b] M. Christ and S. Shao, “On the extremizers of an adjoint Fourier restriction inequality”, *Adv. Math.* **230**:3 (2012), 957–977. [MR](#) [Zbl](#)
- [Dodson et al. 2018] B. Dodson, J. L. Marzuola, B. Pausader, and D. P. Spirn, “The profile decomposition for the hyperbolic Schrödinger equation”, *Illinois J. Math.* **62**:1-4 (2018), 293–320. [MR](#) [Zbl](#)
- [Erdoğan et al. 2011] M. B. Erdoğan, D. Hundertmark, and Y.-R. Lee, “Exponential decay of dispersion managed solitons for vanishing average dispersion”, *Math. Res. Lett.* **18**:1 (2011), 11–24. [MR](#) [Zbl](#)
- [Fanelli et al. 2011] L. Fanelli, L. Vega, and N. Visciglia, “On the existence of maximizers for a family of restriction theorems”, *Bull. Lond. Math. Soc.* **43**:4 (2011), 811–817. [MR](#) [Zbl](#)
- [Fanelli et al. 2012] L. Fanelli, L. Vega, and N. Visciglia, “Existence of maximizers for Sobolev–Strichartz inequalities”, *Adv. Math.* **229**:3 (2012), 1912–1923. [MR](#) [Zbl](#)
- [Farah and Versieux 2018] L. G. Farah and H. Versieux, “On the lack of compactness and existence of maximizers for some Airy–Strichartz inequalities”, *Differential Integral Equations* **31**:1-2 (2018), 27–56. [MR](#) [Zbl](#)
- [Foschi 2007] D. Foschi, “Maximizers for the Strichartz inequality”, *J. Eur. Math. Soc.* **9**:4 (2007), 739–774. [MR](#) [Zbl](#)
- [Foschi 2015] D. Foschi, “Global maximizers for the sphere adjoint Fourier restriction inequality”, *J. Funct. Anal.* **268**:3 (2015), 690–702. [MR](#) [Zbl](#)
- [Foschi and Oliveira e Silva 2017] D. Foschi and D. Oliveira e Silva, “Some recent progress on sharp Fourier restriction theory”, *Anal. Math.* **43**:2 (2017), 241–265. [MR](#) [Zbl](#)
- [Frank and Sabin 2018] R. L. Frank and J. Sabin, “Extremizers for the Airy–Strichartz inequality”, *Math. Ann.* **372**:3-4 (2018), 1121–1166. [MR](#) [Zbl](#)
- [Frank et al. 2016] R. L. Frank, E. H. Lieb, and J. Sabin, “Maximizers for the Stein–Tomas inequality”, *Geom. Funct. Anal.* **26**:4 (2016), 1095–1134. [MR](#) [Zbl](#)

- [Gonçalves 2019] F. Gonçalves, “Orthogonal polynomials and sharp estimates for the Schrödinger equation”, *Int. Math. Res. Not.* **2019**:8 (2019), 2356–2383. [MR](#) [Zbl](#)
- [Hundertmark and Shao 2012] D. Hundertmark and S. Shao, “Analyticity of extremizers to the Airy–Strichartz inequality”, *Bull. Lond. Math. Soc.* **44**:2 (2012), 336–352. [MR](#) [Zbl](#)
- [Hundertmark and Zharnitsky 2006] D. Hundertmark and V. Zharnitsky, “On sharp Strichartz inequalities in low dimensions”, *Int. Math. Res. Not.* **2006** (2006), art. id. 34080. [MR](#) [Zbl](#)
- [Jiang et al. 2010] J.-C. Jiang, B. Pausader, and S. Shao, “The linear profile decomposition for the fourth order Schrödinger equation”, *J. Differential Equations* **249**:10 (2010), 2521–2547. [MR](#) [Zbl](#)
- [Jiang et al. 2014] J. Jiang, S. Shao, and B. Stovall, “Linear profile decompositions for a family of fourth order Schrödinger equations”, preprint, 2014. [arXiv](#)
- [Kenig et al. 1991] C. E. Kenig, G. Ponce, and L. Vega, “Oscillatory integrals and regularity of dispersive equations”, *Indiana Univ. Math. J.* **40**:1 (1991), 33–69. [MR](#) [Zbl](#)
- [Lieb 1983] E. H. Lieb, “Sharp constants in the Hardy–Littlewood–Sobolev and related inequalities”, *Ann. of Math.* (2) **118**:2 (1983), 349–374. [MR](#) [Zbl](#)
- [Lions 1984a] P.-L. Lions, “The concentration-compactness principle in the calculus of variations: the locally compact case, I”, *Ann. Inst. H. Poincaré Anal. Non Linéaire* **1**:2 (1984), 109–145. [MR](#) [Zbl](#)
- [Lions 1984b] P.-L. Lions, “The concentration-compactness principle in the calculus of variations: the locally compact case, II”, *Ann. Inst. H. Poincaré Anal. Non Linéaire* **1**:4 (1984), 223–283. [MR](#) [Zbl](#)
- [Negro 2018] G. Negro, “A sharpened Strichartz inequality for the wave equation”, preprint, 2018. [arXiv](#)
- [Oliveira e Silva and Quilodrán 2018] D. Oliveira e Silva and R. Quilodrán, “On extremizers for Strichartz estimates for higher order Schrödinger equations”, *Trans. Amer. Math. Soc.* **370**:10 (2018), 6871–6907. [MR](#) [Zbl](#)
- [Oliveira e Silva and Quilodrán 2019] D. Oliveira e Silva and R. Quilodrán, “A comparison principle for convolution measures with applications”, *Math. Proc. Cambridge Philos. Soc.* (online publication June 2019).
- [Quilodrán 2012] R. Quilodrán, *On extremizers for adjoint Fourier restriction inequalities and a result in incidence geometry*, Ph.D. thesis, University of California, Berkeley, 2012, available at <https://search.proquest.com/docview/1081722960>.
- [Quilodrán 2013] R. Quilodrán, “On extremizing sequences for the adjoint restriction inequality on the cone”, *J. Lond. Math. Soc.* (2) **87**:1 (2013), 223–246. [MR](#) [Zbl](#)
- [Reed and Simon 1975] M. Reed and B. Simon, *Methods of modern mathematical physics, II: Fourier analysis, self-adjointness*, Academic Press, New York, 1975. [MR](#) [Zbl](#)
- [Segal 1976] I. Segal, “Space-time decay for solutions of wave equations”, *Adv. Math.* **22**:3 (1976), 305–311. [MR](#) [Zbl](#)
- [Shao 2009] S. Shao, “The linear profile decomposition for the Airy equation and the existence of maximizers for the Airy–Strichartz inequality”, *Anal. PDE* **2**:1 (2009), 83–117. [MR](#) [Zbl](#)
- [Shao 2016a] S. Shao, “On existence of extremizers for the Tomas–Stein inequality for S^1 ”, *J. Funct. Anal.* **270**:10 (2016), 3996–4038. [MR](#) [Zbl](#)
- [Shao 2016b] S. Shao, “On smoothness of extremizers of the Tomas–Stein inequality for S^1 ”, preprint, 2016. [arXiv](#)
- [Stein and Weiss 1971] E. M. Stein and G. Weiss, *Introduction to Fourier analysis on Euclidean spaces*, Princeton Math. Series **32**, Princeton Univ. Press, 1971. [MR](#) [Zbl](#)
- [Strichartz 1977] R. S. Strichartz, “Restrictions of Fourier transforms to quadratic surfaces and decay of solutions of wave equations”, *Duke Math. J.* **44**:3 (1977), 705–714. [MR](#) [Zbl](#)
- [Tao 2003] T. Tao, “A sharp bilinear restriction estimate for paraboloids”, *Geom. Funct. Anal.* **13**:6 (2003), 1359–1384. [MR](#) [Zbl](#)
- [Tomas 1975] P. A. Tomas, “A restriction theorem for the Fourier transform”, *Bull. Amer. Math. Soc.* **81** (1975), 477–478. [MR](#) [Zbl](#)
- [Wolff 2001] T. Wolff, “A sharp bilinear cone restriction estimate”, *Ann. of Math.* (2) **153**:3 (2001), 661–698. [MR](#) [Zbl](#)

GIANMARCO BROCCHI: g.brocchi@pgr.bham.ac.uk

School of Mathematics, University of Birmingham, Birmingham, United Kingdom

DIOGO OLIVEIRA E SILVA: d.oliveiraesilva@bham.ac.uk

School of Mathematics, University of Birmingham, Birmingham, United Kingdom

and

Hausdorff Center for Mathematics, Bonn, Germany

RENÉ QUILODRÁN: rquilodr@dim.uchile.cl

EDITORS

EDITOR-IN-CHIEF

Patrick Gérard

patrick.gerard@math.u-psud.fr

Université Paris Sud XI

Orsay, France

BOARD OF EDITORS

Massimiliano Berti	Scuola Intern. Sup. di Studi Avanzati, Italy berti@sissa.it	Gilles Pisier	Texas A&M University, and Paris 6 pisier@math.tamu.edu
Michael Christ	University of California, Berkeley, USA mchrist@math.berkeley.edu	Tristan Rivière	ETH, Switzerland riviere@math.ethz.ch
Charles Fefferman	Princeton University, USA cf@math.princeton.edu	Igor Rodnianski	Princeton University, USA irod@math.princeton.edu
Ursula Hamenstädt	Universität Bonn, Germany ursula@math.uni-bonn.de	Yum-Tong Siu	Harvard University, USA siu@math.harvard.edu
Vadim Kaloshin	University of Maryland, USA vadim.kaloshin@gmail.com	Terence Tao	University of California, Los Angeles, USA tao@math.ucla.edu
Herbert Koch	Universität Bonn, Germany koch@math.uni-bonn.de	Michael E. Taylor	Univ. of North Carolina, Chapel Hill, USA met@math.unc.edu
Izabella Laba	University of British Columbia, Canada ilaba@math.ubc.ca	Gunther Uhlmann	University of Washington, USA gunther@math.washington.edu
Richard B. Melrose	Massachusetts Inst. of Tech., USA rbm@math.mit.edu	András Vasy	Stanford University, USA andras@math.stanford.edu
Frank Merle	Université de Cergy-Pontoise, France Frank.Merle@u-cergy.fr	Dan Virgil Voiculescu	University of California, Berkeley, USA dvv@math.berkeley.edu
William Minicozzi II	Johns Hopkins University, USA minicozz@math.jhu.edu	Steven Zelditch	Northwestern University, USA zelditch@math.northwestern.edu
Clément Mouhot	Cambridge University, UK c.mouhot@dpmms.cam.ac.uk	Maciej Zworski	University of California, Berkeley, USA zworski@math.berkeley.edu
Werner Müller	Universität Bonn, Germany mueller@math.uni-bonn.de		

PRODUCTION

production@msp.org

Silvio Levy, Scientific Editor

See inside back cover or msp.org/apde for submission instructions.

The subscription price for 2020 is US \$340/year for the electronic version, and \$550/year (+\$60, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues from the last three years and changes of subscriber address should be sent to MSP.

Analysis & PDE (ISSN 1948-206X electronic, 2157-5045 printed) at Mathematical Sciences Publishers, 798 Evans Hall #3840, c/o University of California, Berkeley, CA 94720-3840, is published continuously online. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices.

APDE peer review and production are managed by EditFlow® from MSP.

PUBLISHED BY

 **mathematical sciences publishers**

nonprofit scientific publishing

<http://msp.org/>

© 2020 Mathematical Sciences Publishers

ANALYSIS & PDE

Volume 13 No. 2 2020

Regularity estimates for elliptic nonlocal operators BARTŁOMIEJ DYDA and MORITZ KASSMANN	317
On solvability and ill-posedness of the compressible Euler system subject to stochastic forces DOMINIC BREIT, EDUARD FEIREISL and MARTINA HOFMANOVÁ	371
Variable coefficient Wolff-type inequalities and sharp local smoothing estimates for wave equations on manifolds DAVID BELTRAN, JONATHAN HICKMAN and CHRISTOPHER D. SOGGE	403
On the Hölder continuous subsolution problem for the complex Monge–Ampère equation, II NGOC CUONG NGUYEN	435
The Calderón problem for the fractional Schrödinger equation TUHIN GHOSH, MIKKO SALO and GUNTHER UHLMANN	455
Sharp Strichartz inequalities for fractional and higher-order Schrödinger equations GIANMARCO BROCCHI, DIOGO OLIVEIRA E SILVA and RENÉ QUILODRÁN	477
A bootstrapping approach to jump inequalities and their applications MARIUSZ MIREK, ELIAS M. STEIN and PAVEL ZORIN-KRANICH	527
On the trace operator for functions of bounded \mathbb{A} -variation DOMINIC BREIT, LARS DIENING and FRANZ GMEINEDER	559
Optimal constants for a nonlocal approximation of Sobolev norms and total variation CLARA ANTONUCCI, MASSIMO GOBBINO, MATTEO MIGLIORINI and NICOLA PICENNI	595