

# ANALYSIS & PDE

Volume 13      No. 3      2020

NEIL LYALL AND ÁKOS MAGYAR

DISTANCE GRAPHS AND SETS OF POSITIVE UPPER DENSITY IN  
 $\mathbb{R}^d$





## DISTANCE GRAPHS AND SETS OF POSITIVE UPPER DENSITY IN $\mathbb{R}^d$

NEIL LYALL AND ÁKOS MAGYAR

We present a refinement and sharp extension of a result of Bourgain on finding configurations of  $k+1$  points in general position in measurable subset of  $\mathbb{R}^d$  of positive upper density whenever  $d \geq k+1$  to all proper  $k$ -degenerate distance graphs.

### 1. Introduction

**1.1. Background.** A result of Furstenberg, Katznelson, and Weiss [Furstenberg et al. 1990] states that if  $A \subseteq \mathbb{R}^2$  has positive upper Banach density, then its distance set  $\{|x - x'| : x, x' \in A\}$  contains all sufficiently large numbers. Recall that the *upper Banach density* of a measurable set  $A \subseteq \mathbb{R}^d$  is defined by

$$\delta^*(A) = \limsup_{N \rightarrow \infty} \sup_{t \in \mathbb{R}^d} \frac{|A \cap (t + Q_N)|}{|Q_N|}, \quad (1)$$

where  $|\cdot|$  denotes Lebesgue measure on  $\mathbb{R}^d$  and  $Q_N$  denotes the cube  $[-N/2, N/2]^d$ .

Note that the distance set of any set of positive Lebesgue measure in  $\mathbb{R}^d$  automatically contains all sufficiently small numbers (by for example the Lebesgue density theorem) and that it is easy to construct a set of positive upper density which does not contain a fixed distance by placing small balls centered on an appropriate square lattice.

This result was later reproved using Fourier analytic techniques by Bourgain [1986]. In fact he established the following more general result for all finite point configurations  $V = \{v_0, v_1, \dots, v_k\}$  with the property that  $\{v_1 - v_0, \dots, v_k - v_0\}$  forms a linearly independent collection of vectors in  $\mathbb{R}^d$ , namely for all nondegenerate simplices. In the sequel we shall refer to such point configurations as being in *general position*.

**Theorem 1** [Bourgain 1986]. *Let  $\Delta_k \subseteq \mathbb{R}^d$  be a fixed collection of  $k+1$  points in general position.*

*If  $A \subseteq \mathbb{R}^d$  has positive upper Banach density and  $d \geq k+1$ , then there exists a threshold  $\lambda_0 = \lambda_0(A, \Delta_k)$  such that  $A$  contains an isometric copy of  $\lambda \cdot \Delta_k$  for all  $\lambda \geq \lambda_0$ .*

Recall that a point configuration  $\Delta'_k$  is said to be an isometric copy of  $\lambda \cdot \Delta_k$  if there exists a bijection  $\phi : \Delta_k \rightarrow \Delta'_k$  such that  $|\phi(v) - \phi(w)| = \lambda|v - w|$  for all  $v, w \in \Delta_k$ .

Bourgain [1986] further demonstrated that no result along the lines of Theorem 1 can hold for configurations that contain any three points in arithmetic progression on a line, specifically showing that

---

Lyall was partially supported by grant NSF-DMS 1702411 and the Simons Foundation Collaboration Grant for Mathematicians 245792. Magyar was partially supported by grant NSF-DMS 1600840 and ERC-AdG 321104.

MSC2010: 11B30.

Keywords: distance graphs, uniformity norms, geometric Ramsey theory.

for any  $d \geq 1$  there are sets of positive upper Banach density in  $\mathbb{R}^d$  which do not contain an isometric copy of configurations of the form  $\{0, y, 2y\}$  with  $|y| = \lambda$  for all sufficiently large  $\lambda$ . However, Ziegler [2006] established the remarkable result that if  $A \subseteq \mathbb{R}^d$  with  $d \geq 2$  has positive upper density and  $V = \{0, v_1, \dots, v_n\} \subseteq \mathbb{R}^d$ , where  $n$  can notably be taken arbitrarily large with respect to  $d$ , then there does exist a threshold  $\lambda_0 = \lambda_0(A, V)$  such that  $A_\varepsilon$  contains an isometric copy of  $\lambda \cdot V$  for all  $\lambda \geq \lambda_0$  and any  $\varepsilon > 0$ , where  $A_\varepsilon$  denotes the  $\varepsilon$ -neighborhood of  $A$ .

Together these results may be viewed as initial results in *geometric Ramsey theory* where, roughly speaking, one shows that “large” but otherwise arbitrary sets necessarily contain certain geometric configurations. Recently there has been a number of results in this direction in various contexts; see [Bulinski 2018; Bennett et al. 2016; Iosevich and Parshall 2018]. The objective of this article is to present a common extension in the setting of measurable subsets of Euclidean spaces of positive upper Banach density, while simultaneously presenting a new approach to (and refinement of) Theorem 1 based on a simple notion of uniform distribution attached to an appropriate scale. For another instance of this new approach see [Lyall and Magyar 2018] where configurations of points that form the vertices of a rigid geometric square, and more generally the direct product of any two finite point configurations in general position, are addressed.

**1.2. Distance graphs and main result.** A distance graph  $\Gamma = \Gamma(V, E)$  is a connected finite graph with vertex set  $V$  contained in  $\mathbb{R}^d$  for some  $d \geq 1$ . We say that  $\Gamma$  is *k-degenerate* if each of its subgraphs contains a vertex with degree at most  $k$ ; that is, some vertex in the subgraph touches  $k$  or fewer of the subgraphs edges. It is thus straightforward to verify, by induction, that if a given graph is *k-degenerate*, then there exists an ordering of its vertex set  $V = \{v_0, v_1, \dots, v_n\}$  in such a way that  $|V_j| \leq k$  for all  $1 \leq j \leq n$ , where

$$V_j := \{v_i : (v_i, v_j) \in E \text{ with } 0 \leq i < j\} \quad (2)$$

denotes the set of predecessors of the vertex  $v_j$ . In this article we shall always assume that the vertices of any given *k-degenerate* graph have been ordered as such. The *degeneracy* of a graph is defined to be the smallest  $k$  for which it is *k-degenerate*. Finally, we shall refer to a distance graph as *proper* if for every  $1 \leq j \leq n$ , the set of vertices  $v_j \cup V_j$ , namely  $v_j$  together with its predecessors, are in general position.

Given a distance graph  $\Gamma = \Gamma(V, E)$  and  $\lambda > 0$  we will say that  $\Gamma' = \Gamma'(V', E')$  is *isometric* to  $\lambda \cdot \Gamma$  if there exists a bijection  $\phi : V \rightarrow V'$  such that  $(v, w) \in E$  if and only if  $(\phi(v), \phi(w)) \in E'$  and  $|\phi(v) - \phi(w)| = \lambda|v - w|$ , and say that  $\Gamma'$  is a  *$\delta$ -close isometric* copy of  $\lambda \cdot \Gamma$  if one has the additional “angular closeness” property that

$$\frac{(\phi(v) - \phi(w)) \cdot (v - w)}{|\phi(v) - \phi(w)||v - w|} > 1 - \delta \quad (3)$$

for all  $(v, w) \in E$ . Note that if  $\delta > 2$  then a  *$\delta$ -close isometric* copy is merely an isometric copy. Finally, we say that  $A \subseteq \mathbb{R}^d$  contains a distance graph  $\Gamma = \Gamma(V, E)$  if  $V \subseteq A$ .

The main result of this article is the following:

**Theorem 2.** *Let  $\Gamma = \Gamma(V, E)$  be a proper  $k$ -degenerate distance graph and  $\delta > 0$ :*

- (i) *If  $A \subseteq \mathbb{R}^d$  has positive upper Banach density and  $d \geq k + 1$ , then there exists  $\lambda_0 = \lambda_0(A, \Gamma, \delta)$ , which tends to infinity as  $\delta \rightarrow 0^+$ , such that  $A$  contains a  $\delta$ -close isometric copy of  $\lambda \cdot \Gamma$  for all  $\lambda \geq \lambda_0$ .*

(ii) If  $A \subseteq [0, 1]^d$  with  $|A| > 0$  and  $d \geq k + 1$ , then  $A$  will contain a  $\delta$ -close isometric copy of  $\lambda \cdot \Gamma$  for all  $\lambda$  in some interval of length at least  $\exp(-C_{\Gamma, \delta} |A|^{-C|V|})$ , with  $C_{\Gamma, \delta}$  tending to infinity as  $\delta \rightarrow 0^+$ .

Intuitively one should visualize a distance graph with edges made of rigid rods which can freely turn around the vertices. One should further visualize an isometric copy of a distance graph in a set  $A \subseteq \mathbb{R}^d$  as a folding of the graph so that all of its vertices are supported on  $A$ , and a  $\delta$ -close isometric copy of a distance graph in a set  $A \subseteq \mathbb{R}^d$ , with  $\delta > 0$  and small, as a suitably small perturbation of the graph so that all of its vertices are supported on  $A$ .


Part (i) of Theorem 2 already constitutes a refinement of Theorem 1 when  $\Gamma$  is simply taken to be a complete distance graph on  $(k + 1)$  vertices in general position and  $\delta > 0$  is taken sufficiently small. In this special case it establishes that positive upper density subsets of  $\mathbb{R}^d$  not only contain an isometric copy of all sufficiently large dilates of a given nondegenerate simplex, as already guaranteed by Theorem 1, but that these copies can in fact be found as sufficiently large dilates of a “small rotation” of the original simplex. We further note that in both parts of Theorem 2 the dimension  $d$  is restricted only by the “level of degeneracy” of the given distance graph and not on the number of its vertices which could in fact be arbitrarily large. It is important to further observe that the length of the interval of dilations guaranteed by Part (ii) of Theorem 2 depends only on the measure of  $A$  and not on the set  $A$  itself.

Allowing the edges to rotate around the vertices is essential in our arguments. For example, the authors are unaware of any proof that there are  $k$ -equally spaced points along a line in a subset of positive density of  $\mathbb{R}^2$ , with *arbitrary* large gaps that does not invoke Szemerédi’s theorem [1975], and that such a result is in fact not possible for *all* sufficiently large gaps<sup>1</sup>. The reason being that the linear relations between the points of the pattern are no longer there when we allow for rotations of the edges around the vertices. A crucial observation of this note is that in this case the frequency of isometric copies in a given set is controlled by a simple norm, which may be viewed as a Euclidean analogue of the so-called  $U^1$ -seminorm [Tao and Vu 2006, Chapter 11], utilized in additive combinatorics. In the context of finite field geometries, a geometric analogue of the Gowers  $U^2$ -uniformity norm was developed in [Lyll et al. 2018] and used to prove that sets of positive density contain isometric copies of all circular quadrilaterals. We hope to address such problems for subsets of positive upper density of Euclidean spaces in the future.

As mentioned above, various special cases of our main result have been established, albeit in different contexts. Indeed, in [Bulinski 2018] the embedding of large copies of trees (1-degenerate distance graphs) was shown for dense subsets of the integer lattice. In [Bennett et al. 2016] it was shown that measurable subsets  $A \subseteq [0, 1]^d$  of Hausdorff dimension larger than  $(d + 1)/2$  contain an isometric copy of  $\lambda \cdot \Gamma$  for all  $\lambda$  in some interval, in the special case when  $\Gamma$  is a finite path. Very recently, parallel to our work, embedding of bounded degree distance graphs was addressed for subsets of vector spaces over finite fields [Iosevich and Parshall 2018].

*Examples of distance graphs.* (a) A nonempty connected graph is 1-degenerate if and only if it is a tree (contains no cycles). Any tree with vertices in  $\mathbb{R}^d$  with  $d \geq 1$  is isometric to a proper 1-degenerate distance graph in  $\mathbb{R}^2$ .

<sup>1</sup>For  $k = 3$  a result of this type was obtained in [Cook et al. 2017], with the Euclidean distance replaced by the  $\ell^p$ -distance, for all  $p \neq 2$ .



**Figure 1.** An example of a proper 2-degenerate distance graph in  $\mathbb{R}^3$

(b) Cycles with vertices in  $\mathbb{R}^d$  with  $d \geq 1$  form 2-degenerate distance graphs, but these are not necessarily isometric to a proper 2-degenerate distance graph in  $\mathbb{R}^d$  for any  $d \geq 1$ . Indeed, if  $V = \{0, 1, 2\} \subseteq \mathbb{R}$  and  $E = \{(0, 1), (1, 2), (0, 2)\}$ , then this defines just such a distance graph.

(c) If  $V = \{(i, j) : 0 \leq i, j \leq n\} \subseteq \mathbb{R}^2$  and  $E = \{((i, j), (i', j')) : |i - i'| + |j - j'| = 1\}$ , then this “2-dimensional grid” forms a proper 2-degenerate distance graph in  $\mathbb{R}^2$ .

In general, one can construct a proper 2-degenerate distance graph in  $\mathbb{R}^3$  as follows: Start with any proper cycle with vertices in  $\mathbb{R}^3$ , such as a proper triangle (three vertices in general position) or four vertices forming a “nonrigid” square (no diagonal edges), and at every step attach an edge (or vertex) of another proper cycle (or tree) to any of the edges (or vertices) of the graph constructed at the previous step. See Figure 1.

(d) A complete graph with vertices  $\{v_0, \dots, v_k\} \subseteq \mathbb{R}^k$  forms a proper  $k$ -degenerate distance graph if and only if  $\{v_0, \dots, v_k\}$  are in general position. Another example of a proper  $k$ -degenerate distance graph in  $\mathbb{R}^k$  is the “ $k$ -dimensional grid” with vertices  $V = \{(i_1, \dots, i_k) : 0 \leq i_1, \dots, i_k \leq n\} \subseteq \mathbb{R}^k$  and edges  $E = \{((i_1, \dots, i_k), (i'_1, \dots, i'_k)) : |i_1 - i'_1| + \dots + |i_k - i'_k| = 1\}$ .

More generally, one can construct a proper  $k$ -degenerate distance graph in  $\mathbb{R}^{k+1}$  as follows: Start with any known proper  $k$ -degenerate distance graph with vertices in  $\mathbb{R}^{k+1}$  and at every step attach another proper  $\ell$ -degenerate distance graph with  $\ell \leq k$  to any of the faces, edges, or vertices of the graph constructed at the previous step.

*Remark on the sharpness of the dimension condition in Theorem 2.* Let  $e_1, \dots, e_k$  be the standard basis vectors of  $\mathbb{R}^k$  and  $\Delta_+$  and  $\Delta_-$  denote the complete graphs with vertices  $\{0, e_1, e_2, \dots, e_k\}$  and  $\{0, -e_1, e_2, \dots, e_k\}$  respectively. It is clear that  $\Gamma = \Delta_+ \cup \Delta_-$  then defines a proper  $k$ -degenerate distance graph with the property that any isometric copy of  $\lambda \cdot \Gamma$  in  $\mathbb{R}^k$  must contain three collinear points, i.e., a copy of  $\{-\lambda e_1, 0, \lambda e_1\}$  obtained by a translation and a rotation. As mentioned above, it was shown in [Bourgain 1986] that there are sets of positive upper Banach density in  $\mathbb{R}^k$ , for any  $k$ , which do not contain such configurations for all large  $\lambda$ . This example shows the sharpness of the dimension condition  $d \geq k + 1$  in Theorem 2.

**1.3. Outline of the paper.** In Section 2 we introduce a norm which measures the uniformity of distribution with respect to a scale  $L$ . We prove that this norm controls the frequency with which isometric copies of

a given distance graph occur in a subset of the unit cube. This is analogous to the so-called von Neumann type inequalities in additive combinatorics; see for example [Tao and Vu 2006, Chapter 11].

In Section 3 we observe that sets of positive density are uniformly distributed with respect to sufficiently large scales which immediately implies Part (i) of Theorem 2. The proof of Part (ii) is also provided in Section 3 and based on a decomposition of a set into uniformly distributed parts.

Section 4 contains an alternative approach inspired by the argument in [Bourgain 1986]. We include this in order to highlight the simplicity and directness of our approach to Part (i) of Theorem 2, but also with the hope that this will serve to clarify Bourgain’s approach and emphasize that our approach to Part (ii) of Theorem 2 is in essence a physical space reinterpretation of Bourgain’s original.

**2. A counting function and generalized von Neumann inequality**

We now fix  $\delta > 0$  and let  $\Gamma = \Gamma(V, E)$  denote a fixed proper  $k$ -degenerate distance graph with vertex set  $V = \{v_0, v_1, \dots, v_n\}$  with  $v_0 = 0$  in  $\mathbb{R}^d$  with  $d \geq k + 1$ .

As our arguments are analytic, we need to define a measure on, or at least on a local piece of, the configuration space of all isometric copies of  $\Gamma$ . For each  $(v_i, v_j) \in E$  let  $t_{ij} = |v_i - v_j|^2$ . The configuration space of all isometric copies of  $\Gamma$ , with the vertex  $v_0$  remaining fixed at 0, namely

$$S_\Gamma := \{(x_0, x_1, \dots, x_n) \in \mathbb{R}^{d(n+1)} : x_0 = 0 \text{ and } |x_i - x_j|^2 = t_{ij} \text{ for all } i, j \text{ for which } (v_i, v_j) \in E\} \quad (4)$$

is clearly a real subvariety. We now proceed to give an equivalent description of the  $S_\Gamma$ .

For each point  $(x_0, x_1, \dots, x_n) \in \mathbb{R}^{d(n+1)}$  and  $1 \leq j \leq n$  we let

$$X_j := \{x_i : i \text{ has the property that } v_i \in V_j\},$$

with  $V_j = \{v_i : (v_i, v_j) \in E \text{ with } 0 \leq i < j\}$  as in (2) above. Moreover, for each  $(x_0, x_1, \dots, x_n) \in \mathbb{R}^{d(n+1)}$  and  $1 \leq j \leq n$ , we define the sets

$$S_j = S_j(X_j) := \{x \in \mathbb{R}^d : |x - x_i|^2 = t_{ij} \text{ for all } x_i \in X_j\}. \quad (5)$$

Note that for any finite set  $X_j$ ,  $S_j(X_j)$  is the intersection of  $|X_j|$  spheres and hence is itself either a sphere (of some dimension) or empty. It is thus easy to see that

$$(0, x_1, \dots, x_n) \in S_\Gamma \iff x_j \in S_j(X_j) \text{ for all } 1 \leq j \leq n \quad (6)$$

since both sides are defined by the same set of equations given in (4) above.

Since  $\Gamma$  is proper there exists a point  $(x_0, x_1, \dots, x_n) \in S_\Gamma$  with the property that the sets  $\bar{X}_j := \{x_j\} \cup X_j$  are in general position for all  $1 \leq j \leq n$ ; for example one could (and we will) take  $x_j = v_j$  for all  $0 \leq j \leq n$ . We shall refer to such points in  $S_\Gamma$  as *proper*. The following example illustrates that there may exist points of  $S_\Gamma$  that are not proper. Let  $V = \{v_0, v_1, v_2, v_3\} \subseteq \mathbb{R}^3$  and  $E = \{(v_0, v_1), (v_0, v_2), (v_1, v_3), (v_2, v_3)\}$ , where

$$v_0 = (0, 0, 0), \quad v_1 = (1, 0, 0), \quad v_2 = (0, 2, 0), \quad v_3 = (1, 2, 0).$$

Then  $\Gamma = \Gamma(V, E)$  is a proper 2-degenerate distance graph in  $\mathbb{R}^3$ . If we let  $x_0 = v_0$ ,  $x_1 = v_1$ ,  $x_2 = (2, 0, 0)$ , and  $x_3 = (3, 0, 0)$ , then it is easy to see that  $(x_0, x_1, x_2, x_3)$  is a point on  $S_\Gamma$  that is not proper since  $\bar{X}_3 = \{x_1, x_2, x_3\}$  is not in general position. It is however clear in this example, and in fact also true in general (although we do not need this fact), that “generic points” of  $S_\Gamma$  will always be proper.

The basic properties of the spheres  $S_j(X_j)$  and their geometric relationship with the sets  $X_j$ , specifically in the nontrivial case when  $|X_j| \geq 2$ , are collected in Lemma 3 below.

**Lemma 3.** *If  $(x_0, x_1, \dots, x_n)$  is a proper point of  $S_\Gamma$ , then for all  $1 \leq j \leq n$  for which we have  $|X_j| \geq 2$ :*

- (i) *The affine subspaces spanned by  $S_j(X_j)$  and  $X_j$  respectively are orthogonal and hence  $S_j(X_j)$  is a sphere of dimension  $d - |X_j|$ , which is at least 1 since  $|X_j| \leq k$  and  $d \geq k + 1$ .*
- (ii) *The radius  $r_j$  of the sphere  $S_j(X_j)$  is positive, equal to the distance from any point  $x \in S_j$  to the affine subspace spanned by  $X_j$ , and in fact depends continuously on the points in  $X_j$ .*

*Proof.* We first recall the elementary fact that the intersection of any two spheres in  $\mathbb{R}^3$  is either empty or a circle lying in a plane perpendicular to the line joining the centers of the spheres. Since any four points in  $\mathbb{R}^d$  with  $d \geq 3$  span a three-dimensional affine subspace, it follows immediately from the elementary fact above that if we let  $x \in S_j(X_j)$  and  $x_{i_1}$  and  $x_{i_2}$  be any two points in  $X_j$ , then the vectors  $x - x_j$  and  $x_{i_2} - x_{i_1}$  are orthogonal, and hence the affine subspaces spanned by  $S_j(X_j)$  and  $X_j$  respectively are orthogonal.

Let  $c_j$  denote the projection of  $x_j$  onto the affine subspace spanned by  $X_j$ . We claim that  $c_j$  is the center of the sphere  $S_j(X_j)$ . Indeed, since  $c_j$  has the property that the vectors  $x - c_j$  and  $x_i - c_j$  are orthogonal for all  $x \in S_j(X_j)$  and all  $x_i \in X_j$ , it follows that for any fixed  $x_i \in X_j$  we have  $|x_j - x_i|^2 = |x - x_i|^2$  for all  $x \in S_j(X_j)$ , and hence by Pythagoras that  $|x - c_j| = |x_j - c_j|$  for all  $x \in S_j(X_j)$ .

The discussion above implies that the radius  $r_j$  of  $S_j(X_j)$  is positive and equal to the distance from any point  $x \in S_j$ , so in particular  $x_j$ , to the affine subspace spanned by  $X_j$ . Specifically, if  $X_j = \{x_{i_1}, \dots, x_{i_\ell}\}$ , then the fact that  $\bar{X}_j = \{x_{i_1}, \dots, x_{i_\ell}, x_j\}$  is in general position ensures that the volume of the  $\ell$ -dimensional fundamental parallelotope determined by the vectors  $\{x_j - x_{i_1}, \dots, x_j - x_{i_\ell}\}$  is nonzero. It is a basic fact, see for example either Section 8.72 in [Shilov 1971] or Theorem 7 in Chapter X of [Birkhoff and Mac Lane 1941], that the volume of this parallelotope is equal to the square root of the so-called Gram determinant, namely the determinant of the (Gram) inner product matrix

$$\det\{(x_j - x_{i_{m_1}}) \cdot (x_j - x_{i_{m_2}})\}_{1 \leq m_1, m_2 \leq \ell}.$$

It thus follows that

$$r_j = \sqrt{\frac{\det\{(x_j - x_{i_{m_1}}) \cdot (x_j - x_{i_{m_2}})\}_{1 \leq m_1, m_2 \leq \ell}}{\det\{(x_{i_\ell} - x_{i_{m_1}}) \cdot (x_{i_\ell} - x_{i_{m_2}})\}_{1 \leq m_1, m_2 \leq \ell-1}}}, \tag{7}$$

as  $r_j$  is the height of our parallelotope if we take its base to be the  $(\ell - 1)$ -dimensional parallelotope determined by the vectors  $\{x_{i_\ell} - x_{i_1}, \dots, x_{i_\ell} - x_{i_{\ell-1}}\}$ ; see for example Section 8.72 in [Shilov 1971].

Since one can easily see, by expanding  $|(x_j - x_{i_{m_2}}) - (x_j - x_{i_{m_1}})|^2$ , that

$$\det\{(x_j - x_{i_{m_1}}) \cdot (x_j - x_{i_{m_2}})\}_{1 \leq m_1, m_2 \leq \ell} = \frac{1}{2}(t_{i_{m_1}j} + t_{i_{m_2}j} - |x_{i_{m_1}} - x_{i_{m_2}}|^2),$$

it follows that  $r_j$ , in addition to being positive, in fact depends continuously on the points in  $X_j$ . □

Finally we introduce compactly supported functions  $\eta_j \in C^\infty(\mathbb{R}^d)$  with  $0 \leq \eta_j \leq 1$  and  $\eta_j(v_j) = 1$  for each  $0 \leq j \leq n$ . We further assume that the supports of each  $\eta_j$  have been chosen small enough to ensure that every  $(x_0, x_1, \dots, x_n) \in S_\Gamma$  with  $x_j \in \text{supp } \eta_j$  corresponds to a  $\delta$ -close isometric copy of  $\Gamma$  with vertex  $v_0$  remaining fixed at 0.

An important consequence of Part (ii) of Lemma 3 above is that we may also assume that the supports of each  $\eta_j$  have been chosen small enough to ensure that there exists a constant  $r_\Gamma > 0$  such that for each  $(x_0, x_1, \dots, x_n) \in S_\Gamma$ , with  $x_j \in \text{supp } \eta_j$ , the corresponding spheres  $S_j(X_j)$  will all have radius  $r_j \geq r_\Gamma$ .

**Definition 4** (localized counting function). For any  $0 < \lambda \ll 1$  and functions

$$f_0, f_1, \dots, f_n : [0, 1]^d \rightarrow \mathbb{R},$$

with  $d \geq k + 1$ , we define

$$T_{\Gamma, \delta}(f_0, f_1, \dots, f_n)(\lambda) = \iint \dots \int f_0(x) f_1(x - \lambda x_1) \dots f_n(x - \lambda x_n) d\mu_n(x_n) \dots d\mu_1(x_1) dx, \quad (8)$$

where  $d\mu_j(x_j) = \eta_j(x_j) d\sigma_j(x_j)$  and  $\sigma_j$  denotes the normalized surface measure on  $S_j$ .

Note that if  $A \subseteq [0, 1]^d$  and  $T_{\Gamma, \delta}(1_A, 1_A, \dots, 1_A)(\lambda) > 0$ , then  $A$  must contain a point configuration  $\Gamma' = \{x, x + \lambda x_1, \dots, x + \lambda x_n\}$  with each  $x_j \in S_j(X_j)$ , and hence a  $\delta$ -close isometric copy of  $\lambda \cdot \Gamma$ .

The key to showing that  $T_{\Gamma, \delta}(1_A, 1_A, \dots, 1_A)(\lambda)$  is positive for certain sets  $A$  is to estimate (8) in terms of a suitable uniformity norm localized to a scale  $L$  (related to  $\lambda$ ).

**Definition 5** ( $U^1(L)$ -norm). For  $0 < L \ll 1$  and functions  $f : [0, 1]^d \rightarrow \mathbb{R}$  we define

$$\|f\|_{U^1(L)} = \|f * \varphi_L\|_2,$$

where  $\varphi_L(x) = L^{-d} \varphi(L^{-1}x)$ , with  $\varphi = 1_{[-1/2, 1/2]^d}$ .

Note that if  $A \subseteq [0, 1]^d$  with  $\alpha = |A| > 0$  and we define  $f_A := 1_A - \alpha 1_{[0, 1]^d}$ , then

$$\|f_A\|_{U^1(L)}^2 = \int_{\mathbb{R}^d} \left| \frac{|A \cap (t + Q_L)|}{|Q_L|} - \alpha \right|^2 dt, \quad (9)$$

where  $Q_L = [-L/2, L/2]^d$ .

Evidently the  $U^1(L)$ -norm is measuring the mean-square uniform distribution of  $A$  on scale  $L$ . The engine that drives our approach to Theorem 2 is the following:

**Proposition 6** (generalized von Neumann). *Let  $0 < \varepsilon, \lambda \ll 1$ . For any  $L \leq \varepsilon^6 \lambda$ ,  $0 \leq m \leq n$ , and functions*

$$f_0, f_1, \dots, f_m : [0, 1]^d \rightarrow [-1, 1],$$

*we have*

$$|T_{\Gamma, \delta}(f_0, f_1, \dots, f_m, 1, \dots, 1)(\lambda)| \leq \|f_m\|_{U^1(L)} + O_\Gamma(\varepsilon).$$

Here 1 stands for the indicator function of the unit cube  $[0, 1]^d$  and  $O_\Gamma(\varepsilon)$  means a quantity bounded by  $C_\Gamma \varepsilon$  with  $C_\Gamma$  a constant depending only on  $\Gamma$ . We will also use the notation  $f \ll_{\Gamma, \delta} g$  to indicate that  $|f| \leq c_{\Gamma, \delta} g$  with a constant  $c_{\Gamma, \delta} > 0$ , depending on only  $\Gamma$  and  $\delta$ , that is *sufficiently small* for our purposes.

The above proposition immediately implies the following result for uniformly distributed sets from which we will deduce both parts of Theorem 2 in Section 3 below.

**Corollary 7.** *Let  $\delta > 0$  and  $\Gamma$  be a proper  $k$ -degenerate distance graph on  $n + 1$  vertices in  $\mathbb{R}^d$  with  $d \geq k + 1$ .*

*Let  $\alpha \in (0, 1)$  and  $0 < \lambda \leq \varepsilon \ll_{\Gamma, \delta} \alpha^{n+1}$ . If  $A \subseteq [0, 1]^d$  with  $|A| = \alpha$  satisfies  $\|f_A\|_{U^1(\varepsilon^6\lambda)} \ll \varepsilon$ , then*

$$T_{\Gamma, \delta}(1_A, 1_A, \dots, 1_A)(\lambda) \geq \frac{1}{2}c_0\alpha^{n+1},$$

where

$$c_0 = c_0(\Gamma, \delta) = \iint \dots \int d\mu_n(x_n) \dots d\mu_1(x_1) dx.$$

Note that in light of the assumptions that we have placed on the measures  $\mu_j$  (via the functions  $\eta_j$ ), the quantity  $c_0(\Gamma, \delta)$  above clearly tends to zero as  $\delta \rightarrow 0^+$ .

*Proof of Corollary 7.* The result follows immediately from Proposition 6 since

$$T_{\Gamma, \delta}(1_A, \dots, 1_A)(\lambda) = c_0\alpha^{n+1} + \sum_{m=0}^n \alpha^{n-m} T_{\Gamma, \delta}(\underbrace{1_A, \dots, 1_A}_{m \text{ copies}}, f_A, 1, \dots, 1)(\lambda),$$

where  $f_A = 1_A - \alpha 1_{[0, 1]^d}$ . □

We conclude this section with the proof of Proposition 6.

*Proof of Proposition 6.* Fix  $0 \leq m \leq n$ . We have

$$\begin{aligned} &|T_{\Gamma, \delta}(f_0, f_1, \dots, f_m, 1, \dots, 1)(\lambda)| \\ &\leq \int \dots \int \left( \int \left| \int f_m(x - \lambda x_m) c_{m+1}(x_1, \dots, x_m) d\mu_m(x_m) \right| dx \right) d\mu_{m-1}(x_{m-1}) \dots d\mu_1(x_1), \end{aligned}$$

where

$$c_{m+1}(\Gamma, \delta; x_1, \dots, x_m) = \int \dots \int d\mu_n(x_n) \dots d\mu_{m+1}(x_{m+1}) \tag{10}$$

if  $0 \leq m \leq n - 1$  and  $c_{n+1} = 1$ . It follows from an application of Cauchy–Schwarz and Plancherel that

$$|T_{\Gamma, \delta}(f_0, f_1, \dots, f_m, 1, \dots, 1)(\lambda)|^2 \leq \int |\hat{f}_m(\xi)|^2 I_m(\lambda\xi) d\xi, \tag{11}$$

where

$$I_m(\xi) = \int \dots \int |\widehat{c_{m+1}\mu_m}(\xi)|^2 d\mu_{m-1}(x_{m-1}) \dots d\mu_1(x_1), \tag{12}$$

with

$$\widehat{c_{m+1}\mu_m}(\xi) = \int c_{m+1}(x_1, \dots, x_m) \eta_m(x_m) e^{-2\pi i x_m \cdot \xi} d\sigma_m(x_m)$$

if  $2 \leq m \leq n$  and  $I_1 = |\widehat{c_2\mu_1}|^2$ . In light of the trivial uniform bound  $0 \leq I_m(\xi) \leq 1$  and the fact that

$$\|f_m\|_{U^1(L)}^2 = \int |\hat{f}_m(\xi)|^2 |\hat{\varphi}(L\xi)|^2 d\xi,$$

it suffices to establish that

$$I_m(\lambda\xi)(1 - \hat{\varphi}(L\xi)^2) = O_{\Gamma}(\varepsilon^2). \tag{13}$$

Since  $0 \leq \hat{\varphi}(\xi)^2 \leq 1$  for all  $\xi \in \mathbb{R}^d$  and  $\hat{\varphi}(0) = 1$  it follows that  $0 \leq 1 - \hat{\varphi}(L\xi)^2 \leq \min\{1, 4\pi L|\xi|\}$ . The uniform bound (13) thus reduces to establishing the decay estimate

$$I_m(\xi) \leq \min\{1, C_\Gamma |\xi|^{-1/2}\} \tag{14}$$

since this would in turn imply that

$$I_m(\lambda\xi)(1 - \hat{\varphi}(L\xi)^2) \leq C_\Gamma \min\{(\lambda|\xi|)^{-1/2}, \varepsilon^6 \lambda|\xi|\} \leq C_\Gamma \varepsilon^2$$

whenever  $L \leq \varepsilon^6 \lambda$ .

To establish (14) we will use the fact that in addition to being trivially bounded by 1, the Fourier transform of  $c_{m+1}\mu_m$  also decays for large  $\xi$  in certain directions, specifically

$$|\widehat{c_{m+1}\mu_m}(\xi)| \leq \min\{1, (r_\Gamma \cdot (\text{dist}(\xi, \text{span } X_m))^{-1/2})\} \tag{15}$$

uniformly over all  $x_1, \dots, x_{m-1}$  with  $x_j \in \text{supp } \eta_j$ . This estimate is an easy consequence of the well-known asymptotic behavior of the Fourier transform of the measure on the unit sphere  $S^{d-|X_m|} \subseteq \mathbb{R}^{d-|X_m|+1}$  induced by Lebesgue measure; see for example [Stein 1993].

Using the fact that the measure  $d\sigma_{m-1}(x_{m-1}) \cdots d\sigma_1(x_1)$  is invariant under the rotation

$$(x_1, \dots, x_m) \rightarrow (Ux_1, \dots, Ux_m),$$

for any  $U \in \text{SO}(d)$ , together with (15) and the fact that  $0 \leq \eta_j \leq 1$  for  $1 \leq j \leq m$ , then gives

$$\begin{aligned} I_m(\xi) &\leq C \int \cdots \int (1 + r_\Gamma \cdot \text{dist}(\xi, \text{span } X_m))^{-1} d\sigma_{m-1}(x_{m-1}) \cdots d\sigma_1(x_1) \\ &= C \int \cdots \int_{\text{SO}(d)} (1 + r_\Gamma \cdot \text{dist}(\xi, \text{span } UX_m))^{-1} d\mu(U) d\sigma_{m-1}(x_{m-1}) \cdots d\sigma_1(x_1) \\ &= C \int \cdots \int_{S^{d-1}} (1 + r_\Gamma |\xi| \cdot \text{dist}(y, \text{span } X_m))^{-1} d\sigma(y) d\sigma_{m-1}(x_{m-1}) \cdots d\sigma_1(x_1), \end{aligned} \tag{16}$$

where  $\sigma$  denotes normalized measure on the unit sphere  $S^{d-1}$  in  $\mathbb{R}^d$  induced by Lebesgue measure. Estimate (14) then follows from the easy observation that the inner integral above satisfies the uniform estimate

$$\int_{S^{d-1}} (1 + r_\Gamma |\xi| \cdot \text{dist}(y, \text{span } X_m))^{-1} d\sigma(y) = O((1 + r_\Gamma |\xi|)^{-1/2}). \quad \square$$

### 3. Proof of Theorem 2

We will deduce Theorem 2 from Corollary 7 by localizing to cubes on which our set is suitably uniformly distributed. In the case of Part (i) this is achieved as a direct consequence of the definition of upper Banach density, while for Part (ii) this is achieved via an energy increment argument.

**3.1. Direct proof of Part (i) of Theorem 2.** Let  $\varepsilon > 0$  and  $A \subseteq \mathbb{R}^d$  with  $\delta^*(A) > 0$ .

The following two facts follow immediately from the definition of upper Banach density, see (1):

(i) There exist  $M_0 = M_0(A, \varepsilon)$  such that for all  $M \geq M_0$  and all  $t \in \mathbb{R}^d$

$$\frac{|A \cap (t + Q_M)|}{|Q_M|} \leq (1 + \frac{1}{3}\varepsilon^4) \delta^*(A).$$

(ii) There exist arbitrarily large  $N \in \mathbb{R}$  such that

$$\frac{|A \cap (t_0 + Q_N)|}{|Q_N|} \geq (1 - \frac{1}{3}\varepsilon^4) \delta^*(A)$$

for some  $t_0 \in \mathbb{R}^d$ .

Combining (i) and (ii) above we see that for any  $\lambda \geq \lambda_0 := \varepsilon^{-6}M_0$ , there exist  $N \geq \varepsilon^{-6}\lambda$  and  $t_0 \in \mathbb{R}^d$  such that

$$\frac{|A \cap (t + Q_{\varepsilon^6\lambda})|}{|Q_{\varepsilon^6\lambda}|} \leq (1 + \varepsilon^4) \frac{|A \cap (t_0 + Q_N)|}{|Q_N|}$$

for all  $t \in \mathbb{R}^d$ . Consequently, Theorem 2 reduces, via a rescaling of  $A \cap (t_0 + Q_N)$  to a subset of  $[0, 1]^d$ , to establishing that if  $\Gamma$  is a proper  $k$ -degenerate distance graph,  $0 < \lambda \leq \varepsilon \ll 1$ , and  $A \subseteq [0, 1]^d$  is measurable with  $|A| > 0$  and the property that

$$\frac{|A \cap (t + Q_{\varepsilon^6\lambda})|}{|Q_{\varepsilon^6\lambda}|} \leq (1 + \varepsilon^4) |A|$$

for all  $t \in \mathbb{R}^d$ , then  $A$  contains an isometric copy of  $\lambda \cdot \Gamma$ .

Now since  $A \cap (t + Q_{\varepsilon^6\lambda})$  is only supported in  $[-\varepsilon^6\lambda, 1 + \varepsilon^6\lambda]^d$  and

$$|A| = \int_{\mathbb{R}^d} \frac{|A \cap (t + Q_{\varepsilon^6\lambda})|}{|Q_{\varepsilon^6\lambda}|} dt$$

it easily follows that

$$\left| \left\{ t \in \mathbb{R}^d : 0 < \frac{|A \cap (t + Q_{\varepsilon^6\lambda})|}{|Q_{\varepsilon^6\lambda}|} \leq (1 - \varepsilon^2) |A| \right\} \right| = O(\varepsilon^2)$$

and hence that

$$\|f_A\|_{U^1(\varepsilon^6\lambda)}^2 = \int_{\mathbb{R}^d} \left| \frac{|A \cap (t + Q_{\varepsilon^6\lambda})|}{|Q_{\varepsilon^6\lambda}|} - |A| \right|^2 dt = O(\varepsilon^2).$$

The result thus follows from Corollary 7 above provided  $\varepsilon \ll_{\Gamma, \delta} \delta^*(A)^{n+1}$ . □

### 3.2. Proof of Part (ii) of Theorem 2.

**Lemma 8** (localization principle). *Let  $A \subseteq [0, 1]^d$  with  $d \geq k + 1$  and  $|A| = \alpha > 0$ .*

*Let  $\varepsilon > 0$  and  $\varepsilon^7 \gg L_1 \gg L_2 \gg \dots$  be any decreasing sequence with  $L_1^{-1} \in \mathbb{N}$  and  $L_{j+1} \leq c\varepsilon^7 L_j$  with  $L_{j+1} | L_j$  for all  $j \geq 1$ . If we let  $\mathcal{G}_j$  denote the partition of  $[0, 1]^d$  into cubes of side length  $L_j$ , then there exists  $1 \leq j \leq C\varepsilon^{-2}$  such that for all but at most  $\varepsilon L_j^{-d}$  of the cubes  $Q$  in  $\mathcal{G}_j$  the set  $A$  will be uniformly distributed on the smaller scale  $L_{j+1}$  inside  $Q$  in the sense that*

$$\frac{1}{|Q|} \int_Q \left| \frac{|A \cap Q \cap (t + Q_{L_{j+1}})|}{|Q_{L_{j+1}}|} - \frac{|A \cap Q|}{|Q|} \right|^2 dt \leq \varepsilon. \tag{17}$$

Before proving Lemma 8 we first show that it, together with Corollary 7 (after rescaling), is sufficient to establish Part (ii) of Theorem 2. Let  $\varepsilon \ll_{\Gamma, \delta} \alpha^{n+1}$  and  $\{Q_i\}$  denote the cubes of side length  $L_j$  in the partition  $\mathcal{G}_j$  of  $[0, 1]^d$  that we obtain from Lemma 8. If we then let  $A_i = A \cap Q_i$  and set  $\alpha_i = |A \cap Q_i|/|Q_i|$ , it follows from Corollary 7 (after rescaling) and Hölder’s inequality that for any  $\lambda \in (\varepsilon^{-6}L_{j+1}, \varepsilon L_j)$  we have

$$T_{\Gamma, \delta}(1_A, \dots, 1_A)(\lambda) \geq \sum_{i=1}^{L_j^{-d}} T_{\Gamma, \delta}(1_{A_i}, \dots, 1_{A_i})(\lambda) \geq \frac{1}{4}c_0 L_j^d \sum_{i=1}^{L_j^{-d}} \alpha_i^{n+1} \geq \frac{1}{4}c_0 \left( L_j^d \sum_{i=1}^{L_j^{-d}} \alpha_i \right)^{n+1} = \frac{1}{4}c_0 |A|^{n+1}. \tag{18}$$

*Proof of Lemma 8.* Let  $\{Q_i\}$  denote the cubes of side length  $L_j$  in the partition  $\mathcal{G}_j$  of  $[0, 1]^d$  and

$$g_j = 1_A - \mathbb{E}(1_A | \mathcal{G}_j),$$

where

$$\mathbb{E}(1_A | \mathcal{G}_j)(x) = \frac{|A \cap Q_i|}{|Q_i|}$$

for each  $x \in Q_i$ . If  $\|g_j\|_{U^1(L_{j+1})} \geq \varepsilon$ , then by definition

$$\int \left| \frac{1}{|Q_{L_{j+1}}|} \int_{x+Q_{L_{j+1}}} g_j(y) dy \right|^2 dx \geq c\varepsilon^2.$$

It follows that there must exist a  $x_0 \in [0, 1]^d$  for which the shifted grid  $x_0 + \mathcal{G}_{j+1}$  satisfies

$$\int |\mathbb{E}(g_j | x_0 + \mathcal{G}_{j+1})|^2 dx \geq c\varepsilon^2,$$

from which one can easily conclude that the (unshifted) refined grid  $\mathcal{G}_{j+2}$  satisfies

$$\int |\mathbb{E}(g_j | \mathcal{G}_{j+2})|^2 dx \geq c\varepsilon^2 \tag{19}$$

provided  $L_{j+2} \ll \varepsilon^2 L_{j+1}$ . By orthogonality, it follows immediately from (19) and the definition of  $g_j$  that

$$\|\mathbb{E}(1_A | \mathcal{G}_{j+2})\|_2^2 \geq \|\mathbb{E}(1_A | \mathcal{G}_j)\|_2^2 + c\varepsilon^2 \tag{20}$$

and hence that there must exist  $1 \leq j \leq C\varepsilon^{-2}$  such that  $\|g_j\|_{U^1(L_{j+1})} \leq \varepsilon$ . From this it follows that

$$\sum_{i=1}^{L_j^{-d}} \int \left| \frac{1}{|Q_{L_{j+1}}|} \int_{x+Q_{L_{j+1}}} (1_{A_i} - \alpha_i 1_{Q_i})(y) dy \right|^2 dx \leq C\varepsilon^2$$

provided  $L_{j+1} \ll \varepsilon^2 L_j$ . □

#### 4. A second proof of Theorem 2

Let  $\delta > 0$  and  $\Gamma$  be a proper  $k$ -degenerate distance graph in  $[0, 1]^d$  with  $d \geq k + 1$ . We shall make use of the same notation as in Section 2, specifically for the counting function  $T_{\Gamma, \delta}$  as defined in (8), and make the same assumptions on the measures  $\mu_j$  (via the functions  $\eta_j$ ).

**4.1. Reducing Theorem 2 to a dichotomy between randomness and structure.** As we shall see, Theorem 2 is an immediate consequence of the following proposition which reveals that if  $A \subseteq [0, 1]^d$  has positive measure but does not contain an isometric copy of  $\lambda \cdot \Gamma$  for all  $\lambda$  in a given interval, then this “nonrandom” behavior is detected by the Fourier transform of the characteristic function of  $A$  and results in “structural information”, specifically a concentration of its  $L^2$ -mass on appropriate annuli.

**Proposition 9** (dichotomy). *Let  $\delta > 0$  and  $\Gamma$  be a proper  $k$ -degenerate distance graph in  $[0, 1]^d$  with  $d \geq k + 1$ .*

*If  $A \subseteq [0, 1]^d$  with  $|A| > 0$ ,  $0 < a \leq b \ll \varepsilon^4$  with  $0 < \varepsilon \ll_{\Gamma, \delta} |A|^{n+1}$ , and  $A$  does not contain a  $\delta$ -close isometric copy of  $\lambda \cdot \Gamma$  for some  $\lambda$  in  $[a, b]$ , then*

$$\int_{\varepsilon^2/b \leq |\xi| \leq 1/\varepsilon^2 a} |\hat{1}_A(\xi)|^2 d\xi \gg c_1^2 |A|^{2n+2}, \tag{21}$$

with the implied constant above independent of  $a, b$ , and  $\varepsilon$ , and

$$c_1 = c_1(\Gamma, \delta) = \int \cdots \int d\mu_n(x_n) \cdots d\mu_1(x_1).$$

*Proof that Proposition 9 implies Theorem 2.* We shall first establish Part (ii) of Theorem 2, so we start by letting  $A \subseteq [0, 1]^d$  with  $|A| > 0$ . For any fixed  $0 < \varepsilon \ll_{\Gamma, \delta} |A|^{n+1}$ , let  $\{\mathcal{I}_j\}_{j=1}^{J(\varepsilon)}$  denote a sequence of intervals with  $\mathcal{I}_j := [a_j, b_j]$  satisfying

$$b_{j+1} \ll \varepsilon^4 a_j \tag{22}$$

and  $b_1 \ll \varepsilon^4$  with the property that for each  $1 \leq j \leq J(\varepsilon)$  there exists a  $\lambda \in \mathcal{I}_j$  such that

$$x + \lambda \cdot U(\Delta) \not\subseteq A \tag{23}$$

for all  $x \in A$  and  $U \in \text{SO}(d)$ . Proposition 9, together with (22), would then imply

$$J(\varepsilon)\varepsilon^2 \leq \sum_{j=1}^{J(\varepsilon)} \int_{\varepsilon^2/b_j \leq |\xi| \leq 1/\varepsilon^2 a_j} |\hat{1}_A(\xi)|^2 d\xi \leq \int |\hat{1}_A(\xi)|^2 d\xi, \tag{24}$$

a contradiction if  $J(\varepsilon) \gg \varepsilon^{-2}$  since by Plancherel we know that  $\int |\hat{1}_A(\xi)|^2 d\xi = |A| \leq 1$ .

To establish Part (i) of Theorem 2 with this approach we will argue indirectly and thus suppose that  $A \subseteq \mathbb{R}^d$  is a set with  $\delta^*(A) > 0$  for which the conclusion of Part (i) of Theorem 2 fails to hold, namely that there exist arbitrarily large  $\lambda \in \mathbb{R}$  for which  $A$  does not contain an isometric copy of  $\lambda \cdot \Gamma$ .

We now let  $0 < \alpha < \delta^*(A)$ ,  $0 < \varepsilon \ll_{\Gamma, \delta} \alpha^{n+1}$ , and fix  $J \gg \varepsilon^{-2}$  as above. By our indirect assumption we can choose a sequence  $\{\lambda_j\}_{j=1}^J$  with the property that  $\lambda_{j+1} \ll \varepsilon^4 \lambda_j$  for all  $1 \leq j \leq J - 1$  and  $A$  does not contain an isometric copy of  $\lambda_j \cdot \Gamma$  for each  $1 \leq j \leq J$ . It follows from the definition of upper Banach density that there exist  $N \in \mathbb{R}$  with  $N \gg \lambda_1$  and  $t_0 \in \mathbb{R}^d$  for which

$$\frac{|A \cap (t_0 + Q_N)|}{|Q_N|} \geq \alpha.$$

Rescaling  $A \cap (t_0 + Q_N)$  to a subset of  $[0, 1]^d$  and arguing as in the proof of Part (ii) above but this time with  $b_j = \lambda_j/N$  again leads to a contradiction. □

**4.2. Proof of Proposition 9.** Let  $f = 1_A$ . We will utilize the existence of a suitably *smoothed* version of  $f$  with the certain properties, specifically:

**Lemma 10.** *For any  $\varepsilon > 0$  there exists a function  $g : \mathbb{R}^d \rightarrow (0, 1]$ , an appropriate smoothing of  $f$ , such that*

$$|g(x - \lambda z) - g(x)| \ll \varepsilon \tag{25}$$

*uniformly in  $x \in [0, 1]^d$  and  $|z| \leq 1$ . Moreover, if  $\varepsilon \ll |A|^{n+1}$ , then*

$$\int f(x)g(x)^n dx \gg |A|^{n+1}. \tag{26}$$

The proof of Lemma 10 is straightforward and presented in Section 4.3 below. Assuming for now the existence of a function  $g$  with property (25) it follows that

$$\begin{aligned} T_{\Gamma, \delta}(f, f, \dots, f)(\lambda) &= c_1 \int f(x)g(x)^n dx + \sum_{m=1}^n T_{\Gamma, \delta}(fg^{n-m}, f, \dots, f, f - g, \underbrace{1, \dots, 1}_{n-m \text{ copies}})(\lambda) + O(n\varepsilon). \end{aligned} \tag{27}$$

If  $A$  does *not* contain a  $\delta$ -close isometric copy of  $\lambda \cdot \Gamma$  for some  $\lambda$  in  $[a, b]$ , then it clearly follows that

$$T_{\Gamma, \delta}(f, f, \dots, f)(\lambda) = 0.$$

In light of (26) and (27) it follows that if  $\varepsilon \ll c_1|A|^{n+1}/n$  then there must exist  $1 \leq m \leq n$  such that

$$\begin{aligned} \int \cdots \int \left( \int \left| \int [f - g](x - \lambda x_m) c_{m+1}(x_1, \dots, x_m) d\mu_m(x_m) \right| dx \right) d\mu_{m-1}(x_{m-1}) \cdots d\mu_1(x_1) \\ \gg c_1|A|^{n+1}, \end{aligned} \tag{28}$$

with  $c_{m+1}$  defined as before in (10) above. It then follows from an application of Cauchy–Schwarz and Plancherel that

$$\int |\hat{f}(\xi) - \hat{g}(\xi)|^2 I_m(\lambda\xi) d\xi \gg c_1^2|A|^{2n+2}, \tag{29}$$

with  $I_m$  again defined as before in (12) above. The fact that  $g$  will be taken to be a sufficient smoothing of  $f$  ensures that its Fourier transform satisfies

$$|\hat{f}(\xi) - \hat{g}(\xi)| \leq \varepsilon |\hat{f}(\xi)| \tag{30}$$

provided  $|\xi| \leq \varepsilon^2 b^{-1}$ ; see Section 4.3 below. This, together with the fact that  $I_m(\xi)$  is bounded by 1 uniformly in  $\xi$ , and Plancherel, ensures that (29) implies

$$\int_{\varepsilon^2/b \leq |\xi|} |\hat{f}(\xi)|^2 I_m(\lambda\xi) d\xi \gg c_1^2|A|^{2n+2} \tag{31}$$

provided  $\varepsilon \ll_{\Gamma, \delta} |A|^{n+1}$ . Estimate (21), and hence Proposition 9, then follows easily from estimate (31) and our previously established estimates for  $I_m$ , namely (14).

**4.3. A smooth cutoff function and proof of Lemma 10.**

**4.3.1. A smooth cutoff function.** Let  $\psi : \mathbb{R}^d \rightarrow (0, \infty)$  be a Schwartz function that satisfies

$$1 = \hat{\psi}(0) \geq \hat{\psi}(\xi) \geq 0 \quad \text{and} \quad \hat{\psi}(\xi) = 0 \quad \text{for } |\xi| > 1.$$

As usual, for any given  $t > 0$ , we define

$$\psi_t(x) = t^{-d} \psi(t^{-1}x). \tag{32}$$

First we record the trivial observation that

$$\int \psi_t(x) dx = \int \psi(x) dx = \hat{\psi}(0) = 1, \tag{33}$$

as well as the simple, but important, observation that  $\psi$  may be chosen so that

$$|1 - \hat{\psi}_t(\xi)| = |1 - \hat{\psi}(t\xi)| \ll \min\{1, t|\xi|\}. \tag{34}$$

Finally we record a formulation, appropriate to our needs, of the fact that for any given small parameter  $\varepsilon$ , our cutoff function  $\psi_t(x)$  will be essentially supported where  $|x| \leq \varepsilon^{-1}t$  and is approximately constant on smaller scales. More precisely:

**Lemma 11.** *Let  $\varepsilon > 0$  and  $t > 0$ ; then*

$$\int_{|y| \geq \varepsilon^{-1}t} \psi_t(y) dy \ll \varepsilon, \tag{35}$$

$$\int |\psi_t(y - \lambda z) - \psi_t(y)| dy \ll \varepsilon \tag{36}$$

*uniformly for  $|z| \leq 1$ , provided  $t \gg \varepsilon^{-1}\lambda$ .*

*Proof of Lemma 11.* Estimate (35) is easily verified using the fact that  $\psi$  is a Schwartz function on  $\mathbb{R}^d$  as

$$\int_{|y| \geq \varepsilon^{-1}t} \psi_t(y) dy = \int_{|y| \geq \varepsilon^{-1}} \psi(y) dy \ll \int_{|y| \geq \varepsilon^{-1}} (1 + |y|)^{-d-1} dy \ll \varepsilon.$$

To verify estimate (36) we make use of the fact that both  $\psi$  and its derivative are rapidly decreasing, specifically

$$\begin{aligned} \int |\psi_t(y - \lambda z) - \psi_t(x)| dy &\leq \int |\psi(y - \lambda z/t) - \psi(y)| dy \\ &\ll \frac{\lambda}{t} \int (1 + |y|)^{-d-1} dy \ll \frac{\lambda}{t}. \end{aligned} \quad \square$$

**4.3.2. Proof of Lemma 10.** Let  $g = f * \psi_{\varepsilon^{-1}b}$ .

We first note that estimates (30) and (25) follow immediately from (34) and (36) respectively. In order to establish the remaining “main term” estimate (26), we need only establish that if  $\varepsilon \ll |A|^{n+1}$ , then

$$\int f(x)g(x) dx \geq (1 - C\varepsilon) |A|^2 \tag{37}$$

for some constant  $C > 0$ , since by Hölder we would then obtain

$$(1 - C\varepsilon)^n |A|^{2n} \leq \left( \int f(x)g(x) dx \right)^n \leq |A|^{n-1} \int f(x)g(x)^n dx,$$

from which (26) clearly follows for sufficiently small  $\varepsilon > 0$ .

To establish (37) we first note that Parseval, the fact that  $0 \leq \hat{\psi} \leq 1$ , and a final application of Cauchy–Schwarz give

$$\begin{aligned} \int f(x)g(x) dx &= \int |\hat{f}(\xi)|^2 \hat{\psi}(\varepsilon^{-1}b\xi) d\xi \\ &\geq \int |\hat{f}(\xi)|^2 |\hat{\psi}(\varepsilon^{-1}b\xi)|^2 d\xi = \int g(x)^2 dx \geq \left( \int_{[0,1]^d} g(x) dx \right)^2. \end{aligned} \tag{38}$$

Establishing (37) therefore reduces to showing that if  $\varepsilon \ll |A|^3$ , then

$$\int_{[0,1]^d} g(x) dx \geq (1 - C\varepsilon)|A| \tag{39}$$

for some constant  $C > 0$ . To establish (39) we use (33) and write

$$\begin{aligned} |A| &= \int_{\mathbb{R}^d} g(x) dx \\ &= \int_{[0,1]^d} g(x) dx + \int_{\{x \in \mathbb{R}^d : \text{dist}(x, [0,1]^d) \geq \varepsilon^{-2}b\}} g(x) dx + \int_{\{x \in \mathbb{R}^d : 0 < \text{dist}(x, [0,1]^d) < \varepsilon^{-2}b\}} g(x) dx. \end{aligned} \tag{40}$$

The fact that  $b \leq \varepsilon^4$  ensures that

$$|\{x \in \mathbb{R}^d : 0 < \text{dist}(x, [0,1]^d) < \varepsilon^{-2}b\}| \ll \varepsilon^2 \tag{41}$$

and hence, since  $\varepsilon \ll |A|$  and  $0 \leq g \leq 1$ , that

$$\int_{\{x \in \mathbb{R}^d : 0 < \text{dist}(x, [0,1]^d) < \varepsilon^{-2}b\}} g(x) dx \ll \varepsilon^2 \leq \varepsilon|A|,$$

while (35) ensures that

$$\int_{\{x \in \mathbb{R}^d : \text{dist}(x, [0,1]^d) \geq \varepsilon^{-2}b\}} g(x) dx \leq |A| \int_{|y| \gg \varepsilon^{-2}b} \psi_{\varepsilon^{-1}b}(y) dy \ll \varepsilon|A|, \tag{42}$$

which completes the proof. □

### References

[Bennett et al. 2016] M. Bennett, A. Iosevich, and K. Taylor, “Finite chains inside thin subsets of  $\mathbb{R}^d$ ”, *Anal. PDE* **9**:3 (2016), 597–614. MR Zbl

[Birkhoff and Mac Lane 1941] G. Birkhoff and S. Mac Lane, *A survey of modern algebra*, Macmillan, New York, 1941. MR Zbl

[Bourgain 1986] J. Bourgain, “A Szemerédi type theorem for sets of positive density in  $\mathbb{R}^k$ ”, *Israel J. Math.* **54**:3 (1986), 307–316. MR Zbl

- [Bulinski 2018] K. Bulinski, “Spherical recurrence and locally isometric embeddings of trees into positive density subsets of  $\mathbb{Z}^d$ ”, *Math. Proc. Cambridge Philos. Soc.* **165**:2 (2018), 267–278. MR Zbl
- [Cook et al. 2017] B. Cook, Á. Magyar, and M. Pramanik, “A Roth-type theorem for dense subsets of  $\mathbb{R}^d$ ”, *Bull. Lond. Math. Soc.* **49**:4 (2017), 676–689. MR Zbl
- [Furstenberg et al. 1990] H. Furstenberg, Y. Katznelson, and B. Weiss, “Ergodic theory and configurations in sets of positive density”, pp. 184–198 in *Mathematics of Ramsey theory*, edited by J. Nešetřil and V. Rödl, Algorithms Combin. **5**, Springer, 1990. MR Zbl
- [Iosevich and Parshall 2018] A. Iosevich and H. Parshall, “Embedding distance graphs in finite field vector spaces”, preprint, 2018. arXiv
- [Lyall and Magyar 2018] N. Lyall and Á. Magyar, “Product of simplices and sets of positive upper density in  $\mathbb{R}^d$ ”, *Math. Proc. Cambridge Philos. Soc.* **165**:1 (2018), 25–51. MR Zbl
- [Lyall et al. 2018] N. Lyall, Á. Magyar, and H. Parshall, “Spherical configurations over finite fields”, preprint, 2018, available at <https://tinyurl.com/lyallpdf>. To appear in *Amer. J. Math.* **142**: 2 (2020).
- [Shilov 1971] G. E. Shilov, *Linear algebra*, Prentice-Hall, Englewood Cliffs, NJ, 1971. MR Zbl
- [Stein 1993] E. M. Stein, *Harmonic analysis: real-variable methods, orthogonality, and oscillatory integrals*, Princeton Math. Series **43**, Princeton Univ. Press, 1993. MR Zbl
- [Szemerédi 1975] E. Szemerédi, “On sets of integers containing no  $k$  elements in arithmetic progression”, *Acta Arith.* **27** (1975), 199–245. MR Zbl
- [Tao and Vu 2006] T. Tao and V. Vu, *Additive combinatorics*, Cambridge Studies in Adv. Math. **105**, Cambridge Univ. Press, 2006. MR Zbl
- [Ziegler 2006] T. Ziegler, “Nilfactors of  $\mathbb{R}^m$ -actions and configurations in sets of positive upper density in  $\mathbb{R}^m$ ”, *J. Anal. Math.* **99** (2006), 249–266. MR Zbl

Received 23 Mar 2018. Revised 11 Jan 2019. Accepted 13 Mar 2019.

NEIL LYALL: [lyall@math.uga.edu](mailto:lyall@math.uga.edu)

*Department of Mathematics, The University of Georgia, Athens, GA, United States*

ÁKOS MAGYAR: [magyar@math.uga.edu](mailto:magyar@math.uga.edu)

*Department of Mathematics, The University of Georgia, Athens, GA, United States*

# Analysis & PDE

msp.org/apde

## EDITORS

EDITOR-IN-CHIEF

Patrick Gérard

patrick.gerard@math.u-psud.fr

Université Paris Sud XI

Orsay, France

## BOARD OF EDITORS

Massimiliano Berti	Scuola Intern. Sup. di Studi Avanzati, Italy berti@sissa.it	Gilles Pisier	Texas A&M University, and Paris 6 pisier@math.tamu.edu
Michael Christ	University of California, Berkeley, USA mchrist@math.berkeley.edu	Tristan Rivière	ETH, Switzerland riviere@math.ethz.ch
Charles Fefferman	Princeton University, USA cf@math.princeton.edu	Igor Rodnianski	Princeton University, USA irod@math.princeton.edu
Ursula Hamenstaedt	Universität Bonn, Germany ursula@math.uni-bonn.de	Yum-Tong Siu	Harvard University, USA siu@math.harvard.edu
Vadim Kaloshin	University of Maryland, USA vadim.kaloshin@gmail.com	Terence Tao	University of California, Los Angeles, USA tao@math.ucla.edu
Herbert Koch	Universität Bonn, Germany koch@math.uni-bonn.de	Michael E. Taylor	Univ. of North Carolina, Chapel Hill, USA met@math.unc.edu
Izabella Laba	University of British Columbia, Canada ilaba@math.ubc.ca	Gunther Uhlmann	University of Washington, USA gunther@math.washington.edu
Richard B. Melrose	Massachusetts Inst. of Tech., USA rbm@math.mit.edu	András Vasy	Stanford University, USA andras@math.stanford.edu
Frank Merle	Université de Cergy-Pontoise, France Frank.Merle@u-cergy.fr	Dan Virgil Voiculescu	University of California, Berkeley, USA dvv@math.berkeley.edu
William Minicozzi II	Johns Hopkins University, USA minicozz@math.jhu.edu	Steven Zelditch	Northwestern University, USA zelditch@math.northwestern.edu
Clément Mouhot	Cambridge University, UK c.mouhot@dpms.cam.ac.uk	Maciej Zworski	University of California, Berkeley, USA zworski@math.berkeley.edu
Werner Müller	Universität Bonn, Germany mueller@math.uni-bonn.de		

## PRODUCTION

production@msp.org

Silvio Levy, Scientific Editor

---

See inside back cover or [msp.org/apde](http://msp.org/apde) for submission instructions.

---

The subscription price for 2020 is US \$340/year for the electronic version, and \$550/year (+\$60, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues from the last three years and changes of subscriber address should be sent to MSP.

---

Analysis & PDE (ISSN 1948-206X electronic, 2157-5045 printed) at Mathematical Sciences Publishers, 798 Evans Hall #3840, c/o University of California, Berkeley, CA 94720-3840, is published continuously online. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices.

---

APDE peer review and production are managed by EditFlow® from MSP.

PUBLISHED BY

 **mathematical sciences publishers**  
nonprofit scientific publishing

<http://msp.org/>

© 2020 Mathematical Sciences Publishers

# ANALYSIS & PDE

Volume 13 No. 3 2020

---

On the gap between the Gamma-limit and the pointwise limit for a nonlocal approximation of the total variation	627
CLARA ANTONUCCI, MASSIMO GOBBINO and NICOLA PICENNI	
External boundary control of the motion of a rigid body immersed in a perfect two-dimensional fluid	651
OLIVIER GLASS, JÓZSEF J. KOLUMBÁN and FRANCK SUEUR	
Distance graphs and sets of positive upper density in $\mathbb{R}^d$	685
NEIL LYALL and ÁKOS MAGYAR	
Isolated singularities for semilinear elliptic systems with power-law nonlinearity	701
MARIUS GHERGU, SUNGHAN KIM and HENRIK SHAHGOLIAN	
Regularity of the free boundary for the vectorial Bernoulli problem	741
DARIO MAZZOLENI, SUSANNA TERRACINI and BOZHIDAR VELICHKOV	
On the discrete Fuglede and Pompeiu problems	765
GERGELY KISS, ROMANOS DIOGENES MALIKIOSIS, GÁBOR SOMLAI and MÁTÉ VIZER	
Energy conservation for the compressible Euler and Navier–Stokes equations with vacuum	789
IBROKHIMBEK AKRAMOV, TOMASZ DĘBIEC, JACK SKIPPER and EMIL WIEDEMANN	
A higher-dimensional Bourgain–Dyatlov fractal uncertainty principle	813
RUI HAN and WILHELM SCHLAG	
Local minimality results for the Mumford–Shah functional via monotonicity	865
DORIN BUCUR, ILARIA FRAGALÀ and ALESSANDRO GIACOMINI	
The gradient flow of the Möbius energy: $\varepsilon$ -regularity and consequences	901
SIMON BLATT	
Correction to the article The heat kernel on an asymptotically conic manifold	943
DAVID A. SHER	



2157-5045(2020)13:3;1-B