

ANALYSIS & PDE

Volume 13 No. 3 2020

GERGELY KISS, ROMANOS DIOGENES MALIKIOSIS,
GÁBOR SOMLAI AND MATE VIZER

ON THE DISCRETE FUGLEDE AND POMPEIU PROBLEMS

ON THE DISCRETE FUGLEDE AND POMPEIU PROBLEMS

GERGELY KISS, ROMANOS DIOGENES MALIKIOSIS, GÁBOR SOMLAI AND MÁTÉ VIZER

We investigate the discrete Fuglede conjecture and the Pompeiu problem on finite abelian groups and develop a strong connection between the two problems. We give a geometric condition under which a multiset of a finite abelian group has the discrete Pompeiu property. Using this description and the revealed connection we prove that Fuglede's conjecture holds for $\mathbb{Z}_{p^n q^2}$, where p and q are different primes. In particular, we show that every spectral subset of $\mathbb{Z}_{p^n q^2}$ tiles the group. Further, using our combinatorial methods we give a simple proof for the statement that Fuglede's conjecture holds for \mathbb{Z}_p^2 .

1. Introduction

In this article we deal with the discrete version of Fuglede's conjecture and the Pompeiu problem; both originated in analysis. We build a relationship between them that helps us to provide new results for Fuglede's conjecture in the discrete setting.

The following question was asked by Pompeiu [1929]. Take a continuous function f on the plane whose integral is zero on every unit disc. Does it follow that f is constantly zero? The answer to this question is no, but it initiated several different types of investigations in various settings, and in some cases the answer is affirmative for an analogous question. We give an implicit characterization of the non-Pompeiu sets for finite abelian groups.

Fuglede [1974] conjectured that a bounded domain $S \subset \mathbb{R}^d$ tiles the d -dimensional Euclidean space if and only if the set of $L^2(S)$ functions admits an orthogonal basis of exponential functions. This conjecture was disproved by Tao [2004].

A discrete version of Fuglede's conjecture might be formulated in the following way. A subset S of a finite abelian group G tiles G if and only if the character table of G has a submatrix, whose rows are indexed by the elements of S , which is a complex Hadamard matrix. This version of Fuglede's conjecture is not only interesting on its own but also plays a crucial role in the above-mentioned counterexample of Tao. Actually his counterexample (in \mathbb{R}^5) is based on a counterexample for elementary abelian p -groups of finite rank.

Fuglede's conjecture is especially interesting for finite cyclic groups, since, e.g., every tiling of \mathbb{Z} is periodic, so it goes back to a tiling of a finite cyclic group. However, not much is known for cyclic groups. A recent paper of the second author and Kolountzakis [Malikiosis and Kolountzakis 2017] shows that Fuglede's conjecture holds for any cyclic group of order $p^n q$, where p and q are different primes.

MSC2010: primary 39B32, 43A46; secondary 13F20, 20K01.

Keywords: Pompeiu problem, spectral set conjecture, tiling, vanishing sums of roots of unity.

Our main contribution towards Fuglede's conjecture for cyclic groups is to connect this problem with the Pompeiu problem, introduce more combinatorial ideas and verify it for yet unknown cases: cyclic groups of order $p^n q^2$, $n \geq 1$ (see Theorem 2.5).

Further using our techniques we give a neat and combinatorial proof for the previously known fact, proved by Iosevich, Mayeli and Pakianathan [Iosevich et al. 2017], that Fuglede's conjecture holds for \mathbb{Z}_p^2 (see also Theorem A.1).

Structure of the paper. Section 2 is devoted to a detailed introduction to Fuglede's conjecture and the Pompeiu problem, introducing also the discrete versions of them. Further we establish a connection between the two problems. In Section 3 we give some sort of solution for the Pompeiu problem for abelian groups that we apply later in Section 6. Sections 4 and 5 are preparations for the proof of Theorem 2.5. In Section 4 we reduce the cases to a special one partly based on our results concerning the Pompeiu problem. In Section 5 we prove some technical lemmas, which we use later. Section 6 is devoted to the proof of Theorem 2.5. Finally, in the Appendix we give an alternative proof of Theorem A.1.

2. Fuglede and Pompeiu problems

Fuglede's spectral set conjecture. The original conjecture of Fuglede [1974] was formulated as follows. Let Ω be a measurable subset of \mathbb{R}^n of positive Lebesgue measure. A set $\Omega \subseteq \mathbb{R}^n$ is called *spectral* if there is a set $\Lambda \subset \mathbb{R}^n$ such that $\{e^{i\lambda \cdot x} : \lambda \in \Lambda, x \in \Omega\}$ is an orthogonal basis of $L^2(\Omega)$. Then $\Lambda \subseteq \mathbb{R}^n$ is called the *spectrum* of Ω .

We say that S is a *tile* of \mathbb{R}^n if there is a set $T \subset \mathbb{R}^n$ such that almost every point of \mathbb{R}^n can be uniquely written as $s + t$, where $s \in S$ and $t \in T$. In this case, we say that T is the *tiling complement* of S .

Fuglede's spectral set conjecture [1974] (which we just call Fuglede's conjecture) states the following:

Conjecture 1. Ω is spectral if and only if Ω is a tile.

The conjecture was proved by Fuglede [1974] in the special case when the tiling complement or the spectrum is a lattice in \mathbb{R}^n . Also it has been verified by Fuglede that the L^2 -space over a triangle or a disc does not admit an orthogonal basis of exponentials. (The proof for the disc was corrected by Iosevich, Katz and Pedersen [Iosevich et al. 1999].) The conjecture was further verified in some other cases; see, e.g., [Iosevich et al. 2003; Łaba 2001].

Tao [2004] disproved the Spectral \implies Tiling direction of the conjecture by constructing a spectral set in \mathbb{R}^5 that does not tile the 5-dimensional space. As an extension of Tao's work, Matolcsi [2005] proved that (the same direction of) the conjecture fails in dimension 4 as well. Further, Kolountzakis and Matolcsi [2006a; 2006b] and Farkas, Matolcsi and Móra [Farkas et al. 2006] provided counterexamples in dimension 3 for each direction of the conjecture.

Discrete abelian groups. Fuglede's conjecture can be naturally stated for other groups, for example \mathbb{Z} . These cases are not only interesting on their own, but they also have connection with the original case, since, e.g., in his disproof of the 5-dimensional case, Tao constructed a spectral set in \mathbb{Z}_3^5 (containing six elements, hence not a tile, as the cardinality of any tile of a finite abelian group divides the order of the

group), then he lifted this counterexample to \mathbb{R}^5 . A similar strategy was carried out by Kolountzakis and Matolcsi [2006b] in the disproof of the other direction of the original conjecture. We also mention some examples where Fuglede’s conjecture holds. These include finite cyclic p -groups [Łaba 2002], $\mathbb{Z}_p \times \mathbb{Z}_p$ [Iosevich et al. 2017], and \mathbb{Q}_p [Fan et al. 2019], the field of p -adic numbers.

Borrowing the notation from [Dutkay and Lai 2014; Malikiosis and Kolountzakis 2017], we write $S - T(G)$ (resp. $T - S(G)$), if the Spectral \implies Tiling (resp. Tiling \implies Spectral) direction of Fuglede’s conjecture holds in G for every bounded subset. The above-mentioned connection between the conjecture on \mathbb{R} , on \mathbb{Z} and on finite cyclic groups is summarized below [Dutkay and Lai 2014], where $T - S(\mathbb{Z}_{\mathbb{N}})$ means that $T - S(\mathbb{Z}_n)$ holds for every $n \in \mathbb{N}$:

$$\begin{aligned} T - S(\mathbb{R}) &\iff T - S(\mathbb{Z}) \iff T - S(\mathbb{Z}_{\mathbb{N}}), \\ S - T(\mathbb{R}) &\implies S - T(\mathbb{Z}) \implies S - T(\mathbb{Z}_{\mathbb{N}}). \end{aligned}$$

According to this, a counterexample to the Spectral \implies Tiling direction in a finite cyclic group can be lifted to a counterexample in \mathbb{R} ; on the other hand, if the same direction of the conjecture were true for every cyclic group or even in \mathbb{Z} , this would hold no meaning for the original conjecture in \mathbb{R} .

Concerning tiles in discrete groups it was proved in [Newman 1977] that if S is a finite set, which tiles \mathbb{Z} with tiling complement T , then T is periodic; i.e., $T + N = T$ for some $N \in \mathbb{Z}$. This shows that every tiling of the integers reduces to a tiling of a cyclic group \mathbb{Z}_N for some $N \in \mathbb{N}$.

We also mention a related result of [Rédei 1965]. We say that $A_1 + \dots + A_k$ is a *factorization* of the abelian group G if every element of G can uniquely be written as the sum of one element from each A_i .

Theorem 2.1 [Rédei 1965]. *Let $G = A_1 + A_2 + \dots + A_n$ be a factorization of an abelian group G , where each A_i contains 0 and is of prime cardinality. Then at least one of the sets A_i is a subgroup of G .*

Cyclic groups. Surprisingly, despite their previously described role in the discrete version of Fuglede’s conjecture, not much is known for cyclic groups. A recent result of [Malikiosis and Kolountzakis 2017] proved Conjecture 2 (see later) for $\mathbb{Z}_{p^n q}$. They also wrote that most likely, their result might be extended to cyclic groups of order having two different prime divisors but they haven’t succeeded yet.

As we will mainly deal with cyclic groups, let us state the conjecture again in this setting. First let us define spectral sets and tiles in cyclic groups also.

Definition 2.2. For a set $S \subset \mathbb{Z}_N$, we say that S is *spectral* if $L^2(S)$ has an orthogonal basis of exponentials (indexed by Λ). This is equivalent to the following two conditions holding:

- (1) There is $\Lambda \subset \mathbb{Z}_N$ such that any $f : S \rightarrow \mathbb{C}$ can be written as the \mathbb{C} -linear combination of exponentials of the form

$$\xi_N^{\lambda \cdot x}, \quad \lambda \in \Lambda,$$

where the product $\lambda \cdot x$ is taken modulo N and $\xi_N = e^{2\pi i / N}$.

- (2) For any two different $\lambda, \lambda' \in \Lambda$ we have

$$\sum_{x \in S} \xi_N^{(\lambda - \lambda') \cdot x} = 0$$

(i.e., the representations $\chi_\lambda(x) = \xi_N^{\lambda \cdot x}$ and $\chi_{\lambda'}(x) = \xi_N^{\lambda' \cdot x}$ are orthogonal).

We denote $\{\chi_\lambda : \lambda \in \Lambda\}$ by χ_Λ .

Remark 2.3. We note that if S is a spectral set, then $|S| = |\Lambda|$ follows from Definition 2.2. Condition (2) further implies

$$\Lambda - \Lambda \subseteq \{0\} \cup \{x \in \mathbb{Z}_N : \hat{1}_S(x) = 0\}, \quad (1)$$

where 1_S is the characteristic function of S , and $\hat{f}(x) = \sum_{y \in \mathbb{Z}_N} f(y) \xi_N^{-x \cdot y}$ is the discrete Fourier transform of $f : G \mapsto \mathbb{C}$, as usual.

Definition 2.4. Let G be a discrete abelian group. We say that $S \subset G$ *tiles* G if there exists $T \subset G$ such that $S + T = G$, where $S + T$ is the set of elements of G of the form $s + t$, $s \in S, t \in T$, counted with multiplicity, so we have each $g \in G$ exactly once. In this case we say that T is a *tiling complement* of S in G .

For cyclic groups Fuglede's conjecture can be stated as follows.

Conjecture 2. For any N and $S \subset \mathbb{Z}_N$ we have that S is spectral if and only if S tiles \mathbb{Z}_N .

There has been some recent progress on this conjecture over the last few years. The known results for the Tiling \implies Spectral direction follow from [Coven and Meyerowitz 1999; Łaba 2002]. Coven and Meyerowitz proved that if a finite subset A of the integers satisfies two conditions (T1) and (T2) (to be defined below) then it tiles \mathbb{Z} by translations. The inverse holds when the cardinality of A is divisible by at most two primes (corollary to [Coven and Meyerowitz 1999, Theorem B2]). Łaba then connected these properties with Fuglede's conjecture on \mathbb{Z} , proving that if A satisfies (T1) and (T2), then it has a spectrum [Łaba 2002, Theorem 1.5(i)]; therefore, if A tiles \mathbb{Z} and its cardinality is divisible by at most two distinct primes, then it must be spectral [Łaba 2002, Corollary 1.6(i)]. The passage to cyclic groups of order $p^n q^m$ is easily done through [Coven and Meyerowitz 1999, Lemma 2.3], which implies that if A tiles \mathbb{Z} and $|A| = p^a q^b$, then there is a (possibly different) tiling of \mathbb{Z} by translates of A with period $N = p^n q^m$ for some $m, n \in \mathbb{N}$. In other words, A could be considered as a subset of \mathbb{Z}_N , and the Tiling \implies Spectral direction in \mathbb{Z}_N follows verbatim using the above results. For a proof containing all the above arguments strictly in the setting of cyclic groups, we refer the reader to [Malikiosis and Kolountzakis 2017, Section 3].

Concerning the case for N square-free, the Tiling \implies Spectral direction in \mathbb{Z}_N follows easily from the fact that any tile of \mathbb{Z}_N is a set of coset representatives of a subgroup of \mathbb{Z}_N , again from combined arguments from [Coven and Meyerowitz 1999; Łaba 2002]. This fact was posed as a problem in Tao's blog,¹ which was subsequently solved by Łaba and Meyerowitz (see the comments of that post). Their arguments were based on [Coven and Meyerowitz 1999, Lemma 2.3], which implies that a tile A in \mathbb{Z}_N with N square-free, accepts the subgroup $M\mathbb{Z}_N$ as a tiling complement, where $M = |A|$. This is one instance where the properties (T1) and (T2) hold trivially; thus A is also spectral due to [Łaba 2002, Theorem 1.5(i)]. For a self-contained proof of the Tiling \implies Spectral direction in the setting of cyclic groups of square-free order, which is along the same lines, we refer the reader to [Shi 2019].

¹<https://terrytao.wordpress.com/2011/11/19/some-notes-on-the-coven-meyerowitz-conjecture/>

The reverse direction, Spectral \implies Tiling, is considerably harder, and the best results to this date are the proofs for $N = p^n q$ [Malikiosis and Kolountzakis 2017] and $N = pqr$ [Shi 2019], where p, q, r are distinct primes. The main tool that is introduced in the Spectral \implies Tiling direction is the structure of the vanishing sums of roots of unity [Lam and Leung 2000].

In this paper we verify Conjecture 2 for cyclic groups of order $p^n q^2$ by proving the following.

Theorem 2.5. *Let p and q be two different primes. Then we have $S - T(\mathbb{Z}_{p^n q^2})$ for every $n \geq 1$.*

As stated above, $T - S(\mathbb{Z}_{p^n q^m})$ has already been proven [Coven and Meyerowitz 1999; Łaba 2002]. Combining this result and Theorem 2.5 we obtain:

Theorem 2.6. *Let p and q be two different primes. Then Fuglede’s conjecture holds for $\mathbb{Z}_{p^n q^2}$, $n \geq 1$.*

Furthermore, using our method, we give in the Appendix a simple proof of the theorem of Iosevich, Mayeli and Pakianathan [Iosevich et al. 2017], stating that Fuglede’s conjecture holds for \mathbb{Z}_p^2 .

Pompeiu problem. The problem goes back to the seminal paper [Pompeiu 1929], where he asked the following question of integral geometry:

Question 1. Let K be a compact set of positive Lebesgue measure. Is it true that if $f : \mathbb{R}^2 \rightarrow \mathbb{C}$ is a continuous function that satisfies

$$\int_{\sigma(K)} f(x, y) d\lambda_x d\lambda_y = 0 \tag{2}$$

for every rigid motion σ (here λ denotes the Lebesgue measure), then f is identically zero (i.e. $f \equiv 0$)?

If K is the closed disc of radius $r > 0$, then the answer is negative. It was shown in [Chakalov 1944], see also [Garofalo 1989], that (2) holds if $f(x, y) = \sin(a(x + iy))$, where $a > 0$, and $J_1(ra) = 0$, where J_λ denotes the Bessel function of order λ . On the other hand, for every nonempty polygon (moreover, for any convex domain with at least one corner) the answer for Question 1 is affirmative by a result of Brown, Schreiber, and Taylor [Brown et al. 1973]. Recently, Ramm [2017] showed that there exists a function $f \not\equiv 0$ that satisfies the 3-dimensional analogue of (2) for a bounded domain $K \subseteq \mathbb{R}^3$ with C^1 -smooth boundary if and only if K is a closed ball. Extensive literature is concerned with the Pompeiu problem. For the history of the problem see [Ramm 1997] and the bibliographical survey [Zalcman 1992].

Here we investigate the discrete version of the *Pompeiu problem* on finite abelian groups. We note that it was studied on infinite abelian groups in [Kiss et al. 2018; Puls 2013; Zeilberger 1978].

The discrete version of Pompeiu problem for an abelian group G . In the sequel we denote the binary operation acting on an abelian group G by $+$ (as the usual addition).

Definition 2.7. Let G be an abelian group:

- Let S be a nonempty finite subset of G . We say that S has the discrete Pompeiu property (or S is Pompeiu) if whenever $f : G \rightarrow \mathbb{C}$ satisfies

$$\sum_{s \in S} f(s + x) = 0 \quad \text{for every } x \in G, \tag{3}$$

then $f \equiv 0$.

We say that S is a *non-Pompeiu set with respect to f* if $f \not\equiv 0$ and satisfies (3).

One can define the discrete Pompeiu property for multisets similarly.

- We call $w : G \rightarrow \mathbb{Q}$ a *weight function*² defined on G . We say that w is a *Pompeiu weight function* if for any $f : G \rightarrow \mathbb{C}$

$$\sum_{g \in G} w(g)f(g+x) = 0 \quad \text{for every } x \in G \quad (4)$$

implies that $f \equiv 0$.

We say that w is a *non-Pompeiu weight function with respect to f* if $f \not\equiv 0$ and satisfies (4).

Note that S is a Pompeiu set if and only if its characteristic function is a Pompeiu weight function.

Remark 2.8. We can extend the previous definition for arbitrary finite group (G, \cdot) and weight function w as follows.

Let $w : G \rightarrow \mathbb{Q}$. We denote by $\text{Cay}(G, w)$ the *Cayley graph* of G with respect to w . The vertex set of $\text{Cay}(G, w)$ is G and g is connected to h by an edge with weight $w(g^{-1}h)$ for every $g, h \in G$. We denote by A_w the adjacency matrix of $\text{Cay}(G, w)$. Using the adjacency matrix A_w of $\text{Cay}(G, w)$ we may also say w is a Pompeiu weight function if and only if $A_w f = 0$ implies $f \equiv 0$. The equation $A_w f = 0$ implies that if $f \not\equiv 0$, then f is an eigenvector of A_w with eigenvalue 0. So w is a Pompeiu weight function if and only if 0 is not an eigenvalue of A_w . In the finite case this is equivalent to A_w being invertible.

We note that if G is a cyclic group, then A_w is a circulant matrix.

The set of irreducible representations of a finite abelian group G will be denoted by \tilde{G} . Every irreducible representation of an abelian group is 1-dimensional (a character). Thus \tilde{G} is a group which is isomorphic to G . Note that \tilde{G} is usually called the *dual group* of G .

It is well known [Steinberg 2012] that the set of irreducible representations forms an orthogonal basis of $L^2(G)$ with respect to the natural scalar product

$$[\psi, \chi] := \sum_{g \in G} \psi(g)\overline{\chi(g)}$$

for $\psi, \chi \in \tilde{G}$. Thus every function $f : G \rightarrow \mathbb{C}$ can be uniquely written as

$$f(x) = \sum_{\chi \in \tilde{G}} c_\chi \chi(x) \quad \text{for all } x \in G, \quad (5)$$

for some $c_\chi \in \mathbb{C}$.

The following proposition can be deduced from [Székelyhidi 2001]. In order to make our paper self-contained, we provide the proof.

Proposition 2.9. *If w is a non-Pompeiu weight function with respect to a function f , then w is a non-Pompeiu weight function with respect to all irreducible representations χ which have nonzero coefficient c_χ in (5).*

²Every weight function is a rational constant multiple of a weight function with integer coefficients. The Pompeiu property is invariant under scalar multiplication, and thus we may restrict our attention to those weight functions which take their values in \mathbb{Z} .

Proof. Let w be non-Pompeiu with respect to a function f ; then $\sum_{s \in G} w(s)f(s+x) = 0$ for every $x \in G$. Using (5) we get

$$0 = \sum_{s \in S} w(s) \sum_{\chi \in \tilde{G}} c_\chi \chi(s+x) = \sum_{\chi \in \tilde{G}} c_\chi \sum_{s \in S} w(s)\chi(s+x) = \sum_{\chi \in \tilde{G}} \left(c_\chi \sum_{s \in S} w(s)\chi(s) \right) \chi(x),$$

since χ is a character. This statement holds for every $x \in G$ so we can formulate it as

$$\sum_{\chi \in \tilde{G}} \left(c_\chi \sum_{s \in S} w(s)\chi(s) \right) \chi = 0.$$

Since the irreducible representations are linearly independent over \mathbb{C} , the previous equation holds if and only if $\sum_{s \in S} w(s)\chi(s) = 0$ for all χ such that $c_\chi \neq 0$. Multiplying with $\chi(x)$ we obtain $\sum_{s \in S} w(s)\chi(x+s) = 0$. Since this holds for every $x \in G$, this means that w is non-Pompeiu with respect to such χ . □

We note that a stronger result was proved by Babai [1979], who determined the spectrum of Cayley graphs of abelian groups. The set of the eigenvalues of $\text{Cay}(G, S)$ is $\{ \sum_{s \in S} \chi(s) : \chi \in \tilde{G} \}$.

Corollary 2.10. *If S is a non-Pompeiu set in a finite abelian group, then S is non-Pompeiu with respect to some irreducible representation of G .*

Remark 2.11. Since the characters (irreducible representations) play the role of exponential functions over the abelian group G , it seems reasonable that the function $\sin(ax)$ can provide an example on the disk for the original Pompeiu problem. On the other hand, it is surprising that exponential solutions were not found in the literature.

Connection of the problems.

Proposition 2.12. *Let G be a finite abelian group. If $S \subset G$ is a spectral set with $|S| \geq 2$, then S is a non-Pompeiu set.*

Proof. The spectral property of S requires a set of irreducible representations, of the same cardinality of S , whose restrictions to S are pairwise orthogonal. Assume χ and ψ are different irreducible representations of G , whose restrictions to S are orthogonal. Since $[\chi|_S, \psi|_S] = [(\chi\bar{\psi})|_S, \text{id}|_S]$ we obtain a representation $\rho = \chi\bar{\psi}$ such that $\sum_{s \in S} \rho(s) = 0$, which leads us back to the Pompeiu problem. Thus we get that S is a non-Pompeiu set with respect to the irreducible representation ρ . □

3. Pompeiu problem for cyclic groups

In this section we consider the non-Pompeiu sets for abelian groups.

Every representation of a finite abelian group is linear, so it factorizes through a faithful representation of a cyclic group since the finite subgroups of $\mathbb{C} \setminus \{0\}$ are cyclic. This shows that some sort of description for non-Pompeiu sets of finite abelian groups is given by understanding the non-Pompeiu weight functions of cyclic groups with respect to faithful representations.

Let $(\mathbb{Z}_N, +)$ be the cyclic group of order N . Note that for all $k \mid N$ there is a unique normal subgroup $\mathbb{Z}_k \leq \mathbb{Z}_N$ of order k . The group generated this way contains exactly the elements of \mathbb{Z}_N divisible by N/k so this subgroup of $(\mathbb{Z}_N, +)$ will also be denoted by $H_{N/k}$.

We use the following isomorphism between \mathbb{Z}_N and $\tilde{\mathbb{Z}}_N$: Fix a primitive N -th root of unity α and a generator g of \mathbb{Z}_N . Then for any $j \in \mathbb{Z}_N$ the function $\psi_j(g^i) = \alpha^{ji}$ gives a homomorphism from \mathbb{Z}_N to \mathbb{C}^* ; hence it is an irreducible representation. Now $j \rightarrow \psi_j$ gives the isomorphism from \mathbb{Z}_N to $\tilde{\mathbb{Z}}_N$; throughout the text, we will use the isomorphism that arises from $\alpha = \xi_N$. From now on the subgroup of $\tilde{\mathbb{Z}}_N$ isomorphic to $H \leq \mathbb{Z}_N$ will be denoted by \tilde{H} .

Hereinafter we use the notion of mask polynomial.

Definition 3.1. Let G be a cyclic group and $w : G \rightarrow \mathbb{Q}$ be a weight function. We call

$$m_w(x) = \sum_{h \in G} w(h)x^h$$

the *mask polynomial* of w , where $w(h)$ denotes the weight of $h \in G$. This might be considered as an element of $\mathbb{Q}[x]/(x^n - 1)$. For a (multi-)set S of G we define the *mask polynomial* of S by

$$S(x) = \sum_{s \in S} c_s x^s,$$

where c_s denotes the cardinality of $s \in S$.

Let $\Phi_k(x)$ denote the k -th cyclotomic polynomial, which is of degree $\varphi(k)$, where φ denotes the Euler totient function. Note that for fixed N and prime $p \mid N$ the mask polynomial of $\mathbb{Z}_p \leq \mathbb{Z}_N$ is $\Phi_p(x^{N/p})$. The following is one of the key preliminary observations. Basically, this can be considered as a statement on roots of unity. There is a vast literature on vanishing sums of roots of unity. This particular statement gives a generalization of Theorem 3.3 of [Lam and Leung 2000]. Similar results might appear in other papers.

Proposition 3.2. *Let G be a cyclic group of order N and let α be a primitive N -th root of unity. We denote by P_N the set of prime divisors of N . Further let w be a weighted function. Then w is non-Pompeiu with respect to the faithful representation ψ_α if and only if*

$$w = \sum_{g \in G} \sum_{p \in P_N} w_{p,g} 1_{\mathbb{Z}_p+g}$$

for some $w_{p,g} \in \mathbb{Q}$, where $1_{\mathbb{Z}_p+g}$ denotes the characteristic function of the coset $\mathbb{Z}_p + g$.

Proof. The fact that w is a non-Pompeiu weight function with respect to the faithful representation ψ_α means that α is the root of the mask polynomial m_w of w , since $m_w(\alpha) = \sum_{i=0}^{N-1} w(i)\alpha^i = 0$. On the other hand, for a given $N \in \mathbb{N}$, for every $p \in P_N$ we have that α is the root of the mask polynomial of $\mathbb{Z}_p \leq \mathbb{Z}_N$, that is, $\Phi_p(x^{N/p})$. Indeed, α is a primitive N -th root of unity so $\alpha^{N/p} \neq 1$. Clearly, $\alpha^{N/p}\Phi_p(\alpha^{N/p}) = \Phi_p(\alpha^{N/p})$, so it implies $\Phi_p(\alpha^{N/p}) = 0$.

Then α is also a root of the polynomial $m_w(x) + \sum_{p \in P_N} a_p(x)\Phi_p(x^{N/p})$, where $a_p(x) \in \mathbb{Q}[x]$. By using Euclidean division there are polynomial $q(x), r(x) \in \mathbb{Q}[x]$ such that

$$m_w(x) = q(x)\Phi_N(x) + r(x),$$

with either $r(x)$ the constant zero function or $\deg(r(x)) < \varphi(N)$.

The common roots of the polynomials $\Phi_p(x^{N/p})$ ($p \in P_N$) are exactly the primitive N -th roots of unity. The multiplicity of these roots in all of these polynomials is 1. These polynomials are all in $\mathbb{Q}[x]$ so the greatest common divisor in the ring $\mathbb{Q}[x]$ of the polynomials $\Phi_p(x^{N/p})$, $p \in P_N$, is $\Phi_N(x)$. Thus

$$\Phi_N(x) = \sum_{p \in P_N} a_p(x) \Phi_p(x^{N/p})$$

for some $a_p(x) \in \mathbb{Q}[x]$. Substituting this into the previous equation we obtain that

$$m_w(x) - \sum_{p \in P_N} q(x) a_p(x) \Phi_p(x^{N/p})$$

is of degree less than $\varphi(N)$ or is the constant zero function. Since $\Phi_N(x)$ is the minimal polynomial of α over \mathbb{Q} , we have $m_w(x) - \sum_{p \in P_N} q(x) a_p(x) \Phi_p(x^{N/p}) = 0$. Thus

$$m_w(x) = \sum_{p \in P_N} q(x) a_p(x) \Phi_p(x^{N/p}).$$

It is clear that $x^k \Phi_p(x^{N/p})$ is the mask polynomial of a coset of \mathbb{Z}_p for every $0 \leq k < N$. Hence we have

$$w(x) = \sum_{g \in G} \sum_{p \in P_N} w_{p,g} 1_{\mathbb{Z}_p+g}(x)$$

for some $w_{p,g} \in \mathbb{Q}$.

The other direction follows from the fact that $\Phi_p(\alpha^{N/p}) = 0$ for every $p \in P_N$. □

We note that using Proposition 3.2 one can simply construct the asymmetric minimal sums of roots of unity appearing in [Lam and Leung 2000].

In terms of mask polynomials the previous proposition can be stated as follows.

Corollary 3.3. *Let $S(x) \in \mathbb{Z}_{\geq 0}[x]$ with $S(\xi_N) = 0$, where $N = p_1^{m_1} \cdots p_n^{m_n}$ and p_1, \dots, p_n are primes. Then,*

$$S(x) \equiv P_1(x) \Phi_{p_1}(x^{N/p_1}) + \cdots + P_n(x) \Phi_{p_n}(x^{N/p_n}) \pmod{(x^N - 1)}$$

for some $P_1(x), \dots, P_n(x) \in \mathbb{Q}[x]$.

The following is an easy consequence of Proposition 3.2.

Corollary 3.4. *Let G be a cyclic group of order N and Ψ be a faithful representation of G . Assume w is a non-Pompeiu weight function with respect to Ψ . Then the restriction of w to each $\mathbb{Z}_{\text{Rad}(N)}$ -coset is the weighted sum of characteristic functions of \mathbb{Z}_{p_i} -cosets, where $\text{Rad}(N)$ denotes the square-free radical of N .*

We will consider

$$\mathbb{Z}_{\prod_{i=1}^d p_i} \cong \prod_{i=1}^d \mathbb{Z}_{p_i}$$

as a grid in \mathbb{R}^d , whose points have integer coordinates. More precisely for $\mathbb{Z}_{\prod_{i=1}^d p_i}$ we assign

$$\mathcal{G} = \{x \in \mathbb{Z}^d : 0 \leq x_i \leq p_i - 1 \text{ for } 1 \leq i \leq d\},$$

where x_i denotes the i -th coordinate of x . The cosets of \mathbb{Z}_{p_i} coincide with collections of parallel line segments (containing p_i grid points of \mathcal{G}). A d -dimensional grid-cuboid will be a collection of 2^d grid points that resembles a d -dimensional cuboid in \mathbb{R}^d . Let $P \subset \mathcal{G}$ be a d -dimensional grid-cuboid and fix a point $y \in P$. For a point of $z \in P$ let $\pi(z)$ denote the Hamming distance between z and y . Note that w can also be considered as a function from \mathcal{G} to \mathbb{Q} .

The following statement makes the Pompeiu property for weight functions easily recognizable.

Proposition 3.5. *Let w be a non-Pompeiu weight function on the set $\mathbb{Z}_{\prod_{i=1}^d p_i}$, where p_i are mutually different primes. If w is the weighted sum of characteristic functions of \mathbb{Z}_{p_i} -cosets, then for every d -dimensional grid-cuboid P we have*

$$\sum_{c \in P} (-1)^{\pi(c)} w(c) = 0. \tag{6}$$

Proof. It is easy to see that each coset of \mathbb{Z}_{p_i} for any $p_i \mid n$ contains either 2 or 0 elements of the cuboid P . Substituting the characteristic function of any coset of \mathbb{Z}_{p_i} as a weight function into the left-hand side of (6), it is clearly reduced to a sum of at most two elements with different signs; thus (6) holds. \square

Remark 3.6. The converse of the previous statement also holds. We leave it to the reader to work out the details of the proof.

Now we describe a few special cases which will be later used for the proof of Theorem 2.5. In the proof of the next proposition we use the following definition.

Definition 3.7. Let $S \subseteq \mathbb{Z}_N$. For every $j \in \mathbb{Z}$ and $d \mid N$, we define the subsets

$$S_{j \bmod d} = \{s \in S : s \equiv j \pmod{d}\}.$$

Proposition 3.8. (a) *Every non-Pompeiu set in \mathbb{Z}_{pq} with respect to a faithful representation is either the union of cosets of \mathbb{Z}_p or those of \mathbb{Z}_q .*

(b) *Let $N = p^m q^n$ and let S be a non-Pompeiu multiset in \mathbb{Z}_N with respect to a faithful representation. Then there are polynomials $P(x), Q(x) \in \mathbb{Z}_{\geq 0}[x]$ such that*

$$S(x) \equiv P(x)\Phi_p(x^{N/p}) + Q(x)\Phi_q(x^{N/q}) \pmod{x^N - 1}.$$

Proof. (a) Let S be a non-Pompeiu set in \mathbb{Z}_{pq} with respect to a faithful representation and let w be the characteristic function of S . Using Proposition 3.2 we can write $w = \sum_{i=0}^{q-1} a_i 1_{\mathbb{Z}_p+i} + \sum_{j=0}^{p-1} b_j 1_{\mathbb{Z}_q+j}$, where $a_i, b_j \in \mathbb{Q}$. Then the range of w is $\text{Ran}(w) = \{a_i + b_j : 0 \leq i \leq p-1, 0 \leq j \leq q-1\}$. We have $\text{Ran}(w) = \{0, 1\}$. Thus there are at most two different a_i and two different b_j .

One can treat the case when a_i and b_j are constants as functions of i or j , respectively. Thus we may assume that $a_k < a_l$ for some $0 \leq k, l \leq p-1$. Then clearly $a_k + b_j = 0$ and $a_l + b_j = 1$ for all b_j ; in particular all b_j are the same. Therefore, we may write

$$w = b + \sum_{i=0}^{p-1} a_i 1_{\mathbb{Z}_p+i} = \sum_{i=0}^{p-1} (b + a_i) 1_{\mathbb{Z}_p+i},$$

finishing the proof of the statement.

(b) By Corollary 3.3, it is clear that

$$S(x) \equiv P(x)\Phi_p(x^{N/p}) + Q(x)\Phi_q(x^{N/q}) \pmod{x^N - 1}$$

for some $P(x), Q(x) \in \mathbb{Q}[x]$. Now we show that P and Q can be chosen such that $P(x), Q(x) \in \mathbb{Z}_{\geq 0}[x]$.

The subgroups \mathbb{Z}_p and \mathbb{Z}_q generate \mathbb{Z}_{pq} . Thus S can be written as the disjoint union

$$S = \bigcup_{k \in C} S_{k \pmod{N/(pq)}}$$

for $k = 0, \dots, N/(pq) - 1$, where k runs through a set of representatives C of the cosets of \mathbb{Z}_{pq} . Thus we are given

$$S_{k \pmod{N/(pq)}} = \sum_{a \in A} c_a(\mathbb{Z}_p + a) + \sum_{b \in B} d_b(\mathbb{Z}_q + b),$$

where $c_a + d_b \in \mathbb{Z}_{\geq 0}$ and A and B are sets of coset representatives of \mathbb{Z}_p and \mathbb{Z}_q , respectively, in $\mathbb{Z}_{pq} + k$. We want to modify the coefficients c_a and d_b such that they produce the same multiset and all of them remain nonnegative.

Let $e = c_a + d_b$ be one of the minimal weights of the multiset S . Then the values $d'_x = (c_a + d_x) - e$ are nonnegative for every $x \in B$, and let $c'_y = c_y + d_b$, which are nonnegative since these values are given by the multiset S only.

Now $c'_y + d'_x = ((c_a + d_x) - e) + c_y + d_b = c_y + d_x$ for every $x \in B$ and $y \in A$, finishing the proof of the lemma. □

4. Reduction (of Fuglede's conjecture)

Before we proceed to the proof of Theorem 2.5 we make a few general observations.

Lemma 4.1. *Let G be a finite abelian group. Assume that $S \subset G$ is a spectral set having Λ as a spectrum:*

- (a) $S + t$ is spectral with the same spectrum Λ for every $t \in G$.
- (b) $\Lambda + \omega$ is a spectrum for S for every $\omega \in G$.
- (c) S is a spectrum for Λ .

Proof. (a) If $\sum_{s \in S} \chi_\delta(s) = 0$ for some $\delta \in \Lambda - \Lambda$, then since χ_δ is a homomorphism, we have

$$\sum_{u \in (S+t)} \chi_\delta(u) = \chi_\delta(t) \sum_{s \in S} \chi_\delta(s) = 0.$$

Thus the orthogonality of the representations corresponding to the spectrum is preserved under translation.

(b) Similarly, the orthogonality of the representations corresponding to $\Lambda + \omega$ follows from the fact that $\Lambda - \Lambda = (\Lambda + \omega) - (\Lambda + \omega)$.

(c) This follows from the fact that a finite abelian group is canonically isomorphic to its double dual. □

Corollary 4.2. *It is enough to prove Theorem 2.5 for spectral sets S with $0 \in S$ and with spectrum Λ that contains 0.*

From now on we assume $0 \in S$ and $0 \in \Lambda$.

Lemma 4.3. *Let G be a finite abelian group and let S be spectral in G and such that it does not generate G . Assume that for every proper subgroup H of G we have $S - T(H)$. Then S tiles G .*

Proof. Let S be a spectral set with orthogonal basis $\{\chi_\lambda : \lambda \in \Lambda\} = \chi_\Lambda \subset \tilde{G}$ and let $\langle S \rangle = H < G$. Since every χ_λ is 1-dimensional, we have $\{\chi_{\lambda|_H} : \lambda \in \Lambda\} \subseteq \tilde{H}$ and clearly these are still orthogonal on S , since $S \subset H$. Then using that $S - T(H)$ holds, there is a set $T \subset H$ with $S + T = H$. Now let U be a complete set of coset representatives of G/H . Then we have $S + (T + U) = G$. \square

Now we prove a similar lemma reducing the possible structure of Λ .

Lemma 4.4. *Let G be a cyclic group of order N and let us suppose that $S - T(G/H)$ holds on every proper factor G/H . Let S be a spectral set of G and Λ be the corresponding spectrum. Assume that the intersection of the kernels of the elements of χ_Λ contains $H_{N/\ell} \neq 1$ for some $1 < \ell \mid N$. Then S tiles G .*

Proof. By our assumptions the elements of χ_Λ can be considered as irreducible representations of $G/H_{N/\ell}$ since their kernel is contained in $H_{N/\ell}$.

Let S_ℓ denote the multiset obtained as the image of S by the canonical projection π_ℓ of G to $G/H_{N/\ell} \cong H_\ell$. We claim that the multiset S_ℓ is a set in H_ℓ . Indeed there cannot be two elements of S in the same coset of $H_{N/\ell}$ since otherwise each element of χ_Λ would have the same value on them, contradicting the fact that these representations form a basis of the set of complex-valued functions on S . Thus S_ℓ is a set. Now it is easy to derive that $\Lambda/H_{N/\ell}$ is a spectrum with respect to S_ℓ in $G/H_{N/\ell}$ since $\chi_\lambda(\pi_\ell(s)) = \chi_\lambda(s)$ for every $s \in S$ and $\lambda \in \Lambda$.

We know $S - T(G/H_{N/\ell})$ holds. As S_ℓ is a spectral set in $G/H_{N/\ell}$ there is $T_\ell \subset G/H_{N/\ell}$ with $S_\ell + T_\ell = G/H_{N/\ell}$. Then if T is the preimage of T_ℓ under the canonical projection from G to $G/H_{N/\ell}$, then we have $S + T = G$. \square

Observation 4.5. Let us recall that $S(x)$ is the mask polynomial of the spectral set S . Note that for $\chi \in \tilde{G}$ of order k , we have $\sum_{s \in S} \chi(s) = 0$ is equivalent to the fact that a primitive k -th root of unity ξ_k is a root of $S(x)$. Since $\Phi_k(x)$ is irreducible over \mathbb{Q} we have $\Phi_k(x) \mid S(x)$; hence every primitive k -th root of unity is the root of $S(x)$ and $\sum_{s \in S} \chi'(s) = 0$ for every $\chi' \in \tilde{G}$ of the same order. If $\Lambda \subseteq G$ is a spectrum of S , the above can be summarized to

$$S(\xi_{\text{ord}(\lambda - \lambda')}) = 0 \tag{7}$$

for every $\lambda \neq \lambda'$ in a spectrum Λ , using (1).

The question whether our techniques can be generalized naturally arises. We point out here that in the next proposition we heavily use the assumption that the order of cyclic groups is divisible by at most two different primes.

Proposition 4.6. *Let G be a cyclic group of order $p^k q^\ell$ and let $|S| \geq 2$ be a spectral set. Assume further that Λ is a spectrum for S such that the elements of χ_Λ do not have a nontrivial common kernel. Then for every faithful representation ψ of G we have $\sum_{s \in S} \psi(s) = 0$.*

Proof. Note that by Observation 4.5, it is enough to prove the statement for one faithful representation.

Since the elements of χ_Λ do not have a common kernel we have a $\lambda_1 \in \Lambda$ with $p \nmid \lambda_1$. If $q \nmid \lambda_1$, then we are done so we assume $q \mid \lambda_1$. Similarly, we might assume that there exists $\lambda_2 \in \Lambda$ with $q \nmid \lambda_2$ and $p \mid \lambda_2$. In this case $\chi_{\lambda_1 - \lambda_2}$ generates \tilde{G} so we have $\sum_{s \in S} \chi_{\lambda_1 - \lambda_2}(s) = 0$. \square

This has the following interpretation in terms of mask polynomials.

Corollary 4.7. *Let (S, Λ) be a spectral pair in \mathbb{Z}_N , where $N = p^k q^\ell$, such that $0 \in S$, $0 \in \Lambda$, and each of S, Λ generates \mathbb{Z}_N . Then*

$$S(\xi_N) = \Lambda(\xi_N) = 0.$$

Proposition 4.8. *Let S be a spectral set in \mathbb{Z}_N and let p be a prime divisor of N . Assume that for every proper factor group \mathbb{Z}_N/H of \mathbb{Z}_N we have $S - T(\mathbb{Z}_N/H)$. Assume further that S is the disjoint union of cosets of \mathbb{Z}_p . Then S tiles \mathbb{Z}_N .*

Proof. By our assumptions $|S| = pr = |\Lambda|$ for some $r \in \mathbb{N}$ and Λ is a spectrum for S . Thus at least one of the cosets of H_p contains at least r elements of Λ . By Lemma 4.1(b)enumi we may assume that $|H_p \cap \Lambda| \geq r$. The elements $\chi_\Lambda \subseteq \tilde{H}_p$ are representations having a common kernel $\mathbb{Z}_p = H_N/p$. By our assumption S is the disjoint union of \mathbb{Z}_p -cosets, so it can be written as $\mathbb{Z}_p + B$ for some $B \subseteq \mathbb{Z}_N/\mathbb{Z}_p$. The representations in $\tilde{H}_p \cap \chi_\Lambda$ are constant on every coset of \mathbb{Z}_p . Hence for every $\chi_1 \neq \chi_2 \in \tilde{H}_p \cap \chi_\Lambda$ we have

$$\begin{aligned} 0 &= \sum_{s \in S} \chi_1(s) \bar{\chi}_2(s) = \sum_{s \in \mathbb{Z}_p + B} \chi_1(s) \bar{\chi}_2(s) = \sum_{t \in B} \sum_{x \in \mathbb{Z}_p} \chi_1(t+x) \bar{\chi}_2(t+x) \\ &= \sum_{t \in B} \sum_{x \in \mathbb{Z}_p} \chi_1(t) \chi_1(x) \bar{\chi}_2(t) \bar{\chi}_2(x) = \sum_{t \in B} p \chi_1(t) \bar{\chi}_2(t) = p \sum_{t \in B} \chi_1(t) \bar{\chi}_2(t), \end{aligned}$$

since the kernel of χ_1 and χ_2 contains \mathbb{Z}_p . Thus we obtain a set of $r = |B|$ representations of $\mathbb{Z}_N/\mathbb{Z}_p$, which are mutually orthogonal, hence forming a basis of $L^2(B)$. Thus B is a spectral set in $\mathbb{Z}_N/\mathbb{Z}_p$ and using our assumption we obtain that there exists T with $B + T = \mathbb{Z}_N/\mathbb{Z}_p$. So finally we get $S + T = (\mathbb{Z}_p + B) + T = \mathbb{Z}_p + (B + T) = \mathbb{Z}_p + \mathbb{Z}_N/\mathbb{Z}_p = \mathbb{Z}_N$. \square

Before we start to detail the proof of Theorem 2.5 we summarize what we have already proved in the previous sections about the structure of a spectral set S in $\mathbb{Z}_{p^n q^2}$. Note that we may assume by induction on n that $S - T(H)$ holds for every proper subgroup or factor H of $\mathbb{Z}_{p^n q^2}$. Indeed, Fuglede’s conjecture holds for \mathbb{Z}_{pq^2} and for $\mathbb{Z}_{p^n q}$ by [Malikiosis and Kolountzakis 2017], which corresponds to the base case of our induction.

If $|S| = 1$, then S is clearly a spectral set and also a tile. By Lemma 4.4 we might assume that the elements of χ_Λ do not have a common kernel so by Proposition 4.6 we might assume that $|S| \geq 2$ is a non-Pompeiu set with respect to a faithful representation of $\mathbb{Z}_{p^n q^2}$. Hence by Proposition 3.2 we have

$$S = \sum_{g \in A} u_g (\mathbb{Z}_p + g) + \sum_{h \in B} v_h (\mathbb{Z}_q + h),$$

where $u_g, v_h \in \mathbb{Q}$ and A and B are sets of coset representatives of \mathbb{Z}_p and \mathbb{Z}_q , respectively. Thus S is the weighted sum of cosets of \mathbb{Z}_p and \mathbb{Z}_q . Until now we have only seen that the weights are rational numbers. Now we prove that all weights are 0 or 1.

The subgroups \mathbb{Z}_p and \mathbb{Z}_q generate \mathbb{Z}_{pq} , so we write S as the disjoint union

$$S = \bigcup_{k \in C} S_{k \bmod N/(pq)},$$

where k runs through a set of representatives C of the cosets of \mathbb{Z}_{pq} for $k = 0, \dots, N/(pq) - 1$. Now

$$S_{k \bmod N/(pq)} = \sum_{\substack{g \in A \\ g + \mathbb{Z}_p \subset k + \mathbb{Z}_{pq}}} u_g(\mathbb{Z}_p + g) + \sum_{\substack{h \in B \\ h + \mathbb{Z}_q \subset k + \mathbb{Z}_{pq}}} v_h(\mathbb{Z}_q + h) \tag{8}$$

for every $k \in C$, so $S_{k \bmod N/(pq)}$ inherits its weights from S . Now it follows from Proposition 3.8 that in (8) $u_g = 0$ for every $g \in A$, $g + \mathbb{Z}_p \subset k + \mathbb{Z}_{pq}$ or $v_h = 0$ for every $h \in B$, $h + \mathbb{Z}_q \subset k + \mathbb{Z}_{pq}$. Since $S_{k \bmod N/(pq)}$ is a set, the remaining coefficients are 0 or 1. Then $S_{k \bmod N/(pq)}$ is the disjoint nontrivial union of \mathbb{Z}_p -cosets or \mathbb{Z}_q -cosets. Only one type appears for every fixed $k = 0, \dots, N/(pq) - 1$ except in the obvious case as follows:

It can happen that S contains a whole \mathbb{Z}_{pq} -coset, in which case it can be considered as the union of only \mathbb{Z}_p -cosets and only \mathbb{Z}_q -cosets as well. Thus S is the disjoint union of \mathbb{Z}_p -cosets and \mathbb{Z}_q -cosets.

Beside the case when S contains both \mathbb{Z}_p -cosets and \mathbb{Z}_q -cosets, by Proposition 4.8 we are done. Thus, we may assume S contains both \mathbb{Z}_p -cosets and \mathbb{Z}_q -cosets; we shall call such sets *nontrivial unions* of \mathbb{Z}_p - and \mathbb{Z}_q -cosets, to emphasize that they cannot be expressed as unions consisting solely of \mathbb{Z}_p -cosets, or \mathbb{Z}_q -cosets.

The above also follows from Corollary 4.7 and the structure of vanishing sums of roots of unity of order N , where N has at most two distinct prime factors [Lam and Leung 2000]. We added also a condition that shows when such a vanishing sum corresponds to a nontrivial union of \mathbb{Z}_p - and \mathbb{Z}_q -cosets, which is a consequence of Corollary 3.3 and Proposition 3.8(b)enumi or alternatively of Proposition 2.6 in [Malikiosis and Kolountzakis 2017].

Theorem 4.9. *Let $F(x) \in \mathbb{Z}_{\geq 0}[x]$ and $N = p^m q^n$, where p, q are different primes. Then, $F(\xi_N) = 0$ if and only if*

$$F(x) \equiv P(x)\Phi_p(x^{N/p}) + Q(x)\Phi_q(x^{N/q}) \pmod{(x^N - 1)}$$

for some $P(x), Q(x) \in \mathbb{Z}_{\geq 0}[x]$. If $F(\xi_N^{p^k}) \neq 0$ for some $1 \leq k \leq m$, then we cannot have $P(x) \equiv 0 \pmod{(x^N - 1)}$, and if $F(\xi_N^{q^\ell}) \neq 0$ for some $1 \leq \ell \leq n$, then we cannot have $Q(x) \equiv 0 \pmod{(x^N - 1)}$.

We will repeatedly use the above in Section 6 in order to obtain information about the structure of S and Λ from the vanishing of their mask polynomials on various N -th roots of unity. Regarding the case when S is a union of \mathbb{Z}_p -cosets (or \mathbb{Z}_q -cosets), there is a characterization in terms of the mask polynomial. This follows from a special case of Ma’s lemma [1985], see also [Schmidt 2002, Lemma 1.5.1] or [Pott 1995, Corollary 1.2.14], adapted to the cyclic case, using the polynomial notation.

Lemma 4.10. *Suppose that $S(x) \in \mathbb{Z}[x]$, and let \mathbb{Z}_N be a cyclic group such that $p^m \mid N$, but $p^{m+1} \nmid N$. If $S(\xi_d) = 0$ for every $p^m \mid d \mid N$, then*

$$S(x) \equiv P(x)\Phi_p(x^{N/p}) \pmod{(x^N - 1)}.$$

If the coefficients of S are nonnegative, then P can be taken with nonnegative coefficients as well. In particular, if $S \subseteq \mathbb{Z}_N$ satisfies $S(\xi_d) = 0$ for every $p^m \mid d \mid N$, then S is a union of \mathbb{Z}_p -cosets.

We summarize the reductions made so far in the following list.

Reduction 1. We may assume that a spectral set $S \subset \mathbb{Z}_{p^n q^2}$, along with a spectrum Λ , has the following structure:

- (a) $0 \in S$, $0 \in \Lambda$ and each of S and Λ generates $\mathbb{Z}_{p^n q^2}$.
- (b) Both S and Λ can be written as the disjoint nontrivial union of \mathbb{Z}_p -cosets and \mathbb{Z}_q -cosets and this holds for $S \cap (\mathbb{Z}_{pq} + g)$ and $\Lambda \cap (\mathbb{Z}_{pq} + h)$ for every $g, h \in \mathbb{Z}_{p^n q^2}$ as well.
- (c) There is a \mathbb{Z}_{pq} -coset which intersects S and its complement. Further the intersection is the union of \mathbb{Z}_p -cosets. The same holds for another \mathbb{Z}_{pq} -coset with \mathbb{Z}_q -cosets as well.
- (d) Fuglede’s conjecture holds for all \mathbb{Z}_M , with $M \mid p^n q^2$, $M < p^n q^2$ (induction assumption).

Proof. (a) This follows from Lemmas 4.1 and 4.3.

(b) This is an immediate consequence of part (a), Propositions 3.2 and 3.8 and Corollary 4.7.

(c) This follows from Proposition 4.8.

(d) It was proved in [Malikiosis and Kolountzakis 2017] that Fuglede’s conjecture holds for $N = p^n q$, and also for $N = pq^2$, so the given statement certainly holds for $p^2 q^2$, which is the base case for the inductive argument. □

Now we turn to the main tool already used in [Malikiosis and Kolountzakis 2017] to prove that a spectral set tiles $\mathbb{Z}_{p^n q^2}$. Clearly, sets coincide with mask polynomials having only coefficients 0 and 1. The following theorem was proved in [Coven and Meyerowitz 1999]. Let H_S be the set of prime powers r^a dividing N such that $\Phi_{r^a}(x) \mid S(x)$.

Theorem 4.11. *If $S \subset \mathbb{Z}_N$ satisfies the following two conditions (T1) and (T2), then S tiles \mathbb{Z}_N .*

(T1) $S(1) = \prod_{d \in H_S} \Phi_d(1)$.

(T2) *For pairwise relative prime elements s_i of H_S , we have $\Phi_{\prod s_i} \mid S(x)$.*

Note that $\Phi_{p^a}(1) = p$ for a prime p and $\Phi_k(1) = 1$ if k has at least two different prime divisors.

5. Preliminary lemmas

We introduce extra notation for divisibility. Fix $N \in \mathbb{N}$. For a natural number k we write $\ell \parallel_N k$ if ℓ is the largest divisor of N , which divides k . In our case N will be $p^n q^2$ so we simply write $\ell \parallel k$.

We review first (1) and (7) for a spectral pair (S, Λ) in \mathbb{Z}_N . First, we define as usual

$$\mathbb{Z}_N^* = \{g \in \mathbb{Z}_N : \gcd(g, N) = 1\},$$

the group of *reduced residues* mod N . It is precisely the subset of elements of N of order exactly N . Similarly, the subset of \mathbb{Z}_N of elements of order N/d , where $d \mid N$, is

$$d\mathbb{Z}_N^* = \{g \in \mathbb{Z}_N : \gcd(g, N) = d\}.$$

The zero set

$$Z(S) = \{d \in \mathbb{Z}_N : S(\xi_N^d) = 0\}$$

is then a union of subsets of the form $d\mathbb{Z}_N^*$ for some $d \mid N$, and (1) and (7) can be rewritten as

$$\Lambda - \Lambda \subseteq \{0\} \cup \bigcup_{d \mid N, S(\xi_N^d)=0} d\mathbb{Z}_N^*. \tag{9}$$

Of course, by Lemma 4.1(c)enumi, the roles of S and Λ can be reversed.

Definition 5.1. Let $S \subseteq \mathbb{Z}_N$. Recall that for every $j \in \mathbb{Z}$ and $d \mid N$, we define the subsets

$$S_{j \bmod d} = \{s \in S : s \equiv j \pmod{d}\}.$$

We say that S is *equidistributed mod d* if

$$|S_{j \bmod d}| = \frac{1}{d}|S|$$

for every j . Equivalently, every $\mathbb{Z}_{N/d}$ -coset of \mathbb{Z}_N contains the same number of elements of S .

Lemma 5.2. (a) *Assume $\Phi_p(x) \mid S(x)$. Then every $\mathbb{Z}_{N/p}$ -coset of \mathbb{Z}_N contains the same number of elements of S .*

(b) *Assume $\Phi_k(x) \mid S(x)$ for every $1 < k \mid d$. Then every $\mathbb{Z}_{N/d}$ -coset of \mathbb{Z}_N contains the same number of elements of S .*

Proof. (a) $\Phi_p(x) \mid S(x)$ is equivalent to the fact that S is a non-Pompeiu set with respect to an irreducible representation of order p , whose kernel is $\mathbb{Z}_{N/p}$. It is easy to see that a non-Pompeiu multiset on \mathbb{Z}_p has to be constant, and we obtain the result.

(b) Consider the formula

$$S(x) \equiv \sum_{j=0}^{d-1} |S_{j \bmod d}| x^j \pmod{x^d - 1}, \tag{10}$$

which holds for every $S \subseteq \mathbb{Z}_N$. It holds that $S(\xi_k) = 0$ for every $1 < k \mid d$ if and only if

$$1 + x + \dots + x^{d-1} = \prod_{1 < k \mid d} \Phi_k(x) \mid S(x),$$

or equivalently $S(x) = (1 + x + \dots + x^{d-1})G(x)$, where $G(x) \in \mathbb{Z}[x]$. The latter implies

$$S(x) \equiv (1 + x + \dots + x^{d-1})G(1) \pmod{x^d - 1},$$

so by (10) we get $|S_{j \bmod d}| = G(1)$ for all j . Conversely, if $|S_{j \bmod d}| = c$ for all j , then

$$S(x) \equiv c(1 + x + \dots + x^{d-1}) \pmod{x^d - 1},$$

due to (10), which easily gives $S(\xi_k) = 0$ for every $1 < k \mid d$, as desired. □

Let (S, Λ) be a spectral pair in \mathbb{Z}_N satisfying the conditions of Reduction 1, where $N = p^n q^2$. An immediate consequence of Reduction 1(c) is that $S - S$ contains the difference set of both a \mathbb{Z}_p -coset and a \mathbb{Z}_q -coset; thus

$$\frac{N}{p}\mathbb{Z}_N \cup \frac{N}{q}\mathbb{Z}_N \subseteq S - S,$$

whence

$$\Lambda(\xi_p) = \Lambda(\xi_q) = 0, \tag{11}$$

by (7), and we obtain in particular,

$$|\Lambda_{i \bmod p}| = \frac{1}{p}|\Lambda|, \quad |\Lambda_{j \bmod q}| = \frac{1}{q}|\Lambda| \tag{12}$$

for all i, j , by Lemma 5.2. This shows that pq divides $|S| = |\Lambda|$.

6. Proof of Theorem 2.5

A significant special case will be shown first.

Lemma 6.1. *Let $S \subseteq \mathbb{Z}_N$ be spectral. If $q^2 \mid |S|$, then S tiles \mathbb{Z}_N .*

Proof. Let $H_S(p) = \{p^m : S(\xi_{p^m}) = 0, 1 \leq m \leq n\}$, and similarly define $H_\Lambda(p)$ for a spectrum $\Lambda \subseteq \mathbb{Z}_N$. Suppose that

$$H_\Lambda(p) = \{p^{m_1}, \dots, p^{m_k}\},$$

where $1 \leq m_1 < m_2 < \dots < m_k \leq n$. For every j ,

$$S_{j \bmod q^2} - S_{j \bmod q^2} \subseteq (S - S) \cap q^2\mathbb{Z}_N \subseteq \{0\} \cup \bigcup_{i=0}^k \frac{N}{p^{m_i}}\mathbb{Z}_N^*, \tag{13}$$

by (9). Consider the p -adic expansion of every $s \in S$ taken mod p^n , as follows:

$$s \equiv s_0 + s_1 p + \dots + s_{n-1} p^{n-1} \pmod{p^n}, \quad 0 \leq s_i \leq p - 1, \quad 0 \leq i \leq n - 1.$$

Due to (13), the elements of each $S_{j \bmod q^2}$ cannot have the same p -adic digits corresponding to p^{n-m_i} , $1 \leq i \leq k$, yielding

$$|S_{j \bmod q^2}| \leq p^k, \quad 0 \leq j < q^2;$$

thus, $|S| \leq p^k q^2$. On the other hand, we have

$$\prod_{i=1}^k \Phi_{p^{m_i}}(x) \mid \Lambda(x),$$

and putting $x = 1$ we obtain $p^k \mid |\Lambda|$; we then get by hypothesis $p^k q^2 \mid |S|$, whence $|S| = p^k q^2$, and

$$|S_{j \bmod q^2}| = p^k, \quad 0 \leq j < q^2.$$

Since S is equidistributed mod q^2 , we must also have $S(\xi_q) = S(\xi_{q^2}) = 0$ by Lemma 5.2. We note that each element of $S_{j \bmod q^2}$ is unique mod p^n , so the reduction mod p^n map

$$\pi : \mathbb{Z}_N \mapsto \mathbb{Z}_{p^n}$$

is injective on each $S_j \bmod p^n$; fix some j , and let $\pi(S_j \bmod p^n) = S'$. Since $q^2 \mid s - s'$ for every $s, s' \in S_j \bmod q^2$, we conclude that the order of $s - s'$ in \mathbb{Z}_N is the same as the order of $\pi(s - s')$ in \mathbb{Z}_{p^n} , which gives

$$S' - S' \subseteq \{0\} \cup \bigcup_{i=0}^k p^{n-m_i} \mathbb{Z}_{p^n}^*.$$

Consider now the set $\Lambda' \subseteq \mathbb{Z}_{p^n}$ whose mask polynomial is given by

$$\Lambda'(x) \equiv \prod_{i=1}^k \Phi_{p^{m_i}}(x) \bmod (x^{p^n} - 1).$$

We have $|S'| = |\Lambda'| = p^k$ and

$$S' - S' \subseteq \{0\} \cup \{d \in \mathbb{Z}_{p^n} : \Lambda'(\xi_{p^n}^d) = 0\};$$

therefore, (S', Λ') is a spectral pair in \mathbb{Z}_{p^n} by (9). Since

$$\Phi_{p^{m_i}}(x) = 1 + x^{p^{m_i-1}} + x^{2p^{m_i-1}} + \dots + x^{(p-1)p^{m_i-1}},$$

we obtain

$$(\Lambda' - \Lambda') \cap p^{n-m_i+1} \mathbb{Z}_{p^n}^* \neq \emptyset, \quad 1 \leq i \leq k;$$

therefore,

$$\bigcup_{i=0}^k p^{n-m_i+1} \mathbb{Z}_{p^n}^* \subseteq \{d \in \mathbb{Z}_{p^n} : S'(\xi_{p^n}^d) = 0\},$$

by (9), or equivalently

$$\prod_{i=0}^k \Phi_{p^{n-m_i+1}}(x) \mid S_j \bmod q^2(x),$$

since

$$S_j \bmod q^2(x) \equiv S'(x) \bmod (x^{p^n} - 1).$$

Moreover, by $S(x) = \sum_{j=0}^{q^2-1} S_j \bmod q^2(x)$ and $|S| = p^k q^2$, we conclude that

$$H_S = \{p^{n-m_k+1}, \dots, p^{n-m_1+1}, q, q^2\};$$

hence S satisfies (T1).

Consider next the polynomial $F(X)$ satisfying

$$S_j \bmod q^2(x) \equiv x^j F(x^{q^2}) \bmod (x^N - 1)$$

for a fixed j . Since $\Phi_{p^{n-m_i+1}}(x) \mid F(x^{q^2})$ for all $1 \leq i \leq k$ and q^2 is prime to p^{n-m_i+1} , we also get that $\Phi_{p^{n-m_i+1}}(x) \mid F(x)$. Therefore, for $\ell = 1$ or 2 we get

$$S_j \bmod q^2(\xi_{p^{n-m_i+1}q^\ell}) = \xi_{p^{n-m_i+1}q^\ell}^j F(\xi_{p^{n-m_i+1}q^\ell}^{q^2}) = \xi_{p^{n-m_i+1}q^\ell}^j F(\xi_{p^{n-m_i+1}}^{q^{2-\ell}}) = 0$$

for all j , which shows that S satisfies (T2). □

We distinguish now the following cases:

Case 1: $S(\xi_N^q) = S(\xi_N^{q^2}) = 0$. Since $S(\xi_N) = 0$ by Corollary 4.7, S is a union of \mathbb{Z}_p -cosets by Lemma 4.10 and S tiles due to Reduction 1(c).

Case 2: $S(\xi_N^q)S(\xi_N^{q^2}) \neq 0$. Consider the difference sets $\Lambda_{j \bmod q} - \Lambda_{j \bmod q}$. They are always subsets of $(\Lambda - \Lambda) \cap q\mathbb{Z}_N$, but since they avoid $q\mathbb{Z}_N^* \cup q^2\mathbb{Z}_N^*$ in this case by (9), we get

$$\Lambda_{j \bmod q} - \Lambda_{j \bmod q} \subseteq pq\mathbb{Z}_N$$

for all j . This shows that every element of $\Lambda_{j \bmod q}$ has the same remainder mod p , or equivalently, for every j there is an $i = i(j)$ such that

$$\Lambda_{j \bmod q} \subseteq \Lambda_{i(j) \bmod p}.$$

This, in particular, shows that $p < q$, and that every $\Lambda_{i \bmod p}$ is the disjoint union of sets of the form $\Lambda_{j \bmod q}$, namely

$$\Lambda_{i \bmod p} = \bigcup_{i(j)=i} \Lambda_{j \bmod q}.$$

Suppose that the number of sets appearing in the union are ℓ . Then, the above equation along with (12) implies $1/p = \ell/q$, which leads to a contradiction (no such spectrum can exist).

Case 3: $S(\xi_N^q) = 0 \neq S(\xi_N^{q^2})$. We apply Theorem 4.9 to $S(x) \bmod (x^{N/q} - 1)$. We obtain

$$S(x) \equiv P(x)\Phi_p(x^{N/(pq)}) + Q(x)\Phi_q(x^{N/q^2}) \bmod (x^{N/q} - 1),$$

since $S(\xi_{N/q}) = 0$, where $P(x)$ and $Q(x)$ have nonnegative coefficients. Furthermore, since $S(\xi_N^{q^2}) \neq 0$, we cannot have $Q \equiv 0$. Due to the nonnegativity of P and Q , we obtain the existence of $s, s' \in S$ such that

$$s - s' \equiv \frac{N}{q^2} \bmod \frac{N}{q};$$

hence $p^n \mid s - s'$ but $q \nmid s - s'$, yielding $s - s' \in p^n\mathbb{Z}_N^*$ and

$$\Lambda(\xi_{q^2}) = 0,$$

which further gives $q^2 \mid |\Lambda|$, so by Lemma 6.1, S tiles \mathbb{Z}_N .

Case 4: $S(\xi_N^q) \neq 0 = S(\xi_N^{q^2})$. We will prove the following:

Claim 1. $(S - S) \cap \frac{N}{pq^2}\mathbb{Z}_N^* \neq \emptyset$.

Proof of Claim. By Theorem 4.9, the multiset³ q^2S is a union of \mathbb{Z}_p -cosets, or equivalently

$$|S_{i \bmod p^n}| = |S_{i+kp^{n-1} \bmod p^n}| \tag{14}$$

³Here, we consider the elements $q^2s \bmod N$, $s \in S$, counting multiplicities. For example, if $N = 4$ and $S = \{0, 2\}$, then $2S$ is the multiset whose only element is 0, appearing with multiplicity 2.

for every i, k . We partition the above sets mod $p^n q$:

$$S_i \bmod p^n = \bigcup_{\ell=0}^{q-1} S_{i+\ell p^n} \bmod p^n q \quad \text{and} \quad S_{i+kp^{n-1}q} \bmod p^n = \bigcup_{\ell=0}^{q-1} S_{i+kp^{n-1}q+\ell p^n} \bmod p^n q.$$

If for every i there existed some ℓ such that

$$S_{i+kp^{n-1}q} \bmod p^n = S_{i+kp^{n-1}q+\ell p^n} \bmod p^n q$$

for every k , then qS would also be a union of \mathbb{Z}_p -cosets. Indeed, as for every i there is at most one value of $0 \leq \ell \leq q - 1$ such that $S_{i+\ell p^n} \bmod p^n q \neq \emptyset$, and by the above condition the cardinalities of $S_{i+kp^{n-1}q+\ell p^n} \bmod p^n q$ are the same for $0 \leq k \leq p - 1$. Therefore, $S(\xi_{p^n q}) = 0$ by Theorem 4.9 (or equivalently by Proposition 3.2), contradicting the hypothesis. Thus, there exists i such that there are nonempty $S_{i+\ell p^n} \bmod p^n q$ and $S_{i+\ell' p^n} \bmod p^n q$, with $0 \leq \ell < \ell' \leq q - 1$. Clearly, $S_{i+\ell p^n} \bmod p^n q \subseteq S_i \bmod p^n$, so $S_i \bmod p^n$ is nonempty. Using (14) we have $S_{i+p^{n-1}} \bmod p^n$ is nonempty.

Now let $s \in S_{i+p^{n-1}} \bmod p^n$, $s' \in S_{i+\ell p^n} \bmod p^n q$ and $s'' \in S_{i+\ell' p^n} \bmod p^n q$, so that $p^{n-1} \parallel s - s'$ and $p^{n-1} \parallel s - s''$. Since $s'' - s' \equiv (\ell' - \ell)p^n \bmod p^n q$, we get $q \nmid s'' - s'$, so either $q \nmid s - s'$ or $q \nmid s - s''$ would hold, yielding $(S - S) \cap p^{n-1} \mathbb{Z}_N^* \neq \emptyset$, as desired. \square

This implies

$$\Lambda(\xi_{pq^2}) = 0, \tag{15}$$

by (7). If $\Lambda(\xi_{q^2}) = 0$ then we would have $q^2 \mid |S|$ and S would tile \mathbb{Z}_N by virtue of Lemma 6.1. So, we may assume $\Lambda(\xi_{q^2}) \neq 0$.

By (15) and Theorem 4.9 we get

$$\Lambda(x) \equiv \sum_{j=0}^{pq^2-1} |\Lambda_j \bmod pq^2| x^j \equiv P(x)\Phi_p(x^{N/p}) + Q(x)\Phi_q(x^{N/q}) \bmod (x^{pq^2} - 1)$$

for some $P(x), Q(x) \in \mathbb{Z}_{\geq 0}[x]$ and $P(x) \neq 0$ by $\Lambda(\xi_{q^2}) \neq 0$. We note that the function $f(j) = |\Lambda_j \bmod pq^2|$ restricted on a \mathbb{Z}_{pq} -coset of \mathbb{Z}_{pq^2} is supported either on a \mathbb{Z}_p -coset or a \mathbb{Z}_q -coset; otherwise, there would exist $\lambda \in \Lambda_j \bmod pq^2$ and $\lambda' \in \Lambda_{j'} \bmod pq^2$, where j, j' satisfy

$$j - j' \in q\mathbb{Z}_{pq^2}^*.$$

This shows that $q \parallel \lambda - \lambda'$ and $p \nmid \lambda - \lambda'$; thus $\lambda - \lambda' \in q\mathbb{Z}_N^*$ and $S(\xi_N^q) = 0$ by (9), contradicting the hypothesis.

Next, we consider a nonempty subset $\Lambda_j \bmod pq^2$; the polynomials with nonnegative coefficients $P(x)\Phi_p(x^{N/p})$ and $Q(x)\Phi_q(x^{N/q})$ contribute to the coefficient of x^j of $\Lambda(x) \bmod (x^{pq^2} - 1)$. If both contributions are positive, then all subsets $\Lambda_{j+kq^2} \bmod pq^2$ and $\Lambda_{j+\ell pq} \bmod pq^2$ are nonempty for $0 < k < p$ and $0 < \ell < q$. Then, for $\lambda \in \Lambda_{j+q^2} \bmod pq^2$ and $\lambda' \in \Lambda_{j+pq} \bmod pq^2$, we have $q \parallel \lambda - \lambda'$; hence $\lambda - \lambda' \in q\mathbb{Z}_N^*$, which contradicts $S(\xi_N^q) \neq 0$, due to (7).

Let $\Gamma(x)$ be $\Lambda(x) \bmod (x^{pq^2} - 1)$. The previous argument shows that the coefficient of x^j of $\Gamma(x)$ is determined completely either from $P(x)\Phi_p(x^{N/p})$ or $Q(x)\Phi_q(x^{N/q})$. Moreover, if $q \parallel j - j'$, then

we cannot have that both the coefficients of x^j and $x^{j'}$ in $\Gamma(x)$ are nonzero by the same argument. This means that $f(j) = |\Lambda_{j \bmod pq^2}|$ restricted on a \mathbb{Z}_{pq} -coset of \mathbb{Z}_{pq^2} is supported either on a \mathbb{Z}_p -coset or a \mathbb{Z}_q -coset and constant restricted to this coset.

This shows that for each j such that $\Lambda_{j \bmod pq^2} \neq \emptyset$, either

$$|\Lambda_{j+kq^2 \bmod pq^2}| = \frac{1}{p} |\Lambda_{j \bmod q^2}| = \frac{1}{p} |\Lambda_{j \bmod q}|, \quad 0 \leq k < p, \tag{16}$$

or

$$|\Lambda_{j+\ell pq \bmod pq^2}| = |\Lambda_{j+\ell pq \bmod q^2}| = \frac{1}{q} |\Lambda_{j \bmod q}|, \quad 0 \leq \ell < q, \tag{17}$$

holds. If (17) holds for some j , then $q^2 \mid |\Lambda|$, so by Lemma 6.1 we get that S tiles \mathbb{Z}_N . Therefore, we may assume that (16) holds for all j with $\Lambda_{j \bmod pq^2} \neq \emptyset$. For such j , we have

$$\Lambda_{j \bmod q} - \Lambda_{j \bmod q} = \Lambda_{j \bmod q^2} - \Lambda_{j \bmod q^2} \subseteq (\Lambda - \Lambda) \cap q^2 \mathbb{Z}_N;$$

hence $\Lambda - \Lambda$ completely avoids $q\mathbb{Z}_N \setminus q^2\mathbb{Z}_N$. On the other hand, $(S - S) \cap p^n \mathbb{Z}_N^* = \emptyset$ by (9) and the assumption $\Lambda(\xi_{q^2}) \neq 0$; hence the polynomials

$$\begin{aligned} \bar{S}(x) &\equiv S(x)\Phi_q(x^{p^n}) \bmod (x^N - 1), \\ \bar{\Lambda}(x) &\equiv \Lambda(x)\Phi_q(x^{N/q}) \bmod (x^N - 1) \end{aligned}$$

are mask polynomials of subsets of \mathbb{Z}_N , say \bar{S} and $\bar{\Lambda}$, i.e., their coefficients are 0 or 1. We claim that $(\bar{S}, \bar{\Lambda})$ is a spectral pair. They obviously have the same cardinality as $q|S|$, and an element of $\bar{\Lambda} - \bar{\Lambda}$ can be expressed as $\lambda - \lambda' + lN/q$, where $\lambda, \lambda' \in \Lambda$, $|l| < q$.

If $\lambda - \lambda' \in p^k \mathbb{Z}_N^*$, then $q \nmid \lambda - \lambda' + lN/q$; hence $\lambda - \lambda' + lN/q \in p^k \mathbb{Z}_N^*$ as well, yielding $S(\xi_N^{\lambda - \lambda' + lN/q}) = 0$, since $N/q = p^n q$.

The remaining case is $\lambda - \lambda' \in p^k q^2 \mathbb{Z}_N^*$, where $0 \leq k \leq n - 1$, as $\Lambda - \Lambda$ avoids $q\mathbb{Z}_N \setminus q^2\mathbb{Z}_N$. In this case, $\lambda - \lambda' + lN/p \in p^k q \mathbb{Z}_N^*$ if $1 \leq l \leq q - 1$, and $\Phi_q(\xi_{p^k q}^{p^n}) = \Phi_q(\xi_q^{p^{n-k}}) = 0$, so

$$\bar{S}(\xi_N^{\lambda - \lambda' + lN/p}) = 0.$$

If $l = 0$, then clearly $\lambda - \lambda' \in \Lambda - \Lambda$. Considering all of these cases we have $\bar{\Lambda} - \bar{\Lambda} \subseteq \{0\} \cup Z(\bar{S})$, proving that the pair $(\bar{S}, \bar{\Lambda})$ is spectral by virtue of (9). Since $q^2 \mid \bar{S}$ we have \bar{S} tiles \mathbb{Z}_N by Lemma 6.1; thus there is $T \subseteq \mathbb{Z}_N$ such that

$$S(x)\Phi_q(x^{p^n})T(x) \equiv \bar{S}(x)T(x) \equiv 1 + x + \dots + x^{N-1} \bmod (x^N - 1),$$

so $\Phi_q(x^{p^n})T(x)$ is the mask polynomial of a tiling complement of S using Lemma 1.3 in [Coven and Meyerowitz 1999], completing the proof. □

Appendix

Theorem A.1. *Let S be a subset of \mathbb{Z}_p^2 . Then S tiles \mathbb{Z}_p^2 if and only if S is spectral.*

Iosevich et al. [Iosevich et al. 2017] has already proved this theorem, but we provide an easy combinatorial proof for one of the two directions and a short one for the other direction using Theorem 2.1.

Proposition A.2. *Let S be a spectral set of \mathbb{Z}_p^2 . Then S tiles \mathbb{Z}_p^2 .*

Proof. Let S be a spectral set. We may assume $|S| > 1$, since one-element sets clearly tile every group. The corresponding spectrum Λ is also of size at least 2. Then there is a nontrivial irreducible representation ψ of \mathbb{Z}_p^2 such that $\sum_{s \in S} \psi(s) = 0$. We may also assume $|S| = |\Lambda| < p^2$.

Representations of \mathbb{Z}_p^2 can be parametrized by the elements of \mathbb{Z}_p^2 . For $u \in \mathbb{Z}_p^2$ let $\chi_u(v) = e^{2\pi i \langle u, v \rangle / p}$, where the scalar product of $\langle u, v \rangle$ is taken modulo p . This can be written as $\sum_{j=0}^{p-1} a_j e^{(2\pi i / p)j}$, where the a_j are integers which are determined in the following way.

From now on we may also think of \mathbb{Z}_p^2 as a 2-dimensional vector space over \mathbb{Z}_p . Cosets of 1-dimensional subspaces are called lines. Let u' be a nonzero vector orthogonal to u . Then $\langle u, v \rangle$ is constant on every coset of the subgroup generated by u' . Basically, we count the intersection of S with the elements the set of lines parallel with $\langle u' \rangle$. We have $\sum_{j=0}^{p-1} a_j e^{(2\pi i / p)j} = 0$ if and only if a_j is a constant sequence so every element of this class of parallel lines contains the same number of elements of S (i.e., S is equidistributed on these set of parallel lines). As a consequence we get that $p \mid |S|$.

If $|S| = p$, then by the previous argument we have that S intersects each element of a class of parallel lines once. Then S clearly tiles \mathbb{Z}_p^2 .

Thus we may assume $p + 1 < 2p \leq |\Lambda| = |S| < p^2$. It is enough to show that such spectral set does not exist. Each class of parallel lines consists of p lines. Thus we have that for every class of parallel lines, at least one line contains at least two elements of Λ . Thus using the argument above we have that every element of every class of parallel lines contains the same amount of elements of S . This means that every line contains k elements of S for some fixed number $p > k \geq 2$.

Let $x \in \mathbb{Z}_p^2 \setminus S$. Take every line containing x . These lines give a disjoint cover of $\mathbb{Z}_p^2 \setminus \{x\}$. Since each of them contains $k \geq 2$ elements we have $|S| = (p + 1)k$, which is not divisible by p , a contradiction. \square

Proposition A.3. *Let S be a set in \mathbb{Z}_p^2 , which tiles. Then S is spectral.*

Proof. We may assume that $1 < |S| < p^2$. Then S and its tiling complements are of cardinality p . Using Theorem 2.1 we obtain that either S or T is a subgroup of \mathbb{Z}_p^2 .

Subgroups are clearly spectral sets. If T is a subgroup, then S is a complete set of coset representatives of T . Let U denote the subgroup of \mathbb{Z}_p^2 consisting of vectors orthogonal to T . Then clearly S is equidistributed on the orthogonal lines for $u \in U$ so U is a spectrum for S . \square

Acknowledgements

The authors wish to thank the anonymous referees, as well as the editor, Professor Izabella Łaba, for improving the quality of our paper with their suggestions. The research of Kiss was supported by the Hungarian National Foundation for Scientific Research, grant no. K124749, internal research project R-AGR-0500 and the Premium Postdoctoral Fellowship of Hungarian Academy of Science. The research of Somlai was supported by project no. ED 181-2019-0030 (Application-specific highly reliable IT solutions) which has been implemented with the support provided from the National Research, Development and Innovation Fund of Hungary, financed under the Thematic Excellence Programme funding scheme. The research of Vizer was supported by the National Research, Development and

Innovation Office - NKFIH under the grants nos. FK 132060, KH130371 and SNN 129364, by the János Bolyai Research Fellowship and by the New National Excellence Program under the grant number no. ÚNKP-19-4-BME-287.

References

- [Babai 1979] L. Babai, “Spectra of Cayley graphs”, *J. Combin. Theory Ser. B* **27**:2 (1979), 180–189. MR Zbl
- [Brown et al. 1973] L. Brown, B. M. Schreiber, and B. A. Taylor, “Spectral synthesis and the Pompeiu problem”, *Ann. Inst. Fourier (Grenoble)* **23**:3 (1973), 125–154. MR Zbl
- [Chakalov 1944] L. Chakalov, “Sur un problème de D. Pompéiu”, *Ann. [Godishnik] Univ. Sofia. Fac. Phys.-Math. Livre I* **40** (1944), 1–14. In Bulgarian. MR Zbl
- [Coven and Meyerowitz 1999] E. M. Coven and A. Meyerowitz, “Tiling the integers with translates of one finite set”, *J. Algebra* **212**:1 (1999), 161–174. MR Zbl
- [Dutkay and Lai 2014] D. E. Dutkay and C.-K. Lai, “Some reductions of the spectral set conjecture to integers”, *Math. Proc. Cambridge Philos. Soc.* **156**:1 (2014), 123–135. MR Zbl
- [Fan et al. 2019] A. Fan, S. Fan, L. Liao, and R. Shi, “Fuglede’s conjecture holds in \mathbb{Q}_p ”, *Math. Ann.* **375**:1-2 (2019), 315–341. MR Zbl
- [Farkas et al. 2006] B. Farkas, M. Matolcsi, and P. Móra, “On Fuglede’s conjecture and the existence of universal spectra”, *J. Fourier Anal. Appl.* **12**:5 (2006), 483–494. MR Zbl
- [Fuglede 1974] B. Fuglede, “Commuting self-adjoint partial differential operators and a group theoretic problem”, *J. Funct. Anal.* **16**:1 (1974), 101–121. MR Zbl
- [Garofalo 1989] N. Garofalo, “A new result on the Pompeiu problem”, *Rend. Sem. Mat. Univ. Politec. Torino* **47**:special issue (1989), 25–38. MR Zbl
- [Iosevich et al. 1999] A. Iosevich, N. Katz, and S. Pedersen, “Fourier bases and a distance problem of Erdős”, *Math. Res. Lett.* **6**:2 (1999), 251–255. MR Zbl
- [Iosevich et al. 2003] A. Iosevich, N. Katz, and T. Tao, “The Fuglede spectral conjecture holds for convex planar domains”, *Math. Res. Lett.* **10**:5 (2003), 559–569. MR Zbl
- [Iosevich et al. 2017] A. Iosevich, A. Mayeli, and J. Pakianathan, “The Fuglede conjecture holds in $\mathbb{Z}_p \times \mathbb{Z}_p$ ”, *Anal. PDE* **10**:4 (2017), 757–764. MR Zbl
- [Kiss et al. 2018] G. Kiss, M. Laczkovich, and C. Vincze, “The discrete Pompeiu problem on the plane”, *Monatsh. Math.* **186**:2 (2018), 299–314. MR Zbl
- [Kolountzakis and Matolcsi 2006a] M. N. Kolountzakis and M. Matolcsi, “Complex Hadamard matrices and the spectral set conjecture”, *Collect. Math.* **57**:special issue (2006), 281–291. MR Zbl
- [Kolountzakis and Matolcsi 2006b] M. N. Kolountzakis and M. Matolcsi, “Tiles with no spectra”, *Forum Math.* **18**:3 (2006), 519–528. MR Zbl
- [Łaba 2001] I. Łaba, “Fuglede’s conjecture for a union of two intervals”, *Proc. Amer. Math. Soc.* **129**:10 (2001), 2965–2972. MR Zbl
- [Łaba 2002] I. Łaba, “The spectral set conjecture and multiplicative properties of roots of polynomials”, *J. London Math. Soc. (2)* **65**:3 (2002), 661–671. MR Zbl
- [Lam and Leung 2000] T. Y. Lam and K. H. Leung, “On vanishing sums of roots of unity”, *J. Algebra* **224**:1 (2000), 91–109. MR Zbl
- [Ma 1985] S.-L. Ma, *Polynomial addition sets*, Ph.D. thesis, University of Hong Kong, 1985, available at <https://tinyurl.com/slmphd>.
- [Malikiosis and Kolountzakis 2017] R. D. Malikiosis and M. N. Kolountzakis, “Fuglede’s conjecture on cyclic groups of order $p^n q$ ”, *Discrete Anal.* **2017** (2017), art. id. 12. MR Zbl

- [Matolcsi 2005] M. Matolcsi, “Fuglede’s conjecture fails in dimension 4”, *Proc. Amer. Math. Soc.* **133**:10 (2005), 3021–3026. MR Zbl
- [Newman 1977] D. J. Newman, “Tesselation of integers”, *J. Number Theory* **9**:1 (1977), 107–111. MR Zbl
- [Pompeiu 1929] D. Pompeiu, “Sur une propriété intégrale des fonctions de deux variables réelles”, *Bull. Acad. Bruxelles* **5**:15 (1929), 265–269. Zbl
- [Pott 1995] A. Pott, *Finite geometry and character theory*, Lecture Notes in Math. **1601**, Springer, 1995. MR Zbl
- [Puls 2013] M. J. Puls, “The Pompeiu problem and discrete groups”, *Monatsh. Math.* **172**:3-4 (2013), 415–429. MR Zbl
- [Ramm 1997] A. G. Ramm, “The Pompeiu problem”, *Appl. Anal.* **64**:1-2 (1997), 19–26. MR Zbl
- [Ramm 2017] A. G. Ramm, “Solution to the Pompeiu problem and the related symmetry problem”, *Appl. Math. Lett.* **63** (2017), 28–33. MR Zbl
- [Rédei 1965] L. Rédei, “Die neue Theorie der endlichen abelschen Gruppen und Verallgemeinerung des Hauptsatzes von Hajós”, *Acta Math. Acad. Sci. Hungar.* **16**:3-4 (1965), 329–373. MR Zbl
- [Schmidt 2002] B. Schmidt, *Characters and cyclotomic fields in finite geometry*, Lecture Notes in Math. **1797**, Springer, 2002. MR Zbl
- [Shi 2019] R. Shi, “Fuglede’s conjecture holds on cyclic groups \mathbb{Z}_{pqr} ”, *Discrete Anal.* **2019** (2019), art. id. 14. MR Zbl
- [Steinberg 2012] B. Steinberg, *Representation theory of finite groups: an introductory approach*, Springer, 2012. MR Zbl
- [Székelyhidi 2001] L. Székelyhidi, “A Wiener Tauberian theorem on discrete abelian torsion groups”, *Ann. Acad. Paedagog. Crac. Stud. Math.* **4** (2001), 147–150. Zbl
- [Tao 2004] T. Tao, “Fuglede’s conjecture is false in 5 and higher dimensions”, *Math. Res. Lett.* **11**:2 (2004), 251–258. MR Zbl
- [Zalcman 1992] L. Zalcman, “A bibliographic survey of the Pompeiu problem”, pp. 185–194 in *Approximation by solutions of partial differential equations* (Hanstholm, Denmark, 1991), edited by B. Fuglede et al., NATO Adv. Sci. Inst. Ser. C **365**, Kluwer, Dordrecht, 1992. MR Zbl
- [Zeilberger 1978] D. Zeilberger, “Pompeiu’s problem on discrete space”, *Proc. Nat. Acad. Sci. U.S.A.* **75**:8 (1978), 3555–3556. MR Zbl

Received 10 Jul 2018. Revised 2 Mar 2019. Accepted 8 Apr 2019.

GERGELY KISS: kigergo57@gmail.com

Faculty of Science, Mathematics Research Unit, University of Luxembourg, Luxembourg, Luxembourg

ROMANOS DIOGENES MALIKIOSIS: rwwanos@gmail.com

Department of Mathematics, Aristotle University of Thessaloniki, Thessaloniki, Greece

GÁBOR SOMLAI: gsomlai@cs.elte.hu

MTA-ELTE Geometric and Algebraic Combinatorics Research Group NKFIH 115799 and, Faculty of Science, Institute of Mathematics, Eötvös Loránd University, Budapest, Hungary

MÁTÉ VIZER: vizermate@gmail.com

Alfréd Rényi Institute of Mathematics, Hungarian Academy of Sciences, Budapest, Hungary

Analysis & PDE

msp.org/apde

EDITORS

EDITOR-IN-CHIEF

Patrick Gérard

patrick.gerard@math.u-psud.fr

Université Paris Sud XI

Orsay, France

BOARD OF EDITORS

Massimiliano Berti	Scuola Intern. Sup. di Studi Avanzati, Italy berti@sissa.it	Gilles Pisier	Texas A&M University, and Paris 6 pisier@math.tamu.edu
Michael Christ	University of California, Berkeley, USA mchrist@math.berkeley.edu	Tristan Rivière	ETH, Switzerland riviere@math.ethz.ch
Charles Fefferman	Princeton University, USA cf@math.princeton.edu	Igor Rodnianski	Princeton University, USA irod@math.princeton.edu
Ursula Hamenstaedt	Universität Bonn, Germany ursula@math.uni-bonn.de	Yum-Tong Siu	Harvard University, USA siu@math.harvard.edu
Vadim Kaloshin	University of Maryland, USA vadim.kaloshin@gmail.com	Terence Tao	University of California, Los Angeles, USA tao@math.ucla.edu
Herbert Koch	Universität Bonn, Germany koch@math.uni-bonn.de	Michael E. Taylor	Univ. of North Carolina, Chapel Hill, USA met@math.unc.edu
Izabella Laba	University of British Columbia, Canada ilaba@math.ubc.ca	Gunther Uhlmann	University of Washington, USA gunther@math.washington.edu
Richard B. Melrose	Massachusetts Inst. of Tech., USA rbm@math.mit.edu	András Vasy	Stanford University, USA andras@math.stanford.edu
Frank Merle	Université de Cergy-Pontoise, France Frank.Merle@u-cergy.fr	Dan Virgil Voiculescu	University of California, Berkeley, USA dvv@math.berkeley.edu
William Minicozzi II	Johns Hopkins University, USA minicozz@math.jhu.edu	Steven Zelditch	Northwestern University, USA zelditch@math.northwestern.edu
Clément Mouhot	Cambridge University, UK c.mouhot@dpms.cam.ac.uk	Maciej Zworski	University of California, Berkeley, USA zworski@math.berkeley.edu
Werner Müller	Universität Bonn, Germany mueller@math.uni-bonn.de		

PRODUCTION

production@msp.org

Silvio Levy, Scientific Editor

See inside back cover or msp.org/apde for submission instructions.

The subscription price for 2020 is US \$340/year for the electronic version, and \$550/year (+\$60, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues from the last three years and changes of subscriber address should be sent to MSP.

Analysis & PDE (ISSN 1948-206X electronic, 2157-5045 printed) at Mathematical Sciences Publishers, 798 Evans Hall #3840, c/o University of California, Berkeley, CA 94720-3840, is published continuously online. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices.

APDE peer review and production are managed by EditFlow® from MSP.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2020 Mathematical Sciences Publishers

ANALYSIS & PDE

Volume 13 No. 3 2020

On the gap between the Gamma-limit and the pointwise limit for a nonlocal approximation of the total variation	627
CLARA ANTONUCCI, MASSIMO GOBBINO and NICOLA PICENNI	
External boundary control of the motion of a rigid body immersed in a perfect two-dimensional fluid	651
OLIVIER GLASS, JÓZSEF J. KOLUMBÁN and FRANCK SUEUR	
Distance graphs and sets of positive upper density in \mathbb{R}^d	685
NEIL LYALL and ÁKOS MAGYAR	
Isolated singularities for semilinear elliptic systems with power-law nonlinearity	701
MARIUS GHERGU, SUNGHAN KIM and HENRIK SHAHGOLIAN	
Regularity of the free boundary for the vectorial Bernoulli problem	741
DARIO MAZZOLENI, SUSANNA TERRACINI and BOZHIDAR VELICHKOV	
On the discrete Fuglede and Pompeiu problems	765
GERGELY KISS, ROMANOS DIOGENES MALIKIOSIS, GÁBOR SOMLAI and MÁTÉ VIZER	
Energy conservation for the compressible Euler and Navier–Stokes equations with vacuum	789
IBROKHIMBEK AKRAMOV, TOMASZ DĘBIEC, JACK SKIPPER and EMIL WIEDEMANN	
A higher-dimensional Bourgain–Dyatlov fractal uncertainty principle	813
RUI HAN and WILHELM SCHLAG	
Local minimality results for the Mumford–Shah functional via monotonicity	865
DORIN BUCUR, ILARIA FRAGALÀ and ALESSANDRO GIACOMINI	
The gradient flow of the Möbius energy: ε -regularity and consequences	901
SIMON BLATT	
Correction to the article The heat kernel on an asymptotically conic manifold	943
DAVID A. SHER	



2157-5045(2020)13:3;1-B