

ANALYSIS & PDE

Volume 13 No. 4 2020

THOMAS METTLER AND GABRIEL P. PATERNAIN

**CONVEX PROJECTIVE SURFACES
WITH COMPATIBLE WEYL CONNECTION ARE HYPERBOLIC**

CONVEX PROJECTIVE SURFACES WITH COMPATIBLE WEYL CONNECTION ARE HYPERBOLIC

THOMAS METTLER AND GABRIEL P. PATERNAIN

We show that a properly convex projective structure p on a closed oriented surface of negative Euler characteristic arises from a Weyl connection if and only if p is hyperbolic. We phrase the problem as a nonlinear PDE for a Beltrami differential by using that p admits a compatible Weyl connection if and only if a certain holomorphic curve exists. Turning this nonlinear PDE into a transport equation, we obtain our result by applying methods from geometric inverse problems. In particular, we use an extension of a remarkable L^2 -energy identity known as Pestov's identity to prove a vanishing theorem for the relevant transport equation.

1. Introduction

A *projective structure* on a smooth manifold M is an equivalence class p of torsion-free connections on its tangent bundle TM , where two such connections are declared to be projectively equivalent if they share the same unparametrised geodesics. The set of torsion-free connections on TM is an affine space modelled on the sections of $S^2(T^*M) \otimes TM$. By a classical result of Cartan, Eisenhart, Weyl (see [Spivak 1999] for a modern reference), two connections are projectively equivalent if and only if their difference is pure trace. In particular, it follows from the representation theory of $GL(2, \mathbb{R})$ that a projective structure on a surface M is a section of a natural affine bundle of rank 4 whose associated vector bundle is canonically isomorphic to $V = S^3(T^*M) \otimes \Lambda^2(TM)$. Choosing an orientation and Riemannian metric g on M , the bundle V decomposes into irreducible $SO(2)$ -bundles $V \simeq T^*M \oplus S_0^3(T^*M)$, where the latter summand denotes the totally symmetric $(0,3)$ -tensors on M that are trace-free with respect to g , or equivalently, the cubic differentials with respect to the complex structure J induced by g and the orientation. In other words, fixing an orientation and Riemannian metric g on M , a projective structure p may be encoded in terms of a unique triple (g, A, θ) , where A is a cubic differential — and θ a 1-form on M . A conformal change of the metric $g \mapsto e^{2u}g$ corresponds to a change

$$(g, A, \theta) \mapsto (e^{2u}g, e^{2u}A, \theta + du).$$

Consequently, the section $\Phi = A/d\sigma$ of $K^2 \otimes \overline{K}^*$ does only depend on the complex structure J . Here $d\sigma$ denotes the area form of g and K the canonical bundle of M . In addition, we obtain a connection D on the anticanonical bundle K^* inducing the complex structure by taking the Chern connection with respect

MSC2010: primary 32W50, 53A20; secondary 30F30, 37D40.

Keywords: convex projective structures, Weyl connections, transport equations, energy identity.

to g and by subtracting twice the $(1,0)$ -part of θ . Again, the connection D does only depend on J . Fixing a complex structure J on M thus encodes a given projective structure $\mathfrak{p}$ in terms of a unique pair (D, Φ) .

There are two special cases of particular interest. Firstly, we can find a complex structure J so that D is the Chern connection of a metric in the conformal class determined by J . This amounts to finding a complex structure for which θ is exact. Secondly, we can find a complex structure J so that Φ vanishes identically. This turns out to be equivalent to $\mathfrak{p}$ containing a *Weyl connection* for the conformal structure $[g]$ determined by J , that is, a torsion-free connection on TM whose parallel transport maps are angle-preserving with respect to $[g]$.

In [Mettler 2014], it is shown that a two-dimensional projective structure $\mathfrak{p}$ does locally always contain a Weyl connection and moreover, finding the Weyl connection turns out to be equivalent to finding a holomorphic curve into a certain complex surface Z fibering over M . Here we use this observation to rephrase the problem in terms of a nonlinear PDE for a Beltrami differential. More precisely, we think of $\mathfrak{p}$ as being given on a Riemann surface (M, J) in terms of (D, Φ) . We show (see Proposition 4.4) that $\mathfrak{p}$ contains a Weyl connection with respect to the complex structure defined by the Beltrami differential μ on (M, J) if and only if

$$D''\mu - \mu D'\mu = \Phi\mu^3 + \bar{\Phi}, \quad (1-1)$$

where D' and D'' denote the $(1,0)$ - and $(0,1)$ -part of D . Since every two-dimensional projective structure locally contains a Weyl connection, the above PDE for the Beltrami differential μ can locally always be solved. Moreover, on the 2-sphere every solution μ lies in a complex 5-manifold of solutions, whereas on a closed surface of negative Euler characteristic the solution is unique, provided it exists; see [Mettler 2015b] (and Corollary 4.6 below).

Here we address the problem of finding a projective structure $\mathfrak{p}$ for which the above PDE has no global solution. Naturally, one might start by looking at projective structures $\mathfrak{p}$ at “the other end”, that is, those that arise from pairs (D, Φ) where D is the Chern connection of a conformal metric, or equivalently, those for which there exists a metric g so that $\mathfrak{p}$ is encoded in terms of the triple $(g, A, 0)$. This class of projective structures includes the so-called *properly convex projective structures*. A projective surface $(M, \mathfrak{p})$ is called properly convex if it arises as a quotient of a properly convex open set $\Omega \subset \mathbb{RP}^2$ by a free and cocompact action of a group $\Gamma \subset \mathrm{SL}(3, \mathbb{R})$ of projective transformations. In particular, using the Beltrami–Klein model of two-dimensional hyperbolic geometry, it follows that every closed hyperbolic Riemann surface is a properly convex projective surface. Motivated by Hitchin’s generalisation of Teichmüller space [1992], Labourie [2007] and Loftin [2001] have shown independently that on a closed oriented surface M of negative Euler characteristic every properly convex projective structure arises from a unique pair $(g, A, 0)$, where g and A are subject to the equations

$$K_g = -1 + 2|A|_g^2 \quad \text{and} \quad \bar{\partial}A = 0.$$

Using quasilinear elliptic PDE techniques, C. P. Wang [1991] previously showed (see also [Dumas and Wolf 2015]) that the metric g is uniquely determined in terms of $([g], A)$ by the equation for the Gauss curvature K_g of g . Consequently, Labourie and Loftin concluded that on M the properly convex projective

structures are in bijective correspondence with pairs $([g], A)$ consisting of a conformal structure and a cubic holomorphic differential.

Naturally one might speculate that (1-1) does not admit a global solution for a properly convex projective structure $\mathfrak{p}$ unless A vanishes identically, in which case $\mathfrak{p}$ is hyperbolic. This is indeed the case:

Corollary 6.2. *Let $(M, \mathfrak{p})$ be a closed oriented properly convex projective surface with $\chi(M) < 0$ and with $\mathfrak{p}$ containing a Weyl connection D . Then $\mathfrak{p}$ is hyperbolic and moreover D is the Levi-Civita connection of the hyperbolic metric.*

This corollary is an application of the more general vanishing theorem, Theorem 6.1 (see below), whose proof makes use of a remarkable L^2 -energy identity. This energy identity — known for geodesic flows as *Pestov's identity* — is ubiquitous when solving uniqueness problems for X-ray transforms, including tensor tomography. To make the bridge between (1-1) and this circle of ideas, it is necessary to recast the nonlinear PDE in dynamical terms as a transport problem. Given a projective structure $\mathfrak{p}$ captured by the triple (g, A, θ) we associate a dynamical system on the unit tangent bundle $\pi : SM \rightarrow M$ of g as follows. We consider a vector field of the form $F = X + (a - V\theta)V$, where X, V denote the geodesic and vertical vector fields of SM , $a \in C^\infty(SM, \mathbb{R})$ represents the cubic differential A (essentially its imaginary part) and where we think of θ as a function on SM . The flow of the vector field F is a *thermostat* (see Section 3 below for more details) and it has the property that its orbits project to M as unparametrised geodesics of $\mathfrak{p}$. We show that (1-1) is equivalent to the transport equation (see Corollary 5.6)

$$Fu = Va + \beta \tag{1-2}$$

on SM , where the real-valued function u encodes a conformal metric of the sought-after complex structure $\hat{J}$ and β is a 1-form on M , again thought of as a function on SM . Explicitly

$$u = \frac{3}{2} \log \left(\frac{p}{(pq - r^2)^{2/3}} \right),$$

where p, q, r are given in terms of a $\hat{J}$ -conformal metric $\hat{g}$ and the complex structure J of (M, g) by

$$p(x, v) = \hat{g}(v, v), \quad r(x, v) = \hat{g}(v, Jv) \quad \text{and} \quad q(x, v) = \hat{g}(Jv, Jv).$$

The right-hand side in (1-2) has degree 3 in the velocities and the dynamics of F is Anosov when $\mathfrak{p}$ is a properly convex projective structure [Mettler and Paternain 2019]; hence it is natural to think that techniques from tensor tomography might work. Regular tensor tomography involves the geodesic vector field X and the typical question at the level of the transport equation is the following: if $Xu = f$ where f has degree m in the velocities, is it true that u has degree $m - 1$ in the velocities? The case $m = 2$ is perhaps the most important and it is at the core of spectral rigidity of negatively curved manifolds and Anosov surfaces [Croke and Sharafutdinov 1998; Guillemin and Kazhdan 1980; Paternain et al. 2014]. Thermostats introduce new challenges; however we are able to successfully use a general L^2 energy identity developed in [Jane and Paternain 2009] (following earlier results for geodesic flows in [Sharafutdinov and Uhlmann 2000]) together with ideas in [Mettler and Paternain 2019] to show that if (1-2) holds then $a = 0$ and β is exact. Our vanishing theorem, Theorem 6.1, is actually rather general and

it applies to a class of projective structures considerably larger than properly convex projective structures; see Corollary 6.4 below.

For the case of surfaces with boundary, a full solution to the tensor tomography problem was given in [Paternain et al. 2013]; the solution was inspired by the proof of the Kodaira vanishing theorem in complex geometry. In the present paper, we go in the opposite direction; we import ideas from geometric inverse problems to solve an existence question for a nonlinear PDE in complex geometry. These connections were not anticipated, and it is natural to wonder if they are manifestations of something deeper.

2. Preliminaries

Here we collect some standard facts about Riemann surfaces and the unit tangent bundle that will be needed throughout the paper.

2A. The frame bundle. Throughout the article M will denote a connected oriented smooth surface with empty boundary. Unless stated otherwise, all maps are assumed to be smooth, i.e., C^∞ . Let $\pi : P \rightarrow M$ denote the *oriented frame bundle* of M whose fibre at a point $x \in M$ consists of the linear isomorphisms $f : \mathbb{R}^2 \rightarrow T_x M$ that are orientation-preserving, where we equip $\mathbb{R}^2$ with its standard orientation. The Lie group $\mathrm{GL}^+(2, \mathbb{R})$ acts transitively from the right on each fibre by the rule $R_h(f) = f \circ h$ and this action turns $\pi : P \rightarrow M$ into a principal right $\mathrm{GL}^+(2, \mathbb{R})$ -bundle. The bundle P is equipped with a tautological $\mathbb{R}^2$ -valued 1-form $\omega = (\omega^i)$ defined by $\omega_f = f^{-1} \circ d\pi_f$ and which satisfies the equivariance property $R_h^* \omega = h^{-1} \omega$. The components of ω are a basis for the 1-forms on P that are semibasic for the projection $\pi : P \rightarrow M$, i.e., those 1-forms that vanish when evaluated on a vector field that is tangent to the fibres of $\pi : P \rightarrow M$. Therefore, if g is a Riemannian metric on M , there exist unique real-valued functions $g_{ij} = g_{ji}$ on P so that $\pi^* g = g_{ij} \omega^i \otimes \omega^j$. The Levi-Civita connection ${}^g \nabla$ of g corresponds to the unique connection form $\psi = (\psi_j^i) \in \Omega^1(P, \mathfrak{gl}(2, \mathbb{R}))$ satisfying the structure equations

$$\begin{aligned} d\omega^i &= -\psi_j^i \wedge \omega^j, \\ dg_{ij} &= g_{ik} \psi_j^k + g_{kj} \psi_i^k. \end{aligned} \tag{2-1}$$

The curvature $\Psi = (\Psi_j^i)$ of ψ is the 2-form

$$\Psi_j^i = d\psi_j^i + \psi_k^i \wedge \psi_j^k = K_g g_{jk} \omega^i \wedge \omega^k,$$

where K_g denotes (the pullback to P of) the Gauss curvature of g .

2B. Conformal connections. The *conformal frame bundle* of the conformal equivalence class $[g]$ of g is the principal right $\mathrm{CO}(2)$ -subbundle $\pi : P_{[g]} \rightarrow M$ defined by

$$P_{[g]} = \{f \in P : g_{11}(f) = g_{22}(f) \text{ and } g_{12}(f) = 0\}.$$

Here $\mathrm{CO}(2) = \mathbb{R}^+ \times \mathrm{SO}(2)$ denotes the linear conformal group whose Lie algebra $\mathfrak{co}(2)$ is spanned by the matrices

$$\begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \quad \text{and} \quad \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}.$$

A *conformal connection* for $[g]$ is principal $\mathrm{CO}(2)$ connection

$$\kappa = \begin{pmatrix} \kappa_1 & -\kappa_2 \\ \kappa_2 & \kappa_1 \end{pmatrix}, \quad \kappa_i \in \Omega^1(P_{[g]}),$$

on $P_{[g]}$ which is *torsion-free*, that is, satisfies

$$d \begin{pmatrix} \omega^1 \\ \omega^2 \end{pmatrix} = - \begin{pmatrix} \kappa_1 & -\kappa_2 \\ \kappa_2 & \kappa_1 \end{pmatrix} \wedge \begin{pmatrix} \omega^1 \\ \omega^2 \end{pmatrix}. \quad (2-2)$$

The standard identification $\mathbb{R}^2 \simeq \mathbb{C}$ gives an identification $\mathrm{CO}(2) \simeq \mathrm{GL}(1, \mathbb{C})$ and consequently, $\mathfrak{co}(2) \simeq \mathbb{C}$. In particular, (2-2) takes the form $d\omega = -\kappa \wedge \omega$, where we think of κ and ω as being complex-valued. Writing $re^{i\phi}$ for the elements of $\mathrm{CO}(2)$, the equivariance property for ω implies $(R_{re^{i\phi}})^*\omega = \frac{1}{r}e^{-i\phi}\omega$. In particular, we see that the π -semibasic complex-valued 1-form ω is well-defined on M up to complex scale. It follows that there exists a unique complex-structure J on M whose $(1,0)$ -forms are represented by smooth complex-valued functions u on $P_{[g]}$ satisfying the equivariance property $(R_{re^{i\phi}})^*u = re^{i\phi}u$, that is, so that $u\omega$ is invariant under the $\mathrm{CO}(2)$ -right action. Of course, this is the standard complex structure on M obtained by rotation of a tangent vector v counterclockwise by $\frac{\pi}{2}$ with respect to $[g]$. Denoting the canonical bundle of M with respect to J by K , it follows that the sections of $L_{m,\ell} := K^m \otimes \bar{K}^\ell$ are in one-to-one correspondence with the smooth complex-valued functions u on $P_{[g]}$ satisfying the equivariance property $(R_{re^{i\phi}})^*u = r^{m+\ell}e^{i(m-\ell)\phi}u$. Infinitesimally, this translates to the existence of unique smooth complex-valued functions u' and u'' on $P_{[g]}$ so that

$$du = u'\omega + u''\bar{\omega} + mu\kappa + \ell u\bar{\kappa}. \quad (2-3)$$

Recall, if α is a 1-form on M taking values in some complex vector bundle over M , the decomposition $\alpha = \alpha' + \alpha''$ of α into its $(1,0)$ -part α' and $(0,1)$ -part α'' is given by

$$\alpha' = \frac{1}{2}(\alpha - iJ\alpha) \quad \text{and} \quad \alpha'' = \frac{1}{2}(\alpha + iJ\alpha),$$

where we define $(J\alpha)(v) := \alpha(Jv)$ for all tangent vectors $v \in TM$. The principal $\mathrm{CO}(2)$ -connection κ induces a connection on all (real or complex) vector bundles associated to $P_{[g]}$ and — by standard abuse of notation — we use the same letter D to denote the induced connection on the various bundles. If s is the section of $L_{m,\ell}$ represented by the function u satisfying (2-3), then $D's := (Ds)'$ is represented by u' and $D''s := (Ds)''$ is represented by u'' .

Since $dg_{11} = dg_{22}$ and $dg_{12} = 0$ on $P_{[g]}$, it follows from (2-1) that the pullback of the Levi-Civita connection ψ of g to $P_{[g]}$ is a conformal connection. The difference of any two principal $\mathrm{CO}(2)$ -connections is π -semibasic. Therefore, any other torsion-free principal $\mathrm{CO}(2)$ -connection κ on $P_{[g]}$ is of the form $\kappa = \psi - 2\theta_1\omega$ for a unique complex-valued function θ_1 on $P_{[g]}$. Since κ is a connection, it satisfies the equivariance property $(R_{re^{i\phi}})^*\kappa = \frac{1}{r}e^{-i\phi}\kappa re^{i\phi} = \kappa$ and so does ψ . Therefore, $2\theta_1\omega$ is invariant under the $\mathrm{CO}(2)$ -right action as well and hence twice the pullback of a $(1,0)$ -form on M , which we denote by θ' . From (2-3) we see that we may think of κ as being the connection form of the induced connection on the anticanonical bundle K^* . In particular, ψ may be thought of as being the connection form of the Chern connection induced by g on K^* . By the definition of the Chern connection, it induces

the complex structure of K^* . Since ψ and κ differ by a $(1,0)$ -form, κ also induces the complex structure of K^* . Consequently, the conformal connections on $P_{[g]}$ are in one-to-one correspondence with the connections D on K^* inducing the complex structure, that is, $D'' = \bar{\partial}_{K^*}$.

2C. The unit tangent bundle. For what follows it will be necessary to further reduce $P_{[g]}$. The unit tangent bundle

$$SM = \{(x, v) \in TM : g(v, v) = 1\}$$

of g may be interpreted as the principal right $SO(2)$ -subbundle of P defined by

$$SM = \{f \in P : g_{ij}(f) = \delta_{ij}\}.$$

On SM the identities $dg_{ij} \equiv 0$ imply the identities $\psi_1^1 \equiv \psi_2^2 \equiv 0$ and $\psi_2^1 \equiv -\psi_1^2$, so that ψ is purely imaginary.

Abusing notation by henceforth writing ψ instead of ψ_1^2 , the structure equations thus take the form

$$d \begin{pmatrix} \omega_1 \\ \omega_2 \end{pmatrix} = - \begin{pmatrix} 0 & -\psi \\ \psi & 0 \end{pmatrix} \wedge \begin{pmatrix} \omega_1 \\ \omega_2 \end{pmatrix} \quad \text{and} \quad d\psi = -K_g \omega_1 \wedge \omega_2, \quad (2-4)$$

where we write $\omega_i = \delta_{ij} \omega^j$. Note that on SM the 1-forms ω_1, ω_2 take the explicit form

$$\omega_1(\xi) = g(v, d\pi(\xi)) \quad \text{and} \quad \omega_2(\xi) = g(Jv, d\pi(\xi)), \quad \xi \in T_{(x,v)}SM. \quad (2-5)$$

Furthermore, the 1-form ψ becomes

$$\psi(\xi) = g(\gamma''(0), Jv), \quad (2-6)$$

where $\xi \in T_{(x,v)}SM$ and $\gamma : (-\varepsilon, \varepsilon) \rightarrow SM$ is any curve with $\gamma(0) = (x, v)$, $\dot{\gamma}(0) = \xi$ and γ'' denotes the covariant derivative of γ along $\pi \circ \gamma$.

The three 1-forms $(\omega_1, \omega_2, \psi)$ trivialise the cotangent bundle of SM and we denote by (X, H, V) the corresponding dual vector fields. The vector field X is the geodesic vector field of g , V is the infinitesimal generator of the $SO(2)$ -action and H is the horizontal vector field satisfying $H = [V, X]$. The structure equations (2-4) imply the additional commutation relations

$$[V, H] = -X \quad \text{and} \quad [X, H] = K_g V.$$

Following [Guillemin and Kazhdan 1980], we use the volume form $\Theta = \omega_1 \wedge \omega_2 \wedge \psi$ on SM to define an inner product

$$\langle u, v \rangle = \int_{SM} u \bar{v} \Theta$$

for complex-valued functions u, v on SM and we denote by $L^2(SM)$ the corresponding space of square integrable complex-valued functions on SM . The structure equations (2-4) and Cartan's formula imply that all vector fields X, H, V preserve Θ . In particular, $-iV$ is densely defined and self-adjoint with respect to $\langle \cdot, \cdot \rangle$. Consequently, we have an orthogonal direct sum decomposition into the kernels $\mathcal{H}_m$ of the operators $m\text{Id} + iV$

$$L^2(SM) = \bigoplus_{m \in \mathbb{Z}} \mathcal{H}_m. \quad (2-7)$$

2D. Weyl connections. If θ is a 1-form on M , we may write $\pi^*\theta = \theta\omega_1 + V(\theta)\omega_2$, where on the right-hand side we think of θ as being a real-valued function on SM . Therefore, $\pi^*\theta' = \theta_1\omega$, where $\theta_1 = \frac{1}{2}(\theta - iV\theta)$ and likewise $\pi^*\theta'' = \theta_{-1}\bar{\omega}$, where $\theta_{-1} = \frac{1}{2}(\theta + iV\theta)$. On SM the connection form κ of a conformal connection thus becomes $\kappa = i\psi - 2\theta_1\omega$, or in matrix notation

$$\kappa = \begin{pmatrix} 0 & -\psi \\ \psi & 0 \end{pmatrix} + \begin{pmatrix} -\theta\omega_1 - V(\theta)\omega_2 & -V(\theta)\omega_1 + \theta\omega_2 \\ V(\theta)\omega_1 - \theta\omega_2 & -\theta\omega_1 - V(\theta)\omega_2 \end{pmatrix}. \quad (2-8)$$

Finally, without the identification $\mathbb{R}^2 \simeq \mathbb{C}$, we may equivalently think of the connection form κ as the connection form of a torsion-free connection on TM . Writing κ as

$$\kappa = \begin{pmatrix} 0 & -\psi \\ \psi & 0 \end{pmatrix} + \begin{pmatrix} \theta\omega_1 & \theta\omega_2 \\ V(\theta)\omega_1 & V(\theta)\omega_2 \end{pmatrix} - \begin{pmatrix} 2\theta\omega_1 + V(\theta)\omega_2 & V(\theta)\omega_1 \\ \theta\omega_2 & \theta\omega_1 + 2V(\theta)\omega_2 \end{pmatrix},$$

the reader may easily check that κ is the connection form of

$$D = {}^g\nabla + g \otimes \theta^\sharp - \text{Sym}(\theta), \quad (2-9)$$

where the section $\text{Sym}(\theta)$ of $S^2(T^*M) \otimes TM$ is defined by the rule

$$\text{Sym}(\theta)(v_1, v_2) = \theta(v_1)v_2 + \theta(v_2)v_1$$

for all tangent vectors $v_1, v_2 \in T_x M$ and all $x \in M$. Connections of the form (2-9) for $g \in [g]$ and $\theta \in \Omega^1(M)$ are known as *Weyl connections* for the conformal structure $[g]$. By construction, they preserve $[g]$; that is, the parallel transport maps are angle-preserving with respect to $[g]$. Conversely, every torsion-free connection on TM preserving $[g]$ is of the form (2-9) for some $g \in [g]$ and 1-form θ . Summarising, we have the following folklore result:

Proposition 2.1. *On a Riemann surface M with conformal structure $[g]$ the following sets are in one-to-one correspondence:*

- (i) *the conformal connections on $P_{[g]}$,*
- (ii) *the connections on K^* inducing the complex structure,*
- (iii) *the Weyl connections for $[g]$.*

3. Projective thermostats

In this section we show how to associate the triple (g, A, θ) to a given projective structure p . As mentioned in the Introduction, the existence of such a triple is a consequence of some elementary facts about $\text{SO}(2)$ -representation theory and a description of projective structures as sections of a certain affine bundle over M (see [Mettler 2015a] for a construction of (g, A, θ) in that spirit); here instead we obtain the triple as a by-product of a characterisation of *projective thermostats*.

A (generalised) *thermostat* is a flow ϕ on SM which is generated by a vector field of the form $F = X + \lambda V$, where λ is a smooth real-valued function on SM . In this article we are mainly interested in the case where the generalised thermostat is *projective*. By this we mean that there exists a torsion-free

connection ∇ on TM having the property that for every ϕ -orbit $\gamma : I \rightarrow SM$ there exists a reparametrisation $\varphi : I' \rightarrow I$ so that $\pi \circ \gamma \circ \varphi : I' \rightarrow M$ is a geodesic of ∇ .

Phrased more loosely, the orbit projections to M agree with the geodesics of a projective structure $\mathfrak{p}$ on M . By a classical result of Cartan, Eisenhart, Weyl (see for instance [Spivak 1999, Chapter 6, Addendum 1, Proposition 17] for a modern reference), two torsion-free connections ∇ and ∇' on TM are projectively equivalent if and only if there exists a 1-form α on M so that

$$\nabla' - \nabla = \text{Sym}(\alpha).$$

3A. A characterisation of projective thermostats. It turns out that projective thermostats admit a simple characterisation in terms of the vertical Fourier decomposition (2-7) of λ . Towards this end we first show:

Lemma 3.1. *Let ∇ be a torsion-free connection on the tangent bundle TM and $\varphi = (\varphi_j^i) \in \Omega^1(SM, \mathfrak{gl}(2, \mathbb{R}))$ its connection form. Then, up to reparametrisation, the leaves of the foliation $\mathcal{F}$ defined by $\varphi_1^2 = \omega_2 = 0$ project to M to become the geodesics of ∇ . Conversely, every geodesic of ∇ , parametrised with respect to g -arc length, lifts to become a leaf of $\mathcal{F}$.*

Proof. Recall that the set of torsion-free connections on TM is an affine space modelled on the sections of $S^2(T^*M) \otimes TM$. It follows that there exists a 1-form $\tilde{B}$ on M with values in the endomorphisms of TM so that $\nabla = {}^g\nabla + \tilde{B}$. As we have seen, the connection form of the Levi-Civita connection of g on TM is

$$\kappa = \begin{pmatrix} 0 & -\psi \\ \psi & 0 \end{pmatrix}.$$

Hence there exist unique real-valued function $b_{jk}^i = b_{kj}^i$ on SM so that

$$\varphi = \begin{pmatrix} 0 & -\psi \\ \psi & 0 \end{pmatrix} + \begin{pmatrix} b_{11}^1\omega_1 + b_{12}^1\omega_2 & b_{21}^1\omega_1 + b_{22}^1\omega_2 \\ b_{11}^2\omega_1 + b_{12}^2\omega_2 & b_{21}^2\omega_1 + b_{22}^2\omega_2 \end{pmatrix}.$$

Explicitly, $b_{jk}^i(v) = g(\tilde{B}(e_j)e_k, e_i)$, where we write $e_1 = v$ and $e_2 = Jv$ for $v \in SM$.

Let $\delta : I \rightarrow SM$ be a leaf of $\mathcal{F}$, so that $\delta^*\omega_2 = 0$. Writing $\gamma := \pi \circ \delta$ and evaluating $\delta^*\omega_2$ on the standard vector field ∂_t of $\mathbb{R}$, we obtain

$$0 = \partial_t \lrcorner \delta^*\omega_2 = g(d(\pi \circ \delta)(\partial_t), J\delta(t)) = g(\dot{\gamma}(t), J\delta(t)),$$

so that $\delta = f\dot{\gamma}$ for some unique $f \in C^\infty(I)$. Hence without losing generality, we may assume that the leaves of $\mathcal{F}$ are of the form $\dot{\gamma}$ for some smooth curve $\gamma : I \rightarrow M$ having unit-length velocity vector with respect to g .

By the construction of ψ , see (2-6), the pullback 1-form $\dot{\gamma}^*\psi$ evaluated on ∂_t gives the function $g({}^g\nabla_{\dot{\gamma}}\dot{\gamma}, J\dot{\gamma})$; hence $\dot{\gamma}^*\varphi_1^2 = 0$ if and only if

$$0 = g({}^g\nabla_{\dot{\gamma}}\dot{\gamma}, J\dot{\gamma}) + b_{11}^2(\dot{\gamma}) = g({}^g\nabla_{\dot{\gamma}}\dot{\gamma} + \tilde{B}(\dot{\gamma})\dot{\gamma}, J\dot{\gamma}).$$

It follows that there exists a function $f \in C^\infty(I)$ so that

$${}^g\nabla_{\dot{\gamma}}\dot{\gamma} + \tilde{B}(\dot{\gamma})\dot{\gamma} = \nabla_{\dot{\gamma}}\dot{\gamma} = f\dot{\gamma}.$$

By a standard lemma in projective differential geometry [Spivak 1999, Chapter 6, Addendum 1, Proposition 17] a smooth immersed curve $\gamma : I \rightarrow M$ can be reparametrised to become a geodesic of the torsion-free connection ∇ on TM if and only if there exists a smooth function $f : I \rightarrow \mathbb{R}$ so that $\nabla_{\dot{\gamma}} \dot{\gamma} = f \dot{\gamma}$. The claim follows by applying this lemma. $\square$

Lemma 3.2. *Suppose the thermostat $F = X + \lambda V$ is projective. Then*

$$0 = \frac{3}{2}\lambda + \frac{5}{3}VV\lambda + \frac{1}{6}VVVV\lambda.$$

Proof. Using notation as in the proof of Lemma 3.1, we must have $F \lrcorner \varphi_1^2 = 0$ and $F \lrcorner \omega_2 = 0$. The latter condition is trivially satisfied, but the former gives

$$F \lrcorner \varphi_1^2 = (X + \lambda V) \lrcorner (\psi + b_{11}^2 \omega_1 + b_{12}^2 \omega_2) = \lambda + b_{11}^2 = 0,$$

so that $\lambda = -b_{11}^2$. Since the functions b_{jk}^i represent a section of $S^2(T^*M) \otimes TM$, they satisfy the structure equations

$$db_{jk}^i = b_{jl}^i \kappa_k^l + b_{lk}^i \kappa_j^l - b_{jk}^l \kappa_l^i, \quad \text{mod } \omega_i.$$

In particular, from this we compute

$$Vb_{11}^2 = V \lrcorner db_{11}^2 = V \lrcorner (2b_{12}^2 - b_{11}^1)\psi = 2b_{12}^2 - b_{11}^1.$$

Applying V again we obtain

$$VVb_{11}^2 = 2b_{22}^2 - 3b_{11}^2 - 4b_{12}^1,$$

and likewise

$$VVVVb_{11}^2 = 40b_{12}^1 + 21b_{11}^2 - 20b_{22}^2,$$

so that the claim follows from an elementary calculation. $\square$

Lemma 3.3. *For $\lambda \in C^\infty(SM)$ the following statements are equivalent:*

- (i) $0 = \frac{3}{2}\lambda + \frac{5}{3}VV\lambda + \frac{1}{6}VVVV\lambda$.
- (ii) $\lambda \in \mathcal{H}_{-1} \oplus \mathcal{H}_1 \oplus \mathcal{H}_{-3} \oplus \mathcal{H}_3$.

Proof. Let $\lambda \in \mathcal{H}_{-3} \oplus \mathcal{H}_{-1} \oplus \mathcal{H}_1 \oplus \mathcal{H}_3$ so that we may write $\lambda = \lambda_{-3} + \lambda_{-1} + \lambda_1 + \lambda_3$ with $\lambda_m \in \mathcal{H}_m$. Since λ is real-valued we have $\lambda_{-1} = \bar{\lambda}_1$ and $\lambda_{-3} = \bar{\lambda}_3$. Hence setting $v_1 = \lambda_{-1} + \lambda_1$ and $v_3 = \lambda_{-3} + \lambda_3$, we obtain $VVv_1 = -v_1$ and $VVv_3 = -9v_3$ so that

$$\frac{3}{2}\lambda + \frac{5}{3}VV\lambda + \frac{1}{6}VVVV\lambda = \frac{3}{2}(v_3 + v_1) + \frac{5}{3}(-9v_3 - v_1) + \frac{1}{6}(81v_3 + v_1) = 0.$$

Conversely, suppose $\lambda \in C^\infty(SM)$ satisfies $0 = \frac{3}{2}\lambda + \frac{5}{3}VV\lambda + \frac{1}{6}VVVV\lambda$ and write $\lambda = \sum_m \lambda_m$, with $\lambda_m \in \mathcal{H}_m$. Hence we obtain

$$0 = \frac{3}{2}\lambda + \frac{5}{3}VV\lambda + \frac{1}{6}VVVV\lambda = \sum_m \left(\frac{3}{2} - \frac{5}{3}m^2 + \frac{1}{6}m^4 \right) \lambda_m,$$

so that $\lambda_m = 0$ unless

$$0 = \frac{3}{2} - \frac{5}{3}m^2 + \frac{1}{6}m^4 = \frac{1}{6}(m-3)(m-1)(m+1)(m+3).$$

The claim follows. $\square$

Finally, we obtain:

Proposition 3.4. *A thermostat $F = X + \lambda V$ is projective if and only if $\lambda \in \mathcal{H}_{-1} \oplus \mathcal{H}_1 \oplus \mathcal{H}_{-3} \oplus \mathcal{H}_3$.*

Proof. It remains to show that if $\lambda \in \mathcal{H}_{-1} \oplus \mathcal{H}_1 \oplus \mathcal{H}_{-3} \oplus \mathcal{H}_3$, then there exists a torsion-free connection ∇ on TM so that $F \lrcorner \varphi_1^2$ vanishes identically, where $\varphi = (\varphi_j^i)$ denotes the connection form of ∇ . We may write

$$\lambda = a - V\theta,$$

where $a \in C^\infty(SM)$ satisfies $9a + VVa = 0$ and θ is a smooth 1-form on M , thought of as a real-valued function on SM . Since $9a + VVa = 0$, there exists a unique cubic differential A on M so that

$$\pi^* A = \left(\frac{1}{3}Va + ia\right)\omega^3.$$

Hence simple computations show that

$$\begin{aligned} a(v) &= \operatorname{Re} A(Jv, Jv, Jv) = -\operatorname{Re} A(Jv, v, v), \\ \frac{1}{3}Va(v) &= \operatorname{Re} A(v, v, v) = -\operatorname{Re} A(v, Jv, Jv) \end{aligned} \quad (3-1)$$

for all $v \in SM$. Let B be the unique 1-form on M with values in the endomorphisms of TM satisfying

$$g(B(v_1)v_2, v_3) = \operatorname{Re} A(v_1, v_2, v_3) \quad (3-2)$$

for all tangent vectors $v_1, v_2, v_3 \in T_x M$ and all $x \in M$. On TM consider the torsion-free connection $\nabla = D + B$, where D is the Weyl connection

$$D = {}^g\nabla + g \otimes \theta^\sharp - \operatorname{Sym}(\theta).$$

Using (2-8) and (3-1), we compute that the connection form of ∇ is

$$\varphi = \begin{pmatrix} -\theta\omega_1 - V(\theta)\omega_2 & -V(\theta)\omega_1 + \theta\omega_2 - \psi \\ \psi + V(\theta)\omega_1 - \theta\omega_2 & -\theta\omega_1 - V(\theta)\omega_2 \end{pmatrix} + \begin{pmatrix} \frac{1}{3}V(a)\omega_1 - a\omega_2 & -a\omega_1 - \frac{1}{3}V(a)\omega_2 \\ -a\omega_1 - \frac{1}{3}V(a)\omega_2 & -\frac{1}{3}V(a)\omega_1 + a\omega_2 \end{pmatrix}. \quad (3-3)$$

In particular, we have

$$\varphi_1^2 = \psi + (V(\theta) - a)\omega_1 - \left(\theta + \frac{1}{3}V(a)\right)\omega_2,$$

so that $F \lrcorner \varphi_1^2 = 0$. □

3B. The effect of a conformal change. Summarising the previous subsection, we have seen that if ∇ is a torsion-free connection on TM and we fix a Riemannian metric g on M , then we may write $\nabla = {}^g\nabla + \tilde{B}$ for some endomorphism-valued 1-form $\tilde{B}$ on M . The thermostat on SM defined by $\lambda = -b_{11}^2$ has the property that its orbits project to M to become the geodesics of ∇ up to parametrisation. Moreover, we obtain a 1-form $\theta \in \Omega^1(M)$ as well as a cubic differential $A \in \Gamma(K^3)$, so that the connection ∇ shares its geodesics — up to parametrisation — with the projections to M of the orbits of the projective thermostat defined by $\lambda = a - V\theta$, where a represents the imaginary part of A .

Next we compute how θ and A transform under conformal change of the metric. As a consequence, we obtain:

Proposition 3.5. *Let ∇ be a torsion-free connection on TM . Then the choice of a conformal structure $[g]$ on M determines a unique Weyl connection D for $[g]$ and a unique section Φ of $K^2 \otimes \bar{K}^*$ so that $D + \text{Re } \Phi$ is projectively equivalent to ∇ .*

Proof. Let $g \mapsto \hat{g} = e^{2u}g$ be a conformal change of the metric, where $u \in C^\infty(M)$. For the new metric $\hat{g}$ there exists a 1-form $\hat{\theta}$ and a cubic differential $\hat{A}$ on M so that $D + B$ and $\hat{D} + \hat{B}$ are projectively equivalent. Here $\hat{B}$ denotes the 1-form constructed from $\hat{A}$ by using the metric $\hat{g}$. Projective equivalence corresponds to the existence of a 1-form α on M so that

$$D + B = \hat{D} + \hat{B} + \text{Sym}(\alpha).$$

Using (2-9) as well as (see [Besse 1987, Theorem 1.159])

$$\exp(2u)g \nabla = {}^g\nabla - g \otimes {}^g\nabla u + \text{Sym}(du), \quad (3-4)$$

this is equivalent to

$${}^g\nabla + g \otimes \theta^\sharp - \text{Sym}(\theta) + B = {}^g\nabla - g \otimes {}^g\nabla u + \text{Sym}(du) + e^{2u}g \otimes \hat{\theta}^\sharp - \text{Sym}(\hat{\theta}) + \hat{B} + \text{Sym}(\alpha)$$

or

$$g \otimes (\theta^\sharp + {}^g\nabla u - \hat{\theta}^\sharp) + B - \hat{B} = \text{Sym}(\beta),$$

where $\beta = \alpha + \theta + du - \hat{\theta}$. Evaluating this equation on the pair (v, Jv) with v a unit tangent vector with respect to g gives

$$B(v)Jv - \hat{B}(v)Jv = \text{Sym}(\beta)(v, Jv).$$

Computing the inner product with the tangent vector v yields

$$\text{Re } A(v, Jv, v) - e^{-2u} \text{Re } \hat{A}(v, Jv, v) = \beta(Jv).$$

Thought of as an identity for functions on SM , the left-hand side lies in $\mathcal{H}_{-3} \oplus \mathcal{H}_3$, whereas the right-hand side lies in $\mathcal{H}_{-1} \oplus \mathcal{H}_1$, and hence they can only be equal if both sides vanish identically. Consequently, it follows that $\beta = 0$ and that

$$\hat{A} = e^{2u}A.$$

Therefore, $B = \hat{B}$ and

$$\hat{\theta} = \theta + du, \quad (3-5)$$

so that $\alpha = 0$ as well as $D = \hat{D}$.

In particular, we see that both D and B do only depend on the conformal equivalence class of g . We may define a section Φ of $K^2 \otimes \bar{K}^*$ by $\Phi d\sigma = A$, where $d\sigma$ denotes the area form of g . Comparing with (3-2), we see that B is the real part of Φ . $\square$

4. Holomorphic curves

It is natural to ask whether for a given torsion-free connection ∇ on TM one can always (at least locally) choose a conformal structure $[g]$ on M so that Φ vanishes identically, or equivalently, whether every torsion-free connection ∇ on TM is locally projectively equivalent to a Weyl connection D . This question

was answered in the affirmative in [Mettler 2014], where it is also observed that the problem is equivalent to finding a suitable holomorphic curve into a complex surface fibering over M . Here we will briefly review this observation and use it to derive a nonlinear PDE for the Beltrami differential of the sought-after conformal structure.

Remark 4.1. Given that one can locally always find a conformal structure so that Φ vanishes identically, one might wonder whether it is possible to simultaneously pick a conformal metric so that the 1-form θ is closed. Indeed, (3-4) and (3-5) imply that the additional closedness condition corresponds to ∇ being locally projectively equivalent to a Levi-Civita connection of some metric. However, this is not always possible; see [Bryant et al. 2009].

4A. A complex surface. Inspired by the twistorial construction of holomorphic projective structures in [Hitchin 1982], it was shown in [Dubois-Violette 1983; O’Brian and Rawnsley 1985] how to construct a “twistor space” for smooth projective structures. Let ∇ be a torsion-free connection on TM and $\varphi = (\varphi_j^i) \in \Omega^1(P, \mathfrak{gl}(2, \mathbb{R}))$ its connection form on the frame bundle P . We can use φ to construct a complex structure on the quotient $P/\mathrm{CO}(2)$. By definition, an element of $P/\mathrm{CO}(2)$ gives a frame in some tangent space of M , well defined up to rotation and scaling. Therefore, the conformal structures on M are in one-to-one correspondence with the sections of the fibre bundle $P/\mathrm{CO}(2) \rightarrow M$ whose fibre is $\mathrm{GL}^+(2, \mathbb{R})/\mathrm{CO}(2)$, that is, the open disk. We will construct a complex structure on $P/\mathrm{CO}(2)$ in terms of its $(1,0)$ -forms, or more precisely, the pullbacks of the $(1,0)$ -forms to P . Recall that the Lie algebra $\mathfrak{co}(2)$ of $\mathrm{CO}(2)$ is spanned by the matrices

$$\begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \quad \text{and} \quad \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}.$$

Consequently, the complex-valued 1-forms on P that are semibasic for the quotient projection $P \rightarrow P/\mathrm{CO}(2)$ are spanned by the form ω and

$$\zeta = (\varphi_1^1 - \varphi_2^2) + i(\varphi_2^1 + \varphi_1^2),$$

as well as their complex conjugates. Recall that we have $(R_{re^{i\phi}})^*\omega = \frac{1}{r}e^{-i\phi}\omega$ and using that φ satisfies the equivariance property $R_h^*\varphi = h^{-1}\varphi h$ for all $h \in \mathrm{GL}^+(2, \mathbb{R})$, we compute $(R_{re^{i\phi}})^*\zeta = e^{-2i\phi}\zeta$. It follows that there exists a unique almost-complex structure J on $P/\mathrm{CO}(2)$ whose $(1,0)$ -forms pull back to P to become linear combinations of the forms ω, ζ . The almost-complex structure J can be shown to only depend on the projective equivalence class of ∇ , and moreover, an application of the Newlander–Nirenberg theorem shows that J is always integrable; see [Mettler 2014] for details.

4B. Möbius action. In our setting it is convenient to reduce the frame bundle P to the unit tangent bundle SM of some fixed metric g . In order to get a handle on the complex surface $P/\mathrm{CO}(2)$ after having carried out this reduction, we interpret the disk bundle $P/\mathrm{CO}(2) \rightarrow M$ as an associated bundle to the frame bundle P . This requires an action of the structure group $\mathrm{GL}^+(2, \mathbb{R})$ on the open disk and this is what we compute next.

The group $\mathrm{GL}^+(2, \mathbb{R})$ acts from the left on the lower half plane

$$-\mathbb{H} := \{w \in \mathbb{C} : \Im(w) < 0\}$$

by Möbius transformations, where w denotes the standard coordinate on $\mathbb{C}$. We let $\mathbb{D} \subset \mathbb{C}$ denote the open unit disk. Identifying $-\mathbb{H}$ with $\mathbb{D}$ via the Möbius transformation

$$-\mathbb{H} \rightarrow \mathbb{D}, \quad w \mapsto -\left(\frac{w+i}{w-i}\right),$$

we obtain an induced action of $\mathrm{GL}^+(2, \mathbb{R})$ on $\mathbb{D}$ making this transformation equivariant:

$$\begin{pmatrix} a & b \\ c & d \end{pmatrix} \cdot z = \frac{iz(a+d) + z(b-c) - i(a-d) + (b+c)}{-iz(a-d) - z(b+c) + i(a+d) - (b-c)}. \quad (4-1)$$

The stabiliser subgroup of the point $z=0$ consists of elements in $\mathrm{GL}^+(2, \mathbb{R})$ satisfying $a=d$ and $b+c=0$, i.e., the linear conformal group $\mathrm{CO}(2)$. Consequently, we have

$$\mathbb{D} \simeq \mathrm{GL}^+(2, \mathbb{R}) / \mathrm{CO}(2)$$

and we obtain a projection

$$\lambda : \mathrm{GL}^+(2, \mathbb{R}) \rightarrow \mathbb{D}, \quad \begin{pmatrix} a & b \\ c & d \end{pmatrix} \mapsto \begin{pmatrix} a & b \\ c & d \end{pmatrix} \cdot 0 = \frac{-i(a-d) + (b+c)}{i(a+d) - (b-c)}.$$

In particular, a mapping $z : N \rightarrow \mathbb{D}$ from a smooth manifold N into $\mathbb{D}$ is covered by a map

$$\tilde{z} = \begin{pmatrix} \frac{1-|z|^2}{(1+z)(1+\bar{z})} & \frac{i(z-\bar{z})}{(1+z)(1+\bar{z})} \\ 0 & 1 \end{pmatrix}$$

into $\mathrm{GL}^+(2, \mathbb{R})$. Equivalently, we have $\tilde{z} \cdot 0 = z$ or $z \cdot \tilde{z} = 0$, where as usual we turn the left action into a right action by the definition $z \cdot \tilde{z} := \tilde{z}^{-1} \cdot z$.

Let $\rho : Z \rightarrow M$ denote the disk-bundle associated to the above $\mathrm{GL}^+(2, \mathbb{R})$ action on $\mathbb{D}$. Suppose $z : P \rightarrow \mathbb{D}$ represents a section of $Z \rightarrow M$ so that z is a $\mathrm{GL}^+(2, \mathbb{R})$ -equivariant map. For every coframe $u \in P$ the pair $(u, z(u)) \in P \times \mathbb{D}$ lies in the same $\mathrm{GL}^+(2, \mathbb{R})$ orbit as

$$(u \cdot \tilde{z}(u), z(u) \cdot \tilde{z}(u)) = (u \cdot \tilde{z}(u), 0). \quad (4-2)$$

Therefore, the map z gives for every point $p \in M$ a coframe $u \cdot \tilde{z}(u)$ which is unique up to the action of $\mathrm{CO}(2)$. It follows that the bundle $Z \rightarrow M$ is isomorphic to $P / \mathrm{CO}(2) \rightarrow M$, as desired.

Let $\Upsilon : P \times \mathbb{D} \rightarrow P$ be the map defined by (4-2). We will next compute the pullback of ω, ζ under Υ . Note that we may write $\Upsilon = R \circ (\mathrm{Id}_P \times \tilde{z})$, where $R : P \times \mathrm{GL}^+(2, \mathbb{R}) \rightarrow P$ denotes the $\mathrm{GL}^+(2, \mathbb{R})$ right action of P . Recall the standard identities

$$R^* \varphi = h^{-1} \varphi h + h^{-1} dh \quad \text{and} \quad R^* \omega = h^{-1} \omega,$$

where $h : P \times \mathrm{GL}^+(2, \mathbb{R}) \rightarrow \mathrm{GL}^+(2, \mathbb{R})$ denotes the projection onto the latter factor. From this we compute

$$\omega_\Upsilon := \Upsilon^* \omega = \tilde{z}^{-1} \omega = \left(\frac{1 + \bar{z}}{1 - |z|^2} \right) (\omega + z \bar{\omega}), \quad (4-3)$$

$$\varphi_\Upsilon := \Upsilon^* \varphi = \tilde{z}^{-1} \varphi \tilde{z} + \tilde{z}^{-1} d\tilde{z}. \quad (4-4)$$

We also obtain $\zeta_\Upsilon = \Upsilon^* \zeta = (\varphi_\Upsilon)_1^1 - (\varphi_\Upsilon)_2^2 + i((\varphi_\Upsilon)_2^1 + (\varphi_\Upsilon)_1^2)$. Writing

$$\chi = \frac{1}{2}(3(\varphi_1^1 + \varphi_2^2) + i(\varphi_1^2 - \varphi_2^1)),$$

and using (4-4), a tedious but straightforward calculation gives

$$\zeta_\Upsilon = \frac{2(1 + \bar{z})}{(|z|^2 - 1)(z + 1)} \left(dz - \frac{1}{2}\zeta + \frac{1}{2}z^2\bar{\zeta} + z\chi - z\bar{\chi} \right). \quad (4-5)$$

Remark 4.2. The complex-valued 1-form χ is chosen so that $\chi, \bar{\chi}, \omega, \bar{\omega}, \zeta, \bar{\zeta}$ span the complex-valued 1-forms on P . Clearly, this condition does not pin down χ uniquely. The particular choice is so that in the absence of θ the form χ becomes the connection form of the Chern connection on K^* upon reducing to SM ; see (4-6) below.

The complex structure on Z only depends on the projective equivalence class of ∇ . Thus, after possibly replacing φ with a projectively equivalent connection, we can assume that the torsion-free connection on TM corresponding to φ is of the form $D + B$ for some 1-form θ and some cubic differential A on M . On the unit tangent bundle SM of g the connection form of $D + B$ takes the form (3-3). Using this equation and reducing to $SM \subset P$ yields the following identities on SM :

$$\begin{aligned} \zeta &= 2a_{-3}\bar{\omega}, \\ \chi &= i\psi - 4\theta_1\omega - 2\theta_{-1}\bar{\omega}. \end{aligned} \quad (4-6)$$

Recall, we write $a_3 = \frac{1}{3}Va + ia$ and $a_{-3} = \bar{a}_3$ as well as $\theta_1 = \frac{1}{2}(\theta - iV\theta)$ and $\theta_{-1} = \bar{\theta}_1$. Also, the connection form κ of the induced Weyl connection is $\kappa = i\psi - 2\theta_1\omega$; see (2-8). Therefore, we have

$$\chi = 2\kappa + \bar{\kappa}.$$

The $\mathrm{SO}(2)$ -action induced by (4-1) is

$$\begin{pmatrix} \cos \phi & -\sin \phi \\ \sin \phi & \cos \phi \end{pmatrix} \cdot z = \frac{2iz \cos \phi - 2z \sin \phi}{2i \cos \phi + 2 \sin \phi} = e^{2i\phi} z,$$

and hence the equivariance property of a function $z : SM \rightarrow \mathbb{D}$ representing a section of $Z \rightarrow M$ becomes $(R_{e^{i\phi}})^* z = e^{-2i\phi} z$; that is, z represents a section of K^{-2} . Since we have a metric, we have an identification $K^* \simeq \bar{K}$ and hence $K^{-2} \simeq K^* \otimes \bar{K}$. In particular, we may write

$$dz = z'\omega + z''\bar{\omega} + \bar{\kappa}z - \kappa z \quad (4-7)$$

for unique complex-valued functions z' and z'' on SM . Consequently, using (4-5), (4-6) and (4-7) we obtain

$$\begin{aligned} \left(\frac{(|z|^2 - 1)(z + 1)}{2(1 + \bar{z})} \right) \zeta_\gamma &= z' \omega + z'' \bar{\omega} + \bar{\kappa} z - \kappa z - a_{-3} \bar{\omega} + z^2 a_3 \omega + z(2\kappa + \bar{\kappa}) - z(2\bar{\kappa} + \kappa) \\ &= (z' + z^2 a_3) \omega + (z'' - a_{-3}) \bar{\omega}. \end{aligned} \quad (4-8)$$

In order to connect the expressions for ω_γ and ζ_γ to the condition of z representing a conformal structure $[\hat{g}]$ that defines a holomorphic curve into Z , we use the following elementary lemma:

Lemma 4.3. *Let Z be a complex surface and $\omega, \zeta \in \Omega^1(Z, \mathbb{C})$ a basis for the $(1,0)$ -forms of Z . Suppose $M \subset Z$ is a smooth surface on which $\omega \wedge \bar{\omega}$ is nonvanishing. Then M admits the structure of a holomorphic curve — that is, a complex one-dimensional submanifold of Z — if and only if $\omega \wedge \zeta$ vanishes identically on M .*

Proof. Since $\omega \wedge \bar{\omega}$ is nonvanishing on M , the forms ω and $\bar{\omega}$ span the complex-valued 1-forms on M . Since M is a complex submanifold of Z if and only if the pullback of a $(1,0)$ -form on Z is a $(1,0)$ -form on M , the claim follows. $\square$

The reduction of P to SM identifies Z with $SM \times_{SO(2)} \mathbb{D}$. Now suppose the conformal structure $[\hat{g}] : M \rightarrow Z$ is represented by the map $z : SM \rightarrow \mathbb{D}$. If $v : U \rightarrow SM$ is a local section of $\pi : SM \rightarrow M$, then $[\hat{g}]|_U : U \rightarrow Z$ is covered by the map $(\text{Id}_{SM} \times z) \circ v : U \rightarrow SM \times \mathbb{D}$. Recall that the complex structure on Z has the property that its $(1,0)$ -forms pull-back to become linear combinations of ω_γ and ζ_γ . Using the expressions (4-3) and (4-8) for the pullbacks of ω_γ and ζ_γ to SM we obtain

$$\omega_\gamma \wedge \zeta_\gamma = -\frac{2(1 + \bar{z})^2}{(|z|^2 - 1)^2(z + 1)} (z'' - zz' - z^3 a_3 - a_{-3}) \omega \wedge \bar{\omega}.$$

In particular, since $v : U \rightarrow SM$ is a π -section and ω and $\bar{\omega}$ are π -semibasic, the pullback $v^*(\omega_\gamma \wedge \zeta_\gamma)$ vanishes if and only if $\omega_\gamma \wedge \zeta_\gamma$ vanishes on $\pi^{-1}(U)$. Thus, Lemma 4.3 implies that z represents a holomorphic curve if and only if

$$z'' - zz' = z^3 a_3 + a_{-3}. \quad (4-9)$$

4C. The Beltrami differential. So far we have not explicitly tied the conformal structure $[\hat{g}]$ to the function $z : SM \rightarrow \mathbb{D}$ representing it. In order to do this we first recall the Beltrami differential. The choice of a metric $\hat{g}$ on M allows us to define the functions

$$p(x, v) = \hat{g}(v, v), \quad r(x, v) = \hat{g}(v, Jv) \quad \text{and} \quad q(x, v) = \hat{g}(Jv, Jv)$$

on SM . The orientation-compatible complex structure $\hat{J}$ on M induced by the conformal equivalence class of $\hat{g}$ has matrix representation

$$\hat{J} = \frac{1}{\sqrt{pq - r^2}} \begin{pmatrix} -r & -q \\ p & r \end{pmatrix}.$$

In particular, we compute that the $(1,0)$ -forms with respect to $\hat{J}$ pull-back to SM to become complex multiples of

$$\omega_{\hat{J}} := \frac{1}{2}(\omega - i\hat{J}\omega) = \left(\frac{p+q+2\sqrt{pq-r^2}}{4\sqrt{pq-r^2}} \right) (\omega + \mu\bar{\omega}), \quad (4-10)$$

where

$$\mu = \frac{(p-q) + 2ir}{p+q+2\sqrt{pq-r^2}}$$

is the *Beltrami coefficient* of $\hat{J}$. Clearly, μ does only depend on the conformal equivalence class $[\hat{g}]$ of $\hat{g}$. Moreover, the function μ represents a $(0,1)$ -form on M with values in K^* called the *Beltrami differential* of $[\hat{g}]$, which — by abuse of language — we denote by μ as well.

The reduction of P to the unit tangent bundle SM of g turns ω into a basis for the $(1,0)$ -forms with respect to the complex structure induced by g and the orientation. The mapping z represents a conformal structure $[\hat{g}]$ and consequently, induces an orientation-compatible complex structure $\hat{J}$ whose $(1,0)$ -forms we computed in (4-3). Comparing this expression with the formula (4-10) for the Beltrami coefficient shows that we obtain the same $(1,0)$ -forms if and only if $z = \mu$. Remember, z' and z'' represent the $(1,0)$ - and $(0,1)$ -part of the derivative of z with respect to the connection D induced by the Weyl connection D . Furthermore, the function a_3 represents the cubic differential A or equivalently, the form Φ , since $\Phi d\sigma = A$ and $d\sigma$ is represented by the constant function 1 on SM . Using (4-9) and the fact that $\mathfrak{p}$ contains a Weyl connection with respect to $[\hat{g}]$ if and only if $[\hat{g}] : M \rightarrow Z$ is a holomorphic curve [Mettler 2014, Theorem 3], we have thus shown:

Proposition 4.4. *Let $(M, [g])$ be a Riemann surface equipped with a projective structure $\mathfrak{p}$ given in terms of (D, Φ) . Then $\mathfrak{p}$ contains a Weyl connection with respect to the conformal structure defined by the Beltrami differential μ if and only if*

$$D''\mu - \mu D'\mu = \Phi\mu^3 + \bar{\Phi}. \quad (4-11)$$

Remark 4.5. In the special case where $\mathfrak{p}$ is a properly convex projective structure, an equation equivalent to (4-11) was previously obtained by N. Hitchin using the Higgs bundle description of $\mathfrak{p}$.¹

As a corollary, we obtain:

Corollary 4.6. *Let M be a closed oriented surface with $\chi(M) < 0$. Suppose the Weyl connections D and $\hat{D}$ on TM are projectively equivalent. Then $D = \hat{D}$ and they preserve the same conformal structure.*

Proof. Equip M with the Riemann surface structure defined by $[g]$ and the orientation. Let $\mathfrak{p}$ be the projective structure defined by D (or $\hat{D}$). The projective structure $\mathfrak{p}$ is encoded in terms of the pair $(D, 0)$. Moreover, the Beltrami differential μ defined by $[\hat{g}]$ solves (4-11); that is,

$$D''\mu - \mu D'\mu = 0.$$

¹Private communication, August 2014.

Now observe that $\bar{\partial}_\mu = D'' - D'\mu$ defines a del-bar operator on $\bar{K} \otimes K^*$ and hence (4-11) can be written as $\bar{\partial}_\mu \mu = 0$. Therefore, μ is holomorphic with respect to the holomorphic line bundle structure defined by $\bar{\partial}_\mu$ on $\bar{K} \otimes K^*$. However, since $\chi(M) < 0$, the line bundle $\bar{K} \otimes K^*$ has negative degree, so that its only holomorphic section is the zero-section. It follows that $\mu = 0$ and hence $[g] = [\hat{g}]$. Since D and $\hat{D}$ are projectively equivalent and preserve the same conformal structure $[g]$, we conclude exactly as in the proof of Proposition 3.5 that $D = \hat{D}$. $\square$

Remark 4.7. The above corollary was first proved in [Mettler 2015b]. In particular, as a special case, it also shows that on a closed surface with $\chi(M) < 0$, the unparametrised geodesics of a Riemannian metric determine the metric up to rescaling by a positive constant. This was first observed in [Matveev and Topalov 2000].

5. The transport equation

While the PDE (4-11) for the Beltrami differential μ is natural from a complex geometry point of view, it turns out to be advantageous to rephrase it as a transport equation on SM . The relevant transport equation on SM can be derived using (4-11)—see the Appendix—but here we will instead take a different approach, as it leads to a more general result about thermostats having the same unparametrised geodesics; see Proposition 5.2.

Let $g, \hat{g}$ be Riemannian metrics on M . In what follows all objects defined in terms of the metric $\hat{g}$ will be decorated with a hat symbol. There is an obvious scaling map

$$\ell : SM \rightarrow \widehat{SM}, \quad (x, v) \mapsto \left(x, \frac{v}{\sqrt{\hat{g}(v, v)}} \right),$$

which is a fibre-bundle isomorphism covering the identity on M . As before we define

$$p(x, v) = \hat{g}(v, v), \quad r(x, v) = \hat{g}(v, Jv) \quad \text{and} \quad q(x, v) = \hat{g}(Jv, Jv).$$

Lemma 5.1. *The pullback of the volume form $\widehat{\Theta}$ on $\widehat{SM}$ is*

$$\ell^* \widehat{\Theta} = \left(\frac{pq - r^2}{p} \right) \Theta.$$

Proof. Since

$$d\pi(X(x, v)) = v \quad \text{and} \quad d\pi(H(x, v)) = Jv,$$

we obtain

$$\pi^* \hat{g} = p\omega_1 \otimes \omega_1 + 2r\omega_1 \otimes \omega_2 + q\omega_2 \otimes \omega_2,$$

where we write $\omega_1 \otimes \omega_2 := \frac{1}{2}(\omega_1 \otimes \omega_2 + \omega_2 \otimes \omega_1)$. We first compute

$$\begin{aligned} (X \lrcorner \ell^* \widehat{\omega}_1)(x, v) &= \widehat{\omega}_1(d\ell(X(x, v))) = \hat{g}(\ell(x, v), (d\hat{\pi} \circ d\ell)(X(x, v))) \\ &= \frac{1}{\sqrt{\hat{g}(v, v)}} \hat{g}(v, d\pi(X(x, v))) = \sqrt{\hat{g}(v, v)} = \sqrt{p}, \end{aligned}$$

where we have used that $\hat{\pi} \circ \ell = \pi$. Likewise, we obtain

$$\begin{aligned} (H \lrcorner \ell^*_1)(x, v) &= \hat{\omega}_1(d\ell(H(x, v))) = \hat{g}(\ell(x, v), (d\hat{\pi} \circ d\ell)(H(x, v))) \\ &= \frac{1}{\sqrt{\hat{g}(v, v)}} \hat{g}(v, d\pi(H(x, v))) = \frac{\hat{g}(v, Jv)}{\sqrt{\hat{g}(v, v)}} = \frac{r}{\sqrt{p}}. \end{aligned}$$

Since $\hat{\omega}_1$ is semibasic for the projection $\hat{\pi}$, the pullback $\ell^*\hat{\omega}_1$ is semibasic for the projection π ; hence $V \lrcorner \ell^*\hat{\omega}_1 = 0$, so that we have

$$\ell^*\hat{\omega}_1 = \sqrt{p}\omega_1 + \frac{r}{\sqrt{p}}\omega_2. \quad (5-1)$$

The pullback $\ell^*\hat{\omega}_2$ must be a multiple of ω_2 . Indeed, $\ell^*\hat{\omega}_2$ is π -semibasic and we obtain

$$\begin{aligned} (X \lrcorner \ell^*\hat{\omega}_2)(x, v) &= \hat{\omega}_2(d\ell(X(x, v))) = \hat{g}(\hat{J}\ell(x, v), (d\hat{\pi} \circ d\ell)(X(x, v))) \\ &= \frac{1}{\sqrt{\hat{g}(v, v)}} \hat{g}(\hat{J}v, d\pi(X(x, v))) = \frac{\hat{g}(\hat{J}v, v)}{\sqrt{\hat{g}(v, v)}} = 0. \end{aligned}$$

Recall that the area form $d\hat{\sigma}$ of $\hat{g}$ satisfies $\hat{\pi}^*d\hat{\sigma} = \hat{\omega}_1 \wedge \hat{\omega}_2$; hence

$$\ell^*(\hat{\omega}_1 \wedge \hat{\omega}_2) = \pi^*d\hat{\sigma} = \sqrt{pq - r^2} \omega_1 \wedge \omega_2.$$

Thus we must have

$$\ell^*\hat{\omega}_2 = \frac{\sqrt{pq - r^2}}{\sqrt{p}} \omega_2. \quad (5-2)$$

Since the Lie derivative of $\pi^*\hat{g}$ with respect to V vanishes identically, we compute that $V\sqrt{p} = r/\sqrt{p}$. Moreover, since $\sqrt{pq - r^2}$ is the π -pullback of a function on M , we obtain

$$V\left(\frac{\sqrt{pq - r^2}}{\sqrt{p}}\right) = -\frac{r\sqrt{pq - r^2}}{p^{3/2}}.$$

Pulling back the structure equation $d\hat{\omega}_2 = -\hat{\psi} \wedge \hat{\omega}_1$ whilst using (5-1) and (5-2) gives

$$\begin{aligned} \ell^*(d\hat{\omega}_2) &= d(\ell^*\hat{\omega}_2) = d\left(\frac{\sqrt{pq - r^2}}{\sqrt{p}}\omega_2\right) = \left(\hat{a}\omega_1 - \frac{r\sqrt{pq - r^2}}{p^{3/2}}\psi\right) \wedge \omega_2 - \frac{\sqrt{pq - r^2}}{\sqrt{p}}\psi \wedge \omega_1 \\ &= -\ell^*\hat{\psi} \wedge \ell^*\hat{\omega}_1 = -\ell^*\hat{\psi} \wedge \left(\sqrt{p}\omega_1 + \frac{r}{\sqrt{p}}\omega_2\right) \end{aligned}$$

for some unique real-valued function $\hat{a}$ on SM . Comparing the coefficients in the above equations, it follows that

$$\ell^*\hat{\psi} = a\omega_1 + b\omega_2 + \frac{\sqrt{pq - r^2}}{p}\psi \quad (5-3)$$

for some unique real-valued functions a, b on SM . In particular, we obtain

$$\ell^*\hat{\Theta} = \ell^*(\hat{\omega}_1 \wedge \hat{\omega}_2 \wedge \hat{\psi}) = \left(\frac{pq - r^2}{p}\right)\omega_1 \wedge \omega_2 \wedge \psi,$$

as claimed. $\square$

We use this lemma to derive the following observation about general thermostats:

Proposition 5.2. *If two thermostats determined by pairs (g, λ) and $(\hat{g}, \hat{\lambda})$ have the same unparametrised geodesics, then*

$$\sqrt{p} (\hat{V}\hat{\lambda} \circ \ell) = F \log \left(\frac{pq - r^2}{p^{3/2}} \right) + V\lambda.$$

As an immediate application we obtain the following classical fact:

Corollary 5.3. *Let g and $\hat{g}$ be two Riemannian metrics on M having the same unparametrised geodesics. Then $p/(pq - r^2)^{2/3}$ is an integral for the geodesic flow of g .*

Proof. This special case corresponds to $\lambda = \hat{\lambda} = 0$ and hence Proposition 5.2 implies

$$0 = X \log \left(\frac{pq - r^2}{p^{3/2}} \right) = -\frac{3}{2} X \log \left(\frac{p}{(pq - r^2)^{2/3}} \right) = -\frac{3}{2} \frac{(pq - r^2)^{2/3}}{p} X \left(\frac{p}{(pq - r^2)^{2/3}} \right). \quad \square$$

In order to prove Proposition 5.2 we also recall a general lemma whose proof is elementary and thus omitted.

Lemma 5.4. *Let X be a vector field on a manifold M with volume form Ω . Let f and $s > 0$ be smooth functions. Then*

$$\text{Div}_\Omega(fX) = Xf + f \text{Div}_\Omega X \quad \text{and} \quad \text{Div}_{s\Omega}(X) = X \log s + \text{Div}_\Omega X.$$

Proof of Proposition 5.2. This follows from Lemmas 5.1 and 5.4 and the key fact that if the thermostats have the same unparametrised geodesics then

$$\ell^* \hat{F} = \frac{1}{\sqrt{p}} F. \quad (5-4)$$

To see the last equality, note that we can rephrase the hypothesis as follows. There is a smooth function $\tau : SM \times \mathbb{R} \rightarrow \mathbb{R}$ implementing the time change so that

$$\ell \circ \phi_{\tau(x,v,t)}(x, v) = \hat{\phi}_t \circ \ell(x, v).$$

Differentiating this with respect to t and setting $t = 0$ gives

$$d\ell(fF) = \hat{F} \circ \ell,$$

where

$$f(x, v) := \frac{d}{dt} \tau(x, v, t)|_{t=0}.$$

To check that f has the desired form, apply $d\hat{\pi}$ to the last equation to get $fv = v/\sqrt{\hat{g}(v, v)}$.

Writing $s := (pq - r^2)/p$ and taking the divergence of (5-4) with respect to $\ell^*\hat{\Theta} = s\Theta$ gives

$$\begin{aligned} \text{Div}_{s\Theta}(\sqrt{p} \ell^* \hat{F}) &= (\ell^* \hat{F}) \sqrt{p} + \sqrt{p} \text{Div}_{s\Theta}(\ell^* \hat{F}) \\ &= (1/\sqrt{p}) F \sqrt{p} + \sqrt{p} \text{Div}_{\ell^*\hat{\Theta}}(\ell^* \hat{F}) \\ &= F(\log \sqrt{p}) + \sqrt{p}(\text{Div}_{\hat{\Theta}} \hat{F}) \circ \ell \\ &= \text{Div}_{s\Theta} F = F \log s + \text{Div}_\Theta F, \end{aligned}$$

where we have used Lemma 5.4. Since $\text{Div}_\Theta F = V\lambda$ and $\text{Div}_\Theta \widehat{F} = \widehat{V}\widehat{\lambda}$, this last equation is equivalent to

$$\sqrt{p}(\widehat{V}\widehat{\lambda} \circ \ell) = F \log\left(\frac{s}{\sqrt{p}}\right) + V\lambda,$$

which proves the claim. $\square$

Remark 5.5. Note that the crucial identity (5-4) also follows from a different argument. Since the orbits of F and $\widehat{F}$ project onto the same unparametrised curves, there must exist a smooth function w on SM , so that $\ell^*\widehat{F} = wF$. From (5-1), (5-2) and (5-3), we compute

$$\ell^*\widehat{X} = \frac{1}{\sqrt{p}}X - \frac{a\sqrt{p}}{\sqrt{pq-r^2}}V \quad \text{and} \quad \ell^*\widehat{V} = \frac{p}{\sqrt{pq-r^2}}V,$$

from which one immediately obtains $w = 1/\sqrt{p}$.

A special case of Proposition 5.2 is the following:

Corollary 5.6. *Suppose the projective thermostat associated to the pair $(g, \lambda) = (g, a - V\theta)$ has the same unparametrised geodesics as the Weyl connection D defined by $(\hat{g}, \alpha)$. Then*

$$u = \frac{3}{2} \log\left(\frac{p}{(pq-r^2)^{2/3}}\right)$$

satisfies the transport equation

$$Fu = Va + \beta, \tag{5-5}$$

where $\beta = \theta - \alpha$.

Proof. Applying Proposition 5.2 in the special case $\lambda = a - V\theta$ and $\widehat{\lambda} = -\widehat{V}\alpha$ gives

$$-\sqrt{p}(\widehat{V}\widehat{V}\alpha \circ \ell) = \sqrt{p}(\alpha \circ \ell) = F \log\left(\frac{pq-r^2}{p^{3/2}}\right) + V(a - V\theta),$$

the left-hand side of which is simply α , thought of as a function on SM . Hence we obtain

$$-(Va + \theta - \alpha) = F \log\left(\frac{pq-r^2}{p^{3/2}}\right) = F\left(-\frac{3}{2}(\log p - \frac{2}{3}\log(pq-r^2))\right) = -Fu,$$

as claimed. $\square$

6. The tensor tomography result

In this final section we prove a vanishing theorem for the transport equation $Fu = Va + \beta$, provided the triple (g, A, θ) defining F satisfies certain conditions. Recall that every properly convex projective structure p arises from a triple $(g, A, 0)$ satisfying

$$K_g = -1 + 2|A|_g^2 \quad \text{and} \quad \bar{\partial}A = 0.$$

In particular, we would like to conclude that if such a p contains a Weyl connection, then A must vanish identically and hence p is hyperbolic. It turns out that one can prove a more general vanishing theorem for a class of thermostats arising from a triple (g, A, θ) where A is a differential of degree $m \geq 3$ on M ,

that is, a section of K^m . Suppose $A \in \Gamma(K^m)$. Like in the case $m = 3$ there exists a unique smooth real-valued function a on SM lying in $\mathcal{H}_{-m} \oplus \mathcal{H}_m$, so that $\pi^*A = (V(a)/m + ia)\omega^m$. In particular, to a triple (g, A, θ) we may associate the thermostat $F = X + (a - V\theta)V$. We now have:

Theorem 6.1. *Let M be a closed oriented surface and (g, A, θ) be a triple satisfying*

$$\bar{\partial}A = \left(\frac{1}{2}(m-1)\right)(\theta - i \star_g \theta) \otimes A \quad \text{and} \quad K_g - \delta_g \theta + (2-m)|A|_g^2 \leq 0.$$

Let F denote the vector field of the thermostat determined by (g, A, θ) . Suppose there is a 1-form $\beta \in \Omega^1(M)$ and a function $u \in C^\infty(SM)$ such that

$$Fu = Va + \beta.$$

Then $A = 0$ and β is exact.

Let us first verify that this gives the desired statement.

Corollary 6.2. *Let $(M, \mathfrak{p})$ be a closed oriented properly convex projective surface with $\chi(M) < 0$ and with $\mathfrak{p}$ containing a Weyl connection D . Then $\mathfrak{p}$ is hyperbolic and moreover D is the Levi-Civita connection of the hyperbolic metric.*

Proof. By a result of [Calabi 1972], if $m = 3$ and (g, A) satisfy

$$K_g = -1 + 2|A|_g^2 \quad \text{and} \quad \bar{\partial}A = 0,$$

then $K_g \leq 0$. In particular, the triple $(g, A, 0)$ satisfies the assumptions of Theorem 6.1, and Corollary 5.6 implies that we have a solution u to the transport equation $Fu = Va + \beta$. Hence the theorem gives right away that A vanishes identically and hence $\mathfrak{p}$ is hyperbolic. In particular, the Levi-Civita connection ${}^g\nabla$ of the hyperbolic metric and the connection D both lie in $\mathfrak{p}$ and hence are projectively equivalent, but this can happen if and only if ${}^g\nabla = D$, by Corollary 4.6. $\square$

Remark 6.3. In [Mettler 2019] the notion of a minimal Lagrangian connection is introduced. These are torsion-free connections on TM of the form $\nabla = D + B$, where (g, A, θ) defining D and B are subject to the equations

$$K_g - \delta_g \theta = -1 + 2|A|_g^2, \quad \bar{\partial}A = (\theta - i \star_g \theta) \otimes A, \quad d\theta = 0.$$

In particular, on a closed oriented surface of negative Euler characteristic every properly convex projective structure arises from a minimal Lagrangian connection. Another immediate consequence of Theorem 6.1 and Corollary 4.6 thus is:

Corollary 6.4. *Let M be a closed oriented surface of negative Euler characteristic and ∇ a minimal Lagrangian connection arising from the triple (g, A, θ) . Suppose $|A|_g^2 \leq 1$ and that ∇ is projectively equivalent to a Weyl connection D . Then A vanishes identically and hence $\nabla = D$.*

In order to show the theorem we use the following L^2 identity proved in [Jane and Paternain 2009, equation (5)], which is in turn an extension of an identity in [Sharafutdinov and Uhlmann 2000] for

geodesic flows. The identity holds for arbitrary thermostats $F = X + \lambda V$. If we let $H_c := H + cV$, where $c : SM \rightarrow \mathbb{R}$ is any smooth function, then

$$2\langle H_c u, VFu \rangle = \|Fu\|^2 + \|H_c u\|^2 - \langle Fc + c^2 + K_g - H_c \lambda + \lambda^2, (Vu)^2 \rangle, \quad (6-1)$$

where u is any smooth function. All norms and inner products are L^2 with respect to the volume form Θ .

We also need the following lemma whose proof is a straightforward calculation (see [Mettler and Paternain 2019, Lemma 4.1] for a proof).

Lemma 6.5. *We have*

$$\bar{\partial} A = \left(\frac{1}{2}(m-1)\right)(\theta - i \star_g \theta) \otimes A$$

if and only if

$$XVa - mHa - (m-1)(\theta Va - maV\theta) = 0.$$

Proof of Theorem 6.1. Without loss of generality we may assume that β has zero divergence. Indeed if not, a standard application of scalar elliptic PDE theory shows that we can always find a smooth function h on M such that $\beta + dh$ has zero divergence. Now note that $F(u+h) = Va + \beta + dh$.

A calculation shows that if we pick $c = \theta + V(a)/m$, then

$$Fc + c^2 + K_g - H_c \lambda + \lambda^2 = K_g - \delta_g \theta + (1-m)|A|_g^2,$$

where we use that

$$\pi^*|A|_g^2 = (Va)^2/m^2 + a^2 \quad \text{and} \quad \pi^*\delta_g \theta = -(X\theta + HV\theta);$$

hence for this choice of c , (6-1) simplifies to

$$2\langle H_c u, VFu \rangle - \| |A|_g Vu \|^2 = \|Fu\|^2 + \|H_c u\|^2 - \langle K_g - \delta_g \theta + (2-m)|A|_g^2, (Vu)^2 \rangle. \quad (6-2)$$

If $Fu = Va + \beta$, then $VFu = -m^2a + V\beta$. Using that X and H preserve Θ and that $XVa - mHa - (m-1)(\theta Va - maV\theta) = 0$ we compute

$$\begin{aligned} 2\langle H_c u, -m^2a \rangle &= -2m^2\langle Hu, a \rangle - 2m^2\langle cVu, a \rangle \\ &= 2m^2\langle u, Ha \rangle - 2m^2\langle cVu, a \rangle \\ &= -2m^2\langle Xu, V(a)/m \rangle - 2m(m-1)\langle u, \theta Va - maV\theta \rangle - 2m^2\langle cVu, a \rangle \\ &= -2m\|Va\|^2 = -2m^3\|a\|^2, \end{aligned}$$

where the last equation is obtained using that $Xu = \beta + Va - (a - V\theta)Vu$, $\langle \beta, Va \rangle = 0$ and $c = \theta + V(a)/m$.

Using that X and H preserve Θ and that $X\beta + HV\beta = 0$ (β is assumed to have zero divergence) we compute

$$\begin{aligned} 2\langle H_c u, V\beta \rangle &= 2\langle Hu, V\beta \rangle + 2\langle cVu, V\beta \rangle \\ &= -2\langle u, HV\beta \rangle + 2\langle cVu, V\beta \rangle \\ &= -2\langle Xu, \beta \rangle + 2\langle cVu, V\beta \rangle \\ &= -2\|\beta\|^2 + 2\langle (a - V\theta)Vu, \beta \rangle + 2\langle cVu, V\beta \rangle \\ &= -2\|\beta\|^2 + 2\langle aVu, \beta \rangle + 2\langle (VaVu)/m, V\beta \rangle, \end{aligned}$$

where the penultimate equation is obtained using that $Xu = \beta + Va - (a - V\theta)Vu$ and $\langle \beta, Va \rangle = 0$. The last equation uses that $c = \theta + V(a)/m$ and

$$V(\theta V\beta - V\theta\beta) = 0.$$

Inserting these calculations back into (6-2), we derive

$$\begin{aligned} -2m^3 \|a\|^2 - 2\|\beta\|^2 + 2\langle aVu, \beta \rangle + 2\langle (VaVu)/m, V\beta \rangle - \| |A|_g Vu \|^2 \\ = \|Fu\|^2 + \|Hcu\|^2 - \langle K_g - \delta_g \theta + (2-m)|A|_g^2, (Vu)^2 \rangle. \end{aligned}$$

Since $|A|_g^2 = a^2 + (Va)^2/m^2$ this can be rewritten as

$$-2m^3 \|a\|^2 - \|\beta - aVu\|^2 - \|V\beta - VaVu/m\|^2 = \|Fu\|^2 + \|Hcu\|^2 - \langle K_g - \delta_g \theta + (2-m)|A|_g^2, (Vu)^2 \rangle,$$

where we have used that $\|\beta\|^2 = \|V\beta\|^2$. By hypothesis the right-hand side is ≥ 0 , which gives right away that $a = \beta = 0$. $\square$

Appendix: Deriving the transport equation

Here we sketch how to derive the transport equation for the function u starting from the PDE

$$D''\mu - \mu D'\mu = \Phi\mu^3 + \bar{\Phi}$$

for the Beltrami differential μ . Let (g, A, θ) be the triple encoding $\mathfrak{p}$ so that the connection form of D on SM is (see (2-8)) $\kappa = i\psi - 2\theta_1\omega$, where we write $\theta_1 = \frac{1}{2}(\theta - iV\theta)$. Moreover, on SM the section Φ of $K^2 \otimes \bar{K}^*$ is represented by $a_3 = \frac{1}{3}Va + ia$, where $a(v) = \text{Re } A(Jv, Jv, Jv)$, $v \in SM$. Writing μ_{-2} for the complex-valued function on SM representing the Beltrami differential μ and $\mu_2 = \bar{\mu}_{-2}$, the PDE for μ is equivalent to

$$d\mu_{-2} = \mu'_{-2}\omega + (\mu_{-2}\mu'_{-2} + a_3\mu_{-2}^3 + \bar{a}_3)\bar{\omega} + \bar{\kappa}\mu_{-2} - \kappa\mu_{-2},$$

where μ'_{-2} is a complex-valued function on SM . Since μ_{-2} represents a section of $\bar{K} \otimes K^* \simeq K^{-2}$, writing $\eta_{\pm} = \frac{1}{2}(X \mp iH)$ we also have

$$d\mu_{-2} = \eta_+(\mu_{-2})\omega + \eta_-(\mu_{-2})\bar{\omega} - 2i\mu_{-2}\psi.$$

Thus the PDE is equivalent to the system

$$\eta_-\mu_{-2} - \mu_{-2}\eta_+\mu_{-2} = a_3\mu_{-2}^3 - 2\mu_{-2}^2\theta_1 - 2\mu_{-2}\bar{\theta}_1 + \bar{a}_3 \quad (\text{A-1})$$

and $V\mu_{-2} = -2i\mu_{-2}$. The Beltrami differential only defines a conformal equivalence class $[\hat{g}]$. We may fix a metric $\hat{g} \in [\hat{g}]$ by requiring

$$\frac{1}{2}(p+q) = \frac{1+|\mu_2|^2}{(1-|\mu_2|^2)^4},$$

where again we specify the metric $\hat{g}$ in terms of the functions p, q, r . Explicitly, we have

$$\frac{1}{2}(p-q) = \frac{\mu_{-2} + \mu_2}{(1-|\mu_2|^2)^4} \quad \text{and} \quad r = \frac{i(\mu_2 - \mu_{-2})}{(1-|\mu_2|^2)^4}.$$

In particular, this yields

$$h := \frac{p}{(pq - r^2)^{2/3}} = (\mu_{-2} + 1)(\mu_2 + 1).$$

Writing $F = X + (a - V\theta)V$ and using (A-1), a lengthy but straightforward calculation shows that

$$Fh = \frac{2}{3}hVa + 2h \operatorname{Re}(a_3\mu_{-2}^2 - \mu_2a_{-3} - 2\mu_2\theta_{-1} + \eta_+\mu_{-2}).$$

Hence if we define $u = \frac{3}{2} \log h$, then we obtain

$$Fu - Va = 3 \operatorname{Re}(a_3\mu_{-2}^2 - \mu_2a_{-3} - 2\mu_2\theta_{-1} + \eta_+\mu_{-2}).$$

Note that the right-hand side of the last equation lies in $\mathcal{H}_{-1} \oplus \mathcal{H}_1$; hence there exists a 1-form β on M so that

$$Fu = Va + \beta,$$

which is the transport equation (5-5).

Acknowledgements

The authors are grateful to Nigel Hitchin for helpful conversations. A part of the research for this article was carried out while Mettler was visiting FIM at ETH Zürich. Mettler thanks FIM for its hospitality. Mettler was partially funded by the priority programme SPP 2026 “Geometry at Infinity” of DFG. Paternain was partially supported by EPSRC grant EP/R001898/1.

References

- [Besse 1987] A. L. Besse, *Einstein manifolds*, Ergebnisse der Mathematik (3) **10**, Springer, 1987. MR Zbl
- [Bryant et al. 2009] R. Bryant, M. Dunajski, and M. Eastwood, “Metrisability of two-dimensional projective structures”, *J. Differential Geom.* **83**:3 (2009), 465–499. MR Zbl
- [Calabi 1972] E. Calabi, “Complete affine hyperspheres, I”, pp. 19–38 in *Symposia Mathematica, X* (Rome, 1971), Academic Press, London, 1972. MR Zbl
- [Croke and Sharafutdinov 1998] C. B. Croke and V. A. Sharafutdinov, “Spectral rigidity of a compact negatively curved manifold”, *Topology* **37**:6 (1998), 1265–1273. MR Zbl
- [Dubois-Violette 1983] M. Dubois-Violette, “Structures complexes au-dessus des variétés, applications”, pp. 1–42 in *Mathématique et physique* (Paris, 1979/1982), edited by L. Boutet de Monvel et al., Progr. Math. **37**, Birkhäuser, Boston, 1983. MR Zbl
- [Dumas and Wolf 2015] D. Dumas and M. Wolf, “Polynomial cubic differentials and convex polygons in the projective plane”, *Geom. Funct. Anal.* **25**:6 (2015), 1734–1798. MR Zbl
- [Guillemin and Kazhdan 1980] V. Guillemin and D. Kazhdan, “Some inverse spectral results for negatively curved 2-manifolds”, *Topology* **19**:3 (1980), 301–312. MR Zbl
- [Hitchin 1982] N. J. Hitchin, “Complex manifolds and Einstein’s equations”, pp. 73–99 in *Twistor geometry and nonlinear systems* (Primorsko, Bulgaria, 1980), edited by H.-D. Doebner and T. D. Palev, Lecture Notes in Math. **970**, Springer, 1982. MR Zbl
- [Hitchin 1992] N. J. Hitchin, “Lie groups and Teichmüller space”, *Topology* **31**:3 (1992), 449–473. MR Zbl
- [Jane and Paternain 2009] D. Jane and G. P. Paternain, “On the injectivity of the X-ray transform for Anosov thermostats”, *Discrete Contin. Dyn. Syst.* **24**:2 (2009), 471–487. MR Zbl

- [Labourie 2007] F. Labourie, “Flat projective structures on surfaces and cubic holomorphic differentials”, *Pure Appl. Math. Q.* **3**:4 (2007), 1057–1099. MR Zbl
- [Loftin 2001] J. C. Loftin, “Affine spheres and convex $\mathbb{RP}^n$ -manifolds”, *Amer. J. Math.* **123**:2 (2001), 255–274. MR Zbl
- [Matveev and Topalov 2000] V. S. Matveev and P. J. Topalov, “Metric with ergodic geodesic flow is completely determined by unparameterized geodesics”, *Electron. Res. Announc. Amer. Math. Soc.* **6** (2000), 98–104. MR Zbl
- [Mettler 2014] T. Mettler, “Weyl metrisability of two-dimensional projective structures”, *Math. Proc. Cambridge Philos. Soc.* **156**:1 (2014), 99–113. MR Zbl
- [Mettler 2015a] T. Mettler, “Extremal conformal structures on projective surfaces”, preprint, 2015. To appear in *Ann. Scuola Norm. Sup. Pisa Cl. Sci.* arXiv
- [Mettler 2015b] T. Mettler, “Geodesic rigidity of conformal connections on surfaces”, *Math. Z.* **281**:1-2 (2015), 379–393. MR Zbl
- [Mettler 2019] T. Mettler, “Minimal Lagrangian connections on compact surfaces”, *Adv. Math.* **354** (2019), art. id. 106747. MR Zbl
- [Mettler and Paternain 2019] T. Mettler and G. P. Paternain, “Holomorphic differentials, thermostats and Anosov flows”, *Math. Ann.* **373**:1-2 (2019), 553–580. MR Zbl
- [O’Brian and Rawnsley 1985] N. R. O’Brian and J. H. Rawnsley, “Twistor spaces”, *Ann. Global Anal. Geom.* **3**:1 (1985), 29–58. MR Zbl
- [Paternain et al. 2013] G. P. Paternain, M. Salo, and G. Uhlmann, “Tensor tomography on surfaces”, *Invent. Math.* **193**:1 (2013), 229–247. MR Zbl
- [Paternain et al. 2014] G. P. Paternain, M. Salo, and G. Uhlmann, “Spectral rigidity and invariant distributions on Anosov surfaces”, *J. Differential Geom.* **98**:1 (2014), 147–181. MR Zbl
- [Sharafutdinov and Uhlmann 2000] V. Sharafutdinov and G. Uhlmann, “On deformation boundary rigidity and spectral rigidity of Riemannian surfaces with no focal points”, *J. Differential Geom.* **56**:1 (2000), 93–110. MR Zbl
- [Spivak 1999] M. Spivak, *A comprehensive introduction to differential geometry, II*, 3rd ed., Publish or Perish, Houston, 1999. Zbl
- [Wang 1991] C. P. Wang, “Some examples of complete hyperbolic affine 2-spheres in $\mathbb{R}^3$ ”, pp. 271–280 in *Global differential geometry and global analysis* (Berlin, 1990), edited by D. Ferus et al., Lecture Notes in Math. **1481**, Springer, 1991. MR Zbl

Received 14 Jun 2018. Revised 20 Apr 2019. Accepted 1 Jun 2019.

THOMAS METTLER: mettler@math.uni-frankfurt.de

Institut für Mathematik, Goethe-Universität Frankfurt, Frankfurt am Main, Germany

GABRIEL P. PATERNAIN: g.p.paternain@dpms.cam.ac.uk

Department of Pure Mathematics and Mathematical Statistics, University of Cambridge, Cambridge, United Kingdom

Analysis & PDE

msp.org/apde

EDITORS

EDITOR-IN-CHIEF

Patrick Gérard
patrick.gerard@math.u-psud.fr
Université Paris Sud XI
Orsay, France

BOARD OF EDITORS

Massimiliano Berti	Scuola Intern. Sup. di Studi Avanzati, Italy berti@sissa.it	Gilles Pisier	Texas A&M University, and Paris 6 pisier@math.tamu.edu
Michael Christ	University of California, Berkeley, USA mchrist@math.berkeley.edu	Tristan Rivière	ETH, Switzerland riviere@math.ethz.ch
Charles Fefferman	Princeton University, USA cf@math.princeton.edu	Igor Rodnianski	Princeton University, USA irod@math.princeton.edu
Ursula Hamenstaedt	Universität Bonn, Germany ursula@math.uni-bonn.de	Yum-Tong Siu	Harvard University, USA siu@math.harvard.edu
Vadim Kaloshin	University of Maryland, USA vadim.kaloshin@gmail.com	Terence Tao	University of California, Los Angeles, USA tao@math.ucla.edu
Herbert Koch	Universität Bonn, Germany koch@math.uni-bonn.de	Michael E. Taylor	Univ. of North Carolina, Chapel Hill, USA met@math.unc.edu
Izabella Laba	University of British Columbia, Canada ilaba@math.ubc.ca	Gunther Uhlmann	University of Washington, USA gunther@math.washington.edu
Richard B. Melrose	Massachusetts Inst. of Tech., USA rbm@math.mit.edu	András Vasy	Stanford University, USA andras@math.stanford.edu
Frank Merle	Université de Cergy-Pontoise, France Frank.Merle@u-cergy.fr	Dan Virgil Voiculescu	University of California, Berkeley, USA dvv@math.berkeley.edu
William Minicozzi II	Johns Hopkins University, USA minicozz@math.jhu.edu	Steven Zelditch	Northwestern University, USA zelditch@math.northwestern.edu
Clément Mouhot	Cambridge University, UK c.mouhot@dpms.cam.ac.uk	Maciej Zworski	University of California, Berkeley, USA zworski@math.berkeley.edu
Werner Müller	Universität Bonn, Germany mueller@math.uni-bonn.de		

PRODUCTION

production@msp.org
Silvio Levy, Scientific Editor

See inside back cover or msp.org/apde for submission instructions.

The subscription price for 2020 is US \$340/year for the electronic version, and \$550/year (+\$60, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues from the last three years and changes of subscriber address should be sent to MSP.

Analysis & PDE (ISSN 1948-206X electronic, 2157-5045 printed) at Mathematical Sciences Publishers, 798 Evans Hall #3840, c/o University of California, Berkeley, CA 94720-3840, is published continuously online. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices.

APDE peer review and production are managed by EditFlow® from MSP.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2020 Mathematical Sciences Publishers

ANALYSIS & PDE

Volume 13 No. 4 2020

Estimates for the Navier–Stokes equations in the half-space for nonlocalized data	945
YASUNORI MAEKAWA, HIDEYUKI MIURA and CHRISTOPHE PRANGE	
Almost-sure scattering for the radial energy-critical nonlinear wave equation in three dimensions	1011
BJOERN BRINGMANN	
On the existence of translating solutions of mean curvature flow in slab regions	1051
THEODORA BOURNI, MAT LANGFORD and GIUSEPPE TINAGLIA	
Convex projective surfaces with compatible Weyl connection are hyperbolic	1073
THOMAS METTLER and GABRIEL P. PATERNAIN	
Stability of small solitary waves for the one-dimensional NLS with an attractive delta potential	1099
SATOSHI MASAKI, JASON MURPHY and JUN-ICHI SEGATA	
Geometric regularity for elliptic equations in double-divergence form	1129
RAIMUNDO LEITÃO, EDGARD A. PIMENTEL and MAKSON S. SANTOS	
Nonexistence of global characteristics for viscosity solutions	1145
VALENTINE ROOS	
Strichartz estimates for the Schrödinger flow on compact Lie groups	1173
YUNFENG ZHANG	
Parabolic L^p Dirichlet boundary value problem and VMO-type time-varying domains	1221
MARTIN DINDOŠ, LUKE DYER and SUKJUNG HWANG	



2157-5045(2020)13:4;1-A