

ANALYSIS & PDE

Volume 13

No. 5

2020

PAATA IVANISVILI AND CONNOR MOONEY

**SHARPENING THE TRIANGLE INEQUALITY:
ENVELOPES BETWEEN L^2 AND L^p SPACES**

SHARPENING THE TRIANGLE INEQUALITY: ENVELOPES BETWEEN L^2 AND L^p SPACES

PAATA IVANISVILI AND CONNOR MOONEY

Motivated by the inequality $\|f + g\|_2^2 \leq \|f\|_2^2 + 2\|fg\|_1 + \|g\|_2^2$, Carbery (2009) raised the question of what is the “right” analogue of this estimate in L^p for $p \neq 2$. Carlen, Frank, Ivanisvili and Lieb (2018) recently obtained an L^p version of this inequality by providing upper bounds for $\|f + g\|_p^p$ in terms of the quantities $\|f\|_p^p$, $\|g\|_p^p$ and $\|fg\|_{p/2}^{p/2}$ when $p \in (0, 1] \cup [2, \infty)$, and lower bounds when $p \in (-\infty, 0) \cup (1, 2)$, thereby proving (and improving) the suggested possible inequalities of Carbery. We continue investigation in this direction by refining the estimates of Carlen, Frank, Ivanisvili and Lieb. We obtain upper bounds for $\|f + g\|_p^p$ also when $p \in (-\infty, 0) \cup (1, 2)$ and lower bounds when $p \in (0, 1] \cup [2, \infty)$. For $p \in [1, 2]$ we extend our upper bounds to any finite number of functions. In addition, we show that all our upper and lower bounds of $\|f + g\|_p^p$ for $p \in \mathbb{R}$, $p \neq 0$, are the best possible in terms of the quantities $\|f\|_p^p$, $\|g\|_p^p$ and $\|fg\|_{p/2}^{p/2}$, and we characterize the equality cases.

1. Introduction

For any real-valued functions $f, g \in L^p$ on an arbitrary measure space, and any $p \geq 1$, one has the inequality

$$\|f + g\|_p^p \leq 2^{p-1}(\|f\|_p^p + \|g\|_p^p). \quad (1)$$

The estimate (1) follows from the fact that the map $x \mapsto |x|^p$ is convex. If $f = g$ in (1) then the constant 2^{p-1} is sharp and the inequality becomes equality. On the other hand, if f and g have disjoint supports then the constant 2^{p-1} is not needed. We remark that the estimate (1) reflects the convexity of the unit ball in L^p , which is equivalent to the usual L^p triangle (Minkowski) inequality; see, e.g., [Carlen et al. 2020a].

Carbery [2009] asked under what conditions on the sequence of functions $\{f_j\} \subset L^p$ the inequality $\sum \|f_j\|_p^p < \infty$ would imply $\sum f_j \in L^p$. If we try to adapt the inequality (1) to say n functions $f_1, f_2, \dots, f_n$ instead of two, then the constant 2^{p-1} should be replaced by n^{p-1} , which grows with n . To remove dependence on n , Carbery suggested several extensions of inequality (1) which were motivated by the estimate $\|f + g\|_2^2 \leq \|f\|_2^2 + 2\|fg\|_1 + \|g\|_2^2$. All of them involve the extra parameter $\|fg\|_{p/2}^{p/2}$, which measures the “overlap” between the functions, and the strongest one in the case of two functions he could prove only for indicator functions of sets. Recently a sharpened form of the triangle inequality was obtained [Carlen et al. 2020a], which implied the proposed estimates of Carbery. Namely, take any

$p \in \mathbb{R} \setminus \{0\}$, and put

$$\|f\|_p := \left(\int_X |f|^p d\mu \right)^{1/p} \quad \text{and} \quad \Gamma_p := \frac{2\|fg\|_{p/2}^{p/2}}{\|f\|_p^p + \|g\|_p^p}.$$

Then

$$\|f + g\|_p^p \leq (1 + \Gamma_p^{2/p})^{p-1} (\|f\|_p^p + \|g\|_p^p) \quad (2)$$

holds true if $p \in (0, 1] \cup [2, \infty)$, and the inequality reverses if $p \in (-\infty, 0) \cup (1, 2)$, where in the latter case we assume that f, g are positive almost everywhere. Since by Cauchy–Schwarz $\Gamma_p \in [0, 1]$ for all $p \in \mathbb{R} \setminus \{0\}$, we see that (2) improves on the trivial bound (1).

In this paper we continue investigation in this direction and we address the following questions:

Question 1. Can one further sharpen the right-hand side of the estimate (2) if we are allowed to use only the quantities $\|f\|_p, \|g\|_p, \|fg\|_{p/2}$?

Question 2. What is the optimal upper bound on $\|f + g\|_p^p$ in terms of the quantities $\|f\|_p, \|g\|_p, \|fg\|_{p/2}$, also when $p \in (-\infty, 0) \cup (1, 2)$? Additionally we consider the same question about lower bounds on $\|f + g\|_p^p$, also when $p \in (0, 1] \cup [2, \infty)$.

Question 3. Can one extend these estimates to more than two functions?

We will give complete answers to Questions 1 and 2, and we will provide an answer to Question 3 when $p > 0$. In particular we show that, for $p \in [1, 2]$, if $\sum_j \|f_j\|_p^p < \infty$ and $\sum_{i < j} \|f_i f_j\|_{p/2}^{p/2} < \infty$, then $\sum_j f_j \in L^p$.

2. Main results

Let $(X, \mathcal{A}, \mu)$ be an arbitrary measure space. In what follows we consider functions f, g on X that are measurable and nonnegative. Given $p \in \mathbb{R} \setminus \{0\}$ we will always assume that $\|f\|_p^p, \|g\|_p^p < \infty$. When $p < 0$ we allow f, g to take the value $+\infty$, where we understand $f^p, g^p = 0$.

Theorem 2.1. For any $p \in (0, 1] \cup [2, \infty)$, and any nonnegative f, g on any measure space we have

$$\|f + g\|_p^p \leq \left(\left(\frac{1 + \sqrt{1 - \Gamma_p^2}}{2} \right)^{1/p} + \left(\frac{1 - \sqrt{1 - \Gamma_p^2}}{2} \right)^{1/p} \right)^p (\|f\|_p^p + \|g\|_p^p). \quad (3)$$

The inequality reverses if $p \in (-\infty, 0) \cup [1, 2]$. Equality holds if $(fg)^{p/2} = k(f^p + g^p)$ for some constant $k \in [0, \frac{1}{2}]$.

Remark 2.2. The right-hand side of (3) is the best possible in the following sense: consider the measure space $([0, 1], \mathcal{B}, dx)$. Pick any nonnegative numbers x, y and z such that $0 \leq z \leq \sqrt{xy}$. Then, for any $p \in (0, 1] \cup [2, \infty)$ the supremum of the left-hand side of (3) over all nonnegative f, g with fixed $\|f\|_p^p = x, \|g\|_p^p = y, \|fg\|_{p/2}^{p/2} = z$ coincides with the right-hand side of (3). Similarly, for any $p \in (-\infty, 0) \cup [1, 2]$ the infimum of the left-hand side of (3) over all such f, g coincides with the right-hand side of (3). We justify this remark in Section 3.

Remark 2.2 implies in particular that **Theorem 2.1** refines the estimate (2). As a consequence we have the following peculiar estimate:

Corollary 2.3. For any $p \in (0, 1] \cup [2, \infty)$ and any number $\Gamma \in [0, 1]$, we have

$$\left(\left(\frac{1 + \sqrt{1 - \Gamma^2}}{2} \right)^{1/p} + \left(\frac{1 - \sqrt{1 - \Gamma^2}}{2} \right)^{1/p} \right)^p \leq (1 + \Gamma^{2/p})^{p-1}. \quad (4)$$

The inequality reverses if $p \in (-\infty, 0) \cup [1, 2]$.

If we set $\Gamma := 2(ab)^{p/2}/(a^p + b^p)$ for nonnegative a, b , then after a short computation inequality (4) becomes

$$\frac{(a+b)^p}{a^p + b^p} \leq \left(1 + \left(2 \frac{(ab)^{p/2}}{a^p + b^p} \right)^{2/p} \right)^{p-1}. \quad (5)$$

This estimate was previously obtained in [Carlen et al. 2020a] (where it was also shown to be equivalent to the inequality (2)), and the arguments are quite involved.

Remark 2.4. If we let $q := 1/p$ and $x = \sqrt{1 - \Gamma^2}$, then inequality (4) can also be written as the two-point-type inequality

$$\frac{(1+x)^q + (1-x)^q}{2} \leq \left(\frac{1 + (1-x^2)^q}{2} \right)^{1-q} \quad (6)$$

for all $q \in (-\infty, \frac{1}{2}] \cup [1, \infty)$, $x \in [0, 1]$, and the inequality reverses if $q \in [\frac{1}{2}, 1)$. This inequality is reminiscent of Bonami's two-point inequality

$$\left(\frac{|y + u\sqrt{(p-1)/(q-1)}|^q + |y - u\sqrt{(p-1)/(q-1)}|^q}{2} \right)^{1/q} \leq \left(\frac{|y+u|^p + |y-u|^p}{2} \right)^{1/p}, \quad (7)$$

which holds true for all $y, u \in \mathbb{R}$ and $1 \leq p \leq q < \infty$; see [Bonami 1970]. Indeed, if we take $y = 1$, $p = 2$, and $u = x\sqrt{q-1}$ then we get

$$\frac{|1+x|^q + |1-x|^q}{2} \leq (1 + (q-1)x^2)^{q/2}. \quad (8)$$

The right sides of inequalities (6) and (8) are not comparable. For example, when $x = 1$ the estimate (6) gives better upper bounds for $q > 2$, while near $x = 0$ it gives worse upper bounds.

Next, let $p \in \mathbb{R} \setminus \{0\}$, and set¹

$$C_p := \frac{\min\{\|f\|_p^p, \|g\|_p^p, \|fg\|_{p/2}^{p/2}\}}{\|fg\|_{p/2}^{p/2}}.$$

Theorem 2.5. For any $p \in (1, 2)$ and any nonnegative f, g on any measure space we have

$$\|f + g\|_p^p \leq \|f\|_p^p + \|g\|_p^p + ((C_p^{-1/p} + C_p^{1/p})^p - C_p^{-1} - C_p) \|fg\|_{p/2}^{p/2}. \quad (9)$$

¹If $\|fg\|_{p/2}^{p/2} = 0$ then we set $C_p = 1$.

The inequality reverses if $p \in (0, 1] \cup [2, \infty)$. Equality holds in (9) if one of the following three conditions holds: $f = g$ on $\{fg > 0\}$, $g = \lambda f$ on $\{f > 0\}$ for some $\lambda \geq 1$, or $f = \lambda g$ on $\{g > 0\}$ for some $\lambda \geq 1$.

For $p \in (-\infty, 0)$ we have

$$\|f + g\|_p^p \leq (C_p^{-1/p} + C_p^{1/p})^p \|fg\|_{p/2}^{p/2}. \quad (10)$$

Equality holds in (10) if one of the following three conditions holds: $f = g$ on $\{fg < \infty\}$, $g = \lambda f$ on $\{f < \infty\}$ for some $\lambda \leq 1$, or $f = \lambda g$ on $\{g < \infty\}$ for some $\lambda \leq 1$.

Exactly the same remark as before applies to Theorem 2.5; that is, the right-hand sides of (9) and (10) are the best possible. Together, Theorems 2.1 and 2.5, along with the remarks about optimality, answer Questions 1 and 2.

Finally, we state a partial answer to Question 3 in the case $p > 0$.

Corollary 2.6. For any $p \in [1, 2]$, and any sequence of nonnegative functions $\{f_j\}_{j \geq 1}$ we have

$$\left\| \sum_j f_j \right\|_p^p \leq \sum_j \|f_j\|_p^p + (2^p - 2) \sum_{i < j} \|f_i f_j\|_{p/2}^{p/2}.$$

If $p \in (0, 1] \cup [2, \infty)$ the inequality reverses. Equality holds if and only if

$$\left(\sum_j f_j \right)^p = \sum_j f_j^p + (2^p - 2) \sum_{i < j} (f_i f_j)^{p/2}$$

almost everywhere.

In particular, when $p \in [1, 2]$ we have $\sum_j f_j \in L^p$ provided $\sum_j \|f_j\|_p^p < \infty$ and $\sum_{i < j} \|f_i f_j\|_{p/2}^{p/2} < \infty$.

Remark 2.7. After we finished writing this paper we received the preprint [Carlen et al. 2020b], in which the authors obtain an upper bound for the L^p norm of a sum of N functions in the case $p \geq 2$, in terms of a certain analogue for N functions of the quantity Γ_p . Their estimate complements our result Corollary 2.6, which holds for $p \in (1, 2)$, and is obtained using different techniques.

The rest of the paper is organized as follows. In Section 3 we reduce the proofs of Theorems 2.1 and 2.5, as well as the remarks about their optimality, to computing the concave and convex envelopes of a certain function defined on the boundary of a convex cone in $\mathbb{R}^3$. In Section 4 we compute these envelopes. Finally, in Section 5 we prove Corollary 2.6 using an observation about the proof of Theorem 2.5.

3. Reductions

In this section we reduce Theorems 2.1 and 2.5 to computing explicitly the convex and concave envelopes of a certain function defined on the boundary of a convex cone in $\mathbb{R}^3$. Let

$$\Omega := \{x, y \geq 0, 0 \leq z \leq \sqrt{xy}\}$$

be the convex cone in $\mathbb{R}^3$ whose vertical cross-sections $\Omega \cap \{x + y = c > 0\}$ are half-ellipses. For $p \in \mathbb{R} \setminus \{0\}$ define φ_p on $\partial\Omega$ by

$$\varphi_p(x, y, \sqrt{xy}) = (x^{1/p} + y^{1/p})^p, \quad x, y > 0, \quad \varphi_p(x, y, 0) = \begin{cases} x + y, & p > 0, \\ 0, & p < 0. \end{cases}$$

Let f and g be nonnegative functions on an arbitrary measure space $(X, \mathcal{A}, \mu)$ with $\|f\|_p^p, \|g\|_p^p < \infty$. Note that the triple $(\|f\|_p^p, \|g\|_p^p, \|fg\|_{p/2}^{p/2})$ is in Ω by the Cauchy–Schwarz inequality. By the equality case, if the triple is in $\partial\Omega$ we have $\|f + g\|_p^p = \varphi_p(\|f\|_p^p, \|g\|_p^p, \|fg\|_{p/2}^{p/2})$. Our approach is based on the following lemma:

Lemma 3.1. *Let $p \in \mathbb{R} \setminus \{0\}$, and assume that $H \in C(\Omega)$ is a concave, one-homogeneous function on Ω with $H|_{\partial\Omega} = \varphi_p$. Then*

$$\|f + g\|_p^p \leq H(\|f\|_p^p, \|g\|_p^p, \|fg\|_{p/2}^{p/2}).$$

If H is convex, the inequality reverses.

Proof. By the boundary conditions, we have

$$1 = H\left(\frac{f^p}{(f+g)^p}, \frac{g^p}{(f+g)^p}, \frac{(fg)^{p/2}}{(f+g)^p}\right)$$

on the set $X' = \{f + g > 0\}$ when $p > 0$, or $\{f + g < \infty\}$ when $p < 0$. Integrating this identity with respect to the probability measure $(f + g)^p d\mu / \|f + g\|_p^p$ on X' and applying Jensen's inequality gives

$$1 \leq H\left(\frac{\|f\|_p^p}{\|f + g\|_p^p}, \frac{\|g\|_p^p}{\|f + g\|_p^p}, \frac{\|fg\|_{p/2}^{p/2}}{\|f + g\|_p^p}\right)$$

when H is concave, and the other inequality for H convex. The result follows from the one-homogeneity of H . $\square$

Lemma 3.1 reduces our problem to computing the concave and convex envelopes of φ_p on Ω . By concave envelope we mean the infimum of linear functions on Ω that are greater than φ_p on $\partial\Omega$, and by convex envelope we mean the supremum of linear functions on Ω that are smaller than φ_p on $\partial\Omega$. Let $\bar{H}_p$ denote the concave envelope, and $\underline{H}_p$ the convex envelope. For $(x, y, z) \in \Omega$, define

$$w(x, y, z) := \frac{2z}{x + y}, \quad v(x, y, z) := \min\left\{\frac{x}{z}, \frac{y}{z}, 1\right\},$$

where we take $w = 0$ at the origin and $v = 1$ on $\Omega \cap \{z = 0\}$. Define the one-homogeneous functions F_p, G_p on Ω by

$$F_p(x, y, z) := \frac{x + y}{2} \left((1 + \sqrt{1 - w^2})^{1/p} + (1 - \sqrt{1 - w^2})^{1/p} \right)^p, \quad (11)$$

$$G_p(x, y, z) := \begin{cases} x + y + ((v^{1/p} + v^{-1/p})^p - (v + v^{-1}))z, & p > 0, \\ (v^{1/p} + v^{-1/p})^p z, & p < 0. \end{cases} \quad (12)$$

Proposition 3.2. *The concave and convex envelopes $\bar{H}_p, \underline{H}_p$ of φ_p in Ω are in $C(\Omega)$ and are given explicitly by the formulae*

$$\bar{H}_p = \begin{cases} F_p, & p \in (0, 1] \cup [2, \infty), \\ G_p, & p \in (-\infty, 0) \cup (1, 2) \end{cases}$$

and

$$\underline{H}_p = \begin{cases} F_p, & p \in (-\infty, 0) \cup (1, 2), \\ G_p, & p \in (0, 1] \cup [2, \infty). \end{cases}$$

We delay the proof of [Proposition 3.2](#) to [Section 4](#), and immediately note that [Theorems 2.1](#) and [2.5](#) follow quickly:

Proof of Theorems 2.1 and 2.5. To prove the inequalities, just apply [Lemma 3.1](#) to the functions $\bar{H}_p$ and $\underline{H}_p$. To check the equality cases, observe that in the proof of [Lemma 3.1](#), we have equality in Jensen provided $\{(f^p, g^p, (fg)^{p/2})\}$ lie in a set where H is linear.

Since F_p is linear when restricted to the hyperplanes $\{z = k(x + y)\} \cap \Omega$, which are nontrivial when $k \in [0, \frac{1}{2}]$, we obtain the equality case in [Theorem 2.1](#).

We note that G_p is linear on the triangular cone $\{z \leq \min\{x, y\}\} \cap \Omega$, and on the hyperplanes $\{z = \gamma x\} \cap \Omega$ and $\{z = \gamma y\} \cap \Omega$ for each $\gamma \geq 1$. The first condition gives $(fg)^{p/2} \leq \min\{f^p, g^p\}$, so $f = g$ on $\{fg > 0\}$ in the case $p > 0$ and on $\{fg < \infty\}$ in the case $p < 0$. The second condition gives $(fg)^{p/2} = \gamma f^p$, and the third $(fg)^{p/2} = \gamma g^p$. When $p > 0$, the second condition gives that $g = \lambda f$ on $\{f > 0\}$ for some $\lambda \geq 1$, and the third gives that $f = \lambda g$ on $\{g > 0\}$ for some $\lambda \geq 1$; when $p < 0$ the second condition gives $g = \lambda f$ on $\{f < \infty\}$ for some $\lambda \leq 1$, and the third gives that $f = \lambda g$ on $\{g < \infty\}$ for some $\lambda \leq 1$. $\square$

To conclude the section we address the optimality of [Theorems 2.1](#) and [2.5](#) in the measure space $(X, \mathcal{A}, \mu) = ([0, 1], \mathcal{B}, dx)$. We define

$$\begin{aligned}\bar{B}_p(x, y, z) &= \sup\{\|f + g\|_p^p : (\|f\|_p^p, \|g\|_p^p, \|fg\|_{p/2}^{p/2}) = (x, y, z)\}, \\ \underline{B}_p(x, y, z) &= \inf\{\|f + g\|_p^p : (\|f\|_p^p, \|g\|_p^p, \|fg\|_{p/2}^{p/2}) = (x, y, z)\}.\end{aligned}$$

It is easy to see that $\bar{B}_p, \underline{B}_p$ are defined on a cone $\Omega_p \subset \Omega$, are locally bounded by the inequalities $(f + g)^p \leq 2^{p-1}(f^p + g^p)$ for $p \in (-\infty, 0) \cup [1, \infty)$ and $(f + g)^p \leq f^p + g^p$ for $p \in (0, 1]$, are one-homogeneous, and equal φ_p on $\partial\Omega$ (by the equality case of Cauchy–Schwarz). Furthermore, by [Lemma 3.1](#) we have

$$\underline{H}_p \leq \underline{B}_p \leq \bar{B}_p \leq \bar{H}_p$$

on the common domain of definition.

Lemma 3.3. *If $\bar{B}_p$ ($\underline{B}_p$) is defined on all of Ω and is concave (convex), then*

$$\bar{H}_p = \bar{B}_p \quad (\underline{B}_p = \underline{H}_p).$$

Proof. Local boundedness and concavity of $\bar{B}_p$ implies continuity in the interior of Ω , and since $\bar{B}_p$ is trapped between envelopes that attain the data continuously, we have $\bar{B}_p \in C(\Omega)$. Since $\bar{H}_p$ is the smallest such concave function, we conclude that $\bar{B}_p \geq \bar{H}_p$. The argument is similar for $\underline{B}_p$. $\square$

Thus, it just remains to show that when $(X, \mathcal{A}, \mu) = ([0, 1], \mathcal{B}, dx)$, the domain of definition for $\bar{B}_p$ and $\underline{B}_p$ is all of Ω , and that $\bar{B}_p$ is concave and $\underline{B}_p$ is convex.

Lemma 3.4. *For $(X, \mathcal{A}, \mu) = ([0, 1], \mathcal{B}, dx)$ we have $\Omega_p = \Omega$ for all $p \neq 0$, that $\bar{B}_p$ is concave in Ω , and that $\underline{B}_p$ is convex in Ω .*

The optimality of the inequalities in [Theorems 2.1](#) and [2.5](#) follows:

Proof of optimality statements. For either inequality, given

$$(x, y, z) = (\|f\|_p^p, \|g\|_p^p, \|fg\|_{p/2}^{p/2}),$$

the functions $\bar{B}_p(x, y, z)$ and $\underline{B}_p(x, y, z)$ are by definition the best we can do. These are equal to the envelopes $\bar{H}_p, \underline{H}_p$ by Lemmas 3.3 and 3.4. $\square$

Remark 3.5. For given $(x, y, z) \in \Omega$ and $p \in \mathbb{R} \setminus \{0\}$, the supremum (infimum) in the definition of $\bar{B}_p$ ($\underline{B}_p$) is in fact attained.

For equality in (3) consider pairs of the form $(f, g) = (a, b)\chi_{[0,c]} + (b, a)\chi_{[c,1]}$ for a, b, c chosen appropriately.

For equality in (9), consider pairs of the form

$$(f, g) = (a, a)\chi_{[0,1/2]} + (b, 0)\chi_{[1/2,3/4]} + (0, c)\chi_{[3/4,1]}$$

for a, b, c appropriately chosen when $z \leq \min\{x, y\}$, and $(f, g) = (a, b)\chi_{[0,1/2]} + (c, d)\chi_{[1/2,1]}$ when $z > \min\{x, y\}$ for appropriate a, b, c, d , with one of c, d equal to 0.

For equality in (10), consider pairs of the form

$$(f, g) = (a, a)\chi_{[0,1/2]} + (b, \infty)\chi_{[1/2,3/4]} + (\infty, c)\chi_{[3/4,1]}$$

for a, b, c appropriately chosen when $z \leq \min\{x, y\}$, and $(f, g) = (a, b)\chi_{[0,1/2]} + (c, d)\chi_{[1/2,1]}$ when $z > \min\{x, y\}$ for appropriate a, b, c, d , with one of c, d equal to ∞ .

Proof of Lemma 3.4. For the first part, if $p > 0$ take $f_s = (2x)^{1/p}\chi_{[s,s+1/2]}$ for $s \in [0, \frac{1}{2}]$ and let $g = (2y)^{1/p}\chi_{[1/2,1]}$. Then $\|f_s\|_p^p = x$ and $\|g\|_p^p = y$. Furthermore, we have $h(s) := \|f_s g\|_{p/2}^{p/2}$ is continuous, increasing, and $h(0) = 0$, $h(\frac{1}{2}) = \sqrt{xy}$. When $p < 0$, use the same example but set $f_s, g = \infty$ where they were previously zero.

For the second part, let $(x_i, y_i, z_i) \in \Omega$ with $i = 1, 2$, and for $\epsilon > 0$ choose f_i, g_i such that $(x_i, y_i, z_i) = (\|f_i\|_p^p, \|g_i\|_p^p, \|f_i g_i\|_{p/2}^{p/2})$ and

$$\|f_i + g_i\|_p^p \geq \bar{B}_p(x_i, y_i, z_i) - \epsilon, \quad i = 1, 2.$$

Extend f_i, g_i to be zero outside of $[0, 1]$, and define the rescalings

$$\tilde{f}_1(s) = 2^{1/p} f_1(2s), \quad \tilde{g}_1(s) = 2^{1/p} g_1(2s), \quad \tilde{f}_2(s) = 2^{1/p} f_2(2s - 1), \quad \tilde{g}_2(s) = 2^{1/p} g_2(2s - 1),$$

so that $\tilde{f}_i, \tilde{g}_i$ are supported in $[0, \frac{1}{2}]$ for $i = 1$ and in $[\frac{1}{2}, 1]$ for $i = 2$. We then have

$$\begin{aligned} \frac{1}{2}(\bar{B}_p(x_1, y_1, z_1) + \bar{B}_p(x_2, y_2, z_2)) - \epsilon &\leq \frac{1}{2}(\|\tilde{f}_1 + \tilde{g}_1\|_{L^p([0,1/2])}^p + \|\tilde{f}_2 + \tilde{g}_2\|_{L^p([1/2,1])}^p) \\ &= \frac{1}{2}\|\tilde{f}_1 + \tilde{g}_1 + \tilde{f}_2 + \tilde{g}_2\|_p^p \\ &= \left\| \frac{\tilde{f}_1 + \tilde{f}_2}{2^{1/p}} + \frac{\tilde{g}_1 + \tilde{g}_2}{2^{1/p}} \right\|_p^p \\ &\leq \bar{B}_p\left(\frac{1}{2}(x_1 + x_2, y_1 + y_2, z_1 + z_2)\right). \end{aligned}$$

For the last inequality, we used that for $f_0 := 2^{-1/p}(\tilde{f}_1 + \tilde{f}_2)$, $g_0 := 2^{-1/p}(\tilde{g}_1 + \tilde{g}_2)$ we have

$$\|f_0\|_p^p = \frac{1}{2}(x_1 + x_2), \quad \|g_0\|_p^p = \frac{1}{2}(y_1 + y_2), \quad \|f_0 g_0\|_{p/2}^{p/2} = \frac{1}{2}(z_1 + z_2).$$

Taking $\epsilon \rightarrow 0$, we conclude that $\bar{B}_p$ is concave. The convex direction is similar. $\square$

Remark 3.6. Lemma 3.4 holds for any measure space with translation and scaling properties similar to $([0, 1], \mathcal{B}, dx)$, e.g., $(B_1 \subset \mathbb{R}^n, \mathcal{B}, dx)$.

Remark 3.7. The fact that $\bar{B}_p$ is concave also follows from Theorem 1 in [Ivanisvili 2018]. Since the argument is simple, we decided to include it for the reader's convenience.

4. Envelopes

In this section we prove Proposition 3.2. We begin with some simple observations.

First, to check concavity (convexity) in Ω and continuity up to $\partial\Omega$ of $\bar{H}_p(\underline{H}_p)$, by one-homogeneity it suffices to check these properties on the half-ellipse

$$D := \Omega \cap \{x + y = 2\}.$$

More generally, any one-homogeneous function B in a convex cone in $\mathbb{R}^n$ (say contained in $\{x_n > 0\}$) is concave (convex) if it is concave (convex) when restricted to a cross-section of the cone (say $\{x_n = 1\}$). Indeed, by one-homogeneity we have

$$B\left(\frac{x+y}{2}\right) = \frac{x_n+y_n}{2} B\left(\lambda \frac{x}{x_n} + (1-\lambda) \frac{y}{y_n}\right)$$

where $\lambda = x_n/(x_n + y_n)$, and the statement follows by applying concavity/convexity of B on the cross-section and then using one-homogeneity once more.

Second, to prove that $\bar{H}_p(\underline{H}_p)$ is the concave (convex) envelope of φ_p , it suffices to check that each point in the interior of D lies on a segment that connects boundary points of D , on which $\bar{H}_p(\underline{H}_p)$ is linear. Indeed, then any linear function larger (smaller) than φ_p on $\partial\Omega$ will then be larger than $\bar{H}_p$ (smaller than $\underline{H}_p$) in the interior of Ω .

Proof of Proposition 3.2. We first examine F_p , and then G_p .

The function F_p : On D we can write $F_p(1+s, 1-s, t) = u(t)$, where

$$u(t) := [(1 + \sqrt{1-t^2})^{1/p} + (1 - \sqrt{1-t^2})^{1/p}]^p, \quad t \in [0, 1].$$

It is clear that F_p is continuous up to ∂D for each $p \in \mathbb{R} \setminus \{0\}$, and $u(0) = \varphi_p$ (that is, 2 if $p > 0$ and 0 if $p < 0$) on the bottom of D and

$$F_p(1-s, 1+s, \sqrt{1-s^2}) = ((1+s)^{1/p} + (1-s)^{1/p})^p = \varphi_p$$

on the top of D . Since F_p is constant along the horizontal segments in D , it suffices to check that u is concave when $p \in (0, 1] \cup [2, \infty)$, and convex otherwise. To that end, we let $t = \sin(x)$, with $x \in [0, \frac{\pi}{2}]$.

Then

$$u(\sin(x)) = [(1 + \cos(x))^{1/p} + (1 - \cos(x))^{1/p}]^p.$$

Let us rewrite the last equality as

$$\frac{1}{2}u(\sin(2s)) = [\sin^{2/p}(s) + \cos^{2/p}(s)]^p,$$

where $s = \frac{x}{2} \in [0, \frac{\pi}{4}]$. Differentiating both sides of the equality in s , we obtain

$$\begin{aligned} u'(\sin(2s)) \cos(2s) &= p[\sin^{2/p}(s) + \cos^{2/p}(s)]^{p-1} \frac{2}{p} (\sin^{2/p-1}(s) \cos(s) - \cos^{2/p-1}(s) \sin(s)) \\ &= p[\sin^{2/p}(s) + \cos^{2/p}(s)]^{p-1} \frac{2 \cos^{2/p}(s)}{p} (\tan^{2/p-1}(s) - \tan(s)). \end{aligned}$$

Taking the derivative a second time we obtain

$$\begin{aligned} 2u''(\sin(2s)) \cos^2(2s) - 2u'(\sin(2s)) \sin(2s) \\ &= p(p-1)[\sin^{2/p}(s) + \cos^{2/p}(s)]^{p-2} \left[\frac{2 \cos^{2/p}(s)}{p} (\tan^{2/p-1}(s) - \tan(s)) \right]^2 + p[\sin^{2/p}(s) + \cos^{2/p}(s)]^{p-1} \\ &\quad \times \left(-\frac{4 \cos^{2/p}(s) \tan(s)}{p^2} (\tan^{2/p-1}(s) - \tan(s)) + \frac{2 \cos^{2/p}(s)}{p} \left(\left(\frac{2}{p} - 1 \right) \tan^{2/p-2}(s) - 1 \right) (1 + \tan^2(s)) \right). \end{aligned}$$

Therefore

$$\begin{aligned} 2u''(\sin(2s)) \cos^2(2s) \\ &= [\sin^{2/p}(s) + \cos^{2/p}(s)]^{p-2} \frac{4}{p} \cos^{4/p}(s) \\ &\quad \times \left[(p-1)[(\tan^{2/p-1}(s) - \tan(s))]^2 \right. \\ &\quad \left. + [1 + \tan^{2/p}(s)] \left(-\tan(s)(\tan^{2/p-1}(s) - \tan(s)) + \left(\left(1 - \frac{p}{2} \right) \tan^{2/p-2}(s) - \frac{p}{2} \right) (1 + \tan^2(s)) \right) \right. \\ &\quad \left. + p \tan(2s)[1 + \tan^{2/p}(s)](\tan^{2/p-1}(s) - \tan(s)) \right]. \end{aligned}$$

Since $\tan(2s) = 2 \tan(s)/(1 - \tan^2(s))$, after setting $\tan(s) = w \in [0, 1]$ we obtain

$$\begin{aligned} \frac{2u''(\sin(2s)) \cos^2(2s)}{[\sin^{2/p}(s) + \cos^{2/p}(s)]^{p-2} \cos^{4/p}(s)} &= \frac{4(p-1)}{p} (w^{2/p-1} - w)^2 \\ &\quad + \frac{4(1+w^{2/p})}{p} \left(-w^{2/p} + w^2 + \left(\left(1 - \frac{p}{2} \right) w^{2/p-2} - \frac{p}{2} \right) (1 + w^2) \right) \\ &\quad + \frac{8w}{1-w^2} (1 + w^{2/p})(w^{2/p-1} - w) \\ &= \frac{2(1+w^2)^2}{1-w^2} \left(w^{4/p-2} + \left(\frac{2}{p} - 1 \right) w^{2/p-2} (1 - w^2) - 1 \right). \end{aligned}$$

(The last equality is a tedious computation, but can be checked by hand). Since

$$\frac{2(1+w^2)^2}{1-w^2} > 0,$$

we see after defining $x := w^2 \in [0, 1]$ that $\operatorname{sgn}(u'') = \operatorname{sgn}(v(x))$, where

$$v(x) = x^{2/p-1} + \left(\frac{2}{p} - 1\right)x^{1/p-1}(1-x) - 1, \quad x \in [0, 1].$$

Let us study the sign of $v(x)$. Without loss of generality assume that $p \neq 1, 2$, otherwise the claims about concavity/convexity of u are trivial. First notice that $v(1) = 0$, and

$$v'(x) = x^{1/p-2} \left(\frac{2}{p} - 1\right) \left(x^{1/p} - \left(1 + \frac{1}{p}(x-1)\right)\right).$$

Therefore, if $p \in (2, \infty)$ it follows from concavity of $x \mapsto x^{1/p}$ that $v' \geq 0$, and hence $v \leq 0$; i.e., u is concave. Similarly, if $p \in (1, 2)$, then $v \geq 0$; i.e., u is convex. Next, if $p \in (0, 1)$ then $x \mapsto x^{1/p}$ is convex, and hence $v' \geq 0$, i.e., u is concave. Finally, if $p \in (-\infty, 0)$ then $x \mapsto x^{1/p}$ is convex, and therefore $v' \leq 0$; i.e., u is convex.

The function G_p : Let $b_p(s, z) = G_p(1+s, 1-s, z)$, with (s, z) in the upper half-disc. For $p > 0$ we can write b_p explicitly as

$$b_p(s, z) = 2 + \begin{cases} w(1-|s|, z), & z \geq 1-|s|, \\ (2^p-2)z, & z < 1-|s|, \end{cases}$$

where w is the one-homogeneous function given by

$$w(t, z) := \left(t^{1/p} + \left(\frac{z^2}{t}\right)^{1/p}\right)^p - \left(t + \frac{z^2}{t}\right),$$

with $(t, z) \in (0, 1)^2$. It is easy to check that b_p continuously takes the boundary values

$$\begin{aligned} b_p(s, 0) &= 2 = \varphi_p, \\ b_p(s, \sqrt{1-s^2}) &= ((1+s)^{1/p} + (1-s)^{1/p})^p = \varphi_p. \end{aligned}$$

Let

$$h(t) := w(t, 1) = (t^{1/p} + t^{-1/p})^p - (t + t^{-1}), \quad t \in (0, 1).$$

By the one-homogeneity of w and the fact that b_p is linear on the triangle $\{z < 1-|s|\}$ with vertical gradient, if we show that $h'(1) = 0$ and that h is concave/convex on $[0, 1]$, then b_p is C^1 away from $(s, z) = (\pm 1, 0)$ and concave/convex. Furthermore, b_p is linear when restricted to the segments through $(s, z) = (\pm 1, 0)$ that lie outside of the triangle $\{z \leq 1-|s|\}$, so G_p is the concave/convex envelope provided the above conditions on h are confirmed. To that end we compute the first two derivatives of h . The first derivative is

$$h'(t) = (t^{1/p} + t^{-1/p})^{p-1}(t^{1/p-1} - t^{-1/p-1}) - (1 - t^{-2}).$$

This confirms that $h'(1) = 0$. The second derivative is

$$\begin{aligned}
 h''(t) &= \frac{p-1}{p} (t^{1/p} + t^{-1/p})^{p-2} (t^{1/p-1} - t^{-1/p-1})^2 \\
 &\quad + \frac{1}{p} (t^{1/p} + t^{-1/p})^{p-1} ((1-p)t^{1/p-2} + (1+p)t^{-1/p-2}) - 2t^{-3} \\
 &= \frac{1}{p} (t^{1/p} + t^{-1/p})^{p-2} [(p-1)(t^{1/p-1} - t^{-1/p-1})^2 \\
 &\quad + (t^{1/p} + t^{-1/p})((1-p)t^{1/p-2} + (1+p)t^{-1/p-2})] - 2t^{-3} \\
 &= \frac{2}{p} (t^{1/p} + t^{-1/p})^{p-2} [pt^{-2/p-2} + (2-p)t^{-2}] - 2t^{-3} \\
 &= 2t^{-3} [(t^{1/p} + t^{-1/p})^{p-2} (t^{1-2/p} + (2/p-1)t) - 1] \\
 &= 2t^{-3} [(1+t^{2/p})^{p-2} (1 + (2/p-1)t^{2/p}) - 1].
 \end{aligned}$$

Let $x := t^{2/p} \in [0, 1]$. It suffices to show that

$$g_p(x) := \left(1 + \left(\frac{2}{p} - 1\right)x\right) - (1+x)^{2-p}$$

satisfies $g_p \leq 0$ on $[0, 1]$ for $p \in (1, 2)$ and $g_p \geq 0$ on $[0, 1]$ for $p \in (0, 1] \cup [2, \infty)$. Note that $g_p(0) = 0$. The desired inequality for $g_p(1)$ is equivalent to the fact that the linear function p crosses the convex function 2^{p-1} at $p = 1$ and $p = 2$. Finally, we observe that the first term in g_p is linear, and the second term is convex for $p \in (1, 2)$ and concave for $p \in (0, 1) \cup (2, \infty)$. The desired inequality for $g_p(x)$ with $x \in (0, 1)$ follows immediately from this observation and the inequalities at the endpoints $x = 0$ and $x = 1$.

When $p < 0$ we can write b_p explicitly as

$$b_p(s, z) = \begin{cases} \tilde{w}(1 - |s|, z), & z \geq 1 - |s|, \\ 2^p z, & z < 1 - |s|, \end{cases}$$

where $\tilde{w}$ is the one-homogeneous function given by

$$\tilde{w}(t, z) := (t^{1/p} + (z^2/t)^{1/p})^p$$

with $(t, z) \in (0, 1)^2$. The same considerations as above reduce the problem to showing that

$$\tilde{h}(t) := \tilde{w}(t, 1) = (t^{1/p} + t^{-1/p})^p$$

satisfies $\tilde{h}'(1) = 0$ and $\tilde{h}$ is concave on $[0, 1]$. We have

$$\begin{aligned}
 \tilde{h}' &= (t^{1/p} + t^{-1/p})^{p-1} (t^{1/p-1} - t^{-1/p-1}) \implies \tilde{h}'(1) = 0, \\
 \tilde{h}'' &= 2t^{-2} (t^{1/p} + t^{-1/p})^{p-2} [t^{-2/p} + (2/p-1)],
 \end{aligned}$$

and the conclusion follows quickly using $p < 0$. □

Remark 4.1. It follows from the concavity/convexity properties of G_p that

$$G_p(x, y, z) \leq x + y + (2^p - 2)z$$

when $p \in [1, 2]$, and the inequality reverses for $p \in (0, 1] \cup [2, \infty)$. Indeed, G_p agrees with the linear function on the right-hand side on an open set. We conclude from [Theorem 2.5](#) that for any nonnegative numbers a, b , and any $p \in [1, 2]$, we have

$$(a + b)^p \leq a^p + b^p + (2^p - 2)(ab)^{p/2},$$

and the inequality reverses if $p \in (0, 1] \cup [2, \infty)$.

5. Proof of [Corollary 2.6](#)

In this final section we prove [Corollary 2.6](#).

Proof of [Corollary 2.6](#). Recall from [Remark 4.1](#) that for any nonnegative numbers a, b , and any $p \in [1, 2]$, we have

$$(a + b)^p \leq a^p + b^p + (2^p - 2)(ab)^{p/2},$$

and the inequality reverses for $p \in (0, 1] \cup [2, \infty)$. Since for $p \in [0, 2]$ we have $(a + b)^{p/2} \leq a^{p/2} + b^{p/2}$, and the reverse inequality holds if $p \geq 2$, it follows by induction that for any nonnegative numbers $a_j \geq 0$ we have

$$\left(\sum_j a_j \right)^p \leq \sum_j a_j^p + (2^p - 2) \sum_{i < j} (a_i a_j)^{p/2} \quad (13)$$

for $p \in [1, 2]$, and the reverse inequality holds if $p \in (0, 1] \cup [2, \infty)$. Finally it remains to put $a_j = f_j(x)$ and integrate the inequality. $\square$

Remark 5.1. When $p < 0$, inequality (13) does not hold with three or more a_j . Take, e.g., $a_j = 1$ for $j \leq 3$.

6. Concluding remarks on envelopes

An important challenge in this work was to compute the envelopes (11) and (12). In this section we briefly explain how we found them.

We recall from [Section 3](#) that for the measure space $([0, 1], \mathcal{B}, dx)$ we have $\bar{B}_p = \bar{H}_p$ is defined on Ω , one-homogeneous, and equal to φ_p on $\partial\Omega$; that is, $\bar{H}_p(x, y, \sqrt{xy}) = (x^{1/p} + y^{1/p})^p$. We also recall from the discussion at the beginning of [Section 4](#) that by one-homogeneity, to compute $\bar{H}_p$ it is enough to restrict our attention to the cross-section $D = \Omega \cap \{x + y = 2\}$. Writing $D = \{(1 + s, 1 - s, z)\}$ with (s, z) in the upper half-disc, this reduces the problem to understanding how the upper boundary of the convex envelope of the space curve

$$\gamma(s) = (s, \sqrt{1 - s^2}, ((1 - s)^{1/p} + (1 + s)^{1/p})^p), \quad s \in [-1, 1],$$

looks. One can show that the torsion τ_γ of the space curve γ changes sign only once from $-$ to $+$, at $s = 0$, when $p \in (0, 1) \cup (2, \infty)$, and from $+$ to $-$ when $p \in (-\infty, 0) \cup (1, 2)$. Consider the case $p \in (0, 1) \cup (2, \infty)$. Then it follows from Lemma 29 of Section 3.2 in [\[Ivanisvili 2015\]](#) that locally, say

for some $\delta \in (0, 1]$, there exists a function $a(s) : [0, \delta] \rightarrow [-1, 0]$ such that $a(0) = 0$, $a(s)$ is strictly decreasing, and the function $B(u, w)$, defined parametrically by

$$B(\lambda(a(s), \sqrt{1-a(s)^2}) + (1-\lambda)(s, \sqrt{1-s^2})) \\ = \lambda((1-a(s))^{1/p} + (1+a(s))^{1/p})^p + (1-\lambda)((1-s)^{1/p} + (1+s)^{1/p})^p$$

for $\lambda \in [0, 1]$, $s \in [0, \delta]$, is concave. In other words B has the prescribed boundary condition, i.e., $B(s, \sqrt{1-s^2}) = ((1-s)^{1/p} + (1+s)^{1/p})^p$, it is linear along the line segments

$$\ell(s) := [(a(s), \sqrt{1-a(s)^2}), (s, \sqrt{1-s^2})],$$

and B is concave. It follows that “locally” B is a concave envelope. Because of the symmetry in x and y of the boundary data φ_p , one can show that the line segments $\ell(s)$ must be horizontal; i.e., $a(s) = -s$, and in fact $\delta = 1$. This means that B is a global concave envelope

$$B(u, w) = ((1-\sqrt{1-w^2})^{1/p} + (1+\sqrt{1-w^2})^{1/p})^p$$

for all $|u| \leq 1$ and $0 \leq w \leq \sqrt{1-u^2}$. Now it remains to change variables back to recover the envelope (11).

The case $p \in (-\infty, 0) \cup (1, 2)$ is different because τ_γ changes sign from $+$ to $-$, and in this case an “angle” arises with vertex sitting around the point $s = 0$; see Section 3 in [Ivanisvili 2015].

Acknowledgement

We are grateful to the anonymous referee for helpful comments and suggestions. Ivanisvili was supported by NSF grants DMS-1856486 and CAREER-DMS-1945102. Mooney was supported by NSF grant DMS-1854788

References

- [Bonami 1970] A. Bonami, “Étude des coefficients de Fourier des fonctions de $L^p(G)$ ”, *Ann. Inst. Fourier (Grenoble)* **20**:2 (1970), 335–402. [MR](#) [Zbl](#)
- [Carbery 2009] A. Carbery, “Almost-orthogonality in the Schatten–von Neumann classes”, *J. Operator Theory* **62**:1 (2009), 151–158. [MR](#) [Zbl](#)
- [Carlen et al. 2020a] E. A. Carlen, R. L. Frank, P. Ivanisvili, and E. H. Lieb, “Inequalities for L^p -norms that sharpen the triangle inequality and complement Hanner’s Inequality”, *J. Geom. Anal.* (online publication May 2020).
- [Carlen et al. 2020b] E. A. Carlen, R. L. Frank, and E. H. Lieb, “Inequalities that sharpen the triangle inequality for sums of N functions in L^p ”, *Ark. Mat.* **58**:1 (2020), 57–69.
- [Ivanisvili 2015] P. Ivanisvili, “Inequality for Burkholder’s martingale transform”, *Anal. PDE* **8**:4 (2015), 765–806. [MR](#) [Zbl](#)
- [Ivanisvili 2018] P. Ivanisvili, “Bellman function approach to the sharp constants in uniform convexity”, *Adv. Calc. Var.* **11**:1 (2018), 89–93. [MR](#) [Zbl](#)

Received 11 Feb 2019. Revised 2 May 2019. Accepted 11 Jun 2019.

PAATA IVANISVILI: pivanisv@uci.edu

Department of Mathematics, University of California, Irvine, CA, United States

CONNOR MOONEY: mooneycr@math.uci.edu

Department of Mathematics, University of California, Irvine, CA, United States

Analysis & PDE

msp.org/apde

EDITORS

EDITOR-IN-CHIEF

Patrick Gérard

patrick.gerard@math.u-psud.fr

Université Paris Sud XI

Orsay, France

BOARD OF EDITORS

Massimiliano Berti	Scuola Intern. Sup. di Studi Avanzati, Italy berti@sissa.it	Gilles Pisier	Texas A&M University, and Paris 6 pisier@math.tamu.edu
Michael Christ	University of California, Berkeley, USA mchrist@math.berkeley.edu	Tristan Rivière	ETH, Switzerland riviere@math.ethz.ch
Charles Fefferman	Princeton University, USA cf@math.princeton.edu	Igor Rodnianski	Princeton University, USA irod@math.princeton.edu
Ursula Hamenstaedt	Universität Bonn, Germany ursula@math.uni-bonn.de	Yum-Tong Siu	Harvard University, USA siu@math.harvard.edu
Vadim Kaloshin	University of Maryland, USA vadim.kaloshin@gmail.com	Terence Tao	University of California, Los Angeles, USA tao@math.ucla.edu
Herbert Koch	Universität Bonn, Germany koch@math.uni-bonn.de	Michael E. Taylor	Univ. of North Carolina, Chapel Hill, USA met@math.unc.edu
Izabella Laba	University of British Columbia, Canada ilaba@math.ubc.ca	Gunther Uhlmann	University of Washington, USA gunther@math.washington.edu
Richard B. Melrose	Massachusetts Inst. of Tech., USA rhm@math.mit.edu	András Vasy	Stanford University, USA andras@math.stanford.edu
Frank Merle	Université de Cergy-Pontoise, France Frank.Merle@u-cergy.fr	Dan Virgil Voiculescu	University of California, Berkeley, USA dvv@math.berkeley.edu
William Minicozzi II	Johns Hopkins University, USA minicozz@math.jhu.edu	Steven Zelditch	Northwestern University, USA zelditch@math.northwestern.edu
Clément Mouhot	Cambridge University, UK c.mouhot@dpmms.cam.ac.uk	Maciej Zworski	University of California, Berkeley, USA zworski@math.berkeley.edu
Werner Müller	Universität Bonn, Germany mueller@math.uni-bonn.de		

PRODUCTION

production@msp.org

Silvio Levy, Scientific Editor

See inside back cover or msp.org/apde for submission instructions.

The subscription price for 2020 is US \$340/year for the electronic version, and \$550/year (+\$60, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues from the last three years and changes of subscriber address should be sent to MSP.

Analysis & PDE (ISSN 1948-206X electronic, 2157-5045 printed) at Mathematical Sciences Publishers, 798 Evans Hall #3840, c/o University of California, Berkeley, CA 94720-3840, is published continuously online. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices.

APDE peer review and production are managed by EditFlow® from MSP.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2020 Mathematical Sciences Publishers

ANALYSIS & PDE

Volume 13 No. 5 2020

Regularity results for generalized double phase functionals SUN-SIG BYUN and JEHAN OH	1269
Epsilon-regularity for p -harmonic maps at a free boundary on a sphere KATARZYNA MAZOWIECKA, RÉMY RODIAC and ARMIN SCHIKORRA	1301
Uniform Sobolev estimates for Schrödinger operators with scaling-critical potentials and applications HARUYA MIZUTANI	1333
When does a perturbed Moser–Trudinger inequality admit an extremal? PIERRE-DAMIEN THIZY	1371
Well-posedness of the hydrostatic Navier–Stokes equations DAVID GÉRARD-VARET, NADER MASMOUDI and VLAD VICOL	1417
Sharp variation-norm estimates for oscillatory integrals related to Carleson’s theorem SHAOMING GUO, JORIS ROOS and PO-LAM YUNG	1457
Federer’s characterization of sets of finite perimeter in metric spaces PANU LAHTI	1501
Spectral theory of pseudodifferential operators of degree 0 and an application to forced linear waves YVES COLIN DE VERDIÈRE	1521
Global existence for the derivative nonlinear Schrödinger equation with arbitrary spectral singularities ROBERT JENKINS, JIAQI LIU, PETER PERRY and CATHERINE SULEM	1539
Unconditional existence of conformally hyperbolic Yamabe flows MARIO B. SCHULZ	1579
Sharpening the triangle inequality: envelopes between L^2 and L^p spaces PAATA IVANISVILI and CONNOR MOONEY	1591