

ANALYSIS & PDE

Volume 13

No. 7

2020

MARCH T. BOEDIHARDJO

C^* -ALGEBRAS ISOMORPHICALLY REPRESENTABLE ON l^p



C^* -ALGEBRAS ISOMORPHICALLY REPRESENTABLE ON l^p

MARCH T. BOEDIHARDJO

Let $p \in (1, \infty) \setminus \{2\}$. We show that every homomorphism from a C^* -algebra \mathcal{A} into $B(l^p(J))$ satisfies a compactness property where J is any set. As a consequence, we show that a C^* -algebra \mathcal{A} is isomorphic to a subalgebra of $B(l^p(J))$, for some set J , if and only if \mathcal{A} is residually finite-dimensional.

1. Introduction

For $1 \leq p < \infty$ and a set J , let $l^p(J)$ be the space

$$\left\{ f : J \rightarrow \mathbb{C} : \sum_{j \in J} |f(j)|^p < \infty \right\}$$

with norm

$$\|f\| = \left(\sum_{j \in J} |f(j)|^p \right)^{\frac{1}{p}}.$$

Two Banach algebras \mathcal{A}_1 and \mathcal{A}_2 are *isomorphic* if there exist a bijective homomorphism $\phi : \mathcal{A}_1 \rightarrow \mathcal{A}_2$ and $C > 0$ such that

$$\frac{1}{C} \|a\| \leq \|\phi(a)\| \leq C \|a\|$$

for all $a \in \mathcal{A}_1$. The algebras \mathcal{A}_1 and \mathcal{A}_2 are *isometrically isomorphic* if, moreover, ϕ can be chosen so that $\|\phi(a)\| = \|a\|$ for all $a \in \mathcal{A}_1$.

Gardella and Thiel [2020] showed that for $p \in [1, \infty) \setminus \{2\}$, a C^* -algebra \mathcal{A} is isometrically isomorphic to a subalgebra of $B(l^p(J))$, for some set J , if and only if \mathcal{A} is commutative. So it is natural to consider the question of whether this result holds if we relax the condition of isometrically isomorphic to isomorphic. In this paper, we show that for $p \in (1, \infty) \setminus \{2\}$, a C^* -algebra \mathcal{A} is isomorphic to a subalgebra of $B(l^p(J))$, for some set J , if and only if \mathcal{A} is residually finite-dimensional (Corollary 2.2). We prove this by showing that every homomorphism from a C^* -algebra \mathcal{A} into $B(l^p(J))$ satisfies a compactness property (Theorem 2.1).

The proofs of the main results Theorem 2.1 and Corollary 2.2 in this paper are quite different from the proof of Gardella and Thiel's result. Lamperti's characterization [1958] of isometries on L^p , for $p \neq 2$, plays a crucial role in the proof of Gardella and Thiel's result, while uniform convexity of l^p , for $1 < p < \infty$, and an argument in probability that imitates the proof of Khintchine's inequality [Lindenstrauss and Tzafriri 1977, Theorem 2.b.3], for $p = 1$, are used in the proof of Theorem 2.1.

MSC2010: 46H20.

Keywords: l^p space, C^* -algebra.

2. Main results and proofs

Throughout this paper, the scalar field is \mathbb{C} . For algebras \mathcal{A}_1 and \mathcal{A}_2 , a *homomorphism* $\phi : \mathcal{A}_1 \rightarrow \mathcal{A}_2$ is a bounded linear map such that $\phi(a_1 a_2) = \phi(a_1)\phi(a_2)$ for all $a_1, a_2 \in \mathcal{A}$. For an element a of a C^* -algebra, $|a| = \sqrt{a^* a}$. The algebra of bounded linear operators on a Banach space \mathcal{X} is denoted by $B(\mathcal{X})$ and the dual of \mathcal{X} is denoted by \mathcal{X}^* . For $1 \leq p \leq \infty$, the l^p direct sum of Banach spaces \mathcal{X}_α , for $\alpha \in \Lambda$, is denoted by $(\bigoplus_{\alpha \in \Lambda} \mathcal{X}_\alpha)_{l^p}$. Two Banach spaces \mathcal{X}_1 and \mathcal{X}_2 are *isomorphic* if there is an invertible operator $S : \mathcal{X}_1 \rightarrow \mathcal{X}_2$. A C^* -algebra \mathcal{A} is *residually finite-dimensional* if for every $a \in \mathcal{A}$, there is a $*$ -representation ϕ of \mathcal{A} on a finite-dimensional space such that $\phi(a) \neq 0$.

Theorem 2.1. *Let $p \in (1, \infty) \setminus \{2\}$. Let J be a set. Let \mathcal{A} be a C^* -algebra. Let $\phi : \mathcal{A} \rightarrow B(l^p(J))$ be a homomorphism. Then*

- (i) *the norm closure of $\{\phi(a)x : a \in \mathcal{A}, \|a\| \leq 1\}$ in $l^p(J)$ is norm compact for every $x \in l^p(J)$, and*
- (ii) *$\mathcal{A}/\ker \phi$ is a residually finite-dimensional C^* -algebra.*

Corollary 2.2. *Let $p \in (1, \infty) \setminus \{2\}$. A C^* -algebra \mathcal{A} is isomorphic to a subalgebra of $B(l^p(J))$, for some set J , if and only if \mathcal{A} is residually finite-dimensional.*

[Theorem 2.1](#) and [Corollary 2.2](#) will be proved at the end of this section after a series of lemmas are proved. [Theorem 2.1](#) has an easier proof when ϕ is contractive. Indeed, if $\phi : \mathcal{A} \rightarrow B(l^p(J))$ is a contractive homomorphism, then the range of ϕ is in the algebra of diagonal operators on $l^p(J)$ by [[Blecher and Phillips 2019](#), Proposition 2.12] (or by [[Gardella and Thiel 2020](#), Lemma 5.2] when J is countable). Thus, $\{\phi(a)x : a \in \mathcal{A}, \|a\| \leq 1\}$ is norm relatively compact, for every $x \in l^p(J)$, and $\mathcal{A}/\ker \phi$ is commutative.

It is not known if [Theorem 2.1](#) and [Corollary 2.2](#) hold for $p = 1$. However, throughout their proofs, we use, in an essential way, the assumption that p is in the reflexive range. For example, in the proof of [Theorem 2.1\(i\)](#), we use the fact that every bounded sequence in $l^p(J)$ has a weakly convergent subsequence. In the proof of [Corollary 2.2](#), we use a classical result of Pełczyński that the l^p direct sum of finite-dimensional Hilbert spaces is isomorphic to $l^p(J)$ for some set J . This result of Pełczyński holds only when p is in the reflexive range.

The structure of the proof of [Theorem 2.1\(i\)](#) goes as follows: If the closure of $\{\phi(a)x_0 : a \in \mathcal{A}, \|a\| \leq 1\}$ is not compact for some $x_0 \in l^p(J)$, then we can find a bounded sequence in $(b_k)_{k \in \mathbb{N}}$ in \mathcal{A} such that $\phi(b_k)x_0 \rightarrow 0$ weakly, as $k \rightarrow \infty$, and $\inf_{k \in \mathbb{N}} \|\phi(b_k)x_0\| > 0$. Assume that $p > 2$. In [Lemma 2.5](#), we show that $\phi(b_k) \rightarrow 0$ weakly implies that $\omega(b_k^* b_k) \rightarrow 0$ for all positive linear functionals $\omega : \mathcal{A} \rightarrow \mathbb{C}$ of the form $\omega(a) = y_0^*(\phi(a)x_0)$. This is proved by considering $\sum_{k=1}^n \delta_k b_k$ for random $\delta_1, \dots, \delta_n$ in $\{-1, 1\}$ and by exploiting $p > 2$. [Lemma 2.9](#) says that when $y_0^* \in (l^p(J))^*$ is suitably chosen, $\omega(b_k^* b_k) \rightarrow 0$ implies that $\|\phi(b_k)x_0\| \rightarrow 0$, which contradicts $\inf_{k \in \mathbb{N}} \|\phi(b_k)x_0\| > 0$. This is proved by using the uniform convexity of $l^p(J)$.

[Theorem 2.1\(ii\)](#) follows from [Theorem 2.1\(i\)](#) by using a GNS-type construction and a classical result about compact unitary representations of groups on Hilbert spaces.

The following two lemmas are needed for the proof of [Lemma 2.5](#).

Lemma 2.3. *Let \mathcal{A} be a unital C^* -algebra. Let $a \in \mathcal{A}$. Then there exists a sequence $(c_n)_{n \in \mathbb{N}}$ in \mathcal{A} such that $\|c_n\| \leq 1$ for all $n \in \mathbb{N}$ and $|a| = \lim_{n \rightarrow \infty} c_n a$.*

Proof. Without loss of generality, we may assume that $\|a\| \leq 1$. For $n \in \mathbb{N}$, define $g_n \in C[0, 1]$ by

$$g_n(x) = \begin{cases} \frac{1}{\sqrt{x}}, & \frac{1}{n} \leq x \leq 1, \\ n\sqrt{nx}, & 0 \leq x \leq \frac{1}{n}. \end{cases}$$

Take $c_n = g_n(a^*a)a^*$. Then $c_n c_n^* = g_n(a^*a)a^* a g_n(a^*a)$. Note that

$$x g_n(x)^2 = \begin{cases} 1, & \frac{1}{n} \leq x \leq 1, \\ n^3 x^3, & 0 \leq x \leq \frac{1}{n}. \end{cases}$$

Thus, $0 \leq x g_n(x)^2 \leq 1$ for all $x \in [0, 1]$ and so $0 \leq c_n c_n^* \leq 1$. Hence $\|c_n\| \leq 1$.

We have

$$x g_n(x) = \begin{cases} \sqrt{x}, & \frac{1}{n} \leq x \leq 1, \\ n\sqrt{nx}^2, & 0 \leq x \leq \frac{1}{n}, \end{cases}$$

and so

$$|x g_n(x) - \sqrt{x}| \leq \frac{1}{\sqrt{n}} \quad \text{for all } x \in [0, 1].$$

Since $c_n a = g_n(a^*a)a^* a$, it follows that

$$\|c_n a - \sqrt{a^*a} a\| \leq \frac{1}{\sqrt{n}}.$$

Thus, the result follows. □

Lemma 2.4. *Let \mathcal{A} be a unital C^* -algebra. Let ω be a positive linear functional on \mathcal{A} . Let $a \in \mathcal{A}$. If $a \geq 0$ then*

$$\omega(a^2) \leq \omega(a)^{\frac{2}{3}} \omega(a^4)^{\frac{1}{3}}.$$

Proof. There exists a measure μ on $[0, \|a\|]$ such that

$$\omega(f(a)) = \int f(x) d\mu(x),$$

for all $f \in C[0, \|a\|]$. So

$$\omega(a^2) = \int x^2 d\mu(x) \leq \left(\int x d\mu(x) \right)^{\frac{2}{3}} \left(\int x^4 d\mu(x) \right)^{\frac{1}{3}} = \omega(a)^{\frac{2}{3}} \omega(a^4)^{\frac{1}{3}}. \quad \square$$

Lemma 2.5. *Let $2 < p < \infty$. Let J be a set. Let \mathcal{A} be a unital C^* -algebra. Let $\phi : \mathcal{A} \rightarrow B(l^p(J))$ be a unital homomorphism. Let $x_0 \in l^p(J)$. Let y_0^* be a bounded linear functional on $l^p(J)$. Define $\omega : \mathcal{A} \rightarrow \mathbb{C}$ by*

$$\omega(a) = y_0^*(\phi(a)x_0),$$

for $a \in \mathcal{A}$. Assume that ω is a positive linear functional. Let $(b_k)_{k \in \mathbb{N}}$ be a sequence in \mathcal{A} such that $\|b_k\| \leq 1$ for all $k \in \mathbb{N}$ and $\phi(b_k)x_0 \rightarrow 0$ weakly as $k \rightarrow \infty$. Then $\omega(b_k^ b_k) \rightarrow 0$ as $k \rightarrow \infty$.*

Proof. By contradiction, suppose that $\omega(b_k^*b_k)$ does not converge to 0. Passing to a subsequence, we have that there exists $\gamma > 0$ such that $\omega(b_k^*b_k) \geq \gamma$ for all $k \in \mathbb{N}$.

Since $\|\phi(b_k)x_0\| \leq \|\phi\| \|x_0\|$ and $\phi(b_k)x_0 \rightarrow 0$ weakly, passing to a further subsequence, we may assume that there are z_1, z_2, \dots in $l^p(J)$ with disjoint supports such that $\|z_k\| \leq \|\phi\| \|x_0\|$ and $\|\phi(b_k)x_0 - z_k\| \leq 1/2^k$ for all $k \in \mathbb{N}$.

Let $n \in \mathbb{N}$. For each $\delta = (\delta_1, \dots, \delta_n) \in \{-1, 1\}^n$, let

$$a_\delta = \left| \sum_{k=1}^n \delta_k b_k \right| \in \mathcal{A}.$$

By Lemma 2.4,

$$\omega(a_\delta^2) \leq \omega(a_\delta)^{\frac{2}{3}} \omega(a_\delta^4)^{\frac{1}{3}}.$$

Thus,

$$\mathbb{E}\omega(a_\delta^2) \leq [\mathbb{E}\omega(a_\delta)]^{\frac{2}{3}} [\mathbb{E}\omega(a_\delta^4)]^{\frac{1}{3}},$$

where \mathbb{E} denotes expectation over $\delta = (\delta_1, \dots, \delta_n)$ uniformly distributed on $\{-1, 1\}^n$.

Note that

$$\begin{aligned} \mathbb{E}\omega(a_\delta^2) &= \mathbb{E}\omega\left(\left(\sum_{k=1}^n \delta_k b_k\right)^* \left(\sum_{k=1}^n \delta_k b_k\right)\right) \\ &= \mathbb{E}\omega\left(\sum_{1 \leq j, k \leq n} \delta_j \delta_k b_j^* b_k\right) = \sum_{1 \leq j, k \leq n} \mathbb{E}(\delta_j \delta_k) \omega(b_j^* b_k) = \sum_{k=1}^n \omega(b_k^* b_k) \geq n\gamma. \end{aligned}$$

Therefore,

$$n\gamma \leq [\mathbb{E}\omega(a_\delta)]^{\frac{2}{3}} [\mathbb{E}\omega(a_\delta^4)]^{\frac{1}{3}}. \tag{2-1}$$

We have

$$a_\delta^4 = \left[\left(\sum_{k=1}^n \delta_k b_k\right)^* \left(\sum_{k=1}^n \delta_k b_k\right) \right]^2 = \sum_{1 \leq k_1, \dots, k_4 \leq n} \delta_{k_1} \delta_{k_2} \delta_{k_3} \delta_{k_4} b_{k_1}^* b_{k_2} b_{k_3}^* b_{k_4}.$$

Since $\|b_k\| \leq 1$, it follows that

$$\mathbb{E}\omega(a_\delta^4) = \sum_{1 \leq k_1, \dots, k_4 \leq n} \mathbb{E}(\delta_{k_1} \delta_{k_2} \delta_{k_3} \delta_{k_4}) \omega(b_{k_1}^* b_{k_2} b_{k_3}^* b_{k_4}) \leq \sum_{1 \leq k_1, \dots, k_4 \leq n} \mathbb{E}(\delta_{k_1} \delta_{k_2} \delta_{k_3} \delta_{k_4}).$$

Note that $\mathbb{E}(\delta_{k_1} \delta_{k_2} \delta_{k_3} \delta_{k_4}) = 0$ unless the following occurs:

$$(k_1 = k_2 \text{ and } k_3 = k_4) \quad \text{or} \quad (k_1 = k_3 \text{ and } k_2 = k_4) \quad \text{or} \quad (k_1 = k_4 \text{ and } k_2 = k_3).$$

Thus, $\mathbb{E}\omega(a_\delta^4) \leq 3n^2$. So by (2-1), we have $n\gamma \leq 3^{\frac{1}{3}} n^{\frac{2}{3}} [\mathbb{E}\omega(a_\delta)]^{\frac{2}{3}}$. Hence,

$$\mathbb{E}\omega(a_\delta) \geq \frac{\gamma^{\frac{3}{2}} n^{\frac{1}{2}}}{3^{\frac{1}{2}}}. \tag{2-2}$$

Fix $\delta \in \{-1, 1\}^n$. By Lemma 2.3,

$$\omega(a_\delta) = \omega\left(\left|\sum_{k=1}^n \delta_k b_k\right|\right) \leq \sup_{c \in \mathcal{A}, \|c\| \leq 1} \left| \omega\left(c \sum_{k=1}^n \delta_k b_k\right) \right|.$$

For $c \in \mathcal{A}$ with $\|c\| \leq 1$,

$$\begin{aligned} \left| \omega \left(c \sum_{k=1}^n \delta_k b_k \right) \right| &= \left| y_0^* \left(\phi(c) \left(\sum_{k=1}^n \delta_k \phi(b_k) x_0 \right) \right) \right| \\ &\leq \|y_0^*\| \|\phi\| \left\| \sum_{k=1}^n \delta_k \phi(b_k) x_0 \right\| \\ &\leq \|y_0^*\| \|\phi\| \left(\left\| \sum_{k=1}^n \delta_k z_k \right\| + \sum_{k=1}^n \frac{1}{2^k} \right) \leq \|y_0^*\| \|\phi\| (\|\phi\| \|x_0\| n^{\frac{1}{p}} + 1), \end{aligned}$$

where the last two inequalities follow from the fact that z_1, z_2, \dots have disjoint supports, $\|z_k\| \leq \|\phi\| \|x_0\|$ and $\|\phi(b_k)x_0 - z_k\| \leq 1/2^k$. Thus,

$$\omega(a_\delta) \leq \|y_0^*\| \|\phi\| (\|\phi\| \|x_0\| n^{\frac{1}{p}} + 1) \quad \text{for all } \delta \in \{-1, 1\}^n.$$

So by (2-2),

$$\frac{\gamma^{\frac{3}{2}}}{3^{\frac{1}{2}}} n^{\frac{1}{2}} \leq \|y_0^*\| \|\phi\| (\|\phi\| \|x_0\| n^{\frac{1}{p}} + 1).$$

Since n can be chosen to be arbitrarily large and $p > 2$, an absurdity follows. □

For $1 < p < 2$, we have the following result, where the order of b_k^* and b_k is switched, by using the dual l^p space in Lemma 2.5.

Lemma 2.6. *Let $1 < p < 2$. Let J be a set. Let \mathcal{A} be a unital C^* -algebra. Let $\phi : \mathcal{A} \rightarrow B(l^p(J))$ be a unital homomorphism. Let $x_0 \in l^p(J)$. Let y_0^* be a bounded linear functional on l^p . Define $\omega : \mathcal{A} \rightarrow \mathbb{C}$ by*

$$\omega(a) = y_0^*(\phi(a)x_0),$$

for $a \in \mathcal{A}$. Let $(b_k)_{k \in \mathbb{N}}$ be a sequence in \mathcal{A} such that $\|b_k\| \leq 1$ for all $k \in \mathbb{N}$ and such that the sequence $y_0^* \circ \phi(b_k)$ of bounded linear functionals on $l^p(J)$ converges to 0 weakly as $k \rightarrow \infty$. Assume that ω is a positive linear functional. Then $\omega(b_k b_k^*) \rightarrow 0$ as $k \rightarrow \infty$.

Proof. Let \mathcal{A}_1 be the unital C^* -algebra consisting of the same elements as \mathcal{A} but with reverse order multiplication

$$a \cdot b = ba.$$

Define a unital homomorphism $\phi_1 : \mathcal{A}_1 \rightarrow B((l^p(J))^*)$ by

$$\phi_1(a) y^* = y^* \circ \phi(a),$$

for all $a \in \mathcal{A}_1$, $y^* \in (l^p(J))^*$. Define $\omega_1 : \mathcal{A}_1 \rightarrow \mathbb{C}$ by

$$\omega_1(a) = \omega(a) = x_0^{**}(\phi(a)y_0^*),$$

for all $a \in \mathcal{A}_1$, where x_0^{**} is the image of x_0 in the bidual $(l^p)^{**}$. By Lemma 2.5, the result follows. □

The following two lemmas are needed for the proof of Lemma 2.9.

Lemma 2.7 [Clarkson 1936]. *Let $1 < p < \infty$. Let J be a set. For every $\epsilon > 0$, there exists $\gamma > 0$ such that, for all $x, y \in l^p(J)$ satisfying $\|x\|, \|y\| \leq 1$ and $\|x + y\| > 2 - \gamma$, we have $\|x - y\| < \epsilon$.*

Lemma 2.8 [Russo and Dye 1966]. *Let \mathcal{A} be a unital C^* -algebra. Then the closed unital ball of \mathcal{A} is the closed convex hull of the set of all unitary elements of \mathcal{A} .*

Lemma 2.9. *Let $1 < p < \infty$. Let J be a set. Let \mathcal{A} be a unital C^* -algebra. Let $\phi : \mathcal{A} \rightarrow B(l^p(J))$ be a unital homomorphism. Let $x_0 \in l^p(J)$. Then there exists $y_0^* \in (l^p(J))^*$ such that $\omega : \mathcal{A} \rightarrow \mathbb{C}$,*

$$\omega(a) = y_0^*(\phi(a)x_0), \quad a \in \mathcal{A},$$

*defines a positive linear functional and, for every $\epsilon > 0$, there exists $\gamma > 0$ such that whenever $a \in \mathcal{A}$ satisfies $\|a\| \leq 1$ and $\omega(a^*a) < \gamma$, we have $\|\phi(a)x_0\| < \epsilon$.*

Proof. Let $\mathcal{U}(\mathcal{A})$ be the set of all unitary elements of \mathcal{A} . Let $(v_n)_{n \in \mathbb{N}}$ be a sequence in $\mathcal{U}(\mathcal{A})$ such that

$$\lim_{n \rightarrow \infty} \|\phi(v_n)x_0\| = \sup_{u \in \mathcal{U}(\mathcal{A})} \|\phi(u)x_0\|.$$

For each $n \in \mathbb{N}$, let x_n^* be a bounded linear functional on $l^p(J)$ such that $\|x_n^*\| = 1$ and $x_n^*(\phi(v_n)x_0) = \|\phi(v_n)x_0\|$. Then $x_n^* \circ \phi(v_n)$ is a bounded sequence in $(l^p(J))^*$. Passing to a subsequence, we may assume that $x_n^* \circ \phi(v_n)$ converges weakly to a bounded linear functional $y_0^* \in (l^p(J))^*$ as $n \rightarrow \infty$. Thus, $\omega : \mathcal{A} \rightarrow \mathbb{C}$,

$$\omega(a) = y_0^*(\phi(a)x_0) = \lim_{n \rightarrow \infty} x_n^*(\phi(v_n a)x_0),$$

for $a \in \mathcal{A}$, defines a bounded linear functional on \mathcal{A} . Note that

$$\omega(1) = \lim_{n \rightarrow \infty} x_n^*(\phi(v_n)x_0) = \lim_{n \rightarrow \infty} \|\phi(v_n)x_0\| = \sup_{u \in \mathcal{U}(\mathcal{A})} \|\phi(u)x_0\|,$$

and, for every $u_0 \in \mathcal{U}(\mathcal{A})$,

$$|\omega(u_0)| = \lim_{n \rightarrow \infty} |x_n^*(\phi(v_n u_0)x_0)| \leq \sup_{u \in \mathcal{U}(\mathcal{A})} \|\phi(u)x_0\|.$$

So by Lemma 2.8, we have $\|\omega\| \leq \sup_{u \in \mathcal{U}(\mathcal{A})} \|\phi(u)x_0\|$. Thus, $\omega(1) = \|\omega\|$ and hence ω is a positive linear functional.

By contradiction, suppose that there are $\epsilon > 0$ and a sequence $(a_k)_{k \in \mathbb{N}}$ in \mathcal{A} such that $\|a_k\| \leq 1$ and $\|\phi(a_k)x_0\| \geq \epsilon$ for all $k \in \mathbb{N}$ and $\omega(a_k^*a_k) \rightarrow 0$ as $k \rightarrow \infty$. We have

$$\|a_k\| \geq \frac{\|\phi(a_k)x_0\|}{\|\phi\| \|x_0\|} \geq \frac{\epsilon}{\|\phi\| \|x_0\|},$$

for all $k \in \mathbb{N}$. For $k \in \mathbb{N}$, let $b_k = a_k/\|a_k\|$. We have $\|b_k\| = 1$ and $\|\phi(b_k)x_0\| \geq \epsilon$ for all $k \in \mathbb{N}$ and $\omega(b_k^*b_k) \rightarrow 0$ as $k \rightarrow \infty$.

Since $\|x_n^*\| = 1$,

$$\begin{aligned} \liminf_{n \rightarrow \infty} \|\phi(v_n)\phi(1 - |b_k|)x_0 + \phi(v_n)x_0\| &\geq \liminf_{n \rightarrow \infty} [x_n^*(\phi(v_n)\phi(1 - |b_k|)x_0) + x_n^*(\phi(v_n)x_0)] \\ &= \omega(1 - |b_k|) + \omega(1) = 2\omega(1) - \omega(|b_k|). \end{aligned}$$

Thus,

$$\liminf_{n \rightarrow \infty} \|\phi(v_n)\phi(1 - |b_k|)x_0 + \phi(v_n)x_0\| \geq 2\omega(1) - \omega(|b_k|).$$

But

$$\|\phi(v_n)\phi(1 - |b_k|)x_0\| \leq \sup_{b \in \mathcal{A}, \|b\| \leq 1} \|\phi(b)x_0\| \|1 - |b_k|\| \leq \sup_{u \in \mathcal{U}(\mathcal{A})} \|\phi(u)x_0\| = \omega(1)$$

and $\|\phi(v_n)x_0\| \leq \omega(1)$ for all $n \in \mathbb{N}$. Take

$$x = \frac{1}{\omega(1)}\phi(v_n)\phi(1 - |b_k|)x_0 \quad \text{and} \quad y = \frac{1}{\omega(1)}\phi(v_n)x_0$$

in Lemma 2.7 and note that $\omega(|b_k|) \leq \omega(b_k^*b_k)^{\frac{1}{2}}\omega(1)^{\frac{1}{2}} \rightarrow 0$ as $k \rightarrow \infty$. We have

$$\lim_{k \rightarrow \infty} \limsup_{n \rightarrow \infty} \|\phi(v_n)\phi(1 - |b_k|)x_0 - \phi(v_n)x_0\| = 0.$$

Thus,

$$\lim_{k \rightarrow \infty} \limsup_{n \rightarrow \infty} \|\phi(v_n)\phi(|b_k|)x_0\| = 0.$$

So $\|\phi(|b_k|)x_0\| \rightarrow 0$ as $k \rightarrow \infty$. Since

$$b_k = b_k \left(|b_k| + \frac{1}{k}\right)^{-1} \left(|b_k| + \frac{1}{k}\right) \quad \text{and} \quad \left\| b_k \left(|b_k| + \frac{1}{k}\right)^{-1} \right\| \leq 1,$$

it follows that $\|\phi(b_k)x_0\| \rightarrow 0$ as $k \rightarrow \infty$ which contradicts $\|\phi(b_k)x_0\| \geq \epsilon$. □

Proof of Theorem 2.1(i). Without loss generality, we may assume that \mathcal{A} is unital by extending ϕ to a homomorphism from the unitization of \mathcal{A} into $B(l^p(J))$. We may also assume that ϕ is unital since $\phi(1)$ is an idempotent on $l^p(J)$ and the range of every idempotent on $l^p(J)$ is isomorphic to $l^p(J_0)$ for some set J_0 [Pełczyński 1960; Johnson 2012].

Let $x_0 \in l^p$. Let $(a_k)_{k \in \mathbb{N}}$ be a sequence in \mathcal{A} such that $\|a_k\| \leq \frac{1}{2}$ for all $k \in \mathbb{N}$. We need to show that $(\phi(a_k)x_0)_{k \in \mathbb{N}}$ has a norm-convergent subsequence.

Case 1: $p > 2$. Passing to a subsequence, we may assume that $(\phi(a_k)x_0)_{k \in \mathbb{N}}$ converges weakly to an element of $l^p(J)$. Thus, $\phi(a_{k_1} - a_{k_2})x_0 \rightarrow 0$ weakly as $k_1, k_2 \rightarrow \infty$.

By Lemma 2.5, we have

$$\lim_{k_1, k_2 \rightarrow \infty} \omega((a_{k_1} - a_{k_2})^*(a_{k_1} - a_{k_2})) = 0$$

for every positive linear functional $\omega : \mathcal{A} \rightarrow \mathbb{C}$ of the form $\omega(a) = y_0^*(\phi(a)x_0)$ for $a \in \mathcal{A}$. By Lemma 2.9, we have $\lim_{k_1, k_2 \rightarrow \infty} \|\phi(a_{k_1} - a_{k_2})x_0\| = 0$. So $(\phi(a_k)x_0)_{k \in \mathbb{N}}$ is norm-convergent.

Case 2: $p < 2$. Passing to a subsequence, we may assume that $(y_0^* \circ \phi(a_k^*))_{k \in \mathbb{N}}$ converges weakly to an element of $(l^p(J))^*$. Thus, $y^* \circ \phi(a_{k_1}^* - a_{k_2}^*) \rightarrow 0$ weakly as $k_1, k_2 \rightarrow \infty$ for every $y^* \in (l^p(J))^*$.

By Lemma 2.6, we have

$$\lim_{k \rightarrow \infty} \omega((a_{k_1}^* - a_{k_2}^*)(a_{k_1}^* - a_{k_2}^*)^*) = 0$$

for every positive linear functional $\omega : \mathcal{A} \rightarrow \mathbb{C}$ of the form $\omega(a) = y_0^*(\phi(a)x_0)$ for $a \in \mathcal{A}$. By Lemma 2.9, we have $\lim_{k_1, k_2 \rightarrow \infty} \|\phi(a_{k_1} - a_{k_2})x_0\| = 0$. So $(\phi(a_k)x_0)_{k \in \mathbb{N}}$ is norm-convergent. □

Lemma 2.10 [Kerr and Li 2016, Theorem 2.24]. *Let G be a group. Let \mathcal{H} be a Hilbert space. Let $\psi : G \rightarrow B(\mathcal{H})$ be a unital homomorphism such that $\psi(g)$ is unitary for all $g \in G$. If $\{\psi(g)x : g \in G\}$ is norm precompact in \mathcal{H} for all $x \in \mathcal{H}$, then \mathcal{H} is the direct sum of some finite-dimensional subspaces \mathcal{H}_α , for $\alpha \in \Lambda$, such that \mathcal{H}_α is invariant under $\psi(g)$ for all $\alpha \in \Lambda$ and $g \in G$.*

Proof of Theorem 2.1(ii). As in the proof Theorem 2.1(i), we may assume that \mathcal{A} is unital and ϕ is unital. We may also assume that $\ker \phi = \{0\}$. Let $a_0 \neq 0$. There exists $x_0 \in l^p(J)$ such that $\phi(a_0)x_0 \neq 0$. By Lemma 2.9, there exists $y_0^* \in (l^p(J))^*$ such that $\omega : \mathcal{A} \rightarrow \mathbb{C}$,

$$\omega(a) = y_0^*(\phi(a)x_0),$$

for $a \in \mathcal{A}$, defines a positive linear functional and $\omega(a_0^*a_0) \neq 0$.

Equip \mathcal{A} with the positive semidefinite sesquilinear form

$$\langle a, b \rangle = \omega(b^*a),$$

for $a, b \in \mathcal{A}$. Consider the ideal $\mathcal{A}_0 = \{a \in \mathcal{A} : \langle a, a \rangle = 0\}$ of \mathcal{A} . Let \mathcal{H} be the completion of the quotient space $\mathcal{A}/\mathcal{A}_0$. Then \mathcal{H} is a Hilbert space. For each $a \in \mathcal{A}$, we can define a bounded linear operator on \mathcal{H} by sending $b + \mathcal{A}_0$ to $ab + \mathcal{A}_0$ for $b \in \mathcal{A}$. So $\eta : \mathcal{A} \rightarrow B(\mathcal{H})$,

$$\eta(a)(b + \mathcal{A}_0) = ab + \mathcal{A}_0,$$

for $a, b \in \mathcal{A}$, defines a unital $*$ -homomorphism. We have

$$\begin{aligned} \|\eta(a_1)(b + \mathcal{A}_0) - \eta(a_2)(b + \mathcal{A}_0)\| &= \omega(b^*(a_1 - a_2)^*(a_1 - a_2)b) \\ &= y_0^*(\phi(b^*(a_1 - a_2)^*(a_1 - a_2)b)x_0) \\ &\leq \|y_0^*\| \|\phi\| \|b^*\| \|a_1 - a_2\| \|\phi(a_1 - a_2)\phi(b)x_0\|, \end{aligned}$$

for all $a_1, a_2, b \in \mathcal{A}$. By Theorem 2.1(i), we have that $\{\phi(a)x_0 : a \in \mathcal{A}, \|a\| \leq 1\}$ is norm precompact so $\{\eta(a)(b + \mathcal{A}_0) : a \in \mathcal{A}, \|a\| \leq 1\}$ is norm precompact for all $b \in \mathcal{A}$. Let $\mathcal{U}(\mathcal{A})$ be the set of all unitary elements of \mathcal{A} . By Lemma 2.10, we have that \mathcal{H} is the direct sum of some finite-dimensional subspaces \mathcal{H}_α , for $\alpha \in \Lambda$, such that \mathcal{H}_α is invariant under $\eta(u)$ for all $\alpha \in \Lambda$ and $u \in \mathcal{U}(\mathcal{A})$. Note that \mathcal{H}_α is thus invariant under $\eta(a)$ for all $a \in \mathcal{A}$.

Since $\omega(a_0^*a_0) \neq 0$, we have $\eta(a_0) \neq 0$. So $\eta(a_0) \neq 0$ on \mathcal{H}_{α_0} for some $\alpha_0 \in \Lambda$. Thus, \mathcal{A} is residually finite-dimensional. \square

Proof of Corollary 2.2. One direction follows from Theorem 2.1. For the other direction, suppose that \mathcal{A} is a residually finite-dimensional C^* -algebra. Then there is a collection $(\phi_\alpha)_{\alpha \in \Lambda}$ of $*$ -representations of \mathcal{A} on finite-dimensional Hilbert spaces \mathcal{H}_α such that $\|a\| = \sup_{\alpha \in \Lambda} \|\phi_\alpha(a)\|$ for all $a \in \mathcal{A}$. Define $\phi : \mathcal{A} \rightarrow B((\bigoplus_{\alpha \in \Lambda} \mathcal{H}_\alpha)_{l^p})$ by $\phi = \bigoplus_{\alpha \in \Lambda} \phi_\alpha$. Thus ϕ is a norm-preserving homomorphism. However, it is a classical result of [Pełczyński 1960] that for $1 < p < \infty$, the l^p direct sum of finite-dimensional Hilbert spaces is isomorphic to $l^p(J)$ for some set J . Therefore, \mathcal{A} is isomorphic to a subalgebra of $B(l^p(J))$, via the map $a \mapsto S\phi(a)S^{-1}$, where $S : (\bigoplus_{\alpha \in \Lambda} \mathcal{H}_\alpha)_{l^p} \rightarrow l^p(J)$ is any invertible operator. \square

Acknowledgements

The author is grateful to the referee for some suggestions that improved the exposition. The author is supported by NSF DMS-1856221.

References

- [Blecher and Phillips 2019] D. P. Blecher and N. C. Phillips, “ L^p -operator algebras with approximate identities, I”, *Pacific J. Math.* **303**:2 (2019), 401–457. [MR](#) [Zbl](#)
- [Clarkson 1936] J. A. Clarkson, “Uniformly convex spaces”, *Trans. Amer. Math. Soc.* **40**:3 (1936), 396–414. [MR](#) [Zbl](#)
- [Gardella and Thiel 2020] E. Gardella and H. Thiel, “Extending representations of Banach algebras to their biduals”, *Math. Z.* **294**:3-4 (2020), 1341–1354. [MR](#) [Zbl](#)
- [Johnson 2012] B. Johnson, “Complemented subspaces of $\ell_p(I)$ for uncountable I ”, answer on MathOverflow, 2012, available at <https://mathoverflow.net/questions/111882/complemented-subspaces-of-ell-pi-for-uncountable-i>.
- [Kerr and Li 2016] D. Kerr and H. Li, *Ergodic theory: independence and dichotomies*, Springer, 2016. [MR](#) [Zbl](#)
- [Lamperti 1958] J. Lamperti, “On the isometries of certain function-spaces”, *Pacific J. Math.* **8** (1958), 459–466. [MR](#) [Zbl](#)
- [Lindenstrauss and Tzafriri 1977] J. Lindenstrauss and L. Tzafriri, *Classical Banach spaces*, vol. I: Sequence spaces, *Ergebnisse der Mathematik und ihrer Grenzgebiete* **92**, Springer, 1977. [MR](#) [Zbl](#)
- [Pełczyński 1960] A. Pełczyński, “Projections in certain Banach spaces”, *Studia Math.* **19** (1960), 209–228. [MR](#) [Zbl](#)
- [Russo and Dye 1966] B. Russo and H. A. Dye, “A note on unitary operators in C^* -algebras”, *Duke Math. J.* **33** (1966), 413–416. [MR](#) [Zbl](#)

Received 29 Dec 2018. Revised 19 Jun 2019. Accepted 6 Sep 2019.

MARCH T. BOEDIHARDJO: march@math.ucla.edu

Department of Mathematics, University of California, Los Angeles, CA, United States

Analysis & PDE

msp.org/apde

EDITORS

EDITOR-IN-CHIEF

Patrick Gérard

patrick.gerard@math.u-psud.fr

Université Paris Sud XI

Orsay, France

BOARD OF EDITORS

Massimiliano Berti	Scuola Intern. Sup. di Studi Avanzati, Italy berti@sissa.it	Gilles Pisier	Texas A&M University, and Paris 6 pisier@math.tamu.edu
Michael Christ	University of California, Berkeley, USA mchrist@math.berkeley.edu	Tristan Rivière	ETH, Switzerland riviere@math.ethz.ch
Charles Fefferman	Princeton University, USA cf@math.princeton.edu	Igor Rodnianski	Princeton University, USA irod@math.princeton.edu
Ursula Hamenstaedt	Universität Bonn, Germany ursula@math.uni-bonn.de	Yum-Tong Siu	Harvard University, USA siu@math.harvard.edu
Vadim Kaloshin	University of Maryland, USA vadim.kaloshin@gmail.com	Terence Tao	University of California, Los Angeles, USA tao@math.ucla.edu
Herbert Koch	Universität Bonn, Germany koch@math.uni-bonn.de	Michael E. Taylor	Univ. of North Carolina, Chapel Hill, USA met@math.unc.edu
Izabella Laba	University of British Columbia, Canada ilaba@math.ubc.ca	Gunther Uhlmann	University of Washington, USA gunther@math.washington.edu
Richard B. Melrose	Massachusetts Inst. of Tech., USA rhm@math.mit.edu	András Vasy	Stanford University, USA andras@math.stanford.edu
Frank Merle	Université de Cergy-Pontoise, France Frank.Merle@u-cergy.fr	Dan Virgil Voiculescu	University of California, Berkeley, USA dvv@math.berkeley.edu
William Minicozzi II	Johns Hopkins University, USA minicozz@math.jhu.edu	Steven Zelditch	Northwestern University, USA zelditch@math.northwestern.edu
Clément Mouhot	Cambridge University, UK c.mouhot@dpms.cam.ac.uk	Maciej Zworski	University of California, Berkeley, USA zworski@math.berkeley.edu
Werner Müller	Universität Bonn, Germany mueller@math.uni-bonn.de		

PRODUCTION

production@msp.org

Silvio Levy, Scientific Editor

See inside back cover or msp.org/apde for submission instructions.

The subscription price for 2020 is US \$340/year for the electronic version, and \$550/year (+\$60, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues from the last three years and changes of subscriber address should be sent to MSP.

Analysis & PDE (ISSN 1948-206X electronic, 2157-5045 printed) at Mathematical Sciences Publishers, 798 Evans Hall #3840, c/o University of California, Berkeley, CA 94720-3840, is published continuously online. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices.

APDE peer review and production are managed by EditFlow[®] from MSP.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2020 Mathematical Sciences Publishers

ANALYSIS & PDE

Volume 13 No. 7 2020

Refined mass-critical Strichartz estimates for Schrödinger operators CASEY JAO	1955
Scattering for defocusing energy subcritical nonlinear wave equations BENJAMIN DODSON, ANDREW LAWRIE, DANA MENDELSON and JASON MURPHY	1995
New formulas for the Laplacian of distance functions and applications FABIO CAVALLETTI and ANDREA MONDINO	2091
Convex sets evolving by volume-preserving fractional mean curvature flows ELEONORA CINTI, CARLO SINISTRARI and ENRICO VALDINOCI	2149
C^* -algebras isomorphically representable on l^p MARCH T. BOEDIHARDJO	2173
Exponential convergence of parabolic optimal transport on bounded domains FARHAN ABEDIN and JUN KITAGAWA	2183
Nuclear dimension of simple stably projectionless C^* -algebras JORGE CASTILLEJOS and SAMUEL EVINGTON	2205
On the regularity of minimizers for scalar integral functionals with (p, q) -growth PETER BELLA and MATHIAS SCHÄFFNER	2241