

ANALYSIS & PDE

Volume 13

No. 8

2020

BENJAMIN GESS, JONAS SAUER AND EITAN TADMOR

**OPTIMAL REGULARITY IN TIME AND SPACE
FOR THE POROUS MEDIUM EQUATION**

OPTIMAL REGULARITY IN TIME AND SPACE FOR THE POROUS MEDIUM EQUATION

BENJAMIN GESS, JONAS SAUER AND EITAN TADMOR

Regularity estimates in time and space for solutions to the porous medium equation are shown in the scale of Sobolev spaces. In addition, higher spatial regularity for powers of the solutions is obtained. Scaling arguments indicate that these estimates are optimal. In the linear limit, the proven regularity estimates are consistent with the optimal regularity of the linear case.

1. Introduction	2441
2. Preliminaries, notation and function spaces	2445
3. Optimality of estimates via scaling	2449
4. Averaging lemma approach	2452
5. Application to porous medium equations	2467
Appendix A. Kinetic solutions	2474
Appendix B. Fourier multipliers	2476
Acknowledgments	2478
References	2478

1. Introduction

We prove estimates on the time and space regularity of solutions to porous medium equations

$$\begin{cases} \partial_t u - \Delta u^{[m]} = S & \text{in } (0, T) \times \mathbb{R}^d, \\ u(0) = u_0 & \text{in } \mathbb{R}^d, \end{cases} \quad (1-1)$$

where $u^{[m]} := |u|^{m-1}u$ with $m > 1$, $u_0 \in L^1(\mathbb{R}^d)$ and $S \in L^1((0, T) \times \mathbb{R}^d)$. Solutions to porous medium equations are known to exhibit nonlinear phenomena like slow diffusion or filling up of holes at finite rate: If the initial data is compactly supported, then the support of the solution evolves with a free boundary that has finite speed of propagation. The solution close to the boundary is not smooth even for smooth initial data and zero forcing.

Despite many works on the problem of regularity of solutions to porous medium equations, until recently, established regularity results in the literature in terms of Hölder or Sobolev spaces were restricted to spatial differentiability of order less than 1; see [Ebmeyer 2005; Tadmor and Tao 2007]. For $m \searrow 1$ this is in stark contrast to the limiting case $m = 1$, where u is up to twice weakly differentiable in space. Very recently, the first author has proven optimal spatial regularity for (1-1) in [Gess 2020] for initial data

MSC2010: 35K59, 35B65, 35D30, 76S05.

Keywords: porous medium equation, entropy solutions, kinetic formulation, velocity averaging, regularity results.

$u_0 \in (L^1 \cap L^{1+\varepsilon})(\mathbb{R}^d)$ for some $\varepsilon > 0$. This leaves open three main aspects addressed in the present work: first, the derivation of optimal¹ space-time regularity, second, the limit case $u_0 \in L^1(\mathbb{R}^d)$, which is of particular importance since it covers the case of the Barenblatt solution for which the estimates are shown to be optimal, see [Section 3](#) below, and third, higher-order integrability. Solving these three open problems is the purpose of the present paper.

The first main result provides optimal space-time regularity for L^1 data.

Theorem 1.1. *Let $u_0 \in L^1(\mathbb{R}^d)$, $S \in L^1((0, T) \times \mathbb{R}^d)$ and $m \in (1, \infty)$. Let u be the unique entropy solution to (1-1) on $[0, T] \times \mathbb{R}^d$.*

(i) *Let $p \in (1, m]$ and define*

$$\kappa_t := \frac{m-p}{p} \frac{1}{m-1}, \quad \kappa_x := \frac{p-1}{p} \frac{2}{m-1}.$$

Then for all $\sigma_t \in [0, \kappa_t) \cup \{0\}$ and $\sigma_x \in [0, \kappa_x)$ we have

$$u \in W^{\sigma_t, p}(0, T; W^{\sigma_x, p}(\mathbb{R}^d)).$$

Moreover, we have the estimate

$$\|u\|_{W^{\sigma_t, p}(0, T; W^{\sigma_x, p}(\mathbb{R}^d))} \lesssim \|u_0\|_{L_x^1}^m + \|S\|_{L_{t,x}^1}^m + 1. \quad (1-2)$$

(ii) *Suppose $\mathcal{O} \Subset \mathbb{R}^d$. Let $s \in [0, 1]$ and define*

$$p := s(m-1) + 1, \quad \kappa_t := \frac{1-s}{s(m-1) + 1}, \quad \kappa_x := \frac{2s}{s(m-1) + 1}.$$

Then for all $\sigma_t \in [0, \kappa_t) \cup \{0\}$, $\sigma_x \in [0, \kappa_x) \cup \{0\}$ and $q \in [1, p]$ we have

$$u \in W^{\sigma_t, q}(0, T; W^{\sigma_x, q}(\mathcal{O})).$$

Moreover, we have the estimate

$$\|u\|_{W^{\sigma_t, q}(0, T; W^{\sigma_x, q}(\mathcal{O}))} \lesssim \|u_0\|_{L_x^1}^m + \|S\|_{L_{t,x}^1}^m + 1. \quad (1-3)$$

In [\[Tadmor and Tao 2007; Ebmeyer 2005\]](#) initial data in $L^1 \cap L^\infty$ was considered. However, the methods employed in these works did not allow a systematic analysis of the order of integrability of the solutions. For example, the results of [\[Ebmeyer 2005\]](#) are restricted to the particular order of integrability $p = 2/(m+1)$, while [\[Tadmor and Tao 2007\]](#) is restricted to $p = 1$. In the second main result we provide a systematic treatment of higher-order integrability. In particular, this includes and generalizes the corresponding results of [\[Ebmeyer 2005\]](#) in terms of regularity in Sobolev spaces.

Noting that the regularity of $u^{[m]}$ contains information on the time regularity of u in light of (1-1), in addition, we analyze the spatial regularity of powers of the solution u^μ for $\mu \in [1, m]$.

Theorem 1.2. *Let $u_0 \in L^1(\mathbb{R}^d) \cap L^\rho(\mathbb{R}^d)$, $S \in L^1([0, T] \times \mathbb{R}^d) \cap L^\rho([0, T] \times \mathbb{R}^d)$ for some $\rho \in (1, \infty)$ and assume $m \in (1, \infty)$. Let u be the unique entropy solution to (1-1) on $[0, T] \times \mathbb{R}^d$.*

¹Optimality is indicated by scaling arguments in [Section 3](#) below, and the derived estimates are consistent with the optimal space-time regularity in the linear case $m = 1$.

(i) Let $\mu \in [1, m]$. Then for all

$$p \in \left(1, \frac{m-1+\rho}{\mu}\right), \quad \sigma_x \in \left[0, \frac{\mu p - 1}{p} \frac{2}{m-2+\rho}\right)$$

we have

$$u^{[\mu]} \in L^p(0, T; W^{\sigma_x, p}(\mathbb{R}^d)),$$

and we have the estimate

$$\|u^{[\mu]}\|_{L^p(0, T; W^{\sigma_x, p}(\mathbb{R}^d))} \lesssim \|u_0\|_{L_x^1 \cap L_x^\rho}^{\mu\rho} + \|S\|_{L_{t,x}^1 \cap L_{t,x}^\rho}^{\mu\rho} + 1. \quad (1-4)$$

(ii) Let $p \in (\rho, m-1+\rho)$ and define

$$\kappa_t := \frac{m-1+\rho-p}{p} \frac{1}{m-1}, \quad \kappa_x := \frac{p-\rho}{p} \frac{2}{m-1}.$$

Then for all $\sigma_t \in [0, \kappa_t)$ and $\sigma_x \in [0, \kappa_x)$ we have

$$u \in W^{\sigma_t, p}(0, T; W^{\sigma_x, p}(\mathbb{R}^d)).$$

Moreover, we have the estimate

$$\|u\|_{W^{\sigma_t, p}(0, T; W^{\sigma_x, p}(\mathbb{R}^d))} \lesssim \|u_0\|_{L_x^1 \cap L_x^\rho}^\rho + \|S\|_{L_{t,x}^1 \cap L_{t,x}^\rho}^\rho + 1. \quad (1-5)$$

Much as in [Theorem 1.1](#), if one restricts to estimates that are localized in space, the rigid interdependency of the coefficients in [Theorem 1.2](#) can be relaxed.

Corollary 1.3. Under the assumptions of [Theorem 1.2](#), suppose $\mathcal{O} \in \mathbb{R}^d$.

(i) Let $\mu \in [1, m]$. Then for all $\sigma_x \in [0, 2\mu/m)$ and $q \in [1, m/\mu]$ we have

$$u^{[\mu]} \in L^q(0, T; W^{\sigma_x, q}(\mathcal{O})),$$

and we have the estimate

$$\|u^{[\mu]}\|_{L^q(0, T; W^{\sigma_x, q}(\mathcal{O}))} \lesssim \|u_0\|_{L_x^1 \cap L_x^\rho}^{\mu\rho} + \|S\|_{L_{t,x}^1 \cap L_{t,x}^\rho}^{\mu\rho} + 1. \quad (1-6)$$

(ii) Let $s \in [0, 1]$ and define

$$p := s(m-1) + 1, \quad \kappa_t := \frac{1-s}{s(m-1)+1}, \quad \kappa_x := \frac{2s}{s(m-1)+1}.$$

Then for all $\sigma_t \in [0, \kappa_t) \cup \{0\}$, $\sigma_x \in [0, \kappa_x) \cup \{0\}$ and $q \in [1, p]$ we have

$$u \in W^{\sigma_t, q}(0, T; W^{\sigma_x, q}(\mathcal{O})).$$

Moreover, we have the estimate

$$\|u\|_{W^{\sigma_t, q}(0, T; W^{\sigma_x, q}(\mathcal{O}))} \lesssim \|u_0\|_{L_x^1 \cap L_x^\rho}^\rho + \|S\|_{L_{t,x}^1 \cap L_{t,x}^\rho}^\rho + 1. \quad (1-7)$$

The methods employed in this work are inspired by [\[Tadmor and Tao 2007\]](#) and rely on the kinetic form of (1-1), that is, with $f(t, x, v) := 1_{v < u(t, x)} - 1_{v < 0}$,

$$\partial_t f - m|v|^{m-1} \Delta_x f = \partial_v q + S(t, x) \delta_{u(t, x)}(v) \quad (1-8)$$

for a nonnegative measure q , which allows the use of averaging lemmas and real interpolation. There is a relatively short yet intense history of applying such velocity-averaging techniques to deduce regularizing effects in nonlinear PDEs — from the early works [DiPerna, Lions, and Meyer 1991; Golse, Lions, Perthame, and Sentis 1988; Lions, Perthame, and Tadmor 1994a; 1994b; 1996] to the more recent [Arsénio and Masmoudi 2019; DeVore and Petrova 2001; Golse and Perthame 2013; Golse and Saint-Raymond 2004; Jabin 2009; Jabin and Vega 2004; Perthame 2002]. An essential difference to purely spatial regularity consists in the necessity to work with anisotropic fractional Sobolev spaces, which only in their homogeneous form are nicely adapted to the Fourier analytic methods of this work, much in contrast to the purely spatial case in [Gess 2020]. This leads to the so-called dominating mixed anisotropic Besov spaces introduced in [Schmeisser and Triebel 1987]. Passing from these homogeneous anisotropic spaces to standard inhomogeneous fractional Sobolev spaces is delicate and treated in detail below. A main ingredient in the proof of optimal regularity in [Gess 2020] was the existence of singular moments $\int_{t,x,v} |v|^{-\gamma} q$ for $\gamma \in (0, 1)$. This ceases to be true for general L^1 -initial data. This difficulty is overcome in the present work by treating separately the degeneracy at $|v| = 0$ and the singularity at $|v| = \infty$ as they appear in (1-8). This also necessitates making use of (1-8) in the case of small spatial modes ξ in order to obtain optimal time regularity; see Corollary 4.7 below.

Comments on the literature. The (spatial) regularity of solutions to porous medium equations in Sobolev spaces has previously been considered in [Ebmeyer 2005; Gess 2020; Tadmor and Tao 2007]. Since our main focus is on time-space regularity, we refer to [Gess 2020] for a more detailed account on the available literature in this regard.

In the case of nonnegative solutions the spatial regularity of special types of powers of solutions has been investigated in the literature. For example, much work is devoted to the pressure defined by $v := (m/(m-1))u^{m-1}$; see, e.g., [Vázquez 2007]. In the recent work [Gianazza and Schwarzacher 2019] the authors proved higher integrability for nonnegative, local weak solutions to forced porous medium equations in the sense that $u^{(m+1)/2} \in L_{\text{loc}}^{2+\varepsilon}((0, T); W_{\text{loc}}^{1,2+\varepsilon})$ for all $\varepsilon > 0$ small enough. This result was generalized in [Bögelein, Duzaar, Korte, and Scheven 2019].

The analysis of regularity in time of solutions to porous medium equations (without forcing) has a long history initiated in [1979] and continued in [Crandall, Pazy, and Tartar 1979; Bénilan and Crandall 1981], where it was shown that

$$\partial_t u \in L_{\text{loc}}^1((0, \infty); L^1(\mathbb{R}^d)) \quad (1-9)$$

for $u_0 \in L^1(\mathbb{R}^d)$. Subsequently, Crandall and Pierre [1982a; 1982b] devoted considerable effort to relaxing the required assumptions on the nonlinearity ψ in the case of generalized porous medium equations

$$\partial_t u - \Delta \psi(u) = 0 \quad \text{in } (0, T) \times \mathbb{R}^d. \quad (1-10)$$

More precisely, in [Crandall and Pierre 1982a] assuming the global generalized homogeneity condition

$$v \frac{\psi(r)\psi''(r)}{(\psi'(r))^2} \in [m, M] \quad (1-11)$$

for some $0 < m < M$, $v \in \{\pm 1\}$ and all $r \in \mathbb{R}$, (1-9) was recovered.

It should be noted that the methods developed in these works are restricted to the nonforced case $S \equiv 0$. In fact, for $S \not\equiv 0$, the linear case $m = 1$ demonstrates that (1-9) should not be expected. We are not aware of results proving regularity in time in Sobolev spaces for porous medium equations with nonvanishing forcing. In this sense, restricting to regularity in time alone, the results of the present work can be regarded as the (partial) extension of the results of [Aronson and B  nilan 1979; B  nilan and Crandall 1981; Crandall, Pazy, and Tartar 1979; Crandall and Pierre 1982a; 1982b] to nonvanishing forcing.

We are not aware of previous results on mixed time and space regularity in Sobolev spaces for solutions to porous medium equations.

For simplicity of the presentation we restrict to the nonlinearity $\psi(u) = u^{[m]}$ in this work. However, the methods that we present are not restricted to this case, as long as ψ satisfies a nonlinearity condition as in [Gess 2020]. In addition, by means of a velocity decomposition, i.e., writing

$$u(t, x) = \sum_{i=1}^K u^i(t, x) := \sum_{i=1}^K \int_v \varphi^i(v) f(t, x, v) dv,$$

where φ^i , $i = 1, \dots, K$, is a smooth decomposition of the unity, such a nonlinearity condition only needs to be supposed locally at points of degeneracy. This is in contrast to the assumptions, such as (1-11), supposed in the series of works [Aronson and B  nilan 1979; B  nilan and Crandall 1981; Crandall, Pazy, and Tartar 1979; Crandall and Pierre 1982a; 1982b] mentioned above, which can be regarded as *global* generalized homogeneity conditions.

Structure of this work. In Section 2 we collect information on the class of homogeneous and inhomogeneous anisotropic, dominating mixed-derivative spaces employed in this work. The optimality of the obtained estimates is indicated in Section 3 by scaling arguments and by explicit computations in case of the Barenblatt solution. In Section 4 we provide general averaging lemmas (Lemmas 4.2 and 4.4) in the framework of homogeneous dominating mixed-derivative spaces and translate them to more standard inhomogeneous anisotropic fractional Sobolev spaces (Corollaries 4.5, 4.6 and 4.7). In this formulation, they imply the main result by their application to the porous medium equation in Section 5.

2. Preliminaries, notation and function spaces

We use the notation $a \lesssim b$ if there is a universal constant $C > 0$ such that $a \leq Cb$. We introduce $a \gtrsim b$ in a similar manner, and write $a \sim b$ if $a \lesssim b$ and $a \gtrsim b$. For a Banach space X and $I \subset \mathbb{R}$ we denote by $C(I; X)$ the space of bounded and continuous X -valued functions endowed with the norm $\|f\|_{C(I; X)} := \sup_{t \in I} \|f(t)\|_X$. If $X = \mathbb{R}$ we write $C(I)$. For $k \in \mathbb{N} \cup \{\infty\}$, the space of k -times continuously differentiable functions is denoted by $C^k(I; X)$. The subspace of $C^k(I; X)$ consisting of compactly supported functions is denoted by $C_c^k(I; X)$. Moreover, we write \mathcal{M}_{TV} for the space of all measures with finite total variation. Throughout this article we use several types of L^p -based function spaces. For a Banach space X and $p \in [1, \infty]$, we endow the Bochner–Lebesgue space $L^p(\mathbb{R}; X)$ with the usual norm

$$\|f\|_{L^p(\mathbb{R}; X)} := \left(\int_{\mathbb{R}} \|f(t)\|_X^p dt \right)^{\frac{1}{p}},$$

with the standard modification in the case of $p = \infty$. For $k \in \mathbb{N}_0 := \mathbb{N} \cup \{0\}$, the corresponding X -valued Sobolev space is denoted by $W^{k,p}(\mathbb{R}; X)$. If $\sigma \in (0, \infty)$ is noninteger (say $\sigma = k + r$, with $k \in \mathbb{N}_0$ and $r \in (0, 1)$), then we define the X -valued Sobolev–Slobodecki space $W^{\sigma,p}(\mathbb{R}; X)$ as the space of functions in $W^{k,p}(\mathbb{R}; X)$ with

$$\|f\|_{\dot{W}^{\sigma,p}(\mathbb{R}; X)} := \left(\int_{\mathbb{R} \times \mathbb{R}} \frac{\|D^k f(t) - D^k f(s)\|_X^p}{|t-s|^{rp+1}} \, ds \, dt \right)^{\frac{1}{p}} < \infty, \quad (2-1)$$

again with the usual modification in the case of $p = \infty$. Further, let $\dot{W}^{\sigma,p}(\mathbb{R}; X)$ be the space of all locally integrable X -valued functions f for which (2-1) is finite. If we factor out the equivalence relation \sim , where $f \sim g$ if $\|f - g\|_{\dot{W}^{\sigma,p}(\mathbb{R}; X)} = 0$, the space $\dot{W}^{\sigma,p}(\mathbb{R}; X)$ equipped with the norm $\|\cdot\|_{\dot{W}^{\sigma,p}(\mathbb{R}; X)}$ is a Banach space.

Moreover, in order to treat regularity results in both time and space efficiently, we introduce spaces with dominating mixed derivatives set in the framework of Fourier analysis, that is, corresponding Besov spaces. These spaces have a long history in the literature, beginning with [Nikolsky 1962; 1963a; 1963b]. We refer the reader to [Schmeisser and Triebel 1987]. We adopt the notation of [Schmeisser and Triebel 1987] for the nonhomogeneous spaces. Corresponding homogeneous Besov spaces are treated in [Triebel 1977a; 1977b]; we adapt the notation to be consistent with that of [Schmeisser and Triebel 1987]. We recall from [Triebel 1977a] the definition of the spaces \mathcal{Z} and \mathcal{Z}' replacing the standard Schwartz space $\mathcal{S} = \mathcal{S}(\mathbb{R}^{d+1})$ and the space of tempered distributions $\mathcal{S}' = \mathcal{S}'(\mathbb{R}^{d+1})$ in the definition of homogeneous spaces. As we are concerned with function spaces in the time variable $t \in \mathbb{R}$ and the spatial variable $x \in \mathbb{R}^d$, we introduce, in addition to $\mathbb{R}^{d+1} = \mathbb{R}_t \times \mathbb{R}_x^d$, also the subset

$$\dot{\mathbb{R}}^{d+1} := \{(t, x) \in \mathbb{R}^{d+1} : t|x| \neq 0\}.$$

Note that in [Triebel 1977a], the notation \mathbb{R}^{d+1}_+ is used, which gives a better geometrical intuition of the set taken out of \mathbb{R}^{d+1} . However, for typesetting reasons, we have decided on the notation $\dot{\mathbb{R}}^{d+1}$. Then we let $\dot{\mathcal{D}}$ be the subset of the standard space of test functions \mathcal{D} , consisting of functions with compact support in $\dot{\mathbb{R}}^{d+1}$ and view it as a locally convex space equipped with the canonical topology. Its dual space is denoted by $\dot{\mathcal{D}}'$, and is referred to as distributions over $\dot{\mathbb{R}}^{d+1}$. We define \mathcal{Z} as the image of $\dot{\mathcal{D}} \subset \mathcal{S}$ under the Fourier transform \mathcal{F} in time and space, equipped with the inherited topology from $\dot{\mathcal{D}}$. The corresponding dual space is denoted by \mathcal{Z}' . Since $\mathcal{F} : \dot{\mathcal{D}} \rightarrow \mathcal{Z}$, we can define by duality the Fourier transform $\mathcal{F} : \mathcal{Z}' \rightarrow \dot{\mathcal{D}}'$.

It holds $\mathcal{Z} \subset \mathcal{S}$ with a continuous embedding, but the fact that \mathcal{Z} is not densely embedded in \mathcal{S} prevents one from stating $\mathcal{S}' \subset \mathcal{Z}'$. However, we note that for $p \in (1, \infty)$, the space $L^p(\mathbb{R}^{d+1})$ can be viewed both as subspace of \mathcal{S}' and as a subspace of \mathcal{Z}' ; see Theorem 3.3 in [Triebel 1977a].

Let φ be a smooth function supported in the annulus $\{\xi \in \mathbb{R}^d : \frac{1}{2} \leq |\xi| \leq 2\}$ and such that

$$\sum_{j \in \mathbb{Z}} \varphi_j(\xi) := \sum_{j \in \mathbb{Z}} \varphi(2^{-j} \xi) = 1 \quad \text{for all } \xi \in \mathbb{R}^d \setminus \{0\}.$$

Similarly, let η be a smooth function supported in $(-2, -\frac{1}{2}) \cup (\frac{1}{2}, 2)$ with

$$\sum_{l \in \mathbb{Z}} \eta_l(\tau) := \sum_{l \in \mathbb{Z}} \eta(2^{-l} \tau) = 1 \quad \text{for all } \tau \in \mathbb{R} \setminus \{0\}.$$

Moreover, define $\phi_j := \varphi_j$ for $j \geq 1$ and $\phi_0 := 1 - \sum_{j \geq 1} \phi_j$, as well as $\psi_l := \eta_l$ for $l \geq 1$ and $\psi_0 := 1 - \sum_{l \geq 1} \eta_l$. We will use the shorthand notation $\eta_l \varphi_j$ for the function $(\tau, \xi) \mapsto \eta_l(\tau) \varphi_j(\xi)$, and similarly for combinations of ψ_l and ϕ_j .

Definition 2.1. Let $\sigma_i \in (-\infty, \infty)$, $i = t, x$, and $p \in [1, \infty]$. Set $\bar{\sigma} := (\sigma_t, \sigma_x)$.

(i) The homogeneous Besov space with dominating mixed derivatives $S_{p,\infty}^{\bar{\sigma}} \dot{B}(\mathbb{R}^{d+1})$ is given by

$$S_{p,\infty}^{\bar{\sigma}} \dot{B} := S_{p,\infty}^{\bar{\sigma}} \dot{B}(\mathbb{R}^{d+1}) := \{f \in \mathcal{S}' : \|f\|_{S_{p,\infty}^{\bar{\sigma}} \dot{B}} < \infty\},$$

with the norm

$$\|f\|_{S_{p,\infty}^{\bar{\sigma}} \dot{B}} := \sup_{l,j \in \mathbb{Z}} 2^{\sigma_t l} 2^{\sigma_x j} \|\mathcal{F}_{t,x}^{-1} \eta_l \varphi_j \mathcal{F}_{t,x} f\|_{L^p(\mathbb{R}^{d+1})}.$$

Similarly, the space $S_{p,\infty,(\infty)}^{\bar{\sigma}} \dot{B}(\mathbb{R}^{d+1})$ is given via the norm

$$\|f\|_{S_{p,\infty,(\infty)}^{\bar{\sigma}} \dot{B}} := \sup_{l,j \in \mathbb{Z}} 2^{\sigma_t l} 2^{\sigma_x j} \|\mathcal{F}_{t,x}^{-1} \eta_l \varphi_j \mathcal{F}_{t,x} f\|_{L^{p,\infty}(\mathbb{R}^{d+1})}.$$

(ii) The homogeneous Chemin–Lerner spaces $\tilde{L}_t^p \dot{B}_{p,\infty}^{\sigma_x}(\mathbb{R}^{d+1})$ and $\tilde{L}_x^p \dot{B}_{p,\infty}^{\sigma_t}(\mathbb{R}^{d+1})$ are given by

$$\tilde{L}_t^p \dot{B}_{p,\infty}^{\sigma_x} := \tilde{L}_t^p \dot{B}_{p,\infty}^{\sigma_x}(\mathbb{R}^{d+1}) := \{f \in \mathcal{S}' : \|f\|_{\tilde{L}_t^p \dot{B}_{p,\infty}^{\sigma_x}} < \infty\},$$

$$\tilde{L}_x^p \dot{B}_{p,\infty}^{\sigma_t} := \tilde{L}_x^p \dot{B}_{p,\infty}^{\sigma_t}(\mathbb{R}^{d+1}) := \{f \in \mathcal{S}' : \|f\|_{\tilde{L}_x^p \dot{B}_{p,\infty}^{\sigma_t}} < \infty\},$$

with the norms

$$\|f\|_{\tilde{L}_t^p \dot{B}_{p,\infty}^{\sigma_x}} := \sup_{j \in \mathbb{Z}} 2^{\sigma_x j} \|\mathcal{F}_x^{-1} \varphi_j \mathcal{F}_x f\|_{L^p(\mathbb{R}^{d+1})},$$

$$\|f\|_{\tilde{L}_x^p \dot{B}_{p,\infty}^{\sigma_t}} := \sup_{l \in \mathbb{Z}} 2^{\sigma_t l} \|\mathcal{F}_t^{-1} \eta_l \mathcal{F}_t f\|_{L^p(\mathbb{R}^{d+1})},$$

respectively.

(iii) The nonhomogeneous Besov space with dominating mixed derivatives $S_{p,\infty}^{\bar{\sigma}} B(\mathbb{R}^{d+1})$ is given by

$$S_{p,\infty}^{\bar{\sigma}} B := S_{p,\infty}^{\bar{\sigma}} B(\mathbb{R}^{d+1}) := \{f \in \mathcal{S}'(\mathbb{R}^{d+1}) : \|f\|_{S_{p,\infty}^{\bar{\sigma}} B} < \infty\},$$

with the norm

$$\|f\|_{S_{p,\infty}^{\bar{\sigma}} B} := \sup_{l,j \geq 0} 2^{\sigma_t l} 2^{\sigma_x j} \|\mathcal{F}_{t,x}^{-1} \psi_l \phi_j \mathcal{F}_{t,x} f\|_{L^p(\mathbb{R}^{d+1})}.$$

(iv) The nonhomogeneous Chemin–Lerner space $\tilde{L}_t^p B_{p,\infty}^{\sigma_x}(\mathbb{R}^{d+1})$ is given by

$$\tilde{L}_t^p B_{p,\infty}^{\sigma_x} := \tilde{L}_t^p B_{p,\infty}^{\sigma_x}(\mathbb{R}^{d+1}) := \{f \in \mathcal{S}' : \|f\|_{\tilde{L}_t^p B_{p,\infty}^{\sigma_x}} < \infty\},$$

with the norm $\|f\|_{\tilde{L}_t^p B_{p,\infty}^{\sigma_x}} := \sup_{j \geq 0} 2^{\sigma_x j} \|\mathcal{F}_x^{-1} \phi_j \mathcal{F}_x f\|_{L^p(\mathbb{R}^{d+1})}$.

Remark 2.2. All spaces considered in Definition 2.1 are Banach spaces; see [Triebel 1977a]. Note that for $\vartheta \in \mathbb{R}$, we use the notation $\vartheta \bar{\sigma} = (\vartheta \sigma_t, \vartheta \sigma_x)$. In this note, we restrict ourselves to the third index of the Besov-type space being infinity, in which case the spaces $S_{p,\infty}^{\bar{\sigma}} B$ are sometimes called *Nikolsky spaces of dominating mixed derivatives* in the literature. However, there is no conceptual limitation to consider also third indices $q \in [1, \infty]$. By the same token, one could also consider spaces with different indices p and q in different directions. We refer the reader to [Schmeisser and Triebel 1987] for more details concerning such spaces.

Lemma 2.3. *Let $\kappa_x \geq 0$ and $p \in [1, \infty]$. Then*

$$\tilde{L}_t^p B_{p,\infty}^{\kappa_x+\varepsilon}(\mathbb{R}^{d+1}) \subset L^p(\mathbb{R}; W^{\kappa_x,p}(\mathbb{R}^d)) \subset \tilde{L}_t^p B_{p,\infty}^{\kappa_x-\delta}(\mathbb{R}^{d+1}),$$

whenever $\varepsilon > 0$ and $\delta \in (0, \kappa_x]$.

Proof. This follows from [Bahouri, Chemin, and Danchin 2011, p. 98]. \square

Lemma 2.4. *Let $\kappa_t, \kappa_x > 0$ and $p \in [1, \infty)$. Then $S_{p,\infty}^{\bar{\kappa}} B \subset W^{\sigma_t,p}(\mathbb{R}; W^{\sigma_x,p}(\mathbb{R}^d))$ whenever $\sigma_t \in [0, \kappa_t)$ and $\sigma_x \in [0, \kappa_x)$.*

Proof. The proof is a combination of results in [Schmeisser and Triebel 1987], which are written for $\mathbb{R} \times \mathbb{R}$ but also true for $\mathbb{R} \times \mathbb{R}^d$ by an inspection of their respective proofs: Without loss of generality, we can assume that σ_t and σ_x are noninteger. By [loc. cit., Section 2.3.4, Remark 4], we have $W^{\sigma_t,p}(\mathbb{R}; W^{\sigma_x,p}(\mathbb{R}^d)) = SB_{p,p}^{\bar{\sigma}}$; see [loc. cit., Section 2.2.1, Definition 2] for a definition of the latter space. Since by [loc. cit., Section 2.2.3, Proposition 2] we have $S_{p,\infty}^{\bar{\kappa}} B \subset SB_{p,p}^{\bar{\sigma}}$, this yields the claim. \square

Lemma 2.5. *Let $\sigma_t, \sigma_x > 0$ and $p \in [1, \infty]$. Then*

$$(L^p(\mathbb{R}^{d+1}) \cap \tilde{L}_x^p \dot{B}_{p,\infty}^{\sigma_t} \cap \tilde{L}_t^p \dot{B}_{p,\infty}^{\sigma_x} \cap S_{p,\infty}^{\bar{\sigma}} \dot{B}) = S_{p,\infty}^{\bar{\sigma}} B$$

with equivalent norms.

Proof. As smooth and compactly supported functions, ψ_0 and ϕ_0 extend to L^p multipliers for all $p \in [1, \infty]$; see, e.g., [Bergh and Löfström 1976].

For $f \in (L^p(\mathbb{R}^{d+1}) \cap \tilde{L}_x^p \dot{B}_{p,\infty}^{\sigma_t} \cap \tilde{L}_t^p \dot{B}_{p,\infty}^{\sigma_x} \cap S_{p,\infty}^{\bar{\sigma}} \dot{B}) \subset \mathcal{S}'(\mathbb{R}^{d+1})$ we obtain

$$\begin{aligned} \|f\|_{S_{p,\infty}^{\bar{\sigma}} B} &\leq \|\mathcal{F}_{t,x}^{-1} \psi_0 \phi_0 \mathcal{F}_{t,x} f\|_{L_{t,x}^p} + \sup_{l>0} 2^{\sigma_t l} \|\mathcal{F}_{t,x}^{-1} \eta_l \phi_0 \mathcal{F}_{t,x} f\|_{L_{t,x}^p} \\ &\quad + \sup_{j>0} 2^{\sigma_x j} \|\mathcal{F}_{t,x}^{-1} \psi_0 \varphi_j \mathcal{F}_{t,x} f\|_{L_{t,x}^p} + \sup_{l,j>0} 2^{\sigma_t l} 2^{\sigma_x j} \|\mathcal{F}_{t,x}^{-1} \eta_l \varphi_j \mathcal{F}_{t,x} f\|_{L_{t,x}^p} \\ &\lesssim \|f\|_{L_{t,x}^p} + \sup_{l>0} 2^{\sigma_t l} \|\mathcal{F}_t^{-1} \eta_l \mathcal{F}_t f\|_{L_{t,x}^p} \\ &\quad + \sup_{j>0} 2^{\sigma_x j} \|\mathcal{F}_x^{-1} \varphi_j \mathcal{F}_x f\|_{L_{t,x}^p} + \sup_{l,j>0} 2^{\sigma_t l} 2^{\sigma_x j} \|\mathcal{F}_{t,x}^{-1} \eta_l \varphi_j \mathcal{F}_{t,x} f\|_{L_{t,x}^p} \\ &\lesssim \|f\|_{L_{t,x}^p} + \|f\|_{\tilde{L}_x^p \dot{B}_{p,\infty}^{\sigma_t}} + \|f\|_{\tilde{L}_t^p \dot{B}_{p,\infty}^{\sigma_x}} + \|f\|_{S_{p,\infty}^{\bar{\sigma}} \dot{B}}. \end{aligned}$$

Conversely, for $f \in S_{p,\infty}^{\bar{\sigma}} B$, we estimate the four contributions corresponding to $L^p(\mathbb{R}^{d+1})$, $\tilde{L}_x^p \dot{B}_{p,\infty}^{\sigma_t}$, $\tilde{L}_t^p \dot{B}_{p,\infty}^{\sigma_x}$, and $S_{p,\infty}^{\bar{\sigma}} \dot{B}$ separately. We start by noting that due to $\sigma_t, \sigma_x > 0$, the invariance of multiplier norms with respect to dilation, $\eta_l = \eta_l \tilde{\psi}_0$ for $l \leq 0$ and $\varphi_j = \varphi_j \tilde{\phi}_0$ for $j \leq 0$, where $\tilde{\psi}_0 := \psi_0 + \psi_1$ and $\tilde{\phi}_0 := \phi_0 + \phi_1$, we have

$$\begin{aligned} \sup_{l \leq 0} 2^{\sigma_t l} \|\mathcal{F}_t^{-1} \eta_l \mathcal{F}_t f\|_{L_{t,x}^p} &\lesssim \|\mathcal{F}_t^{-1} \tilde{\psi}_0 \mathcal{F}_t f\|_{L_{t,x}^p}, \\ \sup_{j \leq 0} 2^{\sigma_x j} \|\mathcal{F}_x^{-1} \varphi_j \mathcal{F}_x f\|_{L_{t,x}^p} &\lesssim \|\mathcal{F}_x^{-1} \tilde{\phi}_0 \mathcal{F}_x f\|_{L_{t,x}^p}. \end{aligned}$$

Furthermore we use the fact that for $\sigma > 0$ one has the estimate $\sum_{n \geq 0} |a_n| \lesssim \sup_{n \geq 0} 2^{\sigma n} |a_n|$ for any sequence $(a_n) \subset \mathbb{R}$ with a constant depending on σ . With this, we obtain

$$\|f\|_{L_{t,x}^p} \leq \sum_{l,j \geq 0} \|\mathcal{F}_{t,x}^{-1} \psi_l \phi_j \mathcal{F}_{t,x} f\|_{L_{t,x}^p} \lesssim \sup_{l,j \geq 0} 2^{\sigma_l l} 2^{\sigma_x j} \|\mathcal{F}_{t,x}^{-1} \psi_l \phi_j \mathcal{F}_{t,x} f\|_{L_{t,x}^p} \leq \|f\|_{S_{p,\infty}^{\tilde{\sigma}} B}.$$

Next, we compute

$$\begin{aligned} \|f\|_{\tilde{L}_x^p \dot{B}_{p,\infty}^{\sigma_t}} &\leq \sup_{l \leq 0} 2^{\sigma_l l} \|\mathcal{F}_t^{-1} \eta_l \mathcal{F}_t f\|_{L_{t,x}^p} + \sup_{l > 0} 2^{\sigma_l l} \|\mathcal{F}_t^{-1} \psi_l \mathcal{F}_t f\|_{L_{t,x}^p} \\ &\lesssim \|\mathcal{F}_t^{-1} \tilde{\psi}_0 \mathcal{F}_t f\|_{L_{t,x}^p} + \sup_{l > 0} 2^{\sigma_l l} \|\mathcal{F}_t^{-1} \psi_l \mathcal{F}_t f\|_{L_{t,x}^p} \\ &\leq \sum_{j \geq 0} \|\mathcal{F}_{t,x}^{-1} \tilde{\psi}_0 \phi_j \mathcal{F}_{t,x} f\|_{L_{t,x}^p} + \sup_{l > 0} \sum_{j \geq 0} 2^{\sigma_l l} \|\mathcal{F}_{t,x}^{-1} \psi_l \phi_j \mathcal{F}_{t,x} f\|_{L_{t,x}^p} \\ &\lesssim \sup_{j \geq 0} 2^{\sigma_x j} \|\mathcal{F}_{t,x}^{-1} \tilde{\psi}_0 \phi_j \mathcal{F}_{t,x} f\|_{L_{t,x}^p} + \sup_{l > 0, j \geq 0} 2^{\sigma_l l} 2^{\sigma_x j} \|\mathcal{F}_{t,x}^{-1} \psi_l \phi_j \mathcal{F}_{t,x} f\|_{L_{t,x}^p} \lesssim \|f\|_{S_{p,\infty}^{\tilde{\sigma}} B}. \end{aligned}$$

By analogy, $\|f\|_{\tilde{L}_t^p \dot{B}_{p,\infty}^{\sigma_x}} \lesssim \|f\|_{S_{p,\infty}^{\tilde{\sigma}} B}$. Hence, it remains to control $\|f\|_{S_{p,\infty}^{\tilde{\sigma}} \dot{B}}$. We split this term into the four contributions

$$\begin{aligned} \|f\|_{S_{p,\infty}^{\tilde{\sigma}} \dot{B}} &= \sup_{l,j > 0} 2^{\sigma_l l} 2^{\sigma_x j} \|\mathcal{F}_{t,x}^{-1} \psi_l \phi_j \mathcal{F}_{t,x} f\|_{L_{t,x}^p} + \sup_{l > 0, j \leq 0} 2^{\sigma_l l} 2^{\sigma_x j} \|\mathcal{F}_{t,x}^{-1} \psi_l \varphi_j \mathcal{F}_{t,x} f\|_{L_{t,x}^p} \\ &\quad + \sup_{l \leq 0, j > 0} 2^{\sigma_l l} 2^{\sigma_x j} \|\mathcal{F}_{t,x}^{-1} \eta_l \phi_j \mathcal{F}_{t,x} f\|_{L_{t,x}^p} + \sup_{l,j \leq 0} 2^{\sigma_l l} 2^{\sigma_x j} \|\mathcal{F}_{t,x}^{-1} \eta_l \varphi_j \mathcal{F}_{t,x} f\|_{L_{t,x}^p}. \end{aligned}$$

The first contribution is immediately estimated by $\|f\|_{S_{p,\infty}^{\tilde{\sigma}} B}$. For the second contribution, we have

$$\sup_{l > 0, j \leq 0} 2^{\sigma_l l} 2^{\sigma_x j} \|\mathcal{F}_{t,x}^{-1} \psi_l \varphi_j \mathcal{F}_{t,x} f\|_{L_{t,x}^p} \lesssim \sup_{l > 0} 2^{\sigma_l l} \|\mathcal{F}_{t,x}^{-1} \psi_l \tilde{\phi}_0 \mathcal{F}_{t,x} f\|_{L_{t,x}^p} \leq \|f\|_{S_{p,\infty}^{\tilde{\sigma}} B},$$

and a similar estimate holds for the third contribution. For the fourth contribution, we have

$$\sup_{l,j \leq 0} 2^{\sigma_l l} 2^{\sigma_x j} \|\mathcal{F}_{t,x}^{-1} \eta_l \varphi_j \mathcal{F}_{t,x} f\|_{L_{t,x}^p} \lesssim \|\mathcal{F}_{t,x}^{-1} \tilde{\psi}_0 \tilde{\phi}_0 \mathcal{F}_{t,x} f\|_{L_{t,x}^p}. \quad \square$$

3. Optimality of estimates via scaling

It is well known that in the linear case $m = 1$ one has estimates of the form

$$\|u\|_{L_t^1 \dot{W}_x^{\sigma_x, 1}} \leq c(\sigma_x) (\|u_0\|_{L_x^1} + \|S\|_{L_{t,x}^1}), \quad (3-1)$$

for all $\sigma_x < 2$. In the case $m > 1$, such an estimate cannot be true for any $\sigma_x > 0$ anymore. Intuitively, this is due to the linear nature of (3-1) (observe that the integrability exponent is equal on both sides of the inequality), which is not compatible with the nonlinear equation (1-1). We will make this intuition more precise by the following lemma based on a scaling argument.

Lemma 3.1. *Let $T > 0$, $m > 1$, $\mu \in [1, m]$, $p \in [1, \infty)$ and $\sigma_t, \sigma_x \geq 0$. Assume that there is a constant $c = c(m, \mu, p, \sigma_t, \sigma_x) > 0$ such that*

$$\|u^{[\mu]}\|_{\dot{W}^{\sigma_t, p}(0, T; \dot{W}^{\sigma_x, p}(\mathbb{R}^d))}^p \leq c(\|u_0\|_{L^1(\mathbb{R}^d)} + \|S\|_{L^1(0, T; L^1(\mathbb{R}^d))}) \quad (3-2)$$

for all solutions u to (1-1). Then

$$p \leq \frac{m}{\mu + (m-1)\sigma_t} \leq \frac{m}{\mu}, \quad \sigma_t \leq \frac{m - \mu p}{p(m-1)} \leq \frac{m - \mu}{m-1}, \quad \sigma_x = \frac{\mu p - 1}{p} \frac{2}{m-1} \leq \frac{2(\mu - \sigma_t)}{m} \leq \frac{2\mu}{m}. \quad (3-3)$$

In particular, if $\sigma_t = (m - \mu)/(m - 1)$, then $p = 1$ and $\sigma_x = 2(\mu - 1)/(m - 1)$.

Proof. For positive constants $\eta, \gamma \geq 1$ with $\eta^{m-1} = \gamma$ and a fixed triple (u, u_0, S) such that u satisfies (1-1) with initial condition u_0 and forcing S we consider the rescaled quantities $(\tilde{u}, \tilde{u}_0, \tilde{S})$ defined via

$$\tilde{u}(t, x) := \eta u(\gamma t, x), \quad \tilde{u}_0(x) := \eta u_0(x), \quad \tilde{S}(t, x) := \eta^m S(\gamma t, x),$$

where we have tacitly extended S on $(T, \gamma T)$ by 0. Then \tilde{u} satisfies (1-1) with $\tilde{u}_0 \in L^1(\mathbb{R}^d)$ and $\tilde{S} \in L^1(0, T; L^1(\mathbb{R}^d))$, so that (3-2) gives

$$\|\tilde{u}^{[\mu]}\|_{\dot{W}^{\sigma_t, p}(0, T; \dot{W}^{\sigma_x, p}(\mathbb{R}^d))}^p \leq c(\|\tilde{u}_0\|_{L^1(\mathbb{R}^d)} + \|\tilde{S}\|_{L^1(0, T; L^1(\mathbb{R}^d))}). \quad (3-4)$$

We observe

$$\|\tilde{u}^{[\mu]}\|_{\dot{W}^{\sigma_t, p}(0, T; \dot{W}^{\sigma_x, p}(\mathbb{R}^d))}^p = \eta^{\mu p} \gamma^{\sigma_t p - 1} \|u^{[\mu]}\|_{\dot{W}^{\sigma_t, p}(0, \gamma T; \dot{W}^{\sigma_x, p}(\mathbb{R}^d))}^p,$$

as well as $\|\tilde{u}_0\|_{L^1(\mathbb{R}^d)} = \eta \|u_0\|_{L^1(\mathbb{R}^d)}$ and $\|\tilde{S}\|_{L^1(0, T; L^1(\mathbb{R}^d))} = \eta \|S\|_{L^1(0, \gamma T; L^1(\mathbb{R}^d))}$. Thus, it follows from (3-4) that

$$\begin{aligned} \|u^{[\mu]}\|_{\dot{W}^{\sigma_t, p}(0, T; \dot{W}^{\sigma_x, p}(\mathbb{R}^d))}^p &\leq c \eta^{1-\mu p} \gamma^{1-\sigma_t p} (\|u_0\|_{L^1(\mathbb{R}^d)} + \|S\|_{L^1(0, \gamma T; L^1(\mathbb{R}^d))}) \\ &= c \eta^{(m-1)(1-\sigma_t p) + 1 - \mu p} (\|u_0\|_{L^1(\mathbb{R}^d)} + \|S\|_{L^1(0, T; L^1(\mathbb{R}^d))}). \end{aligned} \quad (3-5)$$

As long as u_0 or S are nontrivial and unless

$$(m-1)(1-\sigma_t p) + 1 - \mu p \geq 0, \quad (3-6)$$

this gives the contradiction $u = 0$ by sending $\eta \rightarrow \infty$ (and consequently also $\gamma \rightarrow \infty$). Since $\sigma_t \geq 0$, (3-6) gives

$$p \leq \frac{m}{\mu + (m-1)\sigma_t} \leq \frac{m}{\mu}.$$

By the same token, since $p \geq 1$, (3-6) gives

$$\sigma_t \leq \frac{m - \mu p}{p(m-1)} \leq \frac{m - \mu}{m-1}.$$

Next, we rescale in space. More precisely, for positive constants $\eta, \gamma > 0$ with $\eta^{1-m} = \gamma^2$ and a fixed triple (u, u_0, S) as above we consider the rescaled quantities $(\tilde{u}, \tilde{u}_0, \tilde{S})$ defined via

$$\tilde{u}(t, x) := \eta u(t, \gamma x), \quad \tilde{u}_0(x) := \eta u_0(\gamma x), \quad \tilde{S}(t, x) := \eta S(t, \gamma x).$$

Then \tilde{u} satisfies (1-1) with $\tilde{u}_0 \in L^1(\mathbb{R}^d)$ and $\tilde{S} \in L^1(0, T; L^1(\mathbb{R}^d))$, so that (3-2) gives

$$\|\tilde{u}^{[\mu]}\|_{\dot{W}^{\sigma_t, p}(0, T; \dot{W}^{\sigma_x, p}(\mathbb{R}^d))}^p \leq c(\|\tilde{u}_0\|_{L^1(\mathbb{R}^d)} + \|\tilde{S}\|_{L^1(0, T; L^1(\mathbb{R}^d))}). \quad (3-7)$$

We have

$$\|\tilde{u}^{[\mu]}\|_{\dot{W}^{\sigma_t, p}(0, T; \dot{W}^{\sigma_x, p}(\mathbb{R}^d))}^p = \eta^{\mu p} \gamma^{\sigma_x p - d} \|u^{[\mu]}\|_{\dot{W}^{\sigma_t, p}(0, T; \dot{W}^{\sigma_x, p}(\mathbb{R}^d))}^p,$$

as well as $\|\tilde{u}_0\|_{L^1(\mathbb{R}^d)} = \eta \gamma^{-d} \|u_0\|_{L^1(\mathbb{R}^d)}$ and $\|\tilde{S}\|_{L^1(0, T; L^1(\mathbb{R}^d))} = \eta \gamma^{-d} \|S\|_{L^1(0, T; L^1(\mathbb{R}^d))}$. Thus, it follows from (3-7) and the relation $\eta^{1-m} = \gamma^2$ that

$$\begin{aligned} \|u^{[\mu]}\|_{\dot{W}^{\sigma_t, p}(0, T; \dot{W}^{\sigma_x, p}(\mathbb{R}^d))}^p &\leq c \eta^{1-\mu p} \gamma^{-\sigma_x p} (\|u_0\|_{L^1(\mathbb{R}^d)} + \|S\|_{L^1(0, T; L^1(\mathbb{R}^d))}) \\ &= c \eta^{\frac{\sigma_x p(m-1)}{2} + 1 - \mu p} (\|u_0\|_{L^1(\mathbb{R}^d)} + \|S\|_{L^1(0, T; L^1(\mathbb{R}^d))}). \end{aligned} \quad (3-8)$$

As long as u_0 or S are nontrivial and unless

$$\frac{\sigma_x p(m-1)}{2} + 1 - \mu p = 0 \quad \Longleftrightarrow \quad \sigma_x = \frac{\mu p - 1}{p} \frac{2}{m-1}, \quad (3-9)$$

this gives the contradiction $u = 0$ by sending $\eta \rightarrow 0$ or $\eta \rightarrow \infty$ (and consequently $\gamma \rightarrow \infty$ or $\gamma \rightarrow 0$, respectively). Plugging into (3-9) the restrictions on p and σ_t , we obtain the result. \square

Remark 3.2. If one sets $\mu = 1$, $p = 1$ and $\sigma_t = 0$, Lemma 3.1 tells us that σ_x cannot be positive, which is what we claimed following (3-1). Moreover, we emphasize that Lemma 3.1 shows that in the case of the whole space, the regularity exponent $\sigma_x \in [2(\mu - 1)/(m - 1), 2\mu/m]$ is in a one-to-one correspondence to the integrability exponent $p \in [1, m/\mu]$ via

$$\sigma_x = \frac{\mu p - 1}{p} \frac{2}{m-1} \quad \text{and} \quad p = \frac{2}{2\mu - \sigma_x(m-1)}.$$

The Barenblatt solution. Consider the Barenblatt solution

$$u_{BB}(t, x) := t^{-\alpha} (C - k|x t^{-\beta}|^2)_+^{\frac{1}{m-1}},$$

where

$$m > 1, \quad \alpha := \frac{d}{d(m-1) + 2}, \quad k = \frac{\alpha(m-1)}{2md}, \quad \beta = \frac{\alpha}{d},$$

and $C > 0$ is a free constant. Then, for $\mu \in [1, m]$, $u_{BB}^{[\mu]} \in L^{m/\mu}(0, T; \dot{W}^{s, m/\mu}(\mathbb{R}^d))$ implies $s < 2\mu/m$.

Proof. With $F(x) := (C - k|x|^2)_+^{\mu/(m-1)}$ we have $u_{BB}^{[\mu]}(t, x) = t^{-\alpha\mu} F(x t^{-\beta})$. We next observe that, for $s \in (0, 1)$ and each $t \geq 0$,

$$\begin{aligned} \|u_{BB}^{[\mu]}(t, \cdot)\|_{\dot{W}^{s, m/\mu}(\mathbb{R}^d)}^{\frac{m}{\mu}} &= \int_{\mathbb{R}^d \times \mathbb{R}^d} \frac{|u_{BB}^{[\mu]}(t, x) - u_{BB}^{[\mu]}(t, y)|^{\frac{m}{\mu}}}{|x - y|^{\frac{sm}{\mu} + d}} dx dy \\ &= t^{-\alpha m - \beta(\frac{sm}{\mu} + d) + 2d\beta} \|F\|_{\dot{W}^{s, m/\mu}(\mathbb{R}^d)}^{\frac{m}{\mu}}. \end{aligned}$$

Hence,

$$\|u_{BB}^{[\mu]}\|_{L^{m/\mu}(0, T; \dot{W}^{s, m/\mu}(\mathbb{R}^d))}^{\frac{m}{\mu}} = \|t^{-\alpha m - \beta(\frac{sm}{\mu} + d) + 2d\beta}\|_{L^1(0, T)} \|F\|_{\dot{W}^{s, m/\mu}(\mathbb{R}^d)}^{\frac{m}{\mu}},$$

which is finite if and only if

$$-\alpha m - \beta \left(\frac{sm}{\mu} + d \right) + 2d\beta > -1 \quad \text{and} \quad F \in \dot{W}^{s, \frac{m}{\mu}}(\mathbb{R}^d).$$

Hence, necessarily

$$m + \frac{1}{d} \left(\frac{sm}{\mu} + d \right) - 2 < \frac{1}{\alpha} = \frac{d(m-1)+2}{d},$$

which is equivalent to $s < 2\mu/m$. In the case $s \in (1, 2)$ we observe that it holds

$$\partial_{x_i} u_{BB}^{[\mu]}(t, x) = t^{-\alpha\mu+\beta} \partial_{x_i} F(xt^{-\beta}),$$

so that analogous arguments may be applied. □

4. Averaging lemma approach

In [Gess 2020], an averaging lemma was introduced that can be applied directly to the porous medium equations (1-1) to obtain estimates on the spatial regularity of u , but so far, no corresponding estimates for powers of the solution u^μ or its time regularity could be obtained. In this section, we provide an averaging lemma that gives a comprehensive answer to both of these questions. To this end, we recall the definition of the anisotropic and isotropic truncation properties from [Gess 2020], which extend the truncation property introduced in [Tadmor and Tao 2007, Definition 2.1].

Definition 4.1. (i) Let m be a complex-valued Fourier multiplier. We say that m has the truncation property if, for any locally supported bump function ψ on \mathbb{C} and any $1 \leq p < \infty$, the multiplier with symbol $\psi(m(\xi)/\delta)$ is an L^p -multiplier as well as an \mathcal{M}_{TV} -multiplier uniformly in $\delta > 0$, that is, its L^p -multiplier norm (\mathcal{M}_{TV} -multiplier norm resp.) depends only on the support and C^l size of ψ (for some large l that may depend on m) but otherwise is independent of δ .

(ii) Let $m : \mathbb{R}_\xi^d \times \mathbb{R}_v \rightarrow \mathbb{C}$ be a Carathéodory function such that $m(\cdot, v)$ is radial for all $v \in \mathbb{R}$. Then m is said to satisfy the isotropic truncation property if, for every bump function ψ supported on a ball in \mathbb{C} , every bump function φ supported in $\{\xi \in \mathbb{C} : \frac{1}{2} \leq |\xi| \leq 2\}$ and every $1 < p < \infty$,

$$M_{\psi, J} f(x, v) := \mathcal{F}_x^{-1} \varphi \left(\frac{|\xi|^2}{J^2} \right) \psi \left(\frac{m(\xi, v)}{\delta} \right) \mathcal{F}_x f(x)$$

is an L_x^p -multiplier for all $v \in \mathbb{R}$, $J = 2^j$, $j \in \mathbb{Z}$, and, for all $r \geq 1$,

$$\| \| M_{\psi, J} \|_{\mathcal{M}^p} \|_{L_v^r} \lesssim |\Omega_m(J, \delta)|^{\frac{1}{r}},$$

where

$$\Omega_m(J, \delta) := \left\{ v \in \mathbb{R} : \left| \frac{m(J, v)}{\delta} \right| \in \text{supp } \psi \right\}.$$

Here we use an abuse of notation

$$\left| \frac{m(J, v)}{\delta} \right| := \sup \left\{ \left| \frac{m(\xi, v)}{\delta} \right| : \frac{|\xi|^2}{J^2} \in \text{supp } \varphi \right\}.$$

We recall that for $m(\xi, v) := |\xi|^2|v|$, the anisotropic truncation property is satisfied uniformly in v by Example A.2 in [Gess 2020] and the isotropic truncation property is satisfied by Example 3.2 in [Gess 2020], albeit only in the case $J \geq 1$. However, the proof given there can be used without any changes to obtain the full assertion for general $J \in \mathbb{Z}$.

Lemma 4.2. Assume $m \in (1, \infty)$, $\gamma \in (-\infty, m)$, $\mu \in [1, m+1-\gamma]$ and let $f \in L_{t,x,v}^\beta$, where $\beta' = 1/\rho$ with $\rho \in (0, 1)$, be a solution to

$$\mathcal{L}(\partial_t, \nabla_x, v)f(t, x, v) = g_0(t, x, v) + \partial_v g_1(t, x, v) \text{ on } \mathbb{R}_t \times \mathbb{R}_x^d \times \mathbb{R}_v. \quad (4-1)$$

Here, the differential operator $\mathcal{L}(\partial_t, \nabla_x, v)$ that is given in terms of its symbol

$$\mathcal{L}(i\tau, i\xi, v) := i\tau + |v|^{m-1}|\xi|^2, \quad (4-2)$$

and g_i are Radon measures satisfying

$$|g_0|(t, x, v)|v|^{1-\gamma} + |g_1|(t, x, v)|v|^{-\gamma} \in \mathcal{M}_{TV}(\mathbb{R}_t \times \mathbb{R}_x^d \times \mathbb{R}_v).$$

Suppose

$$s \in \left(\frac{\mu-2+\gamma}{m-1}, 1 \right] \cap [0, 1].$$

Then $\bar{f} \in S_{p,\infty,(\infty)}^{\bar{\kappa}} \dot{B}$, where $\bar{f}(t, x) := \int f(t, x, v)|v|^{\mu-1} dv$, $\bar{\kappa} := (\kappa_t, \kappa_x)$ and

$$p := \frac{s(m-1)+1-\gamma+\rho}{\rho\mu+(1-\rho)(s(m-1)+1-\gamma)}, \quad \kappa_t := \frac{(1-s)(\mu-1+\rho)}{s(m-1)+1-\gamma+\rho}, \quad \kappa_x := \frac{2s(\mu-1+\rho)}{s(m-1)+1-\gamma+\rho}. \quad (4-3)$$

Moreover, we have the estimate

$$\|\bar{f}\|_{S_{p,\infty,(\infty)}^{\bar{\kappa}} \dot{B}} \lesssim \| |v|^{1-\gamma} g_0 \|_{\mathcal{M}_{TV}} + \| |v|^{-\gamma} g_1 \|_{\mathcal{M}_{TV}} + \| f \|_{L_{t,x,v}^\beta}. \quad (4-4)$$

If additionally $\bar{f} \in L_{t,x}^r$, $p \neq r \in [1, \infty]$, then for all $q \in (\min\{p, r\}, \max\{p, r\})$ it holds $\bar{f} \in S_{q,\infty}^{\vartheta\bar{\kappa}} \dot{B}$, where $\vartheta \in (0, 1)$ is such that

$$\frac{1}{q} = \frac{1-\vartheta}{r} + \frac{\vartheta}{p}.$$

In this case we have

$$\|\bar{f}\|_{S_{q,\infty}^{\vartheta\bar{\kappa}} \dot{B}} \lesssim \| |v|^{1-\gamma} g_0 \|_{\mathcal{M}_{TV}} + \| |v|^{-\gamma} g_1 \|_{\mathcal{M}_{TV}} + \| f \|_{L_{t,x,v}^\beta} + \|\bar{f}\|_{L_{t,x}^r}. \quad (4-5)$$

Finally, if $s = 1$ and consequently $\kappa_t = 0$, then (4-5) remains true if we replace the space $S_{q,\infty}^{\vartheta\bar{\kappa}} \dot{B} = S_{q,\infty}^{(0,\vartheta\kappa_x)} \dot{B}$ by $\tilde{L}_t^q \dot{B}_{q,\infty}^{\vartheta\kappa_x}$.

Remark 4.3. Observe that for

$$\rho \in \left(\frac{m+1-\gamma-\mu}{m+1-\gamma}, 1 \right)$$

one may prescribe a specific integrability exponent. More precisely, given

$$\tilde{p} \in \left[\frac{1-\gamma+\rho}{\rho\mu+(1-\rho)(1-\gamma)}, \frac{m+1-\gamma}{\mu} \right] \cap \left(1, \frac{m+1-\gamma}{\mu} \right]$$

choose

$$s := \frac{\mu \tilde{p}\rho + \tilde{p}(1-\rho)(1-\gamma) - 1 + \gamma + \rho}{(m-1)(1-\tilde{p}(1-\rho))} \in \left(\frac{\mu-2+\gamma}{m-1}, 1 \right] \cap [0, 1].$$

Then (4-3) reads $p = \tilde{p}$, as well as

$$\begin{aligned} \kappa_t &:= \frac{m + \rho - \gamma - \mu p\rho + p(1-\rho)(\gamma-m)}{p\rho} \frac{1}{m-1}, \\ \kappa_x &:= \frac{\mu p\rho + p(1-\rho)(1-\gamma) - 1 + \gamma - \rho}{p\rho} \frac{2}{m-1}. \end{aligned}$$

Observe that in the limiting case $\rho \rightarrow 1$ and $\gamma \rightarrow 1$, these orders of differentiability correspond to the ones found in (3-3).

Proof of Lemma 4.2. We first assume that f is compactly supported with respect to the variable v . This condition will enter only qualitatively, and never appears in quantitative form. Therefore, at the end of the proof, we can again remove this additional assumption.

Since we are interested in regularity in terms of homogeneous Besov spaces, we decompose f into Littlewood–Paley blocks with respect to the t -variable and the x -variable. Let $\{\eta_l\}_{l \in \mathbb{Z}}$ be a partition of unity on $\mathbb{R} \setminus \{0\}$ and $\{\varphi_j\}_{j \in \mathbb{Z}}$ a partition of unity on $\mathbb{R}^d \setminus \{0\}$ as in Section 2. Then we define for $l, j \in \mathbb{Z}$

$$f_{l,j} := \mathcal{F}_{t,x}^{-1}[\eta_l \varphi_j \mathcal{F}_{t,x} f],$$

where $\mathcal{F}_{t,x} f_{l,j}(\tau, \xi, v)$ is supported on frequencies $|\xi| \sim 2^j$, $|\tau| \sim 2^l$ for $l, j \in \mathbb{Z}$. Similarly, we define the decompositions $g_{0,l,j}$ and $g_{1,l,j}$ of g_0 and g_1 , respectively. We consider a microlocal decomposition of $f_{l,j}$ connected to the degeneracy of the operator $\mathcal{L}(\partial_t, \nabla_x, v)$. Let $\psi_0 \in C_c^\infty(\mathbb{R})$ be a smooth function supported in $B_2(0)$ and set $\psi_1 := 1 - \psi_0$. For $\delta > 0$ to be specified later we write

$$f_{l,j} = \mathcal{F}_x^{-1} \psi_0 \left(\frac{|v||\xi|^2}{\delta} \right) \mathcal{F}_x f_{l,j} + \mathcal{F}_x^{-1} \psi_1 \left(\frac{|v||\xi|^2}{\delta} \right) \mathcal{F}_x f_{l,j} =: f_{l,j}^0 + f_{l,j}^1.$$

Since f is a solution to (4-1), we have

$$\mathcal{F}_{t,x}^{-1} \mathcal{L}(i\tau, i\xi, v) \mathcal{F}_{t,x} f_{l,j}^1(t, x, v) = \mathcal{F}_x^{-1} \psi_1 \left(\frac{|v||\xi|^2}{\delta} \right) \mathcal{F}_x \left(g_{0,l,j}(t, x, v) + \partial_v g_{1,l,j}(t, x, v) \right) \quad (4-6)$$

and thus

$$\begin{aligned} f_{l,j}^1(t, x, v) &= \mathcal{F}_{t,x}^{-1} \psi_1 \left(\frac{|v||\xi|^2}{\delta} \right) \frac{1}{\mathcal{L}(i\tau, i\xi, v)} \mathcal{F}_{t,x} g_{0,l,j}(t, x, v) \\ &\quad + \mathcal{F}_{t,x}^{-1} \psi_1 \left(\frac{|v||\xi|^2}{\delta} \right) \frac{1}{\mathcal{L}(i\tau, i\xi, v)} \mathcal{F}_{t,x} \partial_v g_{1,l,j}(t, x, v) \\ &=: f_{l,j}^2(t, x, v) + f_{l,j}^3(t, x, v). \end{aligned} \quad (4-7)$$

In conclusion, we have arrived at the decomposition

$$\bar{f}_{l,j} := \int f_{l,j} |v|^{\mu-1} dv = \int f_{l,j}^0 |v|^{\mu-1} dv + \int f_{l,j}^2 |v|^{\mu-1} dv + \int f_{l,j}^3 |v|^{\mu-1} dv =: \bar{f}_{l,j}^0 + \bar{f}_{l,j}^2 + \bar{f}_{l,j}^3.$$

We aim to estimate the regularity of these three contributions separately.

Step 1: f^0 . We note that we have the estimate $\|\mathcal{F}_t^{-1} \eta_l \mathcal{F} f\|_{L_{t,x}^\beta} \lesssim \|f\|_{L_{t,x}^\beta}$ with a constant independent of l , since $\|\eta_l\|_{\mathcal{M}^\beta} = \|\eta_0\|_{\mathcal{M}^\beta} < \infty$. Let $l, j \in \mathbb{Z}$ be arbitrary, fixed. Then, we have that $|v| \leq 2 \cdot 2^{-2j} \delta$ on the support of $\varphi(2^{-j} \xi) \psi_0(|v| |\xi|^2 / \delta)$, so that $|\Omega_m(2^j, \delta)| \lesssim [-2 \cdot 2^{-2j} \delta, 2 \cdot 2^{-2j} \delta] \lesssim 2^{-2j} \delta$. Hence, by the isotropic truncation property and Minkowski's and Hölder's inequality it holds

$$\begin{aligned} \left\| \int f_{l,j}^0 |v|^{\mu-1} dv \right\|_{L_{t,x}^\beta} &= \left\| \int \mathcal{F}_x^{-1} \psi_0 \left(\frac{|v| |\xi|^2}{\delta} \right) |v|^{\mu-1} \mathcal{F}_x f_{l,j} dv \right\|_{L_{t,x}^\beta} \\ &\lesssim \int \left\| \mathcal{F}_x^{-1} \psi_0 \left(\frac{|v| |\xi|^2}{\delta} \right) |v|^{\mu-1} \mathcal{F}_x f_{l,j} \right\|_{L_{t,x}^\beta} dv \\ &\lesssim \left(\frac{\delta}{2^{2j}} \right)^{\mu-1} \int \left\| \mathcal{F}_x^{-1} \psi_0 \left(\frac{|v| |\xi|^2}{\delta} \right) \mathcal{F}_x f_{l,j} \right\|_{L_{t,x}^\beta} dv \\ &\lesssim \left(\frac{\delta}{2^{2j}} \right)^{\mu-1} \int \|M_{\psi_0, 2^{-j}}\|_{\mathcal{M}^\beta} \|f\|_{L_{t,x}^\beta} dv \\ &\leq \left(\frac{\delta}{2^{2j}} \right)^{\mu-1} \|M_{\psi_0, 2^{-j}}\|_{\mathcal{M}^\beta} \|f\|_{L_{t,x,v}^\beta} \\ &\lesssim \left(\frac{\delta}{2^{2j}} \right)^{\mu-1} |\Omega_m(2^j, \delta)|^{\frac{1}{\beta'}} \|f\|_{L_{t,x,v}^\beta} \lesssim \left(\frac{\delta}{2^{2j}} \right)^{\mu-1+\rho} \|f\|_{L_{t,x,v}^\beta}, \end{aligned}$$

where we have used $\beta' = 1/\rho$.

Step 2: f^2 . Let $l, j \in \mathbb{Z}$ be arbitrary, fixed. Since $s \in [0, 1]$, we clearly have

$$|\tau|^{1-s} |v|^{s(m-1)} |\xi|^{2s} \leq |\mathcal{L}(i\tau, i\xi, v)|.$$

Moreover, in light of $s > \mu - 2 + \gamma/(m-1)$ we have on the support of $\eta_l \varphi_j \psi_1(|v| |\xi|^2 / \delta)$ (so that $|\tau| \sim 2^l$, $|\xi| \sim 2^j$, and $|v| \gtrsim 2^{-2j} \delta$)

$$\frac{|v|^{\mu-2+\gamma}}{|\mathcal{L}(i\tau, i\xi, v)|} \lesssim \frac{|v|^{\mu-2+\gamma}}{|\tau|^{1-s} |v|^{s(m-1)} |\xi|^{2s}} \lesssim \frac{(2^{-2j} \delta)^{\mu-2+\gamma-s(m-1)}}{2^{l(1-s)} 2^{2js}} = \frac{2^{2j(s(m-2)-\mu+2-\gamma)}}{\delta^{s(m-1)-\mu+2-\gamma} 2^{l(1-s)}}.$$

Hence, by [Theorem B.1](#) and [Lemma B.4](#), $|v|^{\mu-2+\gamma} / \mathcal{L}(i\tau, i\xi, v)$ acts on the support of $\eta_l \varphi_j \psi_1(|v| |\xi|^2 / \delta)$ as a constant multiplier of order

$$\frac{2^{2j(s(m-2)-\mu+2-\gamma)}}{\delta^{s(m-1)-\mu+2-\gamma} 2^{l(1-s)}}.$$

Consequently, by the anisotropic truncation property

$$\begin{aligned} \left\| \int f_{l,j}^2 |v|^{\mu-1} dv \right\|_{L_{t,x}^1} &= \left\| \int \mathcal{F}_{t,x}^{-1} \psi_1 \left(\frac{|v| |\xi|^2}{\delta} \right) \frac{|v|^{\mu-2+\gamma}}{\mathcal{L}(i\tau, i\xi, v)} \mathcal{F}_{t,x} |v|^{1-\gamma} g_{0,l,j} dv \right\|_{L_{t,x}^1} \\ &\lesssim \frac{2^{2j(s(m-2)-\mu+2-\gamma)}}{\delta^{s(m-1)-\mu+2-\gamma} 2^{l(1-s)}} \| |v|^{1-\gamma} g_0 \|_{\mathcal{M}_{TV}}. \end{aligned}$$

Here, we have used that with $\psi_0(|v||\xi|^2/\delta)$ also

$$\psi_1\left(\frac{|v||\xi|^2}{\delta}\right) = 1 - \psi_0\left(\frac{|v||\xi|^2}{\delta}\right)$$

is a bounded \mathcal{M}_{TV} -multiplier independent of $\delta > 0$.

Step 3: f^3 . Let $l, j \in \mathbb{Z}$ arbitrary, fixed. We observe (recall $\mathcal{L}(i\tau, i\xi, v) = i\tau + |v|^{m-1}|\xi|^2$)

$$\begin{aligned} \int f_{l,j}^3 |v|^{\mu-1} dv &= - \int \mathcal{F}_{t,x}^{-1} \psi_1' \left(\frac{|v||\xi|^2}{\delta} \right) \frac{\operatorname{sgn}(v)|\xi|^2}{\delta} \frac{|v|^{\mu-1}}{\mathcal{L}(i\tau, i\xi, v)} \mathcal{F}_{t,x} g_{1,l,j} dv \\ &\quad - (\mu-1) \int \mathcal{F}_{t,x}^{-1} \psi_1 \left(\frac{|v||\xi|^2}{\delta} \right) \frac{\operatorname{sgn}(v)|v|^{\mu-2}}{\mathcal{L}(i\tau, i\xi, v)} \mathcal{F}_{t,x} g_{1,l,j} dv \\ &\quad + \int \mathcal{F}_{t,x}^{-1} \psi_1 \left(\frac{|v||\xi|^2}{\delta} \right) \frac{|v|^{\mu-1} \partial_v \mathcal{L}(i\tau, i\xi, v)}{\mathcal{L}(i\tau, i\xi, v)^2} \mathcal{F}_{t,x} g_{1,l,j} dv \\ &= - \int \mathcal{F}_{t,x}^{-1} \psi_1' \left(\frac{|v||\xi|^2}{\delta} \right) \frac{|v||\xi|^2}{\delta} \frac{\operatorname{sgn}(v)|v|^{\mu-2+\gamma}}{\mathcal{L}(i\tau, i\xi, v)} \mathcal{F}_{t,x} |v|^{-\gamma} g_{1,l,j} dv \\ &\quad - (\mu-1) \int \mathcal{F}_{t,x}^{-1} \psi_1 \left(\frac{|v||\xi|^2}{\delta} \right) \frac{\operatorname{sgn}(v)|v|^{\mu-2+\gamma}}{\mathcal{L}(i\tau, i\xi, v)} \mathcal{F}_{t,x} |v|^{-\gamma} g_{1,l,j} dv \\ &\quad + (m-1) \int \mathcal{F}_{t,x}^{-1} \psi_1 \left(\frac{|v||\xi|^2}{\delta} \right) \frac{|v|^{\mu+m-3+\gamma} |\xi|^2}{\mathcal{L}(i\tau, i\xi, v)^2} \mathcal{F}_{t,x} |v|^{-\gamma} g_{1,l,j} dv. \end{aligned}$$

Observe that ψ_1' is supported on an annulus. Therefore, we have as before $|\tau| \sim 2^l$, $|\xi| \sim 2^j$ and $|v| \gtrsim 2^{-2j}\delta$ on the support of $\eta_l \varphi_j \psi_1(|v||\xi|^2/\delta)$, and additionally also $|v| \sim 2^{-2j}\delta$ on the support of $\eta_l \varphi_j \psi_1'(|v||\xi|^2/\delta)$. This last observation allows us to estimate the expression $|v||\xi|^2/\delta$ appearing in the first integral on the right-hand side by

$$\frac{|v||\xi|^2}{\delta} \lesssim 1.$$

As in Step 2, we obtain

$$\frac{|v|^{\mu-2+\gamma}}{|\mathcal{L}(i\tau, i\xi, v)|} \lesssim \frac{2^{2j(s(m-2)-\mu+2-\gamma)}}{\delta^{s(m-1)-\mu+2-\gamma} 2^{l(1-s)}},$$

and, similarly,

$$\begin{aligned} \frac{|v|^{\mu+m-3+\gamma} |\xi|^2}{|\mathcal{L}(i\tau, i\xi, v)|^2} &= \frac{|v|^{\mu-2+\gamma}}{|\mathcal{L}(i\tau, i\xi, v)|} \frac{|v|^{m-1} |\xi|^2}{|\mathcal{L}(i\tau, i\xi, v)|} \\ &\lesssim \frac{|v|^{\mu-2+\gamma}}{|\mathcal{L}(i\tau, i\xi, v)|} \lesssim \frac{2^{2j(s(m-2)-\mu+2-\gamma)}}{\delta^{s(m-1)-\mu+2-\gamma} 2^{l(1-s)}}. \end{aligned}$$

In light of these estimates, the expressions

$$\frac{|v||\xi|^2}{\delta} \frac{\operatorname{sgn}(v)|v|^{\mu-2+\gamma}}{\mathcal{L}(i\tau, i\xi, v)}, \quad \frac{\operatorname{sgn}(v)|v|^{\mu-2+\gamma}}{\mathcal{L}(i\tau, i\xi, v)}, \quad \frac{|v|^{\mu+m-3+\gamma} |\xi|^2}{\mathcal{L}(i\tau, i\xi, v)^2}$$

extend by [Theorem B.1](#) and [Lemma B.4](#) to constant multipliers of order

$$\frac{2^{2j(s(m-2)-\mu+2-\gamma)}}{\delta^{s(m-1)-\mu+2-\gamma} 2^{l(1-s)}}$$

on supports of $\eta_l \varphi_j \psi'_1(|v| |\xi|^2 / \delta)$ and $\eta_l \varphi_j \psi_1(|v| |\xi|^2 / \delta)$, respectively. Hence, by the anisotropic truncation property, we obtain

$$\left\| \int f_{l,j}^3 |v|^{\mu-1} dv \right\|_{L_{t,x}^1} \lesssim \frac{2^{2j(s(m-2)-\mu+2-\gamma)}}{\delta^{s(m-1)-\mu+2-\gamma} 2^{l(1-s)}} \| |v|^{-\gamma} g_{1,j} \|_{\mathcal{M}_{TV}}.$$

Step 4: Conclusion. We aim to conclude by real interpolation. We set, for $z > 0$,

$$K(z, \bar{f}_{l,j}) := \inf \{ \| \bar{f}_{l,j}^1 \|_{L_{t,x}^1} + z \| \bar{f}_{l,j}^0 \|_{L_{t,x}^\beta} : \bar{f}_{l,j}^0 \in L_{t,x}^\beta, \bar{f}_{l,j}^1 \in L_{t,x}^1, \bar{f}_{l,j} = \bar{f}_{l,j}^0 + \bar{f}_{l,j}^1 \}.$$

By the above estimates we obtain

$$K(z, \bar{f}_{l,j}) \lesssim \frac{2^{2j(s(m-2)-\mu+2-\gamma)}}{\delta^{s(m-1)-\mu+2-\gamma} 2^{l(1-s)}} (\| |v|^{1-\gamma} g_0 \|_{\mathcal{M}_{TV}} + \| |v|^{-\gamma} g_1 \|_{\mathcal{M}_{TV}}) + z \left(\frac{\delta}{2^{2j}} \right)^{\mu-1+\rho} \| f \|_{L_{t,x,v}^\beta}.$$

We now equilibrate the first and the second term on the right-hand side: we choose $\delta > 0$ such that

$$\frac{2^{2j(s(m-2)-\mu+2-\gamma)}}{\delta^{s(m-1)-\mu+2-\gamma} 2^{l(1-s)}} = z \left(\frac{\delta}{2^{2j}} \right)^{\mu-1+\rho};$$

that is,

$$\delta^{-a} c^{1-s} d^{-a+s} = z \delta^b d^b,$$

with $a := s(m-1) - \mu + 2 - \gamma$, $b := \mu - 1 + \rho$, $c := 2^{-l}$ and $d := 2^{-2j}$. This yields

$$\delta = z^{-\frac{1}{a+b}} c^{\frac{1-s}{a+b}} d^{\frac{s-a-b}{a+b}},$$

and further

$$\delta^{-a} c^{1-s} d^{-a+s} = z^{\frac{a}{a+b}} c^{\frac{(1-s)b}{a+b}} d^{\frac{sb}{a+b}}.$$

Hence, with

$$\theta := \frac{a}{a+b} = \frac{s(m-1) - \mu + 2 - \gamma}{s(m-1) + 1 - \gamma + \rho}$$

we obtain

$$\begin{aligned} z^{-\theta} K(z, \bar{f}_{l,j}) &\lesssim 2^{-l} \frac{(1-s)(\mu-1+\rho)}{s(m-1)+1-\gamma+\rho} 2^{-2j} \frac{s(\mu-1+\rho)}{s(m-1)+1-\gamma+\rho} (\| |v|^{1-\gamma} g_0 \|_{\mathcal{M}_{TV}} + \| |v|^{-\gamma} g_1 \|_{\mathcal{M}_{TV}} + \| f \|_{L_{t,x,v}^\beta}) \\ &= 2^{-l\kappa_t} 2^{-j\kappa_x} (\| |v|^{1-\gamma} g_0 \|_{\mathcal{M}_{TV}} + \| |v|^{-\gamma} g_1 \|_{\mathcal{M}_{TV}} + \| f \|_{L_{t,x,v}^\beta}). \end{aligned}$$

Observe that $1 - \theta + \theta/\beta = 1 - \theta + \theta(1 - \rho) = 1 - \theta\rho$, so that $(L_{t,x}^1, L_{t,x}^\beta)_{\theta,\infty} = L_{t,x}^{p,\infty}$ with

$$p = \frac{1}{1 - \theta\rho} = \frac{a+b}{a(1-\rho)+b} = \frac{s(m-1) + 1 - \gamma + \rho}{\rho\mu + (1-\rho)(s(m-1) + 1 - \gamma)}.$$

Hence, we may take the supremum over $z > 0$ to obtain

$$\| \bar{f}_{l,j} \|_{L_{t,x}^{p,\infty}} \lesssim 2^{-l\kappa_t} 2^{-j\kappa_x} (\| |v|^{1-\gamma} g_0 \|_{\mathcal{M}_{TV}} + \| |v|^{-\gamma} g_1 \|_{\mathcal{M}_{TV}} + \| f \|_{L_{t,x,v}^\beta}). \quad (4-8)$$

Multiplying by $2^{l\kappa_t} 2^{j\kappa_x}$ and taking the supremum over $j, l \in \mathbb{Z}$ yields (4-4).

If we assume additionally $\bar{f} \in L_{t,x}^r$, $r \neq p$, we choose for $q \in (\min\{p, r\}, \max\{p, r\})$ a corresponding $\vartheta \in (0, 1)$ subject to $1/q = (1 - \vartheta)/r + \vartheta/p$. Then using $(L_{t,x}^r, L_{t,x}^{p,\infty})_{\vartheta,q} = L_{t,x}^q$, together with (4-8), we obtain

$$\begin{aligned} \|\bar{f}_{l,j}\|_{L_{t,x}^q} &\lesssim \|\bar{f}_{l,j}\|_{L_{t,x}^r}^{1-\vartheta} \|\bar{f}_{l,j}\|_{L_{t,x}^{p,\infty}}^{\vartheta} \\ &\lesssim \|\bar{f}\|_{L_{t,x}^r}^{1-\vartheta} 2^{-l\vartheta\kappa_t} 2^{-j\vartheta\kappa_x} (\| |v|^{1-\gamma} g_0 \|_{\mathcal{M}_{TV}} + \| |v|^{-\gamma} g_1 \|_{\mathcal{M}_{TV}} + \|f\|_{L_{t,x,v}^{\infty}})^{\vartheta} \\ &\leq 2^{-l\vartheta\kappa_t} 2^{-j\vartheta\kappa_x} (\| |v|^{1-\gamma} g_0 \|_{\mathcal{M}_{TV}} + \| |v|^{-\gamma} g_1 \|_{\mathcal{M}_{TV}} + \|f\|_{L_{t,x,v}^{\beta}} + \|\bar{f}\|_{L_{t,x}^r}). \end{aligned}$$

Multiplying by $2^{l\vartheta\kappa_t} 2^{j\vartheta\kappa_x}$ and taking the supremum over $j, l \in \mathbb{Z}$ yields (4-5).

Finally we note that if $s = 1$ and consequently $\kappa_t = 0$, then the partition of unity $\{\eta_l\}_{l \in \mathbb{Z}}$ in the Fourier space connected to the time variable t is not necessary. Hence, if we set $\alpha_\tau = 0$ whenever Lemma B.4 is invoked and replace Theorem B.1 by its isotropic variant (see Remark B.3), we obtain

$$\|\bar{f}_j\|_{L_{t,x}^q} \lesssim 2^{-j\vartheta\kappa_x} (\| |v|^{1-\gamma} g_0 \|_{\mathcal{M}_{TV}} + \| |v|^{-\gamma} g_1 \|_{\mathcal{M}_{TV}} + \|f\|_{L_{t,x,v}^{\beta}} + \|\bar{f}\|_{L_{t,x}^r}),$$

which shows $\bar{f} \in \tilde{L}^q \dot{B}_{q,\infty}^{\vartheta\kappa_x}$.

It remains to consider the case when f is not localized in v . We observe that for a smooth cut-off function $\psi \in C_c^\infty(\mathbb{R})$, the function $(t, x, v) \rightarrow f(t, x, v)\psi(v) =: f^\psi(t, x, v)$ is a solution to

$$\mathcal{L}(\partial_t, \nabla_x, v) f^\psi(t, x, v) = g_0^\psi(t, x, v) + g_1^{\psi'}(t, x, v) + \partial_v g_1^\psi(t, x, v) \quad \text{on } \mathbb{R}_t \times \mathbb{R}_x^d \times \mathbb{R}_v,$$

where g_0^ψ , $g_1^{\psi'}$ and g_1^ψ are defined analogously. Hence, estimate (4-8) reads in this case

$$\|\bar{f}_{l,j}^\psi\|_{L_{t,x}^{p,\infty}} \leq 2^{-l\vartheta\kappa_t} 2^{-j\vartheta\kappa_x} (\| |v|^{1-\gamma} (g_0^\psi + g_1^{\psi'}) \|_{\mathcal{M}_{TV}} + \| |v|^{-\gamma} g_1^\psi \|_{\mathcal{M}_{TV}} + \|f^\psi\|_{L_{t,x,v}^{\beta}}).$$

Since $|v|^{-\gamma} g_1 \in \mathcal{M}_{TV}$ by assumption, there exists for $\varepsilon_n \downarrow 0$ a sequence $r_n \uparrow \infty$ such that

$$\int_{\mathbb{R}_t \times \mathbb{R}_x^d \times \mathbb{R}_v} \chi_{\{r_n \leq |v|\}} |v|^{-\gamma} g_1 \, dv \, dx \, dt \leq \varepsilon_n$$

for all $n \in \mathbb{N}$. For $n \in \mathbb{N}$ and a smooth cut-off function $\psi \in C_c^\infty(\mathbb{R})$ with $\psi = 1$ on $B_1(0)$ and $\text{supp } \psi \subset B_2(0)$, we define ψ_n via $\psi_n(v) := \psi(v/r_n)$. Hence ψ_n' is supported on $r_n \leq |v| \leq 2r_n$ and takes values in $[0, 1/r_n]$, so that we may estimate

$$\begin{aligned} \| |v|^{1-\gamma} g_1^{\psi_n'} \|_{\mathcal{M}_{TV}} &= \int_{\mathbb{R}_t \times \mathbb{R}_x^d \times \mathbb{R}_v} |\psi_n'(v)| |v| (|v|^{-\gamma} g_1) \, dv \, dx \, dt \\ &= \int_{\mathbb{R}_t \times \mathbb{R}_x^d \times \mathbb{R}_v} \chi_{\{r_n \leq |v| \leq 2r_n\}} |\psi_n'(v)| |v| (|v|^{-\gamma} g_1) \, dv \, dx \, dt \\ &\lesssim \int \chi_{r_n \leq |v| \leq 2r_n} |v|^{-\gamma} g_1 \, dv \leq \varepsilon_n. \end{aligned}$$

Thus, taking the limit $n \rightarrow \infty$ and using Fatou's lemma, we obtain (4-8) also for general f . Multiplying by $2^{l\vartheta\kappa_t} 2^{j\vartheta\kappa_x}$ and taking the supremum over $j, l \in \mathbb{Z}$, we may conclude as before. \square

Lemma 4.4. Assume $\gamma \in (-\infty, 1)$, $m \in (1, \infty)$, $\mu \in [1, 2 - \gamma)$, $\rho \in (0, 1]$, $\beta' = 1/\rho$, and let f, g_0, g_1 , and \bar{f} be as in Lemma 4.2. Define

$$p := \frac{1 - \gamma + \rho}{\rho\mu + (1 - \rho)(1 - \gamma)}, \quad \kappa_t := \frac{\mu - 1 + \rho}{1 - \gamma + \rho}. \quad (4-9)$$

If $\bar{f} \in L^r_{t,x}$, $p \neq r \in [1, \infty]$, then for all $q \in (\min\{p, r\}, \max\{p, r\})$ we have $\bar{f} \in \tilde{L}^q_x \dot{B}^{\vartheta, \kappa_t}_{q, \infty}$, where $\vartheta \in (0, 1)$ is such that

$$\frac{1}{q} = \frac{1 - \vartheta}{r} + \frac{\vartheta}{p}.$$

Moreover,

$$\|\bar{f}\|_{\tilde{L}^q_x \dot{B}^{\vartheta, \kappa_t}_{q, \infty}} \lesssim \| |v|^{1-\gamma} g_0 \|_{\mathcal{M}_{TV}} + \| |v|^{-\gamma} g_1 \|_{\mathcal{M}_{TV}} + \|f\|_{L^{\beta}_{t,x,v}} + \|\bar{f}\|_{L^r_{t,x}}. \quad (4-10)$$

Proof. By the same arguments as in the proof of Lemma 4.2, we may assume that f is localized in v . In fact, the whole proof of Lemma 4.4 is similar to the one of Lemma 4.2, with the modification that here we consider a microlocal decomposition of f depending on the size of v only and do not localize in the Fourier space connected to the spatial variable x . More precisely, let $\{\eta_l\}_{l \in \mathbb{Z}}$ be a partition of unity on $\mathbb{R} \setminus \{0\}$ as in Section 2. Then we define for $l \in \mathbb{Z}$

$$f_l := \mathcal{F}_x^{-1}[\eta_l \mathcal{F}_t f],$$

where $\mathcal{F}_t f_l(\tau, x, v)$ is supported on frequencies $|\tau| \sim 2^l$ for $l \in \mathbb{Z}$. Similarly, we define the decompositions $g_{0,l}$ and $g_{1,l}$ of g_0 and g_1 , respectively. Moreover, we again consider a smooth function $\psi_0 \in C_c^\infty(\mathbb{R})$ supported in $B_2(0)$ and set $\psi_1 := 1 - \psi_0$. For $\delta > 0$ to be specified later we write

$$f_l = \psi_0\left(\frac{|v|}{\delta}\right) f_l + \psi_1\left(\frac{|v|}{\delta}\right) f_l =: f_l^0 + f_l^1.$$

Since f is a solution to (4-1), we have

$$\mathcal{F}_{t,x}^{-1} \mathcal{L}(i\tau, i\xi, v) \mathcal{F}_{t,x} f_l^1(t, x, v) = \psi_1\left(\frac{|v|}{\delta}\right) (g_{0,l}(t, x, v) + \partial_v g_{1,l}(t, x, v))$$

and thus

$$\begin{aligned} f_l^1(t, x, v) &= \mathcal{F}_{t,x}^{-1} \psi_1\left(\frac{|v|}{\delta}\right) \frac{1}{\mathcal{L}(i\tau, i\xi, v)} \mathcal{F}_{t,x} g_{0,l}(t, x, v) + \mathcal{F}_{t,x}^{-1} \psi_1\left(\frac{|v|}{\delta}\right) \frac{1}{\mathcal{L}(i\tau, i\xi, v)} \mathcal{F}_{t,x} \partial_v g_{1,l}(t, x, v) \\ &=: f_l^2(t, x, v) + f_l^3(t, x, v), \end{aligned}$$

so that we arrive at the decomposition

$$\begin{aligned} \bar{f}_l &:= \int f_l |v|^{\mu-1} dv = \int f_l^0 |v|^{\mu-1} dv + \int f_l^2 |v|^{\mu-1} dv + \int f_l^3 |v|^{\mu-1} dv \\ &=: \bar{f}_l^0 + \bar{f}_l^2 + \bar{f}_l^3. \end{aligned}$$

Again, we treat the three contributions separately.

Step 1: f^0 . Let $l \in \mathbb{Z}$ be arbitrary, fixed. Since $|v| \lesssim \delta$ on the support of $\psi_0(|v|/\delta)$, using Minkowski's and Hölder's inequalities, we have

$$\begin{aligned} \left\| \int f_l^0 |v|^{\mu-1} dv \right\|_{L_{t,x}^\beta} &= \left\| \int \psi_0\left(\frac{|v|}{\delta}\right) |v|^{\mu-1} f_l dv \right\|_{L_{t,x}^\beta} \\ &\leq \int |\psi_0\left(\frac{|v|}{\delta}\right)| |v|^{\mu-1} \|f_l\|_{L_{t,x}^\beta} dv \\ &\lesssim \delta^{\mu-1} \int |\psi_0\left(\frac{|v|}{\delta}\right)| \|f_l\|_{L_{t,x}^\beta} dv \\ &\lesssim \delta^{\mu-1} \|f\|_{L_{t,x,v}^\beta} \left(\int |\psi_0\left(\frac{|v|}{\delta}\right)|^{\beta'} dv \right)^{\frac{1}{\beta'}} \\ &\lesssim \delta^{\mu-1+\rho} \|f\|_{L_{t,x,v}^\beta}. \end{aligned}$$

Step 2: f^2 . Let $l \in \mathbb{Z}$ be arbitrary, fixed. Since $\mu \leq 2 - \gamma$, we have on the support of $\eta_l \psi_1(|v|/\delta)$ (so that $|\tau| \sim 2^l$ and $|v| \geq \delta$)

$$\frac{|v|^{\mu-2+\gamma}}{|\mathcal{L}(i\tau, i\xi, v)|} \lesssim \frac{|v|^{\mu-2+\gamma}}{|\tau|} \lesssim \frac{\delta^{\mu-2+\gamma}}{2^l}.$$

By [Lemma B.4](#) applied with $\alpha_\xi = 0$ and the isotropic variant of [Theorem B.1](#) (see [Remark B.3](#)), $|v|^{\mu-2+\gamma}/|\mathcal{L}(i\tau, i\xi, v)|$ acts as a constant multiplier of order $\delta^{\mu-2+\gamma}/2^l$ on the support of $\eta_l \psi_1(|v|/\delta)$. Consequently

$$\begin{aligned} \left\| \int f_l^2 |v|^{\mu-1} dv \right\|_{L_{t,x}^1} &= \left\| \int \mathcal{F}_{t,x}^{-1} \psi_1\left(\frac{|v|}{\delta}\right) \frac{|v|^{\mu-2+\gamma}}{\mathcal{L}(i\tau, i\xi, v)} \mathcal{F}_{t,x} |v|^{1-\gamma} g_{0,l} dv \right\|_{L_{t,x}^1} \\ &\lesssim \frac{\delta^{\mu-2+\gamma}}{2^l} \| |v|^{1-\gamma} g_0 \|_{\mathcal{M}_{TV}}. \end{aligned}$$

Step 3: f^3 . Let $l \in \mathbb{Z}$ be arbitrary, fixed. We observe (recall $\mathcal{L}(i\tau, i\xi, v) = i\tau + |v|^{m-1}|\xi|^2$)

$$\begin{aligned} \int f_l^3 |v|^{\mu-1} dv &= - \int \mathcal{F}_{t,x}^{-1} \psi_1' \left(\frac{|v|}{\delta} \right) \frac{\operatorname{sgn}(v)}{\delta} \frac{|v|^{\mu-1}}{\mathcal{L}(i\tau, i\xi, v)} \mathcal{F}_{t,x} g_{1,l} dv \\ &\quad - (\mu-1) \int \mathcal{F}_{t,x}^{-1} \psi_1 \left(\frac{|v|}{\delta} \right) \frac{\operatorname{sgn}(v)|v|^{\mu-2}}{\mathcal{L}(i\tau, i\xi, v)} \mathcal{F}_{t,x} g_{1,l} dv \\ &\quad + \int \mathcal{F}_{t,x}^{-1} \psi_1 \left(\frac{|v|}{\delta} \right) \frac{|v|^{\mu-1} \partial_v \mathcal{L}(i\tau, i\xi, v)}{\mathcal{L}(i\tau, i\xi, v)^2} \mathcal{F}_{t,x} g_{1,l} dv \\ &= - \int \mathcal{F}_{t,x}^{-1} \psi_1' \left(\frac{|v|}{\delta} \right) \frac{|v|}{\delta} \frac{\operatorname{sgn}(v)|v|^{\mu-2+\gamma}}{\mathcal{L}(i\tau, i\xi, v)} \mathcal{F}_{t,x} |v|^{-\gamma} g_{1,l} dv \\ &\quad - (\mu-1) \int \mathcal{F}_{t,x}^{-1} \psi_1 \left(\frac{|v|}{\delta} \right) \frac{\operatorname{sgn}(v)|v|^{\mu-2+\gamma}}{\mathcal{L}(i\tau, i\xi, v)} \mathcal{F}_{t,x} |v|^{-\gamma} g_{1,l} dv \\ &\quad + (m-1) \int \mathcal{F}_{t,x}^{-1} \psi_1 \left(\frac{|v|}{\delta} \right) \frac{|v|^{\mu+m-3+\gamma} |\xi|^2}{\mathcal{L}(i\tau, i\xi, v)^2} \mathcal{F}_{t,x} |v|^{-\gamma} g_{1,l} dv \end{aligned}$$

Observe that ψ'_1 is supported on an annulus. Therefore, we have as before $|\tau| \sim 2^l$ and $|v| \geq \delta$ on the support of $\eta_l \psi_1(|v|/\delta)$, and additionally also $|v| \sim \delta$ on the support of $\eta_l \psi'_1(|v|/\delta)$. This last observation allows us to estimate the expression $|v|/\delta$ appearing in the first integral on the right-hand side by $|v|/\delta \lesssim 1$. As in Step 2, we obtain

$$\frac{|v|^{\mu-2+\gamma}}{|\mathcal{L}(i\tau, i\xi, v)|} \lesssim \frac{\delta^{\mu-2+\gamma}}{2^l},$$

and, similarly,

$$\frac{|v|^{\mu+m-3+\gamma} |\xi|^2}{|\mathcal{L}(i\tau, i\xi, v)|^2} = \frac{|v|^{\mu-2+\gamma}}{|\mathcal{L}(i\tau, i\xi, v)|} \frac{|v|^{m-1} |\xi|^2}{|\mathcal{L}(i\tau, i\xi, v)|} \lesssim \frac{|v|^{\mu-2+\gamma}}{|\mathcal{L}(i\tau, i\xi, v)|} \lesssim \frac{\delta^{\mu-2+\gamma}}{2^l}.$$

In light of these estimates, [Lemma B.4](#) applied with $\alpha_\xi = 0$ and the isotropic variant of [Theorem B.1](#) (see [Remark B.3](#)) show that the expressions

$$\frac{|v| \operatorname{sgn}(v) |v|^{\mu-2+\gamma}}{\delta \mathcal{L}(i\tau, i\xi, v)}, \quad \frac{\operatorname{sgn}(v) |v|^{\mu-2+\gamma}}{\mathcal{L}(i\tau, i\xi, v)}, \quad \frac{|v|^{\mu+m-3+\gamma} |\xi|^2}{\mathcal{L}(i\tau, i\xi, v)^2}$$

extend to constant multipliers of order $\delta^{\mu-2+\gamma}/2^l$ on the supports of $\eta_l \psi'_1(|v|/\delta)$ and $\eta_l \psi_1(|v|/\delta)$, respectively. Hence, we obtain

$$\left\| \int f_l^3 |v|^{\mu-1} dv \right\|_{L^1_{t,x}} \lesssim \frac{\delta^{\mu-2+\gamma}}{2^l} \| |v|^{-\gamma} g_{1,j} \|_{\mathcal{M}_{TV}}.$$

Step 4: Conclusion. We aim to conclude by real interpolation. We set, for $z > 0$,

$$K(z, \bar{f}_l) := \inf \{ \| \bar{f}_l^1 \|_{L^1_{t,x}} + z \| \bar{f}_l^0 \|_{L^\beta_{t,x}} : \bar{f}_l^0 \in L^\beta_{t,x}, \bar{f}_l^1 \in L^1_{t,x}, \bar{f}_l = \bar{f}_l^0 + \bar{f}_l^1 \}.$$

By the above estimates we obtain

$$K(z, \bar{f}_l) \lesssim \frac{\delta^{\mu-2+\gamma}}{2^l} (\| |v|^{1-\gamma} g_0 \|_{\mathcal{M}_{TV}} + \| |v|^{-\gamma} g_1 \|_{\mathcal{M}_{TV}}) + z \delta^{\mu-1+\rho} \| f \|_{L^\beta_{t,x,v}}.$$

We now equilibrate the first and the second term on the right-hand side: we choose $\delta > 0$ such that

$$\frac{\delta^{\mu-2+\gamma}}{2^l} = z \delta^{\mu-1+\rho},$$

that is,

$$\delta := z^{-\frac{1}{1-\gamma+\rho}} 2^{-\frac{l}{1-\gamma+\rho}}.$$

Hence, with

$$\theta := \frac{-\mu + 2 - \gamma}{1 - \gamma + \rho}$$

we obtain

$$\begin{aligned} z^{-\theta} K(z, \bar{f}_l) &\lesssim 2^{-l \frac{\mu-1+\rho}{1-\gamma+\rho}} (\| |v|^{1-\gamma} g_0 \|_{\mathcal{M}_{TV}} + \| |v|^{-\gamma} g_1 \|_{\mathcal{M}_{TV}} + \| f \|_{L^\beta_{t,x,v}}) \\ &= 2^{-l \kappa_l} (\| |v|^{1-\gamma} g_0 \|_{\mathcal{M}_{TV}} + \| |v|^{-\gamma} g_1 \|_{\mathcal{M}_{TV}} + \| f \|_{L^\beta_{t,x,v}}). \end{aligned}$$

As in Step 4 of the proof of [Lemma 4.2](#) we use $(L_{t,x}^1, L_{t,x}^\beta)_{\theta,\infty} = L_{t,x}^{p,\infty}$ with

$$p = \frac{1}{1-\theta\rho} = \frac{1-\gamma+\rho}{\rho\mu + (1-\rho)(1-\gamma)}$$

to obtain

$$\|\bar{f}_l\|_{L_{t,x}^{p,\infty}} \lesssim 2^{-l\kappa_l} (\| |v|^{1-\gamma} g_0 \|_{\mathcal{M}_{TV}} + \| |v|^{-\gamma} g_1 \|_{\mathcal{M}_{TV}} + \|f\|_{L_{t,x,v}^\beta}). \quad (4-11)$$

For $q \in (\min\{p, r\}, \max\{p, r\})$ we choose a corresponding $\vartheta \in (0, 1)$ subject to $1/q = (1-\vartheta)/r + \vartheta/p$. Then using $(L_{t,x}^r, L_{t,x}^{p,\infty})_{\vartheta,q} = L_{t,x}^q$, together with [\(4-11\)](#), we obtain

$$\begin{aligned} \|\bar{f}_l\|_{L_{t,x}^q} &\lesssim \|\bar{f}_l\|_{L_{t,x}^{r-\vartheta}}^{1-\vartheta} \|\bar{f}_l\|_{L_{t,x}^{p,\infty}}^\vartheta \\ &\lesssim \|\bar{f}\|_{L_{t,x}^{r-\vartheta}}^{1-\vartheta} 2^{-l\vartheta\kappa_l} (\| |v|^{1-\gamma} g_0 \|_{\mathcal{M}_{TV}} + \| |v|^{-\gamma} g_1 \|_{\mathcal{M}_{TV}} + \|f\|_{L_{t,x,v}^\infty})^\vartheta \\ &\leq 2^{-l\vartheta\kappa_l} (\| |v|^{1-\gamma} g_0 \|_{\mathcal{M}_{TV}} + \| |v|^{-\gamma} g_1 \|_{\mathcal{M}_{TV}} + \|f\|_{L_{t,x,v}^\infty} + \|\bar{f}\|_{L_{t,x}^r}). \end{aligned}$$

Multiplying by $2^{l\vartheta\kappa_l}$ and taking the supremum over $l \in \mathbb{Z}$ yields [\(4-10\)](#). \square

Corollary 4.5. *Let $m \in (1, \infty)$, $\gamma \in (-\infty, m)$, $\mu \in [1, m+1-\gamma)$, $f \in L_{t,x,v}^1 \cap L_{t,x,v}^\infty$ be a solution to [\(4-1\)](#), and let g_0, g_1 and \bar{f} be as in [Lemma 4.2](#). Let $q \in (1, (m+1-\gamma)/\mu)$ and define*

$$\tilde{\kappa}_x := \frac{\mu q - 1}{q} \frac{2}{m - \gamma}.$$

If $\bar{f} \in L^1(\mathbb{R}^{d+1}) \cap L^q(\mathbb{R}; L^1(\mathbb{R}^d))$, then $\bar{f} \in L^q(\mathbb{R}; W^{\sigma_x, q}(\mathbb{R}^d))$ for all $\sigma_x \in [0, \tilde{\kappa}_x)$. Furthermore,

$$\|\bar{f}\|_{L_t^q(W_x^{\sigma_x, q})} \lesssim \| |v|^{1-\gamma} g_0 \|_{\mathcal{M}_{TV}} + \| |v|^{-\gamma} g_1 \|_{\mathcal{M}_{TV}} + \|f\|_{L_{t,x,v}^1 \cap L_{t,x,v}^\infty} + \|\bar{f}\|_{L_{t,x}^1 \cap L_t^q L_x^1}. \quad (4-12)$$

Proof. We recall the decomposition $f_j = \mathcal{F}_x^{-1} \varphi_j \mathcal{F}_x f$ introduced in the proof of [Lemma 4.2](#). We argue that it suffices to consider the case when $f_j = 0$ for all $j < 0$. Indeed, the part $f_{<} := \sum_{j < 0} f_j$ can be estimated in view of Bernstein's lemma, see [\[Bahouri, Chemin, and Danchin 2011, Lemma 2.1\]](#), via

$$\|\bar{f}_{<}\|_{L_t^q(W_x^{\sigma_x, q})} \lesssim \|\bar{f}\|_{L_t^q L_x^1}.$$

We aim to control \bar{f} in $\tilde{L}_t^q \dot{B}_{q,\infty}^{\vartheta\kappa_x}$ where $\vartheta \in (0, 1)$ is sufficiently large such that $\sigma_x < \vartheta\kappa_x$, and then use [Lemma 2.3](#) to the effect of

$$\|\bar{f}\|_{L_t^q(W_x^{\sigma_x, q})} \lesssim \|\bar{f}\|_{\tilde{L}_t^q \dot{B}_{q,\infty}^{\vartheta\kappa_x}} = \|\bar{f}\|_{\tilde{L}_t^q \dot{B}_{q,\infty}^{\vartheta\kappa_x}},$$

where the last equality is apparent from the definition of the homogeneous and nonhomogeneous Chemin–Lerner spaces and the fact that the low frequencies of f vanish. Thus, it remains to establish

$$\|\bar{f}\|_{\tilde{L}_t^q \dot{B}_{q,\infty}^{\vartheta\kappa_x}} \lesssim \| |v|^{1-\gamma} g_0 \|_{\mathcal{M}_{TV}} + \| |v|^{-\gamma} g_1 \|_{\mathcal{M}_{TV}} + \|f\|_{L_{t,x,v}^1 \cap L_{t,x,v}^\infty} + \|\bar{f}\|_{L_{t,x}^1}. \quad (4-13)$$

For $\tilde{p} \in (1, (m+1-\gamma)/\mu)$, choose

$$\rho := \frac{(\tilde{p}-1)(m-\gamma)}{1 + \tilde{p}(m-\mu-\gamma)}.$$

We claim that ρ is positive and well-defined: Since the numerator is positive due to $\tilde{p} > 1$ and $m > \gamma$, it remains to check that the denominator is positive. This is obvious for $\mu \leq m - \gamma$. For $\mu > m - \gamma$, we observe that due to $\mu < m + 1 - \gamma$ we have

$$\tilde{p} < \frac{m + 1 - \gamma}{\mu} < \frac{1}{\mu + \gamma - m},$$

which implies $1 + \tilde{p}(m - \mu - \gamma) > 0$. Moreover, $\tilde{p} < (m + 1 - \gamma)/\mu$ can be rewritten as $(\tilde{p} - 1)(m - \gamma) < 1 + \tilde{p}(m - \mu - \gamma)$, so that $\rho \in (0, 1)$. Hence, we may apply [Lemma 4.2](#) with this choice of ρ and with $s = 1$. One checks that in this case the integrability and differentiability exponents in (4-3) read

$$p = \tilde{p}, \quad \kappa_t = 0, \quad \kappa_x = \frac{\mu \tilde{p} - 1}{\tilde{p}} \frac{2}{m - \gamma}.$$

Choose $\tilde{p} \in (q, (m + 1 - \gamma)/\mu)$ so that $\tilde{\kappa}_x < \kappa_x$ and define $\vartheta \in (0, 1)$ through

$$\frac{1}{q} = 1 - \vartheta + \frac{\vartheta}{\tilde{p}}.$$

We may choose $\tilde{p} \in (q, (m + 1 - \gamma)/\mu)$ sufficiently small so that $\vartheta \in (0, 1)$ is so large that $\sigma_x < \vartheta \tilde{\kappa}_x < \vartheta \kappa_x$. In view of (4-5) (with the space $S_{q,\infty}^{\vartheta \tilde{\kappa}} \dot{B} = S_{q,\infty}^{(0,\vartheta \kappa_x)} \dot{B}$ replaced by $\tilde{L}_t^q \dot{B}_{q,\infty}^{\vartheta \kappa_x}$) we obtain

$$\|\bar{f}_j\|_{L_{t,x}^q} \lesssim 2^{-j\vartheta \kappa_x} (\| |v|^{1-\gamma} g_0 \|_{\mathcal{M}_{TV}} + \| |v|^{-\gamma} g_1 \|_{\mathcal{M}_{TV}} + \|f\|_{L_{t,x,v}^\beta} + \|\bar{f}\|_{L_{t,x}^1}),$$

where we recall the notation $\bar{f}_j := \int \mathcal{F}_x^{-1}[\varphi_j \mathcal{F}_x f] |v|^{\mu-1} dv$. If we multiply by $2^{j\vartheta \kappa_x}$ and take the supremum over $j \in \mathbb{Z}$, this yields

$$\|\bar{f}\|_{\tilde{L}_t^q \dot{B}_{q,\infty}^{\vartheta \kappa_x}} \lesssim \| |v|^{1-\gamma} g_0 \|_{\mathcal{M}_{TV}} + \| |v|^{-\gamma} g_1 \|_{\mathcal{M}_{TV}} + \|f\|_{L_{t,x,v}^\beta} + \|\bar{f}\|_{L_{t,x}^1}.$$

By the estimate $\|f\|_{L_{t,x,v}^\beta} \lesssim \|f\|_{L_{t,x,v}^1} + \|f\|_{L_{t,x,v}^\infty}$, this gives (4-13). \square

Corollary 4.6. *Let $m \in (1, \infty)$, $\gamma \in (-\infty, 1)$, $f \in L_{t,x,v}^1 \cap L_{t,x,v}^\infty$ be a solution to (4-1), and let g_0 and g_1 be as in [Lemma 4.2](#). Assume $\bar{f} \in L_{t,x}^r$ for all $r \in [1, m + 1 - \gamma)$, where $\bar{f}(t, x) := \int f(t, x, v) dv$. Let $\tilde{p} \in (2 - \gamma, m + 1 - \gamma)$ and define*

$$\tilde{\kappa}_t := \frac{m + 1 - \gamma - \tilde{p}}{\tilde{p}} \frac{1}{m - 1}, \quad \tilde{\kappa}_x := \frac{\tilde{p} - 2 + \gamma}{\tilde{p}} \frac{2}{m - 1}.$$

Then $\bar{f} \in W^{\sigma_t, \tilde{p}}(\mathbb{R}; W^{\sigma_x, \tilde{p}}(\mathbb{R}^d))$ for all $\sigma_t \in [0, \tilde{\kappa}_t)$ and $\sigma_x \in [0, \tilde{\kappa}_x)$. Furthermore, there is an $r \in (\tilde{p}, m + 1 - \gamma)$ such that

$$\|\bar{f}\|_{W^{\sigma_t, \tilde{p}}(W^{\sigma_x, \tilde{p}})} \lesssim \| |v|^{1-\gamma} g_0 \|_{\mathcal{M}_{TV}} + \| |v|^{-\gamma} g_1 \|_{\mathcal{M}_{TV}} + \|f\|_{L_{t,x,v}^1 \cap L_{t,x,v}^\infty} + \|\bar{f}\|_{L_{t,x}^r}. \quad (4-14)$$

Proof. As we need to pass from homogeneous spaces (the output of [Lemmas 4.2](#) and [4.4](#)) to a nonhomogeneous space, our strategy is to invoke [Lemmas 2.5](#) and [2.4](#). The input to [Lemma 2.5](#) requires four pieces of information, namely control of \bar{f} in $L^{\tilde{p}}(\mathbb{R}^{d+1})$, $\tilde{L}_x^{\tilde{p}} \dot{B}_{\tilde{p},\infty}^{\sigma_t}$, $\tilde{L}_t^{\tilde{p}} \dot{B}_{\tilde{p},\infty}^{\sigma_x}$ and $S_{\tilde{p},\infty}^{\tilde{\sigma}} \dot{B}$. Since the control of \bar{f} in $L^{\tilde{p}}(\mathbb{R}^{d+1})$ is ensured by assumption, we concentrate on the other three contributions. Note that the main difficulty lies in the condition that both the integrability exponent and the orders of differentiability have to match exactly.

Step 1: $\bar{f} \in S_{\tilde{p}, \infty}^{\tilde{\sigma}} \dot{B}$. Let $r \in (\tilde{p}, m+1-\gamma)$ to be chosen in Step 3. We claim that there exist functions $k_t, k_x : (0, \infty) \rightarrow (0, \infty)$ with $k_t(\varepsilon), k_x(\varepsilon) \rightarrow 0$ as $\varepsilon \rightarrow 0$ such that it holds for all $\varepsilon \ll 1$

$$\|\bar{f}\|_{S_{\tilde{p}, \infty}^{\tilde{\sigma}} \dot{B}} \lesssim \| |v|^{1-\gamma} g_0 \|_{\mathcal{M}_{TV}} + \| |v|^{-\gamma} g_1 \|_{\mathcal{M}_{TV}} + \|f\|_{L_{t,x,v}^1 \cap L_{t,x,v}^\infty} + \|\bar{f}\|_{L_{t,x}^r}, \quad (4-15)$$

where we have used the notation $\sigma_t := \tilde{\kappa}_t - k_t(\varepsilon)$ and $\sigma_x := \tilde{\kappa}_x - k_x(\varepsilon)$.

We apply [Lemma 4.2](#) with $\mu = 1$, $\rho = 1 - \varepsilon$, and $s := s_\varepsilon \in (0, 1)$, where s_ε is chosen so that the integrability assertion in (4-3) reads $p = \tilde{p}$; this is possible for ρ close to 1 in view of [Remark 4.3](#). Moreover, we may choose $\vartheta \in (0, 1)$ such that κ_t and κ_x defined through (4-3) satisfy $\vartheta \kappa_t = \tilde{\kappa}_t - k_t(\varepsilon)$ and $\vartheta \kappa_x = \tilde{\kappa}_x - k_x(\varepsilon)$ for some functions k_t and k_x as above. Then for $1 < q_0 < \tilde{p} < q_1 < m+1-\gamma$ so that

$$\frac{1}{q_0} = 1 - \vartheta + \frac{\vartheta}{\tilde{p}}, \quad \frac{1}{q_1} = \frac{1 - \vartheta}{r} + \frac{\vartheta}{\tilde{p}},$$

we obtain in view of (4-5) that

$$\|\bar{f}_{l,j}\|_{L_{t,x}^{q_i}} \lesssim 2^{-l\vartheta\kappa_t} 2^{-j\vartheta\kappa_x} (\| |v|^{1-\gamma} g_0 \|_{\mathcal{M}_{TV}} + \| |v|^{-\gamma} g_1 \|_{\mathcal{M}_{TV}} + \|f\|_{L_{t,x,v}^\beta} + \|\bar{f}\|_{L_{t,x}^1 \cap L_{t,x}^r}),$$

for $i = 0, 1$, where we recall the notation $\bar{f}_{l,j} := \int \mathcal{F}_{t,x}^{-1}[\eta_l \varphi_j \mathcal{F}_{t,x} f] dv$. Since $(L_{t,x}^{q_0}, L_{t,x}^{q_1})_{\theta, \tilde{p}} = L_{t,x}^{\tilde{p}}$ for an appropriate $\theta \in (0, 1)$, we thus obtain

$$\|\bar{f}_{l,j}\|_{L_{t,x}^{\tilde{p}}} \lesssim 2^{-l\vartheta\kappa_t} 2^{-j\vartheta\kappa_x} (\| |v|^{1-\gamma} g_0 \|_{\mathcal{M}_{TV}} + \| |v|^{-\gamma} g_1 \|_{\mathcal{M}_{TV}} + \|f\|_{L_{t,x,v}^\beta} + \|\bar{f}\|_{L_{t,x}^1} + \|\bar{f}\|_{L_{t,x}^r}),$$

which after multiplying by $2^{l\vartheta\kappa_t} 2^{j\vartheta\kappa_x}$ and taking the supremum over $l, j \in \mathbb{Z}$ yields

$$\|\bar{f}\|_{S_{\tilde{p}, \infty}^{\vartheta(\kappa_t, \kappa_x)} \dot{B}} \lesssim \| |v|^{1-\gamma} g_0 \|_{\mathcal{M}_{TV}} + \| |v|^{-\gamma} g_1 \|_{\mathcal{M}_{TV}} + \|f\|_{L_{t,x,v}^\beta} + \|\bar{f}\|_{L_{t,x}^1} + \|\bar{f}\|_{L_{t,x}^r}. \quad (4-16)$$

By the estimate $\|f\|_{L_{t,x,v}^\beta} + \|\bar{f}\|_{L_{t,x}^1} \lesssim \|f\|_{L_{t,x,v}^1} + \|f\|_{L_{t,x,v}^\infty}$, this gives (4-15).

Step 2: $\bar{f} \in \tilde{L}_t^{\tilde{p}} \dot{B}_{\tilde{p}, \infty}^{\sigma_x}$. In this step we establish

$$\|\bar{f}\|_{\tilde{L}_t^{\tilde{p}} \dot{B}_{\tilde{p}, \infty}^{\sigma_x}} \lesssim \| |v|^{1-\gamma} g_0 \|_{\mathcal{M}_{TV}} + \| |v|^{-\gamma} g_1 \|_{\mathcal{M}_{TV}} + \|f\|_{L_{t,x,v}^1 \cap L_{t,x,v}^\infty} + \|\bar{f}\|_{L_{t,x}^r}. \quad (4-17)$$

Choose

$$\rho := \frac{(\tilde{p} - 1)(m - \gamma)}{1 + \tilde{p}(m - 1 - \gamma)}.$$

We claim that ρ is positive and well-defined: Since the numerator is positive due to $\tilde{p} > 1$ and $m > \gamma$, it remains to check that the denominator is positive. This is obvious for $\gamma \leq m - 1$. For $\gamma > m - 1$, we observe that

$$\tilde{p} < m + 1 - \gamma < \frac{1}{1 + \gamma - m},$$

which implies $1 + \tilde{p}(m - 1 - \gamma) > 0$. Moreover, $\tilde{p} < m + 1 - \gamma$ can be rewritten as $(\tilde{p} - 1)(m - \gamma) < 1 + \tilde{p}(m - 1 - \gamma)$, so that $\rho \in (0, 1)$. Hence, we may apply [Lemma 4.2](#) with this choice of ρ and with $s = 1$. One checks that in this case the integrability and differentiability exponents in (4-3) read

$$p = \tilde{p}, \quad \kappa_t = 0, \quad \kappa_x = \frac{p-1}{p} \frac{2}{m-\gamma}.$$

We observe that $\kappa_x \geq \tilde{\kappa}_x$ and hence we find $\vartheta \in (0, 1)$ such that $\vartheta \kappa_x = \tilde{\kappa}_x - k_x(\varepsilon)$. The same interpolation argument as in Step 1 gives now the estimate (4-17).

Step 3: $\bar{f} \in \tilde{L}_x^{\tilde{p}} \dot{B}_{\tilde{p}, \infty}^{\sigma_t}$. In this step we show that there is some $r \in (\tilde{p}, m + 1 - \gamma)$ such that

$$\|\bar{f}\|_{\tilde{L}_x^{\tilde{p}} \dot{B}_{\tilde{p}, \infty}^{\sigma_t}} \lesssim \| |v|^{1-\gamma} g_0 \|_{\mathcal{M}_{TV}} + \| |v|^{-\gamma} g_1 \|_{\mathcal{M}_{TV}} + \|f\|_{L_{t,x,v}^\infty} + \|\bar{f}\|_{L_{t,x}^r}. \quad (4-18)$$

We apply Lemma 4.4 with $\mu = 1$ and $\rho = 1$. In this case, (4-9) reads $p = 2 - \gamma$ and $\kappa_t = 1/(2 - \gamma)$. Since $\tilde{p} > 2 - \gamma$, we have $\tilde{\kappa}_t < \kappa_t$. Hence, we can choose $\vartheta \in (0, 1)$ such that $\vartheta \kappa_t = \tilde{\kappa}_t - k_t(\varepsilon)$. In particular,

$$\vartheta < \frac{\tilde{\kappa}_t}{\kappa_t} = \frac{m + 1 - \gamma - \tilde{p}}{\tilde{p}} \frac{2 - \gamma}{m - 1} < \frac{2 - \gamma}{\tilde{p}},$$

so that

$$r = \frac{\tilde{p}(2 - \gamma)(1 - \vartheta)}{2 - \gamma - \vartheta \tilde{p}}$$

is well-defined. Since r is increasing in ϑ due to $\tilde{p} > 2 - \gamma$, we see that $r \in (\tilde{p}, m + 1 - \gamma)$. We have $1/\tilde{p} = (1 - \vartheta)/r + \vartheta/p$, and hence Lemma 4.4 gives estimate (4-18).

Step 4: Conclusion. Since $\bar{f} \in L_{t,x}^{\tilde{p}}$ by assumption, Lemma 2.5 combined with Lemma 2.4 yields the result. \square

Corollary 4.7. *Let $m \in (1, \infty)$, $\gamma \in (-\infty, m)$, and let $f \in L_{t,x,v}^1 \cap L_{t,x,v}^\infty$ be a solution to (4-1). Let g_0 and g_1 be as in Lemma 4.2 and assume additionally*

$$|g_0|(t, x, v) \in \mathcal{M}_{TV}(\mathbb{R}_t \times \mathbb{R}_x^d \times \mathbb{R}_v).$$

Assume $\tilde{p} \in (2 - \gamma, m + 1 - \gamma) \cap (1, m + 1 - \gamma)$ and define

$$\tilde{\kappa}_t := \frac{m + 1 - \gamma - \tilde{p}}{\tilde{p}} \frac{1}{m - 1}, \quad \tilde{\kappa}_x := \frac{\tilde{p} - 2 + \gamma}{\tilde{p}} \frac{2}{m - 1}.$$

If $\bar{f} \in L^r(\mathbb{R}^{d+1}) \cap L^1(\mathbb{R}; L^{\tilde{p}}(\mathbb{R}^d))$ for all $r \in [1, m + 1 - \gamma)$, where $\bar{f}(t, x) := \int f(t, x, v) dv$, and if $\int |v|^{m-1} f dv \in L^1(\mathbb{R}^{d+1})$, then $\bar{f} \in W^{\sigma_t, \tilde{p}}(\mathbb{R}; W^{\sigma_x, \tilde{p}}(\mathbb{R}^d))$ for all $\sigma_t \in [0, \tilde{\kappa}_t)$ and $\sigma_x \in [0, \tilde{\kappa}_x)$. Furthermore, there is an $r \in (\tilde{p}, m + 1 - \gamma)$ such that

$$\begin{aligned} \|\bar{f}\|_{W^{\sigma_t, \tilde{p}}(W^{\sigma_x, \tilde{p}})} &\lesssim \|g_0\|_{\mathcal{M}_{TV}} + \| |v|^{1-\gamma} g_0 \|_{\mathcal{M}_{TV}} + \| |v|^{-\gamma} g_1 \|_{\mathcal{M}_{TV}} \\ &\quad + \|f\|_{L_{t,x,v}^1 \cap L_{t,x,v}^\infty} + \|\bar{f}\|_{L_t^1 L_x^{\tilde{p}} \cap L_{t,x}^r} + \left\| \int |v|^{m-1} f dv \right\|_{L_{t,x}^1}. \end{aligned} \quad (4-19)$$

Proof. It suffices to adapt Step 3 of the proof of Corollary 4.6, that is, the control of \bar{f} in $\tilde{L}_x^{\tilde{p}} \dot{B}_{\tilde{p}, \infty}^{\sigma_t}$.

Step 3: $\bar{f} \in \tilde{L}_x^{\tilde{p}} \dot{B}_{\tilde{p}, \infty}^{\sigma_t}$. In this step we show that there is some $r \in (\tilde{p}, m + 1 - \gamma)$ such that

$$\begin{aligned} \|\bar{f}\|_{\tilde{L}_x^{\tilde{p}} \dot{B}_{\tilde{p}, \infty}^{\sigma_t}} &\lesssim \|g_0\|_{\mathcal{M}_{TV}} + \| |v|^{1-\gamma} g_0 \|_{\mathcal{M}_{TV}} + \| |v|^{-\gamma} g_1 \|_{\mathcal{M}_{TV}} \\ &\quad + \|f\|_{L_{t,x,v}^1 \cap L_{t,x,v}^\infty} + \|\bar{f}\|_{L_t^1 L_x^{\tilde{p}} \cap L_{t,x}^r} + \left\| \int |v|^{m-1} f dv \right\|_{L_{t,x}^1}. \end{aligned} \quad (4-20)$$

We split f into three contributions

$$\begin{aligned} f &= \mathcal{F}_t^{-1} \psi_0(\tau) \mathcal{F}_t f + \mathcal{F}_{t,x}^{-1} (1 - \psi_0(\tau)) (1 - \phi_0(\xi)) \mathcal{F}_{t,x} f + \mathcal{F}_{t,x}^{-1} (1 - \psi_0(\tau)) \phi_0(\xi) \mathcal{F}_{t,x} f \\ &=: f^1 + f^2 + f^3. \end{aligned}$$

The low time-frequency part f^1 can be estimated in view of [Lemma 2.3](#) and Bernstein's lemma, see [\[Bahouri, Chemin, and Danchin 2011, Lemma 2.1\]](#), via

$$\|\bar{f}^1\|_{\tilde{L}_x^{\tilde{p}} \dot{B}_{\tilde{p},\infty}^{\sigma_t}} \lesssim \|\bar{f}^1\|_{\tilde{L}_x^{\tilde{p}} B_{\tilde{p},\infty}^{\sigma_t}} \lesssim \|\bar{f}^1\|_{W^{\sigma_t+\varepsilon,\tilde{p}}(L_x^{\tilde{p}})} \lesssim \|\bar{f}\|_{L_t^1 L_x^{\tilde{p}}}. \quad (4-21)$$

Next, we apply [Lemma 4.2](#) with $\mu = 1$, sufficiently large $\rho \in (0, 1)$ and sufficiently small $s \in ((\gamma - 1)/(m - 1), 1]$ so that (4-3) implies $p < \tilde{p}$ and $\kappa_t > \tilde{\kappa}_t$. Hence, we can choose $\vartheta \in (0, 1)$ such that $\tilde{\kappa}_t > \vartheta \kappa_t > \tilde{\kappa}_t - k_t(\varepsilon)$. In particular, in light of [Remark 4.3](#)

$$\vartheta < \frac{\tilde{\kappa}_t}{\kappa_t} = \frac{m + 1 - \gamma - \tilde{p}}{m + \rho - \gamma - p\rho + p(1 - \rho)(\gamma - m)} \frac{p\rho}{\tilde{p}} < \frac{p}{\tilde{p}} \quad \text{if } 1 - \rho \ll 1,$$

so that $r = \tilde{p}p(1 - \vartheta)/(p - \vartheta\tilde{p})$ is well-defined. Since r is increasing in ϑ due to $\tilde{p} > p$, we see that $r \in (\tilde{p}, m + 1 - \gamma)$. We have $1/\tilde{p} = (1 - \vartheta)/r + \vartheta/p$, and hence [Lemma 4.2](#) gives

$$\|\bar{f}\|_{S_{\tilde{p},\infty}^{\vartheta\tilde{\kappa}} \dot{B}} \lesssim \| |v|^{1-\gamma} g_0 \|_{\mathcal{M}_{TV}} + \| |v|^{-\gamma} g_1 \|_{\mathcal{M}_{TV}} + \| f \|_{L_{t,x,v}^1 \cap L_{t,x,v}^\infty} + \|\bar{f}\|_{L_{t,x}^r}.$$

Thus, since f^2 is supported only on $\eta_l \varphi_j$ for nonnegative $l, j \in \mathbb{Z}$, [Lemmas 2.3](#) and [2.4](#) show, in view of the definition of the homogeneous and nonhomogeneous Besov spaces and $\sigma_t < \vartheta \kappa_t$ as well as $0 < \vartheta \kappa_x$,

$$\|\bar{f}^2\|_{\tilde{L}_x^{\tilde{p}} \dot{B}_{\tilde{p},\infty}^{\sigma_t}} = \|\bar{f}^2\|_{\tilde{L}_x^{\tilde{p}} B_{\tilde{p},\infty}^{\sigma_t}} \lesssim \|\bar{f}^2\|_{S_{\tilde{p},\infty}^{\vartheta\tilde{\kappa}} B} = \|\bar{f}\|_{S_{\tilde{p},\infty}^{\vartheta\tilde{\kappa}} \dot{B}}.$$

Thus,

$$\|\bar{f}^2\|_{\tilde{L}_x^{\tilde{p}} \dot{B}_{\tilde{p},\infty}^{\sigma_t}} \lesssim \| |v|^{1-\gamma} g_0 \|_{\mathcal{M}_{TV}} + \| |v|^{-\gamma} g_1 \|_{\mathcal{M}_{TV}} + \| f \|_{L_{t,x,v}^1 \cap L_{t,x,v}^\infty} + \|\bar{f}\|_{L_{t,x}^r}. \quad (4-22)$$

It remains to estimate the contribution of f^3 . For $l \in \mathbb{Z}$, we introduce $f_l^3 := \mathcal{F}_t^{-1} \eta_l(\tau) \mathcal{F}_t f^3$. Since $f_l^3 = 0$ for $l < 0$, we may concentrate on the case $l \geq 0$. Observe that f_l^3 solves the equation

$$f_l^3 = -m|v|^{m-1} \mathcal{F}_{t,x}^{-1} \frac{|\xi|^2}{i\tau} \eta_l(\tau) \phi_0(\xi) \mathcal{F}_{t,x} f + \mathcal{F}_{t,x}^{-1} \frac{\phi_0(\xi)}{i\tau} \mathcal{F}_{t,x} g_{0,l} + \mathcal{F}_{t,x}^{-1} \frac{\phi_0(\xi)}{i\tau} \mathcal{F}_{t,x} \partial_v g_{1,l}.$$

Integrating in v , we obtain

$$\bar{f}_l^3 = -m \int |v|^{m-1} \mathcal{F}_{t,x}^{-1} \frac{|\xi|^2}{i\tau} \eta_l(\tau) \phi_0(\xi) \mathcal{F}_{t,x} f \, dv + \mathcal{F}_{t,x}^{-1} \frac{1}{i\tau} \phi_0(\xi) \mathcal{F}_{t,x} \int g_{0,l,j} \, dv.$$

Since $|\xi|^2$ acts as a constant multiplier on the support of ϕ_0 and τ^{-1} acts as a constant multiplier of order 2^{-l} on the support of η_l , it follows by Bernstein's lemma

$$\|\bar{f}_l^3\|_{L_{t,x}^{\tilde{p}}} \lesssim 2^{l(1-\frac{1}{\tilde{p}})} \|\bar{f}_l^3\|_{L_{t,x}^1} \lesssim 2^{-l\frac{1}{\tilde{p}}} \left(\left\| \int |v|^{m-1} f \, dv \right\|_{L_{t,x}^1} + \|g_0\|_{\mathcal{M}_{TV}} \right).$$

Since $\tilde{p} > 2 - \gamma$, we have

$$\sigma_t < \tilde{\kappa}_t = \frac{m+1-\gamma-\tilde{p}}{\tilde{p}} \frac{1}{m-1} < \frac{1}{\tilde{p}}.$$

In view of $l \geq 0$ this yields

$$\|\tilde{f}_l^3\|_{L_{t,x}^{\tilde{p}}} \lesssim 2^{-l\sigma_t} \left(\left\| \int |v|^{m-1} f \, dv \right\|_{L_{t,x}^1} + \|g_0\|_{\mathcal{M}_{TV}} \right).$$

Multiplying by $2^{l\sigma_t}$ and taking the supremum over $l \in \mathbb{Z}$, we conclude

$$\|\tilde{f}^3\|_{\tilde{L}_x^{\tilde{p}} \dot{B}_{\tilde{p},\infty}^{\sigma_t}} \lesssim \left\| \int |v|^{m-1} f \, dv \right\|_{L_{t,x}^1} + \|g_0\|_{\mathcal{M}_{TV}}. \quad (4-23)$$

Collecting (4-21), (4-22) and (4-23), we arrive at (4-20). \square

5. Application to porous medium equations

In this section, we provide proofs of our main results by applying the averaging lemmas obtained in the previous section to entropy solutions to (1-1).

Proof of Theorem 1.2. We first argue that we have $u \in L_{t,x}^s$ for all $s \in [1, m-1+\rho)$. Since $T < \infty$, Theorem A.2 gives

$$\|u\|_{L_{t,x}^1} \lesssim \sup_{t \in [0,T]} \|u(t)\|_{L_x^1} \lesssim \|u_0\|_{L_x^1} + \|S\|_{L_{t,x}^1}, \quad (5-1)$$

so that we may concentrate on $s > 1$. Let f be the kinetic function corresponding to u and solving (1-8). In order to apply Corollary 4.5 with $\mu = 1$ and $\sigma_x = 0$, we need to extend (1-8) to all times $t \in \mathbb{R}$, which can be achieved by multiplication with a smooth cut-off function $\varphi \in C_c^\infty(0, T)$ with $0 \leq \varphi \leq 1$. Hence, we set $g_0 := \delta_{v=u(t,x)} S + \partial_t \varphi f$ and $g_1 := q$. Let $\gamma := 2 - \rho$, so that $s \in (1, m+1-\gamma)$. From (4-12) we obtain

$$\begin{aligned} \|\varphi u\|_{L_{t,x}^s} &\lesssim \| |v|^{\rho-1} g_0 \|_{\mathcal{M}_{TV}} + \| |v|^{\rho-2} g_1 \|_{\mathcal{M}_{TV}} + \|\varphi f\|_{L_{t,x,v}^1 \cap L_{t,x,v}^\infty} + \|\varphi u\|_{L_{t,x}^1 \cap L_t^s L_x^1} \\ &\lesssim \| |v|^{\rho-1} g_0 \|_{\mathcal{M}_{TV}} + \| |v|^{\rho-2} g_1 \|_{\mathcal{M}_{TV}} + \|f\|_{L_{t,x,v}^1 \cap L_{t,x,v}^\infty} + \sup_{t \in [0,T]} \|u(t)\|_{L_x^1}. \end{aligned}$$

We note that since trivially $f \in L_{t,x,v}^\infty$ with norm bounded by 1, estimate (5-1) gives

$$\|f\|_{L_{t,x,v}^1 \cap L_{t,x,v}^\infty} + \sup_{t \in [0,T]} \|u(t)\|_{L_x^1} \lesssim \|u\|_{L_{t,x}^1} + 1 + \sup_{t \in [0,T]} \|u(t)\|_{L_x^1} \lesssim \|u_0\|_{L_x^1} + \|S\|_{L_{t,x}^1} + 1.$$

Next, we check that $|v|^{\rho-1} g_0 \in \mathcal{M}_{TV}$. Indeed, we observe that $(\rho-1)\rho' := \rho$, and hence, applying Lemma A.3,

$$\begin{aligned} \| |v|^{\rho-1} g_0 \|_{\mathcal{M}_{TV}} &= \| |v|^{\rho-1} (\delta_{v=u(t,x)} S + \partial_t \varphi f) \|_{\mathcal{M}_{TV}} \lesssim \| |u|^{\rho-1} S \|_{L_{t,x}^1} + \| \partial_t \varphi |u|^\rho \|_{L_{t,x}^1} \\ &\lesssim \| |u|^{(\rho-1)\rho'} \|_{L_{t,x}^1} + \| |S|^\rho \|_{L_{t,x}^1} + \| \partial_t \varphi |u|^\rho \|_{L_{t,x}^1} \\ &\lesssim \|u_0\|_{L_x^\rho}^\rho + \|S\|_{L_{t,x}^\rho}^\rho + \| \partial_t \varphi |u|^\rho \|_{L_{t,x}^1}. \end{aligned}$$

Utilizing [Lemma A.3](#) once more to the effect of

$$\| |v|^{\rho-2} g_1 \|_{\mathcal{M}_{TV}} = \| |v|^{\rho-2} q \|_{\mathcal{M}_{TV}} \lesssim \| u_0 \|_{L_{t,x}^\rho}^\rho + \| S \|_{L_{t,x}^\rho}^\rho,$$

we obtain

$$\| \varphi u \|_{L_{t,x}^s} \lesssim \| u_0 \|_{L_x^1 \cap L_x^\rho}^\rho + \| S \|_{L_{t,x}^1 \cap L_{t,x}^\rho}^\rho + \| \partial_t \varphi |u|^\rho \|_{L_{t,x}^1} + 1.$$

We may set $\varphi_n(t) = \psi(nt) - \psi(nt - T/2)$, where $\psi \in C^\infty(\mathbb{R})$ with $0 \leq \psi \leq 1$, $\text{supp } \psi \subset (0, \infty)$, $\psi(t) = 1$ for $t > T/2$ and $\| \partial_t \psi \|_{L^1} = 1$. For $n \rightarrow \infty$, φ_n converges to $1_{[0,T]}$ in the supremum norm, while $\partial_t \varphi_n$ is a smooth approximation of $\delta_{\{t=0\}} - \delta_{\{t=T\}}$. Therefore, $\| \varphi_n u \|_{L_{t,x}^s} \rightarrow \| u \|_{L_{t,x}^s}$ and by an application of [Lemma A.3](#)

$$\| \partial_t \varphi_n |u|^\rho \|_{L_{t,x}^1} \rightarrow \| |u|(0)^\rho - |u|(T)^\rho \|_{L_x^1} \lesssim \| u_0 \|_{L_x^\rho}^\rho + \| S \|_{L_{t,x}^\rho}^\rho,$$

so that $u \in L^s([0, T] \times \mathbb{R}^d)$ and

$$\| u \|_{L_{t,x}^s} \lesssim \| u_0 \|_{L_x^1 \cap L_x^\rho}^\rho + \| S \|_{L_{t,x}^1 \cap L_{t,x}^\rho}^\rho + 1. \quad (5-2)$$

(i) We apply [Corollary 4.5](#) once more. Let f , φ , g_0 , g_1 and γ be as above. Then, in particular $p \in (1, (m+1-\gamma)/\mu)$. From (4-12) we obtain

$$\| \varphi u^{[\mu]} \|_{L_t^p(W^{\sigma_x, p})} \lesssim \| |v|^{1-\gamma} g_0 \|_{\mathcal{M}_{TV}} + \| |v|^{-\gamma} g_1 \|_{\mathcal{M}_{TV}} + \| f \|_{L_{t,x,v}^1 \cap L_{t,x,v}^\infty} + \| u^{[\mu]} \|_{L_{t,x}^1 \cap L_t^p L_x^1}.$$

The first three contributions on the right-hand side are estimated as above. For the last contribution, we note $1 \leq \mu < p\mu$ and thus

$$\begin{aligned} \| u^{[\mu]} \|_{L_{t,x}^1 \cap L_t^p L_x^1} &\lesssim \| u^{[\mu]} \|_{L_t^p L_x^1} = \| u \|_{L_t^{p\mu} L_x^\mu}^\mu \lesssim (\| u \|_{L_t^{p\mu} L_x^1} + \| u \|_{L_{t,x}^{p\mu}})^\mu \\ &\lesssim \left(\sup_{t \in [0, T]} \| u(t) \|_{L_x^1} + \| u \|_{L_{t,x}^{p\mu}} \right)^\mu \lesssim \sup_{t \in [0, T]} \| u(t) \|_{L_x^1}^\mu + \| u \|_{L_{t,x}^{p\mu}}^\mu + 1. \end{aligned}$$

Furthermore, (5-1) together with (5-2) applied with $s = p\mu \in (1, m-1+\rho)$ shows

$$\sup_{t \in [0, T]} \| u(t) \|_{L_x^1}^\mu + \| u \|_{L_{t,x}^{p\mu}}^\mu + 1 \lesssim \| u_0 \|_{L_x^1 \cap L_x^\rho}^{\mu\rho} + \| S \|_{L_{t,x}^1 \cap L_{t,x}^\rho}^{\mu\rho} + 1.$$

Hence, arguing as above by taking the limit $\varphi_n \rightarrow 1_{[0,T]}$, we obtain $u^{[\mu]} \in L^p(\mathbb{R}; W^{\sigma_x, p}(\mathbb{R}^d))$ and (1-4).

(ii) The proof is similar to the first part, but we use [Corollary 4.6](#) instead of [Corollary 4.5](#). Again we localize in time by multiplying with a smooth cut-off function $\varphi \in C_c^\infty(0, T)$ with $0 \leq \varphi \leq 1$ and set g_0 and g_1 as before. Choose $\gamma := 2 - \rho$, so that $p \in (2 - \gamma, m + 1 - \gamma)$. From (4-14) in [Corollary 4.6](#) we obtain

$$\| \varphi u \|_{W^{\sigma_t, p}(W^{\sigma_x, p})} \lesssim \| |v|^{1-\gamma} g_0 \|_{\mathcal{M}_{TV}} + \| |v|^{-\gamma} g_1 \|_{\mathcal{M}_{TV}} + \| f \|_{L_{t,x,v}^1 \cap L_{t,x,v}^\infty} + \| u \|_{L_{t,x}^r},$$

where $r \in (\rho, m-1+\rho)$. The terms involving g_0 , g_1 and f can be estimated as above, while the $L_{t,x}^r$ -norm of u can be estimated by (5-2). Choosing φ_n as above, we hence infer that $\varphi_n u$ is bounded in

$W^{\sigma_t, p}(0, T; W^{\sigma_x, p}(\mathbb{R}^d))$ and

$$\sup_{n \in \mathbb{N}} \|\varphi_n u\|_{W^{\sigma_t, p}(W^{\sigma_x, p})} \lesssim \|u_0\|_{L_x^1 \cap L_x^\rho}^\rho + \|S\|_{L_{t,x}^1 \cap L_{t,x}^\rho}^\rho + 1.$$

Since $\varphi_n u \rightarrow u1_{[0,T]}$ in the sense of distributions, we obtain the result by the weak lower semicontinuity of the norm in $W^{\sigma_t, p}(0, T; W^{\sigma_x, p}(\mathbb{R}^d))$. \square

Proof of Corollary 1.3. (i) Let $\sigma_x \in [0, 2\mu/m)$. We apply Theorem 1.2(i) with $p = m/\mu$ for sufficiently small $\eta \in (1, \rho]$ so that

$$\sigma_x < \frac{\mu p - 1}{p} \frac{2}{m - 2 + \eta} = \frac{2\mu}{m} \frac{m - 1}{m - 2 + \eta}$$

and observe that for all $q \in [1, p]$ we have the embedding $L^p(0, T; W^{\sigma_x, p}(\mathbb{R}^d)) \subset L^q(0, T; W^{\sigma_x, q}(\mathcal{O}))$.

(ii) For $s > 0$ we have, with $p = s(m - 1) + 1 \in (1, m]$,

$$\kappa_t = \frac{1 - s}{s(m - 1) + 1} = \frac{m - p}{p} \frac{1}{m - 1}, \quad \kappa_x = \frac{2s}{s(m - 1) + 1} = \frac{p - 1}{p} \frac{2}{m - 1}.$$

Hence, in this case the assertion follows by an application of Theorem 1.2(ii) with sufficiently small $\eta \in (1, \rho]$ such that $p > \rho$ and

$$\sigma_x < \frac{p - \rho}{p} \frac{2}{m - 1}$$

combined with the embedding

$$W^{\sigma_t, p}(0, T; W^{\sigma_x, p}(\mathbb{R}^d)) \subset W^{\sigma_t, q}(0, T; W^{\sigma_x, q}(\mathcal{O})).$$

If $s = 0$ and $\sigma_t \in [0, 1)$, we may choose $s_0 > 0$ such that

$$\sigma_t < \frac{1 - s_0}{s_0(m - 1) + 1} =: \kappa_t(s_0),$$

and the result follows by the embedding

$$W^{\kappa_t(s_0), s_0(m-1)+1}(0, T; L^{s_0(m-1)+1}(\mathcal{O})) \subset W^{\sigma_t, 1}(0, T; L^1(\mathcal{O})). \quad \square$$

Proof of Theorem 1.1. The proof is similar to that of Theorem 1.2(ii), but we discriminate between small and large velocity contributions to the kinetic function. Let f be the kinetic function corresponding to u and solving (1-8). We extend again to all times $t \in \mathbb{R}$ by multiplying with a smooth cut-off function $\varphi \in C_c^\infty(0, T)$ with $0 \leq \varphi \leq 1$. Further, we split $f =: f^< + f^>$ and $q =: q^< + q^>$ into a small-velocity and a large-velocity part by multiplying with a smooth cut-off function ψ_0 respectively $\psi_1 := 1 - \psi_0$ in v . This gives rise to the two equations

$$\begin{aligned} \partial_t(\varphi f^<) - m|v|^{m-1} \Delta_x(\varphi f^<) &= \varphi \psi_0 \delta_{v=u(t,x)} S + \partial_v(\varphi q^<) - \varphi q \partial_v \psi_0 + \partial_t \varphi f^<, \\ \partial_t(\varphi f^>) - m|v|^{m-1} \Delta_x(\varphi f^>) &= \varphi \psi_1 \delta_{v=u(t,x)} S + \partial_v(\varphi q^>) + \varphi q \partial_v \psi_0 + \partial_t \varphi f^>, \end{aligned}$$

Integrating $f^<$ and $f^>$ in v , we obtain a decomposition of $u = u^< + u^>$.

The proof proceeds in several steps: In first the three steps, we argue that $u \in L^s(0, T; L^s(\mathbb{R}^d))$ for all $s \in [1, m + 2/d)$ if $d \geq 2$ and $s \in [1, m + 1)$ if $d = 1$. With this additional bound, we can conclude the

higher-order estimates in the last three steps of the proof. We only detail the proof for $d \geq 2$, the case $d = 1$ being similar.

Step 1: In this step we establish for $\rho \in (m, md/(d-2))$ the bound

$$\|u^<\|_{L_t^m L_x^\rho} \lesssim \|u_0\|_{L_x^1} + \|S\|_{L_{t,x}^1} + 1. \quad (5-3)$$

Set $g_0 := \varphi \psi_0 \delta_{v=u(t,x)} S + \partial_t \varphi f^< - \varphi q \partial_v \psi_0$, $g_1 := \varphi q^<$, and

$$\sigma_x := \frac{d}{m} - \frac{d}{\rho} \in \left(0, \frac{2}{m}\right).$$

Consequently, we may choose $\gamma \in (0, 1)$ so large that

$$\sigma_x \in \left[0, \frac{m-1}{m} \frac{2}{m-\gamma}\right).$$

From [Corollary 4.5](#) applied with $\mu = 1$ and $q = m$ we obtain

$$\begin{aligned} \|\varphi u^<\|_{L_t^m W_x^{\sigma_x, m}} &\lesssim \| |v|^{1-\gamma} g_0 \|_{\mathcal{M}_{TV}} + \| |v|^{-\gamma} g_1 \|_{\mathcal{M}_{TV}} + \|\varphi f^<\|_{L_{t,x,v}^1 \cap L_{t,x,v}^\infty} + \|\varphi u^<\|_{L_{t,x}^1 \cap L_t^m L_x^1} \\ &\lesssim \| |v|^{1-\gamma} g_0 \|_{\mathcal{M}_{TV}} + \| |v|^{-\gamma} g_1 \|_{\mathcal{M}_{TV}} + \|f\|_{L_{t,x,v}^1 \cap L_{t,x,v}^\infty} + \sup_{t \in [0, T]} \|u(t)\|_{L_x^1}. \end{aligned}$$

We note that since trivially $f^< \in L_{t,x,v}^\infty$ with norm bounded by 1 we have by [Theorem A.2](#)

$$\|f\|_{L_{t,x,v}^1 \cap L_{t,x,v}^\infty} + \sup_{t \in [0, T]} \|u(t)\|_{L_x^1} \lesssim \|u\|_{L_{t,x}^1} + 1 + \sup_{t \in [0, T]} \|u(t)\|_{L_x^1} \lesssim \|u_0\|_{L_x^1} + \|S\|_{L_{t,x}^1} + 1.$$

Next, we check that $|v|^{1-\gamma} g_0 \in \mathcal{M}_{TV}$. Indeed, since $|v|^{1-\gamma}$ can be estimated by a constant on the supports of ψ_0 and $\partial_v \psi_0$, we may apply [Lemma A.4](#) to the effect of

$$\begin{aligned} \| |v|^{1-\gamma} g_0 \|_{\mathcal{M}_{TV}} &= \| |v|^{1-\gamma} (\varphi \psi_0 \delta_{v=u(t,x)} S + \partial_t \varphi f^< - \varphi q \partial_v \psi_0) \|_{\mathcal{M}_{TV}} \\ &\lesssim \|S\|_{L_{t,x}^1} + \|\partial_t \varphi |u|\|_{L_{t,x}^1} + \|q \partial_v \psi_0\|_{\mathcal{M}_{TV}} \\ &\lesssim \|\partial_t \varphi |u|\|_{L_{t,x}^1} + \|u_0\|_{L_x^1} + \|S\|_{L_{t,x}^1}. \end{aligned}$$

Utilizing [Lemma A.4](#) once more to the effect of

$$\| |v|^{-\gamma} g_1 \|_{\mathcal{M}_{TV}} \lesssim \| |v|^{-\gamma} q^< \|_{\mathcal{M}_{TV}} \lesssim \|u_0\|_{L_x^1} + \|S\|_{L_{t,x}^1},$$

we obtain by Sobolev embedding

$$\|\varphi u^<\|_{L_t^m L_x^\rho} \lesssim \|\varphi u^<\|_{L_t^m W_x^{\sigma_x, m}} \lesssim \|u_0\|_{L_x^1} + \|\partial_t \varphi |u|\|_{L_{t,x}^1} + \|S\|_{L_{t,x}^1} + 1. \quad (5-4)$$

With the same construction $\varphi_n \rightarrow 1_{[0, T]}$ as in the proof of [Theorem 1.2](#), this gives (5-3).

Step 2: Next, we investigate $u^>$ and establish for $\eta \in (1, m)$ and

$$\eta^* = \frac{\eta d(m-1)}{d(m-1) - 2(\eta-1)}$$

the bound

$$\|u^>\|_{L_t^\eta L_x^{\eta^*}} \lesssim \|u_0\|_{L_x^1} + \|S\|_{L_{t,x}^1} + 1. \quad (5-5)$$

Set $g_0 := \varphi \psi_1 \delta_{v=u(t,x)} S + \partial_t \varphi f^> + \varphi q \partial_v \psi_0$ and $g_1 := \varphi q^>$. Choose $\gamma \in (1, m)$ sufficiently small, so that $\eta \in (1, m + 1 - \gamma)$, and define

$$\sigma_x := \frac{\eta-1}{\eta} \frac{2}{m-1} \in \left(0, \frac{\eta-1}{\eta} \frac{2}{m-\gamma}\right).$$

We apply [Corollary 4.5](#) with $\mu = 1$ and $q = \eta$, which gives

$$\begin{aligned} \|\varphi u^>\|_{L_t^\eta W_x^{\sigma_x, \eta}} &\lesssim \| |v|^{1-\gamma} g_0 \|_{\mathcal{M}_{TV}} + \| |v|^{-\gamma} g_1 \|_{\mathcal{M}_{TV}} + \|\varphi f^>\|_{L_{t,x,v}^1 \cap L_{t,x,v}^\infty} + \|\varphi u^>\|_{L_{t,x}^1 \cap L_t^\eta L_x^1} \\ &\lesssim \| |v|^{1-\gamma} g_0 \|_{\mathcal{M}_{TV}} + \| |v|^{-\gamma} g_1 \|_{\mathcal{M}_{TV}} + \|f\|_{L_{t,x,v}^1 \cap L_{t,x,v}^\infty} + \sup_{t \in [0, T]} \|u(t)\|_{L_x^1}. \end{aligned}$$

The terms involving f and u are estimated as in Step 1. Further, since $|v|^{1-\gamma}$ can be estimated by a constant on the support of ψ_1 and $\partial_v \psi_0$, we have by [Lemma A.4](#)

$$\begin{aligned} \| |v|^{1-\gamma} g_0 \|_{\mathcal{M}_{TV}} &= \| |v|^{1-\gamma} (\varphi \psi_1 \delta_{v=u(t,x)} S + \partial_t \varphi f^> + \varphi q \partial_v \psi_0) \|_{\mathcal{M}_{TV}} \\ &\lesssim \|S\|_{L_{t,x}^1} + \|\partial_t \varphi |u|\|_{L_{t,x}^1} + \|q \partial_v \psi_0\|_{\mathcal{M}_{TV}} \\ &\lesssim \|\partial_t \varphi |u|\|_{L_{t,x}^1} + \|u_0\|_{L_x^1} + \|S\|_{L_{t,x}^1}, \end{aligned}$$

and, again due to [Lemma A.4](#),

$$\| |v|^{-\gamma} g_1 \|_{\mathcal{M}_{TV}} \lesssim \| |v|^{-\gamma} q^> \|_{\mathcal{M}_{TV}} \lesssim \|u_0\|_{L_x^1} + \|S\|_{L_{t,x}^1}.$$

Since $\eta^* = \eta d / (d - \sigma_x \eta)$, we have by Sobolev embedding $W_x^{\sigma_x, \eta} \subset L_x^{\eta^*}$, and hence

$$\|\varphi u^>\|_{L_t^\eta L_x^{\eta^*}} \lesssim \|\varphi u^>\|_{L_t^\eta W_x^{\sigma_x, \eta}} \lesssim \|u_0\|_{L_x^1} + \|\partial_t \varphi |u|\|_{L_{t,x}^1} + \|S\|_{L_{t,x}^1} + 1.$$

With the same construction $\varphi_n \rightarrow 1_{[0, T]}$ as before, this yields [\(5-5\)](#).

Step 3: In this step, we show that for $s \in [1, m + 2/d)$ we have

$$\|u\|_{L_{t,x}^s} \lesssim \|u_0\|_{L_x^1} + \|S\|_{L_{t,x}^1} + 1. \quad (5-6)$$

Observe that it suffices to show the assertion for $s > m$, since $u \in L^1(0, T; L^1(\mathbb{R}^d))$ is already established by [Theorem A.2](#).

Define

$$\rho := \frac{m}{m+1-s} \in \left(m, \frac{md}{d-2}\right).$$

For $\vartheta \in (0, 1)$, it holds $[L_t^\infty L_x^1, L_t^m L_x^\rho]_\vartheta = L_t^{p_\vartheta} L_x^{q_\vartheta}$ with

$$\frac{1}{p_\vartheta} = \frac{\vartheta}{m} \quad \text{and} \quad \frac{1}{q_\vartheta} = 1 - \vartheta + \frac{\vartheta}{\rho}.$$

Choosing

$$\vartheta := \frac{m\rho}{m\rho + \rho - m} \in (0, 1),$$

we obtain $p_\vartheta = q_\vartheta = s$, and hence by [\(5-3\)](#) and [Theorem A.2](#)

$$\|u^<\|_{L_{t,x}^s} \lesssim \|u^<\|_{L_t^\infty L_x^1} + \|u^<\|_{L_t^m L_x^\rho} \lesssim \|u_0\|_{L_x^1} + \|S\|_{L_{t,x}^1} + 1. \quad (5-7)$$

Next, we define

$$\eta := \frac{sd(m-1)+2}{d(m-1)+2} \in (1, m) \quad \text{and} \quad \eta^* = \frac{\eta d}{d-2(\eta-1)/(m-1)}$$

and observe that for $\vartheta \in (0, 1)$, it holds $[L_t^\infty L_x^1, L_t^\eta L_x^{\eta^*}]_{\vartheta} = L_t^{p\vartheta} L_x^{q\vartheta}$ with

$$\frac{1}{p\vartheta} = \frac{\vartheta}{\eta} \quad \text{and} \quad \frac{1}{q\vartheta} = 1 - \vartheta + \frac{\vartheta}{\eta^*}.$$

Choosing

$$\vartheta := \frac{\eta d(m-1)}{\eta d(m-1) + 2(\eta-1)} \in (0, 1),$$

we obtain $p\vartheta = q\vartheta = s$, and hence by (5-5) and Theorem A.2

$$\|u^>\|_{L_{t,x}^s} \lesssim \|u^>\|_{L_t^\infty L_x^1} + \|u^>\|_{L_t^\eta L_x^{\eta^*}} \lesssim \|u_0\|_{L_x^1} + \|S\|_{L_{t,x}^1} + 1. \quad (5-8)$$

Combining (5-7) and (5-8), we obtain (5-6).

Step 4: In this step we argue that

$$\|\varphi u^<\|_{W^{\sigma_t, p}(W^{\sigma_x, p})} \lesssim \|\partial_t \varphi |u|\|_{L_{t,x}^1} + \|u_0\|_{L_x^1}^m + \|S\|_{L_{t,x}^1}^m + 1.$$

Indeed, we choose $\gamma \in (0, 1)$ so large that

$$\sigma_x < \frac{p-2+\gamma}{p} \frac{2}{m-1}$$

and $m+1-\gamma < m+2/d$. Then we apply Corollary 4.7 with $g_0 := \varphi \psi_0 \delta_{v=u(t,x)} S + \partial_t \varphi f^< - \varphi q \partial_v \psi_0$, $g_1 := \varphi q^<$ and $\tilde{p} = p$. We obtain by (4-19) some $r \in (p, m+1-\gamma)$ such that

$$\begin{aligned} \|\varphi u^<\|_{W^{\sigma_t, p}(W^{\sigma_x, p})} &\lesssim \|g_0\|_{\mathcal{M}_{TV}} + \| |v|^{1-\gamma} g_0 \|_{\mathcal{M}_{TV}} + \| |v|^{-\gamma} g_1 \|_{\mathcal{M}_{TV}} \\ &\quad + \|f\|_{L_{t,x,v}^1 \cap L_{t,x,v}^\infty} + \|u\|_{L_t^1 L_x^p \cap L_{t,x}^r} + \| |u|^m \|_{L_{t,x}^1}. \end{aligned}$$

The first four terms on the right-hand side can be estimated as in Step 1 (indeed, we did not use the coefficient $|v|^{1-\gamma}$ in the estimate of g_0) via

$$\|g_0\|_{\mathcal{M}_{TV}} + \| |v|^{1-\gamma} g_0 \|_{\mathcal{M}_{TV}} + \| |v|^{-\gamma} g_1 \|_{\mathcal{M}_{TV}} + \|f\|_{L_{t,x,v}^1 \cap L_{t,x,v}^\infty} \lesssim \|\partial_t \varphi |u|\|_{L_{t,x}^1} + \|u_0\|_{L_x^1} + \|S\|_{L_{t,x}^1} + 1,$$

while the last two terms are estimated in light of $r < m+1-\gamma < m+2/d$ through (5-6) as

$$\|u\|_{L_t^1 L_x^p \cap L_{t,x}^r} + \| |u|^m \|_{L_{t,x}^1} \lesssim \|u\|_{L_{t,x}^p \cap L_{t,x}^r} + \|u\|_{L_{t,x}^m}^m \lesssim \|u_0\|_{L_x^1}^m + \|S\|_{L_{t,x}^1}^m + 1.$$

Step 5: In this step we establish

$$\|\varphi u^>\|_{W^{\sigma_t, p}(W^{\sigma_x, p})} \lesssim \|\partial_t \varphi |u|\|_{L_{t,x}^1} + \|u_0\|_{L_x^1}^m + \|S\|_{L_{t,x}^1}^m + 1. \quad (5-9)$$

Assume first $p < m$. Choose $\gamma \in (1, m)$ so small that $p \in (1, m+1-\gamma)$ and

$$\sigma_t < \frac{m+1-\gamma-p}{p} \frac{1}{m-1}$$

and apply [Corollary 4.7](#) with $g_0 := \varphi \psi_1 \delta_{v=u(t,x)} S + \partial_t \varphi f^> + \varphi q \partial_v \psi_0$, $g_1 := \varphi q^>$ and $\tilde{p} = p$. Estimate (4-19) gives

$$\begin{aligned} \|\varphi u^>\|_{W^{\sigma_t,p}(W^{\sigma_x,p})} &\lesssim \|g_0\|_{\mathcal{M}_{TV}} + \| |v|^{1-\gamma} g_0 \|_{\mathcal{M}_{TV}} + \| |v|^{-\gamma} g_1 \|_{\mathcal{M}_{TV}} \\ &\quad + \|f\|_{L^1_{t,x,v} \cap L^\infty_{t,x,v}} + \|u\|_{L^1_t L^p_x \cap L^r_{t,x}} + \| |u|^m \|_{L^1_{t,x}}. \end{aligned}$$

The first four terms on the right-hand side are estimated as in Step 2 via

$$\|g_0\|_{\mathcal{M}_{TV}} + \| |v|^{1-\gamma} g_0 \|_{\mathcal{M}_{TV}} + \| |v|^{-\gamma} g_1 \|_{\mathcal{M}_{TV}} + \|f\|_{L^1_{t,x,v} \cap L^\infty_{t,x,v}} \lesssim \|\partial_t \varphi |u|\|_{L^1_{t,x}} + \|u_0\|_{L^1_x} + \|S\|_{L^1_{t,x}} + 1,$$

while the last two terms are estimated through (5-6) as

$$\|u\|_{L^1_t L^p_x \cap L^r_{t,x}} + \| |u|^m \|_{L^1_{t,x}} \lesssim \|u\|_{L^p_{t,x} \cap L^r_{t,x}} + \|u\|_{L^m_{t,x}}^m \lesssim \|u_0\|_{L^1_x}^m + \|S\|_{L^1_{t,x}}^m + 1.$$

Hence, we have shown (5-9) in the case $p \in (1, m)$. If $p = m$, we choose $p_0 \in (1, m)$ sufficiently large such that for

$$\kappa_x(p_0) := \frac{p_0 - 1}{p_0} \frac{2}{m - 1}$$

it holds

$$\kappa_x(p_0) - \frac{d}{p_0} > \sigma_x - \frac{d}{m}.$$

We observe that for

$$\kappa_t(p_0) := \frac{m - p_0}{p_0} \frac{1}{m - 1}$$

it holds

$$\kappa_t(p_0) - \frac{1}{p_0} > \sigma_t - \frac{1}{m}$$

due to $p_0 < m$ (indeed, we have necessarily $\sigma_t = 0$). Choosing sufficiently large $\sigma_x(p_0) < \kappa_x(p_0)$ and $\sigma_t(p_0) < \kappa_t(p_0)$, we conclude by Sobolev embedding

$$\begin{aligned} \|\varphi u^>\|_{L^m_t(W^{\sigma_x,m}_x)} &\lesssim \|\varphi u^>\|_{W^{\sigma_t(p_0),p_0}(W^{\sigma_x(p_0),p_0})} \\ &\lesssim \|\partial_t \varphi |u|\|_{L^1_{t,x}} + \|u_0\|_{L^1_x}^m + \|S\|_{L^1_{t,x}}^m + 1, \end{aligned}$$

which is (5-9) in the case $p = m$.

Step 6: Conclusion. With the same construction $\varphi_n \rightarrow 1_{[0,T]}$ as in the proof of [Theorem 1.2](#), Steps 4 and 5 combine to

$$\begin{aligned} \sup_{n \in \mathbb{N}} \|\varphi_n u\|_{W^{\sigma_t,p}(W^{\sigma_x,p})} &\lesssim \sup_{n \in \mathbb{N}} \|\varphi_n u^<\|_{W^{\sigma_t,p}(W^{\sigma_x,p})} + \sup_{n \in \mathbb{N}} \|\varphi_n u^>\|_{W^{\sigma_t,p}(W^{\sigma_x,p})} \\ &\lesssim \|u_0\|_{L^1_x}^m + \|S\|_{L^1_{t,x}}^m + 1. \end{aligned}$$

Since $\varphi_n u \rightarrow u 1_{[0,T]}$ in the sense of distributions, we obtain (1-2) by the weak lower semicontinuity of the norm in $W^{\sigma_t,p}(0, T; W^{\sigma_x,p}(\mathbb{R}^d))$. Estimate (1-3) follows analogously to the proof of [Corollary 1.3\(ii\)](#). \square

Appendix A: Kinetic solutions

In this section we recall some details of the concept of entropy/kinetic solutions and their well-posedness for partial differential equations of the type

$$\begin{aligned} \partial_t u + \operatorname{div} A(u) &= \operatorname{div}(b(u) \nabla u) + S(t, x) \quad \text{on } (0, T) \times \mathbb{R}_x^d \\ u(0) &= u_0 \quad \text{on } \mathbb{R}_x^d, \end{aligned} \quad (\text{A-1})$$

where

$$\begin{aligned} u_0 &\in L^1(\mathbb{R}_x^d), \quad S \in L^1([0, T] \times \mathbb{R}_x^d), \quad T \geq 0, \\ a &:= A' \in C(\mathbb{R}; \mathbb{R}^d) \cap C^1(\mathbb{R} \setminus \{0\}; \mathbb{R}^d), \\ b &= (b_{jk})_{j,k=1,\dots,d} \in C(\mathbb{R}; S_+^{d \times d}) \cap C^1(\mathbb{R} \setminus \{0\}; S_+^{d \times d}). \end{aligned} \quad (\text{A-2})$$

Here, $S_+^{d \times d}$ denotes the space of symmetric, nonnegative definite matrices. For $b = (b)_{i,j=1,\dots,d} \in S_+^{d \times d}$ we set $\sigma = b^{1/2}$, that is, $b_{i,j} = \sum_{k=1}^d \sigma_{i,k} \sigma_{k,j}$. For a locally bounded function $b: \mathbb{R} \rightarrow S_+^{d \times d}$ we let $\beta_{i,k}$ be such that $\beta'_{i,k}(v) = \sigma_{i,k}(v)$. Similarly, for $\psi \in C_c^\infty(\mathbb{R}_v)$ we let $\beta_{i,k}^\psi$ be such that $(\beta_{i,k}^\psi)'(v) = \psi(v) \sigma_{i,k}(v)$. The corresponding kinetic form of (A-1) reads, see [Chen and Perthame 2003],

$$\begin{aligned} \mathcal{L}(\partial_t, \nabla_x, v) f(t, x, v) &= \partial_t f + a(v) \cdot \nabla_x f - \operatorname{div}(b(v) \nabla_x f) \\ &= \partial_v q + S(t, x) \delta_{u(t,x)=v}(v), \end{aligned}$$

where $q \in \mathcal{M}^+$ and \mathcal{L} is identified with the symbol

$$\mathcal{L}(i\tau, i\xi, v) := i\tau + a(v) \cdot i\xi - (b(v)\xi, \xi). \quad (\text{A-3})$$

We will use the terms kinetic and entropy solution synonymously. From [Chen and Perthame 2003] we recall the definition of entropy/kinetic solutions to (A-1).

Definition A.1. We say that $u \in C([0, T]; L^1(\mathbb{R}^d))$ is an entropy solution to (A-1) if the corresponding kinetic function f satisfies:

(i) For any nonnegative $\psi \in \mathcal{D}(\mathbb{R})$, $k = 1, \dots, d$,

$$\sum_{i=1}^d \partial_{x_i} \beta_{ik}^\psi(u) \in L^2([0, T] \times \mathbb{R}^d).$$

(ii) For any two nonnegative functions $\psi_1, \psi_2 \in \mathcal{D}(\mathbb{R})$, $k = 1, \dots, d$,

$$\sqrt{\psi_1(u(t, x))} \sum_{i=1}^d \partial_{x_i} \beta_{ik}^{\psi_2}(u(t, x)) = \sum_{i=1}^d \partial_{x_i} \beta_{ik}^{\psi_1 \psi_2}(u(t, x)) \quad \text{a.e.}$$

(iii) There are nonnegative measures $m, n \in \mathcal{M}^+$ such that, in the sense of distributions,

$$\partial_t f + a(v) \cdot \nabla_x f - \operatorname{div}(b(v) \nabla_x f) = \partial_v(m + n) + \delta_{v=u(t,x)} S \quad \text{on } (0, T) \times \mathbb{R}_x^d \times \mathbb{R}_v,$$

where n is defined by

$$\int \psi(v) n(t, x, v) \, dv = \sum_{k=1}^d \left(\sum_{i=1}^d \partial_{x_i} \beta_{ik}^\psi(u(t, x)) \right)^2$$

for any $\psi \in \mathcal{D}(\mathbb{R})$ with $\psi \geq 0$.

(iv) We have

$$\int (m + n) \, dx \, dt \leq \mu(v) \in L_0^\infty(\mathbb{R}),$$

where L_0^∞ is the space of L^∞ -functions vanishing for $|v| \rightarrow \infty$.

The well-posedness of entropy solutions to (A-1) follows along the same lines as [Chen and Perthame 2003]. In this form, it can be found in [Gess 2020].

Theorem A.2. *Let $u_0 \in L^1(\mathbb{R}^d)$ and $S \in L^1([0, T] \times \mathbb{R}^d)$. Then there is a unique entropy solution u to (A-1) satisfying $u \in C([0, T]; L^1(\mathbb{R}^d))$. For two entropy solutions u^1, u^2 with initial conditions u_0^1, u_0^2 and forcing S^1, S^2 we have*

$$\sup_{t \in [0, T]} \|u^1(t) - u^2(t)\|_{L^1(\mathbb{R}^d)} \leq \|u_0^1 - u_0^2\|_{L^1(\mathbb{R}^d)} + \|S^1 - S^2\|_{L^1([0, T] \times \mathbb{R}^d)}.$$

Furthermore, the following a priori estimate was given in Lemma 2.3 in [Gess 2020].

Lemma A.3. *Let u be the unique entropy solution to (A-1) with $u_0 \in (L^1 \cap L^{2-\gamma})(\mathbb{R}_x^d)$ and $S \in (L^1 \cap L^{2-\gamma})([0, T] \times \mathbb{R}_x^d)$ for some $\gamma \in (-\infty, 1)$. Then, there is a constant $C = C(T, g) \geq 0$ such that*

$$\sup_{t \in [0, T]} \|u(t)\|_{L_x^{2-\gamma}}^{2-\gamma} + (1-\gamma) \int_0^T \int_{\mathbb{R}^{d+1}} |v|^{-\gamma} q \, dv \, dx \, dr \leq C(\|u_0\|_{L_x^{2-\gamma}}^{2-\gamma} + \|S\|_{L_{t,x}^{2-\gamma}}^{2-\gamma}).$$

In the case of L^1 initial data a different proof for the existence of singular moments of the kinetic measure q is needed.

Lemma A.4. *Let u be the unique entropy solution to (A-1) with $u_0 \in L^1(\mathbb{R}_x^d)$ and $S \in L^1([0, T] \times \mathbb{R}_x^d)$. Then, the map*

$$v \mapsto \int_0^T \int_{\mathbb{R}_x^d} q(r, x, v) \, dx \, dr$$

is continuous and, for all $v_0 \in \mathbb{R}_v$, we have

$$\begin{aligned} \int_0^T \int_{\mathbb{R}_x^d} q(r, x, v_0) \, dx \, dr &\leq \int_{\mathbb{R}_x^d} (\operatorname{sgn}(v_0)(u_0 - v_0))_+ \, dx + \int_0^T \int_{\mathbb{R}_x^d} \operatorname{sgn}_+(\operatorname{sgn}(v_0)(u - v_0)) S \, dx \, dr \\ &\leq \int_{\mathbb{R}_x^d} |u_0| \, dx + \int_0^T \int_{\mathbb{R}_x^d} |S| \, dx \, dr. \end{aligned} \tag{A-4}$$

Proof. In the proof, we use the short-hand notation $\bar{g}(v) := \int_0^T \int_{\mathbb{R}_x^d} g(r, x, v) \, dx \, dr$ for a generic $g : (r, x, v) \mapsto g(r, x, v)$. We first argue that \bar{q} has left and right limits. Indeed, by a standard approximation

argument, the kinetic formulation yields, for every $\eta \in C_c^\infty(\mathbb{R}_v)$,

$$\begin{aligned} \int_{\mathbb{R}_v} \eta' \bar{q} \, dv &= - \int_{\mathbb{R}_v} \eta \left(\int_{\mathbb{R}_x^d} f \, dx \Big|_0^T \right) dv + \overline{\eta(u)S} \\ &= - \int_{\mathbb{R}_v} \eta \left(\int_{\mathbb{R}_x^d} f \, dx \Big|_0^T \right) dv + \int_{\mathbb{R}_v} \eta' \overline{1_{v < u} S} \, dv. \end{aligned} \quad (\text{A-5})$$

Since $v \mapsto \int_{\mathbb{R}_x^d} f(r, x, v) \, dx \Big|_0^T$ is in $L^1(\mathbb{R}_v)$, this implies $\bar{q} - \overline{1_{v < u} S} \in \dot{W}^{1,1}(\mathbb{R}_v)$. Since $\overline{1_{v < u} S} \in BV_{\text{loc}}(\mathbb{R}_v)$, this shows $\bar{q} \in BV_{\text{loc}}(\mathbb{R}_v)$ and thus the existence of left and right limits.

Next we claim that (A-5) continues to hold for all $\eta \in C^\infty(\mathbb{R}_v)$ with $\eta' \in C_c^\infty(\mathbb{R}_v)$. For $R > 0$ let $\varphi_R \in C_c^\infty(\mathbb{R}_v)$ be such that $\varphi_R(v) = 1$ for $|v| \leq R$, $\text{supp } \varphi_R \subset [-(R+1), R+1]$ and $|\varphi_R| + |\varphi'_R| \lesssim 1$. Defining $\eta_R := \eta \varphi_R$, we have by (A-5)

$$\int_{\mathbb{R}_v} (\eta' \varphi_R + \eta \varphi'_R) \bar{q} \, dv = - \int_{\mathbb{R}_v} \eta_R \left(\int_{\mathbb{R}_x^d} f \, dx \Big|_0^T \right) dv + \overline{\eta_R(u)S}.$$

Since η_R is uniformly bounded in R , $\eta_R \rightarrow \eta$ locally uniformly, $v \mapsto \int_{\mathbb{R}_x^d} f(r, x, v) \, dx \Big|_0^T$ is in $L^1(\mathbb{R}_v)$ and $S \in L^1([0, T] \times \mathbb{R}_x^d)$, we may take the limit $R \rightarrow \infty$ on the right-hand side by dominated convergence. Again by dominated convergence the contribution from the term $\eta' \varphi_R$ to the left-hand side converges, since η' has compact support and $\bar{q} \in BV_{\text{loc}}(\mathbb{R}_v) \subset L^1_{\text{loc}}(\mathbb{R}_v)$. Moreover, the contribution from the term $\eta \varphi'_R$ vanishes for $R \rightarrow \infty$, since both η and φ'_R are bounded, $\text{supp } \varphi'_R \subset [-(R+1), -R] \cup [R, R+1]$ and $\bar{q} \in L^\infty_0(\mathbb{R}_v)$ by Definition A.1(iv).

We are now in the position to conclude. Assume first $v_0 \in \mathbb{R}_+$. Let $\phi_\pm \in C_c^\infty(\mathbb{R}_v)$ with $\phi_\pm \geq 0$, $\text{supp } \phi_+ \subset [0, 1]$, $\text{supp } \phi_- \subset [-1, 0]$, $\int_{\mathbb{R}_v} \phi_\pm \, dv = 1$ and define $\phi_\pm^\varepsilon(v) = \varepsilon^{-1} \phi_\pm(\varepsilon^{-1}v)$ for $\varepsilon > 0$. Moreover let η_\pm^ε be such that $(\eta_\pm^\varepsilon)'(v) = \phi_\pm^\varepsilon(v - v_0)$ and $(\eta_\pm^\varepsilon)(v_0) = 0$. Observe that $(\eta_\pm^\varepsilon)' \rightarrow \delta_{v=v_0}$ and $\eta_\pm^\varepsilon(v) \rightarrow \text{sgn}_+(v - v_0)$ as $\varepsilon \searrow 0$ independent of the choice of \pm . Choosing now $\eta := \eta_\pm^\varepsilon$ in (A-5) and using dominated convergence to take the limit $\varepsilon \searrow 0$, we obtain

$$\begin{aligned} \bar{q}(v_0 \pm) &= - \int_{\mathbb{R}_x^d} (u - v_0)_+ \, dx \Big|_0^T + \int_0^T \int_{\mathbb{R}_x^d} \text{sgn}_+(u - v_0) S \, dx \, dr \\ &\leq \int_{\mathbb{R}_x^d} (u_0 - v_0)_+ \, dx + \int_0^T \int_{\mathbb{R}_x^d} \text{sgn}_+(u - v_0) S \, dx \, dr. \end{aligned}$$

In particular $\bar{q}(v_0-) = \bar{q}(v_0+)$, so that \bar{q} is continuous. The case $v_0 \in \mathbb{R}_-$ is treated analogously replacing the conditions $\phi_\pm \geq 0$ and $\int_{\mathbb{R}_v} \phi_\pm \, dv = 1$ by $\phi_\pm \leq 0$ and $\int_{\mathbb{R}_v} \phi_\pm \, dv = -1$, respectively, so that $\eta_\pm^\varepsilon(v) \rightarrow \text{sgn}_+(v - v_0)$ is replaced by $\eta_\pm^\varepsilon(v) \rightarrow \text{sgn}_+(-(v - v_0))$. \square

Appendix B: Fourier multipliers

In this section, we provide some Fourier multiplier results well-adapted to our averaging lemma, Lemma 4.2. We recall the definition of \mathbb{R}^{d+1} and of the functions η_l and φ_j given in Section 2, and define $\tilde{\eta}_l := \eta_{l-1} + \eta_l + \eta_{l+1}$ and $\tilde{\varphi}_j := \varphi_{j-1} + \varphi_j + \varphi_{j+1}$. We observe $\tilde{\eta}_l(2^l \cdot) = \tilde{\eta}_0$ and $\tilde{\varphi}(2^j \cdot) = \tilde{\varphi}_0$. Moreover, $\tilde{\eta}_l$ and $\tilde{\varphi}_j$ are identically unity on the support of η_l and φ_j , respectively.

Theorem B.1. Let $k = 2 + 2[1 + d/2]$. Let $m : \dot{\mathbb{R}}^{d+1} \rightarrow \mathbb{C}$ be k -times differentiable and such that for all $\alpha = (\alpha_\tau, \alpha_\xi) \in \mathbb{N}_0 \times \mathbb{N}_0^d$ with $|\alpha| \leq k$ there is a constant C_α such that for all $(\tau, \xi) \in \dot{\mathbb{R}}^{d+1}$

$$|\partial_\tau^{\alpha_\tau} \partial_\xi^{\alpha_\xi} m(\tau, \xi)| \leq C_\alpha |\tau|^{-\alpha_\tau} |\xi|^{-|\alpha_\xi|}. \quad (\text{B-1})$$

Then there is a constant $C > 0$, depending only on the constants C_α , such that for any $p \in [1, \infty]$ and all $l, j \in \mathbb{Z}$, we have

$$\|\tilde{\eta}_l \tilde{\varphi}_j m\|_{\mathcal{M}^p} \leq C; \quad (\text{B-2})$$

i.e., $\tilde{\eta}_l \tilde{\varphi}_j m$ (more precisely the mapping $(\tau, \xi) \mapsto \tilde{\eta}_l(\tau) \tilde{\varphi}_j(\xi) m(\tau, \xi)$) extends to an $L^p_{t,x}$ -multiplier with a norm independent of l and j . Furthermore, this mapping extends to an \mathcal{M}_{TV} -multiplier with the same norm bound.

Proof. Since $\|\cdot\|_{\mathcal{M}^p} \leq \|\cdot\|_{\mathcal{M}^1}$, it suffices to estimate the L^1 multiplier norm of $\tilde{\eta}_l \tilde{\varphi}_j m$ in order to obtain (B-2). Since multiplier norms are invariant under dilation and since $\|m\|_{\mathcal{M}^1}$ is equal to the total mass of $\mathcal{F}^{-1}m$, see [Bergh and Löfström 1976, Theorem 6.1.2], we have

$$\|\tilde{\eta}_l \tilde{\varphi}_j m\|_{\mathcal{M}^1} = \|\tilde{\eta}_0 \tilde{\varphi}_0 m_{l,j}\|_{\mathcal{M}^1} = \|\mathcal{F}_{t,x}^{-1} \tilde{\eta}_0 \tilde{\varphi}_0 m_{l,j}\|_{L^1_{t,x}},$$

where $m_{l,j}(\tau, \xi) := m(2^l \tau, 2^j \xi)$. Let $M := [1 + d/2]$. We observe

$$\begin{aligned} & (1+t^2)(1+|x|^2)^M \mathcal{F}_{t,x}^{-1}[\tilde{\eta}_0 \tilde{\varphi}_0 m_{l,j}](t, x) \\ &= c_d \int_{\mathbb{R}_t \times \mathbb{R}_x^d} (\text{id} - \partial_\tau^2)(\text{id} - \Delta_\xi)^M (e^{it\tau + ix \cdot \xi}) \tilde{\eta}_0(\tau) \tilde{\varphi}_0(\xi) m(2^l \tau, 2^j \xi) d\xi d\tau \\ &= c_d \int_{\mathbb{R}_t \times \mathbb{R}_x^d} e^{it\tau + ix \cdot \xi} (\text{id} - \partial_\tau^2)(\text{id} - \Delta_\xi)^M (\tilde{\eta}_0(\tau) \tilde{\varphi}_0(\xi) m(2^l \tau, 2^j \xi)) d\xi d\tau \\ &= \sum_{\substack{\alpha_\tau + \beta_\tau \leq 2 \\ |\alpha_\xi| + |\beta_\xi| \leq 2M}} c_{d,\alpha,\beta} 2^{l\beta_\tau} 2^{j|\beta_\xi|} \int_{\mathbb{R}_t \times \mathbb{R}_x^d} e^{it\tau + ix \cdot \xi} \partial_\tau^{\alpha_\tau} \tilde{\eta}_0(\tau) \partial_\xi^{\alpha_\xi} \tilde{\varphi}_0(\xi) \partial_\tau^{\beta_\tau} \partial_\xi^{\beta_\xi} m(2^l \tau, 2^j \xi) d\xi d\tau, \end{aligned}$$

where c_d and $c_{d,\alpha,\beta}$ are constants that do not depend on l and j . On $\text{supp } \tilde{\eta}_0 \times \text{supp } \tilde{\varphi}_0$ we have $|\partial_\tau^{\beta_\tau} \partial_\xi^{\beta_\xi} m(2^l \tau, 2^j \xi)| \leq C_\beta 2^{-l\beta_\tau} 2^{-j|\beta_\xi|}$, and hence we obtain

$$(1+t^2)(1+|x|^2)^M |\mathcal{F}_{t,x}^{-1}[\tilde{\eta}_0 \tilde{\varphi}_0 m_{l,j}](t, x)| \leq c.$$

Since $2M > d$, it follows $\|\mathcal{F}_{t,x}^{-1}[\tilde{\eta}_0 \tilde{\varphi}_0 m_{l,j}]\|_{L^1_{t,x}} \leq C$, which yields (B-2). In particular, $\tilde{\eta}_l \tilde{\varphi}_j m$ is an L^1 -multiplier with a norm bound independent of l and j , and as such extends to a multiplier on \mathcal{M}_{TV} with the same norm bound. \square

Remark B.2. In Theorem B.1, the assumptions on the differentiability of m may be relaxed: Indeed, the proof shows that it suffices to assume that m is a continuous function such that $\partial_\tau^{\alpha_\tau} m$, $\partial_\xi^{\alpha_\xi} m$ and $\partial_\tau^{\alpha_\tau} \partial_\xi^{\alpha_\xi} m$ exist for all $\alpha = (\alpha_\tau, \alpha_\xi)$ with $\alpha_\tau \leq 2$ and $|\alpha_\xi| \leq 2[1 + d/2]$, and that (B-1) holds for these choices of α .

Remark B.3. Clearly, Theorem B.1 has an isotropic variant; see [Bahouri, Chemin, and Danchin 2011, Lemma 2.2]. More precisely, a simple adaptation of the proof shows the following: Let $k = 2[1 + d/2]$. Let $m : \mathbb{R}^d \setminus \{0\} \rightarrow \mathbb{C}$ be k -times differentiable and such that for all $\alpha \in \mathbb{N}_0^d$ with $|\alpha| \leq k$ there is a

constant C_α such that for all $\xi \in \mathbb{R}^d \setminus \{0\}$ we have $|\partial^\alpha m(\xi)| \leq C_\alpha |\xi|^{-|\alpha|}$. Then there is a constant $C > 0$, depending only on the constants C_α such that for any $p \in [1, \infty]$ and all $j \in \mathbb{Z}$ we have $\|\tilde{\varphi}_j m\|_{\mathcal{M}^p} \leq C$. Again, $\tilde{\varphi}_j m$ extends to an \mathcal{M}_{TV} -multiplier (in ξ) with the same norm bound.

Lemma B.4. *Let \mathcal{L} be defined as in (4-2) and fix $\alpha = (\alpha_\tau, \alpha_\xi) \in \mathbb{N}_0 \times \mathbb{N}_0^d$. Then we have for all $(\tau, \xi, v) \in \mathbb{R}^{d+1} \times \mathbb{R}$ the estimate*

$$\left| \partial_\tau^{\alpha_\tau} \partial_\xi^{\alpha_\xi} \frac{1}{\mathcal{L}(i\tau, i\xi, v)} \right| \lesssim \frac{1}{|\mathcal{L}(i\tau, i\xi, v)|} |\tau|^{-\alpha_\tau} |\xi|^{-|\alpha_\xi|}.$$

Proof. The proof rests on the identity

$$\partial_\xi^{\alpha_\xi} \frac{1}{\mathcal{L}(i\tau, i\xi, v)} = \sum_\beta c_\beta \frac{\xi^\beta |v|^{(m-1)N_\beta}}{\mathcal{L}(i\tau, i\xi, v)^{1+N_\beta}},$$

where c_β are constants, $N_\beta := (|\alpha_\xi| + |\beta|)/2$, and the sum runs over those $\beta \in \mathbb{N}_0^d$ with $|\beta| \leq |\alpha_\xi|$ such that $|\alpha_\xi| + |\beta|$ is even. The identity can be proven easily by induction on the order of α_ξ . From this and $\partial_\tau \mathcal{L}(i\tau, i\xi, v) = i$, it immediately follows

$$\left| \partial_\tau^{\alpha_\tau} \partial_\xi^{\alpha_\xi} \frac{1}{\mathcal{L}(i\tau, i\xi, v)} \right| \lesssim \sum_\beta \left| \frac{\xi^\beta |v|^{(m-1)N_\beta}}{\mathcal{L}(i\tau, i\xi, v)^{1+\alpha_\tau+N_\beta}} \right|,$$

which in view of

$$\frac{|\xi|^{|\beta|} |v|^{(m-1)N_\beta}}{|\mathcal{L}(i\tau, i\xi, v)|^{N_\beta}} \leq \frac{|\xi|^{|\beta|} |v|^{(m-1)N_\beta}}{(|v|^{m-1} |\xi|^2)^{N_\beta}} = |\xi|^{-(2N_\beta-|\beta|)} = |\xi|^{-|\alpha_\xi|}$$

and

$$\frac{1}{|\mathcal{L}(i\tau, i\xi, v)|^{\alpha_\tau}} \leq |\tau|^{-\alpha_\tau}$$

yields the assertion. □

Acknowledgments

Gess acknowledges financial support by the Max Planck Society through the Max Planck Research Group “Stochastic partial differential equations” and by the DFG through the CRC “Taming uncertainty and profiting from randomness and low regularity in analysis, stochastics and their applications”. Research of Tadmor was supported in part by NSF grants DMS16-13911, RNMS11-07444 (KI-Net) and ONR grant N00014-1812465. The hospitality of Laboratoire Jacques-Louis Lions in Sorbonne University and its support through ERC grant 740623 under the EU Horizon 2020 is gratefully acknowledged.

References

- [Aronson and Bénilan 1979] D. G. Aronson and P. Bénilan, “Régularité des solutions de l’équation des milieux poreux dans \mathbb{R}^N ”, *C. R. Acad. Sci. Paris Sér. A-B* **288**:2 (1979), A103–A105. [MR](#) [Zbl](#)
- [Arsénio and Masmoudi 2019] D. Arsenio and N. Masmoudi, “Maximal gain of regularity in velocity averaging lemmas”, *Anal. PDE* **12**:2 (2019), 333–388. [MR](#) [Zbl](#)

- [Bahouri, Chemin, and Danchin 2011] H. Bahouri, J.-Y. Chemin, and R. Danchin, *Fourier analysis and nonlinear partial differential equations*, Grundlehren der Math. Wissenschaften **343**, Springer, 2011. [MR](#) [Zbl](#)
- [Bénilan and Crandall 1981] P. Bénilan and M. G. Crandall, “Regularizing effects of homogeneous evolution equations”, pp. 23–39 in *Contributions to analysis and geometry* (Baltimore, MD, 1980), edited by D. N. Clark et al., Johns Hopkins Univ. Press, Baltimore, MD, 1981. [MR](#) [Zbl](#)
- [Bergh and Löfström 1976] J. Bergh and J. Löfström, *Interpolation spaces: an introduction*, Grundlehren der Math. Wissenschaften **223**, Springer, 1976. [MR](#) [Zbl](#)
- [Bögelein, Duzaar, Korte, and Scheven 2019] V. Bögelein, F. Duzaar, R. Korte, and C. Scheven, “The higher integrability of weak solutions of porous medium systems”, *Adv. Nonlinear Anal.* **8**:1 (2019), 1004–1034. [MR](#) [Zbl](#)
- [Chen and Perthame 2003] G.-Q. Chen and B. Perthame, “Well-posedness for non-isotropic degenerate parabolic-hyperbolic equations”, *Ann. Inst. H. Poincaré Anal. Non Linéaire* **20**:4 (2003), 645–668. [MR](#) [Zbl](#)
- [Crandall and Pierre 1982a] M. Crandall and M. Pierre, “Regularizing effects for $u_t + A\varphi(u) = 0$ in L^1 ”, *J. Funct. Anal.* **45**:2 (1982), 194–212. [MR](#) [Zbl](#)
- [Crandall and Pierre 1982b] M. G. Crandall and M. Pierre, “Regularizing effects for $u_t = \Delta\varphi(u)$ ”, *Trans. Amer. Math. Soc.* **274**:1 (1982), 159–168. [MR](#) [Zbl](#)
- [Crandall, Pazy, and Tartar 1979] M. G. Crandall, A. Pazy, and L. Tartar, “Remarks on generators of analytic semigroups”, *Israel J. Math.* **32**:4 (1979), 363–374. [MR](#) [Zbl](#)
- [DeVore and Petrova 2001] R. DeVore and G. Petrova, “The averaging lemma”, *J. Amer. Math. Soc.* **14**:2 (2001), 279–296. [MR](#) [Zbl](#)
- [DiPerna, Lions, and Meyer 1991] R. J. DiPerna, P.-L. Lions, and Y. Meyer, “ L^p regularity of velocity averages”, *Ann. Inst. H. Poincaré Anal. Non Linéaire* **8**:3-4 (1991), 271–287. [MR](#) [Zbl](#)
- [Ebmeyer 2005] C. Ebmeyer, “Regularity in Sobolev spaces for the fast diffusion and the porous medium equation”, *J. Math. Anal. Appl.* **307**:1 (2005), 134–152. [MR](#) [Zbl](#)
- [Gess 2020] B. Gess, “Optimal regularity for the porous medium equation”, *J. Eur. Math. Soc.* (online publication October 2020).
- [Gianazza and Schwarzacher 2019] U. Gianazza and S. Schwarzacher, “Self-improving property of degenerate parabolic equations of porous medium-type”, *Amer. J. Math.* **141**:2 (2019), 399–446. [MR](#) [Zbl](#)
- [Golse and Perthame 2013] F. Golse and B. Perthame, “Optimal regularizing effect for scalar conservation laws”, *Rev. Mat. Iberoam.* **29**:4 (2013), 1477–1504. [MR](#) [Zbl](#)
- [Golse and Saint-Raymond 2004] F. Golse and L. Saint-Raymond, “The Navier–Stokes limit of the Boltzmann equation for bounded collision kernels”, *Invent. Math.* **155**:1 (2004), 81–161. [MR](#) [Zbl](#)
- [Golse, Lions, Perthame, and Sentis 1988] F. Golse, P.-L. Lions, B. Perthame, and R. Sentis, “Regularity of the moments of the solution of a transport equation”, *J. Funct. Anal.* **76**:1 (1988), 110–125. [MR](#) [Zbl](#)
- [Jabin 2009] P.-E. Jabin, “Averaging lemmas and dispersion estimates for kinetic equations”, *Riv. Mat. Univ. Parma* (8) **1** (2009), 71–138. [MR](#) [Zbl](#)
- [Jabin and Vega 2004] P.-E. Jabin and L. Vega, “A real space method for averaging lemmas”, *J. Math. Pures Appl.* (9) **83**:11 (2004), 1309–1351. [MR](#) [Zbl](#)
- [Lions, Perthame, and Souganidis 1996] P.-L. Lions, B. Perthame, and P. E. Souganidis, “Existence and stability of entropy solutions for the hyperbolic systems of isentropic gas dynamics in Eulerian and Lagrangian coordinates”, *Comm. Pure Appl. Math.* **49**:6 (1996), 599–638. [MR](#) [Zbl](#)
- [Lions, Perthame, and Tadmor 1994a] P.-L. Lions, B. Perthame, and E. Tadmor, “A kinetic formulation of multidimensional scalar conservation laws and related equations”, *J. Amer. Math. Soc.* **7**:1 (1994), 169–191. [MR](#) [Zbl](#)
- [Lions, Perthame, and Tadmor 1994b] P.-L. Lions, B. Perthame, and E. Tadmor, “Kinetic formulation of the isentropic gas dynamics and p -systems”, *Comm. Math. Phys.* **163**:2 (1994), 415–431. [MR](#) [Zbl](#)
- [Nikolsky 1962] S. M. Nikolsky, “Boundary properties of differentiable functions of several variables”, *Dokl. Akad. Nauk SSSR* **146** (1962), 542–545. In Russian; translated in *Sov. Math. Dokl.* **3** (1962), 1357–1360. [MR](#) [Zbl](#)

- [Nikolsky 1963a] S. M. Nikolsky, “Functions with dominant mixed derivative, satisfying a multiple Hölder condition”, *Sibirsk. Mat. Ž.* **4**:6 (1963), 1342–1364. In Russian; translated in *Amer. Math. Soc. Transl. (2)* **102** (1973), 27–51. [MR](#) [Zbl](#)
- [Nikolsky 1963b] S. M. Nikolsky, “Stable boundary-value problems of a differentiable function of several variables”, *Mat. Sb. (N.S.)* **61**:2 (1963), 224–252. In Russian. [MR](#)
- [Perthame 2002] B. Perthame, *Kinetic formulation of conservation laws*, Oxford Lect. Series Math. Appl. **21**, Oxford Univ. Press, 2002. [MR](#) [Zbl](#)
- [Schmeisser and Triebel 1987] H.-J. Schmeisser and H. Triebel, *Topics in Fourier analysis and function spaces*, Wiley, Chichester, UK, 1987. [MR](#) [Zbl](#)
- [Tadmor and Tao 2007] E. Tadmor and T. Tao, “Velocity averaging, kinetic formulations, and regularizing effects in quasi-linear PDEs”, *Comm. Pure Appl. Math.* **60**:10 (2007), 1488–1521. [MR](#) [Zbl](#)
- [Triebel 1977a] H. Triebel, “General function spaces, III: Spaces $B_{p,q}^{g(x)}$ and $F_{p,q}^{g(x)}$, $1 < p < \infty$: basic properties”, *Anal. Math.* **3**:3 (1977), 221–249. [MR](#) [Zbl](#)
- [Triebel 1977b] H. Triebel, “General function spaces, IV: Spaces $B_{p,q}^{g(x)}$ and $F_{p,q}^{g(x)}$, $1 < p < \infty$: special properties”, *Anal. Math.* **3**:4 (1977), 299–315. [MR](#) [Zbl](#)
- [Vázquez 2007] J. L. Vázquez, *The porous medium equation: mathematical theory*, Oxford Univ. Press, 2007. [MR](#) [Zbl](#)

Received 19 Apr 2019. Revised 30 Jul 2019. Accepted 26 Sep 2019.

BENJAMIN GESS: bgess@mis.mpg.de

Max Planck Institute for Mathematics in the Sciences, Leipzig, Germany

and

Fakultät für Mathematik, Universität Bielefeld, Bielefeld, Germany

JONAS SAUER: jonas.sauer@mis.mpg.de

Max Planck Institute for Mathematics in the Sciences, Leipzig, Germany

EITAN TADMOR: tadmor@umd.edu

Department of Mathematics, Institute for Physical Science and Technology, University of Maryland, College Park, MD, United States

Analysis & PDE

msp.org/apde

EDITORS

EDITOR-IN-CHIEF

Patrick Gérard

patrick.gerard@math.u-psud.fr

Université Paris Sud XI

Orsay, France

BOARD OF EDITORS

Massimiliano Berti	Scuola Intern. Sup. di Studi Avanzati, Italy berti@sissa.it	Werner Müller	Universität Bonn, Germany mueller@math.uni-bonn.de
Michael Christ	University of California, Berkeley, USA mchrist@math.berkeley.edu	Gilles Pisier	Texas A&M University, and Paris 6 pisier@math.tamu.edu
Charles Fefferman	Princeton University, USA cf@math.princeton.edu	Igor Rodnianski	Princeton University, USA irod@math.princeton.edu
Ursula Hamenstaedt	Universität Bonn, Germany ursula@math.uni-bonn.de	Yum-Tong Siu	Harvard University, USA siu@math.harvard.edu
Vadim Kaloshin	University of Maryland, USA vadim.kaloshin@gmail.com	Terence Tao	University of California, Los Angeles, USA tao@math.ucla.edu
Herbert Koch	Universität Bonn, Germany koch@math.uni-bonn.de	Michael E. Taylor	Univ. of North Carolina, Chapel Hill, USA met@math.unc.edu
Izabella Laba	University of British Columbia, Canada ilaba@math.ubc.ca	Gunther Uhlmann	University of Washington, USA gunther@math.washington.edu
Richard B. Melrose	Massachusetts Inst. of Tech., USA rhm@math.mit.edu	András Vasy	Stanford University, USA andras@math.stanford.edu
Frank Merle	Université de Cergy-Pontoise, France Frank.Merle@u-cergy.fr	Dan Virgil Voiculescu	University of California, Berkeley, USA dvv@math.berkeley.edu
William Minicozzi II	Johns Hopkins University, USA minicozz@math.jhu.edu	Steven Zelditch	Northwestern University, USA zelditch@math.northwestern.edu
Clément Mouhot	Cambridge University, UK c.mouhot@dpmms.cam.ac.uk	Maciej Zworski	University of California, Berkeley, USA zworski@math.berkeley.edu

PRODUCTION

production@msp.org

Silvio Levy, Scientific Editor

See inside back cover or msp.org/apde for submission instructions.

The subscription price for 2020 is US \$340/year for the electronic version, and \$550/year (+\$60, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues from the last three years and changes of subscriber address should be sent to MSP.

Analysis & PDE (ISSN 1948-206X electronic, 2157-5045 printed) at Mathematical Sciences Publishers, 798 Evans Hall #3840, c/o University of California, Berkeley, CA 94720-3840, is published continuously online. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices.

APDE peer review and production are managed by EditFlow® from MSP.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2020 Mathematical Sciences Publishers

ANALYSIS & PDE

Volume 13 No. 8 2020

Propagation properties of reaction-diffusion equations in periodic domains	2259
ROMAIN DUCASSE	
An elementary approach to free entropy theory for convex potentials	2289
DAVID JEKEL	
Parametrix for a semiclassical subelliptic operator	2375
HART F. SMITH	
On the propagation of regularity for solutions of the dispersion generalized Benjamin–Ono equation	2399
ARGENIS J. MENDEZ	
Optimal regularity in time and space for the porous medium equation	2441
BENJAMIN GESS, JONAS SAUER and EITAN TADMOR	