

ANALYSIS & PDE

Volume 16

No. 1

2023

MATTHIEU ASTORG, LUKA BOC THALER AND HAN PETERS

**WANDERING DOMAINS ARISING FROM LAVAURS MAPS WITH
SIEGEL DISKS**

WANDERING DOMAINS ARISING FROM LAVAURS MAPS WITH SIEGEL DISKS

MATTHIEU ASTORG, LUCA BOC THALER AND HAN PETERS

The first example of polynomial maps with wandering domains was constructed in 2016 by the first and last authors, together with Buff, Dujardin and Raissy. In this paper, we construct a second example with different dynamics, using a Lavaurs map with a Siegel disk instead of an attracting fixed point. We prove a general necessary and sufficient condition for the existence of a trapping domain for nonautonomous compositions of maps converging parabolically towards a Siegel-type limit map. Constructing a skew-product satisfying this condition requires precise estimates on the convergence to the Lavaurs map, which we obtain by a new approach. We also give a self-contained construction of parabolic curves, which are integral to this new method.

1. Introduction

Rational functions do not have wandering domains, a classical result due to [Sullivan 1985]. Recently in [Astorg et al. 2016] it was shown that there do exist polynomial maps in two complex variables with wandering Fatou components. The maps constructed in [Astorg et al. 2016] are polynomial skew products of the form

$$(z, w) \mapsto (f_w(z), g(w)),$$

where $g(w)$ and $f_w(z) = f(z, w)$ are polynomials in respectively one and two variables. While the construction holds for families of maps with arbitrarily many parameters, the constructed examples are essentially unique: they all arise from similar behavior and cannot easily be distinguished in terms of the geometry of the components or qualitative behavior of the orbits in the components. The goal in this paper is to modify the construction in [Astorg et al. 2016] to obtain quite different examples of wandering Fatou components. Our construction requires much more precise convergence estimates, forcing us to revisit and clarify the original proof, obtaining a better understanding of the methodology.

The maps considered in [Astorg et al. 2016] are of the specific form

$$P : (z, w) \mapsto \left(f(z) + \frac{\pi^2}{4}w, g(w) \right), \quad (1)$$

where $f(z) = z + z^2 + O(z^3)$ and $g(w) = w - w^2 + O(w^3)$. Recall that the constant $\frac{\pi^2}{4}$ is essential to guarantee the following key result in [Astorg et al. 2016]:

This research was partially supported by the ANR grant Fatou ANR-17-CE40-0002-01. Boc Thaler was supported by the SIR grant “NEWHOLITE - New methods in holomorphic iteration” no. RBS114CFME and by the research program P1-0291 from ARRS, Republic of Slovenia.

MSC2010: 32H50, 37F10, 37G15.

Keywords: Fatou set, holomorphic dynamics, parabolic implosion, polynomial mappings, skew-products, wandering Fatou components, parabolic curves, nonautonomous dynamics.

Proposition A. *As $n \rightarrow +\infty$, the sequence of maps*

$$(z, w) \mapsto P^{\circ 2n+1}(z, g^{\circ n^2}(w))$$

converges locally uniformly in $\mathcal{B}_f \times \mathcal{B}_g$ to the map

$$(z, w) \mapsto (\mathcal{L}_f(z), 0).$$

Throughout this paper \mathcal{B}_f and \mathcal{B}_g refer to the *parabolic basins* of, respectively, f and g , and \mathcal{L}_f refers to the *Lavaurs map* of f with phase 0; see for example [Lavaurs 1989; Shishikura 2000]. By carefully choosing the higher-order terms of f , one can select Lavaurs maps with desired dynamical behavior.

In Proposition B of [Astorg et al. 2016] it was shown that \mathcal{L}_f can have an attracting fixed point. The fact that P has a wandering Fatou component is then a quick corollary of Proposition A. It seems very likely that one can similarly construct wandering domains when \mathcal{L}_f has a parabolic fixed point, using the refinement of Proposition A presented here.

We will construct wandering domains arising when \mathcal{L}_f has a Siegel fixed point: an irrationally indifferent fixed point with Diophantine rotation number. Compositions of small perturbations of \mathcal{L}_f behave so subtly that it is far from clear that Lavaurs maps with Siegel disks can produce wandering domains.

In order to control the behavior of successive perturbations, we prove a refinement of Proposition A with precise convergence estimates, showing that the convergence towards the Lavaurs map is “parabolic”. Moreover, we study the behavior of nonautonomous systems given by maps converging parabolically to a limit map with a Siegel fixed point. We introduce an easily computable index characterizing the behavior of the nonautonomous systems.

In the next section we give more precise statements of our results, and prove how the combination of these results provides a new construction of wandering domains.

2. Background and overview of results

2A. Polynomial skew products and Fatou components. There is more than one possible interpretation of Fatou and Julia sets for polynomial skew products; see for example [Jonsson 1999] for a thorough discussion. When we discuss Fatou components of skew products here, we consider open connected sets in \mathbb{C}^2 whose orbits are uniformly bounded, which of course implies equicontinuity. Since the degrees of f and g in (1) are at least 2, the complement of a sufficiently large bidisk is contained in the escape locus, which is connected; all other Fatou components are therefore bounded and have bounded orbits.

Given a Fatou component U of P , normality implies that its projection onto the second coordinate $\pi_w(U)$ is contained in a Fatou component of g , which must therefore be periodic or preperiodic. Without loss of generality we may assume that this component of g is invariant, and thus either an attracting basin, a parabolic basin or a Siegel disk.

The behavior of P inside a Siegel disk of g may be very complicated and has received little attention in the literature, but see [Peters and Raissy 2019] for the treatment of a special case.

There have been a number of results proving the nonexistence of wandering domains inside attracting basins of g . The nonexistence of wandering domains in the superattracting case was proved in [Lilov

2004], but it was shown in [Peters and Vivas 2016] that the arguments from Lilov cannot hold in the geometrically attracting case. The nonexistence of wandering domains under progressively weaker conditions was proved in [Peters and Smit 2018; Ji 2020].

Here, as in [Astorg et al. 2016], we will consider components U for which $\pi_w(U)$ is contained in a parabolic basin of g . We assume that the fixed point of g lies at the origin, and that g is of the form $g(w) = w - w^2 + \text{h.o.t.}$, so that orbits approach 0 tangent to the positive real axis. We will in fact make the stronger assumption $g(w) = w - w^2 + w^3 + \text{h.o.t.}$

2B. Fatou coordinates and Lavaurs' theorem. Consider a polynomial $f(z) = z - z^2 + az^3 + \text{h.o.t.}$ For $r > 0$ small enough we define incoming and outgoing petals

$$P_f^l = \{|z + r| < r\} \quad \text{and} \quad P_f^o = \{|z - r| < r\}.$$

The incoming petal P_f^l is forward invariant, and all orbits in P_f^l converge to 0. Moreover, any orbit which converges to 0 but never lands at 0 must eventually be contained in P_f^l . Therefore we can define the parabolic basin as

$$\mathcal{B}_f = \bigcup f^{-n} P_f^l.$$

The outgoing petal P_f^o is backwards invariant, with backwards orbits converging to 0.

On P_f^l and P_f^o one can define incoming and outgoing Fatou coordinates $\phi_f^l : P_f^l \rightarrow \mathbb{C}$ and $\phi_f^o : P_f^o \rightarrow \mathbb{C}$ solving the functional equations

$$\phi_f^l \circ f(z) = \phi_f^l(z) + 1 \quad \text{and} \quad \phi_f^o \circ f(z) = \phi_f^o(z) + 1,$$

where $\phi_f^l(P_f^l)$ contains a right half-plane and $\phi_f^o(P_f^o)$ contains a left half-plane. By the first functional equation the incoming Fatou coordinates can be uniquely extended to the attracting basin \mathcal{B}_f . On the other hand, the inverse of ϕ_f^o , denoted by ψ_f^o , can be extended to the entire complex plane, still satisfying the functional equation

$$f \circ \psi_f^o(Z) = \psi_f^o(Z + 1).$$

The fact that the exceptional set of f is empty implies that $\psi_f^o : \mathbb{C} \rightarrow \mathbb{C}$ is surjective. We note that both incoming and outgoing Fatou coordinates are (on the corresponding petals) of the form $Z = -1/z + b \log(z) + o(1)$, where the coefficient b vanishes when $a = 1$. This is one reason for working with maps f of the form $f(z) = z + z^2 + z^3 + \text{h.o.t.}$

Let us now consider small perturbations of the map f . For $\epsilon \in \mathbb{C}$ we write $f_\epsilon(z) = f(z) + \epsilon^2$, and consider the behavior as $\epsilon \rightarrow 0$. The most interesting behavior occurs when ϵ approaches 0 tangent to the positive real axis.

Lavaurs' theorem [1989]. *Let $\epsilon_j \rightarrow 0$, $n_j \in \mathbb{N}$ and $\alpha \in \mathbb{C}$ satisfy*

$$n_j - \frac{\pi}{\epsilon_j} \rightarrow \alpha \quad \text{as } j \rightarrow \infty.$$

Then

$$f_{\epsilon_j}^{n_j} \rightarrow \mathcal{L}_f(\alpha) = \psi_f^o \circ \tau_\alpha \circ \phi_f^l,$$

where $\tau_\alpha(Z) = Z + \alpha$.

The map $\mathcal{L}_f(\alpha)$ is called the *Lavaurs map*, and α is called the *phase*. In this paper we will only consider phase $\alpha = 0$, and write \mathcal{L}_f instead of $\mathcal{L}_f(0)$.

2C. Propositions A and B. The construction of wandering domains in [Astorg et al. 2016] follows quickly from two key propositions, the aforementioned Propositions A and B. In this paper we will prove a variation to Proposition B, and a refinement to Proposition A, which we will both state here.

Our main technical result is the following refinement of Proposition A. As before we write $P(z, w) = (f(z) + \frac{\pi^2}{4}w, g(w))$, with $f(z) = z + z^2 + z^3 + bz^4 + \text{h.o.t.}$, and $g(w) = w - w^2 + w^3 + \text{h.o.t.}$

Proposition A'. *There exists a holomorphic function $h : \mathcal{B}_f \times \mathcal{B}_g \rightarrow \mathbb{C}$ such that*

$$P^{2n+1}(z, g^{n^2}(w)) = (\mathcal{L}_f(z), 0) + \left(\frac{h(z, w)}{n}, 0 \right) + O\left(\frac{\log n}{n^2} \right),$$

uniformly on compact subsets of $\mathcal{B}_f \times \mathcal{B}_g$. The function $h(z, w)$ is given by

$$h(z, w) = \frac{\mathcal{L}'_f(z)}{(\phi'_f)^t(z)} \cdot (C + \phi'_f(z) - \phi'_g(w)),$$

where the constant $C \in \mathbb{C}$ depends on b .

Proposition A' will be proved in Section 5; see Theorem 5.33.

Proposition B in [Astorg et al. 2016] states that the Lavaurs map \mathcal{L}_f of a polynomial $f(z) = z + z^2 + az^3 + O(z^4)$ has an attracting fixed point for suitable choices of the constant $a \in \mathbb{C}$. We recall very briefly the main idea in the proof of Proposition B: For $a = 1$ the ‘‘horn map’’ has a parabolic fixed point at infinity. By perturbing $a \simeq 1$, the parabolic fixed point bifurcates, and for appropriate perturbations this guarantees the existence of an attracting fixed point for the horn map, and thus also for the Lavaurs map.

In this paper we will consider a more restrictive family of polynomials of the form $f(z) = z + z^2 + z^3 + O(z^4)$, which means that we cannot use the above bifurcation argument. Using a different line of reasoning, using small perturbations of a suitably chosen degree-7 real polynomial, we will prove a variation to Proposition B, namely Proposition B' below. The proof of Proposition B' will be given in Section 6.

Before stating the proposition we recall that a fixed point $z_0 = \mathcal{L}_f(z_0)$ is said to be of Siegel type if $\lambda = \mathcal{L}'_f(z_0) = e^{2\pi i \zeta}$, where $\zeta \in \mathbb{R} \setminus \mathbb{Q}$ is Diophantine, i.e., if there exist $c, r > 0$ such that $|\lambda^n - 1| \geq cn^{-r}$ for all integers $n > 0$. Recall that neutral fixed points with Diophantine rotation numbers are always locally linearizable:

Theorem 2.1 [Siegel 1942]. *Let $p(z) = e^{2\pi i \zeta} z + O(z^2)$ be a holomorphic germ. If ζ is Diophantine then there exist a neighborhood of the origin Ω_p and a biholomorphic map $\varphi : \Omega_p \rightarrow D_r(0)$ of the form $\varphi(z) = z + a_2 z^2 + O(z^3)$ satisfying*

$$\varphi(p(z)) = e^{2\pi i \zeta} \varphi(z).$$

Proposition B'. *There exist polynomials of the form $f(z) = z + z^2 + z^3 + O(z^4)$ for which the Lavaurs map \mathcal{L}_f has a **Siegel fixed point** z_0 , with $\lambda = \mathcal{L}'_f(z_0)$. Moreover we can guarantee that*

$$\frac{\mathcal{L}''_f(z_0)(\phi_f^l)'(z_0)}{\lambda(1-\lambda)} - (\phi_f^l)''(z_0) \neq 0. \quad (2)$$

Condition (2) is necessary to guarantee the existence of wandering domains; see the discussion of the index κ later in this section, and the discussion in Section 5C.

A more precise description of the derivatives λ for which p is locally linearizable was given in [Bruno 1971; 1972; Yoccoz 1995]. As we are only concerned with constructing examples of maps with wandering Fatou components, we find it convenient to work with the stronger Diophantine condition. Proposition B' will be proved in Section 6.

2D. Perturbations of Siegel disks. A key element in our study is the following question:

Let f_1, f_2, \dots be a sequence of holomorphic germs, converging locally uniformly to a holomorphic function f having a Siegel fixed point at 0. Under which conditions does there exist a trapping region?

By a trapping region we mean the existence of arbitrarily small neighborhoods U, V of 0 and $n_0 \in \mathbb{N}$ such that

$$f_m \circ \dots \circ f_n(z) \in V$$

for all $z \in U$ and $m \geq n \geq n_0$. In other words, any orbit $(z_n)_{n \geq 0}$ that intersects U for sufficiently large n will afterwards be contained in a small neighborhood of the origin. Note that this in particular guarantees normality of the sequence of compositions $f_m \circ \dots \circ f_0$ in a neighborhood of z_0 , which is the reason for our interest in trapping regions.

We are particularly interested in the case where the differences $f_n - f$ are not absolutely summable, i.e., when

$$\sum_{n \geq n_0} \|f_n - f\|_U = \infty$$

for any n_0 and U . In this situation one generally does not expect a trapping region. However, motivated by Proposition A', we will assume that $f_n - f$ is roughly of size $1/n$, and converges to zero along some real direction. More precisely, we assume that

$$f_n(z) - f(z) = \frac{h(z)}{n} + O\left(\frac{1}{z^{1+\epsilon}}\right), \quad (3)$$

where h is a holomorphic germ, defined in a neighborhood of the origin.

Theorem 2.2. *There exists an **index** κ , a rational expression in the coefficients of f and h , such that the following hold:*

- (1) *If $\operatorname{Re}(\kappa) = 0$, then there is a trapping region, and all limit maps have rank 1.*
- (2) *If $\operatorname{Re}(\kappa) < 0$, then there is a trapping region, and all orbits converge uniformly to the origin.*
- (3) *If $\operatorname{Re}(\kappa) > 0$, then there is no trapping region. In fact, there can be at most one orbit that remains in a sufficiently small neighborhood of the origin.*

Theorem 2.2 holds under more general assumptions regarding the convergence towards the limit map, but the above statement is sufficient for our purposes. An example of a more general statement is given in Remark 3.15. An explicit formula for the index κ is given in Section 3, which contains the proof of Theorem 2.2.

Remark 2.3. The nonautonomous dynamics of the functions f_n satisfying (3) is closely related to the autonomous dynamics of the quasiparabolic map

$$F(z, w) = (f(z) + wh(z) + O(w^2), w - w^2 + O(w^2)).$$

The case $\operatorname{Re}(\kappa) < 0$ in Theorem 2.2 corresponds to F being dynamically separating and parabolically attracting, using the terminology of [Bracci and Zaitsev 2013]; by Corollary 6.3 of that work the map F has a connected basin of attraction at the origin. In particular this implies the existence of a trapping region for the sequence (f_n) .

2E. Parabolic curves. An important idea in the proof of Lavaurs’ theorem is that in a sufficiently small neighborhood of the origin, the function $f_\epsilon = f + \epsilon^2$ can be interpreted as a near-translation in the “almost Fatou coordinates”: functions that converge to the ingoing and outgoing Fatou coordinates as $\epsilon \rightarrow \infty$. This idea is especially apparent in the treatment given in [Bedford et al. 2017]. The almost Fatou coordinates are defined using the pair of fixed points $\zeta_\pm(\epsilon)$ “splitting” from the parabolic fixed point.

When iterating two-dimensional skew products $P(z, w) = (f_w(z), g(w))$ it does not make sense to base the almost Fatou coordinates on the pair of fixed points of the maps $f_w(z) = f(z) + \frac{\pi^2}{4}w$, as the parameter w changes after every iteration of P . Instead, the natural idea would be to base these coordinates on a pair of invariant curves $\{z = \zeta_\pm(w)\}$, so-called *parabolic curves*, defined over a forward-invariant parabolic petal in the w -plane. The invariance of these parabolic curves is equivalent to the functional equations

$$\zeta_\pm(g(w)) = f_w(\zeta_\pm(w)).$$

In [Astorg et al. 2016], it is asked whether such parabolic curves exist. Instead, in that work it was shown that there exist *almost parabolic curves*, approximate solutions to the above functional equation with explicit error estimates. The proof of Proposition A relies to a great extent on these almost parabolic curves, and the fact that these are not exact solutions causes significant extra work.

In [López-Hernanz and Rosas 2020] it is shown that the parabolic curves indeed exist, in fact, the authors prove the existence of parabolic curves for any characteristic direction for diffeomorphisms in two complex dimensions. However, to be used in the proof of Proposition A, it is necessary to also obtain control over the domain of definition of the two parabolic curves. The result from [López-Hernanz and Rosas 2020] does not give the needed control.

In Section 4, Proposition 4.1, we give an alternative proof of the existence of parabolic curves, with control over the domains of definition. The availability of these parabolic curves forms an important ingredient in the proof of Proposition A’. The method of proof is a variation to the well-known graph transform method, and can likely be used to prove the existence of parabolic curves in greater generality.

2F. Wandering domains. Let us conclude this section by proving how Propositions A' and B' together imply the existence of wandering Fatou components. As before we let

$$P(z, w) = \left(f(z) + \frac{\pi^2}{4} w, g(w) \right),$$

where $g(w) = w - w^2 + w^3 + \text{h.o.t.}$ and the function $f(z) = z + z^2 + z^3 + \text{h.o.t.}$ is chosen such that \mathcal{L}_f has a neutral fixed point z_0 with Diophantine rotation number. The existence of such f is given by Proposition B'.

Proposition A' states that

$$P^{2n+1}(z, g^{n^2}(w)) = (\mathcal{L}_f(z), 0) + \left(\frac{h(z, w)}{n}, 0 \right) + O\left(\frac{\log n}{n^2} \right),$$

uniformly on compact subsets of $\mathcal{B}_f \times \mathcal{B}_g$.

Recall from Proposition A' that the function $h(z, w)$ is given by

$$h(z, w) = \frac{\mathcal{L}'_f(z)}{(\phi'_f)^l(z)} \cdot (C + \phi'_f(z) - \phi'_g(w)),$$

from which it follows directly that the index κ depends affinely on $\phi'_g(w)$, although it is conceivable that the multiplicative constant in this dependence vanishes.

As will be explained in detail in Section 5C, the index κ is independent from w if and only if, denoting the fixed point of \mathcal{L}_f again by z_0 , we have

$$\frac{\mathcal{L}''_f(z_0)(\phi'_f)^l(z_0)}{\lambda(1-\lambda)} - (\phi'_f)^l(z_0) = 0, \quad (4)$$

in which case κ is constantly equal to $+1$. The second statement in Proposition B' therefore implies that f can be chosen in order to obtain an inequality in (4), which implies that the affine dependence of κ on $\phi'_g(w)$ is nonconstant.

It follows that there exists an open subset of \mathcal{B}_g where the w -values are such that $\text{Re}(\kappa)$ is strictly negative. Let $D_2 \subset \mathcal{B}_g$ be a small disk contained in this open subset, so that $\text{Re}(\kappa)$ is negative for all $w \in D_2$.

Let D_1 be a small disk centered at z_0 , the Siegel-type fixed point of \mathcal{L}_f . We claim that, for $n \in \mathbb{N}$ large enough, the open set $D_1 \times g^{n^2}(D_2)$ is contained in a wandering Fatou component.

Indeed, it follows from Proposition A' that the nonautonomous one-dimensional system given by compositions of the maps $z \mapsto \pi_z \circ P^{2n+1}(z, g^{n^2}(w))$ satisfies case (2) of Theorem 2.2, where π_z is the projection onto the z -coordinate. Thus Theorem 2.2 implies that

$$P^{m^2-n^2}(z, w) \rightarrow (z_0, 0)$$

uniformly for all $(z, w) \in D_1 \times g^{n^2}(D_2)$. The remainder of the proof follows the argument from [Astorg et al. 2016]. Since the complement of the escape locus of P is bounded, it follows that the entire orbits $P^m(z, w)$ must remain uniformly bounded, which implies normality of (P^m) on $D_1 \times g^{n^2}(D_2)$, which is therefore contained in a Fatou component, say U . The fact that on an open subset of U the

subsequence $P^{m^2-n^2}$ converges to the constant $(z_0, 0)$ implies convergence of this subsequence to $(z_0, 0)$ on all of U , since limit maps of convergent subsequences are holomorphic. But if U was periodic or preperiodic, the limit set would have been periodic. The point $(z_0, 0)$ is however not periodic: its orbit converges to $(0, 0)$. Thus U is wandering, which completes the proof.

Remark 2.4. From the above discussion we can conclude that all possible limit maps of the convergent subsequence $P^{n_j}|_U$ are points. In fact these points form (the closure of) a bi-infinite orbit of $(z_0, 0)$, converging to $(0, 0)$ both under backward and forward iteration.

We note however that there are fibers $\{w = w_0\}$, with $w_0 \in \mathcal{B}_g$, for which $\text{Re}(\kappa) = 0$. Let D_1 again be a sufficiently small disk centered at z_0 , the Siegel-type fixed point of \mathcal{L}_f . Proposition A' together with case (1) of Theorem 2.2 implies that for sufficiently large n the disk $D_1 \times \{g^{n^2}(w_0)\}$ is a Fatou disk for P , i.e., the restriction of the iterates P^n to the disk form a normal family. For this Fatou disk the sequence of iterates $P^{m^2-n^2}$ converges to a rank-1 limit map, whose image is a holomorphic disk containing $(z_0, 0)$. All the limit sets together form (the closure of) a bi-infinite sequence of disks, converging under backward and forward iteration to the point $(0, 0)$.

3. Perturbations of Siegel disks

3A. Notation. The following conventions will be used throughout this section:

- (i) Given a holomorphic function f , we will write \hat{f} for the nonlinear part of f .
- (ii) For a sequence of constants $\lambda_n \in \mathbb{C}$ we will write

$$\lambda_{n,m} = \prod_{j=m+1}^n \lambda_j \quad \text{and} \quad \lambda(n) = \lambda_{n,0} = \prod_{j=1}^n \lambda_j,$$

and similarly for a sequence of functions (f_n)

$$f_{n,m} = f_n \circ \cdots \circ f_{m+1}.$$

- (iii) Given two sequences of holomorphic functions (f_n) and (g_n) defined on some uniform neighborhood of the origin, we will write $f_n \asymp g_n$ if the norms of the sequence of differences $(f_n - g_n)$ is summable on some uniform neighborhood of the origin.

3B. Preparation. In this section we introduce nonautonomous analogies of attracting, repelling, and locally linearizable indifferent fixed points and make a few initial observations. In the next subsection we introduce the index κ and show that the local behavior of the nonautonomous systems we consider can be deduced from the real part of the index.

Definition 3.1. Two sequences of functions (f_n) and (g_n) are said to be *nonautonomously conjugate* if there exist a uniformly bounded sequence of local coordinate changes $(\psi_n)_{n \geq n_0}$, all defined in a uniform neighborhood of the origin, satisfying

$$f_n \circ \psi_n = \psi_{n+1} \circ g_n$$

for all $n \geq n_0$.

Definition 3.2. A sequence of functions (f_n) is said to be nonautonomously *linearizable* if there exists a sequence $(\lambda_n)_{n \geq n_0}$ in $\mathbb{C} \setminus \{0\}$ and a sequence of coordinate changes $(\psi_n)_{n \geq n_0}$, defined and uniformly bounded in a uniform neighborhood of the origin, and with derivative $\psi'_n(0)$ uniformly bounded away from zero, so that

$$f_n \circ \psi_n(z) = \psi_{n+1}(\lambda_n \cdot z)$$

for all $n \geq n_0$. If the sequence $|\lambda(n)|$ is bounded, both from above and away from 0, then we say that (f_n) is *rotationally linearizable*.

Definition 3.3. A sequence of functions (f_n) is said to be *collapsing* if there is a neighborhood of the origin U and an $n_0 \in \mathbb{N}$ such that $f_{n,m} \rightarrow 0$ on U as $n \rightarrow \infty$ for any $m \geq n_0$.

An example of a collapsing sequence is given by a sequence of functions f_n converging to a function f with an attracting fixed point at the origin.

Definition 3.4. We say that sequence (f_n) is *expulsive* if there exists $r > 0$ such that for every $m \geq 0$ there exists at most one exceptional point \hat{z} such that for every $z \in D_r(0) \setminus \{\hat{z}\}$ there exist $n > m$ for which $f_{n,m}(z) \notin D_r(0)$. Here $D_r(0)$ denotes the disk of radius r centered at the origin.

An example of an expulsive sequence can be obtained by considering a sequence of maps (f_n) converging locally uniformly to a map with a repelling fixed point. Since f maps a small disk around the origin to a strictly larger holomorphic disk, the same holds for sufficiently small perturbations. A nested sequence argument shows that, starting at a sufficiently large time n_0 , there is a unique orbit which remains in the small disk.

Lemma 3.5. *Consider a sequence (f_n) of univalent holomorphic functions, defined in a uniform neighborhood of the origin. Suppose the compositions $f_{n,0}$ are all defined in a possibly smaller neighborhood of the origin, and form a normal family. Then the sequence (f_n) is either rotationally linearizable, or there exist subsequences (n_j) for which $f_{n_j,0}$ converges to a constant.*

Proof. By normality the orbit $f_{n,0}(0)$ stays bounded. By nonautonomously conjugating with a sequence of translations we may therefore assume that $f_n(0) = 0$ for all n . Note that normality is preserved under nonautonomous conjugation by bounded translations.

Write $\lambda_n = f'_n(0)$. Normality implies that $|\lambda(n)|$ is bounded from above. The functions

$$\psi_{n+1}(z) := f_{n,0}(\lambda(n)^{-1} \cdot z)$$

are tangent to the identity, and they satisfy the functional equation

$$f \circ \psi_n(z) = \psi_{n+1}(\lambda \cdot z).$$

If the sequence $|\lambda(n)|$ is bounded away from the origin then the maps ψ_n are uniformly bounded, and the sequence $(f(n))$ is rotationally linearizable. Suppose that the sequence $\lambda(n)$ is not bounded from below, in which case there is a subsequence $\lambda(n_j)$ converging to 0. By the Hurwitz theorem the sequence of maps $f_{n_j,0}$ converges to a constant. \square

Lemma 3.6. *If the sequence (f_n) is rotationally linearizable, and (ζ_n) is a sequence of absolutely summable holomorphic functions, i.e.,*

$$\sum \|\zeta_n\|_{D_r(0)} < \infty$$

for some $r > 0$, then the sequence $(f_n + \zeta_n)$ is also rotationally linearizable.

Proof. Write $g_n = f_n + \zeta_n$. We consider the errors due to the perturbations in linearization coordinates, i.e.,

$$\psi_{n+1}^{-1} \circ g_n \circ \psi_n(z) - \psi_{n+1}^{-1} \circ f_n \circ \psi_n(z) = \psi_{n+1}^{-1} \circ g_n \circ \psi_n(z) - \lambda_n \cdot z.$$

By the definition of the nonautonomous linearization, it follows that after restricting to a smaller neighborhood of the origin the derivatives of the maps ψ_n and their inverses are uniformly bounded. It follows that the above errors are also absolutely summable, which guarantees normality of the sequence $\psi_{n+1}^{-1} \circ g_n$ in a small neighborhood of the origin, and hence normality of the sequence g_n . It follows from Lemma 3.5 that $(f_n + \zeta_n)$ is either rotationally linearizable or has subsequences converging to the origin. It follows from the summability of the errors that the latter is impossible. \square

3C. Introduction of the index. Let $f(z) = \lambda z + b_2 z^2 + O(z^3)$ be a holomorphic function with $\lambda = e^{2\pi i \zeta}$ and $\zeta \in \mathbb{R} \setminus \mathbb{Q}$ Diophantine. Let $h(z) = c_0 + c_1 z + O(z^2)$ be a holomorphic function defined in a neighborhood of the origin. Let $(\zeta_n(z))$ be a sequence of holomorphic functions that is defined and absolutely summable on some uniform neighborhood of the origin. We consider the nonautonomous dynamical system given by compositions of the maps

$$f_n(z) = f(z) + \frac{1}{n} h(z) + \zeta_n(z).$$

We introduce the index κ , depending rationally on the two-jet of f at the origin and the one-jet of h at the origin, by

$$\kappa := \frac{2b_2 c_0}{\lambda(1-\lambda)} + \frac{c_1}{\lambda}. \quad (5)$$

We claim that the index κ is invariant under local autonomous changes of coordinates, i.e., when all the maps f_n are conjugated by a single analytic transformation. One easily observes that the index is invariant under affine changes of coordinates and is unaffected by terms of order 3 and higher. It is therefore sufficient to only consider local changes of the form $z \mapsto z + \alpha z^2$. It is clear that λ and c_0 are unaffected by such a coordinate change, while computation shows that b_2 is replaced by $b_2 + \alpha\lambda - \alpha\lambda^2$ and c_1 is replaced by $c_1 - \alpha\lambda c_0$. Indeed, κ is invariant under these changes.

Since ζ is Diophantine, the function f is linearizable. Let us write $\phi(z) = z + \text{h.o.t.}$ for the linearization map of f , i.e., $f \circ \phi(z) = \phi(\lambda z)$.

We define

$$\theta_n(z) := z + \frac{1}{n} \frac{c_0}{1-\lambda}.$$

Lemma 3.7. *With the above definitions we can write*

$$f_n := \phi^{-1} \circ \theta_{n+1}^{-1} \circ f_n \circ \theta_n \circ \phi = \lambda \cdot e^{\kappa/n} \cdot z + \frac{1}{n} \sum_{k=2}^{\infty} d_k z^k + \xi_n(z), \quad (6)$$

where (ξ_n) is a sequence of holomorphic functions that are defined and whose norms are summable on a uniform neighborhood of the origin.

Proof. First observe that

$$\begin{aligned}\theta_{n+1}^{-1} \circ f_n \circ \theta_n &\asymp f\left(z + \frac{1}{n} \frac{c_0}{1-\lambda}\right) + \frac{1}{n} h\left(z + \frac{1}{n} \frac{c_0}{1-\lambda}\right) - \frac{1}{n+1} \frac{c_0}{1-\lambda} \\ &\asymp f(z) + f'(z) \frac{1}{n} \frac{c_0}{1-\lambda} + \frac{1}{n} h(z) - \frac{1}{n+1} \frac{c_0}{1-\lambda}.\end{aligned}$$

Using the power series expansions of f' and h we can therefore write

$$\begin{aligned}\theta_{n+1}^{-1} \circ f_n \circ \theta_n &\asymp f(z) + \left(\frac{1}{n} \frac{\lambda c_0}{1-\lambda} + \frac{1}{n} c_0 - \frac{1}{n+1} \frac{c_0}{1-\lambda}\right) + \frac{1}{n} \left(\frac{2b_2 c_0}{1-\lambda} + c_1\right) z + \frac{1}{n} \sum_{k=2}^{\infty} \beta_k z^k \\ &\asymp f(z) + \frac{c_0}{1-\lambda} \left(\frac{1}{n} - \frac{1}{n+1}\right) + \frac{1}{n} \lambda \kappa z + \frac{1}{n} \sum_{k=2}^{\infty} \beta_k z^k \\ &\asymp f(z) + \frac{1}{n} \lambda \kappa z + \frac{1}{n} \sum_{k=2}^{\infty} \beta_k z^k.\end{aligned}$$

It follows that

$$\begin{aligned}f_n &\asymp \phi^{-1}(f(\phi(z))) + (\phi^{-1})'(f(\phi(z))) \left(\frac{1}{n} \lambda \kappa \phi(z) + \frac{1}{n} \sum_{k=2}^{\infty} \beta_k \phi(z)^k\right) \\ &\asymp \lambda \left(1 + \frac{\kappa}{n}\right) z + \frac{1}{n} \sum_{k=2}^{\infty} d_k z^k \\ &\asymp \lambda e^{\kappa/n} z + \frac{1}{n} \sum_{k=2}^{\infty} d_k z^k.\end{aligned}$$

For the last equality we used that

$$1 + \frac{\kappa}{n} = e^{\kappa/n} + O\left(\frac{1}{n^2}\right). \quad \square$$

Corollary 3.8. *If $\operatorname{Re}(\kappa) < 0$ the sequence f_n is collapsing.*

Proof. Observe that $f'_n(z) \asymp \lambda e^{\kappa/n} + O(z/n)$ and note that there is a small disk $D_r(0)$ such that for n sufficiently large

$$\|f'_n\|_{D_r(0)} < e^{\operatorname{Re}(\kappa)/(2n)},$$

and thus

$$|f_n(z) - f_n(w)| \leq e^{\operatorname{Re}(\kappa)/(2n)} |z - w|. \quad (7)$$

Since $\operatorname{Re}(\kappa) < 0$ it follows that $\prod_{n \geq 1} e^{\operatorname{Re}(\kappa)/(2n)} = 0$.

Let us write

$$\varphi_n(z) = \lambda \cdot e^{\kappa/n} \cdot z + \hat{f}_n(z),$$

i.e., we drop the term ξ_n from f_n . By decreasing the radius r if necessary we can choose m_0 such that

$$\sum_{j \geq m_0} \|\xi_j\|_{D_r(0)} < \frac{1}{2}r.$$

By increasing m_0 if necessary we can also guarantee that $\varphi_{n,m}(z) \in D_{r/2}(0)$ for all $z \in D_{r/4}(0)$ and $m \geq m_0$. Using (7) it follows by induction on n that whenever $z \in D_{r/4}(0)$ and $m \geq m_0$ we have

$$\|f_{n,m}(z) - \varphi_{n,m}(z)\| \leq \sum_{j=m}^n \left(\prod_{k=j+1}^n e^{\operatorname{Re}(\kappa)/(2k)} \right) \|\xi_j\|_{D_r(0)}.$$

Indeed, the inequality is trivially satisfied for $n = m$, and assuming the inequality holds for some $n \geq m$ implies

$$\begin{aligned} \|f_{n+1,m}(z) - \varphi_{n+1,m}(z)\| &= \|f_{n+1} \circ f_{n,m}(z) - f_{n+1} \circ \varphi_{n,m}(z) + \xi_{n+1}(\varphi_{n,m}(z))\| \\ &\leq \sum_{j=m}^{n+1} \left(\prod_{k=j+1}^{n+1} e^{\operatorname{Re}(\kappa)/(2k)} \right) \|\xi_j\|_{D_r(0)}. \end{aligned}$$

Note that

$$\sum_{j=m}^n \left(\prod_{k=j+1}^n e^{\operatorname{Re}(\kappa)/(2k)} \right) \|\xi_j\|_{D_r(0)} \rightarrow 0$$

as $n \rightarrow \infty$; hence the fact that the sequence (φ_n) collapses implies the sequence (f_n) collapses as well. \square

Since the sequence (f_n) collapses, it follows immediately that the sequence (f_n) collapses as well, concluding the case $\operatorname{Re}(\kappa) < 0$.

Corollary 3.9. *If $\operatorname{Re}(\kappa) > 0$, the sequence f_n is expulsive.*

Proof. Note that there are $r, n_0 > 0$ such that for every $z, w \in D_r(0)$ and every $n > n_0$ we have

$$|f_n(z) - f_n(w)| = |z - w| \cdot \left| e^{\kappa/n} + \frac{1}{n} O(z, w) \right| > e^{\operatorname{Re}(\kappa)/(2n)} |z - w|.$$

Expulsion of all but one orbit follows immediately. \square

Again it follows that (f_n) is expulsive, completing the case $\operatorname{Re}(\kappa) > 0$.

3D. Rotationally linearizable case ($\operatorname{Re}(\kappa) = 0$). Let us define

$$L_n(z) = e^{\kappa \log n} \cdot z.$$

We obtain

$$\begin{aligned} g_n &= L_{n+1}^{-1} \circ f_n \circ L_n \\ &= \lambda z + e^{-\kappa \log(n+1)} \frac{1}{n} \sum_{\ell=2}^{\infty} d_\ell e^{\kappa \ell \log n} z^\ell + L_{n+1}^{-1} \circ \xi_n \circ L_n \\ &\asymp \lambda z + \frac{1}{n} \sum_{\ell=2}^{\infty} d_\ell e^{\kappa(\ell-1) \log n} z^\ell. \end{aligned}$$

Since $\operatorname{Re}(\kappa) = 0$, the maps L_n are rotations; hence it is sufficient to prove that the sequence (g_n) is rotationally linearizable.

By Lemma 3.6 we may ignore the absolutely summable part of g_n ; hence with slight abuse of notation we may assume that

$$g_n = \lambda z + \frac{1}{n} \sum_{\ell=2}^{\infty} d_{\ell} e^{\kappa(\ell-1) \log n} z^{\ell}.$$

Recall that $\lambda = e^{2\pi i \zeta}$, where ζ is Diophantine.

Lemma 3.10. *There exist constants $C, r > 0$ such that for every integer $\ell \geq 1$ and for every $0 < m < N$ we have*

$$\left| \sum_{j=m}^N \lambda^{\ell j} \right| < C \ell^r.$$

Proof. Since ζ is assumed to be Diophantine, there exist $c, r > 0$ such that $|\lambda^n - 1| \geq cn^{-r}$ for all n . This gives the bound

$$\left| \sum_{j=m}^N \lambda^{\ell j} \right| = \left| \sum_{j=m}^N \frac{\lambda^{\ell(j+1)} - \lambda^{\ell j}}{\lambda^{\ell} - 1} \right| = \left| \frac{1}{\lambda^{\ell} - 1} \sum_{j=m}^N (\lambda^{\ell(j+1)} - \lambda^{\ell j}) \right| < \left| \frac{2}{\lambda^{\ell} - 1} \right| < C \ell^r. \quad \square$$

Lemma 3.11. *There exist $\tilde{C}, r > 0$ such that for all integers $n, \ell > 0$ we have*

$$\left| \sum_{k=n}^{\infty} \frac{e^{\kappa \ell \log k}}{k} \lambda^{k \ell} \right| < \frac{\tilde{C} \ell^{r+1}}{n}.$$

Proof. Summation by parts gives

$$\begin{aligned} \sum_{k=n}^N \frac{e^{\kappa \ell \log k}}{k} \lambda^{k \ell} &= \frac{e^{\kappa \ell \log N}}{N} \sum_{k=n}^N \lambda^{k \ell} - \sum_{k=n}^{N-1} \left(\frac{e^{\kappa \ell \log(k+1)}}{k+1} - \frac{e^{\kappa \ell \log k}}{k} \right) \sum_{j=n}^k \lambda^{j \ell} \\ &= \frac{e^{\kappa \ell \log N}}{N} \sum_{k=n}^N \lambda^{k \ell} - \sum_{k=n}^{N-1} e^{\kappa \ell \log k} \left(\frac{1 + \kappa \ell / k + O(1/k^2)}{k+1} - \frac{1}{k} \right) \sum_{j=n}^k \lambda^{j \ell}. \end{aligned}$$

Observe that

$$\frac{1 + \kappa \ell / k + O(1/k^2)}{k+1} - \frac{1}{k} = O\left(\frac{1}{k^2}\right)$$

is absolutely summable; hence using Lemma 3.10 we obtain

$$\begin{aligned} \left| \sum_{k=n}^N \frac{e^{\kappa \ell \log k}}{k} \lambda^{k \ell} \right| &< \frac{1}{N} \left| \sum_{k=n}^N \lambda^{k \ell} \right| + \sum_{k=n}^{N-1} \left| \frac{1 + \kappa \ell / k + O(1/k^2)}{k+1} - \frac{1}{k} \right| \left| \sum_{j=n}^k \lambda^{j \ell} \right| \\ &< \frac{C \ell^r}{N} + C \ell^r \sum_{k=n}^{N-1} \left| \frac{\ell \kappa - 1}{k(k+1)} + O\left(\frac{1}{k^3}\right) \right| \\ &< \frac{\tilde{C} \ell^{r+1}}{n}. \quad \square \end{aligned}$$

Let us introduce one more change of coordinates

$$S_{n+1}(z) = z - \lambda^{-1} \sum_{\ell=2}^{\infty} \lambda^{(n+1)(1-\ell)} d_{\ell} z^{\ell} \sum_{k=n+1}^{\infty} \frac{e^{\kappa(\ell-1) \log k}}{k} \lambda^{k(\ell-1)}.$$

Lemma 3.12. *Writing $S_n(z) = z + \hat{S}_n(z)$ we obtain*

$$\hat{S}_{n+1}(\lambda z) = \lambda \hat{S}_n(z) + \hat{g}_n(z).$$

Proof. Computing $\hat{S}_{n+1}(\lambda z) - \hat{g}_n(z)$ gives

$$\begin{aligned} & -\lambda^{-1} \sum_{\ell=2}^{\infty} \lambda^{(n+1)(1-\ell)} d_{\ell} \lambda^{\ell} z^{\ell} \sum_{k=n+1}^{\infty} \frac{e^{\kappa(\ell-1) \log k}}{k} \lambda^{k(\ell-1)} - \frac{1}{n} \sum_{\ell=2}^{\infty} e^{\kappa(\ell-1) \log n} d_{\ell} z^{\ell} \\ &= - \sum_{\ell=2}^{\infty} \lambda^{n(1-\ell)} d_{\ell} z^{\ell} \sum_{k=n+1}^{\infty} \frac{e^{\kappa(\ell-1) \log k}}{k} \lambda^{k(\ell-1)} - \sum_{\ell=2}^{\infty} \frac{e^{\kappa(\ell-1) \log n}}{n} \lambda^{n(\ell-1)} \lambda^{n(1-\ell)} d_{\ell} z^{\ell} \\ &= - \sum_{\ell=2}^{\infty} \lambda^{n(1-\ell)} d_{\ell} z^{\ell} \sum_{k=n}^{\infty} \frac{e^{\kappa(\ell-1) \log k}}{k} \lambda^{k(\ell-1)} = \lambda \hat{S}_n(z). \quad \square \end{aligned}$$

Lemma 3.13. *The maps S_n satisfy $S_n = z + O(1/n)$, with uniform bounds.*

Proof.

$$|\hat{S}_n(z)| = \left| \lambda^{-1} \sum_{\ell=2}^{\infty} \lambda^{n(1-\ell)} d_{\ell} z^{\ell} \sum_{k=n}^{\infty} \frac{e^{\kappa(\ell-1) \log k}}{k} \lambda^{k(\ell-1)} \right| < \frac{\tilde{C}}{n} \sum_{\ell=2}^{\infty} |d_{\ell} z^{\ell}| (\ell-1)^{r+1}. \quad \square$$

Let us define

$$h_n := S_{n+1}^{-1} \circ g_n \circ S_n.$$

Lemma 3.14. *The maps h_n are of the form*

$$h_n = \lambda z + O(n^{-2}).$$

Proof. The definition of h_n immediately gives that $h_n(z) = \lambda z + O(1/n)$,

$$g_n \circ S_n = S_{n+1} \circ h_n,$$

and thus

$$\lambda z + \lambda \hat{S}_n(z) + \hat{g}_n(z + \hat{S}_n) = \lambda z + \hat{h}_n(z) + \hat{S}_{n+1}(\lambda z + \hat{h}_n),$$

which gives

$$\lambda \hat{S}_n(z) + \hat{g}_n(z) + \hat{g}'_n(z) \hat{S}_n(z) + O(\hat{S}_n^2) = \hat{h}_n(z) + \hat{S}_{n+1}(\lambda z) + \hat{S}'_{n+1}(\lambda z) \hat{h}_n + O(\hat{h}_n^2).$$

Hence by Lemma 3.12 we obtain

$$\hat{g}'_n(z) \hat{S}_n(z) + O(\hat{S}_n^2) = \hat{h}_n(z) (1 + \hat{S}'_{n+1}(\lambda z)) + O(\hat{h}_n^2).$$

Since $\hat{g}_n = O(1/n)$ and $\hat{S}_n = O(1/n)$, we get

$$\hat{h}_n(z)(1 + \hat{S}'_{n+1}(\lambda z)) + O(\hat{h}_n^2) = O\left(\frac{1}{n^2}\right).$$

Since $h_n(z) = \lambda z + O(1/n)$, it follows that $\hat{h}_n(z) = O(1/n^2)$. \square

Lemma 3.6 implies that the sequence (h_n) is rotationally linearizable; hence the same holds for (g_n) , (f_n) and finally (f_n) , which completes the proof of Theorem 2.2.

Remark 3.15. The proof of Theorem 2.2 also works for more general perturbations, for example

$$f_n(z) \asymp f(z) + \frac{1}{n}h_1(z) + \frac{\log n}{n}h_2(z),$$

where h_1 and h_2 are holomorphic around the origin. In this case we have two indexes κ_j , $j \in \{1, 2\}$, that can be computed using (5), where constants c_0 and c_1 are the coefficients of the linear part of the Taylor series of h_j at the origin. The following is a general version of Theorem 2.2:

- (1) If $\operatorname{Re}(\kappa_2) > 0$ then the sequence (f_n) is expansive.
- (2) If $\operatorname{Re}(\kappa_2) < 0$ then the sequence (f_n) is collapsing.
- (3) If $\operatorname{Re}(\kappa_2) = 0$ and:
 - (a) $\operatorname{Re}(\kappa_1) > 0$, then the sequence (f_n) is expansive.
 - (b) $\operatorname{Re}(\kappa_1) < 0$, then the sequence (f_n) is collapsing.
 - (c) $\operatorname{Re}(\kappa_1) = 0$, then the sequence (f_n) is rotationally linearizable; hence all limit maps have rank 1.

4. Existence of parabolic curves

The purpose of this section is to prove the following proposition.

Proposition 4.1. *Let $P(z, w) := (f(z) + \frac{\pi^2}{4}w, g(w))$, with $f(z) = z + z^2 + bz^3 + O(z^4)$ and $g(w) = w - w^2 + O(w^3)$. Then P has at least three parabolic curves: one is contained in the invariant fiber $w = 0$ and is an attracting petal for f ; the other two are graphs over the same petal \mathcal{P} in the parabolic basin \mathcal{B}_g . Moreover they are of the form*

$$\zeta^\pm(w) = \pm c_1 \sqrt{w} + c_2 w \pm c_3 w^{3/2} + O(w^2),$$

where $c_1 = \frac{\pi}{2}i$ and $c_2 = \frac{\pi^2}{8}b - \frac{1}{4}$.

Proposition 4.1 gives a positive answer to a question posed in [Astorg et al. 2016]. We note that the result does not follow from the results [Hakim 1998], as the two characteristic directions we consider are *degenerate*, in the language used by Hakim. The existence of three parabolic curves can be derived from [López-Hernanz and Rosas 2020]. However, their proof gives no guarantee that the parabolic curves ζ^\pm are graphs over the same petal in \mathcal{B}_g , which is crucial for our purpose.

Let us start by observing that P is semiconjugate to a map Q , holomorphic near the origin, given by

$$Q(z, \epsilon) = (f(z) + \frac{\pi^2}{4}\epsilon^2, \epsilon - \frac{1}{2}\epsilon^3 + O(\epsilon^5))$$

(with $\epsilon^2 = w$). The map Q has three characteristic directions: $\epsilon = 0$, $z = \frac{\pi}{2}i\epsilon$ and $z = -\frac{\pi}{2}i\epsilon$. It is clear that there is a parabolic curve tangent to the characteristic direction $\epsilon = 0$, namely the attracting petal for f in the invariant fiber $\{\epsilon = 0\}$. We call this parabolic curve the trivial curve. For the existence of the two other parabolic curves we will use a graph transform argument.

Let us write $Q(z, \epsilon) = (f_\epsilon(z), \tilde{g}(\epsilon))$, so that $f_\epsilon(z) = f(z) + \frac{\pi^2}{4}\epsilon^2$ and $\tilde{\epsilon} := \tilde{g}(\epsilon) = \sqrt{g(\epsilon^2)} = \epsilon - \frac{1}{2}\epsilon^3 + O(\epsilon^5)$. We are looking for parabolic curves of the form $\epsilon \rightarrow (\zeta(\epsilon), \epsilon)$, hence satisfying the equation

$$Q(\zeta(\epsilon), \epsilon) = (\zeta(\tilde{\epsilon}), \tilde{\epsilon}). \quad (8)$$

Equivalently we are looking for a function ζ , defined for ϵ in a parabolic petal of \tilde{g} , satisfying the functional equation

$$\zeta(\tilde{g}(\epsilon)) = f_\epsilon(\zeta(\epsilon)).$$

We will prove that Q has two parabolic curves ζ^\pm , corresponding to the characteristic directions $z = \pm \frac{\pi}{2}i\epsilon$, which are graphs over the same attracting petal of \tilde{g} in the right half-plane. This will complete the proof of Proposition 4.1, since these two parabolic curves can be lifted to parabolic curves of P satisfying the desired properties.

The key idea in proving the existence of $\zeta(\epsilon)$ is to start with sufficiently high-order jets $\zeta_1(\epsilon)$ of the formal solution to (8), and then apply a graph transform argument, starting with ζ_1 . By starting with higher-order jets, we obtain higher-order error estimates, but the constants in those estimates are likely to deteriorate. However, these estimates can be controlled by dropping the order of the error estimates by 1, and working with $|\epsilon| < \delta$, with δ depending on the order of the jets. It turns out that starting with jets of order 20 is sufficient to obtain convergence of the graph transforms. We do not claim that 20 is the minimal order for which convergence can be obtained, only that the order suffices for our purposes.

Lemma 4.2. *For every integer $n > 0$ there exists $\zeta_1(\epsilon) = c_1\epsilon + c_2\epsilon^2 + c_3\epsilon^3 + \dots + c_n\epsilon^n$ and $\delta > 0$ such that $|\zeta_1(\tilde{\epsilon}) - f_\epsilon(\zeta_1(\epsilon))| < |\epsilon|^n$ for all $|\epsilon| < \delta$. Moreover we have $c_1 = \pm \frac{\pi}{2}i$ and $c_2 = \frac{\pi^2}{8}b - \frac{1}{4}$.*

Proof. Recall from [Astorg et al. 2016] that by choosing $\zeta_1(\epsilon) = c_1\epsilon + c_2\epsilon^2$, with $c_1 = \pm \frac{\pi}{2}i$ and $c_2 = \frac{\pi^2}{8}b - \frac{1}{4}$, we obtain

$$|\zeta_1(\tilde{\epsilon}) - f_\epsilon(\zeta_1(\epsilon))| < O(|\epsilon|^4).$$

Now suppose that c_1, \dots, c_n are found such that for $\zeta(\epsilon) = c_1\epsilon + \dots + c_n\epsilon^n$ we have

$$|\zeta_1(\tilde{\epsilon}) - f_\epsilon(\zeta_1(\epsilon))| < O(|\epsilon|^{n+2}).$$

Let $E_n(\epsilon) := f_\epsilon(\zeta_1(\epsilon)) - \zeta_1(\tilde{\epsilon})$. For $c_{n+1} \in \mathbb{C}$, let

$$E_{n+1}(\epsilon) := f_\epsilon(\zeta_1(\epsilon) + c_{n+1}\epsilon^{n+1}) - \zeta_1(\tilde{\epsilon}) - c_{n+1}\tilde{\epsilon}^{n+1};$$

we shall prove that there exists some c_{n+1} such that $E_{n+1} = O(\epsilon^{n+3})$. Indeed,

$$\begin{aligned} f_\epsilon(\zeta_1(\epsilon) + c_{n+1}\epsilon^{n+1}) &= f_\epsilon(\zeta_1(\epsilon)) + f'(\zeta_n(\epsilon))c_{n+1}\epsilon^{n+1} + O(\epsilon^{2n+2}) \\ &= f_\epsilon(\zeta_1(\epsilon)) + (1 + 2c_1\epsilon)c_{n+1}\epsilon^{n+1} + O(\epsilon^{n+3}). \end{aligned}$$

On the other hand, we have $c_{n+1}\tilde{\epsilon}^{n+1} = c_{n+1}\epsilon^{n+1} + O(\epsilon^{n+3})$; so

$$E_{n+1}(\epsilon) = E_n(\epsilon) + 2c_1c_{n+1}\epsilon^{n+2} + O(\epsilon^{n+3}).$$

Since $E_n(\epsilon) = O(\epsilon^{n+2})$ (and $c_1 \neq 0$), we may therefore find some value of c_{n+1} for which $E_{n+1}(\epsilon) = O(\epsilon^{n+3})$.

We conclude that if δ is small enough then $|\zeta_1(\tilde{\epsilon}) - f_\epsilon(\zeta_1(\epsilon))| < |\epsilon|^n$ for all $|\epsilon| < \delta$. \square

Remark 4.3. The choice of parabolic curve is determined by the choice of c_1 . From now on we will assume that $c_1 = \frac{\pi}{2}i$; for the case $c_1 = -\frac{\pi}{2}i$ the proofs are essentially the same.

For $R \in \mathbb{C}$ we write $\mathbb{H}_R = \{Z \in \mathbb{C} : \arg(Z - R) \in (-\frac{\pi}{2} - \epsilon_0, \frac{\pi}{2} + \epsilon_0)\}$ for some $\epsilon_0 > 0$, and

$$\mathcal{P}_\delta = \{\epsilon \in \mathbb{C} : \epsilon^{-2} \in \mathbb{H}_{\delta^{-2}} \text{ and } \operatorname{Re}(\epsilon) > 0\}.$$

For $\delta > 0$ sufficiently small the petal \mathcal{P}_δ is forward-invariant under \tilde{g} , i.e., $\tilde{g}(\mathcal{P}_\delta) \subset \mathcal{P}_\delta$. Recall the existence of Fatou coordinates on \mathcal{P}_δ : the function \tilde{g} is conjugate to the translation $T_1 : Z \mapsto Z + 1$ via a conjugation of the form

$$Z = \frac{1}{\epsilon^2} + \alpha \log(\epsilon) + o(1),$$

where the constant α depends on g . All forward orbits in \mathcal{P}_δ converge to 0 tangent to the positive real axis, and the conjugation gives the estimates

$$|\operatorname{Re}(\tilde{g}^k(\epsilon))| < \frac{C}{\sqrt{k}} \quad \text{and} \quad |\operatorname{Im}(\tilde{g}^k(\epsilon))| < \frac{C}{k} \quad (9)$$

for a uniform $C > 0$ depending on δ . We note that by choosing δ sufficiently small, the constant C can be chosen arbitrarily small as well.

Lemma 4.4. *Let $n > 0$ and $\zeta_1(\epsilon)$ be as in Lemma 4.2. There exist $\delta, A > 0$ such that for every $|\epsilon| < \delta$ we have*

$$|f^{-1}(f(\zeta_1(\epsilon)) + 3\epsilon^4) - \zeta_1(\epsilon)| \leq A|\epsilon|^4.$$

Proof. The Taylor series expansion of f gives

$$|f^{-1}(f(\zeta_1(\epsilon)) + 3\epsilon^4) - \zeta_1(\epsilon)| \leq \sum_{i=1}^{\infty} \left| \frac{(f^{-1})^{(i)}(f(\zeta_1(\epsilon)))}{i!} \right| 3^i |\epsilon^4|^i,$$

and the desired estimate follows immediately. \square

Lemma 4.5. *Let $n > 0$ and $\zeta_1(\epsilon)$ be as in Lemma 4.2, $A > 0$ and $\delta > 0$ sufficiently small. Let $(\zeta_k(\epsilon))$ be any sequence of holomorphic functions defined on \mathcal{P}_δ and satisfying*

$$|\zeta_k(\epsilon) - \zeta_1(\epsilon)| < A|\epsilon|^4.$$

Then there exists $C_1 > 0$, depending on ζ_1 , such that

$$\left| \prod_{s=\ell}^k f'(\zeta_s(\tilde{g}^{k+1-s}(\epsilon))) \right|^{-1} < C_1 \cdot (k+1-\ell)$$

for all $\epsilon \in \mathcal{P}_\delta$ and every $0 < \ell \leq k$.

Proof. Let us write $x_k = \operatorname{Re}(\tilde{g}^k(\epsilon)) > 0$ and $y_k = \operatorname{Im}(\tilde{g}^k(\epsilon))$. Estimates (9) imply

$$\sum_{k=0}^{\infty} |\tilde{g}^k(\epsilon)|^3 < K < \infty \quad \text{for all } \epsilon \in \mathcal{P}_\delta. \quad (10)$$

Since by assumption $|\zeta_s(\epsilon) - \zeta_1(\epsilon)| < A|\epsilon|^4$ for every $s \geq 1$, it follows that $\zeta_s(\epsilon) = c_1\epsilon + c_2\epsilon^2 + O(\epsilon^3)$ and

$$|f'(\zeta_s(\epsilon)) - f'(\zeta_1(\epsilon))| < B|\epsilon^4|,$$

where $B > 0$ depends only on ζ_1 and A .

Observe that $f'(z) = 1 + 2z + 3bz^2 + O(z^3) = e^{2z + (3b-2)z^2 + O(z^3)}$; hence we obtain

$$f'(\zeta_s(\epsilon)) = e^{\pi i \epsilon + (\frac{1}{2}\pi^2(1-b) - \frac{1}{2})\epsilon^2 + O(\epsilon^3)},$$

where the bound $O(\epsilon^3)$ is uniform with respect to s .

Therefore we can find $C_1 > 0$ such that

$$\begin{aligned} \left| \prod_{s=\ell}^k f'(\zeta_s(\tilde{g}^{k+1-s}(\epsilon))) \right| &> \left| e^{\sum_{s=\ell}^k \operatorname{Re}(\pi i \tilde{g}^{k+1-s}(\epsilon) + (\frac{1}{2}\pi^2(1-b) - \frac{1}{2})(\tilde{g}^{k+1-s}(\epsilon))^2) + O((\tilde{g}^{k+1-s}(\epsilon))^3)} \right| \\ &> \frac{1}{C_1} \left| e^{\sum_{s=\ell}^k -\pi y_{k+1-s} + (\frac{1}{2}\pi^2(1-\operatorname{Re}(b)) - \frac{1}{2})x_{k+1-s}^2} \right| \\ &> \frac{1}{C_1} \left| e^{-\sum_{s=\ell}^k 1/(k+1-s)} \right| \\ &> \frac{1}{C_1(k+1-\ell)}. \end{aligned}$$

In the first inequality we used the fact that $|e^z| = e^{\operatorname{Re}(z)}$. The second inequality follows from estimates (9) and (10). The third inequality depends on the constant C from (9) being sufficiently small, which can be guaranteed by taking sufficiently small δ . \square

Remark 4.6. Note that the estimates in Lemmas 4.2, 4.4 and 4.5 hold regardless of the choice of n in the definition of ζ_1 . If n is increased, then all estimates hold, with the same constants, for δ sufficiently small. It turns out that it will be sufficient for us to work with $n = 20$, and we will work with this choice from now on.

Lemma 4.7. *There exists sufficiently small $\delta > 0$ such that for every $k \geq 2$ and every $\epsilon \in \mathcal{P}_\delta$ we have*

$$|\tilde{g}^k(\epsilon)|^{19}k + |\tilde{g}^k(\epsilon)|^{39}(k-1) + \sum_{\ell=2}^{k-1} \frac{|\tilde{g}^{k+1-\ell}(\epsilon)|^{23}(k-\ell)}{(\ell-1)^4} + \frac{|\tilde{g}(\epsilon)|^{23}}{(k-1)^4} < \frac{4|\epsilon|^{12}}{k^2}.$$

Proof. We will prove that each of the four terms in the left-hand summation is bounded by $|\epsilon|^{12}/k^2$. It follows from (9) that for every $0 \leq \ell \leq 19$ we have

$$|\tilde{g}^k(\epsilon)|^{19}k < \frac{C^{19}k}{|k+1/\epsilon^2|^{19/2}} < \frac{C^{19}|\epsilon|^\ell k}{k^{(19-\ell)/2}}.$$

If we choose $\ell = 13$ and assume that δ is small enough, then we get

$$|\tilde{g}^k(\epsilon)|^{19} k < \frac{|\epsilon|^{12}}{k^2}$$

for $\epsilon \in \mathcal{P}_\delta$. The desired bound for a second term follows immediately from the inequality

$$|\tilde{g}^k(\epsilon)|^{39} (k-1) < |\tilde{g}^k(\epsilon)|^{19} k.$$

Next observe that for every $k \geq 2$ we have

$$\frac{|\tilde{g}(\epsilon)|^{23}}{(k-1)^4} < \frac{2^2 |\epsilon|^{23}}{(2(k-1)^2)^2} < \frac{|\epsilon|^{12}}{k^2},$$

where the last inequality holds for sufficiently small δ . Finally, for the third term in the summation we use (9) to obtain

$$\sum_{\ell=2}^{k-1} \frac{|\tilde{g}^{k+1-\ell}(\epsilon)|^{23} (k-\ell)}{(\ell-1)^4} < \sum_{\ell=2}^{k-1} \frac{C^{10} |\epsilon|^{13} (k-\ell)}{(k+1-\ell)^5 (\ell-1)^4} < C^{10} |\epsilon| \sum_{\ell=2}^{k-1} \frac{|\epsilon|^{12}}{(k-\ell)^4 (\ell-1)^4}.$$

In order to obtain the desired bound it suffices to prove that

$$\sum_{\ell=2}^{k-1} \frac{1}{(k-\ell)^4 (\ell-1)^4} < \frac{4^4}{k^2}.$$

First observe that

$$\frac{1}{(\ell-1)(k-\ell)} \leq \frac{4}{k}$$

for every $k \geq 3$ and $2 \leq \ell \leq k-1$. To see this let us set $s = \ell-1$ and $t = k-1$. The above inequality now translates to

$$\frac{1}{s(t-s)} \leq \frac{4}{t+1}$$

for $t \geq 2$ and $1 \leq s \leq t-1$, and hence to

$$p_t(s) := 4s^2 - 4ts + t + 1 \leq 0.$$

Observe that $p_t(1) < 0$ and that roots of $p_t(s)$ lie outside the closed interval $[1, t-1]$. Therefore we obtain

$$\sum_{\ell=2}^{k-1} \frac{1}{(k-\ell)^4 (\ell-1)^4} < \sum_{\ell=2}^{k-1} \frac{4^4}{k^4} < \frac{4^4}{k^2},$$

and hence for δ sufficiently small

$$\sum_{\ell=2}^{k-1} \frac{|\tilde{g}^{k+1-\ell}(\epsilon)|^{23} (k-\ell)}{(\ell-1)^4} < \frac{|\epsilon|^{12}}{k^2}. \quad \square$$

Proof of Proposition 4.1. As we remarked at the beginning of this section, it is enough to prove that Q has two parabolic curves ζ^\pm corresponding to the characteristic directions $z = \pm \frac{\pi}{2} i \epsilon$, both curves

graph over the same attracting petal of \tilde{g} in the right half-plane. By Lemma 4.2 there exist $\delta > 0$ and $\zeta_1(\epsilon) = c_1\epsilon + c_2\epsilon^2 + \dots + c_{20}\epsilon^{20}$ such that $|\zeta_1(\tilde{\epsilon}) - f_\epsilon(\zeta_1(\epsilon))| < |\epsilon|^{20}$ for $|\epsilon| < \delta$. Let $A > 0$ be as in Lemma 4.4, and let $C_1 > 0$ be the constant defined in Lemma 4.5.

We will show that the sequence of functions defined inductively by

$$\zeta_{k+1}(\epsilon) := f_\epsilon^{-1}(\zeta_k(\tilde{\epsilon}))$$

is convergent and that the limit satisfies the functional equation (8). Let us define

$$E_k(\epsilon) := \zeta_k(\tilde{\epsilon}) - f_\epsilon(\zeta_k(\epsilon))$$

and observe that

$$\zeta_{k+1}(\epsilon) = f_\epsilon^{-1}(\zeta_k(\tilde{\epsilon})) = f_\epsilon^{-1}(f_\epsilon(\zeta_k(\epsilon)) + E_k(\epsilon))$$

and hence

$$f_\epsilon(\zeta_{k+1}(\epsilon)) = f_\epsilon(\zeta_1(\epsilon)) + \sum_{\ell=1}^k E_\ell(\epsilon).$$

Note that we can replace f_ϵ by f on both sides, giving

$$\zeta_{k+1}(\epsilon) = f^{-1}\left(f(\zeta_1(\epsilon)) + \sum_{\ell=1}^k E_\ell(\epsilon)\right),$$

and hence

$$\zeta_{k+1}(\epsilon) = \zeta_1(\epsilon) + \sum_{i=1}^{\infty} \frac{(f^{-1})^{(i)}(f(\zeta_1(\epsilon)))}{i!} \left(\sum_{\ell=1}^k E_\ell(\epsilon)\right)^i.$$

We will prove that $|E_k(\epsilon)| < |\epsilon|^4/|k-1|^2$ for every $k \geq 2$ on some small petal \mathcal{P}_δ . This will imply that the sequence ζ_{k+1} converges to a parabolic curve ζ on \mathcal{P}_δ for sufficiently small δ .

We claim there exists $\delta > 0$ such that for every $\epsilon \in \mathcal{P}_\delta$ and every $k > 1$ the following two statements hold:

$I_k(1)$: $|\zeta_k(\epsilon) - \zeta_1(\epsilon)| < A|\epsilon|^4$, and

$I_k(2)$: $|E_k(\epsilon)| < 4|\epsilon|^{12}/|k-1|^2 < |\epsilon|^4/|k-1|^2$.

We will prove these two statements simultaneously by induction on k .

Step 1: First we prove $I_2(1)$. By definition

$$\zeta_2 = \zeta_1(\epsilon) + \sum_{i=1}^{\infty} \frac{(f^{-1})^{(i)}(f(\zeta_1(\epsilon)))}{i!} (E_1(\epsilon))^i;$$

hence by Lemma 4.4 we obtain the desired inequality.

Next we prove that $I_2(2)$. Observe that for sufficiently small δ we get

$$\begin{aligned} |E_2(\epsilon)| &< \left| \frac{E_1(\tilde{\epsilon})}{f'(\zeta_1(\tilde{\epsilon}))} \right| + C_2|E_1(\tilde{\epsilon})|^2 < C_1|\tilde{g}(\epsilon)|^{20} + C_2|\epsilon||\tilde{g}(\epsilon)|^{40} \\ &< C_1|\epsilon||\tilde{g}(\epsilon)|^{19} + C_2|\epsilon||\tilde{g}(\epsilon)|^{39} < 4|\epsilon|^{12}. \end{aligned}$$

Here C_1 is the constant introduced in Lemma 4.5.

Step 2: Now let us assume that $I_\ell(1)$ and $I_\ell(2)$ hold for every $2 \leq \ell \leq k$. Observe that

$$|\zeta_{k+1}(\epsilon) - \zeta_1(\epsilon)| < \sum_{i=1}^{\infty} \left| \frac{(f^{-1})^{(i)}(f(\zeta_1(\epsilon)))}{i!} \right| \left| \sum_{\ell=1}^k E_\ell(\epsilon) \right|^i.$$

Since $|E_\ell(\epsilon)| < |\epsilon|^4 / |\ell - 1|^2$ for $\ell \geq 2$ and $|E_1(\epsilon)| < |\epsilon|^4$, we get

$$\left| \sum_{\ell=1}^k E_\ell(\epsilon) \right| < 3|\epsilon|^4;$$

hence by Lemma 4.4 inequality $I_{k+1}(1)$ holds.

Observe that

$$\begin{aligned} E_{k+1}(\epsilon) &= \zeta_{k+1}(\tilde{\epsilon}) - f_\epsilon(\zeta_{k+1}(\epsilon)) = \zeta_{k+1}(\tilde{\epsilon}) - \zeta_k(\tilde{\epsilon}) \\ &= f_{\tilde{\epsilon}}^{-1}(f_{\tilde{\epsilon}}(\zeta_k(\tilde{\epsilon})) + E_k(\tilde{\epsilon})) - \zeta_k(\tilde{\epsilon}) \\ &= f^{-1}(f(\zeta_k(\tilde{\epsilon})) + E_k(\tilde{\epsilon})) - \zeta_k(\tilde{\epsilon}) \\ &= (f^{-1})'(f(\zeta_k(\tilde{\epsilon}))) \cdot E_k(\tilde{\epsilon}) + O(E_k(\tilde{\epsilon})^2), \end{aligned}$$

where the constant in the order can be chosen independently from k . It follows that there exists $C_2 > 0$ independent of k such that

$$|E_{k+1}(\epsilon)| < \left| \frac{E_k(\tilde{\epsilon})}{f'(\zeta_k(\tilde{\epsilon}))} \right| + C_2 |E_k(\tilde{\epsilon})|^2. \quad (11)$$

Using the inequality (11) successively we obtain

$$\begin{aligned} |E_{k+1}(\epsilon)| &< \left| \frac{E_k(\tilde{g}(\epsilon))}{f'(\zeta_k(\tilde{g}(\epsilon)))} \right| + C_2 |E_k(\tilde{g}(\epsilon))|^2 \\ &< \left| \frac{E_{k-1}(\tilde{g}^2(\epsilon))}{f'(\zeta_{k-1}(\tilde{g}^2(\epsilon))) \cdot f'(\zeta_k(\tilde{g}(\epsilon)))} \right| + C_2 \frac{|E_{k-1}(\tilde{g}^2(\epsilon))|^2}{|f'(\zeta_k(\tilde{g}(\epsilon)))|} + C_2 |E_k(\tilde{g}(\epsilon))|^2 \\ &< \frac{|E_1(\tilde{g}^k(\epsilon))|}{\prod_{\ell=1}^k |f'(\zeta_\ell(\tilde{g}^{k+1-\ell}(\epsilon)))|} + C_2 \sum_{\ell=1}^{k-1} \frac{|E_\ell(\tilde{g}^{k+1-\ell}(\epsilon))|^2}{\prod_{s=\ell+1}^k |f'(\zeta_s(\tilde{g}^{k+1-s}(\epsilon)))|} + C_2 |E_k(\tilde{\epsilon})|^2. \quad (12) \end{aligned}$$

Combining (12) and Lemma 4.5 gives

$$\begin{aligned} &|E_{k+1}(\epsilon)| \\ &< C_1 |\tilde{g}^k(\epsilon)|^{20} k + C_1 C_2 |\tilde{g}^k(\epsilon)|^{40} (k-1) + 16 C_1 C_2 \sum_{\ell=2}^{k-1} \frac{|\tilde{g}^{k+1-\ell}(\epsilon)|^{24} (k-\ell)}{(\ell-1)^4} + 16 C_1 \frac{|\tilde{g}(\epsilon)|^{24}}{(k-1)^4} \\ &< C_1 |\epsilon| |\tilde{g}^k(\epsilon)|^{19} k + C_1 C_2 |\epsilon| |\tilde{g}^k(\epsilon)|^{39} (k-1) + 16 C_1 C_2 |\epsilon| \sum_{\ell=2}^{k-1} \frac{|\tilde{g}^{k+1-\ell}(\epsilon)|^{23} (k-\ell)}{(\ell-1)^4} + 16 C_1 |\epsilon| \frac{|\tilde{g}(\epsilon)|^{23}}{(k-1)^4}. \end{aligned}$$

If δ is sufficiently small this last inequality together with Lemma 4.7 implies

$$|E_{k+1}(\epsilon)| < \frac{4|\epsilon|^{12}}{k^2} < \frac{|\epsilon|^4}{k^2},$$

completing the proof of $I_{k+1}(2)$ and thus the induction argument. We emphasize that throughout the proof δ can be chosen dependently of k .

To summarize, the equation

$$\zeta_{k+1}(\epsilon) = f^{-1}\left(f(\zeta_1(\epsilon)) + \sum_{\ell=1}^k E_\ell(\epsilon)\right)$$

implies that for sufficiently small δ the sequence ζ_k converges on \mathcal{P}_δ to a parabolic curve ζ satisfying $\zeta(\tilde{\epsilon}) = f_\epsilon(\zeta(\epsilon))$. Recall that we have only proven the existence of a parabolic curve for $c_1 = \frac{\pi}{2}i$. For $c_1 = -\frac{\pi}{2}i$ we can use same arguments as above, but we might get a different value for δ . Since the parabolic petals are nested and forward invariant, both parabolic curves are graphs over the petal with minimal δ . \square

From the proof it follows that

$$\zeta^\pm(\epsilon) = \pm c_1 \epsilon + c_2 \epsilon^2 \pm c_3 \epsilon^3 + O(\epsilon^4),$$

where $c_1 = \frac{\pi}{2}i$ and $c_2 = \frac{\pi^2}{8}b - \frac{1}{4}$.

5. Estimates on convergence towards Lavaurs map

5A. Preliminaries. The goal of this section is to obtain explicit estimates for one of the main objects to appear in our arguments: the functions $A(\epsilon, z)$ and $A_0(z)$, which measure how much the dynamics differ from a translation after a certain change of coordinates. The key difference between this section and the corresponding computations in [Astorg et al. 2016] is that we now know that we have two exactly invariant parabolic curves ζ^\pm , instead of invariant jets. This is used crucially in the proof of Proposition 5.5.

Definition 5.1. Let $f_w(z) := f(z) + \frac{\pi^2}{4}w$, where $f(z) = z + z^2 + z^3 + O(z^4)$ is a degree- d polynomial. Let $g(w) = w - w^2 + O(w^3)$ be a degree- d polynomial.

In what follows, we set $\epsilon := \sqrt{w}$, working throughout with the branch that takes positive values on the positive real axis. We note that this branch is well-defined on the parabolic basin of the polynomial g . Abusing notation, we write $f_\epsilon(z) := f(z) + \frac{\pi^2}{4}\epsilon^2$ and $\zeta^\pm(\epsilon) = \pm i \frac{\pi}{2}\epsilon + c_2 \epsilon^2 + O(\epsilon^3)$, where ζ^\pm are the parabolic curves constructed in the preceding section. Let $\tilde{g}(\epsilon) := \sqrt{g(\epsilon^2)} = \epsilon - \frac{1}{2}\epsilon^3 + O(\epsilon^5)$ (\tilde{g} is analytic near $\epsilon = 0$).

Let us first record here the following lemma for later use:

Lemma 5.2. Let $w_0 \in \mathcal{B}_g$ and let $\epsilon_j := \sqrt{g^{n^2+j}(w_0)}$. For $1 \leq j \leq n$, we have

$$\epsilon_j = \frac{1}{n} - \frac{j}{2n^3} - \frac{\phi_g^l(w_0)}{2n^3} + o\left(\frac{1}{n^3}\right).$$

Proof. Let us write $w_{n^2+j} := g^{n^2+j}(w_0)$. We have

$$\phi_g(w_{n^2+j}) = \phi_g^l(w_0) + n^2 + j = \frac{1}{w_{n^2+j}} + o(1)$$

(note that we assume here $g(w) = w - w^2 + w^3 + O(w^4)$). Therefore

$$w_{n^2+j} = \frac{1}{n^2 + j + \phi_g^t(w_0) + o(1)},$$

and

$$\begin{aligned} \epsilon_j &= \sqrt{w_{n^2+j}} = \frac{1}{n} \left(1 + \frac{j + \phi_g^t(w_0)}{n^2} + o\left(\frac{1}{n^2}\right) \right)^{-1/2} \\ &= \frac{1}{n} \left(1 - \frac{1}{2} \frac{j + \phi_g^t(w_0)}{n^2} + o\left(\frac{1}{n^2}\right) \right). \end{aligned} \quad \square$$

Definition 5.3. Let

$$\begin{aligned} \psi_\epsilon^t(z) &:= \frac{1}{i\pi} \log \left[\frac{\zeta^+(\epsilon) - z}{z - \zeta^-(\epsilon)} \right] + 1, \\ \psi_\epsilon^o(z) &:= \frac{1}{i\pi} \log \left[\frac{\zeta^+(\epsilon) - z}{z - \zeta^-(\epsilon)} \right] - 1, \end{aligned}$$

where \log is the principal branch of the logarithm.

Note with that choice of branch, ψ is defined on $\mathbb{C} \setminus L_\epsilon$, where L_ϵ is the real line through $\zeta^+(\epsilon)$ and $\zeta^-(\epsilon)$ minus the segment $[\zeta^-(\epsilon), \zeta^+(\epsilon)]$. In particular, ψ_ϵ^t and ψ_ϵ^o are both defined in a disk centered at $z = 0$ whose radius is of order ϵ .

It will also be useful to note that

$$(\psi_\epsilon^{t/o})^{-1}(Z) = \frac{\zeta^+(\epsilon) - e^{\pm i\pi Z} \zeta^-(\epsilon)}{1 - e^{\pm i\pi Z}} = -\frac{\pi}{2} \epsilon \cot\left(\pm \frac{\pi Z}{2}\right) + O(\epsilon^2). \quad (13)$$

Definition 5.4. Let

- (1) $A(\epsilon, z) := \psi_{\zeta^-(\epsilon)}^{t/o} \circ f_\epsilon(z) - \psi_\epsilon^{t/o}(z) - \epsilon$,
- (2) $A_0(z) := -1/f(z) + 1/z - 1$.

Note that the formula for $A(\epsilon, z)$ does not depend on whether the ingoing or outgoing coordinate ψ_ϵ is used, and is therefore well-defined.

Proposition 5.5. *We have*

- (1) A_0 is analytic near zero,
- (2) there exists $r > 0$ such that for all $\epsilon \neq 0$ in a neighborhood of zero, $A(\epsilon, \cdot)$ is analytic on $\mathbb{D}(0, r)$.

Proof. (1) A quick computation shows that

$$A_0(z) = \frac{f(z) - z - zf(z)}{zf(z)} = \frac{O(z^2)}{1 + O(z)},$$

from which the conclusion easily follows.

For (2), note that

$$\begin{aligned} A(\epsilon, z) &= \frac{1}{i\pi} \log \left(\frac{\zeta^+(\tilde{g}(\epsilon)) - f_\epsilon(z)}{f_\epsilon(z) - \zeta^-(\tilde{g}(\epsilon))} ; \frac{\zeta^+(\epsilon) - z}{z - \zeta^-(\epsilon)} \right) - \epsilon \\ &= \frac{1}{i\pi} \log \left(\frac{f_\epsilon(\zeta^+(\epsilon)) - f_\epsilon(z)}{\zeta^+(\epsilon) - z} ; \frac{f_\epsilon(z) - f_\epsilon(\zeta^-(\epsilon))}{z - \zeta^-(\epsilon)} \right) - \epsilon. \end{aligned}$$

From the above expression we see that the singularities at $z = \zeta^\pm(\epsilon)$ are in fact removable, unless one of the points coincides with a critical point of f . The fact that these critical points are bounded away from zero completes the proof. \square

Lemma 5.6. *Let K be a compact subset of \mathbb{C}^* . There exists $C = C_K > 0$ such that, for all $z \in K$,*

$$\left| \frac{z - \zeta^+(\epsilon)}{z - \zeta^-(\epsilon)} - \left(1 - \frac{i\pi}{z}\epsilon - \frac{\pi^2}{2z^2}\epsilon^2 \right) \right| \leq C\epsilon^3.$$

Proof. For $z \in K$, we have

$$\begin{aligned} \frac{\zeta^+(\epsilon) - z}{z - \zeta^-(\epsilon)} &= \frac{\zeta^+(\epsilon)}{z - \zeta^-(\epsilon)} - \frac{z}{z - \zeta^-(\epsilon)} \\ &= \frac{\zeta^+(\epsilon)}{z} \left(\frac{1}{1 - \zeta^-(\epsilon)/z} \right) - \frac{1}{1 - \zeta^-(\epsilon)/z} \\ &= \frac{\zeta^+(\epsilon)}{z} \left(1 + \frac{\zeta^-(\epsilon)}{z} + O(\epsilon^2) \right) - \left(1 + \frac{\zeta^-(\epsilon)}{z} + \left(\frac{\zeta^-(\epsilon)}{z} \right)^2 + O(\epsilon^3) \right) \\ &= \frac{c_1\epsilon + c_2\epsilon^2 - \frac{c_1^2}{z}\epsilon^2}{z} - 1 - \frac{-c_1\epsilon + c_2\epsilon^2}{z} - \frac{c_1^2\epsilon^2}{z^2} + O(\epsilon^3) \\ &= -1 + \frac{2c_1}{z}\epsilon - \frac{2c_1^2}{z^2}\epsilon^2 + O(\epsilon^3) \\ &= -1 + \frac{i\pi}{z}\epsilon + \frac{\pi^2}{2z^2}\epsilon^2 + O(\epsilon^3). \end{aligned} \quad \square$$

Lemma 5.7. *Let K be a compact subset of \mathbb{C}^* . Then*

$$\frac{f_\epsilon(z) - f_\epsilon(\zeta^+(\epsilon))}{f_\epsilon(z) - f_\epsilon(\zeta^-(\epsilon))} = 1 - \frac{i\pi}{f(z)}\epsilon - \frac{\pi^2}{2f(z)^2}\epsilon^2 + O(\epsilon^3).$$

As in the previous lemma the constant in the O depends on K .

Proof. The invariance of the parabolic curves gives

$$\begin{aligned} \frac{f_\epsilon(z) - f_\epsilon(\zeta^+(\epsilon))}{f_\epsilon(z) - f_\epsilon(\zeta^-(\epsilon))} &= \frac{f_\epsilon(z) - \zeta^+(\tilde{g}(\epsilon))}{f_\epsilon(z) - \zeta^-(\tilde{g}(\epsilon))} = 1 - \frac{i\pi}{f_\epsilon(z)}\tilde{g}(\epsilon) + \frac{\pi^2}{2f_\epsilon(z)^2}\tilde{g}(\epsilon)^2 + O(\epsilon^3) \\ &= 1 - \frac{i\pi}{f(z)}\epsilon + \frac{\pi^2}{2f(z)^2}\epsilon^2 + O(\epsilon^3). \end{aligned}$$

The last equality uses the fact that $\tilde{g}(\epsilon) = \epsilon + O(\epsilon^3)$. \square

Proposition 5.8. *There exists a constant $C_0 \in \mathbb{C}$ (depending only on f and g) such that*

$$A(\epsilon, z) = \epsilon A_0(z) + \epsilon^3 C_0 + O(\epsilon^4, \epsilon^3 z),$$

where the constants in the O are uniform for $(z, \epsilon) \in \mathbb{C}^2$ near $(0, 0)$ (with $\operatorname{Re}(\epsilon) > 0$).

Proof. Let K be a compact of \mathbb{C}^* . Then by the two previous lemmas, we have

$$\begin{aligned} A(\epsilon, z) &= \frac{1}{i\pi} \log \left(\frac{z - \zeta^-(\epsilon)}{z - \zeta^+(\epsilon)} \cdot \frac{f_\epsilon(z) - f_\epsilon(\zeta^+(\epsilon))}{f_\epsilon(z) - f_\epsilon(\zeta^-(\epsilon))} \right) - \epsilon \\ &= -\frac{1}{i\pi} \log \left(1 - \frac{i\pi}{z} \epsilon - \frac{\pi^2}{2z^2} \epsilon^2 + O(\epsilon^3) \right) + \frac{1}{i\pi} \log \left(1 - \frac{i\pi}{f(z)} \epsilon - \frac{\pi^2}{2f(z)^2} \epsilon^2 + O(\epsilon^3) \right) - \epsilon \\ &= \frac{\epsilon}{z} - \frac{\epsilon}{f(z)} - \epsilon + O(\epsilon^3) = \epsilon A_0(z) + O(\epsilon^3). \end{aligned}$$

Here the constant in the O still depends on $K \subset \mathbb{C}^*$. Let $\phi_\epsilon(z) := (A(\epsilon, z) - \epsilon A_0(z))/\epsilon^3$. By Proposition 5.5, ϕ_ϵ is holomorphic on $\mathbb{D}(0, r)$. We have proved that for all compact $K \subset \mathbb{C}^*$, for all $z \in K$, and for all small $\epsilon \neq 0$ with $\operatorname{Re}(\epsilon) > 0$, we have $|\phi_\epsilon(z)| \leq C_K$. By taking $K = \{|z| = \frac{1}{2}r\}$ we therefore obtain the same estimate $|\phi_\epsilon(z)| \leq C_K$ for all $|z| \leq \frac{1}{2}r$ because of the maximum modulus principle. This gives the desired uniformity. \square

Lemma 5.9. *If $\zeta^\pm(\epsilon) = \pm \frac{\pi}{2}i\epsilon + c_2\epsilon^2 + c_3^\pm\epsilon^3 + O(\epsilon^4)$ and $f(z) = z + z^2 + z^3 + bz^4 + O(z^5)$, then*

$$C_0 = \frac{-3b\pi^3 + 2\pi^3 + 12c_2\pi + 12i(c_3^- - c_3^+)}{12\pi}.$$

Proof. By repeating the computations from Lemmas 5.6 and 5.7 with one additional order of significance, one obtains

$$\begin{aligned} \frac{z - \zeta^+(\epsilon)}{z - \zeta^-(\epsilon)} &= 1 - \frac{i\pi}{z} \epsilon - \frac{\pi^2}{2z^2} \epsilon^2 + \left(\frac{c_3^- - c_3^+}{z} - \frac{i\pi c_2}{z^2} + \frac{i\pi^3}{4z^3} \right) \epsilon^3 + O(\epsilon^3), \\ \frac{f_\epsilon(z) - f_\epsilon(\zeta^+(\epsilon))}{f_\epsilon(z) - f_\epsilon(\zeta^-(\epsilon))} &= 1 - \frac{i\pi}{f(z)} \epsilon - \frac{\pi^2}{2f(z)^2} \epsilon^2 + \left(\frac{i\pi^3}{4f(z)^2} + \frac{c_3^- - c_3^+}{f(z)} - \frac{i\pi c_2}{f(z)^2} + \frac{i\pi^3}{4f(z)^3} \right) \epsilon^3 + O(\epsilon^4). \end{aligned}$$

Plugging these two equations into the formula for $A(\epsilon, z)$, and using the power series expansions of $1/f(z)^j$ for $j = 1, \dots, 3$, one notices again that all terms involving negative powers of z cancel, either by the argument used in the proof of the previous proposition, or by lengthy computations using

$$c_2 = \frac{\pi^2}{8} - \frac{1}{4}.$$

Summing the terms that do not depend on z gives the desired result. \square

Lemma 5.10. *We have*

$$c_3^+ = -c_3^- = \frac{1}{i\pi} \left(\frac{3}{16} + \frac{5\pi^4}{64} - \frac{b\pi^4}{16} - \frac{\pi^2}{4} \right).$$

We will omit the proof, which is a long but direct computation, starting from the functional equation $f_\epsilon \circ \zeta^\pm(\epsilon) = \zeta^\pm \circ \tilde{g}(\epsilon)$ and identifying coefficients in powers of ϵ .

In particular, it follows that

$$\begin{aligned} C_0 &= -\frac{b\pi^2}{4} - \frac{1}{4} + \frac{7\pi^2}{24} + \left(\frac{b\pi^2}{8} + \frac{1}{2} - \frac{5\pi^2}{32} - \frac{3}{8\pi^2} \right) \\ &= \frac{-b\pi^2}{8} + \frac{13\pi^2}{96} - \frac{3}{8\pi^2} + \frac{1}{4}. \end{aligned}$$

5B. Convergence result. For the rest of Section 5 we fix a compact subset $K \times K' \subset \mathcal{B}_f \times \mathcal{B}_g$ and a point $(z_0, w_0) \in K \times K'$. Moreover we assume that n is sufficiently large so that $g^{n^2}(K')$ is contained in a petal \mathcal{P} from Proposition 4.1. Unless otherwise stated, all the constants appearing in estimates depend only on the compact $K \times K'$, but not on the point (z_0, w_0) nor the integer n .

Let $f_j(z) := f(z) + \frac{\pi^2}{4}w_{n^2+j}$, where $w_{n^2+j} := g^{n^2+j}(w_0)$. Let $z_j := f_j \circ f_{j-1} \circ \dots \circ f_1(z_0)$. Let $F_{m,p} := f_m \circ \dots \circ f_{p+1}$, and let $\epsilon_j := \sqrt{w_{n^2+j}}$.

The strategy of the proof of Theorem 5.33 is as follows: we will use *approximate Fatou coordinates* $\phi_n^{i/o}$ and prove that on some appropriate domains $\phi_n^{i/o}$ converges locally uniformly to $\phi_f^{i/o}$ (with a known error term of order $1/n$). Moreover, we will compute $\phi_n^i(z_0)$ and $\phi_n^o(z_{2n+1})$, again at a precision of order $1/n$. This will allow us to compare accurately z_{2n+1} and $\mathcal{L}_f(z_0) = (\phi_f^o)^{-1} \circ \phi_f^i(z_0)$. This approach differs from [Astorg et al. 2016] in that approximate Fatou coordinates in that work were only used at small scale near 0, while here they are defined on a whole petal: this simplifies the comparison with the actual Fatou coordinates $\phi_f^{i/o}$. The approach used here is strongly inspired by [Bedford et al. 2017].

Definition 5.11. Let

$$Z_j^{i/o} := \psi_{\epsilon_j}^{i/o}(z_j) = \frac{1}{i\pi} \log \frac{\zeta^+(\epsilon_j) - z_j}{z_j - \zeta^-(\epsilon_j)} \pm 1.$$

Observe that by definition of $A(\epsilon, z)$,

$$A(\epsilon_j, z_j) = Z_{j+1}^i - Z_j^i - \epsilon_j. \quad (14)$$

Proposition 5.12. *We have*

$$\psi_{\epsilon_j}^{i/o}(z_0) = -\frac{\epsilon_j}{z_0} + O\left(\frac{1}{n^3}\right).$$

Proof. This follows from computations similar to those appearing in the proof of Proposition 5.5 (recall as well that $\epsilon_j = O(1/n)$). \square

We now introduce approximate incoming Fatou coordinates:

Definition 5.13. Let

$$\phi_n^i(z_0) := \frac{1}{\epsilon_n} Z_n - \frac{1}{\epsilon_n} \sum_{j=1}^{n-1} \epsilon_j.$$

Let D_ϵ be the disk of radius $\frac{1}{2}|\zeta^+(\epsilon) - \zeta^-(\epsilon)|$ centered at $\frac{1}{2}(\zeta^+(\epsilon) + \zeta^-(\epsilon))$. Let $S(\epsilon, r)$ be the union of the two disks of radius r that both contain the points $\zeta^+(\epsilon), \zeta^-(\epsilon)$ on their boundary. Here r will be a sufficiently small number, to be fixed in the paragraph before Lemma 5.14. The definition of $S(\epsilon, r)$ of course only makes sense when the distance between $\zeta^+(\epsilon)$ and $\zeta^-(\epsilon)$ is less than $2r$, which once r is fixed will be satisfied for ϵ sufficiently small. We note that the choice of r will depend on the map f , but not on ϵ .




Figure 1. The sets $D_\epsilon \subset S(\epsilon, r)$ and their images under ψ_ϵ^t .

The line L_ϵ through $\zeta^+(\epsilon)$ and $\zeta^-(\epsilon)$ cuts the complex plane into the left half-plane H_ϵ^l and the right half-plane H_ϵ^o . We define $S^{t/o}(\epsilon, r) := S(\epsilon, r) \cap H_\epsilon^{t/o}$. The map ψ_ϵ^t maps the disk D_ϵ to the strip $[\frac{1}{2}, \frac{3}{2}] \times i\mathbb{R}$. The image of $S^t(\epsilon, r)$ is bounded by two vertical lines, intersecting the real line in a point of the form $0 + O(\epsilon)$ and in the point 1; see Figure 1. In particular we can find $0 < \alpha < \beta$ such that

$$[\beta\epsilon, 1] \times i\mathbb{R} \subset \psi_\epsilon^t(S^t(\epsilon, r)) \subset [\alpha\epsilon, 1] \times i\mathbb{R}$$

for all ϵ . We define $S^t(0, r) := \mathbb{D}(-r, r)$.

Recall that $A(\epsilon, \cdot)$ is analytic on a small disk $\mathbb{D}(0, R)$ centered at the origin. Moreover there exists $R > 0$ such that $|A(\epsilon, z)| \leq \frac{1}{12}|\epsilon|$ for all $z \in \mathbb{D}(0, R)$ and ϵ in the petal \mathcal{P}_δ defined in Section 4. By taking smaller R if necessary we may assume that f is 1-Lipschitz on $\mathbb{D}(-R, R)$.

Now let us assume that $r \ll R$ is sufficiently small so that $S(\epsilon, r) \subset \mathbb{D}(0, R)$ for all $\epsilon > 0$, and note that for every compact set $K \subset \mathcal{B}_f$ there exist $n', \epsilon' > 0$ so that $f^n(K) \subset S^t(\epsilon, r) \cap \mathbb{D}(-R, R)$ for all $n \geq n'$ and all $\epsilon \geq \epsilon'$. We now fix this r .

Lemma 5.14. *Let $K \times K' \subset \mathcal{B}_f \times \mathcal{B}_g$ be a compact set. There exist $n_0, m_0 > 0$ such that for all $(z_0, w_0) \in K \times K'$ and all $n > n_0$ we have*

- (1) $z_j \in S^t(\epsilon_j, r) \cup D_{\epsilon_j}$ for all $m_0 \leq j \leq n-1$,
- (2) $z_j \in D_{\epsilon_j}$ for all $\frac{2}{3}n \leq j \leq n-1$,
- (3) If $z_k \in S(\epsilon_k, r)$ for all $m_0 \leq k \leq j$, then $|\text{Im}(\psi_{\epsilon_{j+1}}^t(z_{j+1}))| < 1$,

where $\epsilon_j := \sqrt{w_{n^2+j}} = 1/n + O(j/n^3)$.

Proof. There exists $m_0 > 0$ so that $f^{m_0}(K) \subset S^t(0, r)$. Let n_0 be sufficiently large so that for all $(z_0, w_0) \in K \times K'$ we have:

- (i) $\beta|\epsilon_{m_0}| < \text{Re}(\psi_{\epsilon_{m_0}}^t(z_{m_0})) < \frac{1}{6}$.
- (ii) $|\text{Im}(\psi_{\epsilon_{m_0}}^t(z_{m_0}))| < \frac{1}{2}$.
- (iii) $|\epsilon_j| < 2\text{Re}(\epsilon_j) < R$ for $0 \leq j \leq 2n+1$.

Indeed, (i) and (ii) follow from the equality $\psi_\epsilon^t(z) = -\epsilon/z + O(\epsilon^3)$ and (iii) follows from the fact that $\epsilon_j = 1/n + O(j/n^3)$. Note that the constants in O depend only on the compact $K \times K'$ and not on n or j .

Recall that by our assumption $\mathcal{S}^t(\epsilon_j, r) \subset \mathbb{D}(0, R)$ for all j and that $|A(\epsilon_j, z)| \leq \frac{1}{12}|\epsilon_j|$ for all $z \in \mathbb{D}(0, R)$. By (i) we have $z_{m_0} \in \mathcal{S}^t(\epsilon_{m_0}, r)$ and observe that for $z_j \in \mathcal{S}^t(\epsilon_j, r)$ we have

$$\frac{5}{6} \operatorname{Re}(\epsilon_j) < \operatorname{Re}(\psi_{\epsilon_{j+1}}^t(z_{j+1}) - \psi_{\epsilon_j}^t(z_j)) < \frac{7}{6} \operatorname{Re}(\epsilon_j).$$

It follows that

$$\beta|\epsilon_{m_0}| + \frac{5}{6} \sum_{k=m_0}^{j-1} \operatorname{Re}(\epsilon_k) < \operatorname{Re}(\psi_{\epsilon_j}^t(z_j)) < \frac{1}{6} + \frac{7}{6} \sum_{k=m_0}^{j-1} \operatorname{Re}(\epsilon_k), \quad (15)$$

and since $\operatorname{Re}(\epsilon_k) = 1/n + O(k/n^3)$ we have

$$\beta|\epsilon_{m_0}| < \operatorname{Re}(\psi_{\epsilon_j}^t(z_j)) < \frac{3}{2}$$

for all $m_0 \leq j \leq n-1$ as long as n is sufficiently large. This proves (1).

For (2) observe that

$$-\frac{1}{2} < -1 + \frac{5}{6} \sum_{k=0}^{j-1} \operatorname{Re}(\epsilon_k)$$

for all $\frac{2}{3}n \leq j \leq n-1$ as long as n is sufficiently large.

Finally for (3) observe that (15) implies that $z_k \in \mathcal{S}(\epsilon_k, r)$ for all $0 \leq k \leq j$ can only hold for some $j < 3n$. By (ii) and (iii) we have

$$\operatorname{Im}(\epsilon_j) - \frac{1}{6} \operatorname{Re}(\epsilon_j) < \operatorname{Im}(\psi_{\epsilon_{j+1}}^t(z_{j+1}) - \psi_{\epsilon_j}^t(z_j)) < \operatorname{Im}(\epsilon_j) + \frac{1}{6} \operatorname{Re}(\epsilon_j)$$

for $z_j \in \mathcal{S}^t(\epsilon_j, r)$; hence

$$-\frac{1}{2} + \sum_{k=m_0}^{j-1} \operatorname{Im}(\epsilon_k) - \frac{1}{6} \operatorname{Re}(\epsilon_k) < \operatorname{Im}(\psi_{\epsilon_j}^t(z_j)) < \frac{1}{2} + \sum_{k=m_0}^{j-1} \operatorname{Im}(\epsilon_k) + \frac{1}{6} \operatorname{Re}(\epsilon_k).$$

Since $\operatorname{Im}(\epsilon_k) = O(k/n^3)$ we can conclude that (3) holds as long as n is sufficiently large. \square

Lemma 5.15. *For $0 \leq j \leq n-1$ we have*

$$z_j - f^j(z_0) = O\left(\frac{j}{n^2}\right).$$

Proof. Let m_0 be as in Lemma 5.14. Since m_0 is independent from n it is easy to see that for all $0 \leq j \leq m_0$ we have

$$z_j - f^j(z_0) = \frac{\pi^2}{4} \sum_{k=0}^{j-1} \left(\epsilon_k^2 + O\left(\frac{1}{n^3}\right) \right) = O\left(\frac{j}{n^2}\right).$$

Let $V_\epsilon := \{z \in \mathcal{S}^t(\epsilon, r) : |\operatorname{Im}(\psi_\epsilon^t(z))| < 1\}$ and observe that $V_\epsilon \setminus D_\epsilon \subset \mathbb{D}(-R, R)$ for all sufficiently small ϵ . Since by our assumption f is 1-Lipschitz on $\mathbb{D}(-R, R)$, it follows by (1) in Lemma 5.14 that

for all $m_0 < j < \frac{2}{3}n$ we have

$$\begin{aligned} |z_j - f^j(z_0)| &< |z_{j-1} - f^{j-1}(z_0)| + \frac{\pi^2}{4}\epsilon_{j-1}^2 \\ &< |z_{m_0} - f^{m_0}(z_0)| + \frac{\pi^2}{4} \sum_{k=m_0}^{j-1} \epsilon_k^2 = \frac{\pi^2}{4} \sum_{k=0}^{j-1} \left(\frac{1}{n^2} + O\left(\frac{1}{n^3}\right) \right) = O\left(\frac{j}{n^2}\right). \end{aligned}$$

Finally for $\frac{2}{3}n \leq j \leq n-1$, by item (2) of Lemma 5.14, we have $z_j \in D_{\epsilon_j}$, and in particular $z_j = O(1/n)$. It follows that

$$z_j - f^j(z_0) = O\left(\frac{1}{n}, \frac{1}{j}\right) = O\left(\frac{j}{n^2}\right),$$

where the last equality follows from the fact that $\frac{2}{3}n \leq j \leq n-1$. □

Lemma 5.16. *We have*

$$\sum_{j=0}^{n-1} A_0(z_j) - A_0(f^j(z_0)) = (b-1) \sum_{j=0}^{n-1} z_j^2 - f^j(z_0)^2 + O\left(\frac{\log n}{n^2}\right).$$

Proof. Recall that $f(z) = z + z^2 + z^3 + bz^4 + O(z^5)$ and $A_0(z) = -1/f(z) + 1/z - 1$. An elementary computation gives $A_0(0) = A'_0(0) = 0$ and $A''_0(0) = 2(b-1)$.

To simplify the notation, let $y_j := f^j(z_0)$. We have

$$\begin{aligned} A_0(z_j) - A_0(y_j) &= A'_0(y_j)(z_j - y_j) + \frac{1}{2}A''_0(y_j)(z_j - y_j)^2 + O((z_j - y_j)^3) \\ &= (y_j A''_0(0) + O(y_j^2))(z_j - y_j) + \frac{1}{2}(A''_0(0) + O(y_j))(z_j - y_j)^2 + O((z_j - y_j)^3). \end{aligned}$$

By Lemma 5.15 we have $z_j - y_j = O(j/n^2)$; hence

$$\begin{aligned} A_0(z_j) - A_0(y_j) &= y_j A''_0(0)(z_j - y_j) + \frac{1}{2}A''_0(0)(z_j - y_j)^2 + O\left(\frac{1}{jn^2}, \frac{j}{n^4}, \frac{j^3}{n^6}\right) \\ &= (b-1)(2y_j(z_j - y_j) + (z_j - y_j)^2) + O\left(\frac{1}{jn^2}, \frac{j}{n^4}, \frac{j^3}{n^6}\right) \\ &= (b-1)(z_j^2 - y_j^2) + O\left(\frac{1}{jn^2}, \frac{j}{n^4}, \frac{j^3}{n^6}\right). \end{aligned}$$

It follows that

$$\sum_{j=0}^{n-1} A_0(z_j) - A_0(y_j) = (b-1) \sum_{j=0}^{n-1} z_j^2 - y_j^2 + O\left(\frac{\log n}{n^2}\right). \quad \square$$

Lemma 5.17. *For $0 \leq j \leq n-1$, let*

$$\gamma_j := j + \sum_{k=0}^{j-1} A_0(f^k(z_0)) \quad \text{and} \quad x_j := \sum_{k=0}^{j-1} \epsilon_k + A(\epsilon_k, z_k). \quad (16)$$

Then

$$x_j = \frac{\gamma_j}{n} + O\left(\frac{j^2}{n^3}\right) = \frac{j}{n} + O\left(\frac{1}{n}\right).$$

In particular, there exists $k \in \mathbb{N}$ independent from n such that for all $k \leq j \leq n - k$

$$\alpha_j := \cot\left(\frac{\pi}{2}x_j\right)$$

is well-defined and strictly positive.

Proof. According to Lemma 5.15, for $0 \leq j \leq n - 1$ we have that $z_j - f^j(z_0) = O(j/n^2)$. In particular, $z_j = O(1)$. By Proposition 5.8, we have for every $0 \leq k \leq n - 1$

$$A(\epsilon_k, z_k) = \epsilon_k A_0(z_k) + O(\epsilon_k^3, z_k \epsilon_k^3) = \epsilon_k A_0(z_k) + O\left(\frac{1}{n^3}\right) \quad (17)$$

(indeed, by Lemma 5.15, $z_k = f^k(z_0) + O(k/n^2)$, so in particular $z_k = O(1)$). By Lemma 5.2, $\epsilon_k = 1/n + O(k/n^3)$; hence

$$\begin{aligned} x_j &= \sum_{k=0}^{j-1} \epsilon_k + A(\epsilon_k, z_k) = \sum_{k=0}^{j-1} \frac{1}{n} + \frac{1}{n} A_0(z_k) + O\left(\frac{k}{n^3}\right) \\ &= \frac{j}{n} + \frac{1}{n} \sum_{k=0}^{j-1} A_0(f^k(z_0)) + O\left(\frac{k}{n^2}, A'_0(f^k(z_0))(z_k - f^k(z_0))\right) \\ &= \frac{j}{n} + \frac{1}{n} \sum_{k=0}^{j-1} A_0(f^k(z_0)) + O\left(\frac{k}{n^2}, \frac{1}{k} \cdot \frac{k}{n^2}\right) = \frac{\gamma_j}{n} + O\left(\frac{j^2}{n^3}\right). \end{aligned}$$

Since $\gamma_j = j + O(1)$, we also have

$$\frac{\gamma_j}{n} = \frac{j}{n} + O\left(\frac{1}{n}\right).$$

Finally, the last assertion follows from the preceding equality and the fact that, for $x \in (0, \frac{\pi}{2})$, $\cot(x) > 0$. \square

Lemma 5.18. *Let*

$$u(x) := \frac{2}{\pi} \tan\left(\frac{\pi}{2}x\right), \quad \Phi(x) := \frac{x^2 - u(x)^2}{x^2 u(x)^2} \quad \text{and} \quad \beta_j := \frac{2n}{\pi \alpha_j}.$$

We have

$$\frac{\gamma_j^2 - \beta_j^2}{\gamma_j^2 \beta_j^2} = \frac{1}{n^2} \Phi(x_j) + O\left(\frac{1}{jn^2}\right).$$

Proof. We have

$$\frac{\gamma_j^2 - \beta_j^2}{\gamma_j^2 \beta_j^2} = \frac{n^2 (\gamma_j^2/n^2) - u(x_j)^2}{n^4 u(x_j)^2 (\gamma_j^2/n^2)}.$$

Now recall that by Lemma 5.17, $\gamma_j/n = x_j + O(j^2/n^3)$, so that $\gamma_j^2/n^2 = x_j^2 + O(j^3/n^4)$. So

$$\frac{\gamma_j^2 - \beta_j^2}{\gamma_j^2 \beta_j^2} = \frac{1}{n^2} \frac{x_j^2 - u(x_j)^2 + O(j^3/n^4)}{u(x_j^2)(x_j^2 + O(j^3/n^4))} = \frac{1}{n^2} \frac{x_j^2 - u(x_j)^2}{u(x_j^2)(x_j^2 + O(j^3/n^4))} + O\left(\frac{j^3}{x_j^2 u(x_j)^2 n^6}\right)$$

and note that

$$\frac{1}{x_j^2 u(x_j)^2} = O\left(\frac{1}{x_j^4}\right) = O\left(\frac{n^4}{j^4}\right).$$

Therefore

$$\begin{aligned} \frac{\gamma_j^2 - \beta_j^2}{\gamma_j^2 \beta_j^2} &= \frac{1}{n^2} \frac{x_j^2 - u(x_j)^2}{u(x_j^2)(x_j^2 + O(j^3/n^4))} + O\left(\frac{1}{jn^2}\right) \\ &= \frac{1}{n^2} \frac{x_j^2 - u(x_j)^2}{u(x_j^2)x_j^2(1 + O(j^3/(n^4x_j^2)))} + O\left(\frac{1}{jn^2}\right) \\ &= \frac{\Phi(x_j)}{n^2} \left(1 + O\left(\frac{j}{n^2}\right)\right) + O\left(\frac{1}{jn^2}\right) \\ &= \frac{\Phi(x_j)}{n^2} + O\left(\frac{1}{jn^2}\right). \end{aligned}$$

Note that in the last line, we used the fact that Φ has only removable singularities at $x = 0$ and $x = 1$, so that $\Phi(x_j) = O(1)$. \square

Proposition 5.19. *There exists a universal constant $C_1 \in \mathbb{R}$ such that*

$$\sum_{j=0}^{n-1} A_0(z_j) - A_0(f^j(z)) = \frac{C_1(b-1)}{n} + O\left(\frac{\log n}{n^2}\right).$$

More precisely, $C_1 := \int_0^1 \Phi(x) dx = \frac{1}{4}(4 - \pi^2)$.

Proof. We have, for $0 \leq j \leq n-1$,

$$z_j = \psi_{\epsilon_j}^{-1}(Z_j^t) = -\frac{\pi}{2n} \cot\left(\frac{\pi}{2} Z_j^t\right) + O\left(\frac{1}{n^2}\right)$$

and

$$Z_j^t = Z_0^t + \sum_{k=0}^{j-1} \epsilon_k + A(\epsilon_k, z_k).$$

Recalling the notation $x_j := \sum_{k=0}^{j-1} \epsilon_k + A(\epsilon_k, z_k)$ and $\alpha_j := \cot((\pi/2)x_j)$ from Lemma 5.17, and using the trigonometric formula

$$\cot(a+b) = \frac{\cot a \cot b - 1}{\cot a + \cot b},$$

we therefore obtain

$$z_j = -\frac{\pi}{2n} \frac{\cot((\pi/2)Z_0^t)\alpha_j - 1}{\alpha_j + \cot((\pi/2)Z_0^t)} + O\left(\frac{1}{n^2}\right). \quad (18)$$

Let k be as in Lemma 5.17, so that $\alpha_j > 0$ for $k \leq j \leq n-k$. We have

$$\cot\left(\frac{\pi}{2} Z_0^t\right) = -\frac{2n}{\pi} z_0 + O\left(\frac{1}{n}\right),$$

so that

$$z_j = -\frac{\pi}{2n} \frac{\cot((\pi/2)Z'_0)\alpha_j}{\alpha_j + \cot((\pi/2)Z'_0)} + O\left(\frac{1}{n^2}\right) = \frac{z_0\alpha_j}{\alpha_j - z_0(2n/\pi)} + O\left(\frac{1}{n^2}\right). \quad (19)$$

Finally, with $\beta_j := 2n/(\pi\alpha_j)$, we get

$$z_j = -\frac{1}{-1/z_0 + \beta_j} + O\left(\frac{1}{n^2}\right). \quad (20)$$

On the other hand, from the definition of A_0 it follows that $\sum_{k=0}^{j-1} A_0(f^k(z_0)) = 1/z_0 - 1/f^j(z_0) - j$, which we may rewrite as

$$f^j(z_0) = -\frac{1}{-1/z_0 + \gamma_j}. \quad (21)$$

Therefore

$$z_j - f^j(z_0) = \frac{\beta_j - \gamma_j}{(-1/z_0 + \beta_j)(-1/z_0 + \gamma_j)} + O\left(\frac{1}{n^2}\right). \quad (22)$$

Now note that $j = O(\beta_j)$; indeed,

$$\frac{1}{\beta_j} = O\left(\frac{\cot(j\pi/n + O(1/n))}{n}\right) = O\left(\frac{n/j}{n}\right) = O\left(\frac{1}{j}\right).$$

Therefore

$$\begin{aligned} \frac{1}{(-1/z_0 + \gamma_j)(-1/z_0 + \beta_j)} &= \frac{1}{(\gamma_j + O(1))(\beta_j + O(1))} \\ &= \frac{1}{\gamma_j\beta_j + O(\beta_j)} = \frac{1}{\gamma_j\beta_j} + O\left(\frac{1}{\gamma_j^2\beta_j}\right). \end{aligned}$$

Thus, setting $y_j := f^j(z_0)$,

$$\begin{aligned} z_j^2 - y_j^2 &= (z_j - y_j)(z_j + y_j) \\ &= \left(\frac{\beta_j - \gamma_j}{\beta_j\gamma_j} + O\left(\frac{\beta_j - \gamma_j}{\gamma_j^2\beta_j}\right)\right) \left(-\frac{\beta_j + \gamma_j}{\beta_j\gamma_j} + O\left(\frac{\beta_j + \gamma_j}{\gamma_j^2\beta_j}\right)\right) \\ &= \frac{\gamma_j^2 - \beta_j^2}{\beta_j^2\gamma_j^2} + O\left(\frac{\beta_j^2 - \gamma_j^2}{\beta_j^2\gamma_j^3}\right) \\ &= \frac{1}{n^2} \Phi(x_j) + O\left(\frac{1}{jn^2}\right) \quad \text{by Lemma 5.18.} \end{aligned}$$

Therefore by Lemma 5.16,

$$\begin{aligned} \sum_{j=0}^{n-1} A_0(z_j) - A_0(y_j) &= \left(\frac{b-1}{n^2} \sum_{j=0}^{n-1} \Phi(x_j)\right) + O\left(\frac{\log n}{n^2}\right) \\ &= \frac{b-1}{n} \int_0^1 \Phi(x) dx + O\left(\frac{\log n}{n^2}\right). \end{aligned}$$

In the last equality, we recognize a Riemann sum with subdivision $(x_j)_{0 \leq j \leq n-1}$. Finally, we have

$$\int_0^1 \Phi(x) dx = \frac{\pi}{2} \int_0^{\frac{\pi}{2}} \cot^2 t - \frac{1}{t^2} dt = -\frac{\pi}{2} \left[\left(\cot t - \frac{1}{t} \right) + t \right]_0^{\frac{\pi}{2}} = 1 - \frac{\pi^2}{4}. \quad \square$$

5B1. Incoming part. The following error estimate is one of the two crucial estimates that we will obtain in this section: it measures accurately how close ϕ_n^t is to the incoming Fatou coordinate ϕ_f^t . This estimate differs from those obtained in [Astorg et al. 2016] in that we compare ϕ_n^t with ϕ_f^t on a definite region of \mathcal{B}_f (independent from n), instead of comparing the two at small scale near the origin, compare with [Astorg et al. 2016, Property 1, p. 10]. Moreover, the point of Proposition 5.20 is to push the precision of the estimate further and obtain the first error term $E^t(z_0)/n$, which cannot be easily obtained from the computations in [Astorg et al. 2016].

Proposition 5.20. *We have*

$$\phi_n^t(z_0) = \phi_f^t(z_0) + \frac{E^t(z_0)}{n} + O\left(\frac{\log n}{n^2}\right),$$

where $E^t(z) := C_0 + (C_1 - 1)(b - 1) + \frac{1}{2}\phi^t(z_0)$.

Proof. Recall that by definition,

$$\phi_f^t(z) = \lim_{n \rightarrow \infty} -\frac{1}{f^n(z)} - n = \lim_{n \rightarrow \infty} -\frac{1}{z} + \sum_{j=0}^{n-1} A_0(f^j(z)).$$

Similarly, we have

$$\sum_{j=0}^{n-1} A(\epsilon_j, z_j) = \sum_{j=0}^{n-1} Z_{j+1}^t - Z_j^t - \epsilon_j = Z_n^t - Z_0^t - \sum_{j=0}^{n-1} \epsilon_j,$$

and thus

$$\phi_n^t(z_0) = \frac{Z_n^t}{\epsilon_n} - \frac{1}{\epsilon_n} \sum_{j=0}^{n-1} \epsilon_j = \frac{Z_0^t}{\epsilon_n} + \frac{1}{\epsilon_n} \sum_{j=0}^{n-1} A(\epsilon_j, z_j).$$

Therefore

$$\phi_n^t(z_0) - \phi_f^t(z_0) = E_1 + E_2 + E_3, \quad (23)$$

where

$$\begin{aligned} E_1 &:= \frac{Z_0^t}{\epsilon_n} + \frac{1}{z_0}, \\ E_2 &:= \frac{1}{\epsilon_n} \sum_{j=0}^{n-1} A(\epsilon_j, z_j) - \sum_{j=0}^{n-1} A_0(f^j(z_0)), \\ E_3 &:= - \sum_{j=n}^{\infty} A_0(f^j(z_0)). \end{aligned}$$

We will now estimate each of the error terms E_i separately. For $j \in \mathbb{N}$, we set $y_j := f^j(z_0)$.

Lemma 5.21. *We have*

$$E_1 = -\frac{1}{2nz_0} + O\left(\frac{1}{n^2}\right).$$

Proof of lemma. We have

$$\begin{aligned} \frac{Z_0^l}{\epsilon_n} &= \frac{1}{\epsilon_n} \psi_{\epsilon_0}(z_0) = -\frac{\epsilon_0}{\epsilon_n z_0} + O\left(\frac{\epsilon_0^3}{\epsilon_n}\right) \quad (\text{by Proposition 5.12}) \\ &= -\frac{1}{z_0} \sqrt{\frac{n^2 + n + O(1)}{n^2 + O(1)}} + O\left(\frac{1}{n^2}\right) \\ &= -\frac{1}{z_0} - \frac{1}{2nz_0} + O\left(\frac{1}{n^2}\right). \end{aligned} \quad \square$$

Lemma 5.22. *We have*

$$E_2 = \frac{1}{n} \left(\frac{1}{2z_0} + \frac{1}{2} \phi_f^l(z_0) + C_0 + C_1(b-1) \right) + O\left(\frac{\log n}{n^2}\right).$$

Proof of lemma. Recall that we have

$$A(\epsilon, z) = \epsilon A_0(z) + C_0 \epsilon^3 + O(z\epsilon^3, \epsilon^4),$$

so that

$$\begin{aligned} E_2 &= \frac{1}{\epsilon_n} \sum_{j=0}^{n-1} A(\epsilon_j, z_j) - \sum_{j=0}^{n-1} A_0(y_j) \\ &= \frac{1}{\epsilon_n} \sum_{j=0}^{n-1} \epsilon_j A_0(z_j) + C_0 \epsilon_j^3 + O\left(\frac{z_j}{n^3}\right) - \epsilon_n A_0(y_j). \end{aligned}$$

Therefore

$$E_2 = \left(\sum_{j=0}^{n-1} \frac{\epsilon_j}{\epsilon_n} A_0(z_j) - A_0(y_j) \right) + \left(\sum_{j=0}^{n-1} C_0 \frac{\epsilon_j^3}{\epsilon_n} + O\left(\frac{z_j}{n^2}\right) \right)$$

and

$$\sum_{j=0}^{n-1} C_0 \frac{\epsilon_j^3}{\epsilon_n} + O\left(\frac{z_j}{n^2}\right) = C_0 \sum_{j=0}^{n-1} \frac{1}{n^2} + O\left(\frac{1}{n^2 j}\right) = \frac{C_0}{n} + O\left(\frac{\log n}{n^2}\right).$$

On the other hand, we have

$$\sum_{j=0}^{n-1} \frac{\epsilon_j}{\epsilon_n} A_0(z_j) - A_0(y_j) = \sum_{j=0}^{n-1} \left(\frac{\epsilon_j}{\epsilon_n} - 1 \right) A_0(z_j) + \sum_{j=0}^{n-1} A_0(z_j) - A_0(y_j). \quad (24)$$

Now note that

$$\sum_{j=0}^{n-1} \left(\frac{\epsilon_j}{\epsilon_n} - 1 \right) A_0(z_j) = \sum_{j=0}^{n-1} \left(\frac{\epsilon_j}{\epsilon_n} - 1 \right) A_0(y_j) + \sum_{j=0}^{n-1} \left(\frac{\epsilon_j}{\epsilon_n} - 1 \right) (A_0(z_j) - A_0(y_j)),$$

and that

$$\left| \sum_{j=0}^{n-1} \left(\frac{\epsilon_j}{\epsilon_n} - 1 \right) (A_0(z_j) - A_0(y_j)) \right| \leq \max_{0 \leq j \leq n-1} \left| 1 - \frac{\epsilon_j}{\epsilon_n} \right| \cdot \sum_{j=0}^{n-1} |A_0(z_j) - A_0(y_j)| = O\left(\frac{1}{n^2}\right),$$

by Lemma 5.2 and Proposition 5.19. Another consequence of Lemma 5.2 is that

$$\frac{\epsilon_j}{\epsilon_n} - 1 = \frac{1}{2n} \left(1 - \frac{j}{n} + O\left(\frac{1}{n}\right) \right). \quad (25)$$

Therefore, by (24), (25) and Proposition 5.19,

$$\begin{aligned} \sum_{j=0}^{n-1} \frac{\epsilon_j}{\epsilon_n} A_0(z_j) - A_0(y_j) &= \frac{C_1(b-1)}{n} + \frac{1}{2n} \sum_{j=0}^{n-1} A_0(y_j) + O\left(\frac{\log n}{n^2}\right) \\ &= \frac{C_1(b-1)}{n} + \left(\frac{1/z_0 + \phi_f^t(z_0)}{2n} \right) + O\left(\frac{\log n}{n^2}\right). \end{aligned}$$

Therefore, as announced, we have

$$E_2 = \frac{1}{n} \left(\frac{1}{2z_0} + \frac{1}{2} \phi_f^t(z_0) + C_0 + C_1(b-1) \right) + O\left(\frac{\log n}{n^2}\right). \quad \square$$

Lemma 5.23. *We have*

$$E_3 = \frac{1-b}{n} + O\left(\frac{1}{n^2}\right).$$

Proof. By explicit computations, $A_0(z) = (b-1)z^2 + O(z^3)$, so that $A_0(y_j) = (b-1)j^{-2} + O(j^{-3})$. Therefore

$$E_3 = (1-b) \sum_{j=n}^{\infty} j^{-2} + O(j^{-3})$$

and

$$\sum_{j=n}^{\infty} j^{-3} = O\left(\int_n^{\infty} \frac{dx}{x^3}\right) = O\left(\frac{1}{n^2}\right).$$

Similarly,

$$\sum_{j=n}^{\infty} j^{-2} \sim \int_n^{\infty} \frac{dx}{x^2} = \frac{1}{n},$$

so that $E_3 = (1-b)/n + O(1/n^2)$. □

Finally, putting together the three preceding lemmas, the proof of Proposition 5.20 is finished. □

5B2. Outgoing part. We will now work to obtain estimates for the outgoing part of the orbit, that is, for $n \leq j \leq 2n+1$. The method is largely similar to the incoming case. Recall that the estimates we obtain only depend on the chosen compact set $K \subset \mathcal{B}_f$.

We will first need a rough preliminary estimate on the boundedness of z_{2n+1} . Of course, by [Astorg et al. 2016], we know that z_{2n+1} converges to $\mathcal{L}(z_0)$, and we could deduce this preliminary estimate from there. However, we prefer to present here a direct argument, so that the proof of Theorem 5.33 remains self-contained.

Proposition 5.24. *There exists $k \in \mathbb{N}$ (independent from n) such that z_{2n+1-k} belongs to a repelling petal $\mathbb{D}(r, r)$ for f . In particular, $z_{2n+1} = O(1)$.*

Proof. Recall that by Proposition 5.20, we have that

$$\phi_n^l(z) := \frac{Z_n^l}{\epsilon_n} - \frac{1}{\epsilon_n} \sum_{j=0}^{n-1} \epsilon_j = \phi^l(z_0) + o(1) = O(1).$$

In particular,

$$Z_n^l = \left(\sum_{j=0}^{n-1} \epsilon_j \right) + O(\epsilon_n) = 1 + O\left(\frac{1}{n}\right)$$

and therefore $Z_n^o = -1 + O(1/n)$.

Let R_n denote the rectangle defined by the conditions $-1 - C/n \leq \operatorname{Re}(Z) \leq -3/n$ and $-1 \leq \operatorname{Im}(Z) \leq 1$, where $C > 0$ is a constant chosen large enough that $Z_n^o \in R_n$. Let

$$j_n := \max\{k \leq 2n + 1 : Z_k^o \in R_n\}. \quad (26)$$

Recall that for $j \leq 2n$, we have $Z_{j+1}^o = Z_j^o + A(\epsilon_j, z_j)$, and that by Proposition 5.8, we have

$$A(\epsilon_k, z_k) = \epsilon_k A_0(z_k) + O(\epsilon_k^3, \epsilon_k^3 z_k) = O(\epsilon_k z_k^2). \quad (27)$$

Moreover, for $n \leq j \leq j_n$, we have

$$Z_j^o = -\frac{\pi}{2n} \cot\left(\frac{\pi}{2} Z_j^o\right) + O\left(\frac{1}{n^2}\right),$$

and therefore there exists a constant $C > 0$ such that, for all $n \leq j \leq j_n$,

$$|A(\epsilon_j, z_j)| \leq \frac{C'}{n^3} \left| \cot\left(\frac{\pi}{2} Z_j^o\right) \right|^2 \leq \frac{C}{|Z_j|^2 n^3}, \quad (28)$$

and thus

$$\left| Z_j^o - Z_n^o - \sum_{k=n}^{j-1} \epsilon_k \right| \leq \frac{C}{n^3} \sum_{k=n}^{j-1} \frac{1}{|Z_k|^2}. \quad (29)$$

From (29), we can prove inductively on j that, for $n \leq j \leq j_n$,

$$\left| Z_j^o - Z_n^o - \sum_{k=n}^{j-1} \epsilon_k \right| = O\left(\frac{1}{n}\right)$$

and hence $j_n = 2n + O(1)$.

Let $r > 0$ be small enough such that $\mathbb{D}(r, r)$ is a repelling petal for f . By the argument above and the definition of R_n , we have that $Z_{j_n}^o = O(1/n)$, so that

$$z_{2n+1-k} = -\frac{\pi}{2} \cot\left(\frac{\pi}{2} Z_{2n+1-k}^o\right) + O\left(\frac{1}{n^2}\right) = \frac{1}{k + O(1)}.$$

Therefore, we can find some k bounded independently from n such that $z_{2n+1-k} \in \mathbb{D}(r, r)$. \square

We now introduce approximate outgoing Fatou coordinates:

Definition 5.25. For $n \leq m \leq 2n + 1$, let

$$\phi_n^o(z_m) := \frac{Z_n^o}{\epsilon_n} + \frac{1}{\epsilon_n} \sum_{j=n}^{m-1} \epsilon_j.$$

Lemma 5.26. *We have*

$$\phi_n^o(z_m) = \frac{Z_m^o}{\epsilon_n} - \frac{1}{\epsilon_n} \sum_{j=n}^{m-1} A(\epsilon_j, z_j).$$

Proof. We have

$$\sum_{j=n}^{m-1} A(\epsilon_j, z_j) = \sum_{j=n}^{m-1} Z_{j+1}^o - Z_j^o - \epsilon_j = Z_m^o - Z_n^o - \sum_{j=n}^{m-1} \epsilon_j$$

so that

$$\frac{Z_n^o}{\epsilon_n} + \frac{1}{\epsilon_n} \sum_{j=n}^{m-1} \epsilon_j = \phi_n^o(z_m) = \frac{Z_m^o}{\epsilon_n} - \frac{1}{\epsilon_n} \sum_{j=n}^{m-1} A(\epsilon_j, z_j). \quad \square$$

Proposition 5.27. *Let $k \in \mathbb{N}$ be the integer from Proposition 5.24. Let $y_{2n+1-k} := z_{2n+1-k}$ and $y_{2n+1} = f^k(y_{2n+1-k})$. For $n \leq j \leq 2n$ we define*

$$y_j := f^{-(2n+1-j)}(y_{2n+1}),$$

where f^{-1} is the local inverse of f fixing 0: $f^{-1}(z) = z - z^2 + z^3 - bz^4 + O(z^5)$. We have

$$\sum_{j=n}^{2n} A_0(z_j) - A_0(y_j) = \frac{C_1(b-1)}{n} + O\left(\frac{\log n}{n^2}\right).$$

Proof. The proof mirrors the incoming case, so we will only sketch it and leave the details to the reader. Recall that $y_{2n+1} = O(1)$ by Proposition 5.24 and that z_{2n+1-k} belongs to a repelling petal for f for some $k \in \mathbb{N}$ independent from n , so that the $(y_j)_{n \leq j \leq 2n+1}$ are well-defined.

By a straightforward adaptation of Lemma 5.15,

$$z_j - y_j = O\left(\frac{2n+1-j}{n^2}\right)$$

for $n \leq j \leq 2n+1$. More precisely, this applies for $n \leq j \leq 2n+1-k$; but it is clear from the definition of the y_j that for $2n+1-k \leq j \leq 2n+1$ we have $z_j - y_j = O(1/n^2)$. Therefore the proof of Lemma 5.16 can be repeated to yield that

$$\sum_{j=n}^{2n} A_0(z_j) - A_0(y_j) = (b-1) \sum_{j=n}^{2n} z_j^2 - y_j^2 + O\left(\frac{\log n}{n^2}\right). \quad (30)$$

Next, we have, for $n \leq j \leq 2n$,

$$\begin{aligned} z_j &= (\psi_{\epsilon_j}^o)^{-1}(Z_j^o) = (\psi_{\epsilon_j}^o)^{-1}\left(Z_{2n+1}^o - \sum_{k=j}^{2n} \epsilon_k + A(\epsilon_k, z_k)\right) \\ &= -\frac{\pi}{2n} \cot\left(\frac{\pi}{2} Z_{2n+1}^o - \frac{\pi}{2} \sum_{k=j}^{2n} \epsilon_k + A(\epsilon_k, z_k)\right) + O\left(\frac{1}{n^2}\right). \end{aligned}$$

Through computations similar to those appearing in the proof of Proposition 5.19, we deduce that

$$z_j = -\frac{1}{-1/z_{2n+1} - \beta_j} + O\left(\frac{1}{n^2}\right), \quad (31)$$

with

$$\beta_j := \frac{2n}{\pi} \tan\left(\frac{\pi}{2} x_j\right) = \frac{2n}{\pi} \tan\left(\frac{\pi}{2} \sum_{k=j}^{2n} \epsilon_k + A(\epsilon_k, z_k)\right).$$

On the other hand,

$$-\frac{1}{y_j} = -\frac{1}{y_{2n+1}} - \sum_{k=j}^{2n} A_0(y_j),$$

from which it follows that

$$y_j = -\frac{1}{-1/y_{2n+1} - \gamma_j},$$

with $\gamma_j := \sum_{k=j}^{2n} A_0(y_j)$. Then, again, similar computations show that

$$z_j^2 - y_j^2 = \frac{1}{n^2} \Phi(x_j) + O\left(\frac{1}{n^2(2n+1-j)}\right),$$

and $x_j = (2n - j + O(1))/n$ for $n \leq j \leq 2n$. Therefore, we finally obtain

$$\sum_{j=n}^{2n} A_0(z_j) - A_0(y_j) = \frac{b-1}{n} \int_0^1 \Phi(x) dx + O\left(\frac{\log n}{n^2}\right) = \frac{C_1(b-1)}{n} + O\left(\frac{\log n}{n^2}\right). \quad \square$$

In what follows, a slight technical complication comes from the fact that the expected endpoint of the orbit, z_{2n+1} , needs not lie in a small enough repelling petal in which ϕ_f^o is well-defined. In order to overcome this issue, we stop a few iterations short and work instead with z_{2n+1-k} .

We now come to the main proposition of this subsection:

Proposition 5.28. *We have*

$$\phi_n^o(z_{2n+1-k}) = \phi_f^o(z_{2n+1-k}) + \frac{E^o(z_{2n+1-k})}{n} + O\left(\frac{\log n}{n^2}\right),$$

where $E^o(z) = -\frac{1}{2}\phi_f^o(z) - C_0 - (C_1 - 1)(b - 1)$.

Proof. We proceed similarly to the proof of Proposition 5.20. We have, for z in a small enough repelling petal,

$$\phi_f^o(z) = -\frac{1}{z} - \sum_{j=1}^{\infty} A_0(f^{-j}(z)), \quad (32)$$

where f^{-1} is the inverse branch of f fixing 0. With the same notation as in Proposition 5.27, we set

$$y_j := f^{j-(2n+1-k)}(z_{2n+1-k}).$$

We have

$$\begin{aligned} \phi_n^o(z_{2n+1-k}) - \phi_f^o(z_{2n+1-k}) &= \frac{Z_{2n+1-k}^o}{\epsilon_n} + \frac{1}{z_{2n+1-k}} + \sum_{j=n}^{2n-k} -\frac{1}{\epsilon_n} A(\epsilon_j, z_j) + A_0(y_j) + \sum_{j=-\infty}^{n-1} A_0(y_j) \\ &= E_1 + E_2 + E_3, \end{aligned} \quad (33)$$

where

$$\begin{aligned} E_1 &= \frac{Z_{2n+1-k}^o}{\epsilon_n} + \frac{1}{z_{2n+1-k}}, \\ E_2 &= \sum_{j=n}^{2n-k} -\frac{1}{\epsilon_n} A(\epsilon_j, z_j) + A_0(y_j), \\ E_3 &= \sum_{j=-\infty}^{n-1} A_0(y_j). \end{aligned} \quad (34)$$

Lemma 5.29. *We have*

$$E_1 = \frac{1}{n} \frac{1}{2z_{2n+1-k}} + O\left(\frac{1}{n^2}\right).$$

Proof of the lemma. By Proposition 5.12, we have

$$Z_{2n+1-k}^o = -\frac{\epsilon_{2n+1-k}}{z_{2n+1-k}} + O\left(\frac{1}{n^3}\right)$$

so that

$$\begin{aligned} E_1 &= \frac{1}{z_{2n+1-k}} - \frac{\epsilon_{2n+1-k}}{\epsilon_n} \frac{1}{z_{2n+1-k}} + O\left(\frac{1}{n^2}\right) \\ &= \frac{1}{z_{2n+1-k}} \left(1 - \sqrt{\frac{n^2 + n + O(1)}{n^2 + 2n + O(1)}}\right) + O\left(\frac{1}{n^2}\right) \\ &= \frac{1}{n} \frac{1}{2z_{2n+1-k}} + O\left(\frac{1}{n^2}\right). \end{aligned} \quad \square$$

Lemma 5.30. *We have*

$$E_2 = \frac{1}{n} \left(-\frac{1}{2z_{2n+1-k}} - \frac{1}{2} \phi_f^o(z_{2n+1-k}) - C_0 - C_1(b-1) \right) + O\left(\frac{\log n}{n^2}\right).$$

Proof of the lemma. We have

$$\begin{aligned} E_2 &= \sum_{j=n}^{2n-k} A_0(y_j) - \frac{1}{\epsilon_n} A(\epsilon_j, z_j) \\ &= \left(\sum_{j=n}^{2n-k} A_0(y_j) - \frac{\epsilon_j}{\epsilon_n} A_0(z_j) \right) - \left(\sum_{j=n}^{2n-k} C_0 \epsilon_j^3 + O(z_j \epsilon_j^3) \right). \end{aligned}$$

As before, we have

$$\frac{1}{\epsilon_n} \sum_{j=n}^{2n-k} C_0 \epsilon_j^3 + O(z_j \epsilon_j^3) = \frac{C_0}{n} + O\left(\frac{1}{n^2}\right).$$

On the other hand, we have

$$\sum_{j=n}^{2n-k} A_0(y_j) - \frac{\epsilon_j}{\epsilon_n} A_0(z_j) = \sum_{j=n}^{2n-k} \left(1 - \frac{\epsilon_j}{\epsilon_n}\right) A_0(z_j) + \sum_{j=n}^{2n-k} A_0(y_j) - A_0(z_j).$$

Now note that

$$\sum_{j=n}^{2n-k} \left(1 - \frac{\epsilon_j}{\epsilon_n}\right) A_0(z_j) = \sum_{j=n}^{2n-k} \left(1 - \frac{\epsilon_j}{\epsilon_n}\right) A_0(y_j) + \sum_{j=n}^{2n-k} \left(1 - \frac{\epsilon_j}{\epsilon_n}\right) (A_0(z_j) - A_0(y_j)),$$

and that

$$\left| \sum_{j=n}^{2n-k} \left(1 - \frac{\epsilon_j}{\epsilon_n}\right) (A_0(z_j) - A_0(y_j)) \right| \leq \max_{n \leq j \leq 2n-k} \left|1 - \frac{\epsilon_j}{\epsilon_n}\right| \cdot \sum_{j=n}^{2n-k} |A_0(z_j) - A_0(y_j)| = O\left(\frac{1}{n^2}\right),$$

by Proposition 5.27. Therefore, as in the proof of Proposition 5.20,

$$\begin{aligned} \sum_{j=n}^{2n-k} A_0(y_j) - \frac{\epsilon_j}{\epsilon_n} A_0(z_j) &= -\frac{C_1(b-1)}{n} + \frac{1}{2n} \sum_{j=n}^{2n-k} A_0(y_j) + O\left(\frac{\log n}{n^2}\right) \\ &= -\frac{1}{n} \left(C_1(b-1) + \frac{1}{2z_{2n+1-k}} + \frac{1}{2} \phi_f^o(z_{2n+1-k}) \right) + O\left(\frac{\log n}{n^2}\right) \end{aligned}$$

from which the lemma follows. \square

Lemma 5.31. *We have*

$$E_3 = \frac{b-1}{n} + O\left(\frac{1}{n^2}\right).$$

Proof of the lemma. The proof is the same as in the incoming case: it follows from the fact that $A_0(y) = (b-1)y^2 + O(y^3)$ and

$$y_j = \frac{1}{2n-j} + O\left(\frac{1}{(2n-j)^2}\right). \quad \square$$

This completes the proof of Proposition 5.28. \square

5B3. Conclusion.

Proposition 5.32. *We have*

$$\frac{1}{\epsilon_n} \left(\left(\sum_{j=0}^{2n} \epsilon_j \right) - 2 \right) = -\frac{1}{n} \left(\frac{1}{2} + \phi_g^t(w_0) \right) + O\left(\frac{1}{n^2}\right).$$

Proof. We have

$$\begin{aligned} \sum_{j=0}^{2n} \epsilon_j &= \sum_{j=0}^{2n} \frac{1}{\sqrt{n^2 + j + \phi_g^t(w_0) + o(1)}} \\ &= \sum_{j=0}^{2n} \frac{1}{n} \left(1 - \frac{j}{2n^2} - \frac{\phi_g^t(w_0)}{2n^2} + o\left(\frac{1}{n^2}\right) \right) \\ &= 2 + \frac{1}{n} - \frac{\phi_g^t(w_0)}{n^2} + o\left(\frac{1}{n^2}\right) - \frac{1}{2n^3} \sum_{j=0}^{2n} j \\ &= 2 + \frac{1}{n} - \frac{\phi_g^t(w_0)}{n^2} + o\left(\frac{1}{n^2}\right) - \frac{1}{2n^3} \frac{2n(2n+1)}{2} \\ &= 2 - \frac{\phi_g^t(w_0)}{n^2} - \frac{1}{2n^2} + o\left(\frac{1}{n^2}\right). \end{aligned}$$

On the other hand

$$\frac{1}{\epsilon_n} = \sqrt{n^2 + n + O(1)} = n \left(1 + \frac{1}{2n} + O\left(\frac{1}{n^2}\right) \right) = n + \frac{1}{2} + O\left(\frac{1}{n}\right),$$

and therefore

$$\begin{aligned} \frac{1}{\epsilon_n} \left(\left(\sum_{j=0}^{2n} \epsilon_j \right) - 2 \right) &= \left(-\frac{\phi_g^t(w_0)}{n^2} - \frac{1}{2n^2} + o\left(\frac{1}{n^2}\right) \right) \left(n + \frac{1}{2} + O\left(\frac{1}{n}\right) \right) \\ &= -\frac{1}{n} \left(\frac{1}{2} + \phi_g^t(w_0) \right) + O\left(\frac{1}{n^2}\right). \end{aligned} \quad \square$$

We are now finally ready to prove the following theorem:

Theorem 5.33 (Lavaurs' theorem with an error estimate). *Let $K \subset \mathcal{B}_f \times \mathcal{B}_g$ be a compact set. For all $(z_0, w_0) \in K$ and all sufficiently large n we have*

$$z_{2n+1} = \mathcal{L}_f(z_0) + \frac{h(z_0, w_0)}{n} + O\left(\frac{\log n}{n^2}\right),$$

where

$$h(z, w) = \frac{\mathcal{L}'_f(z)}{(\phi'_f)^t(z)} \left(2C_0 + 2(C_1 - 1)(b - 1) - \frac{1}{2} + \phi_f^t(z) - \phi_g^t(w) \right)$$

is holomorphic on $\mathcal{B}_f \times \mathcal{B}_g$ and the constant in $O((\log n)/n^2)$ is independent of the point (z_0, w_0) and the integer n .

Proof. We have, by definition

$$\phi_n^o(z_{2n+1-k}) = \frac{1}{\epsilon_n} Z_n^o + \frac{1}{\epsilon_n} \sum_{j=n}^{2n-k} \epsilon_j = \frac{Z_n^o}{\epsilon_n} - \frac{2}{\epsilon_n} + \frac{1}{\epsilon_n} \sum_{j=n}^{2n-k} \epsilon_j = \phi_n^t(z_0) - \frac{2}{\epsilon_n} + \frac{1}{\epsilon_n} \sum_{j=0}^{2n-k} \epsilon_j,$$

and therefore

$$\phi_f^o(z_{2n+1-k}) + \frac{E^o(z_{2n+1-k})}{n} = \phi_f'(z_0) + \frac{E'(z_0)}{n} - \frac{\phi_g'(w_0) + \frac{1}{2}}{n} - \frac{1}{\epsilon_n} \sum_{j=2n-k}^{2n} \epsilon_j + O\left(\frac{\log n}{n^2}\right),$$

by Propositions 5.20, 5.28 and 5.32.

On the other hand, we have

$$\begin{aligned} \frac{1}{\epsilon_n} \sum_{j=2n-k}^{2n} \epsilon_j &= \frac{1}{1/n - 1/(2n^2) + O(1/n^3)} \left(\frac{k}{n} - k \frac{2n}{2n^3} + O\left(\frac{1}{n^3}\right) \right) \quad (\text{by Lemma 5.2}) \\ &= \left(1 + \frac{1}{2n} + O\left(\frac{1}{n^2}\right) \right) \left(k - \frac{k}{n} + O\left(\frac{1}{n^2}\right) \right) \\ &= k - \frac{k}{2n} + O\left(\frac{1}{n^2}\right). \end{aligned}$$

Therefore

$$\phi_f^o(z_{2n+1-k}) + k + \frac{E^o(z_{2n+1-k}) - \frac{1}{2}k}{n} = \phi_f'(z_0) + \frac{E'(z_0)}{n} - \frac{\phi_g'(w_0) + \frac{1}{2}}{n} + O\left(\frac{\log n}{n^2}\right).$$

Recall that the outgoing Fatou coordinate ϕ_f^o has a well-defined inverse $\psi_f : \mathbb{C} \rightarrow \mathbb{C}$ satisfying the functional equation $\psi_f(Z+1) = f \circ \psi_f(Z)$. Observe that since $k = O(1)$, we have

$$\psi_f(\phi_f^o(z_{2n+1-k}) + k) = f^k(z_{2n+1-k}) + O\left(\frac{1}{n^2}\right) = z_{2n+1} + O\left(\frac{1}{n^2}\right).$$

Therefore, composing on both sides by ψ_f and setting $E^o(z_{2n+1}) := E^o(z_{2n+1-k}) - \frac{1}{2}k$, we get

$$\begin{aligned} z_{2n+1} &= (\phi_f^o)^{-1} \left(\phi_f'(z_0) + \frac{E'(z_0) - E^o(z_{2n+1}) - \frac{1}{2} - \phi_g'(w_0)}{n} + O\left(\frac{\log n}{n^2}\right) \right) \\ &= \mathcal{L}_f(z_0) + ((\phi_f^o)^{-1})'(\phi_f'(z_0)) \left(\frac{E'(z_0) - E^o(z_{2n+1}) - \frac{1}{2} - \phi_g'(w_0)}{n} \right) + O\left(\frac{\log n}{n^2}\right) \\ &= \mathcal{L}_f(z_0) + \frac{\mathcal{L}'_f(z_0)}{(\phi_f^o)'(z_0)} \left(\frac{E'(z_0) - E^o(z_{2n+1}) - \frac{1}{2} - \phi_g'(w_0)}{n} \right) + O\left(\frac{\log n}{n^2}\right). \end{aligned}$$

In particular, we have proved that $z_{2n+1} = \mathcal{L}_f(z_0) + O(1/n)$. From there, we deduce that

$$\phi_f^o(z_{2n+1-k}) + k = \phi_f^o(z_0) + O\left(\frac{1}{n}\right).$$

Plugging this into the expression for $E^o(z_{2n+1})$, we finally obtain

$$z_{2n+1} = \mathcal{L}_f(z_0) + \frac{1}{n} \frac{\mathcal{L}'_f(z_0)}{(\phi_f^o)'(z_0)} (2C_0 + 2(C_1 - 1)(b - 1) - \frac{1}{2} + \phi_f'(z_0) - \phi_g'(w_0)) + O\left(\frac{\log n}{n^2}\right). \quad \square$$

5C. Choice of index. Assume that z_0 is a Siegel fixed point for the Lavaurs map \mathcal{L}_f , and let λ be its multiplier. Denote by κ_{z_0} the index from Theorem 2.2: it is given by the formula

$$\kappa_{z_0} = \frac{2b_2c_0}{\lambda(1-\lambda)} + \frac{c_1}{\lambda},$$

with $2b_2 = \mathcal{L}_f''(z_0)$, $c_0 = h(z_0)$, $c_1 = h'(z_0)$, and

$$h(z) := \frac{\mathcal{L}_f'(z)}{(\phi_f^t)'(z)} \left(2C_0 + 2(C_1 - 1)(b - 1) - \frac{1}{2} + \phi_f^t(z) - \phi_g^t(w_0) \right). \quad (35)$$

The function h is the error term computed in the previous section.

A straightforward computation gives us that

$$\kappa_{z_0} = 1 + \frac{C + \phi_f^t(z_0) - \phi_g^t(w_0)}{((\phi_f^t)'(z_0))^2} \left(\frac{\mathcal{L}_f''(z_0)(\phi_f^t)'(z_0)}{\lambda(1-\lambda)} - (\phi_f^t)''(z_0) \right) \quad (36)$$

for some universal constant $C \in \mathbb{R}$.

Observe that $\operatorname{Re}(\kappa_{z_0})$ is independent from w_0 if and only if

$$\frac{\mathcal{L}_f''(z_0)(\phi_f^t)'(z_0)}{\lambda(1-\lambda)} - (\phi_f^t)''(z_0) = 0. \quad (37)$$

If condition (37) is satisfied, then $\kappa_{z_0} = 1$, and accordingly, Theorem 2.2 implies that there are no wandering domains for P converging to the bi-infinite orbit of $(z_0, 0)$, since we are then in the expulsion scenario of the trichotomy.

On the other hand, if the equality (37) is not satisfied, then $w_0 \mapsto \kappa_{z_0}(w_0)$ is a nonconstant holomorphic function (defined on the parabolic basin \mathcal{B}_g) of the form $w_0 \mapsto a\phi_g^t(w_0) + b$, with $a, b \in \mathbb{C}$ (independent from w_0) and $a \neq 0$. Therefore, the condition for $\operatorname{Re}(\kappa_{z_0}(w_0))$ to be negative is equivalent to $\phi_g^t(w_0)$ belonging to some half-plane, but $\phi_g^t(\mathcal{B}_g)$ contains a domain of the form

$$U := \{W \in \mathbb{C} : \operatorname{Re}(W) > R - k|\operatorname{Im}(W)|\}$$

for some $R > 0$ and $k \in (0, 1)$; see, e.g., [Shishikura 2000, Proposition 2.2.1, p. 330]. Since U intersects any open half-plane, if condition (37) is not satisfied, then there exists some open subset $U_0 \subset U$ for which $\operatorname{Re}(\kappa_{z_0}(w_0)) < 0$, and so by Theorem 2.2 there is a wandering domain accumulating on $(z_0, 0)$.

6. A Lavaurs map with a Siegel disk

The goal of this section is to construct a polynomial f of the form $f(z) = z + z^2 + z^3 + O(z^4)$, whose Lavaurs map has a Siegel fixed point with Diophantine multiplier λ , which does not satisfy equality (37). The outline of the argument is as follows:

- We start by finding a degree-7 real polynomial whose Lavaurs map has a superattracting fixed point, and for which a suitable reformulation of (37) does not hold.

- We perturb that polynomial to get an attracting but not superattracting fixed point, in a way that equality (37) still does not hold.
- We apply quasiconformal surgery to get a multiplier arbitrarily close to 1.
- We show that in the limit, we get a polynomial whose Lavaurs map has a parabolic fixed point that does not exit the parabolic basin.
- We perturb that last polynomial to get a Siegel fixed point, leaving the family of real polynomials; we prove that condition (37) does not hold for that last polynomial.

Recall that in [Astorg et al. 2016], there are two constructions of a Lavaurs map with an attracting fixed point. One is based on a residue computation near infinity in the Ecalle cylinder and makes use of the fact that in the family $f_a(z) := z + z^2 + az^3$, the multiplier of the horn map e_a of f_a at the ends of the Ecalle cylinder is a nonconstant holomorphic function of a . This method cannot be used in a family of polynomials of the form $f(z) = z + z^2 + z^3 + O(z^4)$, where those fixed points for the horn map are persistently parabolic. This is why we adapt the second strategy for the first two steps described above.

Remark. From now on we will be using slightly different notation than in previous sections. Namely we will drop the subscript f from the Fatou coordinates and the Lavaurs map in order to have space for other indexes in the subscript. It will be clear from the context to which function the Fatou coordinates or Lavaurs maps correspond.

Let ϕ^l be the incoming Fatou coordinate, and ψ^o the outgoing Fatou parametrization. Recall that the Lavaurs map is given by $\mathcal{L} = \psi^o \circ \phi^l : \mathcal{B}_f \rightarrow \mathbb{C}$, the lifted horn map is $\mathcal{E} = \phi^l \circ \psi^o : V \rightarrow \mathbb{C}$, with $V \subset \mathbb{C}$ containing $\{Z : |\operatorname{Im}(Z)| > R\}$ for R large enough. We have $\mathcal{E} \circ \phi^l = \phi^l \circ \mathcal{L}$, and $\mathcal{E}(Z + 1) = \mathcal{E}(Z) + 1$, so \mathcal{E} descends to a self-map of \mathbb{C}/\mathbb{Z} . Conjugating by the isomorphism $Z \mapsto e^{2i\pi Z}$, we obtain a map $e : U - \{0, \infty\} \rightarrow \mathbb{C}^*$, where U is an open set containing 0 and ∞ . The map extends to U , and fixes 0 and ∞ . Since we consider polynomials with $f(z) = z + z^2 + z^3 + O(z^4)$, both of those fixed points have multiplier 1.

6A. Construction a polynomial. Let $f(z) = z + z^2 + O(z^3)$ be a polynomial and $\zeta = \mathcal{L}(\zeta)$ a fixed point of its Lavaurs map, with multiplier λ .

Definition 6.1. If $\lambda \notin \{0, 1\}$ we say that the pair (f, ζ) is degenerate if and only if

$$\frac{\mathcal{L}''(\zeta)(\phi^l)'(\zeta)}{\lambda(1-\lambda)} - (\phi^l)''(\zeta) = 0. \quad (38)$$

Lemma 6.2. *We have*

$$\frac{\mathcal{L}''(\zeta)(\phi^l)'(\zeta)}{\lambda(1-\lambda)} - (\phi^l)''(\zeta) = \frac{\lambda}{1-\lambda} \left[\frac{(\psi^o)''(\phi^l(\zeta))(\phi^l)'(\zeta)}{(\psi^o)'(\phi^l(\zeta))^2} + (\phi^l)''(\zeta) \right]. \quad (39)$$

Proof. Since $\mathcal{L} = \psi^o \circ \phi^l$ we obtain

$$\begin{aligned} \mathcal{L}'(z) &= (\psi^o)'(\phi^l(z))\phi'(z), \\ \mathcal{L}''(z) &= (\psi^o)''(\phi^l(z))\phi'(z)^2 + (\psi^o)'(\phi^l(z))(\phi^l)''(z). \end{aligned}$$

Recalling that $\mathcal{L}'(\zeta) = \lambda$ it follows that

$$\frac{\phi'(\zeta)}{\lambda} = \frac{1}{(\psi^o)'(\phi^t(\zeta))},$$

and so

$$\frac{\mathcal{L}''(\zeta)\phi'(\zeta)}{\lambda} = \frac{(\psi^o)''(\phi^t(\zeta))(\phi^t)'(\zeta)^2}{(\psi^o)'(\phi^t(\zeta))} + (\phi^t)''(\zeta).$$

It follows that

$$\begin{aligned} \frac{\mathcal{L}''(\zeta)(\phi^t)'(\zeta)}{\lambda(1-\lambda)} - (\phi^t)''(\zeta) &= \frac{1}{1-\lambda} \left[\frac{(\psi^o)''(\phi^t(\zeta))(\phi^t)'(\zeta)^2}{(\psi^o)'(\phi^t(\zeta))} + (\phi^t)''(\zeta) \right] - (\phi^t)''(\zeta) \\ &= \frac{(\psi^o)''(\phi^t(\zeta))(\phi^t)'(\zeta)^2}{(1-\lambda)(\psi^o)'(\phi^t(\zeta))} + (\phi^t)''(\zeta) \frac{\lambda}{1-\lambda} \\ &= \frac{\lambda}{1-\lambda} \left[\frac{(\psi^o)''(\phi^t(\zeta))(\phi^t)'(\zeta)}{(\psi^o)'(\phi^t(\zeta))^2} + (\phi^t)''(\zeta) \right]. \quad \square \end{aligned}$$

For the rest of the paper we shall set

$$\mathcal{F}(f, \zeta) := \frac{(\psi^o)''(\phi^t(\zeta))(\phi^t)'(\zeta)}{(\psi^o)'(\phi^t(\zeta))^2} + (\phi^t)''(\zeta), \quad (40)$$

where ψ^o and ϕ^t are the Fatou parametrization and coordinates associated to f . Note that for $\lambda \notin \{0, 1\}$ the pair (f, ζ) is degenerate if and only if $\mathcal{F}(f, \zeta) = 0$.

We record here for later use the following lemma:

Lemma 6.3. *Let $f(z) = z + z^2 + az^3 + O(z^4)$ and let ϕ^t denote its incoming Fatou coordinate. Let c be a critical point in the parabolic basin of f . Then we have $(\phi^t)''(c) = 0$ if and only if either c is multiple critical point of f , or if the orbit of c meets another critical point of f .*

Proof. The sequence of functions

$$\phi_n(z) := -\frac{1}{f^n(z)} - n - (1-a) \log n$$

converges locally uniformly on the parabolic basin to

$$\phi^t(z) := \lim_{n \rightarrow \infty} \phi_n(z).$$

Therefore $(\phi^t)''(c)$ equals $\lim_{n \rightarrow \infty} \phi_n''(c)$. Moreover, $\phi_n'(z) = (f^n)'(z)/[f^n(z)]^2$ and

$$\begin{aligned} \phi_n''(c) &= \frac{d}{dz} \Big|_{z=c} \frac{(f^n)'(z)}{[f^n(z)]^2} \\ &= \frac{(f^n)''(c)[f^n(c)]^2 - 2[(f^n)'(c)]^2 f^n(c)}{[f^n(c)]^4} \\ &= \frac{(f^n)''(c)}{[f^n(c)]^2} = f''(c) \frac{\prod_{k=1}^{n-1} f'(f^k(c))}{[f^n(c)]^2}. \end{aligned}$$

For the third and fourth equalities we used the fact that $f'(c) = 0$. Since c is in the parabolic basin of f , we have $[f^n(c)]^2 \sim 1/n^2$. Moreover, for $k \geq k_0$ with k_0 large enough, $f'(f^k(c)) \neq 0$ and

$$f'(f^k(c)) = 1 - \frac{2}{k} + O\left(\frac{\log k}{k^2}\right) = \exp\left(-\frac{2}{k} + O\left(\frac{\log k}{k^2}\right)\right).$$

Therefore

$$\prod_{k=k_0}^{n-1} f'(f^k(z)) = \prod_{k=k_0}^{n-1} \exp\left(-\frac{2}{k} + O\left(\frac{\log k}{k^2}\right)\right) = \frac{\exp(O(1))}{n^2}.$$

In particular,

$$\lim_{n \rightarrow \infty} \frac{\prod_{k=k_0}^{n-1} f'(f^k(c))}{[f^n(c)]^2} \neq 0,$$

so $(\phi^t)''(c) = 0$ if and only if $f''(c) = 0$ or $(f^k)'(c) = 0$, which concludes the proof. \square

For $t \in \mathbb{R}$, a real polynomial $P(z) = z + z^2 + z^3 + O(z^4)$ and $n > \deg P$ odd, let

$$f_t(z) = P(z) - \frac{P'(t)}{nt^{n-1}}z^n.$$

Note that $f_t'(t) = 0$; the choice of this family ensures that we have a marked critical point in \mathbb{R} . By \mathcal{L}_t we denote the Lavaurs map of phase 0 for the polynomial f_t .

Proposition 6.4. *Assume that there exists P, n and $t_\infty < 0$ as above such that:*

- (1) $f_{t_\infty}(t_\infty) = 0$.
- (2) $(d/dt)|_{t=t_\infty} f_t(t) < 0$.
- (3) f_{t_∞} has negative leading coefficient.
- (4) There exists $x > 0$ in the repelling petal of f_{t_∞} that escapes to infinity.

Then there is a sequence $t_n \rightarrow t_\infty$ such that $\mathcal{L}_{t_n}(t_n) = t_n$.

Proof. We will rely on the following two claims:

Claim 1. *For $t \in (t_\infty, t_\infty + \epsilon)$ with $\epsilon > 0$ small enough, the critical point t is in the parabolic basin of f_t .*

Proof of the claim. It is enough to show that there is $r > 0$ such that $(-r, 0)$ is in the parabolic basin of f_t for all t close enough to t_∞ . Indeed, by (1) and (2), we have that for all $r > 0$ there exists $\epsilon > 0$ such that $f_t(t) \in (-r, 0)$ for all $t \in (t_\infty, t_\infty + \epsilon)$. Let

$$r_t := \sup\{r > 0 : \text{for all } y \in (-r, 0), 0 < f_t(y)/y < 1\}.$$

For all $y \in (-r_t, 0)$, we have $t < f_t(y) < 0$; hence y is in the parabolic basin of f_t . Finally, $t \mapsto r_t$ is continuous and $r_{t_\infty} > 0$. \square

Claim 2. *There exists a sequence $\tilde{t}_n \rightarrow t_\infty$ (with $\tilde{t}_n > t_\infty$) such that $\mathcal{L}_{\tilde{t}_n}(\tilde{t}_n) = f_{\tilde{t}_n}^n(x)$.*

Proof of the claim. We adapt here the argument from [Astorg et al. 2016]. The desired equality $\mathcal{L}_{\tilde{t}_n}(\tilde{t}_n) = f_{\tilde{t}_n}^n(x)$ is equivalent to

$$\psi_{\tilde{t}_n}^o \circ \phi_{\tilde{t}_n}^l(\tilde{t}_n) = \psi_{\tilde{t}_n}^o(\phi_{\tilde{t}_n}^o(x) + n).$$

In particular, it is enough to find \tilde{t}_n such that $\phi_{\tilde{t}_n}^l(\tilde{t}_n) = \phi_{\tilde{t}_n}^o(x) + n$. We look for \tilde{t}_n under the form

$$\tilde{t}_n = t_\infty - \frac{\alpha}{n+u}, \quad \text{with } \alpha = \frac{1}{\frac{d}{dc}\big|_{c=t_\infty} f_c(c)}.$$

By the preceding claim, it is in the parabolic basin for n large enough.

We have $\phi_{\tilde{t}_n}^o(x) + n = n + \phi_{t_\infty}^o(x) + o(1)$ since the map $t \mapsto \phi_t^o$ is continuous. Additionally,

$$\begin{aligned} \phi_{\tilde{t}_n}^l(\tilde{t}_n) &= \phi_{\tilde{t}_n}^l(f_{\tilde{t}_n}(\tilde{t}_n)) - 1 \\ &= -\frac{1}{f_{\tilde{t}_n}(\tilde{t}_n)} - 1 + o(1) \quad (\text{according to the asymptotic expansion of } \phi^l) \\ &= n + u - 1 + o(1). \end{aligned}$$

Therefore, we have reduced the problem to solving the equation $u - 1 + o(1) = \phi_{t_\infty}^o(x)$ for $u \in \mathbb{R}$, where the $o(1)$ term is a continuous function of u . By the intermediate value theorem there is a solution $u = u_n \in (\phi_{t_\infty}^o(x), \phi_{t_\infty}^o(x) + 2)$. We can take $\tilde{t}_n = t_\infty - \alpha/(n + u_n)$, and since $(u_n)_{n \in \mathbb{N}}$ is bounded from below, the sequence (\tilde{t}_n) is well-defined for n large enough and converges to t_∞ . \square

We now come back to the proof of Proposition 6.4. For n large enough, $G_{\tilde{t}_n}(x) > 0$ (by continuity of the Green's function G). Therefore $\mathcal{L}_{\tilde{t}_n}(\tilde{t}_n) = f_{\tilde{t}_n}^n(x)$ tends to ∞ , and more precisely, $+\infty$ or $-\infty$ depending on the parity of n , thanks to condition (3). Therefore the continuous function $F(t) := \mathcal{L}_t(t) - t$ alternates sign between two consecutive \tilde{t}_n , so by the intermediate value theorem must have a zero t_n between them. \square

Proposition 6.5. *Let $P(z) := z + z^2 + z^3 + \frac{23}{7}z^4 + \frac{17}{7}z^5$, let $t_\infty := -1$ and let $n := 7$. Then P , n and t_∞ satisfy conditions (1)–(4) in Proposition 6.4.*

Proof. Observe that $f_{t_\infty}(t_\infty) = 0$ and $P'(t_\infty) = 1$. That second property implies that f_{t_∞} has negative leading coefficient. Therefore, conditions (1) and (3) are satisfied.

Let us check that condition (2) is also satisfied. We have

$$\frac{d}{dt}\bigg|_{t=t_\infty} f_t(t) = \frac{d}{dt}\bigg|_{t=t_\infty} P(t) - \frac{t}{n}P'(t) = \frac{n-1}{n}P'(t_\infty) - \frac{t_\infty}{n}P''(t_\infty) = \frac{6}{7} + \frac{1}{7}P''(-1) = -\frac{50}{49} < 0.$$

Finally, condition (4) is satisfied for $x := 1$. Indeed, recall here that if $f(z) = \sum_{k=0}^n a_k z^k$ is a complex polynomial and

$$R = \max\left\{1, \frac{1 + |a_0| + \dots + |a_{n-1}|}{|a_n|}\right\}$$

then for all $|z| > R$ we have $|f(z)| \geq |z|^n/R$; hence if an orbit at any point leaves the disk of radius R , then it must converge to infinity. Observe that for our polynomial f_{t_∞} we have $R = 68$ and that a straightforward computation yields $f_{t_\infty}(1) = \frac{60}{7}$ and $|f_{t_\infty}^2(1)| > 68$.

This proves rigorously that $x := 1$ has unbounded orbit under f_{t_∞} . \square

Lemma 6.6. *For $\epsilon_0 > 0$ small enough, there exists $t > -1$ such that the following properties hold for f_t :*

- (1) \mathcal{L}_t has a fixed point x_t with multiplier $\epsilon_0 \neq 0$.
- (2) $\mathcal{F}(f_t, x_t) \neq 0$.
- (3) f_t has four real critical points, ordered from left to right, c_1, c_2, c_3, c_4 , with $t = c_2$, and two nonreal complex conjugate critical points c' and \bar{c}' .
- (4) The critical points c_1 and c_4 lie in the basin of infinity; the critical points c_2 and c_3 are in the parabolic basin.
- (5) There is a unique repelling fixed point $\xi \in (c_1, c_2)$, and the intersection of \mathbb{R} and the immediate basin of attraction of 0 is $(\xi, 0)$.
- (6) There is a unique $y \in (\xi, c_2)$ such that $f_t(y) = c_2$.

Proof. We will find t by taking a perturbation of one of the t_{n_0} constructed above, with n_0 large enough.

First, note that properties (3)–(6) hold for

$$f := f_{t_\infty} : z \mapsto z + z^2 + z^3 + \frac{23}{7}z^4 + \frac{17}{7}z^5 - \frac{z^7}{7};$$

we leave the details to the reader; see Figure 2. Therefore, for n_0 large enough, properties (3)–(6) still hold for $f_{t_{n_0}}$, as these properties are clearly open (in \mathbb{R}) near $t = t_\infty$. To lighten the notation, we let $f := f_{t_{n_0}}$ and $c_2 := t_{n_0}$.

We now claim that \mathcal{F} is well-defined at (f, c_2) , and that $\mathcal{F}(f, c_2) \neq 0$. According to Lemma 6.3, since f satisfies conditions (3)–(6), we have $(\phi^t)''(c) \neq 0$. Indeed, c_2 is a simple critical point of f , and we claim that the forward orbit of c_2 does not meet any other critical point of f . To see this, note that the critical point c_2 is simple for f , and real. Since c' and \bar{c}' are not real, the orbit of c_2 cannot land on either of them. Since the critical points c_1 and c_4 do not belong to the parabolic basin, the orbit of c_2 cannot land on them either. Finally, since $f(c_2) > c_3$, and since $f(c_2)$ belongs to a small attracting petal in which the sequence of iterates $(f^n(c_2))_{n \in \mathbb{N}}$ is increasing, the orbit of c_2 cannot land on c_3 either.

Now that we have proved that $(\phi^t)''(c_2) \neq 0$, it is sufficient to prove that

$$\frac{(\psi^o)''(\phi^t(c_2))(\phi^t)'(c_2)}{(\psi^o)'(\phi^t(c_2))^2} = 0.$$

In fact, since $(\phi^t)'(c_2) = 0$, it suffices to prove that $(\psi^o)'(\phi^t(c_2)) \neq 0$. Recall that for any $Z \in \mathbb{C}$, $(\psi^o)'(Z) = 0$ if and only if there exists $n \geq 1$ such that $(\psi^o)'(Z - n)$ is a critical point for f ; here, $Z = \phi^t(c_2)$ and $\psi^o \circ \phi^t(c_2) = c_2$, so we must prove that for all $n \geq 1$ and any critical point c_i of f , $f^n(c_i) \neq c_2$. Since c_1 and c_4 escape, neither of their orbits can land on c_2 , and since c_2 is not periodic under f , its own orbit cannot land on itself either. Since c_3 is in the immediate parabolic basin, the orbit $(f^n(c_3))_{n \in \mathbb{N}}$ is increasing, and so does not contain c_2 since $c_3 > c_2$.

Finally, it remains to argue that the orbits of the two nonreal critical points c' and \bar{c}' do not eventually land on c_2 . To see that it cannot be the case, note that since the horn map e of f has two parabolic fixed points at 0 and ∞ corresponding to the ends of the Ecalle cylinder, each of those fixed points must attract singular values of e distinct from themselves; see [Astorg et al. 2016]. The singular values of e are




Figure 2. The graph of $f := f_{t_\infty}$ (blue), with the line $y = x$ in red. We have $c_1 \approx -2.8$, $c_2 = -1$, $c_3 \approx -0.4$, and $c_4 \approx 4$. The critical values $f(c_1)$ and $f(c_4)$ are out of the picture.

the fixed points at 0 and ∞ , as well as the $\pi(c_i)$, where c_i are the critical points of f in the parabolic basin and $\pi(z) = e^{2i\pi\phi^t(z)}$. If $f^n(c') = c_2$ for some $n \geq 1$, then by real symmetry we would also have $f^n(\bar{c}') = c_2$, and so $\pi(c') = \pi(\bar{c}') = \pi(c_2)$; but then $\pi(c_3)$ would be the only nonfixed singular value of e , which is impossible.

Therefore f has no critical relation, and so $(\psi^o)'(\phi^t(c_2)) \neq 0$, and $\mathcal{F}(f, c_2) \neq 0$ as announced.

To summarize, we have proved that for n_0 large enough, the polynomial $f_{t_{n_0}}$ satisfies properties (2)–(6). Since t_{n_0} is a superattracting fixed point of $\mathcal{L}_{t_{n_0}}$ but persistently fixed, for $\epsilon_0 > 0$ small enough, there exists t close to t_{n_0} such that f_t satisfies (1), and by openness, if ϵ_0 is small enough, f_t still satisfies (2)–(6). \square

The next step is to use quasiconformal deformations to construct an immersed disk D in parameter space passing through f_t , made of polynomials p_u , whose Lavaurs map has an attracting fixed point of multiplier $e^{2i\pi u}$, $u \in \mathbb{H}$. We purposely use the notation p_u instead of f_t to emphasize the fact that except for f_t , the polynomials p_u do not a priori belong to the family $(f_t)_{t \in \mathbb{R}^*}$.

Proposition 6.7. *Let $p := f_t$ and $\epsilon_0 > 0$ be as in Lemma 6.6. There exists a holomorphic map $\Phi : \mathbb{H} \rightarrow \mathcal{P}_7$ such that:*

- (1) $\Phi(u_0) = p$ for some $u_0 \in \mathbb{H}$ with $e^{2i\pi u_0} = \epsilon_0$.
- (2) For all $u \in \mathbb{H}$, the Lavaurs map of $\Phi(u) := p_u$ has a fixed point z_u of multiplier $e^{2i\pi u} \in \mathbb{D}^*$, and $u \mapsto z_u$ is holomorphic.
- (3) All the maps p_u are quasiconformally conjugated to p , the conjugacy being holomorphic outside of the grand orbit under p of the attracting basin of $z_{u_0} := x_t$.
- (4) If $e^{2i\pi u} \in (0, 1)$, then the conjugacy preserves the real line.
- (5) The set $\Phi(\mathbb{H})$ is relatively compact in \mathcal{P}_7 .

Proof. Let $e : U \rightarrow \mathbb{P}^1$ be the horn map of g ; since \mathcal{L} has an attracting fixed point $z_{u_0} := x_t$, so does e (since they are semiconjugated). Denote this attracting fixed point by x .

Let $u \in \mathbb{H}$ and μ be a Beltrami form invariant by e (i.e., $e^*\mu = \mu$) such that the corresponding quasiconformal homeomorphism h_μ conjugates e to some holomorphic map e_μ with an attracting fixed point of multiplier $e^{2i\pi u}$: $h_\mu \circ e = e_\mu \circ h_\mu$ and $e'_\mu(h_\mu(x)) = e^{2i\pi u}$. We recall here briefly how to construct such a Beltrami form, and refer the reader to [Branner and Fagella 2014] for more details. If τ is a linearizing coordinate for the horn map e near x , i.e., a holomorphic map defined near p satisfying the functional equation $\tau \circ e = \epsilon_0 \tau$, we set

$$\mu = \mu(u) := \tau^* \left(\frac{u - u_0}{u + u_0} \frac{z}{\bar{z}} \frac{d\bar{z}}{dz} \right), \quad (41)$$

where $u_0 \in \mathbb{H}$ is any point such that $e^{2i\pi u_0} = \epsilon_0$. Notice that $u \mapsto \mu(u)$ is holomorphic. In the rest of the proof, we fix $u \in \mathbb{H}$ and just use the notation μ instead of $\mu(u)$.

We choose the normalization of h_μ so that it fixes $0, 1$ and ∞ . Let $E(z) := e^{2i\pi z}$ and $T_1(z) := z + 1$. We define

- (1) $\nu := E^*\mu$ so that $\nu = T_1^*\nu$, and $\nu = \mathcal{E}^*\nu$,
- (2) $\sigma := \phi^*\nu$ so that $\sigma = g^*\sigma$ and $\sigma = \mathcal{L}^*\sigma$,
- (3) the quasiconformal homeomorphisms h_ν and h_σ associated to ν, σ respectively.

Since $\nu = T_1^*\nu$, the map $h_\nu \circ T_1 \circ h_\nu^{-1} : \mathbb{C} \rightarrow \mathbb{C}$ is holomorphic; since it is conjugated to T_1 , it is also a translation (distinct from the identity), and we choose the normalization of h_ν so that $h_\nu \circ T_1 \circ h_\nu^{-1} = T_1$ and $h_\nu(0) = 0$. Similarly, since $\sigma = g^*\sigma$, the map $p_u := h_\sigma \circ p \circ h_\sigma^{-1}$ is holomorphic, and hence a polynomial (since it has the same topological degree as f); it also has a parabolic fixed point with one attracting petal at the origin. We choose the unique normalization of h_σ such that $p_u(z) = z + z^2 + O(z^3)$. We set $\Phi(u) := p_u$; the holomorphic dependence $u \mapsto \mu(u)$ and the parametric version of the Ahlfors–Bers theorem imply that Φ is holomorphic on \mathbb{H} .

We now define

- (1) $\phi_\sigma := h_\nu \circ \phi \circ h_\sigma^{-1} : h_\sigma(B) \rightarrow \mathbb{C}$, where B is the parabolic basin of f ,
- (2) $\psi_\nu := h_\sigma \circ \psi \circ h_\nu^{-1} : \mathbb{C} \rightarrow \mathbb{C}$.

Lemma 6.8. *The map ϕ_σ is an incoming Fatou coordinate for p_u , and the map ψ_ν is an outgoing Fatou parametrization for p_u .*

Proof of the lemma. We start with ϕ_σ . First, note that since $\sigma = \phi^*\nu$, the map ϕ_σ is holomorphic on $B_\sigma := h_\sigma(B)$, which is exactly the parabolic basin of p_u . Then, note that

$$\begin{aligned} \phi_\sigma \circ p_u &= h_\nu \circ \phi \circ h_\sigma^{-1} \circ p_u = h_\nu \circ \phi \circ g \circ h_\sigma^{-1} \\ &= h_\nu \circ T_1 \circ \phi \circ h_\sigma^{-1} = T_1 \circ h_\nu \circ \phi \circ h_\sigma^{-1} = T_1 \circ \phi_\sigma. \end{aligned}$$

So ϕ_σ conjugates p_u on the whole parabolic basin to a translation, which means it is a Fatou coordinate.

The proof is completely analogous for ψ_ν : first, to prove that ψ_ν is holomorphic, note that $\nu = \psi^*\sigma$. Indeed, $\nu = \mathcal{E}^*\nu = \psi^*\phi^*\nu = \psi^*\sigma$. To conclude, one can check directly that $\psi_\nu \circ T_1 = p_u \circ \psi_\nu$. \square

As a consequence of the lemma, $\mathcal{E}_v := h_v \circ \mathcal{E} \circ h_v^{-1}$ is a lifted horn map of p_u , and $\mathcal{L}_\sigma := h_\sigma \circ \mathcal{L} \circ h_\sigma^{-1}$ is a Lavaurs map of p_u , and they have the same phase. The phase could a priori be a nonzero, but we will prove that it is not the case. In order to do that, first we will prove that $E \circ \mathcal{E}_v = e_\mu \circ E$, i.e., that e_μ is a horn map that lifts to \mathcal{E}_v .

Since $v = E^* \mu$, the map $E_v := h_\mu \circ E \circ h_v^{-1} : \mathbb{C} \rightarrow \mathbb{C}^*$ is holomorphic. Moreover, since $E : \mathbb{C} \rightarrow \mathbb{C}^*$ is a universal cover, so is E_v . So E_v is of the form $E_v(z) = \lambda e^{\alpha z}$, and with our choices of normalizations we find $E_v(z) = e^{2i\pi z} = E(z)$. So $E \circ h_v = h_\mu \circ E$.

From this, we deduce

$$\begin{aligned} E \circ \mathcal{E}_v &= E \circ h_v \circ \mathcal{E} \circ h_v^{-1} = h_\mu \circ E \circ \mathcal{E} \circ h_v^{-1} \\ &= h_\mu \circ e \circ E \circ h_v^{-1} = h_\mu \circ e \circ h_\mu^{-1} \circ E = e_\mu \circ E. \end{aligned}$$

Finally, it remains to observe that since e_μ is topologically conjugated to e , it also has two parabolic fixed points at 0 and ∞ respectively, each of multiplier 1. Recall that the horn map of phase 0 of a parabolic polynomial $f(z) = z + z^2 + az^3 + O(z^4)$ has multipliers at 0 and ∞ both equal to $e^{2\pi^2(1-a)}$, and that the horn map of phase $\varphi \in \mathbb{C}/\mathbb{Z}$ is obtained from the horn map e of phase 0 by multiplication by $e^{2i\pi\varphi}$. In particular, its multipliers at 0 and ∞ are respectively $e^{2\pi^2(1-a)+2i\pi\varphi}$ and $e^{2\pi^2(1-a)-2i\pi\varphi}$. In this case, since both multipliers are equal to 1, we must have $a = 1$ and $\varphi = 0$. Therefore, \mathcal{L}_σ is the Lavaurs map of phase 0 of p_u , and $p_u(z) = z + z^2 + z^3 + O(z^4)$.

Finally, if $\pi_\sigma(z) := e^{2i\pi\phi_\sigma(z)}$, then $\pi_\sigma \circ \mathcal{L}_\sigma = e_\mu \circ \pi_\sigma$, and π_σ is locally invertible near $z_u := h_\sigma(z_{u_0})$, and $\pi_\sigma(z_u) = h_\mu(x)$. Therefore, z_u as a fixed point of \mathcal{L}_σ has the same multiplier $e^{2i\pi u}$ as $h_\mu(x)$. This proves claims (1)–(3) of the proposition.

To prove claim (4), note that if $e^{2i\pi u} \in (0, 1)$ then the Beltrami form

$$\frac{u - u_0}{u + u_0} \frac{z}{\bar{z}} \frac{d\bar{z}}{dz}$$

has real symmetry (since then $(u - u_0)/(u + u_0) \in \mathbb{R}$). We claim that this implies that σ has real symmetry. Indeed, since $g(\mathbb{R}) = \mathbb{R}$, its Lavaurs map \mathcal{L} maps a small interval $I \subset \mathbb{R}$ centered at x_t into itself. Moreover, the map $\tau \circ \pi$ semiconjugates \mathcal{L} to the multiplication by $\epsilon_0 > 0$; so $\tau \circ \pi$ maps I into \mathbb{R} , which means that the holomorphic map $\tau \circ \pi$ is real: $\tau \circ \pi(\bar{z}) = \overline{\tau \circ \pi(z)}$ for all z in the parabolic basin of g . Therefore

$$\sigma = (\tau \circ \pi)^* \left(\frac{u - u_0}{u + u_0} \frac{z}{\bar{z}} \frac{d\bar{z}}{dz} \right)$$

has real symmetry; hence h_σ restricts to a real homeomorphism.

Finally, $\Phi : \mathbb{H} \rightarrow \mathcal{P}_7$ is bounded in the space of polynomials of degree 7. Indeed, by [Bassanelli and Berteloot 2011, Proposition 4.4] the set of polynomials of given degree with given values of the Green's function at the critical points is bounded, and since the conjugacy between the p_u and p is analytic outside of the parabolic basin, their Green's functions have the same values at critical points. \square

Proposition 6.9. *With the same notation as before, there exists p_0 in the closure of $\Phi(\mathbb{H})$ such that the Lavaurs map of p_0 has a parabolic fixed point of multiplier 1.*

Proof. Applying Proposition 6.7 with $u_n = i/n$, we get a sequence of polynomials p_{u_n} such that $p_{u_n}(z) = z + z^2 + z^3 + O(z^4)$, and the Lavaurs map \mathcal{L}_n of p_{u_n} has a fixed point x_n of multiplier $e^{-2\pi/n}$.

Each of the p_{u_n} are quasiconformally conjugate to the real polynomial f_t from Lemma 6.6 by a homeomorphism whose restriction to the real line is real and increasing, so the p_{u_n} still satisfy the properties (3)–(6) from Lemma 6.6.

By item (5) in Proposition 6.7, the sequence $(p_{u_n})_{n \in \mathbb{N}}$ is bounded in the space of degree-7 polynomials. So up to extracting, we may assume that:

- (1) p_{u_n} converges to a degree 7 polynomial p_0 .
- (2) The critical points $c_{i,n}$ of p_{u_n} converge to critical points c_i of p_0 .
- (3) The repelling fixed point ξ_n converges to a nonattracting fixed point ξ of p_0 .
- (4) x_n converges to $x \in \mathbb{R}$ and y_n to $y \in \mathbb{R}$.

We denote by \mathcal{L} the Lavaurs map of p_0 . If we can prove that x lies in the parabolic basin of p_0 , then we will get that $\mathcal{L}(x) = x$ and $\mathcal{L}'(x) = 1$. To do that, it is enough to prove that $x \in (\xi, 0)$. But for all n , we have

$$\xi_n < y_n < x_n < c_{2,n} < 0;$$

hence $\xi < y \leq x \leq c_2 < 0$. The inequality $\xi < y$ is strict because as a limit of repelling fixed points, we have $|f'(\xi)| \geq 1$, so we cannot have $y = \xi$, for otherwise we would have $\xi = f(\xi) = f(y) = c_2$ and so $f'(\xi) = 0$, a contradiction. Similarly, we cannot have $c_2 = 0$ since $p_0'(0) = 1 \neq 0$. So $x \in (\xi, 0)$ and ξ is in the parabolic basin of f , and so $\mathcal{L}'(x) = 1$ and $\mathcal{L}(x) = x$. Therefore p_0 has the desired property. \square

Proposition 6.10. *There exists a polynomial $g(z) = z + z^2 + z^3 + O(z^4)$ of degree 7 such that*

- (1) \mathcal{L} has a Siegel fixed point ζ with Diophantine multiplier, and
- (2) the pair (g, ζ) is nondegenerate.

Proof. Recall that \mathcal{P}_7 denotes the space of degree-7 polynomials of the form $f(z) = z + z^2 + z^3 + O(z^4)$, and let $V = \{(f, \zeta) \in \mathcal{P}_7 \times \mathbb{C} : \zeta \in \mathcal{B}_f\}$. V may be identified with an open set in \mathbb{C}^5 . Finally, we consider $F := \{(f, \zeta) \in V : \mathcal{L}(\zeta) = \zeta\}$, which is an analytic hypersurface of V .

We consider the functions $\lambda : F \rightarrow \mathbb{C}$ and $\mathcal{F} : F \rightarrow \mathbb{C}$ defined as $\lambda(f, \zeta) = \mathcal{L}'(\zeta)$ and

$$\mathcal{F}(f, \zeta) = \frac{(\psi^o)''(\phi^t(\zeta))(\phi^t)'(\zeta)}{(\psi^o)'(\phi^t(\zeta))^2} + (\phi^t)''(\zeta),$$

where ϕ^t and ψ^o are the Fatou coordinate and parametrization of f . The function λ is analytic on F , and \mathcal{F} is meromorphic on F and analytic on $\lambda^{-1}(\mathbb{C}^*)$, since $(\psi^o)'(\phi^t(z)) = 0$ implies that $\mathcal{L}'(z) = 0$.

Let $\Phi : \mathbb{H} \rightarrow \mathcal{P}_7$ be the map defined in Proposition 6.7, and let $\tilde{\Phi} : \mathbb{H} \rightarrow F$ be the map given by $\tilde{\Phi}(u) = (p_u, z_u)$, where z_u is the fixed point of the Lavaurs map of p_u with multiplier $e^{2i\pi u}$. Then $D := \tilde{\Phi}(\mathbb{H})$ is contained in one irreducible component F_0 of F .

Let p_0 be the polynomial given by Proposition 6.9 such that its Lavaurs map has a parabolic fixed point z_0 . By Proposition 6.9, (p_0, z_0) is in the closure of D in V ; therefore $(p_0, z_0) \in F_0$.

Assume for a contradiction that all pairs $(f, \zeta) \in F_0$ for which $\mathcal{L}'(\zeta)$ has modulus 1 and Diophantine argument are degenerate. Then by the density of Diophantine numbers on the real line, we must have

$\mathcal{F}(f, \zeta) = 0$ on $\lambda^{-1}(S^1) \cap F_0$. Since for all $u \in \mathbb{H}$, we have $\lambda \circ \tilde{\Phi}(u) = e^{2i\pi u}$, the analytic map λ is nonconstant on F_0 . In particular, $\lambda^{-1}(S^1)$ is a real-analytic subset of F_0 of real codimension 1, nonempty since $\lambda(p_0, z_0) = 1$. By Proposition 6.7, D contains (f_t, x_t) , where f_t is the polynomial given by Lemma 6.6, and such that $\mathcal{F}(f_t, z_t) \neq 0$. So the analytic map \mathcal{F} is not identically zero on F_0 , and therefore it cannot vanish identically on $\lambda^{-1}(S^1) \cap F_0$, a contradiction. \square

References

- [Astorg et al. 2016] M. Astorg, X. Buff, R. Dujardin, H. Peters, and J. Raissy, “A two-dimensional polynomial mapping with a wandering Fatou component”, *Ann. of Math. (2)* **184**:1 (2016), 263–313. MR Zbl
- [Bassanelli and Berteloot 2011] G. Bassanelli and F. Berteloot, “Distribution of polynomials with cycles of a given multiplier”, *Nagoya Math. J.* **201** (2011), 23–43. MR Zbl
- [Bedford et al. 2017] E. Bedford, J. Smillie, and T. Ueda, “Semi-parabolic bifurcations in complex dimension two”, *Comm. Math. Phys.* **350**:1 (2017), 1–29. MR Zbl
- [Bracci and Zaitsev 2013] F. Bracci and D. Zaitsev, “Dynamics of one-resonant biholomorphisms”, *J. Eur. Math. Soc.* **15**:1 (2013), 179–200. MR Zbl
- [Branner and Fagella 2014] B. Branner and N. Fagella, *Quasiconformal surgery in holomorphic dynamics*, Cambridge Stud. Adv. Math. **141**, Cambridge Univ. Press, 2014. MR Zbl
- [Bruno 1971] A. D. Bruno, “Analytic form of differential equations, I”, *Tr. Mosk. Mat. Obs.* **25** (1971), 119–262. In Russian; translated in *Trans. Moscow Math. Soc.* **25** (1971), 131–288. MR Zbl
- [Bruno 1972] A. D. Bruno, “Analytic form of differential equations, II”, *Tr. Mosk. Mat. Obs.* **26** (1972), 199–239. In Russian; translated in *Trans. Moscow Math. Soc.* **26** (1972), 199–238. MR Zbl
- [Hakim 1998] M. Hakim, “Analytic transformations of $(\mathbb{C}^P, 0)$ tangent to the identity”, *Duke Math. J.* **92**:2 (1998), 403–428. MR Zbl
- [Ji 2020] Z. Ji, “Non-wandering Fatou components for strongly attracting polynomial skew products”, *J. Geom. Anal.* **30**:1 (2020), 124–152. MR Zbl
- [Jonsson 1999] M. Jonsson, “Dynamics of polynomial skew products on \mathbb{C}^2 ”, *Math. Ann.* **314**:3 (1999), 403–447. MR Zbl
- [Lavaurs 1989] P. Lavaurs, *Systèmes dynamiques holomorphiques: explosion des points périodiques*, Ph.D. thesis, Université Paris-Sud, 1989.
- [Lilov 2004] K. Lilov, *Fatou theory in two dimensions*, Ph.D. thesis, University of Michigan, 2004, available at <https://www.proquest.com/docview/305178400>.
- [López-Hernanz and Rosas 2020] L. López-Hernanz and R. Rosas, “Characteristic directions of two-dimensional biholomorphisms”, *Compos. Math.* **156**:5 (2020), 869–880. MR Zbl
- [Peters and Raissy 2019] H. Peters and J. Raissy, “Fatou components of elliptic polynomial skew products”, *Ergodic Theory Dynam. Systems* **39**:8 (2019), 2235–2247. MR Zbl
- [Peters and Smit 2018] H. Peters and I. M. Smit, “Fatou components of attracting skew-products”, *J. Geom. Anal.* **28**:1 (2018), 84–110. MR Zbl
- [Peters and Vivas 2016] H. Peters and L. R. Vivas, “Polynomial skew-products with wandering Fatou-disks”, *Math. Z.* **283**:1–2 (2016), 349–366. MR Zbl
- [Shishikura 2000] M. Shishikura, “Bifurcation of parabolic fixed points”, pp. 325–363 in *The Mandelbrot set, theme and variations*, edited by L. Tan, Lond. Math. Soc. Lect. Note Ser. **274**, Cambridge Univ. Press, 2000. MR Zbl
- [Siegel 1942] C. L. Siegel, “Iteration of analytic functions”, *Ann. of Math. (2)* **43** (1942), 607–612. MR Zbl
- [Sullivan 1985] D. Sullivan, “Quasiconformal homeomorphisms and dynamics, I: Solution of the Fatou–Julia problem on wandering domains”, *Ann. of Math. (2)* **122**:3 (1985), 401–418. MR Zbl
- [Yoccoz 1995] J.-C. Yoccoz, “Théorème de Siegel, nombres de Bruno et polynômes quadratiques”, pp. 3–88 in *Petits diviseurs en dimension 1*, Astérisque **231**, Soc. Math. France, Paris, 1995. MR

Received 14 Oct 2019. Revised 23 Feb 2021. Accepted 14 Jun 2021.

MATTHIEU ASTORG: matthieu.astorg@univ-orleans.fr

Institut Denis Poisson, Collegium Sciences et Techniques, Université d'Orléans, Orléans, France

LUKA BOC THALER: luka.boc@pef.uni-lj.si

Faculty of Education, University of Ljubljana, Ljubljana, Slovenia

and

Institute of Mathematics, Physics and Mechanics, Ljubljana, Slovenia

HAN PETERS: hanpeters77@gmail.com

Korteweg de Vries Institute for Mathematics, University of Amsterdam, Amsterdam, Netherlands

Analysis & PDE

msp.org/apde

EDITORS-IN-CHIEF

Patrick Gérard Université Paris Sud XI, France
patrick.gerard@universite-paris-saclay.fr

Clément Mouhot Cambridge University, UK
c.mouhot@dpms.cam.ac.uk

BOARD OF EDITORS

Massimiliano Berti	Scuola Intern. Sup. di Studi Avanzati, Italy berti@sissa.it	Werner Müller	Universität Bonn, Germany mueller@math.uni-bonn.de
Zbigniew Błocki	Uniwersytet Jagielloński, Poland zbigniew.blocki@uj.edu.pl	Gilles Pisier	Texas A&M University, and Paris 6 pisier@math.tamu.edu
Charles Fefferman	Princeton University, USA cf@math.princeton.edu	Igor Rodnianski	Princeton University, USA irod@math.princeton.edu
Isabelle Gallagher	Université Paris-Diderot, IMJ-PRG, France gallagher@math.ens.fr	Yum-Tong Siu	Harvard University, USA siu@math.harvard.edu
Colin Guillarmou	Université Paris-Saclay, France colin.guillarmou@universite-paris-saclay.fr	Terence Tao	University of California, Los Angeles, USA tao@math.ucla.edu
Ursula Hamenstaedt	Universität Bonn, Germany ursula@math.uni-bonn.de	Michael E. Taylor	Univ. of North Carolina, Chapel Hill, USA met@math.unc.edu
Vadim Kaloshin	University of Maryland, USA vadim.kaloshin@gmail.com	Gunther Uhlmann	University of Washington, USA gunther@math.washington.edu
Izabella Laba	University of British Columbia, Canada ilaba@math.ubc.ca	András Vasy	Stanford University, USA andras@math.stanford.edu
Anna L. Mazzucato	Penn State University, USA alm24@psu.edu	Dan Virgil Voiculescu	University of California, Berkeley, USA dvv@math.berkeley.edu
Richard B. Melrose	Massachusetts Inst. of Tech., USA rbm@math.mit.edu	Jim Wright	University of Edinburgh, UK j.r.wright@ed.ac.uk
Frank Merle	Université de Cergy-Pontoise, France merle@ihes.fr	Maciej Zworski	University of California, Berkeley, USA zworski@math.berkeley.edu
William Minicozzi II	Johns Hopkins University, USA minicozz@math.jhu.edu		

PRODUCTION

production@msp.org

Silvio Levy, Scientific Editor

See inside back cover or msp.org/apde for submission instructions.

The subscription price for 2023 is US \$405/year for the electronic version, and \$630/year (+\$65, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues from the last three years and changes of subscriber address should be sent to MSP.

Analysis & PDE (ISSN 1948-206X electronic, 2157-5045 printed) at Mathematical Sciences Publishers, 798 Evans Hall #3840, c/o University of California, Berkeley, CA 94720-3840, is published continuously online.

APDE peer review and production are managed by EditFlow[®] from MSP.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2023 Mathematical Sciences Publishers

ANALYSIS & PDE

Volume 16 No. 1 2023

Volume comparison with respect to scalar curvature WEI YUAN	1
Wandering domains arising from Lavaurs maps with siegel disks MATTHIEU ASTORG, LUKA BOC THALER and HAN PETERS	35
Gaussian analytic functions of bounded mean oscillation ALON NISHRY and ELLIOT PAQUETTE	89
Generic KAM Hamiltonians are not quantum ergodic SEÁN GOMES	119
Strichartz estimates for mixed homogeneous surfaces in three dimensions LJUDEVIT PALLE	173
Holomorphic factorization of mappings into $Sp_4(\mathbb{C})$ BJÖRN IVARSSON, FRANK KUTZSCHEBAUCH and ERIK LØW	233
Transversal families of nonlinear projections and generalizations of Favard length ROSEMARIE BONGERS and KRYSTAL TAYLOR	279