

ANALYSIS & PDE

Volume 16

No. 3

2023

EDUARDO GARCÍA-JUÁREZ, YOICHIRO MORI AND ROBERT M. STRAIN

THE PESKIN PROBLEM WITH VISCOSITY CONTRAST



THE PESKIN PROBLEM WITH VISCOSITY CONTRAST

EDUARDO GARCÍA-JUÁREZ, YOICHIRO MORI AND ROBERT M. STRAIN

The Peskin problem models the dynamics of a closed elastic filament immersed in an incompressible fluid. We consider the case when the inner and outer viscosities are possibly different. This viscosity contrast adds further nonlocal effects to the system through the implicit nonlocal relation between the net force and the free interface. We prove the first global well-posedness result for the Peskin problem in this setting. The result applies for *medium-size* initial interfaces in critical spaces and shows instant analytic smoothing. We carefully calculate the medium-size constraint on the initial data. These results are new even without viscosity contrast.

1. Introduction	785
2. Linearization around the steady state	794
3. A priori estimates	806
4. Proof of main theorem	828
Acknowledgements	836
References	836

1. Introduction

Fluid structure interaction (FSI) problems in which an elastic structure interacts with a surrounding fluid are found in many areas of science and engineering. Many numerical algorithms have been developed for such problems, and the scientific computing of FSI problems continues to be a very active area of research [Li and Ito 2006; Peskin 2002; Tryggvason et al. 2001; Richter 2017]. The *Peskin problem*, considered in this paper, is arguably one of the simplest FSI problems and has been used extensively in physical modeling as well as in the development of numerical algorithms as a prototypical test problem.

1A. Formulation. Consider the following fluid problem in \mathbb{R}^2 . A closed elastic string Γ encloses a simply connected bounded domain $\Omega_1 \subset \mathbb{R}^2$ filled with a Stokes fluid with viscosity μ_1 . The outside region $\Omega_2 = \mathbb{R}^2 \setminus (\Omega_1 \cup \Gamma)$ is filled with a Stokes fluid of viscosity μ_2 . The equations satisfied are

$$\mu_1 \Delta \mathbf{u} - \nabla p = \mathbf{0} \quad \text{in } \Omega_1, \quad (1-1)$$

$$\mu_2 \Delta \mathbf{u} - \nabla p = \mathbf{0} \quad \text{in } \Omega_2, \quad (1-2)$$

$$\nabla \cdot \mathbf{u} = 0 \quad \text{in } \mathbb{R}^2 \setminus \Gamma. \quad (1-3)$$

Here \mathbf{u} is the velocity field and p is the pressure.

MSC2020: primary 35Q35, 35C15, 35R11, 35R35, 76D07; secondary 35C10.

Keywords: Peskin problem, fluid-structure interface, viscosity contrast, global regularity, critical regularity, immersed boundary problem, Stokes flow, fractional Laplacian, solvability, stability.

We must specify the interface conditions at Γ . Parametrize Γ by the material or Lagrangian coordinate $\theta \in \mathbb{S} = \mathbb{R}/(2\pi\mathbb{Z})$, and let $\mathcal{X}(\theta, t)$ denote the coordinate position of Γ at time t . The parametrization is in the counterclockwise direction, so that the interior region Ω_1 is on the left-hand side of the tangent vector $\partial\mathcal{X}/\partial\theta$. For any quantity w defined on Ω_1 and Ω_2 , we set

$$[w] = w|_{\Gamma_1} - w|_{\Gamma_2},$$

where $w|_{\Gamma_1}$ and $w|_{\Gamma_2}$ are the trace values of w at Γ evaluated from Ω_1 (interior) and Ω_2 (exterior) sides of Γ . Let \mathbf{n} be the outward-pointing unit normal vector on Γ :

$$\mathbf{n} = -\frac{\partial_\theta \mathcal{X}^\perp}{|\partial_\theta \mathcal{X}|}, \quad \partial_\theta \mathcal{X} = \frac{\partial \mathcal{X}}{\partial \theta}, \quad \partial_\theta \mathcal{X}^\perp = \mathcal{R} \partial_\theta \mathcal{X}, \quad \mathcal{R} = \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix},$$

where \mathcal{R} is the $\frac{\pi}{2}$ -rotation matrix. The interface conditions are

$$\frac{\partial \mathcal{X}}{\partial t} = \mathbf{u}(\mathcal{X}, t), \tag{1-4}$$

$$[\mathbf{u}] = 0, \tag{1-5}$$

$$[\Sigma \mathbf{n}] = \mathbf{F}_{\text{el}} |\partial_\theta \mathcal{X}|^{-1}, \quad \Sigma = \begin{cases} \mu_1 (\nabla \mathbf{u} + (\nabla \mathbf{u})^\text{T}) - pI & \text{in } \Omega_1, \\ \mu_2 (\nabla \mathbf{u} + (\nabla \mathbf{u})^\text{T}) - pI & \text{in } \Omega_2, \end{cases} \tag{1-6}$$

where I is the 2×2 identity matrix. The first condition is the no-slip boundary condition and the second is the stress balance condition, where Σ is the fluid stress and \mathbf{F}_{el} is the elastic force exerted by the string Γ . We let

$$\mathbf{F}_{\text{el}} = k_0 \partial_\theta^2 \mathcal{X}, \quad k_0 > 0, \tag{1-7}$$

where k_0 is the elasticity constant of the string Γ .

In the far field, $\mathbf{x} \rightarrow \infty$, we impose the condition that $\mathbf{u} \rightarrow 0$ and $p \rightarrow 0$. This completes the specification of the Peskin problem.

Let us rewrite the above problem using boundary integral equations. Given some function \mathbf{F} defined on Γ , we express the solution to our problem as the following single-layer potential on $\mathbb{S} = [-\pi, \pi]$:

$$\mathbf{u}(\mathbf{x}, t) = \int_{\mathbb{S}} G(\mathbf{x} - \mathcal{X}(\eta)) \mathbf{F}(\eta) d\eta, \tag{1-8}$$

$$G(\mathbf{x}) = \frac{1}{4\pi} \left(-\log|\mathbf{x}|I + \frac{\mathbf{x} \otimes \mathbf{x}}{|\mathbf{x}|^2} \right), \quad \mathbf{x} = (x_1, x_2)^\text{T} \in \mathbb{R}^2, \tag{1-9}$$

where G is the stokeslet, the fundamental solution of the two-dimensional Stokes problem. Additionally for $\mathbf{y} = (y_1, y_2)^\text{T} \in \mathbb{R}^2$ we use the notation

$$\mathbf{x} \otimes \mathbf{y} = \begin{bmatrix} x_1 y_1 & x_1 y_2 \\ x_2 y_1 & x_2 y_2 \end{bmatrix}.$$

We note that \mathcal{X} and \mathbf{F} (and other variables) depend on t , but we will often suppress this dependence to avoid cluttered notation. We note that the single-layer potential does not have a velocity jump across the

interface, and thus the boundary condition (1-5) is automatically satisfied. We then have

$$\frac{\partial \mathcal{X}}{\partial t}(\theta) = \int_{\mathbb{S}} G(\Delta \mathcal{X}) \mathbf{F}(\eta) d\eta, \tag{1-10}$$

where we use the notation

$$\Delta \mathcal{X} = \mathcal{X}(\theta) - \mathcal{X}(\eta).$$

On the other hand, the stress interface condition (1-6) is not automatically satisfied, and this will lead to an equation for \mathbf{F} . Let us compute the stress associated with the single-layer expression (1-8). The stress Σ in Ω_2 is given by

$$\Sigma_{ij}(\mathbf{x}) = \mu_2 \int_{\mathbb{S}} \mathcal{T}_{ijk}(\mathbf{x} - \mathcal{X}(\eta)) F_k(\eta) d\eta,$$

with

$$\mathcal{T}_{ijk} = -\frac{1}{\pi} \frac{x_i x_j x_k}{|\mathbf{x}|^4}, \tag{1-11}$$

where the subscripts denote the components of the respective tensors/vectors, such as $\mathbf{F} = (F_1, F_2)^T$, and the summation convention is in effect for repeated indices. We refer to Chapter 2 of [Pozrikidis 1992] for further details on the derivation of the stokeslet and the stresslet tensors. In Ω_1 , the stress is given by

$$\Sigma_{ij}(\mathbf{x}, t) = \mu_1 \int_{\mathbb{S}} \mathcal{T}_{ijk}(\mathbf{x} - \mathcal{X}(\eta)) F_k(\eta) d\eta.$$

Thus, the trace values of the normal stresses are given by the equations

$$\begin{aligned} \Sigma_{ij}(\mathcal{X}(\theta)) n_j(\theta)|_{\Gamma_2} &= \mu_2 \left(-\frac{1}{2} F_i |\partial_\theta \mathcal{X}|^{-1} + \text{pv} \int_{\mathbb{S}} \mathcal{T}_{ijk}(\Delta \mathcal{X}) F_k(\eta) n_j(\theta) d\eta \right), \\ \Sigma_{ij}(\mathcal{X}(\theta)) n_j(\theta)|_{\Gamma_1} &= \mu_1 \left(\frac{1}{2} F_i |\partial_\theta \mathcal{X}|^{-1} + \text{pv} \int_{\mathbb{S}} \mathcal{T}_{ijk}(\Delta \mathcal{X}) F_k(\eta) n_j(\theta) d\eta \right). \end{aligned}$$

The stress jump condition (1-6) thus reduces to (for $i = 1, 2$)

$$F_i(\theta) + 2A_\mu \int_{\mathbb{S}} \mathcal{T}_{ijk}(\Delta \mathcal{X}) F_k(\eta) \partial_\theta \mathcal{X}_j^\perp(\theta) d\eta = \frac{2}{\mu_1 + \mu_2} F_{\text{el},i}(\theta),$$

where

$$A_\mu = \frac{\mu_2 - \mu_1}{\mu_1 + \mu_2}. \tag{1-12}$$

We define

$$\mathcal{S}_i(\mathbf{F}, \mathcal{X})(\theta) = -\partial_\theta \mathcal{X}_j^\perp(\theta) \int_{\mathbb{S}} \mathcal{T}_{ijk}(\Delta \mathcal{X}) F_k(\eta) d\eta.$$

We will frequently write it in vector notation as

$$\mathbf{F}(\theta) = 2A_\mu \mathcal{S}(\mathbf{F}, \mathcal{X})(\theta) + 2A_e \tilde{\mathbf{F}}_{\text{el}}(\theta), \tag{1-13}$$

where

$$\mathcal{S}(\mathbf{F}, \mathcal{X})(\theta) = -\partial_\theta \mathcal{X}(\theta)^\perp \cdot \int_{\mathbb{S}} \mathcal{T}(\mathcal{X}(\theta) - \mathcal{X}(\eta)) \cdot \mathbf{F}(\eta) d\eta, \tag{1-14}$$

with

$$\tilde{\mathbf{F}}_{\text{el}} = \frac{1}{k_0} \mathbf{F}_{\text{el}}, \quad A_e = \frac{k_0}{\mu_2 + \mu_1}. \tag{1-15}$$

We point out that the above boundary integral equation has a unique solution \mathbf{F} given \mathbf{F}_{el} for sufficiently smooth \mathcal{X} .

The Peskin problem thus reduces to the integral equations (1-10) and (1-13) for \mathcal{X} , where $G, \mathcal{T}, A_\mu, \mathcal{S}, A_e$, and $\tilde{\mathbf{F}}_{\text{el}}$ are given by (1-9), (1-11), (1-12), (1-14), and (1-15), with \mathbf{F}_{el} given by (1-7). Note also that, when $A_\mu = 0$, i.e., $\mu_1 = \mu_2$, equation (1-13) reduces to $\mathbf{F} = 2A_e \mathbf{F}_{\text{el}}$, and we may just work with the single equation (1-10).

Assuming that the stationary solutions are sufficiently smooth, it can be shown by an easy calculation that the only stationary solutions are those in which \mathcal{X} is a uniformly parametrized circle and the velocity field is $\mathbf{u} = 0$; see Section 5.1 of [Mori et al. 2019]. Thus, all of the equilibrium configurations of (1-10) and (1-13) are spanned by

$$\mathbf{e}_r(\theta) = \begin{bmatrix} \cos \theta \\ \sin \theta \end{bmatrix}, \quad \mathbf{e}_t(\theta) = \begin{bmatrix} -\sin \theta \\ \cos \theta \end{bmatrix}, \quad \mathbf{e}_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}, \quad \mathbf{e}_2 = \begin{bmatrix} 0 \\ 1 \end{bmatrix}. \tag{1-16}$$

1B. Critical regularity and related results. A general guideline for seeking the most natural and largest class of initial data for a given problem is to identify its scaling and consider a function space that is critical (invariant) with respect to this scaling. The Peskin problem given above by (1-10) and (1-13) is invariant under dilation, and thus to make proper sense of scaling one must first fix a reference scale. Consider the scaling parameter $\lambda > 0$. The domain scales accordingly from the torus $\mathbb{S} = [-\pi, \pi]$ to $\mathbb{S}/\lambda = [-\pi/\lambda, \pi/\lambda]$. Then, we choose as the reference scale the length of uniformly parametrized circles, which we pick to be 2π . Given the additional rotation and translation invariance of the problem, let us consider the particular choice

$$\mathbf{X}_{*,\lambda}(\theta) = \lambda^{-1} \mathbf{X}_*(\lambda\theta),$$

where $\mathbf{X}_*(\theta) = \mathbf{e}_r(\theta)$. Then, the system (1-10), (1-13) is written in terms of the difference $\mathbf{X}(\theta, t) = \mathcal{X}(\theta, t) - \mathbf{X}_*(\theta)$. One can check that the following dilation invariance holds: if $\mathbf{X}(\theta, t)$ is a solution, then $\mathbf{X}_\lambda(\theta, t) = \lambda^{-1} \mathbf{X}(\lambda\theta, \lambda t)$ is also a solution.

More generally, if the elastic force \mathbf{F}_{el} is given by (1-7), then (1-10) has an additional scaling invariance given by $\mathbf{X}_{\lambda,\tau}(\theta, t) = \tau \mathbf{X}(\lambda\theta, \lambda t)$ and $\mathbf{X}_{*,\lambda,\tau}(\theta) = \tau \mathbf{X}_*(\lambda\theta)$ for any $\lambda, \tau > 0$. The stress jump condition (1-13) then scales as $\mathbf{F}_{\lambda,\tau}(\theta, t) = \lambda^2 \tau \mathbf{F}(\lambda\theta, \lambda t)$. This more general scaling leaves the equation invariant with τ unrelated to λ . We note however that the chord arc condition, defined below in (1-17), is only invariant under the dilation rescaling where $\tau = \lambda^{-1}$.

The analytical study of the Peskin problem was initiated in [Lin and Tong 2019; Mori et al. 2019], in which the case of equal viscosity $\mu_1 = \mu_2$ was studied. In [Lin and Tong 2019], well-posedness was established in $\mathcal{X} \in C([0, T]; H^{5/2}(\mathbb{S}))$, $T > 0$, with initial data \mathcal{X}_0 in $H^{5/2}(\mathbb{S})$, whereas in [Mori et al. 2019], the solution resides in $\mathcal{X} \in C([0, T]; C^{1,\alpha}(\mathbb{S}))$, $\alpha > 0$, $T > 0$, with initial data \mathcal{X}_0 in $h^{1,\alpha}(\mathbb{S})$, $\alpha > 0$ (this space is the completion of smooth functions in the $C^{1,\alpha}$ norm). These spaces are subcritical with respect to the above scaling. Indeed, in the L^2 Sobolev scale, $H^{3/2}(\mathbb{S})$ (or $C([0, T]; H^{3/2}(\mathbb{S}))$) is the critical space, whereas in the scale of (Hölder) continuous functions, $C^1(\mathbb{S})$ (or $C([0, T]; C^1(\mathbb{S}))$) is the critical scale. In this sense, the results in [Mori et al. 2019] are only barely subcritical. The semilinear parabolic methods [Lunardi 1995] that are used in [Mori et al. 2019] rely crucially on subcriticality, however, and do not seem to be readily extendible to the critical regularity exponent.

In this paper, we consider the Peskin problem in which the viscosities μ_1 and μ_2 are not necessarily equal. Furthermore, we establish a solution theory with initial data \mathcal{X}_0 in the Wiener space $\mathcal{F}^{1,1}(\mathbb{S})$, the space of functions whose derivatives have a Fourier series that is absolutely summable (see Section 1C). This space is critical with respect to the scaling of the Peskin problem identified above.

In contrast to [Lin and Tong 2019; Mori et al. 2019], our theory is restricted to initial data that is sufficiently close to the stationary states, i.e., the uniformly parametrized circles. The papers [Lin and Tong 2019; Mori et al. 2019] establish local-in-time well-posedness in their respective function spaces subject to the following arc-chord condition on the initial data:

$$|\mathcal{X}_0|_* \equiv \inf_{\theta, \eta \in \mathbb{S}, \theta \neq \eta} \frac{|\mathcal{X}_0(\theta) - \mathcal{X}_0(\eta)|}{|\theta - \eta|} > 0. \tag{1-17}$$

In this sense, our results might be better compared to the results on asymptotic stability of the uniformly parametrized circle obtained in [Lin and Tong 2019; Mori et al. 2019]. The uniformly parametrized circle is proved to be exponentially stable in the above L^2 Sobolev and Hölder scales respectively, and in the latter paper, it is proved that the solution is in $C^\infty(\mathbb{S})$ for all positive time. In this paper, we improve upon this result to prove that the solution is analytic for positive time.

Local-in-time well-posedness for initial data in $\mathcal{F}^{1,1}$ merely satisfying condition (1-17) is an open question that we do not address in this paper. It is notable, however, that the arc-chord condition (1-17) is invariant under the dilation scaling described above. In [Mori et al. 2019], it is shown that, if the solution ceases to exist as t approaches $t_* < \infty$, then following must hold:

$$\lim_{t \rightarrow t_*} \varrho_\alpha(\mathcal{X}) = \infty, \quad \varrho_\alpha(\mathcal{X}) = \frac{\|\partial_\theta \mathcal{X}\|_{C^\alpha}}{|\mathcal{X}|_*} \quad \text{for any } \alpha > 0.$$

On the other hand, if $\varrho_\alpha(\mathcal{X})$ remains bounded for all time for some $\alpha > 0$, then \mathcal{X} must converge to a uniformly parametrized circle. A similar criterion, in which the numerator of ϱ_α is replaced with a critical norm such as the $\mathcal{F}^{1,1}$ norm, would be a major improvement that should lead to a better understanding of the global-in-time dynamics of the Peskin problem.

Another extension of the Peskin problem is to consider the following elastic force in place of (1-7):

$$\mathbf{F}_{\text{el}} = \partial_\theta \left(\mathcal{T}(|\partial_\theta \mathcal{X}|) \frac{\partial_\theta \mathcal{X}}{|\partial_\theta \mathcal{X}|} \right), \tag{1-18}$$

where $\mathcal{T}(s)$ is a tension coefficient that must satisfy the structure condition $\mathcal{T} > 0$ and $d\mathcal{T}/ds > 0$. Note that the above expression is reduced to (1-7) if we take $\mathcal{T}(s) = k_0 s$, hence $k_0 = \mathcal{T}(1) = d\mathcal{T}/ds$. In the case of equal viscosity $\mu_1 = \mu_2$, a local-in-time well-posedness theory for initial data satisfying (1-17) under the more general force (1-18) is established in [Rodenberg 2018] in the Hölder scale similarly to [Mori et al. 2019], using nonlinear parabolic methods [Lunardi 1995]. It is expected that the results and methods of this paper can be extended to this more general case.

Finally, we mention [Tong 2021] in which the author considers a regularization of the Peskin problem inspired by the immersed boundary method, extending the techniques in [Lin and Tong 2019]. Such studies may form the basis for numerical analysis of the Peskin problem.

The surface tension problem, in which the interface is not elastic but only exerts a surface tension, may be the most closely related class of problems for which there are extensive analytical studies. We note that our problem is distinct from the surface tension problem; in contrast to an elastic interface considered in the Peskin problem, an interface with surface tension only does not resist stretching. This difference manifests itself in the different energy dissipation laws satisfied by the respective problems; see Section 1.1 of [Mori et al. 2019]. We refer the reader to [Prüss and Simonett 2009; 2016; Shimizu 2009] for an extensive survey of the analytical study of the surface tension problem.

There is also an increasing number of analytical studies on fluid-structure interaction problems in which an elastic structure interacts with a fluid, related to the Peskin problem considered here [Ambrose and Siegel 2017; Cheng et al. 2007; Cheng and Shkoller 2010; Liu and Ambrose 2017; Muha and Canić 2013; Plotnikov and Toland 2011; 2012; Li 2021; Boulakia et al. 2012]. The equations dealt with in these studies are typically more complicated than those of the Peskin problem; the sharp results obtained for the simpler Peskin problem should serve as a guide to what is possibly true for the more complicated model problems.

From an analytical perspective, the Muskat problem is perhaps the closest nonlinear PDE to our problem for which there is a large body of analytical studies. However, it models a very different physical setting: two immiscible and incompressible fluids in a porous media governed by Darcy's law. On the other hand, for a nearly flat interface in the presence of gravity both problems have the same symbol at the linear level. The authors of [Constantin et al. 2013] introduced the use of the Wiener algebra to obtain global well-posedness results for the Muskat problem at critical regularity. Moreover, the size restriction on the initial data was given by an explicit constant that is independent of any physical parameter. These techniques were extended in [Constantin et al. 2016; Gancedo et al. 2019a] to deal with the three-dimensional setting and the case of viscosity jumps, respectively. Other results for the Muskat problem that only require *medium-size* initial data in critical spaces (as opposed to the more standard arbitrarily small data condition) [Cameron 2019; 2020] rely on the maximum principle; these methods have thus far not been shown to be well-suited to deal with viscosity contrasts.

In this paper, we will use spaces related to the Wiener algebra that allow us to perform careful and detailed estimates on the nonlinear terms to control explicitly the size constraint on the initial data (see Figure 1). As opposed to the Muskat problem, here the problem is not only described by the shape of the interface: the parametrization corresponds to the distribution of material points, and thus it matters. As a consequence, we have to develop further techniques to deal with a system of equations (for both components of the curve). Interestingly, a careful understanding of the linear system, together with an appropriate change of framework, allows us to decouple the frequencies associated to the projection of the interface onto the space of equilibria from the others. Indeed, we overcome a major difficulty of the very recent result in [Gancedo et al. 2019b] that deals with the Muskat problem for closed interfaces (i.e., bubbles), and obtain the global existence and uniqueness result for the Peskin problem with viscosity contrast at critical regularity.

1C. Notation and functional spaces. We summarize here the notation and functional spaces that will be used throughout the paper.

For a vector $\mathbf{x} = (x_1, x_2)^T \in \mathbb{C}^2$ we define

$$\mathbf{x}^\perp \stackrel{\text{def}}{=} \mathcal{R}\mathbf{x}, \quad \mathcal{R} \stackrel{\text{def}}{=} \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix}, \quad \mathcal{R}^{-1} = \begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix}.$$

We denote the Euclidean norm as

$$|\mathbf{x}| = \sqrt{\mathbf{x}^T \bar{\mathbf{x}}} = \sqrt{|x_1|^2 + |x_2|^2},$$

and for a matrix $A = (a_{ij})_{1 \leq i, j \leq 2}$ we use the induced matrix norm

$$\|A\| = \sigma_{\max}(A), \tag{1-19}$$

where $\sigma_{\max}(A)$ is the largest singular value of A . For a vector such as \mathbf{X}_* we will write $X_{*,j}$ to be the j -th component of that vector.

We now define the periodic Hilbert transform of a function f with period $2P$ as

$$\mathcal{H}(f)(\theta) \stackrel{\text{def}}{=} \frac{1}{2P} \text{pv} \int_{-P}^P \frac{f(\theta - \eta)}{\tan\left(\frac{\eta}{2P/\pi}\right)} d\eta = \frac{1}{4P} \text{pv} \int_{-P}^P \frac{f(\theta - \eta) - f(\theta + \eta)}{\tan\left(\frac{\eta}{2P/\pi}\right)} d\eta. \tag{1-20}$$

Unless stated otherwise, throughout the paper we will use the case $P = \pi$. In this case, we also define the Fourier transform of a periodic function f with domain $\mathbb{S} = [-\pi, \pi]$ as

$$\mathcal{F}(f)(k) \stackrel{\text{def}}{=} \hat{f}(k) = \frac{1}{2\pi} \int_{-\pi}^{\pi} f(\theta) e^{-ik\theta} d\theta, \quad k \in \mathbb{Z}.$$

Further $\mathcal{F}(\mathcal{H}(f))(k) = -i \operatorname{sgn}(k) \hat{f}(k)$. Then we define the operator Λ using the Fourier transform as $\mathcal{F}(\Lambda f)(k) \stackrel{\text{def}}{=} |k| \hat{f}(k)$. And we observe that $\mathcal{H}(\partial_\theta f)(\theta) = \Lambda f$.

We denote by $f * g$ the standard convolution of f and g . We use the iterated convolution notation

$$*^k f = \underbrace{f * \dots * f}_{k-1 \text{ convolutions of } k \text{ copies of } f} \tag{1-21}$$

Thus for instance $*^2 f = f * f$.

We also use the following notation for the discrete delta function, $\delta_a(k)$, which is the function that is equal to 1 when $k = a$ and equal to 0 elsewhere. Throughout the paper we will further define

$$\delta_{1,-1}(k) = \delta_1(k) + \delta_{-1}(k). \tag{1-22}$$

We further define the high-frequency cut-off operator \mathcal{J}_M for $M \geq 0$ by

$$\widehat{\mathcal{J}_M X}(k) \stackrel{\text{def}}{=} 1_{|k| \leq M} \widehat{X}(k), \tag{1-23}$$

where 1_A is the standard indicator function of the set A , so that $1_A(x) = 1$ if $x \in A$ and $1_A(x) = 0$ if $x \notin A$.

For two vectors $\mathbf{X}(\theta), \mathbf{Y}(\theta) \in \mathbb{R}^2$ we define

$$\langle \mathbf{X}, \mathbf{Y} \rangle = \int_{\mathbb{S}} \mathbf{X}(\theta) \cdot \mathbf{Y}(\theta) d\theta. \tag{1-24}$$

Generalizing the Wiener algebra of functions with absolutely convergent Fourier series as in [Gancedo et al. 2019b], we further define the homogeneous $\dot{\mathcal{F}}_v^{s,1}$ and nonhomogeneous $\mathcal{F}_v^{s,1}$ norms as

$$\|\mathbf{X}\|_{\dot{\mathcal{F}}_v^{s,1}} = \sum_{k \in \mathbb{Z} \setminus \{0\}} e^{\nu(t)|k|} |k|^s |\widehat{\mathbf{X}}(k)|, \quad s \in \mathbb{R}, \tag{1-25}$$

$$\|\mathbf{X}\|_{\mathcal{F}_v^{s,1}} = |\widehat{\mathbf{X}}(0)| + \sum_{k \in \mathbb{Z} \setminus \{0\}} e^{\nu(t)|k|} |k|^s |\widehat{\mathbf{X}}(k)|, \quad s \geq 0, \tag{1-26}$$

with

$$\nu(t) = \nu_\infty \frac{t}{1+t} \geq 0, \tag{1-27}$$

and $\nu_\infty > 0$ is chosen sufficiently small. Note that $\nu(0) = 0$, $\nu(t) > 0$ for all $t > 0$. Further $\nu'(t) \leq \nu_\infty$ and $\nu(t) \leq \nu_\infty$ are bounded for all time. When $\nu \equiv 0$, we write $\dot{\mathcal{F}}_0^{s,1} = \dot{\mathcal{F}}^{s,1}$ and $\mathcal{F}_0^{s,1} = \mathcal{F}^{s,1}$. These are the main norms that we will use in this paper. Note that when $s = 1$, the $\dot{\mathcal{F}}^{s,1}$ norm is critical for the Peskin problem.

In this paper we write $A \lesssim B$ if $A \leq CB$ for some inessential constant $C > 0$. We also write $A \approx B$ if both $A \lesssim B$ and $B \lesssim A$ hold. Throughout the paper, we will define

$$C_i = C_i(\|\mathbf{X}\|_{\dot{\mathcal{F}}_v^{1,1}}) = C_i(\|\mathbf{X}\|_{\dot{\mathcal{F}}_v^{1,1}}; \nu_\infty) > 0, \quad i = 1, 2, \dots, \tag{1-28}$$

as functions that are increasing in $\|\mathbf{X}\|_{\dot{\mathcal{F}}_v^{1,1}} \geq 0$ and might depend on the analyticity constant ν_∞ , with the properties that $C_i(\|\mathbf{X}\|_{\dot{\mathcal{F}}_v^{1,1}}) \approx 1$ for all $\nu_\infty \geq 0$ and $\lim_{\|\mathbf{X}\|_{\dot{\mathcal{F}}_v^{1,1}} \rightarrow 0^+} C_i(\|\mathbf{X}\|_{\dot{\mathcal{F}}_v^{1,1}}; 0) = 1$. We will also define

$$D_i = D_i(\|\mathbf{X}\|_{\dot{\mathcal{F}}_v^{1,1}}) = D_i(\|\mathbf{X}\|_{\dot{\mathcal{F}}_v^{1,1}}; A_\mu, \nu_\infty) > 0, \quad i = 1, 2, \dots,$$

as functions that are increasing in $\|\mathbf{X}\|_{\dot{\mathcal{F}}_v^{1,1}} \geq 0$ and might depend on the physical parameter A_μ and the analyticity constant ν_∞ , with the properties that $D_i(\|\mathbf{X}\|_{\dot{\mathcal{F}}_v^{1,1}}) \approx 1$ for all $A_\mu \in (-1, 1)$ and all $\nu_\infty \geq 0$, and $\lim_{\|\mathbf{X}\|_{\dot{\mathcal{F}}_v^{1,1}} \rightarrow 0^+} D_i(\|\mathbf{X}\|_{\dot{\mathcal{F}}_v^{1,1}}; 0, 0) = 1$.

1D. Main results. In this section we will state the main result of this paper: namely, that membranes whose initial interface has critical regularity (in terms of the scaling of the problem), and that are not too far from an equilibrium configuration, become instantaneously analytic and converge exponentially fast to the equilibrium. Without loss of generality, we assume that the initial area enclosed by the membrane is π . We get the result under an explicit medium-size condition for the initial deviation and for general viscosity contrast $A_\mu \in (-1, 1)$.

Definition 1.1 (strong solution). Let

$$\mathcal{X} \in C([0, T]; \mathcal{F}^{1,1}) \cap C^1((0, T]; \mathcal{F}^{0,1})$$

and

$$|\mathcal{X}|_*(t) = \inf_{\theta, \eta \in \mathbb{S}, \theta \neq \eta} \frac{|\mathcal{X}(\theta, t) - \mathcal{X}(\eta, t)|}{|\theta - \eta|} > 0$$

for $0 \leq t \leq T$. Then, \mathcal{X} is a *strong solution* to the viscosity-contrast Peskin problem with initial value $\mathcal{X}(0) = \mathcal{X}_0$ if it satisfies (1-10), (1-13) for $0 < t \leq T$ and $\mathcal{X}(t) \rightarrow \mathcal{X}_0$ in $\mathcal{F}^{1,1}$ as $t \rightarrow 0$.

Theorem 1.2 (main result). *Let $A_\mu \in (-1, 1)$ and $\mathcal{X}_0 \in \mathcal{F}^{1,1}$. Let $\mathbf{X}_{0,c}$ be the projection of \mathcal{X}_0 onto the vector space spanned by (1-16) and $\mathbf{X}_0 = \mathcal{X}_0 - \mathbf{X}_{0,c}$; thus \mathbf{X}_0 is mean zero and $\widehat{\mathcal{X}}_0(0) = \widehat{\mathbf{X}}_{0,c}(0)$. Assume that initially the deviation \mathbf{X}_0 satisfies the medium-size condition*

$$\|\mathbf{X}_0\|_{\dot{\mathcal{F}}^{1,1}} < k(A_\mu), \tag{1-29}$$

where $k(A_\mu) > 0$ is defined in (4-9) (see also (4-10) and Figure 1), and that the area enclosed by \mathcal{X}_0 is π . Then, for any $T > 0$, there exists a constant $\nu_\infty > 0$ such that there exists a **unique global strong solution** $\mathcal{X}(t)$ to the system (1-10) and (1-13), which lies in the space

$$\mathcal{X} \in C([0, T]; \mathcal{F}_\nu^{1,1}) \cap C^1((0, T]; \mathcal{F}_\nu^{0,1}) \cap L^1([0, T]; \dot{\mathcal{F}}_\nu^{2,1}),$$

with $\nu(t)$ given by (1-27). In particular, it becomes instantaneously analytic. Moreover, the following energy inequality is satisfied for $0 \leq t \leq T$:

$$\|\mathbf{X}\|_{\dot{\mathcal{F}}_\nu^{1,1}}(t) + \frac{A_e}{4} \mathcal{C} \int_0^t \|\mathbf{X}\|_{\dot{\mathcal{F}}_\nu^{2,1}}(\tau) d\tau \leq \|\mathbf{X}_0\|_{\dot{\mathcal{F}}^{1,1}}, \tag{1-30}$$

with $\mathcal{C} = \mathcal{C}(\|\mathbf{X}_0\|_{\dot{\mathcal{F}}^{1,1}}, A_\mu, \nu_\infty) > 0$ defined in (4-12). In addition,

$$\|\mathbf{X}\|_{\dot{\mathcal{F}}_\nu^{1,1}}(t) \leq \|\mathbf{X}_0\|_{\dot{\mathcal{F}}^{1,1}} e^{-(A_e/4)\mathcal{C}t}. \tag{1-31}$$

The zero frequency $\widehat{\mathbf{X}}_c(0)$ remains uniformly bounded for all times as

$$|\widehat{\mathbf{X}}_c(0)| \leq |\widehat{\mathbf{X}}_{0,c}(0)| + \widetilde{\mathcal{C}} \|\mathbf{X}_0\|_{\dot{\mathcal{F}}^{1,1}}^2,$$

with $\widetilde{\mathcal{C}} = \widetilde{\mathcal{C}}(\|\mathbf{X}_0\|_{\dot{\mathcal{F}}^{1,1}}, A_\mu) > 0$ given in (4-15), while

$$1 - \frac{1}{2} \|\mathbf{X}\|_{\dot{\mathcal{F}}_\nu^{1,1}}^2 \leq |\widehat{\mathbf{X}}_c(1)|^2 \leq 1 + \frac{1}{2} \|\mathbf{X}\|_{\dot{\mathcal{F}}_\nu^{1,1}}^2. \tag{1-32}$$

We remark that the decay to zero of the deviation \mathbf{X} in (1-31) together with (1-32) shows the exponentially fast convergence to a uniformly parametrized circle with the same area as the initial one.

Remark 1.3. The size of $\nu_\infty > 0$ is limited by the size of the initial data. This can be seen in (4-7). Because we are only interested in having any fixed but arbitrarily small ν_∞ to ensure analyticity, we stated the size condition as in (1-29).

Remark 1.4. In our results, we assume that both viscosities μ_1 and μ_2 are positive and hence $-1 < A_\mu < 1$. We remark on the endpoint cases of $A_\mu = \pm 1$, which formally correspond to the cases when $\mu_1 = 0$ or $\mu_2 = 0$. As can be seen from Figure 1, the allowed size of the deviation from \mathbf{X}_0 tends to 0 as $A_\mu \rightarrow \pm 1$, which may indicate potential difficulties in formulating a well-posed mathematical problem for the endpoint cases. From a physical standpoint, it does not make sense to set the viscosity to 0 in either Ω_1 or Ω_2 , and thus a proper treatment of these endpoint cases will require a rethinking of the physical situation under consideration. The case $A_\mu = -1$ or $\mu_2 = 0$ may be thought of as corresponding to the problem in which a droplet of Stokesian fluid is floating in vacuum. One significant difference between this and the Peskin problem is that in the former problem a droplet in linear translation or rigid rotation experiences no external forces. The force balance and continuity equations will thus have to be




Figure 1. $10^3 k(A_\mu)$

supplemented by auxiliary conditions that assure uniqueness, after which this problem is likely to be well-posed. In the case $A_\mu = 1$ or $\mu_1 = 0$, Ω_1 might be considered to be vacuum. It is not clear if this problem is well-posed. We will not pursue these issues further.

1E. Outline. The rest of the paper is structured as follows. In [Section 2](#), we first decompose in [Section 2A](#) the system (1-10), (1-13) into zero-order, linear, and nonlinear parts around the equilibrium configuration, and then in [Section 2B](#) we perform the linearization of the problem and show its parabolic structure. [Section 2C](#) shows how this structure leads to dissipation and in [Section 2D](#) we summarize the system of equations in its final form. [Section 3](#) contains the crucial nonlinear estimates needed to prove [Theorem 1.2](#). Finally, [Section 4](#) is dedicated to the proof of [Theorem 1.2](#) via a regularization argument and also shows the uniqueness of the solutions.

2. Linearization around the steady state

We will linearize the system (1-9)-(1-15), with F_{cl} given by (1-7), around a time-dependent uniformly parametrized circle with center $(c(t), d(t))$ and radius $R(t)$:

$$\begin{aligned} X_c(\theta, t) &= a(t)\mathbf{e}_r(\theta) + b(t)\mathbf{e}_t(\theta) + c(t)\mathbf{e}_1 + d(t)\mathbf{e}_2, \\ R^2(t) &= a^2(t) + b^2(t), \end{aligned} \tag{2-1}$$

where $a(t)$, $b(t)$, $c(t)$ and $d(t)$ are arbitrary time-dependent functions and $\mathbf{e}_r(\theta)$, $\mathbf{e}_t(\theta)$, $\mathbf{e}_1(\theta)$, $\mathbf{e}_2(\theta)$ are defined in (1-16). For notational convenience, we will suppress the time dependence of the coefficients.

2A. Nonlinear expansion. We will denote by $X(\theta)$ the deviation from the circle $X_c(\theta)$ as $X(\theta) = \mathcal{X}(\theta) - X_c(\theta)$. We define further

$$\Delta X \stackrel{\text{def}}{=} X(\theta) - X(\eta)$$

and

$$\Delta_\eta \mathbf{X}(\theta) \stackrel{\text{def}}{=} \frac{\mathbf{X}(\theta) - \mathbf{X}(\eta)}{2 \sin\left(\frac{\theta-\eta}{2}\right)}. \tag{2-2}$$

In particular, we have

$$\Delta_\eta \mathbf{X}_c(\theta) = a\mathbf{e}_t\left(\frac{\theta+\eta}{2}\right) - b\mathbf{e}_r\left(\frac{\theta+\eta}{2}\right),$$

since

$$\begin{aligned} \Delta_\eta \mathbf{e}_r(\theta) &= \begin{bmatrix} -\sin\left(\frac{\theta+\eta}{2}\right) \\ \cos\left(\frac{\theta+\eta}{2}\right) \end{bmatrix} = \mathbf{e}_t\left(\frac{\theta+\eta}{2}\right), \\ \Delta_\eta \mathbf{e}_t(\theta) &= \begin{bmatrix} -\cos\left(\frac{\theta+\eta}{2}\right) \\ -\sin\left(\frac{\theta+\eta}{2}\right) \end{bmatrix} = -\mathbf{e}_r\left(\frac{\theta+\eta}{2}\right), \end{aligned}$$

where we have used the trigonometric identities

$$\begin{aligned} \sin(\theta) - \sin(\eta) &= 2 \sin\left(\frac{\theta-\eta}{2}\right) \cos\left(\frac{\theta+\eta}{2}\right), \\ \cos(\theta) - \cos(\eta) &= -2 \sin\left(\frac{\theta-\eta}{2}\right) \sin\left(\frac{\theta+\eta}{2}\right). \end{aligned}$$

Recalling (2-1) and using the identities

$$\begin{aligned} \partial_\theta \mathbf{e}_r(\theta) &= \mathbf{e}_r(\theta)^\perp = \mathbf{e}_t(\theta), \\ \partial_\theta \mathbf{e}_t(\theta) &= \mathbf{e}_t(\theta)^\perp = -\mathbf{e}_r(\theta), \end{aligned}$$

one has

$$\begin{aligned} \partial_\theta \mathbf{X}_c(\theta) &= a\mathbf{e}_t(\theta) - b\mathbf{e}_r(\theta), \\ \partial_\theta \mathbf{X}_c(\theta)^\perp &= -a\mathbf{e}_r(\theta) - b\mathbf{e}_t(\theta). \end{aligned}$$

The trigonometric identities $\cos(a+b) = \cos(a)\cos(b) - \sin(a)\sin(b)$ and $\sin(a+b) = \sin(a)\cos(b) + \cos(a)\sin(b)$ further give

$$\begin{aligned} \mathbf{e}_r(\theta) \cdot \mathbf{e}_t\left(\frac{\theta+\eta}{2}\right) &= \sin\left(\frac{\theta-\eta}{2}\right), & \mathbf{e}_t(\theta) \cdot \mathbf{e}_t\left(\frac{\theta+\eta}{2}\right) &= \cos\left(\frac{\theta-\eta}{2}\right), \\ \mathbf{e}_r(\theta) \cdot \mathbf{e}_r\left(\frac{\theta+\eta}{2}\right) &= \cos\left(\frac{\theta-\eta}{2}\right), & \mathbf{e}_t(\theta) \cdot \mathbf{e}_r\left(\frac{\theta+\eta}{2}\right) &= -\sin\left(\frac{\theta-\eta}{2}\right). \end{aligned}$$

These calculations imply the following computations for a circle that will be used frequently throughout the paper:

$$\partial_\theta \mathbf{X}_c(\theta)^\perp \cdot \Delta_\eta \mathbf{X}_c(\theta) = -R^2 \sin\left(\frac{\theta-\eta}{2}\right), \tag{2-3}$$

$$\partial_\theta \mathbf{X}_c(\theta) \cdot \Delta_\eta \mathbf{X}_c(\theta) = R^2 \cos\left(\frac{\theta-\eta}{2}\right), \tag{2-4}$$

$$\begin{aligned} \Delta_\eta \mathbf{X}_c(\theta) \otimes \Delta_\eta \mathbf{X}_c(\theta) &= \frac{a^2}{2} \begin{bmatrix} 1 - \cos(\theta+\eta) & -\sin(\theta+\eta) \\ -\sin(\theta+\eta) & 1 + \cos(\theta+\eta) \end{bmatrix} \\ &+ \frac{b^2}{2} \begin{bmatrix} 1 + \cos(\theta+\eta) & \sin(\theta+\eta) \\ \sin(\theta+\eta) & 1 - \cos(\theta+\eta) \end{bmatrix} + ab \begin{bmatrix} \sin(\theta+\eta) & -\cos(\theta+\eta) \\ -\cos(\theta+\eta) & -\sin(\theta+\eta) \end{bmatrix}. \end{aligned} \tag{2-5}$$

The matrices in the last line above have been simplified using the identities $\sin^2(a) = (1 - \cos(2a))/2$, $\cos^2(a) = (1 + \cos(2a))/2$, and $\sin(2a) = 2 \sin(a) \cos(a)$.

Next, we perform a Taylor expansion of the nonlinear terms around the a time-dependent uniformly parametrized circle (2-1) under the assumption that $|\Delta_\eta \mathbf{X}(\theta)| < 1$. First, we start with the magnitude of the curve

$$|\Delta \mathbf{X} + \Delta \mathbf{X}_c|^2 = 4R^2 \sin^2 \left(\frac{\theta - \eta}{2} \right) \left(1 + \frac{2}{R^2} \Delta_\eta \mathbf{X}_c(\theta) \cdot \Delta_\eta \mathbf{X}(\theta) + \frac{1}{R^2} |\Delta_\eta \mathbf{X}(\theta)|^2 \right).$$

Recalling the expression for $G(\Delta \mathcal{X})$ in (1-9), we expand each term as

$$\begin{aligned} \log |\Delta \mathbf{X} + \Delta \mathbf{X}_c| &= \log \left(2R \left| \sin \left(\frac{\theta - \eta}{2} \right) \right| \right) + \frac{1}{2} \log \left(1 + \frac{2}{R^2} \Delta_\eta \mathbf{X}_c(\theta) \cdot \Delta_\eta \mathbf{X}(\theta) + \frac{1}{R^2} |\Delta_\eta \mathbf{X}(\theta)|^2 \right) \\ &= \log \left(2R \left| \sin \left(\frac{\theta - \eta}{2} \right) \right| \right) + \frac{1}{R^2} \Delta_\eta \mathbf{X}_c(\theta) \cdot \Delta_\eta \mathbf{X}(\theta) + \mathcal{R}_1(\Delta_\eta \mathbf{X}(\theta)), \end{aligned} \quad (2-6)$$

where

$$\mathcal{R}_1(\Delta_\eta \mathbf{X}(\theta)) = \sum_{n=1}^{\infty} \sum_{\substack{m=0 \\ n+m \geq 2}}^n \binom{n}{m} \frac{(-1)^{n-1}}{2nR^{2n}} (2\Delta_\eta \mathbf{X}_c(\theta) \cdot \Delta_\eta \mathbf{X}(\theta))^{n-m} |\Delta_\eta \mathbf{X}(\theta)|^{2m}. \quad (2-7)$$

We expand the denominator in the second term of (1-9) as

$$\frac{1}{|\Delta \mathbf{X} + \Delta \mathbf{X}_c|^2} = \frac{1}{4R^2 \sin^2 \left(\frac{\theta - \eta}{2} \right)} \left(1 - \frac{2}{R^2} \Delta_\eta \mathbf{X}_c(\theta) \cdot \Delta_\eta \mathbf{X}(\theta) \right) + \frac{1}{4R^2 \sin^2 \left(\frac{\theta - \eta}{2} \right)} \mathcal{R}_2(\Delta_\eta \mathbf{X}(\theta)), \quad (2-8)$$

with the notation

$$\mathcal{R}_2(\Delta_\eta \mathbf{X}(\theta)) = \sum_{n=1}^{\infty} \sum_{\substack{m=0 \\ n+m \geq 2}}^n \binom{n}{m} \frac{(-1)^n}{R^{2n}} (2\Delta_\eta \mathbf{X}_c(\theta) \cdot \Delta_\eta \mathbf{X}(\theta))^{n-m} |\Delta_\eta \mathbf{X}(\theta)|^{2m}. \quad (2-9)$$

Therefore, we can write

$$\frac{(\Delta \mathbf{X} + \Delta \mathbf{X}_c) \otimes (\Delta \mathbf{X} + \Delta \mathbf{X}_c)}{|\Delta \mathbf{X} + \Delta \mathbf{X}_c|^2} = A_0 + A_L + A_N, \quad (2-10)$$

with

$$A_0 = \frac{1}{R^2} \Delta_\eta \mathbf{X}_c(\theta) \otimes \Delta_\eta \mathbf{X}_c(\theta),$$

$$A_L = -\frac{2}{R^4} \Delta_\eta \mathbf{X}_c(\theta) \cdot \Delta_\eta \mathbf{X}(\theta) \Delta_\eta \mathbf{X}_c(\theta) \otimes \Delta_\eta \mathbf{X}_c(\theta) + \frac{1}{R^2} \Delta_\eta \mathbf{X}_c(\theta) \otimes \Delta_\eta \mathbf{X}(\theta) + \frac{1}{R^2} \Delta_\eta \mathbf{X}(\theta) \otimes \Delta_\eta \mathbf{X}_c(\theta),$$

and the nonlinear term is given by

$$\begin{aligned} A_N &= \frac{1}{R^2} \Delta_\eta \mathbf{X}(\theta) \otimes \Delta_\eta \mathbf{X}(\theta) \left(1 - \frac{2}{R^2} \Delta_\eta \mathbf{X}_c(\theta) \cdot \Delta_\eta \mathbf{X}(\theta) + \mathcal{R}_2(\Delta_\eta \mathbf{X}(\theta)) \right) \\ &\quad + \frac{1}{R^2} \Delta_\eta \mathbf{X}_c(\theta) \otimes \Delta_\eta \mathbf{X}(\theta) \left(-\frac{2}{R^2} \Delta_\eta \mathbf{X}_c(\theta) \cdot \Delta_\eta \mathbf{X}(\theta) + \mathcal{R}_2(\Delta_\eta \mathbf{X}(\theta)) \right) \\ &\quad + \frac{1}{R^2} \Delta_\eta \mathbf{X}(\theta) \otimes \Delta_\eta \mathbf{X}_c(\theta) \left(-\frac{2}{R^2} \Delta_\eta \mathbf{X}_c(\theta) \cdot \Delta_\eta \mathbf{X}(\theta) + \mathcal{R}_2(\Delta_\eta \mathbf{X}(\theta)) \right) \\ &\quad + \frac{1}{R^2} \Delta_\eta \mathbf{X}_c(\theta) \otimes \Delta_\eta \mathbf{X}_c(\theta) \mathcal{R}_2(\Delta_\eta \mathbf{X}(\theta)). \end{aligned}$$

Joining the expansions (2-6) and (2-10), we split $G(\Delta \mathcal{X})$ in (1-9) into zero-order, linear, and nonlinear parts in terms of \mathbf{X} as follows:

$$G(\Delta \mathcal{X}) = G_0(\Delta_\eta \mathbf{X}_c(\theta)) + G_L(\Delta_\eta \mathbf{X}_c(\theta), \Delta_\eta \mathbf{X}(\theta)) + G_N(\Delta_\eta \mathbf{X}_c(\theta), \Delta_\eta \mathbf{X}(\theta)), \quad (2-11)$$

where

$$G_0(\Delta_\eta \mathbf{X}_c(\theta)) = \frac{1}{4\pi} \left(-\log \left(2R \left| \sin \left(\frac{\theta - \eta}{2} \right) \right| \right) I + \frac{1}{R^2} \Delta_\eta \mathbf{X}_c(\theta) \otimes \Delta_\eta \mathbf{X}_c(\theta) \right), \quad (2-12)$$

$$G_L(\Delta_\eta \mathbf{X}_c(\theta), \Delta_\eta \mathbf{X}(\theta)) = \frac{1}{4\pi R^2} \left(-\Delta_\eta \mathbf{X}_c(\theta) \cdot \Delta_\eta \mathbf{X}(\theta) I - \frac{2}{R^2} \Delta_\eta \mathbf{X}_c(\theta) \cdot \Delta_\eta \mathbf{X}(\theta) \Delta_\eta \mathbf{X}_c(\theta) \otimes \Delta_\eta \mathbf{X}_c(\theta) + \Delta_\eta \mathbf{X}_c(\theta) \otimes \Delta_\eta \mathbf{X}(\theta) + \Delta_\eta \mathbf{X}(\theta) \otimes \Delta_\eta \mathbf{X}_c(\theta) \right), \quad (2-13)$$

$$G_N(\Delta_\eta \mathbf{X}_c(\theta), \Delta_\eta \mathbf{X}(\theta)) = -\frac{1}{4\pi} \mathcal{R}_1(\Delta_\eta \mathbf{X}(\theta)) I + \frac{1}{4\pi R^2} \Delta_\eta \mathbf{X}(\theta) \otimes \Delta_\eta \mathbf{X}(\theta) \left(1 - \frac{2}{R^2} \Delta_\eta \mathbf{X}_c(\theta) \cdot \Delta_\eta \mathbf{X}(\theta) + \mathcal{R}_2(\Delta_\eta \mathbf{X}(\theta)) \right) + \frac{1}{4\pi R^2} \left(\Delta_\eta \mathbf{X}_c(\theta) \otimes \Delta_\eta \mathbf{X}(\theta) + \Delta_\eta \mathbf{X}(\theta) \otimes \Delta_\eta \mathbf{X}_c(\theta) \right) \times \left(-\frac{2}{R^2} \Delta_\eta \mathbf{X}_c(\theta) \cdot \Delta_\eta \mathbf{X}(\theta) + \mathcal{R}_2(\Delta_\eta \mathbf{X}(\theta)) \right) + \frac{1}{4\pi R^2} \Delta_\eta \mathbf{X}_c(\theta) \otimes \Delta_\eta \mathbf{X}_c(\theta) \mathcal{R}_2(\Delta_\eta \mathbf{X}(\theta)). \quad (2-14)$$

Consider the splitting of the solution $\mathbf{F}(\theta)$ to (1-13) into zero-order, linear, and nonlinear parts as

$$\mathbf{F}(\theta) = \mathbf{F}_0(\theta) + \mathbf{F}_L(\theta) + \mathbf{F}_N(\theta). \quad (2-15)$$

(We will prove bounds for these terms in Section 3B.) Introducing the splittings (2-11) and (2-15) in (1-10), we obtain

$$\mathcal{X}_t(\theta) = \mathcal{O}(\mathbf{X}_c)(\theta) + \mathcal{L}(\mathbf{X}_c, \mathbf{X})(\theta) + \mathcal{N}(\mathbf{X}_c, \mathbf{X})(\theta), \quad (2-16)$$

where we recall that $\mathcal{X}(\theta) = \mathbf{X}(\theta) + \mathbf{X}_c(\theta)$ and we use the notation

$$\begin{aligned} \mathcal{O}(\mathbf{X}_c)(\theta) &= \int_{\mathbb{S}} G_0(\Delta_\eta \mathbf{X}_c(\theta)) \mathbf{F}_0(\eta) d\eta, \\ \mathcal{L}(\mathbf{X}_c, \mathbf{X})(\theta) &= \int_{\mathbb{S}} G_0(\Delta_\eta \mathbf{X}_c(\theta)) \mathbf{F}_L(\eta) d\eta + \int_{\mathbb{S}} G_L(\Delta_\eta \mathbf{X}_c(\theta), \Delta_\eta \mathbf{X}(\theta)) \mathbf{F}_0(\eta) d\eta, \\ \mathcal{N}(\mathbf{X}_c, \mathbf{X})(\theta) &= \int_{\mathbb{S}} \left(G_L(\Delta_\eta \mathbf{X}_c(\theta), \Delta_\eta \mathbf{X}(\theta)) \mathbf{F}_L(\eta) + G_N(\Delta_\eta \mathbf{X}_c(\theta), \Delta_\eta \mathbf{X}(\theta)) \mathbf{F}_0(\eta) \right. \\ &\quad \left. + G_N(\Delta_\eta \mathbf{X}_c(\theta), \Delta_\eta \mathbf{X}(\theta)) \mathbf{F}_L(\eta) + G(\mathcal{X}(\theta) - \mathcal{X}(\eta)) \mathbf{F}_N(\eta) \right) d\eta. \end{aligned}$$

We have thus expanded the evolution equation (1-10) distinguishing the zero-order, linear, and nonlinear in \mathbf{X} contributions.

2B. Linearized system. We proceed to show that the linearized system gives rise to a diffusion operator on \mathbf{X} . Since the linear structure is the same for any uniformly parametrized circle (see [Mori et al. 2019]), we will use now (2-1) with $a = 1$, $b = c = d = 0$ and $R = 1$ to simplify the computations, and for clarity of notation we will denote this circle by \mathbf{X}_\star .

We will now linearize (1-10) and (1-13). We first determine \mathbf{F}_0 , the value of \mathbf{F} at the steady state:

$$0 = \mathcal{O}(\mathbf{X}_\star)(\theta) = \int_{\mathbb{S}} G_0(\Delta_\eta \mathbf{X}_\star(\theta)) \mathbf{F}_0(\eta) d\eta, \quad (2-17)$$

$$\mathbf{F}_0(\theta) - 2A_\mu \mathcal{S}_0(\mathbf{F}_0, \mathbf{X}_\star)(\theta) = 2A_e \tilde{\mathbf{F}}_{\text{el},0}(\theta), \quad (2-18)$$

where

$$\tilde{F}_{el,0}(\theta) = \partial_\theta^2 X_\star(\theta)$$

and

$$\mathcal{S}_0(F_0, X_\star)(\theta) = \frac{1}{\pi} \int_{\mathbb{S}} \partial_\theta X_\star^\perp \cdot \Delta_\eta X_\star(\theta) \Delta_\eta X_\star(\theta) \Delta_\eta X_\star(\theta) \cdot \frac{F_0(\eta)}{2 \sin\left(\frac{\theta-\eta}{2}\right)} d\eta.$$

Rewriting $\Delta_\eta X_\star(\theta) \Delta_\eta X_\star(\theta) \cdot F_0(\eta) = \Delta_\eta X_\star(\theta) \otimes \Delta_\eta X_\star(\theta) F_0(\eta)$, and recalling the computations (2-3) and (2-5), one finds that

$$\mathcal{S}_0(F_0, X_\star)(\theta) = -\frac{1}{2\pi} \int_{\mathbb{S}} M(\theta, \eta) F_0(\eta) d\eta,$$

where from (2-5) we have

$$M(\theta, \eta) = \Delta_\eta X_\star(\theta) \otimes \Delta_\eta X_\star(\theta) = \frac{1}{2} \begin{bmatrix} 1 - \cos(\theta + \eta) & -\sin(\theta + \eta) \\ -\sin(\theta + \eta) & 1 + \cos(\theta + \eta) \end{bmatrix}, \tag{2-19}$$

and therefore F_0 is defined by

$$F_0(\theta) + \frac{A_\mu}{\pi} \int_{\mathbb{S}} M(\theta, \eta) F_0(\eta) d\eta = 2A_e \partial_\theta^2 X_\star(\theta).$$

Since $\partial_\theta^2 X_\star = -X_\star$ and noting that

$$\frac{1}{\pi} \int_{\mathbb{S}} M(\theta, \eta) X_\star(\eta) d\eta = -X_\star(\theta),$$

it is easily seen that

$$F_0(\theta) = \frac{2A_e}{1 - A_\mu} \partial_\theta^2 X_\star(\theta). \tag{2-20}$$

Now, recalling (2-12), it can be checked that (2-20) satisfies in fact (2-17):

$$\begin{aligned} 4\pi \left(\frac{1 - A_\mu}{2A_e} \right) \int_{\mathbb{S}} G_0(\Delta_\eta X_\star(\theta)) F_0(\eta) d\eta \\ = - \int_{\mathbb{S}} \log \left(2 \left| \sin \left(\frac{\theta - \eta}{2} \right) \right| \right) \partial_\eta^2 X_\star(\eta) d\eta + \int_{\mathbb{S}} M(\theta, \eta) \partial_\eta^2 X_\star(\eta) d\eta, \end{aligned}$$

so integration by parts in the first term yields (2-17).

We now proceed to compute the linear term $\mathcal{L}(X_\star, X)(\theta)$ in (2-16). For convenience, we write it as

$$\mathcal{L}(X_\star, X)(\theta) = \int_{\mathbb{S}} G_0(\Delta_\eta X_\star) F_L(\eta) d\eta + \int_{\mathbb{S}} (\nabla G(\Delta X_\star) F_0(\eta)) \Delta X d\eta, \tag{2-21}$$

where G_0 and G are defined in (2-12) and (1-9), respectively. To simplify the second integral above, note

$$\begin{aligned} \frac{\partial G_{ij}}{\partial x_1}(\Delta X_\star) e_{r,j}(\eta) &= \frac{\partial G_{i1}}{\partial x_1}(\Delta X_\star) e_{r,1}(\eta) + \frac{\partial G_{i2}}{\partial x_1}(\Delta X_\star) e_{r,2}(\eta) \\ &= \frac{\partial G_{i1}}{\partial x_1}(\Delta X_\star) \partial_\eta X_{\star,2} - \frac{\partial G_{i2}}{\partial x_1}(\Delta X_\star) \partial_\eta X_{\star,1} \\ &= -\frac{\partial G_{i2}}{\partial x_2}(\Delta X_\star) \partial_\eta X_{\star,2} - \frac{\partial G_{i2}}{\partial x_1}(\Delta X_\star) \partial_\eta X_{\star,1} = \partial_\eta G_{i2}(\Delta X_\star). \end{aligned}$$

Here $e_{r,j}$ is the j -th component of the vector e_r etc. Further, in the third equality above, we used the fact that the stokeslet is divergence-free:

$$\frac{\partial G_{i1}}{\partial x_1} + \frac{\partial G_{i2}}{\partial x_2} = 0.$$

Likewise, we have

$$\frac{\partial G_{ij}}{\partial x_2}(\Delta \mathbf{X}_\star) e_{r,j}(\eta) = -\partial_\eta G_{i1}(\Delta \mathbf{X}_\star).$$

We thus have

$$\begin{aligned} \int_{\mathbb{S}} (\nabla G(\Delta \mathbf{X}_\star) \mathbf{F}_0(\eta)) \Delta \mathbf{X} \, d\eta &= -\frac{2A_e}{1-A_\mu} \int_{\mathbb{S}} (\mathcal{R}^{-1} \partial_\eta G(\mathbf{X}_\star(\theta) - \mathbf{X}_\star(\eta))) \Delta \mathbf{X} \, d\eta \\ &= -\frac{2A_e}{1-A_\mu} \int_{\mathbb{S}} (\mathcal{R}^{-1} G(\mathbf{X}_\star(\theta) - \mathbf{X}_\star(\eta))) \partial_\eta \mathbf{X}(\eta) \, d\eta \\ &= \frac{2A_e}{1-A_\mu} \int_{\mathbb{S}} G(\mathbf{X}_\star(\theta) - \mathbf{X}_\star(\eta)) (\mathcal{R}^{-1} \partial_\eta \mathbf{X}(\eta)) \, d\eta. \end{aligned}$$

Since $G(\mathbf{X}_\star(\theta) - \mathbf{X}_\star(\eta)) \stackrel{\text{def}}{=} G_0(\Delta_\eta \mathbf{X}_\star)$, equation (2-21) simplifies to

$$\mathcal{L}(\mathbf{X}_\star, \mathbf{X})(\theta) = \int_{\mathbb{S}} G_0(\Delta_\eta \mathbf{X}_\star) \left(\mathbf{F}_L(\eta) + \frac{2A_e}{1-A_\mu} \mathcal{R}^{-1} \partial_\eta \mathbf{X} \right) \, d\eta. \tag{2-22}$$

This is our specification of the linearized operator.

We will now determine \mathbf{F}_L as in (2-15), that is, the linear part of \mathbf{F} in (1-13). We find

$$\mathbf{F}_L(\theta) + \frac{A_\mu}{\pi} \int_{\mathbb{S}} M(\theta, \eta) \mathbf{F}_L(\eta) \, d\eta = 2A_e \partial_\theta^2 \mathbf{X} - 2A_\mu (\mathbf{Q} + \mathbf{S}), \tag{2-23}$$

where

$$\begin{aligned} Q_i &= - \int_{\mathbb{S}} \mathcal{T}_{ijk}(\Delta \mathbf{X}_\star) F_{0,k}(\eta) \mathcal{R}_{jl}^{-1} \partial_\theta X_l(\theta) \, d\eta, \\ S_i &= - \int_{\mathbb{S}} \frac{\partial \mathcal{T}_{ijk}}{\partial x_m}(\Delta \mathbf{X}_\star) \Delta X_m F_{0,k}(\eta) \mathcal{R}_{jl}^{-1} \partial_\theta X_{*,l}(\theta) \, d\eta. \end{aligned}$$

Let us compute \mathbf{Q} . We start with

$$-\mathcal{T}_{ijk}(\Delta \mathbf{X}_\star) F_{0,k} = -\frac{2A_e}{1-A_\mu} \frac{\Delta X_{*,i} \Delta X_{*,j} \Delta X_{*,k} e_{r,k}(\eta)}{\pi |\Delta \mathbf{X}_\star|^4} = \frac{2A_e}{1-A_\mu} \frac{\Delta X_{*,i} \Delta X_{*,j}}{2\pi |\Delta \mathbf{X}_\star|^2},$$

where we used

$$\frac{\Delta \mathbf{X}_\star \cdot \mathbf{e}_r(\eta)}{|\Delta \mathbf{X}_\star|^2} = -\frac{1}{2}. \tag{2-24}$$

Therefore, we have

$$\begin{aligned} \mathbf{Q} &= \frac{A_e}{1-A_\mu} \frac{1}{\pi} \int_{\mathbb{S}} \frac{\Delta \mathbf{X}_\star \otimes \Delta \mathbf{X}_\star}{|\Delta \mathbf{X}_\star|^2} \, d\eta \mathcal{R}^{-1} \partial_\theta \mathbf{X}(\theta) \\ &= \frac{A_e}{1-A_\mu} \frac{1}{\pi} \int_{\mathbb{S}} M(\theta, \eta) \, d\eta \mathcal{R}^{-1} \partial_\theta \mathbf{X}(\theta) = \frac{A_e}{1-A_\mu} \mathcal{R}^{-1} \partial_\theta \mathbf{X}(\theta), \end{aligned}$$

where we used (2-19) in the second equality. We next compute S ,

$$\begin{aligned}
 & -\frac{\partial \mathcal{T}_{ijk}}{\partial x_m}(\Delta X_\star)\Delta X_m F_{0,k}(\eta)\mathcal{R}_{jl}^{-1}\partial_\theta X_{\star,l}(\theta) \\
 &= -\frac{1}{\pi}\frac{2A_e}{1-A_\mu}\left(\frac{\Delta X_i\Delta X_{\star,j}\Delta X_{\star,k}e_{r,j}(\theta)e_{r,k}(\eta)}{|\Delta X_\star|^4}\right) \\
 & \quad -\frac{2A_e/\pi}{1-A_\mu}\left(\frac{\Delta X_{\star,i}\Delta X_j\Delta X_{\star,k}e_{r,j}(\theta)e_{r,k}(\eta)}{|\Delta X_\star|^4}+\frac{\Delta X_{\star,i}\Delta X_{\star,j}\Delta X_k e_{r,j}(\theta)e_{r,k}(\eta)}{|\Delta X_\star|^4}\right) \\
 & \quad +\frac{8A_e/\pi}{1-A_\mu}\left(\frac{\Delta X_{\star,i}\Delta X_{\star,j}\Delta X_{\star,k}e_{r,j}(\theta)e_{r,k}(\eta)\Delta X_{\star,m}\Delta X_m}{|\Delta X_\star|^6}\right) \\
 &= -\frac{1}{\pi}\frac{2A_e}{1-A_\mu}I-\frac{2A_e/\pi}{1-A_\mu}II+\frac{8A_e/\pi}{1-A_\mu}III.
 \end{aligned}$$

We simplify each term as follows:

$$\begin{aligned}
 I &= -\frac{1}{4}\Delta X_i, \\
 II &= -\frac{\Delta X_{\star,i}\Delta X_j e_{r,j}(\theta)}{2|\Delta X_\star|^2}+\frac{\Delta X_{\star,i}\Delta X_k e_{r,k}(\eta)}{2|\Delta X_\star|^2}=-\frac{\Delta X_{\star,i}\Delta X_{\star,j}\Delta X_j}{2|\Delta X_\star|^2}, \\
 III &= -\frac{\Delta X_{\star,i}\Delta X_{\star,j}\Delta X_j}{4|\Delta X_\star|^2},
 \end{aligned}$$

where above we made repeated use of (2-24) and

$$\frac{\Delta X_\star \cdot \mathcal{R}^{-1}\partial_\theta X_\star(\theta)}{|\Delta X_\star|^2}=\frac{\Delta X_\star \cdot e_r(\theta)}{|\Delta X_\star|^2}=\frac{1}{2}.$$

Thus

$$-\frac{\partial \mathcal{T}_{ijk}}{\partial x_m}(\Delta X_\star)\Delta X_m F_{0,k}(\eta)\mathcal{R}_{jl}^{-1}\partial_\theta X_{\star,l}=\frac{A_e/2\pi}{1-A_\mu}\left(\Delta X_i-\frac{2\Delta X_{\star,i}\Delta X_{\star,j}\Delta X_j}{|\Delta X_\star|^2}\right).$$

Substituting this back into the expression for S in (2-23), we have

$$\begin{aligned}
 S &= \frac{A_e/2\pi}{1-A_\mu}\int_{\mathbb{S}}(I-2M(\theta,\eta))\Delta X\,d\eta \\
 &= -\frac{A_e/2\pi}{1-A_\mu}\int_{\mathbb{S}}(I-2M(\theta,\eta))\mathbf{X}(\eta)\,d\eta=\frac{A_e/2\pi}{1-A_\mu}(-\langle \mathbf{e}_r, \mathbf{X} \rangle \mathbf{e}_r+\langle \mathbf{e}_t, \mathbf{X} \rangle \mathbf{e}_t),
 \end{aligned}$$

where we used (2-19) in the first equality and we are using the notation (1-24) for the inner product.

Equation (2-23) thus reduces to

$$\begin{aligned}
 \mathbf{F}_L(\theta)+\frac{A_\mu}{\pi}\int_{\mathbb{S}}M(\theta,\eta)\mathbf{F}_L(\eta)\,d\eta \\
 =2A_e\partial_\theta^2\mathbf{X}(\theta)-\frac{2A_eA_\mu}{1-A_\mu}\left(\mathcal{R}^{-1}\partial_\theta\mathbf{X}(\theta)+\frac{1}{2\pi}(-\langle \mathbf{e}_r, \mathbf{X} \rangle \mathbf{e}_r(\theta)+\langle \mathbf{e}_t, \mathbf{X} \rangle \mathbf{e}_t(\theta))\right).
 \end{aligned}$$

We must solve the above equation for \mathbf{F}_L in terms of \mathbf{X} . Suppose $\langle \mathbf{e}_r, \mathbf{X} \rangle = \langle \mathbf{e}_t, \mathbf{X} \rangle = 0$. Then, it is easily checked that

$$\mathbf{F}_L(\theta)=2A_e\partial_\theta^2\mathbf{X}(\theta)-\frac{2A_eA_\mu}{1-A_\mu}\mathcal{R}^{-1}\partial_\theta\mathbf{X}(\theta). \tag{2-25}$$

We may further compute F_L when X is either e_r or e_t . Noting that

$$M(\theta, \eta)e_r(\eta) = \frac{1}{2}(e_r(\eta) - e_r(\theta)), \quad M(\theta, \eta)e_t(\eta) = \frac{1}{2}(e_t(\eta) + e_t(\theta)),$$

we see by an easy calculation that

$$\text{If } X = e_{r,t}, \text{ then } F_L = -\frac{2A_e}{1 - A_\mu}e_{r,t}.$$

Note that

$$2A_e\partial_\theta^2 e_{r,t} - \frac{2A_e A_\mu}{1 - A_\mu} \mathcal{R}^{-1} \partial_\theta e_{r,t} = -\frac{2A_e}{1 - A_\mu} e_{r,t}.$$

This shows that the expression for F_L in (2-25) is in fact valid without the restriction $\langle e_r, X \rangle = \langle e_t, X \rangle = 0$. Substituting (2-25) into (2-22) yields

$$\mathcal{L}(X_\star, X)(\theta) = 2A_e \int_{\mathbb{S}} G_0(\Delta_\eta X_\star)(\partial_\eta^2 X(\eta) + \mathcal{R}^{-1} \partial_\eta X(\eta)) d\eta. \tag{2-26}$$

Finally, since

$$G_0(\Delta_\eta X_\star) = -\frac{1}{4\pi} \left(\log \left| 2 \sin \left(\frac{\theta - \eta}{2} \right) \right| \right) I + M(\theta, \eta)$$

and

$$\int_{\mathbb{S}} M(\theta, \eta) \left(\partial_\eta^2 X(\eta) + \mathcal{R}^{-1} \partial_\eta X(\eta) \right) d\eta = \int_{\mathbb{S}} (\partial_\eta^2 M(\theta, \eta) - \partial_\eta M(\theta, \eta) \mathcal{R}^{-1}) X(\eta) d\eta = 0,$$

we have

$$\begin{aligned} \mathcal{L}(X_\star, X)(\theta) &= -\frac{A_e}{2\pi} \int_{\mathbb{S}} \log \left| 2 \sin \left(\frac{\theta - \eta}{2} \right) \right| (\partial_\eta^2 X(\eta) + \mathcal{R}^{-1} \partial_\eta X(\eta)) d\eta \\ &= -\frac{A_e}{2\pi} \int_{\mathbb{S}} \frac{\partial_\eta X(\eta) + \mathcal{R}^{-1} X(\eta)}{2 \tan \left(\frac{\theta - \eta}{2} \right)} d\eta, \end{aligned}$$

which is given by a Hilbert transform

$$\mathcal{L}(X_\star, X)(\theta) = -\frac{A_e}{2} \mathcal{H}(\partial_\eta X(\eta) + \mathcal{R}^{-1} X(\eta))(\theta). \tag{2-27}$$

Therefore, the system (2-16) can be written as

$$\mathcal{X}_t(\theta) = -\frac{A_e}{2} (\Lambda X(\theta) + \mathcal{H} \mathcal{R}^{-1} X(\theta)) + \mathcal{N}(X_c, X)(\theta). \tag{2-28}$$

Notice that X_c is a uniformly parametrized circle with time-dependent radius $R(t)$, as opposed to the X_\star used in this subsection to obtain the linearization. We will use the system (2-28) to study the global-in-time dynamics of the Peskin problem in the rest of this paper.

2C. Evolution of the $\dot{\mathcal{F}}_v^{1,1}$ norm of X . We first notice that, because $X(\theta)$ is real-valued, it must hold that $\widehat{X}(-k) = \overline{\widehat{X}(k)}$. Therefore, the norm (1-25) can be written in terms of positive frequencies alone

$$\|X\|_{\dot{\mathcal{F}}_v^{1,1}} = 2 \sum_{k \geq 1} e^{\nu(t)k} k |\widehat{X}(k)| = 2 \sum_{k \geq 1} e^{\nu(t)k} k \sqrt{\widehat{X}_1 \overline{\widehat{X}_1}(k) + \widehat{X}_2 \overline{\widehat{X}_2}(k)}. \tag{2-29}$$

The system (2-28) in Fourier variables reads for $k \geq 0$ as

$$\widehat{\mathcal{X}}_t(k) = -\frac{A_e}{2}L(k)\widehat{\mathcal{X}}(k) + \mathcal{F}(\mathcal{N}(\mathbf{X}_c, \mathbf{X}))(k). \tag{2-30}$$

Here we recall that $\mathcal{X} = \mathbf{X} + \mathbf{X}_c$. Further the diffusion matrix is given by

$$L(k) = \begin{bmatrix} k & -i \operatorname{sgn}(k) \\ i \operatorname{sgn}(k) & k \end{bmatrix}, \quad k \geq 1, \quad L(0) = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}.$$

The diagonalization of this matrix for $k \geq 1$ shows that

$$L(k) = P(k)\mathcal{D}(k)P(k)^{-1},$$

where for $k \geq 1$ we have

$$P(k) = \frac{1}{\sqrt{2}} \begin{bmatrix} -i \operatorname{sgn}(k) & 1 \\ 1 & -i \operatorname{sgn}(k) \end{bmatrix}, \quad P(k)^{-1} = \overline{P(k)}, \quad \mathcal{D}(k) = \begin{bmatrix} k+1 & 0 \\ 0 & k-1 \end{bmatrix}.$$

And when $k = 0$ we define

$$P(0) = \frac{1}{\sqrt{2}} \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}, \quad P(0)^{-1} = 2\overline{P(0)}, \quad \mathcal{D}(0) = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}.$$

This leads us to define the change of variables

$$\widehat{\mathcal{Y}}(k) \stackrel{\text{def}}{=} P(k)^{-1}\widehat{\mathcal{X}}(k), \quad \widehat{\mathcal{Y}}_c(k) \stackrel{\text{def}}{=} P(k)^{-1}\widehat{\mathcal{X}}_c(k), \tag{2-31}$$

with $\mathcal{Y} \stackrel{\text{def}}{=} \mathbf{Y} + \mathbf{Y}_c$. The system (2-30) for $k \geq 0$ then becomes

$$\widehat{\mathcal{Y}}_t(k) = -\frac{A_e}{2}\mathcal{D}(k)\widehat{\mathcal{Y}}(k) + P(k)^{-1}\mathcal{F}(\mathcal{N}(\mathbf{X}_c, \mathbf{X}))(k). \tag{2-32}$$

The relationship between \mathbf{X} and \mathbf{Y} in space variables is given by the Hilbert transform (1-20), using also $\mathcal{H}^2(Y_j) = -Y_j$, as follows:

$$\mathbf{X}(\theta) = \frac{1}{\sqrt{2}} \begin{bmatrix} \mathcal{H}Y_1(\theta) + Y_2(\theta) \\ Y_1(\theta) + \mathcal{H}Y_2(\theta) \end{bmatrix}, \quad \mathbf{Y}(\theta) = \frac{1}{\sqrt{2}} \begin{bmatrix} -\mathcal{H}X_1(\theta) + X_2(\theta) \\ X_1(\theta) - \mathcal{H}X_2(\theta) \end{bmatrix}. \tag{2-33}$$

Because, for $k \neq 0$, $P(k)$ is a unitary matrix, it holds that $\|P(k)\| = \|P(k)^{-1}\| = 1$, and therefore

$$|\widehat{\mathcal{Y}}(k)| = |\widehat{\mathcal{X}}(k)|, \quad k \neq 0,$$

and thus

$$\|\mathbf{X}\|_{\dot{\mathcal{F}}_v^{1,1}} = \|\mathbf{Y}\|_{\dot{\mathcal{F}}_v^{1,1}}. \tag{2-34}$$

We will use this norm equivalence several times in the following.

Notice that the first Fourier coefficient of a uniformly parametrized circle (2-1) is given by

$$\widehat{\mathcal{X}}_c(1) = \frac{1}{2} \begin{bmatrix} a+bi \\ -ia+b \end{bmatrix},$$

and in the Y -variable

$$\widehat{Y}_c(1) = \frac{1}{\sqrt{2}} \begin{bmatrix} 0 \\ a+bi \end{bmatrix}. \tag{2-35}$$

Note that $\mathcal{H}(\cos \theta) = \sin \theta$ and $\mathcal{H}(\sin \theta) = -\cos \theta$. Then from the transformation (2-33) uniformly parametrized circles (1-16) in the \mathcal{Y} variable are spanned by

$$\tilde{e}_r(\theta) = \sqrt{2} \begin{bmatrix} 0 \\ \cos \theta \end{bmatrix}, \quad \tilde{e}_t(\theta) = \sqrt{2} \begin{bmatrix} 0 \\ -\sin \theta \end{bmatrix}, \quad \tilde{e}_1 = \frac{1}{\sqrt{2}} \begin{bmatrix} 0 \\ 1 \end{bmatrix}, \quad \tilde{e}_2 = \frac{1}{\sqrt{2}} \begin{bmatrix} 1 \\ 0 \end{bmatrix}. \tag{2-36}$$

Further the second component of $\widehat{Y}(1)$ becomes zero after the operation of $\mathcal{D}(1)$ is applied, which corresponds to the fact that uniformly parametrized circles are steady states. Therefore, we will split the curve $\mathcal{Y}(\theta) = Y_c(\theta) + Y(\theta)$, with

$$\widehat{Y}(0) = \begin{bmatrix} 0 \\ 0 \end{bmatrix}, \quad \widehat{Y}_2(1) = 0,$$

since those frequencies are contained in the time-dependent circle (2-1). In other words, Y is the projection of \mathcal{Y} onto the orthogonal complement of the vector space spanned by (2-36). In fact, the system of equations (2-32) does not provide dissipation for the zero frequency of \mathcal{Y} nor for the second component of its first frequency (i.e., for uniformly parametrized circles). We thus can only expect decay for Y . It is convenient then to write the equations of those frequencies in (2-32) separately:

$$\begin{aligned} \widehat{Y}_t(0) &= \partial_t \widehat{Y}_c(0) = P(0)^{-1} \mathcal{F}(\mathcal{N}(X_c, X))(0), \\ \partial_t \widehat{Y}_1(1) &= \partial_t \widehat{Y}_1(1) = -A_e \widehat{Y}_1(1) + (P(1)^{-1} \mathcal{F}(\mathcal{N}(X_c, X))(1))_1, \\ \partial_t \widehat{Y}_2(1) &= \partial_t \widehat{Y}_{c,2}(1) = (P(1)^{-1} \mathcal{F}(\mathcal{N}(X_c, X))(1))_2, \\ \widehat{Y}_t(k) &= \widehat{Y}_t(k) = -\frac{A_e}{2} \mathcal{D}(k) \widehat{Y}(k) + P(k)^{-1} \mathcal{F}(\mathcal{N}(X_c, X))(k), \quad k \geq 2. \end{aligned} \tag{2-37}$$

Therefore, we study the evolution in time of $\|Y\|_{\dot{Y}_v^{1,1}}$, as in (2-29), which is given by

$$\begin{aligned} \frac{d}{dt} \|Y\|_{\dot{Y}_v^{1,1}} &= \frac{d}{dt} \left(2 \sum_{k \geq 1} e^{\nu(t)k} k \sqrt{\widehat{Y}_1(k) \overline{\widehat{Y}_1(k)} + \widehat{Y}_2(k) \overline{\widehat{Y}_2(k)}} \right) \\ &= 2 \sum_{k \geq 1} \nu'(t) k^2 e^{\nu(t)k} |\widehat{Y}(k)| + 2 \sum_{k \geq 1} e^{\nu(t)k} k \frac{\partial_t \widehat{Y}(k)^T \overline{\widehat{Y}(k)} + \widehat{Y}(k)^T \partial_t \overline{\widehat{Y}(k)}}{2|\widehat{Y}(k)|}, \end{aligned}$$

and introducing the time derivative (2-32), with $\mathcal{N} = \mathcal{N}(X_c, X) = \mathcal{N}(X)$, we have

$$\begin{aligned} \frac{d}{dt} \|Y\|_{\dot{Y}_v^{1,1}} &= 2 \sum_{k \geq 1} \nu'(t) k^2 e^{\nu(t)k} |\widehat{Y}(k)| - 2A_e \sum_{k \geq 1} e^{\nu(t)k} k \frac{(k+1)|\widehat{Y}_1(k)|^2 + (k-1)|\widehat{Y}_2(k)|^2}{2|\widehat{Y}(k)|} \\ &\quad + 2 \sum_{k \geq 1} e^{\nu(t)k} k \frac{(P(k)^{-1} \widehat{\mathcal{N}}(X)(k))^T \overline{\widehat{Y}(k)} + \widehat{Y}(k)^T (P(k)^{-1} \widehat{\mathcal{N}}(X)(k))}{2|\widehat{Y}(k)|}. \end{aligned}$$

Noticing that for $k \geq 1$ we have

$$-A_e k \left((k+1)|\widehat{Y}_1(k)|^2 + (k-1)|\widehat{Y}_2(k)|^2 \right) \frac{1}{|\widehat{Y}(k)|} = -A_e k (k-1) |\widehat{Y}(k)| - 2A_e k \frac{|\widehat{Y}_1(k)|^2}{|\widehat{Y}(k)|},$$

we can then see a diffusion term coming from the linear part:

$$\begin{aligned} \frac{d}{dt} \|\mathbf{Y}\|_{\dot{\mathcal{F}}_v^{1,1}} \leq & -A_e \sum_{k \geq 1} e^{v(t)k} k(k-1) |\widehat{\mathbf{Y}}(k)| - 2A_e \sum_{k \geq 1} e^{v(t)k} k \frac{|\widehat{\mathbf{Y}}_1(k)|^2}{|\widehat{\mathbf{Y}}(k)|} \\ & + 2 \sum_{k \geq 1} v'(t) k^2 e^{v(t)k} |\widehat{\mathbf{Y}}(k)| + 2 \sum_{k \geq 1} e^{v(t)k} k |(P(k)^{-1} \widehat{\mathcal{N}}(\mathcal{X}))(k)|. \end{aligned} \quad (2-38)$$

The balance above does not include the control of $\widehat{\mathbf{Y}}_c$. We will show in [Section 4A](#) that the evolution of $\widehat{\mathbf{Y}}_c(0)$, that is, of the center, can be controlled by all the other frequencies. Moreover, the incompressibility condition (1-3) allows us to control $\widehat{\mathbf{Y}}_{c,2}(1)$ as follows:

$$\begin{aligned} V_0 = \pi &= \frac{1}{2} \int_{-\pi}^{\pi} \mathcal{X}(\theta) \wedge \partial_{\theta} \mathcal{X}(\theta) d\theta = \frac{1}{2} \int_{-\pi}^{\pi} (\mathcal{X}_1 \partial_{\theta} \mathcal{X}_2 - \mathcal{X}_2 \partial_{\theta} \mathcal{X}_1) d\theta \\ &= \frac{1}{4} \int_{-\pi}^{\pi} ((\mathcal{H}\mathcal{Y}_1 + \mathcal{Y}_2)(\partial_{\theta} \mathcal{Y}_1 + \Lambda \mathcal{Y}_2) - (\mathcal{Y}_1 + \mathcal{H}\mathcal{Y}_2)(\Lambda \mathcal{Y}_1 + \partial_{\theta} \mathcal{Y}_2)) d\theta. \end{aligned} \quad (2-39)$$

Performing the products and taking into account the equalities

$$\int_{-\pi}^{\pi} \mathcal{H}\mathcal{Y}_i \Lambda \mathcal{Y}_j d\theta = \int_{-\pi}^{\pi} \mathcal{Y}_i \partial_{\theta} \mathcal{Y}_j d\theta, \quad \int_{-\pi}^{\pi} \mathcal{H}\mathcal{Y}_j \partial_{\theta} \mathcal{Y}_j d\theta = - \int_{-\pi}^{\pi} \mathcal{Y}_j \Lambda \mathcal{Y}_j d\theta,$$

we obtain

$$\begin{aligned} \pi &= \frac{1}{2} \int_{-\pi}^{\pi} (\mathcal{Y}_2 \Lambda \mathcal{Y}_2 - \mathcal{Y}_1 \Lambda \mathcal{Y}_1) d\theta = \pi (\widehat{\mathcal{Y}}_2 \Lambda \widehat{\mathcal{Y}}_2(0) - \widehat{\mathcal{Y}}_1 \Lambda \widehat{\mathcal{Y}}_1(0)) \\ &= \pi \sum_{k \in \mathbb{Z}} (|k| \widehat{\mathcal{Y}}_2(k) \widehat{\mathcal{Y}}_2(-k) - |k| \widehat{\mathcal{Y}}_1(k) \widehat{\mathcal{Y}}_1(-k)) \\ &= \pi \sum_{k \in \mathbb{Z}} |k| (|\widehat{\mathbf{Y}}_{c,2}(k)|^2 + \widehat{\mathbf{Y}}_{c,2}(k) \widehat{\mathbf{Y}}_2(-k) + \widehat{\mathbf{Y}}_2(k) \widehat{\mathbf{Y}}_{c,2}(-k) + |\widehat{\mathbf{Y}}_2(k)|^2 - |\widehat{\mathbf{Y}}_1(k)|^2), \end{aligned}$$

where we have used that $\widehat{\mathbf{Y}}_{c,1}(k) = 0$ for $k \neq 0$. We can also eliminate the terms $\widehat{\mathbf{Y}}_{c,2}(k) \widehat{\mathbf{Y}}_2(-k)$ and $\widehat{\mathbf{Y}}_2(k) \widehat{\mathbf{Y}}_{c,2}(-k)$, since $\widehat{\mathbf{Y}}_2(1) = 0$ and $\widehat{\mathbf{Y}}_{c,2}(k) = 0$ for $k \neq 0, \pm 1$. Therefore,

$$\frac{1}{2} = \frac{a^2 + b^2}{2} + \sum_{k \geq 1} k (|\widehat{\mathbf{Y}}_2(k)|^2 - |\widehat{\mathbf{Y}}_1(k)|^2).$$

And so the incompressibility condition translates to the constraint

$$|\widehat{\mathbf{Y}}_{c,2}(1)|^2 = \frac{a^2 + b^2}{2} = \frac{R^2}{2} = \frac{1}{2} - \sum_{k \geq 1} k (|\widehat{\mathbf{Y}}_2(k)|^2 - |\widehat{\mathbf{Y}}_1(k)|^2). \quad (2-40)$$

Then, we can obtain an upper bound

$$\begin{aligned} |\widehat{\mathbf{Y}}_{c,2}(1)|^2 &\leq \frac{1}{2} + \sum_{k \geq 1} k (|\widehat{\mathbf{Y}}_2(k)|^2 + |\widehat{\mathbf{Y}}_1(k)|^2) = \frac{1}{2} + \sum_{k \geq 1} (k^{1/2} |\widehat{\mathbf{Y}}(k)|)^2 \\ &\leq \frac{1}{2} + \left(\sum_{k \geq 1} k^{1/2} |\widehat{\mathbf{Y}}(k)| \right)^2 = \frac{1}{2} + \frac{1}{4} \|\mathbf{Y}\|_{\dot{\mathcal{F}}^{1/2,1}}^2 \leq \frac{1}{2} + \frac{1}{4} \|\mathbf{Y}\|_{\dot{\mathcal{F}}^{1/2,1}}^2, \end{aligned}$$

and analogously we find the lower bound

$$\frac{R^2}{2} = |\widehat{\mathbf{Y}}_{c,2}(1)|^2 \geq \frac{1}{2} - \frac{1}{4} \|\mathbf{Y}\|_{\dot{\mathcal{F}}^{1/2,1}}^2. \quad (2-41)$$

Recalling the relationship between \mathbf{X} and \mathbf{Y} in (2-34), we finally obtain

$$\frac{1}{2} - \frac{1}{4} \|\mathbf{X}\|_{\dot{\mathcal{F}}_v^{1,1}}^2 \leq |\widehat{\mathbf{Y}}_{c,2}(1)|^2 \leq \frac{1}{2} + \frac{1}{4} \|\mathbf{X}\|_{\dot{\mathcal{F}}_v^{1,1}}^2,$$

and, since $|\widehat{\mathbf{Y}}_c(1)|^2 = R^2/2 = |\widehat{\mathbf{X}}_c(1)|^2/2$,

$$1 - \frac{1}{2} \|\mathbf{X}\|_{\dot{\mathcal{F}}_v^{1,1}}^2 \leq |\widehat{\mathbf{X}}_c(1)|^2 \leq 1 + \frac{1}{2} \|\mathbf{X}\|_{\dot{\mathcal{F}}_v^{1,1}}^2,$$

so using the notation $R^2 = a^2 + b^2$, we have

$$\frac{1}{\sqrt{1 + \frac{1}{2} \|\mathbf{X}\|_{\dot{\mathcal{F}}_v^{1,1}}^2}} \leq \frac{1}{R} \leq \frac{1}{\sqrt{1 - \frac{1}{2} \|\mathbf{X}\|_{\dot{\mathcal{F}}_v^{1,1}}^2}}. \tag{2-42}$$

The upper bound above motivates us to define

$$C_1 \stackrel{\text{def}}{=} C_1(\|\mathbf{X}\|_{\dot{\mathcal{F}}_v^{1,1}}) = \frac{1}{\sqrt{1 - \frac{1}{2} \|\mathbf{X}\|_{\dot{\mathcal{F}}_v^{1,1}}^2}}. \tag{2-43}$$

We will later use (2-42) to control the size of R when $\|\mathbf{X}\|_{\dot{\mathcal{F}}_v^{1,1}}(t) \rightarrow 0$ as $t \rightarrow \infty$.

2D. Complete system. We finally summarize the final form of the system of equations that describes our problem. The system given by (1-10) and (1-13) for \mathcal{X} was replaced by (2-37) on the Fourier coefficients of the associated variable \mathcal{Y} from (2-31). We recall that we decompose \mathcal{Y} into a time-dependent circle Y_c plus the deviation from the circle given by Y . In other words, we decompose \mathcal{Y} into its projection onto the vector space spanned by (2-36) represented by Y_c and its orthogonal complement represented by Y . Therefore, recalling (2-35), we have

$$\widehat{\mathbf{Y}}(0) = 0, \quad \widehat{\mathbf{Y}}_2(1) = 0, \quad \widehat{\mathbf{Y}}_c(k) = 0, \quad k \neq 0, 1, \quad \widehat{\mathbf{Y}}_{c,1}(1) = 0. \tag{2-44}$$

Now, for $k = 1$ and $k \geq 2$ separately, we have

$$\begin{aligned} \partial_t \widehat{\mathbf{Y}}_1(1) &= -A_e \widehat{\mathbf{Y}}_1(1) + (P(1)^{-1} \widehat{\mathcal{N}}(\mathbf{X}_c, \mathbf{X})(1))_1, \\ \partial_t \widehat{\mathbf{Y}}(k) &= -\frac{A_e}{2} \mathcal{D}(k) \widehat{\mathbf{Y}}(k) + P(k)^{-1} \widehat{\mathcal{N}}(\mathbf{X}_c, \mathbf{X})(k), \end{aligned} \tag{2-45}$$

where \mathbf{X}_c and \mathbf{X} are given in terms of Y_c and Y in (2-31). In the following paragraphs, we will write one or the other without distinction for simplicity of notation. The incompressibility condition (2-39) yielded (2-40). Thus in particular

$$\sqrt{\frac{1}{2} - \frac{1}{4} \|\mathbf{Y}\|_{\dot{\mathcal{F}}_v^{1,1}}^2} \leq |\widehat{\mathbf{Y}}_{c,2}(1)| \leq \sqrt{\frac{1}{2} + \frac{1}{4} \|\mathbf{Y}\|_{\dot{\mathcal{F}}_v^{1,1}}^2}. \tag{2-46}$$

To close the system, notice that $\widehat{\mathbf{Y}}_c(0) = P(k)^{-1} \widehat{\mathbf{X}}_c(0)$ and, from (1-10), we have

$$\partial_t \widehat{\mathbf{X}}_c(0) = \frac{1}{2\pi} \int_{\mathbb{S}} \int_{\mathbb{S}} G(\Delta \mathbf{X}_c + \Delta \mathbf{X}) \mathbf{F}(\eta) d\eta d\theta,$$

with \mathbf{F} defined by (1-13). We can also write the equation for $\widehat{\mathbf{X}}_c(0)$ using (2-16) or (2-30) and recalling that the zero frequency of the linear part vanishes,

$$\partial_t \widehat{\mathbf{X}}_c(0) = \widehat{\mathcal{N}}(\mathbf{X}_c, \mathbf{X})(0). \tag{2-47}$$

We notice that the evolution of the zero frequency $\widehat{Y}_c(0)$, corresponding to the center, is decoupled from all the other equations (in terms of the $\widehat{Y}_c(0)$ -variable), because $\widehat{X}_c(0)$ does not appear on the right-hand side of (1-10) and (1-13) and therefore also (2-47). This can be seen from the fact that in (1-10) G only depends on the difference $\Delta\mathcal{X} = \mathcal{X}(\theta) - \mathcal{X}(\eta)$ and in (1-13) the expression for \mathcal{S} only depends on $\partial_\theta\mathcal{X}$ and $\Delta\mathcal{X}$. In summary, the system to determine \mathcal{Y} (equivalently determining \mathcal{X} via (2-31)) consists of (2-44), (2-45), (2-40), and (2-47). That is, all together we have

$$\begin{aligned} \widehat{Y}(0) &= 0, & \widehat{Y}_2(1) &= 0, & \widehat{Y}_c(k) &= 0, \quad k \neq 0, 1, & \widehat{Y}_{c,1}(1) &= 0, \\ \partial_t \widehat{Y}_c(0) &= P(0)^{-1} \widehat{\mathcal{N}}(\mathbf{X}_c, \mathbf{X})(0), \\ \partial_t \widehat{Y}_1(1) &= -A_e \widehat{Y}_1(1) + (P(1)^{-1} \widehat{\mathcal{N}}(\mathbf{X}_c, \mathbf{X})(1))_1, \\ \partial_t \widehat{Y}(k) &= -\frac{A_e}{2} \mathcal{D}(k) \widehat{Y}(k) + P(k)^{-1} \widehat{\mathcal{N}}(\mathbf{X}_c, \mathbf{X})(k), \quad k \geq 2, \\ |\widehat{Y}_{c,2}(1)|^2 &= \frac{1}{2} - \sum_{k \geq 1} k(|\widehat{Y}_2(k)|^2 - |\widehat{Y}_1(k)|^2), \end{aligned} \tag{2-48}$$

with \mathbf{F} given in (1-13), and \mathbf{Y}, \mathbf{X} related by (2-31).

To prove Theorem 1.2 (see Section 4) we will use system (2-48) to obtain the energy balance (2-38) to show the decay of \mathbf{Y} . We will need to perform a priori estimates on the nonlinear terms, which in particular requires us to prove bounds for \mathbf{F} due to the viscosity contrast. Those estimates are performed in the next section. The decay for \mathbf{Y} will allow us to control the evolution of the zero frequency, that is, of the center.

3. A priori estimates

In this section we perform the a priori estimates on \mathbf{X} and \mathbf{F} that will be used in the proof of our main result, Theorem 1.2. First, in Proposition 3.1, we estimate the nonlinear terms in (2-16) in terms of \mathbf{X} and \mathbf{F} . Next, in Section 3B, we obtain the a priori estimates for \mathbf{F} in (1-13) in terms of \mathbf{X} . In order to get the result with critical regularity, we have to get uniform bounds for some Fourier multipliers given by principal values (see Lemma 3.2).

3A. A priori estimates on \mathbf{X} .

Proposition 3.1. *Assume $\mathbf{F}_0, \mathbf{F}_L, \mathbf{F}_N \in \mathcal{F}_v^{0,1}$ and $\mathbf{X} \in \dot{\mathcal{F}}_v^{2,1}$. Then, the nonlinear term $\mathcal{N} = \mathcal{N}(\mathbf{X}_c, \mathbf{X})(\theta) = \mathcal{N}(\mathcal{X})$ in (2-16) satisfies the following estimate in $\dot{\mathcal{F}}_v^{1,1}$:*

$$\|\mathcal{N}\|_{\dot{\mathcal{F}}_v^{1,1}} \leq 11\sqrt{2}D_1 \|\mathbf{X}\|_{\dot{\mathcal{F}}_v^{1,1}} \|\mathbf{F}_L\|_{\mathcal{F}_v^{0,1}} + \frac{147}{2}D_2 \|\mathbf{F}_0\|_{\mathcal{F}_v^{0,1}} \|\mathbf{X}\|_{\dot{\mathcal{F}}_v^{1,1}} \|\mathbf{X}\|_{\dot{\mathcal{F}}_v^{2,1}} + \frac{9}{4}D_3 \|\mathbf{F}_N\|_{\mathcal{F}_v^{0,1}}, \tag{3-1}$$

where $D_i = D_i(\|\mathbf{X}\|_{\dot{\mathcal{F}}_v^{1,1}}, v_\infty) \approx 1$ are increasing functions of $\|\mathbf{X}\|_{\dot{\mathcal{F}}_v^{1,1}}$ and v_∞ such that

$$\lim_{\|\mathbf{X}\|_{\dot{\mathcal{F}}_v^{1,1}} \rightarrow 0^+} D_i(\|\mathbf{X}\|_{\dot{\mathcal{F}}_v^{1,1}}, 0) = 1$$

and are defined in (3-54).

In the proof, the following multiplier will come up frequently:

$$m(k, \eta) \stackrel{\text{def}}{=} \frac{1 - \frac{\sin(k\eta/2)}{k \tan(\eta/2)} e^{-ik\eta/2}}{2 \sin(\eta/2)}, \quad |k| \geq 1, \tag{3-2}$$

and we define $m(0, \eta) = 0$.

Now let $n \geq 1$, $k = k_0, k_1, \dots, k_{2n}$ be integers that further satisfy $|k_j - k_{j+1}| \geq 1$ for all $j = 0, 1, \dots, 2n - 1$. We define the integral of type $I_n = I_n(k, k_1, \dots, k_{2n})$ by

$$I_n \stackrel{\text{def}}{=} \text{pv} \int_{-\pi}^{\pi} m(k - k_1, \eta) \prod_{j=1}^{2n-1} \frac{\sin((k_j - k_{j+1})\eta/2)}{(k_j - k_{j+1}) \sin(\eta/2)} e^{-i(k_1 + k_{2n})\eta/2} d\eta. \tag{3-3}$$

We further define $I_n = 0$ if $k_j = k_{j+1}$ for any $j = 0, 1, \dots, 2n - 1$. We will also consider the integral, $I'_n = I'_n(k_1, \dots, k_{2n})$, under the same conditions

$$I'_n \stackrel{\text{def}}{=} \text{pv} \int_{-\pi}^{\pi} \frac{\sin((k_1 + k_{2n})\eta/2)}{\sin(\eta/2)} \prod_{j=1}^{2n-1} \frac{\sin((k_j - k_{j+1})\eta/2)}{(k_j - k_{j+1}) \sin(\eta/2)} d\eta. \tag{3-4}$$

We again define $I'_n = 0$ if $k_j = k_{j+1}$ for any $j = 1, \dots, 2n - 1$. In the proofs of the a priori estimates in this section we will frequently use the following lemma.

Lemma 3.2. *We recall (3-2), (3-3) and (3-4). Then, the following uniform bounds hold:*

$$\begin{aligned} |I_n(k, k_1, \dots, k_{2n})| &\leq 2\pi, \\ |I'_n(k_1, \dots, k_{2n})| &\leq 2\pi. \end{aligned}$$

This lemma will be proven at the end of this section.

Proof of Proposition 3.1. We first take a derivative of $\mathcal{N}(\mathbf{X}_c, \mathbf{X})(\theta)$ in (2-16) and let

$$\partial_\theta \mathcal{N}(\mathbf{X}_c, \mathbf{X})(\theta) = \mathcal{N}_1(\theta) + \mathcal{N}_2(\theta) + \mathcal{N}_3(\theta) + \mathcal{N}_4(\theta), \tag{3-5}$$

where

$$\begin{aligned} \mathcal{N}_1(\theta) &= \int_{\mathbb{S}} \partial_\theta (G_L(\Delta_\eta \mathbf{X}_c(\theta), \Delta_\eta \mathbf{X}(\theta))) \mathbf{F}_L(\eta) d\eta, \\ \mathcal{N}_2(\theta) &= \int_{\mathbb{S}} \partial_\theta (G_N(\Delta_\eta \mathbf{X}_c(\theta), \Delta_\eta \mathbf{X}(\theta))) \mathbf{F}_0(\eta) d\eta, \\ \mathcal{N}_3(\theta) &= \int_{\mathbb{S}} \partial_\theta (G_N(\Delta_\eta \mathbf{X}_c(\theta), \Delta_\eta \mathbf{X}(\theta))) \mathbf{F}_L(\eta) d\eta, \\ \mathcal{N}_4(\theta) &= \int_{\mathbb{S}} \partial_\theta (G(\mathcal{X}(\theta) - \mathcal{X}(\eta))) \mathbf{F}_N(\eta) d\eta. \end{aligned}$$

We will bound \mathcal{N}_i in $\mathcal{F}_v^{0,1}$ for $i = 1, 2, 3, 4$.

\mathcal{N}_1 estimates: Taking a derivative in (2-13), we obtain

$$\mathcal{N}_1(\theta) = \sum_{i=0}^{10} \mathcal{N}_{1,i}(\theta), \tag{3-6}$$

and we proceed to bound each of these terms in $\mathcal{F}_v^{0,1}$. We note that each term $\mathcal{N}_{1,i}$ corresponds to when the derivative hits a different term inside (2-13). The terms $\mathcal{N}_{1,i}$ are written in (3-8), (3-15), (3-16), (3-19), and (3-20) in the following.

The first term $\mathcal{N}_{1,1}(\theta)$ is given by

$$\mathcal{N}_{1,1}(\theta) = -\frac{1}{4\pi R^2} \int_{\mathbb{S}} \partial_\theta \Delta_\eta \mathbf{X}_c(\theta) \cdot \Delta_\eta \mathbf{X}(\theta) \mathbf{F}_L(\eta) d\eta.$$

We first take the derivative of $\Delta_\eta \mathbf{X}_c(\theta)$ in (2-2) to obtain

$$\partial_\theta \Delta_\eta \mathbf{X}_c(\theta) = \frac{\partial_\theta \mathbf{X}_c(\theta) - \frac{(\mathbf{X}_c(\theta) - \mathbf{X}_c(\eta))}{2 \tan((\theta - \eta)/2)}}{2 \sin((\theta - \eta)/2)}.$$

Further define the operator $\mathcal{D}^2(\mathbf{X}_c)$ (and analogously $\mathcal{D}^2(\mathbf{X})$) to be $\partial_\theta \Delta_\eta \mathbf{X}_c(\theta)$ as above after taking the change of variables $\eta \leftarrow \theta - \eta$ as follows:

$$\mathcal{D}^2(\mathbf{X}_c)(\theta, \eta) \stackrel{\text{def}}{=} \frac{\partial_\theta \mathbf{X}_c(\theta) - \frac{\mathbf{X}_c(\theta) - \mathbf{X}_c(\theta - \eta)}{2 \tan(\eta/2)}}{2 \sin(\eta/2)}. \tag{3-7}$$

Then we make the change of variables $\eta \leftarrow \theta - \eta$ to obtain

$$\mathcal{N}_{1,1}(\theta) = -\frac{1}{4\pi R^2} \int_{\mathbb{S}} \mathcal{D}^2(\mathbf{X}_c)(\theta, \eta)^T \Delta_{\theta-\eta} \mathbf{X}(\theta) \mathbf{F}_L(\theta - \eta) d\eta, \tag{3-8}$$

where we used transpose notation instead of a dot for future convenience in the notation. We will also make extensive use of the identities

$$\begin{aligned} \widehat{\Delta_{\theta-\eta} \mathbf{X}}(k) &= \frac{1 - e^{-ik\eta}}{2 \sin(\eta/2)} \widehat{\mathbf{X}}(k) = \frac{\sin(k\eta/2)}{k \sin(\eta/2)} e^{-ik\eta/2} \widehat{\partial_\theta \mathbf{X}}(k), \\ \widehat{\Delta_{\theta-\eta} \mathbf{X}_c}(k) &= \frac{1 - e^{-ik\eta}}{2 \sin(\eta/2)} \widehat{\mathbf{X}_c}(k) = \frac{\sin(k\eta/2)}{k \sin(\eta/2)} e^{-ik\eta/2} \widehat{\partial_\theta \mathbf{X}_c}(k). \end{aligned} \tag{3-9}$$

We remark that both terms above are equal to 0 when $k = 0$. We further have

$$\widehat{\mathcal{D}^2(\mathbf{X}_c)}(k) = m(k, \eta) \widehat{\partial_\theta \mathbf{X}_c}(k), \tag{3-10}$$

where $m(k, \eta)$ is given by (3-2).

Regarding the Fourier coefficients of the derivative of the circle (2-1) we have

$$\widehat{\partial_\theta \mathbf{X}_c}(k) = \frac{a + ib}{2} \delta_1(k) \begin{bmatrix} i \\ 1 \end{bmatrix} - \frac{a - ib}{2} \delta_{-1}(k) \begin{bmatrix} i \\ -1 \end{bmatrix}. \tag{3-11}$$

Taking Fourier transform in (3-8), we obtain

$$\begin{aligned} \widehat{\mathcal{N}}_{1,1}(k) &= -\frac{1}{4\pi R^2} \int_{\mathbb{S}} \widehat{\mathcal{D}^2(\mathbf{X}_c)}(k)^T * \widehat{\Delta_{\theta-\eta} \mathbf{X}}(k) * e^{-ik\eta} \widehat{\mathbf{F}}_L(k) d\eta \\ &= -\frac{1}{4\pi R^2} \int_{\mathbb{S}} \sum_{k_1 \in \mathbb{Z}} \sum_{k_2 \in \mathbb{Z}} \widehat{\mathcal{D}^2(\mathbf{X}_c)}(k - k_1)^T \widehat{\Delta_{\theta-\eta} \mathbf{X}}(k_1 - k_2) e^{-ik_2\eta} \widehat{\mathbf{F}}_L(k_2) d\eta, \end{aligned}$$

and plugging in (3-9) and (3-10) we have

$$\widehat{\mathcal{N}}_{1,1}(k) = -\frac{1}{4\pi R^2} \sum_{k_1 \in \mathbb{Z}} \sum_{k_2 \in \mathbb{Z}} \widehat{\partial_\theta \mathbf{X}}_c(k - k_1)^T \widehat{\partial_\theta \mathbf{X}}(k_1 - k_2) \widehat{\mathbf{F}}_L(k_2) I_1(k, k_1, k_2),$$

with I_1 given by (3-3). By Lemma 3.2 we have $|I_1(k, k_1, k_2)| \leq 2\pi$. Then we get

$$|\widehat{\mathcal{N}}_{1,1}(k)| \leq \frac{1}{2R^2} \sum_{k_1 \in \mathbb{Z}} \sum_{k_2 \in \mathbb{Z}} |\widehat{\partial_\theta \mathbf{X}}_c(k - k_1)^T \widehat{\partial_\theta \mathbf{X}}(k_1 - k_2)| |\widehat{\mathbf{F}}_L(k_2)|. \tag{3-12}$$

Then, it follows from (3-11) that

$$|\widehat{\partial_\theta \mathbf{X}}_c(k - k_1)| \leq \frac{R}{\sqrt{2}} \delta_{1,-1}(k - k_1). \tag{3-13}$$

We will now also use the notation (1-22). In particular we have

$$|\widehat{\partial_\theta \mathbf{X}}_c(k - k_1)^T \widehat{\partial_\theta \mathbf{X}}(k_1 - k_2)| \leq \frac{\sqrt{2}}{2} R (\delta_1(k - k_1) + \delta_{-1}(k - k_1)) |\widehat{\partial_\theta \mathbf{X}}(k_1 - k_2)|.$$

Therefore, we can write

$$|\widehat{\mathcal{N}}_{1,1}(k)| \leq \frac{\sqrt{2}}{4R} \sum_{k_1 \in \mathbb{Z}} \sum_{k_2 \in \mathbb{Z}} \delta_{1,-1}(k - k_1) |\widehat{\partial_\theta \mathbf{X}}(k_1 - k_2)| |\widehat{\mathbf{F}}_L(k_2)|.$$

We multiply by $e^{\nu(t)k} = e^{\nu(t)(k-k_1)} e^{\nu(t)(k_1-k_2)} e^{\nu(t)k_2}$ to get

$$e^{\nu(t)k} |\widehat{\mathcal{N}}_{1,1}(k)| \leq \frac{\sqrt{2}}{4R} \sum_{k_1 \in \mathbb{Z}} \sum_{k_2 \in \mathbb{Z}} e^{\nu(t)(k-k_1)} \delta_{1,-1}(k - k_1) e^{\nu(t)(k_1-k_2)} |\widehat{\partial_\theta \mathbf{X}}(k_1 - k_2)| e^{\nu(t)k_2} |\widehat{\mathbf{F}}_L(k_2)|,$$

so Young’s inequality for convolutions and the estimate (2-42) yield the bound

$$\|\mathcal{N}_{1,1}\|_{\mathcal{F}_\nu^{0,1}} \leq \frac{e^{\nu_\infty} \sqrt{2} \|\mathbf{X}\|_{\dot{\mathcal{F}}_\nu^{1,1}}}{2\sqrt{1 - \frac{1}{2} \|\mathbf{X}\|_{\dot{\mathcal{F}}_\nu^{1,1}}^2}} \|\mathbf{F}_L\|_{\mathcal{F}_\nu^{0,1}}. \tag{3-14}$$

This is our desired estimate for $\mathcal{N}_{1,1}$.

We now proceed to estimate $\mathcal{N}_{1,2}$ as

$$\mathcal{N}_{1,2}(\theta) = -\frac{1}{4\pi R^2} \int_{\mathbb{S}} \Delta_{\theta-\eta} \mathbf{X}_c(\theta)^T \mathcal{D}^2(\mathbf{X})(\theta, \eta) \mathbf{F}_L(\theta - \eta) d\eta, \tag{3-15}$$

with Fourier transform given by

$$\widehat{\mathcal{N}}_{1,2}(k) = -\frac{1}{4\pi R^2} \int_{\mathbb{S}} \sum_{k_1 \in \mathbb{Z}} \sum_{k_2 \in \mathbb{Z}} \widehat{\Delta_{\theta-\eta} \mathbf{X}}_c(k_1 - k_2)^T \widehat{\mathcal{D}^2(\mathbf{X})}(k - k_1) e^{-ik_2\eta} \widehat{\mathbf{F}}_L(k_2) d\eta.$$

Using again (3-9) and (3-10), we can write it as

$$\widehat{\mathcal{N}}_{1,2}(k) = -\frac{1}{4\pi R^2} \sum_{k_1 \in \mathbb{Z}} \sum_{k_2 \in \mathbb{Z}} \widehat{\partial_\theta \mathbf{X}}_c(k_1 - k_2)^T \widehat{\partial_\theta \mathbf{X}}(k - k_1) \widehat{\mathbf{F}}_L(k_2) I_1(k, k_1, k_2),$$

with I_1 given by (3-3). Using Lemma 3.2, we find that

$$|\widehat{\mathcal{N}}_{1,2}(k)| \leq \frac{1}{2R^2} \sum_{k_1 \in \mathbb{Z}} \sum_{k_2 \in \mathbb{Z}} |\widehat{\partial_\theta \mathbf{X}}_c(k-k_1)^T \widehat{\partial_\theta \mathbf{X}}(k_1-k_2)| |\widehat{\mathbf{F}}_L(k_2)|,$$

so following the steps after (3-12) we conclude that

$$\|\mathcal{N}_{1,2}\|_{\mathcal{F}_v^{0,1}} \leq \frac{e^{\nu_\infty} \sqrt{2} \|\mathbf{X}\|_{\mathcal{F}_v^{1,1}}}{2\sqrt{1 - \frac{1}{2} \|\mathbf{X}\|_{\mathcal{F}_v^{1,1}}^2}} \|\mathbf{F}_L\|_{\mathcal{F}_v^{0,1}}.$$

This completes our bound for $\mathcal{N}_{1,2}$.

The term $\mathcal{N}_{1,3}$ is given by

$$\mathcal{N}_{1,3}(\theta) = -\frac{1}{R^4} \int_{\mathcal{S}} \mathcal{D}^2(\mathbf{X}_c)(\theta, \eta)^T \Delta_{\theta-\eta} \mathbf{X}(\theta) \Delta_{\theta-\eta} \mathbf{X}_c(\theta) \otimes \Delta_{\theta-\eta} \mathbf{X}_c(\theta) \mathbf{F}_L(\theta - \eta) \frac{d\eta}{2\pi}, \quad (3-16)$$

and its Fourier transform by

$$\widehat{\mathcal{N}}_{1,3}(k) = -\frac{1}{2\pi \widehat{\partial_\theta \mathbf{X}}_c(k_2-k_3) R^4} \sum_{k_1 \in \mathbb{Z}} \cdots \sum_{k_4 \in \mathbb{Z}} \widehat{\partial_\theta \mathbf{X}}_c(k-k_1)^T \widehat{\partial_\theta \mathbf{X}}(k_1-k_2) \otimes \widehat{\partial_\theta \mathbf{X}}_c(k_3-k_4) \widehat{\mathbf{F}}_L(k_4) I_2(k, k_1, \dots, k_4),$$

with $I_2(k, k_1, \dots, k_4)$ given by (3-3). Since $|I_2(k, \dots, k_4)| \leq 2\pi$ from Lemma 3.2, we have

$$|\widehat{\mathcal{N}}_{1,3}(k)| \leq \frac{1}{R^4} \sum_{k_1 \in \mathbb{Z}} \cdots \sum_{k_4 \in \mathbb{Z}} |\widehat{\partial_\theta \mathbf{X}}_c(k-k_1)^T \widehat{\partial_\theta \mathbf{X}}(k_1-k_2)| \|\widehat{\partial_\theta \mathbf{X}}_c(k_2-k_3) \otimes \widehat{\partial_\theta \mathbf{X}}_c(k_3-k_4)\| |\widehat{\mathbf{F}}_L(k_4)|. \quad (3-17)$$

Expression (3-11) gives

$$\begin{aligned} & \widehat{\partial_\theta \mathbf{X}}_c(k_2-k_3) \otimes \widehat{\partial_\theta \mathbf{X}}_c(k_3-k_4) \\ &= \frac{(a+ib)^2}{4} \delta_1(k_2-k_3) \delta_1(k_3-k_4) \begin{bmatrix} -1 & i \\ i & 1 \end{bmatrix} + \frac{(a-ib)^2}{4} \delta_{-1}(k_2-k_3) \delta_{-1}(k_3-k_4) \begin{bmatrix} -1 & -i \\ -i & 1 \end{bmatrix} \\ & \quad - \frac{(a+ib)(a-ib)}{4} \delta_1(k_2-k_3) \delta_{-1}(k_3-k_4) \begin{bmatrix} -1 & -i \\ i & -1 \end{bmatrix} - \frac{(a+ib)(a-ib)}{4} \delta_{-1}(k_2-k_3) \delta_1(k_3-k_4) \begin{bmatrix} -1 & i \\ -i & -1 \end{bmatrix}. \end{aligned}$$

All the matrices above have norm equal to 2, so that

$$\|\widehat{\partial_\theta \mathbf{X}}_c(k_2-k_3) \otimes \widehat{\partial_\theta \mathbf{X}}_c(k_3-k_4)\| \leq \frac{R^2}{2} \delta_{1,-1}(k_2-k_3) \delta_{1,-1}(k_3-k_4). \quad (3-18)$$

Introducing this bound, together with (3-13), back to (3-17), we find that

$$|\widehat{\mathcal{N}}_{1,3}(k)| \leq \frac{\sqrt{2}}{4R} \sum_{k_1 \in \mathbb{Z}} \cdots \sum_{k_4 \in \mathbb{Z}} \delta_{1,-1}(k-k_1) |\widehat{\partial_\theta \mathbf{X}}(k_1-k_2)| \delta_{1,-1}(k_2-k_3) \delta_{1,-1}(k_3-k_4) |\widehat{\mathbf{F}}_L(k_4)|;$$

thus multiplication by the exponential $e^{\nu(t)k}$, Young's inequality and (2-42) yield that

$$\|\mathcal{N}_{1,3}\|_{\mathcal{F}_v^{0,1}} \leq \frac{2\sqrt{2}e^{3\nu_\infty} \|\mathbf{X}\|_{\mathcal{F}_v^{1,1}}}{\sqrt{1 - \frac{1}{2} \|\mathbf{X}\|_{\mathcal{F}_v^{1,1}}^2}} \|\mathbf{F}_L\|_{\mathcal{F}_v^{0,1}}.$$

This completes our bound for $\mathcal{N}_{1,3}$.

The term $\mathcal{N}_{1,4}$ is given by

$$\mathcal{N}_{1,4}(\theta) = -\frac{1}{R^4} \int_{\mathbb{S}} \Delta_{\theta-\eta} \mathbf{X}_c(\theta)^T \mathcal{D}^2(\mathbf{X})(\theta, \eta) \Delta_{\theta-\eta} \mathbf{X}_c(\theta) \otimes \Delta_{\theta-\eta} \mathbf{X}_c(\theta) \mathbf{F}_L(\theta - \eta) \frac{d\eta}{2\pi}. \quad (3-19)$$

We take the Fourier transform and write the result as

$$\widehat{\mathcal{N}}_{1,4}(k) = -\frac{1}{2\pi R^4} \sum_{k_1 \in \mathbb{Z}} \cdots \sum_{k_4 \in \mathbb{Z}} \widehat{\partial_\theta \mathbf{X}_c}(k_1 - k_2)^T \widehat{\partial_\theta \mathbf{X}}(k - k_1) \widehat{\partial_\theta \mathbf{X}_c}(k_2 - k_3) \otimes \widehat{\partial_\theta \mathbf{X}_c}(k_3 - k_4) \widehat{\mathbf{F}}_L(k_4) I_2(k, k_1, \dots, k_4),$$

with $I_2(k, k_1, \dots, k_4)$ given by (3-3). Since $|I_2| \leq 2\pi$ by Lemma 3.2, comparing now with (3-17), we conclude that

$$\|\mathcal{N}_{1,4}\|_{\mathcal{F}_v^{0,1}} \leq \frac{2\sqrt{2}e^{3v_\infty} \|\mathbf{X}\|_{\mathcal{F}_v^{1,1}}}{\sqrt{1 - \frac{1}{2} \|\mathbf{X}\|_{\mathcal{F}_v^{1,1}}^2}} \|\mathbf{F}_L\|_{\mathcal{F}_v^{0,1}}.$$

This completes our estimate for $\mathcal{N}_{1,4}$.

The remaining terms from $\mathcal{N}_1(\theta)$ in (3-6) are

$$\begin{aligned} \mathcal{N}_{1,5}(\theta) &= -\frac{1}{R^4} \int_{\mathbb{S}} \Delta_{\theta-\eta} \mathbf{X}_c(\theta)^T \Delta_{\theta-\eta} \mathbf{X}(\theta) \mathcal{D}^2(\mathbf{X}_c)(\theta, \eta) \otimes \Delta_{\theta-\eta} \mathbf{X}_c(\theta) \mathbf{F}_L(\theta - \eta) \frac{d\eta}{2\pi}, \\ \mathcal{N}_{1,6}(\theta) &= -\frac{1}{R^4} \int_{\mathbb{S}} \Delta_{\theta-\eta} \mathbf{X}_c(\theta)^T \Delta_{\theta-\eta} \mathbf{X}(\theta) \Delta_{\theta-\eta} \mathbf{X}_c(\theta) \otimes \mathcal{D}^2(\mathbf{X}_c)(\theta, \eta) \mathbf{F}_L(\theta - \eta) \frac{d\eta}{2\pi}, \\ \mathcal{N}_{1,7}(\theta) &= \frac{1}{4\pi R^2} \int_{\mathbb{S}} \mathcal{D}^2(\mathbf{X}_c)(\theta, \eta) \otimes \Delta_{\theta-\eta} \mathbf{X}(\theta) \mathbf{F}_L(\theta - \eta) d\eta, \\ \mathcal{N}_{1,8}(\theta) &= \frac{1}{4\pi R^2} \int_{\mathbb{S}} \Delta_{\theta-\eta} \mathbf{X}_c(\theta) \otimes \mathcal{D}^2(\mathbf{X})(\theta, \eta) \mathbf{F}_L(\theta - \eta) d\eta, \\ \mathcal{N}_{1,9}(\theta) &= \frac{1}{4\pi R^2} \int_{\mathbb{S}} \mathcal{D}^2(\mathbf{X})(\theta, \eta) \otimes \Delta_{\theta-\eta} \mathbf{X}_c(\theta) \mathbf{F}_L(\theta - \eta) d\eta, \\ \mathcal{N}_{1,10}(\theta) &= \frac{1}{4\pi R^2} \int_{\mathbb{S}} \Delta_{\theta-\eta} \mathbf{X}(\theta) \otimes \mathcal{D}^2(\mathbf{X}_c)(\theta, \eta) \mathbf{F}_L(\theta - \eta) d\eta. \end{aligned} \quad (3-20)$$

It is not hard to see that $\mathcal{N}_{1,5}$ and $\mathcal{N}_{1,6}$ are bounded exactly as $\mathcal{N}_{1,3}$ in (3-16), since the bound (3-18) is also valid for $\mathcal{D}^2(\mathbf{X}_c)(\theta, \eta) \otimes \Delta_{\theta-\eta} \mathbf{X}_c(\theta)$ or $\Delta_{\theta-\eta} \mathbf{X}_c(\theta) \otimes \mathcal{D}^2(\mathbf{X}_c)(\theta, \eta)$.

We proceed then with $\mathcal{N}_{1,7}$. Comparing with $\mathcal{N}_{1,1}$ in (3-8), (3-12), we obtain

$$|\widehat{\mathcal{N}}_{1,7}(k)| \leq \frac{1}{2R^2} \sum_{k_1 \in \mathbb{Z}} \sum_{k_2 \in \mathbb{Z}} \|\widehat{\partial_\theta \mathbf{X}_c}(k - k_1) \otimes \widehat{\partial_\theta \mathbf{X}}(k_1 - k_2)\| |\widehat{\mathbf{F}}_L(k_2)|.$$

Using (3-11), we find that

$$\begin{aligned} \|\widehat{\partial_\theta \mathbf{X}_c}(k - k_1) \otimes \widehat{\partial_\theta \mathbf{X}}(k_1 - k_2)\| &\leq \frac{R}{2} \delta_1(k - k_1) \left\| \begin{bmatrix} -1 \\ i \end{bmatrix} \widehat{\partial_\theta \mathbf{X}}(k_1 - k_2) \right\| + \frac{R}{2} \delta_{-1}(k - k_1) \left\| \begin{bmatrix} -1 \\ -i \end{bmatrix} \widehat{\partial_\theta \mathbf{X}}(k_1 - k_2) \right\| \\ &\leq \frac{\sqrt{2}}{2} R \delta_{1,-1}(k - k_1) |\widehat{\partial_\theta \mathbf{X}}(k_1 - k_2)|, \end{aligned} \quad (3-21)$$

where in the last inequality we have used that the matrix norm (1-19) is bounded by the Frobenius norm. Therefore we conclude that

$$\|\mathcal{N}_{1,7}\|_{\mathcal{F}_v^{0,1}} \leq \frac{\sqrt{2}e^{\nu_\infty} \|\mathbf{X}\|_{\dot{\mathcal{F}}_v^{1,1}}}{2\sqrt{1-\frac{1}{2}}\|\mathbf{X}\|_{\dot{\mathcal{F}}_v^{1,1}}^2} \|\mathbf{F}_L\|_{\mathcal{F}_v^{0,1}}.$$

The bound for $\mathcal{N}_{1,8}$ follows in the same way as that of $\mathcal{N}_{1,7}$:

$$\|\mathcal{N}_{1,8}\|_{\mathcal{F}_v^{0,1}} \leq \frac{\sqrt{2}e^{\nu_\infty} \|\mathbf{X}\|_{\dot{\mathcal{F}}_v^{1,1}}}{2\sqrt{1-\frac{1}{2}}\|\mathbf{X}\|_{\dot{\mathcal{F}}_v^{1,1}}^2} \|\mathbf{F}_L\|_{\mathcal{F}_v^{0,1}}.$$

Finally, the bounds for $\mathcal{N}_{1,9}$ and $\mathcal{N}_{1,10}$ are the same as for $\mathcal{N}_{1,7}$ and $\mathcal{N}_{1,8}$ because

$$\begin{aligned} & \|\widehat{\partial_\theta \mathbf{X}}(k-k_1) \otimes \widehat{\partial_\theta \mathbf{X}}_c(k-k_2)\| \\ & \leq \frac{R}{2} \delta_1(k_1-k_2) \|\widehat{\partial_\theta \mathbf{X}}(k-k_1)[-1 \ i]\| + \frac{R}{2} \delta_{-1}(k_1-k_2) \|\widehat{\partial_\theta \mathbf{X}}(k-k_1)[-1 \ -i]\| \\ & \leq \frac{\sqrt{2}}{2} R \delta_{1,-1}(k_1-k_2) |\widehat{\partial_\theta \mathbf{X}}(k-k_1)|. \end{aligned} \quad (3-22)$$

Joining the bounds for $\mathcal{N}_{1,1}$ to $\mathcal{N}_{1,10}$, we obtain the bound for \mathcal{N}_1 in (3-6) as

$$\|\mathcal{N}_1\|_{\mathcal{F}_v^{0,1}} \leq 11\sqrt{2}e^{3\nu_\infty} C_1 \|\mathbf{X}\|_{\dot{\mathcal{F}}_v^{1,1}} \|\mathbf{F}_L\|_{\mathcal{F}_v^{0,1}}, \quad (3-23)$$

where C_1 is defined in (2-43). This completes our estimates for the \mathcal{N}_1 term.

\mathcal{N}_3 estimates: Taking a derivative in (2-14), we split \mathcal{N}_3 as

$$\mathcal{N}_3(\theta) = \sum_{i=1}^{11} \mathcal{N}_{3,i}, \quad (3-24)$$

where

$$\begin{aligned} \mathcal{N}_{3,1}(\theta) &= -\frac{1}{4\pi} \int_{\mathbb{S}} \partial_\theta \mathcal{R}_1(\Delta_\eta \mathbf{X}(\theta)) \mathbf{F}_L(\eta) d\eta, \\ \mathcal{N}_{3,2}(\theta) &= \frac{1}{4\pi R^2} \int_{\mathbb{S}} \partial_\theta (\Delta_\eta \mathbf{X}(\theta) \otimes \Delta_\eta \mathbf{X}(\theta)) \left(1 - \frac{2}{R^2} \Delta_\eta \mathbf{X}_c(\theta)^T \Delta_\eta \mathbf{X}(\theta)\right) \mathbf{F}_L(\eta) d\eta, \\ \mathcal{N}_{3,3}(\theta) &= -\frac{1}{2\pi R^4} \int_{\mathbb{S}} \Delta_\eta \mathbf{X}(\theta) \otimes \Delta_\eta \mathbf{X}(\theta) \partial_\theta (\Delta_\eta \mathbf{X}_c(\theta)^T \Delta_\eta \mathbf{X}(\theta)) \mathbf{F}_L(\eta) d\eta, \\ \mathcal{N}_{3,4}(\theta) &= \frac{1}{4\pi R^2} \int_{\mathbb{S}} \partial_\theta (\Delta_\eta \mathbf{X}(\theta) \otimes \Delta_\eta \mathbf{X}(\theta)) \mathcal{R}_2(\Delta_\eta \mathbf{X}(\theta)) \mathbf{F}_L(\eta) d\eta, \\ \mathcal{N}_{3,5}(\theta) &= \frac{1}{4\pi R^2} \int_{\mathbb{S}} \Delta_\eta \mathbf{X}(\theta) \otimes \Delta_\eta \mathbf{X}(\theta) \partial_\theta \mathcal{R}_2(\Delta_\eta \mathbf{X}(\theta)) \mathbf{F}_L(\eta) d\eta, \\ \mathcal{N}_{3,6}(\theta) &= -\frac{1}{2\pi R^4} \int_{\mathbb{S}} \partial_\theta (\Delta_\eta \mathbf{X}_c(\theta) \otimes \Delta_\eta \mathbf{X}(\theta) + \Delta_\eta \mathbf{X}(\theta) \otimes \Delta_\eta \mathbf{X}_c(\theta)) \Delta_\eta \mathbf{X}_c(\theta)^T \Delta_\eta \mathbf{X}(\theta) \mathbf{F}_L(\eta) d\eta, \\ \mathcal{N}_{3,7}(\theta) &= -\frac{1}{2\pi R^4} \int_{\mathbb{S}} (\Delta_\eta \mathbf{X}_c(\theta) \otimes \Delta_\eta \mathbf{X}(\theta) + \Delta_\eta \mathbf{X}(\theta) \otimes \Delta_\eta \mathbf{X}_c(\theta)) \partial_\theta (\Delta_\eta \mathbf{X}_c(\theta)^T \Delta_\eta \mathbf{X}(\theta)) \mathbf{F}_L(\eta) d\eta, \\ \mathcal{N}_{3,8}(\theta) &= \frac{1}{4\pi R^2} \int_{\mathbb{S}} \partial_\theta (\Delta_\eta \mathbf{X}_c(\theta) \otimes \Delta_\eta \mathbf{X}(\theta) + \Delta_\eta \mathbf{X}(\theta) \otimes \Delta_\eta \mathbf{X}_c(\theta)) \mathcal{R}_2(\Delta_\eta \mathbf{X}(\theta)) \mathbf{F}_L(\eta) d\eta, \end{aligned}$$

$$\begin{aligned} \mathcal{N}_{3,9}(\theta) &= \frac{1}{4\pi R^2} \int_{\mathbb{S}} (\Delta_\eta \mathbf{X}_c(\theta) \otimes \Delta_\eta \mathbf{X}(\theta) + \Delta_\eta \mathbf{X}(\theta) \otimes \Delta_\eta \mathbf{X}_c(\theta)) \partial_\theta \mathcal{R}_2(\Delta_\eta \mathbf{X}(\theta)) \mathbf{F}_L(\eta) d\eta, \\ \mathcal{N}_{3,10}(\theta) &= \frac{1}{4\pi R^2} \int_{\mathbb{S}} \partial_\theta (\Delta_\eta \mathbf{X}_c(\theta) \otimes \Delta_\eta \mathbf{X}_c(\theta)) \mathcal{R}_2(\Delta_\eta \mathbf{X}(\theta)) \mathbf{F}_L(\eta) d\eta, \\ \mathcal{N}_{3,11}(\theta) &= \frac{1}{4\pi R^2} \int_{\mathbb{S}} \Delta_\eta \mathbf{X}_c(\theta) \otimes \Delta_\eta \mathbf{X}_c(\theta) \partial_\theta \mathcal{R}_2(\Delta_\eta \mathbf{X}(\theta)) \mathbf{F}_L(\eta) d\eta, \end{aligned}$$

where \mathcal{R}_1 and \mathcal{R}_2 were defined in (2-7) and (2-9).

We proceed with $\mathcal{N}_{3,1}$ first. We take the derivative in (2-7) to obtain

$$\mathcal{N}_{3,1}(\theta) = \mathbf{O}_1(\theta) + \mathbf{O}_2(\theta) + \mathbf{O}_3(\theta), \tag{3-25}$$

where

$$\begin{aligned} \mathbf{O}_1(\theta) &= -\frac{1}{8\pi} \int_{\mathbb{S}} \sum_{n \geq 1} \sum_{\substack{m=0 \\ n+m \geq 2}}^{n-1} \binom{n}{m} \frac{(-1)^{n-1}(n-m)}{nR^{2n}} (2\Delta_\eta \mathbf{X}_c(\theta))^T \Delta_\eta \mathbf{X}(\theta)^{n-m-1} \\ &\quad \cdot 2\partial_\theta \Delta_\eta \mathbf{X}_c(\theta)^T \Delta_\eta \mathbf{X}(\theta) |\Delta_\eta \mathbf{X}(\theta)|^{2m} \mathbf{F}_L(\eta) d\eta, \\ \mathbf{O}_2(\theta) &= -\frac{1}{8\pi} \int_{\mathbb{S}} \sum_{n \geq 1} \sum_{\substack{m=0 \\ n+m \geq 2}}^{n-1} \binom{n}{m} \frac{(-1)^{n-1}(n-m)}{nR^{2n}} (2\Delta_\eta \mathbf{X}_c(\theta))^T \Delta_\eta \mathbf{X}(\theta)^{n-m-1} \\ &\quad \cdot 2\Delta_\eta \mathbf{X}_c(\theta)^T \partial_\theta \Delta_\eta \mathbf{X}(\theta) |\Delta_\eta \mathbf{X}(\theta)|^{2m} \mathbf{F}_L(\eta) d\eta, \\ \mathbf{O}_3(\theta) &= -\frac{1}{4\pi} \int_{\mathbb{S}} \sum_{n \geq 1} \sum_{m=1}^n \binom{n}{m} \frac{(-1)^{n-1}m}{nR^{2n}} (2\Delta_\eta \mathbf{X}_c(\theta))^T \Delta_\eta \mathbf{X}(\theta)^{n-m} \\ &\quad \cdot |\Delta_\eta \mathbf{X}(\theta)|^{2(m-1)} \Delta_\eta \mathbf{X}(\theta)^T \partial_\theta \Delta_\eta \mathbf{X}(\theta) \mathbf{F}_L(\eta) d\eta. \end{aligned}$$

After performing the change of variables $\eta \leftarrow \theta - \eta$, we take Fourier transform of $\mathbf{O}_1(\theta)$ to obtain

$$\begin{aligned} \widehat{\mathbf{O}}_1(k) &= -\frac{1}{8\pi} \int_{\mathbb{S}} \sum_{n \geq 1} \sum_{\substack{m=0 \\ n+m \geq 2}}^{n-1} \binom{n}{m} \frac{(-1)^{n-1}(n-m)}{nR^{2n}} *^{n-m-1} \widehat{2\Delta_{\theta-\eta} \mathbf{X}_c(\theta)^T \Delta_{\theta-\eta} \mathbf{X}(\theta)} \\ &\quad * 2\mathcal{D}^2(\mathbf{X}_c)(\theta)^T \Delta_{\theta-\eta} \mathbf{X}(k) *^m \widehat{\Delta_{\theta-\eta} \mathbf{X}(\theta)^T \Delta_{\theta-\eta} \mathbf{X}(\theta)} * \widehat{\mathbf{F}}_L(k) d\eta. \end{aligned}$$

Using (3-9) and (3-10), we rewrite it as

$$\begin{aligned} \widehat{\mathbf{O}}_1(k) &= -\frac{1}{8\pi} \sum_{n \geq 1} \sum_{\substack{m=0 \\ n+m \geq 2}}^{n-1} \binom{n}{m} \frac{(-1)^{n-1}(n-m)}{nR^{2n}} \\ &\quad \sum_{k_1} \dots \sum_{k_{2n}} \prod_{j=0}^{n-m-2} 2\widehat{\partial_\theta \mathbf{X}_c}(k_{2j+1}-k_{2j+2})^T \widehat{\partial_\theta \mathbf{X}}(k_{2j+2}-k_{2j+3}) \\ &\quad \cdot 2\widehat{\partial_\theta \mathbf{X}_c}(k-k_1)^T \widehat{\partial_\theta \mathbf{X}}(k_{2n-2m-1}-k_{2n-2m}) \\ &\quad \prod_{j=n-m}^{n-1} \widehat{\partial_\theta \mathbf{X}}(k_{2j}-k_{2j+1})^T \widehat{\partial_\theta \mathbf{X}}(k_{2j+1}-k_{2j+2}) \widehat{\mathbf{F}}_L(k_{2n}) I_n(k, k_1, \dots, k_{2n}), \tag{3-26} \end{aligned}$$

with $|I_n(k, k_1, \dots, k_{2n})| \leq 2\pi$ given by (3-3) and using Lemma 3.2. Above we are using the convention that $\prod_{j=j_1}^{j_2} f(j) \equiv 1$ if $j_2 < j_1$. Recalling estimate (3-13), distributing the exponential factor $e^{\nu(t)k}$, and applying Young's inequality, we have

$$\|\mathbf{O}_1\|_{\mathcal{F}_v^{0,1}} \leq \frac{1}{4} \left(\sum_{n \geq 1} \sum_{\substack{m=0 \\ n+m \geq 2}}^{n-1} \binom{n}{m} \frac{(n-m)}{nR^{2n}} (2\sqrt{2})^{n-m} e^{(n-m)\nu(t)} R^{n-m} \|\mathbf{X}\|_{\mathcal{F}_v^{1,1}}^{n-m} \|\mathbf{X}\|_{\mathcal{F}_v^{1,1}}^{2m} \right) \|\mathbf{F}_L\|_{\mathcal{F}_v^{0,1}},$$

which can be summed first in m to get

$$\|\mathbf{O}_1\|_{\mathcal{F}_v^{0,1}} \leq \frac{1}{4} \sum_{n \geq 2} (2\sqrt{2})^n e^{nv(t)} \frac{\|\mathbf{X}\|_{\dot{\mathcal{F}}_v^{1,1}}^n}{R^n} \left(1 + \frac{\|\mathbf{X}\|_{\dot{\mathcal{F}}_v^{1,1}}}{2\sqrt{2}e^{v(t)}R}\right)^{n-1} \|\mathbf{F}_L\|_{\mathcal{F}_v^{0,1}}, \quad (3-27)$$

and then summed in n ,

$$\|\mathbf{O}_1\|_{\mathcal{F}_v^{0,1}} \leq \frac{2e^{2v_\infty} \left(1 + \frac{\|\mathbf{X}\|_{\dot{\mathcal{F}}_v^{1,1}}}{2\sqrt{2}e^{v_\infty}R}\right)}{1 - 2\sqrt{2}e^{v_\infty} \frac{\|\mathbf{X}\|_{\dot{\mathcal{F}}_v^{1,1}}}{R} \left(1 + \frac{\|\mathbf{X}\|_{\dot{\mathcal{F}}_v^{1,1}}}{2\sqrt{2}e^{v_\infty}R}\right)} \frac{\|\mathbf{X}\|_{\dot{\mathcal{F}}_v^{1,1}}^2}{R^2} \|\mathbf{F}_L\|_{\mathcal{F}_v^{0,1}}.$$

Using estimate (2-42) and the notation (2-43), we conclude that

$$\|\mathbf{O}_1\|_{\mathcal{F}_v^{0,1}} \leq 2e^{2v_\infty} C_2 C_1^2 \|\mathbf{X}\|_{\dot{\mathcal{F}}_v^{1,1}}^2 \|\mathbf{F}_L\|_{\mathcal{F}_v^{0,1}}, \quad (3-28)$$

with

$$C_2 = \frac{1 + \frac{1}{2\sqrt{2}}e^{-v_\infty} C_1 \|\mathbf{X}\|_{\dot{\mathcal{F}}_v^{1,1}}}{1 - 2\sqrt{2}e^{v_\infty} C_1 \|\mathbf{X}\|_{\dot{\mathcal{F}}_v^{1,1}} \left(1 + \frac{1}{2\sqrt{2}}e^{-v_\infty} C_1 \|\mathbf{X}\|_{\dot{\mathcal{F}}_v^{1,1}}\right)}, \quad (3-29)$$

where C_1 was defined in (2-43).

We proceed with \mathbf{O}_2 in (3-25). We take Fourier transform and, recalling (3-9), we obtain

$$\begin{aligned} \widehat{\mathbf{O}}_2(k) &= -\frac{1}{8\pi} \sum_{n \geq 1} \sum_{\substack{m=0 \\ n+m \geq 2}}^{n-1} \binom{n}{m} \frac{(-1)^{n-1} (n-m)}{nR^{2n}} \\ &\quad \sum_{k_1} \cdots \sum_{k_{2n}} \prod_{j=0}^{n-m-2} 2\widehat{\partial_\theta \mathbf{X}}_c(k_{2j+1} - k_{2j+2})^T \widehat{\partial_\theta \mathbf{X}}(k_{2j+2} - k_{2j+3}) \\ &\quad \cdot 2\widehat{\partial_\theta \mathbf{X}}_c(k_{2n-2m-1} - k_{2n-2m})^T \widehat{\partial_\theta \mathbf{X}}(k - k_1) \\ &\quad \prod_{j=n-m}^{n-1} \widehat{\partial_\theta \mathbf{X}}(k_{2j} - k_{2j+1})^T \widehat{\partial_\theta \mathbf{X}}(k_{2j+1} - k_{2j+2}) \widehat{\mathbf{F}}_L(k_{2n}) I_n(k, k_1, \dots, k_{2n}), \end{aligned} \quad (3-30)$$

again with $|I_n(k, k_1, \dots, k_{2n})| \leq 2\pi$ from (3-3) and Lemma 3.2. Thus, comparing (3-30) with (3-26), we find the estimate for \mathbf{O}_2 ,

$$\|\mathbf{O}_2\|_{\mathcal{F}_v^{0,1}} \leq 2e^{2v_\infty} C_2 C_1^2 \|\mathbf{X}\|_{\dot{\mathcal{F}}_v^{1,1}}^2 \|\mathbf{F}_L\|_{\mathcal{F}_v^{0,1}}, \quad (3-31)$$

with C_2 defined in (3-29) and C_1 in (2-43).

Repeating these steps for \mathbf{O}_3 , we obtain

$$\|\mathbf{O}_3\|_{\mathcal{F}_v^{0,1}} \leq \frac{1}{2} \sum_{n \geq 1} \sum_{m=1}^n \binom{n}{m} \frac{m(2\sqrt{2})^{n-m}}{nR^{2n}} e^{v(t)(n-m)} R^{n-m} \|\mathbf{X}\|_{\dot{\mathcal{F}}_v^{1,1}}^{n-m} \|\mathbf{X}\|_{\dot{\mathcal{F}}_v^{1,1}}^{2(m-1)} \|\mathbf{X}\|_{\dot{\mathcal{F}}_v^{1,1}}^2 \|\mathbf{F}_L\|_{\mathcal{F}_v^{0,1}},$$

which after summation in m the right side above becomes

$$\begin{aligned} \|\mathbf{O}_3\|_{\mathcal{F}_v^{0,1}} &\leq \frac{1}{2} \sum_{n \geq 1} \frac{1}{nR^{2n}} n(2\sqrt{2}e^{v(t)}R \|\mathbf{X}\|_{\dot{\mathcal{F}}_v^{1,1}} + \|\mathbf{X}\|_{\dot{\mathcal{F}}_v^{1,1}}^2)^{n-1} \|\mathbf{X}\|_{\dot{\mathcal{F}}_v^{1,1}}^2 \|\mathbf{F}_L\|_{\mathcal{F}_v^{0,1}} \\ &= \frac{1}{2} \sum_{n \geq 1} \frac{\|\mathbf{X}\|_{\dot{\mathcal{F}}_v^{1,1}}^{n-1}}{R^{n-1}} \left(2\sqrt{2}e^{v(t)} + \frac{\|\mathbf{X}\|_{\dot{\mathcal{F}}_v^{1,1}}}{R}\right)^{n-1} \frac{\|\mathbf{X}\|_{\dot{\mathcal{F}}_v^{1,1}}^2}{R^2} \|\mathbf{F}_L\|_{\mathcal{F}_v^{0,1}}, \end{aligned} \quad (3-32)$$

and after summation in n we have

$$\|\mathbf{O}_3\|_{\mathcal{F}_v^{0,1}} \leq \frac{1}{2} \frac{1}{1 - 2\sqrt{2}e^{\nu_\infty}(\|\mathbf{X}\|_{\dot{\mathcal{F}}_v^{1,1}}/R)(1 + e^{-\nu_\infty}(\|\mathbf{X}\|_{\dot{\mathcal{F}}_v^{1,1}}/2\sqrt{2}R))} \frac{\|\mathbf{X}\|_{\dot{\mathcal{F}}_v^{1,1}}^2}{R^2} \|\mathbf{F}_L\|_{\mathcal{F}_v^{0,1}}.$$

Introducing the bound for R in (2-42), we obtain

$$\|\mathbf{O}_3\|_{\mathcal{F}_v^{0,1}} \leq \frac{1}{2} C_3 C_1^2 \|\mathbf{X}\|_{\dot{\mathcal{F}}_v^{1,1}}^2 \|\mathbf{F}_L\|_{\mathcal{F}_v^{0,1}}, \tag{3-33}$$

and using C_2 in (3-29) and C_1 in (2-43) we have

$$C_3 = \frac{C_2}{1 + \frac{1}{2\sqrt{2}}e^{-\nu_\infty}C_1\|\mathbf{X}\|_{\dot{\mathcal{F}}_v^{1,1}}}. \tag{3-34}$$

Joining the bounds (3-28), (3-31), and (3-33), we find the estimate for $\mathcal{N}_{3,1}$ from (3-25) as

$$\|\mathcal{N}_{3,1}\|_{\mathcal{F}_v^{0,1}} \leq \frac{9}{2} C_4 C_1^2 \|\mathbf{X}\|_{\dot{\mathcal{F}}_v^{1,1}}^2 \|\mathbf{F}_L\|_{\mathcal{F}_v^{0,1}}, \tag{3-35}$$

with

$$C_4 = \frac{2}{9} (4e^{2\nu_\infty} C_2 + \frac{1}{2} C_3). \tag{3-36}$$

This completes our desired estimate for $\mathcal{N}_{3,1}$.

We continue with the next term $\mathcal{N}_{3,2}$ from (3-24), which we split in two:

$$\mathcal{N}_{3,2}(\theta) = \mathbf{O}_4(\theta) + \mathbf{O}_5(\theta),$$

where

$$\begin{aligned} \mathbf{O}_4(\theta) &= \frac{1}{4\pi R^2} \int_{\mathbb{S}} \partial_\theta (\Delta_\eta \mathbf{X}(\theta) \otimes \Delta_\eta \mathbf{X}(\theta)) \mathbf{F}_L(\eta) d\eta, \\ \mathbf{O}_5(\theta) &= -\frac{1}{2\pi R^4} \int_{\mathbb{S}} \partial_\theta (\Delta_\eta \mathbf{X}(\theta) \otimes \Delta_\eta \mathbf{X}(\theta)) \Delta_\eta \mathbf{X}_c(\theta)^T \Delta_\eta \mathbf{X}(\theta) \mathbf{F}_L(\eta) d\eta. \end{aligned}$$

The bounds for these terms follow in a similar way to that of $\mathcal{N}_{1,2}$ from (3-15) and $\mathcal{N}_{1,4}$ from (3-19), respectively. Taking into account that

$$\|\widehat{\partial_\theta \mathbf{X}}(k - k_1) \otimes \widehat{\partial_\theta \mathbf{X}}(k_1 - k_2)\| \leq |\widehat{\partial_\theta \mathbf{X}}(k - k_1)| |\widehat{\partial_\theta \mathbf{X}}(k_1 - k_2)|, \tag{3-37}$$

and Lemma 3.2, it is not hard to find that

$$|\widehat{\mathbf{O}}_4(k)| \leq \frac{1}{R^2} \sum_{k_1 \in \mathbb{Z}} \sum_{k_2 \in \mathbb{Z}} |\widehat{\partial_\theta \mathbf{X}}(k - k_1)| |\widehat{\partial_\theta \mathbf{X}}(k_1 - k_2)| |\widehat{\mathbf{F}}_L(k_2)|,$$

and recalling (3-13), we have

$$|\widehat{\mathbf{O}}_5(k)| \leq \frac{\sqrt{2}}{R^3} \sum_{k_1 \in \mathbb{Z}} \cdots \sum_{k_4 \in \mathbb{Z}} |\widehat{\partial_\theta \mathbf{X}}(k - k_1)| |\widehat{\partial_\theta \mathbf{X}}(k_1 - k_2)| |\delta_{1,-1}(k_2 - k_3)| |\widehat{\partial_\theta \mathbf{X}}(k_3 - k_4)| |\widehat{\mathbf{F}}_L(k_4)|.$$

Therefore,

$$\|\mathbf{O}_4\|_{\mathcal{F}_v^{0,1}} \leq \frac{\|\mathbf{X}\|_{\dot{\mathcal{F}}_v^{1,1}}^2}{R^2} \|\mathbf{F}_L\|_{\mathcal{F}_v^{0,1}}, \quad \|\mathbf{O}_5\|_{\dot{\mathcal{F}}_v^{1,1}} \leq 2\sqrt{2}e^{\nu_\infty} \frac{\|\mathbf{X}\|_{\dot{\mathcal{F}}_v^{1,1}}^3}{R^3} \|\mathbf{F}_L\|_{\mathcal{F}_v^{0,1}};$$

thus

$$\|\mathcal{N}_{3,2}\|_{\mathcal{F}_v^{0,1}} \leq \left(1 + 2\sqrt{2}e^{\nu_\infty} \frac{\|\mathbf{X}\|_{\dot{\mathcal{F}}_v^{1,1}}}{R}\right) \frac{\|\mathbf{X}\|_{\dot{\mathcal{F}}_v^{1,1}}^2}{R^2} \|\mathbf{F}_L\|_{\mathcal{F}_v^{0,1}},$$

so plugging in the estimate (2-42) yields that

$$\|\mathcal{N}_{3,2}\|_{\mathcal{F}_v^{0,1}} \leq C_5 C_1^2 \|\mathbf{X}\|_{\dot{\mathcal{F}}_v^{1,1}}^2 \|\mathbf{F}_L\|_{\mathcal{F}_v^{0,1}}, \quad (3-38)$$

with

$$C_5 = 1 + 2\sqrt{2}e^{\nu_\infty} C_1 \|\mathbf{X}\|_{\dot{\mathcal{F}}_v^{1,1}}. \quad (3-39)$$

This completes our estimate for $\mathcal{N}_{3,2}$.

The Fourier transform of $\mathcal{N}_{3,3}$ in (3-24) can be bounded as

$$|\widehat{\mathcal{N}}_{3,3}(k)| \leq \frac{\sqrt{2}}{2R^3} \sum_{k_1 \in \mathbb{Z}} \cdots \sum_{k_4 \in \mathbb{Z}} |\widehat{\partial_\theta \mathbf{X}}(k - k_1)| |\widehat{\partial_\theta \mathbf{X}}(k_1 - k_2)| |\delta_{1,-1}(k_2 - k_3)| |\widehat{\partial_\theta \mathbf{X}}(k_3 - k_4)| |\widehat{\mathbf{F}}_L(k_4)|,$$

and thus

$$\|\mathcal{N}_{3,3}\|_{\mathcal{F}_v^{0,1}} \leq \sqrt{2}e^{\nu_\infty} \frac{\|\mathbf{X}\|_{\dot{\mathcal{F}}_v^{1,1}}^3}{R^3} \|\mathbf{F}_L\|_{\mathcal{F}_v^{0,1}},$$

which becomes

$$\|\mathcal{N}_{3,3}\|_{\mathcal{F}_v^{0,1}} \leq \sqrt{2}e^{\nu_\infty} C_1^3 \|\mathbf{X}\|_{\dot{\mathcal{F}}_v^{1,1}}^3 \|\mathbf{F}_L\|_{\mathcal{F}_v^{0,1}}. \quad (3-40)$$

Similarly, recalling (2-9), the estimate for $\mathcal{N}_{3,4}$ in (3-24) is

$$\|\mathcal{N}_{3,4}\|_{\mathcal{F}_v^{0,1}} \leq \frac{\|\mathbf{X}\|_{\dot{\mathcal{F}}_v^{1,1}}^2}{R^2} \sum_{n \geq 1} \sum_{\substack{m=0 \\ n+m \geq 2}}^n \binom{n}{m} \frac{(2\sqrt{2})^{n-m} e^{(n-m)\nu(t)} R^{n-m}}{R^{2n}} \|\mathbf{X}\|_{\dot{\mathcal{F}}_v^{1,1}}^{n-m} \|\mathbf{X}\|_{\dot{\mathcal{F}}_v^{1,1}}^{2m} \|\mathbf{F}_L\|_{\mathcal{F}_v^{0,1}},$$

which can be rewritten as

$$\begin{aligned} & \|\mathcal{N}_{3,4}\|_{\mathcal{F}_v^{0,1}} \\ & \leq \frac{\|\mathbf{X}\|_{\dot{\mathcal{F}}_v^{1,1}}^2}{R^2} \left(\sum_{n \geq 1} \frac{(2\sqrt{2}e^{\nu(t)} R \|\mathbf{X}\|_{\dot{\mathcal{F}}_v^{1,1}})^n}{R^{2n}} \left(1 + \frac{\|\mathbf{X}\|_{\dot{\mathcal{F}}_v^{1,1}}}{2\sqrt{2}e^{\nu(t)} R}\right)^n - \frac{2\sqrt{2}e^{\nu(t)} \|\mathbf{X}\|_{\dot{\mathcal{F}}_v^{1,1}}}{R} \right) \|\mathbf{F}_L\|_{\mathcal{F}_v^{0,1}} \\ & = \frac{2\sqrt{2}e^{\nu(t)} \|\mathbf{X}\|_{\dot{\mathcal{F}}_v^{1,1}}^3}{R^3} \left(\sum_{n \geq 1} \left(\frac{2\sqrt{2}e^{\nu(t)} \|\mathbf{X}\|_{\dot{\mathcal{F}}_v^{1,1}}}{R}\right)^{n-1} \left(1 + \frac{\|\mathbf{X}\|_{\dot{\mathcal{F}}_v^{1,1}}}{2\sqrt{2}e^{\nu(t)} R}\right)^n - 1 \right) \|\mathbf{F}_L\|_{\mathcal{F}_v^{0,1}} \\ & = \frac{2\sqrt{2}e^{\nu(t)} \|\mathbf{X}\|_{\dot{\mathcal{F}}_v^{1,1}}^3}{R^3} \left(\frac{\|\mathbf{X}\|_{\dot{\mathcal{F}}_v^{1,1}}}{2\sqrt{2}e^{\nu(t)} R} \right. \\ & \quad \left. + \left(1 + \frac{\|\mathbf{X}\|_{\dot{\mathcal{F}}_v^{1,1}}}{2\sqrt{2}e^{\nu(t)} R}\right) \sum_{n \geq 2} \left(\frac{2\sqrt{2}e^{\nu(t)} \|\mathbf{X}\|_{\dot{\mathcal{F}}_v^{1,1}}}{R}\right)^{n-1} \left(1 + \frac{\|\mathbf{X}\|_{\dot{\mathcal{F}}_v^{1,1}}}{2\sqrt{2}e^{\nu(t)} R}\right)^n \right) \|\mathbf{F}_L\|_{\mathcal{F}_v^{0,1}}. \end{aligned}$$

Performing the sum in n and using estimate (2-42), we conclude that

$$\|\mathcal{N}_{3,4}\|_{\mathcal{F}_v^{0,1}} \leq 9C_6 C_1^4 \|\mathbf{X}\|_{\dot{\mathcal{F}}_v^{1,1}}^4 \|\mathbf{F}_L\|_{\mathcal{F}_v^{0,1}}, \quad (3-41)$$

with

$$C_6 = \frac{1}{9} \left(1 + 8e^{2\nu_\infty} \left(1 + \frac{1}{2\sqrt{2}} e^{-\nu_\infty} C_1 \|\mathbf{X}\|_{\dot{J}_v^{1,1}} \right) C_2 \right), \quad (3-42)$$

where C_1, C_2 were defined in (2-43), (3-29). This completes our estimate for $\mathcal{N}_{3,4}$.

To deal with the term $\mathcal{N}_{3,5}$ in (3-24), we have to take a derivative in \mathcal{R}_2 from (2-9). This gives the splitting

$$\mathcal{N}_{3,5}(\theta) = \mathbf{O}_6(\theta) + \mathbf{O}_7(\theta) + \mathbf{O}_8(\theta), \quad (3-43)$$

where

$$\begin{aligned} \mathbf{O}_6(\theta) = & \frac{1}{4\pi R^2} \int_{\mathbb{S}} \sum_{n \geq 1} \sum_{\substack{m=0 \\ n+m \geq 2}}^{n-1} \binom{n}{m} \frac{(-1)^n (n-m)}{R^{2n}} (\Delta_\eta \mathbf{X}(\theta) \otimes \Delta_\eta \mathbf{X}(\theta)) \\ & \cdot (2\Delta_\eta \mathbf{X}_c(\theta)^T \Delta_\eta \mathbf{X}(\theta))^{n-m-1} 2\partial_\theta \Delta_\eta \mathbf{X}_c(\theta)^T \Delta_\eta \mathbf{X}(\theta) |\Delta_\eta \mathbf{X}(\theta)|^{2m} \mathbf{F}_L(\eta) d\eta, \end{aligned}$$

$$\begin{aligned} \mathbf{O}_7(\theta) = & \frac{1}{4\pi R^2} \int_{\mathbb{S}} \sum_{n \geq 1} \sum_{\substack{m=0 \\ n+m \geq 2}}^{n-1} \binom{n}{m} \frac{(-1)^n (n-m)}{R^{2n}} (\Delta_\eta \mathbf{X}(\theta) \otimes \Delta_\eta \mathbf{X}(\theta)) \\ & \cdot (2\Delta_\eta \mathbf{X}_c(\theta)^T \Delta_\eta \mathbf{X}(\theta))^{n-m-1} 2\Delta_\eta \mathbf{X}_c(\theta)^T \partial_\theta \Delta_\eta \mathbf{X}(\theta) |\Delta_\eta \mathbf{X}(\theta)|^{2m} \mathbf{F}_L(\eta) d\eta, \end{aligned}$$

$$\begin{aligned} \mathbf{O}_8(\theta) = & \frac{1}{2\pi R^2} \int_{\mathbb{S}} \sum_{n \geq 1} \sum_{m=1}^n \binom{n}{m} \frac{(-1)^n m}{R^{2n}} (\Delta_\eta \mathbf{X}(\theta) \otimes \Delta_\eta \mathbf{X}(\theta)) \\ & \cdot (2\Delta_\eta \mathbf{X}_c(\theta)^T \Delta_\eta \mathbf{X}(\theta))^{n-m} |\Delta_\eta \mathbf{X}(\theta)|^{2(m-1)} \Delta_\eta \mathbf{X}(\theta)^T \partial_\theta \Delta_\eta \mathbf{X}(\theta) \mathbf{F}_L(\eta) d\eta. \end{aligned}$$

Comparing \mathbf{O}_6 and \mathbf{O}_8 to \mathbf{O}_1 and \mathbf{O}_3 , respectively, in (3-25), and recalling the bounds (3-27), (3-32), together with (3-37), we find that

$$\begin{aligned} \|\mathbf{O}_6\|_{\mathcal{F}_v^{0,1}} & \leq \frac{\|\mathbf{X}\|_{\dot{J}_v^{1,1}}^2}{2R^2} \sum_{n \geq 2} \frac{(2\sqrt{2})^n e^{n\nu(t)} n}{R^n} \|\mathbf{X}\|_{\dot{J}_v^{1,1}}^n \left(1 + \frac{\|\mathbf{X}\|_{\dot{J}_v^{1,1}}}{2\sqrt{2}e^{\nu(t)}R} \right)^{n-1} \|\mathbf{F}_L\|_{\mathcal{F}_v^{0,1}}, \\ \|\mathbf{O}_8\|_{\mathcal{F}_v^{0,1}} & \leq \frac{\|\mathbf{X}\|_{\dot{J}_v^{1,1}}^2}{R^2} \sum_{n \geq 1} \frac{n \|\mathbf{X}\|_{\dot{J}_v^{1,1}}^{n-1}}{R^{n-1}} \left(2\sqrt{2}e^{\nu(t)} + \frac{\|\mathbf{X}\|_{\dot{J}_v^{1,1}}}{R} \right)^{n-1} \frac{\|\mathbf{X}\|_{\dot{J}_v^{1,1}}^2}{R^2} \|\mathbf{F}_L\|_{\mathcal{F}_v^{0,1}}, \end{aligned}$$

which after summation in n the right side above becomes

$$\begin{aligned} \|\mathbf{O}_6\|_{\mathcal{F}_v^{0,1}} & \leq 4e^{2\nu_\infty} \left(1 + \frac{\|\mathbf{X}\|_{\dot{J}_v^{1,1}}}{2\sqrt{2}e^{\nu_\infty}R} \right) \left(2 - \frac{2\sqrt{2}e^{\nu_\infty} \|\mathbf{X}\|_{\dot{J}_v^{1,1}}}{R} \left(1 + \frac{\|\mathbf{X}\|_{\dot{J}_v^{1,1}}}{2\sqrt{2}e^{\nu_\infty}R} \right) \right) \\ & \quad \times \left(1 - 2\sqrt{2}e^{\nu_\infty} \frac{\|\mathbf{X}\|_{\dot{J}_v^{1,1}}}{R} \left(1 + \frac{\|\mathbf{X}\|_{\dot{J}_v^{1,1}}}{2\sqrt{2}e^{\nu_\infty}R} \right) \right)^{-2} \frac{\|\mathbf{X}\|_{\dot{J}_v^{1,1}}^4}{R^4} \|\mathbf{F}_L\|_{\mathcal{F}_v^{0,1}}, \\ \|\mathbf{O}_8\|_{\mathcal{F}_v^{0,1}} & \leq \left(1 - 2\sqrt{2}e^{\nu_\infty} \frac{\|\mathbf{X}\|_{\dot{J}_v^{1,1}}}{R} \left(1 + \frac{\|\mathbf{X}\|_{\dot{J}_v^{1,1}}}{2\sqrt{2}e^{\nu_\infty}R} \right) \right)^{-2} \frac{\|\mathbf{X}\|_{\dot{J}_v^{1,1}}^4}{R^4} \|\mathbf{F}_L\|_{\mathcal{F}_v^{0,1}}. \end{aligned}$$

It is now clear that, for the same reason that the bound for \mathbf{O}_2 (3-30) was the same as that for \mathbf{O}_1 (3-26), the estimate for \mathbf{O}_7 is the same as the one for \mathbf{O}_6 . Therefore, with (2-42), we conclude that

$$\|\mathcal{N}_{3,5}\|_{\mathcal{F}_v^{0,1}} \leq 17C_7C_1^4 \|\mathbf{X}\|_{\dot{J}_v^{1,1}}^4 \|\mathbf{F}_L\|_{\mathcal{F}_v^{0,1}}, \quad (3-44)$$

with

$$C_7 = \frac{16}{17}e^{2\nu_\infty}C_2C_3\left(1 - C_8^{-1}\sqrt{2}e^{\nu_\infty}\|\mathbf{X}\|_{\dot{\mathcal{F}}_v^{1,1}}\left(1 + \frac{1}{2\sqrt{2}}C_8^{-1}e^{-\nu_\infty}\|\mathbf{X}\|_{\dot{\mathcal{F}}_v^{1,1}}\right)\right) + \frac{C_3^2}{17}, \quad (3-45)$$

where we note that C_7 is indeed increasing in $\|\mathbf{X}\|_{\dot{\mathcal{F}}_v^{1,1}}$ as can be seen because the infinite sums in the upper bounds of $\|\mathbf{O}_6\|_{\mathcal{F}_v^{0,1}}$ and $\|\mathbf{O}_8\|_{\mathcal{F}_v^{0,1}}$ above are indeed increasing. Further above we are also using

$$C_8 \stackrel{\text{def}}{=} \sqrt{1 + \frac{1}{2}\|\mathbf{X}\|_{\dot{\mathcal{F}}_v^{1,1}}^2}, \quad (3-46)$$

and we are further using C_2 and C_3 from (3-29) and (3-34).

Recalling the bounds (3-21) and (3-22), the remaining terms $\mathcal{N}_{3,6} - \mathcal{N}_{3,11}$ can be estimated similarly, using also C_1 , C_6 and C_7 from (2-43), (3-42) and (3-45), to obtain

$$\begin{aligned} \|\mathcal{N}_{3,6}\|_{\dot{\mathcal{F}}_v^{1,1}} &\leq 8e^{2\nu_\infty}C_1^2\|\mathbf{X}\|_{\dot{\mathcal{F}}_v^{1,1}}^2\|\mathbf{F}_L\|_{\mathcal{F}_v^{0,1}}, \\ \|\mathcal{N}_{3,7}\|_{\dot{\mathcal{F}}_v^{1,1}} &\leq 8e^{2\nu_\infty}C_1^2\|\mathbf{X}\|_{\dot{\mathcal{F}}_v^{1,1}}^2\|\mathbf{F}_L\|_{\mathcal{F}_v^{0,1}}, \\ \|\mathcal{N}_{3,8}\|_{\dot{\mathcal{F}}_v^{1,1}} &\leq 18\sqrt{2}e^{\nu_\infty}C_6C_1^3\|\mathbf{X}\|_{\dot{\mathcal{F}}_v^{1,1}}^3\|\mathbf{F}_L\|_{\mathcal{F}_v^{0,1}}, \\ \|\mathcal{N}_{3,9}\|_{\dot{\mathcal{F}}_v^{1,1}} &\leq 34\sqrt{2}e^{\nu_\infty}C_7C_1^3\|\mathbf{X}\|_{\dot{\mathcal{F}}_v^{1,1}}^3\|\mathbf{F}_L\|_{\mathcal{F}_v^{0,1}}, \\ \|\mathcal{N}_{3,10}\|_{\dot{\mathcal{F}}_v^{1,1}} &\leq 18e^{2\nu_\infty}C_6C_1^2\|\mathbf{X}\|_{\dot{\mathcal{F}}_v^{1,1}}^2\|\mathbf{F}_L\|_{\mathcal{F}_v^{0,1}}, \\ \|\mathcal{N}_{3,11}\|_{\mathcal{F}_v^{0,1}} &\leq 34e^{2\nu_\infty}C_7C_1^2\|\mathbf{X}\|_{\dot{\mathcal{F}}_v^{1,1}}^2\|\mathbf{F}_L\|_{\mathcal{F}_v^{0,1}}. \end{aligned} \quad (3-47)$$

Therefore, from the splitting (3-24) and adding all the bounds (3-35), (3-38), (3-40), (3-41), (3-44), and (3-47), we conclude that

$$\|\mathcal{N}_3\|_{\mathcal{F}_v^{0,1}} \leq \frac{147}{2}C_9C_1^2\|\mathbf{X}\|_{\dot{\mathcal{F}}_v^{1,1}}^2\|\mathbf{F}_L\|_{\mathcal{F}_v^{0,1}}, \quad (3-48)$$

where

$$\begin{aligned} C_9 = &\frac{2}{147}\left(\frac{9}{2}C_4 + C_5 + 16e^{2\nu_\infty} + 18e^{2\nu_\infty}C_6 + 34e^{2\nu_\infty}C_7\right. \\ &\left. + (\sqrt{2} + 18\sqrt{2}C_6 + 34\sqrt{2}C_7)e^{\nu_\infty}C_1\|\mathbf{X}\|_{\dot{\mathcal{F}}_v^{1,1}} + (9C_6 + 17C_7)C_1^2\|\mathbf{X}\|_{\dot{\mathcal{F}}_v^{1,1}}^2\right), \end{aligned} \quad (3-49)$$

with C_1 , C_4 , C_5 , C_6 , and C_7 defined in (2-43), (3-36), (3-39), (3-42), and (3-45).

\mathcal{N}_2 estimates: It is clear from (3-5) that the previous estimate for \mathcal{N}_3 in (3-48) is also valid for \mathcal{N}_2 , with $\|\mathbf{F}_L\|_{\mathcal{F}_v^{0,1}}$ replaced by $\|\mathbf{F}_0\|_{\mathcal{F}_v^{0,1}}$. Therefore we have

$$\|\mathcal{N}_2\|_{\mathcal{F}_v^{0,1}} \leq \frac{147}{2}C_9C_1^2\|\mathbf{X}\|_{\dot{\mathcal{F}}_v^{1,1}}^2\|\mathbf{F}_0\|_{\mathcal{F}_v^{0,1}}, \quad (3-50)$$

with C_9 defined above in (3-49).

\mathcal{N}_4 estimates: We split the term \mathcal{N}_4 in (3-5) following the splitting (2-11):

$$\mathcal{N}_4(\theta) = \mathcal{N}_{4,1}(\theta) + \mathcal{N}_{4,2}(\theta) + \mathcal{N}_{4,3}(\theta),$$

where

$$\begin{aligned} \mathcal{N}_{4,1}(\theta) &= \int_{\mathbb{S}} \partial_{\theta}(G_0(\Delta_{\eta} \mathbf{X}_c(\theta))) \mathbf{F}_N(\eta) d\eta, \\ \mathcal{N}_{4,2}(\theta) &= \int_{\mathbb{S}} \partial_{\theta}(G_L(\Delta_{\eta} \mathbf{X}_c(\theta), \Delta_{\eta} \mathbf{X}(\theta))) \mathbf{F}_N(\eta) d\eta, \\ \mathcal{N}_{4,3}(\theta) &= \int_{\mathbb{S}} \partial_{\theta}(G_N(\Delta_{\eta} \mathbf{X}_c(\theta), \Delta_{\eta} \mathbf{X}(\theta))) \mathbf{F}_N(\eta) d\eta. \end{aligned}$$

We notice that the term $\mathcal{N}_{4,2}$ can be bounded exactly as \mathcal{N}_1 in (3-5), with $\|\mathbf{F}_L\|_{\mathcal{F}_v^{0,1}}$ replaced by $\|\mathbf{F}_N\|_{\mathcal{F}_v^{0,1}}$, that is, from (3-23) we have

$$\|\mathcal{N}_{4,2}\|_{\mathcal{F}_v^{0,1}} \leq 11\sqrt{2}e^{3v_{\infty}} C_1 \|\mathbf{X}\|_{\dot{\mathcal{F}}_v^{1,1}} \|\mathbf{F}_N\|_{\mathcal{F}_v^{0,1}},$$

with C_1 from (2-43). Analogously using the similarity between $\mathcal{N}_{4,3}$ and \mathcal{N}_3 in (3-48), we have

$$\|\mathcal{N}_{4,3}\|_{\mathcal{F}_v^{0,1}} \leq \frac{147}{2} C_9 C_1^2 \|\mathbf{X}\|_{\dot{\mathcal{F}}_v^{1,1}}^2 \|\mathbf{F}_N\|_{\mathcal{F}_v^{0,1}},$$

where we recall C_9 from (3-49).

Now taking a derivative in (2-12), the term $\mathcal{N}_{4,1}$ can be written as

$$\begin{aligned} \mathcal{N}_{4,1}(\theta) &= -\frac{1}{4\pi} \int_{\mathbb{S}} \frac{F_N(\eta)}{2 \tan((\theta - \eta)/2)} + \frac{1}{4\pi R^2} \int_{\mathbb{S}} \partial_{\theta} \Delta_{\eta} \mathbf{X}_c(\theta) \otimes \Delta_{\eta} \mathbf{X}_c(\theta) \mathbf{F}_N(\eta) d\eta \\ &\quad + \frac{1}{4\pi R^2} \int_{\mathbb{S}} \Delta_{\eta} \mathbf{X}_c(\theta) \otimes \partial_{\theta} \Delta_{\eta} \mathbf{X}_c(\theta) \mathbf{F}_N(\eta) d\eta, \end{aligned}$$

and therefore, recalling (3-18), we have

$$\|\mathcal{N}_{4,1}\|_{\mathcal{F}_v^{0,1}} \leq \left(\frac{1}{4} + 2e^{2v_{\infty}}\right) \|\mathbf{F}_N\|_{\mathcal{F}_v^{0,1}}.$$

We add the previous bounds to obtain

$$\|\mathcal{N}_4\|_{\mathcal{F}_v^{0,1}} \leq \frac{9}{4} C_{10} \|\mathbf{F}_N\|_{\mathcal{F}_v^{0,1}}, \tag{3-51}$$

with

$$C_{10} = \frac{4}{9} \left(\frac{1}{4} + 2e^{2v_{\infty}} + 11\sqrt{2}e^{3v_{\infty}} C_1 \|\mathbf{X}\|_{\dot{\mathcal{F}}_v^{1,1}} + \frac{147}{2} C_9 C_1^2 \|\mathbf{X}\|_{\dot{\mathcal{F}}_v^{1,1}}^2\right), \tag{3-52}$$

with C_1, C_9 defined in (2-43) and (3-49). Combining the estimates (3-23), (3-50), (3-48), and (3-51), we conclude from (3-5) that

$$\|\mathcal{N}\|_{\dot{\mathcal{F}}_v^{1,1}} \leq \frac{147}{2} C_9 C_1^2 \|\mathbf{X}\|_{\dot{\mathcal{F}}_v^{1,1}} \|\mathbf{F}_0\|_{\mathcal{F}_v^{0,1}} \|\mathbf{X}\|_{\dot{\mathcal{F}}_v^{2,1}} + 11\sqrt{2} C_{11} C_1 \|\mathbf{X}\|_{\dot{\mathcal{F}}_v^{1,1}} \|\mathbf{F}_L\|_{\mathcal{F}_v^{0,1}} + \frac{9}{4} C_{10} \|\mathbf{F}_N\|_{\mathcal{F}_v^{0,1}},$$

where

$$C_{11} = \frac{1}{11\sqrt{2}} \left(11\sqrt{2}e^{3v_{\infty}} + \frac{147}{2} C_9 C_1 \|\mathbf{X}\|_{\dot{\mathcal{F}}_v^{1,1}}\right), \tag{3-53}$$

and C_1, C_9 are defined in (2-43) and (3-49). Rename the constants

$$D_1 = C_{11} C_1, \quad D_2 = C_9 C_1^2, \quad D_3 = C_{10}, \tag{3-54}$$

to get the result (3-1), where C_1, C_9, C_{10} , and C_{11} are given in (2-43), (3-49), (3-52), and (3-53). \square

3B. A priori estimates on F . In this section we will obtain bounds for F_0 , F_L , and F_N in $\mathcal{F}_v^{0,1}$.

Proposition 3.3. *Assume that $X \in \dot{\mathcal{F}}_v^{2,1}$ and that F solves (1-13). Then, the functions F_0 in (3-58), F_L in (2-25) and $F_N = F - F_0 - F_L$, satisfy the estimate*

$$\|F_0\|_{\mathcal{F}_v^{0,1}} \leq \sqrt{2}e^{v_\infty} C_8 \frac{2A_e}{1 - A_\mu}, \tag{3-55}$$

where C_8 is defined in (3-46). Further

$$\|F_L\|_{\mathcal{F}_v^{0,1}} \leq 2A_e \left(1 + \frac{|A_\mu|}{1 - A_\mu}\right) \|X\|_{\dot{\mathcal{F}}_v^{2,1}}, \tag{3-56}$$

$$\|F_N\|_{\mathcal{F}_v^{0,1}} \leq 1000\sqrt{2}A_e \frac{|A_\mu|(1 + |A_\mu|)}{(1 - A_\mu)^2(1 + A_\mu)} D_4 \|X\|_{\dot{\mathcal{F}}_v^{1,1}} \|X\|_{\dot{\mathcal{F}}_v^{2,1}}, \tag{3-57}$$

where $D_4 = D_4(\|X\|_{\dot{\mathcal{F}}_v^{1,1}}; A_\mu, v_\infty)$ is an increasing function of $\|X\|_{\dot{\mathcal{F}}_v^{1,1}}$ as in (1-28) such that

$$\lim_{\|X\|_{\dot{\mathcal{F}}_v^{1,1}} \rightarrow 0^+} D_4(\|X\|_{\dot{\mathcal{F}}_v^{1,1}}; 0, 0) = 1$$

and is defined in (3-78).

Proof. First, for a general circle the expression for F_0 in (2-20) becomes

$$F_0(\theta) = \frac{2A_e}{1 - A_\mu} \partial_\theta^2 X_c. \tag{3-58}$$

Similar to (3-13) using (3-11) we have for (3-55) that

$$\|F_0\|_{\mathcal{F}_v^{0,1}} \leq \sqrt{2}e^{v_\infty} R \frac{2A_e}{1 - A_\mu} \leq \sqrt{2}e^{v_\infty} C_8 \frac{2A_e}{1 - A_\mu},$$

where C_8 is given by (3-46) and we used (2-42).

Further F_L is given by (2-25) and so we have

$$\|F_L\|_{\mathcal{F}_v^{0,1}} \leq 2A_e \|X\|_{\dot{\mathcal{F}}_v^{2,1}} + \frac{2|A_\mu|A_e}{1 - A_\mu} \|X\|_{\dot{\mathcal{F}}_v^{1,1}},$$

which gives (3-56).

We proceed with the expansion of the nonlinear terms in (1-13). First, using (2-8), we write

$$\frac{1}{|\Delta X + \Delta X_c|^4} = \frac{1}{16R^4 \sin^4((\theta - \eta)/2)} \left(1 - \frac{4}{R^2} \Delta_\eta X_c(\theta)^T \Delta_\eta X(\theta) + \mathcal{R}_3(\Delta_\eta X(\theta))\right),$$

where

$$\begin{aligned} \mathcal{R}_3(\Delta_\eta X(\theta)) = & -\frac{4}{R^2} \Delta_\eta X_c(\theta)^T \Delta_\eta X(\theta) \mathcal{R}_2(\Delta_\eta X(\theta)) + 2\mathcal{R}_2(\Delta_\eta X(\theta)) \\ & + \frac{4}{R^4} (\Delta_\eta X_c(\theta)^T \Delta_\eta X(\theta))^2 + (\mathcal{R}_2(\Delta_\eta X(\theta)))^2, \end{aligned} \tag{3-59}$$

and $\mathcal{R}_2(\Delta_\eta X(\theta))$ is given in (2-9). Then, we use the above expansion to rewrite $\mathcal{S}(F, X)(\theta)$ from (1-14) as

$$\mathcal{S}(F, X)(\theta) = \int_{\mathbb{S}} K(X_c, X)(\theta, \eta) \frac{F(\theta - \eta)}{2 \sin(\eta/2)} d\eta, \tag{3-60}$$

where

$$K(\mathbf{X}_c, \mathbf{X})(\theta, \eta) = \frac{1}{\pi R^4} (\partial_\theta \mathcal{X}(\theta)^\perp)^T \Delta_{\theta-\eta} \mathcal{X}(\theta) \Delta_{\theta-\eta} \mathcal{X}(\theta) \otimes \Delta_{\theta-\eta} \mathcal{X}(\theta) \times \left(1 - \frac{4}{R^2} \Delta_\eta \mathbf{X}_c(\theta)^T \Delta_\eta \mathbf{X}(\theta) + \mathcal{R}_3(\Delta_\eta \mathbf{X}(\theta)) \right),$$

and we recall the notation $\mathcal{X}(\theta) = \mathbf{X}_c(\theta) + \mathbf{X}(\theta)$ and (2-2).

We will plug in the splitting for \mathbf{F} in (2-15) into (1-13). We first introduce an analogous splitting for K as

$$K(\mathbf{X}_c, \mathbf{X})(\theta, \eta) = K_0(\mathbf{X}_c)(\theta, \eta) + K_L(\mathbf{X}_c, \mathbf{X})(\theta, \eta) + K_N(\mathbf{X}_c, \mathbf{X})(\theta, \eta). \tag{3-61}$$

After we remove the zero-order (2-18), and linear-order terms (2-23), then (1-13) for the nonlinear-order terms becomes the following equation for \mathbf{F}_N :

$$\mathbf{F}_N(\theta) - 2A_\mu \int_{\mathbb{S}} K_0(\mathbf{X}_c)(\theta, \eta) \frac{\mathbf{F}_N(\theta - \eta)}{2 \sin(\eta/2)} d\eta = \mathbf{J}(\mathbf{X}, \mathbf{F}_N)(\theta), \tag{3-62}$$

with

$$\begin{aligned} \mathbf{J}(\mathbf{X}, \mathbf{F}_N)(\theta) &= 2A_\mu \int_{\mathbb{S}} (K_L(\mathbf{X}_c, \mathbf{X})(\theta, \eta) + K_N(\mathbf{X}_c, \mathbf{X})(\theta, \eta)) \frac{\mathbf{F}_N(\theta - \eta)}{2 \sin(\eta/2)} d\eta \\ &\quad + 2A_\mu \int_{\mathbb{S}} (K_L(\mathbf{X}_c, \mathbf{X})(\theta, \eta) + K_N(\mathbf{X}_c, \mathbf{X})(\theta, \eta)) \frac{\mathbf{F}_L(\theta - \eta)}{2 \sin(\eta/2)} d\eta \\ &\quad + 2A_\mu \int_{\mathbb{S}} K_N(\mathbf{X}_c, \mathbf{X})(\theta, \eta) \mathbf{F}_0(\theta - \eta) \frac{d\eta}{2 \sin(\eta/2)}, \end{aligned} \tag{3-63}$$

where the first term in \mathbf{J} will be treated as a perturbation with \mathbf{F}_0 and \mathbf{F}_L given in (3-58) and (2-25) respectively. Notice that K_0 is given by

$$K_0(\mathbf{X}_c)(\theta, \eta) = \frac{1}{\pi R^4} (\partial_\theta \mathbf{X}_c(\theta)^\perp)^T \Delta_{\theta-\eta} \mathbf{X}_c(\theta) \Delta_{\theta-\eta} \mathbf{X}_c(\theta) \otimes \Delta_{\theta-\eta} \mathbf{X}_c(\theta),$$

where by (2-3) and (2-5) we have

$$\begin{aligned} (\partial_\theta \mathbf{X}_c(\theta)^\perp)^T \Delta_{\theta-\eta} \mathbf{X}_c(\theta) &= -R^2 \sin\left(\frac{\eta}{2}\right), \\ \Delta_{\theta-\eta} \mathbf{X}_c(\theta) \otimes \Delta_{\theta-\eta} \mathbf{X}_c(\theta) &= \frac{a^2}{2} \begin{bmatrix} 1 - \cos(2\theta - \eta) & -\sin(2\theta - \eta) \\ -\sin(2\theta - \eta) & 1 + \cos(2\theta - \eta) \end{bmatrix} \\ &\quad + \frac{b^2}{2} \begin{bmatrix} 1 + \cos(2\theta - \eta) & \sin(2\theta - \eta) \\ \sin(2\theta - \eta) & 1 - \cos(2\theta - \eta) \end{bmatrix} + ab \begin{bmatrix} \sin(2\theta - \eta) & -\cos(2\theta - \eta) \\ -\cos(2\theta - \eta) & -\sin(2\theta - \eta) \end{bmatrix}. \end{aligned}$$

Therefore,

$$\begin{aligned} \int_{\mathbb{S}} K_0(\mathbf{X}_c)(\theta, \eta) \frac{\mathbf{F}_N(\theta - \eta)}{2 \sin(\eta/2)} d\eta &= -\frac{1}{4\pi} \int_{\mathbb{S}} \mathbf{F}_N(\theta - \eta) d\eta \\ &\quad - \frac{a^2 - b^2}{4\pi R^2} \int_{\mathbb{S}} \begin{bmatrix} -\cos(2\theta - \eta) & -\sin(2\theta - \eta) \\ -\sin(2\theta - \eta) & \cos(2\theta - \eta) \end{bmatrix} \mathbf{F}_N(\theta - \eta) d\eta \\ &\quad - \frac{ab}{2\pi R^2} \int_{\mathbb{S}} \begin{bmatrix} \sin(2\theta - \eta) & -\cos(2\theta - \eta) \\ -\cos(2\theta - \eta) & -\sin(2\theta - \eta) \end{bmatrix} \mathbf{F}_N(\theta - \eta) d\eta. \end{aligned}$$

Then taking the Fourier transform we find that

$$\begin{aligned} &\mathcal{F}\left(\int_{\mathbb{S}} K_0(\mathbf{X}_c)(\theta, \eta) \frac{F_N(\theta - \eta)}{2 \sin(\eta/2)} d\eta\right)(k) \\ &= -\frac{1}{2} \widehat{F}_N(0) \delta_0(k) + \frac{(a+ib)^2}{4R^2} \begin{bmatrix} 1 & -i \\ -i & -1 \end{bmatrix} \widehat{F}_N(-1) \delta_1(k) + \frac{(a-ib)^2}{4R^2} \begin{bmatrix} 1 & i \\ i & -1 \end{bmatrix} \widehat{F}_N(1) \delta_{-1}(k). \end{aligned}$$

Equation (3-62) is then given on the Fourier side by the expressions

$$\begin{aligned} \widehat{F}_N(0) &= \frac{1}{1+A_\mu} \widehat{J(\mathbf{X}, F_N)}(0), \\ \widehat{F}_N(k) &= \widehat{J(\mathbf{X}, F_N)}(k), \quad k \geq 2, \end{aligned} \tag{3-64}$$

while for $k = 1$ one has that

$$\begin{aligned} \widehat{F}_N(1) - A_\mu \frac{(a+ib)^2}{2R^2} \begin{bmatrix} 1 & -i \\ -i & -1 \end{bmatrix} \widehat{F}_N(-1) &= \widehat{J(\mathbf{X}, F_N)}(1), \\ \widehat{F}_N(-1) - A_\mu \frac{(a-ib)^2}{2R^2} \begin{bmatrix} 1 & i \\ i & -1 \end{bmatrix} \widehat{F}_N(1) &= \widehat{J(\mathbf{X}, F_N)}(-1), \end{aligned}$$

which gives that

$$\begin{bmatrix} 1-A_\mu^2/2 & -iA_\mu^2/2 \\ iA_\mu^2/2 & 1-A_\mu^2/2 \end{bmatrix} \widehat{F}_N(1) = A_\mu \frac{(a+ib)^2}{2R^2} \begin{bmatrix} 1 & -i \\ -i & -1 \end{bmatrix} \widehat{J(\mathbf{X}, F_N)}(-1) + \widehat{J(\mathbf{X}, F_N)}(1),$$

and thus

$$\widehat{F}_N(1) = \frac{A_\mu}{1-A_\mu^2} \frac{(a+ib)^2}{2R^2} \begin{bmatrix} 1 & -i \\ -i & -1 \end{bmatrix} \widehat{J(\mathbf{X}, F_N)}(-1) + \frac{1}{1-A_\mu^2} \begin{bmatrix} 1-A_\mu^2/2 & iA_\mu^2/2 \\ -iA_\mu^2/2 & 1-A_\mu^2/2 \end{bmatrix} \widehat{J(\mathbf{X}, F_N)}(1).$$

Since we have

$$\left\| \begin{bmatrix} 1 & -i \\ -i & -1 \end{bmatrix} \right\| = 2, \quad \left\| \begin{bmatrix} 1-A_\mu^2/2 & iA_\mu^2/2 \\ -iA_\mu^2/2 & 1-A_\mu^2/2 \end{bmatrix} \right\| = 1,$$

we obtain

$$\begin{aligned} |\widehat{F}_N(1)| &\leq \frac{|A_\mu|}{1-A_\mu^2} |\widehat{J(\mathbf{X}, F_N)}(-1)| + \frac{1}{1-A_\mu^2} |\widehat{J(\mathbf{X}, F_N)}(1)| \\ &= \frac{1+|A_\mu|}{(1-A_\mu)(1+A_\mu)} |\widehat{J(\mathbf{X}, F_N)}(1)|, \end{aligned}$$

which together with (3-64) implies that

$$\|F_N\|_{\mathcal{F}_v^{0,1}} \leq \frac{1+|A_\mu|}{(1-A_\mu)(1+A_\mu)} \|J(\mathbf{X}, F_N)\|_{\mathcal{F}_v^{0,1}}. \tag{3-65}$$

This is our estimate for F_N .

$J(\mathbf{X}, F_N)$ estimate: Notice that $J(\mathbf{X}, F_N)$ corresponds to the nonlinear terms in $\mathcal{S}(F, \mathcal{X})$ except the one in the left-hand side of (3-62). For simplicity in notation, we are going to estimate $\mathcal{S}(F, \mathcal{X})$, and later extract from there the corresponding bounds for $J(\mathbf{X}, F_N)$.

Consider the following splitting for $\mathcal{S}(F, \mathcal{X})$ from (1-14):

$$\mathcal{S}(F, \mathcal{X})(\theta) = \mathcal{S}_1(F, \mathcal{X})(\theta) + \mathcal{S}_2(F, \mathcal{X})(\theta) + \mathcal{S}_3(F, \mathcal{X})(\theta), \tag{3-66}$$

with

$$\begin{aligned} \mathcal{S}_1(F, \mathcal{X})(\theta) &= \frac{1}{\pi R^4} \int_{\mathbb{S}} (\partial_\theta \mathcal{X}(\theta)^\perp)^T \Delta_{\theta-\eta} \mathcal{X}(\theta) \Delta_{\theta-\eta} \mathcal{X}(\theta) \otimes \Delta_{\theta-\eta} \mathcal{X}(\theta) \frac{F(\theta-\eta)}{2 \sin(\eta/2)} d\eta, \\ \mathcal{S}_2(F, \mathcal{X})(\theta) &= -\frac{4}{\pi R^6} \int_{\mathbb{S}} (\partial_\theta \mathcal{X}(\theta)^\perp)^T \Delta_{\theta-\eta} \mathcal{X}(\theta) \Delta_{\theta-\eta} \mathcal{X}(\theta) \\ &\quad \otimes \Delta_{\theta-\eta} \mathcal{X}(\theta) \Delta_\eta \mathbf{X}_c(\theta)^T \Delta_\eta \mathbf{X}(\theta) \frac{F(\theta-\eta)}{2 \sin(\eta/2)} d\eta, \\ \mathcal{S}_3(F, \mathcal{X})(\theta) &= \frac{1}{\pi R^4} \int_{\mathbb{S}} (\partial_\theta \mathcal{X}(\theta)^\perp)^T \Delta_{\theta-\eta} \mathcal{X}(\theta) \Delta_{\theta-\eta} \mathcal{X}(\theta) \otimes \Delta_{\theta-\eta} \mathcal{X}(\theta) \mathcal{R}_3(\Delta_\eta \mathbf{X}(\theta)) \frac{F(\theta-\eta)}{2 \sin(\eta/2)} d\eta. \end{aligned}$$

We take Fourier transform of $\mathcal{S}_1(F, \mathcal{X})$ to obtain

$$\widehat{\mathcal{S}_1(F, \mathcal{X})}(k) = \frac{1}{\pi R^4} \sum_{k_1 \in \mathbb{Z}} \cdots \sum_{k_4 \in \mathbb{Z}} (\widehat{\partial_\theta \mathcal{X}}(k - k_1)^\perp)^T \widehat{\partial_\theta \mathcal{X}}(k_1 - k_2) \widehat{\partial_\theta \mathcal{X}}(k_2 - k_3) \\ \otimes \widehat{\partial_\theta \mathcal{X}}(k_3 - k_4) \widehat{F}(k_4) I'_2(k_1, \dots, k_4),$$

where

$$\begin{aligned} |I'_2(k_1, \dots, k_4)| &= \left| \int_{\mathbb{S}} \prod_{j=1}^3 \frac{\sin((k_j - k_{j+1})\eta/2)}{(k_j - k_{j+1}) \sin(\eta/2)} \frac{e^{-i(k_1+k_4)\eta/2}}{2 \sin(\eta/2)} d\eta \right| \\ &= \left| \int_{\mathbb{S}} \frac{\sin((k_1 + k_4)\eta/2)}{\sin(\eta/2)} \prod_{j=1}^3 \frac{\sin((k_j - k_{j+1})\eta/2)}{(k_j - k_{j+1}) \sin(\eta/2)} d\eta \right|. \end{aligned}$$

The integral I'_2 turns out to be the previously defined integral in (3-4). In Lemma 3.2 we show that $|I'_2| \leq 2\pi$. Using (3-11) and (3-13), we have

$$\begin{aligned} &|(\widehat{\partial_\theta \mathcal{X}}(k - k_1)^\perp)^T \widehat{\partial_\theta \mathcal{X}}(k_1 - k_2)| \\ &\leq \frac{R^2}{2} (\delta_1(k - k_1) \delta_{-1}(k_1 - k_2) + \delta_{-1}(k - k_1) \delta_1(k_1 - k_2)) \\ &\quad + \frac{R}{\sqrt{2}} \delta_{1,-1}(k - k_1) |\widehat{\partial_\theta \mathcal{X}}(k_1 - k_2)| + \frac{R}{\sqrt{2}} \delta_{1,-1}(k_1 - k_2) |\widehat{\partial_\theta \mathcal{X}}(k - k_1)| + |\widehat{\partial_\theta \mathcal{X}}(k - k_1)| |\widehat{\partial_\theta \mathcal{X}}(k_1 - k_2)|, \end{aligned}$$

while recalling (3-18), (3-21) and (3-22), we obtain

$$\begin{aligned} &\|\widehat{\partial_\theta \mathcal{X}}(k_2 - k_3) \otimes \widehat{\partial_\theta \mathcal{X}}(k_3 - k_4)\| \\ &\leq \frac{R^2}{2} \delta_{1,-1}(k_2 - k_3) \delta_{1,-1}(k_3 - k_4) + \frac{\sqrt{2}}{2} R \delta_{1,-1}(k_2 - k_3) |\widehat{\partial_\theta \mathcal{X}}(k_3 - k_4)| \\ &\quad + \frac{\sqrt{2}}{2} R |\widehat{\partial_\theta \mathcal{X}}(k_2 - k_3)| \delta_{1,-1}(k_3 - k_4) + |\widehat{\partial_\theta \mathcal{X}}(k_2 - k_3)| |\widehat{\partial_\theta \mathcal{X}}(k_3 - k_4)|. \end{aligned}$$

Therefore, Young's inequality for convolutions yields that

$$\|\mathcal{S}_1(F, \mathcal{X})\|_{\mathcal{F}_v^{0,1}} \leq 2 \left(e^{2\nu_\infty} + 2\sqrt{2}e^{\nu_\infty} \frac{\|\mathbf{X}\|_{\dot{J}_v^{1,1}}}{R} + \frac{\|\mathbf{X}\|_{\dot{J}_v^{1,1}}^2}{R^2} \right) \left(2e^{2\nu_\infty} + 2\sqrt{2}e^{\nu_\infty} \frac{\|\mathbf{X}\|_{\dot{J}_v^{1,1}}}{R} + \frac{\|\mathbf{X}\|_{\dot{J}_v^{1,1}}^2}{R^2} \right) \|F\|_{\mathcal{F}_v^{0,1}},$$

which by (2-42) and notation (2-43) can be rewritten as

$$\|\mathcal{S}_1(\mathbf{F}, \mathcal{X})\|_{\mathcal{F}_v^{0,1}} \leq 2(2e^{4\nu_\infty} + 6\sqrt{2}e^{3\nu_\infty} C_1 \|\mathbf{X}\|_{\dot{\mathcal{F}}_v^{1,1}} + 11C_{12}C_1^2 \|\mathbf{X}\|_{\dot{\mathcal{F}}_v^{1,1}}^2) \|\mathbf{F}\|_{\mathcal{F}_v^{0,1}}, \quad (3-67)$$

with

$$C_{12} = \frac{1}{11}(11e^{2\nu_\infty} + 4\sqrt{2}e^{\nu_\infty} C_1 \|\mathbf{X}\|_{\dot{\mathcal{F}}_v^{1,1}} + C_1^2 \|\mathbf{X}\|_{\dot{\mathcal{F}}_v^{1,1}}^2), \quad (3-68)$$

and C_1 is defined in (2-43).

Following the same steps, for $\widehat{\mathcal{S}_2(\mathbf{F}, \mathcal{X})}$, one finds that

$$\|\mathcal{S}_2(\mathbf{F}, \mathcal{X})\|_{\mathcal{F}_v^{0,1}} \leq 8\sqrt{2}e^{\nu_\infty} (2e^{4\nu_\infty} + 6\sqrt{2}e^{3\nu_\infty} C_1 \|\mathbf{X}\|_{\dot{\mathcal{F}}_v^{1,1}} + 11C_{12}C_1^2 \|\mathbf{X}\|_{\dot{\mathcal{F}}_v^{1,1}}^2) C_1 \|\mathbf{X}\|_{\dot{\mathcal{F}}_v^{1,1}} \|\mathbf{F}\|_{\mathcal{F}_v^{0,1}}.$$

We define

$$C_{13} = \frac{1}{2}(2e^{4\nu_\infty} + 6\sqrt{2}e^{3\nu_\infty} C_1 \|\mathbf{X}\|_{\dot{\mathcal{F}}_v^{1,1}} + 11C_{12}C_1^2 \|\mathbf{X}\|_{\dot{\mathcal{F}}_v^{1,1}}^2), \quad (3-69)$$

so that

$$\|\mathcal{S}_2(\mathbf{F}, \mathcal{X})\|_{\mathcal{F}_v^{0,1}} \leq 16\sqrt{2}e^{\nu_\infty} C_{13} C_1 \|\mathbf{X}\|_{\dot{\mathcal{F}}_v^{1,1}} \|\mathbf{F}\|_{\mathcal{F}_v^{0,1}}. \quad (3-70)$$

This completes our $\widehat{\mathcal{S}_2(\mathbf{F}, \mathcal{X})}$ estimate.

Next, we proceed with $\widehat{\mathcal{S}_3(\mathbf{F}, \mathcal{X})}$ in (3-66). We split it accordingly to (3-59) as

$$\widehat{\mathcal{S}_3(\mathbf{F}, \mathcal{X})} = \widehat{\mathcal{S}_{3,1}(\mathbf{F}, \mathcal{X})} + \widehat{\mathcal{S}_{3,2}(\mathbf{F}, \mathcal{X})} + \widehat{\mathcal{S}_{3,3}(\mathbf{F}, \mathcal{X})} + \widehat{\mathcal{S}_{3,4}(\mathbf{F}, \mathcal{X})},$$

with

$$\begin{aligned} \mathcal{S}_{3,1}(\mathbf{F}, \mathcal{X}) &= -\frac{4}{\pi R^6} \int_{\mathbb{S}} (\partial_\theta \mathcal{X}(\theta)^\perp)^T \Delta_{\theta-\eta} \mathcal{X}(\theta) \Delta_{\theta-\eta} \mathcal{X}(\theta) \otimes \Delta_{\theta-\eta} \mathcal{X}(\theta) \\ &\quad \cdot \Delta_{\theta-\eta} \mathbf{X}_c(\theta)^T \Delta_{\theta-\eta} \mathbf{X}(\theta) \mathcal{R}_2(\Delta_\eta \mathbf{X}(\theta)) \frac{\mathbf{F}(\theta-\eta)}{2 \sin(\eta/2)} d\eta, \end{aligned}$$

$$\mathcal{S}_{3,2}(\mathbf{F}, \mathcal{X}) = \frac{2}{\pi R^4} \int_{\mathbb{S}} (\partial_\theta \mathcal{X}(\theta)^\perp)^T \Delta_{\theta-\eta} \mathcal{X}(\theta) \Delta_{\theta-\eta} \mathcal{X}(\theta) \otimes \Delta_{\theta-\eta} \mathcal{X}(\theta) \mathcal{R}_2(\Delta_\eta \mathbf{X}(\theta)) \frac{\mathbf{F}(\theta-\eta)}{2 \sin(\eta/2)} d\eta,$$

$$\begin{aligned} \mathcal{S}_{3,3}(\mathbf{F}, \mathcal{X}) &= \frac{4}{\pi R^8} \int_{\mathbb{S}} (\partial_\theta \mathcal{X}(\theta)^\perp)^T \Delta_{\theta-\eta} \mathcal{X}(\theta) \Delta_{\theta-\eta} \mathcal{X}(\theta) \otimes \Delta_{\theta-\eta} \mathcal{X}(\theta) \\ &\quad \cdot (\Delta_{\theta-\eta} \mathbf{X}_c(\theta)^T \Delta_{\theta-\eta} \mathbf{X}(\theta))^2 \frac{\mathbf{F}(\theta-\eta)}{2 \sin(\eta/2)} d\eta, \end{aligned}$$

$$\mathcal{S}_{3,4}(\mathbf{F}, \mathcal{X}) = \frac{1}{\pi R^4} \int_{\mathbb{S}} (\partial_\theta \mathcal{X}(\theta)^\perp)^T \Delta_{\theta-\eta} \mathcal{X}(\theta) \Delta_{\theta-\eta} \mathcal{X}(\theta) \otimes \Delta_{\theta-\eta} \mathcal{X}(\theta) (\mathcal{R}_2(\Delta_\eta \mathbf{X}(\theta)))^2 \frac{\mathbf{F}(\theta-\eta)}{2 \sin(\eta/2)} d\eta.$$

The procedure follows the steps used to bound $\mathcal{N}_{3,4}(\theta)$ in (3-24) and (3-41), where the term \mathcal{R}_2 from (2-9) was also involved. After taking Fourier transform and using Lemma 3.2, Young's inequality for convolutions and summation in m and n gives

$$\|\mathcal{S}_{3,1}(\mathbf{F}, \mathcal{X})\|_{\mathcal{F}_v^{0,1}} \leq 144\sqrt{2}e^{\nu_\infty} C_6 C_{13} C_1^3 \|\mathbf{X}\|_{\dot{\mathcal{F}}_v^{1,1}}^3 \|\mathbf{F}\|_{\mathcal{F}_v^{0,1}},$$

$$\|\mathcal{S}_{3,2}(\mathbf{F}, \mathcal{X})\|_{\mathcal{F}_v^{0,1}} \leq 72C_6 C_{13} C_1^2 \|\mathbf{X}\|_{\dot{\mathcal{F}}_v^{1,1}}^2 \|\mathbf{F}\|_{\mathcal{F}_v^{0,1}},$$

$$\|\mathcal{S}_{3,3}(\mathbf{F}, \mathcal{X})\|_{\mathcal{F}_v^{0,1}} \leq 32e^{2\nu_\infty} C_{13} C_1^2 \|\mathbf{X}\|_{\dot{\mathcal{F}}_v^{1,1}}^2 \|\mathbf{F}\|_{\mathcal{F}_v^{0,1}},$$

$$\|\mathcal{S}_{3,4}(\mathbf{F}, \mathcal{X})\|_{\mathcal{F}_v^{0,1}} \leq 324C_{13} C_6^2 C_1^4 \|\mathbf{X}\|_{\dot{\mathcal{F}}_v^{1,1}}^4 \|\mathbf{F}\|_{\mathcal{F}_v^{0,1}}.$$

Joining the above bounds, we obtain

$$\|\mathcal{S}_3(\mathbf{F}, \mathcal{X})\|_{\mathcal{F}_v^{0,1}} \leq 104C_{13}C_{14}C_1^2\|\mathbf{X}\|_{\dot{\mathcal{J}}_v^{1,1}}^2\|\mathbf{F}\|_{\mathcal{F}_v^{0,1}}, \quad (3-71)$$

with

$$C_{14} = \frac{1}{104}(72C_6 + 32e^{2\nu_\infty} + 144\sqrt{2}e^{\nu_\infty}C_6C_1\|\mathbf{X}\|_{\dot{\mathcal{J}}_v^{1,1}} + 324C_6^2C_1^2\|\mathbf{X}\|_{\dot{\mathcal{J}}_v^{1,1}}^2), \quad (3-72)$$

where C_1 and C_6 previously defined in (2-43) and (3-42), respectively. We combine the bounds (3-67), (3-70), and (3-71), and order them as

$$\|\mathcal{S}(\mathbf{F}, \mathcal{X})\|_{\mathcal{F}_v^{0,1}} \leq 4e^{4\nu_\infty}\|\mathbf{F}\|_{\mathcal{F}_v^{0,1}} + 28\sqrt{2}e^{5\nu_\infty}C_1\|\mathbf{X}\|_{\dot{\mathcal{J}}_v^{1,1}}\|\mathbf{F}\|_{\mathcal{F}_v^{0,1}} + 222C_{15}C_1^2\|\mathbf{X}\|_{\dot{\mathcal{J}}_v^{1,1}}^2\|\mathbf{F}\|_{\mathcal{F}_v^{0,1}},$$

with

$$C_{15} = \frac{1}{222}(104C_{13}C_{14} + 8\sqrt{2}e^{\nu_\infty}(6\sqrt{2}e^{3\nu_\infty} + 11C_{12}C_1\|\mathbf{X}\|_{\dot{\mathcal{J}}_v^{1,1}}) + 22C_{12}), \quad (3-73)$$

and C_1 in (2-43), C_{12} in (3-68), C_{13} in (3-69) and C_{14} in (3-72).

We remark that instead of (3-66), analogously to the splitting for K in (3-61) we can split $\mathcal{S}(\mathbf{F}, \mathcal{X})(\theta)$ as

$$\mathcal{S}(\mathbf{F}, \mathcal{X})(\theta) = \mathcal{S}_0(\mathbf{F}, \mathcal{X})(\theta) + \mathcal{S}_L(\mathbf{F}, \mathcal{X})(\theta) + \mathcal{S}_N(\mathbf{F}, \mathcal{X})(\theta), \quad (3-74)$$

where from (3-60) and (3-61) we have

$$\mathcal{S}_0(\mathbf{F}, \mathcal{X})(\theta) = \int_{\mathbb{S}} K_0(\mathbf{X}_c, \mathbf{X})(\theta, \eta) \frac{\mathbf{F}(\theta - \eta)}{2 \sin(\eta/2)} d\eta.$$

Then $\mathcal{S}_L(\mathbf{F}, \mathcal{X})(\theta)$ analogously contains $K_L(\mathbf{X}_c, \mathbf{X})(\theta, \eta)$ from (3-61) and \mathcal{S}_L is linear in \mathbf{X} . Then $\mathcal{S}_N(\mathbf{F}, \mathcal{X})(\theta)$ similarly contains $K_N(\mathbf{X}_c, \mathbf{X})(\theta, \eta)$ from (3-61) and \mathcal{S}_N is nonlinear in \mathbf{X} . Then it is clear from the above that we have the estimates

$$\begin{aligned} \|\mathcal{S}_0(\mathbf{F}, \mathcal{X})\|_{\mathcal{F}_v^{0,1}} &\leq 4e^{4\nu_\infty}\|\mathbf{F}\|_{\mathcal{F}_v^{0,1}}, \\ \|\mathcal{S}_L(\mathbf{F}, \mathcal{X})\|_{\mathcal{F}_v^{0,1}} &\leq 28\sqrt{2}e^{5\nu_\infty}C_1\|\mathbf{X}\|_{\dot{\mathcal{J}}_v^{1,1}}\|\mathbf{F}\|_{\mathcal{F}_v^{0,1}}, \\ \|\mathcal{S}_N(\mathbf{F}, \mathcal{X})\|_{\mathcal{F}_v^{0,1}} &\leq 222C_{15}C_1^2\|\mathbf{X}\|_{\dot{\mathcal{J}}_v^{1,1}}^2\|\mathbf{F}\|_{\mathcal{F}_v^{0,1}}. \end{aligned}$$

This splits the estimates into zero-order, linear-order, and nonlinear-order which is useful because of (3-63).

Next, recalling the definition of $\mathbf{J}(\mathbf{X}, \mathbf{F}_N)$ from (3-63) and its relation with $\mathcal{S}(\mathbf{F}, \mathcal{X})$ in (3-60) and (3-74), it follows that

$$\begin{aligned} \|\mathbf{J}(\mathbf{X}, \mathbf{F}_N)\|_{\mathcal{F}_v^{0,1}} &\leq 56\sqrt{2}|A_\mu|C_{16}C_1\|\mathbf{X}\|_{\dot{\mathcal{J}}_v^{1,1}}\|\mathbf{F}_N\|_{\mathcal{F}_v^{0,1}} \\ &\quad + 56\sqrt{2}|A_\mu|C_{16}C_1\|\mathbf{X}\|_{\dot{\mathcal{J}}_v^{1,1}}\|\mathbf{F}_L\|_{\mathcal{F}_v^{0,1}} + 444|A_\mu|C_{15}C_1^2\|\mathbf{X}\|_{\dot{\mathcal{J}}_v^{1,1}}^2\|\mathbf{F}_0\|_{\mathcal{F}_v^{0,1}}, \end{aligned} \quad (3-75)$$

with

$$C_{16} = \frac{1}{28\sqrt{2}}(28\sqrt{2}e^{5\nu_\infty} + 222C_{15}C_1\|\mathbf{X}\|_{\dot{\mathcal{J}}_v^{1,1}}). \quad (3-76)$$

Finally, bound (3-75) allows us to estimate \mathbf{F}_N from (3-65),

$$\begin{aligned} &\|\mathbf{F}_N\|_{\mathcal{F}_v^{0,1}} \\ &\leq \frac{56\sqrt{2}|A_\mu|(1 + |A_\mu|)}{(1 - A_\mu)(1 + A_\mu)}C_{17}C_{16}C_1\|\mathbf{X}\|_{\dot{\mathcal{J}}_v^{1,1}}\|\mathbf{F}_L\|_{\mathcal{F}_v^{0,1}} + \frac{444|A_\mu|(1 + |A_\mu|)}{(1 - A_\mu)(1 + A_\mu)}C_{17}C_{15}C_1^2\|\mathbf{X}\|_{\dot{\mathcal{J}}_v^{1,1}}^2\|\mathbf{F}_0\|_{\mathcal{F}_v^{0,1}}, \end{aligned}$$

where

$$C_{17} = \left(1 - \frac{56\sqrt{2}|A_\mu|(1 + |A_\mu|)}{(1 - A_\mu)(1 + A_\mu)} C_{16} C_1 \|X\|_{\dot{X}_v^{1,1}} \right)^{-1}, \tag{3-77}$$

and the bounds for F_0, F_L are given in (3-55) and (3-56). Substituting these bounds we find

$$\begin{aligned} \|F_N\|_{\mathcal{F}_v^{0,1}} &\leq 112\sqrt{2}A_e \frac{|A_\mu|(1 + |A_\mu|)(1 - A_\mu + |A_\mu|)}{(1 - A_\mu)^2(1 + A_\mu)} C_{17} C_{16} C_1 \|X\|_{\dot{X}_v^{1,1}} \|X\|_{\dot{X}_v^{2,1}} \\ &\quad + 888\sqrt{2}A_e \frac{|A_\mu|(1 + |A_\mu|)}{(1 - A_\mu)^2(1 + A_\mu)} e^{v\infty} C_8 C_{17} C_{15} C_1^2 \|X\|_{\dot{X}_v^{1,1}} \|X\|_{\dot{X}_v^{2,1}}. \end{aligned}$$

Defining

$$D_4 = \frac{1}{1000} C_1 C_{17} (112(1 - A_\mu + |A_\mu|) C_{16} + 888e^{v\infty} C_8 C_1 C_{15}), \tag{3-78}$$

where C_1, C_8, C_{15}, C_{16} , and C_{17} are defined in (2-43), (3-46), (3-73), (3-76), and (3-77), we can write the estimate for F_N as (3-57). □

Proof of Lemma 3.2. Recalling (3-3) and (3-2) and using the odd part of the integral we can rewrite $I = I_n$ as

$$\begin{aligned} I &= -i \operatorname{pv} \int_{-\pi}^{\pi} \frac{\sin((k_1 + k_{2n})\eta/2) - \frac{\sin((k - k_1)\eta/2)}{(k - k_1) \tan(\eta/2)} \sin((k + k_{2n})\eta/2)}{2 \sin(\eta/2)} \prod_{j=1}^{2n-1} \frac{\sin((k_j - k_{j+1})\eta/2)}{(k_j - k_{j+1}) \sin(\eta/2)} d\eta \\ &= -\frac{i}{2} (I' - I''), \end{aligned}$$

where

$$I' \stackrel{\text{def}}{=} \operatorname{pv} \int_{-\pi}^{\pi} \frac{\sin((k_1 + k_{2n})\eta/2)}{\sin(\eta/2)} \prod_{j=1}^{2n-1} \frac{\sin((k_j - k_{j+1})\eta/2)}{(k_j - k_{j+1}) \sin(\eta/2)} d\eta, \tag{3-79}$$

$$I'' \stackrel{\text{def}}{=} \operatorname{pv} \int_{-\pi}^{\pi} \cos(\eta/2) \frac{\sin((k + k_{2n})\eta/2)}{\sin(\eta/2)} \prod_{j=0}^{2n-1} \frac{\sin((k_j - k_{j+1})\eta/2)}{(k_j - k_{j+1}) \sin(\eta/2)} d\eta. \tag{3-80}$$

Note that if $k_1 + k_{2n} = 0$ then $I' = 0$ and if $k + k_{2n} = 0$ then $I'' = 0$. We henceforth assume that $|k_1 + k_{2n}| \geq 1$ and $|k + k_{2n}| \geq 1$. We will calculate (3-79) and then (3-80).

Notice that $\sin((k_j - k_{j+1})\eta/2) = \operatorname{sgn}(k_j - k_{j+1}) \sin(|k_j - k_{j+1}|\eta/2)$ and, since $|k_j - k_{j+1}| \geq 1$, we rewrite the quotient in the product form as

$$\begin{aligned} \frac{\sin(|k_j - k_{j+1}|\eta/2)}{\sin(\eta/2)} &= \frac{e^{i|k_j - k_{j+1}|\eta/2} - e^{-i|k_j - k_{j+1}|\eta/2}}{e^{i\eta/2} - e^{-i\eta/2}} = \frac{e^{i|k_j - k_{j+1}|\eta/2} (1 - e^{-i|k_j - k_{j+1}|\eta})}{e^{i\eta/2} (1 - e^{-i\eta})} \\ &= e^{i(|k_j - k_{j+1}| - 1)\eta/2} \sum_{m=0}^{|k_j - k_{j+1}| - 1} e^{-i\eta m} = \sum_{m=0}^{|k_j - k_{j+1}| - 1} e^{i(-2m + |k_j - k_{j+1}| - 1)\eta/2}. \end{aligned}$$

We conclude that

$$\begin{aligned} \prod_{j=1}^{2n-1} \frac{\sin(|k_j - k_{j+1}|\eta/2)}{\sin(\eta/2)} &= \prod_{j=1}^{2n-1} \sum_{m_j=0}^{|k_j - k_{j+1}|-1} e^{i(-2m_j + |k_j - k_{j+1}|-1)\eta/2} \\ &= \sum_{m_1=0}^{|k_1 - k_2|-1} \cdots \sum_{m_{2n-1}=0}^{|k_{2n-1} - k_{2n}|-1} e^{i(|k_1 - k_2| + \cdots + |k_{2n-1} - k_{2n}| - 2(m_1 + \cdots + m_{2n-1}) - 2n)\eta/2}. \end{aligned}$$

In particular following those calculations we can express the integrand of I' in (3-79) as

$$\begin{aligned} \frac{\sin((k_1 + k_{2n})\eta/2)}{\sin(\eta/2)} \prod_{j=1}^{2n-1} \frac{\sin((k_j - k_{j+1})\eta/2)}{(k_j - k_{j+1}) \sin(\eta/2)} \\ = \left(\operatorname{sgn}(k_1 + k_{2n}) \prod_{j=1}^{2n-1} \frac{1}{|k_j - k_{j+1}|} \right) \sum_{\substack{m_j=0 \\ 1 \leq j \leq 2n-1}}^{|k_j - k_{j+1}|-1} \sum_{m_{2n}=0}^{|k_1 + k_{2n}|-1} e^{iB_1\eta/2}, \end{aligned}$$

where to be clear in the sum the m_j indicates a further summation over all $j \in \{1, \dots, 2n - 1\}$. Also we define B_1 above as

$$B_1 = \sum_{j=1}^{2n-1} |k_j - k_{j+1}| + |k_1 + k_{2n}| - 2 \sum_{j=1}^{2n} m_j - 2n.$$

Notice that no matter what the sign of any of the terms inside the absolute values above is we always have $\sum_{j=1}^{2n-1} |k_j - k_{j+1}| + |k_1 + k_{2n}| = 2l$ for some integer l so that B_1 is an even integer. This holds because the sum contains two copies of every k_1, \dots, k_{2n} .

We further integrate as $\int_{-\pi}^{\pi} e^{iB_1\eta/2} d\eta = (4/B_1) \sin(B_1\pi/2)$, and we notice that since B_1 is an even integer, either $(4/B_1) \sin(B_1\pi/2) = 0$ if $B_1 \neq 0$ or $(4/B_1) \sin(B_1\pi/2) = 2\pi$ when $B_1 = 0$. We can then find the following expression for I' :

$$I' = 2\pi \left(\operatorname{sgn}(k_1 + k_{2n}) \prod_{j=1}^{2n-1} \frac{1}{|k_j - k_{j+1}|} \right) \sum_{\substack{m_j=0 \\ 1 \leq j \leq 2n-1}}^{|k_j - k_{j+1}|-1} \sum_{m_{2n}=0}^{|k_1 + k_{2n}|-1} 1_{B_1=0}. \tag{3-81}$$

This is our calculation of the integral (3-79). We note, as a function of the single variable m_{2n} , that B_1 is decreasing and takes the value zero at most one time. Thus $\sum_{m_{2n}=0}^{|k_1 + k_{2n}|-1} 1_{B_1=0} \leq 1$. We conclude that $|I'| \leq 2\pi$. Note that (3-4) is exactly (3-79). So this proves the second estimate for (3-4) in Lemma 3.2.

We now calculate the integral (3-80), which is rather similar. We obtain

$$\prod_{j=0}^{2n-1} \frac{\sin(|k_j - k_{j+1}|\eta/2)}{\sin(\eta/2)} = \sum_{\substack{m_j=0 \\ 0 \leq j \leq 2n-1}}^{|k_j - k_{j+1}|-1} e^{i(|k_0 - k_1| + \cdots + |k_{2n-1} - k_{2n}| - 2(m_0 + \cdots + m_{2n-1}) - 2n)\eta/2}.$$

Then we can express the integrand of I'' as

$$\begin{aligned} \cos(\eta/2) \frac{\sin((k + k_{2n})\eta/2)}{\sin(\eta/2)} \prod_{j=0}^{2n-1} \frac{\sin((k_j - k_{j+1})\eta/2)}{(k_j - k_{j+1}) \sin(\eta/2)} \\ = \frac{1}{2} \left(\operatorname{sgn}(k + k_{2n}) \prod_{j=0}^{2n-1} \frac{1}{|k_j - k_{j+1}|} \right) \sum_{\substack{m_j=0 \\ 0 \leq j \leq 2n-1}}^{|k_j - k_{j+1}|-1} \sum_{m_{2n}=0}^{|k + k_{2n}|-1} (e^{iB_2\eta/2} + e^{iB_3\eta/2}). \end{aligned}$$

We define

$$B_2 = \sum_{j=0}^{2n-1} |k_j - k_{j+1}| + |k + k_{2n}| - 2 \sum_{j=0}^{2n} m_j - 2n,$$

$$B_3 = \sum_{j=0}^{2n-1} |k_j - k_{j+1}| + |k + k_{2n}| - 2 \sum_{j=0}^{2n} m_j - 2n - 2.$$

Similarly since $\sum_{j=0}^{2n} |k_j - k_{j+1}| + |k + k_{2n}|$ contains two copies of every k_0, k_1, \dots, k_{2n} then it is always an even integer. Therefore we conclude that B_2 and B_3 both are even integers.

We can then similarly find the following expression for I'' :

$$I'' = \pi \left(\operatorname{sgn}(k + k_{2n}) \prod_{j=0}^{2n-1} \frac{1}{|k_j - k_{j+1}|} \right) \sum_{\substack{m_j=0 \\ 0 \leq j \leq 2n-1}}^{|k_j - k_{j+1}| - 1} \sum_{m_{2n}=0}^{|k + k_{2n}| - 1} (1_{B_2=0} + 1_{B_3=0}).$$

Then using the same argument as our upper bound estimate for I' we obtain that $|I''| \leq 2\pi$. □

Remark 3.4. One can generally calculate the sum in (3-81) exactly. In particular the value of the sums in (3-81) can be seen as the number of nonnegative integer solutions to the equation

$$m_1 + \dots + m_{2n} = \frac{1}{2} \sum_{j=1}^{2n-1} |k_j - k_{j+1}| + |k_1 + k_{2n}| - n,$$

with the restrictions that $0 \leq m_j \leq |k_j - k_{j+1}| - 1$ for $j = 1, \dots, 2n - 1$ and $0 \leq m_{2n} \leq |k_1 + k_{2n}| - 1$. This value can be calculated exactly using the inclusion-exclusion formula.

Alternatively, if $n = 1$ in (3-4) then one can calculate, on the region where $I'_1 \neq 0$, that we have exactly

$$I'_1 = 2\pi \frac{\min\{|k_1 - k_2|, |k_1 + k_2|\} \operatorname{sgn}(k_1 + k_2)}{|k_1 - k_2|}.$$

And this formula is consistent with our estimate in Lemma 3.2.

4. Proof of main theorem

This section is devoted to the proof of Theorem 1.2. In Section 4A we show the scheme of the proof for existence of solutions via a regularization argument. The main part consists in obtaining the a priori estimates, in particular the energy inequality from (1-30). Uniqueness is later proved in Section 4B.

4A. Existence. The proof follows a standard regularization argument. We will use a regularization of (1-10) and (1-13), written in the form of (2-48), and the a priori estimates of the previous section to find a weak solution in the sense of Definition 4.2 below. The regularity obtained for the solution will imply that the solution found is indeed a strong solution, which we prove later is unique.

Definition 4.1. For fixed $t \in [0, T]$ and $\phi(t) \in W^{2,\infty}(\mathbb{S})$, we say that $\psi(t) \in L^\infty(\mathbb{S})$ is a weak solution of

$$\psi(\theta, t) + 2A_\mu \partial_\theta \phi(\theta, t)^\perp \cdot \int_{\mathbb{S}} \mathcal{T}(\phi(\theta, t) - \phi(\eta, t)) \cdot \psi(\eta, t) d\eta = 2A_e \partial_\theta^2 \phi(\theta, t),$$

with \mathcal{T} given by (1-11), if for any $\varphi \in \mathcal{D}(\mathbb{S})$ it holds that

$$\int_{\mathbb{S}} \psi(\theta, t) \cdot \varphi(\theta) d\theta + 2A_\mu \int_{\mathbb{S}} \varphi_i(\theta) \int_{\mathbb{S}} \partial_\theta \phi_j(\theta, t)^\perp \mathcal{T}_{ijk}(\phi(\theta, t) - \phi(\eta, t)) \psi_k(\eta, t) d\eta d\theta = 2A_e \int_{\mathbb{S}} \partial_\theta^2 \phi(\theta, t) \cdot \varphi(\theta).$$

Definition 4.2. We say that $\mathcal{X} \in L^\infty([0, T]; W^{1,\infty}(\mathbb{S})) \cap L^1([0, T]; W^{2,\infty}(\mathbb{S}))$ is a weak solution of (1-10) if for almost every $t \in [0, T]$ the arc-chord condition (1-17) is satisfied, and if for any $\varphi \in \mathcal{D}(\mathbb{S} \times [0, T])$ it holds that

$$\int_{\mathbb{S}} \mathcal{X}(\theta, t) \cdot \varphi(\theta, t) d\theta - \int_{\mathbb{S}} \mathcal{X}_0(\theta) \cdot \varphi(\theta, 0) d\theta - \int_0^t \int_{\mathbb{S}} \mathcal{X}(\theta, \tau) \cdot \partial_t \varphi(\theta, \tau) d\theta d\tau = \int_0^t \int_{\mathbb{S}} \varphi(\theta, \tau) \cdot \int_{\mathbb{S}} G(\mathcal{X}(\theta, \tau) - \mathcal{X}(\eta, \tau)) \mathbf{F}(\eta, \tau) d\eta d\theta d\tau,$$

where G is defined in (1-9) and $\mathbf{F} \in L^1([0, T]; L^\infty(\mathbb{S}))$ is the solution in the sense of Definition 4.1 of (1-13).

We will write $f_M = \mathcal{J}_M f$ for general f such as $f = X$, $f = X_c$, $f = Y$ or $f = F$, with \mathcal{J}_M the high-frequency cut-off defined in (1-23). We start by considering a regularized version of system (1-10), (1-13) (where (1-10) is written in (2-28) with the linear and nonlinear terms apart). For each positive integer M , consider the regularized initial data $\mathcal{X}_{0,M}$ and the corresponding solution $\mathcal{X} = X_M + X_{M,c}$ to the regularized system

$$\begin{aligned} \partial_t \mathcal{X}_M &= -\frac{A_e}{2} (\Lambda X_M + \mathcal{H}R^{-1} X_M) + \mathcal{J}_M \mathcal{N}(X_{M,c}, X_M), \\ F_M &= 2A_\mu \mathcal{J}_M \mathcal{S}(F_M, \mathcal{X}_M) + 2A_e \partial_\theta^2 \mathcal{X}_M. \end{aligned} \tag{4-1}$$

We define correspondingly $\mathcal{Y}_{0,M}$ and $\mathcal{Y}_M = Y_M + Y_{M,c}$. We recall that (2-28) could be written in \mathcal{Y} -variables as (2-48). The corresponding regularized system in these variables reads as follows:

$$\begin{aligned} \widehat{Y}_M(0) &= 0, \quad \widehat{Y}_{M,2}(1) = 0, \quad \widehat{Y}_{M,c}(k) = 0, \quad k \neq 0, 1, \quad \widehat{Y}_{M,c,1}(1) = 0, \\ \partial_t \widehat{Y}_{M,c}(0) &= P(0)^{-1} \widehat{\mathcal{N}}(X_{M,c}, X_M)(0), \\ \partial_t \widehat{Y}_{M,1}(1) &= -A_e \widehat{Y}_{M,1}(1) + (P(1)^{-1} \widehat{\mathcal{N}}(X_{M,c}, X_M)(1))_1, \\ \partial_t \widehat{Y}_M(k) &= -\frac{A_e}{2} \mathcal{D}(k) \widehat{Y}_M(k) + P(k)^{-1} \widehat{\mathcal{N}}(X_{M,c}, X_M)(k), \quad 2 \leq k \leq M, \\ |\widehat{Y}_{M,c,2}(1)|^2 &= \frac{1}{2} - \sum_{1 \leq k \leq M} k (|\widehat{Y}_{M,2}(k)|^2 - |\widehat{Y}_{M,1}(k)|^2), \end{aligned} \tag{4-2}$$

with F_M given by (4-1). Since $X_{M,c}$ is a circle with radius satisfying (2-42), the chord arc condition (1-17) is clearly satisfied for $\|X_M\|_{\dot{F}_v^{1,1}}$ sufficiently small; we shall soon see that this in fact holds so long as $\|X_M\|_{\dot{F}_v^{1,1}} < k(A_\mu)$, which is defined in (4-9). Then, with the same size condition, F_M is estimated in terms of X_M as in Section 3B. Thus, with F_M solved in terms of Y_M using the transformation (2-31), we obtain an ODE of the form

$$\dot{Y}_M = \mathcal{J}_M \mathcal{G}(Y_M), \quad Y_M(0) = Y_{M,0},$$

for a certain nonlinear function \mathcal{G} . Notice that the ODE for $\widehat{Y}_{M,c}(0)$ is decoupled from the rest because there are no zero modes in $\widehat{\mathcal{N}}(\widehat{X}_{M,c}, \widehat{X}_M)(0)$. Therefore, Picard’s theorem on Banach spaces yields the local existence of regularized solutions $Y_M \in C^1([0, T_M]; H_M^m)$, where $H_M^m = \{f \in H^m(\mathbb{S}) : \text{supp}(\widehat{f}) \subset [-M, M]\}$. Since the a priori energy estimate (4-11) holds for the regularized system, we have uniform bounds for Y_M in the space $L^\infty(\mathbb{R}_+; \dot{F}_\nu^{1,1}) \cap L^1(\mathbb{R}_+; \dot{F}_\nu^{2,1})$. It is not hard to prove that Y_M forms a Cauchy sequence in $L^\infty([0, T]; \mathcal{F}_\nu^{0,1})$, so that we have a candidate for solution. One can then apply a version of the Aubin–Lions lemma (see Corollary 6 of [Simon 1987]) to get the strong convergence, up to a subsequence, of the approximate problems in $L^2([0, T]; \dot{F}_\nu^{1,1})$. Next, since $\widehat{Y}_M(m, t) \rightarrow \widehat{Y}(m, t)$ as $M \rightarrow \infty$, for all $m \in \mathbb{Z}$ and almost every t , Fatou’s lemma allows us to conclude that

$$\begin{aligned} M(t) &= \|Y\|_{\dot{F}_\nu^{1,1}}(t) + \frac{A_e}{4} \mathcal{C} \int_0^t \|Y\|_{\dot{F}_\nu^{2,1}}(\tau) d\tau \\ &\leq \liminf_{M \rightarrow +\infty} \left(\|Y_M\|_{\dot{F}_\nu^{1,1}}(t) + \frac{A_e}{4} \mathcal{C} \int_0^t \|Y_M\|_{\dot{F}_\nu^{2,1}}(\tau) d\tau \right) \leq \|Y_0\|_{\dot{F}_\nu^{1,1}}, \end{aligned}$$

so we obtain that the limit function Y belongs to $L^\infty([0, T]; \dot{F}_\nu^{1,1}) \cap L^1([0, T]; \dot{F}_\nu^{2,1})$. Now, we claim that the strong convergence, up to a subsequence, of $X_M \rightarrow X$ in $L^\infty([0, T]; \dot{F}_\nu^{1,1}) \cap L^1([0, T]; \dot{F}_\nu^{2,1})$ holds. The proof of this claim follows in fact along the lines of the proof of uniqueness (see Section 4B). This strong convergence immediately implies from (4-1), under the size constraint (1-29), the strong convergence $F_M \rightarrow F$ in $L^1([0, T]; \mathcal{F}_\nu^{0,1})$. In fact, it suffices to consider F_{M_1} and F_{M_2} , write their difference as

$$\begin{aligned} F_{M_1} - F_{M_2} &= 2A_\mu (\mathcal{J}_{M_1} \mathcal{S}(F_{M_1}, \mathcal{X}_{M_1}) - \mathcal{J}_{M_2} \mathcal{S}(F_{M_1}, \mathcal{X}_{M_1})) \\ &\quad + 2A_\mu (\mathcal{J}_{M_2} \mathcal{S}(F_{M_1}, \mathcal{X}_{M_1}) - \mathcal{J}_{M_2} \mathcal{S}(F_{M_2}, \mathcal{X}_{M_1})) \\ &\quad + 2A_\mu (\mathcal{J}_{M_2} \mathcal{S}(F_{M_2}, \mathcal{X}_{M_1}) - \mathcal{J}_{M_2} \mathcal{S}(F_{M_2}, \mathcal{X}_{M_2})) \\ &\quad + 2A_e (\partial_\theta^2 \mathcal{X}_{M_1} - \partial_\theta^2 \mathcal{X}_{M_2}), \end{aligned}$$

and perform estimates similar to the ones in Section 3B to find that F_M forms a Cauchy sequence in $L^1([0, T]; \mathcal{F}_\nu^{0,1})$. Since $X_{M,c}$ is given in terms of X_M , the above convergence holds for \mathcal{X}_M . The strong convergence $\mathcal{X}_M \rightarrow \mathcal{X}$ in $L^\infty([0, T]; \dot{F}_\nu^{1,1})$ together with $F_M \rightarrow F$ in $L^1([0, T]; \mathcal{F}_\nu^{0,1})$ yields \mathcal{X} as a solution to (1-10) in the sense of Definition 4.2. (Moreover, it is easy to check in (4-1) the strong convergence of the right-side terms in $L^1([0, T]; \dot{F}_\nu^{1,1})$.)

We refer to Section 5 of [Gancedo et al. 2020] for a similar approximation argument, including the instant generation of analyticity and the continuity in time. We include it here for completeness. From the strong convergence in $L^1([0, T]; \dot{F}_\nu^{1,1})$ of the right-hand side of (1-10), we must have that $\partial_t \mathcal{X}_M \rightarrow \partial_t \mathcal{X}$ in $L^1([0, T]; \dot{F}_\nu^{1,1})$. Consider $0 < t \leq t_1 < t_2$. Then,

$$\|\mathcal{X}(t_2) - \mathcal{X}(t_1)\|_{\dot{F}_\nu^{1,1}(t)} = \left\| \int_{t_1}^{t_2} \partial_t \mathcal{X}(\tau) d\tau \right\|_{\dot{F}_\nu^{1,1}(t)} \leq \int_{t_1}^{t_2} \|\partial_t \mathcal{X}(\tau)\|_{\dot{F}_\nu^{1,1}(\tau)} d\tau,$$

which from the fact that $\partial_t \mathcal{X} \in L^1([0, T]; \dot{F}_\nu^{1,1})$ yields that the solution is analytic for all positive times, and $\mathcal{X} \in C([\varepsilon, T]; \dot{F}_\nu^{1,1})$ for any $\varepsilon > 0$. Moreover, fix $\tilde{\nu}_m \in (0, \nu_\infty)$ and define $\tilde{\nu}(t)$ according to (1-27);

now given any $t_2 > 0$ choose $0 < t_1 < t_2$ close enough to t_2 that $\tilde{\nu}(t_2) < \nu(t_1)$. Thus, it holds that

$$\|\mathcal{X}(t_2) - \mathcal{X}(t_1)\|_{\dot{\mathcal{F}}_{\tilde{\nu}(t_2)}^{1,1}} \rightarrow 0 \quad \text{as } t_1 \rightarrow t_2,$$

and therefore we have $\mathcal{X} \in C([0, T]; \dot{\mathcal{F}}_{\tilde{\nu}}^{1,1})$. Since $\tilde{\nu}_m \in (0, \nu_\infty)$ is an arbitrary number in an open interval, we conclude that $\mathcal{X} \in C([0, T]; \dot{\mathcal{F}}_{\tilde{\nu}}^{1,1})$. Finally, the analyticity in space for all positive times implies that $\mathcal{X} \in C([\varepsilon, T]; \dot{\mathcal{F}}_{\tilde{\nu}}^{s,1})$ for any $s \geq 0$, $\varepsilon > 0$, and $0 < \tilde{\nu} < \nu$. This regularity translates to F as well for $t \geq \varepsilon$. Therefore, one can consider $\partial_t \mathcal{X}(t_2) - \partial_t \mathcal{X}(t_1)$ for arbitrary $t_2, t_1 \geq \varepsilon$ to find in particular that $\partial_t \mathcal{X} \in C((0, T]; \mathcal{F}_{\tilde{\nu}}^{0,1})$.

We have proven that \mathcal{X} is a strong solution in the sense of Definition 1.1 as claimed in Theorem 1.2. In Section 4B we prove that this solution is unique.

We now prove the global-in-time energy inequality in (1-30).

Proof of (1-30). Equations (2-45) show decay of the higher frequencies (2-38) if we are able to control the nonlinear terms, for which we will need the constraint (2-40). Indeed, using (2-38) and the inequality $k(k - 1) \geq k^2/2$ for $k \geq 2$ implies

$$\frac{d}{dt} \|Y\|_{\dot{\mathcal{F}}_{\tilde{\nu}}^{1,1}} \leq -\left(\frac{A_e}{4} - \nu'(t)\right) \|Y\|_{\dot{\mathcal{F}}_{\tilde{\nu}}^{2,1}} + \|\mathcal{N}(X_c, X)\|_{\dot{\mathcal{F}}_{\tilde{\nu}}^{1,1}}, \tag{4-3}$$

where we have used that $\|P(k)^{-1}\| = 1$, and we can choose $\nu'(t)$ as small as we need. The goal is thus to obtain a bound like

$$\|\mathcal{N}(X_c, X)\|_{\dot{\mathcal{F}}_{\tilde{\nu}}^{1,1}} \leq C(\|X\|_{\dot{\mathcal{F}}_{\tilde{\nu}}^{1,1}}) \|Y\|_{\dot{\mathcal{F}}_{\tilde{\nu}}^{2,1}}, \tag{4-4}$$

with $C(\|X\|_{\dot{\mathcal{F}}_{\tilde{\nu}}^{1,1}}) \approx \|X\|_{\dot{\mathcal{F}}_{\tilde{\nu}}^{1,1}}$.

We proceed to complete the nonlinear estimate (4-4) to obtain the adequate sign in the balance (4-3). We insert the a priori bounds on F given by (3-55), (3-56), and (3-57), into the estimate (3-1) to obtain

$$\begin{aligned} \|\mathcal{N}\|_{\dot{\mathcal{F}}_{\tilde{\nu}}^{1,1}} &\leq 22\sqrt{2}A_e \frac{1 - A_\mu + |A_\mu|}{1 - A_\mu} D_1 \|X\|_{\dot{\mathcal{F}}_{\tilde{\nu}}^{1,1}} \|X\|_{\dot{\mathcal{F}}_{\tilde{\nu}}^{2,1}} + 147\sqrt{2} \frac{A_e}{1 - A_\mu} e^{\nu_\infty} D_2 C_8 \|X\|_{\dot{\mathcal{F}}_{\tilde{\nu}}^{1,1}} \|X\|_{\dot{\mathcal{F}}_{\tilde{\nu}}^{2,1}} \\ &\quad + 2250\sqrt{2}A_e \frac{|A_\mu|(1 + |A_\mu|)}{(1 - A_\mu)^2(1 + A_\mu)} D_3 D_4 \|X\|_{\dot{\mathcal{F}}_{\tilde{\nu}}^{1,1}} \|X\|_{\dot{\mathcal{F}}_{\tilde{\nu}}^{2,1}}, \end{aligned}$$

which finally gives the desired estimate

$$\|\mathcal{N}\|_{\dot{\mathcal{F}}_{\tilde{\nu}}^{1,1}} \leq 169\sqrt{2} \frac{A_e}{1 - A_\mu} D_5 \|X\|_{\dot{\mathcal{F}}_{\tilde{\nu}}^{1,1}} \|X\|_{\dot{\mathcal{F}}_{\tilde{\nu}}^{2,1}}, \tag{4-5}$$

where

$$D_5 = \frac{1}{169} \left(22(1 - A_\mu + |A_\mu|) D_1 + 147e^{\nu_\infty} D_2 C_8 + 2250 \frac{|A_\mu|(1 + |A_\mu|)}{(1 - A_\mu)(1 + A_\mu)} D_3 D_4 \right), \tag{4-6}$$

and C_8, D_1, D_2, D_3, D_4 , are given by (3-46), (3-54), and (3-78). Recalling the equivalence (2-34) and inserting the above bound into (4-3), we obtain

$$\frac{d}{dt} \|Y\|_{\dot{\mathcal{F}}_{\tilde{\nu}}^{1,1}} \leq -A_e \left(\frac{1}{4} - \frac{\nu'(t)}{A_e} - 169\sqrt{2} \frac{D_5}{1 - A_\mu} \|X\|_{\dot{\mathcal{F}}_{\tilde{\nu}}^{1,1}} \right) \|Y\|_{\dot{\mathcal{F}}_{\tilde{\nu}}^{2,1}}. \tag{4-7}$$

Since $v'(t) = v_\infty/(1+t)^2$ and $v_\infty > 0$ in (1-27) can be chosen arbitrarily small, if the condition

$$1 - 676\sqrt{2} \frac{D_5(\|\mathbf{X}\|_{\dot{J}_v^{1,1}})}{1 - A_\mu} \|\mathbf{X}\|_{\dot{J}_v^{1,1}} > 0 \tag{4-8}$$

holds initially, where D_5 is defined in (4-6), then the fact that D_5 decreases as $\|\mathbf{X}\|_{\dot{J}_v^{1,1}}$ decreases guarantees that this condition is propagated in time. For the same reasons, this condition can be stated as a smallness condition for $\|\mathbf{X}_0\|_{\dot{J}_v^{1,1}}$ as

$$\|\mathbf{X}_0\|_{\dot{J}_v^{1,1}} < k(A_\mu), \tag{4-9}$$

with k a function defined implicitly via (4-8) (see also Figure 1). Because D_5 is increasing on $\|\mathbf{X}\|_{\dot{J}_v^{1,1}}$, we have the explicit upper bound

$$k(A_\mu) < \frac{1 - A_\mu}{676\sqrt{2}D_5(0)}, \tag{4-10}$$

where

$$\begin{aligned} & \frac{1 - A_\mu}{676\sqrt{2}D_5(0)} \\ &= \left(\frac{588\sqrt{2}}{1 - A_\mu} + 88\sqrt{2} \left(1 + \frac{|A_\mu|}{1 - A_\mu} \right) + \frac{9\sqrt{2}|A_\mu|(1 + |A_\mu|)}{(1 - A_\mu)(1 + A_\mu)} \left(112 \left(1 + \frac{|A_\mu|}{1 - A_\mu} \right) + \frac{888}{1 - A_\mu} \right) \right)^{-1}. \end{aligned}$$

Then, for small enough $\|\mathbf{X}\|_{\dot{J}_v^{1,1}}$, the upper bound in (4-10) approximates the actual value of $k(A_\mu)$. Therefore,

$$\|\mathbf{Y}\|_{\dot{J}_v^{1,1}}(t) + \frac{A_e}{4} \mathcal{C} \int_0^t \|\mathbf{Y}\|_{\dot{J}_v^{2,1}}(\tau) d\tau \leq \|\mathbf{Y}_0\|_{\dot{J}_v^{1,1}}, \tag{4-11}$$

with

$$\mathcal{C} = \mathcal{C}(\|\mathbf{X}_0\|_{\dot{J}_v^{1,1}}, A_\mu, v_\infty) = 1 - 4 \frac{v'(t)}{A_e} - 676\sqrt{2} \frac{D_5(\|\mathbf{X}_0\|_{\dot{J}_v^{1,1}})}{1 - A_\mu} \|\mathbf{X}_0\|_{\dot{J}_v^{1,1}}. \tag{4-12}$$

Moreover, since $\|\mathbf{Y}\|_{\dot{J}_v^{1,1}} \leq \|\mathbf{Y}\|_{\dot{J}_v^{2,1}}$, the inequality (4-7) gives

$$\frac{d}{dt} \|\mathbf{Y}\|_{\dot{J}_v^{1,1}} \leq -\frac{A_e}{4} \mathcal{C} \|\mathbf{Y}\|_{\dot{J}_v^{1,1}},$$

and thus

$$\|\mathbf{Y}\|_{\dot{J}_v^{1,1}} \leq \|\mathbf{Y}_0\|_{\dot{J}_v^{1,1}} e^{-(A_e/4)\mathcal{C}t}. \tag{4-13}$$

This completes the decay estimate.

The control of the zero frequency follows from (2-47) with

$$|\widehat{\mathbf{X}}_c(0)| \leq |\widehat{\mathbf{X}}_{0,c}(0)| + \int_0^t |\widehat{\mathcal{N}}(\mathbf{X}_c, \mathbf{X})(0)| d\tau. \tag{4-14}$$

Notice that the estimates of the nonlinear terms in $\mathcal{F}^{0,1}$ can be done as in Section 3A and yield the bound

$$|\widehat{\mathcal{N}}(\mathbf{X}_c, \mathbf{X})(0)| \leq \|\mathcal{N}(\mathbf{X}_c, \mathbf{X})\|_{\mathcal{F}^{0,1}} \leq A_e \frac{\widetilde{D}_5}{1 - A_\mu} \|\mathbf{X}\|_{\dot{J}_v^{1,1}} \|\mathbf{X}\|_{\dot{J}_v^{2,1}},$$

where \widetilde{D}_5 is a constant that plays the role of D_5 . Recalling (2-34) and the energy balance (4-11), we introduce this bound back to (4-14) to conclude

$$|\widehat{\mathbf{X}}_c(0)| \leq |\widehat{\mathbf{X}}_{0,c}(0)| + \widetilde{\mathcal{C}} \|\mathbf{X}_0\|_{\dot{J}_v^{1,1}}^2,$$

with

$$\tilde{C} = \frac{\tilde{D}_5}{(1 - A_\mu)\mathcal{C}}, \tag{4-15}$$

and D_5, \mathcal{C} given in (4-6) and (4-12) respectively.

Finally, the decay (4-13) applied to (2-46) yields

$$\frac{1}{2}R(t)^2 = |\widehat{Y}_{c,2}(1)|^2 \rightarrow \frac{1}{2} \quad \text{as } t \rightarrow +\infty,$$

showing the exponentially fast convergence to a uniformly parametrized circle of area π . □

4B. Uniqueness. Consider two solutions $\mathcal{X} = X_c + X$ and $\tilde{\mathcal{X}} = \tilde{X}_c + \tilde{X}$ with initial data \mathcal{X}_0 and $\tilde{\mathcal{X}}_0$ in $\mathcal{F}^{1,1}$. Recalling the system in the \mathcal{Y} -variables (2-32), we have

$$\frac{d}{dt} \|\mathcal{Y} - \tilde{\mathcal{Y}}\|_{\dot{\mathcal{F}}_v^{1,1}} \leq -\left(\frac{A_e}{4} - \nu'(t)\right) \|\mathcal{Y} - \tilde{\mathcal{Y}}\|_{\dot{\mathcal{F}}_v^{2,1}} + \sqrt{2} \|\mathcal{N}(X_c, X) - \mathcal{N}(\tilde{X}_c, \tilde{X})\|_{\dot{\mathcal{F}}_v^{1,1}}, \tag{4-16}$$

and

$$|\widehat{Y}_c(0) - \widehat{\tilde{Y}}_c(0)| \leq |\widehat{Y}_{0,c}(0) - \widehat{\tilde{Y}}_{0,c}(0)| + \int_0^t |P(0)^{-1} \widehat{\mathcal{N}(X_c, X)}(0) - P(0)^{-1} \widehat{\mathcal{N}(\tilde{X}_c, \tilde{X})}(0)| d\tau. \tag{4-17}$$

Notice that, in comparison with (4-3), we are including in the left-hand side of (4-16) the terms corresponding to (the first frequency of) the circle part,

$$2|\widehat{Y}_{c,2}(1) - \widehat{\tilde{Y}}_{c,2}(1)|.$$

Although these terms are neutral with respect to the dissipative linear operator, whenever they appear on the right-hand side, we will be able to absorb them by using Grönwall’s lemma and (4-11) (which both Y and \tilde{Y} satisfy). Notice further that since the nonlinear terms do not contain the zero frequency of \mathcal{Y} , i.e., $\widehat{Y}_c(0)$, equation (4-17) implies

$$|\widehat{Y}_c(0) - \widehat{\tilde{Y}}_c(0)| = 0, \tag{4-18}$$

once we show from (4-16) that $\|\mathcal{Y} - \tilde{\mathcal{Y}}\|_{\dot{\mathcal{F}}_v^{1,1}} = 0$. Thus we proceed to deal with (4-16).

The difference between the nonlinear terms in (4-16) is split in four, according to (3-5), so that we have

$$\begin{aligned} \|\mathcal{N}(X_c, X) - \mathcal{N}(\tilde{X}_c, \tilde{X})\|_{\dot{\mathcal{F}}_v^{1,1}} &\leq \|\mathcal{N}_1(X_c, X) - \mathcal{N}_1(\tilde{X}_c, \tilde{X})\|_{\mathcal{F}_v^{0,1}} + \|\mathcal{N}_2(X_c, X) - \mathcal{N}_2(\tilde{X}_c, \tilde{X})\|_{\mathcal{F}_v^{0,1}} \\ &\quad + \|\mathcal{N}_3(X_c, X) - \mathcal{N}_3(\tilde{X}_c, \tilde{X})\|_{\mathcal{F}_v^{0,1}} + \|\mathcal{N}_4(X_c, X) - \mathcal{N}_4(\tilde{X}_c, \tilde{X})\|_{\mathcal{F}_v^{0,1}}. \end{aligned} \tag{4-19}$$

We start by explaining the estimate corresponding to the first subterm $\mathcal{N}_{1,1}$ in detail (see (3-6)), and later we will explain the general procedure. We have

$$\mathcal{N}_{1,1}(X_c, X)(\theta) - \mathcal{N}_{1,1}(\tilde{X}_c, \tilde{X})(\theta) = \mathcal{Q}_1 + \mathcal{Q}_2 + \mathcal{Q}_3 + \mathcal{Q}_4, \tag{4-20}$$

where

$$\begin{aligned} \mathcal{Q}_1 &= \left(-\frac{1}{4\pi R^2} + \frac{1}{4\pi \widetilde{R}^2}\right) \int_{\mathbb{S}} \partial_\theta \Delta_\eta X_c(\theta)^T \Delta_\eta X(\theta) F_L(\eta) d\eta, \\ \mathcal{Q}_2 &= \frac{1}{4\pi \widetilde{R}^2} \int_{\mathbb{S}} (\partial_\theta \Delta_\eta \tilde{X}_c(\theta) - \partial_\theta \Delta_\eta X_c(\theta))^T \Delta_\eta \tilde{X}(\theta) \widetilde{F}_L(\eta) d\eta, \end{aligned}$$

$$\begin{aligned} \mathbf{Q}_3 &= \frac{1}{4\pi \tilde{R}^2} \int_{\mathbb{S}} \partial_\theta \Delta_\eta \mathbf{X}_c(\theta)^T (\Delta_\eta \tilde{\mathbf{X}}(\theta) - \Delta_\eta \mathbf{X}(\theta)) \tilde{\mathbf{F}}_L(\eta) d\eta, \\ \mathbf{Q}_4 &= \frac{1}{4\pi \tilde{R}^2} \int_{\mathbb{S}} \partial_\theta \Delta_\eta \mathbf{X}_c(\theta)^T \Delta_\eta \mathbf{X}(\theta) (\tilde{\mathbf{F}}_L(\eta) - \mathbf{F}_L(\eta)) d\eta. \end{aligned}$$

For the first term, we need to estimate the difference between R and \tilde{R} . Recalling (2-40), where $|\hat{Y}_{c,2}(1)|^2 = R^2/2$ with $R^2 = a^2 + b^2$, we have

$$\begin{aligned} |R^2 - \tilde{R}^2| &= \left| -2 \sum_{k \geq 1} k (|\hat{Y}_2(k)|^2 - |\hat{Y}_1(k)|^2) + 2 \sum_{k \geq 1} k (|\hat{Y}_2(k)|^2 - |\hat{Y}_1(k)|^2) \right| \\ &\leq 2 \sum_{k \geq 1} k (|\hat{Y}_2(k)| - |\tilde{Y}_2(k)|)(|\hat{Y}_2(k)| + |\tilde{Y}_2(k)|) + |\hat{Y}_1(k) - \tilde{Y}_1(k)| (|\hat{Y}_1(k)| + |\tilde{Y}_1(k)|). \end{aligned}$$

Further note for $j = 1, 2$ that

$$|\hat{Y}_j(k) - \tilde{Y}_j(k)| \leq |\hat{Y}_j(k) - \tilde{Y}_j(k)|,$$

and on the \mathbb{S} domain we have $|\hat{Y}_j(k)| \leq \|Y_j\|_{L^\infty(\mathbb{S})} \leq \|Y_j\|_{\dot{J}^{0,1}}$. We conclude

$$|R^2 - \tilde{R}^2| \leq (\|Y\|_{\dot{J}^{0,1}} + \|\tilde{Y}\|_{\dot{J}^{0,1}}) \|Y - \tilde{Y}\|_{\dot{J}^{1,1}}.$$

Therefore, using also (2-41), we obtain

$$\left| \frac{1}{R^2} - \frac{1}{\tilde{R}^2} \right| = \left| \frac{\tilde{R}^2 - R^2}{R^2 \tilde{R}^2} \right| \leq \frac{\|Y\|_{\dot{J}^{0,1}} + \|\tilde{Y}\|_{\dot{J}^{0,1}}}{\sqrt{1 - \frac{1}{2}\|Y\|_{\dot{J}^{\frac{1}{2},1}}} \sqrt{1 - \frac{1}{2}\|\tilde{Y}\|_{\dot{J}^{\frac{1}{2},1}}}} \|Y - \tilde{Y}\|_{\dot{J}^{1,1}}.$$

In particular, for a constant $c(\|Y\|_{\dot{J}^{1,1}}, \|\tilde{Y}\|_{\dot{J}^{1,1}}) > 0$, we can write

$$\left| \frac{1}{R^2} - \frac{1}{\tilde{R}^2} \right| \leq c(\|Y\|_{\dot{J}^{1,1}}, \|\tilde{Y}\|_{\dot{J}^{1,1}}) \|Y - \tilde{Y}\|_{\dot{J}^{1,1}}.$$

Then, the bound for \mathbf{Q}_1 follows as in the estimate for the term (3-8), we obtain

$$\|\mathbf{Q}_1\|_{\mathcal{F}_v^{0,1}} \leq c(\|Y\|_{\dot{J}_v^{1,1}}, \|\tilde{Y}\|_{\dot{J}_v^{1,1}}) \|\mathbf{F}_L\|_{\mathcal{F}_v^{0,1}} \|Y - \tilde{Y}\|_{\dot{J}^{1,1}},$$

and introducing the estimate for \mathbf{F}_L from (3-56) we have

$$\|\mathbf{Q}_1\|_{\mathcal{F}_v^{0,1}} \leq A_e c(\|Y\|_{\dot{J}_v^{1,1}}, \|\tilde{Y}\|_{\dot{J}_v^{1,1}}, A_\mu, \nu_\infty) \|Y\|_{\dot{J}_v^{2,1}} \|Y - \tilde{Y}\|_{\dot{J}^{1,1}},$$

which is trivially bounded by

$$\|\mathbf{Q}_1\|_{\mathcal{F}_v^{0,1}} \leq A_e c(\|Y\|_{\dot{J}_v^{1,1}}, \|\tilde{Y}\|_{\dot{J}_v^{1,1}}, A_\mu, \nu_\infty) \|Y\|_{\dot{J}_v^{2,1}} \|Y - \tilde{Y}\|_{\dot{J}^{1,1}}.$$

It is now clear that (4-11) allows us to control this term by Grönwall’s lemma in (4-16).

We proceed to estimate \mathbf{Q}_2 in (4-20). Following the steps in (3-8), but maintaining the difference between $\tilde{\mathbf{X}}_c$ and \mathbf{X}_c together, we find that

$$\begin{aligned} \|\mathbf{Q}_2\|_{\mathcal{F}_v^{0,1}} &\leq c(\|Y\|_{\dot{J}_v^{1,1}}, \|\tilde{Y}\|_{\dot{J}_v^{1,1}}, A_\mu, \nu_\infty) \|\tilde{Y}\|_{\dot{J}_v^{2,1}} |\hat{Y}_{c,2}(1) - \tilde{Y}_{c,2}(1)| \\ &\leq A_e c(\|Y\|_{\dot{J}_v^{1,1}}, \|\tilde{Y}\|_{\dot{J}_v^{1,1}}, A_\mu, \nu_\infty) \|\tilde{Y}\|_{\dot{J}_v^{2,1}} \|Y - \tilde{Y}\|_{\dot{J}^{1,1}}, \end{aligned}$$

and it is thus controlled in the same way. The bound for \mathcal{Q}_3 follows exactly as in (3-8) and has the same structure as the bound for \mathcal{Q}_2 .

Finally, we are left with \mathcal{Q}_4 , for which we have

$$\mathcal{Q}_4 \leq c(\|Y\|_{\dot{F}_v^{1,1}}, \|\tilde{Y}\|_{\dot{F}_v^{1,1}})\|\tilde{F}_L - F_L\|_{\mathcal{F}_v^{0,1}}.$$

We emphasize that the constant above is given *exactly* by the one for $\mathcal{N}_{1,1}$ in (3-14). The estimate for $\tilde{F}_L - F_L$ follows from (2-25) (compare to (3-56)); we have

$$\|\tilde{F}_L - F_L\|_{\mathcal{F}_v^{0,1}} \leq 2A_e\|\tilde{X} - X\|_{\dot{F}_v^{2,1}} + 2A_e\frac{|A_\mu|}{1 - A_\mu}\|\tilde{X} - X\|_{\dot{F}_v^{1,1}}, \tag{4-21}$$

so, moving to the Y -variable, we obtain

$$\mathcal{Q}_4 \leq A_e c(\|Y\|_{\dot{F}_v^{1,1}}, \|\tilde{Y}\|_{\dot{F}_v^{1,1}})\|\tilde{Y} - Y\|_{\dot{F}_v^{2,1}} + A_e c(\|Y\|_{\dot{F}_v^{1,1}}, \|\tilde{Y}\|_{\dot{F}_v^{1,1}}, A_\mu, \nu_\infty)\|\tilde{Y} - Y\|_{\dot{F}_v^{1,1}},$$

and therefore trivially we have

$$\mathcal{Q}_4 \leq A_e c(\|Y\|_{\dot{F}_v^{1,1}}, \|\tilde{Y}\|_{\dot{F}_v^{1,1}})\|\tilde{Y} - Y\|_{\dot{F}_v^{2,1}} + A_e c(\|Y\|_{\dot{F}_v^{1,1}}, \|\tilde{Y}\|_{\dot{F}_v^{1,1}}, A_\mu, \nu_\infty)\|\tilde{\mathcal{Y}} - \mathcal{Y}\|_{\dot{F}_v^{1,1}}.$$

Although in this section we are denoting by c all constants (possibly depending on $\|Y\|_{\dot{F}_v^{1,1}}, \|\tilde{Y}\|_{\dot{F}_v^{1,1}}, A_\mu, \nu_\infty$), it is important to notice that the constant in front of the high-order term $\|\tilde{Y} - Y\|_{\dot{F}_v^{2,1}}$ is less than or equal to the one appearing in the nonlinear estimates from Section 3A. This will allow us to absorb these terms using the negative sign coming from the dissipative linear term without additional conditions on the initial data other than the one needed for the earlier existence proof.

In summary, so far we have obtained

$$\begin{aligned} \|\mathcal{N}_{1,1}(X_c, X) - \mathcal{N}_{1,1}(\tilde{X}_c, \tilde{X})\|_{\mathcal{F}_v^{0,1}} &\leq A_e c(\|Y\|_{\dot{F}_v^{1,1}}, \|\tilde{Y}\|_{\dot{F}_v^{1,1}})\|\tilde{Y} - Y\|_{\dot{F}_v^{2,1}} \\ &\quad + A_e g(\|Y\|_{\dot{F}_v^{1,1}}, \|\tilde{Y}\|_{\dot{F}_v^{1,1}}, \|Y\|_{\dot{F}_v^{2,1}}, \|\tilde{Y}\|_{\dot{F}_v^{2,1}}, A_\mu, \nu_\infty)\|\tilde{\mathcal{Y}} - \mathcal{Y}\|_{\dot{F}_v^{1,1}}, \end{aligned}$$

where g is a function whose L^1 -in-time norm is bounded independently of time in terms of the initial data $\|Y_0\|_{\mathcal{F}^{0,1}}, \|\tilde{Y}_0\|_{\mathcal{F}^{0,1}}$. Therefore the second term above can be controlled in (4-16) after using the Grönwall inequality.

Following the same steps for all the terms corresponding to \mathcal{N}_1 from (3-6), it is clear that one obtains

$$\begin{aligned} \|\mathcal{N}_1(X_c, X) - \mathcal{N}_1(\tilde{X}_c, \tilde{X})\|_{\mathcal{F}_v^{0,1}} &\leq c(\|Y\|_{\dot{F}_v^{1,1}}, \|\tilde{Y}\|_{\dot{F}_v^{1,1}})\|\tilde{F}_L - F_L\|_{\mathcal{F}_v^{0,1}} \\ &\quad + A_e g(\|Y\|_{\dot{F}_v^{1,1}}, \|\tilde{Y}\|_{\dot{F}_v^{1,1}}, \|Y\|_{\dot{F}_v^{2,1}}, \|\tilde{Y}\|_{\dot{F}_v^{2,1}}, A_\mu, \nu_\infty)\|\tilde{\mathcal{Y}} - \mathcal{Y}\|_{\dot{F}_v^{1,1}}, \end{aligned}$$

where we use the same letter g to denote another L^1 -in-time function as explained above and the constant in front of $\|\tilde{F}_L - F_L\|_{\mathcal{F}_v^{0,1}}$ is exactly given by the one in (3-23). Since the coefficient of the higher-order term in the bound (4-21) is smaller than the one in (3-56), we guarantee that

$$\begin{aligned} \|\mathcal{N}_1(X_c, X) - \mathcal{N}_1(\tilde{X}_c, \tilde{X})\|_{\mathcal{F}_v^{0,1}} &\leq A_e c(\|Y\|_{\dot{F}_v^{1,1}}, \|\tilde{Y}\|_{\dot{F}_v^{1,1}})\|\tilde{Y} - Y\|_{\dot{F}_v^{2,1}} \\ &\quad + A_e g(\|Y\|_{\dot{F}_v^{1,1}}, \|\tilde{Y}\|_{\dot{F}_v^{1,1}}, \|Y\|_{\dot{F}_v^{2,1}}, \|\tilde{Y}\|_{\dot{F}_v^{2,1}}, A_\mu, \nu_\infty)\|\tilde{\mathcal{Y}} - \mathcal{Y}\|_{\dot{F}_v^{1,1}}. \end{aligned} \tag{4-22}$$

Now, we realize that the same idea applies to the other nonlinear terms in (4-19). In \mathcal{N}_2 there are not high-order terms to absorb, since the expression of F_0 (3-58) only depends on the circle part, which has to be controlled via Grönwall. The term \mathcal{N}_3 will provide an estimate like the one above for \mathcal{N}_1 , where the constant in front of $\|\tilde{Y} - Y\|_{\dot{J}_v^{2,1}}$ is smaller than (3-48) for the same reasons given before. Finally, the same can be said for \mathcal{N}_4 , but with an analogous estimate to (4-21) for the difference $F_N - \tilde{F}_N$. It follows in the same way as the estimate (3-57), so we omit details to avoid repetition.

The final estimate for the difference of the nonlinear terms in (4-19) has then the form (4-22), with a coefficient of the highest-order norm smaller than the coefficient of the norm with the highest-order derivative in (4-5). Therefore, under condition (1-29), the highest-regularity terms in the nonlinear upper bound can be absorbed by the dissipation in (4-16) and thus

$$\frac{d}{dt} \|\mathcal{Y} - \tilde{\mathcal{Y}}\|_{\dot{J}_v^{1,1}} \leq A_e g(\|Y\|_{\dot{J}_v^{1,1}}, \|\tilde{Y}\|_{\dot{J}_v^{1,1}}, \|Y\|_{\dot{J}_v^{2,1}}, \|\tilde{Y}\|_{\dot{J}_v^{2,1}}, A_\mu, \nu_\infty) \|\mathcal{Y} - \tilde{\mathcal{Y}}\|_{\dot{J}_v^{1,1}},$$

which provides for all time via Grönwall that

$$\|\mathcal{Y} - \tilde{\mathcal{Y}}\|_{\dot{J}_v^{1,1}} \leq c(\|Y_0\|_{\dot{J}_v^{1,1}}, \|\tilde{Y}_0\|_{\dot{J}_v^{1,1}}, A_e, A_\mu, \nu_\infty) \|\mathcal{Y}_0 - \tilde{\mathcal{Y}}_0\|_{\dot{J}_v^{1,1}}.$$

We conclude that $\|\mathcal{Y} - \tilde{\mathcal{Y}}\|_{\dot{J}_v^{1,1}} = 0$. Together with (4-18), this completes the proof. \square

Acknowledgements

García-Juárez was partially supported by the grant MTM2017-89976-P (Spain), the AMS-Simons Travel Grant, and by the ERC through the Starting Grant projects H2020-EU.1.1-639227 and ERC-StG-CAPA-852741. Mori was partially supported by the NSF grants DMS-1907583 and DMS-2042144 (USA) and the Math+X award from the Simons Foundation. Strain was partially supported by the NSF grants DMS-1764177 and DMS-2055271 (USA).

References

- [Ambrose and Siegel 2017] D. M. Ambrose and M. Siegel, “Well-posedness of two-dimensional hydroelastic waves”, *Proc. Roy. Soc. Edinburgh Sect. A* **147**:3 (2017), 529–570. [MR](#) [Zbl](#)
- [Boulakia et al. 2012] M. Boulakia, E. L. Schwindt, and T. Takahashi, “Existence of strong solutions for the motion of an elastic structure in an incompressible viscous fluid”, *Interfaces Free Bound.* **14**:3 (2012), 273–306. [MR](#) [Zbl](#)
- [Cameron 2019] S. Cameron, “Global well-posedness for the two-dimensional Muskat problem with slope less than 1”, *Anal. PDE* **12**:4 (2019), 997–1022. [MR](#) [Zbl](#)
- [Cameron 2020] S. Cameron, “Global wellposedness for the 3D Muskat problem with medium size slope”, preprint, 2020. [arXiv 2002.00508](#)
- [Cheng and Shkoller 2010] C. H. A. Cheng and S. Shkoller, “The interaction of the 3D Navier–Stokes equations with a moving nonlinear Koiter elastic shell”, *SIAM J. Math. Anal.* **42**:3 (2010), 1094–1155. [MR](#) [Zbl](#)
- [Cheng et al. 2007] C. H. A. Cheng, D. Coutand, and S. Shkoller, “Navier–Stokes equations interacting with a nonlinear elastic biofluid shell”, *SIAM J. Math. Anal.* **39**:3 (2007), 742–800. [MR](#) [Zbl](#)
- [Constantin et al. 2013] P. Constantin, D. Córdoba, F. Gancedo, and R. M. Strain, “On the global existence for the Muskat problem”, *J. Eur. Math. Soc.* **15**:1 (2013), 201–227. [MR](#) [Zbl](#)
- [Constantin et al. 2016] P. Constantin, D. Córdoba, F. Gancedo, L. Rodríguez-Piazza, and R. M. Strain, “On the Muskat problem: global in time results in 2D and 3D”, *Amer. J. Math.* **138**:6 (2016), 1455–1494. [MR](#) [Zbl](#)

- [Gancedo et al. 2019a] F. Gancedo, E. García-Juárez, N. Patel, and R. Strain, “Global regularity for gravity unstable Muskat bubbles”, 2019. To appear in Mem. Amer. Math. Soc. [arXiv 1902.02318](#)
- [Gancedo et al. 2019b] F. Gancedo, E. García-Juárez, N. Patel, and R. M. Strain, “On the Muskat problem with viscosity jump: global in time results”, *Adv. Math.* **345** (2019), 552–597. [MR](#) [Zbl](#)
- [Gancedo et al. 2020] F. Gancedo, R. Granero-Belinchón, and S. Scrobogna, “Surface tension stabilization of the Rayleigh–Taylor instability for a fluid layer in a porous medium”, *Ann. Inst. H. Poincaré C Anal. Non Linéaire* **37**:6 (2020), 1299–1343. [MR](#) [Zbl](#)
- [Li 2021] H. Li, “Stability of the Stokes immersed boundary problem with bending and stretching energy”, *J. Funct. Anal.* **281**:9 (2021), art. id. 109204. [MR](#) [Zbl](#)
- [Li and Ito 2006] Z. Li and K. Ito, *The immersed interface method: numerical solutions of PDEs involving interfaces and irregular domains*, Frontiers in Appl. Math. **33**, Soc. Indust. Appl. Math., Philadelphia, PA, 2006. [MR](#) [Zbl](#)
- [Lin and Tong 2019] F.-H. Lin and J. Tong, “Solvability of the Stokes immersed boundary problem in two dimensions”, *Comm. Pure Appl. Math.* **72**:1 (2019), 159–226. [MR](#) [Zbl](#)
- [Liu and Ambrose 2017] S. Liu and D. M. Ambrose, “Well-posedness of two-dimensional hydroelastic waves with mass”, *J. Differential Equations* **262**:9 (2017), 4656–4699. [MR](#) [Zbl](#)
- [Lunardi 1995] A. Lunardi, *Analytic semigroups and optimal regularity in parabolic problems*, Progr. Nonlinear Differential Equations Appl. **16**, Birkhäuser, Basel, 1995. [MR](#) [Zbl](#)
- [Mori et al. 2019] Y. Mori, A. Rodenberg, and D. Spirn, “Well-posedness and global behavior of the Peskin problem of an immersed elastic filament in Stokes flow”, *Comm. Pure Appl. Math.* **72**:5 (2019), 887–980. [MR](#) [Zbl](#)
- [Muha and Canić 2013] B. Muha and S. Canić, “Existence of a weak solution to a nonlinear fluid-structure interaction problem modeling the flow of an incompressible, viscous fluid in a cylinder with deformable walls”, *Arch. Ration. Mech. Anal.* **207**:3 (2013), 919–968. [MR](#) [Zbl](#)
- [Peskin 2002] C. S. Peskin, “The immersed boundary method”, *Acta Numer.* **11** (2002), 479–517. [MR](#) [Zbl](#)
- [Plotnikov and Toland 2011] P. I. Plotnikov and J. F. Toland, “Modelling nonlinear hydroelastic waves”, *Philos. Trans. R. Soc. Lond. Ser. A Math. Phys. Eng. Sci.* **369**:1947 (2011), 2942–2956. [MR](#) [Zbl](#)
- [Plotnikov and Toland 2012] P. I. Plotnikov and J. F. Toland, “Strain-gradient theory of hydroelastic travelling waves and Young measures of their singular limits”, *Calc. Var. Partial Differential Equations* **44**:1-2 (2012), 153–197. [MR](#) [Zbl](#)
- [Pozrikidis 1992] C. Pozrikidis, *Boundary integral and singularity methods for linearized viscous flow*, Cambridge Univ. Press, 1992. [MR](#) [Zbl](#)
- [Prüss and Simonett 2009] J. Prüss and G. Simonett, “Analysis of the boundary symbol for the two-phase Navier–Stokes equations with surface tension”, pp. 265–285 in *Nonlocal and abstract parabolic equations and their applications* (Będlewo, Poland, 2007), edited by P. B. Mucha et al., Banach Center Publ. **86**, Polish Acad. Sci. Inst. Math., Warsaw, 2009. [MR](#) [Zbl](#)
- [Prüss and Simonett 2016] J. Prüss and G. Simonett, *Moving interfaces and quasilinear parabolic evolution equations*, Monogr. Math. **105**, Birkhäuser, Cham, 2016. [MR](#) [Zbl](#)
- [Richter 2017] T. Richter, *Fluid-structure interactions: models, analysis and finite elements*, Lect. Notes in Computational Sci. Eng. **118**, Springer, 2017. [MR](#) [Zbl](#)
- [Rodenberg 2018] A. Rodenberg, *2D Peskin problems of an immersed elastic filament in Stokes flow*, Ph.D. thesis, University of Minnesota, 2018, available at <https://www.proquest.com/docview/2082409498>.
- [Shimizu 2009] S. Shimizu, “Maximal regularity principle and its applications to free boundary problems for fluids”, *Sūgaku* **61**:4 (2009), 352–375. In Japanese; translated in *Sugaku Expo.* **25**:1 (2012), 105–130. [MR](#)
- [Simon 1987] J. Simon, “Compact sets in the space $L^p(0, T; B)$ ”, *Ann. Mat. Pura Appl.* (4) **146** (1987), 65–96. [MR](#) [Zbl](#)
- [Tong 2021] J. Tong, “Regularized Stokes immersed boundary problems in two dimensions: well-posedness, singular limit, and error estimates”, *Comm. Pure Appl. Math.* **74**:2 (2021), 366–449. [MR](#) [Zbl](#)
- [Tryggvason et al. 2001] G. Tryggvason, B. Bunner, A. Esmaeeli, D. Juric, N. Al-Rawahi, W. Tauber, J. Han, S. Nas, and Y.-J. Jan, “A front-tracking method for the computations of multiphase flow”, *J. Comput. Phys.* **169**:2 (2001), 708–759. [MR](#) [Zbl](#)

Received 16 Dec 2020. Revised 6 Jul 2021. Accepted 23 Aug 2021.

EDUARDO GARCÍA-JUÁREZ: egarcia12@us.es

Departament de Matemàtiques i Informàtica, Universitat de Barcelona, Gran Via de les Corts Catalanes, Barcelona, Spain

YOICHIRO MORI: y1mori@sas.upenn.edu

Department of Mathematics, University of Pennsylvania, David Rittenhouse Lab, Philadelphia, PA, United States

ROBERT M. STRAIN: strain@math.upenn.edu

Department of Mathematics, University of Pennsylvania, David Rittenhouse Lab, Philadelphia, PA, United States

Analysis & PDE

msp.org/apde

EDITORS-IN-CHIEF

Patrick Gérard [Universit  Paris Sud XI, France](mailto:patrick.gerard@universite-paris-saclay.fr)
patrick.gerard@universite-paris-saclay.fr
Cl ment Mouhot [Cambridge University, UK](mailto:c.mouhot@dpmms.cam.ac.uk)
c.mouhot@dpmms.cam.ac.uk

BOARD OF EDITORS

Massimiliano Berti	Scuola Intern. Sup. di Studi Avanzati, Italy berti@sissa.it	Werner M�ller	Universit�t Bonn, Germany mueller@math.uni-bonn.de
Zbigniew Błocki	Uniwersytet Jagielloński, Poland zbigniew.blocki@uj.edu.pl	Gilles Pisier	Texas A&M University, and Paris 6 pisier@math.tamu.edu
Charles Fefferman	Princeton University, USA cf@math.princeton.edu	Igor Rodnianski	Princeton University, USA irod@math.princeton.edu
Isabelle Gallagher	Universit� Paris-Diderot, IMJ-PRG, France gallagher@math.ens.fr	Yum-Tong Siu	Harvard University, USA siu@math.harvard.edu
Colin Guillarmou	Universit� Paris-Saclay, France colin.guillarmou@universite-paris-saclay.fr	Terence Tao	University of California, Los Angeles, USA tao@math.ucla.edu
Ursula Hamenstaedt	Universit�t Bonn, Germany ursula@math.uni-bonn.de	Michael E. Taylor	Univ. of North Carolina, Chapel Hill, USA met@math.unc.edu
Vadim Kaloshin	University of Maryland, USA vadim.kaloshin@gmail.com	Gunther Uhlmann	University of Washington, USA gunther@math.washington.edu
Izabella Laba	University of British Columbia, Canada ilaba@math.ubc.ca	Andr�s Vasy	Stanford University, USA andras@math.stanford.edu
Anna L. Mazzucato	Penn State University, USA alm24@psu.edu	Dan Virgil Voiculescu	University of California, Berkeley, USA dvv@math.berkeley.edu
Richard B. Melrose	Massachusetts Inst. of Tech., USA rbm@math.mit.edu	Jim Wright	University of Edinburgh, UK j.r.wright@ed.ac.uk
Frank Merle	Universit� de Cergy-Pontoise, France merle@ihes.fr	Maciej Zworski	University of California, Berkeley, USA zworski@math.berkeley.edu
William Minicozzi II	Johns Hopkins University, USA minicozz@math.jhu.edu		

PRODUCTION

production@msp.org

Silvio Levy, Scientific Editor


See inside back cover or msp.org/apde for submission instructions.

The subscription price for 2023 is US \$405/year for the electronic version, and \$630/year (+\$65, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues from the last three years and changes of subscriber address should be sent to MSP.

Analysis & PDE (ISSN 1948-206X electronic, 2157-5045 printed) at Mathematical Sciences Publishers, 798 Evans Hall #3840, c/o University of California, Berkeley, CA 94720-3840, is published continuously online.

APDE peer review and production are managed by EditFlow[®] from MSP.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

  2023 Mathematical Sciences Publishers

ANALYSIS & PDE

Volume 16 No. 3 2023

Global regularity for the nonlinear wave equation with slightly supercritical power MARIA COLOMBO and SILJA HAFFTER	613
Subelliptic wave equations are never observable CYRIL LETROUIT	643
Quantitative Alexandrov theorem and asymptotic behavior of the volume preserving mean curvature flow VESA JULIN and JOONAS NIINIKOSKI	679
A simple nuclear C^* -algebra with an internal asymmetry ILAN HIRSHBERG and N. CHRISTOPHER PHILLIPS	711
Partial regularity of Leray–Hopf weak solutions to the incompressible Navier–Stokes equations with hyperdissipation WOJCIECH S. OŻAŃSKI	747
The Peskin problem with viscosity contrast EDUARDO GARCÍA-JUÁREZ, YOICHIRO MORI and ROBERT M. STRAIN	785
Solution of the qc Yamabe equation on a 3-Sasakian manifold and the quaternionic Heisenberg group STEFAN IVANOV, IVAN MINCHEV and DIMITER VASSILEV	839
Garland’s method with Banach coefficients IZHAR OPPENHEIM	861