

# ANALYSIS & PDE

Volume 17

No. 1

2024

HUI ZHU

**PROPAGATION OF SINGULARITIES  
FOR GRAVITY-CAPILLARY WATER WAVES**



# PROPAGATION OF SINGULARITIES FOR GRAVITY-CAPILLARY WATER WAVES

HUI ZHU

We obtain two results of propagation for the gravity-capillary water wave system. The first result shows the propagation of oscillations and the spatial decay at infinity; the second result shows a microlocal smoothing effect under the nontrapping condition of the initial free surface. These results extend the works of Craig, Kappeler and Strauss (1995), Wunsch (1999) and Nakamura (2005) to quasilinear dispersive equations. These propagation results are stated for water waves with asymptotically flat free surfaces, of which we also obtain the existence. To prove these results, we generalize the paradifferential calculus of Bony (1979) to weighted Sobolev spaces and develop a semiclassical paradifferential calculus. We also introduce the quasihomogeneous wavefront sets which characterize, in a general manner, the oscillations and the spatial growth/decay of distributions.

1. Introduction	281
2. Quasihomogeneous microlocal analysis	290
3. Model equations	295
4. Paradifferential calculus	300
5. Asymptotically flat water waves	316
6. Propagation of singularities for water waves	326
List of notation	341
Acknowledgments	341
References	341

## 1. Introduction

We present two results on the propagation of singularities for the gravity-capillary water wave system, including a microlocal smoothing effect. To the best of our knowledge, these results are the first of this type for quasilinear dispersive equations. Before stating the main results, we shall first revisit classical results of propagation for the linear half-wave equation and the linear Schrödinger equation. They lead us to a more generalized concept of singularities which is adaptive to various dispersive equations.

**1A. Wavefront set and the linear half-wave equation.** If  $u \in \mathcal{D}'(M)$ , where  $M$  is a smooth manifold without boundary, then the singular support of  $u$ , denoted by  $\text{sing supp } u$ , is the smallest closed subset of  $M$  outside of which  $u$  is smooth. To study the propagation of singularities when  $u$  solves some partial

---

The author is partially supported by the grant “ANAÉ” ANR-13-BS01-0010-03 of the Agence Nationale de la Recherche, and the Allocation Doctorale of the École Normale Supérieure.

MSC2020: 35A01, 35A18, 35A21, 35S50, 76B15.

Keywords: water wave, smoothing effect, propagation of singularity, wavefront set.

differential equations, the information given by  $\text{sing supp } u$  is usually insufficient. Heuristically, if we consider singularities as accumulations of wavepackets with large wavenumbers, then this is because the propagation direction of a wavepacket is given by its wavenumber rather than its location. It is probably with this mindset that Hörmander [1971] introduced the concept of the wavefront set.

The wavefront set of  $u$ , denoted by  $\text{WF}(u)$ , lifts  $\text{sing supp } u$  to the cotangent bundle  $T^*M \setminus 0$  in the sense that a point  $x_0 \in M$  belongs to  $\text{sing supp } u$  if and only if there exists  $\xi_0 \neq 0$  such that  $(x_0, \xi_0) \in \text{WF}(u)$ . We shall recall an equivalent definition of  $\text{WF}(u)$  essentially due to [Guillemin and Sternberg 1977]: in local coordinates, a point  $(x_0, \xi_0) \in T^*M \setminus 0$  does not belong to  $\text{WF}(u)$  if and only if there exists  $a \in C_c^\infty(\mathbb{R}^{2d})$  with  $a(x_0, \xi_0) \neq 0$  such that  $\|a(x, hD_x)u\|_{L^2} = \mathcal{O}(h^\infty)$  for  $h \in (0, 1]$ . For the definition of the pseudodifferential operator  $a(x, hD_x)$ , see (1-7).

In terms of the wavefront set, Hörmander [1971] proved a propagation result for pseudodifferential equations of real principal type, improving previous works [Courant and Lax 1956; Lax 1957] on wave propagation.

**Theorem 1.1** [Hörmander 1971]. *Let  $M$  be a smooth manifold without boundary. Let  $P \in \Psi^1(M)$  admit a real principal symbol  $\sigma(P) = \sigma(P)(x, \xi) \in C^\infty(T^*M \setminus 0, \mathbb{R})$ , and let  $\Phi = \Phi_t(x, \xi) \in C^\infty(\mathbb{R} \times T^*M \setminus 0, T^*M \setminus 0)$  be the Hamiltonian flow of  $\sigma(P)$ . If  $u$  solves the Cauchy problem*

$$\begin{cases} \partial_t u + iPu = 0, \\ u(0) = u_0 \in L^2(M), \end{cases} \quad (1-1)$$

then for all  $(x_0, \xi_0) \in \text{WF}(u_0)$  and all  $t \in \mathbb{R}$ , we have  $\Phi_t(x_0, \xi_0) \in \text{WF}(u(t))$ .

In particular, if  $P = \sqrt{-\Delta_g}$  where  $g$  is a Riemannian metric on  $M$ , then (1-1) becomes the half-wave equation and  $\Phi$  is the corresponding cogeodesic flow on  $T^*M$ . Therefore, we conclude that, for solutions to the half-wave equation, microlocal singularities travel at speed 1 along cogeodesics. This gives a justification for the Huygens–Fresnel principal of wavefront propagation.

For the propagation of singularities for the semilinear wave equation, we refer to [Bony 1986; Lebeau 1989]. For the propagation and the reflection of singularities for the linear wave equation on manifolds with corners, see [Vasy 2008; Melrose, Vasy and Wunsch 2013].

**1B. The homogeneous wavefront set and the linear Schrödinger equation.** Hörmander’s theorem (Theorem 1.1) is untrue when the order of  $P$  is higher than 1. For example, the Schrödinger propagator  $e^{it\Delta/2}$  on  $\mathbb{R}^d$  sends  $\mathcal{E}'(\mathbb{R}^d)$  to  $C^\infty(\mathbb{R}^d)$  whenever  $t \neq 0$ . We conclude that singularities may appear and disappear along the Schrödinger flow. These phenomena of “microlocal smoothing effect” and “microlocal singularity formation” are due to the infinite speed of propagation of the Schrödinger equation, as wavepackets with large wavenumbers can travel to or back from infinity instantaneously.

The study of the infinite speed of propagation of the Schrödinger equation probably dates back to [Boutet de Monvel 1975; Lascar 1977; 1978]. They proved that space-time singularities, as elements of some space-time wavefront sets, travel along geodesics at an infinite speed. They did not obtain, however, a time-dependent propagation results for wavefront sets with respect to the space variable alone. The study of the smoothing effect for dispersive equations with an infinite speed of propagation was initiated

by Kato [1983], who proved a local smoothing effect for generalized KdV equations. Craig, Kappeler and Strauss [1995] proved microlocal smoothing effects for the linear Schrödinger equation under the nontrapping condition of the geometry. Their results were later refined by Wunsch [1999] who obtained a time-dependent propagation after understanding the transformation between singularities and quadratic oscillations at infinity. The simplest example is the identity

$$e^{it\Delta/2}\delta_{x_0}(x) = \frac{1}{(2\pi it)^{d/2}}e^{i|x-x_0|^2/(2t)},$$

where  $\delta_{x_0}$  is the Dirac measure at  $x_0 \in \mathbb{R}^d$ . Wunsch’s results were stated on Riemannian manifolds endowed with a scattering metric. He introduced the quadratic scattering wavefront set to characterize quadratic oscillations.

Similar results were later obtained, independently, by Nakamura [2005] via a simpler calculus but in a less general geometric setting — asymptotically Euclidean geometries, where he introduced the homogeneous wavefront set. By definition, if  $u \in \mathcal{S}'(\mathbb{R}^d)$ , then the homogeneous wavefront set  $\text{HWF}(u)$  is a subset of  $\mathbb{R}^{2d}$  whose complement consists of all  $(x_0, \xi_0)$  admitting a symbol  $a \in C_c^\infty(\mathbb{R}^{2d})$  with  $a(x_0, \xi_0) \neq 0$  such that  $\|a(hx, hD_x)u\|_{L^2} = \mathcal{O}(h^\infty)$  for  $h \in (0, 1]$ . It was proven by Ito [2006] that the quadratic scattering wavefront set and the homogeneous wavefront set are essentially equivalent in asymptotically Euclidean geometries. In fact, heuristically, if  $x_0 \neq 0$  and  $\xi_0 \neq 0$ , then the pseudodifferential operator  $a(hx, hD_x)$  is a microlocalization in the region of quadratic oscillation:

$$|x| \sim |\xi| \sim h^{-1}.$$

Take for example the free Schrödinger equation in  $\mathbb{R}^d$ , of which the dispersion relation is  $\omega = \frac{1}{2}|\xi|^2$ . A wave packet of frequency  $\xi \sim h^{-1}$  travels at the group velocity  $v = d\omega/d\xi = \xi \sim h^{-1}$ . The homogeneously scaled quantization  $a \mapsto a(hx, hD_x)$  thus allows us to keep up with the infinite speed of propagation and obtain an analogue of Hörmander’s theorem.

**Theorem 1.2** ([Nakamura 2005], similar results in [Wunsch 1999]). *Let  $g$  be an asymptotically Euclidean Riemannian metric on  $\mathbb{R}^d$ , meaning that there exists  $\epsilon > 0$  such that, for all  $\alpha \in \mathbb{N}^d$  and all  $i, j \in \{1, \dots, d\}$ , we have*

$$|\partial_x^\alpha (g_{ij}(x) - \delta_{ij})| \lesssim \langle x \rangle^{-|\alpha|-\epsilon}. \tag{1-2}$$

*Consider the Cauchy problem of the linear Schrödinger equation*

$$\begin{cases} i\partial_t u + \frac{1}{2}\Delta_g u = 0, \\ u(0) = u_0 \in L^2(\mathbb{R}^d). \end{cases}$$

*Then the following propagation results hold:*

- (1) *If  $(x_0, \xi_0) \in \text{HWF}(u_0)$  and  $t_0 \in \mathbb{R}$  such that  $\xi_0 \neq 0$  and  $x_0 + t\xi_0 \neq 0$  for all  $t$  between 0 and  $t_0$ , then  $(x_0 + t_0\xi_0, \xi_0) \in \text{HWF}(u(t_0))$ .*
- (2) *If  $(x_0, \xi_0) \in \text{WF}(u_0)$  is forwardly (resp. backwardly) nontrapping in the sense that the cogeodesic issued from  $(x_0, \xi_0)$ , denoted by  $\{(x_t, \xi_t)\}_{t \in \mathbb{R}}$  (with an abuse of notation), satisfies*

$$\lim_{t \rightarrow +\infty} |x_t| = +\infty \quad (\text{resp.} \quad \lim_{t \rightarrow -\infty} |x_t| = +\infty),$$

then there exists  $\xi_+ \in \mathbb{R}^d$  (resp.  $\xi_- \in \mathbb{R}^d$ ) satisfying  $\xi_{\pm} = \lim_{t \rightarrow \pm\infty} \xi_t$ , and moreover, for all  $t_0 > 0$  (resp.  $t_0 < 0$ ), we have

$$(t_0\xi_+, \xi_+) \in \text{HWF}(u(t_0)) \quad (\text{resp. } (t_0\xi_-, \xi_-) \in \text{HWF}(u(t_0))).$$

**Theorem 1.2(1)** studies the propagation of oscillations and spatial growth/decay for Schrödinger waves at infinity and we thus require the condition  $x_0 + t\xi_0 \neq 0$ . In  $\mathbb{R}^d$ , this result is a consequence of an Egorov-type argument and the commutation relation

$$[i\partial_t + \frac{1}{2}\Delta, a(t, hx, hD_x)] = (i\partial_t a - \xi \cdot \partial_x a)(t, hx, hD_x) + \mathcal{O}(h^2),$$

where  $a \in C_b^\infty(\mathbb{R} \times \mathbb{R}^{2d})$ . A similar argument works in asymptotically Euclidean geometries where we replace the role of the semiclassical quantization  $x \mapsto hx$  with the spatial decay of the metric  $g$ , i.e., the condition (1-2).

**Theorem 1.2(2)** is a microlocal smoothing effect: if  $(t_0\xi_{\pm}, \xi_{\pm})$  does not belong to  $\text{HWF}(u(t_0))$ , then  $(x_0, \xi_0)$  cannot be an element of  $\text{WF}(u_0)$ . This result is a refinement of the result in [Craig, Kappeler and Strauss 1995] and can be proven via a positive commutator estimate. In  $\mathbb{R}^d$ , this estimate has the form

$$[i\partial_t + \frac{1}{2}\Delta, a(t, x, hD_x)] \gtrsim \mathcal{O}(h^\infty),$$

where  $a$  is some well-chosen symbol. For related results, see [Doi 1996; 2000; Burq 2004] for the necessity of the nontrapping condition; see [Robbiano and Zuily 1999] for a microlocal analytic smoothing effect; see [Kenig, Ponce and Vega 1998; Szeftel 2005] for local and microlocal smoothing effects for the semilinear Schrödinger equation. We should also remark that Hörmander [1991] has also introduced an essentially equivalent counterpart of the homogeneous wavefront set to which a similar definition as that of Nakamura was given. See [Rodino and Wahlberg 2014; Schulz and Wahlberg 2017] for more comments. However, **Theorem 1.2(2)** is unable, via simply reversing the time, to show how oscillations at infinity form singularities along the Schrödinger flow. Indeed, the information about the locations of singularities is not contained in quadratic oscillations but rather in linear oscillations at infinity. See [Hassell and Wunsch 2005; Nakamura 2009] for more on this subject.

**1C. Quasihomogeneous wavefront set and the gravity-capillary water wave system.** The gravity-capillary water wave system describes the evolution of inviscid, incompressible and irrotational fluid with a free surface, in the presence of a gravitational field and the surface tension.

**1C1. Formulations of the gravity-capillary water wave system.** We shall first recall the Eulerian formulation of the gravity-capillary water wave system. The area occupied by the fluid is a time-dependent simply connected open subset of  $\mathbb{R}^{d+1}$  and is denoted by  $\Omega$ . The boundary of  $\Omega$  consists of two parts: the free surface  $\Sigma$  and the bottom  $\Gamma$ . The free surface of the fluid is a time-dependent hypersurface which is the graph of a function  $\eta = \eta(t, x)$ , where  $(t, x) \in \mathbb{R} \times \mathbb{R}^d$ , whereas the bottom is independent of time and is of depth  $b \in (0, \infty)$ . Therefore,

$$\Omega = \{-b < y < \eta\}, \quad \Sigma = \{y = \eta\}, \quad \Gamma = \{y = -b\}.$$

The Eulerian formulation describes water waves in the unknowns  $(\eta, v, P)$  where  $v : \Omega \rightarrow \mathbb{R}^d$  is the Eulerian vector field and  $P : \Omega \rightarrow \mathbb{R}$  is the pressure of the fluid:

$$\begin{cases} \partial_t v + v \cdot \nabla_{xy} v = -\nabla_{xy}(P + gy) & \text{(Euler equation),} \\ \nabla_{xy} \cdot v = 0 & \text{(incompressibility),} \\ \nabla_{xy} \times v = 0 & \text{(irrotationality),} \\ (v \cdot \mathbf{n})_{y=\eta} = \partial_t \eta / \langle \nabla \eta \rangle & \text{(kinetic condition at the free surface),} \\ (v \cdot \mathbf{n})_{y=-b} = 0 & \text{(kinetic condition at the bottom),} \\ -P|_{y=\eta} = \kappa H(\eta), & \text{(dynamic condition).} \end{cases} \tag{1-3}$$

Here  $g \in \mathbb{R}$  is the gravitational acceleration,  $\kappa > 0$  is the surface tension,  $\mathbf{n} : \partial\Omega \rightarrow \mathbb{S}^d$  denotes the exterior unit normal vector field of  $\partial\Omega$ , while

$$H(\eta) = \nabla \cdot \left( \frac{\nabla \eta}{\sqrt{1 + |\nabla \eta|^2}} \right) \tag{1-4}$$

is the mean curvature of the free surface. In (1-3), the kinetic condition at the free surface implies that fluid particles which are initially on the free surface will stay on the free surface, whereas the kinetic condition at the bottom is a rephrasing of the impenetrability of the bottom. The dynamic condition is the Laplace–Young equation which expresses the balance between the interior pressure  $P$  and the surface tension  $\kappa$ .

One of the main difficulties in the study of the Eulerian formulation of the system (1-3) is the time-dependence of the domain  $\Omega$ . By [Zakharov 1968; Craig and Sulem 1993], we can reformulate (1-3) as a system in  $\mathbb{R}^d$ . Note that due to the simply connected geometry of  $\Omega$  and the irrotationality of the fluid, there exists a velocity potential  $\phi : \Omega \rightarrow \mathbb{R}$  such that  $\nabla_{xy} \phi = v$ . By the incompressibility of the fluid, the potential  $\phi$  is harmonic. Therefore  $\phi$  satisfies the Laplace equation with Neumann boundary conditions:

$$\Delta_{xy} \phi = 0, \quad \partial_{\mathbf{n}} \phi|_{y=\eta} = \partial_t \eta / \langle \nabla \eta \rangle, \quad \partial_{\mathbf{n}} \phi|_{y=-b} = 0.$$

Define  $\psi = \phi|_{y=\eta}$  and define

$$G(\eta)\psi = \langle \nabla \eta \rangle \partial_{\mathbf{n}} \phi|_{y=\eta}.$$

Here  $G(\eta)$  is the Dirichlet–Neumann operator (see Section 5A for a rigorous definition). Then the system (1-3) can be rewritten in terms of the unknowns  $(\eta, \psi)$ :

$$\begin{cases} \partial_t \eta - G(\eta)\psi = 0, \\ \partial_t \psi + g\eta - \kappa H(\eta) + \frac{1}{2} |\nabla \psi|^2 - \frac{1}{2} \frac{(\nabla \eta \cdot \nabla \psi + G(\eta)\psi)^2}{1 + |\nabla \eta|^2} = 0. \end{cases} \tag{1-5}$$

We shall assume henceforth that  $\kappa = 1$  for simplicity.

**1C2. Quasihomogeneous wavefront set and model equations.** It is known that the linearization of (1-5) about the stationary solution  $(\eta, \psi) = (0, 0)$  can be symmetrized, up to a smoothing remainder, to the fraction Schrödinger equation or order  $\frac{3}{2}$ . Consider the more general model equation

$$\partial_t u + i|D_x|^\gamma u = 0, \quad \gamma \geq 1. \tag{1-6}$$

It is natural to ask ourselves if we can define a new family of wavefront sets and extend the results from Theorems 1.1 and 1.2 to (1-6). Note that a wave packet of (1-6) of frequency  $\xi \sim h^{-1}$  travels at the group velocity

$$v = \frac{d|\xi|^\gamma}{d\xi} = \gamma|\xi|^{\gamma-2}\xi \sim h^{-(\gamma-1)}.$$

It suggests that we need to use pseudodifferential operators of the form  $a(h^{\gamma-1}x, hD_x)$  as test operators. In the following definition, we consider the more general quantization with two parameters.

**Definition 1.3.** If  $u \in \mathcal{S}'(\mathbb{R}^d)$ ,  $\mu \in \mathbb{R} \cup \{\infty\}$ ,  $\delta \geq 0$  and  $\rho \geq 0$  with  $\delta + \rho > 0$ , then the quasihomogeneous wavefront set  $\text{WF}_{\delta,\rho}^\mu(u)$  is a subset of  $\mathbb{R}^{2d}$  defined as follows. A point  $(x_0, \xi_0)$  does not belong to  $\text{WF}_{\delta,\rho}^\mu(u)$  if and only if there exists  $a \in C_c^\infty(\mathbb{R}^{2d})$  with  $a(x_0, \xi_0) \neq 0$  such that  $\|a(h^\delta x, h^\rho D_x)u\|_{L^2} = \mathcal{O}(h^\mu)$  for  $h \in (0, 1]$ . Here,

$$a(h^\delta x, h^\rho D_x)u(x) = (2\pi)^{-d} \iint_{\mathbb{R}^{2d}} e^{i(x-y)\cdot\xi} a(h^\delta x, h^\rho \xi)u(y) dy d\xi. \tag{1-7}$$

Note that  $\text{WF}_{\delta,\rho}^\mu(u)$  is invariant under the scaling  $(x, \xi) \mapsto (\lambda^\delta x, \lambda^\rho \xi)$  for all  $\lambda > 0$ . The existence of  $(x_0, \xi_0) \in \text{WF}_{\delta,\rho}^\mu(u)$  implies an accumulation of mass near the ray  $\{(\lambda^\delta x_0, \lambda^\rho \xi_0)\}_{\lambda>0}$ . By choosing different parameters, we recover the definitions of various wavefront sets from the quasihomogeneous wavefront set: the wavefront set of Hörmander  $(\delta, \rho, \mu) = (0, 1, \infty)$ , the homogeneous wavefront set of Nakamura  $(\delta, \rho, \mu) = (1, 1, \infty)$  and the scattering wavefront set of [Melrose 1994]  $(\delta, \rho, \mu) = (1, 0, \infty)$ .

**Theorem 1.4.** If  $u$  solves (1-6) with initial data  $u(0) = u_0 \in L^2(\mathbb{R}^d)$  and  $\mu \in \mathbb{R} \cup \{\infty\}$ , then the following results of propagation hold:

(1) If  $\rho\gamma = \delta + \rho$ ,  $(x_0, \xi_0) \in \text{WF}_{\delta,\rho}^\mu(u_0) \setminus \{\xi = 0\}$  and  $t_0 \in \mathbb{R}$ , then

$$(x_0 + t_0\gamma|\xi_0|^{\gamma-2}\xi_0, \xi_0) \in \text{WF}_{\delta,\rho}^\mu(u(t_0)).$$

(2) If  $\gamma > 1$ ,  $\rho\gamma > \delta + \rho$ ,  $(x_0, \xi_0) \in \text{WF}_{\delta,\rho}^\mu(u_0) \setminus \{\xi = 0\}$  and  $t_0 \neq 0$ , then

$$(t_0\gamma|\xi_0|^{\gamma-2}\xi_0, \xi_0) \in \text{WF}_{\rho(\gamma-1),\rho}^\mu(u(t_0)).$$

Note that we do not require  $x_0 + t\gamma|\xi_0|^{\gamma-2}\xi_0 \neq 0$  in Theorem 1.4(1), while we require  $x_0 + t\xi_0 \neq 0$  in Theorem 1.2(1). This is because in Theorem 1.2 the geometry is only Euclidean at infinity.

**1C3. Asymptotically flat water waves.** Instead of the linearization at  $(\eta, \psi) = (0, 0)$ , if we parilinearize and symmetrize (1-5) as in [Alazard, Burq and Zuily 2011], then we obtain a quasilinear paradifferential fractional Schrödinger equation of order  $\frac{3}{2}$ . We require the geometry of the free surface to be Euclidean at infinity and the velocity field to be zero at infinity to avoid problems caused by the infinite speed of propagation and the nonlinearity. We shall fulfill this requirement by proving the existence of gravity-capillary water waves in some weighted Sobolev spaces.

**Definition 1.5.** If  $\mu, k \in \mathbb{R}$ , then  $H_k^\mu = H_k^\mu(\mathbb{R}^d)$  is the set of all  $u \in \mathcal{S}'(\mathbb{R}^d)$  such that

$$\|u\|_{H_k^\mu} = \|\langle x \rangle^k \langle D_x \rangle^\mu u\|_{L^2} < +\infty.$$

If in addition  $k \in \mathbb{N}$  and  $\delta \geq 0$ , then define

$$\mathcal{H}_k^{\mu, \delta} = \bigcap_{j=0}^k H_j^{\mu - \delta j}.$$

We are mostly interested in the case where  $\delta = \frac{1}{2}$ . The weighted Sobolev space  $\mathcal{H}_k^{\mu, 1/2}$  is a natural space to apply the energy estimate for the fractional Schrödinger equation of order  $\frac{3}{2}$  and thus also for the gravity-capillary water wave system.

**Theorem 1.6.** *If  $d \geq 1$ ,  $\mu > 3 + \frac{d}{2}$ ,  $k \leq 2\mu - d - 6$  and  $(\eta_0, \psi_0) \in \mathcal{H}_k^{\mu+1/2, 1/2} \times \mathcal{H}_k^{\mu, 1/2}$ , then there exist  $T > 0$  and a unique solution*

$$(\eta, \psi) \in C([-T, T], \mathcal{H}_k^{\mu+1/2, 1/2} \times \mathcal{H}_k^{\mu, 1/2})$$

to the Cauchy problem of (1-5) with initial data  $(\eta_0, \psi_0)$ .

The study of the Cauchy problem for the water wave equation dates back to [Nalimov 1974; Kano and Nishida 1979; Yosihara 1982; 1983]. The local well-posedness in Sobolev spaces with general initial data were achieved in [Wu 1997; 1999; Beyer and Günther 1998]. Our analysis of the water wave equation relies on the paradifferential calculus of [Bony 1986], which was introduced to the study of the water wave equation in [Alazard and Métivier 2009] and later allowed Alazard, Burq and Zuily [2011; 2014] to prove the local well-posedness with low Sobolev regularities. For recent progress of the Cauchy problem, see e.g., [Alazard and Delort 2015; de Poyferré and Nguyen 2016; 2017; Deng, Ionescu, Pausader and Pusateri 2017; Hunter, Ifrim and Tataru 2016; Ifrim and Tataru 2017; Ionescu and Pusateri 2018; Ming, Rousset and Tzvetkov 2015; Rousset and Tzvetkov 2011; Wang 2020].

To prove Theorem 1.6, we shall combine the analysis in [Alazard, Burq and Zuily 2011] and a paradifferential calculus in weighted Sobolev spaces. The latter can be achieved by modifying the definition of paradifferential operators via a spatial dyadic decomposition. More precisely, if  $a$  is a symbol, then we define

$$\mathcal{P}_a = \sum_{j \in \mathbb{N}} \underline{\psi}_j T_{\psi_j a} \underline{\psi}_j,$$

where  $\{\psi_j\}_{j \in \mathbb{N}} \subset C_c^\infty(\mathbb{R}^d)$  is a dyadic partition of unity of  $\mathbb{R}^d$ ,  $\underline{\psi}_j = \sum_{|k-j| \leq N} \psi_k$  for some sufficiently large  $N \in \mathbb{N}$ , and  $T_{\psi_j a}$  is the usual paradifferential operator of Bony. Such dyadic paradifferential calculus inherits the symbolic calculus and the parilinearization of Bony’s calculus, while at the same time allows the spatial polynomial growth/decay of symbols to play their roles in estimates.

We do not attempt to lower  $\mu$  to  $> 2 + \frac{d}{2}$  as it was in [Alazard, Burq and Zuily 2011]. The range of  $k$  is so chosen such that  $\mu - \frac{k}{2} > 3 + \frac{d}{2}$ , enabling us to parilinearize (1-5) in  $\mathcal{H}_k^\mu$ . We should mention that the existence of gravity water waves (water waves without surface tension) in uniformly local weighted Sobolev spaces was obtain by [Nguyen 2016] via a periodic spatial decomposition from [Alazard, Burq and Zuily 2016].

**1C4. Propagation at infinity.** Our first main result concerns the propagation of quasihomogeneous wavefront sets with parameters  $(\delta, \rho) = (\frac{1}{2}, 1)$ , corresponding to Theorem 1.4(1).



**Theorem 1.7.** *Suppose that  $d \geq 1$ ,  $\mu > 3 + \frac{d}{2}$ ,  $3 \leq k < 2\mu - K - d$  for some  $K > 0$ , and*

$$(\eta, \psi) \in C([-T, T], \mathcal{H}_k^{\mu+1/2, 1/2} \times \mathcal{H}_k^{\mu, 1/2}),$$

where  $T > 0$ , solves (1-5). If  $t_0 \in [-T, T]$  and

$$(x_0, \xi_0) \in \text{WF}_{1/2,1}^{\mu+1/2+\sigma}(\eta(0)) \cup \text{WF}_{1/2,1}^{\mu+\sigma}(\psi(0))$$

such that  $\xi_0 \neq 0$ ,  $0 \leq \sigma \leq \frac{k}{2} - \frac{3}{2}$  and

$$x_0 + \frac{3}{2}t|\xi_0|^{-1/2}\xi_0 \neq 0$$

for all  $t$  between 0 and  $t_0$ , then

$$(x_0 + \frac{3}{2}t_0|\xi_0|^{-1/2}\xi_0, \xi_0) \in \text{WF}_{1/2,1}^{\mu+1/2+\sigma}(\eta(t_0)) \cup \text{WF}_{1/2,1}^{\mu+\sigma}(\psi(t_0)).$$

We will see that, by Lemma 2.15, if  $(\eta, \psi) \in \mathcal{H}_k^{\mu+1/2, 1/2} \times \mathcal{H}_k^{\mu, 1/2}$ , then

$$\text{WF}_{1/2,1}^{\mu+1/2}(\eta) \cup \text{WF}_{1/2,1}^{\mu}(\psi) \subset \{x = 0\} \cup \{\xi = 0\}.$$

By [Alazard and Métivier 2009], we expect  $\sigma$  to be at most  $\mu - \alpha - \frac{d}{2}$  for some  $\alpha > 0$ , corresponding to the gain of regularity by the remainder in the parilinearization procedure. Theorem 1.7 does not give the optimal upper bound for  $\sigma$ , as it is not our priority, but when  $k = 2\mu - K - d$ , the parameter  $\sigma$  can still be as large as  $\mu - \frac{K}{2} - \frac{d}{2} - \frac{3}{2}$ , almost reaching the paradifferential threshold.

**1C5. Microlocal smoothing effect.** Our second main result shows that singularities of the initial data which are nontrapped with respect to the initial geometry instantaneously generate an element in the quasihomogeneous wavefront set with parameters  $(\delta, \rho) = (\frac{1}{2}, 1)$ , corresponding to Theorem 1.4(2).

Observe that if  $\eta$  is sufficiently regular, then  $\Sigma$  endowed with the metric inherited from  $\mathbb{R}^{d+1}$  is isometric to  $(\mathbb{R}^d, \varrho)$ , where

$$\varrho = \begin{pmatrix} \text{Id} + (\nabla\eta)'(\nabla\eta) & \nabla\eta \\ '(\nabla\eta) & 1 \end{pmatrix}.$$

Define  $\Sigma_0 = \Sigma|_{t=0}$  and  $\varrho_0 = \varrho|_{t=0}$ . We identify the cogeodesic flow  $\mathcal{G}$  on  $T^*\Sigma_0$  with the Hamiltonian flow on  $\mathbb{R}^{2d}$  of the symbol  $G(x, \xi) = {}^t\xi\varrho_0(x)^{-1}\xi$ . Precisely  $\mathcal{G} = \mathcal{G}_s(x, \xi)$  is defined by the equation

$$\partial_s \mathcal{G}_s = (\partial_\xi G, -\partial_x G)(\mathcal{G}_s), \quad \mathcal{G}_0 = \text{Id}_{\mathbb{R}^{2d}}. \tag{1-8}$$

**Definition 1.8.** A point  $(x_0, \xi_0) \in \mathbb{R}^d \times (\mathbb{R}^d \setminus \{0\})$  is called forwardly (resp. backwardly) nontrapped with respect to  $\mathcal{G}$  if, with an abuse of notation, the cogeodesic  $\{(x_s, \xi_s) = \mathcal{G}_s(x_0, \xi_0)\}_{s \in \mathbb{R}}$  satisfies

$$\lim_{s \rightarrow +\infty} |x_s| = \infty \quad (\text{resp.} \quad \lim_{s \rightarrow -\infty} |x_s| = \infty).$$

**Theorem 1.9.** *If  $d \geq 1$ ,  $\mu > 3 + \frac{d}{2}$ ,  $3 \leq k < \frac{2}{3}(\mu - 1 - \frac{d}{2})$ , and*

$$(\eta, \psi) \in C([-T, T], \mathcal{H}_k^{\mu+1/2, 1/2} \times \mathcal{H}_k^{\mu, 1/2}),$$

where  $T > 0$ , solves (1-5). Let

$$(x_0, \xi_0) \in \text{WF}_{0,1}^{\mu+1/2+\sigma}(\eta(0)) \cup \text{WF}_{0,1}^{\mu+\sigma}(\psi(0)),$$

where  $\xi_0 \neq 0$  and  $0 \leq \sigma \leq \frac{3}{2}k$ . If  $(x_0, \xi_0)$  is forwardly (resp. backwardly) nontrapped, and the cogeodesic  $\{(x_s, \xi_s)\}_{s \in \mathbb{R}}$  is defined as above, then there exists  $\xi_{+\infty}$  (resp.  $\xi_{-\infty}$ ) in  $\mathbb{R}^d \setminus \{0\}$  such that

$$\lim_{s \rightarrow \infty} \xi_s = \xi_{+\infty} \quad (\text{resp. } \lim_{s \rightarrow \infty} \xi_{-s} = \xi_{-\infty}),$$

and moreover, for all  $0 < t_0 \leq T$  (resp.  $-T \leq t_0 < 0$ ), we have

$$\begin{aligned} & \left(\frac{3}{2}t_0|\xi_{+\infty}|^{-1/2}\xi_{+\infty}, \xi_{+\infty}\right) \in \text{WF}_{1/2,1}^{\mu+1/2+\sigma}(\eta(t_0)) \cup \text{WF}_{1/2,1}^{\mu+\sigma}(\psi(t_0)), \\ & (\text{resp. } \left(\frac{3}{2}t_0|\xi_{-\infty}|^{-1/2}\xi_{-\infty}, \xi_{-\infty}\right) \in \text{WF}_{1/2,1}^{\mu+1/2+\sigma}(\eta(t_0)) \cup \text{WF}_{1/2,1}^{\mu+\sigma}(\psi(t_0))). \end{aligned}$$

We remark that the asymptotic directions  $\xi_{\pm\infty}$  are determined solely by the geometry of  $\Sigma_0$ . This is due to the infinite speed of propagation. We can also prove that the nontrapping assumption is, at least in the following two cases, unnecessary: if  $d = 1$ , or if  $\nabla\eta(0) \in L^\infty$  and  $\|\langle x \rangle \nabla^2\eta(0)\|_{L^\infty}$  is sufficiently small. In both cases we obtain the following local smoothing effect.

**Corollary 1.10.** *Suppose  $d, \mu, k, \sigma$  satisfy the hypothesis of the previous theorem,  $T > 0$ ,*

$$(\eta, \psi) \in C([-T, T], \mathcal{H}_k^{\mu+1/2, 1/2} \times \mathcal{H}_k^{\mu, 1/2})$$

*solves (1-5), and both of the following two conditions are satisfied:*

- (1) *Either  $d = 1$  or  $\|\langle x \rangle \nabla^2\eta(0)\|_{L^\infty}$  is sufficiently small.*
- (2)  $\text{WF}_{1/2,1}^{\mu+1/2+\sigma}(\eta(0)) \cup \text{WF}_{1/2,1}^{\mu+\sigma}(\psi(0)) \subset \{x = 0\} \cup \{\xi = 0\}$ .

*Then, for all  $t_0 \in [-T, T] \setminus \{0\}$  and for all  $\epsilon > 0$ ,*

$$(\eta(t_0), \psi(t_0)) \in H_{\text{loc}}^{\mu+1/2+\sigma-\epsilon} \times H_{\text{loc}}^{\mu+\sigma-\epsilon}.$$

The second condition is satisfied if, by Lemma 2.15, there exists  $(k, k') \in \mathbb{R}^2$  such that

$$(\eta(0), \psi(0)) \in H_{2k}^{\mu+1/2+\sigma-k} \times H_{2k'}^{\mu+\sigma-k'}.$$

This is particularly the case if  $(\eta(0), \psi(0)) \in \mathcal{E}'(\mathbb{R}^d) \times \mathcal{E}'(\mathbb{R}^d)$ .

We refer to [Christianson, Hur and Staffilani 2009; Alazard, Burq and Zuily 2011] for local smoothing effects of 2-dimensional capillary-gravity water waves. See also [Alazard, Ifrim and Tataru 2022] for a Morawetz inequality of 2-dimensional gravity water waves.

**1D. Outline of paper.** In Section 2, we present basic properties of weighted Sobolev spaces and the quasihomogeneous wavefront set. In Section 3, we prove Theorem 1.4 by extending the idea of Nakamura. In Section 4, we review the paradifferential calculus of Bony, and extend it to weighted Sobolev spaces by a spatial dyadic decomposition. We also develop a quasihomogeneous semiclassical paradifferential calculus, and study its relations with the quasihomogeneous wavefront set. In Section 5, we study the Dirichlet–Neumann operator in weighted Sobolev spaces and prove the existence of asymptotically flat gravity-capillary water waves, i.e., Theorem 1.6. In Section 6, we prove our main results, i.e., Theorem 1.7, Theorem 1.9 and Corollary 1.10, by extending the proof of Theorem 1.4 to the quasilinear equation using the paradifferential calculus.

## 2. Quasihomogeneous microlocal analysis

In this section we develop the quasihomogeneous semiclassical calculus and discuss its relation with weighted Sobolev spaces and the quasihomogeneous wavefront set.

### 2A. Quasihomogeneous semiclassical calculus.

**Definition 2.1.** For  $(\mu, k) \in \mathbb{R}^2$ , set  $m_k^\mu(x, \xi) = \langle x \rangle^k \langle \xi \rangle^\mu$ . Let  $a_h \in C^\infty(\mathbb{R}^{2d})$ . We say that  $a_h \in S_k^\mu$  if for all  $\alpha, \beta \in \mathbb{N}^d$ , there exists  $C_{\alpha\beta} > 0$ , such that, for all  $(x, \xi) \in \mathbb{R}^{2d}$ ,

$$\sup_{h \in (0,1]} |\partial_x^\alpha \partial_\xi^\beta a_h(x, \xi)| \leq C_{\alpha\beta} m_k^{\mu-|\beta|}(x, \xi). \tag{2-1}$$

We say that  $a_h \in S_k^\mu$  is  $(\mu, k)$ -elliptic if there exist  $R > 0, C > 0$  such that, for  $|x| + |\xi| \geq R$ ,

$$\inf_{h \in (0,1]} |a_h(x, \xi)| \geq C m_k^\mu(x, \xi).$$

Also write  $S_\infty^\infty = \bigcup_{(\mu,k) \in \mathbb{R}^2} S_k^\mu$ , and  $S_{-\infty}^{-\infty} = \bigcap_{(\mu,k) \in \mathbb{R}^2} S_k^\mu$ .

We say that  $a_h \in S_{-\infty}^{-\infty}$  is elliptic at  $(x_0, \xi_0)$  if, for some neighborhood  $\Omega$  of  $(x_0, \xi_0)$ ,

$$\inf_{h \in (0,1]} \inf_{(x,\xi) \in \Omega} |a_h(x, \xi)| > 0.$$

**Definition 2.2.** Let  $\delta, \rho \in \mathbb{R}$  such that  $\delta + \rho > 0$  and, for all  $h \in (0, 1]$ , define the scaling

$$\theta_h^{\delta,\rho} : (x, \xi) \mapsto (h^\delta x, h^\rho \xi), \tag{2-2}$$

which induces a pullback  $\theta_{h,*}^{\delta,\rho}$  on  $S_\infty^\infty$ :  $\theta_{h,*}^{\delta,\rho} a_h = a_h \circ \theta_h^{\delta,\rho}$ . Then define, by (1-7),

$$\text{Op}_h^{\delta,\rho}(a_h) = \text{Op}(\theta_{h,*}^{\delta,\rho} a_h) = a(h^\delta x, h^\rho D_x).$$

The scaling  $\vartheta_h^\delta u(x) = h^{\delta d/2} u(h^\delta x)$  defines an isometry on  $L^2(\mathbb{R}^d)$ . Therefore, by the formula

$$(\vartheta_h^\delta)^{-1} \text{Op}_h^{\delta,\rho}(a) \vartheta_h^\delta = \text{Op}_h^{0,\delta+\rho}(a), \tag{2-3}$$

we deduce the following results from the usual semiclassical calculus, for which we refer to [Zworski 2012].

**Proposition 2.3.** *There exists  $K > 0$  such that, if  $a \in C^\infty(\mathbb{R}^{2d})$  with  $\|\partial_x^\alpha \partial_\xi^\beta a\|_{L^\infty} \leq M$  for all  $|\alpha| + |\beta| \leq d$ , then  $\text{Op}_h^{\delta,\rho}(a) : L^2 \rightarrow L^2$  and  $\|\text{Op}_h^{\delta,\rho}(a)\|_{L^2 \rightarrow L^2} \leq KM$ .*

**Proposition 2.4.** *There exists a bilinear operator  $\sharp_h^{\delta,\rho} : S_\infty^\infty \times S_\infty^\infty \rightarrow S_\infty^\infty$  such that*

$$\text{Op}_h^{\delta,\rho}(a_h) \text{Op}_h^{\delta,\rho}(b_h) = \text{Op}_h^{\delta,\rho}(a_h \sharp_h^{\delta,\rho} b_h).$$

Moreover, if  $a_h \in S_k^\mu$  and  $b_h \in S_\ell^\nu$ , then  $a_h \sharp_h^{\delta,\rho} b_h \in S_{k+\ell}^{\mu+\nu}$ . For all  $r > 0$ , define

$$a_h \sharp_{h,r}^{\delta,\rho} b_h = \sum_{|\alpha| < r} \frac{h^{|\alpha|(\delta+\rho)}}{\alpha!} \partial_\xi^\alpha a_h D_x^\alpha b_h. \tag{2-4}$$

Then we have

$$a_h \sharp_h^{\delta,\rho} b_h - a_h \sharp_{h,r}^{\delta,\rho} b_h = \mathcal{O}(h^{r(\delta+\rho)})_{S_{k+\ell-r}^{\mu+\nu-r}}.$$

**Proposition 2.5.** *There exists a linear operator  $\zeta_h^{\delta,\rho} : S_\infty^\infty \rightarrow S_\infty^\infty$  such that*

$$\text{Op}_h^{\delta,\rho}(a_h)^* = \text{Op}_h^{\delta,\rho}(\zeta_h^{\delta,\rho} a_h).$$

Moreover if  $a_h \in S_k^\mu$ , then  $\zeta_h^{\delta,\rho} a_h \in S_k^\mu$ . For  $r > 0$ , define

$$\zeta_{h,r}^{\delta,\rho} a_h = \sum_{|\alpha| < r} \frac{h^{|\alpha|(\delta+\rho)}}{\alpha!} \partial_\xi^\alpha D_x^\alpha \bar{a}_h. \tag{2-5}$$

Then we have

$$\zeta_h^{\delta,\rho} a_h - \zeta_{h,r}^{\delta,\rho} a_h = \mathcal{O}(h^{r(\delta+\rho)})_{S_{k-r}^{\mu-r}}.$$

**Proposition 2.6** (sharp Gårding inequality). *If  $\delta + \rho > 0$  and  $a_h \in S_0^0$  such that  $\text{Re } a_h \geq 0$ , then there exists  $C > 0$  such that, for all  $u \in L^2(\mathbb{R}^d)$  and  $0 < h < 1$ , we have*

$$\text{Re}(\text{Op}_h^{\delta,\rho}(a_h)u, u)_{L^2} \geq -Ch^{\delta+\rho} \|u\|_{L^2}^2.$$

**2B. Weighted Sobolev spaces.** Recall the weighted Sobolev spaces defined in [Definition 1.5](#).

**Proposition 2.7.** *We have  $\mathcal{S}(\mathbb{R}^d) = \bigcap_{\mu,k \in \mathbb{R}} H_k^\mu$  and  $\mathcal{S}'(\mathbb{R}^d) = \bigcup_{\mu,k \in \mathbb{R}} H_k^\mu$ .*

*Proof.* Clearly  $\mathcal{S}(\mathbb{R}^d) \subset \bigcap_{\mu,k \in \mathbb{R}} H_k^\mu$ . The converse follows by the Sobolev embedding theorems. As for the second statement, clearly  $\bigcup_{(\mu,k) \in \mathbb{R}^2} H_k^\mu \subset \mathcal{S}'(\mathbb{R}^d)$ . Conversely, if  $u \in \mathcal{S}'(\mathbb{R}^d)$ , then there exists  $N > 0$ , such that for all  $\varphi \in \mathcal{S}(\mathbb{R}^d)$  we have

$$\langle u, \varphi \rangle_{\mathcal{S}', \mathcal{S}} \lesssim \sum_{|\alpha|+|\beta| \leq N} \|x^\alpha \partial_x^\beta \varphi\|_{L^\infty} \lesssim \|\text{Op}(m_N^N) \varphi\|_{L^2}.$$

By duality this implies that  $u \in H_{-N}^{-N}$ . □

**Lemma 2.8.** *If  $u \in \mathcal{S}'(\mathbb{R}^d)$ , then there exists  $N > 0$  such that*

$$u = h^{-N} \text{Op}_h^{\delta,\rho}(m_{-N}^{-N}) \mathcal{O}(1)_{L^2}.$$

Therefore, if  $\delta + \rho > 0$ , and  $a_h \in \mathcal{O}(h^\infty)_{S_{-\infty}^\infty}$ , then  $\text{Op}_h^{\delta,\rho}(a_h)u_h = \mathcal{O}(h^\infty)_{\mathcal{S}}$ .

*Proof.* By the proof of [Proposition 2.7](#), there exists  $N > 0$  such that, for all  $\varphi \in \mathcal{S}(\mathbb{R}^d)$ ,

$$\langle u, \varphi \rangle_{\mathcal{S}', \mathcal{S}} \lesssim \sum_{|\alpha|+|\beta| \leq N} \|x^\alpha \partial_x^\beta \varphi\|_{L^\infty} \lesssim h^{-N} \|\text{Op}_h^{\delta,\rho}(m_N^N) \varphi\|_{L^2}.$$

Again we conclude by duality. □

**Definition 2.9.** We say that a linear operator  $\mathcal{A} : \mathcal{S}(\mathbb{R}^d) \rightarrow \mathcal{S}'(\mathbb{R}^d)$  is of order  $(\nu, \ell) \in \mathbb{R}^2$ , and write  $\mathcal{A} \in \mathcal{O}_\ell^\nu$  if for all  $(\mu, k) \in \mathbb{R}^2$  there exists  $C > 0$  such that for all  $u \in \mathcal{S}(\mathbb{R}^d)$  we have

$$\|\mathcal{A}u\|_{H_{k-\ell}^{\mu-\nu}} \leq C \|u\|_{H_k^\mu}.$$

Therefore  $\mathcal{A}$  extends to a bounded linear operator from  $H_k^\mu$  to  $H_{k-\ell}^{\mu-\nu}$ . We write  $\mathcal{A} \in \mathcal{O}_{-\infty}^\nu$  if  $\mathcal{A} \in \mathcal{O}_\ell^\nu$  for all  $(\nu, \ell) \in \mathbb{R}^2$ .

Let  $\mathcal{A}$  be any nonempty set. Let  $\mathcal{A}_\alpha : \mathcal{S}(\mathbb{R}^d) \rightarrow \mathcal{S}'(\mathbb{R}^d)$  and  $C_\alpha > 0$  be indexed by  $\alpha \in \mathcal{A}$ . We say  $\mathcal{A}_\alpha = \mathcal{O}(C_\alpha)_{\mathcal{O}_\ell^v}$  if for all  $(\mu, k) \in \mathbb{R}^2$  there exists  $K > 0$  such that, for all  $\alpha \in \mathcal{A}$ , we have

$$\|\mathcal{A}_\alpha\|_{H_k^\mu \rightarrow H_{k-\ell}^{\mu-v}} \leq KC_\alpha.$$

By Propositions 2.3 and 2.4, we obtain:

**Proposition 2.10.** *The following mapping properties of pseudodifferential operators hold:*

- (1) If  $a_h \in S_\ell^v$  with  $(v, \ell) \in \mathbb{R}^2$ , then  $\text{Op}(a_h) \in \mathcal{O}_\ell^v$ .
- (2) If  $u \in \mathcal{S}'(\mathbb{R}^d)$ , then  $u \in H_k^\mu$  if and only if there exists a  $(\mu, k)$ -elliptic symbol  $a_h \in S_k^\mu$  such that  $\text{Op}(a_h)u = \mathcal{O}(1)_{L^2}$ .

Next, we characterize weighted Sobolev spaces by a dyadic decomposition.

**Definition 2.11.** The set  $\mathcal{P}$  consists of all maps of the form

$$\psi : \mathbb{N} \rightarrow C_c^\infty(\mathbb{R}^d), \quad j \mapsto \psi_j,$$

such that the following conditions are satisfied:

- (1) There exists  $C > 1$  such that for all  $j \geq 1$  we have

$$\text{supp } \psi_j \subset \{x \in \mathbb{R}^d : C^{-1}2^j \leq |x| \leq C2^j\}.$$

- (2) For all  $j \geq 0$ , the function  $\psi_j$  is nonnegative.
- (3) There exists  $C > 1$  such that  $C^{-1} \leq \sum_{j \in \mathbb{N}} \psi_j \leq C$ .
- (4) For all  $\alpha \in \mathbb{N}$  there exists  $C_\alpha$  such that for all  $j \in \mathbb{N}$  we have

$$\|\partial_x^\alpha \psi_j\|_{L^\infty} \leq C_\alpha 2^{-j|\alpha|}.$$

The set  $\mathcal{P}_*$  consists of all  $\psi \in \mathcal{P}$  such that

- (5)  $\sum_{j \in \mathbb{N}} \psi_j = 1$ , and
- (6)  $\text{supp } \psi_j \cap \text{supp } \psi_k = \emptyset$  whenever  $|j - k| > 2$ .

If  $\psi, \tilde{\psi} \in \mathcal{P}$  such that  $\psi_j \tilde{\psi}_j = \psi_j$  for all  $j \in \mathbb{N}$ , then we write  $\psi \Subset \tilde{\psi}$ .

**Proposition 2.12.** *If  $\mu, k \in \mathbb{R}$ ,  $\psi \in \mathcal{P}$  and  $u \in \mathcal{S}'(\mathbb{R}^d)$ , then  $u \in H_k^\mu$  if and only if*

$$\sum_{j \in \mathbb{N}} 2^{2jk} \|\psi_j u\|_{H^\mu}^2 < \infty.$$

Moreover, there exists  $C > 1$  such that, for all  $u \in H_k^\mu$ , we have

$$C^{-1} \|u\|_{H_k^\mu}^2 \leq \sum_{j \in \mathbb{N}} 2^{2jk} \|\psi_j u\|_{H^\mu}^2 \leq C \|u\|_{H_k^\mu}^2.$$

*Proof.* We may assume that  $\psi \in \mathcal{P}_*$  because if  $\phi^1, \phi^2 \in \mathcal{P}$  then

$$\sum_{j \in \mathbb{N}} 2^{2jk} \|\phi_j^1 u\|_{H^\mu}^2 \simeq \sum_{j \in \mathbb{N}} 2^{2jk} \|\phi_j^2 u\|_{H^\mu}^2.$$

Define  $\tilde{\psi} \in \mathcal{D}$  by setting  $\tilde{\psi}_j = \sum_{|k-j| \leq 2} \psi_k$  for all  $j \in \mathbb{N}$ . Then  $\psi \in \tilde{\psi}$ . Note that the family of multiplication operators  $\{2^{-jk} \langle x \rangle^k \tilde{\psi}_j\}_{j \in \mathbb{N}}$  is bounded in  $\mathcal{O}_0^0$ , which implies that, for all  $\mu \in \mathbb{N}$ , the family of pseudodifferential operators  $\{2^{-jk} \langle D_x \rangle^\mu \langle x \rangle^k \tilde{\psi}_j \langle D_x \rangle^{-\mu}\}_{j \in \mathbb{N}}$  is bounded in  $\mathcal{O}_0^0$ . Therefore, for all  $u \in H_k^\mu$ , we have

$$2^{2jk} \|\psi_j u\|_{H^\mu}^2 \lesssim \|\langle D_x \rangle^\mu \psi_j \langle x \rangle^k u\|_{L^2}^2 \lesssim \|\tilde{\psi}_j \langle D_x \rangle^\mu \psi_j \langle x \rangle^k u\|_{L^2}^2 + \|(1 - \tilde{\psi}_j) \langle D_x \rangle^\mu \psi_j \langle x \rangle^k u\|_{L^2}^2. \quad (2-6)$$

Apply Proposition 2.4 with  $(\delta, \rho) = (1, 0)$  and  $h = 2^{-j}$ , we obtain that, for all  $N > 0$ , the estimate

$$(1 - \tilde{\psi}_j) \langle D_x \rangle^\mu \psi_j \langle D_x \rangle^{-\mu} = \mathcal{O}(2^{-jN})_{L^2 \rightarrow L^2} \quad (2-7)$$

holds uniformly for all  $j \in \mathbb{N}$ . Therefore,

$$\sum_{j \in \mathbb{N}} \|(1 - \tilde{\psi}_j) \langle D_x \rangle^\mu \psi_j \langle x \rangle^k u\|_{L^2}^2 \lesssim \sum_{j \in \mathbb{N}} 2^{-2jN} \|u\|_{H_k^\mu}^2 \lesssim \|u\|_{H_k^\mu}^2.$$

For  $r \in \{0, 1, \dots, 9\}$ , set

$$a_r = \sum_{j \in 10\mathbb{N}+r} \tilde{\psi}_j \langle \xi \rangle^\mu \sharp(\langle x \rangle^k \psi_j) \in S_k^\mu,$$

where  $\sharp = \sharp_1^{0,0}$ . Observe that if  $0 \neq j - j' \in 10\mathbb{N}$ , then  $\text{supp } \tilde{\psi}_j \cap \text{supp } \tilde{\psi}_{j'} = \emptyset$ . Therefore,

$$\sum_{j \in \mathbb{N}} \|\tilde{\psi}_j \langle D_x \rangle^\mu \psi_j \langle x \rangle^k u\|_{L^2}^2 = \sum_{r=0}^9 \sum_{j \in 10\mathbb{N}+r} \|\tilde{\psi}_j \langle D_x \rangle^\mu \psi_j \langle x \rangle^k u\|_{L^2}^2 = \sum_{r=0}^9 \|\text{Op}(a_r)u\|_{L^2}^2 \lesssim \|u\|_{H_k^\mu}^2. \quad (2-8)$$

Combining (2-6), (2-7) and (2-8), we prove that if  $u \in H_k^\mu$  then  $\sum_{j \in \mathbb{N}} 2^{2jk} \|\psi_j u\|_{H^\mu}^2 \lesssim \|u\|_{H_k^\mu}^2$ .

Conversely, assume that  $\sum_{j \in \mathbb{N}} 2^{2jk} \|\psi_j u\|_{H^\mu}^2 < \infty$ . Much as above, we have

$$\begin{aligned} \infty > \sum_{j \in \mathbb{N}} 2^{2jk} \|\langle D_x \rangle^\mu \psi_j u\|_{L^2}^2 &\gtrsim \sum_{r=0}^9 \sum_{j \in 10\mathbb{N}+r} \|\tilde{\psi}_j \langle D_x \rangle^\mu \psi_j \langle x \rangle^k u\|_{L^2}^2 \\ &\gtrsim \sum_{r=0}^9 \|\text{Op}(a_r)u\|_{L^2}^2 \gtrsim \|\text{Op}(a)u\|_{L^2}^2, \end{aligned} \quad (2-9)$$

where  $a = \sum_{r=0}^9 a_r$ . Observe  $a$  is  $(\mu, k)$ -elliptic, so  $u \in H_k^\mu$ . By the symbolic calculus, there exists  $r \in S_{-\infty}^-$  such that

$$\|u\|_{H_k^\mu}^2 \lesssim \|\text{Op}(a)u\|_{L^2}^2 + \|\text{Op}(r)u\|_{L^2}^2. \quad (2-10)$$

For the remainder term, we have

$$\|\text{Op}(r)u\|_{L^2}^2 = (u, \text{Op}(r^* \sharp r)u)_{L^2} = \sum_{j \in \mathbb{N}} (u, \text{Op}(r^* \sharp r) \psi_j u)_{L^2}.$$

For each term in the summation, by the analysis above (2-6), we have, for all  $N > 0$  and  $\varepsilon > 0$ ,

$$\begin{aligned} (u, \text{Op}(r^* \sharp r) \psi_j u)_{L^2} &= (\text{Op}(m_k^\mu)u, \text{Op}(m_{-k}^{-\mu} \sharp r^* \sharp r \sharp m_{N-k}^{-\mu}) \langle D_x \rangle^\mu \langle x \rangle^{-N+k} \psi_j u)_{L^2} \\ &\lesssim \|u\|_{H_k^\mu} \|\langle D_x \rangle^\mu \langle x \rangle^{-N+k} \psi_j u\|_{L^2} \\ &\lesssim 2^{-jN} \|u\|_{H_k^\mu} 2^{jk} \|\langle D_x \rangle^\mu \psi_j u\|_{L^2} \\ &\lesssim 2^{-jN} (\varepsilon \|u\|_{H_k^\mu}^2 + \varepsilon^{-1} 2^{2jk} \|\langle D_x \rangle^\mu \psi_j u\|_{L^2}^2), \end{aligned}$$

where the constants are independent of  $\varepsilon$ . Summing up in  $j$ ,

$$\|\text{Op}(r)u\|_{L^2}^2 \lesssim \varepsilon \|u\|_{H_k^\mu}^2 + \varepsilon^{-1} \sum_{j \in \mathbb{N}} 2^{2jk} \|\psi_j u\|_{H^\mu}^2. \tag{2-11}$$

By choosing  $\varepsilon$  sufficiently small, we conclude by (2-9), (2-10) and (2-11) that

$$\|u\|_{H_k^\mu}^2 \lesssim \sum_{j \in \mathbb{N}} 2^{2jk} \|\psi_j u\|_{H^\mu}^2$$

and finishes the proof. □

**2C. The quasihomogeneous wavefront set.** In this section the parameters  $\delta, \rho, \mu$  satisfy the conditions in Definition 1.3 without further specification. By Lemma 2.8, the following characterization of the quasihomogeneous wavefront set is easy to prove by the symbolic calculus.

**Proposition 2.13.** *If  $u \in \mathcal{S}'(\mathbb{R}^d)$ , then  $(x_0, \xi_0) \notin \text{WF}_{\delta, \rho}^\mu(u)$  if and only if there exists  $a_h \in S_{-\infty}^-$  which is elliptic at  $(x_0, \xi_0)$  such that  $\text{Op}_h^{\delta, \rho}(a_h)u = \mathcal{O}(h^\mu)_{L^2}$  for  $h \in (0, 1]$ .*

**Lemma 2.14.** *If  $u \in \mathcal{S}'(\mathbb{R}^d)$  and  $a_h \in S_{-\infty}^-$  is such that*

$$\overline{\bigcup_{h \in (0, 1]} \text{supp } a_h} \in \mathbb{R}^{2d} \setminus \text{WF}_{\delta, \rho}^\mu(u),$$

then  $\langle u, \text{Op}_h^{\delta, \rho}(a_h)u \rangle_{\mathcal{S}', \mathcal{S}} = \mathcal{O}(h^{2\mu})$  and consequently  $\text{Op}_h^{\delta, \rho}(a_h)u = \mathcal{O}(h^\mu)_{L^2}$  for  $h \in (0, 1]$ .

*Proof.* Let  $K = \overline{\bigcup_{h \in (0, 1]} \text{supp } a_h}$  and let  $\{\Omega_i\}_{i \in I}$  be an open cover of  $K$ . Let  $b_h^i \in S_{-\infty}^-$  be elliptic everywhere in  $\Omega_i$  such that  $\text{Op}_h^{\delta, \rho}(b_h^i)u = \mathcal{O}(h^\mu)_{L^2}$ . By a partition of unity, we may assume that  $K \subset \Omega := \Omega_{i_0}$  for some  $i_0 \in I$ , and set  $b_h = b_h^{i_0}$ . By the ellipticity of  $b_h$ , we can find  $c_h \in S_{-\infty}^-$  and  $r_h = \mathcal{O}(h^N)_{S_{-\infty}^-}$  for some large  $N > 0$  such that  $a_h = (\zeta_h^{\delta, \rho} b_h) \sharp_h^{\delta, \rho} c_h \sharp_h^{\delta, \rho} b_h + r_h$ . Therefore, by Lemma 2.8,

$$\begin{aligned} \langle u, \text{Op}_h^{\delta, \rho}(a_h)u \rangle_{\mathcal{S}', \mathcal{S}} &= (\text{Op}_h^{\delta, \rho}(b_h)u, \text{Op}_h^{\delta, \rho}(c_h) \text{Op}_h^{\delta, \rho}(b_h)u)_{L^2} + \langle u, \text{Op}_h^{\delta, \rho}(r_h)u \rangle_{\mathcal{S}', \mathcal{S}} \\ &= \mathcal{O}(h^\mu)^2 + \mathcal{O}(h^\infty) = \mathcal{O}(h^{2\mu}). \end{aligned}$$

Next, observe that there exists  $w_h \in S_{-\infty}^-$  and  $\tilde{r}_h = \mathcal{O}(h^N)_{S_{-\infty}^-}$  such that  $\text{supp } w_h \subset K$  and

$$\text{Op}_h^{\delta, \rho}(a_h)^* \text{Op}_h^{\delta, \rho}(a_h) = \text{Op}_h^{\delta, \rho}(w_h) + \text{Op}_h^{\delta, \rho}(\tilde{r}_h).$$

Therefore,

$$\begin{aligned} \|\text{Op}_h^{\delta, \rho}(a_h)u\|_{L^2}^2 &= \langle u, \text{Op}_h^{\delta, \rho}(a_h)^* \text{Op}_h^{\delta, \rho}(a_h)u \rangle_{\mathcal{S}', \mathcal{S}} \\ &= \langle u, \text{Op}_h^{\delta, \rho}(w_h)u \rangle_{\mathcal{S}', \mathcal{S}} + \mathcal{O}(h^{2\mu}) = \mathcal{O}(h^{2\mu}). \end{aligned} \quad \square$$

**Lemma 2.15.** *If  $u \in \mathcal{S}'(\mathbb{R}^d)$ . Then the following statements hold:*

- (1) *The quasihomogeneous wavefront set  $\text{WF}_{\delta, \rho}^\mu(u)$  is a closed  $(\delta, \rho)$ -cone. To be precise, this means  $\theta_h^{\delta, \rho} \text{WF}_{\delta, \rho}^\mu(u) = \text{WF}_{\delta, \rho}^\mu(u)$  for all  $h > 0$  where the scaling  $\theta_h^{\delta, \rho}$  is defined by (2-2).*
- (2) *If  $\gamma > 0$  then  $\text{WF}_{\delta, \rho}^\mu(u) = \text{WF}_{\delta/\gamma, \rho/\gamma}^{\mu/\gamma}(u)$ . Therefore in all situations we can restrict our discussions to the cases where either  $\delta = 1$  or  $\rho = 1$ .*
- (3) *For all  $(x_0, \xi_0) \in \mathbb{R}^{2d}$ , we have  $(x_0, \xi_0) \in \text{WF}_{\delta, \rho}^\mu(u)$  if and only if  $(\xi_0, -x_0) \in \text{WF}_{\rho, \delta}^\mu(\hat{u})$ .*

- (4) For all  $(x_0, \xi_0) \in \mathbb{R}^{2d}$ , we have  $(x_0, \xi_0) \in \text{WF}_{\delta, \rho}^\mu(u)$  if and only if  $(x_0, -\xi_0) \in \text{WF}_{\delta, \rho}^\mu(\bar{u})$ .
- (5) Define  $\text{WF}_{\delta, \rho}^\mu(u)^\circ = \text{WF}_{\delta, \rho}^\mu(u) \setminus \mathcal{N}_{\delta, \rho}$ , where

$$\mathcal{N}_{\delta, \rho} = \begin{cases} \{x = 0\} \times \mathbb{R}^d, & \delta > 0, \rho = 0, \\ \mathbb{R}^d \times \{\xi = 0\}, & \delta = 0, \rho > 0, \\ \{x = 0\} \times \mathbb{R}^d \cup \mathbb{R}^d \times \{\xi = 0\}, & \delta > 0, \rho > 0. \end{cases} \quad (2-12)$$

If  $u \in H_k^\mu$  with  $(\mu, k) \in \mathbb{R}^2$  and  $a_h \in S_{-\infty}^-$  such that

$$\mathcal{N}_{\delta, \rho} \cap \overline{\bigcup_{0 < h < 1} \text{supp } a_h} = \emptyset, \quad (2-13)$$

then  $\text{Op}_h^{\delta, \rho}(a_h)u = \mathcal{O}(h^{\delta k + \rho \mu})_{L^2}$  and consequently  $\text{WF}_{\delta, \rho}^{\delta k + \rho \mu}(u)^\circ = \emptyset$ .

*Proof.* The statements (1) and (2) are consequences of the quasihomogeneous scaling (2-2) we used to define the pseudodifferential operators. To prove (3), note that if  $a_h \in S_{-\infty}^-$  and  $\mathcal{F}$  is the Fourier transform operator, then

$$\mathcal{F}^{-1} \text{Op}_h^{\rho, \delta}(a_h)\mathcal{F} = \text{Op}_h^{\delta, \rho}(b_h),$$

where  $b_h(x, \xi) = a_h(\xi, -x)$ . To prove (4), we use  $\overline{\text{Op}(a_h)u_h} = \text{Op}(c_h)\bar{u}_h$ , where  $c_h(x, \xi) = \overline{a_h(x, -\xi)}$ . To prove (5), note that if  $a_h$  satisfies the condition (2-13), then

$$(\theta_{h, * }^{\delta, \rho} a_h) \langle \xi \rangle^{-\mu} \#_1^{0, 0} \langle x \rangle^{-k} = \mathcal{O}(h^{\delta k + \rho \mu})_{S_0^0}. \quad \square$$

### 3. Model equations

We prove Theorem 1.4 by combining the ideas of [Nakamura 2005] and simple scaling arguments.

**3A. Proof of Theorem 1.4(1).** If  $a \in W_{\text{loc}}^{1, \infty}(\mathbb{R} \times \mathbb{R}^{2d})$  and  $\mathcal{A} \in W_{\text{loc}}^{1, \infty}(\mathbb{R}, L^2 \rightarrow L^2)$ , then define

$$\mathcal{L}_t a = \partial_t a + \{|\xi|^\gamma, a\}, \quad \mathcal{L}_t \mathcal{A} = \partial_t \mathcal{A} + i[|D_x|^\gamma, \mathcal{A}].$$

Here  $\{ \cdot, \cdot \}$  denotes the Poisson bracket defined by  $\{f, g\} = \partial_\xi f \cdot \partial_x g - \partial_x f \cdot \partial_\xi g$ .

**Lemma 3.1.** If  $a_h \in W_{\text{loc}}^{1, \infty}(\mathbb{R}, S_{-\infty}^-)$  and satisfies the condition

$$\overline{\bigcup_{0 < h < 1} \text{supp } a_h \cap \{\xi = 0\}} = \emptyset,$$

then there exists  $b_h \in L_{\text{loc}}^\infty(\mathbb{R}, S_{-\infty}^-)$ , with  $\text{supp } b_h \subset \text{supp } a_h$ , such that

$$\mathcal{L}_t \text{Op}_h^{\delta, \rho}(a_h) = \text{Op}_h^{\delta, \rho}(\mathcal{L}_t a_h) + h^{\delta + \rho} \text{Op}_h^{\delta, \rho}(b_h) + \mathcal{O}(h^\infty)_{L_{\text{loc}}^\infty(\mathbb{R}, L^2 \rightarrow L^2)}.$$

*Proof.* For all  $T > 0$ , there exists  $\epsilon > 0$  such that

$$\overline{\bigcup_{t \in [-T, T]} \bigcup_{0 < h < 1} \text{supp } a_h(t, \cdot) \cap \{|\xi| \leq \epsilon\}} = \emptyset.$$

Let  $\pi \in C^\infty(\mathbb{R}^d)$  such that  $0 \leq \pi \leq 1$ ,  $\pi(\xi) = 0$  for  $|\xi| \leq \frac{\epsilon}{3}$ , and  $\pi(\xi) = 1$  for  $|\xi| \geq \frac{2\epsilon}{3}$ . Then

$$i[|D_x|^\gamma, \text{Op}_h^{\delta, \rho}(a_h)] = ih^{-\rho\gamma} [h^\rho |D_x|^\gamma \pi(h^\rho D_x), \text{Op}_h^{\delta, \rho}(a_h)] + \mathcal{O}(h^\infty)_{L^\infty([-T, T], L^2 \rightarrow L^2)}.$$

Now that  $|\xi|^\gamma \pi(\xi) \in S_0^\gamma$ , we conclude by Proposition 2.4 and the hypothesis  $\rho\gamma = \delta + \rho$ . □



Assume that  $\mu = \infty$ , as the proof is similar for  $\mu < \infty$ . Let  $(x_0, \xi_0) \notin \text{WF}_{\delta, \rho}^\mu(u_0)$  with  $\xi_0 \neq 0$ . We aim to find  $a_h \in W_{\text{loc}}^{1, \infty}(\mathbb{R}, S_{-\infty}^-)$  of the asymptotic expansion  $a_h \sim \sum_{j \in \mathbb{N}} h^{j(\delta + \rho)} a_h^j$ , where  $a_h^j \in W_{\text{loc}}^{1, \infty}(\mathbb{R}, S_{-\infty}^-)$ , such that, for all  $t \in \mathbb{R}$ ,  $a_h(t, \cdot)$  is elliptic at  $(x_0 + t\gamma|\xi_0|^{\gamma-2}\xi_0, \xi_0)$ , and

$$\mathcal{L}_t \text{Op}_h^{\delta, \rho}(a_h) = \mathcal{O}(h^\infty)_{L_{\text{loc}}^\infty(\mathbb{R}, L^2 \rightarrow L^2)}. \tag{3-1}$$

If such an  $a_h$  is found, we let  $\mathcal{A}_h(t) = \text{Op}_h^{\delta, \rho}(a_h(t))$  and

$$v_h(t) = e^{it|D_x|^\gamma} \mathcal{A}_h(t) e^{-it|D_x|^\gamma} u_0,$$

then by a direct computation and (3-1), we have

$$\partial_t v_h = e^{it|D_x|^\gamma} \mathcal{L}_t \mathcal{A}_h e^{-it|D_x|^\gamma} u_0 = \mathcal{O}(h^\infty)_{L_{\text{loc}}^\infty(\mathbb{R}, L^2 \rightarrow L^2)}. \tag{3-2}$$

If we assume that  $\text{supp } a_h(0)$  is sufficiently close to  $(x_0, \xi_0)$  so that

$$\overline{\bigcup_{h \in (0, 1]} \text{supp } a_h(0)} \Subset \mathbb{R}^{2d} \setminus \text{WF}_{\delta, \rho}^\mu(u_0),$$

then by Lemma 2.14, we have  $v_h(0) = \text{Op}_h^{\delta, \rho}(a_h(0))u_0 = \mathcal{O}(h^\infty)_{L^2}$ . Therefore by (3-2), we have  $v_h \in \mathcal{O}(h^\infty)_{L_{\text{loc}}^\infty(\mathbb{R}, L^2)}$  and thus  $\mathcal{A}_h u \in \mathcal{O}(h^\infty)_{L_{\text{loc}}^\infty(\mathbb{R}, L^2)}$ .

To construct  $a_h$ , let  $\varphi \in C_c^\infty(\mathbb{R}^d \times (\mathbb{R}^d \setminus 0))$  with  $\varphi(x_0, \xi_0) \neq 0$ , such that  $\text{Op}_h^{\delta, \rho}(\varphi)u = \mathcal{O}(h^\infty)_{L^2}$ . Then we can construct  $a_h$  with  $a_h|_{t=0} = \varphi$ , with  $a_h^j \in W_{\text{loc}}^{\infty, \infty}(\mathbb{R}, S_{-\infty}^-)$ , by solving iteratively the transportation equations

$$\begin{cases} \mathcal{L}_t a_h^0 = 0, & \mathcal{L}_t a_h^j + b_h^{j-1} = 0, \\ a_h^0|_{t=0} = \varphi, & a_h^j|_{t=0} = 0, \quad j \geq 1, \end{cases}$$

where  $b_h^j \in W_{\text{loc}}^{\infty, \infty}(\mathbb{R}, S_{-\infty}^-)$  satisfies, by Lemma 3.1, that

$$\mathcal{L}_t \text{Op}_h^{\delta, \rho}(a_h^j) = \text{Op}_h^{\delta, \rho}(\mathcal{L}_t a_h^j) + h^{\delta + \rho} \text{Op}_h^{\delta, \rho}(b_h^j) + \mathcal{O}(h^\infty)_{L_{\text{loc}}^\infty(\mathbb{R}, L^2 \rightarrow L^2)}.$$

Thus we have proved Theorem 1.4(1).

**3B. Proof of Theorem 1.4(2).** Let  $\beta = \rho\gamma - (\delta + \rho) > 0$ . For all  $h > 0$ , introduce the semiclassical time variable  $s = h^{-\beta}t$ , and rewrite (1-6) as

$$\partial_s u + ih^\beta |D_x|^\gamma u = 0. \tag{3-3}$$

If  $a = a(s, x, \xi) \in W_{\text{loc}}^{1, \infty}(\mathbb{R} \times \mathbb{R}^{2d})$  and  $\mathcal{A} = \mathcal{A}(s) \in W_{\text{loc}}^{1, \infty}(\mathbb{R}, L^2 \rightarrow L^2)$ , then define

$$\mathcal{L}_s a = \partial_s a + \{|\xi|^\gamma, a\}, \quad \mathcal{L}_s^h \mathcal{A} = \partial_s \mathcal{A} + ih^\beta [ |D_x|^\gamma, \mathcal{A} ].$$

**Lemma 3.2.** *If  $\phi \in C_c^\infty(\mathbb{R}^d)$  such that  $\phi \geq 0$ ,  $\phi(0) > 0$ , and  $x \cdot \nabla \phi(x) \leq 0$  for all  $x \in \mathbb{R}^d$ , and we define*

$$\chi(s, x, \xi) = \phi\left(\frac{x - s\gamma|\xi|^{\gamma-2}\xi - x_0}{1 + s}\right) \phi\left(\frac{\xi - \xi_0}{\epsilon}\right)$$

for  $s \geq 0$ ,  $\epsilon > 0$ ,  $(x_0, \xi_0) \in \mathbb{R}^d \times (\mathbb{R}^d \setminus 0)$ , then the following statements hold when  $\epsilon$  is sufficiently small and  $|\xi_0|$  is sufficiently large:

- (1)  $\chi \in W^{\infty, \infty}(\mathbb{R}_{\geq 0}, S_0^{-\infty})$ .
- (2)  $\mathcal{L}_s \chi \in W^{\infty, \infty}(\mathbb{R}_{\geq 0}, S_{-1}^{-\infty})$  and  $\mathcal{L}_s \chi \geq 0$ .
- (3) If  $t_0 > 0$  and set  $(\tau u)(s, x, \xi) = u(s, \frac{s}{t_0}x, \xi)$ , then  $\tau \chi \in W^{\infty, \infty}(\mathbb{R}_{\geq 0}, S_{-\infty}^{-\infty})$ .
- (4) If  $s$  is sufficiently large, then  $(\tau \chi)(s, \cdot)$  is elliptic at  $(t_0 \gamma |\xi_0|^{\gamma-2} \xi_0, \xi_0)$ .

*Proof.* Each time we differentiate  $\chi$  with respect to  $x$ , we get a multiplicative factor  $(1+s)^{-1}$ , which is of size  $\langle x \rangle^{-1}$  in  $\text{supp } \chi$  as

$$\text{supp } \chi \subset \{C^{-1}s \leq |x| \leq Cs\} \tag{3-4}$$

for some  $C > 0$  when  $|s|$  and  $|\xi_0|$  are sufficiently large and  $\epsilon$  is sufficiently small. Therefore  $\chi \in W^{\infty, \infty}(\mathbb{R}_{\geq 0}, S_0^{-\infty})$ . Clearly  $\tau \chi(s, \cdot)$  is bounded in  $C_c^\infty(\mathbb{R}^{2d})$ . We write

$$(\tau \chi)(s, x, \xi) = \phi\left(\frac{x - t_0 \gamma |\xi_0|^{\gamma-2} \xi_0}{t_0(1+s)/s} - \frac{\gamma |\xi|^{\gamma-2} \xi - \gamma |\xi_0|^{\gamma-2} \xi_0}{(1+s)/s} - \frac{x_0}{1+s}\right) \phi\left(\frac{\xi - \xi_0}{\epsilon}\right), \tag{3-5}$$

where  $|\xi|^{\gamma-2} \xi - |\xi_0|^{\gamma-2} \xi_0 = o(1)$  as  $\epsilon \rightarrow 0$ , whence  $\tau \chi(s, \cdot)$  is elliptic at  $(t_0 \gamma |\xi_0|^{\gamma-2} \xi_0, \xi_0)$  for sufficiently large  $s$ . To estimate  $\mathcal{L}_s \chi$ , we perform an explicit computation:

$$\begin{aligned} \partial_s \chi(s, x, \xi) &= -(\nabla \phi) \left( \frac{x - s \gamma |\xi|^{\gamma-2} \xi - x_0}{1+s} \right) \phi \left( \frac{\xi - \xi_0}{\epsilon} \right) \frac{(x - s \gamma |\xi|^{\gamma-2} \xi - x_0) + (1+s) \gamma |\xi|^{\gamma-2} \xi}{(1+s)^2}, \\ \{|\xi|^\gamma, \chi\}(s, x, \xi) &= \frac{\gamma |\xi|^{\gamma-2} \xi}{1+s} \cdot (\nabla \phi) \left( \frac{x - s \gamma |\xi|^{\gamma-2} \xi - x_0}{1+s} \right) \phi \left( \frac{\xi - \xi_0}{\epsilon} \right). \end{aligned}$$

Therefore,

$$\mathcal{L}_s \chi(s, x, \xi) = -(\nabla \phi) \left( \frac{x - s \gamma |\xi|^{\gamma-2} \xi - x_0}{1+s} \right) \phi \left( \frac{\xi - \xi_0}{\epsilon} \right) \cdot \frac{x - s \gamma |\xi|^{\gamma-2} \xi - x_0}{(1+s)^2} \geq 0.$$

Note that on  $\text{supp } \mathcal{L}_s \chi$ , we have

$$\frac{x - s \gamma |\xi|^{\gamma-2} \xi - x_0}{(1+s)^2} = \mathcal{O}\left(\frac{1+s}{(1+s)^2}\right) = \mathcal{O}\left(\frac{1}{1+s}\right) = \mathcal{O}\left(\frac{1}{\langle x \rangle}\right).$$

So we prove similarly that  $\mathcal{L}_s \chi \in W^{\infty, \infty}(\mathbb{R}_{\geq 0}, S_{-1}^{-\infty})$ . □

Now fix  $t_0 > 0$  and let  $\mu = \infty$  as the other cases are similar. Let  $\epsilon > 0$  be sufficiently small and let  $\{\lambda_j\}_{j \in \mathbb{N}} \subset [1, 1 + \epsilon)$  be strictly increasing. Choose  $\phi$  as in [Lemma 3.2](#), and set

$$\chi_j(s, x, \xi) = \phi\left(\frac{x - s \gamma |\xi|^{\gamma-2} \xi - x_0}{\lambda_j(1+s)}\right) \phi\left(\frac{\xi - \xi_0}{\lambda_j \epsilon}\right).$$

Then  $\text{supp } \chi_j \subset \{\chi_{j+1} > 0\}$  for all  $j \in \mathbb{N}$ . We aim to construct  $a_h \in W^{\infty, \infty}(\mathbb{R}_{\geq 0}, S_0^{-\infty})$  such that the following statements hold:

- (1) For all  $s \geq 0$  and  $h \in (0, 1]$ , we have  $\text{supp } a_h \subset \bigcup_{j \in \mathbb{N}} \text{supp } \chi_j$ .
- (2) The symbol  $a_h|_{s=0}$  is elliptic at  $(x_0, \xi_0)$ ; more precisely, we have

$$(a_h - (\zeta_h^{\delta, \rho} \chi_0) \sharp_h^{\delta, \rho} \chi_0)|_{s=0} = \mathcal{O}(h^\infty)_{S_{-\infty}^{-\infty}}.$$

(3) For  $t_0 > 0$ , let  $\tau$  be defined as in the lemma. Then  $\tau a_h \in W^{\infty, \infty}(\mathbb{R}_{\geq 0}, S_{-\infty}^-)$  and  $\tau a_h(s, \cdot)$  is elliptic at  $(t_0 \gamma |\xi_0|^{\gamma-2} \xi_0, \xi_0)$  when  $s$  is sufficiently large.

(4)  $\mathcal{L}_s^h \text{Op}_h^{\delta, \rho}(a_h) \geq \mathcal{O}(h^\infty)_{L^\infty(\mathbb{R}_{\geq 0}, L^2 \rightarrow L^2)}$ .

Assume that such an  $a_h$  is found and that

$$(t_0 \gamma |\xi_0|^{\gamma-2} \xi_0, \xi_0) \notin \text{WF}_{\rho(\gamma-1), \rho}^\mu(u(t = t_0)).$$

By (1) and (3-5), if we choose  $\phi$  such that  $\text{supp } \phi$  is sufficiently close to the origin, then for sufficiently small  $h > 0$  we have

$$\text{supp } \theta_{1/h, * }^{\beta, 0} a_h|_{s=h^{-\beta} t_0} \in \mathbb{R}^{2d} \setminus \text{WF}_{\rho(\gamma-1), \rho}^\infty(u|_{t=t_0}).$$

By (3), the symbol  $\theta_{1/h, * }^{\beta, 0} a_h|_{s=h^{-\beta} t_0} \in S_{-\infty}^-$  is elliptic at  $(t_0 \gamma |\xi_0|^{\gamma-2} \xi_0, \xi_0)$ . Therefore, by Lemma 2.14,

$$(u, \text{Op}_h^{\delta, \rho}(a_h)u)_{L^2}|_{s=h^{-\beta} t_0} = (u, \text{Op}_h^{\rho(\gamma-1), \rho}(\theta_{1/h, * }^{\beta, 0} a_h)u)_{L^2}|_{s=h^{-\beta} t_0} = \mathcal{O}(h^\infty).$$

By (3-3), we have

$$\frac{d}{ds}(u, \text{Op}_h^{\delta, \rho}(a_h)u)_{L^2} = (u, \mathcal{L}_s^h \text{Op}_h^{\delta, \rho}(a_h)u)_{L^2},$$

which implies, by (4), that

$$\begin{aligned} (u, \text{Op}_h^{\delta, \rho}(a_h)u)_{L^2}|_{s=0} &= (u, \text{Op}_h^{\delta, \rho}(a_h)u)_{L^2}|_{s=h^{-\beta} t_0} - \int_0^{h^{-\beta} t_0} (u, \mathcal{L}_s^h \text{Op}_h^{\delta, \rho}(a_h)u)_{L^2} ds \\ &\leq \mathcal{O}(h^\infty) + \mathcal{O}(h^{-\beta} \times h^\infty) = \mathcal{O}(h^\infty). \end{aligned}$$

Therefore, by (2), we have

$$\|\text{Op}_h^{\delta, \rho}(\chi_0)u|_{s=0}\|_{L^2}^2 = (u, \text{Op}_h^{\delta, \rho}(a_h)u)_{L^2}|_{s=0} + \mathcal{O}(h^\infty) = \mathcal{O}(h^\infty).$$

We conclude that  $(x_0, \xi_0) \notin \text{WF}_{\delta, \rho}^\infty(u_0)$ .

We shall construct  $a_h$  in the following form of asymptotic expansion:

$$a_h(s, x, \xi) \sim \sum_{j \in \mathbb{N}} h^{j(\delta+\rho)} \varphi^j(s) a_h^j(s, x, \xi),$$

where  $a_h^j \in W^{\infty, \infty}(\mathbb{R}_{\geq 0}, S_0^-)$ , with  $\text{supp } a_h^j \subset \text{supp } \chi_j$ , and  $\varphi^j \in P_j$ , with

$$P_j = \left\{ f(\ln(1+s)) : f(X) = \sum_{k=0}^j c_k X^k, c_k \geq 0 \text{ for all } k \right\}. \tag{3-6}$$

The above asymptotic expansion is in the weak sense that, for some  $\epsilon' > 0$ , and all  $N \in \mathbb{N}$ ,

$$a_h - \sum_{j < N} h^{j(\delta+\rho)} \varphi^j a_h^j \in \mathcal{O}(h^{N(\delta+\rho-\epsilon')})_{W^{\infty, \infty}([0, h^{-\beta} T], S_0^-)}.$$

The following properties for functions in  $\bigcup_{j \in \mathbb{N}} P_j$  will be used in the construction of  $a_h$ .

**Lemma 3.3.** *If  $\psi \in P_j$  for some  $j \in \mathbb{N}$ , then  $\psi$  is smooth and nonnegative on  $[0, +\infty)$  and*

$$((1+s)\partial_s)^{-1} \psi(s) := \int_0^s (1+\sigma)^{-1} \psi(\sigma) d\sigma \in P_{j+1}.$$

*Proof.* The function  $\psi$  is smooth because it is the composition of a polynomial and the smooth function  $s \mapsto \ln(1 + s)$ . The nonnegativity of  $\psi$  is the consequence of the nonnegativity of the coefficients  $c_k$  in (3-6) and the fact that  $\ln(1 + s) \geq 0$  when  $s \geq 0$ . To prove that  $((1 + s)\partial_s)^{-1}\psi \in P_{j+1}$ , note that for all  $n \in \mathbb{N}$  we have

$$((1 + s)\partial_s)^{-1}(\ln(1 + \cdot))^n = (n + 1)^{-1}(\ln(1 + \cdot))^{n+1}.$$

The claim follows by the linearity of the operator  $((1 + s)\partial_s)^{-1}$ . □

To construct  $a_h$ , we begin by setting  $\varphi^0 \equiv 1$  and choosing  $a_h^0$  satisfying

$$\begin{aligned} a_h^0 - (\zeta_h^{\delta,\rho} \chi_0) \sharp_h^{\delta,\rho} \chi_0 &= \mathcal{O}(h^\infty)_{W^\infty(\mathbb{R}_{\geq 0}, S_0^{-\infty})}, \\ (a_h^0 - (\zeta_h^{\delta,\rho} \chi_0) \sharp_h^{\delta,\rho} \chi_0)|_{s=0} &= \mathcal{O}(h^\infty)_{S^{-\infty}}. \end{aligned}$$

By the definition of  $\beta$  and Propositions 2.4 and 2.5, there exists  $r_h^0 \in L^\infty(\mathbb{R}_{\geq 0}, S_{-1}^{-\infty})$  with  $\text{supp } r_h^0 \subset \text{supp } \chi_0$  such that

$$\mathcal{L}_s^h \text{Op}_h^{\delta,\rho}(a_h^0) = 2 \text{Op}_h^{\delta,\rho}(\chi_0 \mathcal{L}_s \chi_0) + h^{\delta+\rho} \text{Op}_h^{\delta,\rho}(r_h^0) + \mathcal{O}(h^\infty)_{L^\infty(\mathbb{R}_{\geq 0}, L^2 \rightarrow L^2)}. \tag{3-7}$$

By (3-4), we have  $\langle s \rangle r_h^0 \in L^\infty(\mathbb{R}_{\geq 0}, S_0^{-\infty})$  and similarly

$$\langle s \rangle \chi_0 \mathcal{L}_s \chi_0 \in L^\infty(\mathbb{R}_{\geq 0}, S_0^{-\infty}). \tag{3-8}$$

By Lemma 3.2, we have

$$\chi_0 \mathcal{L}_s \chi_0 \geq 0. \tag{3-9}$$

Recall that, by the sharp Gårding inequality (Proposition 2.6), if a symbol  $p_h \in S_0^0$  satisfies  $p_h \geq 0$ , then  $\text{Op}_h^{0,1}(p_h) \gtrsim -h$ . By (2-3), we deduce that  $\text{Op}_h^{0,1}(p_h) \gtrsim -h^{\delta+\rho}$ . To be precise, this means there exists  $C > 0$  which only depends on a finite number of seminorms defined by (2-1), such that, for all  $u \in L^2$ ,

$$\langle u, \text{Op}_h^{\delta,\rho}(p_h)u \rangle_{L^2} \geq -Ch^{\delta+\rho} \|u\|_{L^2}^2.$$

Take  $c_h^0 \in L^\infty(\mathbb{R}_{\geq 0}, S_0^{-\infty})$  such that

$$\text{supp } a_h^0 \Subset \{c_h^0 = 1\} \subset \text{supp } c_h^0 \subset \{\chi_1 > 0\}.$$

By (3-7) and (3-9), for all  $u \in L^2$ , we have

$$\begin{aligned} \langle u, \mathcal{L}_s^h \text{Op}_h^{\delta,\rho}(a_h^0)u \rangle_{L^2} &= \langle \text{Op}_h^{\delta,\rho}(c_h)u, \mathcal{L}_s^h \text{Op}_h^{\delta,\rho}(a_h^0) \text{Op}_h^{\delta,\rho}(c_h)u \rangle_{L^2} + \mathcal{O}(h^\infty) \|u\|_{L^2}^2 \\ &\geq -C \langle s \rangle^{-1} h^{\delta+\rho} \|\text{Op}_h^{\delta,\rho}(c_h)u\|_{L^2}^2 + \mathcal{O}(h^\infty), \end{aligned}$$

where the factor  $\langle s \rangle^{-1}$  comes from the estimate (3-8). By the symbolic calculus, there exists  $b_h \in L^\infty(\mathbb{R}_{\geq 0}, S_0^{-\infty})$  such that

$$\text{Op}_h^{\delta,\rho}(b_h) - C \text{Op}_h^{\delta,\rho}(c_h)^* \text{Op}_h^{\delta,\rho}(c_h) = \mathcal{O}(h^\infty)_{L^\infty(\mathbb{R}_{\geq 0}, L^2 \rightarrow L^2)}$$

and  $\text{supp } b_h \subset \text{supp } c_h$ . Therefore,

$$\begin{aligned} \mathcal{L}_s^h \text{Op}_h^{\delta,\rho}(a_h^0) &\geq -C \langle s \rangle^{-1} h^{\delta+\rho} \text{Op}_h^{\delta,\rho}(c_h)^* \text{Op}_h^{\delta,\rho}(c_h) + \mathcal{O}(h^\infty)_{L^\infty(\mathbb{R}_{\geq 0}, L^2 \rightarrow L^2)} \\ &\geq -\langle s \rangle^{-1} h^{\delta+\rho} \text{Op}_h^{\delta,\rho}(b_h^0) + \mathcal{O}(h^\infty)_{L^\infty(\mathbb{R}_{\geq 0}, L^2 \rightarrow L^2)}. \end{aligned} \tag{3-10}$$

Suppose that, for some  $\ell \geq 1$ , we can find  $\varphi^j \in P_j$ ,  $a_h^j$  for  $j = 0, \dots, \ell - 1$  and  $\psi^{\ell-1} \in P_{\ell-1}$ ,  $b_h^{\ell-1} \in L^\infty(\mathbb{R}_{\geq 0}, S_0^{-\infty})$ , with  $\text{supp } b_h^{\ell-1} \subset \{\chi_\ell > 0\}$ , such that

$$\mathcal{L}_s^h \text{Op}_h^{\delta, \rho} \left( \sum_{j=0}^{\ell-1} h^{j(\delta+\rho)} \varphi^j a_h^j \right) \geq -\langle s \rangle^{-1} \psi^{\ell-1} h^{\ell(\delta+\rho)} \text{Op}_h^{\delta, \rho}(b_h^{\ell-1}) + \mathcal{O}(h^\infty)_{L^\infty(\mathbb{R}_{\geq 0}, L^2 \rightarrow L^2)}. \quad (3-11)$$

If we choose  $B_\ell > 0$  sufficiently large and set  $\varphi^\ell = ((1+s)\partial_s)^{-1} \psi^{\ell-1}$  and  $a_h^\ell = B_\ell \chi_\ell$ , then by a direct calculation, we have

$$\begin{aligned} \mathcal{L}_s(\varphi^\ell a_h^\ell) &= B_\ell (1+s)^{-1} \psi^{\ell-1} \chi_\ell + B_\ell \varphi^\ell \mathcal{L}_s \chi_\ell \\ &\geq B_\ell (1+s)^{-1} \psi^{\ell-1} \chi_\ell \geq \langle s \rangle^{-1} \psi^{\ell-1} b_h^{\ell-1}. \end{aligned}$$

Observe that

$$\mathcal{L}_s(\varphi^\ell a_h^\ell) = \mathcal{O}(\langle s \rangle^{-1} (\psi^{\ell-1} + \varphi^\ell))_{S_0^{-\infty}}, \quad \langle s \rangle^{-1} \psi^{\ell-1} b_h^{\ell-1} = \mathcal{O}(\langle s \rangle^{-1} \psi^{\ell-1})_{S_0^{-\infty}}.$$

Much as above, applying the sharp Gårding inequality to the symbol

$$\mathcal{L}_s(\varphi^\ell a_h^\ell) - \langle s \rangle^{-1} \psi^{\ell-1} b_h^{\ell-1} = \mathcal{O}(\langle s \rangle^{-1} (\varphi^\ell + \psi^{\ell-1}))_{S_0^{-\infty}},$$

we can find  $b_h^\ell \in L^\infty(\mathbb{R}_{\geq 0}, S_0^{-\infty})$  with  $\text{supp } b_h^\ell \subset \{\chi_{\ell+1} > 0\}$  such that

$$\mathcal{L}_s^h \text{Op}_h^{\delta, \rho}(\varphi^\ell a_h^\ell) - \langle s \rangle^{-1} \psi^{\ell-1} h^{\ell(\delta+\rho)} \text{Op}_h^{\delta, \rho}(b_h^{\ell-1}) \geq -\langle s \rangle^{-1} \psi^\ell h^{\delta+\rho} \text{Op}_h^{\delta, \rho}(b_h^\ell) + \mathcal{O}(h^\infty)_{L^2 \rightarrow L^2}, \quad (3-12)$$

with  $\psi^\ell = \psi^{\ell-1} + \varphi^\ell \in P_\ell$ . Summing up (3-11) and  $h^{\ell(\delta+\rho)} \times$  (3-12), we close the induction procedure.

Finally we conclude the asymptotic expansion by observing that, for all  $\epsilon' > 0$ , we have

$$\|\varphi^\ell\|_{L^\infty([0, h^{-\beta T}])} = \mathcal{O}(|\log h|^\ell) = \mathcal{O}(h^{-\epsilon' \ell}).$$

Thus we have proved [Theorem 1.4\(2\)](#).

#### 4. Paradifferential calculus

In this section, we develop a paradifferential calculus on weighted Sobolev spaces and a semiclassical paradifferential calculus.

**4A. Classical paradifferential calculus.** We recall some classical results of the paradifferential calculus. We refer to the original work [[Bony 1979](#)] and the books [[Hörmander 1997](#); [Métivier 2008](#); [Bahouri, Chemin and Danchin 2011](#)]. The results and proofs below are mainly based on [[Métivier 2008](#)], so we shall only sketch them. In the meantime, we shall also make some refinements regarding the estimates of the remainder terms, for the sake of the semiclassical paradifferential calculus that will be developed later.

**4A1. Symbol classes and paradifferential operators.**

**Definition 4.1.** For  $m \in \mathbb{R}$ ,  $r \geq 0$ , let  $\Gamma^{m, r}$  be the space of all  $a(x, \xi) \in L_{\text{loc}}^\infty(\mathbb{R}^d \times (\mathbb{R}^d \setminus \{0\}))$  such that:

- (1) For all  $x \rightarrow \mathbb{R}^d$ , the function  $\xi \mapsto a(x, \xi)$  is smooth.
- (2) For all  $\alpha \in \mathbb{N}^d$ , there exists  $C_\alpha > 0$  such that for all  $\xi \in \mathbb{R}^d$  with  $|\xi| \geq \frac{1}{2}$ , we have

$$\|\partial_\xi^\alpha a(\cdot, \xi)\|_{W^{r, \infty}} \leq C_\alpha \langle \xi \rangle^{m-|\alpha|}.$$

If  $a \in \Gamma^{m,r}$ , then we define for all  $n \geq 0$  the seminorm

$$M_n^{m,r}(a) = \sup_{|\alpha| \leq n} \sup_{|\xi| \geq 1/2} \langle \xi \rangle^{|\alpha|-m} \|\partial_\xi^\alpha a(\cdot, \xi)\|_{W^{r,\infty}}.$$

We also define  $M^{m,r}(a) = M_{\tilde{d}+r}^{m,r}(a)$ , where  $\tilde{d} = 1 + \lfloor \frac{d}{2} \rfloor$ .

**Definition 4.2.** A pair of nonnegative functions  $(\chi, \pi) \in C^\infty(\mathbb{R}^{2d} \setminus 0) \times C^\infty(\mathbb{R}^d)$  is called admissible if the following conditions are satisfied:

- (1) The function  $1 - \pi$  is a cutoff function of the origin. To be precise, if  $|\eta| \geq 1$ , then  $\pi(\eta) = 1$ , and if  $|\eta| \leq \frac{1}{2}$ , then  $\pi(\eta) = 0$ .
- (2) The function  $\chi$  is an even and homogeneous of degree 0, and there exist  $\epsilon_1, \epsilon_2 \in (0, 1)$  with  $\epsilon_1 < \epsilon_2$ , such that

$$\begin{cases} \chi(\theta, \eta) = 1, & |\theta| \leq \epsilon_1 |\eta|, \\ \chi(\theta, \eta) = 0, & |\theta| \geq \epsilon_2 |\eta|. \end{cases} \tag{4-1}$$

**Definition 4.3.** If  $m \in \mathbb{R}$  and  $a \in \Gamma^{m,0}$ , then the paradifferential operator  $T_a$  is defined by

$$\widehat{T_a u}(\xi) = (2\pi)^{-d} \int_{\mathbb{R}^d} \chi(\xi - \eta, \eta) \pi(\eta) \hat{a}(\xi - \eta, \eta) \hat{u}(\eta) \, d\eta, \tag{4-2}$$

where  $(\chi, \pi)$  is admissible and  $\hat{a}(\theta, \xi) = \int e^{-ix \cdot \theta} a(x, \xi) \, dx$ . In other words  $T_a = \text{Op}(\sigma_a)$  where

$$\sigma_a(\cdot, \xi) = \pi(\xi) \chi(D_x, \xi) a(\cdot, \xi). \tag{4-3}$$

**Proposition 4.4.** If  $m \in \mathbb{R}$  and  $a \in \Gamma^{m,0}$ , then  $T_a = \mathcal{O}(M^{m,0}(a))_{\mathcal{O}_0^m}$ .

**Remark 4.5.** A symbol  $p$  satisfies the spectral condition if there exists  $\epsilon \in (0, 1)$  such that

$$\text{supp } \hat{p} \subset \{(\eta, \xi) \in \mathbb{R}^{2d} : |\eta| \geq \epsilon \langle \xi \rangle\}.$$

By [Métivier 2008], if  $a \in \Gamma^{m,0}$ , then  $\sigma_a \in \Gamma^{m,0}$  and satisfies the spectral condition. The above Proposition 4.4 is in fact a consequence of the following estimate (4-4) and the mapping property: if  $p \in \Gamma^{m,0}$  satisfies the spectral condition, then  $\text{Op}(p)$  defines a bounded operator from  $H^{\mu+m} \rightarrow H^\mu$  for all  $\mu \in \mathbb{R}$ .

Note the definition (4-2) depends on the choice of admissible pairs of functions. The following lemma and corollary show that if we change the admissible pair, then the error term is regularizing.

**Lemma 4.6.** If  $m \in \mathbb{R}$ ,  $r \geq 0$  and  $a \in \Gamma^{m,r}$ , then, for all  $n \geq 0$ , we have

$$M_n^{m,r}(\sigma_a) \lesssim M_n^{m,r}(a). \tag{4-4}$$

If in addition  $r \in \mathbb{N}$ , then, for all  $\beta \in \mathbb{N}^d$  with  $|\beta| \leq r$ , we have

$$M_n^{m-r+|\beta|,0}(\partial_x^\beta(\sigma_a - a\pi)) \lesssim M_n^{m,0}(\nabla^r a). \tag{4-5}$$

*Proof.* The first statement is proven in [Métivier 2008]. We only prove the second statement. We shall only prove the case where  $\beta = 0$  for the rest is similar. By [Métivier 2008], we have

$$(\sigma_a - a\pi)(x, \xi) = \pi(\xi) \int \rho(x, y, \xi) \Phi(y, \xi) \, dy$$

for all  $(x, \xi) \in \mathbb{R}^d \times (\mathbb{R}^d \setminus 0)$ , where  $\Phi(\cdot, \xi) = \mathcal{F}^{-1}\chi(\cdot, \xi)$  and

$$\rho(x, y, \xi) = \sum_{|\gamma|=r} \frac{(-y)^\gamma}{\gamma!} \int_0^1 r(1-t)^{r-1} \partial_x^\gamma a(x-ty, \xi) dt.$$

Therefore, if  $|\xi| \geq \frac{1}{2}$  and  $|\alpha| \leq n$ , then

$$\|\partial_\xi^\alpha \rho(\cdot, y, \xi)\|_{L^\infty} \lesssim |y|^r \|\partial_\xi^\alpha \nabla^r a(\cdot, \xi)\|_{L^\infty} \lesssim |y|^r |\xi|^{m-|\alpha|} M_n^{m,0}(\nabla^r a). \tag{4-6}$$

Note that the admissibility of  $(\pi, \chi)$  implies that, for all  $\alpha, \beta \in \mathbb{N}$ , there exists  $C_{\alpha,\beta} > 0$  such that, for all  $(x, \xi) \in \mathbb{R}^{2d}$ , we have  $|x^\beta \partial_\xi^\alpha \Phi(x, \xi)| \leq C_{\alpha,\beta} \langle \xi \rangle^{d-|\alpha|-|\beta|}$ . Therefore, for all  $\alpha \in \mathbb{N}$  and all  $\xi \in \mathbb{R}^d$ , there exists  $C_\alpha > 0$  such that

$$\|\partial_\xi^\alpha \Phi(\cdot, \xi)\|_{L^1} \leq C_\alpha \langle \xi \rangle^{-|\alpha|}. \tag{4-7}$$

We conclude (4-5) by estimates (4-6) and (4-7). □

**Corollary 4.7.** *Let  $a \in \Gamma^{m,r}$ , with  $m \in \mathbb{R}$  and  $r \in \mathbb{N}$ . Let  $(\chi, \pi)$  and  $(\chi', \pi')$  be admissible. Denote by  $T_a$  and  $T'_a$  the paradifferential operators respectively defined by these two admissible pairs. Then*

$$T_a - T'_a = \mathcal{O}(M^{m,0}(\nabla^r a))_{\mathcal{O}_0^{m-r}} + \mathcal{O}(M^{m,r}(a))_{\mathcal{O}_0^{-\infty}}.$$

If in addition  $a\pi = a\pi' = a$ , then

$$T_a - T'_a = \mathcal{O}(M^{m,0}(\nabla^r a))_{\mathcal{O}_0^{m-r}}.$$

*Proof.* Let  $T''_a$  be the paradifferential operator defined with respect to  $(\chi', \pi)$ ; then by Lemma 4.6,  $T_a - T''_a = \mathcal{O}(M^{m,0}(\nabla^r a))_{\mathcal{O}_0^{m-r}}$ . Note that  $T''_a - T'_a$  is a composition with a paradifferential operator with a smoothing operator  $\pi(D_x) - \pi'(D_x)$ , which implies  $T''_a - T'_a = \mathcal{O}(M^{m,r}(a))_{\mathcal{O}_0^{-\infty}}$ . This term vanishes if  $a\pi = a\pi' = a$ . □

**Corollary 4.8.** *Let  $\psi \in C_b^\infty(\mathbb{R}^d)$ . Then  $T_\psi - \psi \in \mathcal{O}_0^{-\infty}$ .*

*Proof.* This is a consequence of (4-5) and the Calderón–Vaillancourt theorem. □

**4A2. Symbolic calculus and parilinearization.**

**Proposition 4.9.** *If  $a \in \Gamma^{m,r}$  and  $b \in \Gamma^{m',r}$ , where  $r \in \mathbb{N}$ ,  $m \in \mathbb{R}$  and  $m' \in \mathbb{R}$ , then*

$$T_a T_b - T_{a\sharp b} = \mathcal{O}(M^{m,r}(a)M^{m',0}(\nabla^r b) + M^{m,0}(\nabla^r a)M^{m',r}(b))_{\mathcal{O}_0^{m+m'-r}} + \mathcal{O}(M^{m,r}(a)M^{m',r}(b))_{\mathcal{O}_0^{-\infty}},$$

where the symbol  $a\sharp b = a\sharp_{1,r}^{0,0} b$  is defined by (2-4). If in addition  $a\pi = a$  and  $b\pi = b$ , then

$$T_a T_b - T_{a\sharp b} = \mathcal{O}(M^{m,r}(a)M^{m',0}(\nabla^r b) + M^{m,0}(\nabla^r a)M^{m',r}(b))_{\mathcal{O}_0^{m+m'-r}}.$$

*Proof.* By Corollary 4.7, we may choose an admissible pair  $(\pi, \chi)$  to define paradifferential operators, while assuming that  $\epsilon_2 < \frac{1}{4}$ . We shall only prove the case where  $a\pi = a$  and  $b\pi = b$ , as the general case follows easily. The following proof follows [Métivier 2008]. Take the decomposition  $T_a T_b - T_{a\sharp b} =$  (I) + (II), where

$$(I) = \text{Op}(\sigma_a) \text{Op}(\sigma_b) - \text{Op}(\sigma_a \sharp \sigma_b), \quad (II) = \text{Op}(\sigma_a \sharp \sigma_b) - \text{Op}(\sigma_{a\sharp b}).$$

Write  $\text{Op}(\sigma_a) \text{Op}(\sigma_b) = \text{Op}(\sigma)$ , where

$$\sigma(x, \xi) = \frac{1}{(2\pi)^d} \iint e^{i(x-y)\cdot\eta} \sigma_a(x, \xi + \eta) \theta(\eta, \xi) \sigma_b(y, \xi) dy d\eta.$$

Here  $\theta \in C^\infty(\mathbb{R}^{2d} \setminus \{0\})$  satisfies that  $(\theta, \pi)$  is admissible and  $\theta\chi = \chi$ . By Taylor's formula, we have the decomposition

$$\sigma_a(x, \xi + \eta) = \sum_{|\alpha| < r} \frac{1}{\alpha!} \partial_\xi^\alpha \sigma(x, \xi) \eta^\alpha + \sum_{|\alpha|=r} \rho_\alpha(x, \xi, \eta) \eta^\alpha,$$

where the functions  $\rho_\alpha$  depend on  $\nabla_\xi^r \sigma_a$ . Then write  $\sigma = \sigma_a \sharp \sigma_b + \sum_{|\alpha|=r} q_\alpha$ , where

$$\begin{aligned} q_\alpha(x, \xi) &= \int R_\alpha(x, x-y, \xi) (D_x^\alpha \sigma_b)(y, \xi) dy, \\ R_\alpha(x, y, \xi) &= (2\pi)^{-2} \int e^{iy\eta} \rho_\alpha(x, y, \eta) \theta(\eta, \xi) d\eta. \end{aligned}$$

By the same estimate in [Métivier 2008],

$$\|\partial_\xi^\beta R_\alpha(x, \cdot, \xi)\|_{L^1} \lesssim M^{m,r}(a) \langle \xi \rangle^{m-r-|\beta|}.$$

Using  $D_x^\alpha \sigma_b = \sigma_{D_x^\alpha b}$ , we verify that

$$\|\partial_\xi^\beta q_\alpha(\cdot, \xi)\|_{L^\infty} \lesssim M^{m,r}(a) M^{m',0}(\nabla^r b) \langle \xi \rangle^{m+m'-r-|\beta|},$$

and consequently, by Remark 4.5,

$$\|(I)\|_{H^s \rightarrow H^{s-m-m'+r}} \lesssim \sum_{|\alpha|=r} M^{m+m'-r,0}(q_\alpha) \lesssim M^{m,r}(a) M^{m',0}(\nabla^r b).$$

To estimate (II), for all  $|\alpha| < r$ , take the decomposition  $\partial_\xi^\alpha \sigma_a D_x^\alpha \sigma_b - \sigma_{\partial_\xi^\alpha a} D_x^\alpha b = (i) + (ii) + (iii)$ , where

$$(i) = \partial_\xi^\alpha (\sigma_a - a) D_x^\alpha \sigma_b, \quad (ii) = \partial_\xi^\alpha a D_x^\alpha (\sigma_b - b\pi), \quad (iii) = \partial_\xi^\alpha a D_x^\alpha b - \sigma_{\partial_\xi^\alpha a} D_x^\alpha b.$$

By Lemma 4.6, Leibniz's rule and interpolation,

$$\begin{aligned} M^{m+m'-r,0}(i) &\lesssim M^{m-r,0}(\sigma_a - a) M^{m',0}(D_x^\alpha \sigma_b) \lesssim M^{m,0}(\nabla^r a) M^{m',r}(b), \\ M^{m+m'-r,0}(ii) &\lesssim M^{m,r}(a) M^{m'-r+|\alpha|,0}(D_x^\alpha (\sigma_b - b\pi)) \lesssim M^{m,r}(a) M^{m',0}(\nabla^r b), \\ M^{m+m'-r,0}(iii) &\lesssim M^{m+m'-|\alpha|,0}(\nabla^{r-|\alpha|}(\partial_\xi^\alpha a D_x^\alpha b)) \\ &\lesssim M^{m-|\alpha|,0}(\nabla^r \partial_\xi^\alpha a) M^{m',0}(b) + M^{m-|\alpha|,0}(\partial_\xi^\alpha a) M^{m',0}(\nabla^r b) \\ &\lesssim M^{m,0}(\nabla^r a) M^{m',r}(b) + M^{m,r}(a) M^{m',0}(\nabla^r b). \end{aligned}$$

By Remark 4.5, these estimates imply that

$$(II) = \mathcal{O}(M^{m,r}(a) M^{m',0}(\nabla^r b) + M^{m,0}(\nabla^r a) M^{m',r}(b))_{\mathcal{O}_0^{m+m'-r}}.$$

The proposition follows. □

**Proposition 4.10.** *Let  $a \in \Gamma^{m,r}$  with  $r \in \mathbb{N}$  and  $m \in \mathbb{R}$ . Then*

$$T_a^* - T_{a^*} = \mathcal{O}(M^{m,0}(\nabla^r a))_{\mathcal{O}_0^{m-r}} + \mathcal{O}(M^{m,r}(a))_{\mathcal{O}_0^{-\infty}},$$



where the symbol  $a^* = \zeta_{1,r}^{0,0} a$  is defined by (2-5). If in addition  $a\pi = a$ , then

$$T_a^* - T_{a^*} = \mathcal{O}(M^{m,0}(\nabla^r a))_{\mathcal{O}_0^{m-r}}.$$

*Proof.* Much as in the proposition for the composition, we shall only prove the case where  $a\pi = a$ . Let  $(\theta, \pi)$  be admissible such that  $\theta\chi = \chi$ , then  $T_a^* = \text{Op}(\sigma_a^*)$ , with

$$\sigma_a^*(x, \xi) = (2\pi)^{-d} \int e^{-iy \cdot \eta} \bar{\sigma}_a(x + y, \xi + \eta) \, d\eta \, dy = a^*(x, \xi) + \sum_{|\alpha|=r} r_\alpha(x, \xi),$$

where by Taylor’s formula,

$$r_\alpha(x, \xi) = \frac{2\pi}{\alpha!} \iiint_{\mathbb{R}^{2d} \times [0,1]} r(1-t)^{r-1} e^{-iy \cdot \eta} D_x^\alpha \partial_\xi^\alpha \bar{\sigma}_a(x, \xi + t\eta) \theta(\eta, \xi) \, dt \, d\eta \, dy.$$

The term  $D_x^\alpha \partial_\xi^\alpha \bar{\sigma}_a(x, \xi + t\eta)$  in the integral and the analysis in [Métivier 2008] imply that

$$M^{m-r,0}(\sigma_a^* - \sigma_{a^*}) \leq \sum_{|\alpha|=r} M^{m-r,0}(r_\alpha) + M^{m-r,0}(a^* - \sigma_{a^*}) \lesssim M^{m,0}(\nabla^r a).$$

The proposition follows by Remark 4.5. □

Recall the following results of parilinearization. See, e.g., [Métivier 2008].

**Proposition 4.11.** *If  $a \in H^\alpha$  and  $b \in H^\beta$  with  $\alpha > \frac{d}{2}$  and  $\beta > \frac{d}{2}$ , then*

$$\|ab - T_a b - T_b a\|_{H^{\alpha+\beta-d/2}} \lesssim \|a\|_{H^\alpha} \|b\|_{H^\beta}.$$

**Proposition 4.12.** *If  $F \in C^\infty(\mathbb{R})$  with  $F(0) = 0$ , then for all  $\mu > \frac{d}{2}$ , there exists a monotonically increasing function  $C : \mathbb{R}_{\geq 0} \rightarrow \mathbb{R}_{\geq 0}$  such that, for all  $u \in H^\mu$ , we have*

$$\|F(u)\|_{H^\mu} + \|F(u) - T_{F'(u)}u\|_{H^{2\mu-d/2}} \leq C(\|u\|_{H^s})\|u\|_{H^\mu}.$$

**4B. Dyadic paradifferential calculus.** Now we develop the theory of paradifferential calculus with weighted symbols on weighted Sobolev spaces via a dyadic decomposition of the space.

**4B1. Weighted symbol classes and dyadic paradifferential operators.** We define a family of symbol classes which take into consideration the spacial decay of symbols.

**Definition 4.13.** If  $r \in \mathbb{N}$ ,  $k \in \mathbb{R}$ , and  $\delta \in [0, 1]$ , then  $W_{k,\delta}^{r,\infty}$  is the set of all  $u \in \mathcal{S}'(\mathbb{R}^d)$  such that

$$\|u\|_{W_{k,\delta}^{r,\infty}} = \sum_{|\alpha| \leq r} \|\langle x \rangle^{-k+\delta|\alpha|} \partial_x^\alpha u\|_{L^\infty} < \infty.$$

**Definition 4.14.** If  $m, k \in \mathbb{R}$ ,  $r \in \mathbb{N}$  and  $\delta \in [0, 1]$ , then  $\Gamma_{k,\delta}^{m,r}$  is the set of all  $a(x, \xi) \in L_{\text{loc}}^\infty(\mathbb{R}^d \times (\mathbb{R}^d \setminus \{0\}))$  such that

- (1) for all  $x \in \mathbb{R}^d$  the function  $\xi \mapsto a(x, \xi)$  is smooth, and
- (2) for all  $\alpha \in \mathbb{N}^d$  there exists  $C_\alpha > 0$ , such that

$$\|\partial_\xi^\alpha a(\cdot, \xi)\|_{W_{k,\delta}^{r,\infty}} \leq C_\alpha \langle \xi \rangle^{m-|\alpha|} \quad \text{for all } |\xi| \geq \frac{1}{2}.$$

Moreover, we define

$$M_{k,\delta}^{m,r}(a) = \sup_{|\alpha| \leq r + \tilde{d}} \sup_{|\xi| \geq 1/2} \langle \xi \rangle^{|\alpha| - m} \|\partial_\xi^\alpha a(\cdot, \xi)\|_{W_{k,\delta}^{r,\infty}}.$$

Let  $\Gamma_{k,\delta}^{-\infty,r} = \bigcap_{m \in \mathbb{R}} \Gamma_{k,\delta}^{m,r}$  and  $\Gamma_{-\infty,\delta}^{m,r} = \bigcap_{k \in \mathbb{R}} \Gamma_{k,\delta}^{m,r}$ . Then for  $(m, k) \in (\mathbb{R} \cup \{-\infty\})^2$ , define

$$\Sigma_{k,\delta}^{m,r} = \sum_{0 \leq j \leq r} \Gamma_{k-\delta j,\delta}^{m-j,r-j}.$$

We say that  $a_h = \sum_{0 \leq j \leq r} h^j a_h^j \in {}_h\Sigma_{k,\delta}^{m,r}$  if

$$\sup_{0 < h < 1} \sum_{0 \leq j \leq r} M_{k-\delta j,\delta}^{m-j,r-j}(a_h^j) < \infty.$$

We shall define  $\Sigma^{m,r} = \Sigma_{0,0}^{m,r}$  and  ${}_h\Sigma^{m,r} = {}_h\Sigma_{0,0}^{m,r}$ .

We are mostly interested in the cases where  $\delta \in \{0, 1\}$ . Note that  $W_{k,0}^{r,\infty} = \langle x \rangle^k W^{r,\infty}$  and thus  $\Gamma_{k,0}^{m,r} = \langle x \rangle^k \Gamma^{m,r}$ , whereas  $\Gamma_{k,1}^{m,r}$  is a natural extension of  $S_k^m$  to symbols of finite regularities. We will encounter symbols defined by solutions of the water wave system and thus have coefficients in weighted Sobolev spaces. We need the following lemma.

**Lemma 4.15.** *If  $u \in \mathcal{H}_k^{\mu,\delta}$ , where  $\mu \geq \tilde{d}$ ,  $k \in \mathbb{N}$  and  $\delta \in (0, 1]$ , then, for all  $\alpha \in \mathbb{N}^d$  with  $|\alpha| \leq \min\{(\mu - \tilde{d})/(1 + \delta), k\}$ , we have  $\langle x \rangle^{|\alpha|} \partial_x^\alpha u \in L^\infty$  and consequently we have the inclusion*

$$\mathcal{H}_k^{\mu,\delta} \subset W_{0,1}^{\min\{[(\mu - \tilde{d})/(1 + \delta)], k\}, \infty} \cap \langle x \rangle^{-\min\{[\mu - \tilde{d}]/\delta, k\}} L^\infty.$$

In particular  $\mathcal{H}_k^{\mu,1/2} \subset W_{0,1}^{\min\{[2(\mu - \tilde{d})/3], k\}, \infty} \cap \langle x \rangle^{-\min\{k, 2[\mu - \tilde{d}]\}} L^\infty$ .

*Proof.* The lemma follows directly from the Sobolev injection:

$$\begin{aligned} \|\langle x \rangle^{|\alpha|} \partial_x^\alpha u\|_{L^\infty} &\lesssim \|u\|_{W_{-|\alpha|}^{|\alpha|,\infty}} \lesssim \|u\|_{H_{|\alpha|}^{|\alpha| + \tilde{d}}} \lesssim \|u\|_{\mathcal{H}_k^{\mu,\delta}}, \\ \|\langle x \rangle^n u\|_{L^\infty} &\lesssim \|u\|_{W_{-n,0}^{0,\infty}} \lesssim \|u\|_{H_n^{\tilde{d}}} \lesssim \|u\|_{\mathcal{H}_k^{\mu,\delta}}, \end{aligned}$$

which hold provided  $|\alpha| + \tilde{d} \leq \mu - \delta|\alpha|$ ,  $|\alpha| \leq k$ ,  $\tilde{d} \leq \mu - \delta n$  and  $n \leq k$ , that is,

$$|\alpha| \leq \min\left\{\frac{\mu - \tilde{d}}{1 + \delta}, k\right\}, \quad n \leq \min\left\{\frac{\mu - \tilde{d}}{\delta}, k\right\}. \quad \square$$

**Lemma 4.16.** *Let  $\mathcal{A}$  be a linear operator from  $\mathcal{S}(\mathbb{R}^d)$  to  $\mathcal{S}'(\mathbb{R}^d)$  and let  $m, k \in \mathbb{R}$ . If there exists  $\{A_j\}_{j \in \mathbb{N}} \in \ell^\infty(\mathcal{O}_0^m)$  and  $\psi, \phi \in \mathcal{P}$  such that  $\mathcal{A} = \sum_{j \in \mathbb{N}} 2^{jk} \psi_j \mathcal{A}_j \phi_j$ , then  $\mathcal{A} \in \mathcal{O}_k^m$ .*

*Proof.* The lemma is a consequence of [Proposition 2.12](#). □

**Definition 4.17.** Let  $\psi \in \mathcal{P}_*$  and define  $\underline{\psi} \in \mathcal{P}$  by setting  $\underline{\psi}_j = \sum_{|j-k| \leq 10} \psi_k$ . If  $a \in \Gamma_{k,\delta}^{m,r}$ , where  $m, k \in \mathbb{R}$ ,  $r \in \mathbb{N}$  and  $\delta \in [0, 1]$ , then define the dyadic paradifferential operator

$$\mathcal{P}_a = \sum_{j \in \mathbb{N}} \underline{\psi}_j T_{\psi_j a} \underline{\psi}_j.$$

**Proposition 4.18.** *If  $a \in \Gamma_{k,\delta}^{m,r}$ , then  $\mathcal{P}_a = \mathcal{O}(M_k^{m,r}(a))_{\mathcal{O}_k^m}$ .*

*Proof.* Note that if  $a \in \Gamma_{k,\delta}^{m,r}$  then  $a \in \langle x \rangle^k \Gamma^{m,0}$ . Therefore, by [Proposition 4.4](#), we have

$$\|T_{\psi_j a}\|_{H^v \rightarrow H^{v-m}} \lesssim M^{m,0}(\psi_j a) \lesssim 2^{-jk} M_{k,0}^{m,0}(a).$$

We conclude by [Lemma 4.16](#). □

**4B2. Symbolic calculus.**

**Proposition 4.19.** *Let  $a \in \Gamma_{k,\delta}^{m,r}$ ,  $b \in \Gamma_{k',\delta}^{m',r}$ ,  $r \in \mathbb{N}$ ,  $(m, k), (m', k') \in \mathbb{R}^2$ ,  $0 \leq \delta \leq 1$ , then*

$$\mathcal{P}_a \mathcal{P}_b - \mathcal{P}_{a \sharp b} = \mathcal{O}(M_{k,\delta}^{m,r}(a) M_{k',\delta}^{m',r}(b))_{\mathcal{O}_{k+k'-\delta r}^{m+m'-r} + \mathcal{O}_{k+k'}^{-\infty}},$$

where

$$a \sharp b = \sum_{\alpha \in \mathbb{N}^d} \frac{|\alpha| < r}{\alpha!} \frac{1}{\partial_{\xi}^{\alpha}} a D_x^{\alpha} b \in \Sigma_{k+k',\delta}^{m+m',r}.$$

*Proof.* Let  $\tilde{\psi}_j : \mathbb{N} \rightarrow C_c^{\infty}$ ,  $\tilde{\psi}_j = \sum_{|j-j'| \leq 50} \psi_{j'}$ , so  $\underline{\psi}_{j'} \tilde{\psi}_j = \underline{\psi}_{j'}$  if  $|j-j'| \leq 20$ . Then write

$$\mathcal{P}_a \mathcal{P}_b = \sum_{(j,j') \in \mathbb{N}^2} \sum_{|j-j'| \leq 20} \underline{\psi}_j T_{\psi_j a} \underline{\psi}_{j'} \psi_{j'} T_{\psi_{j'} b} \underline{\psi}_{j'} = \sum_{(j,j') \in \mathbb{N}^2} \sum_{|j-j'| \leq 20} \tilde{\psi}_j T_{\psi_j a} T_{\psi_{j'} b} \tilde{\psi}_j + \tilde{\psi}_j R_{j,j'} \tilde{\psi}_j,$$

the remainder being

$$\begin{aligned} R_{j,j'} &= \underline{\psi}_j T_{\psi_j a} \underline{\psi}_{j'} \psi_{j'} T_{\psi_{j'} b} \underline{\psi}_{j'} - T_{\psi_j a} T_{\psi_{j'} b} \\ &= \mathcal{O}(2^{j(k+k'-\delta r)} M_{k,\delta}^{m,r}(a) M_{k',\delta}^{m',r}(b))_{\mathcal{O}_0^{m+m'-r}} + \mathcal{O}(2^{j(k+k')} M_{k,\delta}^{m,r}(a) M_{k',\delta}^{m',r}(b))_{\mathcal{O}_0^{-\infty}} \end{aligned}$$

by [Propositions 4.4](#) and [4.9](#) and [Corollary 4.8](#). More precisely, when composing  $T_{\psi_j a}$  and  $T_{\psi_{j'} b}$ , we use  $\psi_{j'} \underline{\psi}_j = \underline{\psi}_j$  and have

$$\begin{aligned} T_{\psi_j a} T_{\psi_{j'} b} &= T_{\psi_j a} + \mathcal{O}(M^{m,r}(\psi_j a) M^{0,0}(\nabla^r \underline{\psi}_{j'}))_{\mathcal{O}_0^{m-r}} \\ &\quad + \mathcal{O}(M^{m,0}(\nabla^r(\psi_j a)) M^{0,r}(\underline{\psi}_{j'}))_{\mathcal{O}_0^{m-r}} + \mathcal{O}(M^{m,r}(\psi_j a) M^{0,r}(\underline{\psi}_{j'}))_{\mathcal{O}_0^{-\infty}}, \end{aligned}$$

where  $M^{0,0}(\nabla^r \underline{\psi}_{j'}) = \mathcal{O}(2^{-jr})$ ,  $M^{m,r}(\psi_j a) = \mathcal{O}(2^{jk})$ , and we use  $0 \leq \delta \leq 1$  to induce that

$$M^{m,0}(\nabla^r(\psi_j a)) = \mathcal{O}\left(\max_{0 \leq n \leq r} \{2^{-j(r-n)+j(k-\delta n)}\}\right) = \mathcal{O}(2^{j(k-\delta r)}).$$

Similar arguments work for the composition  $T_{\psi_j} T_{\psi_j a}$ .

Observe that  $\sum_{j':|j-j'| \leq 20} (\psi_j a) \sharp \psi_{j'} b = (\psi_j a) \sharp b$ , for all  $j \in \mathbb{N}$ . Hence

$$\sum_{j':|j-j'| \leq 20} T_{\psi_j a} T_{\psi_{j'} b} = \underline{\psi}_j T_{\psi_j a \sharp b} \underline{\psi}_j + R_j,$$

where the remainder can be estimated much as above:

$$R_j = \mathcal{O}(2^{j(k+k'-\delta r)} M_{k,\delta}^{m,r}(a) M_{k',\delta}^{m',r}(b))_{\mathcal{O}_0^{m+m'-\delta r}} + \mathcal{O}(2^{j(k+k')} M_{k,\delta}^{m,r}(a) M_{k',\delta}^{m',r}(b))_{\mathcal{O}_0^{-\infty}}.$$

We conclude by [Lemma 4.16](#). □

**Proposition 4.20.** *Let  $a \in \Gamma_{k,\delta}^{m,r}$  with  $(m, k) \in \mathbb{R}^2$ , and  $r \in \mathbb{N}$ ,  $0 \leq \delta \leq 1$ , then*

$$\mathcal{P}_a^* - \mathcal{P}_{a^*} = \mathcal{O}(M_{k,\delta}^{m,r}(a))_{\mathcal{O}_{k-\delta r}^{m-r} + \mathcal{O}_k^{-\infty}}, \tag{4-8}$$

where

$$a^* = \sum_{\alpha \in \mathbb{N}^d} \frac{1}{\alpha!} \partial_\xi^\alpha D_x^\alpha \bar{a} \in \Sigma_{k,\delta}^{m,r}.$$

*Proof.* Observe that for any real-valued  $\psi \in C_c^\infty(\mathbb{R}^d)$ ,

$$(\psi a)^* = a^* \sharp \psi. \quad (4-9)$$

More precisely, this means that

$$\begin{aligned} (\psi a)^* &= \sum_{|\gamma| < r} \frac{1}{\gamma!} \partial_\xi^\gamma D_x^\gamma (\psi \bar{a}) = \sum_{|\gamma| < r} \frac{1}{\gamma!} \sum_{\alpha + \beta = \gamma} \frac{\gamma!}{\alpha! \beta!} D_x^\alpha \psi \partial_\xi^\beta D_x^\beta \bar{a} \\ &= \sum_{|\alpha| + |\beta| < r} \frac{1}{\alpha!} \partial_\xi^\alpha \left( \frac{1}{\beta!} \partial_\xi^\beta D_x^\beta \bar{a} \right) D_x^\alpha \psi = \sum_{|\beta| < r} \sum_{|\alpha| < r - |\beta|} \frac{1}{\alpha!} \partial_\xi^\alpha \left( \frac{1}{\beta!} \partial_\xi^\beta D_x^\beta \bar{a} \right) D_x^\alpha \psi \\ &= \sum_{|\beta| < r} \left( \frac{1}{\beta!} \partial_\xi^\beta D_x^\beta \bar{a} \right) \sharp \psi = a^* \sharp \psi. \end{aligned}$$

Then write  $\mathcal{P}_a^* - \mathcal{P}_{a^*} = \sum_{j \in \mathbb{N}} \underline{\psi}_j (R_j^1 + R_j^2) \underline{\psi}_j$ , where, by (4-9),

$$R_j^1 = T_{\psi_j a}^* - T_{(\psi_j a)^*}, \quad R_j^2 = T_{(\psi_j a)^*} - T_{\psi_j a^*} = T_{a^* \sharp \psi_j - \psi_j a^*}.$$

For  $R_j^1$  we use Proposition 4.10,

$$R_j^1 = \mathcal{O}(M^{m,0}(\nabla_x(\psi_j a)))_{\mathcal{O}_0^{m-r}} = \mathcal{O}(2^{j(k-\delta r)} M_{k,\delta}^{m,r}(a))_{\mathcal{O}_0^{m-r}}.$$

By Lemma 4.16,

$$\sum_{j \in \mathbb{N}} \underline{\psi}_j R_j^1 \underline{\psi}_j = \mathcal{O}(M_{k,\delta}^{m,r}(a))_{\mathcal{O}_{k-\delta r}^{m-r} + \mathcal{O}_k^{-\infty}}.$$

Using  $\sum_{j \in \mathbb{N}} \underline{\psi}_j \equiv 1$ , we induce that

$$\sum_{j \in \mathbb{N}} \partial_x^\alpha \underline{\psi}_j \equiv 0 \quad \text{for all } \alpha \in \mathbb{N}^d \setminus \{0\}, \quad \sum_{j \in \mathbb{N}} a^* \sharp \psi_j - \psi_j a^* = 0. \quad (4-10)$$

Then we write

$$a^* \sharp \psi_j - \psi_j a^* = \sum_{\substack{\alpha \neq 0 \\ |\alpha| + |\beta| < r}} D_x^\alpha \psi_j \cdot w_{\alpha\beta}, \quad w_{\alpha\beta} \in \Gamma_{k-|\beta|\delta,\delta}^{m-|\alpha|-|\beta|,r-|\beta|},$$

where the symbols  $w_{\alpha\beta}$  are independent of  $j$ . Write

$$\sum_{j \in \mathbb{N}} \underline{\psi}_j R_j^2 \underline{\psi}_j = \sum_{\alpha,\beta} R_{\alpha\beta}, \quad R_{\alpha\beta} = \sum_{j \in \mathbb{N}} \underline{\psi}_j T_{D_x^\alpha \psi_j \cdot w_{\alpha\beta}} \underline{\psi}_j.$$

By (4-10), we prove similarly to Proposition 4.19 that

$$\begin{aligned} \psi_j R_{\alpha\beta} &= \psi_j \sum_{|j-j'| \leq 20} \underline{\psi}_{j'} T_{D_x^\alpha \psi_{j'} \cdot w_{\alpha\beta}} \underline{\psi}_{j'} \\ &= \mathcal{O}(2^{j(-|\alpha|+k-|\beta|\delta-(r-|\beta|)\delta)}) M_{k-|\beta|\delta,\delta}^{m-|\alpha|-|\beta|,r-|\beta|}(w_{\alpha\beta})_{\mathcal{O}_0^{m-r-|\alpha|}} \\ &\quad + \mathcal{O}(2^{j(-|\alpha|+k-|\beta|\delta)}) M_{k-|\beta|\delta,\delta}^{m-|\alpha|-|\beta|,r-|\beta|}(w_{\alpha\beta})_{\mathcal{O}_0^{-\infty}} \\ &= \mathcal{O}(2^{j(k-\delta r)}) M_{k,\delta}^{m,r}(a)_{\mathcal{O}_0^{m-r}} + \mathcal{O}(2^{jk}) M_{k,\delta}^{m,r}(a)_{\mathcal{O}_0^{-\infty}}. \end{aligned}$$

Setting  $\psi'_j = \sum_{|j'-j| \leq 100} \psi_{j'}$ . We again conclude by [Lemma 4.16](#), and the identity

$$R_{\alpha\beta} = \sum_{j \in \mathbb{N}} \psi_j R_{\alpha\beta} \psi'_j,$$

that  $R_{\alpha\beta} = \mathcal{O}(M_{k,\delta}^{m,r}(a))_{\mathcal{O}_{k-\delta r}^{m-r} + \mathcal{O}_k^{-\infty}}$ . □

### 4B3. Parilinearization.

**Proposition 4.21.** *If  $a \in H_k^\alpha$ ,  $b \in H_\ell^\beta$  with  $\alpha > \frac{d}{2}$ ,  $\beta > \frac{d}{2}$ ,  $k \in \mathbb{R}$ ,  $\ell \in \mathbb{R}$ , then, for all  $\epsilon > 0$ ,*

$$\|ab - \mathcal{P}_a b - \mathcal{P}_b a\|_{H_{k+\ell}^{\alpha+\beta-d/2-\epsilon}} \lesssim \|a\|_{H_k^\alpha} \|b\|_{H_\ell^\beta}.$$

*Consequently if  $a \in \mathcal{H}_m^{\alpha,\delta}$  and  $b \in \mathcal{H}_n^{\beta,\delta}$  with  $\delta \geq 0$ ,  $\alpha - \delta m > \frac{d}{2}$ ,  $\beta - \delta n > \frac{d}{2}$ , then, for all  $\epsilon > 0$ ,*

$$\|ab - \mathcal{P}_a b - \mathcal{P}_b a\|_{\mathcal{H}_{m+n}^{\alpha+\beta-d/2-\epsilon,\delta}} \lesssim \|a\|_{\mathcal{H}_m^{\alpha,\delta}} \|b\|_{\mathcal{H}_n^{\beta,\delta}}.$$

*Proof.* Decompose the product  $ab$  as

$$ab = \sum_{j \in \mathbb{N}} \psi_j (\underline{\psi}_j a) (\underline{\psi}_j b) = \mathcal{P}_a b + \mathcal{P}_b a + R_j^1 + R_j^2,$$

where the remainders  $R_j^1$  and  $R_j^2$  are defined by

$$\begin{aligned} R_j^1 &= \psi_j (\underline{\psi}_j a \cdot \underline{\psi}_j b - T_{\underline{\psi}_j a}(\underline{\psi}_j b) - T_{\underline{\psi}_j b}(\underline{\psi}_j a)), \\ R_j^2 &= \underline{\psi}_j (\psi_j T_{\underline{\psi}_j a} - T_{\psi_j a}) \underline{\psi}_j b + \underline{\psi}_j (\psi_j T_{\underline{\psi}_j b} - T_{\psi_j b}) \underline{\psi}_j a. \end{aligned}$$

By [Proposition 4.11](#),

$$\|R_j^1\|_{H^{\alpha+\beta-d/2}} \lesssim \|\underline{\psi}_j a\|_{H^\alpha} \|\underline{\psi}_j b\|_{H^\beta} \lesssim 2^{-j(k+\ell)} \|a\|_{H_k^\alpha} \|b\|_{H_\ell^\beta}.$$

By [Proposition 4.9](#) and [Corollary 4.8](#) and the Sobolev embedding theorem, for all  $\epsilon > 0$  we have

$$\begin{aligned} \psi_j T_{\underline{\psi}_j a} \underline{\psi}_j - T_{\psi_j a} &= 2^{-jk} \mathcal{O}(\|a\|_{H_k^\alpha}) \mathcal{O}_0^{\alpha-d/2-\epsilon}, \\ \psi_j T_{\underline{\psi}_j b} \underline{\psi}_j - T_{\psi_j b} &= 2^{-j\ell} \mathcal{O}(\|b\|_{H_\ell^\beta}) \mathcal{O}_0^{\beta-d/2-\epsilon}. \end{aligned}$$

We conclude the first statement by [Proposition 2.12](#).

As for the second statement, observe that if  $0 \leq k \leq m$  and  $0 \leq \ell \leq n$ , then

$$\|ab - \mathcal{P}_a b - \mathcal{P}_b a\|_{H_{k+\ell}^{(\alpha-\delta k)+(\beta-\delta \ell)-d/2-\epsilon,\delta}} \lesssim \|a\|_{H_k^{\alpha-\delta k}} \|b\|_{H_\ell^{\beta-\delta \ell}} \lesssim \|a\|_{\mathcal{H}_m^{\alpha,\delta}} \|b\|_{\mathcal{H}_n^{\beta,\delta}}.$$

We conclude by noting that for all  $p \in \mathbb{N} \cap [0, m+n]$ , there exist  $k \in \mathbb{N} \cap [0, m]$  and  $\ell \in \mathbb{N} \cap [0, n]$  such that  $p = k + \ell$ . □

**Proposition 4.22.** *Let  $F \in C^\infty(\mathbb{R})$  with  $F(0) = 0$ . For all  $\mu > \frac{d}{2}$ , there exists some monotonically increasing function  $C : \mathbb{R}_+ \rightarrow \mathbb{R}_+$  such that, for all  $k \geq 0$  and all  $u \in H_k^\mu$ , we have*

$$\|F(u)\|_{H_k^\mu} + \|F(u) - \mathcal{P}_{F'(u)} u\|_{H_k^{2\mu-d/2}} \leq C(\|u\|_{H^\mu}) \|u\|_{H_k^\mu}.$$

*Consequently, if  $u \in \mathcal{H}_k^{\mu,\delta}$  with  $\delta \geq 0$  and  $\mu - \delta k > \frac{d}{2}$ , then*

$$\|F(u)\|_{\mathcal{H}_k^{\mu,\delta}} + \|F(u) - \mathcal{P}_{F'(u)} u\|_{\mathcal{H}_k^{2\mu-d/2,2\delta}} \leq C(\|u\|_{H^\mu}) \|u\|_{\mathcal{H}_k^{\mu,\delta}}.$$

*Proof.* Decompose  $F(u)$  as  $F(u) = \sum_{j \geq 0} \psi_j F(\psi_j u)$ . By [Proposition 4.12](#),

$$\|F(\psi_j u)\|_{H^\mu} \leq C(\|\psi_j u\|_{H^\mu}) \|\psi_j u\|_{H^\mu} \leq C(\|u\|_{H^\mu}) \|\psi_j u\|_{H^\mu}.$$

Then write  $\psi_j F(\psi_j u) = \psi_j T_{\psi_j F'(u)} \psi_j u + \psi_j R_j$ , where

$$R_j = \psi_j(F(\psi_j u) - T_{F'(\psi_j u)} \psi_j u) + \psi_j(\psi_j T_{F'(\psi_j u)} - T_{\psi_j F'(\psi_j u)}) \psi_j u.$$

By [Propositions 4.12](#) and [4.11](#) and [Corollary 4.8](#),

$$\|R_j\|_{H^{2\mu-d/2}} \leq C(\|u\|_{H^\mu}) \|\psi_j u\|_{H^\mu}.$$

We conclude the first statement with [Proposition 2.12](#). To prove the second statement, note that for all  $j \in \mathbb{N} \cap [0, k]$  we have

$$\|F(u)\|_{H_j^{\mu-\delta j}} + \|F(u) - \mathcal{P}_{F'(u)} u\|_{H_j^{2(\mu-\delta j)-d/2}} \leq C(\|u\|_{H^{\mu-\delta j}}) \|u\|_{H_j^{\mu-\delta j}}. \quad \square$$

**4C. Semiclassical paradifferential calculus.** We develop a semiclassical dyadic paradifferential calculus and a quasihomogeneous semiclassical paradifferential calculus, using scaling arguments inspired by [\[Métivier and Zumbrun 2005\]](#).

**4C1. Semiclassical paradifferential operators.**

**Definition 4.23.** For all  $h \in (0, 1]$ , define the scaling operator

$$\tau_h : u(\cdot) \mapsto h^{d/2} u(h \cdot). \quad (4-11)$$

(1) If  $b \in \Gamma^{m,r}$ , then define  $T_b^h = \tau_h^{-1} T_{\theta_{h,*}^{1,0} b} \tau_h$ .

(2) If  $a \in \Gamma_{k,\delta}^{m,r}$ , then define  $\mathcal{P}_a^h = \sum_{j \in \mathbb{N}} \psi_j T_{\psi_j a}^h \psi_j$ .

(3) If  $\epsilon \geq 0$ , then define  $\mathcal{P}_a^{h,\epsilon} = \mathcal{P}_{\theta_{h,*}^{\epsilon,0} a}$ .

**Proposition 4.24.** If  $\epsilon \geq 0$  and  $a \in \Gamma_{k,0}^{m,0}$ , where  $m \leq 0, k \leq 0$ , then  $\sup_{h \in (0,1]} \|\mathcal{P}_a^{h,\epsilon}\|_{L^2 \rightarrow L^2} < \infty$ .

*Proof.* Observe that  $\theta_{h,*}^{1+\epsilon,0} a = \mathcal{O}(1)_{\Gamma^{0,0}}$ . We conclude with [Lemma 4.16](#). □

**4C2. Semiclassical symbolic calculus.**

**Definition 4.25.** If  $a_h \in \mathcal{D}'(\mathbb{R}^{2d})$  and  $\epsilon \geq 0$ , we say that  $a_h \in \sigma_\epsilon$  if

$$\bigcup_{0 < h < 1} \text{supp } a_h \cap \mathcal{N}_{\epsilon,1} = \emptyset.$$

**Proposition 4.26.** Let  $(m, k), (m', k') \in (\mathbb{R} \cup \{-\infty\})^2, r \in \mathbb{N}$ , with  $r \geq m + m', \delta r \geq k + k'$ . Let  $a_h \in \Gamma_{k,\delta}^{m,r} \cap \sigma_0$  and  $b_h \in \Gamma_{k',\delta}^{m',r} \cap \sigma_0$  such that, for some  $R_h \geq 0$  depending on  $h$ ,

$$\text{supp } a_h \cap \text{supp } b_h \subset \{|x| \geq R_h\} \times \mathbb{R}^d. \quad (4-12)$$

Then, for  $h > 0$  sufficiently small,

$$\mathcal{P}_{a_h}^h \mathcal{P}_{b_h}^h - \mathcal{P}_{a_h \sharp_h b_h}^h = \mathcal{O}(h^r (1 + R_h)^{k+k'-\delta r})_{L^2 \rightarrow L^2},$$

where the symbol  $a_h \sharp_h b_h = a_h \sharp_{h,r}^{0,1} b_h \in {}_h \Sigma_{k+k',\delta}^{m+m',r}$  is defined by [\(2-4\)](#).

*Proof.* By (4-12), if  $\psi_j a_h \neq 0$  and  $\psi_j b_h \neq 0$ , then  $j \gtrsim \log_2(1 + R_h)$ . We claim that

$$\begin{aligned} \mathcal{P}_a^h \mathcal{P}_b^h &= \sum_{\substack{j \gtrsim \log_2(1+R_h) \\ |j'-j| \leq 20}} \underline{\psi}_j T_{\psi_j a_h}^h \underline{\psi}_{j'} \underline{\psi}_{j'} T_{\psi_{j'} b_h}^h \underline{\psi}_{j'} \\ &= \sum_{\substack{j \gtrsim \log_2(1+R_h) \\ |j'-j| \leq 20}} \underline{\psi}_j T_{(\psi_j a_h) \sharp_h (\psi_{j'} b_h)}^h \underline{\psi}_{j'} + \mathcal{O}(h^r (1 + R_h)^{k+k'-\delta r})_{L^2 \rightarrow L^2}. \end{aligned} \quad (4-13)$$

Then we conclude by

$$\sum_{j': |j'-j| \leq 20} (\psi_j a_h) \sharp_h (\psi_{j'} b_h) = \psi_j (a_h \sharp_h b_h).$$

It remains to prove (4-13). We use (4-1) to deduce that  $\mathcal{F}(T_{\theta_{h,*}^{1,0}(\psi_{j'} b_h)} u)$  vanishes in a neighborhood of  $\xi = 0$ . By (4-5), for some  $\pi' \in C^\infty(\mathbb{R}^d)$  which vanishes near  $\xi = 0$  and equals 1 outside a neighborhood of  $\xi = 0$ , and, for all  $m + m' \leq N \in \mathbb{N}$ ,

$$\begin{aligned} \tau_h T_{\psi_j a_h}^h \underline{\psi}_j \underline{\psi}_{j'} T_{\psi_{j'} b_h}^h \tau_h^{-1} &= T_{\theta_{h,*}^{1,0}(\psi_j a_h)} \theta_{h,*}^{1,0}(\underline{\psi}_j \underline{\psi}_{j'}) \pi'(D_x) T_{\theta_{h,*}^{1,0}(\psi_{j'} b_h)} \\ &= T_{\theta_{h,*}^{1,0}(\psi_j a_h)} T_{\theta_{h,*}^{1,0}(\underline{\psi}_j \underline{\psi}_{j'})} \otimes \pi' T_{\theta_{h,*}^{1,0}(\psi_{j'} b_h)} \\ &\quad + \mathcal{O}(M^{m,0}(\psi_j a_h))_{\mathcal{O}_0^m} \mathcal{O}(2^{-jN} h^N)_{\mathcal{O}_0^{-N}} \mathcal{O}(M^{m',0}(\psi_{j'} b_h))_{\mathcal{O}_0^{m'}}. \end{aligned} \quad (4-14)$$

Then we use Proposition 4.9 and the fact that  $a_h, b_h \in \mathcal{S}_0$  to deduce

$$\begin{aligned} T_{\theta_{h,*}^{1,0}(\psi_j a_h)} T_{\theta_{h,*}^{1,0}(\underline{\psi}_j \underline{\psi}_{j'})} \otimes \pi' T_{\theta_{h,*}^{1,0}(\psi_{j'} b_h)} \\ &= T_{\theta_{h,*}^{1,0}(\psi_j a_h) \sharp_{h,*}^{1,0}(\underline{\psi}_j \underline{\psi}_{j'} \otimes \pi')} \sharp_{h,*}^{1,0}(\psi_{j'} b_h) \\ &\quad + \mathcal{O}(M^{m,0}(\nabla^r \theta_{h,*}^{1,0}(\psi_j a_h))) M^{0,r}(\theta_{h,*}^{1,0}(\psi_j \psi_{j'})) M^{m',r}(\theta_{h,*}^{1,0}(\psi_{j'} b_h))_{\mathcal{O}_0^{m+m'-r}} \\ &\quad + \mathcal{O}(M^{m,r}(\theta_{h,*}^{1,0}(\psi_j a_h))) M^{0,0}(\nabla^r \theta_{h,*}^{1,0}(\psi_j \psi_{j'})) M^{m',r}(\theta_{h,*}^{1,0}(\psi_{j'} b_h))_{\mathcal{O}_0^{m+m'-r}} \\ &\quad + \mathcal{O}(M^{m,r}(\theta_{h,*}^{1,0}(\psi_j a_h))) M^{0,r}(\theta_{h,*}^{1,0}(\psi_j \psi_{j'})) M^{m',0}(\nabla^r \theta_{h,*}^{1,0}(\psi_{j'} b_h))_{\mathcal{O}_0^{m+m'-r}}. \end{aligned}$$

To estimate the remainders, we see that, for each  $\alpha \in \mathbb{N}^d$  with  $|\alpha| = r$ ,

$$\begin{aligned} \partial_x^\alpha \theta_{h,*}^{1,0}(\psi_j a_h) &= \sum_{\alpha_1 + \alpha_2 = \alpha} \frac{\alpha!}{\alpha_1! \alpha_2!} \partial_x^{\alpha_1} \theta_{h,*}^{1,0} \psi_j \partial_x^{\alpha_2} \theta_{h,*}^{1,0} a_h \\ &= \sum_{\alpha_1 + \alpha_2 = \alpha} \frac{\alpha!}{\alpha_1! \alpha_2!} \mathcal{O}(h^{|\alpha_1|} 2^{-j|\alpha_1|} \times h^{|\alpha_2|} 2^{j(k-\delta|\alpha_2|)})_{L^\infty} = \mathcal{O}(h^r 2^{j(k-\delta r)})_{L^\infty}, \end{aligned}$$

where we use  $0 \leq \delta \leq 1$ . Therefore, the first term in the remainder is

$$\mathcal{O}(h^r 2^{j(k+k'-\delta r)})_{L^2 \rightarrow L^2} = \mathcal{O}(h^r (1 + R_h)^{k+k'-\delta r})_{L^2 \rightarrow L^2}.$$

Similar methods apply to the other two terms and we conclude that

$$T_{\theta_{h,*}^{1,0}(\psi_j a_h)} T_{\theta_{h,*}^{1,0}(\underline{\psi}_j \underline{\psi}_{j'})} \otimes \pi' T_{\theta_{h,*}^{1,0}(\psi_{j'} b_h)} = T_{\theta_{h,*}^{1,0}((\psi_j a_h) \sharp_h (\psi_{j'} b_h))} + \mathcal{O}(h^r (1 + R_h)^{k+k'-\delta r})_{L^2 \rightarrow L^2}. \quad (4-15)$$

The estimate (4-13) follows from (4-14) and (4-15).  $\square$

Combining the analysis of Propositions 4.26 and 4.20, using Proposition 4.9, we obtain a similar result for the adjoint, to the proof of which we shall omit, as it is similar to the above.

**Proposition 4.27.** *Let  $(m, k) \in (\mathbb{R} \cup \{-\infty\})^2$ ,  $r \in \mathbb{N}$ , with  $r \geq m$ ,  $\delta r \geq k$ . Let  $a_h \in \Gamma_{k,\delta}^{m,r} \cap \sigma_0$ , such that, for some  $R_h \geq 0$  depending on  $h$ ,  $\text{supp } a_h \subset \{|x| \geq R_h\} \times \mathbb{R}^d$ , then, for  $h > 0$  sufficiently small,*

$$(\mathcal{P}_{a_h}^h)^* - \mathcal{P}_{a_h^*}^h = \mathcal{O}(h^r(1 + R_h)^{k-\delta r})_{L^2 \rightarrow L^2},$$

where  $a_h^* = \zeta_{h,r}^{0,1} a_h \in {}_h\Sigma_{k,\delta}^{m,r}$  is defined by (2-5).

**Corollary 4.28.** *Let  $\epsilon \geq 0$ ,  $(m, k), (m', k') \in (\mathbb{R} \cup \{-\infty\})^2$ ,  $r \in \mathbb{N}$ , with  $r \geq \max\{m + m', k'\}$ ,  $k \leq 0$ . If  $a_h \in \Gamma_{k,1}^{m,r} \cap \sigma_\epsilon$  and  $b_h \in \Gamma_{k',1}^{m',r} \cap \sigma_0$ , then*

$$\begin{aligned} \mathcal{P}_{a_h}^{h,\epsilon} \mathcal{P}_{b_h}^h - \mathcal{P}_{(\theta_{h,*}^{\epsilon,0} a_h) \sharp_h b_h}^h &= \mathcal{O}(h^{(1+\epsilon)r - \epsilon(k+k')})_{L^2 \rightarrow L^2}, \\ \mathcal{P}_{b_h}^h \mathcal{P}_{a_h}^{h,\epsilon} - \mathcal{P}_{b_h \sharp_h (\theta_{h,*}^{\epsilon,0} a_h)}^h &= \mathcal{O}(h^{(1+\epsilon)r - \epsilon(k+k')})_{L^2 \rightarrow L^2}. \end{aligned}$$

*Proof.* It suffices to observe that, if  $\epsilon > 0$  then  $\text{supp } \theta_{h,*}^{\epsilon,0} a_h \subset \{|x| \gtrsim h^{-\epsilon}\}$  and  $\theta_{h,*}^{\epsilon,0} a_h = \mathcal{O}(h^{-\epsilon k})_{\Gamma_{0,1}^{m,r}}$ . We conclude by Proposition 4.26. □

**Corollary 4.29.** *Let  $\epsilon \geq 0$ ,  $(m, k), (m', k') \in (\mathbb{R} \cup \{-\infty\})^2$ ,  $r \in \mathbb{N}$ , with  $r \geq m + m'$ ,  $k \leq 0$ ,  $k' \leq 0$ . If  $a_h \in \Gamma_{k,1}^{m,r} \cap \sigma_\epsilon$  and  $b_h \in \Gamma_{k',1}^{m',r} \cap \sigma_\epsilon$ , then, for  $h > 0$  sufficiently small,*

$$\mathcal{P}_{a_h}^{h,\epsilon} \mathcal{P}_{b_h}^{h,\epsilon} - \mathcal{P}_{a_h \sharp_h^\epsilon b_h}^{h,\epsilon} = \mathcal{O}(h^{(1+\epsilon)r - \epsilon(k+k')})_{L^2 \rightarrow L^2},$$

where the symbol  $a_h \sharp_h^\epsilon b_h = a_h \sharp_{h,r}^{\epsilon,1} \in h^{1+\epsilon} \Sigma_{k+k',1}^{m+m',r}$  is defined by (2-4).

*Proof.* It suffices to use the identity  $(\theta_{h,*}^{\epsilon,0} a_h) \sharp_h (\theta_{h,*}^{\epsilon,0} b_h) = \theta_{h,*}^{\epsilon,0} (a_h \sharp_h^\epsilon b_h)$ . □

**4C3. Some technical lemmas.** The results above only concerned the high frequency regime as we require the  $\sigma_\epsilon$  condition. The next lemma studies the interaction of high frequencies and low frequencies.

**Lemma 4.30.** *Let  $m \in \mathbb{R}$ ,  $a_h \in \Gamma^{0,0}$ ,  $b_h \in \Gamma^{0,0}$  such that, for some  $R > 0$ ,*

$$\text{supp } a_h \in \{|\xi| \geq R\}, \quad \text{supp } b_h \in \{|\xi| \leq h^{-1}R/4\}.$$

Then  $\mathcal{P}_{a_h}^h \mathcal{P}_{b_h} = \mathcal{O}(h^\infty)_{L^2 \rightarrow L^2}$ .

**Remark 4.31.** This lemma concerns the estimate of  $\mathcal{P}_{a_h}^h \mathcal{P}_{b_h}$ , not  $\mathcal{P}_{a_h}^h \mathcal{P}_{b_h}^h$ . This is not a typo.

*Proof.* By definition

$$\widehat{T_{\psi_j b_h} u}(\xi) = (2\pi)^{-d} \int \chi(\xi - \eta) \pi(\eta) \widehat{\psi_j b_h}(\xi - \eta, \eta) \hat{u}(\eta) \, d\eta.$$

The admissibility of  $\chi$  implies  $\text{supp } \widehat{T_{\psi_j b_h} u} \subset \{|\xi| \leq h^{-1}R/3\}$ . Therefore, for any  $|j' - j| \leq 20$ ,

$$\begin{aligned} \underline{\psi}_j T_{\psi_j a_h}^h \underline{\psi}_j \underline{\psi}_{j'} T_{\psi_{j'} b_h} \underline{\psi}_{j'} &= \underline{\psi}_j T_{\psi_j a_h}^h \pi(hD_x/R) \underline{\psi}_j \underline{\psi}_{j'} (1 - \pi(2hD_x/R)) T_{\psi_{j'} b_h} \underline{\psi}_{j'} \\ &= \underline{\psi}_j \mathcal{O}(h^\infty)_{L^2 \rightarrow L^2} \underline{\psi}_{j'}. \end{aligned}$$

We conclude by Lemma 4.16. □



**Corollary 4.32.** *If  $a \in \Gamma^{m,0}$  is homogeneous of degree  $m$  with respect to  $\xi$ , then, for  $b \in \Gamma^{0,0} \cap \sigma_0$  and  $h > 0$  sufficiently small,*

$$\mathcal{P}_b^h(h^m \mathcal{P}_a - \mathcal{P}_a^h) = \mathcal{O}(h^\infty)_{L^2 \rightarrow L^2}.$$

*Proof.* By a direct verification using (4-2), the homogeneity of  $a$  and the admissible function  $\chi$ , and Corollary 4.7, we see that  $h^m \mathcal{P}_a - \mathcal{P}_a^h = \mathcal{P}'_{\tilde{a}_h}$ , where  $\mathcal{P}'$  denotes the paradifferential quantization with any admissible pair  $(\pi', \chi)$  such that  $\pi \pi' = \pi'$ , and the symbol

$$\tilde{a}_h(x, \xi) = (\pi(h\xi) - \pi(\xi))a(x, h\xi) \in \Gamma^{0,0}$$

satisfies the condition

$$\text{supp } \tilde{a}_h \subset \mathbb{R}^d \times \text{supp}(1 - \pi(h \cdot)) \subset \mathbb{R}^d \times \{|\xi| \leq 2h^{-1}\}.$$

We conclude by Lemma 4.30. □

**Lemma 4.33.** *If  $a_h \in \Gamma^{m,r} \cap \sigma_0$  with  $r \geq \max\{m, 0\} + \tilde{d}$ , then, for  $h > 0$  sufficiently small,*

$$T_{a_h}^h - \text{Op}_h(a_h) = \mathcal{O}(h^r)_{L^2 \rightarrow L^2}.$$

*Proof.* By Calderón–Vaillancourt theorem, we have

$$\begin{aligned} T_{a_h}^h - \text{Op}_h(a_h) &= \tau_h^{-1} (T_{\theta_{h,*}^{1,0} a_h}^{1,0} - \text{Op}(\theta_{h,*}^{1,0} a_h)) \tau_h \\ &= \mathcal{O} \left( \sum_{|\alpha|, |\beta| \leq \tilde{d}} \|\partial_\xi^\alpha \partial_x^\beta (\sigma_{\theta_{h,*}^{1,0} a_h} - \theta_{h,*}^{1,0} a_h)\|_{L^\infty} \right)_{L^2 \rightarrow L^2}. \end{aligned}$$

By hypothesis  $r \geq \max\{m, 0\} + |\beta| \geq |\beta|$ . We use (4-5) to deduce that

$$\begin{aligned} \|\partial_\xi^\alpha \partial_x^\beta (\sigma_{\theta_{h,*}^{1,0} a_h} - \theta_{h,*}^{1,0} a_h)\|_{L^\infty} &\lesssim M^{0,0} (\partial_x^\beta (\sigma_{\theta_{h,*}^{1,0} a_h} - \theta_{h,*}^{1,0} a_h)) \\ &\lesssim M^{\max\{m,0\}-r+|\beta|,0} (\partial_x^\beta (\sigma_{\theta_{h,*}^{1,0} a_h} - \theta_{h,*}^{1,0} a_h)) \\ &\lesssim M^{\max\{m,0\},0} (\nabla_x^r (\theta_{h,*}^{1,0} a_h)) \\ &\lesssim h^r M^{m,0}(a_h). \end{aligned} \quad \square$$

**Lemma 4.34.** *If  $a_h \in \Gamma^{m,\infty} \cap \sigma_0$  with  $m \in \mathbb{R} \cup \{-\infty\}$ , then, for  $h > 0$  sufficiently small,*

$$\mathcal{P}_{a_h}^h - \text{Op}_h(a_h) = \mathcal{O}(h^\infty)_{L^2 \rightarrow L^2}.$$

*Proof.* By Lemmas 4.33 and 4.16,

$$\mathcal{P}_{a_h}^h - \sum_{j \in \mathbb{N}} \underline{\psi}_j \text{Op}_h(\psi_j a_h) \underline{\psi}_j = \mathcal{O}(h^\infty)_{L^2 \rightarrow L^2}.$$

Note that, uniformly in  $j \in \mathbb{N}$ , we have

$$\underline{\psi}_j \sharp_h (\psi_j a_h) \sharp_h \underline{\psi}_j = \underline{\psi}_j a_h + \underline{\psi}_j \mathcal{O}(h^\infty)_{\Gamma^{-\infty,\infty}}.$$

Therefore,

$$\sum_{j \in \mathbb{N}} \underline{\psi}_j \sharp_h (\psi_j a_h) \sharp_h \underline{\psi}_j = a_h + \mathcal{O}(h^\infty)_{\Gamma^{-\infty,\infty}}. \quad \square$$

**4C4.** *Symbols with limited regularities in  $(x, \xi)$ .* The symbols we have encountered so far have limited regularities in the  $x$ -variable but are smooth with respect to the  $\xi$ -variable. When studying the propagation of singularities for nonlinear equations, we need to solve Hamiltonian equations which transfer the limited regularity in the  $x$ -variable to the  $\xi$ -variable. Therefore we need to discuss in this section paradifferential operators with symbols that have limited regularities in both the  $x$ - and  $\xi$ -variables. As we do not intend to obtain optimal regularities, we shall content ourselves with an approach by approximation.

**Definition 4.35.** For all  $r \in \mathbb{N}$ , the symbol class  $\Upsilon^r$  is the set of all  $a \in L^\infty_{\text{loc}}(\mathbb{R}^d \times (\mathbb{R}^d \setminus 0))$  compactly supported in  $\mathbb{R}^d \times (\mathbb{R}^d \setminus 0)$  such that  $N^r(a) < +\infty$ , where

$$N^r(a) = \sum_{\alpha \in \mathbb{N}^{2d}, |\alpha| \leq r} \|\partial_{x,\xi}^\alpha a\|_{L^\infty_x W^\tilde{d}_\xi{}^\infty}.$$

If  $a \in \Upsilon^{r+1}$  with  $r \in \mathbb{N}$ , then the paradifferential operator  $T_a$  is defined via approximating  $a$  by smooth symbols. To be precise, let  $\Omega \Subset \mathbb{R}^d \times (\mathbb{R}^d \setminus 0)$  be an open neighborhood of  $\text{supp } a$  and let  $\{a_n\}_{n \in \mathbb{N}} \subset C_c^\infty(\Omega)$  such that

$$\lim_{n \rightarrow \infty} N^r(a_n - a) = 0.$$

Note that such an approximation is always possible because  $a$  is compactly supported and we only require the convergence with respect to the  $N^r$ -norm (not the  $N^{r+1}$ -norm)! By [Proposition 4.4](#) and [Lemma 4.6](#), for all  $n, m \in \mathbb{N}$  we have

$$\|T_{a_n} - T_{a_m}\|_{L^2 \rightarrow L^2} \lesssim M_{\tilde{d}}^{0,0}(a_n - a_m) \lesssim N^0(a_n - a_m) \leq N^r(a_n - a_m).$$

Therefore, for all  $u \in L^2$ , the sequence  $\{T_{a_n}u\}_{n \in \mathbb{N}}$  is Cauchy in  $L^2$  and we define

$$T_a u = \lim_{n \rightarrow \infty} T_{a_n} u.$$

Clearly this definition is independent of the choice of the sequence  $\{a_n\}_{n \in \mathbb{N}}$  and extends the definition of paradifferential operators with symbols that are smooth with respect to  $\xi$ . Then we define the operators  $T_a^h, \mathcal{P}_a, \mathcal{P}_a^h$  and  $\mathcal{P}_a^h$  exactly as before.

**Proposition 4.36.** *If  $a \in \Upsilon^{r+1}$  with  $r \geq 0$ , then for all  $h \in (0, 1]$ , we have  $T_a^h : L^2 \rightarrow L^2$ . Moreover,*

$$\sup_{h \in (0,1]} \|T_a^h\|_{L^2 \rightarrow L^2} \lesssim N^0(a).$$

Consequently, for all  $\epsilon \geq 0$  we have

$$\sup_{h \in (0,1]} \|\mathcal{P}_a^{h,\epsilon}\|_{L^2 \rightarrow L^2} \lesssim N^0(a).$$

*Proof.* The general case  $h \in (0, 1]$  follows from the case  $h = 1$  and we shall assume  $h = 1$ . Choose a convergent sequence  $\{a_n\}_{n \in \mathbb{N}} \subset C_c^\infty(\Omega)$  as above. For all  $u \in L^2$  with  $\|u\|_{L^2} = 1$ , we have

$$\|T_a u\|_{L^2} \leq \|T_a u - T_{a_n} u\|_{L^2} + \|T_{a_n} u\|_{L^2},$$

where  $\lim_{n \rightarrow \infty} \|T_a u - T_{a_n} u\|_{L^2} = 0$  by the definition of  $T_a u$  and

$$\|T_{a_n} u\|_{L^2} \lesssim N^0(a_n) \lesssim N^0(a - a_n) + N^0(a) \rightarrow N^0(a).$$

Therefore, passing  $n \rightarrow \infty$  we conclude that  $\|T_a\|_{L^2 \rightarrow L^2} \lesssim N^0(a)$ . The estimate for  $\mathcal{P}_a^{h,\epsilon}$  follows similarly to Proposition 4.18.  $\square$

Combining the approximation method above and the analysis in Proposition 4.26, we obtain the following corollaries similarly to Corollaries 4.28 and 4.29.

**Corollary 4.37.** *Let  $\epsilon \geq 0$ ,  $(m, k) \in (\mathbb{R} \cup \{-\infty\})^2$ ,  $r \in \mathbb{N}$ , with  $r \geq 0$ . If  $a_h \in \Upsilon^{r+1} \cap \sigma_\epsilon$  and  $b_h \in \Gamma_{k,1}^{m,r} \cap \sigma_0$ , then, for all  $k' \in \mathbb{R}$  such that  $r \geq k + k'$ , we have*

$$\begin{aligned} \mathcal{P}_{a_h}^{h,\epsilon} \mathcal{P}_{b_h}^h - \mathcal{P}_{(\theta_{h,*}^{\epsilon,0} a_h) \sharp_h b_h}^h &= \mathcal{O}(h^{(1+\epsilon)r - \epsilon(k+k')}) M_{k',1}^{-m,r}(a_h) M_{k,1}^{m,r}(b_h)_{L^2 \rightarrow L^2}, \\ \mathcal{P}_{b_h}^h \mathcal{P}_{a_h}^{h,\epsilon} - \mathcal{P}_{b_h \sharp_h (\theta_{h,*}^{\epsilon,0} a_h)}^h &= \mathcal{O}(h^{(1+\epsilon)r - \epsilon(k+k')}) M_{k',1}^{-m,r}(a_h) M_{k,1}^{m,r}(b_h)_{L^2 \rightarrow L^2}. \end{aligned}$$

*Proof.* Let  $a_h$  be a sequence of approximating symbols  $\{a_h^n\}_{n \in \mathbb{N}} \subset C_c^\infty(\mathbb{R}^{2d}) \cap \sigma_\epsilon$  which is bounded, uniformly in  $h \in (0, 1]$ , with respect to the norm  $N^r(\cdot)$ . Note that for all  $k' \in \mathbb{R}$ , we have  $M_{k',1}^{-m,r}(\cdot) \lesssim N^r(\cdot)$ . And thus, when  $n \in \mathbb{N}$  is sufficiently large, we have  $M_{k',1}^{-m,r}(a_h^n - a_h) \leq 2N^r(a_h^n - a_h) = o(1)$ . By Corollary 4.28, if  $r \geq k + k'$ , we have

$$\begin{aligned} \mathcal{P}_{a_h^n}^{h,\epsilon} \mathcal{P}_{b_h}^h - \mathcal{P}_{(\theta_{h,*}^{\epsilon,0} a_h^n) \sharp_h b_h}^h &= \mathcal{O}(h^{(1+\epsilon)r - \epsilon(k+k')}) M_{k',1}^{-m,r}(a_h^n) M_{k,1}^{m,r}(b_h)_{L^2 \rightarrow L^2} \\ &= \mathcal{O}(h^{(1+\epsilon)r - \epsilon(k+k')}) M_{k',1}^{-m,r}(a_h) M_{k,1}^{m,r}(b_h)_{L^2 \rightarrow L^2} + o(1)_{L^2 \rightarrow L^2}. \end{aligned}$$

In fact, for all  $u \in \mathcal{S}(\mathbb{R}^d)$ , as  $n \rightarrow \infty$ , by Proposition 4.36, we have

$$\begin{aligned} \|\mathcal{P}_{a_h^n}^{h,\epsilon} \mathcal{P}_{b_h}^h u - \mathcal{P}_{(\theta_{h,*}^{\epsilon,0} a_h^n) \sharp_h b_h}^h u\|_{L^2} &= \|(\mathcal{P}_{a_h^n}^{h,\epsilon} - \mathcal{P}_{a_h^n}^{h,\epsilon}) \mathcal{P}_{b_h}^h u\|_{L^2} + \|\mathcal{P}_{(\theta_{h,*}^{\epsilon,0} (a_h^n - a_h)) \sharp_h b_h}^h u\|_{L^2} + \|\mathcal{P}_{a_h^n}^{h,\epsilon} \mathcal{P}_{b_h}^h u - \mathcal{P}_{(\theta_{h,*}^{\epsilon,0} a_h^n) \sharp_h b_h}^h u\| \\ &= o(1)(\|\mathcal{P}_{b_h}^h u\|_{L^2} + \|u\|_{L^2}) + \mathcal{O}(h^{(1+\epsilon)r - \epsilon(k+k')}) M_{k',1}^{-m,r}(a_h) M_{k,1}^{m,r}(b_h) \|u\|_{L^2}. \end{aligned}$$

Passing  $n \rightarrow \infty$  and then use the density of  $\mathcal{S}(\mathbb{R}^d)$  in  $L^2$ , we conclude that for all  $u \in L^2$ , we have

$$\|\mathcal{P}_{a_h^n}^{h,\epsilon} \mathcal{P}_{b_h}^h u - \mathcal{P}_{(\theta_{h,*}^{\epsilon,0} a_h^n) \sharp_h b_h}^h u\|_{L^2} = \mathcal{O}(h^{(1+\epsilon)r - \epsilon(k+k')}) M_{k',1}^{-m,r}(a_h) M_{k,1}^{m,r}(b_h) \|u\|_{L^2}.$$

The estimate for  $\mathcal{P}_{b_h}^h \mathcal{P}_{a_h}^{h,\epsilon} - \mathcal{P}_{b_h \sharp_h (\theta_{h,*}^{\epsilon,0} a_h)}^h$  is similar.  $\square$

**Corollary 4.38.** *If  $\epsilon \geq 0$  and  $a_h, b_h \in \Upsilon^{r+1}$ , where  $r \in \mathbb{N}$ , then*

$$\mathcal{P}_{a_h}^{h,\epsilon} \mathcal{P}_{b_h}^{h,\epsilon} - \mathcal{P}_{a_h \sharp_h^{\epsilon} b_h}^{h,\epsilon} = \mathcal{O}(h^{(1+\epsilon)r})_{L^2 \rightarrow L^2},$$

where the symbol  $a_h \sharp_h^{\epsilon} b_h = a_h \sharp_{h,r}^{0,\epsilon} b_h$  is defined by (2-4).

**4C5. Almost sharp Gårding inequality for paradifferential operators.** We need an almost sharp Gårding inequality for our paradifferential calculus. There are various works on the (almost) sharp Gårding inequality for pseudodifferential operators with limited regularities; see, e.g., [Taylor 1991; Tataru 2002; Hérau 2002].

**Lemma 4.39.** *If  $\epsilon \in (0, 1)$  and  $a_h \in M_{n \times n}(\Gamma^{0,r}) \cap \sigma_0$  is compactly supported, where  $n \in \mathbb{N}$ ,  $r \geq \max\{\tilde{d}, \epsilon^{-1} - 1\}$  and  $\text{Re } a \geq 0$ , then, for all  $\epsilon \in (0, 1)$ , there exists  $C > 0$  such that, for all  $u \in L^2$ ,*

$$\text{Re}(T_{a_h}^h u, u)_{L^2} \geq -Ch^{1-\epsilon} \|u\|_{L^2}^2.$$

*Proof.* By Lemma 4.33 and the condition  $r \geq \tilde{d}$ , we may replace  $T_{a_h}^h$  with  $\text{Op}_h(a_h)$  in the above inequality. As  $a_h \in \sigma_0$  and is compactly supported, we have  $\{b_h(x, \xi) = h^{-1+\epsilon} a_h(x, h\xi)\}_{h \in (0,1]}$  is bounded in  $\Gamma^{1-\epsilon, r}$ . By [Taylor 1991, §2.4 (2.4.6)], as  $r \geq \epsilon^{-1} - 1$ , we have  $1 - \epsilon \leq r/(1+r)$  and thus

$$\text{Re}(\text{Op}(b_h)u, u)_{L^2} \gtrsim -\|u\|_{L^2}^2.$$

We conclude by  $\text{Op}(b_h) = h^{-1+\epsilon} \text{Op}_h(a_h)$ . □

We are mostly interested in the case where  $\epsilon = \frac{1}{2}$ . In this case, the condition for  $r$  is simply  $r \geq \max\{\tilde{d}, (\frac{1}{2})^{-1} - 1\} = \tilde{d}$ . Next we show that the almost sharp Gårding inequality also applies to symbols in  $\Upsilon^{1+r}$ .

**Lemma 4.40.** *If  $\epsilon \in (0, 1)$  and  $a_h \in M_{n \times n}(\Upsilon^{1+r})$ , with  $n \in \mathbb{N}$ ,  $r \geq \max\{\tilde{d}, \epsilon^{-1} - 1\}$ , then there exists  $C > 0$  such that, for all  $u \in L^2$ ,*

$$\text{Re}(T_{a_h}^h u, u)_{L^2} \geq -Ch^{1-\epsilon} \|u\|_{L^2}^2, \quad \text{Re}(\mathcal{P}_{a_h}^h u, u)_{L^2} \geq -Ch^{1-\epsilon} \|u\|_{L^2}^2.$$

*Proof.* Choose a sequence  $a_h^j \in M_{n \times n}(\Gamma^{0, r})$  which converges to  $a_h$  with respect to the norm  $N^r(\cdot)$  and is uniformly compactly supported in  $\mathbb{R}^d \times (\mathbb{R}^d \setminus \{0\})$ . Apply the almost sharp Gårding inequality for  $a_h^j$ ; there exists a constant  $C > 0$  which is independent of  $j$  such that, for all  $u \in L^2$ , we have

$$\text{Re}(T_{a_h}^h u, u)_{L^2} = \text{Re}(T_{a_h - a_h^j}^h u, u)_{L^2} + \text{Re}(T_{a_h^j}^h u, u)_{L^2} \geq o(1) - Ch^{1-\epsilon} \|u\|_{L^2}^2.$$

We conclude the almost sharp Gårding inequality for  $T_{a_h}^h$  by passing  $j \rightarrow \infty$ . Therefore,

$$\begin{aligned} \text{Re}(\mathcal{P}_{a_h}^h u, u)_{L^2} &= \sum_{j \in \mathbb{N}} \text{Re}(\psi_j T_{\psi_j a_h}^h \psi_j u, u)_{L^2} = \sum_{j \in \mathbb{N}} \text{Re}(T_{\psi_j a_h}^h \psi_j u, \psi_j u)_{L^2} \\ &\gtrsim -h^{1-\epsilon} \sum_{j \in \mathbb{N}} \|\psi_j u\|_{L^2}^2 \gtrsim -h^{1-\epsilon} \|u\|_{L^2}^2. \end{aligned} \quad \square$$

#### 4C6. Relation with quasihomogeneous wavefront sets.

**Lemma 4.41.** *If  $r \geq 0$  and  $a_h = \sum_{j=0}^r h^j a_h^j$ , where  $a_h^j \in \Upsilon^{1+r-j}$  such that  $a_h$  is elliptic at  $(x_0, \xi_0) \in \mathbb{R}^d \times (\mathbb{R}^d \setminus \{0\})$  in the sense that, for some neighborhood  $\Omega$  of  $(x_0, \xi_0)$ , we have*

$$\inf_{0 < h < 1} \inf_{(x, \xi) \in \Omega} |a_h(x, \xi)| > 0,$$

*then for all  $u \in L^2$  such that  $T_{a_h}^h u = \mathcal{O}(h^\sigma)_{L^2}$ , where  $0 \leq \sigma \leq r$ , we have  $(x_0, \xi_0) \notin \text{WF}_{0,1}^\sigma(u)$ .*

*Proof.* Assume that  $\Omega \subset \mathbb{R}^d \times (\mathbb{R}^d \setminus \{0\})$ . Let  $b_h \in S_{-\infty}^-$  with  $\text{supp } b_h \subset \Omega$ . Then by the symbolic calculus stated in Corollary 4.38, there exists  $c_h = \sum_{j=0}^r h^j c_h^j$ , where  $c_h^j \in \Upsilon^{1+r-j}$ , such that

$$T_{b_h}^h = T_{c_h}^h T_{a_h}^h + \mathcal{O}(h^r)_{L^2 \rightarrow L^2}.$$

Thus  $T_{b_h}^h u = \mathcal{O}(h^\sigma)_{L^2}$ . By Lemma 4.33 we have  $\text{Op}_h(b_h)u = \mathcal{O}(h^\sigma)_{L^2}$ . We conclude by Lemma 2.14. □

**Lemma 4.42.** *Let  $\epsilon \geq 0$ ,  $e \in \Gamma_{0,0}^{m,r}$  if  $\epsilon = 0$  and  $e \in \Gamma_{0,1}^{m,r}$  if  $\epsilon > 0$ , and suppose that  $e$  is homogeneous of degree  $m$  with respect to  $\xi$ . Then, for  $f \in H^s$  and  $0 \leq \sigma \leq (1+\epsilon)r$ ,*

$$\text{WF}_{\epsilon,1}^{s+\sigma-m}(\mathcal{P}_e f)^\circ \subset \text{WF}_{\epsilon,1}^{s+\sigma}(f)^\circ.$$

If in addition  $e$  is elliptic, i.e., for some  $C > 0$  and  $|\xi|$  sufficiently large,  $|e(x, \xi)| \geq C|\xi|^m$ , then

$$\text{WF}_{\epsilon,1}^{s+\sigma-m}(\mathcal{P}_e f)^\circ = \text{WF}_{\epsilon,1}^{s+\sigma}(f)^\circ.$$

*Proof.* For  $\mu \in \mathbb{R}$ , define  $Z^\mu = \mathcal{P}_{|\xi|^\mu}$ . Then  $Z^{-\mu}Z^\mu - \text{Id} \in \mathcal{O}_{-\infty}^-$ . Therefore,

$$f - Z^{-s}Z^s f \in H_\infty^\infty, \quad \mathcal{P}_e f - \mathcal{P}_{e\sharp|\xi|^{-s}}Z^s f \in H_{\delta r}^{s+r-m} + H^\infty,$$

where  $\delta = 0$  if  $\epsilon = 0$ , while  $\delta = 1$  if  $\epsilon > 0$ . By Lemma 2.15 and the fact that  $Z^{\pm s}$  are pseudodifferential operators with elliptic symbols in  $S_0^{\pm s}$ , we readily have

$$\text{WF}_{\epsilon,1}^{s+\sigma}(f)^\circ = \text{WF}_{\epsilon,1}^\sigma(Z^s f)^\circ, \quad \text{WF}_{\epsilon,1}^{s+\sigma-m}(\mathcal{P}_e f)^\circ = \text{WF}_{\epsilon,1}^{\sigma-(m-s)}(\mathcal{P}_{e|\xi|^{-s}}Z^s f)^\circ.$$

So we may assume that  $s = 0$ . Let  $a, b \in S_{-\infty}^- \cap \mathcal{S}_\epsilon$  such that

$$\text{supp } b \subset \{a = 1\} \subset \text{supp } a \subset \mathbb{R}^{2d} \setminus \text{WF}_{\epsilon,1}^\sigma(f).$$

Then by Lemma 2.14,  $\text{Op}_h^{\epsilon,1}(a)f = \mathcal{O}(h^\sigma)_{L^2}$ . By Corollary 4.32, Lemma 4.34, Proposition 4.26, and Corollary 4.28,

$$\begin{aligned} h^m \text{Op}_h^{\epsilon,1}(b)\mathcal{P}_e f &= \text{Op}_h^{\epsilon,1}(b)\mathcal{P}_e^h f + \mathcal{O}(h^\infty)_{L^2} \\ &= \text{Op}_h^{\epsilon,1}(b)\mathcal{P}_e^h \text{Op}_h^{\epsilon,1}(a)f + \text{Op}_h^{\epsilon,1}(b)\mathcal{P}_e^h \text{Op}_h^{\epsilon,1}(1-a)f + \mathcal{O}(h^\infty)_{L^2} \\ &= \mathcal{O}(1)_{L^2 \rightarrow L^2} \text{Op}_h^{\epsilon,1}(a)f + \mathcal{O}(h^{r(1+\epsilon)})_{L^2} \\ &= \mathcal{O}(h^\sigma)_{L^2}, \end{aligned}$$

proving the first statement. The second statement follows by a construction of parametrix. □

### 5. Asymptotically flat water waves

In this section we prove Theorem 1.6. The idea is to combine the analysis in [Alazard, Burq and Zuily 2011] with the dyadic paradifferential calculus in weighted Sobolev spaces. We shall use the following formal notations for simplicity. Let  $w$  be a function on  $\mathbb{R}^d$  which is nowhere-vanishing. Then for any operator  $\mathcal{A}$  between some function spaces on  $\mathbb{R}^d$  and, for any function  $f$  on  $\mathbb{R}^d$ , we introduce the following notations whenever they are well-defined:

$$\mathcal{A}^{(w)} = w\mathcal{A}w^{-1}, \quad f^{(w)} = wf.$$

Note that  $(\mathcal{A}f)^{(w)} = \mathcal{A}^{(w)}f^{(w)}$ . For  $k \in \mathbb{R}$ , we also define by an abuse of notation

$$\mathcal{A}^{(k)} = \mathcal{A}^{((x)^k)}, \quad f^{(k)} = f^{((x)^k)}.$$

Observe that  $L_k^2 = H_k^0$  is an Hilbert space with the inner product

$$(f, g)_{L_k^2} = (f^{(k)}, g^{(k)})_{L^2}.$$

**5A. Dirichlet–Neumann operator.** We study the Dirichlet–Neumann operator on weighted Sobolev spaces and its parilinearization. The time variable will be temporarily omitted for simplicity.

**5A1. Boundary flattening.** Let  $\eta \in W^{1,\infty}(\mathbb{R}^d)$ , such that

$$\delta := b + \inf_{x \in \mathbb{R}^d} \eta(x) > 0. \quad (5-1)$$

Define  $\tau(x, z) = (x, z + \eta(x))$  and set

$$\begin{aligned} \tilde{\Omega} &= \tau^{-1}(\Omega) = \{-b - \eta(x) < z < 0\}, \\ \tilde{\Sigma} &= \tau^{-1}(\Sigma) = \{z=0\}, \\ \tilde{\Gamma} &= \tau^{-1}(\Gamma) = \{z = -b - \eta(x)\}. \end{aligned}$$

Let  $\tau_*$  be the pullback deduced by  $\tau$ , then

$$\tau_*(dx^2 + dy^2) = (dx \ dz) \varrho \begin{pmatrix} dx \\ dz \end{pmatrix},$$

where

$$\varrho = \begin{pmatrix} \text{Id} + (\nabla \eta)^t (\nabla \eta) & \nabla \eta \\ {}^t(\nabla \eta) & 1 \end{pmatrix}.$$

We verify that

$$\varrho^{-1} = \begin{pmatrix} \text{Id} & -\nabla \eta \\ -{}^t(\nabla \eta) & 1 + |\nabla \eta|^2 \end{pmatrix}.$$

Let  $\nabla_{xz} = (\nabla_x, \partial_z)$ . Then the divergence, gradient and Laplacian with respect to the metric  $\varrho$  are

$$\begin{aligned} \text{div}_\varrho u &= \nabla_{xz} \cdot u, \\ \nabla_\varrho u &= (\nabla u - \nabla \eta \partial_z u, -\nabla \eta \cdot \nabla u + (1 + |\nabla \eta|^2) \partial_z u), \\ \Delta_\varrho u &= \partial_z^2 u + (\nabla - \nabla \eta \partial_z)^2 u. \end{aligned}$$

The exterior unit normal to  $\partial \tilde{\Omega} = \tilde{\Sigma} \cup \tilde{\Gamma}$  is

$$\mathbf{n}_\varrho = \langle (D\tau)^{-1}|_{T\partial \tilde{\Omega}}, \mathbf{n} \rangle = \begin{cases} {}^t(-\nabla \eta, 1 + |\nabla \eta|^2) / \sqrt{1 + |\nabla \eta|^2}, & \tilde{\Sigma}, \\ {}^t(0, 1), & \tilde{\Gamma}. \end{cases}$$

Let  $\psi \in H^{1/2}$ , and suppose that  $\phi$  satisfies the equation

$$\Delta_{xy} \phi = 0, \quad \phi|_\Sigma = \psi, \quad \partial_{\mathbf{n}} \phi|_\Gamma = 0.$$

Then  $v = (\tau|_{\tilde{\Omega}})_* \phi$  satisfies

$$\Delta_\varrho v = 0, \quad v|_{\tilde{\Sigma}} = \psi, \quad \partial_{\mathbf{n}_\varrho} v|_{\tilde{\Gamma}} = 0. \quad (5-2)$$

The Dirichlet–Neumann operator can now be written as

$$\sqrt{1 + |\nabla \eta|^2}^{-1} G(\eta) \psi = \partial_{\mathbf{n}_\varrho} v|_{\tilde{\Sigma}} = \mathbf{n}_\varrho \cdot \nabla_{xz} v|_{z=0}.$$

**5A2. Elliptic estimate.** Let  $\chi_0 \in C^\infty(\mathbb{R})$  with  $\chi_0(z) = 0$  for  $z \leq -\frac{\delta}{2}$  and  $\chi_0(z) = 1$  for  $z \geq 0$ . Take the decomposition  $v = \tilde{v} + \underline{\psi}$ , where

$$\underline{\psi}(x, z) = \chi_0(z) e^{z(D_x)} \psi(x).$$

**Lemma 5.1.** Let  $n \in \mathbb{N}$ ,  $m \in \mathbb{R}$ ,  $\mu \in \mathbb{R}$ ,  $k \in \mathbb{R}$ ,  $a \in S_0^m$ . Then

$$\|\partial_z^n \text{Op}(a) \underline{\psi}\|_{L_z^2(\mathbb{R}_{\leq 0}, H_k^{\mu-n-m+1/2})} \lesssim \|\psi\|_{H_k^\mu}.$$

*Proof.* We only prove the case where  $n = 0$ . The general case follows with a similar argument and the identity

$$\partial_z^n \underline{\psi}(x, z) = \sum_{j=0}^n \binom{n}{j} \chi_0^{(n-j)}(z) \langle D_x \rangle^j e^{z \langle D_x \rangle} \psi(x).$$

Let

$$\begin{aligned} b(x, \xi) &= a(x, \xi) \langle \xi \rangle^{\mu-m} \in S_0^\mu, \\ \lambda(z, \xi) &= \chi_0(z) e^{z \langle \xi \rangle} \langle \xi \rangle^{1/2} \in L_{z \leq 0}^\infty S_0^{1/2}. \end{aligned}$$

Then for all  $N \geq 0$ ,

$$\|\text{Op}(a) \underline{\psi}\|_{L_z^2(\mathbb{R}_{\leq 0}, H_k^{\mu-m+1/2})} \lesssim \|\text{Op}(\lambda) \text{Op}(b) \psi\|_{L_z^2(\mathbb{R}_{\leq 0}, L_k^2)} + \|\psi\|_{H_k^{-N}}.$$

Observe that

$$\begin{aligned} \text{Op}(\lambda)^{(k)} - (\text{Op}(\lambda)^{(k)})^* &\in L_{z \leq 0}^\infty \mathcal{O}_0^{-1/2}, \\ (\text{Op}(\lambda)^{(k)})^2 - \text{Op}(\lambda^2)^{(k)} &\in L_{z \leq 0}^\infty \mathcal{O}_0^0. \end{aligned}$$

Also note that

$$\sigma(\xi) := \int_{-\infty}^0 \lambda^2(z, \xi) dz = \langle \xi \rangle \int_{-\infty}^0 \chi_0^2(z) e^{2 \langle \xi \rangle z} dz \in S_0^0.$$

Therefore,

$$\begin{aligned} \|\text{Op}(\lambda) \text{Op}(b) \psi\|_{L_z^2(\mathbb{R}_{\leq 0}, L_k^2)}^2 &= (\text{Op}(\lambda^2) \text{Op}(b) \psi, \text{Op}(b) \psi)_{L_z^2(\mathbb{R}_{\leq 0}, L_k^2)} + \mathcal{O}(\|\psi\|_{H_k^\mu}^2) \\ &= (\text{Op}(\sigma) \text{Op}(b) \psi, \text{Op}(b) \psi)_{L_k^2} + \mathcal{O}(\|\psi\|_{H_k^\mu}^2) = \mathcal{O}(\|\psi\|_{H_k^\mu}^2). \end{aligned} \quad \square$$

**Lemma 5.2.** *For all  $k \in \mathbb{R}$ , we have  $\|\tilde{v}\|_{H_k^1} \leq C(\|\eta\|_{W^{1,\infty}}) \|\psi\|_{H_k^{1/2}}$ .*

*Proof.* Let  $H_\rho^{1,0}$  be the completion of the space

$$\{f \in C^\infty(\tilde{\Omega}) : f \text{ vanishes in a neighborhood of } \tilde{\Sigma}\},$$

with respect to the norm

$$\|u\|_{H_\rho^{1,0}} := \|\nabla_\rho u\|_{L_\rho^2} = (\nabla_\rho u, \nabla_\rho u)_{L_\rho^2}^{1/2},$$

where  $(X, Y)_{L_\rho^2} := \int_{\tilde{\Omega}} \rho(X, Y) dx dz$ . As  $b < \infty$ , by the Poincaré inequality,

$$\|u\|_{L^2} \leq C(\|\eta\|_{L^\infty}) \|\partial_z u\|_{L^2} \leq C(\|\eta\|_{W^{1,\infty}}) \|u\|_{H_\rho^{1,0}}$$

for all  $u \in H_\rho^{1,0}$ . Let  $0 < \zeta \in C^\infty(\mathbb{R})$  be such that  $\zeta(z) = 1$  for  $|z| \leq 1$ , and  $\zeta(z) = z$  for  $|z| \geq 2$ . For some  $R > 0$  sufficiently large to be determined later, set  $w(x) = R \times \zeta(\langle x \rangle^k / R)$ . Then  $\langle x \rangle^k \lesssim w(x) \lesssim R \langle x \rangle^k$ ,  $\text{supp } \nabla w \subset \{\langle x \rangle \gtrsim R^{1/k}\}$ , and  $|\nabla w(x)| \lesssim R^{(k-1)/k}$ .

As  $\tilde{v}$  satisfies the equation  $\Delta_\rho \tilde{v} = -\Delta_\rho \underline{\psi}$ , we consider  $\tilde{v}^{(w)}$  as the variational solution to the equation  $B(\tilde{v}^{(w)}, \cdot) = -L(\cdot)$ , where, for  $u, \varphi \in H_\rho^{1,0}$ ,

$$B(u, \varphi) = (\nabla_\rho^{(w)} u, \nabla_\rho^{(1/w)} \varphi)_{L^2(\tilde{\Omega})}, \quad L(\varphi) = (\nabla_\rho^{(w)} \underline{\psi}^{(w)}, \nabla_\rho^{(1/w)} \varphi)_{L^2(\tilde{\Omega})}.$$

Observe that  $\nabla_\rho^{(w^\pm)} = \nabla_\rho \mp b_w$ , where  $b_w = (w^{-1}\nabla w, -\nabla\eta \cdot w^{-1}\nabla w) \in L^\infty$ , satisfies  $\|b_w\| \leq C(\|\eta\|_{W^{1,\infty}})R^{-1/k}$ . We verify that  $L$  and  $B$  are continuous linear and bilinear forms on  $H_\rho^{1,0}$ . Moreover  $B$  is coercive when  $R$  is sufficiently large; indeed,

$$B(\varphi, \varphi) = \|\nabla_\rho \varphi\|_{L_\rho^2}^2 - \|b_w \varphi\|_{L_\rho^2}^2 \geq (1 - C(\|\eta\|_{W^{1,\infty}})R^{-2/k})\|\nabla_\rho \varphi\|_{L_\rho^2}^2. \tag{5-3}$$

Therefore, by the Lax–Milgram theorem and Lemma 5.1,

$$\|\tilde{v}\|_{H_k^1} \lesssim \|\tilde{v}^{(w)}\|_{H_\rho^{1,0}} \lesssim \|L\|_{(H_\rho^{1,0})^*} \lesssim \|\underline{\psi}\|_{H^1} \lesssim \|\psi\|_{H_k^{1/2}}. \quad \square$$

**Proposition 5.3.** *Let  $(\eta, \psi) \in W^{1,\infty} \times H_k^{1/2}$ ,  $k \in \mathbb{R}$ . Then  $\|G(\eta)\psi\|_{H_k^{-1/2}} \leq C(\|\eta\|_{W^{1,\infty}})\|\psi\|_{H_k^{1/2}}$ .*

*Proof.* By Lemmas 5.1 and 5.2,  $v \in L_z^2((-\delta, 0), H_k^1) \cap H_z^1((-\delta, 0), L_z^2)$ . By a classical interpolation result (see, e.g., [Alazard, Burq and Zuily 2014, Lemma 2.19]) and the equation satisfied by  $v$ , we deduce that

$$v \in C_z^0([-\delta, 0], H_k^{1/2}) \cap C_z^1([-\delta, 0], H_k^{-1/2}). \quad \square$$

**5A3. Higher regularity.**

**Proposition 5.4.** *Let  $(\eta, \psi) \in H^{\mu+1/2} \times H_k^{\sigma+1/2}$ , where  $k \in \mathbb{R}$ ,  $\mu > \frac{1}{2} + \frac{d}{2}$ ,  $0 \leq \sigma \leq [\mu - \frac{1}{2}]$ . Then*

$$\|G(\eta)\psi\|_{H_k^{\sigma-1/2}} \leq C(\|\eta\|_{H^{\mu+1/2}})\|\psi\|_{H_k^{\sigma+1/2}}.$$

*Consequently, if  $(\eta, \psi) \in H^{\mu+1/2} \times \mathcal{H}_k^{\sigma+1/2,\delta}$ , with  $\delta \geq 0$ ,  $k \in \mathbb{N}$  and  $\sigma - k\delta \geq 0$ , then*

$$\|G(\eta)\psi\|_{\mathcal{H}_k^{\sigma-1/2,\delta}} \leq C(\|\eta\|_{H^{\mu+1/2}})\|\psi\|_{\mathcal{H}_k^{\sigma+1/2,\delta}}.$$

*Proof.* We shall only prove the cases where  $\sigma \in \mathbb{N}$ . The remaining cases follow by interpolation. By Section 5A2, it suffices to prove that, for all  $\sigma \in [0, \mu - \frac{1}{2}] \cap \mathbb{N}$ , there exists  $\delta > 0$  such that

$$\tilde{v} \in L^2((-\delta, 0), H_k^{\sigma+1}) \cap H^1((-\delta, 0), H_k^\sigma).$$

Let  $N_\sigma$  be the corresponding norm of  $\tilde{v}$ , we shall prove that  $N_\sigma < +\infty$ . The case where  $\sigma = 0$  has already been proven by Lemma 5.2. It remains to bound  $N_{\sigma+1}$  by  $N_\sigma$  via a mathematical induction. Note that if  $\chi \in C_c^\infty((-\delta, \delta))$ , then  $\chi \partial_x^\sigma \tilde{v}$  satisfies the equation

$$-\Delta_\rho(\chi \partial_x^\sigma \tilde{v}) + K \tilde{v} = \Delta_\rho(\chi \partial_x^\sigma \underline{\psi}) - K \underline{\psi}. \tag{5-4}$$

where  $K = [\Delta_\rho, \chi \partial_x^\sigma]$ . Note that  $\Delta_\rho = P \cdot P$  with  $P = (\nabla - \nabla\eta \partial_z, \partial_z)$ , so

$$K = P \cdot [P, \chi \partial_x^\sigma] + [P, \chi \partial_x^\sigma] \cdot P.$$

By an explicit calculation

$$[P, \chi \partial_x^\sigma] = (-\chi[\nabla\eta, \partial_x^\sigma] \partial_z - \nabla\eta \chi' \partial_x^\sigma, \chi' \partial_x^\sigma).$$



Integrating the following pairings by parts using  $\tilde{v}|_{z=0} = 0$ , we have by [Lemma 5.1](#) that

$$\begin{aligned} |(\chi \partial_x^\sigma \tilde{v}, K \chi \partial_x^\sigma \tilde{v})_{L^2 L_k^2}| &\lesssim \|P^*(\chi \partial_x^\sigma \tilde{v})\|_{L^2 L_k^2} \| [P, \chi \partial_x^\sigma] \chi \tilde{v} \|_{L^2 L_k^2} + \|P(\chi \partial_x^\sigma \tilde{v})\|_{L^2 L_k^2} \| [P, \chi \partial_x^\sigma]^* \chi \tilde{v} \|_{L^2 L_k^2} \\ &\lesssim N_\sigma N_{\sigma+1}, \end{aligned}$$

$$\begin{aligned} |(\chi \partial_x^\sigma \tilde{v}, K \chi \partial_x^\sigma \underline{\psi})_{L^2 L_k^2}| &\lesssim \|P^*(\chi \partial_x^\sigma \tilde{v})\|_{L^2 L_k^2} \| [P, \chi \partial_x^\sigma] \chi \underline{\psi} \|_{L^2 L_k^2} + \|P(\chi \partial_x^\sigma \underline{\psi})\|_{L^2 L_k^2} \| [P, \chi \partial_x^\sigma]^* \chi \tilde{v} \|_{L^2 L_k^2} \\ &\lesssim \|\psi\|_{H^{\sigma+1/2}} (N_\sigma + N_{\sigma+1}), \end{aligned}$$

$$|(\chi \partial_x^\sigma \tilde{v}, -\Delta_\varrho(\chi \partial_x^\sigma \underline{\psi}))_{L^2 L_k^2}| \lesssim \|P(\chi \partial_x^\sigma \tilde{v})\|_{L^2 L_k^2} \|P(\chi \partial_x^\sigma \underline{\psi})\|_{L^2 L_k^2}^2 \lesssim \|\psi\|_{H^{\sigma+1/2}} N_{\sigma+1}.$$

In the above inequalities, the adjoint operators are taken with respect to  $L^2 L_k^2$ . Using again the structure of  $\Delta_\varrho$ , we have by [\(5-3\)](#) that

$$(\chi \partial_x^\sigma \tilde{v}, -\Delta_\varrho(\chi \partial_x^\sigma \tilde{v}))_{L^2 L_k^2} \gtrsim \|P(\chi \partial_x^\sigma \tilde{v})\|_{L^2 L_k^2}^2 - \|\chi \partial_x^\sigma \tilde{v}\|_{L^2 L_k^2}^2 \gtrsim \|\chi \partial_x^\sigma \tilde{v}\|_{H^1 H_k^1}^2 - N_\sigma^2.$$

Pairing [\(5-4\)](#) with  $\chi \partial_x^\sigma \tilde{v}$  and using the estimates above, for all  $\epsilon > 0$ ,

$$N_{\sigma+1}^2 \lesssim \|\chi \partial_x^\sigma \tilde{v}\|_{H^1 H_k^1}^2 \lesssim N_\sigma N_{\sigma+1} + \|\psi\|_{H^{\sigma+1/2}} (N_\sigma + N_{\sigma+1}) \lesssim \epsilon N_{\sigma+1}^2 + \epsilon^{-1} (N_\sigma^2 + \|\psi\|_{H^{\sigma+1/2}}^2).$$

All the constants hidden by  $\lesssim$  are of the form  $C(\|\eta\|_{H^{\mu+1/2}})$ . We thus conclude the induction by choosing  $\epsilon > 0$  sufficiently small. By interpolation as in [Proposition 5.3](#),

$$v \in C_z^0([-\delta, 0], H_k^{\sigma+1/2}) \cap C_z^1([-\delta, 0], H_k^{\sigma-1/2}).$$

When  $\psi \in \mathcal{H}_k^{\sigma, \delta}$ , we apply the above estimate to  $\psi \in H_j^{\sigma-\delta j}$  and conclude.  $\square$

**5B. Paralinearization.** Now we paralinearize the system of water waves. The following results are immediate consequences of the analysis in [\[Alazard, Burq and Zuily 2011\]](#) and our dyadic paradifferential calculus on weighted Sobolev spaces.

**Proposition 5.5.** *Let  $(\eta, \psi) \in \mathcal{H}_k^{\mu+1/2, \delta} \times \mathcal{H}_k^{\mu, \delta}$  with  $\mu - \frac{1}{2} \in \mathbb{N}$ ,  $k \in \mathbb{N}$  and  $\mu - \delta k > 3 + \frac{d}{2}$ . Let*

$$B = \frac{\nabla \eta \cdot \nabla \psi + G(\eta) \psi}{1 + |\nabla \eta|^2}, \quad V = \nabla \psi - B \nabla \eta,$$

and  $\lambda = \lambda^{(1)} + \lambda^{(0)} \in \Gamma_{0,0}^{3/2, \mu-1/2-\tilde{d}} + \Gamma_{0,0}^{1/2, \mu-3/2-\tilde{d}}$ , where

$$\begin{aligned} \lambda^{(1)}(x, \xi) &= \sqrt{(1 + |\nabla \eta|^2) |\xi|^2 - (\nabla \eta \cdot \xi)^2}, \\ \lambda^{(0)}(x, \xi) &= \frac{1 + |\nabla \eta|^2}{2\lambda^{(1)}} \{ \nabla \cdot (\alpha^{(1)} \nabla \eta) + i \partial_\xi \lambda^{(1)} \cdot \nabla \alpha^{(1)} \}, \end{aligned}$$

and

$$\alpha^{(1)}(x, \xi) = \frac{\lambda^{(1)} + i \nabla \eta \cdot \xi}{1 + |\nabla \eta|^2}.$$

Then

$$G(\eta) \psi = \mathcal{P}_\lambda(\psi - \mathcal{P}_B \eta) - \mathcal{P}_V \cdot \nabla \eta + R(\eta, \psi),$$

where  $R(\eta, \psi) \in \mathcal{H}_k^{\mu+1/2, \delta}$ .

We shall define  $\omega = \psi - \mathcal{P}_B \eta$ , which is called the good unknown of Alinhac.

*Proof.* We only sketch the proof, for the key ingredients are already given in [Alazard, Burq and Zuily 2011]. We simply replace the paradifferential calculus in [loc. cit.] by our dyadic paradifferential calculus. Let  $v$  be defined as in Section 5A. Rewrite (5-2) as

$$\alpha \partial_z^2 v + \Delta v + \beta \cdot \nabla \partial_z v - \gamma \partial_z v = 0,$$

where  $\alpha = 1 + |\nabla \eta|^2$ ,  $\beta = -2\nabla \eta$ ,  $\gamma = \Delta \eta$ . Applying Proposition 4.21, we obtain as in [loc. cit., Lemma 3.17],

$$\mathcal{P}_\alpha \partial_z^2 u + \Delta u + \mathcal{P}_\beta \cdot \nabla \partial_z u - \mathcal{P}_\gamma \partial_z u \in C([- \delta, 0], \mathcal{H}_k^{\mu, \delta}), \quad (5-5)$$

where  $u = v - \mathcal{P}_{\partial_z v} \zeta$  with  $\zeta(x, z) = z + \eta(x)$ . Define  $a_\pm = a_\pm^{(1)} + a_\pm^{(0)} \in \Gamma_{0,0}^{1, \mu-1/2-\tilde{d}} + \Gamma_{0,0}^{0, \mu-3/2-\tilde{d}}$  by

$$\begin{aligned} a_\pm^{(1)}(x, \xi) &= \frac{1}{2\alpha} \left( -\beta \cdot \xi \pm \sqrt{4\alpha|\xi|^2 - (\beta \cdot \xi)^2} \right), \\ a_\pm^{(0)}(x, \xi) &= \pm \frac{1}{a_-^{(1)} - a_+^{(1)}} \left( i \partial_\xi a_-^{(1)} \cdot \partial_x a_+^{(1)} - \frac{\gamma}{\alpha} a_\pm^{(1)} \right). \end{aligned}$$

Then we factorize (5-5) as

$$\mathcal{P}_\alpha (\partial_z - \mathcal{P}_{a_-}) (\partial_z - \mathcal{P}_{a_+}) u \in C([- \delta, 0], \mathcal{H}_k^{\mu, \delta}).$$

Because  $\operatorname{Re} a_-^{(1)} \leq 0$ , a parabolic estimate (see e.g., [loc. cit., Proposition 3.19]) implies that

$$(\partial_z u - \mathcal{P}_{a_+} u)|_{z=0} \in \mathcal{H}_k^{\mu+1/2, \delta}.$$

We conclude by setting  $\lambda = (1 + |\nabla \eta|^2) a_+ - i \nabla \eta \cdot \xi$ . □

The proofs of the following results are in the same spirit and much simpler. Their proofs are exactly the same as in [loc. cit.], simply replacing the usual paradifferential calculus with our dyadic paradifferential calculus, particularly Propositions 4.22 and 4.21. Therefore we shall omit the proofs.

**Proposition 5.6.** *Let  $\eta \in \mathcal{H}_k^{\mu+1/2, \delta}$ , with  $\mu - \frac{1}{2} \in \mathbb{N}$ ,  $\mu - \delta k > 3 + \frac{d}{2}$ , and define  $\ell = \ell^{(2)} + \ell^{(1)} \in \Gamma_{0,0}^{2, \mu-1/2-\tilde{d}} + \Gamma_{0,0}^{1, \mu-3/2-\tilde{d}}$ , where*

$$\ell^{(2)} = \frac{(1 + |\nabla \eta|^2) |\xi|^2 - (\nabla \eta \cdot \xi)^2}{(1 + |\nabla \eta|^2)^{3/2}}, \quad \ell^{(1)} = \frac{1}{2} \partial_\xi \cdot D_x \ell^{(2)}.$$

*Then  $H(\eta) = -\mathcal{P}_\ell \eta + f(\eta)$ , where  $f(\eta) \in \mathcal{H}_k^{2\mu-2-d/2, 2\delta}$ .*

**Proposition 5.7.** *Let  $(\eta, \psi) \in \mathcal{H}_k^{\mu+1/2, \delta} \times \mathcal{H}_k^{\mu, \delta}$ , with  $\mu - \frac{1}{2} \in \mathbb{N}$ ,  $\mu - \delta k > 3 + \frac{d}{2}$ . Then*

$$\frac{1}{2} |\nabla \psi| - \frac{1}{2} \frac{(\nabla \eta \cdot \nabla \psi + G(\eta) \psi)^2}{1 + |\nabla \eta|^2} = \mathcal{P}_V \cdot \nabla \psi - \mathcal{P}_B \mathcal{P}_V \cdot \nabla \eta - \mathcal{P}_B G(\eta) \psi + f(\eta, \psi),$$

*where  $f(\eta, \psi) \in \mathcal{H}_k^{2\mu-2-d/2, 2\delta}$ .*

Note that in the above parilinearization results, we do not use the spatial decay of the symbols, as we only require the symbols to be in the classes  $\Gamma_{0,0}^{m,r}$ . These results will only be used in the proof of the Cauchy theorem, where the spatial decay of the symbols is not important. Later when we study the propagation of singularities, we will heavily use the spatial decay of the symbols.

Combining Propositions 5.5, 5.6 and 5.7, we obtain the parilinearization of the water wave system.

**Proposition 5.8.** *Let  $(\eta, \psi) \in \mathcal{H}_k^{\mu+1/2, \delta} \times \mathcal{H}_k^{\mu, \delta}$ , with  $\mu - \frac{1}{2} \in \mathbb{N}$ ,  $\mu - \delta k > 3 + \frac{d}{2}$ . Then  $(\eta, \psi)$  solves the water wave equation if and only if*

$$(\partial_t + \mathcal{P}_V \cdot \nabla + \mathcal{L}) \begin{pmatrix} \eta \\ \psi \end{pmatrix} = f(\eta, \psi)$$

where

$$\mathcal{L} = Q^{-1} \begin{pmatrix} 0 & -\mathcal{P}_\lambda \\ \mathcal{P}_\ell & 0 \end{pmatrix} Q, \quad \text{with } Q = \begin{pmatrix} \text{Id} & 0 \\ -\mathcal{P}_B & \mathcal{P}_\lambda \end{pmatrix},$$

and  $f(\eta, \psi) = Q^{-1} \begin{pmatrix} f_1 \\ f_2 \end{pmatrix} \in \mathcal{H}_k^{\mu+1/2} \times \mathcal{H}_k^\mu$  is defined by

$$f_1 = G(\eta)\psi - \{\mathcal{P}_\lambda(\psi - \mathcal{P}_B\eta) - \mathcal{P}_V \cdot \nabla \eta\},$$

$$f_2 = -\frac{1}{2}|\nabla \psi|^2 + \frac{1}{2} \frac{(\nabla \eta \cdot \nabla \psi + G(\eta)\psi)^2}{1 + |\nabla \eta|^2} + H(\eta) + \mathcal{P}_V \cdot \nabla \psi - \mathcal{P}_B \mathcal{P}_V \cdot \nabla \eta - \mathcal{P}_B G(\eta)\psi + \mathcal{P}_\ell \eta - g\eta.$$

**5C. Symmetrization.**

**Definition 5.9.** For  $T > 0$ ,  $\gamma \in \mathbb{R}$  and two operators  $\mathcal{A}, \mathcal{B} \in L^\infty([0, T], \mathcal{O}_0^\gamma)$ , we say that  $\mathcal{A} \sim_\gamma \mathcal{B}$ , or simply  $\mathcal{A} \sim \mathcal{B}$  when there is no ambiguity of the choice of  $\gamma$ , if

$$\mathcal{A} - \mathcal{B} \in L^\infty([0, T], \mathcal{O}_0^{\gamma-3/2}).$$

By [Alazard, Burq and Zuily 2011], there exist symbols which depend solely on  $\eta$ ,

$$\gamma = \gamma^{(3/2)} + \gamma^{(1/2)}, \quad p = p^{(1/2)} + p^{(-1/2)}, \quad q = q^{(0)},$$

whose principal symbols are explicitly

$$\gamma^{(3/2)} = \sqrt{\ell^{(2)}\lambda^{(1)}}, \quad p^{(1/2)} = (1 + |\nabla \eta|^2)^{-1/2} \sqrt{\lambda^{(1)}}, \quad q^{(0)} = (1 + |\nabla \eta|^2)^{1/4}$$

such that

$$\mathcal{P}_p \mathcal{P}_\lambda \sim_{3/2} \mathcal{P}_\gamma \mathcal{P}_q, \quad \mathcal{P}_q \mathcal{P}_\ell \sim_2 \mathcal{P}_\gamma \mathcal{P}_p, \quad \mathcal{P}_\gamma \sim_{3/2} (\mathcal{P}_\gamma)^*. \tag{5-6}$$

Define the symmetrizer

$$S = \begin{pmatrix} \mathcal{P}_p & 0 \\ 0 & \mathcal{P}_q \end{pmatrix} Q.$$

Then the first two relations in (5-6) can be rephrased as

$$S\mathcal{L} \sim \begin{pmatrix} 0 & -\mathcal{P}_\gamma \\ \mathcal{P}_\gamma & 0 \end{pmatrix} S. \tag{5-7}$$

where the equivalence relation  $\sim$  is applied separately to each component of the matrices.

**5D. Approximate system.** Set the mollifier  $J_\varepsilon = \mathcal{P}_{j_\varepsilon}$ , where  $j_\varepsilon = j_\varepsilon^{(0)} + j_\varepsilon^{(-1)}$ :

$$j_\varepsilon^{(0)} = \exp(-\varepsilon \gamma^{(3/2)}), \quad j_\varepsilon^{(-1)} = \frac{1}{2} \partial_\xi \cdot D_x j_\varepsilon^{(0)}.$$

Then uniformly for  $\varepsilon > 0$ , we have

$$J_\varepsilon \mathcal{P}_\gamma \sim_{3/2} \mathcal{P}_\gamma J_\varepsilon, \quad J_\varepsilon^* \sim_0 J_\varepsilon.$$

Let  $\tilde{p} = \tilde{p}^{(-1/2)} + \tilde{p}^{(-3/2)}$ , with

$$\tilde{p}^{(-1/2)} = \frac{1}{p^{(1/2)}}, \quad \tilde{p}^{(-3/2)} = \frac{-\left(\tilde{p}^{(-1/2)} p^{(-1/2)} + \frac{1}{i} \partial_{\xi} \tilde{p}^{(-1/2)} \cdot \partial_x p^{(1/2)}\right)}{p^{(1/2)}}.$$

Then we have

$$\mathcal{P}_p \mathcal{P}_{\tilde{p}} \sim_0 \text{Id}, \quad \mathcal{P}_q \mathcal{P}_{1/q} \sim_0 \text{Id}.$$

Let

$$\mathcal{L}_{\varepsilon} = \mathcal{L} Q^{-1} \begin{pmatrix} \mathcal{P}_{\tilde{p}} J_{\varepsilon} \mathcal{P}_p & 0 \\ 0 & \mathcal{P}_{1/q} J_{\varepsilon} \mathcal{P}_q \end{pmatrix} Q.$$

Then as in (5-7) we have

$$S \mathcal{L}_{\varepsilon} \sim \begin{pmatrix} 0 & -\mathcal{P}_{\gamma} \\ \mathcal{P}_{\gamma} & 0 \end{pmatrix} J_{\varepsilon} S. \quad (5-8)$$

We define the approximate system

$$(\partial_t + \mathcal{P}_V \cdot \nabla J_{\varepsilon} + \mathcal{L}_{\varepsilon}) \begin{pmatrix} \eta \\ \psi \end{pmatrix} = f(J_{\varepsilon} \eta, J_{\varepsilon} \psi). \quad (5-9)$$

**5E. A priori estimate.** From now on we restrict ourselves to the case where  $\delta = \frac{1}{2}$ . The weighted Sobolev spaces  $\mathcal{H}_k^{\mu+1/2, 1/2} \times \mathcal{H}_k^{\mu, 1/2}$  are the spaces where we do the energy estimates.

**Proposition 5.10.** *Let  $(\eta, \psi) \in C^1([0, T], \mathcal{H}_k^{\mu+1/2, 1/2} \times \mathcal{H}_k^{\mu, 1/2})$ , with  $\mu - \frac{1}{2} \in \mathbb{N}$ ,  $\mu - \frac{k}{2} > 3 + \frac{d}{2}$ , solve the approximate system (5-9). Define*

$$M_T = \sup_{0 \leq t \leq T} \|(\eta, \psi)(t)\|_{\mathcal{H}_k^{\mu+1/2} \times \mathcal{H}_k^{\mu}}, \quad M_0 = \|(\eta, \psi)(0)\|_{\mathcal{H}_k^{\mu+1/2} \times \mathcal{H}_k^{\mu}}.$$

Then there exists some nondecaying function  $C : \mathbb{R}_{\geq 0} \rightarrow \mathbb{R}_{\geq 0}$  such that

$$M_T \leq C(M_0) + T C(M_T).$$

*Proof.* For  $0 \leq j \leq k$ , set

$$M_T^j = \sup_{0 \leq t \leq T} \|(\eta, \psi)(t)\|_{H_j^{\mu+1/2-j/2} \times H_j^{\mu-j/2}},$$

$$M_0^j = \|(\eta, \psi)(0)\|_{H_j^{\mu+1/2-j/2} \times H_j^{\mu-j/2}}.$$

By [Alazard, Burq and Zuily 2011], we know

$$M_T^0 \leq C(M_0^0) + T C(M_T^0).$$

It remains to prove that, for  $1 \leq j \leq k$ , we have

$$M_T^j \leq C(M_0^j) + T C(M_T^j).$$

To do this, let  $\Lambda_j^{\mu} = \mathcal{P}_{m_j^{\mu-j/2}}$ , and set

$$\Phi = \Lambda_j^{\mu} S \begin{pmatrix} \eta \\ \psi \end{pmatrix}.$$

Then

$$(\partial_t + \mathcal{P}_V \cdot \nabla J_{\varepsilon}) \Phi + \begin{pmatrix} 0 & -\mathcal{P}_{\gamma} \\ \mathcal{P}_{\gamma} & 0 \end{pmatrix} J_{\varepsilon} \Phi = F_{\varepsilon},$$

where  $F_\varepsilon = F_\varepsilon^1 + F_\varepsilon^2 + F_\varepsilon^3$ , with

$$\begin{aligned} F_\varepsilon^1 &= \Lambda_k^\mu S f(J_\varepsilon \eta, J_\varepsilon \psi), \\ F_\varepsilon^2 &= [\partial_t + \mathcal{P}_V \cdot \nabla J_\varepsilon, \Lambda_j^\mu S] \begin{pmatrix} \eta \\ \psi \end{pmatrix}, \\ F_\varepsilon^3 &= \begin{pmatrix} 0 & -\mathcal{P}_\gamma \\ \mathcal{P}_\gamma & 0 \end{pmatrix} J_\varepsilon \Lambda_j^\mu S \begin{pmatrix} \eta \\ \psi \end{pmatrix} - \Lambda_j^\mu S \mathcal{L}_\varepsilon \begin{pmatrix} \eta \\ \psi \end{pmatrix}. \end{aligned}$$

By Propositions 5.8, 5.5, 5.6 and 5.7,

$$\|f(J_\varepsilon \eta, J_\varepsilon \psi)\|_{\mathcal{H}_k^{\mu+1/2, 1/2} \times \mathcal{H}_k^{\mu, 1/2}} \leq C(\|(J_\varepsilon \eta, J_\varepsilon \psi)\|_{\mathcal{H}_k^{\mu+1/2, 1/2} \times \mathcal{H}_k^{\mu, 1/2}}) \leq C(\|(\eta, \psi)\|_{\mathcal{H}_k^{\mu+1/2, 1/2} \times \mathcal{H}_k^{\mu, 1/2}}).$$

Therefore,

$$\|F_\varepsilon^1\|_{L^\infty([0, T], L^2)} \leq C(M_T).$$

As  $\mathcal{P}_V \cdot \nabla J_\varepsilon$  is a scalar operator, Proposition 4.19 gives

$$\|[\partial_t + \mathcal{P}_V \cdot \nabla J_\varepsilon, \Lambda_j^\mu S]\|_{L^\infty([0, T], H_j^{\mu+1/2-j/2} \times H_j^{\mu-j/2} \rightarrow L^2 \times L^2)} \leq C(M_T),$$

which implies

$$\|F_\varepsilon^2\|_{L^\infty([0, T], L^2)} \leq C(M_T).$$

By (5-8), the operator

$$\begin{pmatrix} 0 & -\mathcal{P}_\gamma \\ \mathcal{P}_\gamma & 0 \end{pmatrix} J_\varepsilon S - S \mathcal{L}_\varepsilon$$

sends  $H^{\mu+1/2} \times H^\mu$  to  $H^\mu \times H^\mu$ . Unfortunately,

$$\begin{aligned} R &:= \begin{pmatrix} 0 & -\mathcal{P}_\gamma \\ \mathcal{P}_\gamma & 0 \end{pmatrix} J_\varepsilon \Lambda_k^\mu S - \Lambda_k^\mu S \mathcal{L}_\varepsilon \\ &= \begin{pmatrix} 0 & -\mathcal{P}_\gamma \\ \mathcal{P}_\gamma & 0 \end{pmatrix} J_\varepsilon [\Lambda_k^\mu, S] + [S \mathcal{L}_\varepsilon, \Lambda_k^\mu] + \left( \begin{pmatrix} 0 & -\mathcal{P}_\gamma \\ \mathcal{P}_\gamma & 0 \end{pmatrix} J_\varepsilon S - S \mathcal{L}_\varepsilon \right) \Lambda_k^\mu \\ &=: \text{(I)} + \text{(II)} + \text{(III)} \end{aligned}$$

does not send  $H_j^{\mu+1/2-j/2} \times H_j^{\mu-j/2}$  to  $L^2 \times L^2$  because the subprincipal symbol cannot be canceled out in the symbolic calculus, due to the existence of  $\Lambda_j^\mu$ . Particularly, we need to use Proposition 4.19 to estimate the commutators  $[\Lambda_j^\mu, S]$  and  $[S \mathcal{L}_\varepsilon, \Lambda_j^\mu]$ , and obtain

$$\left\| R \begin{pmatrix} \eta \\ \psi \end{pmatrix} \right\|_{L^2 \times L^2} \lesssim \|(\eta, \psi)\|_{H_{j-1}^{\mu+1-j/2} \times H_{j-1}^{\mu+1/2-j/2}} + \|(\eta, \psi)\|_{H_j^{\mu+1/2-j/2} \times H_j^{\mu-j/2}}.$$

More precisely, the first term on the right-hand side comes from (I) and (II), while the second term comes from (III). When  $j \geq 1$ ,

$$H_{j-1}^{\mu+1-j/2} \times H_{j-1}^{\mu+1/2-j/2} = H_{j-1}^{\mu+1/2-(j-1)/2} \times H_{j-1}^{\mu-(j-1)/2} \supset \mathcal{H}_k^{\mu+1/2, 1/2} \times \mathcal{H}_k^{\mu, 1/2},$$

and we deduce that

$$\|F_\varepsilon^3\|_{L^\infty([0, T], L^2)} \leq C(M_T).$$

Finally by the exact same energy estimate as in [Alazard, Burq and Zuily 2011], we conclude that

$$M_T^j \lesssim \|\Phi\|_{L^\infty([0, T], L^2)} \leq C(M_0^j) + TC(M_T). \quad \square$$

**5F. Existence.**

**Lemma 5.11.** *For all  $(\eta_0, \psi_0) \in \mathcal{H}_k^{\mu+1/2, 1/2} \times \mathcal{H}_k^{\mu, 1/2}$ , where  $\mu - \frac{1}{2} \in \mathbb{N}$  and  $\mu - \frac{k}{2} > 3 + \frac{d}{2}$ , and, for all  $\varepsilon > 0$ , the Cauchy problem of the approximate system (5-9) has a unique maximal solution*

$$(\eta_\varepsilon, \psi_\varepsilon) \in C([0, T_\varepsilon), \mathcal{H}_k^{\mu+1/2, 1/2} \times \mathcal{H}_k^{\mu, 1/2}).$$

Moreover, there exists  $T_0 > 0$  such that

$$\inf_{\varepsilon \in (0, 1]} T_\varepsilon \geq T_0.$$

*Proof.* Following [Alazard, Burq and Zuily 2011], the existence follows from the existence theory of ODEs by writing (5-9) in the compact form

$$\partial_t X = \mathcal{F}_\varepsilon(X),$$

where  $\mathcal{F}_\varepsilon$  is a Lipschitz map on  $\mathcal{H}_k^{\mu+1/2, 1/2} \times \mathcal{H}_k^{\mu, 1/2}$ . Indeed,  $J_\varepsilon \in \mathcal{O}_0^{-\infty}$  is a smoothing operator.<sup>1</sup> The estimates to proving the Lipschitz regularity can be carried out much as in the proof of Proposition 5.10. The only nontrivial term that remains is the Dirichlet–Neumann operator, whose regularity follows by combining Proposition 5.4 and the shape derivative formula (which goes back to [Zakharov 1998],

$$\langle dG(\eta)\psi, \varphi \rangle := \lim_{h \rightarrow 0} \frac{1}{h} (G(\eta + h\varphi) - G(\eta))\psi = -G(\eta)(B\varphi) - \nabla \cdot (V\varphi).$$

A standard abstract argument then shows that  $T_\varepsilon$  has a strictly positive lower bound, we refer to [Alazard, Burq and Zuily 2011] for more details. □

*Proof of Theorem 1.6.* By Lemma 5.11, we obtain a sequence  $\{(\eta_\varepsilon, \psi_\varepsilon)\}_{0 < \varepsilon \leq 1}$  which satisfies (5-9) and is uniformly bounded in  $L^\infty([0, T], \mathcal{H}_k^{\mu+1/2, 1/2} \times \mathcal{H}_k^{\mu, 1/2})$  for some  $T > 0$ . By (5-9), the time derivatives  $\{(\partial_t \eta_\varepsilon, \partial_t \psi_\varepsilon)\}_{0 < \varepsilon \leq 1}$  are uniformly bounded in  $L^\infty([0, T], \mathcal{H}_k^{\mu-1, 1/2} \times \mathcal{H}_k^{\mu-3/2, 1/2})$ . By [Alazard, Burq and Zuily 2011], there exists

$$(\eta, \psi) \in C([0, T], H^{\mu+1/2} \times H^\mu), \tag{5-10}$$

which solves (1-5), such that as  $\varepsilon \rightarrow 0$ , we have  $(\eta_\varepsilon, \psi_\varepsilon) \rightarrow (\eta, \psi)$  weakly in  $L^2([0, T], \mathcal{H}_k^{\mu+1/2, 1/2} \times \mathcal{H}_k^{\mu, 1/2})$ , and strongly in  $C([0, T], \mathcal{H}_k^{\mu-1, 1/2} \times \mathcal{H}_k^{\mu-3/2, 1/2})$ . We then prove that, for  $1 \leq j \leq k$ ,

$$\Phi = \Phi(\eta, \psi) := \Lambda_j^\mu S(\eta, \psi) \begin{pmatrix} \eta \\ \psi \end{pmatrix}$$

lies in  $C([0, T], L^2)$ , where  $\Lambda_j^\mu$  is defined in Proposition 5.10, and  $S = S(\eta, \psi)$  is the symmetrizer. Up to an extraction of a subsequence, we may assume by weak convergence that

$$\begin{aligned} (\eta, \psi) &\in L^\infty([0, T], \mathcal{H}_k^{\mu+1/2, 1/2} \times \mathcal{H}_k^{\mu, 1/2}), \\ (\partial_t \eta, \partial_t \psi) &\in L^\infty([0, T], \mathcal{H}_k^{\mu-1, 1/2} \times \mathcal{H}_k^{\mu-3/2, 1/2}), \end{aligned}$$

---

<sup>1</sup>We do not need  $J_\varepsilon \in \mathcal{O}_{-\infty}^-$  because the operators such as  $\mathcal{P}_V \cdot \nabla, \mathcal{L}$ , etc., are all of nonpositive orders with respect to the spatial decay.

with

$$\|(\eta, \psi)\|_{L^\infty([0, T], \mathcal{H}_k^{\mu+1/2, 1/2} \times \mathcal{H}_k^\mu) \cap W^{1, \infty}([0, T], \mathcal{H}_k^{\mu-1, 1/2} \times \mathcal{H}_k^{\mu-3/2})} \leq C(\|(\eta_0, \psi_0)\|_{\mathcal{H}_k^{\mu+1/2, 1/2} \times \mathcal{H}_k^{\mu, 1/2}}).$$

This already implies that  $(\eta, \psi)$  is weakly continuous in  $\mathcal{H}_k^{\mu+1/2, 1/2} \times \mathcal{H}_k^{\mu, 1/2}$ . By the analysis in the previous section,

$$(\partial_t + \mathcal{P}_V \cdot \nabla)\Phi + \begin{pmatrix} 0 & -\mathcal{P}_\gamma \\ \mathcal{P}_\gamma & 0 \end{pmatrix} \Phi = F,$$

with

$$\|F\|_{L^\infty([0, T], L^2)} \leq C(\|(\eta_0, \psi_0)\|_{\mathcal{H}_k^{\mu+1/2, 1/2} \times \mathcal{H}_k^{\mu, 1/2}}).$$

Let  $J_h = \text{Op}_h(e^{-|x|^2 - |\xi|^2})$ . Now that  $e^{-h^2|x|^2 - h^2|\xi|^2} \in S_0^0$ , we have the commutator estimate

$$[J_h, \mathcal{P}_V \cdot \nabla] = \mathcal{O}(1)_{\mathcal{O}^0_{-1}}, \quad [J_h, \mathcal{P}_\gamma] = \mathcal{O}(1)_{\mathcal{O}^{-1/2}}.$$

Because  $k \geq 1$ , by the same spirit of estimating  $R$  in Proposition 5.10, we obtain the energy estimate

$$\frac{d}{dt} \|J_h \Phi(t)\|_{L^2}^2 \leq C(\|(\eta_0, \psi_0)\|_{\mathcal{H}_k^{\mu+1/2, 1/2} \times \mathcal{H}_k^{\mu, 1/2}}).$$

Therefore,  $t \mapsto \|J_h \Phi(t)\|_{L^2}^2$  are uniformly Lipschitzian. Consequently, by the Arzelà–Ascoli theorem,  $t \mapsto \|\Phi(t)\|_{L^2}^2$  is continuous, because  $J_h \Phi \rightarrow \Phi$  as  $h \rightarrow 0$ . Combining the weak continuity, we deduce by functional analysis that  $\Phi \in C([0, T], L^2)$ . By (5-10), the paradifferential calculus, and the definition of  $\Phi$ , we deduce that

$$(\eta, \psi) \in C([0, T], \mathcal{H}_k^{\mu+1/2, 1/2} \times \mathcal{H}_k^{\mu, 1/2}).$$

Thus we finish the proof of Theorem 1.6. □

### 6. Propagation of singularities for water waves

**6A. Finer parilinearization and symmetrization.** To study the propagation of singularities, we need much finer results of parilinearization and symmetrization than Propositions 5.5 and 5.8 so as to gain regularities in the remainder terms.

**Proposition 6.1.** *If  $(\eta, \psi) \in H^{\mu+1/2} \times H^\mu$ , with  $\mu - \frac{1}{2} \in \mathbb{N}$  and  $\mu > 3 + \frac{d}{2}$ , then there exists  $\lambda = \lambda^{(1)} + \lambda^{(0)} + \dots \in \Sigma^{1, \mu-1/2-d}$  such that*

$$G(\eta)\psi = \mathcal{P}_\lambda(\psi - \mathcal{P}_B \eta) - \mathcal{P}_V \cdot \nabla \eta + R(\eta, \psi),$$

where  $R(\eta, \psi) \in H^{2\mu-K-d/2}$  for some  $K > 0$  independent of the dimension  $d$ . Moreover  $\lambda^{(1-j)}$ , when it is defined, is a function of derivatives  $\partial_x^\alpha \eta$ , where  $|\alpha| \leq 1 + j$ .

*Proof.* This theorem follows by replacing the usual paradifferential calculus with the dyadic paradifferential calculus in the analysis of [Alazard and Métivier 2009]. In that work, the explicit expression for  $\lambda$  is given. We write it down for the sake of later applications:

$$\lambda = (1 + |\nabla \eta|^2)a_+ - i \nabla \eta \cdot \xi,$$

where  $a_{\pm} = \sum_{j \leq 1} a_{\pm}^{(j)} \in \Sigma^{1, \mu-1-d/2}$  is defined as follows. Setting  $c = 1/(1 + |\nabla\eta|^2)$ , we have

$$\begin{aligned} a_{-}^{(1)} &= ic\nabla\eta \cdot \xi - \sqrt{c|\xi|^2 - (c\nabla\eta \cdot \xi)^2}, & a_{+}^{(1)} &= ic\nabla\eta \cdot \xi + \sqrt{c|\xi|^2 - (c\nabla\eta \cdot \xi)^2}, \\ a_{-}^{(0)} &= \frac{i\partial_{\xi} a_{-}^{(1)} \cdot \partial_x a_{+}^{(1)} - c\Delta\eta a_{-}^{(1)}}{a_{+}^{(1)} - a_{-}^{(1)}}, & a_{+}^{(0)} &= \frac{i\partial_{\xi} a_{-}^{(1)} \cdot \partial_x a_{+}^{(1)} - c\Delta\eta a_{+}^{(1)}}{a_{-}^{(1)} - a_{+}^{(1)}}. \end{aligned}$$

Suppose that  $a_{\pm}^{(j)}$  are defined for  $m \leq j \leq 1$ . Then we define

$$\begin{aligned} a_{-}^{(m-1)} &= \frac{1}{a_{-}^{(1)} - a_{+}^{(1)}} \sum_{m \leq k \leq 1} \sum_{m \leq \ell \leq 1} \sum_{|\alpha|=k+\ell-m} \frac{1}{\alpha!} \partial_{\xi}^{\alpha} a_{-}^{(k)} D_x^{\alpha} a_{+}^{(\ell)}, \\ a_{+}^{(m-1)} &= -a_{-}^{(m-1)}. \end{aligned}$$

The principal and subprincipal symbols of  $\lambda$  coincide with the ones given by [Proposition 5.5](#). □

**Proposition 6.2.** *Let  $(\eta, \psi) \in H^{\mu+1/2} \times H^{\mu}$ , with  $\mu - \frac{1}{2} \in \mathbb{N}$  and  $\mu > 3 + \frac{d}{2}$ . Let  $\Lambda^{\mu} = \mathcal{P}_{(\gamma^{(3/2)})^{2\mu/3}}$ , and set*

$$w = \Lambda^{\mu} U S \begin{pmatrix} \eta \\ \psi \end{pmatrix}, \quad U = \begin{pmatrix} -i & 1 \\ i & 1 \end{pmatrix}.$$

*Then there exist  $Q \in M_{2 \times 2}(\Sigma_{0,0}^{0, \mu-1/2-2-\bar{d}})$  and  $\zeta \in \Sigma_{0,0}^{-1/2, \mu-1/2-2-\bar{d}}$  such that, for some  $K > 0$  which is independent of the dimension  $d$ , we have*

$$(\partial_t + \mathcal{P}_V \cdot \nabla + \mathcal{P}_Q)w + i\mathcal{P}_{\gamma} \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} w + \frac{ig}{2} \mathcal{P}_{\zeta} \begin{pmatrix} 1 & -1 \\ 1 & -1 \end{pmatrix} \in H^{\mu-K-d/2}. \tag{6-1}$$

**Remark 6.3.** Because  $\chi$  in the definition of paradifferential operators is an even function, we verify that  $\Lambda^{\mu}, \mathcal{P}_p, \mathcal{P}_q, \mathcal{P}_B$  all map real-valued functions to real-valued functions. Therefore,

$$w = \begin{pmatrix} u \\ \bar{u} \end{pmatrix}, \quad \text{with } u = \Lambda^{\mu}(-i, 1) S \begin{pmatrix} \eta \\ \psi \end{pmatrix} = \Lambda^{\mu} \mathcal{P}_q \omega - i\Lambda^{\mu} \mathcal{P}_p \eta, \tag{6-2}$$

recalling that  $\omega = \psi - \mathcal{P}_B \eta$  is the good unknown of Alinhac.

*Proof.* Combining [Propositions 6.1](#) and [5.8](#), and moving the term  $g\eta$  to the left-hand side,

$$(\partial_t + \mathcal{P}_V \cdot \nabla + \mathcal{L}) \begin{pmatrix} \eta \\ \psi \end{pmatrix} + g \begin{pmatrix} 0 \\ \eta \end{pmatrix} = f(\eta, \psi),$$

where

$$f(\eta, \psi) = Q^{-1} \begin{pmatrix} f_1 \\ f_2 \end{pmatrix} \in H^{2\mu+1/2-K-d/2} \times H^{2\mu-K-d/2}$$

for some  $K > 0$  and

$$\begin{aligned} f_1 &= G(\eta)\psi - \{\mathcal{P}_{\lambda}(\psi - \mathcal{P}_B \eta) - \mathcal{P}_V \cdot \nabla \eta\}, \\ f_2 &= -\frac{1}{2}|\nabla\psi|^2 + \frac{1}{2} \frac{(\nabla\eta \cdot \nabla\psi + G(\eta)\psi)^2}{1 + |\nabla\eta|^2} + H(\eta) + \mathcal{P}_V \cdot \nabla\psi - \mathcal{P}_B \mathcal{P}_V \cdot \nabla\eta - \mathcal{P}_B G(\eta)\psi + \mathcal{P}_{\ell}\eta. \end{aligned}$$



Given two time-dependent operators  $\mathcal{A}, \mathcal{B} : \mathcal{S} \rightarrow \mathcal{S}'$ , we say that  $\mathcal{A} \sim \mathcal{B}$  if

$$\mathcal{A} - \mathcal{B} \in L^\infty([0, T], \mathcal{O}_0^{-\mu+d/2+K}).$$

By the ellipticity of  $\gamma^{(3/2)}$ ,  $p^{(1/2)}$  and  $q^{(0)}$ , we can find paradifferential operators  $\tilde{\Lambda}^\mu$  and  $\tilde{S}$  by a routine construction of a parametrix such that  $\tilde{\Lambda}^\mu \Lambda^\mu \sim \text{Id}$ ,  $\tilde{S}S \sim \text{Id}$ . We can find  $\zeta \in \Sigma^{-1/2, \mu-1/2-2-\tilde{d}}$  with principal symbol  $\zeta^{(-1/2)} = q^{(0)}/p^{(1/2)}$ , which implies (note that the only nonzero entries in the following matrices are in the lower left corners)

$$\begin{pmatrix} 0 & 0 \\ \mathcal{P}_\zeta & 0 \end{pmatrix} \Lambda^\mu S - \Lambda^\mu S \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix} \sim 0.$$

Then by (5-7) and the fact that the Poisson bracket between the symbol of  $\Lambda^\mu$  and  $\gamma$  vanishes, we find by the symbolic calculus two symbols  $A, B \in M_{2 \times 2}(\Sigma^{0, \mu-1/2-2-\tilde{d}})$  such that

$$\begin{aligned} \mathcal{A} &:= [\partial_t + \mathcal{P}_V \cdot \nabla, \Lambda^\mu S] \sim [\partial_t + \mathcal{P}_V \cdot \nabla, \Lambda^\mu S] \tilde{S} \tilde{\Lambda}^\mu \Lambda^\mu S \sim \mathcal{P}_A \Lambda^\mu S, \\ \mathcal{B} &:= \begin{pmatrix} 0 & -\mathcal{P}_\gamma \\ \mathcal{P}_\gamma & 0 \end{pmatrix} \Lambda^\mu S - \Lambda^\mu S \mathcal{L} \sim \left( \begin{pmatrix} 0 & -\mathcal{P}_\gamma \\ \mathcal{P}_\gamma & 0 \end{pmatrix} - \Lambda^\mu S \mathcal{L} \tilde{S} \tilde{\Lambda}^\mu \right) \Lambda^\mu S \sim \mathcal{P}_B \Lambda^\mu S. \end{aligned}$$

In fact, by Proposition 4.19, the symbol  $A$  is a finite sum of symbols which is given by the symbolic calculus of the operator  $[\partial_t + \mathcal{P}_V \cdot \nabla, \Lambda^\mu S] \tilde{S} \tilde{\Lambda}^\mu$ , whereas the symbol  $B$  is given by the symbolic calculus of the operator

$$\begin{pmatrix} 0 & -\mathcal{P}_\gamma \\ \mathcal{P}_\gamma & 0 \end{pmatrix} - \Lambda^\mu S \mathcal{L} \tilde{S} \tilde{\Lambda}^\mu.$$

Clearly  $A$  is of zeroth order. The reason why  $B$  is of zeroth order is the condition (5-7) according to which we constructed the symbols  $\gamma, p, q$ .

Let  $\Phi = \Lambda^\mu S \begin{pmatrix} \eta \\ \psi \end{pmatrix}$ , and write

$$g \begin{pmatrix} 0 \\ \eta \end{pmatrix} = \begin{pmatrix} 0 & 0 \\ g & 0 \end{pmatrix} \begin{pmatrix} \eta \\ \psi \end{pmatrix},$$

we obtain by the analysis above that

$$(\partial_t + \mathcal{P}_V \cdot \nabla) \Phi + \begin{pmatrix} 0 & -\mathcal{P}_\gamma \\ \mathcal{P}_\gamma & 0 \end{pmatrix} \Phi + \begin{pmatrix} 0 & 0 \\ g \mathcal{P}_\zeta & 0 \end{pmatrix} \Phi = \mathcal{P}_A \Phi + \mathcal{P}_B \Phi + F,$$

where

$$F = (\mathcal{A} + \mathcal{B}) \begin{pmatrix} \eta \\ \psi \end{pmatrix} - \mathcal{P}_{A+B} \Phi + \begin{pmatrix} 0 & 0 \\ g \mathcal{P}_\zeta & 0 \end{pmatrix} \Phi - g \Lambda^\mu S \begin{pmatrix} 0 \\ \eta \end{pmatrix} + \Lambda^\mu S f(\eta, \psi) \in H^{\mu-K-d/2}.$$

Finally, observe that

$$\begin{aligned} U \begin{pmatrix} 0 & -\mathcal{P}_\gamma \\ \mathcal{P}_\gamma & 0 \end{pmatrix} U^{-1} &= i \begin{pmatrix} \mathcal{P}_\gamma & 0 \\ 0 & -\mathcal{P}_\gamma \end{pmatrix}, \\ U \begin{pmatrix} 0 & 0 \\ \mathcal{P}_\zeta & 0 \end{pmatrix} U^{-1} &= \frac{i}{2} \begin{pmatrix} \mathcal{P}_\zeta & -\mathcal{P}_\zeta \\ \mathcal{P}_\zeta & -\mathcal{P}_\zeta \end{pmatrix}, \end{aligned}$$

We conclude by setting

$$Q = -\frac{1}{2} U(A + B)U^{-1}. \quad \square$$

**Remark 6.4.** By Proposition 6.1 and the symbolic calculus, the symbols that we have encountered, such as  $\lambda$ ,  $\zeta$  and  $Q$  etc., are of the form  $a = a^{(m)} + a^{(m-1)} + \dots$  such that  $a^{(m-j)}$ , whenever it is defined, is a function of  $(\nabla\eta, \dots, \nabla^{j+1}\eta)$ . To be precise  $a^{(m-j)} = f_j(\nabla\eta, \dots, \nabla^{j+1}\eta, \xi)$ , where  $f_j$  is homogeneous of degree  $m-j$  in  $\xi$  and  $f_0(0, \dots, 0, \xi) = |\xi|^m$ ,  $f_j(0, \dots, 0, \xi) = 0$  for  $j \geq 1$ . Note that if  $\eta \in \mathcal{H}_k^{\mu+1/2, 1/2}$ , then for all  $j \leq \mu + \frac{1}{2} - \tilde{d}$ , we have  $\nabla^j \eta \in \mathcal{H}_k^{\mu+1/2-j, 1/2}$ . Therefore, by Lemma 4.15,

$$\nabla^j \eta \in W_{0,1}^{\min\{[2(\mu+1/2-j-\tilde{d})/3], k\}, \infty} \cap \langle x \rangle^{-\min\{2(\mu+1/2-j-\tilde{d}), k\}} L^\infty, \tag{6-3}$$

and consequently

$$a^{(m)} - |\xi|^m \in \Gamma_{-\min\{2\mu-1-2\tilde{d}, k\}, 0}^{m, 0}, \quad a^{(m-j)} \in \Gamma_{-\min\{2\mu-1-2j-2\tilde{d}, k\}, 0}^{m-j, 0}. \tag{6-4}$$

As another consequence of (6-3), we also have

$$\begin{aligned} a^{(m)} - |\xi|^m &\in \Gamma_{0,1}^{m, \min\{[2(\mu-1/2-\tilde{d})/3], k\}}, \\ a^{(m-j)} &\in \Gamma_{-j,1}^{m-j, \min\{[2(\mu-1/2-j-\tilde{d})/3], k\}} \subset \Gamma_{-j,1}^{m-j, \min\{[2(\mu-1/2-\tilde{d})/3], k\}-j}. \end{aligned} \tag{6-5}$$

**Lemma 6.5.** Let  $u$  be defined as in (6-2). If  $(\eta, \psi) \in H^{\mu+1/2} \times H^\mu$ , with  $\mu - \frac{1}{2} \in \mathbb{N}$ , then, for  $0 \leq \sigma \leq r \in \mathbb{N}$ , with  $r < \mu - \frac{1}{2} - 1 - \tilde{d}$ ,

$$\text{WF}_{0,1}^\sigma(u)^\circ = \text{WF}_{0,1}^{\mu+1/2+\sigma}(\eta)^\circ \cup \text{WF}_{0,1}^{\mu+\sigma}(\psi)^\circ.$$

If  $(\eta, \psi) \in \mathcal{H}_k^{\mu+1/2} \times \mathcal{H}_k^\mu$ , with  $k \leq \frac{2}{3}(\mu - 1 - \tilde{d})$ , then, for  $0 \leq \sigma \leq \frac{3}{2}k$ ,

$$\text{WF}_{1/2,1}^\sigma(u)^\circ = \text{WF}_{1/2,1}^{\mu+1/2+\sigma}(\eta)^\circ \cup \text{WF}_{1/2,1}^{\mu+\sigma}(\psi)^\circ.$$

*Proof.* Clearly if  $\eta \in H^{\mu+1/2}$ , then  $(\gamma^{(3/2)})^{2\mu/3} \in \Gamma^{\mu,r}$ ,  $p^{(1/2)} \in \Gamma^{1/2,r}$ ,  $q^{(0)} \in \Gamma^{0,r}$ ,  $B \in \Gamma^{0,r}$ . By (6-5), if  $\eta \in \mathcal{H}_k^{\mu+1/2}$ , then  $(\gamma^{(3/2)})^{2\mu/3} \in \Gamma_{0,1}^{\mu,k}$ ,  $p^{(1/2)} \in \Gamma_{0,1}^{1/2,k}$ ,  $q^{(0)} \in \Gamma_{0,1}^{0,k}$ ,  $B \in \Gamma_{0,1}^{0,k}$ . By Lemma 4.42 and (6-2), for either  $\epsilon = 0$  or  $\epsilon = \frac{1}{2}$ ,

$$\begin{aligned} \text{WF}_{\epsilon,1}^\sigma(u)^\circ &= \text{WF}_{\epsilon,1}^\sigma(\Lambda^\mu \mathcal{P}_p \eta)^\circ \cup \text{WF}_{\epsilon,1}^\sigma(\Lambda^\mu \mathcal{P}_q (\psi - \mathcal{P}_B \eta))^\circ \\ &= \text{WF}_{\epsilon,1}^{\mu+1/2+\sigma}(\eta)^\circ \cup \text{WF}_{\epsilon,1}^{\mu+\sigma}(\psi - \mathcal{P}_B \eta)^\circ \\ &\subset \text{WF}_{\epsilon,1}^{\mu+1/2+\sigma}(\eta)^\circ \cup (\text{WF}_{\epsilon,1}^{\mu+\sigma}(\psi)^\circ \cup \text{WF}_{\epsilon,1}^{\mu+\sigma}(\mathcal{P}_B \eta)^\circ) \\ &\subset \text{WF}_{\epsilon,1}^{\mu+1/2+\sigma}(\eta)^\circ \cup (\text{WF}_{\epsilon,1}^{\mu+\sigma}(\psi)^\circ \cup \text{WF}_{\epsilon,1}^{\mu+\sigma}(\eta)^\circ) \\ &= \text{WF}_{\epsilon,1}^{\mu+1/2+\sigma}(\eta)^\circ \cup \text{WF}_{\epsilon,1}^{\mu+\sigma}(\psi)^\circ. \end{aligned}$$

Conversely, as  $\text{WF}_{\epsilon,1}^{\mu+\sigma}(\mathcal{P}_B \eta)^\circ \subset \text{WF}_{\epsilon,1}^{\mu+1/2+\sigma}(\eta)^\circ$ , we have

$$\begin{aligned} \text{WF}_{\epsilon,1}^{\mu+1/2+\sigma}(\eta)^\circ \cup \text{WF}_{\epsilon,1}^{\mu+\sigma}(\psi)^\circ &= \text{WF}_{\epsilon,1}^{\mu+1/2+\sigma}(\eta)^\circ \cup (\text{WF}_{\epsilon,1}^{\mu+\sigma}(\psi)^\circ \setminus \text{WF}_{\epsilon,1}^{\mu+1/2+\sigma}(\eta)^\circ) \\ &= \text{WF}_{\epsilon,1}^{\mu+1/2+\sigma}(\eta)^\circ \cup (\text{WF}_{\epsilon,1}^{\mu+\sigma}(\psi - \mathcal{P}_B \eta)^\circ \setminus \text{WF}_{\epsilon,1}^{\mu+1/2+\sigma}(\eta)^\circ) \\ &= \text{WF}_{\epsilon,1}^{\mu+1/2+\sigma}(\eta)^\circ \cup \text{WF}_{\epsilon,1}^{\mu+\sigma}(\psi - \mathcal{P}_B \eta)^\circ \\ &= \text{WF}_{\epsilon,1}^\sigma(u)^\circ. \end{aligned}$$

The lemma follows. □

**6B. Proof of Theorem 1.7.** By Lemma 6.5, it is equivalent to prove the following theorem.

**Theorem 6.6.** Under the hypothesis of Theorem 1.7, let  $u$  be defined by (6-2), and let

$$(x_0, \xi_0) \in \text{WF}_{1/2,1}^\sigma(u_0)^\circ,$$

with  $0 \leq \sigma < \frac{k}{2} - \frac{3}{2}$ . Let  $t_0 \in [0, T]$ , and suppose that

$$x_0 + \frac{3}{2}t|\xi_0|^{-1/2}\xi_0 \neq 0$$

for all  $t \in [0, t_0]$ . Then

$$(x_0 + \frac{3}{2}t_0|\xi_0|^{-1/2}\xi_0, \xi_0) \in \text{WF}_{1/2,1}^\sigma(u(t_0))^\circ.$$

*Proof.* For  $v \in \mathbb{R}$ , define

$$X^v = \sum_{k \in \mathbb{Z}} H_k^{v-k/2}.$$

By Lemma 2.15, if  $f \in X^v$ , then  $\text{WF}_{1/2,1}^v(f)^\circ = \emptyset$ . Also note that if  $f \in X^v$  and  $a \in \Sigma_{0,1}^{m,r}$ , then  $\mathcal{P}_a f \in X^{v-m}$ . As  $k < 2\mu - d$ , we have  $V \in \mathcal{H}_k^\mu \subset \langle x \rangle^k H^{\mu-k/2} \subset \langle x \rangle^{-k} L^\infty$ , which implies

$$\mathcal{P}_V \cdot \nabla w \subset \mathcal{P}_V H^{-1} \subset H_k^{-1} \subset X^{k/2-1}.$$

By Remark 6.4, particularly (6-4),

$$\mathcal{P}_Q w \in \sum_{j < \mu - \tilde{d}} H_{\min\{2\mu-1-2j-2\tilde{d}, k\}}^j \subset \sum_{j < \mu - \tilde{d}} X^{\min\{\mu-1-\tilde{d}, j+k/2\}} \subset X^{k/2}.$$

Similarly

$$\begin{aligned} \mathcal{P}_\gamma w - \mathcal{P}_{|\xi|^{3/2}} w &\in \sum_{j < \mu - \tilde{d}} H_{\min\{2\mu-1-2j-2\tilde{d}, k\}}^{j-3/2} \subset X^{k/2-3/2}, \\ \mathcal{P}_\zeta w - \mathcal{P}_{|\xi|^{-1/2}} w &\in \sum_{j < \mu - \tilde{d}} H_{\min\{2\mu-1-2j-2\tilde{d}, k\}}^{j+1/2} \subset X^{k/2+1/2}. \end{aligned}$$

By the hypothesis on  $m$ , we thus obtain

$$\partial_t w' + i|D_x|^{3/2} \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} w' + \frac{ig}{2}|D_x|^{-1/2} \begin{pmatrix} 1 & -1 \\ 1 & -1 \end{pmatrix} w' \in X^{k/2-3/2}, \tag{6-6}$$

where  $w' = \pi(D_x)w$ , and  $\pi \in C^\infty(\mathbb{R}^d)$ , which vanishes near the origin, and equals 1 outside a neighborhood of the origin. Moreover, we require that  $\text{supp } \pi \subset \{\tilde{\pi} = 1\}$  such that  $1 - \tilde{\pi} \in C_c^\infty(\mathbb{R}^d)$  and  $\tilde{\pi}(\xi) = 0$  if  $|\xi|^2 \leq |g|$ . Observe that the matrix

$$M = |\xi|^{3/2} \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} + \frac{g}{2} |\xi|^{-1/2} \begin{pmatrix} 1 & -1 \\ 1 & -1 \end{pmatrix}$$

is symmetrizable when restricted to  $\text{supp } \pi$ . Indeed, let

$$P = \frac{1}{2} \begin{pmatrix} 1 + \theta & 1 - \theta \\ -(1 - \theta) & -(1 + \theta) \end{pmatrix},$$

where  $\theta = \sqrt{\tilde{\pi}(\xi) \cdot (g|\xi|^{-2} + 1)}$ . Then  $P \in \mathcal{O}_0^0$ . For  $\xi \in \text{supp } \pi$ , we have

$$PMP^{-1} = |\xi|^{3/2}\theta(\xi) \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}.$$

Set

$$\tilde{w} = P(D_x)w' = P(D_x) \begin{pmatrix} u' \\ \bar{u}' \end{pmatrix} = \begin{pmatrix} \text{Re } u' + i\theta(D_x) \text{Im } u' \\ -\text{Re } u' + i\theta(D_x) \text{Im } u' \end{pmatrix},$$

where  $u' = \pi(D_x)u$ , then

$$\partial_t \tilde{w} + |D_x|^{3/2}\theta(D_x) \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \tilde{w} \in X^{k/2-3/2}.$$

Finally, let  $v = \text{Re } u' + i\theta(D_x) \text{Im } u'$ . Then  $\text{WF}_{1/2,1}^\sigma(u)^\circ = \text{WF}_{1/2,1}^\sigma(v)^\circ$ , and

$$\partial_t v + |D_x|^{3/2}\theta(D_x)v \in X^{k/2-3/2}.$$

We are left to prove that if  $(x_0, \xi_0) \in \text{WF}_{1/2,1}^\sigma(v(0))^\circ$ , then

$$(x_0 + \frac{3}{2}t_0|\xi_0|^{-1/2}\xi_0, \xi_0) \in \text{WF}_{1/2,1}^\sigma(v(t_0)).$$

Because  $\theta(\xi) \sim 1$  in the high-frequency regime, a proof similar to that of [Theorem 1.4\(1\)](#) yields the conclusion. □

### 6C. Proof of [Theorem 1.9](#).

**6C1. Hamiltonian flow.** Let  $\Phi = \Phi_s : \mathbb{R}^d \times (\mathbb{R}^d \setminus 0) \rightarrow \mathbb{R}^d \times (\mathbb{R}^d \setminus 0)$  be the Hamiltonian flow of

$$H(x, \xi) = \gamma^{(3/2)}(0, x, \xi) = \left( |\xi|^2 - \frac{(\nabla\eta_0 \cdot \xi)^2}{1 + |\nabla\eta_0|^2} \right)^{3/4}.$$

That is

$$\partial_s \Phi_s(x, \xi) = X_H(\Phi_s(x, \xi)), \quad \Phi|_{s=0} = \text{Id}_{\mathbb{R}^d \times (\mathbb{R}^d \setminus 0)},$$

where  $X_H = (\partial_\xi H, -\partial_x H)$ . We use  $s$  to denote the time variable in accordance to the semiclassical time variable in the following section. Observe that:

**Lemma 6.7.** For  $(x, \xi) \in \mathbb{R}^d \times (\mathbb{R}^d \setminus 0)$ , we have

$$\Phi_s(x, \xi) = \mathcal{G}_{\varphi_s(x, \xi)}(x, \xi),$$

where  $\mathcal{G}$  is the geodesic flow defined in [Section 1C5](#), and

$$\varphi_s(x, \xi) = \frac{3}{4} \int_0^s G(\Phi_\sigma(x, \xi))^{-1/4} d\sigma.$$

*Proof.* We have  $\mathcal{G}_{\varphi_0(x, \xi)}(x, \xi) = \mathcal{G}_0(x, \xi) = (x, \xi) = \Phi_0(x, \xi)$ . Then observe that

$$H(x, \xi) = G(x, \xi)^{3/4} = \varrho_x^{-1}(\xi, \xi)^{3/4}.$$

Therefore,

$$\begin{aligned} \frac{d}{ds} \mathcal{G}_{\varphi_s(x,\xi)}(x, \xi) &= \frac{d}{ds} \varphi_s(x, \xi) \left( \frac{d}{ds} \mathcal{G} \right)_{\varphi_s(x,\xi)}(x, \xi) \\ &= \frac{3}{4} G(\mathcal{G}_{\varphi_s(x,\xi)}(x, \xi))^{-1/4} X_G(\mathcal{G}_{\varphi_s(x,\xi)}(x, \xi)) \\ &= X_H(\mathcal{G}_{\varphi_s(x,\xi)}(x, \xi)). \end{aligned}$$

We conclude by the uniqueness of solutions to Hamiltonian ODEs. □

**Lemma 6.8.** *Suppose that, for some  $\epsilon > 0$ ,  $\nabla \eta_0 \in W_{1/2+\epsilon}^{0,\infty}$ ,  $\nabla^2 \eta_0 \in W_{1+\epsilon}^{0,\infty}$ . Let  $(x_0, \xi_0) \in \mathbb{R}^d \times (\mathbb{R}^d \setminus 0)$  such that the cogeodesic  $\{(x_s, \xi_s) = \Phi_s(x_0, \xi_0)\}_{s \in \mathbb{R}}$  is forwardly nontrapping. Set*

$$z_s = x_s - x_0 - \frac{3}{2} \int_0^s |\xi_\sigma|^{-1/2} \xi_\sigma \, d\sigma.$$

Then there exists  $(z_{+\infty}, \xi_{+\infty}) \in \mathbb{R}^d \times (\mathbb{R}^d \setminus 0)$  such that

$$\lim_{s \rightarrow +\infty} (z_s, \xi_s) = (z_{+\infty}, \xi_{+\infty}).$$

Consequently, by Lemma 6.7, let  $(x'_s, \xi'_s) = \mathcal{G}_s(x_0, \xi_0)$ , and then

$$\lim_{s \rightarrow +\infty} \xi'_s = \xi_{+\infty}.$$

*Proof.* Because  $\{(x_s, \xi_s)\}_{s \in \mathbb{R}}$  is forwardly nontrapping and we only consider the limiting behavior when  $s \rightarrow +\infty$ , we may assume that  $\epsilon_0 := \|\langle x \rangle \nabla^2 \eta_0\|_{L^\infty}$  is sufficiently small. As  $\nabla \eta_0 \in L^\infty$ , we have  $H(\cdot, \xi) \simeq |\xi|^{3/2}$ . Then

$$\frac{d}{ds} (x_s \cdot \xi_s) = \partial_\xi H(x_s, \xi_s) \cdot \xi_s - x_s \cdot \partial_x H(x_s, \xi_s),$$

where

$$\partial_\xi H(x_s, \xi_s) \cdot \xi_s = \frac{3}{2} H(x_s, \xi_s) = \frac{3}{2} H(x_0, \xi_0) \simeq |\xi_0|^{3/2}$$

and

$$\begin{aligned} \partial_x H(x_s, \xi_s) &= \frac{3}{4} H(x_s, \xi_s)^{-1/3} \partial_x G(x_s, \xi_s) \\ &= \frac{3}{4} H(x_s, \xi_s)^{-1/3} \left( \frac{2 \nabla \eta_0 \cdot \xi_s}{1 + |\nabla \eta_0|^2} \nabla^2 \eta_0 \xi_s - \frac{2 (\nabla \eta_0 \cdot \xi_s)^2}{(1 + |\nabla \eta_0|^2)^2} \nabla^2 \eta_0 \nabla \eta_0 \right) \Big|_{x=x_s}. \end{aligned}$$

Therefore

$$x_s \cdot \partial_x H(x_s, \xi_s) = \mathcal{O}(\epsilon |\xi_s|^{3/2}) = \mathcal{O}(\epsilon |\xi_0|^2),$$

and consequently,

$$\frac{d}{ds} (x_s \cdot \xi_s) \gtrsim |\xi_0|^{3/2}. \tag{6-7}$$

So, for any bounded set  $B \subset \mathbb{R}^d$ ,

$$\lambda(s \geq 0 : x_s \in B) \lesssim \frac{\sup\{|x \cdot \xi| : (x, \xi) \in B \times \mathbb{R}^d, H(x, \xi) = H(x_0, \xi_0)\}}{|\xi_0|^{3/2}} \lesssim \sup_{x \in B} |x| |\xi_0|^{-1/2}, \tag{6-8}$$

where  $\lambda$  is the Lebesgue measure on  $\mathbb{R}$ . Let

$$E(x, \xi) = H(x, \xi) - |\xi|^{3/2}.$$

Then by the hypothesis of the decay of  $\eta_0$ , we have  $E \in \Gamma_{-1-\epsilon, 0}^{3/2, 1}$ . By the definition of  $z_s$ , we have

$$\frac{d}{ds}(z_s, \xi_s) = (\partial_\xi E, -\partial_x E)(x_s, \xi_s) = \mathcal{O}(\langle x_s \rangle^{-1-\epsilon}),$$

where we used the conversation of  $H(x_s, \xi_s)$  to deduce the boundedness of  $\xi_s$ . By (6-8),

$$\int_0^\infty \langle x_s \rangle^{-1-\epsilon} ds = (1 + \epsilon) \int_0^\infty t^\epsilon \lambda(s \geq 0 : \langle x_s \rangle^{-1} > t) dt \lesssim \int_0^1 t^\epsilon \sqrt{t^{-2} - 1} dt < \infty.$$

Therefore, for any  $0 < s^- < s^+$  with  $s^- \rightarrow \infty$ ,

$$|(z_{s^+}, \xi_{s^+}) - (z_{s^-}, \xi_{s^-})| \lesssim \int_{s^-}^{s^+} \langle x_\sigma \rangle^{-1-\epsilon} d\sigma \rightarrow 0,$$

implying that  $(x_s, \xi_s)$  is a Cauchy sequence as  $s \rightarrow \infty$ . □

**6C2. Construction of symbol.** For  $h \geq 0$ , and  $h^{1/2}s \leq T$ . Set

$$H_h(s, x, \xi) = \gamma^{(3/2)}(h^{1/2}s, x, \xi),$$

so in particular  $H(x, \xi) \equiv H_0(s, x, \xi)$ . For  $h > 0$ , the semiclassical time variable  $s = h^{-1/2}t$  was inspired by Lebeau [Lebeau 1992]; see also [Zhu 2020] for an application in theory of control for water waves.

For  $a \in C^\infty([0, h^{-1/2}T) \times \mathbb{R}^{2d})$ , set

$$\mathcal{L}_{h,s}^\pm a = \partial_s a \pm \{H_h, a\}.$$

**Lemma 6.9.** *Suppose that, for some  $\epsilon > 0$ ,  $\nabla \eta_0 \in W_{1/2+\epsilon}^{0, \infty}$ ,  $\nabla^2 \eta_0 \in W_{1+\epsilon}^{0, \infty}$ ,  $\nabla^3 \eta_0 \in W_{3/2+\epsilon}^{0, \infty}$ . Let  $(x_0, \xi_0) \in \mathbb{R}^d \times (\mathbb{R}^d \setminus \{0\})$  such that the cogeodesic  $\{(x_s, \xi_s) = \Phi_s(x_0, \xi_0)\}_{s \in \mathbb{R}}$  is forwardly nontrapping. Then there exists  $s_0 > 0$ ,  $K > 0$  and*

$$\chi^\pm \in W^{1, \infty}(\mathbb{R}_{\geq 0}, \Upsilon^{\mu-K-\bar{d}}) \cap W^{1, \infty}(\mathbb{R}_{\geq s_0}, S_0^{-\infty}) \tag{6-9}$$

in the sense that

$$\|N^{\mu-K-\bar{d}}(\chi^\pm)\|_{L^\infty(\mathbb{R}_{\geq 0})} + \|N^{\mu-K-\bar{d}}(\partial_s \chi^\pm)\|_{L^\infty(\mathbb{R}_{\geq 0})} < +\infty,$$

and satisfies the following conditions:

- (1)  $\chi^\pm(0, x, \xi) \in S_{-\infty}^-$  is elliptic at  $(x_0, \pm \xi_0)$ .
- (2) For all  $t_0 > 0$ ,  $\chi^\pm(s, \frac{s}{t_0}x, \xi) \in S_{-\infty}^-$  is elliptic at  $(\frac{3}{2}t_0|\xi_\infty|^{-1/2}\xi_\infty, \pm \xi_\infty)$  for sufficiently large  $s$ .
- (3) If  $\Omega$  is a neighborhood of  $(\frac{3}{2}t_0|\xi_\infty|^{-1/2}\xi_\infty, \pm \xi_\infty)$ , then  $\chi^\pm$  can be chosen such that

$$\text{supp } \chi^\pm \left( s, \frac{s}{t_0}x, \xi \right) \subset \Omega$$

for sufficiently large  $s$ .

Moreover, if  $(\eta, \psi) \in \mathcal{H}_k^{\mu+1/2} \times \mathcal{H}_k^\mu$ , with  $\mu > 3 + \frac{d}{2}$  and  $m \geq 2$ , then

$$\mathcal{L}_{h,s}^\pm \chi^\pm \in L^\infty([0, h^{-1/2}T], \langle x \rangle^{-1} \Upsilon^{\mu-K-\bar{d}-1})$$

and

$$\mathcal{L}_{h,s}^\pm \chi^\pm \geq \mathcal{O}(h^{1/2})_{L^\infty([0, h^{-1/2}T], \langle x \rangle^{-1} \Upsilon^{\mu-K-\bar{d}-1})}. \tag{6-10}$$

*Proof.* Let  $\phi \in C_c^\infty(\mathbb{R}^d)$  such that

- (i)  $\phi \geq 0$ ,  $\phi(x) = 1$  for  $|x| \leq \frac{1}{2}$  and  $\phi(x) = 0$  for  $|x| \geq 1$ ,  $\text{supp } \phi = \{|x| \leq 1\}$ ,
- (ii)  $x \cdot \nabla \phi(x) \leq 0$  for all  $x \in \mathbb{R}^d$ ,
- (iii)  $y \cdot \nabla \phi(x) = 0$  for all  $x, y \in \mathbb{R}^d$ , with  $x \cdot y = 0$ .

Such  $\phi$  can be constructed by setting  $\phi(x) = \varphi(|x|)$ , where  $\varphi : \mathbb{R} \rightarrow \mathbb{R}$  satisfies  $0 \leq \varphi \leq 1$ ,  $\varphi(z) = 1$  if  $z \leq \frac{1}{2}$ ,  $\varphi(z) = 0$  if  $z \geq 1$ . For  $\rho > 0$ ,  $\delta > 0$ ,  $\lambda > 0$ ,  $\nu > 0$  and sufficiently large  $s > 0$ , set

$$\tilde{\chi}^\pm(s, x, \xi) = \phi\left(\frac{x - x_s}{\rho\lambda\delta s}\right)\phi\left(\frac{\xi \mp \xi_s}{\rho(\delta - s^{-\nu})}\right).$$

We verify that  $\mathcal{L}_{0,s}^\pm \tilde{\chi}^\pm(s, \cdot) \geq 0$  for  $s > 0$  sufficient large. Indeed,

$$\begin{aligned} \mathcal{L}_{0,s}^\pm \tilde{\chi}^\pm(s, x, \xi) &= \left(\pm \frac{\partial_\xi H(x, \xi) - \partial_\xi H(x_s, \mp \xi_s)}{\rho\lambda\delta s} - \frac{x - x_s}{\rho\lambda\delta s^2}\right) \nabla \phi\left(\frac{x - x_s}{\rho\lambda\delta s}\right) \phi\left(\frac{\xi \mp \xi_s}{\rho(\delta - s^{-\nu})}\right) \\ &\quad + \left(\pm \frac{\partial_x H(x_s, \mp \xi_s) - \partial_x H(x, \xi)}{\rho(\delta - s^{-\nu})} - \nu \frac{\xi \mp \xi_s}{\rho(\delta - s^{-\nu})^2 s^{\nu+1}}\right) \phi\left(\frac{x - x_s}{\rho\lambda\delta s}\right) \nabla \phi\left(\frac{\xi \mp \xi_s}{\rho(\delta - s^{-\nu})}\right). \end{aligned}$$

By (i),

$$\begin{aligned} \text{supp } \phi\left(\frac{\cdot - x_s}{\rho\lambda\delta s}\right) &\subset \{x \in \mathbb{R}^d : |x - x_s| \leq \rho\lambda\delta s\}, \\ \text{supp } \phi\left(\frac{\cdot \mp \xi_s}{\rho(\delta - s^{-\nu})}\right) &\subset \{\xi \in \mathbb{R}^d : |\xi \mp \xi_s| \leq \rho(\delta - s^{-\nu})\}, \\ \text{supp } \nabla \phi\left(\frac{\cdot - x_s}{\rho\lambda\delta s}\right) &\subset \{x \in \mathbb{R}^d : \frac{1}{2}\rho\lambda\delta s \leq |x - x_s| \leq \rho\lambda\delta s\}, \\ \text{supp } \nabla \phi\left(\frac{\cdot \mp \xi_s}{\rho(\delta - s^{-\nu})}\right) &\subset \{\xi \in \mathbb{R}^d : \frac{1}{2}\rho(\delta - s^{-\nu}) \leq |\xi \mp \xi_s| \leq \rho(\delta - s^{-\nu})\}. \end{aligned}$$

By Lemma 6.8,

$$x_s = x_0 + \frac{3}{2} \int_0^s |\xi_\sigma|^{-1/2} \xi_\sigma \, d\sigma + z_s = \frac{3}{2} s |\xi_\infty|^{-1/2} \xi_\infty + o(s).$$

Therefore, by writing

$$\tilde{\chi}^\pm\left(s, \frac{s}{t_0}x, \xi\right) = \phi\left(\frac{x - \frac{3}{2}t_0|\xi_\infty|^{-1/2}\xi_\infty + o(1)}{\rho\lambda\delta t_0}\right)\phi\left(\frac{\xi \mp \xi_\infty + o(1)}{\rho(\delta - s^{-\nu})}\right),$$

we see that  $\tilde{\chi}^\pm(s, \frac{s}{t_0}x, \xi)$  is elliptic at  $(\frac{3}{2}t_0|\xi_\infty|^{-1/2}\xi_\infty, \pm\xi_\infty)$  for sufficiently large  $s$ . Moreover, if  $\rho\lambda\delta$  is sufficiently small and  $s$  is sufficiently large, then

$$\text{supp } \phi\left(\frac{\cdot - x_s}{\rho\lambda\delta s}\right) \subset \{x \in \mathbb{R}^d : |x| \gtrsim s\}.$$

Therefore, by the hypothesis on  $\eta_0$ , we have, for  $(x, \xi) \in \text{supp } \tilde{\chi}^\pm(s, \cdot)$ ,

$$\nabla_{x\xi}^2 H(x, \xi) = \begin{pmatrix} \nabla_x^2 H & \nabla_x \nabla_\xi H \\ \nabla_\xi \nabla_x H & \nabla_\xi^2 H \end{pmatrix} (x, \xi) = \begin{pmatrix} \mathcal{O}(s^{-2-\epsilon}) & \mathcal{O}(s^{-3/2-\epsilon}) \\ \mathcal{O}(s^{-3/2-\epsilon}) & \mathcal{O}(1) \end{pmatrix},$$

and consequently, by the finite increment formula,

$$\begin{aligned} |\partial_{\xi} H(x_s, \mp \xi_s) - \partial_{\xi} H(x, \xi)| &\lesssim s^{-3/2-\epsilon} |x - x_s| + |\xi \mp \xi_s| \lesssim s^{-1/2-\epsilon} \rho \lambda \delta + \rho \delta, \\ |\partial_x H(x_s, \mp \xi_s) - \partial_x H(x, \xi)| &\lesssim s^{-2-\epsilon} |x - x_s| + s^{-3/2-\epsilon} |\xi \mp \xi_s| \lesssim \rho \lambda \delta s^{-1-\epsilon} + \rho \delta s^{-3/2-\epsilon}. \end{aligned}$$

By (iii) and the estimates above,

$$\begin{aligned} &(\partial_{\xi} H(x, \xi) - \partial_{\xi} H(x_s, \mp \xi_s)) \cdot \nabla \phi \left( \frac{x - x_s}{\rho \lambda \delta s} \right) \\ &= (\partial_{\xi} H(x, \xi) - \partial_{\xi} H(x_s, \mp \xi_s)) \cdot \frac{x - x_s}{|x - x_s|^2} (x - x_s) \cdot \nabla \phi \left( \frac{x - x_s}{\rho \lambda \delta s} \right) \\ &= \mathcal{O}(s^{-3/2-\epsilon} + \lambda^{-1} s^{-1}) (x - x_s) \cdot \nabla \phi \left( \frac{x - x_s}{\rho \lambda \delta s} \right), \\ &(\partial_x H(x_s, \mp \xi_s) - \partial_x H(x, \xi)) \cdot \nabla \phi \left( \frac{\xi \mp \xi_s}{\rho(\delta - s^{-\nu})} \right) \\ &= (\partial_x H(x_s, \mp \xi_s) - \partial_x H(x, \xi)) \cdot \frac{\xi \mp \xi_s}{|\xi \mp \xi_s|^2} (\xi \mp \xi_s) \cdot \nabla \phi \left( \frac{\xi \mp \xi_s}{\rho(\delta - s^{-\nu})} \right) \\ &= \mathcal{O}(\lambda s^{-1-\epsilon} + s^{-3/2-\epsilon}) (\xi \mp \xi_s) \cdot \nabla \phi \left( \frac{\xi \mp \xi_s}{\rho(\delta - s^{-\nu})} \right). \end{aligned}$$

Finally, we fix  $0 < \nu < \epsilon$ ,  $\delta > 0$ . Then, when  $\lambda$  is sufficiently large, and  $s \geq s_0 - 1 > 0$ , with  $s_0$  being sufficiently large, by (ii),

$$\begin{aligned} \mathcal{L}_{0,s}^{\pm} \tilde{\chi}^{\pm} &= -\frac{1 + \mathcal{O}(s^{-1/2-\epsilon} + \lambda^{-1})}{\rho \lambda \delta s^2} (x - x_s) \cdot \nabla \phi \left( \frac{x - x_s}{\rho \lambda \delta s} \right) \phi \left( \frac{\xi \mp \xi_s}{\rho(\delta - s^{-\nu})} \right) \\ &\quad - \frac{\nu - \mathcal{O}(\lambda) s^{\nu-\epsilon}}{\rho(\delta - s^{-\nu})^2 s^{\nu+1}} (\xi \mp \xi_s) \cdot \phi \left( \frac{x - x_s}{\rho \lambda \delta s} \right) \nabla \phi \left( \frac{\xi \mp \xi_s}{\rho(\delta - s^{-\nu})} \right) \geq 0. \end{aligned} \quad (6-11)$$

We verify as in Lemma 3.2 that

$$\tilde{\chi}^{\pm} \in W^{\infty, \infty}(\mathbb{R}_{\geq s_0}, S_0^{-\infty}), \quad \mathcal{L}_{0,s}^{\pm} \tilde{\chi}^{\pm} \in W^{\infty, \infty}(\mathbb{R}_{\geq s_0}, \Gamma_{-1,0}^{-\infty, \mu-K-\bar{d}}).$$

We then choose  $\rho > 0$  sufficiently small such that  $\rho \lambda \delta$  is small and that  $\text{supp } \tilde{\chi}^{\pm}(s, \frac{s}{t_0} x, \xi) \subset \Omega$  when  $s$  is large. Next, we set, for  $s \geq s_0$ ,

$$\chi^{\pm}(s, x, \xi) = \tilde{\chi}^{\pm}(s, x, \xi).$$

To define  $\chi^{\pm}$  for  $s \leq s_0$ , we choose  $\rho \in C^{\infty}(\mathbb{R})$  such that  $0 \leq \rho \leq 1$ ,  $\rho(s) = 1$  for  $s \geq s_0$ , and  $\rho(s) = 0$  for  $s \leq s_0 - \alpha$  for some small  $\alpha > 0$  to be specified later, and solve the transport equation on  $[0, s_0]$ ,

$$\mathcal{L}_{0,s}^{\pm} \chi^{\pm}(s, x, \xi) = \rho(s) \mathcal{L}_{0,s}^{\pm} \tilde{\chi}^{\pm}(s, x, \xi), \quad \chi^{\pm}(s_0, x, \xi) = \tilde{\chi}^{\pm}(s_0, x, \xi).$$

Because the vector field involved in the definition of  $\mathcal{L}_{0,s}^{\pm}$  is in  $W^{\mu-K-\bar{d}, \infty}$  with respect to the  $x$ -variable, we deduce that  $\chi^{\pm} \in W^{1, \infty}(\mathbb{R}_{\geq 0}, \Upsilon^{\mu-K-\bar{d}})$  and thus  $\chi^{\pm}$  satisfies (6-9). Clearly

$$\mathcal{L}_{0,s}^{\pm} \chi^{\pm} \geq 0. \quad (6-12)$$



Moreover, because

$$\chi^\pm(s, x, \xi) = \tilde{\chi}^\pm(s_0, \Phi_{\pm(s_0-s)}(x, \xi)) - \int_s^{s_0} \rho(\sigma) \mathcal{L}_{0,s}^\pm \tilde{\chi}^\pm(\sigma, \Phi_{\pm(\sigma-s)}(x, \xi)) \, d\sigma,$$

if we choose  $\alpha > 0$  sufficiently small, then

$$\begin{aligned} \chi^\pm(0, x_0, \pm\xi_0) &= \tilde{\chi}^\pm(s_0, x_{s_0}, \pm\xi_{s_0}) - \int_{s_0-\alpha}^{s_0} \rho(\sigma) \mathcal{L}_{0,s}^\pm \tilde{\chi}^\pm(\sigma, x_\sigma, \pm\xi_\sigma) \, d\sigma \\ &\geq 1 - \|\mathcal{L}_{0,s}^\pm \tilde{\chi}^\pm(\sigma, x_\sigma, \pm\xi_\sigma)\|_{L^1_\sigma([s_0-\alpha, s_0])} > 0. \end{aligned}$$

Therefore,  $\chi^\pm(0, \cdot)$  is elliptic at  $(x_0, \pm\xi_0)$ .

To estimate  $\mathcal{L}_{h,s}^\pm \chi^\pm$ , we use

$$\begin{aligned} H_h(s, x, \xi) - H_0(s, x, \xi) &= H_h(s, x, \xi) - H_h(0, x, \xi) \\ &= \int_0^s (\partial_s H_h)(\sigma, x, \xi) \, d\sigma = h^{1/2} \int_0^s (\partial_t \gamma^{(3/2)})(h^{1/2}\sigma, x, \xi) \, d\sigma, \end{aligned}$$

and write

$$\begin{aligned} \mathcal{L}_{h,s}^\pm \chi^\pm(s, \cdot) - \mathcal{L}_{0,s}^\pm \chi^\pm(s, \cdot) &= \pm \{H_h - H_0, \chi^\pm\}(s, \cdot) \\ &= \pm h^{1/2} \int_0^s \{\partial_t \gamma^{(3/2)}(h^{1/2}\sigma, \cdot), \chi^\pm(s, \cdot)\} \, d\sigma. \end{aligned}$$

Observe that

$$\partial_t \gamma^{(3/2)} = -\frac{3}{2} \left( |\xi|^2 - \frac{(\nabla\eta \cdot \xi)^2}{1 + |\nabla\eta|^2} \right)^{-1/4} \left( \frac{\nabla\eta \cdot \xi}{1 + |\nabla\eta|^2} \nabla G(\eta) \psi \cdot \xi - \frac{(\nabla\eta \cdot \xi)^2}{(1 + |\nabla\eta|^2)^2} \nabla G(\eta) \psi \cdot \nabla\eta \right).$$

By hypothesis and Proposition 5.4,  $\nabla G(\eta) \psi \in \mathcal{H}_k^{\mu-2, 1/2} \subset H_2^{\mu-3}$  as  $k \geq 2$ . Therefore,

$$\partial_t \gamma^{(3/2)}(h^{1/2} \cdot, \cdot) \in L^\infty([0, h^{-1/2}T], \Gamma_{-2,0}^{3/2, \mu-K-\tilde{d}}).$$

Using  $|x| \sim s$  on  $\text{supp } \chi^\pm(s, \cdot)$ , we have, uniformly for all  $s \in [0, h^{-1/2}T]$ ,

$$\langle s \rangle \{\partial_t \gamma^{(3/2)}(h^{1/2}\sigma, \cdot), \chi^\pm(s, \cdot)\} \in L^\infty([0, h^{-1/2}T], \langle x \rangle^{-1} \Upsilon^{\mu-K-\tilde{d}-1}).$$

Therefore,

$$\begin{aligned} \mathcal{L}_{h,s}^\pm \chi^\pm(s, \cdot) - \mathcal{L}_{0,s}^\pm \chi^\pm(s, \cdot) &= \pm h^{1/2} \langle s \rangle^{-1} \int_0^s \mathcal{O}(1)_{L^\infty([0, h^{-1/2}T], \langle x \rangle^{-1} \Upsilon^{\mu-K-\tilde{d}-1})} \, d\sigma \\ &= \pm h^{1/2} \langle s \rangle^{-1} \mathcal{O}(s)_{\langle x \rangle^{-1} \Upsilon^{\mu-K-\tilde{d}-1}} \\ &= \mathcal{O}(h^{1/2})_{\langle x \rangle^{-1} \Upsilon^{\mu-K-\tilde{d}-1}}, \end{aligned}$$

which, together with (6-12), proves (6-10). □

**6C3. Propagation.** Now we prove Theorem 1.9. By Lemmas 6.5 and 6.7, it suffices to prove the following propagation theorem for  $u$  defined as in (6-2).

**Theorem 6.10.** *Under the hypothesis of Theorem 1.9, let  $u$  be defined as (6-2). Let*

$$(x_0, \xi_0) \in \text{WF}_{0,1}^\sigma(u_0)^\circ,$$

with  $0 \leq \sigma < \min\{(\mu - K - \tilde{d})/2, 3k/2\}$  for some  $K > 0$ , such that the cogeodesic  $\{(x_s, \xi_s) = \Phi_s(x_0, \xi_0)\}_{s \in \mathbb{R}}$  is forwardly nontrapping. Set

$$\xi_\infty = \lim_{s \rightarrow +\infty} \xi_s.$$

Then, for all  $t_0 \in (0, T]$ , we have

$$\left(\frac{3}{2}t_0|\xi_\infty|^{-1/2}\xi_\infty, \xi_\infty\right) \in \text{WF}_{1/2,1}^\sigma(u(t_0)).$$

Under the semiclassical time variable  $s = h^{-1/2}t$ , (6-1) becomes

$$(\partial_s + h^{1/2}\mathcal{P}_V \cdot \nabla + h^{1/2}\mathcal{P}_Q)w + ih^{1/2} \begin{pmatrix} \mathcal{P}_\gamma & 0 \\ 0 & -\mathcal{P}_\gamma \end{pmatrix} w + \frac{ih^{1/2}g}{2}\mathcal{P}_\zeta \begin{pmatrix} 1 & -1 \\ 1 & -1 \end{pmatrix} w = F_h = \mathcal{O}(h^{1/2})_{H^{\mu-K-\tilde{d}}}$$

for some  $K > 0$ . We define  $\mathcal{L}_s^h$ , which applies to time-dependent operators  $\mathcal{A} : \mathcal{S} \rightarrow \mathcal{S}'$ ,

$$\mathcal{L}_s^h \mathcal{A} = \partial_s \mathcal{A} + h^{1/2} \left[ \mathcal{P}_V \cdot \nabla + \mathcal{P}_Q + i\mathcal{P}_\gamma \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} + \frac{ig}{2}\mathcal{P}_\zeta \begin{pmatrix} 1 & -1 \\ 1 & -1 \end{pmatrix}, \mathcal{A} \right].$$

We also define  $\mathcal{L}_s^h$ , which applies to symbols of the diagonal form  $A = \begin{pmatrix} A^+ & 0 \\ 0 & A^- \end{pmatrix}$ :

$$\mathcal{L}_s^h A = \begin{pmatrix} \mathcal{L}_{h,s}^+ A^+ & 0 \\ 0 & \mathcal{L}_{h,s}^- A^- \end{pmatrix}.$$

*Proof of Theorem 6.10.* We shall from now on write  $\rho = \mu - K - \tilde{d}$  for some sufficiently large  $K > 0$ , also define  $I_h = [0, h^{-1/2}T]$  and

$$Y_h^\rho = L^\infty \left( I_h, M_{2 \times 2} \left( \sum_{j=0}^\rho h^j \Upsilon^{\rho-j} \right) \right)$$

for simplicity. More precisely, a symbol  $A_h = \sum_{j=0}^\rho h^j A_h^j \in Y_h^\rho$  if

$$\sup_{h \in (0,1]} \sup_{s \in [0, h^{-1/2}T]} N^{\rho-j}(A_h^j) < +\infty,$$

where the norm  $N^{\rho-j}(A_h^j)$  is applied to every component of  $A_h^j$ . Choose a strictly increasing sequence  $\{\lambda_j\}_{j \geq 0} \subset [1, 1 + \epsilon)$  with  $\epsilon > 0$  being sufficiently small. Define  $\chi_j^\pm$  as in Lemma 6.9, where we replace  $\phi$  with  $\phi(\cdot / \lambda_j)$ . Then

$$\text{supp } \chi_j^\pm \subset \{\chi_{j+1}^\pm > 0\}$$

for all  $j \in \mathbb{N}$ . Set

$$\chi_j = \begin{pmatrix} \chi_j^+ & 0 \\ 0 & \chi_j^- \end{pmatrix}.$$

We shall construct an operator  $\mathcal{A}_h \in L^\infty(I_h, L^2 \rightarrow L^2)$  such that:

- (1)  $\mathcal{A}_h$  is a paradifferential operator; more precisely, there exists

$$A_h^\pm \in W^{1,\infty}(\mathbb{R}_{\geq 0}, \Upsilon^{\rho+1}) \cap W^{1,\infty}(\mathbb{R}_{\geq s_0}, S_0^{-\infty})$$

for some  $s_0 > 0$ , such that

$$\mathcal{A}_h - \mathcal{P}_{A_h}^h = \mathcal{O}(h^\rho)_{L^\infty(I_h, L^2 \rightarrow L^2)}, \quad A_h = \begin{pmatrix} A_h^+ & 0 \\ 0 & A_h^- \end{pmatrix}.$$

Moreover, we require that

$$\text{supp } A_h^\pm \subset \bigcup_{j \geq 0} \text{supp } \chi_j^\pm.$$

- (2)  $A_h^\pm(0, x, \xi)$  is elliptic at  $(x_0, \pm \xi_0)$ .
- (3)  $A_h^\pm(s, \frac{s}{t_0}x, \xi) \in S_{-\infty}^{-\infty}$  is elliptic at  $(\frac{3}{2}t_0|\xi_\infty|^{-1/2}\xi_\infty, \xi_\infty)$  for  $s > 0$  sufficiently large.
- (4)  $\mathcal{L}_s^h \mathcal{A}_h \geq \mathcal{O}(h^\rho)_{L^\infty(I_h, L^2 \rightarrow L^2)}$ .

We shall construct  $\mathcal{A}_h$  of the form

$$\mathcal{A}_h = \sum_{j \geq 0} h^{j/2} \varphi^j \mathcal{A}_h^j,$$

where  $\varphi \in P_j$ , recalling the definition (3-6), and  $\mathcal{A}_h^j \in L^\infty(I_h, L^2 \rightarrow L^2)$ . We begin by setting

$$\mathcal{A}_h^0 = (\mathcal{P}_{\chi_0}^h)^* \mathcal{P}_{\chi_0}^h, \quad \varphi^0 \equiv 1.$$

Therefore, by the symbolic calculus, Lemma 6.9 and Corollary 4.32 (observe that the symbol of  $\mathcal{A}_h^0$  belongs to  $\sigma_0$ , and that  $\gamma$  is a sum of homogeneous symbols),

$$\partial_s \mathcal{A}_h^0 + h^{1/2} \left[ i\mathcal{P}_\gamma \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}, \mathcal{A}_h^0 \right] = 2\mathcal{P}_{\chi_0 \mathcal{L}_s^h \chi_0}^h + h\mathcal{P}_{b_h^0}^h + \mathcal{O}(h^\rho)_{L^\infty(I_h, L^2 \rightarrow L^2)}$$

for some symbol  $b_h^0$  such that  $\langle x \rangle b_h^0 \in Y_h^\rho$ . This  $\langle x \rangle$  factor comes from the spatial decay of  $\partial_{x,\xi} \gamma$ . Moreover, we have  $\text{supp } b_h^0 \subset \text{supp } \chi_0$ , which implies  $\langle s \rangle b_h^0 \in Y_h^\rho$ . Similarly,

$$h^{1/2} [\mathcal{P}_V \cdot \nabla, \mathcal{A}_h^0] = h^{1/2} \mathcal{P}_{b_h^1}^h + \mathcal{O}(h^\rho)_{L^\infty(I_h, L^2 \rightarrow L^2)},$$

where  $\langle s \rangle b_h^1 \in Y_h^\rho$ , with  $\text{supp } b_h^1 \subset \text{supp } \chi_0$ . Be careful that, because  $Q$  and  $\mathcal{P}_\zeta \begin{pmatrix} 1 & -1 \\ 1 & -1 \end{pmatrix}$  are not diagonal matrices, their commutators with  $\mathcal{A}_h^0$  do not gain an extra  $h$ , for the principal symbols do not cancel each other. So,

$$\begin{aligned} h^{1/2} [\mathcal{P}_Q, \mathcal{A}_h^0] &= h^{1/2} \mathcal{P}_{b_h^2}^h + \mathcal{O}(h^\rho)_{L^\infty(I_h, L^2 \rightarrow L^2)}, \\ h^{1/2} \left[ \mathcal{P}_\zeta \begin{pmatrix} 1 & -1 \\ 1 & -1 \end{pmatrix}, \mathcal{A}_h^0 \right] &= h\mathcal{P}_{b_h^3}^h + \mathcal{O}(h^\rho)_{L^\infty(I_h, L^2 \rightarrow L^2)}, \end{aligned}$$

where  $\langle s \rangle b_h^2, \langle s \rangle b_h^3 \in Y_h^\rho$ , with  $\text{supp } b_h^2 \cup \text{supp } b_h^3 \subset \text{supp } \chi_0$ . By Lemma 6.9,

$$\chi_0 \mathcal{L}_s^h \chi_0 \geq h^{1/2} b_h^4,$$

where  $\langle s \rangle b_h^4 \in Y_h^\rho$ , with  $\text{supp } b_h^4 \subset \text{supp } \chi_0$ . Therefore, combining the idea described above (3-10) and the paradifferential Gårding inequality (Lemma 4.40, where we take  $\epsilon = \frac{1}{2}$ ),

$$\mathcal{P}_{\chi_0 \mathcal{L}_s^h \chi_0}^h - h^{1/2} \mathcal{P}_{b_h^4}^h \geq h^{1/2} \mathcal{P}_{b_h^5}^h + \mathcal{O}(h^\rho)_{L^2 \rightarrow L^2}$$

for some  $b_h^5 \in Y_h^\rho$  with  $\text{supp } b_h^5 \subset \{\chi_1 > 0\}$ . In fact, choose  $c_h \in L^\infty(\mathbb{R}_{\geq 0}, S_0^{-\infty})$  such that

$$\text{supp } a_h \subset \{c_h = 1\} \subset \text{supp } c_h \subset \text{supp } \chi_1.$$

Then, for all  $v \in L^2$ , we have

$$\begin{aligned} \langle v, (\mathcal{P}_{\chi_0 \mathcal{L}_s^h \chi_0}^h - h^{1/2} \mathcal{P}_{b_h^4}^4) v \rangle_{L^2} &= \langle \mathcal{P}_{c_h} v, (\mathcal{P}_{\chi_0 \mathcal{L}_s^h \chi_0}^h - h^{1/2} \mathcal{P}_{b_h^4}^4) \mathcal{P}_{c_h} v \rangle_{L^2} + \mathcal{O}(h^\rho) \\ &\gtrsim -Ch^{1/2} \|\mathcal{P}_{c_h} v\|_{L^2}^2 + \mathcal{O}(h^\rho). \end{aligned}$$

Therefore, it suffices to choose  $b_h^5$  such that

$$\mathcal{P}_{b_h^5} - C\mathcal{P}_{c_h}^* \mathcal{P}_{c_h} = \mathcal{O}(h^\rho)_{L^\infty(I_h, L^2 \rightarrow L^2)},$$

which can be achieved by Propositions 4.26 and 4.27. Set

$$\alpha_h^0 = \langle s \rangle (b_h^1 + b_h^2 + 2b_h^4 + 2b_h^5) \in Y_h^\rho, \quad \beta_h^0 = \langle s \rangle (b_h^0 + b_h^3) \in Y_h^\rho.$$

Then

$$\mathcal{L}_s^h \mathcal{A}_h^0 \geq h^{1/2} \langle s \rangle^{-1} \mathcal{P}_{\alpha_h^0 + h^{1/2} \beta_h^0}^h + \mathcal{O}(h^\rho)_{L^\infty(I_h, L^2 \rightarrow L^2)}.$$

Suppose that we have found  $\mathcal{A}_h^j \in L^\infty(I_h, L^2 \rightarrow L^2)$ ,  $\varphi^j \in P_j$  for  $j = 0, \dots, \ell - 1$ , and  $\psi^{\ell-1} \in P_{\ell-1}$ ,  $\alpha_h^{\ell-1}, \beta_h^{\ell-1} \in Y_h^\rho$ , with

$$\text{supp } \alpha_h^{\ell-1} \cup \text{supp } \beta_h^{\ell-1} \subset \{\chi_\ell > 0\},$$

such that

$$\mathcal{L}_s^h \left( \sum_{j=0}^{\ell-1} h^{j/2} \varphi^j \mathcal{A}_h^j \right) \geq h^{\ell/2} \langle s \rangle^{-1} \psi^{\ell-1} \mathcal{P}_{\alpha_h^{\ell-1} + h^{1/2} \beta_h^{\ell-1}}^h + \mathcal{O}(h^\rho)_{L^\infty(I_h, L^2 \rightarrow L^2)}. \quad (6-13)$$

Then as in the proof of Theorem 1.4(2), we set

$$\varphi^\ell(s) = \int_0^s (1 + \sigma)^{-1} \psi^{\ell-1}(\sigma) \, d\sigma, \quad \mathcal{A}_h^\ell = C_\ell \varphi^\ell \mathcal{P}_{\chi_\ell}^h,$$

where the constant  $C_\ell$  is sufficiently large, such that by Lemma 6.9, in the sense of positivity of matrices,

$$\begin{aligned} C_\ell \mathcal{L}_s^s(\varphi^\ell \chi_\ell) &= C_\ell (1 + s)^{-1} \psi^{\ell-1} \chi_\ell + C_\ell \varphi^\ell \mathcal{L}_s^s \chi_\ell \\ &\geq \langle s \rangle^{-1} \psi^{\ell-1} \alpha_h^{\ell-1} + \varphi^\ell h^{1/2} \langle s \rangle^{-1} \tilde{\beta}_h^\ell \end{aligned}$$

for some  $\tilde{\beta}_h^\ell \in Y_h^\rho$ . By the paradifferential Gårding inequality, and a routine construction of a parametrix, we find  $\tilde{\alpha}_h^\ell \in Y_h^\rho$ , with  $\text{supp } \tilde{\alpha}_h^\ell \subset \{\chi_{\ell+1} > 0\}$ , such that

$$\mathcal{P}_{C_\ell \mathcal{L}_s^s(\varphi^\ell \chi_\ell)}^h - \langle s \rangle^{-1} \mathcal{P}_{\psi^{\ell-1} \alpha_h^{\ell-1} + h^{1/2} \varphi^\ell \tilde{\beta}_h^\ell}^h \geq h \langle s \rangle^{-1} \mathcal{P}_{(\psi^{\ell-1} + \varphi^\ell) \tilde{\alpha}_h^\ell}^h + \mathcal{O}(h^\rho)_{L^\infty(I_h, L^2 \rightarrow L^2)}.$$

Similar to the estimate of  $\mathcal{A}_h^0$ , by a symbolic calculus, we find  $\underline{\alpha}_h^\ell, \underline{\beta}_h^\ell \in Y_h^\rho$ , with

$$\text{supp } \underline{\alpha}_h^\ell \cup \text{supp } \underline{\beta}_h^\ell \subset \text{supp } \chi_\ell$$

such that

$$\mathcal{L}_s^h \mathcal{A}_h^\ell = \mathcal{P}_{C_\ell \mathcal{L}_s^s(\varphi^\ell \chi_\ell)}^h + h^{1/2} \langle s \rangle^{-1} \varphi^\ell \mathcal{P}_{\underline{\alpha}_h^\ell + h^{1/2} \underline{\beta}_h^\ell} + \mathcal{O}(h^\rho)_{L^\infty(I_h, L^2 \rightarrow L^2)}.$$

Summing up the two inequalities above,

$$\mathcal{L}_s^h \mathcal{A}_h^\ell - \langle s \rangle^{-1} \psi^{\ell-1} \mathcal{P}_{\alpha_h^{\ell-1}}^h \geq h^{1/2} \langle s \rangle^{-1} \mathcal{P}_{\varphi^\ell (\underline{\alpha}_h^\ell + \tilde{\beta}_h^\ell) + h^{1/2} (\psi^{\ell-1} + \varphi^\ell) \tilde{\alpha}_h^\ell + h^{1/2} \varphi^\ell \underline{\beta}_h^\ell} + \mathcal{O}(h^\rho)_{L^\infty(I_h, L^2 \rightarrow L^2)}. \quad (6-14)$$

Therefore, combining (6-13) and (6-14),

$$\mathcal{L}_s^h \left( \sum_{j=0}^{\ell} h^{j/2} \varphi^j \mathcal{A}_h^j \right) \geq h^{(\ell+1)/2} \langle s \rangle^{-1} \psi^\ell \mathcal{P}_{\alpha_h^\ell + h^{1/2} \beta_h^\ell}^h + \mathcal{O}(h^\rho)_{L^\infty(I_h, L^2 \rightarrow L^2)},$$

where

$$\psi^\ell = 1 + \psi^{\ell-1} + \varphi^\ell, \quad \alpha_h^\ell = \frac{\psi^{\ell-1}}{\psi^\ell} \beta_h^{\ell-1} + \frac{\varphi^\ell}{\psi^\ell} (\underline{\alpha}_h^\ell + \tilde{\beta}_h^\ell), \quad \beta_h^\ell = \frac{\psi^{\ell-1} + \varphi^\ell}{\psi^\ell} \tilde{\alpha}_h^\ell + \frac{\varphi^\ell}{\psi^\ell} \underline{\beta}_h^\ell.$$

Thus we close the induction procedure.

To finish the proof, suppose that

$$\begin{aligned} (\tfrac{3}{2}t_0 |\xi_\infty|^{-1/2} \xi_\infty, \xi_\infty) &\notin \text{WF}_{1/2,1}^\sigma(u(t_0)), \\ (\tfrac{3}{2}t_0 |\xi_\infty|^{-1/2} \xi_\infty, -\xi_\infty) &\notin \text{WF}_{1/2,1}^\sigma(\overline{u(t_0)}). \end{aligned}$$

By Lemma 6.9, we can choose  $\phi$  such that, for sufficiently small  $h > 0$ ,

$$\begin{aligned} \text{supp } \theta_{1/h,*}^{1/2,0} \chi_j^+ |_{s=h^{-1/2}t_0} &\subset \mathbb{R}^{2d} \setminus \text{WF}_{1/2,1}^\sigma(u(t_0)), \\ \text{supp } \theta_{1/h,*}^{1/2,0} \chi_j^- |_{s=h^{-1/2}t_0} &\subset \mathbb{R}^{2d} \setminus \text{WF}_{1/2,1}^\sigma(\overline{u(t_0)}). \end{aligned}$$

So by Lemmas 4.34 and 2.14,

$$(\mathcal{A}_h w, w)_{L^2} |_{s=h^{-1/2}t_0} = \mathcal{O}(h^{2\sigma}).$$

By our construction,  $\varphi^\ell(0) = 0$  for all  $\ell \geq 1$ , so

$$\mathcal{A}_h |_{s=0} = \mathcal{A}_h^0 |_{s=0} = (\mathcal{P}_{\chi_0}^h)^* \mathcal{P}_{\chi_0}^h |_{s=0}.$$

Because  $F_h = \mathcal{O}(h^{1/2})_{H^\rho}$ , we have, by Lemma 2.15, that  $\mathcal{A}_h F_h = \mathcal{O}(h^{\rho+1/2})_{L^2}$ . Therefore, by (4),

$$\begin{aligned} \|\mathcal{P}_{\chi_0}^h w |_{s=0}\|_{L^2}^2 &= \text{Re}(\mathcal{A}_h w, w)_{L^2} |_{s=h^{-1/2}t_0} - \int_0^{h^{-1/2}t_0} \text{Re}(\mathcal{L}_s^h \mathcal{A}_h w, w)_{L^2} ds - \int_0^{h^{-1/2}t_0} \text{Re}(\mathcal{A}_h F_h, w)_{L^2} ds \\ &\leq \mathcal{O}(h^{2\sigma}) + \mathcal{O}(h^{\rho-1/2}) = \mathcal{O}(h^{2\sigma}). \end{aligned}$$

Observe that  $\chi_0 |_{s=0}$  is of compact support with respect to  $x$ , and we have

$$\mathcal{P}_{\chi_0 |_{s=0}}^h = T_{\beta_h}^h + \mathcal{O}(h^\rho)_{L^2 \rightarrow L^2},$$

where

$$\beta_h = \sum_{j \geq 0} \psi_j \chi_0 |_{s=0} \#_h \psi_j \in \sum_{j=0}^{\rho} h^j \Upsilon^{\rho-j}$$

is a finite summation. By Lemma 4.41 and (6-5), we conclude that, if  $(x_0, \xi_0) \notin \text{WF}_{0,1}^\sigma(u_0)$  provided  $\sigma \leq \frac{3}{2}r$ , where

$$r = \min\left\{\left[\frac{2}{3}(\mu - 1 - \tilde{d})\right], k\right\},$$

then under the hypothesis of theorem we have  $r = k$ . □

**6D. Proof of Corollary 1.10.** The case when  $d = 1$  is trivial. For the second case, we shall prove that, on any cogeodesic  $\{(x_t, \xi_t)\}_{t \in \mathbb{R}}$ ,

$$\lim_{t \rightarrow +\infty} x_t \cdot \xi_t = \infty, \quad (6-15)$$

so no geodesics can be trapped. The proof of (6-15) is almost finished by the proof of Lemma 6.8. Indeed, similar calculations imply

$$\frac{d}{dt}(x_t \cdot \xi_t) \gtrsim |\xi_0|^2.$$

### List of notation

$\text{WF}^\mu(u)$	wavefront set
$\text{WF}_{\delta, \rho}^\mu(u)$	quasihomogeneous wavefront set
$\text{Op}(a)$	pseudodifferential operator
$\text{Op}_h(a)$	semiclassical pseudodifferential operator
$\text{Op}_h^{\delta, \rho}(a)$	quasihomogeneous semiclassical pseudodifferential operator
$T_a$	paradifferential operator
$\mathcal{P}_a$	dyadic paradifferential operator
$\mathcal{P}_a^h$	semiclassical dyadic paradifferential operator
$\mathcal{P}_a^{h, \epsilon}$	quasihomogeneous semiclassical dyadic paradifferential operator
$a \sharp_h^{\delta, \rho} b$	composition of symbols
$\zeta_h^{\delta, \rho} a$	adjoint of symbols
$\mathcal{S}, \mathcal{S}'$	Schwartz function space and tempered distribution space
$\mathcal{H}_k^{\mu, \delta}, W_{k, \delta}^{r, \infty}$	weighted Sobolev spaces
$\Gamma^{m, r}$	paradifferential symbol class
$\Gamma_{k, \delta}^{m, r}$	weighted paradifferential symbol class
$\Sigma_{k, \delta}^{m, r}$	weighted paradifferential polysymbol class
$M^{m, r}, M_{k, \delta}^{m, r}$	symbol norm and weighted symbol norm
$\theta_h^{\delta, \rho}$	phase-space scaling operator

### Acknowledgments

The author would like to thank Thomas Alazard, Nicolas Burq and Claude Zuily for their constant support and encouragement. He would like to thank Shu Nakamura for helpful discussions during the early state of this project. He would also like to thank Jean-Marc Delort for his careful reading of the manuscript, and thank Daniel Tataru for his useful comments. Finally the author would like to thank the anonymous referees for their detailed comments which result in a better presentation of the paper.

### References

- [Alazard and Delort 2015] T. Alazard and J.-M. Delort, “Global solutions and asymptotic behavior for two dimensional gravity water waves”, *Ann. Sci. École Norm. Sup. (4)* **48**:5 (2015), 1149–1238. MR Zbl
- [Alazard and Métivier 2009] T. Alazard and G. Métivier, “Paralinearization of the Dirichlet to Neumann operator, and regularity of three-dimensional water waves”, *Comm. Partial Differential Equations* **34**:10-12 (2009), 1632–1704. MR Zbl

- [Alazard, Burq and Zuily 2011] T. Alazard, N. Burq, and C. Zuily, “On the water-wave equations with surface tension”, *Duke Math. J.* **158**:3 (2011), 413–499. [MR](#) [Zbl](#)
- [Alazard, Burq and Zuily 2014] T. Alazard, N. Burq, and C. Zuily, “On the Cauchy problem for gravity water waves”, *Invent. Math.* **198**:1 (2014), 71–163. [MR](#) [Zbl](#)
- [Alazard, Burq and Zuily 2016] T. Alazard, N. Burq, and C. Zuily, “Cauchy theory for the gravity water waves system with non-localized initial data”, *Ann. Inst. H. Poincaré C Anal. Non Linéaire* **33**:2 (2016), 337–395. [MR](#) [Zbl](#)
- [Alazard, Ifrim and Tataru 2022] T. Alazard, M. Ifrim, and D. Tataru, “A Morawetz inequality for water waves”, *Amer. J. Math.* **144**:3 (2022), 607–699. [MR](#) [Zbl](#)
- [Bahouri, Chemin and Danchin 2011] H. Bahouri, J.-Y. Chemin, and R. Danchin, *Fourier analysis and nonlinear partial differential equations*, Grundlehren der Math. Wissenschaften **343**, Springer, 2011. [MR](#) [Zbl](#)
- [Beyer and Günther 1998] K. Beyer and M. Günther, “On the Cauchy problem for a capillary drop, I: Irrotational motion”, *Math. Methods Appl. Sci.* **21**:12 (1998), 1149–1183. [MR](#) [Zbl](#)
- [Bony 1979] J.-M. Bony, “Calcul symbolique et singularités des solutions des équations aux dérivées partielles non linéaires”, art. id. 1 in *Journées: équations aux dérivées partielles* (Saint-Cast, France, 1979), École Polytech., Palaiseau, France, 1979. [MR](#) [Zbl](#)
- [Bony 1986] J.-M. Bony, “Singularités des solutions de problèmes de Cauchy hyperboliques non linéaires”, pp. 15–39 in *Advances in microlocal analysis* (Lucca, Italy, 1985), edited by H. G. Garnir, NATO Adv. Sci. Inst. Ser. C: Math. Phys. Sci. **168**, Reidel, Dordrecht, Netherlands, 1986. [MR](#) [Zbl](#)
- [Boutet de Monvel 1975] L. Boutet de Monvel, “Propagation des singularités des solutions d’équations analogues à l’équation de Schrödinger”, pp. 1–14 in *Fourier integral operators and partial differential equations* (Nice, 1974), edited by J. Chazarain, Lecture Notes in Math. **459**, Springer, 1975. [MR](#) [Zbl](#)
- [Burq 2004] N. Burq, “Smoothing effect for Schrödinger boundary value problems”, *Duke Math. J.* **123**:2 (2004), 403–427. [MR](#) [Zbl](#)
- [Christianson, Hur and Staffilani 2009] H. Christianson, V. M. Hur, and G. Staffilani, “Local smoothing effects for the water-wave problem with surface tension”, *C. R. Math. Acad. Sci. Paris* **347**:3-4 (2009), 159–162. [MR](#) [Zbl](#)
- [Courant and Lax 1956] R. Courant and P. D. Lax, “The propagation of discontinuities in wave motion”, *Proc. Nat. Acad. Sci. USA* **42** (1956), 872–876. [MR](#) [Zbl](#)
- [Craig and Sulem 1993] W. Craig and C. Sulem, “Numerical simulation of gravity waves”, *J. Comput. Phys.* **108**:1 (1993), 73–83. [MR](#) [Zbl](#)
- [Craig, Kappeler and Strauss 1995] W. Craig, T. Kappeler, and W. Strauss, “Microlocal dispersive smoothing for the Schrödinger equation”, *Comm. Pure Appl. Math.* **48**:8 (1995), 769–860. [MR](#) [Zbl](#)
- [Deng, Ionescu, Pausader and Pusateri 2017] Y. Deng, A. D. Ionescu, B. Pausader, and F. Pusateri, “Global solutions of the gravity-capillary water-wave system in three dimensions”, *Acta Math.* **219**:2 (2017), 213–402. [MR](#) [Zbl](#)
- [Doi 1996] S.-i. Doi, “Smoothing effects of Schrödinger evolution groups on Riemannian manifolds”, *Duke Math. J.* **82**:3 (1996), 679–706. [MR](#) [Zbl](#)
- [Doi 2000] S.-i. Doi, “Smoothing effects for Schrödinger evolution equation and global behavior of geodesic flow”, *Math. Ann.* **318**:2 (2000), 355–389. [MR](#) [Zbl](#)
- [Guillemin and Sternberg 1977] V. Guillemin and S. Sternberg, *Geometric asymptotics*, Math. Surv. **14**, Amer. Math. Soc., Providence, RI, 1977. [MR](#) [Zbl](#)
- [Hassell and Wunsch 2005] A. Hassell and J. Wunsch, “The Schrödinger propagator for scattering metrics”, *Ann. of Math.* (2) **162**:1 (2005), 487–523. [MR](#) [Zbl](#)
- [Hérau 2002] F. Hérau, “Melin inequality for paradifferential operators and applications”, *Comm. Partial Differential Equations* **27**:7-8 (2002), 1659–1680. [MR](#) [Zbl](#)
- [Hörmander 1971] L. Hörmander, “Fourier integral operators, I”, *Acta Math.* **127**:1-2 (1971), 79–183. [MR](#) [Zbl](#)
- [Hörmander 1991] L. Hörmander, “Quadratic hyperbolic operators”, pp. 118–160 in *Microlocal analysis and applications* (Montecatini Terme, Italy, 1989), edited by L. Cattabriga and L. Rodino, Lecture Notes in Math. **1495**, Springer, 1991. [MR](#) [Zbl](#)

- [Hörmander 1997] L. Hörmander, *Lectures on nonlinear hyperbolic differential equations*, Math. Appl. (Berlin) **26**, Springer, 1997. [MR](#) [Zbl](#)
- [Hunter, Ifrim and Tataru 2016] J. K. Hunter, M. Ifrim, and D. Tataru, “Two dimensional water waves in holomorphic coordinates”, *Comm. Math. Phys.* **346**:2 (2016), 483–552. [MR](#) [Zbl](#)
- [Ifrim and Tataru 2017] M. Ifrim and D. Tataru, “The lifespan of small data solutions in two dimensional capillary water waves”, *Arch. Ration. Mech. Anal.* **225**:3 (2017), 1279–1346. [MR](#) [Zbl](#)
- [Ionescu and Pusateri 2018] A. D. Ionescu and F. Pusateri, “Recent advances on the global regularity for irrotational water waves”, *Philos. Trans. Roy. Soc. A* **376**:2111 (2018), art. id. 20170089. [MR](#) [Zbl](#)
- [Ito 2006] K. Ito, “Propagation of singularities for Schrödinger equations on the Euclidean space with a scattering metric”, *Comm. Partial Differential Equations* **31**:10-12 (2006), 1735–1777. [MR](#) [Zbl](#)
- [Kano and Nishida 1979] T. Kano and T. Nishida, “Sur les ondes de surface de l’eau avec une justification mathématique des équations des ondes en eau peu profonde”, *J. Math. Kyoto Univ.* **19**:2 (1979), 335–370. [MR](#) [Zbl](#)
- [Kato 1983] T. Kato, “On the Cauchy problem for the (generalized) Korteweg–de Vries equation”, pp. 93–128 in *Studies in applied mathematics*, edited by V. Guillemin, Adv. Math. Suppl. Stud. **8**, Academic Press, New York, 1983. [MR](#) [Zbl](#)
- [Kenig, Ponce and Vega 1998] C. E. Kenig, G. Ponce, and L. Vega, “Smoothing effects and local existence theory for the generalized nonlinear Schrödinger equations”, *Invent. Math.* **134**:3 (1998), 489–545. [MR](#) [Zbl](#)
- [Lascar 1977] R. Lascar, “Propagation des singularités des solutions d’équations pseudo-différentielles quasi homogènes”, *Ann. Inst. Fourier (Grenoble)* **27**:2 (1977), 79–123. [MR](#) [Zbl](#)
- [Lascar 1978] R. Lascar, “Propagation des singularités et hypoellipticité pour des opérateurs pseudo-différentiels à caractéristiques doubles”, *Comm. Partial Differential Equations* **3**:3 (1978), 201–247. [MR](#) [Zbl](#)
- [Lax 1957] P. D. Lax, “Asymptotic solutions of oscillatory initial value problems”, *Duke Math. J.* **24** (1957), 627–646. [MR](#) [Zbl](#)
- [Lebeau 1989] G. Lebeau, “Équations des ondes semi-linéaires, II: Contrôle des singularités et caustiques non linéaires”, *Invent. Math.* **95**:2 (1989), 277–323. [MR](#) [Zbl](#)
- [Lebeau 1992] G. Lebeau, “Contrôle de l’équation de Schrödinger”, *J. Math. Pures Appl.* (9) **71**:3 (1992), 267–291. [MR](#) [Zbl](#)
- [Melrose 1994] R. B. Melrose, “Spectral and scattering theory for the Laplacian on asymptotically Euclidian spaces”, pp. 85–130 in *Spectral and scattering theory* (Sanda, Japan, 1992), edited by M. Ikawa, Lecture Notes in Pure Appl. Math. **161**, Dekker, New York, 1994. [MR](#) [Zbl](#)
- [Melrose, Vasy and Wunsch 2013] R. Melrose, A. Vasy, and J. Wunsch, *Diffraction of singularities for the wave equation on manifolds with corners*, Astérisque **351**, Math. Soc. France, Paris, 2013. [MR](#) [Zbl](#)
- [Métivier 2008] G. Métivier, *Para-differential calculus and applications to the Cauchy problem for nonlinear systems*, Cent. Ricerca Mat. Ennio De Giorgi Ser. **5**, Edizioni della Normale, Pisa, 2008. [MR](#) [Zbl](#)
- [Métivier and Zumbrun 2005] G. Métivier and K. Zumbrun, *Large viscous boundary layers for noncharacteristic nonlinear hyperbolic problems*, Mem. Amer. Math. Soc. **826**, Amer. Math. Soc., Providence, RI, 2005. [MR](#) [Zbl](#)
- [Ming, Rousset and Tzvetkov 2015] M. Ming, F. Rousset, and N. Tzvetkov, “Multi-solitons and related solutions for the water-waves system”, *SIAM J. Math. Anal.* **47**:1 (2015), 897–954. [MR](#) [Zbl](#)
- [Nakamura 2005] S. Nakamura, “Propagation of the homogeneous wave front set for Schrödinger equations”, *Duke Math. J.* **126**:2 (2005), 349–367. [MR](#) [Zbl](#)
- [Nakamura 2009] S. Nakamura, “Semiclassical singularities propagation property for Schrödinger equations”, *J. Math. Soc. Japan* **61**:1 (2009), 177–211. [MR](#) [Zbl](#)
- [Nalimov 1974] V. I. Nalimov, “The Cauchy–Poisson problem”, *Dinamika Sploshn. Sredy* **18** (1974), 104–210. In Russian. [MR](#)
- [Nguyen 2016] Q.-H. Nguyen, “A pseudo-local property of gravity water waves system”, *SIAM J. Math. Anal.* **48**:3 (2016), 1988–2027. [MR](#) [Zbl](#)
- [de Poyferré and Nguyen 2016] T. de Poyferré and Q.-H. Nguyen, “Strichartz estimates and local existence for the gravity-capillary waves with non-Lipschitz initial velocity”, *J. Differential Equations* **261**:1 (2016), 396–438. [MR](#) [Zbl](#)
- [de Poyferré and Nguyen 2017] T. de Poyferré and Q.-H. Nguyen, “A paradifferential reduction for the gravity-capillary waves system at low regularity and applications”, *Bull. Soc. Math. France* **145**:4 (2017), 643–710. [MR](#) [Zbl](#)



- [Robbiano and Zuily 1999] L. Robbiano and C. Zuily, “Microlocal analytic smoothing effect for the Schrödinger equation”, *Duke Math. J.* **100**:1 (1999), 93–129. [MR](#) [Zbl](#)
- [Rodino and Wahlberg 2014] L. Rodino and P. Wahlberg, “The Gabor wave front set”, *Monatsh. Math.* **173**:4 (2014), 625–655. [MR](#) [Zbl](#)
- [Rousset and Tzvetkov 2011] F. Rousset and N. Tzvetkov, “Transverse instability of the line solitary water-waves”, *Invent. Math.* **184**:2 (2011), 257–388. [MR](#) [Zbl](#)
- [Schulz and Wahlberg 2017] R. Schulz and P. Wahlberg, “Equality of the homogeneous and the Gabor wave front set”, *Comm. Partial Differential Equations* **42**:5 (2017), 703–730. [MR](#) [Zbl](#)
- [Szeftel 2005] J. Szeftel, “Microlocal dispersive smoothing for the nonlinear Schrödinger equation”, *SIAM J. Math. Anal.* **37**:2 (2005), 549–597. [MR](#) [Zbl](#)
- [Tataru 2002] D. Tataru, “On the Fefferman–Phong inequality and related problems”, *Comm. Partial Differential Equations* **27**:11–12 (2002), 2101–2138. [MR](#) [Zbl](#)
- [Taylor 1991] M. E. Taylor, *Pseudodifferential operators and nonlinear PDE*, Progr. Math. **100**, Birkhäuser, Boston, 1991. [MR](#) [Zbl](#)
- [Vasy 2008] A. Vasy, “Propagation of singularities for the wave equation on manifolds with corners”, *Ann. of Math. (2)* **168**:3 (2008), 749–812. [MR](#) [Zbl](#)
- [Wang 2020] X. Wang, “Global regularity for the 3D finite depth capillary water waves”, *Ann. Sci. École Norm. Sup. (4)* **53**:4 (2020), 847–943. [MR](#) [Zbl](#)
- [Wu 1997] S. Wu, “Well-posedness in Sobolev spaces of the full water wave problem in 2-D”, *Invent. Math.* **130**:1 (1997), 39–72. [MR](#) [Zbl](#)
- [Wu 1999] S. Wu, “Well-posedness in Sobolev spaces of the full water wave problem in 3-D”, *J. Amer. Math. Soc.* **12**:2 (1999), 445–495. [MR](#) [Zbl](#)
- [Wunsch 1999] J. Wunsch, “Propagation of singularities and growth for Schrödinger operators”, *Duke Math. J.* **98**:1 (1999), 137–186. [MR](#) [Zbl](#)
- [Yosihara 1982] H. Yosihara, “Gravity waves on the free surface of an incompressible perfect fluid of finite depth”, *Publ. Res. Inst. Math. Sci.* **18**:1 (1982), 49–96. [MR](#) [Zbl](#)
- [Yosihara 1983] H. Yosihara, “Capillary-gravity waves for an incompressible ideal fluid”, *J. Math. Kyoto Univ.* **23**:4 (1983), 649–694. [MR](#) [Zbl](#)
- [Zakharov 1968] V. E. Zakharov, “Stability of periodic waves of finite amplitude on the surface of a deep fluid”, *J. Appl. Mech. Tech. Phys.* **9**:2 (1968), 190–194.
- [Zakharov 1998] V. E. Zakharov, “Weakly nonlinear waves on the surface of an ideal finite depth fluid”, pp. 167–197 in *Nonlinear waves and weak turbulence*, edited by V. E. Zakharov, Amer. Math. Soc. Transl. (2) **182**, Amer. Math. Soc., Providence, RI, 1998. [MR](#) [Zbl](#)
- [Zhu 2020] H. Zhu, “Control of three dimensional water waves”, *Arch. Ration. Mech. Anal.* **236**:2 (2020), 893–966. [MR](#) [Zbl](#)
- [Zworski 2012] M. Zworski, *Semiclassical analysis*, Grad. Stud. in Math. **138**, Amer. Math. Soc., Providence, RI, 2012. [MR](#) [Zbl](#)

Received 13 Dec 2021. Accepted 16 Jun 2022.

HUI ZHU: [zhuhui@umich.edu](mailto:zhuhui@umich.edu)

Current address: Department of Mathematics, University of Michigan, Ann Arbor, MI, United States

Laboratoire de Mathématiques d’Orsay, Université Paris-Sud, CNRS, Université Paris-Saclay, Orsay, France

# Analysis & PDE

[msp.org/apde](http://msp.org/apde)

## EDITOR-IN-CHIEF

Clément Mouhot Cambridge University, UK  
[c.mouhot@dpmms.cam.ac.uk](mailto:c.mouhot@dpmms.cam.ac.uk)

## BOARD OF EDITORS

Massimiliano Berti	Scuola Intern. Sup. di Studi Avanzati, Italy <a href="mailto:berti@sissa.it">berti@sissa.it</a>	William Minicozzi II	Johns Hopkins University, USA <a href="mailto:minicozz@math.jhu.edu">minicozz@math.jhu.edu</a>
Zbigniew Blocki	Uniwersytet Jagielloński, Poland <a href="mailto:zbigniew.blocki@uj.edu.pl">zbigniew.blocki@uj.edu.pl</a>	Werner Müller	Universität Bonn, Germany <a href="mailto:mueller@math.uni-bonn.de">mueller@math.uni-bonn.de</a>
Charles Fefferman	Princeton University, USA <a href="mailto:cf@math.princeton.edu">cf@math.princeton.edu</a>	Igor Rodnianski	Princeton University, USA <a href="mailto:irod@math.princeton.edu">irod@math.princeton.edu</a>
David Gérard-Varet	Université de Paris, France <a href="mailto:david.gerard-varet@imj-prg.fr">david.gerard-varet@imj-prg.fr</a>	Yum-Tong Siu	Harvard University, USA <a href="mailto:siu@math.harvard.edu">siu@math.harvard.edu</a>
Colin Guillarmou	Université Paris-Saclay, France <a href="mailto:colin.guillarmou@universite-paris-saclay.fr">colin.guillarmou@universite-paris-saclay.fr</a>	Terence Tao	University of California, Los Angeles, USA <a href="mailto:tao@math.ucla.edu">tao@math.ucla.edu</a>
Ursula Hamenstaedt	Universität Bonn, Germany <a href="mailto:ursula@math.uni-bonn.de">ursula@math.uni-bonn.de</a>	Michael E. Taylor	Univ. of North Carolina, Chapel Hill, USA <a href="mailto:met@math.unc.edu">met@math.unc.edu</a>
Peter Hintz	ETH Zurich, Switzerland <a href="mailto:peter.hintz@math.ethz.ch">peter.hintz@math.ethz.ch</a>	Gunther Uhlmann	University of Washington, USA <a href="mailto:gunther@math.washington.edu">gunther@math.washington.edu</a>
Vadim Kaloshin	Institute of Science and Technology, Austria <a href="mailto:vadim.kaloshin@gmail.com">vadim.kaloshin@gmail.com</a>	András Vasy	Stanford University, USA <a href="mailto:andras@math.stanford.edu">andras@math.stanford.edu</a>
Izabella Laba	University of British Columbia, Canada <a href="mailto:ilaba@math.ubc.ca">ilaba@math.ubc.ca</a>	Dan Virgil Voiculescu	University of California, Berkeley, USA <a href="mailto:dvv@math.berkeley.edu">dvv@math.berkeley.edu</a>
Anna L. Mazzucato	Penn State University, USA <a href="mailto:alm24@psu.edu">alm24@psu.edu</a>	Jim Wright	University of Edinburgh, UK <a href="mailto:j.r.wright@ed.ac.uk">j.r.wright@ed.ac.uk</a>
Richard B. Melrose	Massachusetts Inst. of Tech., USA <a href="mailto:rbm@math.mit.edu">rbm@math.mit.edu</a>	Maciej Zworski	University of California, Berkeley, USA <a href="mailto:zworski@math.berkeley.edu">zworski@math.berkeley.edu</a>
Frank Merle	Université de Cergy-Pontoise, France <a href="mailto:merle@ihes.fr">merle@ihes.fr</a>		

## PRODUCTION

[production@msp.org](mailto:production@msp.org)

Silvio Levy, Scientific Editor

---

See inside back cover or [msp.org/apde](http://msp.org/apde) for submission instructions.

---

The subscription price for 2024 is US \$440/year for the electronic version, and \$690/year (+\$65, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues from the last three years and changes of subscriber address should be sent to MSP.

---

Analysis & PDE (ISSN 1948-206X electronic, 2157-5045 printed) at Mathematical Sciences Publishers, 798 Evans Hall #3840, c/o University of California, Berkeley, CA 94720-3840, is published continuously online.

---

APDE peer review and production are managed by EditFlow<sup>®</sup> from MSP.

PUBLISHED BY

 **mathematical sciences publishers**  
nonprofit scientific publishing

<http://msp.org/>

© 2024 Mathematical Sciences Publishers

# ANALYSIS & PDE

Volume 17 No. 1 2024

---

The prescribed curvature problem for entire hypersurfaces in Minkowski space CHANGYU REN, ZHIZHANG WANG and LING XIAO	1
Anisotropic micropolar fluids subject to a uniform microtorque: the stable case ANTOINE REMOND-TIEDREZ and IAN TICE	41
Strong ill-posedness for SQG in critical Sobolev spaces IN-JEE JEONG and JUNHA KIM	133
Large-scale regularity for the stationary Navier–Stokes equations over non-Lipschitz boundaries MITSUO HIGAKI, CHRISTOPHE PRANGE and JINPING ZHUGE	171
On a family of fully nonlinear integrodifferential operators: from fractional Laplacian to non-local Monge–Ampère LUIS A. CAFFARELLI and MARÍA SORIA-CARRO	243
Propagation of singularities for gravity-capillary water waves HUI ZHU	281
Shift equivalences through the lens of Cuntz–Krieger algebras TOKE MEIER CARLSEN, ADAM DOR-ON and SØREN EILERS	345